

FACTOR VIII POLYPEPTIDE FORMULATIONS

BACKGROUND OF THE INVENTION

Field of the Invention

[0001] The present invention relates generally to the field of therapeutics for hemostatic disorders.

Background Art

[0002] Hemophilia A is an X-linked bleeding disorder caused by mutations and/or deletions in the factor VIII (FVIII) gene resulting in a deficiency of FVIII activity (Peyvandi, F. *et al. Haemophilia* 12:82-89 (2006)). The disease is characterized by spontaneous hemorrhage and excessive bleeding after trauma. Over time, the repeated bleeding into muscles and joints, which often begins in early childhood, results in hemophilic arthropathy and irreversible joint damage. This damage is progressive and can lead to severely limited mobility of joints, muscle atrophy and chronic pain (Rodriguez-Merchan, E.C., *Semin. Thromb. Hemost.* 29:87-96 (2003)), which is herein incorporated by reference in its entirety).

[0003] The A2 domain is necessary for the procoagulant activity of the factor VIII molecule. Studies show that porcine factor VIII has six-fold greater procoagulant activity than human factor VIII (Lollar, P., and E. T. Parker, *J. Biol. Chem.* 266:12481-12486 (1991)), and that the difference in coagulant activity between human and porcine factor VIII appears to be based on a difference in amino acid sequence between one or more residues in the human and porcine A2 domains (Lollar, P., *et al.*, *J. Biol. Chem.* 267:23652-23657 (1992)), incorporated herein by reference in its entirety.

[0004] Treatment of hemophilia A is by replacement therapy targeting restoration of FVIII activity to 1 to 5 % of normal levels to prevent spontaneous bleeding (Mannucci, P.M., *et al.*, *N. Engl. J. Med.* 344:1773-1779 (2001)), which is herein incorporated by reference in its entirety). There are plasma-derived and recombinant FVIII products available to treat bleeding episodes on-demand or to prevent bleeding episodes from occurring by treating prophylactically.

[0005] Reduced mortality, prevention of joint damage, and improved quality of life have been important achievements due to the development of plasma-derived and recombinant

FVIII. Prolonged protection from bleeding would represent another key advancement in the treatment of hemophilia A patients.

BRIEF SUMMARY OF THE INVENTION

[0006] The present invention comprises a pharmaceutical composition comprising: (a) a FVIII polypeptide; (b) one or more stabilizing agents selected from sucrose, trehalose, raffinose, arginine, or mixture thereof; (c) sodium chloride (NaCl); (d) L-histidine; (e) calcium chloride; and (f) polysorbate 20 or polysorbate 80. In one embodiment, mannitol, glycine, alanine, or hydroxyethyl starch is not included in the pharmaceutical composition. In another embodiment, NaCl is the only bulking agent. In other embodiments, the FVIII polypeptide is a long-acting polypeptide or a short-acting polypeptide.

[0007] In other embodiments, the pharmaceutical composition comprises about 1% (w/v) to about 2.5% (w/v) sucrose, about 1.3% (w/v) sucrose to about 2.0% (w/v) sucrose, about 1.33% (w/v) sucrose or about 2.0% (w/v) sucrose, about 10 mg/ml to about 25 mg/ml sucrose, about 13 mg/ml to about 20 mg/ml sucrose, about 13.3 mg/ml sucrose, or about 20.0 mg/ml sucrose. In still other embodiments, the pharmaceutical composition comprises about 150 mM to about 250 mM NaCl, about 175 mM to about 225 mM NaCl, about 200 mM to about 210 mM NaCl, about 205 mM NaCl, about 8.8 mg/ml to about 14.6 mg/ml NaCl, about 10 mg/ml to about 13 mg/ml NaCl, about 12.0 mg/ml NaCl, about 250 mM to about 350 mM NaCl, about 275 mM to about 325 mM NaCl, about 308 mM NaCl, about 14.6 mg/ml to about 20.5 mg/ml NaCl, about 16 mg/ml to about 19 mg/ml NaCl, or about 18.0 mg/ml NaCl.

[0008] In yet other embodiments, the pharmaceutical composition comprises about 5 mM to about 15 mM L-histidine, about 6.64 mM L-histidine to about 9.8 mM L-histidine, about 0.75 mg/ml to about 2.25 mg/ml L-histidine, or about 1.03 mg/ml L-histidine to about 1.55 mg/ml L-histidine. In certain embodiments, the pharmaceutical composition comprises about 5 mM to about 10 mM calcium chloride, about 5.4 mM calcium chloride to about 8 mM calcium chloride, about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate, or about 0.8 mg/ml calcium chloride dihydrate to about 1.18 mg/ml calcium chloride dihydrate.

- [0009] In some embodiments, the pharmaceutical composition comprises about 0.008% (w/v) to about 0.025% (w/v) polysorbate 20 or polysorbate 80, about 0.013% (w/v) polysorbate 20 or polysorbate 80 or about 0.02 % (w/v) polysorbate 20 or polysorbate 80, about 0.08 mg/ml to about 0.25 mg/ml polysorbate 20 or polysorbate 80, about 0.13% mg/ml polysorbate 20 or polysorbate 80 or about 0.20 mg/ml polysorbate 20 or polysorbate 80.
- [0010] In other embodiments, the pharmaceutical composition comprises the rFVIII_{IFc} polypeptide, which comprises a first subunit comprising an amino acid sequence at least 90% or 95% identical to amino acids 20 to 1684 of SEQ ID NO:2 or 20 to 2578 of SEQ ID NO:6, and a second subunit comprising an amino acid sequence at least 90% to 95% identical to amino acids 21 to 247 of SEQ ID NO:4. In certain embodiments, wherein the rFVIII_{IFc} polypeptide comprises a first subunit comprising amino acids 20 to 1684 of SEQ ID NO:2 or 20 to 2578 of SEQ ID NO:6, and a second subunit comprising amino acids 21 to 247 of SEQ ID NO:4.
- [0011] In certain embodiments, a FVIII polypeptide is present in a pharmaceutical composition at a concentration of about 50 IU/ml to about 2500 IU/ml, *e.g.*, 83 IU/ml, 167 IU/ml, 250 IU/ml, 333 IU/ml, 500 IU/ml, 667 IU/ml, 1000 IU/ml, 1333 IU/ml, 1667 IU/ml, or 2000 IU/ml of the FVIII polypeptide. In some embodiments, a FVIII polypeptide is present in a pharmaceutical composition at a concentration of about 100 IU/ml to about 4000 IU/ml, *e.g.*, 150 IU/ml, 287.5 IU/ml, 431.25 IU/ml, 575 IU/ml, 862.5 IU/ml, 1150 IU/ml, 1725 IU/ml, 2300 IU/ml, 2875 IU/ml, or 3450 IU/ml of the FVIII polypeptide.
- [0012] In other embodiments, a pharmaceutical composition comprises: (a) about 50 IU/ml to about 2500 IU/ml of a long-acting FVIII polypeptide; (b) about 1% (w/v) to about 2.5% (w/v) of sucrose; (c) about 150 mM to about 250 mM NaCl; (d) about 5 mM to about 15 mM L-histidine; (e) about 5 mM to about 10 mM calcium chloride; and (f) about 0.008% (w/v) to about 0.025% of polysorbate 20 or polysorbate 80. In other embodiments, the pharmaceutical composition comprises about 175 mM to about 225 mM NaCl or about 200 mM to about 210 mM NaCl.
- [0013] In yet other embodiments, a pharmaceutical composition comprises (a) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, about 1000 IU/ml, about 1333 IU/ml, about 1667 IU/ml, or about 2000 IU/ml of a FVIII polypeptide; (b) about 1.33% (w/v) of sucrose; (c) about 205 mM NaCl; (d) about

6.64 mM L-histidine; (e) about 5.4 mM calcium chloride; and (f) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

[0014] In some embodiments, a pharmaceutical composition comprises: (a) about 100 IU/ml to about 4000 IU/ml of a FVIII polypeptide; (b) about 1% (w/v) to about 2.5% (w/v) of sucrose; (c) about 250 mM to about 350 mM NaCl; (d) about 5 mM to about 15 mM L-histidine; (e) about 5 mM to about 10 mM calcium chloride; and (f) about 0.008% (w/v) to about 0.025% of polysorbate 20 or polysorbate 80. In other embodiments, the pharmaceutical composition comprises about 275 mM to about 325 mM NaCl.

[0015] In certain embodiments, a pharmaceutical composition comprises: (a) about 150 IU/ml, about 287.5 IU/ml, about 431.25 IU/ml, about 575 IU/ml, about 862.5 IU/ml, about 1150 IU/ml, about 1725 IU/ml, about 2300 IU/ml, about 2875 IU/ml, or about 3450 IU/ml of a FVIII polypeptide; (b) about 2.0% (w/v) of sucrose; (c) about 308 mM NaCl; (d) about 9.8 mM L-histidine; (e) about 8 mM calcium chloride; and (f) about 0.020% (w/v) of polysorbate 20 or polysorbate 80. In other embodiments, a pharmaceutical composition comprises: (a) about 50 IU/ml to about 2500 IU/ml of a FVIII polypeptide; (b) about 10 mg/ml to about 25 mg/ml of sucrose; (c) about 8.8 mg/ml to about 14.6 mg/ml NaCl; (d) about 0.75 mg/ml to about 2.25 mg/ml L-histidine; (e) about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate; and (f) about 0.08 mg/ml to about 0.25 mg/ml of polysorbate 20 or polysorbate 80. In some embodiments, a pharmaceutical composition comprises: (a) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, about 1000 IU/ml, about 1333 IU/ml, about 1667 IU/ml, or about 2000 IU/ml of a FVIII polypeptide; (b) about 13.3 mg/ml of sucrose; (c) about 12.0 mg/ml NaCl; (d) about 1.03 mg/ml L-histidine; (e) about 0.8 mg/ml calcium chloride dihydrate; and (f) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

[0016] In other embodiments, a pharmaceutical composition comprises: (a) about 100 IU/ml to about 4000 IU/ml of a FVIII polypeptide; (b) about 10 mg/ml to about 25 mg/ml of sucrose; (c) about 14.6 mg/ml to about 20.5 mg/ml NaCl; (d) about 0.75 mg/ml to about 2.25 mg/ml L-histidine; (e) about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate; and (f) about 0.08 mg/ml to about 0.25 mg/ml of polysorbate 20 or polysorbate 80. In certain embodiments, a pharmaceutical composition comprises: (a) about 150 IU/ml, about 287.5 IU/ml, about 431.25 IU/ml, about 575 IU/ml, about 862.5 IU/ml, about 1150 IU/ml, about 1725 IU/ml, about 2300 IU/ml, about 2875 IU/ml, or about 3450

IU/ml of a FVIII polypeptide; (b) about 20.0 mg/ml of sucrose; (c) about 18.0 mg/ml NaCl; (d) about 1.55 mg/ml L-histidine; (e) about 1.18 mg/ml calcium chloride dihydrate; and (f) about 0.20 mg/ml of polysorbate 20 or polysorbate 80.

[0017] In some embodiments, the present invention includes a pharmaceutical kit comprising: (a) a first container comprising a lyophilized powder, wherein the powder comprises (i) a FVIII polypeptide, (ii) one or more stabilizing agents selected from sucrose, trehalose, raffinose, arginine, or mixture thereof; (iii) sodium chloride (NaCl); (iv) L-histidine; (v) calcium chloride; and (vi) polysorbate 20 or polysorbate 80; and (b) a second container comprising sterilized water for injections to be combined with the lyophilized powder of the first container. In other embodiments, a pharmaceutical kit comprises: (a) a first container comprising a lyophilized powder, where the powder comprises (i) about 250 IU, about 500 IU, about 750 IU, about 1000 IU, about 1500 IU, about 2000 IU, about 3000 IU, about 4000 IU, about 5000 IU, or about 6000 IU of a FVIII polypeptide, (ii) about 40 mg of sucrose; (iii) about 36 mg of sodium chloride; (iv) about 3.1 mg of L-histidine; (v) about 2.40 mg of calcium chloride dihydrate; and (v) about 0.40 mg of polysorbate 20 or polysorbate 80; and (b) a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, about 1000 IU/ml, about 1333 IU/ml, about 1667 IU/ml, or about 2000 IU/ml of a FVIII polypeptide, respectively; (ii) about 1.33% (w/v) of sucrose; (iii) about 205 mM NaCl; (iv) about 6.64 mM L-histidine; (v) about 5.4 mM of calcium chloride; and (vi) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

[0018] In other embodiments, a pharmaceutical kit comprises: (a) a first container comprising a lyophilized powder, where the powder comprises (i) about 250 IU, about 500 IU, about 750 IU, about 1000 IU, about 1500 IU, about 2000 IU, about 3000 IU, about 4000 IU, about 5000 IU, or about 6000 IU of a FVIII polypeptide, (ii) about 40 mg of sucrose; (iii) about 36 mg of sodium chloride; (iv) about 3.1 mg of L-histidine; (v) about 2.40 mg of calcium chloride dihydrate; and (v) about 0.40 mg of polysorbate 20 or polysorbate 80; and (b) a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, about 1000 IU/ml, about 1333

IU/ml, about 1667 IU/ml, or about 2000 IU/ml of a FVIII polypeptide, respectively; (ii) about 13.3 mg/ml of sucrose; (iii) about 12.0 mg/ml of NaCl; (iv) about 1.03 mg/ml of L-histidine; (v) about 0.80 mg/ml of calcium chloride; and (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

[0019] The invention also includes a method of reducing or decreasing a bleeding episode or treating or preventing a bleeding condition comprising administering a pharmaceutical composition comprising a FVIII polypeptide.

BRIEF DESCRIPTION OF THE DRAWINGS

[0020] Figure 1 shows details on the design of the sequential PK subgroup (Arm 1) dosing and PK sampling.

[0021] Figure 2 shows the mean observed FVIII activity with rFVIII-Fc and rFVIII over time: one stage clotting assay (logarithmic scale) in the sequential pharmacokinetic subgroup. rFVIII-Fc demonstrated an approximate 50% longer elimination half-life and mean residence time compared with wild-type Factor VIII (a short-acting FVIII polypeptide) ($P < 0.001$).

DETAILED DESCRIPTION OF THE INVENTION

[0022] The present invention provides a formulation for a Factor VIII polypeptide, *e.g.*, FVIII-Fc, and methods of using thereof.

I. Definitions

[0023] It must be noted that, as used in this specification and the appended claims, the singular forms "a", "an" and "the" include plural referents unless the context clearly dictates otherwise. The terms "a" (or "an"), as well as the terms "one or more," and "at least one" can be used interchangeably herein.

[0024] The term "about" is used herein to mean approximately, roughly, around, or in the regions of. When the term "about" is used in conjunction with a numerical range, it modifies that range by extending the boundaries above and below the numerical values set forth. In general, the term "about" is used herein to modify a numerical value above and below the stated value by a variance of 10 percent, up or down (higher or lower).

[0025] Furthermore, "and/or" where used herein is to be taken as specific disclosure of each of the two specified features or components with or without the other. Thus, the term "and/or" as used in a phrase such as "A and/or B" herein is intended to include "A and B," "A or B," "A" (alone), and "B" (alone). Likewise, the term "and/or" as used in a phrase such as "A, B, and/or C" is intended to encompass each of the following embodiments: A, B, and C; A, B, or C; A or C; A or B; B or C; A and C; A and B; B and C; A (alone); B (alone); and C (alone).

[0026] It is understood that wherever embodiments are described herein with the language "comprising," otherwise analogous embodiments described in terms of "consisting of" and/or "consisting essentially of" are also provided.

[0027] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure is related. For example, the Concise Dictionary of Biomedicine and Molecular Biology, Juo, Pei-Show, 2nd ed., 2002, CRC Press; The Dictionary of Cell and Molecular Biology, 3rd ed., 1999, Academic Press; and the Oxford Dictionary Of Biochemistry And Molecular Biology, Revised, 2000, Oxford University Press, provide one of skill with a general dictionary of many of the terms used in this disclosure.

[0028] Units, prefixes, and symbols are denoted in their Système International de Unites (SI) accepted form. Numeric ranges are inclusive of the numbers defining the range. Unless otherwise indicated, amino acid sequences are written left to right in amino to carboxy orientation. The headings provided herein are not limitations of the various aspects or embodiments of the disclosure, which can be had by reference to the specification as a whole. Accordingly, the terms defined immediately below are more fully defined by reference to the specification in its entirety. Amino acids are referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission. Nucleotides, likewise, are referred to by their commonly accepted single-letter codes.

[0029] The term "polypeptide," "peptide" and "protein" are used interchangeably and refer to a polymeric compound comprised of covalently linked amino acid residues.

[0030] The term "polynucleotide" and "nucleic acid" are used interchangeably and refer to a polymeric compound comprised of covalently linked nucleotide residues. Polynucleotides can be DNA, cDNA, RNA, single stranded, or double stranded, vectors, plasmids, phage, or viruses. Polynucleotides include those in Table 10, which encode the

polypeptides of Table 11 (see Table 10). Polynucleotides also include fragments of the polynucleotides of Table 10, *e.g.*, those that encode fragments of the polypeptides of Table 11, such as the Factor VIII, Fc, signal sequence, propeptide, and other fragments of the polypeptides of Table 11.

[0031] "Administering," as used herein, means to proscribe or give a pharmaceutical composition comprising a Factor VIII polypeptide of the invention to a subject via a pharmaceutically acceptable route. Examples of routes of administration include, but are not limited to, intravenous, *e.g.*, intravenous injection and intravenous infusion, *e.g.*, via central venous access. Additional routes of administration include subcutaneous, intramuscular, oral, nasal, and pulmonary administration. Advantages of the present invention include: improved regimen compliance; reduced break through bleeds; increased protection of joints from bleeds; prevention of joint damage; reduced morbidity; reduced mortality; prolonged protection from bleeding; decreased thrombotic events; and improved quality of life.

[0032] The terms "long-acting" and "long-lasting" are used interchangeably herein. In one embodiment, the term "long-acting" or "long-lasting" indicates that a FVIII activity as a result of administration of the "long-acting" FVIII polypeptide is longer than the FVIII activity of a wild-type FVIII (*e.g.*, ADVATE® or plasma-derived FVIII ("pdFVIII")). The "longer" FVIII activity can be measured by any known methods in the art, *e.g.*, aPTT assay, chromogenic assay, ROTEM®, TGA, etc. In one embodiment, the "longer" FVIII activity can be shown by the $T_{1/2\beta}$ (activity). In another embodiment, the "longer" FVIII activity can be shown by the level of FVIII antigen present in plasma, *e.g.*, by the $T_{1/2\beta}$ (antigen). In other embodiments, the long-acting or long-lasting FVIII polypeptide works longer in a coagulation cascade, *e.g.*, is active for a longer period, compared to a wild-type FVIII polypeptide, *i.e.*, a polypeptide consisting of amino acids 20 to 1457 of SEQ ID NO: 2, *i.e.*, SQ BDD FVIII (REFACTO®) or a polypeptide consisting of amino acids 20 to 2351 of SEQ ID NO: 6 (ADVATE®).

[0033] The terms "short-acting" and "short-lasting" are used interchangeably herein. The term "short-acting" or "short-lasting" indicates that a FVIII activity as a result of administration of the "short-acting" FVIII polypeptide is similar to or the same as the FVIII activity of a wild-type FVIII (*e.g.*, ADVATE® (amino acids 20 to 2351 of SEQ ID NO: 6), REFACTO® (amino acids 20 to 1457 of SEQ ID NO: 2), or plasma-derived FVIII ("pdFVIII")) or shorter than the FVIII activity of a long-acting FVIII polypeptide. The

"shorter" FVIII activity can be measured by any known methods in the art, *e.g.*, aPTT assay, chromogenic assay, ROTEM[®], TGA, etc. Exemplary short-acting FVIII polypeptides include, but are not limited to, wild-type mature FVIII polypeptide (ADVATE[®], RECOMBINATE[®], or HELIXATE[®]) or B-domain deleted FVIII polypeptides such as SQ BDD FVIII (REFACTO[®] and XYNTHA[®]) or FVIII polypeptides containing 21 amino acids from B-domain (*i.e.*, SFSQNSRHPSQNPPVLKRHQR, SEQ ID NO: 17) (*e.g.*, NOVOEIGHT[®]).

[0034] The term "chimeric polypeptide," as used herein, means a polypeptide that includes within it at least two polypeptides (or portions thereof such as subsequences or peptides) from different sources. Chimeric polypeptides can include two, three, four, five, six, seven, or more polypeptides or portions thereof from different sources, such as different genes, different cDNAs, or different animal or other species. Chimeric polypeptides can include one or more linkers joining the different polypeptides or portions thereof. Thus, the polypeptides or portions thereof can be joined directly or they can be joined indirectly, via linkers, or both, within a single chimeric polypeptide. Chimeric polypeptides can include additional peptides such as signal sequences and sequences such as 6His and FLAG that aid in protein purification or detection. In addition, chimeric polypeptides can have amino acid or peptide additions to the N- and/or C-termini. In some embodiments, a chimeric polypeptide is a long-acting FVIII polypeptide. Exemplary chimeric polypeptides of the invention are Factor VIII-FcRn BP chimeric polypeptides, *e.g.*, Factor VIII-Fc polypeptides such as the FVIII-Fc, SEQ ID NO:2 or 6 (Table 11) with or without its signal sequence and propeptide.

[0035] "Hybrid" polypeptides and proteins, as used herein, means a combination of a chimeric polypeptide with a second polypeptide. The chimeric polypeptide and the second polypeptide in a hybrid can be associated with each other via protein-protein interactions, such as charge-charge or hydrophobic interactions. The chimeric polypeptide and the second polypeptide in a hybrid can be associated with each other via disulfide or other covalent bond(s). Hybrids are described in WO 2004/101740 and WO 2006/074199, each of which is incorporated herein by reference in its entirety. See also U.S. Patent Nos. 7,404,956 and 7,348,004, each of which is incorporated herein by reference in its entirety. The second polypeptide can be a second copy of the same chimeric polypeptide or it can be a non-identical chimeric polypeptide. See, *e.g.*, Table 11. In one embodiment, the second polypeptide is a polypeptide comprising an FcRn

binding partner. FcRn binding partners binds to FcRn and protects the FcRn binding partner containing molecule from catabolism, thus extending the plasma half-life. In another embodiment, the chimeric polypeptide is a chimeric Factor VIII-Fc polypeptide and the second polypeptide consists essentially of Fc, *e.g.*, a rFVIII-Fc recombinant fusion protein consisting of a single molecule of recombinant B-domain deleted human FVIII (BDD-rFVIII) fused to the dimeric Fc domain of the human IgG1, with no intervening linker sequence. This hybrid polypeptide is referred to herein as FVIII-Fc monomeric Fc fusion protein, FVIII-Fc monomer hybrid, monomeric FVIII-Fc hybrid, and FVIII-Fc monomer-dimer. See Table 11A. The Examples provide preclinical and clinical data for this hybrid polypeptide.

[0036] The second polypeptide in a hybrid can comprise or consist essentially of a sequence at least 90% or 95% identical to the amino acid sequence shown in Table 11A(ii) without a signal sequence (amino acids 21 to 247 of SEQ ID NO:4) or at least 90% or 95% identical to the amino acid sequence shown in Table 11A(ii) with a signal sequence (amino acids 1 to 247 of SEQ ID NO:4). The second polypeptide can comprise or consist essentially of a sequence identical to the amino acid sequence shown in Table 11A(ii) without a signal sequence (amino acids 21 to 247 of SEQ ID NO:4) or identical to the amino acid sequence shown in Table 11A(ii) with a signal sequence (amino acids 1 to 247 of SEQ ID NO:4).

[0037] Factor VIII coagulant activity is expressed as International Unit(s) (IU). One IU of Factor VIII activity corresponds approximately to the quantity of Factor VIII in one milliliter of normal human plasma. Several assays are available for measuring Factor VIII activity, including the one stage clotting assay (activated partial thromboplastin time; aPTT), thrombin generation time (TGA) and rotational thromboelastometry (ROTEM®).

[0038] The term "lyophilisate" as used herein in connection with the formulation according to the invention denotes a formulation which is manufactured by freeze-drying methods known in the art per se. The solvent (*e.g.* water) is removed by freezing following sublimation under vacuum and desorption of residual water at elevated temperature. In the pharmaceutical field, the lyophilisate has usually a residual moisture of about 0.1 to 5% (w/w) and is present as a powder or a physical stable cake. The lyophilisate is characterized by a fast dissolution after addition of a reconstitution medium.

[0039] The term "reconstituted formulation" as used herein denotes a formulation which is lyophilized and re-dissolved by addition of a diluent. The diluent can contain, without limitation, water for injection (WFI), bacteriostatic water for injection (BWFI), sodium chloride solutions (*e.g.* 0.9% (w/v) NaCl), glucose solutions (*e.g.* 5% glucose), surfactant containing solutions (*e.g.* 0.01% polysorbate 20 or polysorbate 80), a pH-buffered solution (*e.g.* phosphate-buffered solutions) and combinations thereof.

[0040] "Dosing interval," as used herein, means the amount of time that elapses between multiple doses being administered to a subject. Dosing interval can thus be indicated as ranges. The dosing interval in the methods of the invention using a chimeric FVIII-FcRn BP, *e.g.*, a chimeric FVIII -Fc, can be at least about one and one-half to eight times longer than the dosing interval required for an equivalent amount (in IU/kg) of the Factor VIII without the FcRn BP, *e.g.*, Fc portion (*i.e.*, a polypeptide consisting of said FVIII). The dosing interval when administering, *e.g.*, a Factor VIII-Fc chimeric polypeptide (or a hybrid) of the invention can be at least about one and one-half times longer than the dosing interval required for an equivalent amount of the Factor VIII without the FcRn BP, *e.g.*, Fc portion (*i.e.*, a polypeptide consisting of the Factor VIII). The dosing interval can be at least about one and one-half to eight times longer than the dosing interval required for an equivalent amount of the Factor VIII without, *e.g.*, the Fc portion (or a polypeptide consisting of the Factor VIII).

[0041] The term "dosing frequency" as used herein refers to the frequency of administering doses of a FVIII polypeptide in a given time. Dosing frequency can be indicated as the number of doses per a given time, *e.g.*, once a week or once in two weeks.

[0042] The term "prophylaxis of one or more bleeding episode" or "prophylactic treatment" as used herein means administering a FVIII polypeptide in multiple doses to a subject over a course of time to increase the level of FVIII activity in a subject's plasma. In one embodiment, "prophylaxis of one or more bleeding episode" indicates use of a FVIII polypeptide to prevent or inhibit occurrence of one or more spontaneous or uncontrollable bleeding or bleeding episodes or to reduce the frequency of one or more spontaneous or uncontrollable bleeding or bleeding episodes. In another embodiment, the increased FVIII activity level is sufficient to decrease the incidence of spontaneous bleeding or to prevent bleeding in the event of an unforeseen injury. Prophylactic treatment decreases or prevents bleeding episodes, for example, those described under on-

demand treatment. Prophylactic treatment can be individualized, as discussed under "dosing interval", *e.g.*, to compensate for inter-patient variability.

[0043] The term "about twice weekly" as used herein means approximate number, and "about twice weekly" can include twice in one week, *e.g.*, a first dose in three days and a second dose in three days, a first dose in three days and a second dose in four days, a first dose in four days and a second dose in three days, a first dose in four days and a second dose in four days. The term "about twice weekly" can also include every three days, every four days, or every five days.

[0044] The term "about once a week" as used herein means approximate number, and "about once a week" can include every seven days $\pm$ two days, *i.e.*, every five days to every nine days. The dosing frequency of "once a week" thus can be every five days, every six days, every seven days, every eight days, or every nine days.

[0045] The term "individualized prophylaxis" or "prophylactic and individualized" as used herein means use of a FVIII polypeptide for an individualized dosing and/or dosing interval or frequency to prevent or inhibit occurrence of one or more spontaneous and/or uncontrollable bleeding or bleeding episodes or to reduce the frequency of one or more spontaneous and/or uncontrollable bleeding or bleeding episodes. "Individualized" within the context of prophylaxis is used synonymous with "tailored" throughout this application. For example, "individualized prophylaxis" also means "tailored prophylaxis" and "prophylactic and individualized" also means "prophylactic and tailored." In one embodiment, the "individualized interval" includes every 3 days $\pm$ 2 days, *i.e.*, every day to every five days. The dosing frequency of the "individualized interval prophylaxis" thus can be every day, every two days, every three days, every four days, or every five days.

[0046] The term "on-demand treatment," as used herein, means treatment that is intended to take place over a short course of time and is in response to an existing condition, such as a bleeding episode, or a perceived short term need such as planned surgery. The "on-demand treatment" is used interchangeably with "episodic" treatment. Conditions that can require on-demand treatment include a bleeding episode, hemarthrosis, muscle bleed, oral bleed, hemorrhage, hemorrhage into muscles, oral hemorrhage, trauma, trauma capitis, gastrointestinal bleeding, intracranial hemorrhage, intra-abdominal hemorrhage, intrathoracic hemorrhage, bone fracture, central nervous system bleeding, bleeding in the retropharyngeal space, bleeding in the retroperitoneal space, or bleeding in the iliopsoas

sheath. Bleeding episodes other than these are also included. The subject can be in need of surgical prophylaxis, peri-operative management, or treatment for surgery. Such surgeries include minor surgery, major surgery, tooth extraction, tonsillectomy, other dental/thoraco-facial surgeries, inguinal herniotomy, synovectomy, total knee replacement, other joint replacement, craniotomy, osteosynthesis, trauma surgery, intracranial surgery, intra-abdominal surgery, or intrathoracic surgery. Surgeries other than these are also included.

[0047] Additional conditions that can require on-demand treatment include minor hemorrhage, hemarthroses, superficial muscle hemorrhage, soft tissue hemorrhage, moderate hemorrhage, intramuscle or soft tissue hemorrhage with dissection, mucous membrane hemorrhage, hematuria, major hemorrhage, hemorrhage of the pharynx, hemorrhage of the retropharynx, hemorrhage of the retroperitoneum, hemorrhage of the central nervous system, bruises, cuts, scrapes, joint hemorrhage, nose bleed, mouth bleed, gum bleed, intracranial bleeding, intraperitoneal bleeding, minor spontaneous hemorrhage, bleeding after major trauma, moderate skin bruising, or spontaneous hemorrhage into joints, muscles, internal organs or the brain. Additional reasons for on-demand treatment include the need for peri-operative management for surgery or dental extraction, major surgery, extensive oral surgery, urologic surgery, hernia surgery, orthopedic surgery such as replacement of knee, hip, or other major joint.

[0048] The term "treatment" or "treating" as used herein means amelioration or reduction of one or more symptoms of bleeding diseases or disorders including, but not limited to, hemophilia B. In one embodiment, "treatment of" or "treating" a bleeding disease or disorder includes prevention of one or more symptoms of a bleeding disease or disorder. In a bleeding disease or disorder caused by a FVIII deficiency (*e.g.*, a low baseline FVIII activity), the term "treatment" or "treating" means a FVIII replacement therapy. By administering a pharmaceutical composition comprising a FVIII polypeptide to a subject, the subject can achieve and/or maintain a plasma trough level of a FVIII activity at about 1 IU/dl or above 1 IU/dl. In other embodiments, "treatment" or "treating" means reduction of the frequency of one or more symptoms of bleeding diseases or disorders, *e.g.*, spontaneous or uncontrollable bleeding episodes. "Treatment," however, need not be a cure.

[0049] The term "perioperative management" as used herein means use of a pharmaceutical composition comprising a FVIII polypeptide before, concurrently with, or

after an operative procedure, *e.g.*, a surgical operation. The use for "perioperative management" of one or more bleeding episode includes surgical prophylaxis before (*i.e.*, preoperative), during (*i.e.*, intraoperative), or after (*i.e.*, postoperative) a surgery to prevent one or more bleeding or bleeding episode or reducing or inhibiting spontaneous and/or uncontrollable bleeding episodes before, during, and after a surgery.

[0050] "Baseline," as used herein, is the lowest measured plasma Factor VIII level in a subject prior to administering a dose. The FVIII plasma levels can be measured at two time points prior to dosing: at a screening visit and immediately prior to dosing. Alternatively, (a) the baseline in patients whose pretreatment FVIII activity is <1%, who have no detectable FVIII antigen, and have nonsense genotypes can be defined as 0%, (b) the baseline for patients with pretreatment FVIII activity <1% and who have detectable FVIII antigen can be set at 0.5%, (c) the baseline for patients whose pretreatment FVIII activity is between 1 – 2% is C_{min} (the lowest activity throughout the PK study), and (d) the baseline for patients whose pretreatment FVIII activity is ≥2% can be set at 2%. Activity above the baseline pre-dosing can be considered residue drug from prior treatment, and can be decayed to baseline and subtracted from the PK data following rFVIII-Fc dosing.

[0051] " $T_{1/2\beta}$," or " $T_{1/2\text{ beta}}$ " or "Beta HL," as used herein, is half-life associated with elimination phase, $t_{1/2\beta} = (\ln 2) / \text{elimination rate constant}$ associated with the terminal phase. The $T_{1/2\text{ beta}}$ can be measured by FVIII activity or by FVIII antigen level in plasma. The $T_{1/2\text{ beta}}$ based on activity is shown as $T_{1/2\text{ beta}}$ (activity), and the $T_{1/2\text{ beta}}$ based on the FVIII antigen level can be shown as $T_{1/2\text{ beta}}$ (antigen). Both $T_{1/2\text{ beta}}$ (activity) and $T_{1/2\text{ beta}}$ (antigen) can be shown as ranges or a geometric mean.

[0052] "Trough," as used herein, is the lowest plasma FVIII activity level reached after administering a dose of chimeric polypeptide of the invention or another FVIII molecule and before the next dose is administered, if any. Trough is used interchangeably herein with "threshold." Baseline FVIII levels are subtracted from measured FVIII levels to calculate the trough level.

[0053] The term "annualized bleeding rate" ("ABR") as used herein refers to the number of bleeding episodes (including spontaneous and traumatic bleeds) experienced by a subject during a defined time period, extrapolated to 1 year. For example two bleeds in six months would indicate an ABR of four. The median ABR provides the middle value

among all observed ABRs, indicating that half of the subjects had individual ABRs less than or equal to the median and half had ABRs greater than or equal to the median.

[0054] The term "inter quartile range" ("IQR") as used herein refers to a measure of statistical dispersion, being equal to the difference between the upper and lower quartiles. Unlike (total) range, the interquartile range is a robust statistic, having a breakdown point of 25%, and is thus often preferred to the total range. For a symmetric distribution (where the median equals the midline, the average of the first and third quartiles), half the IQR equals the median absolute deviation (MAD). The median is the corresponding measure of central tendency.

[0055] "Subject," as used herein means a human. Subject as used herein includes an individual who is known to have at least one incidence of uncontrolled bleeding episodes, who has been diagnosed with a disease or disorder associated with uncontrolled bleeding episodes, *e.g.*, a bleeding disease or disorder, *e.g.*, hemophilia A, who is susceptible to uncontrolled bleeding episodes, *e.g.*, hemophilia, or any combinations thereof. Subjects can also include an individual who is in danger of one or more uncontrollable bleeding episodes prior to a certain activity, *e.g.*, a surgery, a sport activity, or any strenuous activities. The subject can have a baseline FVIII activity less than 1%, less than 0.5%, less than 2%, less than 2.5%, less than 3%, or less than 4%. Subjects also include pediatric humans. Pediatric human subjects are birth to 20 years, preferably birth to 18 years, birth to 16 years, birth to 15 years, birth to 12 years, birth to 11 years, birth to 6 years, birth to 5 years, birth to 2 years, or 2 to 11 years of age.

[0056] "Therapeutic dose," "dose," "effective dose," or "dosing amount" as used herein, means a dose that achieves a plasma trough level of a FVIII activity at least about 1 IU/dl or above 1 IU/dl in the subject administered with a pharmaceutical composition comprising a FVIII polypeptide. For the purpose of this invention, in one embodiment, the "dose" refers to the amount of the doses that a plasma trough level of a FVIII activity is maintained at least about 1 IU/dl or above 1 IU/dl, at least about 2 IU/dl or above 2 IU/dl, at least about 3 IU/dl or above 3 IU/dl, at least about 4 IU/dl or above 4 IU/dl, or at least about 5 IU/dl or above 5 IU/dl throughout the administration of a pharmaceutical composition comprising a FVIII polypeptide. In another embodiment, the "dose" reduces or decreases frequency of bleeding or bleeding disorder. In other embodiments, the "dose" stops on-going, uncontrollable bleeding or bleeding episodes. In still other embodiments, the "dose" prevents spontaneous bleeding or bleeding episodes in a subject

susceptible to such spontaneous bleeding or bleeding episodes. The "dose" or "therapeutic dose" need not cure hemophilia.

[0057] The term "target joint" is defined as a major joint (*e.g.*, hip, elbow, wrist, shoulder, knee, and ankle) into which repeated bleeding occurs (frequency of ≥ 3 bleeding episodes into the same joint in a consecutive 6-month period).

[0058] The term "bleeding episode" as used herein adopts a standardized definition of a bleeding episode. A bleeding episode started from the first sign of bleeding, and ended 72 hours after the last treatment for the bleeding, within which any symptoms of bleeding at the same location, or injections less than or equal to 72 hours apart, were considered the same bleeding episode. Any injection to treat the bleeding episode, taken more than 72 hours after the preceding one, was considered the first injection to treat a new bleeding episode at the same location. Any bleeding at a different location was considered a separate bleeding episode regardless of time from last injection. This definition has been proposed by the Subcommittee on Standards and Criteria, FVIII/FIX subcommittee of the International Society of Thrombosis and Hemostasis and has been used by the PedNet multicenter study in hemophilia.

[0059] The term "annualized bleeding rates (ABRs) as used herein refers to the number of bleeding episodes that are annualized for each patient using the following formula:

$$\text{ABR} = \frac{\text{Number of bleeding episodes during efficacy period}}{\text{Total number of days during the efficacy period}} \times 365.25$$

[0060] "Variant," as used herein, refers to a polynucleotide or polypeptide differing from the original polynucleotide or polypeptide, but retaining essential properties thereof, *e.g.*, Factor VIII coagulant activity or Fc (FcRn binding) activity. Generally, variants are overall closely similar, and, in many regions, identical to the original polynucleotide or polypeptide. Variants include polypeptide and polynucleotide fragments, deletions, insertions, and modified versions of original polypeptides.

II. Pharmaceutical Composition

[0061] The present invention is directed to a FVIII polypeptide formulated as a pharmaceutical composition. The pharmaceutical composition can be formulated for administration to humans. The FVIII polypeptide can be a short-acting FVIII polypeptide or a long-acting FVIII polypeptide. The pharmaceutical compositions comprise

pharmaceutically acceptable carriers, including, *e.g.*, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat. Various methods of formulating the invention are well known in the art.

[0062] In certain formulations provided herein, a FVIII polypeptide is formulated as a sterile, preservative-free, non-pyrogenic, lyophilized, white to off-white powder to cake, for intravenous (IV) administration. The formulation can be provided in a single-use vial.

[0063] In certain embodiments, a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}) formulation is provided in a single-use vial manufactured to contain, following reconstitution with an appropriate amount of diluent, about 83 IU/ml, 167 IU/ml, 250 IU/ml, 333 IU/ml, 500 IU/ml, 667 IU/ml, 1000 IU/ml, 1333 IU/ml, 1667 IU/ml, or 2000 IU/ml of the long-acting FVIII polypeptide. In certain embodiments in which diluent is added to a final volume of about 3 ml, a single-use vial can nominally contain about 250, about 500, about 750, about 1000, about 1500, about 2000, about 3000, about 4000, about 5000, or about 6000 International Units (IU) of the FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}).

[0064] In certain embodiments the formulation includes, in addition to the active FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}): sucrose (which can act as a stabilizer), sodium chloride (which can act as a bulking agent), L-histidine (which can act as a buffer), calcium chloride, and polysorbate 20 or polysorbate 80 (which can act as a stabilizer). The formulation is provided with a diluent comprising a sterile sodium chloride solution. In certain embodiments, the diluent is provided in a pre-filled syringe.

[0065] Accordingly, provided herein is a pharmaceutical composition comprising a specified amount of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}) (in IU), along with the excipients sucrose, NaCl, L-histidine, calcium chloride, and polysorbate 20 or polysorbate 80. The compositions provided herein comprise various concentrations of the various excipients, and the concentrations

can be expressed in various ways. For example, the concentration of a given excipient can be expressed as a molar concentration (*e.g.*, M or mM), as a weight/volume percent, (*e.g.*, grams per 100 ml diluent), or as milligrams per milliliter (mg/ml). Formulations provided herein can contain specified amounts of the various excipients at a level of precision ranging from approximate, *e.g.*, concentrations expressed only to one significant figure (*e.g.*, about 0.1% (w/v)), or with more precision, *e.g.*, out to 2, 3, 4, 5, or 6 significant figures (*e.g.*, about 3.88 mg/ml, with precision out to three significant figures). The necessary level of precision can vary depending on, *e.g.*, the requirements of a given regulatory agency, or the manufacturing process. In certain embodiments the pharmaceutical composition comprises a reconstituted formulation, which can be provided as a lyophilisate, optionally accompanied by a diluent.

[0066] In certain embodiments, the pharmaceutical composition comprises about 50 IU/ml to about 2500 IU/ml rFVIII^{IFc}, *e.g.*, 83 IU/ml, 167 IU/ml, 250 IU/ml, 333 IU/ml, 500 IU/ml, 667 IU/ml, 1000 IU/ml, 1333 IU/ml, 1667 IU/ml, or 2000 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII^{IFc}). In certain embodiments, the pharmaceutical composition comprises 83 IU/ml, 167 IU/ml, 250 IU/ml, 333 IU/ml, 500 IU/ml, 667 IU/ml, 1000 IU/ml, 1333 IU/ml, 1667 IU/ml, or 2000 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII^{IFc}) in a formulation comprising about 13.3 mg/ml or about 1.33% (w/v) sucrose, about 12.0 mg/ml or about 205 mM NaCl, about 1.03 mg/ml or about 6.64 mM L-histidine, about 0.80 mg/ml or about 5.4 mM calcium chloride dihydrate, and about 0.13 mg/ml or about 0.013% (w/v) polysorbate 20 or polysorbate 80.

[0067] In certain embodiments, the pre-lyophilization pharmaceutical composition comprises about 100 IU/ml to about 4000 IU/ml rFVIII^{IFc}, *e.g.*, 150 IU/ml, 287.5 IU/ml, 431.25 IU/ml, 575 IU/ml, 862.5 IU/ml, 1150 IU/ml, 1725 IU/ml, 2300 IU/ml, 2875 IU/ml, or 3450 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII^{IFc}). In certain embodiments, the pharmaceutical composition comprises 150 IU/ml, 287.5 IU/ml, 431.25 IU/ml, 575 IU/ml, 862.5 IU/ml, 1150 IU/ml, 1725 IU/ml, 2300 IU/ml, 2875 IU/ml, or 3450 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII^{IFc}) in a formulation comprising about 20.0 mg/ml or about 2.0% (w/v) sucrose, about 18.0 mg/ml or about 308 mM NaCl, about 1.55 mg/ml or about 9.8 mM L-histidine, about 1.18 mg/ml or about 8.0 mM

calcium chloride dihydrate, and about 0.2 mg/ml or about 0.02% (w/v) polysorbate 20 or polysorbate 80.

[0068] In certain embodiments, the pharmaceutical composition comprises a pharmaceutically acceptable amount of sucrose. In certain embodiments, the pharmaceutical composition comprises about 1% (w/v) to about 2.5% (w/v) sucrose, preferably about 1.3% (w/v) to about 2% sucrose (w/v) *e.g.*, about 1.33% (w/v) sucrose or about 2.0% (w/v) sucrose. In certain related embodiments the pharmaceutical composition comprises about 10 mg/ml to about 25 mg/ml sucrose, preferably about 13 mg/ml to about 20 mg/ml sucrose, *e.g.*, about 13.3 mg/ml sucrose or about 20.0 mg/ml sucrose. In still other embodiments, stabilizing agents such as trehalose, raffinose and/or arginine can be used at these concentrations in lieu of or in combination with sucrose.

[0069] In certain embodiments, the pharmaceutical composition comprises about 150 mM to about 250 mM NaCl, about 175 mM to about 225 mM NaCl, and about 200 mM to about 210 mM NaCl, *e.g.*, about 205 mM NaCl. In certain related embodiments, the pharmaceutical composition comprises about 8.8 mg/ml to about 14.6 mg/ml NaCl, about 10 mg/ml NaCl to about 13 mg/ml, *e.g.*, about 12.0 mg/ml NaCl. In certain embodiments, NaCl is provided at the desired concentration in a diluent solution in which a lyophilisate comprising a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII^{IFc}) is reconstituted.

[0070] In another embodiment, the compositions of the invention do not include bulking agents such as mannitol, glycine, alanine and/or hydroxyethyl starch. In other embodiments, NaCl is the sole bulking agent.

[0071] In certain embodiments, the pre-lyophilization pharmaceutical composition comprises about 250 mM to about 350 mM NaCl, about 275 mM to about 325 mM NaCl, *e.g.*, about 308 mM NaCl. In certain related embodiments, the pharmaceutical composition comprises about 14.6 mg/ml to about 20.5 mg/ml NaCl, about 16 mg/ml to about 19 mg/ml, *e.g.*, about 18.0 mg/ml NaCl.

[0072] In certain embodiments, the pharmaceutical composition comprises a pharmaceutically acceptable amount of L-histidine. In certain embodiments, the pharmaceutical composition comprises about 5 mM to about 15 mM L-histidine, *e.g.*, about 6.64 mM L-histidine or about 9.8 mM L-histidine. In certain related embodiments the pharmaceutical composition comprises about 0.75 mg/ml to about 2.25 mg/ml L-histidine, *e.g.*, about 1.03 mg/ml L-histidine or about 1.55 mg/ml L-histidine. In certain

embodiments, L-histidine is provided as part of a lyophilisate, which, upon reconstitution with an appropriate amount of diluent provides L-histidine at the desired concentration.

[0073] In certain embodiments, the pharmaceutical composition comprises a pharmaceutically acceptable amount of calcium chloride. In certain embodiments, the pharmaceutical composition comprises about 5 mM to about 10 mM calcium chloride, *e.g.*, about 5.4 mM calcium chloride or about 8 mM calcium chloride. In certain related embodiments the pharmaceutical composition comprises about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate, *e.g.*, about 0.8 mg/ml calcium chloride dihydrate or about 1.18 mg/ml calcium chloride dihydrate. In certain embodiments, calcium chloride is provided as part of a lyophilisate, which, upon reconstitution with an appropriate amount of diluent provides calcium chloride at the desired concentration.

[0074] In certain embodiments, the pharmaceutical composition comprises a pharmaceutically acceptable amount of polysorbate 20 or polysorbate 80. In certain related embodiments the pharmaceutical composition comprises about 0.008% (w/v) to about 0.025% (w/v) polysorbate 20 or polysorbate 80, *e.g.*, about 0.013% (w/v) polysorbate 20 or polysorbate 80 or about 0.02 % (w/v) polysorbate 20 or polysorbate 80. In certain related embodiments the pharmaceutical composition comprises about 0.08 mg/ml to about 0.25 mg/ml polysorbate 20 or polysorbate 80, *e.g.*, about 0.13% mg/ml polysorbate 20 or polysorbate 80 or about 0.20 mg/ml polysorbate 20 or polysorbate 80. In certain embodiments, polysorbate 20 or polysorbate 80 is provided as part of a lyophilisate, which, upon reconstitution with an appropriate amount of diluent provides polysorbate 20 or polysorbate 80 at the desired concentration.

[0075] In certain embodiments, the pharmaceutical composition comprises: about 50 IU/ml to about 2500 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{Fc}); about 1% (w/v) to about 2.5% (w/v) of sucrose; about 150 mM to about 250 mM NaCl; about 5 mM to about 15 mM L-histidine; about 5 mM to about 10 mM calcium chloride; and about 0.008% (w/v) to about 0.025% of polysorbate 20 or polysorbate 80. In certain embodiments the pharmaceutical composition is provided as a lyophilisate and a diluent. In certain embodiments the amount of lyophilizate provides about 3 ml of a pharmaceutical composition with the desired ingredients at the desired concentrations.

[0076] In certain embodiments, the pharmaceutical composition comprises: about 50 IU/ml to about 2500 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting

FVIII polypeptide, *e.g.*, rFVIII_{IFc}); about 10 mg/ml to about 25 mg/ml of sucrose; about 8.8 mg/ml to about 14.6 mg/ml NaCl; about 0.75 mg/ml to about 2.25 mg/ml L-histidine; about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate; and about 0.08 mg/ml to about 0.25 mg/ml of polysorbate 20 or polysorbate 80. In certain embodiments the pharmaceutical composition is provided as a lyophilisate and a diluent. In certain embodiments the amount of lyophilisate provides about 3 ml of a pharmaceutical composition with the desired ingredients at the desired concentrations.

[0077] Exemplary compositions are provided in Table 1 and in Table 2 in the Examples.

[0078] For example, the disclosure provides a pharmaceutical composition comprising: about 83 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride dihydrate, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 167 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride dihydrate, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 250 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride dihydrate, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 333 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride dihydrate, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 500 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride dihydrate, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 667 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium

chloride dihydrate, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 1000 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. The disclosure also provides a pharmaceutical composition comprising: about 1333 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. In some embodiments, the disclosure provides a pharmaceutical composition comprising: about 1667 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. In other embodiments, the disclosure provides a pharmaceutical composition comprising: about 2000 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

[0079] The disclosure further provides a pharmaceutical composition comprising: about 83 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 1673 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 250 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 333

IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 500 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 667 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 1000 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. The disclosure also provides a pharmaceutical composition comprising: about 1333 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. In some embodiments, the disclosure provides a pharmaceutical composition comprising: about 1667 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. In other embodiments, the disclosure provides a pharmaceutical composition comprising: about 2000 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80.

[0080] This disclosure also provides the components of a pharmaceutical kit. Such a kit includes one or more containers and optional attachments. A kit as provided herein facilitates administration of an effective amount of the FVIII polypeptide (*e.g.*, a long-

acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}) to a subject in need thereof. In certain embodiments, the kit facilitates administration of the FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}) via intravenous infusion. In certain embodiments, the kit facilitates self-administration of the FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}) via intravenous infusion.

[0081] In certain embodiments, the disclosure provides a pharmaceutical kit comprising: a first container comprising a lyophilized powder or cake, where the powder or cake comprises: (i) a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}), (ii) sucrose (and/or trehalose, raffinose or arginine); (iii) NaCl; (iv) L-histidine; (v) calcium chloride dihydrate; and (vi) polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections to be combined with the lyophilized powder of the first container. In certain embodiments, sufficient diluent is provided to produce about 3 ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}) formulation with desired properties as disclosed herein. In certain embodiments, the second container is a pre-filled syringe associated with a plunger, to allow addition of the diluent to the first container, reconstitution of the contents of the first container, and transfer back into the syringe. In certain embodiments, the kit further provides an adaptor for attaching the syringe to the first container. In certain embodiments the kit further provides a needle and infusion tubing, to be attached to the syringe containing the reconstituted FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}) formulation to allow IV infusion of the formulation.

[0082] In certain embodiments a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}) is provided in a total amount from about 200 IU to about 6000 IU, *e.g.*, about 250 IU, about 500 IU, about 750 IU, about 1000 IU, about 1500 IU, about 2000 IU, about 3000 IU, about 4000 IU, about 5000 IU, or about 6000 IU.

[0083] In one embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises (i) about 250 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII_{IFc}), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder

of the first container, a solution comprising: (i) about 83 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); (ii) about 1.33% (w/v) of sucrose; (iii) about 205 mM of NaCl; (iv) about 6.64 mM L-histidine; (v) about 5.4 mM of calcium chloride; and (vi) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

[0084] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 500 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 167 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); (ii) about 1.33% (w/v) of sucrose; (iii) about 205 mM of NaCl; (iv) about 6.64 mM L-histidine; (v) about 5.4 mM of calcium chloride; and (vi) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

[0085] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 750 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 250 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); (ii) about 1.33% (w/v) of sucrose; (iii) about 205 mM of NaCl; (iv) about 6.64 mM L-histidine; (v) about 5.4 mM of calcium chloride; and (vi) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

[0086] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 1000 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-

histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 333 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII^{IFc}); (ii) about 1.33% (w/v) of sucrose; (iii) about 205 mM of NaCl; (iv) about 6.64 mM L-histidine; (v) about 5.4 mM of calcium chloride; and (vi) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

[0087] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 1500 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII^{IFc}), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 500 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII^{IFc}); (ii) about 1.33% (w/v) of sucrose; (iii) about 205 mM of NaCl; (iv) about 6.64 mM L-histidine; (v) about 5.4 mM of calcium chloride; and (vi) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

[0088] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 2000 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII^{IFc}), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 667 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII^{IFc}); (ii) about 1.33% (w/v) of sucrose; (iii) about 205 mM of NaCl; (iv) about 6.64 mM L-histidine; (v) about 5.4 mM of calcium chloride; and (vi) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

[0089] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 3000 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 1000 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); (ii) about 1.33% (w/v) of sucrose; (iii) about 205 mM of NaCl; (iv) about 6.64 mM L-histidine; (v) about 5.4 mM of calcium chloride; and (vi) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

[0090] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 4000 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 1333 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); (ii) about 1.33% (w/v) of sucrose; (iii) about 205 mM of NaCl; (iv) about 6.64 mM L-histidine; (v) about 5.4 mM of calcium chloride; and (vi) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

[0091] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 5000 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 1667 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); (ii) about 1.33%

(w/v) of sucrose; (iii) about 205 mM of NaCl; (iv) about 6.64 mM L-histidine; (v) about 5.4 mM of calcium chloride; and (vi) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

[0092] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 6000 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 2000 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); (ii) about 1.33% (w/v) of sucrose; (iii) about 205 mM of NaCl; (iv) about 6.64 mM L-histidine; (v) about 5.4 mM of calcium chloride; and (vi) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

[0093] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 250 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 83 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); (ii) about 13.3 mg/ml of sucrose; (iii) about 12.0 mg/ml of NaCl; (iv) about 1.03 mg/ml L-histidine; (v) about 0.80 mg/ml of calcium chloride dihydrate; and (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

[0094] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 500 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for

injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 167 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); (ii) about 13.3 mg/ml of sucrose; (iii) about 12.0 mg/ml of NaCl; (iv) about 1.03 mg/ml L-histidine; (v) about 0.80 mg/ml of calcium chloride dihydrate; and (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

[0095] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 750 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 250 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); (ii) about 13.3 mg/ml of sucrose; (iii) about 12.0 mg/ml of NaCl; (iv) about 1.03 mg/ml L-histidine; (v) about 0.80 mg/ml of calcium chloride dihydrate; and (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

[0096] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 1000 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 333 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); (ii) about 13.3 mg/ml of sucrose; (iii) about 12.0 mg/ml of NaCl; (iv) about 1.03 mg/ml L-histidine; (v) about 0.80 mg/ml of calcium chloride dihydrate; and (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

[0097] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 1500 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*,

rFVIII(Fc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 500 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII(Fc)); (ii) about 13.3 mg/ml of sucrose; (iii) about 12.0 mg/ml of NaCl; (iv) about 1.03 mg/ml L-histidine; (v) about 0.80 mg/ml of calcium chloride dihydrate; and (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

[0098] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 2000 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII(Fc)), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 667 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII(Fc)); (ii) about 13.3 mg/ml of sucrose; (iii) about 12.0 mg/ml of NaCl; (iv) about 1.03 mg/ml L-histidine; (v) about 0.80 mg/ml of calcium chloride dihydrate; and (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

[0099] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 3000 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII(Fc)), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 1000 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII(Fc)); (ii) about 13.3 mg/ml of sucrose; (iii) about 12.0 mg/ml of NaCl; (iv) about 1.03 mg/ml L-histidine; (v) about 0.80 mg/ml of calcium chloride dihydrate; and (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

[0100] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 4000 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 1333 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); (ii) about 13.3 mg/ml of sucrose; (iii) about 12.0 mg/ml of NaCl; (iv) about 1.03 mg/ml L-histidine; (v) about 0.80 mg/ml of calcium chloride dihydrate; and (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

[0101] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 5000 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 1667 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); (ii) about 13.3 mg/ml of sucrose; (iii) about 12.0 mg/ml of NaCl; (iv) about 1.03 mg/ml L-histidine; (v) about 0.80 mg/ml of calcium chloride dihydrate; and (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

[0102] In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 6000 IU of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 2000 IU/ml of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIIIIFc); (ii) about 13.3

mg/ml of sucrose; (iii) about 12.0 mg/ml of NaCl; (iv) about 1.03 mg/ml L-histidine; (v) about 0.80 mg/ml of calcium chloride dihydrate; and (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

[0103] In certain embodiments the first container of a pharmaceutical kit provided herein is a glass vial comprising a rubber stopper. In certain embodiments, the second container a pharmaceutical kit provided herein is a syringe body, associated with a plunger. In certain embodiments, the syringe is a pre-filled syringe containing the diluent. In certain embodiments, a pharmaceutical kit provided herein further comprises an adaptor to connect the glass vial to the syringe body. In certain embodiments a pharmaceutical kit provided herein further comprises infusion tubing associated with a needle to be connected to the syringe, suitable for intravenous infusion.

[0104] In certain embodiments, a desired dose of a FVIII polypeptide (*e.g.*, a long-acting or short-acting FVIII polypeptide, *e.g.*, rFVIII-Fc) can be achieved through the use of one pharmaceutical kit as provided herein. In certain embodiments, more than one pharmaceutical kit can be used to achieve a desired dose. Provided herein is a method of combining, or pooling the formulations contained in two or more pharmaceutical kits as provided herein in order to achieve a desired dose.

[0105] In some embodiments, the FVIII polypeptide for the pharmaceutical composition is a short-acting FVIII polypeptide. In other embodiments, the FVIII polypeptide for the pharmaceutical composition is a long-acting FVIII polypeptide.

[0106] In some embodiments, a long-acting FVIII polypeptide in a pharmaceutical composition comprises a Factor VIII portion and a non-Factor VIII portion, *e.g.*, a heterologous moiety. In one embodiment, the heterologous moiety is capable of extending *in vivo* or *in vitro* half-life of the FVIII polypeptide. Exemplary non-Factor VIII portions include, but are not limited to, Fc, albumin, a PAS sequence, transferrin, CTP (28 amino acid C-terminal peptide (CTP) of human chorionic gonadotropin (hCG) with its 4 O-glycans), polyethylene glycol (PEG), hydroxyethyl starch (HES), albumin binding polypeptide, albumin-binding small molecules, or any combination thereof. Exemplary long-acting polypeptides of the invention include, *e.g.*, Factor VIII-Fc polypeptides, Factor VIII-albumin polypeptides, Factor VIII-PAS polypeptides, Factor VIII-transferrin polypeptides, Factor VIII-CTP polypeptides, Factor VIII-PEG polypeptides, Factor VIII-HES polypeptides, Factor VIII-albumin binding polypeptide polypeptides, or Factor VIII-albumin-binding small molecule polypeptides.

- [0107]** The Factor VIII (the Factor VIII portion of a long-acting FVIII polypeptide or FVIII of a short-acting FVIII polypeptide) can be at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to a Factor VIII amino acid sequence shown in Table 11 without a signal sequence (amino acids 20 to 1457 of SEQ ID NO:2; and amino acids 20 to 2351 of SEQ ID NO:6), wherein the Factor VIII portion has Factor VIII activity. The Factor VIII (the Factor VIII portion of a chimeric polypeptide) can be identical to a Factor VIII amino acid sequence shown in Table 11 without a signal sequence (amino acids 20 to 1457 of SEQ ID NO:2; and amino acids 20 to 2351 of SEQ ID NO:6).
- [0108]** The Factor VIII (the Factor VIII portion of a long-acting FVIII polypeptide or FVIII of a short-acting FVIII polypeptide) can be at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to a Factor VIII amino acid sequence shown in Table 11 with a signal sequence (amino acids 1 to 1457 of SEQ ID NO:2 and amino acids 1 to 2351 of SEQ ID NO:6), wherein the Factor VIII portion has Factor VIII activity. The Factor VIII (the Factor VIII portion of a long-acting FVIII polypeptide or a short-acting FVIII polypeptide) can be identical to a Factor VIII amino acid sequence shown in Table 11 with a signal sequence (amino acids 1 to 1457 of SEQ ID NO:2 and amino acids 1 to 2351 of SEQ ID NO:6).
- [0109]** The long-acting polypeptide can comprise a sequence at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the Factor VIII and Fc amino acid sequence shown in Table 11A(i) without a signal sequence (amino acids 20 to 1684 of SEQ ID NO:2) or at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the Factor VIII and Fc amino acid sequence shown in Table 11A(i) with a signal sequence (amino acids 1 to 1684 of SEQ ID NO:2), wherein the sequence has Factor VIII activity. The Factor VIII activity can be measured by activated Partial Thromboplastin Time (aPTT) assay, chromogenic assay, or other known methods. The long-acting FVIII polypeptide can comprise the sequence identical to the Factor VIII and Fc amino acid sequence shown in Table 11A(i) without a signal sequence (amino acids 20 to 1684 of SEQ ID NO:2) or identical to the Factor VIII and Fc amino acid sequence shown in Table 11A(i) with a signal sequence (amino acids 1 to 1684 of SEQ ID NO:2).
- [0110]** As discussed above, exemplary long-acting polypeptides also include Factor VIII fused to one or more albumin polypeptides, albumin binding polypeptides, or albumin-binding small molecules. In one embodiment, the albumin is human albumin. The

albumin or albumin binding protein can be fused to either the N-terminal end of FVIII or to the C-terminal end of FVIII or inserted between two amino acids in FVIII. Examples of albumin, *e.g.*, fragments thereof, that can be used in the present invention are known. *e.g.*, U.S. Patent No. 7,592,010; U.S. Patent No. 6,686,179; and Schulte, *Thrombosis Res.* 124 Suppl. 2:S6-S8 (2009), each of which is incorporated herein by reference in its entirety.

[0111] The albumin binding polypeptides can comprise, without limitation, bacterial albumin-binding domains, albumin-binding peptides, or albumin-binding antibody fragments that can bind to albumin. Domain 3 from streptococcal protein G, as disclosed by Kraulis *et al.*, *FEBS Lett.* 378:190-194 (1996) and Linhult *et al.*, *Protein Sci.* 11:206-213 (2002) is an example of a bacterial albumin-binding domain. Examples of albumin-binding peptides include a series of peptides having the core sequence DICLPRWGCLW (SEQ ID NO: 7). See, *e.g.*, Dennis *et al.*, *J. Biol. Chem.* 2002, 277: 35035-35043 (2002). Examples of albumin-binding antibody fragments are disclosed in Muller and Kontermann, *Curr. Opin. Mol. Ther.* 9:319-326 (2007); Rooverset *et al.*, *Cancer Immunol. Immunother.* 56:303-317 (2007), and Holt *et al.*, *Prot. Eng. Design Sci.*, 21:283-288 (2008), which are incorporated herein by reference in their entireties.

[0112] In certain aspects, a long-acting FVIII polypeptide of the invention comprises at least one attachment site for a non-polypeptide small molecule, variant, or derivative that can bind to albumin thereof. An example of such albumin binding moieties is 2-(3-maleimidopropanamido)-6-(4-(4-iodophenyl)butanamido)hexanoate ("Albu" tag) as disclosed by Trusselet *et al.*, *Bioconjugate Chem.* 20:2286-2292 (2009).

[0113] As discussed above, exemplary long-acting polypeptides also include Factor VIII fused to at least one C-terminal peptide (CTP) of the β subunit of human chorionic gonadotropin or fragment, variant, or derivative thereof. The CTP can be fused to Factor VIII either the N-terminal end of FVIII or to the C-terminal end of FVIII or inserted between two amino acids in FVIII. One or more CTP peptides fused to or inserted into a recombinant protein is known to increase the *in vivo* half-life of that protein. See, *e.g.*, U.S. Patent No. 5,712,122, incorporated by reference herein in its entirety. Exemplary CTP peptides include DPRFQDSSSSKAPPSLSPSRLPGPSDTPIL (SEQ ID NO: 8) or SSSSKAPPSLSPSRLPGPSDTPILPQ. (SEQ ID NO: 9). See, *e.g.*, U.S. Patent Application Publication No. US 2009/0087411 A1, incorporated by reference.

[0114] As discussed above, exemplary long-acting FVIII polypeptides also include Factor VII. fused to at least one PAS sequence or fragment, variant, or derivative thereof. The PAS sequence can be fused to either the N-terminal end of FVIII or to the C-terminal end of FVIII or inserted between two amino acids in FVIII. A PAS peptide or PAS sequence, as used herein, means an amino acid sequence comprising mainly alanine and serine residues or comprising mainly alanine, serine, and proline residues, the amino acid sequence forming random coil conformation under physiological conditions. Accordingly, the PAS sequence is a building block, an amino acid polymer, or a sequence cassette comprising, consisting essentially of, or consisting of alanine, serine, and proline which can be used as a part of the heterologous moiety in the chimeric protein. An amino acid polymer also can form random coil conformation when residues other than alanine, serine, and proline are added as a minor constituent in the PAS sequence. By “minor constituent” is meant that that amino acids other than alanine, serine, and proline can be added in the PAS sequence to a certain degree, *e.g.*, up to about 12%, *i.e.*, about 12 of 100 amino acids of the PAS sequence, up to about 10%, up to about 9%, up to about 8%, about 6%, about 5%, about 4%, about 3%, *i.e.* about 2%, or about 1%, of the amino acids. The amino acids different from alanine, serine and proline can be selected from the group consisting of Arg, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Lys, Met, Phe, Thr, Trp, Tyr, and Val. Under physiological conditions, a PAS peptide forms a random coil conformation and thereby can mediate an increased *in vivo* and/or *in vitro* stability to a recombinant protein of the invention, and has procoagulant activity.

[0115] Non-limiting examples of the PAS peptides include ASPAAPAPASPAAPAPSAPA (SEQ ID NO: 10), AAPASPAPAAPSAPAPAAPS (SEQ ID NO: 11), APSSPSPSAPSSPSPASPSS (SEQ ID NO: 12), APSSPSPSAPSSPSPASPS (SEQ ID NO: 13), SSPSAPSPSSPASPSPSSPA (SEQ ID NO: 14), AASPAAPSAPPAAASPAAPSAPPA (SEQ ID NO: 15), ASAAAPAAASAAASAPSAAA (SEQ ID NO: 16) or any variants, derivatives, fragments, or combinations thereof. Additional examples of PAS sequences are known from, *e.g.*, US Pat. Publ. No. 2010/0292130 A1 and PCT Appl. Publ. No. WO 2008/155134 A1. European issued patent EP2173890.

[0116] As discussed above, exemplary long-acting FVIII polypeptides also include Factor VIII fused to at least one transferrin peptide or fragment, variant, or derivative thereof. At least one transferrin peptide can be fused to either the N-terminal end of FVIII or to

the C-terminal end of FVIII or inserted between two amino acids in FVIII. Any transferrin can be fused to or inserted into a recombinant FVIII protein of the invention. As an example, wild-type human Tf (Tf) is a 679 amino acid protein, of approximately 75 KDa (not accounting for glycosylation), with two main domains, N (about 330 amino acids) and C (about 340 amino acids), which appear to originate from a gene duplication. See GenBank accession numbers NM001063, XM002793, M12530, XM039845, XM039847 and S95936 (www.ncbi.nlm.nih.gov), all of which are herein incorporated by reference in their entirety.

[0117] Transferrin transports iron through transferrin receptor (TfR)-mediated endocytosis. After the iron is released into an endosomal compartment and Tf-TfR complex is recycled to cell surface, the Tf is released back extracellular space for next cycle of iron transporting. Tf possesses a long half-life that is in excess of 14-17 days (Li *et al.*, *Trends Pharmacol. Sci.* 23:206-209 (2002)). Transferrin fusion proteins have been studied for half-life extension, targeted deliver for cancer therapies, oral delivery and sustained activation of proinsulin (Brandsma *et al.*, *Biotechnol. Adv.*, 29: 230-238 (2011); Bai *et al.*, *Proc. Natl. Acad. Sci. USA* 102:7292-7296 (2005); Kim *et al.*, *J. Pharmacol. Exp. Ther.*, 334:682-692 (2010); Wang *et al.*, *J. Controlled Release* 155:386-392 (2011)).

[0118] As discussed above, exemplary long-acting FVIII polypeptides also include Factor VIII fused to at least one polyethylene glycol (PEG) moieties.

[0119] PEGylated FVIII can refer to a conjugate formed between FVIII and at least one polyethylene glycol (PEG) molecule. PEG is commercially available in a large variety of molecular weights and average molecular weight ranges. Typical examples of PEG average molecular weight ranges include, but are not limited to, about 200, about 300, about 400, about 600, about 1000, about 1300-1600, about 1450, about 2000, about 3000, about 3000-3750, about 3350, about 3000-7000, about 3500-4500, about 5000-7000, about 7000-9000, about 8000, about 10000, about 8500-11500, about 16000-24000, about 35000, about 40000, about 60000, and about 80000 daltons. These average molecular weights are provided merely as examples and are not meant to be limiting in any way.

[0120] A long-acting FVIII polypeptide useful for the invention can be PEGylated to include mono- or poly-(*e.g.*, 2-4) PEG moieties. PEGylation can be carried out by any of the PEGylation reactions known in the art. Methods for preparing a PEGylated protein product will generally include (i) reacting a polypeptide with polyethylene glycol (such as a reactive ester or aldehyde derivative of PEG) under conditions whereby the peptide of

the invention becomes attached to one or more PEG groups; and (ii) obtaining the reaction product(s). In general, the optimal reaction conditions for the reactions will be determined case by case based on known parameters and the desired result.

[0121] There are a number of PEG attachment methods available to those skilled in the art, for example Malik F *et al.*, *Exp. Hematol.* 20:1028-35 (1992); Francis, *Focus on Growth Factors* 3(2):4-10 (1992); European Pat. Pub. Nos. EP0401384, EP0154316, and EP0401384; and International Pat. Appl. Pub. Nos. WO92/16221 and WO95/34326. As a non-limiting example, FVIII variants can contain cysteine substitutions in one or more insertion sites in FVIII, and the cysteines can be further conjugated to PEG polymer. See Mei *et al.*, *Blood* 116:270-279 (2010) and U.S. Patent No. 7,632,921, which are incorporated herein by reference in their entireties.

[0122] As discussed above, exemplary long-acting FVIII polypeptides also include Factor VIII fused to at least one hydroxyethyl starch (HES) polymer. HES is a derivative of naturally occurring amylopectin and is degraded by alpha-amylase in the body. HES exhibits advantageous biological properties and is used as a blood volume replacement agent and in hemodilution therapy in the clinics. See, *e.g.*, Sommermeyer *et al.*, *Krankenhauspharmazie* 8:271-278 (1987); and Weidler *et al.*, *Arzneim.-Forschung/Drug Res.* 41: 494-498 (1991).

[0123] HES is mainly characterized by the molecular weight distribution and the degree of substitution. HES has a mean molecular weight (weight mean) of from 1 to 300 kD, from 2 to 200kD, from 3 to 100 kD, or from 4 to 70kD. Hydroxyethyl starch can further exhibit a molar degree of substitution of from 0.1 to 3, from 0.1 to 2, from 0.1 to 0.9, or from 0.1 to 0.8, and a ratio between C2:C6 substitution in the range of from 2 to 20 with respect to the hydroxyethyl groups. HES with a mean molecular weight of about 130 kD is VOLUVEN[®] from Fresenius. VOLUVEN[®] is an artificial colloid, employed, *e.g.*, for volume replacement used in the therapeutic indication for therapy and prophylaxis of hypovolaemia. There are a number of HES attachment methods available to those skilled in the art, *e.g.*, the same PEG attachment methods described above.

[0124] In some embodiments, a long-acting FVIII polypeptide comprising a FVIII portion has an increased half-life ($t_{1/2}$) over a polypeptide consisting of the same FVIII portion without the non-FVIII portion. A long-acting FVIII polypeptide with an increased $t_{1/2}$ can be referred to herein as a long-lasting FVIII. Long-acting chimeric Factor VIII polypeptides include, *e.g.*, Factor VIII fused to Fc (including, *e.g.*, chimeric

Factor VIII polypeptides in the form of a hybrid such as a FVIII_h monomer dimer hybrid; see Example, Table 11A; and US Patent Nos. 7,404,956 and 7,348,004), and Factor VIII fused to albumin.

[0125] The Factor VIII polypeptide as used herein is functional factor VIII polypeptide in its normal role in coagulation, unless otherwise specified. Thus, the term Factor VIII includes variant polypeptides that are functional. Factor VIII proteins can be the human, porcine, canine, and murine factor VIII proteins. The full length polypeptide and polynucleotide sequences are known, as are many functional fragments, mutants and modified versions. Examples of human factor VIII sequences are shown as subsequences in SEQ ID NOs: 2 or 6 (Table 11). Factor VIII polypeptides include, *e.g.*, full-length factor VIII, full-length factor VIII minus Met at the N-terminus, mature factor VIII (minus the signal sequence), mature factor VIII with an additional Met at the N-terminus, and/or factor VIII with a full or partial deletion of the B domain. Factor VIII variants include B domain deletions, whether partial or full deletions.

[0126] A great many functional factor VIII variants are known, as is discussed above and below. In addition, hundreds of nonfunctional mutations in factor VIII have been identified in hemophilia patients, and it has been determined that the effect of these mutations on factor VIII function is due more to where they lie within the 3-dimensional structure of factor VIII than on the nature of the substitution (Cutler *et al.*, *Hum. Mutat.* 19:274-8 (2002)), incorporated herein by reference in its entirety. In addition, comparisons between factor VIII from humans and other species have identified conserved residues that are likely to be required for function (Cameron *et al.*, *Thromb. Haemost.* 79:317-22 (1998); US 6,251,632), incorporated herein by reference in its entirety.

[0127] The human factor VIII gene was isolated and expressed in mammalian cells (Toole, J. J., *et al.*, *Nature* 312:342-347 (1984); Gitschier, J., *et al.*, *Nature* 312:326-330 (1984); Wood, W. I., *et al.*, *Nature* 312:330-337 (1984); Vehar, G. A., *et al.*, *Nature* 312:337-342 (1984); WO 87/04187; WO 88/08035; WO 88/03558; U.S. Pat. No. 4,757,006), each of which is incorporated herein by reference in its entirety, and the amino acid sequence was deduced from cDNA. Capon *et al.*, U.S. Pat. No. 4,965,199, incorporated herein by reference in its entirety, discloses a recombinant DNA method for producing factor VIII in mammalian host cells and purification of human factor VIII. Human factor VIII expression in CHO (Chinese hamster ovary) cells and BHK (baby

hamster kidney cells) has been reported. Human factor VIII has been modified to delete part or all of the B domain (U.S. Pat. Nos. 4,994,371 and 4,868,112, each of which is incorporated herein by reference in its entirety), and replacement of the human factor VIII B domain with the human factor V B domain has been performed (U.S. Pat. No. 5,004,803, incorporated herein by reference in its entirety). The cDNA sequence encoding human factor VIII and predicted amino acid sequence are shown in SEQ ID NOs:1 and 2, respectively, of US Application Publ. No. 2005/0100990, incorporated herein by reference in its entirety.

[0128] U.S. Pat. No. 5,859,204, Lollar, J. S., incorporated herein by reference in its entirety, reports functional mutants of factor VIII having reduced antigenicity and reduced immunoreactivity. U.S. Pat. No. 6,376,463, Lollar, J. S., incorporated herein by reference in its entirety, also reports mutants of factor VIII having reduced immunoreactivity. US Application Publ. No. 2005/0100990, Saenko et al., incorporated herein by reference in its entirety, reports functional mutations in the A2 domain of factor VIII.

[0129] A number of functional factor VIII molecules, including B-domain deletions, are disclosed in the following patents US 6,316,226 and US 6,346,513, both assigned to Baxter; US 7,041,635 assigned to In2Gen; US 5,789,203, US 6,060,447, US 5,595,886, and US 6,228,620 assigned to Chiron; US 5,972,885 and US 6,048,720 assigned to Biovitrum, US 5,543,502 and US 5,610,278 assigned to Novo Nordisk; US 5,171,844 assigned to Immuno Ag; US 5,112,950 assigned to Transgene S.A.; US 4,868,112 assigned to Genetics Institute, each of which is incorporated herein by reference in its entirety.

[0130] The porcine factor VIII sequence is published, (Toole, J. J., *et al.*, *Proc. Natl. Acad. Sci. USA* 83:5939-5942 (1986)), incorporated herein by reference in its entirety, and the complete porcine cDNA sequence obtained from PCR amplification of factor VIII sequences from a pig spleen cDNA library has been reported (Healey, J. F. *et al.*, *Blood* 88:4209-4214 (1996), incorporated herein by reference in its entirety). Hybrid human/porcine factor VIII having substitutions of all domains, all subunits, and specific amino acid sequences were disclosed in U.S. Pat. No. 5,364,771 by Lollar and Runge, and in WO 93/20093, incorporated herein by reference in its entirety. More recently, the nucleotide and corresponding amino acid sequences of the A1 and A2 domains of porcine factor VIII and a chimeric factor VIII with porcine A1 and/or A2 domains substituted for

the corresponding human domains were reported in WO 94/11503, incorporated herein by reference in its entirety. U.S. Pat. No. 5,859,204, Lollar, J. S., also discloses the porcine cDNA and deduced amino acid sequences. US Pat. No. 6,458,563, incorporated herein by reference in its entirety assigned to Emory discloses a B-domain deleted porcine Factor VIII.

[0131] A "B domain" of Factor VIII, as used herein, is the same as the B domain known in the art that is defined by internal amino acid sequence identity and sites of proteolytic cleavage by thrombin, *e.g.*, residues Ser741-Arg1648 of full length mature human factor VIII. The other human factor VIII domains are defined by the following amino acid residues: A1, residues Ala1-Arg372; A2, residues Ser373-Arg740; A3, residues Ser1690-Ile2032; C1, residues Arg2033-Asn2172; C2, residues Ser2173-Tyr2332. The A3-C1-C2 sequence includes residues Ser1690-Tyr2332. The remaining sequence, residues Glu1649-Arg1689, is usually referred to as the factor VIII light chain activation peptide. The locations of the boundaries for all of the domains, including the B domains, for porcine, mouse and canine factor VIII are also known in the art. In one embodiment, the B domain of Factor VIII is deleted ("B domain deleted factor VIII" or "BDD FVIII"). An example of a BDD FVIII is REFACTO[®] (recombinant BDD FVIII), which has the sequence as the Factor VIII portion of the sequence in Table 11A(i) (amino acids 1 to 1457 or 20 to 1457 of SEQ ID NO:2). In another embodiment, the B domain deleted Factor VIII contains an intact intracellular processing site, which corresponds to Arginine at residue 754 of B domain deleted Factor VIII, which corresponds to Arginine residue 773 of SEQ ID NO: 2, or residue 1648 of full-length Factor VIII, which corresponds to Arginine residue 1667 of SEQ ID NO: 6. The sequence residue numbers used herein without referring to any SEQ ID Numbers correspond to the Factor VIII sequence without the signal peptide sequence (19 amino acids) unless otherwise indicated. For example, S743/Q1638 of full-length Factor VIII corresponds to S762/Q1657 of SEQ ID NO: 6 due to the 19 amino acid signal peptide sequence. In other embodiments, the B domain deleted FVIII comprises a substitution or mutation at an amino acid position corresponding to Arginine 1645, a substitution or mutation at an amino acid position corresponding to Arginine 1648, or a substitution or mutation at amino acid positions corresponding to Arginine 1645 and Arginine 1648 in full-length Factor VIII. In some embodiments, the amino acid substituted at the amino acid position corresponding to Arginine 1645 is a different amino acid from the amino acid substituted at the amino acid

position corresponding to Arginine 1648. In certain embodiments, the substitution or mutation is an amino acid other than arginine, *e.g.*, alanine.

[0132] A "B domain deleted factor VIII" can have the full or partial deletions disclosed in U.S. Patent Nos. 6,316,226, 6,346,513, 7,041,635, 5,789,203, 6,060,447, 5,595,886, 6,228,620, 5,972,885, 6,048,720, 5,543,502, 5,610,278, 5,171,844, 5,112,950, 4,868,112, and 6,458,563, each of which is incorporated herein by reference in its entirety. In some embodiments, a B domain deleted factor VIII sequence of the present invention comprises any one of the deletions disclosed at col. 4, line 4 to col. 5, line 28 and examples 1-5 of U.S. Patent No. 6,316,226 (also in US 6,346,513). In some embodiments, a B domain deleted factor VIII of the present invention has a deletion disclosed at col. 2, lines 26-51 and examples 5-8 of U.S. Patent No. 5,789,203 (also US 6,060,447, US 5,595,886, and US 6,228,620). In some embodiments, a B domain deleted factor VIII has a deletion described in col. 1, lines 25 to col. 2, line 40 of US Patent No. 5,972,885; col. 6, lines 1-22 and example 1 of U.S. Patent no. 6,048,720; col. 2, lines 17-46 of U.S. Patent No. 5,543,502; col. 4, line 22 to col. 5, line 36 of U.S. Patent no. 5,171,844; col. 2, lines 55-68, figure 2, and example 1 of U.S. Patent No. 5,112,950; col. 2, line 2 to col. 19, line 21 and Table 2 of U.S. Patent No. 4,868,112; col. 2, line 1 to col. 3, line 19, col. 3, line 40 to col. 4, line 67, col. 7, line 43 to col. 8, line 26, and col. 11, line 5 to col. 13, line 39 of U.S. Patent no. 7,041,635; or col. 4, lines 25-53, of U.S. Patent No. 6,458,563. In some embodiments, a B domain deleted factor VIII has a deletion of most of the B domain, but still contains amino-terminal sequences of the B domain that are essential for *in vivo* proteolytic processing of the primary translation product into two polypeptide chain (*i.e.*, intracellular processing site), as disclosed in WO 91/09122, which is incorporated herein by reference in its entirety. In some embodiments, a B domain deleted factor VIII is constructed with a deletion of amino acids 747-1638, *i.e.*, virtually a complete deletion of the B domain. Hoebe R.C., *et al. J. Biol. Chem.* 265 (13): 7318-7323 (1990), incorporated herein by reference in its entirety. A B domain deleted factor VIII can also contain a deletion of amino acids 771-1666 or amino acids 868-1562 of factor VIII. Meulien P., *et al. Protein Eng.* 2(4): 301-6 (1988), incorporated herein by reference in its entirety. Additional B domain deletions that are part of the invention include, *e.g.*, deletion of amino acids 982 through 1562 or 760 through 1639 (Toole *et al., Proc. Natl. Acad. Sci. U.S.A.* 83:5939-5942 (1986)), 797 through 1562 (Eaton *et al., Biochemistry* 25:8343-8347 (1986)), 741 through 1646 (Kaufman (PCT published

application No. WO 87/04187)), 747-1560 (Sarver *et al.*, *DNA* 6:553-564 (1987)), 741 through 1648 (Pasek (PCT application No.88/00831)), 816 through 1598 or 741 through 1689 (Lagner (Behring Inst. Mitt. (1988) No 82:16-25, EP 295597)), each of which is incorporated herein by reference in its entirety. In some embodiments, B domain deleted FVIII comprises a partial deletion in B domain, i.e., having 21 amino acids from B domain (i.e., SFSQNSRHPSQNPPVLKRHRQR, which is SEQ ID NO: 17) disclosed in US Publication No. 20100286067 and US Publication No. US 20120093840, both of which are incorporated herein by reference in their entireties. Each of the foregoing deletions can be made in any Factor VIII sequence.

[0133] In one embodiment, the B domain deleted Factor VIII portion in a FVIII polypeptide is processed into two chains connected (or associated) by a metal bond, the first chain comprising a heavy chain (A1-A2-partial B) and a second chain comprising a light chain (A3-C1-C2). In another embodiment, the B domain deleted Factor VIII portion is a single chain Factor VIII. The single chain Factor VIII can comprise an intracellular processing site, which corresponds to Arginine at residue 754 of B domain deleted Factor VIII (residue 773 of SEQ ID NO: 2) or at residue 1648 of full-length Factor VIII (residue 1657 of SEQ ID NO: 6).

[0134] The metal bond between the heavy chain and the light chain can be any metal known in the art. For example, the metals useful for the invention can be a divalent metal ion. The metals that can be used to associate the heavy chain and light chain include, but not limited to, Ca^{2+} , Mn^{2+} , or Cu^{2+} . Fatouros *et al.*, *Intern. J. Pharm.* 155(1): 121-131 (1997); Wakabayashi *et al.*, *JBC*. 279(13): 12677-12684 (2004).

[0135] A FVIII polypeptide used herein can comprise processed Factor VIII or single chain Factor VIII or a combination thereof. "Processed Factor VIII," as used herein means Factor VIII that has been cleaved at Arginine 1648 (for full-length Factor VIII) or Arginine 754 (for B-domain deleted Factor VIII), i.e., intracellular processing site. Due to the cleavage at the intracellular processing site, processed Factor VIII comprises two polypeptide chains, the first chain being a heavy chain and the second chain being a light chain. For example, the processed Factor VIII-Fc fusion protein (i.e., Heavy chain and Light chain fused to Fc) run at approximately 90 kDa and 130 kDa on a non-reducing SDS-PAGE, respectively, and 90 kDa and 105 kDa on a reducing SDS-PAGE, respectively. Therefore, in one embodiment, at least about 50%, about 60%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, about 96%, about 97%, about

98%, about 99%, or about 100% of the Factor VIII portion in the long-acting polypeptide or in the short-acting FVIII polypeptide is processed Factor VIII. In another embodiment, about 50%, about 60%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% of the Factor VIII portion in the long-acting polypeptide or in the short-acting FVIII polypeptide is processed Factor VIII. In a particular embodiment, the FVIII polypeptide comprising processed Factor VIII is purified (or isolated) from the polypeptide comprising single chain Factor VIII, and at least about 75%, about 80%, about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% of the Factor VIII portion in the long-acting polypeptide or in the short-acting FVIII polypeptide is processed Factor VIII. In some embodiments, the FVIII polypeptide comprises about 15% to 25% of single chain FVIII polypeptide and about 75% to about 85% of processed FVIII polypeptide.

[0136] "Single chain Factor VIII," "SC Factor VIII," or "SCFVIII" as used herein means Factor VIII that has not been cleaved at the Arginine site (residue 1648 for full-length Factor VIII (*i.e.*, residue 1667 of SEQ ID NO: 6) or residue 754 for B-domain deleted Factor VIII (*i.e.*, residue 773 of SEQ ID NO: 2). Therefore, single chain Factor VIII in the FVIII polypeptide used herein comprises a single chain. In one embodiment, the single chain Factor VIII contains an intact intracellular processing site. In another embodiment, the single chain Factor VIII of the invention comprises a substitution or mutation at an amino acid position corresponding to Arginine 1645, a substitution or mutation at an amino acid position corresponding to Arginine 1648, or a substitution or mutation at amino acid positions corresponding to Arginine 1645 and Arginine 1648 in full-length Factor VIII. In other embodiments, the amino acid substituted at the amino acid position corresponding to Arginine 1645 is a different amino acid from the amino acid substituted at the amino acid position corresponding to Arginine 1648. In certain embodiments, the substitution or mutation is an amino acid other than arginine, *e.g.*, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan, valine, alanine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, proline, selenocysteine, serine, tyrosine, histidine, ornithine, pyrrolysine, or taurine. The single chain Factor VIII-Fc fusion protein can run at approximately 220 kDa on a non reducing SDS-PAGE and at approximately 195 kDa on a reducing SDS-PAGE.

[0137] The Factor VIII portion in the FVIII polypeptide used herein has Factor VIII activity. Factor VIII activity can be measured by any known methods in the art. For

example, one of those methods can be a chromogenic assay. The chromogenic assay mechanism is based on the principles of the blood coagulation cascade, where activated Factor VIII accelerates the conversion of Factor X into Factor Xa in the presence of activated Factor IX, phospholipids and calcium ions. The Factor Xa activity is assessed by hydrolysis of a p-nitroanilide (pNA) substrate specific to Factor Xa. The initial rate of release of p-nitroaniline measured at 405 nM is directly proportional to the Factor Xa activity and thus to the Factor VIII activity in the sample. The chromogenic assay is recommended by the Factor VIII and Factor IX Subcommittee of the Scientific and Standardization Committee (SSC) of the International Society on Thrombosis and Hemostasis (ISTH). Since 1994, the chromogenic assay has also been the reference method of the European Pharmacopoeia for the assignment of FVIII concentrate potency. Thus, in one embodiment, the FVIII polypeptide comprising single chain Factor VIII has Factor VIII activity comparable to a FVIII polypeptide comprising processed Factor VIII, when the Factor VIII activity is measured in vitro by a chromogenic assay.

[0138] In certain embodiment, a long-acting FVIII polypeptide is a FVIII monomer-dimer hybrid. To obtain a FVIII monomer-dimer hybrid, the coding sequence of human recombinant B-domain deleted FVIII was obtained by reverse transcription-polymerase chain reaction (RT-PCR) from human liver poly A RNA (Clontech) using FVIII-specific primers. The FVIII sequence includes the native signal sequence for FVIII. The B-domain deletion was from serine 743 (S743; 2287 bp) to glutamine 1638 (Q1638; 4969 bp) for a total deletion of 2682 bp. Then, the coding sequence for human recombinant Fc was obtained by RT-PCR from a human leukocyte cDNA library (Clontech) using Fc specific primers. Primers were designed such that the B-domain deleted FVIII sequence was fused directly to the N-terminus of the Fc sequence with no intervening linker. The FVIII-Fc DNA sequence was cloned into the mammalian dual expression vector pBUDCE4.1 (Invitrogen) under control of the CMV promoter. A second identical Fc sequence including the mouse Igk signal sequence was obtained by RT-PCR and cloned downstream of the second promoter, EF1 α , in the expression vector pBUDCE4.1.

[0139] In one embodiment, an FcRn binding partner can be an Fc region. An FcRn binding partner is any molecule that can be specifically bound by the FcRn receptor with consequent active transport by the FcRn receptor of the FcRn binding partner. Thus, the term Fc includes any variants of IgG Fc that are functional. The region of the Fc portion of IgG that binds to the FcRn receptor has been described based on X-ray crystallography

(Burmeister *et al.*, *Nature* 372:379 (1994), incorporated herein by reference in its entirety). The major contact area of the Fc with the FcRn is near the junction of the CH2 and CH3 domains. Fc-FcRn contacts are all within a single Ig heavy chain. The FcRn binding partners include, *e.g.*, whole IgG, the Fc fragment of IgG, and other fragments of IgG that include the complete binding region of FcRn. The major contact sites include amino acid residues 248, 250-257, 272, 285, 288, 290-291, 308-311, and 314 of the CH2 domain and amino acid residues 385-387, 428, and 433-436 of the CH3 domain. References made to amino acid numbering of immunoglobulins or immunoglobulin fragments, or regions, are all based on Kabat *et al.* 1991, Sequences of Proteins of Immunological Interest, U. S. Department of Public Health, Bethesda; MD, incorporated herein by reference in its entirety. (The FcRn receptor has been isolated from several mammalian species including humans. The sequences of the human FcRn, rat FcRn, and mouse FcRn are known (Story *et al.*, *J. Exp. Med.* 180: 2377 (1994), incorporated herein by reference in its entirety.) An Fc can comprise the CH2 and CH3 domains of an immunoglobulin with or without the hinge region of the immunoglobulin. Exemplary Fc variants are provided in WO 2004/101740 and WO 2006/074199, incorporated herein by reference in its entirety.

[0140] Fc (or Fc portion of a chimeric polypeptide) can contain one or more mutations, and combinations of mutations.

[0141] Fc (or Fc portion of a chimeric polypeptide) can contain mutations conferring increased half-life such as M252Y, S254T, T256E, and combinations thereof, as disclosed in Oganessian *et al.*, *Mol. Immunol.* 46:1750 (2009), which is incorporated herein by reference in its entirety; H433K, N434F, and combinations thereof, as disclosed in Vaccaro *et al.*, *Nat. Biotechnol.* 23:1283 (2005), which is incorporated herein by reference in its entirety; the mutants disclosed at pages 1-2, paragraph [0012], and Examples 9 and 10 of US 2009/0264627 A1, which is incorporated herein by reference in its entirety; and the mutants disclosed at page 2, paragraphs [0014] to [0021] of US 20090163699 A1, which is incorporated herein by reference in its entirety.

[0142] Fc (or Fc portion of a chimeric polypeptide) can also include, *e.g.*, the following mutations: The Fc region of IgG can be modified according to well recognized procedures such as site directed mutagenesis and the like to yield modified IgG or Fc fragments or portions thereof that will be bound by FcRn. Such modifications include, *e.g.*, modifications remote from the FcRn contact sites as well as modifications within the

contact sites that preserve or even enhance binding to the FcRn. For example the following single amino acid residues in human IgG1 Fc (Fcγ1) can be substituted without significant loss of Fc binding affinity for FcRn: P238A, S239A, K246A, K248A, D249A, M252A, T256A, E258A, T260A, D265A, S267A, H268A, E269A, D270A, E272A, L274A, N276A, Y278A, D280A, V282A, E283A, H285A, N286A, T289A, K290A, R292A, E293A, E294A, Q295A, Y296F, N297A, S298A, Y300F, R301A, V303A, V305A, T307A, L309A, Q311A, D312A, N315A, K317A, E318A, K320A, K322A, S324A, K326A, A327Q, P329A, A330Q, A330S, P331A, P331S, E333A, K334A, T335A, S337A, K338A, K340A, Q342A, R344A, E345A, Q347A, R355A, E356A, M358A, T359A, K360A, N361A, Q362A, Y373A, S375A, D376A, A378Q, E380A, E382A, S383A, N384A, Q386A, E388A, N389A, N390A, Y391F, K392A, L398A, S400A, D401A, D413A, K414A, R416A, Q418A, Q419A, N421A, V422A, S424A, E430A, N434A, T437A, Q438A, K439A, S440A, S444A, and K447A, where for example P238A represents wildtype proline substituted by alanine at position number 238. In addition to alanine other amino acids can be substituted for the wildtype amino acids at the positions specified above. Mutations can be introduced singly into Fc giving rise to more than one hundred FcRn binding partners distinct from native Fc. Additionally, combinations of two, three, or more of these individual mutations can be introduced together, giving rise to hundreds more FcRn binding partners. Certain of these mutations can confer new functionality upon the FcRn binding partner. For example, one embodiment incorporates N297A, removing a highly conserved N-glycosylation site. The effect of this mutation is to reduce immunogenicity, thereby enhancing circulating half-life of the FcRn binding partner, and to render the FcRn binding partner incapable of binding to FcγRI, FcγRIIA, FcγRIIB, and FcγRIIIA, without compromising affinity for FcRn (Routledge et al. 1995, Transplantation 60:847, which is incorporated herein by reference in its entirety; Friend et al. 1999, Transplantation 68:1632, which is incorporated herein by reference in its entirety; Shields et al. 1995, J. Biol. Chem. 276:6591, which is incorporated herein by reference in its entirety). Additionally, at least three human Fc gamma receptors appear to recognize a binding site on IgG within the lower hinge region, generally amino acids 234-237. Therefore, another example of new functionality and potential decreased immunogenicity can arise from mutations of this region, as for example by replacing amino acids 233-236 of human IgG1 "ELLG" to the corresponding sequence from IgG2 "PVA" (with one amino acid deletion). It has been

shown that FcγRI, FcγRII, and FcγRIII which mediate various effector functions will not bind to IgG1 when such mutations have been introduced (Ward and Ghetie, *Therapeutic Immunology* 2:77 (1995), which is incorporated herein by reference in its entirety; and Armour *et al.*, *Eur. J. Immunol.* 29:2613 (1999), which is incorporated herein by reference in its entirety). As a further example of new functionality arising from mutations described above affinity for FcRn can be increased beyond that of wild type in some instances. This increased affinity can reflect an increased "on" rate, a decreased "off" rate or both an increased "on" rate and a decreased "off" rate. Mutations believed to impart an increased affinity for FcRn include, *e.g.*, T256A, T307A, E380A, and N434A (Shields *et al.*, *J. Biol. Chem.* 276:6591 (2001), which is incorporated herein by reference in its entirety).

[0143] The Fc (or Fc portion of a chimeric polypeptide) can be at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the Fc amino acid sequence shown in Table 11 (amino acids 1458 to 1684 of SEQ ID NO:2 or amino acids 2352 to 2578 of SEQ ID NO:6). The Fc (or Fc portion of a chimeric polypeptide) can be identical to the Fc amino acid sequence shown in Table 11 (amino acids 1458 to 1684 of SEQ ID NO:2 and amino acids 2352 to 2578 of SEQ ID NO:6).

[0144] As a practical matter, whether any particular nucleic acid molecule or polypeptide is at least 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence or polypeptide of the present invention can be determined conventionally using known computer programs. In one embodiment, a method for determining the best overall match between a query sequence (reference or original sequence) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag *et al.*, *Comp. App. Biosci.* 6:237-245 (1990), which is herein incorporated by reference in its entirety. In a sequence alignment the query and subject sequences are both DNA sequences. An RNA sequence can be compared by converting U's to T's. The result of said global sequence alignment is in percent identity. In another embodiment, parameters used in a FASTDB alignment of DNA sequences to calculate percent identity are: Matrix=Unitary, k-tuple=4, Mismatch Penalty=1, Joining Penalty=30, Randomization Group Length=0, Cutoff Score=1, Gap Penalty=5, Gap Size Penalty 0.05, Window Size=500 or the length of the subject nucleotide sequence, whichever is shorter.

[0145] If the subject sequence is shorter than the query sequence because of 5' or 3' deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for 5' and 3' truncations of the subject sequence when calculating percent identity. For subject sequences truncated at the 5' or 3' ends, relative to the query sequence, the percent identity is corrected by calculating the number of bases of the query sequence that are 5' and 3' of the subject sequence, which are not matched/aligned, as a percent of the total bases of the query sequence. Whether a nucleotide is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This corrected score is what is used for the purposes of the present invention. Only bases outside the 5' and 3' bases of the subject sequence, as displayed by the FASTDB alignment, which are not matched/aligned with the query sequence, are calculated for the purposes of manually adjusting the percent identity score.

[0146] For example, a 90 base subject sequence is aligned to a 100 base query sequence to determine percent identity. The deletions occur at the 5' end of the subject sequence and therefore, the FASTDB alignment does not show a matched/alignment of the first 10 bases at 5' end. The 10 unpaired bases represent 10% of the sequence (number of bases at the 5' and 3' ends not matched/total number of bases in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 bases were perfectly matched the final percent identity would be 90%. In another example, a 90 base subject sequence is compared with a 100 base query sequence. This time the deletions are internal deletions so that there are no bases on the 5' or 3' of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only bases 5' and 3' of the subject sequence which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

[0147] By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a query amino acid sequence of the present invention, it is intended that the amino acid sequence of the subject polypeptide is identical to the query sequence except that the subject polypeptide sequence can include up to five amino acid alterations per

each 100 amino acids of the query amino acid sequence. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a query amino acid sequence, up to 5% of the amino acid residues in the subject sequence can be inserted, deleted, (indels) or substituted with another amino acid. These alterations of the reference sequence can occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

[0148] As a practical matter, whether any particular polypeptide is at least 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequences of SEQ ID NO:2 (the factor VIII portion, the Fc portion, individually or together) or 4, or a known factor VIII or Fc polypeptide sequence, can be determined conventionally using known computer programs. In one embodiment, a method for determining the best overall match between a query sequence (reference or original sequence) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag *et al.*, *Comp. App. Biosci.* 6:237-245(1990), incorporated herein by reference in its entirety. In a sequence alignment the query and subject sequences are either both nucleotide sequences or both amino acid sequences. The result of said global sequence alignment is in percent identity. In another embodiment, parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window Size=500 or the length of the subject amino acid sequence, whichever is shorter.

[0149] If the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity. For subject sequences truncated at the N- and C-termini, relative to the query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the subject sequence, which are not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. Whether a residue is matched/aligned is determined by results of the FASTDB sequence

alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of the present invention. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C-terminal residues of the subject sequence.

[0150] For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C-termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

[0151] The polynucleotide variants can contain alterations in the coding regions, non-coding regions, or both. In one embodiment, the polynucleotide variants contain alterations which produce silent substitutions, additions, or deletions, but do not alter the properties or activities of the encoded polypeptide. In another embodiment, nucleotide variants are produced by silent substitutions due to the degeneracy of the genetic code. In other embodiments, variants in which 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination. Polynucleotide variants can be produced for a variety of reasons, *e.g.*, to optimize codon expression for a particular host (change codons in the human mRNA to others, *e.g.*, a bacterial host such as *E. coli*).

[0152] Naturally occurring variants are called "allelic variants," and refer to one of several alternate forms of a gene occupying a given locus on a chromosome of an organism (Genes II, Lewin, B., ed., John Wiley & Sons, New York (1985)). These allelic variants can vary at either the polynucleotide and/or polypeptide level and are included in the present invention. Alternatively, non-naturally occurring variants can be produced by mutagenesis techniques or by direct synthesis.

[0153] Using known methods of protein engineering and recombinant DNA technology, variants can be generated to improve or alter the characteristics of the polypeptides. For instance, one or more amino acids can be deleted from the N-terminus or C-terminus of the secreted protein without substantial loss of biological function. Ron *et al.*, *J. Biol. Chem.* 268: 2984-2988 (1993), incorporated herein by reference in its entirety, reported variant KGF proteins having heparin binding activity even after deleting 3, 8, or 27 amino-terminal amino acid residues. Similarly, Interferon gamma exhibited up to ten times higher activity after deleting 8-10 amino acid residues from the carboxy terminus of this protein. (Dobeli *et al.*, *J. Biotechnology* 7:199-216 (1988), incorporated herein by reference in its entirety.)

[0154] Moreover, ample evidence demonstrates that variants often retain a biological activity similar to that of the naturally occurring protein. For example, Gayle and coworkers (*J. Biol. Chem* 268:22105-22111 (1993), incorporated herein by reference in its entirety) conducted extensive mutational analysis of human cytokine IL-1a. They used random mutagenesis to generate over 3,500 individual IL-1a mutants that averaged 2.5 amino acid changes per variant over the entire length of the molecule. Multiple mutations were examined at every possible amino acid position. The investigators found that "[m]ost of the molecule could be altered with little effect on either [binding or biological activity]." (See Abstract.) In fact, only 23 unique amino acid sequences, out of more than 3,500 nucleotide sequences examined, produced a protein that significantly differed in activity from wild-type.

[0155] As stated above, polypeptide variants include, *e.g.*, modified polypeptides. Modifications include, *e.g.*, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine,

formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation (Mei *et al.*, *Blood* 116:270-79 (2010), which is incorporated herein by reference in its entirety), proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. In some embodiments, Factor VIII is modified, *e.g.*, pegylated, at any convenient location. In some embodiments, Factor VIII is pegylated at a surface exposed amino acid of Factor VIII, *e.g.*, a surface exposed cysteine, which can be an engineered cysteine. *Id.* In some embodiments, modified Factor VIII, *e.g.*, pegylated Factor VIII, is a long-acting Factor VIII.

III. Method of Administering

[0156] Treatment of hemophilia A is a replacement therapy targeting restoration of FVIII activity to 1 to 5 % of normal levels to prevent spontaneous bleeding (Mannucci, P.M. *et al.*, *N. Engl. J. Med.* 344:1773-9 (2001), herein incorporated by reference in its entirety).

[0157] The present invention also includes methods of administering the pharmaceutical composition of the invention. The present invention provides a method of administering a pharmaceutical composition to a human subject in need thereof (*e.g.*, human patient), comprising administering to the subject a therapeutic dose of a pharmaceutical composition comprising a Factor VIII polypeptide, *e.g.*, a long-acting FVIII polypeptide or a short-acting FVIII polypeptide, *e.g.*, a Factor VIII-Fc polypeptide, or a hybrid of such a polypeptide at a dosing interval.

[0158] In some embodiments, the pharmaceutical composition of the invention is used to reduce or decrease one or more bleeding episodes or frequency (*e.g.*, a bleeding condition) in a subject in need thereof. In other embodiments, the pharmaceutical composition of the invention is used to treat or prevent a bleeding condition in a subject in need thereof.

[0159] The bleeding condition can be caused by a blood coagulation disorder. A blood coagulation disorder can also be referred to as a coagulopathy. In one example, the blood coagulation disorder, which can be treated with a pharmaceutical composition of the current disclosure, is hemophilia. In another example, the blood coagulation disorder, which can be treated with a pharmaceutical composition of the present disclosure is hemophilia A.

[0160] In some embodiments, the type of bleeding associated with the bleeding condition is selected from hemarthrosis, muscle bleed, oral bleed, hemorrhage, hemorrhage into muscles, oral hemorrhage, trauma, trauma capitis, gastrointestinal bleeding, intracranial hemorrhage, intra-abdominal hemorrhage, intrathoracic hemorrhage, bone fracture, central nervous system bleeding, bleeding in the retropharyngeal space, bleeding in the retroperitoneal space, and bleeding in the iliopsoas sheath.

[0161] In other embodiments, the subject suffering from bleeding condition is in need of treatment for surgery, including, *e.g.*, surgical prophylaxis or peri-operative management. In one example, the surgery is selected from minor surgery and major surgery. Exemplary surgical procedures include tooth extraction, tonsillectomy, inguinal herniotomy, synovectomy, craniotomy, osteosynthesis, trauma surgery, intracranial surgery, intra-abdominal surgery, intrathoracic surgery, joint replacement surgery (*e.g.*, total knee replacement, hip replacement, and the like), heart surgery, and caesarean section.

[0162] The present invention also provides a method of reducing or decreasing an annualized bleeding rate (ABR) of a subject having hemophilia comprising administering to the subject an effective dose of a FVIII polypeptide. In one embodiment, a long-acting FVIII polypeptide is administered at a dosing interval of every three days or longer. In another embodiment, the effective dose is between about 20IU/kg and about 90IU/kg. In other embodiments, the effective dose is 20-30 IU/kg, 30-40 IU/kg, 40-50 IU/kg, 50-60 IU/kg, 60-70 IU/kg, 70-80 IU/kg, or 80-90 IU/kg. In yet other embodiments, the effective dose is 20 IU/kg, 25 IU/kg, 30 IU/kg, 35 IU/kg, 40 IU/kg, 45 IU/kg, 50 IU/kg, 55 IU/kg, 60 IU/kg, 65 IU/kg, 70 IU/kg, 75 IU/kg, 80 IU/kg, 85 IU/kg, or 90 IU/kg.

[0163] In certain embodiments, administration of a FVIII polypeptide is for individualized (tailored) prophylaxis and results in an ABR of less than about 5.5, less than about 5.4, less than about 5.3, less than about 5.2, less than about 5.1, less than about 5.0, less than about 4.9, less than about 4.8, less than about 4.7, less than about 4.6, or less than about 4.5. In other embodiments, the administration results in an ABR of about 4.7 to 0. In some embodiments, the median of the ABR is about 1.6. In yet other embodiments, the mean of the ABR is about 2.9. In one aspect of the individualized prophylaxis regimen, the effective dose is about 25IU/kg to about 65IU/kg given every three to five days. For example, the effective dose is about 25IU/kg to about 65IU/kg given every three days. In another aspect, the effective dose is about 50IU/kg to about

65IU/kg every four days or every five days. In other aspects, the effective dose is up to 65IU/kg every three days. In still other embodiments, the effective dose of a FVIII polypeptide is modified based on the patient's pharmacokinetic profile. In one example, the patient is administered with two doses initially, a first dose of 25IU/kg on day 1 and a second dose of 50IU/kg on day 4. If the pharmacokinetic data show less than 1% trough level of normal FVIII activity after the initial two doses, the patient is then administered with about 25IU/kg to about 65IU/kg every three days. In another example, if the pharmacokinetic data show higher than 1% trough level of normal FVIII activity for five days from the dosing, the patient is then administered with about 50IU/kg to about 65IU/kg every five days. If the patient experiences more than two spontaneous bleeds over eight weeks period after administration of about 50IU/kg to about 65IU/kg every five days, the patient is administered one of the followings: (1) up to about 65 IU/kg every three days (for the target trough level of up to 5% of normal), (2) about 50IU/kg to about 65IU/kg every four days (for the target trough level up to 5% of normal), or (3) about 25IU/kg to about 65IU/kg every three days (for the target trough level of 1% to 3% of normal). If the patient receiving about 50IU/kg to about 65IU/kg every four days still experiences more than two spontaneous bleeds, the maximum dose can go up to about 65IU/kg every three days. In some examples, the Interquartile Range (IQR) of the ABR for the individualized prophylaxis regimen is 0 to 4.7.

[0164] In some embodiments, administration of a FVIII polypeptide is for weekly prophylaxis and results in an ABR of less than about 9.0, less than about 8.9, less than about 8.8, less than about 8.7, less than about 8.6, less than about 8.5, or less than about 8.4. In one example, the administration results in an ABR between about 8.4 and 0. In another example, the median of the ABR for weekly prophylaxis is about 3.6. In other examples, the mean of the ABR for weekly prophylaxis is about 8.8. In other embodiments, an effective dose for weekly prophylaxis is about 65IU/kg once every week, *e.g.*, every five days, every six days, every seven days, every eight days, or every nine days. In some examples, the IQR of the ABR for the weekly prophylaxis regimen is 1.9 to 8.4.

[0165] In other embodiments, administration of a FVIII polypeptide is for episodic or on-demand treatment and results in an ABR of less than about 55, less than about 54, less than about 53, less than about 52, less than about 51, less than about 50, less than about 49, less than about 48, or less than about 47. In one example, the administration results in

an ABR between about 49 and 0, *e.g.*, between 48.7 and 0. In another example, the median of the ABR for episodic (on-demand) treatment is about 33.5. In other examples, the mean of the ABR for episodic (on-demand) treatment is about 37.23. In other embodiments, an effective dose for on-demand treatment is about 10IU/kg to 75IU/kg every 12 to 24 hours. In some examples, the IQR of the ABR for the on-demand treatment is 21.1 to 48.7.

[0166] In some embodiments, the effective dose for individualized prophylaxis, weekly prophylaxis, or episodic treatment is a fixed dose or a stratified dose. In one aspect, the fixed dose is about 2,000IU per dose, about 2,500IU per dose, about 3,000IU per dose, about 3,500IU per dose, or about 4,000IU per dose.

[0167] The dosing interval for a long-acting FVIII polypeptide can be at least about one and one-half to six times longer, one and one-half to five times longer, one and one-half to four times longer, one and one-half to three times longer, or one and one-half to two times longer, than the dosing interval required for an equivalent dose of the Factor VIII without the non-Factor VIII portion (a polypeptide consisting of the Factor VIII portion), *e.g.*, without the Fc portion. The dosing interval can be at least about one and one-half, two, two and one-half, three, three and one-half, four, four and one-half, five, five and one-half or six times longer than the dosing interval required for an equivalent dose of the Factor VIII without the non-Factor VIII portion (a polypeptide consisting of the Factor VIII portion), *e.g.*, without the Fc portion. The dosing interval can be about every three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, or fourteen days or longer.

[0168] The methods of the invention can be practiced on a subject in need of prophylactic treatment or on-demand treatment.

[0169] For on-demand treatment, the dosing interval of a pharmaceutical composition comprising a FVIII polypeptide is about once every 24-36, 24-48, 24-72, 24-96, 24-120, 24-144, 24-168, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, or 72 hours or longer.

[0170] In one embodiment, the subject is in need of on-demand (episodic) treatment. In another embodiment, on-demand (episodic) treatment resolves greater than 80% (greater than 80%, greater than 81%, greater than 82%, greater than 83%, greater than 84%, greater than 85%, greater than 86%, greater than 87%, greater than 88%, greater than 89%, greater than 90%, greater than 91%, greater than 92%, greater than 93%, greater

than 94%, greater than 95%, greater than 96%, greater than 97%, greater than 98%, greater than 99%, or 100%) or 80-100%, 80-90%, 85-90%, 90-100%, 90-95%, or 95-100% of bleeds (*e.g.*, spontaneous bleeds) in a single dose. In another embodiment, greater than 80% (greater than 81%, greater than 82%, greater than 83%, greater than 84%, greater than 85%, greater than 86%, greater than 87%, greater than 88%, greater than 89%, greater than 90%, greater than 91%, greater than 92%, greater than 93%, greater than 94%, greater than 95%, greater than 96%, greater than 97%, greater than 98%, or 100%) or 80-100%, 80-90%, 85-90%, 90-100%, 90-95%, or 95-100% of bleeding episodes are rated excellent or good by physicians after on-demand (episodic) treatment. In other embodiments, greater than 5%, (greater than 6%, greater than 7%, greater than 8%, greater than 9%, greater than 10%, greater than 11%, greater than 12%, greater than 13%, greater than 14%, greater than 15%, greater than 16%, greater than 17%, greater than 18%, greater than 19%, greater than 20%), or 5-20%, 5-15%, 5-10%, 10-20%, or 10-15% of bleeding episodes are rated as fair by physicians after on-demand treatment.

[0171] In other embodiments, the pharmaceutical composition comprising a FVIII polypeptide is used for prophylaxis. Prophylaxis can be demonstrated by better C_{max}, better T_{max}, and/or greater mean residence time versus short-acting FVIII. In some embodiments, prophylaxis results in no spontaneous bleeding episodes within about 24, 36, 48, 72, or 96 hours (*e.g.*, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 96, 87, 88, 89, 90, 91, 92, 93, 94, 95, or 96 hours), after injection (*e.g.*, the last injection). In certain embodiments, prophylaxis results in greater than 30% (*e.g.*, greater than 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 96, 87, 88, 89, or 90%, for example, greater than 50%), mean reduction in annualized bleeding episodes with once weekly dosing (*e.g.*, at 65 IU/kg).

[0172] In another example, the subject is concomitantly treated with FIX. Because the compounds of the invention are capable of activating FIXa, they could be used to pre-activate the FIXa polypeptide before administration of the FIXa to the subject.

IV. Method of Making

[0173] A FVIII polypeptide can be manufactured in a host cell comprising a vector encoding the FVIII polypeptide. In one embodiment, the host cell is transformed with one or more vectors comprising a first nucleotide sequence encoding a FVIII polypeptide and a first FcRn polypeptide, a second nucleotide sequence encoding a second FcRn polypeptide, and optionally a third nucleotide sequence encoding a protein convertase, *e.g.*, PC5. As used herein, an expression vector refers to any nucleic acid construct which contains the necessary elements for the transcription and translation of an inserted coding sequence, or in the case of an RNA viral vector, the necessary elements for replication and translation, when introduced into an appropriate host cell. Expression vectors can include plasmids, phagemids, viruses, and derivatives thereof.

[0174] A gene expression control sequence as used herein is any regulatory nucleotide sequence, such as a promoter sequence or promoter-enhancer combination, which facilitates the efficient transcription and translation of the coding nucleic acid to which it is operably linked. The gene expression control sequence can, for example, be a mammalian or viral promoter, such as a constitutive or inducible promoter. Constitutive mammalian promoters include, but are not limited to, the promoters for the following genes: hypoxanthine phosphoribosyl transferase (HPRT), adenosine deaminase, pyruvate kinase, beta-actin promoter, and other constitutive promoters. Exemplary viral promoters which function constitutively in eukaryotic cells include, for example, promoters from the cytomegalovirus (CMV), simian virus (*e.g.*, SV40), papilloma virus, adenovirus, human immunodeficiency virus (HIV), Rous sarcoma virus, cytomegalovirus, the long terminal repeats (LTR) of Moloney leukemia virus, and other retroviruses, and the thymidine kinase promoter of herpes simplex virus. Other constitutive promoters are known to those of ordinary skill in the art. The promoters useful as gene expression sequences of the invention also include inducible promoters. Inducible promoters are expressed in the presence of an inducing agent. For example, the metallothionein promoter is induced to promote transcription and translation in the presence of certain metal ions. Other inducible promoters are known to those of ordinary skill in the art.

[0175] Examples of vectors include, but are not limited to viral vectors or plasmid vectors. Plasmid vectors have been extensively described in the art and are well-known to those of skill in the art. See, *e.g.*, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, 1989. In the last few

years, plasmid vectors have been found to be particularly advantageous for delivering genes to cells *in vivo* because of their inability to replicate within and integrate into a host genome. These plasmids, however, having a promoter compatible with the host cell, can express a peptide from a gene operably encoded within the plasmid. Some commonly used plasmids available from commercial suppliers include pBR322, pUC18, pUC19, various pcDNA plasmids, pRC/CMV, various pCMV plasmids, pSV40, and pBlueScript. Additional examples of specific plasmids include pcDNA3.1, catalog number V79020; pcDNA3.1/hygro, catalog number V87020; pcDNA4/myc-His, catalog number V86320; and pBudCE4.1, catalog number V53220, all from Invitrogen (Carlsbad, CA.). Other plasmids are well-known to those of ordinary skill in the art. Additionally, plasmids can be custom designed using standard molecular biology techniques to remove and/or add specific fragments of DNA.

[0176] The expression vector or vectors are then transfected or co-transfected into a suitable target cell, which will express the polypeptides. Transfection techniques known in the art include, but are not limited to, calcium phosphate precipitation (Wigler *et al.* (1978) *Cell* 14:725), electroporation (Neumann *et al.* (1982) *EMBO J* 1:841), and liposome-based reagents. A variety of host-expression vector systems can be utilized to express the proteins described herein including both prokaryotic and eukaryotic cells. These include, but are not limited to, microorganisms such as bacteria (*e.g.*, *E. coli*) transformed with recombinant bacteriophage DNA or plasmid DNA expression vectors containing an appropriate coding sequence; yeast or filamentous fungi transformed with recombinant yeast or fungi expression vectors containing an appropriate coding sequence; insect cell systems infected with recombinant virus expression vectors (*e.g.*, baculovirus) containing an appropriate coding sequence; plant cell systems infected with recombinant virus expression vectors (*e.g.*, cauliflower mosaic virus or tobacco mosaic virus) or transformed with recombinant plasmid expression vectors (*e.g.*, Ti plasmid) containing an appropriate coding sequence; or animal cell systems, including mammalian cells (*e.g.*, HEK 293, CHO, Cos, HeLa, HKB11, and BHK cells).

[0177] In one embodiment, the host cell is a eukaryotic cell. As used herein, a eukaryotic cell refers to any animal or plant cell having a definitive nucleus. Eukaryotic cells of animals include cells of vertebrates, *e.g.*, mammals, and cells of invertebrates, *e.g.*, insects. Eukaryotic cells of plants specifically can include, without limitation, yeast cells. A eukaryotic cell is distinct from a prokaryotic cell, *e.g.*, bacteria.

[0178] In certain embodiments, the eukaryotic cell is a mammalian cell. A mammalian cell is any cell derived from a mammal. Mammalian cells specifically include, but are not limited to, mammalian cell lines. In one embodiment, the mammalian cell is a human cell. In another embodiment, the mammalian cell is a HEK 293 cell, which is a human embryonic kidney cell line. HEK 293 cells are available as CRL-1533 from American Type Culture Collection, Manassas, VA, and as 293-H cells, Catalog No. 11631-017 or 293-F cells, Catalog No. 11625-019 from Invitrogen (Carlsbad, Calif.). In some embodiments, the mammalian cell is a PER.C6[®] cell, which is a human cell line derived from retina. PER.C6[®] cells are available from Crucell (Leiden, The Netherlands). In other embodiments, the mammalian cell is a Chinese hamster ovary (CHO) cell. CHO cells are available from American Type Culture Collection, Manassas, VA. (*e.g.*, CHO-K1; CCL-61). In still other embodiments, the mammalian cell is a baby hamster kidney (BHK) cell. BHK cells are available from American Type Culture Collection, Manassas, Va. (*e.g.*, CRL-1632). In some embodiments, the mammalian cell is a HKB11 cell, which is a hybrid cell line of a HEK293 cell and a human B cell line. Mei et al., *Mol. Biotechnol.* 34(2): 165-78 (2006).

[0179] The method can further comprise purification steps. Various known purifications steps are well known in the art.

Examples

Example 1. Product Description

[0180] rFVIII^h is a long-acting, fully recombinant fusion protein consisting of human coagulation Factor VIII (FVIII) covalently linked to the Fc domain of human immunoglobulin G1 (IgG1). The Factor VIII portion of rFVIII^h has a primary amino acid sequence and post-translational modifications that are comparable to the 90 + 80 kDa form of Factor VIII (*i.e.*, B-domain deleted). The Fc domain of rFVIII^h contains the hinge, CH2 and CH3 regions of IgG1. rFVIII^h contains 1882 amino acids with an apparent molecular weight of approximately 220 kilodaltons.

[0181] rFVIII^h is produced by recombinant DNA technology in a human embryonic kidney (HEK) cell line, which has been extensively characterized. The cell line expresses rFVIII^h into a defined cell culture medium that does not contain any proteins derived from animal or human sources. rFVIII^h is purified by a series of chromatography steps

that does not require use of a monoclonal antibody. The process includes a detergent viral inactivation step and multiple viral clearance steps including an affinity chromatography step and a 15nm virus-retaining nano-filtration step. No human or animal additives are used in the cell culture, purification, and formulation processes.

[0182] rFVIII^h is in the pharmacotherapeutic group: antihemorrhagics, blood coagulation factor VIII. It is provided as a sterile, preservative-free, non-pyrogenic, lyophilized, white to off-white powder to cake, for intravenous (IV) administration in a single-use vial, accompanied by a liquid diluent in a pre-filled syringe. In addition to rFVIII^h, the pharmaceutical composition comprises in the lyophilizate sucrose, sodium chloride, L-Histidine, calcium chloride, and Polysorbate 20 or polysorbate 80, and comprising in sterilized water for injections. Each single-use vial contains nominally 250, 500, 750, 1000, 1500, 2000, 3000, 4000, 5000, or 6000 International Units (IU) of rFVIII^h. When reconstituted with provided diluent, the product contains the following excipients: sucrose, sodium chloride, L-histidine, calcium chloride, and polysorbate 20 or polysorbate 80, at the concentrations shown in Table 1 or Table 2 below. The pharmaceutical composition is formulated for intravenous administration only after reconstitution.

[0183] Each pack contains a powder vial (type 1 glass) with a stopper (butyl) and a flip-off seal (aluminum), 3 ml solvent in a pre-filled syringe (type 1 glass) with a plunger stopper (butyl), a tip-cap (butyl), and a sterile vial adapter reconstitution device.

TABLE 1: rFVIII^h Formulations Following Reconstitution with Liquid Diluent

rFVIII ^h IU/ml*	Sucrose % (w/v)	NaCl (mM)	L-histidine (mM)	Calcium chloride (mM)	Polysorbate- 20 % (w/v)
83 IU/ml	1.33	205	6.64	5.4	0.013
167 IU/ml	1.33	205	6.64	5.4	0.013
250 IU/ml	1.33	205	6.64	5.4	0.013
333 IU/ml	1.33	205	6.64	5.4	0.013
500 IU/ml	1.33	205	6.64	5.4	0.013
667 IU/ml	1.33	205	6.64	5.4	0.013
1000 IU/ml	1.33	205	6.64	5.4	0.013
1333 IU/ml	1.33	205	6.64	5.4	0.013
1667 IU/ml	1.33	205	6.64	5.4	0.013
2000 IU/ml	1.33	205	6.64	5.4	0.013

TABLE 2: rFVIII Fc Formulations Following Reconstitution with Liquid Diluent

Component	Concentration									
	250 IU/vial	500 IU/vial	750 IU/vial	1000 IU/vial	1500 IU/vial	2000 IU/vial	3000 IU/vial	4000 IU/vial	5000 IU/vial	6000 IU/vial
rFVIII Fc*	83 IU/mL	167 IU/mL	250 IU/mL	333 IU/mL	500 IU/mL	667 IU/mL	1000 IU/mL	1333 IU/mL	1667 IU/mL	2000 IU/mL
Sucrose	13.3 mg/mL	13.3 mg/mL	13.3 mg/mL	13.3 mg/mL	13.3 mg/mL	13.3 mg/mL	13.3 mg/mL	13.3 mg/mL	13.3 mg/mL	13.3 mg/mL
Sodium Chloride	12.0 mg/mL	12.0 mg/mL	12.0 mg/mL	12.0 mg/mL	12.0 mg/mL	12.0 mg/mL	12.0 mg/mL	12.0 mg/mL	12.0 mg/mL	12.0 mg/mL
L-Histidine	1.03 mg/mL	1.03 mg/mL	1.03 mg/mL	1.03 mg/mL	1.03 mg/mL	1.03 mg/mL	1.03 mg/mL	1.03 mg/mL	1.03 mg/mL	1.03 mg/mL
Calcium Chloride Dihydrate	0.80 mg/mL	0.80 mg/mL	0.80 mg/mL	0.80 mg/mL	0.80 mg/mL	0.80 mg/mL	0.80 mg/mL	0.80 mg/mL	0.80 mg/mL	0.80 mg/mL
Polysorbate 20	0.13 mg/mL	0.13 mg/mL	0.13 mg/mL	0.13 mg/mL	0.13 mg/mL	0.13 mg/mL	0.13 mg/mL	0.13 mg/mL	0.13 mg/mL	0.13 mg/mL
*The potency (IU) is determined using the European Pharmacopoeia chromogenic assay against an in-house standard that is referenced to the WHO standard. The specific activity of rFVIII Fc is 4000-10000 IU/mg protein.										

Example 2. Method of Formulation

[0184] The rFVIII Fc drug product is a sterile lyophilized powder for injection intended for intravenous administration. It is supplied in aseptically filled single use vials which contain nominally 250, 500, 750, 1000, 1500, 2000, 3000, 4000, 5000, and 6000 IU per vial. The vials are 10 mL USP/Ph. Eur. Type 1 glass vials sealed with a 20 mm Teflon-coated butyl rubber lyophilization stopper and aluminum flip-off crimp seal. Prior to lyophilization, the nominal fill volume target for 250 through 6000 IU vials is 3 mL. The composition of the formulation excipients prior to lyophilization is the same for all dosage strengths. The powder for injection is reconstituted with 3 mL of diluent comprising sterilized water for injections supplied in a sterile prefilled syringe.

[0185] The compositions of the drug product solutions prior to lyophilization are presented in Table 3. The compositions of the drug products following reconstitution are presented in Table 1 or in Table 2. (Example 1).

TABLE 3: Pre-lyophilization Concentrations of rFVIII Fc Formulation Components

Component	Concentration									
	250	500	750	1000	1500	2000	3000	4000	5000	6000
	IU/vial	IU/vial	IU/vial	IU/vial	IU/vial	IU/vial	IU/vial	IU/vial	IU/vial	IU/vial
rFVIII Fc	150	287.5	431.25	575	862.5	1150	1725	2300	2875	3450
	IU/ml*	IU/ml**	IU/ml**	IU/ml**	IU/ml**	IU/ml**	IU/ml**	IU/ml**	IU/ml**	IU/ml**
Sucrose	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0
	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml
Sodium Chloride	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0
	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml
L-Histidine	1.55	1.55	1.55	1.55	1.55	1.55	1.55	1.55	1.55	1.55
	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml
Calcium Chloride Dihydrate	1.18	1.18	1.18	1.18	1.18	1.18	1.18	1.18	1.18	1.18
	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml
Polysorbate 20	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml
Water for Injection	To make 1 ml ***									
Fill Volume	2.0 ml	2.0 ml	2.0 ml	2.0 ml	2.0 ml	2.0 ml	2.0 ml	2.0 ml	2.0 ml	2.0 ml
* A 20% overage is included for the 250 IU/vial strength to ensure that, after process and testing variations, at least the nominal concentration is present in the vial.										
** A 15% overage is included for the 500-6000 IU/vial strengths to ensure that, after process and testing variations, at least the nominal concentration is present in the vial.										
*** Removed during lyophilization.										

Example 3. Dosage and Method of Administration/ Method of Calculating Initial Estimated Dose

[0186] rFVIII Fc is long-acting anti-hemophilic factor (recombinant) indicated in adults and children (≥ 12 years) with hemophilia A (congenital Factor FVIII deficiency) for, e.g., control and prevention of bleeding episodes, routine prophylaxis to prevent or reduce the frequency of bleeding episodes, and perioperative management (surgical prophylaxis).

[0187] Dosing of rFVIII Fc can be estimated as described in this example, but can also be determined by standard tests such as FVIII activity assays described elsewhere herein.

[0188] 1 IU of rFVIII^{IFc} per kg body weight is expected to increase the circulating level of Factor VIII by 2 [IU/dL]. rFVIII^{IFc} has been shown to have a prolonged circulating half-life.

[0189] Since patients can vary in their pharmacokinetic (*e.g.*, half-life, *in vivo* recovery) and clinical responses to rFVIII^{IFc}, the expected *in vivo* peak increase in Factor VIII level expressed as IU/dL (or % of normal) or the required dose can be estimated using the following formulas:

$$\text{IU/dL (or \% of normal)} = [\text{Total Dose (IU)/body weight (kg)}] \times 2 \text{ (IU/dL per IU/kg)}$$

OR

$$\text{Dose (IU)} = \text{body weight (kg)} \times \text{Desired Factor VIII Rise (IU/dL or \% of normal)} \times 0.5 \text{ (IU/kg per IU/dL)}$$

[0190] The following table (Table 4) can be used to guide dosing in bleeding episodes:

TABLE 4: Guide to rFVIII^{IFc} Dosing for Treatment of Bleeding

Severity of Bleed	Desired Factor VIII Level (IU/dL or % of normal)	Dose (IU/kg)/Frequency of Doses (hrs)
Minor and Moderate For example: joint, superficial muscle/no neurovascular compromise (except iliopsoas), deep laceration and renal, superficial soft tissue, mucous membranes	40-60	20-30 IU/kg Repeat every 24-48 hours until bleeding is resolved
Major For example: iliopsoas and deep muscle with neurovascular injury, or substantial blood loss, retroperitoneum, CNS, throat and neck, gastrointestinal.	80-100	40-50 IU/kg Repeat every 12-24 hours until bleeding is resolved

Adapted from WFH 2012

[0191] Subsequent dosage and duration of treatment depends on the individual clinical response, the severity of the Factor VIII deficiency, and the location and extent of bleeding.

[0192] The following table (Table 5) can be used to guide dosing for perioperative management (surgical prophylaxis):

TABLE 5: Guide to rFVIII Fc Dosing for Perioperative Management (Surgical Prophylaxis)

Type of Surgery	Target Factor VIII Level (IU/dL or % of normal)	Dose (IU/kg)/Frequency of Doses (hrs)
Minor Minor operations including uncomplicated dental extraction	50 to 80	25-40 IU/kg A single infusion can be sufficient. Repeat every 24 hours as needed to control bleeding.
Major Major operations including intra-abdominal, joint replacement surgery	80 to 120	An initial preoperative dose of 40-60 IU/kg followed by a repeat dose of 40-50 IU/kg after 8-24 hours and then every 24 hours to maintain FVIII activity within the target range. rFVIII Fc has a longer half-life than plasma and recombinant FVIII products (see Example 5)

[0193] For routine prophylaxis, the recommended regimen is 50 IU/kg every 3-5 days. The dose can be adjusted based on patient response in the range of 25-65 IU/kg.

[0194] For weekly prophylaxis, the recommended dose is 65 IU/kg.

[0195] rFVIII Fc is contraindicated in patients who have manifested severe hypersensitivity reactions, including anaphylaxis, to the product or its components. Severe hypersensitivity reactions were not observed in clinical trials; however, these have been known to occur with use of other factor VIII replacement factors.

[0196] The clinical response to rFVIII Fc can vary. If bleeding is not controlled with the recommended dose, the plasma level of Factor VIII can be determined, and a sufficient dose of rFVIII Fc can be administered to achieve a satisfactory clinical response. If the patient's plasma Factor VIII level fails to increase as expected or if bleeding is not controlled after rFVIII Fc administration, the presence of an inhibitor (neutralizing antibodies) should be suspected, and appropriate testing performed. Patients using

rFVIII-Fc can be monitored for the development of Factor VIII inhibitors by appropriate clinical observations and laboratory tests known to those of ordinary skill in the art.

- [0197] Patient's plasma can be monitored for Factor VIII activity levels, e.g., the one-stage clotting assay to confirm adequate Factor VIII levels have been achieved and maintained, when clinically indicated. Patient's plasma can further be monitored for the development of Factor VIII inhibitors.

Example 4. Phase 3 Clinical Trial of Extended Half-Life Recombinant Fc Fusion Factor VIII ("A-LONG" Study)

Summary of A-LONG study

- [0198] In the A-LONG study, 165 male patients aged 12 years of age and older were enrolled. The A-LONG study had three treatment arms: individualized prophylaxis, weekly prophylaxis and episodic (on-demand) treatment (Arms 1, 2 and 3, respectively). In a subgroup of patients across treatment arms, rFVIII-Fc was evaluated in the perioperative management of patients who required a major surgical procedure during the study.
- [0199] Overall, 92.7 percent of patients completed the study. Recombinant FVIII-Fc was generally well-tolerated. No inhibitors to rFVIII-Fc were detected and no cases of anaphylaxis were reported in any patients, all of whom switched from commercially-available Factor VIII products. No serious adverse events were assessed to be related to drug by the investigator.
- [0200] The most common adverse events (incidence of ≥ 5 percent) occurring outside of the perioperative management period were nasopharyngitis, arthralgia, headache, and upper respiratory tract infection.
- [0201] The median (mean) annualized bleeding rates (ABR), including spontaneous and traumatic bleeds, were 1.6 (2.9) in the individualized prophylaxis arm, 3.6 (8.8) in the weekly prophylaxis arm and 33.6 (37.2) in the episodic treatment arm. In the individualized prophylaxis arm, the median dosing interval was 3.5 days. During the last 3 months on study, 30 percent of patients in the individualized prophylaxis arm achieved a mean dosing interval of at least 5 days.

- [0202] Control of bleeding was assessed in all patients who experienced a bleeding episode during the study. Overall, 98 percent of bleeding episodes were controlled by one or two injections of rFVIII Fc.
- [0203] In addition, rFVIII Fc was assessed in the perioperative management of 9 patients undergoing 9 major surgical procedures. The treating physicians rated the hemostatic efficacy of rFVIII Fc as excellent or good in 100 percent of these surgeries.
- [0204] A-LONG included pharmacokinetic (PK) analysis of rFVIII Fc in all patients in the study. In a protocol-defined subset of patients with extensive PK sampling, the approximate terminal half-life of rFVIII Fc was 19.0 hours compared to 12.4 hours for ADVATE® [Antihemophilic Factor (Recombinant), Plasma/Albumin-Free Method], consistent with the results obtained in the Phase 1/2 study of rFVIII Fc.

About the A-LONG Study and the rFVIII Fc Program

- [0205] A-LONG was a global, open-label, multi-center Phase 3 study that evaluated the efficacy, safety and pharmacokinetics of intravenously-injected rFVIII Fc. The study was designed to evaluate rFVIII Fc in the control and prevention of bleeding, routine prophylaxis and perioperative management in patients with hemophilia A. A-LONG involved 60 hemophilia treatment centers in 19 countries on 6 continents.
- [0206] The A-LONG study had three treatment arms. In Arm 1 (individualized prophylaxis; n=117), patients were treated with 25-65 IU/kg of rFVIII Fc, at an interval of every three to five days, which was individualized to maintain factor trough levels sufficient to prevent bleeding. In Arm 2 (weekly prophylaxis; n=24), patients were treated with a weekly dose of 65 IU/kg. In Arm 3 (episodic treatment; n=23), patients received rFVIII Fc episodic treatment as needed for bleeding. In a subgroup of patients across treatment arms, rFVIII Fc was evaluated in the surgical setting.
- [0207] The primary efficacy measures were the per-patient annualized bleeding rate in Arm 1 vs. Arm 3, and pharmacokinetics of rFVIII Fc vs. rFVIII (one-stage clotting [activated partial thromboplastin time] assay and the chromogenic assays calibrated to normal human references plasma with potency traceable to World Health Organization standards). The safety endpoints included the incidence of adverse events and inhibitor development (Nijmegen-modified Bethesda assay) in patients studied for up to 54 weeks. Secondary efficacy endpoints included ABR in Arm 2 vs. Arm 3, annualized number of spontaneous and joint bleeding episodes per patient, number of injections and dose per

injections to resolve a bleed, and investigators' assessments of patients' response to surgery with rFVIII-Fc using a bleeding response scale (surgery subgroup only), response to treatment of bleeding episodes and the pharmacokinetics of rFVIII-Fc versus ADVATE®. Non-neutralizing antibodies (NNAs) were assessed with an electrochemiluminescence-based anti-rFVIII-Fc binding-antibody assay.

[0208] Ongoing clinical studies of rFVIII-Fc include the Kids A-LONG and ASPIRE studies. Kids A-LONG is a Phase 3, open-label study in previously-treated children with hemophilia A under age 12, which is actively recruiting patients. ASPIRE is a long-term open-label study for patients who completed the A-LONG study or who complete the Kids A-LONG study.

A-LONG Study Design

[0209] Design: Global, open-label, multicenter, Phase 3 study. The study protocol was approved by local Institutional Review Boards for each participating institution, and the study was conducted in accordance with the International Conference on Harmonisation guidelines for Good Clinical Practice.

[0210] Objectives: To evaluate the efficacy and safety of intravenously-injected recombinant factor VIII Fc fusion protein (rFVIII-Fc) in the control and prevention of bleeding episodes, routine prophylaxis, and perioperative management in individuals with severe hemophilia A.

[0211] **Key inclusion criteria:**

Male

≥12 years of age and at least 40 kg

Diagnosis of severe hemophilia A defined as <1% (<1 IU/dL) endogenous factor VIII (FVIII) activity

Previously Treated Patients (PTPs): history of ≥150 prior documented exposure days (EDs) with any currently marketed FVIII product, been on prophylaxis ≥ 2 times per week with a FVIII product (for Arm 1 only), or experienced ≥ 12 bleeds in the past 12 months on an episodic regimen.

No current/prior FVIII inhibitors of measurable activity.

No history of inhibitors, or history of hypersensitivity or anaphylaxis associated with any FVIII or intravenous immunoglobulin administration

All subjects (or their guardians in the case of minors) gave informed written consent prior to study participation.

[0212] **Treatment arms:** Details of A-LONG study design is described below.

Arm 1 (individualized prophylaxis) was administered 25-65 IU/kg every 3-5 days (maximum 5-day dosing interval). Subjects were treated with an initial dose of 25 IU/kg on Day 1 and 50 IU/kg on Day 4, which was subsequently adjusted to maintain trough factor levels sufficient to prevent bleeding. Further dose adjustments to target trough levels of 3 to 5 IU/dL permitted after week 7 if subject experienced ≥ 2 moderate/severe bleeds over a rolling 8-week period; 10 to 20 IU/kg rFVIIIIFc (target 20–40 IU/dL FVIII) for minor bleeding episodes; 15 to 30 IU/kg rFVIIIIFc (target 30–60 IU/dL FVIII) for moderate to major bleeding episodes; 40 to 50 IU/kg rFVIIIIFc (target 80–100 IU/dL FVIII) for major to life-threatening bleeding episodes.

[0213] Arm 2 (weekly prophylaxis) was administered 65 IU/kg dose. Subjects were treated with 65 IU/kg once weekly with no dose or interval adjustment.

[0214] Arm 3 (episodic [on-demand] treatment) was administered 10-50 IU/kg. Subjects received rFVIIIIFc episodic treatment as needed for bleeding.

A Perioperative Management Subgroup was established. In this subgroup, rFVIIIIFc was administered prior to and following surgery in the subset of patients requiring a major surgical procedure during the study. Subjects in any treatment arm could be enrolled in the surgery subgroup. Eligibility: required major surgery; ≥ 12 exposure days to rFVIIIIFc with negative inhibitor titre following this period and within 4 weeks prior to surgery; and, completed, at minimum, abbreviated PK sampling.

[0215] All subjects (excluding sequential PK subgroup) underwent PK sampling from pre-injection with rFVIIIIFc up to 96-hours post-injection, according to schedule: pre-injection, 30 (± 3) minutes, 3 hours (± 15 minutes), 72 (± 2) hours (Day 3), and 96 (± 2) hours (Day 4) from start of injection.

[0216] Pharmacokinetic (PK) Assessment was performed. All subjects in all arms had an initial PK assessment after their first dose of rFVIIIIFc. A subset of subjects from Arm 1 were assigned to a protocol-specified sequential PK subgroup to compare the stability of PK properties of rFVIIIIFc over time with that of recombinant factor VIII (rFVIII,

ADVATE® [anti-hemophilic factor (recombinant) plasma/albumin-free method], octocog alfa) as follows:

Prior to treatment in Arm 1, PK was assessed after a single dose of ADVATE® 50 IU/kg. PK was then assessed in these same subjects after a single dose of rFVIII Fc 50 IU/kg.

PK of rFVIII Fc was repeated at 12 to 24 weeks.

[0217] Details on the design of the sequential PK subgroup (Arm 1) dosing and PK sampling is described in Figure 1

[0218] Key efficacy outcome measures (included in initial readout):

- a. Annualized bleeding rate (ABR) in Arm 1 versus Arm 3
- b. Individualized prophylaxis arm compared with episodic treatment arm.
- c. Number of injections required to resolve a bleeding episode
- d. Number of injections required to resolve bleeding episodes
- e. Median dose required to resolve bleeding episodes
- f. Treating physicians' assessments of subjects' response to surgery with rFVIII Fc using a 4-point scale

[0219] PK outcome measures included:

- a. The primary PK assessment were based on FVIII activity levels determined at a central laboratory by one-stage clotting assay and by the chromogenic assay against commercially available plasma standards.
- b. PK of rFVIII Fc and ADVATE®
- c. PK properties of rFVIII Fc were compared with rFVIII (ADVATE®; sequential PK subgroup). Overall study duration was ≤ 75 weeks for all subjects. The primary efficacy endpoint was annualized bleeding rate (ABR; Arms 1 and 2 vs. Arm 3). Prophylaxis dose and interval, number of injections required for treatment of bleeding episodes, and perioperative haemostasis were evaluated.

[0220] Key safety outcome measures included:

- a. Incidence of inhibitor development. The study was powered to detect the occurrence of inhibitors with a 2-sided 95% confidence interval using the Clopper-Pearson exact method if 2 cases of inhibitor formation was observed.
- b. Incidence of adverse events (AEs) occurring outside of the perioperative management period

[0221] Adherence with treatment (adherence measures) was assessed using patient electronic diaries.

A-LONG Results

Subjects

[0222] A total of 165 subjects were enrolled in the study. Median age was 30 years (range, 12-65) and 8% were <18 years. The number of patients in each arm of the study were: Arm 1 (individualized prophylaxis), n = 118; Arm 2 (weekly prophylaxis), n=24; Arm 3 (episodic treatment), n = 23; and perioperative management subgroup, n = 9, 9 surgeries (8 subjects from Arm 1 and 1 from Arm 2). Subjects from each treatment arm were eligible to enter the surgery subgroup if they required major surgery, had ≥ 12 EDs to rFVIII-Fc and a negative inhibitor titre following this period and within 4 weeks prior to surgery. Patients on prior episodic treatment had a higher median number of bleeding episodes in the 12 months prior to the study, and a higher proportion of patients in Arms 2 and 3 had target joints. For patients on prior prophylaxis, 87% reported injecting at least three times weekly.

[0223] 92.7% of subjects completed the study. In total, 153 (approximately 93%) subjects completed the study; 112/118 (95%) in Arm 1, 19/24 (79%) in Arm 2, and 22/23 (96%) in Arm 3. Reasons for premature discontinuation were: Arm 1 – subject withdrawal (n=2), physician decision (n=2), other (n=1), death (n=1); Arm 2 – subject withdrawal (n=2; adverse events (AEs) related to study drug can have contributed), AEs (n=2), other (n=1); Arm 3 – other (n=1).

[0224] Age, race, and geography of subjects were representative of the global haemophilia A population who have access to treatment, as summarized in TABLE 6.

TABLE 6. A-LONG subjects demographics

Demographic	Arm 1 (N=118)	Arm 2 (N=24)	Arm 3 (N=23)	Total N=165
Age (years), median (min-max)	29.0 (12-65)	31.5 (18-59)	34.0 (13-62)	30.0 (12-65)
Weight (kg), median (min-max)	71.65 (42.0- 127.4)	75.85 (50.0- 105.0)	70.00 (48.0- 110.4)	71.60 (42.0- 127.4)
BMI (kg/m ²) median (min-max)	23.90 (15.3- 37.1)	24.60 (18.8- 37.4)	22.80 (17.2- 35.6)	23.90 (15.3- 37.4)
Race, n (%)				
White	79 (66.9)	12 (50.0)	16 (69.6)	107 (64.8)
Black	7 (5.9)	1 (4.2)	2 (8.7)	10 (6.1)
Asian	27 (22.9)	11 (45.8)	5 (21.7)	43 (26.1)
Other	5 (4.2)	0	0	5 (3.0)
Geographic location — n (%)				
Europe	34 (28.8)	3 (12.5)	4 (17.4)	41 (24.8)
North America	44 (37.3)	5 (20.8)	7 (30.4)	56 (33.9)
Other*	40 (33.9)	16 (66.7)	12 (52.2)	68 (41.2)
Genotype — n (%)				
Intron 22 inversion	41 (35.0)	7 (33.3)	9 (39.1)	57 (35.4)
Frameshift	24 (20.5)	4 (19.0)	6 (26.1)	34 (21.1)
Missense mutation	22 (18.8)	4 (19.0)	1 (4.3)	27 (16.8)
Nonsense mutation	19 (16.2)	6 (28.6)	1 (4.3)	26 (16.1)
Splice site change	7 (6.0)	0	4 (17.4)	11 (6.8)
Intron 1 inversion	3 (2.6)	0	1 (4.3)	4 (2.5)
Duplication	1 (0.9)	0	0	1 (0.6)
NA	0	0	1 (4.3)	1 (0.6)
von Willebrand factor antigen — median IU/dl (IQR)	118.0 (85, 151)	129.0 (86, 166)	131.0 (83, 155)	118.0 (85, 153)
Pre-study FVIII regimen — n (%)				
Prophylaxis	87 (73.7)	0	0	87 (52.7)
Episodic	31 (26.3)	24 (100)	23 (100)	78 (47.3)
Estimated no. of bleeds prior 12 mo — median (IQR) †				
Prior prophylaxis	6.0 (2, 15)	--	--	6 (2, 15)
Prior episodic	27.0 (17, 41)	29.5 (19, 44)	24.0 (15, 36)	27 (18, 40)

≥1 Target joint – n (%)				
Prior prophylaxis	47 (39.8)	–	–	47 (28.5)
Prior episodic	26 (22.0)	22 (91.7)	18 (78.3)	66 (40.0)
Family history of inhibitor	4 (3.4)	1 (4.2)	2 (8.7)	7 (4.2)
HIV positive	25 (21.2)	4 (16.7)	7 (30.4)	36 (21.8)
HCV positive	55 (46.6)	14 (58.3)	13 (56.5)	82 (49.7)

BMI, body mass index; HIV, human immunodeficiency virus; HCV, hepatitis C virus

IU/dl denotes international units per deciliter, BMI body mass index, HIV human immunodeficiency virus, HCV hepatitis C virus, NA not applicable.

*Other included Australia, New Zealand, Brazil, Hong Kong, India, Japan, Russia, and South Africa.

†Calculation was based on available data.

[0225] Subjects received rFVIIIFc for a median (min-max) of 30.5 (<1-54) weeks with a median (minimum to maximum) in arms 1, 2, and 3 of 32.1 (9, 54), 28.0 (<1, 38), and 28.9 (15, 32) weeks, respectively. In total, 111 subjects (67.7%) had >50 EDs to study drug. A total of 9356 injections were administered during the study, corresponding to 9170 EDs (100.2 patient-years of exposure). Overall, 93.6% of patients were compliant with both the prescribed dose and interval in the prophylaxis arms.

[0226] Overall, 93.6% of subjects were adherent with both the prescribed dose and the prescribed dosing interval in the prophylaxis treatment arms.

Efficacy

[0227] In total, 163 subjects were included in efficacy analyses. One subject was excluded due to only receiving ADVATE® (Arm 1); another subject withdrew before efficacy assessment (Arm 2). Median ABR with the 25th and 75th percentiles (interquartile range [IQR]) were as follows: Arm1 - individualized prophylaxis arm: 1.6 (0.0, 4.7); Arm 2 - weekly prophylaxis arm: 3.6 (1.9, 8.4); and Arm 3 - episodic treatment arm: 33.6 (21.1, 48.7). Most bleeding episodes were spontaneous. In Arms 1, 2, and 3, respectively, 45.3%, 17.4%, and 0% had no bleeding episodes and 13.7%, 34.8%, and 0% had 1 bleeding episode.

[0228] Median dosing interval:

- In the individualized prophylaxis arm, the median dosing interval was 3.5 days, during the last 3 months on study.
- 30 percent of patients in the individualized prophylaxis arm achieved a mean dosing interval of at least 5 days.

- c. Overall, the median (IQR) dosing interval with individualised prophylaxis aimed to achieve a FVIII trough of at least 1 to 3 IU/dL (with the maximum 5-day dosing interval permitted in the protocol) was 3.5 (3.2–4.4) days (Table 7) based on the median (IQR) weekly dose of 78 (72–91) IU/kg.
- d. Approximately 30% of subjects achieved a mean dosing interval of 5 days over the last 3 months on study. 100% of subjects had mean dosing intervals ≥ 2 days throughout the study.

[0229] The median dose per injection for Arm 3 episodic regimen was 26.5 IU/kg (n=23) and the median total dose per bleeding episode was 27.4 IU/kg (n=23).

TABLE 7. Comparative pharmacokinetics for rFVIIIFc vs. rFVIII (n=28)¹

PK parameter	Geometric mean for rFVIIIFc PK (95% CI)	Geometric mean for rFVIII PK (95% CI)	Geometric mean of intra-subject ratio (95% CI) p-value
Elimination $t_{1/2}$ (h)	18.97 (17.03, 21.12)	12.43 (11.14, 13.86)	1.53 (1.36, 1.71) <0.001
CL (mL/h/kg)	1.95 (1.71, 2.22)	3.041 (2.71, 3.41)	0.64 (0.60, 0.69) <0.001
Time to 1 IU/dL (days) (50 IU/kg dose)	4.92 (4.43, 5.46)	3.30 (2.99, 3.65)	1.49 (1.41, 1.57) <0.001

¹Arm 1 sequential PK group, compartmental model, one-stage assay. Estimates and 95% CIs for geometric means and geometric mean ratios; statistical significance assessed at the 2-sided 0.05 level; PK, pharmacokinetics; CI, confidence interval; C_{max}, maximal concentration; AUC, area-under-the-curve; $t_{1/2}$, half-life; CL, clearance; MRT, mean residence time; V_{ss}, volume of distribution at steady state.

[0230] Control of bleeding: approximately 98% (97.7%) of bleeding episodes were controlled by one or two injections of rFVIIIFc. 87.3% of bleeds were controlled with one injection. 1.7% required three injections.

[0231] Perioperative management: Overall, 9 major surgeries were performed in 9 subjects (8 subjects from Arm 1; 1 subject from Arm 2), including knee arthroplasty (n=5), laparoscopic inguinal hernia repair (n=2), appendectomy (n=1), and arthroscopy (n=1). Treating physicians rated the hemostatic efficacy of rFVIIIFc as excellent (8/9) or

good (1/9) in 100% of surgeries. Median (min, max) estimated blood loss available for 7/9 surgeries was 15.0 (0, 600) mL during surgery and 0.0 (0, 1100) mL post-operatively; post-surgical drainage).

PK

[0232] Comparative PK data for rFVIII-Fc and ADVATE® available from 28 subjects in the Arm 1 sequential PK subgroup are summarised in Table 7.

- a. rFVIII-Fc demonstrated an approximate 50% longer elimination half-life (Figure 2) and mean residence time compared with ADVATE® ($P < 0.001$) due to a 35% reduction in rFVIII-Fc clearance with respect to ADVATE®.
- b. Geometric mean time to 1 IU/dL FVIII activity following a 50-IU/kg dose was approximately 5 days for rFVIII-Fc versus approximately 3 days for ADVATE® ($P < 0.001$; Table 7, Figure 2).
- c. Incremental recovery for rFVIII-Fc was clinically comparable with ADVATE®.
- d. There was no shift in the PK properties of rFVIII-Fc as evidenced by comparable PK properties between the baseline and repeat rFVIII-Fc PK profiles.
- e. Analogous PK results were obtained when the analysis was based on the chromogenic FVIII assay.
- f. The one-stage and chromogenic clotting assays accurately and precisely measured both rFVIII-Fc and ADVATE® utilising commercially available plasma FVIII standards.

[0233] The geometric mean (95% confidence interval) terminal half-life of rFVIII-Fc was approximately 19.0 hours (17.0, 21.1) hours, which is 1.53-fold longer than that of ADVATE® (approximately 12.4 (11.1, 13.9) hours).

[0234] Geometric mean time (IQR) to 1% FVIII activity following 50IU/kg of rFVIII-Fc was approximately 5 days (4.92 (4.43, 5.46) days). Comparable PK profiles of rFVIII-Fc were observed at Week 14. The median dosing interval with individualized prophylaxis was 3.5 days and the median dose per week was 78 IU/kg; approximately 30% of subjects achieved a mean dosing interval of ≥ 5 days over the last 3 months on study (subjects

with ≥ 6 months on study) as shown in Table 8. 98% of bleeding episodes were controlled with 1-2 injections.

TABLE 8. Prophylactic dosing summary

	n	Arm 1 Individualised prophylaxis	n	Arm 2 Weekly prophylaxis
Median dose, IU/kg/week (IQR)				
Overall	117	77.9 (72.3, 91.2)	23	65.6 (64.2, 68.2)
Last 3 months ¹	112	77.7 (71.9, 106.2)	16	65.5 ² (64.3, 67.3)
Median dosing interval, days (IQR)				
Overall	117	3.5 (3.2, 4.4)		
Last 3 months ¹	112	3.5 ³ (3.0, 5.0)		

¹Based on last 3 months on study for subjects on study ≥ 6 months

²Fixed dose of 65 IU/kg/wk.

³30% of subjects achieved every 5-day pharmacokinetic-driven dosing to maintain troughs 1% to 3% above baseline.

IQR, interquartile range.

Safety

[0235] No inhibitors were detected to rFVIII Fc, and no cases of anaphylaxis, allergy, or serious thrombotic events were reported.

[0236] rFVIII Fc was generally well tolerated, and no serious adverse events were assessed to be related to rFVIII Fc.

[0237] The most common AEs, regardless of causality, (incidence $\geq 5\%$) occurring outside of the perioperative management period were nasopharyngitis, arthralgia (joint pain), headache, and upper respiratory tract infection, as summarized in Table 9.

[0238] 12 subjects (7.3%) experienced at least one serious AE (SAE) outside of the perioperative management period.

[0239] No SAEs were assessed to be related to drug by the investigator.

[0240] Insignificant incidence of non-neutralizing antibody (NNA) has been observed under NNA assay, which is approximately 20 times more sensitive than Bethesda assay. Five subjects were found positive for NNAs at baseline, but all had at least one negative evaluation during the study. Six subjects, four in Arm 1 study and two in Arm 2 study,

became positive during the study. In almost all cases, however, antibodies were transient and of low titer, and in all cases antibodies were directed against FVIII.

[0241] Overall, 14 (8.5%) subjects reported SAEs. No SAEs were determined by the investigator as related to study drug, and SAEs reported were not experienced in more than 1 subject. There was 1 death during the study, attributed to polysubstance overdose, and assessed as unrelated to rFVIIIIFc by the investigator.

TABLE 9. Summary of adverse events (AEs)

	Total N=164 n (%)
Any AE	108 (65.9)
Most common AEs ($\geq 5\%$ of subjects)	
Nasopharyngitis	20 (12.2)
Arthralgia	13 (7.9)
Headache	9 (5.5)
Upper respiratory infection	9 (5.5)

Summary

[0242] Individualized and weekly prophylactic regimens resulted in low single-digit median annualized bleeding rates

[0243] In the individualized prophylaxis arm, the median dosing interval was 3.5 days. During the last 3 months on study, 30 percent of patients in the individualized prophylaxis arm achieved a mean dosing interval of at least 5 days.

[0244] 98% of bleeding episodes were controlled by one or two injections of rFVIIIIFc. In total, 757 bleeding episodes (Arm 1=209; Arm 2=92; Arm 3=456) were treated in 106 of the 164 subjects. 87.3% required a single injection of rFVIIIIFc for resolution, and 97.8% required ≤ 2 injections. A total of 85.6%, 80.4% and 89.5% of bleeding episodes were resolved with 1 injection of rFVIIIIFc in Arms 1, 2 and 3, respectively.

[0245] Hemostatic efficacy of rFVIIIIFc during surgery was rated by treating physicians as excellent or good in 100% of surgeries.

[0246] The half-life of rFVIIIIFc was approximately 19.0 hours compared to 12.4 hours for ADVATE[®].

- [0247] No subject developed an inhibitor or experienced an anaphylactic reaction to rFVIII-Fc.
- [0248] Recombinant FVIII-Fc was generally well tolerated.
- [0249] A-LONG was the largest registrational global pivotal phase 3 study of long-lasting rFVIII in severe haemophilia A conducted to date. The study showed that rFVIII-Fc offers the potential for a markedly reduced injection frequency, decreased treatment burden, an improvement in clinical outcomes for prevention of bleeds in patients with severe haemophilia A.
- [0250] Goal of the study design: The study was designed to evaluate the efficacy and safety of rFVIII-Fc in the control and prevention of bleeding episodes, routine prophylaxis, and perioperative management in subjects with severe hemophilia A. In addition, the study was designed to assess the effective dose and interval of rFVIII-Fc for prophylaxis, as well as the feasibility of weekly treatment at a dose of 65 IU/Kg.
- [0251] Episodic treatment is the administration of replacement factor only as needed to treat bleeding episodes after they have started. Prophylactic treatment is the regular administration of replacement factor to prevent bleeding episodes.
- [0252] The starting regimen was 25 IU/kg on Day 1 and 50 IU/kg on Day 4. The dose and frequency of treatment could be adjusted based on the subject's PK profile and the goal was to maintain trough factor levels sufficient to prevent bleeding episodes using doses between 25 IU/kg and 65 IU/kg and treatment intervals of 3 to 5 days.
- [0253] More than half of severe hemophilia A patients in the US still do not follow a prophylaxis regimen, due in part to the treatment burden. We therefore included the weekly dosing regimen in the A-LONG study to evaluate whether this regimen would result in a lower ABR compared to an episodic regimen. The median ABR in the weekly regimen was 3.59 (IQR 1.9, 8.4) compared with 33.6 (IQR 21.1, 48.7) for episodic treatment.
- [0254] All subjects had an initial PK evaluation to characterize the PK of rFVIII-Fc in a representative population of patients with Hemophilia A.
- [0255] More extensive PK sampling was conducted in a subset of subjects in the individualized prophylaxis arm (Arm 1) at baseline after a single dose of ADVATE® (50 IU/kg), and after a 4 day washout, followed by a single dose of rFVIII-Fc (50 IU/kg). Blood samples were taken for ADVATE® over a period of 72 hours. Blood samples were

then taken for rFVIII Fc over a period of 120 hours. PK assessment of rFVIII Fc in this subset was repeated at 12 to 24 weeks with the same PK sampling schedule.

[0256] In peri-operative management, a single injection of rFVIII Fc was sufficient to maintain haemostasis to the end date/time of all major surgeries at a median dose of 51.4 IU/kg. Median rFVIII Fc consumption (summarized over all injections during each referenced time period) was 80.6 IU/kg on the day of major surgery, 161.3 IU/kg for Days 1-3 days following surgery, and 387.1 IU/kg for Days 4-14 following surgery. Perioperative haemostasis with rFVIII Fc was rated as excellent or good for all 9 major surgeries. No subjects reported a bleeding episode during the postoperative or rehabilitation periods. Overall, 7 adverse events (AEs) were reported in 4 (44.4%) subjects in the surgery subgroup, of which 6 AEs were of mild or moderate severity, and 1 AE was considered severe. Two serious AEs (inguinal hernia and appendicitis) were reported in 2 subjects. All AEs during the perioperative period were assessed by the investigators as unrelated to rFVIII Fc treatment.

[0257] Treating physicians rated the hemostatic efficacy of rFVIII Fc as excellent or good in 100% of surgeries on a 4 point scale including excellent, good, fair, and poor/none.

[0258] rFVIII Fc resulted in low median ABRs of 3.59 in the weekly prophylaxis arm and 1.60 in the individualized interval prophylaxis arm. In contrast, the episodic treatment arm had a median ABR of 33.57.

[0259] In the individualized prophylaxis arm, the median dosing interval was 3.5 days at a median dose of 77.7 IU/kg during the last 3 months on study.

[0260] The terminal half-life was approximately 18.97 hours for rFVIII Fc and approximately 12.3 hours for ADVATE®.

[0261] 97.8% of bleeding episodes were controlled by one or two injections.

[0262] Hemostatic efficacy of rFVIII Fc for perioperative management was rated by treating physicians as excellent or good in 100% of surgeries.

[0263] Arm 2 was a weekly dosing regimen. It was designed to investigate benefit for patients on prophylaxis therapy, and to compare those results with subjects using an episodic treatment regimen. The median ABR in Arm 2 was 3.59 compared with 33.57 for the episodic treatment regimen (Arm 3).

[0264] The median dose per injection required for resolution of bleeding was 27.4 IU/kg, and the median total dose required was 28.2 IU/kg.

[0265] The median number of injections required for resolution of a bleeding episode was consistently 1.0 when treatment was administered within 8 hours of bleed onset, regardless of type or location of bleed. Overall, 78.8% of rFVIII Fc injections in Arm 1, 64.8% in Arm 2, and 79.7% of injections in Arm 3, were rated by subjects as producing excellent or good response. The median dosing interval of 3.5 days from Arm 1 represents the dosing interval that can be achieved by the majority of patients: 30% of patients were able to achieve a dosing interval of at least 5 days. Some patients were able to achieve a weekly prophylaxis regimen (Arm 2) with a median ABR of 3.59 bleeding episodes.

[0266] Subjects in the individualized prophylaxis arm received an initial dose of 25 IU/kg on Day 1 and 50 IU/kg on Day 4. The study design allowed for adjustment of the dosing interval and dose to maintain targeted trough factor levels and to prevent bleeding. We believe the median dosing interval and dose based on the last 6 months on study is most representative of the individualized prophylaxis regimen.

[0267] 87.3% of bleeding episodes were resolved with one injection of rFVIII Fc and 97.8% were resolved with one or two injections. For the 96 (12.7%) bleeding episodes that required more than 1 injection for resolution, the median interval between the first and second injection was 30.9 hours.

[0268] The investigators' global assessment of subject response to their assigned rFVIII Fc regimen was rated as excellent or effective for 99.4%, 100%, and 98.1% of the subject visits in Arms 1, 2, and 3, respectively.

[0269] In A-LONG, the clotting factor activity was measured using a one-stage (aPTT) clotting assay and a chromogenic assay. The reported half-lives are based on the results of the one-stage clotting assay. While the half-lives of rFVIII Fc and ADVATE® were both slightly longer with the chromogenic assay, the ratios between products were consistent between the two assays.

EMBODIMENTS

[0270] E1. A method of reducing the annualized bleeding rate in a subject having hemophilia comprising administering to the subject an effective dose of a long-acting FVIII polypeptide at a dosing interval of about three days or longer.

[0271] E2. The method of embodiment E1, wherein the administration is prophylactic and individualized for the subject resulting in an annualized bleeding rate less than about

5.0, less than about 4.9, less than about 4.8, less than about 4.7, less than about 4.6, or less than about 4.5.

- [0272] E3. The method of embodiment E2, wherein the median annualized bleeding rate is about 1.6.
- [0273] E4. The method of embodiment E1, wherein the administration is prophylactic and weekly resulting in an annualized bleeding rate less than about 9.0, less than about 8.9, less than about 8.8, less than about 8.7, less than about 8.6, less than about 8.5, or less than about 8.4.
- [0274] E5. The method of embodiment E4, wherein the median annualized bleeding rate is about 3.6.
- [0275] E6. The method of embodiment E1, wherein the administration is on-demand or episodic resulting in an annualized bleeding rate less than about 55, less than about 54, less than about 53, less than about 52, less than about 51, less than about 50, less than about 49, less than about 48, or less than about 47.
- [0276] E7. The method of embodiment E6, wherein the median annualized bleeding rate is about 33.6.
- [0277] E8. The method of any one of embodiments E1 to E7, wherein the effective dose is between about 20 IU/kg to about 90 IU/kg.
- [0278] E9. The method of any one of embodiments E1 to E8, wherein the effective dose is 20-30 IU/kg, 30-40 IU/kg, 40-50 IU/kg, 50-60 IU/kg, 60-70 IU/kg, 70-80 IU/kg, or 80-90 IU/kg.
- [0279] E10. The method of any one of embodiments E1 to E9, wherein the effective dose is 20 IU/kg, 25 IU/kg, 30 IU/kg, 35 IU/kg, 40 IU/kg, 45 IU/kg, 50 IU/kg, 55 IU/kg, 60 IU/kg, 65 IU/kg, 70 IU/kg, 75 IU/kg, 80 IU/kg, 85 IU/kg, or 90 IU/kg.
- [0280] E11. The method of any of embodiments E8 to E10, wherein the administration is prophylactic and individualized at an effective dose of about 25 IU/kg to about 65 IU/kg twice weekly or every three days or about 50 IU/kg to about 65 IU/kg every 4 or 5 days.
- [0281] E12. The method of embodiment E11, wherein a first dose of the long-acting FVIII polypeptide is administered is about 25 IU/kg and a second dose of the long-acting FVIII polypeptide is about 50 IU/kg.
- [0282] E13. The method of any of embodiments E8 to E10, wherein the administration is prophylactic and weekly at an effective dose of 65 IU/kg weekly.

- [0283] E14. The method of any of embodiments E8 to E10, wherein the administration is on-demand or episodic at an effective dose of 10 IU/kg to 75 IU/kg every 12 to 24 hours.
- [0284] E15. The method of any one of embodiments E1 to E7, wherein the effective dose is a fixed dose, which is standard across all body weights.
- [0285] E16. The method of embodiment E15, wherein the fixed dose is about 2,000 IU, about 2,500 IU, about 3,000 IU, about 3,500 IU, or about 4,000 IU per dose.
- [0286] E17. The method of any one of embodiments E1 to E7, wherein the effective dose is a stratified dose.
- [0287] E18. The method of any one of embodiments E1 to E17, wherein a trough level of the long-acting FVIII polypeptide is above 1%, above 2%, or above 3% of normal.
- [0288] E19. The method of any one of embodiments E1 to E18, wherein the long-acting FVIII polypeptide has a $T_{1/2\beta}$ (activity) of about 7 hours to about 48 hours, about 6 hours to about 49 hours, about 5 hours to about 50 hours.
- [0289] E20. The method of embodiment E19, wherein the long-acting FVIII has a $T_{1/2\beta}$ (activity) mean of at least about 15 hours, at least about 16 hours, at least about 17 hours, at least about 18 hours, at least about 19 hours, at least about 20 hours, at least about 21 hours, at least about 22 hours, at least about 23 hours, at least about 24 hours, at least about 25 hours.
- [0290] E21. The method of embodiment E20, wherein the $T_{1/2\beta}$ (activity) mean is about 19 hours.
- [0291] E22. The method of embodiment E19 or E20, wherein the $T_{1/2\beta}$ (activity) mean is at least about 1.5 fold higher than a polypeptide consisting of amino acids 20 to 1457 of SEQ ID NO:2, 20 to 2351 of SEQ ID NO: 6, or ADVATE®.
- [0292] E23. The method of any one of embodiments E1 to E22, wherein the plasma trough level of the long-acting FVIII polypeptide is maintained between about 1% and about 5%, between about 1% and about 6%, between about 1% and about 7%, between about 1% and about 8%, between about 1% and about 9%, between about 1% and about 10%, between about 1% and about 11%, between about 1% and about 12%, between about 1% and about 13%, between about 1% and about 14%, between about 1% and about 15% above the baseline in the subject.

- [0293] E24. The method of any one of embodiments E1 to E23, wherein the long-acting FVIII polypeptide is a long-acting FVIII polypeptide comprising a FVIII polypeptide and a heterologous moiety.
- [0294] E25. The method of embodiment E24, wherein the heterologous moiety is an FcRn binding partner.
- [0295] E26. The method of embodiment E25, wherein the FcRn binding partner comprises an Fc region.
- [0296] E27. The method of embodiment E25 or E26, wherein the long-acting FVIII polypeptide further comprises a second FcRn binding partner.
- [0297] E28. The method of embodiment E27, wherein the second FcRn binding partner comprises a second Fc region.
- [0298] E29. The method of embodiment E27 or E28, wherein the FcRn binding partner and the second FcRn binding partner are associated.
- [0299] E30. The method of embodiment E29, wherein the association is a covalent bond.
- [0300] E31. The method of embodiment E30, wherein the covalent bond is a disulfide bond.
- [0301] E32. The method of any one of embodiments E27 to E31, wherein the second FcRn binding partner is not linked to an amino acid sequence by a peptide bond.
- [0302] E33. The method of any one of embodiments E1 to E32, wherein the long-acting FVIII polypeptide is FVIII monomer dimer hybrid.
- [0303] E34. The method of any one of embodiments E24 to E33, wherein the FVIII polypeptide in the long-acting polypeptide is a human FVIII.
- [0304] E35. The method of any one of embodiments E24 to E34, wherein the FVIII polypeptide in the long-acting polypeptide is a full-length FVIII or a B-domain deleted FVIII.
- [0305] E36. The method of any one of embodiments E24 to E34, wherein the FVIII polypeptide is single chain.
- [0306] E37. The method of any one of embodiments E24 to E34, wherein the FVIII polypeptide is unprocessed.
- [0307] E38. The method of any one of embodiments E24 to E34, wherein the FVIII polypeptide is in two chains, a first chain comprising a heavy chain of the FVIII polypeptide and a second chain comprising a light chain of the FVIII polypeptide.

- [0308] E39. The method of any one of embodiments E25 to E38, wherein the FcRn binding partner in the chimeric polypeptide is a human Fc.
- [0309] E40. The method of any one of embodiments E24 to E39, wherein the FVIII polypeptide is at least 60%, 70%, 80%, 90%, 95%, 95%, 97%, 98%, 99%, or 100% identical to a FVIII amino acid sequence shown in Table 11A or 11B without a signal sequence (amino acids 20 to 1457 of SEQ ID NO: 2 or amino acids 20 to 2351 of SEQ ID NO: 6).
- [0310] E41. The method of any one of embodiments E25 to E40, wherein the FcRn binding partner is at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to a Fc amino acid sequence shown in Table 11B without a signal sequence (amino acids 21 to 247 SEQ ID NO:4).
- [0311] E42. The method of any one of embodiments E27 to E41, wherein the second FcRn binding partner in the chimeric polypeptide is a human Fc.
- [0312] E43 The method of embodiment E42, wherein the second FcRn binding partner is at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to a Fc amino acid sequence shown in Table 11B without a signal sequence (amino acids 21 to 247 SEQ ID NO:4)
- [0313] E44. The method of any one of embodiments E1 to E43, wherein the fixed dose of the clotting factor is for perioperative management of a bleeding episode.
- [0314] E45. The method of any one of embodiments E1 to E44, wherein the subject is in need of controlling or preventing bleeding or bleeding episodes.
- [0315] E46. The method of embodiment E45, wherein the subject is in need of controlling or preventing bleeding in minor hemorrhage, hemarthroses, superficial muscle hemorrhage, soft tissue hemorrhage, moderate hemorrhage, intramuscle or soft tissue hemorrhage with dissection, mucous membrane hemorrhage, hematuria, major hemorrhage, hemorrhage of the pharynx, hemorrhage of the retropharynx, hemorrhage of the retroperitonium, hemorrhage of the central nervous system, bruises, cuts, scrapes, joint hemorrhage, nose bleed, mouth bleed, gum bleed, intracranial bleeding, intraperitoneal bleeding, minor spontaneous hemorrhage, bleeding after major trauma, moderate skin bruising, or spontaneous hemorrhage into joints, muscles, internal organs or the brain.
- [0316] E47. The method of embodiment E46, wherein the subject is in need of management of bleeding associated with surgery or dental extraction.

- [0317] E48. The method of embodiment E47, wherein the subject will undergo, is undergoing, or has undergone major surgery.
- [0318] E49. The method of embodiment E48, wherein the major surgery is orthopedic surgery, extensive oral surgery, urologic surgery, or hernia surgery.
- [0319] E50. The method of embodiment E49, wherein the orthopedic surgery is replacement of knee, hip, or other major joint.
- [0320] E51. The method of any one of embodiments E1 to E50, wherein the subject is in need of long-term treatment.
- [0321] E52. The method of any one of embodiments E1 to E51, wherein the long-acting FVIII polypeptide is administered intravenously or subcutaneously.
- [0322] E53. The method of any one of embodiments E1 to E52, wherein the long-acting FVIII polypeptide is formulated in a pharmaceutical composition comprising:
- (a) the long-acting FVIII polypeptide;
 - (b) one or more stabilizing agents selected from the group consisting of sucrose, trehalose, raffinose, arginine, and mixture thereof;
 - (c) sodium chloride (NaCl);
 - (d) L-histidine;
 - (e) calcium chloride; and
 - (f) polysorbate 20 or polysorbate 80.
- [0323] E54. The method of embodiment E53, wherein the pharmaceutical composition does not include mannitol, glycine, alanine, or hydroxyethyl starch.
- [0324] E55. The method of embodiment E53 or E54, wherein the pharmaceutical composition includes NaCl as the only bulking agent.
- [0325] E56. The method of any one of embodiments E53 to E55, wherein the pharmaceutical composition comprises sucrose.
- [0326] E57. The method of embodiment E56, wherein the pharmaceutical composition comprises about 1% (w/v) to about 2.5% (w/v) sucrose.
- [0327] E58. The method of embodiment E57, wherein the pharmaceutical composition comprises about 1.3% (w/v) sucrose to about 2.0% (w/v) sucrose.

- [0328] E59. The method of embodiment E58, wherein the pharmaceutical composition comprises about 1.33% (w/v) sucrose or about 2.0% (w/v) sucrose.
- [0329] E60. The method of embodiment E56, wherein the pharmaceutical composition comprises about 10 mg/ml to about 25 mg/ml sucrose.
- [0330] E61. The method of embodiment E60, wherein the pharmaceutical composition comprises about 13 mg/ml to about 20 mg/ml sucrose.
- [0331] E62. The method of embodiment E61, wherein the pharmaceutical composition comprises about 13.3 mg/ml sucrose or about 20.0 mg/ml sucrose.
- [0332] E63. The method of any one of embodiments E53 to E62, wherein the pharmaceutical composition comprises about 150 mM to about 250 mM NaCl.
- [0333] E64. The method of embodiment E63, wherein the pharmaceutical composition comprises about 175 mM to about 225 mM NaCl.
- [0334] E65. The method of embodiment E64, wherein the pharmaceutical composition comprises about 200 mM to about 210 mM NaCl.
- [0335] E66. The method of embodiment E65, wherein the pharmaceutical composition comprises about 205 mM NaCl.
- [0336] E67. The method of any one of embodiments E53 to E62, wherein the pharmaceutical composition comprises about 8.8 mg/ml to about 14.6 mg/ml NaCl.
- [0337] E68. The method of embodiment E67, wherein the pharmaceutical composition comprises about 10 mg/ml to about 13 mg/ml NaCl.
- [0338] E69. The method of embodiment E68, wherein the pharmaceutical composition comprises about 12.0 mg/ml NaCl.
- [0339] E70. The method of any one of embodiments E53 to E62, wherein the pharmaceutical composition comprises about 250 mM to about 350 mM NaCl.
- [0340] E71. The method of embodiment E70, wherein the pharmaceutical composition comprises about 275 mM to about 325 mM NaCl.
- [0341] E72. The method of embodiment E71, wherein the pharmaceutical composition comprises about 308 mM NaCl.
- [0342] E73. The method of any one of embodiments E53 to E62, wherein the pharmaceutical composition comprises about 14.6 mg/ml to about 20.5 mg/ml NaCl.
- [0343] E74. The method of embodiment E73, wherein the pharmaceutical composition comprises about 16 mg/ml to about 19 mg/ml NaCl.

- [0344] E75. The method of embodiment E74, wherein the pharmaceutical composition comprises about 18.0 mg/ml NaCl.
- [0345] E76. The method of any one of embodiments E53 to E75, wherein the pharmaceutical composition comprises about 5 mM to about 15 mM L-histidine.
- [0346] E77. The method of embodiment E76, wherein the pharmaceutical composition comprises about 6.64 mM L-histidine or about 9.8 mM L-histidine.
- [0347] E78. The method of any one of embodiments E53 to E75, wherein the pharmaceutical composition comprises about 0.75 mg/ml to about 2.25 mg/ml L-histidine.
- [0348] E79. The method of embodiment E78, the pharmaceutical composition comprises about 1.03 mg/ml L-histidine or about 1.55 mg/ml L-histidine.
- [0349] E80. The method of any one of embodiments E53 to E79, wherein the pharmaceutical composition comprises about 5 mM to about 10 mM calcium chloride.
- [0350] E81. The method of embodiment E80, wherein the pharmaceutical composition comprises about 5.4 mM calcium chloride or about 8 mM calcium chloride.
- [0351] E82. The method of any one of embodiments E53 to E79, wherein the pharmaceutical composition comprises about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate.
- [0352] E83. The method of embodiment E82, wherein the pharmaceutical composition comprises about 0.8 mg/ml calcium chloride dihydrate or about 1.18 mg/ml calcium chloride dihydrate.
- [0353] E84. The method of any one of embodiments E53 to E83, wherein the pharmaceutical composition comprises about 0.008% (w/v) to about 0.025% (w/v) polysorbate 20 or polysorbate 80.
- [0354] E85. The method of embodiment E84, wherein the pharmaceutical composition comprises about 0.013% (w/v) polysorbate 20 or polysorbate 80 or about 0.02 % (w/v) polysorbate 20 or polysorbate 80.
- [0355] E86. The method of any one of embodiments E53 to E83, wherein the pharmaceutical composition comprises about 0.08 mg/ml to about 0.25 mg/ml polysorbate 20 or polysorbate 80.
- [0356] E87. The method of embodiment E86, wherein the pharmaceutical composition comprises about 0.13% mg/ml polysorbate 20 or polysorbate 80 or about 0.20 mg/ml polysorbate 20 or polysorbate 80.

- [0357] E88. The method of any one of embodiments E53 to E87, wherein the rFVIII^{Fc} polypeptide comprises a first subunit comprising an amino acid sequence at least 90% or 95% identical to amino acids 20 to 1684 of SEQ ID NO:2 or 20 to 2578 of SEQ ID NO:6, and a second subunit comprising an amino acid sequence at least 90% to 95% identical to amino acids 21 to 247 of SEQ ID NO:4.
- [0358] E89. The method of embodiment E88, wherein the rFVIII^{Fc} polypeptide comprises a first subunit comprising amino acids 20 to 1684 of SEQ ID NO:2 or 20 to 2578 of SEQ ID NO:6, and a second subunit comprising amino acids 21 to 247 of SEQ ID NO:4.
- [0359] E90. The method of any one of embodiments E53 to E89, wherein the long-acting FVIII polypeptide is present at a concentration of about 50 IU/ml to about 2500 IU/ml.
- [0360] E91. The method of embodiment E90, wherein the pharmaceutical composition comprises 83 IU/ml, 167 IU/ml, 250 IU/ml, 333 IU/ml, 500 IU/ml, 667 IU/ml, 1000 IU/ml, 1333 IU/ml, 1667 IU/ml, or 2000 IU/ml of the long-acting FVIII polypeptide.
- [0361] E92. The method of any one of embodiments E53 to E89, wherein the long-acting FVIII polypeptide is present at a concentration of about 100 IU/ml to about 4000 IU/ml.
- [0362] E93. The method of embodiment E92, wherein the pharmaceutical composition comprises 150 IU/ml, 287.5 IU/ml, 431.25 IU/ml, 575 IU/ml, 862.5 IU/ml, 1150 IU/ml, 1725 IU/ml, 2300 IU/ml, 2875 IU/ml, or 3450 IU/ml of the long-acting FVIII polypeptide.
- [0363] E94. The method of any one of embodiments E53 to E56, wherein the pharmaceutical composition comprises:
- (a) about 50 IU/ml to about 2500 IU/ml of the long-acting FVIII polypeptide;
 - (b) about 1% (w/v) to about 2.5% (w/v) of sucrose;
 - (c) about 150 mM to about 250 mM NaCl;
 - (d) about 5 mM to about 15 mM L-histidine;
 - (e) about 5 mM to about 10 mM calcium chloride; and
 - (f) about 0.008% (w/v) to about 0.025% of polysorbate 20 or polysorbate 80.

[0364] E95. The method of embodiment E94, wherein the pharmaceutical composition comprises about 175 mM to about 225 mM NaCl.

[0365] E96. The method of embodiment E95, wherein the pharmaceutical composition comprises about 200 mM to about 210 mM NaCl.

[0366] E97. The method of embodiment E96, wherein the pharmaceutical composition comprises:

- (a) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, about 1000 IU/ml, about 1333 IU/ml, about 1667 IU/ml, or about 2000 IU/ml of the long-acting FVIII polypeptide;
- (b) about 1.33% (w/v) of sucrose;
- (c) about 205 mM NaCl;
- (d) about 6.64 mM L-histidine;
- (e) about 5.4 mM calcium chloride; and
- (f) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

[0367] E98. The method of any one of embodiments E53 to E56, wherein the pharmaceutical composition comprises:

- (a) about 100 IU/ml to about 4000 IU/ml of the long-acting FVIII polypeptide;
- (b) about 1% (w/v) to about 2.5% (w/v) of sucrose;
- (c) about 250 mM to about 350 mM NaCl;
- (d) about 5 mM to about 15 mM L-histidine;
- (e) about 5 mM to about 10 mM calcium chloride; and
- (f) about 0.008% (w/v) to about 0.025% of polysorbate 20 or polysorbate 80.

[0368] E99. The method of embodiment E98, wherein the pharmaceutical composition comprises about 275 mM to about 325 mM NaCl.

[0369] E100. The method of embodiment E99, wherein the pharmaceutical composition comprises:

- (a) about 150 IU/ml, about 287.5 IU/ml, about 431.25 IU/ml, about 575 IU/ml, about 862.5 IU/ml, about 1150 IU/ml, about 1725 IU/ml, about 2300 IU/ml, about 2875 IU/ml, or 3450 IU/ml of the long-acting FVIII polypeptide;
- (b) about 2.0% (w/v) of sucrose;
- (c) about 308 mM NaCl;
- (d) about 9.8 mM L-histidine;
- (e) about 8 mM calcium chloride; and
- (f) about 0.020% (w/v) of polysorbate 20 or polysorbate 80.

[0370] E101. The method of any one of embodiments E53 to E56, wherein the pharmaceutical composition comprises:

- (a) about 50 IU/ml to about 2500 IU/ml of the long-acting FVIII polypeptide;
- (b) about 10 mg/ml to about 25 mg/ml of sucrose;
- (c) about 8.8 mg/ml to about 14.6 mg/ml NaCl;
- (d) about 0.75 mg/ml to about 2.25 mg/ml L-histidine;
- (e) about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate; and
- (f) about 0.08 mg/ml to about 0.25 mg/ml of polysorbate 20 or polysorbate 80.

[0371] E102. The method of embodiment E101, wherein the pharmaceutical composition comprises about 10 mg/ml to 13 mg/ml NaCl.

[0372] E103. The method of embodiment E102, wherein the pharmaceutical composition comprises:

- (a) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, about 1000 IU/ml, about 1333 IU/ml, about 1667 IU/ml, or about 2000 IU/ml of the long-acting FVIII polypeptide;
- (b) about 13.3 mg/ml of sucrose;
- (c) about 12.0 mg/ml NaCl;

- (d) about 1.03 mg/ml L-histidine;
- (e) about 0.8 mg/ml calcium chloride dihydrate; and
- (f) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

[0373] E104. The method of any one of embodiments E53 to E56, wherein the pharmaceutical composition comprises:

- (a) about 100 IU/ml to about 4000 IU/ml of the long-acting FVIII polypeptide;
- (b) about 10 mg/ml to about 25 mg/ml of sucrose;
- (c) about 14.6 mg/ml to about 20.5 mg/ml NaCl;
- (d) about 0.75 mg/ml to about 2.25 mg/ml L-histidine;
- (e) about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate; and
- (f) about 0.08 mg/ml to about 0.25 mg/ml of polysorbate 20 or polysorbate 80.

[0374] E105. The method of embodiment E104, wherein the pharmaceutical composition comprises about 16 mg/ml to about 19 mg/ml NaCl.

[0375] E106. The method of embodiment E105, wherein the pharmaceutical composition comprises:

- (a) about 150 IU/ml, about 287.5 IU/ml, about 431.25 IU/ml, about 575 IU/ml, about 862.5 IU/ml, about 1150 IU/ml, about 1725 IU/ml, about 2300 IU/ml, about 2875 IU/ml, or 3450 IU/ml of the long-acting FVIII polypeptide;
- (b) about 20.0 mg/ml of sucrose;
- (c) about 18.0 mg/ml NaCl;
- (d) about 1.55 mg/ml L-histidine;
- (e) about 1.18 mg/ml calcium chloride dihydrate; and
- (f) about 0.20 mg/ml of polysorbate 20 or polysorbate 80.

[0376] E107. The method of any one of embodiments E1 to E52, wherein the long-acting FVIII polypeptide is packaged in a pharmaceutical kit comprising:

- (a) a first container comprising a lyophilized powder, where the powder comprises
 - (i) the long-acting FVIII polypeptide,
 - (ii) one or more stabilizing agents selected from the group consisting of sucrose, trehalose, raffinose, arginine, and mixture thereof;
 - (iii) sodium chloride (NaCl);
 - (iv) L-histidine;
 - (v) calcium chloride; and
 - (vi) polysorbate 20 or polysorbate 80; and
- (b) a second container comprising sterilized water for injections to be combined with the lyophilized powder of the first container.

[0377] E108. The method of embodiment E107, wherein the pharmaceutical kit does not include mannitol, glycine, alanine, or hydroxyethyl starch.

[0378] E109. The method of embodiment E107 or E108, wherein the pharmaceutical kit includes NaCl as the only bulking agent.

[0379] E110. The method of any one of embodiments E107 to E109, wherein the pharmaceutical kit comprises:

- (a) a first container comprising a lyophilized powder, where the powder comprises
 - (i) about 250 IU, about 500 IU, about 750 IU, about 1000 IU, about 1500 IU, about 2000 IU, about 3000 IU, about 4000 IU, about 5000 IU, or about 6000 IU of the long-acting FVIII polypeptide,
 - (ii) about 40 mg of sucrose;
 - (iii) about 36 mg of sodium chloride;
 - (iv) about 3.1 mg of L-histidine;

- (v) about 2.40 mg of calcium chloride dihydrate; and
- (v) about 0.40 mg of polysorbate 20 or polysorbate 80; and
- (b) a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising:
 - (i) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, about 1000 IU/ml, about 1333 IU/ml, or about 1667 IU/ml, or about 2000 IU/ml of the long-acting FVIII polypeptide, respectively;
 - (ii) about 1.33% (w/v) of sucrose;
 - (iii) about 205 mM NaCl;
 - (iv) about 6.64 mM L-histidine;
 - (v) about 5.4 mM of calcium chloride; and
 - (vi) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

[0380] E111. The method of any one of embodiments E107 to E109, wherein the pharmaceutical composition comprises:

- (a) a first container comprising a lyophilized powder, where the powder comprises
 - (i) about 250 IU, about 500 IU, about 750 IU, about 1000 IU, about 1500 IU, about 2000 IU, about 3000 IU, about 4000 IU, about 5000 IU, or about 6000 IU of the long-acting FVIII polypeptide,
 - (ii) about 40 mg of sucrose;
 - (iii) about 36 mg of sodium chloride;
 - (iv) about 3.1 mg of L-histidine;
 - (v) about 2.40 mg of calcium chloride dihydrate; and

- (v) about 0.40 mg of polysorbate 20 or polysorbate 80; and
 - (b) a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising:
 - (i) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, about 1000 IU/ml, about 1333 IU/ml, about 1667 IU/ml, or about 2000 IU/ml of the long-acting FVIII polypeptide, respectively;
 - (ii) about 13.3 mg/ml of sucrose;
 - (iii) about 12.0 mg/ml of NaCl;
 - (iv) about 1.03 mg/ml of L-histidine;
 - (v) about 0.80 mg/ml of calcium chloride; and
 - (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.
- [0381]** E112. The method of any one of embodiments E107 to E111, wherein the first container is a glass vial comprising a rubber stopper.
- [0382]** E113. The method of any one of embodiments E107 to E112, wherein the second container is a syringe body, and wherein the syringe body is associated with a plunger.
- [0383]** E114. The method of embodiment E113, the pharmaceutical kit further comprises an adaptor to connect the glass vial to the syringe body.
- [0384]** E115. The method of embodiment E113 or embodiment E114, wherein the pharmaceutical kit further comprises infusion tubing associated with a needle to be connected to the syringe, suitable for intravenous infusion.
- [0385]** E116. The method of any one of embodiments E70-E75, E92, E93, E98-E100, and E104-E106, wherein the pharmaceutical composition is a pre-lyophilization composition.

TABLE 10: Polynucleotide Sequences**A. B-Domain Deleted FVIII_hFc**

(i) B-Domain Deleted FVIII_hFc Chain DNA Sequence (FVIII signal peptide underlined, Fc region in bold) (SEQ ID NO:1, which encodes SEQ ID NO:2)

661				A	TGCAAATAGA	GCTCTCCACC	TGCTTCTTTC
721	TGTGCCTTTT	GGATTCTGC	TTTAGTGCCA	CCAGAAAGATA	CTACCTGGGT	GCAGTGGAAC	
781	TGTCATGGGA	CTATATGCAA	AGTGATCTCG	GTGAGCTGCC	TGTGGACGCA	AGATTTCCTC	
841	CTAGAGTGCC	AAAATCTTTT	CCATTCAACA	CCTCAGTCGT	GTACAAAAAG	ACTCTGTTTG	
901	TAGAATTAC	GGATCACCTT	TTCAACATCG	CTAAGCCAAG	GCCACCCTGG	ATGGGTCTGC	
961	TAGGTCTAC	CATCCAGGCT	GAGGTTTATG	ATACAGTGGT	CATTACACTT	AAGAACATGG	
1021	CTTCCCATCC	TGTCAGTCTT	CATGCTGTTG	GTGTATCCTA	CTGGAAAGCT	TCTGAGGGAG	
1081	CTGAATATGA	TGATCAGACC	AGTCAAAGGG	AGAAAGAAGA	TGATAAAGTC	TTCCCTGGTG	
1141	GAAGCCATAC	ATATGTCTGG	CAGGTCTTGA	AAGAGAATGG	TCCAATGGCC	TCTGACCCAC	
1201	TGTGCCTTAC	CTACTCATAT	CTTCTCATG	TGGACCTGGT	AAAAGACTTG	AATTCAGGCC	
1261	TCATTGGAGC	CCTACTAGTA	TGTAGAGAAG	GGAGTCTGGC	CAAGGAAAAG	ACACAGACCT	
1321	TGCACAAATT	TATACTACTT	TTTGCTGTAT	TTGATGAAGG	GAAAAGTTGG	CACTCAGAAA	
1381	CAAAGAACTC	CTTGATGCAG	GATAGGGATG	CTGCATCTGC	TCGGGCCTGG	CCTAAAATGC	
1441	ACACAGTCAA	TGTTTATGTA	AACAGGTCTC	TGCCAGGTCT	GATTGGATGC	CACAGGAAAT	
1501	CAGTCTATTG	GCATGTGATT	GGAATGGGCA	CCACTCCTGA	AGTGCACCTCA	ATATTCCTCG	
1561	AAGGTCACAC	ATTTCTTGTG	AGGAACCATC	GCCAGGCGTC	CTTGGAATC	TCGCCAATAA	
1621	CTTTCCTTAC	TGCTCAAACA	CTCTTGATGG	ACCTTGGAACA	GTTTCTACTG	TTTTGTCTATA	
1681	TCTCTTCCCA	CCAACATGAT	GGCATGGAAG	CTTATGTCAA	AGTAGACAGC	TGTCCAGAGG	
1741	AACCCCAACT	ACGAATGAAA	AATAATGAAG	AAGCGGAAGA	CTATGATGAT	GATCTTACTG	
1801	ATTCTGAAAT	GGATGTGGTC	AGGTTTGATG	ATGACAACTC	TCCTTCCTTT	ATCCAAATTC	
1861	GCTCAGTTGC	CAAGAAGCAT	CCTAAAACCT	GGGTACATTA	CATTGCTGCT	GAAGAGGAGG	
1921	ACTGGGACTA	TGCTCCCTTA	TCTCTCGCCC	CCGATGACAG	AAGTTATAAA	AGTCAATATT	
1981	TGAACAATGG	CCCTCAGCGG	ATTGGTAGGA	AGTACAAAAA	AGTCCGATTT	ATGGCATACA	
2041	CAGATGAAAC	CTTTAAGACT	CGTGAAGCTA	TTCAGCATGA	ATCAGGAATC	TTGGGACCTT	
2101	TACTTTATGG	GGAAGTTGGA	GACACACTGT	TGATTATATT	TAAGAATCAA	GCAAGCAGAC	
2161	CATATAACAT	CTACCCTCAC	GGAATCACTG	ATGTCCGTCC	TTTGTATTCA	AGGAGATTAC	
2221	CAAAGGTTGT	AAAACATTTG	AAGGATTTTC	CAATTCTGCC	AGGAGAAATA	TTCAAATATA	
2281	AATGGACAGT	GACTGTAGAA	GATGGGCCAA	CTAAATCAGA	TCCTCGGTGC	CTGACCCGCT	
2341	ATTACTCTAG	TTTCGTTAAT	ATGGAGAGAG	ATCTAGCTTC	AGGACTCATT	GGCCCTCTCC	
2401	TCATCTGCTA	CAAAGAATCT	GATAGTCAAA	GAGGAAACCA	GATAATGTCA	GACAAGAGGA	
2461	ATGTCATCCT	GTTTTCTGTA	TTTGATGAGA	ACCGAAGCTG	GTACCTCACA	GAGAATATAC	
2521	AACGCTTTCT	CCCCAATCCA	GCTGGAGTGC	AGCTTGAGGA	TCCAGAGTTC	CAAGCCTCCA	
2581	ACATCATGCA	CAGCATCAAT	GGCTATGTTT	TTGATAGTTT	GCAGTTGTCA	GTTTGTGTTGC	
2641	ATGAGGTGGC	ATACTGGTAC	ATTCTAAGCA	TTGGAGCACA	GACTGACTTC	CTTTCTGTCT	
2701	TCTTCTCTGG	ATATACCTTC	AAACACAAAA	TGGTCTATGA	AGACACACTC	ACCTTATTCC	
2761	CATTCTCAGG	AGAAACTGTC	TTTATGTCGA	TGGAAAACCC	AGGTCTATGG	ATTCTGGGGT	
2821	GCCACAACCT	AGACTTTCGG	AACGAGGCA	TGACCGCCTT	ACTGAAGGTT	TCTAGTTGTG	
2881	ACAAGAACAC	TGGTGATTAT	TACGAGGACA	GTTATGAAGA	TATTTTCAGCA	TACTTGCTGA	
2941	GTAAAAACAA	TGCCATTGAA	CCAAGAAGCT	TCTCTCAAAA	CCCACCAGTC	TTGAAACGCC	
3001	ATCAACGGGA	AATAACTCGT	ACTACTCTTC	AGTCAGATCA	AGAGGAAATT	GACTATGATG	
3061	ATACCATATC	AGTTGAAATG	AAGAAGGAAG	ATTTTGACAT	TTATGATGAG	GATGAAAATC	
3121	AGAGCCCCCG	CAGCTTTCAA	AAGAAAAACAC	GACACTATTT	TATTGCTGCA	GTGGAGAGGC	
3181	TCTGGGATTA	TGGGATGAGT	AGCTCCCCAC	ATGTTCTAAG	AAACAGGGCT	CAGAGTGGCA	
3241	GTGTCCCTCA	GTTCAAGAAA	GTTGTTTTC	AGGAATTTAC	TGATGGCTCC	TTTACTCAGC	
3301	CCTTATACCG	TGGAGAACTA	AATGAACATT	TGGGACTCCT	GGGGCCATAT	ATAAGAGCAG	
3361	AAGTTGAAGA	TAATATCATG	GTAACCTTCA	GAAATCAGGC	CTCTCGTCCC	TATTCCTTCT	
3421	ATTCTAGCCT	TATTTCTTAT	GAGGAAGATC	AGAGGCAAGG	AGCAGAACCT	AGAAAAAACT	
3481	TTGTCAAGCC	TAAAGAAACC	AAAACCTACT	TTTGAAAAGT	GCAACATCAT	ATGGCACCCA	

```

3541 CTAAAGATGA GTTTGA CTGACTGC AAAGCCTGGG CTTATTTCTC TGATGTTGAC CTGGAAAAAG
3601 ATGTGCACTC AGGCCTGATT GGACCCCTTC TGGTCTGCCA CACTAACACA CTGAACCCTG
3661 CTCATGGGAG ACAAGTGACA GTACAGGAAT TTGCTCTGTT TTTCACCATC TTTGATGAGA
3721 CCAAAAGCTG GTACTTCACT GAAAATATGG AAAGAACTG CAGGGCTCCC TGCAATAACC
3781 AGATGGAAGA TCCCCTTTT AAAGAGAATT ATCGCTTCCA TGCAATCAAT GGCTACATAA
3841 TGGATACACT ACCTGGCTTA GTAATGGCTC AGGATCAAAG GATTCGATGG TATCTGCTCA
3901 GCATGGGCAG CAATGAAAAC ATCCATTCTA TTCATTTTCTG TGGACATGTG TTTACTGTAC
3961 GAAAAAAGA GGAGTATAAA ATGGCACTGT ACAATCTCTA TCCAGGTGTT TTTGAGACAG
4021 TGGAAATGTT ACCATCCAAA GCTGGAATTT GGCGGGTGGG ATGCCTTATT GGCGAGCATC
4081 TACATGCTGG GATGAGCACA CTTTTTCTGG TGTACAGCAA TAAGTGTCTG ACTCCCTGG
4141 GAATGGCTTC TGGACACATT AGAGATTTTC AGATTACAGC TTCAGGACAA TATGGACAGT
4201 GGGCCCCAAA GCTGGCCAGA CTTCAATTAT CCGGATCAAT CAATGCCTGG AGCACCAAGG
4261 AGCCCTTTTC TTGGATCAAG GTGGATCTGT TGGCACCAAT GATTATTCAC GGCATCAAGA
4321 CCCAGGGTGC CCGTCAGAAG TTCTCCAGCC TCTACATCTC TCAGTTTATC ATCATGTATA
4381 GTCTTGATGG GAAGAAGTGG CAGACTTATC GAGGAAATTC CACTGGAACC TTAATGGTCT
4441 TCTTTGGCAA TGTGGATTCA TCTGGGATAA AACACAATAT TTTTAACCCCT CCAATTAATTG
4501 CTCGATACAT CCGTTTGAC CCAACTCATT ATAGCATTCG CAGCACTCTT CGCATGGAGT
4561 TGATGGGCTG TGATTTAAAT AGTTGCAGCA TGCCATTGGG AATGGAGAGT AAAGCAATAT
4621 CAGATGCACA GATTACTGCT TCATCCTACT TTACCAATAT GTTTGCCACC TGGTCTCCTT
4681 CAAAAGCTCG ACTTCACCTC CAAGGGAGGA GTAATGCCCG GAGACCTCAG GTGAATAATC
4741 CAAAAGAGTG GCTGCAGTG GACTTCCAGA AGACAATGAA AGTCACAGGA GTAACCTACTC
4801 AGGGAGTAAA ATCTCTGCTT ACCAGCATGT ATGTGAAGGA GTTCCTCATC TCCAGCAGTC
4861 AAGATGGCCA TCAGTGGACT CTCTTTTTTC AGAATGGCAA AGTAAAGGTT TTTGAGGAA
4921 ATCAAGACTC CTTACACCT GTGGTGAAC CTCTAGACCC ACCGTTACTG ACTCGCTACC
4981 TTCGAATTCA CCCCAGAGT TGGGTGCACC AGATTGCCCT GAGGATGGAG GTTCTGGGCT
5041 GCGAGGCACA GGACCTCTAC GACAAAACCT ACACATGCCC ACCGTGCCCC GCTCCAGAAC
5101 TCCTGGGCGG ACCGTCAGTC TTCTCTTCC CCCCAAAACC CAAGGACACC CTCATGATCT
5161 CCCGGACCCC TGAGGTCACA TGCCTGGTGG TGGACGTGAG CCACGAAGAC CCTGAGGTCA
5221 AGTTCAACTG GTACGTGGAC GGCGTGGAGG TGCATAATGC CAAGACAAAG CCGCGGGAGG
5281 AGCAGTACAA CAGCACGTAC CGTGTGGTCA GCGTCCTCAC CGTCTGAC CAGGACTGGC
5341 TGAATGGCAA GGAGTACAAG TGCAAGGTCT CCAACAAAGC CCTCCCAGCC CCCATCGAGA
5401 AAACCATCTC CAAAGCCAAA GGGCAGCCCC GAGAACCACA GGTGTACACC CTGCCCCCAT
5461 CCCGGGATGA GCTGACCAAG AACCAGGTCA GCCTGACCTG CCTGGTCAAA GGCTTCTATC
5521 CCAGCGACAT CGCCGTGGAG TGGGAGAGCA ATGGGCAGCC GGAGAACAAC TACAAGACCA
5581 CGCCTCCCGT GTTGGACTCC GACGGCTCCT TCTTCCCTTA CAGCAAGCTC ACCGTGGACA
5641 AGAGCAGGTG GCAGCAGGGG AACGTCTTCT CATGCTCCGT GATGCATGAG GCTCTGCACA
5701 ACCACTACAC GCAGAAGAGC CTCTCCCTGT CTCCGGGTAA A

```

(ii) Fc DNA sequence (mouse Igk signal peptide underlined) (SEQ ID NO:3, which encodes SEQ ID NO:4)

```

7981 ATGGA GACAGACACA
8041 CTCCTGCTAT GGGTACTGCT GCTCTGGGTT CCAGGTTCCA CTGGTGACAA AACTCACACA
8101 TGCCACCCGT GCCCAGCACC TGAAGTCCCTG GGAGGACCGT CAGTCTTCCT CTTCCCCCA
8161 AAACCCAAGG ACACCCTCAT GATCTCCCGG ACCCTGAGG TCACATGCGT GGTGGTGGAC
8221 GTGAGCCACG AAGACCCTGA GGTCAAGTTC AACTGGTACG TGGACGGCGT GGAGGTGCAT
8281 AATGCCAAGA CAAAGCCGCG GGAGGAGCAG TACAACAGCA CGTACCGTGT GGTCAGCGTC
8341 CTCACCGTCC TGCACCAGGA CTGGCTGAAT GGCAAGGAGT ACAAGTGCAA GGTCTCCAAC
8401 AAAGCCCCCTC CAGCCCCCAT CGAGAAAACC ATCTCCAAAG CCAAAGGGCA GCCCCGAGAA
8461 CCACAGGTGT ACACCCTGCC CCCATCCCGC GATGAGCTGA CCAAGAACCA GGTCAGCCTG
8521 ACCTGCCTGG TCAAAGGCTT CTATCCCAGC GACATCGCCG TGGAGTGGGA GAGCAATGGG
8581 CAGCCGGAGA ACAACTACAA GACCACGCCT CCGGTGTTGG ACTCCGACGG CTCCTTCTTC
8641 CTCTACAGCA AGCTCACCGT GGACAAGAGC AGGTGGCAGC AGGGGAACGT CTTCTCATGC
8701 TCCGTGATGC ATGAGGCTCT GCACAACCAC TACACGCAGA AGAGCCTCTC CCTGTCTCCG
8761 GGTAAA

```

B. Full Length FVIII_h Fc

(i) Full Length FVIII_h Fc DNA Sequence (FVIII signal peptide underlined, Fc region in bold) (SEQ ID NO:5, which encodes SEQ ID NO:6)

```

661                                     ATG CAAATAGAGC TCTCCACCTG
721 CTTCTTTCTG TGCCTTTTGC GATTCTGCTT TAGTGCCACC AGAAGATACT ACCTGGGTGC
781 AGTGGAAGCTG TCATGGGACT ATATGCAAAG TGATCTCGGT GAGCTGCCTG TGGACGCAAG
841 ATTTCTCTCT AGAGTGCCAA AATCTTTTCC ATTCAACACC TCAGTCGTGT ACAAAAAGAC
901 TCTGTTTGTA GAATTCACGG ATCACCTTTT CAACATCGCT AAGCCAAGGC CACCCTGGAT
961 GGGTCTGCTA GGTCTACCA TCCAGGCTGA GGTTTATGAT ACAGTGGTCA TTACACTTAA
1021 GAACATGGCT TCCCATCCTG TCAGTCTTCA TGCTGTTGGT GTATCCTACT GGAAAGCTTC
1081 TGAGGGAGCT GAATATGATG ATCAGACCAG TCAAAGGGAG AAAGAAGATG ATAAAGTCTT
1141 CCCTGGTGGA AGCCATACAT ATGTCTGGCA GGTCCCGAAA GAGAATGGTC CAATGGCCTC
1201 TGACCCACTG TGCCTTACCT ACTCATATCT TTCTCATGTG GACCTGGTAA AAGACTTGAA
1261 TTCAGGCCTC ATTGGAGCCC TACTAGTATG TAGAGAAGGG AGTCTGGCCA AGGAAAAGAC
1321 ACAGACCTTG CACAAATTTA TACTACTTTT TGCTGTATTT GATGAAGGGA AAAGTTGGCA
1381 CTCAGAAACA AAGAACTCCT TGATGCAGGA TAGGGATGCT GCATCTGCTC GGGCCTGGCC
1441 TAAATGCAC ACAGTCAATG GTTATGTAAA CAGGTCTCTG CCAGGTCTGA TTGGATGCCA
1501 CAGGAAATCA GTCTATTGGC ATGTGATTGG AATGGGCACC ACTCCTGAAG TGCACTCAAT
1561 ATTCCTCGAA GGTCACACAT TTCTTGTGAG GAACCATCGC CAGGCGTCCT TGGAAATCTC-
1621 GCCAATAACT TTCTTACTG CTCAAACACT CTTGATGGAC CTTGGACAGT TTCTACTGTT
1681 TTGTCATATC TCTTCCCACC AACATGATGG CATGGAAGCT TATGTCAAAG TAGACAGCTG
1741 TCCAGAGGAA CCCCACCTAC GAATGAAAAA TAATGAAGAA GCGGAAGACT ATGATGATGA
1801 TCTTACTGAT TCTGAAATGG ATGTGGTCAG GTTTGATGAT GACAACTCTC CTTCTTTTAT
1861 CCAAAATCGC TCAGTTGCCA AGAAGCATCC TAAAACTTGG GTACATTACA TTGCTGCTGA
1921 AGAGGAGGAC TGGGACTATG CTCCCTTAGT CCTCGCCCCC GATGACAGAA GTTATAAAAG
1981 TCAATATTTG AACAATGGCC CTCAGCGGAT TGGTAGGAAG TACAAAAAAG TCCGATTTAT
2041 GGCATACACA GATGAAACCT TTAAGACTCG TGAAGCTATT CAGCATGAAT CAGGAATCTT
2101 GGGACCTTTA CTTTATGGGG AAGTTGGAGA CACACTGTTG ATTATATTTA AGAATCAAGC
2161 AAGCAGACCA TATAACATCT ACCCTCACGG AATCACTGAT GTCCGTCTCT TGTATTCAAG
2221 GAGATTACCA AAAGGTGTAA AACATTTGAA GGATTTTCCA ATTCTGCCAG GAGAAATATT
2281 CAAATATAAA TGGACAGTGA CTGTAGAAGA TGGGCCAACT AAATCAGATC CTCGGTGCCT
2341 GACCCGCTAT TACTCTAGTT TCGTTAATAT GGAGAGAGAT CTAGCTTCAG GACTCATTGG
2401 CCCTCTCCTC ATCTGCTACA AAGAATCTGT AGATCAAAGA GGAAACCAGA TAATGTCAGA
2461 CAAGAGGAAT GTCATCCTGT TTTCTGTATT TGATGAGAAC CGAAGCTGGT ACCTCACAGA
2521 GAATATACAA CGCTTTCTCC CCAATCCAGC TGGAGTGCAG CTTGAGGATC CAGAGTTCCA
2581 AGCCTCCAAC ATCATGCACA GCATCAATGG CATAGTTTTT GATAGTTTGC AGTTGTCAGT
2641 TTGTTTGCAT GAGGTGGCAT ACTGGTACAT TCTAAGCATT GGAGCACAGA CTGACTTCCT
2701 TTCTGTCTTC TTCTCTGGAT ATACCTTCAA ACACAAAATG GTCTATGAAG ACACACTCAC
2761 CCTATTCCCA TTCTCAGGAG AAAGTGTCTT CATGTCTGATG GAAAACCCAG GTCTATGGAT
2821 TCTGGGGTGC CACAACCTCAG ACTTTCGGAA CAGAGGCATG ACCGCCTTAC TGAAGGTTTC
2881 TAGTTGTGAC AAGAACACTG GTGATTATTA CGAGGACAGT TATGAAGATA TTTCAGCATA
2941 CTTGCTGAGT AAAAACAATG CCATTGAACC AAGAAGCTTC TCCCAGAATT CAAGACACCC
3001 TAGCACTAGG CAAAAGCAAT TTAATGCCAC CACAATTCCA GAAAATGACA TAGAGAAGAC
3061 TGACCCTTGG TTTGCACACA GAACACCTAT GCCTAAAATA CAAAATGTCT CCTCTAGTGA
3121 TTTGTTGATG CTCTTGCGAC AGAGTCTTAC TCCACATGGG CTATCCTTAT CTGATCTCCA
3181 AGAAGCCAAA TATGAGACTT TTTCTGATGA TCCATCACCT GGAGCAATAG ACAGTAATAA
3241 CAGCTGTCT GAAATGACAC ACTTCAGGCC ACAGCTCCAT CACAGTGGGG ACATGGTATT
3301 TACCCCTGAG TCAGGCCTCC AATTAAGATT AAATGAGAAA CTGGGGACAA CTGCAGCAAC
3361 AGAGTTGAAG AAAGTTGATT TCAAAGTTTC TAGTACATCA AATAATCTGA TTTCAACAAT
3421 TCCATCAGAC AATTTGGCAG CAGGTACTGA TAATACAAAG TCCTTAGGAC CCCCAGTAT
3481 GCCAGTTCAT TATGATAGTC AATTAGATAC CACTCTATTT GGCAAAAAGT CATCTCCCCT
3541 TACTGAGTCT GGTGGACCTC TGAGCTTGAG TGAAGAAAAT AATGATTCAA AGTTGTTAGA
3601 ATCAGGTTTA ATGAATAGCC AAGAAAGTTC ATGGGGAAAA AATGTATCGT CAACAGAGAG
3661 TGGTAGGTTA TTTAAAGGGA AAAGAGCTCA TGGACCTGCT TTGTTGACTA AAGATAATGC

```

3721	CTTATTCAAAA	GTTAGCATCT	CTTTGTAAAA	GACAAACAAA	ACTTCCAATA	ATTCAGCAAC
3781	TAATAGAAAAG	ACTCACATTG	ATGGCCCATC	ATTATTAATT	GAGAATAGTC	CATCAGTCTG
3841	GCAAAATATA	TTAGAAAGTG	ACACTGAGTT	TAAAAAAGTG	ACACCTTTGA	TTCATGACAG
3901	AATGCTTATG	GACAAAAATG	CTACAGCTTT	GAGGCTAAAT	CATATGTCAA	ATAAACTAC
3961	TTCATCAAAA	AACATGGAAA	TGGTCCAACA	GAAAAAGAG	GGCCCCATTC	CACCAGATGC
4021	ACAAAATCCA	GATATGTCGT	TCTTTAAGAT	GCTATTCTTG	CCAGAATCAG	CAAGGTGGAT
4081	ACAAAGGACT	CATGGAAAGA	ACTCTCTGAA	CTCTGGGCAA	GGCCCCAGTC	CAAAGCAATT
4141	AGTATCCTTA	GGACCAGAAA	AATCTGTGGA	AGGTCAGAAT	TTCTTGCTCG	AGAAAAACAA
4201	AGTGGTAGTA	GGAAAGGGTG	AATTTACAAA	GGACGTAGGA	CTCAAAGAGA	TGGTTTTTCC
4261	AAGCAGCAGA	AACCTATTTT	TTACTAACTT	GGATAATTTA	CATGAAAATA	ATACACACAA
4321	TC AAGAAAAA	AAAAATTCAGG	AAGAAATAGA	AAAGAAGGAA	ACATTAATCC	AAGAGAATGT
4381	AGTTTTGCCT	CAGATACATA	CAGTGACTGG	CACTAAGAAT	TTCATGAAGA	ACCTTTTCTT
4441	ACTGAGCACT	AGGCAAAATG	TAGAAGGTTT	ATATGACGGG	GCATATGCTC	CAGTACTTCA
4501	AGATTTTAGG	TCATTAAATG	ATTCAACAAA	TAGAACAAAG	AAACACACAG	CTCATTTCTC
4561	AAAAAAGGG	GAGGAAGAAA	ACTTGGAAGG	CTTGGGAAAT	CAAACCAAGC	AAATTGTAGA
4621	GAAATATGCA	TGCACCACAA	GGATATCTCC	TAATACAAGC	CAGCAGAATT	TTGTCACGCA
4681	ACGTAGTAAG	AGAGCTTTGA	AACAAATCAG	ACTCCCACTA	GAAGAAACAG	AACTTGAAAA
4741	AAGGATAATT	GTGGATGACA	CCTCAACCCA	GTGGTCCAAA	AACATGAAAC	ATTTGACCCC
4801	GAGCACCCCT	ACACAGATAG	ACTACAATGA	GAAGGAGAAA	GGGGCCATTA	CTCAGTCTCC
4861	CTTATCAGAT	TGCCTTACGA	GGAGTCATAG	CATCCCTCAA	GCAAATAGAT	CTCCATTACC
4921	CATTGCAAAAG	GTATCATCAT	TTCCATCTAT	TAGACCTATA	TATCTGACCA	GGGTCCTATT
4981	CCAAGACAAC	TCTTCTCATC	TTCCAGCAGC	ATCTTATAGA	AAGAAAGATT	CTGGGGTCCA
5041	AGAAAGCAGT	CATTTCTTAC	AAGGAGCCAA	AAAAAATAAC	CTTTCCTTAG	CCATTCTAAC
5101	CTTGGAGATG	ACTGGTGATC	AAAGAGAGGT	TGGCTCCCTG	GGGACAAGTG	CCACAAATTC
5161	AGTCACATAC	AAGAAAGTTG	AGAACACTGT	TCTCCCGAAA	CCAGACTTGC	CCAAAACATC
5221	TGGCAAAGTT	GAATTGCTTC	CAAAAAGTTCA	CATTTATCAG	AAGGACCTAT	TCCCTACGGA
5281	AACTAGCAAT	GGGTCTCCTG	GCCATCTGGA	TCTCGTGGAA	GGGAGCCTTC	TTCAGGGAAC
5341	AGAGGGAGCG	ATTAAGTGGA	ATGAAGCAAA	CAGACCTGGA	AAAGTTCCCT	TTCTGAGAGT
5401	AGCAACAGAA	AGCTCTGCAA	AGACTCCCTC	CAAGCTATTG	GATCCTCTTG	CTTGGGATAA
5461	CCACTATGGT	ACTCAGATAC	CAAAAAGAAGA	GTGGAAATCC	CAAGAGAAGT	CACCAGAAAA
5521	AACAGCTTTT	AAGAAAAAGG	ATACCATTTT	GTCCCTGAAC	GCTTGTGAAA	GCAATCATGC
5581	AATAGCAGCA	ATAAATGAGG	GACAAAATAA	GCCCGAAATA	GAAGTCACCT	GGGCAAAGCA
5641	AGGTAGGACT	GAAAGGCTGT	GCTCTCAAAA	CCCACCAGTC	TTGAAACGCC	ATCAACGGGA
5701	AATAACTCGT	ACTACTCTTC	AGTCAGATCA	AGAGGAAATT	GACTATGATG	ATACCATATC
5761	AGTTGAAATG	AAGAAGGAAG	ATTTTGACAT	TTATGATGAG	GATGAAAATC	AGAGCCCCCG
5821	CAGCTTTCAA	AAGAAAACAC	GACACTATTT	TATTGCTGCA	GTGGAGAGGC	TCTGGGATTA
5881	TGGGATGAGT	AGCTCCCCAC	ATGTTCTAAG	AAACAGGGCT	CAGAGTGGCA	GTGTCCCTCA
5941	GTTCAAGAAA	GTTGTTTTCC	AGGAATTTAC	TGATGGCTCC	TTTACTCAGC	CCTTATACCG
6001	TGGAGAACTA	AATGAACATT	TGGGACTCCT	GGGGCCATAT	ATAAGAGCAG	AAGTTGAAGA
6061	TAATATCATG	GTAACCTTCA	GAAATCAGGC	CTCTCGTCCC	TATTCCTTCT	ATTCTAGCCT
6121	TATTTCTTAT	GAGGAAGATC	AGAGGCAAGT	AGCAGAACCT	AGAAAAAATC	TTGTCAAGCC
6181	TAAATGAAACC	AAAACCTACT	TTTGGAAGGT	GCAACATCAT	ATGGCACCCA	CTAAAGATGA
6241	GTTTGACTGC	AAAGCCTGGG	CTTATTTCTC	TGATGTTGAC	CTGGAAAAAG	ATGTGCACTC
6301	AGGCCTGATT	GGACCCCTTC	TGGTCTGCCA	CACTAACACA	CTGAACCCCTG	CTCATGGGAG
6361	ACAAGTGACA	GTACAGGAAT	TTGCTCTGTT	TTTCACCATC	TTTGATGAGA	CCAAAAGCTG
6421	GTACTTCACT	GAAAAATATG	AAAGAAAATG	CAGGGCTCCC	TGCAATATCC	AGATGGAAGA
6481	TCCCACTTTT	AAAGAGAATT	ATCGCTTCCA	TGCATTCAT	GGCTACATAA	TGGATACACT
6541	ACCTGGCTTA	GTAATGGCTC	AGGATCAAAG	GATTCGATGG	TATCTGCTCA	GCATGGGCAG
6601	CAATGAAAAC	ATCCATTCTA	TTCAATTCAG	TGGACATGTG	TTCACTGTAC	GAAAAAAGA
6661	GGAGTATAAA	ATGGCACTGT	ACAATCTCTA	TCCAGGTGTT	TTTGAGACAG	TGGAAATGTT
6721	ACCATCCAAA	GCTGGAATTT	GGCGGGTGGA	ATGCCTTATT	GGCGAGCATC	TACATGCTGG
6781	GATGAGCACA	CTTTTTCTGG	TGTACAGCAA	TAAGTGTCAG	ACTCCCCTGG	GAATGGCTTC
6841	TGGACACATT	AGAGATTTTC	AGATTACAGC	TTCAGGACAA	TATGGACAGT	GGGCCCCAAA
6901	GCTGGCCAGA	CTTCATTATT	CCGGATCAAT	CAATGCCTGG	AGCACCAAGG	AGCCCTTTTC
6961	TTGGATCAAG	GTGGATCTGT	TGGACCAAT	GATTATTCAC	GGCATCAAGA	CCCAGGGTGC
7021	CCGTCAGAA	TTCTCCAGCC	TCTACATCTC	TCAGTTTATC	ATCATGTATA	GTCTTGATGG
7081	GAAGAAGTGG	CAGACTTATC	GAGGAAATTC	CACTGGAACC	TTAATGGTCT	TCTTTGGCAA
7141	TGTGGATTCA	TCTGGGATAA	AACACAATA	TTTTAACCCCT	CCAATTAATTG	CTCGATACAT
7201	CCGTTTGCAC	CCAACCTATT	ATAGCATTCG	CAGCACTCTT	CGCATGGAGT	TGATGGGCTG
7261	TGATTTAAAT	AGTTGCAGCA	TGCCATTGGG	AATGGAGAGT	AAAGCAATAT	CAGATGCACA

7321	GATTACTGCT	TCATCCTACT	TTACCAATAT	GTTTGCCACC	TGGTCTCCTT	CAAAAGCTCG
7381	ACTTACCTC	CAAGGGAGGA	GTAATGCCCTG	GAGACCTCAG	GTGAATAATC	CAAAAGAGTG
7441	GCTGCAAGTG	GACTTCCAGA	AGACAATGAA	AGTCACAGGA	GTAAC TACTC	AGGGAGTAAA
7501	ATCTCTGCTT	ACCAGCATGT	ATGTGAAGGA	GTTCCCTCATC	TCCAGCAGTC	AAGATGGCCA
7561	TCAGTGGACT	CTCTTTTTTC	AGAATGGCAA	AGTAAAGGTT	TTTCAGGGAA	ATCAAGACTC
7621	CTTCACACCT	GTGGTGAAC	CTCTAGACCC	ACCGTTACTG	ACTCGCTACC	TTCGAATTCA
7681	CCCCCAGAGT	TGGGTGCACC	AGATTGCCCT	GAGGATGGAG	GTTCTGGGCT	GCGAGGCACA
7741	GGACCTCTAC	GACAAAAC	ACACATGCCC	ACCGTGCCCA	GCTCCAGAAC	TCC TGGGCGG
7801	ACCGTCAGTC	TTCCTCTTCC	CCCCAAAACC	CAAGGACACC	CTCATGATCT	CCCGGACCCC
7861	TGAGGTCACA	TGCGTGGTGG	TGGACGTGAG	CCACGAAGAC	CCTGAGGTCA	AGTTCAACTG
7921	GTACGTGGAC	GGCGTGGAGG	TGCATAATGC	CAAGACAAAG	CCGCGGGAGG	AGCAGTACAA
7981	CAGCACGTAC	CGTGTGGTCA	GCGTCCTCAC	CGTCCTGCAC	CAGGACTGGC	TGAATGGCAA
8041	GGAGTACAAG	TGCAAGGTCT	CCAACAAAGC	CCTCCCAGCC	CCCATCGAGA	AAACCATCTC
8101	CAAAGCCAAA	GGGCAGCCCC	GAGAACCAACA	GGTGTACACC	CTGCCCCCAT	CCCGGGATGA
8161	GCTGACCAAG	AACCAGGTCA	GCCTGACCTG	CCTGGTCAAA	GGCTTCTATC	CCAGCGACAT
8221	CGCCGTGGAG	TGGGAGAGCA	ATGGGCAGCC	GGAGAACAAC	TACAAGACCA	CGCCTCCCGT
8281	GTTGGACTCC	GACGGCTCCT	TCTTCCTCTA	CAGCAAGCTC	ACCGTGGACA	AGAGCAGGTG
8341	GCAGCAGGGG	AACGTCTTCT	CATGCTCCGT	GATGCATGAG	GCTCTGCACA	ACCACTACAC
8401	GCAGAAGAGC	CTCTCCCTGT	CTCCGGGTAA	A		

(ii) Fc (same sequence as A (ii) (SEQ ID NO:3))

TABLE 11: Polypeptide Sequences

A. B-Domain Deleted FVIII-Fc Monomer Hybrid (BDD FVIII-Fc monomer dimer): created by coexpressing BDD FVIII-Fc and Fc chains.

Construct = HC-LC-Fc fusion. An Fc expression cassette is cotransfected with BDD FVIII-Fc to generate the BDD FVIII-Fc monomer-. For the BDD FVIII-Fc chain, the Fc sequence is shown in bold; HC sequence is shown in double underline; remaining B domain sequence is shown in italics. Signal peptides are underlined.

i) B domain deleted FVIII-Fc chain (19 amino acid signal sequence underlined) (SEQ ID NO:2)

MQIELSTCFFLCLLRFCFS

ATTRYYLGAVELSWDYMQSDLGELPVDARFPFPRVPKSPFNTSVVYKKTLFVEFTDHLF
NIAKPRPPWMGLLGPTIQAEVYDVTVVITLKNMASHPVSLHAVGVSYWKASEGAHEYDDQ
TSQREKEDDKVFPGGSHTYVWQVLKENGPMASDPLCLTYSYLSHVDLVKDINSGLIGA
LLVCREGLAKEKTOTLHKFILLFAVEDEGKSWHSETKNSLMODRDAASARAWPKMHT
VNGYVNRSLPGLIGCHRKSVYWHVIGMGTTPFVHSIFLEGHTFLVRNHROASLEISPITFL
TAQTLLMDLGQFLLFCHISSHOHDGMEAYVKVDSCPEEPQLRMKNNEEAEDYDDDLTD
SEMDVVRFDNNSPFIQIRSVAKKHPKTWVHYIAAEEEDWDYAPLVLPDDRYSYKSOY
LNNGPORIGRKYKKVREMA YDETEKTREAIQHESGILGPLLYGEVGD TLLIIFKNQASRP
YNIYPHGITDVRPLYSRRLPKGVKHLKDFPILPGEIFKYKWTVTVEDGPTKSDPRCLTRY
YSSFVNMERDLASGLIGPLLCYKESVDQRGNOIMSDKRNVLFSVF DENRSWYLTENIQ
RFLPNPAGVQLEDPEFOASNIMHSINGYVFDSLQLSVCLHEVAYWYILSIGAOTDFLSVEF
SGYTFKHKMYEDTLTLPFSGETVFMSMENPGLWILGCHNSDFRNRGMTALLKVSSC
DKNTGDYYEDSYEDISAYLLSKNNAIEPRSF SQNPVLKRHQREITRTTLQSDQEEIDYDD
TISVEMKKEDFDIYDEDENQSPRSFQKKTRHYFIAAVERLWDYGMSSSPHVLNRNRAQSG
SVPQFKKVVFQEFTDGSFTQPLYRGELNEHLGLLGPYIRAEVEDNIMVTFRNQASRPYSF

YSSLISYEEDQRQGAEPKRFVKNPNETKTYFWKVQHMAPTKDEFDCKAWAYFSDVDL
 EKDVHSGGLIGPLL VCHTNTLNPAHGRQVTVQEFALFFTIFDETKSWYFTENMERNCRAP
 CNIQMEDPTFKENYRFHAINGYIMDTLPLGLVMAQDQIRWYLLSMGSNENIHSIHFSGH
 VFTVRKKKEEYKMALYNLYPGVFETVEMLPKAGIWRVECLIGEHLHAGMSTLFLVYSN
 KCQTPLGMASGHIRDFQITASGQYQGWAPKLARLHYSGSINAWSTKEPFSWIKVDLLAP
 MIIHGIKTQGARQKFSSLYISQFIIMYSLDGKKWQTYRGNSTGTLMVFFGNVDSSGIKHNI
 FNPPHARYIRLHPHTHYSIRSTLRMELMGCDLNSCSMPLGMESKAISDAQITASSYFTNMF
 ATWSPSKARLHLQGRSNAWRPQVNNPKEWLQVDFQKTMKVTGVTQGVKSLLTSMY
 VKEFLISSSQDGHQWTLFFQNGKVKVFQGNQDSFTPVVNSLDPPLLTRYLRIHPQSWVH
 QIALRMEVLGCEAQDLYDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTC
 VVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLN
 GKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKG
 FYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVFCSSV
 MHEALHNHYTQKSLSLSPGK

ii) Fc chain (20 amino acid heterologous signal peptide from mouse Igk chain underlined)
 (SEQ ID NO:4)

METDTLLLWVLLLWVPGSTG

DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVD
 GVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISK
 AKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPV
 LDSDGSFFFLYSKLTVDKSRWQQGNVFCSSVMHEALHNHYTQKSLSLSPGK

B. Full length FVIII-Fc monomer hybrid (Full length FVIII-Fc monomer dimer):
 created by coexpressing FVIII-Fc and Fc chains.

Construct = HC-B-LC-Fc fusion. An Fc expression cassette is cotransfected with full length FVIII-Fc to generate the full length FVIII-Fc monomer. For the FVIII-Fc chain, the Fc sequence is shown in bold; HC sequence is shown in double underline; B domain sequence is shown in italics. Signal peptides are underlined.

i) Full length FVIII-Fc chain (FVIII signal peptide underlined) (SEQ ID NO:6)

MQIELSTCFFLCLLRCFCS

ATRRYYLGAVELSWDYMOSDLGELPVDAREPPRVPKSFPFNTSVVYKKTLEVEFTDHLF
NIAPRPPWMGLLGPTIQAEVYDTVVITLKNMASHPVSLHAVGVSYWKASEGAHEYDDQ
TSOREKEDDKVFPGGSHTYVWQVLKENGPMASDPLCLTYSYLSHVDLVKDLNSGLIGA
LLVCREGLAKEKTOTLHKFILLFAVFDEGKSWHSETKNSLMODRDAASARAWPKMHT
VNGYVNRSLPGLIGCHRKSIVYWHVIGMGTTPPEVHSIFLEGHTFLVRNHRQASLEISPITFL
TAQTLLMDLGOFLFCHISSHQHDGMEAYVKVDSCPEEPQLRMKNNEEAEDYDDDLTD
SEMDVVRFDNDDNSPSFIQIRSVAKKHPKTWVHYIAAEEEDWDYAPLV LAPDDRSYKSOY
LNNGPORIGRKYKKVRFEMAYTDETFKTREAIQHESGILGPLLYGEVGDITLLIIFKNQASRP
YNIYPHGITDVRPLYSRRLPKGVKHLKDFILPGEIFKYKWTVTVEDGPTKSDPRCLTRY
YSSFVNMERDLASGLIGPLLCYKESVDORGNOIMSDKRNVLFSVFEDENRSWYLTENIQ
RFLPNPAGVQLEDPEFQASNIMHSINGYVFDSLQSVCLHEVAYWYILSIGAQTDFLSVFE
SGYTFKHKMVYEDTLTLFPFSGETVFMSMENPGLWILGCHNSDFRNRGMTALLKVSSC

DKNTGDYYEDSYEDISAYLLSKNNAIEPRSFSONSRHPSTRQKQFNATTIPENDIEKTDPPWF
 AHRTMPKIQNVSSDLLMLLRQSPTPHGLSLSDLQEAKYETFSDDPSPGAIDSNNLSSEMTHF
 RPQLHHSMDVFTPEGLQLRLNEKLGTTAATELKKLDFKVSSTSNLISTIPSDNLAAGTDNT
 SSLGPPSPMPVHYDSQLDTTLFGKKSSPLTESGGPLSLSEENNDKLLSGLMNSQESSWGKNV
 SITESGRLFKGKRAHGPALLTKDNALFKVSISLLKTNKTSNNSATNRKTHIDGPSLLIENSPSVW
 QNILESDETEFKKVTPLIHDRMLMDKNATALRLNHMSNKTSSKNMEMVQQKKEGPIPPDAQN
 PDMSFFKMLFLPESARWIQORTHGKNSLNSGQGSPKQLVSLGPEKSVEGQNFLSEKNKVVG
 KGEFTKDVGLKEMVFPSSRNFLTNLDNLHENNTHNQEKKIQEEIEKKETLIQENVVLPQIHT
 VTGTKNFMKNLFLSTRQNVESYDGAYAPVLQDFRSLNDSTNRKKHTAHFSKKGEEENLE
 GLGNQTKQIVEKYACTTRISPNTSQONFVTQRSKRALKQFRLPLEETELEKRIIVDDTSTQWSK
 NMKHLTPSTLTQIDYNEKEKGAITQSPSLDCLTRSHSIPQANRSPLPIAKVSSFPSIRPIYLTRVLF
 QDNSSHLPAAASYRKKDSGVQESSHFLQGAKKNNLSLAILTLEMTGDQREVGSGLGTSATNSVTY
 KKVENTVLPKPDLPKTSKGKVELLPKVHIYQKDLFPTETSNGSPGHLDLVEGSLQGTGAIKW
 NEANRPGKVPFLRVATESSAKTPSKLLDPLAWDNHYGTQIPKEEWSQEKSPKTAFFKKKDTI
 LSLNACESNHAIAAINEGQNKPEIEVTWAKQGRTERLCSQNPPVLKRHQREITRTTLQSDQEEI
 DYDDTISVEMKKEDFDIYDEDENQSPRSFQKKTRHYFIAAVERLWDYGMSSSPHVLNRN
 AQSGSVPQFKKVVFQEFTDGSFTQPLYRGELNEHLGLLGPYIRAEVEDNIMVTFRNQASR
 PYSFYSSLISYEEDQRQGAEPKRFVKNPNETKTYFWKVQHMAPTKDEFDCKAWAYFS
 DVDLEKDVHSGLIGPLLCHTNTLNPAHGRQVTVQEFALFFTIFDETKSWYFTENMERN
 CRAPCNIQMEDPTFKENYRFHAINGYIMDTLPGLVMAQDQIRWYLLSMGSNENIHSIH
 SGHVFTVRKKEEYKMALYNLYPGVFETVEMLPKAGIWRVECLIGEHLHAGMSTLFLV
 YSNKCQTPLGMASGHIRDFQITASGQYGQWAPKLARLHYSGSINAWSTKEPFSWIKVDL
 LAPMIIHGIKTQGARQKFSSLYISQFIIMYSLDGKKWQTYRGNSTGTLMVFFGNVDSSGK
 HNIFNPPIIARYIRLHPHTHYSIRSTLRMELMGCDLNSCSMPLGMESKAISDAQITASSYFTN
 MFATWSPSKARLHLQGRSNAWRPQVNNPKEWLQVDFQKTMKVTGVTQGVKSLTS
 MYVKEFLISSSQDGHQWTLFFQNGKVVFQGNQDSFTPVVNSLDPPLLTRYLRIHPQSW
 VHQIALRMEVLGCEAQDLYDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEV
 TCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKFREEQYNSTYRVVSVLTVLHQDW
 LNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLV
 KGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSC
 SVMHEALHNHYTQKSLSLSPGK

ii) Fc chain (20 amino acid heterologous signal peptide from mouse Igk chain underlined)
 (SEQ ID NO:4)

METDTLLLVLLLVWPGSTG

DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVD
 GVEVHNAKTKFREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISK
 AKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPV
 LDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

[0386] The foregoing description of the specific embodiments will so fully reveal the
 general nature of the invention that others can, by applying knowledge within the skill of
 the art, readily modify and/or adapt for various applications such specific embodiments,

without undue experimentation, without departing from the general concept of the present invention. Therefore, such adaptations and modifications are intended to be within the meaning and range of equivalents of the disclosed embodiments, based on the teaching and guidance presented herein. It is to be understood that the phraseology or terminology herein is for the purpose of description and not of limitation, such that the terminology or phraseology of the present specification is to be interpreted by the skilled artisan in light of the teachings and guidance.

[0387] Other embodiments of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein. It is intended that the specification and examples be considered as exemplary only, with a true scope and spirit of the invention being indicated by the following claims.

[0388] All patents and publications cited herein are incorporated by reference herein in their entirety.

WHAT IS CLAIMED IS:

1. A pharmaceutical composition comprising:
 - (a) a FVIII polypeptide;
 - (b) one or more stabilizing agents selected from the group consisting of sucrose, trehalose, raffinose, arginine, and mixture thereof;
 - (c) sodium chloride (NaCl);
 - (d) L-histidine;
 - (e) calcium chloride; and
 - (f) polysorbate 20 or polysorbate 80.
2. The pharmaceutical composition of claim 1, wherein mannitol, glycine, alanine, or hydroxyethyl starch is not included.
3. The pharmaceutical composition of claim 1 or 2, wherein NaCl is the only bulking agent.
4. The pharmaceutical composition of any one of claims 1 to 3, comprising sucrose.
5. The pharmaceutical composition of claim 4, comprising about 1% (w/v) to about 2.5% (w/v) sucrose.
6. The pharmaceutical composition of claim 5, comprising about 1.3% (w/v) sucrose to about 2.0% (w/v) sucrose.
7. The pharmaceutical composition of claim 6, comprising about 1.33% (w/v) sucrose or about 2.0% (w/v) sucrose.
8. The pharmaceutical composition of claim 4, comprising about 10 mg/ml to about 25 mg/ml sucrose.
9. The pharmaceutical composition of claim 8, comprising about 13 mg/ml to about 20 mg/ml sucrose.

10. The pharmaceutical composition of claim 9, comprising about 13.3 mg/ml sucrose or about 20.0 mg/ml sucrose.
11. The pharmaceutical composition of any one of claims 1 to 10, comprising about 150 mM to about 250 mM NaCl.
12. The pharmaceutical composition of claim 11, comprising about 175 mM to about 225 mM NaCl.
13. The pharmaceutical composition of claim 12, comprising about 200 mM to about 210 mM NaCl.
14. The pharmaceutical composition of claim 13, comprising about 205 mM NaCl.
15. The pharmaceutical composition of any one of claims 1 to 10, comprising about 8.8 mg/ml to about 14.6 mg/ml NaCl.
16. The pharmaceutical composition of claim 15, comprising about 10 mg/ml to about 13 mg/ml NaCl.
17. The pharmaceutical composition of claim 16, comprising about 12.0 mg/ml NaCl.
18. The pharmaceutical composition of any one of claims 1 to 10, comprising about 250 mM to about 350 mM NaCl.
19. The pharmaceutical composition of claim 18, comprising about 275 mM to about 325 mM NaCl.
20. The pharmaceutical composition of claim 19, comprising about 308 mM NaCl.

21. The pharmaceutical composition of any one of claims 1 to 10, comprising about 14.6 mg/ml to about 20.5 mg/ml NaCl.
22. The pharmaceutical composition of claim 21, comprising about 16 mg/ml to about 19 mg/ml NaCl.
23. The pharmaceutical composition of claim 22, comprising about 18.0 mg/ml NaCl.
24. The pharmaceutical composition of any one of claims 1 to 23, comprising about 5 mM to about 15 mM L-histidine.
25. The pharmaceutical composition of claim 24, comprising about 6.64 mM L-histidine or about 9.8 mM L-histidine.
26. The pharmaceutical composition of any one of claims 1 to 23, comprising about 0.75 mg/ml to about 2.25 mg/ml L-histidine.
27. The pharmaceutical composition of claim 26, comprising about 1.03 mg/ml L-histidine or about 1.55 mg/ml L-histidine.
28. The pharmaceutical composition of any one of claims 1 to 27, comprising about 5 mM to about 10 mM calcium chloride.
29. The pharmaceutical composition of claim 28, comprising about 5.4 mM calcium chloride or about 8 mM calcium chloride.
30. The pharmaceutical composition of any one of claims 1 to 27, comprising about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate.
31. The pharmaceutical composition of claim 30, comprising about 0.8 mg/ml calcium chloride dihydrate or about 1.18 mg/ml calcium chloride dihydrate.

32. The pharmaceutical composition of any one of claims 1 to 31, comprising about 0.008% (w/v) to about 0.025% (w/v) polysorbate 20 or polysorbate 80.
33. The pharmaceutical composition of claim 32, comprising about 0.013% (w/v) polysorbate 20 or polysorbate 80 or about 0.02 % (w/v) polysorbate 20 or polysorbate 80.
34. The pharmaceutical composition of any one of claims 1 to 31, comprising about 0.08 mg/ml to about 0.25 mg/ml polysorbate 20 or polysorbate 80.
35. The pharmaceutical composition of claim 34, comprising about 0.13% mg/ml polysorbate 20 or polysorbate 80 or about 0.20 mg/ml polysorbate 20 or polysorbate 80.
36. The pharmaceutical composition of any one of claims 1 to 35, wherein the FVIII polypeptide comprises a first subunit comprising an amino acid sequence at least 90% or 95% identical to amino acids 20 to 1684 of SEQ ID NO:2 or 20 to 2578 of SEQ ID NO:6, and a second subunit comprising an amino acid sequence at least 90% to 95% identical to amino acids 21 to 247 of SEQ ID NO:4.
37. The pharmaceutical composition of claim 36, wherein the FVIII polypeptide comprises a first subunit comprising amino acids 20 to 1684 of SEQ ID NO:2 or 20 to 2578 of SEQ ID NO:6, and a second subunit comprising amino acids 21 to 247 of SEQ ID NO:4.
38. The pharmaceutical composition of any one of claims 1 to 37, wherein the FVIII polypeptide is present at a concentration of about 50 IU/ml to about 2500 IU/ml.
39. The pharmaceutical composition of claim 38, comprising 83 IU/ml, 167 IU/ml, 250 IU/ml, 333 IU/ml, 500 IU/ml, 667 IU/ml, 1000 IU/ml, 1333 IU/ml, 1667 IU/ml, or 2000 IU/ml of the FVIII polypeptide.
40. The pharmaceutical composition of any one of claims 1 to 37, wherein the FVIII polypeptide is present at a concentration of about 100 IU/ml to about 4000 IU/ml.

41. The pharmaceutical composition of claim 40, comprising 150 IU/ml, 287.5 IU/ml, 431.25 IU/ml, 575 IU/ml, 862.5 IU/ml, 1150 IU/ml, 1725 IU/ml, 2300 IU/ml, 2875 IU/ml, or 3450 IU/ml of the FVIII polypeptide.
42. The pharmaceutical composition of any one of claims 1 to 4, comprising:
 - (a) about 50 IU/ml to about 2500 IU/ml of the FVIII polypeptide;
 - (b) about 1% (w/v) to about 2.5% (w/v) of sucrose;
 - (c) about 150 mM to about 250 mM NaCl;
 - (d) about 5 mM to about 15 mM L-histidine;
 - (e) about 5 mM to about 10 mM calcium chloride; and
 - (f) about 0.008% (w/v) to about 0.025% of polysorbate 20 or polysorbate 80.
43. The pharmaceutical composition of claim 42, comprising about 175 mM to about 225 mM NaCl.
44. The pharmaceutical composition of claim 43, comprising about 200 mM to about 210 mM NaCl.
45. The pharmaceutical composition of claim 44, comprising:
 - (a) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, about 1000 IU/ml, , 1333 IU/ml, 1667 IU/ml, or 2000 IU/ml of the FVIII polypeptide;
 - (b) about 1.33% (w/v) of sucrose;
 - (c) about 205 mM NaCl;
 - (d) about 6.64 mM L-histidine;
 - (e) about 5.4 mM calcium chloride; and
 - (f) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.
46. The pharmaceutical composition of any one of claims 1 to 4, comprising:
 - (a) about 100 IU/ml to about 4000 IU/ml of the FVIII polypeptide;
 - (b) about 1% (w/v) to about 2.5% (w/v) of sucrose;
 - (c) about 250 mM to about 350 mM NaCl;

- (d) about 5 mM to about 15 mM L-histidine;
 - (e) about 5 mM to about 10 mM calcium chloride; and
 - (f) about 0.008% (w/v) to about 0.025% of polysorbate 20 or polysorbate 80.
47. The pharmaceutical composition of claim 46, comprising about 275 mM to about 325 mM NaCl.
48. The pharmaceutical composition of claim 47, comprising:
- (a) about 150 IU/ml, about 287.5 IU/ml, about 431.25 IU/ml, about 575 IU/ml, about 862.5 IU/ml, about 1150 IU/ml, about 1725 IU/ml, about 2300 IU/ml, about 2875 IU/ml, or about 3450 IU/ml of the FVIII polypeptide;
 - (b) about 2.0% (w/v) of sucrose;
 - (c) about 308 mM NaCl;
 - (d) about 9.8 mM L-histidine;
 - (e) about 8 mM calcium chloride; and
 - (f) about 0.020% (w/v) of polysorbate 20 or polysorbate 80.
49. The pharmaceutical composition of any one of claims 1 to 4, comprising:
- (a) about 50 IU/ml to about 2500 IU/ml of the FVIII polypeptide;
 - (b) about 10 mg/ml to about 25 mg/ml of sucrose;
 - (c) about 8.8 mg/ml to about 14.6 mg/ml NaCl;
 - (d) about 0.75 mg/ml to about 2.25 mg/ml L-histidine;
 - (e) about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate; and
 - (f) about 0.08 mg/ml to about 0.25 mg/ml of polysorbate 20 or polysorbate 80.
50. The pharmaceutical composition of claim 49, comprising about 10 mg/ml to 13 mg/ml NaCl.
51. The pharmaceutical composition of claim 50, comprising:
- (a) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, about 1000 IU/ml, about 1333 IU/ml, about 1667 IU/ml, or about 2000 IU/ml of the FVIII polypeptide;

- (b) about 13.3 mg/ml of sucrose;
 - (c) about 12.0 mg/ml NaCl;
 - (d) about 1.03 mg/ml L-histidine;
 - (e) about 0.8 mg/ml calcium chloride dihydrate; and
 - (f) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.
52. The pharmaceutical composition of any one of claims 1 to 4, comprising:
- (a) about 100 IU/ml to about 4000 IU/ml of the FVIII polypeptide;
 - (b) about 10 mg/ml to about 25 mg/ml of sucrose;
 - (c) about 14.6 mg/ml to about 20.5 mg/ml NaCl;
 - (d) about 0.75 mg/ml to about 2.25 mg/ml L-histidine;
 - (e) about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate; and
 - (f) about 0.08 mg/ml to about 0.25 mg/ml of polysorbate 20 or polysorbate 80.
53. The pharmaceutical composition of claim 52, comprising about 16 mg/ml to about 19 mg/ml NaCl.
54. The pharmaceutical composition of claim 53, comprising:
- (a) about 150 IU/ml, about 287.5 IU/ml, about 431.25 IU/ml, about 575 IU/ml, about 862.5 IU/ml, about 1150 IU/ml, about 1725 IU/ml, about 2300 IU/ml, about 2875 IU/ml, or about 3450 IU/ml of the FVIII polypeptide;
 - (b) about 20.0 mg/ml of sucrose;
 - (c) about 18.0 mg/ml NaCl;
 - (d) about 1.55 mg/ml L-histidine;
 - (e) about 1.18 mg/ml calcium chloride dihydrate; and
 - (f) about 0.20 mg/ml of polysorbate 20 or polysorbate 80.
55. A pharmaceutical kit comprising:
- (a) a first container comprising a lyophilized powder, where the powder comprises
 - (i) the FVIII polypeptide,
 - (ii) one or more stabilizing agents selected from the group consisting of sucrose, trehalose, raffinose, arginine, and mixture thereof;

- (iii) sodium chloride (NaCl);
 - (iv) L-histidine;
 - (v) calcium chloride; and
 - (vi) polysorbate 20 or polysorbate 80; and
 - (b) a second container comprising sterilized water for injections to be combined with the lyophilized powder of the first container.
56. The pharmaceutical kit of claim 55, wherein mannitol, glycine, alanine, or hydroxyethyl starch is not included.
57. The pharmaceutical kit of claim 55 or 56, wherein NaCl is the only bulking agent.
58. The pharmaceutical kit of any one of claims 55 to 57, comprising:
- (a) a first container comprising a lyophilized powder, where the powder comprises
 - (i) about 250 IU, about 500 IU, about 750 IU, about 1000 IU, about 1500 IU, about 2000 IU, about 3000 IU, about 4000 IU, about 5000 IU, or about 6000 IU of the FVIII polypeptide,
 - (ii) about 40 mg of sucrose;
 - (iii) about 36 mg of sodium chloride;
 - (iv) about 3.1 mg of L-histidine;
 - (v) about 2.40 mg of calcium chloride dihydrate; and
 - (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and
 - (b) a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising:
 - (i) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, about 1000 IU/ml, about 1333 IU/ml, about 1667 IU/ml, or about 2000 IU/ml of the FVIII polypeptide, respectively;
 - (ii) about 1.33% (w/v) of sucrose;
 - (iii) about 205 mM NaCl;
 - (iv) about 6.64 mM L-histidine;

- (v) about 5.4 mM of calcium chloride; and
 - (vi) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.
59. The pharmaceutical kit of any one of claims 55 to 57, comprising:
- (a) a first container comprising a lyophilized powder, where the powder comprises
 - (i) about 250 IU, about 500 IU, about 750 IU, about 1000 IU, about 1500 IU, about 2000 IU, or 3000 IU, about 4000 IU, about 5000 IU, or about 6000 IU of the FVIII polypeptide,
 - (ii) about 40 mg of sucrose;
 - (iii) about 36 mg of sodium chloride;
 - (iv) about 3.1 mg of L-histidine;
 - (v) about 2.40 mg of calcium chloride dihydrate; and
 - (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and
 - (b) a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising:
 - (i) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, about 1000 IU/ml, about 1333 IU/ml, about 1667 IU/ml, or about 2000 IU/ml of the FVIII polypeptide, respectively;
 - (ii) about 13.3 mg/ml of sucrose;
 - (iii) about 12.0 mg/ml of NaCl;
 - (iv) about 1.03 mg/ml of L-histidine;
 - (v) about 0.80 mg/ml of calcium chloride; and
 - (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.
60. The pharmaceutical kit of any one of claims 55 to 59, wherein the first container is a glass vial comprising a rubber stopper.
61. The pharmaceutical kit of any one of claims 55 to 60, wherein the second container is a syringe body, and wherein the syringe body is associated with a plunger.

62. The pharmaceutical kit of claim 61, further comprising an adaptor to connect the glass vial to the syringe body.
63. The pharmaceutical kit of claim 61 or 62, further comprising infusion tubing associated with a needle to be connected to the syringe, suitable for intravenous infusion.
64. A method of reducing the annualized bleeding rate in a subject having hemophilia comprising administering to the subject the pharmaceutical composition of any one of claims 1 to 54 at a dosing interval of about three days or longer.
65. The method of claim 64, wherein the administration is prophylactic and individualized for the subject resulting in an annualized bleeding rate less than about 5.0, less than about 4.9, less than about 4.8, less than about 4.7, less than about 4.6, or less than about 4.5.
66. The method of claim 65, wherein the median annualized bleeding rate is about 1.6.
67. The method of claim 64, wherein the administration is prophylactic and weekly resulting in an annualized bleeding rate less than about 9.0, less than about 8.9, less than about 8.8, less than about 8.7, less than about 8.6, less than about 8.5, or less than about 8.4.
68. The method of claim 67, wherein the median annualized bleeding rate is about 3.6.
69. The method of claim 64, wherein the administration is on-demand or episodic resulting in an annualized bleeding rate less than about 55, less than about 54, less than about 53, less than about 52, less than about 51, less than about 50, less than about 49, less than about 48, or less than about 47.
70. The method of claim 69, wherein the median annualized bleeding rate is about 33.6.

71. The method of any one of claims 64 to 70, wherein the pharmaceutical composition comprises an effective dose of the FVIII polypeptide, which is between about 20 IU/kg to about 90IU/kg.
72. The method of claim 71, wherein the effective dose is 20-30 IU/kg, 30-40 IU/kg, 40-50 IU/kg, 50-60 IU/kg, 60-70 IU/kg, 70-80 IU/kg, or 80-90 IU/kg.
73. The method of claim 72, wherein the effective dose is 20 IU/kg, 25 IU/kg, 30 IU/kg, 35 IU/kg, 40 IU/kg, 45 IU/kg, 50 IU/kg, 55 IU/kg, 60 IU/kg, 65 IU/kg, 70 IU/kg, 75 IU/kg, 80 IU/kg, 85 IU/kg, or 90 IU/kg.
74. The method of any of claims 71 to 73, wherein the administration is prophylactic and individualized at an effective dose of about 25 IU/kg to about 65 IU/kg twice weekly or every three days or about 50 IU/kg to about 65 IU/kg every 4 or 5 days.
75. The method of claim 74, wherein a first dose of the FVIII polypeptide is administered is about 25 IU/kg and a second dose of the FVIII polypeptide is about 50 IU/kg.
76. The method of any of claims 71 to 73, wherein the administration is prophylactic and weekly at an effective dose of 65 IU/kg weekly.
77. The method of any of claims 71 to 73, wherein the administration is on-demand or episodic at an effective dose of 10 IU/kg to 75 IU/kg every 12 to 24 hours.
78. The method of any one of claims 64 to 70, wherein the pharmaceutical composition comprises an effective dose, which is a fixed dose that is standard across all body weights.
79. The method of claim 78, wherein the fixed dose is about 2,000 IU, about 2,500 IU, about 3,000 IU, about 3,500 IU, about 4,000 IU, about 5,000 IU, or about 6,000IU per dose.

80. The method of any one of claims 64 to 70, wherein the pharmaceutical composition comprises an effective dose, which is a stratified dose.
81. The method of any one of claims 64 to 70, wherein a trough level of the FVIII polypeptide is above 1%, above 2%, or above 3% of normal after the administration.
82. The method of any one of claims 64 to 81, wherein the FVIII polypeptide has a $T_{1/2\text{beta}}$ (activity) of about 7 hours to about 48 hours, about 6 hours to about 49 hours, about 5 hours to about 50 hours.
83. The method of claim 82, wherein the FVIII polypeptide has a $T_{1/2\text{beta}}$ (activity) mean of at least about 15 hours, at least about 16 hours, at least about 17 hours, at least about 18 hours, at least about 19 hours, at least about 20 hours, at least about 21 hours, at least about 22 hours, at least about 23 hours, at least about 24 hours, or at least about 25 hours.
84. The method of claim 83, wherein the $T_{1/2\text{beta}}$ (activity) mean is about 19 hours.
85. The method of claim 82 or 83, wherein the $T_{1/2\text{beta}}$ (activity) mean is at least about 1.5 fold higher than a polypeptide consisting of amino acids 20 to 1457 of SEQ ID NO:2, 20 to 2351 of SEQ ID NO: 6, or ADVATE[®].
86. The method of any one of claims 64 to 85, wherein the plasma trough level of the FVIII polypeptide is maintained between about 1% and about 5%, between about 1% and about 6%, between about 1% and about 7%, between about 1% and about 8%, between about 1% and about 9%, between about 1% and about 10%, between about 1% and about 11%, between about 1% and about 12%, between about 1% and about 13%, between about 1% and about 14%, between about 1% and about 15% above the baseline in the subject after the administration.
87. The pharmaceutical composition of any one of claims 1 to 35 or 38 to 54, the pharmaceutical kit of any one of claims 55 to 63, or the method of any one of claims 64 to

- 86, wherein the FVIII polypeptide is a long-acting FVIII polypeptide comprising a FVIII polypeptide and a heterologous moiety.
88. The pharmaceutical composition, the pharmaceutical kit, or the method of claim 87, wherein the heterologous moiety is an FcRn binding partner.
89. The pharmaceutical composition, the pharmaceutical kit, or the method of claim 88, wherein the FcRn binding partner comprises an Fc region.
90. The pharmaceutical composition, the pharmaceutical kit, or the method of claim 88 or 89, wherein the long-acting FVIII polypeptide further comprises a second FcRn binding partner.
91. The pharmaceutical composition, the pharmaceutical kit, or the method of claim 90, wherein the second FcRn binding partner comprises a second Fc region.
92. The pharmaceutical composition, the pharmaceutical kit, or the method of claim 90 or 91, wherein the FcRn binding partner and the second FcRn binding partner are associated.
93. The pharmaceutical composition, the pharmaceutical kit, or the method of claim 92, wherein the association is a covalent bond.
94. The pharmaceutical composition, the pharmaceutical kit, or the method of claim 93, wherein the covalent bond is a disulfide bond.
95. The pharmaceutical composition, the pharmaceutical kit, or the method of any one of claims 90 to 94, wherein the second FcRn binding partner is not linked to an amino acid sequence by a peptide bond.

96. The pharmaceutical composition, the pharmaceutical kit, or the method of any one of claims 87 to 95, wherein the long-acting FVIII polypeptide is FVIII monomer dimer hybrid.
97. The pharmaceutical composition, the pharmaceutical kit, or the method of any one of claims 87 to 96, wherein the FVIII polypeptide in the long-acting polypeptide is a human FVIII.
98. The pharmaceutical composition, the pharmaceutical kit, or the method of any one of claims 87 to 97, wherein the FVIII polypeptide in the long-acting polypeptide is a full-length FVIII or a B-domain deleted FVIII.
99. The pharmaceutical composition, the pharmaceutical kit, or the method of any one of claims 87 to 97, wherein the FVIII polypeptide is single chain.
100. The pharmaceutical composition, the pharmaceutical kit, or the method of any one of claims 87 to 97, wherein the FVIII polypeptide is unprocessed.
101. The pharmaceutical composition, the pharmaceutical kit, or the method of any one of claims 87 to 97, wherein the FVIII polypeptide is in two chains, a first chain comprising a heavy chain of the FVIII polypeptide and a second chain comprising a light chain of the FVIII polypeptide.
102. The pharmaceutical composition, the pharmaceutical kit, or the method of any one of claims 88 to 101, wherein the FcRn binding partner in the chimeric polypeptide is a human Fc.
103. The pharmaceutical composition, the pharmaceutical kit, or the method of any one of claims 87 to 102, wherein the FVIII polypeptide is at least 60%, 70%, 80%, 90%, 95%, 95%, 97%, 98%, 99%, or 100% identical to a FVIII amino acid sequence shown in Table 11A or 11B without a signal sequence (amino acids 20 to 1457 of SEQ ID NO: 2 or amino acids 20 to 2351 of SEQ ID NO: 6).

104. The pharmaceutical composition, the pharmaceutical kit, or the method of any one of claims 88 to 103, wherein the FcRn binding partner is at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to a Fc amino acid sequence shown in Table 11B without a signal sequence (amino acids 21 to 247 SEQ ID NO:4).
105. The pharmaceutical composition, the pharmaceutical kit, or the method of any one of claims 90 to 104, wherein the second FcRn binding partner in the chimeric polypeptide is a human Fc.
106. The pharmaceutical composition, the pharmaceutical kit, or the method of claim 105, wherein the second FcRn binding partner is at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to a Fc amino acid sequence shown in Table 11B without a signal sequence (amino acids 21 to 247 SEQ ID NO:4)
107. The pharmaceutical composition of any one of claims 1 to 54, the pharmaceutical kit of any one of claims 55 to 63, or the method of any one of claims 64 to 106, wherein the pharmaceutical composition is for perioperative management of a bleeding episode.
108. The method of any one of claims 64 to 107, wherein the subject is in need of controlling or preventing bleeding or bleeding episodes.
109. The method of claim 108, wherein the subject is in need of controlling or preventing bleeding in minor hemorrhage, hemarthroses, superficial muscle hemorrhage, soft tissue hemorrhage, moderate hemorrhage, intramuscle or soft tissue hemorrhage with dissection, mucous membrane hemorrhage, hematuria, major hemorrhage, hemorrhage of the pharynx, hemorrhage of the retropharynx, hemorrhage of the retroperitoneum, hemorrhage of the central nervous system, bruises, cuts, scrapes, joint hemorrhage, nose bleed, mouth bleed, gum bleed, intracranial bleeding, intraperitoneal bleeding, minor spontaneous hemorrhage, bleeding after major trauma, moderate skin bruising, or spontaneous hemorrhage into joints, muscles, internal organs or the brain.

110. The method of claim 109, wherein the subject is in need of management of bleeding associated with surgery or dental extraction.
111. The method of claim 110, wherein the subject will undergo, is undergoing, or has undergone major surgery.
112. The method of claim 111, wherein the major surgery is orthopedic surgery, extensive oral surgery, urologic surgery, or hernia surgery.
113. The method of claim 112, wherein the orthopedic surgery is replacement of knee, hip, or other major joint.
114. The method of any one of claims 64 to 113, wherein the subject is in need of long-term treatment.
115. The method of any one of claims 64 to 114, wherein the FVIII polypeptide is administered intravenously or subcutaneously.
116. The pharmaceutical composition of any one of claims 18-23, 40, 41, 46-48, and 52-54, wherein the pharmaceutical composition is a pre-lyophilization composition.