


(**I-1**)




(**I-2**)

or a salt thereof; wherein the variables are as defined below.

The present disclosure also provides a pharmaceutical composition comprising a compound of Formula (**I-1**) or (**I-2**), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier or excipient.

The compounds of Formula (**I-1**) or (**I-2**) are useful in the treatment of MASP-2-associated diseases and disorders, and in the manufacture of medicaments for treating MASP-2-associated diseases and disorders. The present disclosure also provides methods of treating a MASP-2-associated disease and disorder comprising administering to a patient a therapeutically effective amount of a compound of Formula (**I-1**) or (**I-2**), or a salt thereof.

In one aspect, the present disclosure refers to a compound of Formula (**I-1**):



(**I-1**)

or a salt thereof, wherein:

Cy^{1A} is substituted C_{6-10} aryl or unsubstituted or substituted 5-10 membered heteroaryl; wherein the ring atoms of the 5-10 membered heteroaryl forming Cy^{1A} consist of carbon atoms and 1, 2, or 3 heteroatoms selected from O, N and S; wherein the substituted C_{6-10} aryl or substituted 5-10 membered heteroaryl forming Cy^{1A} are substituted with 1, 2, 3, 4 or 5 substituents each independently selected from R^{Cy1A}, halogen, C_{1-6} haloalkyl, CN, OR^{a11}, S^{Ra11}, $C(O)R^{b11}$,

$C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)R^{b11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $C(=NOR^{a11})NR^{c11}R^{d11}$, $C(=NOC(O)R^{b11})NR^{c11}R^{d11}$, $C(=NR^{e11})NR^{c11}C(O)OR^{a11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo;

each R^{Cy1A} is independently selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, 5-10 membered heteroaryl, C_{3-10} cycloalkyl and 4-10 membered heterocycloalkyl, wherein the ring atoms of the 5-10 membered heteroaryl or 4-10-membered heterocycloalkyl forming R^{Cy1A} consist of carbon atoms and 1, 2, 3 or 4 heteroatoms selected from O, N and S, wherein each C_{1-6} alkyl, C_{2-6} alkenyl, or C_{2-6} alkynyl forming R^{Cy1A} is independently unsubstituted or substituted with 1, 2 or 3 substituents independently selected from halogen, CN, OR^{a11}, SR^{a11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)R^{b11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo, and wherein each C_{6-10} aryl, 5-10 membered heteroaryl, C_{3-10} cycloalkyl and 4-10 membered heterocycloalkyl forming R^{Cy1A} is independently unsubstituted or substituted with 1, 2 or 3 substituents independently selected from halogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, CN, OR^{a11}, SR^{a11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)R^{b11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo;

R^{11} is H or C_{1-6} alkyl, wherein the C_{1-6} alkyl forming R^{11} is unsubstituted or substituted by 1, 2 or 3 substituents independently selected from halogen, CN, OR^{a11}, SR^{a11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)R^{b11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo;

R^{12} is H or C_{1-6} alkyl;

A^{11} is $CR^{13}R^{15}$ or N;

each R^{13} is independently Cy^{1B}, $(CR^{13A}R^{13B})_{n3}Cy^{1B}$, ($C_{1-6}$ alkylene)Cy^{1B}, (C_{2-6} alkenylene)Cy^{1B}, (C_{2-6} alkynylene)Cy^{1B} or OCy^{1B}, wherein the C_{1-6} alkylene, C_{2-6} alkenylene, or C_{2-6} alkynylene component of R^{13} is unsubstituted or substituted by 1, 2, 3, 4 or 5 substituents each independently selected from the group consisting of halogen, CN, OR^{a11}, SR^{a11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)R^{b11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo;

each R^{14} is independently selected from H and C_{1-6} alkyl;

R^{15} is selected from H, R^{13}, C_{1-6} alkyl and OH;

a pair of R^{14} groups attached to adjacent carbon atoms, or a pairing of R^{14} and R^{15} groups attached to adjacent carbon atoms, may, independently of other occurrences of R^{14}, together be replaced a bond connecting the adjacent carbon atoms to which the pair of R^{14} groups or pairing of R^{14} and R^{15} groups is attached, such that the adjacent carbon atoms are connected by a double bond; or

a pair of R^{14} groups attached to the same carbon atom, or a pairing of R^{13} and R^{15} groups attached to the same carbon atom, may, independently of other occurrences of R^{14}, and together with the carbon atom to which the pair of R^{14} groups or pairing of R^{13} and R^{15} groups is attached together form a spiro-fused C_{3-10} cycloalkyl or 4-10 membered heterocycloalkyl ring, wherein the ring atoms of the 4-10 membered heterocycloalkyl ring formed consist of carbon atoms and 1, 2, or 3 heteroatoms selected from O, N and S, wherein the spiro-fused C_{3-10} cycloalkyl or 4-10 membered heterocycloalkyl ring formed is optionally further substituted with 1, 2 or 3 substituents independently selected from halogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, haloalkyl, CN, OR^{a11}, SR^{a11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)R^{b11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo; or

In one aspect, the present disclosure refers to use of a small molecule compound with MASP-2 inhibitory activity, in the manufacture of a medicament for the treatment of a MASP-2-associated disease or disorder selected from the group consisting of a thrombotic microangiopathy (TMA), a renal condition, an inflammatory reaction resulting from tissue or organ transplantation, an ischemia reperfusion injury, a complication associated with diabetes, a cardiovascular disease or disorder, an inflammatory gastrointestinal disorder, a pulmonary disorder, an ophthalmic disease or disorder, disseminated intravascular coagulation, graft-versus-host disease, veno-occlusive disease and diffuse alveolar hemorrhage, wherein the compound is a compound of Formula (I-1).

In some examples, the compound is a synthetic small molecule MASP-2 inhibitor.

21, or 22 of the pharmacophore elements, preferably combinations of 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 of the pharmacophore elements listed in Table 1. In some examples, the compounds may have pharmacophore elements corresponding to those listed in Table 1, wherein the (x, y, z) coordinates of the pharmacophore elements are within four standard deviations, preferably within three standard deviations, more preferably within two standard deviations and most preferably within one standard deviation as listed in Table 1.

In some examples, a compound is provided that comprises a combination of pharmacophore elements comprising:

(a) an S1 pharmacophore group comprising CA1 and N1 pharmacophore elements or CA1 and C5 pharmacophore elements; and/or

(b) an S2 pharmacophore group comprising H4 and O2 pharmacophore elements; and/or

(c) an S3 pharmacophore group comprising a C2 pharmacophore element and an N2 or H3 pharmacophore element;

wherein:

C2 and C5 are hydrophobic groups;

CA1 is an aromatic ring;

H3 and H4 are hydrogen bond donors;

N1 and N2 are positive ionizable groups; and

O2 is a hydrogen bond acceptor;

wherein C2, C5, CA1, H3, H4, N1, N2, and O2 have coordinates in the ranges given in Table 3, 4 or 5 below.

In one example, the present disclosure refers to use of a small molecule compound in the manufacture of a medicament for the treatment of MASP-2-associated diseases or disorders, wherein the compound comprises a combination of pharmacophore elements comprising:

(a) an S1 pharmacophore group comprising CA1 and N1 pharmacophore elements or CA1 and C5 pharmacophore elements;

(b) an S2 pharmacophore group comprising H4 and O2 pharmacophore elements; and

(c) an S3 pharmacophore group comprising a C2 pharmacophore element and an N2 or H3 pharmacophore element;

wherein:

e) the compound interacts via van der Waals contacts to one or more amino acid residues in the binding site of SEQ ID NO: 1, wherein the compound is not an endogenous ligand or substrate.

In one example, the present disclosure refers to use of a small molecule compound with MASP-2 inhibitory activity in the manufacture of a medicament for the treatment of a MASP-2-associated disease or disorder, wherein the compound interacts with a binding site, the interactions being one or more of (a) to (e):

a) the compound interacts via H-bonds with one or more amino acid residues in the binding site of MASP-2 (SEQ ID NO: 1);

b) the compound interacts via ionic or electrostatic interactions or hydrogen bonding in the binding site of SEQ ID NO: 1;

c) the compound interacts via a water molecule in a binding site of SEQ ID NO: 1;

d) the compound interacts via π-π interactions with one or more amino acid residues in the binding site of SEQ ID NO: 1; and/or

e) the compound interacts via van der Waals contacts to one or more amino acid residues in the binding site of SEQ ID NO: 1, wherein the compound is not an endogenous MASP-2 ligand.

In certain examples, the compound has 1, 2, 3, 4, or 5 of the interactions (a)-(e).

In another example, the present disclosure provides a method for identifying a small molecule capable of inhibiting MASP-2, comprising:

a) screening small molecule libraries using *in silico* docking for candidate small molecules that are selectively identified for their ability to target and bind to MASP-2 at a binding site of a MASP-2 model; and


b) testing/evaluating the candidate agents identified in step (a) through one or more *in vitro* assays for their ability to target and bind to a MASP-2 binding site, to thereby identify the small molecule capable of inhibiting MASP-2.

In certain examples, the candidate small molecules comprise unique chemical scaffolds as identified in step (b) and are optimized for their ability to inhibit MASP-2.

These and other aspects, objects and examples will become more apparent when read with the detailed description and figures which follow.

What is claimed is:

1. A compound of Formula (**I-1**):



(**I-1**)

or a salt thereof, wherein:

Cy^{1A} is substituted C_{6-10} aryl or unsubstituted or substituted 5-10 membered heteroaryl; wherein the ring atoms of the 5-10 membered heteroaryl forming Cy^{1A} consist of carbon atoms and 1, 2, or 3 heteroatoms selected from O, N and S; wherein the substituted C_{6-10} aryl or substituted 5-10 membered heteroaryl forming Cy^{1A} are substituted with 1, 2, 3, 4 or 5 substituents each independently selected from R^{Cy1A}, halogen, C_{1-6} haloalkyl, CN, OR^{a11}, S^{Ra11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)R^{b11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $C(=NOR^{a11})NR^{c11}R^{d11}$, $C(=NOC(O)R^{b11})NR^{c11}R^{d11}$, $C(=NR^{e11})NR^{c11}C(O)OR^{a11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo;

each R^{Cy1A} is independently selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, 5-10 membered heteroaryl, C_{3-10} cycloalkyl and 4-10 membered heterocycloalkyl, wherein the ring atoms of the 5-10 membered heteroaryl or 4-10-membered heterocycloalkyl forming R^{Cy1A} consist of carbon atoms and 1, 2, 3 or 4 heteroatoms selected from O, N and S, wherein each C_{1-6} alkyl, C_{2-6} alkenyl, or C_{2-6} alkynyl forming R^{Cy1A} is independently unsubstituted or substituted with 1, 2 or 3 substituents independently selected from halogen, CN, OR^{a11}, SR^{a11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)R^{b11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo, and wherein each C_{6-10} aryl, 5-10 membered heteroaryl, C_{3-10} cycloalkyl and 4-10 membered heterocycloalkyl forming R^{Cy1A} is independently unsubstituted or substituted with 1, 2 or 3 substituents independently selected from halogen, C_{1-6} alkyl, C_{2-6}

alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, CN, OR^{a11}, SR^{a11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)R^{b11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo;

R^{11} is H or C_{1-6} alkyl, wherein the C_{1-6} alkyl forming R^{11} is unsubstituted or substituted by 1, 2 or 3 substituents independently selected from halogen, CN, OR^{a11}, SR^{a11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)R^{b11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo;

R^{12} is H or C_{1-6} alkyl;

A^{11} is $CR^{13}R^{15}$ or N;

each R^{13} is independently Cy^{1B}, $(CR^{13A}R^{13B})_{n3}Cy^{1B}$, ($C_{1-6}$ alkylene)Cy^{1B}, (C_{2-6} alkenylene)Cy^{1B}, (C_{2-6} alkynylene)Cy^{1B} or OCy^{1B}, wherein the C_{1-6} alkylene, C_{2-6} alkenylene, or C_{2-6} alkynylene component of R^{13} is unsubstituted or substituted by 1, 2, 3, 4 or 5 substituents each independently selected from the group consisting of halogen, CN, OR^{a11}, SR^{a11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)R^{b11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo;

each R^{14} is independently selected from H and C_{1-6} alkyl;

R^{15} is selected from H, R^{13}, C_{1-6} alkyl and OH;

a pair of R^{14} groups attached to adjacent carbon atoms, or a pairing of R^{14} and R^{15} groups attached to adjacent carbon atoms, may, independently of other occurrences of R^{14}, together be replaced a bond connecting the adjacent carbon atoms to which the pair of R^{14} groups or pairing of R^{14} and R^{15} groups is attached, such that the adjacent carbon atoms are connected by a double bond; or

a pair of R^{14} groups attached to the same carbon atom, or a pairing of R^{13} and R^{15} groups attached to the same carbon atom, may, independently of other occurrences of R^{14}, and together with the carbon atom to which the pair of R^{14} groups or pairing of R^{13} and R^{15} groups is attached together form a spiro-fused C_{3-10} cycloalkyl or 4-10 membered heterocycloalkyl ring, wherein the ring atoms of the 4-10 membered heterocycloalkyl ring formed consist of

carbon atoms and 1, 2, or 3 heteroatoms selected from O, N and S, wherein the spiro-fused C_{3-10} cycloalkyl or 4-10 membered heterocycloalkyl ring formed is optionally further substituted with 1, 2 or 3 substituents independently selected from halogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, haloalkyl, CN, OR^{a11}, SR^{a11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)R^{b11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo; or

pairs of R^{14} groups attached to adjacent carbon atoms, or a pairing of R^{14} and R^{15} groups attached to adjacent carbon atoms, may, independently of other occurrences of R^{14}, together with the adjacent carbon atoms to which the pair of R^{14} groups or pairing of R^{14} and R^{15} groups is attached, form a fused C_{3-10} cycloalkyl or 4-10 membered heterocycloalkyl ring, wherein the ring atoms of the 4-10 membered heterocycloalkyl ring formed consist of carbon atoms and 1, 2, or 3 heteroatoms selected from O, N and S, wherein the fused C_{3-10} cycloalkyl or 4-10 membered heterocycloalkyl ring formed is optionally further substituted with 1, 2 or 3 substituents independently selected from halogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, haloalkyl, CN, OR^{a11}, SR^{a11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)R^{b11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo; or

a grouping of four R^{14} groups attached to two adjacent carbon atoms, or a grouping of two R^{14} , one R^{13} and one R^{15} groups attached to two adjacent carbon atoms, may, independently of other occurrences of R^{14}, together with the two adjacent carbon atoms to which the grouping of four R^{14} groups or grouping of two R^{14} , one R^{13} and one R^{15} groups are attached, form a fused C_{6-10} aryl or 5-10 membered heteroaryl, C_{3-10} cycloalkyl or 4-10 membered heterocycloalkyl ring, wherein the ring atoms of the 5-10 membered heteroaryl or 4-10 membered heterocycloalkyl ring formed consist of carbon atoms and 1, 2, or 3 heteroatoms selected from O, N and S, and wherein the fused C_{6-10} aryl or 5-10 membered heteroaryl, C_{3-10} cycloalkyl or 4-10 membered heterocycloalkyl ring formed is optionally further substituted with 1, 2 or 3 substituents independently selected from halogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, haloalkyl, CN, OR^{a11}, SR^{a11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)R^{b11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$,

$NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo;

n1 is 1 or 2;

n2 is 0, 1 or 2;

provided that the sum of n1 and n2 is 1, 2 or 3;

provided that if n1 is 1 or n2 is 0, then A^{11} is $CR^{13}R^{15}$;

n3 is 0, 1 or 2;

each R^{13A} is independently H or C_{1-6} alkyl;

each R^{13B} is independently H or C_{1-6} alkyl; or

or R^{13A} and R^{13B} attached to the same carbon atom, independently of any other R^{13A} and R^{13B} groups, together may form $–(CH_2)_{2-5}$-, thereby forming a 3-6 membered cycloalkyl ring;

Cy^{1B} is unsubstituted or substituted C_{6-10} aryl, unsubstituted or substituted 5-10 membered heteroaryl, unsubstituted or substituted C_{3-10} cycloalkyl, or unsubstituted or substituted 4-10 membered heterocycloalkyl; wherein the ring atoms of the 5-10 membered heteroaryl or 4-10 membered heterocycloalkyl forming Cy^{1B} consist of carbon atoms and 1, 2 or 3 heteroatoms selected from O, N and S; and

wherein the substituted C_{6-10} aryl, substituted 5-10 membered heteroaryl, substituted C_{3-10} cycloalkyl or substituted 4-10 membered heterocycloalkyl forming Cy^{1B} are substituted with 1, 2, 3, 4 or 5 substituents each independently selected from R^{Cy1B}, halogen, C_{1-6} haloalkyl, CN, OR^{a11}, S^{Ra11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)R^{b11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $C(=NOR^{a11})NR^{c11}R^{d11}$, $C(=NOC(O)R^{b11})NR^{c11}R^{d11}$, $C(=NR^{e11})NR^{c11}C(O)OR^{a11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo;

wherein each R^{Cy1B} is independently selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, 5-10 membered heteroaryl, C_{3-10} cycloalkyl and 4-10 membered heterocycloalkyl, wherein the ring atoms of the 5-10 membered heteroaryl or 4-10 membered heterocycloalkyl forming R^{Cy1B} consist of carbon atoms and 1, 2 or 3 heteroatoms selected from O, N and S; wherein each C_{1-6} alkyl, C_{2-6} alkenyl, or C_{2-6} alkynyl forming R^{Cy1B} is independently unsubstituted or substituted with 1, 2 or 3 substituents independently selected from halogen, CN, OR^{a11}, SR^{a11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$,

$OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)^{Rb11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo; and wherein each C_{6-10} aryl, 5-10 membered heteroaryl, C_{3-10} cycloalkyl and 4-10 membered heterocycloalkyl forming R^{Cy1B} is independently unsubstituted or substituted with 1, 2 or 3 substituents independently selected from halogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, CN, OR^{a11}, SR^{a11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)^{Rb11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo;

R^{16} is H, Cy^{1C}, C_{1-6} alkyl, C_{2-6} alkenyl, or C_{2-6} alkynyl, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, or C_{2-6} alkynyl forming R^{16} is unsubstituted or substituted by 1, 2, 3, 4 or 5 substituents selected from the group consisting of Cy^{1C}, halogen, CN, OR^{a11}, SR^{a11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)R^{b11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo, provided that no more than one of the substituents of R^{16} is Cy^{1C};

Cy^{1C} is unsubstituted or substituted C_{6-10} aryl, unsubstituted or substituted 5-10 membered heteroaryl, unsubstituted or substituted C_{3-10} cycloalkyl, or unsubstituted or substituted 4-10 membered heterocycloalkyl; wherein the ring atoms of the 5-10 membered heteroaryl or 4-10 membered heterocycloalkyl forming Cy^{1C} consist of carbon atoms and 1, 2 or 3 heteroatoms selected from O, N and S; and

wherein the substituted C_{6-10} aryl, substituted 5-10 membered heteroaryl, substituted C_{3-10} cycloalkyl or substituted 4-10 membered heterocycloalkyl forming Cy^{1C} are substituted with 1, 2, 3, 4 or 5 substituents each independently selected from R^{Cy1C}, halogen, C_{1-6} haloalkyl, CN, OR^{a11}, S^{Ra11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)R^{b11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $C(=NOR^{a11})NR^{c11}R^{d11}$, $C(=NOC(O)R^{b11})NR^{c11}R^{d11}$, $C(=NR^{e11})NR^{c11}C(O)OR^{a11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo;

wherein each R^{Cy1C} is independently selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, 5-10 membered heteroaryl, C_{3-10} cycloalkyl and 4-10 membered

heterocycloalkyl, wherein the ring atoms of the 5-10 membered heteroaryl or 4-10 membered heterocycloalkyl forming R^{Cy1C} consist of carbon atoms and 1, 2 or 3 heteroatoms selected from O, N and S; wherein each C_{1-6} alkyl, C_{2-6} alkenyl, or C_{2-6} alkynyl forming R^{Cy1C} is independently unsubstituted or substituted with 1, 2 or 3 substituents independently selected from halogen, CN, OR^{a11}, SR^{a11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)^{Rb11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo; and wherein each C_{6-10} aryl, 5-10 membered heteroaryl, C_{3-10} cycloalkyl and 4-10 membered heterocycloalkyl forming R^{Cy1C} is independently unsubstituted or substituted with 1, 2 or 3 substituents independently selected from halogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, CN, OR^{a11}, SR^{a11}, $C(O)R^{b11}$, $C(O)NR^{c11}R^{d11}$, $C(O)OR^{a11}$, $OC(O)R^{b11}$, $OC(O)NR^{c11}R^{d11}$, $NR^{c11}R^{d11}$, $NR^{c11}C(O)^{Rb11}$, $NR^{c11}C(O)NR^{c11}R^{d11}$, $NR^{c11}C(O)OR^{a11}$, $C(=NR^{e11})NR^{c11}R^{d11}$, $NR^{c11}C(=NR^{e11})NR^{c11}R^{d11}$, $S(O)R^{b11}$, $S(O)NR^{c11}R^{d11}$, $S(O)_2R^{b11}$, $NR^{c11}S(O)_2R^{b11}$, $S(O)_2NR^{c11}R^{d11}$ and oxo;

R^{a11}, R^{b11}, R^{c11} and R^{d11} are each independently selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-7} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl-C_{1-3} alkyl, 5-10 membered heteroaryl-C_{1-3} alkyl, C_{3-7} cycloalkyl-C_{1-3} alkyl and 4-10 membered heterocycloalkyl-C_{1-3} alkyl, wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-7} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl-C_{1-3} alkyl, 5-10 membered heteroaryl-C_{1-3} alkyl, C_{3-7} cycloalkyl-C_{1-3} alkyl and 4-10 membered heterocycloalkyl-C_{1-3} alkyl forming R^{a11}, R^{b11}, R^{c11} and R^{d11} are each optionally substituted with 1, 2, 3, 4 or 5 substituents independently selected from C_{1-6} alkyl, halo, CN, OR^{a12}, SR^{a12}, $C(O)R^{b12}$, $C(O)NR^{c12}R^{d12}$, $C(O)OR^{a12}$, $OC(O)R^{b12}$, $OC(O)NR^{c12}R^{d12}$, $NR^{c12}R^{d12}$, $NR^{c12}C(O)R^{b12}$, $NR^{c12}C(O)NR^{c12}R^{d12}$, $NR^{c12}C(O)OR^{a12}$, $C(=NR^{e12})NR^{c12}R^{d12}$, $NR^{c12}C(=NR^{e12})NR^{c12}R^{d12}$, $S(O)R^{b12}$, $S(O)NR^{c12}R^{d12}$, $S(O)_2R^{b12}$, $NR^{c12}S(O)_2R^{b12}$, $S(O)_2NR^{c12}R^{d12}$ and oxo;

or R^{c11} and R^{d11} attached to the same N atom, together with the N atom to which they are both attached, form a 4-, 5-, 6- or 7-membered heterocycloalkyl group or 5-membered heteroaryl group, each optionally substituted with 1, 2 or 3 substituents independently selected from C_{1-6} alkyl, halo, CN, OR^{a12}, SR^{a12}, $C(O)R^{b12}$, $C(O)NR^{c12}R^{d12}$, $C(O)OR^{a12}$, $OC(O)R^{b12}$, $OC(O)NR^{c12}R^{d12}$, $NR^{c12}R^{d12}$, $NR^{c12}C(O)R^{b12}$, $NR^{c12}C(O)NR^{c12}R^{d12}$, $NR^{c12}C(O)OR^{a12}$,

$C(=NR^{e12})NR^{c12}R^{d12}$, $NR^{c12}C(=NR^{e12})NR^{c12}R^{d12}$, $S(O)R^{b12}$, $S(O)NR^{c12}R^{d12}$, $S(O)_2R^{b12}$, $NR^{c12}S(O)_2R^{b12}$, $S(O)_2NR^{c12}R^{d12}$ and oxo;

R^{a12}, R^{b12}, R^{c12} and R^{d12} are each independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl-C_{1-3} alkyl, 5-6 membered heteroaryl-C_{1-3} alkyl, C_{3-7} cycloalkyl-C_{1-3} alkyl and 4-7 membered heterocycloalkyl-C_{1-3} alkyl, wherein said C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl-C_{1-3} alkyl, 5-6 membered heteroaryl-C_{1-3} alkyl, C_{3-7} cycloalkyl-C_{1-3} alkyl and 4-7 membered heterocycloalkyl-C_{1-3} alkyl forming R^{a12}, R^{b12}, R^{c12} and R^{d12} are each optionally substituted with 1, 2 or 3 substituents independently selected from OH, CN, amino, NH(C_{1-6} alkyl), N(C_{1-6} alkyl)$_2$, halo, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy and oxo;


or R^{c12} and R^{d12} attached to the same N atom, together with the N atom to which they are both attached, form a 4-, 5-, 6- or 7-membered heterocycloalkyl group or 5-membered heteroaryl group, each of which is unsubstituted or substituted with 1, 2 or 3 substituents independently selected from OH, CN, amino, NH(C_{1-6} alkyl), N(C_{1-6} alkyl)$_2$, halo, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy and oxo; and

R^{e11} and Re^{12} are each, independently, H, CN or NO_2.

2. Use of a small molecule compound with MASP-2 inhibitory activity, in the manufacture of a medicament for the treatment of a MASP-2-associated disease or disorder selected from the group consisting of a thrombotic microangiopathy (TMA), a renal condition, an inflammatory reaction resulting from tissue or organ transplantation, an ischemia reperfusion injury, a complication associated with diabetes, a cardiovascular disease or disorder, an inflammatory gastrointestinal disorder, a pulmonary disorder, an ophthalmic disease or disorder, disseminated intravascular coagulation, graft-versus-host disease, veno-occlusive disease and diffuse alveolar hemorrhage,

wherein the compound is a compound of Formula (I-1).


3. The compound or salt thereof of claim 1, wherein the compound is of Formula (**I-2**):




(**I-2**).

4. The compound or salt thereof of claim 1, wherein Cy^{1A} is substituted with at least one OR^{a11}, $C(=NR^{e11})NR^{c11}R^{d11}$, $C(=NOR^{a11})NR^{c11}R^{d11}$, $C(=NOC(O)R^{b11})NR^{c11}R^{d11}$, or $C(=NR^{e11})NR^{c11}C(O)OR^{a11}$.


5. The compound or salt thereof of claim 1, wherein Cy^{1A} is of any one of the following formulae:








wherein R^{Cy1A} is C_{1-6} alkyl and each R^{a11} and $R^{b11}$is independently H or C_{1-6} alkyl.


6. The compound or salt thereof of claim 1, wherein Cy^{1A} is of any one of the following formulae:
















wherein each R^{Cy1A} is C_{1-6} alkyl.

7. The compound or salt thereof of claim 1, wherein R^{11} is methyl.


8. The compound or salt thereof of claim 1, wherein the compound is according to any one of the following Formulae (**I-3**) to (**I-9**):




(**I-3**),




(**I-4**),




(**I-5**),




(**I-6**),



(**I-7**),




(**I-8**), or




(**I-9**).


9. The compound or salt thereof of claim 1, wherein the compound is according to any one of the following Formulae (**I-9a**) to (**I-9z**):




(**I-9a**),




(**I-9b**),




(**I-9c**),




(**I-9d**),




(**I-9e**),




(**I-9f**),




(**I-9g**),




(**I-9h**),




(**I-9i**),




(**I-9j**),




(**I-9k**),




(**I-9l**),




(**I-9m**),




(**I-9n**),




(**I-9o**),




(**I-9p**),




(**I-9q**),




(**I-9r**),




(**I-9s**),




(**I-9t**),




(**I-9u**),




(**I-9v**),




(**I-9w**),



(**I-9x**),



(**I-9y**), or




(**I-9z**).


10. The compound or salt thereof of claim 9, wherein R^{13} is Cy^{1B}, CH_2Cy^{1B}, $CH_2CH_2Cy^{1B}$, or OCy^{1B}, and wherein Cy^{1B} is substituted or unsubstituted C_{6-10} aryl or substituted or unsubstituted 5-10 membered heteroaryl.


11. The compound or salt thereof of claim 10, wherein Cy^{1B} is substituted with 1, 2, 3, 4 or 5 substituents each independently selected from R^{Cy1B}, halogen, and C_{1-6} haloalkyl;


wherein each R^{Cy1B} is independently selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl and C_{6-10} aryl or 5-10 membered heteroaryl, wherein each C_{6-10} aryl or 5-10 membered heteroaryl forming R^{Cy1B} is unsubstituted or substituted with 1, 2 or 3 substituents independently selected from halogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and haloalkyl.


12. The compound or salt thereof of claim 1, wherein R^{13} is a group selected from groups of the following formulae:






R
Cy1B
R
R
R



Cl
F



Cl
F



Cl
OMe



Cl
OMe



F



F



Cl
OCF3



Cl
OCF3



Br



Br



Br



Br



Br



Br



Me



Me



Me



Me



Me



Me



CF3



CF3



CF3



CF3



CF3



CF3



OMe



OMe



OMe



OMe



OMe



OMe



S



S



Cl
S



Br
S



Cl
Cl



Cl
Cl



Br
Cl



Br
Cl






















Cl
Br



Cl
Br






CF3
CF3



or ,

wherein R^{Cy1B} is C_{1-6} alkyl.

13. The compound or salt thereof of claim 1, wherein each R^{14} is hydrogen.

14. The compound or salt thereof of claim 1, wherein R^{15} is hydrogen.

15. The compound or salt thereof of claim 1, wherein R^{16} is hydrogen, unsubstituted or substituted C_{1-6} alkyl, unsubstituted or substituted C_{2-6} alkenyl, or unsubstituted or substituted C_{2-6} alkynyl.