
***Compounds and Methods for Inhibiting the Interaction
of BCL Proteins with Binding Partners***

Related Applications

5 This application claims the benefit of priority to United States Provisional Patent Application serial number 60/838,987, filed August 21, 2006.

Background of the Invention

 Apoptosis, or programmed cell death, is important for normal embryological/anatomical development, host defense and suppression of oncogenesis.

10 Faulty regulation of apoptosis has been implicated in cancer and in many other human diseases which result from an imbalance between the process of cell division and cell death. A central check point of apoptosis is the regulation of cytochrome c release from mitochondria. Cytochrome c release is regulated, in part, by Bcl-2 family members. The Bcl-2 family of proteins includes both anti-apoptotic molecules, such as Bcl-2 and Bcl-XL,

15 and pro-apoptotic molecules, such as Bax, Bak, Bid and Bad. Bcl-2 contributes to cancer cell progression by preventing normal cell turnover caused by physiological cell-death mechanisms. Over-expression of Bcl-2 has been observed in 70% of breast cancer and many other forms of cancer.

 Various small molecules have been shown to inhibit the function of Bcl-2.

20 Nevertheless, the need exists for additional small organic molecules that bind to Bcl-2 and block its anti-apoptotic function in cancer and promote cell death in tumors.


Summary of the Invention

 The present invention generally relates to isoxazolidine compounds useful for

25 treating cancer. The isoxazolidine compounds of the invention bind to one or more Bcl proteins and block Bcl anti-apoptotic function in cancer cells and tumor tissue that express the Bcl protein.

 The present invention relates to isoxazolidine compounds (e.g., compounds of formulas 1, 2, and 3), pharmaceutical compositions, and methods of using them. In one

30 embodiment, the present invention relates to a compound of formula 1:



or a pharmaceutically acceptable salt thereof;


wherein independently for each occurrence

5 m is 0, 1, 2, or 3;

n, o, and p are independently for each occurrence 1, 2, 3, 4, or 5;

R₁ is -OH, -OC(O)R₆, -OC(O)N(R₆)(R₇), or -N(R₆)(R₇);

R₂ is -OH, -N(R₈)(R₉), -N(R)C(O)N(R₈)(R₉), or -N(R)C(O)R₁₀; or has the formula
1b;



10

1b

R₃ is alkyl, halide, alkoxy, (cycloalkyl)alkoxy, aralkyloxy, or -O(CH₂)₂-N(R₁₅)(R₁₆);

R₄ is alkyl, alkoxy, hydroxyalkyl, alkoxyalkyl, halide, nitro, amino, acyl, amido,
acylamino, aminoalkyl, acylaminoalkyl, acylaminoalkylamino, sulfonylaminoalkylamino,
15 carboxylate, or -N=C(N(R)₂)₂;

R₅ is -OH or -N(R₁₇)(R₁₈);

R₆ and R₇ are independently for each occurrence H, alkyl, aralkyl, heteroaralkyl, or
-[C(R₁₅)(R₁₆)]_n-R₁₉;

R₈ and R₉ are independently for each occurrence H, alkyl, aralkyl, or heteraralkyl;

20 R₁₀ is alkyl, haloalkyl, or -[C(R₁₅)(R₁₆)]_o-COOR;

R, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, and R₁₆ are independently for each occurrence H or alkyl;

R₁₇ and R₁₈ are independently for each occurrence H, alkyl, aralkyl, heteroaralkyl,
alkoxy, or -[C(R₁₉)(R₂₀)]_p-R₂₁;

R₁₉ and R₂₀ are independently for each occurrence H, hydroxy, alkyl, alkoxy, amino, aminoalkyl, acylamino, sulfonylamino, aryl, aralkyl, cycloalkyl, heterocyclyl, heterocyclalkyl, heteroaryl, or heteroaralkyl;

5 R₂₁ is independently for each occurrence H, alkyl, aryl, heteroaryl, heterocyclyl, heterocyclalkyl, alkoxy, alkylsulfonyl, arylsulfonyl, alkylsulfonamido, arylsulfonamido, amino, amido, or carboxyl;

R₂₂ independently for each occurrence is halide or alkyl;

R₂₃ is selected from the group consisting of alkyl, hydroxyalkyl, alkoxyalkyl, acyloxyalkyl, and haloalkyl;

10 R₂₄ is selected from the group consisting of alkyl, hydroxyalkyl, alkoxyalkyl, acyloxyalkyl, and haloalkyl; and

R₂₅ is selected from the group consisting of alkyl, hydroxyalkyl, alkoxyalkyl, acyloxyalkyl, and haloalkyl.

15 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₁ is selected from the group consisting of -OH, -OC(O)Me, -OC(O)(CH₂)₂Ph, -OC(O)CH₂CHMe₂, -OC(O)NHMe, -OC(O)NMe₂, -OC(O)NHCH₂(4-(OH)-Ph), -OC(O)NHPh, -OC(O)NHCH₂Ph, -OC(O)NH(CH₂)₄Ph, -OC(O)NH(CH₂)₂Ph, -OC(O)NH(CH₂)₂Me, -OC(O)NH(CH₂)₂NMe₂,
20 -OC(O)NH(CH₂)₂NHC(O)Me, -OC(O)NH(CH₂)₂CHMe₂, -NHMe, -NH(CH₂)₂Ph, -NHC(O)Me, and -NH(CH₂)₂NMe₂.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₁ is -OH.

25 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂ is -OH, -OC(O)Me, -OCH₂CO₂H, -OCH₂CO₂Et, -N₃, -N=C(NMe₂)₂, -NH₂, -NMe₂, -NHC(O)Me, -NHC(O)CF₃, -NHC(O)Ph, -NHC(O)NHPh, -NHC(O)CH₂CH₂CO₂H, -NHC(O)CH₂CH₂CO₂Me, -NHCH₂Ph, -NHCH₂(4-pyridyl), -NHCH₂(2-pyridyl), -NHCH₂(4-(CO₂H)Ph), -NHCH₂(3-(CO₂H)Ph), -NH₂Et, -NHCHMe₂, -NHCH₂CHMe₂, -N(CH₂CHMe₂)₂, -
30 NHCH₂(cyclopropyl), or -NHC(O)CH₂CH₂NMe₂.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_2 is -OH or -NH₂.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_2 is -NH₂.

5 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_2 is -OH.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_3 is -OMe, -OEt, -OCH₂(cyclopropyl), F, -O(CH₂)₂NMe₂, -O(CH₂)₂(4-morpholino), -OCH₂(4-(MeO)Ph), or -
10 OCH₂(2-pyridyl).

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_3 is -OMe.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_4 is -NMe₂, -NEt₂, -NHEt, -
15 NHCH₂CHMe₂, -N(Me)CH₂CHMe₂, -N(Me)CH₂CH₂NHC(O)Me, -N(Me)CH₂CH₂NHS(O)₂Me, -N(Me)CH₂CH₂NHS(O)₂CF₃, -NHCH₂CH₂NMe₂, -NHCH₂CH₂NMeCH₂CH₂Cl, -NHCH₂CH₂OH, -N(Me)CH₂CH₂OH, -N(CH₂CH₂OH)₂, -N(Me)CH₂CO₂H, -N(Me)CH₂C(O)NH₂, -N(Me)CH₂C(O)NHMe, -N(Me)CH₂C(O)NMe₂, -NHC(O)Me, -NHC(O)CHMe₂, 1-pyrrolidinyl, 1-piperidinyl, 4-morpholino, (R)-2-(hydroxymethyl)-1-pyrrolidinyl, (S)-2-(hydroxymethyl)-1-pyrrolidinyl, (R)-2-(C(O)NMe₂)-1-pyrrolidinyl, (S)-2-(C(O)NMe₂)-1-pyrrolidinyl, -NH₂, -NO₂, Br, Cl, F, -C(O)Me, -C(O)NMe₂, -C(O)NH₂, -CO₂H, -CHO, -CH₂OH, -CH(Me)OH, -CH₂NH₂, -CH₂NHMe, -CH₂NMe₂, -CH₂NHC(O)Me, -CF₃, or tert-butyl.

25 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_4 is amino.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_4 is -NMe₂, -N(Me)CH₂CH₂OH, -N(Me)CH₂C(O)NMe₂, -N(Me)CH₂C(O)NHMe, (R)-2-(hydroxymethyl)-1-pyrrolidinyl, or (R)-2-(C(O)NMe₂)-1-pyrrolidinyl.

30 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_4 is -NMe₂.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{22} is selected from the group consisting of F, Cl, and tert-butyl; and m is 0 or 1.

5 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{22} is F; and m is 0 or 1.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{22} is selected from the group consisting of F, Cl, and tert-butyl; and m is 1.

10 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{22} is F; and m is 1.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{23} is selected from the group consisting of methyl, hydroxymethyl, alkoxymethyl, acyloxymethyl, and halomethyl.

15 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{24} is selected from the group consisting of methyl, hydroxymethyl, alkoxymethyl, acyloxymethyl, and halomethyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{23} is methyl.

20 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{24} is methyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{23} is methyl; and R_{24} is methyl.

25 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{25} is selected from the group consisting of methyl, hydroxymethyl, alkoxymethyl, acyloxymethyl, and halomethyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{25} is methyl.

30 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{23} is methyl; R_{24} is methyl; and R_{25} is methyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; and R_2 is -OH or -NH₂.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; and
5 R_3 is -OMe.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is -OMe; and R_4 is amino.

In certain embodiments, the present invention relates to the aforementioned
10 compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is -OMe; and R_4 is -NMe₂, -N(Me)CH₂CH₂OH, -N(Me)CH₂C(O)NMe₂, -N(Me)CH₂C(O)NHMe, (R)-2-(hydroxymethyl)-1-pyrrolidinyl, or (R)-2-(C(O)NMe₂)-1-pyrrolidinyl.

In certain embodiments, the present invention relates to the aforementioned
15 compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is -OMe; and R_4 is -NMe₂.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_{23} is methyl; R_{24} is methyl; and R_{25} is methyl.


20 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is -OMe; R_{23} is methyl; R_{24} is methyl; and R_{25} is methyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is
25 -OMe; R_4 is amino; R_{23} is methyl; R_{24} is methyl; and R_{25} is methyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is -OMe; R_4 is -NMe₂, -N(Me)CH₂CH₂OH, -N(Me)CH₂C(O)NMe₂, -N(Me)CH₂C(O)NHMe, (R)-2-(hydroxymethyl)-1-pyrrolidinyl, or (R)-2-(C(O)NMe₂)-1-pyrrolidinyl; R_{23} is methyl;
30 R_{24} is methyl; and R_{25} is methyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is -OMe; R_4 is -NMe₂; R_{23} is methyl; R_{24} is methyl; and R_{25} is methyl.

5 In another embodiment, the present invention relates to a compound of formula 2:



or a pharmaceutically acceptable salt thereof;


wherein independently for each occurrence

10 m is 0, 1, 2, or 3;

n , o , and p are independently for each occurrence 1, 2, 3, 4, or 5;

R_1 is -OH, -OC(O) R_6 , -OC(O) $N(R_6)(R_7)$, or - $N(R_6)(R_7)$;

R_2 is -OH, - $N(R_8)(R_9)$, - $N(R)C(O)N(R_8)(R_9)$, or - $N(R)C(O)R_{10}$; or has the formula 1b;



15

1b

R_3 is alkyl, halide, alkoxy, (cycloalkyl)alkoxy, aralkyloxy, or -O(CH₂)₂- $N(R_{15})(R_{16})$;

R_4 is alkyl, alkoxy, hydroxyalkyl, alkoxyalkyl, halide, nitro, amino, acyl, amido, acylamino, aminoalkyl, acylaminoalkyl, acylaminoalkylamino, sulfonylaminoalkylamino, 20 carboxylate, or -N=C($N(R)_2$)₂;

R_5 is -OH or - $N(R_{17})(R_{18})$;

R_6 and R_7 are independently for each occurrence H, alkyl, aralkyl, heteroaralkyl, or -[C($R_{15})(R_{16})$] _{n} - R_{19} ;

R_8 and R_9 are independently for each occurrence H, alkyl, aralkyl, or heteraralkyl;

R₁₀ is alkyl, haloalkyl, or -[C(R₁₅)(R₁₆)]_o-COOR;

R, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, and R₁₆ are independently for each occurrence H or alkyl;

R₁₇ and R₁₈ are independently for each occurrence H, alkyl, aralkyl, heteroaralkyl, alkoxy, or -[C(R₁₉)(R₂₀)]_p-R₂₁;

5 R₁₉ and R₂₀ are independently for each occurrence H, hydroxy, alkyl, alkoxy, amino, aminoalkyl, acylamino, sulfonylamino, aryl, aralkyl, cycloalkyl, heterocyclyl, heterocyclalkyl, heteroaryl, or heteroaralkyl;

10 R₂₁ is independently for each occurrence H, alkyl, aryl, heteroaryl, heterocyclyl, heterocyclalkyl, alkoxy, alkylsulfonyl, arylsulfonyl, alkylsulfonamido, arylsulfonamido, amino, amido, or carboxyl;

R₂₂ independently for each occurrence is halide or alkyl;

R₂₃ is selected from the group consisting of alkyl, hydroxyalkyl, alkoxyalkyl, acyloxyalkyl, and haloalkyl;

15 R₂₄ is selected from the group consisting of alkyl, hydroxyalkyl, alkoxyalkyl, acyloxyalkyl, and haloalkyl; and

R₂₅ is selected from the group consisting of alkyl, hydroxyalkyl, alkoxyalkyl, acyloxyalkyl, and haloalkyl.

20 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₁ is selected from the group consisting of -OH, -OC(O)Me, -OC(O)(CH₂)₂Ph, -OC(O)CH₂CHMe₂, -OC(O)NHMe, -OC(O)NMe₂, -OC(O)NHCH₂(4-(OH)-Ph), -OC(O)NHPh, -OC(O)NHCH₂Ph, -OC(O)NH(CH₂)₄Ph, -OC(O)NH(CH₂)₂Ph, -OC(O)NH(CH₂)₂Me, -OC(O)NH(CH₂)₂NMe₂, -OC(O)NH(CH₂)₂NHC(O)Me, -OC(O)NH(CH₂)₂CHMe₂, -NHMe, -NH(CH₂)₂Ph, -NHC(O)Me, and -NH(CH₂)₂NMe₂.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₁ is -OH.

30 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂ is -OH, -OC(O)Me, -OCH₂CO₂H, -OCH₂CO₂Et, -N₃, -N=C(NMe₂)₂, -NH₂, -NMe₂, -NHC(O)Me, -NHC(O)CF₃,

-NHC(O)Ph, -NHC(O)NHPH, -NHC(O)CH₂CH₂CO₂H, -NHC(O)CH₂CH₂CO₂Me, -NHCH₂Ph, -NHCH₂(4-pyridyl), -NHCH₂(2-pyridyl), -NHCH₂(4-(CO₂H)Ph), -NHCH₂(3-(CO₂H)Ph), -NHEt, -NHCHMe₂, -NHCH₂CHMe₂, -N(CH₂CHMe₂)₂, -NHCH₂(cyclopropyl), or -NHC(O)CH₂CH₂NMe₂.

5 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂ is -OH or -NH₂.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂ is -NH₂.

10 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂ is -OH.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₃ is -OMe, -OEt, -OCH₂(cyclopropyl), F, -O(CH₂)₂NMe₂, -O(CH₂)₂(4-morpholino), -OCH₂(4-(MeO)Ph), or -OCH₂(2-pyridyl).

15 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₃ is -OMe.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₄ is -NMe₂, -NEt₂, -NHEt, -NHCH₂CHMe₂, -N(Me)CH₂CHMe₂, -N(Me)CH₂CH₂NHC(O)Me, -
 20 N(Me)CH₂CH₂NHS(O)₂Me, -N(Me)CH₂CH₂NHS(O)₂CF₃, -NHCH₂CH₂NMe₂, -NHCH₂CH₂NMeCH₂CH₂Cl, -NHCH₂CH₂OH, -N(Me)CH₂CH₂OH, -N(CH₂CH₂OH)₂, -N(Me)CH₂CO₂H, -N(Me)CH₂C(O)NH₂, -N(Me)CH₂C(O)NHMe, -N(Me)CH₂C(O)NMe₂, -NHC(O)Me, -NHC(O)CHMe₂, 1-pyrrolidinyl, 1-piperidinyl, 4-morpholino, (R)-2-(hydroxymethyl)-1-pyrrolidinyl, (S)-2-(hydroxymethyl)-1-pyrrolidinyl, (R)-2-(C(O)NMe₂)-
 25 1-pyrrolidinyl, (S)-2-(C(O)NMe₂)-1-pyrrolidinyl, -NH₂, -NO₂, Br, Cl, F, -C(O)Me, -C(O)NMe₂, -C(O)NH₂, -CO₂H, -CHO, -CH₂OH, -CH(Me)OH, -CH₂NH₂, -CH₂NHMe, -CH₂NMe₂, -CH₂NHC(O)Me, -CF₃, or tert-butyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₄ is amino.

30 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₄ is -NMe₂, -N(Me)CH₂CH₂OH, -

$N(Me)CH_2C(O)NMe_2$, $-N(Me)CH_2C(O)NHMe$, (R)-2-(hydroxymethyl)-1-pyrrolidinyl, or (R)-2-($C(O)NMe_2$)-1-pyrrolidinyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_4 is $-NMe_2$.

- 5 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{22} is selected from the group consisting of F, Cl, and tert-butyl; and m is 0 or 1.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{22} is F; and m is 0 or 1.

- 10 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{22} is selected from the group consisting of F, Cl, and tert-butyl; and m is 1.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{22} is F; and m is 1.

- 15 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{23} is selected from the group consisting of methyl, hydroxymethyl, alkoxymethyl, acyloxymethyl, and halomethyl.

- In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{24} is selected from the group
20 consisting of methyl, hydroxymethyl, alkoxymethyl, acyloxymethyl, and halomethyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{23} is methyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{24} is methyl.

- 25 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{23} is methyl; and R_{24} is methyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{25} is selected from the group consisting of methyl, hydroxymethyl, alkoxymethyl, acyloxymethyl, and halomethyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂₅ is methyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂₃ is methyl; R₂₄ is methyl; and
5 R₂₅ is methyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₁ is -OH; and R₂ is -OH or -NH₂.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₁ is -OH; R₂ is -OH or -NH₂; and
10 R₃ is -OMe.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₁ is -OH; R₂ is -OH or -NH₂; R₃ is -OMe; and R₄ is amino.

In certain embodiments, the present invention relates to the aforementioned
15 compound and any of the attendant definitions, wherein R₁ is -OH; R₂ is -OH or -NH₂; R₃ is -OMe; and R₄ is -NMe₂, -N(Me)CH₂CH₂OH, -N(Me)CH₂C(O)NMe₂, -N(Me)CH₂C(O)NHMe, (R)-2-(hydroxymethyl)-1-pyrrolidinyl, or (R)-2-(C(O)NMe₂)-1-pyrrolidinyl.

In certain embodiments, the present invention relates to the aforementioned
20 compound and any of the attendant definitions, wherein R₁ is -OH; R₂ is -OH or -NH₂; R₃ is -OMe; and R₄ is -NMe₂.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₁ is -OH; R₂ is -OH or -NH₂; R₂₃ is methyl; R₂₄ is methyl; and R₂₅ is methyl.


In certain embodiments, the present invention relates to the aforementioned
25 compound and any of the attendant definitions, wherein R₁ is -OH; R₂ is -OH or -NH₂; R₃ is -OMe; R₂₃ is methyl; R₂₄ is methyl; and R₂₅ is methyl.

In certain embodiments, the present invention relates to the aforementioned
30 compound and any of the attendant definitions, wherein R₁ is -OH; R₂ is -OH or -NH₂; R₃ is -OMe; R₄ is amino; R₂₃ is methyl; R₂₄ is methyl; and R₂₅ is methyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is -OMe; R_4 is -NMe₂, -N(Me)CH₂CH₂OH, -N(Me)CH₂C(O)NMe₂, -N(Me)CH₂C(O)NHMe, (R)-2-(hydroxymethyl)-1-pyrrolidinyl, or (R)-2-(C(O)NMe₂)-1-pyrrolidinyl; R_{23} is methyl; R_{24} is methyl; and R_{25} is methyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is -OMe; R_4 is -NMe₂; R_{23} is methyl; R_{24} is methyl; and R_{25} is methyl.

In an additional embodiment, the present invention relates to a compound of formula 3:



or a pharmaceutically acceptable salt thereof;


wherein independently for each occurrence

m is 0, 1, 2, or 3;

n , o , and p are independently for each occurrence 1, 2, 3, 4, or 5;

R_1 is -OH, -OC(O) R_6 , -OC(O)N(R_6)(R_7), or -N(R_6)(R_7);

R_2 is -OH, -N(R_8)(R_9), -N(R)C(O)N(R_8)(R_9), or -N(R)C(O) R_{10} ; or has the formula **1b**;



1b

R_3 is alkyl, halide, alkoxy, (cycloalkyl)alkoxy, aralkyloxy, or -O(CH₂)₂-N(R_{15})(R_{16});

R_4 is alkyl, alkoxy, hydroxyalkyl, alkoxyalkyl, halide, nitro, amino, acyl, amido, acylamino, aminoalkyl, acylaminoalkyl, acylaminoalkylamino, sulfonylaminoalkylamino, carboxylate, or $-N=C(N(R)_2)_2$;

R_5 is $-OH$ or $-N(R_{17})(R_{18})$;

5 R_6 and R_7 are independently for each occurrence H, alkyl, aralkyl, heteroaralkyl, or $-[C(R_{15})(R_{16})]_n-R_{19}$;

R_8 and R_9 are independently for each occurrence H, alkyl, aralkyl, or heteroaralkyl;

R_{10} is alkyl, haloalkyl, or $-[C(R_{15})(R_{16})]_o-COOR$;

R , R_{11} , R_{12} , R_{13} , R_{14} , R_{15} , and R_{16} are independently for each occurrence H or alkyl;

10 R_{17} and R_{18} are independently for each occurrence H, alkyl, aralkyl, heteroaralkyl, alkoxy, or $-[C(R_{19})(R_{20})]_p-R_{21}$;

R_{19} and R_{20} are independently for each occurrence H, hydroxy, alkyl, alkoxy, amino, aminoalkyl, acylamino, sulfonylamino, aryl, aralkyl, cycloalkyl, heterocyclyl, heterocyclalkyl, heteroaryl, or heteroaralkyl;

15 R_{21} is independently for each occurrence H, alkyl, aryl, heteroaryl, heterocyclyl, heterocyclalkyl, alkoxy, alkylsulfonyl, arylsulfonyl, alkylsulfonamido, arylsulfonamido, amino, amido, or carboxyl;

R_{22} independently for each occurrence is halide or alkyl;

20 R_{23} is selected from the group consisting of alkyl, hydroxyalkyl, alkoxyalkyl, acyloxyalkyl, and haloalkyl; and

R_{24} is selected from the group consisting of alkyl, hydroxyalkyl, alkoxyalkyl, acyloxyalkyl, and haloalkyl.

In certain embodiments, the present invention relates to the aforementioned
 25 compound and any of the attendant definitions, wherein R_1 is selected from the group consisting of $-OH$, $-OC(O)Me$, $-OC(O)(CH_2)_2Ph$, $-OC(O)CH_2CHMe_2$, $-OC(O)NHMe$, $-OC(O)NMe_2$, $-OC(O)NHCH_2(4-(OH)-Ph)$, $-OC(O)NHPh$, $-OC(O)NHCH_2Ph$, $-OC(O)NH(CH_2)_4Ph$, $-OC(O)NH(CH_2)_2Ph$, $-OC(O)NH(CH_2)_2Me$, $-OC(O)NH(CH_2)_2NMe_2$, $-OC(O)NH(CH_2)_2NHC(O)Me$, $-OC(O)NH(CH_2)_2CHMe_2$, $-NHMe$, $-NH(CH_2)_2Ph$, $-NHC(O)Me$, and $-NH(CH_2)_2NMe_2$.
 30

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₁ is -OH.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂ is -OH, -OC(O)Me, -

5 OCH₂CO₂H, -OCH₂CO₂Et, -N₃, -N=C(NMe₂)₂, -NH₂, -NMe₂, -NHC(O)Me, -NHC(O)CF₃, -NHC(O)Ph, -NHC(O)NHPH, -NHC(O)CH₂CH₂CO₂H, -NHC(O)CH₂CH₂CO₂Me, -NHCH₂Ph, -NHCH₂(4-pyridyl), -NHCH₂(2-pyridyl), -NHCH₂(4-(CO₂H)Ph), -NHCH₂(3-(CO₂H)Ph), -NH₂Et, -NHCHMe₂, -NHCH₂CHMe₂, -N(CH₂CHMe₂)₂, -NHCH₂(cyclopropyl), or -NHC(O)CH₂CH₂NMe₂.

10 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂ is -OH or -NH₂.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂ is -NH₂.

15 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂ is -OH.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₃ is -OMe, -OEt, -

OCH₂(cyclopropyl), F, -O(CH₂)₂NMe₂, -O(CH₂)₂(4-morpholino), -OCH₂(4-(MeO)Ph), or -OCH₂(2-pyridyl).

20 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₃ is -OMe.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₄ is -NMe₂, -NEt₂, -NH₂Et, -

NHCH₂CHMe₂, -N(Me)CH₂CHMe₂, -N(Me)CH₂CH₂NHC(O)Me, -

25 N(Me)CH₂CH₂NHS(O)₂Me, -N(Me)CH₂CH₂NHS(O)₂CF₃, -NHCH₂CH₂NMe₂, -NHCH₂CH₂NMeCH₂CH₂Cl, -NHCH₂CH₂OH, -N(Me)CH₂CH₂OH, -N(CH₂CH₂OH)₂, -N(Me)CH₂CO₂H, -N(Me)CH₂C(O)NH₂, -N(Me)CH₂C(O)NHMe, -N(Me)CH₂C(O)NMe₂, -NHC(O)Me, -NHC(O)CHMe₂, 1-pyrrolidinyl, 1-piperidinyl, 4-morpholino, (R)-2-(hydroxymethyl)-1-pyrrolidinyl, (S)-2-(hydroxymethyl)-1-pyrrolidinyl, (R)-2-(C(O)NMe₂)-1-pyrrolidinyl, (S)-2-(C(O)NMe₂)-1-pyrrolidinyl, -NH₂, -NO₂, Br, Cl, F, -C(O)Me, -

30

C(O)NMe₂, -C(O)NH₂, -CO₂H, -CHO, -CH₂OH, -CH(Me)OH, -CH₂NH₂, -CH₂NHMe, -CH₂NMe₂, -CH₂NHC(O)Me, -CF₃, or tert-butyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₄ is amino.

5 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₄ is -NMe₂, -N(Me)CH₂CH₂OH, -N(Me)CH₂C(O)NMe₂, -N(Me)CH₂C(O)NHMe, (R)-2-(hydroxymethyl)-1-pyrrolidinyl, or (R)-2-(C(O)NMe₂)-1-pyrrolidinyl.

10 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₄ is -NMe₂.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂₂ is selected from the group consisting of F, Cl, and tert-butyl; and m is 0 or 1.

15 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂₂ is F; and m is 0 or 1.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂₂ is selected from the group consisting of F, Cl, and tert-butyl; and m is 1.

20 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂₂ is F; and m is 1.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂₃ is selected from the group consisting of methyl, hydroxymethyl, alkoxymethyl, acyloxymethyl, and halomethyl.

25 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂₄ is selected from the group consisting of methyl, hydroxymethyl, alkoxymethyl, acyloxymethyl, and halomethyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂₃ is methyl.

30 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₂₄ is methyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_{23} is methyl; and R_{24} is methyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; and R_2 is -OH or -NH₂.

5 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; and R_3 is -OMe.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is
10 -OMe; and R_4 is amino.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is -OMe; and R_4 is -NMe₂, -N(Me)CH₂CH₂OH, -N(Me)CH₂C(O)NMe₂, -
15 N(Me)CH₂C(O)NHMe, (R)-2-(hydroxymethyl)-1-pyrrolidinyl, or (R)-2-(C(O)NMe₂)-1-pyrrolidinyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is -OMe; and R_4 is -NMe₂.

In certain embodiments, the present invention relates to the aforementioned
20 compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_{23} is methyl; and R_{24} is methyl.

In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is -OMe; R_{23} is methyl; and R_{24} is methyl.


25 In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is -OMe; R_4 is amino; R_{23} is methyl; and R_{24} is methyl.


In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is
30 -OMe; R_4 is -NMe₂, -N(Me)CH₂CH₂OH, -N(Me)CH₂C(O)NMe₂, -N(Me)CH₂C(O)NHMe,









(R)-2-(hydroxymethyl)-1-pyrrolidinyl, or (R)-2-(C(O)NMe₂)-1-pyrrolidinyl; R₂₃ is methyl; and R₂₄ is methyl.









In certain embodiments, the present invention relates to the aforementioned compound and any of the attendant definitions, wherein R₁ is -OH; R₂ is -OH or -NH₂; R₃ is -OMe; R₄ is -NMe₂; R₂₃ is methyl; and R₂₄ is methyl.










In another embodiment, the present invention relates to a compound selected from the group consisting of:




















5






























5



























5





































































































































































































































































































































































































In another embodiment, the present invention relates to a compound selected from the group consisting of:









Another aspect of the invention relates to a pharmaceutical composition, comprising at least one of the compounds of the present invention; and at least one pharmaceutically acceptable excipient.

The present invention also relates to a method of treating a bcl-mediated disorder, comprising the step of:

administering to a patient in need thereof a therapeutically effective amount of at least one of the compounds of the present invention.

5 In certain embodiments, the bcl-mediated disorder is cancer or neoplastic disease. For example, the bcl-mediated disorder may be acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemia, myeloblastic, promyelocytic, myelomonocytic, monocytic, erythroleukemia, chronic leukemia, chronic myelocytic (granulocytic) leukemia, chronic lymphocytic leukemia, polycythemia Vera, Hodgkin's disease, non-
10 Hodgkin's disease; multiple myeloma, Waldenstrom's macroglobulinemia, heavy chain disease, fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer,
15 ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, stadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, uterine cancer, testicular
20 tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, retinoblastoma, and endometrial cancer.

In addition, the cancer may be follicular lymphoma, diffuse large B-cell lymphoma,
25 mantle cell lymphoma, chronic lymphocytic leukemia prostate cancer, breast cancer, neuroblastoma, colorectal, endometrial, ovarian, lung cancer, hepatocellular carcinoma, multiple myeloma, head and neck or testicular cancer.

In some embodiments the cancer over-expresses a Bcl protein. In addition, the cancer can also be dependent upon a Bcl protein for growth and survival. In some
30 embodiments the Bcl protein specified may be Bcl-2 and/or Bcl-xL. In some embodiments the cancer exhibits a t(14;18) chromosomal translocation.

The present invention also relates to a method of treating a bcl-mediated disorder, comprising the step of:

co-administering to a patient in need thereof a therapeutically effective amount of at least one chemotherapeutic agent; and a therapeutically effective amount of at least one of
5 the compounds of the present invention.

The compound or compounds may be administered parenterally, intramuscularly, intravenously, subcutaneously, orally, topically or intranasally. In some embodiments the compound or compounds are administered systemically.

The patient specified above may be a mammal, primate, or human.

10

Brief Description of the Figures

Figure 1 depicts certain compounds of the invention, some of which are prophetic embodiments.

Figure 2 depicts certain compounds of the invention, some of which are prophetic
15 embodiments.

Figure 3 depicts certain compounds of the invention, some of which are prophetic embodiments.

Figure 4 depicts certain compounds of the invention, some of which are prophetic embodiments.

Figure 5 depicts certain compounds of the invention, some of which are prophetic
20 embodiments.

Figure 6 depicts certain compounds of the invention, some of which are prophetic embodiments.

25

Detailed Description of the Invention

Definitions

For convenience, certain terms employed in the specification, examples, and appended claims are collected here.

The terms "co-administration" and "co-administering" refer to both concurrent administration (administration of two or more therapeutic agents at the same time) and time varied administration (administration of one or more therapeutic agents at a time different from that of the administration of an additional therapeutic agent or agents), as long as the therapeutic agents are present in the patient to some extent at the same time.

The term "heteroatom" as used herein means an atom of any element other than carbon or hydrogen. Preferred heteroatoms are boron, nitrogen, oxygen, phosphorus, sulfur and selenium.

The term "alkyl" refers to the radical of saturated aliphatic groups, including straight-chain alkyl groups, branched-chain alkyl groups, cycloalkyl (alicyclic) groups, alkyl substituted cycloalkyl groups, and cycloalkyl substituted alkyl groups. In preferred embodiments, a straight chain or branched chain alkyl has 30 or fewer carbon atoms in its backbone (e.g., C₁-C₃₀ for straight chain, C₃-C₃₀ for branched chain), and more preferably 20 or fewer. Likewise, preferred cycloalkyls have from 3-10 carbon atoms in their ring structure, and more preferably have 5, 6 or 7 carbons in the ring structure.

Unless the number of carbons is otherwise specified, "lower alkyl" as used herein means an alkyl group, as defined above, but having from one to ten carbons, more preferably from one to six carbon atoms in its backbone structure. Likewise, "lower alkenyl" and "lower alkynyl" have similar chain lengths. Preferred alkyl groups are lower alkyls. In preferred embodiments, a substituent designated herein as alkyl is a lower alkyl.

The term "haloalkyl", as used herein, refers to an alkyl group where anywhere from 1 to all hydrogens have been replaced with a halide. A "perhaloalkyl" is where all of the hydrogens have been replaced with a halide.

The term "aralkyl", as used herein, refers to an alkyl group substituted with an aryl group (e.g., an aromatic or heteroaromatic group).

The terms "alkenyl" and "alkynyl" refer to unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double or triple bond respectively.

The term "aryl" as used herein includes 5-, 6- and 7-membered single-ring aromatic groups that may include from zero to four heteroatoms, for example, benzene, anthracene, naphthalene, pyrene, pyrrole, furan, thiophene, imidazole, oxazole, thiazole, triazole,

pyrazole, pyridine, pyrazine, pyridazine and pyrimidine, and the like. Those aryl groups having heteroatoms in the ring structure may also be referred to as "aryl heterocycles" or "heteroaromatics." The aromatic ring may be substituted at one or more ring positions with such substituents as described above, for example, halogen, azide, alkyl, aralkyl, alkenyl, alkynyl, cycloalkyl, hydroxyl, alkoxyl, amino, nitro, sulfhydryl, imino, amido, phosphonate, phosphinate, carbonyl, carboxyl, silyl, ether, alkylthio, sulfonyl, sulfonamido, ketone, aldehyde, ester, heterocyclyl, aromatic or heteroaromatic moieties, -CF₃, -CN, or the like. The term "aryl" also includes polycyclic ring systems having two or more cyclic rings in which two or more carbons are common to two adjoining rings (the rings are "fused rings") wherein at least one of the rings is aromatic, e.g., the other cyclic rings may be cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and/or heterocyclyls.

The terms *ortho*, *meta* and *para* apply to 1,2-, 1,3- and 1,4-disubstituted benzenes, respectively. For example, the names 1,2-dimethylbenzene and *ortho*-dimethylbenzene are synonymous.


The terms "heterocyclyl" or "heterocyclic group" refer to 3- to 10-membered ring structures, more preferably 3- to 7-membered rings, whose ring structures include one to four heteroatoms. Heterocycles can also be polycycles. Heterocyclyl groups include, for example, thiophene, thianthrene, furan, pyran, isobenzofuran, chromene, xanthene, phenoxathiin, pyrrole, imidazole, pyrazole, isothiazole, isoxazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, indazole, purine, quinolizine, isoquinoline, quinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, pyrimidine, phenanthroline, phenazine, phenarsazine, phenothiazine, furazan, phenoxazine, pyrrolidine, oxolane, thiolane, oxazole, piperidine, piperazine, morpholine, lactones, lactams such as azetidinones and pyrrolidinones, sultams, sultones, and the like. The heterocyclic ring may be substituted at one or more positions with such substituents as described above, as for example, halogen, alkyl, aralkyl, alkenyl, alkynyl, cycloalkyl, hydroxyl, amino, nitro, sulfhydryl, imino, amido, phosphonate, phosphinate, carbonyl, carboxyl, silyl, ether, alkylthio, sulfonyl, ketone, aldehyde, ester, a heterocyclyl, an aromatic or heteroaromatic moiety, -CF₃, -CN, or the like.

The terms "polycyclyl" or "polycyclic group" refer to two or more rings (e.g., cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and/or heterocyclyls) in which two or more

carbons are common to two adjoining rings, e.g., the rings are "fused rings". Rings that are joined through non-adjacent atoms are termed "bridged" rings. Each of the rings of the polycycle may be substituted with such substituents as described above, as for example, halogen, alkyl, aralkyl, alkenyl, alkynyl, cycloalkyl, hydroxyl, amino, nitro, sulfhydryl, imino, amido, phosphonate, phosphinate, carbonyl, carboxyl, silyl, ether, alkylthio, sulfonyl, ketone, aldehyde, ester, a heterocyclyl, an aromatic or heteroaromatic moiety, -CF₃, -CN, or the like.


As used herein, the term "nitro" means -NO₂; the term "halogen" designates -F, -Cl, -Br or -I; the term "sulfhydryl" means -SH; the term "hydroxyl" means -OH; and the term "sulfonyl" means -SO₂-.

The terms "amine" and "amino" are art-recognized and refer to both unsubstituted and substituted amines, e.g., a moiety that may be represented by the general formulas:




wherein R50, R51 and R52 each independently represent a hydrogen, an alkyl, an alkenyl, -(CH₂)_m-R61, or R50 and R51, taken together with the N atom to which they are attached complete a heterocycle having from 4 to 8 atoms in the ring structure; R61 represents an aryl, a cycloalkyl, a cycloalkenyl, a heterocycle or a polycycle; and m is zero or an integer in the range of 1 to 8. In certain embodiments, only one of R50 or R51 may be a carbonyl, e.g., R50, R51 and the nitrogen together do not form an imide. In other embodiments, R50 and R51 (and optionally R52) each independently represent a hydrogen, an alkyl, an alkenyl, or -(CH₂)_m-R61. Thus, the term "alkylamine" includes an amine group, as defined above, having a substituted or unsubstituted alkyl attached thereto, i.e., at least one of R50 and R51 is an alkyl group.

The term "acylamino" is art-recognized and refers to a moiety that may be represented by the general formula:



wherein R50 is as defined above, and R54 represents a hydrogen, an alkyl, an alkenyl or $-(CH_2)_m-R61$, where m and R61 are as defined above.


The term "amido" is art recognized as an amino-substituted carbonyl and includes a moiety that may be represented by the general formula:



wherein R50 and R51 are as defined above. Certain embodiments of the amide in the present invention will not include imides which may be unstable.

The term "alkylthio" refers to an alkyl group, as defined above, having a sulfur radical attached thereto. In certain embodiments, the "alkylthio" moiety is represented by one of -S-alkyl, -S-alkenyl, -S-alkynyl, and -S- $(CH_2)_m-R61$, wherein m and R61 are defined above. Representative alkylthio groups include methylthio, ethyl thio, and the like.

The term "carboxyl" is art recognized and includes such moieties as may be represented by the general formulas:




wherein X50 is a bond or represents an oxygen or a sulfur, and R55 and R56 represents a hydrogen, an alkyl, an alkenyl, $-(CH_2)_m-R61$ or a pharmaceutically acceptable salt, R56 represents a hydrogen, an alkyl, an alkenyl or $-(CH_2)_m-R61$, where m and R61 are defined above. Where X50 is an oxygen and R55 or R56 is not hydrogen, the formula represents an "ester". Where X50 is an oxygen, and R55 is as defined above, the moiety is referred to herein as a carboxyl group, and particularly when R55 is a hydrogen, the formula represents a "carboxylic acid". Where X50 is an oxygen, and R56 is hydrogen, the formula represents a "formate". In general, where the oxygen atom of the above formula is replaced by sulfur, the formula represents a "thiolcarbonyl" group. Where X50 is a sulfur and R55 or R56 is not hydrogen, the formula represents a "thiolester." Where X50 is a sulfur and R55 is hydrogen, the formula represents a "thiolcarboxylic acid." Where X50 is a sulfur and R56 is hydrogen, the formula represents a "thiolformate." On the other hand, where X50 is a

bond, and R55 is not hydrogen, the above formula represents a "ketone" group. Where X50 is a bond, and R55 is hydrogen, the above formula represents an "aldehyde" group.

The terms "alkoxyl" or "alkoxy" as used herein refers to an alkyl group, as defined above, having an oxygen radical attached thereto. Representative alkoxyl groups include methoxy, ethoxy, propyloxy, tert-butoxy and the like. An "ether" is two hydrocarbons covalently linked by an oxygen. Accordingly, the substituent of an alkyl that renders that alkyl an ether is or resembles an alkoxyl, such as may be represented by one of -O-alkyl, -O-alkenyl, -O-alkynyl, -O-(CH₂)_m-R_g, where m and R_g are described above.

The term "sulfonate" is art recognized and includes a moiety that may be represented by the general formula:



in which R₄₁ is an electron pair, hydrogen, alkyl, cycloalkyl, or aryl.

The terms triflyl, tosyl, mesyl, and nonafllyl are art-recognized and refer to trifluoromethanesulfonyl, *p*-toluenesulfonyl, methanesulfonyl, and nonafluorobutanesulfonyl groups, respectively. The terms triflate, tosylate, mesylate, and nonaflate are art-recognized and refer to trifluoromethanesulfonate ester, *p*-toluenesulfonate ester, methanesulfonate ester, and nonafluorobutanesulfonate ester functional groups and molecules that contain said groups, respectively.

The term "carbamoyl" refers to -O(C=O)NRR', where R and R' are independently H, aliphatic groups, aryl groups or heteroaryl groups.

The term "alkylamino" refers to -NHR, where R is an alkyl group.

The term "dialkylamino" refers to -NRR', where both R and R' are alkyl groups.

The term "hydroxyalkyl" refers to -R-OH, where R is an aliphatic group.

The term "aminoalkyl" refers to -R-NH₂, where R is an aliphatic group.

The term "alkylaminoalkyl" refers to -R-NH-R', where both R and R' are aliphatic groups.

The term "dialkylaminoalkyl" refers to -R-N(R')-R'', where R, R', and R'' are aliphatic groups.


The term "arylaminoalkyl" refers to -R-NH-R', where R is an aliphatic and R' is an aryl group.

The term "oxo" refers to a carbonyl oxygen (=O).

The term "thioxo" refers to a carbonyl sulfur (=S).

- 5 The abbreviations Me, Et, Ph, Tf, Nf, Ts, Ms represent methyl, ethyl, phenyl, trifluoromethanesulfonyl, nonafluorobutanesulfonyl, *p*-toluenesulfonyl and methanesulfonyl, respectively. A more comprehensive list of the abbreviations utilized by organic chemists of ordinary skill in the art appears in the first issue of each volume of the *Journal of Organic Chemistry*; this list is typically presented in a table entitled Standard
10 List of Abbreviations. The abbreviations contained in said list, and all abbreviations utilized by organic chemists of ordinary skill in the art are hereby incorporated by reference.


The term "sulfate" is art recognized and includes a moiety that may be represented by the general formula:




15

in which R₄₁ is as defined above.


The term "sulfonylamino" is art recognized and includes a moiety that may be represented by the general formula:



- 20 The term "sulfamoyl" is art-recognized and includes a moiety that may be represented by the general formula:




The term "sulfonyl", as used herein, refers to a moiety that may be represented by the general formula:



in which R₄₄ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl.

The term "sulfoxido" as used herein, refers to a moiety that may be represented by
5 the general formula:



in which R₄₄ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aralkyl, or aryl.

A "selenoalkyl" refers to an alkyl group having a substituted seleno group attached
10 thereto. Exemplary "selenoethers" which may be substituted on the alkyl are selected from one of -Se-alkyl, -Se-alkenyl, -Se-alkynyl, and -Se-(CH₂)_m-R₇, m and R₇ being defined above.

Analogous substitutions may be made to alkenyl and alkynyl groups to produce, for example, aminoalkenyls, aminoalkynyls, amidoalkenyls, amidoalkynyls, iminoalkenyls,
15 iminoalkynyls, thioalkenyls, thioalkynyls, carbonyl-substituted alkenyls or alkynyls.

As used herein, the definition of each expression, e.g., alkyl, m, n, etc., when it occurs more than once in any structure, is intended to be independent of its definition elsewhere in the same structure.

It will be understood that "substitution" or "substituted with" includes the implicit
20 proviso that such substitution is in accordance with permitted valence of the substituted atom and the substituent, and that the substitution results in a stable compound, e.g., which does not spontaneously undergo transformation such as by rearrangement, cyclization, elimination, etc.

As used herein, the term "substituted" is contemplated to include all permissible
25 substituents of organic compounds. In a broad aspect, the permissible substituents include acyclic and cyclic, branched and unbranched, carbocyclic and heterocyclic, aromatic and nonaromatic substituents of organic compounds. Illustrative substituents include, for

example, those described herein above. The permissible substituents may be one or more and the same or different for appropriate organic compounds. For purposes of this invention, the heteroatoms such as nitrogen may have hydrogen substituents and/or any permissible substituents of organic compounds described herein which satisfy the valences
5 of the heteroatoms. This invention is not intended to be limited in any manner by the permissible substituents of organic compounds.

The phrase "protecting group" as used herein means temporary substituents which protect a potentially reactive functional group from undesired chemical transformations. Examples of such protecting groups include esters of carboxylic acids, silyl ethers of
10 alcohols, and acetals and ketals of aldehydes and ketones, respectively. The field of protecting group chemistry has been reviewed (Greene, T.W.; Wuts, P.G.M. *Protective Groups in Organic Synthesis*, 2nd ed.; Wiley: New York, 1991). Protected forms of the inventive compounds are included within the scope of this invention.

Certain compounds of the present invention may exist in particular geometric or
15 stereoisomeric forms. The present invention contemplates all such compounds, including *cis*- and *trans*-isomers, *R*- and *S*-enantiomers, diastereomers, (D)-isomers, (L)-isomers, the racemic mixtures thereof, and other mixtures thereof, as falling within the scope of the invention. Additional asymmetric carbon atoms may be present in a substituent such as an alkyl group. All such isomers, as well as mixtures thereof, are intended to be included in
20 this invention.

If, for instance, a particular enantiomer of a compound of the present invention is desired, it may be prepared by asymmetric synthesis, or by derivation with a chiral auxiliary, where the resulting diastereomeric mixture is separated and the auxiliary group cleaved to provide the pure desired enantiomers. Alternatively, where the molecule
25 contains a basic functional group, such as amino, or an acidic functional group, such as carboxyl, diastereomeric salts are formed with an appropriate optically-active acid or base, followed by resolution of the diastereomers thus formed by fractional crystallization or chromatographic means well known in the art, and subsequent recovery of the pure enantiomers.

30 Contemplated equivalents of the compounds described above include compounds which otherwise correspond thereto, and which have the same general properties thereof, wherein one or more simple variations of substituents are made which do not adversely

affect the efficacy of the compound in binding to sigma receptors. In general, the compounds of the present invention may be prepared by the methods illustrated in the general reaction schemes as, for example, described below, or by modifications thereof, using readily available starting materials, reagents and conventional synthesis procedures.

5 In these reactions, it is also possible to make use of variants which are in themselves known, but are not mentioned here.

The term "subject" as used herein, refers to an animal, typically a mammal or a human, that has been the object of treatment, observation, and/or experiment. When the term is used in conjunction with administration of a compound or drug, then the subject has
10 been the object of treatment, observation, and/or administration of the compound or drug.

The term "therapeutically effective amount" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a cell culture, tissue system, animal, or human that is being sought by a researcher, veterinarian, clinician, or physician, which includes alleviation of the symptoms of the
15 disease, condition, or disorder being treated. In the present invention, such an amount will be sufficient to bind to Bcl-2 in a cell and inhibit at least part of the anti-apoptotic activity of the protein. Such an amount may be sufficient to provide therapeutic effectiveness in a patient or may serve to sensitize the cell to treatment with another anticancer agent.

The term "composition" is intended to encompass a product comprising the
20 specified ingredients in the specified amounts, as well as any product that results, directly or indirectly, from combinations of the specified ingredients in the specified amounts.

The term "pharmaceutically acceptable carrier" refers to a medium that is used to prepare a desired dosage form of a compound. A pharmaceutically acceptable carrier can include one or more solvents, diluents, or other liquid vehicles; dispersion or suspension
25 aids; surface active agents; isotonic agents; thickening or emulsifying agents; preservatives; solid binders; lubricants; and the like. Remington's Pharmaceutical Sciences, Fifteenth Edition, E. W. Martin (Mack Publishing Co., Easton, Pa., 1975) and Handbook of Pharmaceutical Excipients, Third Edition, A. H. Kibbe ed. (American Pharmaceutical Assoc. 2000), disclose various carriers used in formulating pharmaceutical compositions
30 and known techniques for the preparation thereof.

The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial,

intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion.

The phrases "systemic administration," "administered systemically," "peripheral
5 administration" and "administered peripherally" as used herein mean the administration of a compound, drug or other material other than directly into the site of the Bcl mediated disorder, such that it enters the patient's system and, thus, is subject to metabolism and other like processes, for example, subcutaneous administration.

The phrase "therapeutically-effective amount" as used herein means that amount of a
10 compound, material, or composition comprising a compound of the present invention which is effective for producing some desired therapeutic effect in at least a sub-population of cells in an animal at a reasonable benefit/risk ratio applicable to any medical treatment.

The phrase "pharmaceutically acceptable" is employed herein to refer to those
15 compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

The phrase "pharmaceutically-acceptable carrier" as used herein means a
20 pharmaceutically-acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, manufacturing aid (e.g., lubricant, talc magnesium, calcium or zinc stearate, or steric acid), or solvent encapsulating material, involved in carrying or transporting the subject compound from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not injurious to the patient. Some
25 examples of materials which can serve as pharmaceutically-acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil,
30 cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid;

(16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) pH buffered solutions; (21) polyesters, polycarbonates and/or polyanhydrides; and (22) other non-toxic compatible substances employed in pharmaceutical formulations.

The phrases "Bcl-mediated disorder" and "disorder mediated by cells expressing Bcl proteins" refer to pathological and disease conditions in which a Bcl protein plays a role. Such roles may be directly related to the pathological condition or may be indirectly related to the condition. The feature common to this class of conditions is that they may be ameliorated by inhibiting the activity of, function of, or association with Bcl proteins.

As used herein, the terms "Bcl" and "Bcl protein" are intended to encompass one or more of the Bcl-2 subfamily of anti-apoptotic proteins Bcl-2, Bcl-w, Mcl-1, Bcl-XL, A1, Bfl1, Bcl-B, BOO/DIVA, and their homologues.

Synthesis of Heterocyclic Compounds

The isoxazolidine compounds of the invention may be prepared using a [3+2] cycloaddition reaction between a nitron and an alkene. The nitron substrate and alkene may contain functional groups suitable for chemical derivatization following synthesis of the isoxazolidine core. In certain instances, a Lewis acid is added to the reaction. In a preferred embodiment, the Lewis acid is $\text{Ti}(\text{O}i\text{-Pr})_4$. In certain instances, the reaction mixture is subjected to microwave radiation. In general, the subject reactions are carried out in a liquid reaction medium, but may be carried out on a solid support. The reactions may be conducted in an aprotic solvent, preferably one in which the reaction ingredients are substantially soluble. Suitable solvents include ethers, such as diethyl ether, 1,2-dimethoxyethane, diglyme, t-butyl methyl ether, tetrahydrofuran and the like; halogenated solvents, such as chloroform, dichloromethane, dichloroethane, chlorobenzene, carbon tetrachloride, and the like; aliphatic or aromatic hydrocarbon solvents, such as benzene, xylene, toluene, hexane, pentane and the like; esters and ketones, such as ethyl acetate, acetone, and 2-butanone; polar aprotic solvents, such as acetonitrile, dimethylsulfoxide, dimethylformamide, pyridine, and the like; or combinations of two or more solvents. The reactions may be conducted at a variety of temperatures. In certain instances, the cycloaddition reaction is conducted using a substrate attached to a solid support. Following synthesis of the isoxazolidine core, the isoxazolidine compound may be derivatized using a variety of functionalization reactions known in the art. Representative examples include

palladium coupling reactions to alkenylhalides or aryl halides, oxidations, reductions, reactions with nucleophiles, reactions with electrophiles, pericyclic reactions, installation of protecting groups, removal of protecting groups, and the like.

As set out above, certain embodiments of the present compounds may contain a
5 basic functional group, such as amino or alkylamino, and are, thus, capable of forming pharmaceutically-acceptable salts with pharmaceutically-acceptable acids. The term "pharmaceutically-acceptable salts" in this respect, refers to the relatively non-toxic, inorganic and organic acid addition salts of compounds of the present invention. These salts may be prepared *in situ* in the administration vehicle or the dosage form
10 manufacturing process, or by separately reacting a purified compound of the invention in its free base form with a suitable organic or inorganic acid, and isolating the salt thus formed during subsequent purification. Representative salts include the hydrobromide, hydrochloride, sulfate, bisulfate, phosphate, nitrate, acetate, valerate, oleate, palmitate, stearate, laurate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate,
15 tartrate, naphylate, mesylate, glucoheptonate, lactobionate, and laurylsulphonate salts and the like. (See, for example, Berge et al. (1977) "Pharmaceutical Salts", *J. Pharm. Sci.* 66:1-19).

The pharmaceutically acceptable salts of the subject compounds include the conventional nontoxic salts or quaternary ammonium salts of the compounds, e.g., from
20 non-toxic organic or inorganic acids. For example, such conventional nontoxic salts include those derived from inorganic acids such as hydrochloride, hydrobromic, sulfuric, sulfamic, phosphoric, nitric, and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, palmitic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-
25 acetoxibenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isothionic, and the like.

In other cases, the compounds of the present invention may contain one or more acidic functional groups and, thus, are capable of forming pharmaceutically-acceptable salts with pharmaceutically-acceptable bases. The term "pharmaceutically-acceptable salts" in
30 these instances refers to the relatively non-toxic, inorganic and organic base addition salts of compounds of the present invention. These salts can likewise be prepared *in situ* in the administration vehicle or the dosage form manufacturing process, or by separately reacting

the purified compound in its free acid form with a suitable base, such as the hydroxide, carbonate or bicarbonate of a pharmaceutically-acceptable metal cation, with ammonia, or with a pharmaceutically-acceptable organic primary, secondary or tertiary amine. Representative alkali or alkaline earth salts include the lithium, sodium, potassium, calcium, magnesium, and aluminum salts and the like. Representative organic amines useful for the formation of base addition salts include ethylamine, diethylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine and the like. (See, for example, Berge et al., *supra*)

10 Biological Activity Analysis

The following *in vitro* binding and cellular assays may be used to determine the activity and specificity of compounds of the present invention to bind to Bcl-2 and inhibit Bcl-2 function in a cell.

15 Bcl-2 Binding Assay

Bcl-2 and Bcl-xL binding may be determined using a variety of known methods. One such assay is a sensitive and quantitative *in vitro* binding assay using fluorescence polarization (FP) described by Wang, J. -L.; Zhang, Z. -J.; Choksi, S.; Sjam, S.; Lu, Z.; Croce, C. M.; Alnemri, E. S.; Komgold, R.; Huang, Z. Cell permeable Bcl-2 binding peptides: a chemical approach to apoptosis induction in tumor cells. *Cancer Res.* 2000, 60, 1498-1502).

Cell based assays

The ability of isoxazolidine compounds of the present invention to inhibit cell-viability in cancer cells with Bcl-2 protein over-expression was demonstrated. When RL-cells are exposed to isoxazolidine compounds of the present invention, the inhibitors show a dose-dependent cell-killing in Alamar blue cytotoxicity assays. When Panc1 cells are exposed to the isoxazolidine compounds of the present invention in combination with camptothecin, the inhibitors show a synergistic dose-dependent cell killing in propidium iodide exclusion cell survival assays.

30 Pharmaceutical Compositions

In another aspect, the present invention provides pharmaceutically acceptable compositions which comprise a therapeutically-effective amount of one or more of the

compounds described above, formulated together with one or more pharmaceutically acceptable carriers (additives) and/or diluents. As described in detail below, the pharmaceutical compositions of the present invention may be specially formulated for administration in solid or liquid form, including those adapted for the following: (1) oral
5 administration, for example, drenches (aqueous or non-aqueous solutions or suspensions), tablets, e.g., those targeted for buccal, sublingual, and systemic absorption, boluses, powders, granules, pastes for application to the tongue; (2) parenteral administration, for example, by subcutaneous, intramuscular, intravenous or epidural injection as, for example, a sterile solution or suspension, or sustained-release formulation; (3) topical application, for
10 example, as a cream, ointment, or a controlled-release patch or spray applied to the skin; (4) intravaginally or intrarectally, for example, as a pessary, cream or foam; (5) sublingually; (6) ocularly; (7) transdermally; (8) nasally; (9) pulmonary; or (10) intrathecally.

Formulations of the present invention include those suitable for oral, nasal, topical (including buccal and sublingual), rectal, vaginal and/or parenteral administration. The
15 formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. The amount of active ingredient which may be combined with a carrier material to produce a single dosage form will vary depending upon the host being treated, the particular mode of administration. The amount of active ingredient which may be combined with a carrier material to produce a single
20 dosage form will generally be that amount of the compound which produces a therapeutic effect. Generally, out of one hundred per cent, this amount will range from about 0.1 per cent to about ninety-nine percent of active ingredient, preferably from about 5 per cent to about 70 per cent, most preferably from about 10 percent to about 30 percent.

Methods of preparing these formulations or compositions include the step of
25 bringing into association a compound of the present invention with the carrier and, optionally, one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association a compound of the present invention with liquid carriers, or finely divided solid carriers, or both, and then, if necessary, shaping the product.

30 Formulations of the invention suitable for oral administration may be in the form of capsules, cachets, pills, tablets, lozenges (using a flavored basis, usually sucrose and acacia or tragacanth), powders, granules, or as a solution or a suspension in an aqueous or non-

aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion, or as an elixir or syrup, or as pastilles (using an inert base, such as gelatin and glycerin, or sucrose and acacia) and/or as mouth washes and the like, each containing a predetermined amount of a compound of the present invention as an active ingredient. A compound of the present invention may also be administered as a bolus, electuary or paste.

In solid dosage forms of the invention for oral administration (capsules, tablets, pills, dragees, powders, granules, trouches and the like), the active ingredient is mixed with one or more pharmaceutically-acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; (3) humectants, such as glycerol; (4) disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; (5) solution retarding agents, such as paraffin; (6) absorption accelerators, such as quaternary ammonium compounds and surfactants, such as poloxamer and sodium lauryl sulfate; (7) wetting agents, such as, for example, cetyl alcohol, glycerol monostearate, and non-ionic surfactants; (8) absorbents, such as kaolin and bentonite clay; (9) lubricants, such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, zinc stearate, sodium stearate, stearic acid, and mixtures thereof; (10) coloring agents; and (11) controlled release agents such as crospovidone or ethyl cellulose. In the case of capsules, tablets and pills, the pharmaceutical compositions may also comprise buffering agents. Solid compositions of a similar type may also be employed as fillers in soft and hard-shelled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared using binder (for example, gelatin or hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (for example, sodium starch glycolate or cross-linked sodium carboxymethyl cellulose), surface-active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

The tablets, and other solid dosage forms of the pharmaceutical compositions of the present invention, such as dragees, capsules, pills and granules, may optionally be scored or prepared with coatings and shells, such as enteric coatings and other coatings well known in the pharmaceutical-formulating art. They may also be formulated so as to provide slow or controlled release of the active ingredient therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile, other polymer matrices, liposomes and/or microspheres. They may be formulated for rapid release, e.g., freeze-dried. They may be sterilized by, for example, filtration through a bacteria-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which may be dissolved in sterile water, or some other sterile injectable medium immediately before use. These compositions may also optionally contain opacifying agents and may be of a composition that they release the active ingredient(s) only, or preferentially, in a certain portion of the gastrointestinal tract, optionally, in a delayed manner. Examples of embedding compositions which may be used include polymeric substances and waxes. The active ingredient can also be in micro-encapsulated form, if appropriate, with one or more of the above-described excipients.

Liquid dosage forms for oral administration of the compounds of the invention include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredient, the liquid dosage forms may contain inert diluents commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, coloring, perfuming and preservative agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

Pharmaceutical compositions of this invention suitable for parenteral administration comprise one or more compounds of the invention in combination with one or more pharmaceutically-acceptable sterile isotonic aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just prior to use, which may contain sugars, alcohols, antioxidants, buffers, bacteriostats, solutes which render the formulation isotonic with the blood of the intended recipient or suspending or thickening agents.

Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity may be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of the action of microorganisms upon the subject compounds may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of a drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material having poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally-administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microencapsule matrices of the subject compounds in biodegradable polymers such as polylactide-polyglycolide. Depending on the ratio of drug to polymer, and the nature of the particular polymer employed, the rate of

drug release may be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissue.

Dosage forms for the topical or transdermal administration of a compound of this invention include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. The active compound may be mixed under sterile conditions with a pharmaceutically-acceptable carrier, and with any preservatives, buffers, or propellants which may be required.

The ointments, pastes, creams and gels may contain, in addition to an active compound of this invention, excipients, such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.

Powders and sprays can contain, in addition to a compound of this invention, excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder, or mixtures of these substances. Sprays can additionally contain customary propellants, such as chlorofluorohydrocarbons and volatile unsubstituted hydrocarbons, such as butane and propane.

Transdermal patches have the added advantage of providing controlled delivery of a compound of the present invention to the body. Such dosage forms may be made by dissolving or dispersing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate of such flux may be controlled by either providing a rate controlling membrane or dispersing the compound in a polymer matrix or gel.

Formulations of the pharmaceutical compositions of the invention for rectal or vaginal administration may be presented as a suppository, which may be prepared by mixing one or more compounds of the invention with one or more suitable nonirritating excipients or carriers comprising, for example, cocoa butter, polyethylene glycol, a suppository wax or a salicylate, and which is solid at room temperature, but liquid at body temperature and, therefore, will melt in the rectum or vaginal cavity and release the active compound.

Formulations of the present invention which are suitable for vaginal administration also include pessaries, tampons, creams, gels, pastes, foams or spray formulations containing such carriers as are known in the art to be appropriate.

Ophthalmic formulations, eye ointments, powders, solutions and the like, are also contemplated as being within the scope of this invention.

When the compounds of the present invention are administered as pharmaceuticals, to humans and animals, they may be given per se or as a pharmaceutical composition containing, for example, 0.1 to 99% (more preferably, 10 to 30%) of active ingredient in combination with a pharmaceutically acceptable carrier.

Regardless of the route of administration selected, the compounds of the present invention, which may be used in a suitable hydrated form, and/or the pharmaceutical compositions of the present invention, are formulated into pharmaceutically-acceptable dosage forms by conventional methods known to those of skill in the art.

15 Methods of Therapy and Treatment

The present invention further provides methods for treating and reducing the severity of cancer as well as other Bcl-mediated disorders or conditions.

Bcl-2 inhibitors have been shown to be active against a number of cancer cell lines as single agent, including, but not limited to, breast cancer (US 2003/0119894, published PCT applications WO 02/097053 and WO 02/13833), lymphomas (*Nature* (2005) 435, 677-681), small cell lung cancer (*Nature* (2005) 435, 677-681), head and neck cancer (published PCT application WO 02/097053), and leukemias (published PCT application WO 02/13833).

Bcl-2 inhibitors have been shown to be active against a number of cancer cell lines in combination with other anticancer agents and radiation, including, but not limited to, breast cancer (With docetaxel, published PCT application WO 02/097053), prostate cancer (With docetaxel, published PCT application WO 02/097053), head and neck cancer (With docetaxel, published PCT application WO 02/097053), and non small-cell lung cancer (With paclitaxel, *Nature* (2005) 435, 677-681). In addition to the aforementioned combination chemotherapeutics, small molecule inhibitors of Bcl-2 proteins display synergy with other anticancer agents, including, but not limited to etoposide, doxorubicin, cisplatin, paclitaxel, and radiation (*Nature* (2005) 435, 677-681).

Cancers or neoplastic diseases and related disorders that may be treated by administration of compounds and compositions of the present invention, include, but are not limited to those listed in Table 1 (for a review of such disorders, see Fishman et al., 1985, Medicine, 2d Ed., J. B. Lippincott Co., Philadelphia):

TABLE 1

CANCERS AND NEOPLASTIC DISORDERS	
Leukemia	
5	acute leukemia
	acute lymphocytic leukemia
	acute myelocytic leukemia
	myeloblastic
	promyelocytic
10	myelomonocytic
	monocytic
	erythroleukemia
	chronic leukemia
	chronic myelocytic (granulocytic) leukemia
15	chronic lymphocytic leukemia
Polycythemia vera	
Lymphoma	
	Hodgkin's disease
	non-Hodgkin's disease
20	Multiple myeloma
	Waldenstrom's macroglobulinemia
	Heavy chain disease
Solid tumors	
	sarcomas and carcinomas
25	fibrosarcoma
	myxosarcoma
	liposarcoma
	chondrosarcoma
	osteogenic sarcoma
30	chordoma
	angiosarcoma
	lymphangiosarcoma
	lymphangioendotheliosarcoma

synovioma
mesothelioma
Ewing's tumor
leiomyosarcoma
5 rhabdomyosarcoma
colon carcinoma
pancreatic cancer
breast cancer
ovarian cancer
10 prostate cancer
squamous cell carcinoma
basal cell carcinoma
adenocarcinoma
sweat gland carcinoma
15 sebaceous gland carcinoma
papillary carcinoma
papillary adenocarcinomas
cystadenocarcinoma
medullary carcinoma
20 bronchogenic carcinoma
renal cell carcinoma
hepatoma
bile duct carcinoma
choriocarcinoma
25 seminoma
embryonal carcinoma
Wilms' tumor
cervical cancer
uterine cancer
30 testicular tumor
lung carcinoma
small cell lung carcinoma
bladder carcinoma

5 epithelial carcinoma
 glioma
 astrocytoma
 medulloblastoma
 craniopharyngioma
 ependymoma
 pinealoma
 hemangioblastoma
 acoustic neuroma
10 oligodendroglioma
 meningioma
 melanoma
 neuroblastoma
 retinoblastoma

15

 In certain embodiments, the compounds of the present invention are used to treat cancers including, but not limited to lymphomas (preferably follicular lymphoma, diffuse large B-cell lymphoma, mantle cell lymphoma, or chronic lymphocytic leukemia), prostate cancer (more preferably hormone insensitive), breast cancer (preferably estrogen receptor
20 positive), neuroblastoma, colorectal, endometrial, ovarian, lung (preferably small cell), hepatocellular carcinoma, multiple myeloma, head and neck or testicular cancer (preferably germ cell).

 Actual dosage levels of the active ingredients in the pharmaceutical compositions of this invention may be varied so as to obtain an amount of the active ingredient which is
25 effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient.

 The selected dosage level will depend upon a variety of factors including the activity of the particular compound of the present invention employed, or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion
30 or metabolism of the particular compound being employed, the rate and extent of absorption, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compound employed, the age, sex, weight, condition,

general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

A physician or veterinarian having ordinary skill in the art can readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of the compounds of the invention employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved.

In general, a suitable daily dose of a compound of the invention will be that amount of the compound which is the lowest dose effective to produce a therapeutic effect. Such an effective dose will generally depend upon the factors described above. Generally, oral, intravenous, intracerebroventricular and subcutaneous doses of the compounds of this invention for a patient, when used for the indicated analgesic effects, will range from about 0.0001 to about 100 mg per kilogram of body weight per day.

If desired, the effective daily dose of the active compound may be administered as two, three, four, five, six or more sub-doses administered separately at appropriate intervals throughout the day, optionally, in unit dosage forms. In one embodiment, dosing is one administration per day.

The compound of the invention may be administered as such or in admixtures with pharmaceutically acceptable carriers and can also be administered in conjunction with antimicrobial agents such as penicillins, cephalosporins, aminoglycosides and glycopeptides. Conjunctive therapy, thus includes sequential, simultaneous and separate administration of the active compound in a way that the therapeutical effects of the first administered one is not entirely disappeared when the subsequent is administered.

25

Treatment of Cancer in Combination with Chemotherapy or Radiotherapy

In certain embodiments, one or more compounds of the present invention are used to treat or prevent cancer or neoplastic disease in combination with one or more anti-cancer, chemotherapeutic agents including, but not limited to, methotrexate, taxol, mercaptopurine, thioguanine, hydroxyurea, cytarabine, cyclophosphamide, ifosfamide, nitrosoureas, cisplatin, carboplatin, mitomycin, dacarbazine, procarbazine, etoposides, prednisolone, dexamethasone, cytarbine, campathecins, bleomycin, doxorubicin, idarubicin,

daunorubicin, dactinomycin, plicamycin, mitoxantrone, asparaginase, vinblastine, vincristine, vinorelbine, paclitaxel, and docetaxel. In a preferred embodiment, one or more compound of the present invention is used to treat or prevent cancer or neoplastic disease in combination with one or more chemotherapeutic or other anti-cancer agents including, but not limited to those presented in Table 2.

TABLE 2

CHEMOTHERAPEUTICS AND OTHER
ANTI-CANCER AGENTS

10	Radiation:	γ -radiation
	<u>Alkylating agents</u>	
	Nitrogen mustards:	cyclophosphamide
		Ifosfamide
15		trofosfamide
		Chlorambucil
		Estramustine
		melphalan
	Nitrosoureas:	carmustine (BCNU)
20		Lomustine (CCNU)
	Alkylsulphonates	busulfan
		Treosulfan
	Triazenes:	Dacarbazine
	Platinum containing compounds:	Cisplatin
25		carboplatin
		oxaplatin
	<u>Plant Alkaloids</u>	
	Vinca alkaloids:	vincristine
30		Vinblastine
		Vindesine
		Vinorelbine
	Taxoids:	paclitaxel

	Docetaxol
	<u>DNA Topoisomerase Inhibitors</u>
5	Epipodophyllins: etoposide Teniposide Topotecan 9-aminocamptothecin campto irinotecan crisnatol
10	<u>mytomycins</u> mytomycin C Anti-metabolites <u>Anti-folates:</u>
15	DHFR inhibitors: methotrexate Trimetrexate IMP dehydrogenase Inhibitors: mycophenolic acid Tiazofurin
20	Ribavirin EICAR Ribonucleotide reductase Inhibitors: hydroxyurea deferroxamine
25	<u>Pyrimidine analogs:</u> Uracil analogs 5-Fluorouracil Floxuridine Doxifluridine Ratitrexed
30	capecitabine Cytosine analogs cytarabine (ara C) Cytosine arabinoside fludarabine

	Purine analogs:	mercaptopurine Thioguanine
	Hormonal therapies:	
	<u>Receptor antagonists:</u>	
5	Anti-estrogens	Tamoxifen Raloxifene megestrol
	LHRH agonists:	goserelin
10	Anti-androgens:	Leuprolide acetate flutamide bicalutamide
	<u>Retinoids/Deletoids</u>	
15	Vitamin D3 analogs:	EB 1089 CB 1093 KH 1060
	Photodynamic therapies:	verteporfin (BPD-MA) Phthalocyanine
20		photosensitizer Pc4 Demethoxy-hypocrellin A (2BA-2-DMHA)
	Cytokines:	Interferon α Interferon γ
25		Tumor necrosis factor
	<u>Others:</u>	
		Prednisilone Imatinib Thalidomide
30		Lenalidomide Bortezomib Gemcitabine Erlotinib

	Gefitinib
	Sorafenib
	Sutinib
Isoprenylation inhibitors:	Lovastatin
5 Dopaminergic neurotoxins:	1-methyl-4-phenylpyridinium ion
Cell cycle inhibitors:	staurosporine
Actinomycins:	Actinomycin D
	Dactinomycin
Bleomycins:	bleomycin A2
10 Bleomycin B2	
	Peplomycin
Anthracyclines:	daunorubicin
	Doxorubicin (adriamycin)
	Idarubicin
15 Epirubicin	
	Pirarubicin
	Zorubicin
	Mitoxantrone
MDR inhibitors:	verapamil
20 Ca ²⁺ ATPase inhibitors:	thapsigargin
Antibodies	Avastin
	Erbix
	Rituxan

25 The chemotherapeutic agent and/or radiation therapy may be administered according to therapeutic protocols well known in the art. It will be apparent to those skilled in the art that the administration of the chemotherapeutic agent and/or radiation therapy may be varied depending on the disease being treated and the known effects of the chemotherapeutic agent and/or radiation therapy on that disease. Also, in accordance with

30 the knowledge of the skilled clinician, the therapeutic protocols (e.g., dosage amounts and times of administration) may be varied in view of the observed effects of the administered therapeutic agents (i.e., antineoplastic agent or radiation) on the patient, and in view of the observed responses of the disease to the administered therapeutic agents.

Also, in general, compounds of the present invention and the chemotherapeutic agent do not have to be administered in the same pharmaceutical composition, and may, because of different physical and chemical characteristics, have to be administered by different routes. For example, compounds of the present invention may be administered intravenously to generate and maintain good blood levels, while the chemotherapeutic agent may be administered orally. The determination of the mode of administration and the advisability of administration, where possible, in the same pharmaceutical composition, is well within the knowledge of the skilled clinician. The initial administration may be made according to established protocols known in the art, and then, based upon the observed effects, the dosage, modes of administration and times of administration may be modified by the skilled clinician.

The particular choice of chemotherapeutic agent or radiation will depend upon the diagnosis of the attending physicians and their judgment of the condition of the patient and the appropriate treatment protocol.

A compound of the present invention, and chemotherapeutic agent and/or radiation may be administered concurrently (e.g., simultaneously, essentially simultaneously or within the same treatment protocol) or sequentially, depending upon the nature of the proliferative disease, the condition of the patient, and the actual choice of chemotherapeutic agent and/or radiation to be administered in conjunction (i.e., within a single treatment protocol) with a compound of the present invention.

If a compound of the present invention, and the chemotherapeutic agent and/or radiation are not administered simultaneously or essentially simultaneously, then the optimum order of administration of the compound of the present invention, and the chemotherapeutic agent and/or radiation, may be different for different tumors. Thus, in certain situations the compound of the present invention may be administered first followed by the administration of the chemotherapeutic agent and/or radiation; and in other situations the chemotherapeutic agent and/or radiation may be administered first followed by the administration of a compound of the present invention. This alternate administration may be repeated during a single treatment protocol. The determination of the order of administration, and the number of repetitions of administration of each therapeutic agent during a treatment protocol, is well within the knowledge of the skilled physician after evaluation of the disease being treated and the condition of the patient. For example, the chemotherapeutic agent and/or radiation may be administered first, especially if it is a

cytotoxic agent, and then the treatment continued with the administration of a compound of the present invention followed, where determined advantageous, by the administration of the chemotherapeutic agent and/or radiation, and so on until the treatment protocol is complete.


- 5 Thus, in accordance with experience and knowledge, the practicing physician can modify each protocol for the administration of a component (therapeutic agent, i.e., compound of the present invention, chemotherapeutic agent or radiation) of the treatment according to the individual patient's needs, as the treatment proceeds.

Exemplification


- 10 The invention now being generally described, it will be more readily understood by reference to the following examples, which are included merely for purposes of illustration of certain aspects and embodiments of the present invention, and are not intended to limit the invention.

Example 1

15




Part A



- 20 To a solution of phenol **2** (750 mg, 3 mmol, 1 eq) in DMF (5 mL) at 0 °C was added NaH (130 mg, 3.6 mmol, 1.2 eq) followed by MeI (280 μ L, 4.5 mmol, 1.5 eq). The reaction mixture was stirred at rt for 24 h and then quenched with water. The mixture was diluted with EtOAc and washed twice with water then brine. The solution was dried over MgSO_4 , filtered and concentrated to afford 790 mg of crude product **3**. Yield 100%.


Part B



25


Aldehyde 3 (790 mg, 3.0 mmol, 1 eq) and hydroxylamine hydrochloride (250 mg, 3.6 mmol, 1.2 eq) were dissolved in THF/MeOH (3:2, 10 mL). Water (2 mL) was added and the pH was adjusted to 9 with 6 N KOH. The reaction mixture was stirred at rt overnight. After 16 h, NaBH₃CN (380 mg, 6.07 mmol, 2 eq) was added followed by a crystal of methyl orange. The pH was adjusted to 2 and the resulting ruby red color was maintained for the duration of the reaction by the frequent addition of 1 N HCl. After stirring for 2 h, another portion of NaBH₃CN (380 mg, 6.10 mmol, 2 eq) was added. After stirring for a total of 16 h, the pH of the reaction mixture was brought to 7 and DCM was added. The mixture was washed twice with water, brine and then dried over MgSO₄. The crude product was purified by silica gel chromatography (50-100% EtOAc/hexane) to afford 706 mg of hydroxylamine 4. Yield 83%.

Part C




A solution of hydroxylamine 4 (700 mg, 2.5 mmol, 1 eq) and methyl glyoxylate (445 mg, 5.05 mmol, 2 eq) in benzene (15 mL) was heated at reflux with a Dean Stark trap overnight. Excess solvent was removed *in vacuo* and the resulting nitrone 5 was taken on crude in the next step.

Part D




Nitron 5 (880 mg, 2.5 mmol, 1 eq), allylic alcohol 6 (820 mg, 3.80 mmol, 1.5 eq) and Ti(*i*OPr)₄ (1.1 mL, 3.8 mmol, 1.5 eq) were dissolved in toluene (5 mL) and heated in the microwave at 120 °C for 10 min. The reaction mixture was diluted with EtOAc (15 mL) and 3-(dimethylamino)-1,2-propanediol (500 µL) was added. After stirring for 2 h, EtOAc was added and the mixture was washed twice with water and then brine, dried over MgSO₄, filtered over Celite and concentrated *in vacuo* to afford an oil. The crude residue was purified by silica gel chromatography (20% hexanes/EtOAc) to afford 575 mg of lactone 7. Yield 43%.

Part E




To a solution of lactone **7** (225 mg, 42 μmol) in THF (6 mL) was added pyridine (2 mL) and HF/pyridine (2 mL). The mixture was stirred at rt for 4 h then TMSOMe (8 mL) was added. Solvent was removed *in vacuo* and the crude product was purified by silica gel chromatography (EtOAc) to afford 128 mg of **8** as a white foam. Yield 72%.

Part F



To a flame-dried 10-mL round bottom flask containing (+)-isopinocampheylamine (110 μL , 0.61 mmol, 2 eq) and DCM (2 mL) was added trimethylaluminum (300 μL , 2 M in hexane, 0.61 mmol, 2 eq). After stirring for 15 min, a solution of lactone **8** (128 mg, 0.31 mmol, 1 eq) in DCM (4 mL) was added and the mixture was stirred at rt overnight. The reaction was quenched by the addition of a saturated aqueous Rochelle salt solution (5 mL) and the mixture was rapidly stirred for 2 h. DCM was added and the mixture washed three times with water and then brine. The solution was dried over MgSO_4 , filtered and concentrated *in vacuo* to afford a oil. The crude material was purified by silica gel chromatography (EtOAc) to afford 160 mg of **9**. Yield 91%.

Part G




A flask containing aryl iodide **9** (2.0 g, 3.5 mmol, 1 eq), 3-amino-5-carboxyphenylboronic acid (1.2 g, 7.0 mmol, 2 eq), cesium carbonate (3.5 g, 10 mmol, 3 eq), potassium acetate (300 mg, 3.5 mmol, 1 eq) and Pd(dppf)Cl_2 (300 mg, 0.35 mmol, 0.1 eq) was purged with argon and DMSO (35 mL) added. The mixture was heated at 60 $^\circ\text{C}$ for 18 h, then additional Pd(dppf)Cl_2 (300 mg, 0.35 mmol, 0.1 eq) was added and heating

continued for an additional 18 h. The reaction mixture was added to water (300 mL), acidified with 6 N HCl until the aqueous layer attained a pH of 4, and extracted with DCM (3 x 100 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated *in vacuo* to afford a brown oil.


- 5 This crude oil was dissolved in MeOH (50 mL) and treated with AcOH (1 mL), 37% formalin (2 mL, 27 mmol, 8 eq) and NaBH₃CN (600 mg, 10 mmol, 2.7 eq). After stirring at rt for 2 h, the reaction mixture was partitioned between water (300 mL) and DCM (100 mL) and acidified with 6 N HCl until the aqueous layer attained a pH of 4. The layers were separated and the aqueous extracted with DCM (3 x 80 mL). The combined
10 organic layers were dried over Na₂SO₄, filtered, treated with Et₃N (2 mL), and concentrated *in vacuo* to afford a brown oil. The oil was purified by silica gel chromatography (5-7.5% MeOH/DCM) to give 1.34 g of **10** as a brown solid. Yield 63%.

Part H



- 15 A solution of **10** (300 mg, 0.5 mmol, 1 eq) in DCM (10 mL) was treated with DIPEA (150 μ L, 0.86 mmol, 1.8 eq), (S)-N¹,N¹,4,4-tetramethylpentane-1,2-diamine (120 mg, 0.74 mmol, 1.5 eq), and HBTU (230 mg, 0.62 mmol, 1.3 eq). After stirring overnight at rt, the mixture was added to water (90 mL), and extracted with DCM (3 x 50 mL). The organic phases were dried over Na₂SO₄, filtered and concentrated *in vacuo* to give a brown
20 solid. This material was purified by silica gel chromatography (5-10% MeOH/DCM) to give 161 mg **1** as a tan solid. Yield 44%. MS (ESI(+)) *m/z* 750.91 (M+H)⁺.


Example 2



11


Compound **11** was synthesized according to the procedure described in Example 1 using (S)-N¹,N¹,5-trimethylhexane-1,3-diamine in place of (S)-N¹,N¹,4,4-tetramethylpentane-1,2-diamine. 49% yield. MS (ESI(+)) *m/z* 376.23 (M+2H)²⁺.

Example 3



12


Part A



To a flask containing Boc-D-phenylalaninol **13** (100 mg, 0.4 mmol, 1 eq), and triethylbenzylammonium chloride (10 mg, 0.04 mmol, 0.1 eq) was added THF (4 mL), 50% NaOH (3 mL), and MeI (26 μ L, 0.4 mmol, 1 eq). After stirring at rt for 72 h, the reaction mixture was diluted with water (30 mL) and extracted with DCM (3 x 20 mL). The organic layers were dried over Na₂SO₄, concentrated, and purified by silica gel flash chromatography (15% EtOAc/hexanes), providing a clear oil (ca. 50 mg, 50%).


This resulting oil was dissolved in trifluoroacetic acid (5 mL) and stirred for 2 h. The acid was removed *in vacuo* and the residue co-evaporated with toluene to give the trifluoroacetate salt of **27** as a clear oil which was used without further purification.

Part B




Compound 12 was synthesized according to the procedure described in Example 1, using amine 14 in place of (S)-*N*¹,*N*¹,4,4-tetramethylpentane-1,2-diamine. 56% yield. MS (ESI(+)) *m/z* 757.93 (M+H)⁺.

Example 4




15

Part A



To a solution of phenol 16 (2.0 g, 10 mmol, 1 eq) in DMF (5 mL) was added K₂CO₃ (2.1 g, 15 mmol, 1.5 eq) followed by EtI (880 μL, 11 mmol, 1.1 eq). The reaction mixture was stirred at 60 °C for 24 h and then cooled to rt and quenched with water. The mixture was diluted with EtOAc and washed twice with water and then brine. The solution was dried over MgSO₄, filtered and concentrated *in vacuo*. The crude material was purified by silica gel chromatography (10% hexanes/EtOAc) to afford 1.63 g of the desired product 17. Yield 72%.


Part B



Aldehyde 17 (1.6 g, 7.0 mmol, 1 eq) and hydroxylamine hydrochloride (590 mg, 8.5 mmol, 1.2 eq) were dissolved in THF/MeOH (3:2, 10 mL). Water (2 mL) was added and the pH was adjusted to 9 with 6 N KOH. The reaction mixture was stirred at rt for 4 h then NaBH₃CN (890 mg, 14 mmol, 2 eq) was added followed by a crystal of methyl orange. The pH was adjusted to 2 and the resulting ruby red color was maintained for the duration of the reaction by the frequent addition of 1 N HCl. After stirring for 2 h another portion of NaBH₃CN (890 mg, 14 mmol, 2 eq) was added. After stirring for a total of 16 h, the pH of the reaction mixture was brought to 7 and DCM was added. The mixture was washed twice


with water, brine and dried over MgSO_4 . The crude product was used directly in the next reaction without further purification.

Part C




- 5 A solution of hydroxylamine **18** (1.5 g, 6.1 mmol, 1 eq) and methyl glyoxylate (1.07 g, 12.2 mmol, 2 eq) in benzene (30 mL) was heated at reflux with a Dean Stark trap overnight. Excess solvent was removed *in vacuo* and the resulting nitron **19** was taken on crude in the next step.

Part D




- 10 Nitron **19** (2.0 g, 6.3 mmol, 1 eq), allylic alcohol **6** (2.0 g, 9.5 mmol, 1.5 eq) and $\text{Ti}(\text{iOPr})_4$ (2.8 mL, 9.5 mmol, 1.5 eq) were dissolved in toluene (25 mL) and heated in a microwave at 120 °C for 10 min. The reaction mixture was diluted with EtOAc (50 mL) and 3-(dimethylamino)-1,2-propanediol (1 mL) was added. After stirring for 1 h, EtOAc was added and the mixture was washed twice with water, brine, dried over MgSO_4 , filtered and concentrated *in vacuo*. The crude residue was purified by silica gel chromatography (15% hexanes/EtOAc) to afford 944 mg of lactone **20**. Yield 30%.

Part E




- 20 To a solution of **20** (950 mg, 1.9 mmol, 1 eq) in THF (21 mL) was added HF/pyridine (4 mL). The mixture was stirred at rt for 3 h then TMSOMe (25 mL) was added. Solvent was removed *in vacuo* and the crude product was purified by silica gel chromatography (20-100% EtOAc/hexanes) to afford 256 mg of **21**. Yield 35%

Part F



To a flame-dried round bottom flask was added (+)-isopinocampheylamine (235 μ L, 1.33 mmol, 2 eq) and DCM (2.0 mL). Trimethylaluminum (660 μ L, 2.0 M in hexanes, 1.33 mmol, 2 eq) was added and the reaction mixture was stirred at rt for 15 min. Lactone **21** (256 mg, 0.663 mmol, 1 eq) in DCM (8 mL) was added and the solution was stirred at rt. After stirring for 16 h, a saturated solution of Rochelle's salt (5 mL) was added, followed by DCM (5 mL). The mixture was stirred rapidly at rt for 2 h. DCM was added the mixture washed with water, dried over MgSO_4 , filtered and concentrated *in vacuo* to afford an oil. The residue was purified by silica gel chromatography (50-100% EtOAc/hexanes) to afford 200 mg of amide **22** in 56% yield.

Part G




A flask containing aryl bromide **22** (90 mg, 0.17 mmol, 1 eq), 3-amino-5-carboxyphenylboronic acid (60 mg, 0.33 mmol, 2 eq), cesium carbonate (160 mg, 0.50 mmol, 3 eq), potassium acetate (15 mg, 0.17 mmol, 1 eq) and Pd(dppf)Cl_2 (15 mg, 0.017 mmol, 0.1 eq) was purged with argon and DMSO (8 mL) added. The mixture was heated at 60 $^\circ\text{C}$ for 3 h, then additional Pd(dppf)Cl_2 (15 mg, 0.17 mmol, 1 eq) was added and heating continued for 1.5 h. The reaction mixture was partitioned between water (50 mL) and DCM (50 mL) and acidified to pH 4 with 6 M HCl. The layers were separated and the aqueous extracted with DCM (3 x 25 mL). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated *in vacuo* to a brown oil.

The crude oil was dissolved in MeOH (6 mL) and treated with AcOH (30 μ L), 37% formalin (50 μ L, 0.67 mmol, 4 eq) and NaBH_3CN (30 mg, 0.50 mmol, 3 eq). After stirring at rt overnight, the reaction mixture was partitioned between water (40 mL) and DCM (40 mL) and acidified to pH 4 with 6 M HCl. The layers were separated and the aqueous

extracted with DCM (3 x 20 mL). The combined organic layers were dried on Na₂SO₄, filtered and concentrated *in vacuo* to afford a brown oil.

To an aliquot containing 9% of this crude oil (15 mg, 0.024 mmol, 1 eq) in DMF (1 mL) was added (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine (15 μ L, 0.07 mmol, 3 eq) and HBTU (30 mg, 0.07 mmol, 3 eq). The mixture was diluted with MeOH (1 mL) and purified by reverse-phase HPLC (MeCN/water with 40 mM NH₄HCO₃) to give 3 mg of compound **15** as a white solid. Yield 16%. MS (ESI(+)) *m/z* 393.25 (M+2H)²⁺.


Example 5



23

Compound **23** was synthesized according to the procedure described in Example 4, using amine (R)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine. 21% yield. MS (ESI(+)) *m/z* 784.88 (M+H)⁺.


Example 6



24


Compound **24** was synthesized according to the procedure described in Example 4, using amine (S)-*N*¹,*N*¹,4,4-tetramethylpentane-1,2-diamine in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine. 19% yield. MS (ESI(+)) *m/z* 764.95 (M+H)⁺.

Example 7




25

Compound **25** was synthesized according to the procedure described in Example 4, using amine (S)-*N*¹,*N*¹-dimethyl-3-(thiophen-2-yl)propane-1,2-diamine in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine. 16% yield. MS (ESI(+)) *m/z* 396.18 (M+2H)²⁺.

Example 8


26

Part A

10

To a solution of phenol **17** (2.2 g, 12 mmol, 1 eq) in DMF (5 mL) was added K₂CO₃ (2.3 g, 16 mmol, 1.5 eq) followed by (bromomethyl)cyclopropane (1.18 mL, 12.2 mmol, 1.1 eq). The reaction mixture was stirred at 60 °C for 24 h and then cooled to rt and quenched with water. The mixture was diluted with EtOAc and washed twice with water then brine. The solution was dried over MgSO₄, filtered and concentrated *in vacuo*. The crude material was purified by silica gel chromatography (9:1 hexanes/EtOAc) to afford 2.07 g of the desired product **27** as a colorless oil. Yield 73%.

15


Part B

20

Aldehyde **27** (2.07 g, 8.11 mmol, 1 eq) and hydroxylamine hydrochloride (766 mg, 9.74 mmol, 1.1 eq) were dissolved in THF/MeOH (3:2, 10 mL). Water (2 mL) was added and the pH was adjusted to 9 with 6 N KOH. The reaction mixture was stirred at rt for 4 h then NaBH₃CN (1.02 g, 16.2 mmol, 2 eq) was added followed by a crystal of methyl


orange. The pH was adjusted to 2 and the resulting ruby red color was maintained for the duration of the reaction by the frequent addition of 1 N HCl. After stirring for 2 h another portion of NaBH₃CN (1.02 g, 16.2 mmol, 2 eq) was added. After stirring for a total of 16 h, the pH of the reaction mixture was brought to 7 and DCM was added. The mixture was washed twice with water, brine and then dried over MgSO₄, filtered and concentrated *in vacuo*. The crude product was used directly in the next reaction without further purification.

Part C




A solution of hydroxylamine **28** (2.15 g, 7.90 mmol, 1 eq) and methyl glyoxylate (1.39 g, 15.8 mmol, 2 eq) in benzene (30 mL) was heated at reflux with a Dean Stark trap overnight. Excess solvent was removed under reduced pressure and the resulting nitron **29** was taken on crude in the next step.

Part D




Nitron **29** (2.7 g, 7.9 mmol, 1 eq), allylic alcohol **6** (2.6 g, 12 mmol, 1.5 eq) and Ti(iOPr)₄ (3.5 mL, 12 mmol, 1.5 eq) were dissolved in toluene (25 mL) and heated in the microwave at 140 °C for 20 min. The reaction mixture was diluted with EtOAc (50 mL) and 3-(dimethylamino)-1,2-propanediol (1 mL) was added. After stirring for 1 h, EtOAc was added and the mixture was washed twice with water and then brine, dried over MgSO₄, filtered over Celite and concentrated. The crude residue was purified by silica gel chromatography (15% hexanes/EtOAc) to afford 1.35 g of lactone **30**. Yield 33%.

Part E




To a solution of **30** (1.35 g, 2.56 mmol, 1 eq) in THF (21 mL) was added HF/pyridine (4 mL). The mixture was stirred at rt for 3 h then TMSOMe (25 mL) was added. Solvent was removed *in vacuo* and the crude product was purified by flash chromatography (20-100% EtOAc/hexanes) to afford 404 mg of **31**. Yield 38%.

5 Part F



To a flame-dried round bottom flask was added (+)-isopinocampheylamine (325 μ L, 1.83 mmol, 2 eq) and DCM (2.0 mL). Trimethylaluminum (917 μ L, 2.0 M in hexanes, 1.82 mmol, 2 eq) was added and the reaction mixture was stirred at rt for 15 min. The lactone **22** (378 mg, 0.92 mmol, 1 eq) in DCM (8 mL) was added and stirring was continued 16 h. A saturated Rochelle's salt solution (5 mL) was added followed by DCM (5 mL) and the mixture was rapidly stirred at rt for 2 h. DCM was added and the mixture was washed with water, dried over MgSO_4 , filtered and concentrated *in vacuo*. The crude material was purified by silica gel chromatography (50-100% EtOAc/hexanes) to afford 415 mg of amide **32**. Yield 80%.

Part G




A flask containing aryl bromide **32** (90 mg, 0.16 mmol, 1 eq), 3-amino-5-carboxyphenylboronic acid (60 mg, 0.33 mmol, 2 eq), cesium carbonate (160 mg, 0.5 mmol, 3 eq), potassium acetate (15 mg, 0.17 mmol, 1 eq) and Pd(dppf)Cl_2 (15 mg, 0.017 mmol, 0.1 eq) was purged with argon and DMSO (5 mL) added. The mixture was heated at 60 $^{\circ}\text{C}$ for 3 h, then additional Pd(dppf)Cl_2 (15 mg, 0.17 mmol, 1 eq) was added and heating continued for 1.5 h. The reaction mixture was partitioned between water (50 mL) and DCM (50 mL) and acidified to a pH 4 with 6 N HCl. The layers were separated and the aqueous phase was extracted with DCM (3 x 25 mL). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated *in vacuo* to afford a brown oil.

This crude oil was dissolved in MeOH (6 mL) and treated with AcOH (30 μ L), 37% formalin (50 μ L, 0.67 mmol, 4 eq) and NaBH₃CN (32 mg, 0.50 mmol, 3 eq). After stirring at rt for 2 h, the reaction mixture was partitioned between water (40 mL) and DCM (40 mL) and acidified with 6 N HCl until the aqueous layer attained a pH of 4. The layers were separated and the aqueous extracted with DCM (3 x 20 mL). The combined organic layers were dried over Na₂SO₄ and concentrated *in vacuo* to afford a brown oil.

To an aliquot containing 12% of this crude oil (20 mg, 0.031 mmol, 1 eq) in DMF (1 mL) was added (S)-N¹,N¹-dimethyl-3-phenylpropane-1,2-diamine (20 μ L, 0.09 mmol, 3 eq) and HBTU (30 mg, 0.07 mmol, 3 eq). After stirring for 12 h, the mixture was diluted with MeOH (1 mL) and purified by reverse-phase HPLC (MeCN/water with 40 mM NH₅CO₂) to give 3 mg of compound **26** as a white solid. Yield 12%. MS (ESI(+)) *m/z* 406.26 (M+2H)²⁺.


Example 9



33


Compound **33** was synthesized according to the procedure described in Example 8, using amine (S)-N¹,N¹,4,4-tetramethylpentane-1,2-diamine in place of (S)-N¹,N¹-dimethyl-3-phenylpropane-1,2-diamine. Yield 12%. MS (ESI(+)) *m/z* 396.25 (M+2H)²⁺.

Example 10




34


Compound **34** was synthesized according to the procedure described in Example 8, using amine (R)-N¹,N¹,4-trimethylpentane-1,2-diamine in place of (S)-N¹,N¹-dimethyl-3-phenylpropane-1,2-diamine. Yield 13%. MS (ESI(+)) *m/z* 389.19 (M+2H)²⁺.

Example 11**35**

Compound **35** was synthesized according to the procedure described in Example 4, using amine (S)-*N*¹,*N*¹,4-trimethylpentane-1,2-diamine in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine. Yield 17%. MS (ESI(+)) *m/z* 374.24 (M+2H)²⁺.


Example 12**36**

Prepared in an analogous fashion as described in Example 1 using (R)-*N*¹,*N*¹,4,4-tetramethylpentane-1,2-diamine in place of (S)-*N*¹,*N*¹,4,4-tetramethylpentane-1,2-diamine. Yield 4%. MS (ESI(+)) *m/z* 750.5. (M+H)⁺.


Example 13**37**

Prepared in an analogous fashion as described in Example 1 using (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine in place of (S)-*N*¹,*N*¹,4,4-tetramethylpentane-1,2-diamine. Yield 65%. MS (ESI(+)) *m/z* 770.7. (M+H)⁺.


Example 14




38

Part A

5 3-Bromo-2-fluoro-benzoic acid **39** (15 g, 69 mmol, 1 eq) was dissolved in THF (400 mL) followed by the addition of LiAlH₄ (2.9 g, 76 mmol, 1.1 eq) over the course of 5 min. The reaction mixture was stirred vigorously for 12 h, quenched with a saturated Rochelle's salt solution, and stirred for an additional 2 h. The mixture was then partitioned in DCM/saturated NaHCO₃ solution and extracted with DCM (2 x 200 mL). The combined
10 organics were dried over MgSO₄, filtered and concentrated *in vacuo* to afford 11.2 g of alcohol **40**.


Part B

Pyridinium chlorochromate (22.3 g, 103 mmol, 1.91 eq) was dissolved in DCM
15 (140 mL) and 4Å Molecular Sieves (22.3 g) were added to the solution. The reaction mixture was stirred for 30 minutes. A solution of benzyl alcohol **40** (11 g, 54 mmol, 1 eq) in DCM (140 mL) was added dropwise over the course of 5 min. The orange reaction mixture immediately turned dark brown upon addition of alcohol. After stirring for 30 min, the reaction mixture was diluted with 70% Hexane/EtOAc solution (250 mL) and filtered
20 through a plug of silica gel (150 g). The silica gel was washed with Hexane/EtOAc (500 mL), and concentrated *in vacuo* to afford 9.8 g of aldehyde **41** as a white solid. Yield 70% over 2 steps.

Part C


Aldehyde **41** (9.8 g, 48.3 mmol, 1 eq) and hydroxylamine hydrochloride (3.97 g, 57.9 mmol, 1.2 eq) were dissolved in THF/MeOH (3:2, 40 mL) at rt. Water (25 mL) was added and the pH was adjusted to 9 with 6 N KOH. After stirring for 16 h, NaBH₃CN (3.03 g, 48.3 mmol, 1 eq) was added followed by a crystal of methyl orange. The pH was adjusted to 2 and the resulting ruby red color of the reaction mixture was maintained for the duration of the reaction by the frequent addition of 1 N HCl. After stirring for 2 h, another portion of NaBH₃CN (3.03 g, 48.2 mmol, 1 eq) was added. After stirring for a total of 16 h, the reaction mixture was adjusted to pH 7 with 6 N KOH. The mixture was extracted with DCM (2 x 100 mL) and the combined organics were washed with water (50 mL), brine (20 mL), dried over MgSO₄, filtered and concentrated *in vacuo* to afford 8.1 g of hydroxylamine **42**. The crude material was used without further purification. Crude yield 76%.

Part D



To a solution of hydroxylamine **42** (8.1 g, 36.8 mmol, 1 eq) in ether (200 mL) at rt was added methyl glyoxylate methyl acetal (5.3 g, 44.2 mmol, 1.2 eq), followed by the addition of calcium chloride (4.9 g, 44.2 mmol, 1.2 eq). After stirring for 3 h, the reaction mixture was filtered through celite, washed with DCM (2 x 200 mL), and concentrated *in vacuo* to a white solid to afford 10.5 g of **43**. The crude material was used without further purification. Yield 98%.


Part E



Nitron **43** (10.5 g, 36 mmol, 1 eq), allylic alcohol **6** (9.8 g, 45 mmol, 1.2 eq) and Ti(iOPr)₄ (13 g, 45 mmol, 1.2 eq) were dissolved in THF (50 mL) and heated in a microwave at 140 °C for 30 min. The reaction mixture was allowed to cool to rt and then diluted with EtOAc (150 mL), water (150 mL) and 3-(dimethylamino)-1,2-propanediol (5 mL). After stirring for 2 h, the reaction mixture was extracted with EtOAc (2 x 100 mL). The combined organics were washed with water (3 x 50 mL) brine (50 mL), dried over

MgSO₄, filtered through Celite and concentrated *in vacuo*. The crude residue was purified by silica gel chromatography (5–25% EtOAc/hexane) to afford 11.45 g of lactone 44. Yield 67%.


Part F




5

To a 0 °C solution of TBS-protected lactone **44** (11.2 g, 24 mmol, 1 eq) in THF (60 mL) was added 6 N HCl (6 mL). The reaction mixture was stirred for 2 h and then quenched by the addition of a saturated sodium bicarbonate solution (30 mL). The reaction mixture was extracted with EtOAc (2 x 100 mL) and the combined organics were dried over 10 MgSO₄, filtered and concentrated *in vacuo*. The crude residue was purified by silica gel chromatography (30-70% EtOAc/hexane) to afford 7.7 g of alcohol **45**. Yield 91 %.


Part G



To a solution of alcohol **45** (810 mg, 1.99 mmol, 1 eq) in DCM (20 mL) at 0 °C was added trifluoromethanesulfonic anhydride (0.4 mL, 2.4 mmol, 2.4 eq) dropwise over the course of 5 min. After stirring for 30 minutes, the reaction mixture was diluted with DMF (5 mL) followed by the addition of sodium azide (388 mg, 5.97 mmol, 3 eq) in a single portion. The reaction mixture was stirred for 12 h and then quenched by the addition of water, extracted with EtOAc (2 x 100 mL) and the combined organics were dried over MgSO₄, filtered and concentrated *in vacuo* to afford an oil. The crude oil was purified by silica gel chromatography (30-80% EtOAc/hexane) to afford 758 mg of **46**. Yield 88%.

Part H


To a 0 °C solution of (+)-isopinocampheylamine (1.98 mL, 11.8 mmol, 6 eq) in THF (20 mL) was added DIBAL (9.8 mL, 1 M in toluene, 9.8 mmol, 5 eq) and the reaction mixture was stirred for 2 h. A solution of lactone **46** (758 mg, dissolved in 5 mL THF, 1.97 mmol, 1 eq) was then added dropwise over the course of 5 min. After stirring for 2 h, the reaction was diluted with a saturated aqueous solution of Rochelle's salt and EtOAc and then stirred vigorously for 5 h. The mixture was extracted with EtOAc (3 x 100 mL) and the combined organics were dried over MgSO₄, filtered and concentrated *in vacuo* to afford an oil. The crude oil was purified by silica gel chromatography (30-70% EtOAc/Hexane) to afford 500 mg of amide **47**. Yield 47%.

Part I

(3-Amino-5-carboxyphenyl)boronic acid **48** (15 g, 83 mmol, 1 eq) and pinacol (29 g, 249 mmol, 3 eq) were combined with THF (72 mL) and heated to reflux. After heating at reflux for 48 h, the reaction mixture was cooled to rt and then concentrated to an orange oil. The orange oil was dissolved in MeOH (250 mL) and the solution was cooled to 0 °C using an ice bath. Formaldehyde (67 g of a 37% solution in water, 829 mmol, 10 eq) was added followed by AcOH (30 g, 497 mmol, 6 eq) and the portion-wise addition of NaBH₃CN (7.8 g, 124 mmol, 3 eq). The ice bath was removed and the reaction was allowed to warm slowly to rt. After stirring for 12 h, the reaction mixture was filtered through celite, washed with EtOAc (2 x 100 mL) and the combined filtrates were concentrated to an orange oil. The oil was dissolved in EtOAc (200 mL), diluted with water and the pH was adjusted to pH 10 using 6 N NaOH and then extracted with EtOAc (2 x 200 mL). The aqueous phase was adjusted to pH 4 using 6 N HCl and extracted with DCM (2 x 200 mL). The combined DCM extracts were then dried over MgSO₄, filtered and concentrated to


provide 11 g of **49** as an off-white solid. This material was used without further purification.

Part J



5 Isoxazolidine **47** (335 mg, 0.622 mmol, 1 eq), pinacolboronate **49** (362 mg, 1.24 mmol, 2 eq), Pd(dppf)Cl₂ (102 mg, 0.124 mmol, 0.1 eq), potassium acetate (79 mg, 0.81 mmol, 1.3 eq), and cesium carbonate (608 mg, 1.87 mmol, 3 eq) were dissolved in anhydrous DMSO (5 mL) and flushed with argon under positive pressure. The reaction mixture was stirred at 60 °C for 2 h, then allowed to cool to rt. The reaction mixture was
 10 diluted with EtOAc (20 mL) and water (10 mL) and the aqueous phase was adjusted to pH 4 with 6 M HCl. The reaction mixture was extracted with EtOAc (3 x 100 mL) and the combined organics were dried over MgSO₄, filtered and concentrated *in vacuo* to afford an oil. The crude oil was purified by silica gel chromatography (10-35% acetone/hexane) to afford 110 mg of biphenyl acid **50**. Yield 28%.


15 Part K



Biphenyl acid **50** (40 mg, 0.064 mmol, 1 eq) and HBTU (49 mg, 0.13 mmol, 2 eq) were dissolved in DMF (1 mL) followed by the addition of amine (S)-N¹,N¹-dimethyl-3-phenylpropane-1,2-diamine (23 mg, 0.13 mmol, 2 eq). The reaction mixture was stirred for
 20 2 h, diluted with EtOAc and quenched with a saturated aqueous NaHCO₃ solution. The reaction mixture was extracted with EtOAc (3 x 100 mL) and the combined organics were dried over MgSO₄, filtered and concentrated *in vacuo* to afford an oil. The crude oil was


purified by silica gel chromatography (2-5% MeOH/DCM) to afford 23 mg of biphenyl amide **51**. Yield 45%.

Part L




Azide **51** (18 mg, 0.023 mmol, 1 eq) and dithiothreitol (11 mg, 0.07 mmol, 3 eq) were dissolved in DMF (1 mL) followed by the addition of DBU (12 μ L, 0.07 mmol, 3 eq). The reaction mixture was stirred for 1 h, then purified directly by reverse-phase HPLC (MeCN/water with 40 mM NH_4HCO_3), to yield 7 mg of compound **38**. MS ((ESI(+)) m/z 757.4 ($\text{M}+\text{H}$)⁺.

Example 15



Prepared in an analogous fashion as described in Example 13 using 2-amino-N,N-dimethyl-3-(thiazol-4-yl)propanamide in place of (S)-N¹,N¹-dimethyl-3-phenylpropane-1,2-diamine. Yield 29%. MS (ESI(+)) m/z 778.4. ($\text{M}+\text{H}$)⁺.


Example 16




53

Prepared in an analogous fashion as described in Example 13 using *N*¹,*N*¹-dimethylethane-1,2-diamine in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine. Yield 49%. MS (ESI(+)) *m/z* 667.4. (M+H)⁺.


5

Example 17

54


Part A

To a solution of lactone **8** (1.8 g, 4.3 mmol, 1 eq) in DCM (42 mL) at 0 °C was added triflic anhydride (0.87 mL, 5.15 mmol, 1.2 eq) dropwise over the course of 5 min. After 30 min, the reaction mixture was diluted with DMF (10 mL) followed by the addition of sodium azide (0.84 g, 12.9 mmol, 3 eq) in a single portion. The reaction mixture was removed from the ice bath and allowed to warm to rt. After stirring for 12 h, the reaction was quenched by the addition of water, extracted with EtOAc (2 x 100 mL), dried over MgSO₄, filtered and concentrated *in vacuo* to afford an oil. The crude oil was purified by silica gel chromatography (30-80% EtOAc/hexane) to afford 1.7 g of azide **55**. Yield 89%.

Part B


To a 0 °C solution of (+)-isopinocampheylamine (5.75 mL, 34.3 mmol, 6 eq) in THF (60 mL) was added DIBAL (14.3 mL, 2 M in toluene, 28.6 mmol, 5 eq). After stirring for 2 h, the reaction mixture was added to a solution of lactone **55** (2.54 g, 5.72 mmol, 1 eq) dissolved in THF (10 mL). The combined reaction mixture was stirred for 2 h, then diluted with a saturated solution of Rochelle's salt and EtOAc. After stirring for 5 h the reaction mixture was extracted with EtOAc (3 x 100 mL), dried over MgSO₄, filtered and concentrated *in vacuo* to afford an oil. The crude oil was purified by silica gel chromatography (30-70% EtOAc/hexane) to afford 2.79 g of azide **56**. Yield 82%.

Part C



Isoxazolidine **56** (300 mg, 0.50 mmol, 1 eq), pinacol boronate **49** (368 mg, 1.26 mmol, 2.5 eq), Pd(dppf)Cl₂ (82 mg, 0.10 mmol, 0.2 eq), potassium acetate (65 mg, 0.657 mmol, 1.3 eq), and cesium carbonate (494 mg, 1.52 mmol, 3 eq) were dissolved in anhydrous DMSO (5 mL) and the reaction vessel was flushed under positive argon pressure for 15 min. The reaction mixture was stirred at 60 °C for 2 h, then allowed to cool to rt. The reaction mixture was diluted with EtOAc (20 mL) and water (10 mL) and the aqueous phase was adjusted to pH 4 with 6 N HCl. The reaction mixture was extracted with EtOAc (3 x 100 mL) and the combined organics were dried over MgSO₄, filtered and concentrated *in vacuo* to afford an oil. The crude oil was purified by gradient silica gel chromatography (10-35% acetone/hexanes) to afford 170 mg of biphenyl acid **57**. Yield 53%.


Part D



Biphenyl acid **57** (320 mg, 0.50 mmol, 1 eq) and HBTU (287 mg, 0.756 mmol, 1.5 eq) were dissolved in DMF (5 mL) followed by addition of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine (180 mg, 1.0 mmol, 2 eq). After stirring for 2 h, the reaction mixture was diluted with EtOAc (100 mL) and a saturated sodium bicarbonate solution.

- 5 The mixture was extracted with EtOAc (3 x 100 mL), dried over MgSO₄, filtered and concentrated to an oil. The crude oil was purified by silica gel chromatography (2 -5% MeOH/DCM) to afford 197 mg of azide **58**. Yield 49%.


Part E



- 10 Azide **58** (15 mg, 0.018 mmol, 1 eq) and dithiothreitol (8.7 mg, 0.054 mmol, 3 eq) were dissolved in DMF (1 mL) followed by addition of DBU (8.5 μ L, 0.054 mmol, 3 eq). After stirring for 1 h, the reaction mixture was purified directly by reverse-phase HPLC (CH₃CN/water with 40 mM NH₄HCO₃), to yield 7 mg of amine **54**. Yield 47%. MS ((ESI(+)) *m/z* 769.9 (M+H)⁺.

15

Example 18




59

Prepared in an analogous fashion as described in Example 17 using (S)-*N*¹,*N*¹-trimethylpentane-1,2-diamine in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine.


- 20 43 % Yield. MS (ESI(+)) *m/z* 735.1 (M+H)⁺.

Example 19




60

Prepared in an analogous fashion as described in Example 17 using (R)-(1-ethylpyrrolidin-2-yl)methylamine in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine. Yield 43%. MS (ESI(+)) *m/z* 733.1 (*M*+*H*)⁺.

Example 20

61


Prepared in an analogous fashion as described in Example 17 using (S)-*N*¹,*N*¹-4,4-tetramethylpentane-1,2-diamine in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine. Yield 27 %. MS (ESI(+)) *m/z* 749.1 (*M*+*H*)⁺.

Example 21

62

Prepared in an analogous fashion as described in Example 17 using 2-phenylethylamine in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine. Yield 43%. MS (ESI(+)) *m/z* 711.9 (M+H)⁺.


5

Example 22

63

Prepared in an analogous fashion as described in Example 17 using *N*¹,*N*¹-dimethylethane-1,2-diamine in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine. Yield 62%. MS (ESI(+)) *m/z* 679.1 (M+H)⁺.

10


Example 23

64

Prepared in an analogous fashion as described in Example 17 using 2-(pyridin-2-yl)ethylamine in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine. Yield 37%. MS (ESI(+)) *m/z* 712.9 (M+H)⁺.

15

Example 24




65

Prepared in an analogous fashion as described in Example 17 using (S)-methyl 2-amino-3-phenylpropanoate in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine.

5 Yield 37%. MS (ESI(+)) *m/z* 770.1 (M+H)⁺.


Example 25



66

Prepared in an analogous fashion as described in Example 17 using *N*-methoxymethylamine in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine. Yield
10 87%. MS (ESI(+)) *m/z* 652.2 (M+H)⁺.


Example 26



67

Prepared in an analogous fashion as described in Example 17 using (S)-2-amino-*N,N*-dimethyl-3-(thiazol-4-yl)propylamide in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine. Yield 11%. MS (ESI(+)) *m/z* 790.0 (M+H)⁺.


Example 27



68

Prepared in an analogous fashion as described in Example 17 using (R)-*N*¹,*N*¹,4-trimethylpentane-1,2-diamine in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine. Yield 54%. MS (ESI(+)) *m/z* 735.1 (M+H)⁺.


Example 28




69

Prepared in an analogous fashion as described in Example 17 using dimethylamine in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine. Yield 15%. MS (ESI(+)) *m/z* 635.8 (M+H)⁺.


Example 29

**70**

Prepared in an analogous fashion as described in Example 17 using methylamine in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine. Yield 42%. MS (ESI(+)) *m/z* 622.0 (M+H)⁺.


Example 30**71**

Prepared in an analogous fashion as described in Example 17 using compound **57** in place of compound **58**. Yield 33%. MS (ESI(+)) *m/z* 609.1 (M+H)⁺.


Example 31**72**

Prepared in an analogous fashion as described in Example 17 using (S)-*N*¹,*N*¹,4-trimethylpentane-1,2-diamine in place of (S)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine. Yield 54%.. MS (ESI(+)) *m/z* 735.1 (M+H)⁺.

Example 32




73



Compound **61** (4 mg, 0.006 mmol, 1 eq) was dissolved in DCM (1 mL), followed by the addition of pyridine (1.3 μ L, 0.016 mmol, 1.5 eq) and acetic anhydride (1.5 μ L, 0.016 mmol, 1.5 eq). After 1 h, the reaction mixture was quenched with water, extracted with DCM, dried over MgSO_4 , filtered and concentrated *in vacuo* to afford an oil. The crude material was dissolved in MeOH (1 mL) and purified on reverse-phase HPLC ($\text{CH}_3\text{CN}/\text{water}$ 40 mM NH_4HCO_3) to yield 2 mg of acetamide **73**. Yield 47%. MS ((ESI(+)) *m/z* 791.5 (M+H)⁺.


Example 33



74

Prepared in an analogous fashion as described in Example 32 using compound 62 in place of compound 61 and trifluoroacetic anhydride in place of acetic anhydride. Yield 40%. MS (ESI(+)) m/z 808.4 (M+H)⁺.


Example 34



75

Prepared in an analogous fashion as described in Example 31 using compound 70 in place of compound 61 and succinic anhydride in place of acetic anhydride. Yield 28%. MS (ESI(+)) m/z 722.1 (M+H)⁺.


Example 35



76

Prepared in an analogous fashion as described in Example 31 using compound **62** in place of compound **61**. Yield 28%. MS (ESI(+)) m/z 754.4 (M+H)⁺.

Example 36




5


77

Prepared in an analogous fashion as described in Example 31 using compound **70** in place of compound **61**. Yield 47%. MS (ESI(+)) m/z 664.4 (M+H)⁺.

Example 37




10

78

To a solution of mono-methylsuccinate (2.5 mg, 19 μ mol, 3 eq) in DMF (0.5 mL) was added HBTU (7.3 mg, 19 μ mol, 3 eq). After stirring for 15 minutes, compound **70** (4 mg, 64 μ mol, 1 eq) was added as a THF solution (0.5 mL). The crude reaction mixture was


purified directly by reverse-phase HPLC (MeCN/water with 40 mmol NH_4HCO_3) to yield 2 mg of amide 78. Yield 42%. MS ((ESI(+)) m/z 736.1 ($\text{M}+\text{H}$)⁺.

Example 38



79

5




70

79


Compound 70 (4 mg, 6 μmol , 1 eq) was dissolved in DCM (1 mL) followed by the addition of phenyl isocyanate (1 μL , 8 μmol , 1.25 eq). After stirring for 30 min, the reaction mixture was purified directly by reverse-phase HPLC (MeCN/water with 40 mmol NH_4HCO_3) to yield 2 mg of urea 76. Yield 42%. MS ((ESI(+)) m/z 741.2 ($\text{M}+\text{H}$)⁺.

10

Example 39





80




Compound **62** (10 mg, 0.014 mmol, 1 eq) was dissolved in MeOH (1 mL) followed by the addition of formaldehyde (2 mg, 0.07 mmol, 5 eq). After stirring for 30 min, NaBH₃CN (2.6 mg, 0.04 mmol, 3 eq) was then added in a single portion. The reaction mixture was purified by reverse-phase HPLC (MeCN/water with 40 mmol NH₄HCO₃) to yield 5 mg of dimethylamine **77**. Yield 50%. MS ((ESI(+)) *m/z* 740.5 (M+H)⁺.


Example 40




Compound **70** (4 mg, 6 μmol, 1 eq) was dissolved in MeOH (1 mL) followed by the addition of aryl aldehyde (3 mg, 0.02 mmol, 3 eq). After stirring for 3 h, NaBH₃CN (1 mg, 0.02 mmol, 3 eq) was added. The reaction mixture was stirred for an additional 30 min, then purified directly by reverse-phase HPLC (MeCN/water with 40 mmol NH₄HCO₃) to yield 2.5 mg of benzylamine **81**. Yield 51%. MS ((ESI(+)) *m/z* 756.2 (M+H)⁺.

Example 41**82**

Prepared in an analogous fashion as described in Example 40 using
5 isonicotinaldehyde in place of 4-formylbenzoic acid. Yield 37%. MS (ESI(+)) m/z 713.2 (M+H)⁺.


Example 42**83**

10 Prepared in an analogous fashion as described in Example 40 using picolinaldehyde
in place of 4-formylbenzoic acid. Yield 44%. MS (ESI(+)) m/z 713.2 (M+H)⁺.

Example 43**84**

Prepared in an analogous fashion as described in Example 40 using 3-formylbenzoic acid in place of 4-formylbenzoic acid. Yield 37%. MS (ESI(+)) m/z 756.2 (M+H)⁺.

Example 44




5

85


Prepared in an analogous fashion as described in Example 40 using benzaldehyde in place of 4-formylbenzoic acid. Yield 33%. MS (ESI(+)) m/z 712.2 (M+H)⁺.

Example 45




10

86




To a solution of compound **61** (8 mg, 0.01 mmol, 1 eq) in DCM/DMF (4:1, 1 mL), was added Et₃N (5 μ L, 0.03 mmol, 3 eq) and HBTU (10 mg, 0.03 mmol, 3 eq). After stirring for 3 h, the reaction mixture was purified directly by reverse-phase HPLC (MeCN/water with 40 mM NH₄HCO₃) to yield 4 mg of **86**. Yield 44%. MS ((ESI(+)) m/z 810.4 (M+H)⁺.

Example 46




Part A




Prepared in an analogous fashion as described in Example 17 step D using methylamine in place (S)-N¹,N¹-dimethyl-3-phenylpropane-1,2-diamine. Yield 26%.

Part B




Azido alcohol **88** (150 mg, 0.232 mmol, 1 eq) was dissolved in DCM (2.5 mL), followed by the addition of pyridine (55 mL, 0.695 mmol, 3 eq) and para-nitro chloroformate (140 mg, 0.695 mmol, 3 eq). After stirring for 5 h, the reaction mixture was diluted with DCM (20 mL), water (20 mL) and the mixture was extracted with DCM (2 x 20 mL), dried over MgSO_4 , filtered and concentrated *in vacuo* to afford an oil. The crude material was purified by silica gel chromatography (50-90% EtOAc/hexane) to yield 142 mg of *p*-nitrocarbonate **89**. Yield 75%.

10 Part C




To a solution of azido carbonate **89** (15 mg, 0.018 mmol, 1 eq) in THF (0.5 mL) was added dimethylamine (18 μL , 2 M in THF, 0.036 mmol, 2 eq). The reaction mixture immediately turned bright yellow. After stirring for 6 h, the reaction mixture was diluted with EtOAc (10 mL) and washed with water (5 mL). The aqueous layer was extracted with EtOAc (2 x 15 mL) and the combined organics were dried over MgSO_4 , filtered and concentrated *in vacuo* to afford a yellow oil. The crude oil was used directly in next step.

Part D



To a solution of carbamate **90** (13 mg, 0.018 mmol, 1 eq) in THF (0.5 mL) was added dithiothreitol (8 mg, 0.054 mmol, 3 eq) in DMF (0.5 mL) followed by the addition of DBU (8 mg, 0.054 mmol, 3 eq). After stirring for 1 h, the reaction mixture was purified
 5 directly by reverse-phase HPLC (MeCN/water with 40 mmol NH_4HCO_3) to yield 5 mg of compound **87**. Yield 40%. MS ((ESI(+)) m/z 693.4 ($\text{M}+\text{H}$)⁺.


Example 47



91


10 Prepared in an analogous fashion as described in Example 46 using 2-phenylethylamine in place of dimethylamine. Yield 42%. MS (ESI(+)) m/z 769.5 ($\text{M}+\text{H}$)⁺.

Example 48





92

Prepared in an analogous fashion as described in Example 46 using *N*¹,*N*¹-dimethylethane-1,2-diamine in place of dimethylamine. Yield 30%. MS (ESI(+)) *m/z* 736.4 (M+H)⁺.

Example 49**93**


Prepared in an analogous fashion as described in Example 46 using methylamine in place of dimethylamine. Yield 42%. MS (ESI(+)) *m/z* 679.4 (M+H)⁺.

Example 50**94**

To a solution of amine 70 (10 mg, 0.02 mmol, 1 eq) in EtOH (1 mL) was added acetaldehyde (0.5 mg, 0.01 mmol, 0.5 eq) followed by NaBH₃CN (2 mg, 0.03 mmol, 1.5 eq) and finally AcOH (1 mg, 0.02 mmol, 1 eq). After stirring for 12 h, an additional portion of acetaldehyde (1 mg, 0.02 mmol, 1 eq) and NaBH₃CN (2 mg, 0.03 mmol, 1.5 eq) was added and the reaction was stirred at rt for 4 h. The crude material was purified directly by

reverse-phase HPLC (MeCN/water with 40 mM of NH_4HCO_3) to yield 2 mg of **91**. Yield 19%. MS (ESI(+)) m/z 650.81. $(\text{M}+\text{H})^+$.


Example 51



95

Amine **95** was prepared in an analogous fashion as described in Example 50 using isobutyraldehyde in place of acetaldehyde to yield 3 mg of **95**. Yield 28%. MS (ESI(+)) m/z 678.84. $(\text{M}+\text{H})^+$.


Example 52



96

Amine **96** was prepared in an analogous fashion as described in Example 50 using cyclopropylaldehyde in place of acetaldehyde to yield 3 mg of **95**. Yield 28%. MS (ESI(+)) m/z 676.9. $(\text{M}+\text{H})^+$.


Example 53




97

Amine **97** was prepared in an analogous fashion as described in Example 50 using acetone in place of acetaldehyde to yield 3 mg of **97**. Yield 28%. MS (ESI(+)) m/z 664.8. (M+H)⁺.

Example 54




5



To a solution of amine **70** (10 mg, 0.02 mmol, 1 eq) in EtOH (1 mL) was added excess isobutyraldehyde (6 mg, 0.08 mmol, 4 eq) followed by NaBH₃CN (2 mg, 0.03 mmol, 1.5 eq) and finally AcOH (1 mg, 0.02 mmol, 1 eq). After stirring for 2 h, the reaction mixture was purified by reverse-phase HPLC (MeCN/water with 40 mM of NH₄HCO₃) to yield 2 mg of **98**. Yield 18%. MS (ESI(+)) m/z 734.9. (M+H)⁺.

10


Example 55



15


Amine **99** was prepared in an analogous fashion as described in Example 53 using formaldehyde in place of isobutyraldehyde to yield 2 mg of **99**. Yield 38%. MS (ESI(+)) m/z 650.81. (M+H)⁺.

Example 56




100

Part A




Aldehyde **102** (12 g, 33 mmol, 1 eq) and hydroxylamine hydrochloride (2.7 g, 39 mmol, 1.18 eq) were dissolved in THF/MeOH (3:1, 60 mL). Water (2 mL) was added and the pH was adjusted to 9 with 6 N KOH. The reaction mixture was stirred at rt overnight and then NaBH₃CN (3.1 g, 49 mmol) was added followed by a crystal of methyl orange. The pH was adjusted to 3 and the resulting ruby red color was maintained for the duration of the reaction by the frequent addition of 1 N HCl. After stirring for 2 h another portion of NaBH₃CN (1 g, 13 mmol, 0.4 eq) was added. The solution was stirred for 16 h and then neutralized to pH 7 and diluted with DCM. The mixture was washed with water (3 x 10 mL), brine and dried over MgSO₄, filtered and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (50-100% EtOAc/hexane) to afford 8 g of compound **103**. Yield 64%.

Part B




A solution of hydroxylamine **103** (5 g, 13 mmol, 1 eq) and methyl glyoxylate (1 g, 16 mmol) in benzene (15 mL) was heated at reflux with a Dean Stark trap for 3 h. Excess solvent was removed *in vacuo* and the resulting nitrone **104** (6 g) was taken on crude in the next step.

Part C



Nitron **104** (5 g, 11 mmol, 1 eq), allylic alcohol (2 g, 11 mmol, 1 eq) and $\text{Ti}(\text{iOPr})_4$ (4 g, 4 mL, 13 mmol, 1.18 eq) were dissolved in toluene (40 mL) and heated in a microwave at 120 °C for 10 min. The reaction mixture was diluted with EtOAc (10 mL) and 3-(dimethylamino)-1,2-propanediol (4 mL) was added. After stirring for 2 h, EtOAc (10 mL) was added and the mixture was washed with water (3 x 10 mL) then brine (10 mL), dried over MgSO_4 , filtered over Celite® and concentrated *in vacuo*. The crude residue was purified by silica gel chromatography (10-30% Hexane-EtOAc) to afford 2.5 g of compound **105**. Yield 35%.

Part D




To a solution of PMB ether **105** (2 g, 3 mmol, 1 eq) in DCM (150 mL) was added TFA (3 g, 31 mmol, 10.33 eq) dropwise at 0 °C. The solution was stirred for 1.5 h and quenched with saturated NaHCO_3 (60 mL). The organic phase was separated, washed with

brine, dried over Na_2SO_4 , filtered and concentrated *in vacuo* to afford an oil. The resulting oil was purified by silica chromatography (10-30% hexane-EtOAc) to afford 1.2 g of compound **106** Yield 74%.

Part E


5



To a solution of (+)-isopinocampheylamine (0.6 g, 0.7 mL, 4 mmol, 2 eq) in DCM (10 mL) at rt was added trimethylaluminum (0.4 g, 3 mL, 2 M solution in toluene, 6 mmol, 3 eq) dropwise over 2.5 min. The solution was stirred at rt for 10 min prior to the dropwise addition of a solution of lactone **106** (1 g, 2 mmol, 1eq) in DCM (15 mL). The reaction was stirred for 24 h, diluted with DCM (125 mL) and a saturated solution of Rochelle's salt (125 mL). The mixture was vigorously stirred for 2 h until two phases formed. The layers were separated and the organic phase was washed with water, brine, dried over MgSO_4 , filtered and concentrated *in vacuo* to afford a solid. The solid was purified by silica gel chromatography (25% hexane/EtOAc) to afford 0.5 g of compound **107**. Yield 39%.

15

Part F




Phenol **107** (187 mg, 0.27 mmol, 1 eq) was dissolved in DMF (3.5 mL) and treated with K_2CO_3 (111 mg, 0.8 mmol, 3 eq), and allylbromide (49 mg, 0.4 mmol, 1.48 eq). The solution was stirred for 2.5 h, diluted with water and extracted with ether (3×4 mL). The organic phase was separated, dried over Na_2SO_4 , filtered and concentrated *in vacuo* to afford a solid. The resulting solid was dissolved in THF/ Et_3N (1:1, 6 mL) and treated with a solution of HF/pyridine (1 mL) at 0 °C. The solution was stirred for 1 h at rt and quenched with TMSOMe (25 mL) and concentrated *in vacuo* to afford a solid which was purified by

20


silica gel chromatography (20% DCM/hexane) to afford 0.21 g of compound **108** as a white solid. Yield 67%.

Part G



To a solution of the TBS ether **108** (0.25 g, 0.35 mmol, 1 eq) in THF (5 mL) at 0 °C was added pyridine (6 mL) and a solution of HF/pyridine (1.2 mL). The reaction was allowed to warm to rt over 1 h while stirring. The reaction solution was quenched with an excess of TMSOMe (30 mL) and stirred for an additional 30 min. The reaction was then concentrated to an oil and purified by silica gel chromatography to yield 0.21 g of compound **109**. Yield 67%.


Part H



To a solution of the alkene **109** (0.14 g, 0.2 mmol, 1 eq) in *t*-BuOH (16 mL), THF (8 mL) and H₂O (2 mL) was added NMO (80 mg, 0.8 mmol, 4 eq) followed by the dropwise addition of OsO₄ (0.21 g, 2.9 mL, 2.5 % solution in 2-methyl-2-propanol, 0.02 mmol, 0.1 eq). After 3 h, the reaction mixture was diluted with DCM (5 mL), brine and 10% Na₂S₂O₃ and the organic phase was separated. The aqueous phase was extracted with DCM (2 × 60 mL), and the combined organics were dried over Na₂SO₄, filtered and concentrated *in vacuo* to afford a solid. The solid was dissolved in THF/water (10:1, 1.2 mL) and treated with periodic acid (80 mg, 0.4 mmol, 2 eq) in single portion and stirred for 12 h. The solution was diluted with DCM (5 mL) and washed with brine, dried over Na₂SO₄, filtered and *in vacuo* to afford an oil. The resulting oil was suspended in MeOH (5 mL) and treated with acetic acid (100 uL), dimethylamine (20 mg, 2 M in THF, 0.5 mmol, 2.5 eq) and NaBH₃CN (30 mg, 0.5 mmol, 2.5 eq) and stirred for 12 h. The solution was diluted with 0.1 M NaOH (1 mL), saturated NaCl (1 mL) and extracted with EtOAc (3 × 3 mL). The combined organic was concentrated *in vacuo* to afford an oil. The oil was purified


by silica gel chromatography (DCM/MeOH/AcOH, 99.5:0:0.5 to 97.5:2:0.5) to afford 75 mg of compound **110**. Yield 60%.

Part I




- 5 A flask containing aryl iodide **110** (25 mg, 0.04 mmol, 1 eq), pinacolboronate **49** (17 mg, 0.06 mmol, 1.5 eq), potassium acetate (5 mg, 0.048 mmol, 1.2 eq), cesium carbonate (39 mg, 0.12 mmol, 3 eq) and Pd(dppf)Cl₂ (6.5 mg, 8 μmol, 0.2 eq) was purged with argon and DMSO (2 mL) was added. The mixture was heated to 70 °C for 3 h and then cooled to rt. The solution was diluted with DCM (10 mL), water (5 mL) and the pH was adjusted to 6.8 with 0.1 N HCl. The aqueous phase was extracted with DCM (2 x 10 mL) and the combined organics were washed with brine (20 mL), dried over MgSO₄, filtered and concentrated *in vacuo* to afford a dark brown oil. The resulting oil was purified by flash chromatography (2.5-5% CH₂Cl₂/MeOH) to afford 15 mg of compound **111** as a yellow solid. Yield 57%.

15 Part J




- To a solution of **111** (8 mg, 0.02 mmol, 1 eq) in DMF (1.5 mL) was added HBTU (9 mg, 0.02 mmol, 1 eq), amine **112** (4 mg, 0.02 mmol, 1 eq) and Et₃N (4 mg, 5 uL, 0.04 mmol, 2 eq). The solution was stirred for 2 h, diluted with water (0.5 mL) and purified by reverse-phase HPLC (MeCN/water with 40 mM NH₄HCO₂) to yield 7 mg of compound **100**. Yield 67%. MS (ESI(+)) *m/z* 828 (M+H)⁺.

Example 57




113

Prepared in an analogous fashion as described in Example 1 using (R)-*N*¹,*N*¹-dimethyl-3-phenylpropane-1,2-diamine in place of (S)-*N*¹,*N*¹,4,4-tetramethylpentane-1,2-diamine. Yield 65%. MS (ESI(+)) *m/z* 770.7. (M+H)⁺.

Example 58

114


Prepared in an analogous fashion as described in Example 31 using compound 60 in place of compound 72 and succinic anhydride in place of acetic anhydride. 65 % Yield. MS (ESI(+)) *m/z* 770.7 (M+H)⁺.

Example 59



115

Prepared in an analogous fashion as described in Example 37 using compound 61 in place of compound 70 and 3-(dimethylamino)propanoic acid in place of monomethylsuccinate. 28 % Yield. MS (ESI(+)) *m/z* 849.6 (M+H)⁺.

Example 60

**116**


Prepared in an analogous fashion as described in Example 54 using compound 61 in place of compound 70 and formaldehyde in place of isobuteraldehyde. 36 % Yield. MS (ESI(+)) m/z 777.6 (M+H)⁺.

Example 61**117****Part A**

10

To a mixture of MgCl₂ (2.0 g, 20.9 mmol, 2 eq) and paraformaldehyde (0.943 g, 31.4 mmol, 3 eq) in THF (50 mL) under argon, was added triethylamine (2.92 mL, 20.9 mmol, 2 eq). The mixture was stirred at room temperature, under argon, for 10 min and phenol 118 (2.00 g, 10.5 mmol, 1 eq) was added. The reaction was heated to reflux for 2 h then allowed to cool to rt. Diethyl ether (100 mL) was added and the solution was washed with HCl 1 N (3 x 100 mL), dried over Na₂SO₄, filtered and concentrated to dryness to afford crude 119 (2.07 g) which was taken on without further purification. Yield 90%.

15


Part B

20

To a solution of crude phenol 119 (2.1 g, 9.5 mmol, 1 eq) in DMF (50 mL) at rt was added K₂CO₃ (1.88 g, 13.6 mmol, 1.4 eq) followed by MeI (0.848 mL, 13.6 mmol, 1.4 eq).


The reaction mixture was stirred at 40 °C under argon for 5 h and then quenched with 6 N HCl (10ml). The mixture was diluted with 1 N HCl (250 mL) and extracted with DCM (3 x 100 mL). Pooled organics were washed with water (3 x 100mL) then brine (100mL). The solution was dried over Na₂SO₄, filtered and concentrated to dryness. The residue was
 5 purified by flash chromatography (EtOAc in Hexane, gradient 0% to 20%) to afford 1.02g of product **120**. Yield 42%.

Part C



Aldehyde **120** (1.0 g, 4.37 mmol, 1 eq) and hydroxylamine hydrochloride (364 mg,
 10 5.24 mmol, 1.2 eq) were dissolved in THF/MeOH/water (4:2:1, 13 mL). The solution was stirred at rt for 5 min then a crystal of methyl orange and sodium cyanoborohydride (550 mg, 8.74 mmol, 2 eq) were added. The pH was adjusted to 2 and the resulting ruby red color was maintained for the duration of the reaction by the regular addition of 6 N HCl. After stirring for 2 h another portion of sodium cyanoborohydride (380 mg, 6.07 mmol, 2
 15 eq) was added. After stirring for 1 h, the mixture was filtered on paper, rinsing with THF. The filtrate was diluted with 1 N NaOH (100 mL) and extracted with DCM (3 x 50mL). Pooled organics were washed with water (3x), brine and then dried over Na₂SO₄. The crude product was purified by flash chromatography (EtOAc in Hexane, gradient 0% to 70%) to afford 703 mg of hydroxylamine **121**. Yield 64%.

Part D




A solution of allylic alcohol **6** (5.0 g, 23 mmol, 1 eq), pyridine (3.75 mL, 46mmol, 2eq) and DMAP (281 mg, 2.3μmol, 0.1 eq) in DCM (120 mL) was cooled in an ice/ water bath and bromoacetyl chloride (2.47 mL, 29.9 mmol, 1.3 eq) was added. The solution was
 25 stirred at 0 °C for 30 min then allowed to reach rt. After 30 min, the reaction was successively washed with 1 N HCl (200 mL), water (100 mL), 5% NaHCO₃ (100mL), brine (100mL) with 10% Na₂SO₃ (100 mL) then brine (100 mL), dried over Na₂SO₄, filtered then concentrated in vacuo to an amber liquid.

This liquid was diluted with acetone (40 mL) and NaI (3.44 g, 23 mmol, 1 eq) was added. The mixture was stirred at rt for 6 h then diluted with EtOAc (100mL), washed with water (1x) then 10% Na₂SO₃ (2x), dried over Na₂SO₄, filtered and concentrated in vacuo to an amber oil.

- 5 This oil was restored in acetonitrile (50 mL) and AgNO₃ (5.0g, 29 mmol, 1.25 eq) was added. The reaction was stirred overnight at rt then partitioned between water (200 mL) and diethyl ether (150mL). The aqueous layer was separated and washed once more with ether (100 mL). The combined organic layers were washed with brine (2x), dried over Na₂SO₄, filtered and concentrated in vacuo. The residue was purified by flash
- 10 chromatography (EtOAc in Hexane, gradient from 0% to 10%) affording **122** (4.27g) as a clear liquid. Yield 58%.


Part E



- Nitrite **122** (800 mg, 2.50 mmol, 1.05eq) was dissolved in DMSO (5 mL) and
- 15 sodium acetate (308 mg, 3.75 mmol, 1.50 eq) was added. The solution was stirred at rt for 30 min then poured into brine (30 mL) and extracted with diethyl ether (3 x 30 mL). The combined organic layers were washed with saturated NaHCO₃ (1x), water (2x) and brine (1x), dried over Na₂SO₄, filtered and concentrated to dryness.

- The product was then reacted overnight with hydroxylamine **121** (588 mg, 2.35
- 20 mmol, 1 eq) in refluxing toluene (20 mL). The solution was concentrated and the residue purified by flash chromatography (EtOAc in hexane, gradient 0% to 30%) to afford lactone **123** (923 mg). Yield 78%.


Part F



Lactone **123** (923 mg, 1.83 mmol, 1eq) is dissolved in THF (8 mL) and HCl 6N (0.6 mL) is added. The solution is stirred at room temperature for 2 h then diluted with DCM (100 mL) and washed with 5% NaHCO₃ (3x) and brine (1x), dried over Na₂SO₄, filtered and concentrated to dryness.

To a solution of the residue in dry DCM (6 mL) was added a solution of (+)-isopinocampheylamine (614 μ L, 3.66 mmol, 2 eq) previously treated in dry DCM (6 mL) with trimethylaluminum in hexanes (2.0 M, 1.83 mL, 3.66 mmol, 2 eq) for 15 min. The solution was stirred at rt overnight then diluted with dry DCM and quenched with $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ (5.9 g 18.3 mmol, 10 eq). The mixture was stirred vigorously at rt for 5 h then filtered on Celite. The filtrate was concentrated to dryness and the residue purified by flash chromatography (EtOAc in Hexane, gradient 0% to 80%) to afford 994 mg of **124**. Yield 71%.

Part G




15 A flask containing bromide **122** (360 mg, 662 μ mol, 1 eq), 3-amino-5-carboxyphenylboronic acid (240 mg, 1.32 mmol, 2 eq), cesium carbonate (383 mg, 1.99 mmol, 3 eq), potassium acetate (65 mg, 662 μ mol, 1 eq) and Pd(dppf)Cl₂ (48 mg, 66 μ mol, 0.1 eq) was purged with argon and DMSO (5 mL) was added. The reaction mixture was added to water (300 mL), acidified with 6M HCl until the aqueous layer attained a pH of 4,
20 and extracted with DCM (3 x 100 mL). The combined organic layers were washed with water (1 x 100 mL) dried over Na₂SO₄ and concentrated to a brown oil.

This crude oil was dissolved in MeOH (10 mL) and treated with HOAc (38 μ L, 662 μ mol, 1 eq), 37% formalin (493 μ L, 6.62 mol, 10 eq) and sodium cyanoborohydride (333 mg, 5.29 mmol, 8 eq). After stirring at rt for 40 min, the reaction mixture was partitioned between water (50 mL) and DCM (20 mL) and acidified with 6M HCl until the aqueous layer attained a pH of 4. The layers were separated and the aqueous extracted with DCM (3 x 20 mL). The combined organic layers were washed with water (1x), dried over Na₂SO₄, and concentrated to a brown oil. This oil was purified by silica gel flash chromatography


(DCM/MeOH/AcOH 90:10:1 in DCM 0% to 100%) to give 119 mg **123** as a brown solid. Yield 29%.

Part H



- 5 A solution of **123** (33 mg, 53 μmol , 1 eq) in DCM (1 mL) was treated with Et_3N (22 μL , 158 μmol , 3 eq), diamine **124** (11 mg, 79 μmol , 1.5 eq), and HBTU (30 mg, 79 μmol , 1.5 eq). After 3h, the mixture was concentrated to dryness. The residue was purified by HPLC to afford **1** (6 mg) after lyophilization. Yield 15%. MS (ESI(+)) m/z 755.22 ($\text{M}+\text{H}$)⁺.


Example 62



10


Prepared in an analogous fashion as described in Example 1 using 2-phenylethylamine in place of (S)- $N^1,N^1,4,4$ -tetramethylpentane-1,2-diamine. Yield 50%.

Example 63



15


Part A



20


To a solution of hydroxylamine **121** (12.3 g, 49.2 mmol, 1 eq) in anhydrous diethyl ether (270 mL) was added the methyl ester (7.68 g, 63.9 mmol, 1.3 eq) and anhydrous calcium chloride (7.10 g, 63.9 mmol, 1.3 eq). The reaction mixture was stirred at room temperature for 3 hours and then filtered through a Celite plug. The filtrate was washed with DCM and diethyl ether. Pool organics was then concentrated to dryness to obtain crude **128** (15.75 g) which was taken on without further purification.

Part B.



Nitron **128** (15.75 g, 49.2 mmol, 1 eq), allylic alcohol (12.78 g, 59.0 mmol, 1.2 eq) and $\text{Ti}(\text{iOPr})_4$ (21.62 mL, 73.8 mmol, 1.5 eq) were dissolved in anhydrous THF (100 mL) and heated in the microwave at 140 °C for 10 min. The reaction mixture was diluted with EtOAc (30 mL) and 3-(dimethylamino)-1,2-propanediol (30 mL) was added. After stirring overnight, the mixture was washed with Rochelle's salt, water and brine. The mixture was then filtered over Celite and washed with EtOAc, water and then brine. The organic was dried over Na_2SO_4 , and concentrated to dryness to obtain crude **129** (24.82 g) which was taken on without further purification.

Part C.




To a solution of lactone **129** (4.82 g, 49.2 mmol, 1 eq) in THF (270 mL) was added concentrated HCl (24.6 mL, 148 mmol, 3 eq) and stirred at room temperature for 1.5 hours. The reaction was then basified to pH 7 with saturated NaHCO_3 and extracted with EtOAc. Pool organics were washed with brine, dried over Na_2SO_4 , filtered and concentrated to

dryness. The crude product was purified by flash chromatography (EtOAc in Hexane, gradient 20% to 40%) to afford **130** (15.6 g). Yield 81%.


Part D.

5



To a solution of lactone **130** (6.6 g, 16.91 mmol, 1eq) in DCM (71 mL) was added Hunig's base (8.86 mL, 50.7 mmol, 3 eq) and trifluoromethanesulfonic anhydride (3.43 mL, 20.3 mmol, 1.2 eq) at 0 °C and stirred for 15 minutes. The reaction mixture was then warmed to room temperature and DMF (84 mL) and NaN₃ (3.3 g, 50.7 mmol, 3 eq) was added and stirred overnight at room temperature. The reaction mixture was washed with NaHCO₃ and brine, dried over Na₂SO₄, filtered and concentrated to dryness. The crude product was purified by flash chromatography (EtOAc in Hexane, gradient 30% to 50%) to obtain **131** (6.85 g). Yield 98%.

Part E.




20

To a solution of (+)-isopinocampheylamine (1.22 mL, 7.23 mmol, 6 eq) in THF (10 mL) was added DIBAL (6.03 mL, 1 M in DCM, 6.02 mmol, 5 eq) and stirred for 30 minutes at 0 °C. The mixture was then warmed to room temperature and allowed to stir for another 2 hours. A solution of lactone **131** (0.5 g, 1.2 mmol, 1 eq) in THF (5 mL) was added and the mixture was stirred for 30 minutes at room temperature. The reaction was diluted with EtOAc (35 mL) and quenched by the addition of saturated aqueous Rochelle


salt (35 mL) and the mixture was rapidly stirred overnight. The mixture was extracted with diethyl ether and the pooled organics were washed with water and then brine, dried over Na_2SO_4 , filtered and concentrated to dryness. The crude material was purified by flash chromatography (EtOAc in Hexane, gradient 10% to 30%) to afford **132** (0.42 g). Yield 61%.

Part F.




A flask containing aryl bromide **132** (470 mg, 0.83 mmol, 1 eq), 3-amino-5-carboxyphenylboronic acid (36.1 mg, 1.24 mmol, 1.5 eq), cesium carbonate (80.8 mg, 2.48 mmol, 3 eq), potassium acetate (122 mg, 1.24 mmol, 1.5 eq) and Pd(dppf)Cl_2 (121 mg, 0.17 mmol, 0.2 eq) was purged with argon and DMSO (10 mL) added. The mixture was heated at 70 °C for 3 hours. The reaction mixture was added to water (10 mL), acidified with 6 N HCl until the aqueous layer attained a pH of 4, and extracted with EtOAc (3 x 10 mL). The combined organic layers were washed with water, separated, dried over Na_2SO_4 , filtered and concentrated to dryness. The crude material was purified by flash chromatography (EtOAc in Hexane, gradient 30% to 50%) to afford **133** (46 mg). Yield 8.5%.

Part J.




A solution of 133 (64.7 mg, 0.099 mmol, 1 eq) in DCM (5 mL) was treated with Et₃N (41.4 μ L, 0.23 mmol, 3 eq), (S)-N1,N1-dimethyl-3-phenylpropane-1,2-diamine (26.5 mg, 0.149 mmol, 1.5 eq) and HBTU (75 mg, 0.198 mmol, 2 eq). After 2 hours, the mixture was diluted with DCM (15 mL), washed with saturated K₂CO₃ solution (15 mL) and extracted with DCM (2 x 15 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated to dryness. The crude material was purified by flash chromatography (MeOH in DCM, gradient 1% to 5%) to afford 134 (43 mg). Yield 53%.

Part K.




A solution of 134 (43 mg, 0.053 mmol, 1 eq) in DMF (4 mL) was treated with dithiothreitol (24.5 mg, 0.159 mmol, 3 eq) and DBU (23.9 μ L, 0.159 mmol, 3 eq) and stirred at 0 °C. After 15 minutes, the mixture was purified by HPLC to afford 135 (4.1 mg) after lyophilization. Yield 12.7%. MS (ESI(+)) m/e 788.13 (M+H)⁺.

Example 64



136


Part A.



Aminodiol **137** (5.0g, 29.9mmol, 1eq) was dissolved in MeOH (30mL) at 0°C and diterbutyl dicarbonate (7.18g, 32.9mmol, 1.1eq) was added. The solution was stirred at 0°C
 5 for 2h then concentrated to dryness.

The residue was taken in DMF (120mL) and imidazole (4.48g, 65.8mmol, 2.2eq) was added. The solution was cooled in an ice/water bath and ter-butyldiphenylsilyl chloride (9.86g, 32.9mmol, 1.2eq) was added as a solution in DMF (20mL). The solution was stirred at rt for 18h then at 60°C for 2h. The reaction was poured on saturated NaHCO₃ (300mL)
 10 and the solution extracted with EtOAc (3x100mL). The combined organics were dried over Na₂SO₄, filtered, concentrated. The residue was purified by flash chromatography (Hex/EtOAc 0% to 20%) to afford **138** as a colorless oil (6.15g, 12.1mmol). Yield 41%.


Part B.




15

Compound **138** (6.1g, 12mmol, 1eq) was refluxed with 2,2-dimethoxypropane (2.96mL, 24.1mmol, 2eq) in toluene (60mL) using paratoluenesulfonic acid monohydrate (23mg, 0.12mmol, 0.1eq) as a catalyst. After 30min, the reaction was allowed to cool to rt then washed with 5% NaHCO₃ (3x20mL) and brine (1x20L), dried over Na₂SO₄, filtered
 20 and concentrated to dryness.

The crude was dissolved in THF (60mL) and a 1.0M solution of ter-butylammonium fluoride was added (60mL, 60mmol, 5eq). The solution was stirred at 60°C for 2h then its volume was reduced on a rotary evaporator. The residue was partitioned between water (100mL) and CHCl₃ (60mL). The layers were separated and the aqueous extracted with CHCl₃ (2x60mL). The combined organics were washed with brine (1x60mL), dried over Na₂SO₄, filtered and concentrated. The residue was purified by flash chromatography
 25 (Hex/EtOAc 0% to 40%) to afford **139** (3.13g, 10.2mmol) as a colorless oil. Yield 84%.


Part C.

Alcohol 139 (3.13g, 10.2mmol, 1eq) and tosyl chloride (2.13g, 11.2mmol, 1.1eq) are dissolved in DCM (20mL) then triethylamine (2.84mL, 20.3mmol, 2eq) and catalytic DMAP (124mg, 1mmol, 0.1eq) were added. The orange solution was stirred at rt for 3h then diluted with DCM (60mL), washed with HCl 1 N (3x30mL) and brine (1x30mL), dried over Na₂SO₄, filtered, concentrated. The residue was purified by flash chromatography (Hex/EtOAc 0% to 30%) to afford 140 (4.08g, 8.8mmol) as a white solid. Yield 87%.

Part D.

15


In a sealed reactor, tosylate 140 (4.08g, 8.8mmol, 1eq) was heated at 60°C in a 1.0M solution of dimethylamine in THF (88mL, 177mmol, 20eq). After 44h, the solution was concentrated and the residue purified by flash chromatography (DCM and DCM/MeOH/NH₄OH 80:20:0.5 0% to 50%) to afford 141 (2.46g, 7.36mmol) as a clear oil. Yield 83%.

Part E.

25

Compound **141** (487mg, 1.45mmol, 1eq) was treated with a 4.0M solution of HCl in dioxane (5mL) and water (100μL) at rt for 1h30. The solution was then concentrated and dried under high vacuum to obtain the hydrochloric salt of **142** (339mg, 1.45mmol) as an off-white foam in a quantitative yield.

Part F.




10

Acid **10** (30mg, 49μmol, 1eq), amine **142** (14mg, 59μmol, 1.2eq) and HBTU (22mg, 59μmol, 1.2eq) were dispersed in DCM (1mL) and triethylamine (41μL, 295μmol, 6eq) was added. The reaction was stirred at rt for 30min then concentrated to dryness. The residue was purified by HPLC to afford **136** (13mg, 16μmol) as a lyophilized powder. Yield 33%. MS (ESI(+)) m/e 786.27 (M+H)⁺.


15

Example 65




20

Part A.




To a stirred solution of acid **125** (150 mg, 0.24 mmol, 1.0 equiv) in CH_2Cl_2 (3 mL, 0.08M) was added triethylamine (100 μL , 0.717 mmol, 3.0 equiv) followed by benzyl alcohol **142** (45 mg, 0.24 mmol, 1.0 equiv) and HBTU (100 mg, 0.26 mmol, 1.1 equiv.). After 14 hours, the reaction mixture was diluted with CH_2Cl_2 , washed with water, dried with sodium sulfate, and concentrated *in vacuo*. The resulting oil was purified by HPLC (acidic method) to afford **143** as a brown oil (9.2 mg, 5% yield). LCMS (ESI(+)) m/e 804 ($\text{M}+\text{H}$)⁺.

Example 66



Part A.




Aminodiol **137** (4.0g, 23.9mmol, 1eq) was dissolved in MeOH (50mL) at 0°C and di-tert-butyl dicarbonate (6.27g, 28.7mmol, 1.2eq) was added. The solution was stirred at 0°C for 2h then concentrated to dryness.

The residue was taken in DMF/DMSO 2:1 (60mL) and imidazole (3.26g, 47.8mmol, 2eq) then ter-butyltrimethylsilyl chloride (4.33g, 28.7mmol, 1.2eq) were added. The solution was stirred at rt overnight. After 16h, some more imidazole (3.26g, 47.8mmol,


2eq) then ter-butyldimethylsilyl chloride (4.33g, 28.7mmol, 1.2eq) were added. After 3h, the reaction was poured on saturated NaHCO_3 (200mL) and the solution extracted with EtOAc (3x100mL). The combined organics were dried over Na_2SO_4 , filtered, concentrated. The residue was purified by flash chromatography (Hex/EtOAc 0% to 20%) to afford **145** as a clear oil (7.91g, 20.7mmol). Yield 87%.

Part B.



Alcohol **145** (2.08g, 5.45mmol, 1eq), phthalimide (0.96g, 6.54mmol, 1.2eq) and triphenylphosphine (1.71g, 6.54mmol, 1.2eq) were dissolved in dry THF (20mL) under Ar and diethylazodicarboxylate (1.04mL, 6.54mmol, 1.2eq) was added dropwise. The reaction was stirred for 2h then a solution of triphenylphosphine (340mg, 1.3mmol, 0.2eq) in THF (3mL) then diethylazodicarboxylate (0.20mL, 1.3mmol, 0.2eq) were added. The solution was stirred at rt for 1 h then concentrated to dryness. The residue was purified by flash chromatography (Hex/EtOAc 0% to 25%) to afford **146** (1.38g, 5.45mmol). Yield 49%.


Part C.



Compound **146** (1.379g, 2.7mmol, 1eq) was treated with hydrazine hydrate (1.31mL, 27mmol, 10eq) in refluxing ethanol for 2h. The reaction was allowed to reach rt then filtered on paper. The filtrate was concentrated to dryness and the residue taken in DCM. The mixture was filtered on paper and the filtrate concentrated to dryness.

The residue was dissolved in DCM (20mL) and triethylamine (1.13mL, 8.1mmol, 3eq) then mesyl chloride (0.42mL, 5.40mmol, 2eq) were added. The solution was stirred at rt for 1 h then some more triethylamine (1.13mL, 8.1mmol, 3eq) and mesyl chloride (0.42mL, 5.40mmol, 2eq) were added. The solution was stirred at rt for 1 h then diluted with DCM (100mL) and washed with HCl 1 N (3x30mL). The combined organics were dried over Na₂SO₄, filtered and concentrated. The residue was purified by flash chromatography (Hex/EtOAc 0% to 100%) to afford sulfonamide **147** (0.898g, 1.96mmol) as a white solid. Yield 72.5%.


10 Part D.



Compound **147** (0.898g, 1.96mmol, eq) was dissolved in dry THF (20mL) and a 1.0M solution of tertbutylammonium fluoride was added (3.9mL, 3.9mmol, 2eq). The solution was stirred at rt for 2h then its volume was reduced on a rotary evaporator. The residue was partitioned between water (60mL) and CHCl₃ (30mL). The layers were separated and the aqueous extracted with CHCl₃ (2x30mL). The combined organics were washed with brine (1x30mL), dried over Na₂SO₄, filtered and concentrated.

The residue was taken in DCM (10mL) then tosyl chloride (448mg, 2.35mmol, 1.2eq), triethylamine (545μL, 3.9mmol, 2eq) and catalytic DMAP (24mg, 196μmol, 0.1eq) were added. The solution was stirred at rt for 4h then diluted with DCM (60mL), washed with HCl 1 N (3x20mL) and brine (1x20mL), dried over Na₂SO₄, filtered, concentrated. The residue was purified by flash chromatography (Hex/EtOAc 0% to 30%) to afford **148** (713mg, 1.43mmol) as a white solid. Yield 73%.

Part E.




In a sealed reactor, tosylate **148** (713mg, 1.43mmol, 1eq) was heated at 60°C in a 1.0M solution of dimethylamine in THF (28mL, 57.2mmol, 40eq). After 16h, the solution was concentrated and the residue purified by flash chromatography (DCM and DCM/MeOH/NH₄OH 80:20:0.5 0% to 80%) to afford **149** (416mg, 1.12mmol) as a clear oil. Yield 78%.

Part F.



Compound **149** (385mg, 1.04mmol, 1eq) was treated with a 4.0M solution of HCl in dioxane (10mL) at rt for 1. The turbid solution was then concentrated and dried under high vacuum to obtain the hydrochloric salt of **150** (319mg, 1.04mmol) as white solid in a quantitative yield.


Part G.




Acid **10** (30mg, 49μmol, 1eq), amine **150** (17mg, 54μmol, 1.1eq) and HBTU (21mg, 54μmol, 1.1eq) were dispersed in DCM (1mL) and triethylamine (41μL, 295μmol,

6eq) was added. The reaction was stirred at rt for 30min then concentrated to dryness. The residue was purified by HPLC to afford **144** (11.8mg, 13.6 μ mol) as a lyophilized powder. Yield 27.8%. MS (ESI(+)) m/e 863.25 (M+H)⁺.

5

Example 67**151****Part A.**



10



To a stirred solution of acid **125** (59 mg, 0.094 mmol, 1.0 equiv.) in CH₂Cl₂ (3 mL, 0.03M) was added triethylamine (40 μ L, 0.282 mmol, 3.0 equiv.) followed by benzyl sulfonamide **150** (28 mg, 0.103 mmol, 1.1 equiv.) and HBTU (43 mg, 0.113 mmol, 1.2 equiv.). After five minutes, the reaction mixture was diluted with CH₂Cl₂, washed with water, dried with sodium sulfate, and concentrated *in vacuo*. The resulting oil was purified by HPLC (basic method) to afford **151** as a brown oil (15.7 mg, 19% yield). LCMS (ESI(+)) m/e 881 (M+H)⁺.

20


Example 68

Part A.

5

Compound **153** (2.95g, 7.72mmol, 1eq) was dissolved in DCM (30mL) and triethylamine (1.18mL, 8.49mmol, 1.1eq) and DMAP (47mg, 0.38mmol, 0.05eq) were added. The solution was cooled in an ice/water bath and mesyl chloride (0.60mL, 7.72mmol, 1eq) was added. The solution was stirred at rt for 3h then triethylamine (1.18mL, 8.49mmol, 1.1eq) and mesyl chloride (0.60mL, 7.72mmol, 1eq) were added. The solution was stirred for 30min, then partitioned between water (30mL) and DCM (90mL). The layers were separated then the organics were washed with HCl 1 N (3x20mL), dried over Na₂SO₄, filtered, concentrated. The residue was purified by flash chromatography (Hex/EtOAc 0% to 25%) to afford **154** (3.16g, 6.87mmol) as an oil. Yield 89%.

15


Part B.

20

Mesylate **154** (2.619g, 5.70mmol, 1eq) was dissolved in dry THF (40mL) under Ar and potassium tertbutoxide (0.655g, 5.84mmol, 1.025eq) was added dropwise as a solution in dry THF (20mL). The reaction was stirred at rt for 3h then quenched with water (60mL) and extracted with EtOAc (3x40mL). The pooled organics were washed with brine

(1x30mL), dried over Na_2SO_4 , filtered and concentrated. The residue was purified by flash chromatography to obtain aziridine **155** (1.96g, 5.41mmol) as a white wax. Yield 95%.

Part C.



5

In a sealed reactor behind a glass shield, a solution of aziridine **155** (1.87g, 5.14mmol, 1eq) and trimethylsilylazide (2mL, 20.5mmol, 4eq) in methanol (40mL) was heated at 70°C for 5h. The solution was allowed to cool to rt then concentrated. The residue was purified by flash chromatography (Hex/EtOAc 0% to 15%) to afford adduct **156** (1.72g, 4.36mmol). Yield 85%

10

Part D.




15

Compound **156** (1.646g, 4.16mmol, eq) was dissolved in dry THF (20mL) and a 1.0M solution of tertbutylammonium fluoride was added (8.3mL, 8.32mmol, 2eq). The solution was stirred at rt for 2h then its volume was reduced on a rotary evaporator. The residue was partitioned between water (60mL) and CHCl_3 (30mL). The layers were separated and the aqueous extracted with CHCl_3 (2x30mL). The combined organics were washed with brine (1x30mL), dried over Na_2SO_4 , filtered and concentrated.

20

The residue was taken in DCM (10mL) then tosyl chloride (952mg, 4.99mmol, 1.2eq), triethylamine (1.16mL, 8.32mmol, 2eq) and catalytic DMAP (51mg, 416μmol, 0.1eq) were added. The solution was stirred at rt for 4h then diluted with DCM (60mL), washed with HCl 1 N (3x20mL) and brine (1x20mL), dried over Na_2SO_4 , filtered, concentrated. The residue was purified by flash chromatography (Hex/EtOAc 0% to 30%) to afford **157** (886mg, 2.0mmol) as a white solid. Yield 49%.

25


Part E.

In a sealed reactor, tosylate **157** (800mg, 1.84mmol, 1eq) was heated at 60°C in a 1.0M solution of dimethylamine in THF (37mL, 37mmol, 20eq). After 16h, the solution was concentrated and the residue purified by flash chromatography (DCM and DCM/MeOH/NH₄OH 80:20:0.5 0% to 80%) to afford **158** (352mg, 1.14mmol) as a clear oil. Yield 62%.

10

Part F.

Compound **158** (300mg, 0.97mmol, 1eq) was treated with a 4.0M solution of HCl in dioxane (10mL) at rt for 1. The turbid solution was then concentrated and dried under high vacuum to obtain the hydrochloric salt of **159** (238mg, 0.97mmol) as white solid in a quantitative yield.


Part G.

20

Acid **10** (40mg, 66μmol, 1eq), amine **159** (18mg, 72μmol, 1.1eq) and HBTU (27mg, 72μmol, 1.1eq) were dispersed in DCM (1mL) and triethylamine (27μL, 197μmol, 3eq) was added. The reaction was stirred at rt for 30min then concentrated to dryness. The


residue was purified by HPLC to afford **152** (18mg, 23 μ mol) as a lyophilized powder. Yield 34%. MS (ESI(+)) m/e 800.34 (M+H)⁺.

Example 69




160

Part A.



Aminodiol **137** (1.0g, 5.98mmol, 1eq) was dissolved in DCM (50mL) and di-tert-butyl dicarbonate (1.28g, 5.86mmol, 0.98eq) then triethylamine (1.17mL, 12mmol, 2eq) were added. The reaction was stirred at rt. After 1h, the solution was diluted with DCM (50mL), washed with HCl 1 N (3x20mL) and brine (1x20mL), dried over Na₂SO₄, filtered and concentrated to dryness to afford **161** (1.35g, 5.05mmol) which was used without further purification. Yield 84%.


Part B.



Compound **162** (1.13g, 4.2mmol, 1eq) was dissolved in DCM (20mL) then tosyl chloride (0.97g, 5.1mmol, 1.2eq), triethylamine (1.2mL, 8.5mmol, 2eq) and catalytic DMAP (100mg, 0.85mmol, 0.2eq) were added. The solution was stirred at rt for 4h then diluted with DCM (60mL), washed with HCl 1 N (3x20mL) and brine (1x20mL), dried over Na₂SO₄, filtered, concentrated. The residue was purified by flash chromatography

(Hex/EtOAc 0% to 60%) to afford **162** (0.84g, 2.0mmol) as a white solid. Yield 47%.

Part C.




5

Tosylate **162** (0.88g, 2.09mmol, 1eq) was dissolved in DMF (30mL) and pyrrolidine (350μL, 4.17mmol, 2eq) was added. The solution was stirred at 60°C overnight. After 16h, the reaction was allowed to cool to rt, diluted with water (90mL) and extracted with EtOAc (3x30mL). The combined organics were washed with brine, dried over Na₂SO₄, filtered, concentrated. The residue was purified by flash chromatography (DCM and DCM/MeOH/NH₄OH 80:20:0.5 0% to 80%) to afford **163** (224mg, 699μmol) as an oil. Yield 33%.

10

Part D.




15

Compound **163** (224mg, 699μmol, 1eq) was treated with a 4.0M solution of HCl in dioxane (10mL) at rt for 1. The turbid solution was then concentrated and dried under high vacuum to obtain the hydrochloric salt of **164** (154mg, 699μmol) as white solid in a quantitative yield.

20


Part E.



Acid **10** (100mg, 164 μ mol, 1eq), amine **164** (44mg, 197 μ mol, 1.2eq) and HBTU (75mg, 197 μ mol, 1.2eq) were dispersed in DCM (1mL) and triethylamine (69 μ L, 492 μ mol, 3eq) was added. The reaction was stirred at rt for 30min then concentrated to dryness. The residue was purified by HPLC to afford **160** (9.1mg, 11 μ mol) as a lyophilized powder.


5 Yield 6.8%. MS (ESI(+)) m/e 812.56 (M+H)⁺.

Example 70



165


10 Part A.



Boc-(S)-phenylalaninol (2.01g, 8.24mmol, 1eq), phthalimide (1.45g, 9.88mmol, 1.2eq) and triphenylphosphine (3.24g, 12.4mmol, 1.5eq) were dissolved in dry THF (20mL) under Ar.

15 Diisopropylazodicarboxylate (2.43mL, 12.4mmol, 1.5eq) was added dropwise and the reaction stirred at rt. After 2h, the solution was concentrate to dryness and the residue purified by flash chromatography (Hex/EtOAc 15% to 25%) to afford **166** (1.63g, 4.28mmol). Yield 52%.


20 Part B.



Compound **166** (3.46g, 9.09mmol, 1eq) was treated with hydrazine hydrate (2.25mL, 45.5mmol, 5eq) in refluxing ethanol for 2h. The reaction was allowed to reach rt then filtered on paper. The filtrate was concentrated to dryness and the residue taken in DCM. The mixture was filtered on paper and the filtrate concentrated to dryness.


5 The residue was taken in DCM (20mL) and 2,4,6-collidine (0.658mL, 4.99mmol, 1eq) then 2,4-dinitrophenylsulfonyl chloride (1.40g, 5.24mmol, 1.05eq) were added. After 5h, the reaction was diluted with DCM (100mL), washed with HCl 1 N (3x20mL), water (1x20mL) and brine (1x20mL), dried over Na₂SO₄, filtered and concentrated. The residue was purified by flash chromatography (Hex/EtOAc 10% to 25%) to afford **167** (1.63g, 3.99mmol) as a yellow powder. Yield 68%.

Part C.



15 Compound **167** (777mg, 1.62mmol, 1eq) and potassium carbonate (447mg, 3.24mmol, 2eq) were dissolved in acetone (20mL) and iodomethane (204μL, 3.24mmol, 2eq) was added. The reaction was stirred at rt for 20h then diluted with HCl 1 N (100mL) and extracted with EtOAc (3x25mL). The combined organics were washed with water (1x20mL) and brine (1x20mL), dried over Na₂SO₄, filtered and concentrated. The residue was purified by flash chromatography (Hex/EtOAc 10% to 25%) to afford **168** (685mg, 1.39mmol) as a yellow powder. Yield 85%.

Part D.




25

Compound **168** (685mg, 1.4mmol, 1eq) was treated with a 4.0M solution of HCl in dioxane (10mL) at rt for 1. The turbid solution was then concentrated and dried under high

vacuum to obtain the hydrochloric salt of **169** (604mg, 1.4mmol) as yellow foam in a quantitative yield.


Part E.




Acid **10** (40mg, 66 μ mol, 1eq), amine **169** (31mg, 79 μ mol, 1.2eq) and HBTU (30mg, 79 μ mol, 1.2eq) were dispersed in DCM (1mL) and 2,4,6-collidine (26 μ L, 197 μ mol, 3eq) was added. The reaction was stirred at rt for 1 h then mercaptoacetic acid (108mL, 1.32mmol, 20eq) was added. The redish solution was stirred at rt for 1 h then concentrated to dryness. The residue was purified by HPLC to afford **165** (7.3mg, 9.6 μ mol) as a lyophilized powder. MS (ESI(+)) m/e 756.33 (M+H)⁺. Yield 14.6%.

10

Example 71




Part A.



N-Boc-S-terbutyl-(D)-cysteine (2.40g, 8.65mmol, 1eq) was dissolved in dry THF (20mL) under an atmosphere of Argon and the solution cooled down in an ice/ brine bath.


Ethylchloroformate (0.827mL, 8.65mmol, 1eq) then triethylamine (0.876mL, 8.25mmol, 1eq) were added dropwise and the reaction stirred for 10min at -10°C. Sodium borohydride (1.31g, 34.6mmol, 4eq) was added and the reaction stirred at -10°C for 2h. The reaction was then quenched with methanol (40mL) and HCl 1 N (15mL) at 0°C. The methanol was
 5 evaporated off and the solution diluted with HCl 1 N (80mL), extracted with EtOAc (3x30mL). The combined organics were washed with HCl 1 N (3x20mL), NaOH 1 N (3x20mL) and brine (1x20mL), dried over Na₂SO₄, filtered and concentrated to afford **171** (2.08g, 7.90mmol) which was used without further purification. Yield 91%.

10 Part B.




Alcohol **171** (2.0g, 7.6mmol, 1eq), phthalimide (1.3g, 9.1mmol, 1.2eq) and triphenylphosphine (2.4g, 9.1mmol, 1.2eq) were dissolved in dry THF (40mL) under Ar.
 15 Diisopropylazodicarboxylate (2.4mL, 12mmol, 1.6eq) was added dropwise and the reaction stirred at rt. After 2h, the solution was concentrate to dryness and the residue purified by flash chromatography (Hex/EtOAc 0% to 50%) to afford **172** (2.11g, 7.6mmol). Yield 71%.

Part C.




Thioether **172** (2.1g, 5.4mmol, 1eq) was dissolved in methanol (100mL) and Oxone (9.9g, 16mmol, 3eq) was added as a solution in water (20mL). The reaction was stirred at rt for 1 h then quenched with 20% sodium bisulfite (20mL). The solution was diluted with
 25 water (200mL) and extracted with chloroform (3x40mL). The combined organics were washed with brine (1x30mL), dried over Na₂SO₄, filtered and concentrated to afford **173** (2.2g, 5.2mmol) which was used without further purification. Yield 97%.

Part D.


Sulfone **173** (2.20g, 5.2mmol, 1eq) was treated with hydrazine hydrate (2.5mL, 52mmol, 10eq) in refluxing ethanol for 2h. The reaction was allowed to reach rt then filtered on paper. The filtrate was concentrated to dryness and the residue taken in DCM. The mixture was filtered on paper and the filtrate concentrated to dryness.

The residue was taken in methanol (40mL) and formaldehyde 37% (1.1mL, 15mmol, 3eq) was added. The solution was stirred at rt for 10min then sodium cyanoborohydride (0.96g, 15mmol, 3eq) was added. The reaction was stirred at rt for 1 h then quenched with 5% NaHCO_3 (40mL). The volume of the reaction was reduced on a rotary evaporator and the residual aqueous solution extracted with DCM (3x20mL). The combined organics were washed with brine (1x20mL) then dried over Na_2SO_4 , filtered and concentrated to afford **174** (1.416g, 4.4mmol) which was used without further purification. Yield 86%.

Part E.

Compound **174** (1.416g, 4.4mmol, 1eq) was treated with a 4.0M solution of HCl in dioxane (10mL) at rt for 1. The turbid solution was then concentrated and dried under high vacuum to obtain the hydrochloric salt of **175** (1.13g, 4.4mmol) as yellow foam in a quantitative yield.


Part F.




Acid **10** (40mg, 66 μ mol, 1eq), amine **175** (31mg, 79 μ mol, 1.2eq) and HBTU (30mg, 79 μ mol, 1.2eq) were dispersed in DCM (1mL) and triethylamine (27 μ L, 197 μ mol, 3eq) was added. The reaction was stirred at rt for 1 h then concentrated to dryness. The residue was purified by HPLC to afford **170** (9.6mg, 12 μ mol) as a lyophilized powder. Yield 18%.

MS (ESI(+)) m/e 814.47 (M+H)⁺.

Example 72




Part A.



N-Boc-S-methyl-(L)-cysteine (1.50g, 6.37mmol, 1eq) was dissolved in dry THF (20mL) under an atmosphere of Argon and the solution cooled down in an ice/ brine bath. Ethylchloroformate (0.610mL, 6.37mmol, 1eq) then N-methylmorpholine (0.701mL, 6.37mmol, 1eq) were added dropwise and the reaction stirred for 10min at -10°C. Sodium borohydride (0.965g, 25.5mmol, 4eq) was added and the reaction stirred at -10°C for 2h. The reaction was then quenched with methanol (40mL) and HCl 1 N (15mL) at 0°C. The methanol was evaporated off and the solution diluted with HCl 1 N (80mL), extracted with EtOAc (3x30mL). The combined organics were washed with HCl 1 N (3x20mL), NaOH 1

N (3x20mL) and brine (1x20mL), dried over Na₂SO₄, filtered and concentrated to afford 177 (1.11g, 5.04mmol) which was used without further purification. Yield 79%.

Part B.




5

Alcohol 177 (1.11g, 5.04mmol, 1eq), phthalimide (0.889g, 6.05mmol, 1.2eq) and triphenylphosphine (1.98g, 7.56mmol, 1.5eq) were dissolved in dry THF (30mL) under Ar. Diisopropylazodicarboxylate (1.46mL, 7.56mmol, 1.5eq) was added dropwise and the reaction stirred at rt. After 2h, the solution was concentrate to dryness and the residue purified by flash chromatography (Hex/EtOAc 0% to 50%) to afford 178 (1.43g, 4.12mmol). Yield 82%.

10

Part C.


15



Thioether 178 (1.43g, 4.1mmol, 1eq) was dissolved in methanol (100mL) and Oxone (5.1g, 8.2mmol, 2eq) was added as a solution in water (20mL). The reaction was stirred at rt for 1 h then quenched with 20% sodium bisulfite (20mL). The solution was diluted with water (200mL) and extracted with chloroform (3x40mL). The combined organics were washed with brine (1x30mL), dried over Na₂SO₄, filtered and concentrated to afford 179 (1.3g, 3.4mmol) which was used without further purification. Yield 83%.

20

25 Part D.




Sulfone **179** (1.3g, 3.4mmol, 1eq) was treated with hydrazine hydrate (1.06mL, 34mmol, 10eq) in refluxing ethanol for 2h. The reaction was allowed to reach rt then filtered on paper. The filtrate was concentrated to dryness and the residue taken in DCM.

5 The mixture was filtered on paper and the filtrate concentrated to dryness.


The residue was taken in methanol (40mL) and formaldehyde 37% (0.75mL, 10mmol, 3eq) was added. The solution was stirred at rt for 10min then sodium cyanoborohydride (0.64g, 10mmol, 3eq) was added. The reaction was stirred at rt for 1 h then quenched with 5% NaHCO₃ (40mL). The volume of the reaction was reduced on a rotary evaporator and the residual aqueous solution extracted with DCM (3x20mL). The combined organics were washed with brine (1x20mL) then dried over Na₂SO₄, filtered and concentrated to afford **180** (0.85g, 3mmol) which was used without further purification. Yield 90%.

15 Part E.




Compound **180** (0.85g, 3mmol, 1eq) was treated with a 4.0M solution of HCl in dioxane (10mL) at rt for 1. The turbid solution was then concentrated and dried under high vacuum to obtain the hydrochloric salt of **181** (0.21g, 1.17mmol) as a white foam. Yield 39%.

Part F.




Acid described in Example 4 part G (50mg, 80 μ mol, 1eq), amine 181 (17mg, 96 μ mol, 1.2eq) and HBTU (36mg, 96 μ mol, 1.2eq) were dispersed in DCM (1mL) and triethylamine (34 μ L, 240 μ mol, 3eq) was added. The reaction was stirred at rt for 1 h then concentrated to dryness. The residue was purified by HPLC to afford 176 (2.2mg, 2.8 μ mol) as a lyophilized powder. MS (ESI(+)) m/e 786.56 (M+H)⁺. Yield 3.5%.

Example 73




182

Part A.




In a three-neck round-bottom flask topped with a condenser, (L)-Phenylalanine-pyrrolidide (1.00g, 4.6mmol, 1eq) was dissolved in dry THF (20mL) under an atmosphere of Argon. The solution was cooled in an ice/water bath and lithium aluminium hydride was added dropwise as a 2.0M solution in THF (9.2mL, 18mmol, 4eq). The reaction was refluxed overnight under an atmosphere of Argon. After 18h, the reaction was allowed to cool to rt, then cooled in an ice/water bath and quenched by the cautious, sequential, dropwise addition of water (0.7mL), 15% NaOH (0.7mL) and water (2.1mL) to obtain a white precipitate. The reaction was filtered on paper and the cake rinsed with THF (60mL). The filtrate was concentrated to obtain crude 183 as a yellow oil (0.917g, 4.5mmol) which was used without further purification. Yield 98%.

Part B.




Acid prepared as described in Example 4 part G (80mg, 128 μ mol, 1eq), amine **183** (31mg, 154 μ mol, 1.2eq) and HBTU (58mg, 154 μ mol, 1.2eq) were dispersed in DCM (1mL) and triethylamine (54 μ L, 385 μ mol, 3eq) was added. The reaction was stirred at rt for 1 h then concentrated to dryness. The residue was purified by HPLC to afford **182** (16.8mg, 20 μ mol) as a lyophilized powder. Yield 16.1%. MS (ESI(+)) m/e 810.65 (M+H)⁺.


Example 74




Part A.




Sodium hydride 60% (3.1g, 78mmol, 2eq) was dispersed in dry THF (40mL) under an atmosphere of Argon. 3-chlorophenol (4.1mL, 39mmol, 1eq) was added dropwise over 10min as a solution in dry THF (10mL). The solution was stirred for 1 h then diethylchloroformate (9.9mL, 78mmol, 2eq) was added dropwise as a solution in dry THF (10mL). The solution was stirred overnight at rt. After 14h, the reaction was quenched by the addition of water (5mL) dropwise, diluted with diethylether (120mL). The organics were washed with water (2x30mL) then NaOH (2x30mL), dried over Na₂SO₄, filtered and concentrated. The residue was purified by flash chromatography (Hexane/EtOAc 0% then 10%) to obtain **184** as an oil (8.2g, 36mmol). Yield 93%.

Part B.

Dry THF (120mL) was charged in a flame-dried round-bottom flask under an atmosphere of Argon and cooled in a dry ice/ acetone bath. A 1.4M solution of sec-butyl lithium in cyclohexane (31mL, 43mmol, 1.2eq), tetramethylethylenediamine (6.5mL, 43mmol, 1.2eq) and a solution of **184** (8.2g, 36mmol, 1eq) in dry THF (10mL) were added sequentially. The solution was stirred at -78°C for 2h then iodine (11g, 43mmol, 1.2eq) was added as a solution in dry THF (20mL). The solution was stirred at -78°C for 30min then allowed to reach rt. After 2h, the reaction was quenched with 5% $\text{Na}_2\text{S}_2\text{O}_3$ (100mL) then extracted with EtOAc (3x100mL). The combined organics were dried over Na_2SO_4 , filtered and concentrated. The residue was purified by flash chromatography (Hexane/EtOAc 0% to 30%) to obtain **185** an oil (8.92g, 25mmol). Yield 70%.


15 Part C.

Carbamate **185** (10g, 28mmol, 1eq) was treated with sodium hydroxide (11g, 283mmol, 10eq) in refluxing ethanol (140mL). After 14h, the solution was allowed to cool to rt, acidified with HCl 6N then extracted with DCM (3x60mL). The combined organics were dried over Na_2SO_4 , filtered and concentrated. The residue was purified by flash chromatography (Hexane/DCM 0% to 100%) to obtain **186** an oil (3.75g, 15mmol). Yield 52%.

25 Part D.


To a mixture of MgCl_2 (2.25g, 23.6mmol, 2eq) and paraformaldehyde (1.06g, 35.4mmol, 3eq) in THF (60mL) under argon, was added triethylamine (3.29mL, 23.6 mmol, 2eq). The mixture was stirred at room temperature, under argon, for 10min and phenol **186** (3.00g, 11.8mmol, 1eq) was added. The reaction was heated to reflux for 3h then allowed to cool to room temperature. Diethyl ether (100 mL) was added and the solution was washed with HCl 1 N (3x100mL), dried over Na_2SO_4 , filtered and concentrated to dryness to obtain crude **187** (3.33g, 11.8mmol) which was taken on without further purification. Yield 99%.

10 Part E.



To a solution of crude phenol **187** (3.33 g, 11.8mmol, 1eq) in DMF (50 mL) at room temperature was added K_2CO_3 (2.18g, 15.3mmol, 1.30eq) followed by MeI (0.956mL, 15.3mmol, 1.3eq). The reaction mixture was stirred at 40°C under argon for 5h and then quenched with HCl 6N (10ml). The mixture was diluted with HCl 1 N (250 mL) and extracted with DCM (3x100 mL). Pooled organics were washed with water (3x 100mL) then brine (1x100mL). The solution was dried over Na_2SO_4 , filtered and concentrated to dryness. The residue was purified by flash chromatography (EtOAc/Hexane, 0% to 20%) to afford **188** (2.42g, 8.16mmol). Yield 69%.

Part F.




Aldehyde **188** (2.41g, 8.13mmol, 1eq) and hydroxylamine hydrochloride (678mg, 9.75mmol, 1.2eq) were dissolved in THF/MeOH/water (7:4:1, 12mL). The solution was stirred at room temperature for 5 min then a crystal of methyl orange and sodium cyanoborohydride (1.02g, 16.26mmol, 2eq) were added. The pH was adjusted to 2 and the resulting ruby red color was maintained for the duration of the reaction by the regular addition of 6N HCl. After stirring for 1 h, the mixture was filtered on paper, rinsing with

THF. The filtrate was diluted with NaOH 1 N (100mL) and extracted with DCM (3x 50mL). Pooled organics were washed with water (3x20mL), brine (1x20mL) and then dried over Na₂SO₄. The crude product was purified by flash chromatography (Hexane/EtOAc, 0% to 80%) to afford **189** (1.498g, 4.78mmol). Yield 58%.

5


Part G.



Nitrite **122** (850mg, 2.66mmol, 1.05eq) was dissolved in DMSO (5mL) and sodium acetate (327mg, 3.99mmol, 1.50eq) was added. The solution was stirred at room temperature for 30min then poured on brine (30mL) and extracted with diethyl ether (3x30mL). The combined organic layers were washed with saturated NaHCO₃ (1x20mL), water (2x20mL) and brine (1x20mL), dried over Na₂SO₄, filtered and concentrated to dryness. The residue was then reacted overnight with hydroxylamine **189** (588mg, 2.35 mmol, 1 eq) in refluxing toluene (20mL). After 14h, the solution was concentrated and the residue purified by flash chromatography (Hexane/EtOAc, 0% to 30%) to afford **190** (964mg, 1.69mmol). Yield 66%.

15

Part H.



20


Lactone **190** (964mg, 1.69mmol, 1eq) is dissolved in THF (8 mL) and HCl 6N (0.6 mL) is added. The solution is stirred at room temperature for 2 h then diluted with DCM (100 mL) and washed with 5% NaHCO₃ (3x) and brine (1x), dried over Na₂SO₄, filtered and concentrated to dryness.

25

To a solution of the residue in dry DCM (6mL) was added a solution of (+)-isopinocampheylamine (569μL, 3.39mmol, 2eq) previously treated in dry DCM (6mL) with

2.0M trimethylaluminum in hexanes (1.83 mL, 3.39mmol, 2eq) for 15 min. The solution was stirred at room temperature overnight. After 14h, was added a solution of (+)-isopinocampheylamine (285 μ L, 1.69mmol, 1eq) previously treated in dry DCM (3mL) with 2.0M trimethylaluminum in hexanes (850 μ L, 1.69mmol, 1eq) for 15 min. After 5h, the solution was diluted with dry DCM (150mL) and quenched with Na₂SO₄·10H₂O (8.2g, 25.4mmol, 15eq). The mixture was stirred vigorously at room temperature for 15h then filtered on Celite. The filtrate was concentrated to dryness and the residue purified by flash chromatography (Hexane/EtOAc, 0% to 100%) to afford **191** (659mg, 1.09mmol). Yield 64%.

Part I.




A flask containing iodide **191** (659mg, 1.09mmol, 1eq), 3-amino-5-carboxyphenylboronic acid (240mg, 2.17mmol, 2eq), cesium carbonate (628mg, 3.25mmol, 3eq), potassium acetate (107mg, 1.09 μ mol, 1eq) and Pd(dppf)Cl₂ (79mg, 109 μ mol, 0.1eq) was purged with argon and DMSO (5mL) was added. The reaction was heated at 60°C for 24h. The reaction mixture was added to water (100mL), acidified with 6M HCl until the aqueous layer attained a pH of 4, and extracted with DCM (3x50 mL). The combined organic layers were washed with water (1x50mL) dried over Na₂SO₄ and concentrated to a brown oil.

This crude oil was dissolved in MeOH (10mL) and treated with 37% formalin (162 μ L, 2.17mol, 2eq) and sodium cyanoborohydride (205mg, 3.25mmol, 3eq). After stirring at room temperature for 30min, the reaction mixture was quenched with sat NaHCO₃ (4mL) and partitioned between water (50mL) and DCM (20mL). The aqueous layer was acidified to pH 4 with 6M HCl. The layers were separated and the aqueous extracted with DCM (3x20 mL). The combined organic layers were washed with water

(1x20), dried over Na₂SO₄, and concentrated to a brown oil which was used without purification in the next reaction.

Part J.




5

Compound **192** was taken in DCM (2mL) then amine derived from luecine (78mg, 543μmol, 0.5eq), HBTU (206mg, 543μmol, 0.5eq) and triethylamine (151μL, 1086μmol, 1eq) were added. The reaction was stirred at rt for 1 h then concentrated to dryness. The residue was purified by flash chromatography (DCM and DCM/MeOH/NH₄OH 4:1:0.1 0% to 100%) then by HPLC to afford **183** (5.0mg, 6.5μmol) as a lyophilized powder. Yield 0.6% (from **191**). MS (ESI(+)) m/e 770.49 (M+H)⁺.

10


Example 75



193

15

Part A.




20

To a solution of benzoic acid derivate **10** (50 mg, 0.08 mmol, 1.0 eq.) in DMF(3 mL) was added 2M solution of (S)-N⁴-Benzyl-2-Benzylpiperazine (0.04 mg, 0.2 mmol, 2 eq.) in THF followed by triethylamine (10 mg, 0.1 mmol, 1.5 eq.) and HBTU (30 mg, 0.07


mmol, 0.9 eq.). After stirring for 40 h at 23 °C, the reaction mixture was diluted with EtOAc (100 mL) and a saturated NaCl solution (200 mL). The pH of the mixture was adjusted to 12 with 6 N NaOH and extracted with Ethyl acetate (2 x 100 mL). The combined organic extracts were dried (Na₂SO₄), filtered and concentrated *in vacuo* to give of 100 mg of a brown oil. This material was purified using silica gel chromatography (2.5 – 10% MeOH/DCM) to give 33 mg of 194 as brown solid. Yield 47%. MS (ESI(+)) *m/z* 858.62 (M+H)⁺.

Part B.



To a solution of (S)-N⁴-Benzyl-2-Benzylpiperazine amide 194 (27 mg, 0.031 mmol, 1 eq.) in MeOH (3 mL) under H₂ (balloon) was added 44 mg Pd(OH)₂ (44 mg, 0.031 mmol, 1 eq.). After stirring at 23 °C for 18 h, the reaction mixture was filtered through Celite 545 and washed with MeOH (2 x 50 mL). The filtrate was concentrated *in vacuo* to provide crude 195 (14.5 mg) as a light brown solid, which was used directly on the next reaction. Yield: 60%. MS (ESI(+)) *m/z* 768.37 (M+H)⁺.


Part C.



To a solution of 12 mg starting material 195 (0.02 mmol, 1 eq.) in EtOH (1 mL) was added formaldehyde of a 37% aqueous solution (4 mg, 0.05 mmol, 3 eq.), acetic acid (0.9 mg, 0.02 mmol, 1 eq.), and sodium cyanoborohydride (3 mg, 0.05 mmol, 3 eq.). After stirring at 23 °C for 16 h, the reaction mixture was filtered through Celite 545 and

concentrated to dryness to afford an orange solid. The material was diluted with a saturated solution of NaCl, water and EtOAc. The pH of the mixture was adjusted to 10 with 6 N NaOH and extracted with Ethyl acetate (2 x 50 mL). The combined organics were dried over MgSO₄, filtered and concentrated *in vacuo*. The crude product was purified by HPLC to afford 4 mg of (S)-N^d-methyl-2-benzylpiperazine amide product 193. Yield 33%. MS (ESI(+)) *m/z* 782.58 (M+H)⁺.


Example 76



196

10

Part A.



9


197

To a solution of 9 (1 g) in toluene (10 mL) under Argon was added dioxaborolane (1 mL), SPHOS (0.04 g), triethylamine (0.7 mL) and Pd catalyst (0.02 g). The solution was purged with Argon then heated at 80 °C for 2 h. LCMS showed mainly starting material. Another 60 mg SPHOS, 30 mg Pd catalyst and 0.3 mL of dioxaborolane were added. The solution was heated at 80 °C for another 5h, and then cooled to 23 °C and methanol (10 mL) was added. The mixture was push through a short plug of silica gel and concentrated. Flash chromatography on silica gel (Hexane to 30%, 50% ethyl acetate in hexanes) gave 1 g of desired product 197.

15


20

Part B.




To a solution of pinacolboron **197** (1 g) in mixed solvents of acetone and water (1:1, 20 mL) at rt was added sodium periodate (2 g) and ammonium acetate (0.8 g). The mixture was allowed to stir at 23 °C for 12 hr. The reaction was evaporated to dryness and the residue taken up in ethyl acetate (100 mL), 0.1 N HCl (10 mL) was used to wash the organic layer. The organic layer washed with brine, dried over MgSO₄ and evaporated to provide **198** as a yellow solid 750 mg and used directly in the next reaction (Part D of this example).

Part C.




To a solution of dihydro benzofuran carboxylic acid **199** (150 mg) in acetic acid (3 mL) at 0 °C was added an acetic acid solution of bromine (120 µl in 1 mL of acetic acid) drop wise. The mixture was allowed to stir at 23 °C for 12 hr. The reaction was quenched with sodium sulfite solution (2 M) until the red color disappeared. The mixture was concentrated under reduced pressure and residue was taken up in 150 mL DCM, which was washed with 2 M sodium sulfite, brine, dried and concentrated to give 200 mg of crude product **200** which was used directly in the next reaction (Part D of this example).

Part D.



To a solution of boronic acid **198** (100 mg) in DMSO (5 mL) at 23 °C was added aryl bromide **200** (85 mg), cesium carbonate (200 mg), potassium acetate (20 mg) and Pd catalyst (17 mg). The mixture was purged with Argon and heated at 75 °C for 5 h. LCMS showed desired product and reaction was concentrated. HPLC purification gave 20 mg of
5 desired product **201**.

Part E.





10

To a solution of **201** (5 mg) in DCM (1 mL) was added crude amine (10 uL), HATU (4 mg). The mixture was allowed to stir at 23 °C for 2 hrs. The crude mixture was then diluted with methanol (800 uL) and purified by HPLC (basic 10-100) to give desired product **196** 3.5 mg. MS (ESI(+)) m/e 777.58 (M+H)⁺.


15

Example 77

20 Part A.


To a solution of boronic acid **198** (50 mg) in DMSO (1.5 mL) at 23 °C was added aryl bromide **203** (35 mg), cesium carbonate (100 mg), potassium acetate (10 mg) and Pd catalyst (9 mg). The mixture was purged with Argon and heated at 75°C for 5 h. LCMS showed desired product and then reaction concentrated. Purification by HPLC (acidic 10-100) resulted in 30 mg of desired product **204**.

Part B.





10 To a solution of aldehyde **204** (30 mg) in THF (1 mL) was added isoprene (51 μ L), 2.7M phosphate buffer (0.15 mL), and NaClO₂ (18 mg) at 23 °C. The reaction was stopped after 2h by adding water (30 mL), acidifying to pH 1 with 6M HCl, and extraction with DCM (3x15 mL). The organic layers were dried on Na₂SO₄ and concentrated to give **205** as a white foam and used without further purification in part C of this example.

Part C.



To a solution of **205** (7 mg) in DCM (0.6 mL) was added crude amine (10 uL), HATU (4 mg). The mixture was allowed to stir at 23 °C for 2 hr. The crude mixture was diluted with methanol (800 uL) and purified by JPLC (basic 10-100) gave desired product **202** 3 mg. MS (ESI(+)) m/e 751.72 (M+H)⁺.


Example 78

**206**Part A.

5

To a solution of phenylalanine **207** (0.5 g) in mixed solvents of dioxane (5 mL) and 1 N sodium hydroxide (5 mL) was added di-tert-butyl dicarbonate (0.7 g). The reaction mixture was stirred at 23 °C for 1.5 h. KHSO₄ (1 M) was added until the pH of the solution was 5. Ethyl acetate (3X 100 mL) was used to extract the mixture. The combined organic layers were washed with brine, dried with Na₂SO₄ and concentrated to give 583 mg crude product **208** which was used without further purification in part B of this example.


10

Part B.

15


To a solution of Boc protected phenylalanine **208** (583 mg) in DCM (10 mL) at 23 °C was added HATU (544 mg), dimethyl amine (203 mg) and DIPEA (0.8 mL). The reaction mixture was stirred at 23 °C for 2 hr. The reaction mixture was then diluted with DCM (50 mL), washed with sat. sodium bicarbonate, sat. NH₄Cl, brine, dried over MgSO₄ and then concentrated to dryness. Flash chromatography on silica gel (10% ethyl acetate in Hexane, 40%, 60%) gave 500 mg desired product **209** (66% for two steps).

20

Part C.

209 (500 mg) was dissolved in methanol (10 mL). The reaction solution was purged
5 with N₂, palladium on carbon (10%, 0.6 g) was added, and a balloon full of H₂ was
attached to the reaction flask. The reaction was stirred at 23 °C for 45 min and then the
mixture was filtered through a short plug of celite. The Celite plug was washed with 2
portions of ethyl acetate. The organic solutions were combined and concentrated to give
450 mg of **210**.


10

Part D.

210 (100 mg) was dissolved in dry THF (1 mL) at 23 °C followed by the additoion
15 of HCl (4 M, 1 mL). The mixture was stirred at 23 °C for 4 hr then concentrated to
dryness. Ethyl acetate (2 X 5 mL) was used to drive off the excess HCl to give 105 mg
crude **211** which was used without further purification in part E of this example.

Part E.


20



To a solution of **211** (100 mg) in THF (1.5 mL) under argon at 0 °C was added a
solution of LAH (1 M, 1.7 mL) drop wise. The resulting mixture was warmed to 23 °C and
25 stirred 12h. The reaction mixture was quenched by the slow addition of water (0.2 mL),
15% NaOH (0.2 mL), and water (0.6 mL). Subsequent addition of 10ml sodium/potassium


tartrate solution and DCM (3X100 mL) was used to extract the aqueous solution. The combined organic solution was dried with MgSO_4 and concentrated to give crude oily product **212** (70 mg), which was used without further purification in part F of this example.

5 Part F.




To a solution of **10** (14 mg) in DCM (3 mL) was added crude amine **212** (13 mg), HBTU (14 mg) and DIPEA (14 μl). The mixture was allowed to stir at 23 °C for 2 hr at which point the reaction mixture was diluted with DCM (50 mL) washed with sat. Sodium bicarbonate, sat. NH_4Cl , brine, dried over MgSO_4 and concentrated. HPLC purification (acidic method, 10-100) gave 2.8 mg of desired product **206**. MS (ESI(+)) m/e 785.61 ($\text{M}+\text{H}$)⁺.

Example 79




Part A.




To a 0 °C solution of lactone **8** (1.8 g, 4.29 mmol) in DCM (42 mL) at 0 °C was added triflic anhydride (0.867 mL, 5.15 mmol) dropwise over the course of 5 min. After 30 min, the reaction mixture was diluted with DMF (10 mL) followed by the addition of sodium azide (0.837 g, 12.9 mmol) in a single portion. The reaction mixture was removed from the ice bath and was allowed to warm slowly to 23 °C. After stirring for 12 h, the reaction was quenched by the addition of water, extracted with EtOAc (2x100 mL), dried over MgSO₄, filtered and concentrated to an oil. The crude oil was purified by gradient flash chromatography (30-80% EtOAc/hexanes) to afford 1.7 g (89%) of azide **214**.

10 Part B.




To a 0 °C solution of (+)-isopinocampheylamine (5.75 mL, 34.3 mmol) in THF (60 mL) was added 2M DIBAL in toluene (14.3 mL, 28.6 mmol). After stirring for 2 h, the reaction mixture was added to a solution of lactone **214** (2.54 g, 5.72 mmol) dissolved in THF (10 mL). The combined reaction mixture was stirred for 2 h, then poured into an Erlenmeyer flask containing a saturated solution of Rochelle salts and EtOAc. After stirring for 5 h, the reaction mixture was extracted with Ethyl Acetate (3x100 mL), dried (MgSO₄), filtered and concentrated to an oil. The crude oil was purified by gradient flash chromatography (30-70% EtOAc/hexanes) to afford 2.79 g (82%) of azide **215**.

Part C.




Isoxazolidine **215** (302 mg, 0.505 mmol), pinacol boronate **49** (368 mg, 1.26 mmol), Pd(dppf)Cl₂ (82 mg, 0.101 mmol), potassium acetate (65 mg, 0.657 mmol), and cesium carbonate (494 mg, 1.52 mmol) were dissolved in anhydrous DMSO (5 mL) and flushed under positive argon pressure. The reaction mixture was stirred at 60 °C for 2 h, then allowed to cool to ambient temperature. The reaction mixture was diluted with EtOAc (20 mL) and water (10 mL) and the aqueous phase was adjusted to pH 4 with 6M HCl. The reaction mixture was extracted with EtOAc (3x100 mL) and the combined organics were dried (MgSO₄), filtered and concentrated to an oil. The crude oil was purified by gradient flash chromatography (10-35% Acetone/hexanes) to afford 170 mg (53%) of biphenyl acid **216**.

Part D.




Biphenyl acid **216** (320 mg, 0.504 mmol) and HBTU (287 mg, 0.756 mmol) were dissolved in DMF (5 mL) followed by addition of (R)-1-methylpyrrolidin-3-amine (101 mg, 1.01 mmol). After stirring for 2 h, the reaction mixture was diluted with EtOAc (100 mL) and a saturated sodium bicarbonate solution. The mixture was extracted with EtOAc (3x100 mL), dried (MgSO₄), filtered and concentrated to an oil. The crude oil was purified by gradient flash chromatography (2 -5% MeOH/DCM) to afford 176 mg (49%) of azide **217**.

Part E.



Azide **217** (13 mg, 0.018 mmol) and dithiotreitol (8.7 mg, 0.054 mmol) were dissolved in DMF (1 mL), followed by addition of DBU (8.5 μ L, 0.054 mmol). After stirring for 1 h, the reaction mixture was purified directly on HPLC (Acetonitrile/ 40 mmol ammonium bicarbonate aqueous), to yield 6 mg (47%) of amine **213**. MS ((ESI(+)) m/e 691.5 (M+H)⁺.


Example 80




Part A.

Compound **218** was attained using compound **21** and following the protocol outlined in Example 79. MS ((ESI(+)) m/e 783.1 (M+H)⁺.

Example 81




Part A.




To a solution of MgCl_2 (powder 325 mesh, 5.0 g, 52 mmol, 2 eq), paraformaldehyde (3.0 g, 79 mmol, 3 eq) and Et_3N (7.0 mL, 52 mmol, 2 eq) in THF (60 mL) was added **220** (5.0 g, 26 mmol, 1 eq), heated in the microwave at 160 C for 15 min. TLC (3:2 Hexane:DCM) showed complete consumption of **220**. THF was evaporated and the reaction mixture was taken up in EtOAc, washed with brine, dried, filtered and concentrated in *vacuo* to afford 5.2 g of **221** which was used without purification. Yield 93 %.

10 Part B.




To a solution of **221** (6.0 g, 31 mmol, 1 eq) in DMF (38 mL) was added K_2CO_3 (3.0 g, 41 mmol, 1.3 eq), stirred at rt for 10 min, resulting in a suspension. Iodomethane (3.0 mL, 41 mmol, 1.3 eq) was added dropwise and allowed to stir at rt overnight. TLC (9:1 Hexane:EtOAc) showed no remaining **221**. The mixture was diluted into water and extracted with EtOAc. The organic was separated, washed with brine, dried, filtered and concentrated in *vacuo* to afford **222** as an oil. The crude material was purified by column chromatography (silica gel, 50 % DCM in hexane) to afford 2.4 g of **222** as a white solid.

Part C.




To a solution of **222** (2.5 g, 11 mmol, 1 eq) in MeOH-THF (3:1, 20 mL) was added an aqueous solution of hydroxylamine hydrochloride salt (0.75 g, 11 mmol, 1.2 eq in 4 mL of water) in one portion. The pH was adjusted to 9 with NaOH (6N), and stirred at rt for 1 hr where TLC (2:1 Hexane:EtOAc) showed complete consumption of **222**. NaBH₃CN (1.3 g, 21 mmol, 2 eq) was added with a crystal of methyl red and the solution acidified to pH 2-3 using HCl in MeOH (20 V/V). The pH of the reaction solution was maintained at pH 3 over the course of 12 h by addition of small amounts of the methanolic HCl solution, where it was basified to pH 9 with NaOH (2 N). The solution was extracted with EtOAc, washed with brine, dried, filtered and concentrated in *vacuo* to afford an orange oil. The crude material was purified by column chromatography (silica gel, 55 % EtOAc in Hexane) to afford 2.7 g of **223** as a cream solid.

Part D.



To a solution of **223** (1.8 g, 7.0 mmol, 1 eq) and glyoxylate ester **224** (1.0 g, 9.0 mmol, 1.3 eq) in diethyl ether (20 mL) was added anhydrous CaCl₂ (1.0 g, 9.0 mmol, 1.3 eq), left stirring at rt for 3 h, resulting in a suspension. The suspension was filtered through a Celite plug washing with DCM and ether. The resultant yellow solution was concentrated in *vacuo* to afford 2.0 g of **225** as yellow oil and used directly without purification.


Part E.



To a solution of **225** (0.497 g, 1.48 mmol, 1 eq) in anhydrous THF (10 mL) was added (S,Z)-5-tert-butyldimethylsiloxy)pent-3-en-2-ol **6** (0.5 mL, 1.77 mmol, 1.2 eq) and Ti(iOPr)₄ (0.65 mL, 2.22 mmol, 1.5 eq) heated in the microwave at 140 C for 15 min. TLC (30:1 DCM:Et₂O) showed consumption of nitrone **225** and the allylic alcohol **6**. 3-(dimethylamino)-1,2-propanediol (1 mL) in EtOAc (1 mL) was added and the dark brown solution left stirring at rt overnight. The dark brown solution was diluted with EtOAc and

quenched with Rochelle's salt solution, washed with water, brine, dried, and concentrated in *vacuo* to afford a brown oil. The crude material was purified by column chromatography (silica gel, 10 % EtOAc in Hexane) to afford 0.490 g of **226** as a brown oil.

Part F.




5

To a solution of **226** (0.40 g, 0.903 mmol, 1 eq) in THF (5 mL) was added concentrated 6 N HCl (0.45 mL, 2.71 mmol, 3 eq) and stirred at rt for 2 h. TLC (1:2 hexane:EtOAc) of neutralized aliquot showed complete consumption of **226**. The reaction was neutralized with NaHCO₃, extracted with EtOAc, washed with brine, dried, filtered and concentrated in *vacuo* to afford a brown oil. The crude material was purified by column chromatography (silica gel, 50 % EtOAc in Hexane) to afford 0.251 g of **227** as a brown solid. 81% yield

10

Part G.



15


To a solution of (+-) isocamphenylamine (0.314 g, 2.0 mmol, 2 eq) flushed with Ar in anhydrous DCM (5 mL), was added Me₃Al (2 M solution in Hexane, 0.513 mL, 1.02 mmol, 2 eq) dropwise over 20 min and the resultant clear solution stirred for 40 min at rt. A solution of the lactone **227** (0.4 g, 0.1.0 mmol, 1 eq) in DCM (15 mL) was added by dropwise and the reaction mixture was stirred at rt for 12 h until all the **227** was consumed as indicated by TLC (2:1 EtOAc:Hexane). The reaction was diluted with DCM and quenched on dropwise addition to a rapidly stirring solution Rochelle's salt, and stirred at rt for 2 h. The organic layer was separated and washed with water, brine, dried, and concentrated in *vacuo* to afford a yellow oil. The crude material was purified by column

20

25


chromatography (silica gel, 50 % EtOAc in Hexane) to afford 0.33 g of **228** as a yellow oil. Yield 53 %

Part H.




To a solution of isoxazolidine **228** (100 mg, 0.179 mmol, 1 eq) in DMSO (5 mL), flushed with argon, was added the pinacolboronate (68 mg, 0.232 mmol, 1.3 eq), potassium acetate (26 mg, 0.268 mmol, 1.5 eq) and cesium carbonate (175 mg, 0.526 mmol, 3 eq). Upon stirring for 10 min, Pd(dppf)Cl₂ (29 mg, 0.036 mmol, 0.2 eq) was added as a single portion. The mixture was heated to 70 °C for 4 h, then allowed to cool to rt. EtOAc and brine were added and the pH adjusted 3-4 with HCl (2N). The organic layer was separated, washed with water, brine, dried, and concentrated in *vacuo* to afford a brown black oil. The crude material was purified by column chromatography (silica gel, 4% MeOH in DCM) to afford 35 mg of **229** as a brown solid. 30% yield.


Part I.



A solution of **229** (35 mg, 0.055 mmol, 1 eq) in DCM (3 mL) was added Et₃N (23 μL, 0.165 mmol, 3 eq) and HBTU (42 mg, 0.110 mmol, 2 eq) and allowed to stir at rt for 10 min. To this solution was added the amine (9.8 mg, 0.055 mmol, 1 eq) and allowed to stir at rt for 2 h. The reaction was taken up in DCM, washed with K₂CO₃, water, brine, dried, and concentrated in *vacuo* to afford a brown solid. The crude material was diluted with MeOH (2 mL) and purified by reverse-phase HPLC (MeCN/water with 40 mM NH₄HCO₃) to afford 13 mg of **219** as a white solid. 20% yield


Example 82**230**

5

Part A


Aldehyde **231** (12 g, 33 mmol, 1 eq) and hydroxylamine hydrochloride (2.7 g, 39 mmol, 1.18 eq) were dissolved in THF/MeOH (3:1, 60 mL). Water (2 mL) was added and the pH was adjusted to 9 with 6 N KOH. The reaction mixture was stirred at rt overnight and then NaBH₃CN (3.1 g, 49 mmol) was added followed by a crystal of methyl orange. The pH was adjusted to 3 and the resulting ruby red color was maintained for the duration of the reaction by the frequent addition of 1 N HCl. After stirring for 2 h another portion of NaBH₃CN (1 g, 13 mmol, 0.4 eq) was added. The solution was stirred for 16 h and then neutralized to pH 7 and diluted with DCM. The mixture was washed with water (3 x 10 mL), brine and dried over MgSO₄, filtered and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (50-100% EtOAc/hexane) to afford 8 g of compound **232**. Yield 64%.

20

Part B


To a solution of **232** (0.486 g, 1.83 mmol, 1 eq) and glyoxylate ester (0.285g, 2.37 mmol 1.3 eq) in diethyl ether (10 mL) was added anhydrous CaCl_2 (0.263 g, 2.37 mmol, 1.3 eq), left stirring at rt for 3 h, resulting in a suspension. The suspension was filtered through a Celite plug washing with DCM and ether. The resultant yellow solution was concentrated *in vacuo* to afford 0.497 g of **233** as yellow oil and used directly without purification.

Part C



Nitron **233** (5 g, 11 mmol, 1 eq), allylic alcohol **6** (2 g, 11 mmol, 1 eq) and $\text{Ti}(\text{iOPr})_4$ (4 g, 4 mL, 13 mmol, 1.18 eq) were dissolved in toluene (40 mL) and heated in a microwave at 120 °C for 10 min. The reaction mixture was diluted with EtOAc (10 mL) and 3-(dimethylamino)-1,2-propanediol (4 mL) was added. After stirring for 2 h, EtOAc (10 mL) was added and the mixture was washed with water (3 x 10 mL), brine (10 mL), dried over MgSO_4 , filtered and concentrated *in vacuo*. The crude residue was purified by silica gel chromatography (10-30% Hexane-EtOAc) to afford 2.5 g of compound **234**. Yield 35%.


Part D



To a solution of PMB ether **234** (2 g, 3 mmol, 1 eq) in DCM (150 mL) was added TFA (3 g, 31 mmol, 10.33 eq) dropwise at 0 °C. The solution was stirred for 1.5 h and quenched with saturated NaHCO_3 (60 mL). The organic phase was separated, washed with brine, dried over Na_2SO_4 , filtered and concentrated *in vacuo* to afford an oil. The resulting

oil was purified by silica chromatography (10-30% hexane-EtOAc) to afford 1.2 g of compound **235**. Yield 74%.

Part E




5

To a solution of (+) isopinocampheylamine (0.6 g, 0.7 mL, 4 mmol, 2 eq) in DCM (10 mL) at rt was added Me_3Al (0.4 g, 3 mL, 2 M solution in toluene, 6 mmol, 3 eq) dropwise over 2.5 min. The solution was stirred at rt for 10 min prior to the dropwise addition of a solution of lactone **235** (1 g, 2 mmol, 1 eq) in DCM (15 mL). The reaction was stirred for 24 h, diluted with DCM (125 mL) and a saturated solution of Rochelle's salt (125 mL). The mixture was vigorously stirred for 2 h until two phases formed. The layers were separated and the organic phase was washed with water, brine, dried over MgSO_4 , filtered and concentrated *in vacuo* to afford a solid. The solid was purified by silica gel chromatography (25% hexane/EtOAc) to afford 0.5 g of compound **236**. Yield 39%.

15


Part F



Phenol **236** (187 mg, 0.27 mmol, 1 eq) was dissolved in DMF (3.5 mL) and treated with K_2CO_3 (111 mg, 0.8 mmol, 3 eq), allylbromide (49 mg, 0.4 mmol, 1.48 eq). The solution was stirred for 2.5 h, diluted with water and extracted with diethyl ether (3×4 mL). The organic phase was separated, dried over Na_2SO_4 , filtered and concentrated *in vacuo* to afford a solid. The resulting solid was dissolved in THF/ Et_3N (1:1, 6 mL) and treated with a solution of HF/pyridine (1 mL) at 0 °C. The solution was stirred for 1 h at rt and quenched with TMSOMe (25 mL) and concentrated *in vacuo* to afford a solid which was purified by silica gel chromatography (20% DCM/hexane) to afford 0.21 g of compound **237** as a white solid. Yield 67%.


25

Part G



To a solution of the alkene **237** (0.14 g, 0.2 mmol, 1 eq) in *t*-BuOH (16 mL), THF (8 mL) and H₂O (2 mL) was added NMO (80 mg, 0.8 mmol, 4 eq) followed by the dropwise addition of OsO₄ (0.21 g, 2.9 mL, 2.5 % solution in 2-methyl-2-propanol, 0.02 mmol, 0.1 eq). After 3 h, the reaction mixture was diluted with DCM (5 mL), brine and 10% Na₂S₂O₃ and the organic phase was separated. The aqueous phase was extracted with DCM (2 × 60 mL), and the combined organics were dried over Na₂SO₄, filtered and concentrated *in vacuo* to afford a solid. The solid was dissolved in THF/water (10:1, 1.2 mL) and treated with periodic acid (80 mg, 0.4 mmol, 2 eq) in single portion and stirred for 12 h. The solution was diluted with DCM (5 mL) and washed with brine, dried over Na₂SO₄, filtered and *in vacuo* to afford an oil. The resulting oil was suspended in DCM (5 mL) and treated with AcOH (10 μ L), morpholine (40 mg, 0.4 mmol, 2 eq) and sodium triacetoxyborohydride (140 mg, 0.4 mmol, 2 eq) and stirred for 12 h. The solution was diluted with 0.1 M NaOH (1 mL), saturated NaCl (1 mL) and extracted with EtOAc (3 × 3 mL). The combined organic was concentrated *in vacuo* to afford an oil. The oil was purified by silica gel chromatography (DCM/MeOH/AcOH, 99.5:0:0.5 to 97.5:2:0.5) to afford 75 mg of compound **238**.


Part H



A flask containing aryl iodide **238** (25 mg, 0.04 mmol, 1 eq), pinacolboronate (17 mg, 0.06 mmol, 1.5 eq), KOAc (5 mg, 0.048 mmol, 1.2 eq), cesium carbonate (39 mg, 0.12 mmol, 3 eq) and Pd(dppf)Cl₂ (6.5 mg, 8 μ mol, 0.2 eq) was purged with argon and DMSO (2 mL) was added. The mixture was heated to 70 °C for 3 h and then cooled to rt. The solution was diluted with DCM (10 mL), water (5 mL) and the pH was adjusted to 6.8 with 0.1 N HCl. The aqueous phase was extracted with DCM (2 × 10 mL) and the combined organics were washed with brine (20 mL), dried over MgSO₄, filtered and concentrated *in*

vacuo to afford a dark brown oil. The resulting oil was purified by flash chromatography (2.5-5% CH₂Cl₂/MeOH) to afford 16 mg of compound **239** as a yellow solid. Yield 54%.


Part I



- 5 To a solution of **239** (8 mg, 0.02 mmol, 1 eq) in DMF (1.5 mL) was added HBTU (9 mg, 0.02 mmol, 1 eq), amine (4 mg, 0.02 mmol, 1 eq) and Et₃N (4 mg, 5 uL, 0.04 mmol, 2 eq). The solution was stirred for 2 h, diluted with water (0.5 mL) and purified by reverse-phase HPLC (MeCN/water with 40 mM NH₄HCO₂) to yield 7 mg of compound **230**. Yield 67%. MS (ESI(+)) *m/z* 870.1 (M+H)⁺.


10

Example 83




240

Part A




- 15 Methyl (3-Amino-5-carboxyphenyl)boronate **241** (1.5 g, 8.3 mmol, 1 eq) and pinacol (2.9 g, 25 mmol, 3 eq) were combined with THF (7 mL) and heated in a microwave to 140 C for 10 min, the reaction mixture was cooled to rt and then concentrated to an orange oil. The crude material **242** was used without further purification
- 20

Part B




To a solution of aniline **242** (0.4 g, 1.9 mmol) in DMF (5 mL) was added bis(chloroethyl) ether (0.27 g, 1.9 mmol), potassium iodide (0.8 g, 5.7 mmol) and heated to 80 °C for 24 h. The solution was diluted with water (10 mL) and extracted with diethyl ether (2 × 5 mL). The organic layer were washed with water (2 × 5 mL), brine (5 mL), dried over Na_2SO_4 , filtered and concentrated *in vacuo* to afford ester as an oil. To a solution of the ester in THF (2.5 mL) was added a solution of 2N LiOH (0.5 mL). The resultant solution was stirred for 2 h, diluted with water (10 mL) and extracted with EtOAc (2 × 10 mL). The organic layers were washed with brine (10 mL), water (10 mL) and dried (Na_2SO_4), filtered and concentrated *in vacuo* to afford an oil. The oil was subjected to column chromatography (30% EtOAc in hexane) to afford **243** as a white solid. 42% yield

Part C




A flask containing aryl iodide **9** (40 mg, 0.07 mmol, 1 eq), pinacolboronate **243** (30 mg, 0.09 mmol, 1.5 eq), KOAc (6.8 mg, 0.069 mmol, 1.2 eq), cesium carbonate (68 mg, 0.2 mmol, 3 eq) and Pd(dppf)Cl_2 (10 mg, 14 μmol , 0.2 eq) was purged with argon and DMSO (2 mL) was added. The mixture was heated to 70 °C for 3 h and then cooled to rt. The solution was diluted with DCM (10 mL), water (5 mL) and the pH was adjusted to 6.8 with 0.1 N HCl. The aqueous phase was extracted with DCM (2 × 10 mL) and the combined organics were washed with brine (20 mL), dried over MgSO_4 , filtered and concentrated *in vacuo* to afford a dark brown oil. The resulting oil was purified by flash chromatography (2.5-5% $\text{CH}_2\text{Cl}_2/\text{MeOH}$) to afford 15 mg of compound **245** as a yellow solid. Yield 22%.

Part D




A solution of **245** (10 mg, 0.015 mmol, 1 eq) in DCM (1 mL) was added Et₃N (4 uL, 0.031 mmol, 2 eq) and HBTU (8.7 mg, 0.023 mmol, 1.5 eq) and allowed to stir at rt for 10 min. To this solution was added the amine (3.3 mg, 0.023 mmol, 1.5 eq) and allowed to stir at rt for 2 h. The reaction was taken up in DCM, washed with K₂CO₃, water, brine, dried, and concentrated in *vacuo* to afford a brown solid. The crude material was diluted with MeOH (2 mL) and purified by reverse-phase HPLC (MeCN/water with 40 mM NH₄HCO₃) to afford 4 mg of **246** as a white solid. 32% yield MS (ESI(+)) *m/z* 813.0 (M+H)⁺

Example 84




Part A



To a solution of **248** (5 g, 17 mmol, 1 eq) and PyBop (11.6 g, 22 mmol, 1.3) in DCM (60 mL) was added dimethylamine (1.5 g, 17 mL, 34 mmol, 2 eq) at 0 °C. The solution was


stirred for 10 min followed by dropwise addition of DIPEA (4.4 g, 5.9 mL, 34 mmol, 2 eq). After stirring for 6 h, the solution was diluted with sat NaHCO_3 (40 mL) and the aqueous phase was extracted with DCM (2×50 mL). The combined organic phases were washed with brine, separated, dried over MgSO_4 , filtered and concentrated *in vacuo* to afford an oil. The resulting
 5 oil was purified by silica gel chromatography (50-100% hexane/EtOAc) to afford **249** as a clear oil.

Part B



To a solution of amide **249** (4.5 g, 14 mmol, 1 eq) in DCM (100 mL) was added TFA (10 mL) at 0 °C. After stirring for 2 h at rt, the solvent was removed *in vacuo* to afford an oil. The resulting oil was suspended in THF and cooled to 0 °C, to which was added LiAlH_4 (4 g, 108 mmol, 6.8 eq) in portions, and heated at reflux for 12 h under argon. The solution was cooled to rt and quenched with water (4 mL), allowed to stir for 5 min, followed by 15% NaOH (4 mL) and stirred for additional 5 min, and finally water (12 mL) was added and the
 15 suspension was stirred until the white precipitate formed. The solid was filtered and washed with EtOAc and the filtrate was concentrated *in vacuo* to afford **250** as an oil which was used without further purification.

20 Part C




Biphenyl acid **10** (40 mg, 0.064 mmol, 1 eq) and HBTU (49 mg, 0.13 mmol, 2 eq) were dissolved in DMF (1 mL) followed by the addition of *N,N*-dimethyl-1-((2*S*,5*R*)-5-phenylpyrrolidin-2-yl)methanamine **250** (23 mg, 0.13 mmol, 2 eq). The reaction mixture
 25 was stirred for 2 h, diluted with EtOAc and quenched with a saturated aqueous NaHCO_3 solution. The reaction mixture was extracted with EtOAc (3 x 100 mL) and the combined


organics were dried over MgSO_4 , filtered and concentrated *in vacuo* to afford an oil. The crude oil was purified by silica gel chromatography (2 -5% MeOH/DCM , 0.1% Et_3N) to afford 23 mg of biphenyl amide **247**. Yield 45%.

Example 85

5




Part A




10 Biphenyl acid **10** (40 mg, 0.064 mmol, 1 eq) and HBTU (49 mg, 0.13 mmol, 2 eq) were dissolved in DMF (1 mL) followed by the addition of (*S*)-*N*¹,*N*¹-dimethyl-3-(thiophen-2-yl)propane-1,2-diamine (23 mg, 0.13 mmol, 2 eq). The reaction mixture was stirred for 2 h, diluted with EtOAc and quenched with a saturated aqueous NaHCO_3 solution. The reaction mixture was extracted with EtOAc (3 x 100 mL) and the combined organics were dried over MgSO_4 , filtered and concentrated *in vacuo* to afford an oil. The crude oil was purified by silica gel chromatography (2 -5% MeOH/DCM , 0.1% Et_3N) to afford 23 mg of biphenyl amide **251**. Yield 45%.

Example 86

20




Part A.




Biphenyl acid **216** (130 mg, 0.205 mmol, 1 eq), Et₃N (62 μ L, 0.615 mmol, 3 eq) and HBTU (155 mg, 0.410 mmol, 2 eq) were dissolved in DCM (5 mL) followed by addition of amine (74 mg, 0.410 mmol, 2 eq). After stirring for 2 h, the reaction mixture was diluted with EtOAc (100 mL) and a saturated sodium bicarbonate solution. The mixture was extracted with EtOAc (3 x 100 mL), dried over MgSO₄, filtered and concentrated to an oil. The crude oil was purified by silica gel chromatography (2-5% MeOH/DCM) to afford 49 mg of azide **253**. Yield 30 %.

Part B.




Azide **217** (30 mg, 0.038 mmol, 1 eq) and dithiothreitol (17 mg, 0.113 mmol, 3 eq) were dissolved in DMF (1 mL) followed by addition of DBU (17 μ L, 0.113 mmol, 3 eq). After stirring for 1 h, the reaction mixture was purified directly by reverse-phase HPLC (CH₃CN/water with 40 mM NH₄HCO₃), to yield 17 mg of amine **253**. Yield 62 %

Example 87




Part A.




5 A flask containing boronate (20 mg, 0.035 mmol, 1 eq.), 3,4-dibromobenzoic acid (30 mg, 0.11 mmol, 3 eq.), cesium carbonate (50 mg, 1 mmol, 3 eq.), potassium acetate (5 mg, 0.03 mmol, 1 eq.) and Pd(dppf)Cl₂ (5 mg, 0.003 mmol, 0.1 eq.) was purged with argon and DMSO (1 mL) added. The mixture was heated at 60° during 3h, then 45° overnight. The reaction mixture was partitioned between water (30 mL) and DCM (20 mL) and
 10 acidified with 6M HCl until the aqueous layer attained a pH of 1. The layers were separated and the aqueous extracted (2 x 20 mL DCM). The combined organic layers were dried on Na₂SO₄ and concentrated to a brown oil.

To an aliquot containing ca 50% of this crude oil (0.019 mmol, 1 eq.) in DMF (1 mL) was added the amine (8 uL, 0.04 mmol, 2 eq.), DIEA (10 uL) and HBTU (15 mg, 0.04
 15 mmol, 2 eq.). When the reaction appeared complete by HPLC, the mixture was diluted with MeOH (1 mL) and purified by reverse-phase HPLC (MeCN/40 mM ammonium formate 25%→80%) to give 254 as a white solid (4 mg, 27%). MS (ESI(+)) m/e 807.6 (M+H)⁺.

20


Example 88**Part A.**




A flask containing boronate (30 mg, 0.05 mmol, 1 eq.), 3,4-dichlorobenzoic acid (20 mg, 0.1 mmol, 2 eq.), S-Phos (2 mg, 0.005 mmol, 0.1 eq.), potassium phosphate (50 mg, 0.2 mmol, 4 eq.) and a dash of palladium acetate was purged with argon and THF (2 mL) added. The mixture was heated at 40° overnight. The reaction mixture acidified with 6M HCl and extracted with DCM. The combined organic layers were dried on Na₂SO₄ and concentrated to a brown oil.

To an aliquot containing this crude oil (0.05 mmol, 1 eq.) in DMF (1 mL) was added (S)-N1,N1-dimethyl-3-phenylpropane-1,2-diamine (30 μ L, 0.15 mmol, 3 eq.) and HBTU (30 mg, 0.08 mmol, 1.5 eq.). When the reaction appeared complete by HPLC, the mixture was diluted with MeOH (1 mL) and purified by reverse-phase HPLC (MeCN/40 mM ammonium formate 25% $\rightarrow$ 80%) to give 255 as a white solid (3 mg, 8%). MS (ESI(+)) m/e 761.7 (M+H)⁺.



Example 89




Part A.




Compound 256 was synthesized according to the procedure described in Example 88. MS (ESI(+)) m/e 749.6 (M+H)⁺.

Example 905 **Part A.**

A solution of crude compound **1** (8 mg, 0.01 mmol, 1 eq.) in MeCN (500 uL) was treated portionwise with a 1:100 solution of bromine in MeCN until LC showed consumption of **1**. The reaction mixture purified by reverse-phase HPLC (MeCN/40 mM ammonium formate 25%→80%) to give an inseparable mixture that appears by NMR to contain approximately equal amounts of **257** and **258**, as a white solid (3 mg, 30%). MS (ESI(+)) m/e 830.6 (M+H)⁺.

Example 91


15

Part A.

To an aliquot of crude acid (example 3 part G) (0.024 mmol, 1 eq.) in DMF (1 mL) was added amine (11 uL, 0.05 mmol, 2 eq.) and HBTU (30 mg, 0.08 mmol, 3 eq.). When


the reaction appeared complete by HPLC, the mixture was diluted with MeOH (1 mL) and purified by reverse-phase HPLC (MeCN/40 mM ammonium formate 25%→80%) to give **259** as a white solid (3 mg, 17%). MS (ESI(+)) m/e 750.6 (M+H)⁺.

Example 92



5


Part A.




To acid **10** (10 mg, 0.016 mmol, 1 eq.) in DMF (1 mL) was added DIEA (10 μ L), H-L-Leu-NH₂ HCl (8 mg, 0.05 mmol, 3 eq.) and HBTU (20 mg, 0.05 mmol, 3 eq.). When the reaction appeared complete by HPLC, the mixture was diluted with MeOH (1 mL) and purified by reverse-phase HPLC (MeCN/40 mM ammonium formate 25%→80%) to give **260** as a white solid (3 mg, 25%). MS (ESI(+)) m/e 722.6 (M+H)⁺.

15

Example 93





Part A.



A flask containing iodide **9** (200 mg, 0.35 mmol, 1 eq.), boronate (126 mg, 0.7 mmol, 2 eq.), cesium carbonate (340 mg, 1 mmol, 3 eq.), potassium acetate (35 mg, 0.35 mmol, 1 eq.) and Pd(dppf)Cl₂ (20 mg, 0.035 mmol, 0.1 eq.) was purged with argon and DMSO (11 mL) added. The mixture was heated at 60° during 3h, then additional
 5 Pd(dppf)Cl₂ (20 mg, 0.035 mmol, 0.1 eq.) added and stirring continued at ambient temperature overnight. The reaction mixture was treated with tetrabutylammonium bromide (230 mg) and extracted from water (30 mL) 3 x 30 mL DCM. The combined organic layers were washed with brine (20 mL), dried on Na₂SO₄, and concentrated to a brown oil.



10 To an aliquot containing ca. 50% of this crude oil (0.017 mmol, 1 eq.) in DMF (5 mL) was added amine (25 mg, 0.017 mmol, 1 eq.) and HBTU (200 mg, 0.5 mmol, 3 eq.). When the reaction appeared complete by HPLC, the mixture was diluted with MeOH (1 mL) and purified by reverse-phase HPLC (MeCN/40 mM ammonium formate 25%→80%) to give **261** as a white solid. MS (ESI(+)) m/e 708.5 (M+H)⁺.

15


Example 94**Part A.**

20


A flask containing amine **261** (12 mg, 0.017 mmol, 1 eq.) in DCM (1 mL) and MeOH (100 uL) was added DIEA (10 uL, 0.05 mmol, 3 eq.) and acetic anhydride (5 uL, 0.05 mmol, 3 eq.). After shaking overnight, the reaction mixture was diluted with aqueous NaHCO₃ and extracted (3 x 5 mL DCM). The combined organic layers were dried on
 25 Na₂SO₄ and concentrated to give **262** as brownish solid (3 mg 22%). MS (ESI(+)) m/e 750.8 (M+H)⁺.


Example 95**Part A.**

Compound **263** was synthesized according to the procedure described in Example 88 using (R)-N1,N1,4-trimethylpentane-1,2-diamine in place of (S)-N1,N1-dimethyl-3-phenylpropane-1,2-diamine. Yield 9 mg (53%). MS (ESI(+)) m/e 738.5 (M+H)⁺.



Example 96**Part A.**

Compound **264** was synthesized according to the procedure described in Example 88, using (S)-N1,N1-dimethyl-3-phenylpropane-1,2-diamine. Yield 3 mg (16%). MS (ESI(+)) m/e 772.5 (M+H)⁺.


Example 97

Part A.

Compound **265** was synthesized according to the procedure described in Example 88, using amine **266** in place of (S)-N1,N1-dimethyl-3-phenylpropane-1,2-diamine and
 5 intermediate acid described in Example 95 part A.. Yield 3 mg (16%). MS (ESI(+)) m/e 786.5 (M+H)⁺.

Example 9810 Part A.

An aliquot of crude **261** (0.03 mmol, 1 eq.) in MeOH (1 mL) was treated with acetaldehyde (20 uL, 0.3 mmol, 10 eq.) and sodium cyanoborohydride (8 mg, 0.12 mmol, 4 eq.). Acetic acid (5 uL) was added and the solution shaken overnight at 50°. When the reaction appeared complete by HPLC, the mixture was diluted with MeOH (1 mL) and
 15 purified by reverse-phase HPLC (MeCN/40 mM ammonium formate 25%→80%) to give **267** as a white solid (3 mg, 15%). MS (ESI(+)) m/e 764.7 (M+H)⁺.



Example 99

20

Part A.

Compound **268** was synthesized according to the procedure described in Example 98, using glutaraldehyde in place of acetaldehyde. Yield 3 mg (14%). MS (ESI(+)) m/e 776.6 (M+H)⁺.


5

Example 100Part A.

10

An aliquot of crude **261** (0.03 mmol, 1 eq.) in MeOH (1 mL) was treated with acetaldehyde (10 uL, 0.15 mmol, 5 eq.) and sodium cyanoborohydride (6 mg, 0.09 mmol, 3 eq.). The solution was shaken overnight at ambient temperature. The mixture was diluted with MeOH (1 mL) and purified by reverse-phase HPLC (MeCN/40 mM ammonium formate 25%→80%) to give **269** as a white solid (3 mg, 14%). MS (ESI(+)) m/e 736.6 (M+H)⁺.

15



Example 101

20



Part A.

Compound **270** was synthesized according to the procedure described in Example 94, using isobuteraldehyde in place of acetaldehyde. Yield 3 mg (14%). MS (ESI(+)) m/e 764.7 (M+H)⁺.

Example 102

Part A.


- 5 Crude reaction mixture from the production of **270** (0.03 mmol, 1 eq.) was stripped of all volatiles *in vacuo* and taken up in MeOH (1 mL). Acetic acid (5 μL), 37% formalin (10 μL), and sodium cyanoborohydride (4 mg, 0.8 mmol, 3 eq.) were added. After 4h, the mixture was diluted with MeOH (1 mL) and purified by reverse-phase HPLC (MeCN/40 mM ammonium formate 25% $\rightarrow$ 80%) to give **271** as a white solid (3 mg, 14%). MS
- 10 (ESI(+)) m/e 778.7 ($M+H$) $^+$.

Example 10315 Part A.

- To a solution of **216** (45.4 mg, 0.07 mmol, 1 eq) in DCM (5 mL) was added TEA (30 μL , 0.22 mmol, 3 eq) and HBTU (54.2 mg, 0.14 mmol, 2 eq). The solution was stirred at rt for 10 min. Diamine **142** (28.7 mg, 0.11 mmol, 1.5 eq) was then added. After stirring
- 20 at rt for 2h, the mixture was diluted with DCM (15 mL) and washed with saturated


potassium carbonate solution (10 mL). The aqueous layer was extracted with DCM (2 x 10 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated *in vacuo*. The material was purified by silica gel chromatography (80% EtOAc/DCM with 1% triethylamine - 5% MeOH/DCM with 1% triethylamine) to give 40 mg of **273**. Yield 5 69%. MS (ESI(+)) *m/z* 811.28 M⁺.

Part B.




10 To a solution of **273** (40 mg, 0.05 mmol, 1 eq) in DMF (2 mL) at 0 °C was added dithiothreitol (22.8 mg, 0.15 mmol, 3 eq) and DBU (22.3 μL, 0.15 mmol, 3 eq). The solution was stirred at 0 °C for 30 min. The crude reaction was kept at 0 °C and purified by HPLC to give 16 mg of **272**. Yield 41%. MS (ESI(+)) *m/z* 785.33 M⁺.

Example 104




15


Part A.




To a solution of 1-bromo-5-chloro-2-methoxyphenyl **275** (2.5 g, 11 mmol, 1eq) in DCM (100 mL) was added dropwise BBr₃ (1 M solution in DCM, 38.5 mmol, 3.5 eq) over 20 min at - 40 °C. The solution was warmed to rt and stirred for 12 h. TLC (3:2 Hexane:DCM) showed complete consumption of **275**. The solution was quenched with NaHCO₃ and stirred until two phases appeared. The organic was separated, washed with brine, dried, filtered and concentrated *in vacuo* to afford 0.80 g of **276** as a white solid 25 which was used without purification. Yield 32 %

Part B.

To a solution of MgCl_2 (powder 325 mesh, 0.734 g, 7.71 mmol, 2 eq),
 5 paraformaldehyde (0.347 g, 11.57 mmol, 3 eq) and Et_3N (1.08 mL, 7.71 mmol, 2 eq) in
 THF (20 mL) was added **276** (0.800 g, 3.68 mmol, 1 eq), heated in the microwave at 160 C
 for 15 min. TLC (3:2 Hexane:DCM) showed complete consumption of **3**. THF was
 evaporated and the reaction mixture was taken up in EtOAc, washed with brine, dried,
 filtered and concentrated in *vacuo* to afford 0.52 g of **277** which was used without
 10 purification. Yield 47 %

Part C.


To a solution of **277** (0.518 g, 2.2 mmol, 1 eq) in DMF (5 mL) was added K_2CO_3
 (0.456 g, 3.3 mmol, 1.5 eq), stirred at rt for 10 min, resulting in a suspension. Iodomethane
 15 (0.206 mL, 3.3 mmol, 1.5 eq) was added dropwise and allowed to stir at rt overnight. TLC
 (9:1 Hexane:EtOAc) showed no remaining **277**. The mixture was diluted into water and
 extracted with EtOAc. The organic was separated, washed with brine, dried, filtered and
 concentrated in *vacuo* to afford 0.505 g of **278** as an orange oil which was used without
 purification. Yield 92 %

20 Part D.

To a solution of **278** (0.505 g, 2.02 mmol, 1 eq) in MeOH-THF (3:1, 5 mL) was
 added an aqueous solution of hydroxylamine (0.169 g, 2.43 mmol, 1.2 eq in 2.5 mL of
 water) in one portion. The pH was adjusted to 9 with NaOH (6N), and stirred at rt for 1 hr
 25 where TLC (2:1 Hexane:EtOAc) showed complete consumption of **278**. NaBH_3CN (0.254
 g, 2.43 mmol, 2 eq) was added with a fleck of methyl red and the solution acidified to pH 2-
 3 using HCl in MeOH (20 V/V). The pH of the reaction solution was maintained at pH 3
 over the course of 12 h by addition of small amounts of the methanolic HCl solution, where


it was basified to pH 9 with NaOH (2 N). The solution was extracted with EtOAc, washed with brine, dried, filtered and concentrated in *vacuo* to afford an orange oil. The crude material was purified by column chromatography (silica gel, 55 % EtOAc in Hexane) to afford 0.486 g of **279** as a cream solid. Yield 90%.

5

Part E.

To a solution of **279** (0.486 g, 1.83 mmol, 1 eq) and glyoxylate ester (0.285g, 2.37 mmol 1.3 eq) in diethyl ether (10 mL) was added anhydrous CaCl₂ (0.263 g, 2.37 mmol, 1.3 eq), left stirring at rt for 3 h, resulting in a suspension. The suspension was filtered through a Celite plug washing with DCM and ether. The resultant yellow solution was concentrated in *vacuo* to afford 0.497 g of **280** as yellow oil and used directly without purification. MS (ESI(+)) *m/z* 337.8 (M+H)⁺ Yield 81 %.


15

Part F.

To a solution of **280** (0.497 g, 1.48 mmol, 1 eq) in anhydrous THF (10 mL) was added (S,Z)-5-tert-butyldimethylsiloxy)pent-3-en-2-ol **6** (0.5 mL, 1.77 mmol, 1.2 eq) and Ti(O*i*Pr)₄ (0.65 mL, 2.22 mmol, 1.5 eq) heated in the microwave at 140 C for 15 min. TLC (30:1 DCM:Et₂O) showed consumption of nitron **280** and the allylic alcohol **6**. 3-(dimethylamino)-1,2-propanediol (1 mL) in EtOAc (1 mL) was added and the dark brown solution left stirring at rt overnight. The dark brown solution was diluted with EtOAc and quenched with Rochelle's salt solution, washed with water, brine, dried and concentrated in *vacuo* to afford a brown oil. The crude material was purified by column chromatography (silica gel, 10 % EtOAc in Hexane) to afford 0.490 g of **281** as a brown oil. MS (ESI(+)) *m/z* 521.9 (M+H)⁺ Yield 64 %


25

Part G.




To a solution of **281** (0.470 g, 0.903 mmol, 1 eq) in THF (5 mL) was added concentrated 6 N HCl (0.45 mL, 2.71 mmol, 3 eq) and stirred at rt for 2 h. TLC (1:2 hexane:EtOAc) of neutralized aliquot showed complete consumption of **281**. The reaction was neutralized with NaHCO₃, extracted with EtOAc, washed with brine, dried, filtered and concentrated in *vacuo* to afford a brown oil. The crude material was purified by column chromatography (silica gel, 50 % EtOAc in Hexane) to afford 0.251 g of **282** as a brown solid. MS (ESI(+)) *m/z* 407.7 (M+H)⁺ Yield 70 %

10 Part H.




To a solution of (+)-isocamphenylamine (0.291 g, 1.897 mmol, 3 eq) flushed with Ar in anhydrous DCM (3 mL), was added Me₃Al (2 M solution in Hexane, 0.632 mL, 1.26 mmol, 2 eq) dropwise over 20 min and the resultant clear solution stirred for 40 min at rt. A solution of the lactone **282** (0.264 g, 0.632 mmol, 1 eq) in DCM (3 mL) was added by cannula flask slowly, turning the reaction yellow with gas evolution. The reaction mixture was stirred at rt for 12 h TLC (2:1 EtOAc:Hexane) showed a clean reaction with one marginally faster running spot. The reaction was diluted with DCM and quenched on dropwise addition to a rapidly stirring solution Rochelle's salt, and stirred at rt for 2 h. The organic layer was separated and washed with water, brine, dried, and concentrated in *vacuo* to afford a yellow gun. The crude material was purified by column chromatography (silica gel, 50 % EtOAc in Hexane) to afford 0.237 g of **283** as a yellow oil. MS (ESI(+)) *m/z* 560.8 (M+H)⁺ Yield 67 %

25 Part I.




To a solution of isoxazolidine **283** (100 mg, 0.179 mmol, 1 eq) in DMSO (5 mL), flushed with argon, was added the pinacolboronate (68 mg, 0.232 mmol, 1.3 eq), potassium acetate (26 mg, 0.268 mmol, 1.5 eq) and cesium carbonate (175 mg, 0.526 mmol, 3 eq). Upon stirring for 10 min, Pd(dppf)Cl₂ (29 mg, 0.036 mmol, 0.2 eq) was added as a single portion. The mixture was heated to 70 °C for 4 h, then allowed to cool to rt. EtOAc and brine were added and the pH adjusted to 3-4 with HCl (2N). The organics were separated, washed with water, brine, dried, and concentrated in *vacuo* to afford a brown black oil. The crude material was purified by column chromatography (silica gel, 4% MeOH in DCM) to afford 35 mg of **284** as a brown solid. MS (ESI(+)) *m/z* 644.0 (M+H)⁺ Yield 30 %

Part I.




A solution of **284** (35 mg, 0.055 mmol, 1 eq) in DCM (3 mL) was added Et₃N (23 µL, 0.165 mmol, 3 eq) and HBTU (42 mg, 0.110 mmol, 2 eq) and allowed to stir at rt for 10 min. To this solution was added (S)-N1,N1-dimethyl-3-phenylpropane-1,2-diamine (9.8 mg, 0.055 mmol, 1 eq) and allowed to stir at rt for 2 h. The reaction was taken up in DCM, washed with K₂CO₃, water, brine, dried and concentrated in *vacuo* to afford a brown solid. The crude material was diluted with MeOH (2 mL) and purified by reverse-phase HPLC (MeCN/water with 40 mM NH₄HCO₃) to afford 13 mg of **274** as a white solid. MS (ESI(+)) *m/z* 804.2 (M+H)⁺ Yield 30 %

Example 105




Part A.




3-amino-5-boronobenzoic acid (1g, 5.5mmol) was dissolved in 25ml MeOH and 412uL of a 37% solution of formaldehyde was added (0.166mg, 5.5mmol, 1eq). Sodium cyanoborohydride (347mg, 5.5mmol, 1eq) was added and the reaction stirred for 30 minutes at room temperature. The solution was dried via rotary evaporation and the resulting material was purified over silica gel (5-10% MeOH in DCM containing 0.2% acetic acid) to obtain **285** (333mg, 31% yield). The monoalkylated product was carried on to a second reductive alkylation.

Part B.




15 Boronic acid **285** (102mg, 0.5mmol) 2mL MeOH and 100uL of **3** (92mg, 0.5mmol, 1eq) was added. Sodium cyanoborohydride (49mg, 0.75mmol, 1.5eq) was added and the reaction stirred for 90 minutes at room temperature. The solution was dried via rotary evaporation and the resulting material was purified over silica gel (5-10% MeOH in DCM containing 0.2% acetic acid) to obtain **286** (333mg) 31% Yield.

Part C.




Boronic acid **286** (162mg, 0.280mmol) was dissolved in DMSO (7mL) and potassium acetate (28mg, 0.280mmol, 1eq) and cesium carbonate (277mg, 0.850mmol, 3eq) was added. Iodide **9** (100mg, 0.280mmol, 1eq) was added and the stirred solution was degassed by subjecting the flask to three 60 second pulses of alternating vacuum and argon purge cycles. Palladium catalyst **287** (46mg, 0.057mmol, 0.2eq) was added and the solution heated to 60 degrees for 2 hours. The reaction was acidified to pH and extracted with DCM. Organic extracts were combined and dried over sodium sulfate and concentrated to dryness. The crude product was purified by flash (5% MeOH in DCM) to afford **288** (94mg) 44% Yield.

Part D.




Biphenyl acid **288** (50mg, 0.066mmol) was dissolved in DMF (0.5mL) and combined with (S)-N1,N1-dimethyl-3-phenylpropane-1,2-diamine (13mg, 0.073mmol, 1.1eq). Triethylamine (27uL, 0.2mmol, 3eq) was added followed by HBTU (25mg, 0.066mmol, 1eq). After 4 hours the crude reaction was purified by flash (5%MeOH in DCM, containing 0.2% NH₄OH) to yield **289** (42mg). 69% Yield.

Part E.




TBS ether **289** (42mg, 0.046mmol) was dissolved in THF (1mL) and HF-Pyridine added (30uL, 0.092mmol, 2eq). After 6 hours of stirring, the crude material was purified by flash (5%MeOH in DCM, containing 0.2% NH₄OH) to yield **285** (4.5mg). 12% Yield. MS (ESI(+)) m/e 800.5 (M+H)⁺.

Example 106




Part A.





Biphenyl acid **288** (25mg, 0.033mmol) was dissolved in DMF (0.5mL) and combined with amine **150** (10mg, 0.033mmol, 1eq). Triethylamine (14uL, 0.099mmol, 3eq) was added followed by HBTU (13mg, 0.033mmol, 1eq). After 4 hours the crude reaction was purified by flash (5%-10%MeOH in DCM) to yield **291** (9mg). 27% Yield.

Part B.



TBS ether **291** (9mg, 0.009mmol) was dissolved in THF (0.4mL) and HF-Pyridine added (2.7uL, 0.022mmol, 2.5eq). After 12 hours of stirring, the crude material was purified by flash (5%MeOH in DCM, containing 0.2% NH₄OH) to yield **289** (2mg). 25% Yield. MS (ESI(+)) m/e 893.5 (M+H)⁺.

5


Example 107**Part A.**

10

To a solution of iodo-core **9** (1.00 g, 1.74 mmol, 1.0 eq) in DMSO (20 mL) under Ar was added cesium carbonate (1.71 g, 5.24 mmol, 3.0 eq), potassium acetate (171 mg, 1.74 mmol, 1.0 eq) and 3,5-diformylphenylboronic acid (622 mg, 3.49 mmol, 2.0 eq). The solution was degassed by bubbling Ar through it for 5 min. PdCl₂(dppf).CH₂Cl₂ adduct (285 mg, 0.35 mmol, 0.2 eq) was then added and the resulting mixture stirred at 70 °C for 1h. The reaction was allowed to cool to rt, quenched with brine (20 mL) and diluted with AcOEt (20 mL). The layers were separated and the organic one was washed with brine (5 x 10 mL), dried (MgSO₄), filtered and concentrated in vacuo. The residue was purified by flash chromatography (Hex/AcOEt 3:7, 1:4 and 0:1). The product **293** (669 mg, 1.16 mmol, 66% yield) was obtained as a foam.

15

20


Part B.

25

To a solution of **293** (650 mg, 1.12 mmol, 1.0 eq) in THF (12 mL) was added 2 M sodium hydrogenphosphate solution (842 ul, 1.69 mmol, 1.5 eq), isobutylene (788 ul, 11.2

mmol, 10.0 eq) and 1 M sodium chlorite solution (1.35 mL, 1.35 mmol, 1.2 eq). The reaction mixture was stirred at rt for 50 min, quenched with 10% Na₂S₂O₃ (10 mL) and stirred for 15 additional min, diluted with AcOEt (10 mL), the layers separated and the aqueous one extracted with AcOEt (2 x 10 mL). The comb org extracts were washed with brine, dried (MgSO₄), filtered and concentrated in vacuo. The residue was purified by flash chromatography (DCM/MeOH 98:2 and 9:1). **294** was obtained as an off-white foam (204 mg, 0.34 mmol, 31% yield).

Part C.




10

To a stirred solution of acid **294** (200 mg, 0.34 mmol, 1.0 eq) in dry DCM (1 mL) under Ar was added (S)-N1,N1-dimethyl-3-phenylpropane-1,2-diamine (120 mg, 0.67 mmol, 2.0 eq), Hunig's base (117 µL, 0.67 mmol, 2.0 eq) and HBTU (191 mg, 0.50 mmol, 1.5 eq). After 1h, the reaction was quenched with water (5 mL), the layers separated and the aqueous one extracted with DCM (2 x 5 mL). The combined organic extracts were washed with brine (10 mL), dried (MgSO₄), filtered and concentrated in vacuo. The residue was purified by flash chromatography (DCM/MeOH 9:1). Amide **295** (100 mg, 0.13 mmol, 39% yield) was obtained as a foam.

15

20 Part D.





25


To a solution of aldehyde **295** (42 mg, 0.06 mmol, 1.0 eq) in THF (1.5 mL) and MeOH (0.45 mL) was added 0.5 M Ammonia in dioxane (2.23 mL, 1.11 mmol, 20.0 eq) followed by AcOH (32 µL, 0.56 mmol, 10.0 eq) and sodium cyanoborohydride (17 mg, 0.28 mmol, 5.0 eq). The resulting mixture was stirred at rt for 1 h, quenched with saturated sodium bicarbonate solution (10 mL) and extracted with AcOEt (3 x 10 mL). The combined organic extracts were washed with brine (20 mL), dried (MgSO₄), filtered and concentrated in vacuo. A white foam was obtained, which was purified by preparative

HPLC to render **295-amine** (10 mg, 0.013 mmol, 24% yield) as a lyophilized powder. MS (ESI(+)) m/e 756.46 (M+H)⁺.

5

Example 108**Part A.**

10 In a sealed reactor, 3,5-dibromobenzoic acid (1.0g, 3.57mmol, 1eq), N,N-dimethylethanolamine (735 μ l, 7.15mmol, 2eq), potassium triphosphate 1.517g, 7.15mmol, 2eq) and copper(I) iodide (68mg, 0.357mmol, 0.1eq) were dissolved in water (10mL). The solution was heated at 90°C for 16h then allowed to cool to rt and poured on HCl 1 N (100mL). The mixture was brought to pH~4 using NaOH 1 N and HCl 1 N then
 15 extracted with EtOAc (3x30mL). The combined organics were washed with brine, dried over Na₂SO₄, filtered, concentrated. The residue was purified by flash chromatography (DCM and DCM/MeOH/AcOH 90:10:1 0 to 50%) to obtain **297** (318mg, 1.06mmol) as an oil. Yield 30%.


Part B.

20

A flask containing boronate (described in Example 1, part G) (194mg, 396 μ mol, 1eq), bromide **297** (142mg, 475 μ mol, 1.2eq), cesium carbonate (387mg, 1.18mmol, 3eq), potassium acetate (39mg, 396 μ mol, 1eq) and Pd(dppf)Cl₂ (58mg, 79 μ mol, 0.2eq) was

purged with argon and DMSO (2mL) was added. The reaction was heated at 60°C for 5h then allowed to cool to rt, diluted with water (100mL) and acidified to pH~4 with HCl 0.02M. The mixture was extracted with DCM (3x30 mL). The combined organic layers were washed with water (1x30mL) and brine (1x30mL), dried over Na₂SO₄, filtered and concentrated to a brown oil which was purified by flash chromatography (DCM and DCM/MeOH/AcOH 90:10:1 0 to 70%) to obtain **298** (155mg, 233μmol) as an oil. Yield 59%.


Part C.




Acid **298** (43mg, 65μmol, 1eq), (S)-N1,N1-dimethyl-3-phenylpropane-1,2-diamine (14mg, 77μmol, 1.2eq) and HBTU (29mg, 77μmol, 1.2eq) were dispersed in DCM (1mL) and triethylamine (27μL, 194μmol, 3eq) was added. The reaction was stirred at rt for 1 h then concentrated to dryness. The residue was purified by HPLC to afford **296** (3.9mg, 4.7μmol) as a lyophilized powder. MS (ESI(+)) m/e 826.47 (M+H)⁺. Yield 7.3%.

15

Example 109



Part A.




To a solution of **295** (100 mg, 0.13 mmol, 1.0 eq) in THF (2 mL) was added 2M sodium hydrogenphosphate solution (99 ul, 0.20 mmol, 1.5 eq), isobutylene (93 ul, 1.33 mmol, 10.0 eq) and 1M sodium chlorite solution (159 ul, 0.16 mmol, 1.2 eq). The reaction mixture was stirred at rt for 1h, an equal amount of reagents were added and stirring

25

continued for 30 additional min. The reaction was quenched with 10% $\text{Na}_2\text{S}_2\text{O}_3$ (5 mL) and stirred for 15 min, diluted with AcOEt (5 mL), the layers separated and the aqueous one extracted with AcOEt (2 x 5 mL). The combined organic extracts were washed with brine, dried (MgSO_4), filtered and concentrated in vacuo. The residue **300** was used without
 5 further purification.



Part B.

To a stirred solution of crude acid **300** (67 mg, 0.09 mmol, 1.0 eq) in dry DCM (1 mL) under Ar was added *N,O*-dimethyl hydroxylamine hydrochloride **5** (21 mg, 0.22 mmol, 2.5 eq), Hunig's base (56 μL , 0.44 mmol, 5.0 eq) and HBTU (49 mg, 0.13 mmol, 1.5 eq). After 1h, the reaction was quenched with water (5 mL), the layers separated and the aqueous one extracted with DCM (2 x 5 mL). The combined organic extracts were washed with brine (10 mL), dried (MgSO_4), filtered and concentrated in vacuo. The residue **301**,
 15 was used directly in the next reaction.


Part C.

Crude Weinreb amide **301** (35 mg, 0.04 mmol, 1.0 eq) was dissolved in anhydrous THF (1 mL) under Ar and was cooled to -78°C . 1.6 M Methyllithium in Et_2O (54 μL , 0.09 mmol, 2.0 eq) was added and the reaction mixture was stirred for 30 min at -78°C and allowed to warm to rt. The reaction was quenched with saturated NH_4Cl solution (5 mL) and diluted with Et_2O (5 mL), the layers separated, the aqueous one extracted with Et_2O (2
 25 x 5 mL) and the combined organic extracts washed with brine (10 mL), dried (MgSO_4), filtered and concentrated in vacuo. A foam was obtained, which was purified by preparative HPLC to render **299** (6 mg, 0.01 mmol, 18% yield) as a lyophilized powder. MS (ESI(+)) m/e 769.45 ($\text{M}+\text{H}$) $^+$.

Example 110

Part A.


A solution of aryl iodide **215** (113 mg, 0.19 mmol, 1 eq), boronic acid **286** (80 mg, 0.23 mmol, 1.2 eq), cesium carbonate (185 g, 0.57 mmol, 3 eq) and potassium acetate (18.6 mg, 0.19 mmol, 1 eq) in DMSO (4.8 mL) was degassed by bubbling argon through the solution for 10 min. Pd(dppf)Cl₂ (27.7 mg, 0.04 mmol, 0.2 eq) was then added and the flask purged with argon. The mixture was heated at 70 °C for 1 h. The reaction mixture was added to brine (10 mL), acidified with 1 N HCl until the aqueous layer attained a pH of 3, and extracted with EtOAc (2 x 10 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated *in vacuo* to afford a brown oil. The oil was purified by silica gel chromatography (40-60% EtOAc/hexane with 0.25% acetic acid) to give 17.3 mg of **303** as a brown oil. Yield 12%. MS (ESI(+)) *m/z* 779.53 M⁺.

Part B.

To a solution of **303** (17.3 mg, 0.022 mmol, 1 eq) in DCM (2 mL) was added TEA (10 μL, 0.067 mmol, 3 eq) and HBTU (16.8 mg, 0.044 mmol, 2 eq). The solution was stirred at rt for 10 min. (S)-N1,N1-dimethyl-3-phenylpropane-1,2-diamine (6 mg, 0.033 mmol, 1.5 eq) was then added. After stirring at rt for 2h, the mixture was diluted with DCM (10 mL) and washed with saturated potassium carbonate solution (10 mL). The aqueous layer was extracted with DCM (2 x 10 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated *in vacuo*. The material was purified by silica


gel chromatography (60-75% EtOAc/hexane with 0.25% TEA) to give 15.1 mg of **304**. Yield 72%. MS (ESI(+)) m/z 939.68 M^+ .

Part C.




To a solution of **304** (15.1 mg, 0.016 mmol, 1 eq) in THF (1.5 mL) in a polypropylene tube at rt was added a 1:1:1 THF:HF/pyridine:pyridine solution (70 μ L). The solution was stirred at rt for 1 h, quenched with methoxytrimethylsilane (700 μ L) and concentrated *in vacuo*. The crude **305** was used in the next step without further purification. MS (ESI(+)) *m/z* 825.44 M^+ .

Part D.




To a solution of **305** (14.3 mg, 0.017 mmol, 1 eq) in DMF (0.6 mL) at 0 °C was added dithiothreitol (8 mg, 0.052 mmol, 3 eq) and DBU (7.8 μ L, 0.052 mmol, 3 eq). The solution was stirred at 0 °C for 20 min. The crude reaction was kept at 0 °C and purified by HPLC to give 0.8 mg of **302**. Yield 6%. MS (ESI(+)) m/z 799.55 M^+ .

Example 111




20

Part A.




A solution of aryl iodide **9** (403.8 mg, 0.71 mmol, 1 eq), boronic acid **285** (151 mg, 0.78 mmol, 1.1 eq), cesium carbonate (689 g, 2.2 mmol, 3 eq) and potassium acetate (69.2 mg, 0.71 mmol, 1 eq) in DMSO (7.4 mL) was degassed by bubbling argon through the solution for 10 min. Pd(dppf)Cl₂ (103 mg, 0.14 mmol, 0.2 eq) was then added and the flask
 5 purged with argon. The mixture was heated at 70 °C for 1.5 h. The reaction mixture was added to brine (40 mL), acidified with 1 N HCl until the aqueous layer attained a pH of 3, and extracted with EtOAc (2 x 40 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated *in vacuo* to afford a brown oil. The oil was purified by silica gel chromatography (75-95% EtOAc/hexane with 0.25% acetic acid) to give 124.3
 10 mg of **306**. Yield 29.6%. MS (ESI(+)) *m/z* 596.34 (M+H)⁺.

Part B.



To a solution of **306** (25 mg, 0.04 mmol, 1 eq) in DCM (4 mL) was added TEA (17
 15 μL, 0.12 mmol, 3 eq), HBTU (31.8 mg, 0.08 mmol, 2 eq) and (S)-N1,N1-dimethyl-3-phenylpropane-1,2-diamine (15 mg, 0.08 mmol, 2 eq). After stirring at rt for 2h, the mixture was diluted with DCM (15 mL) and washed with saturated potassium carbonate solution (10 mL). The aqueous layer was extracted with DCM (2 x 10 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated *in vacuo*. The material
 20 was purified by silica gel chromatography (60-75% EtOAc/DCM with 0.25% TEA) to give 22.8 mg of **307**. Yield 72%. MS (ESI(+)) *m/z* 756.45 M⁺.


Part C.




To a solution of **307** (6.7 mg, 0.009 mmol, 1 eq) in MeOH (0.5 mL) at rt was added glyoxylic acid monohydrate (1 mg, 0.011 mmol, 1.2 eq). The solution was stirred at rt for

10 min. Sodium cyanoborohydride (0.7 mg, 0.011 mmol, 1.2 eq) was then added. The solution was stirred at rt for 1 h and concentrated *in vacuo*. The crude reaction was purified by HPLC to give 5 mg of **306**. Yield 69%. MS (ESI(+)) m/z 814.53 M^+ .

Example 112




Part A.




10 A solution of **306** (5.5 mg, 0.007 mmol, 1 eq), dimethylamine hydrochloride (1.7 mg, 0.02 mmol, 3 eq), HBTU (5.2 mg, 0.014 mmol, 2 eq) and TEA (4 μL , 0.028 mmol, 4 eq) in DCM (0.5 mL) was stirred at rt for 2h. The crude reaction was concentrated *in vacuo* and purified by HPLC to give 4.4 mg of **308**. Yield 77%. MS (ESI(+)) m/z 841.63 M^+ .

Example 113




Part A.



20 A solution of **306** (14 mg, 0.017 mmol, 1 eq), methylamine (26 μL , 2M in THF, 0.052 mmol, 3 eq), HBTU (13 mg, 0.034 mmol, 2 eq) and TEA (10 μL , 0.069 mmol, 4 eq)

in DCM (1 mL) was stirred at rt for 2h. The crude reaction was concentrated *in vacuo* and purified by HPLC to give 3.3 mg of **309**. Yield 23%. MS (ESI(+)) m/z 827.60 M^+ .


Example 114



310

5

Part A.




308

309

A solution of **306** (13.3 mg, 0.016 mmol, 1 eq), ammonia (7 μ L, 7N in MeOH, 0.049 mmol, 3 eq), HBTU (12.4 mg, 0.033 mmol, 2 eq) and TEA (9 μ L, 0.065 mmol, 4 eq) in DCM (1 mL) was stirred at rt for 2h. The crude reaction was concentrated *in vacuo* and purified by HPLC to give 6 mg of **310**. Yield 45%. MS (ESI(+)) m/z 813.58 M^+ .

10


Example 115



311

15

Part A.



10


311

A solution of **10** (10 mg, 0.02 mmol, 1 eq) in DMF (1 mL) was treated with TEA (9 μ L, 0.07 mmol, 3.0 eq), (S)-N1,N1,N2-trimethyl-3-phenylpropane-1,2-diamine (9 mg, 0.06 mmol, 3.0 eq) (derived from the requisite amino acid analogous to the procedure described

20


in Example 73, part A), and HBTU (10 mg, 0.04 mmol, 2.0 eq). After stirring for 2 h at rt, the reaction mixture was diluted with MeOH (500 μ L) and directly purified by reverse-phase HPLC (MeCN/water with 40 mM NH_4CO_2) to give 5 mg of compound **311** as a white solid. Yield 50%. MS (ESI(+)) m/z 784.4 ($\text{M}+\text{H}$)⁺.

5

Example 116**312****Part A.**


Compound **312** was synthesized according to the procedure described in Example 1, part H, using commercially available (*R*)-2-aminomethyl-1-ethylpyrrolidine in place of (*S*)-*N*¹,*N*¹,4,4-tetramethylpentane-1,2-diamine. Yield 20%. MS (ESI(+)) m/z 720.4 ($\text{M}+\text{H}$)⁺.

10

Example 117**313**15 **Part A.**


Compound **313** was synthesized according to the procedure described in Example 1, part H using (*R*)-1-Benzyl-2-(4-methyl-piperazin-1-yl)-ethylamine (derived from the requisite amino acid analogous to the procedure described in Example 73, part A) in place of (*S*)-*N*¹,*N*¹,4,4-tetramethylpentane-1,2-diamine. Yield 41%. MS (ESI(+)) m/z 825.5 ($\text{M}+\text{H}$)⁺.

20

Example 118**314****Part A.**

Compound **314** was synthesized according to the procedure described in Example 1, part H using (S)-3-Methyl-1-(4-methyl-piperazin-1-ylmethyl)-butylamine (derived from the requisite amino acid analogous to the procedure described in Example 73, part A) in place of amine (S)-*N*¹,*N*¹,4,4-tetramethylpentane-1,2-diamine. Yield 28%. MS (ESI(+)) *m/z* 791.5 (M+H)⁺.

Example 119




315

Part A.


Compound **315** was synthesized according to the procedure described in Example 1, part H using (S)-3-Methyl-1-morpholin-4-ylmethylamine (derived from the requisite amino acid analogous to the procedure described in Example 73, part A) in place of (S)-*N*¹,*N*¹,4,4-tetramethylpentane-1,2-diamine. Yield 25%. MS (ESI(+)) *m/z* 778.6 (M+H)⁺.

Example 120




316

Part A.




Boc-Leucinol (2.20g, 10.1mmol, 1eq), phthalimide (1.79g, 12.2mmol, 1.2eq) and triphenylphosphine (3.98g, 15.2mmol, 1.5eq) were dissolved in dry THF (40mL) under Ar and diisopropyldiazocarbonylate (3.17mL, 15.2mmol, 1.5eq) was added dropwise. The solution was stirred at rt for 1h30 then concentrated to dryness. The residue was purified by flash chromatography (Hexane then Hex/EtOAc 9:1) to afford **317** (2.99g, 8.6mmol) as a white solid. Yield 85%.


Part B.

Compound **317** (1.0g, 2.89mmol, 1eq) was treated with hydrazine hydrate (0.896mL, 28.9mmol, 10eq) in refluxing ethanol for 2h. The reaction was allowed to reach rt then filtered on paper. The filtrate was concentrated to dryness and the residue taken in DCM. The mixture was filtered on paper and the filtrate concentrated to dryness.

The residue was taken in DCM/THF 1:1 (40mL) and HBTU (1.2g, 3.1mmol, 1.1eq) then triethylamine (0.43mL, 3.1mmol, 1.1eq) were added. The solution was stirred at rt for 3h30, then concentrated. The residue was taken in EtOAc (100mL) and the solution washed with NaOH 1 N (3x20mL), dried over Na₂SO₄, filtered, concentrated to dryness to obtain **318** (1.051g, 85% pure) as an oil. Yield 99%.


Part C.

Compound **318** (1.051g, 3.34mmol, 1eq) was treated with HCl 4M in dioxane for 1 h then concentrated to dryness. The residue was precipitated from methanol/diethyl ether and dried under vacuum to obtain in a quantitative yield the bis-hydrochloric salt of **319** as an off-white foam (800mg, 2.89mmol).

Part D.

Compound **316** was synthesized according to the procedure described in Example 1, part H using amine **319** in place of (S)-N¹,N¹,4,4-tetramethylpentane-1,2-diamine. Yield 32%. MS (ESI(+)) *m/z* 806.6 (M+H)⁺.


Example 121




320

Part A.

Compound **320** was synthesized according to the procedure described in Example 1, part H using (*S*)-3-Methyl-1-pyrrolidin-1-ylmethyl-butylamine (derived from the requisite amino acid analogous to the procedure described in Example 73, part A) in place of (*S*)-*N*¹,*N*¹,4,4-tetramethylpentane-1,2-diamine. Yield 22%. MS (ESI(+)) *m/z* 762.6 (M+H)⁺.

Example 122


321

Part A.

322


3-Bromo-5-chloro-2-hydroxybenzaldehyde (2 g, 8.5 mmol, 1 eq) was dissolved in DMF (10 mL) and treated with potassium carbonate (1.76 g, 12.7 mmol, 1.5 eq) and iodomethane (1.06 mL, 17 mmol, 2 eq). The mixture was stirred at 50 °C for 14h. The mixture was poured into water (100 mL) and extracted with ether (2 x 75 mL). The ether layers were washed successively with 15% aqueous NaOH, water, and then brine (50 mL each). Drying over magnesium sulfate and concentration *in vacuo* gave the ether **322** as a white solid (2.07 g, 98% yield).

Part B.



Aldehyde **322** (2.07 g, 8.38 mmol, 1 eq) and hydroxylamine hydrochloride (0.757 g, 10.9 mmol, 1.3 eq) were dissolved in THF/MeOH (1:3, 30 mL) at 23 °C. The pH was adjusted to 9 with 6 N KOH. After stirring for 2 h, NaBH₃CN (0.526 g, 8.4 mmol, 1 eq) was added followed by a crystal of methyl orange. The pH was adjusted to 2 and the resulting ruby red color of the reaction mixture was maintained for the duration of the reaction by the frequent addition of 1 N HCl. After stirring for a total of 16 h, the reaction mixture was adjusted to pH 7 with 6 N KOH. The mixture was diluted with water (100 mL) and extracted with DCM (2 x 50 mL) and the combined organics were washed with water (50 mL), brine (20 mL), dried over MgSO₄, filtered and concentrated *in vacuo* to afford 2.0 g (90% yield) of hydroxylamine **323**. The crude material was used without further purification.


Part C.



A 0 °C dichloromethane solution (120 mL) of (S)-allylic alcohol (5.0 g, 23 mmol, 1 eq) with pyridine (4 mL, 46 mmol, 2 eq) was treated with bromoacetyl chloride (2 mL, 30 mmol, 1.3 eq). After 20 min, the mixture was stirred for 1 h at 23 °C and then poured into 1 N HCl (200 mL). The layers were separated and the organic layer washed successively with water, 5% aqueous NaHCO₃, and brine (50 mL each) and dried over sodium sulfate. Concentration *in vacuo* gave a light amber oil (6.5 g, 83% yield). This oil was restored in acetone (40 mL) with sodium iodide (3.0 g, 19 mmol, 1.0 equiv) and stirred at 23 °C for 6 h. The mixture was then diluted with ethyl acetate (100 mL) and washed successively with water, 10% aqueous sodium sulfite, and brine (50 mL each), and the organic layer was dried over sodium sulfate and dried *in vacuo* to give a light amber oil. This oil was restore in dry acetonitrile (50 mL) and treated with silver nitrate (5.0 g, 29 mmol, 1.5 eq) and stirred 14 h at 23 °C. The mixture was poured into water (200 mL) and extracted twice with

ether (2 x 100 mL). The ether layers were washed with water (200 mL) and brine (100 mL), dried over MgSO_4 and concentrated *in vacuo* to a clear oil. A portion of this oil (500 mg, 1.56 mmol, 1 eq) was dissolved in dry DMSO (6 mL) and treated with sodium acetate (193 mg, 2.35 mmol, 1.5 eq) with stirring at 23 °C. After 25 min, the mixture was poured into ice water (40 mL), extracted with ether (3 x 30 mL), and the ether layers were washed successively with saturated aqueous sodium bicarbonate, water, and brine and dried over sodium sulfate. Concentration *in vacuo* gave 324 as a clear oil (270 mg, 60% yield).

Part D.



Hydroxylamine 323 (350.0 mg, 1.3 mmol, 1 eq) and allylic glyoxylate ester 324 (393 mg, 1.44 mmol, 1.1 eq) were dissolved in toluene (20 mL) and heated in at 80 °C for 2 h. The reaction mixture was concentrated *in vacuo*. The crude residue was purified by silica gel chromatography (10-50% EtOAc/hexane) to afford 420 mg of lactone 325 (61% yield).


Part E.



To a 0 °C solution of TBS-protected lactone 325 (410 mg, 0.8 mmol, 1 eq) in THF (4 mL) was added 6 N HCl (0.3 mL). The reaction mixture was stirred for 4 h and then quenched by the addition of a 5% aqueous sodium bicarbonate solution (30 mL). The reaction mixture was extracted with dichloromethane (2 x 40 mL) and the combined organics were washed with brine (30 mL), dried over MgSO_4 , filtered and concentrated *in vacuo*. The crude residue of alcohol was restored in dry dichloromethane (5 mL). A 0 °C solution of (+)-isopinocampheylamine (0.4 mL, 2.36 mmol, 3 eq) in dichloromethane (5 mL) was treated with trimethylaluminum (1.2 mL, 2 M in toluene, 2.36 mmol, 3 eq) and the reaction mixture was stirred for 30 min. The solution of crude lactone alcohol (was then


added dropwise. After stirring for 14 h at 23 °C, dichloromethane (50 mL) and sodium sulfate-dodecahydrate (1.0 g) was added, and the mixture stirred for 3 h. Filtration and concentrated of the filtrate *in vacuo* afforded an oil that was purified by silica gel chromatography (30-70% EtOAc/Hexane) to afford 335 mg of amide **326** (76% yield).


5

Part F.

Amide **326** (25 mg, 0.045 mmol, 1 eq), 3-(dimethylamino)-5-
 10 (phenethylcarbamoyl)phenylboronic acid (17 mg, 0.054 mmol, 1.2 eq), Pd(OAc)₂ (1.0 mg, 0.0045 mmol, 0.1 eq), potassium carbonate (25 mg, 0.18 mmol, 4 eq), and S-Phos (Kevin W. Anderson and Stephen L. Buchwald, *Angew. Chem. Int. Ed.* 2005, 44, 2–6) (4.7 mg, 0.009 mmol, 0.2 eq) were dissolved in tetrahydrofuran (1 mL) and water (0.5 mL). The mixture was purged with argon and stirred at 50 °C for 5 h. The reaction mixture was
 15 diluted with EtOAc (20 mL) and water (10 mL) and layers separated. The organic layer was washed with brine (10 mL), dried over sodium sulfate, and concentrated *in vacuo*. Purification of the crude residue by HPLC (acetonitrile/30 mM aqueous ammonium bicarbonate) to afford 9 mg of biphenyl **321** (30 % yield) as a white foam. MS (ESI(+)) *m/e* 747.5 (M+H⁺).


20

Example 123Part A.




To a solution of aminoacetaldehyde dimethyl acetal (3.5 g, 33.3 mmol, 1 eq) in DCM (8 ml) under Ar at 0 °C was added triethylamine (9.28 mL, 66.6 mmol, 2 eq) followed by methanesulfonyl chloride (3.89 mL, 49.9 mmol, 1.5 eq). The resulting mixture was stirred at 0 °C for 15 min. The reaction was diluted with DCM (20 mL) and washed with saturated sodium bicarbonate solution (20 mL). The aqueous layer was extracted with DCM (2 x 20 mL). The combined organic layers were dried over MgSO₄, filtered and concentrated *in vacuo*. Compound **328** was obtained as a yellow oil, which was used without further purification.

Part B.



To a solution of **328** (3.0 g, 16.4 mmol, 1 eq) in acetone: water (1:1, 4 ml) was added Amberlyst-15 resins (3.0 g). The resulting mixture was stirred for 3 h. The suspension was filtered through celite and concentrated *in vacuo*. Compound **330** was obtained as a yellow oil and was used without further purification.

Part C.




To a solution of the boronic acid (90 mg, 0.462 mmol, 1 eq) in THF (2 ml) was added **330** (158 mg, 1.154 mmol, 2.5 eq) in MeOH (1 mL), followed by acetic acid (52.8 μL, 0.923 mmol, 2.0 eq) and sodium cyanoborohydride (145 mg, 2.308 mmol, 5.0 eq). The resulting mixture was stirred for 30 min. The reaction mixture was diluted with EtOAc (10 mL) and quenched with brine. The aqueous layer was acidified with 1N HCl to pH 3, and extracted with EtOAc (3 x 20 mL). The combined organic layers were dried over MgSO₄, filtered and concentrated *in vacuo* to afford yellow oil. The oil was purified by silica gel

chromatography (2 – 7 % MeOH/DCM with 0.5 % acetic acid) to give 94 mg of **331**. Yield 64.4%. MS (ESI(+)) m/z 317.24 (M+H)⁺.

Part D.


5



A solution of aryl iodide (140 mg, 0.245 mmol, 1 eq), **331** (85 mg, 0.269 mmol, 1.1 eq), cesium carbonate (159 mg, 0.489 mmol, 2 eq) and potassium acetate (24 mg, 0.245 mmol, 1 eq) in DMSO (3 mL) was degassed by bubbling argon through the solution for 10 min. Pd(dppf)Cl₂ (35.8 mg, 0.049 mmol, 0.2 eq) was then added and the flask purged with argon. The mixture was heated at 70 °C for 4 h. The reaction mixture was added to brine (10 mL), acidified with 1 N HCl until the aqueous layer attained a pH of 3, and extracted with EtOAc (3 x 20 mL). The combined organic layers were dried over MgSO₄, filtered and concentrated *in vacuo* to afford a brown oil. The oil was purified by silica gel chromatography (0 - 15% MeOH/EtOAc) to give 110 mg of **333**. Yield 62.7%. MS (ESI(+)) m/z 717.44 (M+H)⁺.

Part E.


20



To a solution of **333** (30 mg, 0.04 mmol, 1 eq) in DMF (1 mL) was added HBTU (19 mg, 0.05 mmol, 1.2 eq) and (S)-N1,N1-dimethyl-3-phenylpropane-1,2-diamine (15 mg, 0.08 mmol, 2 eq), followed by TEA (17 μ L, 0.12 mmol, 3 eq). After stirring at rt for 30 min, the mixture was diluted with EtOAc (10 mL) and washed with brine (3 x 10 mL). The


organic layer was dried over MgSO_4 , filtered and concentrated *in vacuo*. The material was purified by HPLC to give 11 mg of **327**. Yield 30%. MS (ESI(+)) m/z 877.72 M^+ .

Example 124



334


Part A.



335

To a solution of aminoacetaldehyde dimethyl acetal (1.0 g, 9.51 mmol, 1 eq) in DCM (10 mL) under Ar at 0 °C was added triethylamine (2.64 mL, 19.02 mmol, 2 eq) followed by trifluoromethanesulfonic anhydride (2.40 mL, 14.27 mmol, 1.5 eq) dropwise. The resulting mixture was stirred at 0 °C for 1 h. The reaction was diluted with DCM (10 mL) and washed with saturated ammonium chloride solution (20 mL). The aqueous layer was extracted with DCM (2 x 20 mL). The combined organic layers were dried over MgSO_4 , filtered and concentrated *in vacuo*. Compound **335** was obtained as a pale yellow oil and used without further purification.


Part B.



336

1N HCl (1 mL) was added to 2,2-(dimethoxyethyl)trifluoromethanesulfonamide (500 mg, 2.1 mmol, 1 eq) and the resulting mixture was heated at 100 °C for 2 h. The solution was evacuated to dryness to give **336** as a brown oil, which was used without further purification.


Part C.



To a solution of the boronic acid (60 mg, 0.31 mmol, 1 eq) in THF (2 ml) was added the aldehyde 336 (147 mg, 0.77 mmol, 2.5 eq) in MeOH (1 mL), followed by acetic acid (35 μ L, 0.62 mmol, 2.0 eq) and sodium cyanoborohydride (97 mg, 1.54 mmol, 5.0 eq). The resulting mixture was stirred for 30 min. The reaction mixture was diluted with EtOAc (10 mL) and quenched with brine. The aqueous layer was acidified with 1N HCl to pH 3, and extracted with EtOAc (3 x 20 mL). The combined organic layers were dried over MgSO₄, filtered and concentrated *in vacuo* to afford yellow oil. The oil was purified by silica gel chromatography (2 – 10 % MeOH/DCM with 0.5 % acetic acid) to give 64 mg of the 337.

Yield 56.2%. MS (ESI(+)) m/z 371.22 (M+H)⁺.


Part D.



A solution of aryl iodide (90 mg, 0.157 mmol, 1 eq), boronic acid 337 (64 mg, 0.173 mmol, 1.1 eq), cesium carbonate (102 mg, 0.314 mmol, 2 eq) and potassium acetate (15 mg, 0.157 mmol, 1 eq) in DMSO (2 mL) was degassed by bubbling argon through the solution for 10 min. Pd(dppf)Cl₂ (23 mg, 0.031 mmol, 0.2 eq) was then added and the flask purged with argon. The mixture was heated at 70 °C for 4 h. The reaction mixture was added to brine (10 mL), acidified with 1 N HCl until the aqueous layer attained a pH of 3, and extracted with EtOAc (3 x 20 mL). The combined organic layers were dried over MgSO₄, filtered and concentrated *in vacuo* to afford a brown oil. The oil was purified by silica gel chromatography (0 - 15% MeOH/EtOAc) to give 55 mg of 338. Yield 44.6%.

MS (ESI(+)) m/z 771.45 (M+H)⁺.

Part E.



To a solution of **338** (55 mg, 0.071 mmol, 1 eq) in DMF (1 mL) was added HBTU (32.5 mg, 0.086 mmol, 1.2 eq) and (S)-N1,N1-dimethyl-3-phenylpropane-1,2-diamine (25 mg, 0.143 mmol, 2 eq), followed by TEA (30 μ L, 0.213 mmol, 3 eq). After stirring at rt for 30 min, the mixture was diluted with EtOAc (10 mL) and washed with brine (3 x 10 mL). The organic layer was dried over MgSO_4 , filtered and concentrated *in vacuo*. The material was purified by HPLC to give 16 mg of **334**. Yield 24%. MS (ESI(+)) m/z 931.61 M^+ .

Example 125

Bcl-2 and Bcl-xL binding affinity analysis data is presented below for various compounds of the invention. Note that "*****" indicates that the K_i is < 1 nM, "****" indicates that the K_i is 1-5 nM, "***" indicates that the K_i is 5-9 nM, "**" indicates that the K_i is >9 nM. Note that "++++" indicates that the K_i is <1 μ M, "+++ " indicates that the K_i is 1-5 μ M, "++" indicates that the K_i is 5-10 μ M, and "+" indicates that the K_i is > 10 μ M.

Compound	Bcl2	Bcl-XI	Compound	Bcl2	Bcl-XI	Compound	Bcl2	Bcl-XI
1	****	++++	84	*	+	240	****	+++
12	**	++	85	*	++	247	****	++++
15	****	++++	86	*	+	251	****	++++
23	****	++++	87	*	+	252	****	+++
24	****	++++	91	*	+	253	****	++++
25	****	++++	92	*	+	255	****	++++
26	****	++++	93	*	+	256	****	++++
33	****	++++	94	***	+	257	****	++++
34	****	++++	95	**	++	258	****	++++
35	****	++++	96	**	++	259	****	+++
36	****	++++	97	*	+	260	***	++++
37	****	++++	98	*	++	262	**	+++
38	****	+	99	*	+	263	****	++++
53	***	+	100	****	+++	264	****	++++
54	***	+	111	****	++++	265	****	++++
55	****	+++	112	****	++++	267	****	+++
60	****	+++	113	****	++++	268	***	+++
61	****	++++	114	****	++++	269	****	++++
62	****	++	115	****	++++	270	***	+++
63	****	++	126	****	++++	271	****	++++
64	****	+	127	****	++++	272	****	+++

65	****	†	136	****	††††	274	****	†††
66	****	††	143	****	††††	285	****	††††
67	****	††	144	****	††††	290	****	††††
68	****	††††	151	****	††††	296	****	††††
69	**	†	152	****	††††	299	****	††††
70	***	†	160	****	††††	302	****	†††
71	*	†††	165	****	††††	306	****	††††
72	****	††††	170	***	†††	308	****	††††
73	****	††††	176	***	†††	309	****	††††
74	*	††	182	****	††††	310	****	††††
75	**	†††	183	***	†††	311	****	††††
76	**	†	193	***	†††	312	****	††††
77	**	†††	196	****	††††	313	****	††††
78	*	††	202	****	††††	314	****	††††
79	*	††	206	****	††††	315	****	††††
80	**	††	213	****	†	316	****	††††
81	*	†	218	****	†††	320	****	††††
82	*	†	219	***	†††	321	*	††
83	*	†	230	****	††††	295-amine	****	††††

Incorporation by Reference


Incorporated by reference are all of the U.S. patents, U.S. patent application
5 publications, and PCT patent application publications designating the U.S. cited herein.

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than
routine experimentation, many equivalents to the specific embodiments of the invention
described herein. Such equivalents are intended to be encompassed by the following
10 claims.

What is claimed is:

1. A compound of formula **1**:



1

or a pharmaceutically acceptable salt thereof;


wherein independently for each occurrence

m is 0, 1, 2, or 3;

n, o, and p are independently for each occurrence 1, 2, 3, 4, or 5;

R₁ is -OH, -OC(O)R₆, -OC(O)N(R₆)(R₇), or -N(R₆)(R₇);

R₂ is -OH, -OC(O)Me, -OCH₂CO₂H, -OCH₂CO₂Et, -N₃, -N(R₈)(R₉), -N(R)C(O)N(R₈)(R₉), or -N(R)C(O)R₁₀; or has the formula **1b**;



1b

R₃ is alkyl, halide, alkoxy, (cycloalkyl)alkoxy, aralkyloxy, or -O(CH₂)₂-N(R₁₅)(R₁₆);

R₄ is alkyl, alkoxy, hydroxyalkyl, alkoxyalkyl, halide, nitro, amino, acyl, amido, acylamino, aminoalkyl, acylaminoalkyl, acylaminoalkylamino, sulfonylaminoalkylamino, carboxylate, or -N=C(N(R)₂)₂;

R₅ is -OH or -N(R₁₇)(R₁₈);

R₆ and R₇ are independently for each occurrence H, alkyl, aralkyl, heteroaralkyl, or -[C(R₁₅)(R₁₆)]_n-R₁₉;

R₈ and R₉ are independently for each occurrence H, alkyl, aralkyl, or heteroaralkyl;

R₁₀ is alkyl, haloalkyl, or -[C(R₁₅)(R₁₆)]_o-COOR;

R, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, and R₁₆ are independently for each occurrence H or alkyl;

R₁₇ and R₁₈ are independently for each occurrence H, alkyl, aralkyl, heteroaralkyl, alkoxy, or -[C(R₁₉)(R₂₀)]_p-R₂₁;

R₁₉ and R₂₀ are independently for each occurrence H, hydroxy, alkyl, alkoxy, amino, aminoalkyl, acylamino, sulfonylamino, aryl, aralkyl, cycloalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, or heteroaralkyl;

R₂₁ is independently for each occurrence H, alkyl, aryl, heteroaryl, heterocyclyl, heterocyclylalkyl, alkoxy, alkylsulfonyl, arylsulfonyl, alkylsulfonamido, arylsulfonamido, amino, amido, or carboxyl;

R₂₂ independently for each occurrence is halide or alkyl;

R₂₃ is selected from the group consisting of alkyl, hydroxyalkyl, alkoxyalkyl, acyloxyalkyl, and haloalkyl;

R₂₄ is selected from the group consisting of alkyl, hydroxyalkyl, alkoxyalkyl, acyloxyalkyl, and haloalkyl; and

R₂₅ is selected from the group consisting of alkyl, hydroxyalkyl, alkoxyalkyl, acyloxyalkyl, and haloalkyl.

2. The compound of claim 1, wherein R₁ is selected from the group consisting of -OH, -OC(O)Me, -OC(O)(CH₂)₂Ph, -OC(O)CH₂CHMe₂, -OC(O)NHMe, -OC(O)NMe₂, -OC(O)NHCH₂(4-(OH)-Ph), -OC(O)NHPh, -OC(O)NHCH₂Ph, -OC(O)NH(CH₂)₄Ph, -OC(O)NH(CH₂)₂Ph, -OC(O)NH(CH₂)₂Me, -OC(O)NH(CH₂)₂NMe₂, -OC(O)NH(CH₂)₂NHC(O)Me, -OC(O)NH(CH₂)₂CHMe₂, -NHMe, -NH(CH₂)₂Ph, -NHC(O)Me, and -NH(CH₂)₂NMe₂.

3. The compound of claim 1, wherein R₁ is -OH.

4. The compound of claim 1, wherein R₂ is -OH, -OC(O)Me, -OCH₂CO₂H, -OCH₂CO₂Et, -N₃, -N=C(NMe₂)₂, -NH₂, -NMe₂, -NHC(O)Me, -NHC(O)CF₃, -NHC(O)Ph, -NHC(O)NHPh, -NHC(O)CH₂CH₂CO₂H, -NHC(O)CH₂CH₂CO₂Me, -NHCH₂Ph, -NHCH₂(4-pyridyl), -NHCH₂(2-

pyridyl), -NHCH₂(4-(CO₂H)Ph), -NHCH₂(3-(CO₂H)Ph), -NH₂, -NHCHMe₂, -NHCH₂CHMe₂, -N(CH₂CHMe₂)₂, -NHCH₂(cyclopropyl), or -NHC(O)CH₂CH₂NMe₂.

5. The compound of claim 1, wherein R₂ is -OH or -NH₂.

6. The compound of claim 1, wherein R₂ is -NH₂.

7. The compound of claim 1, wherein R₂ is -OH.

8. The compound of claim 1, wherein R₃ is -OMe, -OEt, -OCH₂(cyclopropyl), F, -O(CH₂)₂NMe₂, -O(CH₂)₂(4-morpholino), -OCH₂(4-(MeO)Ph), or -OCH₂(2-pyridyl).

9. The compound of claim 1, wherein R₃ is -OMe.

10. The compound of claim 1, wherein R₄ is -NMe₂, -NEt₂, -NH₂, -NHCH₂CHMe₂, -N(Me)CH₂CHMe₂, -N(Me)CH₂CH₂NHC(O)Me, -N(Me)CH₂CH₂NHS(O)₂Me, -N(Me)CH₂CH₂NHS(O)₂CF₃, -NHCH₂CH₂NMe₂, -NHCH₂CH₂NMeCH₂CH₂Cl, -NHCH₂CH₂OH, -N(Me)CH₂CH₂OH, -N(CH₂CH₂OH)₂, -N(Me)CH₂CO₂H, -N(Me)CH₂C(O)NH₂, -N(Me)CH₂C(O)NHMe, -N(Me)CH₂C(O)NMe₂, -NHC(O)Me, -NHC(O)CHMe₂, 1-pyrrolidinyl, 1-piperidinyl, 4-morpholino, (R)-2-(hydroxymethyl)-1-pyrrolidinyl, (S)-2-(hydroxymethyl)-1-pyrrolidinyl, (R)-2-(C(O)NMe₂)-1-pyrrolidinyl, (S)-2-(C(O)NMe₂)-1-pyrrolidinyl, -NH₂, -NO₂, Br, Cl, F, -C(O)Me, -C(O)NMe₂, -C(O)NH₂, -CO₂H, -CHO, -CH₂OH, -CH(Me)OH, -CH₂NH₂, -CH₂NHMe, -CH₂NMe₂, -CH₂NHC(O)Me, -CF₃, or tert-butyl.

11. The compound of claim 1, wherein R₄ is amino.

12. The compound of claim 1, wherein R₄ is -NMe₂, -N(Me)CH₂CH₂OH, -N(Me)CH₂C(O)NMe₂, -N(Me)CH₂C(O)NHMe, (R)-2-(hydroxymethyl)-1-pyrrolidinyl, or (R)-2-(C(O)NMe₂)-1-pyrrolidinyl.

13. The compound of claim 1, wherein R₄ is -NMe₂.

14. The compound of claim 1, wherein R₂₂ is selected from the group consisting of F, Cl, and tert-butyl; and m is 0 or 1.

15. The compound of claim 1, wherein R₂₂ is F; and m is 0 or 1.


16. The compound of claim 1, wherein R₂₂ is selected from the group consisting of F, Cl, and tert-butyl; and m is 1.


17. The compound of claim 1, wherein R_{22} is F; and m is 1.
18. The compound of claim 1, wherein R_{23} is selected from the group consisting of methyl, hydroxymethyl, alkoxymethyl, acyloxymethyl, and halomethyl.
19. The compound of claim 1, wherein R_{24} is selected from the group consisting of methyl, hydroxymethyl, alkoxymethyl, acyloxymethyl, and halomethyl.
20. The compound of claim 1, wherein R_{23} is methyl.
21. The compound of claim 1, wherein R_{24} is methyl.
22. The compound of claim 1, wherein R_{23} is methyl; and R_{24} is methyl.
23. The compound of claim 1, wherein R_{25} is selected from the group consisting of methyl, hydroxymethyl, alkoxymethyl, acyloxymethyl, and halomethyl.
24. The compound of claim 1, wherein R_{25} is methyl.
25. The compound of claim 1, wherein R_{23} is methyl; R_{24} is methyl; and R_{25} is methyl.
26. The compound of claim 1, wherein R_1 is -OH; and R_2 is -OH or -NH₂.
27. The compound of claim 1, wherein R_1 is -OH; R_2 is -OH or -NH₂; and R_3 is -OMe.
28. The compound of claim 1, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is -OMe; and R_4 is amino.
29. The compound of claim 1, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is -OMe; and R_4 is -NMe₂, -N(Me)CH₂CH₂OH, -N(Me)CH₂C(O)NMe₂, -N(Me)CH₂C(O)NHMe, (R)-2-(hydroxymethyl)-1-pyrrolidinyl, or (R)-2-(C(O)NMe₂)-1-pyrrolidinyl.
30. The compound of claim 1, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is -OMe; and R_4 is -NMe₂.
31. The compound of claim 1, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_{23} is methyl; R_{24} is methyl; and R_{25} is methyl.
32. The compound of claim 1, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is -OMe; R_{23} is methyl; R_{24} is methyl; and R_{25} is methyl.
33. The compound of claim 1, wherein R_1 is -OH; R_2 is -OH or -NH₂; R_3 is -OMe; R_4 is amino; R_{23} is methyl; R_{24} is methyl; and R_{25} is methyl.









34. The compound of claim 1, wherein R₁ is -OH; R₂ is -OH or -NH₂; R₃ is -OMe; R₄ is -NMe₂, -N(Me)CH₂CH₂OH, -N(Me)CH₂C(O)NMe₂, -N(Me)CH₂C(O)NHMe, (R)-2-(hydroxymethyl)-1-pyrrolidinyl, or (R)-2-(C(O)NMe₂)-1-pyrrolidinyl; R₂₃ is methyl; R₂₄ is methyl; and R₂₅ is methyl.









35. The compound of claim 1, wherein R₁ is -OH; R₂ is -OH or -NH₂; R₃ is -OMe; R₄ is -NMe₂; R₂₃ is methyl; R₂₄ is methyl; and R₂₅ is methyl.







36. A compound selected from the group consisting of:
















































































































































































































































































































































1




2




3




4




5














6













7























37. A compound selected from the group consisting of:

