

COMPOSITIONS AND METHODS FOR SIMULTANEOUSLY MODULATING EXPRESSION OF GENES

CROSS-REFERENCE TO RELATED APPLICATIONS

- 5 [0001] This application claims the benefit of U.S. Provisional Application No. 63/087,643, filed October 5, 2020 and U.S. Provisional Application No. 63/213,841, filed June 23, 2021, each of which is incorporated by reference herein in its entirety.

SEQUENCE LISTING

- 10 [0002] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on September 30, 2021, is named 57623_707_601_SL.txt and is 295,347 bytes in size.

BACKGROUND

- 15 [0003] Many aberrant human conditions are caused by or associated with shifts in gene expression level relative to those protein expression levels in subjects without such aberrant human conditions. This is particularly so in the case of cancer. For example, cancer cells are known to benefit from increasing expression of proteins involved in cell proliferation or angiogenesis and reducing expression of proteins involved in immune response to tumors. Thus, there is a need for therapies that decrease production of one or more target gene products involved in cell proliferation or angiogenesis and concomitantly increase production of others such as proteins involved in immune response to tumors needed to prevent or treat incidents of cancer in a subject.

25

BRIEF SUMMARY

- [0004] Provided herein are compositions and methods for simultaneously modulating expression of two or more proteins or nucleic acid sequences using one recombinant polynucleic acid or RNA construct. In some aspects, provided herein, is a composition comprising a first RNA linked to a second RNA, wherein the first RNA encodes for a cytokine, and wherein the second RNA encodes for a genetic element that modulates expression of a gene associated with tumor proliferation. In some aspects, provided herein, is a composition comprising a first RNA linked to a second RNA, wherein the first RNA encodes for a cytokine, and wherein the second RNA encodes for a genetic element that
- 30

modulates expression of a gene associated with recognition by the immune system. In some aspects, provided herein, is a pharmaceutical composition comprising any of the compositions described herein and a pharmaceutically acceptable excipient.

[0005] In some aspects, provided herein, is a composition comprising a first RNA encoding for interleukin-2 (IL-2), IL-15, a fragment thereof, or a functional variant thereof linked to a second RNA encoding for a genetic element that modulates expression of vascular endothelial growth factor A (VEGFA), an isoform of VEGFA, placental growth factor (PIGF), cluster of differentiation 155 (CD155), programmed cell death-ligand 1 (PD-L1), myc proto-oncogene (c-Myc), a fragment thereof, or a functional variant thereof. In some aspects, provided herein, is a composition comprising a first RNA encoding for interleukin-2 (IL-2), a fragment thereof, or a functional variant thereof linked to a second RNA encoding for a genetic element that modulates expression of MHC class I chain-related sequence A (MICA), MHC class I chain-related sequence B (MICB), endoplasmic reticulum protein (ERp5), a disintegrin and metalloproteinase (ADAM), matrix metalloproteinase (MMP), a fragment thereof, or a functional variant thereof. In some embodiments, the ADAM is ADAM17. In some aspects, provided herein, is a composition comprising a first RNA encoding for interleukin-12 (IL-12), IL-7, a fragment thereof, or a functional variant thereof linked to a second RNA encoding for a genetic element that modulates expression of isocitrate dehydrogenase (IDH1), cyclin-dependent kinase 4 (CDK4), CDK6, epidermal growth factor receptor (EGFR), mechanistic target of rapamycin (mTOR), Kirsten rat sarcoma viral oncogene (KRAS), programmed cell death-ligand 1 (PD-L1), a fragment thereof, or a functional variant thereof. In some aspects, provided herein, is a pharmaceutical composition comprising any of the compositions described herein and a pharmaceutically acceptable excipient.

[0006] In some aspects, provided herein, is a method of treating cancer, comprising administering any of the compositions or the pharmaceutical composition described herein to a subject having a cancer. In some embodiments, the cancer is a solid tumor. In some embodiments, the cancer is melanoma. In some embodiments, the cancer is renal cell carcinoma. In some embodiments, the cancer is a head and neck cancer. In some embodiments, the head and neck cancer is head and neck squamous cell carcinoma. In some embodiments, the head and neck cancer is laryngeal cancer, hypopharyngeal cancer, tonsil cancer, nasal cavity cancer, paranasal sinus cancer, nasopharyngeal cancer, metastatic squamous neck cancer with occult primary, lip cancer, oral cancer, oropharyngeal cancer, salivary gland cancer, brain tumors, esophageal cancer, eye cancer, parathyroid

cancer, sarcoma of the head and neck, or thyroid cancer. In some embodiments, the subject is a human.

[0007] In some aspects, provided herein, is a composition comprising a recombinant polynucleic acid construct comprising a nucleic acid sequence selected from the group
5 consisting of SEQ ID NOs: 1-17 and 125-141.

INCORPORATION BY REFERENCE

[0008] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent,
10 or patent application was specifically and individually indicated to be incorporated by reference.

BRIEF DESCRIPTION OF THE DRAWINGS

[0009] The features of the present disclosure are set forth with particularity in the appended
15 claims. A better understanding of the features and advantages of the present disclosure will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the disclosure are utilized, and the accompanying drawings of which:

[0010] **Figure 1** depicts a schematic representation of construct design. A polynucleic acid
20 construct may comprise a T7 promoter sequence upstream of the gene of interest sequence (IL-2 given as an example) for T7 RNA polymerase binding and successful *in vitro* transcription of both the gene of interest and siRNA in a single transcript. Signal peptide of IL-2 is highlighted in a grey box. Linkers to connect mRNA to siRNA or siRNA to siRNA are indicated with boxes with horizontal stripes or boxes with checkered stripes, respectively. T7:
25 T7 promoter, siRNA: small interfering RNA.

[0011] **Figure 2A** is a plot for induction of IL-2 secretion from human embryonic kidney cells (HEK-293). The X-axis indicates mRNAs used for transfection into HEK-293 cells: Compound (Cpd.) 1, Cpd.2, Cpd.3, or Cpd.4. The Y-axis is a measurement of IL-2 protein secretion fold change compared to IL-2 protein secretion by Cpd.1 using ELISA. Data
30 represent means $\pm$ standard error of the mean of 3 replicates per Cpd. Significance (**, $p < 0.01$) was assessed by one way ANOVA followed by Dunnet's multiple comparing test using Cpd.1 as control.

[0012] **Figure 2B** is a plot for induction of IL-2 secretion from human adult keratinocytes (HaCaT). The X-axis indicates mRNAs used for transfection into HaCaT cells: Compound

(Cpd.) 1, Cpd.2, Cpd.3, or Cpd.4. The Y-axis is a measurement of IL-2 protein secretion fold change compared to IL-2 protein secretion by Cpd.1 using ELISA. Data represent means $\pm$ standard error of the mean of 3 replicates per Cpd. Significance (**, $p < 0.01$) was assessed by one way ANOVA followed by Dunnet's multiple comparing test using Cpd.1 as control.

5 **[0013] Figure 2C** is a plot for induction of IL-2 secretion from human lung epithelial cells (A549). The X-axis indicates mRNAs used for transfection into A549 cells: Compound (Cpd.) 1, Cpd.2, Cpd.3, or Cpd.4. The Y-axis is a measurement of IL-2 protein secretion fold change compared to IL-2 protein secretion by Cpd.1 using ELISA. Data represent means $\pm$ standard error of the mean of 3 replicates per Cpd. Significance (**, $p < 0.01$) was assessed by
10 one way ANOVA followed by Dunnet's multiple comparing test using Cpd.1 as control.

[0014] Figure 3 is a plot for dose-dependent secretion of IL-2 protein and simultaneous interference of VEGFA expression by Compound 5 (Cpd.5) in lung epithelial cells (A549 cells) which overexpresses VEGFA (0.3 μ g VEGFA mRNA). The X-axis indicates concentrations of Cpd.5 (4.4, 8.8, 17.6, 26.4, 35.2 and 44.02 nM that correspond to 0, 150,
15 300, 600, 900, or 1200 ng/well, respectively) used for transfection into A549 cells. The Y-axis is a measurement of VEGFA (left) and IL-2 (right) protein levels (ng/ml) in the same cell culture supernatant by ELISA, 24 hours after transfection with Cpd.5. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0015] Figure 4A is a plot for interference of VEGFA expression by Compound 5 (Cpd.5) in
20 human tongue cell carcinoma cells (SCC-4) transfected with VEGFA mRNA to overexpress VEGFA. The X-axis indicates SCC-4 cells transfected with 9.5 nM (300 ng) of VEGFA mRNA only (VEGFA mRNA) or co-transfected with 9.5 nM (300 ng) of VEGFA mRNA and 26.4 nM (900 ng) of Cpd.5 (Cpd.5). The Y-axis is a measurement of VEGFA protein level (ng/ml) in cell culture supernatant by ELISA, 24 hours after transfection. Data represent
25 means $\pm$ standard error of the mean of 4 replicates.

[0016] Figure 4B is a plot for IL-2 protein level (ng/ml) in the same cell culture supernatant as in **Figure 4A**, measured by ELISA. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0017] Figure 5A is a plot for interference of VEGFA expression by Compound 5 (Cpd.5) in
30 human tongue cell carcinoma cells (SCC-4) that endogenously overexpress VEGFA. The X-axis indicates SCC-4 cells before (Endogenous) and after transfection (Cpd.5) with 26.4 nM (900 ng) of Cpd.5. The Y-axis is a measurement for VEGFA protein level (ng/ml) in cell culture supernatant by ELISA, 24 hours after transfection. Data represent means $\pm$ standard error of the mean of two replicates.

[0018] **Figure 5B** is a plot for IL-2 protein level (ng/ml) in the same cell culture supernatant as in **Figure 5A**, measured by ELISA. Data represent means $\pm$ standard error of the mean of two replicates.

[0019] **Figure 6A** is a plot for interference of VEGFA expression by Compound 5 (Cpd.5) and commercial siRNA in human tongue cell carcinoma cells (SCC-4) transfected with VEGFA mRNA to overexpress VEGFA (9.5 nM or 0.3 μ g VEGFA mRNA). The X-axis indicates SCC-4 cells transfected with increasing concentration of Cpd.5 (4.4 nM to 44.02 nM) or commercial siRNA (0.05 mM to 2.5 mM). The Y-axis indicates a measurement of VEGFA protein level (pg/ml) in cell culture supernatant by ELISA, 24 hours after transfection. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0020] **Figure 6B** is a plot for interference of VEGFA expression by Compound 5 (Cpd.5) and commercial siRNA in human lung epithelial cells (A549) transfected with VEGFA mRNA to overexpress VEGFA (9.5 nM or 0.3 μ g VEGFA mRNA). The X-axis indicates A549 cells transfected with increasing concentration of Cpd.5 (4.4 nM to 44.02 nM) or commercial siRNA (0.05 mM to 2.5 mM). The Y-axis indicates a measurement of VEGFA protein level (pg/ml) in cell culture supernatant by ELISA, 24 hours after transfection. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0021] **Figure 6C** is a table for comparison of IC50 values of Cpd. 5 and commercial siRNAs in SCC-4 and A549 cells.

[0022] **Figure 7A** is a plot for interference of MICB expression by Compound 6 (Cpd.6) in human tongue cell carcinoma cells (SCC-4) that constitutively express soluble and membrane MICB. The X-axis indicates SCC-4 cells before (Endogenous) and after transfection (Cpd.6) with 35.11 nM (900 ng) of Cpd.6. The Y-axis is a measurement for soluble MICB protein level (pg/ml) in cell culture supernatant by ELISA, 24 hours after transfection. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0023] **Figure 7B** is a plot for interference of MICB expression by Compound 6 (Cpd.6) in human tongue cell carcinoma cells (SCC-4) that constitutively express soluble and membrane MICB. The X-axis indicates SCC-4 cells before (Endogenous) and after transfection (Cpd.6) with 35.11 nM (900 ng) of Cpd.6. The Y-axis is a measurement for membrane MICB protein level (pg/ml) in cell culture supernatant by ELISA, 24 hours after transfection. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0024] **Figure 7C** is a plot for IL-2 protein level (ng/ml) in the same cell culture supernatant as in **Figure 7A** and **Figure 7B**, measured by ELISA. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0025] **Figure 8A** is a plot for dose-dependent secretion of IL-2 protein and simultaneous interference of MICA expression by Compound 6 (Cpd.6) in human tongue cell carcinoma cells (SCC-4) that constitutively express soluble MICA. The X-axis indicates concentrations of Cpd.6 (1.58, 2.93, 5.85, 11.7, 23.41, 35.11 and 46.81 nM) used for transfection into SCC-4 cells. The Y-axis is a measurement for soluble MICA protein level (pg/ml) in cell culture supernatant by ELISA, 24 hours after transfection. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0026] **Figure 8B** is a plot for dose-dependent secretion of IL-2 protein and simultaneous interference of MICB expression by Compound 6 (Cpd.6) in the same SCC-4 cells supernatant described in **Figure 8A**. SCC-4 cells constitutively express soluble MICB. The X-axis indicates concentrations of Cpd.6 (1.58, 2.93, 5.85, 11.7, 23.41, 35.11 and 46.81 nM) used for transfection into SCC-4 cells. The Y-axis is a measurement for soluble MICB protein level (pg/ml) in cell culture supernatant by ELISA, 24 hours after transfection. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0027] **Figure 9A** is a plot for IL-2 expression measured at 12, 24 and 48 hours post transfection with Cpd.3 (100 ng) in three-dimensional (3D) spheroid culture of SK-OV-3-NLR cells seeded at 5000 cells/ well into an ultra-low attachment (ULA) plate. IL-2 quantification was performed with TR-FRET assay. Error bars represent mean $\pm$ SEM of three replicates.

[0028] **Figures 9B-9D** shows changes in the total nuclear localized RFP (NLR) integrated intensity of SK-OV-3 NLR spheroids post transfection with Cpd.3 in the presence of peripheral blood mononuclear cells (PBMCs). SK-OV-3 NLR were plated in ULA plates (quadruplicate) at 5000 cells/well and transfected with different doses of Cpd.3 (3ng, 10ng, 30 ng and 100 ng) using Lipofectamine 2000. The cells were then centrifuged to form spheroids and cultured for 48 hrs prior to PBMC addition. PBMCs isolated from 3 donors (**Figures 9B, 9C and 9D**) were added at a density of 200,000 cells/well along with anti-CD3. The co-cultures were imaged every 3 hours for 168 hours (7 days). Total NLR integrated intensity was normalized to the 24 hour time point and analysed using the spheroid module within the IncuCyte software. rhIL2: recombinant human IL-2

[0029] **Figure 9E** shows a set of representative IncuCyte images showing Cpd.3 mediated NLR integrity reduction after PBMC alone control, recombinant human IL-2 (rhIL2) and Cpd.3 treatment (100 ng) in the SK-OV-3 NLR condition at Day-5.

[0030] **Figure 10A** is a plot showing dose-dependent activation of the JAK3/STAT5 pathway in HEK-Blue™ IL-2 reporter cells induced by rh-IL-2 (0.001 ng to 300 ng) or IL-2 (0.001 ng

– 45 ng) derived from supernatant of human embryonic kidney (HEK293) cells that had been transfected with Cpd.5 (0.3 µg/well) and quantified by ELISA. The X-axis indicates different concentration of Cpd.5 derived IL-2 or rh-IL-2. The Y-axis indicates IL-2 signaling activation normalized to rh-IL-2 (lowest SEAP values of rh-IL-2 set to 0 and highest SEAP values of rh-IL-2 set to 100%). Data represent means ± standard error of the mean of 4 replicates per dose.

[0031] Figure 10B is a plot showing dose-dependent activation of the JAK3/STAT5 pathway in HEK-Blue™ IL-2 reporter cells induced by rh-IL-2 (0.001 ng to 300 ng) or IL-2 (0.001 ng – 45 ng) derived from supernatant of human embryonic kidney (HEK293) cells that had been transfected with Cpd.6 (0.3 µg/well) and quantified by ELISA. The X-axis indicates different concentration of Cpd.6 derived IL-2 or rh-IL-2. The Y-axis indicates IL-2 signaling activation normalized to rh-IL-2. Data represent means ± standard error of the mean of 4 replicates per dose.

[0032] Figure 10C is a plot showing a NK cell mediated killing assay measured by luminescent cell viability approach (CellTiter-Glo). SCC-4 cells transfected with different doses of Cpd.5, Cpd.6 and two mock control RNAs (0.1 nM to 2.5 nM). 30 minutes after transfection, NK-92 cells were co-cultured with SCC-4 cells at the 10:1 effector to target (E:T) cell ratio and then incubated for 24 hours at 37°C. Cells were then thoroughly washed to remove NK-92 cells, and survived SCC-4 cells were analyzed by cell viability assay using CellTiter-Glo. Untreated SCC-4 cells were used as control and set to 0%. Data represent mean ± SEM from 4 replicates per dose.

[0033] Figure 11A is a plot showing dose-dependent downregulation of endogenously expressed VEGFA induced by Compound 7 (Cpd.7) and Compound 8 (Cpd.8) in SCC-4 cells. VEGFA levels in the cell culture supernatant were measured by ELISA, 24 hours after transfection. The X-axis indicates concentrations of Cpd.7 (1.1, 2.2, 4.4, 8.8, 17.6, 26.4, 35.2 and 44.04 nM/well) and Cpd.8 (0.47, 0.94, 1.89, 3.79, 7.58, 15.15, 22.73, 30.31 and 37.88 nM/well) used for transfection into SCC-4 cells. VEGFA levels from untransfected cells were set to 100%. The Y-axis indicates down regulation of VEGFA level normalized to untransfected samples (basal level). Data represent means ± standard error of the mean of 4 replicates.

[0034] Figure 11B is a plot showing dose-dependent secretion of IL-2 levels induced by Cpd.7 (3x siRNA) and Cpd.8 (5x siRNA) in SCC-4 cells. IL-2 levels in the cell culture supernatant were measured by ELISA, 24 hours after transfection. The X-axis indicates concentrations of Cpd.7 (1.1, 2.2, 4.4, 8.8, 17.6, 26.4, 35.2 and 44.04 nM/well) and Cpd.8 (0.47, 0.94, 1.89, 3.79, 7.58, 15.15, 22.73, 30.31 and 37.88 nM/well) used for transfection into

SCC-4 cells. The Y-axis is a measurement for IL-2 protein level (nM) in cell culture supernatant, 1nM correspond to dissociation constant (Kd) of IL-2 with its receptor. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0035] **Figure 11C** is a plot showing the time-course of IL-2 secretion induced by Compound 9 (Cpd.9) and Compound (Cpd.10) in SCC-4 cells up to 72 hours. IL-2 levels in the cell culture supernatant were measured by ELISA, from 6 to 72 hours after transfection (30 nM). The X-axis indicates hours after transfection and Y-axis is a measurement for IL-2 protein level (nM) in cell culture supernatant. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0036] **Figure 11D** is a plot for time-dependent downregulation of constitutively expressed VEGFA level by scrambled siRNA (scr. siRNA), commercial VEGFA siRNA, Cpd.9 and Cpd.10 in SCC-4 cells up to 72 hours. VEGFA levels in the cell culture supernatant were measured by ELISA, from 6 hours to 72 hours after transfection (30 nM). VEGFA levels from untransfected cells were set to 100% and down regulation was normalized to this value. The X-axis indicates hours after transfection and Y-axis indicates down regulation of VEGFA level normalized to untransfected samples (basal level). Data represent means $\pm$ standard error of the mean of 4 replicates.

[0037] **Figure 12A and Figure 12C** are plots showing secretion of IL-12 levels induced by compound 11 (Cpd.11) in SCC-4 cells and A549 cells, respectively. IL-12 levels in the cell culture supernatant were measured by ELISA, 24 hours after transfection. The X-axis indicates concentrations of Cpd.11 (7 (10 nM and 30 nM/well) used for transfection into SCC-4 cells. The Y-axis is an IL-12 protein level (pg/ml) in cell culture supernatant. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0038] **Figure 12B and Figure 12D** are plots showing downregulation of IDH1, CDK4 and CDK6 levels resulting from Cpd.11 treatment in SCC-4 cells and A549 cells, respectively. RNA levels of IDH1, CDK4 and CDK6 were measured from cell lysate by qPCR in technical duplicates, 24 hours after transfection. The X-axis indicates concentrations of Cpd.11 (10 nM and 30 nM/well) used for transfection into SCC-4 cells and A549 cells. The Y-axis indicates down regulation of IDH1, CDK4 and CDK6 level normalized to untransfected samples (basal level). Data represent means $\pm$ standard error of the mean of 4 replicates.

[0039] **Figure 12E and Figure 12G** are plots showing secretion of IL-12 levels induced by compound 12 (Cpd.12) in SCC-4 cells and A549 cells, respectively. IL-12 levels in the cell culture supernatant were measured by ELISA, 24 hours after transfection. The X-axis indicates concentrations of Cpd.12 (10 nM and 30 nM/well) used for transfection into SCC-4

cells and A549 cells. The Y-axis is an IL-12 protein level (pg/ml) in cell culture supernatant. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0040] **Figure 12F and Figure 12H** are plots showing downregulation of EGFR, KRAS and mTOR levels resulting from Cpd.12 treatment in SCC-4 cells and A549 cells, respectively.

5 RNA levels of EGFR, KRAS and mTOR were measured from cell lysate by qPCR in technical duplicates, 24 hours after transfection. The X-axis indicates concentrations of Cpd.12 (10 nM and 30 nM/well) used for transfection into SCC-4 cells and A549 cells. The Y-axis indicates down regulation of EGFR, KRAS and mTOR level normalized to untransfected samples (basal level). BQL = below quantification limit of the assay. Data
10 represent means $\pm$ standard error of the mean of 4 replicates.

[0041] **Figure 13A and Figure 13B** are plots showing secretion of IL-12 levels induced by Compound 13 (Cpd.13) in A549 cells and SCC-4 cells, respectively. IL-12 levels in the cell culture supernatant were measured by ELISA, 24 hours after transfection. The X-axis
15 indicates concentrations of Cpd.13 (10 nM and 30 nM/well) used for transfection into A549 cells and SCC-4 cells. The Y-axis is an IL-12 protein level (pg/ml) in cell culture supernatant. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0042] **Figure 13C** is a plot showing secretion of IL-12 levels induced by Compound 14 (Cpd.14) in A549 cells. IL-12 levels in the cell culture supernatant were measured by ELISA, 24 hours after transfection. The X-axis indicates concentrations of Cpd.14 (10 nM and 30
20 nM/well) used for transfection into A549 cells. The Y-axis is an IL-12 protein level (pg/ml) in cell culture supernatant. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0043] **Figure 13D and Figure 13E** are plots showing downregulation of EGFR expression resulting from Cpd.13 treatment in A549 cells and SCC-4 cells, respectively. RNA levels of EGFR were measured from cell lysate by qPCR in technical duplicates, 24 hours after
25 transfection. The X-axis indicates concentrations of Cpd.13 (10 nM and 30 nM/well) used for transfection into A549 cells and SCC-4 cells. The Y-axis indicates down regulation of EGFR level normalized to untransfected samples (basal level). Data represent means $\pm$ standard error of the mean of 4 replicates.

[0044] **Figure 13F** is a plot showing downregulation of mTOR expression resulting from Cpd.14 treatment in A549 cells. RNA levels of mTOR were measured from cell lysate by
30 qPCR in technical duplicates, 24 hours after transfection. The X-axis indicates concentrations of Cpd.14 (10 nM and 30 nM/well) used for transfection into A549 cells. The Y-axis indicates down regulation of mTOR level normalized to untransfected samples (basal level). Data represent means $\pm$ standard error of the mean of 4 replicates.

[0045] **Figure 14A and Figure 14C** are plots showing secretion of IL-15 levels induced by Compound 15 (Cpd.15) in A549 cells and SCC-4 cells, respectively. IL-15 levels in the cell culture supernatant were measured by ELISA, 24 hours after transfection. The X-axis indicates concentrations of Cpd.15 (10 nM and 30 nM/well) used for transfection into A549 cells and SCC-4 cells. The Y-axis is an IL-15 protein level (pg/ml) in cell culture supernatant. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0046] **Figure 14B and Figure 14D** are plots showing downregulation of VEGFA and CD155 expression resulting from Cpd.15 treatment in A549 cells and SCC-4 cells, respectively. RNA levels of VEGFA and CD155 were measured from cell lysate by qPCR in technical duplicates, 24 hours after transfection. The X-axis indicates concentrations of Cpd.15 (10 nM and 30 nM/well) used for transfection into A549 cells and SCC-4 cells. The Y-axis indicates down regulation of VEGFA and CD155 level normalized to untransfected samples (basal level). Data represent means $\pm$ standard error of the mean of 4 replicates.

[0047] **Figure 14E** is a plot showing secretion of IL-15 levels induced by Compound 16 (Cpd.16) in human glioblastoma cell line (U251 MG) cells. IL-15 levels in the cell culture supernatant were measured by ELISA, 24 hours after transfection. The X-axis indicates concentrations of Cpd.16 (10 nM and 30 nM/well) used for transfection into U251 MG cells. The Y-axis is an IL-15 protein level (pg/ml) in cell culture supernatant. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0048] **Figure 14F** is a plot showing downregulation of VEGFA, PD-L1 and c-Myc expression resulting from Cpd.16 treatment in U251 MG cells. RNA levels of VEGFA, PD-L1 and c-Myc were measured from cell lysate by qPCR in technical duplicates, 24 hours after transfection. The X-axis indicates concentrations of Cpd.16 (10 nM and 30 nM/well) used for transfection into U251 MG cells. The Y-axis indicates down regulation of VEGFA, PD-L1 and c-Myc level normalized to untransfected samples (basal level). Data represent means $\pm$ standard error of the mean of 4 replicates.

[0049] **Figure 14G** is a plot showing secretion of IL-7 levels induced by Compound 17 (Cpd.17) in U251 MG cells. IL-7 levels in the cell culture supernatant were measured by ELISA, 24 hours after transfection. The X-axis indicates concentrations of Cpd.17 (10 nM and 30 nM/well) used for transfection into U251 MG cells. The Y-axis is an IL-7 protein level (pg/ml) in cell culture supernatant. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0050] **Figure 14H** is a plot showing downregulation of PD-L1 expression resulting from Cpd.17 treatment in U251 MG cells. RNA levels of PD-L1 were measured from cell lysate by

qPCR in technical duplicates, 24 hours after transfection. The X-axis indicates concentrations of Cpd.17 (10 nM and 30 nM/well) used for transfection into U251 MG cells. The Y-axis indicates down regulation of PD-L1 level normalized to untransfected samples (basal level). Data represent means $\pm$ standard error of the mean of 4 replicates.

5 **[0051]** **Figure 15A** is a plot showing downregulation of endogenously expressed VEGFA induced by Compound 5 (Cpd.5) and Compound 10 (Cpd.10) in SCC-4 cells. VEGFA levels in the cell culture supernatant were measured by ELISA, 24 hours after transfection. The X-axis indicates concentrations of Cpd.5 and Cpd.10 (20 and 30 nM) used for transfection into SCC-4 cells. VEGFA levels from untransfected cells represent the endogenous VEGFA
10 secretion levels of SCC-4 cells and were labelled as '0'. The Y-axis indicates VEGFA levels measured by ELISA. Data represent means $\pm$ standard error of the mean of 2 independent measurements.

[0052] **Figure 15B** is a plot showing the number of branching points induced by VEGFA from different media supernatants in **Figure15A** in the HUVEC in vitro angiogenesis model.
15 Recombinant human VEGFA (VEGF) was used as a control and number of branching points were counted from microscopical pictures at the 6 hours time point. Data represent means $\pm$ standard error of the mean of 6 independent measurements.

DETAILED DESCRIPTION

20 **[0053]** Provided herein are compositions and methods for modulating expression of two or more genes simultaneously, comprising at least one nucleic acid sequence encoding a gene of interest and at least one nucleic acid sequence encoding or comprising a small interfering RNA (siRNA) capable of binding to a target messenger RNA (mRNA). Also provided herein are compositions and methods for treating cancers, comprising recombinant RNA constructs
25 to simultaneously express a cytokine and a genetic element that reduces expression of a gene associated with tumor proliferation, angiogenesis, or recognition by the immune system from a single RNA transcript. Further provided herein are compositions and methods to modulate expression of two or more genes simultaneously. Provided herein are compositions comprising a first RNA linked to a second RNA, wherein the first RNA encodes for a
30 cytokine, and wherein the second RNA encodes for a genetic element that reduces expression of a gene associated with tumor proliferation, angiogenesis, or recognition by the immune system. In one example, the first RNA may be a messenger RNA (mRNA) encoding a cytokine and can increase the protein level of a cytokine. In another example, the second RNA or the genetic element that reduces expression of a gene associated with tumor

proliferation, angiogenesis, or recognition by the immune system can include a small interfering RNA (siRNA) capable of binding to a target mRNA and can downregulate the level of protein encoded by the target mRNA. In some embodiments, target mRNAs can include an mRNA of a gene associated with tumor proliferation, angiogenesis, or recognition by the immune system.

[0054] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, suitable methods, and materials are described below.

Definitions

[0055] Certain specific details of this description are set forth in order to provide a thorough understanding of various embodiments. However, one skilled in the art will understand that the present disclosure may be practiced without these details. In other instances, well-known structures have not been shown or described in detail to avoid unnecessarily obscuring descriptions of the embodiments. Unless the context requires otherwise, throughout the specification and claims which follow, the word “comprise” and variations thereof, such as, “comprises” and “comprising” are to be construed in an open, inclusive sense, that is, as “including, but not limited to.” Further, headings provided herein are for convenience only and do not interpret the scope or meaning of the claimed disclosure.

[0056] As used in this specification and the appended claims, the singular forms “a,” “an,” and “the” include plural referents unless the content clearly dictates otherwise. It should also be noted that the term “or” is generally employed in its sense including “and/or” unless the content clearly dictates otherwise. The terms “and/or” and “any combination thereof” and their grammatical equivalents as used herein, can be used interchangeably. These terms can convey that any combination is specifically contemplated. Solely for illustrative purposes, the following phrases “A, B, and/or C” or “A, B, C, or any combination thereof” can mean “A individually; B individually; C individually; A and B; B and C; A and C; and A, B, and C.” The term “or” can be used conjunctively or disjunctively unless the context specifically refers to a disjunctive use.

[0057] The term “about” or “approximately” can mean within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, i.e. the limitations of the measurement system.

For example, “about” can mean within 1 or more than 1 standard deviation, per the practice in the art. Alternatively, “about” can mean a range of up to 20%, up to 10%, up to 5%, or up to 1% of a given value. Alternatively, particularly with respect to biological systems or processes, the term can mean within an order of magnitude, within 5-fold, or within 2-fold, of a value. Where particular values are described in the application and claims, unless otherwise stated the term “about” meaning within an acceptable error range for the particular value should be assumed.

[0058] As used in this specification and claim(s), the words “comprising” (and any form of comprising, such as “comprise” and “comprises”), “having” (and any form of having, such as “have” and “has”), “including” (and any form of including, such as “includes” and “include”) or “containing” (and any form of containing, such as “contains” and “contain”) are inclusive or open-ended and do not exclude additional, unrecited elements or method steps. It is contemplated that any embodiment discussed in this specification can be implemented with respect to any method or composition of the present disclosure, and vice versa. Furthermore, compositions of the present disclosure can be used to achieve methods of the present disclosure.

[0059] Reference in the specification to “embodiments,” “certain embodiments,” “preferred embodiments,” “specific embodiments,” “some embodiments,” “an embodiment,” “one embodiment” or “other embodiments” mean that a particular feature, structure, or characteristic described in connection with the embodiments is included in at least some embodiments, but not necessarily all embodiments, of the present disclosures. To facilitate an understanding of the present disclosure, a number of terms and phrases are defined below.

[0060] The term “RNA” as used herein includes RNA which encodes an amino acid sequence (e.g., mRNA, etc.) as well as RNA which does not encode an amino acid sequence (e.g., siRNA, shRNA, miRNA etc.). The RNA as used herein may be a coding RNA, i.e., an RNA which encodes an amino acid sequence. Such RNA molecules are also referred to as mRNA (messenger RNA) and are single-stranded RNA molecules. The RNA as used herein may be a non-coding RNA, i.e., an RNA which does not encode an amino acid sequence or is not translated into a protein. A non-coding RNA can include, but is not limited to, a small interfering RNA (siRNA), a short or small harpin RNA (shRNA), a microRNA (miRNA), a piwi-interacting RNA (piRNA), and a long non-coding RNA (lncRNA). siRNAs as used herein may comprise a double-stranded RNA (dsRNA) region, a hairpin structure, a loop structure, or any combinations thereof. In some embodiments, siRNAs may comprise at least one shRNA, at least one dsRNA region, or at least one loop structure. In some embodiments,

siRNAs may be processed from a dsRNA or an shRNA. In some embodiments, siRNAs may be processed or cleaved by an endogenous protein, such as DICER, from an shRNA. In some embodiments, a hairpin structure or a loop structure may be cleaved or removed from an siRNA. For example, a hairpin structure or a loop structure of an shRNA may be cleaved or removed. In some embodiments, RNAs described herein may be made by synthetic, chemical, or enzymatic methodology known to one of ordinary skill in the art, made by recombinant technology known to one of ordinary skill in the art, or isolated from natural sources, or made by any combinations thereof. The RNA may comprise modified or unmodified nucleotides or mixtures thereof, e.g., the RNA may optionally comprise chemical and naturally occurring nucleoside modifications known in the art (e.g., N¹-Methylpseudouridine also referred herein as methylpseudouridine).

[0061] The terms “nucleic acid sequence,” “polynucleic acid sequence,” “nucleotide sequence” are used herein interchangeably and have the identical meaning herein and refer to DNA or RNA. In some embodiments, a nucleic acid sequence is a polymer comprising or consisting of nucleotide monomers, which are covalently linked to each other by phosphodiester-bonds of a sugar/phosphate-backbone. The terms “nucleic acid sequence,” “polynucleic acid sequence,” and “nucleotide sequence” may encompass unmodified nucleic acid sequences, i.e., comprise unmodified nucleotides or natural nucleotides. The terms “nucleic acid sequence,” “polynucleic acid sequence,” and “nucleotide sequence” may also encompass modified nucleic acid sequences, such as base-modified, sugar-modified or backbone-modified etc., DNA or RNA.

[0062] The terms “natural nucleotide” and “canonical nucleotide” are used herein interchangeably and have the identical meaning herein and refer to the naturally occurring nucleotide bases adenine (A), guanine (G), cytosine (C), uracil (U), thymine (T).

[0063] The term “unmodified nucleotide” is used herein to refer to natural nucleotides which are not naturally modified e.g., which are not epigenetically or post-transcriptionally modified *in vivo*. Preferably the term “unmodified nucleotides” is used herein to refer to natural nucleotides which are not naturally modified e.g., which are not epigenetically or post-transcriptionally modified *in vivo* and which are not chemically modified e.g. which are not chemically modified *in vitro*.

[0064] The term “modified nucleotide” is used herein to refer to naturally modified nucleotides such as epigenetically or post-transcriptionally modified nucleotides and to chemically modified nucleotides e.g., nucleotides which are chemically modified *in vitro*.

Recombinant RNA constructs

[0065] Provided herein are compositions and methods for treating cancers, comprising recombinant polynucleic acid or RNA constructs comprising a gene of interest and a genetic element that reduces expression of another gene by binding to a target RNA. Also provided
5 herein are compositions and methods to modulate expression of two or more genes simultaneously using a single RNA transcript. An example of the genetic element that reduces expression of another gene can include a small interfering RNA (siRNA) capable of binding to a target mRNA.

[0066] Further provided herein are recombinant polynucleic acid or RNA constructs
10 comprising a gene of interest and a genetic element that reduces expression of another gene such as siRNA, wherein the gene of interest and the genetic element that reduces expression of another gene such as siRNA may be present in a sequential manner from the 5' to 3' direction, as illustrated in **Fig. 1**, or from 3' to 5' direction. In one example, the gene of interest can be present 5' to or upstream of the genetic element that reduces expression of
15 another gene such as siRNA, and the gene of interest can be linked to siRNA by a linker (mRNA to siRNA/shRNA linker, can be also referred s a "spacer"), as illustrated in **Fig. 1**. In another example, the gene of interest may be present 3' to or downstream of the genetic element that reduces expression of another gene such as siRNA, and siRNA can be linked to the gene of interest by a linker (siRNA/shRNA to mRNA linker, can be also referred s a
20 "spacer"). Recombinant polynucleic acid or RNA constructs provided herein may comprise more than one species of siRNAs and each of more than one species of siRNAs can be linked by a linker (siRNA to siRNA or shRNA to shRNA linker). In some embodiments, the sequence of mRNA to siRNA (or siRNA to mRNA) linker and the sequence of siRNA to siRNA (or shRNA to shRNA) linker may be different. In some embodiments, the sequence of
25 mRNA to siRNA/shRNA (or siRNA/shRNA to mRNA) linker and the sequence of siRNA to siRNA (or shRNA to shRNA) linker may be the same. Recombinant polynucleic acid or RNA constructs provided herein may comprise more than one gene of interest and each of more than one gene of interest can be linked by a linker (mRNA to mRNA linker). As an example of a gene of interest, interleukin 2 (IL-2) is shown in **Fig. 1**. IL-2 comprises a signal peptide sequence at the N-terminus. IL-2 may comprise unmodified (WT) signal peptide sequence or modified signal peptide sequence. Recombinant polynucleic acid constructs provided herein may also comprise a promoter sequence for RNA polymerase binding. As an example, T7
30 promoter for T7 RNA polymerase binding is shown in **Fig. 1**.

[0067] Recombinant RNA constructs provided herein may comprise multiple copies of a gene of interest, wherein each of the multiple copies of a gene of interest encodes the same protein. Also provided herein are compositions comprising recombinant RNA constructs comprising multiple genes of interest, wherein, each of the multiple genes of interest encodes a different protein. Recombinant RNA constructs provided herein may comprise multiple species of siRNAs (e.g., at least two species of siRNAs), wherein each of the multiple species of siRNAs is capable of binding to the same target RNA. In some embodiments, each of the multiple species of siRNAs may bind to the same region of the same target RNA. In some embodiments, each of the multiple species of siRNAs may bind to a different region of the same target RNA. In some embodiments, some of the multiple species of siRNAs may bind to the same target RNA and some of the multiple species of siRNAs may bind to a different region of the same target RNA. Also provided herein are recombinant RNA constructs comprising multiple species of siRNAs, wherein each of the multiple species of siRNAs is capable of binding to a different target RNA. In some embodiments, the target RNA is a messenger (mRNA).

[0068] Provided herein are compositions comprising recombinant RNA constructs comprising a first RNA linked to a second RNA, wherein the first RNA encodes for a cytokine, and wherein the second RNA encodes for a genetic element that reduces expression of a gene associated with tumor proliferation, angiogenesis, or recognition by the immune system. In one example, the first RNA may be an mRNA encoding a cytokine and can increase cytokine protein levels. In another example, the second RNA or the genetic element that reduces expression of a gene associated with tumor proliferation, angiogenesis, or recognition by the immune system in compositions described herein can include a small interfering RNA (siRNA) capable of binding to a target mRNA. In some embodiments, a target mRNA may be an mRNA of a gene associated with tumor proliferation, angiogenesis, or recognition by the immune system, and can downregulate protein expression of the target mRNA.

[0069] A recombinant polynucleic acid or a recombinant RNA can refer to a polynucleic acid or RNA that is not naturally occurring and is synthesized or manipulated *in vitro*. A recombinant polynucleic acid or RNA can be synthesized in a laboratory and can be prepared by using recombinant DNA or RNA technology by using enzymatic modification of DNA or RNA, such as enzymatic restriction digestion, ligation, cloning, and/or *in vitro* transcription. A recombinant polynucleic acid can be transcribed *in vitro* to produce a messenger RNA (mRNA) and recombinant mRNAs can be isolated, purified, and used for transfection into a

cell. A recombinant polynucleic acid or RNA used herein can encode a protein, polypeptide, a target motif, a signal peptide, and/or a non-coding RNA such as small interfering RNA (siRNA). In some embodiments, under suitable conditions, a recombinant polynucleic acid or RNA can be incorporated into a cell and expressed within the cell.

5 [0070] Recombinant RNA constructs provided herein may comprise more than one nucleic acid sequences encoding a gene of interest. For example, recombinant RNA constructs may comprise 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or more nucleic acid sequences encoding a gene of interest. In some instances, each of the two or more nucleic acid sequences may encode the same gene of interest, wherein the mRNA encoded by the same gene of interest is
10 different from the siRNA target mRNA. In some instances, each of the two or more nucleic acid sequences may encode a different gene of interest, wherein the mRNA encoded by the different gene of interest is not a target of siRNA encoded in the same RNA construct. In some instances, recombinant RNA constructs may comprise three or more nucleic acid sequences encoding a gene of interest, wherein each of the three or more nucleic acid
15 sequences may encode the same gene of interest or a different gene of interest, and wherein mRNAs encoded by the same or the different gene of interest are not a target of siRNA encoded in the same RNA construct. For example, recombinant RNA constructs may comprise four nucleic acid sequences encoding a gene of interest, wherein three of the four nucleic acid sequences encode the same gene of interest and one of the four nucleic acid
20 sequences encodes a different gene of interest, and wherein mRNAs encoded by the same or different gene of interest are not a target of siRNA encoded in the same RNA construct.

[0071] Recombinant RNA constructs provided herein may comprise more than one species of siRNA targeting an mRNA of a gene associated with tumor proliferation, angiogenesis, or recognition by the immune system. For example, recombinant RNA constructs provided
25 herein may comprise 1-10 species of siRNA targeting the same mRNA or different mRNAs. In some instances, each of the 1-10 species of siRNA targeting the same mRNA may comprise the same sequence, i.e. each of the 1-10 species of siRNA binds to the same region of the target mRNA. In some instances, each of the 1-10 species of siRNA targeting the same mRNA may comprise different sequences, i.e. each of the 1-10 species of siRNA binds to
30 different regions of the target mRNA. Recombinant RNA constructs provided herein may comprise at least two species of siRNA targeting an mRNA of a gene associated with tumor proliferation, angiogenesis, or recognition by the immune system. For instance, recombinant RNA constructs provided herein, may comprise 3 species of siRNA targeting one mRNA and each of the 3 species of siRNA comprise the same nucleic acid sequence to target the same

region of the mRNA. In this example, each of the 3 species of siRNA may comprise the same nucleic acid sequence to target exon 1. In another example, each of the 3 species of siRNA may comprise different nucleic acid sequence to target different regions of the mRNA. In this example, one of the 3 species of siRNA may comprise a nucleic acid sequence targeting exon 1 and another one of the 3 species of siRNA may comprise a nucleic acid sequence targeting exon 2, etc. In yet another example, each of the 3 species of siRNA may comprise different nucleic acid sequence to target different mRNAs. In all aspects, siRNAs in recombinant RNA constructs provided herein may not affect the expression of the gene of interest such as cytokine, expressed by the mRNA in the same RNA construct compositions.

[0072] Provided herein are compositions comprising recombinant RNA constructs, comprising a first RNA encoding for a cytokine and a second RNA encoding for a genetic element that reduces expression of a gene associated with tumor proliferation, angiogenesis, or recognition by the immune system. The first RNA and second RNA in compositions described herein may be linked by a linker. In some instances, compositions comprising the first RNA and the second RNA further comprises a nucleic acid sequence encoding for the linker. The linker can be from about 6 to about 50 nucleotides in length. For example, the linker can be at least about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or at least about 40 nucleotides in length. For example, the linker can be at most about 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or at most about 50 nucleotides in length. In some instances, a tRNA linker can be used. The tRNA system is evolutionarily conserved cross living organism and utilizes endogenous RNases P and Z to process multicistronic constructs (Dong et al., 2016). In some instances, the tRNA linker described herein may comprise a nucleic acid sequence comprising

AACAAAGCACCAGTGGTCTAGTGGTAGAATAGTACCCTGCCACGGTACAGACCC
GGGTTCGATTCGCGCTGGTGCA (SEQ ID NO: 20). In some instances, a linker comprising a nucleic acid sequence comprising

ATAGTGAGTCGTATTAACGTACCAACAA (SEQ ID NO: 21) may be used to link the first RNA and the second RNA.

[0073] Recombinant RNA constructs provided herein may further comprise a 5' cap, a Kozak sequence, and/or internal ribosome entry site (IRES), and/or a poly(A) tail at the 3' end in a particular in order to improve translation. In some instances, recombinant RNA constructs may further comprise regions promoting translation known to any skilled artisan. Non-limiting

examples of the 5' cap can include an anti-reverse CAP analog, Clean Cap, Cap 0, Cap 1, Cap 2, or Locked Nucleic Acid cap (LNA-cap). In some instances, 5' cap may comprise $m_2^{7,3'}$ - $^oG(5')ppp(5')G$, $m7G$, $m7G(5')G$, $m7GpppG$, or $m7GpppGm$.

[0074] Recombinant RNA constructs provided herein may further comprise a poly(A) tail. In some instances, the poly(A) tail comprises 1 to 220 base pairs of poly(A) (SEQ ID NO: 150). For example, the poly(A) tail comprises 1, 3, 5, 8, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, or 220 base pairs of poly(A) (SEQ ID NO: 150). In some embodiments, the poly(A) tail comprises 1 to 20, 1 to 40, 1 to 60, 1 to 80, 1 to 100, 1 to 120, 1 to 140, 1 to 160, 1 to 180, 1 to 200, 1 to 220, 20 to 40, 20 to 60, 20 to 80, 20 to 100, 20 to 120, 20 to 140, 20 to 160, 20 to 180, 20 to 200, 20 to 220, 40 to 60, 40 to 80, 40 to 100, 40 to 120, 40 to 140, 40 to 160, 40 to 180, 40 to 200, 40 to 220, 60 to 80, 60 to 100, 60 to 120, 60 to 140, 60 to 160, 60 to 180, 60 to 200, 60 to 220, 80 to 100, 80 to 120, 80 to 140, 80 to 160, 80 to 180, 80 to 200, 80 to 220, 100 to 120, 100 to 140, 100 to 160, 100 to 180, 100 to 200, 100 to 220, 120 to 140, 120 to 160, 120 to 180, 120 to 200, 120 to 220, 140 to 160, 140 to 180, 140 to 200, 140 to 220, 160 to 180, 160 to 200, 160 to 220, 180 to 200, 180 to 220, or 200 to 220 base pairs of poly(A) (SEQ ID NO: 150). In some embodiments, the poly(A) tail comprises 1, 20, 40, 60, 80, 100, 120, 140, 160, 180, 200, or 220 base pairs of poly(A) (SEQ ID NO: 150). In some embodiments, the poly(A) tail comprises at least 1, 20, 40, 60, 80, 100, 120, 140, 160, 180, or at least 200 base pairs of poly(A) (SEQ ID NO: 151). In some embodiments, the poly(A) tail comprises at most 20, 40, 60, 80, 100, 120, 140, 160, 180, 200, or at most 220 base pairs of poly(A) (SEQ ID NO: 152). In some embodiments, the poly(A) tail comprises 120 base pairs of poly(A) (SEQ ID NO: 153).

[0075] Recombinant RNA constructs provided herein may further comprise a Kozak sequence. A Kozak sequence may refer to a nucleic acid sequence motif that functions as a protein translation initiation site. Kozak sequences are described at length in the literature, *e.g.*, by Kozak, M., *Gene* 299(1-2):1-34, incorporated herein by reference herein in its entirety. In some embodiments, the Kozak sequence described herein may comprise a sequence comprising GCCACC (SEQ ID NO: 19). In some embodiments, recombinant RNA constructs provided herein may further comprise a nuclear localization signal (NLS).

[0076] Recombinant RNA constructs described herein may include one or more nucleotide variants, including nonstandard nucleotide(s), non-natural nucleotide(s), nucleotide analog(s), and/or modified nucleotides. Examples of modified nucleotides include, but are not limited to diaminopurine, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine,

xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl)uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-methyladenosine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, 2,6-diaminopurine, N1-methylpseudouridine, and the like. In some cases, nucleotides may include modifications in their phosphate moieties, including modifications to a triphosphate moiety. Non-limiting examples of such modifications include phosphate chains of greater length and modifications with thiol moieties. In some embodiments, phosphate chains can comprise 4, 5, 6, 7, 8, 9, 10 or more phosphate moieties. In some embodiments, thiol moieties can include but are not limited to alpha-thiotriphosphate and beta-thiotriphosphates. In some embodiments, a recombinant RNA construct described herein does not comprise 5-methylcytosine and/or N6-methyladenosine.

[0077] Recombinant RNA constructs described herein may be modified at the base moiety, sugar moiety, or phosphate backbone. For example, modifications can be at one or more atoms that typically are available to form a hydrogen bond with a complementary nucleotide and/or at one or more atoms that are not typically capable of forming a hydrogen bond with a complementary nucleotide. In some embodiments, backbone modifications include, but are not limited to, a phosphorothioate, a phosphorodithioate, a phosphoroselenoate, a phosphorodiselenoate, a phosphoroanilothioate, a phosphoraniladate, a phosphoramidate, and a phosphorodiamidate linkage. A phosphorothioate linkage substitutes a sulfur atom for a non-bridging oxygen in the phosphate backbone and delay nuclease degradation of oligonucleotides. A phosphorodiamidate linkage (N3'→P5') allows prevents nuclease recognition and degradation. In some embodiments, backbone modifications include having peptide bonds instead of phosphorous in the backbone structure, or linking groups including carbamate, amides, and linear and cyclic hydrocarbon groups. For example, N-(2-aminoethyl)-glycine units may be linked by peptide bonds in a peptide nucleic acid. Oligonucleotides with modified backbones are reviewed in Micklefield, Backbone modification of nucleic acids: synthesis, structure and therapeutic applications, Curr. Med.

Chem., 8 (10): 1157-79, 2001 and Lyer et al., Modified oligonucleotides-synthesis, properties and applications, Curr. Opin. Mol. Ther., 1 (3): 344-358, 1999.

[0078] Recombinant RNA constructs provided herein may comprise a combination of modified and unmodified nucleotides. In some instances, the adenosine-, guanosine-, and cytidine-containing nucleotides are unmodified or partially modified. In some instances, for modified RNA constructs, 1%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% of uridine nucleotides may be modified. In some embodiments, 5% to 25% of uridine nucleotides are modified in recombinant RNA constructs. Non-limiting examples of the modified uridine nucleotides may comprise pseudouridines, N¹-Methylpseudouridines, or N¹-methylpseudo-UTP and any modified uridine nucleotides known in the art may be utilized. In some embodiments, recombinant RNA constructs may contain a combination of modified and unmodified nucleotides, wherein 1%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% of uridine nucleotides may comprise pseudouridines, N¹-Methylpseudouridines, N¹-methylpseudo-UTP, or any other modified uridine nucleotide known in the art. In some embodiments, recombinant RNA constructs may contain a combination of modified and unmodified nucleotides, wherein 1%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% of the uridine nucleotides may comprise N¹-Methylpseudouridines.

[0079] Recombinant RNA constructs provided herein may be codon-optimized. In general, codon optimization refers to a process of modifying a nucleic acid sequence for expression in a host cell of interest by replacing at least one codon (e.g., more than 1, 2, 3, 4, 5, 10, 15, 20, 25, 50, or more codons) of a native sequence with codons that are more frequently or most frequently used in the genes of that host cell while maintaining the native amino acid sequence. Codon usage tables are readily available, for example, at the "Codon Usage Database," and these tables can be adapted in a number of ways. Computer algorithms for codon optimizing a particular sequence for expression in a particular host cell are also available, such as Gene Forge[®] (Aptagen, PA) and GeneOptimizer[®] (ThermoFischer, MA) which is preferred. In some embodiments, recombinant RNA constructs may not be codon-optimized.

[0080] In some instances, recombinant RNA constructs may comprise a nucleic acid sequence comprising a sequence selected from the group consisting of SEQ ID NOs: 1-17 and 125-141.

RNA interference and small interfering RNA (siRNA)

[0081] RNA interference (RNAi) or RNA silencing is a process in which RNA molecules inhibit gene expression or translation, by neutralizing target mRNA molecules. RNAi process

is described in Mello & Conte (2004) Nature 431, 338-342, Meister & Tuschl (2004) Nature 431, 343-349, Hannon & Rossi (2004) Nature 431, 371-378, and Fire (2007) Angew. Chem. Int. Ed. 46, 6966-6984. Briefly, in a natural process, the reaction initiates with a cleavage of long double-stranded RNA (dsRNA) into small dsRNA fragments or siRNAs with a hairpin structure (i.e., shRNAs) by a dsRNA-specific endonuclease Dicer. These small dsRNA fragments or siRNAs are then integrated into RNA-induced silencing complex (RISC) and guide the RISC to the target mRNA sequence. During interference, the siRNA duplex unwinds, and the antisense strand remains in complex with RISC to lead RISC to the target mRNA sequence to induce degradation and subsequent suppression of protein translation.

Unlike commercially available synthetic siRNAs, siRNAs in the present invention can utilize endogenous Dicer and RISC pathway in the cytoplasm of a cell to get cleaved from recombinant RNA constructs (e.g., recombinant RNA constructs comprising an mRNA and one or more siRNAs) after cellular uptake and follow the natural process detailed above, as siRNAs in the recombinant RNA constructs of the present invention may comprise a hairpin loop structure. In addition, as the rest of the recombinant RNA constructs (i.e., mRNA) is left intact after cleavage of siRNAs by Dicer, the desired protein expression from the gene of interest in the recombinant RNA constructs of the present invention is attained.

[0082] Provided herein are compositions comprising recombinant RNA constructs comprising at least one nucleic acid sequence comprising a siRNA capable of binding to a target RNA. In some instances, the target RNA is an mRNA. In some embodiments, the siRNA is capable of binding to a target mRNA in the 5' untranslated region. In some embodiments, the siRNA is capable of binding to a target mRNA in the 3' untranslated region. In some embodiments, the siRNA is capable of binding to a target mRNA in an exon. In some instances, the target RNA is a noncoding RNA. In some embodiments, recombinant RNA constructs may comprise a nucleic acid sequence comprising a sense siRNA strand. In some embodiments, recombinant RNA constructs may comprise a nucleic acid sequence comprising an anti-sense siRNA strand. In some embodiments, recombinant RNA constructs may comprise a nucleic acid sequence comprising a sense siRNA strand and a nucleic acid sequence comprising an anti-sense siRNA strand. Details of siRNA comprised in the present invention are described in Cheng, et al. (2018) J. Mater. Chem. B., 6, 4638-4644, which is incorporated by reference herein.

[0083] For example, in some instances, recombinant RNA constructs may comprise at least 1 species of siRNA, i.e., a nucleic acid sequence comprising a sense strand of siRNA and a nucleic acid sequence comprising an anti-strand of siRNA. 1 species of siRNA, as described

herein, can refer to 1 species of sense strand siRNA and 1 species of anti-sense strand siRNA. In some instances, recombinant RNA constructs may comprise more than 1 species of siRNA, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more species of siRNA comprising a sense strand of siRNA and an anti-strand of siRNA. In some embodiments, recombinant RNA constructs may
5 comprise 1 to 20 species of siRNA. In some embodiments, recombinant RNA constructs may comprise at least 1, 2, 3, 4, 5, 6, 7, 8, 9, or at least 10 species of siRNA. In some embodiments, recombinant RNA constructs may comprise at most 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or at most 20 species of siRNA. In a preferred embodiment, recombinant RNA constructs described herein comprise at least 2 species of siRNA. In
10 another preferred embodiment, recombinant RNA constructs described herein comprise at least 3 species of siRNA.

[0084] Provided herein are compositions of recombinant RNA constructs comprising 1-20 or more siRNA species, wherein each of the 1-20 or more siRNA species is capable of binding to a target RNA. In some embodiments, a target RNA is an mRNA or a non-coding RNA. In
15 some instances, each of the siRNA species binds to the same target RNA. In one instance, each of the siRNA species may comprise the same sequence and bind to the same region or sequence of the same target RNA. For example, recombinant RNA constructs may comprise 1, 2, 3, 4, 5, or more siRNA species and each of the 1, 2, 3, 4, 5, or more siRNA species comprise the same sequence targeting the same region of a target RNA, *i.e.* recombinant RNA
20 constructs may comprise 1, 2, 3, 4, 5, or more redundant species of siRNA. In another instance, each of the siRNA species may comprise a different sequence and bind to a different region or sequence of the same target RNA. For example, recombinant RNA constructs may comprise 1, 2, 3, 4, 5, or more siRNA species and each of the 1, 2, 3, 4, 5, or more siRNA species may comprise a different sequence targeting a different region of the same target
25 RNA. In this example, one siRNA of the 1, 2, 3, 4, 5, or more siRNA species may target exon 1 and another siRNA of the 1, 2, 3, 4, 5, or more siRNA species may target exon 2 of the same mRNA, *etc.* In some instances, recombinant RNA constructs may comprise 1, 2, 3, 4, 5, or more siRNA species and 2 of the 1, 2, 3, 4, 5, or more siRNA species may comprise the same sequence and bind to the same regions of the target RNA and 3 or more of the 1, 2, 3, 4,
30 5, or more siRNA species may comprise a different sequence and bind to different regions of the same target RNA. In some instances, each of the siRNA species binds to a different target RNA. In some embodiments, a target RNA may be an mRNA or a non-coding RNA, *etc.*

[0085] Provided herein are compositions of recombinant RNA constructs comprising 1-20 or more siRNA species, wherein each of the 1-20 or more siRNA species are connected by a

linker. In some instances, the linker may be a non-cleavable linker. In some instances, the linker may be a cleavable linker such as a self-cleavable linker. In some instances, the linker may be a tRNA linker. The tRNA system is evolutionarily conserved across living organism and utilizes endogenous RNases P and Z to process multicistronic constructs (Dong et al., 5 2016). In some embodiments, the tRNA linker may comprise a nucleic acid sequence comprising

AACAAAGCACCAGTGGTCTAGTGGTAGAATAGTACCCTGCCACGGTACAGACCC
GGGTTCGATTCCCGGCTGGTGCA (SEQ ID NO: 20). In some embodiments, a linker
comprising a nucleic acid sequence comprising

10 TTTATCTTAGAGGCATATCCCTACGTACCAACAA (SEQ ID NO: 22) may be used to
connect different siRNA species.

[0086] In some instances, specific binding of an siRNA to its mRNA target results in
interference with the normal function of the target mRNA to cause a modulation, *e.g.*,
downregulation, of function and/or activity, and wherein there is a sufficient degree of
15 complementarity to avoid non-specific binding of the siRNA to non-target nucleic acid
sequences under conditions in which specific binding is desired, *i.e.* under physiological
conditions in the case of *in vivo* assays or therapeutic treatment, and under conditions in
which assays are performed in the case of *in vitro* assays.

[0087] A protein as used herein can refer to molecules typically comprising one or more
20 peptides or polypeptides. A peptide or polypeptide is typically a chain of amino acid residues,
linked by peptide bonds. A peptide usually comprises between 2 and 50 amino acid residues. A
polypeptide usually comprises more than 50 amino acid residues. A protein is typically folded
into 3-dimensional form, which may be required for the protein to exert its biological
function. A protein as used herein can include a fragment of a protein, a variant of a protein,
25 and fusion proteins. A fragment may be a shorter portion of a full-length sequence of a nucleic
acid molecule like DNA, RNA, or a protein. Accordingly, a fragment, typically, comprises a
sequence that is identical to the corresponding stretch within the full-length sequence. In some
embodiments, a fragment of a sequence may comprise at least 5% to at least 80% of a full-
length nucleotide or amino acid sequence from which the fragment is derived. In some
30 embodiments, a protein can be a mammalian protein. In some embodiments, a protein can be
a human protein. In some embodiments, a protein may be a protein secreted from a cell. In
some embodiments, a protein may be a protein on cell membranes. In some embodiments, a
protein as referred to herein can be a protein that is secreted and acts either locally or
systemically as a modulator of target cell signaling via receptors on cell surfaces, often

involved in immunologic reactions or other host proteins involved in viral infection.

Nucleotide and amino acid sequences of proteins useful in the context of the present invention, including proteins that are encoded by a gene of interest, are known in the art and available in the literature. For example, Nucleotide and amino acid sequences of proteins

5 useful in the context of the present invention, including proteins that are encoded by a gene of interest are available in the UniProt database.

[0088] Provided herein are compositions of recombinant RNA constructs comprising an siRNA capable of binding to a target mRNA to modulate expression of the target mRNA. In some instances, expression of the target mRNA (*e.g.*, the level of protein encoded by the target mRNA) is downregulated by the siRNA capable of binding to the target mRNA. In 10 some embodiments, expression of the target mRNA is inhibited by the siRNA capable of binding to the target mRNA. Inhibition or downregulation of expression of the target mRNA, as described herein, can refer to, but is not limited to, interference with the target mRNA to interfere with translation of the protein from the target mRNA; thus, inhibition or 15 downregulation of expression of the target mRNA can refer to, but is not limited to, a decreased level of proteins expressed from the target mRNA compared to a level of proteins expressed from the target mRNA in the absence of recombinant RNA constructs comprising siRNA capable of binding to the target mRNA. Levels of protein expression can be measured by using any methods well known in the art and these include, but are not limited to Western- 20 blotting, flow cytometry, ELISAs, RIAs, and various proteomics techniques. An exemplary method to measure or detect a polypeptide is an immunoassay, such as an ELISA. This type of protein quantitation can be based on an antibody capable of capturing a specific antigen, and a second antibody capable of detecting the captured antigen. Exemplary assays for detection and/or measurement of polypeptides are described in Harlow, E. and Lane, D. 25 Antibodies: A Laboratory Manual, (1988), Cold Spring Harbor Laboratory Press.

[0089] Provided herein are compositions comprising recombinant RNA constructs comprising at least one nucleic acid sequence comprising siRNA capable of binding to a target mRNA and at least one nucleic acid sequence encoding a gene of interest wherein the target mRNA is different from an mRNA encoded by the gene of interest. Provided herein are 30 compositions comprising recombinant RNA constructs comprising at least one nucleic acid sequence comprising siRNA capable of binding to a target mRNA and at least one nucleic acid sequence encoding a gene of interest wherein the siRNA does not affect expression of the gene of interest. In some instances, the siRNA is not capable of binding to an mRNA encoded by the gene of interest. In some instances, the siRNA does not inhibit the expression

of the gene of interest. In some instances, the siRNA does not downregulate the expression of the gene of interest. Inhibiting or downregulating the expression of the gene of interest, as described herein, can refer to, but is not limited to, interfering with translation of proteins from recombinant RNA constructs; thus, inhibiting or downregulating the expression of the gene of interest can refer to, but is not limited to, a decreased level of protein compared to a level of protein expressed in the absence of recombinant RNA constructs comprising siRNA capable of binding to the target mRNA. Levels of protein expression can be measured by using any methods well known in the art and these include, but are not limited to Western-blotting, flow cytometry, ELISAs, RIAs, and various proteomics techniques. An exemplary method to measure or detect a polypeptide is an immunoassay, such as an ELISA. This type of protein quantitation can be based on an antibody capable of capturing a specific antigen, and a second antibody capable of detecting the captured antigen. Exemplary assays for detection and/or measurement of polypeptides are described in Harlow, E. and Lane, D. Antibodies: A Laboratory Manual, (1988), Cold Spring Harbor Laboratory Press.

[0090] Provided herein are compositions comprising recombinant RNA constructs comprising at least one nucleic acid sequence comprising a siRNA capable of binding to a target mRNA. A list of non-limiting examples of target mRNAs that the siRNA is capable of binding to includes an mRNA of a gene associated with tumor proliferation, angiogenesis, or recognition by the immune system. For example, the target mRNA may be an mRNA encoding vascular endothelial growth factor (VEGF), VEGFA, an isoform of VEGFA, placental growth factor (PIGF), a fragment thereof, or a functional variant thereof. A functional variant as used herein may refer to a full-length molecule, a fragment thereof, or a variant thereof. For example, a variant molecule may comprise a sequence modified by insertion, deletion, and/or substitution of one or more amino acids, in the case of protein sequence, or one or more nucleotides, in the case of nucleic acid sequence. For example, a variant molecule may comprise or encode a mutant protein, including, but not limited to, a gain-of-function or a loss-of-function mutant. A list of non-limiting examples of VEGFA isoforms is shown in **Table A**.

[0091] Table A. List of VEGFA Isoforms

VEGFA Isoforms	UniProt Database #
VEGF111	P15692-10
VEGF121	P15692-9
VEGF145	P15692-6
VEGF148	P15692-5
VEGF165	P15692-4
VEGF165B	P15692-8

VEGFA Isoforms	UniProt Database #
VEGF183	P15692-3
VEGF189	P15692-2
VEGF206	P15692-1
L-VEGF121	P15692-12
L-VEGF165	P15692-11
L-VEGF189	P15692-13
L-VEGF206	P15692-14
Isoform 15	P15692-15
Isoform 16	P15692-16
Isoform 17	P15692-17
Isoform 18	P15692-18

[0092] In some embodiments, VEGFA comprises a sequence listed in SEQ ID NO: 34. An exemplary PIGF sequence is shown below:

[0093] PIGF NCBI Reference Sequence: NM_001207012.1 (SEQ ID NO: 123)

```

5  CCTCGCACGC ACTGCGGGCT CCGGCGCTGC GGGCTGGCCG GGCGCTGCGG
   GCTGACCGGG CGCTCCGGGA ACTCGGCTCG GGAACCTCGT CTGCGGTGGG
   CGGGGCCGGC CCGGAGCCCC GCCCGGCTC AGTCCCTGAA ACCCAGGCGC
   GGACCGGCTG CAGTCTCAGA AGGGAGCTGC TGTCTGCGGA GGAAACTGCA
   TCGACGGACG GCCGCCAGC TACGGGAGGA CCTGGAGTGG CACTGGGCGC
10  CCGACGGACC ATCCCCGGGA CCCGCTGCC CCTCGGCGCC CCGCCCCGCC
   GGGCCGCTCC CCGTCGGGTT CCCAGCCAC AGCCTTACCT ACGGGCTCCT
   GACTCCGCAA GGCTTCCAGA AGATGCTCGA ACCACCGGCC GGGGCCTCGG
   GGCAGCAGTG AGGGAGGCGT CCAGCCCCC ACTCAGCTCT TCTCCTCCTG
   TGCCAGGGGC TCCCCGGGGG ATGAGCATGG TGGTTTTCCC TCGGAGCCCC
15  CTGGCTCGGG ACGTCTGAGA AGATGCCGGT CATGAGGCTG TTCCCTTGCT
   TCCTGCAGCT CCTGGCCGGG CTGGCGCTGC CTGCTGTGCC CCCCAGCAG
   TGGGCCTTGT CTGCTGGGAA CGGCTCGTCA GAGGTGGAAG TGGTACCCTT
   CCAGGAAGTG TGGGGCCGCA GCTACTGCCG GGCCTGGAG AGGCTGGTGG
   ACCTCGTGTG CGAGTACCCC AGCGAGGTGG AGCACATGTT CAGCCCATCC
20  TGTGTCTCCC TGCTGCGCTG CACCGGCTGC TGCAGGATG AGAATCTGCA
   CTGTGTGCCG GTGGAGACGG CCAATGTCAC CATGCAGCTC CTAAAGATCC
   GTTCTGGGGA CCGGCCCTCC TACGTGGAGC TGACGTTCTC TCAGCACGTT
   CGCTGCGAAT GCCGGCCTCT GCGGGAGAAG ATGAAGCCGG AAAGGTGCGG
   CGATGCTGTT CCCCAGAGGT AACCCACCCC TTGGAGGAGA GAGACCCCGC
25  ACCCGGCTCG TGTATTTATT ACCGTCACAC TCTTCAGTGA CTCTGCTGG
   TACCTGCCCT CTATTTATTA GCCAACTGTT TCCCTGCTGA ATGCCTCGCT
   CCCTTCAAGA CGAGGGGCAG GGAAGGACAG GACCCTCAGG AATTCACTGC
   CTTCAACAAC GTGAGAGAAA GAGAGAAGCC AGCCACAGAC CCCTGGGAGC
   TTCCGCTTTG AAAGAAGCAA GACACGTGGC CTCGTGAGGG GCAAGCTAGG
30  CCCCAGAGGC CCTGGAGGTC TCCAGGGGCC TGCAGAAGGA AAGAAGGGGG
   CCCTGCTACC TGTTCTTGGG CCTCAGGCTC TGCACAGACA AGCAGCCCTT
   GCTTTCGGAG CTCTGTCCA AAGTAGGGAT GCGGATCCTG CTGGGGCCGC
   CACGGCCTGG CTGGTGGGAA GGCCGCGAGC GGGCGGAGGG GATCCAGCCA
   CTTCCCCCTC TTCTTCTGAA GATCAGAACA TTCAGCTCTG GAGAACAGTG
35  GTTGCCCTGG GGCTTTTGCC ACTCCTTGTC CCCCCTGATC TCCCCTCACA
   CTTTGCCATT TGCTTGTAAT GGGACATTGT TCTTTCCGGC CAAGGTGCCA
   CCACCCTGCC CCCCCTAAGA GACACATACA GAGTGGGCCC CGGGCTGGAG

```

AAAGAGCTGC CTGGATGAGA AACAGCTCAG CCAGTGGGGA TGAGGTCACC
 AGGGGAGGAG CCTGTGCGTC CCAGCTGAAG GCAGTGGCAG GGGAGCAGGT
 TCCCCAAGGG CCCTGGCACC CCCACAAGCT GTCCCTGCAG GGCCATCTGA
 CTGCCAAGCC AGATTCTCTT GAATAAAGTA TTCTAGTGTG GAAACGCT

5

[0094] For example, the target mRNA may be an mRNA encoding MHC class I chain-related sequence A (MICA), MHC class I chain-related sequence B (MICB), endoplasmic reticulum protein (ERp5), a disintegrin and metalloproteinase (ADAM), matrix metalloproteinase (MMP), a fragment thereof, or a functional variant thereof. A functional variant as used herein may refer to a full-length molecule, a fragment thereof, or a variant thereof. For example, a variant molecule may comprise a sequence modified by insertion, deletion, and/or substitution of one or more amino acids, in the case of protein sequence, or one or more nucleotides, in the case of nucleic acid sequence. For example, a variant molecule may comprise or encode a mutant protein, including, but not limited to, a gain-of-function or a loss-of-function mutant.

10

15

In some embodiments, the ADAM is ADAM 17. In some embodiments, the target mRNA may encode a decoy protein. In some embodiments the decoy protein is a soluble form of a cell receptor. In some embodiments, the decoy protein is soluble MICA, MICB, a fragment thereof, or a functional variant thereof. In some embodiments, the target mRNA may encode a protein involved in shedding of MICA and/or MICB from cell membranes. In some

20

embodiments, the protein involved in shedding of MICA and/or MICB from cell membranes comprises ERp5, ADAM, MMP, a fragment thereof, or a functional variant thereof. In some embodiments, the protein involved in shedding of MICA and/or MICB from cell membranes comprises ADAM17, a fragment thereof, or a functional variant thereof. An exemplary sequence of ADAM17 is shown below:

25

[0095] ADAM17 NCBI Reference Sequence: NM_003183.6 (SEQ ID NO: 124)

AGCGGCGGCC GGAAGCTGGC TGAGCCGGCC TTTGGTAAACG CCACCTGCAC
 TTCTGGGGGC GTCGAGCCTG GCGGTAGAAT CTTCCCAGTA GGCGGCGCGG
 GAGGGAAAAG AGGATTGAGG GGCTAGGCCG GGCGGATCCC GTCCCTCCCC
 GATGTGAGCA GTTTTCCGAA ACCCCGTCAG GCGAAGGCTG CCCAGAGAGG
 30 TGGAGTCGGT AGCGGGGCGG GGAACATGAG GCAGTCTCTC CTATTCCTGA
 CCAGCGTGGT TCCTTTCGTG CTGGCGCCGC GACCTCCGGA TGACCCGGGC
 TTCGGCCCCC ACCAGAGACT CGAGAAAGCTT GATTCTTTGC TCTCAGACTA
 CGATATTCTC TCTTTATCTA ATATCCAGCA GCATTCCGTA AGAAAAAGAG
 ATCTACAGAC TTCAACACAT GTAGAAACAC TACTAACTTT TTCAGCTTTG
 35 AAAAGGCATT TTAAATTATA CCTGACATCA AGTACTGAAC GTTTTTCACA
 AAATTTCAAG GTCGTGGTGG TGGATGGTAA AAACGAAAGC GAGTACACTG
 TAAAATGGCA GGAATTCTTC ACTGGACACG TGGTTGGTGA GCCTGACTCT
 AGGGTTCTAG CCCACATAAG AGATGATGAT GTTATAATCA GAATCAACAC
 AGATGGGGCC GAATATAACA TAGAGCCACT TTGGAGATTT GTTAATGATA
 40 CCAAAGACAA AAGAATGTTA GTTTATAAAT CTGAAGATAT CAAGAATGTT
 TCACGTTTGC AGTCTCCAAA AGTGTGTGGT TATTTAAAAG TGGATAATGA

	AGAGTTGCTC	CCAAAAGGGT	TAGTAGACAG	AGAACCACCT	GAAGAGCTTG
	TTCATCGAGT	GAAAAGAAGA	GCTGACCCAG	ATCCCATGAA	GAACACGTGT
	AAATTATTGG	TGGTAGCAGA	TCATCGCTTC	TACAGATACA	TGGGCAGAGG
	GGAAGAGAGT	ACAACTACAA	ATTACTTAAT	AGAGCTAATT	GACAGAGTTG
5	ATGACATCTA	TCGGAACACT	TCATGGGATA	ATGCAGGTTT	TAAAGGCTAT
	GGAATACAGA	TAGAGCAGAT	TCGCATTCTC	AAGTCTCCAC	AAGAGGTAAA
	ACCTGGTGAA	AAGCACTACA	ACATGGCAAA	AAGTTACCCA	AATGAAGAAA
	AGGATGCTTG	GGATGTGAAG	ATGTTGCTAG	AGCAATTTAG	CTTTGATATA
	GCTGAGGAAG	CATCTAAAGT	TTGCTTGGCA	CACCTTTTCA	CATACCAAGA
10	TTTTGATATG	GGAACCTCTG	GATTAGCTTA	TGTTGGCTCT	CCCAGAGCAA
	ACAGCCATGG	AGGTGTTTGT	CCAAAGGCTT	ATTATAGCCC	AGTTGGGAAG
	AAAAATATCT	ATTTGAATAG	TGTTTTGACG	AGCACAAAGA	ATTATGGTAA
	AACCATCCTT	ACAAAGGAAG	CTGACCTGGT	TACAACTCAT	GAATTGGGAC
	ATAATTTTGG	AGCAGAACAT	GATCCGGATG	GTCTAGCAGA	ATGTGCCCCG
15	AATGAGGACC	AGGGAGGGAA	ATATGTCATG	TATCCCATAG	CTGTGAGTGG
	CGATCACGAG	AACAATAAGA	TGTTTTCAAA	CTGCAGTAAA	CAATCAATCT
	ATAAGACCAT	TGAAAGTAAG	GCCCAGGAGT	GTTTTCAAGA	ACGCAGCAAT
	AAAGTTTGTG	GGAACTCGAG	GGTGGATGAA	GGAGAAGAGT	GTGATCCTGG
	CATCATGTAT	CTGAACAACG	ACACCTGCTG	CAACAGCGAC	TGCACGTTGA
20	AGGAAGGTGT	CCAGTGCAGT	GACAGGAACA	GTCCTTGCTG	TAAAAACTGT
	CAGTTTGAGA	CTGCCCAGAA	GAAGTGCCAG	GAGGCGATTA	ATGCTACTTG
	CAAAGGCGTG	TCCTACTGCA	CAGGTAATAG	CAGTGAGTGC	CCGCCTCCAG
	GAAATGCTGA	AGATGACACT	GTTTGCTTGG	ATCTTGGCAG	GTGTAAGGAT
	GGGAAATGCA	TCCCTTTCTG	CGAGAGGGAA	CAGCAGCTGG	AGTCCTGTGC
25	ATGTAATGAA	ACTGACAACT	CCTGCAAGGT	GTGCTGCAGG	GACCTTCTCTG
	GCCGCTGTGT	GCCCTATGTC	GATGCTGAAC	AAAAGAACTT	ATTTTTGAGG
	AAAGGAAAGC	CCTGTACAGT	AGGATTTTGT	GACATGAATG	GCAAATGTGA
	GAAACGAGTA	CAGCATGTAA	TTGAACGATT	TTGGGATTTT	ATTGACCAGC
	TGAGCATCAA	TACTTTTGGA	AAGTTTTTAG	CAGACAACAT	CGTTGGGTCT
30	GTCCTGGTTT	TCTCCTTGAT	ATTTTGGATT	CCTTTCAGCA	TTCTTGTCCA
	TTGTGTGGAT	AAGAAAATTG	ATAAACAGTA	TGAATCTCTG	TCTCTGTTTC
	ACCCCGAGTAA	CGTCGAAAATG	CTGAGCAGCA	TGGATTCTGC	ATCGGTTTCGC
	ATTATCAAAC	CCTTTCCTGC	GCCCCAGACT	CCAGGCCGCC	TGCAGCCTGC
	CCCTGTGATC	CCTTCGGCGC	CAGCAGCTCC	AAAACITGGAC	CACCAGAGAA
35	TGGACACCAT	CCAGGAAGAC	CCCAGCACAG	ACTCACATAT	GGACGAGGAT
	GGGTTTGAGA	AGGACCCCTT	CCCAAAATAGC	AGCACAGCTG	CCAAGTCATT
	TGAGGATCTC	ACGGACCATC	CGGTCACCAG	AAGTGAAAAG	GCTGCCTCCT
	TTAAACTGCA	GCGTCAGAAT	CGTGTTGACA	GCAAAGAAAAC	AGAGTGCTAA
	TTTAGTTCTC	AGCTCTTCTG	ACTTAAGTGT	GCAAAATATT	TTTATAGATT
40	TGACCTACAA	ATCAATCACA	GCTTGATTTT	TGTGAAGACT	GGGAAGTGAC
	TTAGCAGATG	CTGGTCAIGT	GTTTGAACTT	CCTGCAGGTA	AACAGTTCTT
	GTGTGGTTTG	GCCCTTCTCC	TTTTGAAAAG	GTAAGGTGAA	GGTGAATCTA
	GCTTATTTTG	AGGCTTTCAG	GTTTTAGTTT	TTAAAATATC	TTTTGACCTG
	TGGTGCAAAA	GCAGAAAATA	CAGCTGGATT	GGGTTATGAA	TATTTACGTT
45	TTTGTAATTT	AATCTTTTAT	ATTGATAACA	GCACTGACTA	GGGAAATGAT
	CAGTTTTTTT	TTATACACTG	TAATGAACCG	CTGAATATGA	GGCATTTGGC
	ATTTATTTGT	GATGACAACT	GGAATAGTTT	TTTTTTTTTT	TTTTTTTTTT
	TGCCTTCAAC	TAAAAACAAA	GGAGATAAAT	CTAGTATACA	TTGTCTCTAA
	ATTGTGGGTC	TATTTCTAGT	TATTACCCAG	AGTTTTTATG	TAGCAGGGAA
50	AATATATATC	TAAATTTAGA	AATCATTTGG	GTTAATATGG	CTCTTCATAA
	TTCTAAGACT	AATGCTCTCT	AGAAAACCTAA	CCACCTACCT	TACAGTGAGG
	GCTATACATG	GTAGCCAGTT	GAATTTATGG	AATCTACCAA	CTGTTTAGGG

CCCTGATTG CTGGGCAGTT TTTCTGTATT TTATAAGTAT CTTTCATGTAT
 CCCTGTTACT GATAGGGATA CATGCTCTTA GAAAATTCAC TATTGGCTGG
 GAGTGGTGGC TCATGCCTGT AATCCCAGCA CTTGGAGAGG CTGAGGTTGC
 GCCACTACAC TCCAGCCTGG GTGACAGAGT GAGACTCTGC CTCAAAAAAA
 5 AAAAAAAAAA AAAAAAATTC ACTATCTACA AACCTAGAAT ATTTAAAATA
 CAAAGATTGC CTGTTTTCAA ACACTATTGA ATAAGAGGGT GAGATATTTT
 TTAACAACAA CAACAACAAA AAAAAACAGGT TGTTTTGAAT GTGATGAGCC
 AGCCAGGAGA TAGAATACTA CCTGCCCTTA GGGTTGGGGG CTGTCCCCAC
 AAGACTTGAT ACTTCAGAAA CCCTTTTTTAT TGACCCACAA GCAGATATTT
 10 GAATTACTTC TTTACTTTATT GCTCCAGGAT TCTGGATGGG CTGCATTTAC
 TGTGTGAAGG ATAAAAATCA TTAGCCTGGA TTCTGATTTC TATAAATTGC
 CATTAAAAGC TTTTTTTCCC CTAAGAACTG AAATGTGCTC ACCAGCCAAA
 ACATTTTAAC TTGTAAATTT TGAGGGCAGT TAACCAAACC TGTGACTAAT
 CATATCTCCT CCTACCCCCC ATTTCCAAGG ACATTTGTTA CTCAGATACT
 15 TGTTATACTA ATACTTGAAC TTGTACCTTA TGGTATTTGC TATCTTTTAA
 CTAGTCATGA TATTCTTATA CTTTAGTTAC ACTTTTGGAA TTTGATACAA
 GGTGAGTGG GGTGTGTGGG TGTATGTATG AGTGAAACAG TTCTCAAAG
 AATGTAAGAA AAACCATTTT TATAAAATTG TGACTTTTTA AAAACATAGT
 CTTTGTCAAT TATAGAATTA ACAAGCTGCT CAGGGTATAT TTTATAGCTG
 20 TAGCACTGAT ATCTGCATTA ATAAATACTG TCGAAACACA A
[0096] For example, the target mRNA may be an mRNA encoding isocitrate dehydrogenase (IDH1), cyclin-dependent kinase 4 (CDK4), CDK6, epidermal growth factor receptor (EGFR), mechanistic target of rapamycin (mTOR), Kirsten rat sarcoma viral oncogene (KRAS), cluster of differentiation (CD155), programmed cell death-ligand 1 (PD-L1), or myc
 25 proto-oncogene (c-Myc), a fragment thereof, or a functional variant thereof. A functional variant as used herein may refer to a full-length molecule, a fragment thereof, or a variant thereof. For example, a variant molecule may comprise a sequence modified by insertion, deletion, and/or substitution of one or more amino acids, in the case of protein sequence, or one or more nucleotides, in the case of nucleic acid sequence. For example, a variant
 30 molecule may comprise or encode a mutant protein, including, but not limited to, a gain-of-function or a loss-of-function mutant.
[0097] In some embodiments, the target mRNA may encode a protein selected from the group consisting of VEGFA, an isoform of VEGFA, PIGF, MICA, MICB, ERp5, ADAM17, MMP, IDH1, CDK4, CDK6, EGFR, mTOR, KRAS, CD155, PD-L1, c-Myc, a fragment thereof, a
 35 functional variant thereof, and a combination thereof. In some embodiments, VEGFA mRNA comprises a sequence comprising SEQ ID NO: 36. In some embodiments, MICA mRNA comprises a sequence comprising SEQ ID NO: 39. In some embodiments, MICB mRNA comprises a sequence comprising SEQ ID NO: 42. In some embodiments, IDH1 mRNA comprises a sequence comprising SEQ ID NO: 51. In some embodiments, CDK4 mRNA
 40 comprises a sequence comprising SEQ ID NO: 54. In some embodiments, CDK6 mRNA comprises a sequence comprising SEQ ID NO: 57. In some embodiments, EGFR mRNA

comprises a sequence comprising SEQ ID NO: 60. In some embodiments, mTOR mRNA comprises a sequence comprising SEQ ID NO: 63. In some embodiments, KRAS mRNA comprises a sequence comprising SEQ ID NO: 66. In some embodiments, CD155 mRNA comprises a sequence comprising SEQ ID NO: 72. In some embodiments, PD-L1 mRNA
 5 comprises a sequence comprising SEQ ID NO: 75. In some embodiments, c-Myc mRNA comprises a sequence comprising SEQ ID NO: 78.

Gene of interest

[0098] Provided herein are recombinant RNA constructs comprising one or more copies of nucleic acid sequence encoding a gene of interest. For example, recombinant RNA constructs
 10 may comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more copies of nucleic acid sequence encoding a gene of interest. In some instances, each of the 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more copies of nucleic acid sequence encoding a gene of interest encodes the same gene of interest. In some instances, recombinant RNA constructs may comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more copies of nucleic acid sequence encoding a cytokine.

[0099] Also provided herein are recombinant RNA constructs comprising two or more copies of nucleic acid sequence encoding a gene of interest, wherein each of the two or more nucleic acid sequence may encode a different gene of interest. In some cases, each of the two or more nucleic acid sequences encoding different gene of interest may comprise a nucleic acid sequence encoding a secretory protein. In some cases, each of the two or more nucleic acid
 20 sequences encoding different gene of interest may comprise a nucleic acid sequence encoding a cytokine. In some embodiments, each of the two or more nucleic acid sequences encoding different gene of interest may encode a different cytokine. Further provided herein are recombinant RNA constructs comprising a linker. In some embodiments, the linker may connect each of the two or more nucleic acid sequences encoding a gene of interest. In some
 25 cases, the linker may be a non-cleavable linker. In some cases, the linker may be a cleavable linker. In some cases, the linker may be a self-cleavable linker. Non-limiting examples of the linker comprises a flexible linker, a 2A peptide linker (or 2A self-cleaving peptides) such as T2A, P2A, E2A, or F2A, and a tRNA linker, etc. The tRNA system is evolutionarily conserved across living organism and utilizes endogenous RNases P and Z to process multicistronic constructs (Dong et al., 2016). In some embodiments, the tRNA linker may comprise a nucleic acid sequence comprising
 30 AACAAAGCACCAGTGGTCTAGTGGTAGAATAGTACCCTGCCACGGTACAGACCC
 GGGTTCGATTCCCGGCTGGTGCA (SEQ ID NO: 20).

- [0100] Provided herein are recombinant RNA constructs comprising an RNA encoding for a gene of interest for modulating the expression of the gene of interest. For example, expression of a protein encoded by the mRNA of the gene of interest can be modulated. For example, the expression of the gene of interest is upregulated by expressing a protein encoded by mRNA of the gene of interest in recombinant RNA constructs. For example, the expression of the gene of interest is upregulated by increasing the level of protein encoded by mRNA of the gene of interest in recombinant RNA constructs. The level of protein expression can be measured by using any methods well known in the art and these include, but are not limited to Western-blotting, flow cytometry, ELISAs, RIAs, and various proteomics techniques. An exemplary method to measure or detect a polypeptide is an immunoassay, such as an ELISA. This type of protein quantitation can be based on an antibody capable of capturing a specific antigen, and a second antibody capable of detecting the captured antigen. Exemplary assays for detection and/or measurement of polypeptides are described in Harlow, E. and Lane, D. Antibodies: A Laboratory Manual, (1988), Cold Spring Harbor Laboratory Press.
- 15 [0101] Provided herein are recombinant RNA constructs comprising an RNA encoding for a gene of interest wherein the gene of the interest encodes a protein of interest. In some instances, the protein of interest is a therapeutic protein. In some instances, the protein of interest is of human origin i.e., is a human protein. In some instances, the gene of interest encodes a cytokine. In some embodiments, the cytokine comprises an interleukin. In some
20 embodiments, the protein of interest is an interleukin 2 (IL-2), IL-12, IL-15, IL-7, a fragment thereof, or a functional variant thereof. A functional variant as used herein may refer to a full-length molecule, a fragment thereof, or a variant thereof. For example, a variant molecule may comprise a sequence modified by insertion, deletion, and/or substitution of one or more amino acids, in the case of protein sequence, or one or more nucleotides, in the case of nucleic
25 acid sequence.
- [0102] In some instances, interleukin 2 (IL-2) or IL-2 as used herein may refer to the natural sequence of human IL-2 (Uniprot database: P60568 or Q0GK43 and in the Genbank database: NM_000586.3), a fragment thereof, or a functional variant thereof. The natural DNA sequence encoding human IL-2 may be codon-optimized. The natural sequence of human IL-2 may
30 consist of a signal peptide having 20 amino acids (nucleotides 1-60) and the mature human IL-2 having 133 amino acids (nucleotides 61-459) as shown in SEQ ID NO: 23. In some embodiments, the signal peptide is unmodified IL-2 signal peptide. In some embodiments, the signal peptide is IL-2 signal peptide modified by insertion, deletion, and/or substitution of at least one amino acid. In some embodiments, interleukin 2 (IL-2) or IL-2 as used herein may

refer to the mature human IL-2. In some embodiments, a mature protein can refer to a protein synthesized in the endoplasmic reticulum and secreted via the Golgi apparatus in a cell expressing and secreting the protein. In some embodiments, a mature IL-2 may refer to an IL-2 protein synthesized in the endoplasmic reticulum and secreted via the Golgi apparatus in a cell expressing and secreting IL-2. In some embodiments, a mature human IL-2 may refer to an IL-2 protein synthesized in the endoplasmic reticulum and secreted via the Golgi apparatus in a human cell expressing and secreting human IL-2 and normally contains the amino acids encoded by nucleotide as shown in SEQ ID NO: 24. In some embodiments, IL-2 may comprise an IL-2 fragment, an IL-2 variant, an IL-2 mutein, or an IL-2 mutant. In some embodiments, the IL-2 fragment described herein may be at least partially functional, i.e., can perform an IL-2 activity at a similar or lower level compared to a wildtype or a full length IL-2. In some embodiments, the IL-2 fragment described herein may be fully functional, i.e., can perform an IL-2 activity at the same level compared to a wildtype or a full length IL-2. In some embodiments, the IL-2 variant, an IL-2 mutein, or the IL-2 mutant may comprise an IL-2 amino acid sequence modified by insertion, deletion, and/or substitution of at least one amino acid. In some embodiments, the IL-2 variant, an IL-2 mutein, or the IL-2 mutant may be at least partially functional, i.e., can perform an IL-2 activity at a similar or lower level compared to a wildtype IL-2. In some embodiments, the IL-2 variant, an IL-2 mutein, or the IL-2 mutant may be fully functional, i.e., can perform an IL-2 activity at the same level compared to a wildtype IL-2. In some embodiments, the IL-2 variant, an IL-2 mutein, or the IL-2 mutant may perform an IL-2 activity at a higher level compared to a wildtype IL-2.

[0103] The mRNA encoding IL-2 may refer to an mRNA comprising a nucleotide sequence encoding the propeptide of human IL-2 having 153 amino acids or a nucleotide sequence encoding the mature human IL-2 having 133 amino acids. The nucleotide sequence encoding the propeptide of human IL-2 and the nucleotide sequence encoding the mature human IL-2 may be codon-optimized. In some instances, recombinant RNA constructs, provided herein, may comprise 1 copy of IL-2 mRNA. In some instances, recombinant RNA constructs, provided herein, may comprise 2 or more copies of IL-2 mRNA.

[0104] In some instances, interleukin 12 (IL-12) or IL-12 as used herein may refer to the natural sequence of human IL-12 alpha (Genbank database: NM_000882.4), the natural sequence of human IL-12 beta (Genbank database: NM_002187.2), a fragment thereof, or a functional variant thereof. The natural DNA sequence encoding human IL-12 may be codon-optimized. The natural sequence of human IL-12 alpha may consist of a signal peptide having 22 amino acids and the mature human IL-12 having 197 amino acids as shown in SEQ ID NO:

43. In some embodiments, the signal peptide is unmodified IL-12 alpha signal peptide. In some embodiments, the signal peptide is IL-12 alpha signal peptide modified by insertion, deletion, and/or substitution of at least one amino acid. The natural sequence of human IL-12 beta may consist of a signal peptide having 22 amino acids and the mature human IL-12 having 306 amino acids as shown in SEQ ID NO: 46. In some embodiments, the signal peptide is unmodified IL-12 beta signal peptide. In some embodiments, the signal peptide is IL-12 beta signal peptide modified by insertion, deletion, and/or substitution of at least one amino acid.

[0105] In some embodiments, interleukin 12 (IL-12) or IL-12 as used herein may refer to the mature human IL-12 alpha. In some embodiments, interleukin 12 (IL-12) or IL-12 as used herein may refer to the mature human IL-12 beta. In some embodiments, a mature protein can refer to a protein synthesized in the endoplasmic reticulum and secreted via the Golgi apparatus in a cell expressing and secreting the protein. In some embodiments, a mature IL-12 may refer to an IL-12 alpha protein synthesized in the endoplasmic reticulum and secreted via the Golgi apparatus in a cell expressing and secreting IL-12. In some embodiments, a mature IL-12 may refer to an IL-12 beta protein synthesized in the endoplasmic reticulum and secreted via the Golgi apparatus in a cell expressing and secreting IL-12. In some embodiments, a mature human IL-12 may refer to an IL-12 alpha protein synthesized in the endoplasmic reticulum and secreted via the Golgi apparatus in a human cell expressing and secreting human IL-12 and normally contains the amino acids encoded by nucleotide as shown in SEQ ID NO: 44. In some embodiments, a mature human IL-12 may refer to an IL-12 beta protein synthesized in the endoplasmic reticulum and secreted via the Golgi apparatus in a human cell expressing and secreting human IL-12 and normally contains the amino acids encoded by nucleotide as shown in SEQ ID NO: 47.

[0106] In some embodiments, IL-12 alpha may comprise an IL-12 alpha fragment, an IL-12 alpha variant, an IL-12 alpha mutein, or an IL-12 alpha mutant. In some embodiments, the IL-12 alpha fragment described herein may be at least partially functional, i.e., can perform an IL-12 alpha activity at a similar or lower level compared to a wildtype or a full-length IL-12 alpha. In some embodiments, the IL-12 alpha fragment described herein may be fully functional, i.e., can perform an IL-12 alpha activity at the same level compared to a wildtype or a full-length IL-12 alpha. In some embodiments, the IL-12 alpha variant, an IL-12 alpha mutein, or the IL-12 alpha mutant may comprise an IL-12 alpha amino acid sequence modified by insertion, deletion, and/or substitution of at least one amino acid. In some embodiments, the IL-12 alpha variant, an IL-12 alpha mutein, or the IL-12 alpha mutant may be at least partially functional, i.e., can perform an IL-12 alpha activity at a similar or lower level compared to a

wildtype IL-12 alpha. In some embodiments, the IL-12 alpha variant, an IL-12 alpha mutein, or the IL-12 alpha mutant may be fully functional, i.e., can perform an IL-12 alpha activity at the same level compared to a wildtype IL-12 alpha. In some embodiments, the IL-12 alpha variant, an IL-12 alpha mutein, or the IL-12 alpha mutant may perform an IL-12 alpha activity at a higher level compared to a wildtype IL-12 alpha.

[0107] In some embodiments, IL-12 beta may comprise an IL-12 beta fragment, an IL-12 beta variant, an IL-12 beta mutein, or an IL-12 beta mutant. In some embodiments, the IL-12 beta fragment described herein may be at least partially functional, i.e., can perform an IL-12 beta activity at a similar or lower level compared to a wildtype or a full-length IL-12 beta. In some embodiments, the IL-12 beta fragment described herein may be fully functional, i.e., can perform an IL-12 beta activity at the same level compared to a wildtype or a full-length IL-12 beta. In some embodiments, the IL-12 beta variant, an IL-12 beta mutein, or the IL-12 beta mutant may comprise an IL-12 beta amino acid sequence modified by insertion, deletion, and/or substitution of at least one amino acid. In some embodiments, the IL-12 beta variant, an IL-12 beta mutein, or the IL-12 beta mutant may be at least partially functional, i.e., can perform an IL-12 beta activity at a similar or lower level compared to a wildtype IL-12 beta. In some embodiments, the IL-12 beta variant, an IL-12 beta mutein, or the IL-12 beta mutant may be fully functional, i.e., can perform an IL-12 beta activity at the same level compared to a wildtype IL-12 beta. In some embodiments, the IL-12 beta variant, an IL-12 beta mutein, or the IL-12 beta mutant may perform an IL-12 beta activity at a higher level compared to a wildtype IL-12 beta.

[0108] The mRNA encoding IL-12 may refer to an mRNA comprising a nucleotide sequence encoding the propeptide of human IL-12 alpha having 219 amino acids or a nucleotide sequence encoding the mature human IL-12 alpha having 197 amino acids. The nucleotide sequence encoding the propeptide of human IL-12 alpha and the nucleotide sequence encoding the mature human IL-12 may be codon-optimized. The mRNA encoding IL-12 may refer to an mRNA comprising a nucleotide sequence encoding the propeptide of human IL-12 beta having 328 amino acids or a nucleotide sequence encoding the mature human IL-12 beta having 306 amino acids. The nucleotide sequence encoding the propeptide of human IL-12 beta and the nucleotide sequence encoding the mature human IL-12 may be codon-optimized. In some instances, recombinant RNA constructs, provided herein, may comprise 1 copy of IL-12 mRNA. In some instances, recombinant RNA constructs, provided herein, may comprise 2 or more copies of IL-12 mRNA.

[0109] In some instances, interleukin 15 (IL-15) or IL-15 as used herein may refer to the natural sequence of human IL-15 (Genbank database: NM_000585.4), a fragment thereof, or a functional variant thereof. The natural DNA sequence encoding human IL-15 may be codon-optimized. The natural sequence of human IL-15 may consist of a signal peptide having 29 amino acids and the mature human IL-15 having 133 amino acids as shown in SEQ ID NO: 67. In some embodiments, the signal peptide is unmodified IL-15 signal peptide. In some embodiments, the signal peptide is IL-15 signal peptide modified by insertion, deletion, and/or substitution of at least one amino acid. In some embodiments, interleukin 15 (IL-15) or IL-15 as used herein may refer to the mature human IL-15. In some embodiments, a mature protein can refer to a protein synthesized in the endoplasmic reticulum and secreted via the Golgi apparatus in a cell expressing and secreting the protein. In some embodiments, a mature IL-15 may refer to an IL-15 protein synthesized in the endoplasmic reticulum and secreted via the Golgi apparatus in a cell expressing and secreting IL-15. In some embodiments, a mature human IL-15 may refer to an IL-15 protein synthesized in the endoplasmic reticulum and secreted via the Golgi apparatus in a human cell expressing and secreting human IL-15 and normally contains the amino acids encoded by nucleotide as shown in SEQ ID NO: 68. In some embodiments, IL-15 may comprise an IL-15 fragment, an IL-15 variant, an IL-15 mutein, or an IL-15 mutant. In some embodiments, the IL-15 fragment described herein may be at least partially functional, i.e., can perform an IL-15 activity at a similar or lower level compared to a wildtype or a full-length IL-15. In some embodiments, the IL-15 fragment described herein may be fully functional, i.e., can perform an IL-15 activity at the same level compared to a wildtype or a full-length IL-15. In some embodiments, the IL-15 variant, an IL-15 mutein, or the IL-15 mutant may comprise an IL-15 amino acid sequence modified by insertion, deletion, and/or substitution of at least one amino acid. In some embodiments, the IL-15 variant, an IL-15 mutein, or the IL-15 mutant may be at least partially functional, i.e., can perform an IL-15 activity at a similar or lower level compared to a wildtype IL-15. In some embodiments, the IL-15 variant, an IL-15 mutein, or the IL-15 mutant may be fully functional, i.e., can perform an IL-15 activity at the same level compared to a wildtype IL-15. In some embodiments, the IL-15 variant, an IL-15 mutein, or the IL-15 mutant may perform an IL-15 activity at a higher level compared to a wildtype IL-15.

[0110] The mRNA encoding IL-15 may refer to an mRNA comprising a nucleotide sequence encoding the propeptide of human IL-15 having 162 amino acids or a nucleotide sequence encoding the mature human IL-15 having 133 amino acids. The nucleotide sequence encoding the propeptide of human IL-15 and the nucleotide sequence encoding the mature human IL-15

may be codon-optimized. In some instances, recombinant RNA constructs, provided herein, may comprise 1 copy of IL-15 mRNA. In some instances, recombinant RNA constructs, provided herein, may comprise 2 or more copies of IL-15 mRNA.

[0111] In some instances, interleukin 7 (IL-7) or IL-7 as used herein may refer to the natural sequence of human IL-7 (Genbank database: NM_000880.3), a fragment thereof, or a functional variant thereof. The natural DNA sequence encoding human IL-7 may be codon-optimized. The natural sequence of human IL-7 may consist of a signal peptide having 25 amino acids and the mature human IL-7 having 152 amino acids as shown in SEQ ID NO: 79. In some embodiments, the signal peptide is unmodified IL-7 signal peptide. In some
10 embodiments, the signal peptide is IL-7 signal peptide modified by insertion, deletion, and/or substitution of at least one amino acid. In some embodiments, interleukin 7 (IL-7) or IL-7 as used herein may refer to the mature human IL-7. In some embodiments, a mature protein can refer to a protein synthesized in the endoplasmic reticulum and secreted via the Golgi apparatus in a cell expressing and secreting the protein. In some embodiments, a mature IL-7
15 may refer to an IL-7 protein synthesized in the endoplasmic reticulum and secreted via the Golgi apparatus in a cell expressing and secreting IL-7. In some embodiments, a mature human IL-7 may refer to an IL-7 protein synthesized in the endoplasmic reticulum and secreted via the Golgi apparatus in a human cell expressing and secreting human IL-7 and normally contains the amino acids encoded by nucleotide as shown in SEQ ID NO: 80. In some embodiments,
20 IL-7 may comprise an IL-7 fragment, an IL-7 variant, an IL-7 mutein, or an IL-7 mutant. In some embodiments, the IL-7 fragment described herein may be at least partially functional, i.e., can perform an IL-7 activity at a similar or lower level compared to a wildtype or a full-length IL-7. In some embodiments, the IL-7 fragment described herein may be fully functional, i.e., can perform an IL-7 activity at the same level compared to a wildtype or a full-length IL-7.
25 In some embodiments, the IL-7 variant, an IL-7 mutein, or the IL-7 mutant may comprise an IL-7 amino acid sequence modified by insertion, deletion, and/or substitution of at least one amino acid. In some embodiments, the IL-7 variant, an IL-7 mutein, or the IL-7 mutant may be at least partially functional, i.e., can perform an IL-7 activity at a similar or lower level compared to a wildtype IL-7. In some embodiments, the IL-7 variant, an IL-7 mutein, or the
30 IL-7 mutant may be fully functional, i.e., can perform an IL-7 activity at the same level compared to a wildtype IL-7. In some embodiments, the IL-7 variant, an IL-7 mutein, or the IL-7 mutant may perform an IL-7 activity at a higher level compared to a wildtype IL-7.

[0112] The mRNA encoding IL-7 may refer to an mRNA comprising a nucleotide sequence encoding the propeptide of human IL-7 having 177 amino acids or a nucleotide sequence

encoding the mature human IL-7 having 152 amino acids. The nucleotide sequence encoding the propeptide of human IL-7 and the nucleotide sequence encoding the mature human IL-7 may be codon-optimized. In some instances, recombinant RNA constructs, provided herein, may comprise 1 copy of IL-7 mRNA. In some instances, recombinant RNA constructs, provided herein, may comprise 2 or more copies of IL-7 mRNA.

Target Motif

[0113] Provided herein are compositions comprising recombinant RNA constructs

comprising a target motif. A target motif or a targeting motif as used herein can refer to any short peptide present in the newly synthesized polypeptides or proteins that are destined to

any parts of cell membranes, extracellular compartments, or intracellular compartments, except cytoplasm or cytosol. In some embodiments, a peptide may refer to a series of amino acid residues connected one to the other, typically by peptide bonds between the α -amino and carboxyl groups of adjacent amino acid residues. Intracellular compartments include, but are not limited to, intracellular organelles such as nucleus, nucleolus, endosome, proteasome,

ribosome, chromatin, nuclear envelope, nuclear pore, exosome, melanosome, Golgi apparatus, peroxisome, endoplasmic reticulum (ER), lysosome, centrosome, microtubule, mitochondria, chloroplast, microfilament, intermediate filament, or plasma membrane. In some

embodiments, a signal peptide can be referred to as a signal sequence, a targeting signal, a localization signal, a localization sequence, a transit peptide, a leader sequence, or a leader

peptide. In some embodiments, a target motif is operably linked to a nucleic acid sequence encoding a gene of interest. In some embodiments, the term "operably linked" can refer to a functional relationship between two or more nucleic acid sequences, *e.g.*, a functional relationship of a transcriptional regulatory or signal sequence to a transcribed sequence. For example, a target motif or a nucleic acid encoding a target motif is operably linked to a coding

sequence if it is expressed as a preprotein that participates in targeting the polypeptide

encoded by the coding sequence to a cell membrane, intracellular, or an extracellular

compartment. For example, a signal peptide or a nucleic acid encoding a signal peptide is

operably linked to a coding sequence if it is expressed as a preprotein that participates in the secretion of the polypeptide encoded by the coding sequence. For example, a promoter is

operably linked if it stimulates or modulates the transcription of the coding sequence. Non-

limiting examples of a target motif comprise a signal peptide, a nuclear localization signal (NLS), a nucleolar localization signal (NoLS), a lysosomal targeting signal, a mitochondrial targeting signal, a peroxisomal targeting signal, a microtubule tip localization signal (MtLS), an endosomal targeting signal, a chloroplast targeting signal, a Golgi targeting signal, an

endoplasmic reticulum (ER) targeting signal, a proteasomal targeting signal, a membrane targeting signal, a transmembrane targeting signal, a centrosomal localization signal (CLS) or any other signal that targets a protein to a certain part of cell membrane, extracellular compartments, or intracellular compartments.

- 5 **[0114]** A signal peptide is a short peptide present at the N-terminus of newly synthesized proteins that are destined towards the secretory pathway. The signal peptide of the present invention can be 10-40 amino acids long. A signal peptide can be situated at the N-terminal end of the protein of interest or at the N-terminal end of a pro-protein form of the protein of interest. A signal peptide may be of eukaryotic origin. In some embodiments, a signal peptide
- 10 may be a mammalian protein. In some embodiments, a signal peptide may be a human protein. In some instances, a signal peptide may be a homologous signal peptide (i.e. from the same protein) or a heterologous signal peptide (i.e. from a different protein or a synthetic signal peptide). In some instances, a signal peptide may be a naturally occurring signal peptide of a protein or a modified signal peptide.
- 15 **[0115]** Provided herein are compositions comprising recombinant RNA constructs comprising a target motif, wherein the target motif may be selected from the group consisting of (a) a target motif heterologous to a protein encoded by the gene of interest; (b) a target motif heterologous to a protein encoded by the gene of interest, wherein the target motif heterologous to the protein encoded by the gene of interest is modified by insertion, deletion,
- 20 and/or substitution of at least one amino acid; (c) a target motif homologous to a protein encoded by the gene of interest; (d) a target motif homologous to a protein encoded by the gene of interest, wherein the target motif homologous to the protein encoded by the gene of interest is modified by insertion, deletion, and/or substitution of at least one amino acid; and
- 25 (e) a naturally occurring amino acid sequence which does not have the function of a target motif in nature, wherein the naturally occurring amino acid sequence is optionally modified by insertion, deletion, and/or substitution of at least one amino acid.
- 30 **[0116]** Provided herein are compositions comprising recombinant RNA constructs comprising a target motif, wherein the target motif is a signal peptide. In some embodiments, the signal peptide is selected from the group consisting of: (a) a signal peptide heterologous to a protein encoded by the gene of interest; (b) a signal peptide heterologous to a protein encoded by the gene of interest, wherein the signal peptide heterologous to the protein encoded by the gene of interest is modified by insertion, deletion, and/or substitution of at least one amino acid, with proviso that the protein is not an oxidoreductase; (c) a signal peptide homologous to a protein encoded by the gene of interest; (d) a signal peptide

homologous to a protein encoded by the gene of interest, wherein the signal peptide homologous to the protein encoded by the gene of interest is modified by insertion, deletion, and/or substitution of at least one amino acid; and (e) a naturally occurring amino acid sequence which does not have the function of a signal peptide in nature, wherein the naturally occurring amino acid sequence is optionally modified by insertion, deletion, and/or substitution of at least one amino acid. In some instances, the amino acids 1-9 of the N-terminal end of the signal peptide have an average hydrophobic score of above 2.

[0117] In some instances, a target motif heterologous to a protein encoded by the gene of interest or a signal peptide heterologous to a protein encoded by the gene of interest as used herein can refer to a naturally occurring target motif or signal peptide which is different from the naturally occurring target motif or signal peptide of a protein. For example, the target motif or the signal peptide is not derived from the gene of interest. Usually a target motif or a signal peptide heterologous to a given protein is a target motif or a signal peptide from another protein, which is not related to the given protein. For example, a target motif or a signal peptide heterologous to a given protein has an amino acid sequence that is different from the amino acid sequence of the target motif or the signal peptide of the given protein by more than 50%, 60%, 70%, 80%, 90%, or by more than 95%. Although heterologous sequences may be derived from the same organism, they naturally (in nature) do not occur in the same nucleic acid molecule, such as in the same mRNA. The target motif or the signal peptide heterologous to a protein and the protein to which the target motif or the signal peptide is heterologous can be of the same or different origin. In some embodiments, they are of eukaryotic origin. In some embodiments, they are of the same eukaryotic organism. In some embodiments, they are of mammalian origin. In some embodiments, they are of the same mammalian organism. In some embodiments, they are human origin. For example, an RNA construct may comprise a nucleic acid sequence encoding the human IL-2 gene and a signal peptide of another human cytokine. In some embodiments, an RNA construct may comprise a signal peptide heterologous to a protein wherein the signal peptide and the protein are of the same origin, namely of human origin.

[0118] In some instance, a target motif homologous to a protein encoded by the gene of interest or a signal peptide homologous to a protein encoded by the gene of interest as used herein can refer to a naturally occurring target motif or signal peptide of a protein. A target motif or a signal peptide homologous to a protein is the target motif or the signal peptide encoded by the gene of the protein as it occurs in nature. A target motif or a signal peptide homologous to a protein is usually of eukaryotic origin. In some embodiments, a target motif

or a signal peptide homologous to a protein is of mammalian origin. In some embodiments, a target motif or a signal peptide homologous to a protein is of human origin.

[0119] In some instances, a naturally occurring amino acid sequence which does not have the function of a target motif in nature or a naturally occurring amino acid sequence which does not have the function of a signal peptide in nature as used herein can refer to an amino acid sequence which occurs in nature and is not identical to the amino acid sequence of any target motif or signal peptide occurring in nature. A naturally occurring amino acid sequence which does not have the function of a target motif or a signal peptide in nature can be between 10-50 amino acids long. In some embodiments, a naturally occurring amino acid sequence which does not have the function of a target motif or a signal peptide in nature is of eukaryotic origin and not identical to any target motif or signal peptide of eukaryotic origin. In some embodiments, a naturally occurring amino acid sequence which does not have the function of a target motif or a signal peptide in nature is of mammalian origin and not identical to any target motif or signal peptide of mammalian origin. In some embodiments, a naturally occurring amino acid sequence which does not have the function of a target motif or a signal peptide in nature is of human origin and not identical to any target motif or signal peptide of human origin occurring in nature. A naturally occurring amino acid sequence which does not have the function of a target motif or a signal peptide in nature is usually an amino acid sequence of the coding sequence of a protein. The terms “naturally occurring,” “natural,” and “in nature” as used herein have the equivalent meaning.

[0120] In some instances, amino acids 1-9 of the N-terminal end of the signal peptide as used herein can refer to the first nine amino acids of the N-terminal end of the amino acid sequence of a signal peptide. Analogously, amino acids 1-7 of the N-terminal end of the signal peptide as used herein can refer to the first seven amino acids of the N-terminal end of the amino acid sequence of a signal peptide and amino acids 1-5 of the N-terminal end of the signal peptide can refer to the first five amino acids of the N-terminal end of the amino acid sequence of a signal peptide.

[0121] In some instances, amino acid sequence modified by insertion, deletion, and/or substitution of at least one amino acid can refer to an amino acid sequence which includes an amino acid substitution, insertion, and/or deletion of at least one amino acid within the amino acid sequence. For example, target motif heterologous to a protein encoded by the gene of interest is modified by insertion, deletion, and/or substitution of at least one amino acid or signal peptide heterologous to a protein encoded by the gene of interest is modified by insertion, deletion, and/or substitution of at least one amino acid as used herein can refer to an

amino acid sequence of a naturally occurring target motif or signal peptide heterologous to a protein which includes an amino acid substitution, insertion, and/or deletion of at least one amino acid within its naturally occurring amino acid sequence. For example, target motif homologous to a protein encoded by the gene of interest is modified by insertion, deletion, and/or substitution of at least one amino acid or signal peptide homologous to a protein encoded by the gene of interest is modified by insertion, deletion, and/or substitution of at least one amino acid as used herein can refer to a naturally occurring target motif or signal peptide homologous to a protein which includes an amino acid substitution, insertion, and/or deletion of at least one amino acid within its naturally occurring amino acid sequence. In some embodiments, naturally occurring amino acid sequence may be modified by insertion, deletion, and/or substitution of at least one amino acid and a naturally occurring amino acid sequence can include an amino acid substitution, insertion, and/or deletion of at least one amino acid within its naturally occurring amino acid sequence. An amino acid substitution or a substitution may refer to replacement of an amino acid at a particular position in an amino acid or polypeptide sequence with another amino acid. For example, the substitution R34K refers to a polypeptide, in which the arginine (Arg or R) at position 34 is replaced with a lysine (Lys or K). For the preceding example, 34K indicates the substitution of an amino acid at position 34 with a lysine (Lys or K). In some embodiments, multiple substitutions are typically separated by a slash. For example, R34K/L38V refers to a variant comprising the substitutions R34K and L38V. An amino acid insertion or an insertion may refer to addition of an amino acid at a particular position in an amino acid or polypeptide sequence. For example, insert -34 designates an insertion at position 34. An amino acid deletion or a deletion may refer to removal of an amino acid at a particular position in an amino acid or polypeptide sequence. For example, R34- designates the deletion of arginine (Arg or R) at position 34.

[0122] In some instances, deleted amino acid is an amino acid with a hydrophobic score of below -0.8, -0.7, -0.6, -0.5, -0.4, -0.3, -0.2, -0.1, 0, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, or below 1.9. In some instances, the substitute amino acid is an amino acid with a hydrophobic score which is higher than the hydrophobic score of the substituted amino acid. For example, the substitute amino acid is an amino acid with a hydrophobic score of 2.8 and higher, or 3.8 and higher. In some instances, the inserted amino acid is an amino acid with a hydrophobic score of 2.8 and higher or 3.8 and higher.

[0123] In some instances, an amino acid sequence described herein may comprise 1 to 15 amino acid insertions, deletions, and/or substitutions. In some embodiments, an amino acid

sequence described herein may comprise 1 to 7 amino acid insertions, deletions, and/or substitutions. In some instances, an amino acid sequence described herein may not comprise amino acid insertions, deletions, and/or substitutions. In some instances, an amino acid sequence described herein may comprise 1 to 15 amino acid insertions, deletions, and/or substitutions within the amino acids 1-30 of the N-terminal end of the amino acid sequence of the target motif or the signal peptide. In some embodiments, an amino acid sequence described herein may comprise 1 to 9 amino acid insertions, deletions, and/or substitutions within the amino acids 1-30 of the N-terminal end of the amino acid sequence of the target motif or the signal peptide. In some instances, an amino acid sequence described herein may comprise 1 to 15 amino acid insertions, deletions, and/or substitutions within the amino acids 1-20 of the N-terminal end of the amino acid sequence of the target motif or the signal peptide. In some embodiments, an amino acid sequence described herein may comprise 1 to 9 amino acid insertions, deletions, and/or substitutions within the amino acids 1-20 of the N-terminal end of the amino acid sequence of the target motif or the signal peptide. In some instances, at least one amino acid of an amino acid sequence described herein may be optionally modified by deletion, and/or substitution.

[0124] In some instances, the average hydrophobic score of the first nine amino acids of the N-terminal end of the amino acid sequence of the modified signal peptide is increased 1.0 unit or above compared to the signal peptide without modification. In some instances, hydrophobic score or hydrophobicity score can be used synonymously to hydrophathy score herein and can refer to the degree of hydrophobicity of an amino acid as calculated according to the Kyte-Doolittle scale (Kyte J., Doolittle R.F.; J. Mol. Biol. 157:105-132(1982)). The amino acid hydrophobic scores according to the Kyte-Doolittle scale are as follows:

[0125] Table B. Amino Acid Hydrophobic Scores

Amino Acid	One Letter Code	Hydrophobic Score
Isoleucine	I	4.5
Valine	V	4.2
Leucine	L	3.8
Phenylalanine	F	2.8
Cysteine	C	2.5
Methionine	M	1.9
Alanine	A	1.8
Glycine	G	-0.4
Threonine	T	-0.7
Serine	S	-0.8
Tryptophan	W	-0.9
Tyrosine	Y	-1.3
Proline	P	-1.6

Amino Acid	One Letter Code	Hydrophobic Score
Histidine	H	-3.2
Glutamic acid	E	-3.5
Glutamine	Q	-3.5
Aspartic acid	D	-3.5
Asparagine	N	-3.5
Lysine	K	-3.9
Arginine	R	-4.5

[0126] In some instances, average hydrophobic score of an amino acid sequence can be calculated by adding the hydrophobic score according to the Kyte-Doolittle scale of each of the amino acid of the amino acid sequence divided by the number of the amino acids. For example, the average hydrophobic score of the amino acids 1-9 of the N-terminal end of the amino acid sequence of a signal peptide can be calculated by adding the hydrophobic score or each of the nine amino acids divided by nine.

[0127] The polarity is calculated according to Zimmerman Polarity index (Zimmerman J.M., Eliezer N., Simha R.; J. Theor. Biol. 21:170-201(1968)). In some embodiments, average polarity of an amino acid sequence can be calculated by adding the polarity value calculated according to Zimmerman Polarity index of each of the amino acid of the amino acid sequence divided by the number of the amino acids. For example, the average polarity of the amino acids 1-9 of the N-terminal end of the amino acid sequence of a signal peptide can be calculated by adding the average polarity of each of the nine amino acids of the amino acids 1-9 of the N-terminal end, divided by nine. The polarity of amino acids according to Zimmerman Polarity index is as follows:

[0128] Table C. Amino Acid Polarity

Amino Acid	One Letter Code	Polarity
Isoleucine	I	0.13
Valine	V	0.13
Leucine	L	0.13
Phenylalanine	F	0.35
Cysteine	C	1.48
Methionine	M	1.43
Alanine	A	0
Glycine	G	0
Threonine	T	1.66
Serine	S	1.67
Tryptophan	W	2.1
Tyrosine	Y	1.61
Proline	P	1.58
Histidine	H	51.6
Glutamic acid	E	49.9
Glutamine	Q	3.53

Amino Acid	One Letter Code	Polarity
Aspartic acid	D	49.7
Asparagine	N	3.38
Lysine	K	49.5
Arginine	R	52

[0129] In some instances, a naturally occurring signal peptide of interleukin 2 (IL-2) may be modified by one or more substitutions, deletions, and/or insertions, wherein the naturally occurring signal peptide of IL-2 is referred to the amino acids 1-20 of the IL-2 amino acid sequence in the Uniprot database as P60568 or Q0GK43 and in the Genbank database as NM_000586.3. In some instances, the amino acid sequence of IL-2 signal peptide may be modified by the one or more substitutions, deletions, and/or insertions selected from the group consisting of Y2L, R3K, R3-, M4L, Q5L, S8L, S8A, -13A, L14T, L16A, V17-, and V17A. In some instances, the wild type (WT) IL-2 signal peptide amino acid sequence comprises a sequence comprising SEQ ID NO: 26. In some instances, a modified IL-2 signal peptide has an amino acid sequence comprising a sequence selected from the group consisting of SEQ ID NOs: 27-29. In some instances, a modified IL-2 signal peptide is encoded by a DNA sequence selected from the group consisting of SEQ ID NOs: 31-33.

15 Expression vector and production of RNA constructs

[0130] Provided herein are compositions comprising recombinant polynucleic acid constructs encoding recombinant RNA constructs comprising: (i) an mRNA encoding a gene of interest; and (ii) at least one siRNA capable of binding to a target mRNA. For example, an mRNA encoding a gene of interest can be IL-2, IL-12, IL-15, IL-7, a fragment thereof, or a functional variant thereof. For example, a target mRNA can be VEGF, VEGFA, an isoform of VEGFA, PIGF, MICA, MICB, ERp5, ADAM, MMP, IDH1, CDK4, CDK6, EGFR, mTOR, KRAS, CD155, PD-L1, or c-Myc. In some embodiments, the ADAM is ADAM17. Further provided herein are compositions comprising recombinant polynucleic acid constructs encoding RNA constructs described herein, *e.g.*, an RNA construct comprising a first RNA encoding for a cytokine linked to a second RNA encoding for a genetic element that can reduce expression of a gene associated with tumor proliferation, angiogenesis, or recognition by the immune system. For example, a cytokine can be IL-2, IL-12, IL-15, IL-7, a fragment thereof, or a functional variant thereof. For example, a gene associated with tumor proliferation or angiogenesis can be VEGF, VEGFA, an isoform of VEGFA, PIGF, IDH1, CDK4, CDK6, EGFR, mTOR, KRAS, CD155, PD-L1, c-Myc, a fragment thereof, or a functional variant thereof. Non-limiting examples of an isoform of VEGFA include VEGF111, VEGF121,

VEGF145, VEGF148, VEGF165, VEGF165B, VEGF183, VEGF189, VEGF206, L-VEGF121, L-VEGF165, L-VEGF189, L-VEGF206, Isoform 15, Isoform16, Isoform 17, and Isoform 18. For example, a gene associated with recognition by the immune system can be MICA, MICB, ERp5, ADAM, MMP, a fragment thereof, or a functional variant thereof. In some embodiments, the ADAM is ADAM17. In related aspects, recombinant polynucleic acid constructs encoding recombinant RNA constructs may encode 1, 2, 3, 4, 5, or more siRNA species. In related aspects, recombinant polynucleic acid constructs encoding recombinant RNA constructs may encode 1 siRNA species directed to a target mRNA. In related aspects, recombinant polynucleic acid constructs encoding recombinant RNA constructs may encode 3 siRNAs, each directed to a target mRNA. In related aspects, each of the siRNA species may comprise the same sequence, different sequence, or a combination thereof. For example, recombinant polynucleic acid constructs encoding recombinant RNA constructs may encode 3 siRNAs, each directed to the same region or sequence of the target mRNA. For example, recombinant polynucleic acid constructs encoding recombinant RNA constructs may encode 3 siRNAs, each directed to a different region or sequence of the target mRNA. In some aspects, recombinant polynucleic acid constructs encoding recombinant RNA constructs may encode 3 siRNA species, wherein each of the 3 siRNA species is directed to a different target mRNA. In some embodiments, a target mRNA may be an mRNA of VEGF, VEGFA, an isoform of VEGFA, PIGF, MICA, MICB, ERp5, ADAM17, MMP, IDH1, CDK4, CDK6, EGFR, mTOR, KRAS, CD155, PD-L1, or c-Myc. In related aspects, recombinant polynucleic acid constructs may comprise a sequence selected from the group consisting of SEQ ID NOs: 82-98.

[0131] The polynucleic acid constructs, described herein, can be obtained by any method known in the art, such as by chemically synthesizing the DNA chain, by PCR, or by the Gibson Assembly method. The advantage of constructing polynucleic acid constructs by chemical synthesis or a combination of PCR method or Gibson Assembly method is that the codons may be optimized to ensure that the fusion protein is expressed at a high level in a host cell. Codon optimization can refer to a process of modifying a nucleic acid sequence for expression in a host cell of interest by replacing at least one codon (*e.g.*, more than 1, 2, 3, 4, 5, 10, 15, 20, 25, 50, or more codons) of a native sequence with codons that are more frequently or most frequently used in the genes of that host cell while maintaining the native amino acid sequence. Codon usage tables are readily available, for example, at the “Codon Usage Database,” and these tables can be adapted in a number of ways. Computer algorithms for codon optimizing a particular sequence for expression in a particular host cell are also

available, such as Gene Forge[®] (Aptagen, PA) and GeneOptimizer[®] (ThermoFischer, MA). Once obtained polynucleotides can be incorporated into suitable vectors. Vectors as used herein can refer to naturally occurring or synthetically generated constructs for uptake, proliferation, expression or transmission of nucleic acids *in vivo* or *in vitro*, *e.g.*, plasmids, minicircles, phagemids, cosmids, artificial chromosomes/mini-chromosomes, bacteriophages, viruses such as baculovirus, retrovirus, adenovirus, adeno-associated virus, herpes simplex virus, bacteriophages. Methods used to construct vectors are well known to a person skilled in the art and described in various publications. In particular techniques for constructing suitable vectors, including a description of the functional and regulatory components such as promoters, enhancers, termination and polyadenylation signals, selection markers, origins of replication, and splicing signals, are known to the person skilled in the art. A variety of vectors are well known in the art and some are commercially available from companies such as Agilent Technologies, Santa Clara, Calif.; Invitrogen, Carlsbad, Calif.; Promega, Madison, Wis.; Thermo Fisher Scientific; or Invivogen, San Diego, Calif. A non-limiting examples of vectors for *in vitro* transcription includes pT7CFE1-CHis, pMX (such as pMA-T, pMA-RQ, pMC, pMK, pMS, pMZ), pEVL, pSP73, pSP72, pSP64, and pGEM (such as pGEM[®]-4Z, pGEM[®]-5Zf(+), pGEM[®]-11Zf(+), pGEM[®]-9Zf(-), pGEM[®]-3Zf(+/-), pGEM[®]-7Zf(+/-)). In some instances, recombinant polynucleic acid constructs may be DNA.

[0132] The polynucleic acid constructs, as described herein, can be circular or linear. For example, circular polynucleic acid constructs may include vector system such as pMX, pMA-T, pMA-RQ, or pT7CFE1-CHis. For example, linear polynucleic acid constructs may include linear vector such as pEVL or linearized vectors. In some instances, recombinant polynucleic acid constructs may further comprise a promoter. In some instances, the promoter may be present upstream of the sequence encoding for the first RNA or the sequence encoding for the second RNA. Non-limiting examples of a promoter can include T3, T7, SP6, P60, Syn5, and KP34. In some instances, recombinant polynucleic acid constructs provided herein may comprise a T7 promoter comprising a sequence comprising TAATACGACTCACTATA (SEQ ID NO: 18). In some instances, recombinant polynucleic acid constructs further comprises a sequence encoding a Kozak sequence. A Kozak sequence may refer to a nucleic acid sequence motif that functions as the protein translation initiation site. Kozak sequences are described at length in the literature, *e.g.*, by Kozak, M., Gene 299(1-2):1-34, incorporated herein by reference herein in its entirety. In some embodiments, recombinant polynucleic acid constructs comprises a sequence encoding a Kozak sequence comprising a sequence

comprising GCCACC (SEQ ID NO: 19). In some instances, recombinant polynucleic acid constructs described herein may be codon-optimized.

[0133] Provided herein are compositions comprising recombinant polynucleic acid constructs encoding RNA constructs described herein comprising one or more nucleic acid sequence
5 encoding an siRNA capable of binding to a target RNA and one or more nucleic acid sequence encoding a gene of interest, wherein the siRNA capable of binding to a target RNA is not a part of an intron sequence encoded by the gene of interest. In some instances, the gene of interest is expressed without RNA splicing. In some instances, the siRNA capable of binding to a target RNA binds to an exon of a target mRNA. In some instances, the siRNA
10 capable of binding to a target RNA specifically binds to one target RNA. In some instances, recombinant polynucleic acid constructs may comprise a nucleic acid sequence comprising a sequence selected from the group consisting of SEQ ID NOs: 82-98.

[0134] Provided herein are methods of producing RNA construct compositions described herein. For example, recombinant RNA constructs may be produced by *in vitro* transcription
15 from a polynucleic acid construct comprising a promoter for an RNA polymerase, at least one nucleic acid sequence encoding a gene of interest, at least one nucleic acid sequence encoding an siRNA capable of binding to a target mRNA, and a nucleic acid sequence encoding poly(A) tail. *In vitro* transcription reaction may further comprise an RNA polymerase, a mixture of nucleotide triphosphates (NTPs), and/or a capping enzyme. Details of producing
20 RNAs using *in vitro* transcription as well as isolating and purifying transcribed RNAs is well known in the art and can be found, for example, in Beckert & Masquida ((2011) Synthesis of RNA by In vitro Transcription. RNA. Methods in Molecular Biology (Methods and Protocols), vol 703. Humana Press). A non-limiting list of *in vitro* transcript kits includes
25 MEGAscript™ T3 Transcription Kit, MEGAscript T7 kit, MEGAscript™ SP6 Transcription Kit, MAXIsript™ T3 Transcription Kit, MAXIsript™ T7 Transcription Kit, MAXIsript™ SP6 Transcription Kit, MAXIsript™ T7/T3 Transcription Kit, MAXIsript™ SP6/T7 Transcription Kit, mMESSAGE mMACHINE™ T3 Transcription Kit, mMESSAGE mMACHINE™ T7 Transcription Kit, mMESSAGE mMACHINE™ SP6 Transcription Kit, MEGAscript™ T7 Transcription Kit, HiScribe™ T7 High Yield RNA Synthesis Kit,
30 HiScribe™ T7 In Vitro Transcription Kit, AmpliScribe™ T7-Flash™ Transcription Kit, AmpliScribe™ T7 High Yield Transcription Kit, AmpliScribe™ T7-Flash™ Biotin-RNA Transcription Kit, T7 Transcription Kit, HighYield T7 RNA Synthesis Kit, DuraScribe® T7 Transcription Kit, etc.

[0135] The *in vitro* transcription reaction can further comprise a transcription buffer system, nucleotide triphosphates (NTPs), and an RNase inhibitor. In some embodiments, the transcription buffer system may comprise dithiothreitol (DTT) and magnesium ions. The NTPs can be naturally occurring or non-naturally occurring (modified) NTPs. Non-limiting examples of non-naturally occurring (modified) NTPs include N¹-Methylpseudouridine, Pseudouridine, N¹-Ethylpseudouridine, N¹-Methoxymethylpseudouridine, N¹-Propylpseudouridine, 2-thiouridine, 4-thiouridine, 5-methoxyuridine, 5-methyluridine, 5-carboxymethylesteruridine, 5-formyluridine, 5-carboxyuridine, 5-hydroxyuridine, 5-Bromouridine, 5-Iodouridine, 5,6-dihydrouridine, 6-Azauridine, Thienouridine, 3-methyluridine, 1-carboxymethyl-pseudouridine, 4-thio-1-methyl-pseudouridine, 2-thio-1-methyl-pseudouridine, dihydrouridine, dihydropseudouridine, 2-methoxyuridine, 2-methoxy-4-thio-uridine, 4-methoxy-pseudouridine, 4-methoxy-2-thio-pseudouridine, 5-methylcytidine, 5-methoxycytidine, 5-hydroxymethylcytidine, 5-formylcytidine, 5-carboxycytidine, 5-hydroxycytidine, 5-Iodocytidine, 5-Bromocytidine, 2-thiocytidine, 5-azacytidine, pseudoisocytidine, 3-methyl-cytidine, N⁴-acetylcytidine, 5-formylcytidine, N⁴-methylcytidine, 5-hydroxymethylcytidine, 1-methyl-pseudoisocytidine, 4-methoxy-pseudoisocytidine, and 4-methoxy-1-methyl-pseudoisocytidine, N¹-methyladenosine, N⁶-methyladenosine, N⁶-methyl-2-Aminoadenosine, N⁶-isopentenyladenosine, N⁶,N⁶-dimethyladenosine, 7-methyladenine, 2-methylthio-adenine, and 2-methoxy-adenine. Non-limiting examples of DNA-dependent RNA polymerase include T3, T7, SP6, P60, Syn5, and KP34 RNA polymerases. In some embodiments, the RNA polymerase is selected from the group consisting of T3 RNA polymerase, T7 RNA polymerase, SP6 RNA polymerase, P60 RNA polymerase, Syn5 RNA polymerase, and KP34 RNA polymerase.

[0136] Transcribed RNAs, as described herein, may be isolated and purified from the *in vitro* transcription reaction mixture. For example, transcribed RNAs may be isolated and purified using column purification. Details of isolating and purifying transcribed RNAs from *in vitro* transcription reaction mixture is well known in the art and any commercially available kits may be used. A non-limiting list of RNA purification kits includes MEGAclear kit, Monarch® RNA Cleanup Kit, EasyPure® RNA Purification Kit, NucleoSpin® RNA Cleanup, etc.

Therapeutic applications

[0137] Provided herein are compositions useful in the treatment of a cancer. In some aspects, compositions are present or administered in an amount sufficient to treat or prevent a disease

or condition. Provided herein are compositions comprising a first RNA encoding a cytokine linked to a second RNA encoding a genetic element that can reduce expression of a gene associated with tumor proliferation, angiogenesis, or recognition by the immune system. In some embodiments, a cytokine may comprise IL-2, IL-7, IL-12, IL-15, a fragment thereof, or a functional variant thereof. In some embodiments, a genetic element that can reduce expression of a gene associated with tumor proliferation or angiogenesis may comprise siRNA targeting VEGF, VEGFA, an isoform of VEGFA, PIGF, IDH1, CDK4, CDK6, EGFR, mTOR, KRAS, CD155, PD-L1, c-Myc, a fragment thereof, or a functional variant thereof. In some embodiments, a genetic element that can reduce expression of a gene associated with recognition by the immune system may comprise siRNA targeting MICA, MICB, ERp5, ADAM, MMP, a fragment thereof, or a functional variant thereof. In some embodiments, the ADAM is ADAM17.

[0138] Also provided herein are pharmaceutical compositions comprising any RNA composition described herein and a pharmaceutically acceptable excipient. A pharmaceutical composition can denote a mixture or solution comprising a therapeutically effective amount of an active pharmaceutical ingredient together with one or more pharmaceutically acceptable excipients to be administered to a subject in need thereof. The term “pharmaceutically acceptable” denotes an attribute of a material which is useful in preparing a pharmaceutical composition that is generally safe, non-toxic, and neither biologically nor otherwise undesirable and is acceptable for veterinary as well as human pharmaceutical use. The term “pharmaceutically acceptable” can refer to a material, such as a carrier or diluent, which does not abrogate the biological activity or properties of the compound, and is relatively nontoxic, i.e. the material may be administered to an individual without causing undesirable biological effects or interacting in a deleterious manner with any of the components of the composition in which it is contained. A pharmaceutically acceptable excipient can denote any pharmaceutically acceptable ingredient in a pharmaceutical composition having no therapeutic activity and being non-toxic to the subject administered, such as disintegrators, binders, fillers, solvents, buffers, tonicity agents, stabilizers, antioxidants, surfactants, carriers, diluents, excipients, preservatives or lubricants used in formulating pharmaceutical products. Pharmaceutical compositions can facilitate administration of the compound to an organism and can be formulated in a conventional manner using one or more pharmaceutically acceptable inactive ingredients that facilitate processing of the active compounds into preparations that can be used pharmaceutically. A proper formulation is dependent upon the route of administration chosen and a summary of pharmaceutical compositions can be found,

for example, in Remington: The Science and Practice of Pharmacy, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., Pharmaceutical Dosage Forms, Marcel Decker, New York, N.Y., 1980; and
5 Pharmaceutical Dosage Forms and Drug Delivery Systems, Seventh Ed. (Lippincott Williams & Wilkins 1999), herein incorporated by reference. In some embodiments, pharmaceutical compositions can be formulated by dissolving active substances (e.g., recombinant polynucleic acid or RNA constructs described herein) in aqueous solution for injection into diseased tissues or diseased cells. In some embodiments, pharmaceutical compositions can be
10 formulated by dissolving active substances (e.g., recombinant polynucleic acid or RNA constructs described herein) in aqueous solution for direct injection into diseased tissues or diseased cells. In some embodiments, diseased tissues or diseased cells comprise tumors or tumor cells.

[0139] Also provided herein are methods of treating a cancer in a subject in need thereof,
15 comprising administering to the subject with the cancer a therapeutically effective amount of compositions or pharmaceutical compositions described herein. The terms "effective amount" or "therapeutically effective amount," as used herein, refer to a sufficient amount of an agent or a compound being administered which will relieve to some extent one or more of the symptoms of the disease or the condition being treated; for example a reduction and/or
20 alleviation of one or more signs, symptoms, or causes of a disease, or any other desired alteration of a biological system. For example, an "effective amount" for therapeutic uses can be an amount of an agent that provides a clinically significant decrease in one or more disease symptoms. An appropriate "effective" amount may be determined using techniques, such as a dose escalation study, in individual cases.

25 **[0140]** The terms "treat," "treating" or "treatment," as used herein, include alleviating, abating or ameliorating at least one symptom of a disease or a condition, preventing additional symptoms, inhibiting the disease or the condition, *e.g.*, arresting the development of the disease or the condition, relieving the disease or the condition, causing regression of the disease or the condition, relieving a condition caused by the disease or the condition, or
30 stopping the symptoms of the disease or the condition either prophylactically and/or therapeutically. In some embodiments, treating a disease or condition comprises reducing the size of diseased tissues or diseased cells. In some embodiments, treating a disease or a condition in a subject comprises increasing the survival of a subject. In some embodiments, treating a disease or condition comprises reducing or ameliorating the severity of a disease,

delaying onset of a disease, inhibiting the progression of a disease, reducing hospitalization of or hospitalization length for a subject, improving the quality of life of a subject, reducing the number of symptoms associated with a disease, reducing or ameliorating the severity of a symptom associated with a disease, reducing the duration of a symptom associated with a disease, preventing the recurrence of a symptom associated with a disease, inhibiting the development or onset of a symptom of a disease, or inhibiting of the progression of a symptom associated with a disease. In some embodiments, treating a cancer comprises reducing the size of tumor or increasing survival of a patient with a cancer.

[0141] In some cases, a subject can encompass mammals. Examples of mammals include, but are not limited to, any member of the mammalian class: humans, non-human primates such as chimpanzees, and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, swine; domestic animals such as rabbits, dogs, and cats; laboratory animals including rodents, such as rats, mice and guinea pigs, and the like. In some cases, the mammal is a human. In some cases, the subject may be an animal. In some cases, an animal may comprise human beings and non-human animals. In one embodiment, a non-human animal may be a mammal, for example a rodent such as rat or a mouse. In another embodiment, a non-human animal may be a mouse. In some instances, the subject is a mammal. In some instances, the subject is a human. In some instances, the subject is an adult, a child, or an infant. In some instances, the subject is a companion animal. In some instances, the subject is a feline, a canine, or a rodent. In some instances, the subject is a dog or a cat.

[0142] Further provided herein are methods of treating a cancer comprising administering compositions or pharmaceutical compositions described herein to a subject with a cancer. In some instances, the cancer is a solid tumor. In some instances, a solid tumor may include, but is not limited to, breast cancer, lung cancer, liver cancer, glioblastoma, melanoma, head and neck squamous cell carcinoma, renal cell carcinoma, neuroblastoma, Wilms tumor, retinoblastoma, rhabdomyosarcoma, osteosarcoma, Ewing sarcoma, bladder cancer, cervical cancer, colon cancer, rectal cancer, endometrial cancer, kidney cancer, mesothelioma, non-small cell lung cancer, nonmelanoma skin cancer, ovarian cancer, pancreatic cancer, prostate cancer, small cell lung cancer, colorectal cancer, and thyroid cancer. In some embodiments, a solid tumor may include sarcomas, carcinomas, or lymphomas. In some embodiments, a solid tumor can be benign or malignant.

[0143] In some instances, the cancer is a head and neck cancer. Without wishing to be bound to any theory, the head and neck cancer is the sixth most common cancer worldwide and represent 6% of solid tumors. Approximately 650,000 new patients are diagnosed with head

and neck cancers annually, and there are 350,000 deaths yearly worldwide with 12,000 deaths in the US despite the availability of advanced treatment options. Risk factors that increase the chance of developing head and neck cancers include use of tobacco and/or alcohol, prolonged sun exposure (*e.g.*, in the lip area or skin of the head and neck), human papillomavirus (HPV),
5 Epstein-Barr virus (EBV), gender (*e.g.*, men versus women), age (*e.g.*, people over the age of 40 are at higher risk), poor oral and dental hygiene, and environmental or occupational inhalants (*e.g.*, asbestos, wood dust, paint fumes, and other certain chemicals), marijuana use, poor nutrition, gastroesophageal reflux disease (GERD) and laryngopharyngeal reflux disease (LPRD), weakened immune system, radiation exposure, or previous history of head and neck
10 cancer. Tobacco use is the single largest risk factor for head and neck cancer, and includes smoking cigarettes, cigars, or pipes; chewing tobacco; using snuff; and secondhand smoke. About 85% of head and neck cancers are linked to tobacco use, and the amount of tobacco use may affect prognosis. In addition, nearly 25 % of head and neck cancers are HPV-positive.

[0144] Head and neck cancers can include epithelial malignancies of the upper aerodigestive tract, including the paranasal sinuses, nasal cavity, oral cavity, pharynx, and larynx. Non-limiting examples of the head and neck cancer includes laryngeal cancer, hypopharyngeal cancer, tonsil cancer, nasal cavity cancer, paranasal sinus cancer, nasopharyngeal cancer, metastatic squamous neck cancer with occult primary, lip cancer, oral cancer, oropharyngeal cancer, salivary gland cancer, brain tumors, esophageal cancer, eye cancer, parathyroid
15 cancer, sarcoma of the head and neck, and thyroid cancer. The head and neck cancers described herein may be located at an upper aerodigestive tract. Non-limiting examples of the upper aerodigestive tract include a paranasal sinus, a nasal cavity, an oral cavity, a salivary gland, a tongue, a nasopharynx, an oropharynx, a hypopharynx, and a larynx.

[0145] In some embodiments, the cancer is selected from the group consisting of a head and neck cancer, melanoma, and renal cell carcinoma. In some embodiments, the cancer is a head and neck cancer. In some embodiments, the head and neck cancer is head and neck squamous cell carcinoma. In some embodiments, the head and neck cancer is laryngeal cancer,
25 hypopharyngeal cancer, tonsil cancer, nasal cavity cancer, paranasal sinus cancer, nasopharyngeal cancer, metastatic squamous neck cancer with occult primary, lip cancer, oral cancer, oropharyngeal cancer, salivary gland cancer, brain tumors, esophageal cancer, eye cancer, parathyroid cancer, sarcoma of the head and neck, or thyroid cancer. In some embodiments, the cancer is melanoma. In some embodiments, the cancer is renal cell carcinoma.

30

[0146] Early treatment for cancers described herein may include surgical removal of tumors, radiation therapy, therapies using medications such as chemotherapy, targeted therapy, immunotherapy, or combinations thereof. Targeted therapy is a treatment that target specific genes, proteins, or the tissue environment that can contribute to cancer growth and survival, and the treatment is designed to block the growth and spread of cancer cells while limiting damage to healthy cells. For head and neck cancers, targeted therapies using antibodies may be used to inhibit cell proliferation, tumor proliferation or growth, or suppress tumor angiogenesis. Immunotherapy is a treatment that can improve, target, or restore immune system function to fight cancer. Non-limiting examples of antibodies include anti-epidermal growth factor receptor (EGFR) antibodies and anti-vascular endothelial growth factor (VEGF) antibodies. Non-limiting examples of cancer immunotherapy include immune system modulators, T-cell transfer therapy, immune checkpoint inhibitors, and monoclonal antibodies. Immune system modulators can enhance immune response against cancer and include cytokines such as interleukins and interferon alpha (IFN α). T-cell transfer therapy can refer to a treatment where immune cells are taken from a cancer patient for *ex vivo* manipulation and injected back to the same patient. For example, immune cells are taken from a cancer patient for specific expansion of tumor-recognizing lymphocytes (*e.g.*, tumor-infiltrating lymphocytes therapy) or for modification of cells to express chimeric antigen receptors specifically recognizing tumor antigens (*e.g.*, CAR T-cell therapy). Immune checkpoint inhibitors can block immune checkpoints, restoring or allowing immune responses to cancer cells. Non-limiting examples of immune checkpoint inhibitors include programmed death-ligand 1 (PD-L1) inhibitors, programmed death protein 1 (PD1) inhibitors, and cytotoxic T-lymphocyte-associated protein-4 (CTLA-4) inhibitors. Monoclonal antibodies can be designed to bind to specific target proteins to block the activity of target proteins in cancer cells (*e.g.*, anti-EGFR, anti-VEGF, etc.).

[0147] In cancers, decreasing expression of genes involved in tumor proliferation, angiogenesis, or recognition by the immune system (*e.g.*, VEGF, VEGFA, an isoform of VEGFA, PIGF, MICA, MICB, ERp5, ADAM, MMP, IDH1, CDK4, CDK6, EGFR, mTOR, KRAS, CD155, PD-L1, or c-Myc, etc.) while increasing expression of cytokines (*e.g.*, IL-2, IL-12, IL-15, or IL-7, etc.) to enhance immune response could have a therapeutic effect. In one example, expression of IL-2, that can decrease proliferation rate of cancer cells such as head and neck squamous cell carcinoma (HNSCC) cells, can be increased. IL-2 is a cytokine that regulates lymphocyte activities and is a potent T-cell growth factor. IL-2 is produced by antigen-stimulated CD4⁺ T-cells, natural killer cells, or activated dendritic cells and is

important for maintenance and differentiation of CD4+ regulatory T-cells. Without wishing to be bound by any theory, local IL-2 therapy can cause stagnation of the blood flow inside or near tumors and of the lymph drainage, leading to tumor necrosis and thrombosis. In another example, expression of VEGF, which can promote angiogenesis around tumor, can be

5 decreased to block the supply of blood required for tumor growth. VEGF described herein may be any VEGF family members including VEGFA, an isoform of VEGFA, or PlGF. Non-limiting examples of VEGFA isoforms include, VEGF111, VEGF121, VEGF145, VEGF148, VEGF165, VEGF165B, VEGF183, VEGF189, VEGF206, L-VEGF121, L-VEGF165, L-VEGF189, L-VEGF206, Isoform 15, Isoform16, Isoform 17, and Isoform 18. In yet another

10 example, expression of MICA and/or MICB (MICA/B), cell surface glycoproteins expressed by tumor cells, can be decreased to restore immune response of natural killer (NK) cells and T-cells to enhance tumor regression. MICA/B is recognized by natural killer group 2 member D (NKG2D) receptor expressed on NK cells and lymphocytes to promote recognition and elimination of tumor cells. Cancer cells may evade immune surveillance by shedding

15 MICA/B from cell surface to impair NKG2D recognition. Cancer cells may also release soluble forms of MICA/B that can bind to NKG2D receptor during tumor growth and hypoxia, which may induce NKG2D internalization, to escape immune responses and compromise immune surveillance by NK cells. Shedding or releasing of MICA/B from cell surface may be blocked by inhibiting or reducing the expression of proteins involved in

20 shedding of a membrane protein. Examples of proteins involved in shedding include, but are not limited to, matrix metalloproteinases (MMPs) and a disintegrin and metalloproteinases (ADAMs). Non-limiting examples of MMPs include MMP1, MMP2, MMP3, MMP7, MMP8, MMP9, MMP10, MMP11, MMP12, MMP13, MMP14, MMP15, MMP16, MMP17, and MMP19. Shedding or releasing of MICA/B from cell surface may also be blocked by

25 inhibiting or reducing the expression of factors regulating the proteins involved in shedding such as disulfide isomerase ERp5.

[0148] In some aspects, provided herein, is a method of treating a cancer in a subject, the method comprising administering to the subject RNA compositions or pharmaceutical compositions, described herein, comprising an mRNA encoding a gene of interest and siRNA

30 capable of binding to a target mRNA. In some aspects, provided herein, are any RNA compositions or pharmaceutical compositions, described herein, comprising an mRNA encoding a gene of interest and siRNA capable of binding to a target mRNA for use in a method for the treatment of cancer. In some aspects, provided herein, is the use of RNA compositions or pharmaceutical compositions, described herein, comprising an mRNA

encoding a gene of interest and siRNA capable of binding to a target mRNA for the manufacture of a medicament for treating cancer. In some aspects, provided herein, is the use of RNA compositions or pharmaceutical compositions, described herein, comprising an mRNA encoding a gene of interest and siRNA capable of binding to a target mRNA for
5 treating cancer in a subject. In some embodiments, the siRNA is capable of binding to VEGF, VEGFA, an isoform of VEGFA, PIGF, IDH1, CDK4, CDK6, EGFR, mTOR, KRAS, CD155, PD-L1, c-Myc, a fragment thereof, or a functional variant thereof. In some embodiments, the siRNA is capable of binding to MICA, MICB, both MICA and MICB (MICA/B), ERp5, ADAM, MMP, a fragment thereof, or a functional variant thereof. In some embodiments, the
10 ADAM is ADAM17. In some embodiments, the mRNA encoding the gene of interest encodes a cytokine. In some embodiments, the cytokine is an IL-2, IL-12, IL-15, IL-7, a fragment thereof, or a functional variant thereof.

[0149] In some aspects, provided herein, is a method of treating a cancer in a subject, the method comprising administering to the subject recombinant RNA compositions or
15 pharmaceutical compositions, described herein, comprising siRNA capable of binding to VEGFA, IDH1, CDK4, CDK6, EGFR, mTOR, KRAS, CD155, PD-L1, or c-Myc and an mRNA encoding IL-2, IL-12, IL-15, or IL-7. In some aspects, provided herein, are recombinant RNA compositions or pharmaceutical compositions, described herein, comprising siRNA capable of binding to VEGFA, IDH1, CDK4, CDK6, EGFR, mTOR,
20 KRAS, CD155, PD-L1, or c-Myc and an mRNA encoding IL-2, IL-12, IL-15, or IL-7 for use in a method for the treatment of cancer. In some aspects, provided herein, is the use of recombinant RNA compositions or pharmaceutical compositions, described herein, comprising siRNA capable of binding to VEGFA, IDH1, CDK4, CDK6, EGFR, mTOR, KRAS, CD155, PD-L1, or c-Myc and an mRNA encoding IL-2, IL-12, IL-15, or IL-7 for the
25 manufacture of a medicament for treating cancer. In some aspects, provided herein, is the use of recombinant RNA compositions or pharmaceutical compositions, described herein, comprising siRNA capable of binding to VEGFA, IDH1, CDK4, CDK6, EGFR, mTOR, KRAS, CD155, PD-L1, or c-Myc and an mRNA encoding IL-2, IL-12, IL-15, or IL-7 for treating cancer in a subject. In some aspects, provided herein, is a method of treating a cancer
30 in a subject, the method comprising administering to the subject recombinant RNA compositions or pharmaceutical compositions, described herein, comprising siRNA capable of binding to an mRNA of a VEGFA isoform and an mRNA encoding IL-2. In some aspects, provided herein, is a method of treating a cancer in a subject, the method comprising administering to the subject recombinant RNA compositions or pharmaceutical compositions,

described herein, comprising siRNA capable of binding to a PIGF mRNA and an mRNA encoding IL-2. In some aspects, provided herein, is a method of treating a cancer in a subject, the method comprising administering to the subject recombinant RNA compositions or pharmaceutical compositions, described herein, comprising siRNA capable of binding to an mRNA of MICA or MICB and an mRNA encoding IL-2. In some aspects, provided herein, is a method of treating a cancer in a subject, the method comprising administering to the subject recombinant RNA compositions or pharmaceutical compositions, described herein, comprising siRNA capable of binding to an mRNA of ERp5, ADAM17, or MMP and an mRNA encoding IL-2. In some aspects, provided herein, is a method of treating a cancer in a subject, the method comprising administering to the subject recombinant RNA compositions or pharmaceutical compositions, described herein, comprising siRNA capable of binding to an mRNA of VEGFA, MICA, MICB, IDH1, CDK4, CDK6, EGFR, mTOR, KRAS, CD155, PD-L1, or c-Myc and an mRNA encoding IL-2, IL-7, IL-12, or IL-15.

[0150] In some aspects, compositions or pharmaceutical compositions administered to a subject in need thereof comprise recombinant RNA constructs comprising: (i) an IL-2 mRNA; and (ii) at least one siRNA capable of binding to a VEGFA mRNA. In related aspects, the polynucleic acid construct encodes or comprises at least 1, 2, 3, 4, or 5 siRNAs. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to a VEGFA mRNA. In related aspects, recombinant RNA constructs may comprise at least 3 or at least 5 siRNAs, each directed to a VEGFA mRNA. In related aspects, each of the at least 3 or at least 5 siRNAs is the same, different, or a combination thereof. In related aspects, recombinant RNA constructs may comprise a sequence as set forth in SEQ ID NO: 1-4 or 125-128 (Cpd.1-Cpd.4). In related aspects, recombinant RNA constructs may comprise a sequence as set forth in SEQ ID NO: 5 (Cpd.5), SEQ ID NO: 7 (Cpd.7), SEQ ID NO: 8 (Cpd.8), SEQ ID NO: 9 (Cpd.9), SEQ ID NO: 10 (Cpd.10), SEQ ID NO: 129 (Cpd.5), SEQ ID NO: 131 (Cpd.7), SEQ ID NO: 132 (Cpd.8), SEQ ID NO: 133 (Cpd.9), or SEQ ID NO: 134 (Cpd.10).

[0151] In some aspects, compositions or pharmaceutical compositions administered to a subject in need thereof comprise recombinant RNA constructs comprising: (i) an IL-2 mRNA; and (ii) at least one siRNA capable of binding to a PIGF mRNA. In related aspects, the polynucleic acid construct encodes or comprises at least 1, 2, or 3 siRNAs. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to a PIGF mRNA. In related aspects, recombinant RNA constructs may comprise at least 3 siRNAs, each directed to a PIGF mRNA. In related aspects, each of the at least 3 siRNAs is the same, different, or a combination thereof.

- [0152] In some aspects, compositions or pharmaceutical compositions administered to a subject in need thereof comprise recombinant RNA constructs comprising: (i) an IL-2 mRNA; and (ii) at least one siRNA capable of binding to an mRNA of a VEGFA isoform. In related aspects, the polynucleic acid construct encodes or comprises at least 1, 2, or 3 siRNAs. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to an mRNA of a VEGFA isoform. In related aspects, recombinant RNA constructs may comprise at least 3 siRNAs, each directed to an mRNA of a VEGFA isoform. In related aspects, each of the at least 3 siRNAs is the same, different, or a combination thereof.
- [0153] In some aspects, compositions or pharmaceutical compositions administered to a subject in need thereof comprise recombinant RNA constructs comprising: (i) an IL-2 mRNA; and (ii) at least one siRNA capable of binding to a MICA or MICB mRNA. In related aspects, recombinant RNA constructs may comprise at least 1, 2, or 3 siRNAs. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to a MICA or MICB mRNA. In related aspects, recombinant RNA constructs may comprise at least 3 siRNAs, each directed to a MICA or MICB mRNA. In related aspects, each of the at least 3 siRNAs is the same, different, or a combination thereof. In related aspects, recombinant RNA constructs may comprise a sequence as set forth in SEQ ID NO: 1-4 or 125-128 (Cpd.1-Cpd.4). In related aspects, recombinant RNA constructs may comprise a sequence as set forth in SEQ ID NO: 6 or SEQ ID NO: 130 (Cpd.6).
- [0154] In some aspects, compositions or pharmaceutical compositions administered to a subject in need thereof comprise recombinant RNA constructs comprising: (i) an IL-2 mRNA; and (ii) at least one siRNA capable of binding to an mRNA of ERp5, ADAM17, or MMP. In related aspects, recombinant RNA constructs may comprise at least 1, 2, or 3 siRNAs. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to an mRNA of ERp5, ADAM17, or MMP. In related aspects, recombinant RNA constructs may comprise at least 3 siRNAs, each directed to an mRNA of ERp5, ADAM17, or MMP. In related aspects, each of the at least 3 siRNAs is the same, different, or a combination thereof.
- [0155] In some aspects, compositions or pharmaceutical compositions administered to a subject in need thereof comprise recombinant RNA constructs comprising: (i) an IL-12 mRNA; and (ii) at least one siRNA capable of binding to an mRNA of IDH1, CDK4, and/or CDK6. In related aspects, the polynucleic acid construct encodes or comprises at least 1, 2, or 3 siRNAs. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to an IDH1 mRNA. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to a CDK4 mRNA. In related aspects, recombinant RNA constructs may comprise 1

- siRNA directed to a CDK6 mRNA. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to an IDH1 mRNA, 1 siRNA directed to a CDK4 mRNA, and 1 siRNA directed to a CDK6 mRNA. In related aspects, recombinant RNA constructs may comprise at least 3 siRNAs, each directed to an IDH1 mRNA. In related aspects, recombinant RNA constructs may comprise at least 3 siRNAs, each directed to a CDK4 mRNA. In related aspects, recombinant RNA constructs may comprise at least 3 siRNAs, each directed to a CDK6 mRNA. In related aspects, each of the at least 3 siRNAs is the same, different, or a combination thereof. In related aspects, recombinant RNA constructs may comprise a sequence as set forth in SEQ ID NO: 11 or SEQ ID NO: 135 (Cpd.11).
- 10 **[0156]** In some aspects, compositions or pharmaceutical compositions administered to a subject in need thereof comprise recombinant RNA constructs comprising: (i) an IL-12 mRNA; and (ii) at least one siRNA capable of binding to an mRNA of EGFR, mTOR, and/or KRAS. In related aspects, the polynucleic acid construct encodes or comprises at least 1, 2, or 3 siRNAs. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to an EGFR mRNA. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to an mTOR mRNA. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to a KRAS mRNA. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to an EGFR mRNA, 1 siRNA directed to an mTOR mRNA, and 1 siRNA directed to a KRAS mRNA. In related aspects, recombinant RNA constructs may comprise at least 3 siRNAs, each directed to an EGFR mRNA. In related aspects, recombinant RNA constructs may comprise at least 3 siRNAs, each directed to an mTOR mRNA. In related aspects, recombinant RNA constructs may comprise at least 3 siRNAs, each directed to a KRAS mRNA. In related aspects, each of the at least 3 siRNAs is the same, different, or a combination thereof. In related aspects, recombinant RNA constructs may comprise a sequence as set forth in SEQ ID NO: 12 (Cpd.12), SEQ ID NO: 13 (Cpd.13), SEQ ID NO: 14 (Cpd.14), SEQ ID NO: 136 (Cpd.12), SEQ ID NO: 137 (Cpd.13), or SEQ ID NO: 138 (Cpd.14).
- 25 **[0157]** In some aspects, compositions or pharmaceutical compositions administered to a subject in need thereof comprise recombinant RNA constructs comprising: (i) an IL-15 mRNA; and (ii) at least one siRNA capable of binding to an mRNA of VEGFA and/or CD155. In related aspects, the polynucleic acid construct encodes or comprises at least 1, 2, or 3 siRNAs. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to a VEGFA mRNA. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to a CD155 mRNA. In related aspects, recombinant RNA constructs may comprise 1

siRNA directed to a VEGFA mRNA and 2 siRNAs directed to a CD155 mRNA. In related aspects, recombinant RNA constructs may comprise at least 3 siRNAs, each directed to a VEGFA mRNA. In related aspects, recombinant RNA constructs may comprise at least 3 siRNAs, each directed to a CD155 mRNA. In related aspects, each of the at least 3 siRNAs is the same, different, or a combination thereof. In related aspects, recombinant RNA constructs may comprise a sequence as set forth in SEQ ID NO: 15 or 139 (Cpd.15).

[0158] In some aspects, compositions or pharmaceutical compositions administered to a subject in need thereof comprise recombinant RNA constructs comprising: (i) an IL-15 mRNA; and (ii) at least one siRNA capable of binding to an mRNA of VEGFA, PD-L1, and/or c-Myc. In related aspects, the polynucleic acid construct encodes or comprises at least 1, 2, or 3 siRNAs. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to a VEGFA mRNA. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to a PD-L1 mRNA. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to a c-Myc mRNA. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to a VEGFA mRNA, 1 siRNA directed to a PD-L1 mRNA, and 1 siRNA directed to a c-Myc mRNA. In related aspects, recombinant RNA constructs may comprise at least 3 siRNAs, each directed to a VEGFA mRNA. In related aspects, recombinant RNA constructs may comprise at least 3 siRNAs, each directed to a PD-L1 mRNA. In related aspects, recombinant RNA constructs may comprise at least 3 siRNAs, each directed to a c-Myc mRNA. In related aspects, each of the at least 3 siRNAs is the same, different, or a combination thereof. In related aspects, recombinant RNA constructs may comprise a sequence as set forth in SEQ ID NO: 16 or 140 (Cpd.16).

[0159] In some aspects, compositions or pharmaceutical compositions administered to a subject in need thereof comprise recombinant RNA constructs comprising: (i) an IL-7 mRNA; and (ii) at least one siRNA capable of binding to an mRNA of PD-L1. In related aspects, the polynucleic acid construct encodes or comprises at least 1, 2, or 3 siRNAs. In related aspects, recombinant RNA constructs may comprise 1 siRNA directed to a PD-L1 mRNA. In related aspects, recombinant RNA constructs may comprise at least 3 siRNAs, each directed to a PD-L1 mRNA. In related aspects, each of the at least 3 siRNAs is the same, different, or a combination thereof. In related aspects, recombinant RNA constructs may comprise a sequence as set forth in SEQ ID NO: 17 or 141 (Cpd.17)

[0160] Recombinant RNA construct compositions described herein may be administered as a combination therapy. Combination therapies with two or more therapeutic agents or therapies may use agents and therapies that work by different mechanisms of action. Combination

therapies using agents or therapies with different mechanisms of action can result in additive or synergetic effects. Combination therapies may allow for a lower dose of each agent than is used in monotherapy, thereby reducing toxic side effects and/or increasing the therapeutic index of the agent(s). Combination therapies can decrease the likelihood that resistant cancer cells will develop. In some instances, combination therapies comprise a therapeutic agent or therapy that affects the immune response (*e.g.*, enhances or activates the response) and a therapeutic agent that affects (*e.g.*, inhibits or kills) the tumor/cancer cells. In some instances, combination therapies may comprise (i) recombinant RNA compositions or pharmaceutical compositions described herein; and (ii) one or more additional therapy selected from surgical removal of tumors, radiation therapy, chemotherapy, targeted therapy, and immunotherapy. In some embodiments, recombinant RNA compositions or pharmaceutical compositions described herein may be administered to a subject with a cancer prior to, concurrently with, and/or subsequently to, administration of one or more additional therapy for combination therapies. In some embodiments, the one or more additional therapy comprises 1, 2, 3, or more additional therapeutic agents or therapies.

[0161] Compositions and pharmaceutical compositions described herein can be administered to a subject using any suitable methods known in the art. Suitable formulations for use in the present invention and methods of delivery are generally well known in the art. For example, compositions described herein can be administered to the subject in a variety of ways, including parenterally, intravenously, intradermally, intramuscularly, colonically, rectally, or intraperitoneally. In some embodiments, compositions described herein is administered by an injection to a subject. For example, compositions described herein can be administered by intraperitoneal injection, intramuscular injection, subcutaneous injection, intra-tumoral injection, or intravenous injection of the subject. In some embodiments, compositions described herein can be administered by an injection to a diseased organ or a diseased tissue of a subject. In some embodiments, compositions described herein can be administered by an injection to a tumor or cancer cells in a subject. In some embodiments, compositions described herein can be administered parenterally, intravenously, intramuscularly or orally.

[0162] Any of compositions and pharmaceutical compositions described herein may be provided together with an instruction manual. The instruction manual may comprise guidance for the skilled person or attending physician how to treat (or prevent) a disease or a disorder as described herein (*e.g.*, a cancer such as a head and neck cancer) in accordance with the present invention. In some embodiments, the instruction manual may comprise guidance as to the herein described mode of delivery/administration and delivery/administration regimen,

respectively (*e.g.*, route of delivery/administration, dosage regimen, time of delivery/administration, frequency of delivery/administration, etc.). In some embodiments, the instruction manual may comprise the instruction that how compositions of the present invention is to be administrated or injected and/or is prepared for administration or injection.

- 5 In principle, what has been described herein elsewhere with respect to the mode of delivery/administration and delivery/administration regimen, respectively, may be comprised as respective instructions in the instruction manual.

[0163] Compositions and pharmaceutical compositions described herein can be used in a gene therapy. In certain embodiments, compositions comprising recombinant polynucleic acids or RNA constructs described herein can be delivered to a cell in gene therapy vectors. Gene therapy vectors and methods of gene delivery are well known in the art. Non-limiting examples of these methods include viral vector delivery systems including DNA and RNA viruses, which have either episomal or integrated genomes after delivery to the cell, non-viral vector delivery systems including DNA plasmids, naked nucleic acid, and nucleic acid
10 complexed with a delivery vehicle, transposon system (for delivery and integration into the host genomes; Moriarity, et al. (2013) Nucleic Acids Res 41(8), e92, Aronovich, et al., (2011) Hum. Mol. Genet. 20(R1), R14-R20), retrovirus-mediated DNA transfer (*e.g.*, Moloney Mouse Leukemia Virus, spleen necrosis virus, retroviruses such as Rous Sarcoma Virus, Harvey Sarcoma Virus, avian leukosis virus, gibbon ape leukemia virus, human
15 immunodeficiency virus, adenovirus, Myeloproliferative Sarcoma Virus, and mammary tumor virus; see *e.g.*, Kay et al. (1993) Science 262, 117-119, Anderson (1992) Science 256, 808-813), and DNA virus-mediated DNA transfer including adenovirus, herpes virus, parvovirus and adeno-associated virus (*e.g.*, Ali et al. (1994) Gene Therapy 1, 367-384). Viral vectors also include but are not limited to adeno-associated virus, adenoviral virus, lentivirus,
20 retroviral, and herpes simplex virus vectors. Vectors capable of integration in the host genome include but are not limited to retrovirus or lentivirus.

[0164] In some embodiments, compositions comprising recombinant polynucleic acid or RNA constructs described herein can be delivered to a cell via direct DNA transfer (Wolff et al. (1990) Science 247, 1465-1468). Recombinant polynucleic acid or RNA constructs can be
30 delivered to cells following mild mechanical disruption of the cell membrane, temporarily permeabilizing the cells. Such a mild mechanical disruption of the membrane can be accomplished by gently forcing cells through a small aperture (Sharei et al. PLOS ONE (2015) 10(4), e0118803). In another embodiment, compositions comprising recombinant polynucleic acid or RNA constructs described herein can be delivered to a cell via liposome-

mediated DNA transfer (*e.g.*, Gao & Huang (1991) Biochem. Biophys. Res. Comm. 179, 280-285, Crystal (1995) Nature Med. 1, 15-17, Caplen et al. (1995) Nature Med. 3, 39-46). A liposome can encompass a variety of single and multilamellar lipid vehicles formed by the generation of enclosed lipid bilayers or aggregates. Recombinant polynucleic acid or RNA constructs can be encapsulated in the aqueous interior of a liposome, interspersed within the lipid bilayer of a liposome, attached to a liposome via a linking molecule that is associated with both the liposome and the oligonucleotide, entrapped in a liposome, or complexed with a liposome.

10 **Modulation of gene expression**

[0165] Provided herein are methods of simultaneously expressing an siRNA and an mRNA from a single RNA transcript in a cell, comprising introducing into the cell compositions comprising any recombinant polynucleic acid or RNA constructs described herein. Further provided herein are methods of simultaneously modulating expression of two or more genes in a cell, comprising introducing into the cell compositions comprising recombinant polynucleic acid or RNA constructs encoding or comprising a first RNA linked to a second RNA, wherein the first RNA encodes a gene of interest, and wherein the second RNA encodes a small interfering RNA (siRNA) capable of binding to a target messenger RNA (mRNA); wherein the target mRNA is different from an mRNA encoded by the gene of interest, and wherein the expression of the target mRNA and the gene of interest is modulated simultaneously. In some instances, expression of a polynucleic acid, gene, DNA, or RNA, as used herein, can refer to transcription and/or translation of the polynucleic acid, gene, DNA, or RNA. In some instances, modulating, increasing upregulating decreasing or downregulating expression of a polynucleic acid, gene such as a gene of interest, DNA, or RNA such as a target mRNA, as used herein, can refer to modulating, increasing, upregulating, decreasing, downregulating the level of protein encoded by a polynucleic acid, gene such as a gene of interest, DNA, or RNA such as a target mRNA by affecting transcription and/or translation of the polynucleic acid, gene such as a gene of interest, DNA, or RNA such as a target mRNA. In some instances, inhibiting expression of a polynucleic acid, gene such as a gene of interest, DNA, or RNA such as a target mRNA can refer to affect transcription and/or translation of the polynucleic acid, gene such as a gene of interest, DNA, or RNA such as a target mRNA such that the level of protein encoded by the polynucleic acid, gene such as a gene of interest, DNA, or RNA such as a target mRNA is reduced or abolished.

- [0166] For example, provided herein, are methods of simultaneously modulating expression of two or more genes in a cell, comprising introducing into the cell compositions comprising recombinant polynucleic acid or RNA constructs encoding or comprising a first RNA linked to a second RNA, wherein the first RNA encodes a cytokine, and wherein the second RNA encodes a small interfering RNA (siRNA) capable of binding to an mRNA associated with tumor proliferation, angiogenesis, or recognition by the immune system; wherein the expression of the mRNA of which the protein product is associated with tumor proliferation, angiogenesis, or recognition by the immune system and the cytokine is modulated simultaneously.
- 5 [0167] Provided herein are methods of simultaneously modulating expression of two or more genes in a cell, comprising introducing into the cell compositions comprising recombinant polynucleic acid or RNA constructs comprising a first RNA linked to a second RNA wherein the first RNA encodes IL-2, and wherein the second RNA encodes a small interfering RNA (siRNA) capable of binding to a VEGFA mRNA; wherein the expression of IL-2 and VEGFA is modulated simultaneously, i.e. the expression of IL-2 is upregulated and the expression of VEGFA is downregulated simultaneously. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise at least 1, 2, 3, 4, 5, or more siRNAs. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each directed to the same region of a VEGFA mRNA. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each directed to a different region of a VEGFA mRNA. In related aspects, each of the at least 3 siRNAs is directed to the same, different, or a combination thereof. In related aspects, recombinant polynucleic acid constructs may comprise a sequence comprising in SEQ ID NO: 86 (Cpd.5), SEQ ID NO: 88 (Cpd.7), SEQ ID NO: 89 (Cpd.8), SEQ ID NO: 90 (Cpd.7), or SEQ ID NO: 91 (Cpd.10). In related aspects, recombinant RNA constructs may comprise a sequence comprising in SEQ ID NO: 5 (Cpd.5), SEQ ID NO: 7 (Cpd.7), SEQ ID NO: 8 (Cpd.8), SEQ ID NO: 9 (Cpd.9), SEQ ID NO: 10 (Cpd.10), SEQ ID NO: 129 (Cpd.5), SEQ ID NO: 131 (Cpd.7), SEQ ID NO: 132 (Cpd.8), SEQ ID NO: 133 (Cpd.9), or SEQ ID NO: 134 (Cpd.10).
- 20 [0168] Also provided herein are methods of simultaneously modulating expression of two or more genes in a cell, comprising introducing into the cell compositions comprising recombinant polynucleic acid or RNA constructs encoding or comprising a first RNA linked to a second RNA wherein the first RNA encodes IL-2, and wherein the second RNA encodes a small interfering RNA (siRNA) capable of binding to an mRNA of a VEGFA isoform; wherein the expression of IL-2 and an isoform of VEGFA is modulated simultaneously, i.e.
- 25 30

the expression of IL-2 is upregulated and the expression of an isoform of VEGFA is downregulated simultaneously. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise at least 1, 2, 3, 4, 5, or more siRNAs. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each
5 directed to the same region of an mRNA of a VEGFA isoform. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each directed to a different region of an mRNA of a VEGFA isoform. In related aspects, each of the at least 3 siRNAs is directed to the same, different, or a combination thereof.

[0169] Further provided herein are methods of simultaneously modulating expression of two
10 or more genes in a cell, comprising introducing into the cell compositions comprising recombinant polynucleic acid or RNA constructs encoding or comprising a first RNA linked to a second RNA wherein the first RNA encodes IL-2, and wherein the second RNA encodes a small interfering RNA (siRNA) capable of binding to a PIGF mRNA; wherein the expression of IL-2 and PIGF is modulated simultaneously, i.e. the expression of IL-2 is
15 upregulated and the expression of PIGF is downregulated simultaneously. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise at least 1, 2, 3, 4, 5, or more siRNAs. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each directed to the same region of a PIGF mRNA. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs,
20 each directed to a different region of a PIGF mRNA. In related aspects, each of the at least 3 siRNAs is directed to the same, different, or a combination thereof.

[0170] Provided herein are methods of simultaneously modulating expression of two or more genes in a cell, comprising introducing into the cell compositions comprising recombinant polynucleic acid or RNA constructs encoding or comprising a first RNA linked to a second
25 RNA wherein the first RNA encodes IL-2, and wherein the second RNA encodes a small interfering RNA (siRNA) capable of binding to a MICA and/or MICB (MICA/B) mRNA; wherein the expression of IL-2 and MICA/B is modulated simultaneously, i.e. the expression of IL-2 is upregulated and the expression of MICA/B is downregulated simultaneously. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise at
30 least 1, 2, 3, 4, 5, or more siRNAs. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each directed to the same region of a MICA/B mRNA. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each directed to a different region of a MICA/B mRNA. In related aspects, each of the at least 3 siRNAs is directed to the same, different, or a combination

thereof. In related aspects, recombinant polynucleic acid constructs may comprise a sequence comprising in SEQ ID NO: 87 (Cpd.6). In related aspects, recombinant RNA constructs may comprise a sequence comprising in SEQ ID NO: 6 or 130 (Cpd.6).

[0171] Also provided herein are methods of simultaneously modulating expression of two or more genes in a cell, comprising introducing into the cell compositions comprising recombinant polynucleic acid or RNA constructs encoding or comprising a first RNA linked to a second RNA wherein the first RNA encodes IL-2, and wherein the second RNA encodes a small interfering RNA (siRNA) capable of binding to an mRNA of ERp5, ADAM, or MMP; wherein the expression of IL-2 and ERp5, ADAM, or MMP is modulated simultaneously, i.e. the expression of IL-2 is upregulated and the expression of ERp5, ADAM, or MMP is downregulated simultaneously. In some embodiments, the ADAM is ADAM17. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise at least 1, 2, 3, 4, 5, or more siRNAs. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each directed to the same region of an mRNA of ERp5, ADAM17, or MMP. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each directed to a different region of an mRNA of ERp5, ADAM17, or MMP. In related aspects, each of the at least 3 siRNAs is directed to the same, different, or a combination thereof.

[0172] Provided herein are methods of simultaneously modulating expression of two or more genes in a cell, comprising introducing into the cell compositions comprising recombinant polynucleic acid or RNA constructs comprising a first RNA linked to a second RNA wherein the first RNA encodes IL-12, and wherein the second RNA encodes a small interfering RNA (siRNA) capable of binding to an mRNA of IDH1, CDK4, and/or CDK6; wherein the expression of IL-12, IDH1, CDK4, and/or CDK6 is modulated simultaneously, i.e. the expression of IL-12 is upregulated and the expression of IDH1, CDK4, and/or CDK6 is downregulated simultaneously. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise at least 1, 2, 3, 4, 5, or more siRNAs. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each directed to the same region of an mRNA of IDH1, CDK4, and/or CDK6. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each directed to a different region of an mRNA of IDH1, CDK4, and/or CDK6. In related aspects, each of the at least 3 siRNAs is directed to the same, different, or a combination thereof. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 1 siRNA directed to an mRNA of IDH1, 1 siRNA directed to an mRNA of CDK4, and 1 siRNA

directed to an mRNA of CDK6. In related aspects, recombinant polynucleic acid constructs may comprise a sequence comprising in SEQ ID NO: 92 (Cpd.11). In related aspects, recombinant RNA constructs may comprise a sequence comprising in SEQ ID NO: 11 or 135 (Cpd.11).

5 [0173] Provided herein are methods of simultaneously modulating expression of two or more genes in a cell, comprising introducing into the cell compositions comprising recombinant polynucleic acid or RNA constructs comprising a first RNA linked to a second RNA wherein the first RNA encodes IL-12, and wherein the second RNA encodes a small interfering RNA (siRNA) capable of binding to an mRNA of EGFR, mTOR, and/or KRAS; wherein the
10 expression of IL-12, EGFR, mTOR, and/or KRAS is modulated simultaneously, i.e. the expression of IL-12 is upregulated and the expression of EGFR, mTOR, and/or KRAS is downregulated simultaneously. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise at least 1, 2, 3, 4, 5, or more siRNAs. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each
15 directed to the same region of an mRNA of EGFR, mTOR, and/or KRAS. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each directed to a different region of an mRNA of EGFR, mTOR, and/or KRAS. In related aspects, each of the at least 3 siRNAs is directed to the same, different, or a combination thereof. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 1
20 siRNA directed to an mRNA of EGFR, 1 siRNA directed to an mRNA of mTOR, and 1 siRNA directed to an mRNA of KRAS. In related aspects, recombinant polynucleic acid constructs may comprise a sequence comprising in SEQ ID NO: 93 (Cpd.12), SEQ ID NO: 94 (Cpd.13), or SEQ ID NO: 95 (Cpd.14). In related aspects, recombinant RNA constructs may comprise a sequence comprising in SEQ ID NO: 12 (Cpd.12), SEQ ID NO: 13 (Cpd.13), SEQ
25 ID NO: 14 (Cpd.14), SEQ ID NO: 136 (Cpd.12), SEQ ID NO: 137 (Cpd.13), or SEQ ID NO: 138 (Cpd.14).

[0174] Provided herein are methods of simultaneously modulating expression of two or more genes in a cell, comprising introducing into the cell compositions comprising recombinant polynucleic acid or RNA constructs comprising a first RNA linked to a second RNA wherein
30 the first RNA encodes IL-15, and wherein the second RNA encodes a small interfering RNA (siRNA) capable of binding to an mRNA of VEGFA and/or CD155; wherein the expression of IL-15, VEGFA, and/or CD155 is modulated simultaneously, i.e. the expression of IL-15 is upregulated and the expression of VEGFA and/or CD155 is downregulated simultaneously. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise at

least 1, 2, 3, 4, 5, or more siRNAs. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each directed to the same region of an mRNA of VEGFA and/or CD155. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each directed to a different region of an mRNA of VEGFA and/or CD155. In related aspects, each of the at least 3 siRNAs is directed to the same, different, or a combination thereof. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 1 siRNA directed to an mRNA of VEGFA and 2 siRNAs directed to an mRNA of CD155. In related aspects, recombinant polynucleic acid constructs may comprise a sequence comprising in SEQ ID NO: 96 (Cpd.15). In related aspects, recombinant RNA constructs may comprise a sequence comprising in SEQ ID NO: 15 or 139 (Cpd.15).

[0175] Provided herein are methods of simultaneously modulating expression of two or more genes in a cell, comprising introducing into the cell compositions comprising recombinant polynucleic acid or RNA constructs comprising a first RNA linked to a second RNA wherein the first RNA encodes IL-15, and wherein the second RNA encodes a small interfering RNA (siRNA) capable of binding to an mRNA of VEGFA, PD-L1, and/or c-Myc; wherein the expression of IL-15, VEGFA, PD-L1, and/or c-Myc is modulated simultaneously, i.e. the expression of IL-15 is upregulated and the expression of VEGFA, PD-L1, and/or c-Myc is downregulated simultaneously. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise at least 1, 2, 3, 4, 5, or more siRNAs. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each directed to the same region of an mRNA of VEGFA, PD-L1, and/or c-Myc. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each directed to a different region of an mRNA of VEGFA, PD-L1, and/or c-Myc. In related aspects, each of the at least 3 siRNAs is directed to the same, different, or a combination thereof. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 1 siRNA directed to an mRNA of VEGFA, 1 siRNA directed to an mRNA of PD-L1, and 1 siRNA directed to an mRNA of c-Myc. In related aspects, recombinant polynucleic acid constructs may comprise a sequence comprising in SEQ ID NO: 97 (Cpd.16). In related aspects, recombinant RNA constructs may comprise a sequence comprising in SEQ ID NO: 16 or 140 (Cpd.16).

[0176] Provided herein are methods of simultaneously modulating expression of two or more genes in a cell, comprising introducing into the cell compositions comprising recombinant polynucleic acid or RNA constructs comprising a first RNA linked to a second RNA wherein

the first RNA encodes IL-7, and wherein the second RNA encodes a small interfering RNA (siRNA) capable of binding to a PD-L1 mRNA; wherein the expression of IL-7 and PD-L1 is modulated simultaneously, i.e. the expression of IL-7 is upregulated and the expression of PD-L1 is downregulated simultaneously. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise at least 1, 2, 3, 4, 5, or more siRNAs. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each directed to the same region of a PD-L1 mRNA. In related aspects, recombinant polynucleic acid or RNA constructs may encode or comprise 3 siRNAs, each directed to a different region of a PD-L1 mRNA. In related aspects, each of the at least 3 siRNAs is directed to the same, different, or a combination thereof. In related aspects, recombinant polynucleic acid constructs may comprise a sequence comprising in SEQ ID NO: 98 (Cpd.17). In related aspects, recombinant RNA constructs may comprise a sequence comprising in SEQ ID NO: 17 or 141 (Cpd.17).

[0177] Provided herein are methods of simultaneously upregulating and downregulating expression of two or more genes in a cell, comprising introducing into the cell compositions comprising recombinant polynucleic acid or RNA constructs encoding or comprising a first RNA linked to a second RNA wherein the first RNA encodes a gene of interest (*e.g.*, IL-2, IL-12, IL-15, or IL-7), and wherein the second RNA encodes a small interfering RNA (siRNA) capable of binding to a target mRNA (*e.g.*, VEGFA, a VEGFA isoform, PIGF, MICA, MICB, ERp5, ADAM, MMP, IDH1, CDK4, CDK6, EGFR, mTOR, KRAS, CD155, PD-L1, or c-Myc); wherein the target mRNA is different from an mRNA encoded by the gene of interest, and wherein the expression of the target mRNA is downregulated and the expression of the gene of interest is upregulated simultaneously. In some embodiments, the ADAM is ADAM17. In some embodiments, the expression of the target mRNA is downregulated by the siRNA capable of binding to the target mRNA. In some embodiments, the expression of the gene of interest is upregulated by expressing an mRNA or a protein encoded by the gene of interest.

Illustrative embodiments

[0178] In some aspects, provided herein, is a composition comprising a first RNA linked to a second RNA, wherein the first RNA encodes for a cytokine, and wherein the second RNA encodes for a genetic element that modulates expression of a gene associated with tumor proliferation. In some embodiments, the cytokine is interleukin-2 (IL-2), IL-12, IL-15, IL-7, a fragment thereof, or a functional variant thereof. In some embodiments, the cytokine

comprises a sequence selected from the group consisting of SEQ ID NOs: 24, 44, 47, 68, and 80. In some embodiments, the cytokine comprises a signal peptide. In some embodiments, the signal peptide comprises an unmodified signal peptide sequence or a modified signal peptide sequence. In some embodiments, the unmodified signal peptide sequence comprises a
 5 sequence selected from the group consisting of SEQ ID NOs: 26 and 125-128. In some embodiments, the IL-2 comprises a signal peptide. In some embodiments, the signal peptide comprises an unmodified IL-2 signal peptide sequence. In some embodiments, the unmodified IL-2 signal peptide sequence comprises a sequence listed in SEQ ID NO: 26. In some embodiments, the signal peptide comprises an IL-2 signal peptide sequence modified by
 10 insertion, deletion, or substitution of at least one amino acid. In some embodiments, the IL-2 signal peptide sequence modified by insertion, deletion, or substitution of at least one amino acid comprises a sequence selected from the group consisting of SEQ ID NOs: 27-29.

[0179] In some embodiments, the first RNA is a messenger RNA (mRNA). In some embodiments, the second RNA is a small interfering RNA (siRNA). In some embodiments, the siRNA is capable of binding to an mRNA of the gene associated with tumor proliferation.
 15 In some embodiments, the second RNA comprises 1, 2, 3, 4, 5, or more species of siRNA, wherein each species of siRNA comprises a different sequence targeting a different region of the same mRNA. In some embodiments, the second RNA comprises 1, 2, 3, 4, 5, or more redundant species of siRNA. In some embodiments, each species of the 1, 2, 3, 4, 5, or more
 20 species of siRNA is connected by a linker comprising a sequence listed in SEQ ID NO: 22.

[0180] In some embodiments, the gene associated with tumor proliferation comprises a gene associated with angiogenesis. In some embodiments, the gene associated with angiogenesis encodes vascular endothelial growth factor (VEGF), a fragment thereof, or a functional variant thereof. In some embodiments, the VEGF is VEGFA, a fragment thereof, or a
 25 functional variant thereof. In some embodiments, the VEGFA comprises a sequence listed in SEQ ID NO: 35. In some embodiments, the VEGF is an isoform of VEGFA, a fragment thereof, or a functional variant thereof. In some embodiments, the VEGF is placental growth factor (PIGF), a fragment thereof, or a functional variant thereof. In some embodiments, the gene associated with tumor proliferation comprises isocitrate dehydrogenase (IDH1), cyclin-
 30 dependent kinase 4 (CDK4), CDK6, epidermal growth factor receptor (EGFR), mechanistic target of rapamycin (mTOR), Kirsten rat sarcoma viral oncogene (KRAS), cluster of differentiation (CD155), programmed cell death-ligand 1 (PD-L1), or myc proto-oncogene (c-Myc). In some embodiments, the gene associated with tumor proliferation comprises a

sequence selected from the group consisting of SEQ ID NOs: 50, 53, 56, 59, 62, 65, 71, 74, and 77.

[0181] In some embodiments, the first RNA is linked to the second RNA by a linker. In some embodiments, the linker comprises a tRNA linker or a linker comprising a sequence listed in SEQ ID NO: 21. In some embodiments, the compositions described herein further comprises a poly(A) tail, a 5' cap, or a Kozak sequence. In some embodiments, the first RNA and the second RNA are both recombinant.

[0182] In some aspects, provided herein, is a composition comprising a first RNA linked to a second RNA, wherein the first RNA encodes for a cytokine, and wherein the second RNA encodes for a genetic element that modulates expression of a gene associated with recognition by the immune system. In some embodiments, the cytokine is interleukin-2 (IL-2), a fragment thereof, or a functional variant thereof. In some embodiments, the IL-2 comprises a sequence listed in SEQ ID NO: 24. In some embodiments, the IL-2 comprises a signal peptide. In some embodiments, the signal peptide comprises an unmodified IL-2 signal peptide sequence. In some embodiments, the unmodified IL-2 signal peptide sequence comprises a sequence listed in SEQ ID NO: 26. In some embodiments, the signal peptide comprises an IL-2 signal peptide sequence modified by insertion, deletion, or substitution of at least one amino acid. In some embodiments, the IL-2 signal peptide sequence modified by insertion, deletion, or substitution of at least one amino acid comprises a sequence selected from the group consisting of SEQ ID NOs: 27-29.

[0183] In some embodiments, the first RNA is a messenger RNA (mRNA). In some embodiments, the second RNA is a small interfering RNA (siRNA). In some embodiments, the siRNA is capable of binding to an mRNA of the gene associated with recognition by the immune system encoding for cell surface localizing protein. In some embodiments, the gene associated with recognition by the immune system encodes MHC class I chain-related sequence A (MICA), a fragment thereof, or a functional variant thereof. In some embodiments, the MICA comprises a sequence listed in SEQ ID NO: 38. In some embodiments, the gene associated with immune system surveillance encodes MHC class I chain-related sequence B (MICB), a fragment thereof, or a functional variant thereof. In some embodiments, the MICB comprises a sequence listed in SEQ ID NO: 41. In some embodiments, the gene associated with recognition by the immune system encodes endoplasmic reticulum protein (ERp5), a disintegrin and metalloproteinase (ADAM), matrix metalloproteinase (MMP), a fragment thereof, or a functional variant thereof. In some embodiments, the ADAM is ADAM17. In some embodiments, the second RNA comprises 1,

2, 3, 4, 5, or more species of siRNA, wherein each species of siRNA comprises a different sequence targeting a different region of the same mRNA. In some embodiments, the second RNA comprises 1, 2, 3, 4, 5, or more redundant species of siRNA. In some embodiments, each species of the 1, 2, 3, 4, 5, or more species of siRNA is connected by a linker comprising
5 a sequence listed in SEQ ID NO: 22.

[0184] In some embodiments, the first RNA is linked to the second RNA by a linker. In some embodiments, the linker comprises a tRNA linker or a linker comprising a sequence listed in SEQ ID NO: 21. In some embodiments, the compositions described herein further comprises a poly(A) tail, a 5' cap, or a Kozak sequence. In some embodiments, the first RNA and the
10 second RNA are both recombinant.

[0185] In some aspects, provided herein, is a composition comprising a first RNA encoding for interleukin-2 (IL-2), IL-15, a fragment thereof, or a functional variant thereof linked to a second RNA encoding for a genetic element that modulates expression of vascular endothelial growth factor A (VEGFA), an isoform of VEGFA, placental growth factor (PIGF), cluster of
15 differentiation 155 (CD155), programmed cell death-ligand 1 (PD-L1), myc proto-oncogene (c-Myc), a fragment thereof, or a functional variant thereof. In some embodiments, the first RNA is a messenger RNA (mRNA). In some embodiments, the IL-2 comprises a sequence listed in SEQ ID NO: 24. In some embodiments, the signal peptide comprises an unmodified IL-2 signal peptide sequence. In some embodiments, the unmodified IL-2 signal peptide
20 sequence comprises a sequence listed in SEQ ID NO: 26. In some embodiments, the signal peptide comprises an IL-2 signal peptide sequence modified by insertion, deletion, or substitution of at least one amino acid. In some embodiments, the IL-2 signal peptide sequence modified by insertion, deletion, or substitution of at least one amino acid comprises a sequence selected from the group consisting of SEQ ID NOs: 27-29. In some embodiments,
25 the IL-15 comprises a sequence comprising SEQ ID NO: 68. In some embodiments, the IL-15 comprises a signal peptide. In some embodiments, the signal peptide comprises an unmodified IL-15 signal peptide sequence. In some embodiments, the unmodified IL-15 signal peptide sequence comprises a sequence listed in SEQ ID NO: 144.

[0186] In some embodiments, the second RNA is a small interfering RNA (siRNA). In some
30 embodiments, the siRNA is capable of binding to an mRNA of VEGFA, an isoform of VEGFA, PIGF, CD155, PD-L1, or c-Myc. In some embodiments, the VEGFA comprises a sequence listed in SEQ ID NO: 35. In some embodiments, the CD155 comprises a sequence comprising SEQ ID NO: 71. In some embodiments, the PD-L1 comprises a sequence comprising SEQ ID NO: 74. In some embodiments, the c-Myc comprises a sequence

comprising SEQ ID NO: 77. In some embodiments, the second RNA comprises 1, 2, 3, 4, 5, or more species of siRNA, wherein each species of siRNA comprises a different sequence targeting a different region of the same mRNA. In some embodiments, the second RNA comprises 1, 2, 3, 4, 5, or more redundant species of siRNA. In some embodiments, each
5 species of the 1, 2, 3, 4, 5, or more species of siRNA is connected by a linker comprising a sequence listed in SEQ ID NO: 22.

[0187] In some embodiments, the first RNA is linked to the second RNA by a linker. In some embodiments, the linker comprises a tRNA linker or a linker comprising a sequence listed in SEQ ID NO: 21. In some embodiments, the compositions described herein further comprises
10 a poly(A) tail, a 5' cap, or a Kozak sequence. In some embodiments, the first RNA and the second RNA are both recombinant.

[0188] In some aspects, provided herein, is a composition comprising a first RNA encoding for interleukin-2 (IL-2), a fragment thereof, or a functional variant thereof linked to a second RNA encoding for a genetic element that modulates expression of MHC class I chain-related
15 sequence A (MICA), MHC class I chain-related sequence B (MICB), endoplasmic reticulum protein (ERp5), a disintegrin and metalloproteinase (ADAM), matrix metalloproteinase (MMP), a fragment thereof, or a functional variant thereof. In some embodiments, the ADAM is ADAM17. In some embodiments, the first RNA is a messenger RNA (mRNA). In some embodiments, the IL-2 comprises a sequence listed in SEQ ID NO: 24. In some embodiments,
20 the signal peptide comprises an unmodified IL-2 signal peptide sequence. In some embodiments, the unmodified IL-2 signal peptide sequence comprises a sequence listed in SEQ ID NO: 26. In some embodiments, the signal peptide comprises an IL-2 signal peptide sequence modified by insertion, deletion, or substitution of at least one amino acid. In some embodiments, the IL-2 signal peptide sequence modified by insertion, deletion, or substitution
25 of at least one amino acid comprises a sequence selected from the group consisting of SEQ ID NOs: 27-29. In some embodiments, the second RNA is a small interfering RNA (siRNA). In some embodiments, the siRNA is capable of binding to an mRNA of MICA, MICB, ERp5, ADAM, or MMP. In some embodiments, the MICA comprises a sequence listed in SEQ ID NO: 38. In some embodiments, the MICB comprises a sequence listed in SEQ ID NO: 41. In
30 some embodiments, the ADAM is ADAM17. In some embodiments, the second RNA comprises 1, 2, 3, 4, 5, or more species of siRNA, wherein each species of siRNA comprises a different sequence targeting a different region of the same mRNA. In some embodiments, the second RNA comprises 1, 2, 3, 4, 5, or more redundant species of siRNA. In some embodiments, each species of the 1, 2, 3, 4, 5, or more species of siRNA is connected by a

linker comprising a sequence listed in SEQ ID NO: 22. In some embodiments, the first RNA is linked to the second RNA by a linker. In some embodiments, the linker comprises a tRNA linker or a linker comprising a sequence listed in SEQ ID NO: 21. In some embodiments, the compositions described herein further comprises a poly(A) tail, a 5' cap, or a Kozak

5 sequence. In some embodiments, the first RNA and the second RNA are both recombinant.

[0189] In some aspects, provided herein, is a composition comprising a first RNA encoding for interleukin-12 (IL-12), IL-7, a fragment thereof, or a functional variant thereof linked to a second RNA encoding for a genetic element that modulates expression of isocitrate dehydrogenase (IDH1), cyclin-dependent kinase 4 (CDK4), CDK6, epidermal growth factor
10 receptor (EGFR), mechanistic target of rapamycin (mTOR), Kirsten rat sarcoma viral oncogene (KRAS), programmed cell death-ligand 1 (PD-L1), a fragment thereof, or a functional variant thereof.

[0190] In some embodiments, the first RNA is a messenger RNA (mRNA). In some embodiments, the IL-12 comprises a sequence comprising SEQ ID NO: 44 or SEQ ID NO:
15 47. In some embodiments, the IL-12 comprises a signal peptide. In some embodiments, the signal peptide comprises an unmodified IL-12 signal peptide. In some embodiments, the unmodified IL-12 signal peptide comprises a sequence listed in SEQ ID NO: 142 or SEQ ID NO: 143. In some embodiments, the IL-7 comprises a sequence comprising SEQ ID NO: 80. In some embodiments, the IL-7 comprises a signal peptide. In some embodiments, the signal
20 peptide comprises an unmodified IL-7 signal peptide. In some embodiments, the unmodified IL-7 signal peptide comprises a sequence listed in SEQ ID NO: 128.

[0191] In some embodiments, the second RNA is a small interfering RNA (siRNA). In some embodiments, the siRNA is capable of binding to an mRNA of IDH1, CDK4, CDK6, EGFR, mTOR, KRAS, or PD-L1. In some embodiments, IDH1 comprises a sequence comprising
25 SEQ ID NO: 50. In some embodiments, CDK4 comprises a sequence comprising SEQ ID NO: 53. In some embodiments, CDK6 comprises a sequence comprising SEQ ID NO: 56. In some embodiments, mTOR comprises a sequence comprising SEQ ID NO: 62. In some embodiments, EGFR comprises a sequence comprising SEQ ID NO: 59. In some embodiments, KRAS comprises a sequence comprising SEQ ID NO: 65. In some
30 embodiments, PD-L1 comprises a sequence comprising SEQ ID NO: 74.

[0192] In some embodiments, the second RNA comprises 1, 2, 3, 4, 5, or more species of siRNA, wherein each species of siRNA comprises a different sequence targeting a different region of the same mRNA. In some embodiments, the second RNA comprises 1, 2, 3, 4, 5, or more redundant species of siRNA. The composition of claim 119 or 120, wherein each

species of the 1, 2, 3, 4, 5, or more species of siRNA is connected by a linker comprising a sequence listed in SEQ ID NO: 22.

[0193] In some embodiments, the first RNA is linked to the second RNA by a linker. In some embodiments, the linker comprises a tRNA linker or a linker comprising a sequence

5 comprising SEQ ID NO: 21. In some embodiments, the composition further comprises a poly(A) tail, a 5' cap, or a Kozak sequence. In some embodiments, the first RNA and the second RNA are both recombinant.

[0194] In some aspects, provided herein, is a pharmaceutical composition comprising any of the compositions described herein and a pharmaceutically acceptable excipient. In some aspects, provided herein, is a method of treating cancer, comprising administering any of

10 compositions or pharmaceutical compositions described herein to a subject having a cancer. In some aspects, provided herein, are any of compositions or pharmaceutical compositions described herein for use in a method for the treatment of cancer. In some aspects, provided herein, is the use of any of compositions or pharmaceutical compositions described herein for the manufacture of a medicament for treating cancer. In some aspects, provided herein, is the use of any of compositions or pharmaceutical compositions described herein for treating cancer in a subject. In some embodiments, the cancer is a solid tumor. In some embodiments, the cancer is melanoma. In some embodiments, the cancer is renal cell carcinoma. In some embodiments, the cancer is a head and neck cancer. In some embodiments, the head and neck cancer is head and neck squamous cell carcinoma. In some embodiments, the head and neck cancer is laryngeal cancer, hypopharyngeal cancer, tonsil cancer, nasal cavity cancer, paranasal sinus cancer, nasopharyngeal cancer, metastatic squamous neck cancer with occult primary, lip cancer, oral cancer, oral cancer, oropharyngeal cancer, salivary gland cancer, brain tumors, esophageal cancer, eye cancer, parathyroid cancer, sarcoma of the head and neck, or thyroid cancer. In some embodiments, the cancer is located at an upper aerodigestive tract. In some embodiments, the upper aerodigestive tract comprises a paranasal sinus, a nasal cavity, an oral cavity, a salivary gland, a tongue, a nasopharynx, an oropharynx, a hypopharynx, or a larynx. In some embodiments, the subject has a head and neck cancer. In some embodiments, the subject having the head and neck cancer has a history of tobacco usage. In some embodiments, the subject having the head and neck cancer has a human papillomavirus (HPV) DNA. In some embodiments, the subject is a human.

[0195] In some aspects, provided herein, is a composition comprising a recombinant polynucleic acid construct comprising a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1-17 and 125-141.

[0196] In some aspects, provided herein, is a composition for use in modulating the expression of two or more genes in a cell. In some aspects, provided herein is a cell comprising any one of the compositions described herein. In some aspects, provided herein is a vector comprising a recombinant polynucleic acid construct encoding any one of the compositions described herein.

[0197] In some aspects, provided herein is a method of producing an siRNA and an mRNA from a single RNA transcript in a cell, comprising introducing into the cell any one of the compositions described herein or the vectors described herein. In some aspects, provided herein is a method of modulating protein expression comprising introducing any one of the compositions described herein or the vectors described herein into a cell, wherein the expression of a protein encoded by the second RNA is decreased compared to a cell without the composition or vector. In some aspects, provided herein is a method of modulating protein expression comprising introducing any one of the compositions described herein or the vectors described herein into a cell, wherein the expression of a protein encoded by the first RNA is increased compared to a cell without the composition or vector. In some aspects, provided herein is a method of modulating protein expression comprising introducing any one of the compositions described herein or the vectors described herein into a cell, wherein the expression of a protein encoded by the second RNA is decreased compared to a cell without the composition or vector, and wherein the expression of a protein encoded by the first RNA is increased compared to a cell without the composition or vector.

EXAMPLES

[0198] These examples are provided for illustrative purposes only and not to limit the scope of the claims provided herein.

[0199] Example 1: Construct design, sequence, and synthesis

5 [0200] Construct Design

[0201] Both siRNAs and genes of interest are simultaneously expressed from a single transcript generated by *in vitro* transcription (SEQ ID NOs: 1-17 and 125-141).

Polynucleotide or RNA constructs are engineered to include siRNA designs described in Cheng, et al. (2018) J. Mater. Chem. B., 6, 4638-4644, and further comprising one or more
10 gene of interest downstream or upstream of the siRNA sequence (an example of one orientation is shown in **Fig. 1**). Recombinant constructs may encode or comprise more than one siRNA sequence targeting the same or different target mRNA. Likewise, constructs may comprise nucleic acid sequences of two or more genes of interest. A linker sequence may be present between any two elements of the constructs (*e.g.*, tRNA linker or adapted sequence
15 described by Cheng, et al. 2018).

[0202] A polynucleic acid construct may comprise a T7 promoter sequence (5' TAATACGACTCACTATA 3'; SEQ ID NO: 18) upstream of the gene of interest sequence, for RNA polymerase binding and successful *in vitro* transcription of both the gene of interest and siRNA in a single transcript. An alternative promoter *e.g.*, SP6, T3, P60, Syn5, and KP34
20 may be used. A transcription template is generated by PCR to produce mRNA, using primers designed to flank the T7 promoter, gene of interest, and siRNA sequences. The reverse primer includes a stretch of thymidine (T) base (120) (SEQ ID NO: 154) to add the 120 bp length of poly(A) tail (SEQ ID NO: 153) to the mRNA.

[0203] Construct Synthesis

25 [0204] The constructs as shown in **Table 1** (Compound ID numbers Cpd.1-Cpd.17) were synthesized by GeneArt, Germany (Thermo Fisher Scientific) as vectors containing a T7 RNA polymerase promoter (pMX, *e.g.*, pMA-T, pMK-RQ or pMA-RQ), with codon optimization (GeneOptimizer algorithm). **Table 1** shows, for each compound (Cpd.), protein encoding, signal peptide nature, the number of siRNAs of the construct and the protein to be
30 downregulated through siRNA binding to the corresponding mRNA. The sequences of each construct are shown in **Table 2** and annotated as indicated below the table (SEQ ID 1-17).

[0205] Table 1. Summary of Compounds 1-17

Compound ID	gene of interest	Signal peptide	# of siRNAs	siRNA Target	Mechanism
Cpd.1	IL-2	Endogenous	NA	NA	Anti-tumor activity

Compound ID	gene of interest	Signal peptide	# of siRNAs	siRNA Target	Mechanism
Cpd.2	IL-2	Modified	NA	NA	Anti-tumor activity
Cpd.3	IL-2	Modified	NA	NA	Anti-tumor activity
Cpd.4	IL-2	Modified	NA	NA	Anti-tumor activity
Cpd.5	IL-2	Endogenous	3	VEGFA	Anti-tumor activity, anti-angiogenesis
Cpd.6	IL-2	Endogenous	3	MICA/B	Anti-tumor activity, immune surveillance
Cpd.7	IL-2	Modified'	3	VEGFA	Anti-tumor, anti-angiogenesis
Cpd.8	IL-2	Modified'	5	VEGFA	Anti-tumor, anti-angiogenesis
Cpd.9	IL-2	Modified'	3	VEGFA	Anti-tumor, anti-angiogenesis
Cpd.10	IL-2	Modified'	3	VEGFA	Anti-tumor, anti-angiogenesis
Cpd.11	IL-12	Endogenous	3	IDH1/CDK4/CDK6	Immune-stimulating cytokine, tumor metabolism normalizer, cell cycle inhibitor
Cpd.12	IL-12	Endogenous	3	EGFR/mTOR/KRAS	Immune-stimulating cytokine, tumor growth inhibitor
Cpd.13	IL-12	Endogenous	3	EGFR	immune-stimulating cytokine, tumor growth inhibitor
Cpd.14	IL-12	Endogenous	3	mTOR	Immune-stimulating cytokine, tumor growth inhibitor
Cpd.15	IL-15	Endogenous	3	VEGFA/CD155/CD155	Immune-stimulating cytokine, anti-angiogenesis, inhibition of tumor immune escape
Cpd.16	IL-15	Endogenous	3	VEGFA/PD-L1/c-Myc	Immune-stimulating cytokine, anti-angiogenesis, inhibition of tumor immune escape, inhibition of tumor specific protein transcription
Cpd.17	IL-7	Endogenous	3	PD-L1	Immune-stimulating cytokine, inhibition of tumor immune escape

IL-2: Interleukin-2, VEGFA: vascular endothelial growth factor, MICA: MHC class I chain-related sequence A, MICB: MHC class I chain-related sequence B, IL-12: Interleukin-12, IDH1: Isocitrate dehydrogenase, CDK4: Cyclin-dependent kinase 4, CDK6: Cyclin-dependent kinase 6, EGFR: Epidermal growth factor receptor, mTOR: mechanistic target of rapamycin, KRAS: Kirsten rat sarcoma viral oncogene, IL-15: Interleukin-15, CD155: cluster

5

of differentiation 155 (poliovirus receptor), PD-L1: Programmed cell death - ligand 1, c-Myc: Myc proto-oncogene.

[0206] Table 2. Sequences of Compounds 1-17

SEQ ID NO	Compound	Sequence (5' to 3')
1	Compound 1	<p>GCCACCATGTACAGAATGCAGCTGCTGAGCTGTATCGCCCTGTCTCTGGCC CTGGTCACAAATAGCGCCCTACCAGCAGCAGCACCAAGAAAACACAGCTG CAACTGGAACACCTCCTGCTGGACCTGCAGATGATCCTGAACGGCATCAAC AACTACAAGAACCCCAAGCTGACCCGGATGCTGACCTTCAAGTTCTACATG CCCAAGAAGGCCACCGAGCTGAAGCACCTCCAGTGCCTGGAAGAGGAACTG AAGCCCTTGAAGAAGTGTGAATCTGGCCCAGAGCAAGAACTTCCACCTG AGGCCTAGGGACCTGATCAGCAACATCAACGTGATCGTGCTGGAAGTGA GGCAGCGAGACAACCTTCATGTGCGAGTACGCCGACGAGACAGCTACCATC GTGGAATTTCTGAACCGGTGGATCACCTTCTGCCAGAGCATCATCAGCACC CTGACCTGA</p>
125	Compound 1 RNA sequence	<p>GCCACCAUGUACAGAAUGCAGCUGCUGAGCUGUAUCGCCCUGUCUGGCC CUGGUCACAAAUAGCGCCCUACCAGCAGCAGCACCAAGAAAACACAGCUG CAACUGGAACACCUCCUGCUGGACCUGCAGAUUAUCUGAACGGCAUCAAC AAUACAAGAACCCCAAGCUGACCCGGUAGCUGACCUUACAGUUCUCAUG CCCAAGAAGGCCACCGAGCUGAAGCACCUCCAGUGCCUGGAAGAGGAACUG AAGCCCUUGGAAGAAGUGCUGAAUCUGGCCAGAGCAAGAACUCCACCUG AGGCCUAGGGACCUGAUCAGCAACAUCACGUGAUCGUGUGGAACUGAAA GGCAGCGAGACAACCUCAUGUGCGAGUACGCCGACGAGACAGCUACCAUC GUGGAUUUCUGAACCGGUGGAUCACCUUCUGCCAGAGCAUCAUCAGCACC CUGACCUGA (all Us are modified; N¹-methylpseudouridine)</p>
2	Compound 2*	<p>GCCACCATGCTGAAACTGCTGCTGCTCCTGTGTATCGCCCTGTCTCTGGCC GCCACAAATAGCGCCCTACCAGCAGCTCCACCAAGAAAACACAGCTGCAA CTGGAACATCTGCTGCTGGACCTGCAGATGATCCTGAACGGCATCAACAAC TACAAGAACCCCAAGCTGACCCGGATGCTGACCTTCAAGTTCTACATGCCC AAGAAGGCCACCGAGCTGAAGCACCTCCAGTGCCTGGAAGAGGAACGAAG CCCTTGAAGAAGTGTGAATCTGGCCCAGAGCAAGAACTTCCACCTGAGG CCTAGGGACCTGATCAGCAACATCAACGTGATCGTGCTGGAAGTGAAGGC AGCGAGACAACCTTCATGTGCGAGTACGCCGACGAGACAGCTACCATCGTG GAATTTCTGAACCGGTGGATCACCTTCTGCCAGAGCATCATCAGCACCCTG ACCTGA</p>
126	Compound 2 RNA sequence	<p>GCCACCAUGCUGAAACUGCUGCUGCUCUGUGUAUCGCCCUGUCUGGCC GCCACAAUAGCGCCCUACCAGCAGCUCCACCAAGAAAACACAGCUGCAA CUGGAACAUUCUGCUGCUGGACCUGCAGAUUAUCUGAACGGCAUCAACAAC UACAAGAACCCCAAGCUGACCCGGUAGCUGACCUUACAGUUCUACAUGCCC AAGAAGGCCACCGAGCUGAAGCACCUCCAGUGCCUGGAAGAGGAACUGAAG CCCUUGGAAGAAGUGCUGAAUCUGGCCAGAGCAAGAACUCCACCUGAGG CCUAGGGACCUGAUCAGCAACAUCACGUGAUCGUGUGGAACUGAAAGGC AGCGAGACAACCUCAUGUGCGAGUACGCCGACGAGACAGCUACCAUCGUG GAAUUUCUGAACCGGUGGAUCACCUUCUGCCAGAGCAUCAUCAGCACCUG ACCUGA (all Us are modified; N¹-methylpseudouridine)</p>
3	Compound 3*	<p>GCCACCATGTTGTTGCTGCTGCTCGCCTGTATTGCCCTGGCCTCTACAGCC GCCGCTACAAATTCTGCCCTACCAGCAGCTCCACCAAGAAAACCCAGCTG CAACTGGAACATCTGCTGCTGGACCTGCAGATGATCCTGAACGGCATCAAC AACTACAAGAACCCCAAGCTGACCCGGATGCTGACCTTCAAGTTCTACATG CCCAAGAAGGCCACCGAGCTGAAGCACCTCCAGTGCCTGGAAGAGGAACTG AAGCCCTTGAAGAAGTGTGAATCTGGCCCAGAGCAAGAACTTCCACCTG AGGCCTAGGGACCTGATCAGCAACATCAACGTGATCGTGCTGGAAGTGA GGCAGCGAGACAACCTTCATGTGCGAGTACGCCGACGAGACAGCTACCATC GTGGAATTTCTGAACCGGTGGATCACCTTCTGCCAGAGCATCATCAGCACC CTGACCTGA</p>
127	Compound 3 RNA sequence	<p>GCCACCAUGUUGUUGCUGCUGCUGGCCUGUAUUGCCCUGGCCUCUACAGCC GCCGCUACAAAUUCUGCCCUACCAGCAGCUCCACCAAGAAAACCCAGCUG CAACUGGAACAUUCUGCUGGACCUGCAGAUUAUCUGAACGGCAUCAAC</p>

SEQ ID NO	Compound	Sequence (5' to 3')
		AACUACAAGAACCCCAAGCUGACCCGGAUGCUGACCUUCAAGUUCUACAUG CCCAAGAAGGCCACCCGAGCUGAAGCACCUCAGUGCCUGGAAGAGGAACUG AAGCCCCUGGAAGAAGUGCUGAAUCUGGCCCAGAGCAAGAACUCCACCUG AGGCCUAGGGACCUGAUCAGCAACAUCACGUGAUCGUGCUGGAACUGAAA GGCAGCGAGACAACCUCAUGUGCGAGUACGCCGACGAGACAGCUACCAUC GUGGAAUUCUGAACCGGUGGAUCACCUUCUGCCAGAGCAUCAUCAGCACC CUGACCUGA (all Us are modified; N ¹ -methylpseudouridine)
4	Compound 4*	GCCACCATGTTGTTGCTGCTGCTCGCCTGTATTGCCCTGGCCTCTACAGCC CTGGTCACCAATTCTGCCCTACCAGCAGCTCCACCAAGAAAACCCAGCTG CAACTGGAACATCTGCTGCTGGACCTGCAGATGATCCTGAACGGCATCAAC AACTACAAGAACCCCAAGCTGACCCGGATGCTGACCTTCAAGTTCTACATG CCCAAGAAGGCCACCCGAGCTGAAGCACCCTCCAGTGCCTGGAAGAGGAACTG AAGCCCCTGAAGAAGTGTGAATCTGGCCCAGAGCAAGAACTTCCACCTG AGGCCTAGGGACCTGATCAGCAACATCAACGTGATCGTGTGGAACCTGAAA GGCAGCGAGACAACCTTCATGTGCGAGTACGCCGACGAGACAGCTACCATC GTGGAATTTCTGAACCGGTGGATCACCTTCTGCCAGAGCATCATCAGCACC CTGACCTGA
128	Compound 4 RNA sequence	GCCACCAUGUUGUUGCUGCUGCUGGCCUGUAUUGCCUGGCCUCUACAGCC CUGGUCACCAAUUCUGCCCCUACCAGCAGCUGCCACCAAGAAAACCCAGCUG CAACUGGAACAUCUGCUGCUGGACCUGCAGAUGAUCCUGAACGGCAUCAAC AACUACAAGAACCCCAAGCUGACCCGGAUGCUGACCUUCAAGUUCUACAUG CCCAAGAAGGCCACCCGAGCUGAAGCACCUCAGUGCCUGGAAGAGGAACUG AAGCCCCUGGAAGAAGUGCUGAAUCUGGCCCAGAGCAAGAACUCCACCUG AGGCCUAGGGACCUGAUCAGCAACAUCACGUGAUCGUGCUGGAACUGAAA GGCAGCGAGACAACCUCAUGUGCGAGUACGCCGACGAGACAGCUACCAUC GUGGAAUUCUGAACCGGUGGAUCACCUUCUGCCAGAGCAUCAUCAGCACC CUGACCUGA (all Us are modified; N ¹ -methylpseudouridine)
5	Compound 5	GCCACCATGTACAGAAATGCAGCTGCTGAGCTGTATCGCCCTGTCTCTGGCC CTGGTCACAAATAGCGCCCTACCAGCAGCAGCACCACCAAGAAAACACAGCTG CAACTGGAACACCTCCTGCTGGACCTGCAGATGATCCTGAACGGCATCAAC AACTACAAGAACCCCAAGCTGACCCGGATGCTGACCTTCAAGTTCTACATG CCCAAGAAGGCCACCCGAGCTGAAGCACCCTCCAGTGCCTGGAAGAGGAACTG AAGCCCCTGAAGAAGTGTGAATCTGGCCCAGAGCAAGAACTTCCACCTG AGGCCTAGGGACCTGATCAGCAACATCAACGTGATCGTGTGGAACCTGAAA GGCAGCGAGACAACCTTCATGTGCGAGTACGCCGACGAGACAGCTACCATC GTGGAATTTCTGAACCGGTGGATCACCTTCTGCCAGAGCATCATCAGCACC CTGACCTGAATAGTGAATCGTATTAACGTACCAACAAGAGAAATCATCAGC AAGTGGTACTTGACCACTTCGTGATGATTCTGCTTTATCTTAGAGGCATAT CCCTACGTACCAACAAGAGCTTCCTACAGCACAACAACCTGTTGTTGTGC TGTAGGAAGCTCTTTATCTTAGAGGCATATCCCTACGTACCAACAAGATCC GCAGACGTGTAATGTACTTGACATTTACACGTCTGCGGATCTTTATCTTA GAGGCATATCCCTTTTATCTTAGAGGCATATCCCT
129	Compound 5 RNA sequence	GCCACCAUGUACAGAAUGCAGCUGCUGAGCUGUAUCGCCUGUCUGGCC CUGGUCACAAUAGCGCCCUACCAGCAGCAGCACCACCAAGAAAACACAGCUG CAACUGGAACACCUCUGCUGGACCUGCAGAUGAUCCUGAACGGCAUCAAC AACUACAAGAACCCCAAGCUGACCCGGAUGCUGACCUUCAAGUUCUACAUG CCCAAGAAGGCCACCCGAGCUGAAGCACCUCAGUGCCUGGAAGAGGAACUG AAGCCCCUGGAAGAAGUGCUGAAUCUGGCCCAGAGCAAGAACUCCACCUG AGGCCUAGGGACCUGAUCAGCAACAUCACGUGAUCGUGCUGGAACUGAAA GGCAGCGAGACAACCUCAUGUGCGAGUACGCCGACGAGACAGCUACCAUC GUGGAAUUCUGAACCGGUGGAUCACCUUCUGCCAGAGCAUCAUCAGCACC CUGACCUGAAUAGUGAGUCGUUAUUAACGUACCAACAAGCAGAAUACACG AAGUGGUACUUGACCAUUCUGAUGAUUCUGCUUUUAUCUAGAGGCAUUA CCUACGUACCAACAAGAGCUUCCUACAGCACAACAACUUGUUGUUGUC UGUAGGAAGCUCUUUAUCUAGAGGCAUUAUCCUACGUACCAACAAGAUCC GCAGACGUGUAAUUGUACUUGACAUUUAACGUCUGCGGAUCUUUAUCUUA GAGGCAUUAUCCUUUAUCUAGAGGCAUUAUCCU (all Us are modified; N ¹ -methylpseudouridine)

SEQ ID NO	Compound	Sequence (5' to 3')
6	Compound 6	<p>GCCACCATGTACAGAATGCAGCTGCTGAGCTGTATCGCCCTGTCTCTGGCC CTGGTCACAAATAGCGCCCTACCAGCAGCAGCACCAGAAAACACAGCTG CAACTGGAACACCTCCTGCTGGACCTGCAGATGATCCTGAACGGCATCAAC AACTACAAGAACCCCAAGCTGACCCGGATGCTGACCTTCAAGTTCTACATG CCCAAGAAGGCCACCGAGCTGAAGCACCTCCAGTGCCTGGAAGAGGAACTG AAGCCCCCTGGAAGAAGTGTGAATCTGGCCAGAGCAAGAAGTCCACCTG AGGCCTAGGGACCTGATCAGCAACATCAACGTGATCGTGTGGAAGTGA GGCAGCGAGACAACCTTCATGTGCGAGTACGCCGACGAGACAGCTACCATC GTGGAATTTCTGAACCGGTGGATCACCTTCTGCCAGAGCATCATCAGCACC CTGACCTGAATAGTGTGCTGATTACGTACCAACAAGGAGATTAGGGTCT GTGAGATACTTGATCTCACAGACCCTAATCTCCTTTATCTTAGAGGCATAT CCCTACGTACCAACAAGATGCCATGAAGACCAAGACAACCTGTGTCTTGGT CTTCATGGCATCTTTATCTTAGAGGCATATCCCTACGTACCAACAAGCCTG ATGGGAATGGAACCTAACTTGTAGGTTCCATTCCCATCAGGCTTTATCTTA GAGGCATATCCCTTTATCTTAGAGGCATATCCCT</p>
130	Compound 6 RNA sequence	<p>GCCACCAUGUACAGAAUGCAGCUGCUGAGCUGUAUCGCCCUUGUCUGGCC CUGGUCACAAUAGCGCCCUACCAGCAGCAGCACCAGAAAACACAGCUG CAACUGGAACACCUCUGCUGGACCUGCAGAUAGUCCUGAACGGCAUCAAC AACUACAAGAACCCCAAGCUGACCCGGAUGCUGACCUUCAAGUUCUACAUG CCCAAGAAGGCCACCGAGCUGAAGCACCUCAGUGCCUGGAAGAGGAACUG AAGCCCCUGGAAGAAGUGCUGAAUCUGGCCAGAGCAAGAACUCCACCUG AGGCCUAGGGACCUGAUCAGCAACAUCACGUGAUCGUGUGGAACUGAAA GGCAGCGAGACAACCUCAUGUGCGAGUACGCCGACGAGACAGCUACCAUC GUGGAUUUCUGAACCGGUGGAUCACCUUCUGCCAGAGCAUCAUCAGCACC CUGACCUGAAUAGUGAGUCGUUUAACGUACCAACAAGGAGAUUAGGGUCU GUGAGAUACUUGAUCUCACAGACCCUAAUCUCCUUUAUCUUAGAGGCAUUA CCUACGUACCAACAAGAUGCCAUGAAGACCAAGACAACUUGUGUCUUGGU CUUCAUGGCAUCUUUAUCUUAGAGGCAUUAUCCUACGUACCAACAAGCCUG AUGGGAUUGGAACCUAACUUGUAGGUUCCAUCCCAUCAGGCUUUUAUCUUA GAGGCAUUAUCCCUUUUAUCUUAGAGGCAUUAUCCU (all Us are modified; N¹-methylpseudouridine)</p>
7	Compound 7	<p>GCCACCATGTTGTTGCTGCTGCTCGCCTGTATTGCCCTGGCCTCTACAGCC GCCGCTACAAATTCTGCCCTACCAGCAGCTCCACCAAGAAAACCCAGCTG CAACTGGAACATCTGCTGCTGGACCTGCAGATGATCCTGAACGGCATCAAC AACTACAAGAACCCCAAGCTGACCCGGATGCTGACCTTCAAGTTCTACATG CCCAAGAAGGCCACCGAGCTGAAGCACCTCCAGTGCCTGGAAGAGGAACTG AAGCCCCCTGGAAGAAGTGTGAATCTGGCCAGAGCAAGAAGTCCACCTG AGGCCTAGGGACCTGATCAGCAACATCAACGTGATCGTGTGGAAGTGA GGCAGCGAGACAACCTTCATGTGCGAGTACGCCGACGAGACAGCTACCATC GTGGAATTTCTGAACCGGTGGATCACCTTCTGCCAGAGCATCATCAGCACC CTGACCTGAATAGTGTGCTGATTACGTACCAACAAGCAGAATCATCAGC AAGTGGTACTTGACCACTTCGTGATGATTCTGCTTTATCTTAGAGGCATAT CCCTACGTACCAACAAGAGCTTCCTACAGCACAACAACCTGTGTGTGTC TGTAGGAAGCTCTTTATCTTAGAGGCATATCCCTACGTACCAACAAGATCC GCAGACGTGTAAATGTACTTGACATTTACACGTCTGCGGATCTTTATCTTA GAGGCATATCCCTTTATCTTAGAGGCATATCCCT</p>
131	Compound 7 RNA sequence	<p>GCCACCAUGUUGUUGCUGCUGCUGGCCUGUAUUGCCUGGCCUCUACAGCC GCCGCUACAAUUCUGCCCUACCAGCAGCAGCACCAGAAAACCCAGCUG CAACUGGAACAUUCUGCUGGACCUGCAGAUAGUCCUGAACGGCAUCAAC AACUACAAGAACCCCAAGCUGACCCGGAUGCUGACCUUCAAGUUCUACAUG CCCAAGAAGGCCACCGAGCUGAAGCACCUCAGUGCCUGGAAGAGGAACUG AAGCCCCUGGAAGAAGUGCUGAAUCUGGCCAGAGCAAGAACUCCACCUG AGGCCUAGGGACCUGAUCAGCAACAUCACGUGAUCGUGUGGAACUGAAA GGCAGCGAGACAACCUCAUGUGCGAGUACGCCGACGAGACAGCUACCAUC GUGGAUUUCUGAACCGGUGGAUCACCUUCUGCCAGAGCAUCAUCAGCACC CUGACCUGAAUAGUGAGUCGUUUAACGUACCAACAAGCAGAAUCAUCAGC AAGUGGUACUUGACCACUUCUGAUGAUUCUGCUUUUAUCUUAGAGGCAUUA CCUACGUACCAACAAGAGCUUCCUACAGCACAACAACUUGUGUGUGGC UGUAGGAAGCUCUUUAUCUUAGAGGCAUUAUCCUACGUACCAACAAGAUCC GCAGACGUGUAAUGUACUUGACAUUUACACGUCUGCGGAUCUUUAUCUUA GAGGCAUUAUCCCUUUUAUCUUAGAGGCAUUAUCCU</p>

SEQ ID NO	Compound	Sequence (5' to 3')
		(all Us are modified; N ¹ -methylpseudouridine)
8	Compound 8	<p>GCCACCATGTTGTTGCTGCTGCTCGCCTGTATTGCCCTGGCCTCTACAGCC GCCGCTACAAATTCTGCCCTACCAGCAGCTCCACCAAGAAAACCCAGCTG CAACTGGAACATCTGCTGCTGGACCTGCAGATGATCCTGAACGGCATCAAC AACTACAAGAACCCCAAGCTGACCCGGATGCTGACCTTCAAGTTCTACATG CCCAAGAAGGCCACCGAGCTGAAGCACCTCCAGTGCCTGGAAGAGGAAGT AAGCCCCGGAAGAAGTGTGAATCTGGCCCAGAGCAAGAACTTCCACCTG AGGCTTAGGGACCTGATCAGCAACATCAACGTGATCGTGTGGAAGTGAAA GGCAGCGAGACAACCTTCATGTGCGAGTACGCCGACGAGACAGCTACCATC GTGGAATTTCTGAACCGGTGGATCACCTTCTGCCAGAGCATCATCAGCACC CTGACCTGAATAGTGAGTCGTATTAACTACCAACAAGCAGAAATCATCAGC AAGTGGTACTTGACCACTTCGTGATGATTCTGCTTTATCTTAGAGGCATAT CCCTACGTACCAACAAGAGCTTCCTACAGCACAACTTGTTGTTGTGC TGTAGGAAGCTCTTTATCTTAGAGGCATATCCCTACGTACCAACAAGATCC GCAGACGTGTAATGTACTTGACATTTACACGCTCTGCGGATCTTTATCTTA GAGGCATATCCCTACGTACCAACAAGCGCAAGAAATCCCGGTATAAACTTG TTATACCGGGATTTCTTGCCTTTATCTTAGAGGCATATCCCTACGTACCA ACAAGGGCAGGACGCTTGAGTTAAACTTGTTTAACTCAAGCTGCCTCGCC TTTATCTTAGAGGCATATCCCTTTATCTTAGAGGCATATCCCT</p>
132	Compound 8 RNA sequence	<p>GCCACCAUGUUGUGCUGCUGCUGCCUGUAUUGCCUGGCCUCUACAGCC GCCGCUACAAAUUCUGCCCCUACCAGCAGCUCACCAAGAAAACCCAGCUG CAACUGGAACAUCUGCUGCUGGACCUGCAGAUGAUCCUGAACGGCAUCAAC AACUACAAGAACCCCAAGCUGACCCGGAUGCUGACCUUACAAGUUCUACAUG CCCAAGAAGGCCACCGAGCUGAAGCACCUCAGUGCCUGGAAGAGGAACUG AAGCCCCUGGAAGAAGUGCUGAAUCUGGCCAGAGCAAGAACUCCACCUG AGGCCUAGGGACCUGAUCAGCAACAUCAACGUGAUCUGCUGGAACUGAAA GGCAGCGAGACAACCUCAUGUGCGAGUACGCCGACGAGACAGCUACCAUC GUGGAUUUCUGAACCGGUGGAUACCUUCUGCCAGAGCAUCAUCAGCACC CUGACCUGAUAUGAGUGCUGUAUUAACGUACCAACAAGCAGAAUACAGC AAGUGGUACUUGACCACTUUCGUGAUGAUUCUGCUUUAUCUAGAGGCAU CCCUACGUACCAACAAGAGCUUCCUACAGCACAAACUUGUUGUUGUGC UGUAGGAAGCUCUUUUAUCUAGAGGCAUUAUCCCUACGUACCAACAAGAUCC GCAGACGUGUAAAUGUACUUGACAUUUACAGCUGCGGAUCUUUUAUCUUA GAGGCAUAUCCCUACGUACCAACAAGCGCAAGAAAUCCCGUAUAAACUUG UUUAUACGGGAUUUCUUGCGCUUUAUCUAGAGGCAUUAUCCCUACGUACCA ACAAGGGCAGGACGCUUGAGUUAUAAACUUGUUUAACUCAAGCUGCCUGCC UUUAUCUAGAGGCAUUAUCCCUUUUAUCUAGAGGCAUUAUCCCU (all Us are modified; N¹-methylpseudouridine)</p>
9	Compound 9	<p>GCCACCATGTTGTTGCTGCTGCTCGCCTGTATTGCCCTGGCCTCTACAGCC GCCGCTACAAATTCTGCCCTACCAGCAGCTCCACCAAGAAAACCCAGCTG CAACTGGAACATCTGCTGCTGGACCTGCAGATGATCCTGAACGGCATCAAC AACTACAAGAACCCCAAGCTGACCCGGATGCTGACCTTCAAGTTCTACATG CCCAAGAAGGCCACCGAGCTGAAGCACCTCCAGTGCCTGGAAGAGGAAGT AAGCCCCGGAAGAAGTGTGAATCTGGCCCAGAGCAAGAACTTCCACCTG AGGCTTAGGGACCTGATCAGCAACATCAACGTGATCGTGTGGAAGTGAAA GGCAGCGAGACAACCTTCATGTGCGAGTACGCCGACGAGACAGCTACCATC GTGGAATTTCTGAACCGGTGGATCACCTTCTGCCAGAGCATCATCAGCACC CTGACCTGAATAGTGAGTCGTATTAACTACCAACAAGGAGTACCCTGATG AGATCACTTGATCTCATCAGGGTACTCCTTTATCTTAGAGGCATATCCCT ACGTACCAACAAGGAGTACCCTGATGAGATCACTTGGATCTCATCAGGGTA CTCCTTTATCTTAGAGGCATATCCCTACGTACCAACAAGGAGTACCCTGAT GAGATCACTTGATCTCATCAGGGTACTCCTTTATCTTAGAGGCATATCCCT TTTATCTTAGAGGCATATCCCT</p>
133	Compound 9 RNA sequence	<p>GCCACCAUGUUGUGCUGCUGCUGCCUGUAUUGCCUGGCCUCUACAGCC GCCGCUACAAAUUCUGCCCCUACCAGCAGCUCACCAAGAAAACCCAGCUG CAACUGGAACAUCUGCUGCUGGACCUGCAGAUGAUCCUGAACGGCAUCAAC AACUACAAGAACCCCAAGCUGACCCGGAUGCUGACCUUACAAGUUCUACAUG CCCAAGAAGGCCACCGAGCUGAAGCACCUCAGUGCCUGGAAGAGGAACUG AAGCCCCUGGAAGAAGUGCUGAAUCUGGCCAGAGCAAGAACUCCACCUG AGGCCUAGGGACCUGAUCAGCAACAUCAACGUGAUCUGCUGGAACUGAAA GGCAGCGAGACAACCUCAUGUGCGAGUACGCCGACGAGACAGCUACCAUC </p>

SEQ ID NO	Compound	Sequence (5' to 3')
		<p>GUGGAAUUUCUGAACCGGUGGAUACCUUCUGCCAGAGCAUCAUCAGCACCCUGACCUGAAUAGUGAGUCGUUUUACGUACCAACAAGGAGUACCCUGAUGAGAUACACUUGGAUCUCAUCAGGGUACUCCUUUAUCUUAGAGGCAUAUCCCUACGUACCAACAAGGAGUACCCUGAUGAGAUACUUGGAUCUCAUCAGGGUACUCCUUUAUCUUAGAGGCAUAUCCCUUUUAUCUUAGAGGCAUAUCCCU</p> <p>(all Us are modified; N¹-methylpseudouridine)</p>
10	Compound 10	<p>GCCACCATGTTGTTGCTGCTGCTCGCCTGTATTGCCCTGGCCTCTACAGCCGCCGCTACAAATTTCTGCCCTACCAGCAGCTCCACCAAGAAAACCCAGCTGCAACTGGAACATCTGCTGCTGGACCTGCAGATGATCCTGAACGGCATCAACAACTACAAGAACCCCAAGCTGACCCGGATGCTGACCTTCAAGTTCTACATGCCAAGAAGGCCACCGAGCTGAAGCACCTCCAGTGCCTGGAAGAGGAACTGAAGCCCTGGAAGAAGTGTGAATCTGGCCCAGAGCAAGAAGTTCCACCTGAGGCTTAGGGACCTGATCAGCAACATCAACGTGATCGTGTGGAACTGAAAGGCAGGAGACAACCTTCATGTGCGAGTACGCCGACGAGACAGCTACCATGTGGAATTTCTGAACCGGTGGATCACCTTCTGCCAGAGCATCATCAGCACCTGACCTGAATAGTGAGTGTATTACGTACCAACAAGGAGGGCAGAAATCATCAGGAAGTGGTGAAGTACTTGACTTCACCACTTCGTGATGATTCTGCCCTCTTTATCTTAGAGGCATATCCCTACGTACCAACAAGAGATGAGCTTCCTACAGCACAACAATGTGACTTGACATTTGTTGTGCTGTAGGAAGCTCATCTCTTTATCTTAGAGGCATATCCCTACGTACCAACAAGTACAGATCCGCAGACGTGTAATGTTCCACTTGGGAACATTTACAGCTCTGCGGATCTTGTACTTTATCTTAGAGGCATATCCCTTTTATCTTAGAGGCATATCCCT</p>
134	Compound 10 RNA sequence	<p>GCCACCAUGUUGUUGCUGCUGCUGCCUGUAUUGCCUGGCCUCUACAGCCGCCGCUACAAAUUCUGCCCCUACCAGCAGCUCCACCAAGAAAACCCAGCUGCAACUGGAACAUCUGCUGCUGGACCUGCAGAUGAUCCUGAACGGCAUCAACACUACAAGAACCCCAAGCUGACCCGGAUGCUGACCUUCAAGUUCUACAUGCCCAAGAAGGCCACCGAGCUGAAGCACCUCAGUGCCUGGAAGAGGAACUGAAGCCCUUGGAAGAAGUGCUGAAUCUGGCCAGAGCAAGAACUUCACCUGAGGCCUAGGGACCUGAUCAGCAACAUAACGUGAUCGUGCUGGAACUGAAAGGCAGCGAGACAACCUCAUGUGCGAGUACGCCGACGAGACAGCUACCAUCGUGGAAUUUCUGAACCGGUGGAUACCUUCUGCCAGAGCAUCAUCAGCACCCUGACCUGAAUAGUGAGUCGUUUUACGUACCAACAAGGAGGGCAGAAUCAUCACGAAGUGGUGAAGUACUUGACUUCACCACUUCGUGAUGAUUCUGCCCUUUUAUCUUAGAGGCAUAUCCCUACGUACCAACAAGAGAUAGCUUCCUACAGCACAACAAGUGAGCUUGCACAUUUGUUGUGCUGUAGGAAGCUAUCUUCUUUAUCUUAGAGGCAUAUCCCUACGUACCAACAAGUACAAAGAUCCGCAGACGUGUAAAUGUUCACUUGGGAACAUUUACACGUCUGCGGAGCUUGUACUUAUCUUAGAGGCAUAUCCCUUUUAUCUUAGAGGCAUAUCCCU</p> <p>(all Us are modified; N¹-methylpseudouridine)</p>
11	Compound 11	<p>GCCACCATGTGTACACAGCAGCTGGTTCATCAGCTGGTTTCAGCCTGGTGTCTGGCCTCTCCTCTGGTGGCCATCTGGGAGCTGAAGAAAGAGCTGTACGTGTGGAAGTGGACTGGTATCCCGATGCTCCTGGCGAGATGGTGGTGCTGACCTGCGATACCCCTGAAGAGGACGGCATCACCTGGACACTGGATCAGTCTAGCGAGGTGCTCGGCAGCGGCAAGACCCTGACCATCCAAGTGAAGAGTTTGGCGACGCCGGCCAGTACACCTGTCAAAAGGCGGAGAAGTGTGAGCCACAGCTGCTGCTGCTCCACAAGAAAGAGGATGGCATTGGAGCACCGACATCCTGAAGACCAGAAAGAGCCCAAGAACAAGACCTTCCTGAGATGCGAGGCCAAGAACTACAGCGGCCGTTACATGTTGGTGGCTGACCACCATCAGCACCGACCTGACCTTCAGCGTGAAGTCCAGCAGAGGCAGCAGTGATCCTCAGGGCGTTACATGTGGCGCCGTACACTGTCTGCCGAAAGAGTGGCGGGCGACAACAAAGAATACGAGTACAGCGTGGAATGCCAAGAGGACAGCGCCTGTCCAGCGGCCGAAGAGTCTCTGCCTATCGAAGTGATGGTGGACGCCGTGCACAAGCTGAAGTACGAGAACTACACCTCCAGCTTTTTCATCCGGGACATCATCAAGCCCGATCCTCCAAAGAACCTGCAGCTGAAGCCTCTGAAGAACAGCAGACAGGTGGAA GTGTCTGGGAGTACCCGACACCTGGTCTACACCCACAGCTACTTCAGCTGACCTTTTGCCTGCAAGTGCAGGGCAAGTCCAAGCGCGGAGAAAAGGACCGGGTGTTCACCGACAAGACCAGCGCCACCGTGATCTGCAGAAAGAAGCGCAGCATCAGCGTCAGAGCCAGGACCGGTACTACAGCAGCTCTTGGAGCGAATGGGCCAGCGTGCCATGTTCTGGTGGCGAGGATCTGGCGGAGGTGGAAGC</p>

SEQ ID NO	Compound	Sequence (5' to 3')
		<p>GGCGGAGGCGGATCTAGAAATCTGCCTGTGGCCACTCCTGATCCTGGCATG TTCCCTTGTCTGCACCACAGCCAGAACCCTGCTGAGAGCCGTGTCCAACATG CTGCAGAAGGCCAGACAGACCTTGGAAATCTTACCCTGCACCAGCGAGGAA ATCGACCACGAGGACATCACCAGGATAAGACCAGCACCGTGGAAGCCTGC CTGCCTCTGGAAGTACCAAGAACGAGAGCTGCCGTAACAGCCGGGAAACC AGCTTCATCACCAACGGCTCTTGCCTGGCCAGCAGAAAGACCTCCTTCATG ATGGCCCTGTGCCTGAGCAGCATCTACGAGGACCTGAAGATGTACCAGGTG GAATTCAAGACCATGAACGCCAAGCTGCTGATGGACCCCAAGCGGCAGATC TTCTTGGACCAGAATATGCTGGCCGTGATCGACGAGCTGATGCAGGCCCTG AACTTCAACAGCGAGACAGTGCCTCCAGAAGTCTAGCCTGGGAAGAACCCGAC TTCTACAAGACCAAGATCAAGCTGTGCATCCTGCTGCACGCTTCCGGATC AGAGCCGTGACCATCGACAGAGTGATGAGCTACCTGAACGCTCCTGAATA GTGAGTCGTATTAACTACCAACAAGTTCCTTCCAAATGGCTCTGTACTTG ACAGAGCCATTGGAAGGAACTTTATCTTAGAGGCATATCCCTACGTACCA ACAAGCAATCGTTTACCAGAGATCTGAACTGTTCAGATCTCGGTGAACGATGC TTTATCTTAGAGGCATATCCCTACGTACCAACAAGACCAGCAGCGGACAAA TAAAACTTGTTTATTGTCCGTGCTGGTCTTTTATCTTAGAGGCATATCCC TTTATCTTAGAGGCATATCCCT</p>
135	Compound 11 RNA sequence	<p>GCCACCAUGUGUACACAGCAGCUGGUCAUCAGCUGGUUACAGCCUGGUGUUC CUGGCCUCUCCUCUGGUGGCCAUCUGGGAGCUGAAGAAAGACGUGUACGUG GUGGAACUGGACUGGUAUCCCGAUGUCUCCUGGCAGAUUGGUGGUGCUGACC UGCGAUACCCUGAAGAGGACGGCAUCACCUGGACACUGGAUCAGUACUAGC GAGGUGUCUGGCAGCGGCAAGACCCUGACCAUCCAAGUGAAAGAGUUUGGC GACGCCGGCCAGUACACCCUGUCACAAAGGCGGAGAAGUGCUGAGCCACAGC CUGCUGCUGUCUCCACAAGAAAGAGGAUUGGCAUUUGGAGCACCGACAUCCUG AAGGACCAGAAAGAGCCCAAGAACAAGACCUUCCUGAGAUGCGAGGCCAAG AACUACAGCGGCCCGUUCACAUUGUUGGUGGUGACCAUACAGCACCAGAC CUGACCUUACAGCGUGAAGUCCAGCAGAGGCAGCAGUGAUCCUACAGGGCGU ACAUGUGGGCGCCGCUACACUGUCUGCCGAAAGAGUGCGGGGCGACAAACAA GAAUACGAGUACAGCGUGGAUUGCCAAGAGGACAGCGCCUGUCCAGCCGCC GAAGAGUCUCUGCCUAUCGAAGUGAUGGUGGACGCCGUGCACAAGCUGAAG UACGAGAAUCUACACUCCAGCUUUUUAUCCGGGACAUCAUAAAGCCCGAU CCUCCAAAGAACCUGCAGCUGAAGCCUCUGAAGAACAGCAGACAGGUGGAA GUGUCCUGGAGUAACCCGACACCUUGGUCUACACCCACAGCUACUUCAGC CUGACCUUUUGCGUGCAAGUGCAGGGCAAGUCCAAGCGCGAGAAAAGGAC CGGGUGUACACCGACAAGACACAGCGCCACCGUGAUCUGCAGAAAGAAGCC AGCAUCAGCGUCAGAGCCCAGGACCGGUACUACAGCAGCUCUUGGAGCGAA UGGGCCAGCGUGCCAUGUUCUGGUGGGCGGAGGAUCUGGGCGAGGUGGAAGC GGCGGAGGCGGAUCUAGAAAUUCUGCCUGUGGCCACUCCUGAUCUUGGCAUG UUCCCUUGUCUGCACCACAGCCAGAACCUGCUGAGAGCCGUGUCCAACAUG CUGCAGAAAGGCCAGACAGACCCUGGAAUUCUACCCUUGCACCAGCGAGGAA AUCGACCACGAGGACAUACCAAGGAUAAGACCAGCACCUGGGAAGCCUGC CUGCCUCUGGAACUGACCAAGAAGACGAGAGCUGCCUGAACAGCCGGGAAACC AGCUUCAUCACCAACGGCUCUUGCCUGGCCAGCAGAAAGACCUCCUUAUG AUGGCCUGUGCCUGAGCAGCAUCUACGAGGACCUGAAGAUGUACAGGUG GAAUUAAGACCAUGAAGCCAGCUGCUGAUGGACCCCAAGCGGCAGAU UUCCUGGACCAGAAUUGCUGGCCUGAUCGACGAGCUGAUGCAGGCCCCUG AACUUAACAGCGAGACAGUGCCCCAGAGUCUAGCCUGGAAGAACCAGC UUUUAACAAGACCAAGAUCAAGCUGUGCAUCCUGUGCAGCCUUCGGGAUC AGAGCCGUGACCAUCGACAGAGUGAUGAGCUACCUAAGCCUCCUGAAUA GUGAGUCGUUAUACGUACCAACAAGUCCUUAAGAGGCAUACCCUACGUACCA ACAGAGCCAUUUGGAAGGAACUUUAUCUUAAGAGGCAUACCCUACGUACCA ACAAGCAUCGUUACCCAGAGAUUCUGAACUUGUCAGAUUCUGGUGAACGAGUC UUUAUCUUAAGAGGCAUACCCUACGUACCAACAAGACCAGCAGCGGACAAA UAAAACUUGUUUAUUUGUCCGUGCUGGUCUUUAUCUUAAGAGGCAUACCC UUUUAUCUUAAGAGGCAUACCCU</p> <p>(all Us are modified; N¹-methylpseudouridine)</p>
12	Compound 12	<p>GCCACCATGTGTACACAGCAGCTGGTCATCAGCTGGTTACGCTGGTGTTC CTGGCCTCTCCTCTGGTGGCCATCTGGGAGCTGAAGAAAGACGTGTACGTG GTGGAAGTGGACTGTATCCCAGTCTCCTGGCGAGATGTGTGTGCTGACC TGCGATACCCCTGAAGAGGACGGCATCACCTGGACACTGGATCAGTCTAGC</p>

SEQ ID NO	Compound	Sequence (5' to 3')
		<p>GAGGTGCTCGGCAGCGGCAAGACCCTGACCATCCAAGTGAAGAGTTTGGC GACGCCGGCCAGTACACCTGTACAAAGGCGGAGAAGTGCTGAGCCACAGC CTGCTGCTGCTCCACAAGAAAGAGGATGGCATTTGGAGCACCACATCCTG AAGGACCAGAAAGAGCCCAAGAACAAGACCTTCTGAGATGCGAGGCCAAG AACTACAGCGCGCGTTTACATGTTGGTGGCTGACCACCATCAGCACCGAC CTGACCTTCAGCGTGAAGTCCAGCAGAGGCGAGCAGTGATCCTCAGGGCGTT ACATGTGGCGCCGCTACACTGTCTGCCGAAAGAGTGCGGGGCGACAACAAA GAATACGAGTACAGCGTGGAATGCCAAGAGGACAGCGCCTGTCCAGCCGCC GAAGAGTCTCTGCCATCGAAGTGATGGTGGACGCCGTGCACAAGCTGAAG TACGAGAACTACACCTCCAGCTTTTTCATCCGGGACATCATCAAGCCGAT CCTCCAAAGAACCTGCAGCTGAAGCCTCTGAAGAACAGCAGACAGGTGGAA GTGTCTGGGAGTACCCGACACCTGGTCTACACCCACAGCTACTTCAGC CTGACCTTTTGCCTGCAAGTGCAAGGCAAGTCCAAGCGCGAGAAAAGGAC CGGGTGTTCACCGACAAGACCAGCGCCACCGTGATCTGCAGAAAGAACGCC AGCATCAGCGTCAGAGCCAGGACCGGTACTACAGCAGCTCTTGGAGCGAA TGGGCCAGCGTGCCATGTTCTGGTGGCGGAGGATCTGGCGGAGGTGGAAGC GGCGGAGGCGGATCTAGAAATCTGCCTGTGGCCACTCCTGATCCTGGCATG TTCCCTTGCTGCAACACAGCCAGAACCTGTCTGAGAGCCGTGTCCAACATG CTGCAGAAGGCCAGACAGACCTTGAATTTTACCCCTGCACAGCGAGGAA ATCGACCAGGAGCATCACCAAGGATAAGACCAGCACCCGTGGAAGCCTGC CTGCCTCTGGAAGTACCAAGAACGAGAGCTGCCTGAACAGCCGGGAAACC AGCTTCATCACCAACGGCTCTTGCCTGGCCAGCAGAAAGACCTCCTTCATG ATGGCCCTGTGCCTGACGAGCATCTACGAGGACCTGAAGATGTACCAGGTG GAATTCAAGACCATGAACGCCAAGCTGCTGATGGACCCCAAGCGGCGAGATG TTCTTGGACAGAAATATGCTGGCCGTGATCGACGAGCTGATGCAGGCCCTG AACTTCAACAGCGAGACAGTGCCCCAGAAGTCTAGCCTGGAAGAACCCGAC TTCTACAAGACCAAGATCAAGCTGTGCATCCTGCTGCACGCCTTCCGGATC AGAGCCGTGACCATCGACAGAGTGATGAGCTACCTGAACGCCTCCTGAATA GTGAGTCGTATTAACTACCAACAAATAGTGAGTCGTATTAACTACCAAC AAGAAGGAGCTGCCCATGAGAAAATTGTTTCTCATGGGAGCTCCTTCTT TATCTTAGAGGCATATCCCTACGTACCAACAAGTGAATGAGGGACCAGTA CAACTTGTACTGGTCCCTCATTGCACCTTATCTTAGAGGCATATCCCTA CGTACCAACAAGAGCTGCTGAAGGACTCATCAACTTGTGATGAGTCCTTCA GCAGCTCTTATCTTAGAGGCATATCCCTTTTATCTTAGAGGCATATCCCT</p>
136	Compound 12 RNA sequence	<p>GCCACCAUGUGUACACAGCAGCUGGUCAUCAGCUGGUUACGCCUGGUGUUC CUGGCCUCUCCUCUGGUGGCCAUCUGGGAGCUGAAGAAAGACGUGUACGUG GUGGAACUGGACUGGUAUCCGAUGUCUCCUGGCAGAUUGGUGGUGUGUAC UGCGAUACCCUGAAGAGGACGGCAUACCCUGGACACUGGAUCAGUCUAGC GAGGUGUCUGGCAGCGGCAAGACCCUGACCAUCCAAGUGAAAGAGUUUGGC GACGCCGGCCAGUACACCUUGUCACAAAGGCGGAGAAGUGCUGAGCCACAGC CUGCUGCUGUCUCCACAAGAAAGAGGAUGGCAUUUGGAGCACCACAUCCUG AAGGACCAGAAAGAGCCCAAGAACAAGACCUUCCUGAGAUGCGAGGCCAAG AACUACAGCGGCCGGUUCACAUUUGGUGGUGACCAACCAUACGACCCGAC CUGACCUUCAGCGUGAAGUCCAGCAGAGGCAGCAGUGAUCCUCAGGGCGU ACAUGUGGCGCCGCUACACUGUCUGCCGAAAGAGUGCGGGGCGACAACAAA GAAUACGAGUACAGCGUGGAAUGCCAAGAGGACAGCGCCUGUCAGCCGCC GAAGAGUCUCUGCCUAUCGAAGUGAUGGUGGACGCCGUGCAACAAGCUGAAG UACGAGAACUACACCUCCAGCUUUUUAUCCGGGACAUCAUCAAGCCCGAU CCUCCAAAGAACCUGCAGCUGAAGCCUCUGAAGAACAGCAGACAGGUGGAA GUGUCCUGGGAGUACCCCGACACCUUGGUCUACACCCACAGCUACUUCAGC CUGACCUUUUGCGUGCAAGUGCAGGGCAAGUCCAAGCGCGAGAAAAGGAC CGGGUGUUCACCGACAAGACCAGCGCCACCGUGAUUCAGAGAAAGAACGCC AGCAUCAGCGUCAGAGCCAGGACCGGUACUACAGCAGCUUUGGAGCGAA UGGGCCAGCGUGCAUGUUCUGGUGGCGGAGGAUCUGGCGGAGGUGGAAGC GGCGGAGGCGGAUCUAGAAUUCUGCCUGUGGCCACUCCUGAUCCUGGCAUG UUCCCUUGUCUGCACACAGCCAGAACCUUGCUGAGAGCCGUGUCCAACAU UGCAGAAAGGCCAGACAGACCCUGGAAUUCUACCCUGCACCAGCAGGAGAA AUCGACCACGAGGACAUCACCAAGGAUAAGACCAGCACCUGGAAGCCUGC CUGCCUCUGGAACUGACCAAGAACAGAGCUGCCUGAACAGCCGGGAAACC AGCUUCAUCACCAACGGCUCUUGCCUGGCCAGCAGAAAGACUCCUUCUAG AUGGCCUUGUGCCUGAGCAGCAUCUACGAGGACCUGAAGAUGUACAGGUG</p>

SEQ ID NO	Compound	Sequence (5' to 3')
		<p>GAAUUCAAGACCAUGAACGCCAAGCUGCUGAUGGACCCCAAGCGGCAGAUU UUCUUGGACCAGAAUUGCUGGCCUGAUCGACGAGCUGAUCGAGGCCUG AACUUCAACAGCGAGACAGUGCCCCAGAAGUCUAGCCUGGAAGAACCCGAC UUCUACAAGACCAAGAUCAAGCUGUGCAUCCUGCUGACGCCUUCGGGAUC AGAGCCGUGACCAUCGACAGAGUGAUGAGCUACCUGAACGCCUCCUGAAUA GUGAGUCGUUUAAACGUACCAACAAUAGUGAGUCGUUUAAACGUACCAAC AAGAAGGAGCUGCCCCAUGAGAAAACUUGUUUCUCAUGGGCAGCUCUUCU UAUCUUAGAGGCAUAUCCCUACGUACCAACAAUGGCAUAGAGGGACAGUA CAACUUGUGUACUGGUCCUCAUUGCACUUUAUCUUAGAGGCAUAUCCCUA CGUACCAACAAAGAGCUGCUGAAGGACUCAUACUUGUGAUGAGUCCUUA GCAGCUCUUUAUCUUAGAGGCAUAUCCCUUUUAUCUUAGAGGCAUAUCCCU (all Us are modified; N¹-methylpseudouridine)</p>
13	Compound 13	<p>GCCACCATGTGTCCAGCAGCTGGTCATCAGCTGGTTACGCTGGTGTTC CTGGCCTCTCCTCTGGTGGCCATCTGGGAGCTGAAGAAAGACGTGTACGTG GTGGAACCTGGACTGGTATCCCGATGCTCCTGGCGAGATGGTGGTGTGCTG TGCATACCCCTGAAGAGGACGGCATCACCTGGACACTGGATCAGTCTAGC GAGGTGCTCGGCAGCGCAAGACCCCTGACCATCCAAGTGAAGAGTTTGGC GACGCCGGCCAGTACACCTGTCACAAAGGCGGAGAAGTGTGAGCCACAGC CTGCTGTCTGCTCCACAAGAAAGAGGATGGCATTGAGGACCCGACATCCTG AAGGACCAAGAAAGAGCCCAAGAACAAAGACCTTCCTGAGATGCGAGGCCAAG AACTACAGCGGCCGGTTCACATGTTGGTGGCTGACCACCATCAGCACCAGC CTGACCTTCAGCGTGAAGTCCAGCAGAGGCAGCAGTGTCTCAGGGCGTT ACATGTGGCGCCGTACACTGTCTGCCGAAAGAGTGCAGGGCGACAAACAA GAATACGAGTACAGCGTGAATGCCAAGAGGACAGCGCCTGTCCAGCCGCC GAAGAGTCTCTGCCATCGAAGTGTGGTGGACGCCGTGCACAAGCTGAAG TACGAGAACTACACCTCCAGCTTTTTCATCCGGGACATCATCAAGCCCGAT CCTCCAAAGAACCTGCAGCTGAAGCCTCTGAAGAACAGCAGACAGGTGGAA GTGTCTGGGAGTACCCGACACCTGGTCTACACCCACAGCTACTTCAGC CTGACCTTTTGCCTGCAAGTGCAGGGCAAGTCCAAGCGCGAGAAAAGGAC CGGGTGTTCACCGACAAGACCAGCGCCACCGTGTCTGCAGAAAGAACGCC AGCATCAGCGTCAGAGCCAGGACCGGTACTACAGCAGCTCTTGGAGCGAA TGGGCCAGCGTGCCATGTTCTGGTGGCGGAGGATCTGGCGGAGGTGGAAGC GGCGGAGGCGGATCTAGAAATCTGCCTGTGGCCACTCCTGATCCTGGCATG TTCCCTTGTCTGCACCAAGCCAGCCAGAACCTGCTGAGAGCCGTGTCCAACTG CTGCAGAAAGGCCAGACAGACCTGGAATTCTACCCCTGCACCAGCGAGGAA ATCGACCACGAGGACATCACCAAGGATAAGACCAGCACCCTGGAAGCTGTC CTGCCCTCTGGAACCTGACCAAGAACAGAGCTGCTGAACAGCCGGGAAACC AGCTTCATCACCAACGGCTCTTGCCCTGGCCAGCAGAAAGACCTCCTTCATG ATGGCCCTGTGCCTGAGCAGCATCTACGAGGACCTGAAGATGTACAGGTG GAATTCAAGACCATGAACGCCAAGCTGCTGATGGACCCCAAGCGGCAGATC TTCTTGGACAGAAATATGCTGGCCGTGATCGACGAGCTGATGCAGGCCCTG AACTTCAACAGCGAGACAGTGGCCAGAGTCTAGCCTGGAAGAACCCGAC TTCTACAAGACCAAGATCAAGCTGTGCATCCTGCTGCACGCCTTCCGGATC AGAGCCGTGACCATCGACAGAGTGATGAGCTACCTGAACGCCTCCTGAATA GTGAGTCGTATTACGTACCAACAAGAGGAGCTGCCCATGAGAAAACCTTG TTTCTCATGGGCAGCTCCTTCTTTATCTTAGAGGCATATCCTACGTACCA ACAAGTCCAACGAATGGGCCTAAGAACTTGCTTAGGCCCATTCGTTGGAC TTTATCTTAGAGGCATATCCTACGTACCAACAAGGACAGCATAGACGACA CCTTACTTGAAGGTGTCGTCTATGCTGTCCCTTTATCTTAGAGGCATATCCC TTTATCTTAGAGGCATATCCT</p>
137	Compound 13 RNA sequence	<p>GCCACCAUGUGUACACAGCAGCUGGUCAUCAGCUGGUUCAGCCUGGUGUUC CUGGCCUCUCCUCUGGUGGCCAUUCUGGAGCUGAAGAAAGACGUGUACGUG GUGGAACUGGACUGGUAUCCCGAUGCUCCUGGCAGAUUGGUGUGUGAC UGCGAUACCCUGAAGAGGACGGCAUCACCGGACACUGGAUCAGUCUAGC GAGGUGUCUGGCAGCGGCAAGACCCUGACCAUCCAAGUGAAAGAGUUUGGC GACGCCGGCCAGUACACCGUACACAAAGGCGGAGAAGUGCUGAGCCACAGC CUGCUGCUGCUCCACAAGAAAGAGGAUGGCAUUUGGAGCACCAGACUCCUG AAGGACCAAGAAAGAGCCCAAGAACAAAGACCUUCCUGAGAUCCGAGGCCAAG AACUACAGCGGCCGGUUCACAUUUGGUGGCUGACCACCAUCAGCACCAGC CUGACCUUCAGCGUAGGUCCAGCAGAGGCAGCAGUGAUCCUCAGGGCGUU ACAUGUGGCGCCGUACACUGUCUGCCGAAAGAGUGCGGGGCGACAACAAA</p>

SEQ ID NO	Compound	Sequence (5' to 3')
		<p>GAAUACGAGUACAGCGUGGAAUGCCAAGAGGACAGCGCCUGUCAGCCGCC GAAGAGUCUCUGCCUAUCGAAGUGAUGGUGGACGCCUGGCACAAGCUGAAG UACGAGAACUACACCUCAGCUUUUUCAUCCGGGACAUCAUCAAGCCCGAU CCUCCAAAGAACCUGCAGCUGAAGCCUCUGAAGAACAGCAGACAGGUGGAA GUGUCCUGGGAGUACCCCGACACCUGGUCUACACCCACAGCUACUUCAGC CUGACCUUUUGCGUGCAAGUGCAGGGCAAGUCCAAGCGCGAGAAAAGGAC CGGGUGUUCACCGACAAGACCAGCGCCACCGUGAUCUGCAGAAAGAACGCC AGCAUCAGCGUCAGAGCCCAGGACCGGUACUACAGCAGCUUUGGAGCGAA UGGGCCAGCGUGCCAUGUUCUGGUGGGCGGAGGAUCUGGGCGAGGUGGAAGC GGCGGAGGCGGAUCUAGAAAUCUGCCUGUGGCCACUCCUGAUCUGGCAUG <u>UUCCUUGUCUGCACCACAGCCAGAACCUGCUGAGAGCCGUGUCCAACAU</u> <u>GUGCAGAAGGCCAGACAGACCUGGAAUUCUACCCUGCACCAGCGAGGAA</u> <u>AUCGACCACGAGGACAUCACCAAGGAUAAGACCAGCACCUGGAAGCCUGC</u> <u>CUGCCUCUGGAACUGACCAAGAACGAGAGCUGCCUGAACAGCCGGGAAACC</u> <u>AGCUUCAUCACCAACGCGCUCUUGCCUGGCCAGCAGAAAGACCCUUCU</u> <u>AUGGCCUUGUGCCUGAGCAGCAUCUACGAGGACCUGAAGAUGUACCGGUG</u> <u>GAAUUAAGACCAUGAACGCCAAGCUGCUGAUGGACCCCAAGCGGCAGAU</u> <u>UUCUGGACCCAGAAUUGCUGGCCUGAUCGACGAGCUGAUGCAGGCCUG</u> <u>AACUUAACAGCGAGACAGUGCCCCAGAAGUCUAGCCUGGAAGAACCAGC</u> <u>UUCUACAAGACCAAGAUCAAGCUGUGCAUCCUGCUGCAGCCUUCGGAU</u> <u>AGAGCCGUGACCAUCGACAGAGUGAUGAGCUACCUGAACGCCUCCUGAAU</u> <u>GUGAGUCGUUAUAAACGUACCAACAAGAGGAGCUGCCCAUGAGAAAACU</u> <u>UUUCUUAUGGGCAGCUCUUCUUAUCUUAAGAGGCAUUAUCCUACGUACCA</u> <u>ACAAGUCCAAACGAUUGGGCCUAAGAACUUGUCUUAAGGCCAUUCGUUGGAC</u> <u>UUUAUCUUAAGAGGCAUUAUCCUACGUACCAACAAGGACAGCAUAGACGACA</u> <u>CCUUAUCUUAAGGUGUCGUUAUGCUGUCCUUAUUAUUAAGAGGCAUUAUCC</u> UUUAUCUUAAGAGGCAUUAUCCU (all Us are modified; N¹-methylpseudouridine)</p>
14	Compound 14	<p>GCCACCATGTGTACACAGCAGCTGGTCATCAGCTGGTTCCAGCTGGTGTTC CTGGCCTCTCCTCTGGTGGCCATCTGGGAGCTGAAGAAAGAGCTGTACGTG GTGGAACCTGGACTGGTATCCCGATGCTCCTGGCGAGATGGTGGTGCTGACC TGCGATACCCCTGAAGAGGACGGCATCACCTGGACACTGGATCAGTCTAGC GAGGTGCTCGGCAGCGGCAAGACCCTGACCATCCAAGTGAAGAGTTTGGC GACGCCGGCCAGTACACCTGTCAAAAGCGGAGAAAGTGTGTAGCCACAGC CTGCTGCTGCTCCACAAGAAAGAGGATGGCATTTGGAGCACCAGATCCTG AAGGACCAGAAAGAGCCCAAGAACAAGACCTTCCTGAGATGCGAGGGCAAG AACTACAGCGGCCGGTTCACATGTTGGTGGCTGACCACCATCAGCACCAGC CTGACCTTCAGCGTGAAGTCCAGCAGAGGCAGCAGTGTCTCAGGGCGTT ACATGTGGCGCCGCTACACTGTCTGCCGAAAGAGTGCGGGGCCGACACAAA GAATACGAGTACAGCGTGAATGCCAAGAGGACAGCGCCTGTCCAGCCGCC GAAGAGTCTCTGCCATATCGAAGTGATGGTGGACGCCGTGCAAGAAGTGAAG TACGAGAACTACACCTCCAGCTTTTTCATCCGGGACATCATCAAGCCCGAT CCTCCAAGAACCTGCAGCTGAAGCCTCTGAAGAACAGCAGACAGGTGGAA GTGTCTGGGAGTACCCCGACACCTGGTCTACACCCACAGCTACTTCAGC CTGACCTTTTGGTGCAGTGCAGGGCAAGTCCAAGCGCGAGAAAAAGGAC CGGGTGTTCACCGACAAGACCAGCGCCACCGTGATCTGCAGAAAGAACGCC AGCATCAGCGTCAGAGCCCAGGACCGGTACTACAGCAGCTCTTGGAGCGAA TGGGCCAGCGTGCCATGTTCTGGTGGCGGAGGATCTGGCGGAGGTGGAAGC GGCGGAGGCGGATCTAGAAATCTGCCTGTGGCCACTCCTGATCCTGGCATG <u>TTCCCTTGTCTGCACCACAGCCAGAACCCTGCTGAGAGCCGTGTCCAACATG</u> <u>CTGCAGAAAGGCCAGACAGACCTGGAATTCTACCCCTGCACCAGCGAGGAA</u> <u>ATCGACCACGAGGACATCACCAAGGATAAGACCAGCACCCTGGAAGCCTGC</u> <u>CTGCCTCTGGAACCTGACCAAGAACGAGAGCTGCCTGAACAGCCGGGAAACC</u> <u>AGCTTCATCACCAACGGCTCTTGCCCTGGCCAGCAGAAAGACCTCCTTCATG</u> <u>ATGGCCCTGTGCCTGAGCAGCATCTACGAGGACCTGAAGATGTACCAGGTG</u> <u>GAATTCAAGACCATGAACGCCAAGCTGCTGATGGACCCCAAGCGGCAGATC</u> <u>TTCTTGGACCAAGATATGCTGGCCGTGATCGACGAGCTGATGACGAGCCCTG</u> <u>AACTTCAACAGCGAGACAGTGTGCCCCAGAAGTCTAGCCTGGAGAAGCCGAC</u> <u>TTCTACAAGACCAAGATCAAGCTGTGCATCCTGCTGCACGCCTTCCGGATC</u> <u>AGAGCCGTGACCATCGACAGAGTGATGAGCTACCTGAACGCCTCCTGAATA</u> GTGAGTCGTATTACGTACCAACAAGACCCTGACATTGGCTACTGTACTTG</p>

SEQ ID NO	Compound	Sequence (5' to 3')
		<p>ACAGTAGCGAATGTGAGGGTCTTTATCTTAGAGGCATATCCCTACGTACCA ACAAGAGCTGCTGAAGGACTCATCACTTGTGATGAGTCCTTCAGCAGCTC TTTATCTTAGAGGCATATCCCTACGTACCAACAAGGCCAATGACCCAACAT CTCTACTTGAGAGATGTTGGGTCATTGGCCCTTTATCTTAGAGGCATATCCCT TTTATCTTAGAGGCATATCCCT</p>
138	Compound 14 RNA sequence	<p>GCCACCAUGUGUCACCAGCAGCUGGUAUCAGCUGGUUCAGCCUGGUGUUC CUGGCCUCUCCUCUGGUGGCCAUCUGGGAGCUGAAGAAAGACGUGUACGUG GUGGAACUGGACUGGUAUCCCGAUGCUCCUGGCCGAGAUGGUGGUGCUGACC UGCGAUACCCUGAAGAGGACGGCAUCACCCUGGACACUGGAUCAGUCUAGC GAGGUGCUCGGCAGCGGCAAGACCCUGACCAUCCAAGUGAAAGAGUUUGGC GACGCCGGCCAGUACACCUUGUCACAAAGCGGAGAGUGCUGAGCCACAGC CUGCUGCUGCUCCACAAGAAAGAGGAUGGCAUUUGGAGCACCAGAUCCUG AAGGACCAGAAAGAGCCCAAGAACAGACCUUCCUGAGAUUGCGAGGCCAAG AACUACAGCGGCCGUGUACAUUGUUGGUGGACACCAUACAGCACCAGC CUGACCUUACGCGUGAAGUCCAGCAGAGGCAGCAGUACCUUCCAGGCGUU ACAUGUGGCGCGCUACACUGUCUGCCGAAAGAGUGCGGGCGACACAA GAAUACGAGUACAGCGUGGAAUGCCAAGAGGACAGCGCCUGUCCAGCCGCC GAAGAGUCUCUGCCUAUCGAAGUGAUGGUGGACGCCUGGCAACAGCUGAAG UACGAGAACUACACCUCCAGCUUUUUAUCCGGGACAUCAUAAAGCCCGAU CCUCCAAAGAACCUGCAGCUGAAGCCUCUGAAGAACAGCAGACAGGUGGAA GUGUCCUGGGAGUACCCCGACACCUUGUCUACACCCACAGCUACUUCAGC CUGACCUUUUGCGUGCAAGUGCAGGGCAAGUCCAAGCGCGAGAAAAAGGAC CGGGUGUUCACCGACAAGACCGAGCGCCACCGUGAUCUGCAGAAAGAACGCC AGCAUCAGCGUCAGAGCCCAGGACCGGUACUACAGCAGCUCUUGGAGCGAA UGGGCCAGCGUGGCCAUGUUCUGGUGGGCGAGGAUCUGCGCGAGGUGGAAG GGCGGAGGCGGAUCUAGAAAUCUGCCUGUGGCCACUCCUGAUCCUGGCAUG <u>UUCUUUGUCUGCACCACAGCCAGAACCCUGCUGAGAGCCGUGUCCAACAU</u> <u>UUGCAGAAGGCCAGACAGACCCUGGAAUUCUACCCUGCACCAGCGAGGAA</u> <u>AUCGACCACGAGGACAUACCAAGGAUAAGACCAGCACCCUGGAAGCCUGC</u> <u>UGCCUCUGGAAUCUGACCAAGAACAGAGCUGCCUGAACAGCCGGGAAACC</u> <u>AGCUUCAUACCAACGGCUCUUGCCUGGCCAGCAGAAAGACCUCCUUAUG</u> <u>AUGGCCUCUGCCUGAGCAGCAUCUACGAGGACCGAAGAUUACAGGUG</u> <u>GAAUUAAGACCAUGAACGCCAAGCUGCUGAUGGACCCCAAGCGGCAGAU</u> <u>UUCUUGGACCAAGAAUUGCUGGCCGUGAUCGACGAGCUGAUGCAGGCCUG</u> <u>AACUUAACAGCGAGACAGUGCCCCAGAGUCUAGCCUGGAAGAACCCGAC</u> <u>UUCUACAAGACCAAGAUCAAGCUGUGCAUCCUGCUGCAGCCUUCGGAU</u> <u>AGAGCCGUGACCAUCGACAGAGUGAUGAGCUACCGAACCCUCCUGAAUA</u> <u>GUGAGUCGUUAUACGUACCAACAAGACCCUGACAUCGCUACUGUACUUG</u> ACAGUAGCGAAUGUCAGGGUCUUUAUCUAGAGGCAUAUCCUACGUACCA ACAAGAGCUGCUGAAGGACUCAUAACUUGUGAUGAGUCCUUCAGCAGCUC UUUAUCUAGAGGCAUAUCCUACGUACCAACAAGGCCAUAUGACCAACA CUUCUACUUGAGAGAUUGUGGUCAUUGGCCUUUAUCUAGAGGCAUAUCC UUUAUCUAGAGGCAUAUCCU (all Us are modified; N¹-methylpseudouridine)</p>
15	Compound 15	<p>GCCACCATGAGAATCAGCAAGCCCCACCTGAGATCCATCAGCATCCAGTGC TACCTGTGCTGCTGCTGAACAGCCACTTTCTGACAGAGGCCGGCATCCAC GTGTTTCATCTGGGCTGTTTTCTGCCGGCCTGCCTAAGACCGAGGCCAAC TGGTTTAACGTGATCAGCGACCTGAAGAAGATCGAGGACCTGATCCAGAGC ATGCACATCGACGCCACACTGTACACCGAGAGCGACGTGCACCCTAGCTGT AAAGTGACCGCCATGAAGTGCTTTCTGCTGGAAGTGCAAGTGATCAGCCTG GAAAGCGGCGACGCCAGCATCCACGACACCGTGGAAAACCTGATCATCCTG GCCAACACAGCCTGAGCAGCAACGGCAATGTGACCGAGTCCGGCTGCAAA GAGTGCGAGGAACCTGAAGAGAAGAATATCAAAGAGTTCTGCGAGAGCTTC GTGCACATCGTGAGATGTTTCATCAACACCAGCTGAATAGTGAGTCGTATT AACGTACCAACAAGGAGTACCCTGATGAGATCACTTGGATCTCATAGGGT ACTCCTTTATCTTAGAGGCATATCCCTACGTACCAACAAGGTATCCATCTC TGGCTATGAACTTGTCATAGCCAGAGATGGATACCTTTATCTTAGAGGCAT ATCCCTACGTACCAACAAGTCCCGTTAACGCCATCATCTTACTTGAGATGA TGGCGTTACGGGACTTTATCTTAGAGGCATATCCCTTTTATCTTAGAGGCA TATCCCT</p>

SEQ ID NO	Compound	Sequence (5' to 3')
139	Compound 15 RNA sequence	<p> <i>GCCACCAUGAGAAUCAGCAAGCCCCACCUGAGAUCCAUCAGCAUCCAGUGC</i> <i>UACCUGUGCCUGCUGCUGAACAGCCACUUUCUGACAGAGGCCGGCAUCCAC</i> <i>GUGUUAUCCUGGGCUGUUUUUCUGCCGGCCUGCCUAAGACCGAGGCCAAC</i> <i>UGGGUUAACGUGAUACGCGACCUGAAGAAGAUCCAGGACCUGAUCCAGAGC</i> <i>AUGCACAUACGAGCCACACUGUACACCGAGAGCGACGUGCACCUCUAGCUGU</i> <i>AAAGUGACCGCCAUGAAGUGCUUUUCUGUGGAACUGCAAGUGAUACAGCCUG</i> <i>GAAAGCGGCGACGCCAGCAUCCACGACACCGUGGAAAACCUGAUCAUCCUG</i> <i>GCCAACAACAGCCUGAGCAGCAACGGCAAUGUGACCGAGUCCGGCUGCAAA</i> <i>GAGUGCGAGGAACUGGAAGAGAAGAAUUAUCAAAGAGUUCUGCAGAGCUUC</i> <i>GUGCACAUUCGUGCAGAUUUCAUCAACACCAGCUGAAUAGUGAGUCGUUU</i> <i>AACGUACCAACAAGGAGUACCCUGAUGAGAUCAUUGGAUCUACUACAGGGU</i> <i>ACUCCUUUAUCUAGAGGCAUAUCCUACGUACCAACAAGGUAUCCAUUCUC</i> <i>UGGCUAUGAAUUGUCAUAGCCAGAGAUUGGAUACCUUUUAUCUAGAGGCAU</i> <i>AUCCUACGUACCAACAAGUCCGUAAACGCCAUCAUCUUACUUGAAGAUGA</i> <i>UGGCGUACGGGACUUUAUCUAGAGGCAUAUCCUUUUUAUCUAGAGGCA</i> <i>UAUCCCU</i> (all Us are modified; N¹-methylpseudouridine) </p>
16	Compound 16	<p> <i>GCCACCATGAGAATCAGCAAGCCCCACCTGAGATCCATCAGCATCCAGTGC</i> <i>TACCTGTGCTGCTGCTGAACAGCCACTTTCTGACAGAGGCCGGCATCCAC</i> <i>GTGTTTCATCCTGGGCTGTTTTTCTGCCGGCCTGCCTAAGACCGAGGCCAAC</i> <i>TGGGTAAACGTGATCAGCGACCTGAAGAAGATCGAGGACCTGATCCAGAGC</i> <i>ATGCACATCGACGCCACACTGTACACCGAGAGCGACGTGCACCCTAGCTGT</i> <i>AAAGTGACCGCCATGAAGTGCTTTCTGCTGGAAGTGAAGTGATCAGCCTG</i> <i>GAAAGCGGCGACGCCAGCATCCACGACACCGTGGAAAACCTGATCATCCTG</i> <i>GCCAACAACAGCCTGAGCAGCAACGGCAATGTGACCGAGTCCGGCTGCAAA</i> <i>GAGTGCGAGGAAGTGAAGAGAAGAATATCAAAGAGTTCCTGCAGAGCTTC</i> <i>GTGCACATCGTGAGATGTTTCATCAACACCAGCTGAATAGTGAGTCGTATT</i> <i>AACGTACCAACAAGGAGTACCTTGATGAGATCACTTGATCTCATCAGGGT</i> <i>ACTCCTTTATCTTAGAGGCATATCCCTACGTACCAACAAGAGGTTAGCA</i> <i>TAGTAGCTAACTTGAGCTACTATGCTGAACCTTCTTTATCTTAGAGGCAT</i> <i>ATCCCTACGTACCAACAAGGACGACGAGACCTTCATCAAACTTGTTGATGA</i> <i>AGGTCTCGTCTCCTTTATCTTAGAGGCATATCCCTTTTATCTTAGAGGCA</i> <i>TATCCCT</i> </p>
140	Compound 16 RNA sequence	<p> <i>GCCACCAUGAGAAUCAGCAAGCCCCACCUGAGAUCCAUCAGCAUCCAGUGC</i> <i>UACCUGUGCCUGCUGCUGAACAGCCACUUUCUGACAGAGGCCGGCAUCCAC</i> <i>GUGUUAUCCUGGGCUGUUUUUCUGCCGGCCUGCCUAAGACCGAGGCCAAC</i> <i>UGGGUUAACGUGAUACGCGACCUGAAGAAGAUCCAGGACCUGAUCCAGAGC</i> <i>AUGCACAUACGAGCCACACUGUACACCGAGAGCGACGUGCACCUCUAGCUGU</i> <i>AAAGUGACCGCCAUGAAGUGCUUUUCUGUGGAACUGCAAGUGAUACAGCCUG</i> <i>GAAAGCGGCGACGCCAGCAUCCACGACACCGUGGAAAACCUGAUCAUCCUG</i> <i>GCCAACAACAGCCUGAGCAGCAACGGCAAUGUGACCGAGUCCGGCUGCAAA</i> <i>GAGUGCGAGGAACUGGAAGAGAAGAAUUAUCAAAGAGUUCUGCAGAGCUUC</i> <i>GUGCACAUUCGUGCAGAUUUCAUCAACACCAGCUGAAUAGUGAGUCGUUU</i> <i>AACGUACCAACAAGGAGUACCCUGAUGAGAUCAUUGGAUCUACUACAGGGU</i> <i>ACUCCUUUAUCUAGAGGCAUAUCCUACGUACCAACAAGAGGUUCAGCA</i> <i>UAGUAGCUAAUUGUAGCUACUUGCUUUAUCCUUAUCCUUAUCCUUAUCCU</i> <i>AUCCUACGUACCAACAAGGACGACGAGACCUUUAUCAAACUUGUUGAUGA</i> <i>AGGUCUCGUGUCCUUUAUCCUAGAGGCAUAUCCUUUUUAUCUAGAGGCA</i> <i>UAUCCCU</i> (all Us are modified; N¹-methylpseudouridine) </p>
17	Compound 17	<p> <i>GCCACCATGTTCCACGTGCTCTCCGGTACATCTTCGGCCTGCCTCCACTG</i> <i>ATCCTGGTGCTGCTGCCTGTGGCCAGCAGCGACTGTGATATCGAGGGCAAA</i> <i>GACGGCAAGCAGTACGAGAGCGTGCTGATGGTGTCATCGACCAAGCTGCTG</i> <i>GACAGCATGAAGGAAATCGGCAGCAACTGCCTGAACAACGAGTTCAACTTC</i> <i>TTCAAGCGGCACATCTGCGACGCCAACAAAGAAGGCATGTTCTGTTCAGA</i> <i>GCCGCCAGAAAGCTGCGGCAGTTCTGAGAGTGAACAGCACCAGGCGACTTC</i> <i>GACCTGCATCTGCTGAAAGTGTCTGAGGGCACCACCATCTGCTGAATTGC</i> <i>ACCGGCCAAGTGAAGGGCAGAAAGCCTGCTGCTCTGGGAGAAGCCAGCCT</i> <i>ACCAAGAGCCTGGAAGAGAACAAGTCCCTGAAAGAGCAGAGAAGCTGAAC</i> <i>GACCTCTGCTTCTGAGCGGCTGCTGCAAGAGATCAAGACCTGCTGGAAC</i> <i>AAGATCCTGATGGGCACCAAGAACAAGTGAATAGTGAGTCGTATTACGTA</i> </p>

SEQ ID NO	Compound	Sequence (5' to 3')
		<p>CCAACAAG<u>AGGTT</u>CAGCATAGTAGCTAACTTG<u>AGCTACTATGCTGAACC</u> <u>TTCT</u>TTTATCTTAGAGGCATATCCCTACGTACCAACAAGCGAATTACTGTGA <u>AAGTCAA</u>ACTT<u>TTGACTTT</u>CACAGTAAT<u>TCGC</u>TTTATCTTAGAGGCATAT CCCTACGTACCAACAAG<u>ACCAGCACACTGAGAATCAA</u>ACTTG<u>TTGATTCTC</u> <u>AGTGTGCTGGTCT</u>TTTATCTTAGAGGCATATCCCTTTTATCTTAGAGGCATA TCCCT</p>
141	Compound 17 RNA sequence	<p>GCCACCAUGUCCACGUGUCCUCCGGUACAUCUUCGGCCUGCCUCCACUG AUCUGGUGUGUGCCUGUGGCCAGCAGCGACUGUGAUUUCGAGGGCAAA GACGGCAAGCAGUACGAGAGCGUGUGUUGGUGUCCAUCCAGCAGCUGCUG GACAGCAUGAAGGAAUUCGGCAGCAACUGCCUGAACAGAGUUCACUUC UCCAAGCGGCACAUCUGCGACGCCAACAAAGAAGGCAUGUCCUGUUCAGA GCCGCCAGAAAGCUGCGGCAGUCCUGAAGAUGAACAGCACCAGCGGACUUC GACCUGCAUCUGCGAAAGUGUGUGAGGGCACCACCAUCCUGCUGAAUUGC ACCGGCCAAGUGAAGGGCAGAAAGCCUGCUGCUGGGAGAAGCCAGCCU ACCAAGAGCCUGGAAGAGAACAAGUCCUGAAAGAGCAGAGAAGCUGAAC GACCUCUGCUUCCUGAAGCGGUGUGCAAGAGAUCAAGACCUGCUGGAAC AAGAUCUGAUGGGCACCACAAAGACACUGAAUAGUGAGUGCUAUUAAACGUA CCAACAAG<u>AGAGGUUCAGCAUAGUAGCUA</u>ACUUG<u>UAGCUACUAGUCUGAACC</u> <u>UUCUUUAUCUUAGAGGCAUAUCCCUACGUACCAACAGCGAAUUCUGUGA</u> <u>AAGUCAAA</u>ACUUG<u>UUGACUUUCACAGUAAUUCGCUUUUAUCUUAGAGGCAUAU</u> CCCUACGUACCAACAAG<u>ACCAGCACACUGAGAAUCAAA</u>CUUG<u>UUGAUUCUC</u> <u>AGUGUGCUGGUCUUUAUCUUAGAGGCAUAUCCCUUUUAUCUUAGAGGCAUAU</u> UCCCU (all Us are modified; N¹-methylpseudouridine)</p>

Bold = Sense siRNA strand

Bold and Italics = Anti-Sense siRNA strand

Underline = Signal peptide

Italics = Kozak sequence

- 5 *Bolding within the underlined sequence indicates modified signal peptide.

[0207] Table 3. Table of Sequences Listed

Protein or Nucleic Acid	Sequence	SEQ ID NO:
Compound 1-6 nucleic acid sequences	See Table 2	1-17
T7 promoter	TAATACGACTCACTATA	18
Kozak sequence	GCCACC	19
tRNA linker	AACAAAGCACCAGTGGTCTAGTGGTAGAATAGTACCCTGCCACGGTACA GACCCGGGTTTCGATTCCCGGCTGGTGCA	20
mRNA to siRNA linker	ATAGTGAGTCGTATTAACGTACCAACAA	21
siRNA to siRNA linker	TTTATCTTAGAGGCATATCCCTACGTACCAACAA	22
Human IL-2 amino acid (Genbank NM_000586.3)	MYRMQLLSICIALSLALVTNSAPTSSSTKKTQLQLEHLLLDLQMI L N G I N N Y K N P KLTRMLTFKFYMPKKATELKHLCLEELKPLEEVLNLAQSKNFHLRPRDLISN INVIVLELKGSETTFMCEYADETATIVEFLNRWITFCQSIISTLT	23
Underlined: signal sequence		
Mature Human IL-2 amino acid (Genbank NM_000586.3)	APTSSSTKKTQLQLEHLLLDLQMI L N G I N N Y K N P K L T R M L T F K F Y M P K K A T E L K HLQCLEELKPLEEVLNLAQSKNFHLRPRDLISNINVIVLELKGSETTFMCEYA DETATIVEFLNRWITFCQSIISTLT	24

Protein or Nucleic Acid	Sequence	SEQ ID NO:
<u>Underlined:</u> signal sequence		
Human IL-2 nucleic acid (Genbank NM_000586.3)	AGTTCCCTATCACTCTCTTTAATCACTACTCACAGTAACCTCAACTCCTGCCACA ATGTACAGGATGCAACTCCTGTCTTGCAATTGCACTAAGTCTTGCACTTGTCACAAACAGTGCACCTACTTCAAGTTCTACAAAGAAAACACAGCTACA ACTGGAGCATTTACTGCTGGATTTCAGATGATTTTGAATGGAATTAATAATTACAAGAATCCCAAACCTCACCAGGATGCTCACATTTAAGTTTACATGCCCAAGAAGGCCACAGAAGTGAACATCTTCAGTGTCTAGAGAGAAGTCAAACCTCTGGAGGAAGTGCTAAATTTAGCTCAAAGCAAAAACTTTCACTTAAGACCCAGGGACTTAATCAGCAATATCAACGTAATAGTTCTGGAACTAAAGGGATCTGAAACAACATTCATGTGTGAATAIGCTGTGAGACAGCAACCATTTGTAGAATTTCTGAACAGATGGATTACCTTTTGTCAAAGCATCATCTCAACACTGACTTGATAATTAAGTGCTTCCCACCTTAACATATCAGGCCTTCTATTATTTAAATATTTAAATTTTATATTTATTGTGAATGTATGGTTTGCTACCTATTGTAACATATTATTCTTAATCTTAAACTATAAATATGGATCTTTTATGATTCTTTTGTAAGCCCTAGGGGCTCTAAAATGGTTTCACCTATTATCCCAAATATTTATTATTATGTTGAA TGTTAAATATAGTATCTATGTAGATTGGTTAGTAAAACATTTTAATAAA TTTGATAAATATAAAAAAAAAAAAAAAAAAAAAAAAAAAAA	25
<u>Underlined:</u> coding sequence		
Bold: signal sequence		
IL-2 signal peptide (Genbank NM_000586.3)	MYRMQLLSICIALSLALVTNS	26
Modified IL-2 signal peptide (Cpd.2) amino acid (Y2L/R3K/M4L/Q5L/S8L/L16A / and V17-)	MLKLLLLLLICIALSLAATNS	27
Modified IL-2 signal peptide (Cpd.3) amino acid (Y2L/R3-/M4L/Q5L/S8A/-A13/L14T/L16A and V17A)	MLLLLLLACIALASTAAATNS	28
Modified IL-2 signal peptide (Cpd.4) amino acid (Y2L/R3-/M4L/Q5L/S8A/-A13 and L14T)	MLLLLLLACIALASTALVTNS	29
Endogenous IL-2 signal peptide (Cpd.1) nucleic acid	ATGTACAGAATGCAGCTGCTGAGCTGTATCGCCCTGTCTCTGGCCCTGGTCACAAATAGC	30
Modified IL-2 signal peptide (Cpd.2) nucleic acid	ATGCTGAAACTGCTGCTGCTCCTGTGTATCGCCCTGTCTCTGGCCGCCACAATAGC	31

Protein or Nucleic Acid	Sequence	SEQ ID NO:
Modified IL-2 signal peptide (Cpd.3) nucleic acid	ATGTTGTTGCTGCTGCTCGCCTGTATTGCCCTGGCCTCTACAGCCGCCGCTACAAATTCT	32
Modified IL-2 signal peptide (Cpd.4) nucleic acid	ATGTTGTTGCTGCTGCTCGCCTGTATTGCCCTGGCCTCTACAGCCCTGGTCACCAATTCT	33
VEGFA amino acid (Genbank NM_001171623 .1) (Transcript variant-1; Canonical sequence; Isoform-206)	MNFLLSWVHWSLALLLLYLHHAKWSQAAPMAEGGGQNHHEVVKFMDVYQRSYCHPIETLVDFQEYPDEIEYIFKPSCVPLMRCGGCCNDEGLECVPTESNITMQIMRIKPHQGQHI GEMSFLOHNKCECRPKKDRARQEKKSVRGKGKGQKRKRKKSRYKSWSVYVGARCCCLMPWSLPGPHPCGPCSERRKHLFVQDPQTCKCSCKNTDSRCKARQLELNERTCRCDKPRR	34
VEGFA (SEQ ID NO: 34) encoding DNA sequence (from Genbank NM_001171623 .1) Bold: signal peptide sequence Bold and italicized: siRNA binding regions	ATGAAC TTTCTGCTGTCTTGGGTGCATTGGAGCCTTGCCCTTGCTGCTCTACCTCCACCATGCCAAGTGGTCCAGGCTGCACCCATGGCA GAAGGAGGAGG GCAGAATCATCACGAAGTGGT GAAGTTCATGGATGTCTATCAGCGCAGCTACTGCCATCCAATCGAGACCCTGGTGGACATCTTCCAGGAGTACCCTGATGAGATCGAGTACATCTTCAAGCCA TCCTGTGTGCCCTGATGCGATGCGGGGGCTGCTGCAATGACGAGGGCCTGGAGTGTGTGCCCACTGAGGAGTCCAACATCACCATGCAGATTATGCGGATCAAACCTCACCAAGGCCAGCACATAGGAGAGAT AGCTTCCTACAGCACACA AAATGTGAATGCAGACCAAAGAAAGATAGAGCAAGACAAGAAAAAAATCAGTTCGAGGAAAGGGAAAGGGGCAAAAACGAAAGCGCAAGAAATCCCGGTATAAGTCCTGGAGCGGTGTACGTTGGTGCCCGCTGCTGTCTAATGCCCTGGAGCCTCCCTGGCCCCATCCCTGTGGGCCTTGCTCAGAGCGGAGAAAGCATTTGTTTGTACAA GATCCGCAGACGTGTAAATGT TCTGCAAAAACACAGACTCGCGTTGCAAGGCGAGGCAGCTTGAGTTAAACGAACGTACTTGCAGATGTGACAAGCCGAGGCGGTGA	35
VEGFA (SEQ ID NO: 34) encoding RNA sequence (from Genbank NM_001171623 .1) Bold: signal peptide sequence Bold and italicized: siRNA binding regions	AUGAAC UUUCUGCUGUCUUGGGUGCAUUGGAGCCUUGCCUUGCUGCUCUACCUCACCAUGCCAAGUGGUCCAGGCUGCACCCAUGGCA GAAGGAGGAGG GCAGAAUCAUCAGAAUGGU GAAGUUCAUGGAUGUCUAUCAGCGCAGCUACUGCCAUCCAUCGAGACCCUGGUGGACAUCUCCAGGAGUACCCUGAUGAGAUCAUACAUCUUAAGCCA UCCUGUGUGCCCCUGAUGCGAUGCGGGGGCUGCUGCAAUGACGAGGGCCUGGAGUGUGUGCCACUGAGGAGUCCAACAUCACCAUGCAGAUUAUGCGGAUCAAAACCUCACCAAGGCCAGCACAUAGGAGAGAU AGCUUCCUACAGCACACA AAUGUGAAUGCAGACCAAAGAAAGAUAGAGCAAGACAAGAAAAAAAUCAUUCGAGGAAAGGGAAAGGGGCAAAAACGAAAGCGCAAGAAAUCCCGGUUAUAAGUCCUGGAGCGUGUACGUUGGUGCCCGUGCUGUCUAAUGCCUGGAGCCUCCUGGC CCCC AUCCUGUGGGCCUUGCUCAGAGCGGAGAAAGCAUUUGUUUGUACAA GAUCCGCAGACGUGUAAAUGU UCCUGCAAAAACACAGACUCGCGUUGCAAGGCGAGGCAGCUUGAGUUAACGAACGUACUUGCAGAUGUGACAAGCCGAGGCGGUGA	36

Protein or Nucleic Acid	Sequence	SEQ ID NO:
MICA amino acid (Genbank NM_000247.2) (Transcript variant 1*001)	MGLGPFVFLLLAGIFPFAPPGAAAEPHSLRYNLTVLSWDGVSQSGF LTEVHLDGQPFLRCDRQKCRAPQGGQWAEDVLGNKTDRETRDLT GNGKDLRMTLAHIKDQKEGLHSLQEIRVCEIHEDNSTRSSQHFFYY DGELFLSQNLETKEWTMPQSSRAQTLAMNVRNFLKEDAMKTKTHY HAMHADCLQELRRYLKSGVVLRRTPPMMVNVTRSEASEGNITVTC RASGFYPWNITLSWRQDGVSLSHDTQQWGDVLPDNGTYQTWVAT RICQGEEQRFTCYMEHSGNHSTHPVPSGKVLVLQSHWQTFHVS AV AAAAIFVIIIFYVRCKKKTSAAEGPELVSLQVLDQHPVGTSDHR DATQLGFQPLMSDLGSTGSTEGA	37
MICA (SEQ ID NO: 37) encoding DNA sequence (from Genbank NM_000247.2) <i>Bold and italicized:</i> siRNA binding regions	ATGGGGCTGGGCCCCGGTCTTCTGCTTCTGGCTGGCATCTTCCCT TTTGACCTCCGGGAGCTGCTGCTGAGCCCCACAGTCTTCGTTAT AACCTCACGGTGTCTGCTGGGATGGATCTGTGCAGTCAGGGTTT CTCACTGAGGTACATCTGGATGGTCAGCCCTTCTGCGCTGTGAC AGGCAGAAATGCAGGGCAAAGCCCCAGGGACAGTGGGCAGAAGAT GTCCTGGGAAATAAGACATGGGACAGAGAGACCAGAGACTTGACA GGGAACGGAAAGGACCTCAGGATGACCCTGGCTCATATCAAGGAC CAGAAAGAAGGCTTGCATTCCCTCCAGGAGATTAGGGTCTGTGAG <i>ATCCATGAAGACAACAGCACCAGGAGCTCCAGCATTTCTACTAC</i> <i>GATGGGGAGCTCTTCTCTCTCCAAAACCTGGAGACTAAGGAATGG</i> <i>ACAATGCCCCAGTCTCTCAGAGCTCAGACCTTGGCCATGAACGTC</i> <i>AGGAATTTCTTGAAGGAAGATGCCATGAAGACCAAGACACACTAT</i> <i>CACGCTATGCATGCAGACTGCCTGCAGGAACACGGCGATATCTA</i> <i>AAATCCGGCGTAGTCCTGAGGAGAACAGTGCCCCCATGGTGAAT</i> <i>GTCACCCGCAGCGAGGCCTCAGAGGGCAACATTACCGTGACATGC</i> <i>AGGGCTTCTGGCTTCTATCCCTGGAATATCACACTGAGCTGGCGT</i> <i>CAGGATGGGGTATCTTTGAGCCACGACACCCAGCAGTGGGGGAT</i> <i>GTCCTGCCTGATGGGAATGGAACCTAC</i> <i>CAGACCTGGGTGGCCACC</i> <i>AGGATTTGCCAAGGAGAGGAGCAGAGGTTACCTGCTACATGGAA</i> <i>CACAGCGGGAATCACAGCACTCACCTGTGCCCTCTGGGAAAGTG</i> <i>CTGGTGCTTCAGAGTCATTGGCAGACATTCCATGTTTCTGCTGTT</i> <i>GCTGCTGCTGCTATTTTTGTATTATTATTTCTATGTCCGTTGT</i> <i>TGTAAGAAGAAAACATCAGCTGCAGAGGGTCCAGAGCTCGTGAGC</i> <i>CTGCAGGTCCTGGATCAACACCCAGTTGGGACGAGTGACCACAGG</i> <i>GATGCCACACAGCTCGGATTTAGCCTCTGATGTCAGATCTTGGG</i> <i>TCCACTGGCTCCACTGAGGGCGCCTAG</i>	38
MICA (SEQ ID NO: 37) encoding RNA sequence (from Genbank NM_000247.2) <i>Bold and italicized:</i> siRNA binding regions	AUGGGGCUGGGCCCCGGUCUUCUGCUUCUGGCUUGGCAUCUCCCU UUUGCACCUCCGGGAGCUGCUGCUGAGCCCCACAGUCUUCGUUAU AACCUCACGGUGCUGUCCUGGGAUGGAUCUGUGCAGUCAGGGUUU CUCACUGAGGUACAUCUGGAUGGUCAGCCCUUCCUGCGCUGUGAC AGGCAGAAUUGCAGGGCAAAGCCCCAGGGACAGUGGGCAGAAGAU GUCCUGGGAAAUAGACAUGGGACAGAGAGACCAGAGACUUGACA GGGAACGGAAAGGACCUCAGGAUGACCCUGGCUCAUAUCAAGGAC CAGAAAGAAGGCUUGCAUUCUCCUCCAGGAGAUUAGGGUCUGUGAG <i>AUCCAUGAAGACAACAGCACCAGGAGCUCCAGCAUUCUACUAC</i> <i>GAUGGGGAGCUCUCCUCUCCAAAACCUGGAGACUAAGGAAUGG</i> <i>ACAAUGCCCCAGUCCUCCAGAGCUCAGACCUUGGCCAUGAACGUC</i> <i>AGGAUUUUCUUGAAGGAAUGAUGCCAUGAAGACCAAGACACACUAU</i> <i>CACGCUAUGCAUGCAGACUGCCUGCAGGAACUACGGCGAUUAUCUA</i> <i>AAAUCCGGCGUAGUCCUGAGGAGAACAGUGCCCCCAUGGUGAAU</i>	39

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	GUCACCCGCGAGCGAGGCCUCAGAGGGCAACAUAUACCGUGACAUGC AGGGCUUCUGGCUUCUAUCCCUUGGAAUAUCACACUGAGCUGGCGU CAGGAUGGGGUAUCUUUGAGCCACGACACCCAGCAGUGGGGGGAU GUCCU GCCUGAUGGGAAUGGAACCU ACCAGACCUGGGUGGCCACC AGGAUUUGCCAAGGAGAGGAGCAGAGGUUACCUGCUACAUGGAA CACAGCGGGAAUCACAGCACUCACCCUGUGCCCUCUGGGAAAGUG CUGGUGCUUCAGAGUCAUUGGCAGACAUUCCAUGUUUCUGCUGUU GCUGCUGCUGCUAUUUUUGUUUAUUUAUUUUUCUAUGUCCGUUGU UGUAAGAAGAAAACAUCAGCUGCAGAGGGUCCAGAGCUCGUGAGC CUGCAGGUCCUGGAUCAACACCCAGUUGGGACGAGUGACCACAGG GAUGCCACACAGCUCGGAUUUCAGCCUCUGAUGUCAGAUCUUGGG UCCACUGGCUCCACUGAGGGCGCCUAG	
MICB amino acid (Genbank NM_005931.4) (Transcript variant 1)	MGLGRVLLFLAVAFPFAPPAAAAEPHSLRYNLMVLSQDGSVQSGF LAEGHLDGQPFRLYDRQKRAKFPQGWAENVLGAKTWDTEDELT ENGQDLRRTLTHIKDQKGLHSLQEIRVCEIHEDSSTRGSRHFYY DGELFLSQNLETQESTVPQSSRAQTLAMNVNFWKEDAMKTKTHY RAMQADCLQKLQRYLKSGVAIRRTVPPMVNVTCSEVSEGNITVTC RASSFYPRNITLTWRQDGVSLSHNTQQWGDVLPDGNQTYQTWVAT RIRQGEEQRFTCYMEHSGNHGTHPVPSGKALVLQSQRTDFPYVSA AMPCFVIIILCVPCKKKTSAAEGPPELVSLQVLDQHPVGTGDHR DAAQLGFQPLMSATGSTGSTE	40
MICB (SEQ ID NO: 40) encoding DNA sequence (from Genbank NM_005931.4) <i>Bold and italicized:</i> siRNA binding regions	ATGGGGCTGGGCGGGTCTCTGCTGTTTCTGGCCGTCGCTTCCCT TTTGACCCCCCGGCAGCCGCGCTGAGCCCCACAGTCTTCGTTAC AACCTCATGGTGCTGTCCCAGGATGGATCTGTGCAGTCAGGGTTT CTCGCTGAGGGACATCTGGATGGTCAGCCCTTCTGCGCTATGAC AGGCAGAAACGCAGGGCAAAGCCCCAGGCACAGTGGGCAGAAAAT GTCCTGGGAGCTAAGACCTGGGACACAGAGACCGAGGACTTGACA GAGAATGGGCAAGACCTCAGGAGGACCTGACTCATATCAAGGAC CAGAAAGGAGGCTTGCAATCCCTCCA GGAGATTAGGGTCTGTGAG ATCCATGAAGACAGCAGCACCAGGGGCTCCCGGCATTTCTACTAC GATGGGGAGCTCTTCCTCTCCCAAAACCTGGAGACTCAAGAATCG ACAGTGCCCCAGTCTCCAGAGCTCAGACCTTGGCTATGAACGTC ACAAATTTCTGGAAGGAA GATGCCATGAAGACCAAGACACACTAT CGCGCTATGCAGGCAGACTGCCTGCAGAACTACAGCGATATCTG AAATCCGGGGTGGCCATCAGGAGAACAGTGCCCCCATGGTGAAT GTCACCTGCAGCGAGGTCTCAGAGGGCAACATCACCGTGACATGC AGGGCTTCCAGCTTCTATCCCCGGAATATCACACTGACCTGGCGT CAGGATGGGGTATCTTTGAGCCACAACACCCAGCAGTGGGGGGAT GTCCT GCCTGATGGGAATGGAACCTA CCAGACCTGGGTGGCCACC AGGATTGCGCAAGGAGAGGAGCAGAGGTTACCTGCTACATGGAA CACAGCGGGAATCACGGCACTCACCTGTGCCCTCTGGGAAGGCG CTGGTGCTTTCAGAGTCAACGGACAGACTTTCATATGTTTCTGCT GCTATGCCATGTTTTGTTATTATTATTATTCTCTGTGTCCCTTGT TGCAAGAAGAAAACATCAGCGGCAGAGGGTCCAGAGCTTGTGAGC CTGCAGGTCTTGATCAACACCCAGTTGGGACAGGAGACCACAGG GATGCAGCACAGCTGGGATTTTCAGCCTCTGATGTCAGCTACTGGG TCCACTGGTTCCACTGAGGGCACCTAG	41
MICB (SEQ ID NO: 40)	AUGGGGCUUGGGCCGGGUCCUGCUGUUUCUGGGCGUCGCCUCCCCU UUUGCACCCCCCGGCAGCCGCGCUGAGCCCCACAGUCUUCGUUAC	42

Protein or Nucleic Acid	Sequence	SEQ ID NO:
encoding RNA sequence (from Genbank NM_005931.4) <i>Bold and italicized:</i> siRNA binding regions	AACCUCUAUGGUGCUGUCCCAGGAUGGAUCUGUGCAGUCAGGGUUU CUCGCUGAGGGACAUCUGGAUGGUCAGCCCUUCCUGCGCUAUGAC AGGCAGAAACGCAGGGCAAAGCCCCAGGGACAGUGGGCAGAAAAU GUCCUGGGAGCUAAGACCUGGGACACAGAGACCAGGACUUGACA GAGAAUGGGCAAGACCUCAGGAGGACCCUGACUCAUAUCAAGGAC CAGAAAGGAGGCUUGCAUCCCCUCC <i>AGGAGAUUAGGGUCUGUGAG</i> <i>AUCCAUGAAGACAGCAGCACCAGGGGCUCCCGGCAUUUCUACUAC</i> <i>GAUGGGGAGCUCUCCUCUCCCAAACCCUGGAGACUCAAGAAUCG</i> <i>ACAGUGCCCCAGUCCUCCAGAGCUCAGACCUUGGCUAUGAACGUC</i> <i>ACAAAUUUCUGGAAGGAA</i> <i>GAUGCCAUGAAGACCAAGACACACUAU</i> <i>CGCGCUAUGCAGGCAGACUGCCUGCAGAAACUACAGCGAUUAUCUG</i> <i>AAAUCCGGGGUGGGCAUCAGGAGAACAGUGCCCCCAUGGUGAAU</i> <i>GUCACCUGCAGCGAGGUCUCAGAGGGCAACAUCACCGUGACAUGC</i> <i>AGGGCUUCCAGCUUCUAUCCCCGAAUAUCACACUGACCUGGCGU</i> <i>CAGGAUGGGGUUAUCUUUGAGCCACAACACCCAGCAGUGGGGGGAU</i> <i>GUCCUGCCUGAUGGGAAUGGAACCUA</i> <i>ACCAGACCUGGGUGGCCACC</i> <i>AGGAUUCGCCAAGGAGAGGAGCAGAGGUUACCUUGCUACAUGGAA</i> <i>CACAGCGGGAUUCACGGCACUCACCCUGUGCCCUUGGGAAAGCG</i> <i>CUGGUGCUUCAGAGUCAACGGACAGACUUUCCAUAUGUUUCUGCU</i> <i>GCUAUGCCAUGUUUUGUUUAUUUAUUUAUUCUCUGUGUCCCUUGU</i> <i>UGCAAGAAGAAAACAUCAGCGGCAGAGGGUCCAGAGCUUGUGAGC</i> <i>CUGCAGGUCCUGGAUCAACACCCAGUUGGGACAGGAGACCACAGG</i> <i>GAUGCAGCACAGCUGGGAUUUCAGCCUCUGAUGUCAGCUACUGGG</i> <i>UCCACUGGUUCCACUGAGGGCACCUG</i>	
Human IL-12 alpha amino acid (Genbank NM_000882.4) <u>Underlined:</u> signal sequence	MCPARSLLLVATLVLLDHLISLARNLPVATPDPGMFPC LHHSQNLL RAVSNMLQKARQTLEFYPTSEEIDHEDITKDKTSTVEACLPLEL TKNESCLNSRETSFITNGSCLASRKTSFMMALCLSS IYEDLKMYQ VEFKTMNAKL LMDPKRQIFLDQNM LAVIDELMQALNFNSETVFPQK SSLEEPDFYKTKIKLCIL LHAFRIRAVTIDRVMSYLNAS	43
Mature Human IL-12 alpha amino acid (Genbank NM_000882.4)	RNL PVATPDPGMFPC LHHSQNLLRAVSNMLQKARQTLEFYPTSE EIDHEDITKDKTSTVEACLPLELTKNESCLNSRETSFITNGSCLA SRKTSFMMALCLSS IYEDLKMYQVEFKTMNAKL LMDPKRQIFLDQ NMLAVIDELMQALNFNSETVFPQKSSLEEPDFYKTKIKLCIL LHAF RIRAVTIDRVMSYLNAS	44
Human IL-12 alpha nucleic acid (Genbank NM_000882.4) <u>Underlined:</u> coding sequence Bold: signal sequence	ATTTTCGCTTTTCATTTTGGGCCGAGCTGGAGGCGGCGGGGCCGTCC CGGAACGGCTGCGGCCGGGCACCCCGGGAGTTAATCCGAAAGCGC CGCAAGCCCCGCGGGCCGGCCGCACCGCACGTGTCACCGAGAAGC TGATGTAGAGAGAGACACAGAAGGAGACAGAAAGCAAGAGACCAG AGTCCCGGGAAGTCTCTGCCGCGCCTCGGGACAATTATAAAAAATG TGGCCCCCTGGGT CAGCCTCCAGCCACCGCCCTCACCTGCCGCG GCCACAGGTCTGCATCCAGCGGCTCGCCCTGTGTCCCTGCAGTGC CGGCTCAGC <i>ATGTGTCCAGCGCGCAGCCTCCTCCTTGTGGCTACC</i> <i>CTGGTCCTCCTGGACCACCTCAGTTTGGCCAGAAACCTCCCCGTG</i> <i>GCCACTCCAGACCCAGGAATGTTCCCATGCCTTCACCACTCCCAA</i> <i>AACCTGTCTAGGGCCGTCAGCAACATGCTCCAGAAGGCCAGACAA</i> <i>ACTCTAGAATTTTACCCTTGCACTTCTGAAGAGATTGATCATGAA</i> <i>GATATCACAAAAGATAAAACCAGCACAGTGGAGGCCTGTTTACCA</i> <i>TTGGAATTAACCAAGAATGAGAGTTGCCTAAATTCAGAGAGACC</i>	45

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	<p>TCTTTCATAACTAATGGGAGTTGCCTGGCCTCCAGAAAGACCTCT TTTATGATGGCCCTGTGCCTTAGTAGIATTTATGAAGACTTGAAG ATGTACCAGGTGGAGTTCAAGACCATGAATGCAAAGCTTCTGATG GATCCTAAGAGGCAGATCTTCTAGATCAAAACATGCTGGCAGTT ATTGATGAGCTGATGCAGGCCCTGAATTTCAACAGTGAGACTGTG CCACAAAAATCCTCCCTTGAAGAACCGGATTTTTATAAACTAAA ATCAAGCTCTGCATACTTCTTCATGCTTTCAGAATTCGGGCAGTG ACTATTGATAGAGTGATGAGCTATCTGAATGCTTCCTAAAAAGCG AGGTCCCTCCAAACCGTTGTCATTTTTATAAACTTTGAAATGAG GAACTTTGATAGGATGTGGATTAAAGAACTAGGGAGGGGGGAAAAG AGGATGGGACTATTACATCCACATGATACCTCTGATCAAGTATTT TTGACATTTACTGTGGATAAATTGTTTTTAAGTTTTTCATGAATGA ATTGCTAAGAAGGGGAAAATATCCATCCTGAAGGTGTTTTTCATTC ACTTTAATAGAAGGG</p>	
Human IL-12 beta amino acid (Genbank NM_002187.2) <u>Underlined:</u> signal sequence	<p>MCHQQLVISWFSLVFLASPLVAIWELKKDVYVVELDWYPDAPGEM VVLTCDTPEEDGITWTLDQSSEVLGSGKLTITQVKEFGDAGQYTC HKGGEVLSHSLLLLHKKEDGIWSTDILKDQKEPKNKTFLRCEAKN YSGRFTCWLLTTISTDLTFSVKSSRGSSDPQGVTCGAATLSAERV RGDNKEYEYSVEQCEDSACPAAEESLPIEVMVDAVHKLKYENYTS SFFIRDIIKPDPPKNLQLKPLKNSRQVEVSWEYPDTWSTPHSYFS LTFVQVQGKSKREKKDRVFTDKTSATVICRKNASISVRAQDRYY SSSWSEWASVPCS</p>	46
Mature Human IL-12 beta amino acid (Genbank NM_002187.2)	<p>IWELKKDVYVVELDWYPDAPGEMVVLTCDTPEEDGITWTLDQSSE VLGSGKLTITQVKEFGDAGQYTC HKGGEVLSHSLLLLHKKEDGIW STDILKDQKEPKNKTFLRCEAKNYSGRFTCWLLTTISTDLTFSVK SSRGSSDPQGVTCGAATLSAERV RGDNKEYEYSVEQCEDSACPAA EESLPIEVMVDAVHKLKYENYTS SFFIRDIIKPDPPKNLQLKPLK NSRQVEVSWEYPDTWSTPHSYFSLTFVQVQGKSKREKKDRVFTD KTSATVICRKNASISVRAQDRYYSSSWSEWASVPCS</p>	47
Human IL-12 beta nucleic acid (Genbank NM_002187.2) <u>Underlined:</u> coding sequence Bold: signal sequence	<p>CTGTTTCAGGGCCATTGGACTCTCCGTCCTGCCAGAGCAAGATG TGTACCAGCAGTTGGTCATCTCTTGGTTTTCCCTGGTTTTTCTG GCATCTCCCCTCGTGGCCATATGGGAACTGAAGAAAGATGTTTAT GTCGTAGAATTGGATTGGTATCCGGATGCCCTGGAGAAATGGTG GTCCTCACCTGTGACACCCCTGAAGAAGATGGTATCACCTGGACC TTGGACCAGAGCAGTGAGGTCTTAGGCTCTGGCAAAACCTGACC ATCCAAGTCAAAGAGTTTGGAGATGCTGGCCAGTACACCTGTCAC AAAGGAGGCGAGGTTCTAAGCCATTTCGCTCCTGCTGCTTCACAAA AAGGAAGATGGAATTTGGTCCACTGATATTTTAAAGGACCAGAAA GAACCCAAAAATAAGACCTTTCTAAGATGCGAGGCCAAGAATTAT TCTGGACGTTTCACCTGCTGGTGGCTGACGACAATCAGTACTGAT TTGACATTCACTGTCAAAGCAGCAGAGGCTCTTCTGACCCCCAA GGGGTGACGTGCGGAGCTGCTACACTCTCTGCAGAGAGAGTCAGA GGGGACAACAAGGAGTATGAGTACTCAGTGGAGTGCCAGGAGGAC AGTGCCTGCCAGCTGCTGAGGAGAGTCTGCCCATGAGGTCATG GTGGATGCCGTTTACAAGCTCAAGTATGAAAACCTACACCAGCAGC TTCTTCATCAGGGACATCATCAAACCTGACCCACCCAAGAACTTG CAGCTGAAGCCATTAAAGAATTCTCGGCAGGTGGAGGTGAGCTGG GAGTACCCTGACACCTGGAGTACTCCACATTCTTCTTCTCCCTG ACATTCTGCGTTTCAAGTCCAGGGCAAGAGCAAGAGAGAAAAAGAAA</p>	48

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	GATAGAGTCTTCACGGACAAGACCTCAGCCACGGTCATCTGCCGC AAAAATGCCAGCATTAGCGTGCGGGCCAGGACCGCTACTATAGC TCATCTTGGAGCGAATGGGCATCTGTGCCCTGCAGTTAGGTTCTG ATCCAGGATGAAAATTTGGAGGAAAAGTGAAGATATTAAGCAAA ATGTTTAAAGACACAACGGAATAGACCCAAAAAGATAATTTCTAT CTGATTTGCTTTAAAACGTTTTTTTAGGATCACAATGATATCTTT GCTGTATTTGTATAGTTAGATGCTAAATGCTCATTGAAACAATCA GCTAATTTATGTATAGATTTTCCAGCTCTCAAGTTGCCATGGGCC TTCAIGCTATTTAAATATTTAAGTAATTTATGTATTTATTAGTAT ATTACTGTTATTTAACGTTTGTCTGCCAGGATGTATGGAATGTTT CATACTCTTATGACCTGATCCATCAGGATCAGTCCCTATTATGCA AAATGTGAATTTAAT	
IDH1 amino acid (Genbank NM_005896.3) (Transcript variant 1)	MSKKISGGSVVEMQGDGMTRIIWELIKEKLIFPYVELDLHSYDLG IENRDATNDQVTKDAAEAIKKNHNVGKCATITPDEKRVEEFKLKQ MWKSPNGTIRNILGGTVFREAIICKNIPRLVSGWVKPIIIGRHAY GDQYRATDFVVPGPVKVEITYTPSDGTQKVITYLVHNFEEGGGVAM GMYNQDKSIEDFAHSSFQMALSKGWPLYLSTKNITLKKYDGRFKD IFQEIYDKQYKSQFEAQKIWEHRLIDDMVAQAMKSEGGFIWACK NYDGDVQSDSVAQGYGSLGMMTSVLVCPDGKTVEAEAAHGTVTRH YRMYQKGQETSTNPIASIFAWTRGLAHRAKLDNNKELAFFANALE EVSIEETIEAGFMTKDLAACIKGLPNVQRSDYLNTFEFMDKLGLENL KIKLAQAKL	49
IDH1 amino acid encoding DNA sequence (from Genbank NM_005896.3) <i>Bold and italicized:</i> siRNA binding region	ATGTCCAAAAAATCAGTGCGGTTCTGTGGTAGAGATGCAAGGA GATGAAATGACACGAATCATTTGGGAATTGATTAAAGAGAACTC ATTTTTCCCTACGTGGAATTGGATCTACATAGCTATGATTTAGGC ATAGAGAATCGTGATGCCACCAACGACCAAGTCACCAAGGATGCT GCAGAAGCTATAAAGAAGCATAATGTTGGCGTCAAATGTGCCACT ATCACTCCTGATGAGAAGAGGGTTGAGGAGTTCAAGTTGAAACAA ATGTGGAAATCACCAATGGCACCATACGAAATATTCTGGGTGGC ACGGTCTTCAGAGAAGCCATTATCTGCAAAAAATATCCCCCGGCTT GTGAGTGGATGGGTAAAACCTATCATCATAGGTCGTGATGCTTAT GGGGATCAATACAGAGCAACTGATTTTGTGTCTTCTGGGCTTGGG AAAGTAGAGATAACCTACACACCAAGTGACGGAACCCAAAAGGTG ACATACCTGGTACATAACTTTGAAGAAGGTGGTGGTGTGTCATG GGGATGTATAATCAAGATAAGTCAATTGAAGATTTGCACACAGT <i>TCCTTCCAAATGGCTCTGT</i> CTAAGGGTTGGCCTTTGTATCTGAGC ACCAAAAACACTATTCTGAAGAAATATGATGGGCGTTTTAAAGAC ATCTTTCAGGAGATATATGACAAGCAGTACAAGTCCCAGTTTGAA GCTCAAAAGATCTGGTATGAGCATAGGCTCATCGACGACATGGTG GCCCAAGCTATGAAATCAGAGGGAGGCTTCATCTGGGCTGTAAA AACTATGATGGTGACGTGCAGTCGGACTCTGTGGCCCAAGGGTAT GGCTCTCTCGGCATGATGACCAGCGTGCTGGTTTGTCCAGATGGC AAGACAGTAGAAGCAGAGGCTGCCACGGGACTGTAACCCGTCAC TACCGCATGTACCAGAAAGGACAGGAGACGTCCACCAATCCCAT GCTTCCATTTTTGCTTGGACAGAGGGTTAGCCACAGAGCAAAG CTTGATAACAATAAAGAGCTTGCCTTCTTTGCAATGCTTTGGAA GAAGTCTCTATTGAGACAATTGAGGCTGGCTTCATGACCAAGGAC TTGGCTGCTTGCAATAAAGGTTTACCCAATGTGCAACGTTCTGAC	50

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	TACTTGAATACATTTGAGTTTCATGGATAAACTTGGAGAAAACTTG AAGATCAAACCTAGCTCAGGCCAAACTTTAA	
IDH1 amino acid encoding RNA sequence (from Genbank NM_005896.3) <i>Bold and italicized: siRNA binding region</i>	AUGUCCAAAAAAUUCAGUGGCGGUUCUGUGGUAGAGAUGCAAGGA GAUGAAAUGACACGAAUCAUUUGGGAAUUGAUUAAAAGAGAAACUC AUUUUUCCCUACGUGGAAUUGGAUCUACAUAGCUAUGAUUUAGGC AUAGAGAAUCGUGAUGCCACCAACGACCAAGUCACCAAGGAUGCU GCAGAAGCUAUAAGAAGCAUAAUGUUGGCGUCAAAUGUGCCACU AUCACUCCUGAUGAGAAGAGGGUUGAGGAGUUCAGUUGAAACAA AUGUGGAAAUCACCAAUGGCACCAUACGAAAUAUUCUGGGUGGC ACGGUCUUCAGAGAAGCCAUUAUCUGCAAAAAUAUCCCCCGGCUU GUGAGUGGAUGGGUAAAACCUAUCAUCAUAGGUCGUCUAGCUUUAU GGGGAUCAUACAGAGCAACUGAUUUUGUUGUCCUGGGCCUGGA AAAGUAGAGAUAAACUACACACCAAGUGACCGGAACCCAAAAGGUG ACAUACCUGGUACAUAACUUUGAAGAAGGUGGUGGUGUUGCCAUG GGGAUGUAUAAUCAAGAUAAUGUCAUUGAAGAUUUUGCACACAGU <i>UCCUCCAAAUGGCUCUGUC</i> UAAGGGUUGGCCUUUGUAUCUGAGC ACCAAAAACACUAUUCUGAAGAAAUAUGAUGGGCGUUUAAAAGAC AUCUUUCAGGAGAUUAUGACAAGCAGUACAAGUCCAGUUUGAA GCUCAAAAGAUUCUGGUAUGAGCAUAGGCUCAUCGACGACAUGGUG GCCCAAGCUAUGAAAUCAGAGGGAGGCUUCAUCUGGGCCUGUAAA AACUAUGAUGGUGACGUGCAGUCGGACUCUGUGGCCCAAGGGUUAU GGCUCUCUCGGCAUGAUGACCAGCGUGCUGGUUUUGUCCAGAUGGC AAGACAGUAGAAGCAGAGGCUGCCCACGGGACUGUAACCCGUCAC UACCGCAUGUACCAGAAAGGACAGGAGACGUCCACCAAUCCCAU GCUUCCAUUUUUGCCUGGACCAGAGGGUUAGCCACAGAGCAAAG CUUGAUAAACAAUAAAGAGCUUGCCUUCUUUGCAAAUGCUUUGGAA GAAGUCUCUAUUGAGACAAUUGAGGCUGGCUUCAUGACCAAGGAC UUGGCUGCUUGCAUUAAGGUUUACCCAAUGUGCAACGUUCUGAC UACUUGAAUACAUUUGAGUUAUGGAUAAACUUGGAGAAAAACUUG AAGAUCAAACUAGCUCAGGCCAAACUUUAA	51
CDK4 amino acid (Genbank NM_000075.3)	MATSRYPVAEIGVGAYGTVYKARDPHSGHFVALKSVRVPNGGGG GGGLPISTVREVALLRLEAFEHPNVVRLMDVCATSRTDREIKVT LVFEHVDQDLRTYLDKAPPPGLPAETIKDLMRQFLRGLDFLHANC IVHRDLKPENILVTSGGTVKLADFLGLARIYSYQMALTPVVVTLWY RAPEVLLQSTYATPVDMWVSGCIFAE MFRRKPLFCGNSEADQLGK IFDLIGLPPEDDWPRDVSLPRGAFPPRGPRPVQSVVPEMEESGAQ LLEMLTFNPHKRISAFRALQHSYLHKDEGNPE	52
CDK4 encoding DNA sequence (from Genbank NM_000075.3) <i>Bold and italicized: siRNA binding regions</i>	ATGGCTACCTCTCGATATGAGCCAGTGGCTGAAATTGGTGTCCGT GCCTATGGGACAGTGTAACAAGGCCCGTGATCCCCACAGTGGCCAC TTTGTGGCCCTCAAGAGTGTGAGAGTCCCCAATGGAGGAGGAGGT GGAGGAGGCCTTCCCATCAGCACAGTTCGTGAGGTGGCTTTACTG AGGCGACTGGAGGCTTTTGAGCATCCCAATGTTGTCCGGCTGATG GACGTCTGTGCCACATCCCGAACTGACCGGGAGATCAAGGTAACC CTGGTGTTTGAGCATGTAGACCAGGACCTAAGGACATATCTGGAC AAGGCACCCCCACCAGGCTTGCCAGCCGAAACGATCAAGGATCTG ATGCGCCAGTTTCTAAGAGGCCTAGATTTCCTTCATGCCAATT <i>GC</i> <i>ATCGTTCACCGAGATCTGA</i> AGCCAGAGAACATTCTGTTGACAAGT GGTGGAACAGTCAAGCTGGCTGACTTTGGCCTGGCCAGAATCTAC AGCTACCAGATGGCACTTACACCCGTGGTTGTTACTCTGGTAC	53

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	CGAGCTCCCGAAGTTCTTCTGCAGTCCACATATGCAACACCTGTG GACATGTGGAGTGTGGCTGTATCTTTGCAGAGATGTTTCGTCGA AAGCCTCTCTTCTGTGGAACTCTGAAGCCGACCAGTTGGGCAAA ATCTTTGACCTGATTGGGCTGCCTCCAGAGGATGACTGGCCTCGA GATGTATCCCTGCCCCGTGGAGCCTTTCCCCCAGAGGGCCCCGC CCAGTGCAGTCGGTGGTACCTGAGATGGAGGAGTCGGGAGCACAG CTGCTGCTGGAAATGCTGACTTTTAACCCACACAAGCGAATCTCT GCCTTTCGAGCTCTGCAGCACTCTTATCTACATAAGGATGAAGGT AATCCGGAGTGA	
CDK4 encoding RNA sequence (from Genbank NM_000075.3) Bold and italicized: siRNA binding regions	AUGGCUACCUCUGAUAUGAGCCAGUGGCUGAAAUUGGUGUCGGU GCCUAUGGGACAGUGUACAAGGCCCGUGAUCCCCACAGUGGCCAC UUUGUGGGCCCUCAAGAGUGUGAGAGUCCCCAAUGGAGGAGGAGGU GGAGGAGGCCUUCCCAUCAGCACAGUUCGUGAGGUGGCUUUACUG AGGCGACUGGAGGCUUUUGAGCAUCCCAAUGUUGUCCGGCUGAUG GACGUCUGUGCCACAUCCCGAACUGACCGGGAGAUCAAGGUAACC CUGGUGUUUGAGCAUGUAGACCAGGACCUAAGGACAUUUCUGGAC AAGGCACCCCCACCAGGCUUGCCAGCCGAAACGAUCAAGGAUCUG AUGCGCCAGUUUCUAAGAGGCCUAGAUAUCCUUCUUGCCAAUUGC AUCGUUCACCGAGAUUCUGAAGCCAGAGAACAUCUGGUGACAAGU GGUGGAACAGUCAAGCUGGCUGACUUUGGCCUGGCCAGAAUCUAC AGCUACCAGAUGGCACUUACACCCGUGGUUGUACACUCUGGUAC CGAGCUCCCGAAGUUCUUCUGCAGUCCACAUAUGCAACACCUGUG GACAUGUGGAGUGUUGGCUGUAUCUUUGCAGAGAUUUUCGUCGA AAGCCUCUCUUCUGUGGAAACUCUGAAGCCGACCAGUUGGGCAAA AUCUUUGACCUGAUUGGGCUGCCUCCAGAGGAUGACUGGCCUCGA GAUGUAUCCUGCCCCGUGGAGCCUUUCCCCCAGAGGGCCCCGC CCAGUGCAGUCGGUGGUACCUGAGAUGGAGGAGUCGGGAGCACAG CUGCUGCUGAAAUGCUGACUUUUAACCCACACAAGCGAAUCUCU GCCUUUCGAGCUCUGCAGCACUCUUAUCUACAUAAGGAUGAAGGU AAUCCGGAGUGA	54
CDK6 amino acid (Genbank NM_001259.6)	MEKDGLCRADQQYECVAEIGEGAYGKVFKARDLKNNGRFVALKRV RVQTGEEGMPLSTIREVAVLRLHLETFEHPNVVRLFDVCTVSRTDR ETKLTLVFEHVDQDLTTYLDKVPFPGVPTETIKDMMFQLLRGLDF LHSHRVVHRDLKFPQNILVTSSQIKLADFGGLARIYSFQMALTSVV VTLWYRAPEVLLQSSYATPVDLWSVGCIFAEMFRRKPLFRGSSDV DQLGKILDVIGLPGEEDWPRDVALPRQAFHSKSAQPIEFKFTDID ELGKDLKKLCLTFNPAKRISAYSALSHPHYQDLERCKENLDLHLP PSQNTSELNTA	55
CDK6 encoding DNA sequence (from Genbank NM_001259.6) Bold and italicized: siRNA binding regions	ATGGAGAAGGACGGCCTGTGCCGCGCTGACCAGCAGTACGAATGC GTGGCGGAGATCGGGGAGGGCGCCTATGGGAAGGTGTTCAAGGCC CGCGACTTGAAGAACGGAGGCCGTTTTCGTGGCGTTGAAGCGCGTG CGGGTGCAGACCGGCGAGGAGGGCATGCCGCTCTCCACCATCCGC GAGGTGGCGGTGCTGAGGCACCTGGAGACCTTCGAGCACCCCAAC GTGGTCAGGTTGTTTGATGTGTGCACAGTGTACGAACAGACAGA GAAACCAACTAAGTTAGTGTGTTGAACATGTCGATCAAGACTTG ACCACTTACTTGGATAAAGTTCAGAGCCTGGAGTGCCCACTGAA ACCATAAAGGATATGATGTTTCAGCTTCTCCGAGGTCTGGACTTT CTTCATTCACACCGAGTAGTGATCGCGATCTAAAACACAGAAC ATTCTGGT GACCAGCAGCGGACAAAATAAA ACTCGCTGACTTCGGC	56

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	CTTGCCCGCATCTATAGTTTCCAGATGGCTCTAACCTCAGTGGTC GTCACGCTGTGGTACAGAGCACCCGAAGTCTTGCTCCAGTCCAGC TACGCCACCCCGTGGATCTCTGGAGTGTGGCTGCATATTTGCA GAAATGTTTCGTAGAAAGCCTCTTTTTCGTGGAAGTTCAGATGTT GATCAACTAGGAAAAATCTTGACGTGATTGGACTCCAGGAGAA GAAGACTGGCCTAGAGATGTTGCCCTTCCCAGGCAGGCTTTTCAT TCAAAATCTGCCCAACCAATTGAGAAGTTTGTAAACAGATATCGAT GAACTAGGCAAAGACCTACTTCTGAAGTGTGACATTTAACCCA GCCAAAAGAATATCTGCCTACAGTGCCTGTCTCACCATACTTC CAGGACCTGGAAAGGTGCAAAGAAAACCTGGATTCCCACCTGCCG CCCAGCCAGAACACCTCGGAGCTGAATACAGCCTGA	
CDK6 encoding RNA sequence (from Genbank NM_001259.6) Bold and italicized: siRNA binding regions	AUGGAGAAGGACGGCCUGUGCCGCGCUGACCAGCAGUACGAAUGC GUGGCGGAGAUCCGGGAGGGCGCCUAUGGGAAGGUGUUAAGGCC CGCGACUUGAAGAACGGAGGCGGUUCGUGGCGUUGAAGCGCGUG CGGGUGCAGACCGGCGAGGAGGGCAUGCCGCUCUCCACCAUCCGC GAGGUGGCGGUGCUGAGGCACCUUGAGACCUUCGAGCACCCCAAC GUGGUCAGGUUGUUGAUGUGUGCAGAGUGACGAAACAGACAGA GAAACCAACUAACUUUAGUGUUUGAACAUGUCGAUCAAGACUUG ACCACUUACUUGGAUAAAGUCCAGAGCCUGGAGUGCCACUGAA ACCAUAAAGGAUAUGAUGUUUCAGCUUCUCCGAGGUCUGGACUUU CUUCAUUCACACCGAGUAGUGCAUCGCGAUCUAAAACACAGAAC AUUCUGGU GACCAGCAGCGGACAAAUAAAA CUCGCUGACUUCGGC CUUGCCCGCAUCUAUAGUUUCCAGAUUGGCUUAACCUCAGUGGUC GUCACGCUGUGGUACAGAGCACCCGAAGUCUUGCUCCAGUCCAGC UACGCCACCCCGUGGAUCUCUGGAGUGUUGGUCUGCAUUAUUGCA GAAUUGUUUCGUAGAAAGCCUCUUUUUCUGGAAGUUCAGAUUU GAUCAACUAGGAAAAAUCUUGGACGUGAUUGGACUCCAGGAGAA GAAGACUGGCCUAGAGAUGUUGCCCUUCCAGGCAGGCUUUUCAU UCAAAUUCUGCCCAACCAAUUGAGAAGUUUGUAACAGAUUUCGAU GAACUAGGCAAAGACCUACUUCUGAAGUUUGACAUUUAACCCA GCCAAAAGAAUUCUGCCUACAGUGCCUGUCUACCCAUACUUC CAGGACCUGGAAAGGUGCAAAGAAAACUGGAUUCCACCCUGCCG CCCAGCCAGAACACCUCGGAGCUGAAUACAGCCUGA	57
EGFR amino acid (Genbank NM_005228.4) (Transcript variant 1)	MRPSGTAGAALLALLAALCPASRALEEKKVCQGTSNKLTLQGTFE DHFLSLQRMFNNEVVLGNLEITYVQRNYDLSFLKTIQEVAGYVL IALNTVERIPLLENLQIIRGNMYYENSALAVLSNYDANKTGLKEL PMRNLQEILHGAVRFSNNPALCNVESIQWRDIVSSDFLSNMSMDF QNHLGSCQKCDPSCPNGSCWGAGEENCQKLTKIICAQQCSGRCRG KSPSDCCHNQCAAGCTGPRESCLVCRKFRDEATCKDTCPLMLY NPTTYQMDVNPEGKYSFGATCVKKCPRNYVVTDHGSCVRACGADS YEMEEDGVRKCKKCEGPCKRVCNGIGIGEFKDSLSINATNIKHFK NCTSIISGDLHILPVAFRGDSFTHTPPLDPQELDKTVKEITGFL LIQAWPENRTDLHAFENLEIIRGRKQHGQFSLAVVSLNITSLGL RSLKEISDGDVIIISGNKNLCYANTINWKKLFGTSGQKTKIISNRG ENSCKATGQVCHALCSPEGCWGPEPRDCVSCRNVSRGRECVDKCN LLEGEPRFVENSECIQCHPECLPQAMNITCTGRGPDNCIQCAHY IDGPHCVKTCPAGVMGENNTLVWKYADAGHVCHLCHPNCTYGCTG PGLEGCPNTPGPKIPSIATGMVGALLLLLVVALGIGLFMRRRHIVR KRTLRLQLQERELVEPLTPSGEAPNQALLRLILKETEFKKIKVLGS	58

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	GAFGTVYKGLWIPEGEKVKIPVAIKELREATSPKANKEILDEAYV MASVDNPHVCRLLGICLTSTVQLITQLMPFGCLLDYVREHKDNIG SQYLLNWCVQIAKGMNYLEDRLVHRDLAARNVLVKTPQHVKITD FGLAKLLGAEEKEYHAEGGKVPKWMMALESILHRIYTHQSDVWSY GVTVWELMTFGSKPYDGIPASEISSILEKGERLPQPPICTIDVYM IMVKCWMIDADSRPKFRELIIEFSKMARDPQRYLVIQDERMHLP SPTDSNFYRALMDEEDMDDVVDADLEYLIPQQGFFSSPSTSRTPLL SSLATSNNSTVACIDRNLQSCPIKEDSFLQRYSSDPTGALTED SIDDTFLPVPEYINQSVKRPAGSVQNPVYHNQPLNPAPSRDPHY QDPHSTAVGNPEYLNNTVQPTCVNSTFDSPAHWAKGSHQISLDNP DYQQDFFPKAKPNGIFKGSTAENAEYLRVAPQSSEFIGA	
EGFR encoding DNA sequence (from Genbank NM_005228.4) Bold and italicized: siRNA binding regions	ATGCGACCCCTCCGGGACGGCCGGGCGAGCGCTCCTGGCGCTGCTG GCTGCGCTCTGCCCCGGCAGTCCGGGCTCTGGAGGAAAAGAAAAGTT TGCCAAGGCACGAGTAACAAGCTCACGCAGTTGGGCACCTTTTGAA GATCATTTTCTCAGCCTCCAGAGGATGTTCAATAACTGTGAGGTG GTCCTTGGGAATTTGGAAATTACCTATGTGCAGAGGAATTATGAT CTTTCTTCTTAAAGACCATCCAGGAGGTGGCTGGTTATGTCTC ATTGCCCTCAACACAGTGGAGCGAATTCCTTTGGAAAACCTGCAG ATCATCAGAGGAAATATGTACTACGAAAATTCCTATGCCTTAGCA GTCTTATCTAATACTATGATGCAAATAAAACCGGACT <i>GAAGGAGCTG</i> <i>CCCATGAGAAA</i> TTTACAGGAAATCCTGCATGGCGCCGTGCGGTTC AGCAACAACCCCTGCCCTGTGCAACGTGGAGAGCATCCAGTGGCGG GACATAGTCAGCAGTGACTTTCTCAGCAACATGTGCATGGACTTC CAGAACCACCTGGGCAGCTGCCAAAAGTGTGATCCAAGCTGTCCC AATGGGAGCTGCTGGGGTGCAGGAGAGGAGAACTGCCAGAACTG ACCAAAATCATCTGTGCCCAGCAGTGTCCGGGCGCTGCCGTGGC AAGTCCCCCAGTGACTGCTGCCACAACAGTGTGCTGCAGGCTGC ACAGGCCCCCGGAGAGCGACTGCCTGGTCTGCCGCAAATTCGGA GACGAAGCCACGTGCAAGGACACCTGCCCCCACTCATGCTCTAC AACCCACACAGTACCAGATGGATGTGAACCCCGAGGGCAAATAC AGCTTTGGTGCCACCTGCGTGAAGAAGTGTCCCCGTAATTATGTG GTGACAGATCACGGCTCGTGCGTCCGAGCCTGTGGGGCCGACAGC TATGAGATGGAGGAAGACGGCGTCCGCAAGTGTGAAGAAGTGCAGAA GGGCCTTGCCGCAAAGTGTGTAAACGGAATAGGTATTGGTGAATTT AAAGACTCACTCTCCATAAATGCTACGAATATTAAACACTTCAAA AACTGCACCTCCATCAGTGGCGATCTCCACATCCTGCCGTTGGCA TTTAGGGGTGACTCCTTCACACATACTCCTCCTCTGGATCCACAG GAACTGGATATTCTGAAAACCGTAAAGGAAATCACAGGGTTTTTG CTGATTCAAGGCTTGGCCTGAAAACAGGACGGACCTCCATGCCTTT GAGAACCTAGAAATCATACGCGGCAGGACCAAGCAACATGGTCAG TTTTCTCTTGCAAGTGTGCAACATAACATCCTTGGGATTA CGCTCCCTCAAGGAGATAAGTGTGAGATGTGATAATTTAGGA AACAAAAATTTGTGCTATGCAAATACAATAAACTGGAAAAAACTG TTTGGGACCTCCGGTCAGAAAACCAAATATAAGCAACAGAGGT GAAAACAGCTGCAAGGCCACAGGCCAGGTCTGCCATGCCTTGTGC TCCCCGAGGGCTGTGGGGCCCGAGCCAGGGACTGCGTCTCT TGCCGGAATGTGAGCCGAGGCAGGGAATGCGTGGACAAGTGCAAC CTTCTGGAGGGTGAGCCAAGGGAGTTTGTGGAGAACTCTGAGTGC ATACAGTGCCACCCAGAGTGCCTGCCTCAGGCCATGAACATCACC TGCACAGGACGGGGACCAGACAACCTGTATCCAGTGTGCCCACTAC	59

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	ATTGACGGCCCCCACTGCGTCAAGACCTGCCCGGCAGGAGTCATG GGAGAAAACAACACCCTGGTCTGGAAGTACGCAGACGCCGGCCAT GTGTGCCACCTGTGCCATCCAACTGCACCTACGGATGCACTGGG CCAGGTCTTGAAGGCT GTCCAACGAATGGGCCTAAGAT CCCGTCC ATCGCCACTGGGATGGTGGGGGCCCTCCTCTTGCTGCTGGTGGTG GCCCTGGGGATCGGCCTCTTCATGCGAAGGCGCCACATCGTTCCG AAGCGCACGCTGCGGAGGCTGCTGCAGGAGAGGGAGCTTGTGGAG CCTCTTACACCCAGTGGAGAAGCTCCCAACCAAGCTCTCTTGAGG ATCTTGAAGGAACTGAATTCAAAAAGATCAAAGTGCTGGGCTCC GGTGCGTTTCGGCACGGTGTATAAGGGACTCTGGATCCCAGAAGGT GAGAAAGTTAAATTTCCCGTCGCTATCAAGGAATTAAGAGAAGCA ACATCTCCGAAAGCCAACAAGGAAATCCTCGATGAAGCCTACGTG ATGGCCAGCGTGGACAACCCCCACGTGTGCCGCTGCTGGGCATC TGCCTCACCTCCACCGTGCAGCTCATCACGCAGCTCATGCCCTTC GGCTGCCTCCTGGACTATGTCCGGGAACACAAAGACAATATTGGC TCCCAGTACCTGCTCAACTGGTGTGTGCAGATCGCAAAGGGCATG AACTACTTGGAGGACCGTCGCTTGGTGCACCGCGACCTGGCAGCC AGGAACGTACTGGTGAAAACACCGCAGCATGTCAAGATCAGAGAT TTTGGGCTGGCCAACTGCTGGGTGCGGAAGAGAAAAGAATACCAT GCAGAAGGAGGCAAAGTGCCTATCAAGTGGATGGCATTGGAATCA ATTTTACACAGAATCTATACCCACCAGAGTGATGTCTGGAGCTAC GGGGTGACCGTTTGGGAGTTGATGACCTTTGGATCCAAGCCATAT GACGGAATCCCTGCCAGCGAGATCTCCTCCATCCTGGAGAAAGGA GAACGCCTCCCTCAGCCACCCATATGTACCATCGATGTCTACATG ATCATGGTCAAGTGTCTGGATGATAGACGCAGATAGTCGCCCAAAG TTCCGTGAGTTGATCATCGAATTCTCCAAAATGGCCCCGAGACCCC CAGCGCTACCTTGTCAATTCAGGGGGATGAAAGAATGCATTTGCCA AGTCTACAGACTCCAACCTTCTACCGTGCCCTGATGGATGAAGAA GACATGGACGACGTGGTGGATGCCGACGAGTACCTCATCCCACAG CAGGGCTTCTTCAGCAGCCCCCTCCACGTACCGGACTCCCCCTCTG AGCTCTCTGAGTGCAACCAGCAACAATTCCACCGTGGCTTGCAAT GATAGAAATGGGCTGCAAAGCTGTCCCATCAAGGAAGACAGCTTC TTGCAGCGATACAGCTCAGACCCACAGGCGCCTTGACTGAG GGAC AGCATAGACGACACCTTC CCTCCAGTGCCTGAATACATAAACCAG TCCGTTCCAAAAGGCCCGCTGGCTCTGTGCAGAATCCTGTCTAT CACAAATCAGCCTCTGAACCCCGCGCCCAGCAGAGACCCACACTAC CAGGACCCCCACAGCACTGCAGTGGGCAACCCCGAGTATCTCAAC ACTGTCCAGCCACCTGTGTCAACAGCACATTTCGACAGCCCTGCC CACTGGGCCCAGAAAGGCAGCCACCAAATTAGCCTGGACAACCCCT GACTACCAGCAGGACTTCTTTCCCAAGGAAGCCAAGCCAAATGGC ATCTTTAAGGGCTCCACAGCTGAAAATGCAGAATACCTAAGGGTC GCGCCACAAAGCAGTGAATTTATTGGAGCATGA	
EGFR encoding RNA sequence (from Genbank NM_005228.4) Bold and italicized:	AUGCGACCCUCCGGGACGGCCGGGGCAGCGCUCCUGGCGCUGCUG GCUGCGCUCUGCCCGGCGAGUCGGGCUCUGGAGGAAAAGAAAGUU UGCCAAGGCACGAGUAACAAGCUCACGCAGUUGGGCACUUUUGAA GAUCAUUUUCUCAGCCUCCAGAGGAUGUUAUAACUGUGAGGUG GUCCUUGGGAAUUUGGAAAUUACCUAUGUGCAGAGGAAUUUAUGAU CUUUCUUCUUAAGACCAUCCAGGAGGUGGCUUGGUUAUGUCCUC AUUGCCCUCAACACAGUGGAGCGAAUCCUUUGGAAAACCUGCAG AUCAUCAGAGGAAAUUGUACUACGAAAAUCCUAUGCCUUAGCA	60

Protein or Nucleic Acid	Sequence	SEQ ID NO:
siRNA binding regions	<p>GUCUUUAUCUAACUAUGAUGCAAAUAAAACCGGACUGAAGGAGCUG CCCAUGAGAAAUUUACAGGAAAUCCUGCAUGGCGCCGUGCGGUUC AGCAACAACCCUGCCCUGUGCAACGUGGAGAGCAUCCAGUGGCGG GACAUAGUCAGCAGUGACUUUCUCAGCAACAUGUCGAUGGACUUC CAGAACACCUGGGCAGCUGCCAAAAGUGUGAUCCAAGCUGUCCC AAUGGGAGCUGCUGGGGUGCAGGAGAGGAGAACUGCCAGAAACUG ACCAAAAUCAUCUGUGCCCAGCAGUGCUCCGGGCGCUGCCGUGGC AAGUCCCCCAGUGACUGCUGCCACAACCAGUGUGCUGCAGGCUGC ACAGGCCCCCGGAGAGCGACUGCCUGGUCUGCCGCAAAUCCGA GACGAAGCCACGUGCAAGGACACCUGCCCCCACUCAUGCUCUAC AACCCACCACGUACCAGAUGGAUGUGAACCCCGAGGGCAAAUAC AGCUUUGGUGCCACCUGCGUGAAGAAGUGUCCCCGUAAUUAUGUG GUGACAGAUACGGCUCGUGCGUCCGAGCCUGUGGGGCCGACAGC UAUGAGAUGGAGGAAGACGGCGUCCGCAAGUGUAAGAAGUGCGAA GGGCCUUGCCGCAAAGUGUGUAACGGAAUAGGUUUGGUGAAUUU AAAGACUCACUCUCCAUAUAAUGCUACGAAUUAUAAACACUUCAAA AACUGCACCUCACUAGUGGCGAUCUCCACAUCUCCUGCCGUGGCA UUUAGGGGUGACUCCUUCACACAUACUCCUCCUCUGGAUCCACAG GAACUGGAUAUUCUGAAAACCGUAAAGGAAAUCACAGGGUUUUUG CUGAUUCAGGCUUGGCCUGAAAACAGGACGGACCUCUAGCCUUU GAGAACCUGAAAUCUACGCGGCAGGACCAAGCAACAUGGUCAG UUUUCUCUUGCAGUCGUCAGCCUGAACAUACAUCUUGGGAUUA CGCUCUCCUCAAGGAGAUAGUGAUGGAGAUUGAUAAUUCAGGA AACAAAAUUGUGCUAUGCAAUAACAUAACUGGAAAAAACUG UUUGGGACCUCGCGUCAGAAAACCAAAUUAUAAGCAACAGAGGU GAAAACAGCUGCAAGGCCACAGGCCAGGUCUGCCAUGCCUUGUGC UCCCCCGAGGGCUGCUGGGGCCCGGAGCCAGGGACUGCGUCUCU UGCCGGAUUGUCAGCCGAGGCAGGGAUUGCGUGGACAAGUGCAAC CUUCUGGAGGGUGAGCCAAGGGAGUUUGUGGAGAACUCUGAGUGC AUACAGUGCCACCCAGAGUGCCUGCCUCAGGCCAUGAACAUCACC UGCACAGGACGGGGACCAGACAACUGUAUCCAGUGUGCCACUAC AUUGACGGCCCCCACUGCGUCAAGACCUGCCCGCAGGAGUCAUG GGAGAAAACAACACCCUGGUCUGGAAGUACGCAGACGCCGGCCAU GUGUGCCACCUGGCCAUCCAAACUGCACCUCGGAUGCAGUGGG CCAGGUCUUGAAGGCUGUCCAACGAAUGGGCCUAAGAUCCCGUCC AUCGCCACUGGGAUGGUGGGGGCCUCCUCUUGCUGCUGGUGGUG GCCCUGGGGAUCGGCCUCUUAUGCGAAGGCGCCACAUCGUUCGG AAGCGCACGCUGCGGAGGCUGCUGCAGGAGAGGGAGCUUGUGGAG CCUCUUACACCCAGUGGAGAAGCUCCCAACCAAGCUCUCUUGAGG AUCUUGAAGGAAACUGAAUUCAAAAAGAUCAAAGUGCUGGGCUCC GGUGCGUUCGGCACGGUGUAUAAGGGACUCUGGAUCCAGAAGGU GAGAAAGUAAAAUCCCCGUCGUAUCAAGGAAUUAAGAGAAGCA ACAUCUCCGAAAGCCAACAAGGAAAUCCUCGAUGAAGCCUACGUG AUGGCCAGCGUGGACAACCCCCACGUGUGCCGCCUGCUGGGCAUC UGCCUACCUCCACCGUGCAGCUAUCACGCAGCUAUGCCCUUC GGCUGCCUCCUGGACUAUGUCCGGGAACACAAAGACAAUAUUGGC UCCAGUACCUGCUAACUGGUGUGUGCAGAUCCGAAAGGGCAUG AACUACUUGGAGGACCGUCGUUGGUGCACCAGCGACCUGGCAGCC AGGAACGUACUGGUGAAAACACCGCAGCAUGUCAAGAUCACAGAU UUUGGGCUGGCCAAACUGCGGGUGCGGAAGAGAAAGAAUACCAU</p>	

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	GCAGAAGGAGGCAAAGUGCCUAUCAAGUGGAUGGCAUUGGAAUCA AUUUUACACAGAAUCUAUACCCACCAGAGUGAUGUCUGGAGCUAC GGGGUGACCGUUUGGGAGUUGAUGACCUUUGGAUCCAAGCCAUAU GACGGAAUCCCUGCCAGCGAGAUCCUCCAUCUGGAGAAAGGA GAACGCCUCCCUCAGCCACCCAUUGUACCAUCGAUGUCUACAUG AUCAUGGUCAAGUGCUGGAUGAUAGACGCAGAUAGUCGCCCAAAG UUCGUGAGUUGAUCAUCGAAUUCUCCAAAAUGGCCCCGAGACCCC CAGCGCUACCUUGUCAUUCAGGGGGAUGAAAGAAUGCAUUGCCA AGUCCUACAGACUCCAACUUCUACCGUGCCCUGAUGGAUGAAGAA GACAUGGACGACGUGGUGGAUGCCGACGAGUACCUCAUCCACAG CAGGGCUUCUUCAGCAGCCCCUCCACGUCACGGACUCCCCUCCUG AGCUCUCUGAGUGCAACCAGCAACAAUCCACCGUGGCUUGCAU GAUAGAAAUGGGCUGCAAAGCUGUCCCAUCAAGGAAGACAGCUUC UUGCAGCGAUACAGCUCAGACCCACAGGCGCCUUGACUGA GGAC AGCAUAGACGACACCUUC CUCCCAGUGCCUGAAUACAUAACCAG UCCGUUCCCCAAAAGGCCCGCUGGCUCUGUGCAGAAUCCUGUCU CACAUCAGCCUCUGAACCCCGCGCCCAGCAGAGACCCACACUAC CAGGACCCCCACAGCAGCUGCAGUGGGCAACCCCGAGUAUCUAC ACUGUCCAGCCCACCUGUGUCAACAGCACAUUCGACAGCCCUGCC CACUGGGCCCAGAAAGGCAGCCACCAAAUAGCCUGGACAACCCU GACUACCAGCAGGACUUCUUCUCCAAAGGAAGCCAAGCCAAAUGGC AUCUUUAAGGGCUCCACAGCUGAAAAUGCAGAAUACCUAAGGGUC GCGCCACAAAGCAGUGAAUUUAUUGGAGCAUGA	
mTOR amino acid (Genbank NM_005931.4)	MLGTGPAAATTAATTSSNVSVLQQFASGLKSRNEETRAKAAKELQ HYVTMELREMSQEESTRFYDQLNHHIFELVSSSDANERKGGILAI ASLIGVEGGNATRIGRFANYLRNLLPSNDPVVMMEMASKAIGRLAM AGDTFTAIEYVEFEVKRALEWLGADRNEGRRHA AVLVLRELAISVP TFFFQQVQPFEDNIFVAVWDPKQAIREGAVAALRACLIILTTQREP KEMQKPQWYRHTFEEAEKGFDETLAKEKGMNRDDRIGHALLILNE LVRISSMEGERLREEMEEITQQQLVHDKYCKDLMGFGTKPRHITP FTSFQAVQPPQSNALVGLLGYSSSHQGLMGFGTSPSPAKSTLVESR CCRDLMEEKFDQVCQWVLKCRNSKNSLIQMTILNLLPRLAAFRPS AFTDTQYLQDTMNHVLSVCVKKEKERTAAFQALGLLSVAVRSEFKV YLPRVLDIIRAALPPKDFAHKRQKAMQVDATVFTCISMLARAMGP GIQQDIKELLEPMLAVGLSPALTAVLYDLRSRQIPQLKKDIQDGLL KMLSLVLMHKPLRHPGMPKGLAHQLASPLTTLPEASDVGSITLA LRTLGSFEFEHSLTQFVRHCADHFLNSEHKEIRMEAARTCSRLL TPSIHLISGHAHVVSQTAVQVVADVLSKLLVVGITDPDPDIRYCV LASLDERFDAHLAQAE NLQALFVALNDQVFEIRELAICTVGR LSS MNP AFVMPFLRKMLIQILTELEHSGIGRIKEQSARMLGHLVSNAP RLIRPYMEPILKALILKLKDPDPDPNPGVINNVLATIGELAQVSG LEM RKWVDELFIIMDMLQDSSLLAKRQVALWTLGQLVASTGYVV EPYRKYP TLLLEVLLNFLKTEQNQGTRREAIRVLG LLGALDPYKHK VNIGMIDQSRDASAVSLSESKSSQDSSDYSTSEMLVNMGNLPLDE FYPAVSMVALMRIFRDQSLSHHHTMVVQAITFIFKSLGLKCVQFL PQVMPTFLNVIRVCDGAIREFLFQQLGMLVSFVKSHIRPYMDEIV TLMREFWVMNTSIQSTIILLIEQIVVALGGEFKLYLPQLIPHMLR VFMHDNSPGRIVSIKLLAAIQLFGANLDDYLHLLLPPIVKLFDAP EAPLPSRKAAL ETVDRLTESLDFTDYASRIIHPIVRTL DQSP ELR STAMDTLSSLVFQLGKKYQIFIPMVNKLVRHRINHQRYDVLICR	61

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	IVKGYTLADEEEDPLIYQHRMLRSGQGDALASGPVETGPMKKLHV STINLQKAWGAARRVSKDDWLEWLRRLSLELLKDSSSPSLRSCWA LAQAYNPMARDLFNAAFVSCWSELNEDQQDELIRSIELALTSQDI AEVTQTLLNLAEFMEHSDKGPLPLRDDNGIVLLGERAAKCRAYAK ALHYKELEFQKGPTPAILESLSINNKLQQPEAAAGVLEYAMKHF GELEIQATWYEKLHEWEDALVAYDKKMDTNKDDPELMLGRMRCLE ALGEWGQLHQCCCKWTLVNDETQAKMARMAAAAAWGLGQWDSME EYTCMIPRDTHDGAIFYRAVLALHQDLFSLAQQCIDKARDLLDAEL TAMAGESYSRAYGAMV SCHMLSELEEVIQYKLVPERREIIRQIWW ERLQGCQRIVEDWQKILMVRSLVSPHEDMRTWLKYASLCGKSGR LALAHKTLVLLLGVDP SRQLDHP LPTVHPQV TYAYMKNMWKSARK IDAFQHMQH FVQTMQQQAQHA IATEDQQHKQELHKL MARCFLKLG EWQLNLQGINESTIPKVLQYYSAA TEHDRSWYKAWHAWAVMNFEA VLHYKHQNQARDEKKKL RHASGANITNATTAATTAATATTTASTE GSNSESEAESTENSPTSP LQKKVTEDLSKTL LMYTVP AVQGF FR SISLSRGNNLQD TLRVLT LWF DYGHWPDVNEALVEGVKAIQIDTW LQVIPQLIARIDTPRPLVGR LIHQLLTDIGRYHPQAL IYPLTVAS KSTTTARHNAANKILKNMCEHSNTLVQQAMMVSEELIRVAILWHE MWHEGLEEASRLYFGERNVKGMFEVLEPLHAMMERGPQTLKETS F NQAYGRDLMEAQEWCRKYMKSGNVKDLTQAWDLYYHVFR RISKQL PQLTSLELQYVSPKLLMCRDLELAVPGTYDPNQPIIRIQSIAPSL QVITSKQRPRKLTLMG SNGHEFVFL LKGHEDLRQDERVMQLFGLV NTLLANDPTSLRKNLSIQRYAVIPLSTNSGLIGWVPHCDTLHALI RDYREKKKILLNIEHRIMLRMAPDYDHLTLMQKVEVFEHAVNNTA GDDLAKLLWLKSPSSEVWFDRRTNYTRSLAVMSMVGYILGLDRH PSNLMLDRLSGKILHIDFGDCFEVAMTREKFPEKIPFRLTRMLTN AMEVTGLDGNRYRITCHTVMEVLR EHKDSVM AVLEAFVYDPLL NWR LMDTNTKGNKRSRTRTDSYSAGQSVEILDGV ELGEP AHKKTGTTV PESIHSFIGDGLVKPEALNKKAIQI INVRDKLTGRDFSHDDTLD VPTQVELLIKQATSHENLCQCYIGWCPFW	
mTOR encoding DNA sequence (from Genbank NM_005931.4) Bold and italicized: siRNA binding regions	ATGCTTGGAACCGGACCTGCCGCCGCCACCGCTGCCACCACA TCTAGCAATGTGAGCGTCTCTGCAGCAGTTTGCCAGTGCCCTAAAG AGCCGGAATGAGGAAACCAGGGCCAAAGCCGCCAAGGAGCTCCAG CACTATGTCACCATGGAACCTCCGAGAGATGAGTCAAGAGGAGTCT ACTCGCTTCTATGACCAACTGAACCATCACATTTTGAATTGGTT TCCAGCTCAGATGCCAATGAGAGGAAAGGTGGCATCTTGCCATA GCTAGCCTCATAGGAGTGGAAGGTGGGAATGCCACCCGAATTGGC AGATTTGCCAACTATCTTCGGAACCTCCTCCCCCTCCAATGACCCA GTTGTCATGGAAATGGCATCCAAGGCCATTGGCCGTCTTGCCATG GCAGGGGACACTTTTACCGCTGAGTACGTGGAATTTGAGGTGAAG CGAGCCCTGGAATGGCTGGGTGCTGACCGCAATGAGGGCCGGAGA CATGCAGCTGTCCTGGTTCTCCGTGAGCTGGCCATCAGCGTCCCT ACCTTCTTCTTCCAGCAAGTGCAACCTTCTTTGACAACATTTT GTGGCCGTGTGGGACCCAAACAGGCCATCCGTGAGGGAGCTGTA GCCGCCCTTCGTGCCTGTCTGATTCTCACAACCCAGCGTGAGCCG AAGGAGATGCAGAAGCCTCAGTGGTACAGGCACACATTTGAAGAA GCAGAGAAGGGATTTGATGAGACCTTGGCCAAAGAGAAGGGCATG AATCGGGATGATCGGATCCATGGAGCCTTGTTGATCCTTAACGAG CTGGTCCGAATCAGCAGCATGGAGGGAGAGCGTCTGAGAGAAGAA ATGGAAGAAATCACACAGCAGCAGCTGGTACACGACAAGTACTGC	62

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	AAAGATCTCATGGGCTTCGGAACAAAACCTCGTCACATTACCCCC TTCACCAGTTTCCAGGCTGTACAGCCCCAGCAGTCAAATGCCTTG GTGGGGCTGCTGGGGTACAGCTCTCACCAGGCCTCATGGGATTT GGGACCTCCCCCAGTCCAGCTAAGTCCACCCTGGTGGAGAGCCGG TGTTGCAGAGACTTGATGGAGGAGAAATTTGATCAGGTGTGCCAG TGGGTGCTGAAATGCAGGAATAGCAAGAACTCGCTGATCCAAATG ACAATCCTTAATTTGTTGCCCCGCTTGGCTGCATTCCGACCTTCT GCCTTCACAGATACCCAGTATCTCCAAGATACCATGAACCATGTC CTAAGCTGTGTCAAGAAGGAGAAGGAACGTACAGCGGCCCTTCCAA GCCCTGGGGCTACTTTCTGTGGCTGTGAGGCTGTGAGTTTAAAGTC TATTTGCCTCGCGTGCTGGACATCATCCGAGCGGCCCTGCCCCCA AAGGACTTCGCCCATAAGAGGCAGAAGGCAATGCAGGTGGATGCC ACAGTCTTCACTTGCATCAGCATGCTGGCTCGAGCAATGGGGCCA GGCATCCAGCAGGATATCAAGGAGCTGCTGGAGCCCATGCTGGCA GTGGGACTAAGCCCTGCCCTCACTGCAGTGTCTACGACCTGAGC CGTCAGATTCCACAGCTAAAGAAGGACATTCAAGATGGGCTACTG AAAATGCTGTCCCTGGTCTTATGCACAAACCCCTTCGCCACCCA GGCATGCCCAAGGGCCTGGCCCATCAGCTGGCCCTCTCCTGGCCCTC ACGACCCTCCCTGAGGCCAGCGATGTGGGCAGCATCACTCTTGCC CTCCGAACGCTTGGCAGCTTTGAATTTGAAGGCCACTCTCTGACC CAATTTGTTGCGCACTGTGCGGATCATTTCTGAACAGTGAGCAC AAGGAGATCCGCATGGAGGCTGCCCCGACCTGCTCCCGCCTGCTC ACACCCTCCATCCACCTCATCAGTGGCCATGCTCATGTGGTTAGC CAGACCGCAGTGCAAGTGGTGGCAGATGTGCTTAGCAAACCTGCTC GTAGTTGGGATAACAGATCCT GACCCTGACATTGCTACTGT CTC TTGGCGTCCCTGGACGAGCGCTTTGATGCACACCTGGCCAGGCG GAGAACITGCAGGCCTTGTTTGTGGCTCTGAATGACCAGGTGTTT GAGATCCGGGAGCTGGCCATCTGCACTGTGGGCCGACTCAGTAGC ATGAACCCTGCCTTTGTTCATGCCTTTCTGCGCAAGATGCTCATC CAGATTTTGACAGAGTTGGAGCACAGTGGGATTGGAAGAATCAAA GAGCAGAGTGGCCGCATGCTGGGGCACCTGGTCTCCAATGCCCCC CGACTCATCCGCCCCCTACATGGAGCCTATTCTGAAGGCATTAATT TTGAAACTGAAAGATCCAGACCCTGATCCAAACCCAGGTGTGATC AATAATGTCTGGCAACAATAGGAGAATTGGCACAGGTTAGTGGC CTGGAAATGAGGAAATGGGTTGATGAACTTTTTATTATCATCATG GACATGCTCCAGGATTCTCTTTGTTGGCCAAAAGGCAGGTGGCT CTGTGGACCCTGGGACAGTTGGTGGCCAGCACTGGCTATGTAGTA GAGCCCTACAGGAAGTACCCTACTTTGCTTGAGGTGCTACTGAAT TTTCTGAAGACTGAGCAGAACCAGGGTACACGCAGAGAGGCCATC CGTGTGTTAGGGCTTTTAGGGGCTTTGGATCCTTACAAGCACAAA GTGAACATTGGCATGATAGACCAGTCCCGGGATGCCTCTGCTGTC AGCCTGTGAGAATCCAAGTCAAGTCAGGATTCTCTGACTATAGC ACTAGTGAAATGCTGGTCAACATGGGAACTTGCTCTGGATGAG TTCTACCCAGCTGTGTCCATGGTGGCCCTGATGCGGATCTTCCGA GACCAGTCACTCTCTCATCATCACACCATGGTTGTCCAGGCCATC ACCTTCATCTTCAAGTCCCTGGGACTCAAATGTGTGCAGTCTCTG CCCCAGGTCAATGCCACGTTTCCTTAACGTCATTTCGAGTCTGTGAT GGGGCCATCCGGGAATTTTTGTTCCAGCAGCTGGGAATGTTGGTG TCCTTTGTGAAGAGCCACATCAGACCTTATATGGATGAAATAGTC ACCCTCATGAGAGAATCTGGGTCTGAACACCTCAATTCAGAGC	

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	ACGATCATTCCTTCTCATTGAGCAAATTGTGGTAGCTCTTGGGGGT GAATTTAAGCTCTACCTGCCCCAGCTGATCCACACATGCTGCGT GTCTTCATGCATGACAACAGCCCAGGCCGATTGTCTCTATCAAG TTACTGGCTGCAATCCAGCTGTTTGGCGCCAACCTGGATGACTAC CTGCATTTACTGCTGCCTCCTATTGTAAAGTTGTTTGATGCCCCCT GAAGCTCCACTGCCATCTCGAAAGGCAGCGCTAGAGACTGTGGAC CGCCTGACGGAGTCCCTGGATTTCAGTACTATGCCTCCCGGATC ATTCACCCTATTGTTTCGAACACTGGACCAGAGCCCAGAAGTGCAGC TCCACAGCCATGGACACGCTGTCTTCACTTGTTTTTTCAGCTGGGG AAGAAGTACCAAATTTTTCATTCCAATGGTGAATAAAGTTCTGGTG CGACACCGAATCAATCATCAGCGCTATGATGTGCTCATCTGCAGA ATTGTCAAGGGATACACACTTGCTGATGAAGAGGAGGATCCTTTG ATTTACCAGCATCGGATGCTTAGGAGTGGCCAAGGGGATGCATTG GCTAGTGGACCAAGTGGAAACAGGACCCATGAAGAACTGCACGTC AGCACCATCAACCTCCAAAAGGCCTGGGGCGCTGCCAGGAGGGTC TCCAAAGATGACTGGCTGGAATGGCTGAGACGGCTGAGCCTG GAG CTGCTGAAGGACTCATCAT CGCCCTCCCTGCGCTCCTGCTGGGCC CTGGCACAGGCCTACAACCCGATGGCCAGGGATCTCTTCAATGCT GCATTTGTGTCTGCTGGTCTGAACTGAATGAAGATCAACAGGAT GAGCTCATCAGAAGCATCGAGTTGGCCCTCACCTCACAAGACATC GCTGAAGTCACACAGACCCTCTTAACTTGGCTGAATTCATGGAA CACAGTGACAAGGGCCCCCTGCCACTGAGAGATGACAATGGCATT GTTCTGCTGGGTGAGAGAGCTGCCAAGTGCCGAGCATATGCCAAA GCACTACACTACAAAGAACTGGAGTTCCAGAAAGGCCCCACCCCT GCCATTCTAGAATCTCTCATCAGCATTAATAATAAGCTACAGCAG CCGGAGGCAGCGGCCGGAGTGTTAGAATATGCCATGAAACACTTT GGAGAGCTGGAGATCCAGGCTACCTGGTATGAGAACTGCACGAG TGGGAGGATGCCCTTGTGGCCTATGACAAGAAAATGGACACCAAC AAGGACGACCCAGAGCTGATGCTGGGCCGCTGCGCTGCCTCGAG GCCTTGGGGGAATGGGGTCAACTCCACCAGCAGTGCTGTGAAAAG TGGACCCTGGTTAATGATGAGACCCAAGCCAAGATGGCCCGGATG GCTGCTGCAGCTGCATGGGGTTTAGGTGAGTGGGACAGCATGGAA GAATACACCTGTATGATCCCTCGGGACACCCATGATGGGGCATT TATAGAGCTGTGCTGGCACTGCATCAGGACCTCTTCTCCTTGGCA CAACAGTGCATTGACAAGGCCAGGGACCTGCTGGATGCTGAATTA ACTGCGATGGCAGGAGAGAGTTACAGTCGGGCATATGGGGCCATG GTTTCTTGCCACATGCTGTCCGAGCTGGAGGAGGTTATCCAGTAC AACTTGTCCCCGAGCGACGAGAGATCATCCGCCAGATCTGGTGG GAGAGACTGCAGGGCTGCCAGCGTATCGTAGAGGACTGGCAGAAA ATCCTTATGGTGCGGTCCCTTGTGGTCAGCCCTCATGAAGACATG AGAACCTGGCTCAAGTATGCAAGCCTGTGCGGCAAGAGTGGCAGG CTGGCTCTTGCTCATAAACTTTAGTGTGCTCCTGGGAGTTGAT CCGTCTCGGCAACTTGACCATCCTCTGCCAACAGTTCACCCCTCAG GTGACCTATGCCTACATGAAAAACATGTGGAAGAGTGCCCGCAAG ATCGATGCCTTCCAGCACATGCAGCATTTTGTCCAGACCATGCAG CAACAGGCCCCAGCATGCCATCGCTACTGAGGACCAGCAGCATAAG CAGGAACTGCACAAGCTCATGGCCCCGATGCTTCCTGAAACTTGGA GAGTGGCAGCTGAATCTACAGGGCATCAATGAGAGCACAATCCCC AAAGTGTGTCAGTACTACAGCGCCGCCACAGAGCACGACCGCAGC TGGTACAAGGCCTGGCATGCGTGGGCAGTGATGAACTTCGAAGCT	

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	<p>GTGCTACACTACAAACATCAGAACCAAGCCCGCGATGAGAAGAAG AAACTGCGTCATGCCAGCGGGGCCAACATCACCAACGCCACCACT GCCGCCACCACGGCCGCCACTGCCACCACCACTGCCAGCACCGAG GGCAGCAACAGTGAGAGCGAGGCCGAGAGCACCGAGAACAGCCCC ACCCCATCGCCGCTGCAGAAGAAGGTCCTGAGGATCTGTCCAAA ACCCCTCCTGATGTACACGGTGCCTGCCGTCCAGGGCTTCTTCCGT TCCATCTCCTTGTACGAGGCAACAACCTCCAGGATACACTCAGA GTTCTCACCTTATGGTTTGATTATGGTCACTGGCCAGATGTCAAT GAGGCCITAGTGGAGGGGGTGAAAGCCATCCAGATTGATACCTGG CTACAGGTATATACCTCAGCTCATTGCAAGAATTGATACGCCCAGA CCCTTGGTGGGACGTCTCATTACACAGCTTCTCACAGACATTGGT CGGTACCACCCCCAGGCCCTCATCTACCCACTGACAGTGGCTTCT AAGTCTACCACGACAGCCCCGGCACAATGCAGCCAACAAGATTCTG AAGAACATGTGTGAGCACAGCAACACCCCTGGTCCAGCAGGCCATG ATGGTGAGCGAGGAGCTGATCCGAGTGGCCATCCTCTGGCATGAG ATGTGGCATGAAGGCCTGGAAGAGGCATCTCGTTTGTACTTTGGG GAAAGGAACGTGAAAGGCATGTTTGAGGTGCTGGAGCCCTTGCAAT GCTATGATGGAACGGGGCCCCCAGACTCTGAAGGAAACATCCTTT AATCAGGCCTATGGTTCGAGATTAAATGGAGGCCCAAGAGTGGTGC AGGAAGTACATGAAATCAGGGAATGTCAAGGACCTCACCCAAGCC TGGGACCTCTATTATCATGTGTTCCGACGAATCTCAAAGCAGCTG CCTCAGCTCACATCCTTAGAGCTGCAATATGTTTCCCCAAAACCTT CTGATGTGCCGGGACCTTGAATTGGCTGTGCCAGGAACATATGAC CCCAACCAGCCAATCATTTCGCATTTCAGTCCATAGCACCGTCTTTG CAAGTCATCACATCCAAGCAGAGGCCCGGAAATTGACACTTATG GGCAGCAACGGACATGAGTTTGTCTTCTTCTAAAAGGCCATGAA GAICTGCGCCAGGATGAGCGTGTGATGCAGCTCTTCGGCCTGGTT AACACCCTTCTGGCCAATGACCCAACATCTCTTCGGAAAAACCTC AGCATCCAGAGATACGCTGTATCCCTTTATCGACCAACTCGGGC CTCATTGGCTGGGTTCCTCCACTGTGACACACTGCACGCCCTCATC CGGGACTACAGGGAGAAGAAGAAGATCCTTCTCAACATCGAGCAT CGCATCATGTTGCGGATGGCTCCGGACTATGACCACTTGACTCTG ATGCAGAAGGTGGAGGTGTTTGAGCATGCCGTCAATAATACAGCT GGGGACGACCTGGCCAAGCTGCTGTGGCTGAAAAGCCCCAGCTCC GAGGTGTGGTTTGACCGAAGAACCAATTATACCCGTTCTTTAGCG GTCAITGCAATGGTTGGGTATATTTTAGGCCTGGGAGATAGACAC CCATCCAACCTGATGCTGGACCGTCTGAGTGGGAAGATCCTGCAC ATTGACTTTGGGGACTGCTTTGAGGTGCTATGACCCGAGAGAAG TTTCCAGAGAAGATTCCATTTAGACTAACAAGAATGTTGACCAAT GCTATGGAGGTTACAGGCCTGGATGGCAACTACAGAATCACATGC CACACAGTGATGGAGGTGCTGCGAGAGCACAAAGGACAGTGTCATG GCCGTGCTGGAAGCCTTTGTCTATGACCCCTTGCTGAACTGGAGG CTGATGGACACAAATACCAAAGGCAACAAGCGATCCCGAACGAGG ACGGATTCTACTCTGCTGGCCAGTCAGTCGAAATTTTGGACGGT GTGGAACCTGGAGAGCCAGCCATAAGAAAACGGGGACCACAGTG CCAGAATCTATTCTTTTCAATTGGAGACGGTTTGGTGAAACCA GAGGCCCTAAATAAGAAAGCTATCCAGATTATTAACAGGGTTTCGA GATAAGCTCACTGGTGGGACTTCTCTCATGATGACACTTTGGAT GTTCCAACGCAAGTTGAGCTGCTCATCAAACAAGCGACATCCCAT GAAAACCTCTGCCAGTGCTATATTGGCTGGTGCCCTTTCTGGTAA</p>	

Protein or Nucleic Acid	Sequence	SEQ ID NO:
mTOR encoding RNA sequence (from Genbank NM_005931.4) Bold and italicized: siRNA binding regions	AUGCUUGGAACCGGACCUGCCGCCGCCACCACCGCUGCCACCACA UCUAGCAAUGUGAGCGUCCUGCAGCAGUUUGCCAGUGGCCUAAAAG AGCCGGAAUGAGGAAACCAGGGCCAAAGCCGCCAAGGAGCUCCAG CACUAUGUCACCAUGGAACUCCGAGAGAUGAGUCAAGAGGAGUCU ACUCGCUUCUAUGACCAACUGAACCAUCACAUUUUUGAAUUGGUU UCCAGCUCAGAUGCCAAUGAGAGGAAAGGUGGCAUCUUGGCCAUA GCUAGCCUCAUAGGAGUGGAAGGUGGGAAUGCCACCCGAAUUGGC AGAUUUGCCAACUAUCUUCGGAACCUCCUCCCCUCCAAUGACCCA GUUGUCAUGGAAAUGGCAUCCAAGGCCAUUGGCCGUCUUGCCAUG GCAGGGGACACUUUUACCGCUGAGUACGUGGAAUUUGAGGUGAAG CGAGCCCUGGAAUGGCUGGGUGCUGACCGCAAUGAGGGCCGGAGA CAUGCAGCUGUCCUGGUUCUCCGUGAGCUGGCCAUCAGCGUCCCU ACCUUCUUCUCCAGCAAGUGCAACCCUUCUUUGACAACAUUUUU GUGGCCGUGUGGGACCCCAAACAGGCCAUCCGUGAGGGAGCUGUA GCCGCCUUCGUGCCUGUCUGAUUCUCACAACCCAGCGUGAGCCG AAGGAGAUGCAGAAGCCUCAGUGGUACAGGCACACAUUUGAAGAA GCAGAGAAGGGAAUUGAUGAGACCUUGGCCAAAGAGAAGGGCAUG AAUCGGGAUGAUCGGAUCCAUGGAGCCUUGUUGAUCCUUAACGAG CUGGUCCGAAUCAGCAGCAUGGAGGGAGAGCGUCUGAGAGAAGAA AUGGAAGAAUACACAGCAGCAGCUGGUACACGACAAGUACUGC AAAGAUCUCAUGGGCUUCGGAACAAAACCUUGCUCACAUUACCCCC UUCACCAGUUUCCAGGCUGUACAGCCCCAGCAGUCAAUGCCUUG GUGGGGUGCUGGGGUACAGCUCUCACCAAGGCCUCAUGGGAUUU GGGACCUCCCCAGUCCAGCUAAGUCCACCCUGGUGGAGAGCCGG UGUUGCAGAGACUUGAUGGAGGAGAAAUUGAUCAGGUGUGCCAG UGGGUGCUGAAAUGCAGGAUAGCAAGAACUCGCUGAUCCAAAUG ACAAUCCUUAUUUGUUGCCCCGCUUGGCUGCAUCCGACCUUCU GCCUUCACAGAUACCCAGUAUCUCCAAGAUACCAUGAACCAUGUC CUAAGCUGUGUCAAGAAGGAGAAGGAACGUACAGCGGCCUUCCAA GCCCUGGGGCUACUUUCUGUGGCUGUGAGGUCUGAGUUUAAGGUC UAUUUGCCUCGCGUGCUGGACAUCAUCCGAGCGGCCCUUGCCCCA AAGGACUUCGCCCAUAAGAGGCAGAAGGCAAUGCAGGUGGAUGCC ACAGUCUUCACUUGCAUCAGCAUGCUGGCUCGAGCAAUGGGGCCA GGCAUCCAGCAGGAUAUCAAGGAGCUGCUGGAGCCCAUGCUGGCA GUGGGACUAAGCCCUGCCCUCACUGCAGUCUCUACGACCUGAGC CGUCAGAUUCCACAGCUAAAGAAGGACAUUCAAGAUGGGCUACUG AAAAUGCUGUCCUGGUCCUUAUGCACAAACCCUUCGCCACCCA GGCAUGCCCAAGGGCCUGGCCCAUCAGCUGGCCUCUCCUGGCCUC ACGACCCUCCUGAGGCCAGCGAUGUGGGCAGCAUCACUCUUGCC CUCCGAACGCUUGGCAGCUUUGAAUUUGAAGGCCACUCUCUGACC CAAUUUGUUCGCCACUGUGCGGAUCAUUUCCUGAACAGUGAGCAC AAGGAGAUCCGCAUGGAGGCUGCCCGCACCUGCUCCCGCCUGCUC ACACCCUCCAUCCACCUCAUCAGUGGCCAUGCUCUAGUGGUUAGC CAGACCGCAGUGCAAGUGGUGGCAGAUGUGCUUAGCAAACUGCUC GUAGUUGGGAUAACAGAUCCU GACCCUGACAUUCGCUACUGU GUC UUGGCGUCCUGGACGAGCGCUUUGAUGCACACCUGGCCAGGCG GAGAACUUGCAGGCCUUGUUUGUGGCUCUGAAUGACCAGGUGUUU GAGAUCCGGGAGCUGGCCAUCUGCACUGUGGGCCGACUCAGUAGC AUGAACCCUGCCUUUGUCAUGCCUUUCUGCGCAAGAUGCUCUAC CAGAUUUUGACAGAGUUGGAGCACAGUGGGAUUGGAAGAAUCAA	63

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	GAGCAGAGUGCCCCGAUGCUGGGGCACCUGGUUCCAAUGCCCCC CGACUCAUCCGCCCCUACAUGGAGCCUAUUCUGAAGGCAUUAU UUGAAACUGAAAGAUCCAGACCCUGAUCCAAACCCAGGUGUGAUC AAUAAUGUCCUGGCAACAAUAGGAGAAUUGGCACAGGUUAGUGGC CUGGAAAUGAGGAAAUGGGUUGAUGAACUUUUUAUUAUCAU GACAUGCUCAGGAUUCUCUUGUUGGCCAAAAGGCAGGUGGCU CUGUGGACCCUGGGACAGUUGGUGGCCAGCACUGGCUAUGUAGUA GAGCCCUACAGGAAGUACCCUACUUGCUUGAGGUGCUACUGAAU UUUCUGAAGACUGAGCAGAACCAGGGUACACGCAGAGAGGCCAUC CGUGUGUUAGGGCUUUUAGGGGCUUUGGAUCCUUACAAGCACAAA GUGAACAUUGGCAUGAUAGACCAGUCCCGGAUGCCUCUGCUGUC AGCCUGUCAGAAUCCAAGUCAAGUCAGGAUUCUCUGACUAUAGC ACUAGUGAAAUGCUGGUCAACAUGGGAAACUUGCCUCUGGAUGAG UUCUACCCAGCUGUGUCCAUGGUGGGCCUGAUGCGGAUCUCCGA GACCAGUCACUCUCUCAUCAUCACACCAUGGUUUGUCCAGGCCAUC ACCUUCAUCUUAAGUCCUGGGACUCAAUGUGUGCAGUUCUCUG CCCCAGGUCAUGCCCACGUUCCUUAACGUCAUUCGAGUCUGUGAU GGGGCCAUCGGAUUAUUUUGUUCAGCAGCUGGGAAUGUUGUG UCCUUUGUGAAGAGCCACAUCAGACCUUAUAUGGAUGAAAUAGUC ACCCUCAUGAGAGAAUUCUGGGUCAUGAACACCUCAAUUCAGAGC ACGAUCAUUCUUCUCAUUGAGCAAUUGUGGUAGCUCUUGGGGGU GAAUUUAAGCUCUACCUGCCCCAGCUGAUCCACACAUGCUGCGU GUCUUAUGCAUGACAACAGCCCAGGCCGCAUUGUCUCUAUCAAG UUACUGGCUGCAAUCCAGCUGUUGGGCGCAACCUGGAUGACUAC CUGCAUUUACUGCUGCCUCCUAUUGCUAAGUUGUUGAUGCCCU GAAGCUCCACUGCCAUCUCGAAAGGCAGCGCUAGAGACUGUGGAC CGCCUGACGGAGUCCUGGAUUCACUGACUAUGCCUCCCGGAUC AUUCACCCUAUUGUUCGAACACUGGACCAGAGCCCAGAACUGCGC UCCACAGCCAUGGACACGCUGUCUUCACUUGUUUUCAGCUGGGG AAGAAGUACCAAUUUUCAUUCCAAUGGUGAAUAAAGUUCUGGUG CGACACCGAAUCAUUAUCAGCGCUAUGAUGUCUCAUCUGCAGA AUUGUCAAGGGAUACACACUUGCUGAUGAAGAGGAGGAUCCUUG AUUUACCAGCAUCGGAUGCUUAGGAGUGGCCAAGGGGAUGCAUUG GCUAGUGGACCAGUGGAAACAGGACCCAUGAAGAAACUGCACGUC AGCACCAUCAACCUCCAAAAGGCCUGGGGCGCUGCCAGGAGGUC UCCAAAGAUGACUGGCUGGAAUGGCUGAGACGGCUGAGCCUGGAG CUGCUGAAGGACUCAUCAUCGCCCUCUCCUGCGCUCUUGCUGGGCC CUGGCACAGGCCUACAACCCGAUGGCCAGGAUCUCUUCAAUGCU GCAUUUGUGUCCUGCUGGUCUGAACUGAAUGAAGAUCAACAGGAU GAGCUCAUCAGAAGCAUCGAGUUGGCCCUACCCUCACAAGACAUC GCUGAAGUCACACAGACCCUCUUAACUUGGCUGAAUUCAGGAA CACAGUGACAAGGGCCCCUGCCACUGAGAGAUGACAAUGGCAU GUUCUGCUGGGUGAGAGAGCUGCCAAGUGCCGAGCAUUGCCAAA GCACUACACUACAAGAACUGGAGUCCAGAAAGGCCCCACCCCU GCCAUUCUAGAAUCUCUCAUCAGCAUUAUAUAAGCUACAGCAG CCGGAGGCAGCGGCCGGAGUGUAGAAUAUGCCAUAGAAACACUU GGAGAGCUGGAGAUCCAGGCUACCUGGUUAUGAGAAACUGCACGAG UGGGAGGAUGCCCUUGUGGCCUAUGACAAGAAAAUGGACACCAAC AAGGACGACCCAGAGCUGAUGCUGGGCCGCAUGCGCUGCCUCGAG GCCUUGGGGGAUUGGGGUCAACUCCACCAGCAGUCUGUGAAAAAG	

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	UGGACCCUGGUUAAUGAUGAGACCCAAGCCAAGAUGGCCCGGAUG GCUGCUGCAGCUGCAUGGGGUUUAGGUCAGUGGGACAGCAUGGAA GAAUACACCUGUAUGAUCCUCGGGACACCCAUGAUGGGGCAUUU UAUAGAGCUGUGCUGGCACUGCAUCAGGACCUCUUCUCCUUGGCA CAACAGUGCAUUGACAAGGCCAGGGACCUGCUGGAUGCUGAAUUA ACUGCGAUGGCAGGAGAGAGUUACAGUCGGGCAUUGGGGCCAUG GUUUCUUGCCACAUGCUGUCCGAGCUGGAGGAGGUUAUCCAGUAC AAACUUGUCCCCGAGCGACGAGAGAUCAUCCGCCAGAUUGGUGG GAGAGACUGCAGGGCUGCCAGCGUAUCGUAGAGGACUGGCAGAAA AUCCUUAUGGUGCGGUCCCUUGUGGUCAGCCCUCAUGAAGACAUG AGAACCUGGCUCAAGUAUGCAAGCCUGUGCGGCAAGAGUGGCAGG CUGGCUCUUGCUCUAAAACUUUAGUGUUGCUCUUGGGAGUUGAU CCGUCUCGGCAACUUGACCAUCCUCUGCCAACAGUUCACCCUCAG GUGACCUAUGCCUACAUGAAAAACAUGUGGAAGAGUGCCCGCAAG AUCGAUGCCUCCAGCACAUGCAGCAUUUUGUCCAGACCAUGCAG CAACAGGCCCAGCAUGCCAUUCGUACUGAGGACCAGCAGCAUAAG CAGGAACUGCACAAGCUCUAGGCCCCGAUCUCCUGAAACUUGGA GAGUGGCAGCUGAAUCUACAGGGCAUCAUUGAGAGCACAUAUCCCC AAAGUGCUGCAGUACUACAGCGCCGCCACAGAGCACGACCGCAGC UGGUACAAGGCCUGGCAUGCGUGGGCAGUGAUGAACUUCGAAGCU GUGCUACACUACAAACAUCAGAACCAAGCCCGCGAUGAGAAGAAG AAACUGCGUCAUGCCAGCGGGGCCAACAUACCAACGCCACCACU GCCGCCACCACGGCCGCCACUGCCACCACCACUGCCAGCACCGAG GGCAGCAACAGUGAGAGCGAGGCGGAGAGCACCGAGAACAGCCCC ACCCAUCGCCCGCUGCAGAAGAAGCUCACUGAGGAUCUGUCCAAA ACCCUCCUGAUGUACACGGUGCCUGCCGUCCAGGGCUUCUCCGU UCCAUCUCCUUGUCACGAGGCAACAACCUCCAGGAUACACUCAGA GUUCUCACCUUAUGGUUUGAUUAUGGUCACUGGCCAGAUUGCAAU GAGGCCUUAUGUGGAGGGGGUGAAAGCCAUCCAGAUUGAUACCUUG CUACAGGUUAUACCUCAGCUCUUGCAAGAAUUGAUACGCCCGAGA CCCUUGGUGGGACGUCUCAUUCACCAGCUUCUCACAGACAUUGGU CGGUACCACCCCCAGGCCCUCAUCUACCCACUGACAGUGGCUCU AAGUCUACCACGACAGCCCCGGCACAAUGCAGCCAACAAGAUUCUG AAGAACAUGUGUGAGCACAGCAACACCCUGGUCCAGCAGGCCAUG AUGGUGAGCGAGGAGCUGAUCCGAGUGGCCAUCCUCUGGCAUGAG AUGUGGCAUGAAGGCCUGGAAGAGGCAUCUCGUUUGUACUUUGGG GAAAGGAACGUGAAAGGCAUGUUUGAGGUGCUGGAGCCCUUGCAU GCUAUGAUGGAACGGGGCCCCCAGACUCUGAAGGAAACAUCUUU AAUCAGGCCUAUGGUCGAGAUUUAAUGGAGGCCCAAGAGUGGUGC AGGAAGUACAUGAAAUACAGGGAUUGUCAAGGACCUCACCCAAGCC UGGGACCUCUAUUAUCAUGUGUCCGACGAAUCUCAAAGCAGCUG CCUCAGCUCACAUCUUAAGAGCUGCAUAUGUUUCCCCAAAACU CUGAUGUGCCGGGACCUUGAAUUGGCUGUGCCAGGAACAUAUGAC CCCAACCAGCCAUAUUCGCAUUCAGUCCAUAAGCACCUCUUUG CAAGUCAUCAUCCAAGCAGAGGCCCGGAAAUUGACACUUAUG GGCAGCAACGGACAUGAGUUUGUUUCCUUCUAAAAGGCCAUGAA GAUCUGCGCCAGGAUGAGCGUGUGAUGCAGCUCUUCGGCCUGGUU AACACCCUUCU GGCCAAUGACCCAACAUCUCU UCGGAAAAACUC AGCAUCCAGAGAUACGCUGUCAUCCCUUAUCGACCAACUCGGGC CUCAUUGGCUGGGUCCCCACUGUGACACACUGCACGCCCUCAUC	

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	CGGGACUACAGGGAGAAGAAGAAGAUCCUUCUACAACUAGAGCAU CGCAUCAUGUUGCGGAUGGCUCGCGACUAUGACCACUUGACUCUG AUGCAGAAGGUGGAGGUGUUUGAGCAUGCCGUCAAUAAUACAGCU GGGGACGACCUGGCCAAGCUGCUGUGGCUGAAAAGCCCCAGCUCC GAGGUGUGGUUUGACCGAAGAACCAGUUAUACCCGUUCUUUAGCG GUCAUGUCAUUGGUUGGGUAUUAUUUAGGCCUGGGAGAUAGACAC CCAUCCAACCUGAUGCUGGACCGUCUGAGUGGGAAGAUCUGCAC AUUGACUUUGGGGACUGCUUUGAGGUUGCUAUGACCCGAGAGAAG UUUCCAGAGAAGAUUCCAUAUUUAGACUAACAAGAAUGUUGACCAGU GCUAUGGAGGUUACAGGCCUGGAUGGCAACUACAGAAUCACAUGC CACACAGUGAUGGAGGUGCUGCGAGAGCACAAGGACAGUGUCAUG GCCGUGCUGGAAGCCUUUGUCUAUGACCCUUGCUGAACUGGAGG CUGAUGGACACAAUACCAAAGGCAACAAGCGAUCCCGAACGAGG ACGGAUCCUACUCUGCUGGCCAGUCAGUCGAAAUUUUGGACGGU GUGGAACUUGGAGAGCCAGCCCAUAAGAAAACGGGGACCACAGUG CCAGAAUCUAUUCUAUUCUUAUUGGAGACGGUUUGGUGAAACCA GAGGCCCUAAAUAAGAAAGCUAUCCAGAUUAUUAACAGGGUUCGA GAUAAGCUCACUGGUCGGGACUUCUCUCAUGAUGACACUUUGGAU GUUCCAACGCAAGUUGAGCUGCUCAUCAAACAAGCGACAUCCCAU GAAAACCUCUGCCAGUGCUAUAUUGGCUGGUGCCCUUUCUGGUAA	
KRAS amino acid (Genbank NM_004985.4) (Transcript variant b)	MTEYKLVVVGAGGVGKSALTIQLIQNHFVDEYDPTIEDSYRKQV IDGETCLLDILDITAGQEEYSAMRDQYMRTGEGFLCVFAINNTKSF EDIHHYREQIKRVKDSQVPMVLVGNKCDLPSRTVDTKQAQDLAR SYGIPFIETSAKTRQGVDDAFYTLVREIRKHKEKMSKDGKKKKKK SKTKCVIM	64
KRAS encoding DNA sequence (from Genbank NM_004985.4) Bold and italicized: siRNA binding regions	ATGACTGAATATAAACTTGTGGTAGTTGGAGCTGCTGGCGTAGGC AAGAGTGCCTTGACGATACAGCTAATTCAGAATCATTTTGTGGAC GAATATGATCCAACAATAGAGGATTCTACAGGAAGCAAGTAGTA ATTGATGGAGAAACCTGTCTCTTGGATATTCTCGACACAGCAGGT CAAGAGGAGTACAG <i>GTGCAATGAGGGACCAGTACAT</i> GAGGACTGGG GAGGGCTTTCTTTGTGTATTTGCCATAAATAATACTAAATCATTT GAAGATATTCACCATTATAGAGAACAAATTAAGAGTTAAGGAC TCTGAAGATGTACCTATGGTCCTAGTAGGAAATAAATGTGATTTG CCTTCTAGAACAGTAGACACAAAACAGGCTCAGGACTTAGCAAGA AGTTATGGAATTCCTTTTATTGAAACATCAGCAAAGACAAGACAG GGTGTTGATGATGCCTTCTATACATTAGTTTCGAGAAATTCGAAAA CATAAAGAAAAGATGAGCAAAGATGGTAAAAAGAAGAAAAAGAG TCAAAGACAAAGTGTGTAATTATGTAA	65
KRAS encoding RNA sequence (from Genbank NM_004985.4) Bold and italicized: siRNA binding regions	AUGACUGAAUUAUAAACUUGUGGUAGUUGGAGCUGGUGGCGUAGGC AAGAGUGCCUUGACGAUACAGCUAAUUCAGAAUCAUUUUGUGGAC GAAUAUGAUCCAACAUAAGAGGAUCCUACAGGAAGCAAGUAGUA AUUGAUGGAGAAACCUGUCUCUUGGAUAUUCUCGACACAGCAGGU CAAGAGGAGUACA <i>GUGCAAUGAGGGACCAGUACA</i> UAGGAGACUGGG GAGGGCUUUCUUUGUGUAUUUGCCAUAAAUAAUACUAAAUCAUUU GAAGAUUUCACCAUUAUAGAGAACAAAUUAAAAGAGUUAAGGAC UCUGAAGAUGUACCUAUGGUCCUAGUAGGAAAUAAAUGUGAUUUUG CCUUCUAGAACAGUAGACACAAAACAGGCUCAGGACUUAGCAAGA AGUUAUGGAAUUCUUUUUAUUGAAACAUCAGCAAAGACAAGACAG GGUGUUGAUGAUGCCUUCUAUACAUAUAGUUCGAGAAAAUUCGAAAA	66

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	CAUAAAGAAAAGAUGAGCAAAGAUGGUAAAAAGAAGAAAAAGAAG UCAAGACAAAGUGUGUAAUUAUGUAA	
Human IL-15 amino acid (Genbank NM_000585.4) Underlined: signal sequence	MRISKPHLRISISIQCYLCLLLNSHFLTEAGIHVFILGCF SAGLPK TEANWVNVISDLKKIEDLIQSMHIDATLYTESDVHPSCKVTAMKC FLLELQVISLES GDASIHDTVENLIILANNSLSSNGNVTESGCKE CEELEEKNIKEFLQSFVHIVQMFIN TS	67
Mature Human IL-15 amino acid (Genbank NM_000585.4)	GIHVFILGCF SAGLPKTEANWVNVISDLKKIEDLIQSMHIDATLY TESDVHPSCKVTAMKCFLELQVISLES GDASIHDTVENLIILAN NSLSSNGNVTESGCKECEELEEKNIKEFLQSFVHIVQMFIN TS	68
Human IL-15 nucleic acid (Genbank NM_000585.4) Underlined: coding sequence Bold: signal sequence	ATGTTCCATCATGTTCCATGCTGCTGACGTCACATGGAGCACAGA AATCAATGTTAGCAGATAGCCAGCCCATACAAGATCGTATTGTAT TGTAGGAGGCATTGTGGATGGATGGCTGCTGGAAACCCCTTGCCA TAGCCAGCTCTTCTTCAATACTTAAAGATTIACCGTGGCTTTGAG TAATGAGAATTTTCGAAACCACATTGAGAAAGTATTTCCATCCAGT GCTACTTGTGTTTACTTCTAAACAGTCATTTTCTAACTGAAGCTG GCATTCATGTCTTCATTTTGGGCTGTTTCAGTGCAGGGCTTCCTA AAACAGAAGCCAAC TGGGTGAATGTAATAAGTGATTGAAAAAAA TTGAAGATCTTATTCAATCTATGCATATTGATGCTACTTTATATA CGGAAAGTGATGTTACCCCGAGTTGCAAAGTAACAGCAATGAAGT GCTTTCTCTTGGAGTTACAAGTTATTTCACTTGAGTCCGGAGATG CAAGTATTCATGATACAGTAGAAAATCTGATCATCCTAGCAAACA ACAGTTTGCTCTTCTAATGGGAATGTAACAGAATCTGGATGCAAAG AATGTGAGGAAC TGGAGGAAAAAATATTAAGAATTTTTGCAGA GTTTTGTACATATTGTCCAAATGTTTCATCAACACTTCTTGATTGC AATTGATTCTTTTTAAAGTGTTTCTGTTATTAACAAACATCACTC TGCTGCTTAGACATAACAAAACACTCGGCATTTCAAATGTGCTGT CAAAACAAGTTTTTCTGTCAAGAAGATGATCAGACCTTGGATCAG ATGAAC TCTTAGAAATGAAGGCAGAAAAATGTCATTGAGTAATAT AGT	69
CD155 amino acid (Genbank NM_006505.4) (Transcript variant 1)	MARAMAAWPLLLVALLVLSWPPPGTGDVVVQAPTQVPGFLGDSV TLPCYLQVPNMEVTHVSQLTWARHGESGSMVVFHQ TQG PSYSESK RLEFVAARLGAELRNASLRMFGLRVEDEGNYTCLFVTFPQGSRSV DIWLRVLAKPQNTAEVQKVQLTGEPVPMARCVSTGGRPPAQITWH SDLGMPNTSQVPGFLSGTVTVTSLWILVPSSQVDGKNVTCKVEH ESFEKPQLLTVNLTVYYPPEVSI SGYDNNWYLGQNEATLTCDARS NPEPTGYNWSTTMGPLPPFAVAQGAQLLIRPVDKPINTTLCNVT NALGARQAELTVQVKEGPPSEHSGMSRNAIIFLVLGILVFLILLG IGIYFYWSKCSREVLWHCHLCPSSTEHASASANGHVSYSAVSREN SSSQDPQTEGTR	70
CD155 encoding DNA sequence (from Genbank NM_006505.4) Bold and italicized:	ATGGCCCGAGCCATGGCCGCGCGTGGCCGCTGCTGCTGGTGGCG CTACTGGTGCTGTCCTGGCCACCCCGAGGAACCGGGGACGTCGTC GTGCAGGCGCCACCCAGGTGCCCGGCTTCTTGGGCGACTCCGTG ACGCTGCCCTGCTACCTACAGGTGCCCAACATGGAGGTGACGCAT GTGTACAGCTGACTTGGGCGCGGCATGGTGAATCTGGCAGCATG GCCGCTCTTCCACCAAACGCAGGGCCCCAGCTATTGGAGTCCAAA CGGCTGGAATTCGTGGCAGCCAGACTGGGCGCGGAGCTGCGGAAT	71

Protein or Nucleic Acid	Sequence	SEQ ID NO:
siRNA binding regions	GCCTCGCTGAGGATGTTTCGGGTTGCGCGTAGAGGATGAAGGCAAC TACACCTGCCTGTTTCGTACGTTCCCGCAGGGCAGCAGGAGCGTG GATATCTGGCTCCGAGTGCTTGCCAAGCCCCAGAACACAGCTGAG GTTTCAAGAGGTCCAGCTCACTGGAGAGCCAGTGCCCATGGCCCCG TGCGTCTCCACAGGGGGTGCCTCCAGCCCAAATCACCTGGCAC TCAGACCTGGGCGGGATGCCCAATACGAGCCAGGTGCCAGGGTTC CTGTCTGGCACAGTCACTGTCAACAGCCTCTGGATATTGGTGCC TCAAGCCAGGTGGACGGCAAGAATGTGACCTGCAAGGTGGAGCAC GAGAGCTTTGAGAAGCCTCAGCTGCTGACTGTGAACCTCACCGTG TACTACCCCCCAGAGGTATCCATCTCTGGCTATGATAACAACCTGG TACCTTGGCCAGAATGAGGCCACCTGACCTGCGATGCTCGCAGC AACCAGAGCCACAGGCTATAATTGGAGCACGACCATGGGTCCC CTGCCACCCTTTGCTGTGGCCAGGGCGCCAGCTCCTGATCCGT CCTGTGGACAAACCAATCAACACAACCTTAACTGCAACGTCACC AATGCCCTAGGAGCTCGCCAGGCAGAACTGACCGTCCAGGTCAAA GAGGGACCTCCAGTGAGCACTCAGGCAT GTCCCGTAACGCCATC ATCTTC CCTGGTTCTGGGAATCCTGGTTTTTCTGATCCTGCTGGGG ATCGGGATTTATTTCTATTGGTCCAAATGTTCCCGTGAGGTCCCTT TGGCACTGTCTGTGTCCCTCGAGTACAGAGCATGCCAGCGCC TCAGCTAATGGGCATGTCTCTATTTCAGCTGTGAGCAGAGAGAAC AGCTCTTCCCAGGATCCACAGACAGAGGGCACAAGGTGA	
CD155 encoding RNA sequence (from Genbank NM_006505.4) Bold and italicized: siRNA binding regions	AUGGCCCCGAGCCAUGGCCGCGCGUGGCCGUGCUGCUGGUGGCG CUACUGGUGCUGUCCUGGCCACCCCCAGGAACCGGGGACGUCGUC GUGCAGGCGCCCCACCCAGGUGCCCGCUUCUUGGGCGACUCCGUG ACGCGUCCUGCUACCUACAGGUGCCCAACAUGGAGGUGACGCAU GUGUCACAGCUGACUUGGGCGCGCAUGGUGAAUCUGGCAGCAUG GCCGUCUCCACCAAACGCAGGGCCCCAGCUAUUCGGAGUCCAAA CGGCUGGAAUUCGUGGCAGCCAGACUGGGCGCGGAGCUGCGGAAU GCCUCGCUGAGGAUGUUCGGGUUGCGCGUAGAGGAUGAAGGCAAC UACACCUGCCUGUUCGUCACGUUCCCGCAGGGCAGCAGGAGCGUG GAUAUCUGGCUCCGAGUGCUUGCCAAGCCCCAGAACACAGCUGAG GUUCAGAAGGUCCAGCUCACUGGAGAGCCAGUGCCCAUGGCCCCG UGCGUCUCCACAGGGGGUGCGCCGCCAGCCCAAUACCUUGGCAC UCAGACCUGGGCGGGAUGCCCAAUACGAGCCAGGUGCCAGGGUUC CUGUCUGGCACAGUCACUGUACACAGCCUCUGGAUAUUGGUGCCC UCAAGCCAGGUGGACGGCAAGAAUGUGACCUGCAAGGUGGAGCAC GAGAGCUUUGAGAAGCCUCAGCUGCUGACUGUGAACCUCACCGUG UACUACCCCCCAGAGGUAUCCAUCUCUGGCUAUGAUAAACAACUGG UACCUUGGCCAGAAUGAGGCCACCCUGACCUGCGAUGCUCGCAGC AACCAGAGCCACAGGCUAUAUUGGAGCACGACCAUGGGUCCC CUGCCACCCUUGCUGUGGGCCAGGGCGCCAGCUCCUGAUCCGU CCUGUGGACAAACCAUAACACAACUUAUUCUGCAACGUCACC AAUGCCCUAGGAGCUCGCCAGGCAGAACUGACCGUCCAGGUCAAA GAGGGACCUCCAGUGAGCACUCAGGCAU GUCCCGUAACGCCAUC AUCUUC CUGGUUCUGGGAUCCUGGUUUUUCUGAUCCUGCUGGGG AUCGGGAUUUAUUCUAUUGGUCCAAAUUGUCCCGUGAGGUCCU UGGCACUGUCAUCUGUGUCCUCGAGUACAGAGCAUGCCAGCGCC UCAGCUAAUGGGCAUGUCUCCUAUUCAGCUGUGAGCAGAGAGAAC AGCUCUUCCCAGGAUCCACAGACAGAGGGCACAAGGUGA	72

Protein or Nucleic Acid	Sequence	SEQ ID NO:
PD-L1 amino acid (Genbank NM_014143.3) (Transcript variant 1)	MRIFAVFI FMTYWHLLNAFTVTVPKDLYVVEYGSNMTIECKFPVE KQLDLAALIVYWEMEDKNIIQFVHGEEDLKVOHSSYRQRARLLKD QLSLGNAALQITDVKLQDAGVYRCMISYGGADYKRI TVKVNPYN KINQRILVVDVPTSEHELTCQAEGYPKAEVIWTSSDHQVLSGKTT TTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTAELV IPELPLAHPPNERTHLVILGAILLCLGVALTFIFRLRKGRMMDVK KCGIQDTNSKKQSDTHLEET	73
PD-L1 encoding DNA sequence (from Genbank NM_014143.3) Bold and italicized: siRNA binding regions	ATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATTG CTGAACGCATTTACTGTACGGTTCCCAAGGACCTATATGTGGTA GAGTATGGTAGCAATATGACAATTGAATGCAAAATCCAGTAGAA AAACAATTAGACCTGGCTGCACTAATTGTCTATTGGGAAATGGAG GATAAGAACATTATTCAATTTGTGCATGGAGAGGAAGACCT <i>GAAG GTTTCAGCATAGTAGCTA</i> CAGACAGAGGGCCCCGGCTGTTGAAGGAC CAGCTCTCCCTGGGAAATGCTGCACTTCAGATCACAGATGTGAAA TTGCAGGATGCAGGGGTGTACCGCTGCATGATCAGCTATGGTGGT GCCGACTACAAG <i>GCGAATTACTGTGAAAGTCAAT</i> GCCCCATACAAC AAAATCAACCAAGAATTTTGGTTGTGGATCCAGTCACCTCTGAA CATGAACGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTC ATCTGGACAAGCAGTGACCATCAAGTCCTGAGTGGTAAGACCACC ACCACCAATTCCAAGAGAGAGGAGAAGCTTTTCAATGT <i>GACCAGC ACACTGAGAATCAA</i> CACAACAACATAATGAGATTTTCTACTGCACT TTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTGGTC ATCCCAGAACTACCTCTGGCACATCCTCCAAATGAAAGGACTCAC TTGGTAATTCTGGGAGCCATCTTATTATGCCTTGGTGTAGCACTG ACATTTCATCTTCCGTTTAAGAAAAGGGAGAATGATGGATGTGAAA AAATGTGGCATCCAAGATACAAACTCAAAGAAGCAAAGTGATACA CATTTGGAGGAGACGTAA	74
PD-L1 encoding RNA sequence (from Genbank NM_014143.3) Bold and italicized: siRNA binding regions	AUGAGGAUUAUUUGCUGUCUUUAUUAUUAUGACCUACUGGCAUUUG CUGAACGCAUUUACUGUCACGGUUCCTCAAGGACCUAUUAUGUGGUA GAGUAUGGUAGCAUAUAGACAAUUGAAUGCAAAUUCAGUAGAA AAACAUAUAGACCUGGCUGCACUAAUUGUCUAUUGGGAAAUGGAG GAUAAGAACAUAUUAUUAUUAUUGUGCAUGGAGAGGAAGACCU <i>GAAG GUUCAGCAUAGUAGCUA</i> CAGACAGAGGGCCCCGGCUGUUGAAGGAC CAGCUCUCCUGGGAAAUGCUGCACUUCAGAUACAGAUUGUGAAA UUGCAGGAUGCAGGGGUGUACCGCUGCAUGAUCAGCUAUGGUGGU GCCGACUACAAG <i>GCGAAUUAUCUGUGAAAGUCA</i> AUGCCCCAUACAAC AAAAUCAACCAAGAAUUAUUGGUUGUGGAUCCAGUACCCUCUGAA CAUGAACUGACAUGUCAGGCUGAGGGCUACCCCAAGGCCGAAGUC AUCUGGACAAGCAGUGACCAUCAAGUCCUGAGUGGUAAGACCACC ACCACCAAUUCCAAGAGAGAGGAGAAGCUUUUCAUUGU <i>GACCAGC ACACUGAGAAUCA</i> CACAACAACUAUAGAGAUUUUCUACUGCACU UUUAGGAGAUUAGAUCUGAGGAAAACCAUACAGCUGAAUUGGUC AUCCCAGAACUACCUCUGGCACAUCCUCCAAUUGAAAGGACUCAC UUGGUAUUUCUGGGAGCCAUCUUAUUAUGCCUUGGUGUAGCACUG ACAUUCAUCUCCGUUUAAGAAAAGGGAGAAUGAUGGAUGUGAAA AAUUGUGGAUCCAAGAUACAACUCAAGAAGCAAAGUGAUACA CAUUUGGAGGAGACGUAA	75
c-Myc amino acid (Genbank NM_002467.4)	MDFFRVVENQQPPATMPLNVSFTNRNYDL DYDSVQPYFYCDEEEN FYQQQQQSELQPPAPSEDIWKKFELLPTPPLSPSRRSGLCSPSYV	76

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	AVTPFSLRGDNDGGGGSFSTADQLEMVTELLGGDMVNQSFICDPD DETFIKNII IQDCMWSGFSAAAKLVSEKLASYQAARKDSGSPNPA RGHSVCSTSSLYLQDL SAAASECIDPSVVFYPLNDSSSPKSCAS QDSSAFSPSSDLSLSTESSPQGSPEPLVLHEETPPTTSSDSEEE QEDEEEIDVVSVEKRQAPGKRSESGSPSAGGHSKPPHSPLVLKRC HVSTHQHNYAAPSTRKDYPAKRVLDSVRVLRQISNNRKCTSP RSSDTEENVKRRTHNVLERQRRNELKRSFFALRDQIPELENNEKA PKVVILKKATAYILSVQAEEQKLI SEEDLLRKRREQLKHKLEQLR NSCA	
c-Myc encoding DNA sequence (from Genbank NM_002467.4) Bold and italicized: siRNA binding regions	ATGGATTTTTTTTCGGGTAGTGGAAAAACCAGCAGCCTCCCGCGACG ATGCCCCCTCAACGTTAGCTTCACCAACAGGAACATGACCTCGAC TACGACTCGGTGCAGCCGTATTTCTACTGCGACGAGGAGGAGAAC TTCTACCAGCAGCAGCAGCAGAGCGAGCTGCAGCCCCCGGCGCCC AGCGAGGATATCTGGAAGAAATTCGAGCTGCTGCCACCCCGCCC CTGTCCCCTAGCCGCCGCTCCGGGCTCTGCTCGCCCTCCTACGTT GCGGTACACCCCTTCTCCCTTCGGGGAGACAACGACGGCGGTGGC GGGAGCTTCTCCACGGCCGACCAGCTGGAGATGGTGACCGAGCTG CTGGGAGGAGACATGGTGAACCAGAGTTTCATCTGCGACCC GGAC GACGAGACCTTCATCA AAAAACATCATCATCCAGGACTGTATGTGG AGCGGCTTCTCGGCCGCCGCAAGCTCGTCTCAGAGAAGCTGGCC TCCTACCAGGCTGCGCGCAAAGACAGCGGCAGCCCGAACCCCGCC CGCGGCCACAGCGTCTGCTCCACCTCCAGCTTGTAACCTGCAGGAT CTGAGCGCCGCCGCTCAGAGTGCATCGACCCCTCGGTGGTCTTC CCCTACCCTCTCAACGACAGCAGCTCGCCCAAGTCTGCGCCTCG CAAGACTCCAGCGCCTTCTCTCCGTCTCGGATTCTCTGCTCTCC TCGACGGAGTCTTCCCGCAGGGCAGCCCCGAGCCCTGGTGCTC CATGAGGAGACACCGCCACCACCAGCAGCGACTCTGAGGAGGAA CAAGAAGATGAGGAAGAAATCGATGTTGTTTCTGTGGAAGAGAGG CAGGCTCCTGGCAAAAGGTCAGAGTCTGGATCACCTTCTGCTGGA GGCCACAGCAAACCTCCTCACAGCCCACTGGTCTCAAGAGGTGC CACGTCTCCACACATCAGCACAACCTACGCAGCGCCTCCCTCCACT CGGAAGGACTATCCTGCTGCCAAGAGGGTCAAGTTGGACAGTGTC AGAGTCTTGAGACAGATCAGCAACAACCGAAAATGCACCAGCCCC AGGTCTTCGGACACCGAGGAGAATGTCAAGAGGCGAACACACAAC GTCTTGAGCGCCAGAGGAGGAACGAGCTAAAACGGAGCTTTTTT GCCCTGCGTGACCAGATCCCGGAGTTGGAAAACAATGAAAAGGCC CCCAAGGTAGTTATCCTTAAAAAAGCCACAGCATAACATCCTGTCC GTCCAAGCAGAGGAGCAAAAGCTCATTTCTGAAGAGGACTTGTTG CGGAAACGACGAGAACAGTTGAAACACAACTTGAACAGCTACGG AACTCTTGTCGTAA	77
c-Myc encoding RNA sequence (from Genbank NM_002467.4) Bold and italicized: siRNA binding regions	AUGGAUUUUUUUCGGGUAGUGGAAAAACCAGCAGCCUCCCGCGACG AUGCCCCUCAACGUUAGCUUACCAACAGGAACUAUGACCUCGAC UACGACUCGGUGCAGCCGUUUUUUACUGCGACGAGGAGGAGAAC UUCUACCAGCAGCAGCAGCAGAGCGAGCUGCAGCCCCCGGCGCCC AGCGAGGAUAUCUGGAAGAAAUUCGAGCUGCUGCCACCCCGCCC CUGUCCCCUAGCCGCCGCUCCGGGCUUCUGCUGCCCUCCUACGUU GCGGUCACACCCUUCUCCUUCGGGGAGACAACGACGGCGGUGGC GGGAGCUUCUCCACGGCCGACCAGCUGGAGAUUGGUGACCGAGCUG CUGGGAGGAGACAUGGUGAACCAGAGUUUCAUCUGCGACCC GGAC	78

Protein or Nucleic Acid	Sequence	SEQ ID NO:
	<p><u>GACGAGACCUUCAUCA</u>AAAAACAUCAUCAUCCAGGACUGUAUGUGG AGCGGCUUCUCGGCCGCCGCCAAGCUCGUCUCAGAGAAGCUGGCC UCCUACCAGGCUGCGCGCAAAGACAGCGGCAGCCCCGAACCCCGCC CGCGGCCACAGCGUCUGCUCCACCUCAGCUUGUACCUGCAGGAU CUGAGCGCCGCCGCCUCAGAGUGCAUCGACCCUCGGUGGUCUUC CCCUACCCUCUCAACGACAGCAGCUCGCCCCAAGUCCUGCGCCUCG CAAGACUCCAGCGCCUUCUCUCCGUCCUCGGAUUCUCUGCUUCC UCGACGGAGUCCUCCCCGCAGGGCAGCCCCGAGCCCCUGGUGCUC CAUGAGGAGACACCGCCACCACCAGCAGCGACUCUGAGGAGGAA CAAGAAGAUGAGGAAGAAAUCGAUGUUGUUUCUGUGGAAAAAGAGG CAGGCUCCUGGCAAAAGGUCAGAGUCUGGAUCCAUUCUGUGGA GGCCACAGCAAACCUCUCACAGCCCACUGGUCCUCAAGAGGUGC CAGGUCUCCACACAUCAGCACAACUACGCAGCGCCUCCUCCACU CGGAAGGACUAUCCUGCUGCCAAGAGGGUCAAGUUGGACAGUGUC AGAGUCCUGAGACAGAUCAAGCAACAACCGAAAAUGCACCAGCCCC AGGUCCUCGGACACCGAGGAGAAUGUCAAGAGGCGAACACACAAC GUCUUGGAGCGCCAGAGGAGGAACGAGCUAAAACGGAGCUUUUUU GCCCUGCGUGACCAGAUCCCGGAGUUGGAAAAACAAUGAAAAGGCC CCCAAGGUAGUUAUCCUAAAAAAGCCACAGCAUACAUCUGUCC GUCCAAGCAGAGGAGCAAAAGCUCAUUUCUGAAGAGGACUUGUUG CGGAAACGACGAGAACAGUUGAAACACAAACUUGAACAGCUACGG AACUCUUGUGCGUAA</p>	
Human IL-7 amino acid (Genbank NM_000880.3) Underlined: signal sequence	<p>MFHVSFRYIFGLPPLILVLLPVASSDCDIEGKDGKQYESVLMVSI DQLLDSMKEIGSNCLNNEFNFFKRHICDANKEGMFLFRAARKLRQ FLKMNSTGDFDLHLLKVSEGTILLNCTGQVKGRKPAALGEAQPT KSLEENKSLKEQKKLNDLCFLKRLLQEIKTCWNKILMGTHEH</p>	79
Mature Human IL-7 amino acid (Genbank NM_000880.3)	<p>DCDIEGKDGKQYESVLMVSI DQLLDSMKEIGSNCLNNEFNFFKRH ICDANKEGMFLFRAARKLRQFLKMNSTGDFDLHLLKVSEGTILL NCTGQVKGRKPAALGEAQPTKSLEENKSLKEQKKLNDLCFLKRLL QEIKTCWNKILMGTHEH</p>	80
Human IL-7 nucleic acid (Genbank NM_000880.3) Underlined: coding sequence Bold: signal sequence	<p>ATGTTCCATGTTTCTTTTAGGTATATCTTTGGACTTCCTCCCTG ATCCTTGTTCTGTTGCCAGTAGCATCATCTGATTGTGATATTGAA GGTAAAGATGGCAAACAATATGAGAGTGTTCTAATGGTCAGCATC GATCAATTATTGGACAGCATGAAAGAAATTGGTAGCAATTGCCTG AATAATGAATTTAACTTTTTTAAAGACATATCTGTGATGCTAAT AAGGAAGGTATGTTTTTATTCCGTGCTGCTCGCAAGTTGAGGCAA TTTCTTAAATGAATAGCACTGGTGATTTTGATCTCCACTTATTA AAAGTTTCAGAAGGCACAACAATACTGTTGAACTGCACTGGCCAG GTAAAGGAAGAAAACCAGCTGCCCTGGGTGAAGCCCAACCAACA AAGAGTTTGGAAGAAAATAAATCTTTAAAGGAACAGAAAAAACTG AATGACTTGTGTTTCTTAAAGAGACTATTACAAGAGATAAAAACT TGTTGGAATAAAATTTTGATGGGCACATAAAGAACACTGA</p>	81
Human IL-12 alpha signal peptide (Genbank NM_000882.4)	<p>MCPARSLLLVATLVLLDHLSLA</p>	142

Protein or Nucleic Acid	Sequence	SEQ ID NO:
Human IL-12 beta signal peptide (Genbank NM_002187.2)	MCHQQLVISWFSLVFLASPLVA	143
Human IL-15 signal peptide (Genbank NM_000585.4)	MRISKPHLRISISIQCYLCLLLNSHFLTEA	144
Human IL-7 signal peptide (Genbank NM_000880.3)	MFHVSFRYIFGLPPLILVLLPVASS	145
Endogenous IL-12 alpha signal peptide nucleic acid	ATGTGTCCAGCGCGCAGCCTCCTCCTTGTGGCTACCCTGGTCCCTCCTGGACCACCTCAGTTTGGCC	146
Endogenous IL-12 beta signal peptide nucleic acid	ATGTGTCAACCAGCAGTTGGTCATCTCTTGGTTTTCCCTGGTTTTTCTGGCATCTCCCTCGTGGCC	147
Endogenous IL-15 signal peptide nucleic acid	ATGAGAATTTTCGAAACCACATTTGAGAAGTATTTCCATCCAGTGC TACTTGTGTTTACTTCTAAACAGTCATTTCTAACTGAAGCT	148
Endogenous IL-7 signal peptide nucleic acid	ATGTTCCACGTGTCCTTCCGGTACATCTTCGGCCTGCCTCCACTG ATCCTGGTGCTGCTGCCTGTGGCCAGCAGC	149

[0208] Table 4. Plasmid Vector Sequences for Compounds 1-17

SEQ ID NO	Compound	Sequence (5' to 3')
82	Compound 1 (pMA-T)	CTAAATTGTAAGCGTTAATATTTTGTAAATTCGCGTTAAAT TTTTGTAAATCAGCTCATTTTTTAACCAATAGGCCGAAATCG GCAAATCCCTTATAAATCAAAGAATAGACCGAGATAGGGTT GAGTGGCCGCTACAGGGCGCTCCCATTCGCCATTTCAGGCTGCG CAACTGTTGGGAAGGGCGTTTCGGTGCGGGCCTCTTCGCTATT ACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCGATTAAAGT TGGGTAACGCCAGGGTTTTCCAGTCACGACGTTGTAAACGA CGGCCAGTGAGCGCGACGTAATACGACTCACTATAGGGCGAAT TGGCGGAAGGCCGTCAAGGCCACGTGTCTTGTCCAGAGCTCGC <u>CACCATGTACAGAATGCAGCTGCTGAGCTGTATCGCCCTGTCT</u> <u>CTGGCCCTGGTCACAAATAGCGCCCCCTACCAGCAGCAGCACCA</u> <u>AGAAAACACAGCTGCAACTGGAACACCTCCTGCTGGACCTGCA</u> GATGATCCTGAACGGCATCAACAACATAAGAACCCCAAGCTG ACCCGGATGCTGACCTTCAAGTTCTACATGCCCAAGAAGGCCA CCGAGCTGAAGCACCTCCAGTGCCTGGAAGAGGAAGTGAAGCC CCTGGAAGAAGTGCTGAATCTGGCCCAGAGCAAGAAGTCCAC CTGAGGCCTAGGGACCTGATCAGCAACATCAACGTGATCGTGC TGGAACTGAAAGGCAGCGAGACAACCTTCATGTGCGAGTACGC CGACGAGACAGCTACCATCGTGGAATTTCTGAACCGGTGGATC

SEQ ID NO	Compound	Sequence (5' to 3')
		ACCTTCTGCCAGAGCATCATCAGCACCTGACCTGAGGTACCT GGAGCACAAGACTGGCCTCATGGGCCTTCCGCTCACTGCCCCG TTTCCAGTCGGGAAACCTGTCGTGCCAGCTGCATTAACATGGT CATAGCTGTTTCCTTGCCTATTGGGCGCTCTCCGCTTCCTCGC TCACTGACTCGCTGCGCTCGGTCGTTTCGGGTAAAGCCTGGGGT GCCTAATGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAA AGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGA CGAGCATCACAAAATCGACGCTCAAGTCAGAGGTGGCGAAAC CCGACAGGACTATAAAGATAACAGGCGTTTCCCCCTGGAAGCT CCCTCGTGCCTCTCCTGTTCCGACCTGCCGCTTACCGGATA CCTGTCCGCTTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCAT AGCTCACGCTGTAGGTATCTCAGTTCGGTGTAGGTGCTTCGCT CCAAGCTGGGCTGTGTGCACGAACCCCCCGTTACGCCGACCG CTGCGCCTTATCCGGTAACTATCGTCTTGAGTCCAACCCGGTA AGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGA TTAGCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTTCTTGAA GTGGTGGCCTAACTACGGCTACACTAGAAGAACAGTATTTGGT ATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTG GTAGCTCTTGATCCGGCAAACAAACCACCGCTGGTAGCGGTGG TTTTTTTGTGTTGCAAGCAGCAGATTACGCGCAGAAAAAAGGA TCTCAAGAAGATCCTTTGATCTTTTCTACGGGGTCTGACGCTC AGTGGAACGAAAACCTACGTTAAGGGATTTTGGTCATGAGATT ATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTAAAAATGA AGTTTTAAATCAATCTAAAGTATATATGAGTAAACTTGGTCTG ACAGTTACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGAT CTGTCTATTTTCGTTTCATCCATAGTTGCCTGACTCCCCGTCGTG TAGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTG CTGCAATGATACCGCGAGAACCACGCTCACCGGCTCCAGATTT ATCAGCAATAAACCAGCCAGCCGGAAGGGCCGAGCGCAGAAGT GGTCCTGCAACTTTATCCGCCTCCATCCAGTCTATTAATTGTT GCCGGGAAGCTAGAGTAAGTAGTTTCGCCAGTTAATAGTTTGGC CAACGTTGTTGCCATTGCTACAGGCATCGTGGTGTACGCTCG TCGTTTGGTATGGCTTCATTACAGCTCCGTTCCCAACGATCAA GGCGAGTTACATGATCCCCATGTTGTGCAAAAAGCGGTTAG CTCCTTCGGTCCTCCGATCGTTGTGAGAAGTAAGTTGGCCGCA GTGTTATCACTCATGGTTATGGCAGCACTGCATAATTCTCTTA CTGTCATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTA CTCAACCAAGTCATTCTGAGAATAGTGATGCGGCGACCGAGT TGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATA GCAGAACTTTAAAAGTGCTCATCATTTGGAACCGTTCTTCGGG GCGAAAACCTCTCAAGGATCTTACCGCTGTTGAGATCCAGTTCG ATGTAACCCACTCGTGCACCCAACTGATCTTCAGCATCTTTTA CTTTACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAA TGCCGCAAAAAGGGAATAAGGGCGACACGGAATGTTGAATA CTCATACTCTTCCTTTTCAATATTATTGAAGCATTTATCAGG GTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGAA AAATAAACAAATAGGGGTTCCGCGCACATTTCCCGGAAAAGTG CCAC
83	Compound 2* (pMA-T)	CTAAATTGTAAGCGTTAATATTTTGTAAATTCGCGTTAAAT TTTTGTAAATCAGCTCATTTTTTAACCAATAGGCCGAAATCG

SEQ ID NO	Compound	Sequence (5' to 3')
		GCAAAATCCCTTATAAATCAAAAGAATAGACCGAGATAGGGTT GAGTGGCCGCTACAGGGCGCTCCCATTCGCCATTCAGGCTGCG CAACTGTTGGGAAGGGCGTTTCGGTGCGGGCCTCTTCGCTATT ACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCGATTAAAGT TGGGTAACGCCAGGGTTTTCCAGTCACGACGTTGTAAACGA CGGCCAGTGAGCGCGACGTAATACGACTCACTATAGGGCGAAT TGGCGGAAGGCCGTC AAGGCCACGTGTCTTGTCCAGAGCTC GC CACCATGCTGAAACTGCTGCTGCTCCTGTGTATCGCCCTGTCT CTGGCCGCCACAAATAGCGCCCTACCAGCAGCTCCACCAAGA AAACACAGCTGCAACTGGAACATCTGCTGCTGGACCTGCAGAT GATCCTGAACGGCATCAACAAC TACAAGAACCCCAAGCTGACC CGGATGCTGACCTTCAAGTTCACATGCCCAAGAAGGCCACCG AGCTGAAGCACCTCCAGTGCCTGGAAGAGGAAC TGAAGCCCT GGAAGAAGTGCTGAATCTGGCCAGAGCAAGAACTTCCACCTG AGGCCTAGGGACCTGATCAGCAACATCAACGTGATCGTGCTGG AACTGAAAGGCAGCGAGACAACCTTCATGTGCGAGTACGCCGA CGAGACAGCTACCATCGTGGAATTTCTGAACCGGTGGATCACC TTCTGCCAGAGCATCATCAGCACCTTGACCTGAGGTACCTGGA GCACAAGACTGGCCTCATGGGCCTTCCGCTCACTGCCCGCTTT CCAGTCGGGAAACCTGTCTGTGCCAGCTGCATTAACATCAT AGCTGTTTCCTTGCGTATTGGGCGCTCTCCGCTTCTCTCGCTCA CTGACTCGCTGCGCTCGGTCTGTTCCGGTAAAGCCTGGGGTGCC TAATGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGG CCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGA GCATCACAAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCG ACAGGACTATAAAGATAACAGGCGTTTCCCCCTGGAAGCTCCC TCGTGCGCTCTCCTGTTCCGACCCTGCCGCTTACCGGATACCT GTCCGCTTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCATAGC TCACGCTGTAGGTATCTCAGTTCGGTGTAGGTCTGCTCCCA AGCTGGGCTGTGTGCACGAACCCCCCGTTCAGCCCGACCGCTG CGCCTTATCCGGTAACTATCGTCTTGAGTCCAACCCGGTAAGA CACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTA GCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTCTTGAAGTG GTGGCCTAACTACGGCTACACTAGAAGAACAGTATTTGGTATC TGGCCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTA GCTCTTGATCCGGCAAACAAACCACCGCTGGTAGCGGTGGTTT TTTTGTTTGCAAGCAGCAGATTACGCGCAGAAAAAAGGATCT CAAGAAGATCCTTTGATCTTTTCTACGGGGTCTGACGCTCAGT GGAACGAAAAC TACGTTAAGGGATTTTGGTCATGAGATTATC AAAAAGGATCTTACCTAGATCCTTTTAAATTAAAAATGAAGT TTTAAATCAATCTAAAGTATATATGAGTAAACTTGGTCTGACA GTTACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTG TCTATTTTCGTTTCATCCATAGTTGCCTGACTCCCCGTCGTGTAG ATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTG CAATGATACCGCGAGAACCACGCTCACCGGCTCCAGATTTATC AGCAATAAACCAGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGT CCTGCAACTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCC GGGAAGCTAGAGTAAGTAGTTCGCCAGTTAATAGTTTGCGCAA CGTTGTTGCCATTGCTACAGGCATCGTGGTGTCACGCTCGTCCG TTTGGTATGGCTTCATT CAGCTCCGGTTCCCAACGATCAAGGC

SEQ ID NO	Compound	Sequence (5' to 3')
		GAGTTACATGATCCCCCATGTTGTGCAAAAAAGCGGTTAGCTC CTTCGGTCCTCCGATCGTTGTGAGAAAGTAAGTTGGCCGCAGTG TTATCACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTG TCATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTC AACCAAGTCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGC TCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATAGCA GAACTTTAAAAGTGCTCATCATTGGAAAACGTTCTTCGGGGCG AAACTCTCAAGGATCTTACCGCTGTTGAGATCCAGTTCGATG TAACCCACTCGTGCACCCAAGTATCTTCAGCATCTTTTACTT TCACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGC CGCAAAAAGGGAATAAGGGCGACACGGAAATGTTGAATACTC ATACTCTTCCTTTTCAATATTATTGAAGCATTATCAGGGTT ATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGAAAAA TAAACAAATAGGGGTTCCGCGCACATTTCCCCGAAAAGTGCCA C
84	Compound 3* (pMA-T)	CTAAATTGTAAGCGTTAATATTTTGTAAATTCGCGTTAAAT TTTTGTAAATCAGCTCATTTTTTTAACCAATAGGCCGAAATCG GCAAAATCCCTTATAAATCAAAAGAATAGACCGAGATAGGGTT GAGTGGCCGCTACAGGGCGCTCCCATTCGCCATTACAGGCTGCG CAACTGTTGGGAAGGGCGTTTCGGTGCGGGCCTCTTCGCTATT ACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCATTAAAGT TGGGTAACGCCAGGGTTTTCCAGTCACGACGTTGTAAACGA CGGCCAGTGAGCGCGACGTAATACGACTCACTATAGGGCGAAT TGGCGGAAGGCCGTCAGGCCACGTGTCTTGTCAGAGCTC <u>GC</u> <u>CACCATGTTGTTGCTGCTGCTCGCCTGTATTGCCCTGGCCTCT</u> <u>ACAGCCGCCGCTACAAATTCTGCCCCCTACCAGCAGCTCCACCA</u> <u>AGAAAACCCAGCTGCAACTGGAACATCTGCTGCTGGACCTGCA</u> <u>GATGATCCTGAACGGCATCAACAAC</u> TACAAGAACCCCAAGCTG ACCCGGATGCTGACCTTCAAGTTCTACATGCCCAAGAAGGCCA CCGAGCTGAAGCACCTCCAGTGCCTGGAAGAGGAAGCTGAAGCC CCTGGAAGAAGTGCTGAATCTGGCCCAGAGCAAGAAGCTTCCAC CTGAGGCCTAGGGACCTGATCAGCAACATCAACGTGATCGTGC TGGAAGTGAAGGCAGCGAGACAACCTTCATGTGCGAGTACGC CGACGAGACAGCTACCATCGTGGAATTTCTGAACCGGTGGATC ACCTTCTGCCAGAGCATCATCAGCACCCCTGACCTGAGGTACCT GGAGCACAAGACTGGCCTCATGGGCCTTCCGCTCACTGCCCGC TTTCCAGTCGGGAAACCTGTCGTGCCAGCTGCATTAACATGGT CATAGCTGTTTCCCTTGCGTATTGGGCGCTCTCCGCTTCCCTGC TCACTGACTCGCTGCGCTCGGTTCGTTTCGGGTAAAGCCTGGGGT GCCTAATGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAA AGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGA CGAGCATCACAAAATCGACGCTCAAGTCAGAGGTGGCGAAAC CCGACAGGACTATAAAGATAACAGGCGTTTCCCCCTGGAAGCT CCCTCGTGCCTCTCCTGTTCCGACCCTGCCGCTTACCGGATA CCTGTCCGCCTTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCAT AGCTCACGCTGTAGGTATCTCAGTTCGGTGTAGGTGCTTCGCT CCAAGCTGGGCTGTGTGCACGAACCCCCCGTTCAGCCCGACCG CTGCGCCTTATCCGGTAAGTATCGTCTTGAGTCAACCCGGTA AGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGA TTAGCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTTCTTGAA

SEQ ID NO	Compound	Sequence (5' to 3')
		GTGGTGGCCTAACTACGGCTACACTAGAAGAACAGTATTTGGT ATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTG GTAGCTCTTGATCCGGCAAACAAACCACCGCTGGTAGCGGTGG TTTTTTTGTTTGCAAGCAGCAGATTACGCGCAGAAAAAAGGA TCTCAAGAAGATCCTTTGATCTTTTCTACGGGGTCTGACGCTC AGTGGAACGAAAACACGTTAAGGGATTTTGGTCATGAGATT ATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTAATAATGA AGTTTTAAATCAATCTAAAGTATATATGAGTAAACTTGGTCTG ACAGTTACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGAT CTGTCTATTTTCGTTTCATCCATAGTTGCCTGACTCCCCGTCGTG TAGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTG CTGCAATGATACCGCGAGAACCACGCTCACC GGCTCCAGATTT ATCAGCAATAAACCAGCCAGCCGGAAGGGCCGAGCGCAGAAGT GGTCTTGCAACTTTATCCGCCTCCATCCAGTCTATTAATTGTT GCCGGAAGCTAGAGTAAGTAGTTCGCCAGTTAATAGTTTGCG CAACGTTGTTGCCATTGCTACAGGCATCGTGGTGTACGCTCG TCGTTTGGTATGGCTTCATTCAGCTCCGGTTCCCAACGATCAA GGCGAGTTACATGATCCCCATGTTGTGCAAAAAAGCGGTTAG CTCCTTCGGTCCCTCCGATCGTTGTGAGAAGTAAGTTGGCCGCA GTGTTATCACTCATGGTTATGGCAGCACTGCATAATTCTCTTA CTGTCTATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTA CTCAACCAAGTCATTCTGAGAATAGTGTATGCGGCGACCGAGT TGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATA GCAGAACTTTAAAGTGCTCATCATTTGGAACGTTCTTCGGG GCGAAACTCTCAAGGATCTTACCGCTGTTGAGATCCAGTTCTG ATGTAACCCACTCGTGCACCCAACTGATCTTCAGCATCTTTTA CTTTCACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAA TGCCGCAAAAAGGGAATAAGGGCGACACGGAAATGTTGAATA CTCATACTCTTCCTTTTTCAATATTATTGAAGCATTTATCAGG GTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGAA AAATAAACAAATAGGGGTTCCGCGCACATTTCCCCGAAAAGTG CCAC
85	Compound 4* (pMA-T)	CTAAATTGTAAGCGTTAATATTTTGTAAAATTTCGCGTTAAAT TTTTGTAAATCAGCTCATTTTTTAACCAATAGGCCGAAATCG GCAAAATCCCTTATAAATCAAAAGAATAGACCGAGATAGGGTT GAGTGGCCGCTACAGGGCGCTCCCATTCGCCATTTCAGGCTGCG CAACTGTTGGGAAGGGCGTTTCGGTGCGGGCCTCTTCGCTATT ACGCCAGCTGGCGAAAGGGGATGTGCTGCAAGGCGATTAAAGT TGGGTAACGCCAGGGTTTTCCAGTCACGACGTTGTAAAACGA CGGCCAGTGAGCGCGACGTAATACGACTCACTATAGGGCGAAT TGGCGGAAGGCCGTCAAGGCCACGTGTCTTGCTCCAGAGCTCGC <u>CACCATGTTGTTGCTGCTGCTCGCCTGTATTGCCCTGGCCTCT</u> <u>ACAGCCCTGGTCACCAATTCTGCCCCTACCAGCAGCTCCACCA</u> <u>AGAAAACCCAGCTGCAACTGGAACATCTGCTGCTGGACCTGCA</u> <u>GATGATCCTGAACGGCATCAACAAC</u> TACAAGAACCCCAAGCTG ACCCGGATGCTGACCTTCAAGTTCTACATGCCCAAGAAGGCCA CCGAGCTGAAGCACCTCCAGTGCCTGGAAGAGGAAGCTGAAGCC CCTGGAAGAAGTGTGAATCTGGCCCAGAGCAAGAAGCTTCCAC CTGAGGCCTAGGGACCTGATCAGCAACATCAACGTGATCGTGC TGGAAGTGAAGGCAGCGAGACAACCTTCATGTGCGAGTACGC

SEQ ID NO	Compound	Sequence (5' to 3')
		<p>CGACGAGACAGCTACCATCGTGGAATTTCTGAACCGGTGGATC ACCTTCTGCCAGAGCATCATCAGCACCGTGACCTGAGGTACCT GGAGCACAAGACTGGCCTCATGGGCCTTCCGCTCACTGCCCCG TTTCCAGTCGGGAAACCTGTCGTGCCAGCTGCATTAACATGGT CATAGCTGTTTCTTGGTATTGGGCGCTCTCCGCTTCTTCGC TCACTGACTCGCTGCGCTCGGTCGTTCCGGTAAAGCCTGGGGT GCCTAATGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAA AGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGA CGAGCATCACAAAATCGACGCTCAAGTCAGAGGTGGCGAAAC CCGACAGGACTATAAAGATAACCAGGCGTTTCCCCCTGGAAGCT CCCTCGTGCCTCTCTGTTCCGACCGTGGCGCTTACCGGATA CCTGTCCGCTTTCTCCCTTCGGAAGCGTGGCGCTTTCTCAT AGCTCACGCTGTAGGTATCTCAGTTCGGTGTAGGTGCTTCGCT CCAAGCTGGGCTGTGTGCACGAACCCCCCGTTCAGCCCGACCG CTGCGCCTTATCCGGTAACTATCGTCTTGAGTCAACCCGGTA AGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGA TTAGCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTCTTGAA GTGGTGGCCTAACTACGGCTACACTAGAAGAACAGTATTTGGT ATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAGAGAGTTG GTAGCTCTTGATCCGGCAAACAAACCACCGCTGGTAGCGGTGG TTTTTTTGTGCAAGCAGCAGATTACGCGCAGAAAAAAGGA TCTCAAGAAGATCCTTTGATCTTTTCTACGGGGTCTGACGCTC AGTGGAACGAAAACCTACGTTAAGGGATTTTGGTCATGAGATT ATCAAAAAGGATCTTACCTAGATCCTTTTAAATTAAAAATGA AGTTTTAAATCAATCTAAAGTATATATGAGTAAACTTGGTCTG ACAGTTACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGAT CTGTCTATTTTCGTTTCATCCATAGTTGCCTGACTCCCCGTCGTG TAGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTG CTGCAATGATACCGCGAGAACCACGCTCACC GGCTCCAGATTT ATCAGCAATAAACCAGCCAGCCGGAAGGGCCGAGCGCAGAAGT GGTCCTGCAACTTTATCCGCCTCCATCCAGTCTATTAATTGTT GCCGGGAAGCTAGAGTAAGTAGTTCCGCCAGTTAATAGTTTGCG CAACGTTGTTGCCATTGCTACAGGCATCGTGGTGTACGCTCG TCGTTTGGTATGGCTTCATTTCAGTCCGGTTCCCAACGATCAA GGCGAGTTACATGATCCCCATGTTGTGCAAAAAGCGGTTAG CTCCTTCGGTCCCTCCGATCGTTGTGCAAGTAAGTTGGCCGCA GTGTTATCACTCATGGTTATGGCAGCACTGCATAATTCTCTTA CTGTCATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTA CTCAACCAAGTCATTCTGAGAATAGTGATGCGGCGACCGAGT TGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATA GCAGAACTTTAAAAGTGCTCATCATTTGAAAACGTTCTTCGGG GCGAAAACCTCTCAAGGATCTTACCGCTGTTGAGATCCAGTTTCG ATGTAACCCACTCGTGCACCCAACTGATCTTCAGCATCTTTTA CTTTCACAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAA TGCCGCAAAAAGGGAATAAGGGCGACACGGAAATGTTGAATA CTCATACTCTTCTTTTCAATATTATTGAAGCATTTATCAGG GTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGAA AAATAAACAAATAGGGGTTCCGCGCACATTTCCCCGAAAAGTG CCAC</p>

SEQ ID NO	Compound	Sequence (5' to 3')
86	Compound 5 (pMK-RQ)	CTAAATTGTAAGCGTTAATATTTTGTAAATTCGCGTTAAAT TTTTGTAAATCAGCTCATTATTTTAAACCAATAGGCCGAAATCG GCAAAATCCCTTATAAATCAAAAGAATAGACCGAGATAGGGTT GAGTGGCCGCTACAGGGCGCTCCCATTCGCCATTAGGCTGCG CAACTGTTGGGAAGGGCGTTTCGGTGCGGGCCTCTTCGCTATT ACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCGATTAAAGT TGGGTAAACGCCAGGGTTTTCCCAGTCACGACGTTGTAAACGA CGGCCAGTGAGCGCGACGTAATACGACTCACTATAGGGCGAAT TGAAGGAAGGCCGTCAAGGCCGCATGCCACCATGTACAGAATG <u>CAGCTGCTGAGCTGTATCGCCCTGTCTCTGGCCCTGGTCACAA</u> <u>ATAGCGCCCCCTACCAGCAGCAGCACCAAGAAAACACAGCTGCA</u> <u>ACTGGAACACCTCCTGCTGGACCTGCAGATGATCCTGAACGGC</u> <u>ATCAACAAC TACAAGAACCCCAAGCTGACCCGGATGCTGACCT</u> <u>TCAAGTTCTACATGCCCAAGAAGGCCACCGAGCTGAAGCACCT</u> <u>CCAGTGCTTGAAGAGGAAC TGAAGCCCC TGAAGAAGTGCTG</u> <u>AATCTGGCCCAGAGCAAGAACTTCCACCTGAGGCCTAGGGACC</u> <u>TGATCAGCAACATCAACGTGATCGTGCTGGAAC TGAAGGCAG</u> <u>CGAGACAACCTTCATGTGCGAGTACGCCGACGAGACAGCTACC</u> <u>ATCGTGGAATTTCTGAACCGGTGGATCACCTTCTGCCAGCA</u> <u>TCATCAGCACCCCTGACCTGAATAGTGAGTCGTATTACAGTACC</u> <u>AACAAGCAGAATCATCACGAAGTGGTACTTGACCACTTCGTGA</u> <u>TGATTCTGCTTTATCTTAGAGGCATATCCCTACGTACCAACAA</u> <u>GAGCTTCCTACAGCACAAACAACTTGTTGTTGTGCTGTAGGAA</u> <u>GCTCTTTATCTTAGAGGCATATCCCTACGTACCAACAAGATCC</u> <u>GCAGACGTGTAAATGTACTTGACATTTACACGTCTGCGGATCT</u> <u>TTATCTTAGAGGCATATCCCTTTTATCTTAGAGGCATATCCCT</u> <u>CTGGGCCTCATGGGCCTTCCTTTCACTGCCCCGCTTTCCAGTCG</u> <u>GGAAACCTGTGCGTGCCAGCTGCATTAACATGGTCATAGCTGTT</u> <u>TCCTTGCGTATTGGGCGCTCTCCGCTTCCTCGCTCACTGACTC</u> <u>GCTGCGCTCGGTCGTTTCGGGTAAAGCCTGGGGTGCCATAGAG</u> <u>CAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTT</u> <u>GCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCAC</u> <u>AAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGAC</u> <u>TATAAAGATAACAGGCGTTTTCCCCCTGGAAGCTCCCTCGTGCG</u> <u>CTCTCCTGTTCCGACCTGCCGCTTACCGGATACCTGTCCGCC</u> <u>TTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCATAGCTCACGCT</u> <u>GTAGGTATCTCAGTTCGGTGTAGGTCGTTTCGCTCCAAGCTGGG</u> <u>CTGTGTGCACGAACCCCCGTTTACGCCCCGACCGCTGCGCCTTA</u> <u>TCCGGTAAC TATCGTCTTGAGTCCAACCCGGTAAGACACGACT</u> <u>TATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGC</u> <u>GAGGTATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCT</u> <u>AACTACGGCTACACTAGAAGAACAGTATTTGGTATCTGCGCTC</u> <u>TGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTG</u> <u>ATCCGGCAACAAACCACCGCTGGTAGCGGTGGTTTTTTTTGTT</u> <u>TGCAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAG</u> <u>ATCCTTTGATCTTTTCTACGGGGTCTGACGCTCAGTGGAACGA</u> <u>AACTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGG</u> <u>ATCTTCACCTAGATCCTTTTAAATTAAAAATGAAGTTTTAAAT</u> <u>CAATCTAAAGTATATATGAGTAACTTGGTCTGACAGTTATTA</u> <u>GAAAAATTCATCCAGCAGACGATAAAACGCAATACGCTGGCTA</u>

SEQ ID NO	Compound	Sequence (5' to 3')
		TCCGGTGCCGCAATGCCATACAGCACCAGAAAACGATCCGCC ATTCCGCCGCCAGTTCTTCCGCAATATCACGGGTGGCCAGCGC AATATCCTGATAACGATCCGCCACGCCAGACGGCCGCAATCA ATAAAGCCGCTAAAACGGCCATTTTCCACCATAATGTTCCGGCA GGCACGCATCACCATGGGTACCACCAGATCTTCGCCATCCGG CATGCTCGCTTTCAGACGCGCAAACAGCTCTGCCGGTGCCAGG CCCTGATGTTCTTCATCCAGATCATCCTGATCCACCAGGCCCG CTTCATACGGGTACGCGCACGTTCAATACGATGTTTCGCCTG ATGATCAAACGGACAGGTCGCCGGGTCCAGGGTATGCAGACGA CGCATGGCATCCGCCATAATGCTCACTTTTTCTGCCGGCGCCA GATGGCTAGACAGCAGATCCTGACCCGGCACCTTCGCCCAGCAG CAGCCAATCACGGCCCGCTTCGGTCACCACATCCAGCACCCGCC GCACACGGAACACCGGTGGTGGCCAGCCAGCTCAGACGCGCCG CTTCATCCTGCAGCTCGTTCAGCGCACCGCTCAGATCGGTTTT CACAAACAGCACCGGACGACCCTGCGCGCTCAGACGAAACACC GCCGCATCAGAGCAGCCAATGGTCTGCTGCGCCCAATCATAGC CAAACAGACGTTCCACCCACGCTGCCGGGCTACCCGCATGCAG GCCATCCTGTTCAATCATACTCTTCCTTTTTCAATATTATTGA AGCATTTATCAGGGTTATTGTCTCATGAGCGGATACATATTTG AATGTATTTAGAAAAATAAACAAATAGGGGTTCCGCGCACATT TCCCGAAAAGTGCCAC
87	Compound 6 (pMA-RQ)	CTAAATTGTAAGCGTTAATATTTTGTTAAAATTCGCGTTAAAT TTTTGTTAAATCAGCTCATTTTTTAACCAATAGGCCGAAATCG GCAAAATCCCTTATAAATCAAAGAATAGACCGAGATAGGGTT GAGTGGCCGCTACAGGGCGCTCCCATTCGCCATTTCAGGCTGCG CAACTGTTGGGAAGGGCGTTTCGGTGCGGGCCTCTTCGCTATT ACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCGATTAAGT TGGGTAACGCCAGGGTTTTCCAGTCACGACGTTGTAAACGA CGGCCAGTGAGCGCGACGTAATACGACTCACTATAGGGCGAAT TGGCGGAAGGCCGTCAAGGCCGCATGCCACC <u>ATGTACAGAATG</u> <u>CAGCTGCTGAGCTGTATCGCCCTGTCTCTGGCCCTGGTTCACAA</u> <u>ATAGCGCCCTTACCAGCAGCAGCACCAAGAAAACACAGCTGCA</u> <u>ACTGGAACACCTCCTGCTGGACCTGCAGATGATCCTGAACGGC</u> ATCAACAAC TACAAGAACCCCAAGCTGACCCGGATGCTGACCT TCAAGTTCTACATGCCCAAGAAGGCCACCGAGCTGAAGCACCT CCAGTGCCCTGGAAGAGGAACTGAAGCCCCCTGGAAGAAGTGCTG AATCTGGCCCAGAGCAAGAACTTCCACCTGAGGCCTAGGGACC TGATCAGCAACATCAACGTGATCGTGCTGGAAC TGAAGGCAG CGAGACAACCTTCATGTGCGAGTACGCCGACGAGACAGCTACC ATCGTGGAATTTCTGAACCGGTGGATCACCTTCTGCCAGAGCA TCATCAGCACCTTGACCTGAATAGTGAGTCGTATTAACGTACC AACAAGGAGATTAGGGTCTGTGAGATACTTGATCTCACAGACC CTAATCTCCTTTATCTTAGAGGCATATCCCTACGTACCAACAA GATGCCATGAAGACCAAGACAAC TTGTGTCTTGGTCTTCATGG CATCTTTATCTTAGAGGCATATCCCTACGTACCAACAAGCCTG ATGGGAATGGAACCTAACTTGTAGGTTCCATTCCCATCAGGCT TTATCTTAGAGGCATATCCCTTTTATCTTAGAGGCATATCCCT CTGGGCCTCATGGGCCTTCCGCTCACTGCCCGCTTTCAGTCG GGAAACCTGTCTGTGCCAGCTGCATTAACATGGTCATAGCTGTT TCCTTGCGTATTGGGCGCTCTCCGCTTCCTCGCTCACTGACTC

SEQ ID NO	Compound	Sequence (5' to 3')
		GCTGCGCTCGGTCGTTTCGGGTAAAGCCTGGGGTGCCTAATGAG CAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTT GCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCAC AAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGAC TATAAAGATACCAGGCGTTTTCCCCCTGGAAGCTCCCTCGTGCG CTCTCCTGTTCCGACCCTGCCGCTTACCGGATACCTGTCCGCC TTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCATAGCTCACGCT GTAGGTATCTCAGTTCGGTGTAGGTCGTTTCGCTCCAAGCTGGG CTGTGTGCACGAACCCCCGTTTCAGCCCGACCGCTGCGCCTTA TCCGGTAACATATCGTCTTGAGTCCAACCCGGTAAGACACGACT TATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGC GAGGTATGTAGGCGGTGCTACAGAGTTCTTGAAAGTGGTGGCCT AACTACGGCTACACTAGAAGAACAGTATTTGGTATCTGCGCTC TGCTGAAGCCAGTTACCTTCGGAAGAGAGTTGGTAGCTCTTG ATCCGGCAAACAAACCACCGCTGGTAGCGGTGGTTTTTTTGT TGCAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAG ATCCTTTGATCTTTTCTACGGGGTCTGACGCTCAGTGAACGA AAACTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAG ATCTTCACCTAGATCCTTTTAAATTAAAAATGAAGTTTAAAT CAATCTAAAGTATATATGAGTAAACTTGGTCTGACAGTTACCA ATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTT CGTTCATCCATAGTTGCCTGACTCCCCGTGCTGTAGATAACTA CGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGAT ACCGCGAGAACCACGCTCACC GGCTCCAGATTTATCAGCAATA AACCAGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGTCTGCAA CTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCCGGAAGC TAGAGTAAGTAGTTCGCCAGTTAATAGTTTGC GCAACGTTGTT GCCATTGCTACAGGCATCGTGGTGTACGCTCGTCGTTTGGTA TGGCTTCATT CAGCTCCGGTTC CCAACGATCAAGGCGAGTTAC ATGATCCCCCATGTTGTGCAAAAAGCGGTTAGCTCCTTCGGT CCTCCGATCGTTGT CAGAAGTAAGTTGGCCG CAGTGTTATCAC TCATGGTTATGGCAGCACTGCATAATTCTCTTACTGT CATGCC ATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAG TCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGCC CGGCGTCAATACGGGATAATACCGCGCCACATAGCAGAACTTT AAAAGTGCTCATCATTGGAAAACGTTCTTCGGGGCGAAAAC TCAAGGATCTTACCGCTGTTGAGATCCAGTTTCGATGTAACCCA CTCGTGCACCCAAC TGATCTTCAGCATCTTTTACTTTTACCAG CGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAA AAGGGAATAAGGGCGACACGGAATGTTGAATACTCATACTCT TCCTTTTTTCAATATTATTGAAGCATTTATCAGGGTTATTGTCT CATGAGCGGATACATATTTGAATGTATTTAGAAAAATAAACAA ATAGGGGTTCCGCGCACATTTCCCCGAAAAGTGCCAC
88	Compound 7* (pMA-RQ)	CTAAATTGTAAGCGTTAATATTTTGTAAATTCGCGTTAAAT TTTTGTAAATCAGCTCATTTTTTAACCAATAGGCCGAAATCG GCAAAATCCCTTATAAATCAAAGAATAGACCGAGATAGGGTT GAGTGGCCGCTACAGGGCGCTCCCATTCGCCATT CAGGCTGCG CAACTGTTGGGAAGGGCGTTTCGGTGC GGGCCTCTTCGCTATT ACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCGATTAAAGT TGGGTAACGCCAGGGTTTTCCAGTCACGACGTTGTAAACGA

SEQ ID NO	Compound	Sequence (5' to 3')
		CGGCCAGTGAGCGCGACGTAATACGACTCACTATAGGGCGAAT TGGCGGAAGGCCGTCAAGGCCGCATGCCACCATGTTGTTGCTG CTGCTCGCCTGTATTGCCCTGGCCTCTACAGCCGCGCTACAA ATTCTGCCCCCTACCAGCAGCTCCACCAAGAAAACCCAGCTGCA ACTGGAACATCTGCTGCTGGACCTGCAGATGATCCTGAACGGC ATCAACAACACTACAAGAACCCCAAGCTGACCCGGATGCTGACCT TCAAGTTCTACATGCCCAAGAAGGCCACCGAGCTGAAGCACCT CCAGTGCCCTGGAAGAGGAAGTGAAGCCCCCTGGAAGAAGTGCTG AATCTGGCCCAGAGCAAGAAGTTCACCTGAGGCCTAGGGACC TGATCAGCAACATCAACGTGATCGTGCTGGAAC TGAAAGGCAG CGAGACAACCTTCATGTGCGAGTACGCCGACGAGACAGCTACC ATCGTGGAATTTCTGAACCGGTGGATCACCTTCTGCCAGAGCA TCATCAGCACCCCTGACCTGAATAGTGAGTCGTATTAACGTACC AACAAGCAGAATCATCACGAAGTGGTACTTGACCACTTCGTGA TGATTCTGCTTTATCTTAGAGGCATATCCCTACGTACCAACAA GAGCTTCCTACAGCACAACTTGTGTTGTGCTGTAGGAA GCTCTTTATCTTAGAGGCATATCCCTACGTACCAACAAGATCC GCAGACGTGTAAATGTACTTGACATTTACACGTGCGGATCT TTATCTTAGAGGCATATCCCTTTATCTTAGAGGCATATCCCT CTGGGCCTCATGGGCCTTCCGCTCACTGCCCGCTTTCAGTCG GGAAACCTGTCTGTGCCAGCTGCATTAACATGGTCATAGCTGTT TCCTTGCGTATTGGGCGCTCTCCGCTTCCCTCGCTCACTGACTC GCTGCGCTCGGTCTTTCGGGTAAAGCCTGGGGTGCCTAATGAG CAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTT GCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCAC AAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGAC TATAAAGATAACAGGCGTTTTCCCCCTGGAAGCTCCCTCGTGCG CTCTCCTGTTCCGACCCTGCCGCTTACCGGATACCTGTCCGCC TTTCTCCCTTCGGGAAGCGTGGCGCTTCTCATAGCTCACGCT GTAGGTATCTCAGTTCGGTGTAGGTGTTTCGCTCCAAGCTGGG CTGTGTGCACGAACCCCCCGTTTCAGCCCGACCGCTGCGCCTTA TCCGGTAACTATCGTCTTGAGTCCAACCCGGTAAGACACGACT TATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGC GAGGTATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCT AACTACGGCTACACTAGAAGAACAGTATTTGGTATCTGCGCTC TGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTG ATCCGGCAAACAACCAACCGCTGGTAGCGGTGGTTTTTTTTGTT TGCAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAG ATCCTTTGATCTTTTCTACGGGGTCTGACGCTCAGTGGAACGA AACTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGG ATCTTCACCTAGATCCTTTTAAATTAAAAATGAAGTTTAAAT CAATCTAAAGTATATATGAGTAACTTGGTCTGACAGTTACCA ATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTT CGTTCATCCATAGTTGCCTGACTCCCCGTCGTGTAGATAACTA CGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGAT ACCGCGAGAACCACGCTCACCGGCTCCAGATTTATCAGCAATA AACCAGCCAGCCGAAGGGCCGAGCGCAGAAGTGGTCTCTGCAA CTTTATCCGCTCCATCCAGTCTATTAATTGTTGCCGGGAAGC TAGAGTAAGTAGTTGCGCCAGTTAATAGTTTGGCGAACGTTGTT GCCATTGCTACAGGCATCGTGGTGTACGCTCGTCTGTTGGTA

SEQ ID NO	Compound	Sequence (5' to 3')
		TGGCTTCATTTCAGCTCCGGTTCCCAACGATCAAGGCGAGTTAC ATGATCCCCCATGTTGTGCAAAAAAGCGGTTAGCTCCTTCGGT CCTCCGATCGTTGTCAGAAGTAAGTTGGCCGAGTGTTATCAC TCATGGTTATGGCAGCACTGCATAATTCTCTTACTGTCATGCC ATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAG TCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGCC CGGCGTCAATACGGGATAATACCGCGCCACATAGCAGAACTTT AAAAGTGCTCATCATTGGAAAACGTTCTTCGGGGCGAAAACCTC TCAAGGATCTTACCGCTGTTGAGATCCAGTTCGATGTAACCCA CTCGTGCACCCAACCTGATCTTCAGCATCTTTTACTTTACCCAG CGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAA AAGGGAATAAGGGCGACACGGAAATGTTGAATACTCATACTCT TCCTTTTTCAATATTATTGAAGCATTTATCAGGGTTATTGTCT CATGAGCGGATACATATTTGAATGTATTTAGAAAAATAAACAA ATAGGGGTTCCGCGCACATTTCCCCGAAAAGTGCCAC
89	Compound 8* (pMA-RQ)	CTAAATTGTAAGCGTTAATATTTTGTAAATTCGCGTTAAAT TTTTGTAAATCAGCTCATTTTTTAACCAATAGGCCGAAATCG GCAAAATCCCTTATAAATCAAAAGAATAGACCGAGATAGGGTT GAGTGGCCGCTACAGGGCGCTCCCATTCGCCATTACGGCTGCG CAACTGTTGGGAAGGGCGTTTCGGTGCGGGCCTCTTCGCTATT ACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCGATTAAGT TGGGTAACGCCAGGGTTTTCCAGTCACGACGTTGTAAACGA CGGCCAGTGAGCGCGACGTAATACGACTCACTATAGGGCGAAT TGGCGGAAGGCCGTCAGGCCGCGAT <u>GCCACCATGTTGTTGCTG</u> <u>CTGCTCGCCTGTATTGCCCTGGCCTCTACAGCCGCCGCTACAA</u> <u>ATTCTGCCCCCTACCAGCAGCTCCACCAAGAAAACCCAGCTGCA</u> <u>ACTGGAACATCTGCTGCTGGACCTGCAGATGATCCTGAACGGC</u> <u>ATCAACAAC TACAAGAACCCCAAGCTGACCCGGATGCTGACCT</u> <u>TCAAGTTCTACATGCCCAAGAAGGCCACCGAGCTGAAGCACCT</u> <u>CCAGTGCCCTGGAAGAGGAAC TGAAGCCCCCTGGAAGAAGTGCTG</u> <u>AATCTGGCCCAGAGCAAGAAC TTCCACCTGAGGCCTAGGGACC</u> <u>TGATCAGCAACATCAACGTGATCGTGCTGGAAC TGAAGGCAG</u> <u>CGAGACAACCTTCATGTGCGAGTACGCCGACGAGACAGCTACC</u> <u>ATCGTGGAATTTCTGAACCGGTGGATCACCTTC TGCCAGAGCA</u> <u>TCATCAGCACCTTGACCTGAATAGTGAGTCGTATTAACGTACC</u> <u>AACAAGCAGAATCATCACGAAGTGGTACTTGACCACTTCGTGA</u> <u>TGATTCTGCTTTTATCTTAGAGGCATATCCCTACGTACCAACAA</u> <u>GAGCTTCCTACAGCACAACTTGTGTGTGCTGTAGGAA</u> <u>GCTCTTTATCTTAGAGGCATATCCCTACGTACCAACAAGATCC</u> <u>GCAGACGTGTAAATGTACTTGACATTTACACGTCTGCGGATCT</u> <u>TTATCTTAGAGGCATATCCCTACGTACCAACAAGCGCAAGAAA</u> <u>TCCCGGTATAAATTGTTATACCGGGATTTCTTGCGCTTTATC</u> <u>TTAGAGGCATATCCCTACGTACCAACAAGGCGAGGCAGCTTGA</u> <u>GTTAAAAC TTGTTTAACTCAAGCTGCCTCGCCTTTATCTTAGA</u> <u>GGCATATCCCTTTTATCTTAGAGGCATATCCCTCTGGGCCTCA</u> <u>TGGGCCTTCCGCTCACTGCCCCGCTTCCAGTCGGGAAACCTGT</u> <u>CGTGCCAGCTGCATTAACATGGTCATAGCTGTTTCCTTGCGTA</u> <u>TTGGGCGCTCTCCGCTTCCTCGCTCACTGACTCGCTGCGCTCG</u> <u>GTCGTTCCGGGTAAAGCCTGGGGTGCTAATGAGCAAAAGGCCA</u> <u>GCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTTGCTGGCGTTT</u>

SEQ ID NO	Compound	Sequence (5' to 3')
		<p>TTCCATAGGCTCCGCCCCCTGACGAGCATCACAAAAATCGAC GCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATA CCAGGCGTTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTT CCGACCCTGCCGCTTACCGGATACCTGTCCGCCTTTCTCCCTT CGGGAAGCGTGGCGCTTTCTCATAGCTCACGCTGTAGGTATCT CAGTTCGGTGTAGGTCTGTTGCTCCAAGCTGGGCTGTGTGCAC GAACCCCCGTTTACGCCCCGACGCTGCGCCTTATCCGGTAACT ATCGTCTTGAGTCCAACCCGGTAAGACACGACTTATCGCCACT GGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTA GGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCT ACACTAGAAGAACAGTATTTGGTATCTGCGCTCTGCTGAAGCC AGTTACCTTCGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAA CAAACACCGCTGGTAGCGGTGGTTTTTTTTGTTTGCAAGCAGC AGATTACGCGCAGAAAAAAGGATCTCAAGAAGATCCTTTGAT CTTTTCTACGGGTCTGACGCTCAGTGGAACGAAAACCTCACGT TAAGGGATTTTGGTCATGAGATTATCAAAAAGGATCTTCACCT AGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATCTAAAG TATATATGAGTAACTTGGTCTGACAGTTACCAATGCTTAATC AGTGAGGCACCTATCTCAGCGATCTGTCTATTTTCGTTTCATCCA TAGTTGCCTGACTCCCCGTCGTGTAGATAACTACGATACGGGA GGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGAA CCACGCTCACCGGCTCCAGATTTATCAGCAATAAACAGCCAG CCGGAAGGGCCGAGCGCAGAAGTGGTCCTGCAACTTTATCCGC CTCCATCCAGTCTATTAATTGTTGCCGGGAAGCTAGAGTAAGT AGTTCGCCAGTTAATAGTTTTCGCAACGTTGTTGCCATTGCTA CAGGCATCGTGGTGTACGCTCGTCGTTTGGTATGGCTTCATT CAGCTCCGGTTCCCAACGATCAAGGCGAGTTACATGATCCCCC ATGTTGTGCAAAAAGCGGTTAGCTCCTTCGGTCTCCGATCG TTGTCAGAAGTAAGTTGGCCGAGTGTTATCACTCATGGTTAT GGCAGCACTGCATAATTCTCTTACTGTCTATGCCATCCGTAAGA TGCTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAG AATAGTGTATGCGGCGACCGAGTTGCTCTTGCCCGGCGTCAAT ACGGGATAATACCGCGCCACATAGCAGAACTTTAAAAGTGCTC ATCATTGGAACGTTCTTTCGGGGCGAAAACCTCTCAAGGATCT TACCGCTGTTGAGATCCAGTTCGATGTAACCCACTCGTGCACC CAACTGATCTTCAGCATCTTTTACTTTTACCAGCGTTTCTGGG TGAGCAAAAACAGGAAGGCAAAATGCCGCAAAAAGGGAATAA GGGCGACACGGAATGTTGAATACTCATACTCTTCTTTTCA ATATTATTGAAGCATTTATCAGGGTTATTGTCTCATGAGCGGA TACATATTTGAATGTATTTAGAAAAATAAACAAATAGGGGTTT CGCGCACATTTCCCCGAAAAGTGCCAC</p>
90	Compound 9* (pMA-RQ)	<p>CTAAATTGTAAGCGTTAATATTTTGTAAATTCGCGTTAAAT TTTTGTAAATCAGCTCATTTTTTAACCAATAGGCCGAAATCG GCAAAATCCCTTATAAATCAAAAGAATAGACCGAGATAGGGTT GAGTGGCCGCTACAGGGCGCTCCCATTCGCCATTACAGGCTGCG CAACTGTTGGGAAGGGCGTTTCGGTGCGGGCCTCTTCGCTATT ACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCGATTAAGT TGGGTAACGCCAGGGTTTTCCAGTCACGACGTTGTAAACGA CGGCCAGTGAGCGCGACGTAATACGACTCACTATAGGGCGAAT TGAAGGAAGGCCGTCAAGGCCGCATGCCACCATGTTGTTGCTG</p>

SEQ ID NO	Compound	Sequence (5' to 3')
		<p> CTGCTCGCCTGTATTGCCCTGGCCTCTACAGCCGCCGCTACAA ATTCTGCCCCCTACCAGCAGCTCCACCAAGAAAACCCAGCTGCA ACTGGAACATCTGCTGCTGGACCTGCAGATGATCCTGAACGGC ATCAACAAC TACAAGAACCCCAAGCTGACCCGGATGCTGACCT TCAAGTTCTACATGCCCAAGAAGGCCACCGAGCTGAAGCACCT CCAGTGCCCTGGAAGAGGAACTGAAGCCCCCTGGAAGAAGTGCTG AATCTGGCCCAGAGCAAGAACTTCCACCTGAGGCCTAGGGACC TGATCAGCAACATCAACGTGATCGTGCTGGAAC TGAAGGCAG CGAGACAACCTTCATGTGCGAGTACGCCGACGAGACAGCTACC ATCGTGGAATTTCTGAACCGGTGGATCACCTTCTGCCAGAGCA TCATCAGCACCTGACCTGAATAGTGAGTCGTATTAACGTACC AACAAGGAGTACCCTGATGAGATCACTTGGATCTCATCAGGGT ACTCCTTTATCTTAGAGGCATATCCCTACGTACCAACAAGGAG TACCCTGATGAGATCACTTGGATCTCATCAGGGTACTCCTTTA TCTTAGAGGCATATCCCTACGTACCAACAAGGAGTACCCTGAT GAGATCACTTGGATCTCATCAGGGTACTCCTTTATCTTAGAGG CATATCCCTTTTATCTTAGAGGCATATCCCTCTGGGCCTCATG GGCCTTCCTTTCACTGCCCGCTTTCCAGTCGGGAAACCTGTGCG TGCCAGCTGCATTAACATGGTCATAGCTGTTTCTTGCCTATT GGGCGCTCTCCGCTTCTCGCTCACTGACTCGCTGCGCTCGGT CGTTCGGGTAAAGCCTGGGGTGCCTAATGAGCAAAAGGCCAGC AAAAGGCCAGGAACCGTAAAAAGGCCGCGTTGCTGGCGTTTTT CCATAGGCTCCGCCCCCTGACGAGCATCACAAAAATCGACGC TCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACC AGGCGTTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCC GACCCTGCCGCTTACCGGATACCTGTCCGCCTTTCTCCCTTCG GGAAGCGTGGCGCTTTCTCATAGCTCACGCTGTAGGTATCTCA GTTTCGGTGTAGGTGTTTCGCTCCAAGCTGGGCTGTGTGCACGA ACCCCCCGTTCAGCCCGACCGCTGCGCCTTATCCGGTAACATAT CGTCTTGAGTCCAACCCGGTAAGACACGACTTATCGCCACTGG CAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGG CGGTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTAC ACTAGAAGAACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAG TTACCTTCGGAAGAAAGAGTTGGTAGCTCTTGATCCGGCAAACA AACCACCGCTGGTAGCGGTGGTTTTTTTGTGTTGCAAGCAGCAG ATTACGCGCAGAAAAAAGGATCTCAAGAAGATCCTTTGATCT TTTCTACGGGGTCTGACGCTCAGTGGAACGAAAACCTCACGTTA AGGGATTTTGGTCATGAGATTATCAAAAAGGATCTTCACCTAG ATCCTTTTAAATTAATAATGAAGTTTTAAATCAATCTAAAGTA TATATGAGTAACTTGGTCTGACAGTTATTAGAAAAATTCATC CAGCAGACGATAAAACGCAATACGCTGGCTATCCGGTGCCGCA ATGCCATACAGCACCAGAAAACGATCCGCCCATTCGCCGCCCA GTTCTTCCGCAATATCACGGGTGGCCAGCGCAATATCCTGATA ACGATCCGCCACGCCAGACGGCCGCAATCAATAAAGCCGCTA AAACGGCCATTTTCCACCATAATGTTTCGGCAGGCACGCATCAC CATGGGTACACCACAGATCTTCGCCATCCGGCATGCTCGCTTT CAGACGCGCAAACAGCTCTGCCGGTGCCAGGCCCTGATGTTCT TCATCCAGATCATCCTGATCCACCAGGCCCGCTTCCATACGGG TACGCGCACGTTCAATACGATGTTTCGCCTGATGATCAAACGG ACAGGTCGCCGGGTCCAGGGTATGCAGACGACGCATGGCATCC </p>

SEQ ID NO	Compound	Sequence (5' to 3')
		GCCATAATGCTCACTTTTTCTGCCGGCGCCAGATGGCTAGACA GCAGATCCTGACCCGGCACTTCGCCCAGCAGCAGCCAATCACG GCCCCGTTTCGGTCACCACATCCAGCACCGCCGCACACGGAACA CCGGTGGTGGCCAGCCAGCTCAGACGCGCCGCTTCATCCTGCA GCTCGTTCAGCGCACCGCTCAGATCGGTTTTACAAACAGCAC CGGACGACCCTGCGCGCTCAGACGAAACACCGCCGCATCAGAG CAGCCAATGGTCTGCTGCGCCCAATCATAGCCAAACAGACGTT CCACCCACGCTGCCGGGCTACCCGCATGCAGGCCATCCTGTTT AATCATACTCTTCCTTTTTCAATATTATTGAAGCATTATCAG GGTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGA AAAATAAACAAATAGGGGTTCCGCGCACATTTCCCCGAAAAGT GCCAC
91	Compound 10* (pMA-RQ)	CTAAATTGTAAGCGTTAATATTTTTGTTAAAATTCGCGTTAAAT TTTTGTTAAATCAGCTCATTTTTTAACCAATAGGCCGAAATCG GCAAAATCCCTTATAAATCAAAAGAATAGACCGAGATAGGGTT GAGTGGCCGCTACAGGGCGCTCCCATTCGCCATTCAGGCTGCG CAACTGTTGGGAAGGGCGTTTTCGGTGCGGGCCTCTTCGCTATT ACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCGATTAAAT TGGGTAACGCCAGGGTTTTCCAGTCACGACGTTGTAAAACGA CGGCCAGTGAGCGCGACGTAATACGACTCATAATAGGCGCAAT TGAAGGAAGGCCGTCAAGGCCGCATGCCACCATGTTGTTGCTG CTGCTCGCCTGTATTGCCCTGGCCTCTACAGCCGCCGCTACAA ATTCTGCCCCCTACCAGCAGCTCCACCAAGAAAACCCAGCTGCA ACTGGAACATCTGCTGCTGGACCTGCAGATGATCCTGAACGGC ATCAACAACCTACAAGAACCCCAAGCTGACCCGGATGCTGACCT TCAAGTTCTACATGCCCAAGAAGGCCACCGAGCTGAAGCACCT CCAGTGCCTGGAAGAGGAACTGAAGCCCCCTGGAAGAAGTGCTG AATCTGGCCCAGAGCAAGAACCTCCACCTGAGGCCTAGGGACC TGATCAGCAACATCAACGTGATCGTGCTGGAACGTAAAGGCAG CGAGACAACCTTCATGTGCGAGTACGCCGACGAGACAGCTACC ATCGTGGAATTTCTGAACCGGTGGATCACCTTCTGCCAGAGCA TCATCAGCACCCCTGACCTGAATAGTGAGTCGTATTAACGTACC AACAAGGAGGGCAGAATCATCACGAAGTGGTGAAGTACTTGAC TTCACCACTTCGTGATGATTCTGCCCTCCTTTATCTTAGAGGC ATATCCCTACGTACCAACAAGAGATGAGCTTCTTACAGCACAA CAAATGTGACTTGCACATTTGTTGTGCTGTAGGAAGCTCATCT CTTTATCTTAGAGGCATATCCCTACGTACCAACAAGTACAAGA TCCGCAGACGTGTAAATGTTCCACTTGGGAACATTTACACGTC TGCGGATCTTGTACTTTATCTTAGAGGCATATCCCTTTTATCT TAGAGGCATATCCCTCTGGGCCTCATGGGCCTTCCTTTCACTG CCCGCTTTCCAGTCGGGAAACCTGTGCTGCCAGCTGCATTAAC ATGGTCATAGCTGTTTCCTTGCGTATTGGGCGCTCTCCGCTTC CTCGCTCACTGACTCGCTGCGCTCGGTGCTTCGGGTAAAGCCT GGGGTGCTAATGAGCAAAAGGCCAGCAAAAGGCCAGGAACCG TAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCC CCTGACGAGCATCACAAAATCGACGCTCAAGTCAGAGGTGGC GAAACCCGACAGGACTATAAAGATACCAGGCGTTTCCCCCTGG AAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCTGCCGCTTACC GGATACCTGTCCGCCTTTCTCCCTTCGGGAAGCGTGGCGCTTT CTCATAGCTCACGCTGTAGGTATCTCAGTTCGGTGTAGGTCGT

SEQ ID NO	Compound	Sequence (5' to 3')
		TCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCCGTTCAGCCC GACCGCTGCGCCTTATCCGGTAACTATCGTCTTGAGTCCAACC CGGTAAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAA CAGGATTAGCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTTC TTGAAGTGGTGGCCTAACTACGGCTACACTAGAAGAACAGTAT TTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAG AGTTGGTAGCTCTTGATCCGGCAAACAAACCACCGCTGGTAGC GGTGGTTTTTTTTGTTTGAAGCAGCAGATTACGCGCAGAAAAA AAGGATCTCAAGAAGATCCTTTGATCTTTTCTACGGGGTCTGA CGCTCAGTGAACGAAAACCTCACGTTAAGGGATTTTGGTCATG AGATTATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTAAA AATGAAGTTTTAAATCAATCTAAAGTATATATGAGTAAACTTG GTCTGACAGTTATTAGAAAAATTTCATCCAGCAGACGATAAAAC GCAATACGCTGGCTATCCGGTGCCGCAATGCCATACAGCACCA GAAAACGATCCGCCCATTTCGCCGCCAGTTCTTCGCAATATC ACGGGTGGCCAGCGCAATATCCTGATAACGATCCGCCACGCCC AGACGGCCGCAATCAATAAAGCCGCTAAAACGGCCATTTTCCA CCATAATGTTTCGGCAGGCACGCATCACCATGGGTCAACCACAG ATCTTCGCCATCCGGCATGCTCGCTTTCAGACGCGCAAACAGC TCTGCCGGTGCCAGGCCCTGATGTTCTTCATCCAGATCATCCT GATCCACCAGGCCCGCTTCCATACGGGTACGCGCACGTTCAAT ACGATGTTTCGCCTGATGATCAAACGGACAGGTGCGCGGGTCC AGGGTATGCAGACGACGCATGGCATCCGCCATAATGCTCACTT TTTCTGCCGGCGCCAGATGGCTAGACAGCAGATCCTGACCCGG CACTTCGCCCAGCAGCAGCCAATCACGGCCCGCTTCGGTCACC ACATCCAGCACCGCCGCACACGGAACACCGGTGGTGGCCAGCC AGCTCAGACGCGCCGCTTCATCCTGCAGCTCGTTCAGCGCAC GCTCAGATCGGTTTTTCAAAACAGCACCGGACGACCCTGCGCG CTCAGACGAAACACCGCCGCATCAGAGCAGCCAATGGTCTGCT GCGCCCAATCATAGCCAAACAGACGTTCCACCCACGCTGCCGG GCTACCCGCATGCAGGCCATCCTGTTCAATCATACTCTTCCTT TTTCAATATTATTGAAGCATTTATCAGGGTTATTGTCTCATGA GCGGATACATATTTGAATGTATTTAGAAAAATAAACAAATAGG GGTTCCGCGCACATTTCCCCGAAAAGTGCCAC
92	Compound 11 (pMA-RQ)	CTAAATTGTAAGCGTTAATATTTTGTTAAAATTTCGCGTTAAAT TTTTGTTAAATCAGCTCATTTTTTAACCAATAGGCCGAAATCG GCAAAATCCCTTATAAATCAAAGAATAGACCGAGATAGGGTT GAGTGGCCGCTACAGGGCGCTCCCATTCGCCATTACAGGCTGCG CAACTGTTGGGAAGGGCGTTTCGGTGCGGGCCTCTTCGCTATT ACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCGATTAAAGT TGGGTAACGCCAGGGTTTTCCAGTCACGACGTTGTAAACGA CGGCCAGTGAGCGCGACGTAATACGACTCACTATAGGGCGAAT TGGCGGAAGGCCGTCAGGCGCATGCCACCATGTGTCAACAG <u>CAGCTGGTCATCAGCTGGTTCAGCCTGGTGTCTCTGGCCTCTC</u> <u>CTCTGGTGGCCATCTGGGAGCTGAAGAAAGACGTGTACGTGGT</u> <u>GGAAGTGGACTGGTATCCCGATGCTCCTGGCGAGATGGTGGTG</u> CTGACCTGCGATACCCCTGAAGAGGACGGCATCACCTGGACAC TGGATCAGTCTAGCGAGGTGCTCGGCAGCGGCAAGACCCTGAC CATCCAAGTGAAAGAGTTTGGCGACGCCGGCCAGTACACCTGT CACAAAGGCGGAGAAGTGCTGAGCCACAGCCTGCTGCTGCTCC

SEQ ID NO	Compound	Sequence (5' to 3')
		ACAAGAAAGAGGATGGCATTGAGACACCGACATCCTGAAGGA CCAGAAAGAGCCCAAGAACAAGACCTTCCTGAGATGCGAGGCC AAGAACTACAGCGGCCGGTTACATGTTGGTGGCTGACCACCA TCAGCACCGACCTGACCTTCAGCGTGAAGTCCAGCAGAGGCAG CAGTGATCCTCAGGGCGTTACATGTGGCGCCGCTACACTGTCT GCCGAAAGAGTGCAGGGCGACAACAAGAATACGAGTACAGCG TGGAAATGCCAAGAGGACAGCGCCTGTCCAGCCGCCGAAGAGTC TCTGCCTATCGAAGTGATGGTGGACGCCGTGCACAAGCTGAAG TACGAGAACTACACCTCCAGCTTTTTTCATCCGGGACATCATCA AGCCCGATCCTCCAAAGAACCTGCAGCTGAAGCCTCTGAAGAA CAGCAGACAGGTGGAAGTGTCTGGGAGTACCCCGACACCTGG TCTACACCCACAGCTACTTCAGCCTGACCTTTTGCGTGCAAG TGCAGGGCAAGTCCAAGCGCGAGAAAAAGGACCGGGTGTTCAC CGACAAGACCAGCGCCACCGTGATCTGCAGAAAGAACGCCAGC ATCAGCGTCAGAGCCCAGGACCGGTACTACAGCAGCTCTTGGA GCGAATGGGCCAGCGTGCCATGTTCTGGTGGCGGAGGATCTGG CGGAGGTGGAAGCGCGGAGGCGGATCTAGAAATCTGCCTGTG GCCACTCCTGATCCTGGCATGTTCCCTTGCTGACACACAGCC AGAACCTGCTGAGAGCCGTGTCCAACATGCTGCAGAGGCCAG ACAGACCCTGGAATTCTACCCCTGCACCAGCAGGAAATCGAC CACGAGGACATCACCAAGGATAAGACCAGCACCGTGGAAGCCT GCCTGCCTCTGGAAGTACCAAGAACGAGAGCTGCCTGAACAG CCGGGAACACAGCTTCATCACCAACGGCTCTTGCTGGCCAGC AGAAAGACCTCCTTCATGATGGCCCTGTGCCTGAGCAGCATCT ACGAGGACCTGAAGATGTACCAGGTGGAATTCAAGACCATGAA CGCCAAGCTGCTGATGGACCCCAAGCGGCAGATCTTCTGGAC CAGAATATGCTGGCCGTGATCGACGAGCTGATGCAGGCCCTGA ACTTCAACAGCGAGACAGTGCCCCAGAAGTCTAGCCTGGAAGA ACCCGACTTCTACAAGACCAAGATCAAGCTGTGCATCCTGCTG CACGCCTTCCGGATCAGAGCCGTGACCATCGACAGAGTGATGA GCTACCTGAACGCCTCCTGAATAGTGAGTCGTATTAACGTACC AACAAGTTCTTTCCAAATGGCTCTGTACTTGACAGAGCCATTT GGAAGGAACTTTATCTTAGAGGCATATCCCTACGTACCAACAA GCATCGTTCACCGAGATCTGAACTTGTGAGATCTCGGTGAACG ATGCTTTATCTTAGAGGCATATCCCTACGTACCAACAAGACCA GCAGCGGACAAATAAACTTGTTTATTTGTCCGCTGCTGGTCT TTATCTTAGAGGCATATCCCTTTTATCTTAGAGGCATATCCCT CTGGGCCTCATGGGCCTTCCGCTCACTGCCCGCTTTCCAGTCG GGAAACCTGTGCGTCCAGCTGCATTAACATGGTCATAGCTGTT TCCTTGCGTATTGGGCGCTCTCCGCTTCCGCTCACTGACTC GCTGCGCTCGGTCGTTCCGGTAAAGCCTGGGGTGCCATATGAG CAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTT GCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCAC AAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGAC TATAAAGATAACAGGCGTTTTCCCCCTGGAAGCTCCCTCGTGCG CTCTCCTGTTCCGACCCTGCCGCTTACCGGATACCTGTCCGCC TTTCTCCCTTCGGAAGCGTGGCGCTTTCTCATAGCTCACGCT GTAGGTATCTCAGTTCCGGTGTAGGTCGTTCCGCTCCAAGCTGGG CTGTGTGCACGAACCCCCCGTTCCAGCCCGACCGCTGCGCCTTA TCCGGTAACTATCGTCTTGAGTCCAACCCGGTAAGACACGACT

SEQ ID NO	Compound	Sequence (5' to 3')
		TATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGC GAGGTATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCT AACTACGGCTACACTAGAAGAACAGTATTTGGTATCTGCGCTC TGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTG ATCCGGCAAACAAACCACCGCTGGTAGCGGTGGTTTTTTTGT TGCAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAG ATCCTTTGATCTTTTCTACGGGGTCTGACGCTCAGTGGAACGA AACTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGG ATCTTCACCTAGATCCTTTTAAATTAAAAATGAAGTTTTAAAT CAATCTAAAGTATATATGAGTAACTTGGTCTGACAGTTACCA ATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTT CGTTTCATCCATAGTTGCCTGACTCCCCGTCGTGTAGATAACTA CGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGAT ACCGCGAGAACCACGCTCACC GGCTCCAGATTTATCAGCAATA AACCAGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGTCTTGCAA CTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCCGGGAAGC TAGAGTAAGTAGTTCGCCAGTTAATAGTTTGC GCAACGTTGTT GCCATTGCTACAGGCATCGTGGTGT CACGCTCGTCGTTTGGTA TGGCTTCATT CAGCTCCGGTTCCCAACGATCAAGGCGAGTTAC ATGATCCCCCATGTTGTGCAAAAAAGCGGTTAGCTCCTTCGGT CCTCCGATCGTTGT CAGAAGTAAGTTGGCCGAGTGTTATCAC TCATGGTTATGGCAGCACTGCATAATTCTCTTACTGT CATGCC ATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAG TCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGCC CGGCGTCAATACGGGATAATACCGCGCCACATAGCAGAACTTT AAAAGTGCTCATCATTGGAAAACGTTCTTCGGGGCGAAAAC TC TCAAGGATCTTACCGCTGTTGAGATCCAGTTTCGATGTAACCCA CTCGTGCACCCAAC TGATCTTCAGCATCTTTTACTTTTACCAG CGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAA AAGGGAATAAGGGCGACACGGAAATGTTGAATACTCATACTCT TCCTTTTTTCAATATTATTGAAGCATTTATCAGGGTTATTGTCT CATGAGCGGATACATATTTGAATGTATTTAGAAAAATAAACAA ATAGGGGTTCCGCGCACATTTCCCCGAAAAGTGCCAC
93	Compound 12 (pMA-RQ)	CTAAATTGTAAGCGTTAATATTTTGT TAAAATTTCGCGTTAAAT TTTTGT TAAATCAGCTCATTTTTTTAACCAATAGGCCGAAATCG GCAAAATCCCTTATAAATCAAAAGAATAGACCGAGATAGGGTT GAGTGGCCGCTACAGGGCGCTCCCATTCGCCATT CAGGCTGCG CAACTGTTGGGAAGGGCGTTTCGGTGCGGGCCTCTTCGCTATT ACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCGATTAAAGT TGGGTAAACGCCAGGGTTTTCCAGTCACGACGTTGTAAAACGA CGGCCAGTGAGCGCGACGTAATACGACTCACTATAGGGCGAAT TGGCGGAAGGCCGTC AAGGCCGAT <u>GCCACCATGTGT CACCAG</u> <u>CAGCTGGT CATCAGCTGGTTCAGCCTGGTGTTCCTGGCCTCTC</u> <u>CTCTGGTGGCCATCTGGGAGCTGAAGAAAGACGTGTACGTGGT</u> <u>GGA ACTGGACTGGTATCCCGATGCTCCTGGCGAGATGGTGGTG</u> <u>CTGACCTGCGATAACCCCTGAAGAGGACGGCATCACCTGGACAC</u> <u>TGGATCAGTCTAGCGAGGTGCTCGGCAGCGGCAAGACCCTGAC</u> <u>CATCCAAGTGAAAGAGTTTGGCGACGCCGGCCAGTACACCTGT</u> <u>CACAAAGGCGGAGAAGTGCTGAGCCACAGCCTGCTGCTGCTCC</u> <u>ACAAGAAAGAGGATGGCATTTGGAGCACCGACATCCTGAAGGA</u>

SEQ ID NO	Compound	Sequence (5' to 3')
		CCAGAAAGAGCCCAAGAACAAGACCTTCCTGAGATGCGAGGCC AAGAACTACAGCGGCCGGTTACATGTTGGTGGCTGACCACCA TCAGCACCGACCTGACCTTCAGCGTGAAGTCCAGCAGAGGCAG CAGTGATCCTCAGGGCGTTACATGTGGCGCCGCTACACTGTCT GCCGAAAGAGTGCAGGGCGACAACAAGAATACGAGTACAGCG TGGAATGCCAAGAGGACAGCGCCTGTCCAGCCGCCGAAGAGTC TCTGCCTATCGAAGTGATGGTGGACGCCGTGCACAAGCTGAAG TACGAGAACTACACCTCCAGCTTTTTTCATCCGGGACATCATCA AGCCCGATCCTCCAAAGAACCCTGCAGCTGAAGCCTCTGAAGAA CAGCAGACAGGTGGAAGTGTCTTGGGAGTACCCCGACACCTGG TCTACACCCACAGCTACTTCAGCCTGACCTTTTGCCTGCAAG TGCAGGGCAAGTCCAAGCGCGAGAAAAGGACCGGGTGTTCAC CGACAAGACCAGCGCCACCGTGATCTGCAGAAAAGACGCCAGC ATCAGCGTCAGAGCCCAGGACCGGTACTACAGCAGCTCTTGGA GCGAATGGGCCAGCGTGCCATGTTCTGGTGGCGGAGGATCTGG CGGAGGTGGAAGCGCGGAGGCGGATCTAGAAATCTGCCTGTG GCCACTCCTGATCCTGGCATGTTCCCTTGTCTGCACCACAGCC AGAACCTGCTGAGAGCCGTGTCCAACATGCTGCAGAAGGCCAG ACAGACCCTGGAATTCTACCCCTGCACCAGCAGGAAATCGAC CACGAGGACATACCAAGGATAAGACCAGCACCCTGGAAGCCCT GCCTGCCTCTGGAAGTACCAAGAACGAGAGCTGCCTGAACAG CCGGGAAACCAGCTTCATCACCAACGGCTCTTGCCCTGGCCAGC AGAAAGACCTCCTTCATGATGGCCCTGTGCCTGAGCAGCATCT ACGAGGACCTGAAGATGTACCAGGTGGAATTCAAGACCATGAA CGCCAAGCTGCTGATGGACCCCAAGCGGCAGATCTTCTGGAC CAGAATATGCTGGCCGTGATCGACGAGCTGATGCAGGCCCTGA ACTTCAACAGCGAGACAGTGCCCCAGAAGTCTAGCCTGGAAGA ACCCGACTTCTACAAGACCAAGATCAAGCTGTGCATCCTGCTG CACGCCTTCCGGATCAGAGCCGTGACCATCGACAGAGTGATGA GCTACCTGAACGCCTCCTGAATAGTGAGTCGTATTAACGTACC AACAAGAAGGAGCTGCCCATGAGAAAACCTTGTTTTCTCATGGGC AGCTCCTTCTTTATCTTAGAGGCATATCCCTACGTACCAACAA GTGCAATGAGGGACCAGTACAACCTTGTGTACTGGTCCCTCATT GCACTTTATCTTAGAGGCATATCCCTACGTACCAACAAGAGCT GCTGAAGGACTCATCAACTTGTGATGAGTCCTTCAGCAGCTCT TTATCTTAGAGGCATATCCCTTTTATCTTAGAGGCATATCCCT CTGGGCCTCATGGGCCTTCCGCTCACTGCCCCGCTTTCCAGTCG GGAAACCTGTCGTGCCAGCTGCATTAACATGGTCATAGCTGTT TCCTTGCGTATTGGGCGCTCTCCGCTTCCTCGCTCACTGACTC GCTGCGCTCGGTCGTTCCGGTAAAGCCTGGGGTGCCATATGAG CAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTT GCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCAC AAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGAC TATAAAGATAACAGGCGTTTCCCCCTGGAAGCTCCCTCGTGCG CTCTCCTGTTCCGACCCTGCCGCTTACCGGATACCTGTCCGCC TTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCATAGCTCACGCT GTAGGTATCTCAGTTCGGTGTAGGTGCTTCGCTCCAAGCTGGG CTGTGTGCACGAACCCCCGTTCCAGCCCGACCGCTGCGCCTTA TCCGGTAACTATCGTCTTGAGTCCAACCCGGTAAGACACGACT TATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGC

SEQ ID NO	Compound	Sequence (5' to 3')
		GAGGTATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCT AACTACGGCTACACTAGAAGAACAGTATTTGGTATCTGCGCTC TGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTG ATCCGGCAAACAAACCACCGCTGGTAGCGGTGGTTTTTTTGT TGCAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAG ATCCTTTGATCTTTTCTACGGGGTCTGACGCTCAGTGAACGA AACTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGG ATCTTCACCTAGATCCTTTTAAATTAAAAATGAAGTTTTAAAT CAATCTAAAGTATATATGAGTAACTTGGTCTGACAGTTACCA ATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTT CGTTCATCCATAGTTGCCTGACTCCCCGTCGTGTAGATAACTA CGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGAT ACCGCGAGAACCACGCTCACC GGCTCCAGATTTATCAGCAATA AACCAGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGTCTGCAA CTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCCGGGAAGC TAGAGTAAGTAGTTCGCCAGTTAATAGTTTGC GCAACGTTGTT GCCATTGCTACAGGCATCGTGGTGT CACGCTCGTCGTTTGGTA TGGCTTCATTCAGCTCCGGTTCCCAACGATCAAGGCGAGTTAC ATGATCCCCCATGTTGTGCAAAAAAGCGGTTAGCTCCTTCGGT CCTCCGATCGTTGTGCAAGTAAGTTGGCCGCACTGTTATCAC TCATGGTTATGGCAGCACTGCATAATTCTCTTACTGT CATGCC ATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAG TCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGCC CGGCGTCAATACGGGATAATACCGCGCCACATAGCAGAACTTT AAAAGTGCTCATCATTGGAACGTTCTTCGGGGCGAAACTC TCAAGGATCTTACCGCTGTTGAGATCCAGTTCGATGTAACCCA CTCGTGCACCCAACTGATCTTCAGCATCTTTTACTTT CACCAG CGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAA AAGGGAATAAGGGCGACACGGAAATGTTGAATACTCATACTCT TCCTTTTTCAATATTATTGAAGCATTTATCAGGGTTATTGTCT CATGAGCGGATACATATTTGAATGTATTTAGAAAAATAAACAA ATAGGGGTTCCGCGCACATTTCCCCGAAAAGTGCCAC
94	Compound 13 (pMA-RQ)	CTAAATTGTAAGCGTTAATATTTTGT TAAAATTCGCGTTAAAT TTTTGT TAAATCAGCTCATTTTTTTAACCAATAGGCCGAAATCG GCAAAATCCCTTATAAATCAAAAGAATAGACCGAGATAGGGTT GAGTGGCCGCTACAGGGCGCTCCCATT CGCCATTCAGGCTGCG CAACTGTTGGGAAGGGCGTTTCGGTGCGGGCCTCTTCGCTATT ACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCGATTAAAGT TGGGTAACGCCAGGGTTTTCCAGTCACGACGTTGTAAAACGA CGGCCAGTGAGCGCGACGTAATACGACTCACTATAGGGCGAAT TGGCGGAAGGCCGTC AAGGCCGAT <u>GCCACCATGTGTCACCAG</u> <u>CAGCTGGTCA</u> <u>TACAGCTGGTT</u> <u>CAGCCTGGTGTTCCTGGCCTCTC</u> <u>CTCTGGTGGCCATCTGGGAGCTGAAGAAAGACGTGTACGTGGT</u> <u>GGA</u> <u>ACTGGACTGGTATCCCGATGCTCCTGGCGAGATGGTGGTG</u> <u>CTGACCTGCGATA</u> <u>CCCC</u> <u>TGAAGAGGACGGCATCACCTGGACAC</u> <u>TGGATCAGTCTAGCGAGGTGCTCGGCAGCGGCAAGACCCTGAC</u> <u>CATCCAAGTGAAAGAGTTTGGCGACGCCGCCAGTACACCTGT</u> <u>CACAAAGGCGGAGAAGTGCTGAGCCACAGCCTGCTGCTGCTCC</u> <u>ACAAGAAAGAGGATGGCATT</u> <u>TGGAGCACCGACATCCTGAAGGA</u> <u>CCAGAAAGAGCCCAAGAACAAGACCTTCCTGAGATGCGAGGCC</u>

SEQ ID NO	Compound	Sequence (5' to 3')
		AAGAACTACAGCGGCCGGTTCACATGTTGGTGGCTGACCACCA TCAGCACCGACCTGACCTTCAGCGTGAAGTCCAGCAGAGGCAG CAGTGATCCTCAGGGCGTTACATGTGGCGCCGCTACACTGTCT GCCGAAAGAGTGCAGGGGCGACAACAAAGAATACGAGTACAGCG TGGAATGCCAAGAGGACAGCGCCTGTCCAGCCGCCGAAGAGTC TCTGCCTATCGAAGTGATGGTGGACGCCGTGCACAAGCTGAAG TACGAGAACTACACCTCCAGCTTTTTTCATCCGGGACATCATCA AGCCCGATCCTCCAAAGAACCTGCAGCTGAAGCCTCTGAAGAA CAGCAGACAGGTGGAAGTGTCTTGGGAGTACCCCGACACCTGG TCTACACCCACAGCTACTTCAGCCTGACCTTTTGCCTGCAAG TGCAGGGCAAGTCCAAGCGCGAGAAAAAGGACCGGGTGTTCAC CGACAAGACCAGCGCCACCGTGATCTGCAGAAAGAACGCCAGC ATCAGCGTCAGAGCCCAGGACCGGTACTACAGCAGCTCTTGGA GCGAATGGGCCAGCGTGCCATGTTCTGGTGGCGGAGGATCTGG CGGAGGTGGAAGCGCGGAGGCGGATCTAGAAATCTGCCTGTG GCCACTCCTGATCCTGGCATGTTCCCTTGTCTGCACCACAGCC AGAACCTGCTGAGAGCCGTGTCCAACATGCTGCAGAAGGCCAG ACAGACCCTGGAATTCTACCCCTGCACCAGCGAGGAAATCGAC CACGAGGACATCACCAGGATAAGACCAGCACCCTGGAAGCCT GCCTGCCTCTGGAAGTACCAAGAACGAGAGCTGCCTGAACAG CCGGGAAACCAGCTTCATCACCACGGCTCTTGCCCTGGCCAGC AGAAAGACCTCCTTCATGATGGCCCTGTGCCTGAGCAGCATCT ACGAGGACCTGAAGATGTACCAGGTGGAATTCAAGACCATGAA CGCCAAGCTGCTGATGGACCCCAAGCGGCAGATCTTCCTGGAC CAGAATATGCTGGCCGTGATCGACGAGCTGATGCAGGCCCTGA ACTTCAACAGCGAGACAGTGCCCCAGAAGTCTAGCCTGGAAGA ACCCGACTTCTACAAGACCAAGATCAAGCTGTGCATCCTGCTG CACGCCTTCCGGATCAGAGCCGTGACCATCGACAGAGTGATGA GCTACCTGAACGCCTCCTGAATAGTGAGTCGTATTAACGTACC AACAAGAAGGAGCTGCCCATGAGAAAACCTTGTTTCTCATGGGC AGCTCCTTCTTTATCTTAGAGGCATATCCCTACGTACCAACAA GTCCAACGAATGGGCCTAAGAACTTGTCTTAGGCCCATTCGTT GGACTTTATCTTAGAGGCATATCCCTACGTACCAACAAGGACA GCATAGACGACACCTTACTTGAAGGTGTCGTCTATGCTGTCTT TTATCTTAGAGGCATATCCCTTTTATCTTAGAGGCATATCCCT CTGGGCCTCATGGGCCTTCCGCTCACTGCCCCGCTTTCCAGTCG GGAAACCTGTCGTGCCAGCTGCATTAACATGGTCATAGCTGTT TCCTTGCGTATTGGGCGCTCTCCGCTTCCCTCGCTCACTGACTC GCTGCGCTCGGTGCTTCGGGTAAAGCCTGGGGTGCCTAATGAG CAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTT GCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCAC AAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGAC TATAAAGATACCAGGCGTTTTCCCCCTGGAAGCTCCCTCGTGCG CTCTCCTGTTCCGACCCTGCCGCTTACCGGATACCTGTCCGCC TTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCATAGCTCAGCT GTAGGTATCTCAGTTCGGTGTAGGTGCTTCGCTCCAAGCTGGG CTGTGTGCACGAACCCCCGTTCCAGCCCGACCGCTGCGCCTTA TCCGGTAACATATCGTCTTGAGTCCAACCCGGTAAGACACGACT TATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGC GAGGTATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCT

SEQ ID NO	Compound	Sequence (5' to 3')
		A A C T A C G G C T A C A C T A G A A G A A C A G T A T T T G G T A T C T G C G C T C T G C T G A A G C C A G T T A C C T T C G G A A A A G A G T T G G T A G C T C T T G A T C C G G C A A A C A A A C C A C C G C T G G T A G C G G T G G T T T T T T G T T T G C A A G C A G C A G A T T A C G C G C A G A A A A A A G G A T C T C A A G A A G A T C C T T T G A T C T T T T T A C G G G G T C T G A C G C T C A G T G G A A C G A A A A C T C A C G T T A A G G G A T T T T G G T C A T G A G A T T A T C A A A A A G G A T C T T C A C C T A G A T C C T T T T A A A T T A A A A T G A A G T T T T A A A T C A A T C T A A A G T A T A T A T G A G T A A A C T T G G T C T G A C A G T T A C C A A T G C T T A A T C A G T G A G G C A C C T A T C T C A G C G A T C T G T C T A T T T C G T T C A T C C A T A G T T G C C T G A C T C C C C G T C G T G T A G A T A A C T A C G A T A C G G G A G G G C T T A C C A T C T G G C C C C A G T G C T G C A A T G A T A C C G C G A G A A C C A C G C T C A C C G G C T C C A G A T T T A T C A G C A A T A A A C C A G C C A G C C G G A A G G G C C G A G C G C A G A A G T G G T C C T G C A A C T T T A T C C G C C T C C A T C C A G T C T A T T A A T T G T T G C C G G G A A G C T A G A G T A A G T A G T T C G C C A G T T A A T A G T T T G C G C A A C G T T G T T G C C A T T G C T A C A G G C A T C G T G G T G T C A C G C T C G T C G T T T G G T A T G G C T T C A T T C A G C T C C G G T T C C C A A C G A T C A A G G C G A G T T A C A T G A T C C C C C A T G T T G T G C A A A A A A G C G G T T A G C T C C T T C G G T C C T C C G A T C G T T G T C A G A A G T A A G T T G G C C G C A G T G T T A T C A C T C A T G G T T A T G G C A G C A C T G C A T A A T T C T C T T A C T G A T C A G C C A T C C G T A A G A T G C T T T T C T G T G A C T G G T G A G T A C T C A A C C A A G T C A T T C T G A G A A T A G T G T A T G C G G C G A C C G A G T T G C T C T T G C C C G G C G T C A A T A C G G G A T A A T A C C G C G C C A C A T A G C A G A A C T T T A A A A G T G C T C A T C A T T G G A A A C G T T C T T C G G G G C G A A A A C T C T C A A G G A T C T T A C C G C T G T T G A G A T C C A G T T C G A T G T A A C C C A C T C G T G C A C C C A A C T G A T C T T C A G C A T C T T T T A C T T T C A C C A G C G T T T C T G G G T G A G C A A A A C A G G A A G G C A A A A T G C C G C A A A A A A G G G A A T A A G G G C G A C A C G G A A A T G T T G A A T A C T C A T A C T C T T C C T T T T T C A A T A T T A T T G A A G C A T T T A T C A G G G T T A T T G T C T C A T G A G C G G A T A C A T A T T T G A A T G T A T T T A G A A A A T A A A C A A A T A G G G G T T C C G C G C A C A T T T C C C C G A A A A G T G C C A C
95	Compound 14 (pMA-RQ)	C T A A A T T G T A A G C G T T A A T A T T T T G T T A A A A T T C G C G T T A A A T T T T T G T T A A A T C A G C T C A T T T T T T A A C C A A T A G G C C G A A A T C G G C A A A A T C C C T T A T A A A T C A A A A G A A T A G A C C G A G A T A G G G T T G A G T G G C C G C T A C A G G G C G C T C C C A T T C G C C A T T C A G G C T G C G C A A C T G T T G G G A A G G G C G T T T C G G T G C G G G C C T C T T C G C T A T T A C G C C A G C T G G C G A A A G G G G A T G T G C T G C A A G G C G A T T A A G T T G G G T A A C G C C A G G G T T T T C C A G T C A C G A C G T T G T A A A A C G A C G G C C A G T G A G C G C G A C G T A A T A C G A C T C A C T A T A G G G C G A A T T G G C G G A A G G C C G T C A A G G C C G C A T <u>G C C A C C A T G T G T C A C C A G</u> <u>C A G C T G G T C A T C A G C T G G T T C A G C C T G G T G T T C C T G G C C T C T C</u> <u>C T C T G G T G G C C A T C T G G G A G C T G A A G A A A G A C G T G T A C G T G G T</u> <u>G G A A C T G G A C T G G T A T C C C G A T G C T C C T G G C G A G A T G G T G G T G</u> <u>C T G A C C T G C G A T A C C C T G A A G A G G A C G G C A T C A C C T G G A C A C</u> <u>T G G A T C A G T C T A G C G A G G T G C T C G G C A G C G G C A A G A C C C T G A C</u> <u>C A T C C A A G T G A A A G A G T T T G G C G A C G C C G G C C A G T A C A C C T G T</u> <u>C A C A A A G G C G G A G A A G T G C T G A G C C A C A G C C T G C T G C T G C T C C</u> <u>A C A A G A A A G A G G A T G G C A T T T G G A G C A C C G A C A T C C T G A A G G A</u> <u>C C A G A A A G A G C C C A A G A A C A A G A C C T T C C T G A G A T G C G A G G C C</u> <u>A A G A A C T A C A G C G G C C G G T T C A C A T G T T G G T G G C T G A C C A C C A</u>

SEQ ID NO	Compound	Sequence (5' to 3')
		TCAGCACCGACCTGACCTTCAGCGTGAAGTCCAGCAGAGGCAG CAGTGATCCTCAGGGCGTTACATGTGGCGCCGCTACACTGTCT GCCGAAAGAGTGCGGGGCGACAACAAAGAATACGAGTACAGCG TGGAATGCCAAGAGGACAGCGCCTGTCCAGCCGCCGAAGAGTC TCTGCCTATCGAAGTGATGGTGGACGCCGTGCACAAGCTGAAG TACGAGAACTACACCTCCAGCTTTTTTCATCCGGGACATCATCA AGCCCGATCCTCCAAAGAACCTGCAGCTGAAGCCTCTGAAGAA CAGCAGACAGGTGGAAGTGTCTGGGAGTACCCCGACACCTGG TCTACACCCACAGCTACTTCAGCCTGACCTTTTGCGTGCAAG TGCAGGGCAAGTCCAAGCGCGAGAAAAAGGACCGGGTGTTCAC CGACAAGACCAGCGCCACCGTGATCTGCAGAAAGAACGCCAGC ATCAGCGTCAGAGCCCAGGACCGGTACTACAGCAGCTCTTGGA GCGAATGGGCCAGCGTGCCATGTTCTGGTGGCGGAGGATCTGG CGGAGGTGGAAGCGGCGGAGGCGGATCTAGAAATCTGCCTGTG GCCACTCCTGATCCTGGCATGTTCCCTTGTCTGCACCACAGCC AGAACCTGCTGAGAGCCGTGTCCAACATGCTGCAGAAGGCCAG ACAGACCCTGGAATTCTACCCCTGCACCAGCGAGGAAATCGAC CACGAGGACATCACCAAGGATAAGACCAGCACCGTGGAAGCCT GCCTGCCTCTGGAAGTACCAAGAACGAGAGCTGCCTGGAACAG CCGGGAAACAGCTTCATCACCAACGGCTCTTGCCCTGGCCAGC AGAAAGACCTCCTTCATGATGGCCCTGTGCCTGAGCAGCATCT ACGAGGACCTGAAGATGTACCAGGTGGAATTCAAGACCATGAA CGCCAAGCTGCTGATGGACCCCAAGCGGCAGATCTTCTTGAC CAGAATATGCTGGCCGTGATCGACGAGCTGATGCAGGCCCTGA ACTTCAACAGCGAGACAGTGCCCCAGAAGTCTAGCCTGGAAGA ACCCGACTTCTACAAGACCAAGATCAAGCTGTGCATCCTGCTG CACGCCTTCCGGATCAGAGCCGTGACCATCGACAGAGTGATGA GCTACCTGAACGCCTCCTGAATAGTGAGTCGTATTAACGTACC AACAAGACCCTGACATTGCTACTGTACTTGACAGTAGCGAAT GTCAGGGTCTTTATCTTAGAGGCATATCCCTACGTACCAACAA GAGCTGCTGAAGGACTCATCAACTTGTGATGAGTCCTTCAGCA GCTCTTTATCTTAGAGGCATATCCCTACGTACCAACAAGGCCA ATGACCCAAACATCTCTACTTGAGAGATGTTGGGTCATTGGCCT TTATCTTAGAGGCATATCCCTTTTATCTTAGAGGCATATCCCT CTGGGCCTCATGGGCCTTCCGCTCACTGCCCCGCTTTCCAGTCG GGAAACCTGTCTGTGCCAGCTGCATTAACATGGTCATAGCTGTT TCCTTGCGTATTGGGCGCTCTCCGCTTCCCTCGCTCACTGACTC GCTGCGCTCGGTGCTTCGGGTAAAGCCTGGGGTGCCCTAATGAG CAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTT GCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCAC AAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGAC TATAAAGATACCAGGCGTTTCCCCCTGGAAGCTCCCTCGTGCG CTCTCCTGTTCCGACCCTGCCGCTTACCGGATACCTGTCCGCC TTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCATAGCTCACGCT GTAGGTATCTCAGTTCGGTGTAGGTGTTTCGCTCCAAGCTGGG CTGTGTGCACGAACCCCCGTTTACGCCCCGACCGCTGCGCCTTA TCCGGTAACATATCGTCTTGAGTCCAACCCGGTAAGACACGACT TATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGC GAGGTATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCT AACTACGGCTACACTAGAAGAACAGTATTTGGTATCTGCGCTC

SEQ ID NO	Compound	Sequence (5' to 3')
		TGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTG ATCCGGCAAACAAACCACCGCTGGTAGCGGTGGTTTTTTTGT TGCAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAG ATCCTTTGATCTTTTCTACGGGGTCTGACGCTCAGTGAACGA AACTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGG ATCTTCACCTAGATCCTTTTAAATTAAAAATGAAGTTTTAAAT CAATCTAAAGTATATATGAGTAACTTGGTCTGACAGTTACCA ATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTT CGTTCATCCATAGTTGCCTGACTCCCCGTCGTGTAGATAACTA CGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGAT ACCGCGAGAACCACGCTCACC GGCTCCAGATTTATCAGCAATA AACCAGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGTCTGCAA CTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCCGGGAAGC TAGAGTAAGTAGTTTCGCCAGTTAATAGTTTGGCGAACGTTGTT GCCATTGCTACAGGCATCGTGGTGTACGCTCGTCGTTTGGTA TGGCTTCATTACAGCTCCGGTTCCCAACGATCAAGGCGAGTTAC ATGATCCCCCATGTTGTGCAAAAAAGCGTTAGCTCCTTCGGT CCTCCGATCGTTGTGAGAAGTAAGTTGGCCGCAGTGTTATCAC TCATGGTTATGGCAGCACTGCATAATTCTCTTACTGTCTGCC ATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAG TCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGCC CGGCGTCAATACGGGATAATACCGCGCCACATAGCAGAACTTT AAAAGTGCTCATCATTTGAAAAACGTTCTTCGGGGCGAAAACTC TCAAGGATCTTACCGCTGTTGAGATCCAGTTGATGTAACCCA CTCGTGACCCAACTGATCTTCAGCATCTTTTACTTTACACAG CGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAA AAGGGAATAAGGGCGACACGGAAATGTTGAATACTCATACTCT TCCTTTTTCAATATTATTGAAGCATTTATCAGGGTTATTGTCT CATGAGCGGATACATATTTGAATGTATTTAGAAAAATAACAA ATAGGGGTTCCGCGCACATTTCCCCGAAAAGTGCCAC
96	Compound 15 (pMA-RQ)	CTAAATTGTAAGCGTTAATATTTTGTAAAAATTCGCGTTAAAT TTTTGTAAATCAGCTCATTTTTTAACCAATAGGCCGAAATCG GCAAAATCCCTTATAAATCAAAAGAATAGACCGAGATAGGGTT GAGTGGCCGCTACAGGGCGCTCCCATTGCGCATTCAGGCTGCG CAACTGTTGGGAAGGGCGTTTCGGTGCGGGCCTCTTCGCTATT ACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCGATTAAAGT TGGGTAACGCCAGGGTTTTCCAGTCACGACGTTGTAAAACGA CGGCCAGTGAGCGCGACGTAATACGACTCACTATAGGGCGAAT TGGCGGAAGGCCGTCAAGGCCGCAT <u>GCCACCATGAGAATCAGC</u> <u>AAGCCCCACCTGAGATCCATCAGCATCCAGTGCTACCTGTGCC</u> <u>TGCTGCTGAACAGCCACTTTCTGACAGAGGCCGCGCATCCACGT</u> <u>GTTTCATCCTGGGCTGTTTTTCTGCCGGCCTGCC</u> TAAGACCGAG GCCAACTGGGTAAACGTGATCAGCGACCTGAAGAAGATCGAGG ACCTGATCCAGAGCATGCACATCGACGCCACACTGTACACCGA GAGCGACGTGCACCCTAGCTGTAAAGTGACCGCCATGAAGTGC TTTCTGCTGGAATGCAAGTGATCAGCCTGGAAAGCGGCGACG CCAGCATCCACGACACCGTGGAAAACCTGATCATCCTGGCCAA CAACAGCCTGAGCAGCAACGGCAATGTGACCGAGTCCGGCTGC AAAGAGTGCGAGGAAC TGAAGAGAAGAATATCAAGAGTTCC TGCAGAGCTTCGTGCACATCGTGCAGATGTTTCATCAACACCAG

SEQ ID NO	Compound	Sequence (5' to 3')
		CTGAATAGTGAGTCGTATTAACGTACCAACAAGGAGTACCCCTG ATGAGATCACTTGGATCTCATCAGGGTACTCCTTTATCTTAGA GGCATATCCCTACGTACCAACAAGGTATCCATCTCTGGCTATG AACTTGTCATAGCCAGAGATGGATACCTTTATCTTAGAGGCAT ATCCCTACGTACCAACAAGTCCCGTAACGCCATCATCTTACTT GAAGATGATGGCGTTACGGGACTTTATCTTAGAGGCATATCCC TTTTATCTTAGAGGCATATCCCTCTGGGCCTCATGGGCCTTCC GCTCACTGCCCCGCTTTCCAGTCGGGAAACCTGTCTGCCAGCT GCATTAACATGGTCATAGCTGTTTCCTTGCGTATTGGGCGCTC TCCGCTTCCTCGCTCACTGACTCGCTGCGCTCGGTCTCGGG TAAAGCCTGGGGTGCCTAATGAGCAAAAGGCCAGCAAAAGGCC AGGAACCGTAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGC TCCGCCCCCCTGACGAGCATCACAAAAATCGACGCTCAAGTCA GAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGCGTTT CCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCTGC CGCTTACCGGATACCTGTCCGCCTTTCTCCCTTCGGGAAGCGT GGCGCTTTCTCATAGCTCACGCTGTAGGTATCTCAGTTCGGTG TAGGTCGTTTCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCG TTCAGCCCCGACCGCTGCGCCTTATCCGGTAACTATCGTCTTGA GTCCAACCCGGTAAGACACGACTTATCGCCACTGGCAGCACC ACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGCTA CAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAAG AACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTC GGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAACAAACCACCG CTGGTAGCGGTGGTTTTTTTTGTTTGCAAGCAGCAGATTACGCG CAGAAAAAAGGATCTCAAGAAGATCCTTTGATCTTTTCTACG GGGTCTGACGCTCAGTGAACGAAACTCACGTTAAGGGATTT TGGTCATGAGATTATCAAAAAGGATCTTCACCTAGATCCTTTT AAATTAAAAATGAAGTTTTAAATCAATCTAAAGTATATATGAG TAAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGCAC CTATCTCAGCGATCTGTCTATTTTCGTTTCATCCATAGTTGCCTG ACTCCCCGTCGTGTAGATAACTACGATACGGGAGGGCTTACCA TCTGGCCCCAGTGCTGCAATGATACCGCGAGAACCACGCTCAC CGGCTCCAGATTTATCAGCAATAAACCAGCCAGCCGGAAGGGC CGAGCGCAGAAGTGGTCCTGCAACTTTATCCGCCTCCATCCAG TCTATTAATTGTTGCCGGAAGCTAGAGTAAGTAGTTCGCCAG TTAATAGTTTGCGCAACGTTGTTGCCATTGCTACAGGCATCGT GGTGTCACGCTCGTCGTTTGGTATGGCTTCATTACGCTCCGGT TCCCAACGATCAAGGCGAGTTACATGATCCCCCATGTTGTGCA AAAAAGCGGTTAGCTCCTTCGGTCCTCCGATCGTTGTGAGAAG TAAGTTGGCCGCGAGTGTTATCACTCATGGTTATGGCAGCACTG CATAATTCTCTTACTGTCTATGCCATCCGTAAGATGCTTTTCTG TGAATGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTGTAT GCGGCGACCGAGTTGCTCTTGCCCGGCGTCAATACGGGATAAT ACCGCGCCACATAGCAGAAGTTTAAAAGTGCTCATCATTTGGAA AACGTTCTTCGGGGCGAAACTCTCAAGGATCTTACCGCTGTT GAGATCCAGTTCGATGTAACCCACTCGTGCACCAACTGATCT TCAGCATCTTTTACTTTTACCAGCGTTTCTGGGTGAGCAAAAA CAGGAAGGCAAAATGCCGCAAAAAAGGAATAAGGGCGACACG GAAATGTTGAATACTCATACTCTTCCTTTTTCAATATTATTGA

SEQ ID NO	Compound	Sequence (5' to 3')
		AGCATTTATCAGGGTTATTGTCTCATGAGCGGATACATATTTG AATGTATTTAGAAAAATAAACAAATAGGGGTTCCGCGCACATT TCCCCGAAAAGTGCCAC
97	Compound 16 (pMA-RQ)	CTAAATTGTAAGCGTTAATATTTTGTAAAAATTCGCGTTAAAT TTTTGTAAATCAGCTCATTTTTTAACCAATAGGCCGAAATCG GCAAAATCCCTTATAAATCAAAAGAATAGACCGAGATAGGGTT GAGTGGCCGCTACAGGGCGCTCCCATTCGCCATTAGGGCTGCG CAACTGTTGGGAAGGGCGTTTCGGTGCGGGCCTCTTCGCTATT ACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCGATTAAGT TGGGTAACGCCAGGGTTTTCCAGTCACGACGTTGTAAACGA CGGCCAGTGAGCGCGACGTAATACGACTCACTATAGGGCGAAT TGGCGGAAGGCCGTCAAGGCCGCATGCCACCATGAGAATCAGC <u>AAGCCCCACCTGAGATCCATCAGCATCCAGTGCTACCTGTGCC</u> <u>TGCTGCTGAACAGCCACTTTCTGACAGAGGCCCGGCATCCACGT</u> <u>GTTTCATCCTGGGCTGTTTTTCTGCCGGCCTGCC</u> <u>TAAACCGAG</u> <u>GCCAACTGGGTAAACGTGATCAGCGACCTGAAGAAGATCGAGG</u> <u>ACCTGATCCAGAGCATGCACATCGACGCCACACTGTACACCGA</u> <u>GAGCGACGTGCACCTAGCTGTAAAGTGACCGCCATGAAGTGC</u> <u>TTTCTGCTGGAACGCAAGTGATCAGCCTGGAAGCGGCGACG</u> <u>CCAGCATCCAGCACACCGTGGAACCTGATCATCCTGGCCAA</u> <u>CAACAGCCTGAGCAGCAACGGCAATGTGACCGAGTCCGGCTGC</u> <u>AAAGAGTGCGAGGAAC</u> <u>TGGAAGAGAAGAATATCAAAAGATTCC</u> <u>TGCAGAGCTTCGTGCACATCGTGCAGATGTTTCATCAACACCAG</u> <u>CTGAATAGTGAGTCGTATTAACGTACCAACAAGGAGTACCCTG</u> <u>ATGAGATCACTTGGATCTCATCAGGGTACTCCTTTATCTTAGA</u> <u>GGCATATCCCTACGTACCAACAAGAAGGTT</u> <u>CAGCATAGTAGCT</u> <u>AACCTGTAGCTACTATGCTGAACCTTCTTTATCTTAGAGGCAT</u> <u>ATCCCTACGTACCAACAAGGACGACGAGACCTTCATCAAACTT</u> <u>GTTGATGAAGGTCTCGTCGTCCTTTATCTTAGAGGCATATCCC</u> <u>TTTTATCTTAGAGGCATATCCCTCTGGGCCTCATGGGCCTTCC</u> <u>GCTCACTGCCCCGCTTTCCAGTCGGGAAACCTGTGCTGCCAGCT</u> <u>GCATTAACATGGTCATAGCTGTTTCCTTGCGTATTGGGCGCTC</u> <u>TCCGCTTCCTCGCTCACTGACTCGCTGCGCTCGGTGCTTCGGG</u> <u>TAAAGCCTGGGGTGCC</u> <u>TAATGAGCAAAAGGCCAGCAAAAGGCC</u> <u>AGGAACCGTAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGC</u> <u>TCCGCCCCCTGACGAGCATCACAAAAATCGACGCTCAAGTCA</u> <u>GAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGCGTTT</u> <u>CCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCTGC</u> <u>CGCTTACCGGATACCTGTCCGCCTTTCTCCCTTCGGGAAGCGT</u> <u>GGCGCTTTCTCATAGCTCACGCTGTAGGTATCTCAGTTCGGTG</u> <u>TAGGTGTTTCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCCG</u> <u>TTCAGCCCCGACCGTGCGCCTTATCCGGTAACTATCGTCTTGA</u> <u>GTCCAACCCGTAAGACACGACTTATCGCCACTGGCAGCAGCC</u> <u>ACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGCTA</u> <u>CAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAAG</u> <u>AACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTC</u> <u>GGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAACAAACACCG</u> <u>CTGGTAGCGGTGGTTTTTTTTGTTTGCAAGCAGCAGATTACGCG</u> <u>CAGAAAAAAGGATCTCAAGAAGATCCTTTGATCTTTTCTACG</u> <u>GGGTCTGACGCTCAGTGGAACGAAAACCTCACGTTAAGGGATTT</u>

SEQ ID NO	Compound	Sequence (5' to 3')
		<p>TGGTCATGAGATTATCAAAAAGGATCTTCACCTAGATCCTTTT AAATTAAAAATGAAGTTTTAAATCAATCTAAAGTATATATGAG TAAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGCAC CTATCTCAGCGATCTGTCTATTTTCGTTTCATCCATAGTTGCCTG ACTCCCCGTCGTGTAGATAACTACGATACGGGAGGGCTTACCA TCTGGCCCCAGTGTGCAATGATACCGCGAGAACCACGCTCAC CGGCTCCAGATTTATCAGCAATAAACCAGCCAGCCGGAAGGGC CGAGCGCAGAAGTGGTCTGCAACTTTATCCGCTCCATCCAG TCTATTAATTGTTGCCGGAAGCTAGAGTAAGTAGTTCCGCAG TTAATAGTTTGCACAACGTTGTTGCCATTGCTACAGGCATCGT GGTGTACGCTCGTCGTTTGGTATGGCTTCATTAGCTCCGGT TCCCAACGATCAAGGCGAGTTACATGATCCCCCATGTTGTGCA AAAAAGCGGTTAGCTCCTTCGGTCTCCGATCGTTGTGAGAAG TAAGTTGGCCGAGTGTATCACTCATGGTTATGGCAGCACTG CATAATTCTCTTACTGTTCATGCCATCCGTAAGATGCTTTTCTG TGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTGTAT GCGGCGACCGAGTTGCTCTTGCCCGGCGTCAATACGGGATAAT ACCGCGCCACATAGCAGAACTTTAAAAGTGCTCATCATTGGAA AACGTTCTTCGGGGCGAAAACCTCTCAAGGATCTTACCGCTGTT GAGATCCAGTTCGATGTAACCCACTCGTGACCCCACTGATCT TCAGCATCTTTTACTTTCACCAGCGTTTCTGGGTGAGCAAAAA CAGGAAGGCAAAATGCCGCAAAAAAGGAATAAGGGCGACACG GAAATGTTGAATACTCATACTCTTCCTTTTTCAATATTATTGA AGCATTTATCAGGGTTATTGTCTCATGAGCGGATACATATTTG AATGTATTTAGAAAAATAAACAAATAGGGGTTCCGCGCACATT TCCCGAAAAGTGCCAC</p>
98	Compound 17 (pMA-RQ)	<p>CTAAATTGTAAGCGTTAATATTTTGTAAAAATTCGCGTTAAAT TTTTGTAAATCAGCTCATTTTTTAACCAATAGGCCGAAATCG GCAAAATCCCTTATAAATCAAAAGAATAGACCGAGATAGGGTT GAGTGGCCGCTACAGGGCGCTCCCATTCGCCATTAGGCTGCG CAACTGTTGGGAAGGGCGTTTCGGTGCGGGCCTCTTCGCTATT ACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCGATTAAAGT TGGGTAACGCCAGGGTTTTCCAGTCACGACGTTGTAAACGA CGGCCAGTGAGCGCGACGTAATACGACTCACTATAGGGCGAAT TGGCGGAAGGCCGTCAAGGCCGCATGCCACCATGTTCCACGTC TCCTTCCGGTACATCTTCGGCCTGCCTCCACTGATCCTGGTGC TGCTGCCTGTGGCCAGCAGCGACTGTGATATCGAGGGCAAAGA CGGCAAGCAGTACGAGAGCGTGCTGATGGTGTCATCGACCAG CTGCTGGACAGCATGAAGGAAATCGGCAGCAACTGCCTGAACA ACGAGTTCAACTTCTTCAAGCGGCACATCTGCGACGCCAACAA AGAAGGCATGTTCTGTTCAGAGCCGCCAGAAAGCTGCGGCAG TTCCTGAAGATGAACAGCACCAGGCGACTTCGACCTGCATCTGC TGAAAGTGTCTGAGGGCACCACCATCTGCTGAATTGCACCGG CCAAGTGAAGGGCAGAAAGCCTGCTGCTCTGGGAGAAGCCCAG CCTACCAAGAGCCTGGAAGAGAACAAGTCCCTGAAAGAGCAGA AGAAGCTGAACGACCTCTGCTTCCTGAAGCGGCTGCTGCAAGA GATCAAGACCTGCTGGAACAAGATCCTGATGGGCACCAAAGAA CACTGAATAGTGAGTCGTATTAACGTACCAACAAGAAGGTTCA GCATAGTAGCTAACTTGTAGCTACTATGCTGAACCTTCTTTAT CTTAGAGGCATATCCCTACGTACCAACAAGCGAATTACTGTGA</p>

SEQ ID NO	Compound	Sequence (5' to 3')
		AAGTCAAAC TTGTTGACTTTCACAGTAATTCGCTTTATCTTAG AGGCATATCCCTACGTACCAACAAGACCAGCACACTGAGAATC AAACTTGTTGATTCTCAGTGTGCTGGTCTTTATCTTAGAGGCA TATCCCTTTTATCTTAGAGGCATATCCCTCTGGGCTCATGGG CCTTCCGCTCACTGCCCCTTTCCAGTCGGGAAACCTGTGCTG CCAGCTGCATTAACATGGTCATAGCTGTTTCCTTGCGTATTGG GCGCTCTCCGCTTCCTCGCTCACTGACTCGCTGCGCTCGGTCTG TTCGGGTAAAGCCTGGGGTGCCTAATGAGCAAAAGGCCAGCAA AAGGCCAGGAACCGTAAAAAGGCCGCGTTGCTGGCGTTTTTCC ATAGGCTCCGCCCCCTGACGAGCATCACAAAAATCGACGCTC AAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAG GCGTTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGA CCCTGCCGCTTACCGGATACCTGTCCGCCTTTCTCCCTTCGGG AAGCGTGGCGCTTTCTCATAGCTCACGCTGTAGGTATCTCAGT TCGGTGTAGGTGCTTCGCTCCAAGCTGGGCTGTGTGCACGAAC CCCCCGTTCAGCCCGACCGCTGCGCCTTATCCGGTAACTATCG TCTTGAGTCCAACCCGGTAAGACACGACTTATCGCCACTGGCA GCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCG GTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACAC TAGAAGAACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTT ACCTTCGGA AAAAGAGTTGGTAGCTCTTGATCCGGCAAACAAA CCACCGCTGGTAGCGGTGGTTTTTTTTGTTTGCAAGCAGCAGAT TACGCGCAGAAAAAAGGATCTCAAGAAGATCCTTTGATCTTT TCTACGGGTCTGACGCTCAGTGGAACGAAAACCTCACGTTAAG GGATTTTGGTCATGAGATTATCAAAAAGGATCTTCACCTAGAT CCTTTTAAATTAAAAATGAAGTTTTAAATCAATCTAAAGTATA TATGAGTAACTTGGTCTGACAGTTACCAATGCTTAATCAGTG AGGCACCTATCTCAGCGATCTGTCTATTTTCGTTTCATCCATAGT TGCCTGACTCCCCGTCGTGTAGATAACTACGATACGGGAGGGC TTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGAACCAC GCTCACCGGCTCCAGATTTATCAGCAATAAACCAGCCAGCCGG AAGGGCCGAGCGCAGAAGTGGTCCTGCAACTTTATCCGCTCC ATCCAGTCTATTAATTGTTGCCGGGAAGCTAGAGTAAGTAGTT CGCCAGTTAATAGTTTGCGCAACGTTGTTGCCATTGCTACAGG CATCGTGGTGTACGCTCGTCGTTTGGTATGGCTTCATTACAGC TCCGGTTCCCAACGATCAAGGCGAGTTACATGATCCCCATGT TGTGCAAAAAGCGGTTAGCTCCTTCGGTCTCCGATCGTTGT CAGAAGTAAGTTGGCCGAGTGTTATCACTCATGGTTATGGCA GCACTGCATAATTCTCTTACTGTCATGCCATCCGTAAGATGCT TTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATA GTGTATGCGGCGACCGAGTTGCTCTTGCCCGCGTCAATACGG GATAATACCGCGCCACATAGCAGAACTTTAAAAGTGCTCATCA TTGGAAAACGTTCTTCGGGGCGAAAACCTCTCAAGGATCTTACC GCTGTTGAGATCCAGTTTCGATGTAACCCACTCGTGCACCCAAC TGATCTTCAGCATCTTTTACTTTTACCAGCGTTTCTGGGTGAG CAAAAACAGGAAGGCAAAATGCCGCAAAAAGGGAATAAGGGC GACACGGAAATGTTGAATACTCATACTCTTCCTTTTTCAATAT TATTGAAGCATTTATCAGGGTTATTGTCTCATGAGCGGATACA TATTTGAATGTATTTAGAAAAATAAACAAATAGGGGTTCGCGC CACATTTCCCCGAAAAGTGCCAC

Bold = compound sequence

Bold and underline = compound sequence

Bold Italics = Kozak sequence

*Bolding indicates construct with modified signal peptide.

5

[0209] Example 2: *In vitro* transcription of RNA constructs and data analysis

[0210] PCR-based *in vitro* transcription is carried out using the pMA-T (Cpd.1-Cpd.4), pMK-RQ (Cpd.5) or the pMA-RQ (Cpd.6-Cpd.17) vectors encoding Cpd.1-Cpd.17 to produce mRNA. A transcription template was generated by PCR using the forward and reverse primers in **Table 5**. The poly(A) tail was encoded in the template resulting in a 120 bp poly(A) tail (SEQ ID NO: 153). Optimizations were made as needed to achieve specific amplification given the repetitive sequence of siRNA flanking regions. Optimizations include: 1) decreasing the amount of plasmid DNA of vector, 2) changing the DNA polymerase (Q5 hot start polymerase, New England Biolabs), 3) reducing denaturation time (30 seconds to 10 seconds) and extension time (45 seconds/kb to 10 seconds/kb) for each cycle of PCR, 4) increasing the annealing (10 seconds to 30 seconds) for each cycle of PCR, and 5) increasing the final extension time (up to 15 minutes) for each cycle of PCR. In addition, to avoid non-specific primer binding, the PCR reaction mixture was prepared on ice including thawing reagents, and the number of PCR cycles was reduced to 25.

[0211] For *in vitro* transcription, T7 RNA polymerase (MEGAscript kit, Thermo Fisher Scientific) was used at 37°C for 2 hours. Synthesized RNAs were chemically modified with 100% N1-methylpseudo-UTP and co-transcriptionally capped with an anti-reverse CAP analog (ARCA; [m₂^{7,3'-O}G(5')ppp(5')G]) at the 5' end (Jena Bioscience). After *in vitro* transcription, the mRNAs were column-purified using MEGAclear kit (Thermo Fisher Scientific) and quantified using Nanophotometer-N60 (Implen).

[0212] Table 5. Primers for Template Generation

SEQ ID NO	Primer Direction	Sequence (5' to 3')
99	Forward	GCTGCAAGGCGATTAAGTTG
100	Reverse	U(2'Ome)U(2'Ome)U(2'Ome)TTTTTTTTTTTTTTTTTTTTTTTT TTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTT TTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTT TTTCAGCTATGACCATGTGAATGCAG

[0213] Using *in vitro* transcription, Cpd.1-Cpd.17 were generated as an mRNA and tested in various *in vitro* models specified below for IL-2, IL-7, IL-12, and IL-15 expression and combinatorial effect of respective protein overexpression in parallel to target gene down regulation.

[0214] Determination of Molecular weight of constructs was performed as below. The molecular weight of each construct was determined from each sequence by determining the total number of each base (A, C, G, T or N1-UTP) present in each sequence and multiply the number by respective molecular weight (*e.g.*, A: 347.2 g/mol; C 323.2 g/mol; G 363.2 g/mol; N1-UTP:338.2 g/mol). The molecular weight was determined by the sum of all weights obtained for each base and ARCA molecular weight of 817.4 g/mol. The molecular weight of each construct was used to calculate the amount of mRNA used for transfection in each well to nanomolar (nM) concentration.

[0215] Data were analyzed using GraphPad Prism 8 (San Diego, USA). For the estimation of the protein levels using ELISA in the standard or the sample, the mean absorbance value of the blank was subtracted from the mean absorbance of the standards or the samples. A standard curve was generated and plotted using a four parameters nonlinear regression according to manufacturer's protocol. To determine the concentration of proteins in each sample, the concentration of the different protein was interpolated from the standard curve. The final protein concentration of the sample was calculated by multiplication with the dilution factor. Statistical analysis was carried out using by Student's t-test or one-way ANOVA followed by Dunnet's multiple comparing tests.

[0216] Example 3: *In vitro* transfection of HEK-293 cells

[0217] Human embryonic kidney cells 293 (HEK-293; ATCC CRL-1573, Rockville, MD, USA) were maintained in Dulbecco's Modified Eagle's medium (DMEM, Sigma-Aldrich) supplemented with 10% (v/v) Fetal Bovine Serum (FBS, Thermofischer, Basel, Switzerland cat #10500-064). To assess the IL-2 expression the HEK-293 cells were seeded at 20,000 cell/well in a 96 well culture plate and incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours prior to transfection. Cells were then grown in DMEM growth medium containing 10% of FBS to reach confluency < 80% before transfection. Thereafter, HEK-293 cells were transfected with 300 ng of specific mRNA constructs using Lipofectamine 2000 (Thermo Fisher Scientific) following the manufacturer's instructions with the mRNA to Lipofectamine ratio of 1:1 w/v. 100 µl of DMEM was removed and 50 µl of Opti-MEM (Thermo Fisher Scientific) was added to each well followed by 50 µl mRNA and Lipofectamine 2000 complex in Opti-MEM. After 5 hours of incubation, the medium was replaced by fresh growth medium and the plates were incubated for 24 hours at 37°C in a humidified atmosphere containing 5% CO₂. Cell culture supernatant were collected to measure secreted IL-2 using ELISA (ThermoFisher Cat. # 887025). Significance (**, p<0.01)

was assessed by one way ANOVA followed by Dunnet's multiple comparing test using Cpd.1 as control.

[0218] IL-2 secretion in HEK-293 cells

[0219] Cpd.1-Cpd.4 comprising IL-2 protein coding sequence were tested for IL-2 expression and secretion from HEK-293 cells. Protein levels of secreted IL-2 were measured in the cell culture supernatant using IL-2 ELISA and are represented as fold changed referenced to Cpd.1 (containing WT IL-2 signal peptide) in **Fig. 2A**. The protein levels of secreted IL-2 by cells transfected with Cpd.2-Cpd.4 (containing modified IL-2 signal peptide) were about 2-fold higher than protein level of secreted IL-2 by cells transfected with Cpd.1. Taken together, the data suggest that Cpd.2-Cpd.4 with homologous modified signal peptides can facilitate enhanced cellular exit of produced IL-2 in HEK-293 cells compared to Cpd.1 with endogenous signal peptide. Data represent means $\pm$ standard error of the mean of 3 replicates per Cpd. Significance (**, $p < 0.01$) was assessed by one way ANOVA followed by Dunnet's multiple comparing test using Cpd.1 as control.

[0220] Example 4: *In vitro* transfection of HaCaT cells

[0221] Human keratinocytes (HaCaT; AddexBio Cat. # T0020001) were maintained in Dulbecco's Modified Eagle's medium (DMEM, Sigma-Aldrich) supplemented with 10% (v/v) Fetal Bovine Serum (FBS, ThermoFischer, Basel, Switzerland cat #10500-064). To assess the IL-2 expression the HaCaT cells were seeded at 15,000 cell/well in a 96 well culture plate and incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours prior to transfection. Cells were then grown in DMEM growth medium containing 10% of FBS to reach confluency < 70% before transfection. Thereafter, HaCaT cells were transfected with 300 ng of specific mRNA constructs using Lipofectamine 2000 (Invitrogen) following the manufacturer's instructions with the mRNA to Lipofectamine ratio of 1:1 w/v. 100 μ l of DMEM was removed and 50 μ l of Opti-MEM (Thermo Fisher Scientific) was added to each well followed by 50 μ l mRNA and Lipofectamine 2000 complex in Opti-MEM. After 5 hours of incubation, the medium was replaced by fresh growth medium and the plates were incubated for 24 hours at 37°C in a humidified atmosphere containing 5% CO₂. Cell culture supernatant were collected to measure secreted IL-2 using ELISA (ThermoFisher Cat. # 887025). Significance ($p < 0.01$) was assessed by one way ANOVA followed by Dunnet's multiple comparing test using Cpd.1 as control.

[0222] IL-2 secretion in HaCaT cells

[0223] Cpd.1-Cpd.4 comprising IL-2 protein coding sequence were tested for IL-2 expression and secretion from HaCaT cells. Protein levels of secreted IL-2 were measured in the cell

culture supernatant using IL-2 ELISA and are represented as fold changed referenced to Cpd.1 (containing WT IL-2 signal peptide) in **Fig. 2B**. The protein levels of secreted IL-2 by cells transfected with Cpd.2-Cpd.4 (containing modified IL-2 signal peptide) were about 2.7-fold higher than protein level of secreted IL-2 by cells transfected with Cpd.1. Taken together, the data suggest that Cpd.2-Cpd.4 with homologous modified signal peptides can facilitate enhanced secretion of IL-2 in HaCaT cells compared to Cpd.1 with endogenous signal peptide. Data represent means $\pm$ standard error of the mean of 3 replicates per Cpd. Significance (**, $p < 0.01$) was assessed by one way ANOVA followed by Dunnet's multiple comparing test using Cpd.1 as control.

10 **[0224] Example 5: *In vitro* transfection of A549 cells**

[0225] Human lung epithelial carcinoma cells (A549; Sigma-Aldrich Cat. # 6012804) were maintained in Dulbecco's Modified Eagle's medium high glucose (DMEM, Sigma-Aldrich) supplemented with 10% (v/v) Fetal Bovine Serum (FBS, Thermofischer, Basel, Switzerland cat #10500-064). To assess the IL-2 expression the A549 cells were seeded at 10,000 cell/well in a 96 well culture plate and incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours prior to transfection. Cells were then grown in DMEM growth medium containing 10% of FBS to reach confluency < 70% before transfection. Thereafter, A549 cells were transfected with specific mRNA constructs with varying concentrations 4.4 nM – 35.2 nM (0.15-1.2 μ g) using Lipofectamine 2000 (Invitrogen) following the manufacturer's instructions with the mRNA to Lipofectamine ratio of 1:1 w/v. 100 μ l of DMEM was removed and 50 μ l of Opti-MEM (Thermo Fisher Scientific) was added to each well followed by 50 μ l mRNA and Lipofectamine 2000 complex in Opti-MEM. After 5 hours of incubation, the medium was replaced by fresh growth medium and the plates were incubated for 24 hours at 37°C in a humidified atmosphere containing 5% CO₂. Cell culture supernatant were collected to measure secreted IL-2 using ELISA (ThermoFisher Cat. # 887025). Significance (**, $p < 0.01$) was assessed by one way ANOVA followed by Dunnet's multiple comparing test using Cpd.1 as control.

[0226] IL-2 secretion in A549 cells

[0227] Cpd.1-Cpd.4 comprising IL-2 protein coding sequence were tested for IL-2 expression and secretion from A549 cells. Protein levels of secreted IL-2 were measured in the cell culture supernatant using IL-2 ELISA and are represented as fold changed referenced to Cpd.1 (containing WT IL-2 signal peptide) in **Fig. 2C**. The protein levels of secreted IL-2 by cells transfected with Cpd.2-Cpd.4 (containing modified IL-2 signal peptide) were about 1.6-fold higher than protein level of secreted IL-2 by cells transfected with Cpd.1. Taken together,

the data suggest that Cpd.2-Cpd.4 with homologous modified signal peptides can facilitate enhanced secretion of IL-2 in A549 cells compared to Cpd.1 with endogenous signal peptide. Data represent means $\pm$ standard error of the mean of 3 replicates per Cpd. Significance (**, $p < 0.01$) was assessed by one way ANOVA followed by Dunnet's multiple comparing test using Cpd.1 as control.

[0228] Example 6: Combinatorial effect of IL-2 secretion and VEGFA down regulation in A549 cells: A VEGFA overexpression model

[0229] *In vitro* transfection of A549 cells

[0230] A VEGFA overexpression model was used to evaluate simultaneous VEGFA RNA interference (RNAi) and IL-2 expression by Cpd.5 in A549 cells. The VEGFA overexpression model was established by transfecting A549 cells with 0.3 μ g of VEGFA mRNA. A549 cells were co-transfected with increasing concentration 4.4 nM to 35.2 nM (0.15 to 1.2 μ g) of Cpd.5 to assess dose-dependent response of Cpd.5 for VEGFA interference and IL-2 overexpression. Post transfection, the cells in a growth medium without FBS were incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours, followed by quantification of VEGFA (target mRNA to downregulate; ThermoFisher Cat. #KHG0112) and IL-2 (gene of interest to overexpress; ThermoFisher Cat. # 887025) present in the same cell culture supernatant by ELISA. To assess the potency of Cpd.5 against commercially available siRNA (ThermoFisher Cat. #284703), a dose-dependent response study was performed using commercial VEGFA siRNAs and Cpd.5. A549 cells were co-transfected with VEGFA mRNA (0.3 μ g /well; 9.5 nM) and either commercial VEGFA siRNAs (0.05, 0.125, 0.25, 1.25 and 2.5 mM) or Cpd.5 (4.4, 8.8, 17.6, 26.4, 35.2 and 44.02 nM corresponds to 0.15, 0.3, 0.6, 0.9, 1.2 and 1.5 μ g respectively). Post transfection, the cells in a growth medium without FBS were incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours, followed by quantification of VEGFA (target mRNA to downregulate; ThermoFisher Cat. #KHG0112) and IL-2 (gene of interest to overexpress; ThermoFisher Cat. # 887025) present in the same cell culture supernatant by ELISA.

[0231] Results

[0232] Cpd.5 comprising 3 species of VEGFA-targeting siRNA and IL-2 protein coding sequence was tested for dose-dependent VEGFA downregulation and simultaneous IL-2 expression in A549 cells by co-transfecting A549 cells with an increasing dose of Cpd.5 (4.4 nM to 35.2 nM) and constant dose of VEGFA mRNA (9.5 nM or 300 ng/well) and measuring protein levels in the cell culture supernatant by ELISA. Cpd.5 reduced VEGFA protein level (up to 70%) while increasing IL-2 protein level in a dose-dependent manner (up to above 100

ng/ml), as demonstrated in **Fig. 3**. Taken together, the data suggest that Cpd.5 can downregulate VEGFA without affecting IL-2 expression. Data represent means $\pm$ standard error of the mean of 4 replicates.

[0233] Example 7: Combinatorial effect of IL-2 secretion and VEGFA downregulation

5 in SCC-4 cells: A VEGFA overexpression model

[0234] *In vitro* transfection of SCC-4 cells

[0235] A VEGFA overexpression model was used to evaluate simultaneous VEGFA RNA interference (RNAi) and IL-2 expression by Cpd.5 in SCC-4 cells. The VEGFA overexpression model was established by transfecting SCC-4 cells with 9.5 nM (0.3 μ g) of VEGFA mRNA. SCC-4 cells were co-transfected with increasing concentration 4.4 nM to 35.2 nM (0.15 to 1.2 μ g) of Cpd.5 to assess dose-dependent response of Cpd.5 for VEGFA interference and IL-2 overexpression. Post transfection, the cells in a growth medium without FBS were incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours, followed by quantification of VEGFA (target mRNA to downregulate; ThermoFisher Cat. #KHG0112) and IL-2 (gene of interest to overexpress; ThermoFisher Cat. #887025) present in the same cell culture supernatant by ELISA. To assess the potency of Cpd.5 against VEGFA expression, SCC-4 cells were co-transfected with 9.5 nM (0.3 μ g) VEGFA mRNA and Cpd.5 (4.4, 8.8, 17.6, 26.4, 35.2 and 44.02 nM corresponds to 0.15, 0.3, 0.6, 0.9, 1.2 and 1.5 μ g/well). Post transfection, the cells in a growth medium without FBS were incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours, followed by quantification of VEGFA (target mRNA to downregulate; ThermoFisher Cat. #KHG0112) and IL-2 (gene of interest to overexpress; ThermoFisher Cat. # 887025) present in the same cell culture supernatant by ELISA.

[0236] Results

[0237] Cpd.5, designed to have IL-2 coding sequence and 3 species of siRNA targeting VEGFA, was tested to assess the simultaneous expression of IL-2 and interference of VEGFA expression in an VEGFA overexpression model where SCC-4 cells transfected with VEGFA mRNA. Cpd.5 reduced the level of exogenously overexpressed VEGFA for up to 95% and simultaneously induced IL-2 expression (above 65 ng/ml), as demonstrated in **Fig. 4A** and **Fig. 4B**. In summary, Cpd.5 can reduce exogenously overexpressed VEGFA while simultaneously inducing IL-2 expression and secretion.

[0238] Example 8: Combinatorial effect of IL-2 secretion and VEGFA down regulation in SCC-4 cells: An endogenous VEGFA expression model

[0239] *In vitro* transfection of SCC-4 cells

[0240] SCC-4 cells were used as an endogenous VEGFA overexpression model, as SCC-4 cells endogenously overexpress VEGFA up to 600 pg/mL *in vitro* (Fig. 5A), to evaluate simultaneous VEGFA RNA interference (RNAi) and IL-2 expression by Cpd.5. SCC-4 cells were transfected with 26.4 nM (0.9 µg) of Cpd.5. Cells were incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours, followed by quantification of VEGFA (ThermoFisher Cat. #KHG0112) and IL-2 (ThermoFisher Cat. # 887025) present in the same cell culture supernatant by using specific ELISAs.

[0241] Results

[0242] Cpd.5, designed to have IL-2 coding sequence and 3 species of siRNA targeting VEGFA, was tested to assess the simultaneous expression of IL-2 and interference of VEGFA expression in SCC-4 cells that constitutively express VEGFA up to 600 pg/mL *in vitro*. Cpd.5 reduced the level of endogenous VEGFA expression for up to 90% and simultaneously induced IL-2 expression (up to 12 ng/ml), as demonstrated in Fig. 5A and Fig. 5B. Taken together Cpd.5 can reduce the level of endogenously expressed VEGFA while simultaneously inducing expression and secretion of IL-2.

[0243] Example 9: Comparative analysis of Cpd.5 and commercial siRNA in VEGFA downregulation

[0244] *In vitro* transfection of SCC-4 cells

[0245] Human tongue squamous carcinoma cell line (SCC-4; Sigma-Aldrich, Buchs Switzerland, Cat. # 89062002 CRL-1573) were maintained in Dulbecco's Modified Eagle's high glucose medium (DMEM, Sigma Aldrich) supplemented with HAM F12 (1:1) + 2 mM Glutamine + 10% Fetal Bovine Serum (FBS) + 0.4 µg/ml hydrocortisone. Cells were seeded at 15,000 cell/well in a 96 well culture plate and incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours prior to transfection. Cells were grown in DMEM/HAM F-12 growth medium to reach confluency < 70% before transfection. To assess the potency of Cpd.5 against commercially available siRNA (ThermoFisher Cat. #284703), a dose response study was performed using commercial VEGFA siRNA and Cpd.5. SCC-4 cells were co-transfected with 9.5 nM (0.3 µg) VEGFA mRNA and either commercial VEGFA siRNA (0.05, 0.125, 0.25, 1.25 and 2.5 mM) or Cpd.5 (4.4, 8.8, 17.6, 26.4, 35.2 and 44.02 nM corresponds to 0.15, 0.3, 0.6, 0.9, 1.2 and 1.5 µg/well). SCC-4 cells were transfected with Cpd.5c mRNA or siRNA constructs at specified concentrations using Lipofectamine 2000 (Invitrogen) following the manufacturer's instructions with the mRNA to Lipofectamine ratio of 1:1 w/v. 100 µl of DMEM was removed and replaced with 50 µl of Opti-MEM and 50 µl mRNA and Lipofectamine 2000 complex in Opti-MEM (Thermo Fisher

Scientific). After 5 hours, the medium was replaced by fresh growth medium without FBS and the plates were incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours.

[0246] Results

- 5 **[0247]** To calculate the inhibitory concentration of Cpd.5 against commercially available siRNA in downregulating VEGFA expression, a dose response study was performed in VEGFA overexpression model established in both SCC-4 cells and A549 cells. Both cells were co-transfected with 9.5 nM (0.3 µg) VEGFA mRNA with increasing concentration of either Cpd.5 (4.4 nM to 44.02 nM) or commercial siRNA (0.05 mM to 2.5 mM). In
10 comparison to commercial siRNA, Cpd.5 exhibited 19-fold higher potency in SCC-4 cells and more than 52-fold higher potency in A549 cells in reducing VEGFA expression (**Fig. 6A** and **Fig. 6B**). The IC₅₀ value of Cpd.5 in SCC-4 cells (8 nM) and in A549 cells (11 nM) are shown in **Fig. 6C**.

- [0248] Example 10: Combinatorial effect of IL-2 secretion and MICB down regulation**
15 **in SCC-4 cells – An endogenous MICB expression models**

[0249] *In vitro* transfection of SCC-4 cells

- [0250]** SCC-4 cells were used an endogenous MICB expression model, as SCC-4 cells constitutively express soluble MICB (up to 40 pg/mL) and membrane bound MICB (up to 80 pg/mL) *in vitro*, to evaluate simultaneous MICB RNA interference (RNAi) and IL-2
20 expression by Cpd.6. SCC-4 cells were transfected with 35.11 nM (0.9 µg) of Cpd.6 and were incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours. MICB levels present in the cell culture supernatant and cell lysate were quantified using ELISA (ThermoFisher Cat. #BMS2303). IL-2 levels present in the same cell culture supernatant was measured using ELISA (ThermoFisher Cat. # 887025).

- 25 **[0251] Results**

- [0252]** Cpd.6, designed to have IL-2 coding sequence and 3 species of siRNA targeting MICB, was tested to assess the simultaneous expression of IL-2 and interference of MICB expression in SCC-4 cells that constitutively express soluble MICB (up to 40 pg/mL) and membrane bound MICB (up to 80 pg/mL) *in vitro*. Cpd.6 reduced the level of endogenous
30 expression of both soluble and membrane bound MICB for up to 70% and 90% respectively and simultaneously induced IL-2 expression (up to 65 ng/ml), as demonstrated in **Figs. 7A-7C**. In brief, Cpd.6 can downregulate endogenously expressed MICB (both soluble and membrane bound) while simultaneously inducing expression and secretion of IL-2. Data represent means ± standard error of the mean of four replicates.

[0253] Example 11: Combinatorial effect of IL-2 secretion together with MICA and MICB down regulation in SCC-4 cells – An endogenous MICA & MICB expression model

[0254] *In vitro* transfection of SCC-4 cells

5 **[0255]** In addition to MICB, SCC-4 cells constitutively express soluble MICA (up to 200 pg/mL) *in vitro*, a functional analog to MICB. Due to high genomic homology between MICA and MICB (>90%), siRNAs in Cpd.6 were designed to interfere the expression of both MICA and MICB protein simultaneously. To evaluate synchronized MICA and MICB RNA interference (RNAi) with IL-2 expression and secretion by Cpd.6, SCC-4 cells were
10 transfected with increasing doses of Cpd.6 mRNA (1.58, 2.93, 5.85, 11.7, 23.41, 35.11 and 46.81 nM) and were incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours. MICA levels present in the cell culture supernatant were quantified using ELISA (RayBioech Cat. #ELH-MICA-1). MICB levels present in the same cell culture supernatant were quantified using ELISA (ThermoFisher Cat. #BMS2303). IL-2 levels present in the
15 same cell culture supernatant were measured using ELISA (ThermoFisher Cat. # 887025).

[0256] Results

[0257] Cpd.6, designed to have IL-2 coding sequence and 3 species of siRNA targeting both MICA and MICB, was tested to assess the simultaneous expression of IL-2 and interference of MICA/MICB expression in SCC-4 cells that constitutively express soluble MICA and
20 MICB *in vitro*. Cpd.6 reduced the level of endogenous expression of both soluble MICA and soluble MICB in a dose dependent manner up to 80% and simultaneously induced IL-2 expression (>150 ng/ml), as demonstrated in **Figs. 8A** and **8B**. In brief, Cpd.6 can downregulate endogenously expressed MICA and MICB while simultaneously inducing secretion of IL-2. Data represent means ± standard error of the mean of four replicates for IL-
25 2 level and two replicates for MICA and MICB each.

[0258] Example 12: Bioactivity evaluation of Cpd.3 in a peripheral blood mononuclear cells tumour killing assay in a SK-OV-3 spheroid model

[0259] The anti-tumor activity of Cpd.3 was assessed in immune cell-mediated tumor cell killing, by using nuclear-RFP transduced SK-OV-3 tumor cell lines. For the IL-2 expression
30 and secretion induced by Cpd.3 in spheroids, SK-OV-3-NLR cells from two dimensional (2D) culture were seeded at a single density (5000 cells/ well) into an ultra-low attachment (ULA) plate and transfected with 100 ng of Cpd.3 construct using Lipofectamine 2000, then centrifuged (200 x g for 10 min) to generate spheroids. Conditions were set up in quadruplicates. The supernatants were harvested at 12, 24 and 48 hours following the

transfection to test for IL-2 expression by TR-FRET (PerkinElmer, Cat.# TRF1221C). For experiments with peripheral blood mononuclear cells (PBMCs), the spheroids were generated and transfected with Cpd.3 (3ng, 10ng, 30 ng and 100 ng) as described above and were cultured for 48 hours to allow spheroids to reach between 200-500 μ m in diameter prior to PBMC addition. Following the 48 hour culture period, PBMCs from 3 healthy donors were added to wells (200,000 cells/well) in the presence of anti-CD3 antibody. Recombinant human IL-2 (2000 IU/ml) and PBMCs were added to appropriate wells as the positive control. SK-OV-3-NLR alone conditions did not receive PBMCs. Wells were imaged every 3 hours for 7 days using an IncuCyte (S3), with changes in the total nuclear localized RFP (NLR) integrated intensity measured as the readout for PBMC-mediated SK-OV-3 spheroid tumor killing. Total NLR integrated intensity was normalized to the 24 hour time point and analyzed using the spheroid module within the IncuCyte software. The graphs show data from Day 5 analyzed with an additional smoothing function using GraphPad Prism (averaging 4 values on each side and using a second order smoothing polynomial).

15 [0260] Results

[0261] TR-FRET analysis of the supernatants collected from the spheroids which were formed from cells transfected in 3D suspension cultures with Cpd.3 (100 ng) demonstrated time dependent increase in IL-2 expression and secretion (**Fig. 9A**). No deficiency in spheroid formation and growth was noticed due to lipofectamine transfection. Analysis of the transfected spheroids with Cpd.3 following addition of PBMCs from 3 healthy donors demonstrated clear dose-dependent immune-mediated killing. Across all donors Cpd.3 at 30 ng and 100 ng promoted PBMC-driven tumor killing determined by the reduction in the total integrated NLR intensity measured over the period of the assay (day 6 data is presented in **Figs. 9B, 9C and 9D**). The killing effect induced by Cpd.3 was substantially better than that of recombinant human IL-2 (rhIL-2) added at 6 nM concentration in all the three donors tested. **Fig. 9E** shows a set of representative IncuCyte images showing NLR integrity reduction after Cpd.3 treatment (100 ng) in the SK-OV-3 NLR condition compared to control at Day 5. In summary, transfection of SK-OV-3 NLR spheroids with Cpd.3 mRNA constructs enhanced PBMC-mediated tumor killing in a dose-dependent manner.

30 [0262] Example 13: HEK-Blue™ hIL-2 reporter assay for JAK3-STAT5 activation

[0263] The functional activity of Cpd.5 and Cpd.6 was tested in HEK-Blue™ IL-2 reporter cells (Invivogen, Cat. Code: hkb-il2), which are designed for studying the activation of human IL-2 receptor by monitoring the activation of JAK/STAT pathway. These cells were derived from the human embryonic kidney HEK293 cell line and engineered to express human IL-

2R α , IL-2R β , and IL-2R γ genes, together with the human JAK3 and STAT5 genes to achieve a totally functional IL-2 signaling cascade. In addition, a STAT5-inducible SEAP reporter gene was introduced. Upon IL-2 activation followed by STAT5, produced SEAP can be determined in real-time with HEK-Blue™ Detection cell culture medium in cell culture supernatant. Stimulation of HEK-Blue™ IL-2 cells were achieved by recombinant human IL-2 (rhIL-2, 0.001 ng to 300 ng) or IL-2 derived from cell culture supernatant of HEK293 cells (0.001 ng – 45 ng) which had been transfected with Cpd.5 or Cpd.6 (0.3 μ g/well) with below details.

[0264] HEK-Blue™ hIL-2 cells were maintained in Dulbecco's Modified Eagle's medium (DMEM, Sigma Aldrich) supplemented with 10% (v/v) Fetal Bovine Serum (FBS). The antibiotic Blasticidin (10 μ g/mL) and Zeocin (100 μ g/mL) were added to the media to select cells containing IL-2R α , IL-2R β , IL-2R γ , JAK3, STAT5 and SEAP transgene plasmids. Cells were seeded at 40,000 cell/well in a 96 well culture plate and incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours prior to stimulation. Cells were grown in DMEM growth medium containing 10% of FBS to reach confluency < 80% before stimulation. Defined concentration of IL-2 derived from HEK293 cell culture supernatant were collected, diluted in 20 μ l of media, and added to culture media of HEK-Blue™ IL-2 cells to measure IL-2 receptor recruitment followed by JAK3-STAT5 pathway activation. rhIL-2 (0.001 - 300 ng) or IL-2 derived from Cpd.5 and Cpd.6 (0.001– 45 ng) were tested in parallel. After 2 hours of incubation, SEAP activity was assessed using QUANTI-Blue™ (20 μ l cell culture supernatant + 180 μ l QUANTI-Blue™ solution) and reading the optical density (O.D.) at 620 nm in SpectraMax i3 multi-mode plate reader (Molecular Device). Untransfected samples were used as background control and subtracted from obtained O.D. values in tested samples.

[0265] Results

[0266] Stimulation of HEK-Blue™ IL-2 cells with rhIL-2 or IL-2 derived from cell culture supernatant of HEK293 cells that had been transfected with Cpd.5 or Cpd.6 was functional as all three tested compounds induced SEAP production in a dose-dependent fashion (**Figs. 10A and 10B**). In direct comparison, Cpd.5-derived IL-2 was ~5x more potent (EC₅₀: 0.02 ng/ml) compared to rhIL-2 (EC₅₀: 11 ng/ml), as well as Cpd.6 being ~2x more potent (EC₅₀: 0.08 ng/ml) compared to rhIL-2 (EC₅₀: 0.15 ng/ml). In summary, IL-2 derived from Cpd.5 and Cpd.6 are functional and induce IL-2 signaling cascade at least as potent as rhIL-2.

[0267] Example 14: NK-cell mediated killing assay of Cpd.5 and Cpd.6

[0268] Natural killer cells (NK cells) have the potential to target and eliminate tumor cells and are majorly primed by IL-2 cytokine. To measure the capacity of Cpd.5 and Cpd.6 in activating NK cells through IL-2 mechanism, SCC4 cells (Sigma-Aldrich, Buchs Switzerland, Cat. # 89062002 CRL-1573) and Natural killer 92 cells (NK-92, DSMZ, ACC488, Germany) were used. Dose response study (0.1 nM to 2.5 nM) was performed in SCC4 cells (10,000/well) by transfecting SCC-4 cells with Cpd.5 (IL-2 mRNA + 3x VEGFA siRNA), Cpd.6 (IL-2 mRNA + 3x MICA/B siRNA), mock RNA-1 (IL-4 mRNA + 3x TNF- α siRNA) or mock RNA-2 (MetLuc mRNA, no siRNA) using Lipofectamine MessengerMax (ThermoFisher, Cat.# LMRNA015) in Opti-MEM. The SCC-4 cells were then incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 30 minutes in a black 96 well culture plate. NK-92 effector cells at 100,000 cell/well in Opti-MEM were added to the transfected SCC-4 target cells in Effector to Target ratio of 10:1 (E:T = 10:1). After 24 hours, the black 96 well plate was sealed with a black foil on the bottom and washed 3 times with Dulbecco's Phosphate-Buffered Saline (PBS⁺⁺, BioConcept, Cat.#3-05F001) to remove NK-92 cells which were in suspension. Since SCC-4 cells are adherent in nature, 24 hours incubation led to strong adhesion of cells to the bottom of plate and only NK-92 cells were washed off. The rationale is that if NK cells lead to the killing of SCC-4 cells, there would be less SCC-4 cells survive and attach to the bottom of the plate after washing, which can be quantitatively measured by cell viability assay. After 3X washes, 50 μ l of PBS⁺⁺ and 50 μ l of CellTiter-Glo 2.0 (CTG 2.0, Promega, Cat.# G924B) reagent were added to each well and the 96 well plate was incubated at room temperature in the dark for 10 minutes. The luminescence was measured with the SpectraMax i3x (Molecular Devices) to calculate cell viability using standard settings.

[0269] Results

[0270] NK cell mediated killing assay revealed a dose dependent cell lysis of SCC-4 cells which were transfected with Cpd.5 or Cpd.6, and co-incubated with NK-92 cells. IL-2 secreted from SCC-4 cells promoted targeted killing of SCC-4 tumor cells at E:T ratio of 10:1 (>50% for Cpd.5 and >40% for Cpd.6, **Fig.10C**). NK cell mediated killing was observed for SCC-4 cells transfected with both Cpd.5 and Cpd.6. In brief, Cpd.5 and Cpd.6 demonstrated expected anti-tumor activity by activating NK cells in dose dependent fashion.

[0271] Example 15: Comparative analysis of Cpd.7 and Cpd.8 in IL-2 expression and VEGFA downregulation in SCC-4 cells

[0272] SCC-4 cells were cultured and transfected as described above. To assess the potency of Cpd.7 (IL-2 mRNA + 3x VEGFA siRNA) against Cpd.8 (IL-2 mRNA + 5x VEGFA

siRNA), a dose response study was performed using both compounds. SCC-4 cells were transfected with Cpd.7 (1.1, 2.2, 4.4, 8.8, 17.6, 26.4, 35.2 and 44.04 nM/well) or Cpd.8 (0.47, 0.94, 1.89, 3.79, 7.58, 15.15, 22.73, 30.31 and 37.88 nM/well). After 5 hours, the medium was replaced by fresh growth medium without FBS and the plates were incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours, and supernatant were collected. ELISA was performed to quantify VEGFA (ThermoFisher Cat. #KHG0112) and IL-2 (ThermoFisher Cat. # 887025) levels present in the same cell culture supernatant. 80% downregulation of VEGFA was calculated using a non-linear Hill binding curve with GraphPad prism.

10 [0273] Results

[0274] To calculate the inhibitory concentration of Cpd.7 against Cpd.8 in downregulating VEGFA expression, a dose response study was performed in SCC-4 cells transfected with Cpd.7 or Cpd.8. Cells were transfected with increasing concentrations of either Cpd.7 or Cpd.8 as described above. In comparison to Cpd.7, Cpd.8 exhibited 2.5-fold higher potency in SCC-4 cells in reducing VEGFA expression (**Fig. 11A**). 80% VEGF downregulation was achieved by Cpd.8 in SCC-4 cells at 8 nM whereas by Cpd.7 at 18 nM, demonstrating that increasing copy number of siRNA leads to higher level of VEGFA downregulation. However, IL-2 expression from Cpd.8 was ~2 fold lower than IL-2 expression from Cpd.7 (**Fig. 11B**). In summary, increasing copy number of siRNA in the compounds enhances RNA interference but compromises the expression of mRNA target.

20 [0275] Example 16: Time-course study of Cpd.9 and Cpd.10 in IL-2 expression and VEGFA downregulation

[0276] SCC-4 cells were cultured and transfected as described above. To assess the longitudinal potency of Cpd.9 (IL-2 mRNA + 3x VEGFA siRNA, same siRNA repeated 3 times) against Cpd.10 (IL-2 mRNA + 3x VEGFA siRNA, 3 different siRNAs with 30 bp in length), a time course study was performed using SCC-4 cells transfected with Cpd.9 or Cpd.10. SCC-4 cells were transfected with Cpd.9 or Cpd.10 at 30 nM/well concentration. Commercially available VEGFA siRNA (ThermoFisher Cat. #284703) were added to the experiment for comparison and scrambled siRNA (Sigma, Cat.# SIC002) was used as control. Cells were then incubated at 37°C in a humidified atmosphere containing 5% CO₂. The samples from different wells were collected between 6 hours and 72 hours after transfection. ELISA was performed to quantify VEGFA (ThermoFisher Cat. #KHG0112) and IL-2 (ThermoFisher Cat. # 887025) levels present in the same cell culture supernatant. VEGFA

levels from untransfected cells at each timepoint were set to 100% and the level of VEGFA downregulation was normalized to that level at the respective time point.

[0277] Results

[0278] The time course study showed the accumulation of IL-2 over 72 hours in a similar way for both Cpd.9 and Cpd.10 (**Fig. 11C**). However, Cpd.10 resulted in stronger VEGFA downregulation until 72 hours as higher than 95% RNA interference level was achieved, while Cpd.9 resulted in 85% RNA interference level after 48 hours (**Fig. 11D**). The effect was visible even at the 6 hour time point which showed VEGFA downregulation by Cpd.10 (>30%) was higher than VEGFA downregulation by Cpd.9 (20%) as demonstrated in **Fig. 11D**. As observed previously, commercial VEGFA siRNA resulted in up to 45% downregulation of VEGFA. Universal scrambled siRNA did not alter the VEGFA expression throughout the experiment phase. In summary, Cpd.10 displayed long lasting VEGFA downregulation with slightly improved potency as compared to Cpd. 9.

[0279] Example 17: Targeting multiple signaling pathways in cancer: A combination of multiple siRNA targets and immune stimulating cytokines in *in vitro* tumor models

[0280] Cancer is a complex disease with multiple dysregulated signaling pathways which promote uncontrolled proliferation of cells with reduced apoptosis. The upregulation of tumor growth signals including mammalian target of rapamycin (mTOR), cyclin-dependent kinases (CDK), vascular endothelial growth factor (VEGFA), epidermal growth factor receptor (EGFR), Kirsten rat sarcoma viral oncogene (KRAS), c-Myc proto-oncogene (c-Myc) along with high expression of immune escape proteins such as MHC class I chain-related sequence A/B (MICA/B) and Programmed cell death - ligand 1 (PD-L1) are observed in tumor cells. Moreover, tumor microenvironment displays reduced level of immune stimulating cytokines such as Interleukin-2 (IL-2), Interleukin-12 (IL-12), Interleukin-15 (IL-15) and Interleukin-7 (IL-7). Therefore, downregulation of the key proteins involved in tumor growth along with upregulation of immune stimulating cytokines can be an attractive approach for cancer therapy. To measure the downregulation of multiple pro-tumor targets through RNA interference and upregulation of immune stimulating cytokines, Cpd.11, Cpd.12, Cpd.15 and Cpd.16 were designed to comprise more than one siRNA target along with an anti-tumor interleukin mRNA. The effect of these compounds in targeting multiple signaling pathways were assessed in SCC-4 cells, A549 cells and human glioblastoma cell line (U251 MG) cells.

[0281] Head and Neck cancer *in vitro* model in SCC-4 cells

[0282] Human tongue squamous carcinoma cell line (SCC-4) was derived from the tongue of a 55-year old male and used to simulate a head and neck cancer *in vitro* model in this

example. SCC-4 cells were cultured and transfected as described above. To assess modulation of multiple cancer relevant targets in parallel using Cpd.11 (IL-12 mRNA + 1x IDH1 siRNA + 1x CDK4 siRNA + 1x CDK6 siRNA), Cpd.12 (IL-12 mRNA + 1x EGFR siRNA + 1x mTOR siRNA + 1x KRAS siRNA) and Cpd.15 (IL-15 mRNA + 1x VEGFA siRNA + 2x CD155 siRNA), SCC-4 cells were transfected with these compounds at 10 and 30 nM/well concentration. 5 hours after transfection, the medium was replaced by fresh growth medium without FBS and the plates were incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours, and supernatant were collected. ELISA was performed to quantify human IL-12p70 (ThermoFisher Cat. #88-7126) and human IL-15 (ThermoFisher Cat. #88-7620) levels present in the cell culture supernatant. The respective cell lysates were also processed to measure RNA abundance of siRNA target genes by relative quantification against untransfected samples by RT-qPCR using Cells-to-CT™ 1-Step Power SYBR Green kit (ThermoFisher Cat. #A25599) and primers (primer sequence details are listed in **Table 6**). The human 18s rRNA was used as a reference control.

15 [0283] Results

[0284] The effect of Cpd.11 comprising 1x siRNA of IDH1, CDK4 and CDK6, and IL-12 mRNA and Cpd.12 comprising 1x siRNA of EGFR, mTOR and KRAS and IL-12 mRNA was evaluated for IL-12 expression and simultaneous downregulation of target genes in SCC-4 cells transfected with two different doses (10 nM and 30 nM) of Cpd.11 or Cpd.12. The data demonstrate that both Cpd.11 and Cpd.12 lead to significant IL-12 protein expression and secretion (>7000 pg/ml) as shown in **Figs. 12A** and **12E**. In the same cell lysate, the RNA interference of Cpd.11 against IDH1, CDK4 and CDK6 RNA transcripts was assessed. As demonstrated in **Fig. 12B**, Cpd.11 downregulated endogenous IDH1 (75% for 10 nM, 90% for 30 nM), CDK4 (93% for 10 nM, 98% for 30 nM) and CDK6 (85% for 10 nM, 96% for 30 nM) levels in a dose-dependent manner. The RNA interference of Cpd.12 against EGFR, mTOR and KRAS RNA transcripts was assessed in the same cell lysate of **Fig. 12E**. As shown in **Fig. 12F**, Cpd.12 downregulated endogenous EGFR (80% for 10 nM, 92% for 30 nM), KRAS (92% for 10 nM, 83% for 30 nM) and mTOR (92% for 10 nM, 98% for 30 nM) levels in a dose-dependent manner for KRAS.

30 [0285] In addition, the effect of Cpd.15 comprising 1x VEGFA siRNA, 2x CD155 siRNA. and IL-15 mRNA was evaluated for IL-15 expression and simultaneous downregulation of the target genes in SCC-4 cells transfected with two different doses (10 nM and 30 nM) of Cpd.15. Results showed that Cpd.15 expresses IL-15 protein (>790 pg/ml), as shown in **Fig. 14C**. In the same cell lysate, the RNA interference of Cpd.15 against VEGFA and CD155

RNA transcripts was assessed using qPCR. As demonstrated in **Fig. 14D**, Cpd.15 downregulated endogenous VEGFA (95% for 10 nM, 98% for 30 nM), and CD155 (73% for 10 nM, 71% for 30 nM) levels. In short, multiple signaling pathways can be targeted using Cpd.11, Cpd.12 and Cpd.15 to downregulate multiple oncology targets through siRNAs and upregulate IL-12 or IL-15 cytokine at the same time to provide anti-tumor activity either by promoting infiltration or proliferation of immune cells.

[0286] Lung cancer *in vitro* model in A549 cells

[0287] A549 cells are adenocarcinomic human alveolar basal epithelial cells derived from cancerous lung of a 58-years old male and were used to simulate a lung cancer *in vitro* model in this example. A549 cells were cultured and transfected as described above. To assess modulation of multiple cancer relevant targets in parallel using Cpd.11 (IL-12 mRNA + 1x IDH1 siRNA + 1x CDK4 siRNA + 1x CDK6 siRNA), Cpd.12 (IL-12 mRNA + 1x EGFR siRNA + 1x mTOR siRNA + 1x KRAS siRNA) and Cpd.15 (IL-15 mRNA + 1x VEGFA siRNA + 2x CD155 siRNA), A549 cells were transfected with these compounds at 10 and 30 nM/well concentration. Five hours after transfection, the medium was replaced by fresh growth medium without FBS and the plates were incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours, and supernatant were collected. ELISA was performed to quantify human IL-12p70 (ThermoFisher Cat. #88-7126) and human IL-15 (ThermoFisher Cat. #88-7620) levels present in the cell culture supernatant. The respective cell lysates were also processed to measure RNA abundance of siRNA target genes by relative quantification against untransfected samples by RT-qPCR using Cells-to-CT™ 1-Step Power SYBR Green kit (ThermoFisher Cat. #A25599) and primers (primer sequence details are listed in **Table 6**). The human 18s rRNA used as a reference control.

[0288] Results

[0289] The effect of Cpd.11 comprising 1x siRNA of IDH1, CDK4 and CDK6 and IL-12 mRNA and Cpd.12 comprising 1x siRNA of EGFR, mTOR KRAS, and IL-12 mRNA was evaluated for IL-12 expression and simultaneous downregulation of target genes in A549 cells transfected with two different doses (10 nM and 30 nM) of Cpd.11 or Cpd.12. The data demonstrate that both Cpd.11 and Cpd.12 lead to significant IL-12 protein expression and secretion (> 1925 pg/ml) as shown in **Figs. 12C** and **12G**. In the same cell lysate, the RNA interference of Cpd.11 against IDH1, CDK4 and CDK6 RNA transcripts was assessed. As demonstrated in **Fig. 12D**, Cpd.11 downregulated endogenous IDH1 (88% for 10 nM, 92% for 30 nM), CDK4 (74% for 10 nM, 80% for 30 nM) and CDK6 (58% for 10 nM, 60% for 30 nM) levels. The RNA interference of Cpd.12 against EGFR, mTOR and KRAS RNA

transcripts was assessed in same cell lysate of **Fig. 12G**. As shown in **Fig. 12H**, Cpd.12 downregulated endogenous EGFR levels (up to 58%) in SCC-4 cells transfected with 30 nM of Cpd.12. In this cell line, endogenous KRAS mRNA expression was too low to detect by KRAS qPCR assay, levels were below quantification limit even under control conditions (BQL). As shown in **Fig. 12H**, Cpd.12 downregulated endogenous mTOR levels in a dose-dependent manner (67% for 10 nM and 79% for 30 nM).

[0290] In addition, the effect of Cpd.15 comprising 1x VEGFA siRNA, 2x CD155 siRNA, and IL-15 mRNA was evaluated for IL-15 expression and simultaneous downregulation of target genes in A549 cells transfected with different doses (10 nM and 30 nM) of Cpd.15. As shown in **Fig. 14A**, Cpd.15 lead to significant IL-15 protein expression and secretion (> 715 pg/ml). In the same cell lysate, the RNA interference of Cpd.15 against VEGFA and CD155 RNA transcripts was assessed using qPCR. As demonstrated in **Fig. 14B**, Cpd.15 downregulated endogenous VEGFA (58% for 10 nM, 51% for 30 nM) and CD155 (43% for 10 nM, 42% for 30 nM) levels. In short, multiple signaling pathways can be targeted using Cpd.11, Cpd.12 and Cpd.15 to downregulate multiple oncology targets through siRNAs and upregulate IL-12 or IL-15 cytokine at the same time to provide anti-tumor activity either by promoting infiltration or proliferation of immune cells.

[0291] Glioblastoma cancer *in vitro* model in U251 MG cells

[0292] Human glioblastoma cell line (U251 MG; DSMZ, Germany, Cat. # 09063001) was derived from a human malignant glioblastoma. U251 MG cells were maintained in Dulbecco's Modified Eagle's medium high glucose (DMEM, Sigma Aldrich, Cat # D0822) supplemented with 10% (v/v) Fetal Bovine Serum (FBS). Cells were seeded at 20,000 cell/well in a 96 well culture plate and incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours prior to transfection. Cells were grown in DMEM growth medium to reach confluency < 70% before transfection. Thereafter, U251 MG cells were transfected with Cpd.16 (IL-15 mRNA + 1x VEGFA siRNA + 1x PD-L1 siRNA + 1x c-Myc siRNA) at 10 nM or 30 nM concentration using Lipofectamine MessengerMax (Invitrogen) following the manufacturer's instructions with the compound to Lipofectamine ratio of 1:1 w/v. 100 µl of DMEM was removed and replaced with 90 µl of Opti-MEM (Thermo Fisher Scientific, Switzerland, Cat # 31985-070) and 10 µl compound and Lipofectamine MessengerMax complex in Opti-MEM. After 5 hours, the medium was replaced by fresh growth medium without FBS and the plates were incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours. ELISA was performed to quantify human IL-15 (ThermoFisher Cat. #88-7620) levels present in the cell culture supernatant. The respective cell lysates were also processed to measure RNA

abundance of siRNA target genes by relative quantification against untransfected samples by RT-qPCR using Cells-to-CT™ 1-Step Power SYBR Green kit (ThermoFisher Cat. #A25599) and primers (primer sequence details are listed in **Table 6**). The human 18s rRNA used as a reference control.

5 **[0293] Results**

[0294] The effect of Cpd.16 comprising 1x siRNA of VEGFA, PD-L1 and c-Myc and IL-15 mRNA was evaluated for IL-15 expression and simultaneous downregulation of target genes in U251 MG cells transfected with two different doses (10 nM and 30 nM) of Cpd.16. The data demonstrate that Cpd.16 expresses IL-15 protein (> 300 pg/ml) as shown in **Fig. 14E**. In the same cell lysate, the RNA interference of Cpd.16 against VEGFA, PD-L1 and c-Myc RNA transcripts was assessed. As demonstrated in **Fig. 14F**, Cpd.16 downregulated endogenous VEGFA by 99% for 10 and 30 nM, PD-L1 by >97% for 10 and 30 nM and c-Myc by >99% for 10 and 30 nM levels. In summary, multiple signaling pathways can be targeted using Cpd.16 to downregulate multiple oncology targets through siRNAs and to upregulate the IL-15 cytokine at the same time to provide anti-tumor activity by promoting proliferation of anti-tumor immune cells such as NK-cells and T-cells.

[0295] Example 18: A combination of single siRNA target and immune stimulating cytokines in *in vitro* tumor models

[0296] In this example, the impact of targeting a single pro-tumor gene for down regulation along with over expression of immune stimulating cytokine. The parallel modulation of cancer relevant target and cytokine secretion of Cpd.13 (IL-12 mRNA + 3x EGFR siRNA), Cpd.14 (IL-12 mRNA + 3x mTOR siRNA) and Cpd.17 (IL-7 mRNA + 3x PD-L1 siRNA) in SCC-4 cells, A549 cells and U251MG cells was assessed. All the three cells were cultured and transfected as described above with two different doses (10 nM and 30 nM) of above compounds. 24 hours after transfection, supernatant were collected. ELISA was performed to quantify human IL-12p70 (ThermoFisher Cat. #88-7126) and human IL-7 (ThermoFisher Cat. # EHIL7) levels present in the cell culture supernatant. The respective cell lysates were also processed to measure RNA abundance of siRNA target genes by relative quantification against untransfected samples by RT-qPCR using Cells-to-CT™ 1-Step Power SYBR Green kit (ThermoFisher Cat. #A25599) and primers (primer sequence details are listed in **Table 6**). The human 18s rRNA used as a reference control.

[0297] Results

[0298] The effect of Cpd.13 comprising 3x EGFR siRNA and IL-12 mRNA was evaluated for IL-12 expression and simultaneous EGFR gene downregulation in both A549 cells and

SCC-4 cells transfected with two different doses (10 nM and 30 nM) Cpd.13. As shown in **Figs. 13A and 13B**, Cpd.13 expressed IL-12 protein in both A549 cells (up to 2030 pg/ml) and SCC-4 cells (up to 7420 pg/ml). In the same cell lysate, the RNA interference of Cpd.13 against EGFR RNA transcripts was assessed. As demonstrated in **Fig. 13D and Fig. 13E**,
 5 Cpd.13 downregulated the endogenous EGFR levels (30-40% in A549 cells and 85-92% in SCC-4 cells).

[0299] Likewise, Cpd.14 comprising 3x mTOR siRNA and IL-12 mRNA was evaluated for IL-12 expression and simultaneous mTOR gene downregulation in A549 cells transfected with two different doses (10 nM and 30 nM) of Cpd.14. As shown in **Fig. 13C**, Cpd.14
 10 expressed IL-12 protein (up to 2800 pg/ml in cells transfected with 10 nM of Cpd.14 and 365 pg/ml in cells transfected with 30 nM of Cpd.14 (>7-fold lower compared to 10 nM Cpd.14)). In cells transfected with 30 nM of Cpd.14, a great level of cell death was observed as mTOR is a cell survival marker. In the same cell lysate, the RNA interference of Cpd.14 against mTOR RNA transcripts was evaluated. As demonstrated in **Fig. 13F**, Cpd.14 downregulated
 15 the endogenous mTOR levels (50-73% in A549 cells).

[0300] In U251 MG cells, the effect of Cpd.17 (10 nM and 30 nM concentration) comprising 3x PD-L1 siRNA and IL-7 mRNA was evaluated for IL-7 expression and simultaneous PD-L1 gene downregulation. As shown in **Fig. 14G**, Cpd.17 expressed IL-7 protein (up to 1300 pg/ml). In the same cell lysate, the RNA interference of Cpd.14 against PD-L1 RNA
 20 transcripts was evaluated. As demonstrated in **Fig. 14H**, Cpd.14 downregulated endogenous PD-L1 levels (60-87% in U251 MG cells) in a dose relevant manner.

[0301] **Table 6.** Primers used in qPCR assay

Gene Name	Primer Direction	Sequence (5' to 3')	SEQ ID NO
IDH1	Forward	GCTCTGTCTAAGGGTTGGCC	101
	Reverse	CCATGTCGTCGATGAGCCTA	102
CDK4	Forward	GAGTCCCCAATGGAGGAGGA	103
	Reverse	TCCATCAGCCGACACATT	104
CDK6	Forward	GCAGACCGGCGAGGAG	105
	Reverse	CTGTTCGTGACACTGTGCA	106
EGFR	Forward	TACCTCATCCACAGCAGG	107
	Reverse	GCTGTCTTCCTTGATGGGAC	108
KRAS	Forward	GTACAGTGCAATGAGGGACCA	109
	Reverse	CACAAAGAAAGCCCTCCCCA	110
mTOR	Forward	CATGCATGACAACAGCCCAG	111
	Reverse	AGCTTCAGGGGCATCAAACA	112
VEGFA	Forward	TTGCCTTGCTGCTCTACCTC	113
	Reverse	GGAGGGCAGAATCATCACGA	114
CD155	Forward	CCCAAATCACCTGGCACTCA	115
	Reverse	CTCAAAGCTCTCGTGCTCCA	116
PD-L1	Forward	GTTGAAGGACCAGCTCTCCC	117
	Reverse	CTTGTAAGTCGGCACCACCAT	118

c-Myc	Forward	ACTGTATGTGGAGCGGCTTC	119
	Reverse	CAGGTACAAGCTGGAGGTGG	120
18s	Forward	ACCCGTTGAACCCCATTCGTGA	121
	Reverse	GCCTCACTAAACCATCCAATCGG	122

[0302] Example 19: Human umbilical vein endothelial cells (HUVEC) tube-formation assay: *In vitro* angiogenesis model

[0303] To assess the functional relevance of VEGFA downregulation potency of Cpd.5 and Cpd.10, SCC-4 cells were cultured and transfected with Cpd.5 and Cpd.10 (20 and 30 nM/well) as described above. After 5 hours, the medium was replaced by fresh growth medium without FBS and the plates were incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours to produce and secrete VEGFA into the medium, and supernatants were collected and VEGFA levels quantified by ELISA (ThermoFisher Cat. #KHG0112). The same cell culture supernatant was used to assess the functional ability of the secreted VEGF to induce angiogenesis of human umbilical vein endothelial cells (HUVECs) without treatment or 24 hours post treatment with Cpd.5 and Cpd.10. HUVECs have the ability to form three-dimensional capillary-like tubular structures (also known as pseudo-tube formation) when plated at subconfluent densities with the appropriate extracellular matrix support. The angiogenesis model was established to measure anti-angiogenesis activity of Cpd.5 and Cpd.10 in this *in vitro*. HUVEC cells (ATCC, Cat. #CRL-1730) were maintained in F-12K medium (ATCC Cat. #30-2004) supplemented with 10% FBS (ATCC, #30-2020), 0.1 mg/mL heparin (Sigma, #H3393), and 30 µg/mL ECGS (Corning, #354006) at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours prior to dispensing into Matrigel coated Ibidi plates. 24 hours prior to experiment, pipet tips and µ-slide angiogenesis Ibidi plates (Ibidi, Cat. #81506) were placed at -20°C. Growth factor-reduced BD Matrigel (BD Biosciences, Cat. #354230) was thawed overnight on ice in a refrigerator. On the day of experiment, Matrigel, pipet tips and plate were kept on ice, in the laminar flow, during the Matrigel application. 10 µL of Matrigel was applied into each inner well of Ibidi plates, preventing it from flowing into the upper well. Plates coated with Matrigel were put at 37°C for 1 hour in a humidified chamber. HUVECs were trypsinized and counted using a standard procedure, and the cells were suspended at a concentration of 5000 cells/50µL in cell media either derived from SCC-4 cells supernatant (no treatment) or SCC-4 cells supernatant treated with Cpd.5 or Cpd.10 (20 nM or 30 nM) or media with recombinant VEGFA (0.5 or 5 ng/mL). Fresh HUVEC culture medium used as a baseline control. After Matrigel polymerization, 50µL of cell suspension described above were loaded into each well. Ibidi plates were incubated at 37 °C, 5% CO₂ for 6-hours. Cells were visualized

with a microscope and images were taken (0 hour and 6 hour) and analyzed for tube formation and number of branching points.

[0304] Results

[0305] Cpd.5 and Cpd.10 designed to have IL-2 coding sequence and 3 species of siRNA targeting VEGFA, were tested to assess the interference of VEGFA expression in SCC-4 cells. Under control conditions, SCC-4 cells produced and secreted approximately 0.8 ng/ml VEGFA into the medium (**Fig. 15A**). Transfection with Cpd.5 reduced the VEGFA levels down to 76% and 60% at 20 and 30 nM, respectively, whereas Cpd.10 treatment reduced VEGFA more potently to 30% at both 20 and 30 nM (**Fig. 15A**). 50 µl of these cell culture supernatants were analyzed for their functional ability to induce branching point formation as marker of *in vitro* angiogenesis in HUVEC cells and compared with untreated controls or media with defined rh-VEGFA concentrations (0.5 and 5 ng/mL). **Fig. 15B** shows that the potency to increase branching points as measure for tube formation correlated well with medium VEGFA. SCC-4 cells under control conditions produced VEGFA to induce significant branching point formation similar to the two rh-VEGFA controls. Supernatants from both Cpd.5 and Cpd.10 strongly reduced branching points as result of reduced VEGFA levels, with Cpd.10 supernatant being slightly more potent to reduce branching point formation than Cpd.5 due to lower VEGFA levels.

[0306] The examples and embodiments described herein are for illustrative purposes only and various modifications or changes suggested to persons skilled in the art are to be included within the spirit and purview of this application and scope of the appended claims.

CLAIMS

WHAT IS CLAIMED IS:

1. A composition comprising a first RNA linked to a second RNA, wherein the first RNA encodes for a cytokine, and wherein the second RNA encodes for a genetic element that modulates expression of a gene associated with tumor proliferation.
2. The composition of claim 1, wherein the cytokine is interleukin-2 (IL-2), IL-12, IL-15, IL-7, a fragment thereof, or a functional variant thereof.
3. The composition of claim 1, wherein the cytokine comprises a sequence selected from the group consisting of SEQ ID NOs: 24, 44, 47, 68, and 80.
4. The composition of claim 1, wherein the cytokine comprises a signal peptide.
5. The composition of claim 4, wherein the signal peptide comprises an unmodified signal peptide sequence or a modified signal peptide sequence.
6. The composition of claim 5, wherein the unmodified signal peptide sequence comprises a sequence selected from the group consisting of SEQ ID NOs: 26 and 125-128.
7. The composition of claim 2, wherein the IL-2 comprises a signal peptide.
8. The composition of claim 7, wherein the signal peptide comprises an unmodified IL-2 signal peptide sequence.
9. The composition of claim 8, wherein the unmodified IL-2 signal peptide sequence comprises a sequence listed in SEQ ID NO: 26.
10. The composition of claim 7, wherein the signal peptide comprises an IL-2 signal peptide sequence modified by insertion, deletion, or substitution of at least one amino acid.
11. The composition of claim 10, wherein the IL-2 signal peptide sequence modified by insertion, deletion, or substitution of at least one amino acid comprises a sequence selected from the group consisting of SEQ ID NOs: 27-29.
12. The composition of claim 1, wherein the first RNA is a messenger RNA (mRNA).
13. The composition of claim 1, wherein the second RNA is a small interfering RNA (siRNA).
14. The composition of claim 13, wherein the siRNA is capable of binding to an mRNA of the gene associated with tumor proliferation.
15. The composition of claim 13, wherein the second RNA comprises 1, 2, 3, 4, 5, or more species of siRNA, wherein each species of siRNA comprises a different sequence targeting a different region of the same mRNA.

16. The composition of claim 13, wherein the second RNA comprises 1, 2, 3, 4, 5, or more redundant species of siRNA.
17. The composition of claim 15 or 16, wherein each species of the 1, 2, 3, 4, 5, or more species of siRNA is connected by a linker comprising a sequence listed in SEQ ID NO: 22.
18. The composition of claim 1, wherein the gene associated with tumor proliferation comprises a gene associated with angiogenesis.
19. The composition of claim 18, wherein the gene associated with angiogenesis encodes vascular endothelial growth factor (VEGF), a fragment thereof, or a functional variant thereof.
20. The composition of claim 19, wherein the VEGF is VEGFA, a fragment thereof, or a functional variant thereof.
21. The composition of claim 20, wherein the VEGFA comprises a sequence listed in SEQ ID NO: 35.
22. The composition of claim 19, wherein the VEGF is an isoform of VEGFA, a fragment thereof, or a functional variant thereof.
23. The composition of claim 19, wherein the VEGF is placental growth factor (PIGF), a fragment thereof, or a functional variant thereof.
24. The composition of claim 1, wherein the gene associated with tumor proliferation comprises isocitrate dehydrogenase (IDH1), cyclin-dependent kinase 4 (CDK4), CDK6, epidermal growth factor receptor (EGFR), mechanistic target of rapamycin (mTOR), Kirsten rat sarcoma viral oncogene (KRAS), cluster of differentiation (CD155), programmed cell death-ligand 1 (PD-L1), or myc proto-oncogene (c-Myc).
25. The composition of claim 24, wherein the gene associated with tumor proliferation comprises a sequence selected from the group consisting of SEQ ID NOs: 50, 53, 56, 59, 62, 65, 71, 74, and 77.
26. The composition of claim 1, wherein the first RNA is linked to the second RNA by a linker.
27. The composition of claim 26, wherein the linker comprises a tRNA linker or a linker comprising a sequence listed in SEQ ID NO: 21.
28. The composition of claim 1, further comprising a poly(A) tail, a 5' cap, or a Kozak sequence.
29. The composition of any one of claims 1-28, wherein the first RNA and the second RNA are both recombinant.

30. A composition comprising a first RNA linked to a second RNA, wherein the first RNA encodes for a cytokine, and wherein the second RNA encodes for a genetic element that modulates expression of a gene associated with recognition by the immune system.
- 5 31. The composition of claim 30, wherein the cytokine is interleukin-2 (IL-2), a fragment thereof, or a functional variant thereof.
32. The composition of claim 31, wherein the IL-2 comprises a sequence comprising SEQ ID NO: 24.
33. The composition of claim 31, wherein the IL-2 comprises a signal peptide.
- 10 34. The composition of claim 33, wherein the signal peptide comprises an unmodified IL-2 signal peptide sequence.
35. The composition of claim 34, wherein the IL-2 signal peptide sequence comprises a sequence listed in SEQ ID NO: 26.
36. The composition of claim 33, wherein the signal peptide comprises an IL-2 signal peptide sequence modified by insertion, deletion and/or substitution of at least one amino acid.
- 15 37. The composition of claim 36, wherein the IL-2 signal peptide sequence modified by insertion, deletion, or substitution of at least one amino acid comprises a sequence selected from the group consisting of SEQ ID NOs: 27-29.
- 20 38. The composition of claim 30, wherein the first RNA is a messenger RNA (mRNA).
39. The composition of claim 30, wherein the second RNA is a small interfering RNA (siRNA).
40. The composition of claim 39, wherein the siRNA is capable of binding to an mRNA of the gene associated with recognition by the immune system.
- 25 41. The composition of claim 40, wherein the gene associated with recognition by immune system encodes MHC class I chain-related sequence A (MICA), a fragment thereof, or a functional variant thereof.
42. The composition of claim 41, wherein the MICA comprises a sequence listed in SEQ ID NO: 38.
- 30 43. The composition of claim 40, wherein the gene associated with recognition by the immune system encodes MHC class I chain-related sequence B (MICB), a fragment thereof, or a functional variant thereof.
44. The composition of claim 43, wherein the MICB comprises a sequence listed in SEQ ID NO: 41.

45. The composition of claim 40, wherein the gene associated with recognition by the immune system encodes endoplasmic reticulum protein (ERp5), a disintegrin and metalloproteinase (ADAM), matrix metalloproteinase (MMP), a fragment thereof, or a functional variant thereof.
- 5 46. The composition of claim 45, wherein the ADAM is ADAM17.
47. The composition of claim 39, wherein the second RNA comprises 1, 2, 3, 4, 5, or more species of siRNA, wherein each species of siRNA comprises a different sequence targeting a different region of the same mRNA.
- 10 48. The composition of claim 39, wherein the second RNA comprises 1, 2, 3, 4, 5, or more redundant species of siRNA.
49. The composition of claim 47 or 48, wherein each species of the 1, 2, 3, 4, 5, or more species of siRNA is connected by a linker comprising a sequence listed in SEQ ID NO: 22.
- 15 50. The composition of claim 30, wherein the first RNA is linked to the second RNA by a linker.
51. The composition of claim 50, wherein the linker comprises a tRNA linker or a linker comprising a sequence listed in SEQ ID NO: 21.
52. The composition of claim 30, further comprising a poly(A) tail, a 5' cap, or a Kozak sequence.
- 20 53. The composition of any one of claims 30-52, wherein the first RNA and the second RNA are both recombinant.
54. A composition comprising:
a first RNA encoding for interleukin-2 (IL-2), IL-15, a fragment thereof, or a functional variant thereof linked to a second RNA encoding for a genetic element that
25 modulates expression of vascular endothelial growth factor A (VEGFA), an isoform of VEGFA, placental growth factor (PIGF), cluster of differentiation 155 (CD155), programmed cell death-ligand 1 (PD-L1), myc proto-oncogene (c-Myc), a fragment thereof, or a functional variant thereof.
55. The composition of claim 54, wherein the first RNA is a messenger RNA (mRNA).
- 30 56. The composition of claim 54, wherein the IL-2 comprises a sequence comprising SEQ ID NO: 24.
57. The composition of claim 54, wherein the IL-2 comprises a signal peptide.
58. The composition of claim 57, wherein the signal peptide comprises an unmodified IL-2 signal peptide sequence.

59. The composition of claim 58, wherein the IL-2 signal peptide sequence comprises a sequence listed in SEQ ID NO: 26.
60. The composition of claim 57, wherein the signal peptide comprises an IL-2 signal peptide sequence modified by insertion, deletion and/or substitution of at least one amino acid.
61. The composition of claim 60, wherein the IL-2 signal peptide sequence modified by insertion, deletion, or substitution of at least one amino acid comprises a sequence selected from the group consisting of SEQ ID NOs: 27-29.
62. The composition of claim 54, wherein the IL-15 comprises a sequence comprising SEQ ID NO: 68.
63. The composition of claim 54, wherein the IL-15 comprises a signal peptide.
64. The composition of claim 63, wherein the signal peptide comprises an unmodified IL-15 signal peptide sequence.
65. The composition of claim 64, wherein the unmodified IL-15 signal peptide sequence comprises a sequence listed in SEQ ID NO: 144.
66. The composition of claim 54, wherein the second RNA is a small interfering RNA (siRNA).
67. The composition of claim 66, wherein the siRNA is capable of binding to an mRNA of VEGFA, an isoform of VEGFA, PIGF, CD155, PD-L1, or c-Myc.
68. The composition of claim 67, wherein the VEGFA comprises a sequence comprising SEQ ID NO: 35.
69. The composition of claim 67, wherein the CD155 comprises a sequence comprising SEQ ID NO: 71.
70. The composition of claim 67, wherein the PD-L1 comprises a sequence comprising SEQ ID NO: 74.
71. The composition of claim 67, wherein the c-Myc comprises a sequence comprising SEQ ID NO: 77.
72. The composition of claim 66, wherein the second RNA comprises 1, 2, 3, 4, 5, or more species of siRNA, wherein each species of siRNA comprises a different sequence targeting a different region of the same mRNA.
73. The composition of claim 66, wherein the second RNA comprises 1, 2, 3, 4, 5, or more redundant species of siRNA.

74. The composition of claim 72 or 73, wherein each species of the 1, 2, 3, 4, 5, or more species of siRNA is connected by a linker comprising a sequence listed in SEQ ID NO: 22.
- 5 75. The composition of claim 54, wherein the first RNA is linked to the second RNA by a linker.
76. The composition of claim 75, wherein the linker comprises a tRNA linker or a linker comprising a sequence comprising SEQ ID NO: 21.
77. The composition of claim 54, further comprising a poly(A) tail, a 5' cap, or a Kozak sequence.
- 10 78. The composition of any one of claims 54-77, wherein the first RNA and the second RNA are both recombinant.
79. A composition comprising:
a first RNA encoding for interleukin-2 (IL-2), a fragment thereof, or a functional variant thereof linked to a second RNA encoding for a genetic element that modulates
15 expression of MHC class I chain-related sequence A (MICA), MHC class I chain-related sequence B (MICB), endoplasmic reticulum protein (ERp5), a disintegrin and metalloproteinase (ADAM), matrix metalloproteinase (MMP), a fragment thereof, or a functional variant thereof.
80. The composition of claim 79, wherein the ADAM is ADAM17.
- 20 81. The composition of claim 79, wherein the first RNA is a messenger RNA (mRNA).
82. The composition of claim 79, wherein the IL-2 comprises a sequence comprising SEQ ID NO: 24.
83. The composition of claim 79, wherein the IL-2 comprises a signal peptide.
84. The composition of claim 83, wherein the signal peptide comprises an unmodified
25 IL-2 signal peptide sequence.
85. The composition of claim 84, wherein the IL-2 signal peptide sequence comprises a sequence listed in SEQ ID NO: 26.
86. The composition of claim 83, wherein the signal peptide comprises an IL-2 signal peptide sequence modified by insertion, deletion and/or substitution of at least one
30 amino acid.
87. The composition of claim 86, wherein the IL-2 signal peptide sequence modified by insertion, deletion, or substitution of at least one amino acid comprises a sequence selected from the group consisting of SEQ ID NOs: 27-29.

88. The composition of claim 79, wherein the second RNA is a small interfering RNA (siRNA).
89. The composition of claim 88, wherein the siRNA is capable of binding to an mRNA of MICA, MICB, ERp5, ADAM, or MMP.
- 5 90. The composition of claim 89, wherein the MICA comprises a sequence comprising SEQ ID NO: 38.
91. The composition of claim 89, wherein the MICB comprises a sequence comprising SEQ ID NO: 41.
92. The composition of claim 89, wherein the ADAM is ADAM17.
- 10 93. The composition of claim 88, wherein the second RNA comprises 1, 2, 3, 4, 5, or more species of siRNA, wherein each species of siRNA comprises a different sequence targeting a different region of the same mRNA.
94. The composition of claim 88, wherein the second RNA comprises 1, 2, 3, 4, 5, or more redundant species of siRNA.
- 15 95. The composition of claim 93 or 94, wherein each species of the 1, 2, 3, 4, 5, or more species of siRNA is connected by a linker comprising a sequence listed in SEQ ID NO: 22.
96. The composition of claim 79, wherein the first RNA is linked to the second RNA by a linker.
- 20 97. The composition of claim 96, wherein the linker comprises a tRNA linker or a linker comprising a sequence listed in SEQ ID NO: 21.
98. The composition of claim 79, further comprising a poly(A) tail, a 5' cap, or a Kozak sequence.
99. The composition of any one of claims 79-98, wherein the first RNA and the second RNA are both recombinant.
- 25 100. A composition comprising:
a first RNA encoding for interleukin-12 (IL-12), IL-7, a fragment thereof, or a functional variant thereof linked to a second RNA encoding for a genetic element that modulates expression of isocitrate dehydrogenase (IDH1), cyclin-dependent kinase 4 (CDK4), CDK6, epidermal growth factor receptor (EGFR), mechanistic target of rapamycin (mTOR), Kirsten rat sarcoma viral oncogene (KRAS), programmed cell death-ligand 1 (PD-L1), a fragment thereof, or a functional variant thereof.
- 30 101. The composition of claim 100, wherein the first RNA is a messenger RNA (mRNA).

102. The composition of claim 100, wherein the IL-12 comprises a sequence comprising SEQ ID NO: 44 or SEQ ID NO: 47.
103. The composition of claim 100, wherein the IL-12 comprises a signal peptide.
104. The composition of claim 103, wherein the signal peptide comprises an
5 unmodified IL-12 signal peptide.
105. The composition of claim 104, wherein the unmodified IL-12 signal peptide comprises a sequence listed in SEQ ID NO: 142 or SEQ ID NO: 143.
106. The composition of claim 100, wherein the IL-7 comprises a sequence comprising SEQ ID NO: 80.
107. The composition of claim 100, wherein the IL-7 comprises a signal peptide.
108. The composition of claim 107, wherein the signal peptide comprises an
10 unmodified IL-7 signal peptide.
109. The composition of claim 108, wherein the unmodified IL-7 signal peptide comprises a sequence listed in SEQ ID NO: 128.
110. The composition of claim 100, wherein the second RNA is a small interfering
15 RNA (siRNA).
111. The composition of claim 110, wherein the siRNA is capable of binding to an mRNA of IDH1, CDK4, CDK6, EGFR, mTOR, KRAS, or PD-L1.
112. The composition of claim 111, wherein IDH1 comprises a sequence
20 comprising SEQ ID NO: 50.
113. The composition of claim 111, wherein CDK4 comprises a sequence comprising SEQ ID NO: 53.
114. The composition of claim 111, wherein CDK6 comprises a sequence comprising SEQ ID NO: 56.
115. The composition of claim 111, wherein mTOR comprises a sequence
25 comprising SEQ ID NO: 62.
116. The composition of claim 111, wherein EGFR comprises a sequence comprising SEQ ID NO: 59.
117. The composition of claim 111, wherein KRAS comprises a sequence
30 comprising SEQ ID NO: 65.
118. The composition of claim 111, wherein PD-L1 comprises a sequence comprising SEQ ID NO: 74.

119. The composition of claim 110, wherein the second RNA comprises 1, 2, 3, 4, 5, or more species of siRNA, wherein each species of siRNA comprises a different sequence targeting a different region of the same mRNA.
120. The composition of claim 110, wherein the second RNA comprises 1, 2, 3, 4, 5, or more redundant species of siRNA.
121. The composition of claim 119 or 120, wherein each species of the 1, 2, 3, 4, 5, or more species of siRNA is connected by a linker comprising a sequence listed in SEQ ID NO: 22.
122. The composition of claim 100, wherein the first RNA is linked to the second RNA by a linker.
123. The composition of claim 122, wherein the linker comprises a tRNA linker or a linker comprising a sequence comprising SEQ ID NO: 21.
124. The composition of claim 100, further comprising a poly(A) tail, a 5' cap, or a Kozak sequence.
125. The composition of any one of claims 100-124, wherein the first RNA and the second RNA are both recombinant.
126. A pharmaceutical composition comprising a therapeutically effective amount of the composition of any one of claims 1-125 and a pharmaceutically acceptable excipient.
127. A method of treating a cancer, comprising administering the composition of any one of claims 1-125 of the pharmaceutical composition of claim 126 to a subject having a cancer.
128. The method of claim 127, wherein the cancer is a solid tumor.
129. The method of claim 127, wherein the cancer is melanoma.
130. The method of claim 127, wherein the cancer is renal cell carcinoma.
131. The method of claim 127, wherein the cancer is a head and neck cancer.
132. The method of claim 131, wherein the head and neck cancer is head and neck squamous cell carcinoma.
133. The method of claim 131, wherein the head and neck cancer is laryngeal cancer, hypopharyngeal cancer, nasal cavity cancer, paranasal sinus cancer, nasopharyngeal cancer, oral cancer, oropharyngeal cancer, salivary gland cancer, brain tumors, esophageal cancer, eye cancer, parathyroid cancer, sarcoma of the head and neck, or thyroid cancer.

134. The method of claim 127, wherein the cancer is located at an upper aerodigestive tract.
135. The method of claim 134, wherein the upper aerodigestive tract comprises a paranasal sinus, a nasal cavity, an oral cavity, a salivary gland, a tongue, a nasopharynx, an oropharynx, a hypopharynx, or a larynx.
136. The method of claim 127, wherein the subject has a head and neck cancer.
137. The method of claim 136, wherein the subject having the head and neck cancer has a history of tobacco usage.
138. The method of claim 136, wherein the subject having the head and neck cancer has a human papillomavirus (HPV) DNA.
139. The method of any one of claims 127-138, wherein the subject is a human.
140. A composition comprising a recombinant polynucleic acid construct comprising a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1-17 and 125-141.
141. A composition according to any one of claims 1-125 for use in modulating the expression of two or more genes in a cell.
142. A cell comprising the composition of any one of claims 1-125.
143. A vector comprising a recombinant polynucleic acid construct encoding the composition of any one of claims 1-125.
144. A method of producing an siRNA and an mRNA from a single RNA transcript in a cell, comprising introducing into the cell the composition of any one of claims 1-125 or the vector of claim 143.
145. A method of modulating protein expression comprising introducing the composition of any one of claims 1-125 or the vector of claim 143 into a cell, wherein the expression of a protein encoded by the second RNA is decreased compared to a cell without the composition of any one of claims 1-125 or the vector of claim 143.
146. A method of modulating protein expression comprising introducing the composition of any one of claims 1-125 or the vector of claim 143 into a cell, wherein the expression of a protein encoded by the first RNA is increased compared to a cell without the composition of any one of claims 1-125 or the vector of claim 143.
147. A method of modulating protein expression comprising introducing the composition of any one of claims 1-125 or the vector of claim 143 into a cell,

wherein the expression of a protein encoded by the second RNA is decreased compared to a cell without the composition of any one of claims 1-125 or the vector of claim 143, and wherein the expression of a protein encoded by the first RNA is increased compared to a cell without the composition of any one of claims 1-125 or the vector of claim 143.

5