



We claim:

1. A compound having a structure represented by the following formula (I):



wherein:

X, Y, and Z are independently N or CH;



each A_1 is independently O, S, CR_1R_2 , or NR_1 ;

R is selected from CF_3 , halogen, CN, alkyl, aryl, heteroaryl, NR_1R_2 , $S(O)_{0-2}NR_1R_2$, or $S(O)_{0-2}R_1R_2$;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;


D is selected from a bond, O, CR_1R_2 or NH or NR_1 or NR_1R_2 or $S(O)_{0-2}R_1R_2$;

E is absent or selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂ cycloalkyl or C₃-C₁₂ cycloheteroalkyl or -(CH₂)_x-C₃-C₁₂ cycloalkyl or -(CH₂)_x-C₃-C₁₂ cycloheteroalkyl;


R₁ is absent or selected from H, alkyl, aryl, or heteraryl; and

R₂ is selected from H, alkyl, aryl, heteroaryl, or COR₁, or R₁ and R₂ form a C₃-C₁₂ cycloalkyl or C₃-C₁₂ cycloheteroalkyl containing O, N and/or S; or a pharmaceutically acceptable salt thereof.

2. The compound of claim 1, wherein X, Y and Z are selected to form:



3. The compound of claim 1, wherein X, Y and Z are selected to form:






































4. The compound of claim 1, wherein M is optionally substituted with one or more R, and is selected from C₃-C₁₂ cycloalkyl or C₃-C₁₂ cycloalkenyl or aryl or heteroaryl or C₃-C₁₂

heterocycloalkyl or C₃-C₁₂ heterocycloalkenyl or 3-8 membered ring comprising C, O, S, and/or N.















5. The compound of claim 4, wherein M is optionally substituted phenyl or pyridine.

6. The compound of claim 1, wherein M, D, and E together form:


			
			
			
			
			
			
			


			
			








7. The compound of claim 1, wherein A is chosen from the following:

						
						
						
						
						
 or						

8. A compound of one of the following structures:






or a pharmaceutically acceptable salt thereof.

9. The compound of claim 1, wherein X is N and Y is CH.
10. A compound of any one of claims 1-9 for use in therapy.
11. The compound of claim 10, wherein the therapy is associated with modulating the BMP signaling pathway.
12. Use of a compound of any one of claims 1-9 in the manufacture of a medicament for treating at least one disease state selected from anemia, iron deficiency anemia, anemia of chronic disease, fibrodysplasia ossificans progressiva (FOP), cancer, inflammation, pathologic bone function, ectopic or maladaptive bone formation, a skin disease, hypertension, ventricular hypertrophy, atherosclerosis, spinal cord injury and neuropathy, heart disease, heart damage, liver damage, and liver disease.
13. The use of claim 12, wherein the cancer is breast cancer, prostate cancer, bone cancer, lung cancer, renal cell cancer, or acute megakaryoblastic leukemia.
14. An *in vitro* method for at least one of propagation, engraftment, and differentiation of progenitor cells, the method comprising contacting a progenitor cell with a compound of any one of claims 1-9 in a dosage and amount effective to propagate a cell.

15. A compound having a structure represented by the following formula:



wherein:

X₁ is N or CR₁;

X₂, and X₄ are independently N or CR₁;

X₃ is C;

Y₁, Y₂, and Y₃ are independently N or CR₁;

D is C or N;

W is N or O;

W₁ is N, O or C;


Cy is optionally substituted with one or more R₁, and is selected from C₃-C₁₂ cycloalkyl or C₃-C₁₂ cycloalkenyl or aryl or heteroaryl or C₃-C₁₂ heterocycloalkyl or C₃-C₁₂ heterocycloalkenyl or 3-8 membered ring comprising C, O, S, and/or N.

R₁-R₄ are independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, OC₁₋₁₀ alkyl (which may contain a C₃₋₈ membered ring containing C, O, S or N, optionally

substituted with one or more R^4), NR_1C_{1-10} alkyl (which may contain a C_{3-8} membered ring containing C, O, S or N, optionally substituted with one or more R^4), CN or CF_3 ;

Z is optionally substituted with one or more R_1 , and is selected from C_3-C_{12} cycloalkyl or C_3-C_{12} cycloalkenyl or aryl or heteroaryl or C_3-C_{12} heterocycloalkyl or C_3-C_{12} heterocycloalkenyl or 3-8 membered ring comprising C, O, S, and/or N.

16. A compound having a structure represented by the following formula:



wherein:

X_1 is N or CR_1 ;

X_2 , and X_4 are independently N or CR_1 ;

X_3 is C;


Y_1 , Y_2 , and Y_3 are independently N or CR_1 ;

Cy is a bond, or forms a ring that is optionally substituted with one or more R_1 , and is selected from C_3-C_{12} cycloalkyl or C_3-C_{12} cycloalkenyl or aryl or heteroaryl or C_3-C_{12} heterocycloalkyl or C_3-C_{12} heterocycloalkenyl or 3-8 membered ring comprising C, O, S, and/or N.

R_1-R_4 are independently selected from hydrogen, halogen, C_{1-10} alkyl, C_{3-10} cycloalkyl, OC_{1-10} alkyl (which may contain a C_{3-8} membered ring containing C, O, S or N, optionally substituted with one or more R^4), NR_1C_{1-10} alkyl (which may contain a C_{3-8} membered ring containing C, O, S or N, optionally substituted with one or more R^4), CN or CF_3 ;

Z is optionally substituted with one or more R_1 , and is selected from C_3-C_{12} cycloalkyl or C_3-C_{12} cycloalkenyl or aryl or heteroaryl or C_3-C_{12} heterocycloalkyl or C_3-C_{12} heterocycloalkenyl or 3-8 membered ring comprising C, O, S, and/or N.

17. A compound having a structure represented by the following formula:



wherein:

X₁, X₂, X₃ and X₄ are independently N or CR₁;

X is CR₁;


X₃ is C or N;

Y₁, Y₂, and Y₃ are independently N or CR₁;

R₁-R₄ are absent or independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, OC₁₋₁₀ alkyl (which may contain a C₃₋₈ membered ring containing C, O, S or N, optionally substituted with one or more R⁴), CN or CF₃;

Z is optionally substituted with one or more R₁, and is selected from C₃-C₁₂ cycloalkyl or C₃-C₁₂ cycloalkenyl or aryl or heteroaryl or C₃-C₁₂ heterocycloalkyl or C₃-C₁₂ heterocycloalkenyl or 3-8 membered ring comprising C, O, S, and/or N.

18. The compound of claim 17, wherein the compound is of the following formula:



wherein:

X₁, X₂, X₃ and X₄ are independently N or CR₁;


X is CR₁;

X₃ is C or N;

Y₁, Y₂, and Y₃ are independently N or CR₁;

R₁-R₄ are absent or independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, OC₁₋₁₀ alkyl (which may contain a C₃₋₈ membered ring containing C, O, S or N, optionally substituted with one or more R⁴), CN or CF₃;

19. The compound of claim 17, wherein the compound is of the following formula:



wherein


- (i) X₁ is N, X₄ is N, and X₂, X₃, X₄, Y₁, Y₂, and Y₃ are independently C or CR₁;
- (ii) X₁ is N, X₂ is N, Y₁ is N, and X, X₃, X₄, Y₂, and Y₃ are independently C or CR₁;
- (iii) X₁ is N, X₂ is N, Y₂ is N, and X, X₃, X₄, Y₂, and Y₃ are independently C or CR₁;

or

- (iv) X₁ is N, X₃ is N, and X₂, X₃, X₄, Y₁, Y₂, and Y₃ are independently C or CR₁;
Y₁, Y₂, and Y₃ are independently N or CR₁;


R₁-R₄ are independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, OC₁₋₁₀ alkyl (which may contain a C₃₋₈ membered ring containing C, O, S or N, optionally substituted with one or more R⁴), CN or CF₃.

20. The compound of claim 17, wherein the compound is of the following formula:




wherein R₁-R₄ are independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, OC₁₋₁₀ alkyl (which may contain a C₃₋₈ membered ring containing C, O, S or N, optionally substituted with one or more R⁴), CN or CF₃.

21. The compound of claim 17, wherein the compound is of the following formula:




wherein R₁-R₄ are independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, OC₁₋₁₀ alkyl (which may contain a C₃₋₈ membered ring containing C, O, S or N, optionally substituted with one or more R⁴), CN or CF₃.

22. The compound of claim 17, wherein the compound is of the following formula:



wherein R₁-R₄ are independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, OC₁₋₁₀ alkyl (which may contain a C₃₋₈ membered ring containing C, O, S or N, optionally substituted with one or more R⁴), CN or CF₃.


23. The compound of claim 17, wherein the compound is of the following formula:



wherein Y₁ and Y₃ are independently N or CR₁; R₁-R₄ are independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, OC₁₋₁₀ alkyl (which may contain a C₃₋₈

membered ring containing C, O, S or N, optionally substituted with one or more R⁴), CN or CF₃.

24. The compound of claim 17, wherein the compound is of the following formula:



wherein Y₁ and Y₃ are independently N or CR₁; R₁-R₄ are independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, OC₁₋₁₀ alkyl (which may contain a C₃₋₈ membered ring containing C, O, S or N, optionally substituted with one or more R⁴), CN or CF₃.

25. A compound of any one of claims 15-24 for use in therapy.

26. Use of a compound of any one of claims 15-24 in the manufacture of a medicament for modulating the BMP signaling pathway.


27. Use of a compound of any one of claims 1-9 in the manufacture of a medicament for promoting, inducing, or enhancing modulation of epicardial activation and epithelial-to-mesenchyme transition; or inducing, enhancing, or promoting cardiomyocyte formation.

28. A method for modulating the BMP signaling pathway comprising administering to at least one cell *in vitro* a compound of any one of claims 1-9 in an amount effective to modulate BMP signaling.
29. The method of claim 28, wherein modulating the BMP signaling pathway enhances keratinocyte proliferation.
30. The method of claim 28, wherein the at least one cell is a stem cell.
31. The method of claim 28, wherein modulating the BMP signaling pathway regulates the differentiation and regeneration of precursor and stem cell populations.
32. The method of claim 28, wherein modulating the BMP signaling pathway maintains a pluripotential state in stem cell or multipotent cell populations *in vitro*.

33. The method of claim 28, wherein mutilating the BMP signaling pathway expands stem cell or multipotent cell populations *in vitro*.

We claim:

1. — A compound for use in the treatment of a disease state associated with modulating the BMP signaling pathway in a subject, the compound having a structure represented by the following formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂, S(O)₀₋₂, NR₁R₂, or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;


D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;

E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;

R₁ is selected from H, alkyl, aryl, heteraryl, or COR₁; and

R₂ is selected from H, alkyl, aryl, heteraryl, COR₁, and further, R₁ and R₂ can form a C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

2. — A compound for use as a medicament in modulating the BMP signaling pathway, the compound having a structure represented by the formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂, S(O)₀₋₂NR₁R₂, or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;


D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;

E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;

R₁ is selected from H, alkyl, aryl, heteraryl, or COR₁; and

R₂ is selected from H, alkyl, aryl, heteraryl, COR₁, and further, R₁ and R₂ can form a C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

3. — A compound for use in treating at least one of anemia, including iron deficiency, and anemia of chronic disease, the compound having a structure represented by the following formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂, S(O)₀₋₂NR₁R₂, or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;


D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;

E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;

R₁ is selected from H, alkyl, aryl, heteraryl, or COR₁; and

R_2 is selected from H, alkyl, aryl, heteraryl, COR_1 , and further, R_1 and R_2 can form a C_3 - C_{12} cycloalkyl or C_3 - C_{12} cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

4. — A compound for use in the treatment of fibrodysplasia ossificans progressiva (FOP), the compound having a structure represented by the following formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF_3 , halogen, CN, alkyl, aryl, heteroaryl, NR_1R_2 , COR_1R_2 , CR_1R_2 , $S(O)_{0-2}NR_1R_2$; or $S(O)_{0-2}R_1R_2$;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;


D is selected from a bond, O, CR_1R_2 or NH or NR_1 or NR_1R_2 or $S(O)_{0-2}R_1R_2$;

E is selected from H, CF_3 , halogen, CN, alkyl, aryl, heteroaryl, C_3 - C_{12} cycloalkyl or C_3 - C_{12} cycloheteroalkyl or $(CH_2)_x$ - C_3 - C_{12} cycloalkyl or $(CH_2)_x$ - C_3 - C_{12} cycloheteroalkyl;

R_1 is selected from H, alkyl, aryl, heteraryl, or COR_1 ; and

R_2 is selected from H, alkyl, aryl, heteraryl, COR_1 , and further, R_1 and R_2 can form a C_3 - C_{12} cycloalkyl or C_3 - C_{12} cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

5. — A compound for use as a medicament in inhibiting the ontogenesis, growth, or metastasis of solid tumors, including breast, prostate carcinomas, bone, lung, and renal cell carcinomas, the compound having a structure represented by the following formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF_3 , halogen, CN, alkyl, aryl, heteroaryl, NR_1R_2 , COR_1R_2 , CR_1R_2 , $\text{S(O)}_{0-2}\text{NR}_1\text{R}_2$; or $\text{S(O)}_{0-2}\text{R}_1\text{R}_2$;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;


D is selected from a bond, O, CR_1R_2 or NH or NR_1 or NR_1R_2 or $\text{S(O)}_{0-2}\text{R}_1\text{R}_2$;

E is selected from H, CF_3 , halogen, CN, alkyl, aryl, heteroaryl, $\text{C}_3\text{-C}_{12}$ -cycloalkyl or $\text{C}_3\text{-C}_{12}$ -cycloheteroalkyl or $(\text{CH}_2)_x\text{-C}_3\text{-C}_{12}$ -cycloalkyl or $(\text{CH}_2)_x\text{-C}_3\text{-C}_{12}$ -cycloheteroalkyl;

R_1 is selected from H, alkyl, aryl, heteraryl, or COR_1 ; and

R_2 is selected from H, alkyl, aryl, heteraryl, COR_1 , and further, R_1 and R_2 can form a $\text{C}_3\text{-C}_{12}$ -cycloalkyl or $\text{C}_3\text{-C}_{12}$ -cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

6. — A compound for use as a medicament for augmenting the inflammatory or immune response, the compound having a structure represented by the following formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;


R is selected from H, CF_3 , halogen, CN, alkyl, aryl, heteroaryl, NR_1R_2 , COR_1R_2 , CR_1R_2 , $\text{S(O)}_{0-2}\text{NR}_1\text{R}_2$; or $\text{S(O)}_{0-2}\text{R}_1\text{R}_2$;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;

D is selected from a bond, O, CR_1R_2 or NH or NR_1 or NR_1R_2 or $\text{S(O)}_{0-2}\text{R}_1\text{R}_2$;

E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;
R₁ is selected from H, alkyl, aryl, heteraryl, or COR₁; and
R₂ is selected from H, alkyl, aryl, heteraryl, COR₁, and further, R₁ and R₂ can form a C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

7. — A compound for use as a medicament for treating pathologic bone function, the compound having a structure represented by the following formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂, S(O)₀₋₂NR₁R₂; or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;


D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;

E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;

R₁ is selected from H, alkyl, aryl, heteraryl, or COR₁; and

R₂ is selected from H, alkyl, aryl, heteraryl, COR₁, and further, R₁ and R₂ can form a C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

8. — A compound for use as a medicament for treating ectopic or maladaptive bone formation in a subject in need thereof, the compound having a structure represented by the following formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂, S(O)₀₋₂NR₁R₂; or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;


D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;

E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;

R₁ is selected from H, alkyl, aryl, heteraryl, or COR₁; and

R₂ is selected from H, alkyl, aryl, heteraryl, COR₁, and further, R₁ and R₂ can form a C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

9. — A compound for use as a medicament for treating a skin disease in a subject in need thereof, the compound having a structure represented by the following formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;


R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂, S(O)₀₋₂NR₁R₂; or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;

D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;

E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;
R₁ is selected from H, alkyl, aryl, heteraryl, or COR₄; and
R₂ is selected from H, alkyl, aryl, heteraryl, COR₄, and further, R₁ and R₂ can form a C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

10. — A compound for use as a medicament for treating hypertension in a subject in need thereof, the compound having a structure represented by the following formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂, S(O)₀₋₂NR₁R₂; or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;


D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;

E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;

R₁ is selected from H, alkyl, aryl, heteraryl, or COR₄; and

R₂ is selected from H, alkyl, aryl, heteraryl, COR₄, and further, R₁ and R₂ can form a C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

11. — A compound for use as a medicament for treating ventricular hypertrophy in a subject in need thereof, the compound having a structure represented by the following formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂, S(O)₀₋₂NR₁R₂; or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;


D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;

E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;

R₁ is selected from H, alkyl, aryl, heteraryl, or COR₁; and

R₂ is selected from H, alkyl, aryl, heteraryl, COR₁, and further, R₁ and R₂ can form a C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

12. — A compound for use as a medicament for treating spinal cord injury and neuropathy in a subject in need thereof, the compound having a structure represented by the following formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂, S(O)₀₋₂NR₁R₂; or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;


D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;

E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;

R₁ is selected from H, alkyl, aryl, heteraryl, or COR₄; and

R₂ is selected from H, alkyl, aryl, heteraryl, COR₄, and further, R₁ and R₂ can form a C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

13. — A compound for use as a medicament for treating atherosclerosis in a subject in need thereof, the compound having a structure represented by the following formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂, S(O)₀₋₂NR₁R₂; or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;


D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;

E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;

R₁ is selected from H, alkyl, aryl, heteraryl, or COR₄; and

R₂ is selected from H, alkyl, aryl, heteraryl, COR₄, and further, R₁ and R₂ can form a C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

14. — A compound for use in therapy for at least one of propagation, engraftment, and differentiation of progenitor cells, the compound having a structure represented by the following formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂, S(O)₀₋₂, NR₁R₂; or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;

D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;


E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;

R₁ is selected from H, alkyl, aryl, heteraryl, or COR₁; and

R₂ is selected from H, alkyl, aryl, heteraryl, COR₁, and further, R₁ and R₂ can form a C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.


We claim:

145. A compound having a structure represented by the following formula (I):



wherein:

W, X, Y, and Z are independently N or CH;



each A₁ is independently O, S, CR₁R₂, or NR₁;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂, S(O)₀₋₂NR₁R₂, or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;


D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;

E is absent or selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂ cycloalkyl or C₃-C₁₂ cycloheteroalkyl or -(CH₂)_x-C₃-C₁₂ cycloalkyl or -(CH₂)_x-C₃-C₁₂ cycloheteroalkyl;


R₁ is absent or selected from H, alkyl, aryl, or heteraryl, ~~or COR₁~~; and

R₂ is selected from H, alkyl, aryl, heteroaryl, or COR₁, ~~and further, or~~ R₁ and R₂ can form a C₃-C₁₂ cycloalkyl or C₃-C₁₂ cycloheteroalkyl containing O, N and/or S; or a pharmaceutically acceptable salt thereof, ~~or a pharmaceutically acceptable derivative thereof.~~


~~162.~~ A ~~The~~ compound of claim ~~151~~, wherein W, X, Y and Z ~~are selected together help~~ form:



~~173.~~ A ~~The~~ compound of claim ~~151~~, wherein W, X, Y and Z ~~are selected together help~~ form:



~~18.~~ A compound of claim 15, wherein



wherein A₁ is independently O, CR₁R₂ or NH or NR₁ or NR₁R₂, or can join with another A₁ to form C₃-C₁₂ cycloalkyl or C₃-C₁₂ cycloalkenyl or aryl or heteroaryl or C₃-C₁₂ heterocycloalkyl or C₃-C₁₂ heterocycloalkenyl or 3-8 membered ring comprising C, O, S, and/or N.

419. ~~A~~ The compound of claim ~~451~~, wherein M is optionally substituted with one or more R, and is selected from C₃-C₁₂ cycloalkyl or C₃-C₁₂ cycloalkenyl or aryl or heteroaryl or C₃-C₁₂


heterocycloalkyl or C₃-C₁₂ heterocycloalkenyl or 3-8 membered ring comprising C, O, S, and/or N.


205. A The compound of claim 194, wherein M is optionally substituted phenyl or pyridine.







216. A The compound of claim 151, wherein M, D, and E together form:

227. A The compound of claim 151, wherein A is chosen from the following:

238. A compound of one claim 15, wherein the compound is of the following formulastructures:






or a pharmaceutically acceptable salt thereof.

9. The compound of claim 1, wherein X is N and Y is CH.
10. A compound of any one of claims 1-9 for use in therapy.
11. The compound of claim 10, wherein the therapy is associated with modulating the BMP signaling pathway.
12. Use of a compound of any one of claims 1-9 in the manufacture of a medicament for treating at least one disease state selected from anemia, iron deficiency anemia, anemia of chronic disease, fibrodysplasia ossificans progressiva (FOP), cancer, inflammation, pathologic bone function, ectopic or maladaptive bone formation, a skin disease, hypertension, ventricular hypertrophy, atherosclerosis, spinal cord injury and neuropathy, heart disease, heart damage, liver damage, and liver disease.
13. The use of claim 12, wherein the cancer is breast cancer, prostate cancer, bone cancer, lung cancer, renal cell cancer, or acute megakaryoblastic leukemia.
14. An *in vitro* method for at least one of propagation, engraftment, and differentiation of progenitor cells, the method comprising contacting a progenitor cell with a compound of any one of claims 1-9 in a dosage and amount effective to propagate a cell.
24. ~~A compound having a structure represented by the following formula (II):~~



wherein:

X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂, S(O)₀₋₂NR₁R₂; or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;


D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;

E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;

R₁ is selected from H, alkyl, aryl, heteraryl, or COR₁; and

R₂ is selected from H, alkyl, aryl, heteraryl, COR₁, and further, R₁ and R₂ can form a C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

25. — A compound having a structure represented by the following formula (III):



wherein:

Z is N or CH;

A is substituted or unsubstituted and is selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂, S(O)₀₋₂NR₁R₂; or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and selected from aryl or heteroaryl;


D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;

E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;

R₁ is selected from H, alkyl, aryl, heteraryl, or COR₁; and

R_2 is selected from H, alkyl, aryl, heteroaryl, COR_1 , and further, R_1 and R_2 can form a C_3 - C_{12} cycloalkyl or C_3 - C_{12} cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

2615. A compound for use as a medicament for modulating the BMP signaling pathway, the compound having a structure represented by the following formula:



wherein:

X_1 is N , O , or CR_1 ;

X_2 , and X_4 are independently N or CR_1 ;

X_3 is C or N ;

Y_1 , Y_2 , and Y_3 are independently N or CR_1 ;

D is C or N ;

W is N or O ;

W_1 is N , O or C ;


Cy is optionally substituted with one or more R_1 , and is selected from C_3 - C_{12} cycloalkyl or C_3 - C_{12} cycloalkenyl or aryl or heteroaryl or C_3 - C_{12} heterocycloalkyl or C_3 - C_{12} heterocycloalkenyl or 3-8 membered ring comprising C, O, S, and/or N.

R_1 - R_4 are independently selected from hydrogen, halogen, C_{1-10} alkyl, C_{3-10} cycloalkyl, OC_{1-10} alkyl (which may contain a C_{3-8} membered ring containing C, O, S or N, optionally

substituted with one or more R^4), NR_1C_{1-10} alkyl (which may contain a C_{3-8} membered ring containing C, O, S or N, optionally substituted with one or more R^4), CN or CF_3 ;

Z is optionally substituted with one or more R_1 , and is selected from C_3-C_{12} cycloalkyl or C_3-C_{12} cycloalkenyl or aryl or heteroaryl or C_3-C_{12} heterocycloalkyl or C_3-C_{12} heterocycloalkenyl or 3-8 membered ring comprising C, O, S, and/or N.

2716. A compound for use as a medicament for modulating the BMP signaling pathway, the compound having a structure represented by the following formula:



wherein:

X_1 is N , O , or CR_1 ;

X_2 , and X_4 are independently N or CR_1 ;

X_3 is C or N;


Y_1 , Y_2 , and Y_3 are independently N or CR_1 ;

Cy is a bond, or forms a ring that is optionally substituted with one or more R_1 , and is selected from C_3-C_{12} cycloalkyl or C_3-C_{12} cycloalkenyl or aryl or heteroaryl or C_3-C_{12} heterocycloalkyl or C_3-C_{12} heterocycloalkenyl or 3-8 membered ring comprising C, O, S, and/or N.

R_1-R_4 are independently selected from hydrogen, halogen, C_{1-10} alkyl, C_{3-10} cycloalkyl, OC_{1-10} alkyl (which may contain a C_{3-8} membered ring containing C, O, S or N, optionally substituted with one or more R^4), NR_1C_{1-10} alkyl (which may contain a C_{3-8} membered ring containing C, O, S or N, optionally substituted with one or more R^4), CN or CF_3 ;

Z is optionally substituted with one or more R_1 , and is selected from C_3-C_{12} cycloalkyl or C_3-C_{12} cycloalkenyl or aryl or heteroaryl or C_3-C_{12} heterocycloalkyl or C_3-C_{12} heterocycloalkenyl or 3-8 membered ring comprising C, O, S, and/or N.

~~2817.~~ A compound for use as a medicament for modulating the BMP signaling pathway, the compound having a structure represented by the following formula:



wherein:

X₁, X₂, X₃ and X₄ are independently N or CR₁;

X is ~~N, O, or~~ CR₁;


X₃ is C or N;

Y₁, Y₂, and Y₃ are independently N or CR₁;

R₁-R₄ are absent or independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, OC₁₋₁₀ alkyl (which may contain a C₃₋₈ membered ring containing C, O, S or N, optionally substituted with one or more R⁴), CN or CF₃;

Z is optionally substituted with one or more R₁, and is selected from C₃-C₁₂ cycloalkyl or C₃-C₁₂ cycloalkenyl or aryl or heteroaryl or C₃-C₁₂ heterocycloalkyl or C₃-C₁₂ heterocycloalkenyl or 3-8 membered ring comprising C, O, S, and/or N.

~~1829.~~ The compound of claim ~~1728~~, wherein the compound is of the following formula:



wherein:

X₁, X₂, X₃ and X₄ are independently N or CR₁;


X is ~~N, O, or~~ CR₁;

X₃ is C or N;

Y₁, Y₂, and Y₃ are independently N or CR₁;

R₁-R₄ are absent or independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, OC₁₋₁₀ alkyl (which may contain a C₃₋₈ membered ring containing C, O, S or N, optionally substituted with one or more R⁴), CN or CF₃;

~~30~~19. The compound of claim ~~28~~17, wherein the compound is of the following formula:




wherein

- ~~(i)~~ (i) X₁ is O, X₂ is N, and X, X₃, X₄, Y₁, Y₂, and Y₃ are independently C or CR₁;
- ~~(ii)~~ (ii) X₁ is O, X₂ is N, Y₁ is N, and X, X₃, X₄, Y₂, and Y₃ are independently C or CR₁;
- ~~(iii)~~ (iii) X₁ is N, X₄ is N, and X₂, X₃, X₄, Y₁, Y₂, and Y₃ are independently C or CR₁;
- ~~(iv)~~ (iv) X₁ is N, X₂ is N, Y₁ is N, and X, X₃, X₄, Y₂, and Y₃ are independently C or CR₁;
- ~~(v)~~ (v) X₁ is N, X₂ is N, Y₂ is N, and X, X₃, X₄, Y₂, and Y₃ are independently C or CR₁;
- ~~(vi)~~ (vi) X₁ is N, X₃ is N, and X₂, X₃, X₄, Y₁, Y₂, and Y₃ are independently C or CR₁;
- ~~(vii)~~ (vii) X₁ is N, X₃ is N, and X₂, X₃, X₄, Y₁, Y₂, and Y₃ are independently N or CR₁;


R₁-R₄ are independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, OC₁₋₁₀ alkyl (which may contain a C₃₋₈ membered ring containing C, O, S or N, optionally substituted with one or more R⁴), CN or CF₃.

~~31.~~ The compound of claim ~~28~~, of the following formula:




wherein R_1 - R_4 are independently selected from hydrogen, halogen, C_{1-10} alkyl, C_{3-10} cycloalkyl, OC_{1-10} alkyl (which may contain a C_{3-8} membered ring containing C, O, S or N, optionally substituted with one or more R^4), CN or CF_3 .

32. The compound of claim 28, wherein the compound is of the following formula:




wherein R_1 - R_4 are independently selected from hydrogen, halogen, C_{1-10} alkyl, C_{3-10} cycloalkyl, OC_{1-10} alkyl (which may contain a C_{3-8} membered ring containing C, O, S or N, optionally substituted with one or more R^4), CN or CF_3 .

2033. The compound of claim 1728, wherein the compound is of the following formula:




wherein R_1 - R_4 are independently selected from hydrogen, halogen, C_{1-10} alkyl, C_{3-10} cycloalkyl, OC_{1-10} alkyl (which may contain a C_{3-8} membered ring containing C, O, S or N, optionally substituted with one or more R^4), CN or CF_3 .

3421. The compound of claim 2817, wherein the compound is of the following formula:




wherein R₁-R₄ are independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, OC₁₋₁₀ alkyl (which may contain a C₃₋₈ membered ring containing C, O, S or N, optionally substituted with one or more R⁴), CN or CF₃.

~~3522.~~ The compound of claim ~~2817~~, wherein the compound is of the following formula:




wherein R₁-R₄ are independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, OC₁₋₁₀ alkyl (which may contain a C₃₋₈ membered ring containing C, O, S or N, optionally substituted with one or more R⁴), CN or CF₃.

~~36.~~ The compound of claim ~~28~~, wherein the compound is of the following formula:



wherein Y₁ and Y₃ are independently N or CR₁; R₁-R₄ are independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, OC₁₋₁₀ alkyl (which may contain a C₃₋₈ membered ring containing C, O, S or N, optionally substituted with one or more R⁴), CN or CF₃.


~~3723.~~ The compound of claim ~~2817~~, wherein the compound is of the following formula:



wherein Y₁ and Y₃ are independently N or CR₁; R₁-R₄ are independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, OC₁₋₁₀ alkyl (which may contain a C₃₋₈

membered ring containing C, O, S or N, optionally substituted with one or more R⁴), CN or CF₃.

~~38~~24. The compound of claim ~~28~~17, wherein the compound is of the following formula:




wherein Y₁ and Y₃ are independently N or CR₁; R₁-R₄ are independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, OC₁₋₁₀ alkyl (which may contain a C₃₋₈ membered ring containing C, O, S or N, optionally substituted with one or more R⁴), CN or CF₃.

25. A compound of any one of claims 15-24 for use in therapy.

26. Use of a compound of any one of claims 15-24 in the manufacture of a medicament for modulating the BMP signaling pathway.

39. — A compound for use as a medicament for treating acute megakaryoblastic leukemia, the compound having a structure represented by the following formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂, S(O)₀₋₂NR₁R₂, or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;


D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;

E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;

R₁ is selected from H, alkyl, aryl, heteraryl, or COR₁; and

R₂ is selected from H, alkyl, aryl, heteraryl, COR₁, and further, R₁ and R₂ can form a C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

4027. Use of A-a compound of any one of claims 1-9 in the manufacture of a medicament for use in therapy for at least one of treating heart disease, treating heart damage, promoting, inducing, or enhancing modulation of epicardial activation and epithelial-to-mesenchyme transition; or a method of inducing, enhancing, or promoting cardiomyocyte formation, the compound having a structure represented by the following formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂,
S(O)₀₋₂NR₁R₂; or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;


D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;

E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;

R₁ is selected from H, alkyl, aryl, heteraryl, or COR₁; and

R₂ is selected from H, alkyl, aryl, heteraryl, COR₁; and further, R₁ and R₂ can form a C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

41. — A compound for use in therapy for at least one of promoting or enhancing liver restoration of liver mass, treating liver damage, treating liver disease, or treating acetaminophen overdose in a subject in need thereof, the compound having a structure represented by the following formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂, S(O)₀₋₂NR₁R₂; or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;


D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;

E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;

R₁ is selected from H, alkyl, aryl, heteraryl, or COR₁; and

R₂ is selected from H, alkyl, aryl, heteraryl, COR₁; and further, R₁ and R₂ can form a C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

4228. A method for modulating the BMP signaling pathway comprising administering ~~in~~ *in vitro* administration to at least one cell *in vitro* a compound of any one of claims 1-9 in an amount effective to modulate BMP signaling, the compound having a structure represented by the formula (I):



wherein:

W, X, Y, and Z are independently N or CH;

A is substituted or unsubstituted and selected from cycloalkyl, heterocycloalkyl, aryl or heteroaryl;

R is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, NR₁R₂, COR₁R₂, CR₁R₂, S(O)₀₋₂NR₁R₂, or S(O)₀₋₂R₁R₂;

M is substituted or unsubstituted and is selected from aryl or heteroaryl;

D is selected from a bond, O, CR₁R₂ or NH or NR₁ or NR₁R₂ or S(O)₀₋₂R₁R₂;

E is selected from H, CF₃, halogen, CN, alkyl, aryl, heteroaryl, C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl or (CH₂)_x-C₃-C₁₂-cycloalkyl or (CH₂)_x-C₃-C₁₂-cycloheteroalkyl;

R₁ is selected from H, alkyl, aryl, heteraryl, or COR₁; and

R₂ is selected from H, alkyl, aryl, heteraryl, COR₁, and further, R₁ and R₂ can form a C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloheteroalkyl containing O, N or S; or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable derivative thereof.

4329. The method of claim 4228, wherein modulating the BMP signaling pathway enhances keratinocyte proliferation.

4430. The method of claim 4228, wherein the at least one cell is a stem cell.

4531. The method of claim 4428, wherein modulating the BMP signaling pathway regulates the differentiation and regeneration of precursor and stem cell populations.

4632. The method of claim 4228, wherein mutilating the BMP signaling pathway maintains a pluripotential state in stem cell or multipotent cell populations *in vitro*.

4733. The method of claim 4228, wherein mutilating the BMP signaling pathway expands stem cell or multipotent cell populations *in vitro*.