

Claims

1. A plasmid combination, wherein the plasmid combination comprises plasmids Seq1, PMD2.G, pMDLg-pRRE and pRSV-Rev; and the plasmids Seq1, PMD2.G, pMDLg-pRRE and pRSV-Rev are present at a ratio of 2-6: 1-1.5: 1-3: 1-1.5.
2. The plasmid combination according to claim 1, wherein the plasmid Seq1 comprises a nucleic acid molecule encoding the chimeric antigen receptor, wherein the chimeric antigen receptor comprises an amino acid sequence shown in SEQ ID NO. 1.
3. The plasmid combination according to claim 1, wherein the plasmid Seq1 comprises a nucleic acid molecule encoding a chimeric antigen receptor, wherein the nucleic acid molecule comprises a nucleic acid sequence shown in SEQ ID NO. 2.
4. The plasmid combination according to claim 1, wherein the plasmids Seq1, PMD2.G, pMDLg-pRRE and pRSV-Rev are present at a ratio of 11.8: 3.53: 6.33: 2.3, 13.8: 3.48: 5.31: 2.54 or 14: 4.67: 4.67: 4.67.
5. A method for preparing a lentiviral vector, the method comprising introducing the plasmid combination according to any of claims 1 to 4 into a cell.
6. The method according to claim 5, wherein the cell is 293T.
7. A method for preparing a modified immune effector cell, comprising preparing the lentiviral vector according to the method according to either of claims 5-6.
8. The method according to claim 7, comprising introducing the lentiviral vector into an immune effector cell.
9. The method according to claim 8, wherein the immune effector cell is selected from the group consisting of a T lymphocyte and a natural killer cell.
10. A modified immune effector cell prepared by the method according to any of claims 7 to 9.
11. The modified immune effector cell according to claim 10, wherein the immune effector cell can express the chimeric antigen receptor.
12. A composition comprising the modified immune effector cell according to either of claims 10-11.
13. Use of the immune effector cell according to either of claims 10-11 and/or the composition according to claim 12 in the manufacture of a medicament, wherein the medicament is useful for the treatment of a disease or disorder associated with CD19 expression.
14. The use according to claim 13, wherein the disease or disorder associated with CD19

expression comprises non-solid tumors.

15. The use according to claim 14, wherein the non-solid tumor comprises leukemia and/or lymphoma.

16. The use according to claim 13, wherein the disease or disorder associated with CD19 expression comprises acute lymphoblastic leukemia and/or B-cell lymphoma.

17. The use according to claim 16, wherein the acute lymphoblastic leukemia comprises acute lymphoblastic leukemia in adults and/or acute lymphoblastic leukemia in children.

18. The use according to claim 17, wherein the medicament is administered at a dose of 0.25×10^8 to 0.5×10^8 CAR-positive T cells.

19. The use according to claim 16, wherein the B-cell lymphoma comprises non-Hodgkin's lymphoma.

20. The use according to claim 19, wherein the medicament is administered at a dose of 1×10^8 to 2×10^8 CAR-positive T cells.

21. The immune effector cell according to either of claims 10-11 and/or the composition according to claim 12, for use in the treatment of a disease or disorder associated with CD19 expression.

22. The composition and/or the immune effector cell according to claim 21, wherein the disease or disorder associated with CD19 expression comprises non-solid tumors.

23. The composition and/or the immune effector cell according to claim 22, wherein the non-solid tumor comprises leukemia and/or lymphoma.

24. The composition and/or the immune effector cell according to claim 21, wherein the disease or disorder associated with CD19 expression comprises acute lymphoblastic leukemia and/or B-cell lymphoma.

25. The composition and/or the immune effector cell according to claim 24, wherein the acute lymphoblastic leukemia comprises acute lymphoblastic leukemia in adults and/or acute lymphoblastic leukemia in children.

26. The composition and/or the immune effector cell according to claim 25, wherein the administration is performed at a dose of 0.25×10^8 to 0.5×10^8 CAR-positive T cells.

27. The composition and/or the immune effector cell according to claim 24, wherein the B-cell lymphoma comprises non-Hodgkin's lymphoma.

28. The composition and/or the immune effector cell according to claim 27, wherein the administration is performed at a dose of 1×10^8 to 2×10^8 CAR-positive T cells.