

Description

This invention relates to hair cosmetic compositions and more particularly, to hair cosmetic compositions of the type which comprise extracts obtained by polar solvent extraction of plants and polypeptide compounds and which can impart, to hair, a suitable degree of set retentivity and good feeling to the touch.

Hair style is one of the most important charming points from the standpoint of beauty care and a variety of beauty treatments have been made. For instance, several techniques of suitably waving the hair using hair cosmetics are known including so-called permanent treatments by permanent waving techniques and transient treatments using set lotions or hair sprays. By the treatments, the hair is suitably set or dressed. However, the treatments by the permanent waving techniques may sometimes damage the hair considerably to such an extent that the hair cannot refresh. On the other hand, the treatments using set lotions or hair sprays are disadvantageous in that the hair is merely set transiently and thus the once set hair is broken simply by the action of moisture. Hence, both types of treatments are not satisfactory. Moreover, these techniques have the tendency of stiffening the hair and are not also satisfactory in the touch of the hair.

Shampoos are extensively used for the purpose of washing away soiling attached on the hair. However, currently available shampoos also wash away oil necessary for the hair, so that the shampooed hair become dry and split-ends or broken hairs are undesirably involved upon brushing. To avoid this, hair rinses or pre-shampoos are used, of which some contain oils or fats, causing the hair sticky.

DE—A—2 262 375 discloses a hair shampoo containing a decomposition product of collagen.

In AT—A—175 976 a hair growth composition is used comprising a nettle or burdock root extract.

The hair treatment composition of GB—A—2 088 210 comprises protein per se. The latter also applies to "Kosmetik-Parfum-Drogen Rundschau", page 120, left column of 1970.

In order to obtain hair cosmetic compositions having excellent properties, we have made extensive studies on ingredients of the cosmetics. As a result, it was found that when a combination of a specific type of plant extract and a polypeptide is admixed with hair cosmetic compositions such as shampoo, rinse, set lotion and hair spray, the resulting cosmetic compositions can impart good hair style retentivity and a suitable degree of feeling to the touch with ease in combing.

According to the present invention, there is provided a hair cosmetic composition comprising

(A) an extract obtained by polar solvent extraction of birch or rosemary, and

(B) a polypeptide compound having a molecular weight of 350 to 30,000 selected from decomposition products of silk, decomposition products of keratin and derivatives thereof.

The (A) ingredient useful in the present invention is obtained by extraction of birch or rosemary.

Preferable sites of these plants for the extraction are, for example, bark of birch and an entire grass part of rosemary.

The plant extract is obtained by extracting flowers, leaves, fruit, roots, stems and the like with solvent at a normal temperature or under heating conditions according to any known extraction technique. The extraction solvents are polar organic solvents including, for example, lower alcohols such as methanol, ethanol and the like, propylene glycol, 1,3-butylene glycol and glycerine, and water. These solvents may be used singly or in combination.




The extracts of birch and rosemary are commercially available. The commercially sold birch extracts include Birch Extract made by Novarom Co., Ltd., Boulene MCF787 by Gattefossé Co., Ltd., Birch Extract by Virmin Co., Ltd., Birch Extract by Nowak Co., Ltd., Extrapone Birch Special by Dragoo Co., Ltd., and the like. Mentioned as rosemary extract are Rosemary Extract by Novarom Co. Ltd., Romarin MCF722 by Gattefossé Co. Ltd., Phytelene EG009 by Virmin Co. Ltd., and Rosemary Extract by Nowak Co., Ltd.

When the plant extract is in a liquid form, the ingredient (A) is used in an amount ranging from 0.001 to 10.0 wt%, preferably 0.01 to 1.0 wt% (hereinafter referred to simply as %) of the total composition, calculated as a residue obtained after distillation of extraction solvent therefrom. The ingredient (A) or plant extract may be directly added to hair cosmetic compositions in the form of liquid extract, or may be added after concentration to a desired level or after complete removal of extraction solvent from the extract.

The ingredient (B) used in the present invention are polypeptide compounds as defined hereinbefore.

Keratin is derived from, for example, animal hairs, human hairs, feathers, nails, horns, hooves or scales. These materials may be used as a decomposition derivative thereof. Preferably keratin materials are wool, human hair, and feathers.

The decomposition derivatives of keratin materials are hydrolyzates, oxidative decomposition products, and —SH group-modified compounds of reductive decomposition products. The hydrolysis, oxidation and reduction reactions and a subsequent modification reaction may be suitably combined. The hydrolysis techniques include an acid hydrolysis using hydrochloric acid, sulfuric acid, phosphoric acid or the like; and alkali hydrolysis using sodium hydroxide, sodium carbonate or the like; and an enzyme hydrolysis using protease. The oxidation and/or reduction reaction may be effected according to any known method. The disulfide bond is cleft, by the reduction reaction, into a thiol group (—SH). The thiol group can be modified by a known method to give the following groups:



Of these groups, —SCH₂COOH and —SCHCOOH obtained by the modification are preferably.

$$\begin{array}{c} | \\ \text{CH}_2\text{COOH} \end{array}$$

The chemical modification of the thiol group is conducted according to a method known per se as described, for example, in Textile Progress Vol. 7, page 1 (1975) by N. H. Leon, "Organic Sulfur Compounds" (1968) written by Shigeru Oae and published by Kagaku Dojin Co., Ltd., and "Course of Polymer Experimentation" Vol. 12 (1957) written by Masami Oku and published by Kyoritsu Pub. Co.

The decomposition products of keratin and derivatives thereof have a molecular weight of 350 to 30,000.

Silk fibres are one of the starting materials for the ingredient (B). Preferably, silk fibers are used as a product obtained by hydrolysis with an acid, alkali or enzyme. The hydrolyzate of silk fibres may be obtained by hydrolysis with an acid such as hydrochloric acid, sulfuric acid, phosphoric acid or the like, and an alkali such as sodium hydroxide, sodium carbonate or the like. Moreover, the acid or alkali hydrolysis may be used in combination with an enzyme hydrolysis. The resulting decomposition or hydrolysis product has a molecular weight of 350 to 30,000.

In the practice of the invention, the ingredient (B) is used in an amount of 0.01 to 10 wt%, preferably 0.1 to 5.0 wt%, of the hair cosmetic composition.

The mixing ratio of the ingredients (A) and (B) in the hair cosmetic composition of the invention is 0.1 to 10:1, preferably 1 to 8:1.

The term "hair cosmetic or cosmetics" used herein is intended to mean all cosmetics applicable to the hair. For instance, there are included pre-shampoo treatment agents, shampoos, hair rinses, after shampoo agents, conditioners, hair conditioners, setting lotions, blow styling lotions, hair sprays, hair dyes, bleaches, first permanent waving solutions, second permanent waving solutions, hair liquids or hair tonics.

The hair cosmetic compositions according to the invention are prepared by adding the essential ingredients (A) and (B) to known ingredients for these purposes which may vary depending on the type of hair cosmetic. The hair cosmetic composition may be formulated in various forms such as aqueous solution, ethanolic solution, emulsion, suspension gel, solid, aerosol or powder.

Preparation of the hair cosmetics of the invention are described.

(1) Shampoo:

Prepared by mixing known shampoo constituents by a usual manner along with one or more anionic active agents and the ingredients (A) and (B).

Preferable ionic active agents which are used as a shampoo substrate are mentioned below.

(1) Linear or branched alkylbenzenesulfonates having an alkyl group having from 10 to 16 carbon atoms on average.

(2) Polyoxyalkylene alkylsulfates having a linear or branched alkyl group having from 8 to 20 carbon atoms on average and added with 0.5 to 8 moles, on average, of ethylene oxide and/or propylene oxide in one molecule thereof.


(3) Alkylsulfates having an alkyl group having from 10 to 20 carbon atoms on average.

(4) Olefinsulfonates having from 10 to 20 carbon atoms on average in one molecule thereof.

(5) Alkanesulfonates having from 10 to 20 carbon atoms in one molecule thereof.

(6) Alkylethoxycarboxylates having an alkyl group having from 10 to 20 carbon atoms on average and added with from 0.5 to 8 moles of ethylene oxide on average in one molecule thereof.

(7) Succinic acid derivatives represented by the following formula



(in which R_1 represents an alkyl group or alkenyl group having from 6 to 20 carbon atoms, X_1 and Y_1 independently represent an ion).

Counter ions of these anionic active agents are, for example, alkali metal ions such as sodium, potassium and the like, alkaline earth metal ions such as calcium, magnesium and the like, ammonium ions, and alkanolamines having from 1 to 3 alkanol group having 2 or 3 carbon atoms and including, for example, monoethanolamine, diethanolamine, triethanolamine, triisopropanolamine and the like.

Of these anionic active agents, linear or branched alkylsulfates having from 10 to 16 carbon atoms on average, polyoxyethylene alkylsulfates (an average number of addition moles of 0.5 to 8) having an alkyl group having from 8 to 20 carbon atoms on average, and olefinsulfonates having from 10 to 16 carbon atoms on average are preferably used.

In the shampoo compositions of the invention, anionic active agents are generally used in an amount of from 5 to 30%, preferably 10 to 25%.

Other ingredients may be used in the shampoo composition of the invention in amounts not to impede the effect of the invention, which include amphoteric active agents, nonionic active agents, cationic active agents, solubilizers such as propylene glycol, glycerine, urea and the like, viscosity adjusters such as ethyl alcohol, isopropyl alcohol, hydroxyethyl cellulose, methyl cellulose, higher alcohols and the like, perfumes, colorants, UV absorbers, antioxidants, preservatives, pearling agents and lotionizing agents. These agents may be added as desired.

The thus obtained shampoo composition not only has excellent hair-conditioning and washing effects, but also is less irritative than known shampoos when it comes in contact with eyes by mistake upon washing and rinsing, giving only a mild influence on the conjunctiva and the iris.

(2) Hair rinses, hair conditioners, hair treatments:

The ingredients (A) and (B) are dissolved or dispersed in a suitable solvent or medium such as water, ethanol, glycerin, ethylene glycol, propylene glycol, 1,3-propanediol, isopropanol, polyethylene glycol or the like, thereby obtaining the rinse, conditioner or treatment composition.

To these compositions, may be further added any known ingredients which are ordinarily used in rinses, conditioners or treatments. Preferable ingredients which are added to the compositions are surface active agents such as cationic active agents, anionic active agents, nonionic active agents and amphoteric active agents. Most preferably, cationic active agents should be added.

Specific examples of the surface active agents are mentioned below.

(a) Anionic active agents

(1) Linear or branched alkylbenzenesulfonates having an alkyl group having from 10 to 16 carbon atoms on average.

(2) Alkyl or alkenyl ethoxysulfates having a linear or branched alkyl or alkenyl group having from 8 to 20 carbon atoms on average and having 0.5 to 8 moles, on average, of ethylene oxide added in one molecule thereof.

(3) Alkyl- or alkenylsulfates having an alkyl or alkenyl group having from 10 to 20 carbon atoms on average.


(4) Olefinsulfonates having from 10 to 20 carbon atoms on average in one molecule thereof.

(5) Alkanesulfonates having from 10 to 20 carbon atoms on average in one molecule thereof.

(6) Saturated or unsaturated aliphatic acid salts having from 10 to 20 carbon atoms on average in one molecule thereof.

(7) Alkyl or alkenyl ethoxycarboxylates having an alkyl or alkenyl group having from 10 to 20, preferably 12 to 16, carbon atoms on average and having 0.5 to 8 moles, on average, of ethylene oxide added in one molecule thereof.

(8) Alpha-sulfofatty acid salts or esters represented by the following formula



in which Y_2 represents an alkyl group having from 1 to 3 carbon atoms or a counter ion, M_1 represents a counter ion, R_2 represents an alkyl or alkenyl group having from 10 to 20, preferably 12 to 16, carbon atoms.


The counter ions of the anionic active agents are, for example, alkali metal ions such as sodium, potassium and the like, alkaline earth metal ions such as calcium, magnesium and the like, ammonium ions, and alkanolamines having from 1 to 3 alkanol groups having 2 or 3 carbon atoms and including, for example, monoethanolamine, diethanolamine, triethanolamine, triisopropanolamine and the like.

(b) Nonionic active agents

(1) Polyoxyethylene alkyl or alkenyl ethers having a primary or secondary alkyl or alkenyl group having from 8 to 20 carbon atoms on average and having from 3 to 12 moles of ethylene oxide added thereto.

(2) Polyoxyethylene alkylphenyl ethers having an alkyl group having from 8 to 12 carbon atoms on average and having from 3 to 12 moles of ethylene oxide added thereto.


(3) Higher fatty acid alkanolamides represented by the following formula and alkylene oxide adducts thereof



in which R_3 represents H or CH_3 , R_4 represents an alkyl or alkenyl group having from 10 to 20 carbon atoms, n is an integer of from 1 to 3, and m is an integer of from 0 to 3.

(c) Amphoteric active agents


(1) Alkylamine oxides represented by the following formula



in which R_5 represents an alkyl or alkenyl group having from 10 to 20 carbon atoms, R_6 and R_7 independently represent an alkyl group having from 1 to 3 carbon atoms and may be the same or different.


In the above formula, it is preferred that R_5 represents an alkyl or alkenyl group having from 12 to 16 carbon atoms and R_6 and R_7 independently represent methyl group.

(2) Compounds of the following formula




in which R_8 represents an alkyl or alkenyl group having from 10 to 20 carbon atoms, R_9 and R_{10} independently represent an alkyl group having from 1 to 4 carbon atoms, p is an integer of from 1 to 3, and X_2 represents a $COO-$ or SO_3- group. Preferably, R_8 represents an alkyl or alkenyl group having from 12 to 16 carbon atoms, R_9 and R_{10} independently represent methyl group, and p is an integer of 3.

(3) Imidazoline compounds represented by the following formula



in which R_{11} represents a fatty acid residue having from 10 to 20 carbon atoms on average, R_{12} represents hydrogen, Na or CH_2COOMe in which Me represents H, Na or an organic base, R_{13} represents $COOMe$, CH_2COOMe or



in which Me has the same meaning as defined above, and R_{14} represents a hydroxyl group, an acidic salt, or an anionic active sulfate or sulfonate. Preferably, R_{11} represents a fatty acid residue having from 12 to 16 carbon atoms.

(d) Cationic active agents

In the practice of the invention, any cationic active agents may be used without limitation so far as they

are ordinarily used in hair rinses or the like. Preferable examples of the agents include quaternary ammonium salts represented by the following formula (1)



in which one or two of R_{15} , R_{16} , R_{17} and R_{18} independently represent a long-chained alkyl group or long-chained hydroxyalkyl group having from 8 to 20 carbon atoms, and the others independently represent an alkyl or hydroxyalkyl group having from 1 to 3 carbon atoms or a benzyl group, and X_3 represents a halogen atom or an alkyl sulfate group having from 1 or 2 carbon atoms. Specific examples of the quaternary ammonium salts are distearyldimethylammonium chloride, stearyltrimethylammonium methosulfate, stearyltrimethylammonium chloride, stearyldimethylbenzylammonium chloride, lauryldiethylbenzylammonium chloride, lauryltrimethylammonium bromide, distearylmethylhydroxymethyl chloride, cetyltrimethylammonium chloride and the like.

Good results are obtained using these surface active agents in an amount of 0.01 to 10%, preferably 0.5 to 5%, of the hair rinse composition.

The hair rinse or the like compositions of the invention may further comprise hydrocarbons such as liquid paraffin, Vaseline® or solid paraffin, esters such as isopropyl myristate, lanolin derivatives such as lanolin, refined lanolin or lanolin fatty acids silicone derivatives such as dimethylpolysiloxane, methylphenylpolysiloxane or organo-modified polysiloxane, polyethylene glycol, polypropylene glycol or polymers thereof, oils such as polyoxyalkene alkyl ether or polyoxyalkylene alkyl ether phosphates, polymeric materials such as hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropylmethyl cellulose, methyl cellulose, cationized cellulose or cationized polymers, bactericides, preservatives, perfumes or colorants.

(3) Hair Setting Agents (setting lotions, hair sprays and the like)

The ingredients (A) and (B) are dissolved, as usual, in polar solvents such as water, ethyl alcohol, propyl alcohol and the like, thereby obtaining hair setting agents. Any polymer compounds which are used in ordinary hair setting agents may also be used in the practice of the invention. Examples of such polymer compounds are mentioned below.

(a) Polyvinylpyrrolidone compounds.

Specific examples of these compounds include polyvinylpyrrolidone, copolymers of vinylpyrrolidone and vinyl acetate and alkylaminoacrylates. These compounds are commercially available under the names of Luviskol® K, Liviskol® VA, Luviflex® D410I (Yuka Badische Co., Ltd.), and PVPK, PVP/VA and E-735 (GAF Co., Ltd.).

(b) Acidic vinyl ether polymer compounds

For instance, there are mentioned lower alkyl half-esters of copolymers of methyl vinyl ether and maleic anhydride, which are commercially available under the names of Gantrez® ES-225 and ES-335 (Gaf Co., Ltd.).

(c) Acidic polyvinyl acetate polymer compounds

Examples of these compounds include copolymers of vinyl acetate and crotonic acid and are commercially sold under the names of Resin 28-1310 (National Starch Co., Ltd.), Luviset® CE5055 (Yuka Badische Co., Ltd.).

(d) Acidic acrylic polymer compounds

For instance, copolymers of acrylic acid and/or methacrylic acid and alkyl acrylates and/or alkyl methacrylates, and copolymers of acrylic acid, alkyl acrylates and N-alkylamides are mentioned. Commercial products include Plasize® (Gooh Chem. Co., Ltd.), Ultrahold® (Ciba-Geigy A.G.), and the like.

(e) Amphoteric acrylic polymer compounds

There are mentioned compounds which are obtained by copolymerizing dialkylaminoethyl methacrylates, dialkylaminoethyl arylates or diacetone acrylamides with acrylic acid, methacrylic acid, alkyl acrylates and alkyl methacrylates and subjecting the copolymers to amphoteric reaction with halogenated acetic acid. Yuka Foamer AM-TS (Mitsubishi Petrochemical Co., Ltd.) is typical of commercial products of these polymer compounds.

The hair setting agents of the invention may further comprise, within ranges of amounts not impeding the effect of the invention, arbitrary ingredients which may depend on the purpose. Examples of such arbitrary ingredients include oil substances such as higher alcohols or higher fatty acid esters, non-ionic active agents serving as an emulsifier or solubilizer such as polyoxyethylene lauryl ether, polyoxyethylene sorbitan monolaurate and polyoxyethylene hardened castor oil, humectants such as glycerine or propylene glycol, perfumes and colorants.

The hair setting agents of the invention may be applied to the hair as they are. Alternatively, they may be applied in the form of mist by the use of a pump sprayer or the like or may be applied in the form of mist

or foam after packed in a container along with a jetting agent such as flon gas, liquid hydrocarbons or carbon dioxide gas.

The thus obtained hair setting agents of the invention can form a uniform, tough film after drying, which shows excellent hair setting ability even under high humidity conditions. These hair setting agents can be readily removed from the hair upon washing it with now widely used shampoos comprising anionic active agents. Thus, the hair setting agents of the invention satisfy both requirements of set retentivity and washability.

(4) First permanent waving agents

The agents can be prepared by mixing the ingredients (A) and (B) with an ordinary permanent waving solution comprising a reducing material as a substrate.

The reducing materials which are used as a substrate of the first permanent waving agent may be any known materials which are ordinarily used for the purposes. Preferable examples include ammonium salts of thioglycolic acid and acid and hydrochlorides of cysteine.

The first permanent waving agent may be prepared by mixing the above ingredients together by any known technique. The waving agent may further comprise conventionally used ingredients such as colorants, perfumes, oils, opacifiers, water-soluble silicones, organic salts or urea. These ingredients are used depending on the purpose.

(5) Second permanent waving agents

To ordinary second permanent waving agents comprising oxidative materials as a substrate are added the ingredients (A) and (B).

The amount of an oxidative material which is used as substrate of the second permanent waving agent of the invention may differ depending on whether or not the second agent is dissolved in solvent, or the degree of dilution when such an agent is used as diluted. The amount is generally in the range of from 1 to 30%, preferably from 3 to 20%.

The oxidative materials which are used as a substrate of the second permanent waving agent may be any materials ordinarily used for the purposes and include, for example, alkali metal bromates such as sodium bromate, potassium bromate and the like, hydrogen peroxide, sodium percarbonate, sodium perborate and the like. Of these, alkali metal bromate are preferred.

The second permanent waving agents may further comprise, aside from the above essential ingredients, arbitrary ingredients used for the purposes in amounts not impeding the effect of the invention. Examples of such arbitrary ingredients include anionic active agents, amphoteric active agents, nonionic active agents, cationic polymer compounds, water-soluble silicones, urea, suitable oils, humectants, perfumes or colorants.

Cationic polymer compounds mentioned above are, for example, cationic cellulose derivatives, cationic starch, diallyl quaternary ammonium salts, copolymers of diallyl quaternary ammonium salts and acrylamide, polyglycol and polyamide condensates, methacryloxyethyltrimethylammonium salts, and copolymers of methacryloxyethyltrimethylammonium salts and polyvinyl pyrrolidone. Of these, there are preferably used cationic cellulose a typical product of which is available under the name of "Polymer JR", diallyl quaternary ammonium salts a typical product of which is available under the name of "Merquat 100", and diallyl quaternary ammonium salt/acrylamine copolymer typical of which is a product available under the name of "Merquat 500". These cationic polymer compounds are generally used in an amount of from 0.01 to 5%, preferably 0.05 to 2%.

The thus obtained second agent is adjusted in such a way that an aqueous 5% solution thereof has a pH of below 9, preferably from 3.5 to 6.5.

(6) Hair dyes

The ingredients (A) and (B) are added to dye substrates as usual to give hair dyes.

The dye substrates used in the hair dye composition of the invention are not critical and include any known substrates. For example, oxidation hair dyes and temporary hair dyes are described by way of illustration only.

(i) Oxidation hair dyes

Dye intermediates, oxidizing agents and, if necessary, couplers or modifiers are mixed together.

Examples of the dye intermediates include para or ortho compounds such as p-phenylenediamine, toluene-2,5-diamine, N-phenyl-p-phenylenediamine, 4,4'-diaminodiphenylamine, p-aminophenol, p-methylaminophenol, o-phenylenediamine, toluene-3,4-diamine, o-aminophenol, p-chloro-o-phenylenediamine, p-amino-o-cresol, o-chloro-p-phenylenediamine, phloroglucin pyrogallol, 3,3'-iminodiphenyl, diphenylamine, 2,6-diaminopyridine or p-aminophenylsulfamic acid. Couplers or modifiers include metha compounds or phenols such as m-phenylenediamine, toluene-2,4-diamine, p-methoxy-m-phenylenediamine, m-aminophenol, alpha-naphthol, resorcin, hydroquinone and catechol. Hydrogen peroxide is ordinarily used as the oxidizing agent. Also, there may be used sodium perborate, urea peroxide, sodium percarbonate, sodium peroxytripolyphosphate, sodium peroxyphosphate, sodium peroxyorthophosphate, sodium silicate and hydrogen peroxide adduct and sodium sulfate-sodium chloride-hydrogen peroxide adduct. In order to give an influence on the color of the hair without taking part in the dye

formation reaction, there may be further added direct dyes and particularly nitro dyes such as nitro-p-phenylenediamine, p-nitro-o-phenylenediamine, 2-amino-4-nitrophenol, 2-amino-5-nitrophenol and, 4-amino-2-nitrophenol. If necessary, picramic acid, picric acid and/or 1,4-diaminoanthraquinone may be also added.

Within ranges of amounts not impeding the effect of the invention, there may be also added nonionic active agents, cationic active agents, solvents such as propylene glycol, glycerine and the like, viscosity adjusters including lower alcohols such as ethyl alcohol, isopropyl alcohol and the like, hydroxyethyl cellulose, methyl cellulose, cationic polymer compounds, and higher alcohols, humectants, protein modifiers such as urea, perfumes, colorants, UV absorbers, antioxidants, preservatives, pearling agents or lotionizing agents.

The oxidation hair agents of the invention are prepared by mixing the essential ingredients and the above-mentioned ingredients by a usual manner to give a powder preparation of a cream preparation. In applications, the preparation is added to water or a shampoo substrate. Alternatively, an oxidation dye and an oxidizing agent are separately provided. The essential ingredients are added to either or both of the agents to give powder, cream or liquid preparations. In applications, separately provided systems are combined together.

(ii) Temporary hair dyes

Dyes and pigments are not limited to any specific ones and include, for example, pigments such as titanium oxide, carbon black and the like, and tar-base colorants such as triphenylmethane dyes, azo dyes, quinoline dyes, xanthene dyes, acridine dyes, azine dyes, oxazine dyes, indigoid dyes, stilbene dyes, thiazole dyes and the like.

The resins are, for example, copolymers of acrylic esters or methacrylic esters, copolymers of monochloroacetic acid-amine salt-modified products of N,N'-diethylaminoethyl methacrylate and methacrylic esters, copolymers of vinyl pyrrolidone and vinyl acetate and the like.

The temporary hair dyes are prepared by dissolving or dispersing the essential ingredients, resins, dyes and pigments in a dispersion medium such as water, amyl alcohol, isopropanol, ethanol or acetone. As a matter of course, there may be further added known ingredients, which are ordinarily added to existing temporary hair dye compositions, including anionic active agents, cationic active agents, amphoteric active agents, nonionic active agents, polyhydric alcohols such as propylene glycol, glycerine, polyethylene glycol and the like, higher alcohols such as isostearyl alcohol, oleyl alcohol and the like, fatty acids such as lanolin fatty acids, fatty acids from coconut oil and the like, esters such as isopropyl myristate, hydrocarbons such as liquid paraffin, cationized polymer compounds, amines and perfumes.

(7) Pre-shampoo treatment agents

Pre-shampoo treatment agent is prepared by dissolving or suspending the ingredients (A) and (B) and, optionally, other known ingredients in a medium such as water.

Known ingredients are, for example, oils and fats such as higher alcohols and fatty acid esters, nonionic active agents serving as an emulsifier or solubilizing agent and including polyoxyalkylene alkyl ethers, and humectants such as glycerine and pyrrolidonecarboxylic acid. These ingredients serve to control a degree of styling of the hair when the hair is treated with the pre-shampoo treatment agent of the invention and shampooed thereafter. That is, use of liquid oil or fat ingredients is effective in imparting softness to the hair and humectants can impart moistness to the hair. In addition, use of higher alcohols imparts dryness to the hair.

The present invention is described by way of references and examples, which should not be construed as limiting the present invention thereto.

Reference 1

Ten kilograms of the whole of a birch tree (*Betula alba*) was admixed with 20 liters of 40% 1,3-butylene glycol immersed therein at 20 to 25°C for 10 to 15 days, followed by centrifugal separation. To the residue was further added 10 liters of 1,3-butylene glycol for re-immersion. The resulting immersion solution and the solution obtained by the centrifugal separation were combined and aged in a dark and cold place (0 to 5°C) for about 40 days. As a result, there was obtained a birch extract in the form of a dark brown liquid containing 5% solid matters.

Reference 2

Twenty liters of 50% ethanol was added to 10 kg of the whole of a birch tree (*Betula alba*) for immersion in a cold and dark place for 20 days. As a result, a birch extract in the form of a brown liquid containing 6.5% solid matters was obtained.

Reference 3

Ten liters of an aqueous 50% propylene glycol solution was added to 5 kg of leaves of rosemary (*Rosmarinus Officinalis*), followed by refluxing at 50°C for 12 hours to obtain an extract. The extract was subjected to filtration to remove the residue therefrom to obtain a final product. This product was a rosemary extract in the form of a brown liquid containing 10% solid matters.

Reference 4

Ten kilograms of silk fibers obtained from the silkworm was washed with water, dried, cut into pieces, powdered and passed through a 200 μ m sieve to obtain 9.5 kg of a silk powder.

Reference 5

Sixty grams of silk fibers were dissolved in 600 ml of water containing 48 g of ethylenediamine and 36 g of cupric hydroxide, followed by heating to 60°C for 2 hours. Subsequently, the solution was adjusted in pH with 3N acetic acid to 6.0—7.0, and filtered, followed by placing the resulting filtrate in a cellulose tube C—65 and subjecting to dialysis in running water over one or two days and nights. The resulting dialyzate was filtered and made up to 2000 ml, to which was added 80 g of a strongly acidic cationic exchange resin Dowex® all 50wx—2, followed by agitating for 20 to 30 minutes and filtering. Thereafter, the resulting filtrate was admixed with an 1N sodium hydroxide solution to adjust its pH to 6.0—7.0. As a result, there was obtained 1900 ml of a solution containing 0.5% of the effective ingredient. The thus obtained silk protein had an average molecular weight of 2000.

Reference 6

Preparation of Decomposition Derivatives by oxidation of Keratin Material:

Ten grams of wool fibers were immersed in 700 g of an aqueous 8% peracetic acid solution at room temperature for 1 day to carry out the oxidation reaction. The resulting oxidized wool fibers were filtered, washed with water and immersed in 700 g of a 0.1N ammoniacal solution at room temperature for 1 day, permitting about 90% of the wool to dissolve in the ammoniacal solution. About 1 g of the insoluble matters were removed by filtration and the aqueous ammoniacal solution of keratose to be an oxidized decomposition product of the wool keratin was admixed with 2N hydrochloric acid to adjust the pH to 4.0, whereupon alpha-keratose was settled as precipitate. This precipitate was filtered, washed with acetone and dried to obtain 5.4 g of alpha-keratose.

Reference 7

Keratin Materials:

Ten grams of wool fibers were immersed in 60 ml of an aqueous solution with concentration of 8M urea and 0.01M tris buffer, to which was added 6 ml of 2-mercaptoethanol, followed by adjusting the pH to 10 by means of a 5N potassium hydroxide aqueous solution to effect the reduction reaction in a stream of nitrogen at room temperature. About 3 hours after commencement of the reaction, the wool dissolved in the reaction solution in an amount of about 85% thereof. While the system was adjusted with a 5N potassium hydroxide solution so that the pH was not below 7, 16.5 g of iodoacetic acid was gradually added and the pH of the system was finally adjusted to 8.5 to carry out the carboxymethylation reaction at room temperature for 2 hours. The reaction solution was filtered to remove insoluble matters therefrom and the resultant filtrate was charged into a cellulose tube wherein it was dialyzed against ion-exchanged water to remove low molecular weight impurities including urea. As the urea was dialyzed, the content in the cellulose tube became cloudy since water-insoluble HGT (component with high contents of glycerine and tyrosine) was caused to precipitate. After completion of the dialysis, HGT was centrifugally removed and S-carboxymethyl keratin (SCMKA) was obtained from the neutral transparent solution of SCMKA by the isoelectric precipitation method. That is, 1N hydrochloric acid was added to the system to adjust its pH to 4.4 by which SCMKA became insoluble and separated as precipitate. This precipitate was filtered, washed with ethanol and dried to obtain 4.2 g of SCMKA.

Reference 8

Preparation of Hydrolysis Derivatives of Keratin Materials:

Ten grams of wool fibers were immersed in 300 g of a 1% sodium hydrogensulfite aqueous solution, whose pH was adjusted to 6.7 by the use of a 5N aqueous caustic soda solution. Thereafter, 0.2 g of papain was added to the system to effect the hydrolysis reaction at 60°C for 15 hours, by which about 80% of the wool dissolved. Insoluble matters were removed by filtration and the sulfite in the resulting filtrate was removed by the ultrafiltration technique using a membrane with a fractional molecular weight of 500. The aqueous solution of the hydrolysate was concentrated and freeze dried to obtain 7.7 g of the hydrolysate having a molecular weight of 500—2,000.

Example 1

Shampoo:

Shampoo compositions of the formulations indicated in Table 1—A were prepared and subjected to the performance evaluation test thereof. The results are shown in Table 1—B.

The performance evaluation was conducted according to the following methods in or throughout the examples appearing hereinafter.

(1) Lathering or foaming test

To an aqueous solution of 1% shampoo composition was added 0.1% of lanolin as an artificial dirt, followed by agitating in a cylinder for 5 minutes by means of a flat propeller at 40°C at 1000 r.p.m. in such a way that the propeller was turned in opposite directions every 10 seconds. After completion of the agitation, an amount of lathering after 30 seconds was measured for evaluation.

(2) Touch of lather

Thirty grams of human hair was wetted with water of 40°C to permit 20 g of water to be contained. Thereafter, 1 g of each of the shampoo compositions was used for washing the hair therewith whereupon the touch of the lather was organoleptically evaluated by 20 female panelers.

5

Evaluation item

The degree of ease in passing fingers through the hair upon washing was evaluated as "foam smoothness".

10 Evaluation standard

- ☐ Better foam smoothness than that of a reference
- ☐ Slightly better than a reference
- ☒ Same in foam smoothness as a reference

15 (3) Combing force

Thirty grams of human hair was wetted with water of 40°C to permit 20 g of water to be contained. One gram of each of the shampoo compositions was used to effect washing and rinsing operations two times, after which it was squeezed and set on a strain gauge. Subsequently, the hair was combed to determine a force exerted thereon (wet combing force). The hair was then dried with a dryer and allowed to stand in an air-conditioned room of a temperature of 20°C and a relative humidity of 65% overnight, followed by setting on a strain gauge and combing to determine a force exerted thereon (dry combing force).

20

(4) Hair fly

Upon the measurement of "combing force" under dry conditions, it was observed whether or not there took place an electrostatic hair fly phenomenon.

25

Evaluation

- ☐ Occurrence of hair fly
- ☒ No occurrence of hair fly

30

35

40

50

55

60

65

TABLE 1—A

	A	B	C	D	E	F	G
Sodium polyoxyethylene (2) lauryl sulfate	15%	15%	15%	15%	15%	15%	15%
Coconut fatty acid diethanolamide	3	3	3	3	3	3	3
Rosemary extract (Novarom Co., Ltd.)	—	0.5	0.5	—	0.5	—	0.5
Oxidative decomposition product of keratin (obtained in Reference 6)	—	—	0.1	—	—	—	—
Oxidative decomposition product of collagen*	—	—	—	0.1	0.1	—	—
Oxidative decomposition product of collagen**	—	—	—	—	—	0.1	0.1
Perfume	suitable amount	suitable amount	suitable amount	suitable amount	suitable amount	suitable amount	suitable amount
Water	balance (pH 7.2)	balance (pH 7.2)	balance (pH 7.2)	balance (pH 7.2)	balance (pH 7.2)	balance (pH 7.2)	balance (pH 7.2)

* **, having average molecular weight of about 1,000.

TABLE 1—B

Lathering Characteristics			Hair Finishing	
	Lathering	Lather Smoothness	Combing Force (g)	Hair Fly
A	114	X	210	O
B	105	△	196	O
C (product of invention)	138	⊙	101	X
D	91	X	254	O
E	93	△	247	O
F	87	X	201	O
G	90	X	198	O

Example 2

Shampoo:

Shampoo compositions of the formulation indicated in Table 2—A were prepared and subjected to the performance evaluation test. The results are shown in Table 2—B.

The performance evaluations were effected in the same manner as in Example 1.

TABLE 2—A

	H	I	J
Sodium polyoxyethylene (2) lauryl sulfate	15%	15%	15%
Coconut fatty acid diethanolamide	3	3	3
Birch extract (Novarom Co., Ltd.)	—	0.5	0.5
Hydrolyzate of silk (average molecular weight of 20,000)	—	—	0.1
Perfume	suitable amount	suitable amount	suitable amount
Water	balance (pH 7.2)	balance (pH 7.2)	balance (pH 7.2)

TABLE 2—B

Lathering Characteristics			Hair Finishing	
	Lathering	Lather Smoothness	Combing Force (g)	Hair Fly
H	114	X	210	O
I	107	△	176	O
J (product of invention)	142	⊙	97	X

Example 3

Hair Rinse:

Hair rinses of the formulations indicated in Table 3—A were prepared. Five hundred milliliters of a 50:1 dilution of each hair rinse was used to treat the hair therewith, which was subsequently rinsed with hot water two times and air-dried. The thus dried hair was evaluated by 20 panelers according to the following five-point method.

Evaluation standard:

Good 5, Fair 4, Moderate 3, Rather poor 2, Poor 1

The results are shown in Table 3—B as average values.

TABLE 3—A

	K	L	M	N
Distearyldimethylammonium chloride	2%	2%	2%	2%
Stearyl alcohol	1%	1%	1%	1%
Birch extract (obtained in Reference 1)	—	1%	—	1%
Hydrolyzate of keratin (obtained in Reference 8, average molecular weight 3000)	—	—	1%	1%
Water	balance (pH 5.0)	balance (pH 5.0)	balance (pH 5.0)	balance (pH 5.0)

TABLE 3—B

	Softness	Smoothness	Ease in Combing
K	1.2	1.3	1.2
L	2.1	2.5	3.3
M	3.2	3.4	2.5
N (product of invention)	4.2	4.0	3.9

Example 4

Hair Rinse:

Hair rinses of the formulations indicated in Table 4—A were prepared and evaluated in the same manner as in Example 3. The results are shown in Table 4—B.

TABLE 4—A

	O	P	Q	R
Distearyldimethylammonium chloride	2%	2%	2%	2%
Stearyl alcohol	1%	1%	1%	1%
Rosemary extract (obtained in Reference 3)	—	1%	—	1%
Hydrolyzate of silk average molecular weight 20,000)	—	—	1%	1%
Water	balance (pH 5.0)	balance (pH 5.0)	balance (pH 5.0)	balance (pH 5.0)

TABLE 4—B

	Softness	Smoothness	Ease in Combing
O	1.2	1.3	1.2
P	2.1	2.6	3.4
Q	3.2	3.4	2.5
R (product of invention)	4.5	4.1	4.4

Example 5

Hair Treatment Agent:

Hair treatment agents of formulations (S) through (V) were prepared using the following fundamental composition.

Fundamental Composition

Liquid paraffin	5.0%
White Vaseline®	2.0%
Cetyl alcohol	2.0%
POE (20) sorbitan monostearate	1.0%
Glycerine	10.0%
Water	balance pH 7.0

To the above composition were added:

- (S) no material,
- (T) 1.0% of a birch extract (obtained in Reference 1),
- (U) 2.0% of a hydrolyzate of keratin (an average molecular weight of 20,000), or
- (V) 1.0% of a birch extract (obtained in Reference 1) and 2.0% of a hydrolyzate of keratin (average molecular weight of 20,000).

These hair treatments were evaluated according to the five-point method with respect to the following items. Average values of the respective evaluation tests are shown in Table 5.

TABLE 5

	Organoleptic Evaluation	Combing Force (g)	
		Wet State	Dry State
S	2.0	420	360
T	2.0	405	351
U	2.3	326	265
V (product of invention)	4.6	116	107

Example 6

Hair Treatment Agent:

Based on the fundamental composition of Example 5, the following formulations (W)—(Z) were prepared.

(W) Nothing was added.

(X) 1.0% of a birch extract (obtained in Reference 1) was added.

(Y) 2.0% of a silk hydrolyzate (average molecular weight 20,000) was added.

(Z) 1.0% of a birch extract (obtained in Reference 1) and 2.0% of a silk hydrolyzate were added.

The resulting hair treatments were evaluated in the same manner as in Example 5 with the results shown in Table 6.

TABLE 6

	Organoleptic Evaluation	Combing Force (g) Wet State	Dry State
W	2.0	420	360
X	2.0	405	351
Y	2.3	326	265
Z (product of invention)	4.6	123	115

Example 7

Hair Setting Lotion:

Hair setting lotions of the formulations indicated in Table 7—A were prepared and their set retentivity was evaluated. The results are shown in Table 7—B.

TABLE 7—A

	A	B	C
Ethanol	10%	10%	10%
Birch extract (Vermin Co., Ltd.)	—	2.0%	2.0%
Hydrolyzate of wool keratin (average molecular weight 20,000)	—	—	1.0%
Polyoxyethylene oleyl ether (EO 20)	0.5%	0.5%	0.5%
Perfume	0.1%	0.1%	0.1%
Water	balance	balance	balance

$$\text{Set retentivity (\%)} = \frac{L_o - L_t}{L_o - L_s} \times 100$$

in which, $L_o = 14$ cm,

$L_s =$ a length (cm) of a curled hair immediately after suspension under 95% relative humidity conditions,

$L_t =$ a length (cm) of a curled hair 30 minutes after suspension of the hair under 95% R.H. conditions.

TABLE 7—B

Set Retentivity	
<i>A</i>	53%
<i>B</i>	60%
<i>C</i> (product of invention)	76%

Example 8

Hair Setting Lotion:

Hair setting lotions of the formulations indicated in Table 8—A were prepared to determine set retentivity in the same manner as in Example 7. The results are shown in Table 8—B.

TABLE 8—A

	<i>D</i>	<i>E</i>	<i>F</i>
Ethanol	10%	10%	10%
Birch extract (Vermin Co., Ltd.)	—	2.0%	2.0%
Silk hydrolyzate (average molecular weight 20,000)	—	—	1.0%
Polyoxyethylene oleyl ether (EO 20)	0.5%	0.5%	0.5%
Perfume	0.1%	0.1%	0.1%
Water	balance	balance	balance

TABLE 8—B

Set Retentivity	
<i>D</i>	53%
<i>E</i>	60%
<i>F</i> (product of invention)	84%

Example 9

First Permanent Waving Agent:

Permanent waving treatments were effected using first permanent waving agents of the formulation indicated in Table 9—A and a second permanent waving agent to determine a degree of waving, wave retentivity, adsorptivity, and feeling to the touch. The results are shown in Table 9—B.

Formulation:

(1) First permanent waving agent

TABLE 9—A

	G	H	I
Ammonium thioglycolate	7%	7%	7%
Birch extract (Novarom Co., Ltd.)	—	1%	1%
Silk hydrolyzate (average molecular weight 1,500)	—	—	2%
Water (ammoniacal solution for pH adjustment)	93%	92%	90%

(2) Second permanent waving agent:

Sodium bromate	5.0%
Water	95.0%

Test method:

(1) Measurement of a degree of waving and wave retentivity

(i) A bundle of 20 human hairs were fixed to cylinders of a wave-measuring plate (in which a plurality of cylinders, each having a diameter of 2 mm and a length of 1.5 mm, were arranged in zigzag form in two rows). The bundle was immersed in the first agent of each of the formulations through at 30°C for 10 minutes, followed by immersion in a second agent at 30°C for 10 minutes. Thereafter, the immersed bundle was sufficiently rinsed with water and removed from the plate. A degree of waving in still water was calculated according to the following equation.

It should be noted that the hair used was a 20 cm long virgin hair which was washed with an aqueous solution of 0.5% of sodium lauryl sulfate and dried.

$$\text{Degree of waving (\%)} = \frac{X_o - Z_o}{X_o - Y_o} \times 100$$

X_o : a length of the hair fixed between points a and b of one row of the zigzag cylinders which are distant from each other.

Y_o : a distance between a and b .

Z_o : a distance of the hair, fixed between the point a and b , in still water after removal from the measuring plate.

(ii) The hair used in (i) was slightly moved and washed while immersing in an aqueous solution of 0.5% sodium lauryl sulfate for 1 minute, followed by sufficiently rinsing and air drying for 1 day. The above procedure was repeated four times and, after the fifth washing, the hair was sufficiently rinsed. Thereafter, a distance, Z_o , in still water was measured, from which a degree of wave was determined. A ratio of this degree to a degree of wave prior to washing was calculated to give a wave retentivity.

$$\text{Wave retentivity (\%)} = \frac{\text{degree of wave after fifth washing}}{\text{degree of wave prior to washing}} \times 100$$

(2) Adsorptivity

The hair used for the measurement of the degree of wave was observed through a scanning electron microscope in order to determine the presence or absence of adsorbed matters on the hair surfaces. The evaluation standard is shown below.

Adsorptivity	State of Hair Surfaces
++	Covered with a film and being thus smooth
+	Slight amounts of adsorbed matters observed though the surfaces are smooth
-	Considerable irregularities observed with hair cuticle being partly separated

(3) Evaluation of Feeling to the Touch

A hair bundle made of Japanese virgin hairs was immersed in a first agent of each of formulations through at 30°C for 10 minutes and subsequently immersed in a second agent at 30°C for 10 minutes. The bundle which was sufficiently rinsed with water and air-dried was evaluated by 20 female panelers according to the five-point method with respect to the feeling of the hair bundle to the touch. The evaluation standard was as follows: Good = 5; Fair = 4; Moderate = 3; Rather poor = 2; Poor = 1. The results are indicated by geometric average.

TABLE 9—B

First Permanent	Adsorptivity	Degree of wave (%)	Wave Retentivity (%)	Feeling to the touch
G	-	52	73	2.4
H	+	49	70	2.1
I	++	56	91	3.5

Example 10

Second Permanent Waving Agent:

The hair was permed using a first agent and second agents of the formulations indicated below. A degree of damage of the hair during the perming treatment was determined by measuring weights of the hair prior to and after the treatment. The method of measuring the hair weight and the evaluation standard are described below.

[Composition]

First Agent:

Thioglycolic acid	7.0%
Polyoxyethylene hardened castor oil	1.0%
Perfume	0.2%
Ammoniacal solution, water	balance
(pH adjusted to 9.0 by means of ammoniacal solution)	

TABLE 10

Second Agents

	J	K	L
Sodium bromate	5%	5%	5%
Amphoteric active agent (Miranol® CSM—SF, by Miranol Co., Ltd.)	0.5%	0.5%	0.5%
Perfume	0.1%	0.1%	0.1%
Rosemary extract (Vermin Co., Ltd.)	—	1.0%	1.0%
Silk hydrolyzate (average molecular weight 1,500)	—	—	2.0%
Water	balance	balance	balance

25

[Measurement of Hair Weight]

Ten centimeters long virgin hair was washed with an aqueous solution of 0.5% sodium lauryl sulfate and air-dried to give a hair being tested. About 1 g of the hair was bundled and placed in a desiccator using a phosphorus pentoxide dryer, followed by drying under reduced pressure for one week and measuring the dried hair to give an absolute dry weight of the virgin hair. The thus dried hair was immersed in the first agent or solution at 30°C for 10 minutes and sufficiently washed with water. Thereafter, the hair was further immersed in the respective second agents at 30°C for 10 minutes. After sufficiently rinsing with water, it was air-dried and dried according to the above procedure to give an absolute dry weight of the permed hair.

35

[Evaluation Standard]

	Evaluation	Content
40	⊙	The difference in absolute dry weight between the virgin hair and the permed hair is less than 1%
45	O	The difference in absolute dry weight between the virgin hair and the permed hair is in the range from 1 to 5%
50	X	The difference in absolute dry weight between the virgin hair and the permed hair is over 5%
	<u>Results</u>	
	J	X
55	K	O
	L (product of invention)	⊙

60

Example 11

Hair Dye:

Two-component hair dye compositions of the formulations indicated in Table 11 were prepared and the influence of these compositions on the hair was determined by measuring hair weights prior to and after the hair dyeing treatment. The results are shown below.

Composition:

TABLE 11

First Liquid Component

		<i>M</i>	<i>N</i>	<i>O</i>
10	p-Phenylenediamine	1.0%	1.0%	1.0%
	Birch extract (obtained in Reference 2)	—	1.0%	1.0%
15	Silk hydrolyzate (average molecular weight 20,000)	—	—	0.5%
	Propylene glycol	10.0%	10.0%	10.0%
20	Disodium edetate	0.3%	0.3%	0.3%
	Sodium sulfite	0.5%	0.5%	0.5%
	Perfume	0.1%	0.1%	0.1%
25	Water	balance	balance	balance

(pH was adjusted to 10.0 by means of an ammoniacal solution.)

(Second liquid component)

	Hydrogen peroxide	6.0%
35	Water	94.0%

Evaluation Method:

Ten centimeters long virgin hair was washed with an aqueous solution of 0.5% sodium lauryl sulfate and air-dried to give a hair being tested. About 1 g of the hair was bundled and placed in a desiccator using phosphorus pentoxide dryer, followed by drying under reduced pressure for 1 week and measuring the dried hair weight to give an absolute dry weight of the virgin hair.

Subsequently, the hair was dyed according to the following procedure and air-dried, followed by drying in the same manner as described above and measuring the weight of the dyed hair to give an absolute dry weight of the dyed hair.

The hair weight after the hair dyeing treatment was evaluated by comparison with the virgin hair weight and an influence of the respective dye compositions on the hair was judged.

[Evaluation Standard]

	<u>Evaluation</u>	<u>Content</u>
50	⊙	The dyed hair increased in weight over the virgin hair
55	O	The dyed hair decreased in weight by 0—3% as compared with the virgin hair
60	X	The dyed hair decreases in weight over 3% as compared with the virgin hair

Dyeing Procedure:

A mixture of the first and second liquids in equal amounts was provided as a dyeing solution and was diluted to have a liquor ratio of 1:5. The hair was immersed in the solution at room temperature for 30 minutes and dyed. Thereafter, the hair was washed with city water of 40°C to wash away the deposited dye

therefrom and washed further with an aqueous solution of 0.5% sodium lauryl sulfate, followed by instantaneously immersing in an aqueous 1N acetic acid solution and washing with city water of 40°C.

Results:

5	<i>M</i>	X
	<i>N</i>	O
10	<i>O</i> (product of invention)	⊙

Example 12

Hair Dye:

Temporary hair dye compositions (hair colorants) of the mascara type having formulations indicated in Table 12—A were prepared. 0.5 g of each composition was applied to 1 g of grey hair. After drying in air, the hair was organoleptically evaluated by 10 expert panelers with respect to the gloss, smoothness and feeling. The results are shown in Table 12—B.

Composition:

TABLE 12—A

25		<i>P</i>	<i>Q</i>	<i>R</i>
	Polymer resin*	12.0%	12.0%	12.0%
	Pigment (carbon black)	1.0%	1.0%	1.0%
30	Birch extract (obtained in Reference 1)	—	1.0%	1.0%
35	Silk hydrolyzate (average molecular weight 20,000)	—	—	0.5%
	Perfume	0.1%	0.1%	0.1%
40	Ethanol	balance	balance	balance

* Polymer resin: Copolymer of monochloroacetic acid-amine salt-modified product of N,N'-dimethylaminoethyl methacrylate and methacrylic ester.

Results:

TABLE 12—B

50		Gloss	Smoothness	Feeling to the Touch
	<i>P</i>	X	X	X
	<i>Q</i>	Δ	Δ	O
55	<i>R</i> (product of invention)	O	O	O

In the above table, by the mark "O" is meant "good", by "Δ" meant "fair", and by "X" meant "poor".

0 117 469**Example 13****Hair Liquid:****[Composition]**

5	(a)	Rosemary extract (Novarom Co., Ltd.)	1.0%
	(b)	Oxidative decomposition product of keratin (obtained in Reference 6)	0.1%
10	(c)	Polyoxypropylene (30) butyl ether	15.0%
	(d)	Ethanol	40.0%
15	(e)	Water	44.0%

[Preparation]

20 The above ingredients (a) through (e) were mixed together to completely dissolve the components to obtain a hair liquid.

Example 14**Hair Tonic:****[Composition]**

25	(a)	Rosemary extract (obtained in Reference 3)	1.0%
30	(b)	Hydrolyzate of keratin (obtained in Reference 8)	0.1%
	(c)	PCA—A1	0.5%
35	(d)	Ethanol	55.0%
	(e)	Water	44.0%

[Preparation]

40 The above ingredients (a) through (e) were mixed and agitated until a uniform solution was obtained, thereby giving a hair tonic.

Example 15**Hair Liquid:****[Composition]**

50	(a)	Birch extract (obtained in Reference 1)	1.0%
	(b)	Silk hydrolyzate (average molecular weight 20,000)	0.1%
55	(c)	Polyoxypropylene (30) butyl ether	15.0%
	(d)	Ethanol	40.0%
60	(e)	Water	44.0%

[Preparation]

65 The above ingredients (a) through (e) were mixed for complete dissolution to obtain a hair liquid.

Example 16

Hair Tonic:

[Composition]

5	(a)	Birch extract (Novarom Co. Ltd.)	1.0%
	(b)	Silk hydrolyzate (average molecular weight 20,000)	0.1%
10	(c)	PCA—A1	0.5%
	(d)	Ethanol	55.0%
	(e)	Water	44.0%

15 [Preparation]

The above ingredients (a) through (e) were mixed and agitated until a uniform solution was obtained, thereby giving a hair tonic.

Example 17

Setting Lotion:

20 Setting lotions having formulations indicated in Table 17—A were prepared and applied to the hair to determine the feeling of the hair to the touch and set retentivity. The results are shown in Table 17—B.

TABLE 17—A

25		<i>S</i>	<i>T</i>	<i>U</i>
	Birch extract (Dragoco Co., Ltd.)	—	0.005%	0.005%
30	Oxidative decomposition product of keratin (obtained in Reference 6)	—	—	0.2%
	Cationic polymer (Polymer JR400®)	1.0	1.0	1.0
35	Sodium chloride	1.0	1.0	1.0
	Ethanol	5.0	5.0	5.0
	Water	balance	balance	balance
40	pH	7.0	7.0	7.0

[Test Method]

(1) Feeling of Treated Hair to the Touch

45 Twenty grams of a 20 cm long hair bundle made of Japanese female hair was applied with 2 g of a hair cosmetic throughout the hair and then rinsed with running water of 40°C for 1 minute. The thus treated hair was organoleptically evaluated. The organoleptic evaluation was carried out by paired comparison according to the following evaluation point using, as a reference, a hair bundle treated with a commercial hair rinse comprising as its principal components a quaternary ammonium salt and a hydrocarbon. An average point by 20 expert panelers is indicated in Table appearing hereinafter.

	<u>Evaluation Point</u>	<u>Evaluation</u>
55	+2	Better in feeling than the reference hair bundle
	+1	Slightly better than the reference hair bundle
	0	Equal to the reference hair bundle
60	-1	Slightly poorer than the reference hair bundle
	-2	Poorer than the reference hair bundle
65		

(2) Set-retaining Effect

A 20 cm long hair bundle having a weight of 5 g and made of Japanese female hair was applied with 1 g of a hair cosmetic throughout the hair and rinsed with running water of 40°C for 1 minute, after which an excess of water was removed by means of filter paper. The thus treated hair bundle was wound around a glass tube with a diameter of 1.5 cm so that a winding width was 5 cm, and fixed at opposite ends thereof. The bundle was allowed to stand under conditions of 65 R.H.% and 20°C for 24 hours and curled. After 24 hours, the wound bundle was removed from the tube and vertically suspended, whereupon its length was measured and a degree of curling was calculated according to the following equation.

$$\text{Degree of curling (\%)} = \frac{A_o - (B_o - A_o)}{A_o} \times 100$$

A_o: Length of the hair immediately after removal of the hair.

B_o: Length of the hair 12 hours after removal of the hair.

(3) Measurement of combing force under wetting conditions

After completion of the evaluation of the feeling of the hair, it was fixed on a strain gauge, through which a nylon comb was passed 20 times while recording a force exerted upon the combing. An average value of the 20 measurement was determined as a combining force.

(4) Feeling and measurement of combing force under dry conditions

The hair bundle after the measurement of the combing force under wet conditions was dried in air and then evaluated according to the evaluation and measurement methods described in (1) and (3) above.

Results:

TABLE 17—B

	Feeling of the Hair	Degree of Curling	Combing Force	
			under wet conditions	under dry conditions
S	-0.1	72%	170 (g)	194 (g)
T	+0.1	76%	165	187
U	+1.4	90%	107	160

Example 18

Setting Lotion:

Setting lotions of formulations indicated in Table 18—A were prepared and applied to the hair. The feeling of the hair and the set-retaining effect of the respective lotions were determined in the same manner as in Example 17. The results are shown in Table 18—B.

TABLE 18—A

	V	W	X
Birch extract (Dragoco Co., Ltd.)	—	0.005%	0.005%
Silk hydrolyzate (M.W. 20,000)	—	—	0.2%
Cationic polymer (Polymer JR400®)	1.0	1.0	1.0
Sodium chloride	1.0	1.0	1.0
Ethanol	5.0	5.0	5.0
Water	balance	balance	balance
pH	7.0	7.0	7.0

Results:

TABLE 18—B

		Feeling of the Hair	Degree of Curling	Combing Force	
				under wet conditions	under dry conditions
5					
10	V	-0.1	72%	170 (g)	194 (g)
	W	+0.1	76%	165	187
15	X	+1.3	93%	136	161

20 Claims

1. A hair cosmetic composition comprising
(A) an extract obtained by polar solvent extraction of birch or rosemary, and
(B) a polypeptide compound having a molecular weight of 350 to 30,000 selected from decomposition
25 products of silk, decomposition products of keratin and derivatives thereof.
2. A hair cosmetic composition according to Claim 1, wherein the ingredient (A) is contained in an amount ranging from 0.001 to 10 wt% and the ingredient (B) is in an amount ranging from 0.01 to 10 wt%.
3. A hair cosmetic composition according to Claim 1, wherein the mixing ratio of the ingredient (A) and the ingredient (B) is in the range of 0.1 to 10:1.
- 30 4. A hair cosmetic composition according to Claim 1, wherein the ingredient (B) is a polypeptide compound having a molecular weight of 350 to 30,000 selected from hydrolyzates of silk, hydrolyzates of keratin, oxidative decomposition products of keratin and thiol group-modified compounds of reductive decomposition products of keratin.

35 Patentansprüche

1. Haarkosmetikum-Zusammensetzung, umfassend
(A) einen Extrakt, erhalten durch Extraktion von Birke oder Rosmarin mit einem polaren Lösungsmittel,
40 und
(B) eine Polypeptidverbindung mit einem Molekulargewicht von 350 bis 30000, ausgewählt aus Zersetzungsprodukten von Seide, Zersetzungsprodukten von Keratin und Derivaten derselben.
2. Haarkosmetikum-Zusammensetzung gemäß Anspruch 1, wobei der Bestandteil (A) in einer Menge im Bereich von 0,001 bis 10 Gew.-% enthalten ist und der Bestandteil (B) in einer Menge im Bereich von
45 0,01 bis 10 Gew.-% enthalten ist.
3. Haarkosmetikum-Zusammensetzung gemäß Anspruch 1, wobei das Mischungsverhältnis von dem Bestandteil (A) und dem Bestandteil (B) im Bereich von 0,1 bis 10:1 liegt.
4. Haarkosmetikum-Zusammensetzung gemäß Anspruch 1, wobei der Bestandteil (B) eine Polypeptidverbindung mit einem Molekulargewicht von 350 bis 30000 ist, ausgewählt aus Hydrolysaten
50 von Seide, Hydrolysaten von Keratin, oxidativen Zersetzungsprodukten von Keratin und an der Thiolgruppe modifizierten Verbindungen von reduktiven Zersetzungsprodukten von Keratin.

55 Revendications

1. Composition cosmétique pour cheveux comprenant:
(A) un extrait obtenu moyennant une extraction de bouleau ou de romarin au moyen d'un solvant
polaire, et
(B) un composé polypeptidique ayant un poids moléculaire de 350 à 30,000, choisi parmi les produits
60 de décomposition de la soie, les produits de décomposition de la kératine et leurs dérivés.
2. Composition cosmétique pour cheveux selon la revendication 1, caractérisée en ce que l'ingrédient (A) est présent en une quantité se situant entre 0,001 et 10% en poids, tandis que l'ingrédient (B) est présent en une quantité se situant entre 0,01 et 10% en poids.
3. Composition cosmétique pour cheveux selon la revendication 1, caractérisée en ce que le rapport de
65 mélange de l'ingrédient (A) et de l'ingrédient (B) se situe dans l'intervalle allant de 0,1 à 10:1.



0 117 469

Results:

TABLE 18—B

		Feeling of the Hair	Degree of Curling	Combing Force	
				under wet conditions	under dry conditions
5					
10	V	-0.1	72%	170 (g)	194 (g)
	W	+0.1	76%	165	187
15	X	+1.3	93%	136	161

20 Claims

1. A hair cosmetic composition comprising
(A) an extract obtained by polar solvent extraction of birch or rosemary, and
(B) a polypeptide compound having a molecular weight of 350 to 30,000 selected from decomposition
25 products of silk, decomposition products of keratin and derivatives thereof.
2. A hair cosmetic composition according to Claim 1, wherein the ingredient (A) is contained in an amount ranging from 0.001 to 10 wt% and the ingredient (B) is in an amount ranging from 0.01 to 10 wt%.
3. A hair cosmetic composition according to Claim 1, wherein the mixing ratio of the ingredient (A) and the ingredient (B) is in the range of 0.1 to 10:1.
- 30 4. A hair cosmetic composition according to Claim 1, wherein the ingredient (B) is a polypeptide compound having a molecular weight of 350 to 30,000 selected from hydrolyzates of silk, hydrolyzates of keratin, oxidative decomposition products of keratin and thiol group-modified compounds of reductive decomposition products of keratin.

35

Patentansprüche

1. Haarkosmetikum-Zusammensetzung, umfassend
(A) einen Extrakt, erhalten durch Extraktion von Birke oder Rosmarin mit einem polaren Lösungsmittel,
40 und
(B) eine Polypeptidverbindung mit einem Molekulargewicht von 350 bis 30000, ausgewählt aus Zersetzungsprodukten von Seide, Zersetzungsprodukten von Keratin und Derivaten derselben.
2. Haarkosmetikum-Zusammensetzung gemäß Anspruch 1, wobei der Bestandteil (A) in einer Menge im Bereich von 0,001 bis 10 Gew.-% enthalten ist und der Bestandteil (B) in einer Menge im Bereich von
45 0,01 bis 10 Gew.-% enthalten ist.
3. Haarkosmetikum-Zusammensetzung gemäß Anspruch 1, wobei das Mischungsverhältnis von dem Bestandteil (A) und dem Bestandteil (B) im Bereich von 0,1 bis 10:1 liegt.
4. Haarkosmetikum-Zusammensetzung gemäß Anspruch 1, wobei der Bestandteil (B) eine
50 Polypeptidverbindung mit einem Molekulargewicht von 350 bis 30000 ist, ausgewählt aus Hydrolysaten von Seide, Hydrolysaten von Keratin, oxidativen Zersetzungsprodukten von Keratin und an der Thiolgruppe modifizierten Verbindungen von reduktiven Zersetzungsprodukten von Keratin.

55

Revendications

1. Composition cosmétique pour cheveux comprenant:
(A) un extrait obtenu moyennant une extraction de bouleau ou de romarin au moyen d'un solvant polaire, et
(B) un composé polypeptidique ayant un poids moléculaire de 350 à 30,000, choisi parmi les produits
60 de décomposition de la soie, les produits de décomposition de la kératine et leurs dérivés.
2. Composition cosmétique pour cheveux selon la revendication 1, caractérisée en ce que l'ingrédient (A) est présent en une quantité se situant entre 0,001 et 10% en poids, tandis que l'ingrédient (B) est présent en une quantité se situant entre 0,01 et 10% en poids.
3. Composition cosmétique pour cheveux selon la revendication 1, caractérisée en ce que le rapport de
65 mélange de l'ingrédient (A) et de l'ingrédient (B) se situe dans l'intervalle allant de 0,1 à 10:1.



0 117 469

4. Composition cosmétique pour cheveux selon la revendication 1, caractérisée en ce que l'ingrédient (B) est un composé polypeptidique ayant un poids moléculaire de 350 à 30,000, tandis qu'il est choisi parmi les hydrolysats de la soie, les hydrolysats de la kératine, les produits de décomposition de la kératine par oxydation et les composés (modifiés par un groupe thiol) des produits de décomposition de la kératine par réduction.

10

15

20

25

30

35

40

45

50

55

60

65