INDOLINE DERIVATIVES USEFUL AS PROTEIN KINASE INHIBITORS

Field of the Invention

The present invention relates to novel indolinone derivatives useful as pharmaceuticals, to processes for preparing the compounds, to intermediates useful in the preparation of the compounds, to pharmaceutical compositions comprising the compounds, and to the use of the compounds as pharmaceuticals.

Background of the Invention

WO 96/40116 disclose that certain pyrrole substituted 2-indolinone derivatives are receptor tyrosine kinase inhibitors useful in the treatment of conditions responsive to receptor tyrosine kinase inhibitors, for example proliferative disorders such as cancer. A preferred compound, disclosed on page 17, is 3-(2,3-dimethylpyrrol-5-yl)methylene]-2-indolinone, also known as SU5416. Unfortunately, this compound has been found to exhibit poor solubility in water and low bioavailability upon oral and intravenous administration.

WO 99/61422 discloses further pyrrole substituted 2-indolinone derivatives as receptor tyrosine kinase inhibitors. A preferred compound, disclosed as compound 5 on page 214, is 3-[2,4-dimethyl-5-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrol-3-yl]propionic acid, also known as SU6668. This compound has been found to possess superior oral activity to SU5416, but has been reported to lack the ability of that compound to inhibit the receptor tyrosine kinase Flt-3 (Abstract 497, Anne-Marie O'Farrell et al., America Society of Hematology Meeting, Orlando, Florida, USA, December 7-11, 2001). Flt-3 is an important target for a tyrosine kinase inhibitor, especially for the treatment of Acute Myeloid Leukemia (AML), because about 30% of AML patients have been found to possess mutant forms of Flt-3 which lead to constitutive tyrosine phosphorylation of Flt-3 (Levis et al., *Blood*, 1 August, 2001, Vol. 98. No. 3, pp 885-887).

WO 01/60814 discloses pyrrole substituted 2-indolinone derivatives bearing certain amido substituents directly attached to the pyrrole ring as receptor tyrosine kinase inhibitors.

WO 02/055517 discloses indolinones substituted with aryl substituents at the 4 position which exhibit protein kinase modulating ability.


WO 01/42243 discloses that certain compounds containing two or more pyrrole substituted 2-indolinone groups covalently linked together through the 3 position on each pyrrole by one or more linker groups are also useful as receptor tyrosine kinase inhibitors.

Nonetheless, in view of the severity of conditions responsive to receptor tyrosine kinase inhibitors and of the recent identification of specific kinase inhibitor targets, a need exists for new receptor tyrosine kinase inhibitors with diverse properties.

Summary of the Invention

Pyrrole substituted 2-indolinone derivatives bearing certain carboxamidoethyl groups at the 4 position on pyrrole have now been found that are inhibitors of receptor tyrosine kinases with particularly desirable properties.

Accordingly, the present invention provides a compound of formula (I):



(I)

in which:

(i) R^1 represents a hydrogen atom or a (1-4C)alkyl group; and R^2 represents a group of formula $-A^1-NR^5R^6$ in which each of R^5 and R^6 independently represents a hydrogen atom or a (1-4C)alkyl group and A^1 represents $(CH_2)_m$, $(CH_2)_n-A^2-(CH_2)_p$ or $(CH_2CH_2O)_qCH_2CH_2$ in which *m* is an integer of from 2 to 10, each of *n* and *p* is an integer of from 1 to 6, A^2 is CH=CH, phenylene, biphenylene, cyclohexylene or piperazinylene and *q* is 1, 2 or 3;

(ii) R^1 and R^2 together represent $-A^3-NR^7-A^4-$ in which each of A^3 and A^4 independently represents $(CH_2)_r$ or $(CH_2CH_2O)_sCH_2CH_2$ in which *r* is an integer of from 2 to 6, *s* is 1, 2 or 3, and R^7 represents a hydrogen atom or a (1-4C)alkyl group;

(iii) R^1 and R^2 together with the nitrogen atom to which they are attached represent a piperidinyl group, which piperidinyl group bears a substituent of formula $-A^5-R^8$ at the 4 position, in which A^5 represents (1-4C)alkylene and R^8 represents piperidin-4-yl; or


(iv) R^1 and R^2 together with the nitrogen atom to which they are attached represent a pyrrolidinyl, piperidinyl or morpholino group; and

R^3 and R^4 each independently represents a hydrogen atom, a halogen atom, a (1-4C)alkyl group, a (1-4C)alkoxy group, a phenyl group which is unsubstituted or substituted by one or two substituents selected independently from a halogen atom, a (1-4C)alkyl group and a (1-4C)alkoxy group, a group of formula $R^9S(O)_2NR^{10}-$, a group of formula $R^{11}N(R^{12})S(O)_2-$, a group of formula $R^{13}C(O)N(R^{14})-$ or a group of formula $R^{15}N(R^{16})C(O)-$ in which each of R^9, R^{11}, R^{13} and R^{15} independently represents a (1-4C)alkyl group or a phenyl group which is unsubstituted or substituted by one or two substituents selected independently from a halogen atom, a (1-4C)alkyl group and a (1-4C)alkoxy group, and each of R^{10}, R^{12}, R^{14} and R^{16} independently represents a hydrogen atom or a (1-4C)alkyl group;

or a pharmaceutically-acceptable salt thereof.

Compounds of formula (I) have been found to be potent and selective inhibitors of one or more of the receptor tyrosine kinases PDGFR (platelet-derived growth factor), c-Kit, VEGFR (vascular endothelial growth factor) and Flt-3 in whole cell assays.

In one embodiment, there is provided compounds of formula (Ia):




(Ia)

wherein R is hydrogen, methyl, or ethyl;

or a pharmaceutically-acceptable salt thereof.

The invention provides a process for the preparation of a compound as defined above which comprises:

(a) reacting a compound of formula (II)



(II)


or a reactive derivative thereof, with a compound of formula (III)

$$HNR^1R^2$$

(III)

or a salt thereof, in which R^1, R^2, R^3 and R^4 are as defined in Claim 1, or

(b) for a compound of formula (I) in which R^5 or R^7 represents a hydrogen atom, deprotecting a compound of formula (IV)



(IV)

in which R^{1a} and R^{2a} are as defined in Claim 1 for R^1 and R^2, except in that R^5 or R^7 is replaced with a group R^{5a} or R^{7a} respectively, in which R^{5a} and R^{7a} each represents an amine protecting group, and R^3 and R^4 are as defined above;

followed, if a pharmaceutically-acceptable salt is required, by forming a pharmaceutically-acceptable salt.

The invention also provides pharmaceutical compositions comprising a compound of the invention or a pharmaceutically-acceptable salt thereof and a pharmaceutically-acceptable carrier.

Disclosed herein is a method of treating a condition responsive to a tyrosine kinase inhibitor, the method comprising administering to a patient in need of treatment an effective amount of a compound of the invention.

Further, the invention provides a compound of the invention as described herein for use in medical therapy, as well as the use of a compound of the invention in the manufacture of a formulation or medicament for treating a disease or condition responsive to a tyrosine kinase inhibitor.

Detailed Description

The present invention provides novel pyrrole substituted 2-indoline derivatives which are substituted at the 4 position of the pyrrole ring with carboxamidoethyl substituents.

As used herein, the terms alkyl and alkylene refer to a branched or unbranched group. However, the names of specific groups, such as ethyl, ethylene, propyl, propylene, butyl or butylene, signify unbranched groups or radicals, unless indicated otherwise, such as prop-2-yl. Examples of alkyl groups are methyl, ethyl, propyl, prop-2-yl, and butyl. Examples of alkylene groups are methylene, ethylene, propylene and butylene.

The term halogen atom includes fluorine, chlorine and bromine. The term "therapeutically effective amount' refers to an amount sufficient to effect treatment when administered to a patient hi need of treatment

The term "treatment" as used herein refers to the treatment of a disease or medical condition in a patient, such as a mammal (particularly a human), and includes:

(a) preventing the disease or medical condition from occurring, i.e., prophylactic treatment of a patient;

(b) ameliorating the disease or medical condition, i.e., eliminating or causing regression of the disease or medical condition in a patient;

(c) suppressing the disease or medical condition, i.e., slowing or arresting the development of the disease or medical condition in a patient; or

(d) alleviating the symptoms of the disease or medical condition in a patient

The term "pharmaceutically-acceptable salt" refers to a salt prepared from a base or acid which is acceptable for administration to a patient, such as a mammal. Such salts can be derived from pharmaceutically-acceptable inorganic or organic acids.

Salts derived from pharmaceutically-acceptable acids include acetic, benzenesulfonic, benzoic, camphosulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pantothenic, phosphoric, succinic, sulfuric, tartaric, *p*-toluenesulfonic, xinafoic (1-hydroxy-2-naphthoic acid) and the like. Particularly preferred are salts derived from fumaric, hydrobromic, hydrochloric, acetic, sulfuric, phosphoric, methanesulfonic, *p*-toluenesulfonic, xinafoic, tartaric, citric, malic, maleic, succinic, and benzoic acids.

One preferred sub-group of compounds of formula (I) is that in which:

(i) R^1 represents a hydrogen atom or a (1-4C)alkyl group; and R^2 represents a group of formula $-A^1-NR^5R^6$ in which each of R^5 and R^6 independently represents a hydrogen atom or a (1-4C)alkyl group and A^1 represents $(CH_2)_m$, $(CH_2)_n-A^2-(CH_2)_p$ or $(CH_2CH_2O)_qCH_2CH_2$ in which *m* is an integer of from 2 to 10, each of *n* and *p* is an integer of from 1 to 6, A^2 is CH=CH, phenylene, biphenylene, cyclohexylene or piperazinylene and *q* is 1, 2 or 3;

(ii) R^1 and R^2 together represent $-A^3-NR^7-A^4-$ in which each of A^3 and A^4 independently represents $(CH_2)_r$ or $(CH_2CH_2O)_sCH_2CH_2$ in which *r* is an integer of from 2 to 6, *s* is 1, 2 or 3, and R^7 represents a hydrogen atom or a (1-4C)alkyl group; or

(iii) R^1 and R^2 together with the nitrogen atom to which they are attached represent a piperidinyl group, which piperidinyl group bears a substituent of formula $-A^5-R^8$ at the 4 position, in which A^5 represents (1-4C)alkylene and R^8 represents piperidin-4-yl.

Compounds belonging to the above preferred sub-group have been found to exhibit good solubility in water and good absorption on oral administration.

In this sub-group of compounds, preferably

(i) R^1 represents a hydrogen atom or a (1-4C)alkyl group; and R^2 represents a group of formula $-A^1-NR^5R^6$ in which each of R^5 and R^6 independently represents a hydrogen atom or a (1-4C)alkyl group and A^1 represents $(CH_2)_m$, $(CH_2)_n-A^2-(CH_2)_p$ or $(CH_2CH_2O)_qCH_2CH_2$ in which *m* is an integer of from 2 to 10, each of *n* and *p* is an integer of from 1 to 6, A^2 is CH=CH, phenyl-1,3-ene, phenyl-1,4-ene, biphenyl-2,2'-ene, cyclohex-1,3-ylene or piperazin-1,4-ylene and q is 1, 2 or 3;

(ii) R^1 and R^2 together represent $-A^3-NR^7-A^4-$ in which each of A^3 and A^4 independently represents $(CH_2)_r$ or $(CH_2CH_2O)_sCH_2CH_2$ in which *r* is an integer of from 2 to 6, *s* is 1, 2 or 3, and R^7 represents a hydrogen atom or a (1-4C)alkyl group; or

(iii) R^1 and R^2 together with the nitrogen atom to which they are attached represent a piperidinyl group, which piperidinyl group bears a substituent of formula $-A^5-R^8$ at the 4 position, in which A^5 represents (1-4C)alkylene and R^8 represents piperidin-4-yl.

Preferably

(i) R^1 represents a methyl group; and R^2 represents a group of formula $-A^1-NR^5R^6$ in which R^5 represents a hydrogen atom, R^6 represents a methyl group and A^1 represents $(CH_2)_m$, $(CH_2)_n-A^2-(CH_2)_p$ or $(CH_2CH_2O)_qCH_2CH_2$ in which *m* is an integer of from 2 to 10, each of *n* and *p* is 1 or 2, A^2 is CH=CH, phenyl-1,3-ene, phenyl-1,4-ene, biphenyl-2,2'-ene, cyclohex-1,3-ylene or piperazin-1,4-ylene and q is 1, 2 or 3;

(ii) R^1 and R^2 together represent $-A^3-NR^7-A^4-$ in which each of A^3 and A^4 independently represents $(CH_2)_r$ or $(CH_2CH_2O)_sCH_2CH_2$ in which *r* is an integer of from 2 to 6, *s* is 1 or 2, and R^7 represents a hydrogen atom or a (1-4C)alkyl group; or

(iii) R^1 and R^2 together with the nitrogen atom to which they are attached represent a piperidinyl group, which piperidinyl group bears a substituent of formula $-A^5-R^8$ at the 4 position, in which A^5 represents propylene and R^8 represents piperidin-4-yl.

More preferably,

(i) R^1 represents a methyl group; and R^2 represents a group of formula $-A^1-NR^5R^6$ in which R^5 represents a hydrogen atom, R^6 represents a methyl group and A^1 represents $(CH_2)_m$, in which *m* is 2, 3, 4, 5, 6, 7, 8, 9 or 10; $(CH_2)_n-A^2-(CH_2)_p$ in which n and p are each 1 and A^2 is CH=CH, phenyl-1,3-ene, phenyl-1,4-ene, biphenyl-2,2'-ene or cyclohex-1,3-ylene; $(CH_2)_n-A^2-(CH_2)_p$ in which *n* and *p* are each 2 and A^2 is piperazin-1,4-ylene; or $(CH_2CH_2O)_qCH_2CH_2$ in which *q* is 2 or 3;

(ii) R^1 and R^2 together represent $-(CH_2)_2-NH-(CH_2)_2-$, $-(CH_2)_2-N(CH_3)-(CH_2)_2-$, $-(CH_2)_2-N(CH_2CH_3)-(CH_2)_2-$, $-(CH_2)_2-NH-(CH_2)_3-$, or $-(CH_2CH_2O)_2CH_2CH_2-NH-(CH_2CH_2O)CH_2CH_2-$; or

(iii) R^1 and R^2 together with the nitrogen atom to which they are attached represent a piperidinyl group, which piperidinyl group bears a substituent of formula $-A^5-R^8$ at the 4 position, in which A^5 represents propylene and R^8 represents piperidin-4-yl.

A particularly preferred sub-group of compounds is that in which R^1 represents a methyl group and R^2 represents a group of formula $-A^1-NR^5R^6$ in which R^5 represents a hydrogen atom, R^6 represents a methyl group and A^1 represents $(CH_2)_m$ or $CH_2-CH=CH-CH_2$, in which *m* is an integer of from 2 to 6.

Compounds belonging to this sub-group have been found to exhibit particularly good potency as inhibitors of one or more of the above receptor tyrosine kinases.

Within this sub-group, preferably A^1 represents $(CH_2)_m$ or $CH_2-CH=CH-CH_2$, in which *m* is 2, 3 or 4.

More preferably A^1 represents $(CH_2)_2$,$(CH_2)_3$ or $CH_2-CH=CH-CH_2$.

Especially preferred are compounds in which A^1 represents$(CH_2)_2$.

Another preferred sub-group of compounds is that in which R^1 and R^2 together represent $-A^3-NR^7-A^4-$ in which each of A^3 and A^4 independently represents$(CH_2)_r$ or $(CH_2CH_2O)_sCH_2CH_2$ in which *r* is an integer of from 2 to 6, and *s* is 1, 2 or 3, and R^7 represents a hydrogen atom or a (1-4C)alkyl group.

Compounds belonging to this sub-group have also been found to exhibit particularly good potency.

In this sub-group, preferably R^1 and R^2 together represent $-(CH_2)_2-NR^7-(CH_2)_2-$ or $-(CH_2)_2-NR^7-(CH_2)_3-$, especially $-(CH_2)_2-NR^7-(CH_2)_2-$.

Examples of particular values for R^7 are hydrogen, methyl, ethyl, propyl, prop-2-yl and butyl.

Compounds in which R^7 represents hydrogen are especially preferred.

Referring to R^3 and R^4, examples of particular values are:

hydrogen;

for a halogen atom: fluorine, chlorine or bromine, especially bromine;

for a (1-4C)alkyl group: methyl;

for a (1-4C)alkoxy group: methoxy;

for an unsubstituted or substituted phenyl group: phenyl;

for R^8, R^{10}, R^{12} and R^{14}: methyl or phenyl;

for R^9, R^{11}, R^{13} and R^{15}: hydrogen; and

for a group of formula $R^{12}C(O)N(R^{13})$-: $CH_3C(O)NH$- and $C_6H_5C(O)NH$-.

Preferably R^3 and R^4 each independently represents a hydrogen atom, a bromine atom, $CH_3C(O)NH$-, or $C_6H_5C(O)NH$-. More preferably R^3 and R^4 each independently represents a hydrogen atom.

Another preferred group of compounds of formula (I) are compounds in which:

(i) R^1 represents a methyl group and R^2 represents a group of formula $-A^1-NHCH_3$ in which A^1 represents $(CH_2)_m$, $CH_2CH=CHCH_2$, CH_2-phenylene-CH_2, or CH_2-cyclohexylene-CH_2, in which *m* is an integer of from 2 to 8; or

(ii) R^1 and R^2 together represent $-(CH_2)_2-NH-(CH_2)_2-$, $-(CH_2)_2-N(CH_3)-(CH_2)_2-$, $-(CH_2)_2-N(CH_2CH_3)-(CH_2)_2-$ or $-(CH_2)_2-NH-(CH_2)_3-$; and

R^3 and R^4 are each independently hydrogen.

Compounds of the above sub-group have been found to exhibit particularly good potency as inhibitors of one or more receptor tyrosine kinases. In particular, such compounds have demonstrated IC_{50} values for inhibition of the VEGFR tyrosine kinase of less than 1 μM in the intracellular Ca^{2+} FLIPR or immunoprecipitation assay described below.


A more preferred sub-group of compounds within the above sub-group are compounds in which:

(i) R^1 represents a methyl group and R^2 represents a group of formula $-A^1-NHCH_3$ in which A^1 represents $(CH_2)_m$, $CH_2CH=CHCH_2$, or CH_2-(1,4-phenylene)-CH_2 in which *m* is 2 or 3; or

(ii) R^1 and R^2 together represent $-(CH_2)_2-NH-(CH_2)_2-$, $-(CH_2)_2-N(CH_3)-(CH_2)_2-$ $-(CH_2)_2-N(CH_2CH_3)-(CH_2)_2-$ or $-(CH_2)_2-NH-(CH_2)_3-$.

Compounds belonging to this more preferred sub-group of compounds have demonstrated IC_{50} values for inhibition of both the VEGFR and PDGFR tyrosine kinases of less than 1 μM in the intracellular Ca^{2+} FLIPR or immunoprecipitation assay described below.

A particularly preferred sub-group of compounds of formula (I) are compounds of formula (Ia):



(Ia)

wherein R is hydrogen, methyl, or ethyl, and pharmaceutically-acceptable salts thereof.

Compounds of formula (Ia) that may be given special mention are:

3-[3,5-dimethyl-4-(3-oxo-3-piperazin-1-ylpropyl)-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one and

3-[3,5-dimethyl-4-[3-oxo-3-(4-ethyl)piperazin-1-ylpropyl]-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one.


Especially preferred is the compound 3-[3,5-dimethyl-4-(3-oxo-3-piperazin-1-ylpropyl)-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one, and pharmaceutically-acceptable salts thereof. This compound has been found to be a highly potent and selective inhibitor of PDGFR, c-Kit, VEGFR and Flt-3. It has also been found to have high solubility in water and to possess excellent absorption when administered orally to rats.

Another compound of formula (I) that may be given special mention is 3-[3,5-dimethyl-4-(3-oxo-3-homopiperazin-1-ylpropyl)-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one.

The compounds of formula (I) are useful as receptor tyrosine kinase inhibitors for the treatment of proliferative disorders, such as forms of cancer which include, but are not limited to acute myeloid leukemia, small cell lung cancer, prostate cancer, gastrointestinal cancer, breast cancer and brain cancer, and other proliferative disorders, such as restenosis. The compounds may also be useful in restricting the growth of solid tumors.

According to another aspect, the present invention provides a process for the preparation of a compound of formula (I), which comprises

(a) reacting a compound of formula (II)



(II)


or a reactive derivative thereof, with a compound of formula (III)

HNR^1R^2

(III)

or a salt thereof, in which R^1, R^2, R^3 and R^4 are as defined hereinabove, or

(b) for a compound of formula (I) in which R^5 or R^7 represents a hydrogen atom, deprotecting a compound of formula (IV)



(IV)

in which R^{1a} and R^{2a} are as defined hereinabove for R^1 and R^2, except in that R^5 or R^7 is replaced with a group R^{5a} or R^{7a} respectively, in which R^{5a} and R^{7a} each represents an amine protecting group, and R^3 and R^4 are as defined hereinabove;

followed, if a pharmaceutically-acceptable salt is required, by forming a pharmaceutically-acceptable salt.

In process (a), the reaction of a compound of formula (II) with a compound of formula (III) may conveniently be performed using a conventional amide coupling method. For example, an acid of formula (II) may be treated with a coupling agent, such as benzotriazol-1-yl-oxy-trispyrrolidinophosphonium hexafluorophosphate (PyBOP) or o-(7-azabenzotriazol-1-yl)-N,N,N'N'-tetramethyluronium hexafluorophosphate (HATU), in the presence of a base, such as N,N-diisopropylethylamine, and 1-hydroxy-7-azabenzotriazole (HOAt), followed by addition of the compound of formula (III). Convenient solvents include polar aprotic organic solvents, such as dimethylformamide. The temperature is conveniently in the range of from 0 to 50°C. Alternatively, the compound of formula (II) may be converted into an acid halide, such as the chloride, and then reacted with the compound of formula (III).

In process (b), the amine protecting group represented by R^{5a} or R^{7a} may be a conventional amine protecting group. Examples of amine protecting groups are described in Greene and Wuts, *Protecting Groups in Organic Synthesis*, 2nd Edition, John Wiley & Sons, NY, 1991 and McOmie, *Protecting Groups in Organic Chemistry*, Plenum Press, NY, 1973. Examples of amine protecting groups include acyl groups, for example (1-6C)alkanoyl groups, such as acetyl; (1-6C)alkoxycarbonyl groups, such as

t-butoxycarbonyl; and arylmethoxycarbonyl groups, such as benzyloxycarbonyl; and arylmethyl groups, such as benzyl.

An acyl amine protecting group may conveniently be removed by treatment with an acid, such as trifluoroacetic acid.


Compounds of formula (IV) may be prepared by reacting a compound of formula (II) with an amine of formula (V)

$$HNR^{1a}R^{2a}$$

(V)


in which R^{1a} and R^{2a} are as defined hereinabove, following the method of process step (a).

Compounds of formula (II) are known, for example from WO 99/61422. They may also be prepared by reacting a compound of formula (VI)



(VI)

with a compound of formula (VII)




(VII)

The reaction is conveniently performed in the presence of a base, such as piperidine, in an organic solvent, such as ethanol, and under reflux.

Compounds of formula (VII) are known, for example from WO 99/61422.

Compounds of formula (VI) may be prepared by reacting a compound of formula (VIII)



(VIII)

in which R^{8a} represents a carboxyl protecting group, for example a (1-6C)alkyl group such as methyl, with phosphorus oxychloride and dimethylformamide, followed by removal of the protecting group R^{8a}, for example by alkali hydrolysis.

Compounds of formula (VIII) may be prepared via the corresponding carboxylic acid (R^{8a} is hydrogen) following methods as described in the accompanying examples.

Certain of the intermediates described herein are believed to be novel, for example the compounds of formula (IV). All such novel intermediates are provided as further aspects of the invention.

Pharmaceutical Compositions

When used as pharmaceuticals, the compounds of the invention will usually be administered in a pharmaceutical composition. The compositions comprise a compound of the invention as the active ingredient, together with a pharmaceutically-acceptable diluent or carrier. The compositions may be formulated for any route of administration, in particular for oral, rectal, transdermal, subcutaneous, intravenous, intramuscular or intranasal administration. The compositions may be formulated in any conventional form, for example, as tablets, capsules, solutions, suspensions, dispersions, syrups, sprays, gels, suppositories, patches and emulsions.

The preparation of a suitable pharmaceutical composition for a particular mode of administration is well within the scope of those skilled in the pharmaceutical arts. Additionally, the ingredients for such compositions are commercially available from, for example, Sigma (St. Louis, MO). By way of further illustration, conventional formulation techniques are described in *Remington: The Science and Practice of Pharmacy*, 20th Edition, Lippincott Williams & White, Baltimore, MD (2000); and H. C. Ansel et al., *Pharmaceutical Dosage Forms and Drug Delivery Systems*, 7th Edition, Lippincott Williams & White, Baltimore, MD (1999).

According to another aspect, the present invention provides a pharmaceutical composition, which comprises a therapeutically-effective amount of a compound of formula (I) or a pharmaceutically-acceptable salt thereof, together with a pharmaceutically-acceptable diluent or carrier.

In a preferred embodiment, the pharmaceutical compositions of the invention are suitable for oral administration. Suitable pharmaceutical compositions for oral administration may be in the form of capsules, tablet, pills, lozenges, cachets, dragees, powders, granules, or as a solution or suspension in a liquid, and the like; each containing a predetermined amount of a compound of the present invention as an active ingredient. A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid compositions. Examples of such carriers

include magnesium stearate, starch, lactose, sucrose, microcrystalline cellulose and binders, for example polyvinylpyrrolidone. In addition, the active compound can be formulated in a controlled release dosage form as a tablet comprising a hydrophilic or hydrophobic matrix.

A composition in the form of a capsule can be prepared using routine encapsulation procedures, for example, by incorporation of active compound and excipients into a hard gelatin capsule. Alternatively, a semi-solid matrix of active compound and high molecular weight polyethylene glycol can be prepared and filled into a hard gelatin capsule; or a solution of active compound in polyethylene glycol or a suspension in edible oil, can be prepared and filled into a soft gelatin capsule.

In another preferred embodiment, the compound of the invention can be formulated for injection, for example for intravenous injection. A typical composition for intravenous injection consists of a sterile isotonic aqueous solution containing, for example, active compound and dextrose or sodium chloride, or a mixture of dextrose and sodium chloride. Other examples of suitable excipients include lactated Ringer's injection, lactated Ringer's plus dextrose injection, Normosol-M and dextrose, Isolyte E, acylated Ringer's injection, and the like. Optionally, a co-solvent, for example, polyethylene glycol; a chelating agent, for example, ethylenediamine tetraacetic acid; a stabilizing agent, for example, a cyclodextrin; and an anti-oxidant, for example, sodium metabisulphite, may be included in the formulation.

According to another aspect, the present invention provides a compound of formula (I) or a pharmaceutically-acceptable salt thereof for use in therapy.

The compounds of formula (I) are useful as receptor tyrosine kinase inhibitors. According to another aspect, therefore, the present invention provides the use of a compound of formula (I) or a pharmaceutically-acceptable salt thereof, for the manufacture of a medicament for the treatment of a condition responsive to a tyrosine kinase inhibitor.

According to yet another aspect, the present invention provides a pharmaceutical composition for use in the treatment of a condition responsive to a tyrosine kinase inhibitor, which comprises a compound of formula (I) or a pharmaceutically-acceptable salt thereof.

The present invention also provides a method of treating a condition responsive to a tyrosine kinase inhibitor, which comprises administering to a patient in need of

treatment an effective amount of a compound of formula (I) or a pharmaceutically-acceptable salt thereof.

The patient may be, for example, a mammal, such as a companion animal, and is preferably a human.

The dose (or effective amount) of the compound administered to a patient will depend upon many factors, including the particular compound used, the nature and severity of the condition being treated, the species of the patient, the weight of the patient and the route of administration. In general, a dose in the range of from 0.01 to 100 μM/kg of bodyweight will be administered.

The following non-limiting examples illustrate representative pharmaceutical compositions of the invention.

Formulation Example A

Hard gelatin capsules for oral administration are prepared as follows:

Ingredients	Amount
Compound of the invention	250 mg
Lactose (spray-dried)	200 mg
Magnesium stearate	10 mg

Representative Procedure: The ingredients are thoroughly blended and then loaded into a hard gelatine capsule (460 mg of composition per capsule)

Formulation Example B

Hard gelatin capsules for oral administration are prepared as follows:

Ingredients	Amount
Compound of the invention	20 mg
Starch	89 mg
Microcrystalline cellulose	89 mg
Magnesium stearate	10 mg

Representative Procedure: The ingredients are thoroughly blended and then passed through a No. 45 mesh U.S. sieve and loaded into a hard gelatine capsule (200 mg of composition per capsule)

Formulation Example C

Capsules for oral administration are prepared as follows:

Ingredients	Amount
Compound of the invention	100 mg
Polyoxyethylene sorbitan monooleate	50 mg
Starch powder	250 mg

Representative Procedure: The ingredients are thoroughly blended and then loaded into a hard gelatine capsule (300 mg of composition per capsule)

Formulation Example D

Tablets for oral administration are prepared as follows:

Ingredients	Amount
Compound of the invention	250 mg
Microcrystalline cellulose	400 mg
Silicon dioxide fumed	10 mg
Stearic acid	5 mg

Representative Procedure: The ingredients are thoroughly blended and then loaded into a hard gelatine capsule (460 mg of composition per capsule)

Formulation Example E

An injectable formulation is prepared as follows:

Ingredients	Amount
Compound of the invention	0.2 g
Sodium acetate buffer solution (0.4 M)	400 mg
HCl (0.5 N) or NaOH (0.5 N)	q.s. to pH 4
Water (distilled, sterile)	q.s. to 20 mL

Representative Procedure: The above ingredients are blended and the pH is adjusted to 4 ± 0.5 using 0.5 N HCl or 0.5 N NaOH.

Synthetic Examples

The following synthetic examples are offered to illustrate the invention, and are not to be construed in any way as limiting the scope of the invention.

General

Reagents and solvents were used as received from commercial suppliers unless otherwise noted. All reactions were carried out at room temperature and without rigorous exclusion of ambient atmosphere unless otherwise noted. Ion-spray mass spectra (IS-MS) were obtained using a PE Sciex API 150EX mass spectrometer. Nuclear magnetic resonance (NMR) spectra were recorded at 300 MHz. Chemical shifts (δ) are reported in parts per million downfield of tetramethylsilane. Analytical reversed-phase HPLC (RP-HPLC) was performed on an HP1100 instrument using a 2.1 mm x 50 mm, 3.5 μm C_{18} Zorbax Plus Bonus-RP column. For the analytical separations, a 0.5 minute isocratic period was followed by a 4.5 minute gradient of 0.1% trifluoroacetic acid/acetonitrile (ACN) in 0.1% water at a flow rate of 0.5 mL/minute. Preparative RP-HPLC was performed using trifluoroacetic acid (TFA) buffered ACN/water gradients on a Varian ProStar system using 2.5- or 10 cm x 25 cm, 8 μm C_{18} Rainin Dynamax columns and flow rates of 10- or 50 mL/minute, respectively.

Preparation of Intermediates

Intermediate 1
2-Carboxyethyl-3,5-dimethyl-1H-pyrrole-4-carboxylic acid

3,5-Dimethyl-2,4-pyrrole dicarboxylic acid, diethyl ester (200 g, 836 mmol) was placed in a 1 L beaker and treated with 400 mL concentrated sulfuric acid (H_2SO_4). The mixture was stirred, heated to 45 C with the aid of a heat gun and then maintained at 36-42 C for 25 minutes. The reaction mixture was poured into 3 L crushed ice and stirred for 30 minutes. The yellow solid was recovered by filtration and washed with 200 mL water. The solid was transferred to a 4 L Erlenmeyer flask and treated with 2 L 1 N sodium hydroxide solution (NaOH) followed by 100 mL 10 N NaOH. The basic mixture was filtered and the yellow solid residue was discarded. The filtrate was acidified with H_2SO_4. The resulting solid was recovered by suction filtration and washed with 2 x 500 mL water. After suction drying, the material was transferred to a 6 L Erlenmeyer flask and digested briefly in 4 L acetone. After standing overnight at room temperature (RT), the solid was collected by filtration and dried in a vacuum dessicator to afford 139 g, 659

mmol, 79% of the title compound. ^{1}H NMR (DMSO-d_6) δ 4.22 (q, 2H), 2.44 (s, 3H), 2.38 (s, 3H), 1.27 (t, 3H).

Intermediate 2
2-Carboxyethyl-3,5-dimethyl-1H-pyrrole

Intermediate 1 (137 g, 650 mmol) was placed in a 500 mL Erlenmeyer flask and treated with ethanolamine (80 g, 1.3 mol). The mixture was then heated to 220 C in a heating mantle, producing a brown solution over approximately 30 minutes, at which time gas evolution had essentially ceased. The reaction was heated for 30 minutes more, then poured into 2 L ice water. The crude product was collected by suction filtration and then digested in 700 mL 95% ethanol (EtOH). The mixture was filtered while hot and the filtrate was slowly cooled to RT and then to –20 C. The resulting crystals were collected by suction filtration and dried in a vacuum dessicator to afford 75.6 g, 453 mmol, 70% of the title compound. ^{1}H NMR (DMSO-d_6) δ5.72 (s, 1H), 4.17 (q, 2H), 2.18 (s, 3H), 2.13 (s, 3H), 1.25 (t, 3H).

Intermediate 3
2-Carboxyethyl-3,5-dimethyl-1H-pyrrole-4-carboxaldehyde

Intermediate 2 (75.6 g, 453 mmol) was placed in a dry 1 L, 3-necked round-bottom flask. The solid was treated with anhydrous *N,N*-dimethylformamide (DMF, 43.8 mL , 566 mmol). The flask was shaken to distribute the DMF throughout the solid. The flask was cooled in an ice bath and to the mixture was added phosphorous oxychloride ($POCl_3$, 52.7 mL, 566 mmol) over 30 minutes via an addition funnel. The flask was shaken to evenly distribute the reagents. The flask was then immersed in a 100 C oil bath and heated with magnetic stirring for 6 hours. The resulting deep red mixture was cooled in an ice water bath and treated with 200 mL ice water, resulting in a vigorous, exothermic reaction. After addition of 200 mL additional ice water, the mixture was adjusted to pH 5 with a saturated solution of sodium acetate (NaOAc). The crude product was isolated by suction filtration and recrystallized from 700 mL hot 1:1 EtOH:water to afford 65.8 g, 337 mmol, 74% of the title compound as dark needles. ^{1}H NMR (DMSO-d_6) δ 9.88 (s, 1H), 4.23 (q, 2H), 2.46 (s, 3H), 2.43 (s, 3H), 1.28 (t, 3H).

Intermediate 4
3-(5-Carboxyethyl-2,4-dimethyl-1H-pyrrol-3-yl)propenoic acid

Intermediate 3 (65.8 g, 337 mmol) and malonic acid (39.0 g, 375 mmol) were combined in a 500 mL, 1-necked round bottom flask, treated with 350 mL absolute EtOH, and brought to reflux for 30 minutes. To the resulting dark solution was added aniline (34.0 mL, 375 mmol) and the mixture was refluxed for an additional 5 hours. Solvent was removed under reduced pressure and the residue was treated with 400 mL 2.5 M hydrochloric acid (HCl), warmed, and then allowed to cool to RT. A purple solid was collected by suction filtration, transferred to a 1 L beaker and treated with 250 mL 2 N NaOH with stirring. The resulting slurry was filtered and the solids were washed with 100 mL dilute base. The purple residue was discarded. The red filtrate was cooled in an ice bath, stirred and acidified with approximately 70 mL 6 M HCl. The resulting thick, white paste was collected by suction filtration, washed with water, and dried in air to afford the title compound. ^{1}H NMR (DMSO-d_6) δ7.52 (d, 1H), 5.93 (d, 1H), 4.22 (q, 2H), 2.35 (s, 3H), 2.31 (s, 3H), 1.28 (t, 3H).

Intermediate 5
3-(2-Carboxyethyl-2,4-dimethyl-1H-pyrrol-3-yl)propionic acid

Intermediate 4 was dissolved in 220 mL 2 N NaOH and combined with 3.5 g 10% palladium on activated carbon. The mixture was hydrogenated at 50 psi for 28 hours, then suction filtered over Celite. The Celite was washed with 50 mL water. An aliquot of the filtrate was evaporated to dryness and complete reduction to the title compound was confirmed by ^{1}H NMR (D_2O) δ 4.07 (q, 2H), 2.46 (t, 2H), 2.06 (s, 3H), 2.05 (t, 2H), 2.00 (s, 3H), 1.13 (t, 3H). The remaining filtrate was carried forward to the next step without further manipulation.

Intermediate 6
3-(2,4-Dimethyl-1H-pyrrol-3-yl)propionic acid

The filtrate from the previous step containing Intermediate 6 was treated with 30 mL 10 N NaOH and heated at reflux for 20 hours. An aliquot of the reaction mixture was acidified and evaporated to dryness. Complete hydrolysis/decarboxylation to the title compound was confirmed by ^{1}H NMR (DMSO-d_6) δ9.86 (s, 1H), 6.18 (2, 2H), 2.42 (m, 2H), 2.03 (s, 3H), 1.93 (m, 2H), 1.87 (s, 3H). The reaction mixture was carried forward to the next step without further manipulation.

Intermediate 7
Methyl 3-(2,4-dimethyl-1H-pyrrol-3-yl)propionate

The solution of Intermediate 6 from the previous step was concentrated at 60 C under reduced pressure to approximately 200 mL, cooled in an ice water bath and acidified to pH 2 using approximately 50 ml 50% H_2SO_4. The resulting mixture was filtered through a glass frit. The filtrate was extracted with 2 x 100 mL diethyl ether (Et_2O) and the residue was extracted with 3 x 100 mL Et_2O. The combined red organic extracts were washed with 2 x 100 mL water, transferred to a 2 L Erlenmeyer flask, and treated with stirring with 680 mL of an ethereal solution of diazomethane. After stirring 30 minutes at RT, excess diazomethane was quenched with glacial acetic acid (HOAc). The reaction mixture was extracted with 2 x 200 mL saturated aqueous sodium bicarbonate ($NaHCO_3$), dried over anhydrous magnesium sulfate ($MgSO_4$), filtered, and evaporated to afford 52.6 g of crude red oil. This was purifed by bulb-to-bulb distillation at 145 C and 0.2 mm Hg pressure to afford the title compound (38.4 g, 211 mmol, 63% overall from Intermediate 3). ^{1}H NMR (DMSO-d_6) δ9.92 (s, 1H), 6.22 (s, 1H), 3.54 (s, 3H), 2.52 (t, 2H), 2.32 (t, 3H), 2.02 (s, 3H), 1.87 (s, 3H).

Intermediate 8
3-(5-Formyl-2,4-dimethyl-1H-pyrrol-3-yl)propionic acid

To a dry 250 mL 3-necked, round-bottom flask was added anhydrous DMF (13.8 g mL, 189 mmol). This was cooled in an ice water bath and treated with $POCl_3$ (15.0 mL, 160 mmol) dropwise over 10 minutes. The mixture was diluted with 120 mL anhydrous 1,2-dichloroethane (DCE) and warmed to RT, producing a pale orange solution. The mixture was cooled to –10 C in an ice brine bath, at which point a precipitate formed. Intermediate 7 (14.5 g, 80.0 mmol) dissolved in 30 mL DCE was added dropwise over 10 minutes. The reaction mixture was removed from the cooling bath, stirred at RT for 10 minutes and then evaporated under reduced pressure at 30 C. The residue was transferred to a 2 L beaker using approximately 100 mL methanol (MeOH), treated with 800 mL 2 N NaOH, heated to 90 C and then allowed to cool to RT. The orange solution was extracted with 2 x 200 mL Et_2O, heated to 50 C, treated with activated charcoal, cooled to RT, and filtered through a pad of Celite. The filtrate was cooled in an ice water bath and acidified to pH 3 using approximately 120 mL 6 N HCl. The resulting solid was collected by suction filtration, washed with 3 x 40 mL water, and dried in air to afford 11.2 g, 57.0

mmol, 72% of the title compound as a brownish powder. ^{1}H NMR (DMSO-d_6) δ9.40 (s, 1H), 2.52 (t, 2H), 2.29 (t, 2H), 2.19 (s, 3H), 2.14 (s, 3H).

Intermediate 9
3-[2,4-Dimethyl-5-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrol-3-yl]propionic acid

Intermediate 8 (11 g, 57 mmol) and oxindole (7.6 g, 57 mmol) were combined in a 200 mL round-bottom flask, slurried in 150 mL EtOH, treated with piperidine (8.5 mL, 86 mmol), and heated to reflux for 4 hours. The reaction mixture was cooled to RT, treated with HOAc (14.4 mL, 250 mmol), returned briefly to reflux, cooled again and filtered. The orange solid was collected by suction filtration, washed with 100 mL hot 1:1 HOAc:EtOH followed by 100 mL hot EtOH, and dried in air to afford 15 g, 50 mmol, 87% of the title compound. ^{1}H NMR (DMSO-d_6) δ10.8 (s, 1H), 7.71 (d, 1H), 7.55 (s, 1H), 7.07 (t, 1H), 6.95 (t, 1H), 6.96 (d, 1H), 2.63 (t, 2H), 2.33 (t, 2H), 2.28 (s, 3H), 2.25 (s, 3H).

Intermediate 10
4-{3-[2,4-Dimethyl-5-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrol-3-yl]-propionyl}-piperazine-1-carboxylic acid tert-butyl ester

Intermediate 9 (6.2 g, 20 mmol), *mono*-Boc piperazine (4.1 g, 22 mmol), and 1-hydroxy-7-azabenzotriazole (HOAT, 3.0 g, 22 mmol) were dissolved in 50 mL anhydrous DMF and treated with *N,N*-diisopropylethylamine (DIEA, 3.5 mL, 20 mmol) followed by benzotriazol-1-yl-oxy-trispyrrolidino-phosphonium hexafluorophosphate (PyBOP, 11.4 g, 22 mmol). The resulting mixture was stirred overnight at RT, depositing a yellow solid. The solid was collected by suction filtration, washed with DMF and ACN and dried to afford 1.3 g of Intermediate 10. The filtrate was evaporated and fractionated by chromatography on 700 cc of silica gel using 5% MeOH in dichloromethane (DCM) eluent. Product-containing fractions were combined and evaporated, then digested in 100 mL ACN. After cooling to RT, the fine yellow solid was collected by suction filtration, washed with 2 x 20 mL ACN, and dried to afford an additional 5.7 g of the title compound. In all, 7.0 g, 15 mmol, 75% of product was obtained. ^{1}H NMR (DMSO-d_6) δ10.7 (s, 1H), 7.70 (d, 1H), 7.55 (s, 1H), 7.07 (t, 1H), 6.95 (t, 1H), 6.84 (d, 1H), 3.41-3.19 (m, 8H), 2.62 (t, 2H), 2.43 (t, 2H), 2.28 (s, 3H), 2.24 (s, 3H), 1.35 (s, 9H). IS-MS, calcd. *m/z* for $C_{27}H_{34}N_4O_4$ $[M]^+$: 478; obsd. 478.2.

Example 1
3-[3,5-dimethyl-4-(3-oxo-3-piperazin-1-yl-propyl)-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one, trifluoroacetate

Intermediate 10 (4.78 g, 10.0 mmol) was slurried in 20 mL DCM and treated at room temperature with 20 mL TFA. After 30 minutes the reaction mixture was evaporated under reduced pressure, then dissolved in 20 mL chloroform and re-evaporated two times. The residue was re-dissolved in 20 mL chloroform and added dropwise to 200 mL Et_2O. The resulting yellow solid was collected by suction filtration, washed with 3 x 20 mL Et_2O, and dried to afford 4.8 g, 9.7 mmol, 97% of the title compound. ^1H NMR (DMSO-d_6) δ10.8 (s, 1H), 8.74 (br s, 2H), 7.71 (d, 1H), 7.56 (s, 1H), 7.07 (t, 1H), 6.96 (t, 1H), 6.85 (d, 1H), 3.62 (br s, 4H), 3.02 (br s, 4H), 2.62 (t, 2H), 2.5 (t, 2H), 2.28 (s, 3H), 2.25 (s, 3H). IS-MS, calcd. *m/z* for $C_{22}H_{26}N_4O_2$ $[M+H^+]^+$: 379.2; obsd. 379.0.

Example 1a (Alternative preparation)
3-[3,5-dimethyl-4-(3-oxo-3-piperazin-1-yl-propyl)-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one, trifluoroacetate

Intermediate 9 (0.31 g, 1.0 mmol) was dissolved in 3 mL anhydrous DMF and treated with HOAT (0.14 g, 1.0 mmol) followed by *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU, 0.38 g, 1.0 mmol). After stirring 10 minutes at RT, the reaction mixture was added to a solution of piperazine (0.17 g, 2.0 mmol) and stirred for two days. The reaction mixture was then fractionated by preparative reversed-phase HPLC. The appropriate fractions were combined and lyophilized to afford 0.16 g, 0.34 mmol, 34% of the title compound.

Example 2
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-(2-methylaminoethyl)-propionamide, trifluoroacetate

Intermediate 10 (0.062 g, 0.20 mmol) was dissolved in 0.67 mL anhydrous DMF and treated with HOAT (0.030 g, 0.22 mmol) and HATU (0.084 g, 0.22 mmol). After stirring at RT for 15 minutes, the activated acid was added to a vial containing a solution of *N,N'*-dimethylethylenediamine (0.043 mL, 0.40 mmol) in 0.50 mL anhydrous DMF. The reaction mixture was agitated overnight on an orbiting shaker, then diluted with 0.50 mL of 30% aqueous TFA, filtered, and fractionated by preparative reversed-phase HPLC. The appropriate fractions were combined and lyophilized to afford 0.005 g, 0.010 mmol,

5% of the title compound. IS-MS, calcd. *m/z* for $C_{22}H_{28}N_4O_2$ $[M+H^+]^+$: 381.2; obsd. 381.0.

Examples 3 to 24

In a manner similar to that described in Example 1a and 2, coupling of Intermediate 10 with other amines afforded the compounds of Examples 3 to 24.

Example 3
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-(3-methylaminopropyl)-propionamide, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{23}H_{30}N_4O_2$ $[M+H^+]^+$: 395.2; obsd. 395.0.

Example 4
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-(4-methylaminobutyl)-propionamide, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{24}H_{32}N_4O_2$ $[M+H^+]^+$: 409.3; obsd. 409.0.

Example 5
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-(5-methylaminopentyl)-propionamide, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{25}H_{34}N_4O_2$ $[M+H^+]^+$: 423.3; obsd. 423.2.

Example 6
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-(6-methylaminohexyl)-propionamide, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{26}H_{36}N_4O_2$ $[M+H^+]^+$: 437.3; obsd. 437.2.

Example 7
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-(7-methylaminoheptyl)-propionamide, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{27}H_{38}N_4O_2$ $[M+H^+]^+$: 451.3; obsd. 451.2.

Example 8
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-(8-methylaminooctyl)-propionamide, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{28}H_{40}N_4O_2$ $[M+H^+]^+$: 465.3; obsd. 465.2.

Example 9
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-(9-methylaminononyl)-propionamide, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{29}H_{42}N_4O_2$ $[M+H^+]^+$: 479.3; obsd. 479.2.

Example 10
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-(10-methylaminodecyl)-propionamide, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{30}H_{44}N_4O_2$ $[M+H^+]^+$: 493.4; obsd. 492.8.

Example 11
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-(12-methylaminododecyl)-propionamide, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{32}H_{48}N_4O_2$ $[M+H^+]^+$: 521.4; obsd. 520.8.

Example 12
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-(4-methylaminobut-2-enyl)-propionamide, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{24}H_{30}N_4O_2$ $[M+H^+]^+$: 407.2; obsd. 407.0.

Example 13
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-(8-methylamino-3,6-dioxaoctyl)-propionamide, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{26}H_{36}N_4O_4$ $[M+H^+]^+$: 469.3; obsd. 469.0.

Example 14
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-(11-methylamino-3,6,9-trioxaundecyl)propionamide, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{28}H_{40}N_4O_5$ $[M+H^+]^+$: 513.3; obsd. 512.8.

Example 15
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-(3-methylaminomethylphenyl-methyl)propionamide, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{28}H_{32}N_4O_2$ $[M+H^+]^+$: 457.3; obsd. 457.0.

Example 16
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-(4-methylaminomethylphenyl-methyl)propionamide, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{28}H_{32}N_4O_2$ $[M+H^+]^+$: 457.3; obsd. 457.2.

Example 17
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-(3-methylaminomethylcyclohexyl-methyl)propionamide, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{28}H_{38}N_4O_2$ $[M+H^+]^+$: 463.3; obsd. 463.0.

Example 18
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-[2'-methylaminomethylbiphen-2-ylmethyl]propionamide, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{34}H_{36}N_4O_2$ $[M+H^+]^+$: 533.3; obsd. 533.2.

Example 19
3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrol-3-yl]-N-methyl-N-(3-[4-(3-methylaminopropyl)-piperazin-1-yl]propyl)propionamide, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{30}H_{44}N_6O_2$ $[M+H^+]^+$: 521.4; obsd. 521.2.

Example 20
3-[3,5-dimethyl-4-(3-oxo-3-piperidin-1-ylpropyl)-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{23}H_{27}N_3O_2$ $[M+H^+]^+$: 378.2; obsd. 378.0.

Example 21
3-[3,5-dimethyl-4-[3-oxo-3-(piperidin-4-ylpropyl)piperidin-1-ylpropyl]-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{31}H_{42}N_4O_2$ $[M+H^+]^+$: 503.3; obsd. 503.2.

Example 22
3-[3,5-dimethyl-4-[3-oxo-3-(4-ethyl)piperazin-1-ylpropyl]-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{24}H_{30}N_4O_2$ $[M+H^+]^+$: 407.2; obsd. 407.0.

Example 23
3-[3,5-dimethyl-4-(3-oxo-3-homopiperazin-1-ylpropyl)-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{23}H_{28}N_4O_2$ $[M+H^+]^+$: 393.2; obsd. 393.0.

Example 24
3-[3,5-dimethyl-4-[3-oxo-3-(1,4,10-trioxa-7,13-diazacyclopentadecan-1-yl)-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one, trifluoroacetate

IS-MS, calcd. *m/z* for $C_{28}H_{38}N_4O_5$ $[M+H^+]^+$: 511.3; obsd. 511.0.

Following the methods of Examples 1a and 2, the following compounds are also prepared:

5-Bromo-3-[3,5-dimethyl-4-[3-oxo-3-(4-prop-2-yl)piperazin-1-ylpropyl]-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one, trifluoroacetate and

5-bromo-3-[3,5-dimethyl-4-(3-oxo-3-[4-prop-2-yl]homopiperazin-1-ylpropyl)-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one, trifluoroacetate.

Biological Assays

The ability of test compounds to inhibit receptor tyrosine kinases is demonstrated in the following assays.

Abbreviations

HEPES	4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid
EDTA	Ethylenediaminetetraacetic acid
PDGF	Platelet Derived Growth Factor
PDGFR	Platelet Derived Growth Factor Receptor
VEGF	Vascular Endothelial Growth Factor
VEGFR	Vascular Endothelial Growth Factor Receptor
HEK cells	Human Embryonic Kidney cells
Flt-3	fms-related tyrosine kinase 3
BSA	Bovine Serum Albumin
AML	Acute Myeloid Leukemia
ITD	Internal Tandem Duplication
MTT	3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide.
HUVEC	Human umbilical vein epithelial cells

Ca^{2+} Release Assay (FLIPR Assay)

The binding of growth factors to their respective receptors leads to the autophosphorylation of the receptor. This is the first step in a signaling cascade that results in the release of Ca^{2+} from internal stores and influx of extracellular Ca^{2+}.

The rise in intracelluar Ca^{2+} is quantitated by loading cells with a fluorescent dye prior to stimulation with the growth factor and by subsequently assessing the fluorescent signal in a Fluorometric Image Plate Reader (FLIPR). A kinase inhibitor that can penetrate the cell membrane inhibits receptor autophosphorylation, hence, the Ca^{2+} release is reduced or ablated.

In order to determine the IC_{50} of test compounds against the VEGFR in the FLIPR assay, HUVECs (Walkersville, MD) were used. For the determination of IC_{50}s against the PDGFR, a HEK cell line were used that expresses the human PDGFR. 40-50000 cells per well were plated in a 96 well plate. The cells were incubated for 3-4 hours to adhere to the

plate. Subsequently, the cells were washed twice in FLIPR buffer (1XHBS, 2mM CaCl, 10mM HEPES, pH7.4, 2.5 mM Probenecid, 0.1% BSA). After the second wash 50 µL of the Ca^{2+} sensitive dye FLUO-3 (FLUO-3 (AM) TEF Labs, 50µg in 10mL FLIPR buffer) was added to the remaining 50µL buffer in each well. After loading the cells for one hour, the cells were washed twice and the test compounds were added in 50 µL as a 2X solution to the 50 µL of buffer in each well. The cells were incubated with the compounds for 30 minutes and then VEGF (40 ng/mL, BioSource International) for the VEGFR or PDGF (40 ng/mL, BioSource International) for the PDGFR was added. The change in fluorescence intensity was measured in a Fluorometric Image Plate Reader (FLIPR) (Molecular Devices).

Proliferation and Viability Assay (MTT)

The inhibition of mutant Flt-3 ITD is expected to affect the proliferation and viability of AML cells with this mutation. In order to assess the activity of test compounds an MTT proliferation and viability assay (Roche Molecular Biochemicals, Indianapolis, IN) was performed with an AML cell line called MV4-11. MV4-11 cells express Flt-3 ITD. 50,000 cells per well in 100 µL of media were plated in a 96 well plate and were incubated with increasing concentrations of compound for 48 hours. After this incubation period, 10 µL of the MTT labeling reaction was added for 4 hours. The MTT labeling reagent was metabolized by viable cells to formazan, an insoluble blue salt. To solubilize the formazan salt, 100 µL of solubilization solution was added. The plate was incubated at 37 degrees Celsius for 24 hours and, then, the optical density of the wells was determined spectrophotometrically at 550 nm. The optical density of the solutions in the wells reflects the effect the compound has on the viability of the cells.

Immunoprecipitation/ Western (IP/Western)

The binding of growth factors to receptor tyrosine kinases like Flt-3 or PDGFR leads to the autophosphorylation of the receptors. The inhibition of autophosphorylation is the objective pursued with kinase inhibitors. By performing an IP/Western experiment the level of receptor autophosphorylation can be assessed directly.

$5x10^6$ cells (HEK PDGFR cell line for PDGFR, HEK c-Kit for c-Kit, and THP-1, HL-60 or MV4-11 for Flt-3) were incubated for 30 minutes in 2.5 mL of culture media with a defined concentration of test compound. In order to stimulate receptor

autophosphorylation, growth factor (PDGF, SCF, or Flt-3 ligand respectively, 50 ng/mL, Biosource International, Camarillo, CA) was added for 5 minutes. The cells were then centrifuged and lysed in 500 µL lysis buffer (50mM Tris pH 7.4, 1% NP-40, 150 mM NaCl, 1 mM EDTA, 1 mM Na3V04). The lysate is centrifuged and 10 µL of antibody against the respective receptor was added to the supernatant (anti-PDGFR (P20), anti-c-Kit (C-19), and anti-Flt-3 (S18), Santa Cruz Biotechnology, Inc.). The immunocomplexes were isolated with Protein G beads (Sigma, St. Louis, MO) and PAGE was performed. A Western Blot was done using an antibody against phospho-tyrosine residues (4G10, Upstate Biotechnology, Lake Placid, NY). The intensity of the phospho-tyrosine signal corresponding to various drug concentrations provides a way to determine the IC_{50} of the test compound for the inhibition of autophosphorylation.

In general, the compounds exemplified herein have been found to exhibit an IC_{50} of less than 10 µM in one or more of the above assays.

Pharmacokinetics

To evaluate pharmacokinetics, male Sprague Dawley rats (CD strain, Charles River Laboratories, Wilmington, MA) were dosed with test compounds via intravenous (IV) administration, at 1 mg/kg concentration, and oral (PO) administration, at 10 mg/kg concentration. Blood samples were collected from animals pre-dose, and at 2, 5, 15, and 30 minutes, and at 1, 2, 4, 6, 8, and 24 hours post-dose. Plasma concentrations were determined by liquid chromatography-mass spectrometry (LC-MS) (MDS SCIEX API 4000, Applied Biosystems, Foster City, CA). Standard pharmacokinetic parameters were assessed by non-compartmental methods using the WinNonlin Version 3.2 software package (Pharsight, Mountain View, CA). Oral bioavailability was determined as the ratio of the area under the curve (AUC) in the graph of plasma concentration versus time for PO administration to the corresponding quantity for IV administration. The oral bioavailability in rats, for example, of 3-[3,5-dimethyl-4-(3-oxo-3-piperazin-1-ylpropyl)-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one, determined by this method, was 50.6 %.

Comparative Assay Results

Table 1 lists assay results for two compounds of the invention, the compound of Example 1, 3-[3,5-dimethyl-4-(3-oxo-3-piperazin-1-yl-propyl)-1H-pyrrol-2-ylmethylene]-

1,3-dihydroindol-2-one, trifluoroacetate, and the compound of Example 22, 3-[3,5-dimethyl-4-[3-oxo-3-(4-ethyl)piperazin-1-ylpropyl]-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one, trifluoroacetate. For comparison, Table 1 also lists assay results for two prior art compounds, 3-[2,4-dimethyl-5-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrol-3-yl]propionic acid, identified as SU6668, and 3-(2,3-dimethylpyrrol-5-yl)methylene]-2-indolinone, identified as SU5416. The preparation of the former prior art compound is described above as intermediate 9. The latter compound was prepared as described in Sun et al., *J. Med. Chem.* 1998, Vol. 41, No. 14, pp 2588-2603. The ability of test compounds to inhibit mutant Flt-3 ITD was tested in the cytotoxicity assay. Inhibition of the VEGFR and PDGFR kinases was tested in the Ca^{2+} FLIPR assay, except as indicated. As shown below the compounds of Examples 1 and 22 showed sub-micromolar activity in the Flt-3, VEGFR, and PDGFR assays.

Table 1

	Flt-3 EC_{50} (μM)	VEGFR IC_{50} (μM)	PDGFR IC_{50} (μM)
This Invention			
Example 1	0.24	0.02	0.03#
Example 22	0.22	0.03	0.09
Comparison Compounds			
SU6668	No activity*	No activity**	No activity**#
SU5416	~1-10	0.05	0.06


* Highest concentration tested 10 μm

** Highest concentration tested 1 μm

Immunoprecipitation/Western (IP) assay

CLAIMS

1. A compound of formula (I)



(I)

wherein:

(i) R^1 represents a hydrogen atom or a (1-4C)alkyl group; and R^2 represents a group of formula $-A^1-NR^5R^6$ in which each of R^5 and R^6 independently represents a hydrogen atom or a (1-4C)alkyl group and A^1 represents $(CH_2)_m$, $(CH_2)_n-A^2-(CH_2)_p$ or $(CH_2CH_2O)_qCH_2CH_2$ in which *m* is an integer of from 2 to 10, each of *n* and *p* is an integer of from 1 to 6, A^2 is CH=CH, phenylene, biphenylene, cyclohexylene or piperazinylene and *q* is 1, 2 or 3;

(ii) R^1 and R^2 together represent $-A^3-NR^7-A^4-$ in which each of A^3 and A^4 independently represents $(CH_2)_r$ or $(CH_2CH_2O)_sCH_2CH_2$ in which *r* is an integer of from 2 to 6, *s* is 1, 2 or 3, and R^7 represents a hydrogen atom or a (1-4C)alkyl group;

(iii) R^1 and R^2 together with the nitrogen atom to which they are attached represent a piperidinyl group, which piperidinyl group bears a substituent of formula $-A^5-R^8$ at the 4 position, in which A^5 represents (1-4C)alkylene and R^8 represents piperidin-4-yl; or

(iv) R^1 and R^2 together with the nitrogen atom to which they are attached represent a pyrrolidinyl, piperidinyl or morpholino group; and

R^3 and R^4 each independently represents a hydrogen atom, a halogen atom, a (1-4C)alkyl group, a (1-4C)alkoxy group, a phenyl group which is unsubstituted or substituted by one or two substituents selected independently from a halogen atom, a (1-4C)alkyl group and a (1-4C)alkoxy group, a group of formula $R^8S(O)_2NR^9-$, a group of formula $R^{10}N(R^{11})S(O)_2-$, a group of formula $R^{12}C(O)N(R^{13})-$ or a group of formula $R^{14}N(R^{15})C(O)-$ in which each of R^8, R^{10}, R^{12} and R^{14} independently represents a

(1-4C)alkyl group or a phenyl group which is unsubstituted or substituted by one or two substituents selected independently from a halogen atom, a (1-4C)alkyl group and a (1-4C)alkoxy group, and each of R^9, R^{11}, R^{13} and R^{15} independently represents a hydrogen atom or a (1-4C)alkyl group;

or a pharmaceutically-acceptable salt thereof.

2. A compound as claimed in Claim 1, in which

(i) R^1 represents a methyl group; and R^2 represents a group of formula $-A^1-NR^5R^6$ in which R^5 represents a hydrogen atom, R^6 represents a methyl group and A^1 represents $(CH_2)_m$, in which *m* is 2, 3, 4, 5, 6, 7, 8, 9 or 10; $(CH_2)_n-A^2-(CH_2)_p$ in which *n* and *p* are each 1 and A^2 is CH=CH, phenyl-1,3-ene, phenyl-1,4-ene, biphenyl-2,2'-ene or cyclohex-1,3-ylene; $(CH_2)_n-A^2-(CH_2)_p$ in which *n* and *p* are each 2 and A^2 is piperazin-1,4-ylene; or $(CH_2CH_2O)_qCH_2CH_2$ in which *q* is 2 or 3;

(ii) R^1 and R^2 together represent $-(CH_2)_2-NH-(CH_2)_2-$, $-(CH_2)_2-N(CH_3)-(CH_2)_2-$, $-(CH_2)_2-N(CH_2CH_3)-(CH_2)_2-$, $-(CH_2)_2-NH-(CH_2)_3-$, $-(CH_2CH_2O)_2CH_2CH_2-NH-(CH_2CH_2O)CH_2CH_2-$; or

(iii) R^1 and R^2 together with the nitrogen atom to which they are attached represent a piperidinyl group, which piperidinyl group bears a substituent of formula

-A^5-R^8 at the 4 position, in which A^5 represents propylene and R^8 represents piperidin-4-yl.

3. A compound as claimed in any one of Claim 1 or 2, in which R^3 and R^4 each independently represents a hydrogen atom, a bromine atom, $CH_3C(O)NH$, or $C_6H_5C(O)NH$.

4. A compound as claimed in Claim 1 in which

(i) R^1 represents a methyl group and R^2 represent a group of formula $-A^1$-$NHCH_3$ in which A^1 represents $(CH_2)_m$, $CH_2CH=CHCH_2$, CH_2-phenylene-CH_2, or CH_2-cyclohexylene-CH_2, in which *m* is an integer of from 2 to 8; or

(ii) R^1 and R^2 together represent $-(CH_2)_2-NH-(CH_2)_2-$, $-(CH_2)_2-N(CH_3)-(CH_2)_2-$, $-(CH_2)_2-N(CH_2CH_3)-(CH_2)_2-$ or $-(CH_2)_2-NH-(CH_2)_3-$; and

R^3 and R^4 are each independently hydrogen.

5. A compound as claimed in Claim 4 in which:

(i) R^1 represents a methyl group and R^2 represent a group of formula $-A^1-NHCH_3$ in which A^1 represents $(CH_2)_m$, $CH_2CH=CHCH_2$, or CH_2-(1,4-phenylene)-CH_2, in which *m* is 2 or 3; or

(ii) R^1 and R^2 together represent $-(CH_2)_2-NH-(CH_2)_2-$, $-(CH_2)_2-N(CH_3)-(CH_2)_2-$, $-(CH_2)_2-N(CH_2CH_3)-(CH_2)_2-$ or $-(CH_2)_2-NH-(CH_2)_3-$.

6. A compound as claimed in Claim 1, which is selected from:


3-[3,5-dimethyl-4-(3-oxo-3-piperazin-1-ylpropyl)-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one;

3-[3,5-dimethyl-4-[3-oxo-3-(4-ethyl)piperazin-1-ylpropyl]-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one;

3-[3,5-dimethyl-4-(3-oxo-3-homopiperazin-1-ylpropyl)-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one;

and pharmaceutically-acceptable salts thereof.

7. A compound as claimed in claim 1, which is a compound of formula (Ia):



(Ia)

wherein R is hydrogen, methyl, or ethyl;

or a pharmaceutically-acceptable salt thereof.

8. A compound as claimed in Claim 7, which is selected from:

3-[3,5-dimethyl-4-(3-oxo-3-piperazin-1-ylpropyl)-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one;


3-[3,5-dimethyl-4-[3-oxo-3-(4-ethyl)piperazin-1-ylpropyl]-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one;

and pharmaceutically-acceptable salts thereof.

9. A compound as claimed in Claim 8, which is 3-[3,5-dimethyl-4-(3-oxo-3-piperazin-1-ylpropyl)-1H-pyrrol-2-ylmethylene]-1,3-dihydroindol-2-one, or a pharmaceutically-acceptable salt thereof.

10. A process for the preparation of a compound as claimed in any one of Claims 1 to 9 which comprises:

(a) reacting a compound of formula (II)



(II)


or a reactive derivative thereof, with a compound of formula (III)

HNR^1R^2

(III)

or a salt thereof, in which R^1, R^2, R^3 and R^4 are as defined in Claim 1, or

(b) for a compound of formula (I) in which R^5 or R^7 represents a hydrogen atom, deprotecting a compound of formula (IV)



(IV)

in which R^{1a} and R^{2a} are as defined in Claim 1 for R^1 and R^2, except in that R^5 or R^7 is replaced with a group R^{5a} or R^{7a} respectively, in which R^{5a} and R^{7a} each represents an amine protecting group, and R^3 and R^4 are as defined Claim 1;

followed, if a pharmaceutically-acceptable salt is required, by forming a pharmaceutically-acceptable salt.

11. A pharmaceutical composition, which comprises a therapeutically-effective amount of a compound as claimed in any one of Claims 1-9, together with a pharmaceutically-acceptable diluent or carrier.

12. A compound as claimed in any one of Claims 1 to 9, for use in therapy.

13. Use of a compound as claimed in any one of Claims 1 to 9, for the manufacture of a medicament for the treatment of a condition responsive to a tyrosine kinase inhibitor.