

Description

The present invention relates to novel phenyl carbamates which are useful as pharmaceutical compositions. The invention further relates to pharmaceutical compositions having anticholinesterase activity.

Acetylcholine is a major neurotransmitter which is found in all parts of the body. Any reduction in its activity, either as a result of neuronal damage, degeneration etc. or as induced by drugs or toxins, causes marked changes in the function of the organisms. Acetylcholine itself has an extremely short half life, since it is rapidly hydrolysed at its site of action and in plasma by specific cholinesterase enzymes. Drugs that inhibit acetylcholinesterase, markedly increase and prolong the action of acetylcholine, thereby enhancing cholinergic transmission. Three such agents are used clinically, i.e., physostigmine, a naturally occurring alkaloid, and two synthetic analogues, neostigmine and pyridostigmine. The latter two agents are strongly ionised at physiological pH and therefore are only poorly absorbed from the gastro-intestinal tract, and do not penetrate the central nervous system to any significant extent. Physostigmine is absorbed after oral administration and readily enters the brain. As a therapeutic agent it has several disadvantages. It is chemically unstable and must be prepared in solution with an antioxidant, and protected from light. It has a relatively short half-life (20—40 mins) thereby necessitating frequent administration. The latter is of particular importance when the drug is to be administered chronically. It has a low therapeutic ratio, a value of 3—5 being reported in the majority of studies in laboratory animals, and a small therapeutic window, i.e. small range of dose in which it can be given without the accompaniment of side effects. Although physostigmine is absorbed from the gastro-intestinal tract, this is reported to be irregular and unpredictable, and therefore it is usually preferred to administer the drug parenterally. This is a serious drawback if it is to be used chronically on an outpatient basis.

There are a number of clinical and pathological conditions which are associated with cholinergic under-activity which can be improved by the administration of an anticholinesterase agent. These include reduction in cholinergic transmission induced by a variety of exogenous substances acting in the peripheral, or central nervous system. Peripherally acting agents are gallamine, d-tubocurarine and pancuronium, which are used as muscle relaxants. Their action can readily be overcome by an anticholinesterase drug. Drugs which interfere with central cholinergic transmission are numerous, anticholinergic, atropine-like drugs including antiparkinson drugs, tricyclic antidepressants, neuroleptics, opiate analgesics, benzodiazepines and some types of general anaesthetics. So far the only agent that has proved to be of any value in reversing the effects of the latter group of drugs is physostigmine. In all reported cases of drug overdose or lack of recovery when the agent was used peri-operatively, physostigmine is usually administered parenterally, and administration is repeated every 20—30 minutes as required.

Chronic treatment with neuroleptics often results in tardive dyskinesias. The widespread use of agents having anticholinesterase activity for the treatment of schizophrenia makes this side effect an ever increasing possibility. Physostigmine injected intravenously produces a significant but short lived improvement in a proportion of patients.

A number of pathological and degenerative diseases has also been shown to be associated with a reduction or loss of cholinergic transmission. This includes myasthenia gravis and Eaton Lambert syndrome in which there is an interference with neuromuscular transmission.

A selective loss of choline acetyltransferase (the enzyme that synthesises acetylcholine) has been found in specific brain regions of patients with pre-senile dementia of the Alzheimer type. These include the frontal and temporal cortex, hippocampus, amygdala, caudate nucleus, substantia innominata. Degeneration of cholinergic neurons in some of these areas appears to be associated with the aphasia, apraxia, agnosia and loss of short term memory that occurs in Alzheimer's disease. A similar type of dementia is also found in patients with Down's syndrome that survive to the age of 40 years and show similar cholinergic deficits. There is also a loss of cholinergic transmission in the caudate nucleus and putamen of patients with Huntington's chorea. Physostigmine injections have also been of some benefit in this condition. Treatment with a centrally acting anticholinesterase should also prove to be beneficial in Friedrich's ataxia.

There are two major classes of potent inhibitors of the enzyme cholinesterase. The first group was modelled primarily on the natural alkaloids physostigmine (a carbamate) and an inhibitor of cholinesterase, and d-tubocurarine, an antagonist of acetylcholine. The second group consists of various organophosphorus compounds, such as diisopropylfluorophosphonate, paraxon etc. The vast majority of the compounds of both these series were designed primarily as insecticides. In the first group of carbamate derivatives, almost all of the potent insecticides are monomethyl carbamates lacking a charged nitrogen function (C.A. 71(1969) 111823m). This enables the molecule to penetrate rapidly the insect cuticle and fatty nerve sheath. The dimethyl derivatives are slightly less potent but are particularly toxic to houseflies and aphids. The monomethyl derivatives tend to be unstable in solution and hydrolyse readily at physiological pH. This greatly limits their biological action in mammals and makes them less suitable as pharmaceutical or therapeutic agents. The dimethylcarbamate of o-hydroxybenzyl-diethylamine is known from US-A-1 905 990 as having a physostigmine-like action. The organo-phosphorus group of compound causes irreversible inhibition of cholinesterase and other serine containing enzymes, which, together with their

high relative toxicity, virtually precludes their use in pharmaceutical preparations. The only exception is echothiopate, a quaternary ammonium organophosphorus compound, employed in eye drops for the treatment of glaucoma.

The synthetic anticholinesterase agents currently employed as pharmaceuticals all contain a charged nitrogen function and can be broadly classified in to 3 groups.

1) Reversible inhibitors which contain a charged nitrogen function attached to an aromatic ring, e.g. edrophonium.


2) Dimethyl carbamates with an aromatic or heterocyclic ring containing a charged nitrogen, neostigmine, pyridostigmine.

3) Bisquaternary structures, e.g. Demacarium, Ambenonium. These agents tend to be more selective inhibitors of acetylcholinesterase than butyrylcholinesterase compared with mono-quaternary molecules.

The pharmaceutical application of the quaternary anticholinesterase agents is limited because of their poor penetration through cell membranes. They are therefore used for actions outside the central nervous system, and are usually given parenterally, since they are not reliably absorbed from the gastrointestinal tract. Edrophonium, neostigmine and pyridostigmine and the bisquaternary analogues are used in anaesthetic practice for the reversal of the action of muscle relaxants. They are also used for the treatment of myasthenia gravis, and paralytic ileus.

Physostigmine is the only potent anti-cholinesterase agent which has been used clinically to treat conditions in which an elevation of brain acetylcholine activity is desired. These include, Alzheimer's disease, tardive dyskinesia, Down's syndrome and Huntingdon's chorea. Physostigmine is also used to reverse the effects of overdose of anticholinergic agents, anti-Parkinson drugs, benzodiazepines and opiate analgesics.

Physostigmine is a natural alkaloid extracted from calabar beans and the seeds of the vine *Physostigma venenosum* and has the formula



There is a need to provide new carbamate derivatives which show greater chemical stability than physostigmine.

Furthermore there is a need to provide new compounds which inhibit acetylcholinesterase in the brain for periods exceeding 3 hours but not more than 12 hours after a single administration.

There is also a need to provide new compounds which will be completely and reliably absorbed after oral administration.

There is also a need to provide new compounds which will be relatively less toxic than physostigmine. This means that the therapeutic ratio, defined as

dose to produce therapeutic effect
 dose to produce mortality in 50% of animals

should be significantly higher than those of physostigmine and that the incidence and severity of side effects should be less than those of physostigmine at therapeutic doses.

There is also a need to provide new compounds which can be given orally or parenterally to treat chronic conditions in which it is desired to raise cholinergic activity in the central nervous system. These include, Alzheimer's disease, Down's syndrome, Huntington's chorea, Friedreich's ataxia.

There is also a need to provide compounds that can be given parenterally at the end of operations, and anaesthetic procedures, to restore wakefulness, respiration and cardiovascular parameters to normal, after the use of anticholinesterase, opiates, benzodiazepines, neuroleptics and general anaesthetics, thereby shortening the stay of patients in the recovery room.


There is also a need to provide compounds that can be given together with narcotic analgesics to patients suffering from severe pain, e.g. traumatic, post-operative, or due to carcinomatosis etc. in order to reduce the side effects (respiratory depression, somnolence, constipation and urinary retention) commonly encountered with narcotics, without impairing their analgesic potency.

There is also a need to provide compounds that can be given to patients receiving antipsychotic drugs, which have developed tardive dyskinesias, in order to diminish or abolish the latter syndrome, without

exacerbating the psychosis.

According to the present invention it has now been surprisingly found that certain novel phenyl carbamates also inhibit acetylcholinesterase in the mammalian brain after administration to provide systemic activity, e.g. oral or parenteral administration.

Thus according to the present invention there is now provided a pharmaceutical composition adapted to produce anticholinesterase activity in the central nervous system of mammals comprising a compound of the general formula I



wherein

R₁ is hydrogen, lower alkyl, cyclohexyl, allyl or benzyl,

R₂ is hydrogen, methyl, ethyl or propyl, or

R₁ and R₂ together with the nitrogen to which they are attached form a morpholino or piperidino radical,

R₃ is hydrogen or lower alkyl,

R₄ and R₅ are the same or different and each is a lower alkyl, and the dialkylaminoalkyl group is in the meta, ortho or para position, provided that for compounds wherein R₄ and R₅ are both methyl and having the dialkylaminoalkyl group in the meta position, when R₂ is methyl and R₃ is hydrogen, R₁ is neither hydrogen nor methyl, and when R₂ and R₃ are methyl, R₁ is not hydrogen, and for compounds wherein R₄ and R₅ are both methyl and having the dialkylaminoalkyl group in the ortho or para position when R₁ and R₂ are both hydrogen R₃ is not methyl, or a pharmacologically acceptable salt thereof and a physiologically acceptable carrier therefor. Hereinafter these compounds are called compounds of the invention.

Especially preferred are pharmaceutical compositions having anti-cholinesterase activity in the central nervous system of mammals, wherein the dialkylaminoalkyl group is in the meta position, and R₄ and R₅ are both methyl.


Certain compounds falling within the disclaimer of the above formula have previously been described i.e. the m, disubstituted compound in which R₁ and R₃ = H and R₂, R₄ and R₅ = methyl which is known as Miotine(R) was claimed to be an insecticide and a myopic agent for use in eye drops. The m disubstituted compound in which R₁ and R₂ are methyl, R₃ is H and R₄ and R₅ are methyl has been described as an insecticide. The p and o disubstituted derivatives in which R₁ and R₃ = H and R₂, R₄ and R₅ = CH₃ have been shown to inhibit a preparation of liver cholinesterase. The m disubstituted derivative in which R₁ = H and R₂, R₃, R₄ and R₅ = CH₃ has also been shown to inhibit liver cholinesterase.

The remaining compounds are believed to be novel and thus the present invention also provides novel phenyl carbamate derivatives of the general formula I as defined above.

Preferred compounds of the above formula are N-ethyl-3-[1-(dimethylamino)ethyl]phenyl carbamate, N-propyl-3-[1-(dimethylamino)ethyl]phenyl carbamate, N-allyl-3-[1-(dimethylamino)ethyl]phenyl carbamate, N-ethyl, N-methyl-3-[1-(dimethylamino)ethyl]phenyl carbamate, N,N-diethyl-3-[1-(dimethylamino)ethyl]phenyl carbamate, N-butyl-3-[1-(dimethylamino)ethyl]phenyl carbamate, N-methyl, N-propyl-3-[1-(dimethylamino)ethyl]phenyl carbamate and N-ethyl, N-methyl-3-[1-(dimethylamino)isopropyl]phenyl carbamate.

As indicated, the invention also includes the pharmacologically acceptable salts of these compounds such as the acetate, salicylate, fumarate, phosphate, sulphate, maleate, succinate, citrate, tartrate, propionate and butyrate salts thereof.

The compounds of formula I can be prepared by amidating a compound of formula II




II


wherein R_3 , R_4 and R_5 are as defined above.

The process can be effected in conventional manner, e.g. by reacting the compound of formula II with an appropriate isocyanate if a compound wherein R_1 is hydrogen is desired, or with an appropriate carbamoyl halogenide, e.g. as described below in processes A and B.

PROCESS A:



PROCESS B:



Process A

A stirred suspension of α -m-Hydroxyphenylethyldimethylamine or α -m-hydroxyphenylisopropyl-dimethylamine in benzene (0.2—0.3 g/ml) is treated with 2.5—3 fold molar excess of the isocyanate. After stirring for 15—24 hours at ambient temperature the reaction mixture is connected to a rotovaporator (20 mm Hg). The residue obtained is dissolved in dry ether (25) ml and the solution, which is ice cooled, is saturated with dry HCl (g). The formed precipitate (the anticipated carbamate) is filtered off, washed with dry ether (25) ml and dried to constant weight in a dessicator over KOH pellets under high vacuum (0.1 mm Hg).

Process B

A solution of α -m-Hydroxyphenylethyldimethylamine or α -m-hydroxyphenylisopropyl-dimethylamine in dry acetonitrile (0.1—0.5M) is reacted with 50—70% molar excess of the corresponding carbamoyl chloride in the presence of 200% molar excess of NaH dispersion (50—80% in mineral oil). The reaction mixture is left to stir at ambient temperature for 15—24 hours. Removal of the acetonitrile under reduced pressure (20 mm Hg) is followed by the addition of water (10—25 ml). The pH of the aqueous solution is adjusted to pH = 11 by the addition of the appropriate amount of NaOH 0.1N followed by extraction with ether (3 x 25 ml). The combined organic phases are washed with brine (25 ml) dried over $MgSO_4$ anhydride which is then filtered off. The ice cooled ethereal filtrate is saturated with a stream of HCl (g) resulting in the formation of a heavy precipitate (the anticipated carbamate) which is collected by filtration, washed with dry ether (20 ml) and dried to constant weight in a desiccator under high vacuum (0.1 mm Hg) over KOH pellets.

The compounds of the invention e.g. in free form or salt form can be utilized by formulating one or more

of them in compositions such as tablets, capsules or elixirs for oral administration or in sterile solutions or suspensions for parenteral administration. A compound or mixture of compounds of formula (I) or physiologically acceptable salt(s) thereof is compounded with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, flavor, etc., in a unit dosage form as called for by accepted pharmaceutical practice. The amount of active substance in these compositions or preparations is such that a suitable dosage is obtained.

Illustrative of the adjuvants which may be incorporated in tablets, capsules and the like are the following: a binder such as gum tragacanth, acacia, corn starch or gelatin; an excipient such a dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid an the like; a lubricant such as mangnesium stearate; a sweetening agent such as sucrose, lactose or saccharin; a flavoring agent such as peppermint, oil of wintergreen or cherry. When the dosage unit form is a capsule, it may contain in addition to materials of the above type a liquid carrier such as a fatty oil. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets may be coated with shellac, sugar or both. A syrup or elixir may contain the active compound, sucrose as a sweetening agent, methyl and propyl parabens as preservatives, a dye and a flavoring such as cherry or orange flavour.

Sterile compositions for injection can be formulated according to conventional pharmaceutical practice by dissolving or suspending the active substance in a vehicle such as water for injection.

Buffers, preservatives, antioxidants and the like can be incorporated as required.


Preferred antioxidants for use with the compounds of the present invention include sodium metabisulphite and ascorbic acid.

While the invention will now be described in connection with certain preferred embodiments in the following examples, it will be understood that it is not intended to limit the invention to these particular embodiments. On the contrary, it is intended to cover all alternatives, modifications and equivalents as may be included within the scope of the invention as defined by the appended claims. Thus, the following examples which include preferred embodiments will serve to illustrate the practice of this invention, it being understood that the particulars described are by way of example and for purposes of illustrative discussion of preferred embodiments of the present invention only and are presented in the cause of providing what is believed to be the most useful and readily understood description of procedures as well as of the principles and conceptual aspects of the invention.

Example 1

0.5 g (3.03 mmole) of α -m-hydroxyphenylethyldimethylamine are dissolved in 15 ml of dry acetonitrile and 0.70 g (5.2 mmole) of diethylcarbamyldichloride are added to the mixture with stirring. This is followed by NaH 150 mg (50%) of dispersion. The reaction mixture is stirred overnight at 25–30°C. Removal of acetonitrile under reduced pressure is followed by addition of water (10 ml) and adjustment of the pH to 11. The product is extracted in ether, which is washed by brine, dried over MgSO_4 and filtered. Upon addition of HCl (g) precipitation occurs immediately, the product is filtered off, washed by dry ether and dried in a desiccator under high vacuum over KOH pellets.

The carbamate is obtained as a white powder 640 mg (80%) mp. 137–138° and identified as N,N-diethyl-3-[1-(dimethylamino)ethyl]phenyl carbamate, having the formula



Example 2

0.75 g (4.55 mmol) of α -m-hydroxyphenylethyldimethylamine are suspended in benzene (3 ml) and 0.898 g of ethylisocyanate are added to the mixture with stirring. After stirring 12 hours at room temperature the solvent is removed under reduced pressure.

The residue obtained was dissolved in dry ether. Introduction of dry HCl gas into the reaction mixture causes a heavy precipitation. The product is filtered off, washed with ether and dried in a desiccator over KOH pellets. The carbamate is obtained as a white powder 800 mg (75%) mp. 177–179°C and identified as N-ethyl-3-[1-(dimethylamino)ethyl]phenyl carbamate, having the formula

Table 1**In vitro activity on solubilized mouse brain enzyme**

Compound (R ₄ =R ₅ =CH ₃)	R ₁	R ₂	R ₃	IC ₅₀ (M)	Time of peak activity (mins)
Physiostigmine (Salicylate)	H	CH ₃	H	1.1x10 ⁻⁸	30
Miotine HCl	H	CH ₃	H	1.3x10 ⁻⁸	30
RA ₆ HCl	H	C ₂ H ₅	H	4.0x10 ⁻⁷	120
RA ₁₅ HCl	H	C ₃ H ₇ n-propyl	H	1.1x10 ⁻⁷	120
RA ₁₄ HCl	H	C ₃ H ₅ allyl	H	4.3x10 ⁻⁷	120
RA ₁₃ HCl	H	C ₃ H ₇ isopropyl	H	1.2x10 ⁻⁵	120
RA ₅ HCl	H	C ₄ H ₉ n-butyl	H	7.6x10 ⁻⁸	120
RA ₁₂	H	cyclohexyl	H	9.3x10 ⁻⁸	120
RA ₁₀ HCl	CH ₃	CH ₃	H	2.7x10 ⁻⁸	120
RA ₇ HCl	CH ₃	C ₂ H ₅	H	1.3x10 ⁻⁶	90
RA ₈ HCl	C ₂ H ₅	C ₂ H ₅	H	3.5x10 ⁻⁵	30
RA ₁₁ HCl	morpholino		H	> 2x10 ⁻⁵	30
RA ₄ HCl	CH ₃	propyl	H	1.7x10 ⁻⁶	60

Melting points of compounds (all in hydrochloride form except for RA₁₂, which is in free base form as it precipitated from reaction mixture before addition of hydrogen chloride), in degrees centigrade: RA₆ 167—170, RA₁₅ 141—143, RA₁₄ 147—152, RA₁₃ 146—148, RA₅ 158—162, RA₁₂ 75—77, RA₁₀ 145, RA₇ 135—136, RA₈ 137—138, RA₁₁ amorphous, RA₄ 148—149. Compound RA₁₁ has an R_f value of 0.59 in a system of 95 parts of ethyl acetate and 5 parts of 33% (w/w) dimethylamine in ethanol.

Table 2

Anticholinesterase activity of compounds in mouse brain
compared to that of physostigmine

Compound	Relative potency to physostigmine after subcut. (s.c.) administration	Relative potency to physostigmine after oral administration	% cholinesterase inhibition 3 hours after s.c. administration
Physo- stigmine	100	100	0
Miotine	100	300	5
RA ₆	11	19	35
RA ₁₅	33	32	37
RA ₁₄	15	22	35
RA ₁₃	2	5	-
RA ₅	36	29	30
RA ₁₂	13	17	37
RA ₁₀	81	92	7
RA ₇	25	57	41
RA ₈	2	5	32
RA ₄	13	29	25

Table 3

Acute toxicity of carbamates in mice

Compound	LD ₅₀ umoles/kg s.c.	Degree of protection afforded by pretreatment with atropine	Therapeutic ratio LD ₅₀ /ED ₅₀ s.c.	LD ₅₀ oral
				LD ₅₀ s.c.
Physostigmine	3.0	3.0	3.3	4.1
Miotine	4.5	2.4	4.9	1.2
RA ₆	96	2.6	11.9	2.1
RA ₁₅	31	4.1	11.1	4.5
RA ₁₄	69	8.0	11.5	4.4
RA ₁₃	65	4.5	1.6	1.1
RA ₅	19	5.8	7.6	5.0
RA ₁₂	42	3.8	5.8	3.6
RA ₁₀	14	5.0	12.7	9.7
RA ₇	46	10.4	12.4	1.2
RA ₈	> 568	-	> 10.0	-
RA ₄	72	4.9	10.0	1.7

*Ratio of LD₅₀ after pretreatment with atropine sulphate 5 mg/kg to LD₅₀ of drug alone.

The data in Tables 1 and 2 demonstrate that somewhat larger quantities are required of all the drugs of the RA series than of physostigmine to inhibit the enzyme acetylcholinesterase. However, a comparison of the data in Table 1 with that in Table 2, shows that compounds RA₆, RA₈, RA₁₅, RA₁₄, RA₁₀, RA₇, and RA₅ are all relatively more active *in vitro* compared to physostigmine than one would expect from *in vitro* data. This greater *in vitro* potency is particularly marked when the drugs are administered orally. This relatively greater *in vitro* activity may be due to:

- greater chemical stability
- a slower metabolic degradation or/and excretion
- a higher lipid solubility, enabling a greater proportion of the drug to gain access to the enzyme in the central nervous system.
- more efficient absorption from gastro-intestinal tract.

For the purposes of their therapeutic application it is of little importance if one needs to give the drug (to human subjects) at a dose of 1—2 mg (physostigmine) or 2—50 mg that may be required of the compounds of the RA series. What is important is the safety of the drugs and the presence and severity of side effects that may occur at therapeutic doses. A commonly-used measure of drug safety is the therapeutic index — or LD₅₀/ED₅₀

Dose to kill 50% of animals

Dose to cause the desired therapeutic effect

It is assumed that the therapeutic effect of these anticholinesterase agents results from an elevation of brain cholinergic activity. This in turn, should be related to the degree of inhibition of acetylcholinesterase. For the purpose of the computation of the denominator of the therapeutic ratio, there is used the dose of drug that inhibits the activity of acetylcholinesterase by 50%. This is based on the observation by Thal et al. (Ann Neurology 13, 491, 1983) that the maximum improvement in short term memory obtained in a series of patients with Alzheimer's disease was achieved with a dose of physostigmine which blocked the acetylcholinesterase in the cerebro-spinal fluid by 50%. The numerator is the dose found to kill 50% of the animals within 4 hours of subcutaneous injection.

The therapeutic ratios of compounds RA⁴, ⁵, ⁶, ⁷, ⁸, ⁹, ¹⁰, ¹⁴ and ¹⁵ are all significantly higher than of physostigmine (see Table 3). This indicates that all these compounds have a wider margin of safety than that of physostigmine. Moreover, these RA compounds do not produce any significant undesirable side effects such as defaecation, lachrymation, fasciculations or tremor at the doses which inhibit the brain enzyme by 50%, while the former 3 side effects are clearly evident when physostigmine is given at the appropriate dose (ED₅₀).

The data in Table 3 show that atropine can afford considerably greater protection against the lethality of the derivatives RA⁴, ⁵, ⁷, ¹⁰, ¹³ and ¹⁴. This is particularly important in the treatment of drug overdose since the respiratory muscle paralysis which is not affected by atropine and which is the cause of death induced by excess drug administration in the presence of atropine cannot be satisfactorily reversed by specific antidotes.

The duration of significant brain enzyme inhibition (>30%) induced by physostigmine (ED₅₀ dose) is less than 2 hours. Compounds RA₄, ⁵, ⁶, ⁷, ⁸, ¹², ¹⁴, ¹⁵ all act for more than 3 hours at their respective ED₅₀ doses and RA₆ and RA₇ still causes significant inhibition (36%) after 7 hours. Since none of these drugs caused noticeable side effects at the ED₅₀ doses, an even longer duration of action may be achieved by giving between 50 and 100% larger doses. The longer duration of action is a distinct advantage, particularly if the drugs are to be administered chronically to subjects suffering from neurological and behavioural conditions associated with a deficit in cholinergic transmission in the central nervous system, e.g. Alzheimer's disease, tardive dyskinesias, Huntingdon's chorea, Down's syndrome and Friedrich's ataxia.

The better the absorption of the drug after oral administration the more closely the LD₅₀ given by this route resembles that after subcutaneous injection. Table 3 shows that RA₆, ¹³, ⁷ and ⁴ are more efficiently absorbed from the gastro-intestinal tract than is physostigmine. The ED₅₀ of RA₆ after oral administration is the same as that after S.C. injection, indicating a much better oral bioavailability than that of physostigmine. The higher oral bioavailability of these compounds may be a considerable advantage for their clinical use.

RA₁₀, RA₆, RA₁₄ and RA₁₅ produce significant antagonism of the respiratory depressant effects of morphine in rabbits for periods lasting between 3—5 hours depending on the drug and the dose administered. The analgesic activity of morphine is not reduced by the RA compounds. Muscle fasciculations are not evident at the doses of drugs administered. Physostigmine (0.1—0.2 mg/kg) antagonizes the respiratory depressant effect of morphine for 30—60 mins only and fasciculations are marked at the higher dose.

These findings show that the RA compounds may be given together with morphine to obtain adequate analgesia without significant degrees of respiratory depression.

The most preferred compounds of the RA series are RA₄, RA₅, RA₆, RA₁₅, RA₁₄, RA₇ and RA₈, all of which produce inhibition of brain acetylcholinesterase after parenteral administration of significantly longer duration than that induced by physostigmine or miotine. These compounds also have a greater safety margin (therapeutic ratio) than physostigmine. RA₄, ⁶, ⁷ and ⁸ also show better bioavailability after oral administration than physostigmine. In addition, the acute toxicity (lethality) induced by RA₇ can be decreased more than 10-fold and that of RA₁₄ more than 8-fold by the antidote atropine, compared to only a 3-fold decrease for physostigmine and miotine.

The compounds of the invention are therefore useful for the treatment of senile dementia, Alzheimer's disease, Huntingdon's chorea, tardive dyskinesias, hyperkinesia, mania, acute confusion disorders, Down's syndrome and Friedrich's ataxia.

For these indications, the exact dosage will of course vary depending upon the compound employed, mode of administration and treatment desired. The compounds may be administered by any conventional route, non-oral or preferably orally.

An indicated total daily dosage is in the range from about 0.5 to about 25 mg of the compound, conveniently administered in divided doses 2 to 4 times a day in unit dosage form containing for example from about 0.1 to about 12 mg of the compound or in sustained release form.

The compounds may be administered in similar manner to known standards for use in these utilities. The suitable daily dosage for a particular compound will depend on a number of factors such as its relative potency of activity.


The compounds according to the invention may be administered in free base form or as a pharmaceutically acceptable acid addition salt. Such salts may be prepared in conventional manner and exhibit the same order of activity as the free forms.

It will be evident to those skilled in the art that the invention is not limited to the details of the foregoing illustrative embodiments and examples and that the present invention may be embodied in other specific

forms without departing from the essential attributes thereof, and it is, therefore, desired that the present embodiments and examples be considered in all respects as illustrative and not restrictive, reference being made to the appended claims, rather than to the foregoing description, and all changes which come with the meaning and range of equivalency of the claims are, therefore, intended to be embraced therein.

Claims for the Contracting States: BE CH DE FR GB IT LI LU NL SE

1. A compound of the general formula I



wherein

R₁ is hydrogen, lower alkyl, cyclohexyl, allyl or benzyl,

R₂ is hydrogen, methyl, ethyl or propyl, or

R₁ and R₂ together with the nitrogen to which they are attached form a morpholino or piperidino radical,

R₃ is hydrogen or lower alkyl,

R₄ and R₅ are the same or different and each is a lower alkyl, and the dialkylaminoalkyl group is in the meta, ortho or para position, provided that for compounds wherein R₄ and R₅ are both methyl and having the dialkylaminoalkyl group in the meta position, when R₂ is methyl and R₃ is hydrogen, R₁ is neither hydrogen nor methyl, and when R₂ and R₃ are methyl, R₁ is not hydrogen, and for compounds wherein R₄ and R₅ are both methyl and having the dialkylaminoalkyl group in the ortho or para position when R₁ and R₃ are both hydrogen R₂ is not methyl, or a pharmacologically acceptable salt thereof, for use as a pharmaceutical agent producing anticholinesterase activity in the central nervous system.

2. A compound according to claim 1, for use in the treatment of senile dementia, Alzheimer's disease, Huntingdon's chorea, tardive dyskinesia, hyperkinesia, mania, acute confusion conditions, Friedrich's ataxia and Down's syndrome.

3. A pharmaceutical composition adapted to produce anticholinesterase activity in the central nervous system containing as active agent a compound of formula I according to claim 1 or a pharmaceutically acceptable salt thereof.

4. A compound of the general formula I according to claim 1 or a pharmacologically acceptable salt thereof.

5. A compound according to claim 4 wherein the dialkylaminoalkyl group is in the meta position and R₄ and R₅ are both methyl.

6. N-propyl-3-[1-(dimethylamino)ethyl]phenyl carbamate or a pharmacologically acceptable salt thereof.

7. N-ethyl, N-methyl-3-[1-(dimethylamino)ethyl]phenyl carbamate or a pharmacologically acceptable salt thereof.

8. N,N-diethyl-3-[1-(dimethylamino)ethyl]phenyl carbamate or a pharmacologically acceptable salt thereof.

9. N-cyclohexyl-3-[1-(dimethylamino)ethyl]phenyl carbamate or a pharmacologically acceptable salt thereof.

10. N-allyl-3-[1-(dimethylamino)ethyl]phenyl carbamate or a pharmacologically acceptable salt thereof.

11. N-butyl-3-[1-(dimethylamino)ethyl]phenyl carbamate or a pharmacologically acceptable salt thereof.

12. N-methyl, N-propyl-3-[1-(dimethylamino)ethyl]phenyl carbamate or a pharmacologically acceptable salt thereof.


13. N-methyl, N-ethyl-3-[1-(dimethylamino)ethyl]phenyl carbamate or a pharmacologically acceptable salt thereof.



forms without departing from the essential attributes thereof, and it is, therefore, desired that the present embodiments and examples be considered in all respects as illustrative and not restrictive, reference being made to the appended claims, rather than to the foregoing description, and all changes which come with the meaning and range of equivalency of the claims are, therefore, intended to be embraced therein.

Claims for the Contracting States: BE CH DE FR GB IT LI LU NL SE

1. A compound of the general formula I



wherein

R_1 is hydrogen, lower alkyl, cyclohexyl, allyl or benzyl,

R_2 is hydrogen, methyl, ethyl or propyl, or

R_1 and R_2 together with the nitrogen to which they are attached form a morpholino or piperidino radical,

R_3 is hydrogen or lower alkyl,

R_4 and R_5 are the same or different and each is a lower alkyl, and the dialkylaminoalkyl group is in the meta, ortho or para position, provided that for compounds wherein R_4 and R_5 are both methyl and having the dialkylaminoalkyl group in the meta position, when R_2 is methyl and R_3 is hydrogen, R_1 is neither hydrogen nor methyl, and when R_2 and R_3 are methyl, R_1 is not hydrogen, and for compounds wherein R_4 and R_5 are both methyl and having the dialkylaminoalkyl group in the ortho or para position when R_1 and R_3 are both hydrogen R_2 is not methyl, or a pharmacologically acceptable salt thereof, for use as a pharmaceutical agent producing anticholinesterase activity in the central nervous system.

2. A compound according to claim 1, for use in the treatment of senile dementia, Alzheimer's disease, Huntington's chorea, tardive dyskinesia, hyperkinesia, mania, acute confusion conditions, Friedrich's ataxia and Down's syndrome.

3. A pharmaceutical composition adapted to produce anticholinesterase activity in the central nervous system containing as active agent a compound of formula I according to claim 1 or a pharmaceutically acceptable salt thereof.

4. A compound of the general formula I according to claim 1 or a pharmacologically acceptable salt thereof.

5. A compound according to claim 4 wherein the dialkylaminoalkyl group is in the meta position and R_4 and R_5 are both methyl.

6. N-propyl-3-[1-(dimethylamino)ethyl]phenyl carbamate or a pharmacologically acceptable salt thereof.

7. N-ethyl, N-methyl-3-[1-(dimethylamino)ethyl]phenyl carbamate or a pharmacologically acceptable salt thereof.

8. N,N-diethyl-3-[1-(dimethylamino)ethyl]phenyl carbamate or a pharmacologically acceptable salt thereof.

9. N-cyclohexyl-3-[1-(dimethylamino)ethyl]phenyl carbamate or a pharmacologically acceptable salt thereof.

10. N-allyl-3-[1-(dimethylamino)ethyl]phenyl carbamate or a pharmacologically acceptable salt thereof.


11. N-butyl-3-[1-(dimethylamino)ethyl]phenyl carbamate or a pharmacologically acceptable salt thereof.

12. N-methyl, N-propyl-3-[1-(dimethylamino)ethyl]phenyl carbamate or a pharmacologically acceptable salt thereof.

13. N-methyl, N-ethyl-3-[1-(dimethylamino)ethyl]phenyl carbamate or a pharmacologically acceptable salt thereof.



14. A process for the preparation of a compound according to claim 4, including the step of amidating a compound of formula II




II

wherein R_3 , R_4 and R_5 are as defined in claim 4, and recovering the so obtained compound of formula I in free base form or in form of a pharmacologically acceptable salt.

Claims for the Contracting State: AT

1. A process for the preparation of a compound of the general formula I



I

wherein


R_1 is hydrogen, lower alkyl, cyclohexyl, allyl or benzyl,

R_2 is hydrogen, methyl, ethyl or propyl, or

R_1 and R_2 together with the nitrogen to which they are attached form a morpholino or piperidino radical,

R_3 is hydrogen or lower alkyl,

R_4 and R_5 are the same or different and each is a lower alkyl, and the dialkylaminoalkyl group is in the meta, ortho or para position, provided that for compounds wherein R_4 and R_5 are both methyl and having the dialkylaminoalkyl group in the meta position, when R_2 is methyl and R_3 is hydrogen, R_1 is neither hydrogen nor methyl, and when R_2 and R_3 are methyl, R_1 is not hydrogen, and for compounds wherein R_4 and R_5 are both methyl and having the dialkylaminoalkyl group in the ortho or para position when R_1 and R_2 are both hydrogen R_3 is not methyl, or a pharmacologically acceptable salt thereof, including the step of amidating a compound of formula II



II

wherein R_3 , R_4 and R_5 are as defined above, and recovering the so obtained compound of formula I in free base form or in form of a pharmacologically acceptable salt.


2. A process according to claim 1 for the preparation of a compound of formula I wherein R_1 is hydrogen, including the step of reacting a compound of formula II with a isocyanate.

3. A process according to claim 1, including the step of reacting a compound of formula II with a carbamoyl halogenide.

4. A process according to claim 3, including the step of reacting a compound of formula II with a carbamoyl chloride.

5. A process according to claim 1 for the preparation of a compound wherein the dialkylaminoalkyl group is in the meta position and R_4 and R_5 are both methyl.

6. A compound of the general formula I



wherein

R_1 is hydrogen, lower alkyl, cyclohexyl, allyl or benzyl,

R_2 is hydrogen, methyl, ethyl or propyl, or

R_1 and R_2 together with the nitrogen to which they are attached form a morpholino or piperidino radical,

R_3 is hydrogen or lower alkyl,


R_4 and R_5 are the same or different and each is a lower alkyl, and the dialkylaminoalkyl group is in the meta, ortho or para position, provided that for compounds wherein R_4 and R_5 are both methyl and having the dialkylaminoalkyl group in the meta position, when R_2 is methyl and R_3 is hydrogen, R_1 is neither hydrogen nor methyl, and when R_2 and R_3 are methyl, R_1 is not hydrogen, and for compounds wherein R_4 and R_5 are both methyl and having the dialkylaminoalkyl group in the ortho or para position when R_1 and R_3 are both hydrogen R_2 is not methyl, or a pharmacologically acceptable salt thereof, whenever produced by a process according to any one of claims 1 to 5.

7. A pharmaceutical composition adapted to produce anticholinesterase activity in the central nervous system, comprising as active agent a compound according to claim 6.

8. A process for the production of a pharmaceutical composition according to claim 7, comprising the step of mixing a compound according to claim 6 with a physiologically acceptable carrier.

Patentansprüche für die Vertragsstaaten: BE CH DE FR GB IT LI LU NL SE

1. Verbindung der allgemeinen Formel I



worin

R_1 Wasserstoff, eine Niedrigalkyl-, Cyclohexyl-, Allyl- oder Benzylgruppe ist,



R_1 Wasserstoff, eine Methyl-, Ethyl- oder Propylgruppe ist oder R_1 und R_2 zusammen mit dem Stickstoffatom, an das sie gebunden sind, einen Morpholin- oder Piperidinrest bilden,

R_2 Wasserstoff oder eine Niedrigalkylgruppe ist,

R_4 und R_5 gleich sind oder verschieden und jeweils eine Niedrigalkylgruppe sind und die Dialkylaminoalkylgruppe in Meta-, Ortho- oder Parastellung ist, mit dem Vorbehalt, daß für Verbindungen, worin R_4 und R_5 beide Methylgruppen sind und die Dialkylaminoalkylgruppe in Metastellung ist, dann wenn R_2 eine Methylgruppe ist und R_3 Wasserstoff ist, R_1 weder Wasserstoff noch eine Methylgruppe ist und wenn R_2 und R_3 Methylgruppen sind, R_1 nicht Wasserstoff ist und für Verbindungen, worin R_4 und R_5 beide Methylgruppen sind und die Dialkylaminoalkylgruppe in Ortho- oder Parastellung ist, dann wenn R_1 und R_2 beide Wasserstoff sind, R_3 keine Methylgruppe ist, oder ein pharmakologisch annehmbares Salz davon zur Verwendung als pharmazeutisches Mittel zur Erzeugung von Anticholinesterase-Aktivität im Zentralnervensystem.

2. Verbindung nach Anspruch 1 zur Verwendung für die Behandlung von seniler Demenz, Alzheimer Krankheit, Chorea Huntington, tardiver Dyskinesie, Hyperkinesie, Manie, akuten Konfusionszuständen, Friedreich Ataxie und Down Syndrom.

3. Pharmazeutische Zusammensetzung geeignet eine Anticholinesterase-Aktivität im Zentralnervensystem zu erzeugen, enthaltend eine Verbindung der Formel I gemäß Anspruch 1 oder ein pharmazeutisch annehmbares Salz davon als aktiven Inhaltsstoff.

4. Verbindung allgemeinen Formel I gemäß Anspruch 1 oder ein pharmakologisch annehmbares Salz davon.

5. Verbindung nach Anspruch 4, worin die Dialkylaminoalkylgruppe sich in meta-Stellung befindet und beide Reste R_4 und R_5 Methylgruppen bedeuten.

6. N-Propyl-3-[1-(dimethylamino)ethyl]phenylcarbamat oder ein pharmazeutisch annehmbares Salz davon.

7. N-Ethyl, N-methyl-3-[1-(dimethylamino)ethyl]phenylcarbamat oder ein pharmazeutisch annehmbares Salz davon.

8. N,N-Diethyl-3-[1-(dimethylamino)ethyl]phenylcarbamat oder ein pharmazeutisch annehmbares Salz davon.

9. N-Cyclohexyl-3-[1-(dimethylamino)ethyl]phenylcarbamat oder ein pharmazeutisch annehmbares Salz davon.


10. N-Allyl-3-[1-(dimethylamino)ethyl]phenylcarbamat oder ein pharmazeutisch annehmbares Salz davon.

11. N-Butyl-3-[1-(dimethylamino)ethyl]phenylcarbamat oder ein pharmazeutisch annehmbares Salz davon.

12. N-Methyl, N-propyl-3-[1-(dimethylamino)ethyl]phenylcarbamat oder ein pharmazeutisch annehmbares Salz davon.

13. N-Methyl, N-ethyl-3-[1-(dimethylamino)ethyl]phenylcarbamat oder ein pharmazeutisch annehmbares Salz davon.

14. Verfahren zur Herstellung einer Verbindung gemäß Anspruch 4, einschließlich den Schritt, daß man eine Verbindung der Formel II




II

worin R_3 , R_4 und R_5 wie in Anspruch 4 definiert sind, amidiert und die so erhaltene Verbindung der Formel I in freier Basenform oder in Form eines pharmazeutisch annehmbaren Salzes gewinnt.



Patentanprüche für den Vt. tragetaat: AT

1. Verfahren zur Herstellung einer Verbindung der allgemeinen Formel I




I

20 worin

R₁ Wasserstoff, eine Niedrigalkyl-, Cyclohexyl-, Allyl- oder Benzylgruppe ist,
 R₂ Wasserstoff, eine Methyl-, Ethyl- oder Propylgruppe ist oder
 R₁ und R₂ zusammen mit dem Stickstoffatom, an das sie gebunden sind, einen Morpholin- oder
 Piperidinrest bilden,

25 R₃ Wasserstoff oder eine Niedrigalkylgruppe ist,

R₄ und R₅ gleich sind oder verschieden und jeweils eine Niedrigalkylgruppe sind und die
 Dialkylaminoalkylgruppe in Meta-, Ortho- oder Parastellung ist, mit dem Vorbehalt, daß für Verbindungen,
 worin R₄ und R₅ beide Methylgruppen sind und die Dialkylaminoalkylgruppe in Metastellung ist, dann
 wenn R₂ eine Methylgruppe ist und R₃ Wasserstoff ist, R₁ weder Wasserstoff noch eine Methylgruppe ist
 30 und wenn R₂ und R₃ Methylgruppen sind, R₁ nicht Wasserstoff ist und für Verbindungen, worin R₄ und R₅
 beide Methylgruppen sind und die Dialkylaminoalkylgruppe in Ortho- oder Parastellung ist, dann wenn R₁
 und R₂ beide Wasserstoff sind, R₃ keine Methylgruppe ist, oder ein pharmakologisch annehmbares Salz
 davon, das den Schritt einschließt, daß man eine Verbindung der Formel II



II

35 worin R₃, R₄ und R₅ wie oben definiert sind, amidiert und die so erhaltene Verbindung der Formel I in freier
 Basenform oder in Form eines pharmazeutisch annehmbaren Salzes gewinnt.

50 2. Verfahren nach Anspruch 1 für die Herstellung einer Verbindung der Formel I, worin R₁ Wasserstoff
 ist, das den Schritt einschließt, daß man eine Verbindung der Formel II mit einem Isocyanat umsetzt.


3. Verfahren nach Anspruch 1, das den Schritt einschließt, daß man eine Verbindung der Formel II mit
 einem Carbamoylhalogenid umsetzt.

55 4. Verfahren nach Anspruch 1, das den Schritt einschließt, daß man eine Verbindung der Formel II mit
 einem Carbamoylchlorid umsetzt.

5. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung, worin die Dialkylaminogruppe sich in
 der meta-Stellung befindet und beide Reste R₄ und R₅ Methylgruppen bedeuten.



6. Verbindung der allgemeinen Formel I



worin

R₁ Wasserstoff, eine Niedrigalkyl-, Cyclohexyl-, Alkyl- oder Benzylgruppe ist,

R₂ Wasserstoff, eine Methyl-, Ethyl- oder Propylgruppe ist oder

R₁ und R₂ zusammen mit dem Stickstoffatom an das sie gebunden sind, einen Morpholin- oder Piperidinrest bilden,

R₃ Wasserstoff oder eine Niedrigalkylgruppe ist,


R₄ und R₅ gleich sind oder verschieden und jeweils eine Niedrigalkylgruppe sind und die Dialkylaminoalkylgruppe in Meta-, Ortho- oder Parastellung ist, mit dem Vorbehalt, daß für Verbindungen, worin R₄ und R₅ beide Methylgruppen sind und die Dialkylaminoalkylgruppe in Metastellung ist, dann wenn R₂ eine Methylgruppe ist und R₃ Wasserstoff ist, R₁ weder Wasserstoff noch eine Methylgruppe ist und wenn R₂ und R₃ Methylgruppen sind, R₁ nicht Wasserstoff ist und für Verbindungen, worin R₄ und R₅ beide Methylgruppen sind und die Dialkylaminoalkylgruppe in Ortho- oder Parastellung ist, dann wenn R₁ und R₃ beide Wasserstoff sind, R₂ keine Methylgruppe ist, oder ein pharmakologisch annehmbares Salz, wenn nach einem Verfahren gemäß einem der Ansprüche 1 bis 5 hergestellt.

7. Pharmazeutische Zusammensetzung geeignet Anticholinesterase-Aktivität im Zentralnervensystem zu entwickeln, die als aktives Mittel eine Verbindung gemäß Anspruch 6 enthält.

8. Verfahren zur Herstellung einer pharmazeutischen Zusammensetzung nach Anspruch 7, das den Schritt umfaßt, daß eine Verbindung gemäß Anspruch 6 mit einem physiologisch annehmbaren Träger gemischt wird.

Revendications pour les Etats contractants: BE CH DE FR GB IT LI LU NL SE

1. Un composé de la formule générale I



dans laquelle

R₁ signifie l'hydrogène, alkyle inférieur, cyclohexyle, alkyle ou benzyle,

R₂ signifie l'hydrogène, méthyle, éthyle ou propyle, ou bien

R₁ et R₂ forment ensemble avec l'atome d'azote auquel il sont fixés, un radical morpholino ou pipéridino,

R₃ signifie l'hydrogène ou alkyle inférieur,

R₄ et R₅ sont identiques ou différents et signifient chacun alkyle inférieur et le groupe dialkylaminoalkyle est en position méta, ortho ou para, à condition que pour les composés dans lesquels R₄ et R₅ signifient tous les deux méthyle et ayant le groupe dialkylaminoalkyle en position méta, R₁ ne signifie pas



l'hydrogène ou méthyle lorsque R_1 signifie méthyle et R_2 l'hydrogène, et R_1 ne signifie pas l'hydrogène lorsque R_2 et R_3 signifient méthyle, et pour les composés dans lesquels R_4 et R_5 signifient tous les deux méthyle et ayant le groupe dialkylaminoalkyle en position ortho ou para, R_2 ne signifie pas méthyle lorsque R_1 et R_3 signifient tous les deux l'hydrogène, ou un sel pharmacologiquement acceptable de ce composé, pour l'utilisation comme agent pharmaceutique produisant une activité anticholinestérasique dans le système nerveux central.

2. Un composé selon la revendication 1, pour l'utilisation dans le traitement de la démence sénile, de la maladie de Alzheimer, de la chorée de Huntington, de la diskinésie tardive, de l'hyperkinésie, de la manie, des troubles confusionnels aigus, de l'ataxie de Friedrich et du syndrome de Down.

3. Une composition pharmaceutique adaptée pour produire une activité anticholinestérasique dans le système nerveux central, contenant, comme substance active, un composé de formule I selon la revendication 1 ou un sel pharmaceutiquement acceptable de ce composé.

4. Un composé de la formule générale I selon la revendication 1 ou un sel pharmacologiquement acceptable de ce composé.

5. Un composé selon la revendication 4, dans lequel le groupe dialkylaminoalkyle est en position méta et R_4 et R_5 signifient tous les deux méthyle.

6. Le N-propylcarbamate de 3-[1-(diméthylamino)éthyl]phényle ou un sel pharmacologiquement acceptable de ce composé.

7. Le N-éthyl, N-méthylcarbamate de 3-[1-(diméthylamino)éthyl]phényle ou un sel pharmacologiquement acceptable de ce composé.

8. Le N,N-diéthylcarbamate de 3-[1-(diméthylamino)éthyl]phényle ou un sel pharmacologiquement acceptable de ce composé.

9. Le N-cyclohexylcarbamate de 3-[1-(diméthylamino)éthyl]phényle ou un sel pharmacologiquement acceptable de ce composé.


10. Le N-allylcarbamate de 3-[1-(diméthylamino)éthyl]phényle ou un sel pharmacologiquement acceptable de ce composé.

11. Le N-butylcarbamate de 3-[1-(diméthylamino)éthyl]phényle ou un sel pharmacologiquement acceptable de ce composé.

12. Le N-méthyl, N-propylcarbamate de 3-[1-(diméthylamino)éthyl]phényle ou un sel pharmacologiquement acceptable de ce composé.

13. Le N-méthyl, N-éthylcarbamate de 3-[1-(diméthylamino)éthyl]phényle ou un sel pharmacologiquement acceptable de ce composé.

14. Un procédé de préparation d'un composé selon la revendication 4, comprenant l'amidation d'un composé de formule II




II

dans laquelle R_3 , R_4 et R_5 sont tels que définis à la revendication 4, et la récupération du composé de formule I ainsi obtenu sous forme de base libre ou sous forme d'un sel pharmacologiquement acceptable.



Revendications pour l'Etat contractant: AT

1. Un procédé de préparation d'un composé de la formule générale I



dans laquelle


R_1 signifie l'hydrogène, alkyle inférieur, cyclohexyle, allyle ou benzyle,

R_2 signifie l'hydrogène, méthyle, éthyle ou propyle, ou bien

R_1 et R_2 forment ensemble avec l'atome d'azote auquel il sont fixés, un radical morpholino ou pipéridino,

R_3 signifie l'hydrogène ou alkyle inférieur,

R_4 et R_5 sont identiques ou différents et signifient chacun alkyle inférieur et le groupe dialkylaminoalkyle est en position méta, ortho ou para, à condition que pour les composés dans lesquels R_4 et R_5 signifient tous les deux méthyle et ayant le groupe dialkylaminoalkyle en position méta, R_1 ne signifie pas l'hydrogène ou méthyle lorsque R_2 signifie méthyle et R_3 l'hydrogène, et R_1 ne signifie pas l'hydrogène lorsque R_2 et R_3 signifient méthyle, et pour les composés dans lesquels R_4 et R_5 signifient tous les deux méthyle et ayant le groupe dialkylaminoalkyle en position ortho ou para, R_2 ne signifie pas méthyle lorsque R_1 et R_3 signifient tous les deux l'hydrogène, ou un sel pharmacologiquement acceptable de ce composé, comprenant l'amidation d'un composé de formule II



dans laquelle R_3 , R_4 et R_5 sont tels que définis plus haut, et la récupération du composé de formule I ainsi obtenu sous forme de base libre ou sous forme d'un sel pharmacologiquement acceptable.

2. Un procédé selon la revendication 1, pour la préparation d'un composé de formule I où R_1 signifie l'hydrogène, comprenant la réaction d'un composé de formule II avec un isocyanate.


3. Un procédé selon la revendication 1, comprenant la réaction d'un composé de formule II avec un halogénure de carbamoyle.

4. Un procédé selon la revendication 1, comprenant la réaction d'un composé de formule II avec un chlorure de carbamoyle.

5. Un procédé selon la revendication 1, pour la préparation d'un composé dans lequel le groupe dialkylamino est en position méta et R_4 et R_5 signifient tous les deux méthyle.



6. Un composé de la formule générale I



dans laquelle

R_1 signifie l'hydrogène, alkyle inférieur, cyclohexyle, allyle ou benzyle,

R_2 signifie l'hydrogène, méthyle, éthyle ou propyle, ou bien

R_1 et R_2 forment ensemble avec l'atome d'azote auquel il sont fixés, un radical morpholino ou pipéridino,

R_3 signifie l'hydrogène ou alkyle inférieur,

R_4 et R_5 sont identiques ou différents et signifient chacun alkyle inférieur et le groupe dialkylaminoalkyle est en position méta, ortho ou para, à condition que pour les composés dans lesquels R_4 et R_5 signifient tous les deux méthyle et ayant le groupe dialkylaminoalkyle en position méta, R_1 ne signifie pas l'hydrogène ou méthyle lorsque R_2 signifie méthyle et R_3 l'hydrogène, et R_1 ne signifie pas l'hydrogène lorsque R_2 et R_3 signifient méthyle, et pour les composés dans lesquels R_4 et R_5 signifient tous les deux méthyle et ayant le groupe dialkylaminoalkyle en position ortho ou para, R_2 ne signifie pas méthyle lorsque R_1 et R_3 signifient tous les deux l'hydrogène, ou un sel pharmaceutiquement acceptable de ce composé, lorsqu'il est préparé par un procédé selon l'une quelconque des revendications 1 à 5.

7. Une composition pharmaceutique adaptée pour produire une activité anticholinestérasique dans le système nerveux central, comprenant, comme substance active, un composé selon la revendication 6.

8. Un procédé de préparation d'une composition pharmaceutique selon la revendication 7, comprenant le mélange d'un composé selon la revendication 6 avec un véhicule physiologiquement acceptable.