


Process for hydroformylation of olefins using Pt and DPEphos

The present invention relates to a process for hydroformylation of olefins using Pt and DPEphos.

5

P. Meessen et al., Journal of Organometallic Chemistry, 551, (1998), 165-170 describes the use of DPEphosPtCl₂ for the hydroformylation of methyl 3-pentenoate.




10

The problem addressed by the present invention is that of providing a novel hydroformylation process. The process here is to afford an increased yield compared to the process known from the prior art.

15 This object is achieved by a process according to Claim 1.

Process comprising the process steps of:

- a) initially charging an olefin;
- b) adding a compound of formula (I):



20

where R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸ are selected from: -H, -(C₁-C₁₂)-alkyl, -(C₆-C₂₀)-aryl, and at least two of the R⁵, R⁶, R⁷, R⁸ radicals are -(C₆-C₂₀)-aryl;

- c) adding a Pt compound capable of forming a complex;

- d) adding a bromine compound or an iodine compound;
- e) feeding in CO and H₂;
- f) heating the reaction mixture from steps a) to e), to convert the olefin to an aldehyde.

5 In this process, process steps a) to e) can be effected in any desired sequence. Typically, however, CO and H₂ are added after the co-reactants have been initially charged in steps a) to d).

It is possible here for process steps c) and d) to be effected in one step by adding PtBr₂ or PtI₂.

10 In a preferred variant of the process, the Pt compound and the bromine compound or iodine compound are added in one step by adding PtBr₂ or PtI₂.

The expression (C₁-C₁₂)-alkyl encompasses straight-chain and branched alkyl groups having 1 to 12 carbon atoms. These are preferably (C₁-C₈)-alkyl groups, more preferably (C₁-C₆)-alkyl, most preferably (C₁-C₄)-alkyl.

Suitable (C₁-C₁₂)-alkyl groups are especially methyl, ethyl, propyl, isopropyl, *n*-butyl, *iso*-butyl, *sec*-butyl, *tert*-butyl, *n*-pentyl, 2-pentyl, 2-methylbutyl, 3-methylbutyl, 1,2-dimethylpropyl, 1,1-dimethylpropyl, 2,2-dimethylpropyl, 1-ethylpropyl, *n*-hexyl, 2-hexyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 2,2-dimethylbutyl, 1,3-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethylbutyl, 1-ethyl-2-methylpropyl, *n*-heptyl, 2-heptyl, 3-heptyl, 2-ethylpentyl, 1-propylbutyl, *n*-octyl, 2-ethylhexyl, 2-propylheptyl, nonyl, decyl.

25 The expression (C₆-C₂₀)-aryl encompasses mono- or polycyclic aromatic hydrocarbyl radicals having 6 to 20 carbon atoms. These are preferably (C₆-C₁₄)-aryl, more preferably (C₆-C₁₀)-aryl.

Suitable (C₆-C₂₀)-aryl groups are especially phenyl, naphthyl, indenyl, fluorenyl, anthracenyl, phenanthrenyl, naphthacenyl, chrysenyl, pyrenyl, coronenyl. Preferred (C₆-C₂₀)-aryl groups are phenyl, naphthyl and anthracenyl.


In one variant of the process, R⁵, R⁶, R⁷, R⁸ are -(C₆-C₂₀)-aryl.

In one variant of the process, R⁵, R⁶, R⁷, R⁸ are -Ph.

35 In one variant of the process, R¹ and R⁴ are -H.

In one variant of the process, R^2 and R^3 are -H.

In one variant of the process, the compound of formula (I) has the structure (1):



In one variant of the process, the Pt compound is selected from: Pt(II)I₂, Pt(II)Br₂, Pt(IV)I₄, Pt(IV)Br₄, diphenyl(1,5-COD)Pt(II), Pt(II)(acac)₂, Pt(0)(PPh₃)₄, Pt(0)(DVTS) solution (CAS:68478-92-2), Pt(0)(ethylene)(PPh₃)₂, Pt(II)Br₂(COD), tris(benzylideneacetone)Pt(0),
10 Pt(II)(OAC)₂ solution, Pt(0)(t-Bu)₂, Pt(II)(COD)Me₂, Pt(II)(COD)I₂, Pt(IV)I Me₃, Pt(II)(hexafluoroacetylacetonate)₂.

In one variant of the process, the Pt compound is selected from: Pt(II)I₂, Pt(II)Br₂.

15 In one variant of the process, the iodine compound or the bromine compound is selected from: alkali metal halide, alkaline earth metal halide, NH₄X, alkylammonium halide, dialkyl halide, trialkyl halide, tetraalkyl halide, cycloalkylammonium halide.

In one variant of the process, a bromine compound which is Pt(II)Br₂ is added in process
20 step d).

In one variant of the process, the bromine compound is added in an amount in the range of 0.1 to 10, measured in equivalents based on Pt.

25 In one variant of the process, an iodine compound which is Pt(II)I₂ is added in process step d).

In one variant of the process, the iodine compound is added in an amount in the range of 0.1 to 10, measured in equivalents based on Pt.

30

In one variant of the process, this process comprises the additional process step e')

e') adding a solvent.

In one variant of the process, the solvent is selected from: THF, DCM, ACN, heptane, DMF, toluene, texanol, pentane, hexane, octane, isooctane, decane, dodecane, cyclohexane, benzene, xylene, Marlotherm, propylene carbonate, MTBE, diglyme, triglyme, diethyl ether, dioxane, isopropanol, tert-butanol, isononanol, isobutanol, isopentanol, ethyl acetate.

In one variant of the process, the solvent is selected from: THF, DCM, ACN, heptane, DMF, toluene, texanol.

In one variant of the process, CO and H₂ are fed in at a pressure in a range from 1 MPa (10 bar) to 6 MPa (60 bar).

In one variant of the process, CO and H₂ are fed in at a pressure in a range from 1 MPa (20 bar) to 6 MPa (50 bar).

In one variant of the process, the reaction mixture is heated to a temperature in the range from 25°C to 150°C.

In one variant of the process, the reaction mixture is heated to a temperature in the range from 30°C to 130°C.

In one variant of the process, the olefin is selected from: ethene, propene, 1-butene, cis- and/or trans-2-butene, isobutene, 1,3-butadiene, 1-pentene, cis- and/or trans-2-pentene, 2-methyl-1-butene, 3-methyl-1-butene, 2-methyl-2-butene, hexene, tetramethylethylene, heptene, 1-octene, 2-octene, di-n-butene, or mixtures thereof.

The invention is to be illustrated in more detail hereinafter by a working example.


Experimental description

A vial was charged with PtX₂ (X = halogen), ligand, and an oven-dried stirrer bar. The vial is then sealed with a septum (PTFE-coated styrene-butadiene rubber) and phenolic resin cap. The vial is evacuated and refilled with argon three times. Toluene and 1-octene were added to the vial using a syringe. The vial was placed in an alloy plate, which was transferred to an autoclave of the 4560 series from Parr Instruments under an argon atmosphere. After purging

the autoclave three times with CO/H₂, the synthesis gas pressure was increased to 40 bar at room temperature. The reaction was conducted at 120°C for 20 h. On termination of the reaction, the autoclave was cooled to room temperature and cautiously decompressed. Yield and selectivity were determined by GC analysis.

5

Hydroformylation of 1-octene



Reaction conditions:

- 10 10.0 mmol of 1-octene, 0.1 mol% PtX₂, 2.2 equivalents of ligand, solvent: toluene, p(CO/H₂): 40 bar, T: 120°C, t: 20 h.

Yields:

Ligand	Halogen	Yield [%]
	I / Br / Cl	76 / 45 / 14


15

As the experimental results show, the object is achieved by the process according to the invention.

Claims

5 1. Process comprising the process steps of:

- a) initially charging an olefin;
- b) adding a compound of formula (I):



where $R^1, R^2, R^3, R^4, R^5, R^6, R^7, R^8$ are selected from: -H, $-(C_1-C_{12})$ -alkyl, $-(C_6-C_{20})$ -aryl, and at

10 least two of the R^5, R^6, R^7, R^8 radicals are $-(C_6-C_{20})$ -aryl;

c) adding a Pt compound capable of forming a complex;

d) adding a bromine compound or an iodine compound;

e) feeding in CO and H_2 ;

f) heating the reaction mixture from steps a) to e), to convert the olefin to an aldehyde.

15

2. Process according to Claim 1,

where R^5, R^6, R^7, R^8 are $-(C_6-C_{20})$ -aryl.

3. Process according to either of Claims 1 and 2,

20 where R^5, R^6, R^7, R^8 are -Ph.

4. Process according to any of Claims 1 to 3,


where R^1 and R^4 are -H.

25 5. Process according to any of Claims 1 to 4,

where R^2 and R^3 are -H.

6. Process according to any of Claims 1 to 5,

wherein the compound of formula (I) has the structure (1):



7. Process according to any of Claims 1 to 6,

wherein the Pt compound is selected from: Pt(II)I₂, Pt(II)Br₂, Pt(IV)I₄, Pt(IV)Br₄, diphenyl(1,5-

- 5 COD)Pt(II), Pt(II)(acac)₂, Pt(0)(PPh₃)₄, Pt(0)(DVTs) solution (CAS:68478-92-2),
Pt(0)(ethylene)(PPh₃)₂, Pt(II)Br₂(COD), tris(benzylideneacetone)Pt(0), Pt(II)(OAC)₂ solution,
Pt(0)(t-Bu)₂, Pt(II)(COD)Me₂, Pt(II)(COD)I₂, Pt(IV)I Me₃, Pt(II)(hexafluoroacetylacetonate)₂.

8. Process according to any of Claims 1 to 7,

- 10 wherein a bromine compound which is Pt(II)Br₂ is added in process step d).

9. Process according to Claim 8,

wherein the bromine compound is added in an amount in the range of 0.1 to 10, measured in
equivalents based on Pt.

15

10. Process according to any of Claims 1 to 7,

wherein in process step d) an iodine compound is added, which is Pt(II)I₂.

11. Process according to Claim 10,

- 20 wherein the iodine compound is added in an amount in the range of 0.1 to 10, measured in
equivalents based on Pt.

12. Process according to any of Claims 1 to 11,

comprising the additional process step e')

- 25 e') adding a solvent.

13. Process according to Claim 12,

wherein the solvent is selected from: THF, DCM, ACN, heptane, DMF, toluene, texanol,
pentane, hexane, octane, isooctane, decane, dodecane, cyclohexane, benzene, xylene,
30 Marlotherm, propylene carbonate, MTBE, diglyme, triglyme, diethyl ether, dioxane,
isopropanol, tert-butanol, isononanol, isobutanol, isopentanol, ethyl acetate.

14. Process according to any of Claims 1 to 13,
wherein CO and H₂ are fed in at a pressure in a range from 1 MPa (10 bar) to 6 MPa (60 bar).

5

15. Process according to any of Claims 1 to 14,
wherein the reaction mixture is heated to a temperature in the range from 25°C to 150°C.