

COMPOSITIONS AND METHODS FOR RNA-ENCODED DNA-REPLACEMENT OF ALLELES

STATEMENT REGARDING ELECTRONIC FILING OF A SEQUENCE LISTING

5 A Sequence Listing in ASCII text format, submitted under 37 C.F.R. § 1.821, entitled 1499.57.WO_ST25.txt, 1,043,362 bytes in size, generated on November 5, 2021, and filed via EFS-Web, is provided in lieu of a paper copy. This Sequence Listing is hereby incorporated herein by reference into the specification for its disclosures.

STATEMENT OF PRIORITY

10 This application claims the benefit, under 35 U.S.C. § 119 (e), of U.S. Provisional Application No. 63/110,386 filed on November 6, 2020, the entire contents of which is incorporated by reference herein.

FIELD OF THE INVENTION

15 This invention relates to recombinant nucleic constructs comprising CRISPR-Cas effector proteins, reverse transcriptases and extended guide nucleic acids and methods of use thereof for modifying nucleic acids in plants.

BACKGROUND OF THE INVENTION

20 Base editing has been shown to be an efficient way to change cytosine and adenine residues to thymine and guanine, respectively. These tools, while powerful, do have some limitations such as bystander bases, small base editing windows that give limited accessibility to trait-relevant targets unless enzymes with high PAM density are available to compensate, 25 limited ability to convert cytosines and adenines to residues other than thymine and guanine, respectively, and no ability to edit thymine or guanine residues. Thus, the current tools available for base editing are limited. Therefore, to make nucleic acid editing more useful by increasing the range of possible edits for a greater number of organisms, new editing tools are needed.

SUMMARY OF THE INVENTION

30 In a first aspect, a method of modifying a target nucleic acid is provided, the method comprising: contacting the target nucleic acid with (a) a Type V CRISPR-Cas effector protein or a Type II CRISPR-Cas effector protein; (b) a reverse transcriptase, and (c) an extended guide nucleic acid (e.g., extended Type II or Type V CRISPR RNA, extended Type II or

Type V CRISPR DNA, extended Type II or Type V crRNA, extended Type II or Type V crDNA), thereby modifying the target nucleic acid.

In a second aspect, a method of modifying a target nucleic acid is provided, the method comprising: contacting the target nucleic acid at a first site with (a)(i) a first CRISPR-Cas effector protein; and (ii) a first extended guide nucleic acid (e.g., extended CRISPR RNA, extended CRISPR DNA, extended crRNA, extended crDNA); and (b)(i) a second CRISPR-Cas effector protein, (ii) a first reverse transcriptase; and (ii) a first guide nucleic acid, thereby modifying the target nucleic acid.

In a third aspect, a method of modifying a target nucleic acid in a plant or plant cell is provided, comprising introducing the expression cassette of the invention into the plant or plant cell, thereby modifying the target nucleic acid in the plant or plant cell and producing a plant or plant cell comprising the modified target nucleic acid.

In a fourth aspect, a complex is provided comprising: (a) a Type V CRISPR-Cas effector protein or a Type II CRISPR-Cas effector protein; (b) a reverse transcriptase, and (c) an extended guide nucleic acid (e.g., extended CRISPR RNA, extended CRISPR DNA, extended crRNA, extended crDNA, e.g., targeted allele guide (tag) nucleic acid (i.e., tagDNA, tagRNA)).

In a fifth aspect, an expression cassette codon optimized for expression in an organism is provided, the expression cassette comprising 5' to 3' (a) polynucleotide encoding a plant specific promoter sequence (e.g., ZmUbi1, MtUbi2, RNA polymerase II (Pol II)), (b) a plant codon-optimized polynucleotide encoding a Type V CRISPR-Cas nuclease (e.g., Cpf1 (Cas12a), dCas12a and the like); (c) a linker sequence; and (d) a plant codon-optimized polynucleotide encoding a reverse transcriptase.

In a sixth aspect, an expression cassette codon optimized for expression in an organism is provided, the expression cassette comprising: (a) a polynucleotide encoding a promoter sequence, and (b) an extended RNA guide sequence, wherein the extended guide nucleic acid comprises an extended portion comprising at its 3' end a primer binding site and an edit to be incorporated into the target nucleic acid (e.g., reverse transcriptase template), optionally wherein the extended guide nucleic acid is comprised in an expression cassette, optionally wherein the extended guide nucleic acid is operably linked to a Pol II promoter.

The invention further provides cells, including plant cells, bacterial cells, archaea cells, fungal cells, animal cells comprising target nucleic acids modified by the methods of the invention as well as organisms, including plants, bacteria, archaea, fungi, and animals,

comprising the cells. Additionally, the present invention provides kits comprising the polynucleotides, polypeptides, and expression cassettes of the invention.

These and other aspects of the invention are set forth in more detail in the description of the invention below.

5

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 provides a schematic showing the generation of DNA sequences from reverse transcription off the crRNA and subsequent integration into the nick site. The extended guide crRNA (tagRNA) is bound to the Cpf1 nickase (cas12a nickase) (nCpf1, upper left). Alternatively, the extension encoding the edit template may be located 5' of the crRNA. The 3' end of the crRNA is complimentary to the DNA at the nick site (nonbold pairing lines, upper left). The nCpf1 may be either covalently linked to the reverse transcriptase (RT) or the RT may be recruited to the nCpf1, in which case multiple reverse transcriptase proteins may be recruited to the nCpf1. The RT polymerizes DNA from the 3' end of the DNA nick on the second strand generating a DNA sequence complimentary to the crRNA with nucleotides non-complimentary to the genome (bolded pairing lines, brace, upper right) followed by complimentary nucleotides (non-bold pairing lines, upper right). Upon dissociation, the resultant DNA has an extended ssDNA with a 3' overhang, which is largely the same sequence as the original DNA (non-bolded pairing lines, lower right) but with some non-native nucleotides (bolded pairing lines, brace, lower right). This flap is in equilibrium with a structure having a 5' overhang (lower left) where there are mismatched nucleotides incorporated into the DNA. The equilibrium may be driven toward the structure on the left by reducing mismatch repair, removal of the 5' flap during repair and replication, and also by nicking the first strand as described herein.

Fig. 2 provides a schematic of showing a method for reducing mismatch repair. In order to drive the equilibrium more favorable for forming the final product with the modified nucleotides (bolded, brace), a nickase is directed (via a guide nucleic acid) to cut the first strand (e.g., target strand or bottom strand) of the target nucleic acid in a region outside of the RT-editing region (lightning bolts) - a distance from the nick in the second strand (e.g., target strand or top strand). The nCpf1:crRNA molecules may be on either side or both sides of the editing bubble. Nicking the first strand (dashed line) indicates to the cell that the newly incorporated nucleotides are the correct nucleotides during mismatch repair and replication,

thus favoring a final product with the new nucleotides. Other possible ways of driving the equilibrium toward the desired product can include removal of the 5' flap.

Fig. 3 shows alternative methods of modifying nucleic acids using the compositions of the present invention, wherein in two nicks are introduced in the second strand and the sequence introduced by the RT displaces the double-nicked WT sequence and thereby, is more efficiently incorporated into the genome.

Fig. 4. LbCas12a_R1138A is a nickase as demonstrated *in vitro*, resolved on a 1% TAE-agarose gel. A supercoiled 2.8 kB plasmid ran with an apparent size of 2.0 kB (lane 2) until a double-stranded break was generated by wildtype LbCas12a (lane 3).

Fig. 5 shows configurations of REDRAW editors tested in *E. coli* (see Example 1).

Fig. 6 shows conformations of tagRNAs tested in the first library.

Fig. 7 shows the structure of an example designed hairpin sequence for use in REDRAW editing (**SEQ ID NO:203**).

Fig. 8 shows Sanger sequencing results demonstrating a TGA > CTG edit in a defunct *aadA* gene, restoring antibiotic resistance (**SEQ ID NOs 204-208**). The edit was observed from a colony in Selection 10, with protein configuration SV40-MMLV-RT-XTEN-nLbCas12a-SV40 (**SEQ ID NO:71**).

Fig. 9 shows Sanger sequencing results demonstrating an AAA > CGT edit in the *rpsL* gene in the *E. coli* genome, conferring resistance to the antibiotic streptomycin (**SEQ ID NOs 209-211**). The edit was observed from a colony in Selection 2.5, with protein configuration SV40-MMLV-RT-XTEN-nRVRLbCas12a(H759A)-SV40 (**SEQ ID NO:79**).

Fig. 10 shows Sanger sequencing results demonstrating a TGA > GAT edit in a defunct *aadA* gene, restoring antibiotic resistance (**SEQ ID NOs 212-215**). The edit was observed from a colony in Selection 2.25, with protein configuration SV40-nLbCas12a-XTEN-MMLV-RT-SV40 (**SEQ ID NO:73**).

Fig. 11 shows Sanger sequencing results demonstrating a TGA > GAT edit in a defunct *aadA* gene, restoring antibiotic resistance (**SEQ ID NOs 212-215**). The edit was observed from a colony in Selection 2.31, with protein configuration SV40-MMLV-RT-XTEN-nLbCas12a(H759A)-SV40 (**SEQ ID NO:83**).

Fig. 12 shows an example editing method carried out in human cells (see Example 2). Panel A shows the double stranded target nucleic acid. Cas12a complex (complex includes the extended guide nucleic acid, which is not shown) is recruited to the first strand (target strand, bottom strand) with the 5' flap in the second strand (top strand, non-target strand), optionally being removed with a 5'-3' exonuclease (Panel B). Panel C shows the reverse

transcriptase MMuLV-RT (5M) **SEQ ID NO:53**) extends from the priming site or primer (complementary to the primer binding site) on the target nucleic (dashed line = the extension). Panels D and E show the resolution of DNA intermediates via mismatch repair and DNA ligation and generation of a new edited DNA strand.

5 **Fig. 13** shows precise editing using various guide conformations in HEK293T cells at FANCF1 site. The construct name is Cas12a (H759A) + RT(5M) + RecE FANCF1.

Fig. 14 shows precise editing using various guide conformations in HEK293T cells at DMNT1 site. The construct name is Cas12a (H759A) + RT(5M).

10 **Fig. 15** shows the effect of exonuclease transfection on precise editing activity (normalized to no exonuclease treatment; pUC19 = 1) at DMNT1 site.

Fig. 16 shows various forms of REDRAW architecture (i.e., constructs of the invention) and the percent precise editing of each. The left panel shows the reverse transcriptase (RT) provided in trans (no recruitment). The middle panel shows recruitment of the RT using, as an example, SunTag (e.g., GCN4, e.g., **SEQ ID NO:23**) that is fused to the C-terminus of LbCpf1 (LbCas12a) (LbCpf1-SunTag), which can recruit antibody fused to the N-terminus of RT(5M) (scFv-RT (5M)) (e.g., scFv, **SEQ ID NO:25**). The right panel shows RT and LbCpf1 fusion proteins. The left side of the right panel shows the results with the RT fused to the C-terminus of LbCpf1 and the right side of the right panel shows the results with the RT fused to the N-terminus of LbCpf1.

20 **Fig. 17** provides a schematic of the use of 5'-3' exonuclease to degrade the DNA at both ends of the double-stranded break generated during the REDRAW process.

Fig. 18 shows the percent precise editing of REDRAW using a 5'-3' exonuclease (RecE (**SEQ ID NO:129**), RecJ (**SEQ ID NO:130**), T5_Exo (**SEQ ID NO:131**), T7_Exo (**SEQ ID NO:132**)) that is fused to the C-terminus of the Cas polypeptide (LbCpf1). In this configuration, RT(5M) (**SEQ ID NO:53**) is expressed in trans (no recruitment).

Fig. 19 shows the percent precise editing of REDRAW using either the 5'-3' exonuclease sbcB (**SEQ ID NO:134**) or the 5'-3' exonuclease Exo (**SEQ ID NO:135**) each fused to the C-terminus of a Cas polypeptide (LbCpf1). RT (5M) is expressed in trans (no recruitment).

30 **Fig. 20** shows the percent precise editing of REDRAW using trans expression of exonucleases. The LbCpf1 and RT are provided as fusion proteins. The right side of **Fig. 20** shows results with the RT fused to the N-terminus of the LbCpf1 (RT(5M)-LbCpf1 (H759A)) and the left side of the figure shows the results using an RT fused to the C-terminus of the LbCpf1 (LbCpf1 (H759A)-RT(5M)).

Fig. 21 shows the effect on percent precise editing of REDRAW of example mutations in a Cas12a (LbCpf1) in the REDRAW process. The example mutations tested included K167A, K272A, K349A, K167A+ K272A, K167A+ K349A, K272A+ K349A, and K167A+ K272A + K349A (positions relative to LbCas12a (H759A) **SEQ ID NO:148**).

5 **Fig. 22** shows the percent precise editing of REDRAW in the presence of single stranded DNA binding proteins (ssDNA BP). The ssDNA BP was expressed in trans in the presence of the CRISPR-Cas effector polypeptide (e.g., LbCpf1 (H759A)), RT(5M), and tagRNA1. The RT and LbCpf1 (H759A) were also expressed in trans in this example. The ssDNA BPs tested were hRad51_s208E_A209D, hRad52, BsRecA, EcRecA, and T4SSB.

10 Mock is no ssDNA BP.

Fig. 23 shows the percent precise editing of REDRAW in the presence of single stranded DNA binding proteins (ssDNA BP) when fused to a CRISPR-Cas effector polypeptide (e.g., LbCas12a H759A). ssDNA binding proteins (hRad51, hRad52, BsRecA, EcRecA, T4SSB and Brex27) were fused to N terminus or C-terminus of LbCpf1 (H759A).

15 RT(5M) and the tagRNAs were expressed in trans.

Fig. 24 shows the effect of on the percent of indels produced when REDRAW is carried out in the presence of a polypeptide that prevents NHEJ. In this example, the polypeptide that prevents NHEJ is Gam protein (*Escherichia* phage Mu Gam protein) (**SEQ ID NO:147**), and the reverse transcriptase is expressed in trans, either as a native sequence

20 (e.g., RT(5M)) or with Gam fused to the N-terminus of RT (e.g., Gam-RT(5M)). These constructs are expressed concurrently with either LbCas12a (H759A) or with an LbCas12a (H759A) having a Gam protein fused to its N-terminus (e.g., Gam-LbCas12a H759A).

Fig. 25 shows the percent precise editing of REDRAW in the presence Gam protein. The Gam protein is provided in trans, as a fusion protein with the reverse transcriptase (N-terminal fusion; Gam-RT(5M)) and/or as a fusion protein with the CRISPR-Cas effector polypeptide (e.g., Gam-LbCas12a H759A).

25

Fig. 26 shows the percent precise editing of REDRAW using different length primer binding sites (PBS) and reverse transcriptase templates (RTT). The top and bottom panels show the results using two different spacers (top panel: pwsp143

30 (GCTCAGCAGGCACCTGCCTCAGC) (**SEQ ID NO:136**), bottom panel: pwsp139 (CTGATGGTCCATGTCTGTTACTC)(**SEQ ID NO:137**).

Fig. 27 shows the percent editing depending on the location of the edit in two different reverse transcriptase templates (RTTs). The edit was placed in each RTT at positions varying from position -1 to position 19 (numbering is relative to the protospacer

adjacent motif numbering in the target nucleic acid) (edit in bold font). RTT in the upper panel: TTTGGCTCACTCCTGCTCGGTGAATTT **SEQ ID NO:187**; RTT in the lower panel: TTTCGCGCTTGTCCAATCAGTACGCA **SEQ ID NO:188**.

Fig. 28 shows the percent precise editing of REDRAW using two forms of Cas9, a nuclease (Cas9) and a nickase (nCas9 (D10A mutant)). Both Cas9 and nCas9 were tested using tagRNAs with extensions attached to either the 3' end or the 5' end of the guide RNA (denoted as 3' extension or 5' extension). The lengths of RTT and PBS of the tagRNA extensions were varied and the spacers targeted four different sites (pwspl0: GAGTCCGAGCAGAAGAAGAA (**SEQ ID NO:140**); pwspl621: GCATTTTCAGGAGGAAGCGA (**SEQ ID NO:141**); pwspl15: GTCATCTTAGTCATTACCTG (**SEQ ID NO:142**); pwspl11: GGAATCCCTTCTGCAGCACC (**SEQ ID NO:143**).

Fig. 29 shows the percent precise editing of REDRAW using BhCas12b. The BhCas12b was tested using tagRNAs with extensions attached to either the 3' end or the 5' end of guide RNA (denoted as 3' or 5'). The lengths of RTT and PBS of the tagRNA extensions were varied and the spacers targeted three different sites (PWspl099: ACGTACTGATGTAAACAGCTGA (**SEQ ID NO:144**); PWspl098: GGTCAGCTGTAAACATCAGTAC (**SEQ ID NO:145**); PWspl094: TCCAGCCCCGCTGGCCCTGTAAA (**SEQ ID NO:146**).

Fig. 30 shows the percent precise editing of REDRAW using EnAsCpfI (H800A) (**SEQ ID NO:149**). The left panel shows editing without RT(5M), the middle panel shows editing with an EnAsCpfI (H800A) having a C-terminal fused RT(5M) (EnAsCpfI (H800A)-RT(5M)) and the right panel shows editing with an EnAsCpfI (H800A) having an N-terminal fused RT(5M) (RT(5M)-EnAsCpfI (H800A)). In this example, a single site was targeted with the spacer having the sequence of CCTCACTCCTGCTCGGTGAATTT (**SEQ ID NO:171**).

Fig. 31 shows the editing results for the URA3-I target gene in yeast using the methods of the present invention (REDRAW). The upper panel shows editing results (colony formation upon repair of adenine auxotrophy by editing) using a LbCas12a having a reverse transcriptase (RT) fused to its C-terminus. The lower panel shows editing results (colony formation upon repair of adenine auxotrophy by editing) using a LbCas12a having a RT fused to its N-terminus. The extended guide used for the editing shown in **Fig. 31** either does not have a pseudoknot or includes a pseudoknot at its 3' end. The pseudoknots are referred to either as a decoy hairpin (**SEQ ID NO:95**; **SEQ ID NO:203**), tEvoPreQ1 (**SEQ ID NO:158**)

or EvoPreQ1 (SEQ ID NO:191). The extended guide further includes an RTT having a length of 47, 55 or 63 nucleotides and a PBS having a length of 48 nucleotides.

Fig. 32 shows the editing results for the ADE2 target gene in yeast using the methods of the present invention (REDRAW). The upper panel shows editing results (colony formation upon repair of uracil auxotrophy by editing) using a LbCas12a having a RT fused to its C-terminus. The lower panel shows editing results (colony formation upon repair of uracil auxotrophy by editing) using a LbCas12a having a RT fused to its N-terminus. The extended guide used for the editing shown in **Fig. 32** either does not have a pseudoknot or includes a pseudoknot at its 3' end. The pseudoknots used are referred to either as a decoy hairpin (SEQ ID NO:95, SEQ ID NO:203) tEvoPreQ1 (SEQ ID NO:158) or EvoPreQ1 (SEQ ID NO:191). The extended guide further includes an RTT having a length of 40, 50 or 72 nucleotides and a PBS having a length of 48 nucleotides. In general, the extended guide nucleic acid comprises 5'-3' an RTT, a PBS and when present, a 3' pseudoknot. In the first column of data for the decoy hairpin, in both the upper and lower panels, the tagRNA with 40-bp RTT and decoy hairpin was unable to be synthesized and the condition was not tested.

Fig. 33 shows the percent precise editing results when using the ssRNA binding proteins, defensin (SEQ ID NO:152) and ORF5 (SEQ ID NO:153), each fused to the N-terminus of a RT-LbCas12 fusion protein (e.g., RT-LbCas12a) as compared to the same RT-Cas12a fusion protein that does not comprise a ssRNA binding protein fused at its N-terminus.

Fig. 34 shows the percent precise editing results when using LbCas12a (H759A) fused at its N-terminus to reverse transcriptase (RT) domains having different mutations. The RT included: RT(L139P, D200N, W388R, E607K), RT(L139P, D200N, T306K, W313F, W388R, E607K), RT(5M, F155Y, H638G), RT(5M, Q221R, V223M) and RT(5M, D524N).

Fig. 35 shows the percent precise editing results using four different tagRNAs comprising a structured RNA at the 3' end of each tag RNA. The nucleic acid sequences of the structured RNAs are provided in **Table 16**.

Fig. 36 shows the percent precise editing results using chromatin modulating peptides fused to constructs of the invention in various fusion orientations. The tested chromatin modulating peptides included HN1, HB1, H1G, and CHD1.

Fig. 37 shows the percent precise editing results for fusions using MS2/MCP system. LbCas12a H759A with RT(5M) was transiently expressed without MCP (in trans control), or with MCP-RT(5M) (fusion construct). Two tagRNAs were tested, tagRNA5 and tagRNA6.

The different tagRNA versions tested included the tagRNAs modified with MS2 sequence at their 3' end.

BRIEF DESCRIPTION OF THE SEQUENCES

- 5 **SEQ ID NOs:1-20** and **148-150** are example Cas12a amino acid sequences.
 SEQ ID NO:21 and **SEQ ID NO:22** are exemplary regulatory sequences encoding a promoter and intron.
 SEQ ID NOs:23-25 provide example peptide tags and affinity polypeptides.
 SEQ ID NO:26-36 provide example RNA recruiting motifs and corresponding
 10 affinity polypeptides.
 SEQ ID NOS:37-52 provide example single stranded RNA binding domains (RBDs)
 SEQ ID NOs:53, 97 and **172** provide example reverse transcriptase polypeptide sequences: Moloney Murine Leukemia Virus (M-MuLV)5(M), 5(M) flanked with NLS, and M-MuLV, respectively.
 15 **SEQ ID NOs:54-56** provides an example of a protospacer adjacent motif position for a Type V CRISPR-Cas12a nuclease.
 SEQ ID NO:57 and **SEQ ID NO:58** provide example constructs of the invention.
 SEQ ID NO:59 and **SEQ ID NO:60** provide an example CRISPR RNA and an example protospacer.
 20 **SEQ ID NO:61** and **SEQ ID NO:62** provide example introns.
 SEQ ID NOs:63-86 and **SEQ ID NOs:154-157** provide example REDRAW editor constructs.
 SEQ ID NO:87 provides an example of a tagRNA having an 11 base pair (bp) primer binding sequence and a 96 bp reverse transcriptase template.
 25 **SEQ ID NOs:88-91** provide sequences of example plasmids.
 SEQ ID NOs:92-94 provide sequences of tagRNAs associated with the edits shown in Figs. 9-11, respectively.
 SEQ ID NO:95, SEQ ID NO:158, SEQ ID NO:191 and **SEQ ID NO:203** and provide example pseudoknots sequences.
 30 **SEQ ID NO:96** provides an example LbCas12a having a mutation of H759A and flanked with NLS on both sides.
 SEQ ID NOs:98-101 provide example 5'-3' exonuclease polypeptides.
 SEQ ID NO:102 and **SEQ ID NO:103** provide example DMNT1 target site and target spacer.

SEQ ID NO:104 and SEQ ID NO:105 provide example FANCF1 target site and target spacer.

SEQ ID NO:106 and SEQ ID NO:107 provide example Cas9 polypeptides.

SEQ ID NOs:108-122 provide example Cas9 polynucleotides

5 SEQ ID NOs:123-128 provide example single stranded DNA binding proteins.

SEQ ID NOs:129-135 provide example 5'-3' exonucleases.

SEQ ID NOs:136, 137, 140-146, 159-161 and 171 are example spacers .

SEQ ID NOs:138, 139 and 164-169 provide example reverse transcriptase templates.

SEQ ID NO:140 provides an example Gam protein.

10 SEQ ID NO:151 provides an example Cas12b polypeptide.

SEQ ID NO:152 and SEQ ID NO:153 provide example single stranded RNA binding proteins, defensin and ORF5, respectively.

SEQ ID NO:162 and SEQ ID NO:163 provide example Primer Binding Site (PBS) sequences.

15 SEQ ID NO:170 provides an example LbCas12a crRNA scaffold.

SEQ ID NOs:173-186 provide example tagRNAs (tagRNA 1, tagRNA 2, tagRNA 3, tagRNA 4, tagRNA 5, tagRNA 6, tagRNA 7, tagRNA 8, tagRNA 9, tagRNA 10, tagRNA 11, tagRNA 12, tagRNA 13, and tagRNA 14, respectively).

20 SEQ ID NO:187 and SEQ ID NO:188 are the reverse transcriptase templates shown in Fig. 27.

SEQ ID NOs:95, 189-198, and 203 are example RNA structures.

SEQ ID NOs:199-202 are example chromatin modulating peptides.

SEQ ID NOs:204-215 are sequences found in Figs. 8, 9, 10 and 11.

25 DETAILED DESCRIPTION

The present invention now will be described hereinafter with reference to the accompanying drawings and examples, in which embodiments of the invention are shown. This description is not intended to be a detailed catalog of all the different ways in which the invention may be implemented, or all the features that may be added to the instant invention.

30 For example, features illustrated with respect to one embodiment may be incorporated into other embodiments, and features illustrated with respect to a particular embodiment may be deleted from that embodiment. Thus, the invention contemplates that in some embodiments of the invention, any feature or combination of features set forth herein can be excluded or omitted. In addition, numerous variations and additions to the various embodiments

suggested herein will be apparent to those skilled in the art in light of the instant disclosure, which do not depart from the instant invention. Hence, the following descriptions are intended to illustrate some particular embodiments of the invention, and not to exhaustively specify all permutations, combinations and variations thereof.

5 Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. The terminology used in the description of the invention herein is for the purpose of describing particular embodiments only and is not intended to be limiting of the invention.

10 All publications, patent applications, patents and other references cited herein are incorporated by reference in their entireties for the teachings relevant to the sentence and/or paragraph in which the reference is presented.

Unless the context indicates otherwise, it is specifically intended that the various features of the invention described herein can be used in any combination. Moreover, the
15 present invention also contemplates that in some embodiments of the invention, any feature or combination of features set forth herein can be excluded or omitted. To illustrate, if the specification states that a composition comprises components A, B and C, it is specifically intended that any of A, B or C, or a combination thereof, can be omitted and disclaimed singularly or in any combination.

20 As used in the description of the invention and the appended claims, the singular forms “a,” “an” and “the” are intended to include the plural forms as well, unless the context clearly indicates otherwise.

Also as used herein, “and/or” refers to and encompasses any and all possible combinations of one or more of the associated listed items, as well as the lack of
25 combinations when interpreted in the alternative (“or”).

The term “about,” as used herein when referring to a measurable value such as an amount or concentration and the like, is meant to encompass variations of $\pm 10\%$, $\pm 5\%$, $\pm 1\%$, $\pm 0.5\%$, or even $\pm 0.1\%$ of the specified value as well as the specified value. For example, “about X” where X is the measurable value, is meant to include X as well as
30 variations of $\pm 10\%$, $\pm 5\%$, $\pm 1\%$, $\pm 0.5\%$, or even $\pm 0.1\%$ of X. A range provided herein for a measurable value may include any other range and/or individual value therein.

As used herein, phrases such as “between X and Y” and “between about X and Y” should be interpreted to include X and Y. As used herein, phrases such as “between about X

and Y” mean “between about X and about Y” and phrases such as “from about X to Y” mean “from about X to about Y.”

Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless
5 otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein. For example, if the range 10 to 15 is disclosed, then 11, 12, 13, and 14 are also disclosed.

The term “comprise,” “comprises” and “comprising” as used herein, specify the presence of the stated features, integers, steps, operations, elements, and/or components, but
10 do not preclude the presence or addition of one or more other features, integers, steps, operations, elements, components, and/or groups thereof.

As used herein, the transitional phrase “consisting essentially of” means that the scope of a claim is to be interpreted to encompass the specified materials or steps recited in the claim and those that do not materially affect the basic and novel characteristic(s) of the
15 claimed invention. Thus, the term “consisting essentially of” when used in a claim of this invention is not intended to be interpreted to be equivalent to “comprising.”

As used herein, the terms “increase,” “increasing,” “enhance,” “enhancing,” “improve” and “improving” (and grammatical variations thereof) describe an elevation of at least about 25%, 50%, 75%, 100%, 150%, 200%, 300%, 400%, 500% or more as compared
20 to a control.

As used herein, the terms “reduce,” “reduced,” “reducing,” “reduction,” “diminish,” and “decrease” (and grammatical variations thereof), describe, for example, a decrease of at least about 5%, 10%, 15%, 20%, 25%, 35%, 50%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99%, or 100% as compared to a control. In particular embodiments, the reduction can result
25 in no or essentially no (*i.e.*, an insignificant amount, *e.g.*, less than about 10% or even 5%) detectable activity or amount.

A “heterologous” or a “recombinant” nucleotide sequence is a nucleotide sequence not naturally associated with a host cell into which it is introduced, including non- naturally occurring multiple copies of a naturally occurring nucleotide sequence.

30 A “native” or “wild type” nucleic acid, nucleotide sequence, polypeptide or amino acid sequence refers to a naturally occurring or endogenous nucleic acid, nucleotide sequence, polypeptide or amino acid sequence. Thus, for example, a “wild type mRNA” is an mRNA that is naturally occurring in or endogenous to the reference organism. A

“homologous” nucleic acid sequence is a nucleotide sequence naturally associated with a host cell into which it is introduced.

As used herein, the terms “nucleic acid,” “nucleic acid molecule,” “nucleotide sequence” and “polynucleotide” refer to RNA or DNA that is linear or branched, single or double stranded, or a hybrid thereof. The term also encompasses RNA/DNA hybrids. When dsRNA is produced synthetically, less common bases, such as inosine, 5-methylcytosine, 6-methyladenine, hypoxanthine and others can also be used for antisense, dsRNA, and ribozyme pairing. For example, polynucleotides that contain C-5 propyne analogues of uridine and cytidine have been shown to bind RNA with high affinity and to be potent antisense inhibitors of gene expression. Other modifications, such as modification to the phosphodiester backbone, or the 2'-hydroxy in the ribose sugar group of the RNA can also be made.

As used herein, the term “nucleotide sequence” refers to a heteropolymer of nucleotides or the sequence of these nucleotides from the 5' to 3' end of a nucleic acid molecule and includes DNA or RNA molecules, including cDNA, a DNA fragment or portion, genomic DNA, synthetic (*e.g.*, chemically synthesized) DNA, plasmid DNA, mRNA, and anti-sense RNA, any of which can be single stranded or double stranded. The terms “nucleotide sequence” “nucleic acid,” “nucleic acid molecule,” “nucleic acid construct,” “oligonucleotide” and “polynucleotide” are also used interchangeably herein to refer to a heteropolymer of nucleotides. Nucleic acid molecules and/or nucleotide sequences provided herein are presented herein in the 5' to 3' direction, from left to right and are represented using the standard code for representing the nucleotide characters as set forth in the U.S. sequence rules, 37 CFR §§1.821 - 1.825 and the World Intellectual Property Organization (WIPO) Standard ST.25. A “5' region” as used herein can mean the region of a polynucleotide that is nearest the 5' end of the polynucleotide. Thus, for example, an element in the 5' region of a polynucleotide can be located anywhere from the first nucleotide located at the 5' end of the polynucleotide to the nucleotide located halfway through the polynucleotide. A “3' region” as used herein can mean the region of a polynucleotide that is nearest the 3' end of the polynucleotide. Thus, for example, an element in the 3' region of a polynucleotide can be located anywhere from the first nucleotide located at the 3' end of the polynucleotide to the nucleotide located halfway through the polynucleotide.

As used herein, the term “gene” refers to a nucleic acid molecule capable of being used to produce mRNA, antisense RNA, miRNA, anti-microRNA antisense oligodeoxyribonucleotide (AMO) and the like. Genes may or may not be capable of being

used to produce a functional protein or gene product. Genes can include both coding and non-coding regions (e.g., introns, regulatory elements, promoters, enhancers, termination sequences and/or 5' and 3' untranslated regions). A gene may be “isolated” by which is meant a nucleic acid that is substantially or essentially free from components normally found in association with the nucleic acid in its natural state. Such components include other cellular material, culture medium from recombinant production, and/or various chemicals used in chemically synthesizing the nucleic acid.

The term “mutation” refers to point mutations (e.g., missense, or nonsense, or insertions or deletions of single base pairs that result in frame shifts), insertions, deletions, and/or truncations. When the mutation is a substitution of a residue within an amino acid sequence with another residue, or a deletion or insertion of one or more residues within a sequence, the mutations are typically described by identifying the original residue followed by the position of the residue within the sequence and by the identity of the newly substituted residue.

The terms “complementary” or “complementarity,” as used herein, refer to the natural binding of polynucleotides under permissive salt and temperature conditions by base-pairing. For example, the sequence “A-G-T” (5' to 3') binds to the complementary sequence “T-C-A” (3' to 5'). Complementarity between two single-stranded molecules may be “partial,” in which only some of the nucleotides bind, or it may be complete when total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands.

“Complement” as used herein can mean 100% complementarity with the comparator nucleotide sequence or it can mean less than 100% complementarity (e.g., about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, and the like, complementarity).

A “portion” or “fragment” of a nucleotide sequence of the invention will be understood to mean a nucleotide sequence of reduced length relative (e.g., reduced by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more nucleotides) to a reference nucleic acid or nucleotide sequence and comprising, consisting essentially of and/or consisting of a nucleotide sequence of contiguous nucleotides identical or almost identical (e.g., 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%

identical) to the reference nucleic acid or nucleotide sequence. Such a nucleic acid fragment or portion according to the invention may be, where appropriate, included in a larger polynucleotide of which it is a constituent. As an example, a repeat sequence of a guide nucleic acid of this invention may comprise a portion of a wild type Type V CRISPR-Cas repeat sequence (e.g., a wild Type CRISPR-Cas repeat, e.g., a repeat from the CRISPR Cas system of a Cas12a (Cpf1), Cas12b (C2c1), Cas12c (C2c3), Cas12d (CasY), Cas12e (CasX), Cas12g, Cas12h, Cas12i, C2c4, C2c5, C2c8, C2c9, C2c10, Cas14a, Cas14b, and/or a Cas14c, and the like). In some embodiments, a repeat sequence of a guide nucleic acid of this invention may comprise a portion of a wild type CRISPR-Cas9 repeat sequence.

Different nucleic acids or proteins having homology are referred to herein as “homologues.” The term homologue includes homologous sequences from the same and other species and orthologous sequences from the same and other species. “Homology” refers to the level of similarity between two or more nucleic acid and/or amino acid sequences in terms of percent of positional identity (*i.e.*, sequence similarity or identity).

Homology also refers to the concept of similar functional properties among different nucleic acids or proteins. Thus, the compositions and methods of the invention further comprise homologues to the nucleotide sequences and polypeptide sequences of this invention.

“Orthologous,” as used herein, refers to homologous nucleotide sequences and/ or amino acid sequences in different species that arose from a common ancestral gene during speciation. A homologue of a nucleotide sequence of this invention has a substantial sequence identity (e.g., at least about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or 100%) to said nucleotide sequence of the invention.

As used herein “sequence identity” refers to the extent to which two optimally aligned polynucleotide or polypeptide sequences are invariant throughout a window of alignment of components, *e.g.*, nucleotides or amino acids. “Identity” can be readily calculated by known methods including, but not limited to, those described in: *Computational Molecular Biology* (Lesk, A. M., ed.) Oxford University Press, New York (1988); *Biocomputing: Informatics and Genome Projects* (Smith, D. W., ed.) Academic Press, New York (1993); *Computer Analysis of Sequence Data, Part I* (Griffin, A. M., and Griffin, H. G., eds.) Humana Press, New Jersey (1994); *Sequence Analysis in Molecular Biology* (von Heinje, G., ed.) Academic Press (1987); and *Sequence Analysis Primer* (Gribskov, M. and Devereux, J., eds.) Stockton Press, New York (1991).

As used herein, the term “percent sequence identity” or “percent identity” refers to the percentage of identical nucleotides in a linear polynucleotide sequence of a reference (“query”) polynucleotide molecule (or its complementary strand) as compared to a test (“subject”) polynucleotide molecule (or its complementary strand) when the two sequences are optimally aligned. In some embodiments, “percent identity” can refer to the percentage of identical amino acids in an amino acid sequence as compared to a reference polypeptide.

As used herein, the phrase “substantially identical,” or “substantial identity” in the context of two nucleic acid molecules, nucleotide sequences or protein sequences, refers to two or more sequences or subsequences that have at least about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9% or more nucleotide or amino acid residue identity, when compared and aligned for maximum correspondence, as measured using one of the following sequence comparison algorithms or by visual inspection. In some embodiments of the invention, the substantial identity exists over a region of consecutive nucleotides of a nucleotide sequence of the invention that is about 10 nucleotides to about 20 nucleotides, about 10 nucleotides to about 25 nucleotides, about 10 nucleotides to about 30 nucleotides, about 15 nucleotides to about 25 nucleotides, about 30 nucleotides to about 40 nucleotides, about 50 nucleotides to about 60 nucleotides, about 70 nucleotides to about 80 nucleotides, about 90 nucleotides to about 100 nucleotides, or more nucleotides in length, and any range therein, up to the full length of the sequence. In some embodiments, the nucleotide sequences can be substantially identical over at least about 20 nucleotides (e.g., about 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40 nucleotides). In some embodiments, a substantially identical nucleotide or protein sequence performs substantially the same function as the nucleotide (or encoded protein sequence) to which it is substantially identical.

For sequence comparison, typically one sequence acts as a reference sequence to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

Optimal alignment of sequences for aligning a comparison window are well known to those skilled in the art and may be conducted by tools such as the local homology algorithm of Smith and Waterman, the homology alignment algorithm of Needleman and Wunsch, the

search for similarity method of Pearson and Lipman, and optionally by computerized implementations of these algorithms such as GAP, BESTFIT, FASTA, and TFASTA available as part of the GCG® Wisconsin Package® (Accelrys Inc., San Diego, CA). An “identity fraction” for aligned segments of a test sequence and a reference sequence is the
5 number of identical components which are shared by the two aligned sequences divided by the total number of components in the reference sequence segment, *e.g.*, the entire reference sequence or a smaller defined part of the reference sequence. Percent sequence identity is represented as the identity fraction multiplied by 100. The comparison of one or more polynucleotide sequences may be to a full-length polynucleotide sequence or a portion
10 thereof, or to a longer polynucleotide sequence. For purposes of this invention “percent identity” may also be determined using BLASTX version 2.0 for translated nucleotide sequences and BLASTN version 2.0 for polynucleotide sequences.

Two nucleotide sequences may also be considered substantially complementary when the two sequences hybridize to each other under stringent conditions. In some representative
15 embodiments, two nucleotide sequences considered to be substantially complementary hybridize to each other under highly stringent conditions.

“Stringent hybridization conditions” and “stringent hybridization wash conditions” in the context of nucleic acid hybridization experiments such as Southern and Northern hybridizations are sequence dependent and are different under different environmental
20 parameters. An extensive guide to the hybridization of nucleic acids is found in Tijssen *Laboratory Techniques in Biochemistry and Molecular Biology-Hybridization with Nucleic Acid Probes* part I chapter 2 “Overview of principles of hybridization and the strategy of nucleic acid probe assays” Elsevier, New York (1993). Generally, highly stringent hybridization and wash conditions are selected to be about 5°C lower than the thermal
25 melting point (T_m) for the specific sequence at a defined ionic strength and pH.

The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Very stringent conditions are selected to be equal to the T_m for a particular probe. An example of stringent hybridization conditions for hybridization of complementary nucleotide sequences which have more than
30 100 complementary residues on a filter in a Southern or northern blot is 50% formamide with 1 mg of heparin at 42°C, with the hybridization being carried out overnight. An example of highly stringent wash conditions is 0.1 5M NaCl at 72°C for about 15 minutes. An example of stringent wash conditions is a 0.2x SSC wash at 65°C for 15 minutes (*see*, Sambrook, *infra*, for a description of SSC buffer). Often, a high stringency wash is preceded by a low

stringency wash to remove background probe signal. An example of a medium stringency wash for a duplex of, e.g., more than 100 nucleotides, is 1x SSC at 45°C for 15 minutes. An example of a low stringency wash for a duplex of, e.g., more than 100 nucleotides, is 4-6x SSC at 40°C for 15 minutes. For short probes (e.g., about 10 to 50 nucleotides), stringent conditions typically involve salt concentrations of less than about 1.0 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3, and the temperature is typically at least about 30°C. Stringent conditions can also be achieved with the addition of destabilizing agents such as formamide. In general, a signal to noise ratio of 2x (or higher) than that observed for an unrelated probe in the particular hybridization assay indicates detection of a specific hybridization. Nucleotide sequences that do not hybridize to each other under stringent conditions are still substantially identical if the proteins that they encode are substantially identical. This can occur, for example, when a copy of a nucleotide sequence is created using the maximum codon degeneracy permitted by the genetic code.

The polynucleotide and/or recombinant nucleic acid constructs of this invention can be codon optimized for expression. In some embodiments, the polynucleotides, nucleic acid constructs, expression cassettes, and/or vectors of the invention (e.g., comprising/encoding a CRISPR-Cas effector protein (e.g., a Type V CRISPR-Cas effector protein), a reverse transcriptase, a flap endonuclease, a 5'-3' exonuclease, and the like) are codon optimized for expression in an organism (e.g., in a particular species), optionally an animal, a plant, a fungus, an archaeon, or a bacterium. In some embodiments, the codon optimized nucleic acid constructs, polynucleotides, expression cassettes, and/or vectors of the invention have about 70% to about 99.9% (e.g., 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5%, or 99.9%) identity or more to the nucleic acid constructs, polynucleotides, expression cassettes, and/or vectors that have not been codon optimized.

In any of the embodiments described herein, a polynucleotide or nucleic acid construct of the invention may be operatively associated with a variety of promoters and/or other regulatory elements for expression in a plant and/or a cell of a plant. Thus, in some embodiments, a polynucleotide or nucleic acid construct of this invention may further comprise one or more promoters, introns, enhancers, and/or terminators operably linked to one or more nucleotide sequences. In some embodiments, a promoter may be operably associated with an intron (e.g., Ubi1 promoter and intron). In some embodiments, a promoter associated with an intron may be referred to as a "promoter region" (e.g., Ubi1 promoter and intron).

By “operably linked” or “operably associated” as used herein in reference to polynucleotides, it is meant that the indicated elements are functionally related to each other and are also generally physically related. Thus, the term “operably linked” or “operably associated” as used herein, refers to nucleotide sequences on a single nucleic acid molecule that are functionally associated. Thus, a first nucleotide sequence that is operably linked to a second nucleotide sequence means a situation when the first nucleotide sequence is placed in a functional relationship with the second nucleotide sequence. For instance, a promoter is operably associated with a nucleotide sequence if the promoter effects the transcription or expression of said nucleotide sequence. Those skilled in the art will appreciate that the control sequences (*e.g.*, promoter) need not be contiguous with the nucleotide sequence to which it is operably associated, as long as the control sequences function to direct the expression thereof. Thus, for example, intervening untranslated, yet transcribed, nucleic acid sequences can be present between a promoter and the nucleotide sequence, and the promoter can still be considered “operably linked” to the nucleotide sequence.

As used herein, the term “linked,” in reference to polypeptides, refers to the attachment of one polypeptide to another. A polypeptide may be linked to another polypeptide (at the N-terminus or the C-terminus) directly (*e.g.*, via a peptide bond) or through a linker.

The term “linker” is art-recognized and refers to a chemical group, or a molecule linking two molecules or moieties, *e.g.*, two domains of a fusion protein, such as, for example, a DNA binding polypeptide or domain and peptide tag and/or a reverse transcriptase and an affinity polypeptide that binds to the peptide tag; or a DNA endonuclease polypeptide or domain and peptide tag and/or a reverse transcriptase and an affinity polypeptide that binds to the peptide tag. A linker may be comprised of a single linking molecule or may comprise more than one linking molecule. In some embodiments, the linker can be an organic molecule, group, polymer, or chemical moiety such as a bivalent organic moiety. In some embodiments, the linker may be an amino acid, or it may be a peptide. In some embodiments, the linker is a peptide.

In some embodiments, a peptide linker useful with this invention may be about 2 to about 100 or more amino acids in length, for example, about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 or more amino acids in length (*e.g.*, about 2

to about 40, about 2 to about 50, about 2 to about 60, about 4 to about 40, about 4 to about 50, about 4 to about 60, about 5 to about 40, about 5 to about 50, about 5 to about 60, about 9 to about 40, about 9 to about 50, about 9 to about 60, about 10 to about 40, about 10 to about 50, about 10 to about 60, or about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 amino acids to about 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 or more amino acids in length (e.g., about 105, 110, 115, 120, 130, 140 150 or more amino acids in length). In some embodiments, a peptide linker may be a GS linker.

As used herein, the term "linked," or "fused" in reference to polynucleotides, refers to the attachment of one polynucleotide to another. In some embodiments, two or more polynucleotide molecules may be linked by a linker that can be an organic molecule, group, polymer, or chemical moiety such as a bivalent organic moiety. A polynucleotide may be linked or fused to another polynucleotide (at the 5' end or the 3' end) via a covalent or non-covalent linkage or binding, including e.g., Watson-Crick base-pairing, or through one or more linking nucleotides. In some embodiments, a polynucleotide motif of a certain structure may be inserted within another polynucleotide sequence (e.g., extension of the hairpin structure in guide RNA). In some embodiments, the linking nucleotides may be naturally occurring nucleotides. In some embodiments, the linking nucleotides may be non-naturally occurring nucleotides.

A "promoter" is a nucleotide sequence that controls or regulates the transcription of a nucleotide sequence (e.g., a coding sequence) that is operably associated with the promoter. The coding sequence controlled or regulated by a promoter may encode a polypeptide and/or a functional RNA. Typically, a "promoter" refers to a nucleotide sequence that contains a binding site for RNA polymerase II and directs the initiation of transcription. In general, promoters are found 5', or upstream, relative to the start of the coding region of the corresponding coding sequence. A promoter may comprise other elements that act as regulators of gene expression; e.g., a promoter region. These include a TATA box consensus sequence, and often a CAAT box consensus sequence (Breathnach and Chambon, (1981) *Annu. Rev. Biochem.* 50:349). In plants, the CAAT box may be substituted by the AGGA box (Messing *et al.*, (1983) in *Genetic Engineering of Plants*, T. Kosuge, C. Meredith and A. Hollaender (eds.), Plenum Press, pp. 211-227). In some embodiments, a promoter region may comprise at least one intron (see, e.g., **SEQ ID NO:21**, **SEQ ID NO:22**).

Promoters useful with this invention can include, for example, constitutive, inducible, temporally regulated, developmentally regulated, chemically regulated, tissue-preferred and/or tissue-specific promoters for use in the preparation of recombinant nucleic acid molecules, e.g., “synthetic nucleic acid constructs” or “protein-RNA complex.” These
5 various types of promoters are known in the art.

The choice of promoter may vary depending on the temporal and spatial requirements for expression, and also may vary based on the host cell to be transformed. Promoters for many different organisms are well known in the art. Based on the extensive knowledge present in the art, the appropriate promoter can be selected for the particular host organism of
10 interest. Thus, for example, much is known about promoters upstream of highly constitutively expressed genes in model organisms and such knowledge can be readily accessed and implemented in other systems as appropriate.

In some embodiments, a promoter functional in a plant may be used with the constructs of this invention. Non-limiting examples of a promoter useful for driving
15 expression in a plant include the promoter of the RubisCo small subunit gene 1 (PrbcS1), the promoter of the actin gene (Pactin), the promoter of the nitrate reductase gene (Pnr) and the promoter of duplicated carbonic anhydrase gene 1 (Pdca1) (*See, Walker et al. Plant Cell Rep.* 23:727-735 (2005); Li et al. *Gene* 403:132-142 (2007); Li et al. *Mol Biol. Rep.* 37:1143-1154 (2010)). PrbcS1 and Pactin are constitutive promoters and Pnr and Pdca1 are inducible
20 promoters. Pnr is induced by nitrate and repressed by ammonium (Li et al. *Gene* 403:132-142 (2007)) and Pdca1 is induced by salt (Li et al. *Mol Biol. Rep.* 37:1143-1154 (2010)). In some embodiments, a promoter useful with this invention is RNA polymerase II (Pol II) promoter. In some embodiments, a U6 promoter or a 7SL promoter from *Zea mays* may be useful with constructs of this invention. In some embodiments, the U6c promoter and/or 7SL
25 promoter from *Zea mays* may be useful for driving expression of a guide nucleic acid. In some embodiments, a U6c promoter, U6i promoter and/or 7SL promoter from *Glycine max* may be useful with constructs of this invention. In some embodiments, the U6c promoter, U6i promoter and/or 7SL promoter from *Glycine max* may be useful for driving expression of a guide nucleic acid.

30 Examples of constitutive promoters useful for plants include, but are not limited to, cetrum virus promoter (cmp) (U.S. Patent No. 7,166,770), the rice actin 1 promoter (Wang et al. (1992) *Mol. Cell. Biol.* 12:3399-3406; as well as US Patent No. 5,641,876), CaMV 35S promoter (Odell et al. (1985) *Nature* 313:810-812), CaMV 19S promoter (Lawton et al. (1987) *Plant Mol. Biol.* 9:315-324), nos promoter (Ebert et al. (1987) *Proc. Natl. Acad. Sci*

USA 84:5745-5749), Adh promoter (Walker *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:6624-6629), sucrose synthase promoter (Yang & Russell (1990) *Proc. Natl. Acad. Sci. USA* 87:4144-4148), and the ubiquitin promoter. The constitutive promoter derived from ubiquitin accumulates in many cell types. Ubiquitin promoters have been cloned from

5 several plant species for use in transgenic plants, for example, sunflower (Binet *et al.*, 1991. *Plant Science* 79: 87-94), maize (Christensen *et al.*, 1989. *Plant Molec. Biol.* 12: 619-632), and arabidopsis (Norris *et al.* 1993. *Plant Molec. Biol.* 21:895-906). The maize ubiquitin promoter (*UbiP*) has been developed in transgenic monocot systems and its sequence and vectors constructed for monocot transformation are disclosed in the patent publication EP 0

10 342 926. The ubiquitin promoter is suitable for the expression of the nucleotide sequences of the invention in transgenic plants, especially monocotyledons. Further, the promoter expression cassettes described by McElroy *et al.* (*Mol. Gen. Genet.* 231: 150-160 (1991)) can be easily modified for the expression of the nucleotide sequences of the invention and are particularly suitable for use in monocotyledonous hosts.

15 In some embodiments, tissue specific/tissue preferred promoters can be used for expression of a heterologous polynucleotide in a plant cell. Tissue specific or preferred expression patterns include, but are not limited to, green tissue specific or preferred, root specific or preferred, stem specific or preferred, flower specific or preferred or pollen specific or preferred. Promoters suitable for expression in green tissue include many that regulate

20 genes involved in photosynthesis and many of these have been cloned from both monocotyledons and dicotyledons. In one embodiment, a promoter useful with the invention is the maize PEPC promoter from the phosphoenol carboxylase gene (Hudspeth & Grula, *Plant Molec. Biol.* 12:579-589 (1989)). Non-limiting examples of tissue-specific promoters include those associated with genes encoding the seed storage proteins (such as β -

25 conglycinin, cruciferin, napin and phaseolin), zein or oil body proteins (such as oleosin), or proteins involved in fatty acid biosynthesis (including acyl carrier protein, stearoyl-ACP desaturase and fatty acid desaturases (fad 2-1)), and other nucleic acids expressed during embryo development (such as Bce4, *see, e.g.*, Kridl *et al.* (1991) *Seed Sci. Res.* 1:209-219; as well as EP Patent No. 255378). Tissue-specific or tissue-preferential promoters useful for the

30 expression of the nucleotide sequences of the invention in plants, particularly maize, include but are not limited to those that direct expression in root, pith, leaf or pollen. Such promoters are disclosed, for example, in WO 93/07278, herein incorporated by reference in its entirety. Other non-limiting examples of tissue specific or tissue preferred promoters useful with the invention the cotton rubisco promoter disclosed in US Patent 6,040,504; the rice sucrose

synthase promoter disclosed in US Patent 5,604,121; the root specific promoter described by de Framond (FEBS 290:103-106 (1991); EP 0 452 269 to Ciba-Geigy); the stem specific promoter described in U.S. Patent 5,625,136 (to Ciba-Geigy) and which drives expression of the maize *trpA* gene; the cestrum yellow leaf curling virus promoter disclosed in WO 01/73087; and pollen specific or preferred promoters including, but not limited to, ProOsLPS10 and ProOsLPS11 from rice (Nguyen et al. *Plant Biotechnol. Reports* 9(5):297-306 (2015)), ZmSTK2_USP from maize (Wang et al. *Genome* 60(6):485-495 (2017)), LAT52 and LAT59 from tomato (Twell et al. *Development* 109(3):705-713 (1990)), Zm13 (U.S. Patent No. 10,421,972), PLA2- δ promoter from arabidopsis (U.S. Patent No. 7,141,424), and/or the ZmC5 promoter from maize (International PCT Publication No. WO1999/042587).

Additional examples of plant tissue-specific/tissue preferred promoters include, but are not limited to, the root hair-specific *cis*-elements (RHEs) (Kim et al. *The Plant Cell* 18:2958-2970 (2006)), the root-specific promoters *RCc3* (Jeong et al. *Plant Physiol.* 153:185-197 (2010)) and RB7 (U.S. Patent No. 5459252), the lectin promoter (Lindstrom et al. (1990) *Der. Genet.* 11:160-167; and Vodkin (1983) *Prog. Clin. Biol. Res.* 138:87-98), corn alcohol dehydrogenase 1 promoter (Dennis et al. (1984) *Nucleic Acids Res.* 12:3983-4000), S-adenosyl-L-methionine synthetase (SAMS) (Vander Mijnsbrugge et al. (1996) *Plant and Cell Physiology*, 37(8):1108-1115), corn light harvesting complex promoter (Bansal et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:3654-3658), corn heat shock protein promoter (O'Dell et al. (1985) *EMBO J.* 5:451-458; and Rochester et al. (1986) *EMBO J.* 5:451-458), pea small subunit RuBP carboxylase promoter (Cashmore, "Nuclear genes encoding the small subunit of ribulose-1,5-bisphosphate carboxylase" pp. 29-39 In: *Genetic Engineering of Plants* (Hollaender ed., Plenum Press 1983; and Poulsen et al. (1986) *Mol. Gen. Genet.* 205:193-200), Ti plasmid mannopine synthase promoter (Langridge et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3219-3223), Ti plasmid nopaline synthase promoter (Langridge et al. (1989), *supra*), petunia chalcone isomerase promoter (van Tunen et al. (1988) *EMBO J.* 7:1257-1263), bean glycine rich protein 1 promoter (Keller et al. (1989) *Genes Dev.* 3:1639-1646), truncated CaMV 35S promoter (O'Dell et al. (1985) *Nature* 313:810-812), potato patatin promoter (Wenzler et al. (1989) *Plant Mol. Biol.* 13:347-354), root cell promoter (Yamamoto et al. (1990) *Nucleic Acids Res.* 18:7449), maize zein promoter (Kriz et al. (1987) *Mol. Gen. Genet.* 207:90-98; Langridge et al. (1983) *Cell* 34:1015-1022; Reina et al. (1990) *Nucleic Acids Res.* 18:6425; Reina et al. (1990) *Nucleic Acids Res.* 18:7449; and Wandelt et al. (1989) *Nucleic Acids Res.* 17:2354), globulin-1 promoter (Belanger et al. (1991) *Genetics*

129:863-872), α -tubulin cab promoter (Sullivan *et al.* (1989) *Mol. Gen. Genet.* 215:431-440), PEPCase promoter (Hudspeth & Grula (1989) *Plant Mol. Biol.* 12:579-589), R gene complex-associated promoters (Chandler *et al.* (1989) *Plant Cell* 1:1175-1183), and chalcone synthase promoters (Franken *et al.* (1991) *EMBO J.* 10:2605-2612).

5 Useful for seed-specific expression is the pea vicilin promoter (Czako *et al.* (1992) *Mol. Gen. Genet.* 235:33-40; as well as the seed-specific promoters disclosed in U.S. Patent No. 5,625,136. Useful promoters for expression in mature leaves are those that are switched at the onset of senescence, such as the SAG promoter from *Arabidopsis* (Gan *et al.* (1995) *Science* 270:1986-1988).

10 In addition, promoters functional in chloroplasts can be used. Non-limiting examples of such promoters include the bacteriophage T3 gene 9 5' UTR and other promoters disclosed in U.S. Patent No. 7,579,516. Other promoters useful with the invention include but are not limited to the S-E9 small subunit RuBP carboxylase promoter and the Kunitz trypsin inhibitor gene promoter (Kti3).

15 Additional regulatory elements useful with this invention include, but are not limited to, introns, enhancers, termination sequences and/or 5' and 3' untranslated regions.

 An intron useful with this invention can be an intron identified in and isolated from a plant and then inserted into an expression cassette to be used in transformation of a plant. As would be understood by those of skill in the art, introns can comprise the sequences required
20 for self-excision and are incorporated into nucleic acid constructs/expression cassettes in frame. An intron can be used either as a spacer to separate multiple protein-coding sequences in one nucleic acid construct, or an intron can be used inside one protein-coding sequence to, for example, stabilize the mRNA. If they are used within a protein-coding sequence, they are inserted "in-frame" with the excision sites included. Introns may also be associated with
25 promoters to improve or modify expression. As an example, a promoter/intron combination useful with this invention includes but is not limited to that of the maize Ubi1 promoter and intron.

 Non-limiting examples of introns useful with the present invention include introns from the ADHI gene (e.g., *Adh1-S* introns 1, 2 and 6), the ubiquitin gene (Ubi1), the
30 RuBisCO small subunit (rbcS) gene, the RuBisCO large subunit (rbcL) gene, the actin gene (e.g., *actin-1* intron), the pyruvate dehydrogenase kinase gene (pdk), the nitrate reductase gene (nr), the duplicated carbonic anhydrase gene 1 (Tdca1), the psbA gene, the atpA gene, or any combination thereof. Example intron sequences can include, but are not limited to,
SEQ ID NO:61 and **SEQ ID NO:62**.

In some embodiments, a polynucleotide and/or a nucleic acid construct of the invention can be an “expression cassette” or can be comprised within an expression cassette. As used herein, “expression cassette” means a recombinant nucleic acid molecule comprising, for example, a nucleic acid construct of the invention (e.g., a CRISPR-Cas effector protein, a reverse transcriptase polypeptide or domain, a flap endonuclease polypeptide or domain (e.g., FEN)), and/or a 5'-3' exonuclease), wherein the nucleic acid construct is operably associated with at one or more control sequences (e.g., a promoter, terminator and the like). Thus, some embodiments of the invention provide expression cassettes designed to express, for example, a nucleic acid construct of the invention (e.g., a nucleic acid construct of the invention encoding a CRISPR-Cas effector protein or domain, a reverse transcriptase polypeptide or domain, a flap endonuclease polypeptide or domain and/or 5'-3' exonuclease polypeptide or domain. When an expression cassette of the present invention comprises more than one polynucleotide, the polynucleotides may be operably linked to a single promoter that drives expression of all of the polynucleotides or the polynucleotides may be operably linked to one or more separate promoters (e.g., three polynucleotides may be driven by one, two or three promoters in any combination). When two or more separate promoters are used, the promoters may be the same promoter, or they may be different promoters. Thus, a polynucleotide encoding a CRISPR-Cas effector protein or domain, a polynucleotide encoding a reverse transcriptase polypeptide or domain, a polynucleotide encoding a flap endonuclease polypeptide or domain and/or a polynucleotide encoding a 5'-3' exonuclease polypeptide or domain comprised in an expression cassette may each be operably linked to a separate promoter, or they may be operably linked to two or more promoters in any combination.

An expression cassette comprising a nucleic acid construct of the invention may be chimeric, meaning that at least one of its components is heterologous with respect to at least one of its other components (e.g., a promoter from the host organism operably linked to a polynucleotide of interest to be expressed in the host organism, wherein the polynucleotide of interest is from a different organism than the host or is not normally found in association with that promoter). An expression cassette may also be one that is naturally occurring but has been obtained in a recombinant form useful for heterologous expression.

An expression cassette can optionally include a transcriptional and/or translational termination region (*i.e.*, termination region) and/or an enhancer region that is functional in the selected host cell. A variety of transcriptional terminators and enhancers are known in the art and are available for use in expression cassettes. Transcriptional terminators are responsible

for the termination of transcription and correct mRNA polyadenylation. A termination region and/or the enhancer region may be native to the transcriptional initiation region, may be native, for example, to a gene encoding a CRISPR-Cas effector protein, a gene encoding a reverse transcriptase, a gene encoding a flap endonuclease, and/or a gene encoding a 5'-3' exonuclease, may be native to a host cell, or may be native to another source (*e.g.*, foreign or heterologous to the promoter, to a gene encoding a CRISPR-Cas effector protein, a gene encoding a reverse transcriptase, a gene encoding a flap endonuclease, and/or a gene encoding a 5'-3' exonuclease, to the host cell, or any combination thereof).

An expression cassette of the invention also can include a polynucleotide encoding a selectable marker, which can be used to select a transformed host cell. As used herein, "selectable marker" means a polynucleotide sequence that when expressed imparts a distinct phenotype to the host cell expressing the marker and thus allows such transformed cells to be distinguished from those that do not have the marker. Such a polynucleotide sequence may encode either a selectable or screenable marker, depending on whether the marker confers a trait that can be selected for by chemical means, such as by using a selective agent (*e.g.*, an antibiotic and the like), or on whether the marker is simply a trait that one can identify through observation or testing, such as by screening (*e.g.*, fluorescence). Many examples of suitable selectable markers are known in the art and can be used in the expression cassettes described herein.

In addition to expression cassettes, the nucleic acid molecules/constructs and polynucleotide sequences described herein can be used in connection with vectors. The term "vector" refers to a composition for transferring, delivering, or introducing a nucleic acid (or nucleic acids) into a cell. A vector comprises a nucleic acid construct comprising the nucleotide sequence(s) to be transferred, delivered, or introduced. Vectors for use in transformation of host organisms are well known in the art. Non-limiting examples of general classes of vectors include viral vectors, plasmid vectors, phage vectors, phagemid vectors, cosmid vectors, fosmid vectors, bacteriophages, artificial chromosomes, minicircles, or *Agrobacterium* binary vectors in double or single stranded linear or circular form which may or may not be self-transmissible or mobilizable. In some embodiments, a viral vector can include, but is not limited, to a retroviral, lentiviral, adenoviral, adeno-associated, or herpes simplex viral vector. A vector as defined herein can transform a prokaryotic or eukaryotic host either by integration into the cellular genome or exist extrachromosomally (*e.g.*, autonomous replicating plasmid with an origin of replication). Additionally included are shuttle vectors by which is meant a DNA vehicle capable, naturally or by design, of

replication in two different host organisms, which may be selected from actinomycetes and related species, bacteria and eukaryotic (e.g., higher plant, mammalian, yeast or fungal cells). In some embodiments, the nucleic acid in the vector is under the control of, and operably linked to, an appropriate promoter or other regulatory elements for transcription in a host cell.

- 5 The vector may be a bi-functional expression vector which functions in multiple hosts. In the case of genomic DNA, this may contain its own promoter and/or other regulatory elements and in the case of cDNA this may be under the control of an appropriate promoter and/or other regulatory elements for expression in the host cell. Accordingly, a nucleic acid construct or polynucleotide of this invention and/or expression cassettes comprising the same
10 may be comprised in vectors as described herein and as known in the art.

- As used herein, “contact,” “contacting,” “contacted,” and grammatical variations thereof, refer to placing the components of a desired reaction together under conditions suitable for carrying out the desired reaction (e.g., transformation, transcriptional control, genome editing, nicking, and/or cleavage). As an example, a target nucleic acid may be
15 contacted with a Type II or Type V CRISPR-Cas effector protein, and a reverse transcriptase or a nucleic acid construct encoding the same, under conditions whereby the CRISPR-Cas effector protein and the reverse transcriptase are expressed and the CRISPR-Cas effector protein binds to the target nucleic acid, and the reverse transcriptase is either fused to the CRISPR-Cas effector protein or is recruited to the CRISPR-Cas effector protein (via, for
20 example, a peptide tag fused to the CRISPR-Cas effector protein and an affinity tag fused to the reverse transcriptase) and thus, the reverse transcriptase is positioned in the vicinity of the target nucleic acid, thereby modifying the target nucleic acid. Other methods for recruiting a reverse transcriptase may be used that take advantage of other protein-protein interactions, and also RNA-protein interactions and chemical interactions.

- 25 As used herein, “modifying” or “modification” in reference to a target nucleic acid includes editing (e.g., mutating), covalent modification, exchanging/substituting nucleic acids/nucleotide bases, deleting, cleaving, nicking, and/or transcriptional control of a target nucleic acid. In some embodiments, a modification may include an indel of any size and/or a single base change (SNP) of any type.

- 30 “Introducing,” “introduce,” “introduced” (and grammatical variations thereof) in the context of a polynucleotide of interest means presenting a nucleotide sequence of interest (e.g., polynucleotide, a nucleic acid construct, and/or a guide nucleic acid) to a host organism or cell of said organism (e.g., host cell, e.g., a plant cell) in such a manner that the nucleotide sequence gains access to the interior of a cell.

The terms “transformation” or transfection” may be used interchangeably and as used herein refer to the introduction of a heterologous nucleic acid into a cell. Transformation of a cell may be stable or transient. Thus, in some embodiments, a host cell or host organism may be stably transformed with a polynucleotide/nucleic acid molecule of the invention. In some
5 embodiments, a host cell or host organism may be transiently transformed with a nucleic acid construct of the invention.

“Transient transformation” in the context of a polynucleotide means that a polynucleotide is introduced into the cell and does not integrate into the genome of the cell.

By “stably introducing” or “stably introduced” in the context of a polynucleotide
10 introduced into a cell is intended that the introduced polynucleotide is stably incorporated into the genome of the cell, and thus the cell is stably transformed with the polynucleotide.

“Stable transformation” or “stably transformed” as used herein means that a nucleic acid molecule is introduced into a cell and integrates into the genome of the cell. As such, the integrated nucleic acid molecule is capable of being inherited by the progeny thereof,
15 more particularly, by the progeny of multiple successive generations. “Genome” as used herein includes the nuclear, mitochondrial and the plastid genomes, and therefore includes integration of the nucleic acid into, for example, the chloroplast or mitochondrial genome. Stable transformation as used herein can also refer to a transgene that is maintained extrachromasomally, for example, as a minichromosome or a plasmid.

20 Transient transformation may be detected by, for example, an enzyme-linked immunosorbent assay (ELISA) or Western blot, which can detect the presence of a peptide or polypeptide encoded by one or more transgene introduced into an organism. Stable transformation of a cell can be detected by, for example, a Southern blot hybridization assay of genomic DNA of the cell with nucleic acid sequences which specifically hybridize with a
25 nucleotide sequence of a transgene introduced into an organism (e.g., a plant). Stable transformation of a cell can be detected by, for example, a Northern blot hybridization assay of RNA of the cell with nucleic acid sequences which specifically hybridize with a nucleotide sequence of a transgene introduced into a host organism. Stable transformation of a cell can also be detected by, *e.g.*, a polymerase chain reaction (PCR) or other amplification reactions
30 as are well known in the art, employing specific primer sequences that hybridize with target sequence(s) of a transgene, resulting in amplification of the transgene sequence, which can be detected according to standard methods. Transformation can also be detected by direct sequencing and/or hybridization protocols well known in the art.

Accordingly, in some embodiments, nucleotide sequences, polynucleotides, nucleic acid constructs, and/or expression cassettes of the invention may be expressed transiently and/or they can be stably incorporated into the genome of the host organism. Thus, in some embodiments, a nucleic acid construct of the invention (e.g., one or more expression cassettes
5 encoding a DNA binding polypeptide or domain, an endonuclease polypeptide or domain, a reverse transcriptase polypeptide or domain, a flap endonuclease polypeptide or domain and/or nucleic acid modifying polypeptide or domain) may be transiently introduced into a cell with a guide nucleic acid and as such, no DNA maintained in the cell.

A nucleic acid construct of the invention can be introduced into a cell by any method
10 known to those of skill in the art. In some embodiments of the invention, transformation of a cell comprises nuclear transformation. In other embodiments, transformation of a cell comprises plastid transformation (e.g., chloroplast transformation). In still further embodiments, the recombinant nucleic acid construct of the invention can be introduced into a cell via conventional breeding techniques.

15 Procedures for transforming both eukaryotic and prokaryotic organisms are well known and routine in the art and are described throughout the literature (*See*, for example, Jiang et al. 2013. *Nat. Biotechnol.* 31:233-239; Ran et al. *Nature Protocols* 8:2281-2308 (2013)).

A nucleotide sequence therefore can be introduced into a host organism or its cell in
20 any number of ways that are well known in the art. The methods of the invention do not depend on a particular method for introducing one or more nucleotide sequences into the organism, only that they gain access to the interior of at least one cell of the organism. Where more than one nucleotide sequence is to be introduced, they can be assembled as part of a single nucleic acid construct, or as separate nucleic acid constructs, and can be located on
25 the same or different nucleic acid constructs. Accordingly, the nucleotide sequences can be introduced into the cell of interest in a single transformation event, and/or in separate transformation events, or, alternatively, where relevant, a nucleotide sequence can be incorporated into a plant, for example, as part of a breeding protocol.

Base editing has been shown to be an efficient way to change cytosine and adenine
30 residues to thymine and guanine, respectively. These tools, while powerful, do have some limitations such as bystander bases, small base editing windows, and limited PAMs.

To perform precise templated editing in cells there are several essential steps, each of which has rate limitations that together can severely hamper the ability to effectively perform editing due to low efficiencies. For example, one step requires inducing the cell to initiate a

repair event at the target site. This is typically performed by causing a double-strand break (DSB) or nick by an exogenously provided, sequence-specific nuclease or nickase. Another step requires local availability of a homologous template to be used for the repair. This step requires the template to be in the proximity of the DSB at exactly the right time when the

5 DSB is competent to commit to a templated editing pathway. In particular, this step is widely regarded to be the rate limiting step with current editing technologies. A further step is the efficient incorporation of sequence from the template into the broken or nicked target. Prior to the present invention, this step was typically provided by the cell's endogenous DNA repair enzymes. The efficiency of this step is low and difficult to manipulate. The present

10 invention bypasses many of the major obstacles to the efficiency of the process of templated editing by co-localizing, in a coordinate fashion, the functionalities required to carry out the steps described above.

Fig. 1 shows the generation of DNA sequences from reverse transcription off the crRNA and subsequent integration into the nick site using methods and constructs of the

15 present invention. An extended crRNA is shown in blue and is bound to the second strand nickase Cpf1 (Cas12a) (nCpf1, upper left). As described in more detail herein, the nCpf1 may be either covalently linked via, for example, a peptide to a reverse transcriptase (RT) or the RT may be recruited to the nCpf1 (e.g., via the use of a peptide tag motif/affinity polypeptide that binds to the peptide tag or via chemical interactions as described herein), in

20 which case multiple reverse transcriptase proteins (RTⁿ) may be recruited. The 3' end of the guide RNA is complimentary to the DNA at the nick site (non-bold pairing lines, upper left). The RT then polymerizes DNA from the 3' end of the DNA nick generating a DNA sequence complimentary to the RNA with nucleotides non-complimentary to the genome (bold pairing lines, brackets, upper right) followed by complimentary nucleotides (non-bold pairing lines, upper right). Upon dissociation, the resultant DNA has an extended ssDNA with a 3'

25 overhang which is largely the same sequence as the original DNA (non-bold pairing lines, lower right) but with some non-native nucleotides (bold pairing lines, brackets, lower right). This flap is in equilibrium with a structure having a 5' overhang (lower left) where there are mismatched nucleotides incorporated into the DNA. This equilibrium lies more to the

30 favorable perfect pairing on the right but can be driven may be reduced in a variety of ways including, for example, nicking the second strand (e.g., non-target strand or top strand). The structure on the left may be preferentially cleaved by cellular flap endonucleases involved in DNA lagging strand synthesis, which are highly conserved between mammalian and plant cells (the amino acid sequence of *Homo sapiens* FEN1 is over 50% identical to both *Zea*

mays and *Glycine max* FEN1). In some embodiments, a flap endonuclease may be introduced to drive the equilibrium in the direction of the 3' flap comprising the non-native/mismatched nucleotides. Longer 5' flaps are often removed in eukaryotic cells by the Dna2 protein, again driving the equilibrium to the 3' flap (desired) product (see, e.g., *Nucleic Acids Res.* 2012 Aug;40(14):6774-86).

Further in the process of the present invention, and as exemplified in **Fig. 2**, to reduce mismatch repair and to drive the equilibrium more in favor of forming the final product with the modified nucleotides (bold, brackets), a Cpf1 nickase may be targeted to regions outside of the RT-editing region (lightning bolts) as described herein. The nCpf1:crRNA molecules may be on either side or both sides of the editing bubble. Nicking the first strand (e.g., target strand or bottom strand of **Fig. 2**) (dashed line) indicates to the cell that the newly incorporated nucleotides are the correct nucleotides during mismatch repair and replication, thus favoring a final product with the new nucleotides.

Variants of the reverse transcriptase (RT) enzyme can have significant effects on the temperature-sensitivity and processivity of the editing system. Natural and rationally- and non-rationally engineered (i.e., directed evolution) variants of the RT can be useful in optimizing activity in plant-preferred temperatures and for optimizing processivity profiles.

Protein domain fusions to an RT polypeptide can have significant effects on the temperature-sensitivity and processivity of the editing system. The RT enzyme can be improved for temperature-sensitivity, processivity, and template affinity through fusions to ssRNA binding domains (RBDs). These RBDs may have sequence specificity, non-specificity or sequence preferences (see, e.g., **SEQ ID NOs:37-52**). A range of affinity distributions may be beneficial to editing in different cellular and *in vitro* environments. RBDs can be modified in both specificity and binding free energy through increasing or decreasing the size of the RBD in order to recognize more or fewer nucleotides. Multiple RBDs result in proteins with affinity distributions that are a combination of the individual RBDs. Adding one or more RBD to the RT enzyme can result in increased affinity, increased or decreased sequence specificity, and/or promote cooperativity.

An RT polypeptide for use with this invention may be fused with a single-stranded RNA binding protein (RBD). An RBD useful with this invention may be an RBD obtained from, for example, a human, a mouse or a fly. A single-stranded binding protein can comprise an amino acid sequence that includes, but is not limited to, any one of **SEQ ID NOs:37-52**.

After reverse transcriptase incorporates an edit into the genome, a sequence redundancy exists between the newly synthesized edited sequence and the original WT sequence it is intended to replace. This leads to either a 5' or 3' flap at the target site, which has to be repaired by the cell. The two states exist in equilibrium with binding energy favoring the 3' flap because more base pairs are available when the WT sequence is paired with its complement than when the edited strand is paired with its complement. This is unfavorable for efficient editing because processing (removal) of the 3' flap may remove the edited residues and revert the target back to WT sequence. However, cellular flap endonucleases such as FEN1 or Dna2 can efficiently process 5' flaps. Thus, instead of relying on the function of 5'- flap endonucleases native to the cell, in some embodiments of this invention the concentration of flap endonucleases at the target may be increased to further favor the desirable equilibrium outcome (removal of the WT sequence in the 5' flap so that the edited sequence becomes stably incorporated at the target site). This may be achieved by overexpression of a 5' flap endonuclease as a free protein in the cell.

Alternatively, FEN or Dna2 may be actively recruited to the target site by association with the CRISPR complex, either by direct protein fusion or by non-covalent recruitment such as with a peptide tag and affinity polypeptide pair (e.g., a SunTag antibody/epitope pair) or chemical interactions as described herein.

The present invention further provides method for modifying a target nucleic acid using the proteins/polypeptides, and/or fusion proteins of the invention and polynucleotides and nucleic acid constructs encoding the same, and/or expression cassettes and/or vectors comprising the same. The methods may be carried out in an *in vivo* system (e.g., in a cell or in an organism) or in an *in vitro* system (e.g., cell free). Thus, in some embodiments, a method of modifying a target nucleic acid in a plant cell is provided, the method comprising: contacting the target nucleic acid with (a) a Type V CRISPR-Cas effector protein or a Type II CRISPR-Cas effector protein; (b) a reverse transcriptase, and (c) an extended guide nucleic acid (e.g., extended Type II or Type V CRISPR RNA, extended Type II or Type V CRISPR DNA, extended Type II or Type V crRNA, extended Type II or Type V crDNA; e.g., tagRNA, tagDNA), thereby modifying the target nucleic acid. In some embodiments, the Type V CRISPR-Cas effector protein or Type II CRISPR-Cas effector protein, the reverse transcriptase, and the extended guide nucleic acid may form a complex or may be comprised in a complex, which is capable of interacting with the target nucleic acid. In some embodiments, the method of the invention may further comprise contacting the target nucleic acid with: (a) a second Type V CRISPR-Cas effector protein or a second Type II CRISPR-

Cas effector protein; (b) a second reverse transcriptase, and (c) a second extended guide nucleic acid (e.g., extended CRISPR RNA, extended CRISPR DNA, extended crRNA, extended crDNA; e.g., tagDNA, tagRNA), wherein the second extended guide nucleic acid targets (spacer is substantially complementary to/binds to) a site on the first strand of the target nucleic acid, thereby modifying the target nucleic acid. In some embodiments, the method of the invention may further comprise contacting the target nucleic acid with: (a) a second Type V CRISPR-Cas effector protein or a second Type II CRISPR-Cas effector protein; (b) a second reverse transcriptase, and (c) a second extended guide nucleic acid (e.g., extended CRISPR RNA, extended CRISPR DNA, extended crRNA, extended crDNA; e.g., tagDNA, tagRNA), wherein the second extended guide nucleic acid targets (spacer is substantially complementary to/binds to) a site on the second strand of the target nucleic acid, thereby modifying the target nucleic acid. In some embodiments, the methods of the invention comprise contacting the target nucleic acid at a temperature of about 20°C to 42°C (e.g., about 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, or 42°C, and any value or range therein). In some embodiments, a target nucleic acid may be contacted with additional polypeptides and/or nucleic acid constructs encoding the same in order to improve mismatch repair. In some embodiments, a method of the invention may further comprise contacting the target nucleic acid with (a) a CRISPR-Cas effector protein; and (b) a guide nucleic acid, wherein (i) the CRISPR-Cas effector protein is a nickase (e.g., nCas9, nCas12a) and nicks a site on the first strand of the target nucleic acid that is located about 10 to about 125 base pairs (e.g., about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, or 125 base pairs, or any range or value therein) that is either 5' or 3' from a site on the second strand that has been nicked by the Type II or Type V CRISPR-Cas effector protein, or (ii) the CRISPR-Cas effector protein is a nickase (e.g., nCas9, nCas12a) and nicks a site on the second strand of the target nucleic acid that is located about 10 to about 125 base pairs (either 5' or 3') from a site on the first strand that has been nicked by the Type II or Type V CRISPR-Cas effector protein, thereby improving mismatch repair. In some embodiments, nicking the second strand (non-target strand) of the target nucleic acid comprises contacting the target nucleic acid with a crRNA comprising a spacer having mismatches (e.g., about 1, 2, 3, or 4 mismatches; e.g., about 80-96% complementary

to the second strand (non-target strand)). In this configuration, the nicking does not alter the equilibrium but rather mismatch product species formation will be favored because 5' flap will be preferentially processed in the cell. Nicking of non-target strand enhances the resolution of the next step, which is when the mismatch repair is involved.

5 Thus, in some embodiments, at least two RNAs may be utilized with the methods of the invention: a tagRNA which guides the CRISPR-Cas effector protein to the right spot and makes a double-strand break using a perfect RNA:DNA match and a second RNA (crRNA) which anneals to the DNA very close by on the same strand. This second RNA (crRNA) has a spacer sequence comprising a couple of mismatches (not fully complementary, e.g., about
10 1, 2, 3, or 4 mismatches, e.g., about 80% to about 96% (80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96%) complementarity. This results in the CRISPR-Cas effector protein nicking the non-target strand. Without intending to be limited to any particular theory, it is believed that following nicking, there are cellular nucleases which chew back some of the non-target strand. Without the non-target strand there is no blueprint for “fixing”
15 the edit that we are making with the methods of the invention (REDRAW), resulting in greater editing efficiency. In this configuration, only one fusion enzyme is used, but it is capable of binding the two different RNAs (tagRNA and crRNA).

In some embodiments, an extended guide nucleic acid comprises: (i) a Type V CRISPR nucleic acid or Type II CRISPR nucleic acid (Type II or Type V CRISPR RNA,
20 Type II or Type V CRISPR DNA, Type II or Type V crRNA, Type II or Type V crDNA) and/or a CRISPR nucleic acid and a tracr nucleic acid (e.g., Type II or Type V tracrRNA, Type II or Type V tracrDNA); and (ii) an extended portion comprising a primer binding site and a reverse transcriptase template (RT template). In some embodiments, the extended portion can be fused to either the 5' end or 3' end of the CRISPR nucleic acid (e.g., 5' to 3':
25 repeat-spacer-extended portion, or extended portion-repeat-spacer) and/or to the 5' or 3' end of the tracr nucleic acid. In some embodiments, the extended portion of an extended guide nucleic acid comprises, 5' to 3', an RT template (RTT) and a primer binding site (PBS) (e.g., 5'-crRNA-spacer-RTT(edit encoded)-PBS-3') or comprises 5' to 3' a PBS and RTT, depending on the location of the extended portion relative to the CRISPR RNA of the guide
30 (e.g., 5'-crRNA-spacer-PBS-RTT(edit encoded)-3'). In some embodiments, a target nucleic acid is double stranded and comprises a first strand and a second strand and the primer binding site binds to the second strand (non-target, top strand) of the target nucleic acid. In some embodiments, a target nucleic acid is double stranded and comprises a first strand and a second strand and the primer binding site binds to the first strand (e.g., binds to the target

strand, same strand to which the CRISPR-Cas effector protein is recruited, bottom strand) of the target nucleic acid. In some embodiments, a target nucleic acid is double stranded and comprises a first strand and a second strand and the primer binding site binds to the second strand (non-target strand, opposite strand from that to which the CRISPR-Cas effector protein is recruited) of the target nucleic acid. Thus, in some embodiments, the editing reverse transcriptase (RT) adds to the target strand (the strand to which the spacer of the CRISPR RNA is complementary and to which the CRISPR- Cas effector protein is recruited) and in some embodiments, the editing reverse transcriptase (RT) adds to the non-target strand (the strand that is complementary to the strand to which the spacer of the CRISPR RNA is complementary and to which the CRISPR- Cas effector protein is recruited).

The RT template encodes a modification to be incorporated into the target nucleic acid (the edit). The modification of edit may be located in any position within an RT template (position location relative to the position of a protospacer adjacent motif (PAM) of the target nucleic acid). Thus, for example, **Fig. 27** shows an RT template having edits located at positions -1-19 (-1, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 27, 18, or 19) relative to the position of a protospacer adjacent motif (PAM) (TTTG) in the target nucleic acid. In each case, precise editing was observed. In some embodiments, an RT template may comprise an edit located at nucleotide position -1, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 27, 18, or 19. In some embodiments, an RT template may comprise an edit located at nucleotide position 4 to nucleotide position 17 (e.g., position 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, or 17) of the RT template relative to the position of a protospacer adjacent motif (PAM) of the target nucleic acid. In some embodiments, an RT template may comprise an edit located at nucleotide position 10 to nucleotide position 17 (e.g., position 10, 11, 12, 13, 14, 15, 16, or 17) of the RT template relative to the position of a protospacer adjacent motif (PAM) of the target nucleic acid. In some embodiments, an RT template may comprise an edit located at nucleotide position 12 to nucleotide position 15 (e.g., position 12, 13, 14, or 15) of the RT template relative to the position of a protospacer adjacent motif (PAM) of the target nucleic acid.

In some embodiments, a method of modifying a target nucleic acid having a first strand and a second strand is provided, the method comprising: contacting the target nucleic acid with (a) a Type V CRISPR-Cas effector protein or a Type II CRISPR-Cas effector protein; (b) a reverse transcriptase, and (c) an extended guide nucleic acid (e.g., extended Type II or Type V CRISPR RNA, extended Type II or Type V CRISPR DNA, extended Type II or Type V crRNA, extended Type II or Type V crDNA), wherein the extended guide

nucleic acid comprises: (i) a Type II or Type V CRISPR nucleic acid (Type II or Type V CRISPR RNA, Type II or Type V CRISPR DNA, Type II or Type V crRNA, Type II or Type V crDNA) and/or a CRISPR nucleic acid and a tracr nucleic acid (e.g., Type II or Type V tracrRNA, Type II or Type V tracrDNA); and (ii) an extended portion comprising a primer binding site and a reverse transcriptase template (RT template), and the Type II or Type V CRISPR nucleic acid comprises a spacer that binds to the first strand (e.g., target strand) (i.e., is complementary to a portion of consecutive nucleotides in the first strand of the target nucleic acid) and the primer binding site binds to the first strand (target strand), thereby modifying the target nucleic acid. In some embodiments, a Type II CRISPR-Cas effector protein can be a Cas9 polypeptide, optionally a spCas9. In some embodiments, a Type V CRISPR-Cas effector protein can be a Cas12a polypeptide or a cas12b polypeptide. In some embodiments, a Type II or Type V CRISPR-Cas effector protein, a reverse transcriptase, and an extended guide nucleic acid can form a complex or are comprised in a complex. In some embodiments, contacting can further comprise contacting the target nucleic acid with a 5'-3' exonuclease.

In some embodiments, the target nucleic acid may be additionally contacted with a 5' flap endonuclease (FEN), optionally an FEN1 and/or Dna2 polypeptide, thereby improving mismatch repair by removing the 5' flap that does not comprise the edits to be incorporated into the target nucleic acid. In some embodiments, an FEN and/or Dna2 may be overexpressed in the presence of the target nucleic acid. In some embodiments, an FEN may be a fusion protein comprising an FEN domain fused to a Type V CRISPR-Cas effector protein or domain, thereby recruiting the FEN to the target nucleic acid. In some embodiments, a Dna2 may be a fusion protein comprising a Dna2 domain fused to a Type V CRISPR-Cas effector protein or domain, thereby recruiting the Dna2 to the target nucleic acid.

In some embodiments, a Type II or Type V CRISPR-Cas effector protein may be a Type II or Type V CRISPR-Cas fusion protein comprising a Type V CRISPR-Cas effector protein domain fused (linked) to a peptide tag (e.g., an epitope or a multimerized epitope) and an FEN may be an FEN fusion protein comprising an FEN domain fused to an affinity polypeptide that binds to the peptide tag, thereby recruiting the FEN to the Type II or Type V CRISPR-Cas effector protein domain, and the target nucleic acid. In some embodiments, a Type II or Type V CRISPR-Cas effector protein may be a Type II or Type V CRISPR-Cas fusion protein comprising a Type II or Type V CRISPR-Cas effector protein domain fused (linked) to a peptide tag (e.g., an epitope or a multimerized epitope) and a Dna2 may be a

Dna2 fusion protein comprising a Dna2 domain fused to an affinity polypeptide that binds to the peptide tag, thereby recruiting the Dna2 to the Type II or Type V CRISPR-Cas effector protein domain, and the target nucleic acid. In some embodiments, a Type V CRISPR-Cas effector protein may be a Type II or Type V CRISPR-Cas fusion protein comprising a Type II or Type V CRISPR-Cas effector protein domain fused (linked) to a peptide tag (e.g., an epitope or a multimerized epitope) and an FEN may be an FEN fusion protein comprising an FEN domain fused to an affinity polypeptide that binds to the peptide tag, thereby recruiting the FEN to the Type II or Type V CRISPR-Cas effector protein domain, and the target nucleic acid. In some embodiments, a Type II or Type V CRISPR-Cas effector protein may be a Type II or Type V CRISPR-Cas fusion protein comprising a Type II or Type V CRISPR-Cas effector protein domain fused (linked) to a peptide tag (e.g., an epitope or a multimerized epitope) and a Dna2 may be a Dna2 fusion protein comprising a Dna2 domain fused to an affinity polypeptide that binds to the peptide tag, thereby recruiting the Dna2 to the Type II or Type V CRISPR-Cas effector protein domain, and the target nucleic acid. In some embodiments, a target nucleic acid may be contacted with two or more FEN fusion proteins and/or Dna2 fusion proteins.

In some embodiments, the methods of the invention may further comprise contacting the target nucleic acid with a 5'-3' exonuclease, thereby improving mismatch repair by removing the 5' flap that does not comprise the edits (non-edited strand) to be incorporated into the target nucleic acid. In some embodiments, a 5'-3' exonuclease may be fused to a Type II or Type V CRISPR-Cas effector protein, optionally to a Type II or Type V CRISPR-Cas fusion protein. In some embodiments, a 5'-3' exonuclease may be a fusion protein comprising the 5'-3' exonuclease fused to a peptide tag and a Type II or Type V CRISPR-Cas effector protein may be a fusion protein comprising a Type II or Type V CRISPR-Cas effector protein domain fused to an affinity polypeptide that is capable of binding to the peptide tag, thereby improving mismatch repair. In some embodiments, a 5'-3' exonuclease may be a fusion protein comprising a 5'-3' exonuclease fused to an affinity polypeptide that is capable of binding to the peptide tag and a Type II or Type V CRISPR-Cas effector protein may be a fusion protein comprising a Type II or Type V CRISPR-Cas effector protein domain fused to a peptide tag. In some embodiments, a 5'-3' exonuclease may be a fusion protein comprising a 5'-3' exonuclease fused to an affinity polypeptide that is capable of binding to an RNA recruiting motif and the extended guide nucleic acid is linked to an RNA recruiting motif, thereby recruiting the 5'-3' exonuclease to the target nucleic acid via interaction between the affinity polypeptide and RNA recruiting motif. A 5'-3' exonuclease

may be any known or later discovered 5'-3' exonuclease functional in the organism, cell or *in vitro* system of interest. In some embodiments, a 5'-3' exonuclease can include but is not limited to, a RecE exonuclease (RecE, e.g., **SEQ ID NO:129**), a RecJ exonuclease (RecJ, e.g., **SEQ ID NO:130**), a T5 exonuclease (T5_Exo, e.g., **SEQ ID NO:131**), and/or a T7 exonuclease (T7_Exo, e.g., **SEQ ID NO:132**), Lambda exonuclease (Lambda_Exo, e.g., **SEQ ID NO:133**), *E. coli* exonuclease sbcB (**SEQ ID NO:134**) and/or human exonuclease (Exo, e.g., **SEQ ID NO:135**). In some embodiments, a RecE exonuclease C-terminal fragment flanked on both sides with nuclear localization sequences (NLS) from, for example, *Escherichia coli* (strain K12) may be used (**SEQ ID NO:98**). In some embodiments, a RecJ exonuclease flanked on both sides with nuclear localization sequences (NLS) from, for example, *Escherichia coli* (strain K12) may be used (**SEQ ID NO:99**). In some embodiments, a T5 exonuclease flanked on both sides with nuclear localization sequences (NLS) may be used (**SEQ ID NO:100**). In some embodiments, a T7 exonuclease flanked on both sides with nuclear localization sequences (NLS) from, for example, *Escherichia* phage 7 may be used (**SEQ ID NO:101**). In some embodiments, a 5'-3' exonuclease includes, but is not limited to, a RecE (e.g., **SEQ ID NO:129**), RecJ (e.g., **SEQ ID NO:130**), T5_Exo (e.g., **SEQ ID NO:131**), T7_Exo (e.g., **SEQ ID NO:132**), sbcB (**SEQ ID NO:134**) and/or Exo (**SEQ ID NO:135**).

In some embodiments, the methods of the invention may further comprise reducing double strand breaks. In some embodiments, reducing double strand breaks may be carried out by introducing, in the region of the target nucleic acid, a chemical inhibitor of non-homologous end joining (NHEJ), or by introducing a CRISPR guide nucleic acid, or an siRNA targeting an NHEJ protein to transiently knock-down expression of the NHEJ protein. In some embodiments, an inhibitor of NHEJ may be fused to the reverse transcriptase (RT) or the CRISPR-Cas effector protein of the invention, optionally to the N-terminal end of the RT or CRISPR-Cas effector protein. In some embodiments, an inhibitor of NHEJ includes, but is not limited to, *Escherichia* phage Mu Gam (**SEQ ID NO:147**).

In some embodiments, a Type II or Type V CRISPR-Cas effector protein may be a fusion protein and/or the reverse transcriptase may be a fusion protein, wherein the Type II or Type V CRISPR-Cas fusion protein, the reverse transcriptase fusion protein and/or the extended guide nucleic acid may be fused to one or more components, which allow for the recruiting the reverse transcriptase to the Type II or Type V CRISPR-Cas effector protein. In some embodiments, the one or more components recruit via protein-protein interactions, protein-RNA interactions, and/or chemical interactions.

Thus, in some embodiments, a Type V CRISPR-Cas effector protein may be a Type V CRISPR-Cas effector fusion protein comprising a Type V CRISPR-Cas effector protein domain fused (linked) to a peptide tag (e.g., an epitope or a multimerized epitope) and the reverse transcriptase may be a reverse transcriptase fusion protein comprising a reverse transcriptase domain fused (linked) to an affinity polypeptide that binds to the peptide tag, wherein the Type V CRISPR-Cas effector protein interacts with the guide nucleic acid, which guide nucleic acid binds to the target nucleic acid, thereby recruiting the reverse transcriptase to the Type V CRISPR-Cas effector protein and to the target nucleic acid. In some embodiments, the Type II CRISPR-Cas effector protein is a Type II CRISPR-Cas fusion protein comprising a Type II CRISPR-Cas effector protein domain fused (linked) to a peptide tag (e.g., an epitope or a multimerized epitope) and the FEN is an FEN fusion protein comprising an FEN domain fused to an affinity polypeptide that binds to the peptide tag, and/or wherein the Type II CRISPR-Cas effector protein is a Type II CRISPR-Cas fusion protein comprising a Type II CRISPR-Cas effector protein domain fused to a peptide tag and the Dna2 polypeptide is an Dna2 fusion protein comprising an Dna2 domain fused to an affinity polypeptide that binds to the peptide tag, optionally wherein the target nucleic acid is contacted with two or more FEN fusion proteins and/or two or more Dna2 fusion proteins, thereby recruiting the FEN and/or Dna2 to the Type II CRISPR-Cas effector protein domain, and the target nucleic acid. In some embodiments, two or more reverse transcriptase fusion proteins may be recruited to the Type II or Type V CRISPR-Cas effector protein, thereby contacting the target nucleic acid with two or more reverse transcriptase fusion proteins.

A peptide tag may include, but is not limited to, a GCN4 peptide tag (e.g., Sun-Tag), a c-Myc affinity tag, an HA affinity tag, a His affinity tag, an S affinity tag, a methionine-His affinity tag, an RGD-His affinity tag, a FLAG octapeptide, a strep tag or strep tag II, a V5 tag, and/or a VSV-G epitope. Any epitope that may be linked to a polypeptide and for which there is a corresponding affinity polypeptide that may be linked to another polypeptide may be used with this invention. In some embodiments, a peptide tag may comprise 1 or 2 or more copies of a peptide tag (e.g., epitope, multimerized epitope (e.g., tandem repeats)) (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or more peptide tags. In some embodiments, an affinity polypeptide that binds to a peptide tag may be an antibody. In some embodiments, the antibody may be a scFv antibody. . In some embodiments, an affinity polypeptide that binds to a peptide tag may be synthetic (e.g., evolved for affinity interaction) including, but not limited to, an affibody, an anticalin, a monobody and/or a DARPIn (see, e.g., Sha et al., *Protein Sci.* 26(5):910-924 (2017));

Gilbreth (*Curr Opin Struc Biol* 22(4):413-420 (2013)), U.S. Patent No. 9,982,053, each of which are incorporated by reference in their entireties for the teachings relevant to affibodies, anticalins, monobodies and/or DARPin. Example peptide tag sequences and their affinity polypeptides include, but are not limited to, the amino acid sequences of **SEQ ID NOs:23-**

5 **25.**

In some embodiments, an extended guide nucleic acid may be linked to an RNA recruiting motif, and the reverse transcriptase may be a reverse transcriptase fusion protein, wherein the reverse transcriptase fusion protein may comprise a reverse transcriptase domain fused to an affinity polypeptide that binds to the RNA recruiting motif, wherein the extended
10 guide binds to the target nucleic acid and the RNA recruiting motif binds to the affinity polypeptide, thereby recruiting the reverse transcriptase fusion protein to the extended guide and contacting the target nucleic acid with the reverse transcriptase domain. In some embodiments, two or more reverse transcriptase fusion proteins may be recruited to an extended guide nucleic acid, thereby contacting the target nucleic acid with two or more
15 reverse transcriptase fusion proteins. Example RNA recruiting motifs and their affinity polypeptides include, but are not limited to, the sequences of **SEQ ID NOs:26-36.**

In some embodiments, an RNA recruiting motif may be located on the 3' end of the extended portion of the extended guide nucleic acid (e.g., 5'-3', repeat-spacer-extended portion (RT template-primer binding site)-RNA recruiting motif). In some embodiments, an
20 RNA recruiting motif may be embedded in the extended portion.

In some embodiments of the invention, an extended guide RNA and/or guide RNA may be linked to one or to two or more RNA recruiting motifs (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more motifs, e.g., at least 10 to about 25 motifs), optionally wherein the two or more RNA recruiting motifs may be the same RNA recruiting motif or different RNA recruiting motifs.
25 In some embodiments, an RNA recruiting motif and corresponding affinity polypeptide may include, but is not limited, to a telomerase Ku binding motif (e.g., Ku binding hairpin) and the corresponding affinity polypeptide Ku (e.g., Ku heterodimer), a telomerase Sm7 binding motif and the corresponding affinity polypeptide Sm7, an MS2 phage operator stem-loop and the corresponding affinity polypeptide MS2 Coat Protein (MCP), a PP7 phage operator stem-
30 loop and the corresponding affinity polypeptide PP7 Coat Protein (PCP), an SfMu phage Com stem-loop and the corresponding affinity polypeptide Com RNA binding protein, a PUF binding site (PBS) and the affinity polypeptide Pumilio/fem-3 mRNA binding factor (PUF), and/or a synthetic RNA-aptamer and the aptamer ligand as the corresponding affinity polypeptide. In some embodiments, the RNA recruiting motif and corresponding affinity

polypeptide may be an MS2 phage operator stem-loop and the affinity polypeptide MS2 Coat Protein (MCP). In some embodiments, the RNA recruiting motif and corresponding affinity polypeptide may be a PUF binding site (PBS) and the affinity polypeptide Pumilio/fem-3 mRNA binding factor (PUF).

5 In some embodiments, the components for recruiting polypeptides and nucleic acids may those that function through chemical interactions that may include, but are not limited to, rapamycin-inducible dimerization of FRB – FKBP; Biotin-streptavidin; SNAP tag; Halo tag; CLIP tag; DmrA-DmrC heterodimer induced by a compound; bifunctional ligand (e.g., fusion of two protein-binding chemicals together, e.g., dihydrofolate reductase (DHFR).

10 In some embodiments of the invention, a CRISPR-Cas effector protein (e.g., a CRISPR-Cas effector protein, a first CRISPR-Cas effector protein, a second CRISPR-Cas effector protein, a third CRISPR-Cas effector protein, and/or a fourth CRISPR-Cas effector protein) may be from a Type I CRISPR-Cas system, a Type II CRISPR-Cas system, a Type III CRISPR-Cas system, a Type IV CRISPR-Cas system and/or a Type V CRISPR-Cas
15 system. In some embodiments, the CRISPR-Cas nuclease is from a Type II CRISPR-Cas system or a Type V CRISPR-Cas system.

In some embodiments of the invention, a CRISPR-Cas effector protein may be a Cas9, C2c1, C2c3, Cas12a (also referred to as Cpf1), Cas12b, Cas12c, Cas12d, Cas12e, Cas13a, Cas13b, Cas13c, Cas13d, Cas1, Cas1B, Cas2, Cas3, Cas3', Cas3'', Cas4, Cas5, Cas6,
20 Cas7, Cas8, Cas9 (also known as Csn1 and Csx12), Cas10, Csy1, Csy2, Csy3, Cse1, Cse2, Cse1, Cse2, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx15, Csf1, Csf2, Csf3, Csf4 (dinG), and/or Csf5 nuclease, optionally wherein the CRISPR-Cas nuclease may be a Cas9, Cas12a (Cpf1), Cas12b, Cas12c (C2c3), Cas12d (CasY), Cas12e (CasX), Cas12g,
25 Cas12h, Cas12i, C2c4, C2c5, C2c8, C2c9, C2c10, Cas14a, Cas14b, and/or Cas14c nuclease.

In some embodiments, a CRISPR-Cas effector protein may be a protein that functions as a nickase (e.g., a Cas9 nickase or a Cas12a nickase). In some embodiments, a CRISPR-Cas effector protein useful with the invention may comprise a mutation in its nuclease active site (e.g., RuvC, HNH, e.g., RuvC site of a Cas12a nuclease domain, e.g., RuvC site and/or HNH
30 site of a Cas9 nuclease domain). A CRISPR-Cas effector protein having a mutation in its nuclease active site, and therefore, no longer comprising nuclease activity, is commonly referred to as "dead," or "deactivated" e.g., dCas. In some embodiments, a CRISPR-Cas nuclease domain or polypeptide having a mutation in its nuclease active site may have impaired activity or reduced activity as compared to the same CRISPR-Cas nuclease without

the mutation. In some embodiments, a CRISPR-Cas effector protein useful with the invention may be a double stranded nuclease. In some embodiments, a CRISPR-Cas effector protein having double stranded nuclease activity may be a Type II or a Type V CRISPR-Cas effector protein. In some embodiments, a Type V CRISPR-Cas effector protein having double stranded nuclease activity is a Cas12a polypeptide. In some embodiments, a Type II CRISPR-Cas effector protein having double stranded nuclease activity is a Cas9 polypeptide.

In some embodiments, a CRISPR-Cas effector protein may be a Type V CRISPR-Cas effector protein. In some embodiments, a Type V CRISPR-Cas effector protein may comprise a Cas12a (Cpf1), Cas12b (C2c1), Cas12c (C2c3), Cas12d (CasY), Cas12e (CasX), Cas12g, Cas12h, Cas12i, C2c4, C2c5, C2c8, C2c9, C2c10, Cas14a, Cas14b, and/or Cas14c effector protein and/or domain.

In some embodiments, a Cas12a (Cpf1) can include, but is not limited to, LbCas12a, Lb2Cas12a, Lb3Cas12a, AsCas12a, BpCas12a, CMtCas12a, EeCas12a, FnCas12a, LiCas12a, MbCas12a, PbCas12a, PcCas12a, PdCas12a, PeCas12a, PmCas12a, SsCas12a, enAsCas12a, optionally wherein the Cas12a comprises one or more mutations as described herein. In some embodiments, a Cas12b (C2c1) can include, but is not limited to, BhCas12b, optionally wherein the Cas12b comprises one or more mutations as described herein.

In some embodiments, a Type V CRISPR-Cas effector protein can include, but is not limited to, a Type V CRISPR-Cas effector protein from *Acidaminococcus* sp. (AsCas12a), from Lachnospiraceae bacterium (e.g., LbCas12a) or from *Butyrivibrio hungatei* (BhCas12b) or a modified Type V CRISPR-Cas effector protein thereof. In some embodiments, a Type V CRISPR-Cas effector protein from *Acidaminococcus* sp. may comprise a sequence having at least 80% identity to **SEQ ID NO:2**. In some embodiments, a Type V CRISPR-Cas effector protein from Lachnospiraceae bacterium may comprise an amino acid sequence having at least 80% identity to any one of **SEQ ID NO:1**, **SEQ ID NO:7**, **SEQ ID NO:8**, or **SEQ ID NO:9**. In some embodiments, a Type V CRISPR-Cas effector protein from *Butyrivibrio hungatei* may comprise a sequence having at least 80% identity to the amino acid sequence of **SEQ ID NO:151**. In some embodiments, a modified Type V CRISPR-Cas effector protein from *Acidaminococcus* sp. may comprise an amino acid sequence having at least 80% identity to any one of the **SEQ ID NO:18**, **SEQ ID NO:19**, **SEQ ID NO:20**, **SEQ ID NO:149**, or **SEQ ID NO:150**. In some embodiments, a modified Type V CRISPR-Cas effector protein from Lachnospiraceae bacterium may comprise a sequence having at least 80% identity to **SEQ ID NO:148**.

In some embodiments, a Type II CRISPR-Cas effector protein can include, but is not limited to, a Cas9 effector protein, optionally wherein the Cas9 effector protein may be from *Streptococcus*, optionally from *Streptococcus pyogenes*. In some embodiments, a Cas9 effector protein may be a modified Cas9 effector protein. In some embodiments, a Cas9 effector protein can comprise a polypeptide sequence having at least 80% identity to any one of **SEQ ID NO:106** or **SEQ ID NO:107**. In some embodiments, a Cas9 effector protein can be encoded by a polynucleotide sequence having at least 80% identity to any one of **SEQ ID NOs:108-122**.

In some embodiments, a Type V CRISPR-Cas system may comprise an effector protein that utilizes a Type V CRISPR nucleic acid only. In some embodiments, a Type V CRISPR-Cas system may comprise an effector protein that, similar to Type II CRISPR-Cas systems, utilize both a CRISPR nucleic acid and a trans-activating CRISPR (tracr) nucleic acid. Thus, in some embodiments, a Type V CRISPR-Cas effector protein useful with the present invention may function with a corresponding CRISPR nucleic acid only (e.g., Cas12a, Cas12a, Cas12i, Cas12h, Cas14b, Cas14c, C2c10, C2c9, C2c8, C2c4). In some embodiments, a Type V CRISPR-Cas effector protein useful with the present invention may function with a corresponding CRISPR nucleic acid and tracr nucleic acid (e.g., Cas12b, Cas12c, Cas12e, Cas12g, Cas14a).

A CRISPR nucleic acid useful with this invention may comprise at least one repeat sequence that is capable of interacting with a corresponding Type V CRISPR-Cas effector protein, and at least one spacer sequence, wherein the at least one spacer sequence is capable of binding a target nucleic acid (e.g., a first strand or a second strand of the target nucleic acid). In some embodiments, a repeat sequence of a CRISPR nucleic acid may be located 5' to the spacer sequence. In some embodiments, CRISPR nucleic acid may comprise more than one repeat sequence, wherein the repeat sequence is linked to both the 5' end and the 3' end of the spacer. In some embodiments, a CRISPR nucleic acid useful with this invention may comprise two or more repeat and one or more spacer sequences, wherein each spacer sequence is linked at the 5' end and the 3' end with a repeat sequence.

A tracr nucleic acid useful with this invention may comprises a first portion that is substantially complementary to and hybridizes to the repeat sequence of a corresponding CRISPR nucleic acid and a second portion that interacts with a corresponding Type II or a Type V CRISPR-Cas effector protein.

In some embodiments, a Type V CRISPR-Cas effector protein useful for this invention may function as a double stranded DNA nuclease. In some embodiments, a Type

V CRISPR-Cas effector protein may function as a single stranded DNA nickase, optionally wherein the first strand is nicked. In some embodiments, a Type V CRISPR-Cas effector protein may function as a single stranded DNA nickase, optionally wherein the second strand is nicked. In some embodiments, the Type V CRISPR-Cas effector protein may be a Cas12a effector protein that functions as a nickase, optionally wherein the first strand (target strand) is nicked. In some embodiments, the Type V CRISPR-Cas effector protein may be a Cas12a effector protein that functions as a nickase, optionally wherein the second strand is nicked. In some embodiments, the Type V CRISPR-Cas effector protein may be a Cas12a effector protein that functions as a nickase through the use of crRNAs that contain strategic mismatches. Thus, for example, a crRNA may comprise a spacer having one to about four mismatches (e.g., 1, 2, 3, or 4 mismatches) (e.g., 80-96% complementary).

In some embodiments, a Cas12a effector protein may be a Cas12a nickase having a mutation of the arginine in the LQMRNS motif. A mutation of the arginine in this motif may be to any amino acid, thereby providing a Cas12a nickase. In some embodiments, the mutation may be to an alanine. In some embodiments, the mutation may be to an alanine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, or valine. In some embodiments, the mutation may be a mutation to an alanine. In some embodiments, the mutation does not include a mutation to a lysine or a histidine. In some embodiments, a Cas12a effector protein may be an LbCas12a nickase comprising an R1138, optionally a R1138A mutation (see reference nucleotide sequence **SEQ ID NO:9**), an R1137 mutation, optionally a R1137A mutation (see reference nucleotide sequence **SEQ ID NO:1**), or an R1124 mutation, optionally a R1124A mutation (see reference nucleotide sequence **SEQ ID NO:7**). In some embodiments, a Cas12a effector protein may be an AsCas12a nickase comprising an R1226 mutation, optionally an R1226A mutation (see reference nucleotide sequence **SEQ ID NO:2**). In some embodiments, a Cas12a effector protein may be a FnCas12a nickase comprising an R1218 mutation, optionally an R1218A mutation (see reference nucleotide sequence **SEQ ID NO:6**). In some embodiments, a Cas12a effector protein may be a PdCas12a nickase comprising an R1241 mutation, optionally an R1241A mutation (see reference nucleotide sequence **SEQ ID NO:14**).

In some embodiments, a Type V CRISPR-Cas effector protein useful with this invention may comprise reduced single stranded DNA cleavage activity (ss DNase activity) (e.g., the Type V CRISPR-Cas effector protein may be modified (mutated) to reduce ss DNase activity (e.g., about 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90,

95, 96, 97, 98, 99, or 100% less ss DNase activity than a wild-type or non-modified Type V CRISPR-Cas effector protein).

In some embodiments, a Type V CRISPR-Cas effector protein useful with this invention may comprise reduced self-processing RNase activity (e.g., the Type V CRISPR-Cas effector protein may be modified (mutated) to reduce self-processing RNase activity (e.g., about 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, 99, or 100% less self-processing RNase activity than a wild-type or non-modified Type V CRISPR-Cas effector protein). In some embodiments, a mutation to reduce self-processing RNase activity may be a mutation of a histidine at residue position 759 with reference to nucleotide position numbering of **SEQ ID NO:1** or **SEQ ID NO:9**, optionally a mutation of a histidine to alanine (H759A). An example Type V CRISPR-Cas effector protein having reduced single stranded DNA cleavage activity can include, but is not limited to, LbCas12a (H759A) (**SEQ ID NO:148**). In some embodiments, a Cas12a CRISPR-Cas effector protein having a H759A mutation useful with the invention may comprise a sequence having at least 80% sequence identity to the amino acid sequence of **SEQ ID NO:148**. In some embodiments, a Cas12a CRISPR-Cas effector protein having a H759A mutation may be a LbCas12a CRISPR-Cas effector protein, optionally wherein the LbCas12a CRISPR-Cas effector protein comprises at least 90% sequence identity to the amino acid sequence of **SEQ ID NO:148**.

In some embodiments, a Type V CRISPR-Cas effector protein or domain useful with the invention may comprise a mutation in its nuclease active site (e.g., RuvC of a dType V CRISPR-Cas effector protein or domain, e.g., RuvC site of a Cas12a nuclease domain). A CRISPR-Cas nuclease having a mutation in its nuclease active site, and therefore, no longer comprising nuclease activity, is commonly referred to as "deactivated" or "dead," e.g., dCas, dCas12a. In some embodiments, a CRISPR-Cas nuclease domain or polypeptide having a mutation in its nuclease active site may have impaired activity or reduced activity as compared to the same CRISPR-Cas nuclease without the mutation. In some embodiments, deactivated Type V CRISPR-Cas effector protein may function as a nickase (a first strand nickase and/or a second strand nickase).

In some embodiments, a Type V CRISPR-Cas effector protein or domain useful with the invention may comprise a modification of one or more amino acid residues that reduce(s) the DNA binding affinity of the Type V CRISPR-Cas effector protein. In some embodiments, the modification may be an amino acid substitution. In some embodiments, positively charged residues that interact with DNA backbone may be mutated, optionally

wherein the positively charged residues that interact with DNA backbone may be mutated to an alanine (e.g., substituted with an alanine). Substitution of a positively charged residue for an alanine in a Cas12a effector protein can include, but is not limited to, the amino acid substitution of K167A, K272A, and/or K349A with reference to the amino acid position numbering of **SEQ ID NO:1** or **SEQ ID NO:148**. In some embodiments, the Type V CRISPR-Cas effector protein is a Cas12a CRISPR-Cas effector protein comprising an amino acid substitution of K167A, K272A, K349A, K167A+ K272A, K167A+ K349A, K272A+ K349A, or K167A+ K272A + K349A with reference to the amino acid position numbering of **SEQ ID NO:148**, optionally wherein the Type V CRISPR-Cas effector protein is an LbCas12a.

In some embodiments, a Type V CRISPR-Cas effector protein may be a Type V CRISPR-Cas fusion protein, wherein the Type V CRISPR-Cas fusion protein comprises a Type V CRISPR-Cas effector protein domain fused to a reverse transcriptase. In some embodiments, the reverse transcriptase may be fused to the C-terminus of the Type V CRISPR-Cas effector polypeptide. In some embodiments, the reverse transcriptase may be fused to the N-terminus of the Type V CRISPR-Cas effector polypeptide.

In some embodiments, a Type V CRISPR-Cas effector protein may be a Type V CRISPR-Cas fusion protein, wherein the Type V CRISPR-Cas fusion protein comprises a Type V CRISPR-Cas effector protein domain fused to a nicking enzyme (e.g., FokI, BfiI, e.g., an engineered FokI or BfiI), optionally wherein the Type V CRISPR-Cas effector protein domain may be a deactivated Type V CRISPR-Cas domain fused to the nicking enzyme.

In some embodiments, a Type II CRISPR-Cas effector protein may be a Type II CRISPR-Cas fusion protein, wherein the Type II CRISPR-Cas fusion protein comprises a Type II CRISPR-Cas effector protein domain fused to a reverse transcriptase. In some embodiments, the reverse transcriptase may be fused to the C-terminus of the Type II CRISPR-Cas effector polypeptide. In some embodiments, the reverse transcriptase may be fused to the N-terminus of the Type II CRISPR-Cas effector polypeptide. In some embodiments, a Type II CRISPR-Cas effector protein may be a Type II CRISPR-Cas fusion protein, wherein the Type II CRISPR-Cas fusion protein comprises a Type II CRISPR-Cas effector protein domain fused to a nicking enzyme (e.g., FokI, BfiI, e.g., an engineered FokI or BfiI), optionally wherein the Type II CRISPR-Cas effector protein domain may be a deactivated Type II CRISPR-Cas domain fused to the nicking enzyme.

In some embodiments, a reverse transcriptase useful with this invention may be a wild type reverse transcriptase. In some embodiments, a reverse transcriptase useful with this invention may be a synthetic reverse transcriptase, see, e.g., Heller et al. *Nucleic Acids Research*, 47(7) 3619–3630 (2019)). Example reverse transcriptase polypeptides include, but
5 are not limited to, those having substantial identity (e.g., at least about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or 100% identity) to the amino acid sequence of **SEQ ID NO:53** or **SEQ ID NO:172**.

In some embodiments, the activity of a reverse transcriptase may be modified for
10 (Type V or Type II) gene editing activity to provide optimal activity in association with a Type V or Type II CRISPR-Cas effector polypeptide (e.g., an increase in activity when associated with a Type V CRISPR-Cas effector polypeptide by about 5, 10, 15, 20, 25, 30, 345, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, 99, or 100% as compared to the reference reverse transcriptase that has not been modified). Such mutations include those that
15 affect or improve RT initiation, processivity, enzyme kinetics, temperature sensitivity, and/or error rate.

In some embodiments, a reverse transcriptase useful with this invention may be modified to improve the transcription function of the reverse transcriptase. The transcription function of a reverse transcriptase may be improved by improving the processivity of the
20 reverse transcriptase, e.g., increase the ability of the reverse transcriptase to polymerize more DNA bases during a single binding event to the template (e.g., before it falls off the template) (e.g., increase processivity by about 5, 10, 15, 20, 25, 30, 345, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, 99, or 100% as compared to the reference reverse transcriptase that has not been modified).

25 In some embodiments, transcription function of a reverse transcriptase may be improved by increasing the template affinity of the reverse transcriptase (e.g., increase template affinity by about 5, 10, 15, 20, 25, 30, 345, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, 99, or 100% as compared to the reference reverse transcriptase that has not been modified).

30 In some embodiments, transcription function of a reverse transcriptase may be improved by improving the thermostability of the reverse transcriptase for improved performance at a desired temperature (e.g., increase thermostability by about 5, 10, 15, 20, 25, 30, 345, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, 99, or 100% as compared to the reference reverse transcriptase that has not been modified). In some

embodiments, the improved thermostability is at a temperature of about 20°C to 42°C (e.g., about 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, or 42°C, and any value or range therein). In some embodiments, a reverse transcriptase having improved thermostability may include, but is not limited to, M-MuLV trimutant

- 5 D200N+L603W+T330P or M-MuLV pentamutant (5M)
 D200N+L603W+T330P+T306K+W313F with reference to amino acid position numbering of **SEQ ID NO:172** (e.g., **SEQ ID NO:53**). See, e.g., Baranauskas et al. (*Protein Eng. Des. Sel.* **25**, 657–668 (2012)); Anzalone et al. (*Nature* 576:149–157 (2019)). Additional amino acid modifications in a reverse transcriptase can include the amino acid substitutions of L139P,
 10 D200N, W388R, E607K, T306K, W313F, F155Y, H638G, Q221R, V223M and/or D524N with reference to the amino acid position numbering of **SEQ ID NO:172**.

In some embodiments, a reverse transcriptase useful with this invention can include, but is not limited to, combinations of amino acid substitutions of (1) L139P, D200N, W388R, and E607K, (2) L139P, D200N, T306K, W313F, W388R, and E607K, (3) 5M

- 15 (T355A/Q357M/K358R/A359G/S360A), F155Y, and H638G, (4) 5M
 (T355A/Q357M/K358R/A359G/S360A), Q221R, and V223M; or (5) 5M
 T355A/Q357M/K358R/A359G/S360A) and D524N with reference to the amino acid position numbering of **SEQ ID NO:172**.

- In some embodiments of the invention, a reverse transcriptase may be fused to one or
 20 more single stranded RNA binding domains (RBDs). RBDs useful with the invention may include, but are not limited to, **SEQ ID NOS:37-52** (**SEQ ID NO:37**, **SEQ ID NO:38**, **SEQ ID NO:39**, **SEQ ID NO:40**, **SEQ ID NO:41**, **SEQ ID NO:42**, **SEQ ID NO:43**, **SEQ ID NO:44**, **SEQ ID NO:45**, **SEQ ID NO:46**, **SEQ ID NO:47**, **SEQ ID NO:48**, **SEQ ID NO:49**, **SEQ ID NO:50**, **SEQ ID NO:51**, and/or **SEQ ID NO:52**), thereby improving the
 25 thermostability, processivity and template affinity of the reverse transcriptase.

- The polypeptides/proteins/domains of this invention (e.g., a CRISPR-Cas effector protein e.g., a Type II or Type V CRISPR-Cas effector protein), a reverse transcriptase, a 5' flap endonuclease, and/or a 5'-3' exonuclease) may be encoded by one or more polynucleotides, optionally operably linked to one or more promoters and/or other regulatory
 30 sequences (e.g., terminator, operon, and/or enhancer and the like). In some embodiments, the polynucleotides of this invention may be comprised in one or more expression cassettes and/or vectors. In some embodiments, the at least one regulatory sequence may be, for example, a promoter, an operon, a terminator, or an enhancer. In some embodiments, the at least one regulatory sequence may be a promoter. In some embodiments, the regulatory

sequence may be an intron. In some embodiments, the at least one regulatory sequence may be, for example, a promoter operably associated with an intron or a promoter region comprising an intron. In some embodiments, the at least one regulatory sequence may be, for example a ubiquitin promoter and its associated intron (e.g., *Medicago truncatula* and/or *Zea mays* and their associated introns) (e.g., ZmUbi1 comprising an intron; MtUb2 comprising an intron, e.g., **SEQ ID NOs:21 or 22**.

In some embodiments, the present invention provides a polynucleotide encoding a Type II CRISPR-Cas effector protein or domain or a Type V CRISPR-Cas effector protein or domain, a polynucleotide encoding a CRISPR-Cas effector protein or domain, a polynucleotide encoding a reverse transcriptase polypeptide or domain, a polynucleotide encoding a 5'-3' exonuclease polypeptide or domain and/or a polynucleotide encoding a flap endonuclease polypeptide or domain operably associated with one or more promoter regions that comprise or are associated with an intron, optionally wherein the promoter region may be a ubiquitin promoter and intron (e.g., a *Medicago* or a maize ubiquitin promoter and intron, e.g., **SEQ ID NOs:21 or 22**.

In some embodiments, a polynucleotide encoding a Type II or Type V CRISPR-Cas effector protein and/or a polynucleotide encoding a reverse transcriptase may be comprised in the same or separate expression cassettes, optionally when the polynucleotide encoding the Type II or Type V CRISPR-Cas effector protein and the polynucleotide encoding the reverse transcriptase are comprised in the same expression cassette, the polynucleotide encoding the Type II or Type V CRISPR-Cas effector protein and the polynucleotide encoding the reverse transcriptase may be operably linked to a single promoter or to two or more separate promoters in any combination. In some embodiments, a polynucleotide encoding a CRISPR-Cas effector protein may be comprised in an expression cassette, wherein the polynucleotide encoding the CRISPR-Cas effector protein may be operably linked to a promoter.

In some embodiments, an extended guide nucleic acid and/or guide nucleic acid may be comprised in an expression cassette, optionally wherein the expression cassette is comprised in a vector. In some embodiments, an expression cassette and/or vector comprising the extended guide nucleic acid may be the same or a different expression cassette and/or vector from that comprising the polynucleotide encoding the Type II or Type V CRISPR-Cas effector protein and/or the polynucleotide encoding the reverse transcriptase. In some embodiments, an expression cassette and/or vector comprising the guide nucleic acid may be the same or a different expression cassette and/or vector from that comprising the polynucleotide encoding the CRISPR-Cas effector protein.

In some embodiments, a polynucleotide encoding a 5' flap endonuclease and/or a polynucleotide encoding a 5'-3' exonuclease may be comprised in one or more expression cassettes, which may be the same or different expression cassettes. In some embodiments, an expression cassette comprising a polynucleotide encoding a 5' flap endonuclease and/or a polynucleotide encoding a 5'-3' exonuclease may be the same or different expression cassette from that comprising a polynucleotide encoding a Type II or Type V CRISPR-Cas effector protein, a polynucleotide encoding a Type II or Type V CRISPR-Cas effector protein and/or a polynucleotide encoding a reverse transcriptase.

In some embodiments of the invention, polynucleotides encoding CRISPR-Cas effector proteins (e.g., a Type II CRISPR-Cas effector protein, a Type V CRISPR-Cas effector protein), reverse transcriptase, flap endonucleases, 5'-3' exonucleases, and fusion proteins comprising the same and nucleic acid constructs, expression cassettes and/or vectors comprising the polynucleotides may be codon optimized for expression in an organism (e.g., an animal (e.g., a mammal, an insect, a fish, and the like), a plant (e.g., a dicot plant, a monocot plant), a bacterium, an archaeon, a virus, and the like). In some embodiments, the polynucleotides, expression cassettes, and/or vectors may be codon optimized for expression in a plant, optionally a dicot plant or a monocot plant. Exemplary mammals for which this invention may be useful include, but are not limited to, primates (human and non-human (e.g., a chimpanzee, baboon, monkey, gorilla, *etc.*)), cats, dogs, ferrets, gerbils, hamsters, cows, pigs, horses, goats, donkeys, or sheep. In some embodiments, the polynucleotides, expression cassettes, and/or vectors may be codon optimized for expression in a fungus, including, but not limited to, a Zygomycota, Ascomycota, Basidiomycota, and Deuteromycota (fungi imperfecti), optionally wherein the fungus may be an ascomycete, optionally a yeast (e.g., *Saccharomyces cerevisiae*).

In some embodiments, the polynucleotides, nucleic acid constructs, expression cassettes or vectors of the invention that are optimized for expression in an organism may be about 70% to 100% identical (e.g., about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or 100%) to the nucleic acid constructs, expression cassettes or vectors encoding the same but which have not been codon optimized for expression in a plant.

In some embodiments, polynucleotides, nucleic acid constructs, expression cassettes and vectors may be provided for carrying out the methods of the invention. Thus, in some embodiments an expression cassette is provided that is codon optimized for expression in an

organism, comprising 5' to 3' (a) polynucleotide encoding a promoter sequence, (b) a polynucleotide encoding a Type V CRISPR-Cas nuclease (e.g., Cpf1 (Cas12a), dCas12a and the like) or a Type II CRISPR-Cas nuclease (e.g., Cas9, dCas9 and the like) that is codon-optimized for expression in the organism; (c) a linker sequence; and (d) a polynucleotide encoding a reverse transcriptase that is codon-optimized for expression in the organism. In some embodiments, the organism is an animal, a plant, a fungus, an archaeon, or a bacterium. In some embodiments, the organism is a plant and the polynucleotide encoding a Type V CRISPR-Cas nuclease is codon optimized for expression in a plant, and the promoter sequence is a plant specific promoter sequence (e.g., ZmUbi1, MtUbi2, RNA polymerase II (Pol II)).

In some embodiments, polynucleotides, nucleic acid constructs, expression cassettes and vectors may be provided for carrying out the methods of the invention. Thus, in some embodiments, an expression cassette is provided that is codon optimized for expression in a plant, comprising 5' to 3' (a) polynucleotide encoding a plant specific promoter sequence (e.g. ZmUbi1, MtUbi2, RNA polymerase II (Pol II)), (b) a plant codon-optimized polynucleotide encoding a Type II or Type V CRISPR-Cas effector protein (e.g., Cpf1 (Cas12a), dCas12a and the like), (c) a linker sequence; and (d) a plant codon-optimized polynucleotide encoding a reverse transcriptase.

In some embodiments, polypeptides of the invention may be fusion proteins comprising one or more polypeptides linked to one another via a linker. In some embodiments, the linker may be an amino acid or peptide linker. In some embodiments, a peptide linker may be about 2 to about 100 amino acids (residues) in length, as described herein. In some embodiments, a peptide linker may be, for example, a GS linker.

In some embodiments, the invention provides an expression cassette that is codon optimized for expression in a plant, comprising: (a) a polynucleotide encoding a plant specific promoter sequence (e.g. ZmUbi1, MtUbi2), and (b) an extended guide nucleic acid sequence, wherein the extended guide nucleic acid comprises an extended portion comprising at its 3' end a primer binding site and an edit to be incorporated into the target nucleic acid (e.g., edit in the reverse transcriptase template) (e.g., 5'-3' – crRNA-RTT-PBS) (e.g., tag nucleic acid; e.g., tagRNA), optionally wherein the extended guide nucleic acid is comprised in an expression cassette, optionally wherein the extended guide nucleic acid is operably linked to a Pol II promoter. In some embodiments, when the extended portion of the guide nucleic acid is attached to a CRISPR RNA at the 5' end of the crRNA, the extended portion

comprises at its 5' end a primer binding site and an edit to be incorporated into the target nucleic acid (e.g., reverse transcriptase template) at the 3' end (5'-3' - PBS-RTT-crRNA).

In some embodiments, an expression cassette of the invention may be codon optimized for expression in a dicot plant or in a monocot plant. In some embodiments, the expression cassettes of the invention may be used in a method of modifying a target nucleic acid in a plant or plant cell, the method comprising introducing one or more expression cassettes of the invention into a plant or plant cell, thereby modifying the target nucleic acid in the plant or plant cell to produce a plant or plant cell comprising the modified target nucleic acid. In some embodiments, the method may further comprise regenerating the plant cell comprising the modified target nucleic acid to produce a plant comprising the modified target nucleic acid.

In some embodiments, an expression cassette of the invention may be codon optimized for expression in an animal, e.g., a mammal. In some embodiments, the expression cassettes of the invention may be used in a method of modifying a target nucleic acid in an animal cell (e.g., a mammalian cell), the method comprising introducing one or more expression cassettes of the invention into a animal cell, thereby modifying the target nucleic acid in the animal cell to produce a animal cell comprising the modified target nucleic acid.

A CRISPR Cas9 polypeptide or CRISPR Cas9 domain (e.g., a Type II CRISPR Case effector protein) useful with this invention may be any known or later identified Cas9 nuclease. In some embodiments, a CRISPR Cas9 polypeptide can be a Cas9 polypeptide from, for example, *Streptococcus* spp. (e.g., *S. pyogenes*, *S. thermophilus*) (e.g., spCas9), *Lactobacillus* spp., *Bifidobacterium* spp., *Kandleria* spp., *Leuconostoc* spp., *Oenococcus* spp., *Pediococcus* spp., *Weissella* spp., and/or *Olsenella* spp.

Cas12a is a Type V Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR)-Cas effector protein or domain. Cas12a differs in several respects from the more well-known Type II CRISPR Cas9 effector protein. For example, Cas9 recognizes a G-rich protospacer-adjacent motif (PAM) that is 3' to its guide RNA (gRNA, sgRNA) binding site (protospacer, target nucleic acid, target DNA) (3'-NGG), while Cas12a recognizes a T-rich PAM that is located 5' to the target nucleic acid (5'-TTN, 5'-TTTN. In fact, the orientations in which Cas9 and Cas12a bind their guide RNAs are very nearly reversed in relation to their N and C termini. Furthermore, Cas12a effector proteins use a single guide RNA (gRNA, CRISPR array, crRNA) rather than the dual guide RNA (sgRNA (e.g., crRNA and tracrRNA)) found in natural Cas9 systems, and Cas12a processes its own gRNAs. Additionally, nuclease activity of a Cas12a produces staggered DNA double stranded breaks

instead of blunt ends produced by nuclease activity of a Cas9, and Cas12a relies on a single RuvC domain to cleave both DNA strands, whereas Cas9 utilizes an HNH domain and a RuvC domain for cleavage.

A CRISPR Cas12a effector protein or domain useful with this invention may be any known or later identified Cas12a nuclease (previously known as Cpf1) (see, e.g., U.S. Patent No. 9,790,490, which is incorporated by reference for its disclosures of Cpf1 (Cas12a) sequences). The term "Cas12a", "Cas12a polypeptide" or "Cas12a domain" refers to an RNA-guided effector protein comprising a Cas12a, or a fragment thereof, which comprises the guide nucleic acid binding domain of Cas12a and/or an active, inactive, or partially active DNA cleavage domain of Cas12a. In some embodiments, a Cas12a useful with the invention may comprise a mutation in the nuclease active site (e.g., RuvC site of the Cas12a domain). A Cas12a effector protein or domain having a mutation in its nuclease active site, and therefore, no longer comprising nuclease activity, is commonly referred to as dead or deactivated Cas12a (e.g., dCas12a).

In some embodiments, a Cas12a effector polypeptide that may be optimized or otherwise modified (e.g., deactivate) according to the present invention can include, but is not limited to, the amino acid sequence of any one of **SEQ ID NOs:1-20** (e.g., **SEQ ID NOs: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19** or **20**), or **SEQ ID NOs:148, 149, 150** or **151**, or a polynucleotide encoding the same.

In some embodiments, a Cas9 effector polypeptide that may be optimized or otherwise modified (e.g., deactivate) according to the present invention can include, but is not limited to, the amino acid sequence of any one of **SEQ ID NO:106** or **SEQ ID NO:107**, or a polynucleotide encoding the same. In some embodiments, a Cas9 effector polypeptide that may be optimized or otherwise modified (e.g., deactivate) according to the present invention can comprise an amino acid sequence encoded by any one of the nucleic acid sequences of **SEQ ID NOs:108-122**.

A "guide nucleic acid," "guide RNA," "gRNA," "CRISPR RNA/DNA" "crRNA" or "crDNA" as used herein means a nucleic acid that comprises at least one spacer sequence, which is complementary to (and hybridizes to) a target DNA (e.g., protospacer), and at least one repeat sequence that corresponds to a particular CRISPR-Cas effector protein (e.g., for a Type V CRISPR Cas effector protein, the repeat or a fragment or portion thereof is from a Type V Cas12a CRISPR-Cas system; for a Type II CRISPR Cas effector protein, the repeat or a fragment or portion thereof is from a Type II Cas9 CRISPR-Cas system). Thus, a repeat of a CRISPR-Cas system useful with the present invention may correspond to the CRISPR-

Cas effector protein of, for example, Cas9, C2c3, Cas12a (also referred to as Cpf1), Cas12b, Cas12c, Cas12d, Cas12e, Cas13a, Cas13b, Cas13c, Cas13d, Cas1, Cas1B, Cas2, Cas3, Cas3', Cas3'', Cas4, Cas5, Cas6, Cas7, Cas8, Cas9 (also known as Csn1 and Csx12), Cas10, Csy1, Csy2, Csy3, Cse1, Cse2, Cse3, Cse4, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx15, Csf1, Csf2, Csf3, Csf4 (dinG), and/or Csf5, or a fragment thereof, wherein the repeat sequence may be linked to the 5' end and/or the 3' end of the spacer sequence. The design of a guide nucleic acid of this invention may be based on a Type I, Type II, Type III, Type IV, or Type V CRISPR-Cas system. In some embodiments, the design of a guide nucleic acid of this invention is based on a Type V CRISPR-Cas system. In some embodiments, the design of a guide nucleic acid of this invention is based on a Type II CRISPR-Cas system.

In some embodiments, a guide nucleic acid (e.g., crRNA, e.g., Cas12a crRNA, Cas12b crRNA, Cas9 crRNA, and the like) may comprise, from 5' to 3', a repeat sequence (full length or portion thereof ("handle")); e.g., pseudoknot-like structure) and a spacer sequence. In some embodiments, an extended guide nucleic acid (e.g., tagRNA, e.g., Cas12a extended guide nucleic acid, Cas12b extended guide nucleic acid, Cas9 extended guide nucleic acid, and the like) may comprise, from 5' to 3', a repeat sequence (full length or portion thereof ("handle")); e.g., pseudoknot-like structure) a spacer sequence, plus a 3' or 5' extended portion comprising a primer binding site and a reverse transcriptase template (RT template) (RTT) (e.g., a tagRNA extension).

In some embodiments, a guide nucleic acid may comprise more than one repeat sequence-spacer sequence (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, or more repeat-spacer sequences) (e.g., repeat-spacer-repeat, e.g., repeat-spacer-repeat-spacer-repeat-spacer-repeat-spacer, and the like). The guide nucleic acids of this invention are synthetic, human-made and not found in nature. A guide nucleic acid may be quite long and may be used as an aptamer (like in the MS2 recruitment strategy) or other RNA structures hanging off the spacer. In some embodiments, as described herein, a guide nucleic acid may include a template for editing and a primer binding site. In some embodiments, a guide nucleic acid may include a region or sequence on its 5' end or 3' end that is complementary to an editing template (a reverse transcriptase template), thereby recruiting the editing template to the target nucleic acid (i.e., an extended guide nucleic acid). In some embodiments, a guide nucleic acid may include a region or sequence on its 5' end or 3' end that is complementary to

a primer on the target nucleic acid (a primer binding site), thereby recruiting the primer binding site to the target nucleic acid (i.e., an extended guide nucleic acid).

A “repeat sequence” as used herein, refers to, for example, any repeat sequence of a wild-type CRISPR Cas locus (e.g., a Cas9 locus, a Cas12a locus, a C2c1 locus, etc.) or a repeat sequence of a synthetic crRNA that is functional with the CRISPR-Cas nuclease encoded by the nucleic acid constructs of the invention. A repeat sequence useful with this invention can be any known or later identified repeat sequence of a CRISPR-Cas locus (e.g., Type I, Type II, Type III, Type IV, Type V or Type VI) or it can be a synthetic repeat designed to function in a Type I, II, III, IV, V or VI CRISPR-Cas system. Thus, in some embodiments, a repeat sequence can be identical to or substantially identical to a repeat sequence from wild-type Type I CRISPR-Cas loci, Type II, CRISPR-Cas loci, Type III, CRISPR-Cas loci, Type IV CRISPR-Cas loci, Type V CRISPR-Cas loci and/or Type VI CRISPR-Cas loci. In some embodiments, a repeat sequence useful with this invention can be any known or later identified repeat sequence of a Type V CRISPR-Cas locus or it can be a synthetic repeat designed to function in a Type V CRISPR-Cas system. A repeat sequence may comprise a hairpin structure and/or a stem loop structure. In some embodiments, a repeat sequence may form a pseudoknot-like structure at its 5' end (i.e., “handle”). Thus, in some embodiments, a repeat sequence can be identical to or substantially identical to a repeat sequence from wild type Type V CRISPR-Cas loci or wild type Type II CRISPR-Cas loci. A repeat sequence from a wild-type CRISPR-Cas locus may be determined through established algorithms, such as using the CRISPRfinder offered through CRISPRdb (see, Grissa et al. *Nucleic Acids Res.* 35 (Web Server issue):W52-7 or *BMC Informatics* 8:172 (2007)(doi:10.1186/1471-2105-8-172)). In some embodiments, a repeat sequence or portion thereof is linked at its 3' end to the 5' end of a spacer sequence, thereby forming a repeat-spacer sequence (e.g., guide RNA, crRNA).

In some embodiments, a repeat sequence comprises, consists essentially of, or consists of at least 10 nucleotides depending on the particular repeat and whether the guide RNA comprising the repeat is processed or unprocessed (e.g., about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50 to 100 or more nucleotides, or any range or value therein; e.g., about). In some embodiments, a repeat sequence comprises, consists essentially of, or consists of about 10 to about 20, about 10 to about 30, about 10 to about 45, about 10 to about 50, about 15 to about 30, about 15 to about 40, about 15 to about 45, about 15 to about

50, about 20 to about 30, about 20 to about 40, about 20 to about 50, about 30 to about 40, about 40 to about 80, about 50 to about 100 or more nucleotides.

A repeat sequence linked to the 5' end of a spacer sequence can comprise a portion of a repeat sequence (e.g., 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 or more contiguous nucleotides of a wild type repeat sequence). In some embodiments, a portion of a repeat sequence linked to the 5' end of a spacer sequence can be about five to about ten consecutive nucleotides in length (e.g., about 5, 6, 7, 8, 9, 10 nucleotides) and have at least 90% identity (e.g., at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more) to the same region (e.g., 5' end) of a wild type CRISPR Cas repeat nucleotide sequence. In some embodiments, a portion of a repeat sequence may comprise a pseudoknot-like structure at its 5' end (e.g., "handle").

A "spacer sequence" as used herein is a nucleotide sequence that is complementary to a target nucleic acid (e.g., target DNA) (e.g., protospacer). The spacer sequence can be fully complementary or substantially complementary (e.g., at least about 70% complementary (e.g., about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9% or more)) to a target nucleic acid. Thus, in some embodiments, the spacer sequence can have one, two, three, four, or five mismatches as compared to the target nucleic acid, which mismatches can be contiguous or noncontiguous. In some embodiments, the spacer sequence can have 70% complementarity to a target nucleic acid. In some embodiments, the spacer nucleotide sequence can have 80% complementarity to a target nucleic acid. In still other embodiments, the spacer nucleotide sequence can have 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9% or more complementarity, and the like, to the target nucleic acid (protospacer). In some embodiments, the spacer sequence is 100% complementary to the target nucleic acid. In some embodiments, a spacer sequence may have a length from about 15 nucleotides to about 30 nucleotides (e.g., 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides, or any range or value therein). Thus, in some embodiments, a spacer sequence may have complete complementarity or substantial complementarity over a region of a target nucleic acid (e.g., protospacer) that is at least about 15 nucleotides to about 30 nucleotides in length. In some embodiments, the spacer is about 20 nucleotides in length. In some embodiments, the spacer is about 23 nucleotides in length.

In some embodiments, the 5' region of a spacer sequence of a guide RNA may be identical to a target DNA, while the 3' region of the spacer may be substantially

complementary to the target DNA (e.g., Type V CRISPR-Cas), or the 3' region of a spacer sequence of a guide RNA may be identical to a target DNA, while the 5' region of the spacer may be substantially complementary to the target DNA (e.g., Type II CRISPR-Cas), and therefore, the overall complementarity of the spacer sequence to the target DNA may be less than 100%. Thus, for example, in a guide for a Type V CRISPR-Cas system, the first 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 nucleotides in the 5' region (i.e., seed region) of, for example, a 20-nucleotide spacer sequence may be 100% complementary to the target DNA, while the remaining nucleotides in the 3' region of the spacer sequence are substantially complementary (e.g., at least about 70% complementary) to the target DNA. In some embodiments, the first 1 to 8 nucleotides (e.g., the first 1, 2, 3, 4, 5, 6, 7, 8, nucleotides, and any range therein) of the 5' end of the spacer sequence may be 100% complementary to the target DNA, while the remaining nucleotides in the 3' region of the spacer sequence are substantially complementary (e.g., at least about 50% complementary (e.g., 50%, 55%, 60%, 65%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9% or more)) to the target DNA.

As a further example, in a guide for a Type II CRISPR-Cas system, the first 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 nucleotides in the 3' region (i.e., seed region) of, for example, a 20-nucleotide spacer sequence may be 100% complementary to the target DNA, while the remaining nucleotides in the 5' region of the spacer sequence are substantially complementary (e.g., at least about 70% complementary) to the target DNA. In some embodiments, the first 1 to 10 nucleotides (e.g., the first 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 nucleotides, and any range therein) of the 3' end of the spacer sequence may be 100% complementary to the target DNA, while the remaining nucleotides in the 5' region of the spacer sequence are substantially complementary (e.g., at least about 50% complementary (e.g., at least about 50%, 55%, 60%, 65%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9% or more or any range or value therein)) to the target DNA.

In some embodiments, a seed region of a spacer may be about 8 to about 10 nucleotides in length, about 5 to about 6 nucleotides in length, or about 6 nucleotides in length.

In some embodiments, an extended guide nucleic acid of this invention may be an extended guide nucleic acid, a first extended guide nucleic acid and/or a second extended guide nucleic acid. In some embodiments, an extended guide nucleic acid useful with this

invention may comprise: (a) a CRISPR nucleic acid (e.g., CRISPR RNA, CRISPR DNA, crRNA, crDNA) and/or a CRISPR nucleic acid and a tracr nucleic acid; and (b) an extended portion comprising a primer binding site and a reverse transcriptase template (RT template), wherein the RT template encodes a modification to be incorporated into the target nucleic acid as described herein (e.g., encodes an edit located in any position within an RT template with the position location relative to the position of a protospacer adjacent motif (PAM) of the target nucleic acid, optionally an edit located at nucleotide position -1 to nucleotide position 19, nucleotide position 10 to nucleotide position 17, or nucleotide position 12 to nucleotide position 15). In some embodiments, a CRISPR nucleic acid may be a Type II or Type V CRISPR nucleic acid and/or a tracr nucleic acid may be any tracr corresponding to the appropriate Type II or Type V CRISPR nucleic acid. An extended guide nucleic acid may also be referred to as a targeted allele guide nucleic acid, a targeted allele guide DNA, a targeted allele guide RNA (tagRNA)). In some embodiments, a CRISPR nucleic acid useful with the invention may be a Type V CRISPR nucleic acid. In some embodiments, a tracr nucleic acid useful with the invention may be a Type V CRISPR tracr nucleic acid. In some embodiments, a CRISPR nucleic acid useful with the invention may be a Type II CRISPR nucleic acid. In some embodiments, a tracr nucleic acid useful with the invention may be a Type II CRISPR tracr nucleic acid. In some embodiments, a CRISPR nucleic acid and/or tracr nucleic acid may be from, for example, a Cas9, C2c3, Cas12a (also referred to as Cpf1), Cas12b, Cas12c, Cas12d, Cas12e, Cas13a, Cas13b, Cas13c, Cas13d, Cas1, Cas1B, Cas2, Cas3, Cas3', Cas3'', Cas4, Cas5, Cas6, Cas7, Cas8, Cas9 (also known as Csn1 and Csx12), Cas10, Csy1, Csy2, Csy3, Cse1, Cse2, Cse1, Cse2, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx15, Csf1, Csf2, Csf3, Csf4 (dinG), and/or Csf5 system.

In some embodiments, an extended portion of the extended guide may comprise, 5' to 3', an RT template and a primer binding site (when the extended guide is linked to the 3' end of the CRISPR nucleic acid). In some embodiments, an extended portion of the extended guide may comprise, 5' to 3', a primer binding site and an RT template (RTT) (when the extended guide is linked to the 5' end of the CRISPR nucleic acid). In some embodiments, an RT template may be a length of about 1 nucleotide to about 100 nucleotides (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 or more

nucleotides, and any range or value therein), e.g., about 35 nucleotide to about 100 nucleotides, about 35 nucleotide to about 80 nucleotides, about 35 nucleotide to about 75 nucleotides, about 40 nucleotides to about 75 nucleotides, about 45 nucleotides to about 75 nucleotides, about 45 nucleotides to about 60 nucleotides in length and any range or value

5 therein. In some embodiments, the length of an RT template may be at least 30 nucleotides, optionally about 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 nucleotides in length to about to about 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, or 80 nucleotides in length, or any range or value therein. In some embodiments, the length of an RT template may be

10 about 836, 40, 44, 47, 50, 52, 55, 63, 72 or 74 nucleotides. Within the length of the RTT is comprised an edit. The edit may be located anywhere within the RTT, wherein the position of the edit may be described relative to the position of a protospacer adjacent motif (PAM) of the target nucleic acid. In some embodiments, an RT template may comprise an edit located at nucleotide position -1, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, or 19. In

15 some embodiments, an RT template may comprise an edit located at nucleotide position 4 to nucleotide position 17 (e.g., position 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, or 17) of the RT template relative to the position of a protospacer adjacent motif (PAM) of the target nucleic acid. In some embodiments, an RT template may comprise an edit located at nucleotide position 10 to nucleotide position 17 (e.g., position 10, 11, 12, 13, 14, 15, 16, or

20 17) of the RT template relative to the position of a protospacer adjacent motif (PAM) of the target nucleic acid. In some embodiments, an RT template may comprise an edit located at nucleotide position 12 to nucleotide position 15 (e.g., position 12, 13, 14, or 15) of the RT template relative to the position of a protospacer adjacent motif (PAM) of the target nucleic acid.

25 As used herein, a "primer binding site" (PBS) of an extended portion of an extended guide nucleic acid (e.g., tagRNA) refers to a sequence of consecutive nucleotides that can bind to a region or "primer" on a target nucleic acid, i.e., is complementary to the target nucleic acid primer. As an example, a CRISPR Cas effector protein (e.g., Type II or Type V, e.g., Cas 9 or Cas12a) nicks/cuts the DNA, the 3' end of the cut DNA acts as a primer for the

30 PBS portion of the extended guide nucleic acid. The PBS is designed to be complementary to the 3'end of a strand of the target nucleic acid and can be designed to bind either to the target strand or non-target strand. A primer binding site can be fully complementary to the primer or it may be substantially complementary (e.g., at least 70% complementary (e.g., about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%,

- 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9% or more)) to the primer on the target nucleic acid. In some embodiments, the length of a primer binding site of an extended portion may be about 1 nucleotide to about 100 nucleotides in length (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 or more nucleotides, or any value or range therein), or about 4 nucleotide to about 85 nucleotides, about 10 nucleotide to about 80 nucleotides, about 20 nucleotide to about 80 nucleotides, about 25 nucleotides to about 80 nucleotides about 30 nucleotide to about 80 nucleotides, about 40 nucleotide to about 80 nucleotides, about 45 nucleotide to about 80 nucleotides, about 45 nucleotide to about 75 nucleotides or about 45 nucleotide to about 60 nucleotides, or any range or value therein. In some embodiments, the length of an PBS may be at least 30 nucleotides, optionally about 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 nucleotides to about 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, or 80 nucleotides in length, or any range or value therein. In some embodiments, the length of a PBS may be about 8, 16, 24, 32, 40, 48, 56, 64, 72, or 80 nucleotides.
- 20 In some embodiments, an RTT may have a length of about 35 nucleotides to about 75 nucleotides and a PBS may have a length of about 30 nucleotides to about 80 nucleotides, optionally wherein the PBS may comprise a length of about 8, 16, 24, 32, 40, 48, 56, 64, 72, or 80 nucleotides and the RTT may comprise a length of about 36, 40, 44, 47, 50, 52, 55, 63, 72 or 74 nucleotides, or any combination thereof of the RTT length and/or PBS length.
- 25 In some embodiments, an extended guide nucleic (e.g., extended guide nucleic acid, first extended guide nucleic acid, second extended guide nucleic acid) may comprise a structured RNA motif, optionally wherein the structured RNA motif may be located at the 3' end of the extended guide nucleic acid. In some embodiments, the structured RNA motif can include, but is not limited to, AsCpf1BB (SEQ ID NO:189), BoxB (SEQ ID NO:190), pseudoknot (decoy) (SEQ ID NO:95, SEQ ID NO:203), pseudoknot (tEvoPreQ1) (SEQ ID NO:191), fmpknot (SEQ ID NO:192), mpknot (SEQ ID NO:193), MS2 (SEQ ID NO:194), PP7 (SEQ ID NO:195), SLBP (SEQ ID NO:196), TAR (SEQ ID NO:197), and/or ThermoPh (SEQ ID NO:198). In some embodiments, a structured RNA motif can be a pseudoknot, optionally wherein the pseudoknot is located at the 3' end of the extended guide

nucleic acid. Pseudoknots are RNA structural motifs formed upon base pairing of a single-stranded region of RNA in the loop of a hairpin to a stretch of complementary nucleotides elsewhere in the RNA chain. In some embodiments, a pseudoknot useful with the invention may be a naturally occurring pseudoknot or a synthetic pseudoknot. The term pseudoknot, as used herein, includes, but is not limited to, hairpins, multiloops, kissing loops, coaxial stacking, triplexes, pseudoknot-like structures, a pseudoknotted hairpins and/or a decoy pseudoknotted hairpins or other RNA structural motifs. In some embodiments, the pseudoknot may be located at the 3' end of the extended guide nucleic acid. In some embodiments, when the extended guide comprises 5'-3' crRNA-RTT-PBS, a pseudoknot may be located 5' of the RTT or 3' of the PBS. In some embodiments, the pseudoknot may be located at the 3' end of the extended guide nucleic acid. In some embodiments, when the extended guide comprises the extension (extended portion) at the 5' end of the crRNA, a pseudoknot may be located 3' of the RTT or 5' of the PBS. In some embodiments, a pseudoknot useful with an extended guide can include, but is not limited to, a tEvoPreQ1 Pseudoknot comprising the nucleic acid sequence of UAAUUUCUACUAAGUGUAGAU (SEQ ID NO:158), a pseudoknot EvoPreQ1 comprising the nucleic acid sequence of TTGACGCGTTCTATCTAGTTACGCGTTAAACCAACUAGAAA (SEQ ID NO:191) or a pseudoknot comprising the nucleic acid sequence of TAAGTCTCCATAGAATGGAGG (SEQ ID NO:95) and/or UAAGUCUCCAUAAGAAUGGAGG (SEQ ID NO:203). An extended guide nucleic acid of this invention may be comprised in an expression cassette, optionally wherein the expression cassette is comprised in a vector.

In some embodiments, an extended portion of an extended guide may be fused to either the 5' end or 3' end of a Type II or a Type V CRISPR nucleic acid (e.g., 5' to 3': repeat-spacer-extended portion, or extended portion-repeat-spacer) and/or to the 5' or 3' end of the tracr nucleic acid. In some embodiments, when an extended portion is located 5' of the crRNA, the Type V CRISPR-Cas effector protein is modified to reduce (or eliminate) self-processing RNase activity. In some embodiments, a Type V CRISPR-Cas effector protein that is modified to reduce (or eliminate) self-processing RNase activity may be utilized also when the extended portion is located 3' of the crRNA.

In some embodiments, the extended portion of an extended guide nucleic acid may be linked to the Type II or Type V CRISPR nucleic acid and/or the Type II or Type V tracrRNA via a linker. In some embodiments, a linker may be a length of about 1 to about 100 nucleotides or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44,

45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 or more nucleotides in length, and any range therein (e.g., about 2 to about 40, about 2 to about 50, about 2 to about 60, about 4 to about 40, about 4 to about 50, about 4 to about 60, about 5 to about 40, about 5 to about 50, about 5 to about 60, about 9 to about 40, about 9 to about 50, about 9 to about 60, about 10 to about 40, about 10 to about 50, about 10 to about 60, about 40 to about 100, about 50 to about 100, or about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 nucleotides to about 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 or more nucleotides in length (e.g., about 105, 110, 115, 120, 130, 140 150 or more nucleotides in length).

As used herein, a “target nucleic acid”, “target DNA,” “target nucleotide sequence,” “target region,” or a “target region in the genome” refers to a region of an organism's genome that is fully complementary (100% complementary) or substantially complementary (e.g., at least 70% complementary (e.g., 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more)) to a spacer sequence in a guide RNA of this invention (e.g., the spacer is substantially complementary to the target strand of the target nucleic acid). A target region useful for a CRISPR-Cas system may be located immediately 3' (e.g., Type V CRISPR-Cas system) or immediately 5' (e.g., Type II CRISPR-Cas system) to a PAM sequence in the genome of the organism (e.g., a plant genome). A target region may be selected from any region of at least 15 consecutive nucleotides (e.g., 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 nucleotides, and the like) located immediately adjacent to a PAM sequence on the target strand.

A “protospacer sequence” refers to the target double stranded DNA and specifically to the portion of the target nucleic acid/target DNA (e.g., or target region in the genome (e.g., nuclear genome, plastid genome, mitochondrial genome), or an extragenomic sequence, such as a plasmid, minichromosome, and the like) that is fully or substantially complementary (and hybridizes) to the spacer sequence of the CRISPR repeat-spacer sequences (e.g., guide RNAs, CRISPR arrays, crRNAs). Thus, the protospacer sequences is complementary to the target strand of the target nucleic acid. In some embodiments, a target nucleic acid may have a first strand and a second strand (double stranded DNA). In some embodiments, the term

"first strand" as used herein in reference to a target nucleic acid may refer to a target strand or a bottom strand. In some embodiments, the term "second strand" as used in reference to a target nucleic acid is the strand that is complementary to the first strand (e.g., top strand or non-target strand).

5 As understood in the art and as used herein, a "target strand" refers to the strand of a double stranded DNA to which the spacer is complementary and to which the CRISPR-Cas effector protein is recruited, while the "non-target strand" refers to the strand opposite to the target strand in a double stranded nucleic acid. In some embodiments of the present invention, the non-target strand of a double stranded nucleic acid, the strand opposite of the strand to which the CRISPR-Cas effector protein is recruited, is nicked by the CRISPR-Cas effector protein and is edited by the reverse transcriptase. In some embodiments, the target strand of a double stranded nucleic acid, the same strand to which the CRISPR-Cas effector protein is recruited, is nicked by CRISPR-Cas effector protein and is edited by the reverse transcriptase.

15 In the case of Type V CRISPR-Cas (e.g., Cas12a) systems and Type II CRISPR-Cas (Cas9) systems, the protospacer sequence is flanked by (e.g., immediately adjacent to) a protospacer adjacent motif (PAM). For Type IV CRISPR-Cas systems, the PAM is located at the 5' end on the non-target strand and at the 3' end of the target strand (see below, as an example).

20 5'-NNNNNNNNNNNNNNNNNNNN-3' RNA Spacer (SEQ ID NO:54)
 | | | | | | | | | | | | | | | |
 3'AAANNNNNNNNNNNNNNNNNNNN-5' Target strand (SEQ ID NO:55)
 | | | |
 5'TTTNNNNNNNNNNNNNNNNNNNN-3' Non-target strand (SEQ ID NO:56)

25 In the case of Type II CRISPR-Cas (e.g., Cas9) systems, the PAM is located immediately 3' of the target region. The PAM for Type I CRISPR-Cas systems is located 5' of the target strand. There is no known PAM for Type III CRISPR-Cas systems. Makarova et al. describes the nomenclature for all the classes, types and subtypes of CRISPR systems (Nature Reviews Microbiology 13:722–736 (2015)). Guide structures and PAMs are described in by R. Barrangou (Genome Biol. 16:247 (2015)).

30 Canonical Cas12a PAMs are T rich. In some embodiments, a canonical Cas12a PAM sequence may be 5'-TTN, 5'-TTTN, or 5'-TTTV. In some embodiments, canonical Cas9 (e.g., *S. pyogenes*) PAMs may be 5'-NGG-3'. In some embodiments, non-canonical PAMs may be used but may be less efficient.

Additional PAM sequences may be determined by those skilled in the art through established experimental and computational approaches. Thus, for example, experimental approaches include targeting a sequence flanked by all possible nucleotide sequences and identifying sequence members that do not undergo targeting, such as through the transformation of target plasmid DNA (Esvelt et al. 2013. *Nat. Methods* 10:1116-1121; Jiang et al. 2013. *Nat. Biotechnol.* 31:233-239). In some aspects, a computational approach can include performing BLAST searches of natural spacers to identify the original target DNA sequences in bacteriophages or plasmids and aligning these sequences to determine conserved sequences adjacent to the target sequence (Briner and Barrangou. 2014. *Appl. Environ. Microbiol.* 80:994-1001; Mojica et al. 2009. *Microbiology* 155:733-740).

In some embodiments, the present invention further provides a method of modifying a target nucleic acid, the method comprising: contacting the target nucleic acid at a first site with (a)(i) a first CRISPR-Cas effector protein; and (ii) a first extended guide nucleic acid (e.g., first extended CRISPR RNA, first extended CRISPR DNA, first extended crRNA, first extended crDNA); and (b)(i) a second CRISPR-Cas effector protein, (ii) a first reverse transcriptase; and (ii) a first guide nucleic acid, thereby modifying the target nucleic acid. In some embodiments, the method of the invention may further comprise contacting the target nucleic acid with (a) a third CRISPR-Cas effector protein; and (b) a second guide nucleic acid, wherein the third CRISPR-Cas effector protein nicks a site on the first strand of the target nucleic acid that is located about 10 to about 125 base pairs (either 5' or 3') from the second site on the second strand that has been nicked by the second CRISPR-Cas effector protein, thereby improving mismatch repair. In some embodiments, the method of the invention may further comprise contacting the target nucleic acid with: (a) a fourth CRISPR-Cas effector protein; (b) a second reverse transcriptase, and (c) a second extended guide nucleic acid (e.g., second extended CRISPR RNA, second extended CRISPR DNA, second extended crRNA, second extended crDNA), wherein the second extended guide nucleic acid targets (spacer is substantially complementary to/binds to) a site on the first strand of the target nucleic acid, thereby modifying the target nucleic acid. A CRISPR-Cas effector protein (e.g., a first, second, third, fourth) useful with the invention may be any Type I, Type II, Type III, Type IV, or Type V CRISPR-Cas effector protein as described herein, in any combination. In some embodiments, the CRISPR-Cas effector protein may be Cas9, C2c3, Cas12a (also referred to as Cpf1), Cas12b, Cas12c, Cas12d, Cas12e, Cas13a, Cas13b, Cas13c, Cas13d, Cas1, Cas1B, Cas2, Cas3, Cas3', Cas3'', Cas4, Cas5, Cas6, Cas7, Cas8, Cas9 (also known as Csn1 and Csx12), Cas10, Csy1, Csy2, Csy3, Cse1, Cse2, Cse1, Cse2, Csa5,

Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx15, Csf1, Csf2, Csf3, Csf4 (dinG), and/or Csf5.

In some embodiments, an extended guide nucleic acid useful with the first CRISPR-Cas effector protein may comprise (a) a CRISPR nucleic acid (CRISPR RNA, CRISPR DNA, crRNA, crDNA); and (b) an extended portion comprising a primer binding site and a reverse transcriptase template (RT template), wherein the RT template encodes a modification to be incorporated into the target nucleic acid as described herein (e.g., encodes an edit located in any position within an RT template with the position location relative to the position of a protospacer adjacent motif (PAM) of the target nucleic acid, optionally an edit located at nucleotide position -1 to nucleotide position 19, nucleotide position 10 to nucleotide position 17, or nucleotide position 12 to nucleotide position 15).

In some embodiments, the CRISPR nucleic acid of the extended guide nucleic acid comprises a spacer sequence capable of binding to (having substantial homology to) a first site on the first strand of the target nucleic acid.

In some embodiments, a guide nucleic acid useful with a CRISPR-Cas effector protein comprises a CRISPR nucleic acid (CRISPR RNA, CRISPR DNA, crRNA, crDNA). In some embodiments, the CRISPR nucleic acid of the first guide nucleic acid comprises a spacer sequence that binds to a second site on the first strand of the target nucleic acid that is upstream (3') of the first site on the first strand of the target nucleic acid.

In some embodiments, the second CRISPR-Cas effector protein may be a CRISPR-Cas fusion protein comprising a CRISPR-Cas effector protein domain fused to the reverse transcriptase.

In some embodiments, the second CRISPR-Cas effector protein may be a CRISPR-Cas fusion protein comprising a CRISPR-Cas effector protein domain fused to a peptide tag and the reverse transcriptase may be a reverse transcriptase fusion protein comprising a reverse transcriptase domain that is fused to an affinity polypeptide capable of binding the peptide tag.

In some embodiments, the first guide nucleic acid may be linked to an RNA recruiting motif and the reverse transcriptase may be a reverse transcriptase fusion protein comprising a reverse transcriptase domain that is fused to an affinity polypeptide capable of binding the RNA recruiting motif.

In some embodiments, the target nucleic acid may further be contacted with a 5'-3' exonuclease, optionally wherein the 5'-3' exonuclease is fused to the first CRISPR-Cas

effector protein. In some embodiments, a 5'-3' exonuclease may be a fusion protein comprising a 5'-3' exonuclease fused to a peptide tag and the first CRISPR-Cas effector protein may be a fusion protein comprising a CRISPR-Cas effector protein domain fused to an affinity polypeptide that is capable of binding to the peptide tag. In some embodiments, a 5'-3' exonuclease may be a fusion protein comprising a 5'-3' exonuclease fused to an affinity polypeptide that is capable of binding to the peptide tag and the first CRISPR-Cas effector protein may be a fusion protein comprising a CRISPR-Cas effector protein domain fused to a peptide tag. In some embodiments, a 5'-3' exonuclease may be a fusion protein comprising a 5'-3' exonuclease that is fused to an affinity polypeptide that is capable of binding to an RNA recruiting motif and the extended guide nucleic acid is linked to an RNA recruiting motif.

In some embodiments, the invention further provides contacting a target nucleic acid with one or more single stranded DNA binding proteins (ssDNA BPs). Single-stranded DNA binding proteins (ssDNA BP) may be useful for stabilizing the single stranded DNAs that are generated during the methods of the invention. Without wishing to be bound by any particular theory, ssDNA BPs may protect DNA strands from degradation or otherwise prevent them from becoming unavailable for RT-mediated priming and polymerization. Single stranded DNA binding proteins useful with the invention can include but are not limited to, those obtained from Example ssDNA BPs include, but are not limited to, those from a human, a bacterium or a phage. In some embodiments an ssDNA BP includes, but is not limited to, hRad51 (optionally, hRad51_S208E_A209D)(SEQ ID NO:123), hRad52 (SEQ ID NO:124), BsRecA (SEQ ID NO:125), EcRecA (SEQ ID NO:126), T4ssB (SEQ ID NO:127) and/or Brex27 (SEQ ID NO:128). In some embodiments, a target nucleic acid may be contacted with one or more ssDNA BPs, wherein the ssDNA BPs may be fused to the C-terminus or the N-terminus of a CRISPR-Cas effector protein (e.g., a CRISPR-Cas effector protein, a first CRISPR-Cas effector protein, a second CRISPR-Cas effector protein, a third CRISPR-Cas effector protein and/or a fourth CRISPR-Cas effector protein). A ssDNA BP may be fused to the C-terminus or the N-terminus of the CRISPR-Cas effector protein/domain. In some embodiments, the ssDNA BP is fused to a Type II CRISPR-Cas effector protein/domain and/or a Type V CRISPR-Cas effector protein/domain.

In some embodiments, the methods of the invention may further comprise reducing double strand breaks by introducing a chemical inhibitor of non-homologous end joining (NHEJ), by introducing a CRISPR guide nucleic acid or an siRNA targeting an NHEJ protein to transiently knock-down expression of the NHEJ protein, or by introducing a polypeptide that prevents NHEJ. In some embodiments, the polypeptide that prevents NHEJ can include,

but is not limited to, a Gam protein, optionally wherein the Gam protein is *Escherichia* phage Mu Gam protein (e.g., **SEQ ID NO:147**).

- In some embodiments, an extended guide nucleic acid is provided, the extended guide nucleic acid comprising (i) a Type V CRISPR nucleic acid or Type II CRISPR nucleic acid (Type II or Type V CRISPR RNA, Type II or Type V CRISPR DNA, Type II or Type V crRNA, Type II or Type V crDNA) and/or a Type V CRISPR nucleic acid or Type II CRISPR nucleic acid and a tracr nucleic acid (e.g., Type II or Type V tracrRNA, Type II or Type V tracrDNA); and (ii) an extended portion comprising a primer binding site and a reverse transcriptase template (RT template) (RTT). In some embodiments, the extended guide nucleic acid further comprise a structured RNA motif, optionally wherein the structured RNA motif is located at the 3' end of the extended guide nucleic acid. In some embodiments, the structured RNA motif can include, but is not limited to, AsCpf1BB (**SEQ ID NO:189**), BoxB (**SEQ ID NO:190**), pseudoknot (decoy) (**SEQ ID NO:95**, **SEQ ID NO:203**), pseudoknot (tEvoPreQ1) (**SEQ ID NO:191**), fmpknot (**SEQ ID NO:192**), mpknot (**SEQ ID NO:193**), MS2 (**SEQ ID NO:194**), PP7 (**SEQ ID NO:195**), SLBP (**SEQ ID NO:196**), TAR (**SEQ ID NO:197**), and/or ThermoPh (**SEQ ID NO:198**). In some embodiments, the structured RNA motif is a pseudoknot, optionally wherein the pseudoknot is located at the 3' end of the extended guide nucleic acid. In some embodiments, a pseudoknot useful with the invention may be a naturally occurring pseudoknot or a synthetic pseudoknot. A pseudoknot may also be referred to herein as a pseudoknot-like structure, a pseudoknotted hairpin and/or a decoy pseudoknotted hairpin. In some embodiments, the pseudoknot may be located at the 3' end of the extended guide nucleic acid. In some embodiments, when the extended guide comprises 5'-3' crRNA-RTT-PBS, the pseudoknot may be located 5' of the RTT or 3' of the PBS. . In some embodiments, when the extended guide comprises the extension (extended portion) at the 5' end of the crRNA, a pseudoknot may be located 3' of the RTT or 5' of the PBS. In some embodiments, a pseudoknot may be located at the 5' end of an extended guide nucleic acid followed 5'-3' by the PBS then RTT, the natural pseudoknot in the crRNA (e.g., in the repeat sequence), followed by the complimentary region (e.g., spacer sequence).
- In some embodiments, a pseudoknot useful with the extended guide can include, but is not limited to, a tEvoPreQ1 Pseudoknot comprising the nucleic acid sequence of **SEQ ID NO:158**, an EvoPreQ1 Pseudoknot comprising the nucleic acid sequence of **SEQ ID NO:191** and/or a pseudoknot comprising the nucleic acid sequence of **SEQ ID NO:95** or **SEQ ID**

NO:203. An extended guide nucleic acid of this invention may be comprised in an expression cassette, optionally wherein the expression cassette is comprised in a vector.

In some embodiments, a complex is provided, the complex comprising: (a) a Type II CRISPR-Cas effector protein or a Type V CRISPR-Cas effector protein; (b) a reverse transcriptase, and (c) an extended guide nucleic acid (e.g., extended CRISPR RNA, extended CRISPR DNA, extended crRNA, extended crDNA; e.g., a tagDNA, tagRNA).

In some embodiments, the Type II or Type V CRISPR-Cas effector protein of a complex may be a fusion protein comprising a Type II or Type V CRISPR-Cas effector protein domain fused to a peptide tag. In some embodiments, the Type II or Type V CRISPR-Cas effector protein of the complex may be a fusion protein comprising a Type II or Type V CRISPR-Cas effector protein domain fused to an affinity polypeptide that is capable of binding a peptide tag. In some embodiments, the Type II or Type V CRISPR-Cas effector protein of the complex may be a fusion protein comprising a Type II or Type V CRISPR-Cas effector protein domain fused to an affinity polypeptide that is capable of binding an RNA recruiting motif.

In some embodiments, the reverse transcriptase of the complex may be a fusion protein comprising a reverse transcriptase domain fused to a peptide tag. In some embodiments, the reverse transcriptase of the complex may be a fusion protein comprising reverse transcriptase domain fused to an affinity polypeptide that is capable of binding a peptide tag. In some embodiments, the reverse transcriptase of the complex may be a fusion protein comprising reverse transcriptase domain fused to an affinity polypeptide that is capable of binding an RNA recruiting polypeptide. In some embodiments, the complex may further comprise a guide nucleic acid (e.g., extended CRISPR RNA, extended CRISPR DNA, extended crRNA, extended crDNA). In some embodiments, the complex may further comprise an extended guide nucleic acid (e.g., extended CRISPR RNA, extended CRISPR DNA, extended crRNA, extended crDNA).

In some embodiments, the extended guide nucleic acid of the complex may further comprise a pseudoknot. In some embodiments, the pseudoknot comprised in the extended guide nucleic acid of the complex may be located at the 3' end of the extended guide nucleic acid. In some embodiments, a pseudoknot useful with an extended guide nucleic acid of a complex of the invention may be a naturally occurring pseudoknot or a synthetic pseudoknot. A pseudoknot may also be referred to herein as a pseudoknot-like structure, a pseudoknotted hairpin and/or a decoy pseudoknotted hairpin. In some embodiments, the pseudoknot may be located at the 3' end of the extended guide nucleic acid. In some embodiments, when the

extended guide comprises 5'-3' crRNA-RTT-PBS, the pseudoknot may be located 5' of the RTT or 3' of the PBS. In some embodiments, a pseudoknot can include, but is not limited to, a tEvoPreQ1 Pseudoknot comprising the nucleic acid sequence of **SEQ ID NO:158**, an EvoPreQ1 Pseudoknot comprising the nucleic acid sequence of **SEQ ID NO:191** or a
 5 pseudoknot comprising the nucleic acid sequence of **SEQ ID NO:95** or **SEQ ID NO:203**.

In some embodiments, a complex of the invention may be comprised in an expression cassette, optionally wherein the expression cassette is comprised in a vector. In some embodiment, the expression cassette comprising a complex of the invention may be codon optimized for expression in an organism as described herein, optionally wherein the organism
 10 is wherein the organism is an animal such as a human, a plant, a fungus, an archaeon, a bacterium or a virus.

The present invention further provides an expression cassette codon optimized for expression in an organism, comprising 5' to 3' (a) polynucleotide encoding a promoter sequence, (b) a polynucleotide encoding a Type V CRISPR-Cas nuclease (e.g., Cpf1
 15 (Cas12a), dCas12a and the like) or a Type II CRISPR-Cas nuclease (e.g., Cas9, dCas9 and the like) that is codon optimized for expression in the organism; (c) a linker sequence; and (d) a polynucleotide encoding a reverse transcriptase that is codon-optimized for expression in the organism, optionally wherein the organism is wherein the organism is an animal such as a human, a plant, a fungus, an archaeon, a bacterium or a virus. Further provided is an
 20 expression cassette codon optimized for expression in a plant, comprising 5' to 3' (a) polynucleotide encoding a plant specific promoter sequence (e.g., ZmUbi1, MtUbi2, RNA polymerase II (Pol II)), (b) a plant codon-optimized polynucleotide encoding a Type V CRISPR-Cas nuclease (e.g., Cpf1 (Cas12a), dCas12a and the like); (c) a linker sequence; and (d) a plant codon-optimized polynucleotide encoding a reverse transcriptase. In some
 25 embodiments, a linker sequence may be an amino acid or peptide linker as described herein. In some embodiments, the reverse transcriptase in an expression cassette may be fused to one or more ssRNA binding domains (RBDs).

The present invention further provides an expression cassette codon optimized for expression in a plant, comprising (a) a polynucleotide encoding a plant specific promoter
 30 sequence (e.g. ZmUbi1, MtUbi2), and (b) an extended RNA guide sequence, wherein the extended guide nucleic acid comprises an extended portion comprising at its 3' end a primer binding site and an edit to be incorporated into the target nucleic acid (e.g., reverse transcriptase template), optionally wherein the extended guide nucleic acid is comprised in an

expression cassette, optionally wherein the extended guide nucleic acid is operably linked to a Pol II promoter..

In some embodiments, the expression cassette comprises an extended guide nucleic acid that further comprises a structured RNA motif, optionally wherein the structured RNA motif is located at the 3' end of the extended guide nucleic acid. In some embodiments, the structured RNA motif can include, but is not limited to, AsCpf1BB (SEQ ID NO:189), BoxB (SEQ ID NO:190), pseudoknot (decoy) (SEQ ID NO:95, SEQ ID NO:203), pseudoknot (tEvoPreQ1) (SEQ ID NO:191), fmpknot (SEQ ID NO:192), mpknot (SEQ ID NO:193), MS2 (SEQ ID NO:194), PP7 (SEQ ID NO:195), SLBP (SEQ ID NO:196), TAR (SEQ ID NO:197), and/or ThermoPh (SEQ ID NO:198). In some embodiments, the structured RNA motif is a pseudoknot, optionally wherein the pseudoknot is located at the 3' end of the extended guide nucleic acid. In some embodiments, a pseudoknot useful with the extended guide can include, but is not limited to, a pseudoknot comprising the nucleic acid sequence of SEQ ID NO:158, SEQ ID NO:191, SEQ ID NO:95 and/or SEQ ID NO:203.

In some embodiments, a plant specific promoter useful with an expression cassette of the invention may be associated with an intron or is a promoter region comprising an intron (e.g., ZmUbi1 comprising an intron; MtUb2 comprising an intron).

In some embodiments, the expression cassette may be codon optimized for expression in a dicot plant. In some embodiments, the expression cassette may be codon optimized for expression in a monocot plant.

In some embodiments, the present invention provides methods for modifying a target nucleic acid in a plant or plant cell, comprising introducing one or more expression cassettes of the invention into the plant or plant cell, thereby modifying the target nucleic acid in the plant or plant cell to produce a plant or plant cell comprising the modified target nucleic acid.

In some embodiments, the methods of the invention further comprise regenerating a plant from the plant cell comprising the modified target nucleic acid to produce a plant comprising the modified target nucleic acid. In some embodiments, the methods of the invention comprise contacting the target nucleic acid at a temperature of about 20°C to 42°C (e.g., about 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, or 42°C, and any value or range therein).

In some embodiments, the invention provides cells comprising one or more polynucleotides, guide nucleic acids, nucleic acid constructs, expression cassettes or vectors of the invention.

When used in combination with guide nucleic acids, the polynucleotides/nucleic acid constructs/expression cassettes of the invention of the invention may be used to modify a target nucleic acid. A target nucleic acid may be contacted with a polynucleotide/nucleic acid construct/expression cassette of the invention prior to, concurrently with or after contacting the target nucleic acid with the guide nucleic acid. In some embodiments, the polynucleotides of the invention and a guide nucleic acid may be comprised in the same expression cassette or vector and therefore, a target nucleic acid may be contacted concurrently with the polynucleotides of the invention and guide nucleic acid. In some embodiments, the polynucleotides of the invention and a guide nucleic acid may be in different expression cassettes or vectors and thus, a target nucleic acid may be contacted with the polynucleotides of the invention prior to, concurrently with, or after contact with a guide nucleic acid.

A target nucleic acid of any organism may be modified (e.g., mutated, e.g., base edited, cleaved, nicked, etc.) using the polynucleotides and methods of the invention, including, but not limited to, eukaryotic organisms or prokaryotic organisms, such as for example, a plant, an animal, a bacterium, an archaeon, a fungus and/or a virus. Any animal or cell thereof may be modified (e.g., mutated, e.g., base edited, cleaved, nicked, etc.) using the polynucleotides of the invention including, but not limited to an insect, a fish, a bird, an amphibian, a reptile, and/or a mammal. Exemplary mammals for which this invention may be useful include, but are not limited to, primates (human and non-human (e.g., a chimpanzee, baboon, monkey, gorilla, etc.)), cats, dogs, ferrets, gerbils, hamsters, cows, pigs, horses, goats, donkeys, or sheep. In some embodiments, a fungal target organism can include, but is not limited to, a Zygomycota, Ascomycota, Basidiomycota, and Deuteromycota (fungi imperfecti), optionally wherein the fungal target organism may be an ascomycete, optionally a yeast. In some embodiments, a fungal target organism may be from the genera *Saccharomyces*, optionally *Saccharomyces cerevisiae*.

A target nucleic acid of any plant or plant part may be modified (e.g., mutated, e.g., base edited, cleaved, nicked, etc.) using the polynucleotides of the invention. Any plant (or groupings of plants, for example, into a genus or higher order classification) may be modified using the nucleic acid constructs of this invention including an angiosperm, a gymnosperm, a monocot, a dicot, a C3, C4, CAM plant, a bryophyte, a fern and/or fern ally, a microalgae, and/or a macroalgae. A plant and/or plant part useful with this invention may be a plant and/or plant part of any plant species/variety/cultivar. The term "plant part," as used herein, includes but is not limited to, embryos, pollen, ovules, seeds, leaves, stems, shoots, flowers,

branches, fruit, kernels, ears, cobs, husks, stalks, roots, root tips, anthers, plant cells including plant cells that are intact in plants and/or parts of plants, plant protoplasts, plant tissues, plant cell tissue cultures, plant calli, plant clumps, and the like. As used herein, “shoot” refers to the above ground parts including the leaves and stems. Further, as used herein, “plant cell”
 5 refers to a structural and physiological unit of the plant, which comprises a cell wall and also may refer to a protoplast. A plant cell can be in the form of an isolated single cell or can be a cultured cell or can be a part of a higher-organized unit such as, for example, a plant tissue or a plant organ.

Non-limiting examples of plants useful with the present invention include turf grasses
 10 (e.g., bluegrass, bentgrass, ryegrass, fescue), feather reed grass, tufted hair grass, miscanthus, arundo, switchgrass, vegetable crops, including artichokes, kohlrabi, arugula, leeks, asparagus, lettuce (e.g., head, leaf, romaine), malanga, melons (e.g., muskmelon, watermelon, crenshaw, honeydew, cantaloupe), cole crops (e.g., brussels sprouts, cabbage, cauliflower, broccoli, collards, kale, Chinese cabbage, bok choy), cardoni, carrots, napa, okra, onions,
 15 celery, parsley, chick peas, parsnips, chicory, peppers, potatoes, cucurbits (e.g., marrow, cucumber, zucchini, squash, pumpkin, honeydew melon, watermelon, cantaloupe), radishes, dry bulb onions, rutabaga, eggplant, salsify, escarole, shallots, endive, garlic, spinach, green onions, squash, greens, beet (sugar beet and fodder beet), sweet potatoes, chard, horseradish, tomatoes, turnips, and spices; a fruit crop such as apples, apricots, cherries, nectarines,
 20 peaches, pears, plums, prunes, cherry, quince, fig, nuts (e.g., chestnuts, pecans, pistachios, hazelnuts, pistachios, peanuts, walnuts, macadamia nuts, almonds, and the like), citrus (e.g., clementine, kumquat, orange, grapefruit, tangerine, mandarin, lemon, lime, and the like), blueberries, black raspberries, boysenberries, cranberries, currants, gooseberries, loganberries, raspberries, strawberries, blackberries, grapes (wine and table), avocados,
 25 bananas, kiwi, persimmons, pomegranate, pineapple, tropical fruits, pomes, melon, mango, papaya, and lychee, a field crop plant such as clover, alfalfa, timothy, evening primrose, meadow foam, corn/maize (field, sweet, popcorn), hops, jojoba, buckwheat, safflower, quinoa, wheat, rice, barley, rye, millet, sorghum, oats, triticale, sorghum, tobacco, kapok, a leguminous plant (beans (e.g., green and dried), lentils, peas, soybeans), an oil plant (rape,
 30 canola, mustard, poppy, olive, sunflower, coconut, castor oil plant, cocoa bean, groundnut, oil palm), duckweed, *Arabidopsis*, a fiber plant (cotton, flax, hemp, jute), *Cannabis* (e.g., *Cannabis sativa*, *Cannabis indica*, and *Cannabis ruderalis*), lauraceae (cinnamon, camphor), or a plant such as coffee, sugar cane, tea, and natural rubber plants; and/or a bedding plant such as a flowering plant, a cactus, a succulent and/or an ornamental plant (e.g., roses, tulips,

violets), as well as trees such as forest trees (broad-leaved trees and evergreens, such as conifers; e.g., elm, ash, oak, maple, fir, spruce, cedar, pine, birch, cypress, eucalyptus, willow), as well as shrubs and other nursery stock. In some embodiments, the nucleic acid constructs of the invention and/or expression cassettes and/or vectors encoding the same may be used to modify maize, soybean, wheat, canola, rice, tomato, pepper, sunflower, raspberry, blackberry, black raspberry and/or cherry.

The present invention further comprises a kit or kits to carry out the methods of this invention. A kit of this invention can comprise reagents, buffers, and apparatus for mixing, measuring, sorting, labeling, etc., as well as instructions and the like as would be appropriate for modifying a target nucleic acid.

In some embodiments, the invention provides a kit comprising one or more nucleic acid constructs of the invention and/or expression cassettes and/or vectors comprising the same, with optional instructions for the use thereof. In some embodiments, a kit may further comprise a CRISPR-Cas guide nucleic acid (or extended guide nucleic acid) (corresponding to the CRISPR-Cas effector protein encoded by the polynucleotide of the invention) and/or expression cassette and/or vector comprising the same. In some embodiments, the guide nucleic acid/extended guide nucleic acid may be provided on the same expression cassette and/or vector as one or more polynucleotides of the invention. In some embodiments, a guide nucleic acid/extended guide nucleic acid may be provided on a separate expression cassette or vector from that comprising one or more of the polynucleotides of the invention.

In some embodiments, the kit may further comprise a nucleic acid construct encoding a guide nucleic acid, wherein the construct comprises a cloning site for cloning of a nucleic acid sequence identical or complementary to a target nucleic acid sequence into backbone of the guide nucleic acid.

In some embodiments, a nucleic acid construct of the invention may be an mRNA that may encode one or more introns within the encoded polynucleotide. In some embodiments, an expression cassette and/or vector comprising one or more polynucleotides of the invention, may further encode one or more selectable markers useful for identifying transformants (e.g., a nucleic acid encoding an antibiotic resistance gene, herbicide resistance gene, and the like).

The invention will now be described with reference to the following examples. It should be appreciated that these examples are not intended to limit the scope of the claims to the invention but are rather intended to be exemplary of certain embodiments. Any variations

in the exemplified methods that occur to the skilled artisan are intended to fall within the scope of the invention.

EXAMPLES

RNA-encoded DNA-replacement of alleles (REDRAW) utilizes a type V Cas effector, an enzyme which polymerizes from a DNA:RNA hybrid from a free DNA 3' end (annealing site, AS), and an extended guide nucleic acid (i.e., a targeted allele guide RNA (tagRNA)). These three macromolecules work in tandem to i) locate the CRISPR enzyme to the genomic site of interest using a CRISPR effector and the crRNA portion of the tagRNA, ii) nick or cut the DNA to produce a free 3' end, iii) provide a portion of the tagRNA which anneals to the free 3' end of the DNA, iv) provide a portion of tagRNA which provides a template for the RNA-dependent DNA polymerase, and v) allow the termination of reverse transcription either by enzyme collision, natural termination, or encountering a stable hairpin.

We tested the REDRAW system using a nontarget-stand (NTS) nickase version of LbCas12a_R1138A and a RT from Moloney Murine Leukemia Virus (M-MuLV). LbCas12a_R1138A was expected to be an NTS nickase based on alignment with an the previously described AsCas12a_R1226A mutation. We demonstrate in Figure XXX that LbCas12a_R1138A is, indeed, a nickase. The LbCas12a used was either RNase (+) or had a mutation which prevented RNase activity (H759A). The LbCas12a_R1138A_H759A mutant was used to prevent self-processing of the tagRNA when making the 5' extension or when incorporating a 3' hairpin (e.g., a pseudoknot comprising a hairpin element).

The tagRNAs tested contained crRNAs containing either 5' or 3' extensions. Various annealing site lengths were tested allowing for shorter or longer DNA:RNA hybrids to form from at the nicked non-target strand. Various lengths of RNA template were tested as well. Finally, two different hairpins were also incorporated into a LbCas12a crRNA sequence, a pseudoknotted hairpin design and a decoy pseudoknotted hairpin design.

Example 1

LbCas12a_R1138A nickase assay

A nucleic acid construct was synthesized comprising LbCas12a, followed by a nucleoplasmin NLS, and a 6x histidine tag (GeneWiz) (SEQ ID NO:57) and cloned into a pET28a vector between NcoI and XhoI, generating pWISE450 (SEQ ID NO:58). There was an additional glycine added to the sequence between Met-1 and Ser-2 to facilitate cloning. Numbering presented herein excludes this extra glycine. Then the R1138A mutation was

made using a QuickChange II site-directed mutagenesis kit (Agilent) according to manufacturer's instructions. These expression plasmids were then transformed into BL21 (DE3) Star competent *E. coli* cells (ThermoFisher Scientific).

The BL21(DE3) Star cells were grown in Luria Broth and 50 ug/ml of kanamycin at 37°C until an optical density of A600=0.5 was achieved. Isopropyl β -d-1-thiogalactopyranoside (IPTG) was added to 0.5 mM and protein was induced overnight at 18°C. Cells were pelleted at 5,000 x g. Purification was accomplished using two columns: a HisTrap column followed by a MonoS column (GE Healthcare) according to manufacturer's protocols.

CRISPR RNA (crRNA) was synthesized by Synthego with the sequence AAUUUCUACUAAGUGUAGAUGGAAUCCCUUCUGCAGCACCUGG (**SEQ ID NO:59**) (where the guide portion is in bold font).

The plasmid to be cleaved was pUC19 with the following sequence inserted: **TTTCGGAATCCCTTCTGCAGCACCTGG** (**SEQ ID NO:60**) where the portion of the sequence in bold font is a PAM sequence recognized by LbCas12a and the remainder (regular font) is the protospacer sequence. The pUC19 plasmid was transformed into XL1-Blue (Agilent) (*E. coli*), and subsequently purified using Qiagen plasmid spin minikits.

The nuclease assay was accomplished by mixing 10:10:1 ratios of LbCas12a_R1138:crRNA:plasmid, incubated for 15 minutes at 37°C in New England Biolabs buffer 2.1, heat inactivated for 20 minutes at 80°C, and loaded onto a 1% TAE-agarose gel with SYBR-Safe stain (Invitrogen) embedded to stain the DNA. As shown in **Fig. 4** in an *in vitro* assay, LbCas12a_R1138A is a nickase. As shown in lanes 2 and 3, a supercoiled 2.8 kB plasmid ran with an apparent size of 2.0 kB (lane 2) until a double-stranded break was generated by wildtype LbCas12a (lane 3). The mutant enzyme LbCas12a_R1138A predominantly generated a nicked product running with the apparent size of 5.0 kB. Lanes 4-6 show that increasing concentrations of the mutant enzyme did not alter the ratio until extremely high concentrations of enzyme were used resulting in general nuclease digestion of the plasmid (256 nM).

REDRAW Editor Plasmid Design and Construction- bacterial screen

REDRAW (RNA-encoded DNA-replacement of alleles) expression constructs were synthesized by solid state synthesis and cloned into expression vector pET28a(+) in between the NcoI and XhoI restriction sites. The REDRAW expression vectors contain a ColE1 origin of replication, a kanamycin resistance marker, and a REDRAW editor under control of a T7

promoter and terminator. The REDRAW editors contain either a Cas12a nickase (R1138A) or an Rnase dead Cas12a nickase (R1138A, H759A) fused to Mu-LV reverse transcriptase MuLV(5M) (see, e.g., **SEQ ID NO:97**) (Murine leukemia virus reverse transcriptase with five mutations - D200N+L603W+T330P+T306K+W313F) (Anzalone et al. *Nature* 576 (7785):149-157 (2019)) with an XTEN or 5R linker. All REDRAW editor sequences were *E. coli* codon optimized. The REDRAW editor configurations tested are shown in **Fig. 5**. Two configurations provided in **Fig. 5** had Cas12a N-terminal to the reverse transcriptase, and two configurations had Cas12a C-terminal to the reverse transcriptase. The tested configurations were built with a Cas12a variant that had an additional H759A mutation to prevent processing of tagRNAs that contain a 5' extension.

tagRNA Plasmid Design and Construction-bacterial screen

The sequences of the tagRNA (targeted allele guide RNA) library were designed using an algorithm that assembled a Cas12a spacer and scaffold sequence together with a reverse transcriptase template and primer binding site unique for each target. The design parameters, shown in **Table 1**, span a wide range of primer binding site and reverse transcriptase template lengths. The desired changes, shown in **Table 3**, were designed to confer resistance to antibiotics following successful editing.

Table 1. Conformations of tagRNAs tested in the first library

Type	PBS	RTT	Targets in Library
5' extension	10-20 nt, 1 nt steps	10-150 nt, 5 nt steps	2 genomic, 3 plasmid
3' extension	10-20 nt, 1 nt steps	10-150 nt, 5 nt steps	2 genomic, 3 plasmid

- 10 **Fig. 6** shows the configurations of the tagRNAs in the first library. Both 5' and 3' extensions containing the RTT and PBS were included in the library.

A second library was designed in a similar fashion as the first, while additionally evaluating whether the presence of a hairpin, located just 3' of the spacer in the 3' tagRNA extension configuration, would improve REDRAW editing. The design parameters, shown in **Table 2**, again interrogate a wide range of primer binding site (PBS) and reverse transcriptase template (RTT) lengths, but also focus on the region of RTT length found to be functional from the first library. Both 5' and 3' extensions containing the RTT and PBS were included in the library. Additionally, variants containing a decoy hairpin were also included in the second tagRNA library. As a hairpin was desired that would be similar to the natural LbCas12a scaffold sequence but would not be recognized and cleaved by the Cas12a protein, an existing hairpin with similar architecture to the LbCas12a hairpin was found in the HIV-1 RNA genome and modified by the addition of a UA sequence to form a pseudoknot, as shown in **Fig. 7**.

Table 2. Conformations of tagRNAs tested in the second library

TagRNA Extension	Range of PBS	Range of RTT	Decoy Hairpin	Targets
5' end	10-20 nt, 1-nt steps	10-190 nt, 5-nt steps	None	2 genomic, 3 plasmid
3' end	10-20 nt, 1-nt steps	10-190 nt, 5-nt steps; 65-85 nt, 2-nt steps	With and without	2 genomic, 3 plasmid

tagRNA Plasmid Construction for Bacterial Screen

The base plasmid for the tagRNA library was generated by solid state synthesis and cloning of a holder fragment into pTwist Amp Medium Copy (TWIST BIOSCIENCE®). The plasmid contains a p15A origin of replication and an ampicillin resistance marker. The tagRNAs are constitutively expressed from a synthetic BbaJ23119 promoter and are terminated by a T7 terminator. The first tagRNA library evaluated was synthesized and cloned into the tagRNA base vector by an external vendor (Genewiz). For the second library, oligos were synthesized and then cloned into the tagRNA base vector using a NEB HiFi Assembly kit according to manufacturer's instructions. Library diversity was investigated by colony PCR and Sanger sequencing of 72 clones from the library, to ensure that a wide range of PBS, RTT, and targets were included in the library and that there was not a substantial bias.

Reporter Plasmid Design and Construction

A base reporter plasmid containing a CloDF13 origin of replication, chloramphenicol resistance marker, and spectinomycin resistance marker (*aadA*) was constructed by PCR amplification of the CloDF13 origin of replication and chloramphenicol resistance marker and ligating it with a PCR-amplified *aadA* resistance marker. Three reporter plasmids containing variants of *aadA* were then constructed by cutting out the wild type *aadA* gene in between the BamHI and BglII restriction sites and ligating in gene blocks synthesized that contained a stop codon at residue position Thr61, Leu115, or Asp132. All reporter plasmids were verified by Sanger sequencing after construction. In addition, reporter plasmids containing an *aadA* variant with a stop codon in the coding sequence were verified as both spectinomycin and streptomycin sensitive prior to using them in REDRAW tagRNA screening experiments.

Targets for REDRAW Editing- bacterial screen

Five targets were tested in the REDRAW editing experiments, shown below in **Table 3**. Two genomic and three plasmid targets were used in all cases. Successful REDRAW editing at any of the targets results in resistance to an antibiotic (nalidixic acid or streptomycin), tying survival of the host organism (*E. coli*) to the success of REDRAW editing.

Table 3. Targets for bacterial REDRAW editing

Target	Location of Target	Desired Edit	Successful Editing Result
gyrA	Genome	Ser83>Leu TCG>TTG	Resistance to Nalidixic Acid
rpsL	Genome	Lys44>Arg AAA>CGT	Resistance to Streptomycin
aadA	Plasmid	Stop61>Thr TGA>ACG	Resistance to Streptomycin
aadA	Plasmid	Stop115>Leu TGA>CTG	Resistance to Streptomycin
aadA	Plasmid	Stop132>Asp TGA>GAT	Resistance to Streptomycin

REDRAW tagRNA Experiments - bacterial screen

The host organism for all bacterial REDRAW tagRNA screening experiments was *E. coli* BL21(DE3). Prior to performing the selection experiments, each REDRAW expression construct was transformed into chemically competent BL21(DE3) according to manufacturer's instructions and plated onto LB agar plates with Kanamycin. Single colonies were then picked from the transformation plates, and batches of electrocompetent cells were made following a previously developed method (Sambrook and Russell (*Transformation of E. coli by electroporation*. Cold Spring Harbor Protocols 2006.1 (2006): pdb-prot3933). Competent cells harboring each REDRAW expression construct were then electroporated with 10 ng of each reporter plasmid, recovered for 1 hour in SOC at 37C, 225 rpm, and plated onto LB agar plates with kanamycin and chloramphenicol. Single colonies from these plates were then picked from the transformation plates, and batches of electrocompetent cells were made again (Sambrook and Russell (*Transformation of E. coli by electroporation*. Cold Spring Harbor Protocols 2006.1 (2006): pdb-prot3933). **Table 4** below summarizes the batches of electrocompetent cells made for the first tagRNA library testing.

Table 1 - Electrocompetent Cells prepared for tagRNA Library 1 Selection Experiments

Competent Cell Batch	Constructs Harbored in BL21(DE3)	SEQ ID NO
1	SV40-MMLV-RT-XTEN-nRRLbCas12a-SV40	63
2	SV40-MMLV-RT-5R-nRRLbCas12a-SV40	64
3	SV40-nRRLbCas12a-XTEN-MMLV-RT-SV40	65
4	SV40-nRRLbCas12a-5R-MMLV-RT-SV40	66
5	SV40-MMLV-RT-XTEN-nRVRLbCas12a-SV40	67
6	SV40-MMLV-RT-5R-nRVRLbCas12a-SV40	68

7	SV40-nRVRLbCas12a-XTEN-MMLV-RT-SV40	69
8	SV40-nRVRLbCas12a-5R-MMLV-RT-SV40	70
9	SV40-MMLV-RT-XTEN-nLbCas12a-SV40 + aadA Thr61	71 + Thr61
10	SV40-MMLV-RT-XTEN-nLbCas12a-SV40 + aadA Leu115	71 + Leu115
11	SV40-MMLV-RT-XTEN-nLbCas12a-SV40 + Asp132	71 + Asp132
12	SV40-MMLV-RT-5R-nLbCas12a-SV40 + Thr61	72 + Thr61
13	SV40-MMLV-RT-5R-nLbCas12a-SV40 + Leu115	72 + Leu115
14	SV40-MMLV-RT-5R-nLbCas12a-SV40 + Asp132	72 + Asp132
15	SV40-nLbCas12a-XTEN-MMLV-RT-SV40 + Thr61	73 + Thr61
16	SV40-nLbCas12a-XTEN-MMLV-RT-SV40 + Leu115	73 + Leu115
17	SV40-nLbCas12a-XTEN-MMLV-RT-SV40 + Asp132	73 + Asp132
18	SV40-nLbCas12a-5R-MMLV-RT-SV40+ Thr61	74 + Thr61
19	SV40-nLbCas12a-5R-MMLV-RT-SV40+ Leu115	74 + Leu115
20	SV40-nLbCas12a-5R-MMLV-RT-SV40+ Asp132	74 + Asp132

SV40 = NLS, MMLV-RT = reverse transcriptase, XTEN = linker, nLbCas12a = nickase Cas12

Selection experiments were performed by first electroporating 100 ng of tagRNA library into 50 uL of each batch of electrocompetent cells. Transformations were recovered for 1 hour at 37°C with 225 rpm shaking. After 1 hour of recovery, 1 uL of recovery was removed, mixed with 99 uL of LB, and plated onto LB agar plates with appropriate antibiotics to check for transformation efficiency. The remaining amount of each transformation was then added to 29 mL of LB + Antibiotics (LB Kan/Carb for genomic selections, and LB Kan/Carb/Cam for plasmid selections) and 0.5 mM IPTG. The expression cultures were grown at 37°C, with 225 rpm shaking overnight.

The following day, the OD₆₀₀ of each expression culture was measured. For each expression culture, 1 OD was plated onto 5 plates (about 0.2 OD per plate) containing antibiotics for the REDRAW expression vector (Kan), the tagRNA plasmid (Carb), the reporter plasmid, 0.5 mM IPTG, and an additional selection antibiotic (nalidixic acid or streptomycin). Plates were incubated overnight at 37°C, and growth was observed the following morning. If no colonies were observed, the plates were incubated an additional 24 hours at 37°C.

Colonies that were observed on the selection plates were picked, re-streaked onto plates with appropriate antibiotics, and then subjected to colony PCR to amplify the gene

targeting for editing and the tagRNA for Sanger sequencing. Sanger sequencing was performed on the colony PCR products by Genewiz.

- 5 Evaluation of the second library was performed the same way as the first tagRNA library, with one modification. Instead of preparing 20 batches of electrocompetent cells, one large batch of electrocompetent BL21(DE3) harboring the second tagRNA library was prepared. The REDRAW expression constructs (100 ng) or the REDRAW expression constructs + reporter plasmids (100 ng each) were then transformed into electrocompetent cells harboring the tagRNA library. All subsequent steps were repeated in the same manner.

10 ***Evaluation of REDRAW Editing with the first tagRNA Library-bacterial screen***

The number of colonies obtained from the selection experiments for the first tagRNA library are summarized in **Table 5** below. No colonies were observed for either of the genomic selections (selections 1-8). For each of the plasmid selections, colonies were observed.

15 **Table 5.** First tagRNA library selection experiment results.

Selection Number	REDRAW Editor	Target	Colonies on Selection Plates
1	SV40-MMLV-RT-XTEN-nRRLbCas12a-SV40 (SEQ ID NO:63)	gyrA (genome)	0
2	SV40-MMLV-RT-5R-nRRLbCas12a-SV40 (SEQ ID NO:64)	gyrA (genome)	0
3	SV40-nRRLbCas12a-XTEN-MMLV-RT-SV40 (SEQ ID NO:65)	gyrA (genome)	0
4	SV40-nRRLbCas12a-5R-MMLV-RT-SV40 (SEQ ID NO:66)	gyrA (genome)	0
5	SV40-MMLV-RT-XTEN-nRVRLbCas12a-SV40 (SEQ ID NO:67)	rpsL (genome)	0
6	SV40-MMLV-RT-5R-nRVRLbCas12a-SV40 (SEQ ID NO:68)	rpsL (genome)	0
7	SV40-nRVRLbCas12a-XTEN-MMLV-RT-SV40 (SEQ ID NO:69)	rpsL (genome)	0
8	SV40-nRVRLbCas12a-5R-MMLV-RT-SV40 (SEQ ID NO:70)	rpsL (genome)	0
9	SV40-MMLV-RT-XTEN-nLbCas12a-SV40 (SEQ ID NO:71)	aadA Thr61 (plasmid)	Lawn
10	SV40-MMLV-RT-XTEN-nLbCas12a-SV40 (SEQ ID NO:71)	aadA Leu115 (plasmid)	11
11	SV40-MMLV-RT-XTEN-nLbCas12a-SV40 (SEQ ID NO:71)	aadA Asp132 (plasmid)	9

12	SV40-MMLV-RT-5R-nLbCas12a-SV40 (SEQ ID NO:72)	aadA Thr61 (plasmid)	Lawn
13	SV40-MMLV-RT-5R-nLbCas12a-SV40 (SEQ ID NO:72)	aadA Leu115 (plasmid)	10
14	SV40-MMLV-RT-5R-nLbCas12a-SV40 (SEQ ID NO:72)	aadA Asp132 (plasmid)	9
15	SV40-nLbCas12a-XTEN-MMLV-RT-SV40 (SEQ ID NO:73)	aadA Thr61 (plasmid)	Lawn
16	SV40-nLbCas12a-XTEN-MMLV-RT-SV40 (SEQ ID NO:73)	aadA Leu115 (plasmid)	1
17	SV40-nLbCas12a-XTEN-MMLV-RT-SV40 (SEQ ID NO:73)	aadA Asp132 (plasmid)	1
18	SV40-nLbCas12a-5R-MMLV-RT-SV40 (SEQ ID NO:74)	aadA Thr61 (plasmid)	Lawn
19	SV40-nLbCas12a-5R-MMLV-RT-SV40 (SEQ ID NO:74)	aadA Leu115 (plasmid)	2
20	SV40-nLbCas12a-5R-MMLV-RT-SV40 (SEQ ID NO:74)	aadA Asp132 (plasmid)	0

For selections 9, 12, 15 and 18 (*aadA* Thr61 target), lawns of bacteria were observed. Isolated colonies from these plates were false positives. For selections 10, 11, 13, 14, 16, and 17 (*aadA* Leu115 target and *aadA* Asp132 target), low numbers of colonies were observed on the plates. Colonies on these plates had both the tagRNA and the target amplified by colony PCR and were sent for Sanger sequencing to confirm the edit made and to identify the tagRNA responsible for the edit. All colonies evaluated from selections 11, 14, 17 and 20 (*aadA* Asp132 target) were false positives. Multiple colonies from selection 10 (*aadA* Leu115 target) had the designed edit and an associated tagRNA. The sequencing result of the edited target is shown in **Fig. 8**, demonstrating a TGA → CTG edit in a defunct *aadA* gene, restoring antibiotic resistance.

The identified sequence of the tagRNA responsible for the edit is associated with the edit shown in **Fig. 8**:

5' – GTTTCAAAGATTAAATAATTTCTACTAAGTGTAGATTACGGCTCCGCAGTGGATGGCGGTAA
TTTCTACTAAGTGTAGATGCGGCGCGTTGTTTCATCAAGGCGTACGGTCACCGTAACCAGCAAAT
CAATATCACTGTGTGGCTTCAGGCCGCCATCCACTGCGG – 3' (SEQ ID NO:87).

The protein configuration from selection 10 is the following: SV40-nCas12a-XTEN-MMLV-RT-SV40.

Evaluation of REDRAW Editing with the Second tagRNA Library- genomic selection results

The number of colonies obtained from the genomic selection experiments for the second tagRNA library are summarized in **Table 6** below. Colonies were observed on the *rpsL* selection plates.

5

Table 6. Second tagRNA library experimental results - colonies on selection plates for the genomic selections

Selection Number	REDRAW Editor	Target	Colonies on Selection Plates
2.1	SV40-MMLV-RT-XTEN-nRRLbCas12a(H759A)-SV40 (SEQ ID NO:75)	gyrA (genome)	0
2.2	SV40-MMLV-RT-5R-nRRLbCas12a(H759A)-SV40 (SEQ ID NO:76)	gyrA (genome)	0
2.3	SV40-nRRLbCas12a(H759A)-XTEN-MMLV-RT-SV40 (SEQ ID NO:77)	gyrA (genome)	0
2.4	SV40-nRRLbCas12a(H759A)-5R-MMLV-RT-SV40 (SEQ ID NO:78)	gyrA (genome)	0
2.5	SV40-MMLV-RT-XTEN-nRVRLbCas12a(H759A)-SV40 (SEQ ID NO:79)	rpsL (genome)	5
2.6	SV40-MMLV-RT-5R-nRVRLbCas12a(H759A)-SV40 (SEQ ID NO:80)	rpsL (genome)	8
2.7	SV40-nRVRLbCas12a(H759A)-XTEN-MMLV-RT-SV40 (SEQ ID NO:81)	rpsL (genome)	2
2.8	SV40-nRVRLbCas12a(H759A)-5R-MMLV-RT-SV40 (SEQ ID NO:82)	rpsL (genome)	11
2.9	SV40-MMLV-RT-XTEN-nRRLbCas12a-SV40 (SEQ ID NO:63)	gyrA (genome)	0
2.10	SV40-MMLV-RT-5R-nRRLbCas12a-SV40 (SEQ ID NO:64)	gyrA (genome)	0
2.11	SV40-nRRLbCas12a-XTEN-MMLV-RT-SV40 (SEQ ID NO:65)	gyrA (genome)	0
2.12	SV40-nRRLbCas12a-5R-MMLV-RT-SV40 (SEQ ID NO:66)	gyrA (genome)	0
2.13	SV40-MMLV-RT-XTEN-nRVRLbCas12a-SV40 (SEQ ID NO:67)	rpsL (genome)	3
2.14	SV40-MMLV-RT-5R-nRVRLbCas12a-SV40 (SEQ ID NO:68)	rpsL (genome)	0
2.15	SV40-nRVRLbCas12a-XTEN-MMLV-RT-SV40 (SEQ ID NO:69)	rpsL (genome)	0
2.16	SV40-nRVRLbCas12a-5R-MMLV-RT-SV40 (SEQ ID NO:70)	rpsL (genome)	1

For selections 2.1-2.4 and 2.9-2.12 (*gyrA* genomic target), no colonies were observed on the plates. For selections 2.5-2.8 and 2.13-2.16 (*rpsL* genomic target), low numbers of colonies were observed on these plates. Colonies on these plates were re-streaked to verify resistance to all antibiotics. Colonies from these plates were then used to generate PCR products of the tagRNA and the target for Sanger sequencing. Sanger sequencing was used to confirm the edit made and to identify the tagRNA responsible for the edit. All colonies from selections 2.6-2.8 and 2.13-2.16 were false positives. One colony from selection 2.5 had the designed edit AAA to CGT, which confers Streptomycin resistance (see **Fig. 9**).

The identified sequence of the tagRNA associated with the edit shown in **Fig. 9** is:

5' – TATTTCTATAAGTGTAGATTACTCGTGTATATATACTCCGCACCGAGGTTGGTACGAACAC
CGGGAGTCTTTAACACGACCGCCACGGATCAGGATCACGGAGTGCTCCTGCAGGTTGTGACCTT
CACCACCGATGTAGGAAGTCACTTCGAAACCGTTAGTCAGACGAACACGGCATACTTTACGCAG
CGCGGAGTTCGGTTTACGAGGAGTGGTAGTATATACACGAGT– 3' **SEQ ID NO:92**.

The protein configuration from selection 2.5 is the following: SV40-MMLV-RT-XTEN-nRVRLbCas12a(H759A)-SV40.

Evaluation of REDRAW Editing with the Second tagRNA Library – Plasmid Selection Results

The number of colonies obtained from the plasmid selection experiments for the second tagRNA library are summarized in **Table 7** below.

Table 7.

Selection Number	REDRAW Editor	Target	Colonies on Selection Plates
2.17	SV40-MMLV-RT-XTEN-nLbCas12a-SV40 (SEQ ID NO:71)	aadA Thr61 (plasmid)	0
2.18	SV40-MMLV-RT-XTEN-nLbCas12a-SV40 (SEQ ID NO:71)	aadA Leu115 (plasmid)	4
2.19	SV40-MMLV-RT-XTEN-nLbCas12a-SV40 (SEQ ID NO:71)	aadA Asp132 (plasmid)	2
2.20	SV40-MMLV-RT-5R-nLbCas12a-SV40 (SEQ ID NO:72)	aadA Thr61 (plasmid)	0
2.21	SV40-MMLV-RT-5R-nLbCas12a-SV40 (SEQ ID NO:72)	aadA Leu115 (plasmid)	0
2.22	SV40-MMLV-RT-5R-nLbCas12a-SV40 (SEQ ID NO:72)	aadA Asp132 (plasmid)	1
2.23	SV40-nLbCas12a-XTEN-MMLV-RT-SV40 (SEQ ID NO:73)	aadA Thr61 (plasmid)	0

2.24	SV40-nLbCas12a-XTEN-MMLV-RT-SV40 (SEQ ID NO:73)	aadA Leu115 (plasmid)	0
2.25	SV40-nLbCas12a-XTEN-MMLV-RT-SV40 (SEQ ID NO:73)	aadA Asp132 (plasmid)	9
2.26	SV40-nLbCas12a-5R-MMLV-RT-SV40 (SEQ ID NO:74)	aadA Thr61 (plasmid)	0
2.27	SV40-nLbCas12a-5R-MMLV-RT-SV40 (SEQ ID NO:74)	aadA Leu115 (plasmid)	0
2.28	SV40-nLbCas12a-5R-MMLV-RT-SV40 (SEQ ID NO:74)	aadA Asp132 (plasmid)	2
2.29	SV40-MMLV-RT-XTEN-nLbCas12a(H759A)- SV40 (SEQ ID NO:83)	aadA Thr61 (plasmid)	0
2.30	SV40-MMLV-RT-XTEN-nLbCas12a(H759A)- SV40 (SEQ ID NO:83)	aadA Leu115 (plasmid)	0
2.31	SV40-MMLV-RT-XTEN-nLbCas12a(H759A)- SV40 (SEQ ID NO:83)	aadA Asp132 (plasmid)	12
2.32	SV40-MMLV-RT-5R-nLbCas12a(H759A)-SV40 (SEQ ID NO:84)	aadA Thr61 (plasmid)	0
2.33	SV40-MMLV-RT-5R-nLbCas12a(H759A)-SV40 (SEQ ID NO:84)	aadA Leu115 (plasmid)	0
2.34	SV40-MMLV-RT-5R-nLbCas12a(H759A)-SV40 (SEQ ID NO:84)	aadA Asp132 (plasmid)	0
2.35	SV40-nLbCas12a(H759A)-XTEN-MMLV-RT- SV40 (SEQ ID NO:85)	aadA Thr61 (plasmid)	0
2.36	SV40-nLbCas12a(H759A)-XTEN-MMLV-RT- SV40 (SEQ ID NO:85)	aadA Leu115 (plasmid)	0
2.37	SV40-nLbCas12a(H759A)-XTEN-MMLV-RT- SV40 (SEQ ID NO:85)	aadA Asp132 (plasmid)	0
2.38	SV40-nLbCas12a(H759A)-5R-MMLV-RT-SV40 (SEQ ID NO:85)	aadA Thr61 (plasmid)	0
2.39	SV40-nLbCas12a(H759A)-5R-MMLV-RT-SV40 (SEQ ID NO:86)	aadA Leu115 (plasmid)	1
2.40	SV40-nLbCas12a(H759A)-5R-MMLV-RT-SV40 (SEQ ID NO:86)	aadA Asp132 (plasmid)	2

Colonies were observed on plates for the Leu115 and Asp132 selections. Selections 2.18, 2.19, 2.22, 2.25, 2.28, 2.31, 2.39, and 2.40 had colonies on the selection plates. These colonies were re-streaked to verify resistance to all antibiotics. They were then used to

5 generate PCR products of the tagRNA and the target for Sanger sequencing. Sanger

sequencing was used to confirm the edit made and to identify the tagRNA responsible for the edit. All colonies from selections 2.18, 2.19, 2.22, 2.28, 2.39, and 2.40 were false positives. Four colonies from selection 2.25 and two colonies from selection 2.31 had the designed edit and an associated tagRNA as shown in **Fig. 10** and **Fig. 11**. The four colonies from selection 2.25 had identical edits and tagRNAs. The two colonies from selection 2.31 also had identical edits and tagRNAs.

The identified sequence of the tagRNA associated with the edit in **Fig. 10** from selection 2.25 is:

5' – TAATTTCTACTAAGTGTAGATTACGGCTCCGCAGTGGATGGCGGTAAGTCTCCATAGAATG
GAGGACAGCGCGGAGAATCTCGCTCTCTCCAGGGGAAGCCGAAGTTTCCAAAAGGTCGTTGATC
AAAGCGCGGCGCGTTGTTTCATCAAGGCGTACGGTCACCGTAACCAGCAAATCAATATCACTGT
GTGGCTTCAGGCCGCCATCCACTGCGGAT– 3' **SEQ ID NO:93**.

The protein configuration from selection 2.25 is the following: SV40-nCas12a-XTEN-MMLV-RT-SV40.

The identified sequence of the tagRNA associated with the edit in **Fig. 11** from selection 2.31 is:

5' – TAATTTCAACTAAGTGTAGATTACGGCTCCGCAGTGGATGGCGGTAAGTCTCCATAGAATGG
AGGGCGGAGAATCTCGCTCTCTCCAGGGGAAGCCGAAGTTTCCAAAAGGTCGTTGATCAAAGCG
CGGCGCGTTGTTTCATCAAGGCGTACGGTCACCGTAACCAGCAAATCAATATCACTGTGTGGCTT
CAGGCCGCCATCCACTGCGGAT – 3' **SEQ ID NO:94**.

The protein configuration from selection 2.31 is the following: SV40-MMLV-RT-XTEN-nLbCas12a(H759)-SV40.

Summary of observed REDRAW Editing in Bacterial Cells

Table 8 below provides a summary of the observed instances of REDRAW editing in *E. coli*. Described for each example is the protein configuration (REDRAW Editor), the target that was edited, the location of the tagRNA extension (5' or 3' of the Cas12a hairpin and guide), the PBS length, and the RTT length.

Table 8. Summary of REDRAW editing observed in *E. coli*.

Selection	REDRAW Editor	Target	Extension	PBS length (bp)	RTT length
10	SV40-MMLV-RT-XTEN-nLbCas12a-SV40 (SEQ ID NO:71)	aadA Leu115 (plasmid)	3'	17	96 bp
2.5	SV40-MMLV-RT-XTEN-nRVRLbCas12a(H759A)-SV40	rpsL (genomic)	3'	17	175 bp

	(SEQ ID NO:79)				
2.25	SV40-nLbCas12a-XTEN-MMLV-RT-SV40 (SEQ ID NO:73)	aadA Asp132 (plasmid)	3'	12	140 bp plus 21 bp decoy hairpin*
2.31	SV40-MMLV-RT-XTEN-nLbCas12a(H759A)-SV40 (SEQ ID NO:83)	aadA asp132 (plasmid)	3'	12	140 bp plus 21 bp decoy hairpin*

*Decoy hairpin sequence: TAAGTCTCCATAGAAATGGAGG SEQ ID NO:95.

Example 2. Precise editing activity in human cells

A further approach that uses the active form of Cas12a in conjunction with reverse transcriptase is shown **Fig. 12.** and outlined below.

- Nuclease active Cas12a is recruited to the site via spacer – target site interaction.
- Cas12a makes a double stranded break. Optionally, a 5' to 3' exonuclease is provided to degrade the non-template strand.
- Priming occurs using the tagRNA. The primer binding site (PBS) encodes the sequences to the right of the cleavage site, complementary to the template strand DNA.
- Reverse transcriptase (MMuLV-RT (5M)) extends from the priming site or primer on the target nucleic (dashed line = the extension), encoding the desired change within the newly synthesized strand.
- Resolution of DNA intermediates via mismatch repair and DNA ligation generates an edited, new DNA strand.

Methods:

Extended guide RNAs were designed to target two genomic sites in HEK293T cells, DNMT1 and FANCF1. Varying combinations of primer binding sites (PBS) and reverse transcriptase template (RTT) lengths were assayed. The guide RNAs encoded a two base change in the PAM region of the target guides, corresponding to TT to AA at the -2 and -3 position (counting TTTV PAM as -4 to -1 position). The guide extensions were fused to either the 5' or the 3' end of the guide RNA.

Plasmids encoding an RNase-dead mutant LbCas12a (H758A), reverse transcriptase (MMuLV-RT(5M)), and optionally an exonuclease (one of T5 Exonuclease, T7 Exonuclease, RecE, and RecJ), and an extended guide RNA were transfected into HEK293T cells grown at 70% confluency using Lipofectamine™ 3000 according to manufacturer's protocol. Cells were harvested after 3 days and gene editing was quantified by next generation sequencing.

Results:

We observed intended precise editing for both sites targeted. Depending on the guide design, we observed up to 0.5% editing at the FANCF1 site (**Fig. 13**) and up to 1.7% at the DMNT1 site (**Fig. 14**). Use of exonuclease improved editing efficiency in some guide designs.

Table 9. Guide design used to target the FANCF1 site (**Fig. 13**).

FANCF1				
pWISE	3' or 5'	RTT length (bases)	PBS length (bases)	% Precise Editing
pWISE878	N/A	0	0	0
pWISE2928	3'	74	48	0.17289
pWISE2929	3'	52	48	0.54658
pWISE2930	3'	44	48	0.10525
pWISE2931	3'	36	48	0
pWISE2932	3'	74	24	0.28148
pWISE2934	3'	44	24	0
pWISE2935	3'	36	24	0
pWISE2936	3'	74	16	0
pWISE2937	3'	52	16	0.20349
pWISE2938	3'	44	16	0.12821
pWISE2940	3'	74	8	0
pWISE2941	3'	52	8	0
pWISE2942	3'	44	8	0
pWISE2943	3'	36	8	0
pWISE2945	5'	52	48	0
pWISE2946	5'	44	48	0
pWISE2947	5'	36	48	0.10335
pWISE2948	5'	74	24	0
pWISE2949	5'	52	24	0
pWISE2950	5'	44	24	0
pWISE2951	5'	36	24	0

Table 10. Guide design used to target the DMNT1 site (**Fig. 14**).

DMNT1				
pWise	3' or 5'	RTT length (bases)	PBS length (bases)	% Precise Editing
pWise258	N/A	0	0	0
pWise2960	3'	74	48	0.77529
pWise2961	3'	52	48	0.3139
pWise2963	3'	36	48	1.17854
pWise2966	3'	44	24	0.30752
pWise2967	3'	36	24	0.71539
pWise2971	3'	36	16	0.96806
pWise2973	3'	52	8	0.23422
pWise2975	3'	36	8	0.53485
pWise2976	5'	74	48	0.33196
pWise2977	5'	52	48	0.77164
pWise2978	5'	44	48	1.17289
pWise2979	5'	36	48	1.72435
pWise2980	5'	74	24	0.3538
pWise2981	5'	52	24	0.44055
pWise2982	5'	44	24	0.55662
pWise2983	5'	36	24	1.55194

Table 11. Example extended guide nucleic acids (tagRNAs)

pWise	Name/SEQ ID NO	Cas12a species	Spacer	tagRNA extension sequence	Intended Precise Edit (PAM is denoted as position -4, -3, -2, -1)
pWise 2960	tagRNA 1 173	LbCas12a	CCTCACTC CTGCTCG GTGAATTT	ACAGCAGGCCTTTGGTCAGGTTGGC TGCTGGGCTGGCCCTGGGGCCGTA ACCCTCACTCCTGCTCGGTGAATTT GGCTCAGCAGGCACCTGCCTCAGCT GCTCACTTGAGCCTCTGGGTCTA	TT to AA at position (-3, -2)
pWise 2961	tagRNA 2 174	LbCas12a	CCTCACTC CTGCTCG GTGAATTT	GGCTGCTGGGCTGGCCCTGGGGCC GTAACCCTCACTCCTGCTCGGTGAA TTTGGCTCAGCAGGCACCTGCCTCA GCTGCTCACTTGAGCCTCTGGGTCT A	TT to AA at position (-3, -2)
pWise 2962	tagRNA 3 175	LbCas12a	CCTCACTC CTGCTCG GTGAATTT	GGCTGGCCCTGGGGCCGTAACCCTC ACTCCTGCTCGGTGAATTTGGCTCA GCAGGCACCTGCCTCAGCTGCTCAC TTGAGCCTCTGGGTCTA	TT to AA at position (-3, -2)
pWise 2963	tagRNA 4 176	LbCas12a	CCTCACTC CTGCTCG GTGAATTT	CTGGGGCCGTAACCCTCACTCCTGC TCGGTGAATTTGGCTCAGCAGGCAC CTGCCTCAGCTGCTCACTTGAGCCT CTGGGTCTA	TT to AA at position (-3, -2)
pWise 4673	tagRNA 5 177	LbCas12a	CCTCACTC CTGCTCG GTGAATTT	ACAGCAGGCCTTTGGTCAGGTTGGC TGCTGGGCTGGCCCTGGGGCCGTTT CCCTCACTCCTGGACGGTGAATTTG	CT to GA at position (12, 13)

				GCTCAGCAGGCACCTGCCTCAGCTG CTCACTTGAGCCTCTGGGTCTA	
pWISE 4674	tagRNA 6 178	LbCas12a	CCTCACTC CTGCTCG GTGAATTT	ACAGCAGGCCTTTGGTCAGGTTGGC TGCTGGGCTGGCCCTGGGGCCGTTT CCCTCACTCCTGCTGCGTGAATTTG GCTCAGCAGGCACCTGCCTCAGCTG CTCACTTGAGCCTCTGGGTCTA	CG to GC at position (14, 15)
pWISE 4735	tagRNA 7 179	LbCas12a	GCGGATG TTCCAATC AGTACGC A	AATAGCATTGCAGAGAGGCGTATC ATTTGCGGATGTTGGAATCAGTAC GCAGAGAGTCGCCGTCTCCAAGGT GAAAGCGGAAGTAGGGCCTTCGCG CAC	CC to GG at position (10, 11)
pWISE 4736	tagRNA 8 180	LbCas12a	GCGGATG TTCCAATC AGTACGC A	AATAGCATTGCAGAGAGGCGTATC ATTTGCGGATGTTCTTTTCACTAC GCAGAGAGTCGCCGTCTCCAAGGT GAAAGCGGAAGTAGGGCCTTCGCG CAC	AA to TT at position (12, 13)
pWISE 4906	tagRNA 9 181	AsCas12a	CCTCACTC CTGCTCG GTGAATTT	ACAGCAGGCCTTTGGTCAGGTTGGC TGCTGGGCTGGCCCTGGGGCCGTA ACCCTCACTCCTGCTCGGTGAATTT GGCTCAGCAGGCACCTGCCTCAGCT GCTCACTTGAGCCTCTGGGTCTA	TT to AA at position (-3, -2)
pWISE 4907	tagRNA 10 182	AsCas12a	CCTCACTC CTGCTCG GTGAATTT	GGCTGCTGGGCTGGCCCTGGGGCC GTAACCCTCACTCCTGCTCGGTGAA TTTGGCTCAGCAGGCACCTGCCTCA GCTGCTCACTTGAGCCTCTGGGTCT A	TT to AA at position (-3, -2)
pWISE 4908	tagRNA 11 183	AsCas12a	CCTCACTC CTGCTCG GTGAATTT	GGCTGGCCCTGGGGCCGTAACCCTC ACTCCTGCTCGGTGAATTTGGCTCA GCAGGCACCTGCCTCAGCTGCTCAC TTGAGCCTCTGGGTCTA	TT to AA at position (-3, -2)
pWISE 4909	tagRNA 12 184	AsCas12a	CCTCACTC CTGCTCG GTGAATTT	CTGGGGCCGTAACCCTCACTCCTGC TCGGTGAATTTGGCTCAGCAGGCAC CTGCCTCAGCTGCTCACTTGAGCCT CTGGGTCTA	TT to AA at position (-3, -2)
pWISE 4438	tagRNA 13 185	LbCas12a	CTGATGG TCCATGTC TGTTACTC	GCACTCTGCCACTTATTGGGTCAGC TGTTAACATCAGTACGTTAATGTAA CCTGATGGTCCATGTCTGTTACTCG CCTGTCAAGTGGCGTGACACCGGG CGTGTTCCCCAGAGTGACTTTTC	TT to AA at position (-3, -2)
pWISE 4439	tagRNA 14 186	LbCas12a	CTGATGG TCCATGTC TGTTACTC	AGCTGTTAACATCAGTACGTTAATG TAACCTGATGGTCCATGTCTGTTACT CGCCTGTCAAGTGGCGTGACACCG GGCGTGTTCCCCAGAGTGACTTTTC	TT to AA at position (-3, -2)

The effect of exonuclease transfection on precise editing activity at DMNT1 site is shown in Fig. 15 (normalized to no exonuclease treatment; pUC19 = 1). Exonuclease improves editing with some guide configurations.

Example 3. Variations in REDRAW protein architecture

The methods of the present invention (i.e., REDRAW) were tested using different protein architectures/constructs for LbCas12a and RT(5M) including: (1) where the reverse transcriptase (RT(5M)) is provided by overexpressing the RT in the cell; (2) a construct in which SunTag (GCN4, e.g., **SEQ ID NO:23**, **SEQ ID NO:24**) is fused to the CRISPR-Cas effector protein (e.g., LbCpf1) and the RT (RT(5M)) is recruited to the site of editing by fusing it to an antibody (e.g., single chain variable fragment (scFv) antibody) that binds to the SunTag fused to the CRISPR-Cas effector protein; and (3) where the reverse transcriptase (RT(5M)) is fused to the N-terminus or C-terminus of the CRISPR-Cas effector protein (e.g., LbCpf1 (LbCas12a), e.g., LbCpf1 (H759A)) (e.g., RT(5M)-LbCpf1 (H759A) or LbCpf1 (H759A)-RT(5M)). The results are shown in **Fig. 16**. All constructs showed evidence of RT(5M) and tagRNA dependent REDRAW activity using four different tagRNA constructs. In this example, active recruitment with SunTag did not enhance activity. It may be that overexpression of non-tagged constructs allows sufficient level of protein concentration in the cell and therefore, under these conditions, this is not a rate-limiting aspect for generating a precise edit.

MS2/MCP system

In addition to the architecture tested above, the MS2/MCP system was also evaluated for use with the constructs and methods of the invention. MS2 hairpin RNA structure binds to MCP protein. MS2 hairpin can be added to the tagRNA. In this example, a MS2 hairpin structure was added to the 3' end of the tagRNA, and MCP was fused RT(5M) in order to recruit RT(5M) to the target site.

LbCas12a H759A with RT(5M) was transiently expressed without MCP (in trans control), or with MCP-RT(5M) (fusion construct). This architecture was tested using two tagRNAs, tagRNA5 and tagRNA6. We also compared the different tagRNA versions where tagRNA5 and tagRNA6 were modified with MS2 sequence at its 3' end. The results are shown in **Fig. 37**. Comparing MCP-RT(5M) and RT(5M), the MS2 tagRNAs and MCP-RT(5M) did not result in an increase in precise editing efficiency. The MCP fusion may not be increasing precise editing efficiency under these experimental conditions because RT concentration is not rate limiting. However, an increase in editing efficiency was noted for the tagRNA having MS2 at its 3' end. The MS2 structure at the 3' end of the tagRNA may stabilize the tagRNA and reduce its degradation.

Example 4. 5'-3' exonucleases for use with methods of the invention (REDRAW)

5'-3' exonuclease may be useful with the methods of the invention by degrading the DNA at both ends of the double-stranded break. Thus, a 5'-3' exonuclease may (1) allow a more robust RNA-DNA duplex formation (a substrate for RT-mediated polymerization) by degrading a strand that is normally base paired with the DNA strand that will be elongated and/or (2) allow the cell to favor the use of RT-synthesized DNA for use in DNA repair by degrading the region that will be overwritten by the RT. See, for example, the schematic in **Fig. 17**.

In this example, the exonucleases tested included are those listed in **Table 11**.

Table 12. 5'-3' Exonucleases tested

Name	Full name	Source species	SEQ ID NO
RecE	RecE	E.coli	129
RecJ	RecJ	E.coli	130
T5_Exo	Exonuclease from T5 Phage	T5 Phage	131
T7_Exo	Exonuclease from T7 Phage	T7 Phage	132
Lambda_Exo	Exonuclease from Lambda Phage	Lambda phage	133
sbcB	Exodeoxyribonuclease I	E.coli	134
hExo1	Human exonuclease 1	Human	135

The 5'-3' exonucleases were fused to the C-terminus of LbCas12a (H759A). Fusion constructs were transfected into HEK293T cells along with Reverse transcriptase (5M) construct and a plasmid expressing an appropriate tagRNA encoding a precise mutation.

Cells were harvested 3 days post transfection and DNA was analyzed using High Throughput Sequencing (HTS). The results are shown in **Fig. 18**. Here, RT is expressed in trans (without recruitment), and the 5'-3' exonucleases are fused to the C-terminus of LbCpf1 H759A. Compared to the construct in which exonuclease is not present (LbCpf1 H759A only), fusion of T7_Exo, in particular, improves REDRAW precise editing in three of the four tagRNAs tested.

Fig. 19 provides additional 5'-3' exonuclease testing with the methods of the invention (REDRAW) and under the same conditions noted above. Specifically, **Fig. 19** shows the percent precise editing with REDRAW using either the 5'-3' exonuclease sbcB (**SEQ ID**

NO:134) or the 5'-3' exonuclease Exo (**SEQ ID NO:135**) each fused to the C-terminus of a Cas polypeptide (LbCpf1). RT(5M) (**SEQ ID NO:97**) is expressed in trans (no recruitment). In contrast to T7_Exo (**SEQ ID NO:132**), exonucleases sbcB and Exo did not improve REDRAW.

5 5'-3' exonucleases were also tested in trans with the methods of the invention. The results are provided in **Fig. 20**. The LbCpf1 and RT(5M) (**SEQ ID NO:97**) are provided as fusion proteins. The right side of **Fig. 20** shows results with the RT fused to the N-terminus of the LbCpf1 (RT(5M)-LbCpf1 (H759A)) and the left side of the figure shows the results using an RT fused to the C-terminus of the LbCpf1 (LbCpf1 (H759A)-RT(5M)). **Fig. 20**
10 shows that when 5'-3' exonucleases are expressed in trans, without being fused to LbCpf1, the editing rate does not increase compared to treatment without 5'-3' exonucleases. Thus, a benefit from use of a 5'-3' exonuclease with the methods of the invention is observed when the 5'-3' exonucleases is fused to the CRISPR-Cas effector protein.

15 **Example 5. Mutations modulating DNA binding affinity of Cas12a for use with REDRAW**

Lowering the DNA binding affinity of a CRISPR-Cas effector protein was envisioned to allow better dissociation of the CRISPR-Cas effector protein from the target site, thereby
20 inducing a double-stranded DNA break. This may allow faster formation of intermediates that promote sequence replacement by RT and increase the efficiency of editing with the methods of the invention (REDRAW)

In this example, positively charged residues in Cas12a (LbCas12a) that interact with DNA backbone were mutated to alanine. Specifically, the following three mutations, K167A,
25 K272A, K349A (with reference to the amino acid position numbering of **SEQ ID NO:1** or **SEQ ID NO:148**), were cloned into LbCas12a H759A as single, double or triple mutants (K167A, K272A, K349A, K167A+ K272A, K167A+ K349A, K272A+ K349A, and K167A+ K272A + K349A). In this case, the H759A mutation (**SEQ ID NO:148**) was used to deactivate RNA processing ability of LbCas12a to facilitate 5' tagRNA extensions to the
30 crRNA.

LbCas12a containing the various combinations of binding affinity mutations were transfected into HEK293T cells along with plasmids encoding RT(5M) and a tagRNA encoding a precise edit. Cells were harvested three days post transfection and DNA was analyzed using High Throughput Sequencing (HTS). Certain mutation combinations were
35 shown to improve the precise editing of the methods of the invention (**Fig. 21**).

Example 6. Single-stranded DNA binding proteins (ssDNA BP) for use with REDRAW

Use of single-stranded DNA binding proteins (ssDNA BP) may potentially improve REDRAW by stabilizing the ssDNA that are generated during the reaction, by protecting DNA strands from degradation and make the same available for RT-mediated priming and polymerization. A selection of ssDNA BP were tested with the methods of the invention (see **Table 12**)

Table 13. Single-stranded DNA binding proteins evaluated

Name	Source species	SEQ ID NO
hRad51(S208E_A209D)	human	123
hRad52	human	124
BsRecA	Bacillus subtilis	125
EcRecA	E.coli	126
T4SSB	Escherichia phage P1	127
Brex27	Human	128

- The ssDNA BPs set forth in **Table 12** were expressed in trans or as a fusion with Cas12a, also in the presence of RT(5M) (trans). The ssDNA BPs tested were hRad51_s208E_A209D (**SEQ ID NO:123**), hRad52 (**SEQ ID NO:124**), BsRecA (**SEQ ID NO:125**), EcRecA (**SEQ ID NO:126**), T4SSB (**SEQ ID NO:127**) and Brex27 (**SEQ ID NO:124**). The results are shown in **Fig. 22** and **Fig. 23**. Trans expression of the ssDNA BPs did not improve the percent of precise editing when compared to a control (pUC19) (see **Fig. 22**). The fusion proteins also failed to show an improvement, with the exception of the N-terminal and C-terminal fusion of Brex27 with Cas12 (see, **Fig. 23**). Brex27 is a peptide that is known to recruit Rad51 in situ and stabilize its interaction with ssDNA.

Example 7. Evaluation of Gam protein for use in REDRAW

Gam protein may be helpful in reducing the formation of indels during REDRAW by preventing NHEJ. Gam binds to a double-stranded DNA break, preventing the DNA end from being processed. Gam may be used to reduce indel formation during cytosine base editing.

- To evaluate the usefulness of Gam protein with the methods of the invention, Gam protein (*Escherichia* phage Mu Gam protein) (**SEQ ID NO:147**) was fused to either a CRISPR-Cas effector protein (LbCas12a H759A) (**SEQ ID NO:148**) or to RT(5M) (**SEQ ID NO:53**). Plasmids encoding LbCas12a H759A, RT(5M), and tagRNA encoding a precise mutation were transfected into HEK293T cells. Target DNA was analyzed after three days with high throughput sequencing. The results are shown in **Fig. 24** and **Fig. 25**.

In **Fig. 24**, the reverse transcriptase (RT) is expressed in trans, either as a native sequence (e.g., RT(5M)) or fused at its N-terminus to the Gam protein (e.g., Gam-RT(5M)). These constructs are expressed concurrently with either LbCas12a (H759A) or with an LbCas12a (H759A) having a Gam protein fused to its N-terminus (e.g., Gam-LbCas12a H759A). In **Fig. 25** the Gam protein is provided in trans, as a fusion protein with the reverse transcriptase (N-terminal fusion; Gam-RT(5M)) and/or as a fusion protein with the CRISPR-Cas effector polypeptide (e.g., Gam-LbCas12a H759A). The results show that in some cases Gam protein may be used to reduce indel formation but overall efficiency of editing using methods of the invention is not improved by inclusion of Gam protein.

Example 8. Evaluation of primer binding site (PBS) length and reverse transcriptase template (RTT) length

The length of RTT and PBS in a tagRNA of the invention was varied to evaluate the effect of length on editing. LbCas12a, RT(5M), and tagRNAs having varying lengths of RTT and PBS were transfected into HEK293T cells and analyzed for editing rate three days post transfection using High Throughput Sequencing (HTS). The results are provided in **Fig. 26**. The top and bottom panels of **Fig. 26** show the results using two different spacers (top panel: pwsp143 (GCTCAGCAGGCACCTGCCTCAGC) (**SEQ ID NO:136**), bottom panel: pwsp139 (CTGATGGTCCATGTCTGTACTC) (**SEQ ID NO:137**). While the results varied with the spacer used, and many different lengths for both the RTT and PBS showed good editing efficiency. One optimal combined PBS length and RTT length may be 48 nucleotides and 52 nucleotides, respectively.

Example 9. Evaluation of edit placement in a tagRNA

REDRAW efficiencies can vary depending on where the desired edit is located within the reverse transcriptase template (RTT) of the tag RNA. In this example, the effect of the location of the edit in the RTT on the percent editing was evaluated. The edit location and results are provided in bold in **Fig. 27**. The upper and lower panels provide different RTT sequences in which the edit location was varied (upper panel RTT: **SEQ ID NO:187**; lower panel RTT: **SEQ ID NO:188**). The 'Edit location' column in both the upper and lower panels of **Fig. 27** shows the reverse complement of the first 26 bases of RTT, which corresponds to the PAM sequence (TTTC) and the 23-base spacer sequence. We tiled the double mutation along the RTT such that when the desired edit is introduced to the DNA, the spacer used for REDRAW is no longer complementary. Editing was determined to be

effective with the edit placement in many locations in the RTT. In some cases, placing the edit in a position in the RTT sequence that corresponds to nucleotide 12-15 (TTTN PAM of LbCas12a is defined as position -4,-3,-2,-1, respectively) provided a very high level of editing.

5

Example 10. REDRAW editing using different CRISPR-Cas effector proteins

REDRAW was envisioned to be compatible with alternate CRISPR-Cas effector proteins that are able to generate double-stranded DNA breaks. In this example, LbCas12a with cas9 (SpCas9), BhCas12b, AsCas12a (EnAsCas12a) showing that alternate CRISPR-Cas effector proteins can be used successfully with the methods of this invention (REDRAW).

10

Cas9

RT(5M), tagRNA encoding a precise edit, and two forms of Cas9 (Cas9 (nuclease), nCas9 (D10A) (nickase)) were transformed into HEK293T cells and expressed. The cells were harvested three days after transfection and target amplicons were sequenced using high throughput sequencing (HTS). The lengths of PBS and RTT were varied, and extensions were added to both 3' and 5' end of the guide RNA (denoted as '3' extension' or '5' extension' in **Fig. 28**). The tagRNA extensions that were used targeted four different target sites (spacers: pwsp10: GAGTCCGAGCAGAAGAAGAA (**SEQ ID NO:140**); pwsp621: GCATTTTCAGGAGGAAGCGA (**SEQ ID NO:141**); pwsp15: GTCATCTTAGTCATTACCTG (**SEQ ID NO:142**); pwsp11: GGAATCCCTTCTGCAGCACC (**SEQ ID NO:143**)). The results are provided in **Fig. 28**. Precise RT-mediated editing was observed using both Cas9 and nCas9 (D10A) using multiple different spacer sequences, however, the nuclease version performed best. Further, while both 3' and 5' tagRNA extensions were effective in REDRAW, the 3' extension of the extended guide RNA performed best.

25

BhCas12b

RT(5M), tagRNA encoding a precise edit and BhCas12b v4 (which is an engineered high efficiency version of BhCas12b) were transformed into HEK293T cells and expressed. The cells were harvested three days after transfection and target amplicons were sequenced using high throughput sequencing (HTS). The lengths of PBS and RTT were varied and extensions were added to both 3' and 5' end of the guide RNA (denoted as 3' or 5' in **Fig. 29**).

30

The tagRNA extensions that were used targeted three different target sites (spacers: PWsp1099: ACGTACTGATGTTAACAGCTGA (**SEQ ID NO:144**); PWsp1098: GGTCAGCTGTTAACATCAGTAC (**SEQ ID NO:145**); PWsp1094: TCCAGCCCCTGGCCCTGTAAA (**SEQ ID NO:146**)). The results are provided in **Fig.**

- 5 **29.** Precise RT-mediated editing was observed using BhCas12b v4 and multiple different spacer sequences. Certain combinations of RTT and PBS lengths resulted in higher editing than others when using BhLbCas12b. In general, 3' extension of tagRNA provided more consistent editing than 5' extension when using BhLbCas12b, although editing was detected using both forms of tagRNA.

10

EnAsCas12a

AsCas12a is a homolog of LbCas12a and EnAsCas12a is the engineered version of AsCas12a. The H800A mutation in EnAsCas12a corresponds to H759A mutation in LbCas12a, which is a mutation that inactivates crRNA-processing ability of Cas12a.

- 15 RT(5M), tagRNA encoding a precise edit and EnAsCas12a H800A (EnAsCpf1 H800A) were transformed into HEK293T cells and expressed. In this case, the reverse transcriptase was provided as a fusion protein with the EnAsCas12a (C-terminal fusion (EnAsCas12a-RT) and N-terminal fusion (RT-EnAsCas12a)). The cells were harvested three days after transfection and target amplicons were sequenced using high throughput
20 sequencing (HTS). Precise RT-dependent and tagRNA-dependent edit was observed using EnAsCas12a using multiple different tagRNA sequences. The tagRNA extensions that were used targeted a single site (spacer: CCTCACTCCTGCTCGGTGAATTT (**SEQ ID NO:171**)).

- The results are provided in **Fig. 30**, which shows that in the presence of various
25 tagRNAs, both the N-terminal and C-terminal fusions of RT and EnAsCas12a resulted in precise editing. EnAsCas12a without RT fusion was used as a control and showed no or very low editing.

Example 11. Editing in yeast

- 30 In addition to showing that human cells can be edited using the methods of the invention, the same was also evaluated in *Saccharomyces cerevisiae* (yeast), a eukaryote. *S. cerevisiae* is an attractive organism for evaluating the methods of this invention for several reasons including, for example: (1) *S. cerevisiae* utilizes NHEJ repair processes; double-stranded breaks in the genome are not lethal, unlike in prokaryotic organisms (such as *E. coli*)

that are often used in directed evolution experiments; (2) yeast grow relatively quickly, allowing rapid testing and tuning many of the conditions for the methods of the invention (REDRAW); (3) thousands of yeast strains are readily available; and (4) large libraries of biomolecules (protein, RNA, etc.) may be investigated in yeast.

- 5 The *S. cerevisiae* strain W303-1a (hereinafter "ScW303-1a") was selected for this example. The genotype of ScW303-1a is: MATa ade2-1 ura3-1 his3-11 trp1-1 leu2-112 can1-100. Targets for editing in this strain include *ADE2*, *CAN1*, *HIS3*, *LYS2*, *TRP1*, and *URA3*. Sanger sequencing was used to confirm the loci sequences for each PCR product. All loci that were sequenced were as expected, except for *ADE2*. The *ADE2* locus was expected
10 to have a stop codon at Gln64; however, sequencing showed that instead of a stop codon at Gln64, a tyrosine codon was present. As a consequence, a custom strain with a modified *ADE2* locus was constructed in order to test REDRAW at that locus. The modified strain was named ScDS21.6. **Table 13** provides the genomic targets selected for testing in yeast.

15 **Table 14.** Yeast genomic targets for REDRAW editing.

Strain (Target)	Mutation	Auxotrophy	Additional Phenotype	Comments
ScDS21.6 (<i>ADE2</i>)	Amino acid 156 GGA -> TAA	Adenine	Red colonies	Generated in-house (same effect as <i>ADE2-I</i> mutation)
W303-1a (<i>ura3-1</i>)	Amino acid 234 GGA -> GAA	Uracil	5-FOA resistance	Li et al. unpublished

Example spacers for targeting these sites included:

PWsp1643 (*ADE2* target): 5' – GCATACGATGGAAGAGGTAACCTT – 3' (**SEQ ID NO:159**)

- 20 PWsp1894 (*ADE2* target): 5' – GCATACGATTAAAGAGGTAACCTT – 3' (**SEQ ID NO:160**)

PWsp1665 (*URA3-I* target): 5' – CAAATAGTCCTCTTTCAACAATA – 3' (**SEQ ID NO:161**)

- 25 Example Primer Binding Site Sequences:

48-bp primer binding site for *ADE2* target:

5' – CGTTGTAAAGAATAAGGAAATGATTCCGGAAGCTTTGGAAGTACTGAA – 3' (**SEQ ID NO:162**)

48-bp primer binding site for *URA3-I* target:

- 30 5' – ATAATGTCAGATCCTGTAGAGACCACATCATCCACGGTTCTATACTGT – 3' (**SEQ ID NO:163**)

Example Reverse Transcriptase Template Sequences:

(A) *ADE2* target:

40-bp RTT: 5' – TGAAGTCGAGGACTTTGGCATACGATGGAAGAGGTAAGT – 3' (SEQ ID NO:164)

50-bp RTT: 5' – CCATTCGTCTTGAAGTCGAGGACTTTGGCATACGATGGAAGAGGTAAGT – 3' (SEQ ID NO:165)

72-bp RTT: 5' – TGTTGGAAGAGATTTGGGTTTTCCATTCGTCTTGAAGTCGAGGACTTTGGCATACGATGGAAGAGGTAAGT – 3' (SEQ ID NO:166)

10 (B) *URA3-1* target

47-bp RTT: 5' – CTACCTTAGCATCCCTTCCCTTTGCAAATAGTCCTCTCTCAACAATA – 3' (SEQ ID NO:167)

55-bp RTT: 5' – TTCACCCTCTACCTTAGCATCCCTTCCCTTTGCAAATAGTCCTCTCTCAACAATA – 3' (SEQ ID NO:168)

63-bp RTT: 5' – CTGTAAACGTTACCCCTCTACCTTAGCATCCCTTCCCTTTGCAAATAGTCCTCTCTCAACAATA – 3' (SEQ ID NO:169)

20

Example LbCas12a crRNA scaffold:

5' – TAATTTCTACTAAGTGTAGAT – 3' (SEQ ID NO:170)

The protein expression vector pESC-LEU was used because (1) it includes a yeast selectable marker, LEU2, that is compatible with the ScW303-1a strain, (2) the GAL promoter system in the plasmid provides strong control of protein expression, (3) the yeast origin of replication, 2 μ , is high copy, allowing for high level of protein expression and (4) the *E. coli* origin of replication (pUC origin) and the selectable marker, AmpR, are also present, allowing all vector manipulation and cloning in *E. coli* prior to working in yeast.

30 The following CRISPR-Cas effector protein and reverse transcriptase configurations were used:

- LbCas12a + C-terminally fused MMLV-RT(5M) (SEQ ID NO:155)
- LbCas12a + N-terminally fused MMLV-RT(5M) (SEQ ID NO:157)

The LbCas12a fusions were placed under control of inducible GAL1 promoter (pol II promoter) and the crRNA and tagRNAs were expressed from the constitutive *SNR52* promoter (pol III promoter).

In addition, the following tagRNA configurations were tested with the two LbCas12a and RT configurations: (1) absence of a 3' pseudoknot, (2) presence of a pseudoknot, either (a) a pseudoknot referred to as a "decoy" pseudoknot (see Fig. 7, SEQ ID NO:203) or (b) a pseudoknot referred to as tEvoPreQ1 pseudoknot (SEQ ID NO:158). In addition, three different reverse transcriptase template (RTT) lengths (47, 55, 64 nucleotides, or 40, 50 and

72 nucleotides) were each tested with a primer binding site (PBS) having a fixed length of 48 nucleotides. These configurations are set forth in **Table 15**.

Table 15. Example REDRAW configurations tested in yeast

pWISE	Target	PBS	RTT	REDRAW Editor	3' Pseudoknot
5584 5585 5586	<i>URA3-1</i>	48-bp	47-bp 55-bp 63-bp	C-terminal RT	None
5890 5591 5592	<i>URA3-1</i>	48-bp	47-bp 55-bp 63-bp	N-terminal RT	None
5853 5854 5855 5856 5857 5858	<i>URA3-1</i>	48-bp	47-bp 55-bp 63-bp 47-bp 55-bp 63-bp	C-terminal RT	Decoy Pseudoknot Decoy Pseudoknot Decoy Pseudoknot tEvoPreQ1 tEvoPreQ1 tEvoPreQ1
5865 5866 5867 5868 5869 5870	<i>URA3-1</i>	48-bp	47-bp 55-bp 63-bp 47-bp 55-bp 63-bp	N-terminal RT	Decoy Pseudoknot Decoy Pseudoknot Decoy Pseudoknot tEvoPreQ1 tEvoPreQ1 tEvoPreQ1
5581 5582 5583	<i>ADE2</i>	48-bp	40-bp 50-bp 72-bp	C-terminal RT	None
5587 5588 5589	<i>ADE2</i>	48-bp	40-bp 50-bp 72-bp	N-terminal RT	None
5848 5849 5850 5851 5852	<i>ADE2</i>	48-bp	50-bp 72-bp 40-bp 50-bp 72-bp	C-terminal RT	
5860 5861 5862 5863 5864	<i>ADE2</i>	48-bp	50-bp 72-bp 40-bp 50-bp 72-bp	N-terminal RT	
5082 (Neg. Control) no tagRNA	<i>ADE2</i> <i>URA3-1</i>	N/A	N/A	C- terminally fused RT	N/A

5083 (Neg. Control) no tagRNA	<i>ADE2</i> <i>URA3-1</i>	N/A	N/A	N-terminally fused RT	N/A
5074 (Cutting Control)	<i>ADE2</i>	N/A	N/A	LbCas12a	N/A
5077 (Cutting control)	<i>URA3-1</i>	N/A	N/A	LbCas12a	N/A

REDRAW was tested in *S. cerevisiae* by first transforming the vectors of interest into either yeast strain ScDS21.6 (*ADE2* target site) or yeast strain ScW303-1a (*URA3* target site) via the PEG/LiAc heat shock method. Transformants were plated out onto synthetic complete media lacking leucine, with 2% glucose as the carbon source (SC-LEU + 2%Glu). After approximately 48-72 hours, single colonies were then picked into 3-mL of liquid SC-LEU + 2% raffinose (SC-LEU + 2% Raff). The cultures were grown up at 28°C with shaking at 200 rpm for approximately 36 hours, until the OD₆₀₀ reached ~1.8. 1.5 ODs of cells was then spun down at room temperature in the centrifuge and brought back up in 3mL of protein expression media, SC-LEU + 1% raffinose + 1% galactose (SC-LEU + 1% Raff + 1% Gal). Expression cultures were grown at 28°C, with 200 rpm shaking for 4 hours. The expression cultures were then removed from the shaking incubator and centrifuged. The supernatant was then pipetted off, and 3mL of SC-LEU + 2% Glu was added. The cells were then allowed 90 additional minutes of growth at 28°C and 200 rpm. After 90 minutes, the OD₆₀₀ of the culture was checked. About 1 OD yeast cells (about 1×10⁷ yeast cells) of each culture was pelleted in the centrifuge at room temperature. The supernatant was removed, and each culture was re-suspended in 200 uL of sterile water. About half of the resuspended culture (0.5 OD's) (about 5×10⁶ yeast cells) was plated onto synthetic complete plates lacking either adenine (SC-ADE) or uracil (SC-URA) to select for edited colonies, and the other half was plated onto SC-LEU plates (non-selective, to see how many cells were in about 0.5 measured OD's). The plates were grown at 28°C for approximately 3 days. Colonies were then counted and recorded. Colonies were selected from either SC-ADE / SC-URA plates or SC-LEU (negative control) plates, and the target loci were amplified using colony PCR. Sanger sequencing was used to analyze the target loci, which confirmed that the intended edits were made (2-bp change in *ADE2*: AA156 TAA -> GGA and 1-bp change in *URA3-1*: AA 234 GGA -> GAA).

Each of the LbCas12a and RT configurations / tagRNA combinations were tested at two different target sites in yeast and the results are provided in **Fig. 31** and **Fig. 32**. **Fig. 31** show the results of the editing of the *URA3-1* target gene (*URA3-1*: 1-bp change (AA 234 GGA -> GAG) (edit repairs adenine auxotrophy) with the upper panel showing the results with the LbCas12-RT C-terminal fusion and the lower panel showing the results for the RT-LbCas12 N-terminal fusion. **Fig. 32** show the results of the editing of the *ADE2* target gene (*ADE2*: 2-bp change (AA 156 TAA -> GGA) (edit repairs uracil auxotrophy) with the upper panel showing the results with the LbCas12-RT C-terminal fusion and the lower panel showing the results for the RT-LbCas12 N-terminal fusion. While all configurations were able to edit the *URA3-1* gene to produce viable colonies (repairing adenine auxotrophy), the most efficient configuration included a pseudoknot and the RTT having a length of 55 nucleotides (**Fig. 31**). The RT, LbCas12a C-terminal fusion was most efficient with the "decoy" pseudoknot and the RT, LbCas12a N-terminal fusion was most efficient with the tEvoPreQ1 pseudoknot (**Fig. 31**). Editing of *ADE2* in yeast showed similar results in that the RT, LbCas12a C-terminal fusion was most efficient with the "decoy" pseudoknot and the RT, LbCas12a N-terminal fusion was most efficient with the tEvoPreQ1 pseudoknot (**Fig. 32**). In the case of *ADE2*, editing was most efficient with an RTT having a length of 50 nucleotides.

Thus, this example showed that the methods of the invention are able to precisely edit yeast at both target sites and using either protein fusion configuration with the C-terminally fused RT configuration being slightly more efficient than the N-terminally fused RT for these two targets. The pseudoknots were observed to improve the efficiency of REDRAW editing in each of the configurations tested. Further, in the absence of the tagRNA and REDRAW editor, no growth is observed on the selective plates (SC-ADE or SC-URA), indicating that these REDRAW assays in yeast are very stringent and escape frequency is below the detection limit.

Example 12. Evaluation of ssRNA binding proteins in editing

Single-stranded RNA binding proteins (ssRNA BP) are proteins that interact nonspecifically with ribonucleic acids. Expressing ssRNA binding proteins when editing with the methods of the invention may stabilize the exposed tagRNA component (extended guide nucleic acid) from degradation by endogenous proteins. To test this, we expressed several RNA binding proteins as an N-terminal fusion to RT(5M)-LbCas12a(H759A).

The precise editing results using the ssRNA binding proteins, defensin (SEQ ID NO:152) and ORF5 (SEQ ID NO:153) are provided in **Fig. 33**. The ssRNA BP defensin and

the ssRNA BP ORF5 were each fused to the N-terminus of a RT-LbCas12 fusion protein (e.g., RT-LbCas12a). The editing is shown as compared to the same RT-Cas12a fusion protein that is not fused at its N-terminus to a ssRNA binding protein. Precise editing was shown to improve with the use of a ssRNA binding protein for one of the two tagRNAs (extended guide nucleic acids) tested.

Example 13. Evaluation of reverse transcriptase polypeptides having different mutations

The reverse transcriptase RT(5M) was engineered by introducing five mutations into wildtype RT sequence (Anzalone et al. *Nature* 576:149-157 (2019)). To evaluate whether the methods of the invention can be further optimized by using an RT domain having different or additional mutations compared to that of RT(5M), several reverse transcriptase(RT) proteins having different mutations and combinations of mutations, with or without the RT(5M) core mutations, were fused to LbCas12a (H759A) at the N-terminus. The RT domains tested included: RT(L139P, D200N, W388R, E607K), RT(L139P, D200N, T306K, W313F, W388R, E607K), RT(5M, F155Y, H638G), RT(5M, Q221R, V223M) and RT(5M, D524N). The mutations in RT(M) include D200N+L603W+T330P+T306K+W313F with reference to the amino acid sequence numbering of **SEQ ID NO:172** (see, **SEQ ID NO:53**). The reference RT for amino acid position numbering for those sequences that do not include RT(5M) mutations is **SEQ ID NO:172**. The reference RT for amino acid position numbering for those sequences that include RT(5M) mutations is **SEQ ID NO:53**. In each case, the RT was fused to the N-terminus of LbCas12a (H759A).

Fig. 34 shows the results. Compared to RT(5M) (left), several other RT domains having different combinations of mutations were able to increase the precise editing as compared to RT(5M). This result was influenced by the tagRNA (extended guide nucleic acid) that was used.

Example 14. Evaluation of 3' structured RNA motifs incorporated at the 3' end the tagRNA

Experiments were carried out to evaluate whether a structured RNA incorporated at 3' end of a tagRNA might further stabilize tagRNA and protect it from possible degradation. For this purpose, several RNA sequences known to form 3-D structures, including hairpins and pseudoknots, were appended to different tagRNAs.

Table 16. DNA sequences that correspond to RNA structures when transcribed and appended to the 3' end of tagRNA

AsCpf1BB	TAATTTCTACTCTTGTAGAT SEQ ID NO:189
BoxB	GGGCCCTGAAGAAGGGCCC SEQ ID NO:190
Pseudoknot (decoy)	TAAGTCTCCATAGAATGGAGG SEQ ID NO:95 (see also, SEQ ID NO:203) -
evopreQ1	TTGACGCGGTTCTATCTAGTTACGCGTTAAACCAACUAGAAA SEQ ID NO:191
fmpknot	GGAGGTCAGGGTCAGGAGCCCCCCCCTGAACCCAGGATAACCCTCAAAGTCG GGGGGCAACCC SEQ ID NO:192
mpknot	GGGTCAGGAGCCCCCCCCTGAACCCAGGATAACCCTCAAAGTCGGGGGGGCA ACCC SEQ ID NO:193
MS2	GGCCAACATGAGGATCACCCATGTCTGCAGGGCC SEQ ID NO:194
PP7	CCGGAGCAGACGATATGGCGTCGCTCCGG SEQ ID NO:195
SLBP	CCAAAGGCTCTTCTCAGAGCCACCCA SEQ ID NO:196
TAR	GGCCAGATCTGAGCCTGGGAGCTCTCTGGCC SEQ ID NO:197
ThermoPh	ATATAACCTTCACCATTAGGTTCAAATAATGGTAAT SEQ ID NO:198

Table 17. Sources for the RNA structures in Table 16

Name	Source
AsCpf1BB	Natural crRNA sequence from <i>Acidaminococcus sp.</i>
BoxB	Nat Biotechnol. 2016 May;34(5):528-30.
Pseudoknot (decoy)	Patent: WO2021092130
Pseudoknot (evopreQ1)	Nat Biotechnol. 2021 Oct 4. doi: 10.1038/s41587-021-01039-7.
fmpknot	Nature 480, 561–564 (2011)
mpknot	Nat Biotechnol. 2021 Oct 4. doi: 10.1038/s41587-021-01039-7.
MS2	Nat Biotechnol. 2016 May;34(5):528-30.
PP7	Nat Biotechnol. 2016 May;34(5):528-30.
SLBP	Cell. 2019 Jun 27;178(1):122-134.e12.
TAR	Cell. 2019 Jun 27;178(1):122-134.e12.
ThermoPh	Proc Natl Acad Sci U S A. 1999 Jun 8; 96(12): 6621–6625

5

The results of including various 3' structured RNAs in the compositions of the invention are provided in **Fig. 35**. In this experiment, RT(5M)-LbCas12a H759A with various tagRNAs was expressed with or without 3' RNA structures in HEK293T cells. After 3 days, the cells were harvested, and the precise editing efficiency was analyzed by high throughput sequencing. We observed that almost all 3' RNA structures on tagRNA can accommodate the methods of this invention (e.g., REDRAW). We did not observe an increase in REDRAW efficiency by using the 3' RNA structures in HEK293T cells.

10

Example 15. Evaluation of the use of chromatin modulating peptide fusions

Genome editing proteins can be occluded by nucleosomes that reduce their activity in living cells. Chromatin-modulating proteins/peptides may be helpful in addressing such affects by promoting chromatin exchange, histone modification, and epigenome modifications, thereby enhancing access by such programmable DNA binding proteins as, for example, Cas9 or Cas12a.

To evaluate this possibility, chromatin-modulating peptides, including CHD1 (e.g., **SEQ ID NO:199**), H1G (e.g., **SEQ ID NO:200**), HB1 (e.g., **SEQ ID NO:201**), and HN1 (e.g., **SEQ ID NO:202**) (see, e.g., Ding et al., *CRISPR J.* 2019 Feb;2:51-63) were fused to selected constructs of the invention in various fusion orientations as follows: HN1-RT(5M)-LBCas12a (H759A), HN1-RT(5M)-LBCas12a (H759A)-HB1, HN1-RT(5M)-LBCas12a (H759A)-H1G, HN1-RT(5M)-LBCas12a (H759A)-CHD1, HN1-RT(5M)-H1G-LBCas12a (H759A) and HN1-RT(5M)-CHD1-LBCas12a (H759A).

The precise editing results using chromatin-modulating peptides with constructs of the invention are provided in **Fig. 36**. Compared to the construct without any additional fusions (e.g., RT(5M)-LbCas12a H759A), many of the constructs did not result in an increase in precise editing activity. A slight increase in precise editing activity was observed for HN1-RT(5M)-LbCas12a (H759A)-HB1 with two of the tagRNAs, tagRNA5 and tagRNA6.

20

Example 16. Evaluation of concurrent nicking of the non-template strand of constructs of the invention.

An intermediate during genome editing events including, for example, base editing, Prime editing, and REDRAW, can be a mismatched DNA duplex where one strand of DNA has been edited by the enzyme (desired edit) and the opposite strand contains wild type sequence. Resolution of such a mismatch towards production of the desired edit can be important to ensure that the desired edit becomes permanent in the cell.

Nicking of the DNA strand opposite of the strand containing the edit is thought to promote the process of making the edit permanent by utilizing mismatch repair (MMR) in the cell. In eukaryotes, MMR resolves base mismatches by identifying the DNA strand that contains a nick (which suggests a newly synthesized strand, therefore ‘likely’ to contain a ‘mistake’) and removes that strand and re-synthesize a completely complementary DNA. This way, DNA containing the wildtype sequence is removed, and new DNA is made that is fully complementary to DNA that contains the desired edit. This approach has been used to

improve the editing efficiencies of base editors (Komor et al. *Nature* 533: 420–424 (2016)) and Prime editors (Anzalone et al. *Nature* 576:149–157 (2019)).

We sought to evaluate the same with the constructs and methods of the invention (REDRAW). In REDRAW, the edit is contained in the template strand of DNA (the DNA strand that is hybridized by crRNA). Therefore, we wanted to determine if nicking the non-template strand during the editing process, near the vicinity of the edit, might increase the precise editing efficiency of REDRAW.

Fu et al. previously reported that crRNAs that contain various mismatches in base positions between 12-15 can lead to Cas12a becoming a non-template strand nickase rather than acting as a nuclease (TTTTV PAM is denoted as position -4, -3, -2, -1) (Fu et al, *Nat Microbiol.* 2019 May;4(5):888-897). To determine if such an approach might be effective in increasing the precise editing achieved with the methods and constructs of the invention, a crRNA (in contrast to the extended guide nucleic acids, tagRNAs) comprising various mismatches was prepared and utilized with the constructs of the invention to edit a target DNA. The results are shown in **Table 18**.

Table 18. Precise editing efficiencies (%) with three different tagRNAs. In this experiment, RT(5M)-LbCas12a H759A fusion was used with tagRNA15, tagRNA16, or tagRNA17. The complementarity/mismatches of the spacers in the crRNAs used in this example are described below.

	tagRNA15	tagRNA16	tagRNA17
No nicking crRNA	0.69	0.00	0.51
crRNA; full complementarity	0.00	0.00	0.00
crRNA; mismatch at P12	0.00	0.00	4.62
crRNA; mismatch at P13	0.00	0.00	0.00
crRNA; mismatch at P14	0.00	0.00	0.00
crRNA; mismatch at P15	0.00	0.41	0.36
crRNA; mismatch at P12, 13	3.77	4.42	4.46
crRNA; mismatch at P13, 14	0.15	0.28	1.08
crRNA; mismatch at P12, 13, 14	0.68	0.66	0.98
crRNA; mismatch at P13, 14, 15	0.00	0.00	0.74

Compared to the treatment where no nicking crRNA was used, the crRNAs that contain single, double, or triple mismatches at positions 12-15 led to an increase in editing efficiency. Taken together, concurrent expression of crRNA (in addition to a tagRNA) that

contains appropriate mismatches may be used to induce a nick on the non-template strand and thereby increase the precise editing efficiency of the methods of the invention.

The foregoing is illustrative of the present invention and is not to be construed as limiting thereof. The invention is defined by the following claims, with equivalents of the

5 claims to be included therein.

THAT WHICH IS CLAIMED IS:

1. A method of modifying a target nucleic acid, the method comprising:
contacting the target nucleic acid with
 - 5 (a) a Type V CRISPR-Cas effector protein or a Type II CRISPR-Cas effector protein;
 - (b) a reverse transcriptase, and
 - (c) an extended guide nucleic acid (e.g., extended Type II or Type V CRISPR RNA, extended Type II or Type V CRISPR DNA, extended Type II or Type V crRNA, extended Type II or Type V crDNA), thereby modifying the target nucleic acid.
- 10 2. The method of claim 1, wherein the Type V CRISPR-Cas effector protein or the Type II CRISPR-Cas effector protein, the reverse transcriptase, and the extended guide nucleic acid form a complex or are comprised in a complex.
3. The method of claim 1 or claim 2, wherein the extended guide nucleic acid comprises:
 - (i) a Type V CRISPR nucleic acid or Type II CRISPR nucleic acid (Type II or Type V CRISPR RNA, Type II or Type V CRISPR DNA, Type II or Type V crRNA, Type II or Type V crDNA) and/or a Type V CRISPR nucleic acid or Type II CRISPR nucleic acid and a tracr nucleic acid (e.g., Type II or Type V tracrRNA, Type II or Type V tracrDNA); and
 - 20 (ii) an extended portion comprising a primer binding site and a reverse transcriptase template (RT template) (RTT).
- 25 4. The method of claim 3, wherein the extended portion is fused to either the 5' end or 3' end of the CRISPR nucleic acid (e.g., 5' to 3': repeat-spacer-extended portion, or extended portion-repeat-spacer) and/or to the 5' or 3' end of the tracr nucleic acid.
5. The method of claim 3 or claim 4, wherein the extended portion of the extended guide nucleic acid comprises, 5' to 3', an RT template and a primer binding site.
- 30 6. The method of any one of claims 1-5, wherein the extended guide nucleic acid further comprises a structured RNA motif.

7. The method of claim 6, wherein the structured RNA motif is located at the 3' end of the extended guide nucleic acid.
8. The method of claim 6 or claim 7, wherein the structured RNA motif is AsCpf1BB (SEQ ID NO:189), BoxB (SEQ ID NO:190), pseudoknot (decoy) (SEQ ID NO:95, SEQ ID NO:203), pseudoknot (tEvoPreQ1) (SEQ ID NO:191), fmpknot (SEQ ID NO:192), mpknot (SEQ ID NO:193), MS2 (SEQ ID NO:194), PP7 (SEQ ID NO:195), SLBP (SEQ ID NO:196), TAR (SEQ ID NO:197), and/or ThermoPh (SEQ ID NO:198).
9. The method of any one of claims 6-8, wherein the structured RNA motif is a pseudoknot.
10. The method of claim 9, wherein the pseudoknot is a tEvoPreQ1 pseudoknot comprising the nucleic acid sequence of SEQ ID NO:158 or an EvoPreQ1 Pseudoknot comprising the nucleic acid sequence of SEQ ID NO:191.
11. The method of claim 9, wherein the pseudoknot comprises the nucleic acid sequenced of SEQ ID NO:95 or SEQ ID NO:203.
12. The method of any one of claims 5-11, wherein the target nucleic acid is double stranded and comprises a first strand and a second strand and the primer binding site binds to the second strand (non-target, top strand) of the target nucleic acid.
13. The method of any one of claims 5-11, wherein the target nucleic acid is double stranded and comprises a first strand and a second strand and the primer binding site binds to the first strand (e.g., binds to the target strand, same strand to which the CRISPR-Cas effector protein is recruited, bottom strand) of the target nucleic acid.
14. The method of any one of claims 5-11, wherein the target nucleic acid is double stranded and comprises a first strand and a second strand and the primer binding site binds to the second strand (non-target strand, opposite strand from that to which the CRISPR-Cas effector protein is recruited) of the target nucleic acid.

15. The method of any one of claims 3-14, wherein the primer binding site (PBS) is length of about one nucleotide to about 100 nucleotides, optionally wherein the PBS is about 4 nucleotides to about 80 nucleotides, about 30 nucleotides to about 80 nucleotides, about 40 to about 60 nucleotides in length, or about 8, 16, 24, 32, 40, 48, 56, 64, 72, or 80 nucleotides in length.

16. The method of any one of claims 3-15, wherein the RT template (RTT) is a length of about one to about 100 nucleotides, optionally wherein the RTT is a length of about 35 to about 100, about 35 nucleotides to about 80 nucleotides, optionally about 35 nucleotides to about 75 nucleotides, about 40 nucleotides to about 75 nucleotides, about 45 nucleotides to about 75 nucleotides, about 45 nucleotides to about 60 nucleotides, or a length of about 36, 40, 44, 47, 50, 52, 55, 63, 72 or 74 nucleotides.

17. The method of any one of claims 3 to 16, wherein the RTT is a length of about 35 nucleotides to about 75 nucleotides and the PBS is a length of about 30 nucleotides to about 80 nucleotides, optionally wherein the PBS is a length of about 8, 16, 24, 32, 40, 48, 56, 64, 72, or 80 nucleotides and the RTT is a length of about 36, 40, 44, 47, 50, 52, 55, 63, 72 or 74 nucleotides.

18. The method of any one of claims 3 to 17, wherein the extended portion of the extended guide RNA is linked to the CRISPR nucleic acid and/or the tracrRNA via a linker.

19. The method of claim 18, wherein the linker is 1 to 100 nucleotides in length.

20. The method of any one of claims 3 to 19, wherein when the extended portion is located 5' of the crRNA, the Type V CRISPR-Cas effector protein is modified to reduce (or eliminate) self-processing RNase activity.

21. The method of any of the preceding claims, wherein the Type V CRISPR-Cas effector protein or the Type II CRISPR-Cas effector protein is a fusion protein and/or the reverse transcriptase is a fusion protein, wherein the Type V CRISPR-Cas fusion protein or Type II CRISPR-Cas fusion protein, the reverse transcriptase fusion protein and/or the extended guide nucleic acid is fused to one or more components that recruit the reverse transcriptase to the Type V CRISPR-Cas effector protein or Type II CRISPR-Cas effector protein, optionally

the one or more components recruit via protein-protein interactions, protein-RNA interactions, and/or chemical interactions.

22. The method of any of the preceding claims, wherein the Type V CRISPR-Cas effector protein is a Type V CRISPR-Cas effector fusion protein comprising a Type V CRISPR-Cas effector protein domain fused (linked) to a peptide tag (e.g., an epitope or a multimerized epitope) and the reverse transcriptase is a reverse transcriptase fusion protein comprising a reverse transcriptase domain fused (linked) to an affinity polypeptide that binds to the peptide tag, optionally wherein the target nucleic acid is contacted with two or more reverse transcriptase fusion proteins, or the Type II CRISPR-Cas effector protein is a Type II CRISPR-Cas effector fusion protein comprising a Type II CRISPR-Cas effector protein domain fused (linked) to a peptide tag (e.g., an epitope or a multimerized epitope) and the reverse transcriptase is a reverse transcriptase fusion protein comprising a reverse transcriptase domain fused (linked) to an affinity polypeptide that binds to the peptide tag, optionally wherein the target nucleic acid is contacted with two or more reverse transcriptase fusion protein.

23. The method of claim 22, wherein the peptide tag comprises a GCN4 peptide tag (e.g., Sun-Tag), a c-Myc affinity tag, an HA affinity tag, a His affinity tag, an S affinity tag, a methionine-His affinity tag, an RGD-His affinity tag, a FLAG octapeptide, a strep tag or strep tag II, a V5 tag, and/or a VSV-G epitope.

24. The method of claim 22 or claim 23, wherein the peptide tag comprises two or more copies of the peptide tag.

25

25. The method of any one of claims 22-24, wherein the affinity polypeptide is an antibody, an affibody, an anticalin, a monobody and/or a DARPin.

26. The method of claim 25, wherein the antibody is an scFv antibody.

30

27. The method of any one of the preceding claims, wherein the extended guide nucleic acid is linked to an RNA recruiting motif, and the reverse transcriptase is a reverse transcriptase fusion protein comprising a reverse transcriptase domain fused (linked) to an affinity polypeptide that binds to the RNA recruiting motif, optionally wherein the target

nucleic acid is contacted with two or more reverse transcriptase fusion proteins, optionally wherein the extended guide RNA is linked to two or more RNA recruiting motifs, optionally wherein the two or more RNA recruiting motifs are the same RNA recruiting motif or different RNA recruiting motifs.

5

28. The method of claim 27 wherein the recruiting motif is located on the 3' end of the extended portion of the extended guide nucleic acid or is embedded in the extended portion.

29. The method of claim 27 or claim 28, wherein the RNA recruiting motif and
 10 corresponding affinity polypeptide are a telomerase Ku binding motif (e.g., Ku binding hairpin) and the affinity polypeptide of Ku (e.g., Ku heterodimer); a telomerase Sm7 binding motif and the affinity polypeptide of Sm7; an MS2 phage operator stem-loop and the affinity polypeptide MS2 Coat Protein (MCP), a PP7 phage operator stem-loop and the affinity
 15 polypeptide PP7 Coat Protein (PCP); an SfMu phage Com stem-loop and the affinity polypeptide Com RNA binding protein; a PUF binding site (PBS) and the affinity polypeptide Pumilio/fem-3 mRNA binding factor (PUF); and/or a synthetic RNA-aptamer and the corresponding aptamer ligand.

30. The method of any one of claims 27-29, wherein the RNA recruiting motif and
 20 corresponding affinity polypeptide are an MS2 phage operator stem-loop and the affinity polypeptide MS2 Coat Protein (MCP), and/or a PUF binding site (PBS) and the affinity polypeptide Pumilio/fem-3 mRNA binding factor (PUF).

31. The method of claim 21, wherein the one or more components that recruit chemical
 25 interactions are rapamycin-inducible dimerization of FRB – FKBP; Biotin-streptavidin; SNAP tag; Halo tag; CLIP tag; DmrA-DmrC heterodimer induced by a compound; bifunctional ligand (e.g., fusion of two protein-binding chemicals together, e.g., dihydrofolate reductase (DHFR).

30 32. The method of any one of the preceding claims, further comprising contacting the target nucleic acid with
 (a) a CRISPR-Cas effector protein; and
 (b) a guide nucleic acid, wherein (i) the CRISPR-Cas effector protein nicks or cuts a site on the first strand of the target nucleic acid that is located about 10 to about 125 base

pairs (either 5' or 3') from a site on the second strand that has been nicked by the Type II or Type V CRISPR-Cas effector protein, or (ii) the CRISPR-Cas effector protein nicks or cuts a site on the second strand of the target nucleic acid that is located about 10 to about 125 base pairs (either 5' or 3') from a site on the first strand that has been nicked by the Type II or Type V CRISPR-Cas effector protein, thereby improving mismatch repair, wherein the CRISPR-Cas effector protein is a Type I, Type II, Type III, Type IV, or Type V CRISPR-Cas effector protein.

33. The method of any one of claims 21-32, wherein the Type V CRISPR-Cas fusion protein or Type II CRISPR-Cas fusion protein is fused to a chromatin modulating peptide.

34. The method of any one of claims 21-33, wherein the chromatin modulating peptide is fused to the C-terminus and/or the N-terminus of the Type V CRISPR-Cas fusion protein or Type II CRISPR-Cas fusion protein.

35. The method of any one of claims 21-34, wherein the Type V CRISPR-Cas fusion protein or Type II CRISPR-Cas fusion protein is fused at its N-terminus to a reverse transcriptase and the chromatin modulating peptide is fused to the N-terminus of the reverse transcriptase.

36. The method of any one of claims 33-35, wherein the chromatin modulating peptide is CHD1 (e.g., **SEQ ID NO:199**), H1G (e.g., **SEQ ID NO:200**), HB1 (e.g., **SEQ ID NO:201**), and HN1 (e.g., **SEQ ID NO:202**).

37. The method of any one of the preceding claims, further comprising contacting the target nucleic acid with a Dna2 polypeptide and/or a 5' flap endonuclease (FEN), optionally an FEN1 polypeptide.

38. The method of claim 37, wherein the FEN and/or Dna2 polypeptide is overexpressed (in the presence of the target nucleic acid).

39. The method of claim 37 or claim 38, wherein the FEN is a fusion protein comprising an FEN domain fused to the Type II or Type V CRISPR-Cas effector protein or domain

and/or wherein the Dna2 polypeptide is a fusion protein comprising an Dna2 domain fused to the Type II or Type V CRISPR-Cas effector protein or domain.

40. The method of any one of the preceding claims, wherein the Type V CRISPR-Cas effector protein is a Type V CRISPR-Cas fusion protein comprising a Type V CRISPR-Cas effector protein domain fused (linked) to a peptide tag (e.g., an epitope or a multimerized epitope) and the FEN is an FEN fusion protein comprising an FEN domain fused to an affinity polypeptide that binds to the peptide tag, and/or wherein the Type V CRISPR-Cas effector protein is a Type V CRISPR-Cas fusion protein comprising a Type V CRISPR-Cas effector protein domain fused to a peptide tag and the Dna2 polypeptide is an Dna2 fusion protein comprising an Dna2 domain fused to an affinity polypeptide that binds to the peptide tag, optionally wherein the target nucleic acid is contacted with two or more FEN fusion proteins and/or two or more Dna2 fusion proteins, thereby recruiting the FEN and/or Dna2 to the Type V CRISPR-Cas effector protein domain, and the target nucleic acid.

41. The method of any one of the claims 1-39, wherein the Type II CRISPR-Cas effector protein is a Type II CRISPR-Cas fusion protein comprising a Type II CRISPR-Cas effector protein domain fused (linked) to a peptide tag (e.g., an epitope or a multimerized epitope) and the FEN is an FEN fusion protein comprising an FEN domain fused to an affinity polypeptide that binds to the peptide tag, and/or wherein the Type II CRISPR-Cas effector protein is a Type II CRISPR-Cas fusion protein comprising a Type II CRISPR-Cas effector protein domain fused to a peptide tag and the Dna2 polypeptide is an Dna2 fusion protein comprising an Dna2 domain fused to an affinity polypeptide that binds to the peptide tag, optionally wherein the target nucleic acid is contacted with two or more FEN fusion proteins and/or two or more Dna2 fusion proteins, thereby recruiting the FEN and/or Dna2 to the Type II CRISPR-Cas effector protein domain, and the target nucleic acid.

42. The method of any one of the preceding claims, wherein the Type V CRISPR-Cas effector protein is a Cas12a (Cpf1) domain, Cas12b (C2c1) domain, Cas12c (C2c3) domain, Cas12d (CasY) domain, Cas12e (CasX) domain, Cas12g domain, Cas12h domain, Cas12i domain, C2c4 domain, C2c5 domain, C2c8 domain, C2c9 domain, C2c10 domain, Cas14a domain, Cas14b domain, and/or Cas14c domain, or the Type II CRISPR-Cas effector protein is a Cas9 domain, optionally wherein the Type V CRISPR-Cas effector protein and/or the Type II CRISPR-Cas effector protein comprises one or more mutations as described herein.

43. The method of claim 42, wherein the Type V CRISPR-Cas effector protein is an LbCas12a nickase comprising a R1138A mutation with reference to the amino acid position numbering of **SEQ ID NO:9**, an AsCas12a nickase comprising an R1226A mutation with reference to the amino acid position numbering of **SEQ ID NO:2**, a FnCas12a comprising an R1228A with reference to the amino acid position numbering of **SEQ ID NO:6**, or a PdCas12a nickase comprising an R1241A mutation with reference to the amino acid position numbering of **SEQ ID NO:14**.
44. The method of claim 42, wherein the Type V CRISPR-Cas effector protein is a Type V CRISPR-Cas effector protein from *Acidaminococcus* sp. (AsCas12a) or from Lachnospiraceae bacterium (e.g., LbCas12a) or a modified Type V CRISPR-Cas effector protein therefrom.
45. The method of claim 42, wherein the Type V CRISPR-Cas effector protein is a Type V CRISPR-Cas effector protein from *Butyrivibrio hungatei* (BhCas12) or a modified Type V CRISPR-Cas effector protein therefrom.
46. The method of claim 42, wherein the Type II CRISPR-Cas effector protein is a Cas9 effector protein/domain, optionally from *Streptococcus*.
47. The method of any one of claims 1-40 or 42-46, wherein the Type V CRISPR-Cas effector protein comprises reduced single stranded DNA cleavage activity (ss DNase activity) or is modified (mutated) to reduce (or eliminate) ss DNase activity.
48. The method of any one of claims 1-40 or 42-47, wherein the Type V CRISPR-Cas effector protein is modified (mutated) to reduce (or eliminate) self-processing RNase activity.
49. The method of claim 48, wherein the Type V CRISPR-Cas effector protein is a Cas12a CRISPR-Cas effector protein having a H759A mutation with reference to amino acid position numbering of **SEQ ID NO:1**.

50. The method of claim 49, wherein the Cas12a CRISPR-Cas effector protein having a H759A mutation comprises a sequence having at least 80% sequence identity to the amino acid sequence of **SEQ ID NO:148**.

5 51. The method of claim 50, wherein the Cas12a CRISPR-Cas effector protein having a H759A mutation is a LbCas12a CRISPR-Cas effector protein having at least 90% sequence identity to the amino acid sequence of **SEQ ID NO:148**.

52. The method of any one of the preceding claims, wherein the Type V CRISPR-Cas effector protein is modified to reduce or eliminate nuclease activity (e.g., a mutation in a nuclease active site (e.g., in the RuvC domain), or the Type II CRISPR-Cas effector protein is modified to reduce or eliminate nuclease activity (e.g., a mutation in a nuclease active site (e.g., in the RuvC or HNH domain), to produce a deactivated Type V CRISPR-Cas effector protein or deactivated Type II CRISPR-Cas effector protein (e.g., deadCas (dCas, , dCas12a, dCas9)).

53. The method of claim 52, wherein the deactivated Type V CRISPR-Cas effector protein or the deactivated Type II CRISPR-Cas effector protein comprises nickase activity.

20 54. The method of claim 52 or claim 53, wherein the deactivated Type V CRISPR-Cas effector protein or the deactivated Type II CRISPR-Cas effector protein is fused to a nicking enzyme (e.g., FokI, BfiI, e.g., an engineered FokI or BfiI).

55. The method of any one of the preceding claims, wherein Type V CRISPR-Cas effector protein comprises an amino acid modification to reduce DNA binding affinity of the Type V CRISPR-Cas effector protein.

56. The method of claim 55, wherein the modification is an amino acid substitution.

30 57. The method of claim 55 or claim 56, wherein the modification is an amino acid substitution to an alanine, optionally wherein the alanine replaces a positively charged amino acid residue.

58. The method of claim 57, wherein the amino acid substitution is K167A, K272A, and/or K349A with reference to the amino acid position numbering of **SEQ ID NO:148**.

59. The method of claim 58, wherein Type V CRISPR-Cas effector protein is a Cas12a
5 CRISPR-Cas effector protein comprising an amino acid substitution of K167A, K272A, K349A, K167A+ K272A, K167A+ K349A, K272A+ K349A, or K167A+ K272A + K349A with reference to the amino acid position numbering of **SEQ ID NO:148**, optionally wherein the Type V CRISPR-Cas effector protein is an LbCas12a.

10 60. The method of any one of the preceding claims, wherein reverse transcriptase comprises an amino acid modification.

61. The method of claim 60, wherein amino acid modification is an amino acid substitution.

15

62. The method of claim 61, wherein the amino acid substitution is L139P, D200N, W388R, E607K, T306K, W313F, F155Y, H638G, Q221R, V223M and/or D524N with reference to the amino acid position numbering of **SEQ ID NO:172**.

20 63. The method of claim 61 or claim 62, wherein reverse transcriptase comprises an amino acid substitution of L139P, D200N, W388R, and E607K; L139P, D200N, T306K, W313F, W388R, and E607K; 5M (T355A/Q357M/K358R/A359G/S360A), F155Y, and H638G; 5M (T355A/Q357M/K358R/A359G/S360A), Q221R, and V223M; or 5M T355A/Q357M/K358R/A359G/S360A and D524N with reference to the amino acid position
25 numbering of **SEQ ID NO:172**.

64. The method of any one of the preceding claims, wherein the Type V CRISPR-Cas effector protein is a Type V CRISPR-Cas fusion protein comprising a Type V CRISPR-Cas effector protein domain fused to the reverse transcriptase or the Type II CRISPR-Cas effector
30 protein is a Type II CRISPR-Cas fusion protein comprising a Type II CRISPR-Cas effector protein domain fused to the reverse transcriptase.

65. The method of claim 64, wherein the reverse transcriptase is fused to the C-terminus of the Type V CRISPR-Cas effector polypeptide or the Type II CRISPR-Cas effector polypeptide.

5 66. The method of claim 65, wherein the reverse transcriptase is fused to the N-terminus of the Type V CRISPR-Cas effector polypeptide or the Type II CRISPR-Cas effector polypeptide.

67. The method of any of the preceding claims, wherein the reverse transcriptase is fused
10 to one or more single stranded RNA binding domains (RBDs), thereby improving the thermostability, processivity and template affinity of the reverse transcriptase.

68. The method of claim 67, wherein the one or more single stranded RNA binding domains are fused to the N-terminus of the reverse transcriptase, optionally wherein the
15 reverse transcriptase is further fused at its C-terminus to the N-terminus of the Type II CRISPR-Cas effector protein and/or Type V CRISPR-Cas effector protein.

69. The method of any one of the preceding claims, further comprising contacting the target nucleic acid with a 5'-3' exonuclease.
20

70. The method of claim 69, wherein the 5'-3' exonuclease is fused to a Type V CRISPR-Cas effector protein, optionally to a Type V CRISPR-Cas fusion protein or the 5'-3' exonuclease is fused to a Type II CRISPR-Cas effector protein, optionally to a Type II CRISPR-Cas fusion protein.
25

71. The method of claim 70, wherein the 5'-3' exonuclease is fused to the C-terminus of the Type V CRISPR-Cas effector protein, optionally to the C-terminus of the Type V CRISPR-Cas fusion protein, or is fused to the C-terminus of the Type II CRISPR-Cas effector protein, optionally to the C-terminus of the Type II CRISPR-Cas fusion protein.
30

72. The method of claim 69, wherein the 5'-3' exonuclease is a fusion protein comprising the 5'-3' exonuclease fused to a peptide tag and the Type II or Type V CRISPR-Cas effector protein is a fusion protein comprising a Type II or Type V CRISPR-Cas effector protein domain fused to an affinity polypeptide that is capable of binding to the peptide tag.

73. The method of claim 69, wherein the 5'-3' exonuclease is a fusion protein comprising the 5'-3' exonuclease fused to an affinity polypeptide that is capable of binding to the peptide tag and the Type II or Type V CRISPR-Cas effector protein is a fusion protein comprising a Type II or Type V CRISPR-Cas effector protein domain fused to a peptide tag.

74. The method of claim 69, wherein the 5'-3' exonuclease is a fusion protein comprising the 5'-3' exonuclease fused to an affinity polypeptide that is capable of binding to an RNA recruiting motif and the extended guide nucleic acid is linked to an RNA recruiting motif.

75. The method of any one of claims 69-74, wherein the 5'-3' exonuclease is a RecE exonuclease (RecE), a RecJ exonuclease (RecJ), a T5 exonuclease (T5_Exo), a T7 exonuclease (T7_Exo), a lambda exonuclease (Lambda_exo), a sbcB (exodeoxyribonuclease), or an hExo1 (human exonuclease 1).

76. The method of any one of the preceding claims, further comprising contacting the target nucleic acid with a single stranded DNA binding protein (ssDNA binding protein).

77. The method of claim 76, wherein the ssDNA binding protein is fused to the Type II V CRISPR-Cas effector protein or Type V CRISPR-Cas effector protein.

78. The method of claim 77, wherein the ssDNA binding protein is fused to the C-terminus of the Type II or Type V CRISPR-Cas effector protein.

79. The method of claim 77, wherein the ssDNA binding protein is fused to the N-terminus of the Type II or Type V CRISPR-Cas effector protein.

80. The method of any of claims 76 to 79, wherein the ssDNA binding protein is from a human, a bacterium or a phage.

81. The method of any of claims 76 to 80, wherein the ssDNA binding protein is hRad51 (optionally, hRad51_S208E_A209D), hRad52, BsRecA, EcRecA, T4ssB and/or Brex27.

82. The method of any one of the preceding claims, further comprising reducing double strand breaks by introducing a chemical inhibitor of non-homologous end joining (NHEJ), by introducing a CRISPR guide nucleic acid or an siRNA targeting an NHEJ protein to transiently knock-down expression of the NHEJ protein, or by introducing a polypeptide that prevents NHEJ

83. The method of claim 82, wherein the polypeptide that prevents NHEJ is fused to the reverse transcriptase and/or the CRISPR-Cas effector protein, optionally the polypeptide that prevents NHEJ is fused to the N-terminus of the reverse transcriptase and/or the CRISPR-Cas effector protein.

84. The method of claim 82 or claim 83, wherein the polypeptide that prevents NHEJ is a Gam protein.

85. The method of claim 84, wherein the Gam protein is *Escherichia* phage Mu Gam protein, optionally the Gam protein comprise the amino acid sequence of **SEQ ID NO:147**.

86. The method of any one of the preceding claims, wherein the Type II or Type V CRISPR-Cas effector protein is encoded by a polynucleotide and/or the reverse transcriptase is encoded by a polynucleotide.

87. The method of any one of claims 32 to 86, wherein the CRISPR-Cas effector protein is encoded by a polynucleotide.

88. The method of any one of claims 37-87, wherein the 5' flap endonuclease (FEN) is encoded by a polynucleotide.

89. The method of any one of claims 69-88, wherein the 5'-3' exonuclease is encoded by a polynucleotide.

90. The method of any one of claims 1-89, wherein the target nucleic acid is in a cell, in a cell free system or in a virus.

91. The method of claim 90, wherein the cell is a eukaryotic cell or a prokaryotic cell.

92. The method of claim 90 or claim 91, wherein the cell is from an animal, a plant, a fungus, an archaeon, a bacterium or a virus.

5 93. The method of any one of claims 86 to 92, wherein the polynucleotide encoding the Type II or Type V CRISPR-Cas effector protein and/or the polynucleotide encoding the reverse transcriptase are comprised in the same or separate expression cassettes, optionally when the polynucleotide encoding the Type II or Type V CRISPR-Cas effector protein and the polynucleotide encoding the reverse transcriptase are comprised in the same expression
10 cassette, the polynucleotide encoding the Type II or Type V CRISPR-Cas effector protein and the polynucleotide encoding the reverse transcriptase are operably linked to a single promoter or to two or more separate promoters in any combination.

94. The method of any one of claims 87 to 93, wherein the polynucleotide encoding the
15 CRISPR-Cas effector protein is comprised in an expression cassette, optionally wherein the CRISPR-Cas effector protein is operably linked to a promoter.

95. The method of claim 94, wherein the same or separate expression cassettes are comprised in the same or separate vectors.

20

96. The method of claim 94, wherein the expression cassette is comprised a vector.

97. The method of any one of claims 88 to 96, wherein the polynucleotide encoding the 5' flap endonuclease is comprised in an expression cassette, optionally wherein the expression
25 cassette is comprised in a vector.

98. The method of any one of claims 83 to 91, wherein the polynucleotide encoding the 5'-3' exonuclease is comprised in an expression cassette, optionally wherein the expression cassette is comprised in a vector.

30 99. The method of any one of the preceding claims, wherein the extended guide nucleic acid is comprised in an expression cassette, optionally wherein the expression cassette is comprised in a vector.

100. The method of any one of claims 32 to 99, wherein the guide nucleic acid is comprised in an expression cassette, optionally wherein the expression cassette is comprised in a vector.

5 101. The method of any one of claims 85 to 100, wherein the polynucleotides encoding the Type II or Type V CRISPR-Cas effector proteins, the CRISPR-Cas effector proteins, the reverse transcriptases, the 5' flap endonucleases, the 5'-3' exonucleases and expression cassettes and/or vectors comprising the same are codon optimized for expression in an organism, optionally in a prokaryote or a eukaryote, optionally in an animal, a plant, a
10 fungus, an archaeon, a bacterium or a virus.

102. The method of claim 101, wherein the codon optimization is for expression in a dicot plant.

15 103. The method of claim 101, wherein the codon optimization is for expression in a monocot plant.

104. A method of modifying a target nucleic acid, the method comprising:
contacting the target nucleic acid at a first site with

20 (a)(i) a first CRISPR-Cas effector protein; and
(ii) a first extended guide nucleic acid (e.g., extended CRISPR RNA, extended CRISPR DNA, extended crRNA, extended crDNA); and
(b)(i) a second CRISPR-Cas effector protein,
(ii) a first reverse transcriptase; and
25 (ii) a first guide nucleic acid, thereby modifying the target nucleic acid.

105. The method of claim 104, wherein the first extended guide nucleic acid comprises
(a) a CRISPR nucleic acid (CRISPR RNA, CRISPR DNA, crRNA, crDNA); and
(b) an extended portion comprising a primer binding site and a reverse transcriptase

30 template (RT template).

106. The method of claim 104 or claim 105, wherein the CRISPR nucleic acid of the first extended guide nucleic acid comprises a spacer sequence capable of binding to (having substantial homology to) a first site on the first strand of the target nucleic acid.

107. The method of any one of claims 104-106, wherein the first guide nucleic acid comprises a CRISPR nucleic acid (CRISPR RNA, CRISPR DNA, crRNA, crDNA).

- 5 108. The method of claim 107, wherein the CRISPR nucleic acid of the first guide nucleic acid comprises a spacer sequence that binds to a second site on the first strand of the target nucleic acid that is upstream of the first site on the first strand of the target nucleic acid.

109. The method of any one of claims 104-108, wherein the second CRISPR-Cas effector protein is a CRISPR-Cas fusion protein comprising a CRISPR-Cas effector protein domain fused to the reverse transcriptase.

110. The method of any one of claims 104-108, wherein the second CRISPR-Cas effector protein is a CRISPR-Cas fusion protein comprising a CRISPR-Cas effector protein domain fused to a peptide tag and the first reverse transcriptase is a reverse transcriptase fusion protein comprising a reverse transcriptase domain fused to an affinity polypeptide capable of binding the peptide tag.

111. The method of any one of claims 104-108, wherein the first guide nucleic acid is linked to an RNA recruiting motif and the first reverse transcriptase is a reverse transcriptase fusion protein comprising a reverse transcriptase domain fused to an affinity polypeptide capable of binding the RNA recruiting motif.

112. The method of any one of claims 104-111, further comprising contacting the target nucleic acid with

- (a) a third CRISPR-Cas effector protein; and
- (b) a second guide nucleic acid,

- wherein the third CRISPR-Cas effector protein nicks a site on the first strand of the target nucleic acid that is located about 10 to about 125 base pairs (either 5' or 3') from the second site on the second strand that has been nicked by the second CRISPR-Cas effector protein, thereby improving mismatch repair.

113. The method of any one of claims 104-112, further comprising contacting the target nucleic acid with a 5'-3' exonuclease.

114. The method of claim 113, wherein the 5'-3' exonuclease is fused to the first CRISPR-Cas effector protein.

5 115. The method of claim 114, wherein the 5'-3' exonuclease is fused to the C-terminus of the first CRISPR-Cas effector protein.

116. The method of claim 113, wherein the 5'-3' exonuclease is a fusion protein comprising the 5'-3' exonuclease fused to a peptide tag and the first CRISPR-Cas effector protein is a
10 fusion protein comprising a CRISPR-Cas effector protein domain fused to an affinity polypeptide that is capable of binding to the peptide tag.

117. The method of claim 113, wherein the 5'-3' exonuclease is a fusion protein comprising the 5'-3' exonuclease fused to an affinity polypeptide that is capable of binding to the peptide
15 tag and the first CRISPR-Cas effector protein is a fusion protein comprising a CRISPR-Cas effector protein domain fused to a peptide tag.

118. The method of claim 113, wherein the 5'-3' exonuclease is a fusion protein comprising the 5'-3' exonuclease fused to an affinity polypeptide that is capable of binding to an RNA
20 recruiting motif and extended guide nucleic acid is linked to an RNA recruiting motif.

119. The method of any one of claims 113-118, wherein the 5'-3' exonuclease is a RecE exonuclease (RecE), a RecJ exonuclease (RecJ), a T5 exonuclease (T5_Exo), a T7
exonuclease (T7_Exo), a lambda exonuclease (Lambda_exo), a sbcB
25 (exodeoxyribonuclease), or an hExo1 (human exonuclease 1).

120. The method of any one of claims 104-119, further comprising contacting the target nucleic acid with

- 30 (a) a fourth CRISPR-Cas effector protein;
(b) a second reverse transcriptase, and
(c) a second extended guide nucleic acid (e.g., extended CRISPR RNA, extended CRISPR DNA, extended crRNA, extended crDNA),

wherein the second extended guide nucleic acid targets (spacer is substantially complementary to/binds to) a site on the first strand of the target nucleic acid, thereby modifying the target nucleic acid.

- 5 121. The method of any one of claims 104-120, wherein the first CRISPR-Cas effector protein, the second CRISPR-Cas effector protein, the third CRISPR-Cas effector protein and/or the fourth CRISPR-Cas effector protein are a Type I, Type II, Type III, Type IV, or Type V CRISPR-Cas effector protein, in any combination .
- 10 122. The method of any one of claims 1 to 103, further comprising contacting the target nucleic acid with
- (a) a second CRISPR-Cas effector protein, optionally a Type II or Type V CRISPR-Cas effector protein;
- (b) a second reverse transcriptase, and
- 15 (c) a second extended guide nucleic acid (e.g., extended CRISPR RNA, extended CRISPR DNA, extended crRNA, extended crDNA),
- wherein the second extended guide nucleic acid targets (spacer is substantially complementary to/binds to) a site on the first strand (target strand) or the second strand (non-target strand) of the target nucleic acid, thereby modifying the target nucleic acid.
- 20 123. The method of claim 104-122, wherein the first and/or second extended guide nucleic acid further comprises a pseudoknot.
124. The method of claim 123, wherein the pseudoknot is located at the 3' end of the
- 25 extended guide nucleic acid.
125. The method of claim 123 or claim 124, wherein the pseudoknot is a tEvoPreQ1 Pseudoknot comprising the nucleic acid sequence of **SEQ ID NO:158** or an EvoPreQ1 Pseudoknot comprising the nucleic acid sequence of **SEQ ID NO:191**.
- 30 126. The method of claim 123 or claim 124, wherein the pseudoknot comprises the nucleic acid sequenced **of SEQ ID NO:95** or **SEQ ID NO:203**.

127. The method of any one of claims 104-126, further comprising contacting the target nucleic acid with a single stranded DNA binding protein (ssDNA binding protein).

128. The method of claim 127, wherein the ssDNA binding protein is fused to the first,
5 second, third and/or fourth CRISPR-Cas effector protein.

129. The method of claim 127 or claim 128, wherein the ssDNA binding protein is fused to the C-terminus of the first, second, third and/or fourth CRISPR-Cas effector protein.

10 130. The method of claim 127 or claim 128, wherein the ssDNA binding protein is fused to the N-terminus of the first, second, third and/or fourth CRISPR-Cas effector protein.

131. The method of any of claims 127-130, wherein the ssDNA binding protein is from a human, a bacterium or a phage.

15

132. The method of any of claims 127-131, wherein the ssDNA binding protein is hRad51 (optionally, hRad51_S208E_A209D), hRad52, BsRecA, EcRecA, T4ssB and/or Brex27.

133. The method of any one of claims 104-132, further comprising reducing double strand
20 breaks by introducing a chemical inhibitor of non-homologous end joining (NHEJ), by introducing a CRISPR guide nucleic acid or an siRNA targeting an NHEJ protein to transiently knock-down expression of the NHEJ protein, or by introducing a polypeptide that prevents NHEJ.

25 134. The method of claim 133, wherein the polypeptide that prevents NHEJ is fused to the first and/or second reverse transcriptase and/or the first, second, third and/or fourth CRISPR-Cas effector protein, optionally the polypeptide that prevents NHEJ is fused to the N-terminus of the first and/or second reverse transcriptase and/or the first, second, third and/or fourth CRISPR-Cas effector protein.

30

135. The method of claim 133 or claim 134, wherein the polypeptide that prevents NHEJ is a Gam protein.

136. The method of claim 135, wherein the Gam protein is *Escherichia* phage Mu Gam protein, optionally the Gam protein comprise the amino acid sequence of **SEQ ID NO:147**.

137. An extended guide nucleic acid comprising (i) a Type V CRISPR nucleic acid or Type II CRISPR nucleic acid (Type II or Type V CRISPR RNA, Type II or Type V CRISPR DNA, Type II or Type V crRNA, Type II or Type V crDNA) and/or a Type V CRISPR nucleic acid or Type II CRISPR nucleic acid and a tracr nucleic acid (e.g., Type II or Type V tracrRNA, Type II or Type V tracrDNA); and (ii) an extended portion comprising a primer binding site and a reverse transcriptase template (RT template) (RTT).

10

138. The extended guide nucleic acid of claim 137, wherein the extended guide nucleic acid further comprises a pseudoknot.

139. The extended guide nucleic acid of claim 138, wherein the pseudoknot is located at the 3' end of the extended guide nucleic acid.

15

140. The extended guide nucleic acid of claim 138 or claim 139, wherein the pseudoknot is a tEvoPreQ1 Pseudoknot comprising the nucleic acid sequence of **SEQ ID NO:158** or an EvoPreQ1 Pseudoknot comprising the nucleic acid sequence of **SEQ ID NO:191**.

20

141. The extended guide nucleic acid of claim 138 or claim 139, wherein the pseudoknot comprises the nucleic acid sequenced of **SEQ ID NO:95** or **SEQ ID NO:203**.

142. The extended guide nucleic acid of any one of claims 137-141, comprised in an expression cassette, optionally wherein the expression cassette is comprised in a vector.

25

143. A complex comprising:

- (a) a Type V CRISPR-Cas effector protein or a Type II CRISPR-Cas effector protein;
- (b) a reverse transcriptase, and

30

(c) an extended guide nucleic acid (e.g., extended CRISPR RNA, extended CRISPR DNA, extended crRNA, extended crDNA).

144. The complex of claim 143, wherein the extended guide nucleic acid further comprises a pseudoknot.

145. The complex of claim 144, wherein the pseudoknot is located at the 3' end of the extended guide nucleic acid.

- 5 146. The complex of claim 144 or claim 145, wherein the pseudoknot is a tEvoPreQ1 Pseudoknot comprising the nucleic acid sequence of **SEQ ID NO:158** or an EvoPreQ1 Pseudoknot comprising the nucleic acid sequence of **SEQ ID NO:191**.

147. The complex of claim 144 or claim 145, wherein the pseudoknot comprises the
10 nucleic acid sequence of **SEQ ID NO:95** or **SEQ ID NO:203**.

148. The complex of claim 143-147, wherein the Type II or Type V CRISPR-Cas effector protein is a fusion protein comprising a Type II or Type V CRISPR-Cas effector protein domain fused to a peptide tag.

15

149. The complex of claim 143-147, wherein the Type II or Type V CRISPR-Cas effector protein is a fusion protein comprising a Type V CRISPR-Cas effector protein domain fused to an affinity polypeptide that is capable of binding a peptide tag.

- 20 150. The complex of claim 143-147, wherein the Type II or Type V CRISPR-Cas effector protein is a fusion protein comprising a Type II or Type V CRISPR-Cas effector protein domain fused to an affinity polypeptide that is capable of binding an RNA recruiting motif.

151. The complex of any one of claims 143-150, wherein the reverse transcriptase is a
25 fusion protein comprising reverse transcriptase domain fused to a peptide tag.

152. The complex of any one of claims 143-150, wherein the reverse transcriptase is a fusion protein comprising reverse transcriptase domain fused to an affinity polypeptide that is capable of binding a peptide tag.

30

153. The complex of any one of claims 143-150, wherein the reverse transcriptase is a fusion protein comprising reverse transcriptase domain fused to an affinity polypeptide that is capable of binding an RNA recruiting polypeptide.

154. The complex of any one of claims 143-153, comprised in an expression cassette, optionally wherein the expression cassette is comprised in a vector.

155. An expression cassette codon optimized for expression in an organism, comprising 5'
5 to 3': (a) polynucleotide encoding a promoter sequence ,
(b) a polynucleotide encoding a Type V CRISPR-Cas nuclease (e.g., Cpf1 (Cas12a), dCas12a and the like) or a Type II CRISPR-Cas nuclease (e.g., Cas9, dCas9 and the like) that is codon optimized for expression in the organism;
(c) a linker sequence; and
10 (d) a polynucleotide encoding a reverse transcriptase that is codon-optimized for expression in the organism.

156. The expression cassette of claim 155, wherein the organism is an animal, a plant, a fungus, an archaeon, a bacterium or a virus.

157. The expression cassette of claim 156, wherein the organism is a plant, the polynucleotide encoding a Type V CRISPR-Cas nuclease is codon optimized for expression in a plant, and the promoter sequence is a plant specific promoter sequence (e.g., ZmUbi1, MtUbi2, RNA polymerase II (Pol II)).

158. The expression cassette of any one of claims 155-157, wherein the reverse transcriptase is fused to one or more ssRNA binding domains (RBDs).

159. The method of claim 158, wherein the one or more single stranded RNA binding
25 domains are fused to the N-terminus of the reverse transcriptase, optionally wherein the reverse transcriptase is further fused at its C-terminus to the N-terminus of the Type II CRISPR-Cas effector protein and/or Type V CRISPR-Cas effector protein.

160. The expression cassette of any one of claims 155 to 159, wherein the linker is an
30 amino acid or peptide linker.

161. The expression cassette of claim 160 wherein the peptide linker is about 2 to about 100 amino acids (residues) in length.

162. The expression cassette of claim 160 or claim 161, wherein the peptide linker is a GS linker.

163. An expression cassette codon optimized for expression in a plant, comprising:

- 5 (a) a polynucleotide encoding a plant specific promoter sequence (e.g., ZmUbi1, MtUb2), and
- (b) an extended guide nucleic acid, wherein the extended guide nucleic acid comprises an extended portion comprising at its 3' end a primer binding site and an edit to be incorporated into the target nucleic acid (e.g., reverse transcriptase template), optionally
- 10 wherein the extended guide nucleic acid is comprised in an expression cassette, optionally wherein the extended guide nucleic acid is operably linked to a Pol II promoter.

164. The expression cassette of claim 163, wherein the extended guide nucleic acid further comprises a pseudoknot.

15

165. The expression cassette of claim 164, wherein the pseudoknot is located at the 3' end of the extended guide nucleic acid.

166. The expression cassette of claim 164 or claim 165, wherein the pseudoknot is a tEvoPreQ1 Pseudoknot comprising the nucleic acid sequence of **SEQ ID NO:158** or an EvoPreQ1 Pseudoknot comprising the nucleic acid sequence of **SEQ ID NO:191**.

20

167. The expression cassette of claim 164 or claim 165, wherein the pseudoknot comprises the nucleic acid sequence of **SEQ ID NO:95** or **SEQ ID NO:203**.

25

168. The expression cassette of any one of claims 157-167, wherein the plant specific promoter is associated with an intron or is a promoter region comprising an intron (e.g., ZmUbi1 comprising an intron; MtUb2 comprising an intron).

30 169. The expression cassette of any one of claims 156-168, wherein the codon optimization is for expression in a dicot plant.

170. The expression cassette of any one of claims 156-168, wherein the codon optimization is for expression in a monocot plant.

171. A method of modifying a target nucleic acid in a plant or plant cell, comprising introducing the expression cassette of any one of claims 155-170 into the plant or plant cell, thereby modifying the target nucleic acid in the plant or plant cell.

5

172. The method of claim 171, further comprising regenerating the plant cell comprising the modified target nucleic acid to produce a plant comprising the modified target nucleic acid.

10 173. The method of any one of claims 1 to 132, 171, or 172, wherein the contacting is carried out at a temperature of about 20°C to 42°C.