

FLUOROINDOLE DERIVATIVES AS MUSCARINIC M1 RECEPTOR POSITIVE ALLOSTERIC MODULATORS

Field of Invention

5 The present invention relates to compounds of formula (I), or their isotopic forms, stereoisomers, or pharmaceutically acceptable salts as muscarinic M1 receptor positive allosteric modulators (M1 PAMs). The present invention also describes method of making such compounds, pharmaceutical compositions comprising such compounds and their use.

10

Background of the Invention

 Muscarinic acetylcholine receptors (mAChRs) which belong to the class A family of G protein-coupled receptors (GPCRs), are widely expressed throughout the body. Five subtypes termed M1 through M5 that respond to the endogenous neurotransmitter acetylcholine (ACh) has been identified till date. They play key role in regulating the activity of many important functions of the central and peripheral nervous system including cognitive function. M1, M3 and M5 couple to Gq, whereas M2 and M4 couple via Gi/o to downstream signaling pathways and associated effector systems (*Critical Reviews in Neurobiology*, 1996, 10, 69-99; *Pharmacology & Therapeutics*, 2008, 117, 232-243). M2 and M3 are highly expressed in the periphery and are known to be involved in gastrointestinal (GI) motility and parasympathetic responses such as salivation (*Life Sciences*, 1993, 52, 441-448). The M1 muscarinic receptor is predominantly expressed in the brain regions such as cortex, hippocampus and amygdala which involved in cognition, and therefore selective activation of the M1 receptor would be expected to boost cognitive performance (*Annals of Neurology*, 2003, 54, 144 - 146).

 Xanomeline, a muscarinic acetylcholine receptor agonist with reasonable selectivity for the M1 and M4 subtypes, produced significant effects on cognition in a clinical Alzheimer's disease (AD) trial (*Alzheimer Disease and Associated Disorders*, 1998, 12(4), 304-312) although gastrointestinal side effects led to a high dropout rate in clinical trials. There is a high degree of conservation between muscarinic receptor

30

subtypes at their orthosteric acetylcholine ligand binding sites which makes it difficult to identify a M1 selective agonist.

To circumvent this issue of selectivity and safety, an alternative approach consists of developing M1 PAMs that act at the less conserved allosteric binding site.

5 Merck reported the development of M1 PAM, PQCA (1-([4-cyano-4-(pyridine-2-yl) piperidin-1-yl] methyl)-4-oxo-4H-quinolizine-3-carboxylic acid). This compound is highly selective for M1 over the other muscarinic receptor subtypes and found to be efficacious in several preclinical models of cognition (*Psychopharmacology*, 2013, 225(1), 21-30) with no gastrointestinal side effects at doses equal to or less than a
10 fivefold margin from the minimum effective dose required to improve cognition. In preclinical studies it was demonstrated that M1 activation increases neurotransmitter acetylcholine concentration in brain. Moreover, the M1 activation has potential as disease-modifying therapy for AD by both shifting the APP processing towards the non-amyloidogenic α -secretase pathway and by decreasing the *tau* hyper-
15 phosphorylation. Positive allosteric modulators at M1 receptor have demonstrated to increase the generation of sAPP α *in-vitro* (*The Journal of Neuroscience*, 2009, 29, 14271-14286). Therefore, M1 PAMs provide an approach to target both symptomatic and disease-modifying treatment of cognitive deficits in AD and schizophrenia.

20 PCT patent application publications, WO2015049574A1, WO2015044072A1, WO2015028483, WO2007067489, and WO2011149801 have disclosed some M1 PAM compounds. The PCT patent application, WO2001058869 and US patent US4616009 discloses some of indole compounds useful in medicaments. While several M1 PAMs have been disclosed in the literature till date, no drug acting as M1
25 PAM is launched in the market.


For drugs with an intended action in the central nervous system (CNS), the compound should cross the blood brain barrier or in other words the compounds should possess brain penetration properties. It is a commonly accepted hypothesis that unbound or free drug is available for interaction with pharmacological and
30 toxicological targets in the brain. This hypothesis is referred to as the free drug hypothesis in pharmacokinetics (*Current Opinion in Drug Discovery & Development*, 2005, 8, 505-512; *Expert Opinion on Drug Discovery*, 2007, 2, 51-64;

Pharmaceutical Research, 2008, 25, 1737–1750; *Current Drug Metabolism*, 2008, 9, 46–59; *Journal of Pharmaceutical Sciences*, 2010, 99, 1107–1122).

Although the prior arts disclose M1 PAM compounds that are useful in the treatment of CNS related diseases, there exist an issue of poor brain penetration and free fraction availability. Therefore, there is an un-met need and scope to discover and develop new M1 PAMs with good brain penetration and adequate free fraction. Such compounds may have efficacy at much lower doses thereby increasing the margin of safety over the efficacy dose. The M1 PAM compounds of the instant invention solve the issue of brain penetration as well as the free fraction availability in the brain thereby very effective in the treatment of CNS related disorders.


Summary of the Invention

In first aspect, the present invention relates to muscarinic M1 receptor PAMs of compound of formula (I),




wherein:

R¹ is



R² is



wherein * represents point of attachment;

R³ is fluorine or hydrogen;

5 R⁴ is halogen, -S-CH₃ or hydrogen;

R⁵ is -CH₃, -CH₂CH₂F or hydrogen;

R⁶ is halogen, -O-CH₃ or hydrogen;

a is 1 or 2; and

b is 1 or 2;

10 or an isotopic form, a stereoisomer or a pharmaceutically acceptable salt thereof.

In another aspect, the present invention relates to the processes for preparing the compound of formula (I), or a stereoisomer and a pharmaceutically acceptable salt thereof.

15 In yet another aspect, the present invention relates to pharmaceutical composition containing a therapeutically effective amount of at least one compound of formula (I), or a stereoisomer and a pharmaceutically acceptable salt thereof and pharmaceutically acceptable excipients or carriers.

In yet another aspect, the present invention relates to compound of formula (I), or a stereoisomer and a pharmaceutically acceptable salt thereof, for use as M1 PAM.

5 In yet another aspect, the present invention relates to compound of formula (I), or a stereoisomer and a pharmaceutically acceptable salt thereof, for use in the treatment of various disorders selected from AD, schizophrenia, cognitive disorders, pain or sleep disorders.

In another aspect, the present invention relates to a method for the treatment of disorders related to muscarinic M1 receptor, comprising administering to a patient
10 in need thereof, a therapeutically effective amount of a compound of formula (I), or a stereoisomer and a pharmaceutically acceptable salt thereof.

In yet another aspect, the present invention relates to use of the compound of formula (I), or stereoisomers and pharmaceutically acceptable salts thereof, for the manufacture of a medicament for the treatment of disorders related to muscarinic M1
15 receptor.

In yet another aspect, the present invention relates to compound of formula (I) for use in positive allosteric modulation of muscarinic M1 receptor.

Brief Description of Drawings

20 Figure 1: Effect of test compound on contextual fear conditioning task
Figure 2: Effect of test compound on modulation of cerebral blood flow in the frontal cortex

Detailed Description of the Invention

25 Unless otherwise stated, the following terms used in the specification and claims have the meanings given below:

The term "halogen" means fluorine, chlorine, bromine or iodine.

The phrase, "therapeutically effective amount" is defined as an amount of a compound of the present invention that (i) treats the particular disease, condition or
30 disorder (ii) eliminates one or more symptoms of the particular disease, condition or disorder (iii) delays the onset of one or more symptoms of the particular disease, condition or disorder described herein.

The term, "isotopic form" as used herein refers to the compound of formula (I) wherein one or more atoms of compound of formula (I) are substituted by their respective isotopes. For example, isotopes of hydrogen include ^2H (deuterium) and ^3H (tritium).

5 The term, "stereoisomers" as used herein refers to isomers of compound of formula (I) that differ in the arrangement of their atoms in space. Compounds disclosed herein may exist as single stereoisomers, racemates and/or mixtures of enantiomers and/or diastereomers. All such single stereoisomers, racemates and mixtures thereof are intended to be within the scope of the present invention.

10 The term, "pharmaceutically acceptable salt" as used herein refers to salts of the active compound i.e. the compound of formula I, and are prepared by reaction with the appropriate acid or acid derivative, depending on the particular substituents found on the compounds described herein.



WO2015044072A1 patent application discloses indole derivatives as M1
15 PAM compounds. Based on the available *in vitro* data disclosed in the patent, three of the most potent compounds (Example number 30, 76 and 77) were synthesized in our laboratories and tested for its pharmacokinetic and brain penetration properties in Wistar rats. All the three compounds were found to have poor brain penetration. This makes these compounds less ideal for the treatment of CNS disorders. The M1 PAM
20 compounds of the instant invention possess brain penetration and/or free fraction available in the brain which will make them useful compounds to further develop for the treatment of CNS disorders.

Embodiments

25

In one embodiment, the present invention relates to the compound of formula (I), wherein:


30 R^1 is



R^2 is other than $(R^6)_b$;
 wherein * represents point of attachment; R^6 and b are as defined in the first aspect;
 or an isotopic form, a stereoisomer or a pharmaceutically acceptable salt thereof.

- 5 In another embodiment, the present invention relates to the compound of formula (I), wherein:


R^1 is



- wherein * represents point of attachment; or an isotopic form, a stereoisomer or a
 10 pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to the compound of formula (I), wherein:


R^2 is



- 15 wherein * represents point of attachment; R^4 , R^5 and a are as defined in the first aspect; or an isotopic form, a stereoisomer or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to the compound of formula (I), wherein:


R² is



wherein * represents point of attachment; R⁶ and b are as defined in the first aspect; or an isotopic form, a stereoisomer or a pharmaceutically acceptable salt thereof.

5 In another embodiment, the present invention relates to the compound of formula (I), wherein:


R² is



10 wherein * represents point of attachment; R⁴, R⁵ and a are as defined in the first aspect; or an isotopic form, a stereoisomer or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to the compound of formula (I), wherein:

R² is




15

wherein * represents point of attachment; R⁴ and a are as defined in the first aspect; or an isotopic form, a stereoisomer or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to the compound of formula (I), wherein:

20 R² is




wherein * represents point of attachment; R^5 is as defined in the first aspect; or an isotopic form, a stereoisomer or a pharmaceutically acceptable salt thereof.


In another embodiment, the present invention relates to the compound of formula (I), wherein: R^3 is fluorine; or an isotopic form, a stereoisomer or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to the compound of formula (I), wherein:

R^1 is




R^2 is




wherein * represents point of attachment; R^4 and a are as defined in the first aspect; or an isotopic form, a stereoisomer or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to the compound of formula (I), wherein:

R^1 is




R^2 is




wherein * represents point of attachment; R^5 is as defined in the first aspect; or an isotopic form, a stereoisomer or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to the compound of formula (I), wherein:

R¹ is




5 R² is




wherein * represents point of attachment; R⁴ and a are as defined in the first aspect; or an isotopic form, a stereoisomer or a pharmaceutically acceptable salt thereof.

10 In another embodiment, the present invention relates to the compound of formula (I), wherein:

R¹ is




R² is



15 wherein * represents point of attachment; R⁴ is fluorine; a is as defined in the first aspect; or an isotopic form, a stereoisomer or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to compound of formula (I), wherein:


20 R¹ is



wherein the compound is racemic mixture.

In another embodiment, the present invention relates to compound of formula (I), wherein:

5 R^1 is



wherein the configuration of chiral centers at C3 and C4 atoms are (3R,4R), (3S,4S), (4R,3S) or (4S,3R).

10 In yet another embodiment the representative compounds of the present invention includes but not limited to,

N-[(1S,2S)-2-Hydroxycyclohexyl]-1-(1-methyl-1H-pyrazol-4-ylmethyl)-4-fluoro-1H-indole-3-carboxamide;

N-[(1S,2S)-2-Hydroxycyclohexyl]-1-(2-Chloropyridin-4-ylmethyl)-4-fluoro-1H-indole-3-carboxamide;

15 N-[(1R,2R)-2-Hydroxycyclohexyl]-1-(1-methyl-1H-pyrazol-4-ylmethyl)-4-fluoro-1H-indole-3-carboxamide;

N-[(1S,2S)-2-Hydroxycyclohexyl]-1-(1-methyl-1H-indazol-3-ylmethyl)-4,7-difluoro-1H-indole-3-carboxamide;

20 N-[(1S,2S)-2-Hydroxycyclohexyl]-1-(1-methyl-1H-pyrazol-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxamide;

N-[(1S,2S)-2-Hydroxycyclohexyl]-1-(2-Chloropyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxamide;

N-[(1S,2S)-2-Hydroxycyclohexyl]-1-(2-fluoropyridin-4-ylmethyl)-4-fluoro-1H-indole-3-carboxamide;

25 N-[(1S,2S)-2-Hydroxycyclohexyl]-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxamide;

- N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(3-fluoropyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;
- N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(5-fluoropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide;
- 5 N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2,5-difluoropyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;
- N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2,5-difluoropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide;
- N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2,3-difluoropyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;
- 10 N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(1-pyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;
- N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(1*H*-pyrazol-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;
- 15 N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2-bromothiazol-5-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;
- N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(1-ethyl-5-methyl-1*H*-pyrazol-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;
- N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(benzothiazol-6-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;
- 20 N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(1-(2-fluoroethyl)-1*H*-pyrazol-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;
- N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2-methylsulfanyl-pyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide;
- 25 N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2-methylsulfanyl-pyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide (Isomer-II);
- trans*-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide (Isomer-I);
- 30 *trans*-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide (Isomer-II);

- cis*-N-(3-Fluoropiperidin-4-yl)-1-(5-fluoropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(2-chloropyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide (Isomer-I);
- 5 *cis*-N-(3-Fluoropiperidin-4-yl)-1-(2-chloropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide (Racemate);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(2-chloropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(4-fluorobenzyl)-4-fluoro-1*H*-indole-3-carboxamide
- 10 (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(4-fluorobenzyl)-4,7-difluoro-1*H*-indole-3-carboxamide (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(4-fluorobenzyl)-4,7-difluoro-1*H*-indole-3-carboxamide (Isomer-II);
- 15 *cis*-N-(3-Fluoropiperidin-4-yl)-1-(2-fluorobenzyl)-4-fluoro-1*H*-indole-3-carboxamide (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(4-chlorobenzyl)-4-fluoro-1*H*-indole-3-carboxamide (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-4-fluoro-1-(3-methoxybenzyl)-1*H*-indole-3-
- 20 carboxamide (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-4-fluoro-1-(3-methoxybenzyl)-1*H*-indole-3-carboxamide (Isomer-II);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(4-methoxybenzyl)-4-fluoro-1*H*-indole-3-carboxamide (Isomer-II);
- 25 *cis*-N-(3-Fluoropiperidin-4-yl)-1-(4-methoxybenzyl)-4-fluoro-1*H*-indole-3-carboxamide (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(4-methoxybenzyl)-4,7-difluoro-1*H*-indole-3-carboxamide (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(4-methoxybenzyl)-4,7-difluoro-1*H*-indole-3-
- 30 carboxamide (Isomer-II);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(3,4-difluorobenzyl)-4-fluoro-1*H*-indole-3-carboxamide (Isomer-I);

- cis*-N-(3-Fluoropiperidin-4-yl)-1-(1-methyl-1*H*-pyrazol-4-yl-methyl)-4-fluoro-1*H*-indole-3-carboxamide (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(1-methyl-1*H*-pyrazol-4-yl-methyl)-4,7-difluoro-1*H*-indole-3-carboxamide (Isomer-I);
- 5 *cis*-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(benzothiazol-6-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(1-methyl-1*H*-indazole-3-ylmethyl)-4-fluoro-1*H*-
- 10 indole-3-carboxamide (Isomer-I); and
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide (Isomer-I);
- or their pharmaceutically acceptable salt thereof.
- In yet another embodiment the representative compounds of pharmaceutically
- 15 acceptable salt of the present invention includes but not limited to,
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-II);
- 20 *trans*- N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-I);
- trans*- N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-II);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(5-fluoropyridin-4-ylmethyl)-4,7-difluoro-1*H*-
- 25 indole-3-carboxamide hydrochloride (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(2-chloropyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(2-chloropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide hydrochloride (Racemate);
- 30 *cis*-N-(3-Fluoropiperidin-4-yl)-1-(2-chloropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-I);


- cis*-N-(3-Fluoropiperidin-4-yl)-1-(4-fluorobenzyl)-4-fluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(4-fluorobenzyl)-4,7-difluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-I);
- 5 *cis*-N-(3-Fluoropiperidin-4-yl)-1-(4-fluorobenzyl)-4,7-difluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-II);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(2-fluorobenzyl)-4-fluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(4-chlorobenzyl)-4-fluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-I);
- 10 *cis*-N-(3-Fluoropiperidin-4-yl)-4-fluoro-1-(3-methoxybenzyl)-1*H*-indole-3-carboxamide hydrochloride (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-4-fluoro-1-(3-methoxybenzyl)-1*H*-indole-3-carboxamide hydrochloride (Isomer-II);
- 15 *cis*-N-(3-Fluoropiperidin-4-yl)-1-(4-methoxybenzyl)-4-fluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-II);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(4-methoxybenzyl)-4-fluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(4-methoxybenzyl)-4,7-difluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-I);
- 20 *cis*-N-(3-Fluoropiperidin-4-yl)-1-(4-methoxybenzyl)-4,7-difluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-II);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(3,4-difluorobenzyl)-4-fluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-I);
- 25 *cis*-N-(3-Fluoropiperidin-4-yl)-1-(1-methyl-1*H*-pyrazol-4-yl-methyl)-4-fluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(1-methyl-1*H*-pyrazol-4-yl-methyl)-4,7-difluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-I);
- cis*-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-I);
- 30 *cis*-N-(3-Fluoropiperidin-4-yl)-1-(benzothiazol-6-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-I); and

cis-N-(3-Fluoropiperidin-4-yl)-1-(1-methyl-1*H*-indazole-3-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide hydrochloride (Isomer-I).

In another embodiment, the present invention relates to the process for the preparation of the compound of formula (I) or a pharmaceutically acceptable salt thereof. The process for preparation of compound of formula (I) is given in the general schemes-1 and 2 wherein all the groups are as defined above.

General Scheme-1 depicts processes for the preparation of compound of formula (I), wherein R^1 , R^2 and R^3 are as defined above.

General Scheme-I



10

Step 1: Preparation of compound of formula B

The compound of formula A is reacted with trifluoroacetic anhydride in a solvent selected from DMF at RT for 2-4 hours to obtain the compound of formula B.

Step 2: Preparation of compound of formula C

15

The compound of formula B obtained in step 1 is reacted with R^2CH_2 -halo or R^2CH_2 -O-SO₂-CH₃ in presence of potassium carbonate, sodium hydride, cesium

carbonate or potassium tert-butoxide in a solvent selected from DMF, THF or acetonitrile overnight at RT to obtain the compound of formula C.

Step 3: Preparation of compound of formula D

The compound of formula C obtained in step 2 is reacted with aqueous sodium hydroxide or potassium hydroxide at 50-70 °C for 16-18 hours to obtain the compound of formula D.

Step 4: Preparation of compound of formula (I)

The compound of formula D obtained in step 3 is coupled with amine R^1 -NH₂.HCl in presence of coupling reagent, HATU, DCC, or EDC and a base, DIPEA in a solvent selected from DMF, THF, dichloromethane or 1,4-dioxane at RT overnight to obtain the compound of formula (I).

Step 5: Preparation of compound of formula E


The compound of formula B obtained in step 1 is reacted with aqueous sodium hydroxide at 50-70 °C for 16-18 hours to obtain the compound of formula E.

Step 6: Preparation of compound of formula F


The compound of formula E obtained in step 5 is coupled with amine R^1 -NH₂.HCl in presence of coupling reagent, HATU, DCC, or EDC and a base, DIPEA in a solvent selected from DMF, THF at RT overnight to obtain the compound of formula F.

Step 7: Preparation of compound of formula (I)

The compound of formula F obtained in step 6 is reacted with R^2 CH₂-halo or R^2 CH₂-O-SO₂-CH₃ in presence of potassium carbonate and potassium iodide in a solvent selected from DMF, overnight at RT to obtain the compound of formula (I).


Step 8: Preparation of compound of formula (I) (wherein R^2 is  ; R^4 is -S-CH₃)

The compound of formula (I) obtained from steps 4 and 7 (wherein R^2 is



; R^4 is F) is reacted with sodium thiomethoxide in the solvent selected from

DMF or THF at the temperature range of 55-65 °C for 2-5 hours to obtain the

compound of formula (I) (wherein R^2 is ; R^4 is $-S-CH_3$).


Preparation of pharmaceutically acceptable salt of compound of formula (I)

The compound of formula (I) can optionally be converted into its
5 pharmaceutically acceptable salt by reaction with the appropriate acid or acid derivative.

Suitable pharmaceutically acceptable salts will be apparent to those skilled in the art. The salts are formed with inorganic acids e. g. hydrochloric, hydrobromic, sulfuric, nitric & phosphoric acid or organic acids e.g., oxalic, succinic, maleic, acetic, fumaric, citric, malic, tartaric, benzoic, p-toluic, p-toluenesulfonic, benzenesulfonic
10 acid, methanesulfonic or naphthalenesulfonic acid.

Scheme 2 describes the process for the preparation of compound of formula (Ia) and (Ib), wherein R^2 and R^3 are as defined above.

Scheme 2



Step 1: Preparation of compound of formula G

The compound of formula D (given in scheme 1) is reacted with *tert*-butyl 4-amino-3-fluoropiperidine-1-carboxylate by following the procedure as described in step 4 of scheme-1 to obtain the compound of formula G.

5 Step 2: Preparation of compound of formula (Ia)

The compound of formula G (obtained in above step) is reacted with ethereal HCl in the solvents selected from DCM and the like at the temperature range of 25-30 °C for 2-4 hours to obtain the compound of formula (Ia).

Step 3: Preparation of compound of formula (Ib)

10 The compound of formula (Ia) (obtained in above step) is subjected to pH adjustment to 7-8 using sodium bicarbonate in water at the temperature range of 5-10 °C to obtain the compound of formula (Ib).

In yet another aspect, the present invention relates to the pharmaceutical composition of the compound of formula (I). In order to use the compound of formula
15 (I), or their stereoisomers and pharmaceutically acceptable salts thereof in therapy, they will normally be formulated into a pharmaceutical composition in accordance with standard pharmaceutical practice.

The pharmaceutical compositions of the present invention may be formulated in a conventional manner using one or more pharmaceutically acceptable excipients.
20 The pharmaceutically acceptable excipient is carrier or diluent. Thus, the active compounds of the invention may be formulated for oral dosing. Such pharmaceutical compositions and processes for preparing same are well known in the art.

The dose of the active compounds can vary depending on factors such as age and weight of patient, nature and severity of the disease to be treated and such other
25 factors. Therefore, any reference regarding pharmacologically effective amount of the compounds of general formula (I), stereoisomers and pharmaceutically acceptable salts thereof refers to the aforementioned factors.

In yet another aspect, the present invention relates to method of treatment of disorders related to muscarinic M1 receptors.

30 In another embodiment, the disorders related to muscarinic M1 receptors are selected from the group consisting of AD, schizophrenia, cognitive disorders, pain or sleep disorders.

Commercial reagents were used without further purification. RT is defined as an ambient temperature range, typically from 25 °C to 35 °C. All mass spectra were obtained using ESI conditions unless otherwise stated. ¹H-NMR spectra were recorded at 400 MHz on a Bruker instrument. Deuterated chloroform, methanol or dimethyl sulfoxide was used as solvent. Tetramethylsilane (TMS) was used as internal reference standard. Chemical shift values are expressed in parts per million (δ) values. The following abbreviations are used for the multiplicity of the NMR signals: s=singlet, bs=broad singlet, d=doublet, t=triplet, q=quartet, qui=quintet, h=heptet, dd=double doublet, dt=double triplet, tt=triplet of triplets, m=multiplet.

Chromatography refers to column chromatography performed using 100 - 200 mesh silica gel and executed under nitrogen pressure (flash chromatography) conditions.

The stereoisomers as a rule are generally obtained as racemates that can be separated into the optically active isomers in a manner known per se. In the case of the compounds of formula (I) having an asymmetric carbon atom the present invention relates to the D-form, the L-form and D, L - mixtures and in the case of compound of formula (I) containing a number of asymmetric carbon atoms, the diastereomeric forms and the invention extends to each of these stereoisomeric forms and to mixtures thereof including racemates. Those compounds of general formula (I) which have an asymmetric carbon and as a rule are obtained as racemates can be separated one from the other by the usual methods, or any given isomer may be obtained by stereo specific or asymmetric synthesis. However, it is also possible to employ an optically active compound from the start, a correspondingly optically active enantiomeric or diastereomeric compound then being obtained as the final compound.

The stereoisomers of compounds of general formula (I) may be prepared by one or more ways presented below:

- i) One or more of the reagents may be used in their optically active form.
- ii) Optically pure catalyst or chiral ligands along with metal catalyst may be employed in the reduction process. The metal catalyst may be Rhodium, Ruthenium, Indium and the like. The chiral ligands may preferably be chiral phosphines. (Principles of Asymmetric synthesis, J. E. Baldwin Ed., Tetrahedron series, 14, 311-316).

- iii) The mixture of stereoisomers may be resolved by conventional methods such as forming diastereomeric salts with chiral acids or chiral amines or chiral amino alcohols, chiral amino acids. The resulting mixture of diastereomers may then be separated by methods such as fractional crystallization, chromatography and the like, which is followed by an additional step of isolating the optically active product by hydrolyzing the derivative.
- iv) The mixture of stereoisomers may be resolved by conventional methods such as microbial resolution, resolving the diastereomeric salts formed with chiral acids or chiral bases.
- Chiral acids that can be employed may be tartaric acid, mandelic acid, lactic acid, camphorsulfonic acid, amino acids and the like. Chiral bases that can be employed may be cinchona alkaloids, brucine or a basic amino acid such as lysine, arginine and the like. In the case of the compounds of general formula (I) containing geometric isomerism the present invention relates to all of these geometric isomers.

Chiral HPLC Methods

Method A:

Column: CHIRALPAK AD-H (250X4.6) mm 5 μ m; Solvent A = 50.0% MeOH, B = 49.9% ACN, C = 0.1% DEA; Isocratic Flow = 1.5 mL/min T = 25 °C.

Method B:

Column: CHIRALPAK AD-H (250X4.6) mm 5 μ m; Solvent A = 99.9% MeOH, B = 0.1% DEA; Isocratic Flow = 0.8 mL/min T = 25 °C.

The following abbreviations are used herein:


ACN	:	Acetonitrile
CCl ₄	:	Carbon tetrachloride
CDCl ₃	:	Deuterated chloroform
DCM	:	Dichloromethane
DCC	:	N,N'-Dicyclohexylcarbodiimide
DEA	:	Diethylamine
DIPEA	:	N,N-Diisopropylethylamine
DMF	:	N,N-Dimethylformamide

	DMSO	:	Dimethyl sulfoxide
	EDC	:	Ethylene dichloride
	HATU	:	2-(7-Aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
5	HCl	:	Hydrochloric acid
	K ₂ CO ₃	:	Potassium carbonate
	MeOH	:	Methanol
	NaBH ₄	:	Sodium borohydride
	NaOH	:	Sodium hydroxide
10	Na ₂ SO ₄	:	Sodium sulphate
	RT	:	Room temperature (25-30 °C)
	THF	:	Tetrahydrofuran

Examples

15 The compounds of the present invention were prepared according to the following experimental procedures, using appropriate materials and conditions. The following examples are provided by way of illustration only but not to limit the scope of present invention.

Preparation 1: 4-Chloromethyl-1-methyl-1H-pyrazole hydrochloride (I-1)



20

Step-1: To a solution of 1H-pyrazole-4-carboxylic acid ethyl ester (35.0 g, 0.25 mole) in THF (100 mL) was added suspension of sodium hydride (17.38 g, 0.43 mole) in THF (100 mL) solution under N₂ at 25 °C and stirred for one hour. Methyl iodide (24 mL, 0.38 mole) was added at RT and the reaction mixture was heated to 60-65 °C for 6 hours. Reaction mixture was quenched in to ice water (200 mL) and extracted with ethyl acetate (100 mL x 3). The combined organic phase was washed with water (50 mL), brine solution (50 mL), dried over Na₂SO₄ and concentrated under vacuum to obtain ethyl 1-methyl-1H-pyrazole-4-carboxylate.

25 Yield: 32.36 g (83 %); ¹H - NMR (CDCl₃, 400 MHz) δ ppm: 1.30 - 1.33 (s, 3H), 3.91 (s, 3H), 4.25 - 4.30 (q, 2H), 7.83 (s, 1H), 7.88 (s, 1H); Mass (m/z): 155.0 (M+H)⁺.

30


Step-2: Lithium aluminium hydride (320 mL, 0.32 mole, 1M in THF) was added to a cooled solution of ethyl 1-methyl-1*H*-pyrazole-4-carboxylate (32.34 g, 0.21 mole) in THF (300 mL) under stirring in N₂ atmosphere. The reaction mixture was warmed to RT and stirred further for 3 hours. The reaction mixture was cooled to 0 °C, diluted with ethyl acetate and treated with water (25 mL). The mixture was filtered through celite bed and concentrated under vacuum to obtain the crude compound which was further purified by flash chromatography (ethyl acetate: methanol (98:2)) to afford (1-methyl-1*H*-pyrazol-4-yl)-methanol.

Yield: 14.66 g (62 %); ¹H - NMR (CDCl₃, 400 MHz) δ ppm: 1.98 (bs, 1H), 3.88 (s, 3H), 4.56 (s, 2H), 7.36 (s, 1H), 7.45 (s, 1H); Mass (m/z): 113.1 (M+H)⁺.

Step-3: To a cooled solution of (1-methyl-1*H*-pyrazol-4-yl)-methanol (8.61 g, 0.076 mole) in DCM (100 mL) under N₂ atmosphere, thionyl chloride (8.7 mL, 0.12 mole) was added drop wise. The reaction mixture was warmed to RT and stirred for 2 hours. The reaction mixture was concentrated under vacuum at 23 – 25 °C to obtain the title compound.


Yield: 12.77 g (99 %); ¹H - NMR (DMSO-*d*₆, 400 MHz) δ ppm: 3.85 (s, 3H), 4.67 (s, 2H), 4.76 - 4.79 (t, 1H), 4.88 (bs, 1H), 7.47 (s, 1H), 7.78 (s, 1H).

Preparation 2: 4-Bromomethyl-2-fluoropyridine (I-2)



To a solution of 2-fluoro-4-methylpyridine (75.0 g, 0.675 mole) in CCl₄ (200 mL) under N₂ atmosphere at 25 °C was added N-bromosuccinimide (160 g, 0.90 mole) and benzoyl peroxide (24.52 g, 0.101 mole). The reaction mass was gradually heated to 85 °C and stirred for 5 hours at this temperature. The reaction mass after cooling to RT was filtered under vacuum and washed with CCl₄ (50 mL). The filtrate was concentrated under vacuum to obtain the crude residue, which was purified by flash chromatography using ethyl acetate: n-hexane (02: 98) to afford the title compound.

Yield: 35.2 g (27 %); ¹H - NMR (CDCl₃, 400 MHz) δ ppm: 4.71 (s, 2H), 7.27 (s, 1H), 7.42 - 7.43 (d, J = 4.9 Hz, 1H), 8.24 - 8.25 (d, J = 5.1 Hz, 1H); Mass (m/z): 190.0 (M+H)⁺, 192.1 (M+H)⁺.


Preparation 3: 4-Bromomethyl-2,5-difluoropyridine (I-3)

Step-1: To a 0 °C cooled solution of 2,5-difluoroisonicotinic acid (0.5 g, 0.003 mole) in THF (5 mL) under N₂, was added lithium aluminum hydride (1 M in THF, 3.7 mL, 0.0037 mole) drop wise. The reaction mixture was warmed to RT and stirred further for 1.5 hours. The reaction mixture was cooled to 0 °C, diluted with ethyl acetate and treated with water (0.5 mL). The mixture was filtered through celite bed and concentrated under vacuum to obtain (2,5-difluoropyridin-4-yl)-methanol.

Yield: 0.45 g (98 %); ¹H - NMR (CDCl₃, 400 MHz) δ ppm: 4.60 (s, 2H), 4.96 (bs, 1H), 7.05 (s, 1H), 7.78 (s, 1H); Mass (m/z): 145.9 (M+H)⁺.

Step-2: To a 0 °C cooled solution of (2,5-difluoropyridin-4-yl)-methanol (0.45 g, 0.003 mole) in DCM (10 mL) under N₂, was added phosphorus tribromide (0.44 mL, 0.0037 mole) drop wise. The reaction mixture was warmed to RT and stirred for 1.5 hours. The reaction mixture was diluted with DCM (75 mL), treated with saturated aqueous sodium bicarbonate (20 mL) and partitioned. Organic layer was washed with water (20 mL), brine solution (20 mL) and dried over Na₂SO₄. The organic phase was concentrated under vacuum to obtain the title compound.

Yield: 0.23 g (37 %); ¹H - NMR (DMSO-*d*₆, 400 MHz) δ ppm: 4.68 (s, 2H), 7.20 (s, 1H), 8.16 (s, 1H); Mass (m/z): 208.1 (M+H)⁺, 210.1 (M+H)⁺.

Preparation 4: 4-Bromomethyl-2-chloropyridine (I-4)

Step-1: To a solution of 2-chloroisonicotinic acid (2.0 g, 0.012 mole) in DMF (5 mL) under N₂ at 25 °C, was added sodium hydride (0.73 g, 0.015 mole) and stirred for 0.5 hour. Methyl iodide (1.5 mL, 0.025 mole) was added at RT and the reaction mixture was warmed to 50 °C for 2 hours. The reaction mixture was dissolved in ice water (50 mL) and extracted with ethyl acetate (50 mL x 3). Organic layer was washed with brine solution (50 mL) and dried over Na₂SO₄. The organic phase was concentrated under vacuum to obtain methyl 2-chloroisonicotinate.

Yield: 2.1 g (100 %); Mass (m/z): 172.0 (M+H)⁺, 174.0 (M+H)⁺.


Step-2: To a cooled solution of methyl 2-chloroisonicotinate (1.7 g, 0.009 mole) in THF (30 mL) under N₂, was added lithium borohydride (0.43 g, 0.019 mole) in portions. The reaction mixture was warmed to RT and stirred further for 3 hours. The reaction mixture was concentrated under reduced pressure; residue was dissolved in ice cold water (50 mL) and extracted with ethyl acetate (50 mL x 3). Organic layer was washed with brine solution (50 mL) and dried over Na₂SO₄. The organic phase was concentrated under vacuum to obtain the crude compound which was further purified by flash chromatography using ethyl acetate: n-hexane (40: 60) to afford (2-chloropyridin-4-yl)-methanol.

Yield: 1.0 g (71 %); ¹H - NMR (CDCl₃, 400MHz) δ ppm: 2.29 (bs, 1H), 4.75 (s, 2H), 7.20 - 7.21 (d, J = 4.9 Hz, 1H), 7.31 (s, 1H), 8.32 - 8.33 (d, J = 5.0 Hz, 1H); Mass (m/z): 144.0 (M+H)⁺, 145.9 (M+H)⁺.

Step-3: (2-Chloropyridin-4-yl)-methanol was converted to the title compound using similar procedure as described in step-2 of preparation 3.

Yield: 0.49 g (85 %); ¹H - NMR (CDCl₃, 400 MHz) δ ppm: 4.35 (s, 2H), 7.36 (s, 1H), 8.37 - 8.38 (d, J = 4.9 Hz, 1H); Mass (m/z): 205.9 (M+H)⁺, 208.0 (M+H)⁺.

Preparation 5: 4-Chloromethyl-3-fluoropyridine (I-5)




Step-1: 3-Fluoroisonicotinic acid was converted to (3-fluoropyridin-4-yl)-methanol similar to a procedure described in step-1 of preparation 3.

Yield: 0.19 g (70 %); Mass (m/z): 128.1 (M+H)⁺.


Step-2: To a cooled solution of (3-fluoropyridin-4-yl)-methanol (0.19 g, 0.001 mole) in DCM (5 mL) under N₂, thionyl chloride (0.21 mL, 0.003 mole) was added drop wise. The reaction mixture was warmed to RT and stirred for 2 hours. The reaction mixture was diluted with DCM (50 mL), and treated with saturated aqueous sodium bicarbonate (10 mL). Organic layer was washed with water (20 mL), brine solution (20 mL) and dried over Na₂SO₄ and concentrated under vacuum to obtain the title compound.

Yield: 0.12 g (56 %); Mass (m/z): 146.0 (M+H)⁺, 148.0 (M+H)⁺.

Preparation 6: 1-Chloromethyl-2-fluorobenzene (I-6)

The title compound, 1-chloromethyl-2-fluorobenzene was synthesized from 2-fluorobenzoic acid following the procedure as described in preparation 5.

5 Yield: 0.455 g (100 %); Mass (m/z): 145 (M+H)⁺, 147.0 (M+H)⁺.

Preparation 7: Methanesulfonic acid 2, 3-difluoro-pyridin-4-ylmethyl ester (I-7)


Step-1: (2,3-Difluoropyridin-4-yl)-methanol was synthesized from 2,3-difluoroisonicotinic acid by following the procedure as described in preparation 3.

10 Yield: 0.16 g (29 %); Mass (m/z): 146.0 (M+H)⁺.

Step-2: (2,3-Difluoropyridin-4-yl)-methanol was converted to the title compound by reacting with methanesulfonyl chloride and the product was used as such without any purification.

Yield: 0.37 g (68 %).

15

Preparation 8: 1-Chloromethyl-3-methoxybenzene (I-8)

Step-1: To the solution of 3-hydroxybenzaldehyde (3.0 g, 0.025 mole) in DMF (15 mL) was added K₂CO₃ (10.18 g, 0.073 mole) and methyl iodide (6.93 g, 0.049 mole) at RT and stirred for 12 hours under nitrogen atmosphere. Reaction mixture was quenched in to water (50 mL) and extracted with ethyl acetate (50 mL x 3). The organic layer was dried over Na₂SO₄ and concentrated under vacuum to get 3-methoxybenzaldehyde.

20 Yield : 3.11 g (93 %); ¹H - NMR (CDCl₃, 400 MHz) δ ppm: 3.87 (s, 3H), 7.17 - 7.20 (m, 1H), 7.40 - 7.41 (m, 1H), 7.43 - 7.46 (m, 2H), 9.98 (s, 1H).

25


Step-2: To a solution of 3-methoxybenzaldehyde (3.11 g, 0.022 mole) in THF (10 mL) and methanol (20 mL), NaBH₄ (1.43 g, 0.042 mole) was added portion wise at 0 – 10 °C under nitrogen atmosphere. After addition, reaction mixture was stirred at RT for 12 hours. Reaction mixture was concentrated under vacuum and quenched in to
5 water (50 mL). Aqueous layer was extracted with ethyl acetate (50 mL x 3). The organic layer was dried over Na₂SO₄ and concentrated under vacuum to get (3-methoxy-phenyl)-methanol.

Yield: 2.95 g (93 %); ¹H - NMR (CDCl₃, 400 MHz) δ ppm: 3.85 (s, 3H), 4.69 - 4.71 (m, 2H), 6.85 - 6.88 (m, 1H), 6.96 - 6.97 (m, 2H), 7.28 - 7.32 (m, 1H).


10 **Step-3:** The title compound was synthesized from (3-methoxyphenyl)-methanol by the procedure described in preparation 5.

Yield: 3.03 g (78 %); ¹H - NMR (CDCl₃, 400 MHz) δ ppm: 3.82 (s, 3H), 4.57 (s, 2H), 6.85 - 6.88 (m, 1H), 6.94 - 6.98 (m, 2H), 7.26 - 7.30 (m, 1H).

15 **Preparation 9: *tert*-Butyl 4-amino-3-fluoropiperidine-1-carboxylate**




Step 1: *tert*-Butyl 4-trimethylsilyloxy-3,6-dihydro-2H-pyridine-1-carboxylate



Chlorotrimethylsilane (16.3 g, 0.15 mole) was added drop wise to a mixture
20 of *tert*-butyl 4-oxo-piperidine-1-carboxylate (25.0 g, 0.12 mole), triethylamine (42.6 mL, 0.31 mole) in 35 mL of DMF at 25 °C in 20 minutes under N₂. The reaction mass was heated to 90 - 92 °C and maintained for 20 hours. Reaction mass was allowed to cool to 25 °C. n-Hexane (120 mL) was added to the reaction mass and neutralized with saturated sodium bicarbonate solution (60 mL). Organic layer was separated,
25 washed with saturated sodium bicarbonate solution, water and brine solution. The

organic layer was dried over Na₂SO₄ and concentrated under vacuum at 45 °C to obtain the title compound. Yield: 33.3 g (66 %)


Step 2: *tert*-butyl-3-fluoro-4-oxo-piperidine-1-carboxylate



5 *tert*-Butyl 4-trimethylsilyloxy-3,6-dihydro-2H-pyridine-1-carboxylate (33.3 g, 0.12 mole) was added drop wise to Selectfluor® (30.6 g, 0.086 mole) in 250 mL of acetonitrile at RT in 15 minutes. The reaction was exothermic and the reaction mass temperature was raised to 60 °C and clear a solution was obtained. After addition, the reaction mass was stirred at RT for 10 hours under N₂. The reaction mass was diluted
10 with 200 mL of ethyl acetate and washed with brine solution (100 mL x 2). Organic layer was dried over Na₂SO₄ and concentrated on rota vac at 45 °C under vacuum to obtain residue, which was purified by flash chromatography using ethyl acetate:hexane (40: 60) to afford the title compound.


Yield: 15.2 g (55%);

15 **Step 3: *tert*-Butyl-4-(benzylamino)-3-fluoropiperidine-1-carboxylate**



To a clear solution of *tert*-butyl 3-fluoro-4-oxo-piperidine-1-carboxylate (3.0 g, 0.0138 mole, obtained from step 2) in EDC (50 mL), benzyl amine (1.77 g, 0.016 mole) was added and stirred for 2 hours. Sodium triacetoxyborohydride (5.86 g, 0.276 mole) was added at 10 °C. After addition, reaction mass was stirred for 12 hours
20 under N₂ at RT. The reaction mass was quenched into water, basified with lye solution and extracted with ethyl acetate (50 mL x 3). The organic layer was dried over Na₂SO₄ and concentrated under vacuum to obtain the crude residue, which was purified by flash chromatography using ethyl acetate:hexane (20: 80) to afford *tert*-
25 butyl 4-(benzylamino)-3-fluoropiperidine-1-carboxylate as *cis*- and *trans*-diastereomers separately.


Trans-diastereomer (3a):



Yield: 0.370 g (8.8 %);

¹H - NMR (CDCl₃, 400 MHz) δ ppm: 1.45 (s, 9H), 1.60 - 1.70 (m, 2H), 1.96 - 1.99 (m, 1H), 2.79 - 2.91 (m, 3H), 3.78 - 3.81 (m, 1H), 3.88 - 3.91 (m, 2H), 4.22 - 4.27 (m, 1H),
 5 4.36 - 4.39 (m, 1H), 7.24 - 7.35 (m, 5H); Mass (m/z): 309.2 (M+H)⁺.

Cis-diastereomer (3b):



Yield: 2.02 g (47 %);

¹H - NMR (DMSO-*d*₆, 400 MHz) δ ppm: 1.38 (s, 9H), 1.44 - 1.68 (m, 2H), 2.05 (m, 1H),
 10 2.57 - 2.66 (m, 2H), 2.93 - 3.02 (m, 1H), 3.76 - 3.77 (m, 2H), 3.90 (m, 1H), 4.12 (m, 1H),
 4.71 - 4.83 (m, 1H), 7.19 - 7.35 (m, 5H); Mass (m/z): 309.2 (M+H)⁺.

Trans-diastereomer (3a) obtained from step-3 was subjected to chiral separation using chiral HPLC 'Method A' to afford two enantiomers. *trans*-Enantiomer-I eluting at retention time 6.86 minutes and *trans*-Enantiomer-II eluting
 15 at retention time 11.96 minutes.

Similarly the *cis* diastereomer (3b), obtained from step-3 was subjected to chiral separation using chiral HPLC 'Method B' to afford two enantiomers. *cis*-Enantiomer-I eluting at retention time 5.76 minutes and *cis*-Enantiomer-II eluting at retention time 6.98 minutes.

20 **Step-4: *cis*-*tert*-Butyl 4-amino-3-fluoropiperidine-1-carboxylate (*cis*-Isomer-I) (I-9a)**

To a solution of *cis*-Enantiomer-I (as obtained in step-3, 2.0 g, 0.06 mole, obtained in above step) in methanol (50 mL), 10 % Pd/C (1.0 g, 0.5 v) was added in one portion and stirred for 2 hours under H₂ gas bubbling. Reaction mixture was
 25 filtered through celite and filtrate was concentrated under vacuum to obtain *cis*-*tert*-butyl 4-amino-3-fluoropiperidine-1-carboxylate (*cis*-Isomer-I).

Yield: 1.2 g (85 %); ^1H - NMR (DMSO- d_6 , 400 MHz) δ ppm: 1.38 (s, 9H), 1.50 - 1.54 (m, 2H), 1.69 - 1.70 (m, 2H); 2.76 - 2.80 (m, 2H), 3.00 (m, 1H), 3.85 (m, 1H), 4.07 (m, 1H), 4.45 - 4.63 (m, 1H); Mass (m/z): 219.1 (M+H) $^+$.

Step-5: *cis-tert*-Butyl 4-amino-3-fluoropiperidine-1-carboxylate (*cis*-Isomer-II)
(I-9b)

Cis-Enantiomer-II (obtained from step 3) was debenzylated by following the above procedure described in step 4 to obtain *cis-tert* butyl 4-amino-3-fluoropiperidine-1-carboxylate (*cis*-Isomer II).

Yield: 1.1 g (83.5 %); ^1H - NMR (DMSO- d_6 , 400 MHz) δ ppm: 1.38 (s, 9H), 1.42 - 1.54 (m, 2H), 1.82 (m, 2H), 2.71 - 2.81 (m, 2H), 2.99 (m, 1H), 3.85 (m, 1H), 4.08 (m, 1H), 4.46 - 4.58 (m, 1H); Mass (m/z): 219.1 (M+H) $^+$.

Step-6: *trans-tert*-Butyl 4-amino-3-fluoropiperidine-1-carboxylate (*trans*-Isomer-I)
(I-9c)

trans-Enantiomer-I (obtained from step-3) was debenzylated by following the above procedure described in step 4 to obtain *trans-tert*-butyl 4-amino-3-fluoropiperidine-1-carboxylate (*trans*-Isomer-I).


Yield: 0.33 g (96 %); ^1H - NMR (CDCl $_3$, 400 MHz) δ ppm: 1.45 (s, 9H), 1.55 - 1.57 (m, 2H), 1.86 - 1.89 (m, 2H), 2.70 - 2.78 (m, 2H), 2.89 - 2.91 (m, 1H), 4.01 - 4.05 (m, 1H), 4.13 - 4.14 (m, 1H), 4.20 - 4.28 (m, 1H); Mass (m/z): 219.1 (M+H) $^+$.

Step-7: *trans-tert*-Butyl 4-amino-3-fluoropiperidine-1-carboxylate (*trans*-Isomer-II)
(I-9d)

trans-Enantiomer-II (obtained from step-3) was debenzylated by following the above procedure described in step 4 to obtain *trans-tert*-butyl 4-amino-3-fluoropiperidine-1-carboxylate (*trans*-Isomer-II).

Yield: 0.31 g (95 %); ^1H - NMR (CDCl $_3$, 400 MHz) δ ppm: 1.45 (s, 9H), 1.52 - 1.57 (m, 2H), 1.86 - 1.89 (m, 2H), 2.75 - 2.80 (m, 2H), 2.86 - 2.94 (m, 1H), 3.98 - 4.04 (m, 1H), 4.11 - 4.16 (m, 1H), 4.20 - 4.28 (m, 1H); Mass (m/z): 219.1 (M+H) $^+$.

Preparation 10: 2, 2, 2-Trifluoro-1-(4-fluoro-1*H*-indol-3-yl)-ethanone (I-10)




To a solution of 4-fluoroindole (28.75 g, 0.213 mole (prepared as per *Org. Synth.* 1985, 63, 214) in DMF (200 mL), trifluoroacetic anhydride (73.50 g, 0.349 mole) was added slowly drop wise at 0 °C, under nitrogen atmosphere. After completion of addition, the reaction mixture was stirred at RT for 2 hours. Reaction mixture was cooled to 0 – 10 °C and quenched slowly in to ice cold water (500 mL). Reaction mass was stirred for 30 minutes. The solid thus obtained were filtered, washed with water (500 mL) followed by n- hexane (500 mL).The resulting solids were dried under vacuum to afford the title compound.


Yield: 37.25 g (75 %); ¹H - NMR (DMSO-*d*₆, 400MHz) δ ppm: 7.03 - 7.07 (m, 1H), 7.30 - 7.35 (m, 1H), 7.39 - 7.41 (m, 1H), 8.51 (s, 1H), 12.91 (s, 1H); Mass (m/z): 230 (M-H)⁺. **Preparation of Examples of compound of formula (I)**

Example 1:

N-[(1S,2S)-2-Hydroxycyclohexyl]-1-(1-methyl-1*H*-pyrazol-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide



Step-1: 1-[1-(1-Methyl-1*H*-pyrazol-4-ylmethyl)-4-fluoro-1*H*-indol-3-yl]-2,2,2-trifluoroethanone




To a cooled solution of the 2,2,2-trifluoro-1-(4-fluoro-1*H*-indol-3-yl)-ethanone (I-10, 13.30 g, 0.057 mole) in DMF (100 mL) under N₂ was added K₂CO₃ (47.06 g, 0.34 mole), 4-chloromethyl-1-methyl-1*H*-pyrazole hydrochloride (13.07 g, 0.078 mole) and the contents were stirred overnight at RT. Reaction mixture was

quenched in to ice cold water (1000 mL) and extracted with ethyl acetate (250 mL x 3). The combined organic layers were washed with water (200 mL x 3), brine solution (100 mL) and dried over Na₂SO₄. The organic phase was concentrated under vacuum to obtain a crude compound which was further purified by flash chromatography using (ethyl acetate: n-hexane (80:20)) to afford 1-[1-(1-methyl-1*H*-pyrazol-4-ylmethyl)-4-fluoro-1*H*-indole-3-yl]-2,2,2-trifluoroethanone.

Yield: 17.23 g (92 %); ¹H - NMR (CDCl₃, 400 MHz) δ ppm: 3.88 (s, 3H), 5.26 (s, 2H), 7.01 - 7.05 (m, 1H), 7.21 - 7.26 (m, 1H), 7.30 - 7.34 (m, 2H), 7.48 (s, 1H), 7.91 (s, 1H); Mass (m/z): 326.2 (M+H)⁺.

10 **Step-2: 1-(1-Methyl-1*H*-pyrazol-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxylic acid**



A mixture of 1-[1-(1-methyl-1*H*-pyrazol-4-ylmethyl)-4-fluoro-1*H*-indole-3-yl]-2,2,2-trifluoroethanone (21.39 g, 0.066 mole) and 4N aqueous NaOH (27.07 g, 0.67 mole) was heated to 98 - 100 °C for 5 hours. Reaction mass was cooled to 5 - 10 °C and diluted with 100 mL of ice cold water. Aqueous layer was acidified with acetic acid to pH ~ 5 at 5 - 10 °C. The solids thus obtained were filtered. These solids were extracted from mixture of ethyl acetate: methanol (80:20), dried over Na₂SO₄ and concentrated under vacuum to obtain 1-(1-methyl-1*H*-pyrazol-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxylic acid.

Yield: 16.71 g (93 %); ¹H - NMR (DMSO-*d*₆, 400 MHz) δ ppm: 3.75 (s, 1H), 5.29 (s, 2H), 6.87 - 6.92 (m, 1H), 7.17 - 7.22 (m, 1H), 7.40 - 7.50 (m, 2H), 7.73 (s, 1H), 8.11 (s, 1H); Mass (m/z): 274.3 (M+H)⁺.

25 **Step-3: N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(1-methyl-1*H*-pyrazol-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide**

To a solution of 1-(1-methyl-1*H*-pyrazol-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxylic acid (16.70 g, 0.061 mole) in DMF (120 mL) and HATU (29.21 g, 0.077 mole) stirred for 15 minutes followed by addition of (1*S*,2*S*) 2-amino cyclohexanol


hydrochloride (11.28 g, 0.074 mole) and DIPEA (49 mL, 0.28 mole) in 15 minutes of time interval at RT. During addition of DIPEA reaction mass becomes exothermic. After completion of addition, reaction mixture was stirred overnight at RT. Reaction mass was quenched slowly in to water (800 mL) and stirred for 1 hour. Solid obtained
5 was filtered, washed with water (1000 mL). These solids were extracted from mixture of ethyl acetate: methanol (80:20), dried over Na₂SO₄ and concentrated under vacuum to obtain solids. These solids were dissolved in ethyl acetate (700 mL), stirred at 60 °C and filtered to remove any undissolved particles. The filtrate was concentrated under vacuum to get the title product.

10 Yield: 17.64 g (78 %); ¹H - NMR (DMSO-*d*₆, 400 MHz) δ ppm: 1.17 - 1.24 (m, 4H), 1.59 - 1.65 (m, 2H), 1.86 - 1.88 (m, 1H), 1.96 - 1.98 (m, 1H), 3.59 - 3.61 (m, 1H), 3.76 (s, 3H), 4.70 - 4.71 (d, J = 4.7 Hz, 1H), 5.28 (s, 2H), 6.90 - 6.95 (m, 1H), 7.18 - 7.22 (m, 1H), 7.40 - 7.45 (m, 2H), 7.50 - 7.52 (d, J = 8.27 Hz, 1H), 7.71 (s, 1H), 8.00 (s, 1H); Mass (m/z): 371.2 (M+H)⁺, [α]_D²⁵ = + 34.8°.


15

Example 2:

N-[(1S,2S)-2-Hydroxycyclohexyl]-1-(2-chloropyridin-4-ylmethyl)-4-fluoro-1H-indole-3-carboxamide



20 **Step-1: 4-Fluoro-1H-indole-3-carboxylic acid**




To 2,2,2-trifluoro-1-(4-fluoro-1H-indol-3-yl)-ethanone (I-10, 18.49 g, 0.080 mole), 4N aqueous NaOH (200 mL, 0.80 mole) was added at RT and heated to 100 °C for 3 hours. Reaction mixture was cooled to RT and diluted with ice-cold water
25 (200 mL). The aqueous layer was washed with ethyl acetate (100 mL x 2) and

acidified with dilute HCl to pH ~ 4. Solid obtained was filtered, washed with water and n-hexane of each 100 mL separately. The solids were dried under vacuum.

Yield: 5.43 g (38 %); ¹H - NMR (DMSO-*d*₆, 400 MHz) δ ppm: 6.84 - 6.89 (m, 1H), 7.12 - 7.17 (m, 1H), 7.26 - 7.28 (m, 1H), 8.02 (s, 1H), 11.87 (s, 1H), 12.05 (m, 1H);

5 Mass (m/z): 180.2 (M+H)⁺.

Step-2: N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-4-fluoro-1*H*-indole-3-carboxamide



To a solution of 4-fluoro-1*H*-indole-3-carboxylic acid obtained in above step (0.39 g, 0.002 mole) in DMF (15 mL) under N₂ at 25 °C was added HATU (0.99 g, 0.0026 mole), (1*S*,2*S*)-2-aminocyclohexanol hydrochloride (0.39 gm, 0.0026 mole), DIPEA (1.5 mL, 0.0026 mole) with 5 minutes gap of each addition. The reaction mixture was stirred for overnight at RT. The reaction mixture was quenched in to water (50 mL) and extracted with ethyl acetate (25 mL x 3). Organic layer was washed with brine solution (20 mL), dried over Na₂SO₄ and concentrated under vacuum to obtain the crude residue, which was further purified by flash chromatography (methanol : chloroform (03: 97)) to afford the title compound.

Yield: 0.43 g (73 %); ¹H - NMR (DMSO-*d*₆, 400 MHz) δ ppm: 1.17 - 1.25 (m, 4H), 1.60 - 1.66 (m, 2H), 1.87 - 1.97 (m, 2H), 3.37 - 3.39 (m, 1H), 3.59 - 3.64 (m, 1H), 4.73 - 4.74 (d, J = 4.9 Hz, 1H), 6.87 - 6.92 (m, 1H), 7.12 - 7.17 (m, 1H), 7.28 - 7.30 (d, J = 8.1 Hz, 1H), 7.41 - 7.45 (t, 1H), 7.91 - 7.92 (d, J = 2.4 Hz, 1H), 11.88 (s, 1H);

20 Mass (m/z): 277.1 (M+H)⁺.



Step 3: N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2-chloropyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide




To a solution of N-[(1*S*,2*S*)-2-hydroxycyclohexyl]-4-fluoro-1*H*-indole-3-carboxamide (0.24 g, 0.0008 mole) in DMF (5 mL) under N₂ at 25 °C was added potassium carbonate (0.37 g, 0.0026 mole), potassium iodide (0.014 g, 0.00008 mole), 4-bromomethyl-2-chloropyridine (I-4, 0.25 g, 0.0012). The reaction mixture was stirred for overnight at RT. The reaction mixture was quenched in to ice water (50 mL) and extracted with ethyl acetate (50 mL x 3). Organic layer was washed with




brine solution (50 mL) and dried over Na₂SO₄. The organic phase was concentrated under vacuum to obtain the crude residue, which was further purified by flash chromatography using methanol: chloroform (02: 98) to afford the title compound.




Yield: 1.0 g (76 %); ¹H - NMR (CDCl₃, 400 MHz) δ ppm: 1.27 - 1.44 (m, 4H), 1.79 (m, 2H), 2.11 (m, 2H), 3.49 (m, 1H), 3.87 (m, 1H), 4.29 (s, 1H), 5.34 (s, 2H), 6.88 (s, 1H), 6.99 - 7.02 (m, 3H), 7.21 (m, 2H), 8.02 (s, 1H), 8.33 - 8.34 (d, J = 3.8 Hz, 1H); Mass (m/z): 402.2 (M+H)⁺.




Examples 3 to 19: The compounds of Examples 3 to 19 were prepared by following the experimental procedures as described in the Examples 1 and 2, with some noncritical variations




Example No.	Chemical name and Structure	Characterization data
3	 N-[(1R,2R)-2-Hydroxycyclohexyl]-1-(1-methyl-1H-pyrazol-4-ylmethyl)-4-fluoro-1H-indole-3-carboxamide	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.16 - 1.29 (m, 4H), 1.59 - 1.65 (m, 2H), 1.86 - 1.98 (m, 2H), 3.29 - 3.34 (m, 1H), 3.56 - 3.60 (m, 1H), 3.75 (s, 3H), 4.70 - 4.71 (d, 1H), 5.28 (s, 2H), 6.90 - 6.95 (m, 1H), 7.17 - 7.22 (m, 1H), 7.40 - 7.44 (t, 1H), 7.45 (s, 1H), 7.50 - 7.52 (d, J = 8.3 Hz, 1H), 7.71 (s, 1H), 8.00 (s, 1H); Mass (m/z): 371.3 (M+H) ⁺ .
4	 N-[(1S,2S)-2-Hydroxycyclohexyl]-1-(1-methyl-1H-indazol-3-ylmethyl)-4,7-difluoro-1H-indole-3-	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.16 - 1.23 (m, 4H), 1.59 - 1.65 (m, 2H), 1.86 - 1.98 (m, 2H), 3.29 - 3.34 (m, 1H), 3.56 - 3.60 (m, 1H), 4.01 (s, 3H), 4.70 - 4.71 (d, 1H), 5.81 (s, 2H), 6.89 - 6.95 (m, 1H), 7.09 - 7.13 (m, 1H), 7.16 - 7.21 (m, 1H), 7.35 - 7.39 (t, 1H), 7.43 - 7.47 (t, 1H), 7.53 - 7.55 (d, J = 8.2 Hz, 1H), 7.59 - 7.61 (d, J = 8.5 Hz, 1H), 7.65 - 7.67

	carboxamide	(d, J = 8.1 Hz, 1H), 8.13 (s, 1H); Mass (m/z): 421.3 (M+H) ⁺ .
5	 N-[(1S,2S)-2-Hydroxycyclohexyl]-1-(1-methyl-1H-pyrazol-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxamide	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.19 - 1.24 (m, 4H), 1.59 - 1.62 (m, 2H), 1.85 - 1.88 (m, 1H), 1.95 - 1.97 (m, 1H), 3.27 - 3.29 (m, 1H), 3.57 - 3.59 (m, 1H), 3.76 (s, 3H), 4.65 - 4.66 (d, J = 4.75 Hz, 1H), 5.35 (s, 2H), 6.86 - 6.88 (m, 1H), 7.01 - 7.03 (m, 1H), 7.40 (s, 1H), 7.48 - 7.53 (m, 1H), 7.66 (s, 1H), 8.00 (s, 1H); Mass (m/z): 389.3 (M+H) ⁺ .
6	 N-[(1S,2S)-2-Hydroxycyclohexyl]-1-(2-chloropyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxamide	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.23 (m, 4H), 1.61 - 1.62 (m, 2H), 1.87 - 1.97 (m, 2H), 3.40 (m, 1H), 3.60 (m, 1H), 4.66 - 4.67 (d, 1H), 5.63 (s, 2H), 6.88 - 6.93 (m, 1H), 6.97 - 7.04 (m, 2H), 7.22 (s, 1H), 7.66 - 7.69 (t, 1H), 8.10 (s, 1H), 8.35 - 8.36 (d, J = 5.0 Hz, 1H); Mass (m/z): 420.3 (M+H) ⁺ .
7	 N-[(1S,2S)-2-Hydroxycyclohexyl]-1-(2-fluoropyridin-4-yl-methyl)-4-fluoro-1H-indole-3-carboxamide	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.17 - 1.29 (m, 4H), 1.59 - 1.65 (t, 2H), 1.86 - 1.98 (dd, 2H), 3.36 (m, 1H), 3.60 - 3.61 (m, 1H), 4.69 - 4.70 (d, 1H), 5.61 (s, 2H), 6.92 - 6.97 (t, 2H), 7.05 - 7.06 (d, J = 4.7 Hz, 1H), 7.15 - 7.20 (m, 1H), 7.31 - 7.33 (d, J = 8.2 Hz, 1H), 7.51 - 7.55 (t, 1H), 8.11 (s, 1H), 8.16 - 8.17 (d, J = 5.1 Hz, 1H); Mass (m/z): 386.3 (M+H) ⁺ .

8	 N-[(1S,2S)-2-Hydroxycyclohexyl]-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxamide	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.22 - 1.29 (m, 4H), 1.60 - 1.66 (m, 2H), 1.88 - 1.98 (m, 2H), 3.33 - 3.38 (m, 1H), 3.61 - 3.63 (m, 1H), 4.66 - 4.68 (d, 1H), 5.67 (s, 2H), 6.86 - 6.93 (m, 2H), 6.96 - 7.02 (m, 2H), 7.65 - 7.68 (t, 1H), 8.11 (s, 1H), 8.18 - 8.20 (d, J = 5.11 Hz, 1H); Mass (m/z): 404.2 (M+H) ⁺ .
9	 N-[(1S,2S)-2-Hydroxycyclohexyl]-1-(3-fluoropyridin-4-ylmethyl)-4-fluoro-1H-indole-3-carboxamide	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.18 - 1.25 (m, 4H), 1.60 - 1.66 (m, 2H), 1.87 - 1.89 (m, 1H), 1.97 - 1.99 (m, 1H), 3.36 - 3.37 (m, 1H), 3.60 - 3.62 (m, 1H), 4.69 - 4.71 (d, J = 4.95 Hz, 1H), 5.67 (s, 2H), 6.86 - 6.89 (m, 1H), 6.94 - 6.99 (m, 1H), 7.17 - 7.23 (m, 1H), 7.35 - 7.37 (d, J = 8.2 Hz, 1H), 7.51 - 7.54 (m, 1H), 8.06 (s, 1H), 8.33 - 8.34 (d, J = 4.7 Hz, 1H), 8.59 (s, 1H); Mass (m/z): 386.2 (M+H) ⁺ .
10	 N-[(1S,2S)-2-Hydroxycyclohexyl]-1-(3-fluoropyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxamide	¹ H-NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.24 - 1.29 (m, 4H), 1.60 - 1.66 (m, 2H), 1.87 - 1.90 (m, 1H), 1.95 - 1.97 (m, 1H), 3.34 - 3.35 (m, 1H), 3.58 - 3.62 (m, 1H), 4.66 - 4.68 (d, J = 4.95 Hz, 1H), 5.72 - 5.76 (s, 2H), 6.76 - 6.79 (m, 1H), 6.88 - 6.93 (m, 1H), 6.97 - 7.03 (m, 1H), 7.67 - 7.64 (m, 1H), 8.08 (s, 1H), 8.33 - 8.35 (d, J = 4.8 Hz, 1H), 8.60 (s, 1H); Mass (m/z): 404.2 (M+H) ⁺ .


11	 N-[(1<i>S</i>,2<i>S</i>)-2-Hydroxycyclohexyl]-1-(2,5-difluoropyridin-4-ylmethyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide	¹ H - NMR (CDCl ₃ , 400 MHz) δ ppm: 1.33 - 1.48 (m, 4H), 1.79 (m, 2H), 2.10 - 2.11 (m, 2H), 3.47 - 3.51 (m, 1H), 3.88 (m, 1H), 4.25 (bs, 1H), 5.42 (s, 2H), 6.27 - 6.28 (m, 1H), 6.98 - 7.09 (m, 2H), 7.21 - 7.23 (m, 2H), 8.03 (s, 1H), 8.11 (s, 1H); Mass (m/z): 404.2 (M+H) ⁺ .
12	 N-[(1<i>S</i>,2<i>S</i>)-2-Hydroxycyclohexyl]-1-(2,5-difluoropyridin-4-ylmethyl)-4,7-difluoro-1<i>H</i>-indole-3-carboxamide	¹ H-NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.23 (m, 4H), 1.63 (m, 2H), 1.87 - 1.96 (m, 2H), 3.51 (m, 1H), 3.60 (m, 1H), 4.66 (m, 1H), 5.73 (s, 2H), 6.61 (s, 1H), 6.91 - 7.02 (m, 2H), 7.66 (m, 1H), 8.04 (s, 1H), 8.31 (s, 1H); Mass (m/z): 422.3 (M+H) ⁺ .
13	 N-[(1<i>S</i>,2<i>S</i>)-2-Hydroxycyclohexyl]-1-(2,3-difluoropyridin-4-ylmethyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide	¹ H-NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.23 - 1.24 (m, 4H), 1.60 - 1.65 (m, 2H), 1.86 - 1.89 (m, 1H), 1.96 - 1.98 (m, 1H), 3.43 - 3.46 (m, 1H), 3.57 - 3.60 (m, 1H), 4.69 - 4.71 (d, J = 4.5 Hz, 1H), 5.74 (s, 2H), 6.76 - 6.77 (m, 1H), 6.95 - 7.00 (m, 1H), 7.18 - 7.23 (m, 1H), 7.36 - 7.38 (d, J = 8.1 Hz, 1H), 7.52 - 7.56 (m, 1H), 7.93 - 7.94 (d, J = 4.4 Hz, 1H), 8.06 (s, 1H); Mass (m/z): 404.2 (M+H) ⁺ .

14	 N-[(1<i>S</i>,2<i>S</i>)-2-Hydroxycyclohexyl]-1-(1-pyridin-4-ylmethyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.23 - 1.25 (m, 4H), 1.60 - 1.66 (m, 2H), 1.87 - 2.00 (m, 2H), 3.40 (m, 1H), 3.60 - 3.61 (m, 1H), 4.71 - 4.72 (d, 1H), 5.57 (s, 2H), 6.90 - 6.95 (m, 1H), 7.13 - 7.16 (m, 1H), 7.18 - 7.20 (d, <i>J</i> = 7.9 Hz, 1H), 7.27 - 7.30 (m, 1H), 7.33 - 7.35 (d, <i>J</i> = 8.2 Hz, 1H), 7.46 - 7.49 (t, 1H), 7.73 - 7.77 (t, 1H), 8.09 (s, 1H), 8.50 - 8.52 (d, <i>J</i> = 4.1 Hz, 1H); Mass (<i>m/z</i>): 368.3 (<i>M</i> + <i>H</i>) ⁺ .
15	 N-[(1<i>S</i>,2<i>S</i>)-2-Hydroxycyclohexyl]-1-(1<i>H</i>-pyrazol-4-ylmethyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.16 - 1.24 (m, 4H), 1.59 - 1.65 (m, 2H), 1.85 - 1.97 (m, 2H), 3.29 - 3.34 (m, 1H), 3.58 (m, 1H), 4.70 - 4.71 (d, 1H), 5.31 (s, 2H), 6.90 - 6.95 (m, 1H), 7.17 - 7.20 (m, 1H), 7.40 - 7.44 (t, 1H), 7.51 - 7.54 (m, 2H), 7.81 (s, 1H), 7.99 (s, 1H), 12.79 (bs, 1H); Mass (<i>m/z</i>): 357.3 (<i>M</i> + <i>H</i>) ⁺ .
16	 N-[(1<i>S</i>,2<i>S</i>)-2-Hydroxycyclohexyl]-1-(2-bromothiazol-5-ylmethyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide	¹ H - NMR (CDCl ₃ , 400 MHz) δ ppm: 1.25 - 1.36 (m, 4H), 1.75 - 1.78 (m, 2H), 2.04 - 2.13 (m, 2H), 3.44 - 3.49 (m, 1H), 3.84 - 3.86 (m, 1H), 4.30 - 4.31 (bs, 1H), 5.43 (s, 2H), 6.96 - 7.01 (m, 1H), 7.15 - 7.23 (m, 3H), 7.49 - 7.52 (m, 1H), 7.98 (s, 1H); Mass (<i>m/z</i>): 452.2, 454.2 (<i>M</i> + <i>H</i>) ⁺ .

17	 N-[(1<i>S</i>,2<i>S</i>)-2-Hydroxycyclohexyl]-1-(1-ethyl-5-methyl-1<i>H</i>-pyrazol-4-ylmethyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.17 - 1.24 (m, 4H), 1.48 - 1.51 (t, 3H), 1.59 - 1.65 (m, 2H), 1.86 - 1.88 (m, 1H), 1.96 - 1.98 (m, 1H), 2.78 (s, 3H), 3.59 - 3.61 (m, 1H), 3.69 - 3.73 (m, 2H), 4.70 - 4.71 (d, J = 4.7 Hz, 1H), 5.28 (s, 2H), 6.90 - 6.95 (m, 1H), 7.18 - 7.22 (m, 1H), 7.40 - 7.45 (m, 2H), 7.50 - 7.52 (d, J = 8.27 Hz, 1H), 7.71 (s, 1H), 8.00 (s, 1H); Mass (m/z): 399.3 (M+H) ⁺ .
18	 N-[(1<i>S</i>,2<i>S</i>)-2-Hydroxycyclohexyl]-1-(benzothiazol-6-ylmethyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.16 - 1.33 (m, 4H), 1.60 - 1.66 (m, 2H), 1.87 - 1.99 (m, 2H), 3.29 - 3.34 (m, 1H), 3.58 - 3.62 (m, 1H), 4.70 - 4.71 (d, 1H), 5.65 (s, 2H), 6.90 - 6.96 (m, 1H), 7.14 - 7.19 (m, 1H), 7.42 - 7.51 (m, 3H), 8.04 - 8.09 (m, 2H), 8.16 (s, 1H), 9.37 (s, 1H); Mass (m/z): 424.20 (M+H) ⁺ .
19	 N-[(1<i>S</i>,2<i>S</i>)-2-Hydroxycyclohexyl]-1-(1-(2-fluoroethyl)-1<i>H</i>-pyrazol-4-ylmethyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.25 - 1.39 (m, 4H), 1.75 - 1.77 (m, 2H), 2.06 - 2.13 (m, 2H), 3.43 - 3.48 (m, 2H), 3.83 - 3.85 (m, 1H), 4.31 - 4.33 (t, 1H), 4.38 - 4.40 (t, 1H), 4.65 - 4.68 (t, 1H), 4.77 - 4.79 (t, 1H), 5.20 (s, 2H), 6.92 - 6.98 (m, 1H), 7.17 - 7.24 (m, 3H), 7.39 (s, 1H), 7.48 (s, 1H), 7.96 (s, 1H); Mass (m/z): 403.2 (M+H) ⁺ .

Example 20:

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2-methylsulfanyl-pyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide




5 To a suspension of N-[(1*S*,2*S*)-2-hydroxycyclohexyl]-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide (Example 8, 0.050 g, 0.00012 mole) in THF (10 mL) sodium thiomethoxide (0.020 g, 0.00029 mole) was added and warmed to 60 °C for 3 hours. Reaction mixture was cooled to RT and quenched in to water. Aqueous layer was extracted with ethyl acetate (25 mL x 3). The organic layer
10 was dried over Na₂SO₄ and concentrated under vacuum to obtain the crude residue, which was further purified by preparative TLC using ethyl acetate to provide the title compound.

Yield: 9.0 mg (16.5 %); ¹H - NMR (DMSO-*d*₆, 400 MHz) δ ppm: 1.24 (m, 5H), 1.60 - 1.66 (m, 2H), 1.87 - 1.90 (m, 1H), 1.94 - 1.97 (m, 1H), 2.46 (s, 3H), 3.34 - 3.35 (m, 1H), 3.60 (m, 1H), 5.55 (s, 2H), 6.71 - 6.73 (d, J = 4 Hz, 1H), 6.86 - 6.92 (m, 1H),
15 6.97 - 7.02 (m, 2H), 7.65 - 7.68 (m, 1H), 8.09 (s, 1H), 8.35 - 8.36 (d, J = 5.1 Hz, 1H); Mass (m/z): 432.2 (M+H)⁺.

Example 21:

20 **N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2-methylsulfanyl-pyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide**




The title compound was prepared by the experimental procedure as described in the Example 20 using example 7 with some noncritical variations.

¹H NMR (DMSO-*d*₆, 400 MHz) δ ppm: 1.18 - 1.23 (m, 4H), 1.33 - 1.35 (m, 1H), 1.60 - 1.66 (m, 2H), 1.87 - 1.89 (m, 1H), 1.97 - 1.99 (m, 1H), 3.33 - 3.38 (m, 3H), 3.40 - 3.48 (m, 1H), 4.70 - 4.71 (d, 1H), 5.50 (s, 2H), 6.79 - 6.81 (m, 1H), 6.92 - 6.97 (m, 1H), 7.11 (s, 1H), 7.15 - 7.21 (m, 1H), 7.31 - 7.33 (m, 1H), 7.51 - 7.55 (m, 1H), 8.10 (s, 1H), 8.34 - 8.35 (m, 1H); Mass (m/z): 413.51 (M+H)⁺.

Example 22:

Cis-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxamide hydrochloride (Isomer-I)



Step 1: cis-tert-Butyl 4-[[4,7-Difluoro-1-(2-fluoropyridin-4-ylmethyl)-1H-indole-3-carbonyl]-amino]-3-fluoro-piperidine-1-carboxylate (Isomer-I)

The title compound was synthesized by the procedure described in the Example 1 step (2) using 1-(2-fluoro-pyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxylic acid and cis-tert-butyl 4-amino-3-fluoro-piperidine-1-carboxylate (Isomer-I), (I-9a). The crude product obtained was further purified by flash chromatography (methanol: DCM (1: 99)) to obtain the title compound.

Yield: 1.0 g (83 %). ¹H - NMR (DMSO-*d*₆, 400 MHz) δ ppm: 1.40 (s, 9H), 1.68 - 1.74 (m, 2H), 2.81 - 3.25 (m, 2H), 3.90 - 4.05 (m, 1H), 4.15 - 4.23 (m, 2H), 4.75 - 4.87 (m, 1H), 5.67 (s, 2H), 6.87 - 6.93 (m, 2H), 6.97 - 7.03 (m, 2H), 7.99 - 8.02 (m, 1H), 8.15 (s, 1H), 8.19 - 8.20 (m, 1H); Mass (m/z): 507.2 (M+H)⁺.

Step 2: cis-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxamide hydrochloride (Isomer-I)


To a 0 - 10 °C cooled solution of the above compound (0.675 g, 1.328 moles) in DCM (10 mL) under N₂, ethereal HCl (33 % w/w, 0.243 g, 6.64 moles) was added slowly. After addition, reaction mass was allowed to 25 °C and stirred for 2 hours.

Reaction Mass was concentrated under vacuum. The reaction mass was triturated with diethyl ether (5 mL x 2), decanted the solvent and solids were dried under vacuum to afford the title compound.

Yield: 0.650 g (96.5 %); ^1H - NMR (DMSO- d_6 , 400 MHz) δ ppm: 1.88 - 2.01 (m, 2H), 3.12 - 3.18 (m, 2H), 3.58 - 3.63 (m, 2H), 4.30 - 4.38 (m, 1H), 5.01 - 5.13 (d, 1H), 5.68 (s, 2H), 6.86 - 6.94 (m, 2H), 6.99 - 7.05 (m, 2H), 8.15 (s, 1H), 8.19 - 8.21 (m, 2H), 8.60 - 8.63 (m, 1H), 9.06 - 9.08 (m, 1H); Mass (m/z): 407.2 M+H $^+$.

Example 23:

***cis*-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxamide hydrochloride (Isomer-II)**



Step 1: *cis*-tert-Butyl 4-[[4,7-difluoro-1-(2-fluoropyridin-4-ylmethyl)-1H-indole-3-carbonyl]-amino}-3-fluoro-piperidine-1-carboxylate (Isomer-II)

The title compound was synthesized by the procedure described in the Example 1 step (2) using 1-(2-fluoro-pyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxylic acid and *cis*-tert-butyl 4-amino-3-fluoro-piperidine-1-carboxylate (Isomer-II), (I-9b). The crude product obtained was further purified by flash chromatography (methanol: DCM (1: 99)) to obtain the title compound.

Yield: 0.092 g (85 %). ^1H - NMR (CDCl $_3$, 400 MHz) δ ppm: 1.48 (s, 9H), 1.86 - 1.88 (m, 2H), 2.86 (m, 2H), 3.47 - 3.49 (m, 2H), 4.28 - 4.38 (m, 1H), 4.73 - 4.85 (m, 1H), 5.51 (s, 2H), 6.58 (s, 1H), 6.85 - 6.88 (m, 3H), 7.36 - 7.51 (m, 1H), 7.95 (s, 1H), 8.16 - 8.18 (d, J = 5.1 Hz, 1H); Mass (m/z): 507.2 (M+H) $^+$.

Step 2: *cis*-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxamide hydrochloride (Isomer-II)


The title compound was synthesized by the procedure described in example 22, step 2 using tert-butyl 4-[[4,7-difluoro-1-(2-fluoropyridin-4-ylmethyl)-1H-indole-3-carbonyl]-amino}-3-fluoro-piperidine-1-carboxylate (Isomer-II).

Yield: 0.039 gm (97.2 %). ¹H - NMR (DMSO-*d*₆, 400 MHz) δ ppm: 1.87 - 2.04 (m, 2H), 3.12 - 3.18 (m, 2H), 3.58 - 3.63 (m, 2H), 4.30 - 4.38 (m, 1H), 5.01 - 5.13 (d, 1H), 5.68 (s, 2H), 6.86 - 6.94 (m, 2H), 6.99 - 7.05 (m, 2H), 8.15 (s, 1H), 8.19 - 8.21 (m, 2H), 8.60 - 8.63 (m, 1H), 9.06 - 9.08 (m, 1H); Mass (m/z): 407.2 (M+H)⁺.

5

Example 24:

***trans*- N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxamide hydrochloride (Isomer-I)**



10 **Step 1: *trans*- *tert*-Butyl 4-[[4,7-Difluoro-1-(2-fluoropyridin-4-ylmethyl)-1H-indole-3-carbonyl]-amino]-3-fluoro-piperidine-1-carboxylate (Isomer-I)**

The title compound was synthesized by the procedure described in the Example 1, step (2) using 1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxylic acid and *trans-tert*-butyl 4-amino-3-fluoropiperidine-1-carboxylate (Isomer-I), (I-9c). The crude product obtained was further purified by flash chromatography (methanol: DCM (1: 99) to obtain the title compound.

15 Yield: 0.05 g (60 %). ¹H - NMR (CDCl₃, 400 MHz) δ ppm: 1.47 (s, 9H), 1.58 - 1.62 (m, 2H), 2.22 - 2.26 (m, 1H), 3.18 - 3.23 (m, 2H), 3.79 - 3.82 (m, 1H), 4.38 - 4.40 (m, 1H), 4.41 - 4.54 (m, 1H), 5.51 (s, 2H), 6.58 (s, 1H), 6.85 - 6.89 (m, 3H), 7.14 - 7.21 (m, 1H), 7.98 (s, 1H), 8.16 - 8.18 (d, J = 5.1 Hz, 1H); Mass (m/z): 507.3 (M+H).

20

Step 2: *trans*-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxamide hydrochloride (Isomer-I)

The title compound was synthesized by the procedure described in example 22, step (2) using *trans-tert*-butyl 4-[[4,7-difluoro-1-(2-fluoropyridin-4-ylmethyl)-1H-indole-3-carbonyl]-amino]-3-fluoro-piperidine-1-carboxylate (Isomer-I).


25

Yield: 0.035 g (90 %). ¹H - NMR (DMSO-*d*₆, 400 MHz) δ ppm: 1.78 - 1.82 (m, 1H), 2.11 - 2.13 (m, 1H), 3.12 - 3.18 (m, 1H), 3.26 - 3.29 (m, 2H), 3.56 - 3.59 (m, 1H), 4.33 - 4.34 (m, 1H), 4.79 - 4.92 (m, 1H), 5.68 (s, 2H), 6.85 - 6.93 (m, 2H), 6.99 - 7.05

(m, 2H), 8.14 (s, 1H), 8.19 - 8.21 (d, J = 5.1 Hz, 1H), 8.35 - 8.37 (d, J = 7.6 Hz, 1H), 9.05 (bs, 1H), 9.25 (bs, 1H); Mass (m/z): 407.2 (M+H)⁺.

Example 25:

5 *trans*-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxamide hydrochloride (Isomer-II)



Step 1: *trans-tert*-Butyl 4-[[4,7-difluoro-1-(2-fluoropyridin-4-ylmethyl)-1H-indole-3-carbonyl]-amino}-3-fluoro-piperidine-1-carboxylate (Isomer-II)

10 The title compound was synthesized by the procedure described in the Example 1, step (2) using 1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxylic acid and *trans-tert*-butyl 4-amino-3-fluoro-piperidine-1-carboxylate (Isomer-II), (I-9d). The crude product obtained was further purified by flash chromatography (methanol: DCM (1: 99)) to obtain the title compound.

15 Yield: 0.05 g (90 %). ¹H - NMR (CDCl₃, 400 MHz) δ ppm: 1.48 (s, 9H), 1.86 - 1.88 (m, 2H), 2.86 (m, 2H), 3.47 - 3.49 (m, 2H), 4.28 - 4.38 (m, 1H), 4.73 - 4.85 (m, 1H), 5.51 (s, 2H), 6.58 (s, 1H), 6.85 - 6.88 (m, 3H), 7.36 - 7.51 (m, 1H), 7.95 (s, 1H), 8.16 - 8.18 (d, J = 5.1 Hz, 1H); Mass (m/z): 507.2 (M+H)⁺.



20 Step 2: *trans*-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxamide hydrochloride (Isomer-II)




 The title compound was synthesized by the procedure described in example 22, step (2) using *trans-tert*-butyl 4-[[4,7-difluoro-1-(2-fluoropyridin-4-ylmethyl)-1H-indole-3-carbonyl]-amino}-3-fluoro-piperidine-1-carboxylate (Isomer-II).




25 Yield: 0.035 g (90 %). ¹H - NMR (DMSO-*d*₆, 400 MHz) δ ppm: 1.78 - 1.82 (m, 1H), 2.11 - 2.13 (m, 1H), 3.13 - 3.18 (m, 1H), 3.25 - 3.29 (m, 2H), 3.55 - 3.59 (m, 1H), 4.33 - 4.34 (m, 1H), 4.78 - 4.92 (m, 1H), 5.68 (s, 2H), 6.85 - 6.93 (m, 2H), 6.99 - 7.05 (m, 2H), 8.14 (s, 1H), 8.19 - 8.21 (d, J = 5.1 Hz, 1H), 8.35 - 8.37 (d, J = 7.7 Hz, 1H), 9.05 (bs, 1H), 9.24 (bs, 1H); Mass (m/z): 407.2 (M+H)⁺.




Examples 26 to 46: The compounds of Examples 26 to 46 were prepared by following the experimental procedures as described in the Examples 22-25, with some noncritical variations




5




Example No.	Chemical name and Structure	Characterization data
26	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(3-fluoropyridin-4-ylmethyl)-4,7-difluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-I)	¹ H-NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.85 - 1.90 (m, 1H), 1.96 - 2.05 (m, 1H), 3.05 - 3.16 (m, 1H), 3.27 - 3.30 (m, 1H), 3.34 - 3.37 (m, 1H), 3.55 - 3.61 (m, 1H), 4.30 - 4.38 (m, 1H), 5.0 - 5.12 (m, 1H), 5.74 (s, 2H), 6.79 - 6.82 (m, 1H), 6.88 - 6.94 (m, 1H), 6.99 - 7.04 (m, 1H), 8.13 (s, 1H), 8.17 - 8.20 (m, 1H), 8.35 - 8.36 (d, J = 4.8 Hz, 1H), 8.62 (s, 1H), 8.68 - 8.71 (m, 1H), 9.33 (m, 1H); Mass (m/z): 407.2 (M+H) ⁺ .
27	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(2-chloropyridin-4-ylmethyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-I)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.89 - 2.00 (m, 2H), 3.1 - 3.16 (m, 2H), 3.26 - 3.30 (m, 2H), 4.25 - 4.35 (m, 1H), 5.01 - 5.13 (d, 1H), 5.60 (s, 2H), 6.94 - 6.99 (m, 1H), 7.13 - 7.14 (m, 1H), 7.18 - 7.20 (m, 1H), 7.31 (s, 1H), 7.34 - 7.36 (m, 1H), 8.02 - 8.06 (m, 1H), 8.17 (s, 1H), 8.35 - 8.36 (m, 1H), 8.64 - 8.66 (m, 1H), 9.18 - 9.21 (m, 1H); Mass (m/z): 405.2 (M+H) ⁺ .




28	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(2-chloropyridin-4-ylmethyl)-4,7-difluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Racemate)	¹ H- NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.87 - 2.00 (m, 2H), 3.09 - 3.18 (m, 2H), 3.57 - 3.63 (m, 2H), 4.30 - 4.38 (m, 1H), 5.01 - 5.13 (d, 1H), 5.65 (s, 2H), 6.88 - 6.94 (m, 1H), 6.99 - 7.06 (m, 2H), 7.22 (s, 1H), 8.15 (s, 1H), 8.20 - 8.22 (m, 1H), 8.36 - 8.37 (dd, <i>J</i> = 5.0 Hz, 1H), 8.60 - 8.63 (m, 1H), 9.05 - 9.07 (m, 1H); Mass (m/z): 423.2 (M+H) ⁺ .
29	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(2-chloropyridin-4-ylmethyl)-4,7-difluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-I)	¹ H- NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.86 - 2.02 (m, 2H), 3.08 - 3.16 (m, 2H), 3.57 - 3.63 (m, 2H), 4.30 - 4.37 (m, 1H), 5.00 - 5.12 (d, 1H), 5.65 (s, 2H), 6.88 - 6.94 (m, 1H), 6.99 - 7.06 (m, 2H), 7.22 (s, 1H), 8.15 (s, 1H), 8.20 - 8.22 (m, 1H), 8.36 - 8.37 (dd, <i>J</i> = 5.0 Hz, 1H), 8.68 - 8.71 (m, 1H), 9.32 - 9.35 (m, 1H); Mass (m/z): 423.2 (M+H) ⁺ .
30	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(4-fluorobenzyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-I)	¹ H- NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.92 - 2.02 (m, 2H), 3.08 - 3.17 (m, 2H), 3.27 - 3.38 (m, 2H), 4.30 - 4.38 (m, 1H), 5.01 - 5.13 (d, 1H), 5.48 (s, 2H), 6.91 - 6.96 (m, 1H), 7.14 - 7.21 (m, 3H), 7.31 - 7.35 (m, 2H), 7.40 - 7.42 (d, <i>J</i> = 8.3 Hz, 1H), 7.92 - 7.95 (t, 1H), 8.16 (s, 1H), 8.61 - 8.64 (m, 1H), 9.12 - 9.14 (m, 1H); Mass (m/z): 388.3 (M+H) ⁺ .


31	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(4-fluorobenzyl)-4,7-difluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-I)	¹ H- NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.89 - 2.00 (m, 2H), 3.11 - 3.14 (m, 2H), 3.56 - 3.59 (m, 2H), 4.29 - 4.37 (m, 1H), 5.00 - 5.12 (d, 1H), 5.53 (s, 2H), 6.86 - 6.90 (m, 2H), 7.15 - 7.23 (m, 4H), 8.15 (s, 2H), 8.61 - 8.63 (m, 1H), 9.09 - 9.12 (m, 1H); Mass (m/z): 406.2 (M+H) ⁺ .
32	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(4-fluorobenzyl)-4,7-difluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-II)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.86 - 1.89 (m, 1H), 1.94 - 2.03 (m, 1H), 3.11 - 3.13 (m, 1H), 3.26 - 3.34 (m, 2H), 3.56 - 3.58 (m, 1H), 4.28 - 4.36 (m, 1H), 5.00 - 5.12 (d, 1H), 5.53 (s, 2H), 6.85 - 6.91 (m, 1H), 7.00 - 7.04 (m, 1H), 7.15 - 7.25 (m, 4H), 8.16 (bs, 2H), 8.63 - 8.65 (d, J = 9.24 Hz, 1H), 9.17 - 9.19 (d, J = 9.84 Hz, 1H); Mass (m/z): 406.4 (M+H) ⁺ .
33	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(2-fluorobenzyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-I)	¹ H- NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.91 - 1.99 (m, 2H), 3.09 - 3.17 (m, 2H), 3.27 - 3.34 (m, 2H), 4.29 - 4.39 (m, 1H), 5.01 - 5.13 (d, 1H), 5.56 (s, 2H), 6.93 - 6.98 (m, 1H), 7.13 - 7.27 (m, 4H), 7.34 - 7.39 (m, 1H), 7.40 - 7.43 (d, J = 8.2 Hz, 1H), 7.95 - 7.98 (t, 1H), 8.07 (s, 1H), 8.61 (m, 1H), 9.01 - 9.03 (m, 1H); Mass (m/z): 388.2 (M+H) ⁺ .

34	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(4-chlorobenzyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-I)	¹ H- NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.88 - 1.99 (m, 2H), 3.12 - 3.15 (m, 2H), 3.28 - 3.30 (m, 2H), 4.22 - 4.38 (m, 1H), 5.01 - 5.13 (d, 1H), 5.50 (s, 2H), 6.92 - 6.97 (m, 1H), 7.12 (s, 1H), 7.16 - 7.21 (m, 1H), 7.26 - 7.28 (d, <i>J</i> = 8.3 Hz, 2H), 7.37 - 7.41 (t, 3H), 7.95 - 7.99 (t, 1H), 8.16 (s, 1H), 8.58 - 8.60 (m, 1H), 8.95 - 8.98 (m, 1H); Mass (<i>m/z</i>): 404.2 (<i>M</i> + <i>H</i>) ⁺ .
35	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-4-fluoro-1-(3-methoxybenzyl)-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-I)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.92 - 1.99 (m, 2H), 3.12 - 3.16 (m, 2H), 3.42 - 3.57 (m, 2H), 3.71 (s, 3H), 4.30 - 4.40 (m, 1H), 5.02 - 5.13 (d, 1H), 5.46 (s, 2H), 6.77 - 6.79 (d, <i>J</i> = 7.3 Hz, 1H), 6.84 - 6.85 (m, 2H), 6.91 - 6.96 (m, 1H), 7.16 - 7.26 (m, 2H), 7.39 - 7.41 (d, <i>J</i> = 8.1 Hz, 1H), 7.93 - 7.96 (t, 1H), 8.14 (s, 1H), 8.59 (bs, 1H), 9.04 (bs, 1H); Mass (<i>m/z</i>): 400.3 (<i>M</i> + <i>H</i>) ⁺ .
36	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-4-fluoro-1-(3-methoxybenzyl)-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-II)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.91 - 2.02 (m, 2H), 3.11 - 3.13 (m, 1H), 3.26 - 3.29 (m, 2H), 3.56 - 3.58 (m, 1H), 3.70 (s, 3H), 4.30 - 4.38 (m, 1H), 5.01 - 5.13 (d, 1H), 5.46 (s, 2H), 6.77 - 6.96 (m, 4H), 7.17 - 7.25 (m, 2H), 7.39 - 7.41 (d, <i>J</i> = 8.32 Hz, 1H), 7.94 - 7.97 (t, <i>J</i> = 7.08 Hz, 1H), 8.15 (s, 1H), 8.66 - 8.68 (bs, 3H), 9.27 - 9.30 (s, 3H); Mass (<i>m/z</i>): 400.2 (<i>M</i> + <i>H</i>) ⁺ .

37	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(4-methoxybenzyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-II)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.90 - 2.06 (m, 2H), 2.65 - 2.67 (m, 1H), 3.07 - 3.15 (m, 1H), 3.56 - 3.62 (m, 2H), 3.69 (s, 3H), 4.28 - 4.36 (m, 1H), 4.99 - 5.11 (m, 1H), 5.39 (s, 2H), 6.86 - 6.94 (m, 3H), 7.11 - 7.19 (m, 1H), 7.22 - 7.24 (d, J = 8.3 Hz, 2H), 7.40 - 7.43 (d, J = 8.2 Hz, 1H), 7.89 - 7.93 (t, 1H), 8.11 (s, 1H), 8.61 (m, 1H), 9.04 - 9.06 (m, 1H); Mass (m/z): 400.4 (M+H) ⁺ .
38	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(4-methoxybenzyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-I)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.92 - 1.99 (m, 2H), 3.11 - 3.16 (m, 2H), 3.27 - 3.37 (m, 2H), 3.70 (s, 3H), 4.30 - 4.39 (m, 1H), 5.01 - 5.13 (d, 1H), 5.40 (s, 2H), 6.87 - 6.90 (d, J = 8.57 Hz, 2H), 6.92 - 6.95 (m, 1H), 7.15 - 7.21 (m, 1H), 7.23 - 7.26 (d, J = 8.52 Hz, 2H), 7.41 - 7.44 (d, J = 8.24 Hz, 1H), 7.89 - 7.92 (m, 1H), 8.12 (m, 1H), 8.62 - 8.64 (m, 1H), 9.12 - 9.15 (m, 1H); Mass (m/z): 400.4 (M+H) ⁺ .
39	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(4-methoxybenzyl)-4,7-difluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-I)	¹ H NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.86 - 1.89 (m, 1H), 1.93 - 2.00 (m, 1H), 3.10 - 3.16 (m, 1H), 3.20 - 3.30 (m, 1H), 3.39 - 3.47 (m, 1H), 3.55 - 3.61 (m, 1H), 3.70 (3H, s), 4.28 - 4.36 (m, 1H), 5.00 - 5.11 (m, 1H), 5.46 (s, 2H), 6.85 - 6.90 (m, 3H), 6.98 - 7.03 (m, 1H), 7.15 - 7.17 (d, 2H), 8.08 - 8.12 (m, 2H), 8.63 - 8.65 (m, 1H), 9.18 (m, 1H); Mass (m/z): 418.2 (M+H) ⁺ .


40	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(4-methoxybenzyl)-4,7-difluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-II)	¹ H NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.86 - 1.99 (m, 2H), 3.10 - 3.19 (m, 1H), 3.20 - 3.30 (m, 1H), 3.38 - 3.45 (m, 1H), 3.56 - 3.59 (m, 1H), 3.70 (3H, s), 4.28 - 4.36 (m, 1H), 5.00 - 5.12 (m, 1H), 5.46 (s, 2H), 6.85 - 6.90 (m, 3H), 6.98 - 7.04 (m, 1H), 7.14 - 7.16 (d, 2H), 8.09 - 8.12 (m, 2H), 8.55 - 8.65 (m, 1H), 9.10 - 9.16 (m, 1H); Mass (m/z): 418.2 (M+H) ⁺ .
41	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(3,4-difluorobenzyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-I)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.89 - 2.00 (m, 2H), 3.08 - 3.17 (m, 2H), 3.27 - 3.30 (m, 2H), 4.29 - 4.38 (m, 1H), 5.01 - 5.13 (d, 1H), 5.49 (s, 2H), 6.92 - 6.97 (m, 1H), 7.12 (s, 1H), 7.17 - 7.22 (m, 1H), 7.37 - 7.43 (m, 3H), 7.94 - 7.98 (t, 1H), 8.17 (s, 1H), 8.63 - 8.66 (m, 1H), 9.17 - 9.20 (m, 1H); Mass (m/z): 406.3 (M+H) ⁺ .
42	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(1-methyl-1<i>H</i>-pyrazol-4-yl-methyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-I)	¹ H- NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.91 - 1.98 (m, 2H), 3.11 - 3.17 (m, 2H), 3.57 - 3.60 (m, 2H), 3.76 (s, 3H), 4.29 - 4.37 (m, 1H), 5.00 - 5.12 (d, 1H), 5.30 (s, 2H), 6.91 - 6.96 (m, 1H), 7.19 - 7.24 (m, 1H), 7.46 (s, 1H), 7.52 - 7.54 (d, J = 8.2 Hz, 1H), 7.29 (s, 1H), 7.86 - 7.89 (t, 1H), 8.06 (s, 1H), 8.59 - 8.62 (bs, 1H), 9.04 - 9.06 (bs, 1H); Mass (m/z): 374.3 (M+H) ⁺ .

43	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(1-methyl-1<i>H</i>-pyrazol-4-ylmethyl)-4,7-difluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-I)	¹ H-NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.86 - 1.98 (m, 2H), 3.08 - 3.15 (m, 2H), 3.27 - 3.34 (m, 2H), 3.75 (s, 3H), 4.24 - 4.34 (m, 1H), 4.99 - 5.11 (d, 1H), 5.36 (s, 2H), 6.85 - 6.90 (m, 1H), 7.02 - 7.08 (m, 1H), 7.41 (s, 1H), 7.67 (s, 1H), 8.02 - 8.05 (m, 2H), 8.62 (m, 1H), 9.04 - 9.07 (m, 1H); Mass (m/z): 392.1 (M+H) ⁺ .
44	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-I)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.85 - 2.01 (m, 2H), 3.08 - 3.16 (m, 2H), 3.27 - 3.30 (m, 2H), 4.31 - 4.38 (m, 1H), 5.01 - 5.13 (d, 1H), 5.64 (s, 2H), 6.94 - 6.99 (m, 2H), 7.08 - 7.09 (m, 1H), 7.17 - 7.22 (s, 1H), 7.34 - 7.36 (m, 1H), 8.01 - 8.04 (m, 1H), 8.17 - 8.19 (m, 2H), 8.67 - 8.69 (m, 1H), 9.28 - 9.30 (m, 1H); Mass (m/z): 389.2 (M+H) ⁺ .
45	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(benzothiazol-6-ylmethyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide hydrochloride (Isomer-I)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.85 - 2.02 (m, 2H), 3.12 - 3.17 (m, 1H), 3.27 - 3.35 (m, 2H), 3.55 - 3.63 (m, 1H), 4.31 - 4.39 (m, 1H), 5.02 - 5.13 (d, 1H), 5.66 (s, 2H), 6.91 - 6.96 (m, 1H), 7.16 - 7.21 (m, 1H), 7.44 - 7.46 (d, J = 8.3 Hz, 2H), 7.94 - 7.98 (t, 1H), 8.04 - 8.10 (m, 2H), 8.21 (s, 1H), 8.61 - 8.64 (m, 1H), 9.10 - 9.12 (m, 1H), 9.37 (s, 1H); Mass (m/z): 427.1 (M+H) ⁺ .

46	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(1-methyl-1H-indazole-3-ylmethyl)-4-fluoro-1H-indole-3-carboxamide hydrochloride (Isomer-I)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.82 - 1.95 (m, 1H), 1.98 - 2.05 (m, 1H), 3.08 - 3.16 (m, 1H), 3.26 - 3.30 (m, 1H), 3.56 - 3.62 (m, 2H), 4.01 (s, 3H), 4.28 - 4.37 (m, 1H), 5.00 - 5.12 (m, 1H), 5.82 (s, 2H), 6.90 - 6.95 (m, 1H), 7.09 - 7.13 (t, 1H), 7.18 - 7.23 (m, 1H), 7.36 - 7.40 (m, 1H), 7.54 - 7.61 (m, 2H), 7.66 - 7.68 (m, 1H), 7.89 - 7.93 (t, 1H), 8.19 (s, 1H), 8.60 - 8.62 (m, 1H), 9.08 - 9.10 (m, 1H); Mass (m/z): 424.2(M+H) ⁺ .
----	--	---

Example 47

***cis*-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1H-indole-3-carboxamide (Isomer-II)**






5




The compound of example 23 (0.033 g, 0.000074 moles) was dissolved in ice cold water (5.0 mL) and pH was adjusted to ~ 8.0 using 2M sodium bicarbonate solution (0.5 mL) at 5 – 10 °C. Aqueous layer was extracted with dichloromethane (5 mL x 3). The combined organic phase was washed with water (5 mL), brine solution (5 mL) and dried over Na₂SO₄. The organic phase was concentrated under vacuum to afford the title compound.




Yield: 0.030 g (99.3 %); ¹H - NMR (DMSO-*d*₆, 400 MHz) δ ppm: 1.62 - 1.68 (m, 2H), 2.56 (m, 1H), 2.67 - 2.81 (m, 2H), 2.93 - 2.96 (m, 1H), 3.08 - 3.14 (m, 1H), 4.05 - 4.14 (m, 1H), 4.60 - 4.72 (d, 1H), 5.67 (s, 2H), 6.87 - 6.94 (m, 2H), 6.97 - 7.03 (m, 2H), 7.87 - 7.91 (t, 1H), 8.16 - 8.19 (m, 2H); Mass (m/z): 407.2 (M+H)⁺.




15




Examples 48 to 71: The following compounds of example 48 to 71 can be prepared from the hydrochloride salt compounds of examples 22 to 46 by following the experimental procedure as described in the example 47.




Example No.	Chemical name and Structure	Characterization data
48	 <i>trans</i>-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1<i>H</i>-indole-3-carboxamide (Isomer-I)	¹ H - NMR (CDCl ₃ , 400 MHz) δ ppm: 2.23 - 2.27 (m, 1H), 2.77 - 2.90 (m, 3H), 2.81 - 2.98 (bs, 1H), 3.30 - 3.37 (m, 1H), 4.34 - 4.39 (m, 2H), 4.48 - 4.51 (m, 1H), 5.51 (s, 2H), 6.58 (s, 1H), 6.85 - 6.89 (m, 3H), 7.18 - 7.20 (m, 1H), 7.98 (s, 1H), 8.16 - 8.17 (d, J = 5.1 Hz, 1H); Mass (m/z): 407.2 (M+H) ⁺ .
49	 <i>trans</i>-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1<i>H</i>-indole-3-carboxamide (Isomer-II)	¹ H - NMR (CDCl ₃ , 400 MHz) δ ppm: 2.23 - 2.27 (m, 1H), 2.77 - 2.90 (m, 3H), 2.81 - 2.98 (bs, 1H), 3.30 - 3.38 (m, 1H), 4.34 - 4.40 (m, 2H), 4.49 - 4.52 (m, 1H), 5.51 (s, 2H), 6.58 (s, 1H), 6.83 - 6.88 (m, 3H), 7.18 - 7.22 (m, 1H), 7.98 (s, 1H), 8.16 - 8.17 (d, J = 5.1 Hz, 1H); Mass (m/z): 407.2 (M+H) ⁺ .
50	 <i>cis</i>-N-(3-Fluoropiperidin-	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.61 - 1.68 (m, 2H), 1.85 - 2.05 (m, 1H), 2.54 - 2.59 (m, 1H), 2.67 - 2.80 (m, 1H), 2.92 - 2.99 (m, 1H), 3.08 - 3.14 (m, 1H), 4.05 - 4.13 (m, 1H), 4.59 - 4.72 (m, 1H), 5.73 (s, 2H), 6.77 - 6.80 (m, 1H), 6.89 - 7.03 (m, 2H), 7.86 - 7.89 (m, 1H), 8.14 (s, 1H), 8.33 - 8.34




	4-yl)-1-(3-fluoropyridin-4-ylmethyl)-4,7-difluoro-1 <i>H</i> -indole-3-carboxamide (Isomer-I)	(d, J = 4.8 Hz 1H), 8.60 (s, 1H); Mass (m/z): 407.3 (M+H) ⁺ .
51	 <i>cis</i> -N-(3-Fluoropiperidin-4-yl)-1-(2-chloropyridin-4-ylmethyl)-4-fluoro-1 <i>H</i> -indole-3-carboxamide (Isomer-I)	¹ H - NMR (CDCl ₃ , 400 MHz) δ ppm: 1.72 - 1.83 (m, 1H), 1.89 - 1.92 (m, 1H), 2.74 - 2.82 (m, 1H), 2.83 - 2.90 (bs, 1H), 2.92 - 2.95 (d, J = 13.26 Hz, 1H), 3.15 - 3.19 (m, 1H), 3.35 - 3.41 (m, 1H), 4.30 - 4.38 (m, 1H), 4.69 - 4.81 (d, 1H), 5.34 (s, 2H), 6.88 - 6.97 (d, J = 4.99 Hz, 1H), 7.00 - 7.02 (m, 3H), 7.17 - 7.23 (m, 1H), 7.48 - 7.54 (m, 1H), 7.99 (s, 1H), 8.32 - 8.33 (d, J = 5.1 Hz, 1H); Mass (m/z): 405.2 (M+H) ⁺ .
52	 <i>cis</i> -N-(3-Fluoropiperidin-4-yl)-1-(2-chloropyridin-4-ylmethyl)-4,7-difluoro-1 <i>H</i> -indole-3-carboxamide (Racemate)	¹ H- NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.62 - 1.68 (m, 2H), 2.50 (bs, 1H), 2.93 - 2.96 (m, 2H), 3.08 - 3.11 (m, 2H), 4.05 - 4.14 (m, 1H), 4.60 - 4.72 (d, 1H), 5.64 (s, 2H), 6.89 - 6.92 (m, 1H), 6.94 - 7.05 (m, 2H), 7.24 (s, 1H), 7.88 - 7.91 (t, 1H), 8.16 (m, 1H), 8.35 - 8.36 (d, J = 5.0 Hz, 1H); Mass (m/z): 423.2 (M+H) ⁺ .
53	 <i>cis</i> -N-(3-Fluoropiperidin-4-yl)-1-(2-chloropyridin-4-ylmethyl)-4,7-difluoro-1 <i>H</i> -indole-3-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.62 - 1.71 (m, 2H), 1.93 - 2.00 (m, 1H), 2.55 - 2.58 (bs, 1H), 2.67 - 2.81 (m, 1H), 2.93 - 2.99 (m, 1H), 3.08 - 3.18 (m, 1H), 4.05 - 4.13 (m, 1H), 4.60 - 4.72 (m, 1H), 5.64 (s, 2H), 6.89 - 6.95 (m, 1H), 6.98 - 7.05 (m, 2H), 7.24 (s, 1H), 7.88 - 7.91 (m, 1H), 8.16 (s, 1H), 8.35 - 8.36 (m, 1H); Mass (m/z): 423.2




	ylmethyl)-4,7-difluoro-1 <i>H</i> -indole-3-carboxamide (Isomer-I)	(M+H) ⁺ .
54	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(4-fluorobenzyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide (Isomer-I)	¹ H- NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.61 - 1.65 (m, 2H), 2.56 (bs, 1H), 2.67 - 2.71 (m, 1H), 2.80 (m, 1H), 2.92 - 2.95 (m, 1H), 3.08 - 3.14 (m, 1H), 4.04 - 4.14 (m, 1H), 4.59 - 4.72 (d, 1H), 5.48 (s, 2H), 6.93 - 6.98 (m, 1H), 7.13 - 7.21 (m, 3H), 7.31 - 7.35 (m, 2H), 7.40 - 7.42 (d, J = 8.3 Hz, 1H), 7.64 - 7.68 (t, 1H), 8.17 (s, 1H); Mass (m/z): 388.3 (M+H) ⁺ .
55	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(4-fluorobenzyl)-4,7-difluoro-1<i>H</i>-indole-3-carboxamide (Isomer-I)	¹ H- NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.61 - 1.67 (m, 2H), 2.55 (bs, 1H), 2.73 (m, 1H), 2.89 - 2.92 (m, 1H), 3.11 - 3.15 (m, 2H), 4.04 - 4.12 (m, 1H), 4.59 - 4.72 (d, 1H), 5.53 (s, 2H), 6.86 - 6.92 (m, 1H), 6.97 - 7.03 (m, 1H), 7.14 - 7.18 (m, 2H), 7.21 - 7.24 (m, 2H), 7.80 - 7.84 (t, 1H), 8.16 (s, 1H); Mass (m/z): 406.3 (M+H) ⁺ .
56	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(4-fluorobenzyl)-4,7-difluoro-1<i>H</i>-indole-3-carboxamide (Isomer-II)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.63 - 1.89 (m, 2H), 2.56 (bs, 1H), 2.73 (m, 1H), 2.92 - 2.95 (m, 1H), 3.11 - 3.14 (m, 2H), 4.04 (m, 1H), 4.58 - 4.72 (d, 1H), 5.53 (s, 2H), 6.85 - 6.92 (m, 1H), 6.97 - 7.03 (m, 1H), 7.15 - 7.18 (m, 2H), 7.21 - 7.24 (m, 2H), 7.80 - 7.84 (t, 1H), 8.16 (s, 1H); Mass (m/z): 406.4 (M+H) ⁺ .

57	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(2-fluorobenzyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide (Isomer-I)	¹ H- NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.62 - 1.65 (m, 2H), 2.54 (bs, 1H), 2.67 - 2.71 (m, 1H), 2.77 - 2.80 (m, 1H), 2.92 - 2.95 (m, 2H), 3.08 - 3.14 (m, 1H), 4.04 - 4.14 (m, 1H), 4.59 - 4.72 (d, 1H), 5.59 (s, 2H), 6.94 - 6.99 (m, 1H), 7.14 - 7.26 (m, 3H), 7.34 - 7.42 (m, 2H), 7.64 - 7.68 (t, 1H), 8.09 (s, 1H); Mass (m/z): 388.3 (M+H) ⁺ .
58	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(4-chlorobenzyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide (Isomer-I)	¹ H- NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.61 - 1.62 (m, 2H), 2.58 (bs, 1H), 2.91 - 2.92 (m, 2H), 3.11 (m, 2H), 4.04 - 4.15 (m, 1H), 4.59 - 4.72 (d, 1H), 5.50 (s, 2H), 6.93 - 6.98 (m, 1H), 7.15 - 7.20 (m, 1H), 7.26 - 7.28 (d, J = 8.3 Hz, 2H), 7.36 - 7.40 (t, 3H), 7.65 - 7.69 (t, 1H), 8.18 (s, 1H); Mass (m/z): 404.2 (M+H) ⁺ .
59	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-4-fluoro-1-(3-methoxybenzyl)-1<i>H</i>-indole-3-carboxamide (Isomer-I)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.63 (m, 2H), 2.58 (bs, 1H), 2.67 (m, 1H), 2.81 (m, 1H), 2.92 - 2.95 (m, 1H), 3.11 (m, 1H), 3.70 (s, 3H), 4.05 (m, 1H), 4.59 - 4.72 (d, 1H), 5.45 (s, 2H), 6.77 - 6.79 (d, J = 7.3 Hz, 1H), 6.83 - 6.86 (m, 2H), 6.93 - 6.98 (m, 1H), 7.15 - 7.25 (m, 2H), 7.39 - 7.41 (d, J = 8.1 Hz, 1H), 7.63 - 7.67 (t, 1H), 8.16 (s, 1H); Mass (m/z): 400.3 (M+H) ⁺ .

60	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-4-fluoro-1-(3-methoxybenzyl)-1H-indole-3-carboxamide (Isomer-II)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.63 - 1.68 (m, 2H), 2.59 (bs, 1H), 2.82 (m, 1H), 2.92 - 2.99 (m, 1H), 3.09 - 3.15 (m, 2H), 3.70 (s, 3H), 4.02 - 4.05 (m, 1H), 4.59 - 4.71 (d, 1H), 5.45 (s, 2H), 6.76 - 6.95 (m, 4H), 7.16 - 7.25 (m, 2H), 7.39 - 7.41 (d, J = 8.3 Hz, 1H), 7.64 - 7.68 (t, 1H), 8.16 (s, 1H); Mass (m/z): 400.2 (M+H) ⁺ .
61	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(4-methoxybenzyl)-4-fluoro-1H-indole-3-carboxamide (Isomer-II)	¹ H - NMR (CDCl ₃ , 400 MHz) δ ppm: 1.72 - 1.80 (m, 1H), 1.85 - 1.90 (m, 1H), 2.72 - 2.80 (m, 1H), 2.82 - 2.91 (bs, 1H), 2.90 - 2.94 (d, 1H), 3.14 - 3.17 (m, 1H), 3.33 - 3.39 (m, 1H), 3.70 (s, 3H), 4.29 - 4.36 (m, 1H), 4.67 - 4.78 (d, 1H), 5.25 (s, 2H), 6.83 - 6.86 (d, J = 8.5 Hz, 2H), 6.93 - 6.96 (m, 1H), 7.09 - 7.11 (d, J = 8.4 Hz, 2H), 7.24 (m, 2H), 7.44 - 7.51 (m, 1H), 7.95 (s, 1H); Mass (m/z): 400.4 (M+H) ⁺
62	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(4-methoxybenzyl)-4-fluoro-1H-indole-3-carboxamide (Isomer-I)	¹ H - NMR (CDCl ₃ , 400 MHz) δ ppm: 1.72 - 1.80 (m, 1H), 1.87 - 1.91 (m, 1H), 2.72 - 2.80 (m, 1H), 2.82 - 2.91 (bs, 1H), 2.90 - 2.94 (d, 1H), 3.14 - 3.17 (m, 1H), 3.33 - 3.39 (m, 1H), 3.70 (s, 3H), 4.28 - 4.36 (m, 1H), 4.67 - 4.79 (d, 1H), 5.25 (s, 2H), 6.83 - 6.85 (d, J = 8.5 Hz, 2H), 6.92 - 6.96 (m, 1H), 7.09 - 7.11 (d, J = 8.4 Hz, 2H), 7.23 - 7.24 (m, 2H), 7.44 - 7.51 (m, 1H), 7.95 (s, 1H); Mass (m/z): 400.4 (M+H) ⁺ .

63	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(4-methoxybenzyl)-4,7-difluoro-1<i>H</i>-indole-3-carboxamide (Isomer-I)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.59 - 1.67 (m, 2H), 1.93 - 2.00 (m, 1H), 2.55 - 2.58 (m, 1H), 2.65 - 2.85 (m, 1H), 2.92 - 2.95 (m, 1H), 3.05 - 3.15 (m, 1H), 3.70 (s, 3H), 4.03 - 4.11 (m, 1H), 4.59 - 4.71 (m, 1H), 5.46 (s, 2H), 6.85 - 6.91 (m, 3H), 6.97 - 7.03 (m, 1H), 7.14 - 7.17 (d, <i>J</i> = 8.4 Hz, 2H), 7.75 - 7.79 (m, 1H), 8.13 (s, 1H); Mass (<i>m/z</i>): 418.3 (M+H) ⁺ .
64	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(4-methoxybenzyl)-4,7-difluoro-1<i>H</i>-indole-3-carboxamide (Isomer-II)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.59 - 1.67 (m, 2H), 1.93 - 2.00 (m, 1H), 2.55 - 2.58 (m, 1H), 2.65 - 2.85 (m, 1H), 2.92 - 2.95 (m, 1H), 3.07 - 3.13 (m, 1H), 3.70 (s, 3H), 4.03 - 4.11 (m, 1H), 4.59 - 4.71 (m, 1H), 5.46 (s, 2H), 6.88 - 6.90 (m, 3H), 6.97 - 7.03 (m, 1H), 7.15 - 7.17 (d, <i>J</i> = 8.3 Hz, 2H), 7.76 - 7.80 (m, 1H), 8.14 (s, 1H); Mass (<i>m/z</i>): 418.3 (M+H) ⁺ .
65	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(3,4-difluorobenzyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide (Isomer-I)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.63 - 1.66 (m, 2H), 2.62 (m, 1H), 2.67 - 2.71 (m, 1H), 2.77 - 2.81 (m, 1H), 2.92 - 2.95 (m, 1H), 3.08 - 3.15 (m, 1H), 4.05 - 4.12 (m, 1H), 4.59 - 4.72 (d, 1H), 5.48 (s, 2H), 6.93 - 6.98 (m, 1H), 7.12 (s, 1H), 7.16 - 7.22 (m, 1H), 7.36 - 7.45 (m, 3H), 7.66 - 7.71 (t, 1H), 8.19 (s, 1H); Mass (<i>m/z</i>): 406.3 (M+H) ⁺ .

66	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(1-methyl-1H-pyrazol-4-yl-methyl)-4-fluoro-1H-indole-3-carboxamide (Isomer-I)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.61 - 1.63 (m, 2H), 2.55 (bs, 1H), 2.56 - 2.58 (m, 1H), 2.67 - 2.68 (m, 1H), 2.92 - 2.95 (m, 1H), 3.12 (m, 1H), 3.75 (s, 3H), 4.04 (m, 1H), 4.59 - 4.71 (d, 1H), 5.29 (s, 2H), 6.93 - 6.98 (m, 1H), 7.18 - 7.23 (m, 1H), 7.46 (s, 1H), 7.51 - 7.53 (d, J = 8.2 Hz, 1H), 7.56 - 7.61 (t, 1H), 7.71 (s, 1H), 8.07 (s, 1H); Mass (m/z): 374.4 (M+H) ⁺ .
67	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(1-methyl-1H-pyrazol-4-yl-methyl)-4,7-difluoro-1H-indole-3-carboxamide (Isomer-I)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.61 - 1.63 (m, 2H), 2.55 (bs, 1H), 2.67 - 2.70 (m, 1H), 2.76 - 2.80 (m, 1H), 2.92 - 2.95 (m, 1H), 3.08 - 3.14 (m, 1H), 3.76 (s, 3H), 4.03 - 4.10 (m, 1H), 4.58 - 4.70 (d, 1H), 5.36 (s, 2H), 6.86 - 6.91 (m, 1H), 7.01 - 7.07 (m, 1H), 7.41 (s, 1H), 7.66 (s, 1H), 7.72 - 7.74 (t, 1H), 8.07 (s, 1H); Mass (m/z): 392.2 (M+H) ⁺ .
68	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4-fluoro-1H-indole-3-carboxamide	¹ H - NMR (CDCl ₃ , 400 MHz) δ ppm: 1.72 - 1.78 (m, 1H), 1.89 - 1.97 (m, 1H), 2.73 - 2.81 (m, 1H), 2.83 - 2.91 (bs, 1H), 2.91 - 2.95 (d, 1H), 3.11 - 3.18 (m, 1H), 3.34 - 3.41 (m, 1H), 4.30 - 4.37 (m, 1H), 4.68 - 4.81 (d, 1H), 5.17 (s, 2H), 6.88 - 6.98 (m, 2H), 7.00 - 7.05 (m, 1H), 7.20 - 7.22 (m, 3H), 7.45 - 7.52 (m, 1H), 7.97 (s, 1H); Mass (m/z): 389.3 (M+H) ⁺ .

	(Isomer-I)	
69	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(benzothiazol-6-ylmethyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide (Isomer-I)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.63 - 1.68 (m, 2H), 2.57 (bs, 1H), 2.80 (m, 1H), 2.92 - 2.96 (m, 1H), 3.08 - 3.15 (m, 2H), 4.04 - 4.15 (m, 1H), 4.57 - 4.71 (d, 1H), 5.46 (s, 2H), 6.91 - 6.95 (m, 1H), 7.15 - 7.20 (m, 1H), 7.44 - 7.46 (d, <i>J</i> = 8.3 Hz, 2H), 7.64 (t, 1H), 8.04 - 8.08 (m, 2H), 8.21 (s, 1H), 8.61 - 8.63 (m, 1H); Mass (<i>m/z</i>): 427.1 (M+H) ⁺ .
70	 <i>Cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(1-methyl-1<i>H</i>-indazole-3-ylmethyl)-4-fluoro-1<i>H</i>-indole-3-carboxamide (Isomer-I)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm: 1.45 - 1.47 (m, 1H), 1.95 - 1.98 (m, 1H), 2.44 - 2.49 (m, 1H), 2.54 - 2.59 (m, 2H), 2.83 - 2.86 (m, 1H), 3.16 - 3.20 (m, 1H), 3.83 (s, 3H), 4.10 - 4.12 (m, 1H), 4.35 - 4.49 (m, 1H), 5.56 (s, 2H), 7.26 - 7.37 (m, 3H), 7.43 - 7.46 (m, 1H), 7.88 (s, 1H), 8.08 - 8.10 (m, 1H), 8.16 (s, 1H), 8.50 - 8.52 (m, 1H), 8.84 - 8.86 (m, 1H); Mass (<i>m/z</i>): 424.2 (M+H) ⁺ .
71	 <i>cis</i>-N-(3-Fluoropiperidin-4-yl)-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1<i>H</i>-indole-3-carboxamide (Isomer-I)	¹ H - NMR (DMSO- <i>d</i> ₆ , 400MHz) δ ppm: 1.62 - 1.68 (m, 1H), 2.51 - 2.56 (m, 2H), 2.67 - 2.71 (m, 1H), 2.93 - 2.96 (m, 1H), 3.08 - 3.14 (m, 1H), 4.05 - 4.13 (m, 1H), 4.05 - 4.13 (m, 1H), 4.60 - 4.72 (d, 1H), 5.67 (s, 2H), 6.87 - 6.94 (m, 2H), 6.97 - 7.03 (m, 2H), 7.87 - 7.91 (t, 1H), 8.16 (s, 1H), 8.18 - 8.19 (d, <i>J</i> = 5.1 Hz, 1H); Mass (<i>m/z</i>): 407.2 M+H) ⁺ .

Biological Data**Example 72:****Determination of allosteric potency EC₅₀ values for Muscarinic M1 receptor:**

- A stable CHO cell line expressing recombinant human Muscarinic M1 receptor and pCRE-Luc reporter system was used for cell-based assay. The assay offers a non-radioactive based approach to determine binding of a compound to GPCRs. In this specific assay, the level of intracellular cyclic AMP which is modulated by activation or inhibition of the receptor is measured. The recombinant cells harbor luciferase reporter gene under the control of cAMP response element.
- The above cells were grown in 96 well clear bottom white plates in Hams F12 medium containing 10% fetal bovine serum (FBS). Prior to the addition of compounds or standard agonist, cells were serum starved overnight. Increasing concentrations of test compounds were added along with EC₂₀ of acetylcholine in OptiMEM medium to the cells. The incubation was continued at 37 °C in CO₂ incubator for 4 hours. Medium was removed and cells were washed with phosphate buffered saline. The cells were lysed and luciferase activity was measured in a Luminometer. Luminescence units were plotted against the compound concentrations using Graphpad software. EC₅₀ values of the compounds were defined as the concentration required in stimulating the luciferase activity by 50 % in presence of EC₂₀ of acetylcholine and the results are provided in table 1.

Table 1:

Ex.	hM1 PAM EC ₅₀ (nM)	Ex.	hM1 PAM EC ₅₀ (nM)
1	1264	25	1441
2	681.1	26	1105
3	1637	27	1689
4	1492	28	1568
3	370.5	29	2920
6	122.5	30	1081
7	1360	31	1608
8	192	32	1586
9	1417	33	1374

10	940	34	1387
11	1534	35	1482
12	432.5	36	1349
13	1378	37	1985
18	160.2	38	1448
19	744	39	1485
22	1495	44	1306
23	1074	45	1163
24	1390		

Example 73:**Protein binding Assay**

Unbound fractions of new chemical entities in plasma, brain homogenate and liver microsomes were determined using high-throughput dialysis (HT dialysis). Briefly, dialysis membranes were soaked in deionized water for 20 minutes and then in deionized water with 30% ethanol for 15 minutes and finally in phosphate buffer until use. Membranes were rinsed in phosphate buffer before assembling. The membranes were layered between teflon bars of dialysis assembly. Stock solutions of test compounds were prepared at 10 mM in DMSO, diluted to 1 mM in acetonitrile and further diluted to 100 μ M in mixture of water and acetonitrile (1:1 v/v). Human plasma (pool of 3) was prepared from human blood (3 donors) by centrifuging at 4000 rpm for 10 min at 4 °C. Rat and dog blood were obtained on the day of the study and centrifuged to obtain plasma. Rat brains were isolated, cleaned and homogenized with 2 volumes of buffer (3 fold dilution). Liver microsomes were prepared at 0.5 mg/mL in phosphate buffer (100 mM, pH 7.4). The dialysate chambers were loaded with 150 μ L of 100 mM phosphate buffer (pH 7.4) in triplicates. The matrix chambers were loaded with 150 μ L of the plasma or brain homogenate or microsomal suspension spiked with test compounds at a final concentration of 1 μ M. 50 μ L of the sample was removed from both the chambers at 0 h. The plate was sealed and incubated at 37 °C for 6 h at 100 rpm. After 6 h, 50 μ L of the sample was removed from both the chambers. Equal volumes of buffer or human plasma / microsomal suspension were added to the plasma / microsomal and buffer samples respectively to

create identical sample matrices for analysis. The samples were precipitated with 150 μ L of acetonitrile containing fluoxetine as an internal standard. All the samples were centrifuged at 10000 rpm for 10 minutes at 4 °C. The supernatants were analyzed by LC-MS/MS and the results are provided in table 2.

5 Table-2:

Ex.	Species	Fu (Mean $\pm$ SEM, n=3)		
		Plasma	Brain	Microsomes
1	Rat	0.267 $\pm$ 0.03	0.166 $\pm$ 0.01	0.836 $\pm$ 0.04
	Dog	0.217 $\pm$ 0.21	NA	0.982 $\pm$ 0.01
	Human	0.197 $\pm$ 0.01	NA	1.046 $\pm$ 0.07
2	Rat	0.068 $\pm$ 0.002	0.015 $\pm$ 0.001	0.679 $\pm$ 0.081
	Dog	0.087 $\pm$ 0.008	NA	0.576 $\pm$ 0.079
	Human	0.040 $\pm$ 0.002	NA	0.459 $\pm$ 0.046
5	Rat	0.210 $\pm$ 0.01	0.128 $\pm$ 0.01	0.779 $\pm$ 0.01
	Dog	0.163 $\pm$ 0.02	NA	0.746 $\pm$ 0.004
	Human	0.077 $\pm$ 0.01	NA	0.899 $\pm$ 0.03
6	Rat	0.050 $\pm$ 0.008	0.014 $\pm$ 0.001	0.560 $\pm$ 0.048
	Dog	0.043 $\pm$ 0.002	NA	0.713 $\pm$ 0.041
	Human	0.030 $\pm$ 0.003	NA	0.546 $\pm$ 0.018
7	Rat	0.20 $\pm$ 0.02	0.06 $\pm$ 0.004	0.80 $\pm$ 0.04
	Dog	0.20 $\pm$ 0.01	NA	0.60 $\pm$ 0.03
	Human	0.10 $\pm$ 0.004	NA	0.70 $\pm$ 0.03
8	Rat	0.089 $\pm$ 0.01	0.0348 $\pm$ 0.0001	0.606 $\pm$ 0.008
	Dog	0.064 $\pm$ 0.01	NA	0.679 $\pm$ 0.066
	Human	0.079 $\pm$ 0.006	NA	0.626 $\pm$ 0.037
22	Rat	0.315 $\pm$ 0.03	0.067 $\pm$ 0.002	0.902 $\pm$ 0.006
	Dog	0.310 $\pm$ 0.01	NA	0.866 $\pm$ 0.034
	Human	0.309 $\pm$ 0.008	NA	0.788 $\pm$ 0.031

Example 74:**Rodent Pharmacokinetic Study**

Male Wistar rats (260 ± 50 grams) were used as experimental animals. Animals were housed individually in polypropylene cage. Two days prior to study, male Wistar rats were anesthetized with isoflurane for surgical placement of jugular vein catheter. Rats were randomly divided for oral (3 mg/kg) and intravenous (*i.v.*) (1 mg/kg) dosing ($n = 3/\text{group}$) and fasted overnight before oral dosing (*p.o.*). However, rats allocated to intravenous dosing food and water was provided *ad libitum*.

At pre-determined point, blood was collected through jugular vein and replenished with an equivalent volume of normal saline. Collected blood was transferred into a labeled eppendorf tube containing 10 μL of heparin as an anticoagulant. Typically blood samples were collected at following time points: 0.08, 0.25, 0.5, 1, 2, 4, 6, 8, and 24 hours post dose. Blood was centrifuged at 4000 rpm for 10 minutes. Plasma was separated and stored frozen at -80°C until analysis. The concentrations of the test compounds were quantified in plasma by qualified LC-MS/MS method using suitable extraction technique. The test compounds were quantified in the calibration range around 1-1000 ng/mL in plasma. Study samples were analyzed using calibration samples in the batch and quality control samples spread across the batch.

Pharmacokinetic parameters C_{max} , AUC_t , $T_{1/2}$, Clearance, and Bioavailability (%F) were calculated by non-compartmental model using standard non-compartmental model by using Phoenix WinNonlin 6.0.2 or 6.0.3 version Software package and the results are tabulated in table 3.

Table-3:

Ex.	RO A	Vehicle	C_{max} (ng/mL)	AUC_{0-t} (ng.hr/mL)	$T_{1/2}$ (hr)	Clearance (mL/min/kg)	% F
1	oral	0.25 % Tween 80 + 99.75 % HEC solution	1187 ± 81	5163 ± 464	2.9 ± 0.2	-	76 ± 7

	<i>i.v</i>	5 % Pharmaso lve + 45 % Propylen e glycol + 50 % PEG 400	-	2250 ± 190	3.0±0.3	7.3 ± 0.4	
2	oral	0.25 % Tween 80 + 99.75 % HEC solution	632 ± 167	2161 ± 599	1.9±0.2	-	63 ± 17
	<i>i.v</i>	5 % Pharmaso lve + 45 % Propylen e glycol + 50 % PEG 400	-	1146 ± 245	1.9±0.4	16.1 ± 2.8	
5	oral	0.25 % Tween 80 + 99.75 % HEC solution	2681±66 4	7867±2857	3.3±0.3	-	90 ± 33
	<i>i.v</i>	5 % Pharmaso lve + 45 % Propylen e glycol	-	2921±1432	4.0±1.7	7.1 ± 4.3	

		+ 50 % PEG 400					
7	oral	0.25 % Tween 80 + 99.75 % HEC solution	783 ± 254	2261 ± 471	1.4±0.2	-	67 ± 14
	i.v	5 % Pharmaso lve + 45 % Propylen e glycol + 50 % PEG 400	-	1119 ± 119	2.0±0.4	14.2 ± 2.0	
8	oral	0.25 % Tween 80 + 99.75 % HEC solution	578 ± 170	2497 ± 842	2.3±1.8	-	45 ± 15
	i.v	5 % Pharmaso lve + 45 % Propylen e glycol + 50 % PEG 400	-	1834 ± 562	1.8±0.3	9 ± 3	
12	oral	0.25 % Tween 80 + 99.75	337 ± 141	1140 ± 114	3.0±3.0	-	62 ± 6

		% HEC solution					
	i.v	5 % Pharmaso lve + 45 % Propylene glycol + 50 % PEG 400	-	609 ± 62	1.6±0.5	26.7 ± 3.4	
22	oral	Reagent grade water	207 ± 76	1647 ± 147	4.8±2.8	-	60 ± 5
	i.v	Water for injection	-	908 ± 198	7.6±1.7	17.1 ± 3.3	

ROA – Route of Administration

Example 75:**Rodent Brain Penetration Study**

5 Male Wistar rats (260 ± 40 grams) were used as experimental animals. Three animals were housed in each cage. Animals were given water and food *ad libitum* throughout the experiment and maintained on a 12 hours light/dark cycle.

10 Brain penetration was determined in discrete manner in rats. One day prior to dosing day, male Wistar rats were acclimatized and randomly grouped according to their weight. At each time point (0.5, 1 and 2 hours) $n = 3$ animals were used.

The test compounds were suitably preformulated and administered orally at (free base equivalent) 3 mg/kg. Blood samples were removed via cardiac puncture by using isoflurane anesthesia. The animals were sacrificed to collect brain tissue. Plasma was separated and brain samples were homogenized and stored frozen at -20 °C until analysis. The concentrations of the test compounds in plasma and brain were determined using LC-MS/MS method.

15

The test compounds were quantified in plasma and brain homogenate by qualified LC-MS/MS method using suitable extraction technique. The test

compounds were quantified in the calibration range of 1-500 ng/mL in plasma and brain homogenate. Study samples were analyzed using calibration samples in the batch and quality control samples spread across the batch. Extent of brain-plasma ratio was calculated (Cb/Cp) and the results are tabulated in table 4.

5 Table 4:

Ex.	Single dose Rat Brain Penetration (Cb/Cp) at 3 mg/kg, p.o.
1	0.30 ± 0.05
2	0.95 ± 0.17
5	0.50 ± 0.04
6	2.09 ± 0.46
7	0.61 ± 0.01
8	1.1 ± 0.1
10	2.40 ± 0.30
11	1.59 ± 0.35
12	1.64 ± 0.12
22	0.46 ± 0.11
30	1.05 ± 0.53
35	0.72 ± 0.03
38	1.03 ± 0.17

Example 76:

Object Recognition Task Model

10 The cognition enhancing properties of compounds of this invention were estimated by using this model.

Male Wistar rats (8- 10 weeks old) were used as experimental animals. Four animals were housed in each cage. Animals were kept on 20 % food deprivation from a day prior to experimentation. Water was provided ad libitum throughout the experiment. Animals were maintained on a 12 hours light/dark cycle in temperature and humidity controlled room. The experiment was carried out in an open field made up of acrylic. Rats were habituated to individual arenas (open field) for 1 hour in the absence of any objects on day 1.

One group of 12 rats received vehicle and another set of animals received compound of the formula (I), before the familiar (T_1) and choice (T_2) trials. During the familiarization phase, (T_1), the rats were placed individually in the arena for 3 minutes, in which two identical objects (a_1 and a_2) were positioned 10 cm from the wall. 24 hours after T_1 , trial for long-term memory test was performed. The same rats were placed in the same arena as they were placed in T_1 trial. During the choice phase (T_2) rats were allowed to explore the arena for 3 minutes in presence of a copy of familiar object (a_3) and one novel object (b). During the T_1 and T_2 trial, explorations of each object (defined as sniffing, licking, chewing or having moving vibrissae whilst directing the nose towards the object at a distance of less than 1 cm) were recorded using stopwatch.

T_1 is the total time spent exploring the familiar objects ($a_1 + a_2$).

T_2 is the total time spent exploring the familiar object and novel object ($a_3 + b$).

Discriminative index = Time spent with novel object / (time spent with novel and familiar object).

The object recognition test was performed as described by *Behavioural Brain Research*, 1988, 31, 47-59 and the results are provided in table 5.

Table 5:

Ex.	Dose mg/kg, p.o.	Exploration time mean $\pm$ S.E.M (sec)		Inference
		Familiar object	Novel object	
1	0.3	11.31 $\pm$ 1.07	28.92 $\pm$ 3.54	Active
2	1	9.06 $\pm$ 1.75	15.09 $\pm$ 2.19	Active
5	0.3	11.00 $\pm$ 0.81	19.64 $\pm$ 1.74	Active
6	3	15.75 $\pm$ 1.71	23.67 $\pm$ 2.62	Active
7	1	9.16 $\pm$ 0.83	19.56 $\pm$ 1.96	Active
8	0.3	7.59 $\pm$ 1.02	14.36 $\pm$ 1.56	Active
22	3	8.21 $\pm$ 0.70	15.72 $\pm$ 2.25	Active

20 Example 77:

Object Recognition Task Model – Scopolamine challenge

The cognition enhancing properties of compounds of this invention were estimated by using this model.

Male Wistar rats (8- 10 weeks old) were used as experimental animals. Four animals were housed in each cage. Animals were kept on 20 % food deprivation from a day prior to experimentation. Water was provided ad libitum throughout the experiment. Animals were maintained on a 12 hours light/dark cycle in temperature and humidity controlled room. The experiment was carried out in an open field made up of acrylic. Rats were habituated to individual arenas (open field) for 1 hour in the absence of any objects on day 1.

Rats received vehicle or vehicle and scopolamine or compound of the formula (I) and scopolamine, before the familiar (T_1). During the familiarization phase, (T_1), the rats were placed individually in the arena for 3 minutes, in which two identical objects (a_1 and a_2) were positioned 10 cm from the wall. 3 minutes after T_1 , trial for memory test was performed. The same rats were placed in the same arena as they were placed in T_1 trial. During the choice phase (T_2) rats were allowed to explore the arena for 3 minutes in presence of a copy of familiar object (a_3) and one novel object (b). During the T_1 and T_2 trial, explorations of each object (defined as sniffing, licking, chewing or having moving vibrissae whilst directing the nose towards the object at a distance of less than 1 cm) were recorded using stopwatch.

T_1 is the total time spent exploring the familiar objects ($a_1 + a_2$).

T_2 is the total time spent exploring the familiar object and novel object ($a_3 + b$).

Discriminative index = Time spent with novel object / (time spent with novel and familiar object).

Table 6:

Ex.	Dose mg/kg, p.o.	Exploration time mean $\pm$ S.E.M (sec)		Inference
		Familiar object	Novel object	
1	0.3	7.54 $\pm$ 1.93	13.07 $\pm$ 1.79	Active
2	0.3	14.29 $\pm$ 1.92	20.93 $\pm$ 2.89	Active
5	1	13.61 $\pm$ 1.79	25.72 $\pm$ 3.67	Active
22	1	21.34 $\pm$ 3.65	29.70 $\pm$ 1.72	Active

Example 78:**Contextual fear conditioning task**

Experiment was carried out over a period of two days. On day 1, rats were placed in the operant behavior chamber and allowed to acclimatize for 2 minutes.

- 5 Rats received an unavoidable foot shock (unconditioned stimulus (US): electric shock of 0.5 - 0.7 mA for 3 seconds). Following a 1 minute interval, shocks were repeated to deliver a total of three US. Rats were administered Vehicle or test compound post training. Scopolamine (0.3 mg/kg, s.c.) was administered 120 minutes after training.

- 10 On day 2, rats were placed in the operant behavior chamber and total freezing time was scored for a period of 5 minutes. The results are provided in the figure 1.

Example 79:**Estimation of cortical sAPP α levels in rat brain****Experimental Procedure:**

- 15 Male Wistar rats (250 ± 45 grams) were randomly divided into different treatment groups (n=5 per group). Control group of rats were administered with intraperitoneal (*i.p.*) injection of vehicle (99.75 % of 0.25 % hydroxyethylcellulose HHX + 0.25 % tween 80 for examples 8 and 22; 5 % Pharmasolve + 45 % propylene glycol + 50 % polyethylene glycol-400 for example 1). Rats in treatment groups were
- 20 allocated to one dose of test compound and administered with a single intraperitoneal injection of test compound (dose volume of 2 mL/kg). Rats were sacrificed by cervical dislocation at 60 minutes after treatment. Brains were quickly isolated and the cortex was dissected out at -20 °C. The cortex was immediately kept on a dry ice and weighed before being stored at -80 °C until quantification of sAPP α using
- 25 Enzyme-linked immunosorbent assay (ELISA).

Sample Preparation:

1. Protease inhibitor cocktail tablets (complete mini, Make- Roche; 1 tablet for 8 mL) were added to the Tris Buffer Saline (TBS) prior to using the buffer for tissue processing.
- 30 2. Cortical tissues were thawed and homogenized in five volumes of TBS and the solution was centrifuged at 15,000 rpm at 4 °C for 90 minutes.

3. The supernatant was discarded and homogenized in five volumes of TBS. Samples were centrifuged at 15,000 rpm at 4 °C for 30 minutes.
4. Supernatant was discarded and precipitate was sonicated in ten volumes of 50 mM Tris buffer (pH: 7.6) containing 6 M of Guanidine-HCl. Sonication was repeated four
- 5 times with duration of five seconds every time.
5. Resulting mixture was incubated at the room temperature for 30 minutes and centrifuged at 15,000 rpm at 4°C for 30 minutes. Supernatant was diluted 100 times with EIA buffer prior to addition in the pre-coated ELISA plates.

Measurement of sAPP α by ELISA Kit:

- 10 To investigate the role of an acute treatment of test compound on sAPP α levels, the expression of this protein was measured in samples obtained from brain homogenates of treated and untreated rats by employing ELISA assay. The entire procedure was followed as described in the ELISA kit manual (Mouse/Rat sAPP α (highly sensitive) assay kit, Catalog Number: JP27419, Immuno-Biological
- 15 Laboratories Co. Ltd, Hamburg, Germany).

Statistical analysis:

- Statistical analyses were performed using the Graph Pad Prism (Version 4). Results are expressed as Mean $\pm$ SEM levels of sAPP α expressed as percentage of control values obtained from rats treated with vehicle. Statistical significance after
- 20 treatment was assessed using One-Way ANOVA followed by Dunnett's post test and the significance level was set below p value less than 0.05 and the results are tabulated in table 7.

Table 7: Effect of test compounds on cortical sAPP α levels in male Wistar rats.

Ex.	sAPP α levels (% of vehicle)			Inference
	1.0 mg/kg, i.p.	3.0 mg/kg, i.p.	10.0 mg/kg, i.p.	
1	155 $\pm$ 6**	153 $\pm$ 7**	142 $\pm$ 6**	Active
8	120 $\pm$ 17	137 $\pm$ 14	173 $\pm$ 10**	Active
22	143 $\pm$ 14	164 $\pm$ 12**	176 $\pm$ 10**	Active

Values are mean $\pm$ SEM (n=5/ group). ** p <0.01 Vs Vehicle (Dunnett's post test).

Example 80:**Modulation of cerebral blood flow in frontal cortex:**

The effect of test compound on modulation of cerebral blood flow was evaluated using rats.

5 Rats were acclimatized to the laboratory environment for at least 7 days. Rats (300 – 350 grams) were housed in a group of four in a controlled environment (Temp = 21 ± 3 °C; Humidity = 30-70 %) and maintained on a 12-hour light/dark cycle with lights on at 07:00 AM. Food and water was provided ad libitum.

10 Rats were anaesthetized with 12% urethane (i.p.). Animal's body core temperature was maintained at 37 °C via heating pad and rectal temperature probe. A small incision was made at one of the ventral side of the hind limb and the femoral vein was cannulated with PE10 tubing for drug application. Then animal was placed into a stereotaxic frame and a midline incision was made to expose the skull. A burr hole was drilled over the frontal cortex (stereotaxic coordinates 1 mm anterior and 4
15 mm lateral to bregma). Oxygen was supplied through the nose cone of the stereotaxic apparatus which was connected to the controlled gaseous supplier with a flow of 200 mL/minute. Laser Doppler probe (AD Instruments Inc) was placed over the hole to monitor cerebral blood flow. The Laser Doppler probe was connected to a computerized data acquisition system (PowerLab 16/30, AD Instruments Inc).
20 Vehicle or test compound was administered intravenously after cerebral blood flow was stable for 30 minutes. The cerebral blood flow was collected for further 90 minutes. Data obtained was calculated as percent increase relative to resting basal blood flow level. Test compound data was compared with the control group using one-way ANOVA followed by the Bonferroni post test.

25 **Reference:**


Psychopharmacology (Berl). 2013, 225, 21 - 30.

Result:

Example 1 significantly increased the cerebral blood flow as shown in figure 2.


We Claim

1. A fluoroindole compound of formula (I),




wherein:

R¹ is



R² is



wherein * represents point of attachment;

R³ is fluorine or hydrogen;

R⁴ is halogen, S-CH₃ or hydrogen;

R⁵ is -CH₃, -CH₂CH₂F or hydrogen;


a is 1 or 2; and

b is 1 or 2;

or an isotopic form, a stereoisomer or a pharmaceutically acceptable salt thereof.


2. The compound as claimed in claim 1, wherein:

R¹ is




; and

R² is




3. The compound as claimed in claim 1, wherein:

R¹ is



; and

R² is



4. The compound as claimed in any one of the claims 1 to 3, wherein the compound is selected from the group consisting of:

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(1-methyl-1*H*-pyrazol-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2-Chloropyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;

N-[(1*R*,2*R*)-2-Hydroxycyclohexyl]-1-(1-methyl-1*H*-pyrazol-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(1-methyl-1*H*-indazol-3-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(1-methyl-1*H*-pyrazol-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2-Chloropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2-fluoropyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2-fluoropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(3-fluoropyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(5-fluoropyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2,5-difluoropyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2,5-difluoro-pyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2,3-difluoropyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(1-pyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(1*H*-pyrazol-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2-bromothiazol-5-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(1-ethyl-5-methyl-1*H*-pyrazol-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(benzothiazol-6-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl] 1-(1-(2-fluoroethyl)-1*H*-pyrazol-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2-methylsulfanyl-pyridin-4-ylmethyl)-4,7-difluoro-1*H*-indole-3-carboxamide; and

N-[(1*S*,2*S*)-2-Hydroxycyclohexyl]-1-(2-methylsulfanyl-pyridin-4-ylmethyl)-4-fluoro-1*H*-indole-3-carboxamide;

or a pharmaceutically acceptable salt thereof.

5. A pharmaceutical composition comprising a compound as claimed in any one of the claims 1 to 4, and pharmaceutically acceptable excipients or carriers.

6. The pharmaceutical composition as claimed in claim 5, for use in the treatment of clinical conditions mediated through muscarinic M1 receptor selected from the group consisting of Alzheimer's disease, schizophrenia, cognitive disorders, pain or sleep disorders.

7. Use of the compound as claimed in any one of the claims 1 to 4, in the manufacture of medicament for the treatment of disorder related to Muscarinic M1 receptor.

8. The use of the compound as claimed in the claim 7, wherein the disorder related to muscarinic M1 receptor is selected from the group consisting of Alzheimer's disease, schizophrenia, cognitive disorders, pain or sleep disorders.

9. A compound as claimed in any one of the claims 1 to 4, for use in positive allosteric modulation of muscarinic M1 receptor.

10. The compound as claimed in claim 9, for use in the treatment of a disorder related to muscarinic M1 receptor selected from Alzheimer's disease, schizophrenia, cognitive disorders, pain or sleep disorders.