

ANTIBODIES TO OPGL

[001] This application claims priority to U.S. Provisional Application Serial No. 60/301,172, filed June 26, 2001, which is incorporated herein by reference for any purpose.

FIELD OF THE INVENTION

[002] The present invention relates to antibodies that bind osteoprotegerin ligand (OPGL). Compositions and methods for the treatment of bone diseases, such as osteoporosis, bone loss from arthritis, Paget's disease, and osteopenia, are also described.

BACKGROUND OF THE INVENTION

[003] Bone tissue provides support for the body and includes mineral (including calcium and phosphorous), a matrix of collagenous and noncollagenous proteins, and cells. Living bone tissue exhibits a dynamic equilibrium between formation of bone, which is called deposition, and breakdown of bone, which is called resorption. Three types of cells found in bone, osteocytes, osteoblasts and osteoclasts, are involved in this equilibrium. Osteoblasts promote formation of bone tissue whereas osteoclasts are associated with resorption. Resorption, or the dissolution of bone matrix and mineral, is a fast and efficient process compared to bone formation and can release large amounts of mineral from bone. Osteoclasts are involved in the regulation of the normal remodeling of skeletal tissue and in resorption induced

by hormones. For instance, resorption is stimulated by the secretion of parathyroid hormone in response to decreasing concentrations of calcium ion in extracellular fluids. In contrast, inhibition of resorption is a function of calcitonin. In addition, metabolites of vitamin D alter the responsiveness of bone to parathyroid hormone and calcitonin.

[004] Osteoprotegerin ligand (OPGL), which is a member of the TNF family of cytokines, promotes formation of osteoclasts through binding to the receptor activator of $\text{NF-}\kappa\text{B}$ (RANK, also called osteoclast differentiation and activation receptor, or ODAR). Osteoprotegerin (OPG), on the other hand, inhibits the formation of osteoclasts by sequestering OPGL and preventing OPGL association with ODAR. Thus, the amount of OPGL associated with ODAR correlates with the equilibrium between bone deposition and resorption.

[005] After skeletal maturity, the amount of bone in the skeleton reflects the balance (or imbalance) of bone formation and bone resorption. Peak bone mass occurs after skeletal maturity prior to the fourth decade. Between the fourth and fifth decades, the equilibrium shifts and bone resorption dominates. The inevitable decrease in bone mass with advancing years starts earlier in females than males and is distinctly accelerated after menopause in some females (principally those of Caucasian and Asian descent).

[006] Osteopenia is a condition relating generally to any decrease in bone mass to below normal levels. Such a condition may arise from a decrease in the rate of bone synthesis or an increase in the rate of bone destruction or both. A common form of osteopenia is primary osteoporosis, also

referred to as postmenopausal and senile osteoporosis. This form of osteoporosis is a consequence of the universal loss of bone with age and is often a result of increase in bone resorption with a normal rate of bone formation. Many white females in the United States develop symptomatic osteoporosis. A direct relationship exists between osteoporosis and the incidence of hip, femoral, neck and inter-trochanteric fracture in women 45 years and older. Elderly males may develop symptomatic osteoporosis between the ages of 50 and 70. Osteoporosis may, in certain instances, result from increased levels or activity of OPGL. Thus, it would be useful to have molecules that can regulate the activity of OPGL in osteoclastogenesis.

[007] Several factors have been identified which may contribute to postmenopausal and senile osteoporosis. They include alteration in hormone levels accompanying aging and inadequate calcium consumption attributed to decreased intestinal absorption of calcium and other minerals. Certain treatments have included hormone therapy or dietary supplements in an attempt to retard the process. More recently, anti-resorptive agents such as bisphosphonates and selective estrogen receptor modifiers (SERMs) have emerged for the prevention and treatment of reduced bone mass. Thus, it may be useful to combine those treatments with molecules that can regulate the activity of OPGL in treating certain osteopenic disorders.

SUMMARY OF THE INVENTION

[008] In certain embodiments, the invention provides for an antibody, comprising a heavy chain and a light chain, wherein the heavy chain comprises an amino acid sequence as set forth in SEQ ID NO: 2 or a fragment thereof, and the light chain comprises an amino acid sequence as set forth in SEQ ID NO: 4 or a fragment thereof.

[009] In certain embodiments, the invention provides for an antibody, comprising a heavy chain and a light chain, wherein the heavy chain comprises a variable region comprising an amino acid sequence as set forth in SEQ ID NO: 13 or a fragment thereof, and wherein the light chain comprises a variable region comprising an amino acid sequence as set forth in SEQ ID NO: 14 or a fragment thereof.

[010] In certain embodiments, the invention provides for an antibody comprising a heavy chain and a light chain, wherein the heavy chain comprises an amino acid sequence as set forth in SEQ ID NO: 2 or a fragment thereof.

[011] In certain embodiments, the invention provides for an antibody comprising a heavy chain and a light chain, wherein the heavy chain comprises a variable region comprising an amino acid sequence as set forth in SEQ ID NO: 13 or a fragment thereof.

[012] In certain embodiments, the invention provides for an antibody comprising a heavy chain and a light chain, wherein the light chain

comprises an amino acid sequence as set forth in SEQ ID NO: 4 or a fragment thereof.

[013] In certain embodiments, the invention provides for an antibody comprising a heavy chain and a light chain, wherein the light chain comprises a variable region comprising an amino acid sequence as set forth in SEQ ID NO: 14 or a fragment thereof.

[014] In certain embodiments, the invention provides for an antibody, comprising a heavy chain and a light chain, (a) wherein the heavy chain comprises a first variable region, and wherein the first variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 13, and (b) wherein the light chain comprises a second variable region, and wherein the second variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 14, and (c) wherein the antibody interacts with an osteoprotegerin ligand (OPGL).

[015] In certain embodiments, the first variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 13, and the second variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 14.

[016] In certain embodiments, the first variable region comprises a sequence that has at least 99% identity to the amino acid sequence set forth in SEQ ID NO: 13, and the second variable region comprises a sequence that has at least 99% identity to the amino acid sequence set forth in SEQ ID NO: 14.

[017] In certain embodiments, the invention provides for a heavy chain, comprising an amino acid sequence as set forth in SEQ ID NO:2 or a fragment thereof. In certain embodiments, the invention provides for a heavy chain comprising a variable region and a constant region, wherein the variable region comprises an amino acid sequence as set forth in SEQ ID NO: 13 or a fragment thereof.

[018] In certain embodiments, the invention provides for a light chain, comprising an amino acid sequence as set forth in SEQ ID NO:4 or a fragment thereof. In certain embodiments, the invention provides for a light chain comprising an amino acid sequence as set forth in SEQ ID NO: 14 or a fragment thereof.

[019] In certain embodiments of the invention, single chain antibodies are provided. In certain embodiments of the invention, single chain Fv antibodies are provided. In certain embodiments of the invention, Fab antibodies are provided. In certain embodiments of the invention, Fab' antibodies are provided. In certain embodiments of the invention, (Fab')₂ antibodies are provided.

[020] In certain embodiments, a pharmaceutical composition comprising an antibody of the invention is provided. In certain embodiments, a pharmaceutical composition comprising a therapeutically effective amount of an antibody to OPGL is provided.

[021] In certain embodiments, a pharmaceutical composition comprises an antibody to OPGL and at least one therapeutic agent selected from

a bone morphogenic factor, transforming growth factor- β (TGF- β), an interleukin-1 (IL-1) inhibitor, IL-1ra, Kineret™, a TNF α inhibitor, a soluble TNF α receptor, Enbrel™, an anti-TNF α antibody, Remicade™, a D2E7 antibody, a parathyroid hormone, an analog of a parathyroid hormone, a parathyroid hormone related protein, an analog of a parathyroid hormone related protein, a prostaglandin, a bisphosphonate, an alendronate, fluoride, calcium, a non-steroidal anti-inflammatory drug (NSAID), a COX-2 inhibitor, Celebrex™, Vioxx™; an immunosuppressant, methotrexate, leflunomide, a serine protease inhibitor, a secretory leukocyte protease inhibitor (SLPI), an IL-6 inhibitor, an antibody to IL-6, an IL-8 inhibitor, an antibody to IL-8, an IL-18 inhibitor, an IL-18 binding protein, an IL-18 antibody, an Interleukin-1 converting enzyme (ICE) modulator, a fibroblast growth factor (FGF), an FGF modulator, a PAF antagonist, a keratinocyte growth factor (KGF), a KGF-related molecule, a KGF modulator; a matrix metalloproteinase (MMP) modulator, a nitric oxide synthase (NOS) modulator, a modulator of glucocorticoid receptor, a modulator of glutamate receptor, a modulator of lipopolysaccharide (LPS) levels, a noradrenaline, a noradrenaline mimetic, and a noradrenaline modulator.

[022] In certain embodiments of the invention, a method of treating an osteopenic disorder is provided, comprising administering a pharmaceutically effective amount of an antibody. In certain embodiments, a method of treating an osteopenic disorder comprising administering a pharmaceutical composition is provided.

[023] In certain embodiments, a method of treating an inflammatory condition with attendant bone loss in a patient comprising administering a pharmaceutical composition is provided.

[024] In certain embodiments, a method of treating an autoimmune condition with attendant bone loss in a patient comprising administering a pharmaceutical composition is provided.

[025] In certain embodiments, a method of treating rheumatoid arthritis in a patient, comprising administering a pharmaceutical composition of the invention is provided.

[026] In certain embodiments of the invention, a method of detecting the level of OPGL in a biological sample is provided, comprising contacting the sample with an antibody.

BRIEF DESCRIPTION OF THE FIGURES

[027] Figure 1 shows a cDNA sequence encoding the α OPGL-1 antibody heavy chain (SEQ ID NO: 1).

[028] Figure 2 shows the amino acid sequence of the α OPGL-1 antibody heavy chain (SEQ ID NO: 2).

[029] Figure 3 shows a cDNA sequence encoding the α OPGL-1 antibody light chain (SEQ ID NO: 3).

[030] Figure 4 shows the amino acid sequence of the α OPGL-1 antibody light chain (SEQ ID NO: 4).

[031] Figure 5 shows a schematic diagram of the α OPGL-1 kappa light chain expression plasmid α OPGL-1-Kappa/pDSRa19.

[032] Figure 6 shows a schematic diagram of the α OPGL-1 IgG2 heavy chain expression plasmid, α OPGL-1-IgG2/pDSRa19.

[033] Figure 7 shows dose-dependent binding of α OPGL-1 to OPGL-coated EIA plates.

[034] Figure 8 shows specific binding of α OPGL-1 to membrane-bound OPGL.

[035] Figure 9 shows inhibition of α OPGL-1 binding to OPGL-coated EIA plates by soluble OPGL.

[036] Figure 10 shows specific binding of α OPGL-1 to OPGL-coated EIA plates.

[037] Figure 11 shows dose-dependent inhibition of osteoclast formation by α OPGL-1.

[038] Figure 12 shows dose-dependent inhibition of OPGL binding to ODAR by α OPGL-1.

[039] Figure 13 shows the mean serum concentration time profiles after administering a single dose of α OPGL-1 to Cynomolgus Monkeys.

[040] Figure 14 shows the mean percent change in serum N-Tx concentration after administering a single dose of α OPGL-1 to Cynomolgus Monkeys.

[041] Figure 15 shows the mean percent change in urine N-Tx concentration after administering a single dose of α OPGL-1 to Cynomolgus Monkeys.

[042] Figure 16 shows antibody positive and negative serum-concentration time profiles after administering a single dose of α OPGL-1 to Cynomolgus Monkeys.

[043] Figure 17 shows the amino acid sequence of the α OPGL-1 antibody heavy chain variable region (SEQ ID NO: 13).

[044] Figure 18 shows the amino acid sequence of the α OPGL-1 antibody light chain variable region (SEQ ID NO: 14).

[045] Figure 19 shows a cell culture process for production of α OPGL-1.

[046] Figure 20 shows the serum calcium percent change after administering a single dose of α OPGL-1 to Cynomolgus monkeys.

[047] Figure 21 shows the mean serum Alkaline Phosphatase percent change after administering a single dose of α OPGL-1 to Cynomolgus monkeys.

DETAILED DESCRIPTION OF CERTAIN PREFERRED EMBODIMENTS

[048] The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described. All references cited in this application are expressly incorporated by reference herein for any purpose.

Definitions

[049] Standard techniques may be used for recombinant DNA, oligonucleotide synthesis, and tissue culture and transformation (e.g., electroporation, lipofection). Enzymatic reactions and purification techniques may be performed according to manufacturer's specifications or as commonly accomplished in the art or as described herein. The foregoing techniques and procedures may be generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification. See e.g., Sambrook et al. *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)), which is incorporated herein by reference for any purpose. Unless specific definitions are provided, the nomenclatures utilized in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques may be used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

[050] As utilized in accordance with the present disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

[051] The term "isolated polynucleotide" as used herein shall mean a polynucleotide of genomic, cDNA, or synthetic origin or some

combination thereof, which by virtue of its origin the "isolated polynucleotide" (1) is not associated with all or a portion of a polynucleotide in which the "isolated polynucleotide" is found in nature, (2) is linked to a polynucleotide which it is not linked to in nature, or (3) does not occur in nature as part of a larger sequence.

[052] The term "isolated protein" referred to herein means a protein encoded by cDNA, recombinant RNA, or synthetic origin or some combination thereof, which (1) is free of at least some proteins with which it would normally be found, (2) is essentially free of other proteins from the same source, e.g., from the same species, (3) is expressed by a cell from a different species, or (4) does not occur in nature.

[053] The term "polypeptide" is used herein as a generic term to refer to native proteins, or sequences that have deletions, additions, and/or substitutions of one or more amino acids of the native sequence. The term "polypeptide" also encompasses α OPGL-1 (as described below, SEQ ID NO: 2 and SEQ ID NO: 4), or sequences that have deletions, additions, and/or substitutions of one or more amino acid of α OPGL-1. According to certain embodiments, the invention comprises the human heavy chain immunoglobulin molecule represented by Figure 2 (SEQ ID NO: 2) and the human light chain immunoglobulin molecule represented by Figure 4 (SEQ ID NO: 4), or fragments or analogs thereof.

[054] The term "naturally-occurring" as used herein as applied to an object refers to the fact that an object can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism (including

viruses) that can be isolated from a source in nature and which has not been intentionally modified by man in the laboratory or otherwise is naturally-occurring.

[055] The term "operably linked" as used herein refers to components that are in a relationship permitting them to function in their intended manner. For example, a control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

[056] The term "control sequence" as used herein refers to polynucleotide sequences which may effect the expression and processing of coding sequences to which they are ligated. The nature of such control sequences may differ depending upon the host organism. According to certain embodiments, control sequences for prokaryotes may include promoter, ribosomal binding site, and transcription termination sequence. According to certain embodiments, control sequences for eukaryotes may include promoters and transcription termination sequence. In certain embodiments, "control sequences" can include leader sequences and/or fusion partner sequences.

[057] The term "polynucleotide" as referred to herein means a polymeric form of nucleotides of at least 10 bases in length. In certain embodiments, the bases may be ribonucleotides or deoxyribonucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA.

[058] The term "oligonucleotide" referred to herein includes naturally occurring, and modified nucleotides linked together by naturally

occurring, and/or non-naturally occurring oligonucleotide linkages.

Oligonucleotides are a polynucleotide subset generally comprising a length of 200 bases or fewer. In certain embodiments, oligonucleotides are 10 to 60 bases in length. In certain embodiments, oligonucleotides are 12, 13, 14, 15, 16, 17, 18, 19, or 20 to 40 bases in length. Oligonucleotides may be single stranded or double stranded, e.g. for use in the construction of a gene mutant.

Oligonucleotides of the invention may be sense or antisense oligonucleotides.

[059] The term "naturally occurring nucleotides" includes deoxyribonucleotides and ribonucleotides. The term "modified nucleotides" includes nucleotides with modified or substituted sugar groups and the like. The term "oligonucleotide linkages" includes oligonucleotides linkages such as phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate, phosphoroamidate, and the like. See, e.g., LaPlanche et al. Nucl. Acids Res. 14:9081 (1986); Stec et al. J. Am. Chem. Soc. 106:6077 (1984); Stein et al. Nucl. Acids Res. 16:3209 (1988); Zon et al. Anti-Cancer Drug Design 6:539 (1991); Zon et al. Oligonucleotides and Analogues: A Practical Approach, pp. 87-108 (F. Eckstein, Ed., Oxford University Press, Oxford England (1991)); Stec et al. U.S. Pat. No. 5,151,510; Uhlmann and Peyman Chemical Reviews 90:543 (1990), the disclosures of which are hereby incorporated by reference for any purpose. An oligonucleotide can include a label for detection.

[060] Identity and similarity of related and polypeptides can be readily calculated by known methods. Such methods include, but are not limited

to, those described in Computational Molecular Biology, Lesk, A.M., ed., Oxford University Press, New York (1988); Biocomputing: Informatics and Genome Projects, Smith, D.W., ed., Academic Press, New York (1993); Computer Analysis of Sequence Data, Part 1, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey (1994); Sequence Analysis in Molecular Biology, von Heinjé, G., Academic Press (1987); Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M. Stockton Press, New York (1991); and Carillo *et al.*, *SIAM J. Applied Math.*, 48:1073 (1988).

[061] Preferred methods to determine identity are designed to give the largest match between the sequences tested. Methods to determine identity are described in publicly available computer programs. Preferred computer program methods to determine identity between two sequences include, but are not limited to, the GCG program package, including GAP (Devereux *et al.*, *Nucl. Acid. Res.*, 12:387 (1984); Genetics Computer Group, University of Wisconsin, Madison, WI, BLASTP, BLASTN, and FASTA (Altschul *et al.*, *J. Mol. Biol.*, 215:403-410 (1990)). The BLASTX program is publicly available from the National Center for Biotechnology Information (NCBI) and other sources (*BLAST Manual*, Altschul *et al.* NCB/NLM/NIH Bethesda, MD 20894; Altschul *et al.*, *supra* (1990)). The well-known Smith Waterman algorithm may also be used to determine identity.

[062] Certain alignment schemes for aligning two amino acid sequences may result in the matching of only a short region of the two sequences, and this small aligned region may have very high sequence identity

even though there is no significant relationship between the two full-length sequences. Accordingly, in certain embodiments, the selected alignment method (GAP program) will result in an alignment that spans at least 50 contiguous amino acids of the target polypeptide.

[063] For example, using the computer algorithm GAP (Genetics Computer Group, University of Wisconsin, Madison, WI), two polypeptides for which the percent sequence identity is to be determined are aligned for optimal matching of their respective amino acids (the "matched span", as determined by the algorithm). In certain embodiments, a gap opening penalty (which is calculated as 3X the average diagonal; the "average diagonal" is the average of the diagonal of the comparison matrix being used; the "diagonal" is the score or number assigned to each perfect amino acid match by the particular comparison matrix) and a gap extension penalty (which is usually 1/10 times the gap opening penalty), as well as a comparison matrix such as PAM 250 or BLOSUM 62 are used in conjunction with the algorithm. In certain embodiments, a standard comparison matrix (see Dayhoff *et al.*, *Atlas of Protein Sequence and Structure*, 5(3)(1978) for the PAM 250 comparison matrix; Henikoff *et al.*, *Proc. Natl. Acad. Sci USA*, 89:10915-10919 (1992) for the BLOSUM 62 comparison matrix) is also used by the algorithm.

[064] In certain embodiments, the parameters for a polypeptide sequence comparison include the following:

Algorithm: Needleman *et al.*, *J. Mol. Biol.*, 48:443-453 (1970);
Comparison matrix: BLOSUM 62 from Henikoff *et al.*, *supra* (1992);
Gap Penalty: 12
Gap Length Penalty: 4

Threshold of Similarity: 0

[065] The GAP program may be useful with the above parameters.

In certain embodiments, the aforementioned parameters are the default parameters for polypeptide comparisons (along with no penalty for end gaps) using the GAP algorithm.

[066] As used herein, the twenty conventional amino acids and their abbreviations follow conventional usage. See Immunology--A Synthesis (2nd Edition, E. S. Golub and D. R. Gren, Eds., Sinauer Associates, Sunderland, Mass. (1991)), which is incorporated herein by reference for any purpose. Stereoisomers (e.g., D-amino acids) of the twenty conventional amino acids, unnatural amino acids such as α -, α -disubstituted amino acids, N-alkyl amino acids, lactic acid, and other unconventional amino acids may also be suitable components for polypeptides of the present invention. Examples of unconventional amino acids include: 4-hydroxyproline, γ -carboxyglutamate, ϵ -N,N,N-trimethyllysine, ϵ -N-acetyllysine, O-phosphoserine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine, σ -N-methylarginine, and other similar amino acids and imino acids (e.g., 4-hydroxyproline). In the polypeptide notation used herein, the left-hand direction is the amino terminal direction and the right-hand direction is the carboxy-terminal direction, in accordance with standard usage and convention.

[067] Similarly, unless specified otherwise, the left-hand end of single-stranded polynucleotide sequences is the 5' end; the left-hand direction of double-stranded polynucleotide sequences is referred to as the 5' direction. The

direction of 5' to 3' addition of nascent RNA transcripts is referred to as the transcription direction; sequence regions on the DNA strand having the same sequence as the RNA and which are 5' to the 5' end of the RNA transcript are referred to as "upstream sequences"; sequence regions on the DNA strand having the same sequence as the RNA and which are 3' to the 3' end of the RNA transcript are referred to as "downstream sequences".

[068] Conservative amino acid substitutions may encompass non-naturally occurring amino acid residues, which are typically incorporated by chemical peptide synthesis rather than by synthesis in biological systems. These include peptidomimetics and other reversed or inverted forms of amino acid moieties.

[069] Naturally occurring residues may be divided into classes based on common side chain properties:

- 1) hydrophobic: norleucine, Met, Ala, Val, Leu, Ile;
- 2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln;
- 3) acidic: Asp, Glu;
- 4) basic: His, Lys, Arg;
- 5) residues that influence chain orientation: Gly, Pro; and
- 6) aromatic: Trp, Tyr, Phe.

[070] For example, non-conservative substitutions may involve the exchange of a member of one of these classes for a member from another class. Such substituted residues may be introduced into regions of the human antibody that are homologous with non-human antibodies, or into the non-homologous regions of the molecule.

[071] In making such changes, according to certain embodiments, the hydrophobic index of amino acids may be considered. Each amino acid has been assigned a hydrophobic index on the basis of its hydrophobicity and charge characteristics. They are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

[072] The importance of the hydrophobic amino acid index in conferring interactive biological function on a protein is understood in the art. Kyte *et al.*, *J. Mol. Biol.*, 157:105-131 (1982). It is known that certain amino acids may be substituted for other amino acids having a similar hydrophobic index or score and still retain a similar biological activity. In making changes based upon the hydrophobic index, in certain embodiments, the substitution of amino acids whose hydrophobic indices are within ± 2 is included. In certain embodiments, those which are within ± 1 are included, and in certain embodiments, those within ± 0.5 are included.

[073] It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity, particularly where the biologically functional protein or peptide thereby created is intended for use in immunological embodiments, as in the present case. In certain embodiments, the greatest local average hydrophilicity of a protein, as governed

by the hydrophilicity of its adjacent amino acids, correlates with its immunogenicity and antigenicity, i.e., with a biological property of the protein.

[074] The following hydrophilicity values have been assigned to these amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0 $\pm$ 1); glutamate (+3.0 $\pm$ 1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5 $\pm$ 1); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5) and tryptophan (-3.4). In making changes based upon similar hydrophilicity values, in certain embodiments, the substitution of amino acids whose hydrophilicity values are within ± 2 is included, in certain embodiments, those which are within ± 1 are included, and in certain embodiments, those within ± 0.5 are included. One may also identify epitopes from primary amino acid sequences on the basis of hydrophilicity. These regions are also referred to as "epitopic core regions."

[075] Exemplary amino acid substitutions are set forth in Table 1.

Table 1: Amino Acid Substitutions

Original Residues	Exemplary Substitutions	Preferred Substitutions
Ala	Val, Leu, Ile	Val
Arg	Lys, Gln, Asn	Lys
Asn	Gln	Gln
Asp	Glu	Glu
Cys	Ser, Ala	Ser
Gln	Asn	Asn
Glu	Asp	Asp
Gly	Pro, Ala	Ala
His	Asn, Gln, Lys, Arg	Arg
Ile	Leu, Val, Met, Ala, Phe, Norleucine	Leu

Leu	Norleucine, Ile, Val, Met, Ala, Phe	Ile
Lys	Arg, 1,4 Diamino- butyric Acid, Gln, Asn	Arg
Met	Leu, Phe, Ile	Leu
Phe	Leu, Val, Ile, Ala, Tyr	Leu
Pro	Ala	Gly
Ser	Thr, Ala, Cys	Thr
Thr	Ser	Ser
Trp	Tyr, Phe	Tyr
Tyr	Trp, Phe, Thr, Ser	Phe
Val	Ile, Met, Leu, Phe, Ala, Norleucine	Leu

[076] A skilled artisan will be able to determine suitable variants of the polypeptide as set forth herein using well-known techniques. In certain embodiments, one skilled in the art may identify suitable areas of the molecule that may be changed without destroying activity by targeting regions not believed to be important for activity. In certain embodiments, one can identify residues and portions of the molecules that are conserved among similar polypeptides. In certain embodiments, even areas that may be important for biological activity or for structure may be subject to conservative amino acid substitutions without destroying the biological activity or without adversely affecting the polypeptide structure.

[077] Additionally, one skilled in the art can review structure-function studies identifying residues in similar polypeptides that are important for activity or structure. In view of such a comparison, one can predict the importance of amino acid residues in a protein that correspond to amino acid residues which are important for activity or structure in similar proteins. One

skilled in the art may opt for chemically similar amino acid substitutions for such predicted important amino acid residues.

[078] One skilled in the art can also analyze the three-dimensional structure and amino acid sequence in relation to that structure in similar polypeptides. In view of such information, one skilled in the art may predict the alignment of amino acid residues of an antibody with respect to its three dimensional structure. In certain embodiments, one skilled in the art may choose not to make radical changes to amino acid residues predicted to be on the surface of the protein, since such residues may be involved in important interactions with other molecules. Moreover, one skilled in the art may generate test variants containing a single amino acid substitution at each desired amino acid residue. The variants can then be screened using activity assays known to those skilled in the art. Such variants could be used to gather information about suitable variants. For example, if one discovered that a change to a particular amino acid residue resulted in destroyed, undesirably reduced, or unsuitable activity, variants with such a change may be avoided. In other words, based on information gathered from such routine experiments, one skilled in the art can readily determine the amino acids where further substitutions should be avoided either alone or in combination with other mutations.

[079] A number of scientific publications have been devoted to the prediction of secondary structure. See Moult J., *Curr. Op. in Biotech.*, 7(4):422-427 (1996), Chou et al., *Biochemistry*, 13(2):222-245 (1974); Chou et al., *Biochemistry*, 113(2):211-222 (1974); Chou et al., *Adv. Enzymol. Relat. Areas*

Mol. Biol., 47:45-148 (1978); Chou *et al.*, *Ann. Rev. Biochem.*, 47:251-276 and Chou *et al.*, *Biophys. J.*, 26:367-384 (1979). Moreover, computer programs are currently available to assist with predicting secondary structure. One method of predicting secondary structure is based upon homology modeling. For example, two polypeptides or proteins which have a sequence identity of greater than 30%, or similarity greater than 40% often have similar structural topologies. The recent growth of the protein structural database (PDB) has provided enhanced predictability of secondary structure, including the potential number of folds within a polypeptide's or protein's structure. See Holm *et al.*, *Nucl. Acid. Res.*, 27(1):244-247 (1999). It has been suggested (Brenner *et al.*, *Curr. Op. Struct. Biol.*, 7(3):369-376 (1997)) that there are a limited number of folds in a given polypeptide or protein and that once a critical number of structures have been resolved, structural prediction will become dramatically more accurate.

[080] Additional methods of predicting secondary structure include "threading" (Jones, D., *Curr. Opin. Struct. Biol.*, 7(3):377-87 (1997); Sippl *et al.*, *Structure*, 4(1):15-19 (1996)), "profile analysis" (Bowie *et al.*, *Science*, 253:164-170 (1991); Gribskov *et al.*, *Meth. Enzym.*, 183:146-159 (1990); Gribskov *et al.*, *Proc. Nat. Acad. Sci.*, 84(13):4355-4358 (1987)), and "evolutionary linkage" (See Holm, *supra* (1999), and Brenner, *supra* (1997)).

[081] In certain embodiments, antibody variants include glycosylation variants wherein the number and/or type of glycosylation site has been altered compared to the amino acid sequences of the parent polypeptide. In certain embodiments, protein variants comprise a greater or a lesser number

of N-linked glycosylation sites than the native protein. An N-linked glycosylation site is characterized by the sequence: Asn-X-Ser or Asn-X-Thr, wherein the amino acid residue designated as X may be any amino acid residue except proline. The substitution of amino acid residues to create this sequence provides a potential new site for the addition of an N-linked carbohydrate chain.

Alternatively, substitutions which eliminate this sequence will remove an existing N-linked carbohydrate chain. Also provided is a rearrangement of N-linked carbohydrate chains wherein one or more N-linked glycosylation sites (typically those that are naturally occurring) are eliminated and one or more new N-linked sites are created. Additional preferred antibody variants include cysteine variants wherein one or more cysteine residues are deleted from or substituted for another amino acid (e.g., serine) as compared to the parent amino acid sequence. Cysteine variants may be useful when antibodies must be refolded into a biologically active conformation such as after the isolation of insoluble inclusion bodies. Cysteine variants generally have fewer cysteine residues than the native protein, and typically have an even number to minimize interactions resulting from unpaired cysteines.

[082] According to certain embodiments, amino acid substitutions are those which: (1) reduce susceptibility to proteolysis, (2) reduce susceptibility to oxidation, (3) alter binding affinity for forming protein complexes, (4) alter binding affinities, and/or (4) confer or modify other physicochemical or functional properties on such polypeptides. According to certain embodiments, single or multiple amino acid substitutions (in certain embodiments, conservative amino

acid substitutions) may be made in the naturally-occurring sequence (in certain embodiments, in the portion of the polypeptide outside the domain(s) forming intermolecular contacts). In certain embodiments, a conservative amino acid substitution typically may not substantially change the structural characteristics of the parent sequence (e.g., a replacement amino acid should not tend to break a helix that occurs in the parent sequence, or disrupt other types of secondary structure that characterizes the parent sequence). Examples of art-recognized polypeptide secondary and tertiary structures are described in *Proteins, Structures and Molecular Principles* (Creighton, Ed., W. H. Freeman and Company, New York (1984)); *Introduction to Protein Structure* (C. Branden and J. Tooze, eds., Garland Publishing, New York, N.Y. (1991)); and Thornton et al. *Nature* 354:105 (1991), which are each incorporated herein by reference.

[083] The term "polypeptide fragment" as used herein refers to a polypeptide that has an amino-terminal and/or carboxy-terminal deletion. In certain embodiments, fragments are at least 5 to 467 amino acids long. It will be appreciated that in certain embodiments, fragments are at least 5, 6, 8, 10, 14, 20, 50, 70, 100, 150, 200, 250, 300, 350, 400, or 450 amino acids long.

[084] Peptide analogs are commonly used in the pharmaceutical industry as non-peptide drugs with properties analogous to those of the template peptide. These types of non-peptide compound are termed "peptide mimetics" or "peptidomimetics". Fauchere, J. *Adv. Drug Res.* 15:29 (1986); Veber and Freidinger *TINS* p.392 (1985); and Evans et al. *J. Med. Chem.* 30:1229 (1987), which are incorporated herein by reference for any purpose. Such compounds

are often developed with the aid of computerized molecular modeling. Peptide mimetics that are structurally similar to therapeutically useful peptides may be used to produce a similar therapeutic or prophylactic effect. Generally, peptidomimetics are structurally similar to a paradigm polypeptide (i.e., a polypeptide that has a biochemical property or pharmacological activity), such as human antibody, but have one or more peptide linkages optionally replaced by a linkage selected from: $--CH_2 NH--$, $--CH_2 S--$, $--CH_2 -CH_2 --$, $--CH=CH--$ (cis and trans), $--COCH_2 --$, $--CH(OH)CH_2 --$, and $--CH_2 SO--$, by methods well known in the art. Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (e.g., D-lysine in place of L-lysine) may be used in certain embodiments to generate more stable peptides. In addition, constrained peptides comprising a consensus sequence or a substantially identical consensus sequence variation may be generated by methods known in the art (Rizo and Gierasch Ann. Rev. Biochem. 61:387 (1992), incorporated herein by reference for any purpose); for example, by adding internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

[085] "Antibody" or "antibody peptide(s)" refer to an intact antibody, or a binding fragment thereof that competes with the intact antibody for specific binding. In certain embodiments, binding fragments are produced by recombinant DNA techniques. In certain embodiments, binding fragments are produced by enzymatic or chemical cleavage of intact antibodies. Binding

fragments include, but are not limited to, Fab, Fab', F(ab')₂, Fv, and single-chain antibodies.

[086] The term "heavy chain" includes any polypeptide having sufficient variable region sequence to confer specificity for an OPGL. The term "light chain" includes any polypeptide having sufficient variable region sequence to confer specificity for an OPGL. A full-length heavy chain includes a variable region domain, V_H, and three constant region domains, C_H1, C_H2, and C_H3. The V_H domain is at the amino-terminus of the polypeptide, and the C_H3 domain is at the carboxy-terminus. The term "heavy chain", as used herein, encompasses a full-length heavy chain and fragments thereof. A full-length light chain includes a variable region domain, V_L, and a constant region domain, C_L. Like the heavy chain, the variable region domain of the light chain is at the amino-terminus of the polypeptide. The term "light chain", as used herein, encompasses a full-length light chain and fragments thereof. A Fab fragment is comprised of one light chain and the C_H1 and variable regions of one heavy chain. The heavy chain of a Fab molecule cannot form a disulfide bond with another heavy chain molecule. A Fab' fragment contains one light chain and one heavy chain that contains more of the constant region, between the C_H1 and C_H2 domains, such that an interchain disulfide bond can be formed between two heavy chains to form a F(ab')₂ molecule. The Fv region comprises the variable regions from both the heavy and light chains, but lacks the constant regions. Single-chain antibodies are Fv molecules in which the heavy and light chain variable regions have been connected by a flexible linker to form a single polypeptide chain which

forms an antigen-binding region. Single chain antibodies are discussed in detail in WO 88/01649 and U.S. Patent Nos. 4,946,778 and 5,260,203.

[087] A bivalent antibody other than a "multispecific" or "multifunctional" antibody, in certain embodiments, typically is understood to have each of its binding sites identical.

[088] An antibody substantially inhibits adhesion of a ligand to a receptor when an excess of antibody reduces the quantity of receptor bound to counterreceptor by at least about 20%, 40%, 60%, 80%, 85%, or more (as measured in an *in vitro* competitive binding assay).

[089] The term "epitope" includes any polypeptide determinant capable of specific binding to an immunoglobulin or T-cell receptor. In certain embodiments, epitope determinants include chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl, or sulfonyl, and, in certain embodiments, may have specific three dimensional structural characteristics, and/or specific charge characteristics. An epitope is a region of an antigen that is bound by an antibody. In certain embodiments, an antibody is said to specifically bind an antigen when it preferentially recognizes its target antigen in a complex mixture of proteins and/or macromolecules. In certain embodiments, an antibody is said to specifically bind an antigen when the dissociation constant is $\leq 1 \mu\text{M}$, in certain embodiments, when the dissociation constant is $\leq 100 \text{ nM}$, and in certain embodiments, when the dissociation constant is $\leq 10 \text{ nM}$.

[090] The term "agent" is used herein to denote a chemical compound, a mixture of chemical compounds, a biological macromolecule, or an extract made from biological materials.

[091] As used herein, the terms "label" or "labeled" refers to incorporation of a detectable marker, e.g., by incorporation of a radiolabeled amino acid or attachment to a polypeptide of biotin moieties that can be detected by marked avidin (e.g., streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or colorimetric methods). In certain embodiments, the label or marker can also be therapeutic. Various methods of labeling polypeptides and glycoproteins are known in the art and may be used. Examples of labels for polypeptides include, but are not limited to, the following: radioisotopes or radionuclides (e.g., ^3H , ^{14}C , ^{15}N , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I), fluorescent labels (e.g., FITC, rhodamine, lanthanide phosphors), enzymatic labels (e.g., horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase), chemiluminescent, biotinyl groups, predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In certain embodiments, labels are attached by spacer arms of various lengths to reduce potential steric hindrance.

[092] The term "biological sample", as used herein, includes, but is not limited to, any quantity of a substance from a living thing or formerly living thing. Such living things include, but are not limited to, humans, mice, monkeys, rats, rabbits, and other animals. Such substances include, but are not limited to,

blood, serum, urine, cells, organs, tissues, bone, bone marrow, lymph nodes, and skin.

[093] The term "osteopenic disorder" includes, but is not limited to, osteoporosis, osteopenia, Paget's disease, lytic bone metastases, periodontitis, rheumatoid arthritis, and bone loss due to immobilization. In addition to these bone disorders, certain cancers are known to increase osteoclast activity and induce bone resorption, such as breast, prostate, and multiple myeloma. These cancers are now known to produce factors that result in the over-expression of OPGL in the bone, and lead to increased osteoclast numbers and activity.

[094] The term "pharmaceutical agent or drug" as used herein refers to a chemical compound or composition capable of inducing a desired therapeutic effect when properly administered to a patient.

[095] The term "modulator," as used herein, is a compound that changes or alters the activity or function of a molecule. For example, a modulator may cause an increase or decrease in the magnitude of a certain activity or function of a molecule compared to the magnitude of the activity or function observed in the absence of the modulator. In certain embodiments, a modulator is an inhibitor, which decreases the magnitude of at least one activity or function of a molecule. Certain exemplary activities and functions of a molecule include, but are not limited to, binding affinity, enzymatic activity, and signal transduction. Certain exemplary inhibitors include, but are not limited to, proteins, peptides, antibodies, peptibodies, carbohydrates or small organic molecules. Peptibodies are described, e.g., in WO01/83525.

[096] As used herein, "substantially pure" means an object species is the predominant species present (i.e., on a molar basis it is more abundant than any other individual species in the composition). In certain embodiments, a substantially purified fraction is a composition wherein the object species comprises at least about 50 percent (on a molar basis) of all macromolecular species present. In certain embodiments, a substantially pure composition will comprise more than about 80%, 85%, 90%, 95%, or 99% of all macromolecular species present in the composition. In certain embodiments, the object species is purified to essential homogeneity (contaminant species cannot be detected in the composition by conventional detection methods) wherein the composition consists essentially of a single macromolecular species.

[097] The term patient includes human and animal subjects.

[098] In this application, the use of the singular includes the plural unless specifically stated otherwise. In this application, the use of "or" means "and/or" unless stated otherwise. Furthermore, the use of the term "including", as well as other forms, such as "includes" and "included", is not limiting. Also, terms such as "element" or "component" encompass both elements and components comprising one unit and elements and components that comprise more than one subunit unless specifically stated otherwise.

[099] Osteoprotegerin Ligand (OPGL), a member of the tumor necrosis factor (TNF) family of cytokines, is involved in the formation of osteoclasts. Increased osteoclast activity correlates with a number of osteopenic

disorders, including post-menopausal osteoporosis, Paget's disease, lytic bone metastases, and rheumatoid arthritis. Therefore, a reduction in OPGL activity may result in a decrease in osteoclast activity and may reduce the severity of osteopenic disorders. According to certain embodiments of the invention, antibodies directed to OPGL may be used treat osteopenic disorders, including by not limited to, those mentioned above.

[0100] In certain embodiments of the present invention, there is provided a fully human monoclonal antibody against human osteoprotegerin ligand (OPGL). In certain embodiments, nucleotide sequences encoding, and amino acid sequences comprising, heavy and light chain immunoglobulin molecules, particularly sequences corresponding to the variable regions are provided. In certain embodiments, sequences corresponding to complementarity determining regions (CDR's), specifically from CDR1 through CDR3, are provided. According to certain embodiments, a hybridoma cell line expressing such an immunoglobulin molecule and monoclonal antibody is also provided. In certain embodiments, purified human monoclonal antibody against human OPGL is provided.

[0101] The ability to clone and reconstruct megabase-sized human loci in yeast artificial chromosomes (YACs) and to introduce them into the mouse germline provides an approach to elucidating the functional components of very large or crudely mapped loci as well as generating useful models of human disease. Furthermore, the utilization of such technology for substitution of mouse loci with their human equivalents could provide unique insights into the

expression and regulation of human gene products during development, their communication with other systems, and their involvement in disease induction and progression.

[0102] An important practical application of such a strategy is the "humanization" of the mouse humoral immune system. Introduction of human immunoglobulin (Ig) loci into mice in which the endogenous Ig genes have been inactivated offers the opportunity to study the mechanisms underlying programmed expression and assembly of antibodies as well as their role in B-cell development. Furthermore, such a strategy could provide a source for production of fully human monoclonal antibodies (MAbs). In certain embodiments, fully human antibodies are expected to minimize the immunogenic and allergic responses intrinsic to mouse or mouse-derivatized Mabs, and thus, in certain embodiments, increase the efficacy and safety of the administered antibodies. In certain embodiments, fully human antibodies may be used in the treatment of chronic and recurring human diseases, such as osteoporosis, inflammation, autoimmunity, and cancer, which may involve repeated antibody administrations.

[0103] One can engineer mouse strains deficient in mouse antibody production with large fragments of the human Ig loci in anticipation that such mice would produce human antibodies in the absence of mouse antibodies. Large human Ig fragments may preserve the large variable gene diversity as well as the proper regulation of antibody production and expression. By exploiting the mouse machinery for antibody diversification and selection and the lack of

immunological tolerance to human proteins, the reproduced human antibody repertoire in these mouse strains may yield high affinity antibodies against any antigen of interest, including human antigens. Using the hybridoma technology, antigen-specific human MAbs with the desired specificity may be produced and selected.

[0104] In certain embodiments, one may use constant regions from species other than human along with the human variable region(s).

Naturally Occurring Antibody Structure

[0105] Naturally occurring antibody structural units typically comprise a tetramer. Each such tetramer typically is composed of two identical pairs of polypeptide chains, each pair having one full-length "light" (in certain embodiments, about 25 kDa) and one full-length "heavy" chain (in certain embodiments, about 50-70 kDa). The amino-terminal portion of each chain typically includes a variable region of about 100 to 110 or more amino acids that typically is responsible for antigen recognition. The carboxy-terminal portion of each chain typically defines a constant region that may be responsible for effector function. Human light chains are typically classified as kappa and lambda light chains. Heavy chains are typically classified as mu, delta, gamma, alpha, or epsilon, and define the antibody's isotype as IgM, IgD, IgG, IgA, and IgE, respectively. IgG has several subclasses, including, but not limited to, IgG1, IgG2, IgG3, and IgG4. IgM has subclasses including, but not limited to, IgM1 and IgM2. IgA is similarly subdivided into subclasses including, but not limited to,

IgA1 and IgA2. Within full-length light and heavy chains, typically, the variable and constant regions are joined by a "J" region of about 12 or more amino acids, with the heavy chain also including a "D" region of about 10 more amino acids. See, e.g., *Fundamental Immunology* Ch. 7 (Paul, W., ed., 2nd ed. Raven Press, N.Y. (1989)) (incorporated by reference in its entirety for all purposes). The variable regions of each light/heavy chain pair typically form the antigen binding site.

[0106] The variable regions typically exhibit the same general structure of relatively conserved framework regions (FR) joined by three hyper variable regions, also called complementarity determining regions or CDRs. The CDRs from the two chains of each pair typically are aligned by the framework regions, which may enable binding to a specific epitope. From N-terminal to C-terminal, both light and heavy chain variable regions typically comprise the domains FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. The assignment of amino acids to each domain is typically in accordance with the definitions of Kabat Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, Md. (1987 and 1991)), or Chothia & Lesk *J. Mol. Biol.* 196:901-917 (1987); Chothia et al. *Nature* 342:878-883 (1989).

Bispecific or Bifunctional Antibodies

[0107] A bispecific or bifunctional antibody typically is an artificial hybrid antibody having two different heavy/light chain pairs and two different binding sites. Bispecific antibodies may be produced by a variety of methods

including, but not limited to, fusion of hybridomas or linking of Fab' fragments.

See, e.g., Songsivilai & Lachmann Clin. Exp. Immunol. 79: 315-321 (1990),

Kostelny et al. J. Immunol. 148:1547-1553 (1992).

Preparation of Antibodies

[0108] According to certain embodiments, certain antibodies specifically binding to OPGL are encompassed by the invention. In certain embodiments, the antibodies may be produced by immunization with full-length OPGL, soluble forms of OPGL, or a fragment thereof. In certain embodiments, the antibodies of the invention may be polyclonal or monoclonal, and/or may be recombinant antibodies. In certain embodiments, antibodies of the invention are human antibodies prepared, for example, by immunization of transgenic animals capable of producing human antibodies (see, for example, PCT Published Application No. WO 93/12227).

[0109] In certain embodiments, the complementarity determining regions (CDRs) of the light and heavy chain variable regions of α OPGL-1 may be grafted to framework regions (FRs) from the same, or another, species. In certain embodiments, the CDRs of the light and heavy chain variable regions of α OPGL-1 may be grafted to consensus human FRs. To create consensus human FRs, in certain embodiments, FRs from several human heavy chain or light chain amino acid sequences are aligned to identify a consensus amino acid sequence. In certain embodiments, the FRs of the α OPGL-1 heavy chain or light chain are replaced with the FRs from a different heavy chain or light chain. In

certain embodiments, rare amino acids in the FRs of the heavy and light chains of α OPGL-1 are not replaced, while the rest of the FR amino acids are replaced. Rare amino acids are specific amino acids that are in positions in which they are not usually found in FRs. In certain embodiments, the grafted variable regions from α OPGL-1 may be used with a constant region that is different from the constant region of α OPGL-1. In certain embodiments, the grafted variable regions are part of a single chain Fv antibody. CDR grafting is described, e.g., in U.S. Patent Nos. 6,180,370, 5,693,762, 5,693,761, 5,585,089, and 5,530,101, which are hereby incorporated by reference for any purpose.

[0110] According to certain embodiments, antibodies of the invention are prepared through the utilization of a transgenic mouse that has a substantial portion of the human antibody producing genome inserted but that is rendered deficient in the production of endogenous, murine, antibodies. Such mice, then, are capable of producing human immunoglobulin molecules and antibodies and are deficient in the production of murine immunoglobulin molecules and antibodies. Technologies utilized for achieving this result are disclosed in the patents, applications, and references disclosed in the specification, herein. In certain embodiments, one may employ methods such as those disclosed in PCT Published Application No. WO 98/24893, which is hereby incorporated by reference for any purpose. See also Mendez et al. Nature Genetics 15:146-156 (1997), which is hereby incorporated by reference for any purpose.

[0111] According to certain embodiments, fully human monoclonal antibodies specific for OPGL are produced as follows. Transgenic mice containing human immunoglobulin genes are immunized with the antigen of interest. Lymphatic cells (such as B-cells) from the mice that express antibodies are obtained. Such recovered cells are fused with a myeloid-type cell line to prepare immortal hybridoma cell lines, and such hybridoma cell lines are screened and selected to identify hybridoma cell lines that produce antibodies specific to the antigen of interest. In certain embodiments, the production of a hybridoma cell line that produces antibodies specific to OPGL is provided.

[0112] In certain embodiments, antibodies of the invention are produced by hybridoma lines AMG 6.1, AMG 6.4, AMG 6.5, AMG 7.1, and AMG 7.2. In certain embodiments, antibodies of the invention are produced by hybridoma lines AMG 6.1, AMG 6.4, and AMG 6.5. In certain embodiments, the antibodies of the invention bind to OPGL with a dissociation constant (K_d) of between approximately 0.23 and 0.29 nM. In certain embodiments of the invention, the antibodies bind to OPGL with a K_d of less than 0.23 nM.

[0113] In certain embodiments, the antibodies of the invention are of the IgG2 isotype. In certain embodiments of the invention, the antibodies comprise a human kappa light chain and a human IgG2 heavy chain. In certain embodiments, the antibodies of the invention have been cloned for expression in mammalian cells. In certain embodiments, the variable regions of the antibodies are ligated to a constant region other than the constant region for the IgG2 isotype.

[0114] In certain embodiments, conservative modifications to the heavy and light chains of α OPGL-1 (and corresponding modifications to the encoding nucleotides) will produce antibodies to OPGL having functional and chemical characteristics similar to those of α OPGL-1. In contrast, substantial modifications in the functional and/or chemical characteristics of α OPGL-1 may be accomplished by selecting substitutions in the amino acid sequence of the heavy and light chains that differ significantly in their effect on maintaining (a) the structure of the molecular backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain.

[0115] For example, a "conservative amino acid substitution" may involve a substitution of a native amino acid residue with a nonnative residue such that there is little or no effect on the polarity or charge of the amino acid residue at that position. Furthermore, any native residue in the polypeptide may also be substituted with alanine, as has been previously described for "alanine scanning mutagenesis."

[0116] Desired amino acid substitutions (whether conservative or non-conservative) can be determined by those skilled in the art at the time such substitutions are desired. In certain embodiments, amino acid substitutions can be used to identify important residues of α OPGL-1, or to increase or decrease the affinity of the antibodies to OPGL described herein.

[0117] In certain embodiments, antibodies of the present invention can be expressed in cell lines other than hybridoma cell lines. In certain

embodiments, sequences encoding particular antibodies can be used for transformation of a suitable mammalian host cell. According to certain embodiments, transformation can be by any known method for introducing polynucleotides into a host cell, including, for example packaging the polynucleotide in a virus (or into a viral vector) and transducing a host cell with the virus (or vector) or by transfection procedures known in the art, as exemplified by U.S. Pat. Nos. 4,399,216, 4,912,040, 4,740,461, and 4,959,455 (which patents are hereby incorporated herein by reference for any purpose). In certain embodiments, the transformation procedure used may depend upon the host to be transformed. Methods for introduction of heterologous polynucleotides into mammalian cells are well known in the art and include, but are not limited to, dextran-mediated transfection, calcium phosphate precipitation, polybrene mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei.

[0118] Mammalian cell lines available as hosts for expression are well known in the art and include, but are not limited to, many immortalized cell lines available from the American Type Culture Collection (ATCC), including but not limited to Chinese hamster ovary (CHO) cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), and a number of other cell lines. In certain embodiments, cell lines may be selected through determining which cell lines have high expression levels and produce antibodies with constitutive OPGL binding properties.

[0119] According to certain embodiments, antibodies of the invention are useful for detecting OPGL in biological samples. In certain embodiments, this allows the identification of cells or tissues which produce the protein. In certain embodiments, antibodies which bind to OPGL and block interaction with other binding compounds may have therapeutic use in modulating osteoclast differentiation and bone resorption. In certain embodiments, antibodies to OPGL may block OPGL binding to ODAR, which may result in a block in the signal transduction cascade and loss of NF- κ B mediated transcription activation. Assays for measuring NF- κ B-mediated transcription activation using, e.g., a luciferase reporter assay, are known to those skilled in the art.

[0120] In certain embodiments, methods are provided of treating a bone disorder comprising administering a therapeutically effective amount of an antibody to OPGL. In certain embodiments, methods are provided of treating a bone disorder comprising administering a therapeutically effective amount of an antibody to OPGL and another therapeutic agent. In certain such embodiments, the additional therapeutic agent is administered in a therapeutically effective amount. In certain embodiments, the bone disorder is a disorder characterized by a net bone loss, including but not limited to, osteopenia and osteolysis. In certain embodiments, treatment with an antibody to OPGL is used to suppress the rate of bone resorption. Therefore, in certain embodiments, treatment may be used to reduce the rate of bone resorption where the resorption rate is above normal, or to reduce bone resorption to below normal levels in order to

compensate for below normal levels of bone formation. In certain embodiments, antibodies can be tested for binding to OPGL in the absence or presence of OPG and examined for their ability to inhibit OPGL-mediated osteoclastogenesis and/or bone resorption.

[0121] Conditions which may be treated according to certain embodiments include, but are not limited to, the following:

Osteoporosis, including, but not limited to, primary osteoporosis, endocrine osteoporosis (including, but not limited to, hyperthyroidism, hyperparathyroidism, Cushing's syndrome, and acromegaly), hereditary and congenital forms of osteoporosis (including, but not limited to, osteogenesis imperfecta, homocystinuria, Menkes' syndrome, Riley-Day syndrome), and osteoporosis due to immobilization of extremities;

Paget's disease of bone (osteitis deformans) in adults and juveniles;

Osteomyelitis, i.e., an infectious lesion in bone, leading to bone loss;

Hypercalcemia, including, but not limited to, hypercalcemia resulting from solid tumors (including, but not limited to, breast, lung and kidney) and hematologic malignancies (including, but not limited to, multiple myeloma, lymphoma and leukemia), idiopathic hypercalcemia, and hypercalcemia associated with hyperthyroidism and renal function disorders;

Osteopenia, including but not limited to, osteopenia following surgery, osteopenia induced by steroid administration, osteopenia associated with disorders of the small and large intestine, and osteopenia associated with chronic hepatic and renal diseases;

Osteonecrosis, i.e., bone cell death, including, but not limited to, osteonecrosis associated with traumatic injury, osteonecrosis associated with Gaucher's disease, osteonecrosis associated with sickle cell anemia, osteonecrosis associated with systemic lupus erythematosus, osteonecrosis associated with rheumatoid arthritis, osteonecrosis associated with periodontal disease, osteonecrosis associated with osteolytic metastasis, and osteonecrosis associated with other condition; and

Loss of cartilage and joint erosion associated with rheumatoid arthritis.

[0122] In certain embodiments, an antibody to OPGL may be used alone or with at least one additional therapeutic agents for the treatment of bone disorders. In certain embodiments, an antibody to OPGL is used in conjunction with a therapeutically effective amount of an additional therapeutic agent. Exemplary therapeutic agents that may be administered with an antibody to OPGL include, but are not limited to, the bone morphogenic factors designated BMP-1 through BMP-12; transforming growth factor- β (TGF- β) and TGF- β family members; interleukin-1 (IL-1) inhibitors, including, but not limited to, IL-1ra and derivatives thereof and Kineret™; TNF α inhibitors, including, but not limited to,

soluble TNF α receptors, Enbrel™, anti-TNF α antibodies, Remicade™, and D2E7 antibodies; parathyroid hormone and analogs thereof; parathyroid related protein and analogs thereof; E series prostaglandins; bisphosphonates (such as alendronate and others); bone-enhancing minerals such as fluoride and calcium; non-steroidal anti-inflammatory drugs (NSAIDs), including, but not limited to, COX-2 inhibitors, such as Celebrex™ and Vioxx™; immunosuppressants, such as methotrexate or leflunomide; serine protease inhibitors, including, but not limited to, secretory leukocyte protease inhibitor (SLPI); IL-6 inhibitors (including, but not limited to, antibodies to IL-6), IL-8 inhibitors (including, but not limited to, antibodies to IL-8); IL-18 inhibitors (including, but not limited to, IL-18 binding protein and IL-18 antibodies); Interleukin-1 converting enzyme (ICE) modulators; fibroblast growth factors FGF-1 to FGF-10 and FGF modulators; PAF antagonists; keratinocyte growth factor (KGF), KGF-related molecules, and KGF modulators; matrix metalloproteinase (MMP) modulators; Nitric oxide synthase (NOS) modulators, including, but not limited to, modulators of inducible NOS; modulators of glucocorticoid receptor; modulators of glutamate receptor; modulators of lipopolysaccharide (LPS) levels; and noradrenaline and modulators and mimetics thereof.

[0123] In certain embodiments, an antibody to OPGL is used with particular therapeutic agents to treat various inflammatory conditions, autoimmune conditions, or other conditions with attendant bone loss. In certain embodiments, in view of the condition and the desired level of treatment, two, three, or more agents may be administered. In certain embodiments, such

agents may be provided together by inclusion in the same formulation. In certain embodiments, such agents and an antibody to OPGL may be provided together by inclusion in the same formulation. In certain embodiments, such agents may be provided together by inclusion in a treatment kit. In certain embodiments, such agents and an antibody to OPGL may be provided together by inclusion in a treatment kit. In certain embodiments, such agents may be provided separately. In certain embodiments, when administered by gene therapy, the genes encoding protein agents and/or an antibody to OPGL may be included in the same vector. In certain embodiments, the genes encoding protein agents and/or an antibody to OPGL may be under the control of the same promoter region. In certain embodiments, the genes encoding protein agents and/or an antibody to OPGL may be in separate vectors.

[0124] In certain embodiments, the present invention is directed to therapies comprising an antibody to OPGL and at least one interleukin-1 (IL-1) inhibitor, and methods of treatment using such therapies. In certain embodiments, a therapy comprises an antibody to OPGL and an IL-1 inhibitor and at least one additional molecule described herein. In certain embodiments, methods of treatment use IL-1 inhibitors and/or TNF- α inhibitors in conjunction with an antibody to OPGL. In certain embodiments, an antibody to OPGL in combination with IL-1 inhibitors and/or TNF- α inhibitors may be used for treatment of conditions such as asthma, rheumatoid arthritis, and multiple sclerosis.

[0125] Interleukin-1 (IL-1) is an anti-inflammatory cytokine. In certain instances, IL-1 is a mediator in many diseases and medical conditions. In certain instances, IL-1 is manufactured by cells of the macrophage/monocyte lineage. In certain instances, IL-1 is produced in two forms: IL-1 alpha (IL-1 α) and IL-1 beta (IL-1 β).

[0126] A disease or medical condition is considered to be an "interleukin-1 mediated disease" if the spontaneous or experimental disease or medical condition is associated with elevated levels of IL-1 in bodily fluids or tissue and/or if cells or tissues taken from the body produce elevated levels of IL-1 in culture. In certain embodiments, such interleukin-1 mediated diseases are also recognized by the following additional two conditions: (1) pathological findings associated with the disease or medical condition can be mimicked experimentally in animals by administration of IL-1 or upregulation of expression of IL-1; and (2) a pathology induced in experimental animal models of the disease or medical condition can be inhibited or abolished by treatment with agents that inhibit the action of IL-1. In certain embodiments, one or more of the above conditions are met in an IL-1-mediated disease. In certain embodiments, all three of the conditions are met in an IL-1-mediated disease.

[0127] Acute and chronic interleukin-1 (IL-1) -mediated diseases include, but are not limited to, the following: acute pancreatitis; amyotrophic lateral sclerosis (ALS, or Lou Gehrig's disease); Alzheimer's disease; cachexia/anorexia, including, but not limited to, AIDS-induced cachexia; asthma and other pulmonary diseases; atherosclerosis; autoimmune vasculitis; chronic

fatigue syndrome; Clostridium associated illnesses, including, but not limited to, Clostridium-associated diarrhea; coronary conditions and indications, including, but not limited to, congestive heart failure, coronary restenosis, myocardial infarction, myocardial dysfunction (e.g., related to sepsis), and coronary artery bypass graft; cancer, including, but not limited to, leukemias, including, but not limited to, multiple myeloma leukemia and myelogenous (e.g., AML and CML), and tumor metastasis; diabetes (including, but not limited to, insulin-dependent diabetes); endometriosis; fever; fibromyalgia; glomerulonephritis; graft versus host disease and/or transplant rejection; hemorrhagic shock; hyperalgesia; inflammatory bowel disease; inflammatory conditions of a joint, including, but not limited to, osteoarthritis, psoriatic arthritis, and rheumatoid arthritis; inflammatory eye disease, including, but not limited to, those associated with, for example, corneal transplant; ischemia, including, but not limited to, cerebral ischemia (including, but not limited to, brain injury as a result of, e.g., trauma, epilepsy, hemorrhage or stroke, each of which may lead to neurodegeneration); Kawasaki's disease; learning impairment; lung diseases (including, but not limited to, acute respiratory distress syndrome, or ARDS); multiple sclerosis; myopathies (e.g., muscle protein metabolism, including, but not limited to, muscle protein metabolism in sepsis); neurotoxicity (including, but not limited to, such condition induced by HIV); osteoporosis; pain, including, but not limited to, cancer-related pain; Parkinson's disease; periodontal disease; pre-term labor; psoriasis; reperfusion injury; septic shock; side effects from radiation therapy; temporal mandibular joint disease; sleep disturbance; uveitis; and an

inflammatory condition resulting from, e.g., strain, sprain, cartilage damage, trauma, orthopedic surgery, infection, or other disease processes.

[0128] In certain embodiments, an IL-1 inhibitor may be any protein or molecule capable of specifically preventing activation of cellular receptors to IL-1, which may result from any number of mechanisms. Exemplary mechanisms include, but are not limited to, downregulating IL-1 production, binding free IL-1, interfering with IL-1 binding to its receptor, interfering with formation of the IL-1 receptor complex (i.e., association of IL-1 receptor with IL-1 receptor accessory protein), and interfering with modulation of IL-1 signaling after binding to its receptor.

[0129] Certain interleukin-1 inhibitors include, but are not limited to, IL-1 receptor antagonists, including, but not limited to, Kineret™, IL-1ra, IL-1ra variants, and IL-1ra derivatives, which are collectively termed "IL-1ra proteins;" anti-IL-1 receptor monoclonal antibodies (see, e.g., EP 623674, which is hereby incorporated by reference for any purpose); IL-1 binding proteins, including, but not limited to, soluble IL-1 receptors (see, e.g., U. S. Pat. No. 5,492,888, U. S. Pat. No. 5,488,032, and U. S. Pat. No. 5,464,937, U. S. Pat. No. 5,319,071, and U.S. Pat. No. 5,180,812, which are hereby incorporated by reference for any purpose); anti-IL-1 monoclonal antibodies (see, e.g., WO 9501997, WO 9402627, WO 9006371, U.S.Pat. No. 4,935,343, EP 364778, EP 267611 and EP 220063, which are hereby incorporated by reference for any purpose); IL-1 receptor accessory proteins and antibodies thereto (see, e.g., WO 96/23067 and WO 99/37773, which are hereby incorporated by reference for any purpose);

inhibitors of interleukin-1 beta converting enzyme (ICE) or caspase 1 (see, e.g., WO 99/46248, WO 99/47545, and WO 99/47154, which are hereby incorporated by reference for any purpose), which may be used to inhibit IL-1 beta production and secretion; interleukin-1beta protease inhibitors; and other compounds and proteins that block *in vivo* synthesis or extracellular release of IL-1.

[0130] Exemplary IL-1 inhibitors are disclosed, e.g., in US Pat. Nos. 5,747,444; 5,359,032; 5,608,035; 5,843,905; 5,359,032; 5,866,576; 5,869,660; 5,869,315; 5,872,095; 5,955,480; 5,965,564; International (WO) patent applications 98/21957, 96/09323, 91/17184, 96/40907, 98/32733, 98/42325, 98/44940, 98/47892, 98/56377, 99/03837, 99/06426, 99/06042, 91/17249, 98/32733, 98/17661, 97/08174, 95/34326, 99/36426, 99/36415;. European (EP) patent applications 534978 and 894795; and French patent application FR 2762514. The disclosures of all of the aforementioned references are hereby incorporated by reference for any purpose.

[0131] Interleukin-1 receptor antagonist (IL-1ra) is a human protein that acts as a natural inhibitor of interleukin-1 and is a member of the IL-1 family, which includes IL-1 α and IL-1 β . Certain receptor antagonists, including IL-1ra and variants and derivatives thereof, as well as methods of making and using them, are described in U.S. Patent No. 5,075,222; WO 91/08285; WO 91/17184; AU 9173636; WO 92/16221; WO 93/21946; WO 94/06457; WO 94/21275; FR 2706772; WO 94/21235; DE 4219626, WO 94/20517; WO 96/22793; WO 97/28828; and WO 99/36541, which are incorporated herein by reference for any purpose. In certain embodiments, an IL-1 receptor antagonist may be

glycosylated. In certain embodiments, an IL-1 receptor antagonist may be non-glycosylated.

[0132] Three forms of IL-1ra and variants thereof are described in U.S. Pat. No. 5,075,222 (the '222 patent). The first form, called "IL-1i" in the '222 patent, is characterized as a 22-23 kD molecule on SDS-PAGE with an approximate isoelectric point of 4.8, which elutes from a Mono Q FPLC column at around 52 mM NaCl in Tris buffer, pH 7.6. The second form, IL-1ra β , is characterized as a 22-23 kD protein, which elutes from a Mono Q column at 48 mM NaCl. Both IL-1ra α and IL-1ra β are glycosylated. The third form, IL-1rax, is characterized as a 20 kD protein, which elutes from a Mono Q column at 48 mM NaCl and is non-glycosylated. The '222 patent also describes certain methods for isolating genes that code for the inhibitors, cloning those genes in suitable vectors, transforming and transfecting those genes into certain cell types, and expressing those genes to produce the inhibitors.

[0133] In certain embodiments, deletions, insertions, and/or substitutions (individually or collectively referred to as "variant(s)") are made within the amino acid sequences of IL-1ra. In certain embodiments, an IL-1ra variant is biologically active (e.g., possesses the ability to inhibit IL-1).

[0134] In certain embodiments, the present invention is directed to therapies comprising an antibody to OPGL and at least one TNF α inhibitor, and methods of treatment using such therapies. In certain embodiments, a therapy comprises an antibody to OPGL and a TNF α inhibitor and at least one additional molecule described herein.

[0135] Certain diseases and medical conditions are mediated by TNF and may be categorized as inflammatory conditions. As used herein, a "TNF-mediated disease" includes, but is not limited to, a disease or medical condition that is associated with elevated levels of TNF in bodily fluids or tissue and/or in which cells or tissues taken from the body produce elevated levels of TNF in culture. In certain embodiments, a disease is a TNF-mediated disease if (1) pathological findings associated with the disease or medical condition can be mimicked experimentally in animals by the administration or upregulation of expression of TNF and/or (2) a pathology induced in experimental animal models of the disease or medical condition can be inhibited or abolished by treatment with agents that inhibit the action of TNF.

[0136] Certain acute and chronic TNF-mediated diseases include, but are not limited to: cachexia and anorexia; cancer, including, but not limited to, leukemia; chronic fatigue syndrome; coronary conditions and/or indications, including, but not limited to, congestive heart failure, coronary restenosis, myocardial infarction, myocardial dysfunction (including but not limited to, such condition related to sepsis), and coronary artery bypass graft; depression; diabetes, including, but not limited to, juvenile onset Type 1 diabetes, diabetes mellitus, and insulin resistance (including, but not limited to, insulin resistance associated with obesity); endometriosis, endometritis, and related conditions; fibromyalgia and analgesia; graft versus host rejection; hyperalgesia; inflammatory bowel diseases, including, but not limited to, Crohn's disease and Clostridium difficile-associated diarrhea; ischemia, including, but not limited to,

cerebral ischemia, which includes, but is not limited to, brain injury as a result of trauma, epilepsy, hemorrhage, and/or stroke; lung disease, including, but not limited to, adult respiratory distress syndrome, asthma, and pulmonary fibrosis; multiple sclerosis; neuroinflammatory diseases; ocular diseases and conditions, including, but not limited to, corneal transplant, ocular degeneration and uveitis; pain, including, but not limited to, cancer-related pain; pancreatitis; periodontal diseases; Pityriasis rubra pilaris (PRP); prostatitis, including bacterial and non-bacterial prostatitis, and related conditions; psoriasis and related conditions; pulmonary fibrosis; reperfusion injury; rheumatic diseases, including, but not limited to, rheumatoid arthritis, osteoarthritis, juvenile arthritis (including, but not limited to, juvenile rheumatoid arthritis), seronegative polyarthritis, ankylosing spondylitis, Reiter's syndrome and reactive arthritis, Still's disease, psoriatic arthritis, enteropathic arthritis, polymyositis, dermatomyositis, scleroderma, systemic sclerosis, vasculitis (e.g., Kawasaki's disease), cerebral vasculitis, Lyme disease, staphylococcal-induced ("septic") arthritis, Sjögren's syndrome, rheumatic fever, polychondritis and polymyalgia rheumatica and giant cell arteritis); septic shock; side effects from radiation therapy; systemic lupus erythematosus (SLE); temporal mandibular joint disease; thyroiditis; and tissue transplantation and/or an inflammatory condition, e.g., resulting from strain, sprain, cartilage damage, trauma, orthopedic surgery, infection (e.g., HIV, Clostridium difficile and related species) or other disease process.

[0137] In certain embodiments, TNF inhibitors may act by at least one of downregulating or inhibiting TNF production, binding free TNF, interfering

with TNF binding to its receptor, and interfering with modulation of TNF signaling after binding to its receptor. The term "TNF inhibitor" includes, but is not limited to, solubilized TNF receptors, including, but not limited to, soluble tumor necrosis factor receptor type I (sTNF-RI; also called the p55 receptor), soluble tumor necrosis factor receptor type II (also called the p75 receptor), and Enbrel™; antibodies to TNF, including, but not limited to, Remicade™ and D2E7 (see, e.g., U.S. Patent Nos. 6,090,382 and 6,258,562); antibodies to TNF receptor; sTNF-RI (see, e.g., WO 98/24463), etanercept (Enbrel™), Avakine™; inhibitors of TNF- α converting enzyme (TACE); and other molecules that affect TNF activity.

[0138] Exemplary TNF- α inhibitors are described, e.g., in European patent applications EP 308 378; EP 422 339; EP 393 438; EP 398 327; EP 412 486; EP 418 014, EP 417 563, EP 433 900; EP 464 533; EP 512 528; EP 526 905; EP 568 928; EP 607 776, which describes the use of leflunomide for inhibition of TNF- α ; EP 663 210; EP 542 795; EP 818 439; EP 664 128; EP 542 795; EP 741 707; EP 874 819 ; EP 882 714; EP 880 970; EP 648 783; EP 731 791; EP 895 988; EP 550 376; EP 882 714; EP 853 083; EP 550 376; EP 943 616; EP 939 121; EP 614 984 ; EP 853 083; U.S. Patent Nos. 5,136,021; 5,929,117; 5,948,638; 5,807,862; 5,695,953; 5,834,435; 5,817,822; 5,830,742; 5,834,435; 5,851,556; 5,853,977; 5,359,037; 5,512,544; 5,695,953; 5,811,261; 5,633,145; 5,863,926; 5,866,616; 5,641,673; 5,869,677; 5,869,511; 5,872,146; 5,854,003; 5,856,161; 5,877,222; 5,877,200; 5,877,151; 5,886,010; 5,869,660; 5,859,207; 5,891,883; 5,877,180; 5,955,480; 5,955,476; 5,955,435; 5,994,351; 5,990,119; 5,952,320; 5,962,481; International patent applications WO 90/13575,

WO 91/03553, WO 92/01002, WO 92/13095, WO 92/16221, WO 93/07863, WO 93/21946, WO 93/19777, WO 95/34326, WO 96/28546, WO 98/27298, WO 98/30541, WO 96/38150, WO 96/38150, WO 97/18207, WO 97/15561, WO 97/12902, WO 96/25861, WO 96/12735, WO 96/11209, WO 98/39326, WO 98/39316, WO 98/38859, WO 98/39315, WO 98/42659, WO 98/39329, WO 98/43959, WO 98/45268, WO 98/47863, WO 96/33172, WO 96/20926, WO 97/37974, WO 97/37973, WO 97/47599, WO 96/35711, WO 98/51665, WO 98/43946, WO 95/04045, WO 98/56377, WO 97/12244, WO 99/00364, WO 99/00363, WO 98/57936, WO 99/01449, WO 99/01139, WO 98/56788, WO 98/56756, WO 98/53842, WO 98/52948, WO 98/52937, WO 99/02510, WO 97/43250, WO 99/06410, WO 99/06042, WO 99/09022, WO 99/08688, WO 99/07679, WO 99/09965, WO 99/07704, WO 99/06041, WO 99/37818, WO 99/37625, WO 97/11668, WO 99/50238, WO 99/47672, WO 99/48491; Japanese patent applications 10147531, 10231285, 10259140, and 10130149, 10316570, 11001481, and 127,800/1991; German application no. 19731521; and British application nos. 2 218 101, 2 326 881, 2 246 569. The disclosures of all of the aforementioned references are hereby incorporated by reference for any purpose.

[0139] EP 393 438 and EP 422 339 describe the amino acid and nucleic acid sequences of a soluble TNF receptor type I (also known as sTNFR-I or 30kDa TNF inhibitor) and a soluble TNF receptor type II (also known as sTNFR-II or 40kDa TNF inhibitor), which are collectively termed "sTNFRs". EP 393 438 and EP 422 339 also describe modified forms of sTNFR-I and sTNFR-II,

including, but not limited to fragments, functional derivatives, and variants.

Furthermore, EP 393 438 and EP 422 339 describe methods for isolating genes that code for the inhibitors, cloning the genes into suitable vectors, transforming or transfecting the genes into certain cell types, and expressing the genes to produce the inhibitors.

[0140] sTNFR-I and sTNFR-II are members of the nerve growth factor/TNF receptor superfamily of receptors, which includes the nerve growth factor receptor (NGF), the B cell antigen CD40, 4-1BB, the rat T-cell antigen MRC OX40, the fas antigen, and the CD27 and CD30 antigens (Smith et al. (1990) Science, 248:1019-1023). A conserved feature of that group of cell surface receptors is a cysteine-rich extracellular ligand binding domain, which can be divided into four repeated motifs of about forty amino acids that contain 4-6 cysteine residues at positions that are well conserved (Smith et al. (1990), supra).

[0141] EP 393 438 teaches a 40kDa TNF inhibitor $\Delta 51$ and a 40kDa TNF inhibitor $\Delta 53$, which are truncated versions of the full-length recombinant 40kDa TNF inhibitor protein. $\Delta 51$ and $\Delta 53$ have 51 or 53 amino acids, respectively, deleted from the carboxyl terminus of the mature protein.

[0142] Published PCT Application No. WO 98/01555 describes truncated forms of sTNFR-I and sTNFR-II that do not contain the fourth domain (amino acid residues Thr¹²⁷-Asn¹⁶¹ of sTNFR-I and amino acid residues Pro¹⁴¹-Thr¹⁷⁹ of sTNFR-II); a portion of the third domain (amino acid residues Asn¹¹¹-Cys¹²⁶ of sTNFR-I and amino acid residues Pro¹²³-Lys¹⁴⁰ of sTNFR-II); and,

optionally, do not contain a portion of the first domain (amino acid residues Asp¹-Cys¹⁹ of sTNFR-I and amino acid residues Leu¹-Cys³² of sTNFR-II). In certain embodiments, the truncated sTNFRs include the proteins represented by the formula R₁-[Cys¹⁹-Cys¹⁰³]-R₂ and R₄-[Cys³²-Cys¹¹⁵]-R₅. These proteins are truncated forms of sTNFR-I and sTNFR-II, respectively.

[0143] As used herein, "R₁-[Cys¹⁹-Cys¹⁰³]-R₂" represents one or more proteins wherein [Cys¹⁹-Cys¹⁰³] is residues 19 through 103 of sTNFR-I, the sequence of which is provided in Figure 1 of WO 98/01555; wherein R₁ represents a methionylated or nonmethionylated amine group of Cys¹⁹ or one or more amino-terminal amino acid residues selected from Cys¹⁸ to Asp¹; and wherein R₂ represents a carboxy group of Cys¹⁰³ or one or more carboxy-terminal amino acid residues selected from Phe¹⁰⁴ to Leu¹¹⁰.

[0144] Exemplary truncated sTNFR-I's of the present invention include, but are not limited to, sTNFR-I 2.6D/C105, sTNFR-I 2.6D/C106, sTNFR-I 2.6D/N105, sTNFR-I 2.3D/d8, sTNFR-I 2.3D/d18, sTNFR-I 2.3D/d15, either methionylated or nonmethionylated, and variants and derivatives thereof. Certain exemplary truncated sTNFR-I's are described, e.g., in published PCT Application No. WO 98/01555.

[0145] As used herein, "R₃-[Cys³²-Cys¹¹⁵]-R₄" represents one or more proteins wherein [Cys³²-Cys¹¹⁵] is residues Cys³² through Cys¹¹⁵ of sTNFR-II, the sequence of which is provided in Figure 8 of WO 98/01555; wherein R₃ represents a methionylated or nonmethionylated amine group of Cys³² or one or more amino-terminal amino acid residues selected from Cys³¹ to

Leu¹; and wherein R₄ represents a carboxy group of Cys¹¹⁵ or one or more carboxy-terminal amino acid residues selected from Ala¹¹⁶ to Arg¹²².

[0146] In certain embodiments, the present invention is directed to therapies comprising an antibody to OPGL and at least one serine protease inhibitor, and methods of treatment using such therapies. In certain embodiments, a therapy comprises an antibody to OPGL and a serine protease inhibitor and at least one additional molecule described herein.

[0147] Endogenous proteolytic enzymes may degrade invading organisms, antigen-antibody complexes, and certain tissue proteins that are no longer necessary or useful. Infective agents may introduce additional proteolytic enzymes into the organism. Protease inhibitors may regulate both endogenous and invading proteolytic enzymes.

[0148] In certain embodiments, naturally occurring protease inhibitors serve to control endogenous proteases by limiting their reactions locally and temporally. In certain embodiments, protease inhibitors may inhibit proteases introduced into the body by infective agents. In certain instances, tissues that are particularly prone to proteolytic attack and infection, including, but not limited to, those of the respiratory tract, are rich in protease inhibitors.

[0149] Protease inhibitors comprise approximately 10% of the human plasma proteins. At least eight inhibitors have been isolated from this source and characterized in the literature. These include, but are not limited to, alpha 2-macroglobulin (alpha 2M), alpha 1-protease inhibitor (alpha 1PI), alpha ...

1-antichymotrypsin (alpha 1Achy), alpha 1-anticollagenase (alpha 1AC), and inter-alpha-trypsin inhibitor (I-alpha-I).

[0150] In certain instances, a disturbance of the protease/protease inhibitor balance can lead to protease-mediated tissue destruction, including, but not limited to, emphysema, arthritis, glomerulonephritis, periodontitis, muscular dystrophy, tumor invasion, and various other pathological conditions. In certain situations, e.g. severe pathological processes such as sepsis or acute leukemia, the amount of free proteolytic enzymes present may increase due to the release of enzyme from secretory cells.

[0151] Furthermore, in certain instances, a diminished regulating inhibitor capacity of the organism may also cause alterations in the protease/protease inhibitor balance. A nonlimiting example of such a diminished regulating inhibitor capacity is an alpha 1-protease inhibitor deficiency, which is correlated with the development of pulmonary emphysema.

[0152] In certain instances, serious damage to the organism can occur when such aberrant conditions are present unless measures can be taken to control the proteolytic enzymes. Therefore, protease inhibitors have been sought that can be administered to an organism to control proteolytic enzymes.

[0153] Leukocyte elastase, trypsin, cathepsin G, and pancreatic elastase are nonlimiting examples of a class of proteases known as serine proteases.

[0154] In certain instances, leukocyte elastase, when released extracellularly, degrades connective tissue and other valuable proteins. While a

normally functioning organism degrades a certain amount of connective tissue and other proteins, the presence of an excessive amount of leukocyte elastase may be associated with various pathological states, including, but not limited to, emphysema and rheumatoid arthritis. In certain embodiments, to counteract the effects of leukocyte elastase when it is present in amounts greater than normal, a protease inhibitor has been sought which is specific for leukocyte elastase. Such a protease inhibitor may be useful if it were capable of being isolated or prepared in a purified form and in sufficient quantities to be pharmaceutically useful.

[0155] Certain leukocyte elastase inhibitors are described, e.g., in Schiessler et al., "Acid-Stable Inhibitors of Granulocyte Neutral Proteases in Human Mucous Secretions: Biochemistry and Possible Biological Function", in Neutral Proteases of Human Polymorphonuclear Leucocytes, Havemann et al. (eds), Urban and Schwarzenberg, Inc. (1978), and in Travis and Salvesen, Ann. Rev. Biochem. 52: 655-709 (1983).

[0156] In certain instances, trypsin initiates degradation of certain soft organ tissue, such as pancreatic tissue, during a variety of acute conditions, including, but not limited to, pancreatitis. A trypsin inhibitor may be useful if it could be isolated and prepared in a purified form and in sufficient quantities to be pharmaceutically useful.

[0157] Cathepsin G is another protease present in leukocytes. In certain embodiments, cathepsin G is capable of degrading a variety of proteins *in vitro*, including those of the complement pathway. Pancreatic elastase is another

protease that may have a role in pancreatitis. Thus, inhibitors for these proteases may also be of pharmaceutical value.

[0158] In certain embodiments, the substrate specificity and sensitivity to different inhibitors of serine proteases are believed to result from changes in only a few amino acid residues. By analogy, it may be possible to conceive of a class of serine protease inhibitors in which changes in a relatively few amino acids will result in inhibition of different proteases. In certain embodiments, a member of this class inhibits every serine protease.

[0159] An exemplary serine protease inhibitor is secretory leukocyte protease inhibitor (SLPI) and fragments and analogs thereof. Exemplary serine protease inhibitors also include, but are not limited to, anti-leukoprotease (ALP), mucous protease inhibitor (MPI), human seminal plasma inhibitor-I (HUSI-I), bronchial mucus inhibitor (BMI), and cervical mucus inhibitor (CUSI). In certain embodiments, a serine protease inhibitors also may be LPS modulator. See, e.g., Jin et al. (1997), Cell 88(3): 417-26. In certain embodiments, these molecules are well-suited for use in conditions leading to bone loss because they are preferentially directed to the cartilage.

[0160] Exemplary serine protease inhibitors are described, e.g., in U.S. Pat. No. 4,760,130; U.S. Pat. No. 5,900,400; and U.S. Pat. No. 5,633,227; which are herein incorporated by reference for any purpose. The molecules disclosed in the foregoing references as well as any variants or analogues thereof are collectively termed "serine protease inhibitors."

[0161] IL-18 is a pro-inflammatory cytokine that was found to induce interferon- γ and was previously named interferon gamma inducing factor (IGIF). In certain instances, IL-1 has been shown to upregulate IL-18 production, and IL-18 induces production of a number of proinflammatory cytokines, including IL-6 and MMP-1. See, e.g., Dinarello et al. (1998), J. Leukocyte Biol. 63: 658-64. In certain instances, caspase 1 is also important for IL-18 production. Experiments also suggest that TNF- α regulates IL-18 production, and that simultaneous inhibition of TNF- α and IL-18 protects against liver toxicity. See, e.g., Faggioni et al. (2000), PNAS 97: 2367-72.

[0162] IL-18 acts *in vivo* through a receptor system reminiscent of the IL-1 system. IL-18 interacts with a cell surface receptor (IL-18R), which interacts with an accessory protein (IL-18RAcP). IL-18-mediated signaling proceeds upon formation of the complex of IL-18, IL-18R, and IL-18RAcP. A natural inhibitor for IL-18 is IL-18bp. In certain embodiments, IL-18bp acts as a "decoy receptor" by binding to IL-18 molecules and preventing interaction with IL-18R.

[0163] In certain embodiments, the present invention is directed to therapies comprising an antibody to OPGL and at least one IL-18 inhibitor, and methods of treatment using such therapies. In certain embodiments, a therapy comprises an antibody to OPGL and an IL-18 inhibitor and at least one additional molecule described herein. Exemplary conditions that may be treated according to certain embodiments include, but are not limited to, inflammation, autoimmune diseases, IL-1 mediated diseases, and TNF-mediated diseases. Exemplary

conditions that may be treated with an antibody to OPGL and at least one IL-18 inhibitor according to certain embodiments include, but are not limited to, arthritis, including, but not limited to rheumatoid arthritis; systemic lupus erythematosus (SLE); graft versus host disease (GvHD); hepatitis; sepsis; and the loss of bone and cartilage accompanying these diseases.

[0164] Exemplary IL-18 inhibitors include, but are not limited to, antibodies that bind to IL-18; antibodies that bind to IL-18R; antibodies that bind to IL-18RAcP; IL-18bp; IL-18R fragments (e.g., a solubilized extracellular domain of the IL-18 receptor); peptides that bind to IL-18 and reduce or prevent its interaction with IL-18R; peptides that bind to IL-18R and reduce or prevent its interaction with IL-18 or with IL-18RAcP; peptides that bind to IL-18RAcP and reduce or prevent its interaction with IL-18R; and small molecules that reduce or prevent IL-18 production or the interaction between any of IL-18, IL-18R, and IL-18RAcP.

[0165] Certain IL-18 inhibitors are described, e.g., in US Pat. No. 5,912,324, issued July 14, 1994; EP 0 962 531, published Dec. 8, 1999; EP 712 931, published Nov. 15, 1994; US Pat. No. 5,914,253, issued July 14, 1994; WO 97/24441, published July 10, 1997; US Pat. No. 6,060,283, issued May 9, 2000; EP 850 952, published Dec. 26, 1996; EP 864 585, published Sep. 16, 1998; WO 98/41232, published Sep. 24, 1998; US Pat. No. 6,054,487, issued April 25, 2000; WO 99/09063, published Aug 14, 1997; WO 99/22760, published Nov. 3, 1997; WO 99/37772, published Jan. 23, 1998; WO 99/37773, published March 20, 1998; EP 0 974 600, published Jan. 26, 2000; WO 00/12555, published Mar.

9, 2000; Japanese patent application JP 111,399/94, published Oct. 31, 1997; Israel patent application IL 121554 A0, published Feb. 8, 1998; which are incorporated herein by reference for any purpose.

[0166] In certain embodiments, an antibody to OPGL may be used with at least one therapeutic agent for inflammation. In certain embodiments, an antibody to OPGL may be used with at least one therapeutic agent for an immune disorder. Exemplary therapeutic agents for inflammation and immune disorders include, but are not limited to, corticosteroids, including, but not limited to, prednisolone; nonsteroidal anti-inflammatory drugs (NSAIDs), including, but not limited to, cyclooxygenase type 1 (COX-1) and cyclooxygenase type 2 (COX-2) inhibitors; disease modifying antirheumatic drugs (DMARDs), including, but not limited to, methotrexate, hydroxychloroquine, chloroquine, cyclosporine, gold compounds (such as auranofin, aurothiomalate and aurothioglucose), and leflunomide; type IV phosphodiesterase inhibitors, including, but not limited to, Rolipram and Pentoxifylline; Tacrolimus (FK-506); Sirolimus (rapamycin); mycophenolic acid; 5-lipoxygenase inhibitors, including, but not limited to, Zileuton; modulators of interleukin-6 (IL-6); small molecule modulators of 38 kDa mitogen-activated protein kinase (p38-MAPK); small molecule modulators of intracellular molecules involved in inflammation pathways, wherein such intracellular molecules include, but are not limited to, jnk, IKK, NF- κ B, ZAP70, and Ick. Certain exemplary therapeutic agents for inflammation are described, e.g., in C.A. Dinarello and L.L. Moldawer Proinflammatory and Anti-Inflammatory Cytokines in Rheumatoid Arthritis: A Primer for Clinicians Third Edition (2001)

Amgen Inc. Thousand Oaks, CA. Certain exemplary therapeutic agents for inflammation and autoimmune diseases include, but are not limited to, interferon gamma (IFN- γ) modulators; modulators of OX40/OX40L (including soluble forms of OX40); modulators of 4-1BB/4-1BB ligand (including soluble forms of 4-1BB); and modulators of B cell-T cell costimulatory pathways, including, but not limited to, modulators of the receptor ligand pairs CD28/B7, CD40/CD40L, ICOS/B7RP1, and AGP-3/TACI/BAFFR (AGP-3 binds to both TACI and BAFFR receptors). Certain exemplary modulators of B cell-T cell costimulatory pathways include, but are not limited to, inhibitors of CD28, B7.1, and B7.2 (including soluble forms of B7.1 or B7.2 and soluble forms of CTLA4, both of which may be fused to a heterologous peptide or protein which reduces or prevents degradation and/or increases half-life, reduces toxicity, reduces immunogenicity, or increases biological activity of a therapeutic protein by increasing solubility or circulating half-life); inhibitors of CD40 and CD40L (including soluble forms of CD40 which may be fused to a heterologous peptide or protein); inhibitors of ICOS and B7RP1 (including soluble forms of ICOS which may be fused to a heterologous peptide or protein) and inhibitors of AGP-3, TACI and BAFFR (including soluble forms of TACI and BAFFR). ICOS, B7RP1 and inhibitors thereof are described, e.g., in WO00/46240. AGP-3, TACI and BAFFR and inhibitors thereof are described, e.g., in WO00/47740, WO01/85872, WO02/15273, WO98/39361, and von Bulow and Bram (1997) Science 278:138-140.

[0167] In certain embodiments, an antibody to OPGL is used to treat bone loss, including, but not limited to, bone loss resulting from osteolytic destruction of bone caused by malignant or metastatic tumors. In certain embodiments, an antibody to OPGL may be used to treat bone loss associated with cancer. Exemplary cancers include, but are not limited to, breast, prostate, thyroid, kidney, lung, esophageal, rectal, bladder, cervical, ovarian, and liver cancers, as well as cancer of the gastrointestinal tract. In certain embodiments, an antibody to OPGL may be used to treat bone loss associated with, e.g., certain hematological malignancies, including, but not limited to, multiple myeloma and lymphoma, including Hodgkin's Disease.

[0168] In certain embodiments, an antibody to OPGL is administered alone. In certain embodiments, an antibody to OPGL is administered with at least one other therapeutic agent, including, but not limited to, at least one other cancer therapy agent. Exemplary cancer therapy agents include, but are not limited to, radiation therapy and chemotherapy. In certain embodiments, chemotherapy may involve treatment with one or more of the following: anthracyclines, taxol, tamoxifene, doxorubicin, 5-fluorouracil, and other drugs known in the art. In certain embodiments, a cancer therapy agent is a luteinizing hormone-releasing hormone (LHRH) antagonist. In certain embodiments, a LHRH antagonist is a peptide antagonist.

[0169] In certain embodiments, an LHRH antagonist comprises the peptide: Ac-D-Nal-4-Cl-Phe-D-Pal-Ser-N-Me-Tyr-D-Asn-Leu-Lys(iPr)-Pro-D-Ala-NH₂ (SEQ ID NO: 20), where Nal is 3-(2-naphthyl)alaninyl; 4-Cl-Phe is (4'-

chlorophenyl)alaninyl; Pal is 3-(3'-pyridyl)alaninyl; and Lys(iPr) is N-epsilon-2-propyl-lysiny.

[0170] In certain embodiments, an LHRH antagonist is an LHRH antagonist decapeptide. Certain exemplary decapeptides are described, e.g., in U.S. Patent No. 5,843,901, which is herein incorporated by reference for any purpose.

[0171] Exemplary therapeutic antibodies according to certain embodiments include, but are not limited to, mouse, mouse-human chimeric, CDR-grafted, humanized and fully human antibodies, and synthetic antibodies, including, but not limited to, those selected by screening antibody libraries. Exemplary antibodies include, but are not limited to, those which bind to cell surface proteins Her2, CDC20, CDC33, mucin-like glycoprotein, and epidermal growth factor receptor (EGFR) present on tumor cells, and optionally induce a cytostatic and/or cytotoxic effect on tumor cells displaying these proteins. Exemplary antibodies also include HERCEPTIN™ (trastuzumab), which may be used to treat breast cancer and other forms of cancer, and RITUXAN™ (rituximab), ZEVALIN™ (ibritumomab tiuxetan), and LYMPHOCIDE™ (epratuzumab), which may be used to treat non-Hodgkin's lymphoma and other forms of cancer. Certain exemplary antibodies also include ERBITUX™ (IMC-C225), BEXXAR™ (iodine 131 tositumomab), and Campath.

[0172] In certain embodiments, cancer therapy agents are polypeptides which selectively induce apoptosis in tumor cells, including, but not limited to, the TNF-related polypeptide TRAIL. In certain embodiments, an

antibody to OPGL may be administered at least one of prior to, concurrent with, and subsequent to treatment with a cancer therapy agent. In certain embodiments, an antibody to OPGL may be administered prophylactically to prevent or mitigate the onset of bone loss by metastatic cancer. In certain embodiments, an antibody to OPGL may be administered for the treatment of an existing condition of bone loss due to metastasis.

[0173] In certain embodiments, an antibody to OPGL may be used to prevent and/or treat bone loss associated with multiple myeloma and/or to prevent and/or treat the disease itself. Multiple myeloma is a B cell derived tumor that may result in significant morbidity and/or mortality. In certain instances, a clinical manifestation of multiple myeloma is focal bone loss, which may be due to increased osteoclast activation in localized regions. Many myeloma patients present with bone lesions visible by radiological analysis and suffer from skeletal pain. In certain instances, patients with myeloma are susceptible to pathological fractures of involved bone, which may occur either spontaneously or due to injury. In certain instances, the skeletal lesions that occur during myeloma not only lead to bone fractures, but also deformity and occasionally nerve compression, particularly in the vertebral spine. In some patients, a pathological increase in serum calcium (hypercalcemia) occurs, and may cause significant problems during disease treatment. In certain embodiments, an antibody to OPGL may be administered to patients to reduce or block bone resorption and release of calcium, which may reduce the risk of fractures and spinal deformity.

[0174] In certain instances, myeloma cells do not directly participate in bone destruction, but instead produce extracellular signals that lead to osteoclast differentiation and activation. In certain instances, osteoclasts produce high levels of the cytokine IL-6, particularly when they become activated. IL-6 is a B-cell growth factor, and contributes to the growth of both murine and human myeloma cells *in vitro*. Myeloma cells may also either directly or indirectly produce OPGL, which may result in local bone lysis surrounding the myeloma cells embedded in bone marrow spaces. In certain instances, the normal osteoclasts adjacent to the myeloma cell in turn produce IL-6, which may lead to local expansion of the tumor cells. Myeloma cells expand in a clonal fashion and may occupy bone spaces that are created by inappropriate bone resorption.

[0175] It has been observed that OPG administration in rodents induces rapid death of the osteoclast population (see, e.g., Lacey et al. (2000) Am. J. Pathol. 157:435-448). A reduction in the number of osteoclasts may counteract the effect of increased IL-6 production by those cells and may therefore affect the growth and survival of myeloma cells within trabecular bone. Thus, in certain embodiments, administration of an antibody to OPGL to a myeloma patient may not only block the hyper resorption of bone, but may also affect the expansion and survival of the tumor itself.

[0176] B-cells express the receptor for OPGL, ODAR. Myeloma cells also express ODAR, and in addition may produce OPGL. In certain instances, the expression of both OPGL and ODAR in the same cell population may create an autocrine stimulus that affects survival of the myeloma cell. Thus,

in certain embodiments, administration of an antibody to OPGL may reduce tumor cell survival, thereby decreasing or eliminating the tumor burden seen in myeloma patients.

[0177] In certain embodiments, the invention provides for pharmaceutical compositions comprising a therapeutically effective amount of an antibody to OPGL together with a pharmaceutically acceptable diluent, carrier, solubilizer, emulsifier, preservative and/or adjuvant.

[0178] In certain embodiments, the invention provides for pharmaceutical compositions comprising a therapeutically effective amount of an antibody to OPGL and a therapeutically effective amount of at least one additional therapeutic agents, together with a pharmaceutically acceptable diluent, carrier, solubilizer, emulsifier, preservative and/or adjuvant. In certain embodiments, the at least one additional therapeutic agent is selected from bone morphogenic factors designated BMP-1 through BMP-12; transforming growth factor- β (TGF- β) and TGF- β family members; interleukin-1 (IL-1) inhibitors, including, but not limited to, IL-1ra and derivatives thereof and Kineret™; TNF α inhibitors, including, but not limited to, a soluble TNF α receptor, Enbrel™, anti-TNF α antibodies, Remicade™, and D2E7 antibody; parathyroid hormone and analogs thereof, parathyroid related protein and analogs thereof; E series prostaglandins; bisphosphonates (such as alendronate and others); bone-enhancing minerals such as fluoride and calcium; non-steroidal anti-inflammatory drugs (NSAIDs), including COX-2 inhibitors, such as Celebrex™ and Vioxx™; immunosuppressants, such as methotrexate or leflunomide; serine protease

inhibitors such as secretory leukocyte protease inhibitor (SLPI); IL-6 inhibitors (e.g., antibodies to IL-6), IL-8 inhibitors (e.g., antibodies to IL-8); IL-18 inhibitors (e.g., IL-18 binding protein or IL-18 antibodies); Interleukin-1 converting enzyme (ICE) modulators; fibroblast growth factors FGF-1 to FGF-10 and FGF modulators; PAF antagonists; keratinocyte growth factor (KGF), KGF-related molecules, or KGF modulators; matrix metalloproteinase (MMP) modulators; Nitric oxide synthase (NOS) modulators, including modulators of inducible NOS; modulators of glucocorticoid receptor; modulators of glutamate receptor; modulators of lipopolysaccharide (LPS) levels; and noradrenaline and modulators and mimetics thereof.

[0179] In certain embodiments, acceptable formulation materials preferably are nontoxic to recipients at the dosages and concentrations employed.

[0180] In certain embodiments, the pharmaceutical composition may contain formulation materials for modifying, maintaining or preserving, for example, the pH, osmolarity, viscosity, clarity, color, isotonicity, odor, sterility, stability, rate of dissolution or release, adsorption or penetration of the composition. In certain embodiments, suitable formulation materials include, but are not limited to, amino acids (such as glycine, glutamine, asparagine, arginine or lysine); antimicrobials; antioxidants (such as ascorbic acid, sodium sulfite or sodium hydrogen-sulfite); buffers (such as borate, bicarbonate, Tris-HCl, citrates, phosphates or other organic acids); bulking agents (such as mannitol or glycine); chelating agents (such as ethylenediamine tetraacetic acid (EDTA)); complexing

agents (such as caffeine, polyvinylpyrrolidone, beta-cyclodextrin or hydroxypropyl-beta-cyclodextrin); fillers; monosaccharides; disaccharides; and other carbohydrates (such as glucose, mannose or dextrans); proteins (such as serum albumin, gelatin or immunoglobulins); coloring, flavoring and diluting agents; emulsifying agents; hydrophilic polymers (such as polyvinylpyrrolidone); low molecular weight polypeptides; salt-forming counterions (such as sodium); preservatives (such as benzalkonium chloride, benzoic acid, salicylic acid, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid or hydrogen peroxide); solvents (such as glycerin, propylene glycol or polyethylene glycol); sugar alcohols (such as mannitol or sorbitol); suspending agents; surfactants or wetting agents (such as pluronics, PEG, sorbitan esters, polysorbates such as polysorbate 20, polysorbate 80, triton, tromethamine, lecithin, cholesterol, tyloxapal); stability enhancing agents (such as sucrose or sorbitol); tonicity enhancing agents (such as alkali metal halides, preferably sodium or potassium chloride, mannitol sorbitol); delivery vehicles; diluents; excipients and/or pharmaceutical adjuvants. (*Remington's Pharmaceutical Sciences*, 18th Edition, A.R. Gennaro, ed., Mack Publishing Company (1990).

[0181] In certain embodiments, an antibody to OPGL and/or a therapeutic molecule is linked to a half-life extending vehicle known in the art. Such vehicles include, but are not limited to, the Fc domain, polyethylene glycol, and dextran. Such vehicles are described, e.g., in U.S. Application Serial No. 09/428,082 and published PCT Application No. WO 99/25044, which are hereby incorporated by reference for any purpose.

[0182] In certain embodiments, the optimal pharmaceutical composition will be determined by one skilled in the art depending upon, for example, the intended route of administration, delivery format and desired dosage. See, for example, *Remington's Pharmaceutical Sciences, supra*. In certain embodiments, such compositions may influence the physical state, stability, rate of *in vivo* release and rate of *in vivo* clearance of the antibodies of the invention.

[0183] .In certain embodiments, the primary vehicle or carrier in a pharmaceutical composition may be either aqueous or non-aqueous in nature. For example, in certain embodiments, a suitable vehicle or carrier may be water for injection, physiological saline solution or artificial cerebrospinal fluid, possibly supplemented with other materials common in compositions for parenteral administration. In certain embodiments, neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles. In certain embodiments, pharmaceutical compositions comprise Tris buffer of about pH 7.0-8.5, or acetate buffer of about pH 4.0-5.5, which may further include sorbitol or a suitable substitute therefor. In certain embodiments, a composition comprising an antibody to OPGL, with or without at least one additional therapeutic agents, may be prepared for storage by mixing the selected composition having the desired degree of purity with optional formulation agents (*Remington's Pharmaceutical Sciences, supra*) in the form of a lyophilized cake or an aqueous solution. Further, in certain embodiments, a composition comprising an antibody to OPGL,

with or without at least one additional therapeutic agents, may be formulated as a lyophilizate using appropriate excipients such as sucrose.

[0184] In certain embodiments, the pharmaceutical compositions of the invention can be selected for parenteral delivery. In certain embodiments, the compositions may be selected for inhalation or for delivery through the digestive tract, such as orally. The preparation of such pharmaceutically acceptable compositions is within the skill of the art.

[0185] In certain embodiments, the formulation components are present in concentrations that are acceptable to the site of administration. In certain embodiments, buffers are used to maintain the composition at physiological pH or at a slightly lower pH, typically within a pH range of from about 5 to about 8.

[0186] In certain embodiments, when parenteral administration is contemplated, a therapeutic composition may be in the form of a pyrogen-free, parenterally acceptable aqueous solution comprising the desired antibody to OPGL, with or without additional therapeutic agents, in a pharmaceutically acceptable vehicle. In certain embodiments, a vehicle for parenteral injection is sterile distilled water in which the antibody to OPGL, with or without at least one additional therapeutic agent, is formulated as a sterile, isotonic solution, properly preserved. In certain embodiments, the preparation can involve the formulation of the desired molecule with an agent, such as injectable microspheres, bio-erodible particles, polymeric compounds (such as polylactic acid or polyglycolic acid), beads or liposomes, that may provide for the controlled or sustained

release of the product which may then be delivered via a depot injection. In certain embodiments, hyaluronic acid may also be used, and may have the effect of promoting sustained duration in the circulation. In certain embodiments, implantable drug delivery devices may be used to introduce the desired molecule.

[0187] In certain embodiments, a pharmaceutical composition may be formulated for inhalation. In certain embodiments, an antibody to OPGL, with or without at least one additional therapeutic agent, may be formulated as a dry powder for inhalation. In certain embodiments, an inhalation solution comprising an antibody to OPGL, with or without at least one additional therapeutic agent, may be formulated with a propellant for aerosol delivery. In certain embodiments, solutions may be nebulized. Pulmonary administration is further described in PCT application no. PCT/US94/001875, which describes pulmonary delivery of chemically modified proteins.

[0188] In certain embodiments, it is contemplated that formulations may be administered orally. In certain embodiments, an antibody to OPGL, with or without at least one additional therapeutic agents, that is administered in this fashion may be formulated with or without those carriers customarily used in the compounding of solid dosage forms such as tablets and capsules. In certain embodiments, a capsule may be designed to release the active portion of the formulation at the point in the gastrointestinal tract when bioavailability is maximized and pre-systemic degradation is minimized. In certain embodiments, at least one additional agent can be included to facilitate absorption of the

antibody to OPGL and/or any additional therapeutic agents. In certain embodiments, diluents, flavorings, low melting point waxes, vegetable oils, lubricants, suspending agents, tablet disintegrating agents, and binders may also be employed.

[0189] In certain embodiments, a pharmaceutical composition may involve an effective quantity of antibodies to OPGL, with or without at least one additional therapeutic agents, in a mixture with non-toxic excipients which are suitable for the manufacture of tablets. In certain embodiments, by dissolving the tablets in sterile water, or another appropriate vehicle, solutions may be prepared in unit-dose form. In certain embodiments, suitable excipients include, but are not limited to, inert diluents, such as calcium carbonate, sodium carbonate or bicarbonate, lactose, or calcium phosphate; or binding agents, such as starch, gelatin, or acacia; or lubricating agents such as magnesium stearate, stearic acid, or talc.

[0190] Additional pharmaceutical compositions will be evident to those skilled in the art, including formulations involving antibodies to OPGL, with or without at least one additional therapeutic agents, in sustained- or controlled-delivery formulations. In certain embodiments, techniques for formulating a variety of other sustained- or controlled-delivery means, such as liposome carriers, bio-erodible microparticles or porous beads and depot injections, are also known to those skilled in the art. See for example, PCT Application No. PCT/US93/00829 which describes the controlled release of porous polymeric microparticles for the delivery of pharmaceutical compositions. In certain

embodiments, sustained-release preparations may include semipermeable polymer matrices in the form of shaped articles, e.g. films, or microcapsules. Sustained release matrices may include polyesters, hydrogels, polylactides (U.S. 3,773,919 and EP 058,481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman *et al.*, *Biopolymers*, 22:547-556 (1983)), poly (2-hydroxyethyl-methacrylate) (Langer *et al.*, *J. Biomed. Mater. Res.*, 15:167-277 (1981) and Langer, *Chem. Tech.*, 12:98-105 (1982)), ethylene vinyl acetate (Langer *et al.*, *supra*) or poly-D(-)-3-hydroxybutyric acid (EP 133,988). In certain embodiments, sustained release compositions may also include liposomes, which can be prepared by any of several methods known in the art. See e.g., Eppstein *et al.*, *Proc. Natl. Acad. Sci. USA*, 82:3688-3692 (1985); EP 036,676; EP 088,046 and EP 143,949.

[0191] The pharmaceutical composition to be used for *in vivo* administration typically is sterile. In certain embodiments, this may be accomplished by filtration through sterile filtration membranes. In certain embodiments, where the composition is lyophilized, sterilization using this method may be conducted either prior to or following lyophilization and reconstitution. In certain embodiments, the composition for parenteral administration may be stored in lyophilized form or in a solution. In certain embodiments, parenteral compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

[0192] In certain embodiments, once the pharmaceutical composition has been formulated, it may be stored in sterile vials as a solution, suspension, gel, emulsion, solid, or as a dehydrated or lyophilized powder. In certain embodiments, such formulations may be stored either in a ready-to-use form or in a form (e.g., lyophilized) that is reconstituted prior to administration.

[0193] In certain embodiments, the present invention is directed to kits for producing a single-dose administration unit. In certain embodiments, the kits may each contain both a first container having a dried protein and a second container having an aqueous formulation. In certain embodiments of this invention, kits containing single and multi-chambered pre-filled syringes (e.g., liquid syringes and lyosyringes) are included.

[0194] In certain embodiments, the effective amount of a pharmaceutical composition comprising an antibody to OPGL, with or without at least one additional therapeutic agent, to be employed therapeutically will depend, for example, upon the therapeutic context and objectives. One skilled in the art will appreciate that the appropriate dosage levels for treatment, according to certain embodiments, will thus vary depending, in part, upon the molecule delivered, the indication for which the antibody to OPGL, with or without at least one additional therapeutic agent, is being used, the route of administration, and the size (body weight, body surface or organ size) and/or condition (the age and general health) of the patient. In certain embodiments, the clinician may titer the dosage and modify the route of administration to obtain the optimal therapeutic effect. In certain embodiments, a typical dosage may range from about 0.1 $\mu\text{g/kg}$

to up to about 100 mg/kg or more, depending on the factors mentioned above. In certain embodiments, the dosage may range from 0.1 µg/kg up to about 100 mg/kg; or 1 µg/kg up to about 100 mg/kg; or 5 µg/kg up to about 100 mg/kg.

[0195] In certain embodiments, the frequency of dosing will take into account the pharmacokinetic parameters of the antibody to OPGL and/or any additional therapeutic agents in the formulation used. In certain embodiments, a clinician will administer the composition until a dosage is reached that achieves the desired effect. In certain embodiments, the composition may therefore be administered as a single dose, or as two or more doses (which may or may not contain the same amount of the desired molecule) over time, or as a continuous infusion via an implantation device or catheter. Further refinement of the appropriate dosage is routinely made by those of ordinary skill in the art and is within the ambit of tasks routinely performed by them. In certain embodiments, appropriate dosages may be ascertained through use of appropriate dose-response data.

[0196] In certain embodiments, the route of administration of the pharmaceutical composition is in accord with known methods, e.g. orally, through injection by intravenous, intraperitoneal, intracerebral (intra-parenchymal), intracerebroventricular, intramuscular, intra-ocular, intraarterial, intraportal, or intralesional routes; by sustained release systems or by implantation devices. In certain embodiments, the compositions may be administered by bolus injection or continuously by infusion, or by implantation device.

[0197] In certain embodiments, the composition may be administered locally via implantation of a membrane, sponge or another appropriate material onto which the desired molecule has been absorbed or encapsulated. In certain embodiments, where an implantation device is used, the device may be implanted into any suitable tissue or organ, and delivery of the desired molecule may be via diffusion, timed-release bolus, or continuous administration.

[0198] In certain embodiments, it may be desirable to use a pharmaceutical composition comprising an antibody to OPGL, with or without at least one additional therapeutic agent, in an *ex vivo* manner. In such instances, cells, tissues and/or organs that have been removed from the patient are exposed to a pharmaceutical composition comprising an antibody to OPGL, with or without at least one additional therapeutic agent, after which the cells, tissues and/or organs are subsequently implanted back into the patient.

[0199] In certain embodiments, an antibody to OPGL and/or any additional therapeutic agents can be delivered by implanting certain cells that have been genetically engineered, using methods such as those described herein, to express and secrete the polypeptides. In certain embodiments, such cells may be animal or human cells, and may be autologous, heterologous, or xenogeneic. In certain embodiments, the cells may be immortalized. In certain embodiments, in order to decrease the chance of an immunological response, the cells may be encapsulated to avoid infiltration of surrounding tissues. In certain embodiments, the encapsulation materials are typically biocompatible,

semi-permeable polymeric enclosures or membranes that allow the release of the protein product(s) but prevent the destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissues.

EXAMPLES

[0200] The following examples, including the experiments conducted and results achieved are provided for illustrative purposes only and are not to be construed as limiting the present invention.

Example 1

Cloning the α OPGL-1 Heavy and Light Chains

[0201] CHO cells expressing the full-length human OPGL cDNA are used to immunize transgenic mice containing human immunoglobulin genes. Lymph nodes from the immunized mice are fused to murine myeloma cells to generate hybridomas. Supernatants from the hybridoma lines are tested in an ELISA assay for antibodies that react with human OPGL. Anti-OPGL expressing hybridoma lines AMG 6.5, AMG 6.4, and AMG 6.1 are found to express antibodies with high affinity for OPGL (K_d's of 0.28 nM, 0.29 nM, and 0.23 nM, respectively), and AMG 6.5 is selected for cloning. Heavy and light chain cDNA clones from AMG 6.5 and AMG 6.4 are identical, and AMG 6.5 is used to clone the α OPGL-1 light chain cDNA, while AMG 6.4 is used to clone the α OPGL-1 heavy chain cDNA.

Cloning of the α OPGL-1 light chain

[0202] The α OPGL-1 kappa light chain variable region is obtained using PCR amplification methods from first strand cDNA prepared from AMG 6.5 total RNA. First strand cDNA is prepared from AMG 6.5 total RNA using a random primer with an extended 5'-adapter (5'-GGCCGGATAGGCCTCACNNNNNNT-3' (SEQ ID NO: 15)) and the materials and methods provided by the Gibco SuperScript II TM Preamplification System for First Strand cDNA Synthesis kit (cat. no. 18089-011). The oligonucleotides below are used for the PCR:

5' kappa RACE primer:

5' - GAT GAC CCA GTC TCC AGC CAC CCT G - 3' (SEQ ID NO: 5)

3' kappa RACE primer:

5' - AAG GGT CAG AGG CCA AAG GAT GG - 3' (SEQ ID NO: 6)

[0203] The amplified DNAs are cloned into pCRII-TOPO (Invitrogen) and the resulting plasmids are sequenced. The kappa chain consensus sequence is used to design primers for PCR amplification of the full-length α OPGL-1 kappa chain. The 5' α OPGL-1 kappa primer incorporates a *Xba*I site (TCTAGA) for cloning and a "CCACC" Kozak sequence before the initiator Met codon. The 3' α OPGL-1 kappa primer incorporates a *Sal*I site (GTCGAC) following the stop codon for cloning.

5' α OPGL-1 kappa primer:

5'-CAA CTC TAG A CC ACC ATG GAA ACC CCA GCG-3' (SEQ ID NO: 7)

*Xba*I Site Kozak M E T P A (SEQ ID NO: 16)

3' α OPGL-1 kappa primer:

5'-TTT GAC GTC GAC TTA TCA ACA CTC TCC CCT GTT GAA G -3' (SEQ ID NO: 8)

SaII Site * * C E G R N F (SEQ ID NO: 17)

[0204] The full-length α OPGL-1 kappa chain cDNA clone is obtained using the AMG 6.5 first strand cDNA, described above, by PCR amplification with the 5' and 3' α OPGL-1 kappa primers. The PCR reaction generates a 738 bp fragment encoding the 235 amino acid residues (including the 20 amino acid kappa chain signal sequence) of the α OPGL-1 kappa chain (Figure 4, SEQ ID NO: 4). Following purification using a QIAquick PCR Purification kit (Qiagen cat. no.28104), this fragment is used to construct the kappa light chain expression vector.

[0205] The 738 bp full-length kappa fragment generated above is cut with *Xba*I and *Sa*II, is purified using the Promega Wizard DNA Clean-Up System (Promega cat. no. A7100), and is cloned into pDSR α 19 to generate plasmid α OPGL-1-kappa/pDSR α 19 (Figure 5). pDSR α 19 has been described previously (see WO 90/14363, which is herein incorporated by reference for any purpose (see, e.g., Figure 12)). Briefly, to make pDSR α 19, pDSR α 2 is modified in the following ways: the FSH polyA is shortened by approximately 1400 base pairs, to 885 base pairs, and now ends at the *Nde*I site; the dihydrofolate reductase (DHFR) promoter now contains 209 base pairs, having been shortened from the 5' end by approximately 1 kilobase; and an approximately 550 base pair *Bgl*II fragment from the DHFR polyA sequence is deleted.

[0206] The α OPGL-1 kappa light chain expression clone is sequenced to confirm that it coded for the same peptide that is identified in the AMG 6.5 hybridoma. The final expression vector, α OPGL-1-kappa/pDSR α 19, is 5476 bp and contains the 7 functional regions shown in Table 2.

Table 2: Features of α OPGL-1-kappa/pDSR α 19

Plasmid Base

Pair Number:

2 to 881	A transcription termination/polyadenylation signal from the α -subunit of the bovine pituitary glycoprotein hormone (α -FSH) (Goodwin, et al, <i>Nucleic Acids Res.</i> 1983 11 : 6873-82; Genbank Accession Number X00004)
882 to 2027	A mouse dihydrofolate reductase (DHFR) minigene containing the endogenous mouse DHFR promoter, the cDNA coding sequences, and the DHFR transcription termination/polyadenylation signals (Gasser et al, <i>Proc Natl Acad Sci U S A.</i> 1982 79 :6522-6.; Nunberg et al, <i>Cell</i> 1980 19 :355-64; Setzer et al, <i>J Biol Chem.</i> 1982 257 : 5143-7; McGrogan et al, <i>J Biol Chem.</i> 1985 260 : 2307-14)
2031 to 3947	pBR322 sequences containing the ampicillin resistance marker gene and the origin for replication of the plasmid in <i>E. coli</i> (Genbank Accession Number J01749)
3949 to 4292	An SV40 early promoter, enhancer and origin of replication (Takebe et al, <i>Mol Cell Biol.</i> 1988 8 : 466-72., Genbank Accession Number J02400)
4299 to 4565	A translational enhancer element from the HTLV-1 LTR domain (Seiki et al, <i>Proc Natl Acad Sci U S A.</i> 1983 80 : 3618-22, Genbank Accession Number J02029)
4574 to 4730	An intron from the SV40 16S, 19S splice donor/acceptor signals (Okayama and Berg, <i>Mol Cell Biol.</i> 1983 3 : 280-9, Genbank Accession Number J02400)
4750 to 5476	The α OPGL-1 Kappa light chain cDNA between the <i>Xba</i> I and <i>Sal</i> I sites

[0207] A circular plasmid map of the vector is shown in Figure 5.

Cloning of the α OPGL-1 heavy chain

[0208] The α OPGL-1 IgG2 heavy chain is cloned from AMG 6.4 hybridoma double-stranded cDNA produced with the Clontech Marathon™ cDNA Amplification Kit (cat. no. K1802-1). Amplification of AMG 6.4 heavy chain cDNA is accomplished by 5' and 3' rapid amplification of cDNA ends (RACE) techniques performed with human germline IgG2 heavy chain constant region specific primers (shown below) and RACE primers and other materials and methods provided in the Marathon™ cDNA amplification kit.

5' IgG2 RACE primer

5'- GGC ACG GTC ACC ACG CTG CTG AG -3' (SEQ ID NO: 9)

3' IgG2 RACE primer

5'- CCT CCA CCA AGG GCC CAT CGG TCT -3' (SEQ ID NO: 10)

[0209] The 600 bp 5' RACE product and 1200 bp 3' RACE product are cloned into pCR2.1 (Invitrogen) and are sequenced. This sequence information is used to design α OPGL-1 heavy chain specific primers for the cloning of the full-length sequence. The heavy chain 5' primer (5' α OPGL-1 IgG2 Primer) is directed against the sense strand and has a HindIII site and consensus Kozak sequence before the natural start site. The heavy chain 3' primer (3' α OPGL-1 IgG2 Primer) is an antisense primer that contains a SalI site and stop codon after the last amino acid of the heavy chain IgG2 sequence.

5' α OPGL-1 IgG2 Primer:

5'- CAGAAGCTTGACCACC ATG GAG TTT GGG CTG AGC TGG CTT TTT CTT GTG GC - 3'

(SEQ ID NO: 11)

HindIII Kozak M E F G L S W L F L V A

(SEQ ID NO: 18)

3' α OPGL-1 IgG2 Primer:

5'- GCA TGTCGAC TTA TCA TTT ACC CGG AGA CAG GGA GAG - 3' (SEQ ID NO: 12)

Sall * * K G P S L S L (SEQ ID NO: 19)

[0210] The double-stranded cDNA described above is used to generate the full-length heavy chain cDNA by PCR amplification with the 5'- and 3'- α OPGL-1 IgG2 primers. The PCR generates a 1433 bp fragment encoding the 467 amino acid residues (including the 19 amino acid IgG signal sequence) of the α OPGL-1 IgG2 heavy chain protein (Figure 2, SEQ ID NO: 2). Following purification using a QIAquick PCR Purification kit (Qiagen cat. no.28104), this fragment is used to construct the heavy chain expression vector as follows.

[0211] DNA encoding the full-length IgG2 heavy fragment generated above is cut with *HindIII* and *Sall*, purified using a QIAquick Gel Extraction kit (Qiagen cat. no.28704), and this fragment is cloned into pDSR α 19. The resulting expression plasmid is named α OPGL-1-IgG2/pDSR α 19 (Figure 6). All vector components are identical to α OPGL-1-kappa/pDSR α 19 vector, described above, except the α OPGL-1 IgG2 heavy chain cDNA replaces the α OPGL-1 kappa light chain cDNA between the *XbaI* and *Sall* sites. The α OPGL-

1 IgG2 heavy chain expression clone is sequenced to confirm that it coded for the same polypeptide that is identified in the AMG 6.4 hybridoma.

Example 2

α OPGL-1 expression in CHO cells

[0212] Stable expression of α OPGL-1 antibody is achieved by co-transfection of α OPGL-1-kappa/pDSR α 19 and α OPGL-1 -IgG2/pDSR α 19 plasmids into dihydrofolate reductase deficient (DHFR⁻) Chinese hamster ovary cells (CHO AM-1/D, U.S. Patent No. 6,210,924) followed by isolation and testing of individual clones.

[0213] A 100 mm tissue culture dish is plated with 1.5×10^6 AM-1/D cells grown in CHO d⁻ medium (DMEM-high glucose, 10% fetal bovine serum, 1% penicillin/ streptomycin/glutamine, 1X NaPyruvate, 1% nonessential amino acids (NEAA))(Gibco®) and 1% ht supplement (Gibco®)) the day before transfections (Day 0). On day one, 400 μ l of serum-free RPMI 1640 medium (Gibco®) is aliquoted into a 12x75 mm polypropylene tube. Twenty four microliters of *TransIT*®-LT1 reagent (Mirus Corporation) is added dropwise to the medium and the mixture is allowed to incubate at room temperature for 10 minutes. A total of 15 μ g of linearized plasmid DNA (7.5 μ g of α OPGL-1-kappa/pDSR α 19 and 7.5 μ g of α OPGL-1 -IgG2/pDSR α 19, digested with Pvu1) is then added dropwise to the mixture and is incubated at room temperature for 10 minutes.

[0214] The CHO d⁻ medium is removed from the cells, which are washed with 10 ml of Dulbecco's phosphate buffered saline (Gibco®). Six

milliliters of serum-free MEM medium supplemented with HT, L-glu, NEAA, and Na pyruvate (Gibco®) is added to the cells. The DNA/LT1 complex is added dropwise to the plates, which are gently rocked back and forth to distribute the DNA evenly over the cells. After 6 hours in a tissue culture incubator, the medium is replaced with fresh CHO d- medium. Forty eight hours later the cells are split to ten 100 mm culture dishes in CHO select medium (DMEM high glucose, 10% dialyzed fetal bovine serum (FBS), 1% penicillin /streptomycin /glutamine, 1% nonessential amino acids and 1X Na pyruvate) (Gibco®). Medium is changed twice weekly until colonies appeared.

[0215] After 10-14 days, colonies are picked using 5 mm cloning discs (Labcore®) soaked in 1x trypsin-EDTA (Gibco®) and are cultured in 24 well tissue culture plates with CHO select medium. When the cells become confluent, serum free media (CHO select medium minus FBS) is added and is then collected 48 hours later. These conditioned media are analyzed for antibody expression by Western blot with horse radish peroxidase (HRP)-conjugated goat anti-human IgG Fc antibody (Pierce, Rockford, IL) to detect the α OPGL-1 heavy chain, and goat anti-human kappa chain antibody (Pierce, Rockford, IL) followed by HRP-conjugated rabbit anti-goat IgG(H+L) antibody (Pierce, Rockford, IL) to detect the α OPGL-1 light chain. The highest expressing clones are expanded and stored in liquid nitrogen.

Example 3

Production of α OPGL-1

Preparation and Creation of Cell Line 125Q

[0216] CHO cells producing α OPGL-1 are cloned by two rounds of limiting dilution in 96 well plates under serum-free conditions. The clones are selected based on production and growth characteristics in various suspension vessels. EIAs are performed to select the clone that produces the highest level of α OPGL-1. Growth characteristics, including doubling times and densities are then measured by growing the clones in 100 ml, 250 ml, 500 ml, 1 L, and 3 L spinner flasks, as well as in 3L Applikon bioreactors. The clone with the fastest doubling time that reaches the highest density in culture is selected, and is designated Cell Line 125Q. When the clone has expanded to yield sufficient cells to freeze 360 ampules at approximately 1×10^7 cells/mL, cells are resuspended in a cryopreservative, serum-free medium (90% VM-Soy Batch Medium (see Table 3 for details) supplemented with 10 ml/L nonessential amino acids and 10 ml/L L-Glutamine (Gibco/LTI/Invitrogen), and 10% dimethyl sulfoxide (JT Baker)) and frozen. Ampules are stored in a limited access facility and are submerged in liquid nitrogen in liquid nitrogen dewars.

[0217] Based on growth and production in small-scale spinners and larger scale bioreactors, Cell Line 125Q is chosen as the cell line for manufacturing of α OPGL-1.

Cell Culture

[0218] α OPGL-1 is produced by expression in Cell Line 125Q, a clonal line of CHO cells that expresses α OPGL-1 from plasmids α OPGL-1-kappa/pDSR α 19 and α OPGL-1-IgG2/pDSR α 19. The cell culture process for α OPGL-1 is shown in Figure 19. For each production run, cells from a vial of Cell Line 125Q are initially grown in 50 mL of VM-Soy Batch Medium (see Table 3 for composition) supplemented with 10 ml/L nonessential amino acids and 10 ml/L L-glutamine (Gibco/LTI/Invitrogen) (VM-Soy Supp) in 125 ml erlenmeyer shakers at 100 rpm for 5 days. The entire culture is then used to inoculate VM-Soy Supp in a 500 ml spinner flask to 3×10^5 viable cells/ml ($3E5$ vc/ml), and is grown with 70 rpm spinning for 3-4 days. The entire culture from the 500 ml spinner flask is then used to inoculate VM-Soy Supp in a 3L spinner flask to $3E5$ vc/ml, and is grown with 70 rpm spinning for 3-4 days.

[0219] The culture from the 3L spinner flask is then split to two 3 L spinner flasks at $3E5$ vc/ml in VM-Soy Supp without phenol red and grown under the same conditions. These spinner flask cultures are then used to inoculate four additional spinner flasks at $3E5$ vc/ml in VM-Soy Supp without phenol red, and grown under the same conditions. Four liters of culture from the four 3L spinner flasks is used to inoculate 10 L of VM-Soy Supp without phenol red in a 20 L bioreactor, and the bioreactor is run in fed-batch mode for 7-10 days. In fed-batch mode, a nutrient feed containing concentrated media components ("Feed", as set forth below in Table 3) is added to maintain cell growth and culture viability.

[0220] The entire culture from the 20L bioreactor is then used to inoculate 70 L of VM-Soy Supp without phenol red in a 150L bioreactor, and the bioreactor is run in fed-batch mode for 9-10 days. Finally, the entire culture from the 150L bioreactor is used to inoculate approximately 880 L of VM-Soy (without supplement or phenol red) in a 2000L bioreactor, and the bioreactor is run in fed-batch mode. The rate of feed during fed-batch mode is determined such that the glucose level in the culture is maintained at 0.6 g/L for each bioreactor. The cell density and glucose concentration are measured daily and the rate of feed adjusted accordingly.

[0221] Production in the 2000L bioreactor lasts for approximately two weeks during which time α OPGL-1 is constitutively produced by the cells and secreted into the cell culture medium.

[0222] The production reactor is controlled at set pH, temperature, and dissolved oxygen level: pH is 7.0 and is controlled by carbon dioxide gas and sodium carbonate addition; dissolved oxygen is 120 mmHg and is controlled by air, nitrogen, and oxygen gas flows. Cells are maintained at 37°C throughout the process. All gases are passed through membrane filters of pore size 0.22 μ m or less.

[0223] At the end of production, the cell broth is fed into a disk stack centrifuge and the culture supernatant is separated from the cells. The centrate is further clarified through a Cuno 90SP depth filter followed by a 0.2 μ m Posidyne filter (Pall Co.). The clarified conditioned media is then concentrated by tangential flow ultrafiltration (UF) using 50kD NMWL membranes (Millipore

Biomax 50). The conditioned media is concentrated 15- to 30- fold. The resulting concentrated conditioned medium (CCM) is then either processed through purification or frozen for purification at a later date. The production process is summarized in Figure19.

Cell Culture Medium

[0224] The cell culture medium for use throughout the entire cell culture process is based on Dulbecco's Modified Eagle's Medium /Ham's Nutrient F12 (DMEM/F12, 1:1), and contains supplemental levels of amino acids, additional nutrients and salts, a soy hydrolysate and recombinant human insulin (Nucellin[®]Zn, Eli Lilly). The components are listed in Table 3. This media is referred to as VM-Soy. Media solutions are filtered through membrane filters of 0.2 µm pore size prior to use.

Table 3. Cell Culture Media Components

COMPONENTS FOR BASAL Media and FEEDS		
COMPONENT	VMsOy Batch Medium (mg/L)	FEED (mg/L)
DMEM/F12 COMPONENTS		
Inorganic salts		
CaCl ₂ (anhyd.)	116.60	233.2
CuSO ₄ .5H ₂ O	0.0026	0.0052
Fe(NO ₃) ₃ .9H ₂ O	0.1000	0.2
FeSO ₄ .7H ₂ O	0.8340	1.668
KCl	311.80	623.6
MgCl ₂ (anhyd.)	57.280	114.56
MgSO ₄ (anhyd.)	97.680	195.36
NaCl	905.990	1811.98

NaH ₂ PO ₄ .H ₂ O	125.00	250
Na ₂ HPO ₄	142.040	284.08
ZnSO ₄ .7H ₂ O	0.8640	1.728
Other Components		
D-Glucose	3151.00	12302
Na Hypoxanthine	5.40	10.8
Linoleic acid	0.090	0.18
Lipoic acid	0.2060	0.412
Phenol Red	8.10	16.2
Putrescine.2HCl	0.1620	0.324
Sodium Pyruvate	110.00	220
Amino acids		
L-Alanine	26.70	53.4
L-Arginine HCl	295.00	590
L-Asparagine.H ₂ O	45.00	90
L-Aspartic acid	39.90	79.8
L-Cysteine.HCl.H ₂ O	35.120	70.24
L-Cystine.2HCl	62.580	125.16
L-Glutamic acid	44.10	88.2
L-Glutamine	657.00	1314
Glycine	52.50	105
L-Histidine.HCl.H ₂ O	62.950	125.9
L-Isoleucine	108.940	217.88
L-Leucine	118.10	236.2
L-Lysine HCl	182.50	365
L-Methionine	34.480	68.96
L-Phenylalanine	70.960	141.92
L-Proline	57.50	115
L-Serine	73.50	147
L-Threonine	106.90	213.8
L-Tryptophan	18.040	36.08
L-Tyrosine.2Na.2H ₂ O	111.580	223.16
L-Valine	105.70	211.4
Vitamins		
Biotin	0.0073	0.0146

D-Ca Pantothenate	4.480	8.96
Choline Chloride	17.960	35.92
Folic Acid	5.30	10.6
i-Inositol	25.20	50.4
Niacinamide	4.040	8.08
Pyridoxal HCl	4.00	8
Pyridoxine HCl	0.0620	0.124
Riboflavin	0.4380	0.876
Thiamine HCl	4.340	8.68
Thymidine	0.3635	0.727
Vitamin B12	1.360	2.72
ADDITIONAL COMPONENTS		
Nucellin Zn, (rhu insulin)	5.00	15
Selenous Acid	0.0050	0.015
Ethanolamine	0.0012	0.0037
Triiodothyronine	0.000040	0.00012
Hydrocortisone	0.020	0.06
Ferric Citrate	122.450	122.450
Pluronic F-68	1000.00	500
Soy Hydrolysate	6000.00	6000.00
NaHCO ₃	3000.00	3000.00
NaCl	3500.00	

Purification Process

[0225] α OPGL-1 expressed in CHO cells is secreted into the extracellular medium. A series of steps may be used to generate pure material. The process uses hydrophobic charge induction, cation exchange, and hydrophobic interaction chromatography along with a low pH step and viral filter. These procedures are described below.

A. Hydrophobic Charge Induction Chromatography (HCIC)

[0226] This chromatography step removes the majority of host cell proteins and DNA. The concentrated conditioned media (CCM) is filtered through a Cuno 30SP filter and then through a Cuno VR07 charged cellulose-based filter, and then loaded on to an MEP HyperCel resin. After loading, the column is washed with equilibration buffer (20 mM Tris pH 7.2). The antibody is eluted from the resin using a low pH buffer (20 mM Sodium Acetate, pH 5.0). As it is eluted from the column, the product is collected based on the absorbance at 280 nm of the column effluent.

B. Viral Inactivation

[0227] The MEP pool is titrated to pH 3.7 and is held for approximately 60 minutes to inactivate potentially contaminating retrovirus. Following the hold step, the pH is adjusted to approximately 6.0.

C. Viral Filtration

[0228] The pH-adjusted pool is filtered through a Millipore Viresolve NFR filter or equivalent. The antibody flows through the filter while potentially contaminating viruses ≥ 50 nm are retained.

D. Cation Exchange Chromatography (CEX)

[0229] The antibody is further purified by cation exchange chromatography using SP Sepharose HP (Amersham Pharmacia) or equivalent. The cation exchange chromatography step removes additional CHO cell proteins, DNA, lower molecular weight proteins, and aggregated forms of α OPGL-1. The viral filtered pool is loaded onto the cation exchange resin. After

loading, the column is washed with equilibration buffer (20 mM NaMES pH 6.2). The antibody is then eluted with a linear gradient of increasing salt (20 mM NaMES pH 6.2, 0 M NaCl to 20 mM NaMES pH 6.2, 0.3 M NaCl). As it is eluted from the column, the product is collected based on the absorbance at 280 nm of the column effluent.

E. Hydrophobic Interaction Chromatography (HIC)

[0230] The antibody is further purified by hydrophobic interaction chromatography using Phenyl Toyopearl 650S (Tosoh Biosep) or equivalent. The hydrophobic interaction chromatography step is used as a polishing step and removes additional CHO cell proteins, DNA, lower molecular weight proteins, and aggregated forms of α OPGL-1. The cation exchange pool is conditioned before loading onto the column by addition of ammonium sulfate to a conductivity of >105 mS/cm at 15-25°C. After loading, the column is washed with the equilibration buffer (1M Potassium Phosphate pH 8). The antibody is then eluted with a linear gradient of decreasing salt concentration (1M Potassium Phosphate, 0mM Tris pH 8 to 0M Potassium Phosphate, 20 mM Tris pH 8). As it is eluted from the column, the product is collected based on the absorbance at 280 nm of the column effluent.

F. Concentration and Diafiltration

[0231] The HIC column pool is concentrated and diafiltered into formulation buffer by tangential flow ultrafiltration using 50kD NMWL membranes (Millipore Biomax 50). The formulation buffer includes 10 mM Acetate, 5% Sorbitol, pH 5.2 and α OPGL-1 is at 30 mg/mL.

Final Filtration and Storage

[0232] The purified bulk is passed through a 0.22 μ m PVDF filter (Millipore), is sampled, and stored at approximately -30°C in a secured freezer.

Example 4

Binding Specificity of α OPGL-1

[0233] Antibodies that are produced in CHO cells that are transfected with the two expression vectors as discussed in Examples 1 and 2 may be used in the following examples 4, 5, and 6.

[0234] Human OPG binds and neutralizes OPGL in rats, mice and cynomolgus monkeys, as well as in humans. α OPGL-1 binds human OPGL with high affinity but does not bind significantly to murine OPGL (Table 4).

Table 4: Affinity of α OPGL-1 to Cell Membrane Expressed OPGL of Human, Cynomolgus Monkey, or Mouse Sequence.

OPGL Species	α OPGL-1 ED ₅₀ (ng/ml)
Human	16
Cynomolgus	19
Mouse	No Specific Binding

OPGL of these species is expressed in CHO cells as the full-length, membrane-bound protein. Binding of α OPGL-1 to the cell surface expressed OPGL is assessed by FACS analysis of cells incubated with α OPGL-1 and a FITC-labeled secondary antibody to human IgG2. α OPGL-1 binds human and cynomolgus OPGL but there is no specific binding to mouse OPGL.

[0235] In addition, human OPG has been reported to show weak binding to tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) (Truneh et al, 2000), a related member of the TNF family, which shows DNA and

amino acid sequence homology to OPGL (Lacey et al., 1998). However, OPG does not detectably bind to other TNF-related proteins such as TNF α , TNF β , or CD40 ligand.

[0236] α OPGL-1 binds specifically to OPGL on EIA plates (Figure 7). Recombinant soluble OPGL (2 μ g/ml) is coated onto 96-well EIA plates at room temperature for 16 to 24 hours. After blocking with 1% BSA in PBS, varying concentrations of α OPGL-1 (approximately 2 ng/ml to 1000 ng/ml) diluted in 1% BSA/PBS are added to the wells and the plates are incubated for about 2 hours at room temperature. Bound antibody is detected with goat anti-Human IgG (Fab')-HRP using TMB-H₂O₂ (tetramethylbenzidine and hydrogen peroxide) substrate cocktail. The absorbance is read at 450nm and 650nm.

[0237] α OPGL-1 binds specifically OPGL expressed on the surface of transfected cells (Figure 8). α OPGL-1 (100 ng/ml) diluted in FACS Buffer (PBS, 0.1%BSA, 0.01%Sodium Azide) is preincubated with varying concentrations of OPGL, TNF α , TNF β , TRAIL, or CD40 ligand (approximately 0.1 ng/ml to 1000 ng/ml), and is then added to approximately 200,000 CHO REN 218-9 cells, which are CHO cells stably expressing membrane-bound OPGL on the cell surface. After 1 hour at 2-8°C, unbound antibody is removed by centrifugation and washing. Cells are then incubated for 30 minutes at 2-8°C with FITC-labeled F(ab')₂ Goat anti-Human IgG (Fc γ fragment specific). After centrifugation and washing, cell surface fluorescence is measured using flow cytometry. Figure 8 shows that binding of α OPGL-1 to CHO REN 218-9 cells is

specific, and is competitively reduced by addition of soluble OPGL, but not by addition of TNF α , TNF β , TRAIL, or CD40 ligand.

[0238] In competition experiments, α OPGL-1 binding to OPGL on EIA plates is inhibited by addition of exogenous OPGL (Figure 9), but not by the addition of TNF α , TNF β , TRAIL, or CD40 ligand (Figure 10). This procedure is performed in substantially the same manner as above, for binding of α OPGL-1 to OPGL on EIA plates, except a constant concentration of α OPGL-1 (100ng/mL) is preincubated with varying concentrations of soluble OPGL or other ligands (approximately 1 ng/ml to 1000 ng/ml for each) before it is added to the OPGL-coated plates.

Example 5

α OPGL-1 Neutralizing Activity

Inhibition of Osteoclast Formation

[0239] RAW 264.7 (ATCC No. TIB-71, Manassas, VA) is a murine macrophage cell line that was derived from an Abelson murine leukemia virus-induced tumor. RAW 264.7 cells will differentiate to osteoclast-like cells in the presence of OPGL. The basic assay for generation of osteoclasts in culture from RAW cells in the presence of OPGL has been described in detail in Simonet et al (1997) *Cell* **89** p. 309, and Lacey et al (1998) *Cell* **93** p. 165, which are herein incorporated by reference for any purpose.

[0240] RAW cells are stimulated by ligand to differentiate into osteoclast-like cells, and the differentiation can be measured by TRAP activity, a

property of osteoclasts. Thus, the effect of α OPGL-1 on osteoclastogenesis can be measured.

[0241] RAW cells are incubated for 4 days in the presence of a constant amount of OPGL (40 ng/ml) and varying amounts of α OPGL-1 (6.3 ng/ml to 200 ng/ml) in cell culture medium (DMEM, 10% FBS, 0.292 mg/ml L-Glut, 100 units/ml Penicillin G, 100 μ g/ml Streptomycin sulfate). At the end of 4 days, the cells are stained for tartrate-resistant acid phosphatase (TRAP) activity by permeabilization and acidification, followed by treatment with para-nitrophenylphosphate for 5 minutes. Briefly, the media is aspirated off of the cells, and 100 μ l of citrate buffer (410 ml 0.1M citric acid, 590 ml 0.1 M citrate, trisodium salt, 1 mL triton X-100) is added to each well and the plates are incubated for 3 to 5 minutes at room temperature. One hundred microliters of PNPP solution is then added (157.8 mg acid phosphatase reagent (Sigma 104-100), 7.2 ml tartrate solution (Sigma cat. no. 387-3), and 22.8 ml citrate buffer), and plates are incubated for 3 to 5 minutes at room temperature. The reaction is terminated by addition of 50 μ l 0.5 M NaOH solution.

[0242] TRAP will convert para-nitrophenylphosphate to para-nitrophenol, which can be quantitated by optical density measurement at 405 nm. The TRAP activity, which is a surrogate marker for osteoclast development, therefore correlates with the optical density at 405 nm. A plot of optical density versus α OPGL-1 concentration is shown in Figure 11, and demonstrates that α OPGL-1 inhibits osteoclast formation in this assay.

Inhibition of OPGL Binding to its Receptor

[0243] The potency of α OPGL-1 is demonstrated by its ability to block the binding of OPG ligand to its cognate receptor, the osteoclast differentiation and activating receptor (ODAR, also known as RANK). This assay uses homogeneous time resolved fluorescent resonance (HTRF) to detect binding of α OPGL-1 to Europium-conjugated osteoprotegerin ligand (Eu-OPGL). If α OPGL-1 inhibits Eu-OPGL binding to ODAR, fluorescent output will decrease, and the amount of α OPGL-1 present will be inversely related to the amount of fluorescence.

[0244] OPGL is labeled with europium, which emits light at 620 nm when excited with 337 nm light. ODAR is fused to FLAG and to Fc, and the Fc-ODAR-FLAG fusion protein is labeled with an anti-FLAG antibody linked to allophycocyanin (APC), a fluorophore which emits 665 nm light when excited by light at 620 nm. Therefore, when Eu-labeled OPG ligand binds to the Fc-ODAR-FLAG/anti-FLAG-APC complex, the tertiary complex will emit 665 nm light when excited with light at 337 nm.

[0245] Eu-OPGL at 0.05 μ g/ml is preincubated with various concentrations (0.1 to 150 ng/ml) of α OPGL-1 in assay buffer (50 mM Tris pH 8, 100 mM NaCl, 0.05% NaN₃, 0.1% BSA, and 0.05% Tween 20) at room temperature for approximately one hour (Preincubation mix). A mixture of Fc-ODAR-FLAG (1 μ g/ml) and anti-FLAG-APC (2.5 μ g/ml) is also prepared in assay buffer and incubated at room temperature for one hour (Fluorochrome mix). Equal volumes of Preincubation mix and Fluorochrome mix are then combined

and incubated at room temperature for 3 hours. The fluorescence is measured by reading plates on the Packard Discovery HTRF microplate analyzer using an excitation wavelength of 337 nm and an emission wavelength of 665 nm.

[0246] When α OPGL-1 is preincubated with Eu-OPG ligand and is then mixed with Fc-ODAR-FLAG/anti-FLAG-APC, the fluorescence intensity at 665 nm decreases in a dose-dependent manner, as shown in Figure 12, demonstrating that α OPGL162 can effectively inhibit OPGL binding to ODAR.

Example 6

Pharmacokinetics in Cynomolgus Monkeys

[0247] Six male and six female cynomolgus monkeys, not greater than 4.5 years of age and weighing 2 to 4 kg are assigned to 4 dose groups. Group 1 consists of 3 males and 3 females. Groups 2, 3, and 4 each consists of 1 male and 1 female. Animals in Group 1 are administered a single SC dose of 1 mg/kg α OPGL-1, while animals in Groups 2, 3 and 4 are administered single IV doses of 0.1, 1.0, or 10.0 mg/kg of α OPGL-1, respectively.

[0248] Animals are dosed with α OPGL-1 expressed from transfected chinese hamster ovary (CHO) cells. Serum samples are taken for determination of α OPGL-1 levels, antibody analysis, and analysis of the bone turnover marker serum N-telopeptide (serum N-Tx), alkaline phosphatase (ALP), and serum calcium (serum Ca). Urine is also collected for analysis of N-telopeptide (urine N-Tx) and creatinine.

[0249] The serum concentration-time profiles following IV administration are characterized by a tri-phasic distribution (Figure 13). Initially,

there is a rapid distribution phase, followed by a significantly slower plateau phase, which appears to be concentration-dependent. The third observed phase is a rapid elimination phase.

[0250] Non-compartmental analysis of complete serum concentration-time profiles using WinNonlin Professional (v1.5), and exponential analysis of the data up to 14 days after test article administration and above 10,000 ng/mL using SAAM II (v1.1.2) are utilized to investigate the pharmacokinetics of α OPGL-1 in monkeys. The initial volume of distribution from all IV doses averages 28.9 mL/kg, similar to plasma volume. Steady state volume (V_{ss}) of distribution averages 39 mL/kg across all IV doses. Exponential analysis indicates that α OPGL-1 has an average distribution half-life ($t_{1/2\alpha}$) of 6.02 hours, an extended secondary phase with a half-life ($t_{1/2\beta}$) that increases with dose from 86.9 hours at a dose of 0.1 mg/kg to a maximum of 444 hours at a dose of 10.0 mg/kg. Terminal elimination half-life ($t_{1/2z}$) estimated non-compartmentally averages 31 hours across all IV dose groups. Clearance (CL, CL/F) of α OPGL-1 is found to be non-linear, with animals receiving IV doses of 10 mg/kg having an average clearance (0.120 mL/hr/kg) that is 3.3-fold lower than those receiving 0.1 mg/kg (0.401 mL/hr/kg).

[0251] After administering subcutaneously, absorption is slow, with average peak concentrations (C_{max}) of 11,600 ng/mL at 132 hr. There is high variability in the range of exposure after SC administration, resulting in an average clearance of 0.387 ± 0.281 mL/hr/kg and mean residence time of 202 ± 80.1 hours. Average bioavailability is 89%.

[0252] The preceding data are summarized in Table 5.

Table 5: Mean ($\pm$ SD) Non-Compartmental Pharmacokinetic Parameters^a in Cynomolgus Monkeys After Administering a Single Dose of α OPGL-1 IV and SC

Non-Compartmental Parameter Estimates						
Parameter	Unit	1.0 mg/kg		0.1 mg/kg	1.0 mg/kg	10 mg/kg
		SC (n=6)		IV (n=2)	IV (n=2)	IV (n=2)
		Mean	SD	Mean	Mean	Mean
T _{max}	hr	132	60.2	0	0	0
C _{max}	ng/ml	11600	3410	4330	38200	326000
t _{1/2z}	hr	34.9	11.1	30.7	31.4	ND ^b
AUC _(0-∞)	μg*hr/ml	3520	1750	253	3950	99900
CL, CL/F	ml /hr/kg	0.387	0.281	0.401	0.256	0.120
MRT	hr	202	80.1	84.8	124	519
V _{ss}	ml/kg	N/A ^c	N/A	33.7	31.7	55.9

^aValues are reported to 3 significant figures

^bNot Determined, PK samples end during plateau (β) phase hence terminal phase is not observed

^cNot Applicable

[0253] α OPGL-1 causes a rapid decrease in serum N-Tx levels

within 24 hours post dose (Figure 14). The average time of maximum effect is observed to occur between 12 hours and 7 days post dose as IV doses increase from 0.1 to 10 mg/kg, and between 12 hours and 11 days in animals receiving a SC dose of 1.0 mg/kg. Maximum effect increases with dose from approximately 80 to 91% over the dose range of 0.1 to 1 mg/kg. However, at higher doses no further suppression is observed with maximum inhibition of 91%. Mean levels of serum N-Tx return to baseline by day 28 after administering 0.1 mg/kg IV and by day 70 after administering 1 mg/kg SC. Urine N-Tx shows similar trends to those of serum N-Tx, except that all groups return to baseline values by study day 105 (Figure 15).

[0254] Suppression of serum Ca increases with dose to a mean nadir of 31.6% below the baseline average seven days after IV administration of 10.0 mg/kg. All other dose groups have mean decreases in serum Ca of less than 26.4% from their baseline averages. By day 17 all serum Ca levels in treated animals return to within 10% of their baseline averages (Figure 20).

[0255] As bone resorption and formation are intimately linked, changes in bone formation markers (ALP) are also observed with a much slower decline in ALP levels and a more prolonged suppression than the formation marker, N-Tx (Figure 21). Observing bone resorption markers decrease prior to bone formation markers (ALP) following dosing with α OPGL-1 confirms that the α OPGL-1 is a bone anti-resorptive agent.

[0256] The majority of animals (9 of 12) develop antibodies to α OPGL-1. The incidence of antibodies to α OPGL-1 is not dose or route dependent. It is not possible to assess the effect of antibodies to α OPGL-1 on α OPGL-1 pharmacokinetics above 0.1 mg/kg when no dose group has both antibody negative and positive animals. At 0.1 mg/kg IV, the majority of α OPGL-1 is cleared prior to antibody development and therefore, effects on α OPGL-1 disposition are not observed (Figure 16).

We Claim:

1. An antibody, comprising a heavy chain and a light chain, wherein the heavy chain comprises an amino acid sequence as set forth in SEQ ID NO: 2 or a fragment thereof, and the light chain comprises an amino acid sequence as set forth in SEQ ID NO: 4 or a fragment thereof.

2. An antibody, comprising a heavy chain and a light chain, wherein the heavy chain comprises a variable region comprising an amino acid sequence as set forth in SEQ ID NO: 13 or a fragment thereof, and wherein the light chain comprises a variable region comprising an amino acid sequence as set forth in SEQ ID NO: 14 or a fragment thereof.

3. The antibody of claim 2, wherein the heavy chain and the light chain are connected by a flexible linker to form a single-chain antibody.

4. The antibody of claim 3, which is a single-chain Fv antibody.

5. The antibody of claim 2, which is a Fab antibody.

6. The antibody of claim 2, which is a Fab' antibody.

7. The antibody of claim 2, which is a (Fab')₂ antibody.

8. The antibody of claim 2, which is fully human.

9. The antibody of claim 2, wherein the antibody inhibits binding of OPGL to an osteoclast differentiation and activation receptor (ODAR).

10. A heavy chain comprising an amino acid sequence as set forth in SEQ ID NO: 2 or a fragment thereof.

11. A heavy chain comprising an amino acid sequence as set forth in SEQ ID NO: 13 or a fragment thereof.

12. A light chain comprising an amino acid sequence as set forth in SEQ ID NO: 4 or a fragment thereof.

13. A light chain comprising an amino acid sequence as set forth in SEQ ID NO: 14 or a fragment thereof.

14. An antibody, comprising a heavy chain and a light chain, wherein the heavy chain comprises an amino acid sequence as set forth in SEQ ID NO: 2 or a fragment thereof.

15. An antibody, comprising a heavy chain and a light chain, wherein the heavy chain comprises a variable region comprising an amino acid sequence as set forth in SEQ ID NO: 13 or a fragment thereof.

16. An antibody, comprising a heavy chain and a light chain, wherein the light chain comprises an amino acid sequence as set forth in SEQ ID NO: 4 or a fragment thereof.

17. An antibody, comprising a heavy chain and a light chain, wherein the light chain comprises a variable region comprising an amino acid sequence as set forth in SEQ ID NO: 14 or a fragment thereof.

18. An antibody, comprising a heavy chain and a light chain,
(a) wherein the heavy chain comprises a first variable region, and wherein the first variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 13, and
(b) wherein the light chain comprises a second variable region, and wherein the second variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 14, and

(c) wherein the antibody interacts with an osteoprotegerin ligand (OPGL).

19. The antibody of claim 18, wherein the first variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 13, and wherein the second variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 14.

20. The antibody of claim 18, wherein the first variable region comprises a sequence that has at least 99% identity to the amino acid sequence set forth in SEQ ID NO: 13, and wherein the second variable region comprises a sequence that has at least 99% identity to the amino acid sequence set forth in SEQ ID NO: 14.

21. A pharmaceutical composition comprising an antibody of any of claims 1 to 9 or 14 to 20.

22. The pharmaceutical composition of claim 21, further comprising at least one therapeutic agent selected from a bone morphogenic factor, transforming growth factor- β (TGF- β), an interleukin-1 (IL-1) inhibitor, IL-1ra, KineretTM, a TNF α inhibitor, a soluble TNF α receptor, EnbrelTM, an anti-TNF α antibody, RemicadeTM, a D2E7 antibody, a parathyroid hormone, an analog of a parathyroid hormone, a parathyroid hormone related protein, an analog of a parathyroid hormone related protein, a prostaglandin, a bisphosphonate, an alendronate, fluoride, calcium, a non-steroidal anti-inflammatory drug (NSAID), a COX-2 inhibitor, CelebrexTM, VioxxTM; an immunosuppressant, methotrexate,

leflunomide, a serine protease inhibitor, a secretory leukocyte protease inhibitor (SLPI), an IL-6 inhibitor, an antibody to IL-6, an IL-8 inhibitor, an antibody to IL-8, an IL-18 inhibitor, an IL-18 binding protein, an IL-18 antibody, an Interleukin-1 converting enzyme (ICE) modulator, a fibroblast growth factor (FGF), an FGF modulator, a PAF antagonist, a keratinocyte growth factor (KGF), a KGF-related molecule, a KGF modulator; a matrix metalloproteinase (MMP) modulator, a nitric oxide synthase (NOS) modulator, a modulator of glucocorticoid receptor, a modulator of glutamate receptor, a modulator of lipopolysaccharide (LPS) levels, a noradrenaline, a noradrenaline mimetic, and a noradrenaline modulator.

23. The pharmaceutical composition of claim 22, wherein the at least one therapeutic agent is selected from Kineret™, Enbrel™, Remicade™, D7E7 antibody, and methotrexate.

24. A method of detecting the level of OPGL in a biological sample comprising contacting the sample with the antibody of claim 2.

25. A method of treating an osteopenic disorder in a patient, comprising administering the pharmaceutical composition of claim 21.

26. A method of treating an osteopenic disorder in a patient, comprising administering the pharmaceutical composition of claim 22.

27. A method of treating an inflammatory condition with attendant bone loss in a patient, comprising administering the pharmaceutical composition of claim 22.

28. A method of treating an autoimmune condition with attendant bone loss in a patient, comprising administering the pharmaceutical composition of claim 22.

29. A method of treating rheumatoid arthritis in a patient, comprising administering the pharmaceutical composition of claim 22.

ANTIBODIES TO OPGL SEQ LISTING.txt
SEQUENCE LISTING

<110> Boyle, William J
Martin, Francis H
Corvalan, Jose R
Davis, C. Geoffrey

<120> Antibodies to OPGL

<130> 06843.0049-00000

<150> 60/301,172

<151> 2001-06-26

<160> 20

<170> PatentIn version 3.1

<210> 1

<211> 1426

<212> DNA

<213> Mus musculus

<400> 1
aagcttgacc accatggagt ttgggctgag ctggccttttt. cttgtggcta ttttaaaagg 60
tgtccagtgt gaggtgcagc tggtggagtc tgggggaggc ttggtacagc ctgggggggtc 120
cctgagaactc tcctgtgcag cctctggatt caccttttagc agctatgcca tgagctgggt 180
ccgccagggt ccaggggaagg ggctggagtg ggtctcaggt attactggga gtggtggtag 240
tacatactac gcagactccg tgaagggccg gttcaccatc tccagagaca attccaagaa 300
cacgctgtat ctgcaaataa acagcctgag agccgaggac acggccgtat attactgtgc 360

ANTIBODIES TO OPGL SEQ LISTING.txt

gaaagatcca gggactacgg tgattatgag ttggttcgac cctggggcc agggaaacct 420
 ggtcaccgtc tcctcagcct ccaccaaggg cccatcggtc ttccccctgg cgccttgcgc 480
 caggagcacc tccgagagca cagcggccct gggctgcctg gtcaaggact acttccccga 540
 accggtgacg gtgtcgtgga actcaggcgc tctgaccagc ggctgcaca ccttcccagc 600
 tgtcctacag tcctcaggac tctactccct cagcagcgtg gtgaccgtgc cctccagcaa 660
 cttcggcacc cagacctaca cctgcaacgt agatcacaag cccagcaaca ccaagggtga 720
 caagacagtt gagcgcaaat gttgtgtcga gtgccaccg tgcccagcac cacctgtggc 780
 aggaccgtca gtcttcctct tcccccaaaa acccaaggac accctcatga tctcccggac 840
 ccttgaggtc acgtgcgtgg tgggtggacgt gagccacgaa gaccccaggg tccagttcaa 900
 ctggtacgtg gacggcgtgg aggtgcataa tgccaagaca aagccacggg aggagcagtt 960
 caacagcacg ttccgtgtgg tcagcgtcct caccgttgtg caccaggact ggctgaacgg 1020
 caaggagtac aagtgcagg tctccaacaa aggcctccca gcccocatg agaaaacat 1080
 ctccaaaacc aaagggcagc cccgagaacc acagggtgtac accctgcccc catcccggga 1140
 ggagatgacc aagaaccagg tcagcctgac ctgcctggtc aaaggcttct accccagcga 1200
 catcgccgtg gagtgggaga gcaatgggca gccggagaac aactacaaga ccacacctcc 1260
 catgctggac tccgacggct ccttcttccct ctacagcaag ctcaccgtgg acaagagcag 1320
 gtggcagcag gggaaacgtct tctcatgctc cgtgatgcat gaggtctctgc acaaccacta 1380
 cagcgagaag agcctctccc tgtctccggg taaatgataa gtcgac 1426

<210> 2

<211> 467

<212> PRT

<213> Mus musculus

<400> 2

Met Glu Phe Gly Leu Ser Trp Leu Phe Leu Val Ala Ile Leu Lys Gly
 1 5 10 15

Val Gln Cys Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln
 20 25 30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45

ANTIBODIES TO OPGL SEQ LISTING.txt

Ser	Ser	Tyr	Ala	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	50	55	60	
Glu	Trp	Val	Ser	Gly	Ile	Thr	Gly	Ser	Gly	Gly	Ser	Thr	Tyr	Tyr	Ala	65	70	75	80
Asp	Ser	Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	85	90	95	
Thr	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	100	105	110	
Tyr	Tyr	Cys	Ala	Lys	Asp	Pro	Gly	Thr	Thr	Val	Ile	Met	Ser	Trp	Phe	115	120	125	
Asp	Pro	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	130	135	140	
Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Cys	Ser	Arg	Ser	Thr	Ser	145	150	155	160
Glu	Ser	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	165	170	175	
Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	180	185	190	
Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	195	200	205	
Val	Val	Thr	Val	Pro	Ser	Ser	Asn	Phe	Gly	Thr	Gln	Thr	Tyr	Thr	Cys	210	215	220	
Asn	Val	Asp	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Thr	Val	Glu	225	230	235	240
Arg	Lys	Cys	Cys	Val	Glu	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Pro	Val	Ala	245	250	255	
Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	260	265	270	
Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His				

ANTIBODIES TO OPGL SEQ LISTING.txt
280 285

275

Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val
290 295 300

His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe
305 310 315 320

Arg Val Val Ser Val Leu Thr Val Val His Gln Asp Trp Leu Asn Gly
325 330 335

Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ala Pro Ile
340 345 350

Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu Pro Gln Val
355 360 365

Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser
370 375 380

Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
385 390 395 400

Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
405 410 415

Met Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val
420 425 430

Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met
435 440 445

His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
450 455 460

Pro Gly Lys
465

<210> 3

<211> 728

<212> DNA

<213> Mus musculus

ANTIBODIES TO OPGL SEQ LISTING.txt

<400> 3
tctagaccac catggaaaacc ccagcgcagc ttctcttcct cctgctactc tggctcccag 60
ataccaccgg agaaattgtg ttgacgcagt ctccaggcac cctgtotttg tctccagggg 120
aaagagccac cctctcctgt agggccagtc agagtgttcg cggcaggtag ttagcctggt 180
accagcagaa acctggccag gctcccaggc tcctcatcta tggcgcaccc agcagggcca 240
ctggcatccc agacagggtc agtggcagtg ggtctgggac agacttcact ctcaccatca 300
gcagactgga gcctgaagat ttgacagtgt ttactgtca gcagtatggt agttcacctc 360
ggacgttcgg ccaagggacc aaggtggaaa tcaaacgaac tgtggctgca ccactgtctt 420
tcctcttccc gccatctgat gacgagttga aatctggaac tgctctgtt gtgtgcctgc 480
tgaataactt ctatcccaga gaggccaaag tacagtggaa ggtggataac gccctccaat 540
cgggtaactc ccaggagagt gtcacagagc aggacagcaa ggacagcacc tacagcctca 600
gcagcaccct gacgctgagc aaagcagact acgagaaaca caaagtctac gcctgcgaag 660
tcacccatca gggcctgagc tcgcccgtca caaagagctt caacagggga gagtgttgat 720
aagtcgac 728

<210> 4

<211> 235

<212> PRT

<213> Mus musculus

<400> 4

Met Glu Thr Pro Ala Gln Leu Leu Phe Leu Leu Leu Leu Trp Leu Pro
1 5 10 15

Asp Thr Thr Gly Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser
20 25 30

Leu Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser
35 40 45

Val Arg Gly Arg Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala
50 55 60

Pro Arg Leu Leu Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro

65 70 75 80

Page 6

ANTIBODIES TO OPGL SEQ LISTING.txt

<210> 6

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> 3' kappa RACE primer

<400> 6

aagggtcaga ggccaaagga tgg

23

<210> 7

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> 5' anti-OPGL-1 kappa primer

<400> 7

caactctaga ccaccatgga aaccccagcg

30

<210> 8

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> 3' anti-OPGL-1 kappa primer

<400> 8

tttgacgtcg acttatcaac actctcccct gttgaag

37

<210> 9

<211> 23

ANTIBODIES TO OPGL SEQ LISTING.txt

<212> DNA

<213> Artificial Sequence

<220>

<223> 5' IgG2 RACE primer

<400> 9

ggcacggtca ccacgctgct gag

23

<210> 10

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> 3' IgG2 RACE primer

<400> 10

cctccaccaa gggcccatcg gtct

24

<210> 11

<211> 51

<212> DNA

<213> Artificial Sequence

<220>

<223> 5' anti-OPGL-1 IgG2 Primer

<400> 11

cagaagcttg accaccatgg agtttgggct gagctggctt tttcttgtgg c

51

<210> 12

<211> 37

<212> DNA

<213> Artificial Sequence

ANTIBODIES TO OPGL SEQ LISTING.txt

<220>

<223> 3' anti-OPGL-1 IgG2 Primer

<400> 12

gcatgtcgac ttatcattta cccggagaca gggagag

37

<210> 13

<211> 122

<212> PRT

<213> Mus musculus

<400> 13

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Gly Ile Thr Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Asp Pro Gly Thr Thr Val Ile Met Ser Trp Phe Asp Pro Trp
100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 14

<211> 108

<212> PRT

ANTIBODIES TO OPGL SEQ LISTING.txt

<213> Mus musculus

<400> 14

Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly
1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Arg Gly Arg
20 25 30

Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu
35 40 45

Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser
50 55 60

Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu
65 70 75 80

Pro Glu Asp Phe Ala Val Phe Tyr Cys Gln Gln Tyr Gly Ser Ser Pro
85 90 95

Arg Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
100 105

<210> 15

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> random primer

<220>

<221> misc_feature

<222> (18)..(23)

<223> N is A, C, G, or T

ANTIBODIES TO OPGL SEQ LISTING.txt

<400> 15
ggccggatag gcctcacnnn nnnt

24

<210> 16

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> translation of portion of 5' anti-OPGL-1 kappa primer

<400> 16

Met Glu Thr Pro Ala
1 5

<210> 17

<211> 6

<212> PRT

<213> Artificial Sequence

<220>

<223> translation of portion of 3' anti-OPGL-1 kappa primer

<400> 17

Cys Glu Gly Arg Asn Phe
1 5

<210> 18

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> translation of portion of 5' anti-OPGL-1 IgG2 Primer

<400> 18

ANTIBODIES TO OPGL SEQ LISTING.txt

Met Glu Phe Gly Leu Ser Trp Leu Phe Leu Val Ala
 1 5 10

<210> 19

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> translation of portion of 3' anti-OPGL-1 IgG2 Primer

<400> 19

Lys Gly Pro Ser Leu Ser Leu
 1 5

<210> 20

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> LHRH antagonist peptide

<220>

<221> MISC_FEATURE

<222> (1)..(1)

<223> Xaa is Ac-D-Nal, where Nal is 3-(2-naphthyl)alaninyl

<220>

<221> MISC_FEATURE

<222> (2)..(2)

<223> Xaa is (4'-chlorophenyl)alaninyl

ANTIBODIES TO OPGL SEQ LISTING.txt

<220>

<221> MISC_FEATURE

<222> (3)..(3)

<223> Xaa is D-Pal, where Pal is 3-(3'-pyridyl)alaninyl

<220>

<221> MISC_FEATURE

<222> (5)..(5)

<223> Xaa is N-methyl tyrosine

<220>

<221> MISC_FEATURE

<222> (6)..(6)

<223> Xaa is D-asparagine

<220>

<221> MISC_FEATURE

<222> (8)..(8)

<223> Xaa is N-epsilon-2-propyl-lysiny1

<220>

<221> MISC_FEATURE

<222> (10)..(10)

<223> Xaa is D-alanine-NH2

<400> 20

Xaa Xaa Xaa Ser Xaa Xaa Leu Xaa Pro Xaa
1 5 10