

NOVEL INHIBITORS OF PROLIFERATION AND ACTIVATION OF
SIGNAL TRANSDUCER AND ACTIVATOR OF TRANSCRIPTION (STATS)

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Application Serial No. 60/079,002 filed on July 8, 2008. The application is incorporated by reference herein in its entirety.

STATEMENT REGARDING FEDERALLY
SPONSORED RESEARCH OR DEVELOPMENT

[0002] None.

THE NAMES OF THE PARTIES TO A JOINT RESEARCH AGREEMENT

[0003] None.

REFERENCE TO SEQUENCE LISTING

[0004] None.

FIELD OF THE INVENTION

[0005] Pyridine compounds and compositions and their application as pharmaceuticals for the treatment of disease are provided herein. Methods of inhibiting of STAT3 and/or STAT5 activation in a human or animal subject are also provided for the treatment of diseases such as cancer.

BACKGROUND OF THE INVENTION


[0006] The Jak/STAT signaling pathway is known to be activated in a number of cancers and inflammatory diseases. When activated through phosphorylation of a single tyrosine residue STAT3 dimerizes, translocates to nucleus, and serves as a transcription factor driving the upregulation of a number of proliferative and pro-survival genes including survivin, VEGF, Bcl-2, and Bcl-xL. These gene products are known to promote the growth and metastasis of over 20 tumor types and are also associated with the promotion of several proliferative skin disorders including psoriasis, T-cell lymphoma, and atopic dermatitis.

[0007] Inhibition of STAT3 phosphorylation in tumor bearing animals leads to the induction of apoptotic cell death and in preclinical orthotopic models of several diseases a significant reduction in tumor size. Likewise in inflammatory skin diseases inhibition of STAT3 phosphorylation leads to resolution of plaques, scales, and other associated lesions.

[0100] While the development of inhibitors of this pathway is currently desirable, a variety of inhibition (such as small molecule, antisense, iRNA) approaches have been tried with unsatisfactory results.

SUMMARY OF THE INVENTION

[0008] Novel pyridine compounds and pharmaceutical compositions that inhibit cancer, have been found, together with methods of synthesizing and using the compounds including methods for the treatment of STAT3-mediated diseases and conditions in a patient by administering the compounds. The compounds can also be useful in treating STAT5-mediated diseases and conditions. This class of compounds is defined by structural Formula I as follows:





[0009] or a pharmaceutically acceptable salt or solvate thereof, wherein:

[0010] n is an integer selected from 1, 2 or 3;

[0011] R_0 is R_1 , or R_0 is R_1-Z_1 — wherein Z_1 is alkyl, and specifically may be a lower alkyl such as $-(CH_2)_{m_3}$ — where $m_3=0, 1, 2, 3$, or 4;

[0012] R_1 is:







[0013] wherein X_1 , X_2 , X_3 , and X_4 , are each independently hydrogen, halogen, alkyl, alkoxy, OH, trihalomethyl, or NO_2 , and Y_1 is OH, halogen, specifically including Br and Cl, or O_2N ;

[0014] R_2 is alkyl, alkenyl, alkynyl, alkoxy, alkylaryl, halogen, hydrogen, OH, NO_2 , thioether, amine, SH, or NH_2 ;

[0015] R_3 is:





[0016] wherein m_1 is an integer selected from 1, 2, 3, or 4,


[0017] X_5 and X_6 are each independently hydrogen, alkyl, upper alkyl, lower alkyl, aryl, alkoxy, aryloxy, cyclic alkyl, cycloalkyl, cycloarylalkyl, aralkyl, alkylester, alkylesteralkyl, alkylacetoxyl, hydroxyl, hydroxylalkyl, cyclopropyl, cyclobutyl, $-\text{CH}_3$, $-\text{CH}_2\text{OH}$, cyclopentyl, $-\text{CH}_2\text{OAc}$, $-\text{CH}_2\text{OC}(\text{O})\text{C}(\text{CH}_3)_3$, $-\text{CH}_2\text{C}_6\text{H}_5$, or cyclohexyl,

[0018] Z is NH, S, or O, and

[0019] Z_3 is alkyl or lower alkyl;

[0020] R_4 is CN, substituted amine, CH_2S -alkyl, alkyl, or CH_2N_3 ; and

[0021] R_5 and R_6 are each independently selected from the group consisting of:



[0022] monosaccharide (e.g., glucose, fructose, galactose, etc.), polysaccharide, monosaccharide derivative (e.g., an acetylated monosaccharide such as acetylated galactose,

1,2,3,4-diisopropylideno-D-g- alactose) substituted and unsubstituted aryl, and substituted and unsubstituted alkylaryl.

[0023] The compounds described herein possess useful STAT3 inhibiting activity, and may be used in the treatment or prophylaxis of a disease or condition in which STAT3 plays an active role. Thus, in broad aspect, pharmaceutical compositions comprising one or more compounds of the present invention together with a pharmaceutically acceptable carrier, as well as methods of making and using the compounds and compositions is described.

[0024] Methods for inhibiting STAT3 or STAT5 activation are also provided herein. Methods for treating a STAT3-mediated or STAT5-mediated disorder in a patient in need of such treatment comprising administering to said patient a therapeutically effective amount of a compound or composition described herein. Further, these compounds presented herein can be used in the manufacture of a medicament for the treatment of a disease or condition ameliorated by the modulation of STAT3 or STAT5 activation.

BRIEF DESCRIPTION OF THE DRAWINGS

[0025] The foregoing summary as well as the following detailed description of the preferred embodiment of the invention will be better understood when read in conjunction with the appended drawings. It should be understood, however, that the invention is not limited to the precise arrangements and instrumentalities shown herein.

[0026] For a more complete understanding of the present invention, and the advantages thereof, reference is now made to the following descriptions taken in conjunction with the accompanying drawings, in which:

[0027] Figure 1 is data from experimentation that shows the inhibition of STAT3 phosphorylation by the compound of Example 1 in Colo357FG tumor cell line.

[0028] Figure 2 provides a chart of the data show the lack of cell survival of the U87 brain tumor after administration of the compounds of Example 2 and Example 4.


[0029] Figure 3 depicts the data depicting the inhibition of the SCOV3 ovarian cancer cell growth after administration of the compounds of Example 2 and Example 4.

[0030] Figure 4 is data from the compound of Example 9 constitutively inhibiting the IL-2 phosphorylation of STAT-3 in D54 brain tumor cell lines.

[0031] Figure 5 is a schematic representation of signaling pathways which lead to STAT activation and STAT-mediated gene expression.

DETAILED DESCRIPTION

[0032] Compounds presented herein have the general Formula I:




[0033] or a pharmaceutically acceptable salt or solvate thereof, wherein:

[0034] n is an integer selected from 1, 2 or 3;

[0035] R_0 is R_1 , or R_0 is $\text{R}_1 - \text{Z}_1 -$ wherein Z_1 is alkyl, and specifically may be a lower alkyl such as $-(\text{CH}_2)_{m_3}-$ where $m_3=0, 1, 2, 3$, or 4;


[0036] R_1 is:



[0037] wherein X_1 , X_2 , X_3 , and X_4 , are each independently hydrogen, halogen, alkyl, alkoxy, OH, trihalomethyl, or NO_2 , and Y_1 is OH, halogen, specifically including Br and Cl, or O_2N ;

[0038] R_2 is alkyl, alkenyl, alkynyl, alkoxy, alkylaryl, halogen, hydrogen, OH, NO_2 , thioether, amine, SH, or NH_2 ;

[0039] R_3 is:



[0040] wherein m_1 is an integer selected from 1, 2, 3, or 4,


[0041] X_5 and X_6 are each independently hydrogen, alkyl, upper alkyl, lower alkyl, aryl, alkoxyl, aryloxy, cyclic alkyl, cycloalkyl, cycloarylalkyl, aralkyl, alkylester, alkylesteralkyl, alkylacetoxyl, hydroxyl, hydroxylalkyl, cyclopropyl, cyclobutyl, $-\text{CH}_3$, $-\text{CH}_2\text{OH}$, cyclopentyl, $-\text{CH}_2\text{OAc}$, $-\text{CH}_2\text{OC}(\text{O})\text{C}(\text{CH}_3)_3$, $-\text{CH}_2\text{C}_6\text{H}_5$, or cyclohexyl,

[0042] Z is NH, S, or O, and

[0043] Z_3 is alkyl or lower alkyl;


[0044] R_4 is CN, substituted amine, CH_2S -alkyl, alkyl, or CH_2N_3 ; and

[0045] R_5 and R_6 are each independently selected from the group consisting of:



[0046] monosaccharide (e.g., glucose, fructose, galactose, etc.), polysaccharide, monosaccharide derivative (e.g., an acetylated monosaccharide such as acetylated galactose, 1,2,3,4-diisopropylideno-D-galactose) substituted and unsubstituted aryl, and substituted and unsubstituted alkylaryl.

[0047] More specifically, R_5 may be an alkylaryl having the structure:




[0048] wherein m is an integer selected from 0, 1, 2, 3, 4, 5, 6, or 7,

[0049] X_5 and X_6 are each independently hydrogen, alkyl, upper alkyl, lower alkyl, aryl, alkoxy, aryloxy, cyclic alkyl, cycloalkyl, cycloarylalkyl, aralkyl, alkylester, alkylesteralkyl,


alkylacetoxyl, hydroxyl, hydroxylalkyl, cyclopropyl, cyclobutyl, $-\text{CH}_3$, $-\text{CH}_2\text{OH}$, cyclopentyl, $-\text{CH}_2\text{OAc}$, $-\text{CH}_2\text{OC}(\text{O})\text{C}(\text{CH}_3)_3$, $-\text{CH}_2\text{C}_6\text{H}_5$, or cyclohexyl,


[0050] and R_7 , R_8 , R_9 , R_{10} , and R_{11} , are each independently selected from the group consisting of hydrogen, halogen, alkyl, alkoxy, OH, trihalomethyl, and NO_2 .

[0051] Even more specifically, R_5 can be any one of the following:



[0052] Furthermore, in Formula I, R_1 may be specifically:





[0053] wherein X_1 may be a halogen such as Br or Cl, and

[0054] X_2 , X_3 , and X_4 , are each independently hydrogen, halogen, alkyl, alkoxy, OH, trihalomethyl, or NO_2 .

[0055] As used herein, the terms below have the meanings indicated.

[0056] As used herein, the term "amino" means —NH_2 ; the term "nitro" means —NO_2 ; the term "halo" designates —F , —Cl , —Br or —I ; the term "mercapto" means —SH ; the term "cyano" means —CN ; the term "silyl" means —SiH_3 , and the term "hydroxy" means —OH .

[0057] An "amide" refers to a —C(O)—NH—R , where R is either alkyl, aryl, alkylaryl, or hydrogen.

[0058] A "thioamide" refers to a —C(S)—NH—R , where R is either alkyl, aryl, alkylaryl, or hydrogen.

[0059] An "ester" refers to a —C(O)—OR' , where R' is either alkyl, aryl, or alkylaryl.

[0060] An "amine" refers to a —N(R'')R''' , where R'' and R''' is each independently either hydrogen, alkyl, aryl, or alkylaryl, provided that R'' and R''' are not both hydrogen.

[0061] A "thioether" refers to —S—R , where R is either alkyl, aryl, or alkylaryl.

[0062] A "sulfonyl" refers to $\text{—S(O)}_2\text{—R}$, where R is aryl, C(CN)=C-aryl , $\text{CH}_2\text{—CN}$, alkylaryl, NH-alkyl, NH-alkylaryl, or NH-aryl.

[0063] An "alkane" refers to an acyclic branched or unbranched hydrocarbon, in many cases having the general formula $\text{C}_n\text{H}_{2n+2}$.

[0064] An "alkyl" refers to a univalent group derived from an alkane by removal of a hydrogen atom from any carbon atom thus having the formula $\text{—C}_n\text{H}_{2n+1}$ in many cases. Alkyl groups include either straight-chained or branched chained and/or groups that may be substituted with additional acyclic alkyl, cycloalkyl, or cyclic alkyl groups. An alkyl group may be heteroatom-substituted or heteroatom-unsubstituted, see below. Preferably, an alkyl group has 1 to 12 carbons and may be referred to as a lower alkyl when having 1 to 7 carbons, and/or an upper alkyl when having 8 or more carbon atoms. Alkyl further refers and includes a "divalent alkyl" that refers to a divalent group derived from an alkane by removal of two hydrogen atoms from either the same carbon atom (e.g. methylene, ethylidene, propylidene) or from different carbon atoms (e.g. $\text{—C}_2\text{H}_4\text{—}$).

[0065] A "cycloalkane" refers to saturated monocyclic hydrocarbons with or without side chains.

[0066] A "cycloalkyl" specifically refers to a univalent group derived from cycloalkane by removal of a hydrogen atom from a ring carbon atom.

[0067] A cyclic alkyl and/or alkyl cyclic or alicyclic compound refers to an aliphatic compound having a carbocyclic ring structure that may be saturated or unsaturated, but are not a benzenoid or other aromatic system, and where the univalent group is derived by removal of a hydrogen atom from any carbon of alkane chain.

[0068] The term "heteroatom-substituted," when used to modify a class of organic radicals (e.g. alkyl, aryl, acyl, etc.), means that one, or more than one, hydrogen atom of that radical has been replaced by a heteroatom, or a heteroatom containing group. Examples of heteroatoms and heteroatom containing groups include: hydroxy, cyano, alkoxy, =O , =S , —NO_2 , $\text{—N(CH}_3)_2$, amino, or —SH . Specific heteroatom-substituted organic radicals are defined more fully below.

[0069] The term "heteroatom-unsubstituted," when used to modify a class of organic radicals (e.g. alkyl, aryl, acyl, etc.) means that none of the hydrogen atoms of that radical have been replaced with a heteroatom or a heteroatom containing group. Substitution of a hydrogen atom with a carbon atom, or a group consisting of only carbon and hydrogen atoms, is not sufficient to make a group heteroatom-substituted. For example, the group $\text{—C}_6\text{H}_4\text{C}\equiv\text{CH}$ is an example of a heteroatom-unsubstituted aryl group, while $\text{—C}_6\text{H}_4\text{F}$ is an example of a heteroatom-substituted aryl group. Specific heteroatom-unsubstituted organic radicals are defined more fully below.

[0070] The term "heteroatom-unsubstituted C_n-alkyl" refers to an alkyl, further having a total of n carbon atoms, all of which are nonaromatic, 3 or more hydrogen atoms, and no heteroatoms. For example, a heteroatom-unsubstituted C₁-C₁₀-alkyl has 1 to 10 carbon atoms. The groups, —CH₃, cyclopropylmethyl, —CH₂CH₃, —CH₂CH₂CH₃, —CH(CH₃)₂, —CH₂CH₂CH₂CH₃, —CH(CH₃)CH₂CH₃, —CH₂CH(CH₃)₂, —C(CH₃)₃, and —CH₂C(CH₃)₃ are all examples of heteroatom-unsubstituted alkyl groups.

[0071] The term "heteroatom-substituted C_n-alkyl" refers to an alkyl, further having a total of n carbon atoms, all of which are nonaromatic, 0, 1, or more than one hydrogen atom, at least one heteroatom, wherein each heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₁-C₁₀-alkyl has 1 to 10 carbon atoms. The following groups are all examples of heteroatom-substituted alkyl groups: trifluoromethyl, —CH₂F, —CH₂Cl, —CH₂Br, —CH₂OH, —CH₂OCH₃, —CH₂OCH₂CH₃, —CH₂OCH₂CH₂CH₃, —CH₂OCH(CH₃)₂, —CH₂OCH₂CF₃, —CH₂OCOCH₃, —CH₂NH₂, —CH₂NHCH₃, —CH₂N(CH₃)₂, —CH₂NHCH₂CH₃, —CH₂N(CH₃)CH₂CH₃, —CH₂NHCH₂CH₂CH₃, —CH₂NHCH(CH₃)₂, —CH₂OCH(CH₂)₂, —CH₂NHCH(CH₂)₂, —CH₂CH₂NHCH(CH₂)₂, —CH₂N(CH₂CH₃)₂, —CH₂CH₂F, —CH₂CH₂Cl, —CH₂CH₂Br, —CH₂CH₂I, —CH₂CH₂OH, —CH₂CH₂OCOCH₃, —CH₂CH₂NH₂, —CH₂CH₂N(CH₃)₂, —CH₂CH₂NHCH₂CH₃, —CH₂CH₂N(CH₃)CH₂CH₃, —CH₂CH₂NHCH₂CH₂CH₃, —CH₂CH₂NHCH(CH₃)₂, —CH₂CH₂N(CH₂CH₃)₂, —CH₂CH₂NHCO₂C(CH₃)₃, and —CH₂Si(CH₃)₃.

[0072] The term "heteroatom-unsubstituted C_n-cycloalkyl" refers to a cycloalkyl, further having a total of n carbon atoms, all of which are nonaromatic, 3 or more hydrogen atoms, and no heteroatoms. For example, a heteroatom-unsubstituted C₁-C₁₀-cycloalkyl has 1 to 10 carbon atoms. The groups —CH(CH₂)₂ (cyclopropyl), cyclobutyl, cyclopentyl, and cyclohexyl, are all examples of heteroatom-unsubstituted cycloalkyl groups.

[0073] The term "heteroatom-substituted C_n-cycloalkyl" refers to a cycloalkyl, further having a total of n carbon atoms, all of which are nonaromatic, 0, 1, or more than one hydrogen atom, at least one heteroatom, wherein each heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₁-C₁₀-cycloalkyl has 1 to 10 carbon atoms.

[0074] An "alkenyl" group refers to an unsaturated hydrocarbon group containing at least one carbon-carbon double bond, including straight-chain, branched-chain, and cyclic groups. Preferably, the alkenyl group has 1 to 12 carbons. More preferably it is a lower alkenyl of from

1 to 7 carbons, more preferably 1 to 4 carbons. The alkenyl group may be substituted or unsubstituted. When substituted, the substituted group(s) is preferably hydroxyl, cyano, alkoxy, =O, =S, NO₂, N(CH₃)₂, halogen, amino, or SH.

[0075] The term "heteroatom-unsubstituted C_n-alkenyl" refers to a radical, having a linear or branched, cyclic or acyclic structure, further having at least one nonaromatic carbon-carbon double bond, but no carbon-carbon triple bonds, a total of n carbon atoms, three or more hydrogen atoms, and no heteroatoms. For example, a heteroatom-unsubstituted C₂-C₁₀-alkenyl has 2 to 10 carbon atoms. Heteroatom-unsubstituted alkenyl groups include: —CH=CH₂, —CH=CHCH₃, —CH=CHCH₂CH₃, —CH=CHCH₂CH₂CH₃, —CH=CHCH(CH₃)₂, —CH=CHCH(CH₂)₂, —CH₂CH=CH₂, —CH₂CH=CHCH₃, —CH₂CH=CHCH₂CH₃, —CH₂CH=CHCH₂CH₂CH₃, —CH₂CH=CHCH(CH₃)₂, —CH₂CH=CHCH(CH₂)₂, and —CH=CH—C₆H₅.

[0076] The term "heteroatom-substituted C_n-alkenyl" refers to a radical, having a single nonaromatic carbon atom as the point of attachment and at least one nonaromatic carbon-carbon double bond, but no carbon-carbon triple bonds, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, 0, 1, or more than one hydrogen atom, and at least one heteroatom, wherein each heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₂-C₁₀-alkenyl has 2 to 10 carbon atoms. The groups, —CH=CHF, —CH=CHCl and —CH=CHBr, are examples of heteroatom-substituted alkenyl groups.

[0077] An "alkynyl" group refers to an unsaturated hydrocarbon group containing at least one carbon-carbon triple bond, including straight-chain, branched chain, and cyclic groups. Preferably, the alkynyl group has 1 to 12 carbons. More preferably it is a lower alkynyl of from 1 to 7 carbons, more preferably 1 to 4 carbons. The alkynyl group may be substituted or unsubstituted. When substituted, the substituted group(s) is preferably hydroxyl, cyano, alkoxy, =O, =S, NO₂, N(CH₃)₂, amino, or SH.

[0078] The term "heteroatom-unsubstituted C_n-alkynyl" refers to a radical, having a linear or branched, cyclic or acyclic structure, further having at least one carbon-carbon triple bond, a total of n carbon atoms, at least one hydrogen atom, and no heteroatoms. For example, a heteroatom-unsubstituted C₂-C₁₀-alkynyl has 2 to 10 carbon atoms. The groups, —C≡CH, —C≡CCH₃, and —C≡CC₆H₅ are examples of heteroatom-unsubstituted alkynyl groups.

[0079] The term "heteroatom-substituted C_n-alkynyl" refers to a radical, having a single nonaromatic carbon atom as the point of attachment and at least one carbon-carbon triple bond,

further having a linear or branched, cyclic or acyclic structure, and having a total of n carbon atoms, 0, 1, or more than one hydrogen atom, and at least one heteroatom, wherein each heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₂-C₁₀-alkynyl has 2 to 10 carbon atoms. The group, —C≡CSi(CH₃)₃, is an example of a heteroatom-substituted alkynyl group.

[0080] An "aryl" group refers to an aromatic group which has at least one ring having a conjugated pi electron system, and includes carbocyclic aryl, heterocyclic aryl, and biaryl groups, all of which may be optionally substituted. Preferably, the aryl is a substituted or unsubstituted phenyl or pyridyl. Preferred aryl substituent(s) are halogen, trihalomethyl, hydroxyl, SH, OH, NO₂, amine, thioether, cyano, alkoxy, alkyl, and amino groups.

[0081] An "alkylaryl" group refers to an alkyl (as described above), covalently joined to an aryl group (as described above). Preferably, the alkyl is a lower alkyl.

[0082] "Carbocyclic aryl" groups are groups wherein the ring atoms on the aromatic ring are all carbon atoms. The carbon atoms are optionally substituted with preferred groups as described for aryl groups above.

[0083] "Heterocyclic aryl" groups are groups having from 1 to 3 heteroatoms as ring atoms in the aromatic ring and the remainder of the ring atoms are carbon atoms. Suitable heteroatoms include oxygen, sulfur, and nitrogen, and include furanyl, thienyl, pyridyl, pyrrolyl, N-lower alkyl pyrrolo, pyrimidyl, pyrazinyl, imidazolyl, and the like, all optionally substituted.

[0084] The term "heteroatom-unsubstituted C_n-aryl" refers to a radical, having a single carbon atom as a point of attachment, wherein the carbon atom is part of an aromatic ring structure containing only carbon atoms, further having a total of n carbon atoms, 5 or more hydrogen atoms, and no heteroatoms. For example, a heteroatom-unsubstituted C₆-C₁₀-aryl has 6 to 10 carbon atoms. Examples of heteroatom-unsubstituted aryl groups include phenyl, methylphenyl, (dimethyl)phenyl, —C₆H₄CH₂CH₃, —C₆H₄CH₂CH₂CH₃, —C₆H₄CH(CH₃)₂, —C₆H₄CH(CH₂)₂, —C₆H₃(CH₃)CH₂CH₃, —C₆H₄CH=CH₂, —C₆H₄CH=CHCH₃, —C₆H₄C≡CH, —C₆H₄C≡CCH₃, naphthyl, quinolyl, indolyl, and the radical derived from biphenyl. The term "heteroatom-unsubstituted aryl" includes carbocyclic aryl groups, biaryl groups, and radicals derived from polycyclic fused hydrocarbons (PAHs).

[0085] The term "heteroatom-substituted C_n-aryl" refers to a radical, refers to a radical, having either a single aromatic carbon atom or a single aromatic heteroatom as the point of attachment, further having a total of n carbon atoms, at least one hydrogen atom, and at least one

heteroatom, further wherein each heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-unsubstituted C₁-C₁₀-heteroaryl has 1 to 10 carbon atoms. The term "heteroatom-substituted aryl" includes heteroaryl and heterocyclic aryl groups. It also includes those groups derived from the compounds: pyrrole, furan, thiophene, imidazole, oxazole, isoxazole, thiazole, isothiazole, triazole, pyrazole, pyridine, pyrazine, pyridazine, pyrimidine, and the like. Further examples of heteroatom-substituted aryl groups include the groups: —C₆H₄F, —C₆H₄Cl, —C₆H₄Br, —C₆H₄I, —C₆H₄OH, —C₅H₄OCH₃, —C₆H₄OCH₂CH₃, —C₆H₄OCOCH₃, —C₆H₄OC₆H₅, —C₆H₄NH₂, —C₆H₄NHCH₃, —C₆H₄NHCH₂CH₃, —C₆H₄CH₂Cl, —C₆H₄CH₂Br, —C₆H₄CH₂OH, —C₆H₄CH₂OCOCH₃, —C₆H₄CH₂NH₂, —C₆H₄N(CH₃)₂, —C₆H₄CH₂CH₂Cl, —C₆H₄CH₂CH₂OH, —C₆H₄CH₂CH₂OCOCH₃, —C₆H₄CH₂CH₂NH₂, —C₆H₄CH₂CH=CH₂, —C₆H₄CF₃, —C₆H₄CN, —C₆H₄C—CSi(CH₃)₃, —C₆H₄COH, —C₆H₄COCH₃, —C₆H₄COCH₂CH₃, —C₆H₄COCH₂CF₃, —C₆H₄COC₆H₅, —C₆H₄CO₂H, —C₆H₄CO₂CH₃, —C₆H₄CONH₂, —C₆H₄CONHCH₃, —C₆H₄CON(CH₃)₂, furanyl, thienyl, pyridyl, pyrrolyl, pyrimidyl, pyrazinyl, and imidazolyl.

[0086] The term "heteroatom-unsubstituted C_n-aralkyl" refers to a radical, having a single saturated carbon atom as the point of attachment, further having a total of n carbon atoms, wherein at least 6 of the carbon atoms form an aromatic ring structure containing only carbon atoms, 7 or more hydrogen atoms, and no heteroatoms. For example, a heteroatom-unsubstituted C₇-C₁₀-aralkyl has 7 to 10 carbon atoms. An "aralkyl" includes an alkyl heteroatom-substituted with an aryl group. Examples of heteroatom-unsubstituted aralkyls include phenylmethyl (benzyl) and phenylethyl.

[0087] The term "heteroatom-substituted C_n-aralkyl" refers to a radical, having a single saturated carbon atom as the point of attachment, further having a total of n carbon atoms, 0, 1, or more than one hydrogen atom, and at least one heteroatom, wherein at least one of the carbon atoms is incorporated in an aromatic ring structure, further wherein each heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₂-C₁₀-heteroaralkyl has 2 to 10 carbon atoms.

[0088] The term "heteroatom-unsubstituted C_n-acyl" refers to a radical, having a single carbon atom of a carbonyl group as the point of attachment, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, 1 or more hydrogen atoms, a total of one oxygen atom, and no additional heteroatoms. For example, a heteroatom-unsubstituted C₁-C₁₀-acyl has 1 to 10 carbon atoms. The groups, —COH, —COCH₃, —COCH₂CH₃, —COCH₂CH₂CH₃, —COCH(CH₃)₂, —COCH(CH₂)₂, —COC₆H₅,

$\text{—COC}_6\text{H}_4\text{CH}_3$, $\text{—COC}_6\text{H}_4\text{CH}_2\text{CH}_3$, $\text{—COC}_6\text{H}_4\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{—COC}_6\text{H}_4\text{CH}(\text{CH}_3)_2$, $\text{—COC}_6\text{H}_4\text{CH}(\text{CH}_2)_2$, and $\text{—COC}_6\text{H}_3(\text{CH}_3)_2$, are examples of heteroatom-unsubstituted acyl groups.

[0089] The term "heteroatom-substituted C_n -acyl" refers to a radical, having a single carbon atom as the point of attachment, the carbon atom being part of a carbonyl group, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, 0, 1, or more than one hydrogen atom, at least one additional heteroatom in addition to the oxygen of the carbonyl group, wherein each additional heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C_1 - C_{10} -acyl has 1 to 10 carbon atoms. The term heteroatom-substituted acyl includes carbamoyl, thiocarboxylate, and thiocarboxylic acid groups. The groups, $\text{—COCH}_2\text{CF}_3$, $\text{—CO}_2\text{H}$, $\text{—CO}_2\text{CH}_3$, $\text{—CO}_2\text{CH}_2\text{CH}_3$, $\text{—CO}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{—CO}_2\text{CH}(\text{CH}_3)_2$, $\text{—CO}_2\text{CH}(\text{CH}_2)_2$, —CONH_2 , —CONHCH_3 , $\text{—CONHCH}_2\text{CH}_3$, $\text{—CONHCH}_2\text{CH}_2\text{CH}_3$, $\text{—CONHCH}(\text{CH}_3)_2$, $\text{—CONHCH}(\text{CH}_2)_2$, $\text{—CON}(\text{CH}_3)_2$, $\text{—CON}(\text{CH}_2\text{CH}_3)\text{CH}_3$, $\text{—CON}(\text{CH}_2\text{CH}_3)_2$ and $\text{—CONHCH}_2\text{CF}_3$, are examples heteroatom-substituted acyl groups.

[0090] An "alkoxy" group refers to an " —O-alkyl " group, where "alkyl" is defined above.

[0091] An "aryl" group refers to an aromatic group which has at least one ring having a conjugated pi electron system, and includes carbocyclic aryl, heterocyclic aryl, and biaryl groups, all of which may be optionally substituted. Preferably, the aryl is a substituted or unsubstituted phenyl or pyridyl. Preferred aryl substituent(s) are halogen, trihalomethyl, hydroxyl, SH, OH, NO_2 , amine, thioether, cyano, alkoxy, alkyl, and amino groups.

[0092] The term "heteroatom-unsubstituted C_n -alkoxy" refers to a group, having the structure —OR , in which R is a heteroatom-unsubstituted C_n -alkyl, as that term is defined above. Heteroatom-unsubstituted alkoxy groups include: —OCH_3 , $\text{—OCH}_2\text{CH}_3$, $\text{—OCH}_2\text{CH}_2\text{CH}_3$, $\text{—OCH}(\text{CH}_3)_2$, and $\text{—OCH}(\text{CH}_2)_2$.

[0093] The term "heteroatom-substituted C_n -alkoxy" refers to a group, having the structure —OR , in which R is a heteroatom-substituted C_n -alkyl, as that term is defined above. For example, $\text{—OCH}_2\text{CF}_3$ is a heteroatom-substituted alkoxy group.

[0094] The term "heteroatom-unsubstituted C_n -alkenyloxy" refers to a group, having the structure —OR , in which R is a heteroatom-unsubstituted C_n -alkenyl, as that term is defined above.

[0095] The term "heteroatom-substituted C_n-alkenyloxy" refers to a group, having the structure —OR, in which R is a heteroatom-substituted C_n-alkenyl, as that term is defined above.

[0096] The term "heteroatom-unsubstituted C_n-alkynyloxy" refers to a group, having the structure —OR, in which R is a heteroatom-unsubstituted C_n-alkynyl, as that term is defined above.

[0097] The term "heteroatom-substituted C_n-alkynyloxy" refers to a group, having the structure —OR, in which R is a heteroatom-substituted C_n-alkynyl, as that term is defined above.

[0098] The term "heteroatom-unsubstituted C_n-aryloxy" refers to a group, having the structure —OAr, in which Ar is a heteroatom-unsubstituted C_n-aryl, as that term is defined above. An example of a heteroatom-unsubstituted aryloxy group is —OC₆H₅.

[0099] The term "heteroatom-substituted C_n-aryloxy" refers to a group, having the structure —OAr, in which Ar is a heteroatom-substituted C_n-aryl, as that term is defined above.

[0101] The term "heteroatom-unsubstituted C_n-aralkyloxy" refers to a group, having the structure —OAr, in which Ar is a heteroatom-unsubstituted C_n-aralkyl, as that term is defined above.

[0102] The term "heteroatom-substituted C_n-aralkyloxy" refers to a group, having the structure —OAr, in which Ar is a heteroatom-substituted C_n-aralkyl, as that term is defined above.

[0103] The term "heteroatom-unsubstituted C_n-acyloxy" refers to a group, having the structure —OAc, in which Ac is a heteroatom-unsubstituted C_n-acyl, as that term is defined above. A heteroatom-unsubstituted acyloxy group includes alkylcarbonyloxy and arylcarbonyloxy groups. For example, —OCOCH₃ is an example of a heteroatom-unsubstituted acyloxy group.

[0104] The term "heteroatom-substituted C_n-acyloxy" refers to a group, having the structure —OAc, in which Ac is a heteroatom-substituted C_n-acyl, as that term is defined above. A heteroatom-substituted acyloxy group includes alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, alkoxycarbonyl, aminocarbonyl, and alkylthiocarbonyl groups.

[0105] The term "heteroatom-unsubstituted C_n-alkylamino" refers to a radical, having a single nitrogen atom as the point of attachment, further having one or two saturated carbon atoms attached to the nitrogen atom, further having a linear or branched, cyclic or acyclic structure, containing a total of n carbon atoms, all of which are nonaromatic, 4 or more hydrogen atoms, a

total of 1 nitrogen atom, and no additional heteroatoms. For example, a heteroatom-unsubstituted C_1 - C_{10} -alkylamino has 1 to 10 carbon atoms. The term "heteroatom-unsubstituted C_n -alkylamino" includes groups, having the structure $-NHR$, in which R is a heteroatom-unsubstituted C_n -alkyl, as that term is defined above. A heteroatom-unsubstituted alkylamino group would include $-NHCH_3$, $-NHCH_2CH_3$, $-NHCH_2CH_2CH_3$, $-NHCH(CH_3)_2$, $-NHCH(CH_2)_2$, $-NHCH_2CH_2CH_2CH_3$, $-NHCH(CH_3)CH_2CH_3$, $-NHCH_2CH(CH_3)_2$, $-NHC(CH_3)_3$, $-N(CH_3)_2$, $-N(CH_3)CH_2CH_3$, $-N(CH_2CH_3)_2$, N-pyrrolidinyl, and N-piperidinyl.

[0106] The term "heteroatom-substituted C_n -alkylamino" refers to a radical, having a single nitrogen atom as the point of attachment, further having one or two saturated carbon atoms attached to the nitrogen atom, no carbon-carbon double or triple bonds, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, all of which are nonaromatic, 0, 1, or more than one hydrogen atom, and at least one additional heteroatom, that is, in addition to the nitrogen atom at the point of attachment, wherein each additional heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C_1 - C_{10} -alkylamino has 1 to 10 carbon atoms. The term "heteroatom-substituted C_n -alkylamino" includes groups, having the structure $-NHR$, in which R is a heteroatom-substituted C_n -alkyl, as that term is defined above.

[0107] The term "heteroatom-unsubstituted C_n -alkenylamino" refers to a radical, having a single nitrogen atom as the point of attachment, further having one or two carbon atoms attached to the nitrogen atom, further having a linear or branched, cyclic or acyclic structure, containing at least one nonaromatic carbon-carbon double bond, a total of n carbon atoms, 4 or more hydrogen atoms, a total of one nitrogen atom, and no additional heteroatoms. For example, a heteroatom-unsubstituted C_2 - C_{10} -alkenylamino has 2 to 10 carbon atoms. The term "heteroatom-unsubstituted C_n -alkenylamino" includes groups, having the structure $-NHR$, in which R is a heteroatom-unsubstituted C_n -alkenyl, as that term is defined above. Examples of heteroatom-unsubstituted C_n -alkenylamino groups also include dialkenylamino and alkyl(alkenyl)amino groups.

[0108] The term "heteroatom-substituted C_n -alkenylamino" refers to a radical, having a single nitrogen atom as the point of attachment and at least one nonaromatic carbon-carbon double bond, but no carbon-carbon triple bonds, further having one or two carbon atoms attached to the nitrogen atom, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, 0, 1, or more than one hydrogen atom, and at least one

additional heteroatom, that is, in addition to the nitrogen atom at the point of attachment, wherein each additional heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₂-C₁₀-alkenylamino has 2 to 10 carbon atoms. The term "heteroatom-substituted C_n-alkenylamino" includes groups, having the structure —NHR, in which R is a heteroatom-substituted C_n-alkenyl, as that term is defined above.

[0109] The term "heteroatom-unsubstituted C_n-alkynylamino" refers to a radical, having a single nitrogen atom as the point of attachment, further having one or two carbon atoms attached to the nitrogen atom, further having a linear or branched, cyclic or acyclic structure, containing at least one carbon-carbon triple bond, a total of n carbon atoms, at least one hydrogen atom, a total of one nitrogen atom, and no additional heteroatoms. For example, a heteroatom-unsubstituted C₂-C₁₀-alkynylamino has 2 to 10 carbon atoms. The term "heteroatom-unsubstituted C_n-alkynylamino" includes groups, having the structure —NHR, in which R is a heteroatom-unsubstituted C_n-alkynyl, as that term is defined above. An alkynylamino group includes dialkynylamino and alkyl(alkynyl)amino groups.

[0110] The term "heteroatom-substituted C_n-alkynylamino" refers to a radical, having a single nitrogen atom as the point of attachment, further having one or two carbon atoms attached to the nitrogen atom, further having at least one nonaromatic carbon-carbon triple bond, further having a linear or branched, cyclic or acyclic structure, and further having a total of n carbon atoms, 0, 1, or more than one hydrogen atom, and at least one additional heteroatom, that is, in addition to the nitrogen atom at the point of attachment, wherein each additional heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₂-C₁₀-alkynylamino has 2 to 10 carbon atoms. The term "heteroatom-substituted C_n-alkynylamino" includes groups, having the structure —NHR, in which R is a heteroatom-substituted C_n-alkynyl, as that term is defined above.

[0111] The term "heteroatom-unsubstituted C_n-arylamino" refers to a radical, having a single nitrogen atom as the point of attachment, further having at least one aromatic ring structure attached to the nitrogen atom, wherein the aromatic ring structure contains only carbon atoms, further having a total of n carbon atoms, 6 or more hydrogen atoms, a total of one nitrogen atom, and no additional heteroatoms. For example, a heteroatom-unsubstituted C₆-C₁₀-arylamino has 6 to 10 carbon atoms. The term "heteroatom-unsubstituted C_n-arylamino" includes groups, having the structure —NHR, in which R is a heteroatom-unsubstituted C_n-aryl,

as that term is defined above. A heteroatom-unsubstituted arylamino group includes diarylamino and alkyl(aryl)amino groups.

[0112] The term "heteroatom-substituted C_n-arylamino" refers to a radical, having a single nitrogen atom as the point of attachment, further having a total of n carbon atoms, at least one hydrogen atom, at least one additional heteroatoms, that is, in addition to the nitrogen atom at the point of attachment, wherein at least one of the carbon atoms is incorporated into one or more aromatic ring structures, further wherein each additional heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₆-C₁₀-arylamino has 6 to 10 carbon atoms. The term "heteroatom-substituted C_n-arylamino" includes groups, having the structure —NHR, in which R is a heteroatom-substituted C_n-aryl, as that term is defined above. A heteroatom-substituted arylamino group includes heteroarylamino groups.

[0113] The term "heteroatom-unsubstituted C_n-aralkylamino" refers to a radical, having a single nitrogen atom as the point of attachment, further having one or two saturated carbon atoms attached to the nitrogen atom, further having a total of n carbon atoms, wherein at least 6 of the carbon atoms form an aromatic ring structure containing only carbon atoms, 8 or more hydrogen atoms, a total of one nitrogen atom, and no additional heteroatoms. For example, a heteroatom-unsubstituted C₇-C₁₀-aralkylamino has 7 to 10 carbon atoms. The term "heteroatom-unsubstituted C_n-aralkylamino" includes groups, having the structure —NHR, in which R is a heteroatom-unsubstituted C_n-aralkyl, as that term is defined above. An aralkylamino group includes diaralkylamino groups.

[0114] The term "heteroatom-substituted C_n-aralkylamino" refers to a radical, having a single nitrogen atom as the point of attachment, further having at least one or two saturated carbon atoms attached to the nitrogen atom, further having a total of n carbon atoms, 0, 1, or more than one hydrogen atom, at least one additional heteroatom, that is, in addition to the nitrogen atom at the point of attachment, wherein at least one of the carbon atom incorporated into an aromatic ring, further wherein each heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₇-C₁₀-aralkylamino has 7 to 10 carbon atoms. The term "heteroatom-substituted C_n-aralkylamino" includes groups, having the structure —NHR, in which R is a heteroatom-substituted C_n-aralkyl, as that term is defined above. The term "heteroatom-substituted aralkylamino" includes the term "heteroaralkylamino."

[0115] The term "heteroatom-unsubstituted C_n-amido" refers to a radical, having a single nitrogen atom as the point of attachment, further having a carbonyl group attached via its carbon atom to the nitrogen atom, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, 1 or more hydrogen atoms, a total of one oxygen atom, a total of one nitrogen atom, and no additional heteroatoms. For example, a heteroatom-unsubstituted C₁-C₁₀-amido has 1 to 10 carbon atoms. The term "heteroatom-unsubstituted C_n-amido" includes groups, having the structure —NHR, in which R is a heteroatom-unsubstituted C_n-acyl, as that term is defined above. The term amido includes N-alkyl-amido, N-aryl-amido, N-aralkyl-amido, acylamino, alkylcarbonylamino, arylcarbonylamino, and ureido groups. The group, —NHCOCH₃, is an example of a heteroatom-unsubstituted amido group.

[0116] The term "heteroatom-substituted C_n-amido" refers to a radical, having a single nitrogen atom as the point of attachment, further having a carbonyl group attached via its carbon atom to the nitrogen atom, further having a linear or branched, cyclic or acyclic structure, further having a total of n aromatic or nonaromatic carbon atoms, 0, 1, or more than one hydrogen atom, at least one additional heteroatom in addition to the oxygen of the carbonyl group, wherein each additional heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₁-C₁₀-amido has 1 to 10 carbon atoms. The term "heteroatom-substituted C_n-amido" includes groups, having the structure —NHR, in which R is a heteroatom-unsubstituted C_n-acyl, as that term is defined above. The group, —NHCO₂CH₃, is an example of a heteroatom-substituted amido group.

[0117] The term "heteroatom-unsubstituted C_n-sulfonamido" refers to a radical, having a single nitrogen atom as the point of attachment, further having a sulfonyl group attached via its sulfur atom to the nitrogen atom, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, 1 or more hydrogen atoms, a total of one oxygen atom, a total of one nitrogen atom, and no additional heteroatoms. For example, a heteroatom-unsubstituted C₁-C₁₀-amido has 1 to 10 carbon atoms. The term amido includes N-alkyl-sulfonamido, N-aryl-sulfonamido, N-aralkyl-sulfonamido, sulfonylamino, alkylsulfonamino, and arylsulfonamino groups. The group, —NHS(O)₂CH₃, is an example of a heteroatom-unsubstituted sulfonamido group.

[0118] The term "heteroatom-substituted C_n-sulfonamido" refers to a radical, having a single nitrogen atom as the point of attachment, further having a sulfonyl group attached via its sulfur atom to the nitrogen atom, further having a linear or branched, cyclic or acyclic structure, further having a total of n aromatic or nonaromatic carbon atoms, 0, 1, or more than one

hydrogen atom, at least one additional heteroatom in addition to the sulfur and oxygen atoms of the sulfonyl group, wherein each additional heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₁-C₁₀-sulfonamido has 1 to 10 carbon atoms. The group, —NHS(O)₂OCH₃, is an example of a heteroatom-substituted sulfonamido group.

[0119] The term "heteroatom-unsubstituted C_n-alkylthio" refers to a group, having the structure —SR, in which R is a heteroatom-unsubstituted C_n-alkyl, as that term is defined above. The group, —SCH₃, is an example of a heteroatom-unsubstituted alkylthio group.

[0120] The term "heteroatom-substituted C_n-alkylthio" refers to a group, having the structure —SR, in which R is a heteroatom-substituted C_n-alkyl, as that term is defined above.

[0121] The term "heteroatom-unsubstituted C_n-alkenylthio" refers to a group, having the structure —SR, in which R is a heteroatom-unsubstituted C_n-alkenyl, as that term is defined above.

[0122] The term "heteroatom-substituted C_n-alkenylthio" refers to a group, having the structure —SR, in which R is a heteroatom-substituted C_n-alkenyl, as that term is defined above.

[0123] The term "heteroatom-unsubstituted C_n-alkynylthio" refers to a group, having the structure —SR, in which R is a heteroatom-unsubstituted C_n-alkynyl, as that term is defined above.

[0124] The term "heteroatom-substituted C_n-alkynylthio" refers to a group, having the structure —SR, in which R is a heteroatom-substituted C_n-alkynyl, as that term is defined above.

[0125] The term "heteroatom-unsubstituted C_n-arylthio" refers to a group, having the structure —SAr, in which Ar is a heteroatom-unsubstituted C_n-aryl, as that term is defined above. The group, —SC₆H₅, is an example of a heteroatom-unsubstituted arylthio group.

[0126] The term "heteroatom-substituted C_n-arylthio" refers to a group, having the structure —SAr, in which Ar is a heteroatom-substituted C_n-aryl, as that term is defined above.

[0127] The term "heteroatom-unsubstituted C_n-aralkylthio" refers to a group, having the structure —SAr, in which Ar is a heteroatom-unsubstituted C_n-aralkyl, as that term is defined above. The group, —SCH₂C₆H₅, is an example of a heteroatom-unsubstituted aralkyl group.

[0128] The term "heteroatom-substituted C_n-aralkylthio" refers to a group, having the structure —SAr, in which Ar is a heteroatom-substituted C_n-aralkyl, as that term is defined above.

[0129] The term "heteroatom-unsubstituted C_n-acylthio" refers to a group, having the structure —SAc, in which Ac is a heteroatom-unsubstituted C_n-acyl, as that term is defined above. The group, —SCOCH₃, is an example of a heteroatom-unsubstituted acylthio group.

[0130] The term "heteroatom-substituted C_n-acylthio" refers to a group, having the structure —SAc, in which Ac is a heteroatom-substituted C_n-acyl, as that term is defined above.

[0131] The term "heteroatom-unsubstituted C_n-alkylsilyl" refers to a radical, having a single silicon atom as the point of attachment, further having one, two, or three saturated carbon atoms attached to the silicon atom, further having a linear or branched, cyclic or acyclic structure, containing a total of n carbon atoms, all of which are nonaromatic, 5 or more hydrogen atoms, a total of 1 silicon atom, and no additional heteroatoms. For example, a heteroatom-unsubstituted C₁-C₁₀-alkylsilyl has 1 to 10 carbon atoms. An alkylsilyl group includes dialkylamino groups. The groups, —Si(CH₃)₃ and —Si(CH₃)₂C(CH₃)₃, are examples of heteroatom-unsubstituted alkylsilyl groups.

[0132] The term "heteroatom-substituted C_n-alkylsilyl" refers to a radical, having a single silicon atom as the point of attachment, further having at least one, two, or three saturated carbon atoms attached to the silicon atom, no carbon-carbon double or triple bonds, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, all of which are nonaromatic, 0, 1, or more than one hydrogen atom, and at least one additional heteroatom, that is, in addition to the silicon atom at the point of attachment, wherein each additional heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₁-C₁₀-alkylsilyl has 1 to 10 carbon atoms.

[0133] The term "pharmaceutically acceptable salts," as used herein, refers to salts of compounds of this invention that are substantially non-toxic to living organisms. Typical pharmaceutically acceptable salts include those salts prepared by reaction of a compound of this invention with an inorganic or organic acid, or an organic base, depending on the substituents present on the compounds of the invention.

[0134] Examples of inorganic acids which may be used to prepare pharmaceutically acceptable salts include: hydrochloric acid, phosphoric acid, sulfuric acid, hydrobromic acid, hydroiodic acid, phosphorous acid and the like. Examples of organic acids which may be used to prepare pharmaceutically acceptable salts include: aliphatic mono- and dicarboxylic acids, such as oxalic acid, carbonic acid, citric acid, succinic acid, phenyl-heteroatom-substituted alkanolic acids, aliphatic and aromatic sulfuric acids and the like. Pharmaceutically acceptable salts prepared from inorganic or organic acids thus include hydrochloride, hydrobromide, nitrate,

sulfate, pyrosulfate, bisulfate, sulfite, bisulfate, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, hydroiodide, hydrofluoride, acetate, propionate, formate, oxalate, citrate, lactate, p-toluenesulfonate, methanesulfonate, maleate, and the like. Other suitable salts are known to one of ordinary skill in the art.

[0135] Suitable pharmaceutically acceptable salts may also be formed by reacting the agents of the invention with an organic base such as methylamine, ethylamine, ethanolamine, lysine, ornithine and the like. Other suitable salts are known to one of ordinary skill in the art.

[0136] Pharmaceutically acceptable salts include the salts formed between carboxylate or sulfonate groups found on some of the compounds of this invention and inorganic cations, such as sodium, potassium, ammonium, or calcium, or such organic cations as isopropylammonium, trimethylammonium, tetramethylammonium, and imidazolium.

[0137] It should be recognized that the particular anion or cation forming a part of any salt of this invention is not critical, so long as the salt, as a whole, is pharmacologically acceptable and as long as the anion or cation does not contribute undesired qualities or effects. Further, additional pharmaceutically acceptable salts are known to those skilled in the art, and may be used within the scope of the invention. Additional examples of pharmaceutically acceptable salts and their methods of preparation and use are presented in *Pharmaceutical Salts: Properties, Selection and Use-A Handbook* (2002), which is incorporated herein by reference.

[0138] As used herein, the term "patient" is intended to include living organisms in which certain conditions as described herein can occur. Examples include humans, monkeys, cows, sheep, goats, dogs, cats, mice, rats, and transgenic species thereof. In a preferred embodiment, the patient is a primate. In an even more preferred embodiment, the primate is a human. Other examples of subjects include experimental animals such as mice, rats, dogs, cats, goats, sheep, pigs, and cows. The experimental animal can be an animal model for a disorder, e.g., a transgenic mouse with an Alzheimer's-type neuropathology. A patient can be a human suffering from a neurodegenerative disease, such as Alzheimer's disease, or Parkinson's disease.

[0139] As used herein, the term " IC_{50} " refers to an inhibitory dose which is 50% of the maximum response obtained.

[0140] As used herein, the term "water soluble" means that the compound dissolves in water at least to the extent of 0.010 mole/liter or is classified as soluble according to literature precedence.

[0141] As used herein, "predominantly one enantiomer" means that the compound contains at least 85% of one enantiomer, or more preferably at least 90% of one enantiomer, or even more preferably at least 95% of one enantiomer, or most preferably at least 99% of one enantiomer. Similarly, the phrase "substantially free from other optical isomers" means that the composition contains at most 5% of another enantiomer or diastereomer, more preferably 2% of another enantiomer or diastereomer, and most preferably 1% of another enantiomer or diastereomer.

[0142] As used herein the specification, "a" or "an" may mean one or more. As used herein in the claim(s), when used in conjunction with the word "comprising" or "having," the words "a" or "an" may mean one or more than one. As used herein "another" may mean at least a second or more.

[0143] Other abbreviations used herein are as follows: DMSO, dimethyl sulfoxide; iNOS, inducible nitric oxide synthase; COX-2, cyclooxygenase-2; NGF, nerve growth factor; IBMX, isobutylmethylxanthine; FBS, fetal bovine serum; GPDH, glycerol 3-phosphate dehydrogenase; RXR, retinoid X receptor; TGF- β , transforming growth factor- β ; IFN- γ , interferon- γ ; LPS, bacterial endotoxic lipopolysaccharide; TNF- α , tumor necrosis factor- α ; IL-1 β , interleukin-1 β ; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; MTT, 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide; TCA, trichloroacetic acid; HO-1, inducible heme oxygenase.

[0144] The term "combination therapy" means the administration of two or more therapeutic agents to treat a therapeutic condition or disorder described in the present disclosure. Such administration encompasses co-administration of these therapeutic agents in a substantially simultaneous manner, such as in a single capsule having a fixed ratio of active ingredients or in multiple, separate capsules for each active ingredient. In addition, such administration also encompasses use of each type of therapeutic agent in a sequential manner. In either case, the treatment regimen will provide beneficial effects of the drug combination in treating the conditions or disorders described herein.

[0145] The phrase "therapeutically effective" is intended to qualify the amount of active ingredients used in the treatment of a disease or disorder. This amount will achieve the goal of reducing or eliminating the said disease or disorder.

[0146] The term "therapeutically acceptable" refers to those compounds (or salts, prodrugs, tautomers, zwitterionic forms, etc.) which are suitable for use in contact with the

tissues of patients without undue toxicity, irritation, and allergic response, are commensurate with a reasonable benefit/risk ratio, and are effective for their intended use.

[0147] As used herein, reference to "treatment" of a patient is intended to include prophylaxis. The term "patient" means all mammals including humans. Examples of patients include humans, cows, dogs, cats, goats, sheep, pigs, and rabbits. Preferably, the patient is a human.

[0148] The term "prodrug" refers to a compound that is made more active in vivo. Certain compounds provided herein may also exist as prodrugs, as described in *Hydrolysis in Drug and Prodrug Metabolism: Chemistry, Biochemistry, and Enzymology* (Testa, Bernard and Mayer, Joachim M. Wiley-VHCA, Zurich, Switzerland 2003). Prodrugs of the compounds as described herein are structurally modified forms of the compound that readily undergo chemical changes under physiological conditions to provide the compound. Additionally, prodrugs can be converted to the compound by chemical or biochemical methods in an ex vivo environment. For example, prodrugs can be slowly converted to a compound when placed in a transdermal patch reservoir with a suitable enzyme or chemical reagent. Prodrugs are often useful because, in some situations, they may be easier to administer than the compound, or parent drug. They may, for instance, be bioavailable by oral administration whereas the parent drug is not. The prodrug may also have improved solubility in pharmaceutical compositions over the parent drug.

[0149] Furthermore, a wide variety of prodrug derivatives are known in the art, such as those that rely on hydrolytic cleavage or oxidative activation of the prodrug. An example, without limitation, of a prodrug would be a compound which is administered as an ester (the "prodrug"), but then is metabolically hydrolyzed to the carboxylic acid, the active entity. Additional examples include peptidyl derivatives of a compound.

[0150] While it may be possible for the compounds of the subject invention to be administered as the raw chemical, it is also possible to present them as a pharmaceutical formulation. Accordingly, the subject invention provides a pharmaceutical formulation comprising a compound or a pharmaceutically acceptable salt, ester, prodrug or solvate thereof, together with one or more pharmaceutically acceptable carriers thereof and optionally one or more other therapeutic ingredients. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. Proper formulation is dependent upon the route of administration chosen. Any of the well-known techniques, carriers, and excipients may be used as suitable and as understood in the art; e.g., in Remington's Pharmaceutical Sciences. The pharmaceutical compositions of the

present invention may be manufactured in a manner that is itself known, *e.g.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or compression processes.

[0151] The formulations include those suitable for oral, parenteral (including subcutaneous, intradermal, intramuscular, intravenous, intraarticular, and intramedullary), intraperitoneal, transmucosal, transdermal, rectal and topical (including dermal, buccal, sublingual and intraocular) administration although the most suitable route may depend upon for example the condition and disorder of the recipient. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing into association a compound of the subject invention or a pharmaceutically acceptable salt, ester, prodrug or solvate thereof ("active ingredient") with the carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both and then, if necessary, shaping the product into the desired formulation.

[0152] Formulations suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be presented as a bolus, electuary or paste.

[0153] Pharmaceutical preparations which can be used orally include tablets, push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. Tablets may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with binders, inert diluents, or lubricating, surface active or dispersing agents. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active ingredient therein. All formulations for oral administration should be in dosages suitable for such administration. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable

liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

[0154] The compounds may be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, *e.g.*, in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in powder form or in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, saline or sterile pyrogen-free water, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

[0155] Formulations for parenteral administration include aqueous and non-aqueous (oily) sterile injection solutions of the active compounds which may contain antioxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

[0156] Preferred unit dosage formulations are those containing an effective dose, as herein below recited, or an appropriate fraction thereof, of the active ingredient.

[0157] It should be understood that in addition to the ingredients particularly mentioned above, the formulations may include other agents conventional in the art having regard to the

type of formulation in question, for example those suitable for oral administration may include flavoring agents.

[0158] The compounds provided herein can be administered in various modes, *e.g.* orally, topically, or by injection. The precise amount of compound administered to a patient will be the responsibility of the attendant physician. The specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diets, time of administration, route of administration, rate of excretion, drug combination, the precise disorder being treated, and the severity of the indication or condition being treated. Also, the route of administration may vary depending on the condition and its severity.

[0159] In short, the compounds of the invention may be administered orally, topically, or by injection at a dose of from 0.1 to 500 mg/kg per day. Tablets or other forms of presentation provided in discrete units may conveniently contain an amount of compound of the invention which is effective at such dosage or as a multiple of the same, for instance, units containing 5 mg to 500 mg.

[0160] The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration.

[0161] In certain instances, it may be appropriate to administer at least one of the compounds described herein (or a pharmaceutically acceptable salt, ester, or prodrug thereof) in combination with another therapeutic agent. By way of example only, if one of the side effects experienced by a patient upon receiving one of the compounds herein is hypertension, then it may be appropriate to administer an anti-hypertensive agent in combination with the initial therapeutic agent. Or, by way of example only, the therapeutic effectiveness of one of the compounds described herein may be enhanced by administration of an adjuvant (*i.e.*, by itself the adjuvant may only have minimal therapeutic benefit, but in combination with another therapeutic agent, the overall therapeutic benefit to the patient is enhanced). Or, by way of example only, the benefit experienced by a patient may be increased by administering one of the compounds described herein with another therapeutic agent (which also includes a therapeutic regimen) that also has therapeutic benefit. By way of example only, in a treatment for cancer involving administration of one of the compounds described herein, increased therapeutic benefit may result by also providing the patient with another therapeutic agent for cancer. In any case, regardless of the disease, disorder or condition being treated, the overall benefit experienced by

the patient may simply be additive of the two therapeutic agents or the patient may experience a synergistic benefit.

[0162] In any case, the multiple therapeutic agents (at least one of which is a compound provided herein) may be administered in any order or even simultaneously. If simultaneously, the multiple therapeutic agents may be provided in a single, unified form, or in multiple forms (by way of example only, either as a single pill or as two separate pills). One of the therapeutic agents may be given in multiple doses, or both may be given as multiple doses. If not simultaneous, the timing between the multiple doses may be any duration of time ranging from a few minutes to four weeks.

[0163] Thus, methods for treating STAT3-mediated disorders or conditions (as well as STAT5-mediated disorders or conditions) in a human or animal subject in need of such treatment comprise administering to the subject an amount of a compound presented herein effective to reduce or prevent said disorder in the subject in combination with at least one additional agent for the treatment of said disorder that is known in the art. In a related aspect, therapeutic compositions comprise at least one these compounds in combination with one or more additional agents for the treatment of STAT3- and STAT5-mediated disorders.

[0164] The compounds described herein may be useful for the treatment of a wide variety of disorders or conditions where inhibition or modulation of the STAT3 pathway is needed. The compounds decrease and can therapeutically treat proliferative disease conditions. Uses for these compounds include the compounds as agents to: decrease STAT3 activity; STAT3 phosphorylation; and the expression of proteins controlled by transcriptional activation by activated STAT3. Along these lines, the compounds can be specifically used as agents to decrease VEGF, MMP9, MMP2, survivin, c-Myc, MMP-1, MEK-5, c-FOS, COX-2, Bcl-xl, MMP-10, HSP-27 and Jmjdla.

[0165] Disorders or conditions that can be prevented or treated by compounds and methods described herein include the prevention or treatment of cancer, such as cutaneous T-cell leukemia, head and neck tumors, pancreatic cancer, bladder cancer, high grade gliomas, brain metastasis, melanoma, skin cancer, lung cancer, breast cancer, prostate cancer, colon cancer, leukemia, myelodysplastic syndrome (a pre-leukemia condition), and multiple myeloma. In general, metastasis of any cancer can be prevented or treated with the compounds and methods described herein. The compounds can also be used to prevent or treat proliferative angiogenic conditions including telangiectasia, venous angiomas, hemangioblastoma.

[0166] These compounds and methods can also be used to prevent or treat proliferative diseases or disorders of the skin, including topical dermatitis, psoriasis, and rosacea.

[0167] Furthermore, the compounds and methods that are described herein can be used to prevent or treat Central Nervous System (“CNS”) diseases and conditions such as CNS inflammatory and conditions, *e.g.*, multiple sclerosis and progressive multifocal leukoencephalopathy.

[0168] Moreover, the compounds and methods that are described herein can be used to prevent or treat inflammatory diseases and conditions, such as osteoarthritis, Rheumatoid arthritis, Crohn’s disease, ulcerative colitis, and auto-immune diseases such as lupus and mixed auto-immune disease.

[0169] Diseases and conditions such as telangiectasia, venous angiomas, hemangioblastoma, and polycythemia vera may also be advantageously prevented or treated with the compounds and methods described herein.

[0170] These compounds and methods can affect stem cell survival and differentiation by maintaining stem cell stemness, *e.g.*, preventing the differentiation of stem cells.

[0171] The compounds taught herein may also be used for the augmentation of immune response, particularly where the augmentation of the immune response leads to the expression of costimulatory molecules on the peripheral macrophages and tumor-infiltrating microglia. These compounds are also useful when the immune response leads to proliferation of effector T cells and/or up-regulation of several key intracellular signaling molecules that critically regulate T-cell and monocyte activation. The compounds are useful when the immune responses leads to up-regulation of several key intracellular signaling molecules that critically regulate T-cell and monocyte activation, particularly phosphorylation of Syk (Tyr(352)) in monocytes and ZAP-70 (Tyr (319)) in T cells.

[0172] Table 1 as found immediately below describes certain of the diseases and conditions that may be treated using the compounds described herein:

Table 1
Activation of STATs in Human Cancers

<u>Tumor Type</u>	<u>Activated STAT</u>
<i>Blood Tumors</i>	
Multiple myeloma	STAT1, STAT3
Leukaemias:	
HTLV-1-dependent	STAT3, STAT5
Erythroleukaemia	STAT1, STAT5
Acute myelogenous leukaemia (AML)	STAT1, STAT3, STAT5
Chronic myelogenous leukaemia (CML)	STAT5
Large granular lymphocyte leukaemia (LGL)	STAT3
Lymphomas:	
EBV-related/Burkitt's	STAT3
Mycosis fungoides	STAT3
Cutaneous T-cell lymphoma	STAT3
Non-Hodgkins lymphoma (NHL)	STAT3
Anaplastic large-cell lymphoma (ALCL)	STAT3
<i>Solid Tumors</i>	
Breast Cancer	STAT1, STAT3, STAT5
Head and neck cancer	STAT1, STAT3, STAT5
Melanoma	STAT3
Ovarian cancer	STAT3
Lung cancer	STAT3
Pancreatic cancer	STAT3
Prostate cancer	STAT3
Glioma	STAT3, STAT5

Based on references cited in REFS 12, 17, EBV, Epstein-Barr virus; HTLV-1, human T-lymphotrophic virus-1.

[0173] The STAT ("signal transducer and activator of transcription") protein family includes transcription factors which are specifically activated to regulate gene transcription when cells encounter cytokines and growth factors. In mammals, there are seven known members of the STAT family of proteins: STAT1, STAT2, STAT3, STAT4, STAT5a, STAT5b, and STAT6. Yu, H., Jove, R., *The STATS of Cancer – New Molecular Targets Come of Age*, Nature Reviews, 4:97-105 (2004). These seven proteins range in size from 750 and 850 amino acids. The STAT5a and STAT5b proteins, collectively referred to herein as "STAT5," are closely

related but encoded by different genes. Yu, H., Jove, R., *The STATS of Cancer – New Molecular Targets Come of Age*, Nature Reviews, 4:97-105 (2004).

[0174] STAT proteins act as signal transducers in the cytoplasm and transcription activators in the nucleus. Kisseleva T., Bhattacharya S., Braunstein J., Schindler C.W., *Signaling Through the JAK/STAT Pathway, Recent Advances and Future Challenges*, Gene 285: 1-24 (2002). For example, STAT proteins transduce signals from cytokine receptors, growth factor receptors, and non-receptor, cytoplasmic tyrosine kinases such as Src and Abl to the nucleus of a cell, where they bind DNA and regulate transcription of an assortment of genes. *Id.* As a result, STAT proteins regulate physiological functions such as immune response, inflammation, proliferation, differentiation, survival, metastasis, apoptosis, and immuno tolerance (e.g. tumor immune evasion). Xie, T. et al., *Stat3 Activation Regulates the Expression of Matrix Metalloproteinase-2 and Tumor Invasion and Metastasis*, Oncogene 23: 3550-3560 (2004); Levy, D.E., Inghirami, G., *STAT3: A Multifaceted Oncogene*, PNAS 103: 10151-52 (2006).

[0175] STATs share structurally and functionally conserved domains including: an N-terminal domain that strengthens interactions between STAT dimers on adjacent DNA-binding sites; a coiled-coil STAT domain that is implicated in protein-protein interactions; a DNA-binding domain with an immunoglobulin-like fold similar to p53 tumor suppressor protein; an EF-hand-like linker domain connecting the DNA-binding and SH2 domains; an SH2 domain that acts as a phosphorylation-dependent switch to control receptor recognition and DNA-binding; and a C-terminal transactivation domain. Chen X., Vinkemeier U., Zhao Y., Jeruzalmi D., Darnell J.E., Kuriyan J., *Crystal Structure of a Tyrosine Phosphorylated STAT-1 Dimer Bound to DNA*, Cell 93: 827-839 (1998). In order to bind DNA, STAT proteins must dimerize. Yu, H., Jove, R., *The STATS of Cancer – New Molecular Targets Come of Age*, Nature Reviews, 4:97-105 (2004), Darnell, J. Jr., *Validating Stat3 in Cancer Therapy*, Nature Medicine, 11:595-96 (2005). Dimer formation involves reciprocal interaction between a SRC-homology 2 (SH2) domain of one STAT molecule with a phosphotyrosine residue of a second STAT molecule.

[0176] STAT proteins have been found to be overactive and/or persistently activate in cancer cells, including solid tumors and blood malignancies, proliferative diseases of the skin, and inflammatory diseases. Yu, H., Jove, R., *The STATS of Cancer – New Molecular Targets Come of Age*, Nature Reviews, 4:97-105 (2004). For example, STAT signaling has been implicated in various cancers. Song, J. I. and Grandis, J. R., *STAT Signaling in Head and Neck Cancer*, Oncogene, 19: 2489-2495, 2000; Nikitakis, N.G. et al., *Targeting the STAT Pathway in*

Head and Neck Cancer: Recent Advances and Future Prospects, Current Cancer Drug Targets, 4:639-651. An increased presence of activated STAT3 and/or STAT5 in the nucleus is associated with dysregulation of gene transcription. Yu, H., Jove, R., *The STATS of Cancer – New Molecular Targets Come of Age*, Nature Reviews, 4:97-105. Further, an increased presence of STAT3 and/or STAT5 is associated with increased transcription of genes which contribute to a cell proliferation, survival, angiogenesis, and tumor-induced immunotolerance. Yu, H., Jove, R., *The STATS of Cancer – New Molecular Targets Come of Age*, Nature Reviews, 4:97-105. For example, STAT5 is activated in certain leukemias, breast cancer, and head and neck cancer. Yu, H., Jove, R., *The STATS of Cancer – New Molecular Targets Come of Age*, Nature Reviews, 4:97-105. Similarly, STAT3 activation is known to abrogate growth factor dependence which contributes to certain carcinoma tumor growth. Kijima, T., Niwa, H., Steinman, R. A., Drenning, S. D., Gooding, W. E., Wentzel, A. L., Xi, S., and Grandis, J. R., *STAT3 Activation Abrogates Growth Factor Dependence And Contributes To Head And Neck Squamous Cell Carcinoma Tumor Growth In Vivo*, Cell Growth Differ, 13: 355-362, 2002. Activation of STAT3 is also reported to regulate survival in human non-small cell carcinoma cells. Song, L., Turkson, J., Karras, J. G., Jove, R., and Haura, E. B., *Activation Of Stat3 By Receptor Tyrosine Kinases And Cytokines Regulates Survival In Human Non-Small Cell Carcinoma Cells*, Oncogene, 22: 4150-4165, 2003. Accordingly, STAT3 and STAT5 are useful targets in treating STAT3- and STAT5-mediated disease.

[0177] Both STAT3- and STAT5-signaling pathways involve binding of cytokines or growth factors to cell-surface receptors, which leads to activation of cytoplasmic tyrosine kinases, such as the JAK family, which subsequently leads to phosphorylation of STAT monomers. Gadina, M., Hilton, D., Johnston, J. A., Morinobu, A., Lighvani, A., Zhou, Y. J., Visconti, R., O'Shea, J. J. *Signaling by Type I and Type II Cytokine Receptors: Ten Years After*, Curr. Opin. Immunol. 2001, 13: 363. STAT proteins are activated by phosphorylation causing them to dimerize and translocate to the nucleus, where they bind to specific promoter sequences in target genes. Horvath, C. M., *The Jak-STAT Pathway Stimulated by Interferon Gamma*, Science, STKE, 2004, 260: tr8.

[0178] As shown schematically in Figure 5, growth factor receptors, cytokine receptors and non-receptor tyrosine kinases have signaling pathways that converge on STAT3 and STAT5. For STAT3, these receptors and kinases include IL-2, IL-6, IL-7, IL-9, IL-10, IL-11, IL-15, IL-21, EGF, OSM, G-CSF, TPO, LIF and GH. For STAT5, the receptors and kinases include IL-2, IL-3, IL-5, IL-7, IL-9, IL-15, G-CSF, GM-CSF, EPO, TPO, GH, and PRL.

[0179] More specifically, STAT3 activation is known to be mediated by EGFR, EPO-R, and IL-6 R via c-Src or JAK2. See e.g., Lai, S. Y., Childs, E. E., Xi, S., Coppelli, F. M., Gooding, W. E., Wells, A., Ferris, R. L., and Grandis, J. R., *Erythropoietin-Mediated Activation of JAK-STAT Signaling Contributes to Cellular Invasion in Head and Neck Squamous Cell Carcinoma*, *Oncogene*, 24: 4442-4449, 2005; Siavash, H., Nikitakis, N. G., and Sauk, J. J., *Abrogation of IL-6-Mediated JAK Signalling by the Cyclopentenone Prostaglandin 15d-PGJ(2) in Oral Squamous Carcinoma Cells*, *Br J Cancer*, 91: 1074-1080, 2004; & Quadros, M. R., Peruzzi, F., Kari, C., and Rodeck, U., *Complex Regulation of Signal Transducers and Activators of Transcription 3 Activation in Normal and Malignant Keratinocytes*, *Cancer Res*, 64: 3934-3939, 2004. MAPK activation can lead to decreased STAT3 phosphorylation. In solid tumors, PDGFR and c-Met can also activate STAT3 via c-Src. IGFR1 and EGFR can activate STAT3 in a JAK-independent manner. STAT3 activation can lead to activation of several downstream target genes including Bcl-XL, cyclin D1 and VEGF.

[0180] Ligand binding to a cell-surface cytokine receptor triggers activation of JAKs. With increased kinase activity, JAK phosphorylates tyrosine residues on the receptor and creates sites for interaction with proteins that contain phosphotyrosine-binding SH2 domain. STATs possessing SH2 domains capable of binding these phosphotyrosine residues are recruited to the receptors and are tyrosine-phosphorylated by JAKs. The phosphotyrosine then acts as a docking site for SH2 domains of other STATs, mediating their dimerization. Different STATs form hetero- as well as homodimers. Activated STAT dimers accumulate in the cell nucleus and activate transcription of their target genes. Hebenstreit D. et al. (2005) *Drug News Perspect.* Vol. 18 (4), pages 243-249. STATs can be tyrosine-phosphorylated by other non-receptor tyrosine kinases, such as c-src, as well as receptor tyrosine kinases, such as the epidermal growth factor receptor.

[0181] Likewise, the binding of IL-6 family cytokines (including IL-6, oncostatin M and leukemia inhibitory factor) to the gp130 receptor triggers STAT3 phosphorylation by JAK2. Boulton, TG, Zhong, Z, Wen, Z, Darnell, Jr, JE, Stahl, N, and Yancopoulos, GD, *STAT3 Activation by Cytokines Utilizing gp130 and Related Transducers Involves a Secondary Modification Requiring an H7-Sensitive Kinase* *Proc Natl Acad Sci U S A.* 92(15): 6915-6919. EGF-R and certain other receptor tyrosine kinases, such as c-MET phosphorylate STAT3 in response to their ligands. Yuan ZL et al., (2004) *Mol. Cell Biol.* Vol. 24 (21), pages 9390-9400. STAT3 is also a target of the c-src non-receptor tyrosine kinase. Silva C.M. (2004) *Oncogene* Vol. 23 (48), pages 8017-8023. In addition to STAT3 activation through IL-6 binding, STAT3 is

further activated by binding of IL-2, IL-7, IL-9, IL-10, IL-11, IL-15, IL-21, EGF, OSM, G-CSF, TPO, LIF, or GH to an appropriate receptor. Similarly, STAT5 is activated by binding of IL-2, IL-3, IL-5, IL-7, IL-9, IL-15, G-CSF, GM-CSF, EPO, TPO, GH, or PRL to an appropriate receptor.

[0182] Indeed, Janus kinases (“JAK”) play an important role in the initial steps of cytokine receptor signaling. While the specificity of the four members of the JAK family (Jak1, Jak2, Jak3, and Tyk2) for different cytokine receptors is not fully understood, studies report that certain specific cytokine receptors can activate one or more Jak. O'shea, J. J., Pesu, M., Borie, D. C., Changelian, P. S., *A New Modality for Immunosuppression: Targeting the JAK/STAT Pathway*, Nature Rev. Drug Disc. 2004 (3): 555-564. For example, the growth hormone receptor (GHR) may interact predominantly with Jak2, and has been shown that to induce phosphorylation of Jak1 and Jak3. Hellgren, G., Jansson, J. O., Carlsson, L. M., Carlsson, B., *The Growth Hormone Receptor Associates with Jak1, Jak2 and Tyk2 in Human Liver*, Growth Horm. IGF Res. 1999 9(3):212-8.

[0183] The JAK-STAT pathway can be negatively regulated on multiple levels. Protein tyrosine phosphatases remove phosphates from cytokine receptors as well as activated STATs Hebenstreit D. et al. (2005) Drug News Perspect. Vol. 18 (4), pages 243-249. More recently, identified Suppressors of Cytokine Signaling (SOCS) inhibit STAT phosphorylation by binding and inhibiting JAKs or competing with STATs for phosphotyrosine binding sites on cytokine receptors. Krebs, L. et al. (2001) Stem Cells Vol. 19, pages 378-387. STATs are also negatively regulated by Protein Inhibitors of Activated STATs (PIAS), which act in the nucleus through several mechanisms. Shuai, K. (2006) Vol. 16 (2), pages 196-202. For example, PIAS1 and PIAS3 inhibit transcriptional activation by STAT1 and STAT3 respectively by binding and blocking access to the DNA sequences they recognize.

[0184] The JAK-STAT signaling pathway takes part in the regulation of cellular responses to cytokines and growth factors. Employing Janus kinases (JAKs) and Signal Transducers and Activators of Transcription (STATs), the pathway transduces the signal carried by these extracellular polypeptides to the cell nucleus, where activated STAT proteins modify gene expression. Although STATs were originally discovered as targets of Janus kinases, certain stimuli can activate them independent of JAKs. D W Leaman, S Pisharody, T W Flickinger, M A Commane, J Schlessinger, I M Kerr, D E Levy, and G R Stark *Roles of JAKs in Activation of STATs and Stimulation of c-fos Gene Expression by Epidermal Growth Factor*, Mol Cell Biol. 1996 16(1): 369–375. This pathway plays a central role in principal cell fate decisions,

regulating the processes of cell proliferation, differentiation and apoptosis. STAT activator phosphorylation requires phosphorylation of JAK to phosphorylate. In other words, JAK must first phosphorylate and bind to STAT. STAT is then able to phosphorylate and dimerize with another STAT.

[0185] The compounds described herein can control cell growth and survival via STAT modulation. Cell activity is regulated by “signals.” Signals enter cell walls through receptors, Signals are transduced by STATs. STATs complete a pathway to DNA. Signals activate DNA transcription (otherwise referred to as gene activation). BCL-XL, MCL1, Survivin and p53 are involved in cell survival. MYC, Cyclin D1/D2 and p53 are involved in cell proliferation. Downstream of the phosphorylation of STAT3, signaling pathways cause angiogenesis as a result of regulation of VEGF, HIF1 and p53, and immune evasion as regulated by immune-suppressing factors, pro-inflammatory cytokines and pro-inflammatory chemokines.

[0186] The compounds and methods described herein can directly or indirectly modulate STAT3 and/or STAT5 activation. For example, the compounds and methods can inhibit STAT3 activation by: (1) affecting signaling pathways which lead to STAT3 or STAT5 activation, such that the downstream activation of STAT3 or STAT5 is prevented or reduced; (2) directly preventing or reducing STAT3 or STAT5 activation, e.g., by preventing phosphorylation of STAT3 or STAT5; and (3) disrupting complexation of STAT3 or STAT5 with any number of cofactors and/or macromolecules, where the complexes are responsible for regulation of STAT3 or STAT5 signaling, and likewise STAT5 activation.

[0187] Inhibition of STAT3 and/or STAT5 activation can be accomplished, directly or indirectly, through any of the STAT targeting strategies currently known in the art. By way of example only, STAT3 activity might be decreased by: (1) reducing the availability of STAT3 or STAT5 molecules for recruitment; (2) blocking STAT3 or STAT5 recruitment to receptors, (3) blocking phosphorylation of STAT3 or STAT5 molecules; (4) interfering with the formation of STAT3 or STAT5 dimers; (5) interference with translocation of STAT3 or STAT5 dimers into the nucleus; (6) interference of STAT3 or STAT5 binding to DNA; (7) interference of STAT3 or STAT5 interaction with transcriptional activators; and/or (8) other methods of preventing STAT-mediated transcriptional activation. See e.g., N.G. Nikitakis, *et al.*, *Targeting the STAT Pathway in Head and Neck Cancer: Recent Advances and Future Prospects*, Current Cancer Drug Targets, 4: 637-651 (2004).

[0188] Direct or indirect modulation of STAT3 or STAT5 activity by the compounds presented herein can prevent, reduce, or eliminate STAT3- and/or STAT5-dependent activation

of genes responsible for proliferation, survival, metastasis, angiogenesis, immune response, and tumor immune evasion. As shown schematically, in Figures 5 and 6, STAT3 and/or STAT5 activation have an effect on the transcription of many genes including, but not limited to, VEGF, MMP9, MMP2, survivin, c-Myc, MMP-1, MEK-5, c-FOS, COX-2, Bcl-xL, MMP-10, HSP27, Jmjdla, PGE-2.

[0189] These activated STAT effects on the transcription of certain genes are likewise associated with physiological effects. For example, as reported by Yu and Jove at pages 99-101, which are incorporated by reference herein, activated STATs promote cell survival by up-regulating Bcl-xL, MCL1, and surviving, and down-regulating p53. Yu, H., et al., *The STATS of Cancer – New Molecular Targets Come of Age*, Nature Reviews, 4:97-105 (2004). Activated STATs also support proliferation by up-regulating MYC and cyclin D1/D2 and down-regulating p53. Additionally, activated STATs support angiogenesis through up-regulation of VEGF and HIF1, and down regulation of p53. Furthermore, activated STATs also aid tumor immune evasion by up-regulating immune-suppressing factors and down-regulating pro-inflammatory cytokines and chemokines.

[0190] Besides being useful for human treatment, the compounds and formulations of the present invention are also useful for veterinary treatment of companion animals, exotic animals and farm animals, including mammals, rodents, and the like. More preferred animals include horses, dogs, and cats.

GENERAL SYNTHETIC METHODS FOR PREPARING COMPOUNDS

[0191] The following schemes can be used to practice the present invention.

Synthesis and Characterization

Synthesis of (E)-3-(6-bromopyridin-2-yl)acrylaldehyde

Solution of 6-bromo-2-carboxypyridine (3g, 16.1 mmol) was dissolved in dichloromethane (100 mL). (Triphenylphosphoranylidene) acetaldehyde (4.9 g, 16.1 mmol) was added and obtained mixture was stirred at room temperature for 5 hrs. Dichloromethane was partially evaporated (to the volume about 50 mL) and reaction mixture was applied on chromatography column (SilicaGel 60) Products were eluted with dichloromethane. Fractions containing product were combined and evaporated to give 2.3 g of white powder (Yield 67%).


[0192] ¹HNMR (CDCl₃, δ) ppm: 9.81 (d, 1H, J = 7.6 Hz, CHO), 7.63 (dd, 1H, J = J = 7.7 Hz, H₄), 7.51 (dd, 2H, J = J = 7.7 Hz, H-3,5), 7.45 (d, 1H, J = 15,8 Hz, Ar-CH=CH-CHO), 7.12 (dd, 1H, 1H, J = 15.8 Hz, J = 7.6 Hz, Ar-CH=CH-CHO)

General procedure: Synthesis of (2E,4E)-5-(6-bromopyridin-2-yl)-2-cyano-N-substituted-2,4-dienamides -

Solution of (1) (2.1 g, 9.9 mmol) and piperidine (0.2 mL) in anhydrous ethanol was prepared. Aldehyde (10.6 mmol) was added and reaction mixture was stirred at room temperature. After 3 hrs obtained solid was filtered, washed with ethanol and dried. The compounds as illustrated in the following examples were made.

EXAMPLE 1


(2E,4E)-5-(6-bromopyridin-2-yl)-2-cyano-N-[(1S)-1-phenylethyl]penta-2,4-dienamide



[0193] Yield 67 % (2E,4E)-5-(6-bromopyridin-2-yl)-2-cyano-N-((S)-1-phenylethyl)penta-2,4-dienamide ¹HNMR (CDCl₃, δ) ppm: 8.07 (d, 1H, J = 12.1 Hz, H-3), 7.71 (dd, 1H, J = 12.1 Hz, J = 15.0 Hz, H-4), 7.59 (dd, 1H, J = J = 7.7 Hz, H-4'), 7.47 (d, 1H, J = 7.5 Hz, H-3'), 7.41 (d, 1H, J = 7.63 Hz, H-5'), 7.39 – 7.29(m, 5H, H arom), 7.15 (d, 1H, J = 15.0 Hz, H-5), 6.45 (d, 1 H, J = 7.5 Hz, NH), 5.23 (dt, 1 H, J = 7.0 Hz, J = 7.5 Hz, H-1''), 1.61 (d, 1 H, J = 7.0 Hz, Me).

EXAMPLE 2


(2E,4E)-5-(6-chloropyridin-2-yl)-2-cyano-N-[(1S)-1-phenylethyl]penta-2,4-dienamide



[0194] ¹HNMR (CDCl₃, δ) ppm: 8.07 (d, 1H, J = 12.1 Hz, H-3), 7.73 (dd, 1H, J = 12.0 Hz, J = 15.0 Hz, H-4), 7.70 (dd, 1H, J = J = 7.81 Hz, H-4'), 7.43 – 7.29(m, 7H, H-3', H-5', H arom), 7.18 (d, 1H, J = 15.0 Hz, H-5), 6.45 (d, 1H, J = 7.3 Hz, NH), 5.25 (dt, 1H, J = 7.0 Hz, J = 7.5 Hz, H-1''), 1.61 (d, 1 H, J = 7.0 Hz, Me).

EXAMPLE 3

(2E,4E)-5-(6-chloropyridin-2-yl)-2-cyano-N-[(1R)-1-phenylethyl]penta-2,4-dienamide




[0195] ¹HNMR (CDCl₃, δ) ppm: 8.07 (d, 1H, J = 12.1 Hz, H-3), 7.71 (dd, 1H, J = 12.1 Hz, J = 15.0 Hz, H-4), 7.59 (dd, 1H, J = J = 7.7 Hz, H-4'), 7.47 (d, 1H, J = 7.5 Hz, H-3'), 7.41

(d, 1H, $J = 7.63$ Hz, H-5'), 7.39 – 7.29(m, 5H, H arom), 7.15 (d, 1H, $J = 15.0$ Hz, H-5), 6.45 (d, 1H, $J = 7.5$ Hz, NH), 5.23 (dt, 1H, $J = 7.0$ Hz, $J = 7.5$ Hz, H-1''), 1.61 (d, 1H, $J = 7.0$ Hz, Me).

EXAMPLE 4


(2*E*,4*E*)-*N*-benzyl-5-(6-bromopyridin-2-yl)-2-cyanopenta-2,4-dienamide



[0196] ^1H NMR (CDCl_3 , δ) ppm: 8.10 (d, 1H, $J = 12.1$ Hz, H-3), 7.70 (dd, 1H, $J = 12.1$ Hz, $J = 15.0$ Hz, H-4), 7.57 (d, 1H, $J = 7.7$ Hz, H-4'), 7.46 – 7.29 (m, 7H, H-3', H-5', H arom), 7.16 (d, 1H, $J = 15.0$ Hz, H-5), 6.52 (bs, 1H, NH), 4.58 (d, 1H, $J = 5.8$ Hz, 1'').

EXAMPLE 5


(2*E*,4*E*)-5-(6-chloropyridin-2-yl)-2-cyano-*N*-[(1*R*)-1-phenylethyl]penta-2,4-dienamide



[0197] ^1H NMR (CDCl_3 , δ) ppm: 8.07 (d, 1H, $J = 12.1$ Hz, H-3), 7.73 (dd, 1H, $J = 12.0$ Hz, $J = 15.0$ Hz, H-4), 7.70 (dd, 1H, $J = J = 7.81$ Hz, H-4'), 7.43 – 7.29(m, 7H, H-3', H-5', H arom), 7.18 (d, 1H, $J = 15.0$ Hz, H-5), 6.45 (d, 1H, $J = 7.3$ Hz, NH), 5.25 (dt, 1H, $J = 7.0$ Hz, $J = 7.5$ Hz, H-1''), 1.61 (d, 1H, $J = 7.0$ Hz, Me).

EXAMPLE 6


(2*E*,4*E*)-*N*-benzyl-5-(6-chloropyridin-2-yl)-2-cyanopenta-2,4-dienamide



[0198] ^1H NMR (CDCl_3 , δ) ppm: 8.09 (d, 1H, $J = 12.1$ Hz, H-3), 7.72 (dd, 1H, $J = 12.1$ Hz, $J = 15.0$ Hz, H-4), 7.68 (dd, 1H, $J = J = 7.8$ Hz, H-4'), 7.40 – 7.28 (m, 7H, H-3', H-5', H arom), 7.17 (d, 1H, $J = 15.0$ Hz, H-5), 6.45 (bs, 1H, NH), 4.58 (d, 1H, $J = 5.75$ Hz, H-1'').

EXAMPLE 7


(2*E*,4*E*)-5-(6-bromopyridin-2-yl)-2-cyano-*N*-[(*S*)-cyclopropyl(phenyl)methyl]penta-2,4-dienamide



[0199] ^1H NMR (CDCl_3 , δ) ppm: 8.06 (d, 1H, $J = 12.1$ Hz, H-3), 7.72 (dd, 1H, $J = 12.1$ Hz, $J = 15.0$ Hz, H-4), 7.59 (dd, 1H, $J = 7.9$ Hz, H-4'), 7.47 – 7.29 (m, 7H, H-3', H-5', H arom), 7.15 (d, 1H, $J = 15.0$ Hz, H-5), 6.68 (d, 1H, $J = 7.8$ Hz, NH), 4.49 (dd, 1H, $J = 8.0$ Hz, $J = 8.8$ Hz, H-1''), 1.33-1.24 (m, 1 H, H-2''), 0.69 – 0.41 (m, 4H, 3''-CH₂, 4''-CH₂).

EXAMPLE 8


(2*E*,4*E*)-5-(6-bromopyridin-2-yl)-2-cyano-*N*-[(1*S*)-1-phenylpropyl]penta-2,4-dienamide



[0200] ^1H NMR (CDCl_3 , δ) ppm: 8.03 (dd, 1H, $J = 12.1$ Hz, $J = 0.7$ Hz, H-3'), 7.69 (dd, 1H, $J = 12.1$ Hz, $J = 15.0$ Hz, H-4), 7.57 (dd, 1H, $J = 7.5$ Hz, $J = 7.9$ Hz, H-4'), 7.45 (dd, 1H, $J = 7.9$ Hz, $J = 0.8$ Hz, H-3), 7.39 (dd, 1H, $J = 7.5$ Hz, $J = 0.8$ Hz, H-5), 7.39 – 7.27 (m, 5H, H arom), 7.1 (d, 1H, $J = 15.0$ Hz, H-5), 6.43 (d, 1H, $J = 7.8$ Hz, NH), 4.96 (dd, 1H, $J = 15.2$ Hz, $J = 7.5$ Hz, H-1''), 1.99 – 1.85 (m, 2H, H-2''), 0.93 (t, 3H, $J = 7.4$ Hz, H-3'').

EXAMPLE 9

(2*E*,4*E*)-5-(6-bromopyridin-2-yl)-2-cyano-*N*-[(1*S*)-1-phenylbutyl]penta-2,4-dienamide



[0201] ^1H NMR (CDCl_3 , δ) ppm: 8.03 (dd, 1H, $J = 12.1$ Hz, $J = 0.7$ Hz, H-3), 7.68 (dd, 1H, $J = 15.0$ Hz, $J = 12.1$ Hz, H-4), 7.55 (dd, 1H, $J = 7.7$ Hz, H-4'), 7.45 (dd, 1H, $J = 7.9$ Hz, $J = 0.8$ Hz, H-3'), 7.39 (dd, 1H, $J = 7.5$ Hz, $J = 0.8$ Hz, H-5'), 6.66 – 7.28 (m, 6H, H-4, H arom), 7.13 (d, 1H, $J = 14.9$ Hz, H-5), 6.42 (d, 1H, $J = 8.0$ Hz, NH), 5.05 (dd, 1H, $J = 15.3$, $J = 7.6$ Hz, H-1''), 1.91 – 1.82 (m, 2H, H-2''), 1.41 – 1.24 (m, 2H, H-3''), 0.95 (t, 3H, $J = 7.3$ Hz, H-4'').

[0202] The antiproliferative activity of the Compound of Example 1 and Example 7 have been shown in a Colo357FG pancreatic cancer cell line as follows:

	Colo357FG
Compound	IC50 μ M
Example 1	3.2
Example 7	5.4

[0203] The activity of the compounds as STAT inhibitors in Examples 1, 2, 4, and 9 has been shown by at least one of the following assays. The associated activity data obtained is provided in Figures 1 through 4. The other compounds described above, which have not yet been made, are predicted to have activity in these assays as well.

Biological Activity Assay

MTT Method for Determining Cell Viability and Drug Dose-Response

[0204] A variety of cell types were seeded at a density of 50,000 cells/100 μ l per well onto 96-well plates and cultured for 24–48 h in DMEM medium until they reached confluency. At confluency, 25 μ l of fresh media containing WP1220 at various concentrations (0 to 100 μ M) was added to the wells and incubated for 72 h at 37 °C. The complex was removed, 100 μ l of fresh medium was added and the cells were incubated overnight. Medium was removed and fresh medium (100 μ l) was added to each well along with (20 μ l) MTT solution (5 mg/ml in PBS) before incubating the cells for a further 4 h at 37 °C. The reaction product was solubilised with sodium dodecyl sulfate (100 μ l, 10%, w/v) in 0.01 N HCl overnight before quantifying the product using a microplate reader at 590 nm and comparing with control cells. The experiment was carried out in triplicate (n=3).

Biological Activity Assay

Western Blot Methods for Determining Protein and Phosphoprotein Content

[0205] Using phospho-tyrosine STAT3 (Y705), phospho-tyrosine Jak2 (Y1007/1008), total STAT3, total Jak2, Bcl-xL, actin (Cell Signaling Technology, Danvers, MA), and survivin (R&D Systems, Minneapolis, MN) antibodies, western blots were performed using the following methods. Briefly, (1) Samples were prepared from cells that were homogenized in a buffer protecting the protein of interest from degradation; (2) The sample was then separated using SDS-PAGE (10-15 μ g protein/well) and then transferred to a membrane for detection; and (3) The membrane was incubated with a generic protein (milk proteins) to bind to any remaining sites on the membrane. A primary antibody was then added to the solution which is able to bind


to its specific protein; (4) A secondary antibody-enzyme conjugate, which recognizes the primary antibody was then added to find locations where the primary antibody has bound.

[0206] From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

CLAIMS

What is claimed is:

1. A compound of structural Formula I:




or a pharmaceutically acceptable salt or solvate thereof, wherein:

n is an integer selected from 1, 2 or 3;

R_0 is R_1 , or R_0 is R_1-Z_1 — wherein Z_1 is alkyl, and specifically may be a lower alkyl such as $-(CH_2)_{m_3}-$ where $m_3=0, 1, 2, 3$, or 4;

R_1 is:




wherein X_1 , X_2 , X_3 , and X_4 , are each independently hydrogen, halogen, alkyl,

alkoxy, OH, trihalomethyl, or NO₂, and Y₁ is OH, halogen, specifically including Br and Cl, or O₂N;

R₂ is alkyl, alkenyl, alkynyl, alkoxy, alkylaryl, halogen, hydrogen, OH, NO₂, thioether, amine, SH, or NH₂;

R₃ is:



wherein m₁ is an integer selected from 1, 2, 3, or 4,


X₅ and X₆ are each independently hydrogen, alkyl, upper alkyl, lower alkyl, aryl, alkoxy, aryloxy, cyclic alkyl, cycloalkyl, cycloarylalkyl, aralkyl, alkylester, alkylesteralkyl, alkylacetoxyl, hydroxyl, hydroxylalkyl, cyclopropyl, cyclobutyl, —CH₃, —CH₂OH, cyclopentyl, —CH₂OAc, —CH₂OC(O)C(CH₃)₃, —CH₂C₆H₅, or cyclohexyl,


Z is NH, S, or O, and

Z₃ is alkyl or lower alkyl;

R₄ is CN, substituted amine, CH₂S-alkyl, alkyl, or CH₂N₃; and

R₅ and R₆ are each independently selected from the group consisting of:





monosaccharide (e.g., glucose, fructose, galactose, etc.), polysaccharide, monosaccharide derivative (e.g., an acetylated monosaccharide such as acetylated galactose, 1,2,3,4-diisopropylideno-D-galactose) substituted and unsubstituted aryl, and substituted and unsubstituted alkylaryl.

2. A method of inhibiting STAT3 activation comprising contacting a cancer cell with a compound as recited in Claim 1 (or of Formula I, and define).

3. A method of treatment of a STAT3-mediated disease comprising the administration of a therapeutically effective amount of a compound as recited in Claim 1 (or of Formula I, and define) to a patient in need thereof.

4. A method of inhibiting STAT5 activation comprising contacting a cancer cell with a compound as recited in Claim 1 (or of Formula I, and define).

5. A method of treatment of a STAT5-mediated disease comprising the administration of a therapeutically effective amount of a compound as recited in Claim 1 (or of Formula I, and define) to a patient in need thereof.