


CLAIMS

1. A compound of formula (I),



formula (I),

- 5 or a pharmaceutically acceptable salt thereof, for use in the treatment, prevention, or alleviation of a combination of:
erectile dysfunction and depression,
erectile dysfunction and sexual dysfunction,
erectile dysfunction and anxiety,
10 erectile dysfunction and ejaculation disorder,
sexual dysfunction and depression,
sexual dysfunction and anxiety,
sexual dysfunction and ejaculation disorder,
depression and anxiety,
15 depression and ejaculation disorder, and/or
anxiety and ejaculation disorder,
in a subject.
- 20 2. The compound for use according to any one of the preceding claims, wherein the treatment, prevention, and/or alleviation is the treatment, prevention, and/or alleviation of a combination of erectile dysfunction and depression.
- 25 3. The compound for use according to any one of the preceding claims, wherein the treatment, prevention, and/or alleviation is the treatment, prevention, and/or alleviation of a combination of sexual dysfunction and anxiety.
4. The compound for use according to any one of the preceding claims, wherein the ejaculation disorder is premature ejaculation or ejaculatory disorder.
- 30 5. The compound for use according to any one of the preceding claims, wherein the alleviation of anxiety associated with ejaculatory disorder.


6. The compound for use according to any one of the preceding claims, wherein the erectile dysfunction is treatment-emergent erectile dysfunction.
- 5 7. The compound for use according to any one of the preceding claims, wherein the sexual dysfunction is caused by depression and/or anxiety.
8. The compound for use according to any one of the preceding claims, wherein the erectile dysfunction is caused by depression.
- 10 9. The compound for use according to any one of the preceding claims, or a pharmaceutically acceptable salt thereof, wherein the erectile dysfunction is treatment-emergent erectile dysfunction.
- 15 10. The compound for use according to any one of the preceding claims, or a pharmaceutically acceptable salt thereof, wherein the treatment-emergent erectile dysfunction is an adverse effect originating from treatment of a medicament.
- 20 11. The compound for use according to any one of the preceding claims, wherein the medicament is selected from the group consisting of antidepressants, NSAIDs, finasteride, antiepileptics and neuroleptics.
- 25 12. The compound for use according to any one of the preceding claims, wherein the erectile dysfunction is treatment-emergent erectile dysfunction caused by the treatment by and antidepressant medicament.
- 30 13. The compound for use according to any one of the preceding claims, wherein the erectile dysfunction is treatment-emergent erectile dysfunction caused by the treatment of depression by and antidepressant medicament.
14. The compound for use according to any one of the preceding claims, wherein the depression is caused by sexual dysfunction.

15. The compound for use according to any one of the preceding claims, wherein the depression is caused by erectile dysfunction.
- 5 16. A pharmaceutical composition, comprising a therapeutically effective amount of the compound according to any one of the preceding claims together with at least one pharmaceutically acceptable carrier, excipient or diluent.
- 10 17. The compound according to any one of the preceding claims, wherein the subject is a mammal.
18. The compound according to any one of the preceding claims, wherein the mammal is a human.
- 15 19. The compound according to any one of the preceding claims, wherein the human is male.
20. The compound for use according to any one of the preceding claims, wherein the subject is male.
- 20 21. The compound for use according to any one of the preceding claims, wherein the subject is a young male.
- 25 22. The compound for use according to any one of the preceding claims, wherein the subject is a male below 40 years of age, such as below 35 years of age, such as below 30 years of age, such as below 25 years of age, such as below 20 years of age.
- 30 23. The compound for use according to any one of the preceding claims, wherein the subject is an adult male, such as an adult male over the age of 20, such as an adult male over the age of 25, such as an adult male over the age of 30, such as an adult male over the age of 35, such as an adult male over the age of 40, such as an adult male over the age of 45, such as an adult male over the age of 50, such as an adult male over the age of 55, such as an adult male over the age of 60, such as an adult male over the age of 65, such as an adult male

over the age of 70, such as an adult male over the age of 75.

- 5 24. The compound for use according to any one claims 1 to 18, wherein the treatment, prevention, and/or alleviation, is treatment, prevention, and/or alleviation of sexual dysfunction, depression, and/or anxiety; or any combination thereof; in a female subject.
- 10 25. The compound for use according to any one of claims 1 to 18, wherein the treatment, prevention, and/or alleviation, is treatment, prevention, and/or alleviation of a combination of sexual dysfunction and depression in a female subject.
- 15 26. The compound for use according to any one of the preceding claims, wherein the subject is not simultaneously treated with other anxiolytic and/or antidepressive medication.
27. The compound for use according to any one of the preceding claims, wherein the compound is administered systemically.
- 20 28. The compound for use according to claim 27, wherein the compound is administered orally.
29. The compound for use according to any one of claims 1 to 26, wherein the compound is administered locally.
- 25 30. The compound for use according to claim 29, wherein the compound is administered topically.
31. The compound for use according to claim 30, wherein the topical administration is in the form of a lotion, a cream, an ointment, a gel, or by a transdermal patch.
- 30 32. A method for treatment, prevention, or alleviation of a combination of: erectile dysfunction and depression,


erectile dysfunction and sexual dysfunction,
erectile dysfunction and anxiety,
erectile dysfunction and ejaculation disorder,
sexual dysfunction and depression,
5 sexual dysfunction and anxiety,
sexual dysfunction and ejaculation disorder,
depression and anxiety,
depression and ejaculation disorder, and/or
anxiety and ejaculation disorder,
10 said method comprising administering to a subject in need thereof a
therapeutically effective amount of the compound of formula (I),



formula (I),

or a pharmaceutically acceptable salt thereof.

- 15 33. A method for increasing the magnitude of erectile response in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I),



formula (I),


or a pharmaceutically acceptable salt thereof.

20

34. The method according to claim 33, wherein the subject is a young male.

35. The method according to any one of the preceding claims, wherein the young male is below 40 years old, such as below 35 years old, such as below 30
25 years old, such as below 25 years old, such as below 20 years old.

36. A method for increasing the duration of an erectile response in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I),




formula (I),

or a pharmaceutically acceptable salt thereof.

5

37. A method for increasing the frequency of erectile responses in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I),




formula (I),

or a pharmaceutically acceptable salt thereof.

10

38. A method for inducing relaxation of the corpus cavernosum in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I),




formula (I),

or a pharmaceutically acceptable salt thereof.

15

39. A method for increasing penile blood flow in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I),



formula (I),

20

or a pharmaceutically acceptable salt thereof.

25

40. The method according to any one of the preceding claims, wherein the subject is an adult male, such as an adult male over the age of 20, such as an adult male over the age of 25, such as an adult male over the age of 30, such as an adult male over the age of 35, such as an adult male over the age of 40, such

as an adult male over the age of 45, such as an adult male over the age of 50,
such as an adult male over the age of 50, such as an adult male over the age of
55, such as an adult male over the age of 60, such as an adult male over the
age of 65, such as an adult male over the age of 70, such as an adult male over
5 the age of 75.

41. The method according to any one of the preceding claims, wherein the subject
is suffering from depression and/or anxiety.
- 10 42. The method according to any one of the preceding claims, wherein the subject
is suffering from ejaculatory disorder.