

MAN-PLANTED FOREST, METHOD OF PLANTING TREES AND METHOD OF GROWING PLANTS

BACKGROUND OF THE INVENTION

The present invention relates to a plant growing technique such as planting trees, and more specifically, to a technique of selectively growing young trees or the like which is resistant to ultraviolet rays so that vegetation such as a forest which is resistant to ultraviolet rays can be reliably obtained.

There has been discussed recently on the issue of ultraviolet rays problem on earth. Because especially the ozonosphere of stratosphere surrounding the earth is recently destroyed, the ultraviolet rays come to the earth from outside and has been confirmed to show bad influence on biology on the earth.

For example, as to human body, there has been reported the increment of damage due to ultraviolet rays, and it has been confirmed that the increase of skin cancer cases are due to increment of radiation by ultraviolet rays. Thus damage based on ultraviolet rays was thought to cause the occurrence of active oxygen. Niwa, one of the present inventors, has studied their pharmacological, biochemical properties on several damages caused by active oxygen for long years.

SUMMARY OF THE INVENTION

The real states that damages based on conventional ultraviolet rays are mainly by studying on their effects on human body as the main topics. In contrast to this, the present inventors has reached to the conclusion that ultraviolet rays have also given big damages on forest.

It is disclosed that the biggest reason that damage forwarded to the

forest is acid rain's exertion. For example, at Schwarzwald district in Germany, a report dealt with big damage covered large area where so many trees uprightly withered by acid rain. The acidic oxides such as sulfur-containing oxides existed in the industrial burning, were dissolved in the rain drops and resulting become acidic rain.

The investigation on the damage was performed repeatedly by the present inventors who visited the damaged domestic forest and obtained many results.

The present inventors observed the damage of the domestic forest by repeatedly visiting several places of the damaged forest in several districts. After a detailed investigation about the observed states, the present inventors investigated that it is possible that the factors other than acid rain strongly concerned with abnormal phenomena of withered trees found largely in the forest found. The damages may possibly be due to ultraviolet rays found as the main factor in relating to forest damage to our further research result.

There is not so far examined before on the effect of ultraviolet rays exerting the forest damage. As the present inventors viewed the forest damage state due to ultraviolet rays, it is necessary to take a practical measure rapidly to inhibit such forest damage. Because enlarging the depletion of ozonosphere and the ultraviolet radiation increasing of world scale, it is a top urgent action to be taken for inhibiting the damage caused by ultraviolet rays during the several factors exerting to forest damage. If the damages were dealt with no policy, in the future, the forest trees withered uprightly worldwide by the ultraviolet rays. And the prophecy is forest will be disappeared with global scale.

The object of the present invention is to suppress the damage on forest

caused by ultraviolet rays.

The man-planted forest of the present invention is characterized in that it is constituted by planting seedling having multiploid (TASUSEI in Japanese). The chromosome number of the seedlings is triploid or tetraploid. The seedlings are those of a conifer. The seedlings of a conifer are seedlings selected from the group consisting of Japanese cedar and Japanese cypress. The chromosome number of the seedling is obtained by a means of increasing the number of chromosomes such as breeding and mutation.

The method of planting trees of the present invention is characterized in that it comprises the steps of : selecting seedlings having multiploid chromosome; and planting the selected seedlings. The chromosome number of the seedlings is triploid or tetraploid. The seedlings are those of a conifer. The seedlings of a conifer are seedlings of Japanese cedar or Japanese cypress. The chromosome number of the seedlings is obtained by means of increasing the number of chromosomes such as breeding and mutation. The method of growing plants of the present invention is characterized in that it comprises the step of artificially forming vegetation which is significantly resistant to ultraviolet rays, as compared with vegetation constituted of diploid plants, by selectively growing plants whose chromosome number exceeds the basic number intrinsic of the species.

In this specification, "multiploid"(TASUSEI in Japanese) means those chromosomes with more than 3-fold of fundamental chromosome by multiplying integral number of chromosome, and those chromosomes possessing more or less , by 1 to few, than the basic number is acquired by the species. Furthermore it should be noted that haploid, or diploid which is normally observed in plants, is not included into the scope defined by the

aforementioned "multiploid".

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The following describes the examples state of performance and details.

The seedling used in man-planted forest, method of planting trees and method of growing plants in this invention is ultra-violet-resistant (including "UV-resistant" abbreviation afterwards). The afforestation by choosing UV-resistant seedling and a man-planted forest with capacity of possessed resistant property against ultra-violet rays.

Upon selection of seedling possessing UV-resistant property, a standard of multiploid (TASUSEI) chromosomes acquired by the seedling is adopted for selection. The multiploid stated here means the case that the seedling having more than triploid chromosomes.

Such standard is suggested for the first time by the present inventors. Trees having multiploid chromosomes possess their resistance against ultraviolet rays than the diploid trees normally observed. In the man-planted forest formed by afforesting UV-resistant seedling, the occurrences of damage from ultraviolet rays observed in the trees having diploid chromosomes will be inhibited effectively. The abnormal phenomena such as tree withering uprightly, dying of leaves of evergreens, deciduous trees their leaves fall at the season of not falling. This abnormal phenomena is thought to be caused by ultraviolet rays, and this can be inhibited.

For description of several kinds of tree and ascertain the efficacy of this invention in the examples, the following article and discussion are given in detail.

(Example)

In the Example, those leaves of plants described in items (1) - (3) below,

were collected and subjected to experiment about ultraviolet resistance (UV-resistance).

(1) Rice

For a comparison of samples, the rice leaf was used. Rice, a monocotyledonous plant is well known for its resistance against ultraviolet rays. The rice leaf samples used in the experiment were grown at Tosashimizu city, Kochi Prefecture, Shikoku, Japan . The rice leaf were collected from five places at Tosashimizu city and subjected to measurement of enzyme activity, etc. by the method described later.

(2) Osmunda

Leaves of osmunda which is spontaneously growing in Tosashimizu, Japan , were collected as the object of experiment. As well known that osmunda, a fern, its plant evolutionary is old than other trees. Many reports dealt with the cases that osmunda possesses the adaptability to the environmental conditions by acquiring the multiplying ploid of chromosome of the species.

(3) Gymnosperms such as black pines etc.

As a sensitive plant to ultraviolet rays, the leaves of gymnosperms were included into the objects of experiment. In Tables 1-4 of this Example, KUROMATSU tree (English name: Japanese black pine tree), SUGI tree (English name: Japanese cedar) and HINOKI tree (English name: Japanese cypress) spontaneously growing in mountain of Japan, their leaves were adopted for the object of experiments. These leaf samples of black pine, cedar and cypress trees were collected from the places and numerated as shown in Tables 1-4.

Black pine tree (hereinafter abbreviated as "control black pine ") is known to be sensitive to common UV rays and black pines resistant to UV

rays (hereinafter abbreviated as "UV-resistant black pine) were separately described in Table 1, assays for SOD activity, GSH-Px activity and lipid peroxide levels were performed.

Furthermore, as demonstrated in Tables 5 and 6, measurements for triploid species and tetraploid species resulting from crossing and mutation were also performed.

[Table 1] Comparison of lipid peroxide levels, SOD and glutathione peroxide (GSH-Px) activities between UV resistant Japanese black pines and UV sensitive Japanese black pines.

	Sample name	SOD activity (unit/gr)	GSH-Px activity (unit/gr)	Lipid peroxide Levels (n mol/gr)
UV resistant Japanese black Pines	Nakamikata -73	391±46	0.0215±0.0032	18.0±2.1
	Tosashimizu -63	225±29	0.0171±0.0021	16.7±2.0
	Ooita -8	201±28	0.0269±0.0029	18.2±2.3
	Ei -425	358±46	0.0190±0.0022	20.7±2.8
	Tsuyasaki-50	461±73	0.0236±0.0035	22.2±3.3
	Misaki -90	346±41	0.0230±0.0036	23.4±3.0
	Namikata- 73	456±59	0.0331±0.0052	22.0±3.7
	Tanabe -54	367±44	0.0214±0.0027	20.5±2.8
	Yasu -37	195±21	0.0236±0.0033	18.4±2.9
	Yoshida -2	817±106	0.0246±0.0039	23.5±2.8
	Sendai -290	245±31	0.0233±0.0039	19.9±2.7
	Mitoyo -103	896±107	0.0276±0.0030	24.1±3.8
	Average	423±56*	0.0237±0.0035	20.6±3.6
Control UV sensitive Japanese black pines	No.1	252±30	0.0177±0.0023	20.5±3.2
	No.2	99±11	0.0142±0.0018	22.9±4.1
	No.3	141±18	0.0198±0.0031	21.1±2.9
	No.4	356±46	0.0206±0.0037	21.9±3.5
	No.5	352±42	0.0198±0.0017	18.7±2.4
	Average	240±37*	0.00184±0.0031	21.0±3.8

*P<0.01 between UV resistant and sensitive black pine trees.

[Table 2] Comparison of lipid peroxide levels, SOD and GSH-Px activities in Japanese cypress triploid

		Sample name	Poly ploidy	SOD activity (unit/gr)	GSH-Px activity (unit/gr)	Lipid peroxide Levels (n mol/gr)
Japanese cypress	Control diploids	B-4	2X	472±56	0.0180±0.0023	19.9±2.3
		B-11	2X	614±73	0.0155±0.0020	21.3±2.9
		B-13	2X	552±71	0.0019±0.0003	27.5±3.0
		D-4	2X	874±104	0.0107±0.0016	26.4±3.9
		D-9	2X	549±65	0.0028±0.0003	25.7±3.5
		D-12	2X	683±88	-	28.8±3.7
		D-14	2X	466±60	0.0077±0.0012	22.9±3.6
		Average		601±46 †	0.0094±0.0002	24.6±4.1
	Artificial Triploids	A-8	3X	479±71	0.0123±0.0017	24.7±3.4
		A-13	3X	543±70	0.0119±0.0015	27.8±3.6
		A-24	3X	685±89	0.0014±0.0002	28.1±4.4
		A-54	3X	505±70	0.0058±0.0008	22.7±3.1
		A-58	3X	558±55	0.0079±0.0012	21.1±2.3
		A-72	3X	674±74	-	19.1±2.4
		A-77	3X	482±77	0.0065±0.0011	20.8±3.3
		C-6	3X	559±61	0.0100±0.0012	21.5±2.5
		C-22	3X	548±65	0.0126±0.0017	19.5±2.3
		C-37	3X	580±69	0.0055±0.0008	22.2±2.8
		C-50	3X	589±82	-	21.1±3.1
		C-55	3X	577±75	-	20.9±2.7
		C-60	3X	624±74	-	19.4±2.3
		C-140	3X	646±83	0.0026±0.0004	19.9±2.7
		C-225	3X	1005±120		21.1±2.1
		Average		606±51 †	0.0076±0.0018	22.0±3.9

[Table 3] Comparison of lipid peroxide levels, SOD and GSH-Px activities in plus trees of Japanese cypress

		Sample Name	Poly-ploidy	SOD activity (unit/gr)	GSH-Px (unit/gr)	Lipid Peroxide Levels (n mol/gr)
Japanese Cypress (Plus Trees)	Diploids	Aso-7	2X	459±55	0.0069±0.0009	24.4±2.9
		Saeki-17	2X	423±54	0.0160±0.0020	21.2±2.7
		Hiji-4	2X	544±76	0.0019±0.0002	23.5±2.5
		Kusu-6	2X	459±59	0.0171±0.0027	20.7±2.4
		Taketa-8	2X	655±85	0.0068±0.0009	24.0±2.6
		Average		508 ± 78 †	0.0097±0.0018	22.7±4.3
	Triploids	Misoshi-4	3X	420±46	0.0136±0.0020	23.1±2.3
		Fuji-2	3X	582±75	0.0106±0.0012	14.1±1.5
		Average		501 ± 76 †	0.0121±0.0026	18.6±4.1
	Tetraploid	Sanko-Hinoki	4X	1902±190	0.0103±0.0012	21.5±3.4
		Kiso-4X	4X	905±99	0.0058±0.0007	16.2±2.5
		Average		1403 ± 212 †	0.0081±0.0013	18.8±2.8

† P<0.001 (between tetraploids of plus trees of Japanese and diploids or triploids of Japanese cypress, between tetraploids of plus trees of Japanese and diploids or triploids of plus trees of Japanese cypress)

[Table 4] Comparison of lipid peroxide levels, SOD and GSH-Px activities in plus trees of Japanese cedar, rice leaves and osmund

		Sample Name	Poly-ploidy	SOD activity (unit/gr)	GSH-Px activity (unit/gr)	Lipid Peroxide Levels (n mol/gr)
Japanese Cedar (Plus Trees)	Diploids	Naga-9	2X	12±1.4	0.0134±0.0021	4.6±0.5
		Akita-1	2X	37±4.4	0.0096±0.0015	5.0±0.6
		Saga-3	2X	21±2.5	0.0114±0.0018	3.1±0.5
		Saeki-10	2X	52±6.7	0.0104±0.0015	4.1±0.5
		Kunisaki-3	2X	26±2.8	0.0104±0.0017	4.5±0.7
		Nishikawa-2	2X	17±2.2	0.0093±0.0011	5.0±0.8
		Average		27 ± 3.8 §	0.0108±0.0019	4.3±0.6
	Triploids	Ooi-5	3X	161±19	0.0141±0.0019	7.0±0.8
		Kuji-30	3X	41±5.3	0.0092±0.0014	3.7 ± 0.4
		Onta-2	3X	317±44	0.0150±0.0024	9.5 ± 1.1
		Kyurin-3X	3X	108±11	0.0182±0.0023	8.7 ± 1.1
		Fujitsu-28	3X	236±28	0.0158±0.0022	8.2 ± 1.2
		Miyoshi-10	3X	95±12	0.0109±0.0011	6.1 ± 0.8
		Kuga-1	3X	59±8.2	0.0115±0.0013	8.0 ± 1.2
		Average		145±23 ‡	0.0135±0.0021	7.3 ± 1.2
	Tetraploid	Kanagawa-sugi	4X	3589±430	0.0176±0.0017	8.2 ± 0.9
		Cr-38	4X	2684±295	0.0128±0.0020	9.3 ± 1.2
		Average		3136±40 ‡ §	0.0152±0.0025	8.7 ± 1.3
rice leaves		Tosashimizu		2941±352	1.3841±0.1522	71 ± 8.5
Osmund		Tosashimizu		1394±181	0.0889±0.0115	51 ± 8.1

‡ P<0.0001 between tetraploid and triploids of plus trees of Japanese cedar.

§ P<0.00001 between tetraploids and diploids of plus trees of Japanese cedar.

As shown in Table 1, ultraviolet-resistant (UV-resistant) Japanese black pines was compared with ultraviolet-sensitive Japanese black pines (as control) on SOD activity, glutathione peroxidase (GSH-Px) activity and lipid peroxide levels.

In Table 2, diploids of Japanese cypress were compared with triploids with lipid peroxide levels, SOD activity and GSH-Px activity.

In Table 3, diploids, triploids and tetraploids of plus trees of Japanese cypress were compared with individual SOD activity, GSH-Px activity and lipid peroxide levels.

In Table 4, comparisons of SOD activity, GSH-Px activity, lipid peroxide level for diploids, triploids and tetra-ploids of plus trees of Japanese cedar, rice leaves and osmund. The comparison of Table 4 with Tables 5 and 6 which were described later.

The extracts from the leaves of diploid, triploid and tetraploid of Japanese cedar and Japanese cypress, were compared in Table 5 and 6 as described later. As shown in Table 5, in the column of sample added, for a mark used to description, the sample size with 0.6 gr/ml is termed as "x1" and thus 3.0 gr/ml "x2", 6.0 gr/ml "x10". In Table 6, the data of rice leaves are included for comparison.

The individual enzyme activity and lipid peroxide level for the plants described in items (1)-(3) were assayed. Testing items for measurement of enzyme activity are: SOD activity, glutathione peroxidase (GSH-Px) activity and catalase activity. The results obtained are shown in Tables 1- 4. The figure is shown by mean $\pm$ SD style. On the other hand, catalase activity for those plants other than rice leaves was utterly not detectable and their results were not shown in Tables 1-4.

Regarding to the quantitative analysis of lipid peroxide level, its level

was measured by using TBA reaction method of converting docohexanoic acid into TBA reactive substance.

The leaf homogenates from individual plants described above were prepared as follows and used as samples. Leaves collected were from the plant in (1)-(3). The collected leaves were powdered with a mill. The leaf homogenates from individual plants described above were used as samples. The procedure for preparation for these leaf homogenates are as follows. This powder was suspended in 95% ethanol solution for lipid peroxide formation test or in physiological saline for those tests other than lipid peroxide formation test. Both cases were at concentration of 60 mg/ml. The suspension solution was exposed to sonication (at 15W for 15 second).

After sonication, the solution was centrifuged (at 1300g). procedures as shown in Figure 1, the supernatants after centrifugation were divided into two parts. One part is for the preparations of sample for lipid peroxide level determination. The other one was filtrated through 45 μ m filter. One portion of the filtrate was used as sample for SOD activity assay and the remained filtrates were used as sample for catalase, GSH-Px activities assays.

(SOD 's quantitative analysis)

To add the mixture prepared as described above to 0.2 ml of the assay mixture containing xanthine-xanthine oxidase producing O_2^- , and the resulting O_2^- quantity was determined from reduction quantity of ferri-cytochrome c (type III) by the optical absorbance at 550 nm measured with a spectrophotometer (Beckman).

The quantity of SOD contained in sample is defined as the reduction quantity of reducing cytochrome c at 50% and 1 unit of SOD is expressed by unit/mg protein. In this analysis, sample itself not only inhibits the reaction on cytochrome c without the mediation of O_2^- , a small quantity of

cytochrome c would be directly reduced.

With these factors, the real SOD quantity was calculated by the formula:

$$\text{unit} = \{a - (b - c)\} / (a/2)$$

where

a: absorbance obtained by the addition of xanthine oxidase alone.

b: absorbance obtained by leaf homogenates in the presence of xanthine oxidase.

c: absorbance obtained by the addition of leaf homogenate alone before the addition of xanthine oxidase.

(Catalase Activity Assay)

The catalase activity was determined on the velocity of reduction of hydrogen peroxide (H₂O₂). The velocity of reduction was measured for 12 - 30 seconds in the presence of leaf homogenate samples, using a spectrophotometer at 240 nm. Activity was expressed with the formula shown below.

$$K = (2.3/18) \times \log (A_1/A_2)$$

A₁: Absorbance measured for 12 seconds at 240 nm for the assay mixture of total volume at 3.1 ml consisting 10mmol/L H₂O₂ dissolving in 3 ml of 50 m mol/L potassium phosphate buffer solution and 0.1 ml of leaf homogenate.

A₂: Absorbance measured for 30 seconds at 240 nm for above assay mixture

In this assay, hydrogen peroxide used as substrate may have the possibility of instability and thus substrate, hydrogen peroxide was substituted by sodium perborate. In this case, 0.002-0.05 ml leaf sample

solution was suspended in 2.8 ml of 0.05 mol/L potassium phosphate buffer solution (pH 7.4), the mixture was kept at 30 °C preincubation for 5 minutes. Thereafter, to the preincubated mixture, 0.2 N sodium perborat solution was added in cuvette and the reaction was started . The data was recorded for 2-3 minutes at 220 nm.

(GSH-Px Activity Assay)

For GSH-Px measurement, the method of Lawrence and Burk, connecting to with oxidation of NADPH by glutathione reductase was used. The oxidation potency for NADPH was measured at 37 °C with a spectrophotometer, its wave length at 340 nm.

The assay mixture was consisted of 50 mmol/L potassium phosphate buffer solution (pH 7.0), 1 mmol/L ethylenediaminetetraacetic acid, 1 mmol/L NaN₃, 0.2 mmol/L NADPH, 1 mmol/L glutathione, 2 unit of glutathione reductase, and 1.5 mmol/L cumene hydroperoxide or 10 mmol/L tert-butylperoxide solutions.

To the above mixture, leave homogenate was added to make the resulting solution volume was 1.0 ml. Thus the enzyme activity measured by this method is shown as the quantity of NADPH oxidized per minute. The activity is shown by unit/mg protein according to Lowry's method.

(Lipid Peroxide Assay)

The polyunsaturated fatty acid 4,7,10, 13, 16, 19-docosahexanoic acid was diluted to 200-fold in 94% ethanol. To the test tube previously poured with this substrate, then various concentration of homogenate was added and subjected to radiation with ultraviolet rays. In the experiment, homogenates of 0.6, 3.0 and 6.0 mg/ml concentration were tested. After radiation, the solution was analyzed by fluorospectrometry to measure the

quantity of TBA (thiobarbituric acid). The wave length at 515 nm for excitation and at 553 nm for emission were set in an equipment of Hitachi F-2000 fluorospectrophotometer.

The result is shown with mean $\pm$ S.E. The quantity of TBA substance was adopted as lipid peroxide quantity. Furthermore, significant difference analysis was performed by Student t-test. The result of t-test demonstrates the significant difference between two groups.

(Identification of Ploids of Chromosome)

Poloidy of the previously described trees was determined by cytological analysis by using root ends cells or flowcytometric analysis by using leaf cells.

Discussion on Tables 1-6, was performed as follows. As shown in Tables 1- 4, rice and osmund leaves (ref. Table 4) showed an apparent high SOD activity than those of gymnosperms plants such as Japanese black pines, Japanese cedar, Japanese cypress. Especially, GHS-Px of rice leaf showed as high as 1.384 unit/g and SOD activity as very high as 2941 unit/g.

As can be seen from Table 4, the SOD activity for rice leaf is higher than those of individual trees tested (except Japanese cedar, a tetraploid plant) , and in some cases, the highest SOD activity, showed a 100-fold activity of individual ordinary tree tested. In Table 4, osmund leaf also showed the SOD activity as high as 1394 unit/g.

Gymnosperms plants as the test subject, as shown in Table 1 ($P < 0.01$), Japanese black pines possessing ultraviolet-resistant property, its SOD activity is found to be apparently higher than the ordinary Japanese black pines. For example, the SOD activity of the Japanese black pines possessing ultraviolet-resistant property is 413 ± 56 unit/g on average, and it is much

higher 240 ± 37 unit/g on average of the ordinary Japanese black pines.

From the above results, it is ascertained that in the case the tree possesses the ultraviolet-resistant properties, the tree has high SOD activity. This may suggest that SOD activity play a role to protect them from damage of ultraviolet rays.

When comparing the SOD activity of triploid species tree and tetraploid species tree, as shown in Tables 2-3, especially in the individual trees, the tetraploid species tree showed a higher SOD activity than triploid species tree. The plus tree of the Japanese cedar (sample name Kamikawa cedar, Cr-38c) as shown in Table 4, showed a very high SOD activity than those ordinary trees. In such Japanese cedar, the apparent increase of SOD activity was found in trees possessing tetraploid chromosome. In contrast to this, both diploid and triploid trees showed no apparent increase of SOD activity.

Secondly, as to the lipid peroxide level, it was demonstrated that those cedars with small quantity of resin have a very little lipid peroxide formation ability compared with other trees.

The difference of GSH-Px activity between ultraviolet-resistant Japanese black pines and ultraviolet-sensitive ordinary Japanese black pines as control in Table was not found (refer to Table 1, $P > 0.05$). This may suggest that no relationship of GSH-Px enzymes with ultraviolet resistance.

[Table 5] Comparison of SOD activity and the effect of lipid peroxidation by the extract of leaves among 2X, 3X and 4X in Japanese trees

Samples	Polyploidy	SOD Activity (unit/gr)	Lipid peroxidation	
			average	
			UV(-) control 45	
			UV(+)control 260	
			Sample added	% control
S-20	2X	138±15	x1*(0.6mgr/ml)	124.6±14.9
			x5*(3.0mgr/ml)	73.3±9.5
			x10*(6.0mgr/ml)	33.8±4.7
S-22	2X	149±17	x1	115.3±13.8
			x5	71.4±9.2
			x10	31.5±4.4
S-23	2X	421±54	x1	221.5±24.3
			x5	137.9±17.9
			x10	60.3±7.2
S-25	3X	629±81	x1	174.6±19.2
			x5	112.5±14.6
			x10	63.9±10.2
S-26	3X	2173±260	x1	82.7±10.7
			x5	62.1±9.9
			x10	43.6±5.2
S-28	3X	309±33	x1	150.6±24.0
			x5	89.2±13.3
			x10	41.9±6.7
S-30	4X	1382±193	x1	84.2±10.9
			x5	62.3±8.7
			x10	42.7±4.6
S-31	4X	1447±231	x1	79.4±10.3
			x5	57.3±9.1
			x10	37.1±5.1
S-32	4X	1096±164	x1	72.9±9.4
			x5	61.4±7.9
			x10	51.9±8.3
S-33	4X	1237±160	x1	71.6±12.1
			x5	49.8±6.4
			x10	38.4±6.1
S-34	4X	1645±213	x1	86.0±11.1
			x5	64.1±9.6
			x10	44.1±7.0
S-36	4X	1105±154	x1	140.1±15.4
			x5	94.8±15.1
			x10	53.7±6.9
S-38	4X	1046±135	x1	136.6±19.1
			x5	111.6±15.6
			x10	89.0±10.6

*x1 denotes 0.6 gr/ml as unit and the column "sampl added" shows multiple of the unit

[Table 6] Comparison of SOD activity and the effect of lipid peroxidation by the extract of leaves among 2X, 3X and 4X in Japanese trees

Samples	Polyploidy	SOD activity (unit/gr)	Lipid peroxidation	
			Average	
			UV(-) control 45	
			UV(+)control 260	
			Sample added	% control
H-1	2X	160±19	x1*(0.6mgr/ml)	163.0±17.9
			x5*(3.0mgr/ml)	98.2±11.7
			x10*(6.0mgr/ml)	58.5± 7.6
H-2	2X	352±45	x 1	151.8±21.2
			x 5	103.4±11.3
			x10	56.8± 6.2
H-5	3X	578±80	x 1	170.0±22.1
			x 5	114.3±17.1
			x10	60.8± 7.2
H-6	3X	1066±138	x 1	95.2±13.3
			x 5	53.1± 6.3
			x10	45.2± 5.8
H-7	3X	742±89	x 1	139.1±15.3
			x 5	87.6±10.5
			x10	40.4± 4.4
H-8	3X	821±106	x 1	123.6±14.8
			x 5	78.4± 9.4
			x10	38.1± 4.9
H-9	3X	1280±140	x 1	92.5±12.9
			x 5	62.3± 6.9
			x10	39.0± 4.6
H-10	4X	1773±265	x 1	52.6± 6.8
			x 5	37.8± 6.0
			x10	25.3± 3.5
H-11	4X	1232±172	x 1	90.8±12.7
			x 5	61.4± 7.9
			x10	32.3± 4.8
rice leaves		2941±352	x 1	455.2±54.6
			x 5	298.3±38.7
			x10	165.0±21.4

*x1 denotes 0.6 gr/ml as unit and the column "sampl added" shows multiple of the unit

Lipid peroxide formation was enhanced at low concentration of homogenate (e.g. 0.6 mg/ml) and a high concentration of homogenate inhibited the formation of lipid peroxide. This is clearly shown in Tables 5 and 6.

As shown in Tables 5 and 6, when tetraploid tree leaf samples are compared each other, it is clear that in those samples of high SOD activity whose inhibitory effects is strong in presence of low sample concentration. This may suggest that SOD activity plays a role to inhibit formation of the lipid peroxide as cytotoxic role . This means that the higher SOD activity, the more extensive inhibition on the lipid peroxide formation.

For example, Japanese cedar sample nos. S-26, -30, -32, -34, -36 -38 in Table 5 and Japanese cypress samples nos. H-6, -9, -10, -11 in Table 6, have higher SOD activity than 1,000 unit/g. Among them, nine cases: sample nos. S-26, -30, -32, -34; H-6, -9, -10, showed a lower lipid peroxide formation (indicating in column of x1 of Table 5) .with addition of low sample concentration than controls.

In the case of rice leaf's sample, the formation of lipid peroxide was inhibited regardless of low concentration of homogenate (see "x1 column" of Table 5) or high concentration (see "x10 column" of Table 5). Thus SOD activity demonstrated a high activity as those of GHS-Px activity and catalase activity. Among the leaf samples tested, the rice leaf showed the highest level for lipid peroxide formation. Samples other than rice leaf, their catalase activities were not detectable and thus data are not included in Tables.

On the other hand, the higher SOD activity, the stronger anti-oxidation effects are known as usual. The SOD plays an important role of removal of active oxygen.

Upon consideration on factors in relating to formation of active oxygen, the existence of ultraviolet rays is not neglected. In recent years, destruction of ozonosphere involved with increase of ultraviolet radiation forwarding to earth. Based on this evidence, apparent increase of oxygen radical had been identified. Like human beings, trees

Occur withering uprightly when exposing to ultraviolet rays and succeeding to high concentration of oxygen radical.

But according to the experimental results that SOD activity are high, as obtained by the present inventors for those trees possessing ultraviolet resistance, they are thought to play a function to protect themselves from the damage exerted from environment by using the anti-oxidization agents such as SOD containing in their leaves. Until

now, no reports concerning the self-protection against the environmental damage, by trees using anti-oxidization agent possessed in the trees. This is the new finding by the present inventors.

The anti-oxidization ability of ultraviolet-resistant pine tree (hereinafter abbreviated as "UV-resistant") was compared with that of ordinary pine tree. Usually, reports on induction of GSH-px activity in response to oxygen radical were available. But in the experiments performed by the present inventors, no result on apparent change of GSH-px activity paralleled to SOD activity was not established.

Regarding to catalase, there was a report dealt with the occurrence of catalase induced by the oxygen radicals in tree. But in the present experiment found no occurrence of catalase except for rice leaves. This may be considered that terpene, etc in the homogenates inhibits the activation.

It is well known that like the enzymes such as superoxidase, the low

molecular weight substances for example, vitamin C as oxidizing agent, plays a major role in trees. With the experiment in this invention, study on anti-oxidization function of low molecular weight compound was not performed but those on SOD activity performed. It is concluded that SOD, a strong anti-oxidizing enzyme is occurring in the tree leaves for protecting itself against excessive oxygen radicals.

Succeeding to the above discussions, when consider the relationship of SOD activity and chromosomes of plants, those trees possessing triploid and tetraploid, especially the tetraploid trees have a high SOD activity found by the present inventors first time. There was no discussion on the relationship between extent of SOD activity and ploid of tree done by others, the results obtained by the present inventors as mentioned above are utterly beyond estimation.

When consider on the relationship of SOD activity and multiploid of chromosome, an increase of locus of chromosome encoding SOD, this brings the increase of SOD quantity existing. This consideration may be affirmative as referring to the evidence for example, in human beings, Down's syndrome was found to have three chromosomes (trysomy) containing Cu,Zn-SOD loci and this leading to an increase by 50% of SOD quantity.

Though the increase of SOD activity found in the trees of triploid and of tetraploid individually, in the experiments performed by the present inventors, an promoting level of SOD itself was not confirmed. This is to say that another elucidation may be applicable to consider, however It is concluded that locus of SOD chromosome is not found to increase in the individual sample tested.

And rice leaves are thought to possess the ultraviolet light resistance,

its SOD activity was found to be not only high than those of Japanese cedar, black pine and plus tree of Japanese cypress, but also showed a high GHS-Px acitivity than the trees. Furthermore, catalase was hardly detectable in the trees, rice leaves showed a detectable level.

Namely, with an ability that a strong enzyme activity possessed by the tree in response to the environmental oxygen radical and ultraviolet rays, the monocotyledonous plants can acquire the high adaptability against environmental ultraviolet rays. For example, a fern is more primitive plant than gymensperms and angiosperms, and it is well known that a fern acquire the high adaptability by multiplying their chromosome number to multiploid. Improving adaptability against the environmental perturbation by acquiring multiploid fact is also reported recently in bacteria.

Furthermore, from the experiments described above, in the Japanese cedar, Japanese cypress possessing ultraviolet-resistant properties were proved to have triploid and tetraploid of chromosomes.

While even those ultraviolet-sensitive gymensperms, strengthen their SOD activity by multiplying their ploid of chromosome, to protect themselves in response to the increased exposing time to ultraviolet rays and oxygen radical. This is to say that the species appeared to have adaptability is considerable.

On the other hand, it is considerable that UV-resistant black pine tree increases SOD to protect itself without the relationship of multiploid of chromosome in contrast to the ordinary black pine tree. It is also thought that the UV-resistant black pine tree does not adopt the measure to be taken for multiplying its chromosome into multiploid. As a general thinking, the increased UV-B(B region ultraviolet rays distributes

in 280 nm-320 nm) radiation is known to be not connected to damage of oxidation to the bodies such as plants, etc. Thus the case of black pine tree is corresponding to this hypothesis.

There two types on acquisition of multiploid of chromosome by plants: one type is natural or artificial cross, the other one is nature mutation such as branch change type or the artificial mutation with colchicines treatment.

Osmunda, a fern, comparing to gymnesperms and angiosperms plants, many kinds of fern already acquired multiploid. It is well known that monocotyledonous plants such as rice etc. possess the ultraviolet resistance than the black pine, Japanese cedars and Japanese cypress. The experiments performed by the present inventors indicate that both osmunda and the rice leaves showed a stronger SOD activity than other trees. Gymnesperms plants such as black pine, Japanese cedars, and Japanese cypress, etc., should to demonstrate their actions in response to the real environment by multiplying their chromosome having SOD gene , as the present inventors observed so.

From survey of references on agriculture and gardening field, it is confirmed that in the field of improvement of kind of plants such as green tea, apple, banana, tulip, orchid etc, polyploid is used in many cases. But in the conventional Japanese cedar, Japanese cypress, etc , the multiploid species are found to grow not so good as the ordinary diploid species. This value of multiploid species was utterly not noticed by the forestry industries.

In the plus tree of Japanese cedars and Japanese cypress, the present inventors found out the natural triploid species among the sterile species with cell biological research. The reports together with the results obtained by present inventors, reveal that only 41 clones of Japanese cedars and two

clones of plus tree of Japanese cypress were found to be the triploid species, respectively. By the way, "plus tree" is a kind of trees shows a good growth character.

The present inventors made an effort to breed the species of triploid by using artificial crossing technology. The effort was performed by crossing between diploid and tetraploid species in Japanese cedar and Japanese cypress, respectively. The artificial triploid species bred thus obtained by artificial crossing, were confirmed to grow better than diploid species by the present inventors.

Namely, a better growth of artificial crossing triploid species was confirmed than the conventional triploid one. This is concluded that it is possible and beneficial to use the multiploid to breed Japanese cedar and Japanese cypress.

Furthermore in the lipid peroxide assay, the leave homogenate of tetraploid containing a high SOD activity, apparently inhibited the lipid peroxide formation. This evidence may suggest that trees protect themselves from the damage by oxygen radicals introduced by ultraviolet rays through multiplying their chromosomal number.

Those Japanese cedars except tetraploid cedar, containing comparatively small resin than other trees, showed a low lipid peroxide Level and low SOD activity. Such results were identified in Tables 2-4. Because this Japanese cedar contains small resin, it is clear that the formed lipid peroxide can be estimated to be small. This means that treatment of low lipid peroxide formation, a low SOD activity is considered to be enough.

Namely, it is considered that Japanese cedar not needs to use high SOD activity to treat the low lipid peroxide, because, cedar tree does not produce the quantity of lipid peroxide.

According to the research of present inventors, it was confirmed that trees show an excellent adaptability against the increase of harmful oxygen radicals occurring due to the recent acceleration tendency of ozonosphere damage. A fern, an evolutionary more primitive plant than gymnosperms and angiosperms plants, acquired multiploid as its adaptability against the disadvantage of environmental perturbations by multiplying chromosome number.

Also in those needle-leaved plants such as Japanese cedar and Japanese cypress, it is found that their seedling acquired the ultraviolet resistant property by multiplying their chromosome number in the comparison of the ultraviolet resistant species with the ultraviolet non-resistant species. Especially, triploid, tetraploid trees were found to have this tendency.

On the other hand, although some species with natural cross, grew not good or became sterile, it is confirmed that some species obtain a good growth characteristics by artificial cross.

Namely, for Japanese cedar, Japanese cypress, etc. the means to acquire adaptability by multiplying ploid into multiploid were found to be adoptable. This production of multiploid descendants by artificial crossing etc. can be said as the effective method to protect trees from increasing tendency of oxygen radicals induced by ultraviolet rays.

This invention is not necessary to limit to the form of above examples, all the examples if not separated from the object of this invention is termed as within the range of this invention.

For example, the above needle-leaved trees such as Japanese cedar, Japanese cypress were explained and this will be considered also to adapt to the broad-leaved trees. And this also be considered to adapt to evergreen

trees, deciduous trees. On the other hand, on the evidence that tree acquired the ultraviolet rays resistance by multiplying its chromosome number, though this experiment was not obtained in Japanese black pine tree, there might of course be some species of Japanese black pine trees acquired the ultraviolet rays resistance through obtaining the multiploid means, because there are many kinds of pine trees.

It is considered that the strong plant species of ultraviolet resistance can be selected from the standard of ploid of chromosome. And the selected plants are planted in the plenty sunshine area to keep the strong ultraviolet rays resistance plant ecology, this means are considered to be the positive protection of plant ecological environment in the area.

Though the above explanation is focused on the triploid and tetraploid, Those multiploid trees such as pentaploid and hexaploid chromosomes will be the same result obtainable.

In the man-planted forest of the present invention, damages such as death of trees caused by ultraviolet rays are significantly inhibited . Accordingly, if the scale of ultraviolet radiation on the earth increases due to the progression of destruction of the ozonosphere the man-planted forest of the present invention will reliably be remained. By employing the method of planting trees of the present invention, a man-planted forest having such resistance to ultraviolet can be formed.

What is claimed is:

1. A man-planted forest, which is formed by planting seedling whose chromosomes ploid is multiploid species.
2. The man-planted forest according to Claim 1, wherein the chromosome number of seedling is multiploid such as triploid or tetraploid.
3. The man-planted forest according to Claim 2, wherein the seedlings are those of conifer.
4. The man-panted forest according to Claim 3, wherein the seedling of a conifer are cedar or cypress.
5. The man-planted forest according to Claim 4, wherein the chromosome number of the seedlings is obtained by means of increasing the basic number of chromosomes such as breed and mutation.
6. The man-planted forest according to Claim 1, wherein the seedlings are those of conifer.
7. The man-panted forest according to Claim 1, wherein the seedling of a conifer are cedar or cypress.
8. The method of planting trees, which comprises the steps of selecting seedling having multiploid; and planting the selected seedling.
9. The method of planting trees according to Claim 8, wherein the chromosome number of the seedling is multiploid such as triploid and tetraploid
10. The method of planting trees according to Claims 9, wherein the seedlings are those of a conifer.
11. The method of planting trees according to Claims 10, wherein



the seedling of a conifer are cedar or cypress.

12. The method of planting trees according to Claims 11, wherein the chromosome number of the seedlings is obtained by means of increasing the basic number of chromosomes such as breed and mutation.
13. The method of planting trees according to Claims 8, wherein the seedlings are those of a conifer.
14. The method of planting trees according to Claims 8, wherein the seedling of a conifer are cedar or cypress.
15. The method of growing plants, which comprises the step of artificially forming vegetation which is significantly resistant to ultraviolet rays, as compared with vegetation constituted of diploid plants, by selectively growing plants whose chromosome number exceeds the basic number of the species.

