



159159



G00002

DESCRIPTION

PROCESS FOR PRODUCING CYCLIC COMPOUND

Technical Field

5 The present invention relates to a process for producing cyclic compounds useful as medicines, agricultural chemicals, foods, cosmetics and chemical products, or intermediates thereof.

10 Background Art

 Heretofore, several synthetic methods for cyclic compounds have been described in literature, including, for example, (1) an intramolecular Friedel-Crafts Reaction in the case of cyclic ketone derivatives (JP-A 57-192379, etc.), (2) a Dieckmann-type cyclization reaction of diester
15 derivatives, followed by reduction and dehydration reactions in the case of cyclic acrylic acid derivatives (J. Chem. Res., Synop., (1987), (12), 394-5, etc.), (3) an intramolecular Wittig Reaction in the case of cyclic alkene
20 derivatives (WO98/55475, etc.) and others.

 However, even though a variety of useful intermediates have been described, for example, in J. Heterocyclic Chem., 31, 351-355 (1994), Heterocycles, 49, 215-232 (1998), J. Heterocyclic Chem., 33, 1909-1913 (1996), etc., cyclic
25 compounds (especially, cyclic compounds having 8- or more-

membered ring) are still generally difficult to synthesize and can be obtained only at low yields even if possible.


From the current situation as described above, it is desirable to establish a convenient production method (i.e.,
5 a production method which employs readily available low-cost raw materials, and reagents suitable for large scale synthesis and having no disposal problems, and which comprises relatively short steps and is easier to carry out), of cyclic compounds, especially cyclic alkene
10 derivatives, which can be easily converted to a variety of derivatives including, e.g., cyclic ketone derivatives or cyclic acrylic acid derivatives, due to their characteristic chemical structure.

15 Disclosure of Invention


As a result of intensive studies, the present inventors found that a cyclic alkene derivative can be produced inexpensively and conveniently by using an activated methylene derivative as a reaction intermediate
20 and by conducting a cyclization reaction in the presence of a carbonic diester, and on the basis of this knowledge, after further studies, completed the present invention.

That is, the present invention provides

(1) A process for producing a compound represented by
25 the formula:



[wherein, Z is an electron-withdrawing group; W is an optionally substituted ethylene or an optionally substituted vinylene, and when the ethylene or the vinylene has two substituent groups, said substituent groups may be bound to each other to form an optionally substituted cyclic ring; R³ is a hydrogen atom or an optionally substituted hydrocarbon group ; and X is a divalent group (preferably, a divalent group having 3 or more atoms in the straight-chain moiety thereof) (however, when W is an optionally substituted vinylene, -X-CH₂-Z is not -X¹-X²-CH₂-Z (wherein, X¹ is a sulfur atom or an optionally substituted nitrogen atom, X² is an optionally substituted ethylene))] or a salt thereof, which comprises subjecting a compound represented by the formula:



[wherein, each symbol has the same meaning described as above] or a salt thereof to a cyclization reaction in a solvent containing a carbonic diester;

(2) The process according to the above-mentioned (1), wherein X is a divalent group having 1 to 12 atoms in the straight-chain moiety thereof (preferably, 3 to 12 atoms in the straight-chain moiety thereof);


5 (3) The process according to the above-mentioned (1), wherein Z is a carboxyl group which is esterified;

(4) The process according to the above-mentioned (1), wherein R^3 is a hydrogen atom;

10 (5) The process according to the above-mentioned (1), which comprises the reaction being carried out in the presence of a base;

(6) The process according to the above-mentioned (5), wherein the base is an alcoholate;


15 (7) The process for producing a compound represented by the formula:



[wherein, Z is an electron-withdrawing group; R^1 and R^2 each are a hydrogen atom, a halogen atom, an optionally substituted amino group, an optionally substituted hydroxy group, an optionally substituted thiol group, an optionally substituted hydrocarbon group, or an optionally substituted heterocyclic; R^1 and R^2 may be bound to each other to form

20

an optionally substituted ring; R^3 is a hydrogen atom or an optionally substituted hydrocarbon group; the combined line of a broken line and a solid line is a single bond or a double bond; and X is a divalent group (preferably, a divalent group having 3 or more atoms in the straight-chain moiety thereof) (however, when the combined line of a broken line and a solid line is a double bond, $-X-CH_2-Z$ is not $-X^1-X^2-CH_2-Z$ (wherein , X^1 is a sulfur atom or an optionally substituted nitrogen atom, X^2 is an optionally substituted ethylene))) or a salt thereof, which comprises
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55
 60
 65
 70
 75
 80
 85
 90
 95
 100
 105
 110
 115
 120
 125
 130
 135
 140
 145
 150
 155
 160
 165
 170
 175
 180
 185
 190
 195
 200
 205
 210
 215
 220
 225
 230
 235
 240
 245
 250
 255
 260
 265
 270
 275
 280
 285
 290
 295
 300
 305
 310
 315
 320
 325
 330
 335
 340
 345
 350
 355
 360
 365
 370
 375
 380
 385
 390
 395
 400
 405
 410
 415
 420
 425
 430
 435
 440
 445
 450
 455
 460
 465
 470
 475
 480
 485
 490
 495
 500
 505
 510
 515
 520
 525
 530
 535
 540
 545
 550
 555
 560
 565
 570
 575
 580
 585
 590
 595
 600
 605
 610
 615
 620
 625
 630
 635
 640
 645
 650
 655
 660
 665
 670
 675
 680
 685
 690
 695
 700
 705
 710
 715
 720
 725
 730
 735
 740
 745
 750
 755
 760
 765
 770
 775
 780
 785
 790
 795
 800
 805
 810
 815
 820
 825
 830
 835
 840
 845
 850
 855
 860
 865
 870
 875
 880
 885
 890
 895
 900
 905
 910
 915
 920
 925
 930
 935
 940
 945
 950
 955
 960
 965
 970
 975
 980
 985
 990
 995



[wherein, each symbol has the same meaning described as above] or a salt thereof to a cyclization reaction in a solvent containing a carbonic diester ;
 15
 20
 25
 30
 35
 40
 45
 50
 55
 60
 65
 70
 75
 80
 85
 90
 95
 100
 105
 110
 115
 120
 125
 130
 135
 140
 145
 150
 155
 160
 165
 170
 175
 180
 185
 190
 195
 200
 205
 210
 215
 220
 225
 230
 235
 240
 245
 250
 255
 260
 265
 270
 275
 280
 285
 290
 295
 300
 305
 310
 315
 320
 325
 330
 335
 340
 345
 350
 355
 360
 365
 370
 375
 380
 385
 390
 395
 400
 405
 410
 415
 420
 425
 430
 435
 440
 445
 450
 455
 460
 465
 470
 475
 480
 485
 490
 495
 500
 505
 510
 515
 520
 525
 530
 535
 540
 545
 550
 555
 560
 565
 570
 575
 580
 585
 590
 595
 600
 605
 610
 615
 620
 625
 630
 635
 640
 645
 650
 655
 660
 665
 670
 675
 680
 685
 690
 695
 700
 705
 710
 715
 720
 725
 730
 735
 740
 745
 750
 755
 760
 765
 770
 775
 780
 785
 790
 795
 800
 805
 810
 815
 820
 825
 830
 835
 840
 845
 850
 855
 860
 865
 870
 875
 880
 885
 890
 895
 900
 905
 910
 915
 920
 925
 930
 935
 940
 945
 950
 955
 960
 965
 970
 975
 980
 985
 990
 995

(8) The process according to the above-mentioned (7), wherein X is a divalent group having 1 to 12 atoms in the straight-chain moiety thereof (preferably, 3 to 12 atoms in the straight-chain moiety thereof);

20 (9) The process according to the above-mentioned (7), wherein Z is a carboxyl group which is esterified;


(10) The process according to the above-mentioned (7),

wherein R^3 is a hydrogen atom;

(11) The process according to the above-mentioned (7), which comprises the reaction being carried out in the presence of a base;

5 (12) The process according to the above-mentioned (11), wherein the base is an alcoholate;


(13) The process for producing a compound represented by the formula:



10 [wherein, Z is an electron-withdrawing group; R^3 is a hydrogen atom or an optionally substituted hydrocarbon group; the combined line of a broken line and a solid line is a single bond or a double bond; ring A is an optionally substituted ring; and X is a divalent group (preferably, a

15 divalent group having 3 or more atoms in the straight-chain moiety thereof) (however, when the ring A is an optionally substituted benzene ring and the combined line of a broken line and a solid line is a double bond, $-X-CH_2-Z$ is not $-X^1-X^2-CH_2-Z$ (wherein, X^1 is sulfur atom or an optionally

20 substituted nitrogen atom ; X^2 is an optionally substituted ethylene))] or a salt thereof, which comprises subjecting a compound represented by the formula:



[wherein, each symbol has the same meaning described as above] or a salt thereof to a cyclization reaction in a solvent containing a carbonic diester;

5 (14) The process according to the above-mentioned (13), wherein X is a divalent group having 1 to 12 atoms in the straight-chain moiety thereof (preferably, 3 to 12 atoms in the straight-chain moiety thereof);


10 (15) The process according to the above-mentioned (13), wherein Z is a carboxyl group which is esterified;

 (16) The process according to the above-mentioned (13), wherein R³ is a hydrogen atom;

15 (17) The process according to the above-mentioned (13), which comprises the reaction being carried out in the presence of a base;

 (18) The process according to the above-mentioned (17), wherein the base is an alcoholate;

 (19) The compound represented by the formula:




[wherein, Z is an electron-withdrawing group; W is an optionally substituted ethylene or an optionally substituted vinylene, and when the ethylene or the vinylene has two substituent groups, said substituent groups may be bound to each other to form an optionally substituted ring; R³ is a hydrogen atom or an optionally substituted hydrocarbon group; and X'' is a divalent group having 4 or more atoms in the straight-chain moiety thereof] or a salt thereof;

(20) The compound according to the above-mentioned (19), wherein X'' is a divalent group having 4 to 6 atoms in the straight-chain moiety thereof;

(21) The compound according to the above-mentioned (19), wherein Z is a carboxyl group which is esterified;

(22) The compound according to the above-mentioned (19), wherein R³ is a hydrogen atom;

(23) The compound represented by the formula:



[wherein, Z is an electron-withdrawing group; R¹ and R² each are a hydrogen atom, a halogen atom, an optionally substituted amino group, an optionally substituted hydroxy group, an optionally substituted thiol group, an optionally


substituted hydrocarbon group or an optionally substituted heterocyclic group; R^1 and R^2 may be bound to each other to form an optionally substituted ring; R^3 is a hydrogen atom or an optionally substituted hydrocarbon group; the
 5 combined line of a broken line and a solid line is a single bond or a double bond; and X'' is a divalent group having 4 or more atoms in the straight-chain moiety thereof] or a salt thereof;

(24) The compound according to the above-mentioned
 10 (23), wherein X'' is a divalent group having 4 to 6 atoms in the straight-chain moiety thereof;

(25) The compound according to the above-mentioned (23), wherein Z is a carboxyl group which is esterified;

(26) The compound according to the above-mentioned
 15 (23), wherein R^3 is a hydrogen atom;

(27) The compound represented by the formula:



[wherein, Z is an electron-withdrawing group; R^3 is a hydrogen atom or an optionally substituted hydrocarbon
 20 group; the combined line of a broken line and a solid line is a single bond or a double bond; ring A is an optionally substituted ring; and X'' is a divalent group having 4 or more atoms in the straight-chain moiety thereof] or a salt

thereof;

(28) The compound according to the above-mentioned (27), wherein X" is a divalent group having 4 to 6 atoms in the straight-chain moiety thereof;

5 (29) The compound according to the above-mentioned (27), wherein Z is a carboxyl group which is esterified; and

(30) The compound according to the above-mentioned (27), wherein R³ is a hydrogen atom; and the like.

10

Examples of "electron-withdrawing group" used in the present specification include (i) a carboxyl group which may be optionally esterified or amidated, (ii) a group represented by the formula: $-(CO)R^4$ (wherein, R⁴ is a hydrogen atom or an optionally substituted hydrocarbon group), (iii) nitrile group, (iv) nitro group, (v) a group represented by the formula: $-(SO_m)R^5$ (wherein, m is an integer of 1 or 2, and R⁵ is an optionally substituted hydrocarbon group), (vi) a group represented by the formula: $-PR^6R^7$ (wherein, R⁶ and R⁷ each are an optionally substituted hydrocarbon group), (vii) a group represented by the formula: $-(PO)(OR^8)(OR^9)$ (wherein, R⁸ and R⁹ each are a hydrogen or an optionally substituted hydrocarbon group), (viii) an optionally substituted aryl group,

25 (ix) an optionally substituted alkenyl group, (x) a halogen

atom (e.g., fluorine, chlorine, bromine, iodine, etc.) and
 (xi) a nitroso group and the like, preferably, a carboxyl
 group which may be optionally esterified or amidated, a
 group represented by the formula: $-(CO)R^4$, nitrile group,
 5 nitro group, a group represented by the formula: $-(SO_m)R^5$,
 a group represented by the formula: $-PR^6 R^7$, a group
 represented by the formula: $-(PO)(OR^8)(OR^9)$, more
 preferably, a carboxyl group which is esterified (e.g., a
 carboxyl group which is esterified by a C_{1-4} alkyl such as
 10 methoxycarbonyl, ethoxycarbonyl, t-butoxycarbonyl, etc.)
 and the like.

The "carboxyl group which is esterified" in the
 "carboxyl group which may be optionally esterified or
 amidated" of (i) above include a group represented by the
 15 formula: $-(CO)OR^{10}$ (wherein, R^{10} is an optionally
 substituted hydrocarbon group), while the "carboxyl group
 which is amidated" include a group represented by the
 formula: $-(CO)NR^{11} R^{12}$ (wherein, R^{11} and R^{12} are, respectively,
 a hydrogen atom or an optionally substituted hydrocarbon
 20 group, and R^{11} and R^{12} may be bound to each other to form a
 5- to 7-membered (preferably, 5- to 6-membered) cyclic
 amino group together with a neighboring nitrogen atom (e.g.,
 tetrahydropyrrole, piperazine, piperidine, morpholine,
 thiomorpholine, pyrrole, imidazole, etc.)).

25 Further, in the formula of (vi) or (vii) above, R^6 and

R⁷, or R⁸ and R⁹ may be bound to each other to form, for example, a lower (C₂₋₆) alkylene (e.g., dimethylene, trimethylene, tetramethylene, etc.), a lower (C₂₋₆) alkenylene (e.g., -CH₂ -CH=CH-, -CH₂ -CH₂ -CH=CH-, -CH₂ -CH=CH-CH₂ -, etc.), or a lower (C₄₋₆) alkadienylene (e.g., -CH=CH-CH=CH-, etc.), etc., preferably, a lower (C₁₋₆) alkylene, more preferably, a lower (C₂₋₄) alkylene, and these divalent groups may have optionally substituent groups including, for example, hydroxy group, a halogen atom, a C₁₋₄ alkyl, a C₁₋₄ alkoxy, etc.

Examples of the "aryl group" in the optionally substituted aryl group of (viii) above include, for example, C₆₋₁₄ aryl groups such as phenyl, naphthyl, etc., preferably, C₆₋₁₀ aryl groups, etc., more preferably, phenyl, etc. The aryl group may have 1 to 3 substituent groups similar to those of the "optionally substituted hydrocarbon group" described below.

The "alkenyl groups" in the optionally substituted alkenyl group of (ix) above include, for example, alkenyl groups having 2 to 10 carbon atoms such as vinyl, allyl, crotyl, 2-pentenyl, 3-hexenyl, etc., and the like, preferably, lower (C₂₋₆) alkenyl groups, etc., more preferably, vinyl, etc. The alkenyl group may have 1 to 3 substituent groups similar to those of the "optionally substituted hydrocarbon group" described below.

Examples of the "hydrocarbon group" in the "optionally substituted hydrocarbon group" used in the present specification include;

5 (1) an alkyl group (e.g., C_{1-10} alkyl groups such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, hexyl, heptyl, octyl, nonyl, decyl, etc., preferably, lower (C_{1-6}) alkyl groups, etc.);

10 (2) a cycloalkyl group (e.g., C_{3-7} cycloalkyl groups such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and the like);

 (3) an alkenyl group (e.g., alkenyl groups having 2 to 10 carbon atoms such as vinyl, allyl, crotyl, 2-pentenyl, 15 3- hexenyl, etc., preferably, lower (C_{2-6}) alkenyl groups, etc.);

 (4) a cycloalkenyl group (e.g., cycloalkenyl groups having 3 to 7 carbon atoms such as 2-cyclopentenyl, 2-cyclohexenyl, 2-cyclopentenylmethyl, 2-cyclohexenylmethyl 20 and the like);

 (5) an alkynyl group (e.g., alkynyl groups having 2 to 10 carbon atoms such as ethynyl, 1-propynyl, 2-propynyl, 1-butynyl, 2-pentynyl, 3-hexynyl, etc., preferably, lower (C_{2-6}) alkynyl groups and the like);

25 (6) an aryl group (e.g., C_{6-14} aryl groups such as

phenyl, naphthyl, etc., preferably, C₆₋₁₀ aryl groups, more preferably, phenyl, etc.); and

(7) an aralkyl group (e.g., phenyl-C₁₋₄ alkyl groups (e.g., benzyl, phenethyl, etc.) and the like). The
5 examples of the hydrocarbon groups are preferably, an alkyl group, more preferably, a C₁₋₄ alkyl group such as methyl, ethyl, etc., and particularly preferably, methyl.

Said hydrocarbon group may have substituent groups, including, for example, a halogen atom (e.g., fluorine, chlorine, bromine, iodine, etc.), nitro, cyano, hydroxy, an
10 optionally substituted thiol group (e.g., thiol, C₁₋₄ alkylthio, etc.), an optionally substituted amino group (e.g., amino, monoC₁₋₄ alkylamino, diC₁₋₄alkylamino, 5- to 6-membered cyclic amino groups such as tetrahydropyrrole, piperazine, piperidine, morpholine, thiomorpholine, pyrrole,
15 imidazole and the like), a carboxyl group which may be optionally esterified or amidated (e.g., carboxyl, C₁₋₄ alkoxy carbonyl, carbamoyl, monoC₁₋₄alkylcarbamoyl, diC₁₋₄alkylcarbamoyl, etc.), a C₁₋₄ alkyl group which may be
20 optionally substituted with halogen atoms or C₁₋₄ alkoxy groups (e.g., trifluoromethyl, methyl, ethyl, etc.), a C₁₋₄ alkoxy group which may be optionally substituted with halogen atoms or C₁₋₄ alkoxy groups (e.g., methoxy, ethoxy, trifluoromethoxy, trifluoroethoxy, etc.), formyl, a C₂₋₄
25 alkanoyl group (e.g., acetyl, propionyl, etc.), a C₁₋₄

alkylsulfonyl group (e.g., methanesulfonyl, ethanesulfonyl, etc.), and a C₁₋₄ alkylsulfinyl (e.g., methanesulfinyl, ethanesulfinyl, etc.). The number of the substituent groups is preferably 1 to 3.

5 In the formula above, the "ethylene" and the "vinylene" in the "optionally substituted ethylene " and the "optionally substituted vinylene" represented by W include divalent groups represented by the formula: -CH₂-CH₂- and the formula: -CH=CH-. These divalent groups may have substituent groups at any position where substitution is possible, and examples thereof include substituent groups similar to those of "a halogen atom", "an optionally substituted amino group", "an optionally substituted hydroxy group", "an optionally substituted thiol group",
10 "an optionally substituted hydrocarbon group" and "an optionally substituted heterocyclic group" represented by R¹ and R² described below. When the ethylene or the vinylene has two substituent groups, the substituent groups may be bound to each other to form an optionally substituted ring. Further, R¹ and R² may be bound to each other to form an optionally substituted ring. Here, examples of the optionally substituted ring are those similar to an optionally substituted ring as ring A described below (preferably, an optionally substituted
15 benzene ring and the like).

Examples of the "halogen atom" represented by R^1 and R^2 in the formula above include fluorine, chlorine, bromine, iodine, etc.

5 Examples of the "optionally substituted amino group" represented by R^1 and R^2 in the formula above include amino groups which may be optionally substituted by the "optionally substituted hydrocarbon groups" mentioned above. The number of substituent groups may be any of 0 to 2. When two substituent groups are present, the two substituent
10 groups may be the same or different. Further, the two substituent groups may be bound to each other to form a 5- to 7-membered (preferably, 5- to 6-membered) cyclic amino group together with neighboring nitrogen atoms (e.g., tetrahydropyrrole, piperazine, piperidine, morpholine,
15 thiomorpholine, pyrrole, imidazole, etc.).

Examples of the "optionally substituted hydroxy group" represented by R^1 and R^2 in the formula above include hydroxy groups which may be optionally substituted by the "optionally substituted hydrocarbon groups" mentioned above.

20 Examples of "the optionally substituted thiol group" represented by R^1 and R^2 in the formula above include thiol groups which may be optionally substituted by the "optionally substituted hydrocarbon groups" mentioned above.

In the formula above, examples of the "heterocyclic
25 ring" in the "optionally substituted heterocyclic group"

represented by R^1 and R^2 , include 5- to 7-membered aromatic heterocyclic rings and saturated or unsaturated non-aromatic heterocyclic rings (aliphatic heterocyclic rings), containing at least one (preferably, 1 to 4, more preferably 1 to 2) hetero atoms of 1 to 3 (preferably, 1 to 2) hetero atoms selected from the group consisting of oxygen atom, sulfur atom, nitrogen atom and other atoms.

Herein, examples of the "aromatic heterocyclic ring" include 5- to 6-membered aromatic monocyclic heterocyclic rings (e.g., furan, thiophene, pyrrole, oxazole, isoxazole, thiazole, isothiazole, imidazole, pyrazole, 1,2,3-oxadiazole, 1,2,4-oxadiazole, 1,3,4-oxadiazole, furazan, 1,2,3-thiadiazole, 1,2,4-thiadiazole, 1,3,4-thiadiazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole, pyridine, pyridazine, pyrimidine, pyrazine, triazine, etc.). Examples of the "non-aromatic heterocyclic ring" include 5- to 7-membered (preferably, 5- to 6-membered) saturated or unsaturated (preferably, saturated) non-aromatic heterocyclic rings (aliphatic heterocyclic rings) such as pyrrolidine, tetrahydrofuran, thiolane, piperidine, tetrahydropyran, morpholine, thiomorpholine, piperazine, pyran, etc., and 5- to 6-membered non-aromatic heterocyclic rings wherein all or a part of double bonds in the aforementioned aromatic monocyclic heterocyclic rings are saturated. The heterocyclic ring is preferably a 5- to 6-

membered aromatic ring, and more preferably furan, thiophene, pyrrole, or pyridine (preferably a 6-membered ring).

5 Examples of substituent groups which may be possessed by the heterocyclic ring include substituent groups exemplified as the substituent groups on the "optionally substituted hydrocarbon group" described above, and the number of the substituent groups is 1 to 3.

10 In the formula above, Z is preferably an esterified carboxyl group, R^3 is preferably a hydrogen atom, and R^1 and R^2 each is preferably a hydrogen atom or an optionally substituted hydrocarbon group, more preferably a hydrogen atom, and it is also preferable that R^1 and R^2 are bound to each other to form an optionally substituted benzene ring.

15 The substituent groups which may be possessed by the "optionally substituted benzene ring" used in the present specification include substituent groups exemplified as the substituent groups on the "optionally substituted hydrocarbon group" described above; the afore-described
20 "optionally substituted aryl groups", which may be bound via a spacer (e.g., a divalent group having 1 to 4 atoms in the straight-chain moiety thereof) (preferably, afore-described "optionally substituted aryl groups which is directly bound), and particularly preferably electron-
25 donating groups. The number of the substituent groups is 1

to 4.

Examples of the "spacer" are $-(CH_2)_a-$ [a is an integer of 1 to 4 (preferably, an integer of 1 to 2)], $-(CH_2)_b-X'-$ [b is an integer of 0 to 3 (preferably, an integer of 0 to 1), X' is an optionally substituted imino group (e.g., imino groups which may be substituted with lower (C_{1-6}) alkyl, lower (C_{3-7}) cycloalkyl, formyl, lower (C_{2-7}) alkanoyl, lower (C_{1-6}) alkoxy-carbonyl etc.), a carbonyl group, an oxygen atom or an optionally oxidized sulfur atom (e.g., $-S(O)_n-$ (n is an integer of 0 to 2))], $-CH=CH-$, $-C\equiv C-$, $-CO-NH-$, $-SO_2-NH-$ etc. (preferably $-(CH_2)_b-X'-$, more preferably $-CH_2-O-$). These groups may be bound to the "optionally substituted benzene ring" via either the right or left bond, but they are bound to the "optionally substituted benzene ring" preferably via the right bond.

Examples of the ring formed by binding R^1 and R^2 and the "optionally substituted ring" represented by A in the formula above include 5- to 7-membered (preferably, 5- to 6-membered) saturated or unsaturated alicyclic hydrocarbons such as C_{5-7} cycloalkanes (e.g., cyclopentane, cyclohexane, cycloheptane, etc.), C_{5-7} cycloalkenes (e.g., 1-cyclopentene, 2-cyclopentene, 3-cyclopentene, 2-cyclohexene, 3-cyclohexene, etc.), and C_{5-6} cycloalkadienes (e.g., 2,4-cyclopentadiene, 2,4-cyclohexadiene, 2,5-cyclohexadiene, etc.); 6- to 14-membered aromatic hydrocarbons such as

benzene, naphthalene, azulene, acenaphthylene, etc.; and 5- to 7-membered aromatic heterocyclic rings and saturated or unsaturated non-aromatic heterocyclic rings (aliphatic heterocyclic rings) containing at least one (preferably, 1 to 4, more preferably 1 to 2) hetero atoms of 1 to 3 (preferably, 1 to 2) hetero atoms selected from the group consisting of oxygen atom, sulfur atom, nitrogen atom and other atoms.

Examples of the "aromatic heterocyclic ring" include 5- to 6-membered aromatic monocyclic heterocyclic rings (e.g., furan, thiophene, pyrrole, oxazole, isoxazole, thiazole, isothiazole, imidazole, pyrazole, 1,2,3-oxadiazole, 1,2,3-thiadiazole, 1,2,3-triazole, pyridine, pyridazine, pyrimidine, pyrazine, etc.), and examples of the "non-aromatic heterocyclic ring" include 5- to 7-membered (preferably 5- to 6-membered) saturated or unsaturated non-aromatic monocyclic heterocyclic rings (aliphatic heterocyclic rings) such as pyrrolidine, tetrahydrofuran, thiolane, piperidine, morpholine, thiomorpholine, tetrahydropyran, piperazine, pyran etc., or 5- to 6-membered non-aromatic heterocyclic rings wherein all or a part of the double bonds in the afore-described aromatic monocyclic heterocyclic rings are saturated.

Further, the ring formed by binding R^1 and R^2 , and the "optionally substituted ring" represented by A in the

formula above may form a fused ring wherein 2 to 3 (preferably, 2) monocyclic rings described above are condensed. Preferable examples of the fused ring include fused rings of two same or different heterocyclic rings (preferably, one heterocyclic ring and one aromatic heterocyclic ring, more preferably, two same or different aromatic heterocyclic rings); and fused rings of a heterocyclic ring and a homocyclic ring (preferably, a heterocyclic ring and a benzene ring, more preferably, an aromatic heterocyclic ring and a benzene ring). Specific examples of the fused ring include indole, benzothiophene, benzofuran, benzimidazole, imidazo[1,2 - a]pyridine, quinoline, isoquinoline, cinnoline, etc.


The ring formed by binding R^1 and R^2 , and the "optionally substituted ring" represented by A in the formula above is preferably a 5- to 6-membered aromatic ring, more preferably benzene, furan, thiophene, pyrrole, or pyridine (preferably, a 6-membered ring), and particularly preferably benzene.

The ring formed by binding R^1 and R^2 , and the ring A may have substituent groups, examples thereof include those similar to the substituent groups which may be possessed by the "benzene ring" in the "optionally substituted benzene ring" described above. 1 to 3 same or different substituent groups may substitute at any position of the

ring where substitution is possible.


In the formula above, the "divalent group" represented by X is preferably a divalent group having 3 or more atoms in the straight-chain moiety thereof, and, for example, any group may be employed whichever ring B in the following
 5 formula can form a 5- to 16-membered ring (preferably 5- to 12-membered ring, more preferably 7- to 10-membered ring, particularly preferably 8- to 10-membered ring):

the formula




10

the formula



or the formula



15 That is, X may be any of divalent groups having 1 to 12 atoms in the straight-chain moiety thereof (preferably 1 to 8, more preferably 3 to 6, and particularly preferably 4 to

6), and specific examples of the divalent group include

(1) $-(CH_2)_{a1}-$ ($a1$ is an integer of 1 to 12, preferably 3 to 12), $-(CH_2)_{a2}-(CH=CH)-(CH_2)_{a3}-$ ($a2$ and $a3$ each are the same or different integer of 0 to 10. But, the sum of $a2$ and $a3$ is 10 or lower, preferably 1 to 10), $-(CH_2)_{a4}-(CH=C=CH)-(CH_2)_{a5}-$ ($a4$ and $a5$ each are the same or different integer of 0 to 9. But, the sum of $a4$ and $a5$ is 9 or lower),

(2) $-(CH_2)_{b1}-Y-(CH_2)_{b2}-$ ($b1$ and $b2$ each are the same or different integer of 0 to 11. But, the sum of $b1$ and $b2$ is 11 or lower, preferably 2 to 11. Y is O, S, or NH), $-(CH_2)_{b3}-Y-(CH_2)_{b4}-(CH=CH)-(CH_2)_{b5}-$ or $-(CH_2)_{b5}-(CH=CH)-(CH_2)_{b4}-Y-(CH_2)_{b3}-$ ($b3$, $b4$ and $b5$ are, respectively, the same or different integer of 0 to 9. But, the sum of $b3$, $b4$ and $b5$ is 9 or less. Y is O, S, or NH), $-(CH_2)_{b6}-Y-(CH_2)_{b7}-(CH=C=CH)-(CH_2)_{b8}-$ or $-(CH_2)_{b8}-(CH=C=CH)-(CH_2)_{b7}-Y-(CH_2)_{b6}-$ ($b6$, $b7$, and $b8$ are, respectively, the same or different integer of 0 to 8. But, the sum of $b6$, $b7$ and $b8$ is 8 or less. Y is O, S, or NH),

(3) $-(CH_2)_{c1}-(N=CH)-(CH_2)_{c2}-$ ($c1$ and $c2$ are, respectively, the same or different integer of 0 to 10. But, the sum of $c1$ and $c2$ is 10 or less, preferably 1 to 10), $-(CH_2)_{c3}-(N=CH)-(CH_2)_{c4}-(CH=CH)-(CH_2)_{c5}-$ or $-(CH_2)_{c5}-(CH=CH)-(CH_2)_{c4}-(N=CH)-(CH_2)_{c3}-$ ($c3$, $c4$ and $c5$ are, respectively, the same or different integer of 0 to 8. But, the sum of $c3$, $c4$ and $c5$ is 8 or less), $-(CH_2)_{c6}-(N=CH)-(CH_2)_{c7}-(CH=C=CH)-$

$(\text{CH}_2)_{c8}-$ or $-(\text{CH}_2)_{c8}-(\text{CH}=\text{C}=\text{CH})-(\text{CH}_2)_{c7}-(\text{N}=\text{CH})-(\text{CH}_2)_{c6}-$ ($c6$, $c7$ and $c8$ are, respectively, the same or different integer of 0 to 7. But, the sum of $c6$, $c7$ and $c8$ is 7 or less),

(4) $-(\text{CH}_2)_{d1}-(\text{CH}=\text{N})-(\text{CH}_2)_{d2}-$ ($d1$ and $d2$ are, respectively the same or different integer of 0 to 10. But, the sum of $d1$ and $d2$ is 10 or lower, preferably 1 to 10), $-(\text{CH}_2)_{d3}-(\text{CH}=\text{N})-(\text{CH}_2)_{d4}-(\text{CH}=\text{CH})-(\text{CH}_2)_{d5}-$ or $-(\text{CH}_2)_{d5}-(\text{CH}=\text{CH})-(\text{CH}_2)_{d4}-(\text{CH}=\text{N})-(\text{CH}_2)_{d3}-$ ($d3$, $d4$ and $d5$ are, respectively, the same or different integer of 0 to 8. But, the sum of $d3$, $d4$ and $d5$ is 8 or less), $-(\text{CH}_2)_{d6}-(\text{CH}=\text{N})-(\text{CH}_2)_{d7}-(\text{CH}=\text{C}=\text{CH})-(\text{CH}_2)_{d8}-$ or $-(\text{CH}_2)_{d8}-(\text{CH}=\text{C}=\text{CH})-(\text{CH}_2)_{d7}-(\text{CH}=\text{N})-(\text{CH}_2)_{d6}-$ ($d6$, $d7$ and $d8$ are, respectively, the same or different integer of 0 to 7. But, the sum of $d6$, $d7$ and $d8$ is 7 or less),

(5) $-(\text{CH}_2)_{e1}-(\text{N}=\text{N})-(\text{CH}_2)_{e2}-$ ($e1$ and $e2$ are, respectively, the same or different integer of 0 to 10. But, the sum of $e1$ and $e2$ is 10 or less, preferably 1 to 10), $-(\text{CH}_2)_{e3}-(\text{N}=\text{N})-(\text{CH}_2)_{e4}-(\text{CH}=\text{CH})-(\text{CH}_2)_{e5}-$ or $-(\text{CH}_2)_{e5}-(\text{CH}=\text{CH})-(\text{CH}_2)_{e4}-(\text{N}=\text{N})-(\text{CH}_2)_{e3}-$ ($e3$, $e4$ and $e5$ are, respectively, the same or different integer of 0 to 8. But, the sum of $e3$, $e4$ and $e5$ is 8 or less), $-(\text{CH}_2)_{e6}-(\text{N}=\text{N})-(\text{CH}_2)_{e7}-(\text{CH}=\text{C}=\text{CH})-(\text{CH}_2)_{e8}-$ or $-(\text{CH}_2)_{e8}-(\text{CH}=\text{C}=\text{CH})-(\text{CH}_2)_{e7}-(\text{N}=\text{N})-(\text{CH}_2)_{e6}-$ ($e6$, $e7$ and $e8$ are, respectively, the same or different integer of 0 to 7. But, the sum of $e6$, $e7$ and $e8$ is 7 or less),

(6) $-(\text{CH}_2)_{f1}-\text{Y}-(\text{CH}_2)_{f2}-(\text{N}=\text{CH})-(\text{CH}_2)_{f3}-$ ($f1$, $f2$ and $f3$ are, respectively, the same or different integer of 0 to 9. But,

the sum of f_1 , f_2 and f_3 is 9 or less. Y is O, S or NH), -
 $(CH_2)_{f_3}-(N=CH)-(CH_2)_{f_2}-Y-(CH_2)_{f_1}$ (f_1 , f_2 and f_3 are,
respectively, the same or different integer of 0 to 9. But,
the sum of f_1 , f_2 and f_3 is 9 or less, Y is O, S or NH), -
5 $(CH_2)_{f_4}-(CH=CH)-(CH_2)_{f_5}-Y-(CH_2)_{f_6}-(N=CH)-(CH_2)_{f_7}$ (f_4 , f_5 , f_6
and f_7 are, respectively, the same or different integer of
0 to 7. But, the sum of f_4 , f_5 , f_6 and f_7 is 7 or less. Y
is O, S or NH), - $(CH_2)_{f_8}-Y-(CH_2)_{f_9}-(CH=CH)-(CH_2)_{f_{10}}-(N=CH)-$
 $(CH_2)_{f_{11}}$ (f_8 , f_9 , f_{10} and f_{11} are, respectively, the same
10 or different integer of 0 to 7. But, the sum of f_8 , f_9 , f_{10}
and f_{11} is 7 or less. Y is O, S or NH), - $(CH_2)_{f_{12}}-Y-(CH_2)_{f_{13}}-$
 $(N=CH)-(CH_2)_{f_{14}}-(CH=CH)-(CH_2)_{f_{15}}$ (f_{12} , f_{13} , f_{14} and f_{15} are,
respectively, the same or different integer of 0 to 7. But,
the sum of f_{12} , f_{13} , f_{14} and f_{15} is 7 or less), - $(CH_2)_{f_7}-$
15 $(N=CH)-(CH_2)_{f_6}-Y-(CH_2)_{f_5}-(CH=CH)-(CH_2)_{f_4}$ (f_4 , f_5 , f_6 and f_7
are, respectively, the same or different integer of 0 to 7.
But, the sum of f_4 , f_5 , f_6 and f_7 is 7 or less. Y is O, S
or NH), - $(CH_2)_{f_{11}}-(N=CH)-(CH_2)_{f_{10}}-(CH=CH)-(CH_2)_{f_9}-Y-(CH_2)_{f_8}$ (f_8 ,
 f_9 , f_{10} and f_{11} are, respectively, the same or different
20 integer of 0 to 7. But, the sum of f_8 , f_9 , f_{10} and f_{11} is 7
or less. Y is O, S or NH), - $(CH_2)_{f_{15}}-(CH=CH)-(CH_2)_{f_{14}}-(N=CH)-$
 $(CH_2)_{f_{13}}-Y-(CH_2)_{f_{12}}$ (f_{12} , f_{13} , f_{14} and f_{15} are, respectively,
the same or different integer of 0 to 7. But, the sum of f_{12} ,
 f_{13} , f_{14} and f_{15} is 7 or less),
25 (7) - $(CH_2)_{g_1}-Y-(CH_2)_{g_2}-(CH=N)-(CH_2)_{g_3}$ (g_1 , g_2 and g_3 are,

respectively the same or different integer of 0 to 9. But,
 the sum of g_1 , g_2 and g_3 is 9 or less. Y is O, S or NH), -
 $(CH_2)_{g_3}-(CH=N)-(CH_2)_{g_2}-Y-(CH_2)_{g_1}-$ (g_1 , g_2 and g_3 are,
 respectively, the same or different integer of 0 to 9. But,
 5 the sum of g_1 , g_2 and g_3 is 9 or less. Y is O, S or NH), -
 $(CH_2)_{g_4}-(CH=CH)-(CH_2)_{g_5}-Y-(CH_2)_{g_6}-(CH=N)-(CH_2)_{g_7}-$ (g_4 , g_5 , g_6
 and g_7 are, respectively, the same or different integer of
 0 to 7. But, the sum of g_4 , g_5 , g_6 and g_7 is 7 or less. Y
 is O, S or NH), - $(CH_2)_{g_8}-Y-(CH_2)_{g_9}-(CH=CH)-(CH_2)_{g_{10}}-(CH=N)-$
 10 $(CH_2)_{g_{11}}-$ (g_8 , g_9 , g_{10} and g_{11} are, respectively, the same
 or different integer of 0 to 7. But, the sum of g_8 , g_9 , g_{10}
 and g_{11} is 7 or less. Y is O, S or NH), - $(CH_2)_{g_{12}}-Y-(CH_2)_{g_{13}}-$
 $(CH=N)-(CH_2)_{g_{14}}-(CH=CH)-(CH_2)_{g_{15}}-$ (g_{12} , g_{13} , g_{14} and g_{15} are,
 respectively, the same or different integer of 0 to 7. But,
 15 the sum of g_{12} , g_{13} , g_{14} and g_{15} is 7 or less), - $(CH_2)_{g_7}-$
 $(CH=N)-(CH_2)_{g_6}-Y-(CH_2)_{g_5}-(CH=CH)-(CH_2)_{g_4}-$ (g_4 , g_5 , g_6 and g_7
 are, respectively, the same or different integer of 0 to 7.
 But, the sum of g_4 , g_5 , g_6 and g_7 is 7 or less. Y is O, S
 or NH), - $(CH_2)_{g_{11}}-(CH=N)-(CH_2)_{g_{10}}-(CH=CH)-(CH_2)_{g_9}-Y-(CH_2)_{g_8}-$ (g_8 ,
 20 g_9 , g_{10} and g_{11} are, respectively, the same or different
 integer of 0 to 7. But, the sum of g_8 , g_9 , g_{10} and g_{11} is 7
 or less. Y is O, S or NH), - $(CH_2)_{g_{15}}-(CH=CH)-(CH_2)_{g_{14}}-(CH=N)-$
 $(CH_2)_{g_{13}}-Y-(CH_2)_{g_{12}}-$ (g_{12} , g_{13} , g_{14} and g_{15} are, respectively,
 the same or different integer of 0 to 7. But, the sum of
 25 g_{12} , g_{13} , g_{14} and g_{15} is 7 or less),

(8) $-(CH_2)_{h1}-Y-(CH_2)_{h2}-(N=N)-(CH_2)_{h3}-$ ($h1$, $h2$ and $h3$ are, respectively the same or different integer of 0 to 9. But, the sum of $h1$, $h2$ and $h3$ is 9 or less. Y is O, S or NH), $-(CH_2)_{h3}-(N=N)-(CH_2)_{h2}-Y-(CH_2)_{h1}-$ ($h1$, $h2$ and $h3$ are, respectively, the same or different integer of 0 to 9. But, the sum of $h1$, $h2$ and $h3$ is 9 or less. Y is O, S or NH), $-(CH_2)_{h4}-(CH=CH)-(CH_2)_{h5}-Y-(CH_2)_{h6}-(N=N)-(CH_2)_{h7}-$ ($h4$, $h5$, $h6$ and $h7$ are, respectively, the same or different integer of 0 to 7. But, the sum of $h4$, $h5$, $h6$ and $h7$ is 7 or less. Y is O, S or NH), $-(CH_2)_{h8}-Y-(CH_2)_{h9}-(CH=CH)-(CH_2)_{h10}-(N=N)-(CH_2)_{h11}-$ ($h8$, $h9$, $h10$ and $h11$ are, respectively, the same or different integer of 0 to 7. But, the sum of $h8$, $h9$, $h10$ and $h11$ is 7 or less. Y is O, S or NH), $-(CH_2)_{h12}-Y-(CH_2)_{h13}-(N=N)-(CH_2)_{h14}-(CH=CH)-(CH_2)_{h15}-$ ($h12$, $h13$, $h14$ and $h15$ are, respectively, the same or different integer of 0 to 7. But, the sum of $h12$, $h13$, $h14$ and $h15$ is 7 or less), $-(CH_2)_{h7}-(N=N)-(CH_2)_{h6}-Y-(CH_2)_{h5}-(CH=CH)-(CH_2)_{h4}-$ ($h4$, $h5$, $h6$ and $h7$ are, respectively, the same or different integer of 0 to 7. But, the sum of $h4$, $h5$, $h6$ and $h7$ is 7 or less. Y is O, S or NH), $-(CH_2)_{h11}-(N=N)-(CH_2)_{h10}-(CH=CH)-(CH_2)_{h9}-Y-(CH_2)_{h8}-$ ($h8$, $h9$, $h10$ and $h11$ are, respectively the same or different integer of 0 to 7. But, the sum of $h8$, $h9$, $h10$ and $h11$ is 7 or less. Y is O, S or NH), $-(CH_2)_{h15}-(CH=CH)-(CH_2)_{h14}-(N=N)-(CH_2)_{h13}-Y-(CH_2)_{h12}-$ ($h12$, $h13$, $h14$ and $h15$ are, respectively, the same or different integer of 0 to 7. But, the sum of

h12, h13, h14 and h15 is 7 or less),

(9) $-(\text{CH}_2)_{j1}-\text{Y}^1-(\text{CH}_2)_{j2}-\text{Y}^2-(\text{CH}_2)_{j3}-$ ($j1$, $j2$ and $j3$ are, respectively, the same or different integer of 0 to 10. But, the sum of $j1$, $j2$ and $j3$ is 10 or less, preferably 1 to 10.

5 Y^1 and Y^2 are, respectively, O, S or NH), $-(\text{CH}_2)_{j4}-\text{Y}^4-(\text{CH}_2)_{j5}-\text{Y}^5-(\text{CH}_2)_{j6}-\text{Y}^6-(\text{CH}_2)_{j7}-$ ($j4$, $j5$, $j6$ and $j7$ are, respectively, the same or different integer of 0 to 9. But, the sum of $j4$, $j5$, $j6$ and $j7$ is 9 or lower. Y^4 , Y^5 , Y^6 and Y^7 are, respectively, O, S or NH), and so on. Specific examples of

10 the divalent group include $-\text{O}-$, $-\text{O}-(\text{CH}_2)_{k1}-$ ($k1$ is an integer of 1 to 5, preferably 2 to 5), $-\text{O}-\text{CH}=\text{CH}-$, $-\text{O}-\text{CH}_2-\text{CH}=\text{CH}-$, $-\text{O}-\text{CH}=\text{CH}-\text{CH}_2-$, $-(\text{CH}_2)_{k1}-\text{O}-$ ($k1$ is an integer of 1 to 5, preferably 2 to 5), $-\text{CH}=\text{CH}-\text{O}-$, $-\text{CH}_2-\text{CH}=\text{CH}-\text{O}-$, $-\text{CH}=\text{CH}-\text{CH}_2-\text{O}-$, $-\text{S}-$, $-\text{S}-(\text{CH}_2)_{k2}-$ ($k2$ is an integer of 1 to 5, preferably 2 to 5), $-\text{S}-\text{CH}=\text{CH}-$, $-\text{S}-\text{CH}_2-\text{CH}=\text{CH}-$, $-\text{S}-\text{CH}=\text{CH}-\text{CH}_2-$, $-(\text{CH}_2)_{k1}-\text{S}-$ ($k1$ is an integer of 1 to 5, preferably 2 to 5), $-\text{CH}=\text{CH}-\text{S}-$, $-\text{CH}_2-\text{CH}=\text{CH}-\text{S}-$, $-\text{CH}=\text{CH}-\text{CH}_2-\text{S}-$, $-\text{NH}-$, $-\text{NH}-(\text{CH}_2)_{k2}-$ ($k2$ is an integer of 1 to 5, preferably 2 to 5), $-\text{NH}-\text{CH}=\text{CH}-$, $-\text{NH}-\text{CH}_2-\text{CH}=\text{CH}-$, $-\text{NH}-\text{CH}=\text{CH}-\text{CH}_2-$, $-(\text{CH}_2)_{k1}-\text{NH}-$ ($k1$ is an integer of 1 to 5, preferably 2 to 5), $-\text{CH}=\text{CH}-\text{NH}-$, $-\text{CH}_2-\text{CH}=\text{CH}-\text{NH}-$, $-\text{CH}=\text{CH}-\text{CH}_2-\text{NH}-$, $-(\text{CH}_2)_{k2}-$ ($k2$ is an integer of 1 to 5, preferably 2 to 5), $-\text{CH}=\text{CH}-$, $-\text{CH}_2-\text{CH}=\text{CH}-$, $-\text{CH}=\text{CH}-\text{CH}_2-$, $-\text{N}=\text{CH}-$, $-\text{CH}=\text{N}-$, $-\text{N}=\text{N}-$, $-\text{CH}_2-\text{N}=\text{CH}-$, $-\text{CH}_2-\text{CH}=\text{N}-$, $-\text{CH}_2-\text{N}=\text{N}-$, $-\text{N}=\text{CH}-\text{CH}_2-$, $-\text{CH}=\text{N}-\text{CH}_2-$, $-\text{N}=\text{N}-\text{CH}_2-$ etc., and preferable
20
25 examples include $-\text{O}-(\text{CH}_2)_{k1}-$ ($k1$ is an integer of 1 to 5,

preferably 2 to 5, more preferably 3 to 5), $-\text{S}-(\text{CH}_2)_{k2}-$ ($k2$ is an integer of 1 to 5, preferably 2 to 5, more preferably 3 to 5), $-\text{NH}-(\text{CH}_2)_{k2}-$ ($k2$ is an integer of 1 to 5, preferably 2 to 5, more preferably 3 to 5), $-(\text{CH}_2)_{k2}-$ ($k2$ is an integer of 1 to 5, preferably 2 to 5, more preferably 3 to 5), $-(\text{CH}_2)_{k1}-\text{O}-$ ($k1$ is an integer of 1 to 5, preferably 2 to 5, more preferably 3 to 5), $-(\text{CH}_2)_{k1}-\text{S}-$ ($k1$ is an integer of 1 to 5, preferably 2 to 5, more preferably 3 to 5), and $-(\text{CH}_2)_{k1}-\text{NH}-$ ($k1$ is an integer of 1 to 5, preferably 2 to 5, more preferably 3 to 5), and more preferable examples are $-\text{O}-(\text{CH}_2)_2-$, $-\text{O}-(\text{CH}_2)_3-$, $-\text{O}-(\text{CH}_2)_4-$, $-\text{O}-(\text{CH}_2)_5-$, $-\text{S}-(\text{CH}_2)_3-$, $-\text{S}-(\text{CH}_2)_4-$, $-\text{S}-(\text{CH}_2)_5-$, $-\text{NH}-(\text{CH}_2)_3-$, $-\text{NH}-(\text{CH}_2)_4-$, $-\text{NH}-(\text{CH}_2)_5-$ and the like.

The divalent group may have substituent groups, and said substituent groups may be any group which can be bound to the divalent group, including, for example, a halogen atom, an optionally substituted amino group, an optionally substituted hydroxy group, an optionally substituted thiol group, an optionally substituted hydrocarbon group and an optionally substituted heterocyclic group exemplified as R^1 and R^2 . Further, the substituent groups may be bound to each other to form a ring with the divalent group, and such rings include those exemplified as the rings formed by binding R^1 and R^2 . Further, when an atom constituting the straight chain moiety is a sulfur atom in the "divalent

group" represented by X above, the sulfur atom may be oxidized, and may be any of -S-, -SO- or -SO₂ -.

Further, R¹ and a substituent group on X or R² and R³ may be bound to each other to form an optionally substituted ring, and examples of such ring include those exemplified as the optionally substituted rings formed by binding R¹ and R². When R¹ and a substituent group on X are bound to each other to form an optionally substituted ring, a substituent group on the first to third (preferably the first to second, more preferably the first) atom in the straight chain moiety of X bound to the ethylene or vinylene is preferably bound to R¹.

Alternatively, a substituent group on the "ethylene" or "vinylene" of the "optionally substituted ethylene" or "optionally substituted vinylene" represented by W may be bound to a substituent group of X or R³ to form an optionally substituted ring, and examples of such ring include those exemplified as the optionally substituted rings formed by binding R¹ and R². When a substituent group of the "ethylene" or "vinylene" and a substituent group of X are bound to each other to form a ring, binding between a substituent on the first to third (preferably first to second, more preferably first) atom in the straight chain moiety of X bound to the ethylene or vinylene and a substituent group of the "ethylene" or

"vinylene" is preferable.

In the formula above, the combined line of "broken line and solid line" indicates a single bond or a double bond (preferably double bond).


5 When a compound having a substituent group is a basic compound depending on the kind of the substituent group exemplified above, the compound may be converted to a salt thereof by using an acid according to usual methods. Any acid may be used if the acid does not interfere with the
10 reaction, and examples of the acid include inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, nitric acid, sulfamic acid etc., organic acids such as formic acid, acetic acid, trifluoroacetic acid, tartaric acid, citric acid, fumaric
15 acid, maleic acid, succinic acid, malic acid, p-toluenesulfonic acid, methanesulfonic acid, benzenesulfonic acid etc., and acidic amino acids such as aspartic acid, glutamic acid etc. When an obtained compound is a salt, the compound may be converted to a free base according to
20 usual methods.

 Alternatively, when a compound having a substituent group is an acidic compound depending on the kind of the substituent group exemplified above, the compound may be converted to a salt thereof by using a base according to
25 usual methods. Any base may be used if the base does not


interfere with the reaction, and examples of such salts include salts with inorganic bases, organic bases, basic amino acids, etc. Preferable examples of the salt with an inorganic base include alkali metal salts such as sodium salt, potassium salt, etc.; alkali-earth metal salts such as calcium salt, magnesium salt, etc.; and aluminum salt, ammonium salt, etc. Preferable examples of the salt with an organic base include salts with trimethylamine, triethylamine, pyridine, picoline, ethanolamine, diethanolamine, triethanolamine, dicyclohexylamine, N,N'-dibenzylethylenediamine, etc. Preferable examples of the salt with a basic amino acid include salts with arginine, lysine, ornithine, etc. When an obtained compound is a salt, the compound may be converted to a free acid according to usual methods.

The reaction above-mentioned (1) is conducted, for example, under a following reaction condition.

The compound represented by the formula:



[wherein each symbol has the same meaning described above] or a salt thereof is produced by subjecting a compound represented by the formula:



[wherein each symbol has the same meaning described above] or a salt thereof to cyclization reaction in a solvent containing carbonic diester.

5 The reaction above-mentioned (1) is preferably conducted in the presence of a base, and examples of such base include metal hydride compounds (e.g., alkali metal hydrides such as sodium hydride, potassium hydride, etc.), metal hydrocarbons (e.g., compounds having direct chemical
10 bond between alkali metal and C₁₋₄ alkyl group such as n-butyllithium, etc.), alcoholates (e.g., compounds in which a hydroxy hydrogen of C₁₋₄ alcohols is replaced by a alkali metal such as sodium methoxide, sodium ethoxide, sodium t-butoxide, potassium methoxide, potassium ethoxide,
15 potassium t-butoxide, lithium methoxide, lithium ethoxide, lithium t-butoxide, etc.), alkali metal hydroxides (e.g., NaOH, KOH, etc.), basic carbonates (e.g., alkali metal salts of carbonate such as sodium salt, potassium salt, etc., or alkali-earth metal salts of carbonate such as
20 calcium salt, magnesium salt, etc.), basic bicarbonates (e.g., alkali metal salts of bicarbonate such as sodium salt, potassium salt, etc.), and organic bases (e.g.,

trimethylamine, triethylamine, diisopropylethylamine, pyridine, picoline, N-methylpyrrolidine, N-methylmorpholine, 1,5-diazabicyclo[4.3.0]non-5-ene, 1,4-diazabicyclo[2.2.2]octane, 1,8-diazabicyclo [5.4.0]-7-undecene, etc.), and preferably metal hydride compounds (e.g., sodium hydride, potassium hydride, etc.), alcoholates (e.g., NaOMe, NaOEt, t-BuONa, t-BuOK, etc.), particularly preferably alcoholates (e.g., NaOMe, NaOEt, t-BuONa, t-BuOK, etc.) is used.

10 The amount of base used for the reaction above-mentioned (1) is about 0.1 to 100 equivalents, preferably 1 to 5 equivalents.

 The reaction solvent may be any solvent if only it contains carbonic diester, for example, a carbonic diester alone, a mixture of 2 or more carbonic diesters or a mixed solvent of carbonic diesters and a solvent except carbonic diesters.

 The carbonic diester is a compound represented by the formula, $Z''-O(CO)O-Z'$, [wherein, Z'' and Z' each are an optionally substituted hydrocarbon group (preferably an optionally substituted alkyl group)]. Preferable are carbonic diesters which are liquid at the reaction temperature. In addition, Z'' and Z' are preferably the same, and preferable examples of the carbonic diester are dialkyl carbonates, and di C_{1-4} alkyl carbonates such as

dimethyl carbonate, diethyl carbonate are preferably used.

The "solvent except carbonic diesters" used in combination with one or more kinds of carbonic diesters may be any solvent if the solvent does not interfere with the reaction, and includes, for example, halogenated solvents (e.g., methylene chloride, dichloroethane, chloroform, etc.), aliphatic hydrocarbons (e.g., n-hexane, etc.), aromatic hydrocarbons (e.g., benzene, toluene, etc.), ethers (e.g., tetrahydrofuran (THF), diethylether, etc.), polar solvents (e.g., dimethylformamide (DMF), dimethylsulfoxide (DMSO), etc.), alcohols (e.g., methanol, ethanol, propanol, isopropanol, n-butanol, 2-methoxyethanol, etc.), formic acid esters (e.g., C₁₋₄ alkyl formate, etc.), and oxalic acid diesters (e.g., di C₁₋₄ alkyl oxalate, etc), and preferably dimethylformamide (DMF), alcohols (e.g., methanol, ethanol, etc.) and so on.


The reaction may be conducted in an appropriate mixed solvent, preferably in a solvent containing carbonic diester (preferably, dimethyl carbonate or diethyl carbonate).

The reaction temperature is usually about -20 to 200°C, preferably about 10 to 100°C, and the reaction period is usually about 0.1 to 100 hours, preferably about 0.5 to 50 hours.

The reaction above-mentioned (7) is conducted, for


example, under a following reaction condition.

The compound represented by the formula:



[wherein each symbol has the same meaning described above]

5 or a salt thereof is produced by subjecting a compound represented by the formula:




[wherein each symbol has the same meaning described above]

10 or a salt thereof to a cyclization reaction in a solvent containing a carbonic diester.

The reaction above-mentioned (7) may be conducted under a similar condition to that for the reaction above-mentioned (1).


15 The reaction above-mentioned (13) is conducted for example under a following reaction condition.

The compound represented by the formula:



[wherein each symbol has the same meaning described above]

or a salt thereof is produced by subjecting a compound represented by the formula:




[wherein each symbol has the same meaning described above]

5 or a salt thereof to a cyclization reaction in a solvent containing a carbonic diester.


The reaction above-mentioned (13) may be conducted under a condition similar to those for the reaction above-mentioned (1).

10 Among the compounds represented by the following formula:




[wherein each symbol has the same meaning described above]

15 or the salt thereof obtained in the reaction above-mentioned (1), the compounds represented by the formula:




[wherein, Z is an electron-withdrawing group, W is an optionally substituted ethylene or an optionally substituted vinylene and when the ethylene or vinylene has two substituent groups, said substituent groups may be bound to each other to form an optionally substituted ring, R³ is a hydrogen atom or an optionally substituted hydrocarbon group, and X'' is a divalent group having 4 or more atoms in the straight-chain moiety thereof (preferably, a divalent group having 4, 5 or 6 atoms in the straight-chain moiety thereof)] or the salts thereof are novel compounds which have not been described in literature.

Further, the compounds represented by the formula:



[wherein, each symbol has the same meaning described above] or a salt thereof obtained in the reaction above-mentioned (1) are useful as medicines, agricultural chemicals, foods, cosmetics and chemicals, or intermediates therefor. For example, from the compounds represented by the formula:



[wherein each symbol has the same meaning described above]
 or the salt thereof obtained in the reaction above-
 mentioned (1), useful medicines such as an osteogenesis
 5 promoter, a Na-H exchange inhibitor, treating agent for
 pollakiuria and incontinence of urine, edema treating agent
 and treating agent for central nervous system diseases can
 be produced according to the methods known in the art (e.g.,
 methods described in JP-A 8-73476, WO99/55690, JP-A 11-
 10 302267, JP-A 11-302270, WO99/51242, etc.).

Best Mode for Carrying Out the Invention

Hereinafter, the present invention is described in
 more detail by the following Reference Examples and
 15 Examples, which are not intended to restrict the present
 invention.

EXAMPLES

Preparation of 2-alkoxybenzaldehyde derivatives in general;

20 [REFERENCE EXAMPLE 1] Ethyl 4-(2-Formylphenoxy)butyrate

Ethyl 4-bromobutyrate (6.6ml) was added to a

suspension of salicyl aldehyde (5.0g) and potassium carbonate (6.2g) in DMF(15ml), and the mixture was stirred at 90°C for 1 hour. The mixture was allowed to cool to room temperature, and neutralized with 1N hydrochloric acid.

5 After extraction with ethyl acetate, the organic layer was washed with saturated salt water, dried over anhydrous sodium sulfate, and concentrated. The concentrate was purified by silica gel column chromatography (n-hexane/ethyl acetate=4/1) and subsequent concentration of
10 the appropriate eluate gave colorless oil of Ethyl 4-(2-formylphenoxy)butyrate (9.6g, yield 99%).

¹H-NMR(CDCl₃, δ, 300MHz): 1.27(3H, t, J=7.1Hz), 2.14-2.23(2H, m), 2.54(2H, t, J=7.2Hz), 4.11-4.18(4H, m), 6.96-7.04(2H, m), 7.52(1H, dt, J=1.7, 7.1Hz), 7.82(1H, dd, J=1.7,
15 7.7Hz), 10.49(1H, s).

IR(neat, cm⁻¹): 1733, 1687, 1598, 1243.

Compounds below in [REFERENCE EXAMPLE 2] to [REFERENCE EXAMPLE 4] were prepared according to a method similar to that of [REFERENCE EXAMPLE 1].

20

[REFERENCE EXAMPLE 2] Ethyl 4-(4-Bromo-2-formylphenoxy)butyrate

Colorless oil, yield 98%. ¹H-NMR(CDCl₃, δ, 300MHz):
1.23(3H, t, J=7.1Hz), 2.11-2.20(2H, m), 2.50(2H, t, J=7.2Hz), 4.08-4.16(4H, m), 6.85(1H, d, J=8.9), 7.58(1H, dd,
25

$J=8.9, 2.6\text{Hz}$), 7.88(1H, d, $J=2.6\text{Hz}$), 10.37 (1H, s).

IR(neat, cm^{-1}): 1731, 1683, 1590, 1272.

[REFERENCE EXAMPLE 3] Ethyl 4-(2-Formyl-4-methoxy
5 phenoxy)butyrate

Colorless oil, yield 99%. $^1\text{H-NMR}(\text{CDCl}_3, \delta, 300\text{MHz})$:
1.23(3H, t, $J=7.1\text{Hz}$), 2.09-2.18(2H, m), 2.50(2H, t,
 $J=7.2\text{Hz}$), 3.77(3H, s), 4.05-4.15(4H, m), 6.90(1H, d, $J=9.1$),
7.08(1H, dd, $J=9.1, 3.2\text{Hz}$), 7.30(1H, d, $J=3.2\text{Hz}$), 10.43(1H,
10 s). IR(neat, cm^{-1}): 1731, 1683, 1496, 1218.

[REFERENCE EXAMPLE 4] Ethyl 5-(2-Formyl-4-methoxyphenoxy
)pentanoate

Colorless oil, yield 99%. $^1\text{H-NMR}(\text{CDCl}_3, \delta, 300\text{MHz})$:
15 1.21(3H, t, $J=7.1\text{Hz}$), 1.80-1.87(4H, m), 2.37(2H, t,
 $J=7.1\text{Hz}$), 3.77(3H, s), 4.01-4.14(4H, m), 6.89(1H, d,
 $J=9.1\text{Hz}$), 7.08(1H, dd, $J=9.1, 3.2\text{Hz}$), 7.29(1H, d, $J=3.2\text{Hz}$),
10.44 (1H, s). IR(neat, cm^{-1}): 1731, 1683, 1496, 1218.

20 Preparation of 2-formyl-N-methylaniline derivatives in
general

[REFERENCE EXAMPLE 5] 5-(4-Bromo-2-formyl-N-
methylanilino)pentanoic acid

A suspension of 1-methyl-2-piperidone (5.0g) in 4N
25 sodium hydroxide (22.1ml) was heated and stirred for 8.5

hours under reflux. After the suspension was cooled to room temperature, conc. hydrochloric acid (7.4ml) was added. A solution of sodium carbonate (9.4g) and 5-bromo-4-fluorobenzaldehyde (4.5g) in DMSO (74ml) was added, and the resulting mixture was stirred for 1.5 hours under reflux. The mixture was allowed to cool to room temperature, and adjusted to pH of about 3.3 with 6N hydrochloric acid. After extraction with ethyl acetate, the organic layer was washed with saturated salt water, dried over anhydrous sodium sulfate, and concentrated. The precipitated crystal obtained was redissolved in isopropylether [IPE] (20ml) under heating, and the solution was allowed to cool to room temperature, and stirred for 1 hour at 0°C. The crystal obtained was filtered and washed with IPE. Drying under reduced pressure (40°C, 1 hour) gave yellow crystal of 5-(4-bromo-2-formyl-N-methylanilino)pentanoic acid (4.0g, yield 57%).

mp. 72-73°C. $^1\text{H-NMR}$ (CDCl_3 , δ , 300MHz): 1.61-1.69(4H, m), 2.36(2H, t, $J=6.7\text{Hz}$), 2.87(3H, s), 3.13(2H, t, $J=6.9\text{Hz}$), 6.97(1H, d, $J=8.8\text{Hz}$), 7.54(1H, dd, $J=8.8, 2.5\text{Hz}$), 7.86(1H, d, $J=2.5\text{Hz}$), 10.17(1H, s). Anal. Calcd for $\text{C}_{13}\text{H}_{16}\text{NO}_3\text{Br}$: C, 49.70; H, 5.13; N, 4.46; Br, 25.43, Found: C, 49.75; H, 5.16; N, 4.45; Br, 25.40. IR(KBr, cm^{-1}): 1731, 1648.

Compounds below in [REFERENCE EXAMPLE 6] to [Reference EXAMPLE 8] were prepared according to a method similar to

that of [REFERENCE EXAMPLE 5].

[REFERENCE EXAMPLE 6] 5-(2-Formyl-N-methylanilino)pentanoic acid

5 Yellow oil, yield 37%. $^1\text{H-NMR}(\text{CDCl}_3, \delta, 300\text{MHz})$: 1.64-1.73(4H, m), 2.38(2H, t, $J=6.7\text{Hz}$), 2.90(3H, s), 3.17(2H, t, $J=6.9\text{Hz}$), 7.04-7.13(2H, m), 7.50(1H, dd, $J=1.7, 5.5\text{Hz}$), 7.80(1H, dd, $J=1.7, 7.7\text{Hz}$), 10.28(1H, s). IR(neat, cm^{-1}): 1708, 1683.

10

[REFERENCE EXAMPLE 7] 5-(2-Formyl-4-nitro-N-methylanilino)pentanoic acid

Yellow crystal, yield 93%. mp. 107 - 108°C. $^1\text{H-NMR}(\text{CDCl}_3, \delta, 300\text{MHz})$: 1.60-1.67(2H, m), 1.75-1.81(2H, m), 15 2.38(2H, t, $J=6.7\text{Hz}$), 2.90(3H, s), 3.43(2H, t, $J=7.4\text{Hz}$), 6.98(1H, d, $J=9.4$), 8.20(1H, dd, $J=9.4, 2.8\text{Hz}$), 8.58(1H, d, $J=2.8\text{Hz}$), 9.67(1H, s). IR (KBr, cm^{-1}): 1704, 1677, 1598, 1324, 1307.

20 [REFERENCE EXAMPLE 8] 6-(4-Bromo-2-formyl-N-methylanilino)hexanoic acid.

Yellow crystal, yield 80%. mp. 95-96°C. $^1\text{H-NMR}(\text{CDCl}_3, \delta, 300\text{MHz})$: 1.27-1.32(2H, m), 1.54-1.66(4H, m), 2.31(2H, t, $J=7.4\text{Hz}$), 2.83(3H, s), 3.08(2H, t, $J=7.5\text{Hz}$), 6.93(1H, d, 25 $J=8.8\text{Hz}$), 7.50(1H, dd, $J=8.8, 2.5\text{Hz}$), 7.83(1H, d, $J=2.5\text{Hz}$),

10.12(1H, s). Anal. Calcd for $C_{14}H_{18}NO_3Br$: C, 51.23; H, 5.53; N, 4.27; Br, 24.35. Found: C, 51.25; H, 5.54; N, 4.21; Br, 24.48. IR (KBr, cm^{-1}): 1720, 1644.

5 Preparation of 2-formyl-N-benzylaniline derivatives in general

[REFERENCE EXAMPLE 9] 5-(4-Bromo-2-formyl-N-benzylanilino)pentanoic acid

To a solution of benzaldehyde (6.0g) in methanol
10 (80ml), 4-aminobutyric acid (6.6g) and 1N sodium hydroxide (56.5ml) were added. 20% Pd-C(wet, 0.6g) was added to the reaction apparatus which was previously purged with an argon gas. Subsequently, the apparatus was filled with a hydrogen gas, and the solution was stirred at room
15 temperature overnight. Pd-C was filtered off and washed with methanol. To the filtrate was added 6N hydrochloric acid (9.4ml), and the solvents were evaporated. Subsequently, sodium carbonate (12.0g) and DMSO/water (57ml/37ml) were added to the residue, and the resulting
20 mixture was heated under reflux with dropwise addition of a solution of 5-bromo-4-fluorobenzaldehyde (5.7g) in DMSO (17ml), and stirred further for 5 hours. After cooled to room temperature, the solution was adjusted to pH of about 3.5 by addition of 6N hydrochloric acid. After extraction
25 with ethyl acetate, the organic layer was washed with

saturated salt water, dried over anhydrous sodium sulfate, and concentrated. The concentrate was purified by silica gel column chromatography (n-hexane/ethyl acetate=4/1) and subsequent concentration of the appropriate eluate gave
 5 yellow oil of 5-(4-bromo-2-formyl-N-benzylanilino)pentanoic acid (7.4g, yield 61%).

$^1\text{H-NMR}$ (CDCl_3 , δ , 300MHz): 1.52-1.65(4H, m), 2.27-2.35(2H, m), 3.08-3.15(2H, m), 4.31(2H, s), 7.01(1H, d, $J=8.7\text{Hz}$), 7.16-7.36(5H, m), 7.57(1H, dd, $J=8.7$, 2.4Hz), 7.92(1H, d, $J=2.4\text{Hz}$), 10.36(1H, s). EI-MS(M^+): 391.
 10 IR(neat, cm^{-1}): 1708, 1683.

A compound below in [REFERENCE EXAMPLE 10] was prepared according to a method similar to that of [REFERENCE EXAMPLE 9].

15

[REFERENCE EXAMPLE 10] 5-(4-Bromo-2-formyl-N-benzylanilino)heptanoic acid

Yellow oil, yield 27%. $^1\text{H-NMR}$ (CDCl_3 , δ , 300MHz): 1.21-1.26(4H, m), 1.46-1.56(4H, m), 2.27(2H, t, $J=7.4\text{Hz}$), 3.05(2H, t, $J=7.5\text{Hz}$), 4.26(2H, s), 6.97(1H, d, $J=8.7\text{Hz}$),
 20 7.13-7.28(5H, m), 7.51(1H, dd, $J=8.7$, 2.5Hz), 7.87(1H, d, $J=2.5\text{Hz}$), 10.30(1H, s). IR (neat, cm^{-1}): 1706, 1683.

Cyclization reactions in general (method A)

25 [EXAMPLE 1] 2,3-Dihydro-1-benzoxepin-4-carboxylic acid

ethylester

To a solution of 4-(2-formylphenoxy)butyric acid ethylester (2.4g) in diethyl carbonate (24ml), a solution of 20% sodium ethoxide in ethanol (4.1g) was added at room temperature, and the solution was stirred at 50°C for 1 hour. The solution was allowed to cool to room temperature, and neutralized with 1N hydrochloric acid. After ethyl acetate extraction, the organic layer was washed with saturated salt water, dried over anhydrous sodium sulfate, and concentrated. The concentrate was purified by silica gel column chromatography (n-hexane/ethyl acetate=10/1) and concentration of the appropriate eluate gave colorless oil of 2,3-dihydro-1-benzoxepin-4-carboxylic acid ethylester (1.3g, yield 61%).

¹H-NMR(CDCl₃, δ, 300MHz): 1.36(3H, t, J=7.1Hz), 2.97-3.00(2H, m), 4.26-4.31(4H, m), 6.96-7.04(2H, m), 7.21-7.27(1H, m), 7.32-7.35(1H, m), 7.58(1H, s). IR (neat, cm⁻¹): 1700, 1249.

[REFERENCE EXAMPLE 11] 2,3-Dihydro-1-benzoxepin-4-carboxylic acid ethylester

To a solution of 4-(2-formylphenoxy)butyric acid ethylester (2.4g) in THF (24ml), potassium t-butoxide (1.2g) was added at room temperature, and the mixture was stirred at 50°C for 2 hours. The solution was allowed to

cool to room temperature, and neutralized with 1N hydrochloric acid. After ethyl acetate extraction, the organic layer was washed with saturated salt water. The organic layer was subjected to HPLC analysis.

5 The results of quantitative analyses: 4-(2-formylphenoxy)butyric acid ethylester (0.1g, yield 4%), 4-(2-formylphenoxy)butyric acid (0.3g, yield 14%), 2,3-dihydro-1-benzoxepin-4-carboxylic acid ethylester (0.4g, yield 17%), 2,3-dihydro-1-benzoxepin-4-carboxylic acid
10 (0.3g, yield 15%).

Compounds below in [EXAMPLE 2] to [EXAMPLE 4] were prepared according to a method similar to that of [EXAMPLE 1].

15 [EXAMPLE 2] 7-Bromo-2,3-dihydro-1-benzoxepin-4-carboxylic acid ethylester (method A)

White crystal, yield 34%. mp. 85-86°C. ¹H-NMR(CDCl₃, δ, 300MHz): 1.34(3H, t, J=7.1Hz), 2.92-2.97(2H, m), 4.19-4.25(4H, m), 6.82(1H, d, J=8.6Hz), 7.28(1H, dd, J=8.6,
20 2.4Hz), 7.42(1H, d, J=2.4Hz), 7.43(1H, s).

[EXAMPLE 3] 7-Methoxy-2,3-dihydro-1-benzoxepin-4-carboxylic acid ethylester (method A)

White crystal, yield 82%. mp. 62-63°C. ¹H-NMR(CDCl₃, δ, 300MHz): 1.32(3H, t, J=7.1Hz), 2.92-2.95(2H, m), 3.76(3H,
25

s), 4.17-4.28 (4H, m), 6.77-6.81 (2H, m), 6.87 (1H, dd, $J=1.6$, 7.4Hz), 7.49 (1H, s).

[EXAMPLE 4] 8-Methoxy-3,4-dihydro-2H-1-benzoxocin-5-
5 carboxylic acid ethylester (method A)

Colorless oil, yield 11%. $^1\text{H-NMR}$ (CDCl_3 , δ , 300MHz):
1.44 (3H, t, $J=7.1\text{Hz}$), 1.82-1.88 (2H, m), 2.66-2.70 (2H, m),
3.86 (3H, s), 4.24-4.39 (4H, m), 6.77 (1H, d, $J=3.0\text{Hz}$),
6.92 (1H, dd, $J=3.0$, 8.9Hz), 7.02 (1H, d, $J=8.9\text{Hz}$), 7.70 (1H,
10 s). IR (neat, cm^{-1}): 1704, 1496, 1243.

Cyclization reactions in general (Method B)

[EXAMPLE 5] 8-Bromo-1-methyl-1,2,3,4-tetrahydro-1-
15 benzazocin-5-carboxylic acid methylester

To a solution of 5-(4-bromo-2-formyl-N-methylanilino)pentanoic acid (3.0g) in DMF (8ml), potassium carbonate (1.6g) and a solution of methyl iodide/DMF (1.6g/2ml) were added at room temperature, and the
20 resulting mixture was stirred for 2 hours. Ethyl acetate and water were added to the mixture, and the organic layer was separated, washed with water, dried over anhydrous sodium sulfate, and concentrated to give yellow oil of 5-(4-bromo-2-formyl -N-methylanilino)pentanoic acid
25 methylester (3.0g, yield 97%).

$^1\text{H-NMR}$ (CDCl_3 , δ , 300MHz): 1.58-1.64 (4H, m), 2.28 (2H, t, $J=6.8\text{Hz}$), 2.83 (3H, s), 3.09 (2H, t, $J=6.8\text{Hz}$), 3.63 (3H, s), 6.95 (1H, d, $J=8.8\text{Hz}$), 7.51 (1H, dd, $J=8.8, 2.4\text{Hz}$), 7.83 (1H, d, $J=2.4\text{Hz}$), 10.14 (1H, s). FAB-MS ($[\text{M}+\text{Na}]^+$): 350. IR (neat, 5 cm^{-1}): 1735, 1681.

To a solution of 5-(4-bromo-2-formyl-N-methylanilino)pentanoic acid methylester (3.0g) in dimethyl carbonate (40ml), a solution of 28% sodium methoxide in methanol (2.3g) was added at room temperature, and the 10 resulting mixture was stirred at 50°C for 2 hours. The mixture was allowed to cool to room temperature, and neutralized with 1N hydrochloric acid. After ethyl acetate extraction, the organic layer was washed with saturated salt water, dried over anhydrous sodium sulfate, and 15 concentrated. The concentrate was purified by silica gel column chromatography (h-hexane/ethyl acetate=10/1) and the appropriate eluate was concentrated. The precipitated crystal was washed with IPE and collected by filtration. Drying under reduced pressure (40°C , 1 hour) gave yellow 20 crystal of 8-bromo-1-methyl-1,2,3,4-tetrahydro-1-benzazocin-5-carboxylic acid methylester (0.9g, yield 31%). mp. $88-89^\circ\text{C}$. $^1\text{H-NMR}$ (CDCl_3 , δ , 300MHz): 1.44 (2H, m), 2.54 (2H, t, $J=6.2\text{Hz}$), 2.87 (3H, s), 3.42 (2H, t, $J=5.6\text{Hz}$), 3.78 (3H, s), 6.56 (1H, d, $J=6.0\text{Hz}$), 7.19-7.23 (2H, m), 25 7.68 (1H, s). Anal. Calcd for $\text{C}_{14}\text{H}_{16}\text{NO}_2\text{Br}$: C, 54.21;

H, 5.20; N, 4.52; Br, 25.76. Found: C, 54.19; H, 5.29; N, 4.37; Br, 25.74. IR (KBr, cm^{-1}): 1689, 1189.

Compounds below in [EXAMPLE 6] to [EXAMPLE 9] were prepared according to a method similar to that of [EXAMPLE 5].

[EXAMPLE 6] 1-Methyl-1,2,3,4-tetrahydro-1-benzazocin-5-carboxylic acid methylester (method B)

5-(2-Formyl-N-methylanilino)pentanoic acid methylester

10 Yellow oil, yield 99%. $^1\text{H-NMR}$ (CDCl_3 , δ , 300MHz): 1.58-1.68 (4H, m), 2.30 (2H, t, $J=6.8$), 2.86 (3H, s), 3.12 (2H, t, $J=6.8\text{Hz}$), 3.64 (3H, s), 7.01-7.10 (2H, m), 7.46 (1H, dt, $J=1.7$, 7.3Hz), 7.76 (1H, dd, $J=1.7$, 7.7Hz), 10.25 (1H, s). IR (neat, cm^{-1}): 1735, 1685, 1159.

15 1-methyl-1,2,3,4-tetrahydro-1-benzazocin-5-carboxylic acid methylester

Yellow crystal, yield 30%. mp. 80-82°C. $^1\text{H-NMR}$ (CDCl_3 , δ , 300MHz): 1.42-1.50 (2H, m), 2.57 (2H, t, $J=6.2\text{Hz}$), 2.90 (3H, s), 3.45 (2H, t, $J=5.4\text{Hz}$), 3.79 (3H, s), 6.62-6.73 (2H, m), 20 7.11-7.22 (2H, m), 7.80 (1H, s). Anal. Calcd for $\text{C}_{14}\text{H}_{17}\text{NO}_2$: C, 72.70; H, 7.41; N, 6.06. Found: C, 73.00; H, 7.65; N, 6.06. IR (KBr, cm^{-1}): 1685, 1189.

[EXAMPLE 7] 1-Benzyl-8-bromo-1,2,3,4-tetrahydro-1-benzazocin-5-carboxylic acid methylester (method B)

25

5-(4-Bromo-2-formyl-N-benzylanilino)pentanoic acid
methylester

Yellow oil, yield 100%. $^1\text{H-NMR}(\text{CDCl}_3, \delta, 300\text{MHz})$:
1.57-1.60(4H, m), 2.27(2H, t, $J=6.7$), 3.12(2H, t, $J=6.6\text{Hz}$),
5 3.66(3H, s), 4.31(2H, s), 7.01(1H, d, $J=8.7$), 7.17-7.33(5H,
m), 7.56(1H, dd, $J=8.7, 2.5\text{Hz}$), 7.92(1H, d, $J=2.5\text{Hz}$),
10.35 (1H, s). EI-MS(M^+): 403. IR (neat, cm^{-1}): 1735, 1683.

1-Benzyl-8-bromo-1,2,3,4-tetrahydro-1-benzazocin-5-
carboxylic acid methylester

10 Yellow oil, yield 76%. $^1\text{H-NMR}(\text{CDCl}_3, \delta, 300\text{MHz})$:
1.44-1.50(2H, m), 2.59(2H, t, $J=6.1\text{Hz}$), 3.50(2H, t,
 $J=5.3\text{Hz}$), 3.81(3H, s), 4.45(2H, s), 6.50(1H, d, $J=9.0$),
7.09-7.39(7H, m), 7.75(1H, s). EI-MS(M^+): 385.
IR(neat, cm^{-1}): 1704, 1495, 1087.

15

[EXAMPLE 8] 9-Bromo-1-methyl-2,3,4,5-tetrahydro-1H-1-
benzazonin-6-carboxylic acid methylester (method B)

6-(4-Bromo-2-formyl-N-methylanilino) hexanoic acid
methylester

20 Yellow oil, yield 98%. $^1\text{H-NMR}(\text{CDCl}_3, \delta, 300\text{MHz})$:
1.20-1.30(2H, m), 1.51-1.61(4H, m), 2.24(2H, t, $J=7,4$),
2.81(3H, s), 3.05(2H, t, $J=6.6\text{Hz}$), 3.61(3H, s), 6.91(1H, d,
 $J=8.8\text{Hz}$), 7.47(1H, dd, $J=8.8, 2.3\text{Hz}$), 7.80(1H, d, $J=2.3\text{Hz}$),
10.10(1H, s). FAB-MS($[\text{M}+\text{Na}]^+$): 364. IR(neat, cm^{-1}): 1735,
25 1681.

9-Bromo-1-methyl-2,3,4,5-tetrahydro-1H-1-benzazonin-6-carboxylic acid methylester

Yellow oil, yield 18%. $^1\text{H-NMR}$ -(CDCl_3 , δ , 300MHz):
1.67-1.77(2H, m), 1.87-1.96(2H, m), 2.30(2H, t, $J=6.9\text{Hz}$),
5 2.76(3H, s), 3.12(2H, t, $J=6.9\text{Hz}$), 3.77(3H, s), 6.70(1H, d,
 $J=6.4\text{Hz}$), 7.19-7.22(2H, m), 7.60(1H, s). EI-MS(M^+):323.
IR(neat, cm^{-1}): 1708, 1494.

Cyclization reactions in general (method C)

10 [EXAMPLE 9] 8-Nitro-1-methyl-1,2,3,4-tetrahydro-1-benzazocin-5-carboxylic acid methylester

To a solution of 5-(2-formyl-4-nitro-N-methylanilino)pentanoic acid (3.0g) in DMF (8.1ml),
potassium carbonate (1.6g) and a solution of methyl
15 iodide/DMF (1.8g/1ml) were added at room temperature, and
the resulting mixture was stirred further for 2 hours. To
the mixture, dimethyl carbonate (18ml) and subsequently a
solution of 28% sodium methoxide in methanol (5.0g) were
added and the resulting mixture was stirred at 50°C for 2.5
20 hours. The mixture was allowed to cool to room temperature,
and neutralized with 1N hydrochloric acid. After ethyl
acetate extraction, the organic layer was washed with
saturated salt water, dried over anhydrous sodium sulfate,
and concentrated. The concentrate was redissolved in hot
25 isopropyl alcohol [IPA], and the solution was allowed to

cool to room temperature, stirred at room temperature further for 1 hour, and at ice temperature for 0.5 hours. The precipitated crystal was collected by filtration, and washed with IPA. The crystal was purified by silica gel column chromatography (n-hexane/ethyl acetate=2/1 to 1/1), and subsequent concentration of the appropriate eluate gave yellow crystal of 8-nitro-1-methyl-1,2,3,4-tetrahydro-1-benzazocin-5-carboxylic acid methylester (2.3g, yield 78%). mp. 127-129°C. $^1\text{H-NMR}$ (CDCl_3 , δ , 300MHz): 1.38-1.72(2H, m), 2.35-2.80(2H, m), 3.02(3H, s), 3.42-3.85(2H, m), 3.80(3H, s), 6.65(1H, d, $J=9.4\text{Hz}$), 7.80(1H, s), 7.99-8.06(2H, m). Anal. Calcd for $\text{C}_{14}\text{H}_{16}\text{N}_2\text{O}_2$: C, 60.86; H, 5.84; N, 10.14. Found: C, 60.79; H, 5.81; N, 9.98. IR (KBr, cm^{-1}): 1708, 1255, 1191.

A compound below in [EXAMPLE 10] was prepared according to a method similar to that of [EXAMPLE 9].

[EXAMPLE 10] 1-Benzyl-10-bromo-1,2,3,4,5,6-hexahydro-1-benzazecin-7-carboxylic acid methylester (method C)

Yellow oil, yield 6%. $^1\text{H-NMR}$ (CDCl_3 , δ , 300MHz): 1.24-1.29(2H, m), 1.44-1.50(2H, m), 1.57-1.63(2H, m), 2.13(2H, t, $J=6.5\text{Hz}$), 2.89(2H, t, $J=6.2\text{Hz}$), 3.48(3H, s), 3.90(2H, s), 7.02(1H, d, $J=8.6\text{Hz}$), 7.10-7.25(6H, m), 7.37(1H, dd, $J=8.6, 2.4\text{Hz}$), 7.48(1H, s). IR(KBr, cm^{-1}): 1712, 1279, 1232.

[REFERENCE EXAMPLE 11]

Preparation of 4'-ethoxy-4-hydroxy-1,1'-biphenyl-3-carbaldehyde

To a suspension of magnesium (0.5g) in THF (7ml), a
5 solution of 1-bromo-4-ethoxybenzene (4.0g) in THF (2.5ml)
was added slowly at room temperature under an argon
atmosphere, and the resulting solution was stirred for 1
hour. Subsequently, a solution of trimethoxyborane (2.1g)
in THF (2.5ml) was added dropwise at -10 to 10°C, and after
10 further addition of THF (7ml), the reaction mixture was
stirred at the same temperature for 1 hour. The reaction
mixture was allowed to warm to room temperature and
tetrakis(triphenylphosphine) palladium (115mg), 5-bromo-2-
hydroxybenzaldehyde (2.0g) and an aqueous solution of
15 tripotassium phosphate (11.1g) in water (15ml) were added,
and the resulting mixture was stirred under reflux for 1
hour. After the mixture was allowed to cool to room
temperature, 6N hydrochloric acid (20ml) was added dropwise,
and then toluene (20ml) was added. After separation, the
20 aqueous phase was further extracted with toluene (30ml).
The organic layers were combined and washed with aqueous
10% sodium chloride solutions (10ml×3), dried over
anhydrous magnesium sulfate and concentrated. The
concentrate was purified by silica gel chromatography (n-
25 hexane/ethyl acetate=6/1), and the appropriate eluate was

concentrated. Diisopropylether (15ml) was added to the concentrate and the resulting mixture was heated under reflux, and then allowed to cool to room temperature. n-Hexane (15ml) was added to the solution, and stirred at room temperature for 1 hour and at 0°C for 0.5 hour. The precipitated crystal was collected by filtration and washed with diisopropylether/n-hexane (1/1.4ml) which was previously cooled to 0 to 5°C. Drying under reduced pressure (room temperature, 2 hours) gave yellow crystal of the title compound (1.8g, yield 73%).

mp. 94-95°C.

Anal Calcd. for $C_{15}H_{14}O_3$: C, 74.36; H, 5.82. Found: C, 74.61; H, 5.80.

IR(KBr, cm^{-1}): 1660, 1473, 1276, 1245, 1047, 831.

1H -NMR($CDCl_3$, 300MHz) δ 1.44(3H, t, $J=7.0$ Hz), 4.07(2H, q, $J=7.0$ Hz), 6.96(2H, dd, $J=1.9, 6.7$ Hz), 7.04(1H, d, $J=8.4$ Hz), 7.45(2H, dd, $J=1.9, 6.7$ Hz), 7.68-7.77(2H, m), 9.95(1H, s), 10.95(1H, s).

[REFERENCE EXAMPLE 12]

Preparation of 4-[(4'-ethoxy-3-formyl-1,1'-biphenyl-4-yl)oxy]butyric acid ethylester

To a solution of 4'-ethoxy-4-hydroxy-1,1'-biphenyl-3-carbaldehyde(1.0g), and 4-bromobutyric acid ethylester (0.9g) in DMF (5ml) was added potassium carbonate (1.1g) at

room temperature, and the solution was stirred at room temperature for 14 hours, and at 50°C for 3 hours. After the solution was allowed to cool to room temperature, water (10ml) was added, and extracted with ethyl acetate (30ml).

5 The organic layer was washed with water (10ml×3), and concentrated. To the concentrate was added diisopropylether (14ml), and the resulting mixture was heated under reflux and subsequently cooled to room temperature. The solution was further stirred at room
10 temperature for 1 hour, and at 0°C for 1 hour. The precipitated crystal was collected by filtration and washed with diisopropylether (3ml) which was previously cooled to 0 to 5°C. Drying under vacuum (room temperature, 2 hours) gave white crystal of the title compound (1.3g, yield 91%).
15 mp. 69-70°C.

Anal Calcd. for $C_{21}H_{24}O_5$: C, 70.77; H, 6.79. Found: C, 70.91; H, 7.08.

IR (KBr, cm^{-1}) 1683, 1606, 1471, 1270, 1240, 1187, 1047.

1H -NMR($CDCl_3$, 300MHz) δ 1.26(3H, t, $J=7.1Hz$), 1.43(3H, t, $J=7.0Hz$), 2.18-2.25(2H, m), 2.56(2H, t, $J=7.1Hz$), 4.04-4.20(6H, m), 6.94(2H, dd, $J=1.9, 6.7Hz$), 7.03(1H, d, $J=8.7Hz$), 7.47(2H, dd, $J=1.9, 6.7Hz$), 7.72(1H, dd, $J=8.7, 2.5Hz$), 8.02(1H, d, $J=2.5Hz$), 10.53(1H, s).

20

25 [EXAMPLE 11]

7-(4-Ethoxyphenyl)-2,3-dihydro-1-benzoxepin-4-carboxylic acid ethylester

To a solution of 4-[(4'-ethoxy-3-formyl-1,1'-biphenyl-4-yl)oxy]butyric acid ethylester (0.50g) in diethyl carbonate (5ml) was added a solution of 20% sodium ethoxide in ethanol (0.57g) at room temperature, and the mixture was then heated to 50°C and stirred for 1 hour. The solution was neutralized with 1N hydrochloric acid, and extracted with ethyl acetate. The organic layer was washed, and concentrated. The concentrate was purified by silica gel chromatography (n-hexane/ethyl acetate=10/1) and subsequent concentration of the appropriate eluate gave white crystal of the title compound (0.39g, yield 82%).

mp. 128-129°C. Anal Calcd. for $C_{21}H_{22}O_4$: C, 74.54; H, 6.55.

Found: C, 74.32; H, 6.46.

IR(KBr, cm^{-1}), 1702, 1496, 1251, 1213.


1H -NMR($CDCl_3$, 300MHz) δ 1.37(3H, t, $J=7.1Hz$), 1.44(3H, t, $J=7.0Hz$), 2.99-3.02(2H, m), 4.07(2H, q, $J=7.0Hz$), 4.26-4.32(4H, m), 6.95(2H, dd, $J=1.9, 6.7Hz$), 7.00(1H, d, $J=8.4Hz$), 7.41-7.51(4H, m), 7.65(1H, s).

Industrial Applicability

The present invention provides a process suitable for mass production wherein cyclic compounds can be produced safely through a short step.


CLAIMS

1. A process for producing a compound represented by the formula:



5

[wherein, Z is an electron-withdrawing group; W is an optionally substituted ethylene or an optionally substituted vinylene, and when the ethylene or the vinylene has two substituent groups, said substituent groups may be bound to each other to form an optionally substituted ring; R³ is a hydrogen atom or an optionally substituted hydrocarbon group; and X is a divalent group (however, when W is an optionally substituted vinylene, -X-CH₂-Z is not -X¹-X²-CH₂-Z (wherein, X¹ is a sulfur atom or an optionally substituted nitrogen atom; X² is an optionally substituted ethylene))] or a salt thereof, which comprises subjecting a compound represented by the formula:



15

[wherein, each symbol has the same meaning described as above] or a salt thereof to a cyclization reaction in a solvent containing a carbonic diester.

2. The process according to claim 1, wherein X is a
5 divalent group having 1 to 12 atoms in the straight-chain moiety thereof.


3. The process according to claim 1, wherein Z is a carboxyl group which is esterified.

4. A process according to claim 1, wherein R³ is a
10 hydrogen atom.

5. A process according to claim 1, which comprises the reaction being carried out in the presence of a base.


6. A process according to claim 5, wherein the base is an alcoholate.

7. A process for producing a compound represented by
15 the formula:



[wherein, z is an electron-withdrawing group; R¹ and R²
each are a hydrogen atom, a halogen atom, an optionally
20 substituted amino group, an optionally substituted hydroxy
group, an optionally substituted thiol group, an optionally
substituted hydrocarbon group or an optionally substituted

heterocyclic group; R^1 and R^2 may be bound to each other to form an optionally substituted ring; R^3 is a hydrogen atom or an optionally substituted hydrocarbon group; the combined line of a broken line and a solid line is a single bond or a double bond; and X is a divalent group (however, when the combined line of a broken line and a solid line is a double bond, $-X-CH_2-Z$ is not $-X^1-X^2-CH_2-Z$ (wherein, X^1 is a sulfur atom or an optionally substituted nitrogen atom, X^2 is an optionally substituted ethylene))) or a salt thereof, which comprises subjecting a compound represented by the formula:



[wherein, each symbol has the same meaning described as above] or a salt thereof to a cyclization reaction in a solvent containing a carbonic diester.

8. The process according to claim 7, wherein X is a divalent group having 1 to 12 atoms in the straight-chain moiety thereof.


9. The process according to claim 7, wherein Z is a carboxyl group which is esterified.

10. The process according to claim 7, wherein R^3 is a hydrogen atom.


11. The process according to claim 7, which comprises the reaction being carried out in the presence of a base.

12. The process according to claim 11, wherein the base is an alcoholate.

5 13. A process for producing a compound represented by the formula:



[wherein, Z is an electron-withdrawing group; R³ is a hydrogen atom or an optionally substituted hydrocarbon group; the combined line of a broken line and a solid line is a single bond or a double bond; ring A is an optionally substituted ring; and X is a divalent group (however, when the ring A is an optionally substituted benzene ring and the combined line of a broken line and a solid line is a double bond, -X-CH₂-Z is not -X¹-X²-CH₂-Z (wherein, X¹ is sulfur atom or an optionally substituted nitrogen atom ; X² is an optionally substituted ethylene))] or a salt thereof, which comprises subjecting a compound represented by the formula:



[wherein, each symbol has the same meaning described as above] or a salt thereof to a cyclization reaction in a solvent containing a carbonic diester.

14. The process according to claim 13, wherein X is a
5 divalent group having 1 to 12 atoms in the straight-chain moiety thereof.

15. The process according to claim 13, wherein Z is a carboxyl group which is esterified.


16. The process according to claim 13, wherein R³ is
10 a hydrogen atom.

17. The process according to claim 13, which comprises the reaction being carried out in the presence of a base.

18. The process according to claim 17, wherein the
15 base is an alcoholate.

19. A compound represented by the formula:

[formula]



[wherein, Z is an electron-withdrawing group; W is an
20 optionally substituted ethylene or an optionally substituted vinylene and when the ethylene or the vinylene has two substituent groups, said substituent groups may be

bound to each other to form an optionally substituted ring;
R³ is a hydrogen atom or an optionally substituted
hydrocarbon group; and X'' is a divalent group having 4 or
more atoms in the straight-chain moiety thereof] or a salt
5 thereof.