
PSMA AND CD3 BISPECIFIC T CELL ENGAGING ANTIBODY CONSTRUCTS

BACKGROUND

[1] Bispecific antibody derived molecules such as BiTE[®] (bispecific T cell engager) antibody constructs are recombinant protein constructs made from two flexibly linked antibody derived binding domains. One binding domain of BiTE[®] antibody constructs is specific for a selected tumor-associated surface antigen on target cells; the second binding domain is specific for CD3, a subunit of the T cell receptor complex on T cells. By their particular design, BiTE[®] antibody constructs are uniquely suited to transiently connect T cells with target cells and, at the same time, potentially activate the inherent cytolytic potential of T cells against target cells. An important further development of the first generation of BiTE[®] antibody constructs (see WO 99/54440 and WO 2005/040220) developed into the clinic as AMG 103 and AMG 110 was the provision of bispecific antibody constructs binding to a context independent epitope at the N-terminus of the CD3 ϵ chain (WO 2008/119567). BiTE[®] antibody constructs binding to this elected epitope do not only show cross-species specificity for the human and the *Callithrix jacchus*, *Saguinus oedipus* or *Saimiri sciureus* CD3 ϵ chain, but also, due to recognizing this specific epitope (instead of previously described epitopes of CD3 binders in bispecific T cell engaging molecules), do not unspecifically activate T cells to the same degree as observed for the previous generation of T cell engaging antibodies. This reduction in T cell activation was connected is correlated with less or reduced T cell redistribution in patients, the latter being identified as a risk for side effects.

[2] Antibody constructs as described in WO 2008/119567 are likely to suffer from rapid clearance from the body; thus, while they are able to reach most parts of the body rapidly, and are quick to produce and easier to handle, their in vivo applications may be limited by their brief persistence in vivo. Prolonged administration by continuous intravenous infusion was used to achieve therapeutic effects because of the short in vivo half-life of this small, single chain molecule. However, such continuous intravenous infusions are classified as inconvenient for the patients and, thus, in case of more convenient alternative treatment approaches, hamper the election of the compound demonstrated to be more efficient in the treatment of the respective disease. Hence, there is a need in the art for bispecific therapeutics that retain similar therapeutic efficacy, that have a format which is straightforward to produce, and that have favorable pharmacokinetic properties, including a longer half-life.

[3] An increased half-life is generally useful in *in vivo* applications of immunoglobulins, especially antibodies and most especially antibody fragments of small size. Approaches described in the art to achieve such effect comprise the fusion of the small bispecific antibody construct to larger proteins, which preferably do not interfere with the therapeutic effect of the BiTE[®]. Examples for such further developments of bispecific T cell engagers comprise bispecific Fc-molecules e.g. described in US 2014/0302037, US 2014/0308285, WO 2014/144722, WO 2014/151910 and WO 2015/048272. An alternative strategy is the use of HSA fused to the bispecific molecule or the mere fusion of human albumin binding peptides (see e.g. WO2013/128027, WO2014/140358).

[4] Several markers for prostate cancer have been identified, including e.g. prostate-specific antigen (PSA), the six-transmembrane epithelial antigen of the prostate (STEAP) (Hubert et al., PNAS 96 (1999), 14523-14528), the prostate stem cell antigen (PSCA) (Reiter et al., Proc. Nat. Acad. Sci. 95: 1735-1740, 1998) and the prostate-specific membrane antigen (PSMA; PSM) (Israeli et al., Cancer Res. 53 (1993). PSMA was originally defined by the monoclonal antibody (MAb) 7E11 derived from immunization with a partially purified membrane preparation from the lymph node prostatic adenocarcinoma (LNCaP) cell line (Horoszewicz et al., Anticancer Res. 7 (1987), 927-35). A 2.65-kb cDNA fragment encoding the PSMA protein was cloned and subsequently mapped to chromosome 11p11.2 (Israeli et al., loc. cit.; O'Keefe et al., Biochem. Biophys. Acta 1443 (1998), 113-127). Initial analysis of PSMA demonstrated widespread expression within the cells of the prostatic secretory epithelium. Immunohistochemical staining demonstrated that PSMA was absent to moderately expressed in hyperplastic and benign tissues, while malignant tissues stained with the greatest intensity (Horoszewicz et al., loc. cit.). Subsequent investigations have recapitulated these results and evinced PSMA expression as a universal feature in practically every prostatic tissue examined to date. These reports further demonstrate that expression of PSMA increases precipitously proportional to tumor aggressiveness (Burger et al., Int. J. Cancer 100 (2002), 228-237; Chang et al., Cancer Res. 59 (1999), 3192-98; Chang et al., Urology 57 (2001), 1179-83), Kawakami and Nakayama, Cancer Res. 57 (1997), 2321-24; Liu et al., Cancer Res. 57 (1997), 3629-34; Lopes et al., Cancer Res. 50 (1990), 6423-29; Silver et al., Clin. Cancer Res. 9 (2003), 6357-62; Sweat et al., Urology 52 (1998), 637-40; Troyer et al., Int. J. Cancer 62 (1995), 552-558; Wright et al., Urology 48 (1996), 326-334). Consistent with the correlation between PSMA expression and tumor stage, increased levels of PSMA are associated with androgen-independent prostate cancer (PCa). Analysis of tissue samples from patients with prostate cancer has demonstrated elevated PSMA levels after physical castration or androgen-deprivation therapy. Unlike expression of prostate specific antigen, which is downregulated after androgen ablation, PSMA expression is significantly increased in both primary and

metastatic tumor specimens (Kawakami et al., Wright et al., loc. cit.). Consistent with the elevated expression in androgen-independent tumors, PSMA transcription is also known to be downregulated by steroids, and administration of testosterone mediates a dramatic reduction in PSMA protein and mRNA levels (Israeli et al., Cancer Res. 54 (1994), 1807-11; Wright et al., loc. cit.). PSMA is also highly expressed in secondary prostatic tumors and occult metastatic disease. Immunohistochemical analysis has revealed relatively intense and homogeneous expression of PSMA within metastatic lesions localized to lymph nodes, bone, soft tissue, and lungs compared with benign prostatic tissues (Chang et al. (2001), loc. cit.; Murphy et al., Cancer 78 (1996), 809-818; Sweat et al., loc. cit.). Some reports have also indicated limited PSMA expression in extraprostatic tissues, including a subset of renal proximal tubules, some cells of the intestinal brush-border membrane, and rare cells in the colonic crypts (Chang et al. (1999), Horoszewicz et al., Israeli et al. (1994), Lopes et al., Troyer et al., loc. cit.). However, the levels of PSMA in these tissues are generally two to three orders of magnitude less than those observed in the prostate (Sokoloff et al., Prostate 43 (2000), 150-157). PSMA is also expressed in the tumor-associated neovasculature of most solid cancers examined yet is absent in the normal vascular endothelium (Chang et al. (1999), Liu et al., Silver et al., loc. cit.). Although the significance of PSMA expression within the vasculature is unknown, the specificity for tumor-associated endothelium makes PSMA a potential target for the treatment of many forms of malignancy.

SUMMARY

[5] All the half-life extending formats (HLE formats) of bispecific T cell engaging molecules described in the art, which included the hetero Fc (also designated as heterodimeric Fc, hetFc or hFc) format and the fusion of human serum albumin (also designated as HSA or hALB) had individual disadvantages such as unspecific T cell activation, complement activation, instability, or a pharmacokinetic profile which does not meet the desired half-life prolongation of the molecules. It is thus the object of the present invention to provide a half-life extending format of bispecific T cell engaging molecules, which overcomes at least one and, of course, preferably more than one of these individual defects observed for the state of the art molecules. Accordingly, the present invention provides antibody constructs of a specific Fc modality characterized by comprising a first domain binding to PSMA, a second domain binding to an extracellular epitope of the human and the *Macaca* CD3 ϵ chain, and a third domain, which is the specific Fc modality. Moreover, the invention provides a polynucleotide encoding the antibody construct, a vector comprising this polynucleotide, host cells expressing the construct and a pharmaceutical composition comprising the same.

DESCRIPTION OF THE FIGURES

[6] Figure 1: FIG. 1a shows a diagram of one embodiment of an antibody construct of the invention. Fig. 1b shows a heterodimeric Fc antibody construct and Fig. 1c shows a X-body construct described in the art. The indicated charged pairs are introduced in order to enforce the heterodimerization. Fig 1d shows the fusion of an antibody construct with a human serum albumin (HSA / hALB).

[7] Figure 2: Evaluation of Target-independent T Cell Activation by Target A HLE BiTE[®] antibody constructs. 2(a) antibody construct of the invention in a 48 h activation assay with human PBMC (3x); HLE BiTE[®] serial dilutions (start 20 nM; 1:5, 7x+blank); w/o or with FcR blocking [10 mg/mL hulgG (Kiovog, Baxter)]; FACS measurement of CD69 and CD25 [not shown] expression on CD4⁺, CD8⁺ T cells. 2(b) Hetero-Fc antibody construct in 48 h activation assay with human PBMC and CD14⁺/CD33⁺ cell depleted PBMC (3x); HLE BiTE[®] serial dilutions (start 20 nM; 1:5, 7x+blank); FACS measurement of CD69 and CD25 [not shown] expression on CD4⁺, CD8⁺ T cells.

[8] Figure 3: Evaluation of Target-independent T Cell Activation by Target B HLE BiTE[®] antibody constructs. 3(a) antibody construct of the invention in a 48 h activation assay with human PBMC (3x); HLE BiTE[®] serial dilutions (start 20 nM; 1:5, 7x+blank); w/o or with FcR blocking [10 mg/mL hulgG (Kiovog, Baxter)]; FACS measurement of CD69 and CD25 [not shown] expression on CD4⁺, CD8⁺ T cells. 3(b) Hetero-Fc antibody construct in a 48 h activation assay with human PBMC and CD14⁺/CD33⁺ cell depleted PBMC (3x); HLE BiTE[®] serial dilutions (start 20 nM; 1:5, 7x+blank); FACS measurement of CD69 and CD25 [not shown] expression on CD4⁺, CD8⁺ T cells. 3(c) X-body construct in a 48 h activation assay with human PBMC and CD14⁺/CD33⁺ cell depleted PBMC (3x); HLE BiTE[®] serial dilutions (start 20 nM; 1:5, 7x+blank); FACS measurement of CD69 and CD25 [not shown] expression on CD4⁺, CD8⁺ T cells. 3(d) – 3(f) Isolated PBMC from three different healthy human donors were cultured with increasing concentrations of HLE bispecific antibody constructs specific for Target B antigens for 48 h. The expression of the activation marker CD69 on CD4⁺ and CD8⁺ T cells was determined by flow cytometric analysis using a PE-Cy7 conjugated mab specific for CD69.

[9] Figure 4: Complement C1q Binding of BiTE[®] Fc fusion antibody constructs. BiTE[®] Fc fusion antibody constructs (BiTE[®] single chain Fc (triangle), BiTE[®] hetero Fc (squares), canonical BiTE[®] (circle)) were coated on a Maxisorp plate (in dilution series), prior to incubation with pooled human serum and incubation with polyclonal anti-human CC1q murine antibody, visualized by goat anti-mouse Fc-AP conjugate.

[10] Figure 5: Mean PK profiles of two different PSMA BiTE[®]-HLE antibody constructs after single dose administration in cynomolgus monkeys. For reasons of comparability, serum concentrations were dose-normalized to 15 µg/kg and indicated in nmol.

[11] Figure 6: Bispecific scFc variants D9F (SEQ ID NO: 481), T2G (SEQ ID NO: 482), D3L (SEQ ID NO: 483), T7I (SEQ ID NO: 484) and K6C (SEQ ID NO: 485). A preferred antibody construct of the present invention is shown in SEQ ID NO: 481.

[12] Figure 7: Surface Plasmon Resonance (SPR)-based determination of binding to human FcRn. Constructs D9F, T2G, D3L, T7I and K6C were each tested for their capability of binding against human FcRn in SPR (Biacore) experiments. The maximal binding during the injection phase was measured for all constructs as the respective response units (RU), equivalent to the molecular mass increase on the FcRn coated CM5 chip due to bound construct. All constructs were measured in duplicates. Average values of the duplicate determinations are depicted in Figure 7A and 7B.

[13] Figure 8: The constructs D9F, T2G, D3L, T7I and K6C and a human IgG1-kappa antibody MT201 were each tested for their capability of binding against human FcRn in SPR (Biacore) experiments. The maximal binding during the injection phase was measured for all constructs as the respective response units (RU), equivalent to the molecular mass increase on the FcRn coated CM5 chip due to bound construct. All constructs were measured in duplicates. Average values of the duplicate determinations are depicted including standard deviation error bars

DETAILED DESCRIPTION

[14] In addition to the significantly prolonged half-life of antibody constructs of the invention, which are preferably bispecific, the fusion of the specific Fc modality, i.e. the third domain according to the present invention, is responsible for a surprising significant impact on the first and second binding domain of the antibody construct of the invention. Thus, while other half-life extending modalities of T cell engaging antibody constructs show individual preferred features, the election of the present specific Fc modality allows for the provision of bispecific molecules which typically show a broad spectrum of preferred characteristics of a robust molecular format and, thus, allow for the development of promising pharmaceutical compositions.

[15] Thus, the present invention provides an antibody construct comprising :

- a first domain which binds to PSMA,
- a second domain which binds to an extracellular epitope of the human and the *Macaca* CD3ε chain; and
- a third domain which comprises two polypeptide monomers, each comprising a hinge, a CH2 domain and a CH3 domain, wherein said two polypeptide monomers are fused to each other via a peptide linker.

[16] The term “antibody construct” refers to a molecule in which the structure and/or function is/are based on the structure and/or function of an antibody, *e.g.*, of a full-length or whole immunoglobulin molecule. An antibody construct is hence capable of binding to its specific target or antigen and/or is/are drawn from the variable heavy chain (VH) and/or variable light chain (VL) domains of an antibody or fragment thereof. Furthermore, the binding domain of an antibody construct according to the invention comprises the minimum structural requirements of an antibody which allow for the target binding. This minimum requirement may *e.g.* be defined by the presence of at least the three light chain CDRs (*i.e.* CDR1, CDR2 and CDR3 of the VL region) and/or the three heavy chain CDRs (*i.e.* CDR1, CDR2 and CDR3 of the VH region), preferably of all six CDRs. An alternative approach to define the minimal structure requirements of an antibody is the definition of the epitope of the antibody within the structure of the specific target, respectively, the protein domain of the target protein composing the epitope region (epitope cluster) or by reference to an specific antibody competing with the epitope of the defined antibody. The antibodies on which the constructs according to the invention are based include for example monoclonal, recombinant, chimeric, deimmunized, humanized and human antibodies.

[17] The binding domain of an antibody construct according to the invention may *e.g.* comprise the above referred groups of CDRs. Preferably, those CDRs are comprised in the framework of an antibody light chain variable region (VL) and an antibody heavy chain variable region (VH); however, it does not have to comprise both. Fd fragments, for example, have two VH regions and often retain some antigen-binding function of the intact antigen-binding domain. Additional examples for the format of antibody fragments, antibody variants or binding domains include (1) a Fab fragment, a monovalent fragment having the VL, VH, CL and CH1 domains; (2) a F(ab')₂ fragment, a bivalent fragment having two Fab fragments linked by a disulfide bridge at the hinge region; (3) an Fd fragment having the two VH and CH1 domains; (4) an Fv fragment having the VL and VH domains of a single arm of an antibody, (5) a dAb fragment (Ward et al., (1989) Nature 341 :544-546), which has a VH domain; (6) an isolated complementarity determining region (CDR), and (7) a single chain Fv (scFv) , the latter being preferred (for example, derived from an scFV-library). Examples for embodiments of antibody constructs according to the invention are *e.g.* described in WO 00/006605, WO 2005/040220, WO 2008/119567, WO 2010/037838, WO 2013/026837, WO 2013/026833, US 2014/0308285, US 2014/0302037, WO 2014/144722, WO 2014/151910, and WO 2015/048272.

[18] Also within the definition of “binding domain” or “domain which binds” are fragments of full-length antibodies, such as VH, VHH, VL, (s)dAb, Fv, Fd, Fab, Fab', F(ab')₂ or “r IgG” (“half antibody”). Antibody constructs according to the invention may also comprise modified

fragments of antibodies, also called antibody variants, such as scFv, di-scFv or bi(s)-scFv, scFv-Fc, scFv-zipper, scFab, Fab₂, Fab₃, diabodies, single chain diabodies, tandem diabodies (Tandab's), tandem di-scFv, tandem tri-scFv, "multibodies" such as triabodies or tetrabodies, and single domain antibodies such as nanobodies or single variable domain antibodies comprising merely one variable domain, which may be VHH, VH or VL, that specifically bind an antigen or epitope independently of other V regions or domains.

[19] As used herein, the terms "single-chain Fv," "single-chain antibodies" or "scFv" refer to single polypeptide chain antibody fragments that comprise the variable regions from both the heavy and light chains, but lack the constant regions. Generally, a single-chain antibody further comprises a polypeptide linker between the VH and VL domains which enables it to form the desired structure which would allow for antigen binding. Single chain antibodies are discussed in detail by Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds. Springer-Verlag, New York, pp. 269-315 (1994). Various methods of generating single chain antibodies are known, including those described in U.S. Pat. Nos. 4,694,778 and 5,260,203; International Patent Application Publication No. WO 88/01649; Bird (1988) *Science* 242:423-442; Huston *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883; Ward *et al.* (1989) *Nature* 334:54454; Skerra *et al.* (1988) *Science* 242:1038-1041. In specific embodiments, single-chain antibodies can also be bispecific, multispecific, human, and/or humanized and/or synthetic.

[20] Furthermore, the definition of the term "antibody construct" includes monovalent, bivalent and polyvalent / multivalent constructs and, thus, bispecific constructs, specifically binding to only two antigenic structure, as well as polyspecific / multispecific constructs, which specifically bind more than two antigenic structures, e.g. three, four or more, through distinct binding domains. Moreover, the definition of the term "antibody construct" includes molecules consisting of only one polypeptide chain as well as molecules consisting of more than one polypeptide chain, which chains can be either identical (homodimers, homotrimers or homo oligomers) or different (heterodimer, heterotrimer or heterooligomer). Examples for the above identified antibodies and variants or derivatives thereof are described *inter alia* in Harlow and Lane, *Antibodies a laboratory manual*, CSHL Press (1988) and *Using Antibodies: a laboratory manual*, CSHL Press (1999), Kontermann and Dübeler, *Antibody Engineering*, Springer, 2nd ed. 2010 and Little, *Recombinant Antibodies for Immunotherapy*, Cambridge University Press 2009.

[21] The term "bispecific" as used herein refers to an antibody construct which is "at least bispecific", *i.e.*, it comprises at least a first binding domain and a second binding domain, wherein the first binding domain binds to one antigen or target (here: PSMA), and the second binding domain binds to another antigen or target (here: CD3). Accordingly, antibody

constructs according to the invention comprise specificities for at least two different antigens or targets. For example, the first domain does preferably not bind to an extracellular epitope of CD3 ϵ of one or more of the species as described herein. The term “target cell surface antigen” refers to an antigenic structure expressed by a cell and which is present at the cell surface such that it is accessible for an antibody construct as described herein. It may be a protein, preferably the extracellular portion of a protein, or a carbohydrate structure, preferably a carbohydrate structure of a protein, such as a glycoprotein. It is preferably a tumor antigen. The term “bispecific antibody construct” of the invention also encompasses multispecific antibody constructs such as trispecific antibody constructs, the latter ones including three binding domains, or constructs having more than three (e.g. four, five...) specificities.

[22] Given that the antibody constructs according to the invention are (at least) bispecific, they do not occur naturally and they are markedly different from naturally occurring products. A “bispecific” antibody construct or immunoglobulin is hence an artificial hybrid antibody or immunoglobulin having at least two distinct binding sides with different specificities. Bispecific antibody constructs can be produced by a variety of methods including fusion of hybridomas or linking of Fab' fragments. See, e.g., Songsivilai & Lachmann, Clin. Exp. Immunol. 79:315-321 (1990).

[23] The at least two binding domains and the variable domains (VH / VL) of the antibody construct of the present invention may or may not comprise peptide linkers (spacer peptides). The term “peptide linker” comprises in accordance with the present invention an amino acid sequence by which the amino acid sequences of one (variable and/or binding) domain and another (variable and/or binding) domain of the antibody construct of the invention are linked with each other. The peptide linkers can also be used to fuse the third domain to the other domains of the antibody construct of the invention. An essential technical feature of such peptide linker is that it does not comprise any polymerization activity. Among the suitable peptide linkers are those described in U.S. Patents 4,751,180 and 4,935,233 or WO 88/09344. The peptide linkers can also be used to attach other domains or modules or regions (such as half-life extending domains) to the antibody construct of the invention.

[24] The antibody constructs of the present invention are preferably “*in vitro* generated antibody constructs”. This term refers to an antibody construct according to the above definition where all or part of the variable region (e.g., at least one CDR) is generated in a non-immune cell selection, e.g., an *in vitro* phage display, protein chip or any other method in which candidate sequences can be tested for their ability to bind to an antigen. This term thus preferably excludes sequences generated solely by genomic rearrangement in an

immune cell in an animal. A “recombinant antibody” is an antibody made through the use of recombinant DNA technology or genetic engineering.

[25] The term “monoclonal antibody” (mAb) or monoclonal antibody construct as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations and/or post-translation modifications (e.g., isomerizations, amidations) that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic side or determinant on the antigen, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (or epitopes). In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized by the hybridoma culture, hence uncontaminated by other immunoglobulins. The modifier “monoclonal” indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method.

[26] For the preparation of monoclonal antibodies, any technique providing antibodies produced by continuous cell line cultures can be used. For example, monoclonal antibodies to be used may be made by the hybridoma method first described by Koehler *et al.*, *Nature*, 256: 495 (1975), or may be made by recombinant DNA methods (see, e.g., U.S. Patent No. 4,816,567). Examples for further techniques to produce human monoclonal antibodies include the trioma technique, the human B-cell hybridoma technique (Kozbor, *Immunology Today* 4 (1983), 72) and the EBV-hybridoma technique (Cole *et al.*, *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc. (1985), 77-96).

[27] Hybridomas can then be screened using standard methods, such as enzyme-linked immunosorbent assay (ELISA) and surface plasmon resonance (BIAcore™) analysis, to identify one or more hybridomas that produce an antibody that specifically binds with a specified antigen. Any form of the relevant antigen may be used as the immunogen, *e.g.*, recombinant antigen, naturally occurring forms, any variants or fragments thereof, as well as an antigenic peptide thereof. Surface plasmon resonance as employed in the BIAcore system can be used to increase the efficiency of phage antibodies which bind to an epitope of a target cell surface antigen (Schier, *Human Antibodies Hybridomas* 7 (1996), 97-105; Malmborg, *J. Immunol. Methods* 183 (1995), 7-13).

[28] Another exemplary method of making monoclonal antibodies includes screening protein expression libraries, e.g., phage display or ribosome display libraries. Phage display is described, for example, in Ladner *et al.*, U.S. Patent No. 5,223,409; Smith (1985) *Science*

228:1315-1317, Clackson *et al.*, Nature, 352: 624-628 (1991) and Marks *et al.*, J. Mol. Biol., 222: 581-597 (1991).

[29] In addition to the use of display libraries, the relevant antigen can be used to immunize a non-human animal, *e.g.*, a rodent (such as a mouse, hamster, rabbit or rat). In one embodiment, the non-human animal includes at least a part of a human immunoglobulin gene. For example, it is possible to engineer mouse strains deficient in mouse antibody production with large fragments of the human Ig (immunoglobulin) loci. Using the hybridoma technology, antigen-specific monoclonal antibodies derived from the genes with the desired specificity may be produced and selected. See, *e.g.*, XENOMOUSE™, Green *et al.* (1994) Nature Genetics 7:13-21, US 2003-0070185, WO 96/34096, and WO 96/33735.

[30] A monoclonal antibody can also be obtained from a non-human animal, and then modified, *e.g.*, humanized, deimmunized, rendered chimeric etc., using recombinant DNA techniques known in the art. Examples of modified antibody constructs include humanized variants of non-human antibodies, "affinity matured" antibodies (see, *e.g.* Hawkins *et al.* J. Mol. Biol. 254, 889-896 (1992) and Lowman *et al.*, Biochemistry 30, 10832- 10837 (1991)) and antibody mutants with altered effector function(s) (see, *e.g.*, US Patent 5,648,260, Kontermann and Dübel (2010), *loc. cit.* and Little (2009), *loc. cit.*).

[31] In immunology, affinity maturation is the process by which B cells produce antibodies with increased affinity for antigen during the course of an immune response. With repeated exposures to the same antigen, a host will produce antibodies of successively greater affinities. Like the natural prototype, the *in vitro* affinity maturation is based on the principles of mutation and selection. The *in vitro* affinity maturation has successfully been used to optimize antibodies, antibody constructs, and antibody fragments. Random mutations inside the CDRs are introduced using radiation, chemical mutagens or error-prone PCR. In addition, the genetic diversity can be increased by chain shuffling. Two or three rounds of mutation and selection using display methods like phage display usually results in antibody fragments with affinities in the low nanomolar range.

[32] A preferred type of an amino acid substitutional variation of the antibody constructs involves substituting one or more hypervariable region residues of a parent antibody (*e. g.* a humanized or human antibody). Generally, the resulting variant(s) selected for further development will have improved biological properties relative to the parent antibody from which they are generated. A convenient way for generating such substitutional variants involves affinity maturation using phage display. Briefly, several hypervariable region sides (*e. g.* 6-7 sides) are mutated to generate all possible amino acid substitutions at each side. The antibody variants thus generated are displayed in a monovalent fashion from

filamentous phage particles as fusions to the gene III product of M13 packaged within each particle. The phage-displayed variants are then screened for their biological activity (e. g. binding affinity) as herein disclosed. In order to identify candidate hypervariable region sides for modification, alanine scanning mutagenesis can be performed to identify hypervariable region residues contributing significantly to antigen binding. Alternatively, or additionally, it may be beneficial to analyze a crystal structure of the antigen-antibody complex to identify contact points between the binding domain and, *e.g.*, human PSMA. Such contact residues and neighbouring residues are candidates for substitution according to the techniques elaborated herein. Once such variants are generated, the panel of variants is subjected to screening as described herein and antibodies with superior properties in one or more relevant assays may be selected for further development.

[33] The monoclonal antibodies and antibody constructs of the present invention specifically include “chimeric” antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is/are identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567; Morrison *et al.*, Proc. Natl. Acad. Sci. USA, 81: 6851-6855 (1984)). Chimeric antibodies of interest herein include “primitized” antibodies comprising variable domain antigen-binding sequences derived from a non-human primate (*e.g.*, Old World Monkey, Ape *etc.*) and human constant region sequences. A variety of approaches for making chimeric antibodies have been described. See *e.g.*, Morrison *et al.*, Proc. Natl. Acad. Sci. U.S.A. 81:6851, 1985; Takeda *et al.*, Nature 314:452, 1985, Cabilly *et al.*, U.S. Patent No. 4,816,567; Boss *et al.*, U.S. Patent No. 4,816,397; Tanaguchi *et al.*, EP 0171496; EP 0173494; and GB 2177096.

[34] An antibody, antibody construct, antibody fragment or antibody variant may also be modified by specific deletion of human T cell epitopes (a method called “deimmunization”) by the methods disclosed for example in WO 98/52976 or WO 00/34317. Briefly, the heavy and light chain variable domains of an antibody can be analyzed for peptides that bind to MHC class II; these peptides represent potential T cell epitopes (as defined in WO 98/52976 and WO 00/34317). For detection of potential T cell epitopes, a computer modeling approach termed “peptide threading” can be applied, and in addition a database of human MHC class II binding peptides can be searched for motifs present in the VH and VL sequences, as described in WO 98/52976 and WO 00/34317. These motifs bind to any of the 18 major MHC class II DR allotypes, and thus constitute potential T cell epitopes. Potential T cell epitopes

detected can be eliminated by substituting small numbers of amino acid residues in the variable domains, or preferably, by single amino acid substitutions. Typically, conservative substitutions are made. Often, but not exclusively, an amino acid common to a position in human germline antibody sequences may be used. Human germline sequences are disclosed *e.g.* in Tomlinson, *et al.* (1992) *J. Mol. Biol.* 227:776-798; Cook, G.P. *et al.* (1995) *Immunol. Today* Vol. 16 (5): 237-242; and Tomlinson *et al.* (1995) *EMBO J.* 14: 14:4628-4638. The V BASE directory provides a comprehensive directory of human immunoglobulin variable region sequences (compiled by Tomlinson, LA. *et al.* MRC Centre for Protein Engineering, Cambridge, UK). These sequences can be used as a source of human sequence, *e.g.*, for framework regions and CDRs. Consensus human framework regions can also be used, for example as described in US Patent No. 6,300,064.

[35] “Humanized” antibodies, antibody constructs, variants or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) are antibodies or immunoglobulins of mostly human sequences, which contain (a) minimal sequence(s) derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region (also CDR) of the recipient are replaced by residues from a hypervariable region of a non-human (*e.g.*, rodent) species (donor antibody) such as mouse, rat, hamster or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, “humanized antibodies” as used herein may also comprise residues which are found neither in the recipient antibody nor the donor antibody. These modifications are made to further refine and optimize antibody performance. The humanized antibody may also comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, *Nature*, 321: 522-525 (1986); Reichmann *et al.*, *Nature*, 332: 323-329 (1988); and Presta, *Curr. Op. Struct. Biol.*, 2: 593-596 (1992).

[36] Humanized antibodies or fragments thereof can be generated by replacing sequences of the Fv variable domain that are not directly involved in antigen binding with equivalent sequences from human Fv variable domains. Exemplary methods for generating humanized antibodies or fragments thereof are provided by Morrison (1985) *Science* 229:1202-1207; by Oi *et al.* (1986) *BioTechniques* 4:214; and by US 5,585,089; US 5,693,761; US 5,693,762; US 5,859,205; and US 6,407,213. Those methods include isolating, manipulating, and expressing the nucleic acid sequences that encode all or part of immunoglobulin Fv variable domains from at least one of a heavy or light chain. Such nucleic acids may be obtained from a hybridoma producing an antibody against a predetermined

target, as described above, as well as from other sources. The recombinant DNA encoding the humanized antibody molecule can then be cloned into an appropriate expression vector.

[37] Humanized antibodies may also be produced using transgenic animals such as mice that express human heavy and light chain genes, but are incapable of expressing the endogenous mouse immunoglobulin heavy and light chain genes. Winter describes an exemplary CDR grafting method that may be used to prepare the humanized antibodies described herein (U.S. Patent No. 5,225,539). All of the CDRs of a particular human antibody may be replaced with at least a portion of a non-human CDR, or only some of the CDRs may be replaced with non-human CDRs. It is only necessary to replace the number of CDRs required for binding of the humanized antibody to a predetermined antigen.

[38] A humanized antibody can be optimized by the introduction of conservative substitutions, consensus sequence substitutions, germline substitutions and/or back mutations. Such altered immunoglobulin molecules can be made by any of several techniques known in the art, (e.g., Teng *et al.*, Proc. Natl. Acad. Sci. U.S.A., 80: 7308-7312, 1983; Kozbor *et al.*, Immunology Today, 4: 7279, 1983; Olsson *et al.*, Meth. Enzymol., 92: 3-16, 1982, and EP 239 400).

[39] The term "human antibody", "human antibody construct" and "human binding domain" includes antibodies, antibody constructs and binding domains having antibody regions such as variable and constant regions or domains which correspond substantially to human germline immunoglobulin sequences known in the art, including, for example, those described by Kabat *et al.* (1991) (*loc. cit.*). The human antibodies, antibody constructs or binding domains of the invention may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis *in vitro* or by somatic mutation *in vivo*), for example in the CDRs, and in particular, in CDR3. The human antibodies, antibody constructs or binding domains can have at least one, two, three, four, five, or more positions replaced with an amino acid residue that is not encoded by the human germline immunoglobulin sequence. The definition of human antibodies, antibody constructs and binding domains as used herein also contemplates fully human antibodies, which include only non-artificially and/or genetically altered human sequences of antibodies as those can be derived by using technologies or systems such as the Xenomouse. Preferably, a "fully human antibody" does not include amino acid residues not encoded by human germline immunoglobulin sequences.

[40] In some embodiments, the antibody constructs of the invention are "isolated" or "substantially pure" antibody constructs. "Isolated" or "substantially pure", when used to describe the antibody constructs disclosed herein, means an antibody construct that has

been identified, separated and/or recovered from a component of its production environment. Preferably, the antibody construct is free or substantially free of association with all other components from its production environment. Contaminant components of its production environment, such as that resulting from recombinant transfected cells, are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. The antibody constructs may e.g constitute at least about 5%, or at least about 50% by weight of the total protein in a given sample. It is understood that the isolated protein may constitute from 5% to 99.9% by weight of the total protein content, depending on the circumstances. The polypeptide may be made at a significantly higher concentration through the use of an inducible promoter or high expression promoter, such that it is made at increased concentration levels. The definition includes the production of an antibody construct in a wide variety of organisms and/or host cells that are known in the art. In preferred embodiments, the antibody construct will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Ordinarily, however, an isolated antibody construct will be prepared by at least one purification step.

[41] The term "binding domain" characterizes in connection with the present invention a domain which (specifically) binds to / interacts with / recognizes a given target epitope or a given target side on the target molecules (antigens), here: PSMA and CD3, respectively. The structure and function of the first binding domain (recognizing PSMA), and preferably also the structure and/or function of the second binding domain (recognizing CD3), is/are based on the structure and/or function of an antibody, e.g. of a full-length or whole immunoglobulin molecule, and/or is/are drawn from the variable heavy chain (VH) and/or variable light chain (VL) domains of an antibody or fragment thereof. Preferably the first binding domain is characterized by the presence of three light chain CDRs (i.e. CDR1, CDR2 and CDR3 of the VL region) and/or three heavy chain CDRs (i.e. CDR1, CDR2 and CDR3 of the VH region). The second binding domain preferably also comprises the minimum structural requirements of an antibody which allow for the target binding. More preferably, the second binding domain comprises at least three light chain CDRs (i.e. CDR1, CDR2 and CDR3 of the VL region) and/or three heavy chain CDRs (i.e. CDR1, CDR2 and CDR3 of the VH region). It is envisaged that the first and/or second binding domain is produced by or obtainable by phage-display or library screening methods rather than by grafting CDR sequences from a pre-existing (monoclonal) antibody into a scaffold.

[42] According to the present invention, binding domains are in the form of one or more polypeptides. Such polypeptides may include proteinaceous parts and non-proteinaceous parts (e.g. chemical linkers or chemical cross-linking agents such as glutaraldehyde). Proteins (including fragments thereof, preferably biologically active fragments, and peptides, usually having less than 30 amino acids) comprise two or more amino acids coupled to each other via a covalent peptide bond (resulting in a chain of amino acids).

[43] The term "polypeptide" as used herein describes a group of molecules, which usually consist of more than 30 amino acids. Polypeptides may further form multimers such as dimers, trimers and higher oligomers, *i.e.*, consisting of more than one polypeptide molecule. Polypeptide molecules forming such dimers, trimers etc. may be identical or non-identical. The corresponding higher order structures of such multimers are, consequently, termed homo- or heterodimers, homo- or heterotrimers etc. An example for a heteromultimer is an antibody molecule, which, in its naturally occurring form, consists of two identical light polypeptide chains and two identical heavy polypeptide chains. The terms "peptide", "polypeptide" and "protein" also refer to naturally modified peptides / polypeptides / proteins wherein the modification is effected *e.g.* by post-translational modifications like glycosylation, acetylation, phosphorylation and the like. A "peptide", "polypeptide" or "protein" when referred to herein may also be chemically modified such as pegylated. Such modifications are well known in the art and described herein below.

[44] Preferably the binding domain which binds to PSMA and/or the binding domain which binds to CD3 ϵ is/are human binding domains. Antibodies and antibody constructs comprising at least one human binding domain avoid some of the problems associated with antibodies or antibody constructs that possess non-human such as rodent (*e.g.* murine, rat, hamster or rabbit) variable and/or constant regions. The presence of such rodent derived proteins can lead to the rapid clearance of the antibodies or antibody constructs or can lead to the generation of an immune response against the antibody or antibody construct by a patient. In order to avoid the use of rodent derived antibodies or antibody constructs, human or fully human antibodies / antibody constructs can be generated through the introduction of human antibody function into a rodent so that the rodent produces fully human antibodies.

[45] The ability to clone and reconstruct megabase-sized human loci in YACs and to introduce them into the mouse germline provides a powerful approach to elucidating the functional components of very large or crudely mapped loci as well as generating useful models of human disease. Furthermore, the use of such technology for substitution of mouse loci with their human equivalents could provide unique insights into the expression and regulation of human gene products during development, their communication with other systems, and their involvement in disease induction and progression.

[46] An important practical application of such a strategy is the “humanization” of the mouse humoral immune system. Introduction of human immunoglobulin (Ig) loci into mice in which the endogenous Ig genes have been inactivated offers the opportunity to study the mechanisms underlying programmed expression and assembly of antibodies as well as their role in B-cell development. Furthermore, such a strategy could provide an ideal source for production of fully human monoclonal antibodies (mAbs) – an important milestone towards fulfilling the promise of antibody therapy in human disease. Fully human antibodies or antibody constructs are expected to minimize the immunogenic and allergic responses intrinsic to mouse or mouse-derivatized mAbs and thus to increase the efficacy and safety of the administered antibodies / antibody constructs. The use of fully human antibodies or antibody constructs can be expected to provide a substantial advantage in the treatment of chronic and recurring human diseases, such as inflammation, autoimmunity, and cancer, which require repeated compound administrations.

[47] One approach towards this goal was to engineer mouse strains deficient in mouse antibody production with large fragments of the human Ig loci in anticipation that such mice would produce a large repertoire of human antibodies in the absence of mouse antibodies. Large human Ig fragments would preserve the large variable gene diversity as well as the proper regulation of antibody production and expression. By exploiting the mouse machinery for antibody diversification and selection and the lack of immunological tolerance to human proteins, the reproduced human antibody repertoire in these mouse strains should yield high affinity antibodies against any antigen of interest, including human antigens. Using the hybridoma technology, antigen-specific human mAbs with the desired specificity could be readily produced and selected. This general strategy was demonstrated in connection with the generation of the first XenoMouse mouse strains (see Green et al. *Nature Genetics* 7:13-21 (1994)). The XenoMouse strains were engineered with yeast artificial chromosomes (YACs) containing 245 kb and 190 kb-sized germline configuration fragments of the human heavy chain locus and kappa light chain locus, respectively, which contained core variable and constant region sequences. The human Ig containing YACs proved to be compatible with the mouse system for both rearrangement and expression of antibodies and were capable of substituting for the inactivated mouse Ig genes. This was demonstrated by their ability to induce B cell development, to produce an adult-like human repertoire of fully human antibodies, and to generate antigen-specific human mAbs. These results also suggested that introduction of larger portions of the human Ig loci containing greater numbers of V genes, additional regulatory elements, and human Ig constant regions may recapitulate substantially the full repertoire that is characteristic of the human humoral response to infection and immunization. The work of Green et al. was recently extended to the introduction of greater than approximately 80% of the human antibody repertoire through introduction of megabase

sized, germline configuration YAC fragments of the human heavy chain loci and kappa light chain loci, respectively. See Mendez *et al.* Nature Genetics 15:146-156 (1997) and U.S. patent application Ser. No. 08/759,620.

[48] The production of the XenoMouse mice is further discussed and delineated in U.S. patent applications Ser. No. 07/466,008, Ser. No. 07/610,515, Ser. No. 07/919,297, Ser. No. 07/922,649, Ser. No. 08/031,801, Ser. No. 08/112,848, Ser. No. 08/234,145, Ser. No. 08/376,279, Ser. No. 08/430,938, Ser. No. 08/464,584, Ser. No. 08/464,582, Ser. No. 08/463,191, Ser. No. 08/462,837, Ser. No. 08/486,853, Ser. No. 08/486,857, Ser. No. 08/486,859, Ser. No. 08/462,513, Ser. No. 08/724,752, and Ser. No. 08/759,620; and U.S. Pat. Nos. 6,162,963; 6,150,584; 6,114,598; 6,075,181, and 5,939,598 and Japanese Patent Nos. 3 068 180 B2, 3 068 506 B2, and 3 068 507 B2. See also Mendez *et al.* Nature Genetics 15:146-156 (1997) and Green and Jakobovits J. Exp. Med. 188:483-495 (1998), EP 0 463 151 B1, WO 94/02602, WO 96/34096, WO 98/24893, WO 00/76310, and WO 03/47336.

[49] In an alternative approach, others, including GenPharm International, Inc., have utilized a “minilocus” approach. In the minilocus approach, an exogenous Ig locus is mimicked through the inclusion of pieces (individual genes) from the Ig locus. Thus, one or more VH genes, one or more DH genes, one or more JH genes, a mu constant region, and a second constant region (preferably a gamma constant region) are formed into a construct for insertion into an animal. This approach is described in U.S. Pat. No. 5,545,807 to Surani *et al.* and U.S. Pat. Nos. 5,545,806; 5,625,825; 5,625,126; 5,633,425; 5,661,016; 5,770,429; 5,789,650; 5,814,318; 5,877,397; 5,874,299; and 6,255,458 each to Lonberg and Kay, U.S. Pat. Nos. 5,591,669 and 6,023,010 to Krimpenfort and Berns, U.S. Pat. Nos. 5,612,205; 5,721,367; and 5,789,215 to Berns *et al.*, and U.S. Pat. No. 5,643,763 to Choi and Dunn, and GenPharm International U.S. patent application Ser. No. 07/574,748, Ser. No. 07/575,962, Ser. No. 07/810,279, Ser. No. 07/853,408, Ser. No. 07/904,068, Ser. No. 07/990,860, Ser. No. 08/053,131, Ser. No. 08/096,762, Ser. No. 08/155,301, Ser. No. 08/161,739, Ser. No. 08/165,699, Ser. No. 08/209,741. See also EP 0 546 073 B1, WO 92/03918, WO 92/22645, WO 92/22647, WO 92/22670, WO 93/12227, WO 94/00569, WO 94/25585, WO 96/14436, WO 97/13852, and WO 98/24884 and U.S. Pat. No. 5,981,175. See further Taylor *et al.* (1992), Chen *et al.* (1993), Tuailon *et al.* (1993), Choi *et al.* (1993), Lonberg *et al.* (1994), Taylor *et al.* (1994), and Tuailon *et al.* (1995), Fishwild *et al.* (1996).

[50] Kirin has also demonstrated the generation of human antibodies from mice in which, through microcell fusion, large pieces of chromosomes, or entire chromosomes, have been introduced. See European Patent Application Nos. 773 288 and 843 961. Xenerex Biosciences is developing a technology for the potential generation of human antibodies. In

this technology, SCID mice are reconstituted with human lymphatic cells, e.g., B and/or T cells. Mice are then immunized with an antigen and can generate an immune response against the antigen. See U.S. Pat. Nos. 5,476,996; 5,698,767; and 5,958,765.

- 5 **[51]** Human anti-mouse antibody (HAMA) responses have led the industry to prepare chimeric or otherwise humanized antibodies. It is however expected that certain human anti-chimeric antibody (HACA) responses will be observed, particularly in chronic or multi-dose utilizations of the antibody. Thus, it would be desirable to provide antibody constructs comprising a human binding domain against PSMA and a human binding domain against CD3 ϵ in order to vitiate concerns and/or effects of HAMA or HACA response.
- 10 **[52]** The terms “(specifically) binds to”, “(specifically) recognizes”, “is (specifically) directed to”, and “(specifically) reacts with” mean in accordance with this invention that a binding domain interacts or specifically interacts with a given epitope or a given target side on the target molecules (antigens), here: PSMA and CD3 ϵ , respectively.
- 15 **[53]** The term “epitope” refers to a side on an antigen to which a binding domain, such as an antibody or immunoglobulin, or a derivative, fragment or variant of an antibody or an immunoglobulin, specifically binds. An “epitope” is antigenic and thus the term epitope is sometimes also referred to herein as “antigenic structure” or “antigenic determinant”. Thus, the binding domain is an “antigen interaction side”. Said binding/interaction is also understood to define a “specific recognition”.
- 20 **[54]** “Epitopes” can be formed both by contiguous amino acids or non-contiguous amino acids juxtaposed by tertiary folding of a protein. A “linear epitope” is an epitope where an amino acid primary sequence comprises the recognized epitope. A linear epitope typically includes at least 3 or at least 4, and more usually, at least 5 or at least 6 or at least 7, for example, about 8 to about 10 amino acids in a unique sequence.
- 25 **[55]** A “conformational epitope”, in contrast to a linear epitope, is an epitope wherein the primary sequence of the amino acids comprising the epitope is not the sole defining component of the epitope recognized (*e.g.*, an epitope wherein the primary sequence of amino acids is not necessarily recognized by the binding domain). Typically a conformational epitope comprises an increased number of amino acids relative to a linear epitope. With
30 regard to recognition of conformational epitopes, the binding domain recognizes a three-dimensional structure of the antigen, preferably a peptide or protein or fragment thereof (in the context of the present invention, the antigenic structure for one of the binding domains is comprised within the target cell surface antigen protein). For example, when a protein molecule folds to form a three-dimensional structure, certain amino acids and/or the

polypeptide backbone forming the conformational epitope become juxtaposed enabling the antibody to recognize the epitope. Methods of determining the conformation of epitopes include, but are not limited to, x-ray crystallography, two-dimensional nuclear magnetic resonance (2D-NMR) spectroscopy and site-directed spin labelling and electron
 5 paramagnetic resonance (EPR) spectroscopy.

[56] A method for epitope mapping is described in the following: When a region (a contiguous amino acid stretch) in the human PSMA protein is exchanged / replaced with its corresponding region of a non-human and non-primate PSMA (*e.g.*, mouse PSMA, but others like chicken, rat, hamster, rabbit etc. may also be conceivable), a decrease in the
 10 binding of the binding domain is expected to occur, unless the binding domain is cross-reactive for the non-human, non-primate PSMA used. Said decrease is preferably at least 10%, 20%, 30%, 40%, or 50%; more preferably at least 60%, 70%, or 80%, and most preferably 90%, 95% or even 100% in comparison to the binding to the respective region in the human PSMA protein, whereby binding to the respective region in the human PSMA
 15 protein is set to be 100%. It is envisaged that the aforementioned human PSMA / non-human PSMA chimeras are expressed in CHO cells. It is also envisaged that the human PSMA / non-human PSMA chimeras are fused with a transmembrane domain and/or cytoplasmic domain of a different membrane-bound protein such as EpCAM.

[57] In an alternative or additional method for epitope mapping, several truncated versions
 20 of the human PSMA extracellular domain can be generated in order to determine a specific region that is recognized by a binding domain. In these truncated versions, the different extracellular PSMA domains / sub-domains or regions are stepwise deleted, starting from the N-terminus. It is envisaged that the truncated PSMA versions may be expressed in CHO cells. It is also envisaged that the truncated PSMA versions may be fused with a
 25 transmembrane domain and/or cytoplasmic domain of a different membrane-bound protein such as EpCAM. It is also envisaged that the truncated PSMA versions may encompass a signal peptide domain at their N-terminus, for example a signal peptide derived from mouse IgG heavy chain signal peptide. It is furthermore envisaged that the truncated PSMA versions may encompass a v5 domain at their N-terminus (following the signal peptide)
 30 which allows verifying their correct expression on the cell surface. A decrease or a loss of binding is expected to occur with those truncated PSMA versions which do not encompass any more the PSMA region that is recognized by the binding domain. The decrease of binding is preferably at least 10%, 20%, 30%, 40%, 50%; more preferably at least 60%, 70%, 80%, and most preferably 90%, 95% or even 100%, whereby binding to the entire human
 35 PSMA protein (or its extracellular region or domain) is set to be 100.

[58] A further method to determine the contribution of a specific residue of PSMA to the recognition by an antibody construct or binding domain is alanine scanning (see e.g. Morrison KL & Weiss GA. *Cur Opin Chem Biol.* 2001 Jun;5(3):302-7), where each residue to be analyzed is replaced by alanine, e.g. via site-directed mutagenesis. Alanine is used because of its non-bulky, chemically inert, methyl functional group that nevertheless mimics the secondary structure references that many of the other amino acids possess. Sometimes bulky amino acids such as valine or leucine can be used in cases where conservation of the size of mutated residues is desired. Alanine scanning is a mature technology which has been used for a long period of time.

[59] The interaction between the binding domain and the epitope or the region comprising the epitope implies that a binding domain exhibits appreciable affinity for the epitope / the region comprising the epitope on a particular protein or antigen (here: PSMA and CD3, respectively) and, generally, does not exhibit significant reactivity with proteins or antigens other than the PSMA or CD3. "Appreciable affinity" includes binding with an affinity of about 10^{-6} M (KD) or stronger. Preferably, binding is considered specific when the binding affinity is about 10^{-12} to 10^{-8} M, 10^{-12} to 10^{-9} M, 10^{-12} to 10^{-10} M, 10^{-11} to 10^{-8} M, preferably of about 10^{-11} to 10^{-9} M. Whether a binding domain specifically reacts with or binds to a target can be tested readily by, *inter alia*, comparing the reaction of said binding domain with a target protein or antigen with the reaction of said binding domain with proteins or antigens other than the PSMA or CD3. Preferably, a binding domain of the invention does not essentially or substantially bind to proteins or antigens other than PSMA or CD3 (*i.e.*, the first binding domain is not capable of binding to proteins other than PSMA and the second binding domain is not capable of binding to proteins other than CD3). It is an envisaged characteristic of the antibody constructs according to the present invention to have superior affinity characteristics in comparison to other HLE formats. Such a superior affinity, in consequence, suggests a prolonged half-life in vivo. The longer half-life of the antibody constructs according to the present invention may reduce the duration and frequency of administration which typically contributes to improved patient compliance. This is of particular importance as the antibody constructs of the present invention are particularly beneficial for highly weakened or even multimorbid cancer patients.

[60] The term "does not essentially / substantially bind" or "is not capable of binding" means that a binding domain of the present invention does not bind a protein or antigen other than the PSMA or CD3, *i.e.*, does not show reactivity of more than 30%, preferably not more than 20%, more preferably not more than 10%, particularly preferably not more than 9%, 8%, 7%, 6% or 5% with proteins or antigens other than PSMA or CD3, whereby binding to the PSMA or CD3, respectively, is set to be 100%.

[61] Specific binding is believed to be effected by specific motifs in the amino acid sequence of the binding domain and the antigen. Thus, binding is achieved as a result of their primary, secondary and/or tertiary structure as well as the result of secondary modifications of said structures. The specific interaction of the antigen-interaction-side with its specific antigen may result in a simple binding of said side to the antigen. Moreover, the specific interaction of the antigen-interaction-side with its specific antigen may alternatively or additionally result in the initiation of a signal, *e.g.* due to the induction of a change of the conformation of the antigen, an oligomerization of the antigen, etc.

[62] The term “variable” refers to the portions of the antibody or immunoglobulin domains that exhibit variability in their sequence and that are involved in determining the specificity and binding affinity of a particular antibody (i.e., the “variable domain(s)”). The pairing of a variable heavy chain (VH) and a variable light chain (VL) together forms a single antigen-binding site.

[63] Variability is not evenly distributed throughout the variable domains of antibodies; it is concentrated in sub-domains of each of the heavy and light chain variable regions. These sub-domains are called “hypervariable regions” or “complementarity determining regions” (CDRs). The more conserved (i.e., non-hypervariable) portions of the variable domains are called the “framework” regions (FRM or FR) and provide a scaffold for the six CDRs in three dimensional space to form an antigen-binding surface. The variable domains of naturally occurring heavy and light chains each comprise four FRM regions (FR1, FR2, FR3, and FR4), largely adopting a β -sheet configuration, connected by three hypervariable regions, which form loops connecting, and in some cases forming part of, the β -sheet structure. The hypervariable regions in each chain are held together in close proximity by the FRM and, with the hypervariable regions from the other chain, contribute to the formation of the antigen-binding side (see Kabat *et al.*, *loc. cit.*).

[64] The terms “CDR”, and its plural “CDRs”, refer to the complementarity determining region of which three make up the binding character of a light chain variable region (CDR-L1, CDR-L2 and CDR-L3) and three make up the binding character of a heavy chain variable region (CDR-H1, CDR-H2 and CDR-H3). CDRs contain most of the residues responsible for specific interactions of the antibody with the antigen and hence contribute to the functional activity of an antibody molecule: they are the main determinants of antigen specificity.

[65] The exact definitional CDR boundaries and lengths are subject to different classification and numbering systems. CDRs may therefore be referred to by Kabat, Chothia, contact or any other boundary definitions, including the numbering system described herein. Despite differing boundaries, each of these systems has some degree of overlap in what

constitutes the so called “hypervariable regions” within the variable sequences. CDR definitions according to these systems may therefore differ in length and boundary areas with respect to the adjacent framework region. See for example Kabat (an approach based on cross-species sequence variability), Chothia (an approach based on crystallographic studies of antigen-antibody complexes), and/or MacCallum (Kabat *et al.*, *loc. cit.*; Chothia *et al.*, J. Mol. Biol, 1987, 196: 901-917; and MacCallum *et al.*, J. Mol. Biol, 1996, 262: 732). Still another standard for characterizing the antigen binding side is the AbM definition used by Oxford Molecular's AbM antibody modeling software. See, *e.g.*, Protein Sequence and Structure Analysis of Antibody Variable Domains. In: Antibody Engineering Lab Manual (Ed.: Duebel, S. and Kontermann, R., Springer-Verlag, Heidelberg). To the extent that two residue identification techniques define regions of overlapping, but not identical regions, they can be combined to define a hybrid CDR. However, the numbering in accordance with the so-called Kabat system is preferred.

[66] Typically, CDRs form a loop structure that can be classified as a canonical structure. The term “canonical structure” refers to the main chain conformation that is adopted by the antigen binding (CDR) loops. From comparative structural studies, it has been found that five of the six antigen binding loops have only a limited repertoire of available conformations. Each canonical structure can be characterized by the torsion angles of the polypeptide backbone. Correspondent loops between antibodies may, therefore, have very similar three dimensional structures, despite high amino acid sequence variability in most parts of the loops (Chothia and Lesk, J. Mol. Biol., 1987, 196: 901; Chothia *et al.*, Nature, 1989, 342: 877; Martin and Thornton, J. Mol. Biol, 1996, 263: 800). Furthermore, there is a relationship between the adopted loop structure and the amino acid sequences surrounding it. The conformation of a particular canonical class is determined by the length of the loop and the amino acid residues residing at key positions within the loop, as well as within the conserved framework (*i.e.*, outside of the loop). Assignment to a particular canonical class can therefore be made based on the presence of these key amino acid residues.

[67] The term “canonical structure” may also include considerations as to the linear sequence of the antibody, for example, as catalogued by Kabat (Kabat *et al.*, *loc. cit.*). The Kabat numbering scheme (system) is a widely adopted standard for numbering the amino acid residues of an antibody variable domain in a consistent manner and is the preferred scheme applied in the present invention as also mentioned elsewhere herein. Additional structural considerations can also be used to determine the canonical structure of an antibody. For example, those differences not fully reflected by Kabat numbering can be described by the numbering system of Chothia *et al.* and/or revealed by other techniques, for example, crystallography and two- or three-dimensional computational modeling.

Accordingly, a given antibody sequence may be placed into a canonical class which allows for, among other things, identifying appropriate chassis sequences (e.g., based on a desire to include a variety of canonical structures in a library). Kabat numbering of antibody amino acid sequences and structural considerations as described by Chothia *et al.*, *loc. cit.* and their implications for construing canonical aspects of antibody structure, are described in the literature. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known in the art. For a review of the antibody structure, see Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, eds. Harlow *et al.*, 1988.

[68] The CDR3 of the light chain and, particularly, the CDR3 of the heavy chain may constitute the most important determinants in antigen binding within the light and heavy chain variable regions. In some antibody constructs, the heavy chain CDR3 appears to constitute the major area of contact between the antigen and the antibody. *In vitro* selection schemes in which CDR3 alone is varied can be used to vary the binding properties of an antibody or determine which residues contribute to the binding of an antigen. Hence, CDR3 is typically the greatest source of molecular diversity within the antibody-binding side. H3, for example, can be as short as two amino acid residues or greater than 26 amino acids.

[69] In a classical full-length antibody or immunoglobulin, each light (L) chain is linked to a heavy (H) chain by one covalent disulfide bond, while the two H chains are linked to each other by one or more disulfide bonds depending on the H chain isotype. The CH domain most proximal to VH is usually designated as CH1. The constant ("C") domains are not directly involved in antigen binding, but exhibit various effector functions, such as antibody-dependent, cell-mediated cytotoxicity and complement activation. The Fc region of an antibody is comprised within the heavy chain constant domains and is for example able to interact with cell surface located Fc receptors.

[70] The sequence of antibody genes after assembly and somatic mutation is highly varied, and these varied genes are estimated to encode 10^{10} different antibody molecules (Immunoglobulin Genes, 2nd ed., eds. Jonio *et al.*, Academic Press, San Diego, CA, 1995). Accordingly, the immune system provides a repertoire of immunoglobulins. The term "repertoire" refers to at least one nucleotide sequence derived wholly or partially from at least one sequence encoding at least one immunoglobulin. The sequence(s) may be generated by rearrangement *in vivo* of the V, D, and J segments of heavy chains, and the V and J segments of light chains. Alternatively, the sequence(s) can be generated from a cell in response to which rearrangement occurs, e.g., *in vitro* stimulation. Alternatively, part or all of the sequence(s) may be obtained by DNA splicing, nucleotide synthesis, mutagenesis, and other methods, see, e.g., U.S. Patent 5,565,332. A repertoire may include only one

sequence or may include a plurality of sequences, including ones in a genetically diverse collection.

[71] The term "Fc portion" or "Fc monomer" means in connection with this invention a polypeptide comprising at least one domain having the function of a CH2 domain and at least one domain having the function of a CH3 domain of an immunoglobulin molecule. As apparent from the term "Fc monomer", the polypeptide comprising those CH domains is a "polypeptide monomer". An Fc monomer can be a polypeptide comprising at least a fragment of the constant region of an immunoglobulin excluding the first constant region immunoglobulin domain of the heavy chain (CH1), but maintaining at least a functional part of one CH2 domain and a functional part of one CH3 domain, wherein the CH2 domain is amino terminal to the CH3 domain. In a preferred aspect of this definition, an Fc monomer can be a polypeptide constant region comprising a portion of the Ig-Fc hinge region, a CH2 region and a CH3 region, wherein the hinge region is amino terminal to the CH2 domain. It is envisaged that the hinge region of the present invention promotes dimerization. Such Fc polypeptide molecules can be obtained by papain digestion of an immunoglobulin region (of course resulting in a dimer of two Fc polypeptide), for example and not limitation. In another aspect of this definition, an Fc monomer can be a polypeptide region comprising a portion of a CH2 region and a CH3 region. Such Fc polypeptide molecules can be obtained by pepsin digestion of an immunoglobulin molecule, for example and not limitation. In one embodiment, the polypeptide sequence of an Fc monomer is substantially similar to an Fc polypeptide sequence of: an IgG₁ Fc region, an IgG₂ Fc region, an IgG₃ Fc region, an IgG₄ Fc region, an IgM Fc region, an IgA Fc region, an IgD Fc region and an IgE Fc region. (See, e.g., Padlan, Molecular Immunology, 31(3), 169-217 (1993)). Because there is some variation between immunoglobulins, and solely for clarity, Fc monomer refers to the last two heavy chain constant region immunoglobulin domains of IgA, IgD, and IgG, and the last three heavy chain constant region immunoglobulin domains of IgE and IgM. As mentioned, the Fc monomer can also include the flexible hinge N-terminal to these domains. For IgA and IgM, the Fc monomer may include the J chain. For IgG, the Fc portion comprises immunoglobulin domains CH2 and CH3 and the hinge between the first two domains and CH2. Although the boundaries of the Fc portion may vary an example for a human IgG heavy chain Fc portion comprising a functional hinge, CH2 and CH3 domain can be defined e.g. to comprise residues D231 (of the hinge domain— corresponding to D234 in Table 1 below) to P476, respectively L476 (for IgG₄) of the carboxyl-terminus of the CH3 domain, wherein the numbering is according to Kabat. The two Fc portion or Fc monomer, which are fused to each other via a peptide linker define the third domain of the antibody construct of the invention, which may also be defined as scFc domain.

[72] In one embodiment of the invention it is envisaged that a scFc domain as disclosed herein, respectively the Fc monomers fused to each other are comprised only in the third domain of the antibody construct.

In line with the present invention an IgG hinge region can be identified by analogy using the Kabat numbering as set forth in Table 1. In line with the above, the minimal requirement it is envisaged that for a hinge domain/region of the present invention in line with the present invention are comprises the amino acid residues corresponding to the IgG1 sequence stretch of D231 D234 to P243 according to the Kabat numbering. It is likewise envisaged that a hinge domain/region of the present invention comprises or consists of the IgG1 hinge sequence DKHTHTCPPCP (SEQ ID NO: 477) (corresponding to the stretch D234 to P243 as shown in Table 1 below – variations of said sequence are also envisaged provided that the hinge region still promotes dimerization). In a preferred embodiment of the invention the glycosylation site at Kabat position 314 of the CH2 domains in the third domain of the antibody construct is removed by a N314X substitution, wherein X is any amino acid excluding Q. Said substitution is preferably a N314G substitution. In a more preferred embodiment, said CH2 domain additionally comprises the following substitutions (position according to Kabat) V321C and R309C (these substitutions introduce the intra domain cysteine disulfide bridge at Kabat positions 309 and 321).

It is also envisaged that the third domain of the antibody construct of the invention comprises or consists in an amino to carboxyl order: DKHTHTCPPCP (SEQ ID NO: 477) (i.e. hinge) - CH2-CH3-linker- DKHTHTCPPCP (SEQ ID NO: 477) (i.e. hinge) -CH2-CH3. The peptide linker of the aforementioned antibody construct is in a preferred embodiment characterized by the amino acid sequence Gly-Gly-Gly-Gly-Ser, i.e. Gly4Ser (SEQ ID NO: 1), or polymers thereof, i.e. (Gly4Ser)_x, where x is an integer of 5 or greater (e.g. 5, 6, 7, 8 etc. or greater), 6 being preferred ((Gly4Ser)₆). Said construct may further comprise the aforementioned substitutions N314X, preferably N314G, and/or the further substitutions V321C and R309C. In a preferred embodiment of the antibody constructs of the invention as defined herein before, it is envisaged that the second domain binds to an extracellular epitope of the human and/or the *Macaca* CD3ε chain. Table 1: Kabat numbering of the amino acid residues of the hinge region

IMGT numbering for the hinge	IgG ₁ amino acid translation	Kabat numbering
1	(E)	226
2	P	227
3	K	228
4	S	232
5	C	233

6	D	234
7	K	235
8	T	236
9	H	237
10	T	238
11	C	239
12	P	240
13	P	241
14	C	242
15	P	243

In further embodiments of the present invention, the hinge domain/region comprises or consists of the IgG2 subtype hinge sequence ERKCCVECPPCP (SEQ ID NO: 478), the IgG3 subtype hinge sequence ELKTPLDTTHTCPRCP (SEQ ID NO: 479) or ELKTPLGDTTHTCPRCP (SEQ ID NO: 486), and/or the IgG4 subtype hinge sequence ESKYGPPCPSCP (SEQ ID NO: 480). The IgG1 subtype hinge sequence may be the following one EPKSCDKTHTCPPCP (as shown in Table 1 and SEQ ID NO: 487). These core hinge regions are thus also envisaged in the context of the present invention.

[73] The location and sequence of the IgG CH2 and IgG CD3 domain can be identified by analogy using the Kabat numbering as set forth in Table 2:

Table 2: Kabat numbering of the amino acid residues of the IgG CH2 and CH3 region

IgG subtype	CH2 aa translation	CH2 Kabat numbering	CH3 aa translation	CH3 Kabat numbering
IgG₁	APE... ...KAK	244... ...360	GQP..... PGK	361... ...478
IgG₂	APP... ...KTK	244... ...360	GQP..... PGK	361... ...478
IgG₃	APE... ...KTK	244... ...360	GQP..... PGK	361... ...478
IgG₄	APE... ...KAK	244... ...360	GQP..... LGK	361... ...478

[74] In one embodiment of the invention the emphasized bold amino acid residues in the CH3 domain of the first or both Fc monomers are deleted.

[75] The peptide linker, by whom the polypeptide monomers ("Fc portion" or "Fc monomer") of the third domain are fused to each other, preferably comprises at least 25 amino acid residues (25, 26, 27, 28, 29, 30 etc.). More preferably, this peptide linker comprises at least 30 amino acid residues (30, 31, 32, 33, 34, 35 etc.). It is also preferred that the linker comprises up to 40 amino acid residues, more preferably up to 35 amino acid residues, most preferably exactly 30 amino acid residues. A preferred embodiment of such peptide linker is characterized by the amino acid sequence Gly-Gly-Gly-Gly-Ser, i.e. Gly₄Ser

(SEQ ID NO: 1), or polymers thereof, i.e. (Gly₄Ser)_x, where x is an integer of 5 or greater (e.g. 6, 7 or 8). Preferably the integer is 6 or 7, more preferably the integer is 6.

[76] In the event that a linker is used to fuse the first domain to the second domain, or the first or second domain to the third domain, this linker is preferably of a length and sequence sufficient to ensure that each of the first and second domains can, independently from one another, retain their differential binding specificities. For peptide linkers which connect the at least two binding domains (or two variable domains) in the antibody construct of the invention, those peptide linkers are preferred which comprise only a few number of amino acid residues, e.g. 12 amino acid residues or less. Thus, peptide linkers of 12, 11, 10, 9, 8, 7, 6 or 5 amino acid residues are preferred. An envisaged peptide linker with less than 5 amino acids comprises 4, 3, 2 or one amino acid(s), wherein Gly-rich linkers are preferred. A preferred embodiment of the peptide linker for a fusion the first and the second domain is depicted in SEQ ID NO:1. A preferred linker embodiment of the peptide linker for fusing the second and the third domain is a (Gly)₄-linker, also called G₄-linker.

[77] A particularly preferred "single" amino acid in the context of one of the above described "peptide linker" is Gly. Accordingly, said peptide linker may consist of the single amino acid Gly. In a preferred embodiment of the invention a peptide linker is characterized by the amino acid sequence Gly-Gly-Gly-Gly-Ser, i.e. Gly₄Ser (SEQ ID NO: 1), or polymers thereof, i.e. (Gly₄Ser)_x, where x is an integer of 1 or greater (e.g. 2 or 3). Preferred linkers are depicted in SEQ ID NOs: 1 to 12. The characteristics of said peptide linker, which comprise the absence of the promotion of secondary structures, are known in the art and are described e.g. in Dall'Acqua et al. (Biochem. (1998) 37, 9266-9273), Cheadle et al. (Mol Immunol (1992) 29, 21-30) and Raag and Whitlow (FASEB (1995) 9(1), 73-80). Peptide linkers which furthermore do not promote any secondary structures are preferred. The linkage of said domains to each other can be provided, e.g., by genetic engineering, as described in the examples. Methods for preparing fused and operatively linked bispecific single chain constructs and expressing them in mammalian cells or bacteria are well-known in the art (e.g. WO 99/54440 or Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 2001).

[78] In a preferred embodiment of the antibody construct or the present invention the first and second domain form an antibody construct in a format selected from the group consisting of (scFv)₂, scFv-single domain mAb, diabody and oligomers of any of the those formats.

[79] According to a particularly preferred embodiment, and as documented in the appended examples, the first and the second domain of the antibody construct of the

invention is a “bispecific single chain antibody construct”, more preferably a bispecific “single chain Fv” (scFv). Although the two domains of the Fv fragment, VL and VH, are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker – as described hereinbefore – that enables them to be made as a single protein chain in which the VL and VH regions pair to form a monovalent molecule; see *e.g.*, Huston et al. (1988) Proc. Natl. Acad. Sci. USA 85:5879-5883). These antibody fragments are obtained using conventional techniques known to those with skill in the art, and the fragments are evaluated for function in the same manner as are whole or full-length antibodies. A single-chain variable fragment (scFv) is hence a fusion protein of the variable region of the heavy chain (VH) and of the light chain (VL) of immunoglobulins, usually connected with a short linker peptide of about ten to about 25 amino acids, preferably about 15 to 20 amino acids. The linker is usually rich in glycine for flexibility, as well as serine or threonine for solubility, and can either connect the N-terminus of the VH with the C-terminus of the VL, or *vice versa*. This protein retains the specificity of the original immunoglobulin, despite removal of the constant regions and introduction of the linker.

[80] Bispecific single chain antibody constructs are known in the art and are described in WO 99/54440, Mack, J. Immunol. (1997), 158, 3965-3970, Mack, PNAS, (1995), 92, 7021-7025, Kufer, Cancer Immunol. Immunother., (1997), 45, 193-197, Löffler, Blood, (2000), 95, 6, 2098-2103, Brühl, Immunol., (2001), 166, 2420-2426, Kipriyanov, J. Mol. Biol., (1999), 293, 41-56. Techniques described for the production of single chain antibodies (see, *inter alia*, US Patent 4,946,778, Kontermann and Dübel (2010), *loc. cit.* and Little (2009), *loc. cit.*) can be adapted to produce single chain antibody constructs specifically recognizing (an) elected target(s).

[81] Bivalent (also called divalent) or bispecific single-chain variable fragments (bi-scFvs or di-scFvs having the format (scFv)₂ can be engineered by linking two scFv molecules (*e.g.* with linkers as described hereinbefore). If these two scFv molecules have the same binding specificity, the resulting (scFv)₂ molecule will preferably be called bivalent (*i.e.* it has two valences for the same target epitope). If the two scFv molecules have different binding specificities, the resulting (scFv)₂ molecule will preferably be called bispecific. The linking can be done by producing a single peptide chain with two VH regions and two VL regions, yielding tandem scFvs (see *e.g.* Kufer P. *et al.*, (2004) Trends in Biotechnology 22(5):238-244). Another possibility is the creation of scFv molecules with linker peptides that are too short for the two variable regions to fold together (*e.g.* about five amino acids), forcing the scFvs to dimerize. This type is known as diabodies (see *e.g.* Hollinger, Philipp *et al.*, (July 1993) Proceedings of the National Academy of Sciences of the United States of America 90 (14): 6444-8).

[82] In line with this invention either the first, the second or the first and the second domain may comprise a single domain antibody, respectively the variable domain or at least the CDRs of a single domain antibody. Single domain antibodies comprise merely one (monomeric) antibody variable domain which is able to bind selectively to a specific antigen, independently of other V regions or domains. The first single domain antibodies were engineered from heavy chain antibodies found in camelids, and these are called V_HH fragments. Cartilaginous fishes also have heavy chain antibodies (IgNAR) from which single domain antibodies called V_{NAR} fragments can be obtained. An alternative approach is to split the dimeric variable domains from common immunoglobulins e.g. from humans or rodents into monomers, hence obtaining V_H or V_L as a single domain Ab. Although most research into single domain antibodies is currently based on heavy chain variable domains, nanobodies derived from light chains have also been shown to bind specifically to target epitopes. Examples of single domain antibodies are called sdAb, nanobodies or single variable domain antibodies.

[83] A (single domain mAb)₂ is hence a monoclonal antibody construct composed of (at least) two single domain monoclonal antibodies, which are individually selected from the group comprising V_H, V_L, V_HH and V_{NAR}. The linker is preferably in the form of a peptide linker. Similarly, an “scFv-single domain mAb” is a monoclonal antibody construct composed of at least one single domain antibody as described above and one scFv molecule as described above. Again, the linker is preferably in the form of a peptide linker.

[84] Whether or not an antibody construct competes for binding with another given antibody construct can be measured in a competition assay such as a competitive ELISA or a cell-based competition assay. Avidin-coupled microparticles (beads) can also be used. Similar to an avidin-coated ELISA plate, when reacted with a biotinylated protein, each of these beads can be used as a substrate on which an assay can be performed. Antigen is coated onto a bead and then precoated with the first antibody. The second antibody is added and any additional binding is determined. Possible means for the read-out includes flow cytometry.

[85] T cells or T lymphocytes are a type of lymphocyte (itself a type of white blood cell) that play a central role in cell-mediated immunity. There are several subsets of T cells, each with a distinct function. T cells can be distinguished from other lymphocytes, such as B cells and NK cells, by the presence of a T cell receptor (TCR) on the cell surface. The TCR is responsible for recognizing antigens bound to major histocompatibility complex (MHC) molecules and is composed of two different protein chains. In 95% of the T cells, the TCR consists of an alpha (α) and beta (β) chain. When the TCR engages with antigenic peptide and MHC (peptide / MHC complex), the T lymphocyte is activated through a series of

biochemical events mediated by associated enzymes, co-receptors, specialized adaptor molecules, and activated or released transcription factors.

[86] The CD3 receptor complex is a protein complex and is composed of four chains. In mammals, the complex contains a CD3 γ (gamma) chain, a CD3 δ (delta) chain, and two CD3 ϵ (epsilon) chains. These chains associate with the T cell receptor (TCR) and the so-called ζ (zeta) chain to form the T cell receptor CD3 complex and to generate an activation signal in T lymphocytes. The CD3 γ (gamma), CD3 δ (delta), and CD3 ϵ (epsilon) chains are highly related cell-surface proteins of the immunoglobulin superfamily containing a single extracellular immunoglobulin domain. The intracellular tails of the CD3 molecules contain a single conserved motif known as an immunoreceptor tyrosine-based activation motif or ITAM for short, which is essential for the signaling capacity of the TCR. The CD3 epsilon molecule is a polypeptide which in humans is encoded by the *CD3E* gene which resides on chromosome 11. The most preferred epitope of CD3 epsilon is comprised within amino acid residues 1-27 of the human CD3 epsilon extracellular domain. It is envisaged that antibody constructs according to the present invention typically and advantageously show less unspecific T cell activation, which is not desired in specific immunotherapy. This translates to a reduced risk of side effects.

[87] The redirected lysis of target cells via the recruitment of T cells by a multispecific, at least bispecific, antibody construct involves cytolytic synapse formation and delivery of perforin and granzymes. The engaged T cells are capable of serial target cell lysis, and are not affected by immune escape mechanisms interfering with peptide antigen processing and presentation, or clonal T cell differentiation; see, for example, WO 2007/042261.

[88] Cytotoxicity mediated by antibody constructs of the invention can be measured in various ways. Effector cells can be e.g. stimulated enriched (human) CD8 positive T cells or unstimulated (human) peripheral blood mononuclear cells (PBMC). If the target cells are of macaque origin or express or are transfected with macaque PSMA which is bound by the first domain, the effector cells should also be of macaque origin such as a macaque T cell line, e.g. 4119LnPx. The target cells should express (at least the extracellular domain of) PSMA, e.g. human or macaque PSMA. Target cells can be a cell line (such as CHO) which is stably or transiently transfected with PSMA, e.g. human or macaque PSMA. Usually EC₅₀ values are expected to be lower with target cell lines expressing higher levels of PSMA on the cell surface. The effector to target cell (E:T) ratio is usually about 10:1, but can also vary. Cytotoxic activity of PSMAxCD3 bispecific antibody constructs can be measured in a ⁵¹Cr-release assay (incubation time of about 18 hours) or in a FACS-based cytotoxicity assay (incubation time of about 48 hours). Modifications of the assay incubation time (cytotoxic reaction) are also possible. Other methods of measuring cytotoxicity are well-known to the

skilled person and comprise MTT or MTS assays, ATP-based assays including bioluminescent assays, the sulforhodamine B (SRB) assay, WST assay, clonogenic assay and the ECIS technology.

[89] The cytotoxic activity mediated by PSMAxCD3 bispecific antibody constructs of the present invention is preferably measured in a cell-based cytotoxicity assay. It may also be measured in a ^{51}Cr -release assay. It is represented by the EC_{50} value, which corresponds to the half maximal effective concentration (concentration of the antibody construct which induces a cytotoxic response halfway between the baseline and maximum). Preferably, the EC_{50} value of the PSMAxCD3 bispecific antibody constructs is ≤ 5000 pM or ≤ 4000 pM, more preferably ≤ 3000 pM or ≤ 2000 pM, even more preferably ≤ 1000 pM or ≤ 500 pM, even more preferably ≤ 400 pM or ≤ 300 pM, even more preferably ≤ 200 pM, even more preferably ≤ 100 pM, even more preferably ≤ 50 pM, even more preferably ≤ 20 pM or ≤ 10 pM, and most preferably ≤ 5 pM.

[90] The above given EC_{50} values can be measured in different assays. The skilled person is aware that an EC_{50} value can be expected to be lower when stimulated / enriched CD8^+ T cells are used as effector cells, compared with unstimulated PBMC. It can furthermore be expected that the EC_{50} values are lower when the target cells express a high number of PSMA compared with a low target expression rat. For example, when stimulated / enriched human CD8^+ T cells are used as effector cells (and either PSMA transfected cells such as CHO cells or PSMA positive human cell lines are used as target cells), the EC_{50} value of the PSMAxCD3 bispecific antibody construct is preferably ≤ 1000 pM, more preferably ≤ 500 pM, even more preferably ≤ 250 pM, even more preferably ≤ 100 pM, even more preferably ≤ 50 pM, even more preferably ≤ 10 pM, and most preferably ≤ 5 pM. When human PBMCs are used as effector cells, the EC_{50} value of the PSMAxCD3 bispecific antibody construct is preferably ≤ 5000 pM or ≤ 4000 pM (in particular when the target cells are PSMA positive human cell lines), more preferably ≤ 2000 pM (in particular when the target cells are PSMA transfected cells such as CHO cells), more preferably ≤ 1000 pM or ≤ 500 pM, even more preferably ≤ 200 pM, even more preferably ≤ 150 pM, even more preferably ≤ 100 pM, and most preferably ≤ 50 pM, or lower. When a macaque T cell line such as LnPx4119 is used as effector cells, and a macaque PSMA transfected cell line such as CHO cells is used as target cell line, the EC_{50} value of the PSMAxCD3 bispecific antibody construct is preferably ≤ 2000 pM or ≤ 1500 pM, more preferably ≤ 1000 pM or ≤ 500 pM, even more preferably ≤ 300 pM or ≤ 250 pM, even more preferably ≤ 100 pM, and most preferably ≤ 50 pM.

[91] Preferably, the PSMAxCD3 bispecific antibody constructs of the present invention do not induce / mediate lysis or do not essentially induce / mediate lysis of PSMA negative cells such as CHO cells. The term “do not induce lysis”, “do not essentially induce lysis”, “do not

mediate lysis” or “do not essentially mediate lysis” means that an antibody construct of the present invention does not induce or mediate lysis of more than 30%, preferably not more than 20%, more preferably not more than 10%, particularly preferably not more than 9%, 8%, 7%, 6% or 5% of PSMA negative cells, whereby lysis of a PSMA positive human cell line is set to be 100%. This usually applies for concentrations of the antibody construct of up to 500 nM. The skilled person knows how to measure cell lysis without further ado. Moreover, the present specification teaches specific instructions how to measure cell lysis.

[92] The difference in cytotoxic activity between the monomeric and the dimeric isoform of individual PSMAxCD3 bispecific antibody constructs is referred to as “potency gap”. This potency gap can e.g. be calculated as ratio between EC_{50} values of the molecule's monomeric and dimeric form. Potency gaps of the PSMAxCD3 bispecific antibody constructs of the present invention are preferably ≤ 5 , more preferably ≤ 4 , even more preferably ≤ 3 , even more preferably ≤ 2 and most preferably ≤ 1 .

[93] The first and/or the second (or any further) binding domain(s) of the antibody construct of the invention is/are preferably cross-species specific for members of the mammalian order of primates. Cross-species specific CD3 binding domains are, for example, described in WO 2008/119567. According to one embodiment, the first and/or second binding domain, in addition to binding to human PSMA and human CD3, respectively, will also bind to PSMA / CD3 of primates including (but not limited to) new world primates (such as *Callithrix jacchus*, *Saguinus Oedipus* or *Saimiri sciureus*), old world primates (such as baboons and macaques), gibbons, and non-human *homininae*.

[94] In one embodiment of the antibody construct of the invention the first domain binds to human PSMA and further binds to macaque PSMA, such as PSMA of *Macaca fascicularis*, and more preferably, to macaque PSMA expressed on the surface macaque cells. The affinity of the first domain for PSMA, preferably for human PSMA, is preferably ≤ 100 nM or ≤ 50 nM, more preferably ≤ 25 nM or ≤ 20 nM, more preferably ≤ 15 nM or ≤ 10 nM, even more preferably ≤ 5 nM, even more preferably ≤ 2.5 nM or ≤ 2 nM, even more preferably ≤ 1 nM, even more preferably ≤ 0.6 nM, even more preferably ≤ 0.5 nM, and most preferably ≤ 0.4 nM. The affinity can be measured for example in a BIAcore assay or in a Scatchard assay. Other methods of determining the affinity are also well-known to the skilled person. The affinity of the first domain for macaque PSMA is preferably ≤ 15 nM, more preferably ≤ 10 nM, even more preferably ≤ 5 nM, even more preferably ≤ 1 nM, even more preferably ≤ 0.5 nM, even more preferably ≤ 0.1 nM, and most preferably ≤ 0.05 nM or even ≤ 0.01 nM.

[95] Preferably the affinity gap of the antibody constructs according to the invention for binding macaque PSMA versus human PSMA [ma PSMA : hu PSMA] (as determined e.g. by

BiaCore or by Scatchard analysis) is <100, preferably <20, more preferably <15, further preferably <10, even more preferably <8, more preferably <6 and most preferably <2. Preferred ranges for the affinity gap of the antibody constructs according to the invention for binding macaque PSMA versus human PSMA are between 0.1 and 20, more preferably
 5 between 0.2 and 10, even more preferably between 0.3 and 6, even more preferably between 0.5 and 3 or between 0.5 and 2.5, and most preferably between 0.5 and 2 or between 0.6 and 2.

[96] The second domain of the antibody construct of the invention binds to human CD3 epsilon and/or to *Macaca* CD3 epsilon. In a preferred embodiment the second domain further
 10 binds to *Callithrix jacchus*, *Saguinus Oedipus* or *Saimiri sciureus* CD3 epsilon. *Callithrix jacchus* and *Saguinus oedipus* are both new world primate belonging to the family of *Callitrichidae*, while *Saimiri sciureus* is a new world primate belonging to the family of *Cebidae*.

[97] It is preferred for the antibody construct of the present invention that the second
 15 domain which binds to an extracellular epitope of the human and/or the *Macaca* CD3 epsilon chain comprises a VL region comprising CDR-L1, CDR-L2 and CDR-L3 selected from:

(a) CDR-L1 as depicted in SEQ ID NO: 27 of WO 2008/119567, CDR-L2 as depicted in SEQ ID NO: 28 of WO 2008/119567 and CDR-L3 as depicted in SEQ ID NO: 29 of WO 2008/119567;

20 (b) CDR-L1 as depicted in SEQ ID NO: 117 of WO 2008/119567, CDR-L2 as depicted in SEQ ID NO: 118 of WO 2008/119567 and CDR-L3 as depicted in SEQ ID NO: 119 of WO 2008/119567; and

(c) CDR-L1 as depicted in SEQ ID NO: 153 of WO 2008/119567, CDR-L2 as depicted in SEQ ID NO: 154 of WO 2008/119567 and CDR-L3 as depicted in SEQ ID NO: 155 of
 25 WO 2008/119567.

[98] In a furthermore preferred embodiment of the antibody construct of the present invention, the second domain which binds to an extracellular epitope of the human and/or the *Macaca* CD3 epsilon chain comprises a VH region comprising CDR-H 1, CDR-H2 and CDR-H3 selected from:

30 (a) CDR-H1 as depicted in SEQ ID NO: 12 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 13 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 14 of WO 2008/119567;

(b) CDR-H1 as depicted in SEQ ID NO: 30 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 31 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 32 of
 35 WO 2008/119567;

(c) CDR-H1 as depicted in SEQ ID NO: 48 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 49 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 50 of WO 2008/119567;

(d) CDR-H1 as depicted in SEQ ID NO: 66 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 67 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 68 of WO 2008/119567;

(e) CDR-H1 as depicted in SEQ ID NO: 84 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 85 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 86 of WO 2008/119567;

(f) CDR-H1 as depicted in SEQ ID NO: 102 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 103 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 104 of WO 2008/119567;

(g) CDR-H1 as depicted in SEQ ID NO: 120 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 121 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 122 of WO 2008/119567;

(h) CDR-H1 as depicted in SEQ ID NO: 138 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 139 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 140 of WO 2008/119567;

(i) CDR-H1 as depicted in SEQ ID NO: 156 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 157 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 158 of WO 2008/119567; and

(j) CDR-H1 as depicted in SEQ ID NO: 174 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 175 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 176 of WO 2008/119567.

[99] In a preferred embodiment of the antibody construct of the invention the above described three groups of VL CDRs are combined with the above described ten groups of VH CDRs within the second binding domain to form (30) groups, each comprising CDR-L 1-3 and CDR-H 1-3.

[100] It is preferred for the antibody construct of the present invention that the second domain which binds to CD3 comprises a VL region selected from the group consisting of those depicted in SEQ ID NOs: 17, 21, 35, 39, 53, 57, 71, 75, 89, 93, 107, 111, 125, 129, 143, 147, 161, 165, 179 or 183 of WO 2008/119567 or as depicted in SEQ ID NO: 13.

[101] It is also preferred that the second domain which binds to CD3 comprises a VH region selected from the group consisting of those depicted in SEQ ID NO: 15, 19, 33, 37, 51, 55, 69, 73, 87, 91, 105, 109, 123, 127, 141, 145, 159, 163, 177 or 181 of WO 2008/119567 or as depicted in SEQ ID NO: 14.

[102] More preferably, the antibody construct of the present invention is characterized by a second domain which binds to CD3 comprising a VL region and a VH region selected from the group consisting of:

- (a) a VL region as depicted in SEQ ID NO: 17 or 21 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 15 or 19 of WO 2008/119567;
- (b) a VL region as depicted in SEQ ID NO: 35 or 39 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 33 or 37 of WO 2008/119567;
- (c) a VL region as depicted in SEQ ID NO: 53 or 57 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 51 or 55 of WO 2008/119567;
- (d) a VL region as depicted in SEQ ID NO: 71 or 75 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 69 or 73 of WO 2008/119567;
- (e) a VL region as depicted in SEQ ID NO: 89 or 93 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 87 or 91 of WO 2008/119567;
- (f) a VL region as depicted in SEQ ID NO: 107 or 111 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 105 or 109 of WO 2008/119567;
- (g) a VL region as depicted in SEQ ID NO: 125 or 129 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 123 or 127 of WO 2008/119567;
- (h) a VL region as depicted in SEQ ID NO: 143 or 147 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 141 or 145 of WO 2008/119567;
- (i) a VL region as depicted in SEQ ID NO: 161 or 165 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 159 or 163 of WO 2008/119567; and
- (j) a VL region as depicted in SEQ ID NO: 179 or 183 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 177 or 181 of WO 2008/119567.

[103] Also preferred in connection with the antibody construct of the present invention is a second domain which binds to CD3 comprising a VL region as depicted in SEQ ID NO: 13 and a VH region as depicted in SEQ ID NO: 14.

[104] According to a preferred embodiment of the antibody construct of the present invention, the first and/or the second domain have the following format: The pairs of VH regions and VL regions are in the format of a single chain antibody (scFv). The VH and VL regions are arranged in the order VH-VL or VL-VH. It is preferred that the VH-region is positioned N-terminally of a linker sequence, and the VL-region is positioned C-terminally of the linker sequence.

[105] A preferred embodiment of the above described antibody construct of the present invention is characterized by the second domain which binds to CD3 comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 23, 25, 41, 43, 59, 61, 77,

79, 95, 97, 113, 115, 131, 133, 149, 151, 167, 169, 185 or 187 of WO 2008/119567 or as depicted in SEQ ID NO: 15.

[106] It is also envisaged that the first binding domain of the antibody construct of the invention comprises a VL region comprising CDR-L1, CDR-L2 and CDR-L3, and a VH region comprising CDR-H1, CDR-H2 and CDR-3 selected from the group consisting of:

(a) CDR-L1 as depicted in SEQ ID NO: 45, CDR-L2 as depicted in SEQ ID NO: 46, CDR-L3 as depicted in SEQ ID NO: 47, CDR-H1 as depicted in SEQ ID NO: 42, CDR-H2 as depicted in SEQ ID NO: 43, and CDR-H3 as depicted in SEQ ID NO: 44

(b) CDR-L1 as depicted in SEQ ID NO: 63, CDR-L2 as depicted in SEQ ID NO: 64, CDR-L3 as depicted in SEQ ID NO: 65, CDR-H1 as depicted in SEQ ID NO: 60, CDR-H2 as depicted in SEQ ID NO: 61, and CDR-H3 as depicted in SEQ ID NO: 62;

(c) CDR-L1 as depicted in SEQ ID NO: 81, CDR-L2 as depicted in SEQ ID NO: 82, CDR-L3 as depicted in SEQ ID NO: 83, CDR-H1 as depicted in SEQ ID NO: 78, CDR-H2 as depicted in SEQ ID NO: 79, and CDR-H3 as depicted in SEQ ID NO: 80;

(d) CDR-L1 as depicted in SEQ ID NO: 99, CDR-L2 as depicted in SEQ ID NO: 100, CDR-L3 as depicted in SEQ ID NO: 101, CDR-H1 as depicted in SEQ ID NO: 96, CDR-H2 as depicted in SEQ ID NO: 97, and CDR-H3 as depicted in SEQ ID NO: 98;

(e) CDR-L1 as depicted in SEQ ID NO: 117, CDR-L2 as depicted in SEQ ID NO: 118, CDR-L3 as depicted in SEQ ID NO: 119, CDR-H1 as depicted in SEQ ID NO: 114, CDR-H2 as depicted in SEQ ID NO: 115, and CDR-H3 as depicted in SEQ ID NO: 116;

(f) CDR-L1 as depicted in SEQ ID NO: 135, CDR-L2 as depicted in SEQ ID NO: 136, CDR-L3 as depicted in SEQ ID NO: 137, CDR-H1 as depicted in SEQ ID NO: 132, CDR-H2 as depicted in SEQ ID NO: 133, and CDR-H3 as depicted in SEQ ID NO: 134;

(g) CDR-L1 as depicted in SEQ ID NO: 153, CDR-L2 as depicted in SEQ ID NO: 154, CDR-L3 as depicted in SEQ ID NO: 155, CDR-H1 as depicted in SEQ ID NO: 150, CDR-H2 as depicted in SEQ ID NO: 151, and CDR-H3 as depicted in SEQ ID NO: 152;

(h) CDR-L1 as depicted in SEQ ID NO: 171, CDR-L2 as depicted in SEQ ID NO: 172, CDR-L3 as depicted in SEQ ID NO: 173, CDR-H1 as depicted in SEQ ID NO: 168, CDR-H2 as depicted in SEQ ID NO: 169, and CDR-H3 as depicted in SEQ ID NO: 170;

- (i) CDR-L1 as depicted in SEQ ID NO: 189, CDR-L2 as depicted in SEQ ID NO: 190
CDR-L3 as depicted in SEQ ID NO: 191, CDR-H1 as depicted in SEQ ID NO: 186, CDR-H2
as depicted in SEQ ID NO: 187, and CDR-H3 as depicted in SEQ ID NO: 188;
- (j) CDR-L1 as depicted in SEQ ID NO: 207, CDR-L2 as depicted in SEQ ID NO: 208
5 CDR-L3 as depicted in SEQ ID NO: 209, CDR-H1 as depicted in SEQ ID NO: 204, CDR-H2
as depicted in SEQ ID NO: 205, and CDR-H3 as depicted in SEQ ID NO: 206;
- (k) CDR-L1 as depicted in SEQ ID NO: 225, CDR-L2 as depicted in SEQ ID NO: 226
CDR-L3 as depicted in SEQ ID NO: 227, CDR-H1 as depicted in SEQ ID NO: 222, CDR-H2
as depicted in SEQ ID NO: 223, and CDR-H3 as depicted in SEQ ID NO: 224;
- 10 (l) CDR-L1 as depicted in SEQ ID NO: 243, CDR-L2 as depicted in SEQ ID NO: 244
CDR-L3 as depicted in SEQ ID NO: 245, CDR-H1 as depicted in SEQ ID NO: 240, CDR-H2
as depicted in SEQ ID NO: 241, and CDR-H3 as depicted in SEQ ID NO: 242;
- (m) CDR-L1 as depicted in SEQ ID NO: 261, CDR-L2 as depicted in SEQ ID NO: 262
CDR-L3 as depicted in SEQ ID NO: 263, CDR-H1 as depicted in SEQ ID NO: 258, CDR-H2
15 as depicted in SEQ ID NO: 259, and CDR-H3 as depicted in SEQ ID NO: 260;
- (n) CDR-L1 as depicted in SEQ ID NO: 279, CDR-L2 as depicted in SEQ ID NO: 280
CDR-L3 as depicted in SEQ ID NO: 281, CDR-H1 as depicted in SEQ ID NO: 276, CDR-H2
as depicted in SEQ ID NO: 277, and CDR-H3 as depicted in SEQ ID NO: 278; and
- (o) CDR-L1 as depicted in SEQ ID NO: 297, CDR-L2 as depicted in SEQ ID NO: 298,
20 CDR-L3 as depicted in SEQ ID NO: 299, CDR-H1 as depicted in SEQ ID NO: 294, CDR-H2
as depicted in SEQ ID NO: 295, and CDR-H3 as depicted in SEQ ID NO: 296.
- (p) CDR-L1 as depicted in SEQ ID NO: 315, CDR-L2 as depicted in SEQ ID NO: 316,
CDR-L3 as depicted in SEQ ID NO: 317, CDR-H1 as depicted in SEQ ID NO: 312, CDR-H2
as depicted in SEQ ID NO: 313, and CDR-H3 as depicted in SEQ ID NO: 314.
- 25 (q) CDR-L1 as depicted in SEQ ID NO: 330, CDR-L2 as depicted in SEQ ID NO: 331,
CDR-L3 as depicted in SEQ ID NO: 332, CDR-H1 as depicted in SEQ ID NO: 327, CDR-H2
as depicted in SEQ ID NO: 328, and CDR-H3 as depicted in SEQ ID NO: 329.
- (r) CDR-L1 as depicted in SEQ ID NO: 345, CDR-L2 as depicted in SEQ ID NO: 346,
CDR-L3 as depicted in SEQ ID NO: 347, CDR-H1 as depicted in SEQ ID NO: 342, CDR-H2
30 as depicted in SEQ ID NO: 343, and CDR-H3 as depicted in SEQ ID NO: 344.

- (s) CDR-L1 as depicted in SEQ ID NO: 360, CDR-L2 as depicted in SEQ ID NO: 361, CDR-L3 as depicted in SEQ ID NO: 362, CDR-H1 as depicted in SEQ ID NO: 357, CDR-H2 as depicted in SEQ ID NO: 358, and CDR-H3 as depicted in SEQ ID NO: 359.
- (t) CDR-L1 as depicted in SEQ ID NO: 375, CDR-L2 as depicted in SEQ ID NO: 376, CDR-L3 as depicted in SEQ ID NO: 377, CDR-H1 as depicted in SEQ ID NO: 372, CDR-H2 as depicted in SEQ ID NO: 373, and CDR-H3 as depicted in SEQ ID NO: 374.
- (u) CDR-L1 as depicted in SEQ ID NO: 390, CDR-L2 as depicted in SEQ ID NO: 391, CDR-L3 as depicted in SEQ ID NO: 392, CDR-H1 as depicted in SEQ ID NO: 387, CDR-H2 as depicted in SEQ ID NO: 388, and CDR-H3 as depicted in SEQ ID NO: 389.
- (v) CDR-L1 as depicted in SEQ ID NO: 405, CDR-L2 as depicted in SEQ ID NO: 406, CDR-L3 as depicted in SEQ ID NO: 407, CDR-H1 as depicted in SEQ ID NO: 402, CDR-H2 as depicted in SEQ ID NO: 403, and CDR-H3 as depicted in SEQ ID NO: 404.
- (w) CDR-L1 as depicted in SEQ ID NO: 420, CDR-L2 as depicted in SEQ ID NO: 421, CDR-L3 as depicted in SEQ ID NO: 422, CDR-H1 as depicted in SEQ ID NO: 417, CDR-H2 as depicted in SEQ ID NO: 418, and CDR-H3 as depicted in SEQ ID NO: 419.
- (x) CDR-L1 as depicted in SEQ ID NO: 435, CDR-L2 as depicted in SEQ ID NO: 436, CDR-L3 as depicted in SEQ ID NO: 437, CDR-H1 as depicted in SEQ ID NO: 432, CDR-H2 as depicted in SEQ ID NO: 433, and CDR-H3 as depicted in SEQ ID NO: 434.
- (y) CDR-L1 as depicted in SEQ ID NO: 450, CDR-L2 as depicted in SEQ ID NO: 451, CDR-L3 as depicted in SEQ ID NO: 452, CDR-H1 as depicted in SEQ ID NO: 447, CDR-H2 as depicted in SEQ ID NO: 448, and CDR-H3 as depicted in SEQ ID NO: 449.
- (z) CDR-L1 as depicted in SEQ ID NO: 465, CDR-L2 as depicted in SEQ ID NO: 466, CDR-L3 as depicted in SEQ ID NO: 467, CDR-H1 as depicted in SEQ ID NO: 462, CDR-H2 as depicted in SEQ ID NO: 463 and CDR-H3 as depicted in SEQ ID NO: 464.
- [107]** It is furthermore envisaged that the first binding domain of the antibody construct of the invention comprises a VH region and a VL region selected from the group consisting of:
- (a) a VH region as depicted in SEQ ID NO: 48 and a VL region as depicted in SEQ ID NO: 49;
- (b) a VH region as depicted in SEQ ID NO: 66 and a VL region as depicted in SEQ ID NO: 67;
- (c) a VH region as depicted in SEQ ID NO: 72 and a VL region as depicted in SEQ ID NO: 73;

- (d) a VH region as depicted in SEQ ID NO: 84 and a VL region as depicted in SEQ ID NO: 85;
- (e) a VH region as depicted in SEQ ID NO: 90 and a VL region as depicted in SEQ ID NO: 91;
- 5 (f) a VH region as depicted in SEQ ID NO: 102 and a VL region as depicted in SEQ ID NO: 103;
- (g) a VH region as depicted in SEQ ID NO: 108 and a VL region as depicted in SEQ ID NO: 109;
- 10 (h) a VH region as depicted in SEQ ID NO: 120 and a VL region as depicted in SEQ ID NO: 121;
- (i) a VH region as depicted in SEQ ID NO: 126 and a VL region as depicted in SEQ ID NO: 127;
- (j) a VH region as depicted in SEQ ID NO: 138 and a VL region as depicted in SEQ ID NO: 139;
- 15 (k) a VH region as depicted in SEQ ID NO: 144 and a VL region as depicted in SEQ ID NO: 145;
- (l) a VH region as depicted in SEQ ID NO: 156 and a VL region as depicted in SEQ ID NO: 157;
- 20 (m) a VH region as depicted in SEQ ID NO: 162 and a VL region as depicted in SEQ ID NO: 163;
- (n) a VH region as depicted in SEQ ID NO: 180 and a VL region as depicted in SEQ ID NO: 181;
- (o) a VH region as depicted in SEQ ID NO: 192 and a VL region as depicted in SEQ ID NO: 193;
- 25 (p) a VH region as depicted in SEQ ID NO: 198 and a VL region as depicted in SEQ ID NO: 199;
- (q) a VH region as depicted in SEQ ID NO: 210 and a VL region as depicted in SEQ ID NO: 211;
- 30 (r) a VH region as depicted in SEQ ID NO: 216 and a VL region as depicted in SEQ ID NO: 217;

- (s) a VH region as depicted in SEQ ID NO: 228 and a VL region as depicted in SEQ ID NO: 229;
- (t) a VH region as depicted in SEQ ID NO: 234 and a VL region as depicted in SEQ ID NO: 235;
- 5 (u) a VH region as depicted in SEQ ID NO: 246 and a VL region as depicted in SEQ ID NO: 247;
- (v) a VH region as depicted in SEQ ID NO: 252 and a VL region as depicted in SEQ ID NO: 253;
- (w) a VH region as depicted in SEQ ID NO: 264 and a VL region as depicted in SEQ ID NO: 265;
- 10 (x) a VH region as depicted in SEQ ID NO: 270 and a VL region as depicted in SEQ ID NO: 271;
- (y) a VH region as depicted in SEQ ID NO: 282 and a VL region as depicted in SEQ ID NO: 283;
- 15 (z) a VH region as depicted in SEQ ID NO: 288 and a VL region as depicted in SEQ ID NO: 289;
- (aa) a VH region as depicted in SEQ ID NO: 300 and a VL region as depicted in SEQ ID NO: 301;
- (ab) a VH region as depicted in SEQ ID NO: 306 and a VL region as depicted in SEQ ID NO: 307;
- 20 (ac) a VH region as depicted in SEQ ID NO: 54 and a VL region as depicted in SEQ ID NO: 55
- (ad) a VH region as depicted in SEQ ID NO: 174 and a VL region as depicted in SEQ ID NO: 175.
- 25 (ae) a VH region as depicted in SEQ ID NO: 318 and a VL region as depicted in SEQ ID NO: 319;
- (af) a VH region as depicted in SEQ ID NO: 333 and a VL region as depicted in SEQ ID NO: 334;

(ag) a VH region as depicted in SEQ ID NO: 348 and a VL region as depicted in SEQ ID NO: 349;

(ah) a VH region as depicted in SEQ ID NO: 363 and a VL region as depicted in SEQ ID NO: 364;

5 (ai) a VH region as depicted in SEQ ID NO: 378 and a VL region as depicted in SEQ ID NO: 379;

(aj) a VH region as depicted in SEQ ID NO: 393 and a VL region as depicted in SEQ ID NO: 394;

10 (ak) a VH region as depicted in SEQ ID NO: 408 and a VL region as depicted in SEQ ID NO: 409;

(al) a VH region as depicted in SEQ ID NO: 423 and a VL region as depicted in SEQ ID NO: 424;

(am) a VH region as depicted in SEQ ID NO: 438 and a VL region as depicted in SEQ ID NO: 439;

15 (an) a VH region as depicted in SEQ ID NO: 453 and a VL region as depicted in SEQ ID NO: 454; and

(ao) a VH region as depicted in SEQ ID NO: 468 and a VL region as depicted in SEQ ID NO: 469.

20 **[108]** It is furthermore envisaged that the first binding domain of the antibody construct of the invention comprises an amino acid sequence selected from the group consisting of those depicted in SEQ ID NOs: 50, 56, 68, 74, 86, 92, 104, 110, 122, 128, 140, 146, 158, 164, 176, 182, 194, 200, 212, 218, 230, 236, 248, 254, 266, 272, 284, 290, 302, 308, 320, 335, 350, 365, 380, 395, 410, 425, 440, 455, and 470.

25 **[109]** The invention further provides an antibody construct comprising or having an amino acid sequence selected from the group consisting of SEQ ID NO: 51, 57, 69, 75, 87, 93, 105, 111, 123, 129, 141, 147, 159, 165, 177, 183, 195, 201, 213, 219, 231, 237, 249, 255, 267, 273, 285, 291, 303, 309, 321, 324, 336, 339, 351, 354, 366, 369, 381, 384, 396, 399, 411, 414, 426, 429, 441, 444, 456, 459, 471 and 474.

30 **[110]** Covalent modifications of the antibody constructs are also included within the scope of this invention, and are generally, but not always, done post-translationally. For example,

several types of covalent modifications of the antibody construct are introduced into the molecule by reacting specific amino acid residues of the antibody construct with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C-terminal residues.

5 [111] Cysteinyl residues most commonly are reacted with α -haloacetates (and corresponding amines), such as chloroacetic acid or chloroacetamide, to give carboxymethyl or carboxyamidomethyl derivatives. Cysteinyl residues also are derivatized by reaction with
10 bromotrifluoroacetone, α -bromo- β -(5-imidozoyl)propionic acid, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl 2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4-nitrophenol, or chloro-7-nitrobenzo-2-oxa-1,3-diazole.

[112] Histidyl residues are derivatized by reaction with diethylpyrocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain. Para-bromophenacyl bromide also is useful; the reaction is preferably performed in 0.1 M sodium cacodylate at
15 pH 6.0. Lysinyl and amino terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysinyl residues. Other suitable reagents for derivatizing alpha-amino-containing residues include imidoesters such as methyl picolinimate; pyridoxal phosphate; pyridoxal; chloroborohydride; trinitrobenzenesulfonic acid; O-methylisourea; 2,4-pentanedione; and
20 transaminase-catalyzed reaction with glyoxylate.

[113] Arginyl residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3-butanedione, 1,2-cyclohexanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pKa of the guanidine functional group. Furthermore, these
25 reagents may react with the groups of lysine as well as the arginine epsilon-amino group.

[114] The specific modification of tyrosyl residues may be made, with particular interest in introducing spectral labels into tyrosyl residues by reaction with aromatic diazonium compounds or tetranitromethane. Most commonly, N-acetylimidazole and tetranitromethane are used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively. Tyrosyl
30 residues are iodinated using ^{125}I or ^{131}I to prepare labeled proteins for use in radioimmunoassay, the chloramine T method described above being suitable.

[115] Carboxyl side groups (aspartyl or glutamyl) are selectively modified by reaction with carbodiimides ($\text{R}'\text{---N}=\text{C}=\text{N}\text{---R}'$), where R and R' are optionally different alkyl groups, such as 1-cyclohexyl-3-(2-morpholinyl-4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-

dimethylpentyl) carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginyl and glutaminyl residues by reaction with ammonium ions.

5 **[116]** Derivatization with bifunctional agents is useful for crosslinking the antibody constructs of the present invention to a water-insoluble support matrix or surface for use in a variety of methods. Commonly used crosslinking agents include, *e.g.*, 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), and bifunctional maleimides such as bis-N-maleimido-1,8-octane. Derivatizing agents such as methyl-3-[(p-azidophenyl)dithio]propioimide yield
10 photoactivatable intermediates that are capable of forming crosslinks in the presence of light. Alternatively, reactive water-insoluble matrices such as cyanogen bromide-activated carbohydrates and the reactive substrates as described in U.S. Pat. Nos. 3,969,287; 3,691,016; 4,195,128; 4,247,642; 4,229,537; and 4,330,440 are employed for protein immobilization.

15 **[117]** Glutaminyl and asparaginyl residues are frequently deamidated to the corresponding glutamyl and aspartyl residues, respectively. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues falls within the scope of this invention.

20 **[118]** Other modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains (T. E. Creighton, *Proteins: Structure and Molecular Properties*, W. H. Freeman & Co., San Francisco, 1983, pp. 79-86), acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

25 **[119]** Another type of covalent modification of the antibody constructs included within the scope of this invention comprises altering the glycosylation pattern of the protein. As is known in the art, glycosylation patterns can depend on both the sequence of the protein (*e.g.*, the presence or absence of particular glycosylation amino acid residues, discussed below), or the host cell or organism in which the protein is produced. Particular expression systems are discussed below.

30 **[120]** Glycosylation of polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tri-peptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tri-

peptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylgalactosamine, galactose, or xylose, to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

5 **[121]** Addition of glycosylation sites to the antibody construct is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tri-peptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the starting sequence (for O-linked glycosylation sites). For ease, the amino acid sequence of an
10 antibody construct is preferably altered through changes at the DNA level, particularly by mutating the DNA encoding the polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

[122] Another means of increasing the number of carbohydrate moieties on the antibody construct is by chemical or enzymatic coupling of glycosides to the protein. These
15 procedures are advantageous in that they do not require production of the protein in a host cell that has glycosylation capabilities for N- and O-linked glycosylation. Depending on the coupling mode used, the sugar(s) may be attached to (a) arginine and histidine, (b) free carboxyl groups, (c) free sulfhydryl groups such as those of cysteine, (d) free hydroxyl groups such as those of serine, threonine, or hydroxyproline, (e) aromatic residues such as
20 those of phenylalanine, tyrosine, or tryptophan, or (f) the amide group of glutamine. These methods are described in WO 87/05330, and in Aplin and Wriston, 1981, CRC Crit. Rev. Biochem., pp. 259-306.

[123] Removal of carbohydrate moieties present on the starting antibody construct may be accomplished chemically or enzymatically. Chemical deglycosylation requires exposure of
25 the protein to the compound trifluoromethanesulfonic acid, or an equivalent compound. This treatment results in the cleavage of most or all sugars except the linking sugar (N-acetylglucosamine or N-acetylgalactosamine), while leaving the polypeptide intact. Chemical deglycosylation is described by Hakimuddin *et al.*, 1987, *Arch. Biochem. Biophys.* 259:52 and by Edge *et al.*, 1981, *Anal. Biochem.* 118:131. Enzymatic cleavage of carbohydrate
30 moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura *et al.*, 1987, *Meth. Enzymol.* 138:350. Glycosylation at potential glycosylation sites may be prevented by the use of the compound tunicamycin as described by Duskin *et al.*, 1982, *J. Biol. Chem.* 257:3105. Tunicamycin blocks the formation of protein-N-glycoside linkages.

[124] Other modifications of the antibody construct are also contemplated herein. For example, another type of covalent modification of the antibody construct comprises linking the antibody construct to various non-proteinaceous polymers, including, but not limited to, various polyols such as polyethylene glycol, polypropylene glycol, polyoxyalkylenes, or copolymers of polyethylene glycol and polypropylene glycol, in the manner set forth in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337. In addition, as is known in the art, amino acid substitutions may be made in various positions within the antibody construct, e.g. in order to facilitate the addition of polymers such as PEG.

[125] In some embodiments, the covalent modification of the antibody constructs of the invention comprises the addition of one or more labels. The labelling group may be coupled to the antibody construct *via* spacer arms of various lengths to reduce potential steric hindrance. Various methods for labelling proteins are known in the art and can be used in performing the present invention. The term “label” or “labelling group” refers to any detectable label. In general, labels fall into a variety of classes, depending on the assay in which they are to be detected – the following examples include, but are not limited to:

- a) isotopic labels, which may be radioactive or heavy isotopes, such as radioisotopes or radionuclides (e.g., ^3H , ^{14}C , ^{15}N , ^{35}S , ^{89}Zr , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I)
- b) magnetic labels (e.g., magnetic particles)
- c) redox active moieties
- d) optical dyes (including, but not limited to, chromophores, phosphors and fluorophores) such as fluorescent groups (e.g., FITC, rhodamine, lanthanide phosphors), chemiluminescent groups, and fluorophores which can be either “small molecule” fluoeres or proteinaceous fluoeres
- e) enzymatic groups (e.g. horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase)
- f) biotinylated groups
- g) predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sides for secondary antibodies, metal binding domains, epitope tags, etc.)

[126] By “fluorescent label” is meant any molecule that may be detected *via* its inherent fluorescent properties. Suitable fluorescent labels include, but are not limited to, fluorescein, rhodamine, tetramethylrhodamine, eosin, erythrosin, coumarin, methyl-coumarins, pyrene, Malacite green, stilbene, Lucifer Yellow, Cascade BlueJ, Texas Red, IAEDANS, EDANS, BODIPY FL, LC Red 640, Cy 5, Cy 5.5, LC Red 705, Oregon green, the Alexa-Fluor dyes (Alexa Fluor 350, Alexa Fluor 430, Alexa Fluor 488, Alexa Fluor 546, Alexa Fluor 568, Alexa Fluor 594, Alexa Fluor 633, Alexa Fluor 660, Alexa Fluor 680), Cascade Blue, Cascade

Yellow and R-phycoerythrin (PE) (Molecular Probes, Eugene, OR), FITC, Rhodamine, and Texas Red (Pierce, Rockford, IL), Cy5, Cy5.5, Cy7 (Amersham Life Science, Pittsburgh, PA). Suitable optical dyes, including fluorophores, are described in Molecular Probes Handbook by Richard P. Haugland.

5 **[127]** Suitable proteinaceous fluorescent labels also include, but are not limited to, green fluorescent protein, including a Renilla, Ptilosarcus, or Aequorea species of GFP (Chalfie *et al.*, 1994, *Science* 263:802-805), EGFP (Clontech Laboratories, Inc., Genbank Accession Number U55762), blue fluorescent protein (BFP, Quantum Biotechnologies, Inc. 1801 de
10 Maisonneuve Blvd. West, 8th Floor, Montreal, Quebec, Canada H3H 1J9; Stauber, 1998, *Biotechniques* 24:462-471; Heim *et al.*, 1996, *Curr. Biol.* 6:178-182), enhanced yellow fluorescent protein (EYFP, Clontech Laboratories, Inc.), luciferase (Ichiki *et al.*, 1993, *J. Immunol.* 150:5408-5417), β galactosidase (Nolan *et al.*, 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85:2603-2607) and Renilla (WO92/15673, WO95/07463, WO98/14605, WO98/26277, WO99/49019, U.S. Patent Nos. 5,292,658; 5,418,155; 5,683,888; 5,741,668; 5,777,079;
15 5,804,387; 5,874,304; 5,876,995; 5,925,558).

[128] The antibody construct of the invention may also comprise additional domains, which are *e.g.* helpful in the isolation of the molecule or relate to an adapted pharmacokinetic profile of the molecule. Domains helpful for the isolation of an antibody construct may be selected from peptide motives or secondarily introduced moieties, which can be captured in
20 an isolation method, *e.g.* an isolation column. Non-limiting embodiments of such additional domains comprise peptide motives known as Myc-tag, HAT-tag, HA-tag, TAP-tag, GST-tag, chitin binding domain (CBD-tag), maltose binding protein (MBP-tag), Flag-tag, Strep-tag and variants thereof (*e.g.* StreptII-tag) and His-tag. All herein disclosed antibody constructs may comprise a His-tag domain, which is generally known as a repeat of consecutive His
25 residues in the amino acid sequence of a molecule, preferably of five, and more preferably of six His residues (hexa-histidine). The His-tag may be located *e.g.* at the N- or C-terminus of the antibody construct, preferably it is located at the C-terminus. Most preferably, a hexa-histidine tag (HHHHHH) (SEQ ID NO:16) is linked via peptide bond to the C-terminus of the antibody construct according to the invention. Additionally, a conjugate system of PLGA-
30 PEG-PLGA may be combined with a poly-histidine tag for sustained release application and improved pharmacokinetic profile.

[129] Amino acid sequence modifications of the antibody constructs described herein are also contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody construct. Amino acid sequence variants of the
35 antibody constructs are prepared by introducing appropriate nucleotide changes into the antibody constructs nucleic acid, or by peptide synthesis. All of the below described amino

acid sequence modifications should result in an antibody construct which still retains the desired biological activity (binding to PSMA and to CD3) of the unmodified parental molecule.

[130] The term “amino acid” or “amino acid residue” typically refers to an amino acid having its art recognized definition such as an amino acid selected from the group consisting of: alanine (Ala or A); arginine (Arg or R); asparagine (Asn or N); aspartic acid (Asp or D); cysteine (Cys or C); glutamine (Gln or Q); glutamic acid (Glu or E); glycine (Gly or G); histidine (His or H); isoleucine (Ile or I); leucine (Leu or L); lysine (Lys or K); methionine (Met or M); phenylalanine (Phe or F); proline (Pro or P); serine (Ser or S); threonine (Thr or T); tryptophan (Trp or W); tyrosine (Tyr or Y); and valine (Val or V), although modified, synthetic, or rare amino acids may be used as desired. Generally, amino acids can be grouped as having a nonpolar side chain (e.g., Ala, Cys, Ile, Leu, Met, Phe, Pro, Val); a negatively charged side chain (e.g., Asp, Glu); a positively charged side chain (e.g., Arg, His, Lys); or an uncharged polar side chain (e.g., Asn, Cys, Gln, Gly, His, Met, Phe, Ser, Thr, Trp, and Tyr).

[131] Amino acid modifications include, for example, deletions from, and/or insertions into, and/or substitutions of, residues within the amino acid sequences of the antibody constructs. Any combination of deletion, insertion, and substitution is made to arrive at the final construct, provided that the final construct possesses the desired characteristics. The amino acid changes also may alter post-translational processes of the antibody constructs, such as changing the number or position of glycosylation sites.

[132] For example, 1, 2, 3, 4, 5, or 6 amino acids may be inserted, substituted or deleted in each of the CDRs (of course, dependent on their length), while 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or 25 amino acids may be inserted, substituted or deleted in each of the FRs. Preferably, amino acid sequence insertions into the antibody construct include amino- and/or carboxyl-terminal fusions ranging in length from 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 residues to polypeptides containing a hundred or more residues, as well as intra-sequence insertions of single or multiple amino acid residues. Corresponding modifications may also be performed within the third domain of the antibody construct of the invention. An insertional variant of the antibody construct of the invention includes the fusion to the N-terminus or to the C-terminus of the antibody construct of an enzyme or the fusion to a polypeptide.

[133] The sites of greatest interest for substitutional mutagenesis include (but are not limited to) the CDRs of the heavy and/or light chain, in particular the hypervariable regions, but FR alterations in the heavy and/or light chain are also contemplated. The substitutions are preferably conservative substitutions as described herein. Preferably, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acids may be substituted in a CDR, while 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12,

13, 14, 15, 16, 17, 18, 19, 20, or 25 amino acids may be substituted in the framework regions (FRs), depending on the length of the CDR or FR. For example, if a CDR sequence encompasses 6 amino acids, it is envisaged that one, two or three of these amino acids are substituted. Similarly, if a CDR sequence encompasses 15 amino acids it is envisaged that one, two, three, four, five or six of these amino acids are substituted.

[134] A useful method for identification of certain residues or regions of the antibody constructs that are preferred locations for mutagenesis is called “alanine scanning mutagenesis” as described by Cunningham and Wells in *Science*, 244: 1081-1085 (1989). Here, a residue or group of target residues within the antibody construct is/are identified (e.g. charged residues such as arg, asp, his, lys, and glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine) to affect the interaction of the amino acids with the epitope.

[135] Those amino acid locations demonstrating functional sensitivity to the substitutions are then refined by introducing further or other variants at, or for, the sites of substitution. Thus, while the site or region for introducing an amino acid sequence variation is predetermined, the nature of the mutation *per se* needs not to be predetermined. For example, to analyze or optimize the performance of a mutation at a given site, alanine scanning or random mutagenesis may be conducted at a target codon or region, and the expressed antibody construct variants are screened for the optimal combination of desired activity. Techniques for making substitution mutations at predetermined sites in the DNA having a known sequence are well known, for example, M13 primer mutagenesis and PCR mutagenesis. Screening of the mutants is done using assays of antigen binding activities, such as PSMA or CD3 binding.

[136] Generally, if amino acids are substituted in one or more or all of the CDRs of the heavy and/or light chain, it is preferred that the then-obtained “substituted” sequence is at least 60% or 65%, more preferably 70% or 75%, even more preferably 80% or 85%, and particularly preferably 90% or 95% identical to the “original” CDR sequence. This means that it is dependent of the length of the CDR to which degree it is identical to the “substituted” sequence. For example, a CDR having 5 amino acids is preferably 80% identical to its substituted sequence in order to have at least one amino acid substituted. Accordingly, the CDRs of the antibody construct may have different degrees of identity to their substituted sequences, e.g., CDRL1 may have 80%, while CDRL3 may have 90%.

[137] Preferred substitutions (or replacements) are conservative substitutions. However, any substitution (including non-conservative substitution or one or more from the “exemplary substitutions” listed in Table 3, below) is envisaged as long as the antibody construct retains

its capability to bind to PSMA via the first domain and to CD3 epsilon via the second domain and/or its CDRs have an identity to the then substituted sequence (at least 60% or 65%, more preferably 70% or 75%, even more preferably 80% or 85%, and particularly preferably 90% or 95% identical to the "original" CDR sequence).

- 5 **[138]** Conservative substitutions are shown in Table 3 under the heading of "preferred substitutions". If such substitutions result in a change in biological activity, then more substantial changes, denominated "exemplary substitutions" in Table 3, or as further described below in reference to amino acid classes, may be introduced and the products screened for a desired characteristic.

10 Table 3: Amino acid substitutions

Original	Exemplary Substitutions	Preferred Substitutions
Ala (A)	val, leu, ile	val
Arg (R)	lys, gln, asn	lys
Asn (N)	gln, his, asp, lys, arg	gln
Asp (D)	glu, asn	glu
Cys (C)	ser, ala	ser
Gln (Q)	asn, glu	asn
Glu (E)	asp, gln	asp
Gly (G)	Ala	ala
His (H)	asn, gln, lys, arg	arg
Ile (I)	leu, val, met, ala, phe	leu
Leu (L)	norleucine, ile, val, met, ala	ile
Lys (K)	arg, gln, asn	arg
Met (M)	leu, phe, ile	leu
Phe (F)	leu, val, ile, ala, tyr	tyr
Pro (P)	Ala	ala
Ser (S)	Thr	thr
Thr (T)	Ser	ser
Trp (W)	tyr, phe	tyr
Tyr (Y)	trp, phe, thr, ser	phe
Val (V)	ile, leu, met, phe, ala	leu

[139] Substantial modifications in the biological properties of the antibody construct of the present invention are accomplished by selecting substitutions that differ significantly in their

effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties: (1) hydrophobic: norleucine, met, ala, val, leu, ile; (2) neutral hydrophilic: cys, ser, thr; asn, gln (3) acidic: asp, glu; (4) basic: his, lys, arg; (5) residues that influence chain orientation: gly, pro; and (6) aromatic : trp, tyr, phe.

[140] Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Any cysteine residue not involved in maintaining the proper conformation of the antibody construct may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking. Conversely, cysteine bond(s) may be added to the antibody to improve its stability (particularly where the antibody is an antibody fragment such as an Fv fragment).

[141] For amino acid sequences, sequence identity and/or similarity is determined by using standard techniques known in the art, including, but not limited to, the local sequence identity algorithm of Smith and Waterman, 1981, *Adv. Appl. Math.* 2:482, the sequence identity alignment algorithm of Needleman and Wunsch, 1970, *J. Mol. Biol.* 48:443, the search for similarity method of Pearson and Lipman, 1988, *Proc. Nat. Acad. Sci. U.S.A.* 85:2444, computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Drive, Madison, Wis.), the Best Fit sequence program described by Devereux *et al.*, 1984, *Nucl. Acid Res.* 12:387-395, preferably using the default settings, or by inspection. Preferably, percent identity is calculated by FastDB based upon the following parameters: mismatch penalty of 1; gap penalty of 1; gap size penalty of 0.33; and joining penalty of 30, "Current Methods in Sequence Comparison and Analysis," Macromolecule Sequencing and Synthesis, Selected Methods and Applications, pp 127-149 (1988), Alan R. Liss, Inc.

[142] An example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments. It can also plot a tree showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle, 1987, *J. Mol. Evol.* 35:351-360; the method is similar to that described by Higgins and Sharp, 1989, *CABIOS* 5:151-153. Useful PILEUP parameters including a default gap weight of 3.00, a default gap length weight of 0.10, and weighted end gaps.

[143] Another example of a useful algorithm is the BLAST algorithm, described in: Altschul *et al.*, 1990, *J. Mol. Biol.* 215:403-410; Altschul *et al.*, 1997, *Nucleic Acids Res.* 25:3389-

3402; and Karin *et al.*, 1993, *Proc. Natl. Acad. Sci. U.S.A.* 90:5873-5787. A particularly useful BLAST program is the WU-BLAST-2 program which was obtained from Altschul *et al.*, 1996, *Methods in Enzymology* 266:460-480. WU-BLAST-2 uses several search parameters, most of which are set to the default values. The adjustable parameters are set with the following values: overlap span=1, overlap fraction=0.125, word threshold (T)=11. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched; however, the values may be adjusted to increase sensitivity.

[144] An additional useful algorithm is gapped BLAST as reported by Altschul *et al.*, 1993, *Nucl. Acids Res.* 25:3389-3402. Gapped BLAST uses BLOSUM-62 substitution scores; threshold T parameter set to 9; the two-hit method to trigger ungapped extensions, charges gap lengths of k a cost of 10+k; Xu set to 16, and Xg set to 40 for database search stage and to 67 for the output stage of the algorithms. Gapped alignments are triggered by a score corresponding to about 22 bits.

[145] Generally, the amino acid homology, similarity, or identity between individual variant CDRs or VH / VL sequences are at least 60% to the sequences depicted herein, and more typically with preferably increasing homologies or identities of at least 65% or 70%, more preferably at least 75% or 80%, even more preferably at least 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, and almost 100%. In a similar manner, "percent (%) nucleic acid sequence identity" with respect to the nucleic acid sequence of the binding proteins identified herein is defined as the percentage of nucleotide residues in a candidate sequence that are identical with the nucleotide residues in the coding sequence of the antibody construct. A specific method utilizes the BLASTN module of WU-BLAST-2 set to the default parameters, with overlap span and overlap fraction set to 1 and 0.125, respectively.

[146] Generally, the nucleic acid sequence homology, similarity, or identity between the nucleotide sequences encoding individual variant CDRs or VH / VL sequences and the nucleotide sequences depicted herein are at least 60%, and more typically with preferably increasing homologies or identities of at least 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%, and almost 100%. Thus, a "variant CDR" or a "variant VH / VL region" is one with the specified homology, similarity, or identity to the parent CDR / VH / VL of the invention, and shares biological function, including, but not limited to, at least 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% of the specificity and/or activity of the parent CDR or VH / VL.

[147] In one embodiment, the percentage of identity to human germline of the antibody constructs according to the invention is $\geq 70\%$ or $\geq 75\%$, more preferably $\geq 80\%$ or $\geq 85\%$, even more preferably $\geq 90\%$, and most preferably $\geq 91\%$, $\geq 92\%$, $\geq 93\%$, $\geq 94\%$, $\geq 95\%$ or even $\geq 96\%$. Identity to human antibody germline gene products is thought to be an important feature to reduce the risk of therapeutic proteins to elicit an immune response against the drug in the patient during treatment. Hwang & Foote ("Immunogenicity of engineered antibodies"; Methods 36 (2005) 3-10) demonstrate that the reduction of non-human portions of drug antibody constructs leads to a decrease of risk to induce anti-drug antibodies in the patients during treatment. By comparing an exhaustive number of clinically evaluated antibody drugs and the respective immunogenicity data, the trend is shown that humanization of the V-regions of antibodies makes the protein less immunogenic (average 5.1 % of patients) than antibodies carrying unaltered non-human V regions (average 23.59 % of patients). A higher degree of identity to human sequences is hence desirable for V-region based protein therapeutics in the form of antibody constructs. For this purpose of determining the germline identity, the V-regions of VL can be aligned with the amino acid sequences of human germline V segments and J segments (<http://vbase.mrc-cpe.cam.ac.uk/>) using Vector NTI software and the amino acid sequence calculated by dividing the identical amino acid residues by the total number of amino acid residues of the VL in percent. The same can be for the VH segments (<http://vbase.mrc-cpe.cam.ac.uk/>) with the exception that the VH CDR3 may be excluded due to its high diversity and a lack of existing human germline VH CDR3 alignment partners. Recombinant techniques can then be used to increase sequence identity to human antibody germline genes.

[148] In a further embodiment, the bispecific antibody constructs of the present invention exhibit high monomer yields under standard research scale conditions, *e.g.*, in a standard two-step purification process. Preferably the monomer yield of the antibody constructs according to the invention is ≥ 0.25 mg/L supernatant, more preferably ≥ 0.5 mg/L, even more preferably ≥ 1 mg/L, and most preferably ≥ 3 mg/L supernatant.

[149] Likewise, the yield of the dimeric antibody construct isoforms and hence the monomer percentage (*i.e.*, monomer : (monomer+dimer)) of the antibody constructs can be determined. The productivity of monomeric and dimeric antibody constructs and the calculated monomer percentage can *e.g.* be obtained in the SEC purification step of culture supernatant from standardized research-scale production in roller bottles. In one embodiment, the monomer percentage of the antibody constructs is $\geq 80\%$, more preferably $\geq 85\%$, even more preferably $\geq 90\%$, and most preferably $\geq 95\%$.

[150] In one embodiment, the antibody constructs have a preferred plasma stability (ratio of EC50 with plasma to EC50 w/o plasma) of ≤ 5 or ≤ 4 , more preferably ≤ 3.5 or ≤ 3 , even

more preferably ≤ 2.5 or ≤ 2 , and most preferably ≤ 1.5 or ≤ 1 . The plasma stability of an antibody construct can be tested by incubation of the construct in human plasma at 37°C for 24 hours followed by EC50 determination in a ^{51} chromium release cytotoxicity assay. The effector cells in the cytotoxicity assay can be stimulated enriched human CD8 positive
5 T cells. Target cells can *e.g.* be CHO cells transfected with human PSMA. The effector to target cell (E:T) ratio can be chosen as 10:1. The human plasma pool used for this purpose is derived from the blood of healthy donors collected by EDTA coated syringes. Cellular components are removed by centrifugation and the upper plasma phase is collected and subsequently pooled. As control, antibody constructs are diluted immediately prior to the
10 cytotoxicity assay in RPMI-1640 medium. The plasma stability is calculated as ratio of EC50 (after plasma incubation) to EC50 (control).

[151] It is furthermore preferred that the monomer to dimer conversion of antibody constructs of the invention is low. The conversion can be measured under different conditions and analyzed by high performance size exclusion chromatography. For example,
15 incubation of the monomeric isoforms of the antibody constructs can be carried out for 7 days at 37°C and concentrations of *e.g.* 100 µg/ml or 250 µg/ml in an incubator. Under these conditions, it is preferred that the antibody constructs of the invention show a dimer percentage that is $\leq 5\%$, more preferably $\leq 4\%$, even more preferably $\leq 3\%$, even more preferably $\leq 2.5\%$, even more preferably $\leq 2\%$, even more preferably $\leq 1.5\%$, and most
20 preferably $\leq 1\%$ or $\leq 0.5\%$ or even 0%.

[152] It is also preferred that the bispecific antibody constructs of the present invention present with very low dimer conversion after a number of freeze/thaw cycles. For example, the antibody construct monomer is adjusted to a concentration of 250 µg/ml *e.g.* in generic formulation buffer and subjected to three freeze/thaw cycles (freezing at -80°C for 30 min
25 followed by thawing for 30 min at room temperature), followed by high performance SEC to determine the percentage of initially monomeric antibody construct, which had been converted into dimeric antibody construct. Preferably the dimer percentages of the bispecific antibody constructs are $\leq 5\%$, more preferably $\leq 4\%$, even more preferably $\leq 3\%$, even more preferably $\leq 2.5\%$, even more preferably $\leq 2\%$, even more preferably $\leq 1.5\%$, and most
30 preferably $\leq 1\%$ or even $\leq 0.5\%$, for example after three freeze/thaw cycles.

[153] The bispecific antibody constructs of the present invention preferably show a favorable thermostability with aggregation temperatures $\geq 45^\circ\text{C}$ or $\geq 50^\circ\text{C}$, more preferably $\geq 52^\circ\text{C}$ or $\geq 54^\circ\text{C}$, even more preferably $\geq 56^\circ\text{C}$ or $\geq 57^\circ\text{C}$, and most preferably $\geq 58^\circ\text{C}$ or $\geq 59^\circ\text{C}$. The thermostability parameter can be determined in terms of antibody aggregation
35 temperature as follows: Antibody solution at a concentration 250 µg/ml is transferred into a single use cuvette and placed in a Dynamic Light Scattering (DLS) device. The sample is

heated from 40°C to 70°C at a heating rate of 0.5°C/min with constant acquisition of the measured radius. Increase of radius indicating melting of the protein and aggregation is used to calculate the aggregation temperature of the antibody.

5 **[154]** Alternatively, temperature melting curves can be determined by Differential Scanning Calorimetry (DSC) to determine intrinsic biophysical protein stabilities of the antibody constructs. These experiments are performed using a MicroCal LLC (Northampton, MA, U.S.A) VP-DSC device. The energy uptake of a sample containing an antibody construct is recorded from 20°C to 90°C compared to a sample containing only the formulation buffer. The antibody constructs are adjusted to a final concentration of 250 µg/ml *e.g.* in SEC
10 running buffer. For recording of the respective melting curve, the overall sample temperature is increased stepwise. At each temperature *T* energy uptake of the sample and the formulation buffer reference is recorded. The difference in energy uptake C_p (kcal/mole/°C) of the sample minus the reference is plotted against the respective temperature. The melting temperature is defined as the temperature at the first maximum of energy uptake.

15 **[155]** The PSMAxCD3 bispecific antibody constructs of the invention are also envisaged to have a turbidity (as measured by OD340 after concentration of purified monomeric antibody construct to 2.5 mg/ml and over night incubation) of ≤ 0.2 , preferably of ≤ 0.15 , more preferably of ≤ 0.12 , even more preferably of ≤ 0.1 , and most preferably of ≤ 0.08 .

20 **[156]** In a further embodiment the antibody construct according to the invention is stable at physiologic or slightly lower pH, *i.e.* about pH 7.4 to 6.0. The more tolerant the antibody construct behaves at unphysiologic pH such as about pH 6.0, the higher is the recovery of the antibody construct eluted from an ion exchange column relative to the total amount of loaded protein. Recovery of the antibody construct from an ion (*e.g.*, cation) exchange column at about pH 6.0 is preferably $\geq 30\%$, more preferably $\geq 40\%$, more preferably $\geq 50\%$,
25 even more preferably $\geq 60\%$, even more preferably $\geq 70\%$, even more preferably $\geq 80\%$, even more preferably $\geq 90\%$, even more preferably $\geq 95\%$, and most preferably $\geq 99\%$.

[157] It is furthermore envisaged that the bispecific antibody constructs of the present invention exhibit therapeutic efficacy or anti-tumor activity. This can *e.g.* be assessed in a study as disclosed in the following example of an advanced stage human tumor xenograft
30 model:

[158] On day 1 of the study, 5×10^6 cells of a human target cell antigen (here: PSMA) positive cancer cell line are subcutaneously injected in the right dorsal flank of female NOD/SCID mice. When the mean tumor volume reaches about 100 mm³, *in vitro* expanded human CD3 positive T cells are transplanted into the mice by injection of about 2×10^7 cells

into the peritoneal cavity of the animals. Mice of vehicle control group 1 do not receive effector cells and are used as an untransplanted control for comparison with vehicle control group 2 (receiving effector cells) to monitor the impact of T cells alone on tumor growth. The antibody treatment starts when the mean tumor volume reaches about 200 mm³. The mean
 5 tumor size of each treatment group on the day of treatment start should not be statistically different from any other group (analysis of variance). Mice are treated with 0.5 mg/kg/day of a PSMAxCD3 bispecific antibody construct by intravenous bolus injection for about 15 to 20 days. Tumors are measured by caliper during the study and progress evaluated by intergroup comparison of tumor volumes (TV). The tumor growth inhibition T/C [%] is
 10 determined by calculating TV as $T/C\% = 100 \times (\text{median TV of analyzed group}) / (\text{median TV of control group 2})$.

[159] The skilled person knows how to modify or adapt certain parameters of this study, such as the number of injected tumor cells, the site of injection, the number of transplanted human T cells, the amount of bispecific antibody constructs to be administered, and the
 15 timelines, while still arriving at a meaningful and reproducible result. Preferably, the tumor growth inhibition T/C [%] is ≤ 70 or ≤ 60 , more preferably ≤ 50 or ≤ 40 , even more preferably ≤ 30 or ≤ 20 and most preferably ≤ 10 or ≤ 5 or even ≤ 2.5 .

[160] In a preferred embodiment of the antibody construct of the invention the antibody construct is a single chain antibody construct.

20 **[161]** Also in a preferred embodiment of the antibody construct of the invention said third domain comprises in an amino to carboxyl order:

hinge-CH2-CH3-linker-hinge-CH2-CH3.

[162] In one embodiment of the invention each of said polypeptide monomers of the third domain has an amino acid sequence that is at least 90% identical to a sequence selected
 25 from the group consisting of: SEQ ID NO: 17-24. In a preferred embodiment of the invention each of said polypeptide monomers has an amino acid sequence selected from SEQ ID NO: 17-24.

[163] Also in one embodiment of the invention the CH2 domain of one or preferably each (both) polypeptide monomers of the third domain comprises an intra domain cysteine
 30 disulfide bridge. As known in the art the term "cysteine disulfide bridge" refers to a functional group with the general structure $R-S-S-R$. The linkage is also called an SS-bond or a disulfide bridge and is derived by the coupling of two thiol groups of cysteine residues. It is particularly preferred for the antibody construct of the invention that the cysteines forming the

cysteine disulfide bridge in the mature antibody construct are introduced into the amino acid sequence of the CH2 domain corresponding to 309 and 321 (Kabat numbering).

[164] In one embodiment of the invention a glycosylation site in Kabat position 314 of the CH2 domain is removed. It is preferred that this removal of the glycosylation site is achieved by a N314X substitution, wherein X is any amino acid excluding Q. Said substitution is preferably a N314G substitution. In a more preferred embodiment, said CH2 domain additionally comprises the following substitutions (position according to Kabat) V321C and R309C (these substitutions introduce the intra domain cysteine disulfide bridge at Kabat positions 309 and 321).

[165] It is assumed that the preferred features of the antibody construct of the invention compared e.g. to the bispecific heteroFc antibody construct known in the art (figure 1b) may be inter alia related to the introduction of the above described modifications in the CH2 domain. Thus, it is preferred for the construct of the invention that the CH2 domains in the third domain of the antibody construct of the invention comprise the intra domain cysteine disulfide bridge at Kabat positions 309 and 321 and/or the glycosylation site at Kabat position 314 is removed by a N314G N314X substitution as above, preferably by a N314G substitution.

[166] In a further preferred embodiment of the invention the CH2 domains in the third domain of the antibody construct of the invention comprise the intra domain cysteine disulfide bridge at Kabat positions 309 and 321 and the glycosylation site at Kabat position 314 is removed by a N314G substitution. Most preferably, the polypeptide monomer of the third domain of the antibody construct of the invention has an amino acid sequence selected from the group consisting of SEQ ID NO: 17 and 18.

[167] In one embodiment the invention provides an antibody construct, wherein:

- (i) the first domain comprises two antibody variable domains and the second domain comprises two antibody variable domains;
- (ii) the first domain comprises one antibody variable domain and the second domain comprises two antibody variable domains;
- (iii) the first domain comprises two antibody variable domains and the second domain comprises one antibody variable domain; or
- (iv) the first domain comprises one antibody variable domain and the second domain comprises one antibody variable domain.

[168] Accordingly, the first and the second domain may be binding domains comprising each two antibody variable domains such as a VH and a VL domain. Examples for such

binding domains comprising two antibody variable domains where described herein above and comprise e.g. Fv fragments, scFv fragments or Fab fragments described herein above. Alternatively either one or both of those binding domains may comprise only a single variable domain. Examples for such single domain binding domains where described herein above and comprise e.g. nanobodies or single variable domain antibodies comprising merely one variable domain, which may be VHH, VH or VL, that specifically bind an antigen or epitope independently of other V regions or domains.

[169] In a preferred embodiment of the antibody construct of the invention first and second domain are fused to the third domain via a peptide linker. Preferred peptide linker have been described herein above and are characterized by the amino acid sequence Gly-Gly-Gly-Gly-Ser, i.e. Gly₄Ser (SEQ ID NO: 1), or polymers thereof, i.e. (Gly₄Ser)_x, where x is an integer of 1 or greater (e.g. 2 or 3). A particularly preferred linker for the fusion of the first and second domain to the third domain is depicted in SEQ ID NOs: 1.

[170] In a preferred embodiment the antibody construct of the invention is characterized to comprise in an amino to carboxyl order:

- (a) the first domain;
- (b) a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: **1-3**;
- (c) the second domain;
- (d) a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NO: **1, 2, 3, 9, 10, 11 and 12**;
- (e) the first polypeptide monomer of the third domain;
- (f) a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: **5, 6, 7 and 8**; and
- (g) the second polypeptide monomer of the third domain.

[171] The antibody construct of the present invention comprises a first domain which binds to PSMA, preferably to the extracellular domain (ECD) of PSMA. It is understood that the term “binding to the extracellular domain of PSMA”, in the context of the present invention, implies that the binding domain binds to PSMA expressed on the surface of a target cell. The first domain according to the invention hence preferably binds to PSMA when it is expressed by naturally expressing cells or cell lines, and/or by cells or cell lines transformed or (stably / transiently) transfected with PSMA. In a preferred embodiment the first binding domain also binds to PSMA when PSMA is used as a “target” or “ligand” molecule in an *in vitro* binding assay such as BIAcore or Scatchard. The “target cell” can be any prokaryotic or eukaryotic cell expressing PSMA on its surface; preferably the target cell is a cell that is part of the human or animal body, such as a specific PSMA expressing cancer or tumor cell.

[172] Preferably, the first binding domain binds to human PSMA / PSMA ECD. In a further preferred embodiment, it binds to macaque PSMA / PSMA ECD. According to the most preferred embodiment, it binds to both the human and the macaque PSMA / PSMA ECD. The "PSMA extracellular domain" or "PSMA ECD" refers to the PSMA region or sequence which is essentially free of transmembrane and cytoplasmic domains of PSMA. It will be understood by the skilled artisan that the transmembrane domain identified for the PSMA polypeptide of the present invention is identified pursuant to criteria routinely employed in the art for identifying that type of hydrophobic domain. The exact boundaries of a transmembrane domain may vary but most likely by no more than about 5 amino acids at either end of the domain specifically mentioned herein.

[173] Preferred binding domains which bind to PSMA are disclosed in WO 2010/037836, and WO 2011/121110. Any binding domain for PSMA described in these applications may be used in the context of the present invention.

[174] In one aspect of the invention the antibody construct comprises in an amino to carboxyl order:

- (a) the first domain having an amino acid sequence selected from the group consisting of SEQ ID NOs: 50, 56, 68, 74, 86, 92, 104, 110, 122, 128, 140, 146, 158, 164, 176, 182, 194, 200, 212, 218, 230, 236, 248, 254, 266, 272, 284, 290, 302, 308, 320, 335, 350, 365, 380, 395, 410, 425, 440, 455, 470;
- (b) a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-3;
- (c) the second domain having an amino acid sequence selected from the group consisting of SEQ ID NOs: 23, 25, 41, 43, 59, 61, 77, 79, 95, 97, 113, 115, 131, 133, 149, 151, 167, 169, 185 or 187 of WO 2008/119567 or as depicted in SEQ ID NO: 15;
- (d) a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1, 2, 3, 9, 10, 11 and 12;
- (e) the first polypeptide monomer of the third domain having a polypeptide sequence selected from the group consisting of SEQ ID NOs: 17-24;
- (f) a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 5, 6, 7 and 8; and
- (g) the second polypeptide monomer of the third domain having a polypeptide sequence selected from the group consisting of SEQ ID NOs: 17-24.

[175] In line with this preferred embodiment, the first and second domain which are fused via a peptide linker to a single chain polypeptide comprise a sequence selected from the group consisting of: SEQ ID NO: 51, 57, 69, 75, 87, 93, 105, 111, 123, 129, 141, 147, 159, 165, 177, 183, 195, 201, 213, 219, 231, 237, 249, 255, 267, 273, 285, 291, 303, 309, 321,

324, 336, 339, 351, 354, 366, 369, 381, 384, 396, 399, 411, 414, 426, 429, 441, 444, 456, 459, 471 and 474.

[176] In one aspect the antibody construct of the invention is characterized by having an amino acid sequence selected from the group consisting of: SEQ ID NOs: 52, 53, 58, 59, 70, 71, 76, 77, 88, 89, 94, 95, 106, 107, 112, 113, 124, 125, 130, 131, 142, 143, 148, 149, 160, 161, 166, 167, 178, 179, 184, 185, 196, 197, 202, 203, 214, 215, 220, 221, 232, 233, 238, 239, 250, 251, 256, 257, 268, 269, 274, 275, 286, 287, 292, 293, 304, 305, 310, 311, 322, 323, 325, 326, 337, 338, 340, 341, 352, 353, 355, 356, 367, 368, 370, 371, 382, 383, 385, 386, 397, 398, 400, 401, 412, 413, 415, 416, 427, 428, 430, 431, 442, 443, 445, 446, 457, 458, 460, 461, 472, 473, 475 and 476.

[177] The invention further provides a polynucleotide / nucleic acid molecule encoding an antibody construct of the invention. A polynucleotide is a biopolymer composed of 13 or more nucleotide monomers covalently bonded in a chain. DNA (such as cDNA) and RNA (such as mRNA) are examples of polynucleotides with distinct biological function. Nucleotides are organic molecules that serve as the monomers or subunits of nucleic acid molecules like DNA or RNA. The nucleic acid molecule or polynucleotide can be double stranded and single stranded, linear and circular. It is preferably comprised in a vector which is preferably comprised in a host cell. Said host cell is, *e.g.* after transformation or transfection with the vector or the polynucleotide of the invention, capable of expressing the antibody construct. For that purpose the polynucleotide or nucleic acid molecule is operatively linked with control sequences.

[178] The genetic code is the set of rules by which information encoded within genetic material (nucleic acids) is translated into proteins. Biological decoding in living cells is accomplished by the ribosome which links amino acids in an order specified by mRNA, using tRNA molecules to carry amino acids and to read the mRNA three nucleotides at a time. The code defines how sequences of these nucleotide triplets, called codons, specify which amino acid will be added next during protein synthesis. With some exceptions, a three-nucleotide codon in a nucleic acid sequence specifies a single amino acid. Because the vast majority of genes are encoded with exactly the same code, this particular code is often referred to as the canonical or standard genetic code. While the genetic code determines the protein sequence for a given coding region, other genomic regions can influence when and where these proteins are produced.

[179] Furthermore, the invention provides a vector comprising a polynucleotide / nucleic acid molecule of the invention. A vector is a nucleic acid molecule used as a vehicle to transfer (foreign) genetic material into a cell. The term “vector” encompasses – but is not

restricted to – plasmids, viruses, cosmids and artificial chromosomes. In general, engineered vectors comprise an origin of replication, a multicloning site and a selectable marker. The vector itself is generally a nucleotide sequence, commonly a DNA sequence that comprises an insert (transgene) and a larger sequence that serves as the “backbone” of the vector.

5 Modern vectors may encompass additional features besides the transgene insert and a backbone: promoter, genetic marker, antibiotic resistance, reporter gene, targeting sequence, protein purification tag. Vectors called expression vectors (expression constructs) specifically are for the expression of the transgene in the target cell, and generally have control sequences.

10 **[180]** The term “control sequences” refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

15 **[181]** A nucleic acid is “operably linked” when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is
20 operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, “operably linked” means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in
25 accordance with conventional practice.

[182] “Transfection” is the process of deliberately introducing nucleic acid molecules or polynucleotides (including vectors) into target cells. The term is mostly used for non-viral methods in eukaryotic cells. Transduction is often used to describe virus-mediated transfer of nucleic acid molecules or polynucleotides. Transfection of animal cells typically involves
30 opening transient pores or “holes” in the cell membrane, to allow the uptake of material. Transfection can be carried out using calcium phosphate, by electroporation, by cell squeezing or by mixing a cationic lipid with the material to produce liposomes, which fuse with the cell membrane and deposit their cargo inside.

[183] The term “transformation” is used to describe non-viral transfer of nucleic acid
35 molecules or polynucleotides (including vectors) into bacteria, and also into non-animal

eukaryotic cells, including plant cells. Transformation is hence the genetic alteration of a bacterial or non-animal eukaryotic cell resulting from the direct uptake through the cell membrane(s) from its surroundings and subsequent incorporation of exogenous genetic material (nucleic acid molecules). Transformation can be effected by artificial means. For transformation to happen, cells or bacteria must be in a state of competence, which may occur as a time-limited response to environmental conditions such as starvation and cell density.

[184] Moreover, the invention provides a host cell transformed or transfected with the polynucleotide / nucleic acid molecule or with the vector of the invention. As used herein, the terms "host cell" or "recipient cell" are intended to include any individual cell or cell culture that can be or has/have been recipients of vectors, exogenous nucleic acid molecules, and polynucleotides encoding the antibody construct of the present invention; and/or recipients of the antibody construct itself. The introduction of the respective material into the cell is carried out by way of transformation, transfection and the like. The term "host cell" is also intended to include progeny or potential progeny of a single cell. Because certain modifications may occur in succeeding generations due to either natural, accidental, or deliberate mutation or due to environmental influences, such progeny may not, in fact, be completely identical (in morphology or in genomic or total DNA complement) to the parent cell, but is still included within the scope of the term as used herein. Suitable host cells include prokaryotic or eukaryotic cells, and also include but are not limited to bacteria, yeast cells, fungi cells, plant cells, and animal cells such as insect cells and mammalian cells, e.g., murine, rat, macaque or human.

[185] The antibody construct of the invention can be produced in bacteria. After expression, the antibody construct of the invention is isolated from the *E. coli* cell paste in a soluble fraction and can be purified through, e.g., affinity chromatography and/or size exclusion. Final purification can be carried out similar to the process for purifying antibody expressed e.g., in CHO cells.

[186] In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for the antibody construct of the invention. *Saccharomyces cerevisiae*, or common baker's yeast, is the most commonly used among lower eukaryotic host microorganisms. However, a number of other genera, species, and strains are commonly available and useful herein, such as *Schizosaccharomyces pombe*, Kluyveromyces hosts such as *K. lactis*, *K. fragilis* (ATCC 12424), *K. bulgaricus* (ATCC 16045), *K. wickerhamii* (ATCC 24178), *K. waltii* (ATCC 56500), *K. drosophilae* (ATCC 36906), *K. thermotolerans*, and *K. marxianus*; *Yarrowia* (EP 402 226); *Pichia pastoris* (EP 183 070); *Candida*; *Trichoderma reesei* (EP 244 234); *Neurospora crassa*; *Schwanniomyces*

such as *Schwanniomyces occidentalis*; and filamentous fungi such as *Neurospora*, *Penicillium*, *Tolypocladium*, and *Aspergillus* hosts such as *A. nidulans* and *A. niger*.

[187] Suitable host cells for the expression of glycosylated antibody construct of the invention are derived from multicellular organisms. Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as *Spodoptera frugiperda* (caterpillar), *Aedes aegypti* (mosquito), *Aedes albopictus* (mosquito), *Drosophila melanogaster* (fruit fly), and *Bombyx mori* have been identified. A variety of viral strains for transfection are publicly available, e.g., the L-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses may be used as the virus herein according to the present invention, particularly for transfection of *Spodoptera frugiperda* cells.

[188] Plant cell cultures of cotton, corn, potato, soybean, petunia, tomato, *Arabidopsis* and tobacco can also be used as hosts. Cloning and expression vectors useful in the production of proteins in plant cell culture are known to those of skill in the art. See e.g. Hiatt *et al.*, Nature (1989) 342: 76-78, Owen *et al.* (1992) Bio/Technology 10: 790-794, Artsaenko *et al.* (1995) The Plant J 8: 745-750, and Fecker *et al.* (1996) Plant Mol Biol 32: 979-986.

[189] However, interest has been greatest in vertebrate cells, and propagation of vertebrate cells in culture (tissue culture) has become a routine procedure. Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham *et al.*, J. Gen Virol. 36 : 59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub *et al.*, Proc. Natl. Acad. Sci. USA 77: 4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod. 23: 243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, 1413 8065); mouse mammary tumor (MMT 060562, ATCC CCL5 1); TRI cells (Mather *et al.*, Annals N. Y Acad. Sci. (1982) 383: 44-68); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

[190] In a further embodiment the invention provides a process for the production of an antibody construct of the invention, said process comprising culturing a host cell of the invention under conditions allowing the expression of the antibody construct of the invention and recovering the produced antibody construct from the culture.

[191] As used herein, the term “culturing” refers to the *in vitro* maintenance, differentiation, growth, proliferation and/or propagation of cells under suitable conditions in a medium. The term “expression” includes any step involved in the production of an antibody construct of the invention including, but not limited to, transcription, post-transcriptional modification, translation, post-translational modification, and secretion.

[192] When using recombinant techniques, the antibody construct can be produced intracellularly, in the periplasmic space, or directly secreted into the medium. If the antibody construct is produced intracellularly, as a first step, the particulate debris, either host cells or lysed fragments, are removed, for example, by centrifugation or ultrafiltration. Carter *et al.*, Bio/Technology 10: 163-167 (1992) describe a procedure for isolating antibodies which are secreted to the periplasmic space of *E. coli*. Briefly, cell paste is thawed in the presence of sodium acetate (pH 3.5), EDTA, and phenylmethylsulfonylfluoride (PMSF) over about 30 min. Cell debris can be removed by centrifugation. Where the antibody is secreted into the medium, supernatants from such expression systems are generally first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. A protease inhibitor such as PMSF may be included in any of the foregoing steps to inhibit proteolysis and antibiotics may be included to prevent the growth of adventitious contaminants.

[193] The antibody construct of the invention prepared from the host cells can be recovered or purified using, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography. Other techniques for protein purification such as fractionation on an ion-exchange column, ethanol precipitation, Reverse Phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSE™, chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation are also available depending on the antibody to be recovered. Where the antibody construct of the invention comprises a CH3 domain, the Bakerbond ABX resin (J.T. Baker, Phillipsburg, NJ) is useful for purification.

[194] Affinity chromatography is a preferred purification technique. The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly(styrenedivinyl) benzene allow for faster flow rates and shorter processing times than can be achieved with agarose.

[195] Moreover, the invention provides a pharmaceutical composition comprising an antibody construct of the invention or an antibody construct produced according to the process of the invention. It is preferred for the pharmaceutical composition of the invention that the homogeneity of the antibody construct is $\geq 80\%$, more preferably $\geq 81\%$, $\geq 82\%$, $\geq$

83%, ≥ 84%, or ≥ 85%, further preferably ≥ 86%, ≥ 87%, ≥ 88%, ≥ 89%, or ≥ 90%, still further preferably, ≥ 91%, ≥ 92%, ≥ 93%, ≥ 94%, or ≥ 95% and most preferably ≥ 96%, ≥ 97%, ≥ 98% or ≥ 99%.

5 **[196]** As used herein, the term “pharmaceutical composition” relates to a composition which is suitable for administration to a patient, preferably a human patient. The particularly preferred pharmaceutical composition of this invention comprises one or a plurality of the antibody construct(s) of the invention, preferably in a therapeutically effective amount. Preferably, the pharmaceutical composition further comprises suitable formulations of one or more (pharmaceutically effective) carriers, stabilizers, excipients, diluents, solubilizers, 10 surfactants, emulsifiers, preservatives and/or adjuvants. Acceptable constituents of the composition are preferably nontoxic to recipients at the dosages and concentrations employed. Pharmaceutical compositions of the invention include, but are not limited to, liquid, frozen, and lyophilized compositions.

15 **[197]** The inventive compositions may comprise a pharmaceutically acceptable carrier. In general, as used herein, “pharmaceutically acceptable carrier” means any and all aqueous and non-aqueous solutions, sterile solutions, solvents, buffers, e.g. phosphate buffered saline (PBS) solutions, water, suspensions, emulsions, such as oil/water emulsions, various types of wetting agents, liposomes, dispersion media and coatings, which are compatible with pharmaceutical administration, in particular with parenteral administration. The use of 20 such media and agents in pharmaceutical compositions is well known in the art, and the compositions comprising such carriers can be formulated by well-known conventional methods.

25 **[198]** Certain embodiments provide pharmaceutical compositions comprising the antibody construct of the invention and further one or more excipients such as those illustratively described in this section and elsewhere herein. Excipients can be used in the invention in this regard for a wide variety of purposes, such as adjusting physical, chemical, or biological properties of formulations, such as adjustment of viscosity, and or processes of the invention to improve effectiveness and or to stabilize such formulations and processes against degradation and spoilage due to, for instance, stresses that occur during manufacturing, 30 shipping, storage, pre-use preparation, administration, and thereafter.

[199] In certain embodiments, the pharmaceutical composition may contain formulation materials for the purpose of modifying, maintaining or preserving, e.g., the pH, osmolarity, viscosity, clarity, color, isotonicity, odor, sterility, stability, rate of dissolution or release, adsorption or penetration of the composition (see, REMINGTON'S PHARMACEUTICAL

SCIENCES, 18th Edition, (A.R. Genrmo, ed.), 1990, Mack Publishing Company). In such embodiments, suitable formulation materials may include, but are not limited to:

- amino acids such as glycine, alanine, glutamine, asparagine, threonine, proline, 2-phenylalanine, including charged amino acids, preferably lysine, lysine acetate, arginine, glutamate and/or histidine
- antimicrobials such as antibacterial and antifungal agents
- antioxidants such as ascorbic acid, methionine, sodium sulfite or sodium hydrogen-sulfite;
- buffers, buffer systems and buffering agents which are used to maintain the composition at physiological pH or at a slightly lower pH,; examples of buffers are borate, bicarbonate, Tris-HCl, citrates, phosphates or other organic acids, succinate, phosphate, and histidine; for example Tris buffer of about pH 7.0-8.5;
- non-aqueous solvents such as propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate;
- aqueous carriers including water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media;
- biodegradable polymers such as polyesters;
- bulking agents such as mannitol or glycine;
- chelating agents such as ethylenediamine tetraacetic acid (EDTA);
- isotonic and absorption delaying agents;
- complexing agents such as caffeine, polyvinylpyrrolidone, beta-cyclodextrin or hydroxypropyl-beta-cyclodextrin)
- fillers;
- monosaccharides; disaccharides; and other carbohydrates (such as glucose, mannose or dextrans); carbohydrates may be non-reducing sugars, preferably trehalose, sucrose, octasulfate, sorbitol or xylitol;
- (low molecular weight) proteins, polypeptides or proteinaceous carriers such as human or bovine serum albumin, gelatin or immunoglobulins, preferably of human origin;
- coloring and flavouring agents;
- sulfur containing reducing agents, such as glutathione, thiocetic acid, sodium thioglycolate, thioglycerol, [alpha]-monothioglycerol, and sodium thio sulfate
- diluting agents;
- emulsifying agents;
- hydrophilic polymers such as polyvinylpyrrolidone)
- salt-forming counter-ions such as sodium;
- preservatives such as antimicrobials, anti-oxidants, chelating agents, inert gases and the like; examples are: benzalkonium chloride, benzoic acid, salicylic acid, thimerosal,

phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid or hydrogen peroxide);

- metal complexes such as Zn-protein complexes;
- solvents and co-solvents (such as glycerin, propylene glycol or polyethylene glycol);
- 5 • sugars and sugar alcohols, such as trehalose, sucrose, octasulfate, mannitol, sorbitol or xylitol stachyose, mannose, sorbose, xylose, ribose, myoinisitol, galactose, lactitol, ribitol, myoinisitol, galactitol, glycerol, cyclitols (e.g., inositol), polyethylene glycol; and polyhydric sugar alcohols;
- suspending agents;
- 10 • surfactants or wetting agents such as pluronics, PEG, sorbitan esters, polysorbates such as polysorbate 20, polysorbate, triton, tromethamine, lecithin, cholesterol, tyloxapal; surfactants may be detergents, preferably with a molecular weight of >1.2 KD and/or a polyether, preferably with a molecular weight of >3 KD; non-limiting examples for preferred detergents are Tween 20, Tween 40, Tween 60, Tween 80 and Tween 85;
- 15 non-limiting examples for preferred polyethers are PEG 3000, PEG 3350, PEG 4000 and PEG 5000;
- stability enhancing agents such as sucrose or sorbitol;
- tonicity enhancing agents such as alkali metal halides, preferably sodium or potassium chloride, mannitol sorbitol;
- 20 • parenteral delivery vehicles including sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils;
- intravenous delivery vehicles including fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose).

[200] It is evident to those skilled in the art that the different constituents of the pharmaceutical composition (e.g., those listed above) can have different effects, for example, and amino acid can act as a buffer, a stabilizer and/or an antioxidant; mannitol can act as a bulking agent and/or a tonicity enhancing agent; sodium chloride can act as delivery vehicle and/or tonicity enhancing agent; etc.

[201] It is envisaged that the composition of the invention may comprise, in addition to the polypeptide of the invention defined herein, further biologically active agents, depending on the intended use of the composition. Such agents may be drugs acting on the gastro-intestinal system, drugs acting as cytostatica, drugs preventing hyperurikemia, drugs inhibiting immunoreactions (e.g. corticosteroids), drugs modulating the inflammatory response, drugs acting on the circulatory system and/or agents such as cytokines known in the art. It is also envisaged that the antibody construct of the present invention is applied in a co-therapy, i.e., in combination with another anti-cancer medicament.

[202] In certain embodiments, the optimal pharmaceutical composition will be determined by one skilled in the art depending upon, for example, the intended route of administration, delivery format and desired dosage. See, for example, REMINGTON'S PHARMACEUTICAL SCIENCES, *supra*. In certain embodiments, such compositions may influence the physical state, stability, rate of in vivo release and rate of in vivo clearance of the antibody construct of the invention. In certain embodiments, the primary vehicle or carrier in a pharmaceutical composition may be either aqueous or non-aqueous in nature. For example, a suitable vehicle or carrier may be water for injection, physiological saline solution or artificial cerebrospinal fluid, possibly supplemented with other materials common in compositions for parenteral administration. Neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles. In certain embodiments, the antibody construct of the invention compositions may be prepared for storage by mixing the selected composition having the desired degree of purity with optional formulation agents (REMINGTON'S PHARMACEUTICAL SCIENCES, *supra*) in the form of a lyophilized cake or an aqueous solution. Further, in certain embodiments, the antibody construct of the invention may be formulated as a lyophilizate using appropriate excipients such as sucrose.

[203] When parenteral administration is contemplated, the therapeutic compositions for use in this invention may be provided in the form of a pyrogen-free, parenterally acceptable aqueous solution comprising the desired antibody construct of the invention in a pharmaceutically acceptable vehicle. A particularly suitable vehicle for parenteral injection is sterile distilled water in which the antibody construct of the invention is formulated as a sterile, isotonic solution, properly preserved. In certain embodiments, the preparation can involve the formulation of the desired molecule with an agent, such as injectable microspheres, bio-erodible particles, polymeric compounds (such as polylactic acid or polyglycolic acid), beads or liposomes, that may provide controlled or sustained release of the product which can be delivered via depot injection. In certain embodiments, hyaluronic acid may also be used, having the effect of promoting sustained duration in the circulation. In certain embodiments, implantable drug delivery devices may be used to introduce the desired antibody construct.

[204] Additional pharmaceutical compositions will be evident to those skilled in the art, including formulations involving the antibody construct of the invention in sustained- or controlled-delivery / release formulations. Techniques for formulating a variety of other sustained- or controlled-delivery means, such as liposome carriers, bio-erodible microparticles or porous beads and depot injections, are also known to those skilled in the art. See, for example, International Patent Application No. PCT/US93/00829, which describes controlled release of porous polymeric microparticles for delivery of

pharmaceutical compositions. Sustained-release preparations may include semipermeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules. Sustained release matrices may include polyesters, hydrogels, polylactides (as disclosed in U.S. Pat. No. 3,773,919 and European Patent Application Publication No. EP 058481),
 5 copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman et al., 1983, Biopolymers 2:547-556), poly (2-hydroxyethyl-methacrylate) (Langer et al., 1981, J. Biomed. Mater. Res. 15:167-277 and Langer, 1982, Chem. Tech. 12:98-105), ethylene vinyl acetate (Langer et al., 1981, supra) or poly-D(-)-3-hydroxybutyric acid (European Patent Application Publication No. EP 133,988). Sustained release compositions may also include liposomes
 10 that can be prepared by any of several methods known in the art. See, e.g., Eppstein et al., 1985, Proc. Natl. Acad. Sci. U.S.A. 82:3688-3692; European Patent Application Publication Nos. EP 036,676; EP 088,046 and EP 143,949.

[205] The antibody construct may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization (for example,
 15 hydroxymethylcellulose or gelatine-microcapsules and poly (methylmethacrylate) microcapsules, respectively), in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nanoparticles and nanocapsules), or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, 16th edition, Oslo, A., Ed., (1980).

[206] Pharmaceutical compositions used for in vivo administration are typically provided as sterile preparations. Sterilization can be accomplished by filtration through sterile filtration membranes. When the composition is lyophilized, sterilization using this method may be conducted either prior to or following lyophilization and reconstitution. Compositions for parenteral administration can be stored in lyophilized form or in a solution. Parenteral
 25 compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

[207] Another aspect of the invention includes self-buffering antibody construct of the invention formulations, which can be used as pharmaceutical compositions, as described in
 30 international patent application WO 06138181A2 (PCT/US2006/022599). A variety of expositions are available on protein stabilization and formulation materials and methods useful in this regard, such as Arakawa et al., "Solvent interactions in pharmaceutical formulations," Pharm Res. 8(3): 285-91 (1991); Kendrick et al., "Physical stabilization of proteins in aqueous solution" in: RATIONAL DESIGN OF STABLE PROTEIN
 35 FORMULATIONS: THEORY AND PRACTICE, Carpenter and Manning, eds. Pharmaceutical Biotechnology. 13: 61-84 (2002), and Randolph et al., "Surfactant-protein interactions",

Pharm Biotechnol. 13: 159-75 (2002), see particularly the parts pertinent to excipients and processes of the same for self-buffering protein formulations in accordance with the current invention, especially as to protein pharmaceutical products and processes for veterinary and/or human medical uses.

5 **[208]** Salts may be used in accordance with certain embodiments of the invention to, for example, adjust the ionic strength and/or the isotonicity of a formulation and/or to improve the solubility and/or physical stability of a protein or other ingredient of a composition in accordance with the invention. As is well known, ions can stabilize the native state of proteins by binding to charged residues on the protein's surface and by shielding charged
10 and polar groups in the protein and reducing the strength of their electrostatic interactions, attractive, and repulsive interactions. Ions also can stabilize the denatured state of a protein by binding to, in particular, the denatured peptide linkages (--CONH) of the protein. Furthermore, ionic interaction with charged and polar groups in a protein also can reduce intermolecular electrostatic interactions and, thereby, prevent or reduce protein aggregation
15 and insolubility.

[209] Ionic species differ significantly in their effects on proteins. A number of categorical rankings of ions and their effects on proteins have been developed that can be used in formulating pharmaceutical compositions in accordance with the invention. One example is the Hofmeister series, which ranks ionic and polar non-ionic solutes by their effect on the
20 conformational stability of proteins in solution. Stabilizing solutes are referred to as "kosmotropic". Destabilizing solutes are referred to as "chaotropic". Kosmotropes commonly are used at high concentrations (e.g., >1 molar ammonium sulfate) to precipitate proteins from solution ("salting-out"). Chaotropes commonly are used to denature and/or to solubilize proteins ("salting-in"). The relative effectiveness of ions to "salt-in" and "salt-out" defines their
25 position in the Hofmeister series.

[210] Free amino acids can be used in the antibody construct of the invention formulations in accordance with various embodiments of the invention as bulking agents, stabilizers, and antioxidants, as well as other standard uses. Lysine, proline, serine, and alanine can be used for stabilizing proteins in a formulation. Glycine is useful in lyophilization to ensure correct
30 cake structure and properties. Arginine may be useful to inhibit protein aggregation, in both liquid and lyophilized formulations. Methionine is useful as an antioxidant.

[211] Polyols include sugars, e.g., mannitol, sucrose, and sorbitol and polyhydric alcohols such as, for instance, glycerol and propylene glycol, and, for purposes of discussion herein, polyethylene glycol (PEG) and related substances. Polyols are kosmotropic. They are useful
35 stabilizing agents in both liquid and lyophilized formulations to protect proteins from physical

and chemical degradation processes. Polyols also are useful for adjusting the tonicity of formulations. Among polyols useful in select embodiments of the invention is mannitol, commonly used to ensure structural stability of the cake in lyophilized formulations. It ensures structural stability to the cake. It is generally used with a lyoprotectant, e.g., sucrose.

5 Sorbitol and sucrose are among preferred agents for adjusting tonicity and as stabilizers to protect against freeze-thaw stresses during transport or the preparation of bulks during the manufacturing process. Reducing sugars (which contain free aldehyde or ketone groups), such as glucose and lactose, can glycate surface lysine and arginine residues. Therefore, they generally are not among preferred polyols for use in accordance with the invention. In
10 addition, sugars that form such reactive species, such as sucrose, which is hydrolyzed to fructose and glucose under acidic conditions, and consequently engenders glycation, also is not among preferred polyols of the invention in this regard. PEG is useful to stabilize proteins and as a cryoprotectant and can be used in the invention in this regard.

[212] Embodiments of the antibody construct of the invention formulations further comprise
15 surfactants. Protein molecules may be susceptible to adsorption on surfaces and to denaturation and consequent aggregation at air-liquid, solid-liquid, and liquid-liquid interfaces. These effects generally scale inversely with protein concentration. These deleterious interactions generally scale inversely with protein concentration and typically are exacerbated by physical agitation, such as that generated during the shipping and handling
20 of a product. Surfactants routinely are used to prevent, minimize, or reduce surface adsorption. Useful surfactants in the invention in this regard include polysorbate 20, polysorbate 80, other fatty acid esters of sorbitan polyethoxylates, and poloxamer 188. Surfactants also are commonly used to control protein conformational stability. The use of surfactants in this regard is protein-specific since, any given surfactant typically will stabilize
25 some proteins and destabilize others.

[213] Polysorbates are susceptible to oxidative degradation and often, as supplied, contain sufficient quantities of peroxides to cause oxidation of protein residue side-chains, especially methionine. Consequently, polysorbates should be used carefully, and when used, should be employed at their lowest effective concentration. In this regard, polysorbates exemplify the
30 general rule that excipients should be used in their lowest effective concentrations.

[214] Embodiments of the antibody construct of the invention formulations further comprise one or more antioxidants. To some extent deleterious oxidation of proteins can be prevented in pharmaceutical formulations by maintaining proper levels of ambient oxygen and temperature and by avoiding exposure to light. Antioxidant excipients can be used as well to
35 prevent oxidative degradation of proteins. Among useful antioxidants in this regard are reducing agents, oxygen/free-radical scavengers, and chelating agents. Antioxidants for use

in therapeutic protein formulations in accordance with the invention preferably are water-soluble and maintain their activity throughout the shelf life of a product. EDTA is a preferred antioxidant in accordance with the invention in this regard. Antioxidants can damage proteins. For instance, reducing agents, such as glutathione in particular, can disrupt intramolecular disulfide linkages. Thus, antioxidants for use in the invention are selected to, among other things, eliminate or sufficiently reduce the possibility of themselves damaging proteins in the formulation.

[215] Formulations in accordance with the invention may include metal ions that are protein co-factors and that are necessary to form protein coordination complexes, such as zinc necessary to form certain insulin suspensions. Metal ions also can inhibit some processes that degrade proteins. However, metal ions also catalyze physical and chemical processes that degrade proteins. Magnesium ions (10-120 mM) can be used to inhibit isomerization of aspartic acid to isoaspartic acid. Ca^{+2} ions (up to 100 mM) can increase the stability of human deoxyribonuclease. Mg^{+2} , Mn^{+2} , and Zn^{+2} , however, can destabilize rhDNase. Similarly, Ca^{+2} and Sr^{+2} can stabilize Factor VIII, it can be destabilized by Mg^{+2} , Mn^{+2} and Zn^{+2} , Cu^{+2} and Fe^{+2} , and its aggregation can be increased by Al^{+3} ions.

[216] Embodiments of the antibody construct of the invention formulations further comprise one or more preservatives. Preservatives are necessary when developing multi-dose parenteral formulations that involve more than one extraction from the same container. Their primary function is to inhibit microbial growth and ensure product sterility throughout the shelf-life or term of use of the drug product. Commonly used preservatives include benzyl alcohol, phenol and m-cresol. Although preservatives have a long history of use with small-molecule parenterals, the development of protein formulations that includes preservatives can be challenging. Preservatives almost always have a destabilizing effect (aggregation) on proteins, and this has become a major factor in limiting their use in multi-dose protein formulations. To date, most protein drugs have been formulated for single-use only. However, when multi-dose formulations are possible, they have the added advantage of enabling patient convenience, and increased marketability. A good example is that of human growth hormone (hGH) where the development of preserved formulations has led to commercialization of more convenient, multi-use injection pen presentations. At least four such pen devices containing preserved formulations of hGH are currently available on the market. Norditropin (liquid, Novo Nordisk), Nutropin AQ (liquid, Genentech) & Genotropin (lyophilized--dual chamber cartridge, Pharmacia & Upjohn) contain phenol while Somatropo (Eli Lilly) is formulated with m-cresol. Several aspects need to be considered during the formulation and development of preserved dosage forms. The effective preservative concentration in the drug product must be optimized. This requires testing a given

preservative in the dosage form with concentration ranges that confer anti-microbial effectiveness without compromising protein stability.

[217] As may be expected, development of liquid formulations containing preservatives are more challenging than lyophilized formulations. Freeze-dried products can be lyophilized without the preservative and reconstituted with a preservative containing diluent at the time of use. This shortens the time for which a preservative is in contact with the protein, significantly minimizing the associated stability risks. With liquid formulations, preservative effectiveness and stability should be maintained over the entire product shelf-life (about 18 to 24 months). An important point to note is that preservative effectiveness should be demonstrated in the final formulation containing the active drug and all excipient components.

[218] The antibody constructs disclosed herein may also be formulated as immuno-liposomes. A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes. Liposomes containing the antibody construct are prepared by methods known in the art, such as described in Epstein et al., Proc. Natl. Acad. Sci. USA, 82: 3688 (1985); Hwang et al. , Proc. Natl Acad. Sci. USA, 77: 4030 (1980); US Pat. Nos. 4,485,045 and 4,544,545; and WO 97/38731. Liposomes with enhanced circulation time are disclosed in US Patent No. 5,013, 556. Particularly useful liposomes can be generated by the reverse phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody construct of the present invention can be conjugated to the liposomes as described in Martin et al. J. Biol. Chem. 257: 286-288 (1982) via a disulfide interchange reaction. A chemotherapeutic agent is optionally contained within the liposome. See Gabizon et al. J. National Cancer Inst. 81 (19) 1484 (1989).

[219] Once the pharmaceutical composition has been formulated, it may be stored in sterile vials as a solution, suspension, gel, emulsion, solid, crystal, or as a dehydrated or lyophilized powder. Such formulations may be stored either in a ready-to-use form or in a form (e.g., lyophilized) that is reconstituted prior to administration.

[220] The biological activity of the pharmaceutical composition defined herein can be determined for instance by cytotoxicity assays, as described in the following examples, in WO 99/54440 or by Schlereth et al. (Cancer Immunol. Immunother. 20 (2005), 1-12). "Efficacy" or "in vivo efficacy" as used herein refers to the response to therapy by the

pharmaceutical composition of the invention, using e.g. standardized NCI response criteria. The success or in vivo efficacy of the therapy using a pharmaceutical composition of the invention refers to the effectiveness of the composition for its intended purpose, i.e. the ability of the composition to cause its desired effect, i.e. depletion of pathologic cells, e.g. tumor cells. The in vivo efficacy may be monitored by established standard methods for the respective disease entities including, but not limited to white blood cell counts, differentials, Fluorescence Activated Cell Sorting, bone marrow aspiration. In addition, various disease specific clinical chemistry parameters and other established standard methods may be used. Furthermore, computer-aided tomography, X-ray, nuclear magnetic resonance tomography (e.g. for National Cancer Institute-criteria based response assessment [Cheson BD, Horning SJ, Coiffier B, Shipp MA, Fisher RI, Connors JM, Lister TA, Vose J, Grillo-Lopez A, Hagenbeek A, Cabanillas F, Klippensten D, Hiddemann W, Castellino R, Harris NL, Armitage JO, Carter W, Hoppe R, Canellos GP. Report of an international workshop to standardize response criteria for non-Hodgkin's lymphomas. NCI Sponsored International Working Group. J Clin Oncol. 1999 Apr;17(4):1244]), positron-emission tomography scanning, white blood cell counts, differentials, Fluorescence Activated Cell Sorting, bone marrow aspiration, lymph node biopsies/histologies, and various lymphoma specific clinical chemistry parameters (e.g. lactate dehydrogenase) and other established standard methods may be used.

[221] Another major challenge in the development of drugs such as the pharmaceutical composition of the invention is the predictable modulation of pharmacokinetic properties. To this end, a pharmacokinetic profile of the drug candidate, i.e. a profile of the pharmacokinetic parameters that affect the ability of a particular drug to treat a given condition, can be established. Pharmacokinetic parameters of the drug influencing the ability of a drug for treating a certain disease entity include, but are not limited to: half-life, volume of distribution, hepatic first-pass metabolism and the degree of blood serum binding. The efficacy of a given drug agent can be influenced by each of the parameters mentioned above. It is an envisaged characteristic of the antibody constructs of the present invention provided with the specific FC modality that they comprise, for example, differences in pharmacokinetic behavior. A half-life extended targeting antibody construct according to the present invention preferably shows a surprisingly increased residence time in vivo in comparison to "canonical" non-HLE versions of said antibody construct.

[222] "Half-life" means the time where 50% of an administered drug are eliminated through biological processes, e.g. metabolism, excretion, etc. By "hepatic first-pass metabolism" is meant the propensity of a drug to be metabolized upon first contact with the liver, i.e. during its first pass through the liver. "Volume of distribution" means the degree of retention of a

drug throughout the various compartments of the body, like e.g. intracellular and extracellular spaces, tissues and organs, etc. and the distribution of the drug within these compartments. "Degree of blood serum binding" means the propensity of a drug to interact with and bind to blood serum proteins, such as albumin, leading to a reduction or loss of biological activity of the drug.

[223] Pharmacokinetic parameters also include bioavailability, lag time (Tlag), Tmax, absorption rates, more onset and/or Cmax for a given amount of drug administered. "Bioavailability" means the amount of a drug in the blood compartment. "Lag time" means the time delay between the administration of the drug and its detection and measurability in blood or plasma. "Tmax" is the time after which maximal blood concentration of the drug is reached, and "Cmax" is the blood concentration maximally obtained with a given drug. The time to reach a blood or tissue concentration of the drug which is required for its biological effect is influenced by all parameters. Pharmacokinetic parameters of bispecific antibody constructs exhibiting cross-species specificity, which may be determined in preclinical animal testing in non-chimpanzee primates as outlined above, are also set forth e.g. in the publication by Schlereth et al. (Cancer Immunol. Immunother. 20 (2005), 1-12).

[224] In a preferred aspect of the invention the pharmaceutical composition is stable for at least four weeks at about -20°C. As apparent from the appended examples the quality of an antibody construct of the invention vs. the quality of corresponding state of the art antibody constructs may be tested using different systems. Those tests are understood to be in line with the "ICH Harmonised Tripartite Guideline: *Stability Testing of Biotechnological/Biological Products Q5C* and *Specifications: Test procedures and Acceptance Criteria for Biotech Biotechnological/Biological Products Q6B*" and, thus are elected to provide a stability-indicating profile that provides certainty that changes in the identity, purity and potency of the product are detected. It is well accepted that the term purity is a relative term. Due to the effect of glycosylation, deamidation, or other heterogeneities, the absolute purity of a biotechnological/biological product should be typically assessed by more than one method and the purity value derived is method-dependent. For the purpose of stability testing, tests for purity should focus on methods for determination of degradation products.

[225] For the assessment of the quality of a pharmaceutical composition comprising an antibody construct of the invention may be analyzed e.g. by analyzing the content of soluble aggregates in a solution (HMWS per size exclusion). It is preferred that stability for at least four weeks at about -20°C is characterized by a content of less than 1.5% HMWS, preferably by less than 1% HMWS.

[226] A preferred formulation for the antibody construct as a pharmaceutical composition may e.g. comprise the components of a formulation as described below:

- Formulation:
potassium phosphate, L-arginine hydrochloride, trehalose dihydrate, polysorbate 80
at pH 6.0

[227] Other examples for the assessment of the stability of an antibody construct of the invention in form of a pharmaceutical composition are provided in the appended examples 4-12. In those examples embodiments of antibody constructs of the invention are tested with respect to different stress conditions in different pharmaceutical formulations and the results compared with other half-life extending (HLE) formats of bispecific T cell engaging antibody construct known from the art. In general, it is envisaged that antibody constructs provided with the specific FC modality according to the present invention are typically more stable over a broad range of stress conditions such as temperature and light stress, both compared to antibody constructs provided with different HLE formats and without any HLE format (e.g. "canonical" antibody constructs). Said temperature stability may relate both to decreased (below room temperature including freezing) and increased (above room temperature including temperatures up to or above body temperature) temperature. As the person skilled in the art will acknowledge, such improved stability with regard to stress, which is hardly avoidable in clinical practice, makes the antibody construct safer because less degradation products will occur in clinical practice. In consequence, said increased stability means increased safety.

[228] One embodiment provides the antibody construct of the invention or the antibody construct produced according to the process of the invention for use in the prevention, treatment or amelioration of a cancer correlating with PSMA expression or PSMA overexpression, such as prostate cancer.

[229] The formulations described herein are useful as pharmaceutical compositions in the treatment, amelioration and/or prevention of the pathological medical condition as described herein in a patient in need thereof. The term "treatment" refers to both therapeutic treatment and prophylactic or preventative measures. Treatment includes the application or administration of the formulation to the body, an isolated tissue, or cell from a patient who has a disease/disorder, a symptom of a disease/disorder, or a predisposition toward a disease/disorder, with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve, or affect the disease, the symptom of the disease, or the predisposition toward the disease.

[230] The term “amelioration” as used herein refers to any improvement of the disease state of a patient having a disease as specified herein below, by the administration of an antibody construct according to the invention to a subject in need thereof. Such an improvement may also be seen as a slowing or stopping of the progression of the patient's disease. The term “prevention” as used herein means the avoidance of the occurrence or re-occurrence of a patient having a tumor or cancer or a metastatic cancer as specified herein below, by the administration of an antibody construct according to the invention to a subject in need thereof.

[231] The term “disease” refers to any condition that would benefit from treatment with the antibody construct or the pharmaceutical composition described herein. This includes chronic and acute disorders or diseases including those pathological conditions that predispose the mammal to the disease in question.

[232] A “neoplasm” is an abnormal growth of tissue, usually but not always forming a mass. When also forming a mass, it is commonly referred to as a “tumor”. Neoplasms or tumors or can be benign, potentially malignant (pre-cancerous), or malignant. Malignant neoplasms are commonly called cancer. They usually invade and destroy the surrounding tissue and may form metastases, i.e., they spread to other parts, tissues or organs of the body. Hence, the term “metastatic cancer” encompasses metastases to other tissues or organs than the one of the original tumor. Lymphomas and leukemias are lymphoid neoplasms. For the purposes of the present invention, they are also encompassed by the terms “tumor” or “cancer”.

[233] The term “viral disease” describes diseases, which are the result of a viral infection of a subject.

[234] The term “immunological disorder” as used herein describes in line with the common definition of this term immunological disorders such as autoimmune diseases, hypersensitivities, immune deficiencies.

[235] In one embodiment the invention provides a method for the treatment or amelioration of a cancer correlating with PSMA expression or PSMA overexpression, comprising the step of administering to a subject in need thereof the antibody construct of the invention, or the antibody construct produced according to the process of the invention. The PSMAxCD3 bispecific single chain antibody is particularly advantageous for the therapy of cancer, preferably solid tumors, more preferably carcinomas and prostate cancer.

[236] The terms “subject in need” or those “in need of treatment” includes those already with the disorder, as well as those in which the disorder is to be prevented. The subject in

need or "patient" includes human and other mammalian subjects that receive either prophylactic or therapeutic treatment.

[237] The antibody construct of the invention will generally be designed for specific routes and methods of administration, for specific dosages and frequencies of administration, for specific treatments of specific diseases, with ranges of bio-availability and persistence, among other things. The materials of the composition are preferably formulated in concentrations that are acceptable for the site of administration.

[238] Formulations and compositions thus may be designed in accordance with the invention for delivery by any suitable route of administration. In the context of the present invention, the routes of administration include, but are not limited to

- topical routes (such as epicutaneous, inhalational, nasal, ophthalmic, auricular / aural, vaginal, mucosal);
- enteral routes (such as oral, gastrointestinal, sublingual, sublabial, buccal, rectal); and
- parenteral routes (such as intravenous, intraarterial, intraosseous, intramuscular, intracerebral, intracerebroventricular, epidural, intrathecal, subcutaneous, intraperitoneal, extra-amniotic, intraarticular, intracardiac, intradermal, intralesional, intrauterine, intravesical, intravitreal, transdermal, intranasal, transmucosal, intrasynovial, intraluminal).

[239] The pharmaceutical compositions and the antibody construct of this invention are particularly useful for parenteral administration, e.g., subcutaneous or intravenous delivery, for example by injection such as bolus injection, or by infusion such as continuous infusion. Pharmaceutical compositions may be administered using a medical device. Examples of medical devices for administering pharmaceutical compositions are described in U.S. Patent Nos. 4,475,196; 4,439,196; 4,447,224; 4,447, 233; 4,486,194; 4,487,603; 4,596,556; 4,790,824; 4,941,880; 5,064,413; 5,312,335; 5,312,335; 5,383,851; and 5,399,163.

[240] In particular, the present invention provides for an uninterrupted administration of the suitable composition. As a non-limiting example, uninterrupted or substantially uninterrupted, i.e. continuous administration may be realized by a small pump system worn by the patient for metering the influx of therapeutic agent into the body of the patient. The pharmaceutical composition comprising the antibody construct of the invention can be administered by using said pump systems. Such pump systems are generally known in the art, and commonly rely on periodic exchange of cartridges containing the therapeutic agent to be infused. When exchanging the cartridge in such a pump system, a temporary interruption of the otherwise uninterrupted flow of therapeutic agent into the body of the patient may ensue. In such a case, the phase of administration prior to cartridge replacement and the phase of

administration following cartridge replacement would still be considered within the meaning of the pharmaceutical means and methods of the invention together make up one “uninterrupted administration” of such therapeutic agent.

5 [241] The continuous or uninterrupted administration of the antibody constructs of the invention may be intravenous or subcutaneous by way of a fluid delivery device or small pump system including a fluid driving mechanism for driving fluid out of a reservoir and an actuating mechanism for actuating the driving mechanism. Pump systems for subcutaneous administration may include a needle or a cannula for penetrating the skin of a patient and delivering the suitable composition into the patient’s body. Said pump systems may be
10 directly fixed or attached to the skin of the patient independently of a vein, artery or blood vessel, thereby allowing a direct contact between the pump system and the skin of the patient. The pump system can be attached to the skin of the patient for 24 hours up to several days. The pump system may be of small size with a reservoir for small volumes. As a non-limiting example, the volume of the reservoir for the suitable pharmaceutical composition
15 to be administered can be between 0.1 and 50 ml.

[242] The continuous administration may also be transdermal by way of a patch worn on the skin and replaced at intervals. One of skill in the art is aware of patch systems for drug delivery suitable for this purpose. It is of note that transdermal administration is especially amenable to uninterrupted administration, as exchange of a first exhausted patch can
20 advantageously be accomplished simultaneously with the placement of a new, second patch, for example on the surface of the skin immediately adjacent to the first exhausted patch and immediately prior to removal of the first exhausted patch. Issues of flow interruption or power cell failure do not arise.

[243] If the pharmaceutical composition has been lyophilized, the lyophilized material is first
25 reconstituted in an appropriate liquid prior to administration. The lyophilized material may be reconstituted in, e.g., bacteriostatic water for injection (BWFI), physiological saline, phosphate buffered saline (PBS), or the same formulation the protein had been in prior to lyophilization.

[244] The compositions of the present invention can be administered to the subject at a
30 suitable dose which can be determined e.g. by dose escalating studies by administration of increasing doses of the antibody construct of the invention exhibiting cross-species specificity described herein to non-chimpanzee primates, for instance macaques. As set forth above, the antibody construct of the invention exhibiting cross-species specificity described herein can be advantageously used in identical form in preclinical testing in non-chimpanzee
35 primates and as drug in humans. The dosage regimen will be determined by the attending

physician and clinical factors. As is well known in the medical arts, dosages for any one patient depend upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently.

5 **[245]** The term "effective dose" or "effective dosage" is defined as an amount sufficient to achieve or at least partially achieve the desired effect. The term "therapeutically effective dose" is defined as an amount sufficient to cure or at least partially arrest the disease and its complications in a patient already suffering from the disease. Amounts or doses effective for this use will depend on the condition to be treated (the indication), the delivered antibody
10 construct, the therapeutic context and objectives, the severity of the disease, prior therapy, the patient's clinical history and response to the therapeutic agent, the route of administration, the size (body weight, body surface or organ size) and/or condition (the age and general health) of the patient, and the general state of the patient's own immune system. The proper dose can be adjusted according to the judgment of the attending physician such
15 that it can be administered to the patient once or over a series of administrations, and in order to obtain the optimal therapeutic effect.

[246] A typical dosage may range from about 0.1 $\mu\text{g/kg}$ to up to about 30 mg/kg or more, depending on the factors mentioned above. In specific embodiments, the dosage may range from 1.0 $\mu\text{g/kg}$ up to about 20 mg/kg , optionally from 10 $\mu\text{g/kg}$ up to about 10 mg/kg or from
20 100 $\mu\text{g/kg}$ up to about 5 mg/kg .

[247] A therapeutic effective amount of an antibody construct of the invention preferably results in a decrease in severity of disease symptoms, an increase in frequency or duration of disease symptom-free periods or a prevention of impairment or disability due to the disease affliction. For treating diseases correlating with PSMA expression as described
25 herein above, a therapeutically effective amount of the antibody construct of the invention, here: an anti-PSMA/anti-CD3 antibody construct, preferably inhibits cell growth or tumor growth by at least about 20%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, or at least about 90% relative to untreated patients. The ability of a compound to inhibit tumor growth may be evaluated in an animal model predictive
30 of efficacy

[248] The pharmaceutical composition can be administered as a sole therapeutic or in combination with additional therapies such as anti-cancer therapies as needed, e.g. other proteinaceous and non-proteinaceous drugs. These drugs may be administered simultaneously with the composition comprising the antibody construct of the invention as

defined herein or separately before or after administration of said antibody construct in timely defined intervals and doses.

[249] The term “effective and non-toxic dose” as used herein refers to a tolerable dose of an inventive antibody construct which is high enough to cause depletion of pathologic cells, tumor elimination, tumor shrinkage or stabilization of disease without or essentially without major toxic effects. Such effective and non-toxic doses may be determined e.g. by dose escalation studies described in the art and should be below the dose inducing severe adverse side events (dose limiting toxicity, DLT).

[250] The term “toxicity” as used herein refers to the toxic effects of a drug manifested in adverse events or severe adverse events. These side events may refer to a lack of tolerability of the drug in general and/or a lack of local tolerance after administration. Toxicity could also include teratogenic or carcinogenic effects caused by the drug.

[251] The term “safety”, “in vivo safety” or “tolerability” as used herein defines the administration of a drug without inducing severe adverse events directly after administration (local tolerance) and during a longer period of application of the drug. “Safety”, “in vivo safety” or “tolerability” can be evaluated e.g. at regular intervals during the treatment and follow-up period. Measurements include clinical evaluation, e.g. organ manifestations, and screening of laboratory abnormalities. Clinical evaluation may be carried out and deviations to normal findings recorded/coded according to NCI-CTC and/or MedDRA standards. Organ manifestations may include criteria such as allergy/immunology, blood/bone marrow, cardiac arrhythmia, coagulation and the like, as set forth e.g. in the Common Terminology Criteria for adverse events v3.0 (CTCAE). Laboratory parameters which may be tested include for instance hematology, clinical chemistry, coagulation profile and urine analysis and examination of other body fluids such as serum, plasma, lymphoid or spinal fluid, liquor and the like. Safety can thus be assessed e.g. by physical examination, imaging techniques (i.e. ultrasound, x-ray, CT scans, Magnetic Resonance Imaging (MRI), other measures with technical devices (i.e. electrocardiogram), vital signs, by measuring laboratory parameters and recording adverse events. For example, adverse events in non-chimpanzee primates in the uses and methods according to the invention may be examined by histopathological and/or histochemical methods.

[252] The above terms are also referred to e.g. in the Preclinical safety evaluation of biotechnology-derived pharmaceuticals S6; ICH Harmonised Tripartite Guideline; ICH Steering Committee meeting on July 16, 1997.

[253] Finally, the invention provides a kit comprising an antibody construct of the invention or produced according to the process of the invention, a pharmaceutical composition of the invention, a polynucleotide of the invention, a vector of the invention and/or a host cell of the invention.

5 **[254]** In the context of the present invention, the term “kit” means two or more components – one of which corresponding to the antibody construct, the pharmaceutical composition, the vector or the host cell of the invention – packaged together in a container, recipient or otherwise. A kit can hence be described as a set of products and/or utensils that are sufficient to achieve a certain goal, which can be marketed as a single unit.

10 **[255]** The kit may comprise one or more recipients (such as vials, ampoules, containers, syringes, bottles, bags) of any appropriate shape, size and material (preferably waterproof, *e.g.* plastic or glass) containing the antibody construct or the pharmaceutical composition of the present invention in an appropriate dosage for administration (see above). The kit may additionally contain directions for use (*e.g.* in the form of a leaflet or instruction manual),
 15 means for administering the antibody construct of the present invention such as a syringe, pump, infuser or the like, means for reconstituting the antibody construct of the invention and/or means for diluting the antibody construct of the invention.

[256] The invention also provides kits for a single-dose administration unit. The kit of the invention may also contain a first recipient comprising a dried / lyophilized antibody construct
 20 and a second recipient comprising an aqueous formulation. In certain embodiments of this invention, kits containing single-chambered and multi-chambered pre-filled syringes (*e.g.*, liquid syringes and lyosyringes) are provided.

[257] It must be noted that as used herein, the singular forms “a”, “an”, and “the”, include
 25 plural references unless the context clearly indicates otherwise. Thus, for example, reference to “a reagent” includes one or more of such different reagents and reference to “the method” includes reference to equivalent steps and methods known to those of ordinary skill in the art that could be modified or substituted for the methods described herein.

[258] Unless otherwise indicated, the term “at least” preceding a series of elements is to be
 30 understood to refer to every element in the series. Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the present invention.

[259] The term "and/or" wherever used herein includes the meaning of "and", "or" and "all or any other combination of the elements connected by said term".

[260] The term "about" or "approximately" as used herein means within 20%, preferably within 10%, and more preferably within 5% of a given value or range. It includes, however,
5 also the concrete number, e.g., about 20 includes 20.

[261] The term "less than" or "greater than" includes the concrete number. For example, less than 20 means less than or equal to. Similarly, more than or greater than means more than or equal to, or greater than or equal to, respectively.

[262] Throughout this specification and the claims which follow, unless the context requires
10 otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integer or step. When used herein the term "comprising" can be substituted with the term "containing" or "including" or sometimes when used herein with the term "having".

[263] When used herein "consisting of" excludes any element, step, or ingredient not
15 specified in the claim element. When used herein, "consisting essentially of" does not exclude materials or steps that do not materially affect the basic and novel characteristics of the claim.

[264] In each instance herein any of the terms "comprising", "consisting essentially of" and
20 "consisting of" may be replaced with either of the other two terms.

[265] It should be understood that this invention is not limited to the particular methodology, protocols, material, reagents, and substances, etc., described herein and as such can vary. The terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention, which is defined solely by the
25 claims.

[266] All publications and patents cited throughout the text of this specification (including all patents, patent applications, scientific publications, manufacturer's specifications, instructions, etc.), whether supra or infra, are hereby incorporated by reference in their entirety. Nothing herein is to be construed as an admission that the invention is not entitled to
30 antedate such disclosure by virtue of prior invention. To the extent the material incorporated by reference contradicts or is inconsistent with this specification, the specification will supersede any such material.

[267] A better understanding of the present invention and of its advantages will be obtained from the following examples, offered for illustrative purposes only. The examples are not
35 intended to limit the scope of the present invention in any way.

[268] Example 1: BiTE[®] induced CD69 Expression on T Cells in Absence of Target Cells

Isolated PBMC from healthy human donors were cultured with increasing Target B/CD3 or Target A/CD3 HLE bispecific antibody constructs for 48 h. The expression of the activation marker CD69 on T cells was determined by immunostaining and flow cytometry and antigen specific conjugates mAb.

Target-independent T cell activation in terms of CD69 upregulation was observed for all anti-CDH 19 constructs but was most pronounced for heteroFc and crossbody molecules. Upregulation of CD69 by anti-Target B-scFc occurred at higher concentrations and the amplitude was in part lower compared to the other two Fc-based constructs.

For the anti-Target A almost no target-independent T cell activation was observed for the scFc-containing molecule, while the heteroFc construct induced a strong upregulation of CD69 on the cell surface T cells in the absence of target cells.

Target-independent T cell activation induced by BiTE[®] antibody constructs containing a single chain-Fc, or hetero-Fc fusion at the C-terminus was evaluated for the following constructs:

BiTE[®] antibody constructs (serial dilutions: 0.1 pM – 2 μM)

a. Target A-I2C scFc; 1.14 mg/mL;

b. Target A Hetero Fc; 1.02 mg/

Human PBMC effector cells (3 donors; #065, #823, #836 (scFc) #401, #415, #433 (heteroFc); #590, #595, 598, #605 (X-body)).

48 h incubation time.

Determination of CD69 expression on CD4⁺ and CD8⁺ T cells with flow cytometer and antigen-specific conjugates mAb. Results see Figure 2.

Target-independent T cell activation induced by BiTE[®] antibody constructs containing a single chain-Fc, hetero-Fc or crossbody fusion at the C-terminus was evaluated for the following constructs:

BiTE[®] antibody constructs (serial dilutions: 0.1 pM – 2 μM)

c. Target B x I2C-scFc; 245.3 μg/mL (

d. Target B Hetero Fc; 1 mg/mL

e. Target B crossbody; 6.3 mg/mL

Human PBMC effector cells (3 to 4 donors; #386, #392, #401 (scFc) #282, #284, #287 (heteroFc)). 48 h incubation time.

Determination of CD69 expression on CD4+ and CD8+ T cells with flow cytometer and antigen-specific conjugates mAb. Results see Figure 3.

- 5 Target-independent T cell activation in terms of CD69 upregulation was observed for several bispecific antibody constructs tested in these assays. The CD69 upregulation was in general more pronounced for the canonical BiTE[®] antibody constructs, heteroFc and crossbody antibody constructs when compared to the respective scFc antibody constructs. Upregulation of CD69 by the scFc constructs occurred in general at slightly higher concentrations and the
- 10 amplitude was in part lower compared to the other two Fc-based constructs.
- For the anti-Target B scFc antibody construct, no target-independent T cell activation was observed, while the heteroFc and X-Body antibody constructs induced a strong upregulation of CD69 on the cell surface of T cells in the absence of target cells. Hence, the scFc antibody construct according to the present invention shows an advantage as unspecific T cell
- 15 activation, as exemplified here by CD69 upregulation, is not desired in specific immunotherapy.

Materials and methods

20 Target B

Target-independent T cell activation induced by a BiTE[®] antibody constructs containing a single chain-Fc for the following construct:

1. BiTE[®] antibody construct (serial dilutions: 1.3 pM – 20 nM)
 1. Target B-scFc
- 25 2. Human PBMC effector cells (3 donors)
3. 48 h incubation time
4. Flow cytometric analysis of CD69 expression on CD4+ and CD8+ T cells using a PE-Cy7 conjugated mAb specific for CD69.

30 [269] Example 2:

- Purified BiTE[®] antibody constructs were coated on a Maxisorb Plate in decreasing concentration (100nM, 1:4 dilutions). After 3x washing with PBS-T and blocking with PBS/3% (w/v) BSA (60 min, 37°C), pooled human plasma was incubated for 60 min, 80 rpm at room temperature. After 3x washing a mouse monoclonal antibody specific for human C1q subunit
- 35 A (CC1q) was added (Thermo MA1-83963, 1:500) for 60 min, 80 rpm, room temperature,

after described washing steps a goat anti mouse Fc-POX mAb (1:5,000) was incubated for 60 min, 80 rpm, room temperature. After additional washing, TMB substrate was incubated and stopped after colorimetric reaction by addition of H₂SO₄. The absorption was determined at 450 nm.

- 5 Result: As shown in figure 4 at high concentrations, the BiTE® hetero Fc antibody construct (squares) showed higher binding signals for human CC1q compared to a BiTE® single chain Fc antibody construct (triangle). As a negative control a canonical BiTE® antibody construct (circle) was used, which showed no significant CC1q binding.

10 **[270] Example 3: Pharmacokinetics of BiTE® antibody constructs fused to half-life extension modalities**

Two PSMA-targeting BiTE® antibody constructs were tested in the cynomolgus monkey in the context of pharmacokinetic (PK) studies. One BiTE® antibody construct was fused to a scFc half-life extending (HLE) moiety whereas the other one was used as single, non-half-life extended (non-HLO) “canonical” BiTE® antibody construct. The corresponding nomenclature of both molecules is briefly summarized in Table 4 below.

Table 4 Compound nomenclature of two single dosed BiTE® antibody constructs

compound synonyme	test compound name	target
Compound 11a	PM 76-B10-scFc	PSMA
Compound 11b	MTR112-5	

20 The BiTE®-HLE antibody construct (compound 11a) was administered as short intravenous bolus injection, the canonical antibody construct as a continuous intravenous infusion (compound 11b). To compare pharmacokinetic parameters for both antibody constructs, only the terminal phase starting directly after end of infusion is shown for the canonical PSMA-BiTE® antibody construct. The BiTE® antibody constructs were administered in a dose-linear, pharmacokinetic relevant range of 15 µg/kg (compound 11a) and 15.4 µg/kg/day (compound 11b), respectively. For reasons of comparability the serum concentrations shown are dose-normalized and molecular weight-normalized (indicated in nmol).

30 For each of the above named compounds a group of two animals was used. Blood samples were collected and serum was prepared for determination of serum concentrations. Serum BiTE® antibody construct levels were measured using an immunoassay. This sandwich ELISA assay is performed by capturing via a specific BiTE® targeting antibody, while an

antibody directed against the CD3-binding part of the construct was used for detection. The serum concentration-time profiles were used to determine PK parameters.

Blood sampling time points are listed for both study designs in Table 5 below.

Table 5 Blood sampling time points during PK studies.

blood sampling time points: HLE antibody construct; [h] after start of infusion	blood sampling time points: canonical antibody construct; [h] after start of infusion
0.085	167.76
0.25	168.96
0.5	172.08
1	175.92
4	180.00
8	192.00
24	
48	
72	
168	
240	
336	

The pharmacokinetics of the two BiTE® antibody constructs are shown exemplarily. Each compound group stands for the same BiTE® antibody construct either fused to a scFc half-life extending moiety or left as canonical molecule. For all proteins serum levels were quantifiable for all time points in all animals after BiTE®-HLE antibody construct administration. (**Figure 5**).

The pharmacokinetic parameters were determined using standard non-compartmental analysis (NCA) methods. Using non compartmental analysis, the following PK parameters were estimated: AUC_{inf} (Area under the serum concentration- time curve), V_{ss} (volume of distribution at steady state), CL (systemic clearance) and terminal $t_{1/2}$ (terminal half-life).

The PK parameter for each tested compound are summarized as mean of n=2 in Table 6 below.

Table 6 Pharmacokinetic parameter of two PSMA targeting BiTE® antibody constructs tested in cynomolgus monkeys.

test item	terminal $t_{1/2}$ [h]	AUC_{inf} [normalized to 15 $\mu\text{g/kg}$] [h*ng/mL]	Cl [mL/h/kg]	V_{ss} [mL/kg]
-----------	------------------------------	--	-----------------	---------------------

test item	terminal $t_{1/2}$ [h]	AUC_{inf} [normalized to 15 $\mu\text{g/kg}$] [h*ng/mL]	Cl [mL/h/kg]	Vss [mL/kg]
Compound 11a	214	43014	0.4	98
Compound 11b	6	7763	13.5	1132

Typically the PK profile for the canonical PSMA-BiTE® antibody constructs describes a very steeply declining serum concentration profile related to the clearance mechanism of these canonical proteins. The half-life extended PSMA-targeting BiTE®-scFc antibody construct shows a biphasic, exponential decline after single test item administrations in cynomolgus monkeys.

Overall, the PSMA-BiTE® antibody construct (Compound 11a) fused to a scFc HLE-modality shows a mean AUC_{inf} of 43014 h*ng/mL, a systemic clearance value of 0.4 mL/h/kg, as well as a corresponding volume of distribution of 98 mL/kg. Compound 11b, the canonical, i.e. non-half-life extended PSMA targeting BiTE® antibody construct shows a high clearance of 13.5 mL/h/kg leading to a low serum exposure of 7763 h*ng/mL.

The differences in pharmacokinetic behavior of the two different BiTE® antibody constructs tested exemplifies the general advantage of the half-life extended PSMA targeting BiTE®-scFc antibody construct over the corresponding canonical non-HLE version, especially in terms of residence time of the substance in the body.

[271] Example 4:

Preformulated drug substances containing purified Target A-hALB, Target A-hFc, and Target A-scFc antibody constructs respectively were buffer exchanged via ultrafiltration / diafiltration using membranes with a molecular weight cut-off (MWCO) of 10 kDa. Final formulation was achieved by adding concentrated stock solutions. Resulting formulations for each construct are listed in Table 7. The target protein concentration was 1.0 mg/mL. Formulated Target A antibody constructs were filled to 1 mL in type I glass vials which were stoppered with butyl rubber stoppers and crimped with aluminum seals. Filled vials were incubated at -20, 5, 25 and 37°C. One vial of each version was subjected to five freeze and thaw (F/T) cycles. Target freezing temperature was -29°C. Target thawing temperature was 2°C. The ramp rate was approximately 0.3 K/min.

Visual particles were assessed in accordance to the method described by Ph Eur 2.9.20 by trained operators. Visual particle counts per vial are depicted in Table 7. The number of

visual (larger than 125 μm) proteinaceous particles was higher for Target A-hFc if compared to both Target A-hALB and Target A-scFc.

Table 7: Number of visual proteinaceous particles per vial for stressed and unstressed (T0) samples containing different half-life extended anti-Target A BiTE® antibody constructs.

Construct	hALB	hFc		scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
Number of visible (>125 μm) proteinaceous particles per vial					
T0	0	0	1	0	0
5 F/T cycles	0	2	2	0	1
2w 5°C	0	2	2	0	0
2w 25°C	0	2	1	0	0
2w 37°C	0	2	2	0	0
4w -20°C	0	2	1	0	0
4w 5°C	0	1	2	0	0
4w 25°C	0	2	2	0	0
4w 37°C	0	2	2	0	0

The samples described above were also analyzed by size exclusion ultra-high performance chromatography (SE-UPLC) in order to quantify the percentaged content of high molecular weight species (HMWS). SE-UPLC was performed on an AcquityH-Class UPLC system (Waters) using an Acquity UPLC BEH200 SEC 150 mm column (Waters). Column temperature was set to 25°C. Separation of size variants was achieved by applying an isocratic method with a flow rate of 0.4 mL/min. The mobile phase was composed of 100 mM sodium phosphate, 250 mM NaCl at pH 6.8. The run time totals 6.0 minutes. Samples were held at 8°C within the autosampler until analysis. A total amount of 3 μg protein was injected. In order to avoid carry over an intermediate injection with 40% acetonitrile was performed after each sample. Detection was based on fluorescence emission (excitation at 280 nm, emission at 325 nm). Peak integration was performed using Empower® software. Relative area under the curve of HMWS was reported (Table 8).

Fc-based antibody constructs exhibited lower HMWS contents in the formulation variant G40MSuT than in K60RTrT independent on the stress condition. It became evident that Target A-scFc contained less HMWS than Target A-hFc in both G40MSuT as well as K60RTrT preparations. Target A-scFc in its preferred formulation (G40MSuT) was less prone to HMWS formation than Target A-hALB formulated in K60RTrT. In previous experiments this buffer showed improved stabilizing potential for hALB based constructs.

Table 8: Overview on HMWS contents in stressed and unstressed (T0) Target A-hALB, -hFc, and -scFc antibody construct preparations determined via SE-UPLC

Construct	hALB	hFc		scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
%HMWS					
T0	1.8	6.7	3.3	2.5	1.3
5 F/T cycles	2.0	7.2	4.1	3.0	1.5
2w 5°C	n.t.	n.t.	n.t.	2.9	1.1
2w 25°C	n.t.	6.6	2.7	2.4	0.5
2w 37°C	2.6	6.3	2.1	2.7	0.3
4w -20°C	5.9	8.7	1.6	6.6	0.3
4w 5°C	2.0	8.2	2.8	3.6	0.6
4w 25°C	2.2	6.8	2.6	2.7	0.4
4w 37°C	3.5	7.6	1.9	4.3	0.3

n.t. = not tested

- 5 The abundance of chemical modifications upon heat stress (incubation at 37°C) was monitored by peptide mapping. Protein samples were enzymatically digested and the resulting peptides were separated using reversed phase chromatography. The column eluate was directly injected into the ion source of a mass spectrometer for identification and quantitation of the peptides.
- 10 In order to achieve maximum coverage, two separate enzyme digests were performed: once with trypsin and once with chymotrypsin. In each case, the proteins were denatured with guanidinium chloride and then reduced with dithiothreitol (DTT). After incubation in DTT, free cysteine residues were alkylated by the addition of iodoacetic acid. Samples were then buffer exchanged into 50 mM Tris pH 7.8 for digestion. Trypsin and chymotrypsin were added to
- 15 separate reaction tubes at a ratio of 1:10 (sample:enzyme) each. Samples were digested for 30 min at 37°C and the reaction was quenched by adding trifluoroacetic acid.

A load of 5 µg of each digest was separately injected onto a Zorbax SB-C18 (Agilent #859700-902) reversed phase column equilibrated in 0.1% (V/V) formic acid (FA). A 156 minutes gradient of up to 90% acetonitrile containing 0.1% FA was used to elute the peptides

20 directly into the electrospray ion source of a Q-Exactive Plus mass spectrometer (Thermo Scientific). Data was collected in data dependent mode using a top 12 method in which a full scan (resolution 70 000; scan range 200-2000 m/z) was followed by high energy collision dissociation (HCD) of the 12 most abundant ions (resolution 17 500).

Peptides were identified based on accurate mass and tandem mass spectrum using in-house

25 software. Identifications were manually verified. Relative quantities of modified and unmodified peptides were calculated based on ion abundance using Pinpoint software (Thermo Scientific).

Percentages of chemical modifications of the complement determining regions (CDRs) and of the half-life extending portion (either hALB or Fc) detected in Target A-hALB, -hFc, and -scFc antibody construct preparations are given by Table 9. When comparing similar formulation conditions, it became obvious that overall, chemical modifications were least abundant in scFc constructs.

Table 9: Overview on chemical modifications in stressed and unstressed (T0) Target A-hALB, -hFc, and -scFc antibody construct preparations determined via peptide mapping

Construct	hALB	hFc		scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
%N101 deamidation (CDR)					
T0	0.1	0.2	0.2	0.2	0.2
2w 37°C	0.7	0.8	3.0	0.7	3.2
4w 37°C	1.3	n.t.	8.5	n.t.	6.4
%N162 deamidation (CDR)					
T0	3.0	1.7	1.9	2.3	2.5
2w 37°C	15.9	11.6	2.7	15.0	3.3
4w 37°C	26.8	n.t.	3.7	n.t.	4.1
%M279 oxidation (CDR)					
T0	0.6	1.4	1.6	0.6	1.0
2w 37°C	1.2	0.8	0.8	0.6	1.0
4w 37°C	0.9	n.t.	0.8	n.t.	0.6
%N348 deamidation (CDR)					
T0	0.5	3.2	3.3	0.5	0.9
2w 37°C	20.5	21.6	1.9	9.4	1.3
4w 37°C	22.8	n.t.	2.0	n.t.	2.9
%N351 deamidation (CDR)					
T0	0.2	2.9	2.6	0.5	1.0
2w 37°C	6.6	12.7	0.9	3.8	0.4
4w 37°C	8.7	n.t.	0.8	n.t.	0.8
%M530 oxidation (Fc)					
T0	n.a.	3.9	4.1	2.6	3.2
2w 37°C	n.a.	9.0	3.1	4.0	4.3
4w 37°C	n.a.	n.t.	3.4	n.t.	3.5
% N603 deamidation (Fc)					
T0	n.a.	1.3	1.9	1.3	1.4
2w 37°C	n.a.	7.9	4.6	7.0	5.6
4w 37°C	n.a.	n.t.	6.9	n.t.	8.1
%M706 oxidation (Fc)					
T0	n.a.	3.2	3.6	1.5	2.1
2w 37°C	n.a.	6.0	2.8	2.1	2.5
4w 37°C	n.a.	n.t.	2.6	n.t.	2.0
%M587 oxidation (hALB)					
T0	1.0	n.a.	n.a.	n.a.	n.a.
2w 37°C	2.2	n.a.	n.a.	n.a.	n.a.
4w 37°C	2.3	n.a.	n.a.	n.a.	n.a.
%M623 oxidation (hALB)					
T0	1.9	n.a.	n.a.	n.a.	n.a.
2w 37°C	2.4	n.a.	n.a.	n.a.	n.a.
4w 37°C	3.0	n.a.	n.a.	n.a.	n.a.

Construct	hALB	hFc		scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
%M798 oxidation (hALB)					
T0	1.4	n.a.	n.a.	n.a.	n.a.
2w 37°C	3.3	n.a.	n.a.	n.a.	n.a.
4w 37°C	3.5	n.a.	n.a.	n.a.	n.a.
%M829 oxidation (hALB)					
T0	8.9	n.a.	n.a.	n.a.	n.a.
2w 37°C	42.9	n.a.	n.a.	n.a.	n.a.
4w 37°C	44.1	n.a.	n.a.	n.a.	n.a.

n.a. = not applicable; n.t. = not tested

[272] Example 5:

Target A-hALB, -hFc, -scFc antibody construct formulated as described under Example 4 were subjected to a pH jump experiment. The concentration of the starting materials was 1.0 mg/mL. A volume of 0.38 mL of each starting material was filled in a glass vial. After preconditioning at 37°C the solutions were spiked with 20 fold phosphate buffered saline (PBS) which was composed of 0.090 M potassium phosphate, 0.480 M sodium phosphate (both dibasic), 0.052 M potassium chloride and 2.76 M NaCl. The spiked samples were incubated at 37°C for two weeks. After incubation they were analyzed by SE-UPLC using the method described under Example 4 and the percentaged content of HMWS was reported (Table 10). When comparing all constructs formulated in K60RTrT the HMWS content increased in the following order: hALB < scFc < hFc. Target A-scFc also showed a lower HMWS content than Target A-hFc when formulated in G40MSuT.

Table 10: Overview on HMWS contents in stressed (pH jump + 2w 37°C) Target A-hALB, -hFc, and -scFc antibody construct preparations determined via SE-UPLC

Construct	hALB	hFc		scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
%HMWS					
2w 37°C	1.5	8.3	7.1	5.4	5.1

[273] Example 6:

Target A-hALB, -hFc, and -scFc antibody construct formulated as described under Example 4 were subjected to agitation stress. The concentration of the starting materials was 1.0 mg/mL. A volume of 0.5 mL of each solution was filtered through an appropriate 0.22 µm filter and filled into 3cc glass vials. The vials were placed in a plastic box ensuring that the vials were not displaced within the box during agitation. The box was placed onto an orbital shaker. The samples were agitated at 500 rpm for 65 hours. Visual particles were assessed in accordance to the method described by Ph Eur 2.9.20. The method was conducted by

trained operators. Visual particle counts per vial are depicted in Table 11. Visible proteinaceous particles were only observed in Target A-hFc preparations.

Table 11: Number of visual proteinaceous particles per vial in agitated samples

Construct	hALB	hFc		scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
Number of visible (>125 µm) proteinaceous particles per vial					
65h, 500 rpm	0	1	1	0	0

- 5 Above samples were also analyzed by size exclusion ultra-high performance chromatography (SE-UPLC) in order to quantify the percentaged content of high molecular weight species (HMWS). The same method as described in Example 4 was applied. The HMWS contents of agitated samples are outlined by Table 12. The formation of HMWS was most pronounced in Target A-hFc antibody construct when comparing K60RTrT
10 preparations. HMWS were more abundant in Target A-hFc than in Target A-scFc.

Table 12: Overview on HMWS contents in stressed (pH jump + 2w 37°C) Target A-hALB, -hFc, and -scFc preparations determined via SE-UPLC

Construct	hALB	hFc		scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
%HMWS					
65h, 500 rpm	1.8	5.8	2.4	1.8	0.3

[274] Example 7:

- 15 Target A-hALB, -hFc, and -scFc antibody construct formulated as described under Example 4 were exposed to visible and UVA light (photo stress). Protein concentration totaled 1 mg/mL in all preparations. Protein solutions were filtered through a filter with 0.22 µm pore size and filled to 0.5 mL in type I glass vials. Target A-hALB and -scFc were subjected to two different tests including 0.2 MLux visible light / 25 W*h/m² UVA light and 1.2MLux visible light
20 / 173 W*h/m² respectively. Target A-hFc was subjected to two different tests including 0.2 MLux visible light without UVA light and 1.2 MLux visible light / 30 W*h/m² UVA light respectively. Chamber temperatures were adjusted to 25°C. After light exposure samples were analyzed by visible inspection (Table 13), SE-UPLC (Table 14) and peptide map (Table 15). Aforementioned methods were performed according to the procedures described under
25 Example 4. Although Target A-hALB, and -scFc antibody construct were exposed to higher doses of UVA light, no visible proteinaceous particles was observed whereas Target A-hFc antibody construct samples exhibited one visible proteinaceous particle per vial for both tests irrespective of the formulation.

Table 13: Overview on the number of visible proteinaceous particles per vial in Target A-hALB, -hFc, and -scFc antibody construct preparations determined after light exposure

Construct	hALB	hFc		scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
Number of visible (>125 µm) proteinaceous particles per vial					
T0	0	0	1	0	0
Test 1	0 ¹⁾	1 ²⁾	1 ²⁾	0 ¹⁾	0 ¹⁾
Test 2	0 ³⁾	1 ⁴⁾	1 ⁴⁾	0 ³⁾	0 ³⁾

¹⁾ 0.2 MLux visible light / 25 W*h/m² UVA light, ²⁾ 0.2 MLux visible light without UVA light,

³⁾ 1.2 MLux visible light / 173 W*h/m², ⁴⁾ 1.2 MLux visible light / 30 W*h/m²

HMWS increased in the following order Target A-hALB < -scFc < -hFc antibody construct when the protein was formulated in K60RTrT. HMWS could be reduced for Fc based constructs when formulated in G40MSuT. However HMWS were again less pronounced for Target A-scFc. Target A-hFc antibody construct revealed to be especially sensitive towards UVA light exposure.

Table 14: Overview on HMWS contents in Target A-hALB, -hFc, and -scFc antibody construct preparations determined after light exposure via SE-UPLC

Construct	hALB	hFc		scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
%HMWS					
T0	1.8	6.7	3.3	2.5	1.3
Test 1	1.8 ¹⁾	6.3 ²⁾	2.5 ²⁾	2.1 ¹⁾	0.4 ¹⁾
Test 2	2.0 ³⁾	11.0 ⁴⁾	2.1 ⁴⁾	2.4 ³⁾	0.3 ³⁾

¹⁾ 0.2 MLux visible light / 25 W*h/m² UVA light, ²⁾ 0.2 MLux visible light without UVA light,

³⁾ 1.2 MLux visible light / 173 W*h/m², ⁴⁾ 1.2 MLux visible light / 30 W*h/m²

Percentages of chemical modifications of the complement determining regions (CDRs) and of the half-life extending portion (either hALB or Fc) detected in Target A-hALB, -hFc, and -scFc antibody construct preparations are given by Table 15. When comparing similar formulation conditions, it became obvious that overall, chemical modifications were least abundant in scFc constructs.

Table 15: Overview on chemical modifications in Target A-hALB, -hFc, and -scFc antibody construct preparations determined after light exposure via peptide mapping

Construct	hALB	hFc		scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
%N101 deamidation (CDR)					
T0	0.1	0.2	0.2	0.2	0.2
Test 1	0.2 ¹⁾	n.t.	0.3 ²⁾	n.t.	0.5 ¹⁾
Test 2	0.2 ³⁾	n.t.	0.6 ⁴⁾	n.t.	0.7 ³⁾
%N162 deamidation (CDR)					
T0	3.0	1.7	1.9	2.3	2.5
Test 1	3.0 ¹⁾	n.t.	2.1 ²⁾	n.t.	2.7 ¹⁾
Test 2	3.6 ³⁾	n.t.	3.1 ⁴⁾	n.t.	2.8 ³⁾

Construct	hALB	hFc		scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
%M279 oxidation (CDR)					
T0	0.6	1.4	1.6	0.6	1.0
Test 1	0.8 ¹⁾	n.t.	2.6 ²⁾	n.t.	0.6 ¹⁾
Test 2	1.0 ³⁾	n.t.	6.3 ⁴⁾	n.t.	0.7 ³⁾
%N348 deamidation (CDR)					
T0	0.5	3.2	3.3	0.5	0.9
Test 1	0.4 ¹⁾	n.t.	2.7 ²⁾	n.t.	0.2 ¹⁾
Test 2	0.9 ³⁾	n.t.	3.9 ⁴⁾	n.t.	0.2 ³⁾
%N351 deamidation (CDR)					
T0	0.2	2.9	2.6	0.5	1.0
Test 1	0.4 ¹⁾	n.t.	2.0 ²⁾	n.t.	0.3 ¹⁾
Test 2	0.5 ³⁾	n.t.	2.6 ⁴⁾	n.t.	0.3 ³⁾
%M530 oxidation (Fc)					
T0	n.a.	3.9	4.1	2.6	3.2
Test 1	n.a.	n.t.	7.6 ²⁾	n.t.	3.1 ¹⁾
Test 2	n.a.	n.t.	21.8 ⁴⁾	n.t.	4.1 ³⁾
%M706 oxidation (Fc)					
T0	n.a.	3.2	3.6	1.5	2.1
Test 1	n.a.	n.t.	6.5 ²⁾	n.t.	1.8 ¹⁾
Test 2	n.a.	n.t.	17.8 ⁴⁾	n.t.	2.7 ³⁾
%M587 oxidation (hALB)					
T0	1.0	n.a.	n.a.	n.a.	n.a.
Test 1	1.5	n.a.	n.a.	n.a.	n.a.
Test 2	2.4	n.a.	n.a.	n.a.	n.a.
%M623 oxidation (hALB)					
T0	1.9	n.a.	n.a.	n.a.	n.a.
Test 1	4.0	n.a.	n.a.	n.a.	n.a.
Test 2	4.1	n.a.	n.a.	n.a.	n.a.
%M798 oxidation (hALB)					
T0	1.4	n.a.	n.a.	n.a.	n.a.
Test 1	2.1	n.a.	n.a.	n.a.	n.a.
Test 2	3.1	n.a.	n.a.	n.a.	n.a.
%M829 oxidation (hALB)					
T0	8.9	n.a.	n.a.	n.a.	n.a.
Test 1	31.0	n.a.	n.a.	n.a.	n.a.
Test 2	25.2	n.a.	n.a.	n.a.	n.a.

n.a. = not applicable; n.t. = not tested

[275] Example 8:

Target A-hALB antibody construct was formulated in K60RTrT and Target A-scFc antibody construct was formulated in G40MSuT according to the procedure described in Example 4.

- 5 Protein concentrations totaled 0.05 mg/mL. Glass (borosilicate, type I, 13 mm 3cc vial from West, Art. No. 68000375) and polypropylene test containers (2 mL with O-ring, e.g. from Sarstedt, Art No. 72.694.005) are filled with 500 µL of the test solution. The test solution was left for five minutes in the first test container. Then a 150 µL aliquot was sampled for analysis. The remaining test solution (350 µL) was transferred sequentially from one test container to the next (five containers in total). In each vial, the solution was left for five

10

minutes before the next transfer. The same pipette tip was used for each transfer step. The same test was performed using 30 mL polycarbonate bottles (Nalgene, PCS-000295 with closure, PP/20-415/ZTPE). For this container type the first container was filled with 5 mL. After a 150 μ L aliquot was sampled, the residual volume was transferred from one test container to the next (according to the procedure described above). Samples pulled from container #1 and #5 were analyzed by SE-UPLC (method as described under Example 4). In addition protein detection was carried out with a PDA detector (280 nm) in order to determine protein concentrations. Percentaged protein recovery from each test container is given by Table 16. It was shown that protein recovery was more pronounced for Target A-scFc antibody construct than for Target A-hALB antibody construct irrespective of the container type.

Table 16: Protein recovery from different container types for Target A-hALB, and -scFc antibody construct determined by SE-UPLC

Construct	hALB	scFc
Formulation	K60RTrT	G40MSuT
%Protein recovery (from nominal)		
Type I glass	80.0	92.0
Polypropylene	87.0	97.3
Polycarbonate	87.0	96.0

15 [276] Example 9:

[277] Target A-hALB antibody construct was formulated in K60RTrT and Target A-scFc antibody construct was formulated in K60RTrT and G40MSuT according to the procedure described in Example 4. The protein concentration totaled 1.0 mg/mL. 1950 μ L of each test solution was spiked with 50 μ L of a 1000 ppm silicon standard solution (Specpure from AlfaAesar, Art.No. 38717) resulting in a 25 ppm spike. An unspiked test solution served as control sample. The spiked test solution as well as the control sample were filled into 3cc type I glass vials and were incubated at 37°C for 24 hours. All samples were analyzed by SE-UPLC according to the method described in Example 4 in order to quantify the amount of HMWS (Table 17).

Table 17: Overview on HMWS contents in Target A-hALB, and -scFc preparations determined via SE-UPLC after spiking with 25 ppm silicon

Construct	hALB	scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT
Δ %HMWS (compared to unspiked control)			
25 ppm spike	1.0	1.0	0.2

[278] Example 10:

Preformulated drug substances containing purified Target C (cc)-hALB, Target C (cc)-hFc, and Target C (cc)-scFc antibody constructs respectively were buffer exchanged via ultrafiltration / diafiltration using membranes with a molecular weight cut-off (MWCO) of 10 kDa. Final formulation was achieved by adding concentrated stock solutions. Resulting formulations for each construct are listed in Table 18. The target protein concentration was 1.0 mg/mL. Formulated Target C (cc)- constructs were filled to 1 mL in type I glass vials which were stoppered with butyl rubber stoppers and crimped with aluminum seals. Filled vials were incubated at -20, 5, 25 and 37°C. One vial of each version was subjected to five freeze and thaw (F/T) cycles. Target freezing temperature was -29°C. Target thawing temperature was 2°C. The ramp rate was approximately 0.3 K/min. The samples described above were also analyzed by size exclusion ultra-high performance chromatography (SE-UPLC) in order to quantify the percentage content of high molecular weight species (HMWS). SE-UPLC was performed according to the method described under Example 4. When formulated in K60RTrT, HMWS increased in the following order in unstressed samples: scFc < hALB < hFc. The least pronounced increase in HMWS upon freeze thaw stress was observed for the scFc-construct. The hFc-antibody construct revealed to be most prone to HMWS formation at -20°C. HMWS contents increased after four weeks storage at 5°C. The HMWS formation under these conditions was more pronounced for Fc based constructs than for albumin based constructs. In K60RTrT no significant increases in HMWS were observed at elevated storage temperatures (25 and 37°C). When formulated in G40MSuT, all constructs revealed similar HMWS contents in unstressed samples. The increase during freeze thaw was more distinct for Fc based constructs if compared to the albumin based construct. In G40MSuT, the hFc-construct was least stable during storage at -20°C. Considerable increases in HMWS during liquid storage were only observed for the hALB-construct.

Table 18: Overview on HMWS contents in stressed and unstressed (T0) Target C (cc)-hALB, -hFc, and -scFc antibody construct preparations determined via SE-UPLC

Construct Formulation	hALB		hFc		scFc	
	K60RTrT	G40MSuT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
%HMWS						
T0	1.5	0.3	2.7	0.3	1.3	0.3
5 F/T cycles	2.0	0.5	3.1	0.7	1.6	0.7
2w -20°C	n.t	n.t	n.t	n.t	1.5	0.5
2w 5°C	n.t	n.t	n.t	n.t	1.8	0.2
2w 25°C	1.7	0.6	2.3	0.2	1.3	0.2
2w 37°C	1.9	0.7	1.8	0.2	1.2	0.2
4w -20°C	1.6	0.4	4.2	1.5	1.7	0.9
4w 5°C	1.9	0.3	3.3	0.3	2.1	0.4
4w 25°C	1.4	0.6	2.2	0.2	1.4	0.4
4w 37°C	1.3	0.7	2.0	0.1	1.4	0.3

n.t. = not tested

The abundance of chemical modifications upon heat stress (incubation at 37°C) was monitored by peptide mapping according to the method described in Example 4. Percentages of chemical modifications of the complement determining regions (CDRs) detected in Target C (cc)-hALB, -hFc, and -scFc preparations are given by Table 19. Overall, Target C (cc)-scFc exhibited the lowest amount of chemical modifications in the CDRs. It became evident that especially deamidations of the CDRs were least pronounced for the scFc construct.

Table 19: Overview on chemical modifications in stressed and unstressed (T0) Target C (cc)-hALB, -hFc, and -scFc antibody construct preparations determined via peptide mapping

Construct	hALB		hFc		scFc	
Formulation	K60RTrT	G40MSuT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
%M34 oxidation (CDR)						
T0	1.0	1.8	1.0	1.4	1.7	1.9
2w 37°C	0.9	1.3	0.9	1.1	1.0	1.7
4w 37°C	n.t.	n.t.	n.t.	1.6	n.t.	1.8
%D103 isomerization (CDR)						
T0	0.8	0.8	0.8	0.8	0.6	0.6
2w 37°C	4.0	4.6	4.5	4.4	5.8	7.3
4w 37°C	n.t.	n.t.	n.t.	8.0	n.t.	12.4
%M290 oxidation (CDR)						
T0	0.7	1.4	0.8	1	1.3	1.4
2w 37°C	0.7	1.0	0.8	0.8	0.8	1.3
4w 37°C	n.t.	n.t.	n.t.	1.2	n.t.	1.6
%N359 deamidation (CDR)						
T0	5.8	11.4	5.3	6.3	0.4	0.5
2w 37°C	19.3	5.8	11.2	2.8	7.0	0.9
4w 37°C	n.t.	n.t.	n.t.	2.9	n.t.	2.2
%N362 deamidation (CDR)						
T0	5.4	8.7	3.9	4.0	0.2	0.3
2w 37°C	13.5	3.6	6.7	1.2	3.1	0.3
4w 37°C	n.t.	n.t.	n.t.	1.4	n.t.	0.7

n.a. = not applicable; n.t. = not tested

[279] Example 11:

Target C (cc)-hALB, -hFc, and -scFc antibody constructs formulated as described under Example 4 were subjected to a pH jump experiment. The concentration of the starting materials was 1.0 mg/mL. A volume of 0.38 mL of each starting material was filled in a glass vial. After preconditioning at 37°C the solutions were spiked with 20 fold phosphate buffered saline (PBS) which was composed of 0.090 M potassium phosphate, 0.480 M sodium phosphate (both dibasic), 0.052 M potassium chloride and 2.76 M NaCl. The spiked samples were incubated at 37°C for two weeks. After incubation they were analyzed by SE-UPLC using the method described under Example 4 and the percentaged content of HMWS was

reported (Table 20). Target C (cc)-scFc constructs showed the lowest HMWS content after pH jump if compared to Target C (cc)-hALB and -hFc irrespective of the formulation.

Table 20: Overview on HMWS contents in stressed (pH jump + 2w 37°C) Target C (cc)-hALB, -hFc, and -scFc antibody construct preparations determined via SE-UPLC

Construct	hALB		hFc		scFc	
Formulation	K60RTrT	G40MSuT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
%HMWS						
2w 37°C	1.7	4.8	1.6	1.8	1.1	1.5

5

[280] Example 12:

Target C (cc)-hALB, -hFc, and -scFc antibody constructs formulated as described under Example 10 were subjected to agitation stress. The concentration of the starting materials was 1.0 mg/mL. A volume of 0.5 mL of each solution was filter through an appropriate 0.22 µm filter and filled into 3cc type I glass vials. The vials were placed in a plastic box ensuring that the vials were not displaced within the box during agitation. The box was placed onto an orbital shaker. The samples were agitated at 500 rpm for 65 hours. Samples were analyzed by SE-UPLC in order to quantify the percentaged content of high molecular weight species (HMWS). The same method as described in Example 4 was applied. The HMWS contents of agitated samples are outlined by Table 21. The formation of HMWS was least pronounced for Target C (cc)-scFc in either formulation.

Table 21: Overview on HMWS contents in stressed (pH jump + 2w 37°C) Target C (cc)-hALB, -hFc, and -scFc antibody construct preparations determined via SE-UPLC

Construct	hALB		hFc		scFc	
Formulation	K60RTrT	G40MSuT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
%HMWS						
65h, 500 rpm	2.1	0.8	2.3	0.4	1.3	0.2

[281] Example 13:

Target C (cc)-hALB, -hFc, and -scFc antibody constructs formulated as described under Example 4 were exposed to visible and UVA light (photo stress). Protein concentration totaled 1 mg/mL in all preparations. Protein solutions were filtered through a filter with 0.22 µm pore size and filled to 0.5 mL in type I glass vials. Target C (cc)-hALB and -scFc were subjected to two different tests including 0.2 MLux visible light / 25 W*h/m² UVA light and 1.2MLux visible light / 173 W*h/m² respectively. Target C (cc)-hFc was subjected to two different tests including 0.2 MLux visible light without UVA light and 1.2 MLux visible light / 30 W*h/m² UVA light respectively. Chamber temperatures were adjusted to 25°C. After light

20

25

Construct	hALB		hFc		scFc	
Formulation	K60RTrT	G40MSuT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
%HMWS						
T0	1.5	0.3	2.7	0.3	1.3	0.3
Test 1	1.8 ¹	0.3 ¹	2.5 ²	0.3 ²	1.4 ¹	0.3 ¹⁾
Test 2	4.6 ³⁾	1.1 ³⁾	6.0 ⁴⁾	0.7 ⁴⁾	1.5 ³⁾	0.3 ³⁾

Overall chemical modifications upon light exposure were least pronounced for Target C (cc)-scFc. Especially deamidations of the CDRs were formed to a higher extent in Target C (cc)-hALB and Target C (cc)-hFc. When comparing Fc based constructs it was revealed that Target C (cc)-scFc was less prone to chemical modifications of the Fc portion although the scFc construct was exposed to higher UVA light doses than the hFc-construct. Table 23 also lists the most abundant chemical modifications of the albumin portion in Target C (cc)-hALB demonstrating that the half-life extending portion of this construct was chemically more degraded than the Fc portions of Target C (cc)-hFc and -scFc antibody construct.

Construct	hALB		hFc		scFc	
Formulation	K60RTrT	G40MSuT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
%M34 oxidation (CDR)						
T0	1.0	1.8	1.0	1.4	1.7	1.9
Test 1	1.5 ¹⁾	n.t.	0.7 ²⁾	4.2 ²⁾	1.4 ¹⁾	1.2 ¹⁾
Test 2	1.7 ³⁾	n.t.	1.1 ⁴⁾	4.2 ⁴⁾	1.3 ³⁾	1.7 ³⁾
%D103 isomerization (CDR)						
T0	0.8	0.8	0.8	0.8	0.6	0.6
Test 1	0.8 ¹⁾	n.t.	0.9 ²⁾	0.9 ²⁾	0.8 ¹⁾	1.0 ¹⁾
Test 2	1.1 ³⁾	n.t.	1.2 ⁴⁾	1.4 ⁴⁾	1.0 ³⁾	1.3 ³⁾
%M290 oxidation (CDR)						
T0	0.7	1.4	0.8	1	1.3	1.4
Test 1	1.1 ¹⁾	n.t.	0.5 ²⁾	3.3 ²⁾	1.0 ¹⁾	0.9 ¹⁾
Test 2	1.4 ³⁾	n.t.	1.1 ⁴⁾	4.2 ⁴⁾	1.0 ³⁾	1.4 ³⁾
%N359 deamidation (CDR)						
T0	5.8	11.4	5.3	6.3	0.4	0.5
Test 1	10.8 ¹⁾	n.t.	4.5 ²⁾	5.6 ²⁾	0.4 ¹⁾	0.2 ¹⁾
Test 2	12.4 ³⁾	n.t.	10.3 ⁴⁾	3.6 ⁴⁾	0.6 ³⁾	0.2 ³⁾
%N362 deamidation (CDR)						

Construct	hALB		hFc		scFc	
Formulation	K60RTrT	G40MSuT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
T0	5.4	8.7	3.9	4.0	0.2	0.3
Test 1	8.8 ¹⁾	n.t.	3.4 ²⁾	3.5 ²⁾	0.3 ¹⁾	0.3 ¹⁾
Test 2	9.8 ³⁾	n.t.	6.4 ⁴⁾	2.3 ⁴⁾	0.5 ³⁾	0.3 ³⁾
%D510 isomerization (Fc)						
T0	n.a.	n.a.	0.4	0.4	0.5	0.4
Test 1	n.a.	n.a.	0.4 ²⁾	0.6 ²⁾	0.6 ¹⁾	0.5 ¹⁾
Test 2	n.a.	n.a.	0.7 ⁴⁾	0.7 ⁴⁾	0.7 ³⁾	0.7 ³⁾
%M541 oxidation (Fc)						
T0	n.a.	n.a.	2.8	3	3.7	4.1
Test 1	n.a.	n.a.	5.0 ²⁾	0.7 ²⁾	3.8 ¹⁾	3.8 ¹⁾
Test 2	n.a.	n.a.	17.5 ⁴⁾	18.4 ⁴⁾	4.8 ³⁾	5.5 ³⁾
%N614 deamidation (Fc)						
T0	n.a.	n.a.	1.2	1.2	1.6	1.5
Test 1	n.a.	n.a.	1.3 ²⁾	1.7 ²⁾	2.8 ¹⁾	2.2 ¹⁾
Test 2	n.a.	n.a.	6.1 ⁴⁾	1.9 ⁴⁾	1.9 ³⁾	2.3 ³⁾
%N673 deamidation (Fc)						
T0	n.a.	n.a.	0.3	0.3	0.0	0.0
Test 1	n.a.	n.a.	0.5 ²⁾	0.6 ²⁾	0.5 ¹⁾	0.6 ¹⁾
Test 2	n.a.	n.a.	0.5 ⁴⁾	0.6 ⁴⁾	0.5 ³⁾	1.5 ³⁾
%M717 oxidation (Fc)						
T0	n.a.	n.a.	2.1	2.4	2.5	2.8
Test 1	n.a.	n.a.	4.1 ²⁾	7.3 ²⁾	2.2 ¹⁾	2.3 ¹⁾
Test 2	n.a.	n.a.	13.7 ⁴⁾	13.5 ⁴⁾	2.8 ³⁾	3.8 ³⁾
%M598 oxidation (hALB)						
T0	1.0	n.t.	n.a.	n.a.	n.a.	n.a.
Test 1	2.3 ¹⁾	n.t.	n.a.	n.a.	n.a.	n.a.
Test 2	6.4 ³⁾	n.t.	n.a.	n.a.	n.a.	n.a.
%M809 oxidation (hALB)						
T0	1.8	n.t.	n.a.	n.a.	n.a.	n.a.
Test 1	3.5 ¹⁾	n.t.	n.a.	n.a.	n.a.	n.a.
Test 2	8.3 ³⁾	n.t.	n.a.	n.a.	n.a.	n.a.
%M840 oxidation (hALB)						
T0	12.8	n.t.	n.a.	n.a.	n.a.	n.a.
Test 1	32.0 ¹⁾	n.t.	n.a.	n.a.	n.a.	n.a.
Test 2	61.7	n.t.	n.a.	n.a.	n.a.	n.a.
%K1036 glycation (hALB)						
T0	10.1	n.t.	n.a.	n.a.	n.a.	n.a.
Test 1	10.2 ¹⁾	n.t.	n.a.	n.a.	n.a.	n.a.
Test 2	9.9 ³⁾	n.t.	n.a.	n.a.	n.a.	n.a.

¹⁾ 0.2 MLux visible light / 25 W*h/m² UVA light, ²⁾ 0.2 MLux visible light without UVA light,

³⁾ 1.2 MLux visible light / 173 W*h/m², ⁴⁾ 1.2 MLux visible light / 30 W*h/m²

[282] Example 14

5 Different BiTE® antibody constructs designed for targeting Target C including Target C-non half-life extended (non HLE, “canonical”), Target C-hALB (comprising human serum albumin), and Target C-scFc (comprising a scFC modality as third domain as described above) were formulated in a solution comprising 100 mM L-arginine hydrochloride, 8% (w/v) Trehalose dihydrate and Polysorbate 80. The target protein concentration was 1.0 mg/mL for

the hALB and scFc and 0.4 mg/mL for the non HLE version. Formulated BiTE® antibody constructs were filled to 1 mL in type I glass vials which were stoppered with butyl rubber stoppers and crimped with aluminum seals. Filled vials were incubated at -20°C and 37°C (w/o and with 25 ppm silicon which is known for its potential to induce aggregation of proteins) for 4 weeks. Above constructs were also exposed to light (1.2 MLux visible light / 173 W*h/m² UVA light). For light stress, chamber temperature was set to 25°C. Samples stored at -70°C served as controls (T0).

The samples described above were analyzed in duplicates by size exclusion ultra-high performance chromatography (SE-UPLC) in order to quantify the percentaged content of high molecular weight species (HMWS). SE-UPLC was performed on an Aquity H-Class UPLC system (Waters) using an Acquity UPLC BEH200 SEC 150 mm column (Waters). Column temperature was set to 25°C. Separation of size variants was achieved by applying an isocratic method with a flow rate of 0.4 mL/min. The mobile phase was composed of 100 mM sodium phosphate, 250 mM NaCl pH 6.8. The run time totals 6.0 minutes. Samples were held at 8°C within the autosampler until analysis. A total amount of 3 µg protein was injected. In order to avoid carry over an intermediate injection with 40% ACN was performed after each sample. Detection was based on fluorescence (excitation at 280 nm, emission at 325 nm). Peak integration was performed using Empower® software. Relative area under the curve of HMWS was reported (24).

Within non-stressed samples, HMWS were least pronounced for the scFc-construct. HMWS formation was exclusively observed during 4 weeks storage at -20°C. The HMWS contents under these conditions increase in the following order scFc < hALB < non HLE.

Table 24: Overview on HMWS contents in stressed and unstressed (T0) Target C -non HLE, -hALB, and -scFc preparations determined via SE-UPLC.

Construct	Non HLE (canonical)	hALB	scFc
T0	1.3%	1.3%	1.0%
4w -20°C	4.6%	1.8%	1.6%
4w 37°C	0.9%	0.6%	0.5%
4w 37°C (25 ppm silicon)	1.1%	0.8%	0.8%
Light exposure	1.0%	0.9%	0.5%

[283] Additionally, samples derived from heat stress in absence and presence of silicon were assessed for the abundance of subvisible particles by Microfluid Imaging (MFI) using a Flowcam from Fluid Imaging Technologies, Inc. The instrument was equipped with a FC80FV flow cell. A tenfold optical magnification was applied. System suitability was verified with particle free water. An autoimage rate of 20 frames per second was applied. Dark and light thresholds were set to 25 and 20 pixels respectively. Sample volume for a single measurement totals 0.25 mL. Samples were measured in triplicates. Prior to each triplicate the system was flushed of 0.5 mL of the respective sample solutions. At the beginning and between each triplicate a wash with 1.0 mL particle free water was performed. Data evaluation was performed with Visual Spreadsheet software. Samples were measured in triplicates. Results are outlined in Table 15.

Heat stress resulted in subvisible particle formation in preparations containing non HLE and hALB constructs. In contrast, the scFc construct remained stable. Subvisible particle formation was not promoted by the addition of silicon independent on the nature of the BiTE® antibody construct.

Table 15: Assessment of subvisible particles by MFI in Target F-non HLE (canonical), -hALB, and -scFc preparations after heat stress in absence and presence of silicon.

Construct	Non HLE (canonical)				hALB				scFc			
Particle size [µm]	≥ 2	≥ 5	≥ 10	≥ 25	≥ 2	≥ 5	≥ 10	≥ 25	≥ 2	≥ 5	≥ 10	≥ 25
T0	146	35	12	0	281	71	35	0	298	150	33	0
4w 37°C	410	163	23	0	742	225	11	0	110	44	0	0
4w 37°C (25 ppm silicon)	69	35	11	0	272	91	34	0	146	55	11	0

Samples from heat stress were also analyzed by Weak Cation Exchange (WCX) chromatography in order to quantify the percentaged content of charge variants using a UPLC Aquity H class from Waters. A Protein-Pak Hi Res CM 7µm 4.6 x 100 mm column (Waters, cat No. 186004929) was applied. The column temperature was adjusted to 30°C. The flow rate was set to 0.65 mL/min. The applied gradient was designed as follows (**Table 26**). The temperature of the autosampler was kept at 2-8°C.

Table 26: Gradient applied for WCX chromatography

Time [min:sec]	% A 20 mM sodium phosphate, pH 6.5	% B 20 mM sodium phosphate, 250 mM sodium chloride, pH 6.5
00:00	100	0
04:00	100	0
25:00	50	50
25:01	0	100
29:00	0	100
29:01	100	0
33:00	100	0

[284] A total amount of 3 µg of protein was injected. Detection was based on fluorescence (excitation at 280 nm, emission at 325 nm). Peak integration was performed using Empower® software. Relative areas under the curve of the main peak as well as of acidic and basic charge variants was reported (**Table27**).

Heat stress resulted in a reduced main peak percentage which had to be attributed to a predominant formation of acidic charge variants. The loss in main peak percentage was least pronounced for the scFc construct (7.5%). Basic charge variants were formed in both constructs with extended half-life upon light exposure. The increase in basic charge variants ranged between 5 and 6% in hALB and scFc constructs.

Table 27: Assessment of charge variants by WCX chromatography in Target F-non HLE (canonical), -hALB, and –scFc preparations after heat and light induced stress.

Construct	Non HLE (canonical)			hALB			scFc		
Fraction	% main	% acidics	% basics	% main	% acidics	% basics	% main	% acidics	% basics
T0	89.9	3.6	6.5	83.3	0.7	16.0	74.5	3.4	22.1
4w 37°C	79.3	11.1	9.5	75.6	9.8	14.6	67.0	11.2	21.8

In addition, sample purity was quantified in heat and light stressed samples using a microfluidic capillary electrophoresis sodium dodecylsulphate (CE-SDS) assay based on the

LabChip GXII system (Perkin Elmer). The sample denaturing solution was composed of the HT Protein Express Sample Buffer (provided by Perkin Elmer) supplemented with 34 mM dithiothreitol. Each sample was diluted 1:8 with the denaturing solution and heated up to 70°C for 10 minutes together with the protein express ladder. 35 µL of water for injection (WFI) were added to 40 µL of the denatured sample. 120 µL WFI were added to 12 µL of the ladder. Samples, ladder, protein express wash buffer, gel dye and destain solution are transferred to the respective reservoirs. Samples are electrokinetically loaded from a microtiter plate onto the chip integrating the separation, staining, destaining, and detection of the protein and its size variants. The resulting electropherograms were evaluated and changes in purity were reported. An overview on the percentaged purity detected post stress is given by **Table 28** and compared to unstressed samples (T0).

Higher purities were observed for hALB and scFc constructs if compared to the non HLE construct under all conditions. Slight decreases in purity if compared to T0 were detected for hALB and scFc constructs upon heat and light stress. The loss in purity after 4 weeks storage at 37°C totals 8.4% for the hALB construct and 6.6% for the scFc constructs. The losses upon light exposure were comparable between hALB and scFc.

Table 28: Overview on percentaged purity in stressed and unstressed (T0) Target F-non HLE, -hALB, and -scFc preparations determined via LabChip GXII (Caliper).

Construct	Non HLE (canonical)	hALB	scFc
T0	57.4	96.0	92.2
4w 37°C	60.6	87.6	85.6
Light exposure	61.5	90.1	86.4

[285] Example 15

Different BiTE® antibody constructs designed for targeting Target D including Target D-hALB and Target D-scFc were formulated at pH 7.0. The target protein concentration was 1.0 mg/mL for both constructs. Formulated BiTE® antibody constructs were filled to 1 mL in type I glass vials which were stoppered with butyl rubber stoppers and crimped with aluminum seals. Filled vials were incubated at 37°C (Target D-hALB) and 40°C (Target D-scFc) for 4 weeks. Samples stored at -70°C served as controls (T0). Samples were analyzed by SE-UPLC according to the method described under **Example 14**. Results are outlined in **Table 29**.

The scFc construct exhibited a reduced monomer loss (2.3%) upon heat stress if compared to the hALB construct (4.0%) although the incubation temperature was slightly higher.

Table 29: Overview on monomer peak percentage in stressed and unstressed (T0) Target D-hALB and –scFc preparations determined via SE-UPLC.

Construct	hALB	scFc
T0	97.6%	99.8%
4w	93.6%	97.5%

5

[286] Example 16

Different BiTE® antibody constructs designed for targeting PSMA including PSMA-non HLE (canonical) and PSMA-scFc were examined. The target protein concentration was 1.0 mg/mL. Formulated BiTE® antibody constructs were filled to 1 mL in type I glass vials which were stoppered with butyl rubber stoppers and crimped with aluminum seals. Filled vials were incubated at -20°C and 37°C (w/o and with 25 ppm silicon) for 4 weeks. Above constructs were also exposed to light (1.2 MLux visible light / 173 W*h/m² UVA light). For light stress chamber temperature was set to 25°C. Samples stored at -70°C served as controls (T0). All samples were analyzed by SE-UPLC according to the method described under **Example 14**. Results are outlined in **Table 30**.

15

The scFc construct showed higher monomer contents than the non HLE construct under all tested stress conditions indicating a lower propensity to the formation of high (aggregates) or low molecular weight species (fragments) of the scFc construct.

Table 30: Overview on monomer contents in stressed and unstressed (T0) PSMA-non HLE and -scFc preparations determined via SE-UPLC.

20

Construct	Non-HLE (canonical)	scFc
T0	97.0%	98.6%
4w -20°C	96.7%	98.9%
4w 37°C	88.3%	94.1%
4w 37°C (25 ppm silicon)	90.4%	96.1%
Light exposure	96.1%	99.0%

Additionally, samples derived from heat stress in absence and presence of silicon were assessed for the abundance of subvisible particles by Microfluid Imaging (MFI) using the method described under **Example 14**. Results are outlined in **Table 31**.

5 The scFc construct revealed a significantly lower abundance of subvisible particles than the non HLE construct when subjected to heat stress in absence and presence of silicon.

Table 31: Assessment of subvisible particles by MFI in PSMA-non HLE (canonical) and –scFc preparations after heat stress in absence and presence of silicon.

Construct	Non-HLE (canonical)				scFc			
Particle size [µm]	≥ 2	≥ 5	≥ 10	≥ 25	≥ 2	≥ 5	≥ 10	≥ 25
4w 37°C	4675	2334	1017	22	2518	1364	655	24
4w 37°C (25 ppm silicon)	8249	4243	1906	34	965	473	179	11

10 Samples from heat and light stress were also analyzed by Weak Cation Exchange (WCX) chromatography in order to quantify the percentaged content of charge variants using a UPLC Aquity H class from Waters according to the method described under **Example 14**. Relative areas under the curve of the main peak as well as of acidic and basic charge variants was reported (**Table 32**).

15 The scFc construct showed higher main peak percentages (lower amount of charge variants) than the non HLE construct. This difference was most pronounced when comparing samples subjected to heat stress indicating a higher chemical stability of the scFc construct.

Table 32: Assessment of charge variants by WCX chromatography in PSMA-non HLE (canonical) and –scFc preparations after heat and light induced stress.

Construct	Non HLE (canonical)			scFc		
Fraction	% main	% acidics	% basics	% main	% acidics	% basics
T0	66.1	18.4	15.6	70.4	14.5	15.1
4w 37°C	32.1	51.8	16.1	41.5	45.6	12.9
Light exposure	68.0	24.6	7.4	69.9	12.6	17.5

In addition, sample purity was quantified in heat and light stressed samples using a microfluidic capillary electrophoresis sodium dodecylsulphate (CE-SDS) assay based on the LabChip GXII system (Perkin Elmer) according to the method described under **Example 14**.

- 5 An overview on the percentaged purity detected post stress is given by **Table33** and compared to unstressed samples (T0).

Higher purities were observed for the scFc construct if compared to the non HLE construct irrespective of the stress conditions. Hence, as exemplified hereby, scFc antibody constructs are more stable over a broad array of stress conditions than non-HLE antibody constructs and, thus, safer and better suited for versatile clinical employment with ever changing stress conditions in practice.

10

Table 33: Overview on percentaged purity in stressed and unstressed (T0) PSMA-non HLE, and -scFc preparations determined via LabChip GXII.

Construct	Non-HLE (canonical)	scFc
T0	89.2	100.0
4w 37°C	85.4	87.3
Light exposure	89.5	93.8

15 **[287] Example 17**

Different BiTE® antibody constructs designed for targeting Target B including Target B-Xbody and Target B-scFc were examined. The target protein concentration was 1.0 mg/mL. Formulated BiTE® antibody constructs were filled to 1 mL in type I glass vials which were stoppered with butyl rubber stoppers and crimped with aluminum seals. Filled vials were incubated at -20°C and 37°C for 4 weeks. Additionally, all samples were exposed to 1.2 MLux visible light and 173 W*h/m² UVA light. Chamber temperature was adjusted to 25°C. Samples stored at -70°C served as controls (T0). Samples stored at -20 and -37°C were analyzed by SE-UPLC according to the method described under **Example 14**. Results are outlined in **Table34**.

20

- 25 The scFc construct preserved a higher monomer content when stored for four weeks at -20 and 37°C respectively if compared to the Xbody.

Table 34: Overview on monomer contents in stressed and unstressed (T0) Target B-Xbody and -scFc preparations determined via SE-UPLC.

Construct	Xbody	scFc
T0	100.0	98.8
4w -20°C	97.1	97.9
4w 37°C	94.5	95.7

Additionally, unstressed samples were assessed for the abundance of subvisible particles by Microfluid Imaging (MFI) using the method described under **Example 14**. Results are outlined in **Table35**. The CD19-scFc preparation exhibited significantly lower amounts of subvisible particles if compared to the CD19-Xbody preparation. This applies to all included size fractions.

Table 35: Assessment of subvisible particles by MFI in unstressed Target B-Xbody and -scFc

Construct	Xbody				scFc			
Particle size [µm]	≥ 2	≥ 5	≥ 10	≥ 25	≥ 2	≥ 5	≥ 10	≥ 25
T0	2648	688	192	32	160	64	43	11

Samples from light stress were also analyzed by Weak Cation Exchange (WCX) chromatography in order to quantify the percentaged content of charge variants using a UPLC Aquity H class from Waters according to the method described under **Example 14**. Relative areas under the curve of the main peak as well as of acidic and basic charge variants was reported (**Table36**).

Table 36: Assessment of charge variants by WCX chromatography in Target B-Xbody and -scFc preparations after heat and light induced stress.

Construct	Xbody			scFc		
Fraction	% main	% acidics	% basics	% main	% acidics	% basics
T0	51.4	30.3	18.3	83.5	1.3	15.2
Light exposure	45.9	33.2	20.9	82.1	1.2	16.7

The scFc construct showed surprisingly enhanced stability against light exposure if compared to the Xbody indicated by a less pronounced loss in main peak which totaled 1.4%

compared to 5.5% for the Xbody construct. Hence, it can be derived from said exemplarily setting that an anti-PSMA antibody construct provided with a scFc domain as HLE is superior to other antibody constructs comprising different HLE in terms of stress resistance and, in consequence, features surprisingly improved stability.

5

[288] Example 18 Size exclusion chromatography of bispecific scFc variants

The constructs D9F, T2G, D3L, T7I and K6C (see Figure 6) were each tested for their running behavior by size exclusion chromatography according to standard procedures. In detail, a defined amount of 25 µg of each construct were run (at 750µl/min) in Citrate Lysin Buffer (10 mM and 75 mM, pH7) on a Superdex 200 increase 10/300GL column at room temperature and the OD 280 nm was recorded. Subsequently, constructs have been compared by their retention times. As a result, construct D9F shows significantly delayed elution (Table 37) as compared to T2G, D3L, T7I and K6C, which indicates a difference in the structure/arrangement of the Fc domains. This difference in retention time was most significant with construct T7I having unpaired cysteines in the hinge region and the linkage of CH2 and CH2CH3 to CH3 (18.98 min vs. 18.62 min, difference of 0.36 min). However, also the difference in retention time of 0.16 min between D9F and T2G is significant taking the respective retention time of the BSA control into consideration. The BSA control showed a retention time of 19.07 min for the monomer and 16.82 min for the dimer displaying a difference of 2.25 min in retention time for a doubled molecular weight. Hence, as the constructs having only structural differences in the Fc part, 0.16 min difference in retention time are significant. In summary, construct D9F showed the longest retention time indicating the strongest binding. This conclusion leads to the expectation of D9F also has the longest half live *in vivo*.

Table 37

Construct	Retention time in min
D9F	18.98
T2G	18.82
D3L	18.78
K6C	18.77
T7I	18.62

BSA monomer	19.07
BSA dimer	16.82

[289] Example 19: Surface Plasmon Resonance -based determination of binding to human FcRn (FCGRT/B2M)

The constructs D9F, T2G, D3L, T7I and K6C (Figure 6) were each tested for their capability of binding against human FcRn in SPR (Biacore) experiments according to standard procedures. In detail, CM5 Sensor Chips (GE Healthcare) were immobilized with 450-500 RU of FCGRT/B2M (ACRO Biosystems) by using Na acetate buffer pH 4.5 and a running buffer consisting of 200 mM HEPES, 150 mM NaCl, 3 mM EDTA pH 6.0. The constructs were then injected in subsequent runs in two concentrations of 250 nM and 125 nM diluted in 200 mM HEPES, 150 mM NaCl, 3 mM EDTA, pH 6.0 and 36°C. Association was done for 90 seconds with a 30 µl/min flow rate followed by the dissociation phase for 90 seconds at a 30 µl/min flow rate in 200 mM HEPES, 150 mM NaCl, 3 mM EDTA, pH 6.0 at 36°C. Subsequent regeneration was done for 10 sec with 30 µl/min with 10 mM HEPES, 150 mM NaCl, 3 mM EDTA pH 7.4.

The maximal binding during the injection phase was measured for all constructs as the respective response units (RU), equivalent to the molecular mass increase on the FcRn coated CM5 chip due to bound construct. All constructs were measured in duplicates. Average values of the duplicate determinations are depicted in Figure 7A and 7B, respectively.

As a result, construct D9F shows significantly higher mass increase on the FcRn coated CM5 chip, as compared to T2G, D3L, T7I and K6C, which indicates stronger binding affinity of D9F to human FcRn. This observation was seen for both concentrations of the respective constructs.

The binding against FcRn is mediated through the Fc portion within the constructs. Stronger binding against human FcRn as described in the literature is an indicator for longer half-life in vivo due to a higher intracellular rescue of the respective protein and a therefore reduced degradation rate. For this reason, stronger binding of D9F to human FcRn as compared to the other constructs makes this molecule clearly superior as a basis for therapeutic molecules to allow for longer exposure of the potential drug in the patient and a lower frequency of drug administration.

[290] Example 20: Surface Plasmon Resonance-based determination of binding to human FcRn (FCGRT/B2M)

The constructs D9F, T2G, D3L, T7I and K6C and a human IgG1-kappa antibody MT201 were each tested for their capability of binding against human FcRn in SPR (Biacore) experiments according to standard procedures. In detail, CM5 Sensor Chips (GE Healthcare) were immobilized with around 350 RU of FCGRT/B2M (ACRO Biosystems) by using Na acetate buffer pH 4.5 and a running buffer consisting of 200 mM HEPES, 150 mM NaCl, 3 mM EDTA pH 6.0. The constructs and the human IgG1-kappa control (MT201) were then injected at a concentration of 125 nM diluted in 200 mM HEPES, 150 mM NaCl, 3 mM EDTA, pH 6.0 and 36°C. Association was done for 90 seconds with a 30 µl/min flow rate followed by the dissociation phase for 60 seconds at a 30 µl/min flow rate in 200 mM HEPES, 150 mM NaCl, 3 mM EDTA, pH 6.0 at 36°C. Subsequent regeneration was done for 10 sec with 30 µl/min with 10 mM HEPES, 150 mM NaCl, 3 mM EDTA pH 7.4.

The maximal binding during the injection phase was measured for all constructs as the respective response units (RU), equivalent to the molecular mass increase on the FcRn coated CM5 chip due to bound construct. All constructs were measured in duplicates. Average values of the duplicate determinations are depicted in Figure 8 including standard deviation error bars.

As a result, construct D9F shows significantly higher mass increase on the FcRn coated CM5 chip, as compared to T2G, D3L, T7I and K6C, which indicates stronger binding affinity of D9F to human FcRn. The mass increase on the FcRn-coated CM5 chip for D9F is well comparable to the mass increase of the human IgG1-kappa control antibody MT201, indicating a comparable binding of construct D9F to human FcRn.

The binding against FcRn is mediated through the human IgG1 Fc portion within the constructs. Stronger binding against human FcRn as described in the field is an indicator for longer half-life in vivo due to a higher intracellular rescue of the respective protein and a therefore reduced degradation rate. For this reason, stronger binding of D9F to human FcRn in the range of a human IgG1-kappa antibody (MT201), as compared to the other constructs makes this molecule clearly superior as a basis for therapeutic molecules to allow for longer exposure of the potential drug in the patient, presumably in the range of a full human IgG1 antibody, and a lower frequency of drug administration.

Table 38: Sequence Table

SEQ ID NO:	Designation	Format / Source	Sequence
------------	-------------	-----------------	----------

1.	G4S linker		GGGGS
2.	(G4S)2 linker		GGGGS GGGGS
3.	(G4S)3 linker		GGGGS GGGGS GGGGS
4.	(G4S)4 linker		GGGGS GGGGS GGGGS GGGGS
5.	(G4S)5 linker		GGGGS GGGGS GGGGS GGGGS GGGGS
6.	(G4S)6 linker		GGGGS GGGGS GGGGS GGGGS GGGGS GGGGS
7.	(G4S)7 linker		GGGGS GGGGS GGGGS GGGGS GGGGS GGGGS GGGGS
8.	(G4S)8 linker		GGGGS GGGGS GGGGS GGGGS GGGGS GGGGS GGGGS GGGGS
9.	Peptide linker		P GGGGS
10.	Peptide linker		P GGDGS
11.	Peptide linker		S GGGGS
12.	Peptide linker		G GGG
13.	CD3ε binder VL		QTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNYPNWWQQKPGQAPR GLIGGTKFLAPGTPARFSGSL LGGKAALTL SGVQPEDEAEYYC VLWY SNRWVFGGGTKLTVL
14.	CD3ε binder VH		EVQLVESGGGLVQP GGSRLRLSCAASGFTFNSYAMNWRQAPGKGLEW VARIRSKYNNYATYYADSVKGRFTISRDDSKNTAYLQMNSLKTEDTA VYYCVRHGNFGNSYVSWWAYWGQGLTVTVSS
15.	CD3ε binder scFv		EVQLVESGGGLVQP GGSRLRLSCAASGFTFNSYAMNWRQAPGKGLEW VARIRSKYNNYATYYADSVKGRFTISRDDSKNTAYLQMNSLKTEDTA VYYCVRHGNFGNSYVSWWAYWGQGLTVTVSSGGGGSGGGSGGGGSQ TVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNYPNWWQQKPGQAPRG LIGGTKFLAPGTPARFSGSL LGGKAALTL SGVQPEDEAEYYC VLWYS NRWVFGGGTKLTVL
16.	hexa-histidine tag		HHHHHH
17.	Fc monomer-1 +c/-g		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVDS HEDPEVKFNWYVDGVEVHNAKTKPCEEQYGYSTYRCVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYS KLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
18.	Fc monomer-2 +c/-g/delGK		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVDS HEDPEVKFNWYVDGVEVHNAKTKPCEEQYGYSTYRCVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYS KLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSP
19.	Fc monomer-3 -c/+g		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVDS HEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYS KLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
20.	Fc monomer-4 -c/+g/delGK		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVDS HEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYS KLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSP
21.	Fc monomer-5 -c/-g		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVDS HEDPEVKFNWYVDGVEVHNAKTKPREEQYGYSTYRVVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYS KLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
22.	Fc monomer-6 -c/-g/delGK		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVDS HEDPEVKFNWYVDGVEVHNAKTKPREEQYGYSTYRVVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN

			QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYS KLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSP
23.	Fc monomer- 7 +c/+g		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYNSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYS KLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
24.	Fc monomer- 8 +c/+g/delGK		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYNSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYS KLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSP
25.	scFc-1		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
26.	scFc-2		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSP
27.	scFc-3		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
28.	scFc-4		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSP
29.	scFc-5		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYGSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK

			KPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPR EEQYGSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKA KGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQ PENNYKTTTPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALH NHYTQKSLSLSPGK
30.	scFc-6		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDV HEDPEVKFNWYVDGVEVHNAKTKPREEQYGSTYRVVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYS KLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGGGSGGGGS GGGSGGGSGGGSGGGSGGGSDKTHTCPPCPAPELLGGPSVFLFPPK KDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREE QYGSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKG QPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTTPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALHNH YTQKSLSLSP
31.	scFc-7		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDV HEDPEVKFNWYVDGVEVHNAKTKPCEEQYNSTYRCVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYS KLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGKGGGSGGGG SGGGSGGGSGGGSGGGSGGGSDKTHTCPPCPAPELLGGPSVFLFPP KPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPC EEQYNSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKA KGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQ PENNYKTTTPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALH NHYTQKSLSLSPGK
32.	scFc-8		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDV HEDPEVKFNWYVDGVEVHNAKTKPCEEQYNSTYRCVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYS KLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGGGSGGGGS GGGSGGGSGGGSGGGSGGGSDKTHTCPPCPAPELLGGPSVFLFPPK KDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEE QYNSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKG QPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTTPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALHNH YTQKSLSLSP
33.	Target A-HLE	Hetero Fc chain 1	QVQLVESGGGLVKPGGSLRLSCAASGFTTFSDYYMTWIRQAPGKGLEW LSYISSSGSTIYYADSVKGRFTISRDNANKSLFLQMNSLRAEDTAVY YCARDRNSHFDYWQGTLVTVSSGGGSGGGGSGGGGSDIQMTQSPS SVSASVGRVTITCRASQGINTWLAWYQQKPKAPKLLIYGASGLQS GVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQAKSFPRFTFGQGTK VEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWV RQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQ MNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQTLVTVSSGGGSG GGGSGGGGSGTVVTQEPSTVSPGGTVTLTCSGSTGAVTSGNYPNWV QQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLGVPQPEDE AEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPS VFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE KTISKAKGQPREPQVYTLPPSRKEMTKNQVSLTCLVKGFYPSDIAVE WESNGQPENNYKTTTPVLKSDGSFFFLYSKLTVDKSRWQQGNVSCSV MHEALHNHYTQKSLSLSPGK
34.	Target A -HLE	Hetero Fc chain 2	DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDV HEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN QVSLTCLVKGFYPSDIAVEWESNGQPENNYDTTPVLDSDGSFFLYS DLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
35.	Target A -HLE	hALB fusion	QVQLVESGGGLVKPGGSLRLSCAASGFTTFSDYYMTWIRQAPGKGLEW LSYISSSGSTIYYADSVKGRFTISRDNANKSLFLQMNSLRAEDTAVY YCARDRNSHFDYWQGTLVTVSSGGGSGGGGSGGGGSDIQMTQSPS

			SVSASVGDVRTITCRASQGINTWLAWYQQKPGKAPKLLIYGASGLQS GVPSRFSGSGSGTDFTLTISSSQPEDFATYYCQQAQSFPRFTFGQGTK VEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWW RQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQ MNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSG GGGSGGGGSQTIVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNYPNWV QQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDE AEYYCVLWYSNRWVFGGGTKLTVLPGGDGS DAHKSEVAHRFKDLGEE NFKALVLI AFAQYLQQCPFEDHVKL VNEVTEFAKTCVADESAENCDK SLHTLFGDKLCTVATLRETYGEMADCCAKQEPERNECF LQHKDDNP LPRLVRPEVDVMCTAFHDNEETFLKKYLYEIARRHPYFYAPELLFFA KRYKAAFTECCQAADKAACLLPKLDELRLDEGKASSAKQRLKCA SLQK FGERAFKAWAVARLSQRFPKAEFAEVSKLVTDLT KVHTECCHGDLLE CADDRADLAKYICENQDSISSKLKECCEKPLLEKSHCIAEVENDEMP ADLP SLAADFVESKDVCKNYAEAKDVFLGMFLYEYARRHPDYSVVL LRLAKTYETTLEKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNC E LFEQLGEYKFQNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKH PEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVTCKCTESLVNRRPC FSALEVDETYVPKEFNAETFTFHADICTLSEKERQIKKQTALVELVK HKPKATKEQLKAVMDDFAAFVEKCKADDKETCF AEEGKKLVAASQA ALGLHHHHHH
36.	Target B-HLEa	X-body chain 1	EVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWWRQAPGKGLEW VARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTA VYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGSSYELTQ PPSVSVSPGQTASITCSGDRLEGEKYSWYQQRPQGSPLLVIYQDTR K PSGIPERFSGSNSGNTATLTISGTQAMDEADYYCQAWESSTVVF GGG TKLTVLASTKGPSVFP LAPSSKSTSGGTAALGCLVKDYFPEPVT VSW NSGALTSQVHTFPAVLQSSGLYSLSVVTVPSSSLGTQTYICNVN HK PSENTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKD TLM ISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQY GST YRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQ PREP QVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPEN NYDT TPPVLDSGDSFFLYSDLTVDKSRWQQGNVFCSSVMHEALHNHY TQKS LSLSPGK
37.	Target B-HLEb	X-body chain 2	QVQLVESGGGVVQPGGSLRLSCAASGFTFSSYGMHWVRQAPGKGLEW VAFIWIYEGSNKYAAESVKDRFTISRDN SKNTLYLQMNSLRAEDTAVY YCARRAGIIGTIGYYYGMDVWGQGT TVTVSSGGGGSGGGGSQTIVVT Q EPSTLVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIG GT KFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCVLWYSNR WVF GGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYP GAV TVAWKADSSPVKAGVETTTTPSKQSNKYAASSYLSLTPEQWKSH RSY SCQVTHEGSTVEKTVAPTECSDKTHTCPPCPAPELLGGPSVFL FPPK PKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAK TKPCE EQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKT ISKAK GQPREPQVYTLPPSRKEMTKNQVSLTCLVKGFYPSDIAVEW ESNGQP ENNYKTTTPVLKSDGSFFLYSKLTVDKSRWQQGNVFCSSVM HEALHN HYTQKSLSLSPGK
38.	Target B-HLE	Hetero Fc chain 1	QVQLVESGGGVVQPGGSLRLSCAASGFTFSSYGMHWVRQAPGKGLEW VAFIWIYEGSNKYAAESVKDRFTISRDN SKNTLYLQMNSLRAEDTAVY YCARRAGIIGTIGYYYGMDVWGQGT TVTVSSGGGGSGGGGS YELTQPPSVSVSPGQTASITCSGDRLEGEKYSWYQQRPQGSPLLVIY QDTRKPSGIPERFSGSNSGNTATLTISGTQAMDEADYYCQAWESSTV VFGGGTKLTVLSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFN KYAMNWWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD S KNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTV S SGGGGSGGGSGGGGSQTIVVTQEPSTLVSPGGTVTLTCGSSTGAVT S GNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTL S GVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCP AP ELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWY V DGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNK ALPAPIEKTISKAKGQPREPQVYTLPPSRKEMTKNQVSLTCLVKGFY PSDIAVEWESNGQPENNYKTTTPVLKSDGSFFLYSKLTVDKSRWQQG

			NVFSCSVMHEALHNHYTQKSLSLSPGK
39.	Target B -HLE	Hetero Fc chain 2	DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYDTPPVLDSDGSFFFLYSDLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
40.	Target C-HLE	Hetero Fc chain 1	QVQLVQSGAEVKKPGESVKVSCKASGYTFTNYGMNWVKQAPGQCLEWMGWINTYTGPTYADKFKQGRVTMTTDTSTSTAYMEIRNLGGDDTAVYYCARWSWSDGYVYFDYWGQGTSTVTVSSGGGGSGGGSGGGSDIVMTQSPDSLTVSLGERTTINCKSSQSVLDSSTNKNLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDSPQPEDSATYYCQQSAHFPITFGCGTRLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLVTVSSGGGGSGGGSGGGSGQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAA LTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRKEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLKSDGSFFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
41.	Target C -HLE	Hetero Fc chain 2	DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYDTPPVLDSDGSFFFLYSDLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
42.	PM_1xCD3-scFc	VH CDR1	DYYMN
43.	PM_1xCD3-scFc	VH CDR2	IISDGGYYTYYSDIKIG
44.	PM_1xCD3-scFc	VH CDR3	GFPLLRHGAMDY
45.	PM_1xCD3-scFc	VL CDR1	KASQNVDTNVA
46.	PM_1xCD3-scFc	VL CDR2	SASYRYS
47.	PM_1xCD3-scFc	VL CDR3	QQYDSYPYT
48.	PM_1xCD3-scFc	VH	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKGLEWVAIISDGGYYTYYSDIKGRFTISRDNANKSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMDYWGQGTLLVTVSS
49.	PM_1xCD3-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSASYRYSVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDSYPYTFGGGKLEIK
50.	PM_1xCD3-scFc	scFv	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKGLEWVAIISDGGYYTYYSDIKGRFTISRDNANKSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMDYWGQGTLLVTVSSGGGGSGGGSGGGSDIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSASYRYSVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDSYPYTFGGGKLEIK
51.	PM_1xCD3-scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKGLEWVAIISDGGYYTYYSDIKGRFTISRDNANKSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMDYWGQGTLLVTVSSGGGGSGGGSGGGSDIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSASYRYSVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDSYPYTFGGGKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGSGGGSGQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAA LTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRKEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLKSDGSFFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

			PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
52.	PM_1xCD3- scFc	bispecific HLE molecule	QVQLVESGGGLVQPGESLRLSKAASGFTFSDYYMNWVRQAPGKGLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSKAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGGG GGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
53.	PM_1xCD3- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVQPGESLRLSKAASGFTFSDYYMNWVRQAPGKGLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSKAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
54.	PM_1_CCxCD 3-scFc	VH	QVQLVESGGGLVQPGESLRLSKAASGFTFSDYYMNWVRQAPGKCLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSS
55.	PM_1_CCxCD 3-scFc	VL	DIQMTQSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSL IYSASYRYSVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDSY PYTFGCGTKLEIK
56.	PM_1_CCxCD 3-scFc	scFv	QVQLVESGGGLVQPGESLRLSKAASGFTFSDYYMNWVRQAPGKCLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDSYPYTFG CGTKLEIK
57.	PM_1_CCxCD 3-scFc	bispecific molecule	QVQLVESGGGLVQPGESLRLSKAASGFTFSDYYMNWVRQAPGKCLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSKAASGFTFNKYA

			MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGSGGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
58.	PM_1_CCxCD 3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLR LSCAASGFTFSDYYMNWVRQAPGKCLEW VAIISDGGYYTYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISSLQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGSGGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGGG GGSGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
59.	PM_1_CCxCD 3-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLR LSCAASGFTFSDYYMNWVRQAPGKCLEW VAIISDGGYYTYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISSLQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGSGGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGGGGSGGGSGGGSGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
60.	PM_2xCD3- scFc	VH CDR1	DYYMY
61.	PM_2xCD3- scFc	VH CDR2	IISDAGGLTYSDIIKG
62.	PM_2xCD3- scFc	VH CDR3	GFPLLRHGAMDY
63.	PM_2xCD3- scFc	VL CDR1	KASQNVDTNVA
64.	PM_2xCD3- scFc	VL CDR2	SASYRYS
65.	PM_2xCD3- scFc	VL CDR3	QQYDSYPYT

66.	PM_2xCD3-scFc	VH	QVQLVESGGGLVKPAGESLRSLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDAGGLTYYSDI IKGRFTISRDNANKNSLYLQINSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLVTVSS
67.	PM_2xCD3-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSL IYSASYRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDSY PYTFGGGTGLEIK
68.	PM_2xCD3-scFc	scFv	QVQLVESGGGLVKPAGESLRSLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDAGGLTYYSDI IKGRFTISRDNANKNSLYLQINSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLVTVSSGGGSGGGSGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDSYPYTFG GGTKLEIK
69.	PM_2xCD3-scFc	bispecific molecule	QVQLVESGGGLVKPAGESLRSLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDAGGLTYYSDI IKGRFTISRDNANKNSLYLQINSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLVTVSSGGGSGGGSGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGG GGSGGGSGGGGSQTVVTQEP SLTVSPGGTVTLT CGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
70.	PM_2xCD3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPAGESLRSLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDAGGLTYYSDI IKGRFTISRDNANKNSLYLQINSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLVTVSSGGGSGGGSGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGG GGSGGGSGGGGSQTVVTQEP SLTVSPGGTVTLT CGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGKGGGSGGGSGGGSGGGSGGGG GGSGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
71.	PM_2xCD3-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPAGESLRSLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDAGGLTYYSDI IKGRFTISRDNANKNSLYLQINSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLVTVSSGGGSGGGSGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGG GGSGGGSGGGGSQTVVTQEP SLTVSPGGTVTLT CGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGGGGSGGGSGGGSGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL

			HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNNHYTQKSLSLSPGK
72.	PM_2_CCxCD3-scFc	VH	QVQLVESGGGLVLPKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKCLEWVAIIISDAGGLTYYSIDIIGKRFTISRDNANKSLYLQINSLKAEDTAVYYCARGFPLLRHGAMDYWGQGTLLVTVSS
73.	PM_2_CCxCD3-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVDNTNVAWYQQKPGQAPKSLIYSASYRYSADVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDSYPYTFGCGTKLEIK
74.	PM_2_CCxCD3-scFc	scFv	QVQLVESGGGLVLPKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKCLEWVAIIISDAGGLTYYSIDIIGKRFTISRDNANKSLYLQINSLKAEDTAVYYCARGFPLLRHGAMDYWGQGTLLVTVSSGGGGSGGGSGGGSDIQMTQSPSSLSASVGDRTITCKASQNVDNTNVAWYQQKPGQAPKSLIYSASYRYSADVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDSYPYTFGCGTKLEIK
75.	PM_2_CCxCD3-scFc	bispecific molecule	QVQLVESGGGLVLPKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKCLEWVAIIISDAGGLTYYSIDIIGKRFTISRDNANKSLYLQINSLKAEDTAVYYCARGFPLLRHGAMDYWGQGTLLVTVSSGGGGSGGGSGGGSDIQMTQSPSSLSASVGDRTITCKASQNVDNTNVAWYQQKPGQAPKSLIYSASYRYSADVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDSYPYTFGCGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGSGGGSGQTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGKTF LAPGTPARFSGSLLGGKAALTL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
76.	PM_2_CCxCD3-scFc	bispecific HLE molecule	QVQLVESGGGLVLPKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKCLEWVAIIISDAGGLTYYSIDIIGKRFTISRDNANKSLYLQINSLKAEDTAVYYCARGFPLLRHGAMDYWGQGTLLVTVSSGGGGSGGGSGGGSDIQMTQSPSSLSASVGDRTITCKASQNVDNTNVAWYQQKPGQAPKSLIYSASYRYSADVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDSYPYTFGCGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGSGGGSGQTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGKTF LAPGTPARFSGSLLGGKAALTL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGGSGGGSGDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNNHYTQKSLSLSPGK
77.	PM_2_CCxCD3-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVLPKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKCLEWVAIIISDAGGLTYYSIDIIGKRFTISRDNANKSLYLQINSLKAEDTAVYYCARGFPLLRHGAMDYWGQGTLLVTVSSGGGGSGGGSGGGSDIQMTQSPSSLSASVGDRTITCKASQNVDNTNVAWYQQKPGQAPKSLIYSASYRYSADVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDSYPYTFGCGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGSGGGSGQTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGKTF LAPGTPARFSGSLLGGKAALTL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNNHYTQKSLSLSPGK

			SCSVMHEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK
78.	PM_3xCD3-scFc	VH CDR1	DYYMY
79.	PM_3xCD3-scFc	VH CDR2	IISDAGGLTYYSDI IKG
80.	PM_3xCD3-scFc	VH CDR3	GFPLLRHGAMDY
81.	PM_3xCD3-scFc	VL CDR1	KASQNVDTNVA
82.	PM_3xCD3-scFc	VL CDR2	SASYRYS
83.	PM_3xCD3-scFc	VL CDR3	QQYDDL RPT
84.	PM_3xCD3-scFc	VH	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIISDAGGLTYYSDI IKG RFTISRDNAKNSLYLQINSLKAEDTAVY YCARGFPLLRHGAMDYWGQGT LTVTVSS
85.	PM_3xCD3-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSL IYSASYRYS DVP SRFSGSASGTDFTLTISSLQSEDFATYYCQQYDDL RPTFGGGTKLEIK
86.	PM_3xCD3-scFc	scFv	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIISDAGGLTYYSDI IKG RFTISRDNAKNSLYLQINSLKAEDTAVY YCARGFPLLRHGAMDYWGQGT LTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLTISSLQSEDFATYYCQQYDDL RPTFG GGTKLEIK
87.	PM_3xCD3-scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIISDAGGLTYYSDI IKG RFTISRDNAKNSLYLQINSLKAEDTAVY YCARGFPLLRHGAMDYWGQGT LTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLTISSLQSEDFATYYCQQYDDL RPTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LTVTVSSGG GGSGGGGSGGGGSGTQVTVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
88.	PM_3xCD3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIISDAGGLTYYSDI IKG RFTISRDNAKNSLYLQINSLKAEDTAVY YCARGFPLLRHGAMDYWGQGT LTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLTISSLQSEDFATYYCQQYDDL RPTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LTVTVSSGG GGSGGGGSGGGGSGTQVTVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDG VEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNV SCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGG GGGGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS

			REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK
89.	PM_3xCD3-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKCLEWVAIIISDAGGLTYYSDDIIKGRFTISRDNANKNSLYLQINSLKAEDTAVYYCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSASRYSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDDLRTFGGGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLTSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK
90.	PM_3_CCxCD 3-scFc	VH	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKCLEWVAIIISDAGGLTYYSDDIIKGRFTISRDNANKNSLYLQINSLKAEDTAVYYCARGFPLLRHGAMDYWGQGTLLTVTVSS
91.	PM_3_CCxCD 3-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSASRYSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDDLRTFGCGTKLEIK
92.	PM_3_CCxCD 3-scFc	scFv	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKCLEWVAIIISDAGGLTYYSDDIIKGRFTISRDNANKNSLYLQINSLKAEDTAVYYCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSASRYSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDDLRTFGCGTKLEIK
93.	PM_3_CCxCD 3-scFc	bispecific molecule	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKCLEWVAIIISDAGGLTYYSDDIIKGRFTISRDNANKNSLYLQINSLKAEDTAVYYCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSASRYSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDDLRTFGCGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLTSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
94.	PM_3_CCxCD 3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKCLEWVAIIISDAGGLTYYSDDIIKGRFTISRDNANKNSLYLQINSLKAEDTAVYYCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSASRYSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDDLRTFGCGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLTSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGG

			GSGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
95.	PM_3_CCxCD3-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVQKPGESLRLSAASGFTFSDYYMYWVRQAPGKCLEWVAIIISDAGGLTYYSDDIIKGRFTISRDNANKNSLYLQINSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLTISSLQSEDFATYYCQQYDDL RPTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
96.	PM_4xCD3-scFc	VH CDR1	DYYMY
97.	PM_4xCD3-scFc	VH CDR2	II SDHYSHTYYSDIIKD
98.	PM_4xCD3-scFc	VH CDR3	GFPLLRHGAMDY
99.	PM_4xCD3-scFc	VL CDR1	KASQNVD TNVA
100.	PM_4xCD3-scFc	VL CDR2	SASYRYS
101.	PM_4xCD3-scFc	VL CDR3	QQYDSYPYT
102.	PM_4xCD3-scFc	VH	QVQLVESGGGLVQKPGESLRLSAASGFTFSDYYMYWVRQAPGKGLEWVAIIISDHYSHTYYSDIIKDRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSS
103.	PM_4xCD3-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSASYRYS DVP SRFSGSASGTDFTLTISSVQSEDFATYYCQQYDSYPYTFGGG TKLEIK
104.	PM_4xCD3-scFc	scFv	QVQLVESGGGLVQKPGESLRLSAASGFTFSDYYMYWVRQAPGKGLEWVAIIISDHYSHTYYSDIIKDRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLTISSVQSEDFATYYCQQYDSYPYTFGGG TKLEIK
105.	PM_4xCD3-scFc	bispecific molecule	QVQLVESGGGLVQKPGESLRLSAASGFTFSDYYMYWVRQAPGKGLEWVAIIISDHYSHTYYSDIIKDRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLTISSVQSEDFATYYCQQYDSYPYTFGGG TKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ

			PEDEAEYYCVLWYSNRWVFGGGTKLTVL
106.	PM_4xCD3-scFc	bispecific HLE molecule	<p>QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAII SDHYSHTYYSDIIKDRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISVVQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSQTIVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK</p>
107.	PM_4xCD3-scFc_delGK	bispecific HLE molecule	<p>QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAII SDHYSHTYYSDIIKDRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISVVQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSQTIVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK</p>
108.	PM_4_CCxCD3-scFc	VH	<p>QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAII SDHYSHTYYSDIIKDRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSS</p>
109.	PM_4_CCxCD3-scFc	VL	<p>DIQMTQSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSL IYSASYRYSDVPSRFSGSASGTDFTLTISVVQSEDFATYYCQQYDSY PYTFGCGTKLEIK</p>
110.	PM_4_CCxCD3-scFc	scFv	<p>QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAII SDHYSHTYYSDIIKDRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISVVQSEDFATYYCQQYDSYPYTFG CGTKLEIK</p>
111.	PM_4_CCxCD3-scFc	bispecific molecule	<p>QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAII SDHYSHTYYSDIIKDRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISVVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT</p>

			AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
112.	PM_4_CCxCD 3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIISDHYSHTYYSDIIKDRFTISRDNANSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLTISVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGKGGGSGGGGSGGGGSGGGGSGGG GGSGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
113.	PM_4_CCxCD 3-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIISDHYSHTYYSDIIKDRFTISRDNANSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLTISVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
114.	PM_5xCD3- scFc	VH CDR1	DYYMY
115.	PM_5xCD3- scFc	VH CDR2	IISDKRHHTYYSDIIKG
116.	PM_5xCD3- scFc	VH CDR3	GFPLLRHGAMDY
117.	PM_5xCD3- scFc	VL CDR1	KASQNVDTNVA
118.	PM_5xCD3- scFc	VL CDR2	SASYRYS
119.	PM_5xCD3- scFc	VL CDR3	QQYDSYPYT
120.	PM_5xCD3-	VH	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIISDKRHHTYYSDIIKGRFTISRDNANSLYLQMNSLKAEDTAVY

			FFLYSKLTVDKSRWQQGNVFSOSVMHEALHNHYTQKSLSLSPGK
126.	PM_5_CCxCD 3-scFc	VH	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIIISDKRHHTYYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSS
127.	PM_5_CCxCD 3-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVDNTNAWYQQKPGQAPKSL IYSASYRSDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDSY PYTFGCGTKLEIK
128.	PM_5_CCxCD 3-scFc	scFv	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIIISDKRHHTYYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDNTNAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDSYPYTFG CGTKLEIK
129.	PM_5_CCxCD 3-scFc	bispecific molecule	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIIISDKRHHTYYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDNTNAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSGTQVTVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
130.	PM_5_CCxCD 3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIIISDKRHHTYYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDNTNAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSGTQVTVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGGG SGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD GSFFLYSKLTVDKSRWQQGNVFSOSVMHEALHNHYTQKSLSLSPGK
131.	PM_5_CCxCD 3-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIIISDKRHHTYYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDNTNAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSGTQVTVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGGGGSGGGSGGGSGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV

			VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK
132.	PM_6xCD3-scFc	VH CDR1	DYYMY
133.	PM_6xCD3-scFc	VH CDR2	IISDKRHHTYYSDI IKG
134.	PM_6xCD3-scFc	VH CDR3	GFPLLRHGAMDY
135.	PM_6xCD3-scFc	VL CDR1	KASQNVD TNVA
136.	PM_6xCD3-scFc	VL CDR2	SASYRYS
137.	PM_6xCD3-scFc	VL CDR3	QQYDDL RPT
138.	PM_6xCD3-scFc	VH	QVQLVESGGGLVKPGESLR LSCAASGFTTFSDYYMYWVRQAPGKGLEWVAIISDKRHHTYYSDI IKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSS
139.	PM_6xCD3-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSASYRSDVPSRFSGSASGTDFTLTIS SLQSEDFATYYCQQYDDL RPTFGGGTKLEIK
140.	PM_6xCD3-scFc	scFv	QVQLVESGGGLVKPGESLR LSCAASGFTTFSDYYMYWVRQAPGKGLEWVAIISDKRHHTYYSDI IKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGSGGGSGGGSDIQMTQSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSASYRSDVPSRFSGSASGTDFTLTIS SLQSEDFATYYCQQYDDL RPTFGGGTKLEIK
141.	PM_6xCD3-scFc	bispecific molecule	QVQLVESGGGLVKPGESLR LSCAASGFTTFSDYYMYWVRQAPGKGLEWVAIISDKRHHTYYSDI IKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGSGGGSGGGSDIQMTQSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSASYRSDVPSRFSGSASGTDFTLTIS SLQSEDFATYYCQQYDDL RPTFGGGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGSGGGSGGGSGTQVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
142.	PM_6xCD3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLR LSCAASGFTTFSDYYMYWVRQAPGKGLEWVAIISDKRHHTYYSDI IKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGSGGGSGGGSDIQMTQSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSASYRSDVPSRFSGSASGTDFTLTIS SLQSEDFATYYCQQYDDL RPTFGGGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGSGGGSGGGSGTQVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSGFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGKGGGSGGGSGGGSGGGSGGGSGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSGFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK

143.	PM_6xCD3-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDKRHHTYYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDDLRLPTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLGKKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
144.	PM_6_CCxCD3-scFc	VH	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIIISDKRHHTYYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSS
145.	PM_6_CCxCD3-scFc	VL	DIQMTQSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSL IYSASYRYSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDDL RPTFGCGTKLEIK
146.	PM_6_CCxCD3-scFc	scFv	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIIISDKRHHTYYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDDLRLPTFG CGTKLEIK
147.	PM_6_CCxCD3-scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIIISDKRHHTYYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDDLRLPTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLGKKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
148.	PM_6_CCxCD3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIIISDKRHHTYYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDDLRLPTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLGKKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGG GGGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT

			VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD GSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
149.	PM_6_CCxCD 3-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPAGESLRLSAASGFTTFSDYYMYWVRQAPGKCLEW VAIISDKRHHTYYSDIIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVITITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDDLRTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNHLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNV SCSVMHEALHNHYTQKSLSLSPGGGGSGGGSGGGSGGGSGGGG GGGGSDKHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGS FFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
150.	PM_7xCD3- scFc	VH CDR1	DYYMY
151.	PM_7xCD3- scFc	VH CDR2	IISDGGYYTYSDIIKG
152.	PM_7xCD3- scFc	VH CDR3	GFPLLRHGAMDY
153.	PM_7xCD3- scFc	VL CDR1	KASQNVDTNVA
154.	PM_7xCD3- scFc	VL CDR2	SASYRYS
155.	PM_7xCD3- scFc	VL CDR3	QQYDQQLIT
156.	PM_7xCD3- scFc	VH	QVQLVESGGGLVKPAGESLRLSAASGFTTFSDYYMYWVRQAPGKGLEW VAIISDGGYYTYSDIIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSS
157.	PM_7xCD3- scFc	VL	DIQMTQSPSSLSASVGDRVITITCKASQNVDTNVAWYQQKPGQAPKSL IYSASYRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLIT FTGGGKLEIK
158.	PM_7xCD3- scFc	scFv	QVQLVESGGGLVKPAGESLRLSAASGFTTFSDYYMYWVRQAPGKGLEW VAIISDGGYYTYSDIIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVITITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFTG GGTKLEIK
159.	PM_7xCD3- scFc	bispecific molecule	QVQLVESGGGLVKPAGESLRLSAASGFTTFSDYYMYWVRQAPGKGLEW VAIISDGGYYTYSDIIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVITITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFTG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNHLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
160.	PM_7xCD3-	bispecific	QVQLVESGGGLVKPAGESLRLSAASGFTTFSDYYMYWVRQAPGKGLEW

	scFc	HLE molecule	VAIIISDGGYYTYYSDDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISSVQSEDFATYYCQQYDQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGSGGGGSGTQVTVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGG GGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
161.	PM_7xCD3- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSDDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISSVQSEDFATYYCQQYDQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGSGGGGSGTQVTVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCTV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
162.	PM_7_CCxCD 3-scFc	VH	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKCLEW VAIIISDGGYYTYYSDDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSS
163.	PM_7_CCxCD 3-scFc	VL	DIQMTQSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSL IYSASYRYSDVPSRFSGSASGTDFTLTISSVQSEDFATYYCQQYDQQL LITFGCGTKLEIK
164.	PM_7_CCxCD 3-scFc	scFv	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKCLEW VAIIISDGGYYTYYSDDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISSVQSEDFATYYCQQYDQQLITFG CGTKLEIK
165.	PM_7_CCxCD 3-scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKCLEW VAIIISDGGYYTYYSDDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISSVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGSGGGGSGTQVTVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY

			PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
166.	PM_7_CCxCD 3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYSIDIIGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQTLVTVSSGG GGSGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGGG GGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
167.	PM_7_CCxCD 3-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYSIDIIGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQTLVTVSSGG GGSGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGGGGSGGGSGGGSGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
168.	PM_8xCD3- scFc	VH CDR1	DYYMY
169.	PM_8xCD3- scFc	VH CDR2	IISDGGYYTYYSIDIIGK
170.	PM_8xCD3- scFc	VH CDR3	GFPLLRHGAMDY
171.	PM_8xCD3- scFc	VL CDR1	KASQNVDTNVA
172.	PM_8xCD3- scFc	VL CDR2	SASYRYS
173.	PM_8xCD3- scFc	VL CDR3	QQYDQVRLT
174.	PM_8xCD3- scFc	VH	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKGLEW VAIISDGGYYTYYSIDIIGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLVTVSS

175.	PM_8xCD3-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSASYRSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDQVRLTFGGGKLEIK
176.	PM_8xCD3-scFc	scFv	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKGLEWVAIISDGGYYTYSDIIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVVSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDQVRLTFG GGTLEIK
177.	PM_8xCD3-scFc	bispecific molecule	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKGLEWVAIISDGGYYTYSDIIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVVSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDQVRLTFG GGTLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVVSSGG GSGGGGSGGGGSGTVVTVQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
178.	PM_8xCD3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKGLEWVAIISDGGYYTYSDIIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVVSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDQVRLTFG GGTLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVVSSGG GSGGGGSGGGGSGTVVTVQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNV FSCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGG GSGGGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNV FSCSVMEALHNHYTQKSLSLSPGK
179.	PM_8xCD3-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKGLEWVAIISDGGYYTYSDIIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVVSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDQVRLTFG GGTLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVVSSGG GSGGGGSGGGGSGTVVTVQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNV FSCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGG GGGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNV FSCSVMEALHNHYTQKSLSLSPGK

180.	PM_8_CCxCD 3-scFc	VH	QVQLVESGGGLVKPAGESLRSLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSS
181.	PM_8_CCxCD 3-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVDNTNVAWYQQKPGQAPKSL IYSASYRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQV RLTFGCGTKLEIK
182.	PM_8_CCxCD 3-scFc	scFv	QVQLVESGGGLVKPAGESLRSLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDNTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQVRLTFG CGTKLEIK
183.	PM_8_CCxCD 3-scFc	bispecific molecule	QVQLVESGGGLVKPAGESLRSLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDNTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQVRLTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
184.	PM_8_CCxCD 3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPAGESLRSLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDNTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQVRLTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGGG GGGGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
185.	PM_8_CCxCD 3-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPAGESLRSLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDNTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQVRLTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGGGGSGGGSGGGSGGGSGGGG GGGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL

			HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHYTEQKSLSLSPGK
186.	PM_9xCD3- scFc	VH CDR1	DYYMN
187.	PM_9xCD3- scFc	VH CDR2	IISDGGYYTYYSDIKIG
188.	PM_9xCD3- scFc	VH CDR3	GFPLLRHGAMDY
189.	PM_9xCD3- scFc	VL CDR1	KASQNVDTNVA
190.	PM_9xCD3- scFc	VL CDR2	SASYRYS
191.	PM_9xCD3- scFc	VL CDR3	QQYDDL RPT
192.	PM_9xCD3- scFc	VH	QVQLVESGGGLVKPGESLRLSAASGFTTFSDYYMNWVRQAPGKGLEW VAIISDGGYYTYYSDIKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSS
193.	PM_9xCD3- scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSL IYSASYRYSVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDDL RPTFGGGTKLEIK
194.	PM_9xCD3- scFc	scFv	QVQLVESGGGLVKPGESLRLSAASGFTTFSDYYMNWVRQAPGKGLEW VAIISDGGYYTYYSDIKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDDL RPTFG GGTKLEIK
195.	PM_9xCD3- scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSAASGFTTFSDYYMNWVRQAPGKGLEW VAIISDGGYYTYYSDIKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDDL RPTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
196.	PM_9xCD3- scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSAASGFTTFSDYYMNWVRQAPGKGLEW VAIISDGGYYTYYSDIKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDDL RPTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFC SCVMHEALHNNHYTEQKSLSLSPGKGGGGSGGGSGGGSGGGSGGG GSGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHYTEQKSLSLSPGK
197.	PM_9xCD3-	bispecific	QVQLVESGGGLVKPGESLRLSAASGFTTFSDYYMNWVRQAPGKGLEW

	scFc_delGK	HLE molecule	VAIIISDGGYYTYYSDI I KGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLT ISSLQSEDFATYYCQQYDDL RPTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGSGGGGSGTVVTVQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTCPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKHTCPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
198.	PM_9_CCxCD 3-scFc	VH	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMNWVRQAPGKCLEW VAIIISDGGYYTYYSDI I KGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSS
199.	PM_9_CCxCD 3-scFc	VL	DIQMTQSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSL IYSASYRYSDVPSRFSGSASGTDFTLT ISSLQSEDFATYYCQQYDDL RPTFGCGTKLEIK
200.	PM_9_CCxCD 3-scFc	scFv	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMNWVRQAPGKCLEW VAIIISDGGYYTYYSDI I KGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLT ISSLQSEDFATYYCQQYDDL RPTFG CGTKLEIK
201.	PM_9_CCxCD 3-scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMNWVRQAPGKCLEW VAIIISDGGYYTYYSDI I KGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLT ISSLQSEDFATYYCQQYDDL RPTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGSGGGGSGTVVTVQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
202.	PM_9_CCxCD 3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMNWVRQAPGKCLEW VAIIISDGGYYTYYSDI I KGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLT ISSLQSEDFATYYCQQYDDL RPTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGSGGGGSGTVVTVQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTCPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGG GGGGSDKHTCPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS

			REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK
203.	PM_9_CCxCD3-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKCLEWVAIISDGGYYTYYSDI IKGRFTISRDNANKNSLYLQMNSLKAEDTAVYYCARGFPLL RHGAMDYWGQGTLVTVSSGGGGSGGGSGGGSGDIQMTQSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSLQSEDFATYYCQQYDDL RPTFGCGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLVTVSSGGGGSGGGSGGGSGQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGGGGSGGGSGGGSGGGSGGGSGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK
204.	PM_10xCD3-scFc	VH CDR1	DYYMY
205.	PM_10xCD3-scFc	VH CDR2	IISDH HHWTYYSDI IKG
206.	PM_10xCD3-scFc	VH CDR3	GFPLL RHGAMDY
207.	PM_10xCD3-scFc	VL CDR1	KASQNVD TNVA
208.	PM_10xCD3-scFc	VL CDR2	SASYRYS
209.	PM_10xCD3-scFc	VL CDR3	QQYDSYPYT
210.	PM_10xCD3-scFc	VH	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKGLEWVAIISDH HHWTYYSDI IKGRFTISRDNANKNSLYLQMNSLKAEDTAVYYCARGFPLL RHGAMDYWGQGTLVTVSS
211.	PM_10xCD3-scFc	VL	DIQMTQSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSASYRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDSYPYTFGGG TKLEIK
212.	PM_10xCD3-scFc	scFv	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKGLEWVAIISDH HHWTYYSDI IKGRFTISRDNANKNSLYLQMNSLKAEDTAVYYCARGFPLL RHGAMDYWGQGTLVTVSSGGGGSGGGSGGGSGDIQMTQSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDSYPYTFGGG TKLEIK
213.	PM_10xCD3-scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKGLEWVAIISDH HHWTYYSDI IKGRFTISRDNANKNSLYLQMNSLKAEDTAVYYCARGFPLL RHGAMDYWGQGTLVTVSSGGGGSGGGSGGGSGDIQMTQSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDSYPYTFGGG TKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLVTVSSGGGGSGGGSGGGSGQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
214.	PM_10xCD3-	bispecific	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKGLEWVAIISDH HHWTYYSDI IKGRFTISRDNANKNSLYLQMNSLKAEDTAVY

	scFc	HLE molecule	YCARGFPLL RHGAMDYWGQGT LVT VSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQP GGS LKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVT VSSGG GGSGGGGSGGGGSQT VVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMH EALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGG GGSGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMH EALHNHYTQKSLSLSPGK
215.	PM_10xCD3- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLR LSCAASGFTFSDYYMYWVRQAPGKGLEW VAIISDHHHWTYYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGT LVT VSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQP GGS LKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVT VSSGG GGSGGGGSGGGGSQT VVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMH EALHNHYTQKSLSLSPGSGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMH EALHNHYTQKSLSLSPGK
216.	PM_10_CCxC D3-scFc	VH	QVQLVESGGGLVKPGESLR LSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDHHHWTYYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGT LVT VSS
217.	PM_10_CCxC D3-scFc	VL	DIQMTQSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSL IYSASYRYSDVPSRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDSY PYTFGCGTKLEIK
218.	PM_10_CCxC D3-scFc	scFv	QVQLVESGGGLVKPGESLR LSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDHHHWTYYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGT LVT VSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDSYPYTFG CGTKLEIK
219.	PM_10_CCxC D3-scFc	bispecific molecule	QVQLVESGGGLVKPGESLR LSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDHHHWTYYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGT LVT VSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQP GGS LKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVT VSSGG GGSGGGGSGGGGSQT VVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ

			PEDEAEYYCVLWYSNRWVFGGGTKLTVL
220.	PM_10_CCxC D3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGEISLRLSAASGFTTFSDYYMYWVRQAPGKCLEW VAIIISDHHHWTYYSDIIKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRLHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSQTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGG GGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
221.	PM_10_CCxC D3- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGEISLRLSAASGFTTFSDYYMYWVRQAPGKCLEW VAIIISDHHHWTYYSDIIKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRLHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSQTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
222.	PM_11xCD3- scFc	VH CDR1	DYYMY
223.	PM_11xCD3- scFc	VH CDR2	IISDGGYYTYYSDDIKG
224.	PM_11xCD3- scFc	VH CDR3	GFPLLRLHGAMDY
225.	PM_11xCD3- scFc	VL CDR1	KASQNVDTNVA
226.	PM_11xCD3- scFc	VL CDR2	SASYRYS
227.	PM_11xCD3- scFc	VL CDR3	QQYDRQPLT
228.	PM_11xCD3- scFc	VH	QVQLVESGGGLVKPGEISLRLSAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSDDIKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRLHGAMDYWGQGTLLTVTVSS
229.	PM_11xCD3-	VL	DIQMTQSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSL IYSASYRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDRQ

	scFc		PLTFGGGKLEIK
230.	PM_11xCD3-scFc	scFv	QVQLVESGGGLVKPGEISRLSCAASGFTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSDI IKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDRQPLTFG GGTKLEIK
231.	PM_11xCD3-scFc	bispecific molecule	QVQLVESGGGLVKPGEISRLSCAASGFTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSDI IKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDRQPLTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNVWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
232.	PM_11xCD3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGEISRLSCAASGFTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSDI IKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDRQPLTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNVWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGYSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHAEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGYSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMHAEALHNHYTQKSLSLSPGK
233.	PM_11xCD3-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGEISRLSCAASGFTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSDI IKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDRQPLTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNVWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGYSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHAEALHNHYTQKSLSLSPGSGGGSGGGSGGGSGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGYSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMHAEALHNHYTQKSLSLSPGK
234.	PM_11_CcxC D3-scFc	VH	QVQLVESGGGLVKPGEISRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIIISDGGYYTYYSDI IKGRFTISRDNAKNSLYLQMNSLKAEDTAVY

			YCARGFPLLRHGAMDYWGQGTLLTVSS
235.	PM_11_CCxC D3-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSL IYSASYRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDRQ PLTFGCGTKLEIK
236.	PM_11_CCxC D3-scFc	scFv	QVQLVESGGGLVQPGESLRLSAASGFTTFSDYYMYWVRQAPGKCLEW VAIIISDGGYYTYSDIIKGRFTISRDNANKSLYLQMNLSKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVSSGGGGSGGGSGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDRQPLTFG CGTKLEIK
237.	PM_11_CCxC D3-scFc	bispecific molecule	QVQLVESGGGLVQPGESLRLSAASGFTTFSDYYMYWVRQAPGKCLEW VAIIISDGGYYTYSDIIKGRFTISRDNANKSLYLQMNLSKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVSSGGGGSGGGSGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDRQPLTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNLLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVSSGG GGSGGGSGGGGSGTQVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
238.	PM_11_CCxC D3-scFc	bispecific HLE molecule	QVQLVESGGGLVQPGESLRLSAASGFTTFSDYYMYWVRQAPGKCLEW VAIIISDGGYYTYSDIIKGRFTISRDNANKSLYLQMNLSKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVSSGGGGSGGGSGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDRQPLTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNLLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVSSGG GGSGGGSGGGGSGTQVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNV SCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGG GGSGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEV TCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVL TVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
239.	PM_11_CCxC D3- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVQPGESLRLSAASGFTTFSDYYMYWVRQAPGKCLEW VAIIISDGGYYTYSDIIKGRFTISRDNANKSLYLQMNLSKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVSSGGGGSGGGSGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDRQPLTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNLLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVSSGG GGSGGGSGGGGSGTQVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNV SCSVMHEALHNHYTQKSLSLSPGGGGSGGGSGGGSGGGSGGGSG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS

			FFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
240.	PM_12xCD3-scFc	VH CDR1	DYYMN
241.	PM_12xCD3-scFc	VH CDR2	IISDGGYYTYYSDI IKG
242.	PM_12xCD3-scFc	VH CDR3	GFPLLRHGAMDY
243.	PM_12xCD3-scFc	VL CDR1	KASQNVD TNVA
244.	PM_12xCD3-scFc	VL CDR2	SASYRYS
245.	PM_12xCD3-scFc	VL CDR3	QQYDLLLLT
246.	PM_12xCD3-scFc	VH	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKGLEW VAIISDGGYYTYYSDI IKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLVTVSS
247.	PM_12xCD3-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSL IYSASYRSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDLL LLTFGGGKLEIK
248.	PM_12xCD3-scFc	scFv	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKGLEW VAIISDGGYYTYYSDI IKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLVTVSSGGGSGGGSGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDLLLLTFG GGTKLEIK
249.	PM_12xCD3-scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKGLEW VAIISDGGYYTYYSDI IKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLVTVSSGGGSGGGSGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDLLLLTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTVSPGGTVTLTCSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
250.	PM_12xCD3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKGLEW VAIISDGGYYTYYSDI IKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLVTVSSGGGSGGGSGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDLLLLTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTVSPGGTVTLTCSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGSGGGSGGGSGGGSGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
251.	PM_12xCD3-scFc_delGK	bispecific HLE	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKGLEW VAIISDGGYYTYYSDI IKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLVTVSSGGGSGGGSGGGSDIQMT

		molecule	QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSSQSEDFATYYCQQYDLLLFTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
252.	PM_12_CCxC D3-scFc	VH	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMNWVRQAPGKCLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSS
253.	PM_12_CCxC D3-scFc	VL	DIQMTQSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSL IYSASYRYSDVPSRFSGSASGTDFTLTITSSSQSEDFATYYCQQYDLL LLTFGCGTKLEIK
254.	PM_12_CCxC D3-scFc	scFv	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMNWVRQAPGKCLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSSQSEDFATYYCQQYDLLLFTFG CGTKLEIK
255.	PM_12_CCxC D3-scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMNWVRQAPGKCLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSSQSEDFATYYCQQYDLLLFTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
256.	PM_12_CCxC D3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMNWVRQAPGKCLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSSQSEDFATYYCQQYDLLLFTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

257.	PM_12_CCxC D3- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSAASGFTTFSDYYMNWVRQAPGKCLEW VAIIISDGGYYTYYSDDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDLLLLTFTG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCIVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNV SCSVMHEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
258.	PM_13xCD3- scFc	VH CDR1	DYYMY
259.	PM_13xCD3- scFc	VH CDR2	IISDSTLHTYYSDIIKG
260.	PM_13xCD3- scFc	VH CDR3	GFPLLRHGAMDY
261.	PM_13xCD3- scFc	VL CDR1	KASQNVDTNVA
262.	PM_13xCD3- scFc	VL CDR2	SASYRYS
263.	PM_13xCD3- scFc	VL CDR3	QQYDSYPYT
264.	PM_13xCD3- scFc	VH	QVQLVESGGGLVKPGESLRLSAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDSTLHTYYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSS
265.	PM_13xCD3- scFc	VL	DIQMTQSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSL IYSASYRYSVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDSY PYTFGGGKLEIK
266.	PM_13xCD3- scFc	scFv	QVQLVESGGGLVKPGESLRLSAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDSTLHTYYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDSYPYTFG GGTKLEIK
267.	PM_13xCD3- scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDSTLHTYYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCIVLWYSNRWVFGGGTKLTVL
268.	PM_13xCD3- scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDSTLHTYYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS

			YRYS DVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTVQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGG GGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
269.	PM_13xCD3- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKGLEW VAIIISDSTLHTYYSDIIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYS DVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTVQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGSGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
270.	PM_13_CCxC D3-scFc	VH	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKCLEW VAIIISDSTLHTYYSDIIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSS
271.	PM_13_CCxC D3-scFc	VL	DIQMTQSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSL IYSASYRYS DVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDSY PYTFGCGTKLEIK
272.	PM_13_CCxC D3-scFc	scFv	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKCLEW VAIIISDSTLHTYYSDIIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYS DVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDSYPYTFG CGTKLEIK
273.	PM_13_CCxC D3-scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKCLEW VAIIISDSTLHTYYSDIIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYS DVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTVQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
274.	PM_13_CCxC	bispecific	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKCLEW

	D3-scFc	HLE molecule	VAIIISDSTLHTYYSDIIKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRLHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTIISSVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGG GGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
275.	PM_13_CcxC D3- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKCLEW VAIIISDSTLHTYYSDIIKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRLHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTIISSVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCTV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
276.	PM_14xCD3- scFc	VH CDR1	DYYMY
277.	PM_14xCD3- scFc	VH CDR2	IISDGGYYTYSDIIKG
278.	PM_14xCD3- scFc	VH CDR3	GFPLLRLHGAMDY
279.	PM_14xCD3- scFc	VL CDR1	KASQNVDTNVA
280.	PM_14xCD3- scFc	VL CDR2	SASYRYS
281.	PM_14xCD3- scFc	VL CDR3	QQYDAPMGT
282.	PM_14xCD3- scFc	VH	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYSDIIKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRLHGAMDYWGQGTLLTVTVSS
283.	PM_14xCD3- scFc	VL	DIQMTQSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSL IYSASYRYSDVPSRFSGSASGTDFTLTIISSVQSEDFATYYCQQYDAP MGTFGGGKLEIK

284.	PM_14xCD3-scFc	scFv	QVQLVESGGGLVKPGEISRLRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSIDIIGKRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDAPMGTFG GGTKLEIK
285.	PM_14xCD3-scFc	bispecific molecule	QVQLVESGGGLVKPGEISRLRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSIDIIGKRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDAPMGTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
286.	PM_14xCD3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGEISRLRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSIDIIGKRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDAPMGTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGG GGSGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
287.	PM_14xCD3-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGEISRLRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSIDIIGKRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDAPMGTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GSGGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGSGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
288.	PM_14_CCxC D3-scFc	VH	QVQLVESGGGLVKPGEISRLRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIIISDGGYYTYYSIDIIGKRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSS

289.	PM_14_CCxC D3-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSL IYSASYRSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDAP MGTFGCGTKLEIK
290.	PM_14_CCxC D3-scFc	scFv	QVQLVESGGGLVKPGE SLRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAII SDGGYYTYYS DI I I KGRFTISRDN AKNSLYLQMN SLKAEDTAVY YCARGFPLLRHGAMDYWGQGT LVT VSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDAPMGTFG CGTKLEIK
291.	PM_14_CCxC D3-scFc	bispecific molecule	QVQLVESGGGLVKPGE SLRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAII SDGGYYTYYS DI I I KGRFTISRDN AKNSLYLQMN SLKAEDTAVY YCARGFPLLRHGAMDYWGQGT LVT VSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDAPMGTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVT VSSGG GGSGGGSGGGGSGT VVTQEP SLTVSPGGTVTLT CGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYC VLVWYSNRWVFGGGTKLTVL
292.	PM_14_CCxC D3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGE SLRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAII SDGGYYTYYS DI I I KGRFTISRDN AKNSLYLQMN SLKAEDTAVY YCARGFPLLRHGAMDYWGQGT LVT VSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDAPMGTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVT VSSGG GGSGGGSGGGGSGT VVTQEP SLTVSPGGTVTLT CGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYC VLVWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGG GGGGSGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFCSSVMHEALHNHYTQKSLSLSPGK
293.	PM_14_CCxC D3- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGE SLRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAII SDGGYYTYYS DI I I KGRFTISRDN AKNSLYLQMN SLKAEDTAVY YCARGFPLLRHGAMDYWGQGT LVT VSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDAPMGTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVT VSSGG GGSGGGSGGGGSGT VVTQEP SLTVSPGGTVTLT CGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYC VLVWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGGSGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFCSSVMHEALHNHYTQKSLSLSPGK

294.	PM_15xCD3-scFc	VH CDR1	DYYMN
295.	PM_15xCD3-scFc	VH CDR2	IISDNKHQTYYSDIKIG
296.	PM_15xCD3-scFc	VH CDR3	GFPLLRHGAMDY
297.	PM_15xCD3-scFc	VL CDR1	KASQNVDTNVA
298.	PM_15xCD3-scFc	VL CDR2	SASYRYS
299.	PM_15xCD3-scFc	VL CDR3	QQYDSYPYT
300.	PM_15xCD3-scFc	VH	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKGLEW VAIISDNKHQTYYSDIKGRFAISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLVTVSS
301.	PM_15xCD3-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLI YASARYSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDSY PYTFGGGTKLEIK
302.	PM_15xCD3-scFc	scFv	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKGLEW VAIISDNKHQTYYSDIKGRFAISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYAS YRYSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDSYPYTFG GGTKLEIK
303.	PM_15xCD3-scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKGLEW VAIISDNKHQTYYSDIKGRFAISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYAS YRYSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGG GGSGGGSGGGGSGTQVTVTQEPSTVSPGGTVTLTCSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
304.	PM_15xCD3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKGLEW VAIISDNKHQTYYSDIKGRFAISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYAS YRYSDVPSRFSGSASGTDFTLTISLQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGG GGSGGGSGGGGSGTQVTVTQEPSTVSPGGTVTLTCSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGGG GGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
305.	PM_15xCD3-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKGLEW VAIISDNKHQTYYSDIKGRFAISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYAS

			YRYSDVPSRFSGSASGTDFTLTITSSSQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
306.	PM_15_CCxC D3-scFc	VH	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKCLEW VAIIISDNKHQTYYSIDIIGRFAISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSS
307.	PM_15_CCxC D3-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSL IYSASYRYSDVPSRFSGSASGTDFTLTITSSSQSEDFATYYCQQYDSY PYTFGCGTKLEIK
308.	PM_15_CCxC D3-scFc	scFv	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKCLEW VAIIISDNKHQTYYSIDIIGRFAISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSSQSEDFATYYCQQYDSYPYTFG CGTKLEIK
309.	PM_15_CCxC D3-scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKCLEW VAIIISDNKHQTYYSIDIIGRFAISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSSQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
310.	PM_15_CCxC D3-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKCLEW VAIIISDNKHQTYYSIDIIGRFAISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSSQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGG GSGGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
311.	PM_15_CCxC	bispecific	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMNWVRQAPGKCLEW

	D3-scFc_delGK	HLE molecule	VAIIISDNKHQTYYSDDIIKGRFAISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRRHGAMDYWGQGTLLVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYS DVPSRFSGSASGTDFTLTISSVQSEDFATYYCQQYESYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNVWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGG GGSGGGGSGGGGSGTVVTVQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
312.	PM 76-B10.02 CC x I2C0-scFc	VH CDR1	DYYMY
313.	PM 76-B10.02 CC x I2C0-scFc	VH CDR2	IISDAGYYTYYSDDIIKG
314.	PM 76-B10.02 CC x I2C0-scFc	VH CDR3	GFPLLRRHGAMDY
315.	PM 76-B10.02 CC x I2C0-scFc	VL CDR1	KASQNVDTNVA
316.	PM 76-B10.02 CC x I2C0-scFc	VL CDR2	SASYRYS
317.	PM 76-B10.02 CC x I2C0-scFc	VL CDR3	QQYESYPYT
318.	PM 76-B10.02 CC x I2C0-scFc	VH	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIIISDAGYYTYYSDDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRRHGAMDYWGQGTLLVTVSS
319.	PM 76-B10.02 CC x I2C0-scFc	VL	DIQMTQSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSL IYSASYRYS DVPSRFSGSASGTDFTLTISSVQSEDFATYYCQQYESY PYTFGCGTKLEIK
320.	PM 76-B10.02 CC x I2C0-scFc	scFv	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIIISDAGYYTYYSDDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRRHGAMDYWGQGTLLVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYS DVPSRFSGSASGTDFTLTISSVQSEDFATYYCQQYESYPYTFG CGTKLEIK
321.	PM 76-B10.02 CC x I2C0-scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIIISDAGYYTYYSDDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRRHGAMDYWGQGTLLVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYS DVPSRFSGSASGTDFTLTISSVQSEDFATYYCQQYESYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNVWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGG

			GGSGGGGSGGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
322.	PM 76- B10.02 CC x I2C0-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGE SLRLSCAASGFTFSDYYMYWVRQAPGKC LEW VAIISDAGYYTYYS DI I KGRFTISRDN AKNSLYLQMN SLKAEDTAVY YCARGFPLL RHGAMDYWGQGLVTVSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGD RVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYESYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGG GGSGGGGSGGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSV MHEALHNHYTQKSLSLSPGKGGGSGGGGSGGGGSGGGGSGGG GGSGGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSV MHEALHNHYTQKSLSLSPGK
323.	PM 76- B10.02 CC x I2C0- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGE SLRLSCAASGFTFSDYYMYWVRQAPGKC LEW VAIISDAGYYTYYS DI I KGRFTISRDN AKNSLYLQMN SLKAEDTAVY YCARGFPLL RHGAMDYWGQGLVTVSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGD RVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYESYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGG GGSGGGGSGGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSV MHEALHNHYTQKSLSLSPGGGSGGGGSGGGGSGGGGSGGGGSG GGGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSV MHEALHNHYTQKSLSLSPGK
324.	PM 76- B10.02 CC x I2C0 CC (103/43)-scFc	bispecific molecule	QVQLVESGGGLVKPGE SLRLSCAASGFTFSDYYMYWVRQAPGKC LEW VAIISDAGYYTYYS DI I KGRFTISRDN AKNSLYLQMN SLKAEDTAVY YCARGFPLL RHGAMDYWGQGLVTVSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGD RVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYESYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLVTVSSGG GGSGGGGSGGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCP RGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
325.	PM 76- B10.02 CC x I2C0 CC (103/43)-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGE SLRLSCAASGFTFSDYYMYWVRQAPGKC LEW VAIISDAGYYTYYS DI I KGRFTISRDN AKNSLYLQMN SLKAEDTAVY YCARGFPLL RHGAMDYWGQGLVTVSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGD RVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYESYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA

			MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGTTLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTLVSPGGTTLTTCGSSTGAVTSGNY PNWVQQKPGQCPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGG GGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
326.	PM 76- B10.02 CC x I2C0 CC (103/43)- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDAGYYTYYSDIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTTLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSQVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYESYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGTTLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTLVSPGGTTLTTCGSSTGAVTSGNY PNWVQQKPGQCPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGGSG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
327.	PM 76- B10.08 x I2C0-scFc	VH CDR1	DYYMY
328.	PM 76- B10.08 x I2C0-scFc	VH CDR2	IISDGGYYTYYSDIKIG
329.	PM 76- B10.08 x I2C0-scFc	VH CDR3	GFPLLRHGAMDY
330.	PM 76- B10.08 x I2C0-scFc	VL CDR1	KASQNVDTNVA
331.	PM 76- B10.08 x I2C0-scFc	VL CDR2	SASYRYS
332.	PM 76- B10.08 x I2C0-scFc	VL CDR3	QQYDQQQLIT
333.	PM 76- B10.08 x I2C0-scFc	VH	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKGLEW VAIISDGGYYTYYSDIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTTLTVTVSS

334.	PM 76-B10.08 x I2C0-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSASYRSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDQQLITFGGGGKLEIK
335.	PM 76-B10.08 x I2C0-scFc	scFv	QVQLVESGGGLVQPGESLRLSAASGFTTFSDYYMYWVRQAPGKGLEWVAIISDGGYYTYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDQQLITFG GGTLEIK
336.	PM 76-B10.08 x I2C0-scFc	bispecific molecule	QVQLVESGGGLVQPGESLRLSAASGFTTFSDYYMYWVRQAPGKGLEWVAIISDGGYYTYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDQQLITFG GGTLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GSGGGGSGGGGSGTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
337.	PM 76-B10.08 x I2C0-scFc	bispecific HLE molecule	QVQLVESGGGLVQPGESLRLSAASGFTTFSDYYMYWVRQAPGKGLEWVAIISDGGYYTYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDQQLITFG GGTLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GSGGGGSGGGGSGTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGG GSGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
338.	PM 76-B10.08 x I2C0-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVQPGESLRLSAASGFTTFSDYYMYWVRQAPGKGLEWVAIISDGGYYTYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDQQLITFG GGTLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GSGGGGSGGGGSGTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGSGGGGSGGGGSGGGGSGGGGSGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK

339.	PM 76- B10.08 x I2C0 CC (103/43)- scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSDI IKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLVTVSSGG GGSGGGSGGGGSGTQVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCP RGLIGGTKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
340.	PM 76- B10.08 x I2C0 CC (103/43)- scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSDI IKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLVTVSSGG GGSGGGSGGGGSGTQVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCP RGLIGGTKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGG GGSGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
341.	PM 76- B10.08 x I2C0 CC (103/43)- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSDI IKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGLVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFLTITSSVQSEDFATYYCQQYDQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLVTVSSGG GGSGGGSGGGGSGTQVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCP RGLIGGTKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
342.	PM 76- B10.08 CC x I2C0-scFc	VH CDR1	DYYMY
343.	PM 76- B10.08 CC x I2C0-scFc	VH CDR2	IISDGGYYTYYSDI IKG
344.	PM 76- B10.08 CC x	VH CDR3	GFPLL RHGAMDY

	I2C0-scFc		
345.	PM 76-B10.08 CC x I2C0-scFc	VL CDR1	KASQNVD TNVA
346.	PM 76-B10.08 CC x I2C0-scFc	VL CDR2	SASYRYS
347.	PM 76-B10.08 CC x I2C0-scFc	VL CDR3	QQYDQQ LIT
348.	PM 76-B10.08 CC x I2C0-scFc	VH	QVQLVESGGGLVKPGESLR LSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYS DIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGT LVTVSS
349.	PM 76-B10.08 CC x I2C0-scFc	VL	DIQMTQSPSSLSASV GDRVTITCKASQNVD TNVAWYQQKPGQAPKSL IYSASYRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDQQ LITFGCGTKLEIK
350.	PM 76-B10.08 CC x I2C0-scFc	scFv	QVQLVESGGGLVKPGESLR LSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYS DIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGT LVTVSSGGGSGGGSGGGSGDIQMT QSPSSLSASV GDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDQQ LITFG CGTKLEIK
351.	PM 76-B10.08 CC x I2C0-scFc	bispecific molecule	QVQLVESGGGLVKPGESLR LSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYS DIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGT LVTVSSGGGSGGGSGGGSGDIQMT QSPSSLSASV GDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDQQ LITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVTVSSGG GGSGGGSGGGGSQT VVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYC VLVWSNRWVFGGGTKLTVL
352.	PM 76-B10.08 CC x I2C0-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLR LSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYS DIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGT LVTVSSGGGSGGGSGGGSGDIQMT QSPSSLSASV GDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDQQ LITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVTVSSGG GGSGGGSGGGGSQT VVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYC VLVWSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNV SCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGG GGSGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEV TCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVL TVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
353.	PM 76-B10.08 CC x I2C0-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLR LSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYS DIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGT LVTVSSGGGSGGGSGGGSGDIQMT QSPSSLSASV GDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDQQ LITFG

			CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVVSSGG GGSGGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGYSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGYSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
354.	PM 76- B10.08 CC x I2C0 CC (103/43)-scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYSDIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLVTVVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISSVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLVTVVSSGG GGSGGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
355.	PM 76- B10.08 CC x I2C0 CC (103/43)-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYSDIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLVTVVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISSVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLVTVVSSGG GGSGGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGYSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGYSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
356.	PM 76- B10.08 CC x I2C0 CC (103/43)- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYSDIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLVTVVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISSVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLVTVVSSGG GGSGGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGYSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD

			I AVEWESNGQPENNYKTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFS SCSVMEALHNHYTQKSLSLSPGGGGSGGGSGGGSGGGSGGGSG GGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
357.	PM 76- B10.11 x I2C0-scFc	VH CDR1	DYYMY
358.	PM 76- B10.11 x I2C0-scFc	VH CDR2	IISDGGYYTYYSDIKIG
359.	PM 76- B10.11 x I2C0-scFc	VH CDR3	GFPLLRHGAMDY
360.	PM 76- B10.11 x I2C0-scFc	VL CDR1	KASQNVDTNVA
361.	PM 76- B10.11 x I2C0-scFc	VL CDR2	SASYVYW
362.	PM 76- B10.11 x I2C0-scFc	VL CDR3	QQYDQQLIT
363.	PM 76- B10.11 x I2C0-scFc	VH	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIISDGGYYTYYSDIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSS
364.	PM 76- B10.11 x I2C0-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSL IYSASYVYWDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQ LITFGGGTKLEIK
365.	PM 76- B10.11 x I2C0-scFc	scFv	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIISDGGYYTYYSDIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFG GGTKLEIK
366.	PM 76- B10.11 x I2C0-scFc	bispecific molecule	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIISDGGYYTYYSDIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
367.	PM 76- B10.11 x I2C0-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIISDGGYYTYYSDIKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGSGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVTVSSGG GGSGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ

			PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
368.	PM 76-B10.11 x I2C0-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVQPGESLRLSAASGFTTFSDYYMYWVRQAPGKGLEWVAIIISDGGYYTYYSDI IKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGSGGGSGGGSDIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFGGGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGSGGGSGQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
369.	PM 76-B10.11 x I2C0 CC (103/43)-scFc	bispecific molecule	QVQLVESGGGLVQPGESLRLSAASGFTTFSDYYMYWVRQAPGKGLEWVAIIISDGGYYTYYSDI IKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGSGGGSGGGSDIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFGGGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLTVTVSSGGGGSGGGSGGGSGQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
370.	PM 76-B10.11 x I2C0 CC (103/43)-scFc	bispecific HLE molecule	QVQLVESGGGLVQPGESLRLSAASGFTTFSDYYMYWVRQAPGKGLEWVAIIISDGGYYTYYSDI IKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGSGGGSGGGSDIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFGGGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLTVTVSSGGGGSGGGSGGGSGQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK

			REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHYTQKSLSLSPGK
371.	PM 76-B10.11 x I2C0 CC (103/43)-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVLPKPGESLRLSAASGFTTFSDYYMYWVRQAPGKGLEWVAIISDGGYYTYYSDDIIKGRFTISRDNANKSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMDYWGQGTLVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSASYVYWDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFGGGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGTTLTVTVSSGGGGSGGGGSGGGGSGTIVVTQEPSTLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQCPRLIGGTFKFLAPGTPARFSGSLLGGKAALTLISGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHYTQKSLSLSPGK
372.	PM 76-B10.11 CC x I2C0-scFc	VH CDR1	DYYMY
373.	PM 76-B10.11 CC x I2C0-scFc	VH CDR2	IISDGGYYTYYSDDIIKG
374.	PM 76-B10.11 CC x I2C0-scFc	VH CDR3	GFPLLRHGAMDY
375.	PM 76-B10.11 CC x I2C0-scFc	VL CDR1	KASQNVDTNVA
376.	PM 76-B10.11 CC x I2C0-scFc	VL CDR2	SASYVYW
377.	PM 76-B10.11 CC x I2C0-scFc	VL CDR3	QQYDQQLIT
378.	PM 76-B10.11 CC x I2C0-scFc	VH	QVQLVESGGGLVLPKPGESLRLSAASGFTTFSDYYMYWVRQAPGKCLEWVAIISDGGYYTYYSDDIIKGRFTISRDNANKSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMDYWGQGTLVTVSS
379.	PM 76-B10.11 CC x I2C0-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSASYVYWDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFGCGTKLEIK
380.	PM 76-B10.11 CC x I2C0-scFc	scFv	QVQLVESGGGLVLPKPGESLRLSAASGFTTFSDYYMYWVRQAPGKCLEWVAIISDGGYYTYYSDDIIKGRFTISRDNANKSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMDYWGQGTLVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSASYVYWDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFGCGTKLEIK
381.	PM 76-B10.11 CC x I2C0-scFc	bispecific molecule	QVQLVESGGGLVLPKPGESLRLSAASGFTTFSDYYMYWVRQAPGKCLEWVAIISDGGYYTYYSDDIIKGRFTISRDNANKSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMDYWGQGTLVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSASYVYWDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFG

			CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
382.	PM 76- B10.11 CC x I2C0-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPFLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGG GGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
383.	PM 76- B10.11 CC x I2C0- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPFLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
384.	PM 76- B10.11 CC x I2C0 CC (103/43)-scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPFLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
385.	PM 76- B10.11 CC x I2C0 CC	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPFLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT

	(103/43)-scFc		QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTISSVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNVWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGG GGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
386.	PM 76- B10.11 CC x I2C0 CC (103/43)- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYSDIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTISSVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNVWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
387.	PM 76- B10.14 x I2C0-scFc	VH CDR1	DYYMY
388.	PM 76- B10.14 x I2C0-scFc	VH CDR2	IISDAGYYTYYSDIKKG
389.	PM 76- B10.14 x I2C0-scFc	VH CDR3	GFPLLRRHGAMDY
390.	PM 76- B10.14 x I2C0-scFc	VL CDR1	KASQNVDANVA
391.	PM 76- B10.14 x I2C0-scFc	VL CDR2	SASYRYS
392.	PM 76- B10.14 x I2C0-scFc	VL CDR3	QQYDQQLIT
393.	PM 76-	VH	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIISDAGYYTYYSDIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY

	B10.14 x I2C0-scFc		YCARGFPLLRHGAMDYWGQGLVTVSS
394.	PM 76- B10.14 x I2C0-scFc	VL	DIQMTQSPSSLSASVGDRVTITCKASQNVDANVAWYQQKPGQAPKSL IYSASYRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQ LITFGGGTKLEIK
395.	PM 76- B10.14 x I2C0-scFc	scFv	QVQLVESGGGLVKPAGESLRLSAASGFTFSDYYMYWVRQAPGKGLEW VAIIISDAGYYTYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDANVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQQLITFG GGTKLEIK
396.	PM 76- B10.14 x I2C0-scFc	bispecific molecule	QVQLVESGGGLVKPAGESLRLSAASGFTFSDYYMYWVRQAPGKGLEW VAIIISDAGYYTYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDANVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
397.	PM 76- B10.14 x I2C0-scFc	bispecific HLE molecule	QVQLVESGGGLVKPAGESLRLSAASGFTFSDYYMYWVRQAPGKGLEW VAIIISDAGYYTYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDANVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGG GGSGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
398.	PM 76- B10.14 x I2C0- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPAGESLRLSAASGFTFSDYYMYWVRQAPGKGLEW VAIIISDAGYYTYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDANVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGGGGSGGGSGGGSGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL

			HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGS FFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
399.	PM 76- B10.14 x I2C0 CC (103/43)- scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKGLEW VAIISDAGYYTYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGTLLTVSSGG GGSGGGSGGGGSGTQVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCPRLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
400.	PM 76- B10.14 x I2C0 CC (103/43)- scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKGLEW VAIISDAGYYTYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGTLLTVSSGG GGSGGGSGGGGSGTQVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCPRLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNV SCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSD GSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
401.	PM 76- B10.14 x I2C0 CC (103/43)- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKGLEW VAIISDAGYYTYSDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YRYSVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGTLLTVSSGG GGSGGGSGGGGSGTQVVTQEPSTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCPRLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNV SCSVMHEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGS FFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
402.	PM 76- B10.14 CC x I2C0-scFc	VH CDR1	DYYMY
403.	PM 76- B10.14 CC x	VH CDR2	IISDAGYYTYSDIIKG

	I2C0-scFc		
404.	PM 76-B10.14 CC x I2C0-scFc	VH CDR3	GFPLLRHGAMDY
405.	PM 76-B10.14 CC x I2C0-scFc	VL CDR1	KASQNV DANVA
406.	PM 76-B10.14 CC x I2C0-scFc	VL CDR2	SASYRYS
407.	PM 76-B10.14 CC x I2C0-scFc	VL CDR3	QQYDQQLIT
408.	PM 76-B10.14 CC x I2C0-scFc	VH	QVQLVESGGGLVKPGE SLRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAII SDAGYYTYYS DI I KGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSS
409.	PM 76-B10.14 CC x I2C0-scFc	VL	DIQMTQSPSSLSASV GDRVTITCKASQNV DANVAWYQQKPGQAPKSL IYSASYRYS DVP SRFSGSASGTDFTLTIS SVQSEDFATYYCQQYDQQ LITFGCGTKLEIK
410.	PM 76-B10.14 CC x I2C0-scFc	scFv	QVQLVESGGGLVKPGE SLRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAII SDAGYYTYYS DI I KGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGSGGGSGGGSGGGSDI QMT QSPSSLSASV GDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLTIS SVQSEDFATYYCQQYDQQLITFG CGTKLEIK
411.	PM 76-B10.14 CC x I2C0-scFc	bispecific molecule	QVQLVESGGGLVKPGE SLRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAII SDAGYYTYYS DI I KGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGSGGGSGGGSGGGSDI QMT QSPSSLSASV GDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLTIS SVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGSGGGGSQT VVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
412.	PM 76-B10.14 CC x I2C0-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGE SLRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAII SDAGYYTYYS DI I KGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGSGGGSGGGSGGGSDI QMT QSPSSLSASV GDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLTIS SVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGSGGGGSQT VVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTCPPCPAPELL GGPSVFLFPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNV SCSVMH EALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGG GGGGGSDKHTCPPCPAPELLGGPSVFLFPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVSCSVMH EALHNHYTQKSLSLSPGK
413.	PM 76-	bispecific	QVQLVESGGGLVKPGE SLRLSCAASGFTTFSDYYMYWVRQAPGKCLEW VAII SDAGYYTYYS DI I KGRFTISRDN AKNSLYLQMNSLKAEDTAVY

	B10.14 CC x I2C0-scFc_delGK	HLE molecule	YCARGFPLL RHGAMDYWGQGT LVT VSSGGGSGGGSGGGSDIQMT QSPSSLSASVGDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFLTITISSVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVT VSSGG GSGGGGSGGGGSGT VVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDNLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGGGSGGGSGGGSGGGSGGGGS GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDNLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
414.	PM 76-B10.14 CC x I2C0 CC (103/43)-scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDAGYYTYYSDI IKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGT LVT VSSGGGSGGGSGGGSDIQMT QSPSSLSASVGDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFLTITISSVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGT LVT VSSGG GSGGGGSGGGGSGT VVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCP RGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
415.	PM 76-B10.14 CC x I2C0 CC (103/43)-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDAGYYTYYSDI IKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGT LVT VSSGGGSGGGSGGGSDIQMT QSPSSLSASVGDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFLTITISSVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGT LVT VSSGG GSGGGGSGGGGSGT VVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCP RGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDNLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGKGGGSGGGSGGGSGGGSGGGG GSGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDNLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
416.	PM 76-B10.14 CC x I2C0 CC (103/43)-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIISDAGYYTYYSDI IKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGT LVT VSSGGGSGGGSGGGSDIQMT QSPSSLSASVGDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFLTITISSVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGT LVT VSSGG GSGGGGSGGGGSGT VVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCP RGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL

			GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGGSG GGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
417.	PM 76- B10.17 x I2C0-scFc	VH CDR1	DYYMY
418.	PM 76- B10.17 x I2C0-scFc	VH CDR2	IISDAGYYTYYSDIKKG
419.	PM 76- B10.17 x I2C0-scFc	VH CDR3	GFPLLRHGAMDY
420.	PM 76- B10.17 x I2C0-scFc	VL CDR1	KASQNV DANVA
421.	PM 76- B10.17 x I2C0-scFc	VL CDR2	SASYVYW
422.	PM 76- B10.17 x I2C0-scFc	VL CDR3	QQYDQQLIT
423.	PM 76- B10.17 x I2C0-scFc	VH	QVQLVESGGGLVKPGESLR LSCAASGFTFSDYYMYWVRQAPGKGLEW VAIISDAGYYTYYSDIKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSS
424.	PM 76- B10.17 x I2C0-scFc	VL	DIQMTQSPSSLSASVGD RVTITCKASQNV DANVAWYQQKPGQAPKSL IYSASYVYWDVPSRFSGSASGTDFTLTIS SVQSEDFATYYCQQYDQQ LITFGGGTKLEIK
425.	PM 76- B10.17 x I2C0-scFc	scFv	QVQLVESGGGLVKPGESLR LSCAASGFTFSDYYMYWVRQAPGKGLEW VAIISDAGYYTYYSDIKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGD RVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTIS SVQSEDFATYYCQQYDQQLITFG GGTKLEIK
426.	PM 76- B10.17 x I2C0-scFc	bispecific molecule	QVQLVESGGGLVKPGESLR LSCAASGFTFSDYYMYWVRQAPGKGLEW VAIISDAGYYTYYSDIKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGD RVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTIS SVQSEDFATYYCQQYDQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQP GGS LKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSGTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
427.	PM 76- B10.17 x I2C0-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLR LSCAASGFTFSDYYMYWVRQAPGKGLEW VAIISDAGYYTYYSDIKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGD RVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTIS SVQSEDFATYYCQQYDQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQP GGS LKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT

			<p>AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGGG GGSGGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK</p>
428.	PM 76- B10.17 x I2C0- scFc_delGK	bispecific HLE molecule	<p>QVQLVESGGGLVKPGE SLRLSCAASGFTFSDYYMYWVRQAPGKGLEW VAII SDAGYYTYSDIIKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGGGGSGGGSGGGSGGGSGGGGSGGGG GGGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK</p>
429.	PM 76- B10.17 x I2C0 CC (103/43)- scFc	bispecific molecule	<p>QVQLVESGGGLVKPGE SLRLSCAASGFTFSDYYMYWVRQAPGKGLEW VAII SDAGYYTYSDIIKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCP RGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL</p>
430.	PM 76- B10.17 x I2C0 CC (103/43)- scFc	bispecific HLE molecule	<p>QVQLVESGGGLVKPGE SLRLSCAASGFTFSDYYMYWVRQAPGKGLEW VAII SDAGYYTYSDIIKGRFTISRDNAKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCP RGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGGG</p>

			GSGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALHNNHYTQKSLSLSPGK
431.	PM 76-B10.17 x I2C0 CC (103/43)-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKGLEWVAIIISDAGYYTYYSDDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDQQLITFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGTLLTVTVSSGG GSGGGGSGGGGSGTQVTVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCPRLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG SGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVSCSVMHEALHNNHYTQKSLSLSPGK
432.	PM 76-B10.17 CC x I2C0-scFc	VH CDR1	DYYMY
433.	PM 76-B10.17 CC x I2C0-scFc	VH CDR2	IISDAGYYTYYSDDIIKG
434.	PM 76-B10.17 CC x I2C0-scFc	VH CDR3	GFPLLRHGAMDY
435.	PM 76-B10.17 CC x I2C0-scFc	VL CDR1	KASQNV DANVA
436.	PM 76-B10.17 CC x I2C0-scFc	VL CDR2	SASYVYW
437.	PM 76-B10.17 CC x I2C0-scFc	VL CDR3	QQYDQQLIT
438.	PM 76-B10.17 CC x I2C0-scFc	VH	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKCLEWVAIIISDAGYYTYYSDDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSS
439.	PM 76-B10.17 CC x I2C0-scFc	VL	DIQMTQSPSSLSASVGDRVTITCKASQNV DANVAWYQQKPGQAPKSL IYSASYVYWDVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDQQ LITFGCGTKLEIK
440.	PM 76-B10.17 CC x I2C0-scFc	scFv	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKCLEWVAIIISDAGYYTYYSDDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDQQLITFG CGTKLEIK
441.	PM 76-	bispecific	QVQLVESGGGLVKPGESLRLSCAASGFTTFSDYYMYWVRQAPGKCLEWVAIIISDAGYYTYYSDDIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY

	B10.17 CC x I2C0-scFc	molecule	YCARGFPLL RHGAMDYWGQGT LVT VSSGGGSGGGSGGGSGDIQMT QSPSSLSASVGDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVT VSSGG GGSGGGSGGGGSQT VVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
442.	PM 76- B10.17 CC x I2C0-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKCLEW VAIISDAGYYTYYSDI IKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGT LVT VSSGGGSGGGSGGGSGDIQMT QSPSSLSASVGDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVT VSSGG GGSGGGSGGGGSQT VVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGGG GGGGSGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
443.	PM 76- B10.17 CC x I2C0- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKCLEW VAIISDAGYYTYYSDI IKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGT LVT VSSGGGSGGGSGGGSGDIQMT QSPSSLSASVGDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVT VSSGG GGSGGGSGGGGSQT VVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGSGGGSGGGSGGGSGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
444.	PM 76- B10.17 CC x I2C0 CC (103/43)-scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYWVRQAPGKCLEW VAIISDAGYYTYYSDI IKGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLL RHGAMDYWGQGT LVT VSSGGGSGGGSGGGSGDIQMT QSPSSLSASVGDRVTITCKASQNV DANVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVT VSSGG GGSGGGSGGGGSQT VVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL

445.	PM 76-B10.17 CC x I2C0 CC (103/43)-scFc	bispecific HLE molecule	QVQLVESGGGLVQKPGESLRLSAASGFTTFSDYYMYWVRQAPGKCLEW VAIISDAGYYTYYSDDIIKGRFTISRDNANKSLYLQMNLSKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDANVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFNGNSYISYWAYCGQGTLLTVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCPRLIGGKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGG GGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
446.	PM 76-B10.17 CC x I2C0 CC (103/43)-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVQKPGESLRLSAASGFTTFSDYYMYWVRQAPGKCLEW VAIISDAGYYTYYSDDIIKGRFTISRDNANKSLYLQMNLSKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDANVAWYQQKPGQAPKSLIYSAS YVYWDVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDQQLITFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFNGNSYISYWAYCGQGTLLTVTVSSGG GGSGGGSGGGGSGTQVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCPRLIGGKFLAPGTPARFSGSLLGGKAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
447.	PM 76-B10 x I2C0-scFc	VH CDR1	DYYMY
448.	PM 76-B10 x I2C0-scFc	VH CDR2	IISDGGYYTYYSDDIIKG
449.	PM 76-B10 x I2C0-scFc	VH CDR3	GFPLLRHGAMDY
450.	PM 76-B10 x I2C0-scFc	VL CDR1	KASQNVDTNVA
451.	PM 76-B10 x I2C0-scFc	VL CDR2	SASYRYS
452.	PM 76-B10 x I2C0-scFc	VL CDR3	QQYDSYPYT
453.	PM 76-B10 x I2C0-scFc	VH	QVQLVESGGGLVQKPGESLRLSAASGFTTFSDYYMYWVRQAPGKGLEW VAIISDGGYYTYYSDDIIKGRFTISRDNANKSLYLQMNLSKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVTVSS
454.	PM 76-B10 x I2C0-scFc	VL	DIQMTQSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSL IYSASYRYSVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDSY PYTFGGGKLEIK

455.	PM 76-B10 x I2C0-scFc	scFv	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSIDI I KGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGT LVT VSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDSYPYTFG GGTKLEIK
456.	PM 76-B10 x I2C0-scFc	bispecific molecule	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSIDI I KGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGT LVT VSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQP GGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVT VSSGG GGSGGGSGGGGSGTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
457.	PM 76-B10 x I2C0-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSIDI I KGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGT LVT VSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQP GGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVT VSSGG GGSGGGSGGGGSGTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGG GGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
458.	PM 76-B10 x I2C0-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSIDI I KGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGT LVT VSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTITSSVQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQP GGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVT VSSGG GGSGGGSGGGGSGTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
459.	PM 76-B10 x I2C0 CC (103/43)-scFc	bispecific molecule	QVQLVESGGGLVKPGEISRLSCAASGFTTFSDYYMYWVRQAPGKGLEW VAIIISDGGYYTYYSIDI I KGRFTISRDN AKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGT LVT VSSGGGSGGGGSGGGGSDIQMT

			QSPSSLSASVGDVRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISSVQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
460.	PM 76-B10 x I2C0 CC (103/43)-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKGLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDVRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISSVQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGG GGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
461.	PM 76-B10 x I2C0 CC (103/43)- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYWVRQAPGKGLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDVRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRYSDVPSRFSGSASGTDFTLTISSVQSEDFATYYCQQYDSYPYTFG GGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGSGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
462.	PM 76-B10 CC x I2C0- scFc	VH CDR1	DYYMY
463.	PM 76-B10 CC x I2C0- scFc	VH CDR2	IISDGGYYTYYSDI IKG
464.	PM 76-B10 CC x I2C0- scFc	VH CDR3	GFPLLRHGAMDY
465.	PM 76-B10	VL CDR1	KASQNVDTNVA

	CC x I2C0-scFc		
466.	PM 76-B10 CC x I2C0-scFc	VL CDR2	SASYRYS
467.	PM 76-B10 CC x I2C0-scFc	VL CDR3	QQYDSYPYT
468.	PM 76-B10 CC x I2C0-scFc	VH	QVQLVESGGGLVKPGEISRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIIISDGGYYTYYSIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVSS
469.	PM 76-B10 CC x I2C0-scFc	VL	DIQMTQSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSL IYSASYRSDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDSY PYTFGCGTKLEIK
470.	PM 76-B10 CC x I2C0-scFc	scFv	QVQLVESGGGLVKPGEISRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIIISDGGYYTYYSIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDSYPYTFG CGTKLEIK
471.	PM 76-B10 CC x I2C0-scFc	bispecific molecule	QVQLVESGGGLVKPGEISRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIIISDGGYYTYYSIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVSSGG GGSGGGSGGGGSGTQVTVTQEPSTVSPGGTVTLTCSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLGGAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
472.	PM 76-B10 CC x I2C0-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGEISRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIIISDGGYYTYYSIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVSSGG GGSGGGSGGGGSGTQVTVTQEPSTVSPGGTVTLTCSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLGGAALTLGSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNV SCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGG GGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEV TCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD GSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK
473.	PM 76-B10 CC x I2C0-scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGEISRLSCAASGFTFSDYYMYWVRQAPGKCLEW VAIIISDGGYYTYYSIIKGRFTISRDNANKNSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGTLLTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQAPKSLIYSAS YRSDVPSRFSGSASGTDFTLTISVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVSSGG

			GGSGGGGSGGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
474.	PM 76-B10 CC x 12C0 CC (103/43)-scFc	bispecific molecule	QVQLVESGGGLVKPGESLRLSAASGFTTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCP RGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVL
475.	PM 76-B10 CC x 12C0 CC (103/43)-scFc	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSAASGFTTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCP RGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGG GGGGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK
476.	PM 76-B10 CC x 12C0 CC (103/43)- scFc_delGK	bispecific HLE molecule	QVQLVESGGGLVKPGESLRLSAASGFTTFSDYYMYWVRQAPGKCLEW VAIISDGGYYTYYSDI IKGRFTISRDNANKSLYLQMNSLKAEDTAVY YCARGFPLLRHGAMDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCKASQNVD TNVAWYQQKPGQAPKSLIYSAS YRYS DVP SRFSGSASGTDFTLT ISSVQSEDFATYYCQQYDSYPYTFG CGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYA MNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNT AYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYCGQGLTVTVSSGG GGSGGGGSGGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNY PNWVQQKPGQCP RGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMEALHNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGG GGGGSDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV

			VVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSSFLLYSKLTVDKSRWQQGNVSCFSVMHEALHNHYTQKSLSLSPGK
477.	IgG1 hinge		DKTHTCPPCP
478.	IgG2 hinge		ERKCCVECP
479.	IgG3 hinge		ELKPTPLDTHHTCPPCP
480.	IgG4 hinge		ESKYGPPCPSCP
481.	EGFRvIIIccx12 C-Hinge-CH2-CH3-linker-hinge-CH2-CH3 (DF9)	bispecific HLE molecule	QVQLVESGGGVVQSGRSLRLSCAASGFTFRNYGMHWVRQAPGKCLEWVAIVIWDGSDKYIADSVRGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARDGYDILTG NPRDFDYWGQGLTVTVSSGGGGSGGGGSGGGGSDT VMTQTPLSSSHVTLGQPASISCRSSQSLVHSDGNTYLSWLQQRPGQPPRLLIYRISRFRSGVPDRFSGSGAGTDFTLEISRVEAEDVGVIYCMQS THVPRTFGCGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEP SLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKALTL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCP PCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSSFLLYSKLTVDKSRWQQGNVSCFSVMHEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSDKTHHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSSFLLYSKLTVDKSRWQQGNVSCFSVMHEALHNHYTQKSLSLSPGK
482.	EGFRvIIIccx12 C-Hinge-CH2-CH3-linker-CH2-CH3 (T2G)	bispecific HLE molecule	QVQLVESGGGVVQSGRSLRLSCAASGFTFRNYGMHWVRQAPGKCLEWVAIVIWDGSDKYIADSVRGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARDGYDILTG NPRDFDYWGQGLTVTVSSGGGGSGGGGSGGGGSDT VMTQTPLSSSHVTLGQPASISCRSSQSLVHSDGNTYLSWLQQRPGQPPRLLIYRISRFRSGVPDRFSGSGAGTDFTLEISRVEAEDVGVIYCMQS THVPRTFGCGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEP SLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKALTL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCP PCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSSFLLYSKLTVDKSRWQQGNVSCFSVMHEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSSFLLYSKLTVDKSRWQQGNVSCFSVMHEALHNHYTQKSLSLSPGK
483.	EGFRvIIIccx12 C- Hinge-CH2-linker-Hinge-CH2-CH3-linker-CH3 (D3L)	bispecific HLE molecule	QVQLVESGGGVVQSGRSLRLSCAASGFTFRNYGMHWVRQAPGKCLEWVAIVIWDGSDKYIADSVRGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARDGYDILTG NPRDFDYWGQGLTVTVSSGGGGSGGGGSGGGGSDT VMTQTPLSSSHVTLGQPASISCRSSQSLVHSDGNTYLSWLQQRPGQPPRLLIYRISRFRSGVPDRFSGSGAGTDFTLEISRVEAEDVGVIYCMQS THVPRTFGCGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEP SLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKALTL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCP PCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSSFLLYSKLTVDKSRWQQGNVSCFSVMHEALHNHYTQKSLSLSPGK

			<p>ALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTTCP</p> <p>PCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVK</p> <p>FNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKC</p> <p>KVSNKALPAPIEKTISKAKGSGGGSGGGSGGGSGGGSGGGSDKHTCPP</p> <p>CPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKF</p> <p>NWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCK</p> <p>VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLV</p> <p>KGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSR</p> <p>WQQGNVSCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGG</p> <p>GGSGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWES</p> <p>NGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHE</p> <p>ALHNHYTQKSLSLSPGK</p>
484.	EGFRvIIIccx12 C- Hinge- CH2-linker- CH2-CH3- linker-CH3 (T7I)	bispecific HLE molecule	<p>QVQLVESGGGVVQSGRSLRLSCAASGFTFRNYGMHWVRQAPGKCLEW</p> <p>VAVIWDGSDKYYADSVRGRFTISRDN SKNTLYLQMNSLRAEDTAVY</p> <p>YCARDGYDILTG NPRDFDYWGQGT LVTVSSGGGGSGGGSGGGSDT</p> <p>VMTQTPLSSHVT LGQPASISCRSSQSLVHSDGNTYLSWLQQRPGQPP</p> <p>RLLIYRISR RFGV PDRFSGSGAGTDFTLEISRVEAEDVGVYYCMQS</p> <p>THVPRTFGCGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAAS</p> <p>GFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTI</p> <p>SRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT</p> <p>LVTVSSGGGGSGGGSGGGSGQTVVTQEP SLTVSPGGTVTLTCGSST</p> <p>GAVTSGNYPNWWQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKA</p> <p>ALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTTCP</p> <p>PCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVK</p> <p>FNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKC</p> <p>KVSNKALPAPIEKTISKAKGSGGGSGGGSGGGSGGGSGGSAPELLGGP</p> <p>SVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVH</p> <p>NAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPI</p> <p>EKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAV</p> <p>EWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCS</p> <p>VMHEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGQPREP</p> <p>QVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKT</p> <p>TPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKS</p> <p>LSLSPGK</p>
485.	EGFRvIIIccx12 C-CH2-linker- CH2-CH3- linker-CH3 (K6C)	bispecific HLE molecule	<p>QVQLVESGGGVVQSGRSLRLSCAASGFTFRNYGMHWVRQAPGKCLEW</p> <p>VAVIWDGSDKYYADSVRGRFTISRDN SKNTLYLQMNSLRAEDTAVY</p> <p>YCARDGYDILTG NPRDFDYWGQGT LVTVSSGGGGSGGGSGGGSDT</p> <p>VMTQTPLSSHVT LGQPASISCRSSQSLVHSDGNTYLSWLQQRPGQPP</p> <p>RLLIYRISR RFGV PDRFSGSGAGTDFTLEISRVEAEDVGVYYCMQS</p> <p>THVPRTFGCGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAAS</p> <p>GFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTI</p> <p>SRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT</p> <p>LVTVSSGGGGSGGGSGGGSGQTVVTQEP SLTVSPGGTVTLTCGSST</p> <p>GAVTSGNYPNWWQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKA</p> <p>ALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGAPPELLGG</p> <p>PSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEV</p> <p>HNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAP</p> <p>IEKTISKAKGSGGGSGGGSGGGSGGGSGGSAPELLGGPSVFLFPPKPK</p> <p>DTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQ</p> <p>YGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQ</p> <p>PREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPEN</p> <p>NYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHY</p> <p>TQKSLSLSPGKGGGGSGGGSGGGSGGGSGGQPREPQVYTLPPSRE</p> <p>EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS</p> <p>FFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK</p>
486.	IgG3 hinge		ELKTPLGDTTHTCPRCP
487.	IgG1 hinge		EPKSCDKTHTCPPCP

Claims

1. An antibody construct comprising:
 - a first domain which binds to PSMA,
 - a second domain which binds to an extracellular epitope of the human and the *Macaca* CD3 ϵ chain; and
 - a third domain which comprises two polypeptide monomers, each comprising a hinge, a CH2 domain and a CH3 domain, wherein said two polypeptide monomers are fused to each other via a peptide linker.
2. The antibody construct of claim 1, wherein the antibody construct is a single chain antibody construct.
3. The antibody construct of claim 1 or 2, wherein said third domain comprises in an amino to carboxyl order:

hinge-CH2-CH3-linker-hinge-CH2-CH3.
4. The antibody construct of any one of the preceding claims, wherein each of said polypeptide monomers has an amino acid sequence that is at least 90% identical to a sequence selected from the group from the group consisting of: SEQ ID NO: 17-24.
5. The antibody construct of claim 4, wherein each of said polypeptide monomers has an amino acid sequence selected from SEQ ID NO: 17-24.
6. The antibody construct of any one of the preceding claims, wherein the CH2 domain comprises an intra domain cysteine disulfide bridge.
7. The antibody construct of any one of the preceding claims, wherein
 - (i) the first domain comprises two antibody variable domains and the second domain comprises two antibody variable domains;
 - (ii) the first domain comprises one antibody variable domain and the second domain comprises two antibody variable domains;
 - (iii) the first domain comprises two antibody variable domains and the second domain comprises one antibody variable domain; or

- (iv) the first domain comprises one antibody variable domain and the second domain comprises one antibody variable domain.
8. The antibody construct of any one of the preceding claims, wherein the first and second domain are fused to the third domain via a peptide linker.
9. The antibody construct according to any one of the preceding claims, wherein the antibody construct comprises in an amino to carboxyl order:
- (a) the first domain;
 - (b) a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-3;
 - (c) the second domain;
 - (d) a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1, 2, 3, 9, 10, 11 and 12;
 - (e) the first polypeptide monomer of the third domain;
 - (f) a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 5, 6, 7 and 8; and
 - (g) the second polypeptide monomer of the third domain.
10. The antibody construct according to any one of the preceding claims, wherein the antibody construct comprises in an amino to carboxyl order:
- (a) the first domain having an amino acid sequence selected from the group consisting of SEQ ID NOs: 50, 56, 68, 74, 86, 92, 104, 110, 122, 128, 140, 146, 158, 164, 176, 182, 194, 200, 212, 218, 230, 236, 248, 254, 266, 272, 284, 290, 302, 308, 320, 335, 350, 365, 380, 395, 410, 425, 440, 455, 470;
 - (b) a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-3;
 - (c) the second domain having an amino acid sequence selected from the group consisting of SEQ ID NOs: 23, 25, 41, 43, 59, 61, 77, 79, 95, 97, 113, 115, 131, 133, 149, 151, 167, 169, 185 or 187 of WO 2008/119567 or as depicted in SEQ ID NO: 15;
 - (d) a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1, 2, 3, 9, 10, 11 and 12;
 - (e) the first polypeptide monomer of the third domain having a polypeptide sequence selected from the group consisting of SEQ ID NOs: 17-24;

- (f) a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 5, 6, 7 and 8; and
 - (g) the second polypeptide monomer of the third domain having a polypeptide sequence selected from the group consisting of SEQ ID NOs: 17-24.
11. The antibody construct according to claim 10, having an amino acid sequence selected from the group consisting of:
- SEQ ID NOs: 52, 53, 58, 59, 70, 71, 76, 77, 88, 89, 94, 95, 106, 107, 112, 113, 124, 125, 130, 131, 142, 143, 148, 149, 160, 161, 166, 167, 178, 179, 184, 185, 196, 197, 202, 203, 214, 215, 220, 221, 232, 233, 238, 239, 250, 251, 256, 257, 268, 269, 274, 275, 286, 287, 292, 293, 304, 305, 310, 311, 322, 323, 325, 326, 337, 338, 340, 341, 352, 353, 355, 356, 367, 368, 370, 371, 382, 383, 385, 386, 397, 398, 400, 401, 412, 413, 415, 416, 427, 428, 430, 431, 442, 443, 445, 446, 457, 458, 460, 461, 472, 473, 475 and 476.
12. A polynucleotide encoding an antibody construct as defined in any one of the preceding claims.
13. A vector comprising a polynucleotide as defined in claim 12.
14. A host cell transformed or transfected with the polynucleotide as defined in claim 12 or with the vector as defined in claim 13.
15. A process for the production of an antibody construct according to any one of claims 1 to 11, said process comprising culturing a host cell as defined in claim 12 under conditions allowing the expression of the antibody construct as defined in any one of claims 1 to 11 and recovering the produced antibody construct from the culture.
16. A pharmaceutical composition comprising an antibody construct according to any one of claims 1 to 11, or produced according to the process of claim 15.
17. The pharmaceutical composition of claim 16, which is stable for at least four weeks at about -20°C.

18. The antibody construct according to any one of claims 1 to 13, or produced according to the process of claim 15, for use in the prevention, treatment or amelioration of a disease selected from a proliferative disease, a tumorous disease, cancer or an immunological disorder.
19. The use according to claim 18, wherein the disease is prostate cancer.
20. A method for the treatment or amelioration of a proliferative disease, a tumorous disease, cancer, or an immunological disorder, comprising the step of administering to a subject in need thereof the antibody construct according to any one of claims 1 to 11, or produced according to the process of claim 15.
21. A kit comprising an antibody construct according to any one of claims 1 to 11, or produced according to the process of claim 15, a polynucleotide as defined in claim 12, a vector as defined in claim 13, and/or a host cell as defined in claim 14.