

PATENT SPECIFICATION

NO DRAWINGS

1.058.557

Date of Application and filing Complete Specification: March 31, 1964.

No. 13171/64.

Application made in United States of America (No. 269,751) on April 1, 1963.

Complete Specification Published: Feb. 15, 1967.

© Crown Copyright 1967.



Index at acceptance:—C2 C(1H2A7, 1H2C2, 1Q3, 1Q6B1, 1Q8C, 1Q9E, 1Q11B); A5 E(1A4A2, 1A4A4, 1C4A2, 1C4A4)

Int. Cl.:—C 07 c 121/54 // A 01 n, A 61 l

COMPLETE SPECIFICATION

Improvements in or relating to Isophthalonitriles


We, DIAMOND ALKALI COMPANY, a Corporation organised and existing under the laws of the State of Delaware, United States of America, of 300 Union Commerce Building, Cleveland 14, Ohio, United States of America, (Assignees of ROBERT DEAN BATTERSHELL, and HENRY BLUESTONE), do hereby declare the invention, for which we pray that a patent may be granted to us, and the method by which it is to be performed, to be particularly described in and by the following statement:—

This invention relates to novel halogenated aromatic nitriles and, more particularly, relates to ring halogenated isophthalonitriles.

This application is an improvement upon the invention described and claimed in British Specification No. 36051/62 (Specification No. 980,523).

The term "halogenated aromatic nitriles" as employed in the specification refers to di-cyano substituted benzenes having at least one halogen atom attached to a ring carbon atom.

These di-cyano substituted halogenated benzenes are isophthalonitriles of the general formula:



wherein X¹, X², X³ and X⁴ are hydrogen or halogen, preferably chlorine, fluorine and bromine, with at least one X being halogen. Typical isophthalonitrile derivatives of this invention are tetrachloroisophthalonitrile, 5-chloro-2,4,6-trifluoroisophthalonitrile and 4,6-dichloroisophthalonitrile.

[Price 4s. 6d.]

The novel halogenated isophthalonitriles of this invention may be prepared from the corresponding ring-halogenated e.g. chlorinated acid chlorides. The acid chloride is converted to the corresponding amide by dissolving the acid chloride in a suitable organic solvent and introducing ammonia. A wide range of solvents may be employed in this reaction with the presently preferred solvents being xylene or dioxane. Either anhydrous or aqueous ammonia may be employed, depending upon whether the solvent used is miscible with water.

The reaction between the acid chloride and ammonia is exothermic and the reactor may be cooled to maintain a maximum temperature of 40° to 60° C. The aromatic amide which is formed is insoluble in most organic solvents and this material precipitates from the solution. Ammonium chloride is formed as a by-product and is separated from the desired amide. If a water miscible solvent such as dioxane is used, and aqueous ammonia is employed as the source of ammonia, sufficient water may be added to the reaction mixture to dissolve the ammonium chloride, and the solid precipitate is primarily the desired halogenated aromatic amide. If a solvent such as xylene, which is not miscible with water, is used or if anhydrous ammonia is employed, the precipitate comprising the amide and the ammonium chloride is filtered and then washed with water to remove the ammonium chloride and again filtered to recover the desired amide.

The ring-chlorinated aromatic amide is dried to remove all traces of water and is reacted with dehydrating agent such as phosphorus pentoxide or phosphorus oxychloride with the preferred dehydrating agent being phosphorus oxychloride. The reaction time and temperature will be governed by the reactivity of the dehydrating agent employed. Typical conditions when phosphorus oxychloride is used are a temperature of 75° to

110° C. and a reaction time of 1 to 6 hours. The product of the dehydration reaction is an aromatic nitrile, having chlorine atoms occupying the same positions on the aromatic ring as they did in the acid chloride starting material.

The fluorinated compound may be prepared by a halogen interchange reaction whereby chlorine is replaced by fluorine. Chlorine atoms which are in position, either ortho or para to a cyano group, are activated by the cyano group and, accordingly, are readily removed or replaced by fluorine atoms. The preferred reactant is an alkali metal fluoride wherein the alkali metal is below sodium in the Periodic Table, e.g., potassium fluoride, rubidium fluoride, cesium fluoride, or cuprous fluoride, with the particularly preferred material being potassium fluoride. In order to prepare tetrafluoroterephthalonitrile an excess of the metal fluoride is employed; for example, tetrachloroterephthalonitrile is treated with 4 to 10, typically about 5, molar equivalents of potassium fluoride. The reaction preferably is carried out in a solvent such as dimethyl formamide, N-methyl-2-pyrrolidinone, dimethyl sulfoxide, dimethyl sulfone or tetramethylene sulfone, at a temperature range of 80° to 200° C. and for periods of 1/2 to 24 hours, commonly 2 to 4 hours.

The halogenated isophthalonitriles of this invention exhibit outstanding biological activity as fungicides, bactericides and nematocides.

While it is possible to apply the compounds of the present invention in undiluted form to the plant or other material to be protected, it is frequently desirable to apply them in admixture with either solid or liquid, inert, pesticidal adjuvants. Thus, the nitriles can be applied to the plants for fungicidal purposes, for example, by spraying the plants with aqueous or organic solvent dispersions of the nitriles. The choice of an appropriate solvent is determined largely by the concentration of active ingredient which it is desired to employ, by the volatility required in a solvent, the cost of the solvent and the nature of the material being treated. Among the many suitable solvents which can be employed as carriers for the present pesticides, there may be mentioned hydrocarbons such as benzene, toluene, xylene, kerosene, diesel oil, fuel oil, petroleum, naphtha; ketones such as acetone, methyl ethyl ketone and cyclohexanone; chlorinated hydrocarbons, such as carbon tetrachloride, chloroform, trichloroethylene, perchlorethylene; esters such as ethyl acetate, amyl acetate and butyl acetate; the monoalkyl ethers of ethylene glycol, e.g., the monomethyl ethers; and the monoalkyl ethers of diethylene glycol, e.g., the monoethyl ether; alcohols such as ethanol, isopropanol and amyl alcohol.

The nitriles can also be applied to plants

and other materials along with inert solid fungicidal adjuvants or carriers such as talc, pyrophyllite, Attaclay, kieselguhr, chalk, diatomaceous earth, calcium carbonate, bentonite, fuller's earth, cottonseed hulls, wheat flour, soybean flour, pumice, tripoli, wood flour, walnut shell flour and lignin.

It is frequently desirable to incorporate a surface active agent in the pesticidal compositions of this invention. Such surface active agents are advantageously employed in both the solid and liquid compositions. The surface active agent can be anionic, cationic or nonionic in character.

Typical classes of surface active agents include alkyl sulfonates, alkylaryl sulfonates, alkyl sulfates, alkylamide sulfonates, alkylaryl polyether alcohols, fatty acid esters of polyhydric alcohols, ethylene oxide addition products of such esters; addition products of long-chain mercaptans and ethylene oxide; sodium alkyl benzene sulfonates having 14 to 18 carbon atoms, alkylphenolethylene oxides, e.g., *p*-isooctyl phenol condensed with 10 ethylene oxide units; and soaps, e.g., sodium stearate and sodium oleate. A typical surface active agent is Aerosol OS (sodium salt of propylated naphthalenesulfonic acid).

The solid and liquid formulations can be prepared by any suitable method. Thus, the active ingredients, in finely divided form if a solid, may be tumbled together with finely divided solid carrier. Alternatively, the active ingredient in liquid form, including solutions, dispersions, emulsions and suspensions thereof, may be admixed with the solid carrier in finely divided form in amounts small enough to preserve the free-flowing property of the final dust compositions.

When solid compositions are employed, in order to obtain a high degree of coverage with a minimum dosage of the formulation, it is desirable that the formulation be in finely divided form. The dust containing active ingredient usually should be sufficiently fine that substantially all will pass through a 20-mesh Tyler sieve. A dust which passes through a 200-mesh Tyler sieve also is satisfactory.

For dusting purposes, preferably formulations are employed in which the active ingredient is present in an amount of 5 to 50 percent of the total by weight. However, concentrations outside this range are operative and compositions containing from 1 to 99 percent of active ingredient by weight are contemplated, the remainder being carrier and/or any other additive or adjuvant material which may be desired. It is often advantageous to add small percentages of surface active agents, e.g., 0.5 to 1 percent of the total composition by weight, to dust formulations, such as the surface active agents previously set forth.

For spray application, the active ingredient

may be dissolved or dispersed in a liquid carrier, such as water or other suitable liquid. The active ingredient can be in the form of a solution, suspension, dispersion or emulsion in aqueous or non-aqueous medium. Desirably, 0.5 to 1.0 percent of a surface active agent, by weight, is included in the liquid composition.

For adjuvant purposes, any desired quantity of surface active agent may be employed, such as up to 250-percent of the active ingredient by weight. If the surface active agent is used only to impart wetting qualities, for example, to the spray solution, as little as 0.05 percent, by weight, or less of the spray solution need be employed. The use of larger amounts of surface active agent is not based upon wetting properties but is a function of the physiological behavior of the surface active agent. These considerations are particularly applicable in the case of the treatment of plants. In liquid formulations the active ingredient often constitutes not over 30 percent, by weight, of the total and may be 10 percent, or even as low as 0.01 percent.

The novel isophthalonitriles of the present invention can be employed in compositions containing other pesticides, more especially, fungicides, insecticides and bactericides.

In order that those skilled in the art may more completely understand the present invention and the preferred methods by which

the same may be carried into effect, the following specific examples may be offered.

EXAMPLE 1

Preparation of Tetrachloroisophthaloyl Chloride

In a 2-liter, Morton (creased) flask are placed 276 g. (1.0 mole) of 2,4,5,6-tetrachloro-*m*-xylylene diol and 1 liter of iron-free carbon tetrachloride. The slurry is heated and irradiated with two 250-watt heat lamps and one 100-watt mercury vapor lamp. Into the well-stirred irradiated refluxing mixture is bubbled 775 g. (10.6 moles) of chlorine gas over an 18-hour period. The amount of carbon tetrachloride employed is sufficient to dissolve the acid chloride product so that the disappearance of the nearly insoluble diol can be used as a convenient end point for the chlorination. Air is drawn through the hot mixture to remove the dissolved chlorine and hydrochloric acid, after which the solution is evaporated to dryness, leaving the white crystalline tetrachloroisophthaloyl chloride. After one recrystallization from heptane, the product weighs 291 g. (86% yield) and has a melting point of 99° to 102° C. A sample, which is further purified by being washed with methanol and recrystallized from heptane, has a melting point of 100° to 101° C. and the following elemental analysis:

Element	Percent Calculated	Percent Actual
Carbon	28.1	28.2
Hydrogen	0.0	0.2

By the procedure of this example, other ring-chlorinated acid chlorides are prepared from *m*-xylylene diols having chlorine atoms on appropriate carbon atoms of the benzene ring.

EXAMPLE 2

Preparation of Tetrachloroisophthalamide

In a 1-liter Morton flask are placed 170 g. (0.5 mole) of tetrachloroisophthaloyl chloride and 500 ml. of dioxane. To this stirred mixture is added, dropwise, 200 ml. (3.0 moles) of 28-percent ammonium hydroxide. The reaction is exothermic and, when the reaction temperature reaches about 50° C., the

flask is cooled by means of a water bath and the temperature maintained at about 40° to 50° C. throughout the remainder of the addition period. The resulting heavy slurry is stirred for 2 hours. The mixture is diluted with an equal volume of cold water, the precipitate collected, washed with water and dried in an oven at 110° to 115° C. The dry white amorphous powder weighs 155 g. (quantitative yield). A portion of the product after recrystallization from methanol has a melting point of 398–400° C. and the assigned formula $C_6H_4Cl_4N_2O_2$, is confirmed by the following elemental analysis:

Element	Percent Calculated	Percent Actual
Carbon	31.8	31.8
Hydrogen	1.3	1.2

EXAMPLE 3

Preparation of 4,6-dichloroisophthalamide

Into a 2-liter Morton flask are placed 120 g. (0.44 mole) of 4,6-dichloroisophthaloyl chloride and 1 liter of dioxane. Into this stirred solution is bubbled 75 g. (4.4 moles) of anhydrous ammonia over a period of about 2-1/2 hours causing the reaction temperature to increase from 20° to 40° C. The mixture is heated to 60° C. with continued stirring and is then allowed to cool slowly to about

30° C. The reaction mass is suction filtered and the clear light precipitate transferred to a beaker and washed with 2 liters of cold water to dissolve out the ammonium chloride by-product. The resulting aqueous mixture is suction filtered and the precipitate is air dried to produce 78 g. (76% yield) of 4,6-dichloroisophthalamide, $C_8H_4Cl_2N_2O_2$, having a melting point of 276° to 279° C. The assigned structure is confirmed by the following elemental analysis:

Element	Percent Calculated	Percent Actual
Carbon	41.2	40.8
Hydrogen	2.6	2.5

EXAMPLE 4

Preparation of tetrachloroisophthalonitrile

Into a 300-ml. flask are placed 30 g. (0.1 mole) of tetrachloroisophthalamide and 62 g. (0.4 mole) of phosphorus oxychloride. The mixture is heated with rapid stirring at 75° to 110° C. for 3 hours, during which time there occurs a brisk evolution of HCl. The

mixture is allowed to cool, then poured over crushed ice. The tan precipitate is filtered off, air-dried, washed three times with ether and finally dried in an oven at 110° C. The yield of tetrachloroisophthalonitrile, $C_6Cl_4N_2$, is 25 g. (94% yield). A sample, after recrystallization from benzene, melts at 250°—251° C. and analyzes as follows:

Element	Percent Calculated	Percent Actual
Carbon	36.1	36.2
Hydrogen	0.0	0.2

4,6-Dichloroisophthalonitrile m.p. 185.5°—186° C. has also been prepared by reacting phosphorus oxychloride with the corresponding amide in accordance with the above process. Analysis:

	%C	%H
Calculated:	48.8	1.0
Found:	48.8	1.0

EXAMPLE 5

Conversion of Chlorine-Containing Aromatic Nitriles to the Corresponding Fluorine-Containing or Chlorine-Fluorine Containing Nitriles

By heating a mixture of dry potassium fluoride and dimethyl formamide, with stirring, and then adding to this mixture the chlorine-containing nitrile, and by employing less than the theoretical amount of potassium fluoride required for replacement of all chlorine atoms which are ortho or para to a cyano group, or by interrupting the reaction before it reaches completion, a mixture of chlorofluoronitriles is obtained. Starting with tetrachloroisophthalonitrile, for example, a mixture of dichlorodifluoroisophthalonitrile and chlorotrifluoroisophthalonitrile is obtained

and the components are separated by fractional distillation.

For instance, a mixture was made up of KF and dimethyl formamide in a mole ratio of 6 moles of KF per mole of nitrile and tetrachloroisophthalonitrile was then added. The reaction was carried out for 4 hours at 100° C. The resultant 5-chloro-2,4,6-trifluoroisophthalonitrile has a melting point of 109°—110° C. and the following elemental analysis:

	%C	%H
Calculated:	44.3	0.0
Found:	44.7	0.2

EXAMPLE 6

Inhibition of Sclerotial Germination

This test is used to measure the ability of compounds of this invention to inhibit the germination of sclerotia of *Sclerotium rolfsii*.

Sterilized soil is infested with *Sclerotium rolfsii* and subsamples placed into a series of 4-oz. Dixie cups. An appropriate amount of the test formulation is drenched onto the soil at a dosage of 64 pounds per acre. This test formulation contains 0.4 g. of test compound, viz. 5-chloro-2,4,6-trifluoroisophthalonitrile.

5 nitrile, 8 ml. acetone, 4 ml. stock emulsifier solution (0.5% "Triton" (Registered Trade Mark) X-155 in water by volume) and 187.6 ml. distilled water. The concentration of toxicant in this formulation is 2000 parts per million. Lower concentrations of toxicant are obtained by diluting the formulation with distilled water. The surface area of the soil in the cups is 5.73 sq. in.; therefore, a rate of 1 pound per acre requires 0.414 mg. of chemical per cup. After drenching, the Dixie cups are then incubated for several days at 70°F. in a controlled temperature cabinet. The test chemical was rated for its ability to inhibit mycelial growth of the organism on the surface of the soil. Untreated checks and a standard material were included in the test in addition to a check planted in sterilized soil. Using this procedure, 100 percent control was obtained at a concentration of 64 lbs./acre.

EXAMPLE 7

Spore Inhibition Test on Glass Slides

10 Inhibition of spore germination on glass slides by the test tube dilution method is adopted from the procedure recommended by the American Phytopathological Society's Committee on Standardization of Fungicidal Tests. In this screen, chemicals at 1000, 100, 10 and 1.0 p.p.m. are tested for ability to inhibit germination of spores from 7- to 10-day-old cultures of *Alternaria oleracea* Mil. and *Monilinia fructicola* (Wint.) Honey. These concentrations refer to the actual concentrations after diluting the test preparations with spore stimulant and spore suspension. A formulation containing 0.1 g. of the test compound, 4 ml. acetone, 2 ml. stock emulsifier solution (0.5% Triton X-155 in water by volume) and 74 ml. distilled water is used for this test. The concentration of toxicant in this formulation is 1250 parts per million. The concentrations given above are diluted from this original formulation. Germination records are taken after 20 hours of incubation at 22° C. by observing several microscope fields so that at least 100 spores of each fungus have been examined at each concentration. Test compounds are given alphabetical ratings which correspond to the concentration that inhibits germination of half the spores (ED50) in the test drops: AAA = 0.01 to 0.1 p.p.m.; AA = 0.1 to 1.0 p.p.m.; A = 1.0 to 10 p.p.m.; B = 10 to 100 p.p.m.; C = 100 to 1000 p.p.m.; and

D = 1000 p.p.m. Using this procedure, the following results were obtained:

Compound Tested	Rating
Tetrachloroisophthalonitrile	AA
5-Chloro-2,4,6-trifluoroisophthalonitrile	AA

EXAMPLE 8

Foliage Protectant Fungicide Test

60 The tomato foliage disease test measures the ability of the test compound to protect tomato foliage against infection by the early blight fungus *Alternaria solani* (Ell. and Mart.) Jones and Grouet and the late blight fungus *Phytophthora infestans* (Mont.) deBary. The method used is a modification of that described by McCallan and Wellman and employs tomato plants (var. Bonny Best) 5 to 7 inches high which are 4 to 6 weeks old. Duplicate plants, one set for each test fungus, are sprayed with various dosages of the test formulation at 4 lbs./sq. in. air pressure while being rotated on a turntable in a hood. The center of the turntable is 45 inches from the nozzle of the spray gun. The test formulation containing 0.2 g. of test compound, viz. tetrachloroisophthalonitrile, 8 ml. acetone, 4 ml. stock emulsifier solution (0.5% Triton X-155 in water by volume) and 187.6 ml. distilled water is applied at dosages equivalent to 1000 and 200 p.p.m. of the test chemical. Lower concentrations of toxicant are obtained by dilution, maintaining the same concentration of acetone and emulsifier.

After the spray deposit is dry treated plants and controls (sprayed with formulation without toxicant) are sprayed while being rotated on a turntable with a spore suspension containing approximately 20,000 conidia of *A. solani* per ml., or 150,000 sporangia of *P. infestans* per ml. The atomizer used delivers 20 ml. in the 30-second exposure period. The plants are held in a saturated atmosphere for 24 hours at 70° F. for early blight and 60° F. for late blight, to permit spore germination and infection before removal to the greenhouse.

After two days from the start of the test for early blight and three days for late blight, lesion counts are made on the three uppermost fully expanded leaves. The data are converted to percentage disease control based on the number of lesions obtained on the control plants. Dosages and percent disease control are given in the following table.

Compound	Dosage p.p.m.	% Control	
		E.B.	L.B.
Tetrachloroisophthalonitrile	60	100	100
	40	98	99
	30	92	99
	20	72	94
	10	0	70
	5	0	8

EXAMPLE 9

Soil Mycelial Growth Inhibition Test

5 This test is used to determine activity against seed decay and damping-off fungi, primarily *Pythium* and *Rhizoctonia* species.

10 Sterilized soil is infested with *Pythium* and *Rhizoctonia* and subsamples were then treated as described in Example 6. Using this procedure, 100 percent control was found with a concentration of 64 lbs./acre.

EXAMPLE 10

Bean Rust Systemic

15 Pinto bean plants, at a growth stage when the trifoliate leaves are just beginning to emerge from the axil of the seed leaves, are used as the test species. These plants are grown in 4-inch pots and thinned to three plants per pot. Usually the plants are about 20 10 to 14 days old from time of planting. There are, therefore, six primary seed leaves per pot for each test unit. In the test, an appropriate amount of the test formulation is drenched on each pot. This test formula-
25 tion contains 0.4 g. of test compound, 8 ml.

acetone, 4 ml. stock emulsifier solution (0.5% Triton X-155 in water by volume) and 187.6 ml. distilled water. The concentration of toxicant in this formulation is 2000 parts per million. A dosage of 56 ml. of the test formulation is equivalent to 112 mg. of chemical or 128 pounds per acre. Lower concentrations of toxicant are obtained by diluting the formulation with distilled water. About two or three hours after treatment the plants are exposed to a spore suspension of bean rust from a culture maintained in the greenhouse. After exposure, the plants are immediately placed in a moist chamber in a saturated atmosphere at 60°F., for 24 hours after which they are removed to the greenhouse. Counts are made about 10 days after spore exposure and the mean number of rust pustules per leaf is determined. These counts are calculated against the check counts to arrive at the percentage disease control. Using this procedure, the following results are obtained:

Compound	Dosage	Percent Disease Control
5-Chloro-2,4,6-trifluoroisophthalonitrile	64 lbs.	92 St 5, CH

St = stunting (1 — slight to 9 — severe)

Ch = chlorosis.

EXAMPLE 11

Post-Emergence Pan Test

50 This test measures the post-emergent herbicidal activity of test chemicals when applied to the foliage of seedling plants as well as to the soil in which they are growing. Two mixtures of seed are planted in soil contained in metal half flats, which are 8" x 12" at the soil surface. One-half of each pan is planted with a mixture containing the broadleaf species, buckwheat, aster and turnip. The other half of the pan is seeded to a mixture of three grasses, millet, perennial ryegrass, and sorghum. One-half level teaspoon of each seed mixture is planted in each pan.

65 The flats are then removed to the greenhouse and the test species are allowed to grow until one true leaf is present on the slowest growing broadleaf (aster), which requires 9 to 14 days. At this point the surface of the soil in the pans is uniformly covered by spraying, at 10 pounds per square inch air pressure, with 40 ml. of a test formulation containing the toxicant, 5-chloro-2,4,6-trifluoroisophthalonitrile. Formulations containing 165.6 mg. and 82.8 mg. of the test chemical are equivalent to a coverage of 24 and 12 pounds per acre, respectively. Each test formulation contains the toxicant, equal volumes of water and acetone, and 2 drops of Triton X-155.

One and two weeks after treatment, estimates are made on the fresh weight of seedling stand in the test pans, as well as that in untreated control pans. The percent con-

trol obtained with the test chemical is calculated. Using this procedure, the following results are obtained:

5

Compound Tested	Concentration lbs./acre	Percent Control			
		Broadleaf weeds		Grassy weeds	
		1 week	2 weeks	1 week	2 weeks
5-Chloro-2,4,6-trifluoroisophthalonitrile	24	—	65	—	85
	12	95 ¹	95 ²	70 Ph1	90
	6	90 Ph2	90 Ph2	85	80 Ph2
	3	40 Ph1	20 Ph1	10 Ph1	10 Ph2

1 = Buckwheat left.
2 = No turnip.

Ph = Phytotoxicity rating, 0 for no injury to 11 for killing the plants.

EXAMPLE 12

- 10 Soil-Watering Tests—Tomato and Bean
To measure the growth regulating and herbicidal responses when chemicals are absorbed by roots and translocated, tomato plants, var. Bonny Best, 5 to 7 inches tall, and beans, var. Tendergreen, just as the trifoliolate leaves are beginning to unfold, are treated by drenching appropriate portions of a formulation containing 0.4 g. of the test chemical, viz. 5-chloro-2,4,6-trifluoroisophthalonitrile, 8 ml. acetone, 4 ml. stock emulsifier solution (0.5% Triton X-155 in water) and 187.6 ml. distilled water (concentration of the test chemical = 2000 p.p.m.) on soil in clay pots containing the test plants. Tomatoes (1 plant per 4-inch

pot) and bean plants (4 plants per 3-1/2-inch pot) are treated with appropriate dosages of the 2000 p.p.m. formulation. Dosages of 56 ml. of the formulation in the 4-inch pot and 42 ml. in the 3-1/2-inch pot are equivalent to 112 mg. and 84 mg. of chemical per pot, respectively, or 128 pounds per acre. Plants are held in a greenhouse for two weeks before records are taken. Phytotoxicity is rated on the scale from 0, indicating no plant injury, to 11, plant kill. Additionally, all other responses such as stunting, formative effects, defoliant activity, growth regulant properties, chlorosis and the like are recorded. Using this procedure, the following results are obtained:

30

35

40

Compound	Concentration lbs./acre	Phytotoxicity Rating	
		Tomato	Beans
5-Chloro-2,4,6-trifluoroisophthalonitrile	64	11 St3, Fe	2
	32	2 St9, Ch	0 St9
	16	0	0

Ch = Chlorosis

St = Stunting (1 slight to 9 severe)

Fe = Formative effects to the plants.

EXAMPLE 13

Cucumber Anthracnose

- 45 This test measures the ability of the test compound to control anthracnose of cucumber incited by the fungus *Colletotrichum lagenarium*.

- 50 The method used is a modification of that described by McCallan and Wellman and employs cucumber plants, (var. National

Pickling) having one fully expanded leaf. Duplicate plants are sprayed with various dosages of the test formulation at 20 lbs./sq. in. air pressure while being rotated on a turntable in a hood. The center of the turntable is 45 inches from the nozzle of the spray gun. The test formulation containing 0.2 g. of test compound, 8 ml. acetone, 4 ml. stock emulsifier solution (0.5% Triton X-155 in

55

60

water by volume) and 187.6 ml. distilled water is applied at dosages equivalent to 1000 and 200 p.p.m. of the test chemical. Lower concentrations of toxicant are obtained by employing less toxicant and more water, thereby maintaining the same concentration of acetone and emulsifier.

- 5 After the spray deposit is dry, treated plants and controls (sprayed with formulation without toxicant) are sprayed while being rotated on a turntable with a spore suspension containing approximately 400,000 spores per ml. of *Colletotrichum lagenarium*. The atomizer

used delivers 20 ml. in the 30-second exposure period. The plants are held in a saturated atmosphere for 24 hours at 70°F. to permit spore germination and infection before removal to the greenhouse.

After four to six days from the start of the test lesion, counts are made. The data are converted to percentage disease control based on the number of lesions obtained on the control plants. Dosages and percent disease control are given in the following table:

Compound	Dosage p.p.m.	Percent Disease Control
Tetrachloroisophthalonitrile	64	100
	32	99
	16	95
	8	80
	4	68

EXAMPLE 14

Control of Apple Scab

- 30 The apple scab disease test measures the ability of the test compound to protect apple seedlings against infection by the fungus *Venturia inaequalis*.

- 35 The method used is a modification of that described by McCallan and Wellman and employs McIntosh apple seedlings, 5 to 8 weeks old with 4 to 6 leaves. Duplicate seedlings are sprayed with various dosages of the test formulation at 40 lbs./sq. in. air pressure while being rotated on a turntable in a hood. The center of the turntable is 45 inches from the nozzle of the spray gun. The test formulation containing 0.2 g. of test compound, 8 ml. acetone, 4 ml. stock emulsifier solution (0.5% Triton X-155 in water by volume) and 187.6 ml. distilled water is applied at dosages equivalent to 1000 and 200 p.p.m. of the test chemical. Lower con-

centrations of toxicant are obtained by employing less toxicant and more water, thereby maintaining the same concentration of acetone and emulsifier.

After the spray deposit is dry, treated seedlings and controls (sprayed with formulation without toxicant) are sprayed while being rotated on a turntable with a conidial suspension of the fungus. The atomizer used delivers 20 ml. in the 30-second exposure period. The seedlings are held in a saturated atmosphere for 48 hours at 60°F. to permit spore germination and infection before removal to the greenhouse. After 10 to 14 days from the start of the test lesion counts are made. The data are converted to percentage disease control based on the number of lesions obtained on the control plants. Dosages and percent disease control are given in the following table:

Compound	Dosage p.p.m.	Percent Disease Control
Tetrachloroisophthalonitrile	1	100
	0.5	100
	0.25	100
	0.125	50

EXAMPLE 15

- 70 Protection of Cabbage from *Sclerotium Rolfsii* by Soil Treatment

- 75 Air-dry sterile soil is infested with the organism *Sclerotium rolfsii*, grown on a mixture of corn meal and sand. The desired inoculum level is achieved by adding an appropriate amount of the corn meal-sand

culture to the soil. The soil is placed into 3-1/4" x 3-1/4" x 2-1/2" plastic boxes. An appropriate amount of test formulation is poured onto the soil to give the desired rate of application, e.g. 25 ml. is equivalent to a dosage of 64 pounds per acre. This test formulation contains 0.4 g. of test compound, 8 ml. acetone, 4 ml. stock emulsifier solution

(0.5% Triton X-155 in water by volume) and 187.6 ml. distilled water. The concentration of toxicant in this formulation is 2000 parts per million. The test formulation is thoroughly incorporated into the soil. An identical box of soil is prepared which has been treated with the test formulation, but has not been inoculated with the organism. The non-inoculated treated control is used to determine the phytotoxicity of each test

compound. Twenty-five cabbage seeds (Golden Acres) are planted in the treated soil in each box. Each box receives 50 ml. of water and is placed in a humidity chamber maintained at 70°C. for 2 days. The boxes are then transferred to the greenhouse. Effectiveness of the test chemical is determined by the percentage of plants growing 14 days after treatment and planting.

Compound Tested	Concentration lbs./acre	Percent Stand	
		Non-inoculated	Inoculated
Tetrachloroisophthalonitrile	16	100	52
5-Chloro-2,4,6-trifluoroisophthalonitrile	32	92	68
	16	—	36
	8	—	10

EXAMPLE 16

Soil Mycelial Growth Inhibition Test


The following test measures the ability of compounds of this invention to inhibit mycelial growth in soil. Sterilized soil is infested with *Rhizoctonia solani* (grown on a corn meal sand medium) by mixing two 250-ml. flasks of a ten-day-old culture of the organism per level flat of sterile soil. The infested soil is then placed in small Dixie cups (4-oz. squat). Treatment of the soil is accomplished by drenching the appropriate amount of a diluted formulation containing 0.4 g. of test compound, 8 ml. acetone, 4 ml. stock emulsifier solution (0.5% Triton X-155 in water by volume) and 187.6 ml. distilled water on the surface of the soil in the test cup. The concentration of toxicant in this formulation

is 2000 parts per million. Lower concentrations of toxicant are obtained by diluting the formulation with distilled water. Surface area of the soil in the cups is 5.73 square inches; therefore, a rate of one pound per acre requires 0.414 mg. of chemical per cup. After drenching, the cups are placed in a saturated atmosphere at 70°F. for 48 hours. By this time the fungus mycelium has completely overgrown the surface of the soil in the control cups. Inhibition of mycelial growth in the treated cups is estimated on a scale from zero, complete inhibition of growth, to ten which is equivalent to controls. These grades are expressed as percent control. Using this procedure, the following results are obtained:

Compound	Concentration lbs./acre	Percent Control
Tetrachloroisophthalonitrile	64	90
5-Chloro-2,4,6-trifluoroisophthalonitrile	64	90

WHAT WE CLAIM IS:—

1. Halogenated isophthalonitriles of the structure:



wherein each of X¹, X², X³ and X⁴ is hydrogen or halogen, at least one X being halogen.

2. Tetrachloroisophthalonitrile.

3. 5-Chloro-2,4,6-trifluoroisophthalonitrile.

4. 4,6-Dichloroisophthalonitrile.

5. A method of preparing a ring-chlorinated isophthalonitrile, which comprises reacting the corresponding ring-chlorinated isophthalamide with a dehydrating agent comprising phosphorus pentoxide or phosphorus oxychloride and separating the desired ring-chlorinated isophthalonitrile.



1,058,557

9

(0.5% Triton X-155 in water by volume) and 187.6 ml. distilled water. The concentration of toxicant in this formulation is 2000 parts per million. The test formulation is thoroughly incorporated into the soil. An identical box of soil is prepared which has been treated with the test formulation, but has not been inoculated with the organism. The non-inoculated treated control is used to determine the phytotoxicity of each test

compound. Twenty-five cabbage seeds (Golden Acres) are planted in the treated soil in each box. Each box receives 50 ml. of water and is placed in a humidity chamber maintained at 70°C. for 2 days. The boxes are then transferred to the greenhouse. Effectiveness of the test chemical is determined by the percentage of plants growing 14 days after treatment and planting.

Compound Tested	Concentration lbs./acre	Percent Stand	
		Non-inoculated	Inoculated
Tetrachloroisophthalonitrile	16	100	52
5-Chloro-2,4,6-trifluoroisophthalonitrile	32	92	68
	16	—	36
	8	—	10

EXAMPLE 16

Soil Mycelial Growth Inhibition Test


The following test measures the ability of compounds of this invention to inhibit mycelial growth in soil. Sterilized soil is infested with *Rhizoctonia solani* (grown on a corn meal sand medium) by mixing two 250-ml. flasks of a ten-day-old culture of the organism per level flat of sterile soil. The infested soil is then placed in small Dixie cups (4-oz. squat). Treatment of the soil is accomplished by drenching the appropriate amount of a diluted formulation containing 0.4 g. of test compound, 8 ml. acetone, 4 ml. stock emulsifier solution (0.5% Triton X-155 in water by volume) and 187.6 ml. distilled water on the surface of the soil in the test cup. The concentration of toxicant in this formulation

is 2000 parts per million. Lower concentrations of toxicant are obtained by diluting the formulation with distilled water. Surface area of the soil in the cups is 5.73 square inches; therefore, a rate of one pound per acre requires 0.414 mg. of chemical per cup. After drenching, the cups are placed in a saturated atmosphere at 70°F. for 48 hours. By this time the fungus mycelium has completely overgrown the surface of the soil in the control cups. Inhibition of mycelial growth in the treated cups is estimated on a scale from zero, complete inhibition of growth, to ten which is equivalent to controls. These grades are expressed as percent control. Using this procedure, the following results are obtained:

Compound	Concentration lbs./acre	Percent Control
Tetrachloroisophthalonitrile	64	90
5-Chloro-2,4,6-trifluoroisophthalonitrile	64	90

WHAT WE CLAIM IS:—

1. Halogenated isophthalonitriles of the structure:



wherein each of X¹, X², X³ and X⁴ is hydrogen or halogen, at least one X being halogen.

2. Tetrachloroisophthalonitrile.

3. 5-Chloro-2,4,6-trifluoroisophthalonitrile.

4. 4,6-Dichloroisophthalonitrile.

5. A method of preparing a ring-chlorinated isophthalonitrile, which comprises reacting the corresponding ring-chlorinated isophthalamide with a dehydrating agent comprising phosphorus pentoxide or phosphorus oxychloride and separating the desired ring-chlorinated isophthalonitrile.

- 5 6. A method of preparing a chloro-fluoro-substituted isophthalonitrile, which comprises reacting a ring-chlorinated isophthalonitrile with a metal fluoride in the presence of a solvent.
7. A method of preparing halogenated isophthalonitriles, as described with reference to the foregoing Examples 4 and 5.
- 10 8. Halogenated isophthalonitriles, when prepared by methods according to claim 5, 6 or 7.
- 15 9. A biologically-active composition, comprising as an essential active ingredient a biologically active amount of a compound having the structure defined in claim 1, a minor amount of a surface-active agent and the balance a carrier.
- 20 10. A method of inhibiting fungus growth which comprises applying to the fungus, foliage or soil containing it a fungicidal amount of a compound according to any of claims 1 to 4 and 8 or a composition according to claim 9.
11. A method of inhibiting the growth of undesirable plants, which comprises applying to the plants or to soil containing them a herbicidal amount of a compound according to any of claims 1 to 4 and 8 or a composition according to claim 9. 25
12. A method of controlling bacteria, which comprises contacting the bacteria with a bactericidal amount of a compound according to any of claims 1 to 4 and 8 or a composition according to claim 9. 30
13. A method of killing nematodes, which comprises applying to the soil a nematocidal amount of a compound according to any of claims 1 to 4 and 8 or a composition according to claim 9. 35

POLLAK, MERCER & TENCH,
Chartered Patent Agents,
Audrey House, Ely Place,
London, E.C.1.
Agents for the Applicants.

Leamington Spa: Printed for Her Majesty's Stationery Office, by the Courier Press (Leamington) Ltd.—1967. Published by The Patent Office, 25 Southampton Buildings, London, W.C.2, from which copies may be obtained.



Pollak Mercer