PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

2000608 4-8
IPDS copy!

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07K 7/56

A1

(11) International Publication Number: WO 99/55727

(43) International Publication Date: 4 November 1999 (04.11.99)

(21) International Application Number: PCT/EP99/02715

(22) International Filing Date: 22 April 1999 (22.04.99)

(30) Priority Data:
98107397.6 23 April 1998 (23.04.98) EP

(71) Applicant *(for all designated States except US)*: HOECHST MARION ROUSSEL DEUTSCHLAND GMBH [DE/DE]; Brüningstrasse 50, D-65929 Frankfurt am Main (DE).

(72) Inventors; and
(75) Inventors/Applicants *(for US only)*: MUKHOPADHYAY, Triptikumar [IN/IN]; M-24, Hoechst Quarters Amar Nagar Mulund (West), Mumbai 400 082 (IN). JAYVANTI, Kenia [IN/IN]; 11, Dev Aashish Ganesh Gawde Road Mulund (West), Mumbai 400 080 (IN). KUMAR, Erra, Koteswara, Satya, Vijaya [IN/IN]; K-3 Hoechst Quarters Amar Nagar Mulund (West), Mumbai 400 080 (IN).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published
With international search report.
Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: A PROCESS FOR THE CONVERSION OF ECHINOCANDIN CLASS OF PEPTIDES TO THEIR C4-HOMOTYROSINE MONODEOXY ANALOGUES

(57) Abstract

The invention relates to a process for the conversion of echinocandin class of peptides to their C4-homotyrosine monodeoxy analogues, particularly mulundocandin to deoxy-mulundocandin, which consists of a single step selective reduction of C4-htyr (homotyrosine) hydroxyl group of echinocandins to their monodeoxy analogues under neutral conditions without prior protection/deprotection of the equally facile C5-Orn (ornithine) hydroxyl group and purification of the monodeoxy compound from the crude reaction mixture.