



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: ⁶ A61K 38/00, 38/16, 38/17, C07K 1/08, 14/435, C12P 21/02	A1	(11) International Publication Number: WO 98/48822 ✓
(21) International Application Number: PCT/US98/06384 ✓	(43) International Publication Date: 5 November 1998 (05.11.98)	(81) Designated States: AL, AM, AU, AZ, BA, BE, BG, BR, BY, CA, CN, CU, CZ, EE, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, RO, RU, SD, <u>SG</u> , SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 24 April 1998 (24.04.98) ✓	(30) Priority Data: ① 60/045,255 ✓ 28 April 1997 (28.04.97) ✓ US ✓	
(71) Applicant (for all designated States except US): ^① ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indi- anapolis, IN 46285 (US). ✓	(72) Inventors; and (75) Inventors/Applicants (for US only): ^① BAKER, Jeffrey, Clayton [US/US]; 114 East 44th Street, Indianapolis, IN 46205 (US). ^② CARLSON, Andrew, David [US/US]; 8026 Castle Cove, Indianapolis, IN 46256 (US). HUANG, Lihua [CN/US]; ^③ 13138 Penneagle Drive, Carmel, IN 46033 (US). SHELIGA ^④ Theodore, Arsay [US/US]; 7901 Venetian Way, Indianapo- lis, IN 46217 (US). ✓	Published With international search report. ✓
(74) Agents: CALTRIDER, Steven, P. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).		1 appl ¹ 4 Invs Inv. 1, address changed, PF 23 redged. chk, correct

(54) Title: IMPROVED METHODS FOR PROCESSING ACTIVATED PROTEIN C

(57) Abstract

The present invention is broadly directed to a method for reducing autodegradation of activated protein C during processing and purification. The present invention provides aqueous activated protein C solutions and an improved method of processing of such solutions, comprising conducting such processing at an ionic strength of greater than 150 mM and at pH of about 5.5 to less than 6.3.

