

1705 Copy
9806228-2

THIS SHEET HAS BEEN
REPLACED


-1-

THIOXANTHENE ANTITUMOR AGENTS

The present invention relates to novel
1-[[[(dialkylamino)-alkyl]amino]-4-substituted-
thioxanthen-9-ones, to pharmaceutical compositions
5 containing the thioxanthenones, to methods of
treating tumors with the thioxanthenones and to
methods of treating cancer in mammals with the
compositions containing the thioxanthenones.

Rosi U.S. Patent 3,577,558 discloses 1-[[2-
10 (diethylamino)ethyl]amino]-4-(alkoxymethyl)-
thioxanthen-9-one ethers.

15




R = methyl, ethyl,
n-propyl, isopropyl
and n-hexyl

20

The compounds are stated to be schistosomacidal
agents.

Blanz and French [J. Med. Chem. 6, 185-191 (1963)]
disclose the synthesis of a series of thioxanthenones
related to lucanthone and the results of the testing
25 of the compounds against a leukaemia and two solid
tumors. Among the compounds disclosed are


30




wherein R is methyl, methoxyl, and ethoxyl.

35

This Sheet Has Been Replaced



The present invention relates to compounds of the formula I



10 or acid addition salt or solvate thereof wherein n is two or three, preferably two; R¹ and R² are independently lower-alkyl, preferably both are ethyl; R³ is lower-alkyl and R⁴ is hydrogen, lower-alkyl, lower-alkoxy or halogen. The compounds are useful for
15 the treatment of tumors in mammals.

Lower-alkyl as used herein describes linear, branched or cyclic hydrocarbons containing four or fewer carbon atoms. Halogen describes bromine, chlorine or fluorine.

20 The invention further relates to compositions for treating tumors and cancer in mammals which comprise compounds of formula I together with pharmaceutically acceptable excipients or diluents.

25 The invention further relates to a method of use of a compound of formula I for treating tumors in mammals which comprises administering to the mammal a compound of formula I.


30 The invention further relates to a method of use of a compound of formula I for treating cancer in a mammal which comprises administering to the mammal a composition of a compound of formula I together with pharmaceutically acceptable excipients or diluents.

35 The synthesis of compounds of the invention may be outlined as shown in Scheme A:




-3-


SCHEME A



sulfonylating agent
or
halogenating agent



$R^3S^- M^+$



The compounds of formula I may be synthesized by reacting the appropriate alcohol (II) with about one equivalent of an appropriate sulfonylating agent, for example p-toluenesulfonyl chloride, methanesulfonyl chloride, trifluoromethanesulfonyl chloride and the like, preferably p-toluenesulfonyl chloride, in pyridine at 0° to 80°, preferably about 25°, to produce a sulfonate ester (III, X = p-toluenesulfonate, methanesulfonate, trifluoromethanesulfonate and the like), which is not isolated but, rather, is reacted immediately with an excess, preferably about 5-7 fold excess, of an alkali metal salt of the appropriate thiol. The preferred salt is the sodium salt and the reaction is carried out in a suitable solvent, preferably DMSO.

Alternatively, the alcohol (II) can be converted to the corresponding halide (III, X = Cl, Br, or I) by treatment with a phosphorous trihalide, phosphorous pentahalide, thionyl halide or tetrahalomethane with trialkylphosphine. The group X can then be displaced by reaction with an alkali metal salt of the appropriate thiol as described above.

The appropriate alcohols may be obtained by procedures well known in the art, particularly as described in U.S. Patents 3,711,512 and 3,294,803 which are incorporated herein by reference.

The compounds of formula I are useful both in the free base form and in the form of acid-addition salts, and both forms are within the purview of the invention. The acid-addition salts are in some cases a more convenient form for use, and in practice the use of the salt form inherently amounts to the use of the base form. The acids which can be used to prepare the acid-addition salts include preferably those which produce, when



combined with the free base, medicinally acceptable salts, that is, salts whose anions are relatively innocuous to the animal organism in medicinal doses of the salts so that the beneficial properties n
5 the free base are not vitiated by side effects ascribable to the anions. In practising the present invention it is convenient to form the hydrochloride, fumarate, toluenesulfonate, methanesulfonate, or maleate salts. However, other
10 appropriate medicinally acceptable salts within the scope of the invention are those derived from other mineral acids and organic acids. The acid-addition salts of the basic compounds are prepared either by dissolving the free base in aqueous alcohol
15 solution containing the appropriate acid and isolating the salt by evaporating the solution, or by reacting the free base and an acid in an organic solvent, in which case the salt separates directly, is precipitated with a second organic solvent, or
20 can be obtained by concentration of the solution. Although medicinally acceptable salts of the basic compounds are preferred, all acid-addition salts are within the scope of the present invention. All acid-addition salts are useful as sources of the
25 free base form even if the particular salt per se is desired only as an intermediate product, as, for example, when the salt is formed only for purposes of purification or identification, or when it is used as an intermediate in preparing a medicinally
30 acceptable salt by ion exchange procedures.

The structures of the compounds of the invention were established by the mode of synthesis, by elemental analysis, and by infrared, ultraviolet and nuclear magnetic resonance spectroscopy. The course
35 of the reactions and the identity and homogeneity of



the products were assessed by thin layer chromatography (TLC) or gas-liquid chromatography (GLC). The melting points are given in degrees C and are uncorrected. The starting materials are either
5 commercially available or may be prepared by procedures well known in the art.

The pharmaceutical compositions of the present invention include one or more of the compounds of this invention formulated into compositions together with
10 one or more non-toxic physiologically acceptable carriers, adjuvants or vehicles which are collectively referred to herein as carriers, for parenteral injection, for oral administration in solid or liquid form, for rectal or topical administration, and the
15 like.

The compositions can be administered to humans and animals either orally, rectally, parenterally (intravenously, intramuscularly or subcutaneously), intracisternally, intravaginally, intraperitoneally,
20 locally (powders, ointments or drops), or as a buccal or nasal spray.

Compositions suitable for parenteral injection may comprise physiologically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or
25 emulsions and sterile powders for reconstitution into sterile injectable solutions or dispersions. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (propylene glycol, polyethyleneglycol, glycerol, and
30 the like), suitable mixtures thereof, vegetable oils (such as olive oil) and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in
35 the case of dispersions and by the use of surfactants.



-7-

These compositions may also contain adjuvants such as preserving, wetting, emulsifying, and dispensing agents. Prevention of the action of microorganisms can be ensured by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, for example sugars, sodium chloride and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin.

If desired, and for more effective distribution, the compounds can be incorporated into slow release or targeted delivery systems such as polymer matrices, liposomes, and microspheres. They may be sterilized, for example, by filtration through a bacteria-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved in sterile water, or some other sterile injectable medium immediately before use.

Solid dosage forms for oral administration include capsules, tablets, pills, powders and granules. In such solid dosage forms, the active compound is admixed with at least one inert customary excipient (or carrier) such as sodium citrate or dicalcium phosphate or (a) fillers or extenders, as for example, starches, lactose, sucrose, glucose, mannitol and silicic acid, (b) binders, as for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose and acacia, (c) humectants, as for example, glycerol, (d) disintegrating agents, as for example, agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain complex silicates and sodium carbonate, (e) solution retarders, as for example, paraffin,



(f) absorption accelerators, as for example, quaternary ammonium compounds, (g) wetting agents, as for example, cetyl alcohol and glycerol monostearate, (h) adsorbents, as for example, kaolin and bentonite, and (i) lubricants, as for example, talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate or mixtures thereof. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents.

The percentage of active component in the composition and method for treating tumors or cancer can be varied so that a suitable dosage is obtained. The dosage administered to a particular patient is variable depending upon the clinician's judgement using as the criteria: the route of administration, the duration of treatment, the size and condition of the patient, the potency of the active component, and the patient's response thereto. An effective dosage amount of active component can thus be determined by the clinician considering all criteria and utilizing his best judgement on the patient's behalf.


The invention will now be more particularly described with reference to the following Example which in no way limits the scope of the invention.

Example 1

1-[[2-(Diethylamino)ethyl]amino]-4-(methylthiomethyl)
thioxanthen-9-one

(I: $R^1=R^2=Et$; $R^3=$ methyl; $R^4=H$; $n=2$)

A mixture of 17.9 g (50 mmol) of 1-[[2-(diethylamino)-ethyl]amino]thioxanthen-4-methanol and 9.6 g (50 mmol) of p-toluenesulfonyl chloride in 125 ml of pyridine was stirred for 2 hours at room temperature. The reaction was diluted with 100 ml of DMSO and the pyridine was removed on a



rotary evaporator. A slurry of 300 mmol of the sodium salt of methanethiol in 100 ml of DMSO, preformed from 16.8 g of methanethiol and 12.0 g of 60% NaH in
5 mineral oil, was added. The reaction was stirred for 2 hours and poured into 1.5l of water. The resulting gum was removed, dissolved in chloroform, washed with 5% aqueous NaOH, dried over Na₂SO₄, filtered and
10 stripped. The residue was digested with ethyl acetate and filtered to remove unreacted starting material. The solution was chromatographed on alumina eluting with a gradient from 0 to 1% isopropylamine in 95:5 pentane/methylene chloride then 2% isopropylamine in 90:10 pentane/methylene chloride to obtain 7 g of
15 product which was recrystallized from ethyl acetate to yield 6.6 g of product, mp 125-127°.

Other representative members of the genus I may be made by a method analogous to that of Example 1, substituting the appropriate 1-[[2-(dialkyl-
20 amino)ethyl]amino]- or 1-[[3-(dialkylamino)-propyl]amino]thioxanthen-4-methanol for 1-[[2-(diethylamino)-ethyl]amino]thiioxanthen-4-methanol and the appropriate alkylthiol for methanethiol.

25 A representative example of the invention was tested for antitumor activity in mice according to the following procedure:

The animals were pooled, implanted subcutaneously
30 with 30 to 60 mg tumor fragments by 12-gauge trocar, and again pooled before unselective distribution to the various treatment and control groups. For early-stage treatment, chemotherapy was started within 1 to 5 days after tumor implantation while the number of
35 cells was relatively small (10^7 to 10^8 cells). For



THIS SHEET HAS BEEN
REPLACED

advanced-stage treatment, chemotherapy was delayed until the tumor became relatively large (200 to 300 mg in size). A 300-mg tumor contains approximately 3×10^8 total cells. Tumors within a given advanced-stage trial were within a 2.5-fold size range for 90% of the animals. Tumors were measured with a caliper weekly (or twice weekly for the more rapidly growing tumors). Mice were sacrificed when their tumors reached 1500 mg (i.e., before they could cause the animal discomfort). Tumor weights were estimated from two-dimensional measurements.

The treatment and control groups were measured when the control group tumors reached approximately 700 to 1200 mg in size (Median of Group). The median tumor weight of each group was determined (including zeros). The T/C value (weight of treated tumours over the weight of control tumours) in percent is an indication of antitumor effectiveness: A T/C equal to or less than 42% is considered significant antitumor activity by the Drug Evaluation Branch of the Division of Cancer Treatment (NCI). A T/C value <10% is considered to indicate highly significant antitumor activity. A body weight loss nadir (mean of group) of greater than 20% or greater than 20% drug-deaths is considered to indicate an excessively toxic dosage.

The results are shown in Table I for pancreatic ductal adenocarcinoma #03.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

TABLE 1

5	Example #	T/C	Weight	Drug	Total
		(%)	Loss		Dose
			(g)*	Deaths	(mg/kg)
					i.v.
	1	0	2.0	0	1830

*The average body weight was 25 g.

10

15

20

25

30

35

This Sheet Has Been Replaced






* C L M S *

- 12 -

THIS SHEET HAS BEEN
REPLACED

CLAIMS:

- 5 1. A compound of formula



wherein n is 2 or 3;

- 15 R^1 and R^2 are independently lower-alkyl;
 R^3 is lower-alkyl; and
 R^4 is hydrogen, lower-alkyl, lower-alkoxy or
halogen.

- 20 2. A compound according to claim 1 wherein n is
2 and both of R^1 and R^2 are ethyl, and R^4 is hydrogen.

3. 1-[[2-(Diethylamino)ethyl]amino]-4-
[(methylthio)methyl]-thioxanthen-9-one.

- 25 4. A pharmaceutical composition which comprises
a compound of claim 1 and a pharmaceutically
acceptable carrier or diluent.

5. A pharmaceutical composition which comprises
the compound of claim 3 and a pharmaceutically
acceptable carrier or diluent.

- 30 6. The use of a compound of claim 1 for the
preparation of a medicament for treating a tumor in a
mammal.

7. The use of a compound of claim 3 for the
preparation of a medicament for treating a tumor in a
mammal.

35

This Sheet Has Been Replaced





- 13 -

8. The use of a composition of claim 4 for the preparation of a medicament for treating cancer in a mammal.

5

9. The use of a composition of claim 5 for the preparation of a medicament for treating cancer in a mammal.

10

THIS SHEET HAS BEEN
REPLACED

15

20

25

30

35

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

ABSTRACT

THIOXANTHENONE ANTITUMOR AGENTS

5 1-[[(Dialkylamino)alkyl]amino]-4-substituted-
thioxanthen-9-ones are disclosed as antitumor agents.
Compositions containing the thioxanthenones and the
method of use of thioxanthenones for treating tumors and
cancer in mammals are also disclosed.

10

15

20

25

30

35

This Sheet Has Been Replaced



12 FEB 1998


-1-

THIOXANTHENONE ANTITUMOR AGENTS

Description

The present invention relates to novel 1-[[[(dialkylamino)-alkyl]amino]-4-substituted-thioxanthen-9-ones, to pharmaceutical compositions containing the thioxanthenones, to the use of thioxanthenones for the preparation of pharmaceuticals for the treatment of tumors and cancer in mammals.


Rosi U.S. Patent 3,577,558 discloses 1-[2-(diethylamino)ethyl]amino]-4-(alkoxymethyl)-thioxanthen-9-one ethers.



R = methyl, ethyl, n-propyl, isopropyl and n-hexyl


The compounds are stated to be schistosomacidal agents.

Blanz and French [*J. Med. Chem.* 6, 185-191 (1963)] disclose the synthesis of a series of thioxanthenones related to lucanthone and the results of the testing of the compounds against a leukaemia and two solid tumors. Among the compounds disclosed are



wherein R is methyl, methoxyl, and ethoxyl.

The present invention relates to compounds of the formula I



or acid addition salt or solvate thereof wherein n is two or three, preferably two; R¹ and R² are independently lower-alkyl, preferably both are ethyl; R³ is lower-alkyl and R⁴ is hydrogen, lower-alkyl, lower-alkoxy or halogen. The compounds are useful for the treatment of tumors in mammals.

Lower-alkyl as used herein describes linear, branched or cyclic hydrocarbons containing four or fewer carbon atoms. Halogen describes bromine, chlorine or fluorine.

As preferred compounds of formula I, those in which n is 2 and both of R¹ and R² are ethyl, and R⁴ is hydrogen are preferred.

The invention further relates to compositions for treating tumors and cancer in mammals which comprise compounds of formula I together with pharmaceutically acceptable excipients or diluents.


The compounds of the invention may be used for the preparation of pharmaceuticals intended for the treatment of tumors or cancer in mammals.

The synthesis of compounds of the invention may be outlined as shown in Scheme A:

SCHEME A

REPLACEMENT SHEET

12 FEB 1998



The compounds of formula I may be synthesized by reacting the appropriate alcohol (II) with about one equivalent of an appropriate sulfonylating agent, for example p-toluenesulfonyl chloride, methanesulfonyl chloride, trifluorometh-

12 FEB 1998

- 3 -

anesulfonyl chloride and the like, preferably p-toluenesulfonyl chloride, in pyridine at 0° to 80°, preferably about 25°, to produce a sulfonate ester (III, X = p-toluenesulfonate, methanesulfonate, trifluoromethanesulfonate and the like), which is not isolated but, rather, is reacted immediately with an excess, preferably about 5-7 fold excess, of an alkali metal salt of the appropriate thiol. The preferred salt is the sodium salt and the reaction is carried out in a suitable solvent, preferably DMSO.

Alternatively, the alcohol (II) can be converted to the corresponding halide (III, X = Cl, Br, or I) by treatment with a phosphorous trihalide, phosphorous pentahalide, thionyl halide or tetrahalomethane with trialkylphosphine. The group X can then be displaced by reaction with an alkali metal salt of the appropriate thiol as described above.

The appropriate alcohols may be obtained by procedures well known in the art, particularly as described in U.S. Patents 3,711,512 and 3,294,803 which are incorporated herein by reference.

The compounds of formula I are useful both in the free base form and in the form of acid-addition salts, and both forms are within the purview of the invention. The acid-addition salts are in some cases a more convenient form for use, and in practice the use of the salt form inherently amounts to the use of the base form. The acids which can be used to prepare the acid-addition salts include preferably those which produce, when combined with the free base, medicinally acceptable salts, that is, salts whose anions are relatively innocuous to the animal organism in medicinal doses of the salts so that the beneficial properties of the free base are not vitiated by side effects ascribable to the anions. In practicing the present invention it is convenient to form the hydrochloride, fumarate, toluenesulfonate, methanesulfonate, or maleate salts. However, other appropriate medicinally acceptable salts within the scope of the invention are those derived from other mineral acids and organic acids. The acid-addition salts of the basic compounds are prepared either by dissolving the free base in aqueous alcohol solution containing the appropriate acid and isolating the salt by evaporating the solution, or by reacting the free base and an acid in an organic solvent, in which case the salt separates directly, is precipitated with a second organic solvent, or can be obtained by concentration of the solution. Although medicinally acceptable salts of the basic compounds are preferred, all acid-addition salts are within the scope of the present invention. All acid-addition salts are useful as sources of the free base form even if the particular salt *per se* is desired only as an intermediate product, as, for example, when the salt is formed only for purposes of purification or identification, or when it is used as an intermediate in preparing a medicinally acceptable salt by ion exchange procedures.

The structures of the compounds of the invention were established by the mode of synthesis, by elemental analysis, and by infrared, ultraviolet and nuclear magnetic resonance spectroscopy. The course of the reactions and the identity and homogeneity of the products were assessed by thin layer chromatography (TLC) or gas-liquid chromatography (GLC). The melting points are given in degrees C and are uncorrected. The starting materials are either commercially available or may be prepared by procedures well known in the art.

The pharmaceutical compositions of the present invention include one or more of the compounds of this invention formulated into compositions together with one or more non-toxic physiologically acceptable carriers, adjuvants or vehicles which are collectively referred to herein as carriers, for parenteral injection, for oral administration in solid or liquid form, for rectal or topical administration, and the like.

The compositions can be administered to humans and animals either orally, rectally, parenterally (intravenously, intramuscularly or subcutaneously), intracisternally, intravaginally, intraperitoneally, locally (powders, ointments or drops), or as a buccal or nasal spray.

Compositions suitable for parenteral injection may comprise physiologically acceptable sterile aqueous or non-aqueous solutions, dispersions, suspensions or emulsions and sterile powders for reconstitution into sterile injectable solutions or dispersions. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (propylene glycol, polyethyleneglycol, glycerol, and the like), suitable mixtures thereof, vegetable oils (such as olive oil) and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants.

These compositions may also contain adjuvants such as preserving, wetting, emulsifying, and dispensing agents. Prevention of the action of microorganisms can be ensured by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, for example sugars, sodium chloride and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin.

If desired, and for more effective distribution, the compounds can be incorporated into slow release or targeted delivery systems such as polymer matrices, liposomes, and microspheres. They may be sterilized, for example, by filtration through a bacteria-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved in sterile water, or some other sterile injectable medium immediately before use.

Solid dosage forms for oral administration include capsules, tablets, pills, powders and granules. In such solid dosage forms, the active compound is admixed with at least one inert customary excipient (or carrier) such as sodium citrate or dicalcium phosphate or (a) fillers or extenders, as for example, starches, lactose, sucrose, glucose, mannitol and silicic acid, (b) binders, as for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose and

12 FEB 1998

- 4 -

acacia,

(c) humectants, as for example, glycerol,

(d) disintegrating agents, as for example, agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain complex silicates and sodium carbonate,

(e) solution retarders, as for example, paraffin,

(f) absorption accelerators, as for

example, quaternary ammonium compounds, (g) wetting agents, as for example, cetyl alcohol and glycerol monostearate, (h) adsorbents, as for example, kaolin and bentonite, and (i) lubricants, as for example, talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate or mixtures thereof. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents.

The percentage of active component in the composition of the invention can be varied so that a suitable dosage is obtained. The dosage administered to a particular patient is variable depending upon the clinician's judgement using as the criteria: the route of administration, the duration of treatment, the size and condition of the patient, the potency of the active component, and the patient's response thereto. An effective dosage amount of active component can thus be determined by the clinician considering all criteria and utilizing his best judgement on the patient's behalf.

The invention will now be more particularly described with reference to the following Example which in no way limits the scope of the invention.

Example 1

1-[[2-(Diethylamino)ethyl]amino]-4-(methylthiomethyl) thioxanthen-9-one

(1: $R^1 = R^2 = Et$; $R^3 = methyl$; $R^4 = H$; $n=2$)

A mixture of 17.9 g (50 mmol) of 1-[[2-(diethylamino)-ethyl]amino]thioxanthen-4-methanol and 9.6 g (50 mmol) of p-toluenesulfonyl chloride in 125 ml of pyridine was stirred for 2 hours at room temperature. The reaction was diluted with 100 ml of DMSO and the pyridine was removed on a rotary evaporator. A slurry of 300 mmol of the sodium salt of methanethiol in 100 ml of DMSO, prepared from 16.8 g of methanethiol and 12.0 g of 60% NaH in mineral oil, was added. The reaction was stirred for 2 hours and poured into 1.5 l of water. The resulting gum was removed, dissolved in chloroform, washed with 5% aqueous NaOH, dried over Na_2SO_4 , filtered and stripped. The residue was digested with ethyl acetate and filtered to remove unreacted starting material. The solution was chromatographed on alumina eluting with a gradient from 0 to 1% isopropylamine in 95:5 pentane/methylene chloride then 2% isopropylamine in 90:10 pentane/methylene chloride to obtain 7 g of product which was recrystallized from ethyl acetate to yield 6.6 g of product, mp 125-127°.

Other representative members of the genus I may be made by a method analogous to that of Example 1, substituting the appropriate 1-[[2-(dialkylamino)ethyl]amino]- or 1-[[3-(dialkylamino)propyl]amino]thioxanthen-4-methanol for 1-[[2-(diethylamino)-ethyl]amino]thioxanthen-4-methanol and the appropriate alkylthiol for methanethiol.

A representative example of the invention was tested for antitumor activity in mice according to the following procedure:

The animals were pooled, implanted subcutaneously with 30 to 60 mg tumor fragments by 12-gauge trocar, and again pooled before unselective distribution to the various treatment and control groups. For early-stage treatment, chemotherapy was started within 1 to 5 days after tumor implantation while the number of cells was relatively small (10^7 to 10^8 cells). For advanced-stage treatment, chemotherapy was delayed until the tumor became relatively large (200 to 300 mg in size). A 300-mg tumor contains approximately 3×10^8 total cells. Tumors within a given advanced-stage trial were within a 2.5-fold size range for 90% of the animals. Tumors were measured with a caliper weekly (or twice weekly for the more rapidly growing tumors). Mice were sacrificed when their tumors reached 1500 mg (i.e., before they could cause the animal discomfort). Tumor weights were estimated from two-dimensional measurements.


The treatment and control groups were measured when the control group tumors reached approximately 700 to 1200 mg in size (Median of Group). The median tumor weight of each group was determined (including zeros). The T/C value (weight of treated tumors over the weight of control tumors) in percent is an indication of antitumor effectiveness: A T/C equal to or less than 42% is considered significant antitumor activity by the Drug Evaluation Branch of the Division of Cancer Treatment (NCI). A T/C value <10% is considered to indicate highly significant antitumor activity. A body weight loss nadir (mean of group) of greater than 20% or greater than 20% drug-deaths is considered to indicate an excessively toxic dosage.

The results are shown in Table I for pancreatic ductal adenocarcinoma #03.

TABLE 1

Example #	T/C (%)	Weight Loss (g)*	Drug Deaths	Total Dose (mg/kg) i.v.
1	0	2.0	0	1830

*The average body weight was 25 g.





- 14 -


REPLACEMENT SHEET

12 FEB 1998

Claims

Claims for the following Contracting States : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, MC, NL, PT, SE

1. A compound of formula



wherein

n is 2 or 3;
R¹ and R² are independently lower-alkyl;
R³ is lower-alkyl; and
R⁴ is hydrogen, lower-alkyl, lower-alkoxy or halogen, it being understood that lower-alkyl designates linear, branched or cyclic hydrocarbons containing four or fewer carbon atoms.

- A compound according to claim 1 wherein n is 2 and both of R¹ and R² are ethyl, and R⁴ is hydrogen.
- 1-[[2-(Diethylamino)ethyl]amino]-4-[(methylthio)methyl]-thioxanthene-9-one.
- A pharmaceutical composition which comprises a compound of claim 1 and a pharmaceutically acceptable carrier or diluent.
- A pharmaceutical composition which comprises the compound of claim 3 and a pharmaceutically acceptable carrier or diluent.
- The use of a compound of claim 1 for the preparation of a medicament for treating a tumor in a mammal.
- The use of a compound of claim 3 for the preparation of a medicament for treating a tumor in a mammal.
- The use of a composition of claim 4 for the preparation of a medicament for treating cancer in a mammal.
- The use of a composition of claim 5 for the preparation of a medicament for treating cancer in a mammal.

Claims for the following Contracting States : ES, GR

1. A process for preparing a compound having the formula



THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced






This Sheet Has Been Replaced



wherein


- n is 2 or 3;
 R^1 and R^2 are independently lower-alkyl;
 R^3 is lower-alkyl; and
 R^4 is hydrogen, lower-alkyl, lower-alkoxy or halogen, it being understood that lower-alkyl designates linear, branched or cyclic hydrocarbons containing four or fewer carbon atoms;

which comprises reacting a compound of the formula III



with a compound of the formula $R^3S^+M^-$, wherein M is an alkali metal and X is a suitable leaving group, so as to obtain the corresponding compound of formula I.

2. A process according to claim 1 which comprises the additional step of reacting a compound of the formula II



with an agent capable of converting the hydroxyl group to a suitable leaving group to prepare the corresponding compound of formula III.

3. A process according to claim 2 wherein X is a p-toluenesulfonate ester and M is sodium.
4. A process according to claim 3 wherein n is 2 and both of R^1 and R^2 are ethyl, and R^4 is hydrogen.
5. A process according to claim 4, for preparing 1-[[2-(diethylamino)ethyl]amino]-4-[[methylthio)methyl]-thioxanthene-9-one.

THIS SHEET HAS BEEN
REPLACED



-6-


REPLACEMENT SHEET

17 AUG 1998

Claims

Claims for the following Contracting States : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, MC, NL, PT, SE

1. A compound of formula



wherein

- n is 2 or 3;
R¹ and R² are independently lower-alkyl;
R³ is lower-alkyl; and
R⁴ is hydrogen, lower-alkyl, lower-alkoxy or halogen, it being understood that lower-alkyl designates linear, branched or cyclic hydrocarbons containing four or fewer carbon atoms.

2. A compound according to claim 1 wherein n is 2 and both of R¹ and R² are ethyl, and R⁴ is hydrogen.
3. 1-[[2-(Diethylamino)ethyl]amino]-4-[(methylthio)methyl]-thioxanthene-9-one.
4. A pharmaceutical composition which comprises a compound of claim 1 and a pharmaceutically acceptable carrier or diluent.
5. A pharmaceutical composition which comprises the compound of claim 3 and a pharmaceutically acceptable carrier or diluent.
6. The use of a compound of claim 1 for the preparation of a medicament for treating a tumor in a mammal.
7. The use of a compound of claim 3 for the preparation of a medicament for treating a tumor in a mammal.
8. The use of a composition of claim 4 for the preparation of a medicament for treating cancer in a mammal.
9. The use of a composition of claim 5 for the preparation of a medicament for treating cancer in a mammal.

Claims for the following Contracting States : ES, GR

1. A process for preparing a compound having the formula

This Sheet Has Been Replaced





THIS SHEET HAS BEEN
REPLACED



-7-


This Sheet Has Been Replaced



wherein


- n is 2 or 3;
 R^1 and R^2 are independently lower-alkyl;
 R^3 is lower-alkyl; and
 R^4 is hydrogen, lower-alkyl, lower-alkoxy or halogen, it being understood that lower-alkyl designates linear, branched or cyclic hydrocarbons containing four or fewer carbon atoms;

which comprises reacting a compound of the formula III



with a compound of the formula R^3SM^+ , wherein M is an alkali metal and X is a suitable leaving group, so as to obtain the corresponding compound of formula I.

2. A process according to claim 1 which comprises the additional step of reacting a compound of the formula II



with an agent capable of converting the hydroxyl group to a suitable leaving group to prepare the corresponding compound of formula III.

3. A process according to claim 2 wherein X is a p-toluenesulfonate ester and M is sodium.
4. A process according to claim 3 wherein n is 2 and both of R^1 and R^2 are ethyl, and R^4 is hydrogen.
5. A process according to claim 4, for preparing 1-[[2-(diethylamino)ethyl]amino]-4-[(methylthio)methyl]-thioxanthene-9-one.

THIS SHEET HAS BEEN
REPLACED

ABSTRACT

THIOXANTHENONE ANTITUMOR AGENTS

1-[[(Dialkylamino)alkyl]amino]-4-substituted-thioxanthen-9-ones are disclosed as antitumor agents. Compositions containing the thioxanthenones and the method of use of thioxanthenones for treating tumors and cancer in mammals are also disclosed.

REPLACEMENT SHEET

17 AUG 1998

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED




REPLACEMENT SHEET

02 OCT 1999

Claims

1. A compound of formula



wherein

n is 2 or 3;
R¹ and R² are independently lower-alkyl;
R³ is lower-alkyl; and
R⁴ is hydrogen, lower-alkyl, lower-alkoxy or halogen, it being understood that lower-alkyl designates linear, branched or cyclic hydrocarbons containing four or fewer carbon atoms.

2. A compound according to claim 1 wherein n is 2 and both of R¹ and R² are ethyl, and R⁴ is hydrogen.
3. 1-[[2-(Diethylamino)ethyl]amino]-4-[(methylthio)methyl]-thioxanthene-9-one.
4. A pharmaceutical composition which comprises a compound of claim 1 and a pharmaceutically acceptable carrier or diluent.
5. A pharmaceutical composition which comprises the compound of claim 3 and a pharmaceutically acceptable carrier or diluent.
6. The use of a compound of claim 1 for the preparation of a medicament for treating a tumor in a mammal.
7. The use of a compound of claim 3 for the preparation of a medicament for treating a tumor in a mammal.
8. The use of a composition of claim 4 for the preparation of a medicament for treating cancer in a mammal.
9. The use of a composition of claim 5 for the preparation of a medicament for treating cancer in a mammal.

-7-

ABSTRACT

THIOXANTHENONE ANTITUMOR AGENTS

1-[[(Dialkylamino)alkyl]amino]-4-substituted-thioxanthen-9-ones are disclosed as antitumor agents. Compositions containing the thioxanthenones and the method of use of thioxanthenones for treating tumors and cancer in mammals are also disclosed.

REPLACEMENT SHEET

02 OCT 1999