# Methods for the identification of LRRK2 interacting molecules and for the purification of LRRK2

The present invention relates to methods for the identification of LRRK2 interacting molecules and for the purification of LRRK2 using indol ligand 91 as a ligand for LRRK2. Furthermore, the present invention relates to pharmaceutical compositions comprising said interacting molecules e.g. for the treatment of Parkinson's disease.

Parkinson's disease (PD) [OMIM168600, Online Mendelian Inheritance in Man (OMM), http://www.ncbi.nlm.nih.gov/entrez/dispomim.cgi?id=168600] is a heterogeneous movement disorder characterized by the degeneration of dopaminergic neurons within the substancia nigra of the basal ganglia. It affects 2% of the population over 60 years.

Current therapeutic strategies for PD focus primarily on reducing the severity of its symptoms using dopaminergic medications (e.g. Levodopa/Carbidopa). Although providing benefit to patients, these medications display adverse side effects and may become ineffective after prolonged treatment (wearing off effect). None of these treatments addresses the underlying problem, the progressive loss of dopaminergic neurons.

Broadly, there are two clinicopathological components of PD. First, there is clinically defined parkinsonism, a syndromic term which comprises the cardinal features of the Parkinsonian movement disorder. These are a resting tremor, bradykinesia (slowness of movement), rigidity and postural instability, all problems in initiating or stopping movements. Their pathological correlate is the loss of dopaminergic neurons in the substancia nigra. Second, Parkinson's disease is marked postmortem by the presence of Lewy bodies and Lewy neurites in surviving neurons. These are intracellular aggregations of lipids and proteins including ubiquitin and alpha-synuclein. Pathological definitions of PD require the presence of alpha-synuclein-positive Lewy pathology in surviving nigral neurons, combined with nigral cell loss and intact striatal neurons (Cookson, 2005. Annual Reviews in Biochemistry 74, 29-52).

The alpha-synuclein-centric theory of protein aggregation has in recent years been complemented by the discovery of the proteins DJ1, PINK1 and OMI/HTRA2, all of which are associated with the mitochondria and have been implicated in cellular protection against oxidative damage (Abou-Sleiman et al., 2006. Nature Reviews Neuroscience 7, 207-219).

Genetic evidence has recently been presented that mutations of the Leucine-rich repeat kinase 2 (LRRK2, synonym Dardarin) cause autosomal dominant PD previously linked to the PARK8 locus. LRRK2 mutations are estimated to account for 5 to 6% of PD cases with a positive family history, and were also identified in sporadic cases. LRRK2 encodes a large multi-domain protein that consists of N-terminal leucine-rich repeats, a GTPase ROC/COR domain, a mitogen-activated protein kinase kinase kinase (MAPKKK) and C-terminal WD40 repeats (Paisan-Ruiz et al., 2004. Neuron 44, 595-600; Zimprich et al., 2004. Neuron 44, 601-607). Paisan-Ruz and colleagues named the protein product dardarin, derived from the Basque word "dardara" meaning tremor, but the protein was later renamed LRRK2.

Currently, little is known about LRRK2 function but preliminary observations have been made in *in vitro* overexpression systems. LRRK2 encodes a protein kinase and is capable of autophosphorylation (West et al., 2005. PNAS 102, 16842-16847, Gloeckner, et al., 2005. Hum. Mol. Genet. 15, 223-232). The publication West et al. describes the purification of recombinant LRRK2 with the help of a specific antibody. However, the amount of protein obtained was very small.

Significantly, three PD-associated LRRK2 mutations, two in the kinase domain (G2019S and I2020T) and one in the ROC/COR GTPase domain (R1441C) increase LRRK2 autophosphorylation, suggesting a dominant gain-of-function mechanism. Indeed, overexpression of R1441C, Y1699C or G2019S mutants of LRRK2 is sufficient to induce neuronal degeneration in mouse primary cortical neurons (Smith et al., 2005. PNAS 102, 18676-18681).

Moreover, manipulating the kinase activity of LRRK2 by replacing the kinase domain with a "kinase-dead" mutated version blocks the formation of inclusion bodies and delays cell death, two cellular phenotypes of PD (Greggio et al., 2006. Neurobiol. Dis. 23(2), 329-341). This observation suggets that kinase inhibitors will be useful therapeutic agents for patients with LRRK2 mutations and presuambly also sporadic PD.

The recent discovery that LRRK2 mutations lead to late-onset PD with pleomorphic pathology, including alpha-synuclein, tau and ubiquitin pathology places LRRK2 upstream in a common neurodegenerative pathway for parkinsonism. Thus, LRRK2 has become an attractive therapeutic target for intervention and neuroprotection in Parkinson's disease (Taylor et al., 2006. Trends in Molecular Medicine 12, 76-82).

In view of the above, there is a need for providing effective methods for the identification of LRRK2 interacting compounds as well as for methods for the purification of LRRK2.

In a first aspect, there is provided a method for the identification of an LRRK2 interacting compound, comprising the steps of

a) providing a protein preparation containing LRRK2,
b) contacting the protein preparation with (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide) immobilized on a solid support under conditions allowing the formation of an (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex,
c) incubating the (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex with a given compound, and
d) determining whether the compound is able to separate LRRK2 from the immobilized (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide).

In a second aspect, there is provided a method for the identification of an LRRK2 interacting compound, comprising the steps of

a) providing a protein preparation containing LRRK2,
b) contacting the protein preparation with (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide) immobilized on a solid support and with a given compound under conditions allowing the formation of an (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex, and
c) detecting the (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex formed in step b).

In a third aspect, there is provided a method for the identification of an LRRK2 interacting compound, comprising the steps of:

a) providing two aliquots of a protein preparation containing LRRK2,
b) contacting one aliquot with (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide) immobilized on a solid support under conditions allowing the formation of an (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex,
c) contacting the other aliquot with (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide) immobilized on a solid support and with a given compound under conditions allowing the formation of an (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex, and
d) determining the amount of (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex formed in steps b) and c).

In a fourth aspect, there is provided a method for the purification of LRRK2, comprising the steps of

a) providing a protein preparation containing LRRK2,
b) contacting the protein preparation with (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide) immobilized on a solid support under conditions allowing the formation of an (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex, and
c) separating LRRK2 from the immobilized (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide).

In a fifth aspect, there is provided the use of (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide) for the identification of LRRK2 interacting compounds.

In a sixth aspect, there is provided the

use of (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide) for the purification of LRRK2.

Disclosed herein is a method for the identification of an LRRK2 interacting compound, comprising the steps of

a) providing a protein preparation containing LRRK2,

b) contacting the protein preparation with indol ligand 91 immobilized on a solid support under conditions allowing the formation of an indol ligand 91-LRRK2 complex,

c) incubating the indol ligand 91-LRRK2 complex with a given compound, and

d) determining whether the compound is able to separate LRRK2 from the immobilized indol ligand 91.

Furthermore, also disclosed is a method for the identification of an LRRK2 interacting compound, comprising the steps of

a) providing a protein preparation containing LRRK2,

b) contacting the protein preparation with indol ligand 91 immobilized on a solid support and with a given compound under conditions allowing the formation of an indol ligand 91-LRRK2 complex, and

c) detecting the indol ligand 91-LRRK2 complex formed in steps b).

Also disclosed is a method for the identification of an LRRK2 interacting compound, comprising the steps of:

a) providing two aliquots of a protein preparation containing LRRK2,

b) contacting one aliquot with indol ligand 91 immobilized on a solid support under conditions allowing the formation of an indol ligand 91-LRRK2 complex,

c) contacting the other aliquot with indol ligand 91 immobilized on a solid support and with a given compound under conditions allowing the formation of an indol ligand 91-LRRK2 complex, and

d) determining the amount of indol ligand 91-LRRK2 complex formed in steps b) and c).

In the context of the present invention, it has been surprisingly found that indol ligand 91 is a ligand for LRRK2. This enables the use of indol ligand 91 in screening assays, i.e. in competitive screening assays as well as in methods for the purification of LRRK2.

The structure of indol ligand 91, (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide), is given in Fig. 1. This compound is a molecule structurally similar to the kinase inhibitor Sutent (SU11248; Sun et al., 2003. J. Med. Chem. 46, 1116-1119). Indol ligand 91 can be covalently coupled to a suitable solid support material via the primary amino group and be used for the isolation of binding proteins. The synthesis of indol ligand 91 is described in Example 1. According to the invention, the expression "indol ligand 91" also includes compounds comprising the identical core but which have another linker, preferably coupled to the NH group not being part of the cyclic structures, for linkage to the solid support. Typically linkers may have a backbone of 8, 9 or 10 atoms and may contain either a carboxy- or amino-active group.

According to the present invention, the expression "LRRK2" does not only mean the protein as shown in Figure 3 but also a functionally active derivative thereof, or a functionally active fragment thereof, or a homologue thereof, or a variant encoded by a nucleic acid that hybridizes to the nucleic acid encoding said protein under low stringency conditions. Preferably, these low stringency conditions include hybridization in a buffer comprising 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.02% PVP, 0.02% BSA, 100 ug/ml denatured salmon sperm DNA, and 10% (wt/vol) dextran sulfate for 18-20 hours at 40°C, washing in a buffer consisting of 2X SSC, 25 mM Tris-HCl (pH 7.4), 5 mM EDTA, and 0.1% SDS for 1-5 hours at 55°C, and washing in a buffer consisting of 2X SSC, 25 mM Tris-HCl (pH 7.4) 5 mM EDTA, and 0.1% SDS for 1.5 hours at 60°C.

Moreover, according to the present invention, the expression "LRRK2" includes mutant forms of LRRK2, preferably such mutant forms which observed in PD (in familial forms as well as sporadic cases). Mutations in the LRRK2 gene have been shown to cause

familial autosomal dominant PD (West et al., 2005. PNAS 102, 16842-16847). More preferred, these mutant forms include single amino acid mutations.

The single amino acid substitution G2019S is one of the most frequently observed mutations Taylor et al, 2006. Trends in Molecular Medicine 12, 76-82). This mutation is located in the kinase domain within a conserved region of the activation loop suggesting that modulation of the kinase activity may be involved in the pathogenic mechanism. Other mutations observed in PD include R1441C, Y1699C, and I2020T (see Table 2 of Taylor et al.).

Therefore, in a preferred embodiment, the expression "LRRK2" also includes an LRRK2 protein having a G2019S, R1441C, Y1699C, or I2020T mutation. More preferably, the expression "LRRK2" also includes an LRKK2 protein having one of the following single mutations: G2019S, R1441C, Y1699C, or I2020T.

Furthermore, 15 additional putatively pathogenic amino acid substitutions have been identified in LRRK2 (Taylor et al, 2006. Trends in Molecular Medicine 12, 76-82, see especially Fig. 2).

Therefore, in a preferred embodiment, the expression "LRRK2" also includes an LRRK2 protein having a R793M, Q930R, R1067Q, S1096C, I1122V, S1228T, I1371V, R1441G, R1441H, R1514Q, M1869T, R1941H, I2012T, T2356I or G2385R mutation. More preferably, the expression "LRRK2" also includes an LRKK2 protein having one of the following single mutations: R793M, Q930R, R1067Q, S1096C, I1122V, S1228T, I1371V, R1441G, R1441H, R1514Q, M1869T, R1941H, I2012T, T2356I or G2385R.

In the methods of the invention, first a protein preparation containing LRRK2 is provided. The methods of the present invention can be performed with any protein preparation as a starting material, as long as the LRRK2 is solubilized in the preparation. Examples include a liquid mixture of several proteins, a partial cell lysate which contains not all proteins present in the original cell or a combination of several cell lysates.

Partial cell lysates can be obtained by isolating cell organelles (e.g. nucleus, mitochondria, ribosomes, golgi etc.) first and then preparing protein preparations derived from these organelles. Methods for the isolation of cell organelles are known in the art (Chapter 4.2 Purification of Organelles from Mammalian Cells in ""Current Protocols in Protein Science", Editors: John.E. Coligan, Ben M. Dunn, Hidde L. Ploegh, David W. Speicher, Paul T. Wingfield; Wiley, ISBN: 0-471-14098-8).

In addition, protein preparations can be prepared by fractionation of cell extracts thereby enriching specific types of proteins such as cytoplasmic or membrane proteins (Chapter 4.3 Subcellular Fractionation of Tissue Culture Cells in "Current Protocols in Protein Science", Editors: John.E. Coligan, Ben M. Dunn, Hidde L. Ploegh, David W. Speicher, Paul T. Wingfield; Wiley, ISBN: 0-471-14098-8).

Furthermore protein preparations from body fluids can be used (e.g. blood, cerebrospinal fluid, peritoneal fluid and urine).

For example whole embryo lysates derived from defined development stages or adult stages of model organisms such as C. elegans can be used. In addition, whole organs such as heart dissected from mice can be the source of protein preparations. These organs can also be perfused in vitro in order to obtain a protein preparation.

Furthermore, the protein preparation may be a preparation containing LRRK2 which has been recombinantely produced. Methods for the production of recombinant proteins in prokaryotic and eukaryotic cells are widely established (Chapter 5 Production of Recombinant Proteins *in* "Current Protocols in Protein Science", Editors: John. E. Coligan, Ben M. Dunn, Hidde L. Ploegh, David W. Speicher, Paul T. Wingfield; Wiley, 1995, ISBN: 0-471-14098-8).

In a preferred embodiment of the methods of the invention, the provision of a protein preparation includes the steps of harvesting at least one cell containing LRRK2 and lysing the cell.

In a preferred embodiment, the cell is part of a cell culture system and methods for the harvest of a cell out of a cell culture system are known in the art (literature supra).

The choice of the cell will mainly depend on the expression of LRRK2, since it has to be ensured that the protein is principally present in the cell of choice. In order to determine whether a given cell is a suitable starting system for the methods of the invention, methods like Westernblot, PCR-based nucleic acids detection methods, Northernblots and DNA-microarray methods ("DNA chips") might be suitable in order to determine whether a given protein of interest is present in the cell.

The choice of the cell will also be influenced by the purpose of the study. If the in vivo efficacy for a given drug needs to be analyze then cells or tissues will be selected in which

the desired therapeutic effect occurs (e.g. brain tissue for anti-neurodegenerative drugs). By contrast, for the elucidation of protein targets mediating unwanted side effects the cell or tissue will be analysed in which the side effect is observed (e.g. liver tissue for drug metabolism).

Furthermore, it is envisaged within the present invention that the cell containing LRRK2 may be obtained from an organism, e. g. by biopsy. Corresponding methods are known in the art. For example, a biopsy is a diagnostic procedure used to obtain a small amount of tissue, which can then be examined miscroscopically or with biochemical methods. Biopsies are important to diagnose, classify and stage a disease, but also to evaluate and monitor drug treatment.

It is encompassed within the present invention that by the harvest of the at least one cell, the lysis is performed simultaneously. However, it is equally preferred that the cell is first harvested and then separately lysed.

Methods for the lysis of cells are known in the art (Karwa and Mitra: Sample preparation for the extraction, isolation, and purification of Nuclei Acids; chapter 8 in "Sample Preparation Techniques in Analytical Chemistry", Wiley 2003, Editor: Somenath Mitra, print ISBN: 0471328456; online ISBN: 0471457817). Lysis of different cell types and tissues can be achieved by homogenizers (e.g. Potter-homogenizer), ultrasonic desintegrators, enzymatic lysis, detergents (e.g. NP-40, Triton X-100, CHAPS, SDS), osmotic shock, repeated freezing and thawing, or a combination of these methods.

According to the methods of the invention, the protein preparation is contacted with indol ligand 91 immobilized on a solid support under conditions allowing the formation of a indol ligand 91-LRRK2 complex.

In the present invention, the term "a indol ligand 91-LRRK2 complex" denotes a complex where indol ligand 91 interacts with LRRK2, e.g. by covalent or, most preferred, by non-covalent binding.

The skilled person will know which conditions can be applied in order to enable the formation of the indol ligand 91-LRRK2 complex.

In the context of the present invention, the term "under conditions allowing the formation of the complex" includes all conditions under which such formation, preferably such binding is possible. This includes the possibility of having the solid support on an

immobilized phase and pouring the lysate onto it. In another preferred embodiment, it is also included that the solid support is in a particulate form and mixed with the cell lysate.

In the context of non-covalent binding, the binding between indol ligand 91 and LRRK2 is, e.g., via salt bridges, hydrogen bonds, hydrophobic interactions or a combination thereof.

In a preferred embodiment, the steps of the formation of the indol ligand 91-LRRK2 complex are performed under essentially physiological conditions. The physical state of proteins within cells is described in Petty, 1998 (Howard R. Petty, Chapter 1, Unit 1.5 in: Juan S. Bonifacino, Mary Dasso, Joe B. Harford, Jennifer Lippincott-Schwartz, and Kenneth M. Yamada (eds.) *Current Protocols in Cell Biology* Copyright © 2003 John Wiley & Sons, Inc. All rights reserved. DOI: 10.1002/0471143030.cb0101s00Online Posting Date: May, 2001Print Publication Date: October, 1998).

The contacting under essentially physiological conditions has the advantage that the interactions between the ligand, the cell preparation (i. e. the enzyme to be characterized) and optionally the compound reflect as much as possible the natural conditions. "Essentially physiological conditions" are *inter alia* those conditions which are present in the original, unprocessed sample material. They include the physiological protein concentration, pH, salt concentration, buffer capacity and post-translational modifications of the proteins involved. The term "essentially physiological conditions" does not require conditions identical to those in the original living organism, wherefrom the sample is derived, but essentially cell-like conditions or conditions close to cellular conditions. The person skilled in the art will, of course, realize that certain constraints may arise due to the experimental set-up which will eventually lead to less cell-like conditions. For example, the eventually necessary disruption of cell walls or cell membranes when taking and processing a sample from a living organism may require conditions which are not identical to the physiological conditions found in the organism. Suitable variations of physiological conditions for practicing the methods of the invention will be apparent to those skilled in the art and are encompassed by the term "essentially physiological conditions" as used herein. In summary, it is to be understood that the term "essentially physiological conditions" relates to conditions close to physiological conditions, as e. g. found in natural cells, but does not necessarily require that these conditions are identical.

Preferably, "essentially physiological conditions" may comprise 50-200 mM NaCl or KCl, pH 6.5-8.5, 20-45°C, and 0.001-10 mM divalent cation (e.g. Mg++, Ca++,); more preferably about 150 m NaCl or KCl, pH7.2 to 7.6, 5 mM divalent cation and often include 0.01-1.0 percent non-specific protein (e.g. BSA). A non-ionic detergent (Tween, NP-40,

Triton-X100) can often be present, usually at about 0.001 to 2%, typically 0.05-0.2% (volume/volume). For general guidance, the following buffered aequous conditions may be applicable: 10-250 mM NaCl, 5-50 mM Tris HCl, pH5-8, with optional addition of divalent cation(s) and/or metal chelators and/or non-ionic detergents.

In the context of the present invention, indol ligand 91 is immobilized on a solid support. Throughout the invention, the term "solid support" relates to every undissolved support being able to immobilize a small molecule ligand on its surface.

According to a further preferred embodiment, the solid support is selected from the group consisting of agarose, modified agarose, sepharose beads (e. g. NHS-activated sepharose), latex, cellulose, and ferro- or ferromagnetic particles.

Indol ligand 91 may be coupled to the solid support either covalently or non-covalently. Non-covalent binding includes binding via biotin affinity ligands binding to steptavidin matrices.

Preferably, indol ligand 91 is covalently coupled to the solid support.

Before the coupling, the matrixes can contain active groups such as NHS, Carbodimide etc. to enable the coupling reaction with indol ligand 91. indol ligand 91 can be coupled to the solid support by direct coupling (e.g. using functional groups such as amino-, sulfhydryl-, carboxyl-, hydroxyl-, aldehyde-, and ketone groups) and by indirect coupling (e.g. via biotin, biotin being covalently attached to indol ligand 91 and non-covalent binding of biotin to streptavidin which is bound to solid support directly).

The linkage to the solid support material may involve cleavable and non-cleavable linkers. The cleavage may be achieved by enzymatic cleavage or treatment with suitable chemical methods.

Preferred binding interfaces for binding indol ligand 91 to solid support material are linkers with a C-atom backbone. Typically linkers have backbone of 8, 9 or 10 atoms. Equally preferred are polyethylenglycol (PEG) linkers.

The skilled person will appreciate that between the individual steps of the methods of the invention, washing steps may be necessary. Such washing is part of the knowledge of the person skilled in the art. The washing serves to remove non-bound components of the cell lysate from the solid support. Nonspecific (e.g. simple ionic) binding interactions can be

minimized by adding low levels of detergent or by moderate adjustments to salt concentrations in the wash buffer.

According to the identification methods of the invention, the read-out sytem is either the detection or determination of LRRK2 (first aspect of the invention), the detection of the indol ligand 91-LRRK2 complex (second aspect of the invention), the determination of the amount of the indol ligand 91-LRRK2 complex (third aspect of the invention).

The detection of the indol ligand 91-LRRK2 complex according to the second aspect of the invention can be performed by using labeled antibodies directed against LRRK2 and a suitable readout system.

Antibodies directed against LRRK2 and methods of their use are known in the art, also methods for labeling of antibodies (for example with fluorescent dyes). Anti-LRRK2 antibodies are commercially available (Sant Cruz Biotechnology Inc., Santa Cruz, CA, USA). In addition, antibodies directed at mutated forms of LRRK2 (for example the G2019S and I2020T amino acid substitutions observed in familial forms of PD) are commercially available (Abgent, San Diego, CA, USA).

However, in the course of the second and third aspect of the invention, it is preferred that LRRK2 is separated from the immobilized indol ligand 91.

According to the invention, separating means every action which destroys the interactions between indol ligand 91 and LRRK2. This includes in a preferred embodiment the elution of LRRK2 from the immobilized indol ligand 91.

The elution can be achieved by using non-specific reagents as described in detail below (ionic strength, pH value, detergents). In addition, it can be tested whether a compound of interest can specifically elute LRRK2. Such LRRK2 interacting compounds are described further in the following sections.

Such non-specific methods for destroying the interaction are principally known in the art and depend on the nature of the ligand enzyme interaction. Principally, change of ionic strength, the pH value, the temperature or incubation with detergents are suitable methods to dissociate the target enzymes from the immobilized ligand. The application of an elution buffer can dissociate binding partners by extremes of pH value (high or low pH; e.g. lowering pH by using 0.1 M citrate, pH2-3), change of ionic strength (e.g. high salt concentration using NaI, KI, MgCl2, or KCl), polarity reducing agents which disrupt

hydrophobic interactions (e.g. dioxane or ethylene glycol), or denaturing agents (chaotropic salts or detergents such as Sodium-docedyl-sulfate, SDS; Review: Subramanian A., 2002, Immunoaffinty chromatography).

In some cases, the solid support has preferably to be separated from the released material. The individual methods for this depend on the nature of the solid support and are known in the art. If the support material is contained within a column the released material can be collected as column flowthrough. In case the support material is mixed with the lysate components (so called batch procedure) an additional separation step such as gentle centrifugation may be necessary and the released material is collected as supernatant. Alternatively magnetic beads can be used as solid support so that the beads can be eliminated from the sample by using a magnetic device.

In step d) of the method according to the first aspect of the invention, it is determined if LRRK2 has been separated from immobilized LRRK2. This may include the detection of LRRK2 or the determination of the amount of LRRK2.

Consequently, in preferred embodiments of all identification methods of the invention, methods for the detection of LRRK2 or for the determination of its amount are used. Such methods are known in the art and include physico-chemical methods such as protein sequencing (e.g. Edmann degradation), analysis by mass spectrometry methods or immunodetection methods employing antibodies directed against LRRK2.

Preferably, LRRK2 is detected or the amount of LRRK2 is determined by mass spectrometry or immunodetection methods.

The identification of proteins with mass spectrometric analysis (mass spectrometry) is known in the art (Shevchenko et al., 1996, Analytical Chemistry 68: 850-858; Mann et al., 2001, Analysis of proteins and proteomes by mass spectrometry, Annual Review of Biochemistry 70, 437-473) and is further illustrated in the example section.

Preferably, the mass spectrometry analysis is performed in a quantitative manner, for example by using iTRAQ technology (isobaric tags for relative and absolute quantification) or cICAT (cleavable isotope-coded affinity tags) (Wu et al., 2006. J. Proteome Res. 5, 651-658).

According to a further preferred embodiment of the present invention, the characterization by mass spectrometry (MS) is performed by the identification of proteotypic peptides of

LRRK2. The concept of proteotypic peptides is described in detail in the example section. The idea is that LRRK2 is digested with proteases and the resulting peptides are determined by MS. As a result, peptide frequencies for peptides from the same source protein differ by a great degree, the most frequently observed peptides that "typically" contribute to the identification of this protein being termed "proteotypic peptide". Therefore, a proteotypic peptide as used in the present invention is an experimentally well observable peptide that uniquely identifies a specific protein or protein isoform.

According to a preferred embodiment, the characterization is performed by comparing the proteotypic peptides obtained in the course of practicing the methods of the invention with known proteotypic peptides. Since, when using fragments prepared by protease digestion for the identification of a protein in MS, usually the same proteotypic peptides are observed for a given enzyme, it is possible to compare the proteotypic peptides obtained for a given sample with the proteotypic peptides already known for enzymes of a given class of enzymes and thereby identifying the enzyme being present in the sample.

As an alternative to mass spectrometry analysis, the eluted LRRK2 (including coeluted binding partners or scaffold proteins), can be detected by using specific antibodies directed against LRRK2.

Furthermore, in another preferred embodiment, once the identity of the coeluted binding partner has been established by mass spectrometry analysis, each binding partner can be detected with specific antibodies directed against this protein.

Suitable antibody-based assays include but are not limited to Western blots, ELISA assays, sandwich ELISA assays and antibody arrays or a combination thereof. The establishment of such assays is known in the art (Chapter 11, Immunology, pages 11-1 to 11-30 in: Short Protocols in Molecular Biology. Fourth Edition, Edited by F.M. Ausubel et al., Wiley, New York, 1999).

These assays can not only be configured in a way to detect and quantify an enzyme of interest, but also to analyse posttranslational modification patterns such as phosphorylation. For example, the activation state of a kinase can be determined by probing its phosphorylation status with specific anti-phosphotyrosine, anti-phosphoserine or anti-phosphothreonine antibodies. It is known in the art how to select and use such anti-phospho antibodies (Zhang et al., 2002. Journal of Biological Chemistry 277, 43648-43658).

Furthermore, the identification methods of the invention involve the use of compounds which are tested for their ability to be an LRRK2 interacting compound.

Principally, according to the present invention, such a compound can be every molecule which is able to interact with LRRK2. Preferably, the compound has an effect on LRRK2, e. g. a stimulatory, activating or inhibitory effect.

Preferably, said compound is selected from the group consisting of synthetic or naturally occurring chemical compounds or organic synthetic drugs, more preferably small molecules, organic drugs or natural small molecule compounds. Preferably, said compound is identified starting from a library containing such compounds. Then, in the course of the present invention, such a library is screened.

Such small molecules are preferably not proteins or nucleic acids. Preferably, small molecules exhibit a molecular weight of less than 1000 Da, more preferred less than 750 Da, most preferred less than 500 Da.

A "library" according to the present invention relates to a (mostly large) collection of (numerous) different chemical entities that are provided in a sorted manner that enables both a fast functional analysis (screening) of the different individual entities, and at the same time provide for a rapid identification of the individual entities that form the library. Examples are collections of tubes or wells or spots on surfaces that contain chemical compounds that can be added into reactions with one or more defined potentially interacting partners in a high-throughput fashion. After the identification of a desired "positive" interaction of both partners, the respective compound can be rapidly identified due to the library construction. Libraries of synthetic and natural origins can either be purchased or designed by the skilled artisan.

Examples of the construction of libraries are provided in, for example, Breinbauer R, Manger M, Scheck M, Waldmann H. Natural product guided compound library development. Curr Med Chem. 2002 Dec;9(23):2129-45, wherein natural products are described that are biologically validated starting points for the design of combinatorial libraries, as they have a proven record of biological relevance. This special role of natural products in medicinal chemistry and chemical biology can be interpreted in the light of new insights about the domain architecture of proteins gained by structural biology and bioinformatics. In order to fulfill the specific requirements of the individual binding pocket within a domain family it may be necessary to optimise the natural product structure by chemical variation. Solid-phase chemistry is said to become an efficient tool for this

optimisation process, and recent advances in this field are highlighted in this review article. Other related references include Edwards PJ, Morrell AI. Solid-phase compound library synthesis in drug design and development. Curr Opin Drug Discov Devel. 2002 Jul;5(4):594-605.; Merlot C, Domine D, Church DJ. Fragment analysis in small molecule discovery. Curr Opin Drug Discov Devel. 2002 May;5(3):391-9. Review; Goodnow RA Jr. Current practices in generation of small molecule new leads. J Cell Biochem Suppl. 2001;Suppl 37:13-21; which describes that the current drug discovery processes in many pharmaceutical companies require large and growing collections of high quality lead structures for use in high throughput screening assays. Collections of small molecules with diverse structures and "drug-like" properties have, in the past, been acquired by several means: by archive of previous internal lead optimisation efforts, by purchase from compound vendors, and by union of separate collections following company mergers. Although high throughput/combinatorial chemistry is described as being an important component in the process of new lead generation, the selection of library designs for synthesis and the subsequent design of library members has evolved to a new level of challenge and importance. The potential benefits of screening multiple small molecule compound library designs against multiple biological targets offers substantial opportunity to discover new lead structures.

In a preferred embodiment of the present disclosure, LRRK2 is first incubated with the compound and then with the immobilized indol ligand 91. However, the simultaneous incubation of the compound and the immobilized indol ligand 91 (coincubation) with the LRRK2 containing protein is equally preferred (competitive binding assay).

Preferably the LRRK2 is first incubated with the compound for 10 to 60 minutes, more preferred 30 to 45 minutes at a temperature of 4°C to 37°C, more preferred 4°C to 25°C, most preferred 4°C. Preferably compounds are used at concentrations ranging from 1 µM to 1 mM, preferably from 10 to 100 µM. The second step, contacting with the immobilized ligand, is preferably performed for 10 to 60 minutes at 4°C.

In the case of simultaneous incubation, the LRRK2 is preferably simultaneously incubated with the compound and immobilized indol ligand 91 for 10 to 120 minutes, more preferred 20 to 60 minutes at a temperature of 4°C to 37°C, more preferred 4°C to 25°C, most preferred 4°C to 8°C. Preferably compounds are used at concentrations ranging from 1 nM to 100 µM, preferably from 10 nM to 10 µM.

Furthermore, steps a) to c) of the disclosure herein may be performed with several protein preparations in order to test different compounds. This embodiment is especially interesting in the context of medium or high troughput screenings (see below).

In a preferred embodiment of the method of the present disclosure, a reduced amount of indol ligand 91-LRRK2 formed in step c) in comparison to step b) indicates that LRRK2 is a target of the compound. This results from the fact that in step c) of this method of the invention, the compound competes with the ligand for the binding of LRRK2. If less LRRK2 is present in the aliquot incubated with the compound, this means preferably that the compound has competed with the inhibitor for the interaction with the enzyme and is, therefore, a direct target of the enzyme and vice versa.

Preferably, the identification methods of the invention are performed as a medium or high throughput screening. Such assays are known to the person skilled in the art (Mallari et al., 2003, A generic high-throughput screening assay for kinases: protein kinase A as an example, Journal of Biomolecular Screeing 8, 198-204; Rodems et al., 2002, A FRET-based assay platform for ultra-high density screening of protein kinases and phosphatases, Assay and Drug Development Technologies 1 (1PT1), 9-19).

The interaction compound identified according to the present invention may be further charcterized by determining whether it has an effect on LRRK2 activity, for example on its kinase activity (West et al., 2005. PNAS 102, 16842-16847). Such assays are known in the art, also in a format that allows medium to high throughput screening.

Briefly, a fluorescein-labeled peptide substrate may be incubated with the tyrosine kinase (e.g. LRRK2), ATP and an anti-phosphotyrosine antibody. As the reaction proceeds, the phosphorylated peptide binds to the anti-phosphotyrosine antibody, resulting in an increase in the polarization signal. Compounds that inhibit the kinase result in a low polarization signal.

Alternatively, the assay can be configured in a modified indirect format. A fluorescent phosphopeptide is used as a tracer for complex formation with the anti-phospho-tyrosine antibody yielding a high polarization signal. When unlabeled substrate is phosphorylated by the kinase, the product competes with the fluorescent phosphorylated peptide for the antibody. The fluorescent peptide is then released from the antibody into solution resulting in a loss of polarization signal. Both the direct and indirect assays can be used to identify inhibitors of protein tyrosine kinase activity (Seethala, 2000, Methods 22, 61-70; Seethala

and Menzel, 1997, Anal. Biochem. 253, 210-218; Seethala and Menzel, 1998, Anal. Biochem. 255, 257-262).

This fluorescence polarization assay can be adapted for the use with protein serine/threonine kinases by replacing the antiphophotyrosine antibody with an anti-phosphoserine or anti-phosphothreonine antibody (Turek et al., 2001, Anal. Biochem. 299, 45-53, PMID 11726183; Wu et al., 2000, J. Biomol. Screen. 5, 23-30, PMID 10841597).

The compounds identified according to the present invention may further be optimized (lead optimisation). This subsequent optimisation of such compounds is often accelerated because of the structure-activity relationship (SAR) information encoded in these lead generation libraries. Lead optimisation is often facilitated due to the ready applicability of high-throughput chemistry (HTC) methods for follow-up synthesis.

One example of such a library and lead optimization is described in Wakeling AE, Barker AJ, Davies DH, Brown DS, Green LR, Cartlidge SA, Woodburn JR. Specific inhibition of epidermal growth factor receptor tyrosine kinase by 4-anilinoquinazolines. Breast Cancer Res Treat. 1996;38(1):67-73.

The invention further relates to a method for the preparation of a pharmaceutical composition comprising the steps of

a) identifying a LRRK2 interacting compound as described above, and

b) formulating the interacting compound to a pharmaceutical composition.

Therefore, the invention provides a method for the preparation of pharmaceutical compositions, which may be administered to a subject in an effective amount. In a preferred aspect, the therapeutic is substantially purified. The subject to be treated is preferably an animal including, but not limited to animals such as cows, pigs, horses, chickens, cats, dogs, etc., and is preferably a mammal, and most preferably human. In a specific embodiment, a non-human mammal is the subject.

In general, the pharmaceutical compositions of the present invention comprise a therapeutically effective amount of a therapeutic, and a pharmaceutically acceptable carrier. In a specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more

particularly, in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, including but not limited to peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered orally. Saline and aqueous dextrose are preferred carriers when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions are preferably employed as liquid carriers for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsions, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin. Such compositions will contain a therapeutically effective amount of the therapeutic, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

In a preferred embodiment, the composition is formulated, in accordance with routine procedures, as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lidocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water-free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water or saline for injection can be provided so that the ingredients may be mixed prior to administration.

The therapeutics of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with free carboxyl groups such as

those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., those formed with free amine groups such as those derived from isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc., and those derived from sodium, potassium, ammonium, calcium, and ferric hydroxides, etc..

The amount of the therapeutic of the invention which will be effective in the treatment of a particular disorder or condition will depend on the nature of the disorder or condition, and can be determined by standard clinical techniques. In addition, in vitro assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient's circumstances. However, suitable dosage ranges for intravenous administration are generally about 20-500 micrograms of active compound per kilogram body weight. Suitable dosage ranges for intranasal administration are generally about 0.01 pg/kg body weight to 1 mg/kg body weight. Effective doses may be extrapolated from dose-response curves derived from in vitro or animal model test systems. In general, suppositories may contain active ingredient in the range of 0.5% to 10% by weight; oral formulations preferably contain 10% to 95% active ingredient.

Various delivery systems are known and can be used to administer a therapeutic of the invention, e.g., encapsulation in liposomes, microparticles, and microcapsules: use of recombinant cells capable of expressing the therapeutic, use of receptor-mediated endocytosis (e.g., Wu and Wu, 1987, J. Biol. Chem. 262:4429-4432); construction of a therapeutic nucleic acid as part of a retroviral or other vector, etc. Methods of introduction include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The compounds may be administered by any convenient route, for example by infusion, by bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral, rectal and intestinal mucosa, etc.), and may be administered together with other biologically active agents. Administration can be systemic or local. In addition, it may be desirable to introduce the pharmaceutical compositions of the invention into the central nervous system by any suitable route, including intraventricular and intrathecal injection; intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir. Pulmonary administration can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

In a specific embodiment, it may be desirable to administer the pharmaceutical compositions of the invention locally to the area in need of treatment. This may be

achieved by, for example, and not by way of limitation, local infusion during surgery, topical application, e.g., in conjunction with a wound dressing after surgery, by injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. In one embodiment, administration can be by direct injection at the site (or former site) of a malignant tumor or neoplastic or pre-neoplastic tissue.

In another embodiment, the therapeutic can be delivered in a vesicle, in particular a liposome (Langer, 1990, Science 249:1527-1533).

In yet another embodiment, the therapeutic can be delivered via a controlled release system. In one embodiment, a pump may be used (Langer, supra). In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose

The invention further relates to a method for the purification of LRRK2, comprising the steps of

a) providing a protein preparation containing LRRK2,

b) contacting the protein preparation with indol ligand 91 immobilized on a solid support under conditions allowing the formation of an indol ligand 91-LRRK2 complex, and

c) separating LRRK2 from the immobilized indol ligand 91.

As mentioned above, it has been surprisingly found that indol ligand 91 is an LRRK2 ligand. This enables efficient LRRK2 purification methods.

With respect to LRRK2, the protein peparation containing LRRK2, the conditions for contacting with indol ligand 91, immobilized indol ligand 91, the indol ligand 91-LRRK2 complex, the separation of LRRK2 from the immobilized indol ligand 91, and the detection of LRRK2 or the determination of its amount, the embodiments as defined above for the identification methods of the invention also apply to the purification method of the invention.

In a preferred embodiment, the purification method of the invention further comprises after step c) the identification of proteins being capable of binding to LRRK2. This is especially interesting when the formation of the complex is performed under essentially physiological conditions, because it is then possible to preserve the natural condition of the enzyme which includes the existence of binding partners, enzyme subunits or post-translational modifications, which can then be identified with the help of mass spectrometry (MS).

Consequently, in a preferred embodiment, the purification method of the invention further comprises after step c) the determination whether LRRK2 is posttranslationally modified.

The invention further relates to the use of indol ligand 91 for the identification of LRRK2 interacting compounds and for the purification of LRRK-2. The embodiments as defined above also apply to the uses of the invention.

The invention is further illustrated by the following figures and examples, which are not considered as being limiting for the scope of protection conferred by the claims of the present application.

**Short description of the figures**

**Figure 1:** Structure of linkable indol ligand 91

The figure shows the structure of linkable indol ligand 91. The free primary amino group can be used for covalent coupling to a solid support material.

**Figure 2:** Drug pulldown experiment with immobilized indol ligand 91

The figure shows the picture of a protein gel after staining with Coomassie blue. Proteins bound to immobilized indol ligand 91 were eluted with SDS sample buffer and analyzed by SDS-polyacrylamide gel electrophoresis (SDS-PAGE). The indicated gel areas were cut out as gel slices and proteins were subjected to analysis by mass spectrometry. The position of the band containing LRRK2 is in sample 01 in the upper part of the gel between 00b and 01a.

**Figure 3:** Peptides identified of LRRK2

The peptides that were identified by mass spectrometry analysis of the mouse LRRK2 sequence (IPI00227565.2, 2527 amino acids) are underlined.

**Example 1: Synthesis of linkable indol ligand 91**

**Synthesis of indol ligand 91: 5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide**

**Step 1:** 5-Fluoro-1,3-dihydro-indol-2-one

A solution of 5-Fluoroisatin (1g) in Hydrazine hydrate (55%, 10ml) was heated at 110°C for 30minutes. Once the suspension has gone into solution, the reaction was heated at 110°C for 4 hours, then cooled at 0°C. The precipitate was filtered and washed with water. The solid was suspended in water (10ml), the pH was lowered to pH2 by addition of HCl conc, and the solution stirred at room temperature for 5 hours. The precipitate was collected, the solid washed with water (2x15ml) and dried in the vacuum oven at 40°C (0.26g,30%). $^1$H NMR (400 MHz, DMSO-$d_6$) δ 10.2 (s, 1H); 6.8-7.0 (dd, 2H); 6.6 (m, 1H); 3.2 (s, 2H);. LCMS: method D, RT=1.736min, [$MH^+$=152].

**Step 2:** {3-[(5-Formyl-2,4-dimethyl-1H-pyrrole-3-carbonyl)-amino]-propyl}-carbamic acid tert-butyl ester

To a solution of 5-formyl-2,4-dimethyl-1H-pyrazol-3-carboxylic acid (0.300g, 1.79mmol) , N-(3-Dimethylaminopropyl)-N'-ethylcarbodiimide (0.516g, 2.69mmol), 1-Hydroxybenzotriazole hydrate (0.364g, 2.69mmol), triethylamine (0.502ml, 3.56mmol) in dimethylformamide (3ml) was added N-Boc-1,3-diaminopropane (0.375ml, 2.15mmol). The solution was stirred at room temperature for 15 hours. A mixture of brine (1.5ml), water (1.5ml) and saturated aqueous sodium bicarbonate (1.5ml) was added and the pH of the solution adjusted to 12 by addition of 10N sodium hydroxide. The solution was extracted 3 times with a mixture of dichloromethane: Methanol (9:1). The organic layer was dried with anhydrous magnesium sulfate. The solvent was removed and the residue purified by flash chromatography (Hexane : Ethyl acetate(50 to100%)) to yield the desired compound as a yellow solid (0.30g, 52%). $^1$H NMR (400 MHz, DMSO-$d_6$) δ 11.90 (s, 1H); 9.50 (s, 1H); 7.50 (m, 1H); 6.90 (m, 1H); 3.20 (q, 2H); 3.00 (q, 2H); 2.30 (s, 3H) ); 2.20 (s, 3H) ); 1.60 (m, 2H) ); 1.40 (s, 9H). LCMS: method D, RT=2.103min, [$M+Na^+$=346], and [$M-Boc+Na^+$=246].

**Step 3:** [3-({5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carbonyl}-amino)-propyl]-carbamic acid tert-butyl ester

A solution of {3-[(5-Formyl-2,4-dimethyl-1H-pyrrole-3-carbonyl)-amino]-propyl}-carbamic acid tert-butyl ester (0.200g, 0.62mmol) and 5-Fluoro-1,3-dihydro-indol-2-one (0.011g, 0.62mmol), pyrrolidine (0.003ml) in ethanol (2ml) was heated at 78°C for 3 hours. The reaction was cooled to 0°C and the resulting precipitate filtered, washed with

cold ethanol. The product was suspended in ethanol (4ml) and stirred at 72°C for 30minutes. The reaction was filtered, the precipitate dried in a vacuum oven at 40°C to yield the desired compound as a solid (0.265g, 94%). $^1$H NMR (400 MHz, DMSO-$d_6$) δ 13.8 (s, 1H); 11.00 (s, 1H); 7.80 (m, 2H); 7.70 (m, 1H); 7.0 (m, 1H); 6.9 (m, 2H); 3.30 (q, 2H); 3.10 (q, 2H); 2.50 (dd, 6H) 1.60 (m, 2H) ); 1.40 (s, 9H). LCMS: method D, RT=2.86min, [M+Na$^+$=479], and [M-Boc+Na$^+$=379].

**Step 4:** 5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide

[3-({5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carbonyl}-amino)-propyl]-carbamic acid tert-butyl ester was suspended in methanol. 2ml of HCl (4N) in dioxane was added and the reaction stirred at room temperature overnight. The solvent was removed to yield the desired compound (0.098g, 91 %). $^1$H NMR (400 MHz, DMSO-$d_6$) δ 13.8 (s, 1H); 11.00 (s, 1H); 7.90-7.70 (m, 5H); 6.9 (m, 2H); 3.30 (q, 2H); 2.80 (q, 2H); 2.50 (dd, 6H) 1.80 (m, 2H). LCMS: inconclusive due to fluorescence.

All reactions were carried out under inert atmosphere. NMR spectra were obtained on a Bruker dpx400. LCMS was carried out on an Agilent 1100 using a zorbax SBC-18, 4.6mmx150mm-5μ column or a Small column: ZORBAX® SB-C18, 4.6 x 75 mm, 3.5 microns ("short column"). Column flow was 1ml/min and solvents used were water and acetonitrile (0.1%TFA) with an injection volume of 10ul. Wavelengths were 254 and 210nm. Methods are described below.

**Table 1:** Analytical methods

| **Method** | **Easy Access Method Name** | **ChemStation Method Name** | **Flow Rate** | **Solvent** | **Run Time** |
|---|---|---|---|---|---|
| A | Analytical positive 7mn | ANL_POS7.M | 1ml/min | 0-2.5min<br>5-95% MeCN<br><br>2.5-6min<br>95% MeCN | 7 min |
| B | Analytical positive Ion | ANAL_POS.M | 1ml/min | 0-11min<br>5-95% MeCN<br><br>11-13min<br>95% MeCN | 15 min |
| C | Loop injection, Positive | | 1ml/min | 95% MeCN | 1min |
| D | Analytical positive Ion | Short column ANL Positive | 1ml/mn | 0-4.5min<br>30-95% MeCN | 5min |
| E | Analytical High pH | Analytical High pH | 3ml/min | 0 to 8min<br>5-95% MeCN<br>8 to 9min<br>95% MeCN | 10min |

**Table 2:** Abbreviations used in chemistry protocols

| | |
|---|---|
| aq | aqueous |
| D | doublet |
| DMSO | dimethyl sulfoxide |
| G | gram |
| HCl | Hydrochloric acid |
| HPLC | high pressure liquid chromatography |
| LCMS | liquid chromatography – mass spectrometry |
| M | multiplet |
| mins | minute |
| mmol | millimole |
| N | Normal |
| NMR | nuclear magnetic resonance |
| Q | quartet |
| RT | retention time |
| S | singlet |
| sat | saturated |
| T | triplet |

**Example 2: Drug pulldown with immobilized indol ligand 91 and identification of LRRK2**

This example demonstrates the use of the immobilzed indol ligand 91 ligand for the identification of LRRK2 from mouse brain lysate. Proteins binding to immobilized indol ligand 91 were identified using a chemical proteomics procedure. Binding proteins were purified on a resin to which indol ligand 91 was covalently attached. Bound proteins were eluted and subsequently separated by SDS-Polyacrylamide gel elecrophoresis (see Figure

2), suitable gel bands were cut out and proteins were analyzed by LC-MS/MS mass spectrometry.

The results show that immobilized indol ligand 91 can be used to enrich and characterize LRRK2 from mouse brain lysates. The twelve LRRK2 peptides identified by mass spectrometry are listed in Table 4.

1. Immobilization of small molecule amine compounds

NHS-activated Sepharose 4 Fast Flow (Amersham Biosciences, 17-0906-01) was equilibrated with anhydrous DMSO (Dimethylsulfoxid, Fluka, 41648, H20 <= 0.005%). 1 ml of settled beads were placed in a 15 ml Falcon tube, compound stock solution (usually 100 mM in DMF or DMSO) was added (final conc. 0.2-2 µmol/ml beads) as well as 15 µl of TEA. Beads were incubated at room temperature in darkness on an end-over-end shaker (Roto Shake Genie, Scientific Industries Inc.) for 16 - 20 hours. Coupling efficiency was determined by HPLC. Non-reacted NHS-groups were blocked by incubation with aminoethanol at room temperature on the end-over-end shaker over night. Beads were washed with 10 ml of DMSO. Washed beads were stored in isopropanol at 4°C.

2. Preparation of cell lysates

Mouse brain tissue was homogenized in a Potter S homogenizer (5 ml per brain): 50 mM Tris-HCl, 1% CHAPSO, 5% glycerol, 150 mM NaCl, 1.5 mM MgCl2, 25 mM NaF, 1 mM sodium vanadate, 1 mM DTT, pH 7.5 . One complete EDTA-free tablet (protease inhibitor cocktail, Roche Diagnostics, 1 873 580) per 25 ml buffer was added. The material was dounced 10x using a mechanized POTTER S, transferred to 50 ml falcon tubes, incubated for 30 minutes on ice and spun down for 10 min at 20,000 g at 4°C (10,000 rpm in Sorvall SLA600, precooled). The supernatant was transfered to an ultracentrifuge (UZ)-polycarbonate tube (Beckmann, 355654) and spun for 1 hour at 100.000 g at 4°C (33.500 rpm in Ti50.2, precooled). The supernatant was transferred again to a fresh 50 ml falcon tube, the protein concentration was determined by a Bradford assay (BioRad) and samples containing 50 mg of protein per aliquot were prepared. The samples were immediately used for experiments or frozen in liquid nitrogen and stored frozen at -80°C.

3. Compound pull-down experiment

Sepharose-beads with immobilized compound (100 µl beads per pull-down experiment) were equilibrated in lysis buffer and incubated with a cell lysate sample containing 50 mg

of protein on an end-over-end shaker (Roto Shake Genie, Scientific Industries Inc.) for 2 hours at 4°C. Beads were collected, transfered to Mobicol-columns (MoBiTech 10055) and washed with 10 ml lysis buffer containing 0.5% detergent (NP40), followed by 5 ml lysis buffer with 0.25 % detergent. To elute the bound protein, 60 µl 2 x SDS sample buffer was added, the column was heated for 30 minutes at 50°C and the eluate was transferred to a microfuge tube by centrifugation. Proteins were then separated by SDS-Polyacrylamide electrophoresis (SDS-PAGE).

4. Protein Identification by Mass Spectrometry

4.1 Protein digestion prior to mass spectrometric analysis

Gel-separated proteins were reduced, alkylated and digested in gel essentially following the procedure described by Shevchenko et al., 1996, Anal. Chem. 68:850-858. Briefly, gel-separated proteins were excised from the gel using a clean scalpel, reduced using 10 mM DTT (in 5mM ammonium bicarbonate, 54°C, 45 min) and subsequently alkylated with 55 mM iodoacetamid (in 5 mM ammonium bicarbonate) at room temperature in the dark (30 minutes). Reduced and alkylated proteins were digested in gel with porcine trypsin (Promega) at a protease concentration of 12.5 ng/µl in 5mM ammonium bicarbonate. Digestion was allowed to proceed for 4 hours at 37°C and the reaction was subsequently stopped using 5 µl 5% formic acid.

4.2 Sample preparation prior to analysis by mass spectrometry

Gel plugs were extracted twice with 20 µl 1% TFA and pooled with acidified digest supernatants. Samples were dried in a a vaccuum centrifuge and resuspended in 13 µl 1% TFA.

4.3. Mass spectrometric data acquisition

Peptide samples were injected into a nano LC system (CapLC, Waters or Ultimate, Dionex) which was directly coupled either to a quadrupole TOF (QTOF2, QTOF Ultima, QTOF Micro, Micromass) or ion trap (LCQ Deca XP) mass spectrometer. Peptides were separated on the LC system using a gradient of aqueous and organic solvents (see below). Solvent A was 5% acetonitrile in 0.5% formic acid and solvent B was 70% acetonitrile in 0.5% formic acid.

**Table 3:** Peptides eluting off the LC system were partially sequenced within the mass spectrometer.

| Time (min) | % solvent A | % solvent B |
|---|---|---|
| 0 | 95 | 5 |
| 5.33 | 92 | 8 |
| 35 | 50 | 50 |
| 36 | 20 | 80 |
| 40 | 20 | 80 |
| 41 | 95 | 5 |
| 50 | 95 | 5 |

4.4. Protein identification

The peptide mass and fragmentation data generated in the LC-MS/MS experiments were used to query fasta formatted protein and nucleotide sequence databases maintained and updated regularly at the NCBI (for the NCBInr, dbEST and the human and mouse genomes) and European Bioinformatics Institute (EBI, for the human, mouse, D. melanogaster and C. elegans proteome databases). Proteins were identified by correlating the measured peptide mass and fragmentation data with the same data computed from the entries in the database using the software tool Mascot (Matrix Science; Perkins et al., 1999. Probability-based protein identification by searching sequence databases using mass spectrometry data. Electrophoresis 20, 3551-3567). Search criteria varied depending on which mass spectrometer was used for the analysis.

**Table 4:** Peptide identification by mass spectrometry of LRRK2 from mouse brain (Experiment P11927B)

| Peptide number | Peptide position | Peptide sequence |
|---|---|---|
| 1 | 172 - 180 | YSANDEVQK |
| 2 | 454 - 466 | HAHAPEVAESGCK |
| 3 | 553 - 561 | FIGNPGIQK |
| 4 | 630 - 637 | ILASTLQR |
| 5 | 740 - 751 | GATSLIYQVCEK |
| 6 | 782 - 792 | GDSQVISLLLR |
| 7 | 832 - 841 | QTNTGSVLAR |
| 8 | 1016 - 1035 | LELHQNSLTSFPQQLCETLK |
| 9 | 1712 - 1723 | LLEISPFMLSGR |
| 10 | 1765 - 1771 | ITVPSC |
| 11 | 2031 - 2039 | TSEGTPGFR |
| 12 | 2406 - 2413 | HQLSYSGR |

**Example 3: Screening assay for the identification of LRRK2 interacting compounds**

This examples describes a competitive binding assay in which test compunds are added directly into a cell lysate. Test compounds and the affinity matrix with the immobilized indol ligand 91 are added to lysate aliquots and allowed to bind to the proteins contained in the lysate sample. After the inbutation time the beads with captured proteins are separated from the lysate. Bound proteins are then eluted and the presence of LRRK2 in the eluate is detected and quantified using a specific antibody in a dot blot procedure and the Odyssey infrared detection system. By using various concentrations of a test compound a dose response curve is generated. Further experimental information is disclosed in WO2006/134056.

Washing of the affinity matrix

The affinity matrix as described in example 1 is washed two times with 1x DP buffer containing 0.4% NP40 and then resupended in 1x DP buffer containing 0.4% NP40 (20% beads slurry).

**Table 5:** Preparation of 5x DP buffer

| **Substance:** | **Stock solution** | **Final conc. in 1 x lysis buffer** | **Add for 1l 5 x lysis buffer** |
|---|---|---|---|
| **Tris/HCl pH 7.5** | 1 M | 50 mM | 250 ml |
| **Glycerol** | 87 % | 5 % | 288 ml |

| | | | |
|---|---|---|---|
| $MgCl_2$ | 1 M | 1.5 mM | 7.5 ml |
| NaCl | 5 M | 150 mM | 150 ml |
| $Na_3VO_4$ | 100 mM | 1 mM | 50 ml |

The five-fold concentrated (5x) DP buffer is filtered through a 0.22 µm filter and stored at -80°C. In order to obtain 1xDP buffer the 5xDP buffer is diluted 1:5 with distilled water. Stock solutions are obtained from the following suppliers: 1.0 M Tris/HCl pH 7.5 (Sigma, T-2663), 87% Glycerol (Merck, catalogue number 04091.2500); 1.0 M $MgCl_2$ (Sigma, M-1028); 5.0 M NaCl (Sigma, S-5150).

Preparation of test compounds

Stock solutions of test compounds are prepared in DMSO corresponding to a 100fold higher concentration compared to the final desired test concentration (e.g. a 400 µM stock solution is prepared for a final test concentration of 4 µM). This dilution scheme results in a final DMSO concentration of 1% in the assay. For control experiments (no test compound) a buffer containing 1% DMSO is used so that all samples contain 1% DMSO.

Preparation and dilution of cell lysate

The cell lysate is prepared as described in example 2. For a typical experiment one lysate aliquot containing 50 mg of protein is thawed in a 37°C water bath and then kept at 4°C. To the lysate one volume of 1xDP buffer is added so that a final NP40 concentration of 0.4% is achieved. Then, 1/50 of the final volume of a 50fold concentrated protease inhibitor solution is added (1 tablet of protease inhibitor dissolved in 0.5 ml of 1x DP buffer containing 0.4% NP40; EDTA-free tablet protease inhibitor cocktail from Roche Diagnostics, catalogue number 41647). The lysate is further diluted by adding 1x DP buffer containing 0.4% NP40 so that a final protein concentration of 5 mg/ml is achieved.

Incubation of lysate with test compound and affinity matrix

A volume of 100 µl of diluted lysate is dispensed into each well of a 96 well filter plate. Then 1.5 µl of test compound diluted in DMSO is added. For control reactions 1.5 µl DMSO without test compound is used. Then 50 µl of affinity matrix (20% slurry) per well are added. The plate is incubated for two hours at 4°C on a shaker. The plate is washed using a vacuum manifold instrument (Millipore, MAVM 096 0R). Each well is washed four times with 400 µl of 1x DP buffer containing 0.4% NP-40 and two times with 400 µl 1x DP buffer containing 0.2% NP-40.

For elution the filter plate is placed on a collection plate and 40 µl of 2x sample buffer (100 mM TrisHCl, pH6.8; 4% SDS; 20% glycerol; 0.02% Bromphenol blue) with DTT (50 mM final concentration) is added to each well. The plates are incubated for 30 minutes at room

temperature on a shaker (750 rpm on a Thermomixer, Eppendorf). Subsequently the plates are centrifuged for 2 minutes at 1100 rpm (Heraeus centrifuge) and the eluate is collected in the wells of the collection plate.

Detection and quantification of eluted LRRK2

The LRRK2 protein in the eluates is detected and quantified by a dot blot procedure using a first antibody directed against LRRK2 (Santa Cruz Biotechnolgy, Santa Cruz, CA, USA) and a fluorescently labeled secondary antibody (anti-mouse IRDye™ 800, from Rockland, 610-732-124). The Odyssey Infrared Imaging system from LI-COR Biosciences (Lincoln, Nebraska, USA) is operated according to instructions provided by the manufacturer (Schutz-Geschwendener et al., 2004. Quantitative, two-color Western blot detection with infrared fluorescence. Published May 2004 by LI-COR Biosciences, www.licor.com).

The dot blot apparatus is used according to the instructions of the supplier (Bio-Dot microfiltration apparatus, BioRad 170-65). Nitrocellulose membranes (BioTrace NT Nitrocellulose, PALL BTNT30R) are treated with 20% ethanol and subsequently washed with 1xPBS buffer. Per dot 30 µl of eluate sample are applied .

For the detection of LRRK2 the membranes are first blocked by incubation with Odyssey blocking buffer (LICOR, 927-40000) for one hour at room temperature. Blocked membranes are then incubated for 16 hours at 4°C with the anti-LRRK2 antibody which is diluted in Odyssey blocking buffer containing 0.2% Tween-20. After washing the membrane four times with 1xPBS buffer containing 0.1% Tween 20 the membrane is incubated for 40 minutes with the detection antibody (anti-mouse IRDye™ 800 from Rockland, 610-732-124) diluted in Odyssey blocking buffer containing 0.2% Tween-20. Afterwards the membrane is washed four times for five minutes each with 1x PBS buffer/0.1% Tween 20 and once for 5 minutes with 1x PBS buffer. The membrane is kept in PBS buffer at 4°C and then scanned with the Odyssey instrument and signals are recorded and analysed according to the instructions of the manufacturer.

**Example 4: LRRK2 kinase assay**

This example describes an assay that can be used to determine whether a LRRK2 interacting compound is a LRRK2 kinase inhibitor. The assay is performed as described previously (West et al., 2005. PNAS 102, 16842-16847).

A full-length LRRK2 cDNA clone with a Myc epitope tag is cloned into a mammalian expression vector. The expression plasmid is transiently transfected into HEK-293T cells by using the FuGENE 6.0 reagent (Roche Applied Science) according to the manufacturer's instructions. HEK-293T cells are cultured in Opti-MEM media (Invitrogen) supplemented with 10% FBS, penicillin (100 units/ml), and streptomycin (100 µg/ml). To prepare recombinant LRRK2 protein for kinase assays HEK-293T cells are harvested in immunoprecipitation buffer (0.5% Triton-X-100, 1 x complete mini protease inhibitor cocktail(Roche) in PBS) and the lysates are incubated at 4°C for one hour followed by centrifugation at 17,500 x g for 15 minutes. The supernatant fraction is combined with Protein G sepharose 4 Fast Flow (Amersham Pharmacia Biotech) precomplexed with the mouse monoclonal anti-c-myc antibody (clone 9E10; Roche) followed by incubation overnight with rotating the tube. The protein G Sepharose complex is pelleted by centrifugation and washed five times with buffer supplemented with 500 mM NaCl and once with 1x PBS buffer.

Myc-tagged LRRK2 protein bound to G sepharose is resuspended in kinase buffer (20 mM Hepes, pH7.4; 15 mM $MgCl_2$; 5 mM EGTA; 20 mM beta-glycerol phosphate) on ice together with 25 µM biotinylated myelin basic protein (MBP; Upstate Biotechnology, Lake Placid, NY). The kinase reaction is initiated by adding 0.5 µCi of [gamma$^{32}$P]ATP (Perkin Elmer) and 15 µM ATP. The mixture is incubated at 30°C for 15 minutes. The reaction is placed on ice and terminated by adding 2.5 M guanidine-HCl. The universal kinase assay kit (Calbiochem) is used to quantify incorporated radioisotope in biotinylated MBP by adding avidin solution (Calbiochem) to the reaction and incubating at 25°C for 10 minutes. The reaction is centrifuged at 800 x g for two minutes and the supernatant is placed on a column containing an avidin binding membrane and washed according to the manufacturer's protocol. The amount of incorporated radioisotope is measured by liquid scintillation counting.

The LRRK2 autophosphorylation acitivity is determined in a kinase reaction as described above except that no MBP is added. The reaction is incubated for 15 minutes at 30°C and terminated by adding Laemmli sample buffer containing 5% 2-mercaptoethanol. The reactions are heated for 10 minutes at 75°C and then placed on ice. After brief centrifugation the samples (supernatants) are separated in a 6%SDS-Polyacrylamide gel, transferred to a poly-vinylidinedifluoride (PVDF) membrane and exposed to BioMax film (Kodak). The membrane is probed with an anti-c-myc antibody (9E10; Roche) to determine the amount of LRRK2 in each reaction. The autoradiography and Western blot bands are quantified by using IMAGEQUANT 5.0 thereby normalizing the amount of phosphorylated MBP to the amount of LRRK2 in each experiment.

CLAIMS

1. A method for the identification of an LRRK2 interacting compound, comprising the steps of
   a) providing a protein preparation containing LRRK2,
   b) contacting the protein preparation with (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide) immobilized on a solid support under conditions allowing the formation of an (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex,
   c) incubating the (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex with a given compound, and
   d) determining whether the compound is able to separate LRRK2 from the immobilized (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide).

2. The method of claim 1, wherein step d) includes the detection of LRRK2 or the determination of the amount of LRRK2.

3. The method of claim 2, wherein LRRK2 is detected or the amount of LRRK2 is determined by mass spectrometry or immunodetection methods.

4. A method for the identification of an LRRK2 interacting compound, comprising the steps of
   a) providing a protein preparation containing LRRK2,
   b) contacting the protein preparation with (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide) immobilized on a solid support and with a given compound under conditions allowing the formation of an (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex, and

c) detecting the (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex formed in step b).

5. The method of claim 4, wherein in step c) said detecting is performed by determining the amount of the (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex.

6. The method of any of claims 4 or 5, wherein steps a) to c) are performed with several protein preparations in order to test different compounds.

7. A method for the identification of an LRRK2 interacting compound, comprising the steps of:

a) providing two aliquots of a protein preparation containing LRRK2,

b) contacting one aliquot with (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide) immobilized on a solid support under conditions allowing the formation of an (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex,

c) contacting the other aliquot with (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide) immobilized on a solid support and with a given compound under conditions allowing the formation of an (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex, and

d) determining the amount of (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex formed in steps b) and c).

8. The method of claim 7, wherein a reduced amount of (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 formed in step c) in comparison to step b) indicates that LRRK2 is a target of the compound.

9. The method of any of claims 5 to 8, wherein the amount of (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex is determined by separating LRRK2 from

the immobilized (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide) and subsequent determination of the amount of LRRK2.

10. The method of claim 9, wherein the amount of LRRK2 is determined by mass spectrometry or immunodetection methods.

11. The method of any of claims 1 to 10, performed as a medium or high throughput screening.

12. The method of any of claims 1 to 11, wherein said compound is selected from the group consisting of synthetic compounds, or organic synthetic drugs, more preferably small molecule organic drugs, and natural small molecule compounds.

13. The method of any of claims 1 to 12, wherein the LRRK2 interacting compound is an LRRK2 inhibitor.

14. The method of any of claims 1 to 13, wherein the solid support is selected from the group consisting of agarose, modified agarose, sepharose beads (e.g. NHS-activated sepharose), latex, cellulose, and ferro- or ferrimagnetic particles.

15. The method of any of claims 1 to 14, wherein (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide) is covalently coupled to the solid support.

16. A method for the purification of LRRK2, comprising the steps of

a) providing a protein preparation containing LRRK2,

b) contacting the protein preparation with (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide) immobilized on a solid support under conditions allowing the formation of an (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex, and

c) separating LRRK2 from the immobilized (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide).

17. The method of claim 16, further comprising after step c) the identification of proteins being capable of binding to LRRK2.

18. The method of claim 16, further comprising after step c) the determination of post-translational modifications of LRRK2.

19. The method of any of claims 16 to 18, wherein after step c) either the LRRK2 and/or the proteins being capable of binding to LRRK2 are characterized by mass spectrometry or immunodetection methods.

20. The method of any of the preceding claims, wherein the provision of a protein preparation includes the steps of harvesting at least one cell containing LRRK2 and lysing the cell.

21. The method of any of the preceding claims, wherein the steps of the formation of the (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide)-LRRK2 complex are performed under essentially physiological conditions.

22. Use of (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide) for the identification of LRRK2 interacting compounds.

23. Use of (5-[5-Fluoro-2-oxo-1,2-dihydro-indol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (3-amino-propyl)-amide) for the purification of LRRK2.

SEQUENCE LISTING

<110> CellZome AG

<120> Methods for the identification of LRRK2 interacting molecules and for the purification of LRRK2

<130> CEL64132PC

<150> US 60/782,119
<151> 2006-03-14

<160> 13

<170> PatentIn version 3.3

<210> 1
<211> 2527
<212> PRT
<213> Mus musculus

<400> 1

```
Met Ala Ser Gly Ala Cys Gln Gly Cys Glu Glu Glu Glu Glu Glu Glu
1               5                  10                  15

Ala Leu Lys Lys Leu Ile Val Arg Leu Asn Asn Val Gln Glu Gly Lys
            20                  25                  30

Gln Ile Glu Thr Leu Leu Gln Leu Leu Glu Asp Met Leu Val Phe Thr
        35                  40                  45

Tyr Ser Asp Arg Ala Ser Lys Leu Phe Glu Asp Lys Asn Phe His Val
    50                  55                  60

Pro Leu Leu Ile Val Leu Asp Ser Tyr Met Arg Val Ala Ser Val Gln
65                  70                  75                  80

Gln Ala Gly Trp Ser Leu Leu Cys Lys Leu Ile Glu Val Cys Pro Gly
                85                  90                  95

Thr Leu Gln Ser Leu Ile Gly Pro Gln Asp Ile Gly Asn Asp Trp Glu
            100                 105                 110

Val Leu Gly Ile His Arg Leu Ile Leu Lys Met Leu Thr Val His His
        115                 120                 125

Ala Asn Val Asn Leu Ser Ile Val Gly Leu Lys Ala Leu Asp Leu Leu
    130                 135                 140

Leu Asp Ser Gly Lys Leu Thr Leu Leu Ile Leu Asp Glu Glu Cys Asp
145                 150                 155                 160

Ile Phe Leu Leu Ile Phe Asp Ala Met His Arg Tyr Ser Ala Asn Asp
                165                 170                 175

Glu Val Gln Lys Leu Gly Cys Lys Ala Leu His Val Leu Phe Glu Arg
            180                 185                 190
```

Val Ser Glu Glu Gln Leu Thr Glu Phe Val Glu Asn Lys Asp Tyr Thr
195 200 205

Ile Leu Leu Ser Thr Phe Gly Ser Phe Arg Arg Asp Lys Glu Ile Val
210 215 220

Tyr His Val Leu Cys Cys Leu His Ser Leu Ala Val Thr Cys Ser Asn
225 230 235 240

Val Glu Val Leu Met Ser Gly Asn Val Arg Cys Tyr Asn Leu Val Val
245 250 255

Glu Ala Met Lys Ala Phe Pro Thr Asn Glu Asn Ile Gln Glu Val Ser
260 265 270

Cys Ser Leu Phe Gln Lys Leu Thr Leu Gly Asn Phe Phe Asn Ile Leu
275 280 285

Val Leu Asn Glu Val His Val Phe Val Val Lys Ala Val Arg Gln Tyr
290 295 300

Pro Glu Asn Ala Ala Leu Gln Ile Ser Ala Leu Ser Cys Leu Ala Leu
305 310 315 320

Leu Thr Glu Thr Ile Phe Leu Asn Gln Asp Leu Glu Glu Arg Ser Glu
325 330 335

Thr Gln Glu Gln Ser Glu Glu Glu Asp Ser Glu Lys Leu Phe Trp Leu
340 345 350

Glu Pro Cys Tyr Lys Ala Leu Val Arg His Arg Lys Asp Lys His Val
355 360 365

Gln Glu Ala Ala Cys Trp Ala Leu Asn Asn Leu Leu Met Tyr Gln Asn
370 375 380

Ser Leu His Glu Lys Ile Gly Asp Glu Asp Gly Gln Phe Pro Ala His
385 390 395 400

Arg Glu Val Met Leu Ser Met Leu Met His Ser Ser Ser Lys Asp Val
405 410 415

Phe Gln Ala Ala Ala His Ala Leu Ser Thr Leu Leu Glu Gln Asn Val
420 425 430

Asn Phe Arg Lys Ile Leu Leu Ala Lys Gly Val Tyr Leu Asn Val Leu
435 440 445

Glu Leu Met Gln Lys His Ala His Ala Pro Glu Val Ala Glu Ser Gly
450 455 460

Cys Lys Met Leu Ser His Leu Phe Glu Gly Ser Asn Pro Ser Leu Asp
465 470 475 480

Thr Met Ala Ala Val Val Pro Lys Ile Leu Thr Val Met Lys Ala His
485 490 495

Gly Thr Ser Leu Ser Val Gln Leu Glu Ala Leu Arg Ala Ile Leu His
500 505 510

Phe Val Val Pro Gly Leu Leu Glu Glu Ser Arg Glu Asp Ser Gln Cys
515 520 525

Arg Pro Asn Val Leu Arg Lys Gln Cys Phe Arg Thr Asp Ile His Lys
530 535 540

Leu Val Leu Val Ala Leu Asn Arg Phe Ile Gly Asn Pro Gly Ile Gln
545 550 555 560

Lys Cys Gly Leu Lys Val Ile Ser Ser Leu Ala His Leu Pro Asp Ala
565 570 575

Thr Glu Thr Leu Ser Leu Gln Gly Ala Val Asp Ser Val Leu His Thr
580 585 590

Leu Gln Met Tyr Pro Asp Asp Gln Glu Ile Gln Cys Leu Gly Leu His
595 600 605

Leu Met Gly Cys Leu Met Thr Lys Lys Asn Phe Cys Ile Gly Thr Gly
610 615 620

His Leu Leu Ala Lys Ile Leu Ala Ser Thr Leu Gln Arg Phe Lys Asp
625 630 635 640

Val Ala Glu Val Gln Thr Thr Gly Leu Gln Thr Thr Leu Ser Ile Leu
645 650 655

Glu Leu Ser Val Ser Phe Ser Lys Leu Leu Val His Tyr Ser Phe Asp
660 665 670

Val Val Ile Phe His Gln Met Ser Ser Ser Val Val Glu Gln Lys Asp
675 680 685

Glu Gln Phe Leu Asn Leu Cys Cys Lys Cys Phe Ala Lys Val Ala Val
690 695 700

Asp Asp Glu Leu Lys Asn Thr Met Leu Glu Arg Ala Cys Asp Gln Asn
705 710 715 720

Asn Ser Ile Met Val Glu Cys Leu Leu Leu Leu Gly Ala Asp Ala Asn
725 730 735

Gln Val Lys Gly Ala Thr Ser Leu Ile Tyr Gln Val Cys Glu Lys Glu
740 745 750

Ser Ser Pro Lys Leu Val Glu Leu Leu Leu Asn Gly Gly Cys Arg Glu
755 760 765

Gln Asp Val Arg Lys Ala Leu Thr Ile Ser Ile Gln Lys Gly Asp Ser
770 775 780

Gln Val Ile Ser Leu Leu Leu Arg Lys Leu Ala Leu Asp Leu Ala Asn
785 790 795 800

Asn Ser Ile Cys Leu Gly Gly Phe Gly Ile Gly Lys Ile Asp Pro Ser
805 810 815

Trp Leu Gly Pro Leu Phe Pro Asp Lys Ser Ser Asn Leu Arg Lys Gln
820 825 830

Thr Asn Thr Gly Ser Val Leu Ala Arg Lys Val Leu Arg Tyr Gln Met
835 840 845

Arg Asn Thr Leu Gln Glu Gly Val Ala Ser Gly Ser Asp Gly Asn Phe
850 855 860

Ser Glu Asp Ala Leu Ala Lys Phe Gly Glu Trp Thr Phe Ile Pro Asp
865 870 875 880

Ser Ser Met Asp Ser Val Phe Gly Gln Ser Asp Asp Leu Asp Ser Glu
885 890 895

Gly Ser Glu Ser Ser Phe Leu Val Lys Arg Lys Ser Asn Ser Ile Ser
900 905 910

Val Gly Glu Val Tyr Arg Asp Leu Ala Leu Gln Arg Tyr Ser Pro Asn
915 920 925

Ala Gln Arg His Ser Asn Ser Leu Gly Pro Val Phe Asp His Glu Asp
930 935 940

Leu Leu Arg Arg Lys Arg Lys Ile Leu Ser Ser Asp Glu Ser Leu Arg
945 950 955 960

Ser Ser Arg Leu Pro Ser His Met Arg Gln Ser Asp Ser Ser Ser Ser
965 970 975

Leu Ala Ser Glu Arg Glu His Ile Thr Ser Leu Asp Leu Ser Ala Asn
980 985 990

Glu Leu Lys Asp Ile Asp Ala Leu Ser Gln Lys Cys Cys Leu Ser Ser
995 1000 1005

His Leu Glu His Leu Thr Lys Leu Glu Leu His Gln Asn Ser Leu

1010 1015 1020

Thr Ser Phe Pro Gln Gln Leu Cys Glu Thr Leu Lys Cys Leu Ile
1025 1030 1035

His Leu Asp Leu His Ser Asn Lys Phe Thr Ser Phe Pro Ser Phe
1040 1045 1050

Val Leu Lys Met Pro Arg Ile Thr Asn Leu Asp Ala Ser Arg Asn
1055 1060 1065

Asp Ile Gly Pro Thr Val Val Leu Asp Pro Ala Met Lys Cys Pro
1070 1075 1080

Ser Leu Lys Gln Leu Asn Leu Ser Tyr Asn Gln Leu Ser Ser Ile
1085 1090 1095

Pro Glu Asn Leu Ala Gln Val Val Glu Lys Leu Glu Gln Leu Leu
1100 1105 1110

Leu Glu Gly Asn Lys Ile Ser Gly Ile Cys Ser Pro Leu Ser Leu
1115 1120 1125

Lys Glu Leu Lys Ile Leu Asn Leu Ser Lys Asn His Ile Pro Ser
1130 1135 1140

Leu Pro Gly Asp Phe Leu Glu Ala Cys Ser Lys Val Glu Ser Phe
1145 1150 1155

Ser Ala Arg Met Asn Phe Leu Ala Ala Met Pro Ala Leu Pro Ser
1160 1165 1170

Ser Ile Thr Ser Leu Lys Leu Ser Gln Asn Ser Phe Thr Cys Ile
1175 1180 1185

Pro Glu Ala Ile Phe Ser Leu Pro His Leu Arg Ser Leu Asp Met
1190 1195 1200

Ser His Asn Asn Ile Glu Cys Leu Pro Gly Pro Ala His Trp Lys
1205 1210 1215

Ser Leu Asn Leu Arg Glu Leu Ile Phe Ser Lys Asn Gln Ile Ser
1220 1225 1230

Thr Leu Asp Phe Ser Glu Asn Pro His Val Trp Ser Arg Val Glu
1235 1240 1245

Lys Leu His Leu Ser His Asn Lys Leu Lys Glu Ile Pro Pro Glu
1250 1255 1260

Ile Gly Cys Leu Glu Asn Leu Thr Ser Leu Asp Val Ser Tyr Asn
1265 1270 1275

Leu Glu Leu Arg Ser Phe Pro Asn Glu Met Gly Lys Leu Ser Lys
1280 1285 1290

Ile Trp Asp Leu Pro Leu Asp Gly Leu His Leu Asn Phe Asp Phe
1295 1300 1305

Lys His Val Gly Cys Lys Ala Lys Asp Ile Ile Arg Phe Leu Gln
1310 1315 1320

Gln Arg Leu Lys Lys Ala Val Pro Tyr Asn Arg Met Lys Leu Met
1325 1330 1335

Ile Val Gly Asn Thr Gly Ser Gly Lys Thr Thr Leu Leu Gln Gln
1340 1345 1350

Leu Met Lys Met Lys Lys Pro Glu Leu Gly Met Gln Gly Ala Thr
1355 1360 1365

Val Gly Ile Asp Val Arg Asp Trp Ser Ile Gln Ile Arg Gly Lys
1370 1375 1380

Arg Arg Lys Asp Leu Val Leu Asn Val Trp Asp Phe Ala Gly Arg
1385 1390 1395

Glu Glu Phe Tyr Ser Thr His Pro His Phe Met Thr Gln Arg Ala
1400 1405 1410

Leu Tyr Leu Ala Val Tyr Asp Leu Ser Lys Gly Gln Ala Glu Val
1415 1420 1425

Asp Ala Met Lys Pro Trp Leu Phe Asn Ile Lys Ala Arg Ala Ser
1430 1435 1440

Ser Ser Pro Val Ile Leu Val Gly Thr His Leu Asp Val Ser Asp
1445 1450 1455

Glu Lys Gln Arg Lys Ala Cys Ile Ser Lys Ile Thr Lys Glu Leu
1460 1465 1470

Leu Asn Lys Arg Gly Phe Pro Thr Ile Arg Asp Tyr His Phe Val
1475 1480 1485

Asn Ala Thr Glu Glu Ser Asp Ala Leu Ala Lys Leu Arg Lys Thr
1490 1495 1500

Ile Ile Asn Glu Ser Leu Asn Phe Lys Ile Arg Asp Gln Pro Val
1505 1510 1515

Val Gly Gln Leu Ile Pro Asp Cys Tyr Val Glu Leu Glu Lys Ile
1520 1525 1530

Ile Leu Ser Glu Arg Lys Ala Val Pro Thr Glu Phe Pro Val Ile
1535 1540 1545

Asn Arg Lys His Leu Leu Gln Leu Val Asn Glu His Gln Leu Gln
1550 1555 1560

Leu Asp Glu Asn Glu Leu Pro His Ala Val His Phe Leu Asn Glu
1565 1570 1575

Ser Gly Val Leu Leu His Phe Gln Asp Pro Ala Leu Gln Leu Ser
1580 1585 1590

Asp Leu Tyr Phe Val Glu Pro Lys Trp Leu Cys Lys Val Met Ala
1595 1600 1605

Gln Ile Leu Thr Val Lys Val Asp Gly Cys Leu Lys His Pro Lys
1610 1615 1620

Gly Ile Ile Ser Arg Arg Asp Val Glu Lys Phe Leu Ser Lys Lys
1625 1630 1635

Lys Arg Phe Pro Lys Asn Tyr Met Met Gln Tyr Phe Lys Leu Leu
1640 1645 1650

Glu Lys Phe Gln Ile Ala Leu Pro Ile Gly Glu Glu Tyr Leu Leu
1655 1660 1665

Val Pro Ser Ser Leu Ser Asp His Arg Pro Val Ile Glu Leu Pro
1670 1675 1680

His Cys Glu Asn Ser Glu Ile Ile Ile Arg Leu Tyr Glu Met Pro
1685 1690 1695

Tyr Phe Pro Met Gly Phe Trp Ser Arg Leu Ile Asn Arg Leu Leu
1700 1705 1710

Glu Ile Ser Pro Phe Met Leu Ser Gly Arg Glu Arg Ala Leu Arg
1715 1720 1725

Pro Asn Arg Met Tyr Trp Arg Gln Gly Ile Tyr Leu Asn Trp Ser
1730 1735 1740

Pro Glu Ala Tyr Cys Leu Val Gly Ser Glu Val Leu Asp Asn Arg
1745 1750 1755

Pro Glu Ser Phe Leu Lys Ile Thr Val Pro Ser Cys Arg Lys Gly
1760 1765 1770

Cys Ile Leu Leu Gly Arg Val Val Asp His Ile Asp Ser Leu Met
1775 1780 1785

Glu Glu Trp Phe Pro Gly Leu Leu Glu Ile Asp Ile Cys Gly Glu
1790 1795 1800

Gly Glu Thr Leu Leu Lys Lys Trp Ala Leu Tyr Ser Phe Asn Asp
1805 1810 1815

Gly Glu Glu His Gln Lys Ile Leu Leu Asp Glu Leu Met Lys Lys
1820 1825 1830

Ala Glu Glu Gly Asp Leu Leu Ile Asn Pro Asp Gln Pro Arg Leu
1835 1840 1845

Thr Ile Pro Ile Ser Gln Ile Ala Pro Asp Leu Ile Leu Ala Asp
1850 1855 1860

Leu Pro Arg Asn Ile Met Leu Asn Asn Asp Glu Leu Glu Phe Glu
1865 1870 1875

Glu Ala Pro Glu Phe Leu Leu Gly Asp Gly Ser Phe Gly Ser Val
1880 1885 1890

Tyr Arg Ala Ala Tyr Glu Gly Glu Glu Val Ala Val Lys Ile Phe
1895 1900 1905

Asn Lys His Thr Ser Leu Arg Leu Leu Arg Gln Glu Leu Val Val
1910 1915 1920

Leu Cys His Leu His His Pro Ser Leu Ile Ser Leu Leu Ala Ala
1925 1930 1935

Gly Ile Arg Pro Arg Met Leu Val Met Glu Leu Ala Ser Lys Gly
1940 1945 1950

Ser Leu Asp Arg Leu Leu Gln Gln Asp Lys Ala Ser Leu Thr Arg
1955 1960 1965

Thr Leu Gln His Arg Ile Ala Leu His Val Ala Asp Gly Leu Arg
1970 1975 1980

Tyr Leu His Ser Ala Met Ile Ile Tyr Arg Asp Leu Lys Pro His
1985 1990 1995

Asn Val Leu Leu Phe Thr Leu Tyr Pro Asn Ala Ala Ile Ile Ala
2000 2005 2010

Lys Ile Ala Asp Tyr Gly Ile Ala Gln Tyr Cys Cys Arg Met Gly
2015 2020 2025

Ile Lys Thr Ser Glu Gly Thr Pro Gly Phe Arg Ala Pro Glu Val
2030 2035 2040

Ala Arg Gly Asn Val Ile Tyr Asn Gln Gln Ala Asp Val Tyr Ser

2045 2050 2055

Phe Gly Leu Leu Leu His Asp Ile Trp Thr Thr Gly Ser Arg Ile
2060 2065 2070

Met Glu Gly Leu Arg Phe Pro Asn Glu Phe Asp Glu Leu Ala Ile
2075 2080 2085

Gln Gly Lys Leu Pro Asp Pro Val Lys Glu Tyr Gly Cys Ala Pro
2090 2095 2100

Trp Pro Met Val Glu Lys Leu Ile Thr Lys Cys Leu Lys Glu Asn
2105 2110 2115

Pro Gln Glu Arg Pro Thr Ser Ala Gln Val Phe Asp Ile Leu Asn
2120 2125 2130

Ser Ala Glu Leu Ile Cys Leu Met Arg His Ile Leu Ile Pro Lys
2135 2140 2145

Asn Ile Ile Val Glu Cys Met Val Ala Thr Asn Leu Asn Ser Lys
2150 2155 2160

Ser Ala Thr Leu Trp Leu Gly Cys Gly Asn Thr Glu Lys Gly Gln
2165 2170 2175

Leu Ser Leu Phe Asp Leu Asn Thr Glu Arg Tyr Ser Tyr Glu Glu
2180 2185 2190

Val Ala Asp Ser Arg Ile Leu Cys Leu Ala Leu Val His Leu Ala
2195 2200 2205

Ala Glu Lys Glu Ser Trp Val Val Cys Gly Thr Gln Ser Gly Ala
2210 2215 2220

Leu Leu Val Ile Asn Val Glu Glu Glu Thr Lys Arg His Thr Leu
2225 2230 2235

Glu Lys Met Thr Asp Ser Val Thr Cys Leu His Cys Asn Ser Leu
2240 2245 2250

Ala Lys Gln Ser Lys Gln Ser Asn Phe Leu Leu Val Gly Thr Ala
2255 2260 2265

Asp Gly Asn Leu Met Ile Phe Glu Asp Lys Ala Val Lys Cys Lys
2270 2275 2280

Gly Ala Ala Pro Leu Lys Thr Leu His Ile Gly Asp Val Ser Thr
2285 2290 2295

Pro Leu Met Cys Leu Ser Glu Ser Leu Asn Ser Ser Glu Arg His
2300 2305 2310

```
Ile Thr  Trp Gly Gly Cys Gly  Thr Lys Val Phe Ser  Phe Ser Asn
    2315                 2320                 2325

Asp Phe  Thr Ile Gln Lys Leu  Ile Glu Thr Lys Thr  Asn Gln Leu
    2330                 2335                 2340

Phe Ser  Tyr Ala Ala Phe Ser  Asp Ser Asn Ile Ile  Ala Leu Ala
    2345                 2350                 2355

Val Asp  Thr Ala Leu Tyr Ile  Ala Lys Lys Asn Ser  Pro Val Val
    2360                 2365                 2370

Glu Val  Trp Asp Lys Lys Thr  Glu Lys Leu Cys Glu  Leu Ile Asp
    2375                 2380                 2385

Cys Val  His Phe Leu Lys Glu  Val Met Val Lys Leu  Asn Lys Glu
    2390                 2395                 2400

Ser Lys  His Gln Leu Ser Tyr  Ser Gly Arg Val Lys  Ala Leu Cys
    2405                 2410                 2415

Leu Gln  Lys Asn Thr Ala Leu  Trp Ile Gly Thr Gly  Gly Gly His
    2420                 2425                 2430

Ile Leu  Leu Leu Asp Leu Ser  Thr Arg Arg Val Ile  Arg Thr Ile
    2435                 2440                 2445

His Asn  Phe Cys Asp Ser Val  Arg Ala Met Ala Thr  Ala Gln Leu
    2450                 2455                 2460

Gly Ser  Leu Lys Asn Val Met  Leu Val Leu Gly Tyr  Lys Arg Lys
    2465                 2470                 2475

Ser Thr  Glu Gly Ile Gln Glu  Gln Lys Glu Ile Gln  Ser Cys Leu
    2480                 2485                 2490

Ser Ile  Trp Asp Leu Asn Leu  Pro His Glu Val Gln  Asn Leu Glu
    2495                 2500                 2505

Lys His  Ile Glu Val Arg Thr  Glu Leu Ala Asp Lys  Met Arg Lys
    2510                 2515                 2520

Thr Ser  Val Glu
    2525
```

<210> 2
<211> 9
<212> PRT
<213> artificial sequence

<220>
<223> peptide

<400> 2

Tyr Ser Ala Asn Asp Glu Val Gln Lys
1 5

<210> 3
<211> 13
<212> PRT
<213> artificial sequence

<220>
<223> peptide

<400> 3

His Ala His Ala Pro Glu Val Ala Glu Ser Gly Cys Lys
1 5 10

<210> 4
<211> 9
<212> PRT
<213> artificial sequence

<220>
<223> peptide

<400> 4

Phe Ile Gly Asn Pro Gly Ile Gln Lys
1 5

<210> 5
<211> 8
<212> PRT
<213> artificial sequence

<220>
<223> peptide

<400> 5

Ile Leu Ala Ser Thr Leu Gln Arg
1 5

<210> 6
<211> 12
<212> PRT
<213> artificial sequence

<220>
<223> peptide

<400> 6

Gly Ala Thr Ser Leu Ile Tyr Gln Val Cys Glu Lys
1 5 10

<210> 7
<211> 11
<212> PRT
<213> artificial sequence

<220>

<223> peptide

<400> 7

Gly Asp Ser Gln Val Ile Ser Leu Leu Leu Arg
1 5 10

<210> 8
<211> 10
<212> PRT
<213> artificial sequence

<220>
<223> peptide

<400> 8

Gln Thr Asn Thr Gly Ser Val Leu Ala Arg
1 5 10

<210> 9
<211> 20
<212> PRT
<213> artificial sequence

<220>
<223> peptide

<400> 9

Leu Glu Leu His Gln Asn Ser Leu Thr Ser Phe Pro Gln Gln Leu Cys
1 5 10 15

Glu Thr Leu Lys
20

<210> 10
<211> 12
<212> PRT
<213> artificial sequence

<220>
<223> peptide

<400> 10

Leu Leu Glu Ile Ser Pro Phe Met Leu Ser Gly Arg
1 5 10

<210> 11
<211> 6
<212> PRT
<213> artificial sequence

<220>
<223> peptide

<400> 11

Ile Thr Val Pro Ser Cys
1 5

<210> 12

<211> 9
<212> PRT
<213> artificial sequence

<220>
<223> peptide

<400> 12

Thr Ser Glu Gly Thr Pro Gly Phe Arg
1 5

<210> 13
<211> 8
<212> PRT
<213> artificial sequence

<220>
<223> peptide

<400> 13

His Gln Leu Ser Tyr Ser Gly Arg
1 5