



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07B 35/08, C07C 213/02, 217/18	A1	(11) International Publication Number: ✓ WO 97/26234 (43) International Publication Date: 24 July 1997 (24.07.97)
(21) International Application Number: PCT/GB97/00134 ✓ (22) International Filing Date: 20 January 1997 (20.01.97) ✓ (30) Priority Data: 9601167.1 20 January 1996 (20.01.96) ✓ GB 9618775.2 9 September 1996 (09.09.96) GB (71) Applicant (for all designated States except US): BRADFORD UNIVERSITY [GB/GB]; The Clinical Oncology Unit, Bradford, West Yorkshire BD7 1DP (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): DOUBLE, John [GB/GB]; Bradford University, Bradford, West Yorkshire BD7 1DP (GB). MAITLAND, Derek [GB/GB]; Bradford University, Bradford, West Yorkshire BD7 1DP (GB). POPA, Ioana [RO/GB]; Bradford University, Bradford, West Yorkshire BD7 1DP (GB). (74) Agents: BRIERLEY, Anthony, Paul et al.; Appleyard Lees, 15 Clare Road, Halifax HX1 2HY (GB).	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: TAMOXIFEN AND ANALOGUES THEREOF (57) Abstract A method of preparing a preferred geometric isomer of tamoxifen or an analogue comprises converting a first geometric isomer of a precursor to a second geometric isomer of the same precursor by contact with methanol and/or isopropanol and derivatising the second precursor to form tamoxifen or an antiestrogenic analogue thereof.		