



150150

4 - (3,4 - DISUBSTITUTED PHENYL) - PYRROLIDIN-2-ONE COMPOUNDS AS PHOSPHODIESTERASE 4 INHIBITORS

FIELD OF THE INVENTION

5 This application claims the benefit of U.S. Provisional Application Serial No. 60/463,054, filed April 16, 2003, the entire disclosure of which is hereby incorporated.

 This application is related to copending applications Serial No. 10/270,724, filed October 16, 2002, which claims the benefit of Serial No. 60/329,314, filed October 16,
10 2001, the entire disclosures of which are hereby incorporated.

 The present invention relates generally to the field of phosphodiesterase 4 (PDE4) enzyme inhibition. More specifically this invention relates to selective PDE4 inhibition by novel compounds, e.g., 4-(substituted-phenyl)-2-pyrrolidinone compounds, methods
15 of preparing such compounds, compositions containing such compounds, and methods of use thereof.

BACKGROUND OF THE INVENTION

20 The cyclic nucleotide specific phosphodiesterases (PDEs) represent a family of enzymes that catalyze the hydrolysis of various cyclic nucleoside monophosphates (including cAMP and cGMP). These cyclic nucleotides act as second messengers within cells, and as messengers, carry impulses from cell surface receptors having bound various hormones and neurotransmitters. PDEs act to regulate the level of cyclic nucleotides within
25 cells and maintain cyclic nucleotide homeostasis by degrading such cyclic mononucleotides resulting in termination of their messenger role.



 PDE enzymes can be grouped into eleven families according to their specificity toward hydrolysis of cAMP or cGMP, their sensitivity to regulation by calcium,
30 calmodulin or cGMP, and their selective inhibition by various compounds. For example, PDE 1 is stimulated by Ca^{2+} /calmodulin. PDE 2 is cGMP-dependent, and is found in the



heart and adrenals. PDE 3 is cGMP-inhibited, and inhibition of this enzyme creates positive inotropic activity. PDE 4 is cAMP specific, and its inhibition causes airway relaxation, anti-inflammatory and antidepressant activity. PDE 5 appears to be important in regulating cGMP content in vascular smooth muscle, and therefore PDE 5 inhibitors may have cardiovascular activity. Since the PDEs possess distinct biochemical properties, it is likely that they are subject to a variety of different forms of regulation.

PDE4 is distinguished by various kinetic properties including low Michaelis constant for cAMP and sensitivity to certain drugs. The PDE4 enzyme family consists of four genes, which produce 4 isoforms of the PDE4 enzyme designated PDE4A, PDE4B, PDE4C, and PDE4D [See: Wang et al., Expression, Purification, and Characterization of human cAMP-Specific Phosphodiesterase (PDE4) Subtypes A, B, C, and D, *Biochem. Biophys. Res. Comm.*, 234, 320-324 (1997)]. In addition, various splice variants of each PDE4 isoform have been identified.

PDE4 isoenzymes are localized in the cytosol of cells and are unassociated with any known membranous structures. PDE4 isoenzymes specifically inactivate cAMP by catalyzing its hydrolysis to adenosine 5'-monophosphate (AMP). Regulation of cAMP activity is important in many biological processes, including inflammation and memory. Inhibitors of PDE4 isoenzymes such as rolipram, piclamilast, CDP-840 and ariflo are powerful anti-inflammatory agents and therefore may be useful in treating diseases where inflammation is problematic such as asthma or arthritis. Further, rolipram improves the cognitive performance of rats and mice in learning paradigms.



rolipram

piclamilast

In addition to such compounds as rolipram, xanthine derivatives such as pentoxifylline, denbufylline, and theophylline inhibit PDE4 and have received attention of late for their cognition enhancing effects. cAMP and cGMP are second messengers that mediate cellular responses to many different hormones and neurotransmitters. Thus, therapeutically significant effects may result from PDE inhibition and the resulting increase in intracellular cAMP or cGMP in key cells, such as those located in the nervous system and elsewhere in the body.

Rolipram, previously in development as an anti-depressant, selectively inhibits the PDE4 enzyme and has become a standard agent in the classification of PDE enzyme subtypes. Early work in the PDE4 field focused on depression and inflammation, and has subsequently been extended to include indications such as dementia. [see "The PDE IV Family Of Calcium-Phosphodiesterases Enzymes," John A. Lowe, III, et al., *Drugs of the Future* 1992, 17(9):799-807 for a general review]. Further clinical developments of rolipram and other first-generation PDE4 inhibitors were terminated due to the side effect profile of these compounds. The primary side effect in primates is emesis, while the primary side effects in rodents are testicular degranulation, weakening of vascular smooth muscle, psychotropic effects, increased gastric acid secretion and stomach erosion. In humans, the primary side effect is nausea and emesis.

SUMMARY OF THE INVENTION


The present invention relates to novel compounds that inhibit, preferably selectively, PDE4 enzymes, and especially have improved side effect profiles, e.g., are relatively non-emetic (e.g., as compared to the previously discussed prior art compounds). In particular, the present invention relates to novel rolipram analogs. The compounds of this invention at the same time facilitate entry into cells, especially cells of the nervous system.

Still further, the present invention provides methods for synthesizing compounds with such activity and selectivity as well as methods of (and corresponding pharmaceutical compositions for) treating a patient, e.g., mammals, including humans, in need of PDE inhibition, especially PDE4 inhibition, for a disease state that involves elevated intracellular PDE 4 levels or decreased cAMP levels, e.g., involving neurological syndromes, especially those states associated with memory impairment, most especially long term memory impairment, as where such memory impairment is due at least in part to catabolism of intracellular cAMP levels by PDE 4 enzymes, or where such an impaired condition can be improved by increasing cAMP levels.

In a preferred aspect, the compounds of the invention improve such diseases by inhibiting PDE4 enzymes at doses that do not induce emesis or other side effects.

Upon further study of the specification and appended claims, further aspects, objects and advantages of this invention will become apparent to those skilled in the art.

The present invention includes compounds of Formula I:



wherein

R^1 is alkyl having 1 to 8 carbon atoms wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,

alkyl having 1 to 8 carbon atoms which is substituted one or more times by halogen, oxo or combinations thereof wherein optionally one or more

-CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,

5 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl having 1 to 4 carbon atoms or combinations thereof,

10 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom (e.g., 3-thienyl, 2-thienyl, 3-tetrahydrofuranyl), which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, or combinations thereof,

15 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,

20

25 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,

30 a partially unsaturated carbocyclic group having 5 to 14 carbon atoms, (e.g., cyclohexenyl, cyclohexadienyl, indanyl, and tetrahydronaphthenyl),

- which is unsubstituted or substituted one or more times by halogen, alkyl, alkoxy, nitro, cyano, oxo, or combinations thereof ,
- 5 arylalkenyl having 8 to 16 carbon atoms, wherein the alkenyl portion has up to 5 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., 10 acetamido), and acyloxy (e.g., acetoxy), or combinations thereof;
- 15 a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof, or
- 20 cycloalkylalkyl having 4 to 16 carbon atoms (e.g., cyclopentylethyl and cyclopropylmethyl), which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 25 R^2 is alkyl having 1 to 4 carbon atoms, which is unsubstituted or substituted one or more times by halogen;
- R^3 is H,
- 30 alkyl having 1 to 8 carbon atoms wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,

- alkyl having 1 to 8 carbon atoms which is substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,
- cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- arylalkyl having 7 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenyl, phenoxy, and acyloxy (e.g., acetoxy), or combinations thereof,
- heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof,
- alkoxyalkyl having 3 to 8 carbon atoms,
- C(O)R⁴, -(CH₂)_nC(O)R⁴, -(CH₂)_nOR⁵, -(CH₂)_nSR⁵, -(CH₂)_nSO₂R⁴, -(CH₂)_nNR⁵R⁶, -CH₂CO₂R⁵, -CH₂CONR⁶R⁵, -CH₂CONHR⁵, -(CH₂)_nNR⁶SO₂R⁴, -(CH₂)_nNR⁶COR⁴, or -CH₂CONHSO₂R⁴;

- R⁴ is alkyl having 1 to 12 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,
- 5 alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,
- 10 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- 15 cycloalkylalkyl having 4 to 16 carbon atoms (e.g., cyclopentylethyl and cyclopropylmethyl), which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 20 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, alkylsulphonamido (e.g., CH₃SO₂NH-, C₂H₅SO₂NH-), arylsulphonamido (e.g., C₆H₅SO₂NH-),
- 25 halogenated arylsulphonamido (e.g., 4-F-C₆H₅SO₂NH-), phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,
- 30 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy,

amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, aminosulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetox), or combinations thereof,

5

a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom (e.g., 3-thienyl, 2-thienyl, 3-tetrahydrofuranyl, 1,3,4-thiadiazolyl), which is unsubstituted or substituted one or more times by halogen, aryl (e.g., phenyl, methylphenyl), alkyl, cycloalkyl (e.g., cyclopropyl), cycloalkylalkyl (e.g., cyclopropylmethyl), alkoxy, alkoxyalkyl (e.g., methoxymethyl), cyano, halogenated alkyl (e.g., trifluoromethyl), halogenated alkoxy (e.g., trifluoromethoxy), nitro, oxo, amino, alkylamino, dialkylamino, aminosulphonyl, heterocycle (e.g., pyridyl, piperidinyl, thienyl, tetrahydrofuranyl, pyrazinyl), heterocyclic-alkyl (e.g., thienylmethyl, piperidinylcarbonylmethyl), or combinations thereof, or

10

15

a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof; and

20

25

R⁵ is H,

alkyl having 1 to 12 carbon atoms wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,

30

- alkyl having 1 to 12 carbon atoms which is substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,
- alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,
- cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- cycloalkylalkyl having 4 to 16 carbon atoms (e.g., cyclopentylethyl and cyclopropylmethyl), which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, alkylsulphonamido (e.g., CH₃SO₂NH-, C₂H₅SO₂NH-), arylsulphonamido (e.g., C₆H₅SO₂NH-), halogenated arylsulphonamido (e.g., 4-F-C₆H₅SO₂NH-), phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,
- arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen,

- 5 CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, aminosulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,
- 10 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom (e.g., 3-thienyl, 2-thienyl, 3-tetrahydrofuranyl, 1,3,4-thiadiazolyl), which is unsubstituted or substituted one or more times by halogen, aryl (e.g., phenyl, methylphenyl), alkyl, cycloalkyl (e.g., cyclopropyl), cycloalkylalkyl (e.g., cyclopropylmethyl), alkoxy, alkoxyalkyl (e.g., methoxymethyl), cyano, halogenated alkyl (e.g., trifluoromethyl), halogenated alkoxy (e.g., trifluoromethoxy), nitro, oxo,
- 15 amino, alkylamino, dialkylamino, aminosulphonyl, heterocycle (e.g., pyridyl, piperidiny, thienyl, tetrahydrofuranyl, pyrazinyl), heterocyclic-alkyl (e.g., thienylmethyl, piperidinylcarbonylmethyl), or combinations thereof, or
- 20 a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino,
- 25 dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof;

R⁶ is H,

- alkyl having 1 to 12 carbon atoms wherein optionally one or more $\text{-CH}_2\text{CH}_2\text{-}$ groups are replaced in each case by -CH=CH- or $\text{-C}\equiv\text{C-}$ groups,
- 5 alkyl having 1 to 12 carbon atoms which is substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $\text{-CH}_2\text{CH}_2\text{-}$ groups are replaced in each case by -CH=CH- or $\text{-C}\equiv\text{C-}$ groups,
- 10 alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $\text{-CH}_2\text{CH}_2\text{-}$ groups are replaced in each case by -CH=CH- or $\text{-C}\equiv\text{C-}$ groups,
- 15 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- 20 cycloalkylalkyl having 4 to 16 carbon atoms (e.g., cyclopentylethyl and cyclopropylmethyl), which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 25 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,
- 30 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy,

amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof;

5

a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom (e.g., 3-thienyl, 2-thienyl, 3-tetrahydrofuranyl), which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, alkoxy, alkoxycarbonyl, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, or combinations thereof, or

10

a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof;

15

20

n is 0 to 4 (e.g., 1 to 4 or 2 to 4); and

pharmaceutically acceptable salts thereof;

25

with the proviso that:

- (a) when R^2 is CH_3 and R^3 is H, then R^1 is not methyl, ethyl, n-propyl, isopropyl, sec-butyl, n-butyl, isobutyl, neopentyl, n-pentyl, 2-methylbutyl, isopentyl, n-hexyl, phenyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopentenyl, methylcyclopentyl, cyclopropylmethyl,

30

- 5 cyclopentylmethyl, 2-propenyl, 2-propynyl, 3-methyl-2-butenyl, N-substituted 2-piperazinylethyl, norbornyl, 3-tetrahydrofuryl, 2-tetrahydrofuryl, 3-tetrahydrothienyl, 2-oxacyclopropyl, 2-oxacyclopentyl, 3-oxacyclopentyl, 2-chloroethyl, 2-bromoethyl, 2,2,2-trifluoroethyl, 3-bromopropyl, 3-chloropropyl, or 4-bromobutyl;
- 10 (b) when R^1 is cyclopentyl, and R^2 is methyl, then R^3 is not H, acetyl, benzyl, 4-hydroxybenzyl, 4-acetoxybenzyl, 4-bromobenzyl, 3,4-dimethoxybenzyl, 4-methylthiobenzyl, 4-cyanobenzyl, 2-aminobenzyl, 3-aminobenzyl, 4-aminobenzyl, 4-dimethylaminobenzyl, 2,4-diaminobenzyl, 4-amino-3,5-dimethoxybenzyl, 3-
- 15 carboxybenzyl, 3-methoxycarbonylbenzyl, 4-methoxycarbonylbenzyl, 4-methylsulfinylbenzyl, 4-methylsulfonylbenzyl, 2-nitrobenzyl, 3-nitrobenzyl, 4-nitrobenzyl, 2,4-dinitrobenzyl, 2-nitro-4-aminobenzyl, 2-amino-4-nitrobenzyl, morpholinoethyl, 2-
- 20 pyridylmethyl, 3-pyridylmethyl, 4-pyridylmethyl, 4-(6-fluoroquinolyl)methyl, 2-(7-chloroquinolyl)methyl, 2-imidazolylmethyl, or substituted imidazolylmethyl;
- 25 (c) when R^1 is CH_3 , and R^3 is H, then R^2 is not methyl, ethyl, or butyl;
- (d) when R^3 is H, then R^1 and R^2 are not both ethyl or isobutyl; and
- 30 (e) When R^1 and R^2 are both difluoromethyl, then R^3 is not 4-aminobenzyl, or 4-amino-3,5-dimethoxybenzyl.

- aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,
- arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,
- a partially unsaturated carbocyclic group having 5 to 14 carbon atoms, (e.g., cyclohexenyl, cyclohexadienyl, indanyl, and tetrahydronaphthenyl), which is unsubstituted or substituted one or more times by halogen, alkyl, alkoxy, nitro, cyano, oxo, or combinations thereof ,
- arylalkenyl having 8 to 16 carbon atoms, wherein the alkenyl portion has up to 5 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof;
- a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an

N, O or S atom, which is unsubstituted or substituted one or more times in
 the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano,
 halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino,
 dialkylamino, carboxy or combinations thereof and/or substituted in the
 5 alkyl portion by halogen, oxo, cyano, or combinations thereof, or

 cycloalkylalkyl having 4 to 16 carbon atoms (e.g., cyclopentylethyl and
 cyclopropylmethyl), which is unsubstituted or substituted one or more
 times by halogen, oxo, alkyl or combinations thereof,
 10

 R^2 is alkyl having 1 to 4 carbon atoms, which is unsubstituted or
 substituted one or more times by halogen;

 R^3 is $-(CH_2)_nOR^5$, $-(CH_2)_nSR^5$, $-(CH_2)_nSO_2R^4$, $-(CH_2)_nNR^5R^6$, $-CH_2CO_2R^5$, -
 15 $CH_2CONR^6R^5$, $-(CH_2)_nNR^6SO_2R^4$, $-(CH_2)_nNR^6COR^4$, or -
 $CH_2CONHSO_2R^4$;

 R^4 is alkyl having 1 to 12 carbon atoms which is unsubstituted or substituted
 one or more times by halogen, oxo, or combinations thereof wherein
 20 optionally one or more $-CH_2CH_2-$ groups are replaced in each case by
 $-CH=CH-$ or $-C\equiv C-$ groups,

 alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or
 substituted one or more times by halogen, oxo, or combinations thereof
 25 wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each
 case by $-CH=CH-$ or $-C\equiv C-$ groups,

 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or
 substituted one or more times by halogen, oxo, alkyl, or combinations
 30 thereof,

- cycloalkylalkyl having 4 to 16 carbon atoms (e.g., cyclopentylethyl and cyclopropylmethyl), which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 5 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, alkylsulphonamido (e.g.,
- 10 CH₃SO₂NH-, C₂H₅SO₂NH-), arylsulphonamido (e.g., C₆H₅SO₂NH-), halogenated arylsulphonamido (e.g., 4-F-C₆H₅SO₂NH-), phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,
- 15 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl,
- 20 aminosulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,
- a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N,
- 25 O or S atom (e.g., 3-thienyl, 2-thienyl, 3-tetrahydrofuranyl, 1,3,4-thiadiazolyl), which is unsubstituted or substituted one or more times by halogen, aryl (e.g., phenyl, methylphenyl), alkyl, cycloalkyl (e.g., cyclopropyl), cycloalkylalkyl (e.g., cyclopropylmethyl), alkoxy, alkoxyalkyl (e.g., methoxymethyl), cyano, halogenated alkyl (e.g.,
- 30 trifluoromethyl), halogenated alkoxy (e.g., trifluoromethoxy), nitro, oxo, amino, alkylamino, dialkylamino, aminosulphonyl, heterocycle (e.g.,

pyridyl, piperidinyl, thienyl, tetrahydrofuranyl, pyrazinyl), heterocyclic-alkyl (e.g., thienylmethyl, piperidinylcarbonylmethyl), or combinations thereof, or

5 a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, 10 dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof; and

R⁵ is H,

15 alkyl having 1 to 12 carbon atoms wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,

20 alkyl having 1 to 12 carbon atoms which is substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,

25 alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,

30 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,

- cycloalkylalkyl having 4 to 16 carbon atoms (e.g., cyclopentylethyl and cyclopropylmethyl), which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 5 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, alkylsulphonamido (e.g., CH₃SO₂NH-, C₂H₅SO₂NH-), arylsulphonamido (e.g., C₆H₅SO₂NH-), halogenated arylsulphonamido (e.g., 4-F-C₆H₅SO₂NH-), phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,
- 10 15 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, aminosulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,
- 20 25 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom (e.g., 3-thienyl, 2-thienyl, 3-tetrahydrofuranyl, 1,3,4-thiadiazolyl), which is unsubstituted or substituted one or more times by halogen, aryl (e.g., phenyl, methylphenyl), alkyl, cycloalkyl (e.g., cyclopropyl), cycloalkylalkyl (e.g., cyclopropylmethyl), alkoxy, alkoxyalkyl (e.g., methoxymethyl), cyano, halogenated alkyl (e.g., trifluoromethyl), halogenated alkoxy (e.g., trifluoromethoxy), nitro, oxo,
- 30

- amino, alkylamino, dialkylamino, aminosulphonyl, heterocycle (e.g., pyridyl, piperidinyl, thienyl, tetrahydrofuranyl, pyrazinyl), heterocyclic-alkyl (e.g., thienylmethyl, piperidinylcarbonylmethyl), or combinations thereof, or
- 5 a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano,
- 10 halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof;
- R^6 is H,
- 15 alkyl having 1 to 12 carbon atoms wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,
- 20 alkyl having 1 to 12 carbon atoms which is substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,
- 25 alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,

- cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- 5 cycloalkylalkyl having 4 to 16 carbon atoms (e.g., cyclopentylethyl and cyclopropylmethyl), which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 10 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,
- 15 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof;
- 20 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom (e.g., 3-thienyl, 2-thienyl, 3-tetrahydrofuranyl), which is
- 25 unsubstituted or substituted one or more times by halogen, aryl, alkyl, alkoxy, alkoxycarbonyl, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, or combinations thereof, or
- 30 a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an

5 N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof,

wherein if R^3 is $-\text{CH}_2\text{CONR}^6\text{R}^5$, R^6 is other than H; and

10 n is 0 to 4 (e.g., 1 to 4 or 2 to 4); and

pharmaceutically acceptable salts thereof.

15 In accordance with a further aspect of the invention, the R^4 group of the compound of formula I is an alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-\text{CH}_2\text{CH}_2-$ groups are replaced in each case by $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$ groups.

20 In accordance with a further aspect of the invention, the R^5 group of the compound of formula I is an alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-\text{CH}_2\text{CH}_2-$ groups are replaced in each case by $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$ groups.

25 In accordance with a further aspect of the invention, at least one of R^1 , R^4 or R^5 in formula I is a heterocyclic group which is substituted by halogenated alkyl other than trifluoromethyl, or at least one of R^1 , R^3 , R^4 or R^5 in formula I is a heterocyclic-alkyl group which is substituted by halogenated alkyl other than trifluoromethyl.

30

In accordance with a further aspect of the invention, R^3 in formula I is a arylalkyl group which is substituted by phenyl.


In accordance with a further aspect of the invention, at least one of R^4 and R^5 in formula I is aryl substituted by alkylsulphonamido (e.g., $\text{CH}_3\text{SO}_2\text{NH-}$, $\text{C}_2\text{H}_5\text{SO}_2\text{NH-}$), arylsulphonamido (e.g., $\text{C}_6\text{H}_5\text{SO}_2\text{NH-}$), or halogenated arylsulphonamido, arylalkyl which is substituted by aminosulphonyl, or a heterocyclic group which is substituted one or more times by alkoxyalkyl (e.g., methoxymethyl), cycloalkyl (e.g., cyclopropyl), cycloalkylalkyl (e.g., cyclopropylmethyl), halogenated alkoxy, aminosulphonyl, heterocycle (e.g., pyridyl, piperidiny, thienyl, tetrahydrofuranyl, pyrazinyl), heterocyclic-alkyl (e.g., thienylmethyl), or combinations thereof.

In accordance with a further aspect of the invention, n in formula I is 0 or 1.

In accordance with the method aspect of the invention, there is provided a method of treating a patient (e.g., a mammal such as a human) suffering from a disease state (e.g., memory impairment) involving decreased cAMP levels and/or increased intracellular PDE4 levels, comprising administering to the patient a compound selected from the compounds according to formula I':

20

25



wherein

R^1 is alkyl having 1 to 8 carbon atoms wherein optionally one or more $-\text{CH}_2\text{CH}_2-$ groups are replaced in each case by $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$ groups,

- alkyl having 1 to 8 carbon atoms which is substituted one or more times by halogen, oxo or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,
- cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl having 1 to 4 carbon atoms or combinations thereof,
- a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom (e.g., 3-thienyl, 2-thienyl, 3-tetrahydrofuranyl), which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, or combinations thereof,
- aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,
- arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,

- 5 a partially unsaturated carbocyclic group having 5 to 14 carbon atoms,
(e.g., cyclohexenyl, cyclohexadienyl, indanyl, and tetrahydronaphthenyl),
which is unsubstituted or substituted one or more times by halogen, alkyl,
alkoxy, nitro, cyano, oxo, or combinations thereof ,
- 10 arylalkenyl having 8 to 16 carbon atoms, wherein the alkenyl portion has
up to 5 carbon atoms, which is unsubstituted or substituted preferably in
the aryl portion, one or more times by halogen, alkyl, hydroxy, alkoxy,
nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino,
hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl,
alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g.,
acetamido), and acyloxy (e.g., acetoxy), or combinations thereof;
- 15 a heterocyclic-alkyl group, which is saturated, partially saturated or fully
unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an
N, O or S atom, which is unsubstituted or substituted one or more times in
the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano,
halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino,
20 dialkylamino, carboxy or combinations thereof and/or substituted in the
alkyl portion by halogen, oxo, cyano, or combinations thereof, or
- 25 cycloalkylalkyl having 4 to 16 carbon atoms (e.g., cyclopentylethyl and
cyclopropylmethyl), which is unsubstituted or substituted one or more
times by halogen, oxo, alkyl or combinations thereof,
- R^2 is alkyl having 1 to 4 carbon atoms, which is unsubstituted or
substituted one or more times by halogen;
- 30 R^3 is H,

- alkyl having 1 to 8 carbon atoms wherein optionally one or more $-\text{CH}_2\text{CH}_2-$ groups are replaced in each case by $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$ groups,
- 5 alkyl having 1 to 8 carbon atoms which is substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-\text{CH}_2\text{CH}_2-$ groups are replaced in each case by $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$ groups,
- 10 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- 15 arylalkyl having 7 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenyl, phenoxy, and acyloxy (e.g., acetoxy), or combinations thereof,
- 20 heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano,
- 25 halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof,
- 30 alkoxyalkyl having 3 to 8 carbon atoms,

-C(O)R⁴, -(CH₂)_nC(O)R⁴, -(CH₂)_nOR⁵, -(CH₂)_nSR⁵, -(CH₂)_nSO₂R⁴, -(CH₂)_nNR⁵R⁶, -CH₂CO₂R⁵, -CH₂CONR⁶R⁵, -CH₂CONHR⁵, -(CH₂)_nNR⁶SO₂R⁴, -(CH₂)_nNR⁶COR⁴, or -CH₂CONHSO₂R⁴;

- 5 R⁴ is alkyl having 1 to 12 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,
- 10 alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,
- 15 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- 20 cycloalkylalkyl having 4 to 16 carbon atoms (e.g., cyclopentylethyl and cyclopropylmethyl), which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 25 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, alkylsulphonamido (e.g., CH₃SO₂NH-, C₂H₅SO₂NH-), arylsulphonamido (e.g., C₆H₅SO₂NH-), halogenated arylsulphonamido (e.g., 4-F-C₆H₅SO₂NH-), phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,
- 30

- 5 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, aminosulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,
- 10 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom (e.g., 3-thienyl, 2-thienyl, 3-tetrahydrofuranyl, 1,3,4-thiadiazolyl), which is unsubstituted or substituted one or more times by halogen, aryl (e.g., phenyl, methylphenyl), alkyl, cycloalkyl (e.g., cyclopropyl), cycloalkylalkyl (e.g., cyclopropylmethyl), alkoxy, alkoxyalkyl (e.g., methoxymethyl), cyano, halogenated alkyl (e.g., trifluoromethyl), halogenated alkoxy (e.g., trifluoromethoxy), nitro, oxo, amino, alkylamino, dialkylamino, aminosulphonyl, heterocycle (e.g., pyridyl, piperidinyl, thienyl, tetrahydrofuranyl, pyrazinyl), heterocyclic-alkyl (e.g., thienylmethyl, piperidinylcarbonylmethyl), or combinations thereof, or
- 20 a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof; and
- 30 R⁵ is H,

- alkyl having 1 to 12 carbon atoms wherein optionally one or more
-CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C-
groups,
- 5
- alkyl having 1 to 12 carbon atoms which is substituted one or more times
by halogen, oxo, or combinations thereof wherein optionally one or more
-CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C-
groups,
- 10
- alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or
substituted one or more times by halogen, oxo, or combinations thereof
wherein optionally one or more -CH₂CH₂- groups are replaced in each
case by -CH=CH- or -C≡C- groups,
- 15
- cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or
substituted one or more times by halogen, oxo, alkyl, or combinations
thereof,
- 20
- cycloalkylalkyl having 4 to 16 carbon atoms (e.g., cyclopentylethyl and
cyclopropylmethyl), which is unsubstituted or substituted one or more
times by halogen, oxo, alkyl or combinations thereof,
- 25
- aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one
or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro,
methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino,
hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl,
alkylthio, alkylsulphinyl, alkylsulphonyl, alkylsulphonamido (e.g.,
CH₃SO₂NH-, C₂H₅SO₂NH-), arylsulphonamido (e.g., C₆H₅SO₂NH-),
30 halogenated arylsulphonamido (e.g., 4-F-C₆H₅SO₂NH-), phenoxy,

acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,

5 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, aminosulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g.,
10 acetoxy), or combinations thereof,

a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom (e.g., 3-thienyl, 2-thienyl, 3-tetrahydrofuranyl, 1,3,4-
15 thiadiazolyl), which is unsubstituted or substituted one or more times by halogen, aryl (e.g., phenyl, methylphenyl), alkyl, cycloalkyl (e.g., cyclopropyl), cycloalkylalkyl (e.g., cyclopropylmethyl), alkoxy, alkoxyalkyl (e.g., methoxymethyl), cyano, halogenated alkyl (e.g., trifluoromethyl), halogenated alkoxy (e.g., trifluoromethoxy), nitro, oxo,
20 amino, alkylamino, dialkylamino, aminosulphonyl, heterocycle (e.g., pyridyl, piperidinyl, thienyl, tetrahydrofuranyl, pyrazinyl), heterocyclic-alkyl (e.g., thienylmethyl, piperidinylcarbonylmethyl), or combinations thereof, or

25 a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino,
30 dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof;

R⁶ is H,

5 alkyl having 1 to 12 carbon atoms wherein optionally one or more
-CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C-
groups,

10 alkyl having 1 to 12 carbon atoms which is substituted one or more times
by halogen, oxo, or combinations thereof wherein optionally one or more
-CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C-
groups,

15 alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or
substituted one or more times by halogen, oxo, or combinations thereof
wherein optionally one or more -CH₂CH₂- groups are replaced in each
case by -CH=CH- or -C≡C- groups,

20 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or
substituted one or more times by halogen, oxo, alkyl, or combinations
thereof,

25 cycloalkylalkyl having 4 to 16 carbon atoms (e.g., cyclopentylethyl and
cyclopropylmethyl), which is unsubstituted or substituted one or more
times by halogen, oxo, alkyl or combinations thereof,
aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one
or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro,
methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino,
hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl,
alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g.,
30 acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,

5 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof;

10 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom (e.g., 3-thienyl, 2-thienyl, 3-tetrahydrofuranyl), which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, alkoxy, alkoxycarbonyl, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, or combinations thereof, or

15 a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, 20 halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof;

25 n is 0 to 4 (e.g., 1 to 4 or 2 to 4); and
pharmaceutically acceptable salts thereof.

30 According to a further method aspect, in Formula I', when R² is CH₃ and R³ is H, then R¹ is not cyclopentyl.

In accordance with a further method aspect of the invention, the R⁴ group of the compound of formula I is an alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups.


In accordance with a further method aspect of the invention, the R⁵ group of the compound of formula I is an alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups.

In accordance with a further method aspect of the invention, at least one of R¹, R⁴ or R⁵ in formula I is a heterocyclic group which is substituted by halogenated alkyl other than trifluoromethyl, or at least one of R¹, R³, R⁴ or R⁵ in formula I is a heterocyclic-alkyl group which is substituted by halogenated alkyl other than trifluoromethyl.

In accordance with a further method aspect of the invention, R³ in formula I is a arylalkyl group which is substituted by phenyl.

In accordance with a further method aspect of the invention, at least one of R⁴ and R⁵ in formula I is aryl substituted by alkylsulphonamido (e.g., CH₃SO₂NH-, C₂H₅SO₂NH-), arylsulphonamido (e.g., C₆H₅SO₂NH-), or halogenated arylsulphonamido, arylalkyl which is substituted by aminosulphonyl, or a heterocyclic group which is substituted one or more times by alkoxyalkyl (e.g., methoxymethyl), cycloalkyl (e.g., cyclopropyl), cycloalkylalkyl (e.g., cyclopropylmethyl), halogenated alkoxy, aminosulphonyl, heterocycle (e.g., pyridyl, piperidinyl, thienyl, tetrahydrofuranyl, pyrazinyl), heterocyclic-alkyl (e.g., thienylmethyl), or combinations thereof.

In accordance with the method aspect of the invention, there is also provided a method of treating a patient (e.g., a mammal such as a human) suffering from a disease state (e.g., memory impairment) involving decreased cAMP levels and/or increased intracellular PDE4 levels, comprising administering to the patient a compound according to formula II':



wherein

R¹ is alkyl having 1 to 8 carbon atoms wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,

alkyl having 1 to 8 carbon atoms which is substituted one or more times by halogen, oxo or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,

cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl having 1 to 4 carbon atoms or combinations thereof,

a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom (e.g., 3-thienyl, 2-thienyl, 3-tetrahydrofuranyl), which is unsubstituted or substituted one or more times by halogen, aryl, alkyl,

alkoxy, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, or combinations thereof,

5 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,

10 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, 15 cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,

20 a partially unsaturated carbocyclic group having 5 to 14 carbon atoms, (e.g., cyclohexenyl, cyclohexadienyl, indanyl, and tetrahydronaphthenyl), which is unsubstituted or substituted one or more times by halogen, alkyl, alkoxy, nitro, cyano, oxo, or combinations thereof ,

25 arylalkenyl having 8 to 16 carbon atoms, wherein the alkenyl portion has up to 5 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., 30 acetamido), and acyloxy (e.g., acetoxy), or combinations thereof;

- a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano,
- 5 halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof, or
- cycloalkylalkyl having 4 to 16 carbon atoms (e.g., cyclopentylethyl and cyclopropylmethyl), which is unsubstituted or substituted one or more
- 10 times by halogen, oxo, alkyl or combinations thereof,
- R^2 is alkyl having 1 to 4 carbon atoms, which is unsubstituted or substituted one or more times by halogen;
- 15 R^3 is $-(CH_2)_nOR^5$, $-(CH_2)_nSR^5$, $-(CH_2)_nSO_2R^4$, $-(CH_2)_nNR^5R^6$, $-CH_2CO_2R^5$, $-CH_2CONR^6R^5$, $-(CH_2)_nNR^6SO_2R^4$, $-(CH_2)_nNR^6COR^4$, or $-CH_2CONHSO_2R^4$;
- 20 R^4 is alkyl having 1 to 12 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,
- 25 alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,

- cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- 5 cycloalkylalkyl having 4 to 16 carbon atoms (e.g., cyclopentylethyl and cyclopropylmethyl), which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 10 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, alkylsulphonamido (e.g., CH₃SO₂NH-, C₂H₅SO₂NH-), arylsulphonamido (e.g., C₆H₅SO₂NH-),
- 15 halogenated arylsulphonamido (e.g., 4-F-C₆H₅SO₂NH-), phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,
- 20 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, aminosulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g.,
- 25 acetoxy), or combinations thereof,
- 30 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom (e.g., 3-thienyl, 2-thienyl, 3-tetrahydrofuranyl, 1,3,4-thiadiazolyl), which is unsubstituted or substituted one or more times by halogen, aryl (e.g., phenyl, methylphenyl), alkyl, cycloalkyl (e.g.,

cyclopropyl), cycloalkylalkyl (e.g., cyclopropylmethyl), alkoxy,
 alkoxyalkyl (e.g., methoxymethyl), cyano, halogenated alkyl (e.g.,
 trifluoromethyl), halogenated alkoxy (e.g., trifluoromethoxy), nitro, oxo,
 amino, alkylamino, dialkylamino, aminosulphonyl, heterocycle (e.g.,
 5 pyridyl, piperidinyl, thienyl, tetrahydrofuranyl, pyrazinyl), heterocyclic-
 alkyl (e.g., thienylmethyl, piperidinylcarbonylmethyl), or combinations
 thereof, or

a heterocyclic-alkyl group, which is saturated, partially saturated or fully
 10 unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an
 N, O or S atom, which is unsubstituted or substituted one or more times in
 the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano,
 halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino,
 dialkylamino, carboxy or combinations thereof and/or substituted in the
 15 alkyl portion by halogen, oxo, cyano, or combinations thereof; and

R⁵ is H,

alkyl having 1 to 12 carbon atoms wherein optionally one or more
 20 -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C-
 groups,

alkyl having 1 to 12 carbon atoms which is substituted one or more times
 by halogen, oxo, or combinations thereof wherein optionally one or more
 25 -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C-
 groups,

alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or
 substituted one or more times by halogen, oxo, or combinations thereof
 30 wherein optionally one or more -CH₂CH₂- groups are replaced in each
 case by -CH=CH- or -C≡C- groups,

- cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- 5 cycloalkylalkyl having 4 to 16 carbon atoms (e.g., cyclopentylethyl and cyclopropylmethyl), which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 10 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, alkylsulphonamido (e.g.,
- 15 CH₃SO₂NH-, C₂H₅SO₂NH-), arylsulphonamido (e.g., C₆H₅SO₂NH-), halogenated arylsulphonamido (e.g., 4-F-C₆H₅SO₂NH-), phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,
- 20 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl,
- 25 aminosulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,
- 30 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom (e.g., 3-thienyl, 2-thienyl, 3-tetrahydrofuranyl, 1,3,4-thiadiazolyl), which is unsubstituted or substituted one or more times by

- 5 halogen, aryl (e.g., phenyl, methylphenyl), alkyl, cycloalkyl (e.g., cyclopropyl), cycloalkylalkyl (e.g., cyclopropylmethyl), alkoxy, alkoxyalkyl (e.g., methoxymethyl), cyano, halogenated alkyl (e.g., trifluoromethyl), halogenated alkoxy (e.g., trifluoromethoxy), nitro, oxo, amino, alkylamino, dialkylamino, aminosulphonyl, heterocycle (e.g., pyridyl, piperidinyl, thienyl, tetrahydrofuranyl, pyrazinyl), heterocyclic-alkyl (e.g., thienylmethyl, piperidinylcarbonylmethyl), or combinations thereof, or
- 10 a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the
- 15 alkyl portion by halogen, oxo, cyano, or combinations thereof;
- R^6 is H,
- 20 alkyl having 1 to 12 carbon atoms wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,
- 25 alkyl having 1 to 12 carbon atoms which is substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,
- 30 alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof

wherein optionally one or more $\text{-CH}_2\text{CH}_2\text{-}$ groups are replaced in each case by -CH=CH- or $\text{-C}\equiv\text{C-}$ groups,

5 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,

10 cycloalkylalkyl having 4 to 16 carbon atoms (e.g., cyclopentylethyl and cyclopropylmethyl), which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,

15 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof,

20 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted preferably in the aryl portion, one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido (e.g., acetamido), and acyloxy (e.g., acetoxy), or combinations thereof;

25 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom (e.g., 3-thienyl, 2-thienyl, 3-tetrahydrofuranyl), which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, 30 alkoxy, alkoxycarbonyl, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, or combinations thereof, or

a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano,
5 halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof,
10 wherein if R^3 is $-CH_2CONR^6R^5$, R^6 is other than H;
n is 0 to 4 (e.g., 1 to 4 or 2 to 4) ; and
pharmaceutically acceptable salts thereof.

15 The compounds of the present invention are effective in inhibiting, or modulating the activity of PDE4 in animals, e.g., mammals, especially humans. These compounds exhibit neurological activity, especially where such activity affects cognition, including long term memory. These compounds will also be effective in treating diseases where
20 decreased cAMP levels are involved. This includes but is not limited to inflammatory diseases. These compounds may also function as antidepressants, or be useful in treating cognitive and negative symptoms of schizophrenia.

Assays for determining PDE 4 inhibiting activity, selectivity of PDE4 inhibiting
25 activity, and selectivity of inhibiting PDE 4 isoenzymes are known within the art. See, e.g., US 6,136,821, the disclosure of which is incorporated herein by reference.

Halogen herein refers to F, Cl, Br, and I. Preferred halogens are F and Cl.

30 Alkyl means a straight-chain or branched-chain aliphatic hydrocarbon radical. In the case of R^1 and R^3 suitable alkyl groups have preferably 1 to 8 carbon atoms,

especially 1 to 4 carbon atoms. In the case of R^2 , suitable alkyl groups have preferably 1 to 4 carbon atoms such as methyl, ethyl, propyl, isopropyl, and butyl. In the case of R^4 , R^5 , and R^6 , suitable alkyl groups have preferably 1 to 12 carbon atoms, especially 1 to 8 carbon atoms, in particular 1 to 4 carbon atoms.

5

Suitable alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, tert-butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, and dodecyl. Other examples of suitable alkyl groups include 1-, 2- or 3-methylbutyl, 1,1-, 1,2- or 2,2-dimethylpropyl, 1-ethylpropyl, 1-, 2-, 3- or 4-methylpentyl, 1,1-, 1,2-, 1,3-, 2,2-, 2,3- or 3,3-dimethylbutyl, 1- or 2-ethylbutyl, ethylmethylpropyl, trimethylpropyl, methylhexyl, dimethylpentyl, ethylpentyl, ethylmethylbutyl, dimethylbutyl, and the like.

These alkyl radicals can optionally have one or more $-CH_2CH_2-$ groups replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups. Suitable alkenyl or alkynyl groups are 1-propenyl, 2-propenyl, 1-propynyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-butyne, 1,3-butadienyl and 3-methyl-2-butenyl. Also, these alkyl radicals can optionally have one or more $-CH_2-$ groups replaced in each case by O, S, or NH.

The substituted alkyl groups for R^1 , R^3 , R^4 , R^5 , and R^6 are alkyl groups as described above which are substituted in one or more positions by halogens and/or oxo. Halogens are preferred substituents, especially F and Cl. For R^2 , the substituted alkyl groups have 1 to 4 carbon atoms and are substituted one or more times by halogen, especially F and Cl, e.g., C_1-C_4 alkyl substituted by up to five F atoms.

In the arylalkyl groups, heterocyclic-alkyl groups, cycloalkyl-alkyl groups and alkoxyalkyl groups, "alkyl" refers to a divalent alkylene group having in general up to about 13 carbon atoms. In the case of the arylalkyl group for R^1 , the "alkyl" portion has preferably 2 to 10 carbon atoms. In the case of other arylalkyl groups, the "alkyl" portion has preferably 1 to 10. In the heterocyclic-alkyl groups, the "alkyl" portion preferably has 1 to 12 carbon atoms. In the alkoxyalkyl groups, the "alkyl" portion preferably has 2

to 7 carbon atoms. In the cycloalkylalkyl groups, the “alkyl” portion preferably has 1 to 13 carbon atoms.

In the cases where alkyl is a substituent (e.g., alkyl substituents on aryl and heterocyclic groups) or is part of a substituent (e.g., in the alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, alkylthio, alkylsulphinyl, and alkylsulphonyl substituents for aryl), the alkyl portion preferably has 1 to 12 carbon atoms, especially 1 to 8 carbon atoms, in particular 1 to 4 carbon atoms. The alkyl groups, as described above, can be substituted in one or more positions by halogens and/or oxo. Halogens are preferred substituents, especially F and Cl.

Alkoxy means alkyl-O- groups in which the alkyl portion has 1 to 8 carbon atoms and which, for example, can be substituted by halogens. Suitable alkoxy groups include methoxy, ethoxy, propoxy, isopropoxy, isobutoxy, sec-butoxy, pentoxy, hexoxy, heptoxy, octoxy, and trifluoromethoxy. Preferred alkoxy groups are methoxy and ethoxy. The alkoxyalkyl group for R³ preferably has 3 to 8 carbon atoms, e.g., methoxyethyl.

Similarly, alkoxycarbonyl means an alkyl-O-CO- group in which the alkyl portion has 1 to 8 carbon atoms.

Alkenyl refers to straight-chain or branched-chain aliphatic radicals containing 2 to 12 carbon atoms in which one or more -CH₂-CH₂- structures is replaced by -CH=CH-. Suitable alkenyl groups are ethenyl, 1-propenyl, 2-methylethenyl, 1-butene, 2-butene, 1-pentenyl, and 2-pentenyl. In the arylalkenyl groups, alkenyl refers to a divalent alkenylene group having preferably 2 to 5 carbon atoms.

Cycloalkyl means a monocyclic, bicyclic or tricyclic saturated hydrocarbon radical having 3 to 8 carbon atoms, preferably 4 to 6 carbon atoms, especially 5 carbon atoms. Suitable cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, and norbornyl. Other suitable cycloalkyl groups

include spiropentyl, bicyclo[2.1.0]pentyl, bicyclo[3.1.0]hexyl, spiro[2.4]heptyl, spiro[2.5]octyl, bicyclo[5.1.0]octyl, spiro[2.6]nonyl, bicyclo[2.2.0]hexyl, spiro[3.3]heptyl, and bicyclo[4.2.0]octyl. The preferred cycloalkyl group is cyclopentyl.

- 5 The cycloalkyl group can be substituted by halogens, oxo and/or alkyl. Halogens and/or alkyl groups are preferred substituents.

- Cycloalkylalkyl refers to a cycloalkyl-alkyl-radical in which the cycloalkyl and alkyl portions are in accordance with the previous descriptions. Suitable examples
10 include cyclopentylethyl and cyclopropylmethyl.

- Aryl, as a group or substituent per se or as part of a group or substituent, refers to an aromatic carbocyclic radical containing 6 to 14 carbon atoms, preferably 6 to 12 carbon atoms, especially 6 to 10 carbon atoms. Suitable aryl groups include phenyl,
15 naphthyl and biphenyl. Substituted aryl groups include the above-described aryl groups which are substituted one or more times by, for example, halogen, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenyl, phenoxy, and acyloxy (e.g., acetoxy).

20

 Arylalkyl refers to an aryl-alkyl-radical in which the aryl and alkyl portions are in accordance with the previous descriptions. Suitable examples include 1-phenethyl, 2-phenethyl, phenpropyl, phenbutyl, phenpentyl, and naphthylenemethyl. Also, in the case of R^3 , arylalkyl can be benzyl.

25

 Arylalkenyl refers to an aryl-alkenyl-radical in which the aryl and alkenyl portions are in accordance with the previous descriptions of aryl and alkenyl. Suitable examples include 3-aryl-2-propenyl.

- 30 Heterocyclic groups refer to saturated, partially saturated and fully unsaturated heterocyclic groups having one or two rings and a total number of 5 to 10 ring atoms

- wherein at least one of the ring atoms is an N, O or S atom. Preferably, the heterocyclic group has one ring and has 5, 6, or 7 ring atoms or has two rings and is 5,6 fused or 6,6 fused. Preferably, the heterocyclic group contains 1 to 3, especially 1 or 2, hetero-ring atoms selected from N, O and S. Suitable saturated and partially saturated heterocyclic groups include, but are not limited to tetrahydrofuranyl, tetrahydrothienyl, pyrrolidinyl, isoxazolinyl and the like. Suitable heteroaryl groups include but are not limited to triazolyl (e.g., 1,2,4-triazolyl), thiadiazolyl (e.g., 1,3,4-thiadiazolyl), thiazolyl, oxazolyl, azatriazolyl, furyl, thienyl, pyrrolyl, pyrazolyl, imidazolyl, pyridyl, pyrimidinyl, indolyl, quinolinyl, naphthyridinyl, benzimidazolyl, benzthiazolyl, benzoxazolyl and the like.
- 10 Preferred heterocyclic and heteroaryl groups include tetrahydrofuranyl, tetrahydropyranyl, 2-thienyl, 3-thienyl, 2-, 3- or 4-pyridyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, and 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolinyl. In addition, heterocyclic groups include bicycloheterocyclic groups such as quinuclidinyl, 1-azabicyclo[2.2.1]heptyl, and 1-azabicyclo[3.2.1]octyl.
- 15 Substituted heterocyclic groups refer to the heterocyclic groups described above which are substituted in one or more places by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl (e.g., trifluoromethyl), nitro, oxo, amino, alkylamino, dialkylamino, and carboxy.
- 20 Heterocyclic-alkyl refers to a heterocyclic-alkyl-group wherein the heterocyclic and alkyl portions are in accordance with the previous discussions. Suitable examples are pyridylmethyl, thienylmethyl, pyrimidinylmethyl, pyrazinylmethyl, isoquinolinylmethyl, pyridylethyl and thienylethyl.
- 25 Partially unsaturated carbocyclic structures are non-aromatic monocyclic or bicyclic structures containing 5 to 14 carbon atoms, preferably 6 to 10 carbon atoms, wherein the ring structure(s) contains at least one C=C bond. Suitable examples are cyclopentenyl, cyclohexenyl, tetrahydronaphthenyl and indan-2-yl.

Acyl refers to alkanoyl radicals having 1 to 13 carbon atoms in which the alkyl portion can be substituted by halogen, hydroxy, carboxy, alkyl, aryl and/or alkoxy; or aroyl radicals having 7 to 15 carbon atoms in which the aryl portion can be substituted by halogen, alkyl, alkoxy, nitro, carboxy and/or hydroxy. Suitable acyl groups include
5 formyl, acetyl, propionyl, butanoyl and benzoyl.

Substituted radicals preferably have 1 to 3 substituents, especially 1 or 2 substituents.

10 R^1 is preferably optionally substituted cycloalkyl, cycloalkylalkyl, aryl, heterocyclic, arylalkyl, or a partially unsaturated carbocyclic group, or is CHF_2 ; for example, CHF_2 , cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopentylethyl, cyclopropylmethyl, cyclohexylpropyl, phenyl, naphthyl, phenethyl, phenpropyl, furanyl, pyrazinyl, pyrimidinyl, pyridyl, quinolinyl, isoquinolinyl, thienyl, indanyl,
15 tetrahydrofuranyl, phenylpropenyl, substituted phenyl, and substituted phenethyl. Preferred substituents are oxo, F, Cl, CF_3 , alkyl (such as methyl or ethyl), alkoxy (such as methoxy and ethoxy), CN, vinyl, methylenedioxy, COOH, and combinations thereof. When R^3 is other than H, R^1 is preferably cycloalkyl, cycloalkylalkyl or aryl, as well as CHF_2 , heterocyclic and arylalkyl. In such a case, R^1 is especially optionally substituted
20 cyclopentyl and phenethyl as well as 3-tetrahydrofuranyl, CHF_2 , and cyclopropylmethyl. More preferably, R^1 is 3-tetrahydrofuranyl, in particular, 3-(R)-tetrahydrofuranyl.

R^2 is preferably substituted or unsubstituted alkyl having 1 to 4 C atoms, most preferably CHF_2 and CH_3 .
25

R^3 is preferably H, substituted or unsubstituted alkyl, alkoxyalkyl, substituted or unsubstituted arylalkyl, CH_2CONHR^5 or COR^4 , especially H, substituted benzyl or CH_2CONHR^5 . Examples of suitable R^3 groups include methyl, ethyl, propyl, n-butyl, methoxyethyl, phenpropyl (e.g., 3-phenpropyl), methylbenzyl (e.g., 2-methylbenzyl), butylbenzyl (e.g., 4-tert-butylbenzyl), fluorobenzyl (e.g., 2-fluorobenzyl), difluorobenzyl (e.g., 2,3- or 2,6-difluorobenzyl), chlorobenzyl, methoxybenzyl, cyanobenzyl, dichlorobenzyl, chlorofluorobenzyl, trifluoromethylbenzyl, and $(CF_3)_2$ benzyl. R^3 can also be $-(CH_2)_nOR^5$ (such as $-CH_2CH_2OR^5$), $-(CH_2)_nSR^5$ (such as $-CH_2CH_2SR^5$), $-(CH_2)_nSO_2R^4$, $-(CH_2)_nNR^5R^6$ (such as $-CH_2CH_2NR^6R^5$, $-CH_2CO_2R^5$, $-CH_2CH_2CO_2R^5$, $-CH_2CONR^6R^5$, $-CH_2CONHR^5$, $-(CH_2)_nNR^6SO_2R^4$, $-(CH_2)_nNR^6COR^4$ (such as $CH_2CH_2NHCOR^4$), or $-CH_2CONHSO_2R^4$, for example, 2-phenoxyethyl, 2-phenthioethyl, 2-phenylsulfonyl ethyl, 4-methoxyphenyloxyethyl, phenylsulfonylaminocarbonylmethyl, and 2-methylphenylsulfonylaminocarbonylmethyl. More preferably, R^3 is CH_2CONHR^5 or $CH_2CONR^6R^5$.

15

R^4 , if present, is preferably optionally substituted aryl such as phenyl, or is optionally substituted heterocyclic (e.g., phenoxyphenyl, difluorophenyl, methoxyphenyl). Preferred substituents for R^4 are F, Cl, CH_3 , C_2H_5 , OCH_3 , $COOH$, and CN. In addition, in formula I R^4 can also be alkoxyalkyl having 3 to 8 carbon atoms which is substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups.

20

R⁵, if present, is preferably 2-methylphenyl, 2,6-dimethylphenyl, 2,3-difluorophenyl, 4-fluorophenyl, 4-pyridyl, 2-pyridyl, 6-methyl-2-pyridyl, 6-amino-2-pyridyl, 6-ethyl-2-pyridyl, 4,6-dimethyl-2-pyridyl, 6-bromo-2-pyridyl, 6-methyl-5-bromo-2-pyridyl, 2-methoxy-3-pyridyl, 4-CH₃O-CO-3-pyridyl, and 2-cyano-3-pyridyl.

5 More preferably, R⁵ is 4,5-dimethyltriazole, 4-isopropylphenyl, or 3,4-methylenedioxyphenyl. In addition, in formula I R⁵ can also be alkoxyalkyl having 3 to 8 carbon atoms which is substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups.

10

Other suitable R⁵ groups, for example when R³ is CH₂CONHR⁵, include 3-chlorophenyl, 4-chloro-2-fluorophenyl, 3-methylphenyl, 4-methylphenyl, 4-methoxyphenyl, 4-nitrophenyl, 3-bromo-2-methyl-6-pyridyl, 4-methoxycarbonyl-3-pyridyl, 4,5-dimethylthiazol-2-yl, 3-methoxycarbonylphenyl, 4-carboxyphenyl, 2-15 thiazolyl, 4-isopropylphenyl, 3,4-methylenedioxyphenyl, 4-trifluoromethylpyridyl, 3-carboxyphenyl, 5-chlorobenzoxazolyl, benzthiazolyl, 6-fluorobenzthiazolyl, and benzimidazolyl.

When R³ is CH₂CONHR⁵, preferred R⁵ groups are, for example, 2-thiazolyl or 20 1,3,4-thiadiazolyl, which in each case is unsubstituted or substituted by F, Cl, CF₃, methyl, methoxymethyl, isopropyl, isopbutyl, t-butyl, , carboxy, alkoxycarbonyl, cyclopropyl, cyclopropylmethyl, phenyl, pyridyl, piperidinyl, 3,4-methylenedioxyphenyl, thienyl, pyrazinyl, tetrahydrofuranyl, morpholinyl, pyrrolidinyl, or thienylmethyl. If substituted, the 2-thiazolyl group is preferably substituted in the 4 or 5 position and the 25 1,3,4-thiadiazolyl group is preferably substituted in the 5 position.

R⁶ is preferably H, C₁-C₅ alkyl (such as CH₃ or 2-methylpropyl) or cycloalkylalkyl. More preferably, R⁶ is cyclopropylmethyl.

In addition, preferred PDE4 inhibitors in accordance with the invention are compounds described by subformulas Ia-Iy, IIa, IId, and IIj-IIy, which correspond to formulas I and II respectively, but exhibit the following preferred groups:

5	Ia, IIa	R^2 is CH_3 or CHF_2 .
	Ib	R^2 is CH_3 or CHF_2 , and R^3 is H.
10	Ic	R^3 is H; and R^1 is optionally substituted cycloalkyl, heterocyclic group, heterocyclic alkyl group, arylalkyl or cycloalkylalkyl, or is CHF_2 .
15	Id, IId	R^2 is CH_3 or CHF_2 ; R^1 is cyclopentyl, tetrahydrofuranyl, or cyclopropylmethyl, as well as CHF_2 and phenethyl; and R^3 is not H.
20	Ie	R^2 is CH_3 or CHF_2 R^3 is H; and R^1 is phenethyl which is unsubstituted or the phenyl portion is substituted by F, Cl, Br, I, CN, C_{1-4} -alkyl and/or C_{1-4} -alkoxy,
25		phenpropyl which is unsubstituted or the phenyl portion is substituted by F, Cl, Br, I, CN, C_{1-4} -alkyl and/or C_{1-4} -alkoxy, phenylbutyl which is unsubstituted or the phenyl portion is substituted by F, Cl, Br, I, CN, C_{1-4} -alkyl and/or C_{1-4} -alkoxy, or 3-phenyl-2-propenyl.
30	If	R^1 is tetrahydrofuranyl (e.g., 3(R)- tetrahydrofuranyl);

R^2 is CH_3 or CHF_2 ; and

R^3 is arylalkyl in which the aryl group is substituted by F, Br, Cl, I, alkyl or alkoxy.

5	Ig	R^1 is cycloalkyl; R^2 is CH_3 or CHF_2 ; and R^3 is arylalkyl in which the aryl group is substituted by F, Br, Cl, I, alkyl or alkoxy.
10	Ih	R^1 is cyclopentyl, tetrahydrofuranyl, or cyclopropylmethyl, as well as CHF_2 and phenethyl; R^2 is CH_3 or CHF_2 ; and R^3 is arylalkyl in which the aryl group is substituted by F, Br, Cl, I, alkyl or alkoxy.
15	Ii	R^1 is cyclopentyl, tetrahydrofuranyl, cyclopropylmethyl or CHF_2 , as well as phenethyl; R^2 is CH_3 or CHF_2 ; and R^3 is methylbenzyl, methoxybenzyl, chlorobenzyl, fluorobenzyl, trifluorobenzyl, difluorobenzyl, dichlorobenzyl, fluorochlorobenzyl, or bis(trifluoromethyl)benzyl.
20	Ij, IIj	R^1 is cycloalkyl, heterocyclic group, or heterocyclic alkyl group, as well as CHF_2 , cycloalkylalkyl, and arylalkyl; R^2 is CH_3 or CHF_2 ; and R^3 is $\text{CH}_2\text{CONHR}^5$.
25		
30	Ik, Ilk	R^1 is cyclopentyl, tetrahydrofuranyl, or cyclopropylmethyl, as well as CHF_2 and phenethyl; R^2 is CH_3 or CHF_2 ; and R^3 is $\text{CH}_2\text{CONHR}^5$.

5	II, III	R^1 is cycloalkyl, heterocyclic group, or heterocyclicalkyl group, as well as CHF_2 and cycloalkylalkyl; R^2 is CH_3 or CHF_2; R^3 is $\text{CH}_2\text{CONHR}^5$; and R^5 is substituted or unsubstituted phenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-thiazolyl, or 1,3,4-thiadiazolyl.
10	Im, IIIm	R^1 is cycloalkyl, heterocyclic group, or heterocyclicalkyl group, as well as CHF_2 and cycloalkylalkyl; R^2 is CH_3 or CHF_2; R^3 is $\text{CH}_2\text{CONHR}^5$; and R^5 is 2-methylphenyl, 2,6-dimethylphenyl, 2,3-difluorophenyl, 4-fluorophenyl, 4-pyridyl, 2-pyridyl, 6-methyl-2-pyridyl, 6-amino-2-pyridyl, 6-ethyl-2-pyridyl, 4,6-dimethyl-2-pyridyl, 6-bromo-2-pyridyl, 6-methyl-5-bromo-2-pyridyl, 2-methoxy-3-pyridyl, 4-$\text{CH}_3\text{O-CO-3-pyridyl}$, 2-cyano-3-pyridyl, 4-isopropylphenyl, 3-fluorophenyl, 3-chlorophenyl, 2-thiazolyl, 2-(4,5-dimethyl)thiazolyl, 2-benzthiazolyl, 2-(6-fluoro)-benzthiazolyl, 2-thiazolyl, 2-thiazolyl which is substituted by F, Cl, CF_3, methyl, methoxymethyl, isopropyl, isopbutyl, t-butyl, carboxy, alkoxycarbonyl, cyclopropyl, cyclopropylmethyl, phenyl, pyridyl, piperidinyl, 3,4-methylenedioxyphenyl, thienyl, pyrazinyl, tetrahydrofuranyl, morpholinyl, pyrrolidinyl, or thienylmethyl, 1,3,4-thiadiazolyl, or 1,3,4-thiadiazolyl which is substituted by F, Cl, CF_3, methyl, methoxymethyl, isopropyl, isopbutyl, t-butyl, carboxy, alkoxycarbonyl, cyclopropyl, cyclopropylmethyl, phenyl, pyridyl, piperidinyl, 3,4-methylenedioxyphenyl, thienyl, pyrazinyl, tetrahydrofuranyl, morpholinyl, pyrrolidinyl, or thienylmethyl.
30		

5	In, IIn	R^1 is cyclopentyl, tetrahydrofuran, or cyclopropylmethyl as well as CHF_2; R^2 is CH_3 or CHF_2; R^3 is $\text{CH}_2\text{CONHR}^5$; and R^5 is substituted or unsubstituted phenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl, 2-thiazolyl or 1,3,4-thiadiazolyl.
10	Io, IIo	R^1 is cyclopentyl, tetrahydrofuranyl, or cyclopropylmethyl, as well as CHF_2; R^2 is CH_3 or CHF_2; R^3 is $\text{CH}_2\text{CONHR}^5$; and R^5 is 2-methylphenyl, 2,6-dimethylphenyl, 2,3-difluorophenyl, 4-fluorophenyl, 4-pyridyl, 2-pyridyl, 6-methyl-2-pyridyl, 6-amino-2-pyridyl, 6-ethyl-2-pyridyl, 4,6-dimethyl-2-pyridyl, 6-bromo-2-pyridyl, 6-methyl-5-bromo-2-pyridyl, 2-methoxy-3-pyridyl, 4-$\text{CH}_3\text{O-CO-3-pyridyl}$, 2-cyano-3-pyridyl, 4-isopropylphenyl, 3-fluorophenyl, 3-chlorophenyl, 2-thiazolyl, 2-(4,5-dimethyl)thiazolyl, 2-benzthiazolyl, 2-(6-fluoro)-benzthiazolyl, 2-thiazolyl, 2-thiazolyl which is substituted by F, Cl, CF_3, methyl, methoxymethyl, isopropyl, isopbutyl, t-butyl, carboxy, alkoxycarbonyl, cyclopropyl, cyclopropylmethyl, phenyl, pyridyl, piperidinyl, 3,4-methylenedioxyphenyl, thienyl, pyrazinyl, tetrahydrofuranyl, morpholinyl, pyrrolidinyl, or thienylmethyl, 1,3,4-thiadiazolyl, or 1,3,4-thiadiazolyl which is substituted by F, Cl, CF_3, methyl, methoxymethyl, isopropyl, isopbutyl, t-butyl, carboxy, alkoxycarbonyl, cyclopropyl, cyclopropylmethyl, phenyl, pyridyl, piperidinyl, 3,4-methylenedioxyphenyl, thienyl, pyrazinyl, tetrahydrofuranyl, morpholinyl, pyrrolidinyl, or thienylmethyl.
30	Ip, IIp	R^1 is cyclopentyl, tetrahydrofuranyl, or cyclopropylmethyl, as well as CHF_2;

		R^2 is CH_3 or CHF_2 ; and R^3 is $-(\text{CH}_2)_n\text{OR}^5$, $-(\text{CH}_2)_n\text{SR}^5$, $-(\text{CH}_2)_n\text{SO}_2\text{R}^4$, $-(\text{CH}_2)_n\text{NR}^5\text{R}^6$, $-\text{CH}_2\text{CO}_2\text{R}^5$, $-\text{CH}_2\text{CONR}^6\text{R}^5$, $-\text{CH}_2\text{NR}^6\text{SO}_2\text{R}^4$, $-\text{CH}_2\text{NR}^6\text{COR}^4$, or $\text{CH}_2\text{CONHSO}_2\text{R}^4$.
5	Iq, IIq	R^1 is cyclopentyl, tetrahydrofuran, or cyclopropylmethyl, as well as CHF_2 ; R^2 is CH_3 or CHF_2 ; R^3 is $-(\text{CH}_2)_n\text{OR}^5$, $-(\text{CH}_2)_n\text{SR}^5$, $-(\text{CH}_2)_n\text{SO}_2\text{R}^4$, $-(\text{CH}_2)_n\text{NR}^5\text{R}^6$, $-\text{CH}_2\text{CO}_2\text{R}^5$, $-\text{CH}_2\text{CONR}^6\text{R}^5$, $-\text{CH}_2\text{NR}^6\text{SO}_2\text{R}^4$, $-\text{CH}_2\text{NR}^6\text{COR}^4$, or $\text{CH}_2\text{CONHSO}_2\text{R}^4$. R^4 is substituted or unsubstituted phenyl; R^5 is H, alkyl, substituted or unsubstituted phenyl or pyridyl, or a substituted or unsubstituted heterocyclic group; and R^6 is alkyl, cycloalkyl, or cycloalkylalkyl.
10		
15		
	Ir, IIr	R^1 is cycloalkyl, heterocyclic group, or heterocyclic alkyl group, as well as CHF_2 , cycloalkylalkyl, and arylalkyl; R^2 is CH_3 or CHF_2 ; and R^3 is $\text{CH}_2\text{CONHR}^5$ or $-\text{CH}_2\text{CONR}^6\text{R}^5$.
20		
	Is, IIs	R^1 is cyclopentyl, tetrahydrofuranyl, or cyclopropylmethyl, as well as CHF_2 and phenethyl; R^2 is CH_3 or CHF_2 ; and R^3 is $\text{CH}_2\text{CONHR}^5$ or $-\text{CH}_2\text{CONR}^6\text{R}^5$.
25		
	It, IIIt	R^1 is cycloalkyl, heterocyclic group, or heterocyclicalkyl group, as well as CHF_2 and cycloalkylalkyl; R^2 is CH_3 or CHF_2 ; R^3 is $\text{CH}_2\text{CONHR}^5$ or $-\text{CH}_2\text{CONR}^6\text{R}^5$; and
30		

R⁵ is substituted or unsubstituted phenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-thiazolyl, or 1,3,4-thiadiazolyl.

Iu, Iiu

5

R¹ is cycloalkyl, heterocyclic group, or heterocyclicalkyl group, as well as CHF₂ and cycloalkylalkyl;

R² is CH₃ or CHF₂;

R³ is CH₂CONHR⁵ or -CH₂CONR⁶R⁵; and

10

R⁵ is 2-methylphenyl, 2,6-dimethylphenyl, 2,3-difluorophenyl, 4-fluorophenyl, 4-pyridyl, 2-pyridyl, 6-methyl-2-pyridyl, 6-amino-2-pyridyl, 6-ethyl-2-pyridyl, 4,6-dimethyl-2-pyridyl, 6-bromo-2-pyridyl, 6-methyl-5-bromo-2-pyridyl, 2-methoxy-3-pyridyl, 4-CH₃O-CO-3-pyridyl, 2-cyano-3-pyridyl, 4-isopropylphenyl, 3-fluorophenyl, 3-chlorophenyl, 2-thiazolyl, 2-(4,5-dimethyl)thiazolyl, 2-benzthiazolyl, 2-(6-fluoro)-benzthiazolyl, 2-thiazolyl, 2-thiazolyl which is substituted by F, Cl, CF₃, methyl, methoxymethyl, isopropyl, isopbutyl, t-butyl, carboxy, alkoxycarbonyl, cyclopropyl, cyclopropylmethyl, phenyl, pyridyl, piperidiny, 3,4-methylenedioxyphenyl, thienyl, pyrazinyl, tetrahydrofuranyl, morpholinyl, pyrrolidinyl, or thienylmethyl, 1,3,4-thiadiazolyl, or 1,3,4-thiadiazolyl which is substituted by F, Cl, CF₃, methyl, methoxymethyl, isopropyl, isopbutyl, t-butyl, carboxy, alkoxycarbonyl, cyclopropyl, cyclopropylmethyl, phenyl, pyridyl, piperidiny, 3,4-methylenedioxyphenyl, thienyl, pyrazinyl, tetrahydrofuranyl, morpholinyl, pyrrolidinyl, or thienylmethyl.

15

20

25

Iv, Iiv

R¹ is cyclopentyl, tetrahydrofuran, or cyclopropylmethyl as well as CHF₂;

R² is CH₃ or CHF₂;

R³ is CH₂CONHR⁵ or -CH₂CONR⁶R⁵; and

30

R⁵ is substituted or unsubstituted phenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl, 2-thiazolyl, or 1,3,4-thiadiazolyl.

Iw, IIw	R^1 is cyclopentyl, tetrahydrofuranyl, or cyclopropylmethyl, as well as CHF_2; R^2 is CH_3 or CHF_2; R^3 is $\text{CH}_2\text{CONHR}^5$ or $-\text{CH}_2\text{CONR}^6\text{R}^5$; and R^5 is 2-methylphenyl, 2,6-dimethylphenyl, 2,3-difluorophenyl, 4-fluorophenyl, 4-pyridyl, 2-pyridyl, 6-methyl-2-pyridyl, 6-amino-2-pyridyl, 6-ethyl-2-pyridyl, 4,6-dimethyl-2-pyridyl, 6-bromo-2-pyridyl, 6-methyl-5-bromo-2-pyridyl, 2-methoxy-3-pyridyl, 4-$\text{CH}_3\text{O-CO-3-pyridyl}$, 2-cyano-3-pyridyl, 4-isopropylphenyl, 3-fluorophenyl, 3-chlorophenyl, 2-thiazolyl, 2-(4,5-dimethyl)thiazolyl, 2-benzthiazolyl, 2-(6-fluoro)-benzthiazolyl, 2-thiazolyl, 2-thiazolyl which is substituted by F, Cl, CF_3, methyl, methoxymethyl, isopropyl, isopbutyl, t-butyl, carboxy, alkoxycarbonyl, cyclopropyl, cyclopropylmethyl, phenyl, pyridyl, piperidinyl, 3,4-methylenedioxyphenyl, thienyl, pyrazinyl, tetrahydrofuranyl, morpholinyl, pyrrolidinyl, or thienylmethyl, 1,3,4-thiadiazolyl, or 1,3,4-thiadiazolyl which is substituted by F, Cl, CF_3, methyl, methoxymethyl, isopropyl, isopbutyl, t-butyl, carboxy, alkoxycarbonyl, cyclopropyl, cyclopropylmethyl, phenyl, pyridyl, piperidinyl, 3,4-methylenedioxyphenyl, thienyl, pyrazinyl, tetrahydrofuranyl, morpholinyl, pyrrolidinyl, or thienylmethyl.
Ix, IIx	R^1 is cyclopentyl, tetrahydrofuranyl, or cyclopropylmethyl, as well as CHF_2; R^2 is CH_3 or CHF_2; and R^3 is $\text{CH}_2\text{CONHR}^5$ or $-\text{CH}_2\text{CONR}^6\text{R}^5$.
Iy, IIy	R^1 is cyclopentyl, tetrahydrofuran, or cyclopropylmethyl, as well as CHF_2; R^2 is CH_3 or CHF_2;

- 5 R^3 is CH_2CONHR^5 or $-CH_2CONR^6R^5$;
 R^4 is substituted or unsubstituted phenyl;
 R^5 is H, alkyl, substituted or unsubstituted phenyl or pyridyl, or a substituted or unsubstituted heterocyclic group; and
 R^6 is alkyl, cycloalkyl, or cycloalkylalkyl.

According to a further preferred compound aspect of the invention, the compound of formula I is selected from:

- 4-[4-Methoxy-3-(4-methoxyphenoxy)phenyl]-2-pyrrolidone,
 10 4-[4-Methoxy-3-(3-thienyloxy)phenyl]-2-pyrrolidone,
 4-[3-(4-Fluorophenoxy)-4-methoxyphenyl]-2-pyrrolidone,
 4-(3-(3-Cyclohexyl-1-propyloxy)-4-methoxyphenyl)-2-pyrrolidone,
 4-(4-Methoxy-3-(2-phenylethoxy)phenyl)-2-pyrrolidone,
 4-(4-Methoxy-3-(3-phenyl-1-propoxy)phenyl)-2-pyrrolidone,
 15 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-chloro-4-fluorobenzyl)-2-pyrrolidone,
 4-(3-cyclopentyloxy-4-methoxyphenyl)-1-methoxycarbonylmethyl-2-pyrrolidone,
 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-cyanomethyl-2-pyrrolidone,
 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-cyclopentyl-2-pyrrolidone,
 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-methoxybenzoyl)-2-pyrrolidone,
 20 4(S)-(3-Cyclopentyloxy-4-difluoromethoxyphenyl)-1-(N-(2-(6-methylpyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
 1-(N-(2,3-Difluorophenyl)-aminocarbonylmethyl)-4(S)-(4-methoxy-3-(3(R)-tetrahydrofuranyloxy)phenyl)-2-pyrrolidone,
 4(S)-(4-Methoxy-3-(3(R)-tetrahydrofuranyloxyphenyl)-1-(N-(2-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
 25 4(S)-(4-Methoxy-3-(3(R)-tetrahydrofuranyloxyphenyl)-1-(N-(2-(6-methylpyridinyl))-aminocarbonylmethyl)-2-pyrrolidone, or

- physiologically acceptable salts thereof, wherein in each case the compound can be in the form of a mixture of enantiomers such as the racemate or a mixture of
 30 diastereomers, or can be in the form of a single enantiomer or a single diastereomer.

According to a further preferred compound aspect of the invention, the compound of formula I is selected from:

- (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-methylbenzyl)-2-pyrrolidone,
5 (4*S*)-1-(2-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
(4*S*)-1-(4-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
(4*S*)-1-(3-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
(4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-methoxybenzyl)-2-pyrrolidone,
(4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-fluorobenzyl)-2-pyrrolidone,
10 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-fluorobenzyl)-2-pyrrolidone,
(4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-fluorobenzyl)-2-pyrrolidone,
(4*S*)-1-(4-Cyanobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
(4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-trifluoromethylbenzyl)-2-pyrrolidone,
(4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-trifluoromethylbenzyl)-2-pyrrolidone,
15 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-trifluoromethylbenzyl)-2-pyrrolidone,
(4*S*)-1-(3,5-bis(trifluoromethyl)benzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
(4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,4-difluorobenzyl)-2-pyrrolidone,
(4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,5-difluorobenzyl)-2-pyrrolidone,
20 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,4-difluorobenzyl)-2-pyrrolidone,
(4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,6-difluorobenzyl)-2-pyrrolidone,
(4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,3-difluorobenzyl)-2-pyrrolidone,
(4*S*)-1-(2-Chloro-4-fluorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
(4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,4-dichlorobenzyl)-2-pyrrolidone,
25 (4*S*)-1-(4-*tert*-Butylbenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
(4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-ethyl-2-pyrrolidone,
(4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-propyl-2-pyrrolidone,
(4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-butyl-2-pyrrolidone,
(4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-methoxyethyl)-2-pyrrolidone,
30 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-phenylbenzyl)-2-pyrrolidone,
(4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-methylbenzyl)-2-pyrrolidone,

(4*R*)-1-(2-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*R*)-1-(4-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*R*)-1-(3-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-methoxybenzyl)-2-pyrrolidone,
 5 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-fluorobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-fluorobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-fluorobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-trifluoromethylbenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-trifluoromethylbenzyl)-2-pyrrolidone,
 10 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-trifluoromethylbenzyl)-2-pyrrolidone,
 (4*R*)-1-(3,5-Bis(trifluoromethyl)benzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-
 pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,4-difluorobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,5-difluorobenzyl)-2-pyrrolidone,
 15 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,4-difluorobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,6-difluorobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,3-difluorobenzyl)-2-pyrrolidone,
 (4*R*)-1-(2-Chloro-4-fluorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,4-dichlorobenzyl)-2-pyrrolidone,
 20 (4*R*)-1-(4-*tert*-Butylbenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-ethyl-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-propyl-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-butyl-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-methoxyethyl)-2-pyrrolidone,
 25 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-phenylbenzyl)-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2,6-dimethylphenyl)-
 aminocarbonylmethyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2,6-dimethylphenyl)-
 aminocarbonylmethyl)-2-pyrrolidone,
 30 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2-methylphenyl)-aminocarbonylmethyl)-
 2-pyrrolidone,

- (4R)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- (4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 5 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2,3-difluorophenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3-chlorophenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(4-pyridyl)-aminocarbonylmethyl)-2-
- 10 pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(4-methoxyphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(4-chloro-2-fluorophenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 15 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(4-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(4-nitrophenyl)-aminocarbonylmethyl)-2-
- 20 pyrrolidone,
- 4-(3-Cyclopentyloxy-4-difluoromethoxyphenyl)-1-(N-(2,3-difluorophenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4(S)-(3-Cyclopentyloxy-4-difluoromethoxyphenyl)-1-(N-(2-(6-methylpyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
- 25 4(S)-(4-Methoxy-3-(3(R)-tetrahydrofuran-2-yl)phenyl)-1-(N-(2-(6-methylpyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
- 1-(N-(2,3-Difluorophenyl)-aminocarbonylmethyl)-4(S)-(4-methoxy-3-(3(R)-tetrahydrofuran-2-yl)phenyl)-2-pyrrolidone,
- 1-(N-(2-(6-Aminopyridyl))-aminocarbonylmethyl)-4(S)-(3-cyclopentyloxy-4-
- 30 methoxyphenyl)-2-pyrrolidone,

- 4(S)-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-(6-ethylpyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
 4(S)-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-(4,6-dimethylpyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
 5 1-(N-(2-(6-Bromopyridyl))-aminocarbonylmethyl)-4(S)-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 1-(N-(2-(6-Bromopyridyl))-aminocarbonylmethyl)-4(S)-(4-methoxy-3-(3(R)-tetrahydrofuranyloxy)phenyl)-2-pyrrolidone,
 4(S)-(4-Methoxy-3-(3(R)-tetrahydrofuranyloxy)phenyl)-1-(N-(2-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
 10 4(S)-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3-(2-methoxypyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
 1-(N-(6-(3-Bromo-2-methylpyridyl))-aminocarbonylmethyl)-4(S)-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 15 4(S)-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3-(4-methoxycarbonyl)-pyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
 4(S)-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-(6-methylpyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
 1-(N-(3-(2-Cyanopyridyl))-aminocarbonylmethyl)-4(S)-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone, or
 20 4(S)-(4-Methoxy-3-(3(R)-tetrahydrofuranyloxy)phenyl)-1-(N-(2-(6-methylpyridinyl))-aminocarbonylmethyl)-2-pyrrolidone, or
 physiologically acceptable salts thereof, wherein in each case the compound can be in the form of a mixture of enantiomers such as the racemate, or a mixture of
 25 diastereomers, or can be in the form of a single enantiomer or a single diastereomer.

According to a further preferred compound aspect of the invention, the compound of formula I is selected from:
 (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone-1-acetic acid,
 30 (4S)-1-(N-Methoxycarbonylmethyl)-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,

- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-methylphenyl)-(N-methyl)aminocarbonylmethyl]-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-(6-methylpyridyl)-(N-methyl)aminocarbonylmethyl)-2-pyrrolidone,
- 5 (4S)-1-[N-(2,3-Difluorophenyl)-(N-methyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(phenylaminocarbonylmethyl)-2-pyrrolidone,
- (4S)-1-[N-(4,5-Dimethylthiazol)-2-yl]aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 10 (4S)-1-[N-(3-Chlorophenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(3-methoxycarbonylphenyl)aminocarbonylmethyl]-2-pyrrolidone,
- 15 (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(3-phenpropyl)]-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-phenoxyethyl)]-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-phenthioethyl)]-2-pyrrolidone,
- 20 (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-(N-phensulfonylaminocarbonylmethyl)-2-pyrrolidone,
- (4S)-1-[N-(2,3-Difluorophenyl)-(N-ethyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 25 (4S)-1-[N-(2,3-Difluorophenyl)-(N-isopropyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-1-[N-(2,3-Difluorophenyl)-(N-cyclopropylmethyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-1-[N-(4-Carboxyphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 30

- (4S)-1-[N-(3-Fluorophenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-thiazolyl)aminocarbonylmethyl]-2-pyrrolidone,
- 5 (4S)-1-[N-(4-Methoxyphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-1-[N-(2,6-Dimethylphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-1-[N-(4-Isopropylphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 10 (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(3,4-methylenedioxyphenyl)aminocarbonylmethyl]-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-(4-trifluoromethyl)pyridyl)aminocarbonylmethyl]-2-pyrrolidone,
- 15 (4S)-1-[N-(3-Carboxyphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-phenylsulfonyl)ethyl]-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-methylphenyl)sulfonylaminocarbonylmethyl]-2-pyrrolidone,
- 20 (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-(4-methoxyphenyl)oxyethyl)]-2-pyrrolidone,
- (4S)-1-[N-(2-(5-Chlorobenzoxazolyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 25 (4S)-1-[N-(2-(Benzthiazolyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-1-[N-(2-(6-Fluorobenzthiazolyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-1-[N-(2-(Benzimidazolyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 30 tetrahydrofuranyloxyphenyl)-2-pyrrolidone,

physiologically acceptable salts thereof, wherein in each case the compound can be in the form of a mixture of enantiomers such as the racemate, or a mixture of diastereomers, or can be in the form of a single enantiomer or a single diastereomer.

- 5 According to a further preferred compound aspect of the invention, the compound of formula I is selected from:
- (4S)-1-[2-(3-Chlorophenoxy)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [431.91 MW; 432.0 M+H],
- (4S)-1-[2-(4-Isopropylphenoxy)ethyl]-4-[4-methoxy-3-(3R)-
- 10 tetrahydrofuranyloxyphenyl]-2-pyrrolidone[439.55 MW; 440.1 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methylbenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone[481.57 MW; 481.9 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-methylthiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone[431.51 MW; 432.0 M+H],
- 15 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(6-methylbenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone[481.57 MW; 481.9 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methoxybenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone[497.67 MW; 498.0 M+H],
- (4S)-1-[N-(6-Ethoxycarbonylbenzothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-
- 20 (3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [539.61 MW; 540.0 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(6-trifluoromethoxybenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone[551.54 MW; 551.9 M+H],
- (4S)-1-[N-(4-tert-Butylthiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-
- 25 tetrahydrofuranyloxyphenyl]-2-pyrrolidone [473.591 MW; 474.0 M+H],
- (4S)-1-[2-(4-Isopropylphenylthio)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [455.615 MW; 456.0 M+H],
- (4S)-1-[2-(3-Chlorophenylthio)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[447.98 MW; 448.1 M+H],
- 30 (4S)-1-[2-(2,3-Difluorophenoxy)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[433.45 MW; 434.0 M+H],

- (4S)-1-[N-(2,3-Difluorophenyl)-N-(2-methylpropyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[502.56 MW; 503.0 M+H],
- (4S)-1-[N-(4-Isopropoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[468.55 MW; 469.1 M+H],
- 5 (4S)-1-[N-(4-Fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[428.46 MW; 429.1 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[2-(1,2,3,4-tetrahydroisoquinoliny)carbonylmethyl]-2-pyrrolidone[450.53 MW; 451.2 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[1-(1,2,3,4-tetrahydroquinoliny)carbonylmethyl]-2-pyrrolidone[450.53 MW; 451.2 M+H],
- 10 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-trifluoromethoxyphenyl)aminocarbonylmethyl]-2-pyrrolidone[494.46 MW; 495.1 M+H],
- (4S)-1-[N-(6-Fluorobenzothiazol-2-yl)-N-(methyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[499.56 MW; 499.9 M+H],
- 15 (4S)-1-[2-(Benzothiazol-2-yl)oxyethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[454.54 MW; 455.0 M+H],
- (4S)-1-[2-(6-Fluorobenzothiazol-2-yl)thioethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[488.60 MW; 489.0 M+H],
- (4S)-1-[N-(6-Fluorobenzothiazol-2-yl)aminoethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[471.55 MW; 472.0 M+H],
- 20 (4S)-1-[N-(Benzothiazol-2-yl)aminoethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [453.56 MW; 454.0 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-N-[2-(2-oxopyrrolidin-1-yl)ethyl]-4-phenoxybenzamide[516.59 MW; 517.0 M+H],
- 25 (4S)-1-[N-(3-Fluorophenyl)-N-(methyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[442.48 MW; 443.0 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methoxyphenyl)-N-(methyl)aminocarbonylmethyl]-2-pyrrolidone[454.52 MW; 455.02 M+H],
- (4S)-1-[N-(4-Isopropylphenyl)-N-(methyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[466.57 MW; 467.1 M+H],
- 30

- (4S)-1-[N-(3,4-Methylenedioxyphenyl)-N-(methyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [468.5 MW; 469.0 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methylthiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [431.51 MW; 432.0 M+H],
- 5 (4S)-1-[N-(4-tert-Butylphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [466.57 MW; 467.1 M+H],
- (4S)-1-[N-(6-Chlorobenzothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [501.99 MW; 501.9, 503.9 M+H]
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-methyl-N-(thiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [431.51 MW; 432.0 M+H]
- 10 (4S)-1-[N-(Benzothiazol-2-yl)-N-(cyclopropylmethyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [521.63 MW; 522.0 M+H]
- (4S)-1-[N-(Indol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [449.5 MW; 450.0 M+H]
- 15 (4S)-1-[N-(Indan-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [450.53 MW; 451.0 M+H]
- (4S)-1-[N-(5-Chlorothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [451.91 MW; 451.9, 453.9 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-phenylthiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [493.48 MW; 494.0 M+H],
- 20 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(6-methoxybenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [497.57 MW; 498.0 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-N-[2-(2-oxopyrrolidin-1-yl)ethyl]benzamide [424.49 MW; 425.0 M+H],
- 25 (4S)-2,3-Difluoro-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-N-[2-(2-oxopyrrolidin-1-yl)ethyl]-benzamide [460.47 MW; 461.0 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-N-[2-(2-oxopyrrolidin-1-yl)ethyl]-4-methoxybenzamide [454.52 MW; 455.0 M+H],
- (4S)-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[2-(4-trifluoromethylphenoxy)ethyl]-2-pyrrolidone [465.46 MW; 466.0 M+H],
- 30

- (4S)-1-[N-(5-Cyclopropyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [458.54 MW; 459.20 M+H],
- (4S)-1-[N-(Benzothiazol-6-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [467.54 MW; 468.0 M+H],
- 5 (4S)-1-[N-(4-Ethoxycarbonylthiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [489.55 MW; 490.0 M+H],
- (4S)-1-[N-(5-tert-Butyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [474.58 MW; 475.0 M+H],
- (4S)-1-[N-Cyclopropylmethyl-N-(6-fluorobenzothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [539.63 MW; 540.0 M+H],
- 10 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-(2-oxo-2-phenylethyl)-2-pyrrolidone [395.45 MW; 396.1 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-(2-oxo-2-(4-methoxyphenyl)ethyl)-2-pyrrolidone [425.48 MW; 426.1 M+H],
- 15 (4S)-1-[N-(2,4-Dimethoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [470.52 MW; 471.0 M+H],
- (4S)-1-[N-(3,5-Dimethoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [470.52 MW; 471.0 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(2,2,3,3-tetrafluorobenzo-1,4-dioxan-6-yl)aminocarbonylmethyl]-2-pyrrolidone [540.46 MW; 541.0 M+H],
- 20 (4S)-1-[N-(3,4-(Difluoromethylene)dioxyphenyl)-N-methylaminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [490.46 MW; 491.0 M+H],
- (4S)-1-[N-(3-Fluoro-4-methoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [458.48 MW; 459.1 M+H],
- 25 (4S)-1-[N-(1,4-Benzodioxan-6-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [468.50 MW; 469.1 M+H],
- (4S)-1-[N-(2-Fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [428.46 MW; 429.1 M+H],
- (4S)-1-[N-(3,4-Dimethoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [470.52 MW; 471.1 M+H],
- 30

- (4S)-1-[N-(3,4-Difluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [446.45 MW; 447.1 M+H],
- (4S)-1-[N-(4-Methanesulfonamidophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [503.57 MW; 504.0 M+H],
- 5 (4S)-1-[N-(4-(4-Fluorophenyl)thiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [511.57 MW; 512.1 M+H],
- (4S)-1-[N-(3-Fluoro-4-methylphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [442.48 MW; 443.1 M+H],
- (4S)-1-[N-(4,6-Difluorobenzothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [503.52 MW; 504.1 M+H],
- 10 (4S)-1-[N-(4-Carboxythiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [461.49 MW; 462.1 M+H],
- (4R)-1-[N-(3-Fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3S)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [428.46 MW; 429.1 M+H],
- 15 (4S)-1-[2-(2-Fluorophenylthio)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [431.52 MW; 432.1 M+H],
- (4S)-1-[2-(3-Fluorophenylthio)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [431.52 MW; 432.1 M+H],
- (4S)-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[2-(4-methoxyphenylthio)ethyl]-2-pyrrolidone [443.56 MW; 444.1 M+H],
- 20 (4S)-1-[N-(4-Carboxy-3-fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [472.47 MW; 473.1 M+H],
- (4S)-1-[N-(4-Ethanesulfonamidophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [517.60 MW; 518.1 M+H],
- 25 (4S)-1-[N-(4-Benzenesulfonamidophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [565.64 MW; 566.1 M+H],
- (4S)-1-[N-(4-(4-Fluorobenzene)sulfonamidophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [583.63 MW; 584.1 M+H],
- (4S)-1-[N-(2,3-Difluorobenzyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [460.47 MW; 461.1 M+H],
- 30

- (4S)-1-[N-(5-Cyclopropylmethyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [472.56 MW; 473.2 M+H],
- (4S)-1-[N-(6-Ethylpyridin-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [439.51 MW; 441.4 M+H],
- 5 (4S)-1-[N-(3-Fluorobenzyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [442.48 MW; 443.1 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(2-methylbenzyl)aminocarbonylmethyl]-2-pyrrolidone [438.52 MW; 439.1 M+H],
- (4S)-1-[N-(4-Methanesulfonylbenzyl)aminocarbonylmethyl]-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [502.58 MW; 503.1 M+H],
- 10 (4S)-1-[N-(4-Aminosulfonylbenzyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [503.57 MW; 504.1 M+H],
- (4S)-1-[N-(Benzothiazol-2-yl)methylaminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [481.57 MW; 482.1 M+H],
- 15 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(3-methylpyridin-2-yl)methylaminocarbonylmethyl]-2-pyrrolidone [439.51 MW; 440.2 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-trifluoromethyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [486.47 MW; 487.1 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-pyridyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [495.56 MW; 496.0 M+H],
- 20 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-(3-pyridyl)thiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [494.57 MW; 495.1 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-(2-pyridyl)thiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [494.57 MW; 495.1 M+H],
- 25 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-(4-pyridyl)thiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [494.57 MW; 495.1 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-pyridyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 30 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-ethoxycarbonyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone

- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-methoxycarbonyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 5 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(3,4-methylenedioxyphenyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-thienyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 10 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-thienylmethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-propyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 15 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-pyrazinyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-methoxymethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 20 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-tetrahydrofuranyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-tetrahydrofuranyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 25 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-aminosulfonyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-methoxyphenyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 30 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-methoxyphenyloxymethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-morpholinylcarbonylmethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 35 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-piperidinylcarbonylmethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(1-pyrrolidinylcarbonylmethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 40 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(1-pyrrolidinylcarbonylmethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-piperidinyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 45

(4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-(N-tertbutyloxycarbonyl)piperidinyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone

- 5 (4S)-1-[N-(2,3-Difluorophenylaminocarbonylmethyl)-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone

(4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(4-ethoxycarbonylphenyl)aminocarbonylmethyl]-2-pyrrolidone,

10

(4S)-1-[N-(4-tert-butyloxycarbonyl-3-fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone, and

- physiologically acceptable salts thereof, wherein in each case the compound can
15 be in the form of a mixture of enantiomers such as the racemate, or a mixture of diastereomers, or can be in the form of a single enantiomer or a single diastereomer.

Accordingly to a preferred method aspect of the invention, the compound of formula I is selected from:

- 20 4-[4-Methoxy-3-(4-methoxyphenoxy)phenyl]-2-pyrrolidone,
4-[4-Methoxy-3-(3-thienyloxy)phenyl]-2-pyrrolidone,
4-[3-(4-Fluorophenoxy)-4-methoxyphenyl]-2-pyrrolidone,
4-(3-(3-Cyclohexyl-1-propyloxy)-4-methoxyphenyl)-2-pyrrolidone,
4-(4-Methoxy-3-(2-phenylethoxy)phenyl)-2-pyrrolidone,
25 4-(4-Methoxy-3-(3-phenyl-1-propoxy)phenyl)-2-pyrrolidone,
4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-chloro-4-fluorobenzyl)-2-pyrrolidone,
4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-methoxycarbonylmethyl-2-pyrrolidone,
4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-cyanomethyl-2-pyrrolidone,
4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-cyclopentyl-2-pyrrolidone,
30 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-methoxybenzoyl)-2-pyrrolidone,
(4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-methylbenzyl)-2-pyrrolidone,
(4S)-1-(2-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
(4S)-1-(4-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
(4S)-1-(3-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
35 (4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-methoxybenzyl)-2-pyrrolidone,

- (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-nitrobenzyl)-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-nitrobenzyl)-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-fluorobenzyl)-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-fluorobenzyl)-2-pyrrolidone,
 5 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-fluorobenzyl)-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-benzyl-2-pyrrolidone,
 (4*S*)-1-(4-Cyanobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-trifluoromethylbenzyl)-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-trifluoromethylbenzyl)-2-pyrrolidone,
 10 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-trifluoromethylbenzyl)-2-pyrrolidone,
 (4*S*)-1-(3,5-bistrifluoromethylbenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,4-difluorobenzyl)-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,5-difluorobenzyl)-2-pyrrolidone,
 15 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,4-difluorobenzyl)-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,6-difluorobenzyl)-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,3-difluorobenzyl)-2-pyrrolidone,
 (4*S*)-1-(2-Chloro-4-fluorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,4-dichlorobenzyl)-2-pyrrolidone,
 20 (4*S*)-1-(4-*tert*-Butylbenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-ethyl-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-propyl-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-butyl-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-methoxyethyl)-2-pyrrolidone,
 25 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-phenylbenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-methylbenzyl)-2-pyrrolidone,
 (4*R*)-1-(2-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*R*)-1-(4-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*R*)-1-(3-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 30 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-methoxybenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-nitrobenzyl)-2-pyrrolidone,

- (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-nitrobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-fluorobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-fluorobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-fluorobenzyl)-2-pyrrolidone,
 5 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-benzyl-2-pyrrolidone,
 (4*R*)-1-(4-Cyanobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-trifluoromethylbenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-trifluoromethylbenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-trifluoromethylbenzyl)-2-pyrrolidone,
 10 (4*R*)-1-(3,5-bis(trifluoromethyl)benzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-
 pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,4-difluorobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,5-difluorobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,4-difluorobenzyl)-2-pyrrolidone,
 15 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,6-difluorobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,3-difluorobenzyl)-2-pyrrolidone,
 (4*R*)-1-(2-Chloro-4-fluorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,4-dichlorobenzyl)-2-pyrrolidone,
 (4*R*)-1-(4-*tert*-Butylbenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 20 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-ethyl-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-propyl-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-butyl-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-methoxyethyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-phenylbenzyl)-2-pyrrolidone,
 25 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2,6-dimethylphenyl)-
 aminocarbonylmethyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2,6-dimethylphenyl)-
 aminocarbonylmethyl)-2-pyrrolidone,
 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2-methylphenyl)-aminocarbonylmethyl)-
 30 2-pyrrolidone,

- (4R)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- (4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 5 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2,3-difluorophenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3-chlorophenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(4-pyridyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 10 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(4-methoxyphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(4-chloro-2-fluorophenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 15 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(4-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(4-nitrophenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 20 4-(*R*)-[4-Methoxy-3-(3-(*R*)-tetrahydrofuranyloxy)phenyl]-2-pyrrolidone,
- 4-(*S*)-[4-Methoxy-3-(3-(*R*)-tetrahydrofuranyloxy)phenyl]-2-pyrrolidone,
- 4-(*R*)-[4-Methoxy-3-(3-(*S*)-tetrahydrofuranyloxy)phenyl]-2-pyrrolidone,
- 4-(*S*)-[4-Methoxy-3-(3-(*S*)-tetrahydrofuranyloxy)phenyl]-2-pyrrolidone,
- 25 4-[4-Difluoromethoxy-3-(3-(*R*)-tetrahydrofuranyloxy)phenyl]-2-pyrrolidone,
- 4-[4-Difluoromethoxy-3-(2-cyclohexylethoxy)phenyl]-2-pyrrolidone,
- 4-[4-Difluoromethoxy-3-(2-phenylethoxy)phenyl]-2-pyrrolidone,
- 4-[4-Difluoromethoxy-3-(cyclopropylmethoxy)phenyl]-2-pyrrolidone,
- 4-[4-Difluoromethoxy-3-cyclopentoxypheyl]-2-pyrrolidone,
- 30 4-[4-Difluoromethoxy-3-cyclobutylmethoxyphenyl]-2-pyrrolidone,

- (4*R*)-1-(2,3-Difluorobenzyl)-4-(3-(3-(*S*)-tetrahydrofuryl)oxy-4-methoxyphenyl)-2-pyrrolidone,
- (4*S*)-1-(2,3-Difluorobenzyl)-4-(3-(3-(*S*)-tetrahydrofuryl)oxy-4-methoxyphenyl)-2-pyrrolidone,
- 5 (4*R*)-1-(2,3-Difluorobenzyl)-4-(3-(3-(*R*)-tetrahydrofuryl)oxy-4-methoxyphenyl)-2-pyrrolidone,
- (4*S*)-1-(2,3-Difluorobenzyl)-4-(3-(3-(*R*)-tetrahydrofuryl)oxy-4-methoxyphenyl)-2-pyrrolidone,
- (4*S*)-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2,3-difluorophenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 10 (4*R*)-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2,3-difluorophenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(3-(2-chloropyridinyl))-aminocarbonylmethyl)-2-pyrrolidone,
- 15 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2-(2-pyridylethyl))-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2-(2-(*N*-methylpyrrolidinyl)ethyl))-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(4-pyridylmethyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 20 2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(1-imidazolylpropyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(4-(4-methylpiperazinyl))-hydrazinocarbonylmethyl)-2-pyrrolidone,
- 25 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2-(*N*-methylpyrrolidinyl)methyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(4-pyrimidinyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(3-(5-methylisoxazolyl))-aminocarbonylmethyl)-2-pyrrolidone,
- 30 aminocarbonylmethyl)-2-pyrrolidone,

- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2,6-dimethylpiperazinyl)-hydrazinocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-(4,5-dimethylthiazolyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 5 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-(2-methylpiperidinyloxy)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3,4-dimethoxyphenethyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-pyridinyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 10 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3-pyridinyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3-(2,4-dimethoxypyridinyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 15 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3-(4-methoxypyridinyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(4-aniline)-aminocarbonylmethyl)-2-pyrrolidone,
- 4-(3-Cyclopentyloxy-4-difluoromethoxyphenyl)-1-(N-(2,3-difluorophenyl)-aminocarbonylmethyl)-2-pyrrolidone,
- 20 4(S)-(3-Cyclopentyloxy-4-difluoromethoxyphenyl)-1-(N-(2-(6-methylpyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
- 4(S)-(4-Methoxy-3-(3(R)-tetrahydrofuryloxy)phenyl)-1-(N-(2-(6-methylpyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
- 25 1-(N-(2,3-Difluorophenyl)-aminocarbonylmethyl)-4(S)-(4-methoxy-3-(3(R)-tetrahydrofuryloxy)phenyl)-2-pyrrolidone,
- 1-(N-(2-(6-Aminopyridyl))-aminocarbonylmethyl)-4(S)-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
- 4(S)-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-(6-ethylpyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
- 30

- 4(S)-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-(4,6-dimethylpyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
 1-(N-(2-(6-Bromopyridyl))-aminocarbonylmethyl)-4(S)-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 5 1-(N-(2-(6-Bromopyridyl))-aminocarbonylmethyl)-4(S)-(4-methoxy-3-(3(R)-tetrahydrofuranyloxy)phenyl)-2-pyrrolidone,
 4(S)-(4-Methoxy-3-(3(R)-tetrahydrofuranyloxy)phenyl)-1-(N-(2-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
 4(S)-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3-(2-methoxypyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
 10 1-(N-(6-(3-Bromo-2-methylpyridyl))-aminocarbonylmethyl)-4(S)-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 4(S)-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3-(4-methoxycarbonyl)-pyridyl)-aminocarbonylmethyl)-2-pyrrolidone,
 15 4(S)-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-(6-methylpyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
 1-(N-(3-(2-Cyanopyridyl))-aminocarbonylmethyl)-4(S)-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone, or
 4(S)-(4-Methoxy-3-(3(R)-tetrahydrofuranyloxy)phenyl)-1-(N-(2-(6-methylpyridinyl))-aminocarbonylmethyl)-2-pyrrolidone, or
 20 physiologically acceptable salts thereof, wherein in each case the compound can be in the form of a mixture of enantiomers such as the racemate, or a mixture of diastereomers, or can be in the form of a single enantiomer or a single diastereomer.
- 25 According to a further preferred method aspect of the invention, the compound of formula I is selected from:
 (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone-1-acetic acid,
 (4S)-1-(N-Methoxycarbonylmethyl)-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
 30 (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-methylphenyl)-(N-methyl)aminocarbonylmethyl]-2-pyrrolidone,

- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-(6-methylpyridyl)-(N-methyl)aminocarbonylmethyl)]-2-pyrrolidone,
- (4S)-1-[N-(2,3-Difluorophenyl-(N-methyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 5 (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(phenylaminocarbonylmethyl)]-2-pyrrolidone,
- (4S)-1-[N-(4,5-Dimethylthiazol)-2-yl]aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-1-[N-(3-Chlorophenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 10 (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(3-methoxycarbonylphenyl)aminocarbonylmethyl]-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(3-phenpropyl)]-2-pyrrolidone,
- 15 (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-phenoxyethyl)]-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-phenthioethyl)]-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-(N-phensulfonylaminocarbonylmethyl)-2-pyrrolidone,
- 20 (4S)-1-[N-(2,3-Difluorophenyl-(N-ethyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-1-[N-(2,3-Difluorophenyl-(N-isopropyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 25 (4S)-1-[N-(2,3-Difluorophenyl-(N-cyclopropylmethyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-1-[N-(4-Carboxyphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-1-[N-(3-Fluorophenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 30

- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-thiazolyl)aminocarbonylmethyl]-2-pyrrolidone,
- (4S)-1-[N-(4-Methoxyphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 5 (4S)-1-[N-(2,6-Dimethylphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-1-[N-(4-Isopropylphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(3,4-methylenedioxyphenyl)aminocarbonylmethyl]-2-pyrrolidone,
- 10 (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-(4-trifluoromethyl)pyridyl)aminocarbonylmethyl]-2-pyrrolidone,
- (4S)-1-[N-(3-Carboxyphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 15 (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-phenylsulfonyl)ethyl]-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-methylphenyl)sulfonylaminocarbonylmethyl]-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-(4-methoxyphenyl)oxyethyl)]-2-pyrrolidone,
- 20 (4S)-1-[N-(2-(5-Chlorobenzoxazolyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-1-[N-(2-(Benzthiazolyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 25 (4S)-1-[N-(2-(6-Fluorobenzthiazolyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-1-[N-(2-(Benzimidazolyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,

physiologically acceptable salts thereof, wherein in each case the compound can be in the form of a mixture of enantiomers such as the racemate, or a mixture of diastereomers, or can be in the form of a single enantiomer or a single diastereomer.

Accordingly to a further method aspect of the invention, the compound of formula I is selected from:

- (4S)-1-[2-(3-Chlorophenoxy)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [431.91 MW; 432.0 M+H],
- 5 (4S)-1-[2-(4-Isopropylphenoxy)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[439.55 MW; 440.1 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methylbenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone[481.57 MW; 481.9 M+H],
- 10 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-methylthiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone[431.51 MW; 432.0 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(6-methylbenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone[481.57 MW; 481.9 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methoxybenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone[497.67 MW; 498.0 M+H],
- 15 (4S)-1-[N-(6-Ethoxycarbonylbenzothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [539.61 MW; 540.0 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(6-trifluoromethoxybenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone[551.54 MW;
- 20 551.9 M+H],
- (4S)-1-[N-(4-tert-Butylthiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [473.591 MW; 474.0 M+H],
- (4S)-1-[2-(4-Isopropylphenylthio)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [455.615 MW; 456.0 M+H],
- 25 (4S)-1-[2-(3-Chlorophenylthio)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[447.98 MW; 448.1 M+H],
- (4S)-1-[2-(2,3-Difluorophenoxy)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[433.45 MW; 434.0 M+H],
- (4S)-1-[N-(2,3-Difluorophenyl)-N-(2-methylpropyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[502.56 MW; 503.0 M+H],
- 30

- (4S)-1-[N-(4-Isopropoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[468.55 MW; 469.1 M+H],
- (4S)-1-[N-(4-Fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[428.46 MW; 429.1 M+H],
- 5 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[2-(1,2,3,4-tetrahydroisoquinolinyl)carbonylmethyl]-2-pyrrolidone[450.53 MW; 451.2 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[1-(1,2,3,4-tetrahydroquinolinyl)carbonylmethyl]-2-pyrrolidone[450.53 MW; 451.2 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-trifluoromethoxyphenyl)aminocarbonylmethyl]-2-pyrrolidone[494.46 MW; 495.1 M+H],
- 10 (4S)-1-[N-(6-Fluorobenzothiazol-2-yl)-N-(methyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[499.56 MW; 499.9 M+H],
- (4S)-1-[2-(Benzothiazol-2-yl)oxyethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[454.54 MW; 455.0 M+H],
- 15 (4S)-1-[2-(6-Fluorobenzothiazol-2-yl)thioethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[488.60 MW; 489.0 M+H],
- (4S)-1-[N-(6-Fluorobenzothiazol-2-yl)aminoethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[471.55 MW; 472.0 M+H],
- (4S)-1-[N-(Benzothiazol-2-yl)aminoethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [453.56 MW; 454.0 M+H],
- 20 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-N-[2-(2-oxopyrrolidin-1-yl)ethyl]-4-phenoxybenzamide[516.59 MW; 517.0 M+H],
- (4S)-1-[N-(3-Fluorophenyl)-N-(methyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[442.48 MW; 443.0 M+H],
- 25 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methoxyphenyl)-N-(methyl)aminocarbonylmethyl]-2-pyrrolidone[454.52 MW; 455.02 M+H],
- (4S)-1-[N-(4-Isopropylphenyl)-N-(methyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[466.57 MW; 467.1 M+H],
- (4S)-1-[N-(3,4-Methylenedioxyphenyl)-N-(methyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[468.5 MW; 469.0 M+H],
- 30

- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methylthiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [431.51 MW; 432.0 M+H],
- (4S)-1-[N-(4-tert-Butylphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [466.57 MW; 467.1 M+H],
- 5 (4S)-1-[N-(6-Chlorobenzothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [501.99 MW; 501.9, 503.9 M+H]
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-methyl-N-(thiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [431.51 MW; 432.0 M+H]
- (4S)-1-[N-(Benzothiazol-2-yl)-N-(cyclopropylmethyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [521.63 MW; 522.0 M+H]
- 10 (4S)-1-[N-(Indol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [449.5 MW; 450.0 M+H]
- (4S)-1-[N-(Indan-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [450.53 MW; 451.0 M+H]
- 15 (4S)-1-[N-(5-Chlorothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [451.91 MW; 451.9, 453.9 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-phenylthiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [493.48 MW; 494.0 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(6-methoxybenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [497.57 MW; 498.0 M+H],
- 20 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-N-[2-(2-oxopyrrolidin-1-yl)ethyl]benzamide [424.49 MW; 425.0 M+H],
- (4S)-2,3-Difluoro-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-N-[2-(2-oxopyrrolidin-1-yl)ethyl]-benzamide [460.47 MW; 461.0 M+H],
- 25 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-N-[2-(2-oxopyrrolidin-1-yl)ethyl]-4-methoxybenzamide [454.52 MW; 455.0 M+H],
- (4S)-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[2-(4-trifluoromethylphenoxy)ethyl]-2-pyrrolidone [465.46 MW; 466.0 M+H],
- (4S)-1-[N-(5-Cyclopropyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [458.54 MW; 459.20 M+H],
- 30

- (4S)-1-[N-(Benzothiazol-6-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [467.54 MW; 468.0 M+H],
- (4S)-1-[N-(4-Ethoxycarbonylthiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [489.55 MW; 490.0 M+H],
- 5 (4S)-1-[N-(5-tert-Butyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [474.58 MW; 475.0 M+H],
- (4S)-1-[N-Cyclopropylmethyl-N-(6-fluorobenzothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [539.63 MW; 540.0 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-(2-oxo-2-phenylethyl)-2-pyrrolidone [395.45 MW; 396.1 M+H],
- 10 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-(2-oxo-2-(4-methoxyphenyl)ethyl)-2-pyrrolidone [425.48 MW; 426.1 M+H],
- (4S)-1-[N-(2,4-Dimethoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [470.52 MW; 471.0 M+H],
- 15 (4S)-1-[N-(3,5-Dimethoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [470.52 MW; 471.0 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(2,2,3,3-tetrafluorobenzo-1,4-dioxan-6-yl)aminocarbonylmethyl]-2-pyrrolidone [540.46 MW; 541.0 M+H],
- (4S)-1-[N-(3,4-(Difluoromethylene)dioxyphenyl)-N-methylaminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [490.46 MW; 491.0 M+H],
- 20 (4S)-1-[N-(3-Fluoro-4-methoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [458.48 MW; 459.1 M+H],
- (4S)-1-[N-(1,4-Benzodioxan-6-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [468.50 MW; 469.1 M+H],
- 25 (4S)-1-[N-(2-Fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [428.46 MW; 429.1 M+H],
- (4S)-1-[N-(3,4-Dimethoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [470.52 MW; 471.1 M+H],
- (4S)-1-[N-(3,4-Difluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [446.45 MW; 447.1 M+H],
- 30

- (4S)-1-[N-(4-Methanesulfonamidophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]phenyl]-2-pyrrolidone [503.57 MW; 504.0 M+H],
- (4S)-1-[N-(4-(4-Fluorophenyl)thiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]phenyl]-2-pyrrolidone [511.57 MW; 512.1 M+H],
- 5 (4S)-1-[N-(3-Fluoro-4-methylphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]phenyl]-2-pyrrolidone [442.48 MW; 443.1 M+H],
- (4S)-1-[N-(4,6-Difluorobenzothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]phenyl]-2-pyrrolidone [503.52 MW; 504.1 M+H],
- (4S)-1-[N-(4-Carboxythiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]phenyl]-2-pyrrolidone [461.49 MW; 462.1 M+H],
- 10 (4R)-1-[N-(3-Fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3S)-tetrahydrofuran-2-yl]phenyl]-2-pyrrolidone [428.46 MW; 429.1 M+H],
- (4S)-1-[2-(2-Fluorophenylthio)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]phenyl]-2-pyrrolidone [431.52 MW; 432.1 M+H],
- 15 (4S)-1-[2-(3-Fluorophenylthio)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]phenyl]-2-pyrrolidone [431.52 MW; 432.1 M+H],
- (4S)-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]phenyl]-1-[2-(4-methoxyphenylthio)ethyl]-2-pyrrolidone [443.56 MW; 444.1 M+H],
- (4S)-1-[N-(4-Carboxy-3-fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]phenyl]-2-pyrrolidone [472.47 MW; 473.1 M+H],
- 20 (4S)-1-[N-(4-Ethanesulfonamidophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]phenyl]-2-pyrrolidone [517.60 MW; 518.1 M+H],
- (4S)-1-[N-(4-Benzenesulfonamidophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]phenyl]-2-pyrrolidone [565.64 MW; 566.1 M+H],
- 25 (4S)-1-[N-(4-(4-Fluorobenzene)sulfonamidophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]phenyl]-2-pyrrolidone [583.63 MW; 584.1 M+H],
- (4S)-1-[N-(2,3-Difluorobenzyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]phenyl]-2-pyrrolidone [460.47 MW; 461.1 M+H],
- (4S)-1-[N-(5-Cyclopropylmethyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]phenyl]-2-pyrrolidone [472.56 MW; 473.2 M+H],
- 30

- (4S)-1-[N-(6-Ethylpyridin-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [439.51 MW; 441.4 M+H],
- (4S)-1-[N-(3-Fluorobenzyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [442.48 MW; 443.1 M+H],
- 5 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(2-methylbenzyl)aminocarbonylmethyl]-2-pyrrolidone [438.52 MW; 439.1 M+H],
- (4S)-1-[N-(4-Methanesulfonylbenzyl)aminocarbonylmethyl]-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [502.58 MW; 503.1 M+H],
- (4S)-1-[N-(4-Aminosulfonylbenzyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [503.57 MW; 504.1 M+H],
- 10 (4S)-1-[N-(Benzothiazol-2-yl)methylaminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [481.57 MW; 482.1 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(3-methylpyridin-2-yl)methylaminocarbonylmethyl]-2-pyrrolidone [439.51 MW; 440.2 M+H],
- 15 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-trifluoromethyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [486.47 MW; 487.1 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-pyridyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [495.56 MW; 496.0 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-(3-pyridyl)thiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [494.57 MW; 495.1 M+H],
- 20 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-(2-pyridyl)thiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [494.57 MW; 495.1 M+H],
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-(4-pyridyl)thiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone [494.57 MW; 495.1 M+H],
- 25 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-pyridyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-ethoxycarbonyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 30 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-methoxycarbonyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone

- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(3,4-methylenedioxyphenyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 5 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-thienyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-thienylmethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 10 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-propyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-pyrazinyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 15 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-methoxymethyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-tetrahydrofuranyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 20 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-aminosulfonyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-methoxyphenyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 25 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-methoxyphenyloxymethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 30 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-morpholinylcarbonylmethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(1-piperidinylcarbonylmethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 35 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(1-pyrrolidinylcarbonylmethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-piperidinyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 40 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-(N-tertbutyloxycarbonyl)piperidinyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 45

(4*S*)-1-[N-(2,3-Difluorophenylaminocarbonylmethyl)-4-(4-methoxy-3-(3*R*)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone

5 (4*S*)-4-(4-Methoxy-3-(3*R*)-tetrahydrofuranyloxyphenyl)-1-[N-(4-ethoxycarbonylphenyl)aminocarbonylmethyl]-2-pyrrolidone,

(4*S*)-1-[N-(4-*tert*-butyloxycarbonyl-3-fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3*R*)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone, and

10 physiologically acceptable salts thereof, wherein in each case the compound can be in the form of a mixture of enantiomers such as the racemate, or a mixture of diastereomers, or can be in the form of a single enantiomer or a single diastereomer.

Preferred aspects include pharmaceutical compositions comprising a compound of
15 this invention and a pharmaceutically acceptable carrier and, optionally, another active agent as discussed below; a method of inhibiting a PDE4 enzyme, especially an isoenzyme, e.g., as determined by a conventional assay or one described herein, either *in vitro* or *in vivo* (in an animal, e.g., in an animal model, or in a mammal or in a human); a method of treating a neurological syndrome, e.g., loss of memory, especially long-term
20 memory, cognitive impairment or decline, memory impairment, etc.; a method of treating a disease state modulated by PDE4 activity, in a mammal, e.g., a human, e.g., those disease states mentioned herein.

The compounds of the present invention may be prepared conventionally. Some
25 of the known processes that can be used are described below. All starting materials are known or can be conventionally prepared from known starting materials.

Preparation of starting material (Scheme I)

malonate to provide the corresponding methyl cinnamyl esters after decarboxylation. Conjugate addition of nitromethane using tetramethyl guanidine as base provided methyl 4-nitro-3-(substituted-phenyl)butyrate. Racemic 4-(3-benzyloxy-4-methoxyphenyl)-2-pyrrolidone was produced in 85% yield by selective reduction of the nitro group to the
5 corresponding amine using NiCl_2 and NaBH_4 (Osby, J.O.; Ganem, B., Rapid and efficient reduction of aliphatic nitro compounds to amines, *Tetrahedron Lett.*, 1985, 26, 6413-6416) and subsequent treatment with K_2CO_3 .


Methyl 4-nitro-3-(substituted-phenyl)butyrate was enantiomerically resolved by
10 synthesizing diastereomeric phenethylamine amides as shown in scheme I. Nitro group reduction by catalytic hydrogenation as described by Demnitz, J. *Molecules*, 1998, 3, 107-119 was not successful. However, reduction of the diastereomerically pure nitro amides using NiCl_2 and NaBH_4 produced 90% yield of the corresponding diastereomerically pure amines. Cyclization by warming to reflux in xylene yielded 51%
15 of (*S*)-(+)-rolipram and 24% of (*R*)-(-)-rolipram, both in better than 99% ee as determined by chiral HPLC (Kusters, E.; Spondlin, C.; Influence of temperature on the enantioseparation of rolipram and structurally related racemates on Chiralcel-OD, *J. Chromatography A*, 1996, 737, 333-337). Enantiomerically pure 3-benzyloxy derivatives 8 ($\text{R}=\text{benzyl}$) were produced in similar yield and purity after cyclization.

20


Alternatively, enantiomerically pure pyrrolidones 7 and 8 can be prepared according to the methods of D.M. Barnes et.al. (Ref. D.M. Barnes, et.al., *J. Am. Chem. Soc.*, 2002, 124, 13097-13105.)

25 3-Aryloxy rolipram derivatives were prepared by cross coupling reaction with aryl boronic acids using a copper catalyst in the presence of an amine base. Suitable copper catalysts include copper diacetate, copper (II) chloride, etc. Generally, halogenated solvents are utilized such as chloroform, dichloromethane, 1,2-dichloroethane, and the like. Commonly used bases include triethylamine,
30 diisopropylamine, and pyrrolidine. Alternatively, 3-phenyloxyrolipram can be prepared by Ullman type coupling starting with iodobenzene and 3-hydroxyrolipram as described

previously (Schmiechen, R.; Horowski, R.; Palenschat, D.; Paschelke, G.; Wachtel, H.; Kehr, W., 4-(polyalkoxyphenyl)-2-pyrrolidones., US Patent 4,193,926, filed Mar. 18, 1980). Compounds substituted at the 3-position with arylalkyl, alkyl, cycloalkyl, heteroalkyl or cycloalkylalkyl groups can be prepared by either Mitsunobu reaction
 5 between phenol **9** ($R^1 = H$) and alcohol or by an alkylation reaction between phenol **9** ($R^1 = H$) and R^1X , where X is a suitable leaving group such as Cl, Br, methanesulfonyl, tosyloxy, etc.



- 10 A series of enantiomerically pure *N*-substituted rolipram derivatives were synthesized by methods common to the art (Christensen, S.B., et.al, 1,4-Cyclohexanecarboxylates: Potent and selective inhibitors of Phosphodiesterase 4 for the treatment of asthma, *J. Med. Chem.*, 1998, *41*, 821-835.). Thus, the target compounds



- can be provided by alkylation reaction with alkyl halides, cycloalkyl halides, arylalkyl
 15 chlorides, alpha bromoacetates, appropriately substituted alpha-bromides, and arylalkyl halides (i.e., benzylbromides) in polar aprotic solvents (i.e., DMF, THF, DMSO) using a non-nucleophilic base such as NaH or LDA, and a phase transfer catalyst such as 15-crown-5 ether.

- 20 Reaction of methyl alpha-bromoacetate with a rolipram derivative as described above produces methyl *N*-acetates **12** ($R = CH_3$), which can be saponified to the acid **12**

5

10



15

20

In the compounds of formulas I, when R^3 is H, the 4(R) enantiomers are preferred, and when R^3 is other than H, the 4(S)-enantiomers are preferred. When R^1 is 3-tetrahydrofuran, the 3(R) enantiomers are preferred over the 3(S) enantiomers.

5

The optical isomers can be obtained by resolution of the racemic mixtures according to conventional processes, for example, by the formation of diastereoisomeric salts using an optically active acid or base or formation of covalent diastereomers. Examples of appropriate acids are tartaric, diacetyltartaric, dibenzoyltartaric, ditoluoyltartaric and camphorsulfonic acid. Mixtures of diastereoisomers can be separated into their individual diastereomers on the basis of their physical and/or chemical differences by methods known to those skilled in the art, for example, by chromatography or fractional crystallization. The optically active bases or acids are then liberated from the separated diastereomeric salts. A different process for separation of optical isomers involves the use of chiral chromatography (e.g., chiral HPLC columns), with or without conventional derivatization, optimally chosen to maximize the separation of the enantiomers. Suitable chiral HPLC columns are manufactured by Diacel, e.g., Chiracel OD and Chiracel OJ among many others, all routinely selectable. Enzymatic separations, with or without derivitization, are also useful. The optically active compounds of Formulas I-II can likewise be obtained by utilizing optically active starting materials in chiral syntheses processes under reaction conditions which do not cause racemization.

In addition, one of ordinary skill in the art will recognize that the compounds can be used in different enriched isotopic forms, e.g., enriched in the content of 2H , 3H , ^{11}C , ^{13}C and/or ^{14}C . In one particular embodiment, the compounds are deuterated. Such deuterated forms can be made the procedure described in U.S. Patent Nos. 5,846,514 and 6,334,997. As described in U.S. Patent Nos. 5,846,514 and 6,334,997, deuteration can improve the efficacy and increase the duration of action of drugs.

30

Deuterium substituted compounds can be synthesized using various methods such as described in: Dean, Dennis C.; Editor. Recent Advances in the Synthesis and Applications of Radiolabeled Compounds for Drug Discovery and Development. [In: Curr., Pharm. Des., 2000; 6(10)] (2000), 110 pp. CAN 133:68895 AN 2000:473538
5 CAPLUS; Kabalka, George W.; Varma, Rajender S. The synthesis of radiolabeled compounds VIA organometallic intermediates. Tetrahedron (1989), 45(21), 6601-21, CODEN: TETRAB ISSN:0040-4020. CAN 112:20527 AN 1990:20527 CAPLUS; and Evans, E. Anthony. Synthesis of radiolabeled compounds, J. Radioanal. Chem. (1981), 64(1-2), 9-32. CODEN: JRACBN ISSN:0022-4081, CAN 95:76229 AN 1981:476229
10 CAPLUS.

The present invention also relates to useful forms of the compounds as disclosed herein, such as pharmaceutically acceptable salts or prodrugs of all the compounds of the present invention for which salts or prodrugs can be prepared. Pharmaceutically
15 acceptable salts include those obtained by reacting the main compound, functioning as a base, with an inorganic or organic acid to form a salt, for example, salts of hydrochloric acid, sulfuric acid, phosphoric acid, methane sulfonic acid, camphor sulfonic acid, oxalic acid, maleic acid, succinic acid and citric acid. Pharmaceutically acceptable salts also include those in which the main compound functions as an acid and is reacted with an
20 appropriate base to form, e.g., sodium, potassium, calcium, magnesium, ammonium, and choline salts. Those skilled in the art will further recognize that acid addition salts of the claimed compounds may be prepared by reaction of the compounds with the appropriate inorganic or organic acid via any of a number of known methods. Alternatively, alkali and alkaline earth metal salts are prepared by reacting the compounds of the invention
25 with the appropriate base via a variety of known methods.

The following are further examples of acid salts that can be obtained by reaction with inorganic or organic acids: acetates, adipates, alginates, citrates, aspartates, benzoates, benzenesulfonates, bisulfates, butyrates, camphorates, digluconates,
30 cyclopentanepropionates, dodecylsulfates, ethanesulfonates, glucoheptanoates, glycerophosphates, hemisulfates, heptanoates, hexanoates, fumarates, hydrobromides,

hydroiodides, 2-hydroxy-ethanesulfonates, lactates, maleates, methanesulfonates, nicotines, 2-naphthalenesulfonates, oxalates, palmoates, pectinates, persulfates, 3-phenylpropionates, picrates, pivalates, propionates, succinates, tartrates, thiocyanates, tosylates, mesylates and undecanoates.

5

Preferably, the salts formed are pharmaceutically acceptable for administration to mammals. However, pharmaceutically unacceptable salts of the compounds are suitable as intermediates, for example, for isolating the compound as a salt and then converting the salt back to the free base compound by treatment with an alkaline reagent. The free
10 base can then, if desired, be converted to a pharmaceutically acceptable acid addition salt.

The compounds of the invention can be administered alone or as an active ingredient of a formulation. Thus, the present invention also includes pharmaceutical compositions of compounds of Formulas I-II, containing, for example, one or more
15 pharmaceutically acceptable carriers.

Numerous standard references are available that describe procedures for preparing various formulations suitable for administering the compounds according to the invention. Examples of potential formulations and preparations are contained, for
20 example, in the Handbook of Pharmaceutical Excipients, American Pharmaceutical Association (current edition); Pharmaceutical Dosage Forms: Tablets (Lieberman, Lachman and Schwartz, editors) current edition, published by Marcel Dekker, Inc., as well as Remington's Pharmaceutical Sciences (Arthur Osol, editor), 1553-1593 (current edition).

25

In view of their high degree of selective PDE4 inhibition, the compounds of the present invention can be administered to anyone requiring PDE4 inhibition. Administration may be accomplished according to patient needs, for example, orally, nasally, parenterally (subcutaneously, intravenously, intramuscularly, intrasternally and
30 by infusion) by inhalation, rectally, vaginally, topically and by ocular administration.

Various solid oral dosage forms can be used for administering compounds of the invention including such solid forms as tablets, gelcaps, capsules, caplets, granules, lozenges and bulk powders. The compounds of the present invention can be administered alone or combined with various pharmaceutically acceptable carriers, diluents (such as sucrose, mannitol, lactose, starches) and excipients known in the art, including but not limited to suspending agents, solubilizers, buffering agents, binders, disintegrants, preservatives, colorants, flavorants, lubricants and the like. Time release capsules, tablets and gels are also advantageous in administering the compounds of the present invention.

Various liquid oral dosage forms can also be used for administering compounds of the inventions, including aqueous and non-aqueous solutions, emulsions, suspensions, syrups, and elixirs. Such dosage forms can also contain suitable inert diluents known in the art such as water and suitable excipients known in the art such as preservatives, wetting agents, sweeteners, flavorants, as well as agents for emulsifying and/or suspending the compounds of the invention. The compounds of the present invention may be injected, for example, intravenously, in the form of an isotonic sterile solution. Other preparations are also possible.

Suppositories for rectal administration of the compounds of the present invention can be prepared by mixing the compound with a suitable excipient such as cocoa butter, salicylates and polyethylene glycols. Formulations for vaginal administration can be in the form of a pessary, tampon, cream, gel, paste, foam, or spray formula containing, in addition to the active ingredient, such suitable carriers as are known in the art.

For topical administration the pharmaceutical composition can be in the form of creams, ointments, liniments, lotions, emulsions, suspensions, gels, solutions, pastes, powders, sprays, and drops suitable for administration to the skin, eye, ear or nose. Topical administration may also involve transdermal administration via means such as transdermal patches.

Aerosol formulations suitable for administering via inhalation also can be made. For example, for treatment of disorders of the respiratory tract, the compounds according to the invention can be administered by inhalation in the form of a powder (e.g., micronized) or in the form of atomized solutions or suspensions. The aerosol formulation
5 can be placed into a pressurized acceptable propellant.

The compounds can be administered as the sole active agent or in combination with other pharmaceutical agents such as other agents used in the treatment of cognitive impairment and/or in the treatment of psychosis, e.g., other PDE4 inhibitors, calcium
10 channel blockers, cholinergic drugs, adenosine receptor modulators, amphotericin NMDA-R modulators, mGluR modulators, and cholinesterase inhibitors (e.g., donepezil, rivastigmine, and galanthamine). In such combinations, each active ingredient can be administered either in accordance with their usual dosage range or a dose below their usual dosage range.

15 The present invention further includes methods of treatment that involve inhibition of PDE4 enzymes. Thus, the present invention includes methods of selective inhibition of PDE4 enzymes in animals, e.g., mammals, especially humans, wherein such inhibition has a therapeutic effect, such as where such inhibition may relieve conditions
20 involving neurological syndromes, such as the loss of memory, especially long-term memory. Such methods comprise administering to an animal in need thereof, especially a mammal, most especially a human, an inhibitory amount of a compound, alone or as part of a formulation, as disclosed herein.

25 The condition of memory impairment is manifested by impairment of the ability to learn new information and/or the inability to recall previously learned information. Memory impairment is a primary symptom of dementia and can also be a symptom associated with such diseases as Alzheimer's disease, schizophrenia, Parkinson's disease, Huntington's disease, Pick's disease, Creutzfeldt-Jakob disease, HIV, cardiovascular
30 disease, and head trauma as well as age-related cognitive decline.

Dementias are diseases that include memory loss and additional intellectual impairment separate from memory. The present invention includes methods for treating patients suffering from memory impairment in all forms of dementia. Dementias are classified according to their cause and include: neurodegenerative dementias
5 (Alzheimer's, Parkinson's disease, Pick's disease), vascular (Infarcts, Hemorrhage, Cardiac Disorders), mixed vascular and Alzheimer's, bacterial meningitis, Creutzfeld-Jacob Disease, and multiple sclerosis), traumatic (subdural hematoma or traumatic brain injury), infectious (HIV), toxic (heavy metals, alcohol, medications), metabolic (Vitamin B₁₂ or folate deficiency), CNS hypoxia, Cushing's disease, psychiatric (depression and
10 schizophrenia) and hydrocephalus.

The present invention also includes methods for treating memory loss separate from dementias, including mild cognitive impairment (MCI) and age-related cognitive decline. The present invention includes methods of treatment for memory impairment as
15 a result of disease including Huntington's disease and Down's syndrome. According to another aspect, the invention includes methods for treating memory loss from anesthetics, chemotherapy, radiation treatment, and post-surgical trauma.

The compounds of the invention can also be used to treat schizophrenia, bipolar
20 or manic depression, major depression, and drug addiction. PDE4 inhibitors can be used to raise cAMP levels and prevent neurons from undergoing apoptosis. PDE4 inhibitors are also known to be anti-inflammatory. The combination of preventing neuronal apoptosis and inhibiting inflammatory responses make these compounds useful to treat neurodegeneration resulting from any disease or injury, including stroke, Alzheimer's
25 disease, multiple sclerosis, amyotrophic lateral sclerosis (ALS), and multiple systems atrophy (MSA).

Thus, in accordance with a preferred embodiment, the present invention includes methods of treating patients suffering from memory impairment due to, for example, mild
30 cognitive impairment due to aging, Alzheimer's disease, schizophrenia, Parkinson's disease, Huntington's disease, Pick's disease, Creutzfeld-Jakob disease, depression,

aging, head trauma, stroke, CNS hypoxia, cerebral senility, multiinfarct dementia and other neurological conditions, as well as HIV and cardiovascular diseases, comprising administering an effective amount of a compound according to formulas I-II or a pharmaceutically acceptable salt thereof.

5

As mentioned, the compounds of the invention also exhibit anti-inflammatory activity. As a result, the inventive compounds are useful in the treatment of a variety of allergic and inflammatory diseases, particularly disease states characterized by decreased cyclic AMP levels and/or elevated phosphodiesterase 4 levels. Thus, in accordance with

10 a further embodiment of the invention, there is provided a method of treating allergic and inflammatory disease states, comprising administering an effective amount of a compound according to formulas I-II or a pharmaceutically acceptable salt thereof. Such disease states include: asthma, chronic bronchitis, chronic obstructive pulmonary disease (COPD), atopic dermatitis, urticaria, allergic rhinitis, allergic conjunctivitis, vernal

15 conjunctivitis, eosinophilic granuloma, psoriasis, inflammatory arthritis, rheumatoid arthritis, septic shock, ulcerative colitis, Crohn's disease, reperfusion injury of the myocardium and brain, chronic glomerulonephritis, endotoxic shock, adult respiratory distress syndrome, cystic fibrosis, arterial restenosis, arteriosclerosis, keratosis, rheumatoid spondylitis, osteoarthritis, pyresis, diabetes mellitus, pneumoconiosis, chronic

20 obstructive airways disease, chronic obstructive pulmonary disease, toxic and allergic contact eczema, atopic eczema, seborrheic eczema, lichen simplex, sunburn, pruritis in the anogenital area, alopecia areata, hypertrophic scars, discoid lupus erythematosus, follicular and wide-area pyodermias, endogenous and exogenous acne, acne rosacea, Beghet's disease, anaphylactoid purpura nephritis, inflammatory bowel disease,

25 leukemia, multiple sclerosis, gastrointestinal diseases, autoimmune diseases, osteoporosis, and the like. The compounds can also be used in a method of treating patients suffering from disease states characterized by decreased NMDA function, such as schizophrenia. The compounds can also be used to treat psychosis characterized by elevated levels of PDE 4, for example, various forms of depression, such as manic

30 depression, major depression, and depression associated with psychiatric and neurological disorders.

The use of trisubstituted phenyl derivatives for treating asthma, chronic bronchitis, psoriasis, allergic rhinitis, and other inflammatory diseases, and for inhibiting tumor necrosis factor is known within the art. See, e.g., WO 98/58901, JP 11-189577, JP 10-072415, WO 93/25517, WO 94/14742, US 5,814,651, and US 5,935,978. These
5 references describe 1,3,4-trisubstituted phenyl compounds said to exhibit PDE4 inhibition activity. They also describe assays for determining PDE4 inhibition activity, and methods for synthesizing such compounds. The entire disclosures of these documents are hereby incorporated by reference.

10

The invention is also suitable for use in the treatment of a class of disorders known as polyglutamine-repeat diseases. These diseases share a common pathogenic mutation. The expansion of a CAG repeat, which encodes the amino acid glutamine, within the genome leads to production of a mutant protein having an expanded
15 polyglutamine region. For example, Huntington's disease has been linked to a mutation of the protein huntingtin. In individuals who do not have Huntington's disease, huntingtin has a polyglutamine region containing about 8 to 31 glutamine residues. For individuals who have Huntington's disease, huntingtin has a polyglutamine region with over 37 glutamine residues. Aside from Huntington's disease (HD), other known
20 polyglutamine-repeat diseases and the associated proteins are: dentatorubral-pallidoluysian atrophy, DRPLA (atrophin-1); spinocerebellar ataxia type-1 (ataxin-1); spinocerebellar ataxia type-2 (ataxin-2); spinocerebellar ataxia type-3 also called Machado-Joseph disease, MJD (ataxin-3); spinocerebellar ataxia type-6 (alpha 1a-voltage dependent calcium channel); spinocerebellar ataxia type-7 (ataxin-7); and spinal and
25 bulbar muscular atrophy, SBMA, also known as Kennedy disease (androgen receptor).

Thus, in accordance with a further aspect of the invention, there is provided a method of treating a polyglutamine-repeat disease or CAG repeat expansion disease comprising administering to a patient, especially a human, a therapeutically effective
30 amount of a compound according to Formulas I-II. In accordance with a further embodiment, there is provided a method of treating Huntington's disease (HD),

dentatorubral-pallidoluysian atrophy (DRPLA), spinocerebellar ataxia type-1, spinocerebellar ataxia type-2, spinocerebellar ataxia type-3 (Machado-Joseph disease), spinocerebellar ataxia type-6, spinocerebellar ataxia type-7, or spinal and bulbar muscular atrophy, comprising administering to a patient, especially a human, a therapeutically effective amount of a compound according to Formulas I-II.

PDE4 inhibitors may be used to prevent or ameliorate osteoporosis, as an antibiotic, for treatment of cardiovascular disease by mobilizing cholesterol from atherosclerotic lesions, to treat rheumatoid arthritis (RA), for long-term inhibition of mesenchymal-cell proliferation after transplantation, for treatment of urinary obstruction secondary to benign prostatic hyperplasia, for suppression of chemotaxis and reduction of invasion of colon cancer cells, for treatment of B cell chronic lymphocytic leukemia (B-CLL), for inhibition of uterine contractions, to attenuate pulmonary vascular ischemia-reperfusion injury (IRI), for corneal hydration, for inhibition of IL-2R expression and thereby abolishing HIV-1 DNA nuclear import into memory T cells, for augmentation of glucose-induced insulin secretion, in both the prevention and treatment of colitis, and to inhibit mast cell degranulation.

The compounds of the present invention can be administered as the sole active agent or in combination with other pharmaceutical agents such as other agents used in the treatment of cognitive impairment and/or in the treatment of psychosis, e.g., other PDE4 inhibitors, calcium channel blockers, chloinegic drugs, adenosine receptor modulators, amphotericin NMDA-R modulators, mGluR modulators, and cholinesterase inhibitors (e.g., donepezil, rivastigmine, and glanthanamine). In such combinations, each active ingredient can be administered either in accordance with their usual dosage range or a dose below their usual dosage range.

The dosages of the compounds of the present invention depend upon a variety of factors including the particular syndrome to be treated, the severity of the symptoms, the route of administration, the frequency of the dosage interval, the particular compound

utilized, the efficacy, toxicology profile, pharmacokinetic profile of the compound, and the presence of any deleterious side-effects, among other considerations.

5 The compounds of the invention are typically administered at dosage levels and in a mammal customary for PDE4 inhibitors such as those known compounds mentioned above. For example, the compounds can be administered, in single or multiple doses, by oral administration at a dosage level of generally 0.001-100mg/kg/day, for example, 0.01-100 mg/kg/day, preferably 0.1-70 mg/kg/day, especially 0.5-10 mg/kg/day. Unit dosage forms can contain generally 0.01-1000 mg of active compound, for example, 0.1-10 50 mg of active compound. For intravenous administration, the compounds can be administered, in single or multiple dosages, at a dosage level of, for example, 0.001-50 mg/kg/day, preferably 0.001-10 mg/kg/day, especially 0.01-1 mg/kg/day. Unit dosage forms can contain, for example, 0.1-10 mg of active compound.

15 In carrying out the procedures of the present invention it is of course to be understood that reference to particular buffers, media, reagents, cells, culture conditions and the like are not intended to be limiting, but are to be read so as to include all related materials that one of ordinary skill in the art would recognize as being of interest or value in the particular context in which that discussion is presented. For example, it is often 20 possible to substitute one buffer system or culture medium for another and still achieve similar, if not identical, results. Those of skill in the art will have sufficient knowledge of such systems and methodologies so as to be able, without undue experimentation, to make such substitutions as will optimally serve their purposes in using the methods and procedures disclosed herein.

25 The present invention will now be further described by way of the following non-limiting examples. In applying the disclosure of these examples, it should be kept clearly in mind that other and different embodiments of the methods disclosed according to the present invention will no doubt suggest themselves to those of skill in the relevant art.

30

In the foregoing and in the following examples, all temperatures are set forth uncorrected in degrees Celsius; and, unless otherwise indicated, all parts and percentages are by weight.

5 The entire disclosures of all applications, patents and publications, cited above and below, are hereby incorporated by reference.

EXAMPLES

10 **Intermediate A**

4-(3-Benzyloxy-4-methoxyphenyl)-2-pyrrolidone

To a mixture of 8.3 g (34.8 mmol) of $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ in 750 mL of MeOH was slowly added 14 g (104.5 mmol) of NaBH_4 . This mixture was stirred for 30 minutes at 0 °C and a solution of 25 g of methyl 3-(3-benzyloxy-4-methoxyphenyl)-4-nitrobutanoate in 500
15 mL of MeOH was added. Then, 8.3 g (34.8 mmol) of $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ was added to the reaction mixture, followed by the slow portion-wise addition of 9.2 g (243 mmol) of NaBH_4 . The mixture stirred at 0 °C for 1 hour and then 150 g of K_2CO_3 was added in one portion. The mixture was allowed to warm to ambient temperature and stirring continued for 18 hours. The suspension was filtered through a pad of celite, washed
20 with 2 x 1000 mL of MeOH and concentrated. The residue was taken up in 2000 mL of EtOAc and the organic fraction was successively washed with 200 mL of H_2O , 250 mL of brine, dried (Na_2SO_4) and concentrated to a solid. Trituration with hexanes/EtOAc provided 13 g of the desired product. An additional 2.3 g of product was obtained by extracting the aqueous fractions with EtOAc and combining this with the trituration
25 solvent, concentrating and triturating with EtOAc/hexanes. Total yield was 15.3 g (74% yield). ^1H NMR (400 MHz, CDCl_3) δ 7.44–7.26 (m, 5H), 6.87–6.70 (m, 3H), 5.75 (bs, 1H), 5.14 (s, 2H), 3.87 (s, 3H), 3.70–3.50 (m, 2H), 3.28 (t, 1H), 2.70–2.63 (m, 1H), 2.42–2.35 (m, 1H).

4-(3-Benzyloxy-4-difluoromethoxyphenyl)-2-pyrrolidone was prepared in a
30 similar manner starting with methyl 3-(3-benzyloxy-4-difluoromethoxyphenyl)-4-nitrobutanoate.

Intermediate B


4-(3-Hydroxy-4-methoxyphenyl)-2-pyrrolidone

A mixture of 3.5 g (11.6 mmol) 4-(3-Benzyloxy-4-methoxyphenyl)-2-pyrrolidone and 350 mg of 10% Pd/C in 50 mL of MeOH and 10 mL of CH₂Cl₂ was shaken on a Paar apparatus under 20 psi H₂ for 8 hours. The mixture was filtered through celite and concentrated leaving 2.4 g (99% yield) of crude material used as such for Examples 1 through 3. ¹H NMR (300 MHz, CDCl₃) δ 6.82 (s, 1H), 6.79 (d, J=7.6 Hz, 1H), 6.70 (d, J=7.6Hz, 1H), 6.50 (s, 1H), 3.87 (s, 3H), 3.77 (t, J=8.4Hz, 1H), 3.62 (t, J=8.0Hz, 1H), 3.40 (t, J=8.4Hz, 1H), 2.71 (dd, J=16.8, 8.8 Hz, 1H), 2.49 (dd, J= 16.8, 9.0 Hz, 1H).

4-(4-Difluoromethoxy-3-hydroxyphenyl)-2-pyrrolidone was prepared in a similar fashion starting with 4-(3-benzyloxy-4-difluoromethoxyphenyl)-2-pyrrolidone

Example 1

4-[3-(4-Chlorophenoxy)-4-methoxyphenyl]-2-pyrrolidone



A mixture of 4-(3-hydroxy-4-methoxyphenyl)-2-pyrrolidone (63 mg, 0.3 mmol), 4-chlorophenylboronic acid (61 mg, 0.45 mmol), copper (II) acetate (54 mg, 0.3 mmol), triethylamine (152 mg, 1.5 mmol), dichloromethane (3 ml) and small amount of molecular sieves is allowed to stir at ambient temperature. After 18 hours, the mixture is filtered over celite and the filtrate is concentrated under vacuum. The resulting residue is dissolved in 30 ml of ethyl acetate and washed successively with saturated aqueous sodium bicarbonate solution and brine. Purification by flash column chromatography (ethyl acetate/hexane 1:1 to methanol/dichloromethane 3:97) gives 26 mg (27%) of 4-(3-(4-chlorophenoxy)-4-methoxyphenyl)-2-pyrrolidone: ¹H NMR (300 MHz, CDCl₃) δ 7.24 (d, J = 9.2 Hz, 2 H), 7.03 (d, J = 8.4 Hz, 1H), 6.95(d, J = 8.4 Hz, 1H), 6.87 (s, 1H),

6.83(d, $J = 9.2$ Hz, 2H), 5.83 (b, 1H), 3.80 (s, 3H), 3.72 (t, $J = 8.4$ Hz, 1H), 3.65-3.62 (m, 1H), 3.34 (t, $J = 8.4$ Hz, 1H), 2.68 (dd, $J = 16.8, 8.8$ Hz, 1H), 2.41 (dd, $J = 16.8, 9.2$ Hz, 1H).


5 The following compounds were prepared in a similar fashion starting with different arylboronic acids and aryl vinyl boronic acids:

- A. 4-(4-Methoxy-3-phenoxyphenyl)-2-pyrrolidone
- B. 4-[4-Methoxy-3-(4-methoxyphenoxy)phenyl]-2-pyrrolidone
- C. 4-[4-Methoxy-3-(3-thienyloxy)phenyl]-2-pyrrolidone
- 10 D. 4-[3-(4-Fluorophenoxy)-4-methoxyphenyl]-2-pyrrolidone
- E. 4-[3-Cyanophenoxy-4-methoxyphenyl]-2-pyrrolidone
- F. 4-[4-Methoxy-3-naphthyloxyphenyl]-2-pyrrolidone
- G. 4-[4-Methoxy-3-((2-phenyl)ethenyloxy)phenyl]-2-pyrrolidone
- H. 4-[4-Methoxy-3-((2-(4-chlorophenyl)ethenyloxy)phenyl]-2-pyrrolidone

15 The separate enantiomers of racemic compounds listed above can be obtained through the use of optically active starting materials or by conventional resolution techniques such as chiral HPLC.

Example 2

20 **4-(3-Cinnamyloxy-4-methoxyphenyl)-2-pyrrolidone**



25 Diisopropylazodicarboxylate (61 mg, 0.6 mmol) is added to a solution of 4-(3-hydroxy-4-methoxyphenyl)-2-pyrrolidone (63 mg, 0.3 mmol), cinnamyl alcohol (43 mg, 0.3 mmol) and triphenyl phosphine (157 mg, 0.6 mmol) in 3 ml of tetrahydrofuran and the reaction mixture is stirred at 70° C. After 12 hours, the same work-up procedure as in

Example 1 yields 70 mg (70%) of 4-(3-cinnamyloxy-4-methoxyphenyl)-2-pyrrolidone:

¹H NMR (300 MHz, CDCl₃) δ 7.40 (d, *J* = 6.8 Hz, 2H), 7.31 (t, *J* = 6.8 Hz, 2H), 7.25 (d, *J* = 7.2 Hz, 1H), 6.86-6.81 (m, 2H), 6.71 (d, *J* = 15.6 Hz, 1H), 6.42 (m, 1H), 5.92 (b, 1H), 4.76 (dd, *J* = 6.0, 1.2 Hz, 2H), 3.87 (s, 3H), 3.71 (t, *J* = 8.8 Hz, 1H), 3.64-3.56 (m, 1H),
5 3.35 (t, *J* = 8.8 Hz, 1H).

The following compounds were prepared in a similar fashion starting with different alcohols:


- A. 4-(3-(3-Cyclohexyl-1-propyloxy)-4-methoxyphenyl)-2-pyrrolidone
- 10 B. 4-(4-Methoxy-3-(3-phenyl-1-butyloxy)phenyl)-2-pyrrolidone
- C. 4-[4-Methoxy-3-indanyloxyphenyl]-2-pyrrolidone
- D. 4-[4-Methoxy-3-(2-(4-chlorophenethyl)ethoxy)phenyl]-2-pyrrolidone
- E. 4-[4-Methoxy-3-(2-(4-methylphenyl)ethoxy)phenyl]-2-pyrrolidone
- F. 4-[4-Methoxy-3-(2-(2-thienyl)ethoxy)phenyl]-2-pyrrolidone
- 15 G. 4-[4-Methoxy-3-(2-(3-chlorophenyl)ethoxy)phenyl]-2-pyrrolidone
- H. 4-[4-Methoxy-3-(2-(4-fluorophenyl)ethoxy)phenyl]-2-pyrrolidone
- I. 4-[4-Methoxy-3-(2-(4-pyridyl)ethoxy)phenyl]-2-pyrrolidone
- J. 4-[4-Methoxy-3-(3-(*R*)-tetrahydrofuryloxy)phenyl]-2-pyrrolidone
- K. 4-[4-Methoxy-3-(3-(*S*)-tetrahydrofuryloxy)phenyl]-2-pyrrolidone
- 20 L. 4-[4-Methoxy-3-(2-(4-cyanophenyl)ethoxy)phenyl]-2-pyrrolidone
- M. 4-[4-Methoxy-3-(3-(4-chlorophenyl)propoxy)phenyl]-2-pyrrolidone
- N. 4-[4-Methoxy-3-(2-(3-methylphenyl)ethoxy)phenyl]-2-pyrrolidone
- O. 4-(*R*)-[4-Methoxy-3-(3-(*R*)-tetrahydrofuryloxy)phenyl]-2-pyrrolidone
- P. 4-(*S*)-[4-Methoxy-3-(3-(*R*)-tetrahydrofuryloxy)phenyl]-2-pyrrolidone
- 25 Q. 4-(*R*)-[4-Methoxy-3-(3-(*S*)-tetrahydrofuryloxy)phenyl]-2-pyrrolidone
- R. 4-(*S*)-[4-Methoxy-3-(3-(*S*)-tetrahydrofuryloxy)phenyl]-2-pyrrolidone
- S. 4-[4-Methoxy-3-cyclopropylmethoxyphenyl]-2-pyrrolidone
- T. 4-[4-Methoxy-3-cyclobutylmethoxyphenyl]-2-pyrrolidone
- U. 4-[4-Methoxy-3-cycloheptyloxyphenyl]-2-pyrrolidone
- 30 V. 4-[4-Methoxy-3-cyclohexylmethoxyphenyl]-2-pyrrolidone
- W. 4-[4-Methoxy-3-(2-cyclohexyl)ethoxyphenyl]-2-pyrrolidone

- X. 4-[4-Methoxy-3-cyclopentylmethoxyphenyl]-2-pyrrolidone
 Y. 4-[4-Methoxy-3-cyclohexyloxyphenyl]-2-pyrrolidone
 Z. 4-[4-Methoxy-3-(3-cyclopentyl)propoxyphenyl]-2-pyrrolidone
 AA. 4-[4-Difluoromethoxy-3-(3-(*R*)-tetrahydrofuryloxy)phenyl]-2-pyrrolidone
 5 BB. 4-[4-Difluoromethoxy-3-(2-cyclohexylethoxy)phenyl]-2-pyrrolidone
 CC. 4-[4-Difluoromethoxy-3-(2-phenylethoxy)phenyl]-2-pyrrolidone
 DD. 4-[4-Difluoromethoxy-3-(cyclopropylmethoxy)phenyl]-2-pyrrolidone

The separate enantiomers of racemic compounds listed above can be obtained
 10 through the use of optically active starting materials or by conventional resolution
 techniques such as chiral HPLC.

Example 3

15 4-(4-Methoxy-3-phenpropyloxyphenyl)-2-pyrrolidone



A mixture of 4-(3-hydroxy-4-methoxyphenyl)-2-pyrrolidone (42 mg, 0.2 mmol),
 20 3-phenylpropyl bromide (44 mg, 0.22 mmol), potassium carbonate (83 mg, 0.6 mmol)
 and *N,N*-dimethylformamide (3 ml) is allowed to stir at 100 °C. After 18 hours, the
 reaction mixture is cooled to 25 °C. The solid is filtered off and the filtrate is
 concentrated under vacuum. The same working up procedure gives 47 mg (72%) of 4-(3-
 (3-phenylpropoxy)-4-methoxyphenyl)-2-pyrrolidone: ¹H NMR (300 MHz, CDCl₃) δ
 25 7.34-7.17 (m, 5H), 6.82 (d, *J* = 8.4 Hz, 1H), 6.78 (d, *J* = 12 Hz, 1H), 6.70 (s, 1H), 6.02 (b,
 1H), 4.03 (t, *J* = 7.2 Hz, 2 H), 3.84 (s, 3H), 3.75 (t, *J* = 8.8 Hz, 1H), 3.63-3.55 (m, 1H),

3.36 (t, $J = 8.8$ Hz, 1H), 2.82 (t, $J = 8.0$ Hz, 2H), 2.71 (dd, $J = 16.8, 8.8$ Hz, 2H), 2.47 (dd, $J = 16.6, 8.8$ Hz), 2.20-2.13 (m, 2H).


The following compounds were prepared in a similar fashion starting with different arylalkyl bromides and alkyl bromides:

- A. 4-(4-Methoxy-3-(2-phenylethoxy)phenyl)-2-pyrrolidone
- B. 4-(4-Methoxy-3-(3-phenyl-1-propoxy)phenyl)-2-pyrrolidone
- C. 4-[4-methoxy-3-[1-(3-phenyl-2-propenyl)oxyphenyl]-2-pyrrolidone
- D. 4-[4-methoxy-3-(2-cyclopropylethoxy)phenyl]-2-pyrrolidone
- 10 E. 4-[4-methoxy-3-(2-cyclopentylethoxy)phenyl]-2-pyrrolidone
- F. 4-[4-Difluoromethoxy-3-cyclopentoxypheyl]-2-pyrrolidone
- G. 4-[4-Difluoromethoxy-3-cyclobutylmethoxyphenyl]-2-pyrrolidone

The separate enantiomers of racemic compounds listed above can be obtained through the use of optically active starting materials or by conventional resolution techniques such as chiral HPLC.

Example 4

(4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,3-difluorobenzyl)-2-pyrrolidone



20

4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone (110 mg, 0.4 mmol) in 2 ml of *N, N*-dimethylformamide and 110 μ l of 15-crown-5 is treated with sodium hydride (20 mg, 0.5 mmol) under nitrogen at ambient temperature. After 3 hours, the mixture is cooled to 0°C and treated with 2,3-difluorobenzyl bromide (166 mg, 0.8 mmol) in 2 ml of tetrahydrofuran. The resulting mixture is allowed to stir at ambient temperature for 6 hours. 100 ml of ethyl acetate is added, followed by 100 ml of ice cold water. After

usual aqueous work up, the crude product is purified by flash column chromatography (ethyl acetate/hexane 2:1) to give 112 mg (70%) of 1-(2,3-difluorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone: ¹H NMR (300 MHz, CDCl₃) δ 7.18-6.98 (m, 3H), 6.79 (d, *J* = 8.1 Hz, 1H), 6.90 (d, *J* = 8.1 Hz, 1H), 6.68 (s, 1H) 4.71-4.69 (m, 1H), 4.60 (s, 2H), 3.81 (s, 3H), 3.68 (t, *J* = 9.0 Hz, 1H), 3.66-3.54 (m, 1H), 3.29 (dd, *J* = 9.6, 7.2 Hz, 1H), 2.84 (dd, *J* = 16.8, 9.0 Hz, 1H), 2.57 (dd, *J* = 16.8, 8.4 Hz, 1H), 2.00-1.75 (b, 6H), 1.70-1.52 (b, 2H).

The following compounds were prepared in a similar fashion starting with different 4-(3-Substituted)-oxy-4-methoxyphenyl)-2-pyrrolidone and reacting it with different arylalkyl bromides and alkyl bromides:

- A. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-methylbenzyl)-2-pyrrolidone
- B. (4*S*)-1-(2-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone
- C. (4*S*)-1-(4-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone
- 15 D. (4*S*)-1-(3-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone
- E. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-methoxybenzyl)-2-pyrrolidone
- F. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-nitrobenzyl)-2-pyrrolidone
- G. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-nitrobenzyl)-2-pyrrolidone
- H. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-fluorobenzyl)-2-pyrrolidone
- 20 I. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-fluorobenzyl)-2-pyrrolidone
- J. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-fluorobenzyl)-2-pyrrolidone
- K. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-benzyl-2-pyrrolidone
- L. (4*S*)-1-(4-Cyanobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone
- M. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-trifluoromethylbenzyl)-2-pyrrolidone
- 25 N. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-trifluoromethylbenzyl)-2-pyrrolidone
- O. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-trifluoromethylbenzyl)-2-pyrrolidone
- 30 P. (4*S*)-1-(3,5-Bistrifluoromethylbenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone

- Q. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,4-difluorobenzyl)-2-pyrrolidone
- R. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,5-difluorobenzyl)-2-pyrrolidone
- S. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,4-difluorobenzyl)-2-pyrrolidone
- T. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,6-difluorobenzyl)-2-pyrrolidone
- 5 U. (4*S*)-1-(2-Chloro-4-fluorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone
- V. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,4-dichlorobenzyl)-2-pyrrolidone
- W. (4*S*)-1-(4-*tert*-Butylbenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone
- X. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-ethyl-2-pyrrolidone
- 10 Y. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-propyl-2-pyrrolidone
- Z. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-butyl-2-pyrrolidone
- AA. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-methoxyethyl)-2-pyrrolidone
- BB. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-phenylbenzyl)-2-pyrrolidone
- CC. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-methylbenzyl)-2-pyrrolidone
- 15 DD. (4*R*)-1-(2-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone
- EE. (4*R*)-1-(4-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone
- FF. (4*R*)-1-(3-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone
- GG. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-methoxybenzyl)-2-pyrrolidone
- 20 HH. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-nitrobenzyl)-2-pyrrolidone
- II. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-nitrobenzyl)-2-pyrrolidone
- JJ. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-fluorobenzyl)-2-pyrrolidone
- KK. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-fluorobenzyl)-2-pyrrolidone
- 25 LL. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-fluorobenzyl)-2-pyrrolidone
- MM. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-benzyl-2-pyrrolidone
- NN. (4*R*)-1-(4-Cyanobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone
- OO. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-trifluoromethylbenzyl)-2-pyrrolidone
- 30 PP. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-trifluoromethylbenzyl)-2-pyrrolidone

- QQ. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-trifluoromethylbenzyl)-2-pyrrolidone
- RR. (4*R*)-1-(3,5-bis(trifluoromethyl)benzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone
- 5 SS. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,4-difluorobenzyl)-2-pyrrolidone
- TT. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,5-difluorobenzyl)-2-pyrrolidone
- UU. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,4-difluorobenzyl)-2-pyrrolidone
- 10 VV. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,6-difluorobenzyl)-2-pyrrolidone
- WW. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,3-difluorobenzyl)-2-pyrrolidone
- 15 XX. (4*R*)-1-(2-Chloro-4-fluorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone
- YY. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,4-dichlorobenzyl)-2-pyrrolidone
- ZZ. (4*R*)-1-(4-*tert*-Butylbenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone
- 20 AAA. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-ethyl-2-pyrrolidone
- BBB. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-propyl-2-pyrrolidone
- CCC. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-butyl-2-pyrrolidone
- DDD. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-methoxyethyl)-2-pyrrolidone
- 25 EEE. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-phenylbenzyl)-2-pyrrolidone
- FFF. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-chloro-4-fluorobenzyl)-2-pyrrolidone
- GGG. (4*R*)-(3-cyclopentyloxy-4-methoxyphenyl)-1-methoxycarbonylmethyl-2-pyrrolidone
- 30 HHH. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-cyanomethyl-2-pyrrolidone
- III. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-cyclopentyl-2-pyrrolidone


- JJJ. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-methoxybenzoyl)-2-pyrrolidone
 KKK. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-chlorobenzoyl)-2-pyrrolidone
 LLL. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-nitrobenzoyl)-2-pyrrolidone
 MMM. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-benzoyl-2-pyrrolidone
 5 NNN. (4*R*)-1-(2,3-Difluorobenzyl)-4-(3-(3-(*S*)-tetrahydrofuryl)oxy-4-methoxyphenyl)-
 2-pyrrolidone
 OOO. (4*S*)-1-(2,3-Difluorobenzyl)-4-(3-(3-(*S*)-tetrahydrofuryl)oxy-4-methoxyphenyl)-2-
 pyrrolidone
 PPP. (4*R*)-1-(2,3-Difluorobenzyl)-4-(3-(3-(*R*)-tetrahydrofuryl)oxy-4-methoxyphenyl)-
 10 2-pyrrolidone
 QQQ. (4*S*)-1-(2,3-Difluorobenzyl)-4-(3-(3-(*R*)-tetrahydrofuryl)oxy-4-methoxyphenyl)-
 2-pyrrolidone
 RRR. (4*S*)-1-[2-(3-Chlorophenoxy)ethyl]-4-[4-methoxy-3-(3*R*)-
 tetrahydrofuranyloxyphenyl]-2-pyrrolidone
 15 SSS. (4*S*)-4-[4-Methoxy-3-(3*R*)-tetrahydrofuranyloxyphenyl]-1-(2-oxo-2-phenylethyl)-
 2-pyrrolidone
 TTT. (4*S*)-4-[4-Methoxy-3-(3*R*)-tetrahydrofuranyloxyphenyl]-1-(2-oxo-2-(4-
 methoxyphenyl)ethyl)-2-pyrrolidone

20

The separate enantiomers of racemic compounds listed above can be obtained through the use of optically active starting materials or by conventional resolution techniques such as chiral HPLC.

25 **Example 5 (intermediate)**

(4*S*)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone-1-acetic acid




Methyl (4S)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone-1-acetate (250 mg, 0.72 mmol) was treated with 8 ml of a 1M solution of potassium hydroxide in 95% methanol. After 10 hours, the reaction solution was acidified with 2N HCl to pH 3 and 30 ml of water was added and the resulting mixture was extracted with 60 ml of ethyl acetate twice. The organic layer was concentrated and chromatographed (methanol/dichloromethane 5:100) to give 242 mg (quantitative yield) of 4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone-1-acetic acid: ¹H NMR (300 MHz, CDCl₃) δ 6.85-7.3 (m, 3H), 4.84-4.74 (m, 1H), 4.20 (d, *J* = 12.0 Hz, 1H), 4.12 (d, *J* = 12.0, 1H), 3.82 (s, 4H), 3.66-3.46 (m, 2H), 2.89 (dd, *J* = 16.8, 8.7 Hz, 1H), 2.58 (dd, *J* = 16.8, 8.4 Hz, 1H), 2.00-1.72 (b, 6H), 1.70-1.50 (b, 2H).

The following compounds were formed in a similar fashion with different starting materials:

A. (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone-1-acetic acid

Example 6

(4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone-1-acetyl chloride



20

(4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone-1-acetic acid (50 mg, 0.15 mmol) was dissolved in 3 mL of CH₂Cl₂ and treated with oxalylchloride (0.17 mmol) at 4°C. The reaction mixture was stirred at ambient temperature for 2 hours and concentrated in vacuo.

25


The following compounds were formed in a similar fashion with different starting materials:

4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone-1-acetyl chloride,
(4R)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone-1-acetyl chloride.

5

Example 7

4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2,6-dimethylphenyl)-aminocarbonylmethyl)-2-pyrrolidone



10

4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone-1-acetyl chloride was taken up in 3 ml of CH₂Cl₂ and added to a solution of 1,6-dimethylaniline (39 mg, 0.3 mmol) in tetrahydrofuran (2 ml) containing diisopropylethylamine (39 mg, 0.3 mmol).

15 After 3 hours, the mixture is concentrated in vacuo, the residue was dissolved in 50 ml of EtOAc, washed with 50 ml of 1 N HCl, 50 mL of water, and 50 mL of aqueous saturated sodium bicarbonate solution. The organic phase was concentrated and chromatographed over SiO₂ to give 49 mg (75% yield) of the target acetamide: ¹H NMR (300 MHz, CDCl₃) δ 7.68 (s, 1H), 7.18-7.03 (m, 3H), 6.84-6.73 (m, 3H), 4.84-4.70 (m, 1H), 4.16 (d, J = 11.4 Hz, 1H), 4.13 (d, J = 11.4 Hz, 1H), 3.94 (t, J = 8.0 Hz, 1H), 3.85-3.75 (m, 4H), 3.68-3.46 (m, 2H), 2.86 (dd, J = 16.8, 8.4 Hz, 1H), 2.61 (dd, J = 16.8, 8.4 Hz, 1H), 2.19 (s, 6H), 1.98-1.72 (b, 6H), 1.70-1.53 (b, 2H).

25 The following compounds were formed in a similar fashion with different starting materials:

A. (4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2,6-dimethylphenyl)-aminocarbonylmethyl)-2-pyrrolidone

- B. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2,6-dimethylphenyl)-aminocarbonylmethyl)-2-pyrrolidone
- C. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone
- 5 D. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone
- E. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone
- F. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2,3-difluorophenyl)-aminocarbonylmethyl)-2-pyrrolidone
- 10 G. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(3-chlorophenyl)-aminocarbonylmethyl)-2-pyrrolidone
- H. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(4-pyridyl)-aminocarbonylmethyl)-2-pyrrolidone
- 15 I. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(4-methoxyphenyl)-aminocarbonylmethyl)-2-pyrrolidone
- J. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(4-chloro-2-fluorophenyl)-aminocarbonylmethyl)-2-pyrrolidone
- K. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(3-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone
- 20 L. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(4-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone
- M. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(4-nitrophenyl)-aminocarbonylmethyl)-2-pyrrolidone
- 25 N. (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2,3-difluorophenyl)-aminocarbonylmethyl)-2-pyrrolidone
- O. (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2,3-difluorophenyl)-aminocarbonylmethyl)-2-pyrrolidone
- P. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(3-(2-chloropyridinyl))-aminocarbonylmethyl)-2-pyrrolidone
- 30

- Q. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-(2-pyridylethyl))-aminocarbonylmethyl)-2-pyrrolidone
- R. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-(2-(*N*-methylpyrrolidinyl)ethyl))-aminocarbonylmethyl)-2-pyrrolidone
- 5 S. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(4-pyridylmethyl))-aminocarbonylmethyl)-2-pyrrolidone
- T. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(1-imidazolylpropyl))-aminocarbonylmethyl)-2-pyrrolidone
- U. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(4-(4-methylpiperazinyl))-10 hydrazinocarbonylmethyl)-2-pyrrolidone
- V. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-(*N*-methylpyrrolidinyl)methyl))-aminocarbonylmethyl)-2-pyrrolidone
- W. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(4-pyrimidinyl))-aminocarbonylmethyl)-2-pyrrolidone
- 15 X. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3-(5-methylisoxazolyl))-aminocarbonylmethyl)-2-pyrrolidone
- Y. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2,6-dimethylpiperazinyl))-hydrazinocarbonylmethyl)-2-pyrrolidone
- Z. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-(4,5-dimethylthiazolyl))-20 aminocarbonylmethyl)-2-pyrrolidone
- AA. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-(2-methylpiperidinyl)ethyl))-aminocarbonylmethyl)-2-pyrrolidone
- BB. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3,4-dimethoxyphenethyl))-aminocarbonylmethyl)-2-pyrrolidone
- 25 CC. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-pyridinyl))-aminocarbonylmethyl)-2-pyrrolidone
- DD. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3-pyridinyl))-aminocarbonylmethyl)-2-pyrrolidone
- EE. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3-(2,4-dimethoxypyridinyl))-30 aminocarbonylmethyl)-2-pyrrolidone

FF. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3-(4-methoxypyridinyl))-aminocarbonylmethyl)-2-pyrrolidone


GG. 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(4-tert-butylloxycarbonylaminophenyl)-aminocarbonylmethyl)-2-pyrrolidone

5

The separate enantiomers of racemic compounds listed above can be obtained through the use of optically active starting materials or by conventional resolution techniques such as chiral HPLC.

10 **Example 8**

4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(4-anilino)-aminocarbonylmethyl)-2-pyrrolidone



15


A mixture of 4-(3-cyclopentyloxy-4-methoxyphenyl)-1-(N-(4-tert-butylloxycarbonylaminophenyl)-aminocarbonylmethyl)-2-pyrrolidone (see Serial No. 10/270,724, filed October 16, 2002) was stirred in a 1:2 mixture of TFA and dichloromethane for 2 hours at room temperature. The material was concentrated in vacuo and purified by column chromatography over SiO₂ to provide the pure aniline. ¹H NMR (300 MHz, CDCl₃) δ 7.90 (s, 1H), 7.24 (d, J = 6.3 Hz, 2H), 6.83-6.75 (m, 2H), 6.64 (d, J = 6.3 Hz, 2H), 4.74 (m, 1H), 4.13 (d, J = 15.6 Hz, 1H), 4.03 (d, J = 15.6 Hz, 1H), 3.92 (m, 1H), 3.83 (s, 3H), 3.70-3.50 (m, 3H), 2.84 (dd, J = 16.5, 7.8 Hz, 1H), 2.61 (dd, J = 16.5, 7.8 Hz, 1H), 1.95-1.80 (m 6H), 1.59 (m, 2H).

25 The following compounds were formed in a similar fashion with different starting materials:

(4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuryloxyphenyl]-1-[N-(5-(4-piperidinyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone.

Example 9

5 (4*S*)-1-[N-(2,3-Difluorophenyl-(N-methyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3*R*)-tetrahydrofuran-2-ylphenoxy)-2-pyrrolidone



10 (4*S*)-1-[N-(2,3-Difluorophenylaminocarbonylmethyl)]-4-(4-methoxy-3-(3*R*)-tetrahydrofuran-2-ylphenoxy)-2-pyrrolidone (50 mg, 0.112 mmol) was dissolved in 2 mL of DMF in a N₂ purged flask, cooled to 0 °C with stirring, and 4.9 mg (0.12 mmol) of NaH was added. The reaction mixture stirred at 0 °C for 1 hour followed by the addition of 10.5 μL (0.168 mmol) of iodomethane. The reaction mixture was warmed to room
15 temperature and stirred over night, then concentrated and purified by preparative TLC using MeOH/EtOAc/Hexanes (7/46/46) as eluant to provide 34.5 mg (67% yield) of (4*S*)-1-[N-(2,3-Difluorophenyl-(N-methyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3*R*)-tetrahydrofuran-2-ylphenoxy)-2-pyrrolidone. ¹H NMR (CDCl₃) δ 7.25-7.17 (m, 3H), 6.90-6.78 (m, 3H), 5.02 (broad d, 1H), 4.05-4.02 (m, 3H), 3.97-3.89 (m, 2H), 3.83 (s, 3H), 3.73-3.39 (m, 4H), 3.27 (s, 3H), 2.83-2.78 (m, 1H), 2.52-2.48 (m, 1H), 2.17 (broad s, 2H).
20

The following compounds were made using a similar procedure with different starting materials:


- 25 A. (4*S*)-4-(4-Methoxy-3-(3*R*)-tetrahydrofuran-2-ylphenoxy)-1-[N-(2-methylphenyl-(N-methyl)aminocarbonylmethyl)]-2-pyrrolidone
B. (4*S*)-4-(4-Methoxy-3-(3*R*)-tetrahydrofuran-2-ylphenoxy)-1-[N-(2-(6-methylpyridyl)-(N-methyl)aminocarbonylmethyl)]-2-pyrrolidone

- C. (4S)-1-[N-(2,3-Difluorophenyl-(N-ethyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone
- D. (4S)-1-[N-(2,3-Difluorophenyl-(N-isopropyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone
- 5 E. (4S)-1-[N-(2,3-Difluorophenyl-(N-cyclopropylmethyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone
- F. (4S)-1-[N-(2,3-Difluorophenyl)-N-(2-methylpropyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- G. (4S)-1-[N-(6-Fluorobenzothiazol-2-yl)-N-(methyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- 10 H. (4S)-1-[N-(3-Fluorophenyl)-N-(methyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- I. (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methoxyphenyl)-N-(methyl)aminocarbonylmethyl]-2-pyrrolidone
- 15 J. (4S)-1-[N-(4-Isopropylphenyl)-N-(methyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- K. (4S)-1-[N-(3,4-Methylenedioxyphenyl)-N-(methyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- L. 4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-methyl-N-(thiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 20 M. (4S)-1-[N-(Benzothiazol-2-yl)-N-(cyclopropylmethyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- N. (4S)-1-[N-Cyclopropylmethyl-N-(6-fluorobenzothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone

25

Example 10A

(4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(phenylaminocarbonylmethyl)]-2-pyrrolidone



To a flask containing (4S)-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone-1-acetic acid (108 mg; 0.32 mmol) in 5 mL of dichloromethane was added 1-hydroxybenzotriazole (54 mg; 0.35 mmol) with stirring. After 5 minutes N,N-diisopropylethylamine (61 μ L; 0.35 mmol), aniline (32 μ L; 0.35 mmol) and O-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU) (132 mg; 0.35 mmol) were added. The reaction was stirred at room temperature overnight, diluted with ethyl acetate (50 mL) and washed with 25 mL portions of water, 1M HCl, saturated K_2CO_3 and brine. After drying over sodium sulfate the solvent was removed under reduced pressure to afford 132 mg (100%) of (4S)-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(phenylaminocarbonylmethyl)]-2-pyrrolidone as an off white solid. 1H NMR ($CDCl_3$; 300 MHz) 8.2 (br s, 1H); 7.5 (d, 2H); 7.3 (t, 2H); 7.1 (t, 2H); 6.9 (s, 2H); 6.7 (s, 1H); 4.9 (br s, 1H); 3.8-4.2 (m, 7 H); 3.8 (s, 3H); 3.6 (m, 2H), 2.9 (m, 2H); 2.5 (m, 1H) 2.1 (m, 2H) ppm; MS $[M+H]=411$; $[M+Na]=433$.


The following compounds were made using a similar procedure with different starting materials:

- A. (4S)-1-[N-(4,5-Dimethylthiazol-2-yl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone
- B. (4S)-1-[N-(3-Chlorophenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone
- C. (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(3-methoxycarbonylphenyl)aminocarbonylmethyl]-2-pyrrolidone
- D. (4S)-1-[N-(3-Fluorophenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone

- E. (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-thiazolyl)aminocarbonylmethyl]-2-pyrrolidone
- F. (4S)-1-[N-(4-Methoxyphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone
- 5 G. (4S)-1-[N-(2,6-Dimethylphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone
- H. (4S)-1-[N-(4-Isopropylphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone
- I. (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(3,4-methylenedioxyphenyl)aminocarbonylmethyl]-2-pyrrolidone
- 10 J. (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-(4-trifluoromethyl)pyridyl)aminocarbonylmethyl]-2-pyrrolidone
- K. (4S)-1-[N-(2-(5-Chlorobenzoxazolyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone
- 15 L. (4S)-1-[N-(2-(Benzthiazolyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone
- M. (4S)-1-[N-(2-(6-Fluorobenzthiazolyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone
- N. (4S)-1-[N-(2-(Benzimidazolyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone
- 20 O. (4S)-1-[N-(2,3-Difluorophenylaminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone
- P. (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(4-ethoxycarbonylphenyl)aminocarbonylmethyl]-2-pyrrolidone
- 25

Example 10B

(4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methylbenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone



(4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone-1-acetic acid (700 mg; 2.09 mmol) was stirred in dichloromethane (35 mL) and HOBT (543 mg; 3.55 mmol) was added in one portion followed by addition of N,N-diisopropylethylamine (0.62 mL; 3.55 mmol), 2-amino-4-methylbenzothiazole (583 mg; 3.55 mmol) and EDCI HCl (680 mg; 3.55 mmol). Stirring continued at room temperature for 48 hours and the reaction monitored by LC-MS analysis (peak with $[M+H]^+ = 482.1$ corresponding to the desired product). A precipitate was observed in the reaction. The solvent was evaporated under reduced pressure and the residue was partitioned between ethyl acetate (100 mL) and 1M HCl (50 mL). The organic layer was washed with aqueous sodium bicarbonate (50 mL), brine (25mL) and dried over sodium sulfate. Evaporation afforded (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methylbenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone as a white solid (360 mg; 36%). ^1H NMR (CDCl_3 ; 300 MHz) δ 2.2 (m, 2H); 2.6 (s, 3H); 2.7 (m, 1H); 3.0 (m, 1H); 3.6 (m, 2H); 3.8 (s, 3H); 3.9-4.1 (m, 5H); 4.3 (s, 2H); 5.0 (m, 1H); 6.8 (s, 1H); 6.9 (s, 2H); 7.3 (m, 3H) 7.7 (m, 1H). ES-MS $[M+H]^+ = 482.2$ Elemental analysis calculated for $\text{C}_{25}\text{H}_{27}\text{N}_3\text{O}_5\text{S}$; %C 62.35 %H 5.65 %N 8.73; found %C 62.08 %H 5.89 %N 8.65

The following compounds were made using the same method with different starting materials:

- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methylbenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-methylthiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(6-methylbenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone


- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methoxybenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-1-[N-(6-Ethoxycarbonylbenzothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- 5 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(6-trifluoromethoxylbenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-1-[N-(4-tert-Butylthiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(4-Isopropoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- 10 (4S)-1-[N-(4-Fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[2-(1,2,3,4-tetrahydroisoquinoliny)carbonylmethyl]-2-pyrrolidone
- 15 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[1-(1,2,3,4-tetrahydroquinoliny)carbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-trifluoromethoxyphenyl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methylthiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 20 (4S)-1-[N-(4-tert-Butylphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(6-Chlorobenzothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- 25 (4S)-1-[N-(Indan-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(5-Chlorothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-phenylthiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 30

- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(6-methoxybenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-1-[N-(5-Cyclopropyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- 5 (4S)-1-[N-(Benzothiazol-6-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(4-Ethoxycarbonylthiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(5-tert-Butyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-
- 10 (3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(2,4-Dimethoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(3,5-Dimethoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- 15 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(2,2,3,3-tetrafluorobenzo-1,4-dioxan-6-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-1-[N-(3,4-(Difluoromethylene)dioxyphenyl)-N-methylaminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(3-Fluoro-4-methoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-
- 20 tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(1,4-Benzodioxan-6-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(2-Fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- 25 (4S)-1-[N-(3,4-Dimethoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(3,4-Difluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(4-Methanesulfonamidophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-
- 30 tetrahydrofuranyloxyphenyl]-2-pyrrolidone

- (4S)-1-[N-(4-(4-Fluorophenyl)thiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(3-Fluoro-4-methylphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- 5 (4S)-1-[N-(4,6-Difluorobenzothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4R)-1-[N-(3-Fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3S)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(4-tert-Butyloxycarbonyl-3-fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- 10 (4S)-1-[N-(4-Ethanesulfonamidophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(4-Benzenesulfonamidophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- 15 (4S)-1-[N-(4-(4-Fluorobenzene)sulfonamidophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(2,3-Difluorobenzyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(5-Cyclopropylmethyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- 20 (4S)-1-[N-(6-Ethylpyridin-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(3-Fluorobenzyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- 25 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(2-methylbenzyl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-1-[N-(4-Methanesulfonylbenzyl)aminocarbonylmethyl]-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(4-Aminosulfonylbenzyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- 30

- (4S)-1-[N-(Benzothiazol-2-yl)methylaminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(3-methylpyridin-2-yl)methylaminocarbonylmethyl]-2-pyrrolidone
- 5 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-trifluoromethyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-pyridyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-(3-pyridyl)thiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 10 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-(2-pyridyl)thiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-(4-pyridyl)thiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 15 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-pyridyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-ethoxycarbonyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 20 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-methoxycarbonyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(3,4-methylenedioxyphenyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 25 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-thienyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 30 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-thienylmethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-propyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 35 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-pyrazinyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone

- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-methoxymethyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 5 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-tetrahydrofuranyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-aminosulfonyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 10 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-methoxyphenyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-methoxyphenyloxymethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 15 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-morpholinylcarbonylmethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(1-piperidinylcarbonylmethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 20 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(1-pyrrolidinylcarbonylmethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-(N-tertbutyloxycarbonyl)piperidinyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 25 (4S)-1-[N-(4-Ethoxycarbonylthiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- 30 (4S)-1-[N-(4-tert-butyloxycarbonyl-3-fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone
- 35 **Example 11**
- (4S)-1-[N-(4-Carboxyphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone




(4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(4-ethoxycarbonylphenyl)aminocarbonylmethyl]-2-pyrrolidone (18 mg; 0.037 mmol) was stirred in methanol (1 mL) and water (0.5 mL). A solution of sodium hydroxide (0.5M; 148 uL) was added and stirred at room temperature for 7 days. The solvent was removed and the residue was partitioned between ethyl acetate and 1M HCl (10 mL each). Evaporation of the organic solvent afforded 14 mg of crude product, which was reconstituted in 0.75 mL of 60% acetonitrile/water (containing 0.1 % formic acid) and purified on a reverse phase hplc column (C18- 5 um; 30x100 mm) using a 20-80% acetonitrile/water (0.1% formic acid) gradient over 6 minutes. A fraction at 3.4 minutes was collected and concentrated to afford 8.6 mg of (4S)-1-[N-(4-carboxyphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone as a white solid (51%). ¹H NMR (CDCl₃+trace DMSO-d₆; 300 MHz) 9.5 (s, 1H); 7.9 (d, 2H); 7.6 (d, 2H) 6.8 (s, 2H) 4.9 (m,1H), 3.9-4.2 (q, 2H) 3.7-3.9 (m, 5H); 3.7 (s, 3H); 3.5 (m, 2H); 2.8 (m,1H); 2.3-2.6 (m, 3H); 2.2 (m, 2H); MS [M+H]=455; [M+Na]=477.

(4S)-1-[N-(3-Carboxyphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone was made with a similar procedure using different starting materials.

Example 12

(4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-methylphenyl)sulfonylaminocarbonylmethyl]-2-pyrrolidone




(4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone-1-acetic acid (50 mg; 0.15 mmol) was stirred in DMF (3 mL) followed by addition of o-tolylsulfonamide (51 mg; 0.30 mmol), DMAP (22 mg; 0.18 mmol), and EDCI (58 mg; 0.30 mmol). After stirring for 90 minutes at room temperature the reaction was diluted with ethyl acetate (60 mL) and washed with 1M HCl, 30 mL of water and 30 mL of brine. The organic solvent was dried over sodium sulfate and evaporated to provide 92 mg of an amber oil. The compound was purified by hplc using a C18 column (5 μ m; 30x100 mm) and a gradient of 20-80% acetonitrile/water (0.1% formic acid) over 6 minutes. A fraction at 4 minutes was concentrated to afford 25 mg (35% yield) of (4S)-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-methylphenyl)sulfonylaminocarbonylmethyl]-2-pyrrolidone as a white solid. ¹H NMR (CDCl₃; 300 MHz) 9.6 (br s, 1H); 8.2 (d, 1H); 7.5 (t, 1H); 7.2-7.4 (m, 2H); 6.7-6.9 (m, 3H); 5.0 (br s, 1H); 3.8-4.1 (m, 9H); 5.5-3.7 (m, 2H); 2.9 (m, 2H); 2.5-2.7 (m, 4H); 2.2 (m, 2H). MS [M+H]=489.

(4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-(N-phenylsulfonylaminocarbonylmethyl)-2-pyrrolidone is made by a similar procedure using different starting materials.

Example 13

(4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(3-phenpropyl)]-2-pyrrolidone




A solution of 100 mg (0.36 mmol) of (4*S*)-4-(4-methoxy-3-(3*R*)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone in 2.6 mL of DMF was added to a flask containing 30 mg (0.75 mmol) of NaH (60% dispersion in mineral oil) and 1 mL of DMF. This was stirred at room temperature for 3 hours. Then, a solution of 0.16 mL (1.08 mmol) of 1-bromo-3-phenylpropane in 1 mL of DMF was added and the resulting mixture was stirred at room temperature overnight. This was then poured into a mixture of 15 mL of ethyl acetate and 15 mL of water and washed with 3 x 15 mL of water, 1 x 15 mL of brine, dried over Na₂SO₄, and purified by HPLC to give 101 mg (0.26 mmol, 71% yield) of (4*S*)-4-(4-methoxy-3-(3*R*)-tetrahydrofuranyloxyphenyl)-1-[N-3-phenylpropyl]-2-pyrrolidone as a clear oil. ¹H-NMR (CDCl₃) δ 7.32-7.27 (m, 2H), 7.22-7.18 (m, 3H), 6.84 (t, *J* = 8.3 Hz, 1H), 6.80 (dd, *J* = 1.9 Hz, *J* = 8.3 Hz, 1H), 6.69 (d, *J* = 1.9 Hz), 4.48 (m, 1H), 3.99-3.89 (m, 4H), 3.84 (s, 3H), 3.69 (dd, *J* = 8 Hz, *J* = 9 Hz, 1H), 3.43-3.28 (m, 4H), 2.78 (dd, *J* = 8.8 Hz, *J* = 16.8 Hz, 1H), 2.66 (m, 2H), 2.50 (dd, *J* = 8.0 Hz, *J* = 16.8 Hz, 1H), 2.20-2.13 (m, 2H), 1.88 (m, 2H).

The following compounds were made using the same method with different starting materials:

- A. (4*S*)-4-(4-Methoxy-3-(3*R*)-tetrahydrofuranyloxyphenyl)-1-[N-(2-phenoxyethyl)]-2-pyrrolidone
- B. (4*S*)-4-(4-methoxy-3-(3*R*)-tetrahydrofuranyloxyphenyl)-1-[N-(2-tert-butyltrimethylsilyloxyethyl)]-2-pyrrolidone

Example 14A

(4*S*)-4-(4-Methoxy-3-(3*R*)-tetrahydrofuranyloxyphenyl)-1-[N-(2-hydroxyethyl)]-2-pyrrolidone (Intermediate)




The crude (4*S*)-4-(4-methoxy-3-(3*R*)-tetrahydrofuranyloxyphenyl)-1-[N-(2-tert-
 5 butyldimethylsilyloxyethyl)-2-pyrrolidone] was dissolved in 12 mL of THF and 3.7 mL
 (3.7 mmol, 2.5 equivalents) of TBAF (1.0 M in THF) was added. The reaction was
 stirred for one hour at room temperature. It was then poured into 50 mL of water and
 extracted 2 x 50 mL of CH₂Cl₂ and dried over Na₂SO₄. The solvent was removed and
 residue purified by column chromatography eluting with CH₂Cl₂ to 10% MeOH:CH₂Cl₂
 10 over 20 minutes to yield 387 mg (78% yield) of (4*S*)-4-(4-methoxy-3-(3*R*)-
 tetrahydrofuranyloxyphenyl)-1-[N-(2-hydroxyethyl)]-2-pyrrolidone as a clear oil. ¹H-
 NMR (CDCl₃) δ 6.80 (s, 2H), 6.72 (s, 1H), 4.96-4.91 (m, 1H), 4.02-3.81 (m, 5H), 3.79
 (s, 3H), 3.53-3.41 (m, 4H), 2.98-2.92 (m, 2H), 2.78 (dd, J = 8.8 Hz, J = 16.8 Hz, 1H),
 2.58 (bs, 1H), 2.49 (dd, J = 8.1 Hz, J = 16.8 Hz, 1H), 2.16 (m, 2H).

15

Example 14B

(4*S*)-4-(4-Methoxy-3-(3*R*)-tetrahydrofuranyloxyphenyl)-1-[N-(2-phenthioethyl)]-2-pyrrolidone



20


Methanesulfonyl chloride, 40 μL (0.52 mmol), and 0.11 mL (0.65) of
 diisopropylethylamine were added to a solution of 83 mg (0.26 mmol) of (4*S*)-4-(4-
 methoxy-3-(3*R*)-tetrahydrofuranyloxyphenyl)-1-[N-(2-hydroxyethyl)]-2-pyrrolidone in 3

mL of CH₂Cl₂. This was stirred at room temperature for 1 hour. It was then washed with 10 mL of water and dried over Na₂SO₄, filtered and the solvent was removed. The crude (4*S*)-(4-methoxy-(3*R*)-tetrahydrofuranlylphenyl)-1-[N-(2-methanesulfonylethanol)]-2-pyrrolidone was dissolved in 3 mL of DMF along with 108 mg (0.78 mmol) of K₂CO₃ and 53 μL of thiophenol. The mixture was heated to 60 °C for 2 hours. It was then added to a mixture of 20 mL of saturated K₂CO₃ and 20 mL of ethyl acetate. The organic layer was washed with 2 x 20 mL of water and 1 x 20 mL of brine, dried over Na₂SO₄, filtered and the solvent was removed. Purification was accomplished by column chromatography using 50% ethyl acetate to 100 % ethyl acetate gradient over 20 minutes to yield 34.7 mg (32% yield) of (4*S*)-4-(4-methoxy-3-(3*R*)-tetrahydrofuranlyloxyphenyl)-1-[N-(2-phenthiolethyl)]-2-pyrrolidone as a clear oil. ¹H-NMR δ 7.41-7.2 (m, 5H), 6.73-6.86 (m, 3H), 4.95 (m, 1H), 3.87-4.05 (m, 4H), 3.84 (s, 3H), 3.76 (m, 1H), 3.58 (dt, J = 2 Hz, J = 6.8, 2H), 3.40 (m, 2H), 3.15 (t, J 6.6 Hz, 2H), 2.75 (dd, J = 8.6 Hz, J = 17 Hz, 1H), 2.47 (dd, J = 8.6, J = 17 Hz, 1H), 2.12 (m, 2H).

(4*S*)-4-(4-Methoxy-3-(3*R*)-tetrahydrofuranlyloxyphenyl)-1-[N-(2-(4-methoxyphenyl)oxyethyl)]-2-pyrrolidone was made using the same method with different starting materials.

20 **Example 14C**

(4*S*)-1-[2-(4-Isopropylphenylthio)ethyl]-4-[4-methoxy-3-(3*R*)-tetrahydrofuranlyloxyphenyl]-2-pyrrolidone



Into a flask containing 0.17 mmol crude (4*S*)-4-[4-methoxy-3-(3*R*)-tetrahydrofuranlyloxyphenyl]-(2-oxo-pyrrolidin-1-yl)ethyl methanesulfonate was added 2 mL of DMF, 111 mg (0.34 mmol) of Cs₂CO₃ and 52 μL (0.34 mmol) of 4-

isopropylthiophenol. The reaction mixture was heated to 60 °C overnight, cooled to room temperature and 20 mL of ethyl acetate was added. The organic layer was washed with 10 mL of saturated K₂CO₃, 10 mL of water and 10 mL of brine, dried over Na₂SO₄, filtered and concentrated. Purification by silica gel column chromatography using a
 5 gradient elution from 50% to 100% ethyl acetate in hexanes yielded 55 mg of (4S)-1-[2-(4-isopropylphenylthio)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]oxyphenyl]-2-pyrrolidone. ¹H-NMR δ (CDCl₃) 7.3 (d, 2H), 7.2 (d, 2H), 6.8 (m, 2H), 6.7 (s, 1H), 4.9 (m, 1H), 4.0-3.8 (M, 8H), 3.5 (m, 2H), 3.4 (m, 2H), 3.1 (t, 2H), 2.8 (m, 1H), 2.7 (dd, 1H), 2.4 (dd, 1H), 2.2 (m, 2H), 1.2 (d, 6H). M+1=456.0

10

The following compounds were made using the same method with different starting materials:

- (4S)-1-[2-(4-Isopropylphenoxy)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]oxyphenyl]-2-pyrrolidone
- 15 (4S)-1-[2-(4-Isopropylphenylthio)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]oxyphenyl]-2-pyrrolidone
- (4S)-1-[2-(3-Chlorophenylthio)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]oxyphenyl]-2-pyrrolidone
- (4S)-1-[2-(2,3-Difluorophenoxy)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]oxyphenyl]-2-pyrrolidone
- 20 (4S)-1-[2-(Benzothiazol-2-yl)oxyethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]oxyphenyl]-2-pyrrolidone
- (4S)-1-[2-(6-Fluorobenzothiazol-2-yl)thioethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]oxyphenyl]-2-pyrrolidone
- 25 (4S)-1-[N-(6-Fluorobenzothiazol-2-yl)aminoethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]oxyphenyl]-2-pyrrolidone
- (4S)-1-[N-(Benzothiazol-2-yl)aminoethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]oxyphenyl]-2-pyrrolidone
- (4S)-4-[4-methoxy-3-(3R)-tetrahydrofuran-2-yl]oxyphenyl]-1-[2-(4-trifluoromethylphenoxy)ethyl]-2-pyrrolidone
- 30

(4S)-1-[2-(2-Fluorophenylthio)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone


(4S)-1-[2-(3-Fluorophenylthio)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone

5 (4S)-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[2-(4-methoxyphenylthio)ethyl]-2-pyrrolidone

(S)-1-(2-Azido-ethyl)-4-{4-methoxy-3-[(R)-(tetrahydro-furan-3-yl)oxy]-phenyl}-pyrrolidin-2-one

10 **Example 15**

(4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-phenylsulfonyl)ethyl]-2-pyrrolidone



15

The crude (4S)-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-phenylthioethyl)]-2-pyrrolidone was dissolved in CH₂Cl₂ and 3 equivalent of m-CPBA were added. The reaction was stirred overnight at room temperature. It was then added to mixture of a saturated solution of NaHCO₃ and CH₂Cl₂. It was then extracted with 2 x 10 mL of CH₂Cl₂. The combined organics were then washed with 1 x 10 mL of water and 1 x 10 mL of brine. Purification was accomplished by column chromatography using a 50% ethyl acetate in hexanes to 100 % ethyl acetate gradient over 10 minutes to yield 82 mg (57% yield over three steps) of (4S)-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-phenylsulfonyl)ethyl]-2-pyrrolidone as a clear oil.

25


¹H-NMR δ 7.94 (d, J = 7.5 Hz, 2 H), 7.75-7.59 (m, 3H), 6.83 (m, 2H), 6.73 (s, 1H), 4.97 (m, 1H), 4.05-3.87, (m, 4H), 3.85 (s, 3H), 3.83-3.73 (m, 3H), 3.5-3.38 (m, 4H), 2.72 (dd, J = 9 Hz, J = 17 Hz, 1H), 2.48 (dd, J = 9 Hz, J = 17 Hz, 1H), 2.21-2.15 (m, 2H)

5

Example 16:

(4S)-2,3-Difluoro-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-N-[2-(2-oxopyrrolidin-1-yl)ethyl]-benzamide

10



Into a flask containing 100 mg (0.63 mmol) of 2,3-difluorobenzoic acid and 85 mg (0.63 mmol) of HOBt was added 6.5 mL of THF. This was cooled to 0 °C under argon and 0.10 mL (0.63 mmol) of DIC was added and stirring continued for 30 minutes. This solution was then added to a flask containing 55 mg (0.16 mmol) of (S)-1-(2-azido-ethyl)-4-{4-methoxy-3-[(R)-(tetrahydro-furan-3-yl)oxy]-phenyl}-pyrrolidin-2-one in 3 mL of THF at room temperature. Then, 0.12 mL of PBu₃ was added and the reaction was stirred at room temperature overnight. The solvent was removed under reduced pressure and the residue was dissolved in 10 mL of ethyl acetate. The organic layer was washed with 10 mL of 1N HCl, 10 mL of K₂CO₃, 10 mL of water, and 10 mL of brine, dried over Na₂SO₄, filtered and concentrated. Purification was accomplished by silica gel column chromatography using a gradient elution from 0% to 10% methanol in dichloromethane to yield 27 mg (4S)-2,3-Difluoro-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-N-[2-(2-oxopyrrolidin-1-yl)ethyl]-benzamide. ¹H-NMR δ (CDCl₃) 7.7 (t, 1H), 7.3 (m, 1H), 7.2 (m, 2H), 6.8 (s, 2H), 6.7 (s, 1H), 4.9 (m, 1H), 4.0 (m, 3H), 3.8 (m, 5H), 3.7 (m, 2H), 3.0 (m, 2H), 2.3 (m, 2H), 2.7 (dd, 1H), 2.5 (dd, 1H), 2.1 (m, 2H). M+1=461.0

15

20

25


The following compounds were made using the same method with different starting materials:

4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-N-[2-(2-oxopyrrolidin-1-yl)ethyl]benzamide

- 5 (4S)-2,3-Difluoro-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-N-[2-(2-oxopyrrolidin-1-yl)ethyl]-benzamide

Example 17

- 10 **(4S)-1-[N-(4-Carboxythiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone**



- (4S)-1-[N-(4-Ethoxycarbonylthiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone (95 mg; 0.19 mmol) was stirred in dioxane (5 mL) and water (1mL) and treated with 1M NaOH (0.46 mL; 0.46 mmol). The reaction was heated to 100 °C for 2 hours. LC-MS shows $[M+H]^+ = 462.2$ with a peak indicating amide cleavage ($[M+H]^+ = 336.1$). Purification was performed by reverse phase liquid chromatography on a C18 column (30x100 mm) using a gradient of 20-80% acetonitrile/water (containing 0.1% formic acid) in 6 minutes with a flow of 45 mL/min.
- 15 Ultraviolet detection at 277 nm was utilized and fractions containing the target were evaporated to afford the compound as a white solid (18 mg; 20%). ^1H NMR (CDCl_3 ; 300 MHz) δ 2.2 (m, 2H); 2.6 (m, 1H); 2.9 (m, 1H); 3.5 (m, 2H); 3.8 (s, 3H); 3.9-4.1 (m, 5H); 4.2-4.4 (q, 2H); 5.0 (m, 1H); 6.85 (s, 2H); 6.9 (s, 1H); 7.8 (s, 1H). ES-MS $[M+H]^+ = 462.2$

25


The following compounds were made using the same method with different starting materials:

(4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-N-[2-(2-oxopyrrolidin-1-yl)ethyl]-4-methoxybenzamide

5

Example 18

(4S)-1-[N-(4-Carboxy-3-fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone



- 10 A solution of (4S)-1-[N-(4-tert-butyloxycarbonyl-3-fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone (55mg; 0.104 mmol) in dichloromethane (1 mL) was treated with trifluoroacetic acid (1 mL) and the solution stirred at room temperature for 45 minutes. The solvent was evaporated under reduced pressure and the residue was taken up in ethyl acetate (50 mL) and washed with water
- 15 twice (10mL each) and brine (10 mL). The organic layer was dried over sodium sulfate and evaporated to provide a tan solid. The product was purified by reverse phase liquid chromatography using a C18 column (30x100 mm) and a gradient of 20-80% acetonitrile/water (plus 0.1% formic acid) over six minutes at a flow rate of 45 mL/min. Ultraviolet detection at 269 nm was utilized and fractions containing the target compound
- 20 were pooled and evaporated to afford (4S)-1-[N-(4-Carboxy-3-fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone as a white foam (3.6 mg; 7%). ES-MS [M+H]⁺= 473.1

Example 19

In Vitro Measurement of Type 4 Phosphodiesterase

Enzyme Preparation:

Human PDE4 was obtained from baculovirus-infected Sf9 cells that expressed the recombinant enzyme. The cDNA encoding hPDE-4D6 was subcloned into a baculovirus vector. Insect cells (Sf9) were infected with the baculovirus and cells were cultured until protein was expressed. The baculovirus-infected cells were lysed and the lysate was used as source of hPDE-4D6 enzyme. The enzyme was partially purified using a DEAE ion exchange chromatography. This procedure can be repeated using cDNA encoding other PDE-4 enzymes.

Assay:

Type 4 phosphodiesterases convert cyclic adenosine monophosphate (cAMP) to 5'-adenosine monophosphate (5'-AMP). Nucleotidase converts 5'-AMP to adenosine. Therefore the combined activity of PDE4 and nucleotidase converts cAMP to adenosine. Adenosine is readily separated from cAMP by neutral alumina columns. Phosphodiesterase inhibitors block the conversion of cAMP to adenosine in this assay; consequently, PDE4 inhibitors cause a decrease in adenosine.

Cell lysates (40 ul) expressing hPDE-4D6 were combined with 50 ul of assay mix and 10 ul of inhibitors and incubated for 12 min at room temperature. Final concentrations of assay components were: 0.4 ug enzyme, 10mM Tris-HCl (pH 7.5), 10mM MgCl₂, 3 uM cAMP, 0.002 U 5'-nucleotidase, and 3 x 10⁴ cpm of [3H]cAMP. The reaction was stopped by adding 100 µl of boiling 5mN HCl. An aliquot of 75 µl of reaction mixture was transferred from each well to alumina columns (Multiplate; Millipore). Labeled adenosine was eluted into an OptiPlate by spinning at 2000 rpm for 2 min; 150 µl per well of scintillation fluid was added to the OptiPlate. The plate was sealed, shaken for about 30 min, and cpm of [3H]adenosine was determined using a Packard Topcount 96 counter.

All test compounds are dissolved in 100% DMSO and diluted into the assay such that the final concentration of DMSO is 0.1%. DMSO does not affect enzyme activity at this concentration.

5 **Example 20**

Passive Avoidance in Rats, an in vivo Test for Learning and Memory

The test was performed as previously described (Zhang, H.-T., Crissman, A.M., Dorairaj, N.R., Chandler, L.J., and O'Donnell, J.M., *Neuropsychopharmacology*, 2000, 10 23, 198-204.). The apparatus (Model E10-16SC, Coulbourn Instruments, Allentown, PA) consisted of a two-compartment chamber with an illuminated compartment connected to a darkened compartment by a guillotine door. The floor of the darkened compartment consisted of stainless steel rods through which an electric foot-shock could be delivered from a constant current source. All experimental groups were first habituated to the 15 apparatus the day before the start of the experiment. During the training, the rat (Male Sprague-Dawley (Harlan) weighing 250 to 350 g) was placed in the illuminated compartment facing away from the closed guillotine door for 1 minute before the door was raised. The latency for entering the darkened compartment was recorded. After the rat entered the darkened compartment, the door was closed and a 0.5 mA electric shock 20 was administered for 3 seconds. Twenty-four hours later, the rat was administered 0.1 mg/kg of the test compound or saline, 30 minutes prior to the injection of saline or test compound (dosed from 0.1 to 2.5 mg/kg, i.p.), which was 30 minutes before the retention test started. The rat was again placed in the illuminated compartment with the guillotine door open. The latency for entering the darkened compartment was recorded for up to 25 180 seconds, at which time the trial was terminated.

All data were analyzed by analyses of variance (ANOVA); individual comparisons were made using Kewman-Keuls tests. Naïve rats required less than 30 seconds, on average, to cross from the illuminated compartment to the darkened compartment.

30

Example 21

Radial arm maze task in Rats, an in vivo Test for Learning and Memory

The test was performed as previously described (Zhang, H.-T., Crissman, A.M., Dorairaj, N.R., Chandler, L.J., and O'Donnell, J.M., *Neuropsychopharmacology*, 2000, 23, 198-204.). Five days after initial housing, rats (male Sprague-Dawley (Harlan) weighing 250 to 350 g) were placed in the eight-arm radial maze (each arm was 60x10x12 cm high; the maze was elevated 70 cm above the floor) for acclimation for two days. Rats were then placed individually in the center of the maze for 5 minutes with food pellets placed close to the food wells, and then, the next day, in the wells at the end of the arms; 2 sessions a day were conducted. Next, four randomly selected arms were then baited with one pellet of food each. The rat was restricted to the center platform (26 cm in diameter) for 15 seconds and then allowed to move freely throughout the maze until it collected all pellets of food or 10 minutes passed, whichever came first. Four parameters were recorded: 1) working memory errors, i.e., entries into baited arms that had already been visited during the same trial; 2) reference memory errors, i.e., entries into unbaited arms; 3) total arm entries; and 4) the test duration (seconds), i.e., the time spent in the collection of all the pellets in the maze. If the working memory error was zero and the average reference memory error was less than one in five successive trials, the rats began the drug tests. The test compound or saline was injected 15 minutes prior to vehicle or test agent, which was given 45 minutes before the test. Experiments were performed in a lighted room, which contained several extra-maze visual cues.

All data were analyzed by analyses of variance (ANOVA); individual comparisons were made using Kewman-Keuls tests.

A decrease in adenosine concentration is indicative of inhibition of PDE activity. This procedure was used to screen compounds of the present invention for their ability to inhibit PDE4. pIC₅₀ values were determined by screening 6 to 12 concentrations of compound ranging from 0.1 nM to 10,000 nM and then plotting drug concentration versus ³H-adenosine concentration. Prism® was used to estimate pIC₅₀ values.


The pIC₅₀ values of preferred compounds according to the invention are less than 100 nM, especially less than 10 nM.

5 The preceding examples can be repeated with similar success by substituting the generically or specifically described reactants and/or operating conditions of this invention for those used in the preceding examples.

10 While the invention has been illustrated with respect to the production and of particular compounds, it is apparent that variations and modifications of the invention can be made without departing from the spirit or scope of the invention.

WE CLAIM:

1. A compound of Formula I:



15 wherein

R¹ is alkyl having 1 to 8 carbon atoms wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,

20 alkyl having 1 to 8 carbon atoms which is substituted one or more times by halogen, oxo or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,

25 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl having 1 to 4 carbon atoms or combinations thereof,

30 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, or combinations thereof,

35 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino,

- hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,
- 5 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,
- 10 a partially unsaturated carbocyclic group having 5 to 14 carbon atoms which is unsubstituted or substituted one or more times by halogen, alkyl, alkoxy, nitro, cyano, oxo, or combinations thereof ,
- 15 arylalkenyl having 8 to 16 carbon atoms, wherein the alkenyl portion has up to 5 carbon atoms, which is unsubstituted or substituted one or more times by halogen, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof;
- 20 a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof, or
- 25 30

cycloalkylalkyl having 4 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,

5 R^2 is alkyl having 1 to 4 carbon atoms, which is unsubstituted or substituted one or more times by halogen;

R^3 is $-C(O)R^4$, $-(CH_2)_n C(O)R^4$, $-(CH_2)_n OR^5$, $-(CH_2)_n SR^5$, $-(CH_2)_n SO_2R^4$, $-(CH_2)_n NR^5R^6$, $-CH_2CO_2R^5$, $-CH_2CONR^6R^5$, $-CH_2CONHR^5$, $-(CH_2)_n NR^6SO_2R^4$, $-(CH_2)_n NR^6COR^4$, or $-CH_2CONHSO_2R^4$;

10

R^4 is alkyl having 1 to 12 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,

15

 alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,

20

 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,

25

 cycloalkylalkyl having 4 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,

30 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro,

- methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, alkylsulphonamido, arylsulphonamido, halogenated arylsulphonamido, phenoxy, acylamido, and acyloxy, or combinations thereof,
- arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, aminosulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,
- a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom, which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, cycloalkyl, cycloalkylalkyl, alkoxy, alkoxyalkyl, cyano, halogenated alkyl, halogenated alkoxy, nitro, oxo, amino, alkylamino, dialkylamino, aminosulphonyl, heterocycle, heterocyclic-alkyl, or combinations thereof, or
- a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof; and
- R⁵ is H,

- alkyl having 1 to 12 carbon atoms wherein optionally one or more $\text{-CH}_2\text{CH}_2\text{-}$ groups are replaced in each case by -CH=CH- or $\text{-C}\equiv\text{C-}$ groups,
- 5
- alkyl having 1 to 12 carbon atoms which is substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $\text{-CH}_2\text{CH}_2\text{-}$ groups are replaced in each case by -CH=CH- or $\text{-C}\equiv\text{C-}$ groups,
- 10
- alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $\text{-CH}_2\text{CH}_2\text{-}$ groups are replaced in each case by -CH=CH- or $\text{-C}\equiv\text{C-}$ groups,
- 15
- cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- 20
- cycloalkylalkyl having 4 to 16 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 25
- aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, alkylsulphonamido, arylsulphonamido, halogenated arylsulphonamido, phenoxy, acylamido,
- 30
- and acyloxy, or combinations thereof,

- 5 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, aminosulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,
- 10 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom, which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, cycloalkyl, cycloalkylalkyl, alkoxy, alkoxyalkyl, cyano, halogenated alkyl, halogenated alkoxy, nitro, oxo, amino, alkylamino, dialkylamino, aminosulphonyl, heterocycle, heterocyclic-
- 15 alkyl, or combinations thereof, or
- 20 a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof;
- 25 R⁶ is H,
- 30 alkyl having 1 to 12 carbon atoms wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,

- alkyl having 1 to 12 carbon atoms which is substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $\text{-CH}_2\text{CH}_2\text{-}$ groups are replaced in each case by -CH=CH- or $\text{-C}\equiv\text{C-}$ groups,
- 5 alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $\text{-CH}_2\text{CH}_2\text{-}$ groups are replaced in each case by -CH=CH- or $\text{-C}\equiv\text{C-}$ groups,
- 10 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- 15 cycloalkylalkyl having 4 to 16 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 20 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,
- 25 arylalkyl having 8 to 16 carbon atoms which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof;
- 30

5 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom, which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, alkoxy, alkoxycarbonyl, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, or combinations thereof, or

10 a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof;

15

n is 0 to 4; and

20 pharmaceutically acceptable salts thereof.


20

with the proviso that:

25 at least one R⁴ or R⁵ is an alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups.

2. A compound of Formula I:

5



wherein

15

R^1 is alkyl having 1 to 8 carbon atoms wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,

20

alkyl having 1 to 8 carbon atoms which is substituted one or more times by halogen, oxo or combinations thereof wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,

25

cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl having 1 to 4 carbon atoms or combinations thereof,

30

a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, or combinations thereof,

35

aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl,

alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,

5 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,

10

a partially unsaturated carbocyclic group having 5 to 14 carbon atoms which is unsubstituted or substituted one or more times by halogen, alkyl, alkoxy, nitro, cyano, oxo, or combinations thereof ,

15

arylalkenyl having 8 to 16 carbon atoms, wherein the alkenyl portion has up to 5 carbon atoms, which is unsubstituted or substituted one or more times by halogen, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof;

20

a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof, or

25

30

cycloalkylalkyl having 4 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,


5 R^2 is alkyl having 1 to 4 carbon atoms, which is unsubstituted or substituted one or more times by halogen;

10 R^3 is arylalkyl having 7 to 16 carbon atoms which is substituted in the aryl portion by phenyl and optionally further substituted one or more times by halogen, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, and acyloxy, or combinations thereof; and

15 n is 0 to 4; and

pharmaceutically acceptable salts thereof.

20 3. A compound of Formula I:



wherein

35 R^1 is alkyl having 1 to 8 carbon atoms wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,

- alkyl having 1 to 8 carbon atoms which is substituted one or more times by halogen, oxo or combinations thereof wherein optionally one or more $\text{-CH}_2\text{CH}_2\text{-}$ groups are replaced in each case by -CH=CH- or $\text{-C}\equiv\text{C-}$ groups,
- 5 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl having 1 to 4 carbon atoms or combinations thereof,
- 10 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, or combinations thereof,
- 15 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and
- 20 acyloxy, or combinations thereof,
- 25 arylalkyl having 8 to 16 carbon atoms which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acylox, or combinations thereof,

- a partially unsaturated carbocyclic group having 5 to 14 carbon atoms which is unsubstituted or substituted one or more times by halogen, alkyl, alkoxy, nitro, cyano, oxo, or combinations thereof ,
- 5 arylalkenyl having 8 to 16 carbon atoms, wherein the alkenyl portion has up to 5 carbon atoms, which is unsubstituted or substituted one or more times by halogen, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, 10 alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof;
- a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an 15 N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof, or
- 20 cycloalkylalkyl having 4 to 16 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 25 R^2 is alkyl having 1 to 4 carbon atoms, which is unsubstituted or substituted one or more times by halogen;
- R^3 is

-C(O)R⁴, -(CH₂)_nC(O)R⁴, -(CH₂)_nOR⁵, -(CH₂)_nSR⁵, -(CH₂)_nSO₂R⁴, -
 (CH₂)_nNR⁵R⁶, -CH₂CO₂R⁵, -CH₂CONR⁶R⁵, -CH₂CONHR⁵, -
 (CH₂)_nNR⁶SO₂R⁴, -(CH₂)_nNR⁶COR⁴, or -CH₂CONHSO₂R⁴;

- 5 R⁴ is alkyl having 1 to 12 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,
- 10 alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,
- 15 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- 20 cycloalkylalkyl having 4 to 16 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 25 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, alkylsulphonamido, arylsulphonamido, halogenated arylsulphonamido, phenoxy, acylamido, and acyloxy, or combinations thereof,
- 30

- 5 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxy carbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, aminosulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,
- 10 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom, which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, cycloalkyl, cycloalkylalkyl, alkoxy, alkoxyalkyl, cyano, halogenated alkyl, halogenated alkoxy, nitro, oxo, amino, alkylamino, dialkylamino, aminosulphonyl, heterocycle, heterocyclic-
- 15 alkyl, or combinations thereof, or
- 20 a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof; and
- 25 R⁵ is H,
- 30 alkyl having 1 to 12 carbon atoms wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,

- alkyl having 1 to 12 carbon atoms which is substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-\text{CH}_2\text{CH}_2-$ groups are replaced in each case by $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$ groups,
- 5 alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-\text{CH}_2\text{CH}_2-$ groups are replaced in each case by $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$ groups,
- 10 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- 15 cycloalkylalkyl having 4 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 20 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, alkylsulphonamido, arylsulphonamido, halogenated arylsulphonamido, phenoxy, acylamido, and acyloxy, or combinations thereof,
- 25 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl,
- 30

aminosulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,

a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom, which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, cycloalkyl, cycloalkylalkyl, alkoxy, alkoxyalkyl, cyano, halogenated alkyl, halogenated alkoxy, nitro, oxo, amino, alkylamino, dialkylamino, aminosulphonyl, heterocycle, heterocyclic-alkyl, or combinations thereof, or

a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof;

R^6 is H,

alkyl having 1 to 12 carbon atoms wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,

alkyl having 1 to 12 carbon atoms which is substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,

- alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $\text{-CH}_2\text{CH}_2\text{-}$ groups are replaced in each case by -CH=CH- or $\text{-C}\equiv\text{C-}$ groups,
- 5
- cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- 10
- cycloalkylalkyl having 4 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 15
- aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,
- 20
- arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof;
- 25
- 30
- a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom (e.g., 3-thienyl, 2-thienyl, 3-tetrahydrofuran), which is unsubstituted or substituted one or more times by halogen, aryl, alkyl,

alkoxy, alkoxycarbonyl, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, or combinations thereof, or

a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof;


n is 0 to 4; and

pharmaceutically acceptable salts thereof

with the proviso that:

at least one of R^4 and R^5 in formula I is aryl substituted by alkylsulphonamido, arylsulphonamido, or halogenated arylsulphonamido, arylalkyl which is substituted by aminosulphonyl, or a heterocyclic group which is substituted one or more times by alkoxyalkyl, cycloalkyl, cycloalkylalkyl, halogenated alkoxy, aminosulphonyl, heterocycle, heterocyclic-alkyl, or combinations thereof.

4. A compound of Formula I:



wherein

5 R¹ is alkyl having 1 to 8 carbon atoms wherein optionally one or more
 -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C-
 groups,

10 alkyl having 1 to 8 carbon atoms which is substituted one or more times
 by halogen, oxo or combinations thereof wherein optionally one or more
 -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C-
 groups,

15 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or
 substituted one or more times by halogen, oxo, alkyl having 1 to 4 carbon
 atoms or combinations thereof,

20 a heterocyclic group, which is saturated, partially saturated or fully
 unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an
 N, O or S atom which is unsubstituted or substituted one or more times by
 halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino,
 alkylamino, dialkylamino, or combinations thereof,

25 aryl having 6 to 14 carbon atoms which is unsubstituted or substituted one
 or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro,
 methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino,
 hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl,
 alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and
 acyloxy, or combinations thereof,

30 arylalkyl having 8 to 16 carbon atoms which is unsubstituted or
 substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy,
 alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino,

- dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxy carbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,
- 5 a partially unsaturated carbocyclic group having 5 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, alkyl, alkoxy, nitro, cyano, oxo, or combinations thereof ,
- 10 arylalkenyl having 8 to 16 carbon atoms, wherein the alkenyl portion has up to 5 carbon atoms, which is unsubstituted or substituted one or more times by halogen, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxy carbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or
- 15 combinations thereof;
- a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in
- 20 the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof, or
- 25 cycloalkylalkyl having 4 to 16 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 30 R^2 is alkyl having 1 to 4 carbon atoms, which is unsubstituted or substituted one or more times by halogen;

R³ is H,

alkyl having 1 to 8 carbon atoms wherein optionally one or more
-CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C-
groups,

alkyl having 1 to 8 carbon atoms which is substituted one or more times
by halogen, oxo, or combinations thereof wherein optionally one or more
-CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C-
groups,

cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or
substituted one or more times by halogen, oxo, alkyl, or combinations
thereof,

arylalkyl having 7 to 16 carbon atoms, which is unsubstituted or
substituted one or more times by halogen, alkyl, hydroxy, alkoxy, nitro,
methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino,
hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl,
alkylthio, alkylsulphinyl, alkylsulphonyl, phenyl, phenoxy, and acyloxy,
or combinations thereof,

heterocyclic-alkyl group, which is saturated, partially saturated or fully
unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an
N, O or S atom, which is unsubstituted or substituted one or more times in
the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano,
halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy
or combinations thereof and/or substituted in the alkyl portion by halogen,
oxo, cyano, or combinations thereof,

alkoxyalkyl having 3 to 8 carbon atoms,

-C(O)R⁴, -(CH₂)_nC(O)R⁴, -(CH₂)_nOR⁵, -(CH₂)_nSR⁵, -(CH₂)_nSO₂R⁴, -
 (CH₂)_nNR⁵R⁶, -CH₂CO₂R⁵, -CH₂CONR⁶R⁵, -CH₂CONHR⁵, -
 (CH₂)_nNR⁶SO₂R⁴, -(CH₂)_nNR⁶COR⁴, or -CH₂CONHSO₂R⁴;

5

R⁴ is alkyl having 1 to 12 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,

10

alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,

15

cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,

20

cycloalkylalkyl having 4 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,

25

aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, alkylsulphonamido, arylsulphonamido, halogenated arylsulphonamido, phenoxy, acylamido, and acyloxy, or combinations thereof,

30

- 5 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxy-carbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, aminosulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,
- 10 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom, which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, cycloalkyl, cycloalkylalkyl, alkoxy, alkoxyalkyl, cyano, halogenated alkyl, halogenated alkoxy, nitro, oxo, amino, alkylamino, dialkylamino, aminosulphonyl, heterocycle, heterocyclic-
- 15 alkyl, or combinations thereof, or
- 20 a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof; and
- 25 R⁵ is H,
- 30 alkyl having 1 to 12 carbon atoms wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,

- alkyl having 1 to 12 carbon atoms which is substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-\text{CH}_2\text{CH}_2-$ groups are replaced in each case by $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$ groups,
- 5 alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-\text{CH}_2\text{CH}_2-$ groups are replaced in each case by $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$ groups,
- 10 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- 15 cycloalkylalkyl having 4 to 16 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 20 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, alkylsulphonamido, arylsulphonamido, halogenated arylsulphonamido, phenoxy, acylamido, and acyloxy, or combinations thereof,
- 25 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl,
- 30

aminosulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,

5 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom, which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, cycloalkyl, cycloalkylalkyl, alkoxy, alkoxyalkyl, cyano, halogenated alkyl, halogenated alkoxy, nitro, oxo, amino, alkylamino, dialkylamino, aminosulphonyl, heterocycle, heterocyclic-
10 alkyl, or combinations thereof, or

a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in
15 the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof;

20 R^6 is H,

alkyl having 1 to 12 carbon atoms wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,
25

alkyl having 1 to 12 carbon atoms which is substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,
30

- alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-\text{CH}_2\text{CH}_2-$ groups are replaced in each case by $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$ groups,
- 5
- cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- 10
- cycloalkylalkyl having 4 to 16 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 15
- aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,
- 20
- arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof;
- 25
- a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom, which is unsubstituted or substituted one or more times by
- 30

halogen, aryl, alkyl, alkoxy, alkoxycarbonyl, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, or combinations thereof, or

a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof;

n is 0 or 1; and

pharmaceutically acceptable salts thereof;

with the proviso that:

(a) when R^2 is CH_3 and R^3 is H, then R^1 is not methyl, ethyl, n-propyl, isopropyl, sec-butyl, n-butyl, isobutyl, neopentyl, n-pentyl, 2-methylbutyl, isopentyl, n-hexyl, phenyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopentenyl, methylcyclopentyl, cyclopropylmethyl, cyclopentylmethyl, 2-propenyl, 2-propynyl, 3-methyl-2-butenyl, N-substituted 2-piperazinylethyl, norbornyl, 3-tetrahydrofuryl, 2-tetrahydrofuryl, 3-tetrahydrothienyl, 2-oxacyclopropyl, 2-oxacyclopentyl, 3-oxacyclopentyl, 2-chloroethyl, 2-bromoethyl, 2,2,2-trifluoroethyl, 3-bromopropyl, 3-chloropropyl, or 4-bromobutyl;

(b) when R^1 is cyclopentyl, and R^2 is methyl, then R^3 is not H, acetyl, benzyl, 4-hydroxybenzyl, 4-acetoxybenzyl, 4-bromobenzyl, 3,4-dimethoxybenzyl, 4-methylthiobenzyl, 4-


5 cyanobenzyl, 2-aminobenzyl, 3-aminobenzyl, 4-aminobenzyl, 4-dimethylaminobenzyl, 2,4-diaminobenzyl, 4-amino-3,5-dimethoxybenzyl, 3-carboxybenzyl, 3-methoxycarbonylbenzyl, 4-methoxycarbonylbenzyl, 4-methylsulfinylbenzyl, 4-methylsulfonylbenzyl, 2-nitrobenzyl, 3-nitrobenzyl, 4-nitrobenzyl, 2,4-dinitrobenzyl, 2-nitro-4-aminobenzyl, 2-amino-4-nitrobenzyl, morpholinoethyl, 2-pyridylmethyl, 3-pyridylmethyl, 4-pyridylmethyl, 4-(6-fluoroquinolyl)methyl, 2-(7-chloroquinolyl)methyl, 2-imidazolylmethyl, or substituted imidazolylmethyl;

15 (c) when R^1 is CH_3 , and R^3 is H, then R^2 is not methyl, ethyl, or butyl;

(d) when R^3 is H, then R^1 and R^2 are not both ethyl or isobutyl; and

20 (e) when R^1 and R^2 are both difluoromethyl, then R^3 is not 4-aminobenzyl, or 4-amino-3,5-dimethoxybenzyl.

5. A compound of Formula II:



35 wherein

- R¹ is alkyl having 1 to 8 carbon atoms wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,
- 5 alkyl having 1 to 8 carbon atoms which is substituted one or more times by halogen, oxo or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,
- 10 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl having 1 to 4 carbon atoms or combinations thereof,
- 15 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom, which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, or combinations thereof,
- 20 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and
- 25 acyloxy, or combinations thereof,
- 30 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl,

	alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,
5	a partially unsaturated carbocyclic group having 5 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, alkyl, alkoxy, nitro, cyano, oxo, or combinations thereof ,
10	arylalkenyl having 8 to 16 carbon atoms, wherein the alkenyl portion has up to 5 carbon atoms, which is unsubstituted or substituted one or more times by halogen, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof;
15	a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano,
20	halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof, or
25	cycloalkylalkyl having 4 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
30	R ² is alkyl having 1 to 4 carbon atoms, which is unsubstituted or substituted one or more times by halogen;

- R^3 is $-(CH_2)_nOR^5$, $-(CH_2)_nSR^5$, $-(CH_2)_nSO_2R^4$, $-(CH_2)_nNR^5R^6$, $-CH_2CO_2R^5$, $-CH_2CONR^6R^5$, $-(CH_2)_nNR^6SO_2R^4$, $-(CH_2)_nNR^6COR^4$, or $-CH_2CONHSO_2R^4$;
- 5 R^4 is alkyl having 1 to 12 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,
- 10 alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $-CH_2CH_2-$ groups are replaced in each case by $-CH=CH-$ or $-C\equiv C-$ groups,
- 15 cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- 20 cycloalkylalkyl having 4 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 25 aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, alkylsulphonamido, arylsulphonamido, halogenated arylsulphonamido, phenoxy, acylamido, and acyloxy, or combinations thereof,
- 30

- 5 arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF₃, OCF₃, alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxy carbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, aminosulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,
- 10 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom, which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, cycloalkyl, cycloalkylalkyl, alkoxy, alkoxyalkyl, cyano, halogenated alkyl, halogenated alkoxy, nitro, oxo, amino, alkylamino, dialkylamino, aminosulphonyl, heterocycle, heterocyclic-
- 15 alkyl, or combinations thereof, or
- 20 a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof; and
- 25 R⁵ is H,
- 30 alkyl having 1 to 12 carbon atoms wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,

- alkyl having 1 to 12 carbon atoms which is substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $\text{-CH}_2\text{CH}_2\text{-}$ groups are replaced in each case by -CH=CH- or $\text{-C}\equiv\text{C-}$ groups,
- 5
- alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $\text{-CH}_2\text{CH}_2\text{-}$ groups are replaced in each case by -CH=CH- or $\text{-C}\equiv\text{C-}$ groups,
- 10
- cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- 15
- cycloalkylalkyl having 4 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- 20
- aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, alkylsulphonamido, arylsulphonamido, halogenated arylsulphonamido, phenoxy, acylamido, and acyloxy, or combinations thereof,
- 25
- arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl,
- 30

aminosulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,

5 a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom, which is unsubstituted or substituted one or more times by halogen, aryl, alkyl, cycloalkyl, cycloalkylalkyl, alkoxy, alkoxyalkyl, cyano, halogenated alkyl, halogenated alkoxy, nitro, oxo, amino, alkylamino, dialkylamino, aminosulphonyl, heterocycle, heterocyclic-
10 alkyl, or combinations thereof, or

a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in
15 the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof;

20 R^6 is H,

alkyl having 1 to 12 carbon atoms wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,

25 alkyl having 1 to 12 carbon atoms which is substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more -CH₂CH₂- groups are replaced in each case by -CH=CH- or -C≡C- groups,

30

- alkoxyalkyl having 3 to 8 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, or combinations thereof wherein optionally one or more $\text{-CH}_2\text{CH}_2\text{-}$ groups are replaced in each case by -CH=CH- or $\text{-C}\equiv\text{C-}$ groups,
- 5
- cycloalkyl having 3 to 8 carbon atoms, which is unsubstituted or substituted one or more times by halogen, oxo, alkyl, or combinations thereof,
- 10
- cycloalkylalkyl having 4 to 16 carbon atoms which is unsubstituted or substituted one or more times by halogen, oxo, alkyl or combinations thereof,
- aryl having 6 to 14 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro,
- 15
- methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy, acylamido, and acyloxy, or combinations thereof,
- 20
- arylalkyl having 8 to 16 carbon atoms, which is unsubstituted or substituted one or more times by halogen, CF_3 , OCF_3 , alkyl, hydroxy, alkoxy, nitro, methylenedioxy, ethylenedioxy, amino, alkylamino, dialkylamino, hydroxyalkyl, hydroxyalkoxy, carboxy, cyano, acyl, alkoxycarbonyl, alkylthio, alkylsulphinyl, alkylsulphonyl, phenoxy,
- 25
- acylamido, and acyloxy, or combinations thereof;
- a heterocyclic group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is a N, O or S atom, which is unsubstituted or substituted one or more times by
- 30
- halogen, aryl, alkyl, alkoxy, alkoxycarbonyl, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, or combinations thereof, or

a heterocyclic-alkyl group, which is saturated, partially saturated or fully unsaturated, having 5 to 10 ring atoms in which at least 1 ring atom is an N, O or S atom, which is unsubstituted or substituted one or more times in the heterocyclic portion by halogen, aryl, alkyl, alkoxy, cyano, halogenated alkyl, nitro, oxo, amino, alkylamino, dialkylamino, carboxy or combinations thereof and/or substituted in the alkyl portion by halogen, oxo, cyano, or combinations thereof,

wherein if R^3 is $-\text{CH}_2\text{CONR}^6\text{R}^5$, R^6 is other than H;

n is 0 to 4; and

pharmaceutically acceptable salts thereof.

6. A compound selected from:

4-[4-Methoxy-3-(4-methoxyphenoxy)phenyl]-2-pyrrolidone,
 4-[4-Methoxy-3-(3-thienyloxy)phenyl]-2-pyrrolidone,
 4-[3-(4-Fluorophenoxy)-4-methoxyphenyl]-2-pyrrolidone,
 4-(3-(3-Cyclohexyl-1-propyloxy)-4-methoxyphenyl)-2-pyrrolidone,
 4-(4-Methoxy-3-(2-phenylethoxy)phenyl)-2-pyrrolidone,
 4-(4-Methoxy-3-(3-phenyl-1-propoxy)phenyl)-2-pyrrolidone,
 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-chloro-4-fluorobenzyl)-2-pyrrolidone,
 4-(3-cyclopentyloxy-4-methoxyphenyl)-1-methoxycarbonylmethyl-2-pyrrolidone,
 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-cyanomethyl-2-pyrrolidone,
 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-cyclopentyl-2-pyrrolidone,
 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-methoxybenzoyl)-2-pyrrolidone,
 4(S)-(3-Cyclopentyloxy-4-difluoromethoxyphenyl)-1-(N-(2-(6-methylpyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
 1-(N-(2,3-Difluorophenyl)-aminocarbonylmethyl)-4(S)-(4-methoxy-3-(3(R)-tetra

hydrofuranyloxy)phenyl)-2-pyrrolidone,

4(S)-(4-Methoxy-3-(R)-tetrahydrofuranyloxyphenyl)-1-(N-(2-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,

4(S)-(4-Methoxy-3-(R)-tetrahydrofuranyloxyphenyl)-1-(N-(2-(6-methylpyridinyl))-aminocarbonylmethyl)-2-pyrrolidone, and

physiologically acceptable salts thereof, wherein in each case the compound can be in the form of a mixture of enantiomers such as the racemate or a mixture of diastereomers, or can be in the form of a single enantiomer or a single diastereomer.

7. A compound selected from:

(4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-methylbenzyl)-2-pyrrolidone,

(4S)-1-(2-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,

(4S)-1-(4-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,

(4S)-1-(3-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,

(4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-methoxybenzyl)-2-pyrrolidone,

(4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-fluorobenzyl)-2-pyrrolidone,

(4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-fluorobenzyl)-2-pyrrolidone,

(4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-fluorobenzyl)-2-pyrrolidone,

(4S)-1-(4-Cyanobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,

(4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-trifluoromethylbenzyl)-2-pyrrolidone,

(4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-trifluoromethylbenzyl)-2-pyrrolidone,

(4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-trifluoromethylbenzyl)-2-pyrrolidone,

(4S)-1-(3,5-bis(trifluoromethyl)benzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,

(4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,4-difluorobenzyl)-2-pyrrolidone,

(4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,5-difluorobenzyl)-2-pyrrolidone,

(4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,4-difluorobenzyl)-2-pyrrolidone,

(4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,6-difluorobenzyl)-2-pyrrolidone,

(4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,3-difluorobenzyl)-2-pyrrolidone,

(4S)-1-(2-Chloro-4-fluorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,

(4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,4-dichlorobenzyl)-2-pyrrolidone,

- (4*S*)-1-(4-*tert*-Butylbenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-ethyl-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-propyl-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-butyl-2-pyrrolidone,
 5 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-methoxyethyl)-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-phenylbenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-methylbenzyl)-2-pyrrolidone,
 (4*R*)-1-(2-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*R*)-1-(4-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 10 (4*R*)-1-(3-Chlorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-methoxybenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-fluorobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-fluorobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-fluorobenzyl)-2-pyrrolidone,
 15 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-trifluoromethylbenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3-trifluoromethylbenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(4-trifluoromethylbenzyl)-2-pyrrolidone,
 (4*R*)-1-(3,5-Bis(trifluoromethyl)benzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-
 pyrrolidone,
 20 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,4-difluorobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,5-difluorobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,4-difluorobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,6-difluorobenzyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2,3-difluorobenzyl)-2-pyrrolidone,
 25 (4*R*)-1-(2-Chloro-4-fluorobenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(3,4-dichlorobenzyl)-2-pyrrolidone,
 (4*R*)-1-(4-*tert*-Butylbenzyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-ethyl-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-propyl-2-pyrrolidone,
 30 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-butyl-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-methoxyethyl)-2-pyrrolidone,

- (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(2-phenylbenzyl)-2-pyrrolidone,
 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2,6-dimethylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2,6-dimethylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
 5 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
 (4*R*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
 10 (4*S*)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(2,3-difluorophenyl)-aminocarbonylmethyl)-2-pyrrolidone,
 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(3-chlorophenyl)-aminocarbonylmethyl)-2-pyrrolidone,
 15 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(4-pyridyl)-aminocarbonylmethyl)-2-pyrrolidone,
 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(4-methoxyphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
 20 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(4-chloro-2-fluorophenyl)-aminocarbonylmethyl)-2-pyrrolidone,
 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(3-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(4-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
 25 4-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(*N*-(4-nitrophenyl)-aminocarbonylmethyl)-2-pyrrolidone,
 4-(3-Cyclopentyloxy-4-difluoromethoxyphenyl)-1-(*N*-(2,3-difluorophenyl)-aminocarbonylmethyl)-2-pyrrolidone,
 30 4(*S*)-4-(3-Cyclopentyloxy-4-difluoromethoxyphenyl)-1-(*N*-(2-(6-methylpyridyl))-aminocarbonylmethyl)-2-pyrrolidone,

- 4(S)-(4-Methoxy-3-(3(R)-tetrahydrofuranyloxy)phenyl)-1-(N-(2-(6-methylpyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
 1-(N-(2,3-Difluorophenyl)-aminocarbonylmethyl)-4(S)-(4-methoxy-3-(3(R)-tetrahydrofuranyloxy)phenyl)-2-pyrrolidone,
 5 1-(N-(2-(6-Aminopyridyl))-aminocarbonylmethyl)-4(S)-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 4(S)-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-(6-ethylpyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
 4(S)-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-(4,6-dimethylpyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
 10 1-(N-(2-(6-Bromopyridyl))-aminocarbonylmethyl)-4(S)-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 1-(N-(2-(6-Bromopyridyl))-aminocarbonylmethyl)-4(S)-(4-methoxy-3-(3(R)-tetrahydrofuranyloxy)phenyl)-2-pyrrolidone,
 15 4(S)-(4-Methoxy-3-(3(R)-tetrahydrofuranyloxy)phenyl)-1-(N-(2-methylphenyl)-aminocarbonylmethyl)-2-pyrrolidone,
 4(S)-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3-(2-methoxypyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
 1-(N-(6-(3-Bromo-2-methylpyridyl))-aminocarbonylmethyl)-4(S)-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 20 4(S)-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(3-(4-methoxycarbonyl)-pyridyl)-aminocarbonylmethyl)-2-pyrrolidone,
 4(S)-(3-Cyclopentyloxy-4-methoxyphenyl)-1-(N-(2-(6-methylpyridyl))-aminocarbonylmethyl)-2-pyrrolidone,
 25 1-(N-(3-(2-Cyanopyridyl))-aminocarbonylmethyl)-4(S)-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidone,
 4(S)-(4-Methoxy-3-(3(R)-tetrahydrofuranyloxy)phenyl)-1-(N-(2-(6-methylpyridinyl))-aminocarbonylmethyl)-2-pyrrolidone, and

physiologically acceptable salts thereof, wherein in each case the compound can
 30 be in the form of a mixture of enantiomers such as the racemate, or a mixture of diastereomers, or can be in the form of a single enantiomer or a single diastereomer.

8. A compound selected from:

- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone-1-acetic acid,
(4S)-1-(N-Methoxycarbonylmethyl)-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-
5 2-pyrrolidone,
(4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-methylphenyl)-(N-methyl)aminocarbonylmethyl]-2-pyrrolidone,
(4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-(6-methylpyridyl)-(N-methyl)aminocarbonylmethyl)-2-pyrrolidone,
10 (4S)-1-[N-(2,3-Difluorophenyl)-(N-methyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
(4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(phenylaminocarbonylmethyl)-2-pyrrolidone,
(4S)-1-[N-(4,5-Dimethylthiazol)-2-yl]aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-
15 tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
(4S)-1-[N-(3-Chlorophenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
(4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(3-methoxycarbonylphenyl)aminocarbonylmethyl]-2-pyrrolidone,
20 (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(3-phenpropyl)-2-pyrrolidone,
(4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-phenoxyethyl)-2-pyrrolidone,
(4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-phenthioethyl)-2-
25 pyrrolidone,
(4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-(N-phensulfonylaminocarbonylmethyl)-2-pyrrolidone,
(4S)-1-[N-(2,3-Difluorophenyl)-(N-ethyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
30 (4S)-1-[N-(2,3-Difluorophenyl)-(N-isopropyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,

- (4S)-1-[N-(2,3-Difluorophenyl-(N-cyclopropylmethyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-1-[N-(4-Carboxyphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 5 (4S)-1-[N-(3-Fluorophenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-thiazolyl)aminocarbonylmethyl]-2-pyrrolidone,
- (4S)-1-[N-(4-Methoxyphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 10 (4S)-1-[N-(2,6-Dimethylphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-1-[N-(4-Isopropylphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 15 (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(3,4-methylenedioxyphenyl)aminocarbonylmethyl]-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-(4-trifluoromethyl)pyridyl)aminocarbonylmethyl]-2-pyrrolidone,
- (4S)-1-[N-(3-Carboxyphenyl)aminocarbonylmethyl]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 20 (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-phenylsulfonyl)ethyl]-2-pyrrolidone,
- (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-methylphenyl)sulfonylaminocarbonylmethyl]-2-pyrrolidone,
- 25 (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(2-(4-methoxyphenyl)oxyethyl)]-2-pyrrolidone,
- (4S)-1-[N-(2-(5-Chlorobenzoxazolyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- (4S)-1-[N-(2-(Benzthiazolyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
- 30

(4S)-1-[N-(2-(6-Fluorobenzthiazolyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone,
(4S)-1-[N-(2-(Benzimidazolyl)aminocarbonylmethyl)]-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone, and

5 physiologically acceptable salts thereof, wherein in each case the compound can be in the form of a mixture of enantiomers such as the racemate, or a mixture of diastereomers, or can be in the form of a single enantiomer or a single diastereomer.

9. A compound selected from:

10

(4S)-1-[2-(3-Chlorophenoxy)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,

(4S)-1-[2-(4-Isopropylphenoxy)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,

15 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methylbenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone,

(4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-methylthiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone,

20 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(6-methylbenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone,

(4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methoxybenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone,

(4S)-1-[N-(6-Ethoxycarbonylbenzothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,

25 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(6-trifluoromethoxybenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone,

(4S)-1-[N-(4-tert-Butylthiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,

30 (4S)-1-[2-(4-Isopropylphenylthio)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,

- (4S)-1-[2-(3-Chlorophenylthio)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[2-(2,3-Difluorophenoxy)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 5 (4S)-1-[N-(2,3-Difluorophenyl)-N-(2-méthylpropyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(4-Isopropoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(4-Fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 10 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[2-(1,2,3,4-tetrahydroisoquinoliny)carbonylmethyl]-2-pyrrolidone,
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[1-(1,2,3,4-tetrahydroquinoliny)carbonylmethyl]-2-pyrrolidone,
- 15 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-trifluoromethoxyphenyl)aminocarbonylmethyl]-2-pyrrolidone,,
- (4S)-1-[N-(6-Fluorobenzothiazol-2-yl)-N-(methyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[2-(Benzothiazol-2-yl)oxyethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 20 (4S)-1-[2-(6-Fluorobenzothiazol-2-yl)thioethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(6-Fluorobenzothiazol-2-yl)aminoethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone[,
- 25 (4S)-1-[N-(Benzothiazol-2-yl)aminoethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-N-[2-(2-oxopyrrolidin-1-yl)ethyl]-4-phenoxybenzamide,
- (4S)-1-[N-(3-Fluorophenyl)-N-(methyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 30

- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methoxyphenyl)-N-(methyl)aminocarbonylmethyl]-2-pyrrolidone,
- (4S)-1-[N-(4-Isopropylphenyl)-N-(methyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 5 (4S)-1-[N-(3,4-Methylenedioxyphenyl)-N-(methyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-methylthiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone,
- (4S)-1-[N-(4-tert-Butylphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 10 (4S)-1-[N-(6-Chlorobenzothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-methyl-N-(thiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone,
- 15 (4S)-1-[N-(Benzothiazol-2-yl)-N-(cyclopropylmethyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(Indol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(Indan-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 20 (4S)-1-[N-(5-Chlorothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-phenylthiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone,
- 25 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(6-methoxybenzothiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone,
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-N-[2-(2-oxopyrrolidin-1-yl)ethyl]benzamide,
- (4S)-2,3-Difluoro-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-N-[2-(2-oxopyrrolidin-1-yl)ethyl]-benzamide,
- 30

- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-N-[2-(2-oxopyrrolidin-1-yl)ethyl]-4-methoxybenzamide,
- (4S)-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[2-(4-trifluoromethylphenoxy)ethyl]-2-pyrrolidone,
- 5 (4S)-1-[N-(5-Cyclopropyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(Benzothiazol-6-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(4-Ethoxycarbonylthiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 10 (4S)-1-[N-(5-tert-Butyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-Cyclopropylmethyl-N-(6-fluorobenzothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 15 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-(2-oxo-2-phenylethyl)-2-pyrrolidone,
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-(2-oxo-2-(4-methoxyphenyl)ethyl)-2-pyrrolidone,
- (4S)-1-[N-(2,4-Dimethoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 20 (4S)-1-[N-(3,5-Dimethoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(2,2,3,3-tetrafluorobenzo-1,4-dioxan-6-yl)aminocarbonylmethyl]-2-pyrrolidone,
- 25 (4S)-1-[N-(3,4-(Difluoromethylene)dioxyphenyl)-N-methylaminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(3-Fluoro-4-methoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(1,4-Benzodioxan-6-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 30

- (4S)-1-[N-(2-Fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(3,4-Dimethoxyphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 5 (4S)-1-[N-(3,4-Difluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(4-Methanesulfonamidophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(4-(4-Fluorophenyl)thiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 10 (4S)-1-[N-(3-Fluoro-4-methylphenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(4,6-Difluorobenzothiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 15 (4S)-1-[N-(4-Carboxythiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4R)-1-[N-(3-Fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3S)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[2-(2-Fluorophenylthio)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 20 (4S)-1-[2-(3-Fluorophenylthio)ethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone [431.52 MW; 432.1 M+H],
- (4S)-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[2-(4-methoxyphenylthio)ethyl]-2-pyrrolidone,
- 25 (4S)-1-[N-(4-Carboxy-3-fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,,
- (4S)-1-[N-(4-Ethanesulfonamidophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(4-Benzenesulfonamidophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 30

- (4S)-1-[N-(4-(4-Fluorobenzene)sulfonamidophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(2,3-Difluorobenzyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 5 (4S)-1-[N-(5-Cyclopropylmethyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(6-Ethylpyridin-2-yl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(3-Fluorobenzyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 10 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(2-methylbenzyl)aminocarbonylmethyl]-2-pyrrolidone,
- (4S)-1-[N-(4-Methanesulfonylbenzyl)aminocarbonylmethyl]-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- 15 (4S)-1-[N-(4-Aminosulfonylbenzyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-1-[N-(Benzothiazol-2-yl)methylaminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone,
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(3-methylpyridin-2-yl)methylaminocarbonylmethyl]-2-pyrrolidone,
- 20 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-trifluoromethyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone,
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-pyridyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone,
- 25 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-(3-pyridyl)thiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone,
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-(2-pyridyl)thiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone,
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(4-(4-pyridyl)thiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone,
- 30

- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-pyridyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-ethoxycarbonyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 5 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-methoxycarbonyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(3,4-methylenedioxyphenyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-thienyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 10 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-thienylmethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-propyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 15 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-pyrazinyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-methoxymethyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(2-tetrahydrofuranyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 20 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-aminosulfonyl-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-methoxyphenyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 25 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-methoxyphenyloxymethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-morpholinylcarbonylmethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(1-piperidinylcarbonylmethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 30

- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(1-pyrrolidinylcarbonylmethyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-piperidinyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- 5 (4S)-4-[4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-1-[N-(5-(4-(N-tertbutyloxycarbonyl)piperidinyl)-1,3,4-thiadiazol-2-yl)aminocarbonylmethyl]-2-pyrrolidone
- (4S)-1-[N-(2,3-Difluorophenylaminocarbonylmethyl)-4-(4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-2-pyrrolidone
- 10 (4S)-4-(4-Methoxy-3-(3R)-tetrahydrofuranyloxyphenyl)-1-[N-(4-ethoxycarbonylphenyl)aminocarbonylmethyl]-2-pyrrolidone,
- (4S)-1-[N-(4-tert-butyloxycarbonyl-3-fluorophenyl)aminocarbonylmethyl]-4-[4-methoxy-3-(3R)-tetrahydrofuranyloxyphenyl]-2-pyrrolidone, and
- physiologically acceptable salts thereof, wherein in each case the compound can
- 15 be in the form of a mixture of enantiomers such as the racemate, or a mixture of diastereomers, or can be in the form of a single enantiomer or a single diastereomer.

10. A pharmaceutical composition comprising a compound of any one of claims 1 to 9 and a pharmaceutically acceptable carrier.
- 20 11. A composition of claim 10, wherein the compound is provided in a unit dosage of 0.1 - 50 mg.
12. A method for effecting PDE4 enzyme inhibition, enhancing cognition and/or treating psychosis in a patient comprising administering to said patient an effective amount of a compound according to any one of claims 1 to 9.
- 25 13. A method according to claim 12, wherein said compound is administered in an amount of 0.01-100 mg/kg of body weight/day.

14. A method according to claim 12, wherein said patient is a human.
15. A method of claim 12, wherein the patient is suffering from cognition impairment or decline.
16. A method according to claim 12, wherein said patient is suffering from
5 memory impairment.
17. A method according to claim 16, wherein said patient is suffering from memory impairment due to Alzheimer's disease, schizophrenia, Parkinson's disease, Huntington's disease, Pick's disease, Creutzfeld-Jakob disease, HIV, cardiovascular disease, head trauma or age-related cognitive decline.
- 10 18. A method according to claim 16, wherein said patient is suffering from memory impairment due to dementia.
19. A method according to claim 16, wherein said patient is suffering from memory impairment Alzheimer's disease, schizophrenia, Parkinson's disease, Huntington's disease, Pick's disease, Creutzfeld-Jakob disease, depression, aging, head
15 trauma, stroke, CNS hypoxia, cerebral senility, multiinfarct dementia, an acute neuronal disease, HIV or a cardiovascular disease.
20. A method according to claim 12, wherein said patient is suffering from a psychosis.
21. The method of claim 20, wherein the psychosis is schizophrenia, bipolar
20 or manic depression, major depression, drug addiction or morphine dependence.

22. A method for treating a patient having a disease involving decreased cAMP levels comprising administering to said patient an effective amount of a compound according to any one of claims 1 to 9.

23. A method of claim 12, wherein the patient is treated to effect PDE4
5 enzyme inhibition.

24. A method of treating a patient suffering from an allergic or inflammatory disease comprising administering to said patient an effective amount of a compound according to any one of claims 1 to 9.

25. A method of claim 24, wherein the patient is suffering from chronic
10 obstructive pulmonary disease.

26. A method of treating a patient suffering from neurodegeneration resulting from a disease or injury comprising administering to said patient an effective amount of a compound according to any one of claims 1 to 9.

27. The method of claim 26, wherein the disease or injury is stroke, spinal
15 cord injury, neurogenesis, Alzheimer's disease, multiple sclerosis, amyotrophic lateral sclerosis (ALS), or multiple system atrophy (MSA).