

AMINOESTER DERIVATIVES

FIELD OF THE INVENTION

The present invention relates to novel compounds which are both phosphodiesterase 4 (PDE4) enzyme inhibitors and muscarinic M3 receptor antagonists. More particularly, the invention relates to compounds of formula (I) as below described, methods of preparing such compounds, compositions containing them and therapeutic use thereof.

BACKGROUND OF THE INVENTION

Chronic obstructive pulmonary disease (COPD) is a respiratory disorder characterized by progressive, not fully reversible, airflow limitation associated with an abnormal pulmonary inflammatory response to noxious particles or gases.

For this reason, bronchial relaxation and inflammatory response suppression represent a mechanistic approach to the treatment of COPD that might improve symptoms such as dyspnea, wheezing, chest tightness, cough and mucus secretion, improve health status and reduce exacerbations.

Nowadays, the drug therapy options for COPD fall into 2 general classes: bronchodilators, (β 2-adrenoceptor agonists, antimuscarinic agents and methylxanthines) and anti-inflammatory agents (glucocorticosteroids and selective phosphodiesterase-4 (PDE4) inhibitors).

Bronchodilator drugs are the current mainstay of treatment for symptoms' relief.

As anticholinergic bronchodilators, the efficacy of muscarinic M3 antagonists is based on the fact that the major reversible component of airflow narrowing in COPD patients is the increase of acetylcholine (ACh) released to airway smooth muscle, by the bronchial postganglionic vagal efferent in some pathological conditions. Therefore, compounds that antagonize the action of ACh at muscarinic receptors are able to counteract the bronchoconstriction and thus improve lung function in these patients.

Muscarinic antagonists block the effects of ACh at muscarinic receptors.

Currently, there are five known muscarinic receptor subtypes (M1 – M5); human

airway smooth muscle contains M1, M2 and M3 receptors. M1 receptors facilitate neurotransmission through parasympathetic ganglia and are weakly expressed on submucosal glands in human airways. The M2 receptors are located on the smooth-muscle fibers. Some studies have suggested a small role of M2 mediating the inhibition of airway smooth-muscle relaxation caused by adenylyl cyclase activation by compounds such as beta agonists. In addition, presynaptic M2 receptors are found on postganglionic parasympathetic nerves that project to airway smooth muscle and mucus-producing cells.

These presynaptic M2 autoreceptors provide a negative feedback mechanism, which, when stimulated, inhibit further release of ACh. Postsynaptic M3 receptors are known to mediate both contraction of smooth muscle in the respiratory tract and mucus secretion, making them a major target for symptomatic relief of COPD. Consequently, in the airways, the major effects of muscarinic antagonists are bronchodilation and reduction of mucus secretion via blockage of ACh-induced effects in the parasympathetic nervous system.

Given the distribution of muscarinic receptors, systemically available agents that bind to muscarinic receptors outside of the respiratory tract have the potential to produce unwanted side effects such as tachycardia, dry mouth, urinary retention and constipation. Whereas dry mouth is the most common systemic anticholinergic side effect associated with the use of antimuscarinic antagonists as a result of the systemic blockade of M1 and M3 receptors the most potentially serious systemic effect is tachycardia, which results from the blockade of cardiac M2 receptors.

Inhaled anticholinergic antimuscarinic drugs approved for the treatment of COPD include ipratropium bromide (Atrovent®), oxitropium bromide (Oxivent®) and tiotropium bromide (Spiriva®). Both ipratropium and oxitropium are short-acting agents. In contrast, tiotropium bromide is the only long-acting antimuscarinic agent (LAMA) currently marketed for COPD, proved to be suitable for once-daily administration as a dry powder. Several others newer LAMAs are newly registered for the treatment of COPD, including aclidinium bromide and glycopyrrolate bromide, or are currently in phase III development,

including umeclidinium.

Although bronchodilators are quite effective to improve symptoms, they do not address the underlying chronic inflammation or the changes in airway structure.

Standard treatment with glucocorticosteroids as anti-inflammatory agents has
5 demonstrated limited efficacy. However, among the anti-inflammatory agents currently being developed, PDE4 inhibitors proved to be effective in attenuating the responses of various inflammatory cells, through their ability to elevate cAMP levels.

PDE4 is the predominant PDE expressed in neutrophils and T cells, suggesting that PDE4 inhibitors would be effective in controlling inflammation in COPD. Inhibition of
10 PDE4 in inflammatory cells influences various specific responses, such as the production and/or release of pro-inflammatory mediators including cytokines and reactive oxygen species, with a well-documented efficacy in animal models mimicking certain aspects of asthma and COPD, as well as inflammatory bowel disease, atopic dermatitis, psoriasis and rheumatoid arthritis.

15 The selective PDE4 inhibitor, roflumilast (Daxas®) is an approved phosphodiesterase-4 inhibitor for the treatment of COPD associated with chronic bronchitis and a history of exacerbations. Roflumilast inhibits lung inflammation and emphysema in a smoking model of COPD in mice. In COPD patients, oral roflumilast given over 4 weeks significantly reduces the numbers of neutrophils (by 36%) and CXCL8 concentrations in
20 sputum. In clinical trials roflumilast (500 mg once daily) given over 12 months improved lung function in COPD patients to a small extent but had little effect in reducing exacerbations or improving quality of life. More recently roflumilast has been shown to significantly improve FEV₁ (by approximately 50 mL) and reduce exacerbation (by about 15%) in patients with severe disease who have frequent exacerbations and mucus
25 hypersecretion. Roflumilast provides clinical benefit when added to salmeterol or tiotropium and so may be used as an additional treatment in patients with severe disease.

However, the clinical utility of PDE4 inhibitors has so far been compromised by the occurrence of mechanism-associated side effects, including headache, nausea and emesis,

which often limited the maximally tolerated dose. This problem could be overcome by inhaled delivery and designing compounds with a potentially more advantageous therapeutic window.


Since bronchial relaxation and inflammatory response suppression represent a mechanistic approach to the treatment of COPD, the combination of muscarinic M3 antagonism with selective PDE4 inhibition may lead to a new class of drugs, combining both bronchodilating and anti-inflammatory properties in one molecule, which may open new perspectives in the management of COPD.

Our co-pending applications n. PCT/EP2013/075526, n. PCT/EP2013/075520 and the present invention address the above mentioned need by providing the compounds of the invention.

SUMMARY OF THE INVENTION

The invention is directed to compounds acting both as inhibitors of the phosphodiesterase 4 (PDE4) enzyme and as muscarinic M3 receptor antagonists, methods of preparing said compounds, compositions containing them and therapeutic use thereof.

In particular the invention is directed to compounds of formula (I),



wherein

each R₁ is hydrogen or is independently selected in the group consisting of: halogen,

(C₁-C₄) alkyl, (C₁-C₄) alkoxy, (C₁-C₄) haloalkyl, hydroxy, -SO₂NR^IR^{II}, -CN, -NR^ISO₂R^{III}, -NR^IR^{II}, -(CO)NR^IR^{II} and -NR^I(CO)R^{III}, and wherein said (C₁-C₄) alkyl is optionally substituted by one or more groups selected from (C₃-C₇) cycloalkyl, hydroxy and -NR^IR^{II} and wherein said (C₁-C₄) alkoxy is optionally substituted by one or more halogens or groups

5 (C₃-C₇) cycloalkyl wherein,

R^I is hydrogen or (C₁-C₆) alkyl;

R^{II} is hydrogen or (C₁-C₆) alkyl;

R^{III} is hydrogen or (C₁-C₆) alkyl;

n is an integer ranging from 1 to 3;

10 each R₂ is hydrogen or is independently selected in the group consisting of: halogen, (C₁-C₄) alkyl, (C₁-C₄) alkoxy, (C₁-C₄) haloalkyl, hydroxy, -SO₂NR^IR^{II}, -CN, -NR^ISO₂R^{III}, -NR^IR^{II}, -(CO)NR^IR^{II} and -NR^I(CO)R^{III} and wherein said (C₁-C₄) alkyl is optionally substituted by one or more groups selected from (C₃-C₇) cycloalkyl, hydroxy and -NR^IR^{II} and wherein said (C₁-C₄) alkoxy is optionally substituted by one or more halogens or groups

15 (C₃-C₇) cycloalkyl wherein,

R^I is hydrogen or (C₁-C₆) alkyl;


R^{II} is hydrogen or (C₁-C₆) alkyl;

R^{III} is hydrogen or (C₁-C₆) alkyl;

m is an integer ranging from 1 to 3;

20 R₃ and R₄ are different or the same and are independently selected from the group consisting of: H, (C₃-C₇) cycloalkylcarbonyl, (C₁-C₆) alkyl optionally substituted by one or more substituents selected from (C₃-C₇) cycloalkyl and (C₅-C₇) cycloalkenyl, (C₁-C₆) haloalkyl, (C₃-C₇) cycloalkyl, (C₅-C₇) cycloalkenyl, (C₂-C₆) alkenyl, and (C₂-C₆) alkynyl; or R₃ and R₄, together with the interconnecting atoms, form a 2,2-difluoro-1,3-dioxolane
25 ring of formula (r) fused to the phenyl moiety which bears groups -OR₃ and -OR₄, wherein asterisks indicate carbon atoms shared with such phenyl ring:

6



(r);

each R_5 is selected from the group consisting of: CN, NO_2 , CF_3 and halogen atoms;

k is 0 or an integer ranging from 1 to 3;

5 x' is 0 or 1;

L_1 is selected from a bond and $-(CH_2)_p-$ wherein p is an integer ranging from 1 to 4;

W_1 is selected from a divalent arylene, heteroarylene and saturated monocyclic heterocycloalkylene group;

10 L_2 is a group selected from: a bond, $-(CH_2)_q-$ wherein q is 1 or 2, $[1]-(CO)-[X]-(CH_2)_t-[2]$, and $[1]-(SO_2)-[X]-(CH_2)_t-[2]$, wherein $[1]$ and $[2]$ represent, respectively the point of attachment of group L_2 to the ring W_1 and to the chain nitrogen atom, and wherein $[X]$ is a bond or a substituted or unsubstituted arylene group;

t is an integer ranging from 1 to 4;

W_2 is selected from an aryl and heteroaryl group;

15 L is a bond or a $-(CH_2)-$ group;

R_6 is selected in the group consisting of (C_1-C_4) alkyl, (C_1-C_4) alkoxy, (C_1-C_4) haloalkyl, and $-CN$, wherein said (C_1-C_4) alkyl is optionally substituted by one or more groups selected from (C_3-C_7) cycloalkyl, (C_1-C_4) alkoxy and hydroxyl, or, in alternative, when R_6 is (C_1-C_4) alkyl, W_2 is a phenyl ring, one of R_1 is an alkyl in ortho position with respect to L , both R_1 and R_6 may be connected to form with W_2 a condensed ring radical selected from at least 1H-cyclopropabenzene-1,1-diyl, indane-1,1-diyl (also named as 2,3-dihydro-1H-indene-1,1-diyl), indane-2,2-diyl (also named as 2,3-dihydro-1H-indene-2,2-diyl), 1,2,3,4-tetrahydronaphthalene-1,1-diyl, and 1,2,3,4-tetrahydronaphthalene-2,2-diyl;


25 R_7 is selected from hydrogen and (C_1-C_4) alkyl optionally substituted by hydroxy or $-NR_{11}R_{12}$ and wherein R_{11} and R_{12} are independently selected from hydrogen, (C_1-C_4) alkyl, or, together with the nitrogen atom they are linked to may form a saturated

heterocycloalkyl group having an additional heteroatom selected from O, S and NH;

A is a nitrogen containing group which may be selected from:

- a group (a) which is $-(CH_2)_s-NR_8R_9$ wherein s is an integer ranging from 1 to 4 and R_8 and R_9 are independently hydrogen or (C_1-C_4) alkyl; and
 - 5 - a group (b) which is a saturated monocyclic, bicyclic or tricyclic heterocyclic ring system optionally substituted by one or two groups R_{10} which are at each occurrence independently selected from (C_1-C_4) alkyl and benzyl;
- deuterated derivatives, and pharmaceutically acceptable salts, or solvates thereof.

The invention further concerns compounds of formula (I) wherein x' is 1 which are
 10 represented by the formula (IA) wherein a negative charge may be present in the oxygen atom and a corresponding positive charge is on the adjacent nitrogen atom.



15 wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , L , L_1 , W_1 , L_2 , W_2 , A , m , n , and k are as described above.

The invention further concerns the corresponding deuterated derivatives of compounds of formula (I) wherein at least one hydrogen atom is substituted by corresponding atoms of deuterium.

The invention also encompasses the pharmaceutically acceptable salts and/or solvates thereof.

The term “pharmaceutically acceptable salts” refers to derivatives of compounds of formula (I) wherein the parent compound is suitably modified by converting any of the free
5 acid or basic group, if present, into the corresponding addition salt with any base or acid conventionally intended as being pharmaceutically acceptable.

Suitable examples of said salts may thus include mineral or organic acid addition salts of basic residues such as amino groups, as well as mineral or organic acid residues such as carboxylic groups.

10 Cations of inorganic bases which can be suitably used to prepare salts within the invention comprise ions of alkali or alkaline earth metals such as potassium, sodium, calcium or magnesium.

Those obtained by reacting the main compound, functioning as a base, with an inorganic or organic acid to form a salt comprise, for example, salts of hydrochloric acid,
15 hydrobromic acid, sulfuric acid, phosphoric acid, acetic acid, methane sulfonic acid, camphor sulfonic acid, oxalic acid, maleic acid, fumaric acid, succinic acid and citric acid.

The skilled person will appreciate that many organic compounds can form complexes with solvents in which they are reacted or from which they are precipitated or crystallized. These complexes are known as “solvates”. Pharmaceutically acceptable
20 solvates of compound of the invention are within the scope of the invention.

Included within the scope of the present invention are also polymorphs and crystalline forms of compounds of formula (I), or of pharmaceutically acceptable salts, or solvates thereof.

Hereinafter, compounds of formula (I), (IA), (IB), (IC), (ID), (Ia), (Ib), (Ic), (Id) and
25 (I)', enantiomers, diastereoisomers thereof, their pharmaceutically acceptable salts and solvates, and polymorphs or crystalline forms thereof defined in any aspect of the invention (except intermediate compounds described in the chemical processes) are referred to as “compounds of the invention”.

The invention further comprises a process for the preparation of compounds of the invention.

The invention also provides pharmaceutical compositions of compounds of the invention either alone or in combination, in admixture with one or more pharmaceutically acceptable carriers.

In a further aspect the invention provides the use of the compounds of the invention as a medicament.

In one aspect the invention provides the use of the compounds of the invention for the manufacture of a medicament.

10 In particular the invention provides the use of the compounds of the invention for the prevention and/or treatment of any disease wherein an inhibition of PDE4 activity along with muscarinic M3 receptor antagonism is desirable.

In particular the compounds of the invention alone or combined with other active ingredients may be administered for the prevention and/or treatment of a disease the respiratory tract characterized by airway obstruction such as asthma and COPD. In one embodiment, the compounds of the invention may be administered for the prevention and/or treatment of COPD.

15 In a further aspect the invention provides the use of compounds of the invention for the preparation of a medicament for the prevention and/or treatment of any disease wherein an inhibition of PDE4 activity along with muscarinic M3 receptor antagonism is desirable.

Moreover the invention provides a method for prevention and/or treatment of any disease wherein an inhibition of PDE4 activity along with muscarinic M3 receptor antagonism is desirable, said method comprises administering to a patient in need of such treatment a therapeutically effective amount of a compound of the invention.

25 A further aspect of the invention provides a suitable inhalation device, comprising a pharmaceutical composition of a compound of the invention, which may be respectively selected from a single- or multi-dose dry powder inhaler, a pressurized metered dosed inhaler or a nebulizer and in particular a soft mist nebulizer.

A further aspect of the invention provides a kit comprising the pharmaceutical compositions of a compound of the invention either alone or in combination with one or more active ingredient and a device which may be a single- or multi-dose dry powder inhaler, a metered dose inhaler or a nebulizer.

5 **DEFINITIONS**

The term “halogen atoms” includes fluorine, chlorine, bromine, and iodine, preferably chlorine.

The term “(C₁-C_x) alkyl” where x is an integer greater than 1, refers to straight-chained and branched alkyl groups wherein the number of constituent carbon atoms is in
10 the range 1 to x. Particular alkyl groups are methyl, ethyl, n-propyl, i-propyl, n-butyl, sec-butyl, i-butyl and t-butyl.

By analogy, the term “(C₁-C_x)alkylene” refers to a divalent (C₁-C_x)alkyl radical, wherein (C₁-C_x)alkyl is as above defined.

The term “(C₁-C_x) alkoxy” or “(C₁-C_x) alkoxy” where x is an integer greater than
15 1 refers to straight-chained and branched alkoxy groups wherein the number of constituent carbon atoms is in the range 1 to x. Particular alkoxy groups are methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, sec-butoxy, i-butoxy and t-butoxy.

The expressions “(C₁-C_x)haloalkyl” refer to the above defined “(C₁-C_x)alkyl” groups wherein one or more hydrogen atoms are replaced by one or more halogen atoms,
20 which can be the same or different from each other. Examples of said (C₁-C₆)haloalkyl groups may thus include halogenated, poly-halogenated and fully halogenated alkyl groups wherein all of the hydrogen atoms are replaced by halogen atoms, e.g. trifluoromethyl or difluoro methyl groups.

The term “(C₃-C_y) cycloalkyl”, where y is an integer greater than or equal to 3, refers
25 to saturated cyclic hydrocarbon groups containing from 3 to y ring carbon atoms. Examples include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

The expression “(C₃-C_y)cycloalkylcarbonyl” refers to (C₃-C_y)cycloalkylCO- groups wherein the group “(C₃-C_y)cycloalkyl” has the meaning above defined.

The term “(C₂-C₆)alkenyl” refers to straight or branched, conjugated or not conjugated, carbon chains with one or more double bonds, in cis or trans configuration, wherein the number of atoms is in the range 2 to 6.

The term “(C₅-C_z) cycloalkenyl”, where z is an integer greater than or equal to 5,
5 refers to cyclic hydrocarbon groups containing from 5 to z ring carbon atoms and one or more double bonds.

The term “(C₂-C₆)alkynyl” refers to straight or branched carbon chains with one or more triple bonds wherein the number of atoms is in the range 2 to 6.

The term “arylene” refers to divalent mono- or bi-cyclic systems which have 6 to
10 ring carbon atoms, wherein at least one ring is aromatic. Examples of suitable arylene include, for instance phenylenediyl, naphthalenediyl, tetrahydronaphthalenediyl, indanediyl, indenediyl radicals at any suitable position, and the like.

The expression “heteroarylene” refers to divalent monocyclic ring systems with 5 to 6 ring atoms, and in which at least one ring atom is a heteroatom (e.g. N, NH, S or O).
15 Not limiting examples of suitable heteroarylene systems include, for instance, thiophenediyl, furanediyl, pyrrolediyl, pyrazolediyl, imidazolediyl, triazolediyl, tetrazolediyl, isoxazolediyl, oxazolediyl, isothiazolediyl, thiazolediyl, pyridinediyl radicals at any suitable position, and the like.

The expression “heterocycloalkyl” refers to monocyclic cycloalkyl groups with 3 to
20 6 ring atoms, in which at least one ring carbon atom is replaced by a heteroatom (e.g. N, NH, S or O). Not limiting examples of heterocycloalkyl are represented by: pyrrolidinyl, thiazolidinyl, imidazolidinyl, oxazolidinyl, piperazinyl, piperidinyl, morpholinyl, thiomorpholinyl, azetidiny.

The expression “saturated monocyclic heterocycloalkylene” refers to divalent saturated
25 monocyclic cycloalkyl groups with 3 to 6 ring atoms in which at least one ring carbon atom is replaced by a heteroatom (e.g. N, NH, S or O). Not limiting examples of “saturated monocyclic heterocycloalkylene” are represented by: pyrrolidinediyl, thiazolidinediyl, imidazolidinediyl, oxazolidinediyl, piperazinediyl, piperidinediyl, morpholinediyl, thiomorpholinediyl,

azetidinediyl radicals at any suitable position, and the like.

The term “aryl” refers to mono or bi-cyclic systems which have 6 to 10 ring carbon atoms, wherein at least one ring is aromatic.

The expression “heteroaryl” refers to mono or bi-cyclic systems with 5 to 11 ring atoms, in which at least one ring is aromatic and in which at least one ring atom is a heteroatom (e.g. N, NH, S or O).

Examples of suitable aryl or heteroaryl monocyclic systems with 5 to 6 ring atoms include, for instance, phenyl, thienyl, pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, isoxazolyl, oxazolyl, isothiazolyl, thiazolyl, pyridyl, furanyl derived radicals and the like.

Examples of suitable aryl or heteroaryl bicyclic systems with more than 6 ring atoms include naphthalenyl, biphenylenyl, tetrahydronaphthalenyl, purinyl, pteridinyl, benzimidazolyl, benzotriazolyl, quinolinyl, isoquinolinyl, indolyl, isoindolyl, indazolyl, benzothienyl, benzofuranyl, benzoxazolyl, dihydrobenzo dioxinyl, dihydrobenzo dioxepinyl, benzo-oxazinyl radicals and the like.

The expression “heterocyclic ring system” refers to optionally substituted mono-, bi- or tri-cyclic ring systems which may be saturated, partially unsaturated or unsaturated, such as heterocycloalkyl or heteroaryl having 5 to 11 ring atoms in which at least one ring atom is a heteroatom (e.g. N, S or O). Examples of “heterocyclic ring system” are represented by: pyrrolidinyl, imidazolidinyl, piperazinyl, piperidinyl, quinuclidinyl, 8-azabicyclo[3.2.1]octanyl or dehydroxy scopine radical all optionally substituted by oxygen, (C₁-C_x) alkyl or benzyl on a nitrogen atom.


As used in the present description an oxo moiety is represented by (O) as an alternative to the other common representation, e.g. (=O). Thus, in terms of general formula, the carbonyl group is herein preferably represented as -C(O)- as an alternative to the other common representations such as -CO- , -(CO)- or -C(=O)- . In general the bracketed group is a lateral group, not included into the chain, and brackets are used, when deemed useful, to help disambiguating linear chemical formulas; e.g. the sulfonyl group -SO₂- might be also represented as -S(O)₂- to disambiguate with respect to the sulfinic

group -S(O)O-.

DETAILED DESCRIPTION OF THE INVENTION

The invention is directed to a class of compounds acting both as inhibitors of the phosphodiesterase 4 (PDE4) enzyme and as muscarinic M3 receptor antagonists.


- 5 The present invention relates to derivatives of general formula (I), deuterated derivatives and pharmaceutically acceptable salts or solvates thereof,




(I)

wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, L₁, W₁, L₂, W₂, L, A, n, m, k and x' are as above defined.


- 10 Preferred compounds of formula (I) are those wherein the "saturated heterocyclic ring system" A is represented by a group of formula (i),(ii), (iii) or (iv):




(i)



(ii)



(iii)



(iv)

wherein


$f = 1, 2 \text{ or } 3$;

$g = 1, 2 \text{ or } 3$.


and the asterisk (*) represents the point of attachment to the oxygen atom of formula (I).

More preferably A is represented by a group of formula (i) or (ii):

5



(i)




(ii)

wherein f is 1, g is 2 and the asterisk (*) represents the point of attachment to the oxygen atom of formula (I).

10

It will be apparent to those skilled in the art that compounds of general formula (I) at least contain one stereogenic center, namely represented by the carbon atom (1), and therefore exist as optical stereoisomers.



(I)

15


It will be apparent to the skilled person that compounds according to the invention may have two stereogenic centers (for instance at carbon atoms (1) and (2)), thus they may accordingly exist as four diastereoisomers. Where the compounds according to the invention possess more than two stereogenic centers, they will exist as 2^n diastereoisomers (wherein n here refers to the number of stereogenic centers). It is to be understood that all such isomers and mixtures thereof in any proportion are encompassed within the scope of

20

the present invention.

In a preferred embodiment, the invention is directed to compounds of formula (I)', which are compounds of formula (I) as above defined where the absolute configuration of carbon (1) is that shown herebelow:


5



The absolute configuration for carbon (1) is assigned on the basis of Cahn-Ingold-Prelog nomenclature based on groups' priorities.

10 In one preferred embodiment, for compounds of formula (I), absolute configuration at carbon (1) is (S).

In one embodiment, when A is a group of formula (i) as previously defined, compounds of formula (I), wherein (2) is an asymmetric center, may exist at least as four couples of diastereoisomers (Ia), (Ib), (Ic) and (Id) herebelow reported, which are
15 comprised within the scope of the present invention.




It will be apparent to the skilled person that compounds of formula (Ia), (Ib), (Ic), (Id) may be also obtained as single diastereoisomers wherein the configuration at the stereogenic centre at carbon atom (2) is defined as (R) or (S).

5 In one embodiment, compounds of formula (Ia) are provided as above reported, or single diastereoisomers thereof.

It is to be understood that all preferred groups or embodiments described herebelow and hereabove for compounds of formula (I) may be combined among each other and apply to compounds of formula (IA), (IB), (IC), (ID), (Ia), (Ib), (Ic), (Id) and (I)' as well *mutatis*
 10 *mutandis*.

In one embodiment, the invention provides compounds of formula (IA), which are N-oxides on the pyridine ring of compounds of formula (I) wherein x' is 1, deuterated derivatives and pharmaceutically acceptable salts and solvates thereof:



(IA)

wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, L, L₁, W₁, L₂, W₂, A, m, n, and k are as described above.


- 5 In a preferred embodiment, k is 2 and R₅ are halogen atoms. In a further preferred embodiment such R₅ are two chlorine atoms at positions 3 and 5 of the pyridine ring.

In one preferred embodiment R₄ is selected from a (C₁-C₆) alkyl and R₃ is selected from (C₃-C₇) cycloalkyl or (C₁-C₆) alkyl which is optionally substituted by (C₃-C₇) cycloalkyl.

- 10 In another preferred embodiment, R₃ and R₄ are both methyl.

A preferred compound of formula (I) is that wherein W₂ is a phenyl ring, L is a bond, and one of R₁, in ortho position with respect to L, and R₆ may be connected to form a ring radical, according to the general formula (IB)

18




(IB)

wherein R₁, R₂, R₃, R₄, R₅, R₇, A, L₁, W₁, L₂, m, k and x' are as defined above for compounds of formula (I); deuterated derivatives and pharmaceutically acceptable salts and solvates thereof.

In a more preferred compound of formula (IB) L₁ is a bond, W₁ is a divalent group selected from thiophene-2,5-diyl, thiophene-2,4-diyl, phenylene-1,4-diyl, phenylene-1,3-diyl and phenylene-1,2-diyl, L₂ is -(CH₂)-, R₇ is H and R₁, R₂, R₃, R₄, R₅, A, m and k are as defined above for compounds of formula (I); deuterated derivatives and pharmaceutically acceptable salts and solvates thereof.

Another preferred compound of formula (I) is that wherein W₂ is a phenyl ring and L is a bond according to the general formula (IC):




(IC)

wherein R_6 is selected from methyl, ethyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, methoxymethyl, trifluoromethyl and difluoromethyl and $R_1, R_2, R_3, R_4, R_5, R_7, A, L_1, W_1, L_2, m, n, k$ and x' are as defined above for compounds of formula (I); and pharmaceutically acceptable salts and solvates thereof.

5 In a more preferred compound of formula (IC) L_1 is a bond, W_1 is a divalent group selected from thiophene-2,5-diyl, thiophene-2,4-diyl, phenylene-1,4-diyl, phenylene-1,3-diyl and phenylene-1,2-diyl, L_2 is $-(CH_2)-$, R_7 is H or methyl, R_6 is selected from methyl, ethyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, methoxymethyl, trifluoromethyl and difluoromethyl and $R_1, R_2, R_3, R_4, R_5, A, m, n, k$ and x' are as defined above for
10 compounds of formula (I); and pharmaceutically acceptable salts and solvates thereof.

Another preferred compound of formula (I) is that wherein L_1 is a bond, W_1 is selected from a divalent saturated monocyclic heterocycloalkylene group, represented by the general formula (ID),



(ID)

and wherein

Y is S or CH_2 ;

L_2 is a group selected from $-(CH_2)_q-$ wherein q is 1 or 2, $[1]-(CO)-[X]-(CH_2)_t-[2]$ and $[1]-(SO_2)-[X]-(CH_2)_t-[2]$, wherein $[1]$ and $[2]$ represent, respectively the point of
20 attachment of group L_2 to the saturated monocyclic heterocycloalkylene ring (W_1) and to the chain nitrogen atom, and wherein

$[X]$ is a bond or a substituted or unsubstituted arylene group selected from

phenylene-1,4-, -1,3- and -1,2-diyl ;

t is an integer ranging from 1 to 4;

and wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, A, W₂, m, n, k and x' are as defined above for compounds of formula (I), and pharmaceutically acceptable salts and solvates thereof.

5 According to a preferred embodiment, the present invention provides the compounds reported below:

[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate formate salt

10 [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 4-[[[1-methyl-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]benzoate

[(3R)-quinuclidin-3-yl] 1-[[4-[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethoxy] carbonylphenyl] methylamino]indane-1-carboxylate

15 [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(3R)-quinuclidin-3-yl]oxycarbonylindan-1-yl]amino]methyl]thiophene-2-carboxylate

[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenyl-ethyl]amino]methyl]thiophene-2-carboxylate

20 [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-phenyl-1-[(3R)-quinuclidin-3-yl]oxycarbonyl-propyl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate

25 Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 4-[[[1-methyl-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxyethyl]amino]methyl]benzoate

Single diastereoisomer of [(3R)-quinuclidin-3-yl] 1-[[4-[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethoxy] carbonylphenyl] methylamino]indane-1-carboxylate

Single diastereoisomer of [(3R)-quinuclidin-3-yl] 1-[[4-[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethoxy] carbonylphenyl] methylamino]indane-1-carboxylate

10 Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-[(3R)-quinuclidin-3-yl]oxycarbonylindan-1-yl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-[(3R)-quinuclidin-3-yl]oxycarbonylindan-1-

15 yl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenylethyl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenylethyl]amino]methyl]thiophene-2-carboxylate

[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-[(3R)-1-oxidoquinuclidin-1-ium-3-yl]oxy-2-oxo-1-phenylethyl]amino]methyl]thiophene-2-carboxylate

20 [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(2-dimethylaminoethyloxycarbonyl)indan-1-yl]amino]methyl]thiophene-2-carboxylate

[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-[(1-methyl-4-piperidyl)oxycarbonyl]indan-1-yl]amino]methyl]thiophene-2-carboxylate

[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl]-5-[[[1-(hydroxymethyl)-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenylethyl]amino]methyl]thiophene-2-carboxylate

5 [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]-methylamino]methyl]thiophene-2-carboxylate

[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl]-5-[[[1-(hydroxymethyl)-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate

10 [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(methoxymethyl)-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(2-dimethylaminoethyloxycarbonyl)indan-1-

15 yl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(2-dimethylaminoethyloxycarbonyl)indan-1-

yl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-[(1-methyl-4-piperidyl)oxycarbonyl]indan-1-

20 yl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-[(1-methyl-4-piperidyl)oxycarbonyl]indan-1-

yl]amino]methyl]thiophene-2-carboxylate

25 Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-phenyl-1-[(3R)-quinuclidin-3-yl]oxycarbonyl-

propyl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-phenyl-1-[(3R)-quinuclidin-3-yl]oxycarbonyl-propyl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-

5 dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenyl-ethyl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenyl-ethyl]amino]methyl]thiophene-2-carboxylate

10 Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-

15 yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]-methyl-amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-
20 dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]-methyl-amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(methoxymethyl)-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate

25 and a pharmaceutically acceptable salt or solvate thereof.

The expression “single diastereoisomer” was reported near the chemical name of each compound of formula (I) isolated as single diastereoisomer whose absolute configuration at the stereogenic carbon atom (2) at which R6 is linked was not determined.

The invention also concerns processes for the preparation of compounds of the invention.


Compounds of formula (I) can be obtained according to general synthetic route of Scheme A and Scheme B below reported or following slightly modified procedures that the skilled person can easily apply.

In the following Scheme A and Scheme B only compounds of formula (I) in which x' is 1, corresponding to compounds of formula (IA), are described, starting from the pyridine N-oxides (III) described in the co-pending international application n. PCT/EP2013/075520. Any corresponding compound of formula (I) in which x' is 0 may be similarly obtained starting from the non-oxidized pyridines, analogues of compounds (V) described in the co-pending international application n. PCT/EP2013/075520.


Processes of preparation described below and reported in the following Scheme A should not be viewed as limiting the scope of the synthetic methods available for the preparation of the compounds of the invention.

In the following Scheme A and Scheme B, for compounds of formula (IA) and for compounds of formula (II) to (XXI), unless otherwise indicated, R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , L_1 , W_1 , L_2 , W_2 , L , A , n , m , k and x' are as above defined, PG is a protective group and Y is a group selected from a bond, $-\text{CH}_2-$, $[1]-(\text{CO})-[X]-(\text{CH}_2)_{t'}-[2]$, and $[1]-(\text{SO}_2)-[X]-(\text{CH}_2)_{t'}-[2]$, wherein [1] and [2] represent, respectively the point of attachment of group Y to the ring W_1 and to a formyl group, and wherein [X] is a bond or a substituted or unsubstituted arylene group and t' is 0 or an integer ranging from 1 to 3.

Scheme A




Compounds of formula (IA) may be prepared according to Scheme 1/ (S1) below
 5 by reaction of a compound of formula (II) with a compound of formula (V) as below reported.

Scheme 1 (S1)

Typical reaction conditions comprise reacting a compound of formula (II) with a compound of formula (V) in a suitable solvent, such as acetonitrile, DCM or ethanol in the presence of an acid, such as acetic acid, and an optional base, such as triethylamine, and a reducing agent, such as $NaB(OAc)_3H$ or $NaBH_3CN$, at an appropriate temperature, such as room (or ambient) temperature or $0^\circ C$ or $40^\circ C$.

Compounds of formula (II) may be prepared according to Scheme 2/ (S2) below by reaction of a compound of formula (IV) with a compound of formula (III) as below reported.

Scheme 2 (S2)


Typical reaction conditions comprise reacting a compound of formula (IV) with a compound of formula (III) in a suitable solvent, such as DCM, in the presence of a suitable coupling agent, such as EDC/DMAP or HATU, at an appropriate temperature, such as room (or ambient) temperature.

Compounds of formula (III) may be prepared as described in the co-pending international application n. PCT/EP2013/075520.

Compounds of formula (V) may be prepared according to Scheme 3/ (S3) below by

deprotection of a compound of formula (VI).

Scheme 3 (S3)




5 Typical reaction conditions comprise reacting a compound of formula (VI) with a source of hydrogen, such as ammonium formate or gaseous hydrogen over a suitable catalyst, such as 10% Pd/C, in a suitable solvent, such as EtOAc, at an appropriate temperature, such as room (or ambient) temperature.

Compounds of formula (V) may also be prepared by reacting a compound of
 10 formula (VI) in a suitable solvent, such as 1,4-dioxane, in the presence of a suitable acid, such as hydrochloric acid, at an appropriate temperature, such as room (or ambient) temperature.

Compounds of formula (VI) may be prepared according to Scheme 4/ (S4) below
 by reaction of a compound of formula (VIII) with a compound of formula (VII) as below
 15 reported.

Scheme 4 (S4)




Typical reaction conditions comprise reacting a compound of formula (VIII) with a
 compound of formula (VII) in a suitable solvent, such as THF in the presence of a suitable
 20 coupling agent, such as DCC/HOBt or EDC/DMAP or HATU, at an appropriate temperature, such as room (or ambient) temperature or 40°C.

Compounds of formula (VIII) may be prepared according to Scheme 5/ (S5) below

by reaction of a compound of formula (IX) as below reported.


Scheme 5 (S5)



5 Typical reaction conditions comprise reacting a compound of formula (IX) with benzyl chloroformate or di-*tert*-butyl dicarbonate, in a suitable solvent, such as THF/water or 1,4-dioxane/water, in the presence of a suitable base such as sodium hydroxide, at an appropriate temperature, such as 0°C or room (or ambient) temperature.

Compounds of formula (V) may be prepared according to Scheme 6/ (S6) below by
 10 reaction of a compound of formula (X) as below reported.

Scheme 6 (S6)




Typical reaction conditions comprise reacting a compound of formula (X) in a
 15 suitable solvent, such as 1,4-dioxane, in the presence of a suitable acid, such as hydrochloric acid, at an appropriate temperature, such as room (or ambient) temperature.

Compounds of formula (X) may be prepared according to Scheme 7/ (S7) below by
 reaction of a compound of formula (XI) with a compound of formula (VII) as below
 reported.

20


25

Scheme 7 (S7)

Typical reaction conditions comprise reacting a compound of formula (XI) with a
 5 compound of formula (VII) in a suitable solvent, such as THF in the presence of a suitable coupling agent, such as DCC/HOBt or EDC/DMAP or HATU, at an appropriate temperature, such as room (or ambient) temperature or 40°C.

Compounds of formula (XI) may be prepared according to Scheme 8/ (S8) below by reaction of a compound of formula (XII) as below reported.

10


Scheme 8 (S8)

Typical reaction conditions comprise reacting compound of formula (XII) with a suitable base, such as NaOH in a suitable solvent, such as MeOH at an appropriate temperature, such as room (or ambient) temperature or 50 °C

15

Compounds of formula (XII) may be prepared according to Scheme 9/ (S9) below by reaction of a compound of formula (XIII) as below reported.

Scheme 9 (S9)




Typical reaction conditions comprise reacting compound of formula (XIII) with a suitable Grignard reagent, such as methyl magnesium bromide or ethyl magnesium bromide, in a suitable solvent, such as THF or diethyl ether at a suitable temperature such

5 at -78 °C, 0 °C or room (or ambient) temperature.

Compounds of formula (V) may also be prepared according to Scheme 10 / (S10) below by reaction of a compound of formula (XIV) as below reported.


Scheme 10 (S10)



10 Typical reaction conditions comprise reacting a compound of formula (XIV) in a suitable solvent, such as THF or 1,4-dioxane, in the presence of a suitable acid, such as hydrochloric acid, at an appropriate temperature, such as room (or ambient) temperature.

Compounds of formula (XIV) may be prepared according to Scheme 11 / (S11) below by reaction of a compound of formula (XV) as below reported.


15 **Scheme 11 (S11)**



Typical reaction conditions comprise reacting a compound of formula (XV) with an alkylating agent, such as *para*-formaldehyde or iodomethane, in a suitable solvent, such as THF or 1,4-dioxane, in the presence of a suitable base such as DBU or sodium hydride, at an appropriate temperature, such as 0 °C or room (or ambient) temperature.

- 5 Compounds of formula (XV) may be prepared according to Scheme 12 / (S12) below by reaction of a compound of formula (XVI) as below reported.


Scheme 12 (S12)



- Typical reaction conditions comprise reacting a compound of formula (XVI) with an arylcarboxaldehyde, such as benzaldehyde or 4-methoxybenzaldehyde, in a suitable solvent, such as acetonitrile, DCM or ethanol in the presence of an optional base, such as triethylamine, at an appropriate temperature, such as room (or ambient) temperature or 0 °C or 40 °C.

- Compounds of formula (XVI) may be prepared according to Scheme 13 / (S13) below by reaction of a compound of formula (XVII) as below reported.

Scheme 13 (S13)




- Typical reaction conditions comprise reacting a compound of formula (XVII) with a source of hydrogen, such as ammonium formate or gaseous hydrogen over a suitable catalyst, such as 10% Pd/C, in a suitable solvent, such as EtOAc, at an appropriate

temperature, such as room (or ambient) temperature.

Compounds of formula (XVI) may also be prepared by reacting a compound of formula (XVII) in a suitable solvent, such as 1,4-dioxane, in the presence of a suitable acid, such as hydrochloric acid, at an appropriate temperature, such as room (or ambient) temperature.

Compounds of formula (XVII) may be prepared according to Scheme 14 / (S14) below by reaction of a compound of formula (XVIII) with a compound of formula (VII) as below reported.


Scheme 14 (S14)



Typical reaction conditions comprise reacting a compound of formula (XVIII) with a compound of formula (VII) in a suitable solvent, such as THF in the presence of a suitable coupling agent, such as DCC/HOBt or EDC/DMAP or HATU, at an appropriate temperature, such as room (or ambient) temperature or 40 °C.

Compounds of formula (XVIII) may be prepared according to Scheme 15 / (S15) below by reaction of a compound of formula (XIX) as below reported.

Scheme 15 (S15)




Typical reaction conditions comprise reacting a compound of formula (XIX) with benzyl chloroformate or di-*tert*-butyl dicarbonate, in a suitable solvent, such as THF/water or 1,4-dioxane/water, in the presence of a suitable base such as sodium hydroxide, at an

appropriate temperature, such as 0°C or room (or ambient) temperature.

5


Scheme B



Compounds of formula (IA) may be prepared according to Scheme 16 / (S16) below

10 by reaction of a compound of formula (XX) as below reported.


Scheme 16 (S16)



Typical reaction conditions comprise reacting a compound of formula (XX) in a suitable solvent, such as acetonitrile, DCM or ethanol in the presence of an optional acid, such as acetic acid, and a reducing agent, such as NaB(OAc)₃H or NaBH₃CN, at an appropriate temperature, such as room (or ambient) temperature or 0 °C or 40 °C.

Compounds of formula (XX) may be prepared according to Scheme 17 / (S17) below by reaction of a compound of formula (XXI) as below reported.


Scheme 17 (S17)



Typical reaction conditions comprise reacting a compound of formula (XXI) with an alkylating agent, such as *para*-formaldehyde, in a suitable solvent, such as THF or 1,4-dioxane, in the presence of a suitable base such as DBU or sodium hydride, at an appropriate temperature, such as 0 °C or room (or ambient) temperature.

Compounds of formula (XXI) may be prepared according to Scheme 18 / (S18) below by reaction of a compound of formula (II) with a compound of formula (XVI) as below reported.

Scheme 18 (S18)



Typical reaction conditions comprise reacting a compound of formula (II) with a compound of formula (XVI) in a suitable solvent, such as acetonitrile, DCM or ethanol in the presence of an acid, such as acetic acid, and an optional base, such as triethylamine, at an appropriate temperature, such as room (or ambient) temperature or 0 °C or 40 °C.

The processes described are particularly advantageous as they are susceptible of being properly modulated, through any proper variant known to the skilled person, so as to obtain any of the desired compounds of the invention. Such variants are comprised within the scope of the present invention.

From all of the above, it should be clear to the skilled person that any of the described groups may be present as such or in any properly protected form.

In particular, functional groups present in the compounds of formula II to XIII and which could generate unwanted side reactions and by-products, need to be properly protected before the alkylation, acylation, coupling, oxidation or sulfonylation takes place. Likewise, subsequent deprotection of those same protected groups may follow upon completion of the said reactions.

According to the invention, unless otherwise indicated, the term “protecting group” designates a protective group adapted to preserve the function of the group it is bound to. Typically, protective groups are used to preserve amino, hydroxy, or carboxyl functions. Appropriate protecting groups may thus include, for example, benzyl, benzyloxycarbonyl, t-butoxycarbonyl, alkyl or benzyl esters or the like, which are well known to those skilled in the art [see, for a general reference, T.W. Green; *Protective Groups in Organic Synthesis* (Wiley, N.Y. 1999)].

Likewise, selective protection and deprotection of any of the said groups, for instance including carbonyl, hydroxy or amino groups, may be accomplished according to very well-known methods commonly employed in organic synthetic chemistry.

Optional salification of the compounds of formula (I) may be carried out by properly
5 converting any of the free acidic or amino groups into the corresponding pharmaceutically acceptable salts. In this case too, the operative conditions being employed for the optional salification of the compounds of the invention are all within the ordinary knowledge of the skilled person.

From all of the above, it should be clear to the skilled person that the above process,
10 comprehensive of any variant thereof for the preparation of suitable compounds of the invention, may be conveniently modified so that to adapt the reaction conditions to the specific needs, for instance by choosing appropriate condensing agents, solvents and protective groups, as the case may be.

The present invention also provides pharmaceutical compositions of compounds of
15 the invention in admixture with one or more pharmaceutically acceptable carriers, for example those described in Remington's Pharmaceutical Sciences Handbook, XVII Ed., Mack Pub., N.Y., U.S.A.

Administration of the compounds of the invention or may be accomplished according to patient needs, for example, orally, nasally, parenterally (subcutaneously,
20 intravenously, intramuscularly, intrasternally and by infusion), by inhalation, rectally, vaginally, topically, locally, transdermally, and by ocular administration. Various solid oral dosage forms may be used for administering compounds of the invention including such solid forms as tablets, gelcaps, capsules, caplets, granules, lozenges and bulk powders. The compounds of the invention may be administered alone or combined with various
25 pharmaceutically acceptable carriers, diluents (such as sucrose, mannitol, lactose, starches) and excipients known in the art, including but not limited to suspending agents, solubilizers, buffering agents, binders, disintegrants, preservatives, colorants, flavorants, lubricants and the like. Time release capsules, tablets and gels are also advantageous in administering the

compounds of the present invention.

Various liquid oral dosage forms may also be used for administering compounds of the invention, including aqueous and non-aqueous solutions, emulsions, suspensions, syrups, and elixirs. Such dosage forms can also contain suitable inert diluents known in the art such as water and suitable excipients known in the art such as preservatives, wetting agents, sweeteners, flavorants, as well as agents for emulsifying and/or suspending the compounds of the invention. The compounds of the invention may be injected, for example, intravenously, in the form of an isotonic sterile solution. Other preparations are also possible.

Suppositories for rectal administration of the compounds of the present invention may be prepared by mixing the compound with a suitable excipient such as cocoa butter, salicylates and polyethylene glycols.

Formulations for vaginal administration may be in the form of cream, gel, paste, foam, or spray formula containing, in addition to the active ingredient, such suitable carriers as are known in the art.

For topical administration the pharmaceutical composition may be in the form of creams, ointments, liniments, lotions, emulsions, suspensions, gels, solutions, pastes, powders, sprays, and drops suitable for administration to the skin, eye, ear or nose. Topical administration may also involve transdermal administration via means such as transdermal patches.

For the treatment of the diseases of the respiratory tract, the compounds according to the invention are preferably administered by inhalation.

Inhalable preparations include inhalable powders, propellant-containing metered aerosols or propellant-free inhalable formulations and may be administered through a suitable inhalation device which may be respectively selected from dry powder inhaler, pressurized metered dosed inhaler, or a nebulizer.

For administration as a dry powder, single- or multi-dose inhalers known from the prior art may be utilized. In that case the powder may be filled in gelatine, plastic or other

capsules, cartridges or blister packs or in a reservoir.

A diluent or carrier, generally non-toxic and chemically inert to the compounds of the invention, e.g. lactose or any other additive suitable for improving the respirable fraction may be added to the powdered compounds of the invention.

5 Inhalation aerosols containing propellant gas such as hydrofluoroalkanes may contain the compounds of the invention either in solution or in dispersed form. The propellant-driven formulations may also contain other ingredients such as co-solvents, stabilizers and optionally other excipients.

10 The propellant-free inhalable formulations comprising the compounds of the invention may be in form of solutions or suspensions in an aqueous, alcoholic or hydroalcoholic medium and they may be delivered by jet or ultrasonic nebulizers known from the prior art or by soft-mist nebulizers such as Respimat®.

15 The compounds of the invention may be administered as the sole active agent or in combination with other pharmaceutical active ingredients including those currently used in the treatment of respiratory disorders, e.g. beta2-agonists, antimuscarinic agents, corticosteroids, mitogen-activated protein kinases (P38 MAP kinase) inhibitors, nuclear factor kappa-B kinase subunit beta (IKK2) inhibitors, human neutrophil elastase (HNE) inhibitors, phosphodiesterase 4 (PDE4) inhibitors, leukotriene modulators, non-steroidal anti-inflammatory agents (NSAIDs) and mucus regulators.

20 The present invention also provides combinations of a compound of the invention, with a β_2 -agonist selected from the group consisting of carmoterol, vilanterol (GSK-642444), indacaterol, milveterol, arformoterol, formoterol, salbutamol, levalbuterol, terbutaline, AZD-3199, olodaterol (BI-1744-CL), abediterol (LAS-100977), bambuterol, isoproterenol, procaterol, clenbuterol, reproterol, fenoterol and ASF-1020 and salts thereof.

25 The present invention also provides combinations of a compound of the invention, with a corticosteroid selected from the group consisting of fluticasone propionate, fluticasone furoate, mometasone furoate, beclometasone dipropionate, ciclesonide, budesonide, GSK 685698, GSK 870086.

The present invention also provides combinations of a compound of the invention, with an antimuscarinic agent selected from the group consisting of aclidinium, tiotropium, ipratropium, trospium, glycopyrronium and oxitropium salts.

The present invention also provides combinations of a compound of the invention, with a PDE4 inhibitor selected from the group consisting of AN-2728, AN-2898, CBS-3595, apremilast, ELB-353, KF-66490, K-34, LAS-37779, IBFB-211913, AWD-12-281, cipamfylline, cilomilast, roflumilast, BAY19-8004 and SCH-351591, AN-6415, indus-82010, TPI-PD3, ELB-353, CC-11050, GSK-256066, oglemilast, OX-914, tetomilast, MEM-1414 and RPL-554.

10 The present invention also provides combinations of a compound of the invention, with a P38 MAP kinase inhibitor selected from the group consisting of semapimod, talmapimod, pirfenidone, PH-797804, GSK-725, minokine and losmapimod and salts thereof.

15 In a preferred embodiment, the present invention provides combinations of a compound of the invention with an IKK2 inhibitor.

The invention also provides combinations of a compound of the invention with a HNE inhibitor selected from the group consisting of AAT, ADC-7828, Aeriva, TAPI, AE-3763, KRP-109, AX-9657, POL-6014, AER-002, AGTC-0106, respriva, AZD-9668, zemaira, AAT IV, PGX-100, elafin, SPHD-400, prolactin C and prolactin inhaled.

20 The invention also provides combinations of a compound of the invention with a leukotriene modulator selected from the group consisting of montelukast, zafirlukast and pranlukast.

The invention also provides combinations of a compound of the invention with a NSAID selected from the group consisting of ibuprofen and ketoprofen.

25 The invention also provides combinations of a compound of the invention with a mucus regulator selected from the group consisting of INS-37217, diquafosol, sibenadet, CS-003, talnetant, DNK-333, MSI-1956 and gefitinib.

The dosages of the compounds of the invention depend upon a variety of factors

including the particular disease to be treated, the severity of the symptoms, the route of administration, the frequency of the dosage interval, the particular compound utilized, the efficacy, toxicology profile, and pharmacokinetic profile of the compound.

Advantageously, the compounds of the invention may be administered for example,
5 at a dosage comprised between 0.001 and 1000 mg/day, preferably between 0.1 and 500 mg/day.

When they are administered by inhalation route, the dosage of the compounds of the invention is advantageously comprised between 0.01 and 20 mg/day, preferably between 0.1 and 10 mg/day.

10 Preferably, the compounds of the invention alone or combined with other active ingredients may be administered for the prevention and/or treatment of any obstructive respiratory disease such as asthma, chronic bronchitis and chronic obstructive pulmonary disease (COPD).

The compounds of the invention may be administered for the prevention and/or
15 treatment of any disease wherein PDE4 inhibition or M3 antagonism is required. Said disease include: allergic disease states such as atopic dermatitis, urticaria, allergic rhinitis, allergic conjunctivitis, vernal conjunctivitis, eosinophilic granuloma, psoriasis, inflammatory arthritis, rheumatoid arthritis, septic shock, ulcerative colitis, Crohn's disease, reperfusion injury of the myocardium and brain, chronic glomerulonephritis,
20 endotoxic shock, cystic fibrosis, arterial restenosis, arteriosclerosis, keratosis, rheumatoid spondylitis, osteoarthritis, pyresis, diabetes mellitus, pneumoconiosis, toxic and allergic contact eczema, atopic eczema, seborrheic eczema, lichen simplex, sunburn, pruritus in the anogenital area, alopecia areata, hypertrophic scars, discoid lupus erythematosus, systemic lupus erythematosus, follicular and wide-area pyodermias, endogenous and exogenous
25 acne, acne rosacea, Behçet's disease, anaphylactoid purpura nephritis, inflammatory bowel disease, leukemia, multiple sclerosis, gastrointestinal diseases, autoimmune diseases and the like.

They also include neurological and psychiatric disorders such as Alzheimer's

disease, multiple sclerosis, amyloidosis, amyloid laterosclerosis (ALS), multiple systems atrophy (MSA), schizophrenia, Parkinson's disease, Huntington's disease, Pick's disease, depression, stroke, and spinal cord injury.

The invention will now be further described by way of the following examples.

5

EXAMPLES

Abbreviations

DCC = N,N'-Dicyclohexylcarbodiimide; HOBt = hydroxybenzotriazole; HATU = (Dimethylamino)-N,N-dimethyl(3H-[1,2,3]triazolo[4,5-b]pyridin-3-yloxy) methaniminium hexafluorophosphate; EDC = 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride; DBU = 1,8-Diazabicyclo[5.4.0]undec-7-ene; DMAP = 4-dimethylaminopyridine; DMF = dimethylformamide; DMSO = dimethyl sulfoxide EtOAc = ethyl acetate; RT = room temperature; THF = tetrahydrofuran; DCM = dichloromethane; MeOH = methyl alcohol; EtOH = ethyl alcohol; LHMDs = lithium bis(trimethylsilyl)amide; m-CPBA = meta-Chloroperoxybenzoic acid; TFA = trifluoroacetic acid; LC-MS = Liquid Chromatography/Mass Spectrometry; NMR = nuclear magnetic resonance; HPLC = high pressure liquid chromatography; MPLC = medium pressure liquid chromatography; SFC = Supercritical Fluid Chromatography

10

15

General Experimental details

Analytical Methods

20

Liquid Chromatography-Mass Spectrometry

Method 1

25

LC-MS was performed on a Waters 2795 Alliance HT HPLC with Waters 2996 Diode Array Detector coupled to a Micromass ZQ, single quadrupole mass spectrometer using a Phenomenex Luna C18 (2) column (5 μ m, 100 x 4.6mm plus guard cartridge) with a linear gradient of 5-95% acetonitrile/water (with 0.1% formic acid in each mobile phase) within 3.5 minutes and held at 95% for 2.0 minutes.

Method 2

LC-MS was performed on a Waters 2795 Alliance HT HPLC with Waters 2996

Diode Array Detector coupled to a Micromass ZQ, single quadrupole mass spectrometer using a Waters Xterra MS C18 column (5 μ m, 100 x 4.6mm plus guard cartridge) being initially held at 5% acetonitrile/water (with 10mM ammonium bicarbonate in the aqueous mobile phase) for 0.5 minutes, followed by a linear gradient of 5-95% within 3.5 minutes
5 and then held at 95% for 1.5 minutes.

Method 3

UPLC-MS was performed on a Waters Acquity I-Class with Waters Diode Array Detector coupled to a Waters SQD2 single quadrupole mass spectrometer using an Waters HSS C18 column (1.8 μ m, 100 $\times$ 2.1 mm) being initially held at 5% acetonitrile/water (with
10 0.1% formic acid in each mobile phase) for 1.2 minutes, followed by a linear gradient of 5-100% within 3.5 minutes and then held at 100% for 1.5 minutes (F = 0.5 mL/min).

Method 4

UPLC-MS was performed on a Waters Acquity I-Class with Waters Diode Array Detector coupled to a Waters SQD2 single quadrupole mass spectrometer using a Waters
15 BEH Shield RP18 column (1.7 μ m, 100 $\times$ 2.1 mm) being initially held at 5% acetonitrile/water (with 10 mM ammonium bicarbonate in each mobile phase) for 1.2 minutes, followed by a linear gradient of 5-100% within 3.5 minutes and then held at 100% for 1.5 minutes (F = 0.5 mL/min).

**Supercritical Fluid Chromatography--Mass Spectrometry analytical
20 conditions**

Method 5

SFC-MS was performed on a Waters/Thar SFC systems with Waters SQD using a Chiralpak IA column with a 55% *iso*-propyl alcohol/CO₂ (with 0.1% diethylamine) isocratic run at 5 mL/min, 120 Bar backpressure, 40 °C column temperature.

25 Method 6

SFC-MS was performed on a Waters/Thar SFC systems with Waters SQD using a Chiralpak IA column with a 5% methanol/CO₂ (with 0.1% diethylamine) isocratic run at 5 mL/min, 120 bar backpressure, 40 °C column temperature.

Method 7

SFC-MS was performed on a Waters/Thar SFC systems with Waters SQD using a YMC Amylose-C column with a 55% *iso*-propyl alcohol/CO₂ (with 0.1% diethylamine) isocratic run at 5 mL/min, 120 Bar backpressure, 40 °C column temperature.

5

Method 8

SFC-MS was performed on a Waters/Thar SFC systems with Waters SQD using a YMC Amylose-C column with a 50% *iso*-propyl alcohol/CO₂ (with 0.1% diethylamine) isocratic run at 5 mL/min, 120 Bar backpressure, 40 °C column temperature.

Method 9

10

SFC-MS was performed on a Waters/Thar SFC systems with Waters SQD using a YMC Amylose-C column with a 40% ethanol/CO₂ (with 0.1% diethylamine) isocratic run at 5 mL/min, 120 Bar backpressure, 40 °C column temperature.

Method 10

15

SFC-MS was performed on a Waters/Thar SFC systems with Waters SQD using a YMC Amylose-C column with a 40% *iso*-propyl alcohol/CO₂ (with 0.1% diethylamine) isocratic run at 5 mL/min, 120 Bar backpressure, 40 °C column temperature.

Chiral HPLC- analytical conditions**Method 11**

20

Chiral HPLC was performed on an Agilent 1200 series HPLC system using a Chiralpak IA column with 50% *iso*-propyl alcohol /heptane (with 0.1% diethylamine) at 1 mL/min.

Method 12

25

Chiral HPLC was performed on an Agilent 1200 series HPLC system using a YMC Cellulose-C column with 50% *iso*-propyl alcohol /heptane (with 0.1% diethylamine) at 1 mL/min.

NMR

¹H Nuclear magnetic resonance (NMR) spectroscopy was carried out using a Bruker instrument operating at 400 MHz using the stated solvent at around room temperature

unless otherwise stated. In all cases, NMR data were consistent with the proposed structures. Characteristic chemical shifts (δ) are given in parts-per-million using conventional abbreviations for designation of major peaks: e.g. s, singlet; d, doublet; t, triplet; q, quartet; dd, doublet of doublets; dt, doublet of triplets; m, multiplet; br, broad.

5 **Preparative reverse-phase HPLC conditions**

Preparative HPLC purification was performed by reverse phase HPLC using a Waters Fractionlynx preparative HPLC system (2525 pump, 2996/2998 UV/VIS detector, 2767 liquid handler) or an equivalent HPLC system such as a Gilson Trilution UV directed system. The Waters 2767 liquid handler acted as both auto-sampler and fraction collector.

10 The columns used for the preparative purification of the compounds were a Waters Sunfire OBD Phenomenex Luna Phenyl Hexyl or Waters Xbridge Phenyl at 10 μ m 19 $\times$ 150 mm or Waters CSH Phenyl Hexyl, 19 $\times$ 150, 5 μ m column.

Appropriate focused gradients were selected based on acetonitrile and methanol solvent systems under either acidic or basic conditions.

15 The modifiers used under acidic/basic conditions were formic acid or trifluoroacetic acid (0.1% V/V) and ammonium bicarbonate (10 mM) respectively.

The purification was controlled by Waters Fractionlynx software through monitoring at 210-400 nm, and triggered a threshold collection value at 260 nm and, when using the Fractionlynx, the presence of target molecular ion as observed under APi
20 conditions. Collected fractions were analysed by LCMS (Waters Acquity systems with Waters SQD).

Compound preparation

Where the preparation of starting materials is not described, these are commercially available, known in the literature, or readily obtainable by those skilled in the art using
25 standard procedures. Where it is stated that compounds were prepared “analogously” or “similarly” to earlier examples or intermediates, it will be appreciated by the skilled person that the reaction time, number of equivalents of reagents and temperature can be modified for each specific reaction and that it may be necessary or desirable to employ different

work-up or purification techniques.

Flash chromatography refers to silica gel chromatography and is carried out using an Isolera MPLC system (manufactured by Biotage); pre-packed silica gel cartridges (supplied by Biotage); or using conventional glass column chromatography.

5 In the procedures that follow, after each starting material, reference to a compound number may be provided. This is provided merely for assistance to the skilled chemist. The starting material may not necessarily have been prepared from the batch referred to.

Many of the compounds described in the following Examples have been prepared from stereochemically pure starting materials, for example 95% enantiomeric excess (ee).

10 The stereochemistry of the compounds in the Examples, where indicated, has been assigned on the assumption that absolute configuration at resolved stereogenic centers of starting materials is maintained throughout any subsequent reaction conditions.

Compounds isolated as single diastereoisomers whose absolute configuration at stereogenic center (2) in general formula (I) or (I') was not determined, are herebelow
15 referred to as *Single Diastereoisomers* without mention in their chemical name of absolute configuration for the unknown stereogenic centre.

Chiral separation protocol

The diastereomeric separation of compounds was achieved either by chiral High Performance Liquid Chromatography (HPLC) using a Gilson Trilution preparative HPLC
20 system (322 pump, 155 UV/VIS, GX281 liquid handler and fraction collector) or by Supercritical Fluid Chromatography (SFC) using a Waters Thar Prep100 preparative SFC system (P200 CO₂ pump, 2545 modifier pump, 2998 UV/VIS detector, 2767 liquid handler with Stacked Injection Module). The Waters 2767 liquid handler acted as both auto-sampler and fraction collector.

25 The column used for the preparative purification of the compounds was a Diacel Chiralpak IA/IB/IC, a Phenomenex Lux Cellulose-4, an YMC Amylose-C or an YMC Cellulose-C at 5 μ m 250 $\times$ 20 – 21.2 mm ID.

Appropriate isocratic methods were selected based on methanol, ethanol or

isopropanol solvent systems under un-modified or basic conditions.

The standard SFC method used was modifier, CO₂, 100 mL/min, 120 Bar backpressure, 40 °C column temperature. The standard HPLC method used was modifier, heptane, 5 mL/min and room temperature.

5 The modifier used under basic conditions was diethylamine (0.1% V/V). The modifier used under acidic conditions was either formic acid (0.1% V/V) or trifluoroacetic acid (0.1% V/V).


The SFC purification was controlled by Waters Fractionlynx software through monitoring at 210-400 nm and triggered at a threshold collection value, typically 260 nm.

10 Collected fractions were analysed by SFC (Waters/Thar SFC systems with Waters SQD). The fractions that contained the desired product were concentrated by vacuum centrifugation.

HPLC purification was controlled by Gilson Trilution software monitoring two wavelengths and triggered at a threshold collection value, typically 260 nm. Collected
15 fractions were analysed by HPLC (Agilent 1200 series HPLC system). The fractions that contained the desired product were concentrated by vacuum centrifugation.

Intermediate 1

[(1S)-2-(3,5-Dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-formyl-thiophene-2-carboxylate



20


A stirred solution of 5-formyl-2-thiophenecarboxylic acid (400 mg, 2.56 mmol) in dichloromethane (20 mL) was added with (1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethanol (compound I-1/A described in the co-pending international application n. PCT/EP2013/075520) (881 mg, 2.56 mmol) followed by 4-
25 (dimethylamino)-pyridine (156 mg, 1.28 mmol) and *N*-(3-dimethylaminopropyl)-*N'*-

ethylcarbodiimide hydrochloride (983 mg, 5.12 mmol). The resulting mixture was stirred at room temperature for 18 h. The reaction was partitioned between dichloromethane and saturated aqueous sodium bicarbonate solution. The organic layer was washed with brine, passed through a hydrophobic frit and the solvent was removed *in vacuo*. The crude material was purified by silica gel column chromatography, eluting with 0-100% EtOAc in DCM, to afford the title compound (488 mg, 39%) as yellow oil.

¹H NMR (400 MHz, CDCl₃): δ 9.97 (s, 1 H), 8.15 (s, 2 H), 7.81 (d, J = 3.6 Hz, 1 H), 7.72 (d, J = 3.6 Hz, 1 H), 7.03-6.99 (m, 2 H), 6.87 (d, J = 8.7 Hz, 1 H), 6.26 (dd, J = 4.4, 10.0 Hz, 1 H), 3.91 (s, 3 H), 3.88 (s, 3 H), 3.72 (dd, J = 10.0, 14.0 Hz, 1 H), 3.33 (dd, J = 4.4, 14.0 Hz, 1 H).


LCMS (Method 2): [MH⁺] = 482 at 3.38 min.

The following intermediate was synthesised *via* a similar method to Intermediate 1:

Structure	Intermediate number	Analytical Data
	Intermediate 2	¹ H NMR (400MHz, CDCl ₃): δ 10.09 (s, 1 H), 8.45 (s, 2 H), 8.19 (d, J = 8.1 Hz, 2 H), 7.94 (d, J = 8.1 Hz, 2 H), 7.08 (dd, J = 8.2, 2.1 Hz, 1 H), 6.99 (d, J = 2.0 Hz, 1 H), 6.87 (d, J = 8.3 Hz, 1 H), 6.36 (dd, J = 10.0, 4.3 Hz, 1 H), 3.90 (s, 3 H), 3.88 (s, 3 H), 3.83 (dd, J = 13.7, 10.0 Hz, 1 H), 3.43 (dd, J = 13.7, 4.4 Hz, 1 H). LCMS (Method 1): [MH ⁺] = 460 at 4.45 min.

Intermediate 3

2-(Benzyloxycarbonylamino)-2-phenyl-propanoic acid




A solution of 2-amino-2-phenylpropanoic acid (1.65 g, 10 mmol) in a mixture of THF and water (1:1, 180 mL) at 0 °C was added with benzyl chloroformate (1.42 mL, 10 mmol) and NaOH (1 N, 10 mL, 10 mmol) simultaneously. The mixture was stirred at 0 °C for 1 hour. The ice bath was removed and the reaction mixture was stirred at room temperature for 16 hours. The solvent was removed in vacuo and the aqueous layer was basified with 10% aqueous sodium hydroxide to pH≈13. The aqueous phase was back-extracted with EtOAc (2 × 50 mL). The aqueous phase was then acidified with 6 N HCl to pH≈3-2. EtOAc (200 mL) was added and the reaction mixture stirred at room temperature for 16 hours. The layers were separated and the aqueous phase re-extracted with EtOAc (2 × 50 mL). The combined organic fractions were dried over MgSO₄, filtered and the solvent was removed in vacuo to yield the title compound (1.34 g, 48%) as a clear oil. This oil was used in the next step without further purification.

¹H NMR (400 MHz, CDCl₃): δ 7.46 (d, J = 7.4 Hz, 2 H), 7.37-7.25 (m, 8 H), 6.86-6.54 (m, 1 H), 6.23-6.01 (m, 1 H),)HH5.14-4.99 (m, 2 H), 2.02 (s, 3 H).).

LCMS (Method 1): [MH⁺] = 300 at 2.68 min.

Intermediate 4:

(R)-Quinuclidin-3-yl 2-amino-2-phenyl-propanoate



A solution of 2-(benzyloxycarbonylamino)-2-phenyl-propanoic acid (1.34 g, 4.8 mmol) and N,N'-dicyclohexylcarbodiimide (1.2 g, 4.8 mmol) in dry THF (24 mL) was stirred at room temperature for 1 hour. 1-Hydroxybenzotriazole hydrate (777 mg, 5.8 mmol) and (R)-quinuclidin-3-ol (1.03 g, 8.11 mmol) were subsequently added and the resulting slurry was stirred at room temperature for 3 days. After this time, the reaction mixture was filtered through a pad of Celite® and the solvent was removed in vacuo. The residue was partitioned between EtOAc (100 mL) and saturated aqueous Na₂CO₃ (2 × 50 mL). The aqueous fraction was back-extracted with EtOAc (2 x 50 mL). The combined

organic fractions were washed with brine (30 mL), dried over MgSO₄, filtered and the solvent was removed in vacuo. The solid residue (844 mg) obtained was used in the next step without further purification. This residue was taken up in EtOAc (15 mL) and ammonium formate (750 mg, 11.91 mmol) and 10% Pd/C (550 mg) were added. The mixture was heated to 60 °C for 16 hours. After cooling the slurry to room temperature and filtration through a pad of Celite®, the solids were washed with EtOAc (150 mL).


The solvent was removed in vacuo to yield the title compound as a mixture of two diastereoisomers (606 mg, 46% yield over two steps). The white solid was used in the next step without further purification.

¹H NMR (400 MHz, CDCl₃): δ 7.51-7.47 (m, 2 H), 7.37-7.31 (m, 2 H), 7.30-7.21 (m, 2 H), 7.19-7.10 (m, 1 H), 4.81-4.76 (m, 1 H), 3.24-3.10 (m, 1 H), 2.79-2.61* (m, 4 H), 2.61-2.43† (m, 4 H), 2.30-2.18 (m, 1 H), 2.02-1.96* (m, 1 H), 1.95-1.87† (m, 1 H), 1.73† (s, 3 H), 1.72* (s, 3 H), 1.70-1.58 (m, 1 H), 1.58-1.42 (m, 1 H), 1.42-1.20 (m, 1 H), 1.18-1.01 (m, 1 H), † and * refer to different isomers (arbitrarily assigned).

LCMS (Method 2): [MH⁺] = 275 at 2.70 min.

Intermediate 5:

[(3*R*)-Quinuclidin-3-yl] 1-(tert-butoxycarbonylamino)indane-1-carboxylate







A solution of 1-(tert-butoxycarbonylamino)indane-1-carboxylic acid (1.0 g, 3.6 mmol) and N,N'-dicyclohexylcarbodiimide (1.1 g, 4.3 mmol) in THF (24 mL) was stirred at room temperature for 1 hour. 1-Hydroxybenzotriazole hydrate (577 mg, 4.3 mmol) and (*R*)-quinuclidin-3-ol (914 mg, 7.2 mmol) were subsequently added and the resulting slurry was stirred at room temperature for 3 days. After this time, the reaction mixture was filtered through a pad of Celite® and the solvent was removed in vacuo. The residue was partitioned between EtOAc (100 mL) and saturated aqueous Na₂CO₃ (2 × 50 mL), and the

resulting aqueous fractions were re-extracted with EtOAc (2 x 50 mL). The combined organic fractions were washed with brine (30 mL), dried over Na₂SO₄, filtered and the solvent was removed in vacuo to yield a white foam (1.18 g, 85 % yield) which was used in the next step without further purification.


¹H NMR (400 MHz, CDCl₃): δ 7.36-7.25 (m, 2 H), 7.25-7.18 (m, 2 H), 5.47-5.40 (m, 1 H), 4.79-4.74 (m, 1 H), 3.20-3.05 (m, 4 H), 2.81-2.63 (m, 4 H), 2.54† (d, J = 14.9 Hz, 1 H), 2.47-2.39* (m, 1 H), 2.02-1.96† (m, 1 H), 1.94-1.89* (m, 1 H), 1.75-1.72† (m, 1 H), 1.70-1.58 (m, 1 H), 1.55-1.46 (m, 2 H), 1.43 (bs, 9 H), 1.17-0.99* (m, 1 H), † and * refer to different isomers (arbitrarily assigned). LCMS (Method 2): [MH⁺] = 287 at 3.08 min.

The following intermediates were synthesised *via* a similar method :

Structure	Intermediate number	Analytical Data
	Intermediate 12	LCMS (Method 1): [MH ⁺] = 349 at 2.51 min.
	Intermediate 13	LCMS (Method 2): [MH ⁺] = 375 at 3.65 min.
	Intermediate 14	LCMS (Method 2): [MH ⁺] = 349 at 3.63 min.
	Intermediate 15	LCMS (Method 2): [MH ⁺] = 361 at 3.30 min.

Intermediate 16

(1-Methyl-4-piperidyl)-2-amino-2-phenyl-acetate *bis* hydrochloride




A solution of (1-methyl-4-piperidyl) 2-(*tert*-butoxycarbonylamino)-2-phenylacetate (2.60 g, 7.47 mmol) in 4 N HCl in dioxane (9.34 mL, 37.3 mmol) was stirred

at room temperature for 18 hours. The solvent was removed by evaporation under reduced pressure and co-evaporated with diethyl ether to give the title compound as a yellow gum (2.80 g, quantitative yield).

LCMS (Method 2): [MH⁺] = 249 at 1.93 min.


5 The following intermediate was synthesised *via* a similar method:

Structure	Intermediate number	Analytical Data
	Intermediate 17	LCMS (Method 2): [MH ⁺] = 261 at 2.05 min.

Intermediate 6

Isopropyl 2-(*tert*-butylsulfinylamino)-2-phenyl-propanoate

10





A solution of isopropyl-2-*tert*-butylsulfinylimino-2-phenyl-acetate (0.71 g, 2.40 mmol) in dry DCM (15 mL) was stirred under N₂ and cooled to -78 °C. A 3.0M solution of methyl magnesium bromide in diethyl ether (1.2 mL, 3.60 mmol) was added drop wise over a period of 10 minutes then stirred at -78 °C for 1 hour. The mixture was quenched with drop wise addition of saturated NaHCO₃ solution (6 mL) followed by water (10 mL) and extracted with DCM (2x30 mL). The organic extracts were combined and washed with brine (20 mL) and filtered through a phase separator. The solvent was removed in vacuo to give the title compound as a mobile yellow oil (0.68 g, 91%).

20 LCMS (Method 2): [MH⁺] = 312 at 2.87 min.

The following intermediate was synthesised *via* a similar method ::


Structure	Intermediate number	Analytical Data

	Intermediate 7	¹ H NMR (400 MHz, CDCl ₃): δ 7.48-7.27 (m, 5 H), 5.09-5.02 (m, 1 H), 4.72 (s, 1 H), 2.55-2.30 (m, 2 H), 1.24-1.20 (m, 15 H), 0.97-0.88 (m, 3 H).
---	----------------	---

Intermediate 8**2-(*tert*-Butylsulfinylamino)-2-phenyl-propanoic acid**


- 5 A solution of isopropyl 2-(*tert*-butylsulfinylamino)-2-phenyl-propanoate (1.06 g, 3.41mmol) and aqueous NaOH (2M, 5.1 mL, 10.2mmol) in methanol (5 mL) was heated at 50 °C for 3 hours. After cooling to room temperature the mixture was diluted with EtOAc (30 mL) and extracted with water (2x 20 mL). The aqueous phase was acidified to pH 5/6 with 1M HCl and removed the solvent *in vacuo* to give the title compound as a light brown solid (0.54 g). This was used in the next step without further purification. LCMS (Method
- 10 2): [MH⁺] = 270 at 2.05 and 2.18 min.

The following intermediate was synthesised *via* a similar method:

Structure	Intermediate number	Analytical Data
	Intermediate 9	¹ H NMR (400 MHz, DMSO): δ 7.42 - 7.13 m, 5 H), 5.82 (d, J = 7.5 Hz, 1 H), 2.37-2.26 (m, 1 H), 2.13-2.03 (m, 1 H), 1.03 (s, 9 H), 0.91-0.80 (m, 3 H).

Intermediate 10


- 15 **(1-Methyl-4-piperidyl) 2-(*tert*-butylsulfinylamino)-2-phenyl-propanoate**



A mixture of 2-(*tert*-butylsulfinylamino)-2-phenyl-propanoic acid (0.27 g, 1.0mmol), 1-methylpiperidin-4-ol (0.23 g, 2.0mmol), *N,N'*-dicyclohexylcarbodiimide (0.41 g, 2.0 mmol) and 1-hydroxybenzotriazole hydrate (0.27 g, 2.0 mmol) in THF (25 mL) was stirred at room temperature for 42 hours. The mixture was filtered through a bed of Celite® and the solvent was removed in *vacuo*. The residue was partitioned between EtOAc (25 mL) and 2M aqueous Na₂CO₃ (2 × 50 mL), The organic phase was washed with brine (20 mL), separated and filtered through a phase separator. The solvent was removed in *vacuo* to yield a brown solid (0.23 g, 62 % yield) which was used in the next step without further purification.


10 LCMS (Method 2): [MH⁺] = 367 at 2.67 min.

The following intermediate was synthesised *via* a similar method:

Structure	Intermediate number	Analytical Data
	Intermediate 11	¹ H NMR (400 MHz, CDCl ₃): δ 7.68 (br s, 1 H), 7.48-7.42 (m, 2 H), 7.37-7.29 (m, 3 H), 4.88-4.81 (m, 1 H), 3.32-3.09 (m, 1 H), 2.98-2.33 (m, 6 H), 2.20-1.86 (m, 2 H), 1.81-1.39 (m, 4 H), 1.25 (s, 9 H), 1.0-0.85 (m, 3 H)

Example 1

15 [(1*S*)-2-(3,5-Dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-oxo-1-phenyl-2-[(3*R*)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate formate salt




A suspension of (R)-quinuclidin-3-yl 2-amino-2-phenylpropanoate (160 mg, 0.58 mmol) in CH₃CN (5 mL) was added with [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-formylthiophene-2-carboxylate (234 mg, 0.49 mmol) followed by acetic acid (0.07 mL, 1.0 mmol) and oven dried 3Å crushed molecular sieves (200 mg). The resulting slurry was stirred at room temperature for 2 days. NaBH(OAc)₃ (513 mg, 2.23 mmol) was added in one portion and the reaction mixture was stirred at room temperature for a further 3.5 hours. The reaction mixture was filtered over a pad of Celite® and the solids were washed with EtOAc (100 mL). The filtrate was then concentrated in vacuo and the residue was purified by preparative HPLC, using formic acid as acidic modifiers of the chromatographic solvent, to give the title compound (1:1 mixture of diastereoisomers) as a pale yellow solid (128 mg, 35% yield).

¹H NMR (400 MHz, CD₃CN): δ 8.27 (s, 1 H), 8.19 (s, 2 H), 7.69 (dd, J = 3.8, 1.8 Hz, 1 H), 7.59-7.54 (m, 2 H), 7.42 (t, J = 7.5 Hz, 2 H), 7.38-7.30 (m, 1 H), 7.08-7.02 (m, 2 H), 6.98-6.93 (m, 2 H), 6.18 (dd, J = 9.6, 4.5 Hz, 1 H), 4.98-4.90 (m, 1 H), 3.94-3.86 (m, 2 H), 3.84 (m, 3 H), 3.84-3.79 (m, 4 H), 3.68 (dd, J = 14.2, 9.6 Hz, 1 H), 3.35 (dd, J = 13.9, 4.8 Hz, 3 H), 2.90-2.79 (m, 3 H), 2.80-2.60 (m, 2 H), 2.09-2.00 (m, 1 H), 1.86-1.70 (m, 2 H), 1.72† (s, 3 H), 1.70* (s, 3 H), 1.82-1.48 (m, 2 H), 1.55-1.33 (m, 1 H), † and * refer to different isomers (arbitrarily assigned). LCMS (Method 1): [MH⁺] = 740 at 2.67 min.


The following compound was synthesised as mixture of diastereoisomers via the same method:

Structure	Reference	Analytical Data
[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-	Example 2	¹ H NMR (400 MHz, CD ₃ CN): δ 8.25 (s, 1 H), 8.18 (s, 2 H), 8.02

dimethoxyphenyl)ethyl] 4-[[[1-methyl-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxyethyl]amino]methyl]benzoate formate salt 		(d, J = 8.1 Hz, 2 H), 7.56 (d, J = 7.8 Hz, 2 H), 7.53 (d, J = 8.1 Hz, 2 H), 7.40 (t, J = 7.6 Hz, 2 H), 7.32 (t, J = 7.2 Hz, 1 H), 7.10 (d, J = 2.0 Hz, 1 H), 7.06 (dd, J = 8.3, 2.0 Hz, 1 H), 6.95 (d, J = 8.2 Hz, 1 H), 6.26 (dd, J = 9.6, 4.5 Hz, 1 H), 4.91-4.86 (m, 1 H), 3.84 (s, 3 H), 3.82 (s, 3 H), 3.76 (d, J = 3.8 Hz, 2 H), 3.71 (dd, J = 14.2, 9.6 Hz, 1 H), 3.37 (dd, J = 14.1, 4.6 Hz, 1 H), 3.23 (ddd, J = 14.6, 8.3, 2.4 Hz, 1 H), 2.88-2.72 (m, 3 H), 2.71-2.58 (m, 2 H), 2.04-1.98 (m, 1 H), 1.77-1.71 (m, 1 H), 1.70 (s, 3 H), 1.66-1.55 (m, 2 H), 1.44-1.35 (m, 1 H), NH not observed. LCMS (Method 1): [MH⁺] = 734 at 2.47 min.
---	--	--

Example 3

[(3R)-Quinuclidin-3-yl] 1-[[4-[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethoxy] carbonylphenyl] methylamino]indane-1-carboxylate



5

A solution of [(3R)-quinuclidin-3-yl] 1-(*tert*-butoxycarbonylamino)indane-1-carboxylate (190 mg, 0.49 mmol) in EtOAc (5 mL) was added with a solution of hydrogen chloride in dioxane, (4 N, 5 mL, 20 mmol) at 0 °C. The resulting mixture was stirred at room temperature for 18 hours. The solvent was removed in vacuo and the crude residue was diluted in EtOH (8 mL). [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 4-formylbenzoate (113 mg, 0.33 mmol), Et₃N (0.19 mL, 1.33 mmol) and acetic acid (0.03 mL, 0.5 mmol) were added and the mixture stirred at room


10

temperature for 1 hour. NaBH₃CN (42 mg, 0.66 mmol) was added in one portion and stirring was continued for 18 hours. The solvent was removed and the residue was partitioned between EtOAc (20 mL) and water (20 mL), and the aqueous fraction was back-extracted with EtOAc (2 x 20 mL). The combined organic fractions were washed with 0.2 N HCl (2 X 20 mL). The combined aqueous phases were saturated with NaCl and back-extracted with CHCl₃ (4 X 50 mL). The combined organic phases were passed through a hydrophobic frit and the solvent was removed in vacuo. The residue was then purified by preparative HPLC to yield the title compound as a white solid (149 mg, 61% over two steps).

¹H NMR (400 MHz, DMSO) : δ 8.56 (s, 2 H), 7.91 (d, J=7.8 Hz, 2 H), 7.50-7.43 (m, 3 H), 7.31-7.20 (m, 3 H), 7.06-6.99 (m, 2 H), 6.97 (d, J=8.3 Hz, 1 H), 6.21 (dd, J=4.2, 9.7 Hz, 1 H), 4.70-4.64 (m, 1 H), 3.78 (s, 3 H), 3.75 (s, 3 H), 3.73-3.70 (m, 2 H), 3.62 (dd, J=9.7, 14.1 Hz, 1 H), 3.32-3.22 (m, 2 H), 3.06-2.96 (m, 3 H), 2.68-2.55 (m, 4 H), 2.40 (d, J=14.6 Hz, 1 H), 2.31 (d, J=16.6 Hz, 1 H), 2.15 (td, J=8.5, 13.0 Hz, 1 H), 1.86-1.83† (m, 1 H), 1.81-1.76* (m, 1 H), 1.59-1.50 (m, 1 H), 1.45-1.40 (m, 1 H), 1.27-1.14 (m, 1 H) † and * refer to different isomers (arbitrarily assigned). LCMS (Method 1): [MH⁺] = 746 at 2.46 min.

The following compound was synthesised as mixture of diastereoisomers *via* the same method:

Structure	Reference	Analytical Data
[(1S)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-[(3R)-quinuclidin-3-yl]oxycarbonylindan-1-yl]amino]methyl]thiophene-2-carboxylate	Example 4	¹ H NMR (400 MHz, CD ₃ CN) : δ 8.19† (s, 2 H), 8.19* (s, 2 H), 7.67† (s, 1 H), 7.66* (s, 1 H), 7.38 (dd, J=3.4, 7.5 Hz, 1 H), 7.35-7.28 (m, 2 H), 7.27-7.22 (m, 1 H), 7.06-7.01 (m, 2 H), 6.96-6.93 (m, 2 H), 6.17 (dd, J=4.5, 9.6 Hz, 1 H), 4.80-4.75 (m, 1 H), 3.99-3.86 (m, 2 H), 3.83† (s, 3 H), 3.83* (s, 3 H), 3.82 (s, 3 H), 3.70-3.63 (m, 2 H), 3.34 (dd, J = 4.5, 14.1 Hz, 1 H), 3.17-3.03 (m, 3 H), 2.77-2.60 (m, 5 H), 2.55 (d, J=15.7 Hz, 1 H), 2.44 (d, J=14.8 Hz, 1 H),


		2.00-1.91 (m, 1 H), 1.88-1.81 (m, 1 H), 1.69-1.59 (m, 1 H), 1.55-1.49 (m, 1 H), 1.37-1.27 (m, 1 H) † and * refer to different isomers (arbitrarily assigned). LCMS (Method 1): [MH⁺] = 752 at 2.63 min.
---	--	---

5

10

Example 5

[(1S)-2-(3,5-Dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenylethyl]amino]methyl]thiophene-2-carboxylate



15

A solution of (1-methyl-4-piperidyl) 2-(*tert*-butylsulfinylamino)-2-phenylpropanoate (0.30 g, 0.84mmol) in 4 N hydrogen chloride in dioxane (1.1 mL, 4.18mmol) was stirred at room temperature for 2 hours. The solvent was removed *in vacuo* and the



residue was mixed with a solution of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-formylthiophene-2-carboxylate (0.27 g, 0.56 mmol) in DCM (20 mL). Et₃N (0.23 mL, 1.68mmol) followed by acetic acid (0.048 mL, 0.84mmol) were added and the mixture was stirred at room temperature for 24 hours. NaBH(OAc)₃ (0.70 g, 3.36 mmol) was added and the reaction mixture was stirred at room temperature for a further 5 days. The mixture was diluted with DCM (10 mL), washed with saturated NaHCO₃ solution (2 x 15 mL) and saturated NaCl (15 mL), filtered through a phase separator and the solvent was removed in *vacuo*. Purification by preparative HPLC gave the title compound as a pale yellow solid, (0.16 g, 39%).

¹H NMR (400 MHz, DMSO): δ 8.63 (s, 2 H), 8.62 (s, 2 H), 8.22 (s, 2 H), 7.73 (d, J = 3.8 Hz, 1 H), 7.54-7.51 (m, 2 H), 7.43 (dd, J = 7.6, 7.6 Hz, 2 H), 7.34 (dd, J = 7.2, 7.2 Hz, 1 H), 7.09-7.03 (m, 4 H), 6.19 (dd, J = 4.3, 9.9 Hz, 1 H), 4.84 (dd, J = 3.4, 3.4 Hz, 1 H), 3.94-3.86 (m, 2 H), 3.84 (s, 3 H), 3.81 (s, 3 H), 3.68-3.60 (m, 1 H), 3.36 (d, J = 18.4 Hz, 1 H), 2.49-2.36 (m, 2 H), 2.29 (d, J = 47.3 Hz, 2 H), 2.19 (s, 3 H), 2.13 (s, 1 H), 1.81 (d, J = 5.1 Hz, 2 H), 1.64 (s, 3 H), 1.62-1.57 (m, 2 H).

LCMS (Method 1): [MH⁺] = 728 at 2.59 min.


The following compound was synthesised as mixture of diastereoisomers via the same method:

Structure	Reference	Analytical Data
[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-phenyl-1-[(3R)-quinuclidin-3-yl]oxycarbonyl-propyl]amino]methyl]thiophene-2-carboxylate	Example 6	¹ H NMR (400 MHz, CD ₃ CN): δ 8.19 (s, 2 H), 7.68 (dd, J = 2.0, 3.8 Hz, 1 H), 7.58-7.54 (m, 2 H), 7.44-7.38 (m, 2 H), 7.36-7.30 (m, 1 H), 7.09-7.03 (m, 2 H), 6.98-6.94 (m, 2 H), 6.21-6.16 (m, 1 H), 4.82-4.79 (m, 1 H), 3.87-3.80 (m, 6 H), 3.80-3.65 (m, 4 H), 3.35 (dd, J = 4.5, 14.1 Hz, 1 H), 3.19-3.08 (m, 1 H), 2.93-2.86 (m, 1 H), 2.72-2.61 (m, 4 H), 2.58-2.52 (m, 1 H), 2.51-2.38 (m, 1 H), 2.37-2.28 (m, 1 H), 2.14-2.08 (m, 3 H),

		1.92-1.89 (m, 1 H), 1.70-1.41 (m, 5 H), 1.33-1.24 (m, 1 H), 0.90-0.82 (m, 3 H). LCMS (Method 2): [MH ⁺] = 754 at 3.50 min.
[(1<i>S</i>)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(2-dimethylaminoethoxy)carbonyl]indan-1-yl]amino]methyl]thiophene-2-carboxylate 	Example 17	¹ H NMR (400 MHz, CDCl ₃): δ 8.13 ^{*or†} (s, 2 H), 8.12 ^{*or†} (s, 2 H), 7.64-7.58 (m, 1 H), 7.34-7.17 (m, 4 H), 7.00-6.94 (m, 2 H), 6.89-6.81 (m, 2 H), 6.21 (dd, J = 4.2, 9.5 Hz, 1 H), 4.25-4.23 (m, 2 H), 3.91-3.82 (m, 9 H), 3.65 (dd, J = 9.7, 13.8 Hz, 1 H), 3.30 (dd, J = 4.3, 13.9 Hz, 1 H), 3.10-3.08 (m, 2 H), 2.78-2.68 (m, 1 H), 2.57-2.47 (m, 2 H), 2.22-2.19 (m, 7 H). * and † refer to different isomers. LCMS (Method 2): [MH ⁺] = 714 at 3.77 min.

Example 18

[(1*S*)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-[(1-methyl-4-piperidyl)oxycarbonyl]indan-1-yl]amino]methyl]thiophene-2-carboxylate




A solution of (1-methyl-4-piperidyl)-1-(*tert*-butoxycarbonylamino)indane-1-carboxylate (0.25 g, 0.67 mmol) in 4 N HCl in dioxane (1.0 mL, 4.0 mmol) was stirred at room temperature for 18 hours. The solvent was removed *in vacuo* to give a white solid.

The previously obtained solid was dissolved in acetonitrile (5 mL) and [(1*S*)-2-(3,5-dichloro-1-oxidopyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-formylthiophene-2-carboxylate (322 mg, 0.67 mmol) was added followed by acetic acid (90 μ L, 1.34 mmol). The resulting mixture was stirred at room temperature for 18 hours. The solvent was removed *in vacuo* and the residue azeotroped with toluene. The residue was taken up with acetonitrile (5 mL) and sodium triacetoxyborohydride (457 mg, 2.07 mmol) was added. The resulting mixture was stirred at room temperature for 18 hours. Additional sodium triacetoxyborohydride (457 mg, 2.07 mmol) was added and the stirring was maintained at room temperature for 4 hours. The solvent was removed *in vacuo* and the residue was taken up in water (30 mL) and ethyl acetate (100 mL). The layers were separated and the aqueous phase was extracted with ethyl acetate (2 $\times$ 20 mL). The combined organic phases were passed through a hydrophobic frit and the solvent was removed *in vacuo*. Purification of the crude material by preparative HPLC afforded the title compound (130 mg, 26%) as a white solid.


¹H NMR (400 MHz, CDCl₃): δ 8.14^{*or†} (s, 2 H), 8.13^{*or†} (s, 2 H), 7.63 (d, J = 3.8 Hz, 1 H), 7.30-7.27 (m, 3 H), 7.24-7.18 (m, 1 H), 7.00-6.95 (m, 2 H), 6.88 (d, J = 3.8 Hz, 1 H), 6.84 (d, J = 8.1 Hz, 1 H), 6.21 (dd, J = 4.4, 9.5 Hz, 1 H), 4.85-4.81 (m, 1 H), 3.91^{*or†} (s, 3 H), 3.90^{*or†} (s, 3 H), 3.87 (s, 3 H), 3.88-3.81 (m, 2 H), 3.65 (dd, J = 10.0, 13.8 Hz, 1 H), 3.30 (dd, J = 4.5, 13.9 Hz, 1 H), 3.11 (dd, J = 7.2, 7.2 Hz, 2 H), 2.76-2.68 (m, 1 H), 2.50-2.42 (m, 1 H), 3.35-2.14 (m, 3 H), 2.22 (s, 3 H), 1.92-1.85 (m, 1 H), 1.83-1.77 (m, 1 H), 1.75-1.68 (m, 1 H), 1.56-1.52 (m, 2 H). NH not visible, † and * refer to different isomers (arbitrarily assigned). LCMS (Method 1): [MH⁺] = 740 at 2.59 min.

Example 19

[(1*S*)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl]-5-[[[1-(hydroxymethyl)-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenyl-ethyl]amino]methyl]thiophene-2-carboxylate



Step 1: Preparation of (1-methyl-4-piperidyl) 2-benzylideneamino-2-phenyl-acetate




5

To a solution of (1-methyl-4-piperidyl) 2-amino-2-phenyl-acetate, *bis* hydrochloride (650 mg, 2.02 mmol) in dry DCM (10 mL) was added Et₃N (0.57 mL, 4.05 mmol) followed by MgSO₄. After stirring at room temperature for one hour, benzaldehyde (0.21 mL, 2.02 mmol) was added and the mixture was stirred at room temperature for 36 hours. The mixture was filtered, the solid was washed through with DCM, water (50 mL) was added to the filtrate, the biphasic mixture was filtered through a phase separator and the solvent was removed *in vacuo* to give the title compound as a yellow oil (586 mg, 86%). LCMS (Method 2): [MH⁺] = 337 at 3.25 min.

Step 2: Preparation of (1-methyl-4-piperidyl) 2-amino-3-hydroxy-2-phenyl-propanoate *bis* hydrochloride


15



To a mixture of (1-methyl-4-piperidyl) 2-benzylideneamino-2-phenyl-acetate (0.28 g, 0.83 mmol) and *para*-formaldehyde (50 mg, 1.67 mmol) in dry dioxane (10 mL) was added DBU (0.15 mL, 1.0 mmol) and the mixture was stirred at room temperature for 3 hours. The solvent was removed by evaporation, the residue was dissolved in EtOAc (20 mL) and washed with water (2 × 10 mL). The aqueous phases were combined, back-extracted with EtOAc (10 mL). The organic extracts were combined, filtered through a phase separator frit and evaporated to dryness under reduced pressure to give a yellow gum.

The obtained gum was treated with 1 N HCl (1 mL) in THF (1 mL) and stirred at room temperature for 18 hours. The solvent was removed *in vacuo*, the residue was taken up in 1 N HCl (3 mL) and washed with EtOAc (2 × 5 mL), the combined organic phases were back-extracted with 1 N HCl (3 mL). The combined aqueous extracts were evaporated to dryness under reduced pressure, co-evaporated with CH₃CN to give the title compounds as an off-white solid (155 mg, 53%). LCMS (Method 2): [MH⁺] = 279 at 1.84 min.

Step 3: Preparation of [(1*S*)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenyl-ethyl]amino]methyl]thiophene-2-carboxylate




To a stirred mixture of [(1*S*)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-formylthiophene-2-carboxylate (0.21 g, 0.43 mmol) and (1-methyl-4-piperidyl) 2-amino-3-hydroxy-2-phenyl-propanoate *bis* hydrochloride (0.15 g, 0.43 mmol) in DCM (5 mL) at room temperature was added Et₃N (0.12 mL, 0.85 mmol) followed by AcOH (24 µL, 0.43 mmol). The resulting mixture was stirred at room temperature for 20 hours. Sodium triacetoxyborohydride (0.27 g, 1.28 mmol) was added and the mixture was stirred at room temperature for 24 hours. Additional sodium triacetoxyborohydride (0.27 g, 1.28 mmol) and AcOH (24 µL, 0.43 mmol) were added and the mixture was stirred at room temperature for 72 hours. The reaction mixture was diluted with DCM (10 mL) and washed with saturated NaHCO₃ solution (2 × 20 mL), brine (10 mL), the organic phase was filtered through a phase separator frit and the solvent was removed *in vacuo*. Purification by preparative HPLC gave the title compound as an off-white solid (24.9 mg, 7.7%).

¹H NMR (400 MHz, CD₃CN): δ 8.20 (s, 2H), 7.68 (d, J=3.8 Hz, 1H), 7.52 - 7.48 (m, 2H), 7.40 (dd, J=7.3, 7.3 Hz, 2H), 7.34 (dd, J=7.2, 7.2 Hz, 1H), 7.08 - 7.02 (m, 2H), 6.97 - 6.92 (m, 2H), 6.18 (dd, J=4.7, 9.7 Hz, 1H), 4.91 - 4.87 (m, 1H), 4.17 (d, J=11.1 Hz, 1H), 3.95 - 3.94 (m, 2H), 3.84 (s, 3H), 3.82 (s, 3H), 3.80 - 3.72 (m, 1H), 3.68 (dd, J=9.6, 14.1 Hz, 1H), 3.39 - 3.32 (m, 1H), 3.25 - 3.13 (m, 2H), 2.51 - 2.31 (m, 2H), 2.40 - 2.20 (m, 2H), 2.15 (s, 3H), 1.86 - 1.81 (m, 2H), 1.69 - 1.60 (m, 2H).

LCMS (Method 1): [MH⁺] = 744 at 2.7 min.

Example 20

[(1*S*)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-oxo-1-phenyl-2-[(3*R*)-quinuclidin-3-yl]oxy-ethyl]-methyl-amino]methyl]thiophene-2-carboxylate




To a stirred mixture of [(1*S*)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-formylthiophene-2-carboxylate (480 mg, 1.0 mmol) and [(3*R*)-quinuclidin-3-yl] 2-amino-2-phenyl-acetate, *bis* hydrochloride (480 mg, 1.4 mmol) in
 5 trifluoroethanol (10 mL) at room temperature was added Et₃N (0.4 mL, 2.9 mmol) followed by AcOH (115 μ L, 2.0 mmol) and the mixture was stirred at room temperature for 72 hours. The solvent was removed *in vacuo* and the residue was azeotroped with toluene (3 $\times$ 10 mL) to give a yellow gum. This was suspended in dioxane (10 mL) and *para*-formaldehyde (300 mg, 10.0 mmol) was added. The resulting suspension was stirred
 10 at room temperature and DBU (0.18 mL, 1.2 mmol) was added and the mixture was stirred at room temperature for 3 hours. The solvent was removed *in vacuo*. The obtained residue was suspended in dry MeCN (20 mL) and sodium triacetoxyborohydride (1.05 g, 5.0 mmol) was added. The resulting mixture was stirred at room temperature for 18 hours. The solvent was removed *in vacuo* and the residue was partitioned between EtOAc (35 mL) and 1 N
 15 HCl (30 mL). The aqueous phase was washed again with EtOAc (35 mL) then basified with solid NaHCO₃ and extracted with EtOAc (2 $\times$ 25 mL). The combined organic extracts were filtered through a phase separator frit and the solvent was removed *in vacuo*. Purification by preparative HPLC gave the title compound as an off-white solid (73 mg, 9.5%).

¹H NMR (400 MHz, CD₃CN): δ 8.21^{*or†} (s, 2 H), 8.21^{*or†} (s, 2 H), 7.69^{*or†} (d, *J* = 1.2 Hz, 1 H), 7.68^{*or†} (d, *J* = 1.4 Hz, 1 H), 7.63-7.58 (m, 2 H), 7.45-7.39 (m, 2 H), 7.37-7.33 (m, 1 H), 7.10-7.04 (m, 2 H), 6.99-6.95 (m, 2 H), 6.19 (ddd, *J* = 1.8, 4.4, 9.7 Hz, 1 H), 4.98-4.91 (m, 1 H), 4.29-4.16 (m, 2 H), 4.08-3.93 (m, 2 H), 3.86 (s, 3 H), 3.83 (s, 3 H), 3.70 (dd, *J* = 9.7, 14.0 Hz, 1 H), 3.40-3.33 (m, 1 H), 3.24-3.16 (m, 1 H), 3.10-2.89 (m, 1 H), 2.78-2.60 (m, 5 H), 2.38^{*or†} (s, 3 H), 2.37^{*or†} (s, 3 H), 2.14-2.11 (m, 1 H), 1.85-1.53 (m, 3 H), 1.42-

1.29 (m, 1 H). † and * refer to different isomers (arbitrarily assigned). LCMS (Method 4): [MH⁺] = 770 at 3.4 min.

Example 21

[(1*S*)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl]-5-[[[1-(hydroxymethyl)-2-oxo-1-phenyl-2-[(3*R*)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate




To a stirred mixture of [(1*S*)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-formylthiophene-2-carboxylate (480 mg, 1.0 mmol) and [(3*R*)-quinuclidin-3-yl] 2-amino-2-phenyl-acetate, *bis* hydrochloride (480 mg, 1.4 mmol) in trifluoroethanol (10 mL) at room temperature was added Et₃N (0.4 mL, 2.9 mmol) followed by AcOH (115 μL, 2.0 mmol) and the mixture was stirred at room temperature for 72 hours. The solvent was removed *in vacuo* and the residue was azeotroped with toluene (3 × 10 mL) to give a yellow gum. This was suspended in dioxane (10 mL) and *para*-formaldehyde (300 mg, 10.0 mmol) was added. The resulting suspension was stirred at room temperature and DBU (0.18 mL, 1.2 mmol) was added and the mixture was stirred at room temperature for 3 hours. The solvent was removed *in vacuo* and the residue was partitioned between EtOAc (25 mL) and water (20 mL). The aqueous phase was further extracted with EtOAc (3 × 20 mL) and CHCl₃ (2 × 25 mL). The combined organic extracts were filtered through a phase separator frit and the solvent was removed *in vacuo*. The residue was dissolved in dry CH₃CN (10 mL), sodium triacetoxymethylborohydride (0.63 g, 3.0 mmol) was added and the mixture was stirred at room temperature overnight. The solvent was removed *in vacuo*, the residue was partitioned between EtOAc (40 mL) and 1N HCl

(35 mL). The aqueous phase was further washed with EtOAc (35 mL) then basified with solid NaHCO₃ and extracted with EtOAc (20 mL) and CHCl₃ (3 × 30 mL). The combined organic extracts were filtered through a phase separator frit and the solvent was removed *in vacuo*. Purification by preparative HPLC gave the title compound as a light brown solid (150 mg, 20%).


¹H NMR (400 MHz, CD₃CN): δ 8.20^{*or†} (s, 2H), 8.20^{*or†} (s, 2 H), 7.68^{*or†} (d, J = 2.0 Hz, 1H), 7.67^{*or†} (d, J = 2.0 Hz, 1 H), 7.55-7.50 (m, 2 H), 7.44-7.34 (m, 3 H), 7.09-7.02 (m, 2 H), 6.98-6.92 (m, 2 H), 6.18 (dd, J = 4.5, 9.9 Hz, 1 H), 4.88-4.83 (m, 1 H), 4.27-4.21 (m, 1 H), 3.99-3.90 (m, 2 H), 3.85 (s, 3 H), 3.83 (s, 3 H), 3.77 (d, J = 13.9 Hz, 1 H), 3.69 (dd, J = 10.9, 14.7 Hz, 1 H), 3.39-3.32 (m, 1 H), 3.21-3.09 (m, 1 H), 2.75-2.45 (m, 5 H), 1.95-1.87 (m, 1 H), 1.71-1.63 (m, 2 H), 1.58-1.44 (m, 1 H), 1.38-1.28 (m, 1 H). † and * refer to different isomers (arbitrarily assigned) OH and NH not seen. LCMS (Method 4): [MH⁺] = 756 at 3.31 min.

Example 22

[(1*S*)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(methoxymethyl)-2-oxo-1-phenyl-2-[(3*R*)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate



Step 1: Preparation of [(3*R*)-quinuclidin-3-yl] 2-amino-3-methoxy-2-phenylpropanoate *bis* hydrochloride



5


Di-*tert*-butyl dicarbonate (1.12 g, 5.12 mmol) was added to a solution of 2-amino-3-methoxy-2-phenylpropanoic acid (0.50 g, 2.56 mmol) and 2 N sodium hydroxide (3.85 mL, 7.70 mmol) in 2:1 dioxane:water (10 mL). The mixture was stirred for 18 hours then the solvents were removed under reduced pressure. The residue was cooled in an ice/water bath then 1M KHSO₄ was added carefully to pH 2. The mixture was extracted with ethyl acetate (3 × 10 mL). The combined organic extracts were dried over MgSO₄, filtered then concentrated under reduced pressure. The obtained residue was dissolved in THF (15 mL) and 3-(*R*)-quinuclidinol (0.36 g, 2.81 mmol), hydroxybenzotriazole (0.38 g, 2.81 mmol) and dicyclohexylcarbodiimide (0.58 g, 2.81 mmol) were added. The mixture was stirred for 8 days then filtered through celite ®. The filtrate was concentrated under reduced pressure then partitioned between ethyl acetate (20 mL) and saturated aqueous NaHCO₃ (20 mL). The combined organic extracts were washed with water, dried over MgSO₄, filtered then concentrated under reduced pressure to give an off-white foam which was suspended in 4 N HCl in dioxane (10 mL). The mixture was stirred for 20 hours then concentrated under reduced pressure to give the crude product as an off-white solid (0.56

20

g) in 58% yield.

LCMS (Method 3): $[MH^+] = 304$ at 1.02 min.

Step 2: Preparation of [(1*S*)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(methoxymethyl)-2-oxo-1-phenyl-2-[(3*R*)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate




[(1*S*)-2-(3,5-Dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-formylthiophene-2-carboxylate (282 mg, 0.59 mmol), and pyridine (0.066 mL, 0.82 mmol) was added to a solution of [(3*R*)-quinuclidin-3-yl] 2-amino-3-methoxy-2-phenylpropanoate *bis* hydrochloride (0.31 g, 0.82 mmol) in ethanol (10 mL) at room temperature. The suspension was heated to 60 °C then the resulting solution was stirred for 10 minutes. Sodium cyanoborohydride (0.052 g, 0.82 mmol) was added and the mixture was stirred for 4 hours then concentrated under reduced pressure. The residue was partitioned between ethyl acetate (5 mL) and 2 N hydrochloric acid (5 mL). The aqueous was extracted with ethyl acetate (5 mL) then the combined organics were washed with 2 N hydrochloric acid (5 mL). The combined aqueous layers were basified with saturated aqueous NaHCO₃ then the mixture was extracted with ethyl acetate (3 × 10 mL). The combined organic extracts were dried over MgSO₄, filtered then concentrated under reduced pressure to give the crude product which was purified by preparative HPLC. This gave two fractions of differing purity which contained formic acid. Both fractions were individually basified with saturated aqueous NaHCO₃ then extracted with ethyl acetate (3 × 10 mL). The combined organic extracts were washed with 2 N hydrochloric acid (5 mL). The combined aqueous layers

were basified with saturated aqueous NaHCO_3 then extracted with ethyl acetate (3×10 mL). Both fractions were determined to contain only the desired product so were combined. The organics were dried over MgSO_4 , filtered then concentrated under reduced pressure to give the desired product as a white solid (28 mg, 4%).

- 5 ^1H NMR (400 MHz, DMSO): δ 8.57 (s, 2 H), 7.67 (dd, $J = 3.2, 3.2$ Hz, 1 H), 7.52 (dd, $J = 8.0, 8.0$ Hz, 2 H), 7.44-7.33 (m, 3 H), 7.07-6.96 (m, 4 H), 6.19-6.13 (m, 1 H), 4.83-4.76 (m, 1 H), 4.03-3.97 (m, 1 H), 3.90 (dt, $J = 4.8, 12.2$ Hz, 2 H), 3.81 (s, 3 H), 3.78 (s, 3 H), 3.65-3.58 (m, 1 H), 3.34^{*or†} (s, 3 H), 3.32^{*or†} (s, 3 H), 3.14-3.05 (m, 1 H), 2.73-2.58 (m, 3 H), 2.42-2.34^{*or†} (m, 1 H), 1.95-1.89^{*or†} (m, 1 H), 1.88-1.83^{*or†} (m, 1 H), 1.63-1.58 (m, 2 H),
10 1.54-1.48 (m, 1 H), 1.35-1.19 (m, 2 H). † and * refer to different isomers (arbitrarily assigned). NH not seen. LCMS (Method 3): $[\text{MH}^+] = 770$ at 2.82 min.

Example 7 and Example 8

- Single diastereoisomers of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate
15 (Ex. 7, diast 1 and Ex. 8, diast 2)



- Purification of the 1:1 mixture of diastereoisomers of Example 1 by chiral preparative SFC afforded the single diastereoisomers. The absolute configurations of these
20 were not determined.

Title compound (Example 7, single diastereoisomer 1) was obtained as a white solid (15 mg, 31%).



¹H NMR (400 MHz, CD₃CN): δ 8.20 (s, 2 H), 7.68 (d, J = 3.8 Hz, 1 H), 7.59-7.52 (m, 2 H), 7.41 (t, J = 7.6 Hz, 2 H), 7.33 (t, J = 7.3 Hz, 1 H), 7.07 (d, J = 2.0 Hz, 1 H), 7.04 (dd, J = 8.2, 2.1 Hz, 1 H), 6.97-6.92 (m, 2 H), 6.18 (dd, J = 9.6, 4.5 Hz, 1 H), 4.87-4.82 (m, 1 H), 3.88 (s, 2 H), 3.84 (s, 3 H), 3.82 (s, 3 H), 3.68 (dd, J = 14.1, 9.7 Hz, 1 H), 3.35 (dd, J = 14.1, 4.6 Hz, 1 H), 3.24-3.13 (m, 1 H), 3.05-2.85 (m, 1 H), 2.82-2.64 (m, 4 H), 2.60 (d, J = 13.7 Hz, 1 H), 1.93-1.89 (m, 1 H), 1.71 (s, 3 H), 1.69-1.62 (m, 1 H), 1.70-1.44 (m, 2 H), 1.38-1.27 (m, 1 H). LCMS (Method 1): [MH⁺] = 740 at 2.63 min.

Title compound (Example 8, single diastereoisomer 2) was obtained as a white solid (14 mg, 30%).

¹H NMR (400 MHz, CD₃CN): δ 8.20 (s, 2 H), 7.69 (d, J = 3.8 Hz, 1 H), 7.59-7.54 (m, 2 H), 7.44-7.39 (m, 2 H), 7.37-7.30 (m, 1 H), 7.08-7.02 (m, 2 H), 7.00-6.94 (m, 2 H), 6.18 (dd, J = 9.7, 4.5 Hz, 1 H), 5.00-4.94 (m, 1 H), 3.91 (dd, J = 20.8, 15.3 Hz, 2 H), 3.84 (s, 3 H), 3.82 (s, 3 H), 3.73-3.63 (m, 1 H), 3.34 (dd, J = 13.8, 4.0 Hz, 1 H), 3.38-3.27 (m, 1 H), 3.05-2.80 (m, 4 H), 2.80-2.67 (m, 2 H), 2.12-2.07 (m, 1 H), 1.86-1.74 (m, 1 H), 1.84-1.55 (m, 2 H), 1.69 (s, 3 H), 1.53-1.42 (m, 1 H). LCMS (Method 2): [MH⁺] = 740 at 3.94 min.



Compounds reported in the table herebelow were obtained as single diastereoisomers according to the procedure described in Examples 7 and 8 by chiral preparative SFC or chiral preparative HPLC.

Structure	Reference	Analytical Data
[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 4-[[[1-methyl-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]benzoate	Example 9 (Diastereoisomer 1)	¹ H NMR (400 MHz, CD ₃ CN): δ 8.15 (s, 2 H), 7.99 (d, J = 8.1 Hz, 2 H), 7.53 (dd, J = 7.9, 1.4 Hz, 2 H), 7.49 (d, J = 8.0 Hz, 2 H), 7.39-7.34 (m, 2 H), 7.29 (tt, J = 7.3, 1.5 Hz, 1 H), 7.07 (d, J

		= 2.1 Hz, 1 H), 7.03 (dd, J = 8.2, 2.1 Hz, 1 H), 6.92 (d, J = 8.2 Hz, 1 H), 6.23 (dd, J = 9.6, 4.6 Hz, 1 H), 4.81-4.74 (m, 1 H), 3.81 (s, 3 H), 3.79 (s, 3 H), 3.71 (d, J = 5.0 Hz, 2 H), 3.71-3.65 (m, 1 H), 3.35 (dd, J = 14.1, 4.6 Hz, 1 H), 3.12 (ddd, J = 14.6, 8.2, 2.3 Hz, 1 H), 2.71-2.57 (m, 4 H), 2.50 (dt, J = 14.7, 2.6 Hz, 1 H), 1.86 (dd, J = 6.7, 3.7 Hz, 1 H), 1.68 (s, 3 H), 1.65-1.57 (m, 1 H), 1.54-1.45 (m, 2 H), 1.30-1.21 (m, 3 H). LCMS (Method 1): [MH⁺] = 734 at 2.50 min.
[(3R)-quinuclidin-3-yl] 1-[[4-[(1S)-2-(3,5-dichloro-1-oxo-1H-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethoxy]carbonylphenyl]methylamino]indane-1-carboxylate 	Example 10 (Diastereoisomer 1)	¹H-NMR (400 MHz, DMSO) : δ 8.56 (s, 2 H), 7.91 (d, J = 8.3 Hz, 2 H), 7.48 (d, J = 8.4 Hz, 2 H), 7.45 (d, J = 7.1 Hz, 1 H), 7.30-7.20 (m, 3 H), 7.06-6.99 (m, 2 H), 6.97 (d, J = 8.3 Hz, 1 H), 6.21 (dd, J = 4.3, 9.6 Hz, 1 H), 4.67-4.64 (m, 1 H), 3.78 (s, 3 H), 3.75 (s, 3 H), 3.71 (d, J = 5.8 Hz, 2 H), 3.62 (dd, J = 9.7, 14.0 Hz, 1 H), 3.36-3.30 (m, 1 H), 3.29-3.22 (m, 1 H), 3.06 - 2.92 (m, 3 H), 2.68-2.53 (m, 4 H), 2.53-2.42 (m, 1 H), 2.30 (d, J = 14.9 Hz, 1 H), 2.15 (td, J = 8.1, 13.1 Hz, 1 H), 1.87-1.82 (m, 1 H), 1.60-1.50 (m, 2 H), 1.47-1.40 (m, 1 H), 1.30-1.21 (m, 1 H). LCMS (Method 2): [MH⁺] = 746 at 3.28 min.

(continue)

[(3R)-quinuclidin-3-yl] 1-[[4-[(1S)-2-(3,5-dichloro-1-oxo-1H-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethoxy]carbonylphenyl]methylamino]indane-1-carboxylate	Example 11 (Diastereoisomer 2)	¹H-NMR (400 MHz, DMSO) : δ 8.56 (s, 2 H), 7.91 (d, J = 8.3 Hz, 2 H), 7.48 (d, J = 8.1 Hz, 2 H), 7.44 (d, J = 7.5 Hz, 1 H), 7.31-7.20 (m, 3 H), 7.06 - 7.00 (m, 2 H), 6.97 (d, J = 8.3 Hz, 1 H), 6.21 (dd, J = 4.2, 9.7 Hz, 1 H),
---	---	--



		H), 4.71 - 4.65 (m, 1 H), 3.78 (s, 3 H), 3.75 (s, 3 H), 3.73-3.68 (m, 2 H), 3.62 (dd, J = 9.8, 14.0 Hz, 1 H), 3.37-3.28 (m, 1 H), 3.28-3.20 (m, 1 H), 3.08-2.97 (m, 3 H), 2.69-2.54 (m, 5 H), 2.40 (d, J = 14.1 Hz, 1 H), 2.15 (td, J = 8.2, 13.1 Hz, 1 H), 1.81-1.77 (m, 1 H), 1.58 - 1.49 (m, 1 H), 1.47-1.37 (m, 2 H), 1.25-1.15 (m, 1 H). LCMS (Method 2): [MH⁺] = 746 at 3.27 min.
[(1S)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-[(3R)-quinuclidin-3-yl]oxycarbonyl]indan-1-yl]amino]methyl]thiophene-2-carboxylate 	Example 12 (Diastereoisomer 1)	¹H NMR (400 MHz, DMSO): δ 8.61 (s, 2 H), 7.71 (d, J = 3.8 Hz, 1 H), 7.48 (d, J = 7.1 Hz, 1 H), 7.37-7.26 (m, 3 H), 7.07-7.02 (m, 4 H), 6.18 (dd, J = 4.3, 9.6 Hz, 1 H), 4.77-4.72 (m, 1 H), 3.96-3.86 (m, 2 H), 3.81 (d, J = 6.8 Hz, 6 H), 3.75 (dd, J = 7.6, 7.6 Hz, 2 H), 3.68-3.57 (m, 1 H), 3.38-3.30 (m, 1 H), 3.15-2.99 (m, 3 H), 2.74-2.59 (m, 4 H), 2.48-2.35 (m, 1 H), 2.22-2.13 (m, 1 H), 1.92 (d, J = 2.5 Hz, 1 H), 1.67-1.47 (m, 3 H), 1.39-1.27 (m, 1 H). LCMS (Method 1): [MH⁺] = 752 at 2.61 min.

(continue)

5




10



[(1S)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-[(3R)-quinuclidin-3-yl]oxycarbonyl]indan-1-yl]amino]methyl]thiophene-2-carboxylate	Example 13 (Diastereoisomer 2)	¹H NMR (400 MHz, DMSO): δ 8.55 (s, 2 H), 7.65 (d, J = 3.8 Hz, 1 H), 7.40 (d, J = 6.8 Hz, 1 H), 7.30-7.18 (m, 3 H), 7.01-6.95 (m, 4 H), 6.12 (dd, J = 4.3, 9.6 Hz, 1 H), 4.70-4.66
--	---	---

		(m, 1 H), 3.84 (d, J = 7.3 Hz, 2 H), 3.76 (s, 3 H), 3.73 (s, 3 H), 3.66 (dd, J = 7.6, 7.6 Hz, 1 H), 3.60-3.51 (m, 1 H), 3.31-3.21 (m, 1 H), 3.07-2.96 (m, 3 H), 2.67-2.56 (m, 4 H), 2.41 (d, J = 14.4 Hz, 1 H), 2.15-2.06 (m, 1 H), 1.80-1.78 (m, 1 H), 1.56-1.41 (m, 3 H), 1.26-1.15 (m, 1 H). LCMS (Method 1): [MH⁺] = 752 at 2.60 min.
[(1S)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenylethyl]amino]methyl]thiophene-2-carboxylate 	Example 14 (Diastereoisomer 1)	¹H NMR (400 MHz, CDCl₃) 8.15 (2H, s), 7.64 (1H, d, J = 3.8 Hz), 7.52-7.48 (2H, m), 7.36 (2H, dd, J = 7.7, 7.7 Hz), 7.32-7.28 (1H, m), 7.02-6.98 (2H, m), 6.91-6.84 (2H, m), 6.23 (1H, dd, J = 4.4, 9.5 Hz), 4.90-4.89 (1H, m), 3.92 (3H, s), 3.89 (3H, s), 3.86 (2H, s), 3.68 (1H, dd, J = 10.0, 13.8 Hz), 3.33 (1H, dd, J = 4.5, 13.9 Hz), 2.66-2.32 (3H, m), 2.23 (5H, s), 1.89-1.86 (2H, m), 1.75-1.70 (5H, m); LCMS (Method 1): [MH⁺] = 728 at 2.59 min.

(continue)



[(1S)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenylethyl]amino]methyl]thiophene-2-carboxylate	Example 15 (Diastereoisomer 2)	¹H NMR (400 MHz, CDCl₃) 8.14 (2H, s), 7.64 (1H, d, J = 3.8 Hz), 7.52-7.48 (2H, m), 7.36 (2H, dd, J = 7.5, 7.5 Hz), 7.31-7.27 (1H, m), 6.99 (2H, d, J = 24.2 Hz), 6.90-6.84 (2H, m), 6.23 (1H, dd, J = 4.4,
---	---	---

		9.7 Hz), 4.89 (1H, dd, J = 3.4, 7.2 Hz), 3.91 (3H, s), 3.88 (3H, s), 3.85 (2H, s), 3.67 (1H, dd, J = 9.9, 13.9 Hz), 3.32 (1H, dd, J = 4.7, 14.0 Hz), 2.62-2.60 (1H, m), 2.45-2.45 (2H, m), 2.23 (5H, s), 1.94-1.86 (2H, m), 1.72 (5H, s); LCMS (Method 1): [MH ⁺] = 728 at 2.61 min.
[(1<i>S</i>)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(2-dimethylaminoethoxy)carbonyl]indan-1-yl]amino]methyl]thiophene-2-carboxylate 	Example 23 (Diastereoisomer 1)	¹H NMR (400 MHz, CDCl₃): δ 8.12 (s, 2 H), 7.62 (d, J = 1.8 Hz, 1 H), 7.30-7.24 (m, 4 H), 7.01-6.94 (m, 2 H), 6.88-6.86 (m, 2 H), 6.25-6.18 (m, 1 H), 4.28-4.22 (m, 2 H), 3.89-6.31 (m, 9 H), 3.68-3.60 (m, 1 H), 3.33-3.31 (m, 1 H), 3.11-3.09 (m, 2 H), 2.75-2.71 (m, 1 H), 2.57-2.55 (m, 2 H), 2.26-2.21 (m, 7 H). LCMS (Method 2): [MH⁺] = 714 at 3.32 min.
[(1<i>S</i>)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(2-dimethylaminoethoxy)carbonyl]indan-1-yl]amino]methyl]thiophene-2-carboxylate 	Example 24 (Diastereoisomer 2)	¹H NMR (400 MHz, CDCl₃): δ 8.13 (s, 2 H), 7.61-7.60 (m, 1 H), 7.36-7.31 (m, 4 H), 6.98-6.93 (m, 2 H), 6.90-6.83 (m, 2 H), 6.21-6.20 (m, 1 H), 4.27-4.24 (m, 2 H), 3.89-3.81 (m, 9 H), 3.68-3.62 (m, 1 H), 3.32-3.29 (m, 1 H), 3.11-3.09 (m, 2 H), 2.75-2.71 (m, 1 H), 2.59-2.56 (m, 2 H), 2.24-2.20 (m, 7 H). LCMS (Method 2): [MH⁺] = 714 at 3.32 min.
(continue)		
[(1<i>S</i>)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-[(1-methyl-4-piperidyl)oxycarbonyl]indan-1-yl]amino]methyl]thiophene-2-carboxylate	Example 25 (Diastereoisomer 1)	¹H NMR (400 MHz, CDCl₃): δ 8.12 (s, 2 H), 7.62 (d, J = 3.8 Hz, 1 H), 7.29-7.26 (m, 3 H), 7.23-7.20 (m, 1 H), 7.00-6.95 (m, 2 H), 6.88 (d, J = 3.7 Hz, 1 H), 6.84 (d, J = 8.8 Hz, 1 H), 6.22 (dd, J = 4.7,

		9.5 Hz, 1 H), 4.86-4.81 (m, 1 H), 3.90 (s, 3 H), 3.88-3.85 (m, 2 H), 3.87 (s, 3 H), 3.65 (dd, J = 9.7, 14.0 Hz, 1 H), 3.31 (dd, J = 4.8, 13.9 Hz, 1 H), 3.10 (t, J = 7.1 Hz, 2 H), 2.72 (dt, J = 6.1, 10.0 Hz, 1 H), 2.50-2.45 (m, 1 H), 2.35-2.28 (m, 1 H), 2.27-2.18 (m, 2 H), 2.22 (s, 3 H), 1.92-1.87 (m, 1 H), 1.83-1.62 (m, 4 H), NH not visible. LCMS (Method 1): [MH⁺] = 740 at 2.55 min..
[(1S)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(1-methyl-4-piperidyl)oxycarbonyl]indan-1-yl]amino]methyl]thiophene-2-carboxylate 	Example 26 (Diastereoisomer 2)	¹H NMR (400 MHz, CDCl₃): δ 8.12 (s, 2 H), 7.62 (d, J = 3.8 Hz, 1 H), 7.29-7.26 (m, 3 H), 7.23-7.20 (m, 1 H), 7.00-6.95 (m, 2 H), 6.88 (d, J = 3.7 Hz, 1 H), 6.84 (d, J = 8.8 Hz, 1 H), 6.22 (dd, J = 4.7, 9.5 Hz, 1 H), 4.86-4.81 (m, 1 H), 3.90 (s, 3 H), 3.88-3.85 (m, 2 H), 3.87 (s, 3 H), 3.65 (dd, J = 9.7, 14.0 Hz, 1 H), 3.31 (dd, J = 4.8, 13.9 Hz, 1 H), 3.10 (t, J = 7.1 Hz, 2 H), 2.72 (dt, J = 6.1, 10.0 Hz, 1 H), 2.50-2.45 (m, 1 H), 2.35-2.28 (m, 1 H), 2.27-2.18 (m, 2 H), 2.22 (s, 3 H), 1.92-1.87 (m, 1 H), 1.83-1.62 (m, 4 H), NH not visible. LCMS (Method 1): [MH⁺] = 740 at 2.55 min.



(continue)

[(1S)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-phenyl-1-[(3R)-quinuclidin-3-yl]oxycarbonyl-propyl]amino]methyl]thiophene-2-carboxylate	Example 27 (Diastereoisomer 1)	¹H NMR (400 MHz, CD₃CN): δ 8.20 (s, 2 H), 7.68 (d, J = 3.8 Hz, 1 H), 7.57-7.54 (m, 2 H), 7.40 (dd, J = 7.6, 7.6 Hz, 2 H), 7.34-7.30 (m, 1 H), 7.08-7.02 (m, 2 H), 6.97-6.94 (m, 2 H), 6.18 (dd, J = 4.5,
---	---	---

		9.6 Hz, 1 H), 4.82-4.78 (m, 1 H), 3.84 (s, 3 H), 3.82 (s, 3 H), 3.77-3.65 (m, 3 H), 3.35 (dd, J = 4.5, 14.1 Hz, 1 H), 3.14 (ddd, J = 2.1, 8.2, 14.7 Hz, 1 H), 2.88 (s, 1 H), 2.74-2.51 (m, 5 H), 2.38-2.28 (m, 1 H), 2.16-2.05 (m, 1 H), 1.87-1.83 (m, 1 H), 1.67-1.43 (m, 3 H), 1.31-1.23 (m, 1 H), 0.87 (t, J = 7.4 Hz, 3 H). LCMS (Method 1): [MH⁺] = 754 at 2.85 min.
[(1S)-2-(3,5-dichloro-1-oxo-1,2,3,4-tetrahydropyridin-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-phenyl-1-[(3R)-quinuclidin-3-yl]oxycarbonylpropyl]amino]methyl]thiophene-2-carboxylate 	Example 28 (Diastereoisomer 2)	¹H NMR (400 MHz, CD₃CN): δ 8.19 (s, 2 H), 7.69 (d, J = 4.6 Hz, 1 H), 7.59-7.53 (m, 2 H), 7.44-7.38 (m, 2 H), 7.36-7.29 (m, 1 H), 7.09-7.02 (m, 2 H), 6.99-6.92 (m, 2 H), 6.19 (dd, J = 4.5, 9.8 Hz, 1 H), 4.82-4.77 (m, 1 H), 3.85 (s, 3 H), 3.82 (s, 3 H), 3.77 (t, J = 8.2 Hz, 2 H), 3.73-3.63 (m, 1 H), 3.35 (dd, J = 4.7, 14.1 Hz, 1 H), 3.14-3.06 (m, 1 H), 2.90 (t, J = 8.0 Hz, 1 H), 2.66 (dt, J = 12.0, 20.7 Hz, 3 H), 2.59-2.47 (m, 1 H), 2.47-2.39 (m, 1 H), 2.35-2.24 (m, 1 H), 1.91-1.87 (m, 1 H), 1.70-1.46 (m, 3 H), 1.33-1.22 (m, 2 H), 0.85 (t, J = 7.3 Hz, 3 H). LCMS (Method 2): [MH⁺] = 754 at 3.55 min.



(continue)

[(1S)-2-(3,5-dichloro-1-oxo-1,2,3,4-tetrahydropyridin-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenylethyl]amino]methyl]thiophene-2-carboxylate	Example 29 (Diastereoisomer 1)	¹H NMR (400 MHz, CD₃CN): δ 8.20 (s, 2 H), 7.67 (d, J = 3.8 Hz, 1 H), 7.52-7.48 (m, 2 H), 7.43-7.37 (m, 2 H), 7.36-7.32 (m, 1 H), 7.08-7.02 (m, 2 H), 6.97-6.91 (m, 2 H), 6.18 (dd, J = 4.5, 9.6 Hz, 1
--	---	--

		H), 4.92-4.85 (m, 1 H), 4.17 (d, J = 10.9 Hz, 1 H), 4.00-3.90 (m, 2 H), 3.84 (s, 3 H), 3.83 (s, 3 H), 3.79-3.64 (m, 2 H), 3.35 (dd, J = 4.5, 14.1 Hz, 1 H), 3.11-3.11 (m, 2 H), 2.51-2.44 (m, 1 H), 2.32 (s, 1 H), 2.23 (d, J = 10.9 Hz, 2 H), 2.14 (s, 3 H), 1.90-1.79 (m, 2 H), 1.72-1.58 (m, 2 H). LCMS (Method 2): [MH⁺] = 744 at 3.13 min.
[(1<i>S</i>)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenylethyl]amino]methyl]thiophene-2-carboxylate 	Example 30 (Diastereoisomer 2)	¹H NMR (400 MHz, CD₃CN): δ 8.20 (s, 2 H), 7.68 (d, J = 4.7 Hz, 1 H), 7.52-7.48 (m, 2 H), 7.43-7.38 (m, 2 H), 7.37-7.33 (m, 1 H), 7.08-7.02 (m, 2 H), 6.97-6.93 (m, 2 H), 6.19 (dd, J = 5.7, 9.3 Hz, 1 H), 4.92-4.85 (m, 1 H), 4.21-4.14 (m, 1 H), 3.97-3.93 (m, 2 H), 3.84 (s, 3 H), 3.83 (s, 3 H), 3.80-3.73 (m, 1 H), 3.72-3.65 (m, 1 H), 3.36 (dd, J = 5.2, 14.5 Hz, 1 H), 3.24-3.03 (m, 2 H), 2.73 (d, J = 145.3 Hz, 1 H), 2.51-2.40 (m, 1 H), 2.35 (d, J = 42.4 Hz, 1 H), 2.18-2.05 (m, 4 H), 1.90-1.79 (m, 2 H), 1.70-1.58 (m, 2 H). LCMS (Method 2): [MH⁺] = 744 at 3.12 min.



(continue)

[(1<i>S</i>)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-oxo-1-phenyl-2-[(3<i>R</i>)-quinuclidin-3-yl]oxyethyl]amino]methyl]thiophene-2-carboxylate	Example 31 (Diastereoisomer 1)	¹H NMR (400 MHz, CD₃CN): δ 8.21 (s, 2 H), 7.68 (d, J = 3.8 Hz, 1 H), 7.55-7.51 (m, 2 H), 7.45-7.40 (m, 2 H), 7.39-7.33 (m, 1 H), 7.09-7.03 (m, 2 H), 6.98-6.94 (m, 2 H), 6.18 (dd, J = 4.4, 9.7 Hz, 1 H), 4.86-4.82 (m, 1 H),
---	---	--

		4.23 (d, $J = 10.9$ Hz, 1 H), 3.99-3.77 (m, 3 H), 3.85 (s, 3 H), 3.83 (s, 3 H), 3.69 (dd, $J = 9.6, 14.1$ Hz, 1 H), 3.36 (dd, $J = 4.5, 14.1$ Hz, 1 H), 3.19-3.08 (m, 2 H), 2.75-2.44 (m, 5 H), 2.01-1.89 (m, 1 H), 1.70-1.62 (m, 2 H), 1.58-1.49 (m, 1 H), 1.37-1.27 (m, 1 H). NH not seen. LCMS (Method 3): $[MH^+] = 756$ at 2.74 min.
[(1<i>S</i>)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-oxo-1-phenyl-2-[(3<i>R</i>)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate 	Example 32 (Diastereoisomer 2)	^1H NMR (400 MHz, CD_3CN): δ 8.20 (s, 2 H), 7.68 (d, $J = 3.8$ Hz, 1 H), 7.54-7.51 (m, 2 H), 7.43-7.39 (m, 2 H), 7.37-7.32 (m, 1 H), 7.08-7.03 (m, 2 H), 6.98-6.92 (m, 2 H), 6.18 (dd, $J = 4.5, 9.9$ Hz, 1 H), 4.88-4.83 (m, 1 H), 4.25 (d, $J = 10.9$ Hz, 1 H), 3.99-3.91 (m, 2 H), 3.85 (s, 3 H), 3.83 (s, 3 H), 3.79-3.73 (m, 1 H), 3.69 (dd, $J = 9.7, 13.9$ Hz, 1 H), 3.35 (dd, $J = 4.5, 14.1$ Hz, 1 H), 3.20-3.12 (m, 2 H), 2.75-2.61 (m, 5 H), 1.93-1.85 (m, 1 H), 1.68-1.45 (m, 3 H), 1.31-1.23 (m, 1 H). NH not seen. LCMS (Method 3): $[MH^+] = 756$ at 2.74 min.


(continue)

[(1<i>S</i>)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-oxo-1-phenyl-2-[(3<i>R</i>)-quinuclidin-3-yl]oxy-ethyl]-methyl-amino]methyl]thiophene-2-carboxylate	Example 33 (Diastereoisomer 1)	^1H NMR (400 MHz, CD_3CN): δ 8.21 (s, 2 H), 7.68 (d, $J = 3.8$ Hz, 1 H), 7.62-7.58 (m, 2 H), 7.44-7.39 (m, 2 H), 7.37-7.33 (m, 1 H), 7.10-7.04 (m, 2 H), 6.98-6.95 (m, 2 H), 6.19 (dd, $J = 4.4, 9.7$ Hz, 1 H), 4.96-4.92 (m, 1 H), 4.27-4.17 (m, 2 H), 4.07-
--	---	---

		3.93 (m, 2 H), 3.86 (s, 3 H), 3.83 (s, 3 H), 3.70 (dd, J = 9.9, 14.1 Hz, 1 H), 3.36 (dd, J = 4.5, 14.1 Hz, 1 H), 3.22 (dd, J = 8.2, 14.8 Hz, 1 H), 2.99 (br s, 1 H), 2.77-2.60 (m, 5 H), 2.37 (s, 3 H), 2.03-1.99 (m, 1 H), 1.81-1.53 (m, 3 H), 1.42-1.33 (m, 1 H). LCMS (Method 3): [MH⁺] = 770 at 2.81 min.
[(1<i>S</i>)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-oxo-1-phenyl-2-[(3<i>R</i>)-quinuclidin-3-yl]oxy-ethyl]-methyl-amino]methyl]thiophene-2-carboxylate 	Example 34 (Diastereoisomer 2)	¹H NMR (400 MHz, CD₃CN): δ 8.21 (s, 2 H), 7.69 (d, J = 3.8 Hz, 1 H), 7.63-7.59 (m, 2 H), 7.44-7.40 (m, 2 H), 7.37-7.33 (m, 1 H), 7.10-7.04 (m, 2 H), 6.99-6.95 (m, 2 H), 6.19 (dd, J = 4.5, 9.6 Hz, 1 H), 4.98-4.93 (m, 1 H), 4.26 (d, J = 12.2 Hz, 1 H), 4.18 (d, J = 11.6 Hz, 1 H), 4.08-3.97 (m, 2 H), 3.86 (s, 3 H), 3.83 (s, 3 H), 3.70 (dd, J = 9.9, 14.1 Hz, 1 H), 3.37 (dd, J = 4.5, 14.1 Hz, 1 H), 3.24-3.16 (m, 1 H), 3.05 (br s, 1 H), 2.79-2.61 (m, 5 H), 1.99-1.91 (m, 1 H), 2.38 (s, 3 H), 1.75-1.53 (m, 3 H), 1.40-1.30 (m, 1 H). LCMS (Method 3): [MH⁺] = 770 at 2.81 min.


(continue)

[(1<i>S</i>)-2-(3,5-dichloro-1-oxo-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(methoxymethyl)-2-oxo-1-phenyl-2-[(3<i>R</i>)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate	Example 35 (Diastereoisomer 2)	¹H NMR (400 MHz, DMSO): δ 8.57 (s, 2 H), 7.66 (d, J = 3.8 Hz, 1 H), 7.53-7.49 (m, 2 H), 7.43-7.37 (m, 2 H), 7.33 (dd, J = 7.1, 7.1 Hz, 1 H), 7.04 (s, 1 H), 7.01 (s, 2 H), 6.97 (d, J = 3.8 Hz, 1 H), 6.16 (dd, J = 4.3, 9.6 Hz, 1 H), 4.84-4.80 (m, 1 H), 4.01-3.84 (m, 3 H), 3.80 (s, 3 H),
--	---	---

		3.77 (s, 3 H), 3.76-3.57 (m, 3 H), 3.35-3.33 (m, 4 H), 3.20-3.13 (m, 1 H), 2.76-2.63 (m, 4 H), 1.94-1.86 (m, 1 H), 1.65-1.57 (m, 1 H), 1.53-1.48 (m, 2 H), 1.31-1.23 (m, 2 H). LCMS [(Method 72)]: [MH ⁺] = 770 at 3.52 min.
---	--	--

Example 16

[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-[(3R)-1-oxidoquinuclidin-1-ium-3-yl]oxy-2-oxo-1-phenyl-ethyl]amino]methyl]thiophene-2-carboxylate



5

The title compound was obtained as a side product during the purification by SFC of Example 7 and Example 8.

¹H NMR (400 MHz, CD₃CN): δ 8.20 (s, 2 H), 7.68 (d, J = 3.8 Hz, 1 H), 7.58-7.53 (m, 2 H), 7.44-7.39 (m, 2 H), 7.37-7.30 (m, 1 H), 7.07 (d, J = 2.1 Hz, 1 H), 7.03 (dd, J = 8.4, 2.2 Hz, 1 H), 6.97-6.92 (m, 2 H), 6.18 (dd, J = 9.6, 4.5 Hz, 1 H), 5.15-5.09 (m, 1 H), 3.88 (d, J = 7.5 Hz, 2 H), 3.83 (s, 3 H), 3.82 (s, 3 H), 3.67 (dd, J = 14.3, 10 Hz, 1 H), 3.69-3.61 (m, 1 H), 3.34 (dd, J = 14.1, 4.6 Hz, 1 H), 3.26-2.92 (m, 6 H), 2.14-2.09 (m, 1 H), 2.04-1.98 (m, 1 H), 1.89-1.84 (m, 1 H), 1.83-1.80 (m, 1 H), 1.80-1.75 (m, 1 H), 1.73 (s, 3 H). LCMS (Method 2): [MH⁺] = 756 at 2.80 min.

15

PHARMACOLOGICAL ACTIVITY OF THE COMPOUNDS OF THE INVENTION

In vitro determination of PDE4 inhibitory activity

In vitro determination of PDE4 inhibitory activity for compounds of the invention may be determined according to one of the protocols herebelow reported:

PDE4B2 HTRF assay:

PDE4B2 activity is detected using the LANCE Ultra cAMP homogeneous time
5 resolved fluorescence resonance energy transfer (TR-FRET) assay from Perkin Elmer. The assay is based on the competition between the europium (Eu) chelate-labeled cAMP tracer and sample cAMP for binding sites on cAMP-specific monoclonal antibodies (mAb) labelled with the ULight™ dye. The assay is carried out in 384-well low volume plates in a volume of 10 µl. Human recombinant PDE4B2 (80 pM) is incubated for 2 h with 3 nM
10 cAMP in buffer containing 1 × HBSS, 5 mM HEPES, 3 mM MgCl₂, 0.1% BSA, pH 7.4 with or without test compounds. The enzymatic reactions are efficiently stopped by the addition of 500 µM IBMX present in the combined Stop/Detection buffer containing europium (Eu) chelate-labeled cAMP tracer and cAMP-specific monoclonal antibodies (mAb) labelled with the ULight™ dye. Samples are then further incubated for 1 h before
15 plates are read at ex 340 nm and em at 665 nm and 615 nm on an EnVision reader. IC₅₀ values are determined from competition curves using a non-linear curve fitting program.

PDE4 cell free assay protocol

PDE4 activity is determined in U937 human monocytic supernatants cells lysate. Cells are cultured, harvested and supernatant fraction prepared essentially as described in
20 Torphy TJ et al J. Pharmacol. Exp. Ther. 1992; 263:1195-1205.

U937 cells are grown at 37°C, 5% CO₂ in RPMI 1640 with GlutaMAX™-I medium supplemented with 10% fetal bovine serum and 100 µg/mL Pen-strep (Gibco).

Cells are harvested and washed twice by centrifugation (150 × g, 8 min) in cold PBS. Washed cells are re-suspended in cold Krebs-Ringer-Henseleit buffer at a final
25 concentration 20 × 10⁶ cells /mL and sonicated. After centrifugation at 15000 x g for 20 min, the supernatants are pooled, divided in aliquots and stored at -80°C.

PDE4 activity is determined in cells supernatants by assaying cAMP disappearance from the incubation mixtures.

The concentration of the test compounds ranges between 10^{-12} M and 10^{-6} M. Reactions are stopped by enzyme heat inactivation (2.5 minutes at 100°C) and residual cAMP content is determined using the 'LANCE cAMP Assay' from PerkinElmer following the provider instructions.

5 The results, expressed as mean $\pm$ standard deviation of the molar concentration of the test compound producing 50% inhibition of cAMP disappearance (IC_{50}).

Percentage of inhibition of PDE4 activity is calculated, assuming cAMP disappearance in the absence of inhibitors as 100% and cAMP disappearance in heat inactivated samples as 0%.

10 Representative compounds of the invention, when tested in one of the above reported protocols, displayed an IC_{50} lower than 100 nM.

In vitro determination of M3 antagonism

In vitro determination of M3 antagonism for compounds of the invention may be determined according to one of the protocols herebelow reported:

15 **M3 Receptor radioligand binding assay:**

Human M_3 receptor membranes (15 $\mu\text{g}/\text{well}$) from Perkin Elmer are incubated with 0.52 nM Scopolamine Methyl Chloride, [N-methyl- ^3H] with or without test compounds, or a saturating concentration of Atropine (5 μM) for the determination of non-specific binding. The assay is carried out in 96-well polypropylene plates in a volume of 250 μl .


20 The assay buffer used is 50 mM Tris-HCl, 154 mM NaCl (pH 7.4). The final assay concentration of DMSO is 0.5% (v/v). The plates are sealed and incubated for 2 h at room temperature on an orbital shaker (slow speed). Membranes are harvested onto 96-well unfilter GF/C filter plates pre-treated with 0.5% polyethyleneimine (v/v), using a filter manifold, washed four times with 200 μl of assay buffer. The plates are dried before

25 addition of 50 μl of microscint-0, sealed then read in a Trilux Microbeta scintillation counter. IC_{50} values are determined from competition curves using a non-linear curve fitting program. K_i values are calculated from IC_{50} values by the Cheng and Prusoff equation.

CLAIMS

1. A compound of general formula (I)

5



(I)

wherein

each R₁ is hydrogen or is independently selected in the group consisting of: halogen, (C₁-C₄) alkyl, (C₁-C₄) alkoxy, (C₁-C₄) haloalkyl, hydroxy, -SO₂NR^IR^{II}, -CN, -NR^ISO₂R^{III},
 10 -NR^IR^{II}, -(CO)NR^IR^{II} and -NR^I(CO)R^{III}, and wherein said (C₁-C₄) alkyl is optionally substituted by one or more groups selected from (C₃-C₇) cycloalkyl, hydroxy and -NR^IR^{II} and wherein said (C₁-C₄) alkoxy is optionally substituted by one or more halogens or groups (C₃-C₇) cycloalkyl wherein,

R^I is hydrogen or (C₁-C₆) alkyl;

15 R^{II} is hydrogen or (C₁-C₆) alkyl;

R^{III} is hydrogen or (C₁-C₆) alkyl;

n is an integer ranging from 1 to 3;

each R₂ is hydrogen or is independently selected in the group consisting of: halogen, (C₁-C₄) alkyl, (C₁-C₄) alkoxy, (C₁-C₄) haloalkyl, hydroxy, -SO₂NR^IR^{II}, -CN, -NR^ISO₂R^{III}, -
 20 NR^IR^{II}, -(CO)NR^IR^{II} and -NR^I(CO)R^{III} and wherein said (C₁-C₄) alkyl is optionally substituted by one or more groups selected from (C₃-C₇) cycloalkyl, hydroxy and -NR^IR^{II} and wherein said (C₁-C₄) alkoxy is optionally substituted by one or more halogens or groups (C₃-C₇) cycloalkyl wherein,


R^I is hydrogen or (C₁-C₆) alkyl;

R^{II} is hydrogen or (C₁-C₆) alkyl;

R^{III} is hydrogen or (C₁-C₆) alkyl;

m is an integer ranging from 1 to 3;

- 5 R_3 and R_4 are different or the same and are independently selected from the group consisting of: H, (C₃-C₇) cycloalkylcarbonyl, (C₁-C₆) alkyl optionally substituted by one or more substituents selected from (C₃-C₇) cycloalkyl and (C₅-C₇) cycloalkenyl, (C₁-C₆) haloalkyl, (C₃-C₇) cycloalkyl, (C₅-C₇) cycloalkenyl, (C₂-C₆) alkenyl, and (C₂-C₆) alkynyl; or R_3 and R_4 , together with the interconnecting atoms, form a 2,2-difluoro-1,3-dioxolane
10 ring of formula (r) fused to the phenyl moiety which bears groups $-OR_3$ and $-OR_4$, wherein asterisks indicate carbon atoms shared with such phenyl ring:



(r);

- 15 each R_5 is selected from the group consisting of: CN, NO₂, CF₃ and halogen atoms;
k is 0 or an integer ranging from 1 to 3;
 x' is 0 or 1;
 L_1 is selected from a bond and $-(CH_2)_p-$ wherein p is an integer ranging from 1 to 4;
 W_1 is selected from a divalent arylene, heteroarylene and saturated monocyclic
20 heterocycloalkylene group;
 L_2 is a group selected from: a bond, $-(CH_2)_q-$ wherein q is 1 or 2, [1]-(CO)-[X]-
(CH₂)_t-[2], and [1]-(SO₂)-[X]-(CH₂)_t-[2], wherein [1] and [2] represent, respectively the
point of attachment of group L_2 to the ring W_1 and to the chain nitrogen atom, and wherein
[X] is a bond or a substituted or unsubstituted arylene group;
25 t is an integer ranging from 1 to 4;
 W_2 is selected from an aryl and heteroaryl group;
L is a bond or a $-(CH_2)-$ group;

R₆ is selected in the group consisting of (C₁-C₄) alkyl, (C₁-C₄) alkoxy, (C₁-C₄) haloalkyl, and -CN, wherein said (C₁-C₄) alkyl is optionally substituted by one or more groups selected from (C₃-C₇) cycloalkyl, (C₁-C₄) alkoxy and hydroxyl, or, in alternative, when R₆ is (C₁-C₄) alkyl, W₂ is a phenyl ring, one of R₁ is an alkyl in ortho position with respect to L, both R₁ and R₆ may be connected to form with W₂ a condensed ring radical selected from at least 1H-cyclopropabenzene-1,1-diyl, indane-1,1-diyl (also named as 2,3-dihydro-1H-indene-1,1-diyl), indane-2,2-diyl (also named as 2,3-dihydro-1H-indene-2,2-diyl), 1,2,3,4-tetrahydronaphthalene-1,1-diyl, and 1,2,3,4-tetrahydronaphthalene-2,2-diyl;


R₇ is selected from hydrogen and (C₁-C₄) alkyl optionally substituted by hydroxy or -NR₁₁R₁₂ and wherein R₁₁ and R₁₂ are independently selected from hydrogen, (C₁-C₄) alkyl, or, together with the nitrogen atom they are linked to may form a saturated heterocycloalkyl group having an additional heteroatom selected from O, S and NH;

A is a nitrogen containing group which may be selected from:


- a group (a) which is -(CH₂)_s-NR₈R₉ wherein s is an integer ranging from 1 to 4 and R₈ and R₉ are independently hydrogen or (C₁-C₄) alkyl; and

- a group (b) which is a saturated monocyclic, bicyclic or tricyclic heterocyclic ring system optionally substituted by one or two groups R₁₀ which are at each occurrence independently selected from (C₁-C₄) alkyl and benzyl; deuterated derivative, and pharmaceutically acceptable salts, or solvates thereof.


2. A compound according to claim 1 wherein A is represented by a group of formula (i), (ii), (iii) or (iv):




(i)



(ii)



(iii)



(iv)

5

and wherein


$f = 1, 2$ or 3 ;

$g = 1, 2$ or 3 ;

and the asterisk (*) represents the point of attachment to the oxygen atom of formula

10 (I).

3. A compound according to claim 1 or 2 wherein x' is 1, represented by the formula (IA)




(IA)

15

wherein $R_1, R_2, R_3, R_4, R_5, R_6, R_7, L, L_1, W_1, L_2, W_2, A, m, n$, and k are as described above, deuterated derivative and pharmaceutically acceptable salt and solvate thereof:

4. A compound according to claim 1 wherein W_2 is a phenyl ring, L is a bond, and one

of R₁, in ortho position with respect to L, and R₆ may be connected to form a ring radical, represented by the general formula (IB)




(IB)

wherein R₁, R₂, R₃, R₄, R₅, R₇, A, L₁, W₁, L₂, m, k and x' are as defined above for compounds of formula (I), deuterated derivative and pharmaceutically acceptable salt and solvate thereof.

5. A compound according to claim 4 wherein L₁ is a bond, W₁ is a divalent group selected from thiophene-2,5-diyl, thiophene-2,4-diyl, phenylene-1,4-diyl, phenylene-1,3-diyl and phenylene-1,2-diyl, L₂ is -(CH₂)-, R₇ is H and R₁, R₂, R₃, R₄, R₅, A, m and k are as defined above for compounds of formula (I), deuterated derivative and pharmaceutically acceptable salt and solvate thereof.

6. A compound according to claim 1 wherein W₂ is a phenyl ring and L is a bond represented by the general formula (IC):




(IC)

wherein R_6 is selected from methyl, ethyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, methoxymethyl, trifluoromethyl and difluoromethyl and $R_1, R_2, R_3, R_4, R_5,$
 5 $R_7, A, L_1, W_1, L_2, m, n, k$ and x' are as defined above for compounds of formula (I),
 pharmaceutically acceptable salt and solvate thereof.

7. A compound according to claim 6 wherein L_1 is a bond, W_1 is a divalent group
 selected from thiophene-2,5-diyl, thiophene-2,4-diyl, phenylene-1,4-diyl, phenylene-1,3-
 diyl and phenylene-1,2-diyl, L_2 is $-(CH_2)-$, R_7 is H or methyl, R_6 is selected from methyl,
 10 ethyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, methoxymethyl, trifluoromethyl
 and difluoromethyl and $R_1, R_2, R_3, R_4, R_5, A, m, n, k$ and x' are as defined above for
 compounds of formula (I), pharmaceutically acceptable salt and solvate thereof

8. A compound according to claim 1 wherein L_1 is a bond, W_1 is selected from a
 divalent saturated monocyclic heterocycloalkylene group, represented by the general
 15 formula (ID)

90



(ID)

and wherein

Y is S or CH₂;


- 5 L₂ is a group selected from: -(CH₂)_q- wherein q is 1 or 2, [1]-(CO)-[X]-(CH₂)_t-[2], and [1]-(SO₂)-[X]-(CH₂)_t-[2], wherein [1] and [2] represent, respectively the point of attachment of group L₂ to the saturated monocyclic heterocycloalkylene ring (W₁) and to the chain nitrogen atom, and wherein

[X] is a bond or a substituted or unsubstituted arylene group selected from
 10 phenylene-1,4-, -1,3- and -1,2-diyl ;

t is an integer ranging from 1 to 4

and wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, A, W₂, L, m, n, k and x' are as defined above for compounds of formula (I), and pharmaceutically acceptable salt and solvate thereof.

9. A compound according to any one of claims 1 to 8, represented by formula (I)'
- 15 wherein the absolute configuration of carbon (1) is that shown herebelow



(I)',

wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, L₁, W₁, L₂, W₂, L, A, n, m, k and x' are as defined above for compounds of formula (I), and pharmaceutically acceptable salt and solvate thereof.

10. A compound according to claim 1 which is selected in the list consisting of:

[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate formate salt

10 [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 4-[[[1-methyl-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]benzoate
[(3R)-quinuclidin-3-yl] 1-[[4-[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethoxy] carbonylphenyl] methylamino]indane-1-carboxylate

15 [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(3R)-quinuclidin-3-yl]oxycarbonylindan-1-yl]amino]methyl]thiophene-2-carboxylate

[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenyl-ethyl]amino]methyl]thiophene-2-carboxylate

20 [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-phenyl-1-[(3R)-quinuclidin-3-yl]oxycarbonyl-propyl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxyethyl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxyethyl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 4-[[[1-methyl-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxyethyl]amino]methyl]benzoate

10 Single diastereoisomer of [(3R)-quinuclidin-3-yl] 1-[[4-[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethoxy] carbonylphenyl] methylamino]indane-1-carboxylate

Single diastereoisomer of [(3R)-quinuclidin-3-yl] 1-[[4-[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethoxy] carbonylphenyl]

15 methylamino]indane-1-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-[(3R)-quinuclidin-3-yl]oxycarbonylindan-1-yl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-[(3R)-quinuclidin-3-yl]oxycarbonylindan-1-yl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenyl-ethyl]amino]methyl]thiophene-2-carboxylate

25 Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenyl-ethyl]amino]methyl]thiophene-2-carboxylate

[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-methyl-2-[(3R)-1-oxidoquinuclidin-1-ium-3-yl]oxy-2-oxo-1-phenyl-ethyl]amino]methyl]thiophene-2-carboxylate

5 [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(2-dimethylaminoethyloxycarbonyl)indan-1-yl]amino]methyl]thiophene-2-carboxylate

[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-[(1-methyl-4-piperidyl)oxycarbonyl]indan-1-yl]amino]methyl]thiophene-2-carboxylate

[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl]-5-[[[1-(hydroxymethyl)-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenyl-

10 ethyl]amino]methyl]thiophene-2-carboxylate

[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]-methyl-amino]methyl]thiophene-2-carboxylate

[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl]-5-[[[1-

15 (hydroxymethyl)-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate

[(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(methoxymethyl)-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate

20 Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(2-dimethylaminoethyloxycarbonyl)indan-1-yl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(2-dimethylaminoethyloxycarbonyl)indan-1-

25 yl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-[(1-methyl-4-piperidyl)oxycarbonyl]indan-1-yl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-[(1-methyl-4-piperidyl)oxycarbonyl]indan-1-yl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-phenyl-1-[(3R)-quinuclidin-3-yl]oxycarbonyl-propyl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-phenyl-1-[(3R)-quinuclidin-3-yl]oxycarbonyl-propyl]amino]methyl]thiophene-2-carboxylate

10 Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenyl-ethyl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-[(1-methyl-4-piperidyl)oxy]-2-oxo-1-phenyl-ethyl]amino]methyl]thiophene-2-carboxylate

15 Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate

20 Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]-methyl-amino]methyl]thiophene-2-carboxylate

25 Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(hydroxymethyl)-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]-methyl-amino]methyl]thiophene-2-carboxylate

Single diastereoisomer of [(1S)-2-(3,5-dichloro-1-oxido-pyridin-1-ium-4-yl)-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[1-(methoxymethyl)-2-oxo-1-phenyl-2-[(3R)-quinuclidin-3-yl]oxy-ethyl]amino]methyl]thiophene-2-carboxylate

and a pharmaceutically acceptable salt or solvate thereof.

- 5 11. A pharmaceutical composition comprising a compound as defined in any one of claims 1 to 10, in admixture with one or more pharmaceutically acceptable carrier.
12. A pharmaceutical composition according to claim 11 further comprising another active ingredient,
13. A compound according to any one of claims 1 to 10, for use as a medicament.
- 10 14. A compound according to any one of claims 1 to 10, for use in the prevention and/or treatment of a disease of the respiratory tract characterized by airway obstruction.
15. A compound for use as defined in claim 14 wherein the disease of the respiratory tract is selected from asthma and COPD.