

**Soluble immunoreactive *Treponema pallidum* TpN47 antigens**Field of the invention

5 The invention concerns soluble, stable and immunoreactive variants of *Treponema pallidum* antigen 47 (TpN47) comprising at least amino acid residues 63 to 174 (*i.e.* domain B) of the TpN47 protein molecule with the proviso that the antigens lack amino acids 224 to 351 (*i.e.* domain C) of TpN47, wherein the TpN47 antigen is fused with a chaperone. The invention further concerns a soluble *Treponema pallidum* antigen 47 (TpN47 antigen) comprising domains  
10 B and D or domains A+B and D, also with the proviso that these antigens lack domain C. Moreover, the invention concerns a method of producing these soluble, stable and immune-reactive TpN47 variants as well as the use of these antigens in an immunodiagnostic assay aiming at the detection of antibodies against *Treponema pallidum* in an isolated sample.

15

Background of the invention

*Treponema pallidum* belongs to the bacterial family of spirochetes and is the causative agent of syphilis. Syphilis, also called Lues, is mainly transmitted by sexual contact. *Treponema pallidum*  
20 can also pass from mother to baby during pregnancy. The disease is characterized by distinct clinical stages and long periods of latent, asymptomatic infection. Many people do not notice symptoms and thus are unaware of their syphilis infection for years. The primary infection is confined and usually causes a small painless ulcer (primary stage, "Lues I"). If left untreated by penicillin, the disease proceeds to the secondary stage Lues II (about eight weeks after infection),  
25 which entails flu-like symptoms, non-itchy skin rash and swollen lymph nodes. After some years, at stage Lues III, syphilitic nodes appear throughout the body. The final stage (Lues IV) is characterized by destruction of the central nervous system eventually leading to neurological and cardiological disorders, general paralysis, ataxia, dementia and blindness.

30 Although effective therapies have been available since the introduction of penicillin in the mid-20th century, syphilis still remains an important global health problem. It is mandatory to identify patients with *Treponema* infection, to support antibiotic therapy and thus to prevent spread of syphilis. As a consequence, it is necessary to provide reliable diagnostic tools such as

immunoassays for the detection of antibodies against *Treponema pallidum*. Yet, in order to be used as specific compounds in serological applications, recombinant proteins have to meet several requirements such as solubility, stability and antigenicity.

- 5 One of the membrane-associated polypeptides of *Treponema pallidum* (the causative agent of Syphilis infections) is TpN47, a large protein that consists of 434 amino acid residues. TpN47 has been ascribed immunodominance in the humoral immune response against *Treponema* (Folia Microbiol. (2003) 48 (4), 549-553), and antibodies towards TpN47 are frequently found in human sera from *Treponema*-infected individuals. Thus, a soluble and antigenic variant of recombinant
- 10 TpN47 would be an invaluable ingredient to establish an immunoassay for detection of *Treponema* antibodies that combines high sensitivity and specificity. In the best case, such an antigen should enable the detection of both IgG and IgM molecules.

Recombinant variants of TpN47 have been described in literature. In the Journal of Immunology

15 (1996) Jul 15;157(2):720-31, Baughn *et al.* report on an epitope scan of 12-mer peptides (overlap 8 amino acid residues, offset 4 amino acid residues) encompassing the entire sequence of TpN47. Based on this scan, the authors describe as many as ten immunodominant TpN47 fragments, ranging from 9 to 29 amino acid residues in length. The crystal structure and the domain topology of TpN47 has also been described (Journal of Biological Chemistry (2002), 277 (4), 41857-41864, Deka *et al.*).

20

Immunoassays for detecting *Treponema* antibodies are known in the art. For example, Rostopira *et al.* (Folia Microbiol.48(4), 549-553, 2003) describe an immunoassay for diagnosing syphilis using a combination of TpN17 and TpN47 antigens, identifying TpN47 as one of the immunodominant antigens. In this publication full-length TpN47 was used as an antigen.

25

We overproduced a full-length variant of TpN47 in an *E. coli* host (BL21/DE3) and succeeded in preparing the TpN47 antigen to homogeneity. Yet, our initial experiments with the full-length version of TpN47 unambiguously revealed that this protein tends to aggregate at temperatures above 35°C. Despite the fusion of tandem chaperone fusions such as *EcSlyD-EcSlyD* or even (the more thermostable) *EcSlpA-EcSlpA*, full-length TpN47 inevitably aggregated into a high-molecular-

30 weight associate at temperatures above 35 °C. Since *Treponema pallidum* is known as a rather temperature-sensitive pathogen, the finding that one of its major membrane proteins shares temperature sensitivity may not seem too surprising. Anyway, thermally induced aggregation

processes of proteinaceous ingredients in immunoassays usually result either in a loss of signal or in a loss of specificity. Thus, the fact that full-length TpN47 (even in fusion with solubility-enhancing chaperones such as SlyD or SlpA) tends to aggregate even at moderately elevated temperatures (> 35 °C), clearly precludes this molecule from simple and straightforward applications in a sensitive immunoassay of the DAGS format.

Despite the detailed structural knowledge on TpN47 (Deka *et al.* Journal of Biological Chemistry (2002), 277 (4), 41857-41864), the prior art does neither disclose the pronounced thermolability of TpN47 nor does it disclose TpN47 antigen variants that overcome the problem of thermally induced aggregation and the concomitant loss of sensitivity in immunoassays aiming at the detection of antibodies against *Treponema pallidum* in a sample.

Guo *et al.* (Xiamen Daxue Xuebao -Ziran Kexue Ban (2008), 47(6), 874-878) describe specific soluble TpN47 N- or C-terminal truncated mutants that are recombinantly expressed in *E. coli*.

However, the problem of thermal instability and tendency to aggregation of full-length TpN47 protein is not addressed. In addition, the data of Guo *et al.* suggest that a combination of the domains C and D (C190) of TpN47 almost equals the antigenicity of the full-length TpN47 protein.

The stability problem has been solved in the current invention by generating soluble, stable and immunoreactive variants of *Treponema pallidum* antigen 47 (TpN47 antigen) comprising at least domain B (aa 63-174) of the TpN47 protein molecule with the proviso that the antigens lack domain C (aa 224-351) of TpN47, wherein the TpN47 antigen is fused with a chaperone.

A further solution of this problem is a soluble TpN47 antigen comprising domains B and D (aa 63-174 and 352 to 434 of SEQ ID NO. 1) or a TpN47 antigen comprising domains A+B and D (aa 26 to 223 and 352 to 434 of SEQ ID NO. 1). Both variants lack domain C of TpN47.

### Summary

The disclosure relates to soluble *Treponema pallidum* antigens, *i.e.* to TpN47 antigen that comprises at least domain B (amino acid residues 63 to 174 of SEQ ID NO. 1) or at least domains A+B (amino acid residues 26 to 223 of SEQ ID NO. 1) with the proviso that the TpN47 antigen lacks domain C (amino acid residues 224 to 351 of SEQ ID NO. 1), wherein the TpN47

antigen is fused with a chaperone. In a first aspect, there is provided a soluble *Treponema pallidum* antigen 47 (TpN47 antigen) comprising amino acid residues 63 to 174 (domain B) of Swiss Prot P29723 with the proviso that the antigen lacks amino acid residues 224 to 351 (domain C) of Swiss Prot P29723, wherein the TpN47 antigen is fused with a chaperone. In a second aspect, there is provided a soluble *Treponema pallidum* antigen 47 (TpN47 antigen) comprising amino acid residues 26 to 223 (domain A+B) of Swiss Prot P29723 with the proviso that the antigen lacks amino acid residues 224 to 351 (domain C) of Swiss Prot P29723, wherein the TpN47 antigen is fused with a chaperone. The disclosure also relates to soluble TpN47 antigens that comprise at least domain B (amino acid residues 63 to 174 of SEQ ID NO. 1) and domain D (amino acid residues 352 to 434 of SEQ ID NO. 1), or domains A+B (amino acid residues 26 to 223 of SEQ ID NO. 1) and domain D (amino acid residues 352 to 434 of SEQ ID NO. 1) with the proviso that these antigens also lack domain C (amino acid residues 224 to 351 of SEQ ID NO. 1). In a third aspect, there is provided a soluble *Treponema pallidum* antigen 47 (TpN47 antigen) comprising amino acid residues 63 to 174 (domain B) of Swiss Prot P29723 with the proviso that the antigen lacks amino acid residues 224 to 351 (domain C) of Swiss Prot P29723, additionally comprising amino acid residues 352 to 434 of Swiss Prot P29723 (domain D). In a fourth aspect, there is provided a soluble *Treponema pallidum* antigen 47 (TpN47 antigen) comprising amino acid residues 26 to 223 (domain A+B) of Swiss Prot P29723 with the proviso that the antigen lacks amino acid residues 224 to 351 (domain C) of Swiss Prot P29723, additionally comprising amino acid residues 352 to 434 of Swiss Prot P29723 (domain D). These antigens may be fused to a chaperone or to another fusion partner in order to further increase its solubility.

In a fifth aspect, there is provided a recombinant DNA molecule encoding the TpN47 antigen as described herein. The disclosure also concerns an expression vector containing operably linked or integrated the above-described DNA encoding a TpN47 antigen. In a sixth aspect, there is provided an expression vector comprising operably linked a recombinant DNA molecule as described herein. In a seventh aspect, there is provided a host cell transformed with said expression vector as described herein. The invention also concerns a method of producing said TpN47 antigen. In an eighth aspect, there is provided a method of producing a soluble and immunoreactive *Treponema pallidum* antigen 47 (TpN47 antigen), the method comprising the steps of culturing host cells as described herein, expression of the TpN47 antigen and purification of the TpN47 antigen.

Moreover, the disclosure relates to *in vitro* diagnostic methods for the detection of Syphilis, *i.e.* to a method of detecting antibodies against TpN47 using said TpN47 antigen variants and it also relates to a reagent test kit comprising a TpN47 antigen according to the disclosure. The disclosure also relates to a composition of at least two *Treponema pallidum* antigens comprising, e.g. a TpN47 antigen and a TpN17 or a TpN15 antigen. In a ninth aspect, there is provided a composition of at least two *Treponema pallidum* antigens comprising a TpN47 antigen as described herein and at least one additional *Treponema pallidum* antigen selected from the group consisting of a TpN17 antigen and a TpN15 antigen. In another embodiment said composition comprises a TpN47 antigen and both TpN17 and TpN15 antigens. Additionally, the disclosure concerns a method of producing these antigens as well as the use of these antigens in an immunodiagnostic assay for the detection of antibodies against *Treponema pallidum* in an isolated sample. In a tenth aspect, there is provided a method for detecting antibodies specific for *Treponema pallidum* in an isolated sample wherein a TpN47 antigen as described herein or a combination or composition of antigens as described herein is used as a capture reagent and/or as a binding partner for the *Treponema pallidum* antibodies.

In an eleventh aspect, there is provided a method for detecting antibodies specific for *Treponema pallidum* in an isolated sample, the method comprising (a) forming an immunoreactive admixture by admixing a body fluid sample with TpN47 antigen as described herein or with a composition of at least two *Treponema pallidum* antigens as described herein; (b) maintaining the immunoreaction admixture for a time period sufficient for allowing antibodies against the *Treponema* antigen or composition of *Treponema* antigens present in the body fluid sample to immunoreact with the *Treponema* antigen or composition of *Treponema* antigens to form an immunoreaction product; and (c) detecting the presence and/or the concentration of any of the immunoreaction product.

In a twelfth aspect, there is provided use of a TpN47 antigen as described herein or of a composition of *Treponema pallidum* antigens as described herein in an *in vitro* diagnostic test for the detection of anti-*Treponema pallidum* antibodies.

In a thirteenth aspect, there is provided a reagent kit for the detection of anti-*Treponema pallidum* antibodies, comprising at least a TpN47 antigen as described herein or an antigen composition as described herein.

Brief description of the figures

Figures 1-6 show analytical gel filtration profiles of distinct *EcSlyD-EcSlyD*-TpN47 domain constructs on a Superdex 200 HR 10/30 column, see also example 5. Ca. 200  $\mu$ l of a 1.3 mg/ml protein solution (fusion protein dissolved in 150 mM potassium phosphate pH 8.0, 100 mM KCl, 0.5 mM EDTA) was applied on the SEC column, and elution was monitored at 280 nm at a flow rate of 0.8 ml/min. The FPLC standard (light gray dotted line) contains  $\beta$ -galactosidase (465 000 Da), sheep IgG (150 000 Da), sheep IgG Fab fragment (50 000 Da), horse myoglobin (17 000 Da), and the dipeptide glycine-tyrosine (238 Da).

All of the protein constructs were subjected to elevated temperatures (35°C, 40°C) in an overnight incubation (18 h) under identical buffer concentrations (150 mM potassium phosphate pH 8.0, 100 mM KCl, 0.5 mM EDTA) at a concentration of 1.3 mg/ml, equivalent to a molar concentration of 15.2  $\mu$ M – 26.5  $\mu$ M. Following thermal stress, the protein samples were subjected to centrifugation and then were assessed for their tendency to aggregate by means of FPLC analysis as described above.

Figure 1 exhibits the thermostability of *EcSlyD-EcSlyD-TpN47/AB* (26-223) as assessed by analytical gel filtration on a Superdex 200 column. 230  $\mu$ l were applied to the column, corresponding to 300  $\mu$ g protein. Elution profiles of TpN47/AB after incubation at 4°C, 35°C and 40°C coincide very nicely. There is no hint to aggregation or association processes. The elution profiles as monitored by absorption at 280 nm point to a soluble protein fragment TpN47/AB even at elevated temperatures such as 40°C.

Figure 2 shows the thermostability of *EcSlyD-EcSlyD-TpN47/B* (63-174) as assessed by analytical gel filtration on a Superdex 200 column. 270  $\mu$ l were applied to the column, corresponding to 350  $\mu$ g protein. Elution profiles of TpN47/B after incubation at 4°C, 35°C and 40°C coincide almost perfectly. There is no hint to aggregation or association processes. The elution profiles as monitored by absorption at 280 nm deliver compelling evidence that protein fragment TpN47/B is soluble and stable even at elevated temperatures such as 40°C.

Figure 3 illustrates the thermostability of *EcSlyD-EcSlyD-TpN47/C* (224-351) as assessed by analytical gel filtration on a Superdex 200 column. 154  $\mu$ l were applied to the column, corresponding to 200  $\mu$ g protein. Elution profiles of TpN47/C after incubation at 4°C and 35°C coincide almost perfectly. However, even at 35°C, a very small portion of TpN47 C elutes in the void volume of the gel filtration column (continuous dark gray line), pointing to the onset of aggregation processes. After incubation at 40°C, almost one third of the fusion protein elutes in the void volume of the SEC column, indicative of the formation of large aggregated protein particles. It becomes evident from figure 3 that TpN47 domain C possesses a substantial tendency to aggregate at temperatures above 35°C.

Figure 4 shows the thermostability of *EcSlyD-EcSlyD-TpN47/CD* (224-434) as assessed by analytical gel filtration on a Superdex 200 column. 195  $\mu$ l were applied to the column, corresponding to 250  $\mu$ g protein. The elution profile of *TpN47/CD* after incubation at 4°C exhibits a nice symmetrical peak, indicative of a soluble and homogeneous protein. When incubated at 35°C, *EcSlyD-EcSlyD-TpN47/CD* is forming large aggregated protein particles, as judged by the corresponding profile, which shows a large portion of the protein eluting in the void volume of the SEC column (continuous dark gray line). After incubation at 40°C, the lion's share of the CD fusion protein elutes in the void volume of the gel filtration column (continuous black line). It is obvious from figure 4 that the C-terminal part of *TpN47*, namely the fusion protein CD, possesses a high intrinsic tendency to aggregate even at a moderately elevated temperature such as 35 °C.

Figure 5 illustrates the thermostability of *EcSlyD-EcSlyD-TpN47/D* (352-434) as assessed by analytical gel filtration on a Superdex 200 column. 331  $\mu$ l were applied to the column, corresponding to 430  $\mu$ g protein. Elution profiles of *TpN47/D* after incubation at 4°C, 35°C and 40°C are almost indistinguishable and coincide perfectly. There is no hint whatsoever to aggregation or association processes. The elution profiles as monitored by absorption at 280 nm deliver compelling evidence that protein fragment *TpN47/D* is perfectly soluble and stable even at elevated temperatures such as 40°C.

Figure 6 shows the thermostability of the full-length *TpN47* variant *EcSlyD-EcSlyD-TpN47/ABCD* (21-434) as assessed by analytical gel filtration on a Superdex 200 column. 90  $\mu$ l were applied to the column, corresponding to 117  $\mu$ g protein. The elution profile of *TpN47/ABCD* after incubation at 4°C exhibits a nice peak at an elution volume of 13.5 ml, indicative of a soluble and homogeneous protein. When incubated at 35°C, *EcSlyD-EcSlyD-TpN47/ABCD* is forming large aggregated protein particles, as judged by the corresponding profile, which shows more than 66 % of the protein eluting in the void volume of the SEC column (continuous dark gray line). After incubation at 40°C, almost 96 % of full-length *TpN47* elute in the void volume of the gel filtration column (continuous black line). It is obvious from figure 6 that full-length *TpN47* possesses a tremendous intrinsic tendency to aggregate even at a moderately elevated temperature such as 35 °C.

Due to a scarcity of full-length protein, only 117  $\mu$ g *EcSlyD-EcSlyD* were applied on the Superdex column (leading to a lower absorption signal), which is only a fractional amount when compared with the other TpN47 variants. The thermal stress assessment has, however, been conducted under identical conditions for each TpN47 variant.

5

Figure 7 shows our strategy of B-cell epitope mapping in the search for immunodominant TpN47 antigen variants. At the top: full length TpN47 (21-434, comprising domains ABCD) with *EcSlyD-EcSlyD* fused to the TpN47 N-terminus. Middle section: individual TpN47 polypeptide fragments (fusion to *EcSlyD* not shown), in ladder-like order: TpN47 polypeptide fragments 30-66, 106-132, 137-170, 197-219, 225-250, 273-296, 321-362, 368-388, 391-434 according to SEQ ID NOs. 8-16; experimental results for immunoreactivity of these peptides according to example 4 are shown in table 2 (Figure 8a-c). Bottom: location of TpN47 domains B, AB, C, D and CD (fusion to *EcSlyD-EcSlyD* not shown) relative to the full length TpN47; experimental results for immunoreactivity (*i.e.* antigenicity) of these TpN47 fusion variants according to example 4 are shown in table 3 (Figure 9).

15

Figure 8a-c shows table 2, containing the results obtained in example 4: Immunological reactivity of short (linear) unstructured TpN47 fragments fused to *EcSlyD* chaperone. The immunoassays were performed by using an Elecsys<sup>®</sup> 2010 analyzer. The signal dynamics are normalized relative to the average value obtained for the *Treponema*-negative samples. The *Treponema*-positive sera were purchased from Boca Biolistics (Coconut Creek, FL, USA), the *Treponema*-negative controls were internal blood donors. Please note that the right column (experiment 10, full length TpN47) is identical in each of figures 8a, b and c, respectively.

20

Figure 9 shows table 3, containing the results obtained in example 4: Immunological reactivity (*i.e.* antigenicity) of large TpN47 fragments (domains) fused to *EcSlyD-EcSlyD* tandem chaperone. The immunoassays were performed by using an Elecsys<sup>®</sup> 2010 analyzer. The signal dynamics (SD) are normalized relative to the average value obtained for the *Treponema*-negative samples. The *Treponema*-positive sera were purchased from Boca Biolistics (Coconut Creek, FL, USA), the negative controls were internal blood donors.

25

30

Figure 10 shows table 4: Residual antigenicity of TpN47 domains AB (26-223) & B (63-174) subsequent to thermal stress (72h at 42°C). Both domains AB and domain B were fused to the

solubility-enhancing tandem chaperone module *EcSlyD-EcSlyD* as described. The immunoassays were performed by using an Elecsys® 2010 analyzer. The signal dynamics (SD) are normalized relative to the average value obtained for the *Treponema*-negative samples. The *Treponema*-positive sera were purchased from SeraCare (MA, USA), the *Treponema*-negative controls were purchased from Trina Bioreactives AG (Nänikon, Switzerland).

Figure 11 shows table 5: Residual antigenicity of TpN47 domains C (224-351) & D (352-434) subsequent to thermal stress (72h at 42°C). Both domain C and D were fused to the solubility-enhancing tandem chaperone module *EcSlyD-EcSlyD* as described. The immunoassays were performed by using an Elecsys® 2010 analyzer. The signal dynamics (SD) are normalized relative to the average value obtained for the *Treponema*-negative samples. The *Treponema*-positive sera were purchased from SeraCare (MA, USA), the *Treponema*-negative controls were purchased from Trina Bioreactives AG (Nänikon, Switzerland).

Figure 12 shows table 6: Residual antigenicity of TpN47/CD (224-434) and full-length TpN47 (21-434) subsequent to thermal stress (72h at 42°C). Both domains CD and full-length TpN47 were fused to the solubility-enhancing tandem chaperone module *EcSlyD-EcSlyD* as described. The immunoassays were performed by using an Elecsys® 2010 analyzer. The signal dynamics (SD) are normalized relative to the average value obtained for the *Treponema*-negative samples. The *Treponema*-positive sera were purchased from SeraCare (MA, USA), the *Treponema*-negative controls were purchased from Trina Bioreactives AG (Nänikon, Switzerland).

As further detailed in the sequence listing the following protein sequences are used within this specification:

SEQ ID NO. 1 shows full length TpN47 fused to two *E. coli* SlyD molecules:

*EcSlyD-EcSlyD-TpN47* (aa 21-434 TpN47 of Swiss Prot P29723 is underlined); in position 315, a cysteine has been replaced by alanine, and for purification purposes a hexa-histidine tag has been added to the C-terminal end.

```

MKVAKDLVVS LAYQVRTEDG VLVDESPVSA PLDYLGHGHS LISGLETALE GHEVGDKFDV
AVGANDAYGQ YDENLVQRVP KDVFMGVDEL QVGMRFLEET DQGPVPVEIT AVEDDHVVVD
GNHMLAGQNL KFNVEVVAIR EATEEELAHG HVHGAHDHHH DHDHDGGGSG GSGGGGSGGG

```

SGGGSGGGKV AKDLVVS LAY QVRTEDGVLV DESPVSA PLDYLHGHGS LISGLETAL E GHEVGD KFDV  
 VGD KFDVAVG ANDAYGQYDE NLVQRVPKDV FMGVDELQVG MRFLAETDQG PVPVEITAVE  
 DDHVVDGNH MLAGQNLKFN VEVVAIREAT EEELAHGHVH GAHDHHDHD HDGGSGGGG  
 GGGSGGGSGG GSGGGGSSHH ETHYGYATLS YADYWAGELG QSRDVLLAGN AEADRAGDLD  
 5 AGMFDVSR THGHGAFRQQ FQYAVEVLGE KVLSKQETED SRGRKKWEYE TDPSVTKMVR  
ASASFQDLGE DGEIKFEAVE GAVALADRAS SFMVDSEYK ITNVKVHGMK FVPVAVPHEL  
KGIAKEKFHF VEDSRVTENT NGLKTMLTED SFSARKVSSM ESPHDLVVD VTGYHSRFG  
SDAEASVMLK RADGSELHR EFIDYVMNFN TVRYDYYGDD ASYTNLMASY GTKHSADSWW  
KTGRVPRISA GINYGFDRFK GSGPGYYRLT LIANGYRDVV ADVRFPLKYE GNIDIGLKKG  
 10 VLTIIGDAE TLMDAAVDVF ADGQPKLVSD QAVSLGQNVL SADFTPGTEY TVEVRFKEFG  
SVRAKVVAQL EHHHHHH

SEQ ID NO. 2 shows domains AB of TpN47 fused to two *E. coli* SlyD molecules: *EcSlyD*-  
 15 *EcSlyD*-TpN47/AB (aa 26-223 TpN47 of Swiss Prot P29723 is underlined); for purification  
 purposes a hexa-histidine tag has been added to the C-terminal end.

MKVAKDLVVS LAYQVRTEDG VLVDSPVSA PLDYLHGHGS LISGLETAL E GHEVGD KFDV  
 AVGANDAYGQ YDENLVQRVP KDVFMDGDEL QVGMFLAET DQGPVPVEIT AVEDDHVVVD  
 20 GNHMLAGQNL KFNVEVVAIR EATEEELAHG HVHGAHDHHD DHDHGGGGSG GSGGGSGGGG  
 SGGSGGGGKV AKDLVVS LAY QVRTEDGVLV DESPVSA PLDYLHGHGS LISGLETAL E GHEVGD KFDV  
 VGD KFDVAVG ANDAYGQYDE NLVQRVPKDV FMGVDELQVG MRFLAETDQG PVPVEITAVE  
 DDHVVDGNH MLAGQNLKFN VEVVAIREAT EEELAHGHVH GAHDHHDHD HDGGSGGGG  
 GGGSGGGSGG GSGGGGETHY GATLSYADYW AGELGQSRDV LLAGNAEADR AGDLDA GDMFD  
 25 AVSRAHGHGA FRQQFQYAVE VLGEKVLSKQ ETEDSRGRKK WEYETDPSVT KMVRASASFQ  
DLGEDGEIKF EAVEGAVALA DRASSFMVDS EYKITNVKV HGMKFVPVAV PHELKGIAKE  
KHFVEDSRV TENTNGLKTM LTEDSFSARK VSLEHHHHHH

30 SEQ ID NO. 3 shows domain B of TpN47 fused to two *E. coli* SlyD molecules:  
*EcSlyD*-*EcSlyD*-TpN47/B (aa 63-174 TpN47 of Swiss Prot P29723 is underlined); for  
 purification purposes a hexa-histidine tag has been added to the C-terminal end.

MKVAKDLVVS LAYQVRTEDG VLVDSPVSA PLDYLHGHGS LISGLETAL E GHEVGD KFDV  
 35 AVGANDAYGQ YDENLVQRVP KDVFMDGDEL QVGMFLAET DQGPVPVEIT AVEDDHVVVD  
 GNHMLAGQNL KFNVEVVAIR EATEEELAHG HVHGAHDHHD DHDHGGGGSG GSGGGSGGGG  
 SGGSGGGGKV AKDLVVS LAY QVRTEDGVLV DESPVSA PLDYLHGHGS LISGLETAL E GHEVGD KFDV  
 VGD KFDVAVG ANDAYGQYDE NLVQRVPKDV FMGVDELQVG MRFLAETDQG PVPVEITAVE  
 DDHVVDGNH MLAGQNLKFN VEVVAIREAT EEELAHGHVH GAHDHHDHD HDGGSGGGG  
 40 GGSGGGSGG GSGGGDLDA G MFDVSRATH GHGAFRQQFQ YAVEVLGEKV LSKQETEDSR  
GRKKWEYETD PSVTKMVRAS ASFQDLGEDG EIKFEAVEGA VALADRASSF MVDSEYKIT  
NVKVHGMLH HHHHH

45 SEQ ID NO. 4 shows domain C of TpN47 fused to two *E. coli* SlyD molecules:  
*EcSlyD*-*EcSlyD*-TpN47/C (aa 224-351 TpN47 of Swiss Prot P29723 is underlined); in position

315, a cysteine has been replaced by alanine and for purification purposes a hexa-histidine tag has been added to the C-terminal end.

```

5  MKVAKDLVVS LAYQVRTEDG VLVDESPVSA PLDYLHGHGS LISGLETALE GHEVGDKFDV
   AVGANDAYGQ YDENLVQRVP KDVFMGVDEL QVGMRFLEET DQGPVPVEIT AVEDDHVVVD
   GNHMLAGQNL KFNVEVVAIR EATEEELAHG HVHGAHDHHH DHDHDGGGSG GSGGGSGGGG
   SGGGSGGGKV AKDLVVSLAY QVRTEDGVLV DESPVSAPLD YLHGHGSLIS GLETALEGHE
   VGDKFDVAVG ANDAYGQYDE NLVQRVPKDV FMGVDELQVG MRFLAETDQG PVPVEITAVE
   DDHVVDGNH MLAGQNLKFN VEVVAIREAT EEELAHGHVH GAHDHHDHD HDGGGSGGGG
10 GGGSGGGSGG GSGGGSMESP HDLVVDTVGT GYHSRFGSDA EASVMLKRAD GSELSHREFI
   DYVMNFNTVR YDYYGDDASY TNLMASYGTK HSADSWWKTG RVPRISAGIN YGFDRFKGSG
   PGYYRLTLIA NGYRDVVADV RFLLEHHHHH H

```

15 SEQ ID NO. 5 shows domains CD of TpN47 fused to two *E. coli* SlyD molecules: *EcSlyD-EcSlyD-TpN47/CD* (aa 224-434 TpN47 of Swiss Prot P29723); in position 315, a cysteine has been replaced by alanine, and for purification purposes a hexa-histidine tag has been added to the C-terminal end.

```

20 MKVAKDLVVS LAYQVRTEDG VLVDESPVSA PLDYLHGHGS LISGLETALE GHEVGDKFDV
   AVGANDAYGQ YDENLVQRVP KDVFMGVDEL QVGMRFLEET DQGPVPVEIT AVEDDHVVVD
   GNHMLAGQNL KFNVEVVAIR EATEEELAHG HVHGAHDHHH DHDHDGGGSG GSGGGSGGGG
   SGGGSGGGKV AKDLVVSLAY QVRTEDGVLV DESPVSAPLD YLHGHGSLIS GLETALEGHE
   VGDKFDVAVG ANDAYGQYDE NLVQRVPKDV FMGVDELQVG MRFLAETDQG PVPVEITAVE
25 DDHVVDGNH MLAGQNLKFN VEVVAIREAT EEELAHGHVH GAHDHHDHD HDGGGSGGGG
   GGGSGGGSGG GSGGGSMESP HDLVVDTVGT GYHSRFGSDA EASVMLKRAD GSELSHREFI
   DYVMNFNTVR YDYYGDDASY TNLMASYGTK HSADSWWKTG RVPRISAGIN YGFDRFKGSG
   PGYYRLTLIA NGYRDVVADV RFLPKYEGNI DIGLKGKVL IGGADAETLM DAAVDVFADG
   QPKLVSDQAV SLGQNVLSAD FTPGTEYTV VRFKEFGSVR AKVVAQLEHH HHHH
30

```

SEQ ID NO. 6 shows domain D of TpN47 fused to two *E. coli* SlyD molecules: *EcSlyD-EcSlyD-TpN47/D* (aa 352-434 TpN47 of Swiss Prot P29723)

```

35 MKVAKDLVVS LAYQVRTEDG VLVDESPVSA PLDYLHGHGS LISGLETALE GHEVGDKFDV
   AVGANDAYGQ YDENLVQRVP KDVFMGVDEL QVGMRFLEET DQGPVPVEIT AVEDDHVVVD
   GNHMLAGQNL KFNVEVVAIR EATEEELAHG HVHGAHDHHH DHDHDGGGSG GSGGGSGGGG
   SGGGSGGGKV AKDLVVSLAY QVRTEDGVLV DESPVSAPLD YLHGHGSLIS GLETALEGHE
   VGDKFDVAVG ANDAYGQYDE NLVQRVPKDV FMGVDELQVG MRFLAETDQG PVPVEITAVE
40 DDHVVDGNH MLAGQNLKFN VEVVAIREAT EEELAHGHVH GAHDHHDHD HDGGGSGGGG
   GGGSGGGSGG GSGGGPKYEG NIDIGLKGKV LTIGGADAET LMDAAVDVFA DGQPKLVSDQ
   AVSLGQNVLS ADEFTPGETYT VEVRFKEFGS VRAKVVAQLE HHHHHH

```

SEQ ID No. 7 shows the Linker sequence (GGGS)<sub>5</sub>GGG depicted in example 1. The glycine-rich flexible linker sequence is inserted between the two *E. coli* SlyD molecules and also between SlyD and the TpN47 antigen.

GGGSGGGSGG GSGGGSGGGS GGG

5

SEQ ID No. 8 shows TpN47 peptide *EcSlyD*-TpN47/p02-1, amino acids 30-66 of the *Treponema pallidum* 47 antigen sequence according to Swiss Prot P29723:

GYATLSYADY WAGELGQSRD VLLAGNAEAD RAGDLDA

10 SEQ ID No. 9 shows TpN47 peptide *EcSlyD*-TpN47/p03-1, amino acids 106-132 of the *Treponema pallidum* 47 antigen sequence according to Swiss Prot P29723:

SRGRKKWEYE TDPSTVKMVR ASASFQD

15 SEQ ID No. 10 shows TpN47 peptide *EcSlyD*-TpN47/p04-1, amino acids 137-170 of the *Treponema pallidum* 47 antigen sequence according to Swiss Prot P29723:

GEIKFEAVEG AVALADRASS FMVDSEYKI TNVK

SEQ ID No. 11 shows TpN47 peptide *EcSlyD*-TpN47/p05-1, amino acids 197-219 of the *Treponema pallidum* 47 antigen sequence according to Swiss Prot P29723:

20 EDSRVTTENTN GLKTMLTEDS FSA

SEQ ID No. 12 shows TpN47 peptide *EcSlyD*-TpN47/p06-1, amino acids 225-250 of the *Treponema pallidum* 47 antigen sequence according to Swiss Prot P29723:

MESPHDLVVD TVGTGYHSRF GSDAEA

25

SEQ ID No. 13 shows TpN47 peptide *EcSlyD*-TpN47/p07-1, amino acids 273-296 of the *Treponema pallidum* 47 antigen sequence according to Swiss Prot P29723:

NFNTVRYDYY GDDASYTNLM ASYG

30 SEQ ID No. 14 shows TpN47 peptide *EcSlyD*-TpN47/p08-1, amino acids 321-362 of the *Treponema pallidum* 47 antigen sequence according to Swiss Prot P29723:

FDRFKGSGPG YYRLTLIANG YRDVVADVRF LPKYEGNIDI GL

35 SEQ ID No. 15 shows TpN47 peptide *EcSlyD*-TpN47/p09-1, amino acids 368-388 of the *Treponema pallidum* 47 antigen sequence according to Swiss Prot P29723:

TIGGADAETL MDAAVDVFAD G

SEQ ID No. 16 shows TpN47 peptide *EcSlyD*-TpN47/p10-1, amino acids 391-434 of the *Treponema pallidum* 47 antigen sequence according to Swiss Prot P29723:

5 KLVSDQAVSL GQNVLSADFT PGTEYTVFVR FKEFGSVRAK VVAQ

#### Detailed description of the invention

- 10 The invention concerns soluble *Treponema pallidum* antigens, more precisely soluble variants of the *Treponema* antigen TpN47. The TpN47 antigens according to the invention lack domain C (amino acid residues 224 to 351 of SEQ ID NO.1 and are both stable and immunoreactive in an *in vitro* diagnostic immunoassay.
- 15 We succeeded in purifying to homogeneity full-length TpN47 from transformed prokaryotic host cells. Our initial experiments with the full-length version of TpN47 clearly revealed that this protein tends to aggregate when exposed to moderately elevated temperatures. Full-length TpN47 inevitably aggregated into a high-molecular-weight associate at temperatures above 35 °C, despite the fusion of highly solubilizing tandem chaperone fusions such as *EcSlyD*-*EcSlyD* or even (the more
- 20 thermostable) *EcSlpA*-*EcSlpA* to the full length antigen. In order to settle the stability problems posed by the full-length TpN47, we cloned and overproduced short TpN47 fragments as chaperone fusion proteins in *E.coli*, purified the fragments to homogeneity (as judged by Coomassie-stained SDS-PAGE gels) and assessed them for their respective antigenicity. In short, this approach was a complete failure. To our surprise, only one out of the ten fragments exhibited significant (albeit
- 25 rather poor) antigenicity (see table 2/ Fig. 8a, TpN47 30-66). The remainder of the promising short TpN47 fragments was not active at all. This finding was astonishingly at odds with the literature data given in Baughn *et al.*, J. Immunol. (1996) 157 (2), 720-731. As a consequence, we had to search for another way to circumvent the thermally induced aggregation of the full-length TpN47.
- 30 Instead of focusing on short and presumably unstructured peptide fragments of TpN47, we sought to design conformationally folded parts of TpN47. In the Journal of Biological Chemistry (2002), 277 (4), 41857-41864, Deka *et al.* present the crystal structure of TpN47 and reveal the domain topology of this protein. According to this work, TpN47 consists of four domains, *i.e.* TpN47 comprises four

autonomous folding units. However, Deka *et al.* are silent with regard to immunological features of the identified domains.

Surprisingly, we were able to successfully express the TpN47 domains and domain combinations  
5 AB, B, C, CD and D in an *E. coli* host. In one embodiment of the invention, the TpN47 antigens  
were produced in fusion with chaperone modules such as SlyD, FkpA, SlpA and Skp. All of these  
constructs were purified to homogeneity and assessed for their antigenicity with human anti-Syphilis  
sera in an automated Elecsys analyzer. The result was quite clear-cut: antigenicity was pretty high  
and increased in the order  $C < D < CD < B \ll AB$ . Interestingly, domain C could be identified as  
10 precarious in temperature stress assays (domain C and domain combination CD strongly aggregated  
upon incubation at temperatures  $> 35^{\circ}\text{C}$ , whereas AB, B and D remained perfectly soluble). Briefly,  
AB, B and D were identified as TpN47 fragments with slightly reduced antigenicity but markedly  
improved solubility when compared to full-length TpN47. Thus, the data of our design experiments  
provide compelling evidence that TpN47 variants lacking the C domain are significantly improved  
15 with respect to solubility and stability.

In one embodiment of the invention, the antigen therefore comprises domain B of the *Treponema pallidum* antigen 47 (TpN47) and lacks domain C of TpN47, *i.e.* the invention concerns a soluble  
*Treponema pallidum* antigen 47 (TpN47 antigen) comprising amino acid residues 63 to 174  
20 (domain B) of SEQ ID NO. 1 with the proviso that said antigen lacks amino acid residues 224 to  
351 (domain C) of SEQ ID NO. 1 and wherein the TpN47 antigen is fused with a chaperone

A further embodiment of the invention is a TpN47 antigen that comprises domains A and B and  
lacks domain C. This means that the invention covers a soluble TpN47 antigen comprising  
25 amino acid residues 26 to 223 (domain A+B) of SEQ ID NO. 1 with the proviso that said antigen  
lacks amino acid residues 224 to 351 (domain C) of SEQ ID NO. 1. In a further embodiment the  
TpN47 antigen comprises domain B or domains A+B, wherein domain D is an additional part of  
the TpN47 antigen. However, also in these embodiments domain C is missing. Domain D  
comprises amino acid residues 352 to 434 of SEQ ID NO. 1. Also in these embodiments the  
30 TpN47 antigen is fused with a chaperone.

An additional embodiment refers to a soluble *Treponema pallidum* antigen 47 (TpN47 antigen)  
comprising amino acid residues 63 to 174 (domain B) of SEQ ID NO. 1 with the proviso that

said antigen lacks amino acid residues 224 to 351 (domain C) of SEQ ID NO. 1, additionally comprising amino acid residues 352 to 434 of SEQ ID NO. 1 (domain D).

A further embodiment relates to a soluble *Treponema pallidum* antigen 47 (TpN47 antigen) comprising amino acid residues 26 to 223 (domain A+B) of SEQ ID NO. 1 with the proviso that said antigen lacks amino acid residues 224 to 351 (domain C) of SEQ ID NO. 1, additionally comprising amino acid residues 352 to 434 of SEQ ID NO. 1 (domain D).

According to the invention the term *Treponema* is the short form of the complete term for the organism *Treponema pallidum* – the pathogen causing Syphilis - and both terms can be used interchangeably.

A “TpN47 antigen” is a protein containing a TpN47 amino acid sequence that is suitable as an antigen for use in an immunological assay. This means that the antigens according to the invention are soluble under physiological buffer conditions, for example in a phosphate buffer system at ambient temperature without addition of detergents. The antigens are also capable of binding to or being recognized and bound by antibodies specific for TpN47, like e.g. anti-TpN47 antibodies present in an isolated sample such as human sera.

According to the invention also variants of the TpN47 antigens are included. The term “variants” in this context relates to a protein or a protein fragment (i.e. a polypeptide or peptide) substantially similar to said protein. In particular, a variant may be an isoform which shows amino acid exchanges, deletions or insertions compared to the amino acid sequence of the most prevalent protein isoform. In one embodiment, such a substantially similar protein has a sequence similarity to the most prevalent isoform of the protein of at least 80%, in another embodiment at least 85% or at least 90%, in yet another embodiment at least 95%. The term “variant” also relates to a post-translationally modified protein such as a glycosylated or phosphorylated protein. According to the invention a variant classifies as a TpN47 antigen variant as long as the immunoreactivity in an *in vitro* diagnostic immunoassay is maintained, i.e. the variant is still able to bind and detect anti-TpN47 antibodies present in a sample. A “variant” is also a protein or antigen which has been modified for example by covalent or non-covalent attachment of a label or carrier moiety to the protein or antigen. Possible labels are radioactive,

fluorescent, chemiluminescent, electrochemiluminescent, enzymes or others *e.g.* like digoxigenin or biotin. These labels are known to a person skilled in the art.

The antigens according to the invention can be fused to another protein. The term “fusion protein” as used in the present invention refers to a protein comprising at least one protein part corresponding to a TpN47 antigen according to the invention and at least one protein part derived from another protein that serves the role of a fusion partner.

According to another embodiment of the current invention the TpN47 antigens can be fused to chaperones. Chaperones, which are known as classical folding helpers, are proteins that assist the folding and maintenance of the structural integrity of other proteins. Examples of folding helpers are described in detail in WO 03/000877. According to the invention chaperones of the peptidyl prolyl isomerase class such as chaperones of the FKBP family can be used for fusion to the TpN47 antigen. Examples of FKBP chaperones suitable as fusion partners for TpN47 antigens are FkpA, SlyD and SlpA. A further chaperone suitable as a fusion partner for TpN47 is Skp, a periplasmic trimeric chaperone from *E.coli*, not belonging to the FKBP family. It is not always necessary to use the complete sequence of a chaperone. Functional fragments of chaperones (so-called binding-competent modules) which still possess the required abilities and functions may also be used (cf. WO 98/13496).

According to a further embodiment of the invention at least one or at least two modules of an FKBP chaperone such as *e.g.* *E. coli* SlyD, SlpA or FkpA are used as fusion moieties for expression of the TpN47 antigens. The chaperone Skp may be used as a fusion partner as well. The fusion of two FKBP-chaperone domains results in improved solubility of the resulting fusion polypeptide. The fusion moieties may be located N-terminally or C-terminally or at both ends (sandwich-like) of the TpN47 antigen.

The TpN47 antigens according to the invention can be generated and prepared by means of recombinant DNA techniques. Another aspect of the invention therefore is a recombinant DNA molecule encoding a TpN47 antigen and variants thereof as defined further above.

The term “recombinant DNA molecule” refers to a molecule which is made by the combination of two otherwise separated segments of sequence accomplished by the artificial manipulation of

isolated segments of polynucleotides by genetic engineering techniques or by chemical synthesis. In doing so one may join together polynucleotide segments of desired functions to generate a desired combination of functions. Recombinant DNA techniques for expression of proteins in prokaryotic or lower or higher eukaryotic host cells are well known in the art. They have been described *e.g.* by Sambrook *et al.*, (1989, Molecular Cloning: A Laboratory Manual)

A further subject matter of the present invention relates to a recombinant DNA molecule, encoding a *Treponema pallidum* antigen 47 (TpN47), comprising a nucleotide sequence coding for a TpN47 antigen comprising an amino acid sequence according to amino acid residues 63 to 174 (domain B) of SEQ ID NO. 1 with the proviso that said recombinant DNA molecule lacks the coding region for amino acid residues 224 to 351 (domain C) of SEQ ID NO. 1.

In another embodiment of the invention a recombinant DNA molecule encodes a TpN47 antigen comprising an amino acid sequence according to amino acid residues 26 to 223 (domain A+B) of SEQ ID NO. 1 with the proviso that said recombinant DNA molecule lacks the coding region for amino acid residues 224 to 351 (domain C) of SEQ ID NO. 1.

In a further embodiment of the invention a recombinant DNA molecule encodes a TpN47 antigen comprising an amino acid sequence according to amino acid residues 63 to 174 (domain B) or amino acid residues 26 to 223 (domain A+B) of SEQ ID NO. 1. In addition, the recombinant DNA molecule encodes an amino acid sequence comprising domain D, *i.e.* amino acid residues 352 to 434 of SEQ ID NO. 1. As described before, also in these embodiments the proviso applies that said recombinant DNA molecules lack the coding region for amino acid residues 224 to 351 (domain C) of SEQ ID NO. 1

The recombinant DNA molecules according to the invention may also contain sequences encoding linker peptides of 10 to 100 amino acid residues in between the TpN47 antigen and the fusion moieties and also between several fusion moieties. Such a linker sequence may for example harbor a proteolytic cleavage site.

A further aspect of the invention is an expression vector comprising operably linked a recombinant DNA molecule according to the present invention, *i.e.*, a recombinant DNA molecule encoding a TpN47 antigen or a recombinant DNA molecule encoding a fusion protein

comprising a TpN47 antigen and a peptidyl prolyl isomerase chaperone, such as an FKBP-chaperone, wherein the FKBP-chaperone is selected from FkpA, SlyD and SlpA. In an alternative embodiment the recombinant DNA molecule encodes a fusion protein comprising a TpN47 antigen and Skp. The expression vector comprising a recombinant DNA according to the present invention may be used to express the TpN47 antigen in a cell free translation system or may be used to transform a host cell for expression of the TpN47 antigen according to methods well known in the art. Another aspect of the invention therefore relates to a host cell transformed with an expression vector according to the present invention. In one embodiment of the current invention the recombinant TpN47 antigens are produced in *E. coli* cells.

Also contemplated is a method for producing a soluble, stable and immunoreactive TpN47 antigen, that can also be produced as a fusion protein containing the TpN47 antigen and a chaperone such as Skp or a peptidyl prolyl isomerase class chaperone such as an FKBP chaperone. In a further embodiment of the invention said FKBP chaperone is selected from the group consisting of SlyD, FkpA and SlpA.

This method comprises the steps of

- a) culturing host cells transformed with the above-described expression vector containing a gene encoding a TpN47 antigen
- b) expression of the gene encoding said TpN47 antigen
- c) purification of said TpN47 antigen.

Optionally, as an additional step d), functional solubilization needs to be carried out so that the TpN47 antigen is brought into a soluble and immunoreactive conformation by means of refolding techniques known in the art.

An additional aspect of the present invention concerns a method for the detection of anti-*Treponema* antibodies in an isolated human sample wherein a TpN47 antigen according to the invention is used as a binding partner for the antibodies. The invention thus covers a method for the detection of antibodies specific for *Treponema* in an isolated sample, said method comprising

- a) forming an immunoreaction admixture by admixing a body fluid sample with a TpN47 antigen according to the invention
- b) maintaining said immunoreaction admixture for a time period sufficient for allowing antibodies against said TpN47 antigen present in the body fluid sample to immunoreact with said

TpN47 antigen to form an immunoreaction product; and

c) detecting the presence and/or the concentration of any of said immunoreaction product.

In a further aspect said method is suitable for detecting *Treponema* antibodies of the IgG and the  
5 IgM subclass.

Immunoassays for detection of antibodies are well known in the art, and so are methods for carrying out such assays and practical applications and procedures. The TpN47 antigens according to the invention can be used to improve assays for the detection of anti-*Treponema*  
10 antibodies independently of the labels used and independently of the mode of detection (*e.g.*, radioisotope assay, enzyme immunoassay, electrochemiluminescence assay, *etc.*) or the assay principle (*e.g.*, test strip assay, sandwich assay, indirect test concept or homogenous assay, *etc.*). All biological liquids known to the expert can be used as samples for the detection of anti-*Treponema* antibodies. The samples usually used are bodily liquids like whole blood, blood sera,  
15 blood plasma, urine or saliva.

A further embodiment of the invention is an immunoassay for detecting anti-*Treponema* antibodies in an isolated sample performed according to the so-called double antigen sandwich concept (DAGS). Sometimes this assay concept is also termed double antigen bridge concept,  
20 because the two antigens are bridged by an antibody analyte. In such an assay the ability of an antibody to bind at least two different molecules of a given antigen with its two (IgG, IgA, IgE) or ten (IgM) paratopes is required and utilized.

In more detail, an immunoassay for the determination of anti-*Treponema* antibodies according to  
25 the double antigen bridge format is carried out by incubating a sample containing the anti-*Treponema* antibodies with two different TpN47 antigens, *i.e.* a first ("solid phase") TpN47 antigen and a second TpN47 ("detection") antigen, wherein each of the said antigens binds specifically to said anti-*Treponema* antibodies. The first antigen can be bound directly or indirectly to a solid phase and usually carries an effector group which is part of a bioaffine  
30 binding pair like, *e.g.*, biotin and avidin. For example, if the first antigen is conjugated to biotin the solid phase is coated with either avidin or streptavidin. The second antigen carries a label. Thus an immunoreaction admixture is formed comprising the first antigen, the sample antibody and the second antigen. A solid phase to which the first antigen can be bound is added either

before the addition of the sample to said antigens or after the immunoreaction admixture is formed. This immunoreaction admixture is maintained for a time period sufficient for allowing anti- *Treponema* antibodies against said TpN47 antigens in the body fluid sample to immunoreact with said TpN47 antigens to form an immunoreaction product. Next step is a separation step wherein the liquid phase is separated from the solid phase. Finally, the presence of any of said immunoreaction product is detected in the solid or liquid phase or both.

In said DAGS immunoassay the basic structures of the “solid phase antigen” and the “detection antigen” are essentially the same. It is also possible to use, in a double antigen bridge assay, similar but different TpN47 antigens, which are immunologically cross-reactive. The essential requirement for performing such assays is that the relevant epitope or the relevant epitopes are present on both antigens. According to the invention it is possible to use different fusion moieties for each TpN47 antigen (e.g. SlyD fused to TpN47 on the solid phase side and FkpA fused to TpN47 on the detection side) as such variations significantly alleviate the problem of non-specific binding and thus mitigate the risk of false-positive results.

A further embodiment of the present invention is therefore an immunoassay according to the double antigen bridge concept wherein a first TpN47 antigen or fusion protein according to the present invention, and a second TpN47 antigen or fusion protein according to the present invention are used.

The present invention further relates to the use of at least one antigen of TpN47 in a diagnostic test for the detection of anti-*Treponema* antibodies.

An additional subject matter of the invention is a reagent kit for the detection of antibodies against *Treponema*, containing, in addition to the usual test additives for immunoassays, at least one antigen of the TpN47 antigens suitable for specifically binding to *Treponema* antibodies to be determined and possibly carrying a label as well as other usual additives if necessary. In particular the reagent kit contains a TpN47 antigen comprising amino acid residues 63 to 174 (domain B) of SEQ ID NO. 1 or a TpN47 antigen comprising amino acid residues 26 to 223 (domain A+B) of SEQ ID NO. 1, with the proviso that each of said antigens lacks sequences corresponding to amino acid residues 224 to 351 (domain C) of SEQ ID NO. 1. The antigens being part of said reagent kit are fused to a chaperone.

In a further embodiment said reagent kit comprises a TpN47 antigen comprising domains B or A+B as defined before and additionally comprising domain D, *i.e.* amino acid residues 352-434 of SEQ ID NO. 1. Also in this embodiment the TpN47 antigen lacks domain C, *i.e.* amino acid residues 224-351 are not present in this TpN47 antigen.

In addition, the reagent kits defined above contain controls and standard solutions as well as reagents in one or more solutions with the common additives, buffers, salts, detergents etc. as used by the average man skilled in the art.

10

Another aspect is the use of the TpN47 antigens according to the invention as vaccines. The preparation of vaccines which contain an immunogenic polypeptide as active ingredient is known in the art. Such vaccines are commonly prepared as injectables, either as liquid solutions or suspensions. The active ingredient, *i.e.* the TpN47 antigen or its fusion protein is mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient like for example water, aqueous physiological buffers, saline, dextrose, glycerol, ethanol. The vaccines are conventionally administered parentally, by injection.

Another embodiment is a composition of at least two *Treponema pallidum* antigens comprising a TpN47 antigen according to the current invention and at least one additional *Treponema pallidum* antigen selected from the group consisting of a TpN17 antigen and a TpN15 antigen so that said composition comprises a TpN47 antigen, a TpN17 antigen or a TpN15 antigen or both TpN15 and TpN17 antigens. Yet a further embodiment is a composition of at least three *Treponema pallidum* antigens comprising a TpN47 antigen according to the current invention and both a TpN17 and a TpN15 antigen.

The invention also concerns the use of a TpN47 antigen according to the invention in an *in vitro* diagnostic test for the detection of anti-*Treponema pallidum* antibodies.

The examples section further illustrates the invention. In particular, the examples illustrate that we have developed and generated variants of TpN47 that are more soluble and significantly less thermolabile than the full-length TpN47 protein molecule. Both the solubility and the stability are improved. Our TpN47 variants can be abundantly overexpressed for example in *E. coli*, are

easily purified and refolded via immobilized metal chelate chromatography (IMAC), exhibit satisfying stability properties and may be used to reliably detect anti-*Treponema* antibodies in human sera (in a further embodiment in combination with TpN17 and/or TpN15, two other immunodominant membrane proteins from *Treponema pallidum*). It is noteworthy that the

5 FkpA-TpN47/AB and Skp-TpN47/AB fusion proteins form natural oligomers with epitope densities that are sufficient to detect even IgM molecules. Since we aim at developing an immunoassay for detection of total immunoglobulin (*i.e.* detection of both IgG and IgM), the oligomeric species FkpA-TpN47/AB and Skp-TpN47/AB may be used advantageously as

10 specifiers on both sides of a DAGS format (e.g. FkpA-TpN47/AB-biotin and Skp-TpN47/AB-ruthenium, or vice versa).

### Example 1

#### 15 Cloning and purification of TpN47 and TpN47 chaperone fusion polypeptides

##### Cloning of expression cassettes

On the basis of the pET24a expression plasmid of Novagen (Madison, WI, USA) expression

20 cassettes encoding *EcSlyD-EcSlyD-TpN47* fusion proteins were obtained essentially as described (Scholz, C. *et al.*, J. Mol. Biol. (2005) 345, 1229-1241). The sequence of the TpN47 antigen was retrieved from the SwissProt database (SwissProt ID P29723). A synthetic gene encoding mature TpN47 aa 21-434 (the signal peptide spanning amino acid residues 1-20 was omitted) with a glycine-rich linker region fused in frame to the N-terminus was purchased from

25 Medigenomix (Martinsried, Germany). The unique cysteine residue of TpN47 at position 315 was changed to alanine in order to prevent unwanted side-effects such as oxidation or intermolecular disulfide bridging. *BamHI* and *XhoI* restriction sites were at the 5' and the 3' ends of the TpN47-coding region, respectively. A further synthetic gene encoding two *EcSlyD* units (residues 1-165 according to SEQ ID NO. 1, SwissProt accession no. [P0A9K9](#)) connected via a

30 glycine-rich linker region and encompassing part of a further linker region at the C-terminus were likewise purchased from Medigenomix. *NdeI* and *BamHI* restriction sites were at the 5' and 3' ends of this cassette, respectively. The genes and the restriction sites were designed to enable the in frame fusion of the chaperone part *EcSlyD-EcSlyD* and the TpN47 antigen part by simple

ligation. In order to avoid inadvertent recombination processes and to increase the genetic stability of the expression cassette in the *E. coli* host, the nucleotide sequences encoding the *EcSlyD* units were degenerated as were the nucleotide sequences encoding the extended linker regions. *i.e.*, different codon combinations were used to encode identical amino acid sequences.

5


The pET24a vector was digested with *NdeI* and *XhoI* and the cassette comprising tandem-SlyD fused in frame to the *Treponema* TpN47 fragment 21-434 (Cys 315 Ala) was inserted.

Expression cassettes comprising SlyD or Skp or FkpA were constructed accordingly, as well as expression cassettes comprising target polypeptides different from full-length TpN47, notably the domains and domain combinations B (TpN47 63-174), AB (TpN47 26-223), C (TpN47 224-351), CD (TpN47 224-434) and D (TpN47 352-434). All recombinant fusion polypeptide variants contained a C-terminal hexahistidine tag to facilitate Ni-NTA-assisted purification and refolding. QuikChange (Stratagene, La Jolla, CA, USA) and standard PCR techniques were used to generate point mutations, deletion, insertion and extension variants or restriction sites in the

15 respective expression cassettes.

The drawing below shows a scheme of the *Treponema* TpN47 full length antigen 21-434 bearing two SlyD chaperone units fused in frame to its N-terminal end. To denote the *E. coli* origin of the SlyD fusion partner, the depicted fusion polypeptide has been named *EcSlyD-EcSlyD-*

20 TpN47 (21-434).



25

L = (GGGS)<sub>5</sub>GGG-Linker

The insert of the resulting plasmid was sequenced and found to encode the desired fusion protein.

The complete amino acid sequence of each individual TpN47 antigen is shown in SEQ ID NOs.

1 to 6. The amino acid sequence of the linker L is shown in SEQ ID NO. 7. Purification of fusion proteins comprising TpN47

30

All TpN47 fusion protein variants were purified by using virtually identical protocols. *E. coli* BL21 (DE3) cells harboring the particular pET24a expression plasmid were grown at 37°C in

LB medium plus kanamycin (30 µg/ml) to an OD<sub>600</sub> of 1.5, and cytosolic overexpression was induced by adding 1 mM isopropyl-β-D-thiogalactoside. Three hours after induction, cells were harvested by centrifugation (20 min at 5000 g), frozen and stored at -20°C. For cell lysis, the frozen pellet was resuspended in chilled 50 mM sodium phosphate pH 8.0, 7.0 M GdmCl, 5 mM imidazole and the suspension was stirred for 2 h on ice to complete cell lysis. After centrifugation and filtration (0.45 µm/0.2 µm), the crude lysate was applied onto a Ni-NTA column equilibrated with the lysis buffer including 5.0 mM TCEP. The subsequent washing step was tailored for the respective target protein and ranged from 5 to 15 mM imidazole (in 50 mM sodium phosphate pH 8.0, 7.0 M GdmCl, 5.0 mM TCEP). At least 10-15 volumes of the washing buffer were applied. Then, the GdmCl solution was replaced by 50 mM potassium phosphate pH 8.0, 100 mM KCl, 10 mM imidazole, 5.0 mM TCEP to induce conformational refolding of the matrix-bound protein. In order to avoid reactivation of copurifying proteases, a protease inhibitor cocktail (Complete® EDTA-free, Roche) was included in the refolding buffer. A total of 15-20 column volumes of refolding buffer were applied in an overnight reaction. Then, both TCEP and the Complete® EDTA-free inhibitor cocktail were removed by washing with 3-5 column volumes 50 mM potassium phosphate pH 8.0, 100 mM KCl, 10 mM imidazole. Subsequently, the imidazole concentration – still in 50 mM potassium phosphate pH 8.0, 100 mM KCl – was raised to 25 mM in order to remove unspecifically bound protein contaminants. The native protein was then eluted by 500 mM imidazole in the same buffer. Protein-containing fractions were assessed for purity by Tricine-SDS-PAGE and pooled. Finally, the proteins were subjected to size-exclusion-chromatography (Superdex HiLoad, Amersham Pharmacia) and the protein-containing fractions were pooled and concentrated to 10-20 mg/ml in an Amicon cell (YM10).

After the coupled purification and refolding protocol, protein yields of roughly 5-20 mg could be obtained from 1 g of *E. coli* wet cells, depending on the respective target protein.

Example 2Spectroscopic measurements

- 5 Protein concentration measurements were performed with an Uvikon XL double-beam spectrophotometer. The molar extinction coefficients ( $\epsilon_{M280}$ ) were determined by using the procedure described by Pace (1995), Protein Sci. 4, 2411-2423. The molar extinction coefficients ( $\epsilon_{M280}$ ) used for the distinct Tpn47 fusion polypeptides are specified in table 1.

10

Table 1: Protein parameters of the Tpn47 fusion variants used in this study

| fusion protein                | TpN47<br>Fragment<br>(aa residues) | molecular<br>weight<br>(Da) | pI  | $\epsilon_{M280}$<br>$M^{-1}cm^{-1}$ | Abs 0.1%<br>(= 1 mg/ml) |
|-------------------------------|------------------------------------|-----------------------------|-----|--------------------------------------|-------------------------|
| <b>TpN47 domain variants</b>  |                                    |                             |     |                                      |                         |
| <i>EcSlyD-EcSlyD-TpN47</i>    | 21-434                             | 85390                       | 4.9 | 63720                                | 0.746                   |
| <i>EcSlyD-EcSlyD-TpN47/AB</i> | 26-223                             | 61641                       | 4.8 | 33350                                | 0.541                   |
| <i>EcSlyD-EcSlyD-TpN47/B</i>  | 63-174                             | 52267                       | 4.8 | 21890                                | 0.419                   |
| <i>EcSlyD-EcSlyD-TpN47/D</i>  | 352-434                            | 48561                       | 4.7 | 14900                                | 0.307                   |
| <i>EcSlyD-EcSlyD-TpN47/C</i>  | 224-351                            | 52249                       | 4.9 | 39310                                | 0.725                   |
| <i>EcSlyD-EcSlyD-TpN47/CD</i> | 224-434                            | 62976                       | 4.8 | 42290                                | 0.672                   |
| <b>TpN47 peptide variants</b> |                                    |                             |     |                                      |                         |
| <i>EcSlyD-TpN47/p02-1</i>     | 30-66                              | 24382                       | 4.7 | 15930                                | 0.653                   |
| <i>EcSlyD-TpN47/p03-1</i>     | 106-132                            | 23667                       | 5.1 | 12950                                | 0.547                   |
| <i>EcSlyD-TpN47/p04-1</i>     | 137-170                            | 24211                       | 4.8 | 7450                                 | 0.308                   |
| <i>EcSlyD-TpN47/p05-1</i>     | 197-219                            | 23052                       | 4.8 | 5960                                 | 0.259                   |
| <i>EcSlyD-TpN47/p06-1</i>     | 225-250                            | 23284                       | 4.8 | 7450                                 | 0.320                   |
| <i>EcSlyD-TpN47/p07-1</i>     | 273-296                            | 23307                       | 4.8 | 13410                                | 0.575                   |
| <i>EcSlyD-TpN47/p08-1</i>     | 321-362                            | 25274                       | 5.0 | 11920                                | 0.472                   |
| <i>EcSlyD-TpN47/p09-1</i>     | 368-388                            | 22545                       | 4.7 | 5960                                 | 0.264                   |
| <i>EcSlyD-TpN47/p10-1</i>     | 391-434                            | 25279                       | 5.0 | 7450                                 | 0.295                   |

The amino acid sequences of the TpN47 domain variants are shown in SEQ ID No. 1 to 6. The TpN47 specific sequences derived from Swiss Prot P29723 of the TpN47 peptide variants p02-1 to p10-1 are summarized in SEQ ID NOs. 8 to 16.

5

### Example 3

#### Coupling of biotin and ruthenium moieties to the fusion proteins

10 The lysine  $\epsilon$ -amino groups of the fusion polypeptides were modified at protein concentrations of 10-30 mg/ml with N-hydroxy-succinimide activated biotin and ruthenium label molecules, respectively. The label/protein ratio varied from 2:1 to 5:1 (mol:mol), depending on the respective fusion protein. The reaction buffer was 150 mM potassium phosphate pH 8.0, 100 mM KCl, 1 mM EDTA. The reaction was carried out at room temperature for 15 min and  
15 stopped by adding buffered L-lysine to a final concentration of 10 mM. To avoid hydrolytic inactivation of the labels, the respective stock solutions were prepared in dried DMSO (seccosolv quality, Merck, Germany). DMSO concentrations up to 20% in the reaction buffer were well tolerated by all fusion proteins studied. After the coupling reaction, unreacted free label was removed by passing the crude protein conjugate over a gel filtration column (Superdex  
20 200 HiLoad).

### Example 4

#### Immunological reactivity of the polypeptide fusion proteins

The immunological reactivity (*i.e.* the antigenicity) of the different fusion proteins was assessed in an automated Elecsys® 2010 analyzer (Roche Diagnostics GmbH). Elecsys® is a registered trademark of the Roche group. Measurements were carried out in the double antigen sandwich  
30 format.

Signal detection in Elecsys® 2010 is based on electrochemoluminescence. The biotin-conjugate (*i.e.* the capture-antigen) is immobilized on the surface of a streptavidin coated magnetic bead

whereas the detection-antigen bears a complexed Ruthenium cation (switching between the redox states 2+ and 3+) as the signaling moiety. In the presence of a specific immunoglobulin analyte, the chromogenic ruthenium complex is bridged to the solid phase and emits light at 620 nm after excitation at a platinum electrode. The signal output is in arbitrary light units.

5

The recombinant *Treponema* TpN47 variants were assessed pairwise in a double antigen sandwich (DAGS) immunoassay format. For instance, an *EcSlyD-EcSlyD*-TpN47/AB (26-223)-biotin conjugate was assessed together with an *EcSlyD-EcSlyD*-TpN47/AB (26-223)-ruthenium complex conjugate at a concentration of 70 ng/ml each. As well, an *EcSlyD-EcSlyD*-TpN47/B (63-174)-biotin conjugate was applied together with an *EcSlyD-EcSlyD*-TpN47/B (63-174)-ruthenium complex conjugate at a concentration of 70 ng/ml each .

The biotin and the ruthenium conjugates of the fusion polypeptide variants of TpN47 were applied at concentrations of 70 ng/ml each. In all measurements, chemically polymerized and unlabeled *EcSlyD-EcSlyD* was implemented in large excess (~ 10 µg/ml) in the reaction buffer as an anti-interference substance to avoid immunological cross reactions via the chaperone fusion unit. Anti-*Treponema* negative human sera were used as controls.

In table 2 (Fig. 8a-c), the immunological activity of the TpN47 peptide fusion variants (listed in table 1) is shown. It is obvious at first glance that the antigenicity of the short TpN47 fragments is very poor when compared to the full-length TpN47 molecule. Only *EcSlyD*-TpN47/p02-1 (TpN47 30-66) exhibits significant antigenicity, albeit to a very low extent. From our results we conclude that some weak linear epitopes may reside in the very N-terminal part of TpN47, whereas the remainder of the TpN47 molecule does not harbor any linear epitopes detectable in our DAGS setup. This finding is strikingly at odds with literature data reporting on immunodominant short epitopes within the *Treponema* antigen TpN47 (Baughn *et al.*, Journal of Immunology (1996) Jul 15;157(2):720-31). Contrasting with these literature data, our experimental findings with short TpN47 fragments suggest that linear epitopes do play a rather subordinate role in the humoral immune response following *Treponema* infection.

30

As a direct consequence, we abandoned further attempts to identify linear epitopes and focused on the identification of conformational epitopes instead. In order to attain this goal, we targeted the TpN47 domains disclosed in Ranjit *et al.*, J. Biol. Chem. (2002) 277 (44), pp 41857-41864).

Different from unstructured short peptides, isolated domains (*i.e.* autonomous folding units) of a protein are supposed to adopt a defined conformation and thus are expected to present conformational epitopes. It was, however, totally unclear whether the isolated TpN47 domains would be able to adopt a native-like conformation when excised from the structural context of the full-length protein. Indeed, it turned out that the isolated TpN47 domains exhibit a tremendously high immunological activity when compared to the small unstructured TpN47 fragments (for results see table 3, figure 9). From these data, we infer that the isolated domains are indeed able to adopt a well-ordered, native-like conformation. As judged from our immunological assessments, the antigenicity of the domain fragments increases in the order C<D<CD<B<AB<ABCD; notably, the TpN47 domain combination AB yields about 50% of the signal level of the full-length protein with anti-*Treponema*-positive human sera.

#### Example 5

##### Thermostability of TpN47 domain fusions as assessed by FPLC analysis

Having gathered compelling evidence that the TpN47 domains (*i.e.* the well-ordered autonomous folding units with defined conformation) exhibit significant antigenicity, we wondered whether the distinct domains would possess different stability when exposed to thermal stress. To address this question, we incubated all of our TpN47 domain fusion proteins under identical conditions and subjected them to elevated temperatures (overnight incubation in 150 mM potassium phosphate pH 8.0, 100 mM KCl, 0.5 mM EDTA at 30°C, 35°C and 40°C, at a protein concentration of 1.3 mg/ml each). Then, we assessed all of the samples by analytical size exclusion chromatography (Superdex 200) and checked both the signal recovery (peak area under absorption curve) and the tendency to aggregate (*i.e.* elution of large particles in the void volume of the gel filtration column) for each TpN47 fusion protein. The results are shown in Figures 1-6. It turned out that the tendency to form temperature-induced aggregates significantly decreases in the order full-length-TpN47> CD > C > D, B, AB. In brief, the TpN47 domains D, B and AB are much less prone to aggregation than is full-length TpN47. Upon thermal stress, they invariably show excellent signal recovery in their elution profiles and exhibit only a negligible tendency to form associates or aggregates. Conversely, all TpN47 variants comprising

domain C (*i.e.* full-length TpN47, C, CD) exhibit a strong tendency to aggregate even at a moderately elevated temperature such as 35°C.

## 5 Example 6

### Thermostability of TpN47 domain fusions in immunoassays as assessed by Elecsys® measurements.

- 10 To ascertain the thermotolerance of the distinct TpN47 fusion proteins by means of Elecsys® measurements, the *EcSlyD-EcSlyD-TpN47* variants were subjected to elevated temperature conditions as follows: *EcSlyD-EcSlyD-TpN47* biotin and ruthenium conjugates were incubated, separately, at 42 °C for three days. The concentration of the conjugates in this stress assay was 70 ng/ml each (~ 1 nM), the assay buffer was 100 mM MES pH 6.5, 300 mM NaCl, 2 mM
- 15 EDTA. Subsequently, the thermally stressed samples were assessed for their residual immunological reactivity (*i.e.* their residual antigenicity) in an Elecsys® 2010 automated analyzer under the experimental conditions described above. Unchallenged samples (stored at 2-8 °C) of *EcSlyD-EcSlyD-TpN47* were used as a reference.
- 20 The outcome of the experiments is shown in tables 4-6 (figures 10-12).

Table 4 (figure 10) depicts the immunological reactivity of *EcSlyD-EcSlyD-TpN47/AB* (26-223) and *EcSlyD-EcSlyD-TpN47/B* (63-174) with human anti-*Treponema* positive and anti-*Treponema* negative sera in an automated Elecsys® analyzer as described. Shown is the

25 performance of both antigen variants before and after a harsh three-days-incubation at 42 °C. Table 5 (figure 11) depicts the antigenicity of *EcSlyD-EcSlyD-TpN47 C* (224-351) and *EcSlyD-EcSlyD-TpN47 D* (352-434), and table 6 (figure 12) displays the antigenicity of *EcSlyD-EcSlyD-TpN47 CD* (224-434) and of full-length TpN47 (21-434).

- 30 The outcome of the experiments clearly demonstrates the superiority of heat-stressed TpN47 domains over the full-length TpN47 protein in terms of signal recovery.

Upon thermal challenge, signal recovery of full-length TpN47 (21-434) drops to roughly 50 % of the initial values, whereas the signal recovery of TpN47/B, TpN47/AB and TpN47/D amounts to ~93 %, ~88 % and 100 % of the initial values, respectively. Thus, the signal recovery is markedly enhanced when using domains of TpN47 instead of the full-length molecule.

5

It is noteworthy that the C domain of TpN47 (224-351) does not exhibit any immunological activity (for results see table 5, figure 11). This finding clearly contrasts with the antigenicity found for the other TpN47 domains and the domain combinations B, AB, CD and D. Seemingly, the C domain does – at least in isolation - contribute little or nothing to the astonishing  
10 antigenicity of the TpN47 protein molecule.

The CD fusion variant of TpN47 (224-434) is remarkable in that its signal recovery following thermal stress amounts to roughly 70% and is clearly inferior to the other TpN47 domains and domain combinations (for results see table 6, figure 12). The signal recovery of the D domain  
15 alone is very high and almost unchanged after thermal stress (see table 5, figure 11).

To sum up, the TpN47 domains B, AB and D show a clearly enhanced signal recovery upon thermal challenge when compared to full-length TpN47.

20 According to our invention, domain C (224-351) is dispensable for immunodiagnostic purposes since it does not conspicuously contribute to TpN47 antigenicity. Furthermore, domain C, when fused to domain D, weakens the stability of the construct CD (224-434), which exhibits – exempt from full-length TpN47– the lowest signal recovery of all domains and domain combinations tested (~ 70 %, for results see table 6, figure 12).

25

The relative signal yield (upon thermal stress) of the distinct TpN47 domains as assessed by an automated immunoassay such as Elecsys® correlates nicely with our findings in FPLC analysis. This is remarkable all the more since both experiments have been carried out at very different concentrations: The protein concentrations in the FPLC analyses were in the medium  
30 micromolar range (15.2  $\mu$ M – 26.7  $\mu$ M), whereas the protein concentrations in the immunological analyses were in the very low nanomolar range (0.82 nM – 1.44 nM). It is expected that removal of an aggregation-inducing domain such as TpN47 domain C should yield best results (*i.e.* alleviate aggregation effects) under conditions of high protein concentration.

Our immunological data unequivocally show that removal of domain C clearly improves both the stability and the solubility of the remainder TpN47 molecule, even under conditions of very low protein concentration. This finding enables the development of more robust immunoassay kits and constitutes a major achievement in TpN47-based serological detection of anti-

5 *Treponema* antibodies.

Our experiments provide compelling evidence that full-length TpN47 is extremely prone to aggregation when exposed to moderately elevated temperatures above 35 °C. From these observations, we infer that the use of full-length recombinant TpN47 is critical to the specificity  
10 and the sensitivity of any *Treponema* immunoassay, unless precautions are taken to avoid the thermally-induced loss of this aggregation-prone molecule from the assay mixture. A simple and convenient way to circumvent (or at least to mitigate) thermally-induced aggregation of TpN47 is disclosed in this patent application: it consists in simply omitting the TpN47 domain C, which apparently does not contribute directly to antigenicity and which, on top of that, seems to  
15 constitute a generally destabilizing factor within the TpN47 molecule. As soon as the TpN47 domain C is left out (as in AB, B and D), thermolability of the TpN47 protein molecule is significantly mitigated.

Claims

1. A soluble *Treponema pallidum* antigen 47 (TpN47 antigen) comprising amino acid residues 63 to 174 (domain B) of Swiss Prot P29723 with the proviso that said antigen lacks amino acid residues 224 to 351 (domain C) of Swiss Prot P29723, wherein the TpN47 antigen is fused with a chaperone.  
5
2. A soluble *Treponema pallidum* antigen 47 (TpN47 antigen) comprising amino acid residues 26 to 223 (domain A+B) of Swiss Prot P29723 with the proviso that said antigen lacks amino acid residues 224 to 351 (domain C) of Swiss Prot P29723, wherein the TpN47 antigen is fused with a chaperone.  
10
3. A soluble *Treponema pallidum* antigen 47 (TpN47 antigen) according to claim 1 or 2 (domain B or A+B) additionally comprising amino acid residues 352 to 434 of Swiss Prot P29723 (domain D).  
15
4. A soluble *Treponema pallidum* antigen 47 (TpN47 antigen) comprising amino acid residues 63 to 174 (domain B) of Swiss Prot P29723 with the proviso that said antigen lacks amino acid residues 224 to 351 (domain C) of Swiss Prot P29723, additionally comprising amino acid residues 352 to 434 of Swiss Prot P29723 (domain D).  
20
5. A soluble *Treponema pallidum* antigen 47 (TpN47 antigen) comprising amino acid residues 26 to 223 (domain A+B) of Swiss Prot P29723 with the proviso that said antigen lacks amino acid residues 224 to 351 (domain C) of Swiss Prot P29723, additionally comprising amino acid residues 352 to 434 of Swiss Prot P29723 (domain D).  
25
6. A recombinant DNA molecule encoding a *Treponema pallidum* antigen 47 (TpN47 antigen) according to any of claims 1 to 5.
7. An expression vector comprising operably linked a recombinant DNA molecule according to claim 6.  
30

8. A host cell transformed with an expression vector according to claim 7.
9. A method of producing a soluble and immunoreactive *Treponema pallidum* antigen 47 (TpN47 antigen), said method comprising the steps of
  - 5 - culturing host cells according to claim 8
  - expression of said TpN47 antigen and
  - purification of said TpN47 antigen.
10. A composition of at least two *Treponema pallidum* antigens comprising a TpN47 antigen  
10 according to claims 1 to 5 and at least one additional *Treponema pallidum* antigen  
selected from the group consisting of a TpN17 antigen and a TpN15 antigen.
11. A method for detecting antibodies specific for *Treponema pallidum* in an isolated sample  
15 wherein a TpN47 antigen according to claims 1 to 5 or a combination or composition of  
antigens according to claim 10 is used as a capture reagent and/or as a binding partner for  
said *Treponema pallidum* antibodies.
12. A method for detecting antibodies specific for *Treponema pallidum* in an isolated sample  
20 said method comprising
  - a) forming an immunoreaction admixture by admixing a body fluid sample with TpN47  
antigen according to any of claims 1 to 5 or with a composition of at least two  
*Treponema pallidum* antigens according to claim 10
  - b) maintaining said immunoreaction admixture for a time period sufficient for allowing  
25 antibodies against said *Treponema* antigen or composition of *Treponema* antigens present  
in the body fluid sample to immunoreact with said *Treponema* antigen or composition of  
*Treponema* antigens to form an immunoreaction product; and
  - c) detecting the presence and /or the concentration of any of said immunoreaction product.
13. Use of a TpN47 antigen according to any of claims 1 to 5 or of a composition of  
30 *Treponema pallidum* antigens according to claim 10 in an *in vitro* diagnostic test for the  
detection of anti-*Treponema pallidum* antibodies.

14. A reagent kit for the detection of anti-*Treponema pallidum* antibodies, comprising at least a TpN47 antigen according to any one of claims 1 to 5 or an antigen composition according to claim 10.