

4-Phenyl-1-piperazinyl, -piperidinyl and -tetrahydropyridyl derivatives

Field of the Invention

The present invention relates to a novel class of halogen substituted 4-phenyl-1-piperazinyl,
 5 -piperidinyl and -tetrahydropyridyl derivatives having affinity for dopamine D₄ receptors
 and D₃ receptors. The compounds of the invention are considered useful in the treatment of
 certain psychiatric and neurologic disorders, including psychosis.

Background of the Invention.


10

US patent No. 3.188.313 relates to certain 1-(1-, 2-, and 3-indolylalkyl)piperazines, which
 are said to have CNS depressant and tranquillising effect.

15

Other compounds related to the compounds of the invention, which are said to interact with
 the dopamine and/or the serotonin system, are known in the art.

Thus, EP-B1-496 222, claims compounds having the formula




20

wherein Ar is a phenyl group, which may be substituted with halogen, alkyl, cyano, hydroxy
 etc. and Ind is 3-indolyl, which may be substituted with cyano, aminocarbonyl and
 aminocarbonylamino. The compounds disclosed in EP-B1-496 222 are said to be serotonin
 antagonists and agonists. It is also mentioned that the compounds have effect on dopamine
 accumulation in striatum and 5-HTP accumulation in N. Raphe. The compounds are said to
 25 be useful as anxiolytica, antidepressiva, neuroleptica and antihypertonica.

30

WO 99/09025 claims certain 2-(4-aryl-piperazin-1-yl)methyl-1*H*-indole derivatives. The
 compounds are said to be dopamine D₄ receptor agonists. Further, WO 94/24105 relates to
 2-(2-(4-aryl-piperazin-1-yl)ethyl)-1*H*-indole derivatives, which are said to have selective
 affinity for the dopamine D₄ receptor subtype.

EP-B1-354 094 relates to certain oxindoles having the formula



wherein R^1 is hydrogen, halogen or alkyl, R^2 is hydrogen or alkyl, R^3 is hydrogen, alkyl or -S-alkyl and Ar may be chlorophenyl and other substituted aryl groups. The compounds bind
5 to the 5-HT_{1A} receptor and are said to be agonists, partial agonists or antagonists at this receptor. Certain of the compounds are said to possess activity at 5-HT₂ receptors.

WO 98/08816 also describes oxindoles, which are said to be psychotropic drugs, and the application contains data showing the activity of certain of the compounds at the D₄ receptor.

Pharmazie, 1997, 52, 423-428 describes N-[3-(4-aryl-1-piperazinyl)alkyl] derivatives of
10 indolin-2(1H)-one, quinolin-2(1H)-one and isoquinolin-1(2H)-one and their receptor affinities at the 5-HT_{1A} and the 5-HT_{2A} receptor. The compound 1-(3-(4-phenyl-1-piperazinyl)propyl)indolin-2(1H)-one is described as a 5-HT_{2A} antagonist with weak 5-HT_{1A} agonistic properties. The compound is suggested as a potential antidepressant and/or anxiolytic agent.

15

Subramanian et al., *Heterocyclic Communications* 1999, 5, 63-68 describes certain piperazinyl indolyl propanones claimed to show antagonism at dopamine D₁/D₂ receptors.

Further, Böttcher et al., *J. Med. Chem.* 1992, 35, 4020-4026, describes certain 3-(1,2,3,6-
20 tetrahydro-1-pyridylalkyl)indoles having dopaminergic activity.

Finally, *Pol. J. Pharmacol. Pharm.* 1984, 36, 697-703 describes the compound 1-(3-(4-(3-chlorophenyl)-1-piperazinyl)propyl)indane as having serotinolytic properties.

25 Dopamine D₄ receptors belong to the dopamine D₂ subfamily of receptors, which is considered to be responsible for the antipsychotic effects of neuroleptics. The side effects of

neuroleptic drugs, which primarily exert their effect via antagonism of D₂ receptors, are known to be due to D₂ receptor antagonism in the striatal regions of the brain. However, dopamine D₄ receptors are primarily located in areas of the brain other than striatum, suggesting that antagonists of the dopamine D₄ receptor will be devoid of extrapyramidal side effects. This is illustrated by the antipsychotic clozapine, which exerts higher affinity for D₄ than D₂ receptors, and is lacking extrapyramidal side effects (Van Tol et al. *Nature* 1991, 350, 610; Hadley *Medicinal Research Reviews* 1996, 16, 507-526 and Sanner *Exp. Opin. Ther. Patents* 1998, 8, 383-393).

- 10 A number of D₄ ligands, which were postulated to be selective D₄ receptor antagonists (L-745,879 and U-101958) have been shown to possess antipsychotic potential (Mansbach et al. *Psychopharmacology* 1998, 135, 194-200). However, recently it has been reported that these compounds are partial D₄ receptor agonists in various *in vitro* efficacy assays (Gazi et al. *Br. J. Pharmacol.* 1998, 124, 889-896 and Gazi et al. *Br. J. Pharmacol.* 1999, 128, 613-620).
- 15 Furthermore, it was shown that clozapine, which is an effective antipsychotic, is a silent antagonist (Gazi et al. *Br. J. Pharmacol.* 1999, 128, 613-620).

Consequently, D₄ ligands, which are partial D₄ receptor agonists or antagonists, may have beneficial effects against psychoses.

20

Dopamine D₄ antagonists may also be useful for the treatment of cognitive deficits (Jentsch et al. *Psychopharmacology* 1999, 142, 78-84).

- It has also been suggested that dopamine D₄ antagonists may be useful to reduce dyskinesia occurring as a result of the treatment of Parkinson's disease with L-dopa (Tahar et al. *Eur. J. Pharmacol.* 2000, 399, 183-186).

- Dopamine D₃ receptors also belong to the dopamine D₂ subfamily of receptors, and they are preferentially located in limbic regions of the brain (Sokoloff et al. *Nature* 1990, 347, 146-151), such as the nucleus accumbens, where dopamine receptor blockade has been associated with antipsychotic activity (Willner *Int. Clinical Psychopharmacology* 1997, 12, 297-308). Furthermore, an elevation of the level of D₃ receptors in the limbic part of schizophrenic brains has been reported (Gurevich et al. *Arch. Gen. Psychiatry* 1997, 54, 225-32). Therefore, D₃ receptor antagonists may offer the potential for an effective antipsychotic

therapy, free of the extrapyramidal side effects of the classical antipsychotic drugs, which primarily exert their effect by blockade of D₂ receptors (Shafer et al. *Psychopharmacology* 1998, 135, 1-16; Schwartz et al. *Brain Research Reviews* 2000, 31, 277-287).

5 Moreover, D₃ receptor blockade results in a slight stimulation in the prefrontal cortex (Merchant et al. *Cerebral Cortex* 1996, 6, 561-570), which could be beneficial against negative symptoms and cognitive deficits associated with schizophrenia. In addition, dopamine D₃ antagonists can reverse D₂ antagonist-induced EPS (Millan et al. *Eur. J. Pharmacol.* 1997, 321, R7-R9) and do not cause changes in prolactin (Reavill et al. *J. Pharmacol. Exp. Ther.* 2000, 294, 1154-1165).
10 Consequently, D₃ antagonistic properties of an antipsychotic drug could reduce the negative symptoms and cognitive deficits and result in an improved side effect profile with respect to EPS and hormonal changes.


Dopamine D₃ agonists have also been considered relevant in the treatment of schizophrenia
15 (Wustow et al. *Current Pharmaceutical Design* 1997, 3, 391-404).

According to the present invention, a novel class of dopamine D₄ receptor ligands is provided. Most of the compounds of the invention also have high affinity for the dopamine D₃ receptor and as mentioned above, dopamine D₃ antagonistic properties of an
20 antipsychotic drug may reduce the negative symptoms and cognitive deficits of schizophrenia and result in an improved side effect profile.

Moreover, certain of the compounds of the invention have the further advantage of having only very weak effect at adrenergic alpha-1-receptors which imply a low propensity to cause
25 orthostatic hypotension.

Summary of the Invention

Accordingly, the present invention relates to the novel compounds of formula I



(I)

wherein W is C, CH or N and the dotted line emanating from W indicate a bond when W is C and no bond when W is CH or N;

5

R^1 and R^2 are independently selected from hydrogen and halogen provided that at least one of R^1 and R^2 is a halogen atom;


R^3 is selected from hydrogen, halogen, C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl,

10 trifluoromethyl, C_{1-6} -alkoxy, aryloxy, aralkoxy, hydroxy, amino, C_{1-6} -alkylamino, di(C_{1-6} -alkyl)amino, nitro and cyano

n is 2, 3, 4, or 5;

15 X is CH_2 , O, S, CO, CS, SO or SO_2 ; and

Q is a group of formula



(II)

wherein Z is a chain of 3 to 4 chain members, wherein the chain members are selected from C, CH, CH_2 , CO, N and NH, provided that only one of the chain members may be N or NH, said chain Z optionally containing one or two double bonds

20

R^4 , R^5 , R^6 , R^7 , R^8 and R^9 are independently selected from hydrogen, halogen, trifluoromethyl, nitro, cyano, C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{1-6} -alkoxy, C_{1-6} -alkylthio, C_{1-6} -alkylsulfonyl, hydroxy, hydroxy- C_{1-6} alkyl, amino, C_{1-6} -alkylamino, di(C_{1-6} -alkyl)amino, acyl, aminocarbonyl, C_{1-6} -alkylaminocarbonyl and

di(C₁₋₆-alkyl)aminocarbonyl;

provided that X is not O or S when the group Q is attached via an N atom;

- 5 and any of its enantiomers and acid addition salts thereof.

According to a preferred embodiment of the invention, R¹ and R² are both halogen, in particular chloro.

- 10 In another embodiment of the invention, one of R¹ and R² is halogen and the other is hydrogen. In particular, the invention relates to such compounds wherein R² is halogen and R¹ is hydrogen.

- If R² is hydrogen and R¹ is chloro, Q is preferably selected from indolinyl, 1,2,3,4-
15 tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl or 1-indanyl.

In one particular embodiment of the invention, when R² is hydrogen and R¹ is halogen, R¹ is not chloro.

- 20 In a further embodiment of the invention, R¹ and R² are independently selected from hydrogen and chloro.

In another particular embodiment of the invention W is N.

- 25 In a preferred embodiment of the invention X is CO or CH₂.

In a further embodiment of the invention, Z is a chain of 3 to 4 chain members, wherein the chain members are selected from C, CH, CH₂, N and NH, provided that only one of the chain members may be N or NH.

30

Q may preferably be selected from optionally substituted 1-indolinyl, 3-indolyl, 1-indanyl, 2-oxo-1, 2, 3, 4-tetrahydroquinolinyl and 1,2,3,4-tetrahydroisoquinolinyl.

In particular, Q is unsubstituted or substituted with halogen.

R³ is preferably hydrogen or halogen and if R³ is halogen it is preferably attached in the para position on the phenyl ring.

- 5 The compounds of the invention have been found to show high affinity for dopamine D₄ receptors and dopamine D₃ receptors, in certain cases combined with a very low effect at adrenergic alpha-1- receptors.

The compounds of the invention are therefore considered useful for the treatment of
10 psychosis, including the positive and negative symptoms of schizophrenia.

Moreover, certain of the compounds have the further advantage of having only very weak effect at adrenergic alpha-1-receptors which imply a low propensity to cause orthostatic hypotension.

15

Some of the compounds interact with central serotonergic receptors, e.g. the 5-HT_{1A} or 5-HT_{2A} receptors and/or they act as 5-HT reuptake inhibitors.

These compounds of the invention may therefore also be useful for the treatment of
20 disorders caused by imbalances in the serotonergic system including affective disorders such as generalised anxiety disorder, panic disorder, and obsessive compulsive disorder, depression and aggression.

In particular, compounds with combined effects at dopamine D₄ and 5-HT receptors and/or
25 the 5-HT transporter may have the benefit of improved effect on other psychiatric symptoms associated with schizophrenia, such as depressive and anxiety symptoms.

Thus, in another aspect, the present invention provides a pharmaceutical composition comprising at least one compound of Formula I as defined above or a pharmaceutically
30 acceptable acid addition salt thereof in combination with one or more pharmaceutically acceptable carriers or diluents.

The invention also relates to the use of a compound of the invention for the manufacture of a medicament useful in the treatment of psychosis including the positive and negative symptoms

of schizophrenia, affective disorders such as generalised anxiety disorder, panic disorder, and obsessive compulsive disorder, depression, aggression, cognitive disorders and dyskinesia induced by treatment with L-dopa.

5 Detailed Description of the Invention

Some of the compounds of general Formula I exist as optical isomers thereof and such optical isomers are also embraced by the invention.

- 10 The term C₁₋₆-alkyl refers to a branched or unbranched alkyl group having from one to six carbon atoms inclusive, such as methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, 2-methyl-2-propyl and 2-methyl-1-propyl.









- The terms C₁₋₆-alkoxy, C₁₋₆-alkylthio, C₁₋₆-alkylamino, di(C₁₋₆-alkyl)amino etc. designate
15 such groups in which the alkyl group is C₁₋₆-alkyl as defined above.

The term aryl refers to a carbocyclic aromatic group, such as phenyl, naphthyl, in particular phenyl, including methyl substituted naphthyl, or phenyl.

- 20 The term aralkyl means aryl-C₁₋₆-alkyl, wherein aryl and C₁₋₆-alkyl is as defined above.

The terms aralkoxy and aryloxy means aryl-C₁₋₆-alkyl-O- and aryl-O- where aryl and C₁₋₆-alkyl are as defined above.

- 25 Halogen means fluoro, chloro, bromo or iodo. A group Q wherein Z is as defined above includes groups such as:



wherein $R^4 - R^9$ are as defined above and the dotted line indicates an optional bond.

Preferred compounds of the invention are the compounds selected from

5

3-[4-(2,3-Dichlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1*H*-indol-1-yl)propan-1-one,
 4-[4-(2,3-Dichlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1*H*-indol-1-yl)butan-1-one,
 5-[4-(2,3-Dichlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1*H*-indol-1-yl)pentan-1-one,
 4-[4-(2-Chlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1*H*-indol-1-yl)butan-1-one,

- 4-[4-(3-Chlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1*H*-indol-1-yl)butan-1-one,
 1-{3-[4-(2,3-Dichlorophenyl)piperazin-1-yl]propyl}-2,3-dihydro-1*H*-indole,
 1-{4-[4-(2,3-Dichlorophenyl)piperazin-1-yl]butyl}-2,3-dihydro-1*H*-indole,
 1-{5-[4-(2,3-Dichlorophenyl)piperazin-1-yl]pentyl}-2,3-dihydro-1*H*-indole,
 5 1-{4-[4-(2-Chlorophenyl)piperazin-1-yl]butyl}-2,3-dihydro-1*H*-indole,
 1-{4-[4-(3-Chlorophenyl)piperazin-1-yl]butyl}-2,3-dihydro-1*H*-indole,
 1-(2,3-Dichlorophenyl)-4-[4-(indan-1-yl)butyl]piperazine,
 6-Chloro-3-{2-[4-(2,3-dichlorophenyl)piperazin-1-yl]ethylsulfanyl}-1*H*-indole,
 3-{3-[4-(2,3-Dichlorophenyl)piperazin-1-yl]propyl}-1*H*-indole,
 10 3-{4-[4-(2,3-Dichlorophenyl)piperazin-1-yl]butyl}-1*H*-indole,
 3-{3-[4-(2,3-Dichlorophenyl)piperazin-1-yl]propyl}-5-fluoro-1*H*-indole,
 3-{4-[4-(2,3-Dichlorophenyl)piperazin-1-yl]butyl}-5-fluoro-1*H*-indole,
 6-Chloro-3-{3-[4-(2,3-dichlorophenyl)piperazin-1-yl]propyl}-1*H*-indole,
 1-{4-[4-(2,3-Dichlorophenyl)piperazin-1-yl]butyl}-3,4-dihydroquinolin-2(1*H*)-one,
 15 3-[4-(2,3-Dichlorophenyl)piperazin-1-yl]-1-(5-fluoro-2,3-dihydro-1*H*-indol-1-yl)propan-1-one,
 4-[4-(2,3-Dichlorophenyl)piperazin-1-yl]-1-(5-fluoro-2,3-dihydro-1*H*-indol-1-yl)butan-1-one,
 5-[4-(2,3-Dichlorophenyl)piperazin-1-yl]-1-(5-fluoro-2,3-dihydro-1*H*-indol-1-yl)pentan-1-
 20 one,
 3-[4-(2,3-Dichlorophenyl)piperazin-1-yl]-1-(3,4-dihydro-1*H*-isoquinolin-2-yl)-propan-1-one,
 4-[4-(2,3-Dichlorophenyl)piperazin-1-yl]-1-(3,4-dihydro-1*H*-isoquinolin-2-yl)-butan-1-one,
 5-[4-(2,3-Dichloro-phenyl)-piperazin-1-yl]-1-(3,4-dihydro-1*H*-isoquinolin-2-yl)-pentan-1-
 25 one,
 3-[4-(2-Chlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1*H*-indol-1-yl)propan-1-one,
 5-[4-(2-Chlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1*H*-indol-1-yl)pentan-1-one,
 3-[4-(3-Chlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1*H*-indol-1-yl)propan-1-one,
 5-[4-(3-Chlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1*H*-indol-1-yl)pentan-1-one,
 30 3-[4-(3-Chlorophenyl)piperazin-1-yl]-1-(5-fluoro-2,3-dihydro-1*H*-indol-1-yl)propan-1-one,
 5-[4-(3-Chlorophenyl)piperazin-1-yl]-1-(5-fluoro-2,3-dihydro-1*H*-indol-1-yl)pentan-1-one,
 3-[4-(2-Chlorophenyl)piperazin-1-yl]-1-(5-fluoro-2,3-dihydro-1*H*-indol-1-yl)propan-1-one,
 4-[4-(2-Chlorophenyl)piperazin-1-yl]-1-(5-fluoro-2,3-dihydro-1*H*-indol-1-yl)butan-1-one,
 3-[4-(3-Chlorophenyl)piperazin-1-yl]-1-(3,4-dihydro-1*H*-isoquinolin-2-yl)-propan-1-one,


- 5-[4-(3-Chlorophenyl)piperazin-1-yl]-1-(3,4-dihydro-1*H*-isoquinolin-2-yl)-pentan-1-one,
 3-[4-(2-Chlorophenyl)piperazin-1-yl]-1-(3,4-dihydro-1*H*-isoquinolin-2-yl)-propan-1-one,
 4-[4-(2-Chlorophenyl)piperazin-1-yl]-1-(3,4-dihydro-1*H*-isoquinolin-2-yl)-butan-1-one,
 1-(2,3-Dihydro-1*H*-indol-1-yl)-3-[4-(2-fluoro-phenyl)-piperazin-1-yl]-propan-1-one,
 5 1-(2,3-Dihydro-1*H*-indol-1-yl)-4-[4-(2-fluoro-phenyl)-piperazin-1-yl]-butan-1-one,
 1-(2,3-Dihydro-1*H*-indol-1-yl)-5-[4-(2-fluoro-phenyl)-piperazin-1-yl]-pentan-1-one,
 1-(5-Fluoro-2,3-dihydro-1*H*-indol-1-yl)-3-[4-(2-fluoro-phenyl)-piperazin-1-yl]-propan-1-one,
 1-(5-Fluoro-2,3-dihydro-1*H*-indol-1-yl)-4-[4-(2-fluoro-phenyl)-piperazin-1-yl]-butan-1-one,
 10 1-(5-Fluoro-2,3-dihydro-1*H*-indol-1-yl)-5-[4-(2-fluoro-phenyl)-piperazin-1-yl]-pentan-1-one,
 3-[4-(2,4-Difluoro-phenyl)-piperazin-1-yl]-1-(2,3-dihydro-1*H*-indol-1-yl)propan-1-one,
 4-[4-(2,4-Difluoro-phenyl)-piperazin-1-yl]-1-(2,3-dihydro-1*H*-indol-1-yl)butan-1-one, and
 5-[4-(2,4-Difluoro-phenyl)-piperazin-1-yl]-1-(2,3-dihydro-1*H*-indol-1-yl)pentan-1-one
 15 and pharmaceutically acceptable acid addition salts thereof.

The acid addition salts of the compounds of the invention may be pharmaceutically acceptable salts formed with non-toxic acids. Exemplary of such organic salts are those with maleic, fumaric, benzoic, ascorbic, succinic, oxalic, bis-methylenesalicylic, methanesulfonic, ethanedisulfonic, acetic, propionic, tartaric, salicylic, citric, gluconic, lactic, malic, mandelic,
 20 cinnamic, citraconic, aspartic, stearic, palmitic, itaconic, glycolic, p-aminobenzoic, glutamic, benzenesulfonic and theophylline acetic acids, as well as the 8-halothephyllines, for example 8-bromothephylline. Exemplary of such inorganic salts are those with hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric and nitric acids.

25

The compounds of the invention may be prepared as follows:


- a) Reducing the carbonyl group of a compound of formula III



30

wherein R^1 , R^2 , R^3 , W, n, Q and the dotted line are as previously defined;


- 5 b) alkylating an amine of formula IV with a reagent of formula V



wherein R^1 , R^2 , R^3 , X, W, n, Q and the dotted line are as previously defined, and G is a suitable leaving group such as halogen, mesylate or tosylate;


10

- c) reductive alkylation of an amine of the formula IV with a reagent of formula VII




wherein R^1 , R^2 , R^3 , X, n, W, Q and the dotted line are as previously defined, and E is either
15 an aldehyde or an activated carboxylic acid group;

- d) reducing the amide group of a compound of formula VIII




wherein $R^1, R^2, R^3, X, n, W, Q$ and the dotted line are as previously defined;

e) acylation or reductive alkylation of an amine of the formula II'



- 5 wherein Z^1 is a chain of 3 to 4 chain members, wherein the chain members are selected from C, CH, CH_2 , CO and NH, provided that one of the chain members is NH, and said chain optionally containing one or two double bonds, with a reagent of formula IX




10

wherein R^1, R^2, R^3, n, W and the dotted line are as previously defined, and E is either an aldehyde or an activated carboxylic acid group


- f) Lewis-acid catalyzed cleavage of a resin-bound ester of formula X by an amine of the formula Q'

15



wherein R^1 , R^2 , R^3 , Q' , W , n and the dotted line are as previously defined, or

g) reduction of the double bond in a compound of formula XI



wherein R^1 , R^2 , R^3 , n , X and Q are as previously defined, whereupon the compound of formula I is isolated as the free base or an acid addition salt thereof.

The reduction according to method a) is preferably carried out in an inert organic solvent such as diethyl ether or tetrahydrofuran in the presence of alane or lithium aluminium hydride from 0 °C to reflux temperature. Method b) may be used to prepare starting materials of formula III.

The alkylation according to method b) is conveniently performed in an inert organic solvent such as a suitably boiling alcohol or ketone, preferably in the presence of an organic or inorganic base (potassium carbonate, diisopropylethylamine or triethylamine) at reflux temperature. Alternatively, the alkylation can be performed at a fixed temperature, which is different from the boiling point, in one of the above-mentioned solvents or in dimethyl formamide (DMF), dimethylsulfoxide (DMSO) or *N*-methylpyrrolidin-2-one (NMP), preferably in the presence of a base.

Amines of formula IV are either commercially available or known from the literature (e.g. Oshiro et al. *J. Med. Chem.* 1991, 34, 2014-2023, Oshiro et al. *J. Med. Chem.* 1998, 41, 658-667, and Oshiro et al. *J. Med. Chem.* 2000, 43, 177-189). Alkylating reagents of formula V are known from the literature, or they can be prepared by methods obvious to a chemist skilled in the art by an analogous synthetic sequence. Thus, key intermediates such as halo-(2,3-dihydro-1*H*-indol-1-yl)alkan-1-one are prepared by addition of a haloalkanoyl chloride to a 2,3-dihydro-1*H*-indole in the presence of base. Similarly, halo-(3,4-dihydro-1*H*-isoquinolin-2-yl)alkan-1-one may be prepared from a haloalkanoyl chloride and a 3,4-

dihydro-1*H*-isoquinoline. The 2,3-dihydro-1*H*-indoles are either commercially available or prepared from the corresponding 1*H*-indoles by reduction of the 1*H*-indole with e.g. sodium cyanoborohydride in acetic acid or in trifluoroacetic acid whereas the 3,4-dihydro-1*H*-isoquinolines are commercially available or described in the literature. Alkylating haloalkyl-
 5 1*H*-indoles were prepared by literature methods (Benghiat et al. *J. Med. Chem.* 1983, 26, 1470-1477) or analogues to methods described in the literature (Brodfehrer et al. *J. Org. Chem.* 1997, 62, 9192 and WO 00/35872). Alkylating 3-haloalkylindanes or other types of alkylating alkylindanes can be prepared from known indanylalkancarboxylic acids (Mukhopadhyay et al. *J. Indian Chem. Soc.* 1985, 62, 690-692 and Tanaka et al. *J. Med.*
 10 *Chem.* 1994, 37, 2071-2078) by well known procedures. Alkylating 1-(haloalkyl)-3,4-dihydroquinolin-2(1*H*)-one can be prepared as described in EP-B1-512525.

The reductive alkylation according to methods c) and e) can be performed in two steps, e.g. coupling of amines of formula VI/II' with reagent of formula VII/IX by standard methods
 15 via the carboxylic acid chloride, activated esters or by use of carboxylic acids in combination with coupling reagents such as e.g. dicyclohexyl carbodiimide followed by reduction of the resulting amide with lithium aluminium hydride or alane. The reaction can also be performed by a standard one-pot procedure, e.g. by the use of reductive amination of amines of formula VI/II' with aldehydes of formula VII/IX. Carboxylic acids or aldehydes of
 20 formula VII are either commercially available or described in the literature. Thus, key intermediates have been described in the literature such as indanylalkancarboxylic acids (Mukhopadhyay et al. *J. Indian Chem. Soc.* 1985, 62, 690-692 and Tanaka et al. *J. Med. Chem.* 1994, 37, 2071-2078), substituted 3-(1*H*-indol-3-yl)propionic acids (Carbommelle et al. *Tetrahedron* 1998, 39, 4471-4472) and (2,3-dihydro-1*H*-indol)alkancarboxylic acids
 25 (WO 98/28293 and Ly et al. *Tetrahedron Letts.* 1999, 40, 2533-2536). Other substituted (1*H*-indol-3-yl)alkancarboxylic acids can be prepared by chain elongation of readily accessible 3-indoleglyoxylyl chlorides (Speeter et al. *J. Am. Chem. Soc.* 1954, 76, 6208-6210 and Nichols et al. *Synthesis* 1999, 6, 935-938). The 3-indoleglyoxylyl chlorides may be prepared from commercially available 1*H*-indoles. Various substituted (1*H*-indol-3-
 30 ylsulfanyl)alkancarboxylic acids can be prepared in an analogous manner as described for aminoalkylsulfanyl-1*H*-indoles (Zelesko et al. *J. Med. Chem.* 1983, 26, 230-237 or WO 91/04973) by *in situ* alkylation of substituted sodium 3-indolylthiolate with alkyl haloalkanoates and subsequent hydrolysis of the ester group.

Reduction of amide groups according to method d) is most conveniently performed with lithium aluminium hydride or alane in an inert organic solvent such as e.g. tetrahydrofuran or diethylether from 0 °C to reflux temperature.

- 5 Acylation according to method e) can be performed by standard literature methods, e.g. coupling of amines of formula Q' with reagent of formula IX by standard methods via the carboxylic acid chloride, activated esters or by use of carboxylic acids in combination with coupling reagents such as e.g. dicyclohexyl carbodiimide.
- 10 The Lewis acid catalyzed conversion of an ester to an amide according to method f) can be performed by standard literature methods (Barn et al. *Tet. Lett.* 1996, 37, 3213-3216). The Resin bound ester X can likewise be synthesised according to the literature (see e.g. Barn et al. *Tet. Lett.* 1996, 37, 3213-3216).
- 15 The reduction of the double bond according to method g) is generally performed by catalytic hydrogenation at low pressure (< 3 atm.) in a Parr apparatus, or by using reducing agents such as diborane or hydroboric derivatives as produced *in situ* from NaBH₄ in trifluoroacetic acid in inert solvents such as tetrahydrofuran (THF), dioxane, or diethyl ether.

20 Experimental Section

- Melting points were determined on a Büchi B-535 apparatus and are uncorrected. Mass spectra were obtained on a Quattro MS-MS system from VG Biotech, Fisons Instruments or on a Sciex API 150EX from Perkin Elmer. Spectra were obtained at two sets of operating conditions by the use of either electrospray ionisation or ACPI: one set to obtain molecular weight information and the other set to induce fragmentation patterns. ¹H NMR spectra were recorded at 250.13 MHz on a Bruker AC 250 or at 500.13 MHz on a Bruker DRX 500. Deuterated chloroform (99.8% D) or dimethylsulfoxide (99.9% D) were used as solvents. TMS was used as internal reference standard. Chemical shifts are expressed as ppm values. The following abbreviations are used for multiplicity of NMR signals: s=singlet, d=doublet, t=triplet, q=quartet, qv=quintet, h=heptet, dd=double doublet, dt=double triplet, dq=double quartet, tt=triplet of triplets, m= multiplet, b=broad. NMR signals corresponding to acidic protons are to some extent omitted. Content of water in crystalline compounds were

determined by Karl Fischer titration. For column chromatography silica gel of type Kieselgel 60, 40-60 mesh ASTM was used.

Preparation of intermediates

5

A. Alkylating reagents

3-Chloro-1-(2,3-dihydro-1*H*-indol-1-yl)propan-1-one

A mixture of 2,3-dihydro-1*H*-indole (50 g), triethylamine (132 g) and tetrahydrofuran (1000
10 mL) was cooled down to 10 °C followed by the addition (over a period of 60 min) of a solution of 3-chloropropanoyl chloride (55 g) in tetrahydrofuran (400 mL). The mixture was filtered, and the remaining solution was evaporated *in vacuo* to dryness. The residue was purified by flash chromatography (eluent: ethyl acetate/heptane 1:3) giving the title compound as a crystalline white material (31 g).

15

The following compounds were prepared in a similar manner

4-Chloro-1-(2,3-dihydro-1*H*-indol-1-yl)butan-1-one

from 2,3-dihydro-1*H*-indole and 4-chlorobutanoyl chloride

20

5-Bromo-1-(2,3-dihydro-1*H*-indol-1-yl)pentan-1-one

from 2,3-dihydro-1*H*-indole and 5-bromopentanoyl chloride

The following two compounds were prepared as described in Benghiat et al. *J. Med. Chem.*

25 1983, 26, 1470-1477

3-(3-Bromopropyl)-1*H*-indole

3-(4-Bromobutyl)-1*H*-indole

30

3-(3-Chloropropyl)-5-fluoro-1*H*-indole

The compound 5-chloropentan-1-ol (16.2 mL) was dissolved in cold 5 mM 2,2,6,6-tetramethylpiperidine-1-yloxy (tempo) in dichloromethane (240 mL) and cooled down to 0°C with an ice bath. Potassium bromide (0.5 M in water, 24 mL) was added, and this was
35 followed by the addition (in one portion at 5 °C under vigorous stirring) of a solution of

sodium hydrogencarbonate (24 g) in aqueous sodium hypochlorite (0.3 M, 500 mL). The resulting mixture was stirred at 5 °C for 20 min, and the phases were separated. The water phase was extracted with dichloromethane (200 mL), and the combined organic phases were evaporated *in vacuo* giving 5-chloropentanal as a clear oil (16 g). Subsequently, 5-chloropentanal was suspended in water (100 mL) followed by addition of 4-fluorophenylhydrazine hydrochloride (19.5 g) and toluene (800 mL), and the mixture was stirred at room temperature for 15 min. Phosphoric acid (85%, 100 mL) was added and the mixture was boiled under reflux for 2 h. The mixture was cooled to room temperature and the phases were separated. The organic phase was washed with saturated aqueous sodium hydrogencarbonate, dried (MgSO₄) and evaporated *in vacuo* to yield an orange oil. The crude product was purified by flash chromatography on silicagel (eluent: ethylacetate/heptane 1:4) to give the title compound as an orange oil (14 g).

The following compound was prepared in a similar manner

15

3-(4-Chlorobutyl)-5-fluoro-1H-indole

from 6-chlorohexan-1-ol and 4-fluorophenylhydrazine hydrochloride

6-Chloro 3-(3-iodopropyl)-1H-indole

20 The alcohol 2-(6-chloro-1H-indol-3-yl)ethanol (25 g, prepared analogues to compounds described in Demerson et al. *J. Med. Chem.* 1976, 19, 391-395 from 6-chloro-1H-indole and oxalyl chloride) was dissolved in tetrahydrofuran (300 mL) followed by the addition of triethylamine (17.7 mL). The resulting mixture was cooled to 5-6 °C followed by the addition of a solution of methanesulfonic acid chloride (14.6 g) in tetrahydrofuran (100 mL). The mixture was stirred at room temperature for 2 h, filtered and evaporated to dryness *in vacuo*. The residue was dissolved in acetone followed by addition of sodium iodide (96.2 g), and the resulting mixture was boiled under reflux for 4 h. The mixture was poured onto brine and the aqueous phase was extracted with ethyl acetate. The combined organic phases were dried (MgSO₄), filtered and concentrated *in vacuo* (38.2 g). The residue (30 g) was dissolved in dimethylsulfoxide (DMSO, 200 mL) and added drop-wise to a suspension of NaCN (15 g) and DMSO (250 mL) at 80 °C. The resulting mixture was stirred at 100 °C for 1 h, cooled down to room temperature and poured onto brine. The aqueous phase was extracted with diethyl ether, and the combined organic phase was dried (MgSO₄), filtered and concentrated *in vacuo* to yield crude intermediate (22.5 g). The residue was dissolved in methanol (750

mL) and added a mixture of HCl/methanol resulting in a combined solution of about 1 M HCl in methanol. The mixture was stirred at room temperature for 24 h followed by heating at 40 °C for an additional 3 h. The solvent was removed *in vacuo*, and the residue was dissolved in a mixture of diethyl ether and water. The resulting mixture was stirred at room temperature for 30 min. and the phases were separated. The aqueous phase was extracted an additional two times with diethyl ether and the combined organic phase was washed with brine, dried (MgSO₄), filtered and concentrated *in vacuo* (18.2 g). The residue was dissolved in tetrahydrofuran (300 mL) and added drop-wise to a suspension of LiAlH₄ (11.6 g) in tetrahydrofuran (1000 mL). The resulting mixture was boiled under reflux for 3 h, cooled down to 10 °C and worked up by the use of an equivalent amount of water. The organic phase was dried (MgSO₄), filtered and concentrated *in vacuo* (16.6 g). The residue (8 g) was dissolved in tetrahydrofuran (100 mL) and triethylamine (3.9 g) and cooled down to 10 °C followed by the addition of a solution of methanesulfonic acid chloride (4.4 g) in tetrahydrofuran (50 mL). The mixture was stirred at room temperature for 2 h and then evaporated to dryness *in vacuo*. The residue was dissolved in acetone followed by the addition of NaI (28.6 g), and the resulting mixture was boiled under reflux for 3 h. The mixture was poured onto brine, and the aqueous phase was extracted with tetrahydrofuran. The combined organic phase was dried (MgSO₄), filtered and concentrated *in vacuo* (17.4 g).

20

The following compound was prepared in a similar manner

4-(Indan-1-yl)butyl methanesulfonate

from 4-(indan-1-yl)butanoic acid prepared as described by Mukhopadhyay et al. *J. Indian Chem. Soc.* 1985, 62, 690-692.

25

1-(4-Bromobutyl)-3,4-dihydroquinolin-2(1H)-one

A suspension of sodium hydride (6.8 g, 60% dispersion in mineral oil) and dimethyl formamide (200 mL) was kept at 20-30 °C followed by the addition of a solution of 3,4-dihydroquinolin-2(1H)-one (25 g) in dimethyl formamide (100 mL). The resulting mixture was stirred at room temperature for 30 min followed by the addition of a solution of 1,4-dibromobutane (184 g) in dimethyl formamide (200 mL) at a temperature of 20-40 °C. The reaction mixture was stirred at room temperature for 30 min and evaporated *in vacuo*. The remaining oil was poured into ice water and extracted with ethyl acetate. The combined

30

organic phases were washed with water and brine, treated with charcoal, dried (MgSO₄) and evaporated *in vacuo*. The remaining oil was purified by flash chromatography (eluent: ethyl acetate/heptane 1:1) giving the title compound as a red oil (36 g).

5 B. Acylating reagents

(6-Chloro-1*H*-indol-3-ylsulfanyl)acetic acid

The compounds 6-chloro-1*H*-indole (15.1 g) and thiourea (7.6 g) were dissolved in methanol (150 mL) followed by the addition of a solution of iodine/potassium iodide (1 M, 100 mL) under stirring. The solution was stirred at room temperature for 2 h and then evaporated *in vacuo* to give an oil. Sodium hydroxide (1.5 M, 200 mL) was added, and the solution was heated at 90 °C for 90 min. This solution was cooled to room temperature and extracted with diethyl ether (discarded). The aqueous phase was added diethyl ether (100 mL) and ethyl chloroacetate (10 mL), and the resulting mixture was stirred at room temperature for 16 h.

The phases were separated, and the aqueous phase was extracted with diethyl ether. The combined organic phases were collected and dried (MgSO₄). The suspension was filtered, and the organic phase was evaporated to dryness to give a brown oil (18.1 g). The oil was dissolved in ethanol (50 mL) followed by the addition of a solution of water (50 mL) and potassium hydroxide (4.0 g). The resulting mixture was boiled under reflux for 2 h and cooled to room temperature. The pH of the mixture was adjusted to 3-4 by the addition of hydrochloride acid (1 M). Water was added (100 mL), and the aqueous phase was extracted with ethyl acetate. The combined organic phases were dried (MgSO₄), filtered and evaporated *in vacuo* to give the title compound as an oil (12.3 g).

25 Preparation of the compounds of the invention

Example 1

1a, 3-[4-(2,3-Dichlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1*H*-indol-1-yl)propan-1-one

A mixture of 1-(2,3-dichlorophenyl)piperazine, hydrochloride (8.0 g) and potassium carbonate (15 g) in a mixture of butanone (50 mL) and dimethyl formamide (5 mL) was heated to 50 °C followed by the addition of 3-chloro-1-(2,3-dihydro-1*H*-indol-1-yl)propan-1-one (6.0 g). The resulting mixture was boiled under reflux for 40 h and filtered hot. The remaining organic phase was left to crystallise, and a white crystalline material was collected by filtration and washed with acetone (8.5 g). Mp 157-158 °C. ¹H NMR (DMSO-*d*₆): 2.60 (s,

4H); 2.60-2.80 (m, 4H); 3.00 (s, 4H); 3.15 (t, 2H); 4.10 (t, 2H); 6.95 (t, 1H); 7.10-7.15 (m, 2H); 7.20 (d, 1H); 7.25-7.35 (m, 2H); 8.10 (d, 1H). MS m/z : 404 (MH⁺), 243.

1b, 4-[4-(2,3-Dichlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1H-indol-1-yl)butan-1-one

5 A mixture of 1-(2,3-dichlorophenyl)piperazine, hydrochloride (8.0 g) and diisopropylethylamine (10 mL) in dimethyl formamide (50 mL) was heated to 45 °C followed by the addition of 4-chloro-1-(2,3-dihydro-1H-indol-1-yl)butan-1-one (6.7 g). The resulting mixture was heated at 100 °C for 6 h, cooled to room temperature and poured into water. The aqueous phase was extracted with diethyl ether, and the combined organic phases
10 were dried (MgSO₄), filtered and evaporated *in vacuo* to give a black oil (14.2 g). The oil was crystallised from acetone, and the formed crystals were recrystallised from ethanol to give a white crystalline material (3.8 g). Mp 134-136 °C. ¹H NMR (CDCl₃): 1.90-2.05 (m, 2H); 2.45-2.60 (m, 4H); 2.65 (s, 4H); 3.00 (s, 4H); 3.20 (t, 2H); 4.10 (t, 2H); 6.90 (d, 1H); 7.00 (t, 1H); 7.05-7.25 (m, 4H); 8.25 (d, 1H). MS m/z : 418 (MH⁺), 299, 228, 188.

1c, 5-[4-(2,3-Dichlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1H-indol-1-yl)pentan-1-one

A mixture of 1-(2,3-dichlorophenyl)piperazine hydrochloride (8.0 g) and diisopropylethylamine (15 mL) in butanone (50 mL) was heated to 45 °C followed by the addition of 5-bromo-1-(2,3-dihydro-1H-indol-1-yl)pentan-1-one (5.4 g). The resulting
20 mixture was boiled under reflux for 40 h and filtered hot. The remaining organic phase was left to crystallise, and a white crystalline material was collected by filtration and washed with acetone (3.8 g). Mp 121-123 °C. ¹H NMR (DMSO-*d*₆): 1.50-1.70 (m, 4H); 2.30-2.65 (m, 8H); 3.00 (s, 4H); 3.15 (t, 2H); 4.10 (t, 2H); 6.95 (t, 1H); 7.10-7.15 (m, 2H); 7.20 (d, 1H); 7.25-7.35 (m, 2H); 8.10 (d, 1H). MS m/z : 432 (MH⁺), 315, 202.

25 The following compounds were prepared in a similar manner

1d, 4-[4-(2-Chlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1H-indol-1-yl)butan-1-one

from 1-(2-chlorophenyl)piperazine, hydrochloride and 4-chloro-1-(2,3-dihydro-1H-indol-1-yl)butan-1-one. Mp 119-121 °C. ¹H NMR (DMSO-*d*₆): 1.75-1.85 (m, 2H); 2.35-2.50 (m, 4H); 2.55 (s, 4H); 3.95 (s, 4H); 3.15 (t, 2H); 4.10 (t, 2H); 6.95 (t, 1H); 7.05 (t, 1H); 7.10 (d, 1H); 7.15 (t, 1H); 7.20 (d, 1H); 7.25 (t, 1H); 7.40 (d, 1H); 8.10 (d, 1H). MS m/z : 384 (MH⁺), 265, 188.

1e, 4-[4-(3-Chlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1H-indol-1-yl)butan-1-one

from 1-(3-chlorophenyl)piperazine, dihydrochloride and 4-chloro-1-(2,3-dihydro-1H-indol-1-yl)butan-1-one. Mp 102-107 °C. ¹H NMR (DMSO-d₆): 1.75-1.85 (m, 2H); 2.35 (t, 2H); 2.45-2.55 (m, 6H); 3.10-3.20 (m, 6H); 4.10 (t, 2H); 6.75 (d, 1H); 6.85 (d, 1H); 6.90 (s, 1H); 6.95 (t, 1H); 7.10 (t, 1H); 7.15-7.25 (m, 2H); 8.10 (d, 1H). MS m/z: 384 (MH⁺), 265, 188.

Example 22a, 1-{3-[4-(2,3-Dichlorophenyl)piperazin-1-yl]propyl}-2,3-dihydro-1H-indole,10 hydrochloride

Lithium aluminium hydride (1.8 g) was suspended in tetrahydrofuran (30 mL) at 0 °C, and the suspension was added a solution of aluminium trichloride (1.8 g) in tetrahydrofuran (30 mL) at 0-5 °C over 15 min. To this mixture, a solution of 1a, 3-[4-(2,3-dichlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1H-indol-1-yl)propan-1-one (5 g) in tetrahydrofuran (50 mL) was added at a temperature of 0-10 °C. The resulting mixture was stirred for 30 min at 5 °C and then for 2 h at room temperature. The reaction mixture was quenched with water and sodium hydroxide (28%) and filtered. The organic phase was evaporated to dryness *in vacuo*, and the title compound was precipitated as the hydrochloride salt and recrystallised from ethanol (3.8 g). Mp 214-226 °C. ¹H NMR (DMSO-d₆): 2.05-2.20 (m, 2H); 2.95 (t, 2H); 3.10-3.35 (m, 8H); 3.35-3.50 (m, 4H); 3.60 (d, 2H); 6.70 (b s, 2H); 7.05 (t, 1H); 7.10 (d, 1H); 7.20 (d, 1H); 7.30-7.40 (m, 2H); 11.45 (b s). MS m/z: 390 (MH⁺), 271, 132.

The following compounds were prepared in a similar manner

25 2b, 1-{4-[4-(2,3-Dichlorophenyl)piperazin-1-yl]butyl}-2,3-dihydro-1H-indole, oxalate

from 1b, 4-[4-(2,3-dichlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1H-indol-1-yl)butan-1-one. Mp 157-160 °C. ¹H NMR (DMSO-d₆): 1.55-1.65 (m, 2H); 1.65-1.75 (m, 2H); 2.90 (t, 2H); 2.95 (t, 2H); 3.05 (t, 2H); 3.05-3.25 (m, 8H); 3.30 (t, 2H); 6.50 (d, 1H); 6.55 (t, 1H); 6.95 (t, 1H); 7.00 (d, 1H); 7.20 (d, 1H); 7.30-7.40 (m, 2H). MS m/z: 404 (MH⁺), 285, 174, 132.

2c, 1-{5-[4-(2,3-Dichlorophenyl)piperazin-1-yl]pentyl}-2,3-dihydro-1H-indole,
hydrochloride

from **1c**, 5-[4-(2,3-dichlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1*H*-indol-1-yl)pentan-1-one. Mp 219-228 °C. ¹H NMR (DMSO-*d*₆): 1.35-1.45 (m, 2H); 1.60-1.70 (m, 2H); 1.80-1.90 (m, 2H); 2.95-3.05 (m, 2H); 3.10-3.30 (m, 8H); 3.40-3.65 (m, 6H); 6.85 (b s, 2H); 7.05-7.25 (m, 3H); 7.30-7.40 (m, 2H); 11.20 (b s). MS *m/z*: 418 (MH⁺), 299, 188.

5

2d, 1-[4-[4-(2-Chlorophenyl)piperazin-1-yl]butyl]-2,3-dihydro-1*H*-indole, oxalate

from **1d**, 4-[4-(2-chlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1*H*-indol-1-yl)butan-1-one. Mp 146-148 °C. ¹H NMR (DMSO-*d*₆): 1.55-1.60 (m, 2H); 1.65-1.75 (m, 2H); 2.90 (t, 2H); 3.00 (t, 2H); 3.05 (t, 2H); 3.15 (b s, 8H); 3.30 (t, 2H); 6.50 (d, 1H); 6.55 (t, 1H); 7.00 (t, 1H); 7.05 (d, 1H); 7.10 (t, 1H); 7.20 (d, 1H); 7.35 (t, 1H); 7.45 (d, 1H). MS *m/z*: 370 (MH⁺), 251, 174.

10

2e, 1-[4-[4-(3-Chlorophenyl)piperazin-1-yl]butyl]-2,3-dihydro-1*H*-indole, oxalate

from **1e**, 4-[4-(3-chlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1*H*-indol-1-yl)butan-1-one. Mp 172-176 °C. ¹H NMR (DMSO-*d*₆): 1.55-1.60 (m, 2H); 1.65-1.75 (m, 2H); 2.85 (t, 2H); 2.95 (t, 2H); 3.00-3.20 (m, 6H); 3.30 (t, 2H); 3.40 (b s, 4H); 6.50 (d, 1H); 6.55 (t, 1H); 6.85 (d, 1H); 6.90-7.05 (m, 4H); 7.25 (t, 1H). MS *m/z*: 370 (MH⁺), 251, 174.

15

Example 3

20

3, 1-(2,3-Dichlorophenyl)-4-[4-(indan-1-yl)butyl]piperazine, oxalate

A mixture of 1-(2,3-dichlorophenyl)piperazine, hydrochloride (3.5 g) and diisopropylethylamine in a mixture of methyl isobutyl ketone (50 mL) and dimethyl formamide (5 mL) was heated to 60 °C followed by the addition of 4-(indan-1-yl)butyl methanesulfonate (3.5 g) in methyl isobutyl ketone (10 mL). The resulting mixture was boiled under reflux for 5 h and reduced *in vacuo*. The product was purified by flash chromatography on silicagel (eluent ethylacetate) to give the crude product that subsequently was precipitated as the oxalate salt (0.7 g). Mp 171-176 °C. ¹H NMR (DMSO-*d*₆): 1.35-1.45 (m, 3H); 1.55-1.75 (m, 3H); 1.80-1.90 (m, 1H); 2.20-2.30 (m, 2H); 2.75-2.90 (m, 2H); 2.95 (t, 1H); 3.10 (t, 1H); 3.20 (b s, 8H); 7.10-7.15 (m, 2H); 7.15-7.25 (m, 3H); 7.30-7.40 (m, 2H). MS *m/z*: 403 (MH⁺).

25

30

Example 4**4, 6-Chloro-3-{2-[4-(2,3-dichlorophenyl)piperazin-1-yl]ethylsulfanyl}-1H-indole, oxalate**

A solution of (6-chloro-1H-indol-3-ylsulfanyl)acetic acid (1.75 g) in tetrahydrofuran (30 mL) was added carbonyldiimidazole (1.2 g), stirred at room temperature for 30 min and cooled to 5 °C. To this mixture was added 1-(2,3-dichlorophenyl)piperazine (1.8 g) dissolved in tetrahydrofuran (20 mL), and the resulting mixture was stirred at room temperature for 2 h. The mixture was poured into water, and the aqueous phase was extracted with ethyl acetate. The combined organic phase was dried (MgSO₄), filtered and evaporated *in vacuo* to give an oil (3.6 g). The oil was subjected to the same reaction conditions (reduction with alane) as described in example 2, and the product was purified by flash chromatography on silicagel (eluent: ethylacetate/heptane 5:1) to give an oil. The title compound was isolated as the oxalate salt (0.8 g). Mp 137-141 °C. ¹H NMR (DMSO-d₆): 2.75-2.95 (m, 8H); 2.95-3.15 (m, 4H); 7.10-7.20 (m, 2H); 7.25-7.35 (m, 2H); 7.50 (s, 1H); 7.60-7.70 (m, 2H). 11.60 (b s, 1H). MS m/z: 442 (MH⁺), 291, 182.

Example 5**5a, 3-{3-[4-(2,3-Dichlorophenyl)piperazin-1-yl]propyl}-1H-indole, hydrochloride**

A mixture of 3-(3-bromopropyl)-1H-indole (1.19 g), potassium carbonate (1.4 g) and 1-(2,3-dichlorophenyl)piperazine (1.27 g) in anhydrous acetonitrile (10 mL) was boiled under reflux for 5 h and cooled to room temperature. The mixture was added silicagel (7 g), and the solvent was evaporated *in vacuo*. The compound was purified by flash chromatography on silicagel (eluent: ethylacetate/heptane/triethylamine 49:49:2). Fractions containing the compound were combined and evaporated *in vacuo*. Recrystallisation from acetonitrile gave the title compound as a white crystalline material. The compound was precipitated as the hydrochloride salt (1 g). Mp 241-242°C. ¹H NMR (DMSO-d₆): 2.10-2.25 (m, 2H); 2.75 (t, 2H); 3.10-3.30 (m, 6H); 3.40 (t, 2H); 3.60 (d, 2H); 7.00 (t, 1H); 7.05 (t, 1H); 7.15 (d, 1H); 7.25 (s, 1H); 7.30-7.40 (m, 3H); 7.55 (d, 1H); 10.90 (b s, 1H); 11.40 (b s, 1H). Ms m/z: 388 (MH⁺).

The following compounds were prepared in a similar manner

5b, 3-{4-[4-(2,3-Dichlorophenyl)piperazin-1-yl]butyl}-1H-indole, hydrochloride

from 1-(2,3-dichlorophenyl)piperazine and 3-(4-bromobutyl)-1*H*-indole. Mp 121-122°C. ¹H NMR (DMSO-*d*₆): 1.45-1.55 (m, 2H); 1.65-1.75 (m, 2H); 2.35 (t, 2H); 2.50 (b s, 4H); 2.70 (t, 2H); 2.95 (b s, 4H); 6.95 (t, 1H); 7.05 (t, 1H); 7.10-7.15 (m, 2H); 7.25-7.30 (m, 2H); 7.35 (d, 1H); 7.50 (d, 1H); 10.75 (b s, 1H). Ms m/z: 402 (MH⁺).

5

5c, 3-{3-[4-(2,3-Dichlorophenyl)piperazin-1-yl]propyl}-5-fluoro-1*H*-indole

from 1-(2,3-dichlorophenyl)piperazine and 3-(3-chloropropyl)-5-fluoro-1*H*-indole. Mp 147-148°C. ¹H NMR (DMSO-*d*₆): 1.75-1.85 (m, 2H); 2.30-2.45 (t, 2H); 2.45-2.60 (m, 2H); 2.70 (t, 2H); 3.00 (b s, 4H); 3.35 (b s, 2H); 6.85-6.95 (m, 1H); 7.05-7.15 (m, 1H); 7.20 (s, 1H);

10 7.20-7.35 (m, 4H); 10.85 (b s, 1H). Ms m/z: 406 (MH⁺).

5d, 3-{4-[4-(2,3-Dichlorophenyl)piperazin-1-yl]butyl}-5-fluoro-1*H*-indole

from 1-(2,3-dichlorophenyl)piperazine and 3-(4-chlorobutyl)-5-fluoro-1*H*-indole. Mp 147-148°C. ¹H NMR (DMSO-*d*₆): 1.45-1.55 (m, 2H); 1.60-1.70 (m, 2H); 2.35 (t, 2H); 2.50 (b s, 4H); 2.65 (t, 2H); 2.95 (b s, 4H); 6.85-6.95 (m, 1H); 7.10-7.15 (m, 1H); 7.20 (s, 1H); 7.25-7.35 (m, 4H); 10.85 (b s, 1H). Ms m/z: 420 (MH⁺).

15

5e, 6-Chloro-3-{3-[4-(2,3-dichlorophenyl)piperazin-1-yl]propyl}-1*H*-indole

from 1-(2,3-dichlorophenyl)piperazine and 6-chloro 3-(3-iodopropyl)-1*H*-indole. ¹H NMR (CDCl₃): 1.95 (t, 2H); 2.55 (t, 2H); 2.65 (b s, 4H); 2.80 (t, 2H); 3.10 (b s, 4H); 6.95-7.05 (m, 2H); 7.10 (d, 1H); 7.10-7.20 (m, 2H); 7.35 (s, 1H); 7.55 (d, 1H); 7.95 (b s, 1H). Ms m/z: 422 (MH⁺), 424.

20

Example 6

25

6, 1-{4-[4-(2,3-Dichlorophenyl)piperazin-1-yl]butyl}-3,4-dihydroquinolin-2(1*H*)-one

The free base of 1-(2,3-dichlorophenyl)piperazine, hydrochloride (6.0 g) was liberated by the use ethyl acetate and aqueous ammonia. The remaining oil was dissolved in butanone (500 mL) followed by the addition of potassium carbonate (9.7 g), and the mixture was heated to reflux temperature. To this mixture was added a solution of 1-(4-bromobutyl)-3,4-dihydroquinolin-2(1*H*)-one (7.9 g) in butanone (150 mL), and the resulting mixture was boiled under reflux for 10 h. The mixture was filtered hot and purified by flash chromatography (eluent: ethyl acetate/triethylamine 100:4) giving the title compound which was precipitated as the hydrochloride salt (2.5 g). Mp 234-235 °C. ¹H NMR (DMSO-*d*₆):

30

1.55-1.65 (m, 2H); 1.75-1.85 (m, 2H); 2.55 (t, 2H); 2.85-2.90 (m, 2H); 3.05-3.20 (m, 4H); 3.25 (t, 2H); 3.40 (d, 2H); 3.55 (d, 2H); 3.95 (t, 2H); 7.00 (t, 1H); 7.15 (d, 1H); 7.20-7.30 (m, 3H); 7.30-7.40 (m, 2H); 11.35 (b s). MS m/z: 432 (MH⁺).

5 Example 7

7a, 3-[4-(2,3-Dichlorophenyl)piperazin-1-yl]-1-(5-fluoro-2,3-dihydro-1H-indol-1-yl)propan-1-one

3-Bromopropanoyl chloride (1 g in 10 ml dry dichloromethane) was added with stirring at 0°
 10 C to a suspension of 1 g Wang resin (Rapp polymere, loading 0.95 mmol/g) in 10 ml dry dichloromethane containing 5 equivalents of diisopropylethyl amine. The mixture was stirred overnight at room temperature, filtered and washed with dry dichloromethane (6 x 100 ml). A solution of 1-(2,3-dichlorophenyl)piperazine (2.5 equivalents) in dry acetonitrile containing diisopropylethyl amine (5 equivalents) was added to the dried resin and the
 15 mixture heated to 70°C for 3 hours. The mixture was cooled to room temperature and the resin washed with dry acetonitrile and dichloromethane and dried. A solution of AlCl₃ (1.1 equivalent) in dry acetonitrile (5 ml) was added to the resin followed by a solution of 5-fluoro-2,3-dihydro-1H-indole (3 equivalents) in dry acetonitrile (5 ml) and the mixture agitated for 3 hours. The reaction was quenched by addition of 2N NaOH (1.2 equivalent),
 20 filtrated and the product purified by solid phase ion exchange chromatography (Varian SCX columns) using Gilson ASPEC 232 XL. Further purification was performed on a PE Sciex API 150EX instrument equipped with IonSpray source and Shimadzu LC-8A/SLC-10A LC system. The LC conditions (50 x 20 mm YMC ODS-A with 5 µm particle size) were linear gradient elution with water/acetonitrile/trifluoroacetic acid (80:20:0.05) to
 25 water/acetonitrile/trifluoroacetic acid (10:90:0.03) in 7 min at 22.7 mL/min. Fraction collection was performed by split-flow MS detection. Purity was determined by integration of the UV trace (254 nm). The retention times RT are expressed in minutes.

LC/MS (m/z) 422 (MH⁺), RT = 2.49, purity: 70.57%

30 The following compounds were prepared in a similar manner

7b, 4-[4-(2,3-Dichlorophenyl)piperazin-1-yl]-1-(5-fluoro-2,3-dihydro-1H-indol-1-yl)butan-1-one: LC/MS (m/z) 436 (MH⁺), RT = 2.58, purity: 96.23%

7c, 5-[4-(2,3-Dichlorophenyl)piperazin-1-yl]-1-(5-fluoro-2,3-dihydro-1H-indol-1-yl)pentan-1-one: LC/MS (m/z) 450 (MH⁺), RT = 2.56, purity: 81.68%

7d, 3-[4-(2,3-Dichlorophenyl)piperazin-1-yl]-1-(3,4-dihydro-1H-isoquinolin-2-yl)propan-1-one: LC/MS (m/z) 418 (MH⁺), RT = 2.43, purity: 72.99%

7e, 4-[4-(2,3-Dichlorophenyl)piperazin-1-yl]-1-(3,4-dihydro-1H-isoquinolin-2-yl)butan-1-one: LC/MS (m/z) 432 (MH⁺), RT = 2.49, purity: 81.86%

7f, 5-[4-(2,3-Dichloro-phenyl)-piperazin-1-yl]-1-(3,4-dihydro-1H-isoquinolin-2-yl)pentan-1-one: LC/MS (m/z) 446 (MH⁺), RT = 2.49, purity: 98.39%

8a, 3-[4-(2-Chlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1H-indol-1-yl)propan-1-one: LC/MS (m/z) 370 (MH⁺), RT = 2.29, purity: 92.49%

8b, 5-[4-(2-Chlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1H-indol-1-yl)pentan-1-one: LC/MS (m/z) 398 (MH⁺), RT = 2.37, purity: 70.1%

8c, 3-[4-(3-Chlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1H-indol-1-yl)propan-1-one: LC/MS (m/z) 370 (MH⁺), RT = 2.33, purity: 81.15%

8d, 5-[4-(3-Chlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1H-indol-1-yl)pentan-1-one: LC/MS (m/z) 398 (MH⁺), RT = 2.41, purity: 96.58%

8e, 3-[4-(3-Chlorophenyl)piperazin-1-yl]-1-(5-fluoro-2,3-dihydro-1H-indol-1-yl)propan-1-one: LC/MS (m/z) 388 (MH⁺), RT = 2.37, purity: 92.8%

8f, 5-[4-(3-Chlorophenyl)piperazin-1-yl]-1-(5-fluoro-2,3-dihydro-1H-indol-1-yl)pentan-1-one: LC/MS (m/z) 416 (MH⁺), RT = 2.45, purity: 96.43%

8g, 3-[4-(2-Chlorophenyl)piperazin-1-yl]-1-(5-fluoro-2,3-dihydro-1H-indol-1-yl)propan-1-one: LC/MS (m/z) 388 (MH⁺), RT = 2.33, purity: 93.11%

- 8h, 4-[4-(2-Chlorophenyl)piperazin-1-yl]-1-(5-fluoro-2,3-dihydro-1H-indol-1-yl)butan-1-one: LC/MS (m/z) 402 (MH⁺), RT = 2.43, purity: 89.76%
- 5 8i, 3-[4-(3-Chlorophenyl)piperazin-1-yl]-1-(3,4-dihydro-1H-isoquinolin-2-yl)propan-1-one:
LC/MS (m/z) 384 (MH⁺), RT = 2.31, purity: 92.21%
- 8j, 5-[4-(3-Chlorophenyl)piperazin-1-yl]-1-(3,4-dihydro-1H-isoquinolin-2-yl)pentan-1-one:
LC/MS (m/z) 412 (MH⁺), RT = 2.37, purity: 95.37%
- 10 8k, 3-[4-(2-Chlorophenyl)piperazin-1-yl]-1-(3,4-dihydro-1H-isoquinolin-2-yl)propan-1-one: LC/MS (m/z) 384 (MH⁺), RT = 2.27, purity: 91.51%
- 8l, 4-[4-(2-Chlorophenyl)piperazin-1-yl]-1-(3,4-dihydro-1H-isoquinolin-2-yl)butan-1-one:
LC/MS (m/z) 398 (MH⁺), RT = 2.35, purity: 97.56%
- 15 9a, 1-(2,3-Dihydro-1H-indol-1-yl)-3-[4-(2-fluoro-phenyl)-piperazin-1-yl]propan-1-one:
LC/MS (m/z) 354 (MH⁺), RT = 2.14, purity: 91.64%
- 9b, 1-(2,3-Dihydro-1H-indol-1-yl)-4-[4-(2-fluoro-phenyl)-piperazin-1-yl]butan-1-one:
20 LC/MS (m/z) 368 (MH⁺), RT = 2.24, purity: 76.25%
- 9c, 1-(2,3-Dihydro-1H-indol-1-yl)-5-[4-(2-fluoro-phenyl)-piperazin-1-yl]pentan-1-one:
LC/MS (m/z) 382 (MH⁺), RT = 2.22, purity: 87.9%
- 25 9d, 1-(5-Fluoro-2,3-dihydro-1H-indol-1-yl)-3-[4-(2-fluoro-phenyl)-piperazin-1-yl]propan-1-one: LC/MS (m/z) 372 (MH⁺), RT = 2.22, purity: 76.87%
- 9e, 1-(5-Fluoro-2,3-dihydro-1H-indol-1-yl)-4-[4-(2-fluoro-phenyl)-piperazin-1-yl]butan-1-one:
LC/MS (m/z) 386 (MH⁺), RT = 2.31, purity: 86.01%
- 30 9f, 1-(5-Fluoro-2,3-dihydro-1H-indol-1-yl)-5-[4-(2-fluoro-phenyl)-piperazin-1-yl]pentan-1-one: LC/MS (m/z) 400 (MH⁺), RT = 2.31, purity: 97.52%

9g, 3-[4-(2,4-Difluoro-phenyl)-piperazin-1-yl]-1-(2,3-dihydro-1H-indol-1-yl)propan-1-one:

LC/MS (m/z) 372 (MH⁺), RT = 2.2, purity: 94.79%

9h, 4-[4-(2,4-Difluoro-phenyl)-piperazin-1-yl]-1-(2,3-dihydro-1H-indol-1-yl)butan-1-one:

5 LC/MS (m/z) 386 (MH⁺), RT = 2.29, purity: 79.75%

9i, 5-[4-(2,4-Difluoro-phenyl)-piperazin-1-yl]-1-(2,3-dihydro-1H-indol-1-yl)pentan-1-one:

LC/MS (m/z) 400 (MH⁺), RT = 2.29, purity: 99.06%

10 9j, 3-[4-(2,4-Difluoro-phenyl)-piperazin-1-yl]-1-(5-fluoro-2,3-dihydro-1H-indol-1-

yl)propan-1-one: LC/MS (m/z) 390 (MH⁺), RT = 2.27, purity: 87.99%

9k, 1-(3,4-Dihydro-1H-isoquinolin-2-yl)-4-[4-(2-fluoro-phenyl)-piperazin-1-yl]-butan-1-

one: LC/MS (m/z) 382 (MH⁺), RT = 2.22, purity: 87.75%

15

9l, 1-(3,4-Dihydro-1H-isoquinolin-2-yl)-5-[4-(2-fluoro-phenyl)-piperazin-1-yl]-pentan-1-

one: LC/MS (m/z) 396 (MH⁺), RT = 2.22, purity: 85.52%

9m, 3-[4-(2,4-Difluoro-phenyl)-piperazin-1-yl]-1-(3,4-dihydro-1H-isoquinolin-2-yl)-

20 propan-1-one: LC/MS (m/z) 385 (MH⁺), RT = 2.22, purity: 87.01%

9n, 5-[4-(2,4-Difluoro-phenyl)-piperazin-1-yl]-1-(3,4-dihydro-1H-isoquinolin-2-yl)-pentan-

1-one: LC/MS (m/z) 414 (MH⁺), RT = 2.31, purity: 87.84%

25 10a, 3-[4-(3,4-Dichlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1H-indol-1-yl)propan-1-one:

LC/MS (m/z) 404 (MH⁺), RT = 2.47, purity: 76.03%

10b, 4-[4-(3,4-Dichlorophenyl)piperazin-1-yl]-1-(2,3-dihydro-1H-indol-1-yl)butan-1-one:

LC/MS (m/z) 418 (MH⁺), RT = 2.58, purity: 99.32%

30

10c, 3-[4-(3,4-Dichlorophenyl)piperazin-1-yl]-1-(5-fluoro-2,3-dihydro-1H-indol-1-

yl)propan-1-one: LC/MS (m/z) 422 (MH⁺), RT = 2.52, purity: 80.99%

10d, 3-[4-(3,4-Dichlorophenyl)piperazin-1-yl]-1-(3,4-dihydro-1H-isoquinolin-2-yl)-propan-1-one: LC/MS (m/z) 418 (MH⁺), RT = 2.45, purity: 83.31%

10e, 5-[4-(3,4-Dichlorophenyl)piperazin-1-yl]-1-(3,4-dihydro-1H-isoquinolin-2-yl)-pentan-1-one: LC/MS (m/z) 446.1 (MH⁺), RT = 2.52, purity: 98.79%

Pharmacological Testing

The compounds of the invention were tested by the use of well-recognised and reliable methods. The tests were as follows:

Inhibition of the binding of [³H]YM-09151-2 to human dopamine D₄ receptors

By this method the inhibition by drugs of the binding of [³H]YM-09151-2 (0.06 nM) to membranes of human cloned dopamine D_{4.2} receptors expressed in CHO-cells is determined *in vitro*. Method modified from NEN Life Science Products, Inc., technical data certificate PC2533-10/96. The results are given in the following Table 1 as IC₅₀-values.

Inhibition of the binding of [³H]Spiperone to human D₃ receptors

By this method the inhibition by drugs of the binding [³H]Spiperone (0.3 nM) to membranes of human cloned dopamine D₃ receptors expressed in CHO-cells is determined *in vitro*.

Method modified from MacKenzie et al. *Eur. J. Pharm.-Mol. Pharm. Sec.* 1994, 266, 79-85. The results are given in the following Table 1.

Inhibition of binding of [³H]Prazosine to rat alpha-1 receptors

By this method the inhibition by drugs of the binding of [³H]Prazosine (0.25 nM) to alpha-1 receptors in membranes from rat brain is determined *in vitro*. Method modified from Hyttel et al. *J. Neurochem.* 1985, 44, 1615-1622. The results are given in the following Table 1.

Comp. No.	D ₄ -bind.	D ₃ -bind.	Alpha-1
1a	3.3	100	59
1b	2.8	3.0	1100
1c	39	10	160
1d	0.92	20	97
1e	2.1	50	17
2a	1.8	31	68
2b	12	3.1	10 %
2c	18	22	190
2d	1.2	4.0	31
2e	1.6	17	40
3	11	6.8	-3 %
4	500	40	4800
5a	2.2	2.8	410
5b	14	1.1	570
5c	3.9	6.8	960
5d	8.6	1.0	720
5e	27	93%	470
6	16	1.8	43
7a	25	73%	38%
7b	53	65%	6%
7c	61	92%	45%
7d	6.0	85%	44%
7e	10	94%	31%
7f	26	95%	34%
8k	8.0	73%	92%
8l	4.0	88%	74%
9k	9.0	92%	69%

Table 1: Binding Data (IC₅₀ values in nM or as % inhibition of binding at 100 nM)

The compounds were also tested in in the following test:

Inhibition of the binding of [³H]Spiroperidol to D₂ receptors

The compounds were tested with respect to affinity for the dopamine D₂ receptor by determining their ability to inhibit the binding of [³H]-spiroperidol to D₂ receptors by the
5 method of Hyttel et al. *J. Neuroche.* 1985, 44, 1615.

In general, the compounds of the invention have been found to have high affinity for dopamine D₄ receptors and dopamine D₃ receptors. The compounds have only weak or no affinity for the dopamine D₂ receptor.

10

One important effects of adrenergic alpha-1-receptor blockade is postural hypotension resulting from a fall in central venous pressure due to dilation of small capacitance vessels. This effect may further be accompanied by a decrease in cardiac output. Certain compounds of the invention have the further advantage of having only very weak effect at adrenergic
15 alpha-1-receptors which imply a low propensity to cause orthostatic hypotension.

Certain of the compounds interact with central serotonergic receptors, such as the 5-HT_{1A} and/or the 5-HT_{2A} receptors and/or they act as 5-HT reuptake inhibitors.

20 Accordingly, the compounds of the invention are considered useful in the treatment of psychosis including the positive and negative symptoms of schizophrenia, affective disorders such as generalised anxiety disorder, panic disorder, and obsessive compulsive disorder, depression, aggression cognitive disorders and dyskinesia induced by treatment with L-dopa comprising.

25

The pharmaceutical compositions of this invention or those which are manufactured in accordance with this invention may be administered by any suitable route, for example orally in the form of tablets, capsules, powders, syrups, etc., or parenterally in the form of solutions for injection. For preparing such compositions, methods well known in the art may
30 be used, and any pharmaceutically acceptable carriers, diluents, excipients or other additives normally used in the art may be used.

Conveniently, the compounds of the invention are administered in unit dosage form containing said compounds in an amount of about 0.01 to 100 mg.

The total daily dose is usually in the range of about 0.05 - 500 mg, and most preferably about 0.1 to 50 mg of the active compound of the invention.

Formulation Examples

The pharmaceutical formulations of the invention may be prepared by conventional methods in the art.

For example: Tablets may be prepared by mixing the active ingredient with ordinary adjuvants and/or diluents and subsequently compressing the mixture in a conventional tableting machine. Examples of adjuvants or diluents comprise: corn starch, potato starch, talcum, magnesium stearate, gelatine, lactose, gums, and the like. Any other adjuvants or additives usually used for such purposes such as colourings, flavourings, preservatives etc. may be used provided that they are compatible with the active ingredients.

Solutions for injections may be prepared by dissolving the active ingredient and possible additives in a part of the solvent for injection, preferably sterile water, adjusting the solution to desired volume, sterilisation of the solution and filling in suitable ampules or vials. Any suitable additive conventionally used in the art may be added, such as tonicity agents, preservatives, antioxidants, etc.

Typical examples of recipes for the formulation of the invention are as follows:

1) Tablets containing 5.0 mg of active compound calculated as the free base:

Compound of formula I	5.0 mg
Lactose	60 mg
Maize starch	30 mg
Hydroxypropylcellulose	2.4 mg
Microcrystalline cellulose	19.2 mg
Croscarmellose Sodium Type A	2.4 mg
Magnesium stearate	0.84 mg

2) Tablets containing 0.5 mg of active compound calculated as the free base:

	Compound of formula I	0.5 mg
	Lactose	46.9 mg
5	Maize starch	23.5 mg
	Povidone	1.8 mg
	Microcrystalline cellulose	14.4 mg
	Croscarmellose Sodium Type A	1.8 mg
	Magnesium stearate	0.63 mg
10		

3) Syrup containing per millilitre:


	Compound of formula i	25 mg
15	Sorbitol	500 mg
	Hydroxypropylcellulose	15 mg
	Glycerol	50 mg
	Methyl-paraben	1 mg
	Propyl-paraben	0.1 mg
20	Ethanol	0.005 ml
	Flavour	0.05 mg
	Saccharin sodium	0.5 mg
	Water	ad 1 ml

25 4) Solution for injection containing per millilitre:

	Compound of formula I	0.5 mg
	Sorbitol	5.1 mg
	Acetic Acid	0.05 mg
30	Saccharin sodium	0.5 mg
	Water	ad 1 ml

Patent Claims

1. A halogen substituted 4-phenyl-1-piperazinyl derivative of formula I



- 5 wherein W is C, CH or N, and the dotted line emanating from W indicates a bond when W is C and no bond when W is N or CH;

R¹ and R² are independently selected from hydrogen and halogen, provided at least one of R¹ and R² is a halogen atom;


10

R³ is selected from hydrogen, halogen, C₁-₆-alkyl, C₂-₆-alkenyl, C₂-₆-alkynyl, trifluoromethyl, C₁-₆-alkoxy, aryloxy, aralkoxy, hydroxy, amino, C₁-₆-alkylamino, di(C₁-₆-alkyl)amino, nitro and cyano

- 15 n is 2, 3, 4 or 5;

X is CH₂, O, S, CO, CS, SO or SO₂; and

Q is a group of formula



20

wherein Z is a chain of 3 to 4 chain members, wherein the chain members are selected from C, CH, CH₂, CO, N and NH, provided that only one of the chain members may be N or NH, and said chain Z optionally containing one or two double bonds;

5

R⁴, R⁵, R⁶, R⁷, R⁸ and R⁹ are independently selected from R⁴, R⁵, R⁶, R⁷, R⁸ and R⁹ are independently selected from hydrogen, halogen, trifluoromethyl, nitro, cyano, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₁₋₆-alkoxy, C₁₋₆-alkylthio, C₁₋₆-alkylsulfonyl, hydroxy, hydroxy-C₁₋₆ alkyl, amino, C₁₋₆-alkylamino,

10 di(C₁₋₆-alkyl)amino, acyl,aminocarbonyl, C₁₋₆-alkylaminocarbonyl and di(C₁₋₆-alkyl)aminocarbonyl;

provided that X is not O or S when the group Q is attached via an N atom;

and any of its enantiomers and acid addition salts thereof.

15

2. A compound according to claim 1 wherein R¹ and R² are both halogen.

3. A compound according to claim 2 wherein R¹ and R² are both chloro.

20

4. A compound according to claim 1 wherein one of R¹ and R² is halogen and the other is hydrogen.

5. A compound according to claim 4 wherein R² is halogen.

25

6. A compound according to claim 4 wherein R¹ is halogen.

7. A compound according to claim 1 wherein R¹ and R² are independently selected from hydrogen or chloro.

30

8. A compound according to claims 1-7 wherein W is N.

9. A compound according to claims 1 to 8 wherein X is CO or CH₂.

10. A compound according to claims 1-9 wherein Z is a chain of 3 to 4 chain members,
wherein the chain members are selected from C, CH, CH₂, N and NH, provided that only
5 one of the chain members may be N or NH.
11. A compound according to claim 10 wherein Q is optionally substituted 1-indoliny.
- 10 12. A compound according to claim 10 wherein Q is optionally substituted 3-indolyl.
13. A compound according to claim 10 wherein Q is optionally substituted 1-indanyl.
- 15 14. A compound according to claim 10 wherein Q is optionally substituted 2-oxo-1, 2, 3, 4-tetrahydroquinoliny.
15. A compound according to claim 10 wherein Q is optionally substituted 1,2,3,4-tetrahydroisoquinoliny.
- 20 16. A compound according to claims 9 to 15 wherein Q is unsubstituted or substituted with halogen.
17. A compound according to claim 1 wherein R³ is hydrogen or R³ is attached in the
25 para position on the phenyl ring.
18. A compound according to claim 17 wherein R³ is halogen.
19. A pharmaceutical composition characterised in that it comprises a compound of any of
30 claims 1 to 18 in a therapeutically effective amount together with one or more pharmaceutically acceptable carriers or diluents.
20. Use of a compound of any of Claims 1 to 18 for the manufacture of a medicament useful in the treatment of psychosis including the positive and negative symptoms of

schizophrenia, affective disorders such as generalised anxiety disorder, panic disorder, and obsessive compulsive disorder, depression, aggression, cognitive disorders and dyskinesia induced by treatment with L-dopa.

- 5 21. A method of treating psychosis including the positive and negative symptoms of schizophrenia, affective disorders such as generalised anxiety disorder, panic disorder, and obsessive compulsive disorder, depression, aggression, cognitive disorders and dyskinesia induced by treatment with L-dopa comprising administration of a therapeutically acceptable amount of a compound according to any of claims 1 to 18.