




The invention relates to Vitamin D₃ analogs, particularly 16-ene-23-yne-trifluoro analogs of Vitamin D₃ of the formula



I

5


wherein R is hydrogen, fluorine or hydroxy, and X is =CH₂ or when R is hydroxy, X is hydrogen or =CH₂.

The compounds of formula I induce differentiation and
 10 inhibition of proliferation in various skin and cancer cell lines.
 Accordingly, the compounds of formula I are useful as agents for the
 treatment of hyperproliferative skin diseases, such as, psoriasis.
 Compounds of formula I are also useful in the treatment of neoplastic
 diseases, such as, leukemia and sebaceous gland diseases, such as,
 15 acne or seborrheic dermatitis.

The invention also relates to a pharmaceutical composition
 comprising a compound of formula I and a method of treating the
 above mentioned disease states by administration of the compound of
 20 formula I.

In a preferred embodiment, R is hydroxy.


An embodiment of the invention is a mixture comprising epimers of the formula



wherein R and X are as described above.

The compounds of formula I are prepared as hereafter
10 described in Schemes I-II and the Examples.

SCHEME I




In the above Scheme I, the compound of formula II, [3aS,[3(S*), 3a α ,7 α ,7a β]]-[[3a,4,5,6,7,7a-hexahydro-3 α -methyl-3-(1-methyl-3-butynyl)-1H-inden-7-yl]oxy]-trimethylsilane, a known compound, is converted to a compound of formula III, by reaction with n-butyl-
5 lithium and trifluoroacetone in a ether solvent such as tetrahydrofuran. The reaction is carried out at -100°C to 0°C , preferably -78°C.

The compound of formula III is converted to the compound of formula IV by reaction with tetrabutyl-ammonium fluoride in a ether
10 solvent, such as tetrahydrofuran. The reaction is carried out at 0-50°C, preferably room temperature, preferably under an argon atmosphere.

The compound of formula IV is converted to the compound of
15 formula V by reaction with pyridinium dichromate and pyridinium p-toluene sulfonate in a chlorinated hydrogen solvent such as methylene chloride.

The compound of formula V is converted to the compound of
20 formula VI by reaction with trimethylsilyl-imidazole in a chlorinated solvent such as methylene chloride. Preferably, the reaction is carried out under an argon atmosphere.

SCHEME II



In the above Scheme II, the compound of formula VI is converted to the compound of formula Ib by reaction with [3S-(1Z,3 α ,5 β)]-[2-[3,5-bis[[1,1-dimethylethyl]dimethylsilyl]oxy]-2-methylenecyclohexylidene]ethyl]diphenylphosphine oxide in tetrahydrofuran, preferably under argon in the presence of n-butyl lithium as a base. This reaction is followed by removal of the protecting silyl groups using tetrabutylammonium fluoride in tetrahydrofuran as a solvent.

10 Alternatively, the compound of formula VI is converted to the compound of formula Ic by reaction with [3R-(3 α ,5 β ,Z)-3,5-bis[[1,1-dimethylethyl]dimethylsilyl]oxy] cyclohexylidene]ethyl]diphenylphosphine oxide in tetrahydrofuran, preferably under argon, followed by removal of the protecting silyl groups.

15 Alternatively, the compound of formula VI is converted to the compound of formula Id by reaction with [3S-(3 α ,5 β ,Z)-2-[2-methylene-3-fluoro-5-[[1,1 dimethylethyl]dimethyl-silyl]oxy] cyclohexylidene]ethyl]diphenyl phosphine oxide in tetrahydrofuran, preferably under argon, followed by removal of the protecting silyl groups.

20 Alternatively, the compound of formula VI is converted to the compound of formula Ie by reaction with [5S,Z)-2-[2-[2-methylene-5-[[1,1-dimethylethyl]dimethylsilyl] oxy]cyclo-hexylidene]ethyl] diphenyl phosphine oxide in tetrahydrofuran, preferably under argon, followed by removal of the protecting silyl groups.

30 Any conventional separation method known to those skilled in the art can be used at any point in the preparation of a compound of formula I to separate the epimeric mixture to either the (R) or (S) epimer.

35 Thus, the invention is concerned with compounds of formula I representing mixtures of the (25R) and (25S) epimers as well as with the individual (25R) and (25S) epimers or diastereomers.

The compound of formula I can be administered orally, for example, in the form of tablets, coated tablets, dragees, hard or soft gelatin capsules, solutions, emulsions or suspensions. The
5 administration can, however, also be effected rectally, for example, in the form of suppositories or parenterally, for example, in the form of injection solutions. A composition in accordance with the invention can be processed with pharmaceutically inert, inorganic or organic
10 excipients for the manufacture of tablets, coated tablets, dragees and hard gelatin capsules. Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts, and the like, can be used as such excipients, for example, for tablets, dragees and hard gelatin capsules. Suitable
excipients for soft gelatin capsules are, for example, vegetable oils, waxes, fats, semi-solid and liquid polyols, and the like; depending on
15 the nature of the active ingredient. No excipients are, however, usually required in the case of soft gelatin capsules. Suitable excipients for the preparation of solutions and syrups, are, for example, water, polyols, saccharose, invert sugar and glucose.

20 Suitable excipients for injection solutions are, for example, water, alcohols, polyols, glycerol, vegetable oils, and the like. Suitable excipients for suppositories are, for example, natural or hardened oils, waxes, fats, semi-liquid or liquid polyols and the like.

25 Moreover, the pharmaceutical preparations can contain preservatives, solubilizers, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, masking agents or antioxidants.

30 The compounds of formula I as described above can be administered orally or by injection, for the treatment of neoplastic diseases such as leukemia, to warm-blooded animals which need such treatment. More specifically, the compounds of formula I as described
above can be administered orally to an adult human in dosages that
35 are in the range of about 0.25 to 50 μg per day for the treatment of neoplastic diseases such as leukemia.

The compounds of formula I as described above can be administered orally, for the treatment of hyperproliferative skin diseases such as psoriasis, basal cell carcinomas, disorders of
5 keratinization, and keratosis, to warm-blooded animals which need such treatment. More specifically, the compounds of formula I as described above can be administered orally to an adult human in dosages that are in the range of about 0.25 to 50 μg per day for the treatment of hyperproliferative skin diseases such as psoriasis, basal
10 cell carcinomas, disorders of keratinization, and keratosis. These compounds can be administered orally for the treatment of acne in humans at a dosage of about 0.25 to 50 μg per day; preferably 0.5 to 5 μg per day.

15 The compounds of formula I as described above can be administered topically, for the treatment of hyperproliferative skin diseases such as psoriasis, basal cell carcinomas, disorders of keratinization, and keratosis, to warm-blooded animals which need such treatment. More specifically, the compounds of formula I as
20 described above can be administered topically in dosages that are in the range of about 0.5 to 100 μg per gram of topical formulation per day, for the treatment of hyperproliferative skin diseases such as psoriasis, basal cell carcinomas, disorders of keratinization, and keratosis.

25 The compounds of formula I as described above can also be administered topically for the treatment of sebaceous gland diseases such as acne or seborrheic dermatitis in dosages that are in the range of 0.5 to 100 mg per gram of topical formulation per day.

30 The useful activity of the compounds of formula I as agents for the treatment of neoplastic diseases can be demonstrated by the following test procedures.

HL-60 Cell Differentiation

The induction of differentiation of HL-60 cells was assayed by measuring their oxidative burst potential via the reduction of NBT (Nitrobluetetrazolium).

HL-60 cells were maintained in RPMI 1640 medium supplemented with 10% FCS, 2mM L-glutamine, 1mM sodium pyruvate, 1% non-essential amino acids, 50 U/ml penicillin and 50 μ g/ml streptomycin (=RPMI/FCS). 30,000 cells/90 μ l of RPMI/FCS were seeded into flat-bottomed microtiter wells. 10 μ l of vitamin D derivatives diluted in complete medium were added at the same time to yield final concentrations between 10^{-11} and 10^{-6} M (stock solutions of 10^{-2} M in ethanol were kept at -20°C and protected from light).

After 3 days, the medium was removed with a multichannel pipette and replaced with 100 μ l of NBT solution [1 mg/ml in PBS with 200 mM phorbol myristate acetate (PMA)]. Following an additional hour incubation at 37°C , the NBT solution was removed and 100 μ l of 10% SDS in 0.01 N HCl was added. The amount of the reduced NBT was quantified photometrically at 540 nm using an automated plate reader. The mean of 3 wells was calculated. S.E.M. were between 5 and 10%. Values were expressed as percent of maximal differentiation achieved with 100 - 1000 nM calcitriol in the same experiment. The concentration (nM) leading to 50% of this maximal value is determined graphically and given as ED_{50} in Table I below.

TABLE I

COMPOUND	ED₅₀(nM)
1,25-Dihydroxycholecalciferol (calcitriol)	8.0
1,25-(R,S)Dihydroxy-16-ene-23-yne-26-trifluoro-cholecalciferol	0.15
1,25(R,S)-Dihydroxy-16-ene-23-yne-26-trifluoro-19-nor-cholecalciferol	0.37
1 α -Fluoro-25(R,S)-hydroxy-16-ene-23-yne-26-trifluoro-cholecalciferol	3.70
25(R,S)-Hydroxy-16-ene-23-yne-26-trifluoro-cholecalciferol	160.00

The useful activity of compounds of formula I as agents for the
5 treatment of hyperproliferative skin disease can be determined by
the following.

Inhibition of Keratinocytes Proliferation

10 **HaCaT cell line** - The immortalized human cell line HaCaT was
used. ³H-thymidine incorporation was measured in exponentially
growing cultures after 6 days of culture in presence of the test
compound.

15 **Cell culture** - HaCaT cells were cultured in Dulbecco's Modified Eagle
Medium (DMEM) and Nutrient Mixture Ham's F12 (F12), 3:1 (v/v, ICN)
containing 4.5 g/l glucose and supplemented with 10% fetal calf serum
(Gibco, FCS), L-glutamine (Gibco, 2mM), penicillin (Gibco, 50 UI/ml),
streptomycin (Gibco, 50 μ g/ml), EGF (10 ng/ml), hydrocortisone (400
20 ng/ml), cholera toxin (8.5 ng/ml) and insulin (5 ng/ml). The cells
were maintained in a humidified atmosphere containing 5% CO₂ and
95% air and passaged every 3-4 days.

Inhibition of ³H-thymidine uptake - HaCaT cells (250 cells in
25 complete culture medium) were seeded into 96-well culture dishes

and incubated at 37°C with 5% CO₂ and 95% air for 6 days. Inhibitors, dissolved at 10 x concentration in 1% ethanol, were added immediately at the beginning of the assay. ³H-thymidine (5 Ci/mmol, Amersham) was added at a concentration of 1 µCi/well and cells were pulse-labelled for the last 6 hours of the growth period. Cells were then trypsinized for 10 minutes at 37°C under a vigorous agitation and harvested on to a 96-well GF/C filter plate (Uni Filter, Packard) using a Micro Mate 196 cell harvester (Packard). After drying at 40°C under vacuum for 20-30 minutes, 2 µl of Micro Scint 0 scintillator (Packard) were added and the radioactivity bound to the filters was counted on a TOP COUNT (Packard).

The results measured as IC₅₀ are set forth in Table II below.

15

TABLE II

COMPOUND	IC ₅₀ (nM)
1,25(R,S)-Dihydroxy-16-ene-23-yne-26-trifluorocholecalciferol	10.0
1,25(R,S)-Dihydroxy-16-ene-23-yne-26-trifluoro-19-nor-cholecalciferol	5.1
1α-Fluoro-25(R,S)-hydroxy-16-ene-23-yne-26-trifluorocholecalciferol	11.0
25(R,S)-Hydroxy-16-ene-23-yne-26-trifluoro-cholecalciferol	650.00

The useful activity of compounds of formula I as agents for the treatment of sebaceous gland diseases can be demonstrated by the following.

20

Inhibition of Human Sebocyte Proliferation In Vitro

Sebaceous cells were isolated from adult human sebaceous glands by a combination of enzymatic and mechanical methods (Doran et al., Characterization of Human Cells In Vitro, J. Invest. Dermatol. 96:34-8 (1991)). The cells were cultured in Iscove's medium containing 10% fetal calf serum and 4 µg/ml dexamethasone on a

25

layer of growth-arrested 3T3 mouse fibroblasts. Cells were plated in medium without the test compound and then given test compound in fresh medium, containing the test compound, every 48 hours. On the day of harvesting, the cultures were rinsed with 0.03% EDTA in PBS, to
5 remove only the 3T3 fibroblasts, followed by incubation in 0.05% trypsin/0.03% EDTA. The cells were suspended, mixed vigorously to prepare a single cell suspension and counted in a hemocytometer.

Stock solutions of compounds were made up as 10^{-2} M solutions
10 in degassed 100% ethanol and stored at -20°C in the dark. During experimental use, the solutions, which have been aliquoted, were brought to room temperature and used by diluting directly into complete medium to appropriate concentration.

15 The compounds were tested for the inhibition of proliferation of sebaceous cell growth in vitro at 10^{-6} , 10^{-7} and 10^{-8} M.

The results are summarized in Table III below as the amount of compound necessary to inhibit the proliferation of sebaceous cells by
20 50% (ED_{50}) in nM as compared to a vehicle-treated culture.

TABLE III

COMPOUND	$\text{ED}_{50}(\text{nM})$
1,25(R,S)-Dihydroxy-16-ene-23-yne-26-trifluoro-cholecalciferol	1.0

25 **Calcium liability (tolerance test in mice)**

Profound changes in calcium homeostasis strongly affect the weight development of the animals. This parameter was used as a primary test for tolerance.

30

Mice (25-30 g body weight) received daily subcutaneous administrations of the vitamin D derivative for 4 consecutive days.

Body weight was registered just before and at the end of a 5 day treatment period. The "highest tolerated dose" (HTD) is the dose which results in zero weight gain during this treatment period.

- 5 The results are set forth in Table IV below.

TABLE IV

COMPOUND	HTD(μ g/kg)
1,25(R,S)-Dihydroxy-16-ene-23-yne-26-trifluoro-cholecalciferol	1.0
1,25(R,S)-Dihydroxy-16-ene-23-yne-26-trifluoro-19-nor-cholecalciferol	2.0
1 α -Fluoro-25(R,S)-hydroxy-16-ene-23-yne-26-trifluoro-cholecalciferol	20.0
25(R,S)-Hydroxy-16-ene-23-yne-26-trifluoro-cholecalciferol	600.00

- 10 The following Examples are provided to further describe the invention and are not intended to limit it in any way.

Example 1

- 15 [3aS-[3(1S*),3aa,7a,7ab]]-1,1,1-Trifluoro-6-[3a,4,5,6,7,7a-hexahydro-3a-methyl-7-[trimethylsilyl]oxy-1H-inden-3-yl]-2-methyl-3-heptyn-2-ol (epimers)

- To the solution of 1.1 g (3.80 mmol) of [3aS,[3(S*),3aa,7a,7ab]]-
20 [[3a,4,5,6,7,7a-hexahydro-3a-methyl-3-(1-methyl-3-butynyl)-1H-inden-7-yl]oxy]-trimethylsilane in 20 ml anhydrous tetrahydrofuran at -78°C was added with stirring 2.61 ml (4.18 mmol) of 1.6M n-butyllithium. After stirring for one hour, 0.68 ml (7.6 mmol) of 1,1,1-trifluoroacetone was added, and the reaction mixture was stirred for
25 an additional 1 hr at 78°C. The reaction was quenched with saturated brine and warmed up to room temperature. After dilution with water, it was extracted with ethyl acetate. The combined extracts

were washed with water, dried over sodium sulfate, and evaporated to dryness. The crude product was purified by FLASH chromatography on silica gel column with hexane-ethyl acetate 20:1, to give 1.47 g (96.5%) of the amorphous title compound.

5

Example 2

[3aS-[3(1S*),3aa,7a,7ab]]-3a,4,5,6,7,7a-Hexahydro-3a-methyl-3-(6,6,6-trifluoro-5-hydroxy-1,5-dimethyl -3-hexynyl)-1H-inden-7-ol
10 (epimers)

To a stirred solution of 1.47 g (3.65 mmol) of [3aS,[3(1S*),3aa,7a,7ab]]-1,1,1-trifluoro-6-[3a,4,5,6,7,7a-hexahydro-3a-methyl-7-[(trimethylsilyl)oxy]-1H-inden-3-yl]-2-methyl-3-heptyn-2-ol (epimers) in 15 ml of anhydrous tetrahydrofuran at room temperature under argon was added 8 ml (8.0 mmol) of 1M tetrabutyl ammonium fluoride. The reaction mixture was stirred for 1.5 hours, and then quenched by the addition of ice. It was then diluted with water and extracted with ethyl acetate. The combined extracts were washed with 2N potassium bicarbonate, water until neutral pH, and brine, dried over sodium sulfate, and evaporated to dryness. The crude product was purified by FLASH chromatography on a silica gel column with hexane-ethyl acetate 2:1, to give 1.2 g (100%) of crystalline title compound; m.p. 78-80°C, $[\alpha]_D^{25} + 20.4^\circ$ (c 0.5, ethanol ("EtOH")); $^1\text{H-NMR}$ (CDCl_3): d 1.07 (s, 3H, CH_3), 1.09 (d, 3H, J + 6.5 Hz, CH_3), 1.40 (dt, 1H, $J_{\text{vic}} = 3$ and 12.5 Hz, $J_{\text{gem}} = 12.5$ Hz, CH of CH_2), 1.58 (s, 3H, CH_3), 1.71 - 1.98 (m, 5H), 2.00 (m, 1H, $J_{\text{vic}} = 3$ and 7 Hz, $J_{\text{gem}} = 14.5$ Hz, CH of CH_2), 2.23 - 2.45 (m, 5H); 4.19 (brs, 1H, CH), 5.40 (brm, 1H, CH); Calcd. for $\text{C}_{18}\text{H}_{25}\text{F}_3\text{O}_2$: C 65.49, H 7.63; Found: C 65.59, H 7.77.
30

Example 3

[3aR-[1(R*),3aa,7ab]]-3a,4,5,6,7,7a-Hexahydro-7a-methyl-1-(6,6,6-trifluoro-5-hydroxy-1,5-dimethyl-3-hexynyl)-4H-inden-4-one
5 (epimers)

To a stirred solution of 300 mg (0.91 mmol) of [3aS,[3(1S*),3aa,7a,7ab]]-3a,4,5,6,7,7a-Hexahydro-7a-methyl-1-(6,6,6-trifluoro-5-hydroxy)-1,5-dimethyl-3-hexynyl)-1H-inden-7-ol (epimers) in 8 ml
10 of anhydrous methylene chloride was added 1.402 g (3.73 mmol) of pyridinium dichromate and 70 mg pyridinium-p-toluene sulfonate, and the reaction mixture was stirred for 4 hours. 20 ml of ether was added, stirred for 20 min and filtered over Celite. The Celite plug was washed with 3 x 50 ml of ether. The combined filtrates were washed
15 with 20 ml of ice cold 1N HCl, water, 2N KHCO₃ (40 ml) and water and brine. The aqueous layers were extracted with 2 x 100 ml ether-ethyl acetate (1:1). The organic layers were dried over sodium sulfate and evaporated to dryness. The crude product was purified by FLASH chromatography on silica gel with hexane-ethyl acetate (3:1), to give
20 272 mg (91.2%) of amorphous title compound.

Example 4

[3aR-[1(1R*),3aa,7ab]]-3,3a,5,6,7,7a-Hexahydro-7a-methyl-1-(6,6,6-trifluoro-1,5-dimethyl-5-[(trimethylsilyl)oxy]-3-hexynyl)-4H-inden-
25 4-one (epimers)

To a stirred solution of 272 mg (0.828 mmol) of [3aR-[1(R*),3aa,7ab]]-3,3a,5,6,7,7a-hexahydro-7a-methyl-1-(6,6,6-trifluoro-5-hydroxy-1,5-dimethyl-3-hexynyl)-4H-inden-4-one (epimers) in 6 ml
30 of anhydrous methylene chloride was added 0.79 mg (5.38 mmol) of trimethylsilyl-imidazole under argon. The reaction mixture was stirred at room temperature for 2.5 hours, and then quenched with 7 ml water. After continuous stirring for 30 min, it was extracted with
35 3 x 120 ml ethyl acetate. The organic layers were washed five times with a mixture of water and brine, dried over sodium sulfate and

evaporated to dryness. The crude product was purified by FLASH chromatography on a silica gel column with hexane-ethylacetate 10:1, to give 314 mg (94.6%) of the title compound.

5

Example 5

1,25(R,S)-Dihydroxy-16-ene-23-yne-26-trifluoro-cholecalciferol

To a stirred solution of 730 mg (1.25 mmol) of [3S-(1Z,3a,5b)]-[2-
10 [3,5-bis[[1,1-dimethylethyl]dimethylsilyl]oxy]-2-methylenecyclo-
hexylidene]ethyl]diphenylphosphine oxide in 7 ml of anhydrous
tetrahydrofuran at -78°C was added 0.758 ml (1.21 mmole) of 1.6M
n-butyl lithium in hexane dropwise under argon. After stirring for 5
min, to thus formed red solution was added dropwise over a 10 min
15 period a solution of 314 mg (0.784 mmole) of [3aR-[1(1R*),3aa,7ab]]-
3,3a,5,6,7,7a-hexahydro-7a-methyl-1-[6,6,6-trifluoro-1,5-dimethyl-5-
[(trimethyl-silyl)oxy]-3-hexynyl]-4H-inden -4-one (epimers) in 5 ml
anhydrous tetrahydrofuran. The reaction mixture was stirred at
-78°C for 1.5 hours, and then quenched by addition of 10 ml of a 1:1
20 mixture of 2N Rochelle salt and 2N potassium bicarbonate. After
warm-up to room temperature, an additional 30 ml of Rochelle salt -
potassium bicarbonate solution was added and extracted with 3 x 130
ml ethyl acetate. The organic layers were washed with brine, dried
over sodium sulfate and evaporated to dryness. The residue was
25 purified by FLASH chromatography on a silica gel column with
hexane-ethyl acetate 40:1, to give 475 mg of the trisilylated title
compound.

To the solution of 475 mg (0.621 mmole) of the trisilylated
30 intermediate in 7 ml of anhydrous tetrahydrofuran was added 4 ml (4
mmole) of a 1M tetrabutyl ammonium fluoride in tetrahydrofuran
under argon. The reaction mixture was stirred at room temperature
for 17 hrs, additional 2 ml of 1M tetrabutyl ammonium fluoride was
added and stirred for additional 5 hrs. It was then quenched with 5
35 ml of water, and after stirring for 20 min tetrahydrofuran was
removed by distillation. The residue was extracted with 3 x 120 ml

ethyl acetate. The organic layers were washed with water and brine, dried over sodium sulfate and evaporated to dryness. The crude product was purified by FLASH chromatography on a silica gel column with hexane-ethyl acetate 1:35 to give 270 mg (74.1%) of the title
5 compound as white foam; $[\alpha]_D^{26} + 26^\circ$ (c 0.2, EtOH); UV (EtOH); λ_{max} : 203nm (ϵ 17500), 263 (14120); $^1\text{H-NMR}$ (CDCl_3): δ 0.72 (s, 3H, CH_3), 1.13 (d, 3H, $J = 6.5$ Hz, CH_3), 1.58 (s, 3H, CH_3), 1.92 (ddd, ^1H , $J_{vic} = 3.5$ and 8.5 Hz, $J_{gem} = 13$ Hz, CH of CH_2), 2.21 (dd, ^1H , $J_{vic} = 12$ Hz, $J_{gem} = 14$ Hz, CH of CH_2), 2.61 (dd, ^1H , $J_{vic} = 3$ Hz, $J_{gem} = 13.5$ Hz, CH of CH_2),
10 2.82 (brm, ^1H , CH of CH_2), 4.24 (brm, ^1H , CH), 4.45 (brm, ^1H , CH), 5.02, 5.34 (2s, 2H, CH_2), 5.39 (brs, ^1H , CH), 6.11, 6.38 (AB, 2H, $J = 11.2$ Hz, 2CH).

Example 6

15 1,25(R,S)-Dihydroxy-16-ene-23-yne-26-trifluoro-19-nor-cholecalciferol

To a stirred solution of 630 mg (1.1 mmol) of [3R-(3a,5b,Z)-3,5-bis[[1,1-dimethylethyl)dimethylsilyl]oxy]cyclohexylidene]ethyl]
20 diphenylphosphine oxide in 7 ml of anhydrous tetrahydrofuran at -78°C was added dropwise 0.685 ml (1.1 mmol) of 1.6M n-butyl lithium in hexane under argon. After stirring for 5 min, to the thus formed red solution was added dropwise in the course of 5 min a
25 solution of 232 mg (0.579 mmol) of [3aR-[1(1R*),3aa,7ab]]-3,3a,5,6,7,7a-hexahydro-7a-methyl-1-[6,6,6-trifluoro-1,5-dimethyl-5-[(trimethylsilyl)oxy]-3-hexynyl]-4H-inden-4-one (epimers) in 2 ml anhydrous tetrahydrofuran. The reaction mixture was stirred at -78°C for 1.75 hours. It was then quenched by addition of 10 ml of a
30 1:1 mixture of 2N potassium bicarbonate and 2N Rochelle salt, warmed up to room temperature, and addition of 30 ml of potassium bicarbonate - Rochelle salt mixture, extracted with 3 x 100 ml ethyl acetate. The combined extracts were washed with brine, dried over sodium sulfate and evaporated to dryness. The crude product was
35 purified by FLASH chromatography on a silica gel column with

hexane-ethyl acetate, to give 145 mg of the trisilylated title compound.

To the stirred solution of 145 mg (0.19 mmol) of the trisilylated intermediate in 2.5 ml anhydrous tetrahydrofuran was added under argon 3 ml (3 mmol) of 1M tetrabutyl ammonium fluoride in tetrahydrofuran, and the reaction mixture was stirred at room temperature for 65 hours. It was then quenched with 5 ml water, stirred 15 min., 20 ml of brine was added and extracted with 3 x 90 ml of ethyl acetate. The combined extracts were washed with a mixture of water and brine, dried over sodium sulfate and evaporated to dryness. The crude product was purified by FLASH chromatography on silica gel to give 89 mg (33.9%) of the title compound as white foam; $[\alpha]_D^{25} + 76^\circ$ (c 0.2, EtOH); UV (EtOH); λ_{max} : 242-243 nm (e 27,100), 250-251 (31,800), 260 (21,300); $^1\text{H-NMR}$: (CDCl_3): δ 0.72 (s, 3H, CH_3), 1.13 (d, 3H, $J = 6.3$ Hz, CH_3), 1.59 (s, 3H, CH_3), 1.97 (m, 1H), 2.07 (m, 1H, CH of CH_2), 2.15-2.55 (m, 8H), 2.72-2.84 (m, 2H, 2 CH of CH_2), 4.06 (brm, 1H, CH), 4.13 (brm, 1H, CH), 5.42 (brs, 1H, CH), 5.96, 6.31 (AB, 2H, $J = 11.1$ Hz, CH CH).

Example 7

1 α -fluoro-25(R,S)-hydroxy-16-ene-23-yne-26-trifluoro-cholecalciferol

To a solution of 536 mg (1.14 mmol) of [3S-(3a,5b,Z)-2-[2-methylene-3-fluoro-5-[[1,1-dimethylethyl]dimethylsilyl]oxy]cyclohexylidene]ethyl] diphenyl phosphine oxide in 6 ml of anhydrous tetrahydrofuran at -78°C was added 0.71 ml (1.14 mmol) of 1.6M n-butyl lithium in hexane, dropwise, under argon. After stirring for 5 min, to the red solution was added 282 mg (0.704 mmol) of [3aR-[1(1R*),3aa,7ab]]-3,3a,5,6,7,7a-hexahydro-7a-methyl-1-[6,6,6-trifluoro-1,5-di-methyl-5 [(trimethylsilyl) oxy] -3-hexynyl]-4H-inden-4-one (epimers) in 4.5 ml anhydrous tetrahydrofuran dropwise over 10 min.. The reaction mixture was stirred at -78°C for 2.5 hours. It was then quenched with 10 ml of a 2N Rochelle salt and warmed up

to room temperature. It was further diluted with 25 ml of 2N Rochelle salt, and then extracted with 3 x 100 ml ethyl acetate. The combined organic layers were washed with brine, dried over sodium sulfate and evaporated to dryness. The residue was purified by
5 FLASH chromatography on silica gel with hexane-ethyl acetate 30:1 and then with 1:4. It gave 380 mg of the disilylated title compound.

To a solution of 380 mg (0.582 mmol) of the disilyl intermediate in 4 ml of anhydrous tetrahydrofuran was added under argon 4 ml (4
10 mmol) of 1M tetrabutyl ammonium fluoride in tetrahydrofuran. The reaction mixture was stirred at room temperature for 17 hours. It was then quenched with 5 ml water, stirred for 15 minutes, 20 ml of brine was added, and extracted with 3 x 90 ml ethyl acetate. The combined organic layers were washed with water and brine, dried
15 over sodium sulfate and evaporated to dryness. The crude product was first purified by FLASH chromatography on silica gel with hexane-ethyl acetate 2:1 and then by HPLC on a YMC-50 mm x 50 cm silica column with hexane-ethyl acetate 3:2 to give 233 mg (70.9%) of the title compound as white foam; $[\alpha]_D^{25} +22.5^\circ$ (c 0.2, EtOH); UV (EtOH):
20 λ_{max} : 242-243 nm (e 11,400), 268-269 (11,050); $^1\text{H-NMR}$ (CDCl_3): δ 0.71 (s, 3H, CH_3), 1.16 (d, 3H, $J = 6.5\text{ Hz}$, CH_3), 1.43-1.58 (m, 2H), 1.60 (s, 3H, CH_3), 1.60 - 1.88 (m, 4H), 2.02 (m, 2H, CH_2), 2.14-2.47 (m, 8H), 2.64 (dd, $J_{vic} = 3.8\text{ Hz}$, $J_{gem} = 13.5\text{ Hz}$, CH of CH_2), 2.82 (m, 1H, CH of CH_2), 4.24 (m, 1H, CH), 5.12 (s, 1H, CH of CH_2), 5.15 (ddd, 1H, $J = 3.9$,
25 6.6 Hz, $J_{HF} = 49.5\text{ Hz}$, CH), 5.41 (s, 2H, CH and CH of CH_2), 6.12, 6.40 (AB, 2H, $J = 11.3\text{ Hz}$, CH CH).

Example 8

30 25(R,S)-Hydroxy-16-ene-23-yne-26-trifluoro-cholecalciferol

To a solution of 520 mg (1.15 mmol) of [5S,Z)-2-[2-[2-methylene-5-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]cyclohexylidene]ethyl] diphenyl phosphine oxide in 6 ml of anhydrous tetrahydrofuran at
35 -78°C was added dropwise under argon 0.715 ml (1.14 mmol) of 1.6M butyl lithium in hexane. After stirring for 5 min, to the red solution

thus formed was added dropwise over 10 min 287 mg (0.716 mmol) of [3aR-[1(1R*),3aa,7ab]]-3,3a,5,6,7,7a-hexahydro-7a-methyl-1-[6,6,6-trifluoro-1,5-dimethyl-5-[(trimethylsilyl)oxy]-3-hexynyl]-4H-inden-4-one (epimers) in 4 ml anhydrous tetrahydro-furan. The reaction mixture was then stirred at -78°C for 2 hours. It was quenched with 10 ml of a 2N Rochelle salt and warmed up to room temperature. The resulting mixture was diluted with 25 ml 2N Rochelle salt and extracted with 3 x 100 ml ethyl acetate. The combined organic layers were washed with brine, dried over sodium sulfate and evaporated to dryness. The residue was purified by FLASH chromatography on silica gel with hexane-ethyl acetate 30:1, to give 364 mg of disilylated title compound.

To a solution of 364 mg (0.573 mmol) of the disilyl intermediate in 4 ml of anhydrous tetrahydrofuran was added 4 ml (4 mmol) of 1M tetrabutyl ammonium fluoride in tetrahydrofuran under argon. The reaction mixture was stirred at room temperature for 18 hours. It was then quenched with 10 ml water, stirred for 10 min, 20 ml of brine was added and extracted with 3 x 90 ml ethyl acetate. The organic layers were combined and washed with water and brine, dried over sodium sulfate and evaporated to dryness. The crude product was purified by FLASH chromatography on silica gel with hexane-ethyl acetate 2:1, and then by HPLC on a YMC 50 mm x 50 cm silica column with hexane-ethyl acetate 3:2, to give 238 mg (74%) of the title compound as white foam; $[\alpha]_D^{25} + 47^\circ$ (c 0.2, EtOH); UV (EtOH): λ_{max} 206 nm (e 16,000), 263 (15,500); $^1\text{H-NMR}$ (CDCl_3): δ 0.71 (s, 3H, CH_3), 1.13 (d, 3H, $J = 6.4$ Hz, CH_3), 1.58 (s, 3H, CH_3), 1.93 (m, 1H), 2.03 (ddd, 1H, $J_{vic} = 3$ and 6 Hz, $J_{gem} = 14$ Hz, CH of CH_2), 2.12 - 2.52 (m, 9H), 2.58 (dd, 1H, $J_{vic} = 3.5$ Hz, $J_{gem} = 13$ Hz, CH of CH_2), 2.82 (m, 1H, CH of CH_2), 3.96 (brm, 1H, CH), 4.83 (d, 1H, $J = 1.7$ Hz, CH of CH_2), 5.06 (s, 1H, CH of CH_2), 5.39 (brs, 1H, CH), 6.12, 6.23 (AB, 2H, $J = 11.3$ Hz, CH CH).

The following pharmaceutical compositions can be prepared in a manner known per se:

EXAMPLE A

Soft Gelatin Capsule

5

Item	Ingredients	mg/Capsule
1	Compound of formula I	0.0001-1.0
2	Butylated Hydroxytoluene (BHT)	0.016
3	Butylated Hydroxyanisole (BHA)	0.016
4	Miglyol 812 q.s.	160.0

EXAMPLE B

Soft Gelatin Capsule

10

Item	Ingredients	mg/Capsule
1	Compound of formula I	0.0001-1.0
2	α -Tocopherol	0.016
3	Miglyol 812 q.s.	160.0

EXAMPLE C

Topical Cream

15


	mg/gm
Compound of formula I	0.0005-0.10
Cetyl Alcohol	1.5
Stearyl Alcohol	2.5
Span 60 (Sorbitan monostearate)	2.0
Arlacel 165 (Glyceryl monostearate and polyoxyethylene glycol stearate blend)	4.0
Tween 60 (polysorbate 60)	1.0
Mineral Oil	4.0
Propylene Glycol	5.0
Propylparaben	0.05
BHA	0.05

Sorbitol Solution	2.0
Edetate Disodium	0.01
Methylparaben	0.18
Distilled Water	q.s. to 100 gm



Claims


1. A compound of the formula



I

wherein R is hydrogen, fluorine or hydroxy, and X is =CH₂ or when R is hydroxy, X is hydrogen or =CH₂.

2. The compound of Claim 1, wherein R is hydroxy.
3. A mixture comprising epimers of the formula



Ia



wherein R is hydrogen, fluorine or hydroxy, and X is $=CH_2$ or when R is hydroxy, X is hydrogen or $=CH_2$.

4. 1,25(R,S)-Dihydroxy-16-ene-23-yne-26-trifluoro-
5 cholecalciferol.

5. 1,25(R,S)-Dihydroxy-16-ene-23-yne-26-trifluoro-19-nor-
cholecalciferol.

10 6. 1α -fluoro-25(R,S)-hydroxy-16-ene-23-yne-26-trifluoro-
cholecalciferol.

7. 25(R,S)-Hydroxy-16-ene-23-yne-26-trifluoro-
cholecalciferol.

15 8. A pharmaceutical composition comprising an effective
amount of the compound of formula I as in claim 1, and an inert
carrier.

20 9. Use of the compounds of formula I as in claim 1, for the
manufacture of a pharmaceutical composition for the treatment of
hyperproliferative disorders, neoplastic diseases and sebaceous gland
diseases.
