




FIG.1




FIG.2




FIG.3A




FIG. 3B




FIG. 3C

ID:8619 364 310
 [Example 1] let one primitive function of $f(x)$ is: $F(x) = \ln^2(x + \sqrt{x^2 + 1})$, solve $I = \int xf''(x) dx$
 Solution $I = \int xdf(x) = xf(x) - \int f(x) dx = xF'(x) - F(x) + C$
 $= \frac{2x}{\sqrt{x^2 + 1}} \ln(x + \sqrt{x^2 + 1}) - \ln^2(x + \sqrt{x^2 + 1}) + C$
 [Example 2] Let $F'(x) = f(x)$, if $x \geq 0$, $f(x)F'(x) = \frac{x^2}{2(1+x)^3}$, and $F(0) = 1$,
 $F(x) > 0$, solve $f(x) | x \geq 0$.
 Solution $2 \int f(x)F'(x) dx = 2 \int F'(x) dF(x) = F^2(x) + C_1$
 teacher 788
 Floor control mode
 1. User A
 2. User B
 3. User C
 DiscussionQuestion(0)
 User A 11:32:05
 I think the teacher's lecture is good, and easy to understand.
 User D 11:34:19
 Yes, it is good.
 Classroom list
 pressF2talk
 Recording disabled
 My app My course

FIG. 3D




FIG. 4




FIG. 5




FIG. 6




FIG. 7

5/5




FIG. 8




FIG. 9