

DESCRIPTION

PYRAZOLE COMPOUND AND PHARMACEUTICAL USE THEREOF

5 TECHNICAL FIELD

[0001] The present invention relates to a pyrazole compound having SGLT1 inhibitory activity or a pharmaceutically acceptable salt thereof, a pharmaceutical composition comprising the same, and its pharmaceutical use.

10 BACKGROUND ART

[0002] It is known that SGLT1 (i.e., Sodium-Glucose Co-transporter 1) contributes to a great portion of absorption of glucose and galactose in the small intestine. It is reported that human SGLT1-deficiency in patients causes glucose-galactose malabsorption. Furthermore, it is confirmed that the expression of SGLT1 in the small intestine increases in diabetic patients and it is thought that
15 increased sugar absorption in diabetic patients is caused by the high expression of SGLT1 in the small intestine.

[0003] From these knowledge, a SGLT1 inhibitor is expected to normalize the blood glucose level, since it blocks glucose absorption in the small intestine. Therefore, a SGLT1 inhibitor is considered to be effective against diabetes and diabetic complications associated with hyperglycemia, specifically
20 retinopathy, nephropathy and neuropathy which are known as microangiopathy, and cerebrovascular disease, ischemic heart disease and membrum-inferius arteriosclerosis obliterans which are known as macroangiopathy. Moreover, it is thought to be effective against obesity by inhibiting the inflow of glucose into the body (non-patent literatures 1 and 2).

[0004] In addition, SGLT1 is expressed in cardiac muscle cells. It is known that GLUT1 and
25 GLUT4 (Glucose Transporter Type 4) usually have a role in uptake of glucose to cardiac muscle cells and the contribution of SGLT1 is small. However, the expression of SGLT1 is induced in the cardiac muscles of mice into which was introduced mutated genes of PRKAG2 (gamma 2 subunit of AMPK (AMP-Activated Protein Kinase)) which is a responsible gene of familial hypertrophic cardiomyopathy (glycogen accumulation-type myocardosis), or mice which underwent myocardial
30 ischemia treatment, and SGLT1 is reported to have contributed to the uptake of glucose to cardiac muscle cells in these pathologies. Glucose incorporated by SGLT1 is thought to be excessively accumulated or metabolized within cardiac muscle cells and impaire the cells. It is reported in the former mouse model that accumulation of glycogen in the cardiac muscle is actually inhibited by the treatment of phlorizin which is a non-selective SGLT inhibitor.

35 From these knowledge, a SGLT1 inhibitor is thought to be effective against hypertrophic cardiomyopathy and ischemic heart disease by inhibiting uptake of excess glucose into cardiac muscle cells (non-patent literatures 3 and 4).

[0005] SGLT1 is stabilized by epidermal growth factor receptors (i.e., surface proteins on many kinds of cancer cells) in cancer cells. It is known that transporters of glucose, lactic acid, and amino
40 acid, etc. are involved in nutrition supply to cancer cells, and especially, regarding the transportation of glucose, SGLT1 and GLUT1 supply glucose to cancer cells, continuously. When glucose is not supplied over a long period of time, cells are destroyed by autophagy.

From these knowledge, a SGLT1 inhibitor is thought to inhibit supply of glucose to cancer cells and show anticancer activity (non-patent literatures 5 and 6).

[0006] Since carbohydrate in diet is degraded to monosaccharide in the gastrointestinal tracts and is absorbed in the upper gastrointestinal tracts, not so many sugar would reach the lower gastrointestinal tracts. However, when drugs which delay and/or inhibit glucose absorption are administered, or a large amount of resistant polysaccharide are ingested, undigested sugar would be retained in the lower gastrointestinal tracts and the undigested sugar retained in the lower gastrointestinal tracts would cause osmotic diarrhea.

The amount of monosaccharide in the lower gastrointestinal tracts is increased by the inhibition of glucose absorption by a SGLT1 inhibitor. Therefore, it is believed that a SGLT1 inhibitor is effective against constipation.

[Non-patent document]

[0007]

[Non-patent literature 1] Am J Physiol Gastrointest Liver Physiol. 2002; 282(2):G241-8

[Non-patent literature 2] Nature. 1991; 350(6316): 354-6

[Non-patent literature 3] J Mol Cell Cardiol. 2010; 49(4):683-92

[Non-patent literature 4] Cardiovasc Res. 2009; 84(1):111-8

[Non-patent literature 5] Cancer Cell. 2008, 13: 385-93

[Non-patent literature 6] Pharmacol Ther. 2009, 121: 29-40

SUMMARY OF INVENTION


PROBLEMS TO BE SOLVED BY THE INVENTION

[0008] An object of the present invention is to provide a pyrazole compound useful as a medicine which has SGLT1 inhibitory activity, or a pharmaceutically acceptable salt thereof, and a pharmaceutical composition comprising the same.

MEANS TO SOLVE THE PROBLEMS

[0009] The present invention is as follows.

Paragraph 1: A compound or a pharmaceutically acceptable salt thereof represented by the following general formula [I]:



, wherein

Ring Cy is

(1) C₆₋₁₀ aryl,

(2) C₃₋₈ cycloalkyl or

(3) C₃₋₈ cycloalkenyl,

n₁ is 0, 1, 2, 3 or 4,

R¹ is

- (1) a halogen atom,
 (2) hydroxy,
 (3) carboxy,
 (4) a C₁₋₆ alkyl group,
 5 (5) a C₁₋₆ alkoxy group,
 (6) a C₃₋₆ cycloalkyl group,
 (7) a halo C₁₋₆ alkyl group,
 (8) a hydroxy C₁₋₆ alkyl group,
 (9) a C₁₋₆ alkoxy C₁₋₆ alkyl group,
 10 (10) a halo C₁₋₆ alkoxy group or
 (11) a carboxy C₁₋₆ alkoxy group,
 , provided when n₁ is 2, 3 or 4, R¹ each is the same or different,
 R² is
 (1) a C₁₋₈ alkyl group,
 15 (2) a C₃₋₈ cycloalkyl C₁₋₆ alkyl group,
 (3) a C₆₋₁₀ aryl C₁₋₆ alkyl group,
 (4) a saturated heterocyclo C₁₋₆ alkyl group (the saturated heterocycle is of 5-membered to 6-membered
 saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur),
 (5) a C₃₋₈ cycloalkyl group,
 20 (6) a halo C₁₋₆ alkyl group,
 (7) a C₁₋₆ alkoxy C₁₋₆ alkyl group,
 (8) a halo C₃₋₈ cycloalkyl group,
 (9) a C₆₋₁₀ aryl group (the C₆₋₁₀ aryl group is optionally substituted by 1 to 4 substituents selected from
 (a) a halogen atom,
 25 (b) a C₁₋₆ alkyl group,
 (c) a C₁₋₆ alkoxy group and
 (d) a carboxy C₁₋₆ alkoxy group), or
 (10) a saturated heterocycle group (the saturated heterocycle is of 5-membered to 6-membered saturated
 heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur, and is optionally
 30 substituted by 1 to 4 C₁₋₆ alkyl groups),
 n₂ is 0, 1, 2, 3 or 4,
 m is 0, 1, 2, 3 or 4,
 R³ is
 (1) a C₁₋₆ alkyl group or
 35 (2) a hydroxy C₁₋₆ alkyl group,
 , provided when n₂ is 2, 3 or 4, R³ each is the same or different,
 X^a is
 (1) a bond or
 (2) N-R⁴ (the R⁴ is
 40 (a) a hydrogen atom or
 (b) a C₁₋₆ alkyl group), and
 X^b is
 (1) NH or
 (2) an oxygen atom,

provided, when X^a is $N-R^4$, X^b is not an oxygen atom.

[0010] Paragraph 2: The compound or a pharmaceutically acceptable salt thereof according to paragraph 1, wherein X^b is NH.

5 [0011] Paragraph 3: The compound or a pharmaceutically acceptable salt thereof according to paragraph 1 or 2, wherein the ring Cy is C_{6-10} aryl.

[0012] Paragraph 4: The compound or a pharmaceutically acceptable salt thereof according to paragraph 3, wherein the ring Cy is phenyl.

[0013] Paragraph 5: The compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 1 to 4, wherein R^1 is

10 (1) a halogen atom,

(2) a C_{1-6} alkyl group,

(3) a C_{1-6} alkoxy group,

(4) a halo C_{1-6} alkyl group,

(5) a C_{1-6} alkoxy C_{1-6} alkyl group or

15 (6) a halo C_{1-6} alkoxy group,

, provided when $n1$ is 2, 3 or 4, R^1 each is the same or different.

[0014] Paragraph 6: The compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 1 to 5

20 , wherein R^2 is a C_{6-10} aryl group (the C_{6-10} aryl group is optionally substituted by 1 to 4 substituents selected from

(a) a halogen atom,

(b) a C_{1-6} alkyl group,

(c) a C_{1-6} alkoxy group and

(d) a carboxy C_{1-6} alkoxy group).

25 [0015]

Paragraph 7: The compound or a pharmaceutically acceptable salt thereof according to paragraph 6

, wherein R^2 is a phenyl group (the phenyl group is optionally substituted by 1 to 4 substituents selected from

(a) a halogen atom,

30 (b) a C_{1-6} alkyl group,

(c) a C_{1-6} alkoxy group and

(d) a carboxy C_{1-6} alkoxy group).

[0016] Paragraph 8: A pharmaceutical composition which comprises a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 1 to 7, and a pharmaceutically acceptable carrier.

35 [0017] Paragraph 9: A SGLT1 inhibitor which comprises a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 1 to 7

[0018] Paragraph 10: An agent for treating or preventing diabetes which comprises a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 1 to 7.

40 [0019] Paragraph 11: The agent for treating or preventing diabetes according to paragraph 10, wherein diabetes is type II diabetes mellitus.

[0020] Paragraph 12: A method for inhibiting of SGLT1 which comprises administering to mammals a therapeutically effective amount of a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 1 to 7

[0021] Paragraph 13: A method for treating or preventing diabetes which comprises administering to mammals a therapeutically effective amount of a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 1 to 7.

[0022] Paragraph 14: The method according to paragraph 13, wherein the diabetes is type II diabetes mellitus.

[0023] Paragraph 15: Use of a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 1 to 7 for manufacturing a SGLT1 inhibitor.

[0024] Paragraph 16: The use of a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 1 to 7, for manufacturing an agent for treating or preventing diabetes.

[0025] Paragraph 17: The use according to paragraph 16, wherein the diabetes is type II diabetes mellitus.

[0026] Paragraph 18: A kit for commerce comprising

(a) a pharmaceutical composition which comprises a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 1 to 7 as an active ingredient, and


(b) a package insert about the pharmaceutical composition which indicates that the pharmaceutical composition can be used or should be used for treatment or prevention of type II diabetes mellitus.

[0027] Paragraph 19: A package for commerce comprising

(a) a pharmaceutical composition which comprises a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 1 to 7 as an active ingredient, and

(b) a package insert about the pharmaceutical composition which indicates that the pharmaceutical composition can be used or should be used for treatment or prevention of type II diabetes mellitus.

[0028] Paragraph 20: A compound or a pharmaceutically acceptable salt thereof represented by the general formula [Ib]:



[Ib]

, wherein

Ring Cy is

(1) C_{6-10} aryl,

(2) C_{3-8} cycloalkyl or

(3) C_{3-8} cycloalkenyl,

$n1$ is 0, 1, 2, 3 or 4,


R^{1a} is

(1) a halogen atom,

(2) hydroxy,

(3) carboxy,

- (4) a C₁₋₆ alkyl group,
 (5) a C₂₋₈ alkenyl group,
 (6) a C₂₋₈ alkynyl group,
 (7) a C₁₋₆ alkoxy group,
 5 (8) a C₃₋₆ cycloalkyl group,
 (9) a halo C₁₋₆ alkyl group,
 (10) a hydroxy C₁₋₆ alkyl group,
 (11) a C₁₋₆ alkoxy C₁₋₆ alkyl group,
 (12) a halo C₁₋₆ alkoxy C₁₋₆ alkyl group,
 10 (13) a C₁₋₆ alkylsulfonyl C₁₋₆ alkyl group,
 (14) a halo C₁₋₆ alkylsulfonyl C₁₋₆ alkyl group,
 (15) a halo C₁₋₆ alkylamino C₁₋₆ alkyl group,
 (16) a C₃₋₆ cycloalkyl C₂₋₆ alkynyl group
 (17) a halo C₁₋₆ alkoxy group,
 15 (18) a carboxy C₁₋₆ alkoxy group,
 (19) a C₁₋₆ alkylsulfanyl group,
 (20) a C₁₋₆ alkylsulfonyl group,
 (21) a halo C₁₋₆ alkylsulfonyl group,
 (22) a C₁₋₈ alkylcarbonyl group,
 20 (23) a C₁₋₆ alkyloxycarbonyl group,
 (24) a group represented by the Formula:



, wherein

R₅ is

- 25 (a) a hydrogen atom or
 (b) a C₁₋₆ alkyl group, and

R₆ is

- (a) a C₁₋₆ alkyl group or
 (b) a halo C₁₋₆ alkyl group,
 30 (25) a saturated heterocyclo C₁₋₆ alkyl group (the saturated heterocycle is of 4-membered to 6-membered saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur) or
 (26) a saturated heterocyclo oxy C₁₋₆ alkyl group (the saturated heterocycle is of 4-membered to 6-membered saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur),
 35 , provided when n1 is 2, 3 or 4, R^{1a} each is the same or different,

R^{2a} is

- (1) a C₁₋₈ alkyl group,
 (2) a C₃₋₈ cycloalkyl C₁₋₆ alkyl group,
 (3) a C₆₋₁₀ aryl C₁₋₆ alkyl group,
 40 (4) a saturated heterocyclo C₁₋₆ alkyl group (the saturated heterocycle is of 4-membered to 6-membered

saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur),

(5) a C₃₋₈ cycloalkyl group,

(6) a halo C₁₋₆ alkyl group,

(7) a C₁₋₆ alkoxy C₁₋₆ alkyl group,

5 (8) a halo C₃₋₈ cycloalkyl group,

(9) a C₆₋₁₀ aryl group (the C₆₋₁₀ aryl group is optionally substituted by 1 to 4 substituents selected from

(a) a halogen atom,

(b) hydroxy,

(c) a C₁₋₆ alkyl group,

10 (d) a C₂₋₈ alkenyl group,

(e) a C₂₋₆ alkynyl group,

(f) a C₁₋₆ alkoxy group,

(g) a halo C₁₋₆ alkyl group,

(h) a C₁₋₆ alkoxy C₁₋₆ alkyl group,

15 (i) a halo C₁₋₆ alkoxy group,

(j) a hydroxy C₁₋₆ alkoxy group,

(k) a C₁₋₆ alkoxy C₁₋₆ alkoxy group,

(l) a carboxy C₁₋₆ alkoxy group,

(m) a C₆₋₁₀ aryl C₁₋₆ alkoxy group,

20 (n) a C₁₋₆ alkylsulfanyl C₁₋₆ alkoxy group,

(o) a C₁₋₆ alkylsulfonyl C₁₋₆ alkoxy group,

(p) a C₆₋₁₀ aryl C₁₋₆ alkoxy C₁₋₆ alkoxy group,

(q) a saturated heterocyclo C₁₋₆ alkoxy group (the saturated heterocycle is of 4-membered to 6-membered saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur, and is optionally substituted by 1 to 2 C₁₋₆ alkyl groups),

25 (r) a saturated heterocyclo oxy group (the saturated heterocycle is a 4-membered to 6-membered saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur),

(s) a C₃₋₈ cycloalkyloxy group,


(t) a C₁₋₆ alkoxycarbonyloxy group and

30 (u) a C₁₋₆ alkylsulfonyl group), or

(10) a saturated heterocycle group (the saturated heterocycle is of 4-membered to 6-membered saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur, and is optionally substituted by 1 to 4 C₁₋₆ alkyl groups), and

Ring Cy^a is a group selected from

35 (1) a group represented by the Formula:




, wherein

m2 is 1, 2 or 3,

m3 is 1, 2 or 3,

n_3 is 1 or 2,

(2) a group represented by the Formula:




5 , wherein

m_4 is 0, 1, 2 or 3,

m_5 is 0, 1 or 2,

, provided sum of m_4 and m_5 is 1 or more,

10 (3) a group represented by the Formula:




, wherein

m_4 is 0, 1, 2 or 3,

m_5 is 0, 1 or 2,

15 , provided sum of m_4 and m_5 is 1 or more,

(4) a group represented by the Formula:




, wherein

20 m_6 is 0, 1 or 2,

m_7 is 0, 1 or 2,

, provided sum of m_6 and m_7 is 1 or more,

(5) a group represented by the Formula:




, wherein

m6 is 0, 1 or 2,

m7 is 0, 1 or 2,

5 , provided sum of m6 and m7 is 1 or more,

(6) a group represented by the Formula:




, wherein

10 m8 is 1 or 2,

m9 is 1 or 2, and

(7) a group represented by the Formula:



15 , wherein

m8 is 1 or 2,

m9 is 1 or 2,

n2 is 0, 1, 2, 3 or 4,

20 R^{3a} is

(1) hydroxy,

(2) a C₁₋₆ alkyl group or

(3) a hydroxy C₁₋₆ alkyl group,

, provided when n_2 is 2, 3 or 4, R^{3a} each is the same or different,

R^{3b} is

(1) hydroxy,

5 (2) a C_{1-6} alkyl group (which optionally form a C_{3-6} cycloalkyl group together with the carbon to which it is attached and the carbon adjacent thereto) or

(3) a hydroxy C_{1-6} alkyl group, or

(4) when two R^{3b} are attached to the same carbon, they optionally form a C_{3-6} cycloalkyl group together with the carbon to which they are attached,

10 provided, when n_2 is 2, 3 or 4, each R^{3b} is the same or different, and

R^{4b} is

(1) a hydrogen atom,

(2) a C_{1-6} alkyl group,

15 (3) a carboxy C_{1-6} alkyl group,


(4) a halo C_{1-6} alkyl group or

(5) a C_{1-6} alkoxy C_{1-6} alkyl group.

[0029] Paragraph 21: The compound or a pharmaceutically acceptable salt thereof according to paragraph 20, wherein

Ring Cy^a is selected from

(1) a group represented by the Formula:




, wherein

25 m_4 is 0, 1, 2 or 3,

m_5 is 0, 1 or 2,

, provided sum of m_4 and m_5 is 1 or more,

(2) a group represented by the Formula:



30


, wherein

m_6 is 0, 1 or 2,

m_7 is 0, 1 or 2,

, provided sum of m_6 and m_7 is 1 or more,


(3) a group represented by the Formula:



, wherein

- 5 m6 is 0, 1 or 2,
 m7 is 0, 1 or 2,
 , provided sum of m6 and m7 is 1 or more,


(4) a group represented by the Formula:



, wherein

- 10 m8 is 1 or 2,
 m9 is 1 or 2, and

(5) a group represented by the Formula:



, wherein

- 20 m8 is 1 or 2,
 m9 is 1 or 2,
 n2 is 0, 1, 2, 3 or 4,
 R^{3a} is
 (1) hydroxy,
 (2) a C₁₋₆ alkyl group or
 (3) a hydroxy C₁₋₆ alkyl group,
 , provided when n2 is 2, 3 or 4, each R^{3a} is the same or different,

R^{3b} is

- (1) hydroxy,

(2) a C₁₋₆ alkyl group (which optionally form a C₃₋₆ cycloalkyl group together with the carbon to which it is attached and the carbon adjacent thereto) or

(3) a hydroxy C₁₋₆ alkyl group, or

(4) when two R^{3b} are attached to the same carbon, they optionally form a C₃₋₆ cycloalkyl group together with the carbon to which they are attached,

provided, when n₂ is 2, 3 or 4, each R^{3b} is the same or different, and

R^{4b} is

(1) a hydrogen atom,

(2) a C₁₋₆ alkyl group,

(3) a carboxy C₁₋₆ alkyl group,


(4) a halo C₁₋₆ alkyl group or

(5) a C₁₋₆ alkoxy C₁₋₆ alkyl group.

[0030] Paragraph 22: The compound or a pharmaceutically acceptable salt thereof according to paragraph 20, wherein

Ring Cy^a is selected from

(1) a group represented by the Formula:




, wherein,

m₄ is 0, 1, 2 or 3,

m₅ is 0, 1 or 2,

, provided sum of m₄ and m₅ is 1 or more, and

(2) a group represented by the Formula:



, wherein

m₆ is 0, 1 or 2,

m₇ is 0, 1 or 2,

, provided sum of m₆ and m₇ is 1 or more,

n₂ is 0, 1, 2, 3 or 4,

R^{3a} is

(1) hydroxy,

(2) a C₁₋₆ alkyl group or

(3) a hydroxy C₁₋₆ alkyl group,

, provided, when n₂ is 2, 3 or 4, each R^{3a} is the same or different,

R^{3b} is

- (1) hydroxy,
 - (2) a C_{1-6} alkyl group (which optionally form a C_{3-6} cycloalkyl group together with the carbon to which it is attached and the carbon adjacent thereto) or
 - (3) a hydroxy C_{1-6} alkyl group, or
 - (4) when two R^{3b} are attached to the same carbon, they optionally form a C_{3-6} cycloalkyl group together with the carbon to which they are attached,
- provided, when n_2 is 2, 3 or 4, each R^{3b} is the same or different, and


R^{4b} is

- (1) a hydrogen atom,
- (2) a C_{1-6} alkyl group,
- (3) a carboxy C_{1-6} alkyl group,
- (4) a halo C_{1-6} alkyl group or
- (5) a C_{1-6} alkoxy C_{1-6} alkyl group.

[0031] Paragraph 23: The compound or a pharmaceutically acceptable salt thereof according to paragraph 20, wherein

Ring Cy^a is selected from


- (1) a group represented by the Formula:



, wherein

n_2 is 0, 1, 2, 3 or 4, and

- (2) a group represented by the Formula:



, wherein

n_2 is 0 or 1,

R^{3a} is

- (1) hydroxy,
- (2) a C_{1-6} alkyl group or
- (3) a hydroxy C_{1-6} alkyl group, and

R^{3b} is

- (1) hydroxy,
 - (2) a C₁₋₆ alkyl group (which optionally form a C₃₋₆ cycloalkyl group together with the carbon to which it is attached and the carbon adjacent thereto) or
 - (3) a hydroxy C₁₋₆ alkyl group, or
 - 5 (4) when two R^{3b} are attached to the same carbon, they optionally form a C₃₋₆ cycloalkyl group together with the carbon to which they are attached,
- provided, when n₂ is 2, 3 or 4, each R^{3b} is the same or different, and

R^{4b} is

- (1) a hydrogen atom,
- (2) a C₁₋₆ alkyl group,
- (3) a carboxy C₁₋₆ alkyl group,
- (4) a halo C₁₋₆ alkyl group or
- (5) a C₁₋₆ alkoxy C₁₋₆ alkyl group.

[0032] Paragraph 24: The compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 20 to 23, wherein the ring Cy is C₆₋₁₀ aryl.

[0033] Paragraph 25: The compound or a pharmaceutically acceptable salt thereof according to paragraph 24, wherein the ring Cy is phenyl.

[0034] Paragraph 26: The compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 20 to 25 wherein R^{1a} is

- (1) a halogen atom,
 - (2) a C₁₋₆ alkyl group,
 - (3) a C₁₋₆ alkoxy group,
 - 25 (4) a halo C₁₋₆ alkyl group,
 - (5) a C₁₋₆ alkoxy C₁₋₆ alkyl group,
 - (6) a halo C₁₋₆ alkoxy C₁₋₆ alkyl group or
 - (7) a halo C₁₋₆ alkoxy group,
- , provided when n₁ is 2, 3, or 4, R^{1a} each is the same or different.

[0035] Paragraph 27: The compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 20 to 26
, wherein R^{2a} is a C₆₋₁₀ aryl group (the C₆₋₁₀ aryl group is optionally substituted by 1 to 4 substituents selected from

- (a) a halogen atom,
- 35 (b) a C₁₋₆ alkyl group,
- (c) a C₁₋₆ alkoxy group and
- (d) a carboxy C₁₋₆ alkoxy group).





[0036] Paragraph 28: The compound or a pharmaceutically acceptable salt thereof according to paragraph 27





, wherein R^{2a} is a phenyl group (the phenyl group is optionally substituted by 1 to 4 substituents selected from





- (a) a halogen atom,
- (b) a C₁₋₆ alkyl group,
- (c) a C₁₋₆ alkoxy group and
- 45 (d) a carboxy C₁₋₆ alkoxy group).





[0037] Paragraph 29: The compound or pharmaceutically acceptable salt thereof of paragraph 20,

wherein the compound is selected from the following Formulae:










and



[0038] Paragraph 30: A pharmaceutical composition which comprises a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 20 to 29 and a pharmaceutically acceptable carrier.

[0039] Paragraph 31: A SGLT1 inhibitor which comprises a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 20 to 29.

[0040] Paragraph 32: An agent for treating or preventing diabetes which comprises a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 20 to 29.

[0041] Paragraph 33: The agent for treating or preventing diabetes according to paragraph 32, wherein diabetes is type II diabetes mellitus.

[0042] Paragraph 34: A method for inhibiting SGLT1 which comprises administering to mammals a therapeutically effective amount of a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 20 to 29.

[0043] Paragraph 35: A method for treating or preventing diabetes which comprises administering to mammals a therapeutically effective amount of a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 20 to 29.

[0044] Paragraph 36: The method according to paragraph 35, wherein the diabetes is type II diabetes mellitus.

[0045] Paragraph 37: Use of a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 20 to 29 for manufacturing a SGLT1 inhibitor.

[0046] Paragraph 38: Use of a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 20 to 29 for manufacturing an agent for treating or preventing diabetes.

[0047] Paragraph 39: The use according to paragraph 38 wherein the diabetes is type II diabetes mellitus.

[0048] Paragraph 40: A kit for commerce comprising

(a) a pharmaceutical composition which comprises a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 20 to 29 as an active ingredient, and

(b) a package insert about the pharmaceutical composition which indicates that the pharmaceutical composition can be used or should be used for treatment or prevention of type II diabetes mellitus.

[0049] Paragraph 41: A package for commerce comprising

(a) a pharmaceutical composition which comprises a compound or a pharmaceutically acceptable salt thereof according to any one of paragraphs 20 to 29 as an active ingredient, and

(b) a package insert about the pharmaceutical composition which indicates that the pharmaceutical composition can be used or should be used for treatment or prevention of type II diabetes mellitus.

EFFECT OF THE INVENTION

[0050] Since the pyrazole compound or a pharmaceutically acceptable salt thereof in the present invention has SGLT1 inhibitory activity, it is useful for the treatment and/or prevention of diabetes, obesity, diabetic complications (for example, retinopathy, nephropathy and neuropathy which are known as microangiopathy, as well as cerebrovascular disease, ischemic heart disease and membrum-inferius arteriosclerosis obliterans which are known as macroangiopathy), hypertrophic cardiomyopathy, ischemic heart disease, cancer and constipation.

BEST MODE FOR CARRYING OUT THE INVENTION

[0051] The definitions of the terms in this description are as follows.

The term "optionally substituted" includes both of cases where the replaceable positions of an intended group are substituted and not substituted (non-substituted). Here, the term "non-substituted" means the case where all the replaceable positions of an intended group are occupied with hydrogen atoms.

For example, the term "optionally substituted by 1 to 4 C₁₋₆ alkyl groups" includes both of cases wherein the replaceable positions of an intended group are substituted by 1-4 C₁₋₆ alkyl groups and not substituted (non-substituted), and each substituent may be the same or different.

Unless otherwise specified, an explanation of each group is applied to the case wherein the group is a moiety of other groups or a substituent.

[0052] A "halogen atom" includes, for example, fluorine atom, chlorine atom, bromine atom and iodine atom.

Preferable one is fluorine atom or chlorine atom.

[0053] A "C₁₋₈ alkyl" group means a saturated hydrocarbon group of a straight chain or a branched chain having 1-8 carbon atoms, and the term "C₁₋₆ alkyl" group means the saturated hydrocarbon group of a straight chain or a branched chain having 1-6 carbon atoms.

Among this group, for example, methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, 2-methylpropyl, 1,1-dimethylpropyl, 1-ethylpropyl, hexyl, isohexyl, 1,1-dimethylbutyl, 2,2-dimethylbutyl, 3,3-dimethylbutyl, 2-ethylbutyl are included.

Preferable one is C₁₋₆ alkyl group. More preferable are methyl, ethyl, propyl, isopropyl, n-butyl, tert-butyl, isopentyl, 2-methylpropyl, 1,1-dimethylpropyl, hexyl, and 3,3-dimethylbutyl.

Especially preferable are methyl, ethyl, propyl, n-butyl, and isopentyl.

[0054] A "C₂₋₈ alkenyl" group means a hydrocarbon group having at least one double bond of a straight chain or a branched chain having 2-8 carbon atoms.

As this group, for example vinyl, 1-propenyl, 2-propenyl, isopropenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 1-methyl-2-propenyl, 2-methyl-1-propenyl, 2-methyl-2-propenyl, 1-ethylvinyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1,2-dimethyl-1-propenyl, 1,2-dimethyl-2-propenyl, 1-ethyl-1-propenyl, 1-ethyl-2-propenyl, 1-methyl-1-butenyl, 1-methyl-2-butenyl, 2-methyl-1-butenyl, 1-isopropylvinyl, 2,4-pentadienyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 2,4-hexadienyl, 1-methyl-1-pentenyl, 3,3-dimethylbutenyl (namely, 3,3-dimethyl but-1-enyl), are mentioned.

Preferable one is C₂₋₆ alkenyl group. More preferable are 1-propenyl, 2-methyl-1-propenyl, and 3,3-dimethyl but-1-enyl.

[0055]

A "C₂₋₆ alkynyl" group means a hydrocarbon group which has at least one triple bond of a straight chain or a branched chain having 2-6 carbon atoms, and "C₂₋₈ alkynyl group" means a hydrocarbon group having at least one triple bond of a straight chain or a branched chain having 2-8 carbon atoms. Specifically, ethynyl, 1-propynyl, 2-propynyl, isopropynyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, 1-methyl-1-propynyl, 1-methyl-2-propynyl, 2-methyl-2-propynyl, 1-ethylethynyl, 3,3-dimethylbutynyl (namely, 3,3-dimethyl but-1-ynyl), are mentioned.

Preferable one is C₂₋₆ alkynyl group. More preferable is 3,3-dimethyl but-1-ynyl.

[0056] A "C₁₋₆ alkoxy" group means an alkoxy group with a straight chain or a branched chain of 1-6 carbon atoms.

Among this group, for example, methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, tert-butoxy, pentyloxy, isopentyloxy, 2-methylbutoxy, 1,1-dimethylpropoxy, neopentyl oxy, 3,3-dimethylbutoxy, 1-ethylpropoxy and hexyloxy are included.

Preferable one is C₁₋₄ alkoxy group. More preferable are methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, tert-butoxy, isopentyloxy (namely, 3-methylbutoxy), neopentyl oxy (namely, 2,2-dimethylpropoxy), 1,1-dimethylpropoxy, and 3,3-dimethylbutoxy.

[0057] A "C₆₋₁₀ aryl" group means an aromatic hydrocarbon group having 6-10 carbon atoms.

Among this group, for example, phenyl, 1-naphthyl and 2-naphthyl are included.

Preferable one is phenyl.

[0058] A "C₃₋₈ cycloalkyl" group means a monocycle saturated hydrocarbon group having 3-8 carbon atoms, and the "C₃₋₆ cycloalkyl" group means the monocycle saturated hydrocarbon group having 3-6 carbon atoms.

Among this group, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl are included.

Preferable one is C₃₋₇ cycloalkyl group. More preferable is cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, or cycloheptyl.

[0059] A "C₃₋₈ cycloalkenyl" group means a monocycle unsaturated hydrocarbon group having 3-8 carbon atoms including one or more double bonds.

Among this group, for example, cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl and cyclooctenyl are included. Preferable one is C₃₋₆ cycloalkenyl group. More preferable is cyclohexenyl. Especially preferable is 1-cyclohexenyl.

[0060] A "saturated heterocycle" group means a group having a 4 to 6-membered ring containing 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur in addition to a carbon atom, and whose atomic numbers constituting the ring are 4 to 6, preferably 5 to 6.

Among this group, for example, pyrrolidinyl, pyrrolidino (1-pyrrolidinyl etc.), piperidinyl, piperidino (1-piperidinyl etc.), morpholinyl, morpholino (4-morpholinyl etc.), thiomorpholinyl, thiomorpholino (4-thiomorpholinyl etc.), piperazinyl, piperazino (1-piperazinyl etc.), tetrahydropyranyl ((R)-tetrahydropyran-3-yl, (S)-tetrahydropyran-3-yl etc.), tetrahydrofuranyl ((R)-tetrahydrofuran-3-yl, (S)-tetrahydrofuran-3-yl, etc.), oxetanyl, etc. are included.

Preferable one is a saturated heterocycle group with 5 to 6 members having in its ring containing one heteroatom selected from a nitrogen atom or an oxygen atom in addition to a carbon atom. More preferable are tetrahydropyranyl, tetrahydrofuranyl, oxetanyl and pyrrolidinyl. Especially preferable are (R)-tetrahydropyran-3-yl, (S)-tetrahydropyran-3-yl, 4-tetrahydropyranyl, (R)-tetrahydrofuran-3-yl, (S)-tetrahydrofuran-3-yl, oxetane-3-yl, and 1-pyrrolidinyl.

[0061] A "halo C₁₋₆ alkyl" group means "C₁₋₆ alkyl" group of the above-mentioned definition substituted by 1 to 13 "halogen atoms" of the above-mentioned definition. When multiple halogen atoms substitute, each halogen atom is the same or different. Preferable one is a group which is substituted with 1 to 6 "halogen atoms". More preferable is the group which is substituted with 1 to 6 fluorine atoms.

Among this group, for example, 2-fluoroethyl, 2-chloroethyl, 2-bromoethyl, 3-fluoropropyl, 3-chloropropyl, 4-fluorobutyl, 4-chlorobutyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-difluoro-2-methylpropyl, trifluoromethyl, 2,2,2-trifluoroethyl, 3,3,3-trifluoropropyl, 4,4,4-trifluorobutyl, pentafluoroethyl and 2,2,2-trifluoro-1-trifluoromethylethyl are included.

Preferable one is halo C₁₋₄ alkyl group. More preferable is 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-difluoro-2-methylpropyl, trifluoromethyl, 2,2,2-trifluoroethyl, and 3,3,3-trifluoropropyl.

[0062] A "hydroxy C₁₋₆ alkyl" group means a group in which the hydroxy group(s) is mono- or di-substituted on the "C₁₋₆ alkyl" group of the above-mentioned definition. Preferable one is a group which is mono-substituted by hydroxyl group.

Among this group, for example, hydroxymethyl, 2-hydroxyethyl, 1-hydroxy-1-methylethyl, 1,2-dihydroxyethyl, 3-hydroxypropyl, 1-hydroxy-2,2-dimethylpropyl, 4-hydroxybutyl, 1-hydroxy-2,2-dimethylbutyl, 5-hydroxypentyl and 6-hydroxyhexyl are included.

Preferable one is hydroxy C₁₋₄ alkyl group. More preferable are hydroxymethyl, 2-hydroxyethyl, and 1-hydroxy-1-methylethyl, 1-hydroxy-2,2-dimethylpropyl and 1-hydroxy-2,2-dimethylbutyl.

[0063] A "C₁₋₆ alkoxy C₁₋₆ alkyl" group means a group in which the "C₁₋₆ alkoxy" group(s) of the above-mentioned definition is mono- or di-substituted on the "C₁₋₆ alkyl" group of the above-mentioned definition. When the C₁₋₆ alkoxy groups are di-substituted, each C₁₋₆ alkoxy group is the same or different.

Among this group, for example, methoxymethyl, ethoxymethyl, n-propoxymethyl, isopropoxymethyl, isobutoxymethyl, tert-butoxymethyl, (R)-sec-butoxymethyl, (S)-sec-butoxymethyl, 2-methoxyethyl, (R)-1-propoxyethyl, (S)-1-propoxyethyl, 1-ethylpropoxymethyl, 1-methoxy-1-methylethyl, 1,2-dimethoxyethyl, 2,2-dimethylpropoxymethyl, 2-methoxy-1,1-dimethylethyl, 1-methyl-1-propoxy-ethyl, 2-isopropoxyethyl, 3-methoxypropyl, 3-ethoxypropyl, 2,3-diethoxypropyl, 4-methoxybutyl, 5-methoxypentyl, 5-ethoxypentyl, 6-methoxyhexyl, 6-ethoxyhexyl, pentyloxymethyl and hexyloxymethyl are included.

Preferable one is C₁₋₄ alkoxy C₁₋₃ alkyl group. More preferable are methoxymethyl, ethoxymethyl, n-propoxymethyl, isopropoxymethyl, isobutoxymethyl, 1-methoxy-1-methylethyl, 2-isopropoxyethyl, tert-butoxymethyl, (R)-sec-butoxymethyl, (S)-sec-butoxymethyl, 1-ethylpropoxymethyl, (R)-1-propoxyethyl, (S)-1-propoxyethyl, 1-ethoxy-1-methylethyl, 1-methoxy-1-methylethyl, 2,2-dimethylpropoxymethyl, 2-methoxy-1,1-dimethylethyl, and 1-methyl-1-propoxy-ethyl.

[0064] A "C₃₋₈ cycloalkyl C₁₋₆ alkyl" group means a group in which "C₃₋₈ cycloalkyl" group(s) of the above-mentioned definition is mono- or di-substituted on the "C₁₋₆ alkyl" group of the above-mentioned definition.

When the C₃₋₈ cycloalkyl groups are di-substituted, each C₃₋₈ cycloalkyl group is the same or different.

Among this group, for example, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl,

cyclohexylmethyl, cycloheptylmethyl, cyclooctylmethyl, 2-cyclopentylethyl, 2-cyclohexylethyl, 2-cycloheptylethyl, 2-cyclooctylethyl, 3-cyclopropylpropyl, 3,3-dicyclopropylpropyl, 3-cyclobutylpropyl, 3-cyclopentylpropyl, 2-cyclohexylpropyl, 3-cyclohexylpropyl, 3-cycloheptylpropyl, 3-cyclooctylpropyl, 4-cyclopropylbutyl, 4-cyclobutylmethyl, 4-cyclopentylbutyl, 4-cyclohexylbutyl, 4-cycloheptylbutyl, 4-cyclooctylbutyl, 5-cyclopropylpentyl, 5-cyclobutylpentyl, 3-cyclopentylpentyl, 5-cyclopentylpentyl, 5-cyclohexylpentyl, 5-cycloheptylpentyl, 5-cyclooctylpentyl, 6-cyclopropylhexyl, 6-cyclobutylhexyl, 6-cyclohexylhexyl, 6-cycloheptylhexyl and 6-cyclooctylhexyl are included.

Preferable one is C₃₋₆ cycloalkyl C₁₋₃ alkyl group. More preferable are cyclohexylmethyl, 2-cyclopentylethyl and 2-cyclohexylethyl.

[0065] A "C₃₋₆ cycloalkyl C₂₋₆ alkynyl" group means a group to which "C₃₋₆ cycloalkyl" group(s) of the above-mentioned definition is mono- or di-substituted on the "C₂₋₆ alkynyl" group of the above-mentioned definition. When the C₃₋₆ cycloalkyl groups are di-substituted, each C₃₋₆ cycloalkyl group is the same or different. A Preferable group is a C₂₋₆ alkynyl group mono-substituted by the C₃₋₆ cycloalkyl group.

As this group, for example, cyclopropylethynyl, cyclobutylethynyl, cyclopentylethynyl, cyclohexylethynyl, cyclopropyl-1-propynyl, are mentioned.

Preferable one is C₃₋₆ cycloalkyl C₂₋₄ alkynyl. More preferable is cyclopropylethynyl.

[0066] A "C₆₋₁₀ aryl C₁₋₆ alkyl" group means a group in which the "C₆₋₁₀ aryl" group(s) of the above-mentioned definition is mono- or di-substituted on the "C₁₋₆ alkyl" group of the above-mentioned definition.

When the C₆₋₁₀ aryl groups are di-substituted, each C₆₋₁₀ aryl group is the same or different.

Among this group, for example, benzyl, naphthylmethyl, 1-phenylethyl, 2-phenylethyl, 2-naphthylethyl and 3,3-diphenylpropyl are included.

Preferable one is C₆₋₁₀ aryl C₁₋₃ alkyl group. More preferable are benzyl and 2-phenylethyl.

[0067] A "saturated heterocyclo C₁₋₆ alkyl" group means a group in which "saturated heterocyclo" group(s) of the above-mentioned definition is mono- or di-substituted on the "C₁₋₆ alkyl" group of the above-mentioned definition. When the saturated heterocycle groups are di-substituted, each saturated heterocycle group is the same or different.

Among this group, for example, (1-pyrrolidinyl)methyl (namely pyrrolidine-1-ylmethyl), (1-piperidinyl)methyl, (4-morpholinyl)methyl, (4-thiomorpholinyl)methyl, (1-piperazinyl)methyl, (4-tetrahydropyranyl)methyl, 2-(1-pyrrolidinyl)ethyl, 2-(1-piperidinyl)ethyl, 2-(4-morpholinyl)ethyl, 2-(4-thiomorpholinyl)ethyl, 2-(1-piperazinyl)ethyl, 2-(4-tetrahydropyranyl)ethyl and 2-(4-tetrahydropyranyl)propyl are included.

Preferable one is a saturated heterocyclo C₁₋₃ alkyl group. More preferable are (4-tetrahydropyranyl)methyl, pyrrolidine-1-ylmethyl, 2-(1-pyrrolidinyl)ethyl and 2-(4-tetrahydropyranyl)ethyl.

[0068] A "halo C₁₋₆ alkoxy" group means a group having 1 to 13 "halogen atoms" of the above-mentioned definition substituted on the "C₁₋₆ alkoxy" group of the above-mentioned definition. When multiple halogen atoms are substituted, each halogen atom is the same or different. Preferable one is a group which is substituted by 1 to 6 "halogen atoms". More preferable is a group which is substituted by 1 to 6 fluorine atoms.

Among this group, for example, fluoromethoxy, chloromethoxy, bromomethoxy, 2-fluoroethoxy, 2-chloroethoxy, 2-bromoethoxy, 3-fluoropropoxy, 3-chloropropoxy, 4-fluorobutoxy, 4-

chlorobutoxy, 1,1-difluoroethoxy, 2,2-difluoroethoxy, 1,1-difluoropropoxy, 2,2-difluoropropoxy, 3,3-difluoropropoxy, 1,1-difluoro-2-methylpropoxy, trifluoromethoxy, 2,2,2-trifluoroethoxy, 3,3,3-trifluoropropoxy, 4,4,4-trifluorobutoxy, pentafluoroethoxy and 2,2,2-trifluoro-1-trifluoromethylethoxy are included.

5 Preferable one is halo C_{1-4} alkoxy group. Another preferable group is halo C_{1-6} alkoxy group whose halogens are 1 to 3 fluorine atoms. More preferable are trifluoromethoxy, 2,2,2-trifluoroethoxy and 3,3,3-trifluoropropoxy, 4,4,4-trifluorobutoxy, and 2,2,2-trifluoro-1-trifluoromethylethoxy.

[0069] A "hydroxy C_{1-6} alkoxy" group means a group in which the hydroxy group(s) is mono- or di-substituted on the " C_{1-6} alkoxy" group of the above-mentioned definition. Preferable one is a group
10 which is mono- substituted by hydroxy group.

As this group, for example, hydroxymethoxy, 2-hydroxyethoxy, 1-hydroxy-1-methylethoxy, 1,2-dihydroxyethoxy, 3-hydroxypropoxy, 1-hydroxy-2,2-dimethylpropoxy, 4-hydroxybutoxy, 1-hydroxy-2,2-dimethylbutoxy, 5-hydroxypentyloxy, 6-hydroxyhexyloxy, are mentioned.

Preferable one is hydroxy C_{1-4} alkoxy group. More preferable is 2-hydroxyethoxy.

15 [0070] A "carboxy C_{1-6} alkoxy" group means a group in which carboxy(s) is mono- or di-substituted on the " C_{1-6} alkoxy" group of the above-mentioned definition. Preferable one is a group which is mono-substituted by a carboxy.

Among this group, for example, carboxymethoxy, 2-carboxyethoxy, 3-carboxypropoxy, 2-carboxy-1-methylethoxy, 4-carboxybutoxy, 5-carboxypentyloxy and 6-carboxyhexyloxy are included.

20 Preferable one is carboxy C_{1-4} alkoxy group. More preferable is carboxymethoxy.

[0071] A " C_{1-6} alkoxy C_{1-6} alkoxy" group means a group in which the " C_{1-6} alkoxy" of the above-mentioned definition is mono- or di-substituted on the " C_{1-6} alkoxy" group of the above-mentioned definition. When the C_{1-6} alkoxy groups are di-substituted, each C_{1-6} alkoxy group is the same or different.

25 As this group, for example, methoxymethoxy, ethoxymethoxy, n-propoxymethoxy, isobutoxymethoxy, tert-butoxymethoxy, (R)-sec-butoxymethoxy, (S)-sec-butoxymethoxy, 2-methoxyethoxy, (R)-1-propoxyethoxy, (S)-1-propoxyethoxy, 1-ethylpropoxyethoxy, 1-methoxy-1-methylethoxy, 1,2-dimethoxyethoxy, 2,2-dimethylpropoxymethoxy, 2-methoxy-1,1-dimethylethoxy, 1-methyl-1-propoxy-ethoxy, 2-isopropoxyethoxy, 3-methoxypropoxy, 3-ethoxypropoxy, 2,3-diethoxypropoxy, 4-methoxybutoxy, 5-methoxypentyloxy, 5-ethoxypentyloxy, 6-methoxyhexyloxy, 6-ethoxyhexyloxy, pentyloxymethoxy, hexyloxymethoxy, are mentioned.
30 Preferable one is C_{1-3} alkoxy C_{2-4} alkoxy group. More preferable is 2-methoxyethoxy.

[0072] A " C_{6-10} aryl C_{1-6} alkoxy" group means a group in which the " C_{6-10} aryl" group of the above-mentioned definition is mono- or di-substituted on the " C_{1-6} alkoxy" group of the above-mentioned definition. When the C_{6-10} aryl groups are di-substituted, each C_{6-10} aryl group is the same or different.
35 As this group, benzyloxy, naphthylmethyloxy, 1-phenylethyloxy, 2-phenylethyloxy, 2-naphthylethyloxy, 3,3-diphenylpropoxy, are mentioned, for example.

Preferable one is C_{6-10} aryl C_{1-3} alkyl group. More preferably it is benzyloxy.

[0073] A "halo C_{1-6} alkoxy C_{1-6} alkyl" group means a group in which the "halo C_{1-6} alkoxy" group of the above-mentioned definition is mono- or di-substituted on the " C_{1-6} alkyl" group of the above-mentioned definition. When the halo C_{1-6} alkoxy groups are di-substituted, each halo C_{1-6} alkoxy group is the same or different.
40 Preferable one is halo C_{1-4} alkoxy group. Another preferable group is halo C_{1-6} alkoxy group whose halogens are 1 to 3 fluorine atoms. More preferable are trifluoromethoxy, 2,2,2-trifluoroethoxy and 3,3,3-trifluoropropoxy, 4,4,4-trifluorobutoxy, and 2,2,2-trifluoro-1-trifluoromethylethoxy.

As this group, for example fluoromethoxymethyl, chloromethoxymethyl, bromomethoxymethyl, 2-fluoroethoxymethyl, 2-chloroethoxymethyl, 2-bromoethoxymethyl, 3-fluoropropoxymethyl, 3-chloropropoxymethyl, 4-fluorobutoxymethyl, 4-chlorobutoxymethyl, 1,1-difluoroethoxymethyl, 2,2-difluoroethoxymethyl, 1,1-difluoropropoxymethyl, 2,2-difluoropropoxymethyl, 3,3-difluoropropoxymethyl, 1,1-difluoro-2-methylpropoxymethyl, trifluoromethoxymethyl, 2,3-bis trifluoromethoxypropyl, 2,2,2-trifluoroethoxymethyl, 3,3,3-trifluoropropoxymethyl, 4,4,4-trifluorobutoxymethyl, pentafluoroethoxymethyl, 2,2,2-trifluoro-1-trifluoromethyl-ethoxymethyl, 1-(trifluoromethoxy)ethyl, 1-methyl-2,3-bis-trifluoromethoxypropyl, 1-(2,2,2-trifluoroethoxy)ethyl, 1-(3,3,3-trifluoropropoxy)ethyl, 1-(4,4,4-trifluorobutoxy)ethyl, 1-(2,2,2-trifluoro-1-trifluoromethylethoxy)ethyl, 2-(trifluoromethoxy)ethyl, 2-(2,2,2-trifluoroethoxy)ethyl, 2-(3,3,3-trifluoropropoxy)ethyl, 2-(4,4,4-trifluorobutoxy)ethyl, 2-(2,2,2-trifluoro-1-trifluoromethyl-ethoxy)ethyl, 1-methyl-1-(trifluoromethoxy)ethyl, 1-methyl-1-(2,2,2-trifluoroethoxy)ethyl, 1-methyl-1-(3,3,3-trifluoropropoxy)ethyl, 1-methyl-1-(4,4,4-trifluorobutoxy)ethyl, 1-methyl-1-(2,2,2-trifluoro-1-trifluoromethyl-ethoxy)ethyl, (R)-2,2,2-trifluoro-1-methylethoxymethyl, (S)-2,2,2-trifluoro-1-methylethoxymethyl, 2,2,2-trifluoro-1,1-dimethyl-ethoxymethyl, 2,2,2-trifluoro-1-trifluoromethyl-ethoxymethyl, 2,2,2-trifluoro-1-methyl-1-trifluoromethyl-ethoxymethyl, 3,3,3-trifluoro-2-(trifluoromethyl)propoxymethyl, 4,4,4-trifluoro-3-(trifluoromethyl)butoxymethyl, 1,1-dimethyl-3,3,3-trifluoropropoxymethyl, 2,2-dimethyl-3,3,3-trifluoropropoxymethyl, 4,4,4-trifluoro-3-(trifluoromethyl)butoxymethyl, 1,1-dimethyl-4,4,4-trifluorobutoxymethyl, 2,2-dimethyl-4,4,4-trifluorobutoxymethyl, 3-methyl-4,4,4-trifluoro-3-(trifluoromethyl)-butoxymethyl, 2-methyl-3,3,3-trifluoro-2-trifluoromethyl-propoxymethyl, are mentioned.

Preferable one is halo C₁₋₄ alkoxy C₁₋₃ alkyl group. More preferable are trifluoromethoxymethyl, 2,2,2-trifluoroethoxymethyl, 3,3,3-trifluoropropoxymethyl, (R)-2,2,2-trifluoro-1-methylethoxymethyl, (S)-2,2,2-trifluoro-1-methylethoxymethyl, 2,2,2-trifluoro-1,1-dimethyl-ethoxymethyl, 2,2,2-trifluoro-1-trifluoromethyl-ethoxymethyl and 2,2,2-trifluoro-1-methyl-1-trifluoromethyl-ethoxymethyl.

[0074] A "halo C₃₋₈ cycloalkyl" group means a group in which 1 to 15 "halogen atoms" of the above-mentioned definition is substituted on the "C₃₋₈ cycloalkyl" group of the above-mentioned definition. A preferable group is a group which is substituted by 1 to 2 "halogen atoms".

When multiple halogen atoms are substituted, each halogen atom may be the same or different.

Among this group, for example, 2-fluorocyclopropyl, 2-chlorocyclopropyl, 3-fluorocyclobutyl, 3-chlorocyclobutyl, 3-fluorocyclopentyl, 3-chlorocyclopentyl, 4-fluorocyclohexyl, 4-fluorocycloheptyl, 5-fluorocyclooctyl, 2,2-difluorocyclopropyl, 2,2-dichlorocyclopropyl, 3,3-difluorocyclobutyl, 3,3-dichlorocyclobutyl, 3,3-difluorocyclopentyl, 3,3-dichlorocyclopentyl, 4,4-difluorocyclohexyl and 4,4-dichlorocyclohexyl are included.

Preferable one is halo C₃₋₆ cycloalkyl group. Another preferable group is halo C₃₋₈ cycloalkyl group whose halogen atoms are 1 to 3 fluorine atoms. More preferable is 4,4-difluorocyclohexyl.

[0075] A "C₁₋₆ alkyl sulfanyl" group means the sulfanyl group substituted by the "C₁₋₆ alkyl" group of the above-mentioned definition.

As this group, for example methylsulfanyl, ethylsulfanyl, propylsulfanyl, isopropylsulfanyl,

n-butylsulfanyl, isobutylsulfanyl, sec-butylsulfanyl, tert-butylsulfanyl, pentylsulfanyl, 1,1-dimethylpropylsulfanyl, 2,2-dimethylpropylsulfanyl, hexylsulfanyl, are mentioned.

Preferable one is isobutylsulfanyl.

[0076] A "C₁₋₆ alkyl sulfonyl" group means a sulfonyl group substituted by the "C₁₋₆ alkyl" group of the above-mentioned definition.

As this group, for example, methylsulfonyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl (namely, propane-2-sulfonyl), n-butylsulfonyl, isobutylsulfonyl (namely, 2-methylpropane-1-sulfonyl), sec-butylsulfonyl, tert-butylsulfonyl (namely, 2-methyl-propane-2-sulfonyl), pentylsulfonyl, 1,1-dimethylpropylsulfonyl, hexylsulfonyl, 2,2-dimethylpropylsulfonyl (namely, 2,2-dimethylpropane-1-sulfonyl), are mentioned.

Preferable are propane-2-sulfonyl, 2-methylpropane-1-sulfonyl, 2-methyl-propane-2-sulfonyl, and 2,2-dimethylpropane-1-sulfonyl.

[0077] A "halo C₁₋₆ alkyl sulfonyl" group means a sulfonyl group substituted by the "halo C₁₋₆ alkyl" group of the above-mentioned definition.

As this group, for example, 2-fluoroethanesulfonyl, 2-chloroethanesulfonyl, 2-bromoethanesulfonyl, 3-fluoropropanesulfonyl, 3-chloropropanesulfonyl, 4-fluorobutanesulfonyl, 4-chlorobutanesulfonyl, 1,1-difluoroethanesulfonyl, 1,1-difluoropropanesulfonyl, 1,1-difluoro-2-methylpropanesulfonyl, trifluoromethanesulfonyl, 2,2,2-trifluoroethanesulfonyl, 3,3,3-trifluoropropanesulfonyl, 4,4,4-trifluorobutanesulfonyl, pentafluoroethanesulfonyl, 2,2,2-trifluoro-1-trifluoromethyl ethanesulfonyl, are mentioned.

Preferable are 2,2,2-trifluoro ethanesulfonyl and 3,3,3-trifluoropropanesulfonyl.

[0078] A "C₁₋₈ alkyl carbonyl" group means a carbonyl group substituted by the "C₁₋₈ alkyl" group of the above-mentioned definition.

As this group, acetyl, propionyl, 2,2-dimethylpropionyl, butyryl, 3-methylbutyryl, 2,2-dimethylbutyryl, pentanoyl, 4-methylpentanoyl, 3,4-dimethylpentanoyl, heptanoyl, nonanoyl, are mentioned, for example.

Preferable are 2,2-dimethylbutyryl and 2,2-dimethylpropionyl.

[0079] A "C₁₋₆ alkoxy carbonyl" group means a carbonyl group substituted by the "C₁₋₆ alkoxy" group of the above-mentioned definition.

As this group, for example methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, sec-butoxycarbonyl, tert-butoxycarbonyl, pentyloxycarbonyl, isopentyloxycarbonyl, 2-methylbutoxycarbonyl, 1,1-dimethylpropoxycarbonyl, neopentyloxycarbonyl, 3,3-dimethylbutoxycarbonyl, 1-ethylpropoxycarbonyl, hexyloxycarbonyl, are mentioned.

[0080] A "C₁₋₆ alkoxy carbonyloxy" group means a carbonyloxy group substituted by the "C₁₋₆ alkoxy" group of the above-mentioned definition.

As this group, for example methoxycarbonyloxy, ethoxycarbonyloxy, propoxycarbonyloxy, isopropoxycarbonyloxy, butoxycarbonyloxy, isobuthoxycarbonyloxy, sec-buthoxycarbonyloxy, tert-buthoxycarbonyloxy, pentyloxycarbonyloxy, isopentyloxycarbonyloxy, 2-methylbuthoxycarbonyloxy, 1,1-dimethylpropoxycarbonyloxy, neopentyloxycarbonyloxy, 3,3-dimethylbuthoxycarbonyloxy, 1-ethylpropoxycarbonyloxy, hexyloxycarbonyloxy, are mentioned.

Preferable one is C₁₋₄ alkoxy carbonyloxy group. More preferable is tert-butoxycarbonyloxy. Preferable one is tert-butoxycarbonyl.

[0081] A "C₁₋₆ alkyl sulfonyl C₁₋₆ alkyl" group means a group in which "C₁₋₆ alkyl sulfonyl" group of the above-mentioned definition is mono- or di-substituted on the "C₁₋₆ alkyl" group of the above-mentioned definition. When the C₁₋₆ alkyl sulfonyl groups are di-substituted, each C₁₋₆ alkyl sulfonyl group is the same or different. A preferable group is a group which is mono-substituted by C₁₋₆ alkyl sulfonyl group.

As this group, for example, methylsulfonylmethyl and ethylsulfonylmethyl (namely, ethanesulfonylmethyl), propylsulfonylmethyl, isopropylsulfonylmethyl (namely, propane-2-sulfonylmethyl), n-butylsulfonylmethyl, isobutylsulfonylmethyl (namely, 2-methylpropane-1-sulfonylmethyl), sec-butylsulfonylmethyl, tert-butylsulfonylmethyl (namely, 2-methylpropane-2-sulfonylmethyl), pentylsulfonylmethyl, 1,1-dimethylpropylsulfonylmethyl, hexylsulfonylmethyl, 2,2-dimethylpropylsulfonylmethyl (namely, 2,2-dimethylpropane-1-sulfonylmethyl), 1-ethylsulfonylethyl (namely, 1-ethanesulfonylethyl), 1-isopropylsulfonylethyl (namely, 1-(propane-2-sulfonyl)-ethyl), 2-ethylsulfonylethyl (namely, 2-ethanesulfonylethyl), 2-isopropylsulfonylethyl (namely, 2-(propane-2-sulfonyl)-ethyl), 2,3-bis-methanesulfonylpropyl, 2,3-bis-methanesulfonyl-1-methylpropyl etc. are mentioned.

Preferable one is C₁₋₄ alkyl sulfonyl C₁₋₃ alkyl group. More preferable are ethanesulfonylmethyl and propane-2-sulfonylmethyl.

[0082] A "halo C₁₋₆ alkyl sulfonyl C₁₋₆ alkyl" group means a group in which "halo C₁₋₆ alkyl sulfonyl" group of the above-mentioned definition is mono- or di-substituted on the "C₁₋₆ alkyl" group of the above-mentioned definition. When the halo C₁₋₆ alkyl sulfonyl groups are di-substituted, each halo C₁₋₆ alkyl sulfonyl group is the same or different. A preferable group is a group which is mono-substituted by halo C₁₋₆ alkyl sulfonyl group.

As this group, for example 2-fluoroethanesulfonylmethyl, 2-chloroethanesulfonylmethyl, 2-bromoethanesulfonylmethyl, 3-fluoropropanesulfonylmethyl, 3-chloropropanesulfonylmethyl, 4-fluorobutanesulfonylmethyl, 4-chlorobutanesulfonylmethyl, 1,1-difluoroethanesulfonylmethyl, 1,1-difluoropropanesulfonylmethyl, 1,1-difluoro-isobutanesulfonylmethyl, trifluoromethanesulfonylmethyl, 2,2,2-trifluoroethanesulfonylmethyl, 3,3,3-trifluoropropanesulfonylmethyl, 4,4,4-trifluorobutanesulfonylmethyl, pentafluoroethanesulfonylmethyl, 2,2,2-trifluoro-1-trifluoromethyl-ethanesulfonylmethyl, 1-trifluoromethanesulfonylethyl, 1-(2,2,2-trifluoroethanesulfonyl)-ethyl, 1-(3,3,3-trifluoropropanesulfonyl)-ethyl, 1-(4,4,4-trifluorobutanesulfonyl)-ethyl, 1-(2,2,2-trifluoro-1-trifluoromethyl-ethanesulfonyl)-ethyl, 2-trifluoromethanesulfonylethyl, 2-(2,2,2-trifluoroethanesulfonyl)-ethyl, 2-(3,3,3-trifluoropropanesulfonyl)-ethyl, 2-(4,4,4-trifluorobutanesulfonyl)-ethyl, 2-(2,2,2-trifluoro-1-trifluoromethylethanesulfonyl)-ethyl, 1-methyl-1-trifluoromethanesulfonylethyl, 1-methyl-1-(2,2,2-trifluoro ethanesulfonyl)-ethyl, 1-methyl-1-(3,3,3-trifluoropropanesulfonyl)-ethyl, 1-methyl-1-(4,4,4-trifluorobutanesulfonyl)-ethyl, 1-methyl-1-(2,2,2-trifluoro-1-trifluoromethyl ethanesulfonyl)-ethyl, 2,3-bis(2,2,2-trifluoroethanesulfonyl)propyl, 1-methyl-2,3-bis(2,2,2-trifluoroethanesulfonyl)propyl, are mentioned.

Preferable one is halo C₁₋₄ alkyl sulfonyl C₁₋₃ alkyl group. Preferable one is 2,2,2-trifluoro

ethanesulfonylmethyl.

[0083] A "C₁₋₆ alkyl sulfanyl C₁₋₆ alkoxy" group means a group in which "C₁₋₆ alkyl sulfanyl" group of the above-mentioned definition is mono- or di-substituted on the "C₁₋₆ alkoxy" group of the above-mentioned definition. When the C₁₋₆ alkyl sulfanyl groups are di-substituted, each C₁₋₆ alkyl sulfanyl group is the same or different. A preferable group is a group which is mono-substituted by a C₁₋₆ alkyl sulfanyl group.

As this group, for example, 2-methylsulfanylethoxy, 2-ethylsulfanylethoxy, 2-propylsulfanylethoxy, 2-isopropylsulfanylethoxy, 2-n-butylsulfanylethoxy, 2-isobutylsulfanylethoxy, 2-sec-butyl sulfanylethoxy, 2-tert-butyl sulfanylethoxy, 2-pentylsulfanylethoxy, 2-(1,1-dimethylpropylsulfanyl)ethoxy, 2-(2,2-dimethylpropylsulfanyl)ethoxy, 2-hexylsulfanylethoxy, 3-methylsulfanylpropoxy, 3-ethylsulfanylpropoxy, 3-propylsulfanylpropoxy, 3-isopropylsulfanylpropoxy, 3-n-butylsulfanylpropoxy, 3-isobutylsulfanylpropoxy, 3-sec-butylsulfanylpropoxy, 3-tert-butyl sulfanylpropoxy, 3-pentylsulfanylpropoxy, 3-(1,1-dimethylpropylsulfanyl)propoxy, 3-(2,2-dimethylpropylsulfanyl)propoxy, 3-hexylsulfanylpropoxy, 2-methyl-3-methylsulfanylpropoxy, 2-methyl-3-isopropylsulfanylpropoxy, 2-methyl-3-(2,2-dimethylpropylsulfanyl)-propoxy, 2,3-bis-methylsulfanylpropoxy, 1-methyl-2,3-bis-methylsulfanylpropoxy are mentioned.

Preferable one is C₁₋₃ alkyl sulfanyl C₂₋₄ alkoxy. More preferable is 3-methylsulfanylpropoxy. [0084] A "C₁₋₆ alkyl sulfonyl C₁₋₆ alkoxy" group means a group in which the "C₁₋₆ alkylsulfonyl" group of the above-mentioned definition is mono- or di-substituted on the "C₁₋₆ alkoxy" group of the above-mentioned definition. When the C₁₋₆ alkyl sulfonyl groups are di-substituted, each C₁₋₆ alkyl sulfonyl group may be the same or different.

A preferable group is a group which is mono-substituted by a C₁₋₆ alkyl sulfonyl group.

As this group, for example, 2-methylsulfonylethoxy, 2-ethylsulfonylethoxy, 2-propylsulfonylethoxy, 2-isopropylsulfonylethoxy, 2-n-butylsulfonylethoxy, 2-isobutylsulfonylethoxy, 2-sec-butylsulfonylethoxy, 2-tert-butylsulfonylethoxy, 2-pentylsulfonylethoxy, 2-(1,1-dimethylpropylsulfonyl)ethoxy, 2-(2,2-dimethylpropylsulfonyl)ethoxy, 2-hexylsulfonylethoxy, 3-methylsulfonylpropoxy, 3-ethylsulfonylpropoxy, 3-propylsulfonylpropoxy, 3-isopropylsulfonylpropoxy, 3-n-butylsulfonylpropoxy, 3-isobutylsulfonylpropoxy, 3-sec-butylsulfonylpropoxy, 3-tert-butylsulfonylpropoxy, 3-pentylsulfonylpropoxy, 3-(1,1-dimethylpropylsulfonyl)propoxy, 3-(2,2-dimethylpropylsulfonyl)propoxy, 3-hexylsulfonylpropoxy, 2-methyl-3-methylsulfonylpropoxy, 2-methyl-3-isopropylsulfonylpropoxy, 2-methyl-3-(2,2-dimethylpropylsulfonyl)-propoxy, 2,3-bis-methylsulfonylpropoxy, 1-methyl-2,3-bis-methylsulfonylpropoxy are mentioned.

Preferable one is C₁₋₃ alkyl sulfonyl C₂₋₄ alkoxy. More preferable is 3-methanesulfonylpropoxy.

[0085] A "halo C₁₋₆ alkylamino C₁₋₆ alkyl" group means a group in which an amino group, on which the "halo C₁₋₆ alkyl" group of the above-mentioned definition is mono- or di-substituted, is mono- or di-substituted on the "C₁₋₆ alkyl" group of the above-mentioned definition. A preferable group is a group which is mono-substituted by an amino group on which the halo C₁₋₆ alkyl group is mono-substituted.

As this group, for example, 2-fluoroethylaminomethyl, 2-chloroethylaminomethyl, 2-bromoethylaminomethyl, 3-fluoropropylaminomethyl, 3-chloropropylaminomethyl, 4-fluorobutylaminomethyl, 4-chlorobutylaminomethyl, 1,1-difluoroethylaminomethyl, 1,1-difluoropropylaminomethyl, 1,1-difluoro-2-methylpropylaminomethyl, 2,2,2-trifluoroethylaminomethyl, 3,3,3-trifluoropropylaminomethyl, 4,4,4-trifluorobutylaminomethyl, 2,2,3,3,3-pentafluoropropylaminomethyl, 1-(2,2,2-trifluoroethylamino)-ethyl, 2-(2,2,2-trifluoroethylamino)-ethyl, 1-methyl-1-(2,2,2-trifluoroethylamino)-ethyl, bis-(2,2,2-trifluoroethyl)aminomethyl, are mentioned.

Preferable one is halo C_{1-3} alkylamino C_{1-3} alkyl group. More preferable is 2,2,2-trifluoroethyl aminomethyl.

[0086] As " C_{3-8} cycloalkyloxy" group, cyclopropyloxy, cyclobutoxy, cyclopentyloxy, cyclohexyloxy, cycloheptyloxy, cyclooctyloxy, are mentioned, for example.

Preferable one is C_{3-6} cycloalkyloxy group. More preferable is cyclopentyloxy and cyclobutoxy.

[0087] A "saturated heterocyclo oxy" group means a group in which a chemically acceptable carbon atom in the "saturated heterocycle" group of the above-mentioned definition, and an oxygen atom are combined.

As this group, for example, oxetan-3-yloxy, tetrahydrofuran-3-yloxy, (R)-tetrahydrofuran-3-yloxy, (S)-tetrahydrofuran-3-yloxy, tetrahydropyran-3-yloxy, (R)-tetrahydropyran-3-yloxy, (S)-tetrahydropyran-3-yloxy, tetrahydropyran-4-yloxy, pyrrolidin-3-yloxy, piperidin-3-yloxy, piperidin-4-yloxy, tetrahydrothiopyran-4-yloxy, are mentioned.

Preferable one is oxetan-3-yloxy.

[0088] A "saturated heterocyclo oxy C_{1-6} alkyl" group means a group in which the "saturated heterocyclo oxy" group of the above-mentioned definition is mono- or di-substituted on the " C_{1-6} alkyl" group of the above-mentioned definition. When the saturated heterocyclo oxy groups are di-substituted, each saturated heterocyclo oxy group is the same or different.

As this group, for example, oxetan-3-yloxymethyl, tetrahydrofuran-3-yloxymethyl, (R)-tetrahydrofuran-3-yloxymethyl, (S)-tetrahydrofuran-3-yloxymethyl, tetrahydropyran-3-yloxymethyl, (R)-tetrahydropyran-3-yloxymethyl, (S)-tetrahydropyran-3-yloxymethyl, tetrahydropyran-4-yloxymethyl, pyrrolidin-3-yloxymethyl, piperidine-3-yloxymethyl, piperidin-4-yloxymethyl, tetrahydrothiopyran-4-yloxymethyl, 1-(oxetan-3-yloxy)-ethyl, 1-(tetrahydropyran-4-yloxy)-ethyl, 2-(oxetan-3-yloxy)-ethyl, 2-(tetrahydropyran-4-yloxy)-ethyl, etc. are mentioned.

Preferable one is saturated heterocyclo oxy C_{1-3} alkyl group. More preferable are oxetan-3-yloxymethyl, (R)-tetrahydrofuran-3-yloxymethyl, (S)-tetrahydrofuran-3-yloxymethyl, (R)-tetrahydropyran-3-yloxymethyl, (S)-tetrahydropyran-3-yloxymethyl, and tetrahydropyran-4-yloxymethyl.

[0089] A "saturated heterocyclo C_{1-6} alkoxy" group means a group in which the "saturated heterocyclo" group of the above-mentioned definition is mono- or di-substituted on the " C_{1-6} alkoxy" group of the above-mentioned definition. When the saturated heterocycle groups are di-substituted, each saturated heterocycle group is the same or different.

As this group, for example, (3-pyrrolidinyl)methoxy (namely, pyrrolidin-3-ylmethoxy), (4-

piperidinyl)methoxy, oxetan-3-ylmethoxy, (4-tetrahydropyranyl)methoxy, 2-(1-pyrrolidinyl)ethoxy, 2-(1-piperidinyl)ethoxy, 2-(4-morpholinyl)ethoxy, 2-(4-thio morpholinyl)ethoxy, 2-(1-piperazinyl)ethoxy, 2-oxetan-3-ylethoxy, 2-(4-tetrahydropyranyl) ethoxy, 2-(4-tetrahydropyranyl)propoxy are mentioned.

5 Preferable one is saturated heterocyclo C_{1-3} alkoxy group. More preferable is oxetan-3-ylmethoxy.

[0090] A " C_{6-10} aryl C_{1-6} alkoxy C_{1-6} alkoxy" group means a group in which the " C_{6-10} aryl C_{1-6} alkoxy" group of the above-mentioned definition is mono- or di-substituted on the " C_{1-6} alkoxy" group of the above-mentioned definition. When the C_{6-10} aryl C_{1-6} alkoxy group is di-substituted, each
10 C_{6-10} aryl C_{1-6} alkoxy group is the same or different.

As this group, for example, 2-benzyloxyethoxy, 2-naphthylmethyloxyethoxy, 2-(1-phenylethoxy)-ethoxy, 2-(2-phenylethoxy)-ethoxy, 2-(2-naphthylethoxy)-ethoxy, 2-(3,3-diphenylpropoxy)-ethoxy, 3-benzyloxypropoxy, 3-naphthylmethyloxypropoxy, 3-(1-phenylethoxy)-propoxy, 3-(2-phenylethoxy)-propoxy, 3-(2-naphthylethoxy)-propoxy, 3-(3,3-diphenylpropoxy)-
15 propoxy, 3-benzyloxy-2-methylpropoxy, 2,3-bis-benzyloxypropoxy are mentioned.


Preferable one is C_{6-10} aryl C_{1-3} alkyl C_{1-3} alkoxy group. More preferable is 2-benzyloxyethoxy.

[0091] The case as used in " $-(CH_2)_m-X^a-(C=O)-$ " where X^a is a "bond" means " $-(CH_2)_m-(C=O)-$ ".

[0092] In the general Formula [I], R^3 is optionally substituted on carbon atoms in heterocycle within
20 the chemically acceptable range, and is not substituted on heteroatoms.


[0093] In the general formula [Ib], R^{3a} and R^{3b} on heterocycles denoted by seven kinds of formulae for the ring Cy^a are optionally substituted on carbon atoms in the heterocycle within the chemically acceptable range, and are not substituted on heteroatoms.

[0094] In the case that the ring Cy^a is of the Formula:




25

, the following Formula:




is exemplified for the compound represented by "when two R^{3b} are attached to the same carbon, they optionally form C_{3-6} cycloalkyl group together with the carbon to which they are attached".

30 [0095] In the case that the ring Cy^a is of the Formula:



and R^{3b} is C_{1-6} alkyl, the following Formula:



is exemplified for the compound represented by " R^{3b} optionally form C_{3-6} cycloalkyl group together with the carbon to which it is attached and the carbon adjacent thereto",.

[0096] The preferable embodiments about each group of the compound (it is also henceforth called a "compound of the present invention".) represented by the general formula [I] are explained below.

[0097] The preferable embodiment of the ring Cy is C_{6-10} aryl. The preferable examples of the ring Cy are phenyl, cyclopentyl, cyclohexyl, cycloheptyl, 1-cyclohexenyl. A more preferable ring Cy is phenyl.

The preferable example of $n1$ is 1, 2, 3 or 4. More preferable $n1$ is 1.

[0098] The preferable embodiment of R^1 is

(1) a halogen atom,

(2) hydroxy,

(3) carboxy,

(4) a C_{1-5} alkyl group,

(5) a C_{1-4} alkoxy group,

(6) a C_{3-6} cycloalkyl group,

(7) a halo C_{1-4} alkyl group,

(8) a hydroxy C_{1-4} alkyl group,

(9) a C_{1-4} alkoxy C_{1-3} alkyl group,

(10) a halo C_{1-4} alkoxy group or

(11) a carboxy C_{1-4} alkoxy group,

provided, when $n1$ is 2, 3 or 4, R^1 each is the same or different.

[0099] Another preferable embodiment of R^1 is

(1) a halogen atom,

(2) a C_{1-6} alkyl group,

(3) a C_{1-6} alkoxy group,

(4) a halo C_{1-6} alkyl group,

(5) a C_{1-6} alkoxy C_{1-6} alkyl group or

(6) a halo C_{1-6} alkoxy group,

provided, when $n1$ is 2, 3 or 4, R^1 each is the same or different.

[0100] The preferable example of R^1 is fluoro, chloro, hydroxy, carboxy, methyl, ethyl, propyl, isopropyl, n-butyl, tert-butyl, 2-methylpropyl, 1,1-dimethylpropyl, ethoxy, cyclopropyl, trifluoromethyl,

1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-difluoro-2-methylpropyl, hydroxymethyl, 1-hydroxy-1-methylethyl, 1-methoxy-1-methylethyl, trifluoromethoxy, 2,2,2-trifluoroethoxy, 3,3,3-trifluoropropoxy or carboxymethoxy.

[0101] The preferable embodiment of R^2 is

- 5 (1) a C_{1-6} alkyl group,
- (2) a C_{3-6} cycloalkyl C_{1-3} alkyl group,
- (3) a C_{6-10} aryl C_{1-3} alkyl group,
- (4) a saturated heterocyclo C_{1-3} alkyl group (the saturated heterocycle is of 5-membered to 6-membered saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur),
- 10 (5) a C_{3-7} cycloalkyl group,
- (6) a halo C_{1-4} alkyl group (the halos are 1 to 3 fluorine atoms),
- (7) a C_{1-4} alkoxy C_{1-3} alkyl group,
- (8) a halo C_{3-8} cycloalkyl group (the halos are 1 to 3 fluorine atoms),
- (9) a phenyl group (the phenyl group is optionally substituted by the 1 to 4 substituents selected from
- 15 (a) a halogen atom,
- (b) a C_{1-3} alkyl group,
- (c) a C_{1-3} alkoxy group and
- (d) a carboxy C_{1-3} alkoxy group), or
- (10) a saturated heterocycle group (the saturated heterocycle is of 5-membered to 6-membered saturated
- 20 heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur, and is optionally substituted by 1 to 4 C_{1-3} alkyl groups).

[0102] Another preferable embodiment of R^2 is a C_{6-10} aryl group (the C_{6-10} aryl group is optionally substituted by the 1 to 4 substituents selected from

- (a) a halogen atom,
- 25 (b) a C_{1-6} alkyl group,
- (c) a C_{1-6} alkoxy group, and
- (d) a carboxy C_{1-6} alkoxy group).

[0103] The preferable examples of R^2 are ethyl, propyl, isopropyl, 2-methylpropyl, tert-butyl, 3-methylbutyl, 3,3-dimethylbutyl, hexyl, cyclohexylmethyl, cyclohexylethyl, benzyl, 2-phenylethyl, 4-tetrahydropyranylmethyl, 2-(4-tetrahydropyranyl)ethyl, 2-(1-pyrrolidinyl)ethyl, cyclopentyl, cyclohexyl, 30 cycloheptyl, trifluoromethyl, 2,2,2-trifluoroethyl, 3,3,3-trifluoropropyl, 4,4-difluorocyclohexyl, 2-isopropoxyethyl, 2-cyclopentylethyl, phenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2,4-difluorophenyl, 2-chloro-4-fluorophenyl, 3-fluoro-4-chlorophenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 2-methyl-4-fluorophenyl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 3-carboxy methoxyphenyl, 4-tetrahydropyranyl and 2,6-dimethyl-4-tetrahydropyranyl.

More preferable R^2 is phenyl group.

[0104] The preferable example of n_2 is 0, 1 or 2.

More preferable n_2 is 0 or 1.

40 [0105] The preferable example of m is 0, 1 or 2.

[0106] The preferable example of R^3 is methyl, ethyl or 2-hydroxyethyl.

[0107] The preferable example of X^a is a bond, NH or N-methyl.

More preferable X^a is a bond.

[0108] The preferable example of X^b is NH.

[0109] The preferable example of R^4 is a hydrogen atom or methyl.

[0110] The preferable embodiments of each group of the compound (it is also henceforth called a "compound of the present invention".) represented by the general formula [Ib] are explained below.

[0111] The preferable embodiment of the ring Cy is C_{6-10} aryl. The preferable examples of the ring Cy are phenyl, cyclopentyl, cyclohexyl, cycloheptyl, 1-cyclohexenyl. A more preferable ring Cy is phenyl.

The preferable example of n1 is 1, 2, 3 or 4. More preferable n1 is 1 or 2.

[0112] The preferable embodiment of R^{1a} is

- (1) a halogen atom,
- (2) a C_{1-6} alkyl group,
- 10 (3) a C_{1-6} alkoxy group,
- (4) a C_{3-6} cycloalkyl group,
- (5) a halo C_{1-6} alkyl group,
- (6) a hydroxy C_{1-6} alkyl group,
- (7) a C_{1-6} alkoxy C_{1-6} alkyl group,
- 15 (8) a halo C_{1-6} alkoxy C_{1-6} alkyl group,
- (9) a C_{1-6} alkylsulfonyl C_{1-6} alkyl group,
- (10) a halo C_{1-6} alkylamino C_{1-6} alkyl group,
- (11) a halo C_{1-6} alkoxy group,
- (12) a carboxy C_{1-6} alkoxy group,
- 20 (13) a saturated heterocyclo C_{1-6} alkyl group (the saturated heterocycle is of 4-membered to 6-membered saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur) or
- (14) a saturated heterocyclo oxy C_{1-6} alkyl group (the saturated heterocycle is of 4-membered to 6-membered saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur),
- 25 , provided when n1 is 2, 3 or 4, R^{1a} each is the same or different.

[0113] Another preferable embodiment of R^{1a} is

- (1) a halogen atom,
 - (2) a C_{1-6} alkyl group,
 - (3) a C_{1-6} alkoxy group,
 - 30 (4) a halo C_{1-6} alkyl group,
 - (5) a C_{1-6} alkoxy C_{1-6} alkyl group,
 - (6) a halo C_{1-6} alkoxy group or
 - (7) a halo C_{1-6} alkoxy C_{1-6} alkyl group,
- provided, when n1 is 2, 3 or 4, R^{1a} each is the same or different.

35 [0114] Another preferable embodiment of R^{1a} is

- (1) a halogen atom,
- (2) a C_{1-6} alkyl group,
- (3) a halo C_{1-6} alkoxy group or
- (4) a halo C_{1-6} alkoxy C_{1-6} alkyl group,
- 40 provided, when n1 is 2, 3 or 4, R^{1a} each is the same or different.

[0115] Preferable examples of R^{1a} are chloro, fluoro, hydroxy, carboxy, methyl, ethyl, n-butyl, tert-butyl, propyl, isopropyl, 1,1-difluoropropyl, 1,1-dimethylpropyl, 2-methylpropyl, 3,3-dimethylbut-1-enyl, 3,3-dimethyl but-1-ynyl ethoxy, propoxy, 1,1-dimethylpropoxy, 2,2-dimethylpropoxy, isopropoxy, butoxy, isobutoxy, 3,3-dimethylbutoxy, tert-butoxy, 3-methylbutoxy, cyclopropyl, trifluoromethyl, 1,1-

- difluoroethyl, 1,1-difluoro-2-methylpropyl, 1,1-difluoropropyl, hydroxymethyl, 1-hydroxy-1-methylethyl, 1-hydroxy-2,2-dimethylpropyl, 1-hydroxy-2,2-dimethylbutyl, methoxymethyl, ethoxymethyl, propoxymethyl, isopropoxymethyl, isobutoxymethyl, tert-butoxymethyl, (R)-sec-butoxymethyl, (S)-sec-butoxymethyl, 1-ethylpropoxymethyl, (R)-1-propoxyethyl, (S)-1-propoxyethyl, 1-ethoxy-1-methylethyl, 1-methoxy-1-methylethyl, 2,2-dimethylpropoxymethyl, 2-methoxy-1,1-dimethylethyl, 1-methyl-1-propoxy-ethyl, trifluoromethoxymethyl, 2,2,2-trifluoroethoxymethyl, 3,3,3-trifluoro propoxymethyl, (R)-2,2,2-trifluoro-1-methylethoxymethyl, (S)-2,2,2-trifluoro-1-methylethoxymethyl, 2,2,2-trifluoro-1,1-dimethyl-ethoxymethyl, 2,2,2-trifluoro-1-trifluoromethylethoxymethyl, 2,2,2-trifluoro-1-methyl-1-trifluoromethyl-ethoxymethyl, ethanesulfonylmethyl, propane-2-sulfonylmethyl, 2,2,2-trifluoro ethanesulfonylmethyl, 2,2,2-trifluoroethyl aminomethyl, cyclopropylethynyl, trifluoromethoxy, 2,2,2-trifluoroethoxy, 3,3,3-trifluoropropoxy, 4,4,4-trifluorobutoxy, carboxymethoxy, isobutylsulfanyl, propane-2-sulfonyl, 2-methylpropane-1-sulfonyl, 2-methyl-propane-2-sulfonyl, 2,2-dimethylpropane-1-sulfonyl, 2,2,2-trifluoroethanesulfonyl, 3,3,3-trifluoropropanesulfonyl, 2,2-dimethylbutyryl, 2,2-dimethylpropionyl, tert-butoxycarbonyl, propylcarbamoyl, isopropylcarbamoyl, methylpropylcarbamoyl, 2,2,2-trifluoroethylcarbamoyl, isopropylmethylcarbamoyl, pyrrolidine-1-ylmethyl, (R)-tetrahydrofuran-3-yloxymethyl, (S)-tetrahydrofuran-3-yloxymethyl, (R)-tetrahydropyran-3-yloxymethyl, (S)-tetrahydropyran-3-yloxymethyl, oxetane-3-yloxymethyl and tetrahydropyran-4-yloxymethyl.
- [0116] More preferable examples of R^{1a} are n-butyl, trifluoromethoxy, 2,2,2-trifluoroethoxymethyl, 2,2,2-trifluoro-1-trifluoromethyl-ethoxymethyl, (R)-2,2,2-trifluoro-1-methylethoxymethyl, (S)-2,2,2-trifluoro-1-methylethoxymethyl and 2,2,2-trifluoro-1,1-dimethylethoxymethyl.
- [0117] A preferable embodiment of R^{2a} is
- (1) a C₆₋₁₀ aryl group (the C₆₋₁₀ aryl group is optionally substituted by 1 to 4 substituents selected from
 - (a) a halogen atom,
 - (b) hydroxy,
 - (c) a C₁₋₆ alkyl group,
 - (d) a C₂₋₈ alkenyl group,
 - (e) a C₂₋₆ alkynyl group,
 - (f) a C₁₋₆ alkoxy group,
 - (g) a halo C₁₋₆ alkyl group,
 - (h) a C₁₋₆ alkoxy C₁₋₆ alkyl group,
 - (i) a halo C₁₋₆ alkoxy group,
 - (j) a hydroxy C₁₋₆ alkoxy group,
 - (k) a C₁₋₆ alkoxy C₁₋₆ alkoxy group,
 - (l) a carboxy C₁₋₆ alkoxy group,
 - (m) a C₆₋₁₀ aryl C₁₋₆ alkoxy group,
 - (n) a C₁₋₆ alkylsulfanyl C₁₋₆ alkoxy group,
 - (o) a C₁₋₆ alkyl sulfonyl C₁₋₆ alkoxy group,
 - (p) a C₆₋₁₀ aryl C₁₋₆ alkoxy C₁₋₆ alkoxy group,
 - (q) a saturated heterocyclo C₁₋₆ alkoxy group (the saturated heterocycle is 4-membered to 6-membered saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur and is optionally substituted by 1 to 2 C₁₋₆ alkyl groups),
 - (r) a saturated heterocyclo oxy group (the saturated heterocycle is 4-membered to 6-membered saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen and sulfur,

- (s) a C₃₋₈ cycloalkyloxy group,
- (t) a C₁₋₆ alkoxy-carbonyloxy group and
- (u) a C₁₋₆ alkylsulfonyl group), or

(2) A saturated heterocycle group (the saturated heterocycle is 4-membered to 6-membered saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen and sulfur, and is optionally substituted by 1 to 4 C₁₋₆ alkyl groups).

[0118] Preferable examples of R^{2a} are ethyl, propyl, isopropyl, 2-methylpropyl, tert-butyl, 3-methylbutyl, 3,3-dimethylbutyl, hexyl, cyclohexylmethyl, cyclohexylethyl, benzyl, 2-cyclopentylethyl, 2-phenylethyl, 4-tetrahydropyranylmethyl, 2-(4-tetrahydropyranyl)ethyl, 2-(1-pyrrolidinyl)ethyl, cyclopentyl, cyclohexyl, cycloheptyl, trifluoromethyl, 2,2,2-trifluoroethyl, 3,3,3-trifluoropropyl, 2-isopropoxyethyl, 4,4-difluorocyclohexyl, phenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2,4-difluorophenyl, 2-chloro-4-fluorophenyl, 3-fluoro-4-chlorophenyl, 5-chloro-2-fluorophenyl, 3-hydroxyphenyl, 4-hydroxyphenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 3,5-dimethylphenyl, 2,3-dimethylphenyl, 2,5-dimethylphenyl, 2,6-dimethylphenyl, 2-ethylphenyl, 3-ethylphenyl, 4-ethylphenyl, 3-propylphenyl, 2-isopropylphenyl, 3-isopropylphenyl, 4-isopropylphenyl, 3-isobutylphenyl, 3-(3,3-dimethyl-butyl)phenyl, 3-tert-butylphenyl, 3-(2-methylpropenyl)-phenyl, ((E)-3-propenyl)-phenyl, 3-(3,3-dimethylbut-1-ynyl)-phenyl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 2-ethoxyphenyl, 3-ethoxyphenyl, 4-ethoxyphenyl, 3-propoxyphenyl, 4-propoxyphenyl, 2-isopropoxyphenyl, 3-isopropoxyphenyl, 4-isopropoxyphenyl, 3-isobutoxyphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 3-methoxymethylphenyl, 3-isopropoxymethyl-phenyl, 3-trifluoromethoxy-phenyl, 4-trifluoromethoxy-phenyl, 4-(2,2,2-trifluoro-ethoxy)-phenyl, 4-(3,3,3-trifluoro-propoxy)-phenyl, 3-(2,2,2-trifluoro-ethoxy)-phenyl, 3-(2-hydroxyethoxy)-phenyl, 3-(2-methoxyethoxy)-phenyl, 3-carboxymethoxyphenyl, 3-benzyloxyphenyl, 4-benzyloxyphenyl, 3-(3-methylsulfonylpropoxy)-phenyl, 3-(3-methanesulfonylpropoxy)-phenyl, 3-(2-benzyloxyethoxy)-phenyl, 3-(3-methyloxetan-3-ylmethoxy)-phenyl, 3-(oxetan-3-yloxy)-phenyl, 3-cyclopentylloxyphenyl, 3-cyclobutoxyphenyl, 3-tert-butoxycarbonyloxyphenyl, 3-methanesulfonylphenyl, 4-tetrahydropyranyl, 2,6-dimethyl-4-tetrahydropyranyl, 4-fluoro-2-methyl-phenyl, 4-fluoro-3-(2-hydroxyethoxy)-phenyl, 4-fluoro-3-(2-methoxyethoxy)-phenyl, 2-fluoro-3-(2-methoxyethoxy)-phenyl, 2-fluoro-5-(2-methoxyethoxy)-phenyl, 2-fluoro-3-(2-methoxyethoxy)-phenyl, 2-fluoro-3-methoxy-phenyl, 4-fluoro-3-methoxy-phenyl, 2-fluoro-5-methoxy-phenyl, 3-fluoro-5-methoxy-phenyl, 3-ethoxy-4-fluoro-phenyl, 3-ethoxy-2-fluoro-phenyl, 5-ethoxy-2-fluoro-phenyl, 3-ethoxy-5-fluoro-phenyl, 4-fluoro-3-isopropoxyphenyl, 2-fluoro-5-isopropoxyphenyl, 3-fluoro-5-isopropoxyphenyl, 2-fluoro-3-isopropoxyphenyl, and 4-fluoro-3-propoxy-phenyl.

[0119] More preferable examples of R^{2a} are phenyl, 4-chlorophenyl, 2-fluorophenyl, 3-fluorophenyl and 4-fluorophenyl.

[0120] A preferable example of n₂ is 0, 1 or 2.

More preferable n₂ is 0 or 1.

[0121] A preferable example of m₂ is 1 or 2, and

a preferable example of m₃ is 1 or 2,

provided, the sum of m₂ and m₃ is 3 or 4.

[0122] A preferable example of m₄ is 1 or 2, and

a preferable example of m₅ is 0, 1 or 2,


provided, the sum of m₄ and m₅ is 2, 3 or 4.


[0123] A preferable example of m6 is 0 or 1, and
a preferable m7 is 0, 1 or 2,
provided, the sum of m6 and m7 is 1 or 2.


[0124] A preferable example of m8 is 1.


5 A preferable example of m9 is 1.

[0125] Preferable examples of the ring Cy^a are













[0126] Other preferable examples of the ring Cy^a are








[0127] More preferable examples of the ring Cy^a are





[0128] A preferable example of R^{3a} is hydroxy, methyl, ethyl or 2-hydroxyethyl.

[0129] A more preferable example of R^{3a} is methyl.

5 [0130] A preferable example of R^{3b} is hydroxy, methyl, ethyl, 2-hydroxyethyl, cyclopropyl group which is formed by combining two methyl groups attached on the same carbon with the carbon to which they are attached, or cyclopropyl group which is formed by joining two methyl groups together with the adjacent carbons to which they are attached.


[0131] A more preferable example of R^{3b} is methyl.

10 [0132] A preferable example of R^{4b} is hydrogen atom, methyl, ethyl, propyl, isopropyl, carboxymethyl, 2,2,2-trifluoroethyl or 2-methoxyethyl.

[0133] A more preferable example of R^{4b} is hydrogen atom.


[0134] A certain preferable embodiment of the compound of the present invention is a compound in

general formula [I] wherein the ring Cy is phenyl, n_1 is 1, 2, 3 or 4 and X^b is NH, and the compound is represented by the general formula [III]:




wherein n_1' is 0, 1, 2 or 3, and each of other symbols is the same as those defined above.

- 5 [0135] Another preferable embodiment of the compound of the present invention is a compound in general formula [I] wherein the ring Cy is phenyl, n_1 is 1, 2, 3 or 4, n_2 is 0, m is 1, X^a is a bond and X^b is NH, and the compound is represented by the general formula [III]:




wherein n_1' is 0, 1, 2 or 3, and each of other symbols is the same as those defined above.

- 10 [0136] Another preferable embodiment of the compound of the present invention is a compound in general formula [I] wherein the ring Cy is phenyl, n_1 is 1, 2, 3 or 4, n_2 is 1, m is 1, X^a is a bond and X^b is NH, and the compound is represented by the general formula [IV]:



wherein n_1' is 0, 1, 2 or 3, and each of other symbols is the same as those defined above.


[0137] Another preferable embodiment of the compound of the present invention is a compound in general formula [I] wherein the ring Cy is phenyl, n_1 is 1, 2, 3 or 4, n_2 is 0, m is 2, X^a is a bond and X^b is NH, and the compound is represented by the general formula [V]:



[V]


5 wherein n_1' is 0, 1, 2 or 3, and each of other symbols is the same as those defined above.

[0138] A certain preferable embodiment of the compound of the present invention is a compound represented by the general Formula [VI]:




[VI]

10 wherein, n_1' is 0, 1, 2 or 3, and other symbols are the same as defined above, and the compound is that in the general Formula [Ib], wherein the ring Cy is phenyl and the ring Cy^a is represented by the Formula:




15 [0139] Another preferable embodiment of the compound of the present invention is a compound represented by the general Formula [VII]:




[VII]

wherein, $n1'$ is 0, 1, 2 or 3, and other symbols are the same as defined above, and the compound is that in the general Formula [Ib], wherein the ring Cy is phenyl and the ring Cy^a is represented by the Formula:

5




[0140] Another preferable embodiment of the compound of the present invention is a compound represented by the general Formula [VIII]:




[VIII]

wherein, $n1'$ is 0, 1, 2 or 3, and other symbols are the same as defined above, and the compound is that in the general Formula [Ib], wherein the ring Cy is phenyl and the ring Cy^a is represented by the Formula:

10




[0141] Another preferable embodiment of the compound of the present invention is a compound represented by the general Formula [IX]:



5


[IX]

wherein, $n1'$ is 0, 1, 2 or 3, and other symbols are the same as defined above, and the compound is that in the general Formula [Ib], wherein the ring Cy is phenyl and the ring Cy^a is represented by the Formula:




10

[0142] Another preferable embodiment of the compound of the present invention is a compound represented by the general Formula [X]:




[X]

wherein, $n1'$ is 0, 1, 2 or 3, and other symbols are the same as defined above, and the compound is that in the general Formula [Ib], wherein the ring Cy is phenyl and the ring Cy^a is represented by the Formula:




[0143] Another preferable embodiment of the compound of the present invention is a compound represented by the general Formula [XI]:




[XI]

wherein, $n1'$ is 0, 1, 2 or 3, and other symbols are the same as defined above, and the compound is that in the general Formula [Ib], wherein the ring Cy is phenyl and the ring Cy^a is represented by the Formula:




[0144] Another preferable embodiment of the compound of the present invention is a compound represented by the general Formula [XII]:




[XII]

wherein, $n1'$ is 0, 1, 2 or 3, and other symbols are the same as defined above, and the compound is that in the general Formula [Ib], wherein the ring Cy is phenyl and the ring Cy^a is represented by the Formula:




[0145] Another preferable embodiment of the compound of the present invention is a compound represented by the general Formula [XIII]:




[XIII]

wherein, $n1'$ is 0, 1, 2 or 3, and other symbols are the same as defined above, and

the compound is that in the general Formula [Ib], wherein the ring Cy is phenyl and the ring Cy^a is represented by the Formula:



[0146] Another preferable embodiment of the compound of the present invention is a compound represented by the general Formula [XIV]:




[XIV]


wherein, n1' is 0, 1, 2 or 3, and other symbols are the same as defined

above, and

the compound is that in the general Formula [Ib], wherein the ring Cy is phenyl and the ring Cy^a is represented by the Formula:




[0147] A certain preferable embodiment of the compound of the present invention is a compound represented by the general Formula [XV]:



[XV]

wherein, $n1'$ is 0, 1, 2 or 3, and other symbols are the same as defined above.


[0148] A certain preferable embodiment of the compound of the present invention is a compound
 5 represented by the general Formula [XVI]:



[XVI]

wherein, $n1'$ is 0, 1, 2 or 3, and other symbols are the same as defined above.


[0149] Another preferable embodiment of the compound of the present invention is a compound
 10 represented by the general Formula [XVII]:



[XVII]

wherein, $n1'$ is 0, 1, 2 or 3, and other symbols are the same as defined above.


[0150] A certain preferable embodiment of the compound of the present invention is a compound represented by the general Formula [XVIII]:



[XVIII]

5 wherein, $n1'$ is 0, 1, 2 or 3, and other symbols are the same as defined above.

[0151] Another preferable embodiment of the compound of the present invention is a compound represented by the general Formula [XIX]:



[XIX]

10 wherein, $n1'$ is 0, 1, 2 or 3, and other symbols are the same as defined above.

[0152] A "pharmaceutically acceptable salt" of a compound of the present invention may be any nonpoisonous salt of a compound of the present invention and include any kind of salts, for example, a salt with an inorganic acid, a salt with an organic acid, a salt with an inorganic base, a salt with an organic base and a salt with amino acid.

15 [0153] As the salt with an inorganic acid, salt with hydrochloric acid, nitric acid, sulfuric acid, phosphoric acid, hydrobromic acid are included, for example.

As the salt with an organic acid, for example, salt with oxalic acid, maleic acid, citric acid, fumaric acid, lactic acid, malic acid, succinic acid, tartaric acid, acetic acid, trifluoroacetic acid, gluconic acid, ascorbic acid, methanesulfonic acid, benzenesulfonic acid and p-toluenesulfonic acid are included.

20

[0154] As the salt with an inorganic base, for example, sodium salt, potassium salt, calcium salt, magnesium salt and ammonium salt are included.

As the salt with an organic base, for example, salts with methylamine, diethylamine, trimethylamine, triethylamine, ethanolamine, diethanolamine, triethanolamine, ethylenediamine, tris

(hydroxymethyl) methylamine, dicyclohexylamine, N,N'-dibenzylethylenediamine, guanidine, pyridine, picoline, choline, cinchonine and meglumine are included.

As the salt with amino acid, for example, salts with lysine, arginine, aspartic acid and glutamic acid are included.

[0155] In accordance with known methods, each salt may be obtained by reacting a compound of the present invention with an inorganic base, an organic base, an inorganic acid, an organic acid or amino acid.

[0156] Various "isomers" exist in the compound of the present invention. For example, E isomer and Z isomer exist as geometrical isomers. When asymmetric carbon atoms exist, the enantiomer and the diastereoisomer as stereoisomers based on these in compounds of the present invention exist. When axial asymmetries exist, the stereoisomers based on these exist. By a case, a compound of the present invention may exist in tautomer. Therefore, these all isomers and mixtures thereof are included in the scope of the present invention.

[0157] The compounds represented by the general formulae [I] and [Ib] or pharmaceutically acceptable salts thereof may be in a form of "solvate" thereof. The "solvate" means the compounds represented by the general formulae [I] and [Ib] or pharmaceutically acceptable salts thereof to which is coordinated by a solvent molecules. The solvate which is pharmaceutically acceptable is preferred. The solvate includes, for example, hydrate, ethanolate, dimethyl sulfoxide solvate of the compounds represented by the general formulae [I] and [Ib] or pharmaceutically acceptable salts thereof. Specifically, hemihydrate, monohydrate, dihydrate or mono ethanolate of the compound represented by general formulae [I] and [Ib], or monohydrate of sodium salt of the compound represented by general formulae [I] and [Ib], or 2/3-ethanolate of dihydrochloride of them is mentioned. The solvate may be obtained in accordance with known methods.

[0158] A compound of the present invention may be labeled with isotopes (for example, ^2H , ^3H , ^{14}C , ^{35}S).

As a compound or a pharmaceutically acceptable salt thereof in the present invention, a compound or a pharmaceutically acceptable salt thereof in the present invention purified substantially is preferred. A compound or a pharmaceutically acceptable salt thereof in the present invention over 80% or more of purities is more preferable.

[0159] In the present invention, prodrugs of the compounds represented by the general formulae [I] and [Ib] can also serve as a useful medicine. The "prodrug" means a derivative of the compound of the present invention which has a chemically or metabolically decomposable group, and upon administered to the living body, the derivative restores to the original compound and exerts the original drug action after decomposition by, for example, hydrolysis, solvolysis or decomposition under physiological condition. In the derivatives, a complex and a salt which are not formed by a covalent bond are also included. The prodrug is used for the absorption improvement in oral administration, or for targeting to a targeted position, for example. As a modification site to form a prodrug, functional groups with high reactivity such as hydroxy group, carboxy group, and amino group in the compound of the present invention are included.

[0160] A modifying group for hydroxy group includes, specifically, acetyl group, propionyl group, isobutyryl group, pivaloyl group, palmitoyl group, benzoyl group, 4-methylbenzoyl group, dimethylcarbamoyl group, dimethylaminomethylcarbonyl group, sulfo group, alanyl group, fumaryl group, etc. 3-Carboxybenzoyl group and 2-carboxyethylcarbonyl group which are in the form of sodium-salt are also included.

[0161] A modifying group for carboxy group includes, specifically, methyl group, ethyl group, propyl group, isopropyl group, butyl group, iso-butyl group, tert-butyl group, pivaloyloxymethyl group, carboxymethyl group, dimethylaminomethyl group, 1-(acetyloxy)ethyl group, 1-(ethoxycarbonyloxy)ethyl group, 1-(isopropylloxycarbonyloxy)ethyl group, 1-(cyclohexylloxycarbonyloxy)ethyl group, (5-methyl-2-oxo-1,3-dioxol-4-yl) methyl group, benzyl group, phenyl group, o-tolyl group, morpholinoethyl group, N,N-diethylcarbamoylmethyl group, phthalidyl group, etc.

[0162] A modifying group for amino group includes, specifically, tert-butyl group, docosanoyl group, pivaloylmethyloxy group, alanyl group, hexylcarbamoyl group, pentylcarbamoyl group, 3-methylthio-1-(acetilamino)propylcarbonyl group, 1-sulfo-1-(3-ethoxy-4-hydroxyphenyl)methyl group, (5-methyl-2-oxo-1,3-dioxol-4-yl)methyl group, (5-methyl-2-oxo-1,3-dioxol-4-yl)methoxycarbonyl group, tetrahydrofuranyl group and pyrrolidylmethyl group, etc.

[0163] Since a compound or a pharmaceutically acceptable salt thereof in the present invention has SGLT1 inhibitory activity, it is useful for the treatment and/or prevention of various diseases or conditions of which may be expected to be improved by regulation of SGLT1 activity, for example, diabetes, obesity, diabetic complication (for example, retinopathy, nephropathy and neuropathy which are known as microangiopathy, as well as cerebrovascular disease, ischemic heart disease and membrum-inferius arteriosclerosis obliterans which are known as macroangiopathy), hypertrophic cardiomyopathy, ischemic heart disease, cancer and constipation. Diabetes is type II diabetes mellitus preferably.

[0164] "Inhibition of SGLT1" means inhibiting the function of SGLT1, and disappearing or decreasing the activity, and, for example, means inhibiting the function of SGLT1 based on the conditions of the examples of examinations mentioned below. A preferable "inhibition of SGLT1" is "inhibition of human SGLT1". A preferable "SGLT1 inhibitor" is "human SGLT1 inhibitor".

[0165] "Treatment" means alleviating or healing symptoms or disease, and/or its accompanying symptoms.

"Prevention" means a method of delaying or preventing the onset of symptoms or disease, and/or its accompanying symptoms, a method of keeping a patient from acquiring symptoms or disease, or a method of reducing a risk of a patient acquiring symptoms or disease.

[0166] The pharmaceutical composition of the present invention is manufactured by suitably mixing a compound or a pharmaceutically acceptable salts thereof in the present invention with at least one or more sorts of pharmaceutically acceptable carriers etc. in suitable amounts in accordance with known methods in the technical field of medicinal preparation. Content amounts of a compound or a pharmaceutically acceptable salt thereof in the present invention in the pharmaceutical composition change depending on dosage forms, dose, etc., but are 0.1 to 100 % of the weight to the whole pharmaceutical composition, for example.

[0167] A "pharmaceutical compositions" include oral preparations such as tablets, capsules, granules, powders, trochiscus, syrups, emulsion, and suspension, and parenteral preparations such as external preparations, suppositories, injections, ophthalmic solutions, intranasal agents, and pulmonary agents.

[0168] A "pharmaceutically acceptable carriers" includes various conventional organic or inorganic carrier substances, for example, substances in solid preparations such as excipients, disintegrators, binders, glidants and lubricants, and substances in liquid preparations such as solvents, solubilizing agents, suspending agents, isotonicizing agents, buffers and soothing agents. Additives such as preservatives, antioxidants, colorants, and edulcorants, are used if needed.

[0169] The "excipients" includes, for example, lactose, white soft sugar, D-mannitol, D-sorbitol, corn starch, dextrin, microcrystalline cellulose, crystalline cellulose, carmellose, carmellose calcium, carboxy-methyl-starch sodium, low substituted hydroxypropylcellulose and acacia.

The "disintegrators" include for example, carmellose, carmellose calcium, carmellose sodium, carboxy-methyl-starch sodium, cross-carmellose sodium, crospovidone, low substituted hydroxypropylcellulose, hydroxypropylmethyl cellulose and crystalline cellulose.

The "binders" include, for example, hydroxypropylcellulose, hydroxypropylmethyl cellulose, povidone, crystalline cellulose, white soft sugar, dextrin, starch, gelatin, carmellose sodium and acacia.

The "glidants" include, for example, light anhydrous silicic acid and magnesium stearate.

The "lubricants" include, for example, magnesium stearate, calcium stearate and talc.

[0170] The "solvents" include, for example, purified water, ethanol, propylene glycol, macrogol, sesame oil, corn oil and olive oil.

The "solubilizing agents" include, for example, propylene glycol, D-mannitol, benzyl benzoate, ethanol, triethanolamine, sodium carbonate and sodium citrate.

The "suspending agents" include, for example, benzalkonium chloride, carmellose, hydroxypropylcellulose, propylene glycol, povidone, methylcellulose and glyceryl monostearate.

The "isotonizing agents" include, for example, glucose, D-sorbitol, sodium chloride and D-mannitol.

The "buffers" include, for example, disodium hydrogen phosphate, sodium acetate, sodium carbonate and sodium citrate.

The "soothing agents" include, for example, benzyl alcohol.

[0171] The "preservatives" include, for example, ethyl parahydroxybenzoate, chlorobutanol, benzyl alcohol, sodium dehydroacetate, and sorbic acid.

The "antioxidants" include, for example, sodium sulfite and ascorbic acid.

The "colorants" include, for example, food colors (for example, Food Red No.2 or No. 3, Food Yellow No.4, or No. 5) and beta-carotene.

The "edulcorant" include, for example, saccharin sodium, glycyrrhizinate dipotassium and aspartame.

[0172] The pharmaceutical composition of the present invention can be administered to mammals other than human (for example, mice, rats, hamsters, guinea pigs, rabbits, cats, dogs, pig, cows, horses, sheeps, monkeys) and to human, in oral or parenteral (for example, topical, rectum, intravenous administration) in a therapeutically effective amount. Although the "therapeutically effective amount" changes depending on patients, disease, symptoms, dosage forms, routes of administration, for example, the dose in the case of administering orally to the adult patient (weight: about 60 kg) suffering from diabetes (type II diabetes mellitus etc.) ranges usually from about 1 mg to 1 g per day using, as an active ingredient, a compound or a pharmaceutically acceptable salt thereof in the present invention. Such quantity can be administered to the patient once or in several times a day.

[0173] The pharmaceutical composition comprising a compound or a pharmaceutically acceptable salt thereof in the present invention as an active ingredient or an activator, and the kits (administration, treatment and/or prevention kit), packages (packaging goods etc.), and medicine set (and/or, container) containing the package insert about the pharmaceutical composition which indicate that the pharmaceutical composition can be used or should be used for treatment and/or prevention are also useful. Such kit, the package, and the medicine set may be provided with one or more containers filled with one or more active ingredients and other medicines, or the medicine (or the component) for the

above-mentioned pharmaceutical compositions. As examples of such a kit, a package, and a medicine set, the kit for commerce appropriately directed to the treatment and/or prevention of an object disease and the package for commerce are included. As a package insert comprised in such a kit, a package, and a medicine set, notes by the government organization which regulates manufacture, use or sale of a pharmaceutical or biological products, and notes which show the approval of the government organization about manufacture, use or sale of the product relevant to medication to a human are included. In the above-mentioned kit, package and medicine set, the packed product may also be included, and the structure constituted by adopting a suitable medication step (step) may be included, and the structure constituted as could attain the treatment and/or prevention on more preferable medicine including treatment, prevention of an object disease may be included.

[0174] A compound or a pharmaceutically acceptable salt thereof in the present invention may be used by the general method currently performed in the medicinal field in combination (henceforth "combination therapy") with one or multiple other drugs (henceforth, a "concomitant drug").

[0175] A timing for administration of a compound or a pharmaceutically acceptable salt thereof in the present invention and a concomitant drug is not limited. They may be administrated to the patient as a combination drug, or they may be administrated to the patient simultaneously or in a constant interval. A pharmaceutical kit which is characterized in consisting of a pharmaceutical composition of the present invention and a concomitant drug can be used. The dose of a concomitant drug should comply with the dose in clinical use, and it can be selected suitably depending on patients, disease, symptoms, dosage forms, routes of administration, administration time, combination. The administration method of a concomitant drug is not limited in particular, and a compound or a salt thereof in the present invention and a concomitant drug should just be put together.

[0176] The concomitant drug includes, for example,

- (1) a therapeutic agent and/or prophylactic of dyslipidemia,
- (2) a therapeutic agent and/or prophylactic of obesity,
- (3) a therapeutic agent and/or prophylactic of diabetes,
- (4) a therapeutic agent and/or prophylactic of diabetic complication, and
- (5) a therapeutic agent and/or prophylactic of hypertension,

and any one or multiple of these agents and a compound or a pharmaceutically acceptable salt thereof in the present invention may be used in combination.

[0177] As an example, a method for preparing compounds for working the present invention is explained as follows, and the method for preparing a compound or a pharmaceutically acceptable salt thereof in the present invention is not intended to be limited thereto.


Unless otherwise specified, effective preparation methods may be carried out by devising such as: introducing any protecting groups into any functional groups, if needed, and then deprotecting at a later step; in each step, treating any functional groups in the forms of their precursor, and converting the same into the corresponding desirable functional groups at an appropriate step; interchanging the order of each preparation method and step; optionally using reagents other than those illustrated below in order to accelerate the progress of reaction, etc.

[0178] In each step, aftertreatment of reaction may be performed in a conventional manner, and isolation and purification may be optionally performed by selection from or combination of the conventional procedures, such as crystallization, re-crystallization, distillation, liquid separation, silica gel column chromatography and preparative HPLC, if needed. A reaction may be proceeded to the sequential step without isolation and purification depending on the situation.

[0179]

[Preparation Method 1]

A method for preparing a compound of the general formula [I] is illustrated as "Preparation Method 1-1" and "Preparation Method 1-2" below. Preparation Method 1-1



wherein each symbol has the same meaning as defined above.

(Step 1)

Compound [I] may be obtained by reacting Compound [1] or a salt thereof with Compound [2] in a solvent in the presence of a condensing agent and an additive.

A preferable condensing agent is dicyclohexylcarbodiimide (DCC), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (WSC·HCl), diisopropylcarbodiimide, 1,1'-carbonyldiimidazole (CDI), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU), (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP), diphenylphosphoryl azide, etc.

A preferable additive is 1-hydroxybenzotriazole (HOBt), 1-hydroxy-7-azabenzotriazole (HOAt), N-hydroxysuccinimide (HOSu). Optionally, 4-dimethylaminopyridine, etc. may be added as an additive, etc.

A preferable solvent is a halogenated hydrocarbon solvent such as chloroform; an ether solvent such as tetrahydrofuran; a polar solvent such as pyridine, acetonitrile and N,N-dimethylformamide; or a mixed solvent thereof.

The reaction temperature is usually about 0°C to 100°C, preferably about 0°C to 40°C.


When a salt of Compound [1] is used, the reaction may be done in the presence of a base such as an organic base such as triethylamine; or an alkali metal salt such as sodium carbonate.

Alternatively, Compound [I] may be obtained by preliminarily converting Compound [2] into an acid chloride thereof, etc., and then reacting the resulting compound with Compound [1] in the presence of a base.

Compound [2] may be commercially available, or may be obtained by using Preparation Method 3 below or a conventional method.

[0180]

Preparation Method 1-2



wherein, P^{N1} is a protecting group of an amino group; a preferable P^{N1} is tert-butoxycarbonyl group; and each of other symbols has the same meaning as defined above.

5 (Step 1)

Compound [4] may be obtained by reacting Compound [1] or a salt thereof with Compound [3] in a solvent in the presence of a condensing agent and an additive.

A preferable condensing agent is dicyclohexylcarbodiimide (DCC), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (WSC·HCl), diisopropylcarbodiimide, 1,1'-carbonyldiimidazole (CDI), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU), (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP), diphenylphosphoryl azide, etc.

A preferable additive is 1-hydroxybenzotriazole (HOBt), 1-hydroxy-7-azabenzotriazole (HOAt), N-hydroxysuccinimide (HOSu), etc. Optionally, 4-dimethylaminopyridine, etc. may be added as an additive, etc.

A preferable solvent is a halogenated hydrocarbon solvent such as chloroform; an ether solvent such as tetrahydrofuran; a polar solvent such as pyridine, acetonitrile and N,N-dimethylformamide; or a mixed solvent thereof.

The reaction temperature is usually about 0°C to 100°C, preferably about 0°C to 40°C.

When a salt of Compound [1] is used, the reaction may be done in the presence of a base such as an organic base such as triethylamine; or an alkali metal salt such as sodium carbonate.

Alternatively, Compound [4] may be obtained by preliminarily converting Compound [3] into an acid chloride thereof, etc., and then reacting the resulting compound with Compound [1] in the presence of a base.

Compound [3] may be commercially available, or may be obtained by a conventional method. [0181]

(Step 2)

Compound [5] or a salt thereof may be obtained by removing P^{N1} from Compound [4] in a

conventional deprotection reaction. The deprotection reaction may be done by using suitable conditions for each of kinds of P^{N1} . For example, when P^{N1} is tert-butoxycarbonyl group, Compound [5] or a salt thereof may be obtained by treating Compound [4] in a solvent under an acidic condition at room temperature. Compound [5] may be isolated as a salt, or as a free form after neutralization.

A preferable solvent is, for example, an alcoholic solvent such as ethanol; an ether solvent such as 1,4-dioxane; an ester solvent such as ethyl acetate; a halogenated hydrocarbon solvent such as chloroform; or a mixed solvent thereof.

A preferable acid used in the reaction is hydrochloric acid or trifluoroacetic acid.

[0182]

(Step 3)

Compound [Ia] may be obtained by transforming diamino group of Compound [5] or a salt thereof into urea group.

The reagents used in the reaction include, for example, diethyl carbonate, diethyl pyrocarbonate, 1,1'-carbonyldiimidazole (CDI), chloroformate, etc., preferably 1,1'-carbonyldiimidazole (CDI).

A preferable solvent is a halogenated hydrocarbon solvent such as chloroform; an ether solvent such as tetrahydrofuran; a hydrocarbon solvent such as toluene; a polar solvent such as pyridine, acetonitrile and N,N-dimethylformamide; or a mixed solvent thereof.


The reaction temperature is usually about 0°C to 100°C, preferably about 0°C to 40°C.

When a salt of Compound [5] is used, the reaction may be done in the presence of a base such as an organic base such as triethylamine; or an alkali metal salt such as sodium carbonate.

[0183] When Compound [I] or Compound [Ia] has an enantiomer or a geometric isomer thereof, the reaction may be done by using a commercially available optically active compound or a single diastereomer compound or such a compound obtained by a conventional method. Alternatively, the optically active compound or the single diastereomer of Compound [I] or Compound [Ia] may be obtained by separating and purifying appropriately during the preparation process. The compound which separation and purification undergoes includes Compound [I] or Compound [Ia], etc. A method for separation and purification includes, for example, column chromatography or preparative HPLC.

[0184] A method for preparing the compound represented by the general Formula [Ib] is illustrated in the following Preparation Method 1-3.

Preparation Method 1-3



wherein, each symbol has the same meaning as defined above.

(Step 1)

Similarly to Preparation Method 1-1 Step 1, Compound [Ib] or a salt thereof may be obtained. Compound [2d] or a salt thereof may be commercially available, or it can be obtained by the following Preparation Method 3 or a conventional method.

When Compound [Ib] has an enantiomer or a geometric isomer thereof, the reaction may be


done by using a commercially available optically active compound or a single diastereomer compound or such a compound obtained by a conventional method. Alternatively, the optically active compound or single diastereomer of Compound [Ib] may be obtained by separating and purifying during the preparation process. The compound which undergoes separation and purification includes Compound [Ib], etc. A method for separation and purification includes, for example, column chromatography, preparative HPLC.

[0185]

[Preparation Method 2]

A method for preparing Compound [1] is illustrated as Preparation Method 2-1 to 2-3" below.

Preparation Method 2-1



wherein, each symbol has the same meaning as defined above.

(Step 1)

Compound [9] may be obtained by subjecting Compound [8] to Wittig reaction or Horner-Wadsworth-Emmons reaction in a solvent in the presence of a base.

A preferable reagent used in the Wittig Reaction is (cyanomethyl)triphenylphosphonium chloride, and a preferable reagent used in the Horner-Wadsworth-Emmons reaction is diethyl cyanomethylphosphonate.

The base includes, for example, an organic base such as triethylamine; or an alkali metal base such as sodium hydride, potassium tert-butoxide, sodium carbonate and lithium hexamethyldisilazide (LiHMDS), preferably potassium tert-butoxide.

A preferable solvent is, for example, an ether solvent such as tetrahydrofuran; a hydrocarbon solvent such as toluene; a polar solvent such as N,N-dimethylformamide; or a mixed solvent thereof.

The reaction temperature is usually about $-78^{\circ}C$ to $100^{\circ}C$, preferably about $0^{\circ}C$ to $40^{\circ}C$.

Compound [8] may be commercially available, or obtained by a conventional method.

[0186]

(Step 2)

Compound [11] or a salt thereof may be obtained by reacting Compound [9] with Compound [10] or a salt thereof in a solvent in the presence of a base.

The base used in the reaction includes, for example, an organic base such as triethylamine; and an alkali metal base such as sodium hydride, sodium ethoxide, potassium tert-butoxide, sodium carbonate or lithium hexamethyldisilazide (LiHMDS), preferably sodium ethoxide.

A preferable solvent is an alcoholic solvent such as ethanol; an ether solvent such as tetrahydrofuran; a hydrocarbon solvent such as toluene; a polar solvent such as N,N-dimethylformamide; or a mixed solvent thereof.

The reaction temperature is usually about 0°C to 150°C, preferably about 20°C to 100°C.

Compound [10] or a salt thereof may be commercially available, or may be obtained by a conventional method.

[0187]

5 (Step 3)

Compound [1] or a salt thereof may be obtained by reacting Compound [11] or a salt thereof with an oxidizing agent in a solvent.


A preferable oxidizing agent is 2,3-dichloro-5,6-dicyano-p-benzoquinone (DDQ), p-chloranil, Oxone (Registered Trademark), manganese dioxide, activated carbon, oxygen gas, etc.

10 A preferable solvent is, for example, an ether solvent such as dioxane; a polar solvent such as N,N-dimethylformamide; a hydrocarbon solvent such as toluene; a halogenated hydrocarbon solvent such as chloroform; or a mixed solvent thereof.

The reaction temperature is usually about 0°C to 150°C, preferably about 20°C to 150°C.

[0188]

15 Preparation Method 2-2



wherein, P^{N^2} is hydrogen atom or a protecting group of amino group; preferably, two P^{N^2} s are combined together with the nitrogen atom to which they are attached to form 2,5-dimethylpyrrole; L^1 is a leaving group, preferably bromine atom or iodine atom.; each of the other symbols has the same meaning as defined above.

20

(Step 1)

Compound [13] or a salt thereof may be obtained by the same procedure as Step 3 of Preparation Method 2-1.

Compound [12] or a salt thereof may be commercially available, or may be obtained by a conventional method.

25

[0189]

(Step 2)

Compound [14] may be obtained by introducing P^{N^2} into amino group of Compound [13] or a salt thereof by a conventional protection reaction. The protection reaction may be done by using

suitable conditions for each of kinds of P^{N2} . For example, when two P^{N2} s together with the nitrogen atom to which they are attached to form 2,5-dimethylpyrrole, Compound [14] may be obtained by reacting with 2,5-hexanedione in a solvent under an acidic condition.

The acid used in the reaction includes, for example, conc. hydrochloric acid, conc. sulfuric acid, amidosulfuric acid, p-toluenesulfonic acid or acetic acid, preferably acetic acid.

A preferable solvent is, for example, an alcoholic solvent such as ethanol; an ether solvent such as tetrahydrofuran; a hydrocarbon solvent such as toluene; a polar solvent such as N,N-dimethylformamide; a halogenated hydrocarbon solvent such as dichloroethane; or a mixed solvent thereof. Alternatively, an organic acid such as acetic acid may be used as the solvent.

The reaction temperature is usually about 0°C to 150°C, preferably about 20°C to 120°C.

[0190]

(Step 3)

Compound [15] may be obtained by introducing L^1 into Compound [14] in a solvent in the presence of a base. For example, when L^1 is iodine atom, Compound [15] may be obtained by iodination in a solvent in the presence of a base.

The base used in the reaction includes, for example, n-butyllithium, lithium hexamethyldisilazide, or lithium tetramethylpiperidide, preferably n-butyllithium.

The iodination agent includes, for example, iodine, iodine monochloride, N-iodosuccinimide or 1-chloro-2-iodoethane, preferably iodine.

A preferable solvent is, for example, an ether solvent such as tetrahydrofuran; a hydrocarbon solvent such as toluene; or a mixed solvent thereof.

The reaction temperature is usually about -100°C to 40°C, preferably about -78°C to 20°C.

[0191]

(Step 4)

Compound [16] or a salt thereof may be obtained by removing P^{N2} of Compound [15] by a conventional deprotection reaction. The deprotection reaction may be done by using suitable conditions for each of kinds of P^{N2} . For example, when two P^{N2} s are together with the nitrogen atom to which they are attached to form 2,5-dimethylpyrrole, Compound [16] or a salt thereof may be obtained by treating Compound [15] in a solvent in the presence of hydroxylamine. Compound [16] may be isolated as a salt, or as a free form after neutralization.

A preferable solvent is an alcoholic solvent such as ethanol; water; or a mixed solvent thereof.

The reaction temperature is usually about 40°C to 150°C, preferably about 80°C to 130°C.

[0192]

(Step 5)

Compound [1] or a salt thereof may be obtained by treating Compound [16] or a salt thereof and Compound [17] in Suzuki coupling reaction. For example, Compound [1] or a salt thereof may be obtained by reacting Compound [16] or a salt thereof with Compound [17] in a solvent under heating in the presence of a base and a palladium catalyst.

The palladium catalyst used in the reaction includes, for example, tetrakis(triphenylphosphine)palladium or (bis(diphenylphosphino)ferrocene)palladium dichloride-methylene chloride complex.


The base used in the reaction includes potassium phosphate, sodium carbonate, sodium hydrogen carbonate, potassium carbonate and triethylamine, etc.

A preferable solvent is an ether solvent such as 1,4-dioxane, tetrahydrofuran, diethyl ether

and 1,2-dimethoxyethane; an alcoholic solvent such as methanol, ethanol, 1-propanol and 2-propanol; a hydrocarbon solvent such as toluene, n-hexane and xylene; a polar solvent such as N,N-dimethylformamide, dimethylsulfoxide, acetonitrile; or a mixed solvent thereof with water.

Compound [17] may be commercially available, or obtained by a conventional method. Alternatively, the reactions in this step may be done by using boronic acid ester.

By using Compound [12a] and Compound [17a] respectively in place of Compound [12] and Compound [17], Compound [1a] may be obtained similarly to the Preparation Method 2-2.



wherein, Compound [17a] is boronic acid ester, and each symbol has the same meaning as defined above.

Compound [12a] or a salt thereof may be commercially available, or may be obtained by a conventional method.

Compound [17a] may be commercially available, or may be obtained by a conventional method. The reactions in this step may be carried out by using boronic acid.


A reagent used for obtaining Compound [17a] includes the following reagents.

For example, when the ring Cy is C₆₋₁₀ aryl group substituted by a leaving group (preferably chlorine atom, bromine atom, iodine atom, p-toluenesulfonyloxy group, methanesulfonyloxy group or trifluoromethanesulfonyloxy group) in addition to R^{1a}, palladium catalysts such as palladium acetate, palladium chloride, and tris(dibenzylideneacetone)dipalladium; organic phosphorous compounds such as triphenylphosphine, tricyclohexylphosphine, 1,1'-bis(diphenylphosphino)ferrocene, 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl, 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl and 2-dicyclohexylphosphino-2'-(N,N-dimethylamino)biphenyl; boron compounds such as bis(pinacolate)diboron; bases such as potassium acetate, sodium carbonate, and potassium carbonate, are included.

For example, when the ring Cy is C₆₋₁₀ aryl group which does not have any substituent other than R^{1a}, iridium reagents such as di-μ-methoxybis(1,5-cyclo)(octadiene)diiridium; bipyridine compounds such as 4,4'-di-tert-butyl-2,2'-bipyridine and 4,4'-dimethoxy-2,2'-bipyridine; boron compounds such as bis(pinacolate)diboron are included.

[0193]

Preparation Method 2-3



wherein P^{N2} is hydrogen atom or a protecting group of amino group, preferably, two P^{N2} s combine together with the nitrogen atom to which they are attached to form 2,5-dimethylpyrrole; L^1 is a leaving group, preferably bromine atom or iodine atom; L^2 is a leaving group, preferably chlorine atom, bromine atom, iodine atom, dihydroxyboryl group, p-toluenesulfonyloxy group or methanesulfonyloxy group; each of the other symbols has the same meaning as defined above.

(Step 1)

Compound [19] may be obtained by the same procedure as Step 2 of Preparation Method 2-2.

[0194]

(Step 2)

Compound [14] may be obtained by reacting Compound [19] with Compound [20] in a solvent.

For example, when R^2 is C_{1-8} alkyl group or C_{3-8} cycloalkyl group and L^2 is chlorine atom, bromine atom, iodine atom, p-toluenesulfonyloxy group or methanesulfonyloxy group, Compound [14] may be obtained by reacting Compound [19] with Compound [20] in a solvent in the presence of a base.

The base used in the reaction includes, for example, potassium hydroxide, potassium carbonate, sodium ethoxide, potassium tert-butoxide, sodium hydride, preferably sodium hydride.

A preferable solvent is, for example, an ether solvent such as tetrahydrofuran; a polar solvent such as N,N-dimethylformamide; an alcoholic solvent such as ethanol; or a mixed solvent thereof.

The reaction temperature is usually about 0°C to 100°C , preferably about 0°C to 80°C .

For example, when R^2 is C_{6-10} aryl group and L^2 is bromine atom or iodine atom, Compound [14] may be obtained by reacting Compound [19] with Compound [20] in a solvent in the presence of a metal reagent, a ligand and a base.

The metal reagent used in the reaction includes, for example, copper (I) iodide, copper (I) oxide, cobalt (II) chloride, manganese (II) chloride or iron (III) chloride, preferably copper (I) oxide.

The ligand used in the reaction includes, for example, salicylaldehyde, trans-1,2-cyclohexanediamine, 8-quinolinol, 1,10-phenanthroline, L-proline, and preferably 8-quinolinol.

The base used in the reaction includes, for example, potassium hydroxide, tripotassium phosphate, potassium carbonate, cesium carbonate or sodium tert-butoxide, preferably cesium carbonate.

A preferable solvent is, for example, a hydrocarbon solvent such as toluene and xylene; a polar solvent such as acetonitrile and N,N-dimethylformamide; or a mixed solvent thereof.

The reaction temperature is usually about 20°C to 150°C , preferably about 80°C to 130°C .

For example, when R^2 is C_{6-10} aryl group and L^2 is dihydroxyboryl group, Compound [14]

may be obtained by reacting Compound [19] with Compound [20] in a solvent in the presence of a copper reagent and a base in air at room temperature.

A preferable copper reagent used in the reaction is copper (II) acetate and copper (I) oxide.

A preferable base used in the reaction is an organic base such as pyridine and triethylamine.

A preferable solvent is, for example, an alcoholic solvent such as methanol and ethanol; a halogenated hydrocarbon solvent such as dichloromethane and chloroform; or a mixed solvent thereof..

Compound [20] may be commercially available, or may be obtained by a conventional method.

[0195]

(Step 3)

Compound [15] may be obtained by the same procedure as Step 3 of Preparation Method 2-2.

[0196]

(Step 4)


Compound [16] or a salt thereof may be obtained by the same procedure as Step 4 of Preparation Method 2-2.

[0197]

(Step 5)

Compound [1] or a salt thereof may be obtained by the same procedure as Step 5 of Preparation Method 2-2.

By using Compound [20a] and Compound [17a] respectively in place of Compound [20] and Compound [17], Compound [1a] may be obtained similarly to the Preparation Method 2-3.



wherein, each symbol has the same meaning as defined above.


Compound [20a] or a salt thereof may be commercially available, or it may be obtained by a conventional method.

The reaction in this step may be carried out using boronic acid in place of Compound [17a].

[0198]

[Preparation Method 3]

A method for preparing Compound [2] is illustrated as Preparation Methods 3-1 to 3-3 below.

Preparation Method 3-1

wherein, P^{Cl} is a protecting group of carboxy group, preferably methyl group, ethyl group, tert-butyl group or benzyl group; $\text{P}^{\text{N}3}$ is a protecting group of amino group, preferably 2,4-dimethoxybenzyl group; carbon atoms with * may be optionally substituted by R^3 within the chemically acceptable range; each of the other symbols has the same meaning as defined above.

(Step 1)


Compound [23] may be obtained by reacting Compound [21] with Compound [22] in a solvent.

A preferable solvent is, for example, a hydrocarbon solvent such as toluene; an alcoholic solvent such as methanol and ethanol; or a mixed solvent thereof.

The reaction temperature is usually about 20°C to 150°C , preferably about 80°C to 130°C .

Compound [21] may be commercially available, or may be obtained by Auxiliary Step 1 below or a conventional method. Compound [22] may be commercially available, or obtained by a conventional method.

[0199]

Auxiliary Step 1

wherein, P^{Cl} is a protecting group of carboxy group, preferably methyl group, ethyl group, tert-butyl group or benzyl group; m_1 is 0, 1, 2 or 3; L^3 is a leaving group, preferably, bromine atom; carbon atoms with * may be optionally substituted by R^3 within the chemically acceptable range; each of the other symbols has the same meaning as defined above.

(Auxiliary Step 1-1)

Compound [26] may be obtained by introducing L^3 into Compound [25]. For example, when L^3 is bromine atom, Compound [26] may be obtained by brominating in a solvent in the presence of a

radical reaction initiator.

A preferable radical reaction initiator is azobisisobutyronitrile (AIBN).

A preferable brominating agent is N-bromosuccinimide (NBS).

A preferable solvent is a halogenated hydrocarbon solvent such as dichloromethane and carbon tetrachloride.

The reaction temperature is usually about 20°C to 130°C, preferably about 40°C to 100°C.

Compound [25] may be commercially available, or may be obtained by a conventional method.

[0200]

(Auxiliary Step 1-2)

Compound [21a] may be obtained by introducing R³ into Compound [26]. For example, Compound [21a] may be obtained by reacting Compound [26] with a Grignard reagent in a solvent in the presence of an additive.

A preferable additive is hexamethylphosphoric triamide (HMPA).

A preferable solvent is, for example, an ether solvent such as diethyl ether and tetrahydrofuran; a polar solvent such as N,N-dimethylformamide; or a mixed solvent thereof.

The reaction temperature is usually about -78°C to 40°C, preferably about -20°C to 25°C.

The Grignard reagent may be commercially available, or may be obtained by a conventional method.

[0201]

(Step 2)

Compound [24] or a salt thereof may be obtained by removing P^{C1} of Compound [23] by a conventional deprotection reaction. The deprotection reaction may be done by using suitable conditions for each of kinds of P^{C1}. For example, when P^{C1} is methyl group, Compound [24] or a salt thereof may be obtained by hydrolyzing Compound [23] in a solvent in the presence of a base.

The base includes, for example, lithium hydroxide, sodium hydroxide, potassium hydroxide, sodium methoxide, preferably sodium hydroxide.

A preferable solvent is, for example, an alcoholic solvent such as methanol; an ether solvent such as tetrahydrofuran; water; or a mixed solvent thereof.

The reaction temperature is usually about 0°C to 100°C, preferably about 0°C to 40°C.

[0202]

(Step 3)

Compound [2a] or a salt thereof may be obtained by removing P^{N3} of Compound [24] or a salt thereof by a conventional deprotection reaction. The deprotection reaction may be done by using suitable conditions for each of kinds of P^{N3}. For example, when P^{N3} is 2,4-dimethoxybenzyl group, Compound [2a] or a salt thereof may be obtained by treating with an acid in a solvent in the presence of an additive.

The acid includes, for example, methanesulfonic acid, p-toluenesulfonic acid, trifluoroacetic acid, preferably trifluoroacetic acid.

The additive includes, for example, anisole and triethylsilane; preferably anisole.

A preferable solvent is, for example, a halogenated hydrocarbon solvent such as dichloromethane; a hydrocarbon solvent such as toluene; water; or a mixed solvent thereof. Alternatively, an organic acid such as trifluoroacetic acid may be used as the solvent.


The reaction temperature is usually about 0°C to 130°C, preferably about 25°C to 80°C.

When Compound [2a] has geometric isomers, a single diastereomer may be obtained by separating and purifying appropriately during the preparation process. The compound which undergoes separation and purification includes Compound [23], etc. A method for separation and purification includes, for example, column chromatography, or preparative HPLC. Alternatively,

Enantiomers resulted from an asymmetric carbon atom in the α -position of carboxy group of Compound [2a] may be separated and purified as a single diastereomer by introducing a chiral auxiliary to give an optically active compound thereof. For example, Compounds [2a_chiral_1] and [2a_chiral_2] may be obtained by using (R)-4-benzyl-2-oxazolidinone as the chiral auxiliary for Compound [24] or a salt thereof as shown in Auxiliary Step 2 below.

[0203]

Auxiliary Step 2



wherein, carbon atoms with * may be optionally substituted by R^3 within the chemically acceptable range; each of the other symbols has the same meaning as defined above.

(Auxiliary Step 2-1)

Compound [27_chiral_1] and Compound [27_chiral_2] may be obtained by reacting Compound [24] or a salt thereof with (R)-4-benzyl-2-oxazolidinone in a solvent in the presence of a condensing agent and an additive. At this stage, Compound [27_chiral_1] and Compound [27_chiral_2] may be separated and purified by silica gel column chromatography.

A preferable condensing agent is dicyclohexylcarbodiimide (DCC), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (WSC·HCl), etc.

A preferable additive is 4-dimethylaminopyridine, etc.

A preferable solvent is a halogenated hydrocarbon solvent such as chloroform, etc.

The reaction temperature is usually about 0°C to 100°C, preferably about 0°C to 40°C.

Alternatively, Compound [27_chiral_1] and Compound [27_chiral_2] may be obtained by preliminarily converting Compound [24] into an acid chloride thereof, etc., and then reacting the resulting compound with (R)-4-benzyl-2-oxazolidinone in the presence of a base.

[0204]

(Auxiliary Step 2-2)

Compound [24_chiral_1] or a salt thereof (or Compound [24_chiral_2] or a salt thereof) may be obtained by reacting Compound [27_chiral_1] (or Compound [27_chiral_2]) in a solvent in the

presence of a base and an aqueous hydrogen peroxide.

The base used in the reaction includes, for example, sodium hydroxide, lithium hydroxide, preferably lithium hydroxide.

A preferable solvent is, for example, an ether solvent such as tetrahydrofuran; water; or a mixed solvent thereof.

The reaction temperature is usually about 0°C to 50°C, preferably about 0°C to 25°C.


[0205]

(Auxiliary Step 2-3)

Compound [2a_chiral_1] or a salt thereof (or Compound [2a_chiral_2] or a salt thereof) may be obtained by the same procedure as Step 3 of Preparation Method 3-1.

[0206]

Preparation Method 3-2



wherein, carbon atoms with * may be optionally substituted by R³ within the chemically acceptable range; each of the other symbols has the same meaning as defined above.

(Step 1)

Compound [29] or a salt thereof may be obtained by treating Compound [28] or a salt thereof with bromine, followed by a base in a solvent.

The base includes, for example, potassium hydroxide, sodium hydrogen carbonate, sodium carbonate, triethylamine, preferably sodium hydrogen carbonate or sodium carbonate.

A preferable solvent is, for example, a halogenated hydrocarbon solvent such as dichloromethane, carbon tetrachloride; water.

The reaction temperature is usually about 0°C to 100°C, preferably about 25°C to 80°C.

Compound [28] or a salt thereof may be commercially available, or may be obtained by a conventional method.

[0207]

(Step 2)

Compound [2b] or a salt thereof may be obtained by reducing Compound [29] or a salt thereof by a catalytic hydrogenation in a solvent in the presence of a metal catalyst under hydrogen atmosphere.


The metal catalyst includes, for example, palladium on carbon, alumina-supported rhodium, Raney nickel, Adams' catalyst, preferably palladium on carbon.

A preferable solvent is, for example, an alcoholic solvent such as methanol and ethanol; an ether solvent such as tetrahydrofuran; an ester solvent such as ethyl acetate; water, or a mixed solvent thereof.

The reaction temperature is usually about 25°C to 80°C, preferably about 25°C to 50°C.

[0208]

Preparation Method 3-3



wherein, P^{N4} is a protecting group of amino group, preferably benzyloxycarbonyl group; P^{C3} is a protecting group of carboxy group, preferably methyl group, ethyl group, tert-butyl group or benzyl group; R^{4a} is C_{1-6} alkyl group; L^4 is a leaving group, preferably halogen atom; carbon atoms with * may be optionally substituted by R^3 within the chemically acceptable range; each of the other symbols has the same meaning as defined above.

(Step 1)

Compound [31] or a salt thereof may be obtained by subjecting Compound [30] or a salt thereof to Hofmann rearrangement reaction in the presence of a base and bromine, followed by intramolecular cyclization reaction in a solvent.

The base includes, for example, sodium hydroxide, potassium hydroxide, sodium methoxide, preferably sodium hydroxide.

A preferable solvent is, for example, an alcoholic solvent such as methanol and ethanol; an ether solvent such as 1,4-dioxane; water; or a mixed solvent thereof.

The reaction temperature is usually about -78°C to 100°C , preferably about -40°C to 80°C .

Compound [30] or a salt thereof may be commercially available, or may be obtained by a conventional method.

[0209]

(Step 2)

Compound [32] may be obtained by introducing P^{C3} into carboxy group of Compound [31] or a salt thereof by a conventional protection reaction. The protection reaction may be done by using suitable conditions for each of kinds of P^{C3} . For example, P^{C3} is tert-butyl group, Compound [32] may be obtained by preliminarily converting Compound [31] into an acid chloride thereof in a solvent, etc., and then reacting the resulting compound with tert-butyl alcohol in the presence of a base

The reagent used in the reaction is, for example, thionyl chloride, oxalyl chloride, phosphorous oxychloride, preferably phosphorous oxychloride.

The base includes, for example, an organic amine such as triethylamine and pyridine; an alkali metal carbonate such as sodium carbonate, potassium carbonate and sodium hydrogen

carbonate, preferably pyridine.

A preferable solvent is, for example, a hydrocarbon solvent such as toluene; an ether solvent such as tetrahydrofuran; a halogenated hydrocarbon solvent such as chloroform; or a mixed solvent thereof.

The reaction temperature is usually about 0°C to 130°C, preferably about 25°C to 80°C.

[0210]

(Step 3)

Compound [34] may be obtained by rearranging P^{N4} of Compound [32], followed by introducing R^{4a} into the rearranged compound in a solvent in the presence of a base.

A preferable base is sodium hydride.

A preferable solvent is, for example, an ether solvent such as tetrahydrofuran; a polar solvent such as N,N-dimethylformamide; or a mixed solvent thereof.

The reaction temperature is usually about 0°C to 80°C, preferably about 0°C to 25°C.

Compound [33] may be commercially available, or may be obtained by a conventional method.

[0211]

(Step 4)

Compound [35] may be obtained by removing P^{N4} from Compound [34] by a conventional deprotection reaction. The deprotection reaction may be done by using suitable conditions for each of kinds of P^{N4} . For example, when P^{N4} is benzyloxycarbonyl group, Compound [35] may be obtained by reducing Compound [34] by a catalytic hydrogenation in a solvent in the presence of a palladium catalyst under hydrogen atmosphere.

The palladium catalyst includes, for example, palladium on carbon, palladium (II) hydroxide, preferably palladium on carbon.

A preferable solvent is, for example, an alcoholic solvent such as methanol; an ether solvent such as tetrahydrofuran; an ester solvent such as ethyl acetate; or a mixed solvent thereof.

The reaction temperature is usually about 25°C to 80°C, preferably about 25°C to 50°C.

[0212]

(Step 5)

Compound [2c] or a salt thereof may be obtained by removing P^{C3} from Compound [35] by a conventional deprotection reaction. The deprotection reaction may be done by using suitable conditions for each of kinds of P^{C3} . For example, when P^{C3} is tert-butyl group, Compound [2c] may be obtained by treating Compound [35] in a solvent under an acidic condition.

The acid includes, for example, hydrochloric acid, sulfuric acid, trifluoroacetic acid, preferably trifluoroacetic acid.


A preferable solvent is, for example, an ether solvent such as 1,4-dioxane; a halogenated hydrocarbon solvent such as chloroform; an ester solvent such as ethyl acetate; an alcoholic solvent such as methanol; water; or a mixed solvent thereof.

The reaction temperature is usually about 0°C to 80°C, preferably about 0°C to 40°C.

When Compound [2c] has an enantiomer thereof, the reaction may be done by using a commercially available optically active compound or using a product obtained therefrom by a conventional method.

[0213]

Preparation Method 3-4



wherein, carbon atoms indicated by * in the Formula may be optionally substituted by R^{3a} within the chemically acceptable range, each of the other symbols has the same meaning as defined above.

(Step 1)

Compound [37] or a salt thereof may be obtained by the same procedure as Step 1 of Preparation Method 3-3.

Compound [36] or a salt thereof may be commercially available, or may be obtained by a conventional method.


(Step 2)

Compound [2d] or a salt thereof may be obtained by the same procedure as Step 4 of Preparation Method 3-3.

When Compound [2d] has an enantiomer thereof, the reaction may be carried out by using a commercially available optically active compound or using a product obtained therefrom by a conventional method.

[0214]

Preparation Method 3-5



wherein, P^{N5} is a protecting group of amino group, preferably benzyloxycarbonyl group; carbon atoms indicated by * may be optionally substituted by R^{3a} within the chemically acceptable range; each of the other symbols has the same meaning as defined above.

(Step 1)

Compound [2e] or a salt thereof may be obtained by reacting Compound [38] or a salt thereof in the presence of a base in a solvent.

The base includes, for example, sodium hydroxide, potassium hydroxide, sodium methoxide, preferably sodium hydroxide.

A preferable solvent is, for example, an alcoholic solvent such as methanol and ethanol; an ether solvent such as 1,4-dioxane; water; or a mixed solvent thereof.

The reaction temperature is usually about 0 to 100°C, preferably about 25 to 50°C.


Compound [38] or a salt thereof may be commercially available, or may be obtained by a conventional method.

When Compound [2e] has an enantiomer thereof, the reaction may be carried out using a commercially available optically active compound or using a product obtained therefrom by a

conventional method.

[0215]

Preparation Method 3-6



wherein, P^{C4} is a protecting group of carboxy group, preferably methyl group, ethyl group, tert-butyl group, or benzyl group; carbon atoms indicated by * may be optionally substituted by R^{3a} within the chemically acceptable range; each of the other symbols has the same meaning as defined above.

(Step 1)

Compound [40] may be obtained by subjecting Compound [39] or a salt thereof to Curtius rearrangement reaction, followed by an intramolecular cyclization reaction in a solvent.

The reagent used for the reaction includes, for example, diphenylphosphoryl azide.

A preferable solvent is, for example, a hydrocarbon solvent such as toluene; an alcoholic solvent such as ethanol and tert-butanol; an ether solvent such as 1,4-dioxane; or a mixed solvent thereof.

The reaction temperature is usually about 40 to 140°C, preferably about 80 to 120°C.

Compound [39] or a salt thereof may be commercially available, or may be obtained by a conventional method.

[0216]

(Step 2)

Compound [2e] or a salt thereof may be obtained by removing P^{C4} of Compound [40] by a conventional deprotection reaction. The deprotection reaction may be carried out using suitable conditions for each of kinds of P^{C4} , for example, when P^{C4} is benzyl group, Compound [2e] or a salt thereof may be obtained by reducing Compound [40] by a catalytic hydrogenation in the presence of a palladium catalyst in a solvent under hydrogen atmosphere.

The palladium catalyst includes, for example, palladium on carbon, palladium (II) hydroxide, preferably palladium on carbon.

A preferable solvent is an alcoholic solvent such as methanol; an ether solvent such as tetrahydrofuran; an ester solvent such as ethyl acetate; or a mixed solvent thereof.

The reaction temperature is usually about 25 to 80°C, preferably about 25 to 50°C.

When Compound [2e] has an enantiomer thereof, the reaction may be carried out using a commercially available optically active compound or using a product obtained therefrom by a conventional method.

[0217]

Preparation Method 3-7

5 (Step 1)

10 The reaction temperature is usually about 20 to 150°C, preferably about 80 to 130°C.

15 [0218]

25 The reaction temperature is usually about 0 to 100°C, preferably about 0 to 40°C.

(Step 3).

The acid includes, for example, methanesulfonic acid, p-toluenesulfonic acid, trifluoroacetic acid, preferably trifluoroacetic acid.


The additive includes, for example, anisole, triethylsilane, preferably anisole.

A preferable solvent is, for example, a halogenated hydrocarbon solvent such as dichloromethane; a hydrocarbon solvent such as toluene; water; or a mixed solvent thereof. The organic acid itself such as trifluoroacetic acid may be used as a solvent.

5 The reaction temperature is usually about 0 to 130°C, preferably about 25 to 80°C.

[0220]

Preparation Method 3-8



wherein, $\text{P}^{\text{C}6}$ is a protecting group of carboxy group, preferably methyl group, ethyl group, tert-butyl group or benzyl group; L^6 is a leaving group, preferably chlorine atom or bromine atom; carbon atoms indicated by * may be optionally substituted by R^{3a} within the chemically acceptable range; each of the other symbols has the same meaning as defined above.

(Step 1)

Compound [45] may be obtained by reacting Compound [44] with Compound [22] in a solvent. Compound [22] itself may be used as a solvent.

A preferable solvent is, for example, a hydrocarbon solvent such as toluene; an alcoholic solvent such as ethanol and tert-butanol; a polar solvent such as acetonitrile and N,N-dimethylformamide; or a mixed solvent thereof.

The reaction temperature is usually about 20 to 150°C, preferably about 80 to 130°C.

Compound [44] may be commercially available, or may be obtained by a conventional method.

[0221]

(Step 2)

Compound [47] may be obtained by reacting Compound [45] with Compound [46] in the presence of a base in a solvent.

The base includes, for example, an organic amine such as triethylamine and pyridine; an alkali metal carbonate such as sodium carbonate, potassium carbonate and sodium hydrogen carbonate, preferably triethylamine.

A preferable solvent is, for example, a hydrocarbon solvent such as toluene; an ether solvent such as tetrahydrofuran; a halogenated hydrocarbon solvent such as chloroform; a polar solvent such as acetonitrile; or a mixed solvent thereof.

The reaction temperature is usually about -20 to 130°C, preferably about 0 to 80°C.

5 Compound [48] may be commercially available, or may be obtained by a conventional method.
[0222]
(Step 3)

10 Compound [48] may be obtained by reacting Compound [47] in the presence of a base in a solvent.

The base includes, for example, an alkali metal carbonate such as sodium carbonate, potassium carbonate and cesium carbonate; potassium tert-butoxide, sodium hydride, preferably potassium tert-butoxide.

15 The solvent includes, for example, a hydrocarbon solvent such as toluene; an ether solvent such as tetrahydrofuran; an alcoholic solvent such as tert-butanol; a polar solvent such as N,N-dimethylformamide; or a mixed solvent thereof.

The reaction temperature is usually about 0 to 130°C, preferably about 25 to 80°C.

[0223]
(Step 4)

20 Compound [59] or a salt thereof may be obtained by the same procedure as Step 2 of Preparation Method 3-1.


[0224]
(Step 5)

25 Compound [2g] or a salt thereof may be obtained by the same procedure as Step 3 of Preparation Method 3-1.

An enantiomer derived from an asymmetric carbon atom of the alpha position of carboxy group of Compound [2g] may be obtained by the same procedure as Auxiliary step 2 of Preparation Method 3-1.

[0225]

30 Preparation Method 3-9



- wherein, P^{C7} is a protecting group of carboxy group, preferably methyl group, ethyl group, tert-butyl group or benzyl group; P^{C8} is a protecting group of carboxy group, preferably methyl group, ethyl group, tert-butyl group or benzyl group; P^{C9} is a protecting group of carboxy group, preferably methyl group, ethyl group, tert-butyl group or benzyl group; P^{N6} is a protecting group of amino group, preferably tert-butoxycarbonyl group; P^{N7} is a protecting group of amino group, preferably benzyloxycarbonyl group; L⁷ is a leaving group, preferably chlorine atom or bromine atom; carbon atoms indicated by * may be optionally substituted by R^{3a} within the chemically acceptable range; each of the other symbols has the same meaning as defined above.
- 5
- 10 (Step 1)

Compound [52] or a salt thereof may be obtained by reacting Compound [50] or a salt thereof with Compound [51] in the presence of a base in a solvent.

The base includes, for example, an organic amine such as triethylamine and pyridine; an alkali metal carbonate such as sodium carbonate, potassium carbonate and sodium hydrogen

carbonate, preferably potassium carbonate.

A preferable solvent is, for example, a hydrocarbon solvent such as toluene; an ether solvent such as tetrahydrofuran; a halogenated hydrocarbon solvent such as chloroform; a polar solvent such as acetonitrile; or a mixed solvent thereof.

5 The reaction temperature is usually about 0 to 130°C, preferably about 25 to 80°C.

Compound [50] or a salt thereof may be commercially available, or may be obtained by a conventional method. Compound [51] may be commercially available, or may be obtained by a conventional method.

[0226]

10 (Step 2)

Compound [53] or a salt thereof may be obtained by removing P^{C8} of Compound [52] or a salt thereof by a conventional deprotection reaction. The deprotection reaction may be carried out using suitable conditions for each of kinds of P^{C8} , for example, when P^{C3} is tert-butyl group, Compound [53] or a salt thereof may be obtained by the same procedure as Step 5 of Preparation Method 3-3.

15 [0227]

(Step 3)

Compound [54] or a salt thereof may be obtained by introducing P^{N6} into amino group of Compound [53] or a salt thereof by a conventional protection reaction. The protection reaction may be carried out using suitable conditions for each of kinds of P^{N6} , for example, when P^{N6} is tert-butoxycarbonyl group, Compound [54] or a salt thereof may be obtained by reacting with di-tert-butyl dicarbonate in the presence of a base in a solvent.

20

The base includes, for example, an organic amine such as triethylamine and pyridine; an alkali metal carbonate such as sodium carbonate, potassium carbonate and sodium hydrogen carbonate, preferably triethylamine.

25

A preferable solvent is, for example, a hydrocarbon solvent such as toluene; an ether solvent such as tetrahydrofuran; a halogenated hydrocarbon solvent such as chloroform; a polar solvent such as acetonitrile; water; or a mixed solvent thereof.

The reaction temperature is usually about 0 to 130°C, preferably about 25 to 50°C.

[0228]

30 (Step 4)

Compound [55] may be obtained by subjecting Compound [54] or a salt thereof to Curtius rearrangement reaction in the presence of an alcohol in a solvent.

The reagent used for the reaction includes, for example, diphenylphosphoryl azide.

The alcohol used for the reaction includes, for example, benzyl alcohol, tert-butanol, preferably benzyl alcohol.

35

A preferable solvent is, for example, a hydrocarbon solvent such as toluene; an ether solvent such as 1,4-dioxane; or a mixed solvent thereof.

The reaction temperature is usually about 25 to 140°C, preferably about 80 to 120°C.

[0229]

40 (Step 5)

Compound [56] or a salt thereof may be obtained by removing P^{N7} and P^{C9} of Compound [55] by a conventional deprotection reaction. The deprotection reaction may be carried out using suitable conditions for each of kinds of P^{N7} and P^{C9} , for example, when P^{N7} is benzyloxycarbonyl group and P^{C9} is benzyl group, Compound [58] or a salt thereof may be obtained by reducing Compound [57]

by a catalytic hydrogenations in the presence of a metal catalyst in a solvent under hydrogen atmosphere.

The metal catalyst includes, for example, palladium on carbon, palladium black, palladium (II) hydroxide, preferably palladium on carbon.

A preferable solvent is, for example, an alcoholic solvent such as methanol and ethanol; an ether solvent such as tetrahydrofuran an ester solvent such as ethyl acetate water or a mixed solvent thereof.

The reaction temperature is usually about 25 to 80°C, preferably about 25 to 50°C.

[0230]

(Step 6)

Compound [57] may be obtained by reacting Compound [56] or a salt thereof in the presence of a condensing agent and an additive in a solvent.

A preferable condensing agent is dicyclohexylcarbodiimide (DCC), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (WSC·HCl), diisopropylcarbodiimide, 1,1'-carbonyldiimidazole (CDI), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU), (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP) or diphenylphosphoryl azide, etc.

A preferable additive is 1-hydroxybenzotriazol (HOBt), 1-hydroxy-7-azabenzotriazol(HOAt), N-hydroxysuccinimide (HOSu), etc. 4-Dimethylaminopyridine etc. may be optionally added as an additive.

A preferable solvent is a halogenated hydrocarbon solvent such as chloroform; an ether solvent such as tetrahydrofuran; a polar solvent such as pyridine, acetonitrile and N,N-dimethylformamide; or a mixed solvent thereof.

The reaction temperature is usually about 0 to 100°C, and preferably is about 0 to 40°C.

When an acidic salt of Compound [56] is used, the reaction may be carried out in the presence of a base including an organic base such as triethylamine; an alkali metal salt such as sodium carbonate; etc.

[0231]

(Step 7)

Compound [2h] or a salt thereof may be obtained by removing P^{C7} of Compound [57] by a conventional deprotection reaction. The deprotection reaction may be carried out using suitable conditions for each of kinds of P^{C7}. For example, when P^{C7} is methyl group, Compound [2h] or a salt thereof may be obtained by hydrolyzing Compound [57] in the presence of a base in a solvent.

The base includes, for example, lithium hydroxide, sodium hydroxide, potassium hydroxide, preferably sodium hydroxide.

A preferable solvent is, for example, an alcoholic solvent such as methanol; an ether solvent such as tetrahydrofuran; water; or a mixed solvent thereof.

The reaction temperature is usually about 0 to 100°C, preferably about 0 to 40°C.

When Compounds [2h] has an enantiomer thereof, the reaction may be carried out using a commercially available optically active compound or using a product obtained therefrom by a conventional method.

When Compound [2h] or a salt thereof is used, Compound [Ib] or a salt thereof may be obtained by treating by the same procedure as Preparation Method 1-3, followed by removing P^{N6} by a conventional deprotection reaction. The deprotection reaction may be carried out using suitable

conditions for each of kinds of P^{N6} . For example, when P^{N6} is tert-butoxycarbonyl group, Compound [Ib] or a salt thereof may be obtained by the same procedure as Step 2 of Preparation Method 1-2.

EXAMPLES

- 5 [0232] Preparations of a compound or a pharmaceutically acceptable salt thereof in the present invention are specifically illustrated by Examples. However, the present invention is not intended to be limited thereto.

Stereochemistry in chemical structural formulae of compounds are partially abbreviated in the Examples.

- 10 [0233] Measurement apparatuses and conditions used in the Examples are as follows.

HPLC analysis condition 1

Preparation for solvent A: Sodium dihydrogen phosphate dihydrate (2.34 g) was dissolved in water (3000 ml) to be adjusted to pH 2.6 by using phosphoric acid (1.02 ml).

Measurement instrument: HPLC system SHIMADZU CORPORATION High Performance Liquid

- 15 Chromatograph Prominence

Column: DAICEL CHIRALPAK AD-3R 4.6 mmφ × 150 mm

Column temperature: 40°C

Mobile phase: (solvent A) 10 mM phosphate (sodium) buffer (pH 2.6), (solvent B) acetonitrile
solvent A: solvent B = 60:40 constant (12 minutes flow)

- 20 Flow rate: 0.5 ml/min

Detection: UV (220 nm)

HPLC analysis condition 2

Preparation for solvent A: Sodium dihydrogenphosphate dihydrate (2.34 g) was dissolved in water (3000 ml), and to be adjusted to pH 2.6 by using phosphoric acid (1.02 ml).

- 25 Measurement instrument: HPLC system SHIMADZU CORPORATION High Performance Liquid Chromatograph Prominence

Column: DAICEL CHIRALPAK AD-3R 4.6 mmφ × 150 mm

Column temperature: 40°C

- 30 Mobile phase: (solvent A) 10 mM phosphate (sodium) buffer (pH 2.6), (solvent B) acetonitrile
Solvent A : Solvent B = 50:50 constant (12 minutes flow)

Flow rate: 0.5 ml/min

Detection: UV (220 nm)

- 35 HPLC analysis condition 3

Measurement instrument: HPLC system SHIMADZU CORPORATION High Performance Liquid Chromatograph Prominence

Column: DAICEL CHIRALCEL OD 4.6 mmφ × 250 mm

Column temperature: 40°C

- 40 Mobile phase: (solvent A) n-hexane, (solvent B) 2-propanol

Solvent A: Solvent B = 10:90 constant (30 minutes flow)


Flow rate: 0.3 ml/min

Detection: UV (254 nm)


[0234] NMR was used with 400 MHz.

[0235]

[Preparation 1]: Synthesis of 4-methyl-5-oxopyrrolidine-3-carboxylic acid



(1) (Z)-2-Bromomethyl-2-butenedioic acid dimethyl ester




To a solution of (Z)-2-methyl-2-butenedioic acid dimethyl ester (15.3 g) in carbon tetrachloride (300 ml) were added N-bromosuccinimide (25.8 g) and 2,2'-azobis(isobutyronitrile) (319 mg), and the mixture was stirred for 36 hours at reflux. After cooling the mixture to room temperature, the insoluble substance was filtered off, and the filtrate was concentrated. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/0 to 18/1) to give the titled compound (19.6 g).

¹H-NMR (CDCl₃) δ: 3.83 (s, 3H), 3.88 (s, 3H), 4.72 (s, 2H), 6.84 (s, 1H).

[0236]

(2) 2-Methyl-3-methylenesuccinic acid dimethyl ester




To a solution of (Z)-2-bromomethyl-2-butenedioic acid dimethyl ester (19.3 g) and hexamethylphosphoric acid triamide (70 ml) in diethyl ether (200 ml) was added dropwise methyl magnesium bromide (1.12 M solution in tetrahydrofuran, 100 ml) over 1 hour at -20°C and stirred for additional 1 hour. To this reaction mixture were added a 6M aqueous solution of hydrochloric acid (18 ml) and a saturated aqueous solution of ammonium chloride (100 ml), and then the mixture was warmed to room temperature and extracted with diethyl ether (200 ml). The resulting organic layer was washed with water twice, followed by a saturated aqueous solution of sodium chloride once, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/chloroform = 1/1 to 0/1) to give the titled compound (9.92 g).

¹H-NMR (CDCl₃) δ: 1.38 (d, 3H, J = 7.2 Hz), 3.62 (q, 1H, J = 7.2 Hz), 3.69 (s, 3H), 3.77 (s, 3H), 5.72 (s, 1H), 6.34 (s, 1H).

[0237]

(3) Mixture of (cis)-1-(2,4-dimethoxybenzyl)-4-methyl-5-oxopyrrolidine-3-carboxylic acid methyl ester and (trans)-1-(2,4-dimethoxybenzyl)-4-methyl-5-oxopyrrolidine-3-carboxylic acid methyl ester




To a solution of 2-methyl-3-methylenesuccinic acid dimethyl ester (9.81 g) in toluene (100 ml) was added 2,4-dimethoxybenzylamine (9.72 g) and the mixture was stirred for 3 days at reflux. This reaction solution was cooled to room temperature and then purified by silica gel column chromatography (eluent: n-hexane/chloroform = 1/1 to 0/1) to give the titled compound (14.2 g).

¹H-NMR (CDCl₃) cis δ: 1.13 (d, 3H, J = 7.5 Hz), 2.75-2.86 (m, 1H), 3.22-3.34 (m, 2H), 3.48-3.56 (m, 1H), 3.69 (s, 3H), 3.80 (bs, 6H), 4.41 (d, 1H, J = 14.4 Hz), 4.47 (d, 1H, J = 14.4 Hz), 6.42-6.48 (m, 2H), 7.13-7.17 (m, 1H). trans δ: 1.30 (d, 3H, J = 6.6 Hz), 2.66-2.80 (m, 2H), 3.31-3.43 (m, 2H), 3.71 (s, 3H), 3.80 (s, 6H), 4.41 (d, 1H, J = 14.4 Hz), 4.45 (d, 1H, J = 14.4 Hz), 6.41-6.47 (m, 2H), 7.10-7.15 (m, 1H).

[0238]

(4) (Trans)-1-(2,4-dimethoxybenzyl)-4-methyl-5-oxopyrrolidine-3-carboxylic acid




To a solution of a mixture of (cis)-1-(2,4-dimethoxybenzyl)-4-methyl-5-oxopyrrolidine-3-carboxylic acid methyl ester and (trans)-1-(2,4-dimethoxybenzyl)-4-methyl-5-oxopyrrolidine-3-carboxylic acid methyl ester (14.2 g) in methanol (150 ml) was added sodium methoxide (about 5M solution in methanol, 28 ml), and the mixture was stirred at room temperature for 16 hours, and then heated to 55°C and stirred for 3 hours. To this reaction solution was added sodium methoxide (about 5M solution in methanol, 10 ml), and the mixture was stirred at 55°C for additional 2 hours. This reaction solution was cooled to room temperature, and then water (50 ml) was added thereto, and the mixture was stirred overnight. The reaction solution was concentrated under reduced pressure to remove methanol. To the resulting residue was added a 2M aqueous solution of hydrochloric acid (105 ml), and the mixture was extracted with ethyl acetate (300 ml). The resulting organic layer was washed with a saturated aqueous solution of sodium chloride, and then dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a crude product of the titled compound (14.0 g).

¹H-NMR (CDCl₃) δ: 1.32 (d, 3H, J = 6.6 Hz), 2.69-2.84 (m, 2H), 3.35-3.46 (m, 2H), 3.80 (s, 6H), 4.43 (s, 2H), 6.41-6.47 (m, 2H), 7.10-7.15 (m, 1H).

[0239]

(5) Optically active compound of (R)-4-benzyl-3-[(trans)-1-(2,4-dimethoxybenzyl)-4-methyl-5-oxopyrrolidine-3-carbonyl]-2-oxazolidinone




To a solution of the crude product of (trans)-1-(2,4-dimethoxybenzyl)-4-methyl-5-oxopyrrolidine-3-carboxylic acid (14.0 g) in chloroform (60 ml) were sequentially added (R)-4-benzyl-2-oxazolidinone (8.15 g), WSC·HCl (9.70 g), 4-dimethylaminopyridine (2.81 g), and the mixture was stirred overnight at room temperature. This reaction solution was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 4/1 to 1/3) to give the titled compound (low-polarity component 11.4 g, high-polarity component 10.1 g).

¹H-NMR (CDCl₃) low-polarity component δ: 1.31 (d, 3H, J = 7.3 Hz), 2.77-2.85 (m, 1H), 2.94-3.03 (m, 1H), 3.19-3.30 (m, 2H), 3.51-3.58 (m, 1H), 3.79 (s, 3H), 3.80 (s, 3H), 3.96-4.03 (m, 1H), 4.19-4.23 (m, 2H), 4.41 (d, 1H, J = 15.8 Hz), 4.49 (d, 1H, J = 15.8 Hz), 4.63-4.70 (m, 1H), 6.42-6.46 (m, 2H), 7.10-7.21 (m, 3H), 7.27-7.37 (m, 3H). high-polarity component δ: 1.26 (d, 3H, J = 7.3 Hz), 2.67-2.75 (m, 1H), 2.97-3.08 (m, 1H), 3.15-3.24 (m, 1H), 3.63-3.93 (m, 8H), 4.17-4.28 (m, 3H), 4.36-4.56 (m, 2H), 4.63-4.73 (m, 1H), 6.39-6.56 (m, 2H), 7.11-7.21 (m, 3H), 7.27-7.39 (m, 3H).

[0240]

(6) Optically active compound of (trans)-1-(2,4-dimethoxybenzyl)-4-methyl-5-oxopyrrolidine-3-carboxylic acid



To a solution of lithium hydroxide monohydrate (1.26 g) in water (30 ml) was added dropwise 30 wt% aqueous solution of hydrogen peroxide (7.0 ml) under ice-cooling, and the mixture was stirred for 10 minutes. To this reaction solution was added tetrahydrofuran 30 ml, and then a solution of the low-polarity component of (R)-4-benzyl-3-[(trans)-1-(2,4-dimethoxybenzyl)-4-methyl-5-oxopyrrolidine-3-carbonyl]-2-oxazolidinone (11.4 g) in tetrahydrofuran (90 ml) was added dropwise thereto, and then the mixture was stirred for additional 1 hour. To this reaction solution was added dropwise a solution of sodium hydrogen sulfite (7.18 g) in water (50 ml), and then the mixture was warmed to room temperature, and stirred for 1 hour. This mixture was extracted with ethyl acetate, and the resulting organic layer was washed with a saturated aqueous solution of sodium chloride, and then dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: chloroform/ethyl acetate = 1/1 to 1/2, chloroform/methanol = 5/1) to give a crude product of the titled compound (8.11 g). An analysis of the solid by HPLC analysis condition 1 showed that an isomer with shorter retention time was a main product.


An isomer with shorter retention time (retention time 5.7 minutes)

An isomer with longer retention time (retention time 6.5 minutes)

Alternatively, the titled compound can be prepared according to the following method.

[0241]

(6)-(1) Diastereomeric salt of an optically active compound of (trans)-1-(2,4-dimethoxybenzyl)-4-methyl-5-oxopyrrolidine-3-carboxylic acid and (1R,2R)-(-)-2-amino-1-(4-nitrophenyl)-1,3-propanediol




(Trans)-1-(2,4-dimethoxybenzyl)-4-methyl-5-oxopyrrolidine-3-carboxylic acid (1.0 g), (1R,2R)-(-)-2-amino-1-(4-nitrophenyl)-1,3-propanediol (434 mg), acetonitrile (10 ml) and water (0.9 ml) were mixed, and stirred at reflux to dissolve. This mixed solution was stirred at 50°C for 6 hours, and then stirred at room temperature overnight. A solid precipitated from this mixed solution was collected by filtration, washed with acetonitrile (6 ml), and dried under reduced pressure to give a solid 1 of the titled compound (572 mg). An analysis of the solid 1 by HPLC analysis condition 1 showed that an isomer with shorter retention time was a main product.

An isomer with shorter retention time (retention time 5.7 minutes)

An isomer with longer retention time (retention time 6.5 minutes)


¹H-NMR (DMSO-D₆) δ: 1.11 (d, 3H, J = 7.3 Hz), 2.43-2.53 (m, 1H), 2.55-2.64 (m, 1H), 2.85-2.92 (m, 1H), 3.14-3.22 (m, 2H), 3.27 (dd, 1H, J = 9.3, 9.3 Hz), 3.38 (dd, 1H, J = 11.3, 4.8 Hz), 3.73 (s, 3H), 3.76 (s, 3H), 4.19 (d, 1H, J = 14.5 Hz), 4.29 (d, 1H, J = 14.5 Hz), 4.73 (d, 1H, J = 6.0 Hz), 6.46 (dd, 1H, J = 8.1, 2.4 Hz), 6.54 (d, 1H, J = 2.4 Hz), 6.97 (d, 1H, J = 8.1 Hz), 7.61 (d, 2H, J = 8.5 Hz), 8.19 (d, 2H, J = 8.5 Hz).

An X ray crystal structural analysis of the crystal obtained by recrystallization of the solid 1 from methyl isobutyl ketone showed that this crystal was a diastereomeric salt of (3R,4R)-1-(2,4-dimethoxybenzyl)-4-methyl-5-oxopyrrolidine-3-carboxylic acid and (1R,2R)-(-)-2-amino-1-(4-nitrophenyl)-1,3-propanediol.



[0242]

(6)-(2) (3R,4R)-1-(2,4-Dimethoxybenzyl)-4-methyl-5-oxopyrrolidine-3-carboxylic acid



A diastereomeric salt of (3R,4R)-1-(2,4-dimethoxybenzyl)-4-methyl-5-oxopyrrolidine-3-carboxylic acid and (1R,2R)-(-)-2-amino-1-(4-nitrophenyl)-1,3-propanediol (10.0 g) was mixed with


ethyl acetate (38 ml), potassium hydrogen sulfate (2.4 g) and water (38 ml), and stirred at room temperature for 30 minutes. An organic layer was separated, and a separated aqueous layer was again extracted twice with ethyl acetate (15 ml). The combined organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. To the resulting residue were added 2-propanol (2.2 ml) and diisopropyl ether (45 ml), and the mixture was stirred. The insoluble substance was collected by filtration, washed with a small amount of diisopropyl ether, and dried under reduced pressure to give the titled compound (4.3 g). An analysis of this solid by HPLC analysis condition 1 showed that an isomer with shorter retention time was a main product.

An isomer with shorter retention time (retention time 5.7 minutes)

An isomer with longer retention time (retention time 6.5 minutes)

[0243]

(7) (3R,4R)-4-Methyl-5-oxopyrrolidine-3-carboxylic acid




To a crude product of (3R,4R)-1-(2,4-dimethoxybenzyl)-4-methyl-5-oxopyrrolidine-3-carboxylic acid (8.11 g) were added anisole (3.76 ml) and trifluoroacetic acid (40 ml), and the mixture was stirred at 80°C for 5 hours. This reaction solution was cooled to room temperature, and concentrated under reduced pressure. To the resulting residue was added diisopropyl ether, and the mixture was stirred. The insoluble substance was collected by filtration, and dried under reduced pressure to give a crude product of the titled compound (2.23 g). To the resulting solid was added ethanol (8 ml), and the mixture was heated at reflux to dissolve, and then the solution was stirred at room temperature. A solid precipitated from the mixed solution was collected by filtration, washed with a small amount of cooled ethanol, and dried under reduced pressure to give the titled compound (1.36 g).

¹H-NMR (DMSO-D₆) δ: 1.10 (d, 3H, J = 7.3 Hz), 2.40 (dq, 1H, J = 9.0, 7.3 Hz), 2.83 (ddd, 1H, J = 9.0, 8.6, 7.6 Hz), 3.22 (dd, 1H, J = 10.0, 7.6 Hz), 3.37 (dd, 1H, J = 10.0, 8.6 Hz), 7.66 (s, 1H), 12.58 (s, 1H).


Also, the high-polarity component of (R)-4-benzyl-3-[(trans)-1-(2,4-dimethoxybenzyl)-4-methyl-5-oxopyrrolidine-3-carbonyl]-2-oxazolidinone (10.1 g) was subjected to the similar reactions and post-treatment as described above to give (3S,4S)-4-methyl-5-oxopyrrolidine-3-carboxylic acid (1.53 g) as the optical isomer of the titled compound.

¹H-NMR (DMSO-D₆) δ: 1.10 (d, 3H, J = 7.3 Hz), 2.40 (dq, 1H, J = 9.0, 7.3 Hz), 2.83 (ddd, 1H, J = 9.0, 8.6, 7.6 Hz), 3.22 (dd, 1H, J = 10.0, 7.6 Hz), 3.37 (dd, 1H, J = 10.0, 8.6 Hz), 7.66 (s, 1H), 12.58 (s, 1H).



[0244]

[Preparation 2]: Example 1: Synthesis of ((3R,4R)-4-methyl-5-oxopyrrolidine-3-carboxylic acid [1-phenyl-5-(3-trifluoromethoxyphenyl)-1H-pyrazol-3-yl]amide)



5 (1) 1-Phenyl-1H-pyrazol-3-ylamine




To a solution of 1-phenyl-4,5-dihydro-1H-pyrazol-3-ylamine (50.0 g) in N,N-dimethylformamide (150 ml) and 1,4-dioxane (500 ml) was added 3,4,5,6-tetrachloro-1,4-benzoquinone (84.0 g) under ice-cooling over 20 minutes, and the mixture was stirred at room temperature for 4.5 hours. To the reaction mixture was added a 2M aqueous solution of sodium hydroxide (400 ml) under ice-cooling over 25 minutes, and the mixture was stirred at room temperature for 1 hour. The mixture was filtered through Celite to remove the insoluble substance, and eluted with ethyl acetate (250 ml×3), and then the filtrate was extracted with ethyl acetate (300 ml). The resulting organic layer was sequentially washed with water (300 ml) and a saturated aqueous solution of sodium chloride (300 ml). The separated aqueous layer was extracted twice with ethyl acetate (300 ml). To the combined organic layer were added anhydrous sodium sulfate (50 g) and silica gel (50 g), and the mixture was stirred at room temperature for 1 hour. This mixture was filtered with silica gel (100 g) on Celite, and subjected to elution with ethyl acetate (250 ml×3). The filtrate was concentrated, and diisopropyl ether (500 ml) was added to the resulting residue, and the mixture was stirred at room temperature. The insoluble substance was collected by filtration, washed twice with diisopropyl ether (100 ml), and dried under reduced pressure to give the titled compound (12.2 g).

¹H-NMR (CDCl₃) δ: 3.81 (br s, 2H), 5.85 (d, 1H, J = 2.4 Hz), 7.16-7.19 (m, 1H), 7.38-7.40 (m, 2H), 7.55-7.57 (m, 2H), 7.69 (d, 1H, J = 2.4 Hz).

[0245]

(2) 3-(2,5-Dimethylpyrrol-1-yl)-1-phenyl-1H-pyrazole




To a solution of 1-phenyl-1H-pyrazol-3-ylamine (32.7 g) in acetic acid (330 ml) was added 2,5-hexanedione (25 ml), and the mixture was stirred for 3 hours at reflux. This reaction solution was cooled to room temperature, and then acetic acid was concentrated under reduced pressure, and then toluene (100 ml) was added thereto, and the mixture was further concentrated. The resulting residue

was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate 100/1 to 20/1) to give the titled compound (41.8 g).

¹H-NMR (CDCl₃) δ: 2.19 (s, 6H), 5.90 (s, 2H), 6.39 (d, 1H, J = 2.4 Hz), 7.29-7.31 (m, 1H), 7.46-7.48 (m, 2H), 7.72-7.73 (m, 2H), 7.98 (d, 1H, J = 2.4 Hz).

[0246]

(3) 3-(2,5-Dimethylpyrrol-1-yl)-5-iodo-1-phenyl-1H-pyrazole




To a solution of 3-(2,5-dimethylpyrrol-1-yl)-1-phenyl-1H-pyrazole (14.6 g) in tetrahydrofuran (80 ml) cooled to -78°C was added n-butyllithium (1.6M solution in n-hexane, 24 ml) over 5 minutes, and then the mixture was stirred for 1 hour. To this mixture was added a solution of iodine (15.7 g) in tetrahydrofuran (30 ml) over 10 minutes, and the mixture was stirred for additional 1.5 hours. To this mixture were sequentially added water (5 ml) and a saturated aqueous solution of sodium hydrogen sulfite (30 ml), and then the mixture was warmed to 0°C, and additional water (150 ml) was added thereto, and the mixture was stirred at room temperature for 1 hour. A precipitated solid was collected by filtration, washed with water and a small amount of methanol, and dried under reduced pressure to give the titled compound (13.0 g).

¹H-NMR (CDCl₃) δ: 2.19 (s, 6H), 5.88 (s, 2H), 6.55 (s, 1H), 7.44-7.53 (m, 3H), 7.57-7.60 (m, 2H).

[0247]

(4) 5-Iodo-1-phenyl-1H-pyrazol-3-ylamine




To a solution of 3-(2,5-dimethylpyrrol-1-yl)-5-iodo-1-phenyl-1H-pyrazole (12.2 g) in ethanol/water (2/1, 180 ml) were sequentially added hydroxylammonium chloride (46.6 g) and triethylamine (9.3 ml), and the mixture was stirred at 95°C for 72 hours. This reaction mixture was cooled to room temperature, and concentrated under reduced pressure. To the resulting residue was added a 8M aqueous solution of sodium hydroxide (80 ml), and then the mixture was extracted with ethyl acetate (50 ml). This organic layer was sequentially washed with water and a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 3/1) to give the titled compound (9.1 g).

¹H-NMR (CDCl₃) δ: 3.77 (br s, 2H), 6.01 (s, 1H), 7.35-7.51 (m, 5H).

[0248]

(5) 1-Phenyl-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazol-3-ylamine




To a solution of 5-iodo-1-phenyl-1H-pyrazol-3-ylamine (605 mg) in ethylene glycol dimethyl ether (12 ml) were sequentially added 3-(trifluoromethoxy)phenyl boronic acid (481 mg), a 2M aqueous solution of sodium carbonate (6 ml), tricyclohexylphosphine (119 mg), palladium acetate (48 mg), and the mixture was stirred for 2 hours at reflux. This reaction mixture was cooled to room temperature, and a saturated aqueous solution of sodium hydrogen carbonate was added thereto, and then the mixture was filtered through Celite. The filtrate was extracted with ethyl acetate. The resulting organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 2/1) to give the titled compound (298 mg).

¹H-NMR (CDCl₃) δ: 3.79 (br s, 2H), 5.95 (s, 1H), 7.04 (s, 1H), 7.13-7.34 (m, 8H).

[0249]

(6) (3R,4R)-4-Methyl-5-oxopyrrolidine-3-carboxylic acid [1-phenyl-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazol-3-yl]amide



To a solution of 1-phenyl-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazol-3-ylamine (79 mg) in N,N-dimethylformamide (0.8 ml) were sequentially added (3R,4R)-4-methyl-5-oxopyrrolidine-3-carboxylic acid (46 mg) prepared in Preparation 1, HOBt·H₂O (57 mg) and WSC·HCl (71 mg), and the mixture was stirred at room temperature for 2 hours. To this reaction solution were added water and a saturated aqueous solution of sodium hydrogen carbonate, and the mixture was extracted with ethyl acetate. The separated organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 10/1) to give the titled compound (63 mg).

¹H-NMR (DMSO-D₆) δ: 1.08 (d, 3H, J = 7.3 Hz), 2.55 (ddd, 1H, J = 9.3, 8.8, 8.8 Hz), 3.04 (dq, 1H, J = 9.3, 7.3 Hz), 3.23 (dd, 1H, J = 8.8, 8.8 Hz), 3.45 (dd, 1H, J = 8.8, 8.8 Hz), 7.03 (s, 1H), 7.09 (s, 1H), 7.26 (t, 2H, J = 6.7 Hz), 7.35-7.45 (m, 5H), 7.52 (t, 1H, J = 8.0 Hz), 7.68 (s, 1H), 10.99 (s, 1H).

[0250]


(7) Crystals of (3R,4R)-4-methyl-5-oxopyrrolidine-3-carboxylic acid [1-phenyl-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazol-3-yl]amide

The titled compound was dissolved in ethyl acetate at 90°C, and recrystallized by an addition of heptane thereto at the same temperature to give a crystal of the titled compound. Furthermore, the


corresponding crystal as above was suspended in water at 100°C to give another crystal of the titled compound.

[0251]

[Preparation 3]: Example 2: Synthesis of ((3R,4R)-4-methyl-5-oxopyrrolidine-3-carboxylic acid [1-(4-fluorophenyl)-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazol-3-yl]amide)



(1) Mixture of (E)-3-(3-(trifluoromethoxy)phenyl)acrylonitrile and (Z)-3-(3-(trifluoromethoxy)phenyl)acrylonitrile




To a suspension of (cyanomethyl) triphenylphosphonium chloride (2.89 g) in tetrahydrofuran (10 ml) was added potassium tert-butoxide (942 mg) under ice-cooling, and the mixture was stirred at room temperature for 1 hour. To this reaction solution was added 3-trifluoromethoxybenzaldehyde (1.33 g) under ice-cooling, and then the mixture was stirred at room temperature overnight. To this reaction solution was added n-hexane (20 ml), and the insoluble substance was filtered off, and then the filtrate was concentrated. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/0 to 10/1) to give the titled compound (1.28 g).

¹H-NMR (CDCl₃) (E)-isomer δ: 5.92 (1H, d, J = 16.8 Hz), 7.27-7.49 (5H, m).

(Z)-isomer δ: 5.55 (1H, d, J = 12.0 Hz), 7.12 (1H, d, J = 12.0 Hz), 7.35-7.52 (2H, m), 7.60 (1H, s), 7.79 (1H, d, J = 8.0 Hz).

[0252]

(2) 1-(4-Fluorophenyl)-5-(3-(trifluoromethoxy)phenyl)-4,5-dihydro-1H-pyrazol-3-ylamine




To a solution of a mixture of (E)-3-(3-(trifluoromethoxy)phenyl)acrylonitrile and (Z)-3-(3-(trifluoromethoxy)phenyl)acrylonitrile (1.28 g) in ethanol (13 ml) were sequentially added 4-fluorophenylhydrazine hydrochloride (979 mg) and sodium ethoxide (20 wt% in ethanol, 5.0 ml), and the mixture was stirred overnight at reflux. This reaction solution was concentrated under reduced pressure, and then water was added thereto, and the mixture was extracted with ethyl acetate. This organic layer was sequentially washed with water and a saturated aqueous solution of sodium chloride, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 2/1 to 1/1) to give the titled compound

(1.29 g).

¹H-NMR (CDCl₃) δ: 2.79 (dd, 1H, J = 16.3, 8.8 Hz), 3.41 (dd, 1H, J = 16.3, 11.0 Hz), 3.98 (s, 2H), 4.83 (dd, 1H, J = 11.0, 8.8 Hz), 6.72-6.87 (m, 4H), 7.14 (d, 1H, J = 7.9 Hz), 7.25 (s, 1H), 7.32 (d, 1H, J = 7.7 Hz), 7.38 (dd, 1H, J = 7.9, 7.7 Hz).

5 [0253]

(3) 1-(4-Fluorophenyl)-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazol-3-ylamine




10 To a solution of 1-(4-fluorophenyl)-5-(3-(trifluoromethoxy)phenyl)-4,5-dihydro-1H-pyrazol-3-ylamine (1.29 g) in toluene (10 ml) was added activated carbon (pH5 to 8, 0.64 g), and the mixture was stirred for 2 hours under oxygen atmosphere under cooling to reflux. After cooling to room temperature, the activated carbon was filtered off through Celite eluting with ethyl acetate. This filtrate was concentrated under reduced pressure. To the resulting residue, a mixed solvent of diisopropyl ether /n-hexane was added and the mixture was stirred at room temperature. The insoluble substance was collected by filtration, and then dried under reduced pressure to give the titled compound (652 mg).

15 ¹H-NMR (CDCl₃) δ: 3.78 (bs, 1H), 5.95 (s, 1H), 6.97-7.05 (m, 3H), 7.13-7.21 (m, 4H), 7.334 (t, 1H, J = 8.0 Hz).

[0254]

20 (4) (3R,4R)-4-Methyl-5-oxopyrrolidine-3-carboxylic acid [1-(4-fluorophenyl)-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazol-3-yl]amide




25 To a solution of 1-(4-fluorophenyl)-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazol-3-ylamine (90 mg) in N,N-dimethylformamide (0.9 ml) were added (3R,4R)-4-methyl-5-oxopyrrolidine-3-carboxylic acid (50 mg) prepared in Preparation 1, HOBt·H₂O (61 mg) and WSC·HCl (77 mg), and the mixture was stirred at room temperature for 2 hours. To this mixture were added water and a saturated aqueous solution of sodium hydrogen carbonate, and the mixture was extracted with ethyl acetate. The separated organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 10/1) to give the titled compound (80 mg).

30 ¹H-NMR (DMSO-D₆) δ: 1.08 (d, 3H, J = 7.1 Hz), 2.54 (dq, 1H, J = 9.3, 7.1 Hz), 3.04 (ddd, 1H, J =


9.3, 9.2, 8.4 Hz), 3.23 (dd, 1H, J = 9.2, 8.6 Hz), 3.44 (dd, 1H, J = 8.6, 8.4 Hz), 7.03 (s, 1H), 7.11 (s, 1H), 7.25-7.38 (m, 6H), 7.53 (dd, 1H, J = 8.2, 7.9 Hz), 7.68 (s, 1H), 10.98 (s, 1H).

[0255]

[Preparation 4]: Example 3: Synthesis of ((3R, 4R)-4-methyl-5-oxopyrrolidine-3-carboxylic acid [1-(3-fluorophenyl)-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazol-3-yl]amide)



(1) 3-(2,5-Dimethylpyrrol-1-yl)-1H-pyrazole




To a solution of 3-aminopyrazole (100 g) in acetic acid (1000 ml) was added 2,5-hexanedione (148ml), and the mixture was stirred for 2.4 hours at reflux. This reaction solution was cooled to room temperature, water (1000 ml) was added thereto, and the precipitated solid was collected by filtration. The resulting solid was washed with water (1000 ml), and dried under reduced pressure to give the titled compound (183 g).

¹H-NMR (DMSO-D₆) δ: 2.00 (s, 6H), 5.74 (s, 2H), 6.27 (d, 1H, J = 2.1 Hz), 7.84 (d, 1H, J = 2.1 Hz), 12.90 (s, 1H).

[0256]

(2) 3-(2,5-Dimethylpyrrol-1-yl)-1-(3-fluorophenyl)-1H-pyrazole




To a solution of 3-(2,5-dimethylpyrrol-1-yl)-1H-pyrazole (6.8 g) in N-methyl-2-pyrrolidone (68 ml) were sequentially added cesium carbonate (27.4 g), 8-quinolinol (1.2 g), copper oxide (I) (0.6 g) and 1-fluoro-3-iodobenzene (7.4 ml), and the mixture was stirred at 110°C for 2 hours. This reaction mixture was cooled to room temperature, and toluene (80 ml) was added thereto, and then the mixture was filtered through Celite. After adding a 1M aqueous solution of sodium hydroxide (80 ml) to the filtrate, the mixture was extracted with toluene. The resulting organic layer was sequentially washed with water and a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 9/1) to give the titled compound (10.4 g).

¹H-NMR (CDCl₃) δ: 2.19 (s, 6H), 5.90 (s, 2H), 6.41 (d, 1H, J = 2.6 Hz), 6.97-7.02 (m, 1H), 7.39-7.45 (m, 1H), 7.48-7.49 (m, 1H), 7.49-7.52 (m, 1H), 7.97 (d, 1H, J = 2.6 Hz).

[0257]

(3) 3-(2,5-Dimethylpyrrol-1-yl)-1-(3-fluorophenyl)-5-iodo-1H-pyrazole




To a solution of 3-(2,5-dimethylpyrrol-1-yl)-1-(3-fluorophenyl)-1H-pyrazole (10.4 g) in tetrahydrofuran (100 ml) cooled to -78°C was added n-butyllithium (1.6M solution in n-hexane, 31 ml) over 10 minutes, and then the mixture was stirred for 30 minutes. To the mixture was added dropwise a solution of iodine (12.4 g) in tetrahydrofuran (20 ml) over 10 minutes, and the mixture was stirred for additional 30 minutes. To the mixture were sequentially added a 20 wt% aqueous solution of sodium thiosulfate and a saturated aqueous ammonia, and then the mixture was warmed to room temperature, and extracted with ethyl acetate. The separated organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. To the resulting residue was added a mixed solvent of n-hexane/ethyl acetate (2/1), and the mixture was stirred. The insoluble substance was collected by filtration, and dried under reduced pressure to give the titled compound (4.5 g).

¹H-NMR (CDCl₃) δ: 2.19 (s, 6H), 5.88 (s, 2H), 6.56 (s, 1H), 7.15-7.19 (m, 1H), 7.35-7.37 (m, 1H), 7.41-7.50 (m, 2H).

[0258]

(4) 1-(3-Fluorophenyl)-5-iodo-1H-pyrazol-3-ylamine




To a solution of 3-(2,5-dimethylpyrrol-1-yl)-1-(3-fluorophenyl)-5-iodo-1H-pyrazole (4.5 g) in ethanol/water (2/1, 81 ml) were sequentially added hydroxylammonium chloride (16.3 g) and triethylamine (3.3 ml), and the mixture was stirred at 95°C for 17 hours. This reaction mixture was cooled to room temperature and concentrated under reduced pressure. To the resulting residue was added an 8M aqueous solution of sodium hydroxide, and the mixture was extracted with ethyl acetate. This organic layer was sequentially washed with water and a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 2/1) to give the titled compound (3.2 g).

¹H-NMR (CDCl₃) δ: 3.78 (br s, 2H), 6.03 (s, 1H), 7.05-7.09 (m, 1H), 7.26-7.29 (m, 1H), 7.32-7.43 (m, 2H).

[0259]

(5) 1-(3-Fluorophenyl)-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazol-3-ylamine




To a solution of 1-(3-fluorophenyl)-5-iodo-1H-pyrazol-3-ylamine (263 mg) in ethylene glycol dimethyl ether (5 ml) were sequentially added 3-(trifluoromethoxy)phenylboronic acid (197 mg), a 2M aqueous solution of sodium carbonate (2.5 ml), tricyclohexylphosphine (49 mg), palladium acetate (20 mg), and the mixture was stirred for 2 hours at reflux. This reaction mixture was cooled to room temperature, a saturated aqueous solution of sodium hydrogen carbonate was added thereto, and then the mixture was filtered through Celite. The filtrate was extracted with ethyl acetate, the resulting organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 2/1) to give the titled compound (234 mg).

¹H-NMR (CDCl₃) δ: 3.82 (br s, 2H), 5.96 (s, 1H), 6.92-7.02 (m, 3H), 7.05-7.07 (m, 1H), 7.17-7.38 (m, 4H).

[0260]

(6) (3R,4R)-4-Methyl-5-oxopyrrolidine-3-carboxylic acid [1-(3-fluorophenyl)-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazol-3-yl]amide




To a solution of 1-(3-fluorophenyl)-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazol-3-ylamine (68 mg) in N,N-dimethylformamide (0.7 ml) were sequentially added (3R,4R)-4-methyl-5-oxopyrrolidine-3-carboxylic acid (38 mg) prepared in Preparation 1, HOBT·H₂O (46 mg) and WSC·HCl (58 mg), and the mixture was stirred at room temperature for 2 hours. To this reaction solution were added water and a saturated aqueous solution of sodium hydrogen carbonate, and the mixture was extracted with ethyl acetate. The separated organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 10/1) to give the titled compound (33 mg).


¹H-NMR (DMSO-D₆) δ: 1.08 (d, 3H, J = 7.3 Hz), 2.54 (dq, 1H, J = 9.3, 7.3 Hz), 3.04 (ddd, 1H, J = 9.3, 8.9, 8.5 Hz), 3.23 (dd, 1H, J = 8.9, 8.7 Hz), 3.45 (dd, 1H, J = 8.7, 8.5 Hz), 7.04 (s, 1H), 7.05-7.07 (m, 1H), 7.14-7.18 (m, 2H), 7.22-7.26 (m, 1H), 7.39-7.48 (m, 3H), 7.56 (t, 1H, J = 8.0 Hz), 7.68 (s, 1H), 11.03 (s, 1H).

[0261]

[Preparation 5]: Synthesis of (R) and (S)-5-oxopyrrolidine-3-carboxylic acid




(1) 1-(2,4-Dimethoxybenzyl)-5-oxopyrrolidine-3-carboxylic acid



To a solution of itaconic acid (65.3 g) in toluene (400 ml) was added a solution of 2,4-dimethoxybenzylamine (84.0 g) in toluene (100 ml), and the mixture was stirred for 15 hours at reflux. This reaction solution was cooled to room temperature, and concentrated under reduced pressure. To the resulting residue was added diethyl ether (600 ml), and the mixture was stirred. The precipitated solid was collected by filtration, washed with diethyl ether and a small amount of ethyl acetate, and then dried under reduced pressure to give the titled compound (127 g).

¹H-NMR (CDCl₃) δ: 2.66-2.82 (m, 2H), 3.20 (dt, 1H, J = 17.3, 7.6 Hz), 3.49-3.53 (m, 2H), 3.80 (s, 6H), 4.40 (d, 1H, J = 14.6 Hz), 4.46 (d, 1H, J = 14.6 Hz), 6.42-6.47 (m, 2H), 7.14 (t, 1H, J = 4.5 Hz). [0262]


(2) Optically active compound of (R)-4-benzyl-3-[1-(2,4-dimethoxybenzyl)-5-oxopyrrolidine-3-carboxyl]-2-oxazolidinone



To a solution of 1-(2,4-dimethoxybenzyl)-5-oxopyrrolidine-3-carboxylic acid (31.7 g) in chloroform (300 ml) were sequentially added (R)-4-benzyl-2-oxazolidinone (20.0 g), WSC·HCl (23.8 g) and 4-dimethylaminopyridine (6.90 g) under ice-cooling, the mixture was stirred for 15 minutes, warmed to room temperature and stirred for 18 hours. This reaction mixture was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/1 to 1/9) to give the titled compound (low-polarity component 26.4 g, high-polarity component 13.3 g).

¹H-NMR (CDCl₃) low-polarity component δ: 2.72-2.83 (m, 3H), 3.27 (dd, 1H, J = 13.4, 3.5 Hz), 3.46-3.57 (m, 2H), 3.80 (s, 3H), 3.80 (s, 3H), 4.15-4.29 (m, 3H), 4.40-4.48 (m, 2H), 4.61-4.69 (m, 1H), 6.42-6.47 (m, 2H), 7.12-7.21 (m, 3H), 7.27-7.37 (m, 3H). high-polarity component δ: 2.62-2.72 (m, 2H), 2.86 (dd, 1H, J = 16.9, 5.8 Hz), 3.17 (dd, 1H, J = 13.4, 3.2 Hz), 3.40 (dd, 1H, J = 10.4, 5.1 Hz), 3.66 (dd, 1H, J = 10.4, 8.8 Hz), 3.72 (s, 3H), 3.81 (s, 3H), 4.16-4.26 (m, 3H), 4.40 (d, 1H, J = 14.6 Hz), 4.50 (d, 1H, J = 14.6 Hz), 4.61-4.69 (m, 1H), 6.40-6.45 (m, 2H), 7.11-7.17 (m, 3H), 7.27-7.34 (m, 3H). [0263]

(3) Optically active compound of 1-(2,4-dimethoxybenzyl)-5-oxopyrrolidine-3-carboxylic acid




To a solution of lithium hydroxide monohydrate (3.29 g) in water (50 ml) was added dropwise 30 wt% aqueous solution of hydrogen peroxide (17.5 ml) under cooling with an ice--sodium chloride bath, and the mixture was stirred for 15 minutes. To this reaction solution was added tetrahydrofuran (30 ml), and then a solution of the low-polarity component of (R)-4-benzyl-3- [1-(2,4-dimethoxybenzyl)-5-oxopyrrolidine-3-carbonyl]-2-oxazolidinone (26.4 g) in tetrahydrofuran (150 ml) was added dropwise thereto, and the mixture was stirred for additional 1 hour. To this reaction solution was slowly added a solution of sodium hydrogen sulfite (18.7 g) in water (60 ml), and the mixture was warmed to room temperature by removing the bath, and then extracted with ethyl acetate (100 ml). The resulting organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: chloroform/ethyl acetate = 1/1 to 1/2, chloroform/methanol = 5/1) to give the titled compound (14.3 g). An analysis of this solid by HPLC analysis condition 2 showed that an isomer with shorter retention time was a main product.

An isomer with shorter retention time (retention time 4.1 minutes)

An isomer with longer retention time (retention time 4.6 minutes)

¹H-NMR (CDCl₃) δ: 2.66-2.82 (m, 2H), 3.20 (dt, 1H, J = 17.3, 7.6 Hz), 3.49-3.53 (m, 2H), 3.80 (s, 6H), 4.40 (d, 1H, J = 14.6 Hz), 4.46 (d, 1H, J = 14.6 Hz), 6.42-6.47 (m, 2H), 7.14 (t, 1H, J = 4.5 Hz). [0264]

(4) Optically active compound of 5-oxopyrrolidine-3-carboxylic acid




To an optically active compound of 1-(2,4-dimethoxybenzyl)-5-oxopyrrolidine-3-carboxylic acid (14.3 g) were added anisole (8.4 ml) and trifluoroacetic acid (100 ml), and the mixture was stirred at 80°C for 5 hours. This reaction solution was cooled to room temperature, and concentrated under reduced pressure. To the resulting residue was added diisopropyl ether (100 ml), and the mixture was stirred at room temperature. The insoluble substance was collected by filtration, washed with diisopropyl ether, and dried under reduced pressure to give the titled compound (4.96 g).


¹H-NMR (DMSO-D₆) δ: 2.26-2.40 (m, 2H), 3.18-3.28 (m, 1H), 3.30-3.35 (m, 1H), 3.40-3.47 (m, 1H), 7.64 (s, 1H), 12.58 (s, 1H); [α]_D²⁵ +37 (c 0.10, CH₃OH).

It was shown that the resulting compound was (R)-5-oxopyrrolidine-3-carboxylic acid by

comparing a specific rotation of the compound with those described in the literature (Tetrahedron: Asymmetry 12 (2001) 3241-3249).




Also, the high-polarity component of (R)-4-benzyl-3-[1-(2,4-dimethoxybenzyl)-5-oxopyrrolidine-3-carbonyl]-2-oxazolidinone was subjected to the same reactions and post-treatment as described above to give (S)-5-oxopyrrolidine-3-carboxylic acid (2.83 g) as the optical isomer .
¹H-NMR (DMSO-D₆) δ: 2.26-2.40 (m, 2H), 3.18-3.28 (m, 1H), 3.30-3.35 (m, 1H), 3.40-3.47 (m, 1H), 7.64 (s, 1H), 12.58 (s, 1H); [α]_D²⁵ -40 (c 0.10, CH₃OH).




[0265]

[Preparation 6]: Example 4: Synthesis of ((R)-5-oxopyrrolidine-3-carboxylic acid [1-phenyl-5-(3-trifluoromethoxyphenyl)-1H-pyrazol-3-yl]amide)



(1) (R)-5-Oxopyrrolidine-3-carboxylic acid [1-phenyl-5-(3-trifluoromethoxyphenyl)-1H-pyrazol-3-yl]amide




To a solution of 1-phenyl-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazol-3-ylamine (40 mg) prepared according to the same procedures as Preparation 2 in N,N-dimethylformamide (0.3 ml) were sequentially added (R)-5-oxopyrrolidine-3-carboxylic acid (24 mg) prepared in Preparation 5, HOBT·H₂O (29 mg) and WSC·HCl (36 mg), and the mixture was stirred at room temperature overnight. To this reaction solution were added a saturated aqueous solution of sodium hydrogen carbonate and water, and the mixture was stirred. The insoluble substance was collected by filtration, washed with water, and dried under reduced pressure to give the titled compound (45 mg).


¹H-NMR (DMSO-D₆) δ: 2.35-2.42 (m, 2H), 3.31-3.54 (m, 3H), 7.01 (s, 1H), 7.08 (s, 1H), 7.22-7.28 (m, 2H), 7.32-7.46 (m, 5H), 7.52 (t, 1H, J = 8.0 Hz), 7.64 (s, 1H), 10.90 (s, 1H).

[0266]

[Preparation 7]: Synthesis of optically active compound of 6-oxopiperidine-3-carboxylic acid



(1) 2-Methylenepentanedioic acid dimethyl ester




To a solution of 2,8,9-triisopropyl-2,5,8,9-tetraaza-1-phosphabicyclo(3,3,3) undecane (4.62 g) in tetrahydrofuran (180 ml) was added dropwise a solution of methyl acrylate (40.5 ml) in tetrahydrofuran (20 ml) at room temperature, and then the mixture was stirred overnight. This reaction solution was concentrated under reduced pressure, and then diisopropyl ether, n-hexane and chloroform were added to the resulting residue, and the mixture was again concentrated under reduced pressure. To the precipitated solid was added diisopropyl ether, the mixture was stirred, and then the insoluble substance was filtered off. The filtrate was concentrated under reduced pressure, and the resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 9/1 to 4/1) to give the titled compound (19.2 g).

¹H-NMR (CDCl₃) δ: 2.53 (td, 2H, J = 7.6, 1.3 Hz), 2.65 (t, 2H, J = 7.6 Hz), 3.67 (s, 3H), 3.76 (s, 3H), 5.61 (q, 1H, J = 1.3 Hz), 6.20 (d, 1H, J = 1.3 Hz).

[0267]

(2) 1-(2,4-Dimethoxybenzyl)-6-oxopiperidine-3-carboxylic acid methyl ester




To a solution of 2-methylenepentanedioic acid dimethyl ester (19.2 g) in toluene (195 ml) was added 2,4-dimethoxybenzylamine (16.6 ml), and the mixture was stirred for 3 days at reflux. This reaction solution was cooled to room temperature, and then washed with a 1M aqueous solution of hydrochloric acid, water and a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a crude product of the titled compound (26.8 g).

¹H-NMR (CDCl₃) δ: 1.93-2.05 (m, 1H), 2.08-2.17 (m, 1H), 2.40-2.50 (m, 1H), 2.52-2.62 (m, 1H), 2.74-2.83 (m, 1H), 3.45 (d, 2H, J = 7.4 Hz), 3.68 (s, 3H), 3.81 (s, 3H), 3.81 (s, 3H), 4.54 (d, 1H, J = 14.6 Hz), 4.62 (d, 1H, J = 14.6 Hz), 6.44-6.48 (m, 2H), 7.15-7.19 (m, 1H).

[0268]

(3) 1-(2,4-Dimethoxybenzyl)-6-oxopiperidine-3-carboxylic acid




To a solution of the crude product of 1-(2,4-dimethoxybenzyl)-6-oxopiperidine-3-carboxylic acid methyl ester (26.8 g) in methanol/tetrahydrofuran (1/1, 174 ml) was added a 1M aqueous solution of sodium hydroxide (174 ml) at room temperature, and the mixture was stirred overnight.

To this reaction solution was added a 1M aqueous solution of hydrochloric acid (174 ml), the mixture was stirred. The precipitated solid was collected by filtration, washed with water, and dried under reduced pressure to give the titled compound (15.1 g).

¹H-NMR (DMSO-*d*₆) δ: 1.77-1.90 (m, 1H), 1.95-2.06 (m, 1H), 2.25-2.41 (m, 2H), 2.75-2.86 (m, 1H), 3.31 (d, 2H, *J* = 7.1 Hz), 3.74 (s, 3H), 3.78 (s, 3H), 4.34 (d, 1H, *J* = 15.2 Hz), 4.43 (d, 1H, *J* = 15.2 Hz), 6.47 (dd, 1H, *J* = 8.4, 2.4 Hz), 6.56 (d, 1H, *J* = 2.4 Hz), 7.00 (d, 1H, *J* = 8.4 Hz), 12.56 (br s, 1H). [0269]


(4) Optically active compound of 5-((*R*)-4-benzyl-2-oxooxazolidine-3-carbonyl)-1-(2,4-dimethoxybenzyl)piperidin-2-one



To a solution of 1-(2,4-dimethoxybenzyl)-6-oxopiperidine-3-carboxylic acid (3.39 g) in chloroform (15 ml) were sequentially added (*R*)-4-benzyl-2-oxazolidinone (2.04 g), WSC·HCl (2.44 g) and 4-dimethylaminopyridine (706 mg) at room temperature, and the mixture was stirred overnight. This reaction mixture was purified by silica gel column chromatography (eluent: chloroform/ethyl acetate = 4/1 to 1/1) to give the titled compound (low-polarity component 3.37 g, high-polarity component 2.72 g).

¹H-NMR (CDCl₃) low-polarity component δ: 1.91-2.02 (m, 1H), 2.14-2.24 (m, 1H), 2.50-2.66 (m, 2H), 2.73-2.81 (m, 1H), 3.18-3.25 (m, 1H), 3.34-3.41 (m, 1H), 3.51-3.60 (m, 1H), 3.78 (s, 3H), 3.80 (s, 3H), 3.84-3.93 (m, 1H), 4.05-4.20 (m, 2H), 4.53-4.65 (m, 3H), 6.42-6.47 (m, 2H), 7.14-7.38 (m, 6H). high-polarity component ¹H-NMR (CDCl₃) δ: 1.92-2.14 (m, 2H), 2.47-2.67 (m, 3H), 3.14-3.23 (m, 1H), 3.44-3.59 (m, 2H), 3.77 (s, 3H), 3.81 (s, 3H), 3.91-4.01 (m, 1H), 4.14-4.26 (m, 2H), 4.54-4.68 (m, 3H), 6.42-6.49 (m, 2H), 7.15-7.21 (m, 3H), 7.25-7.35 (m, 3H). [0270]

(5) Optically active compound of 1-(2,4-dimethoxybenzyl)-6-oxopiperidine-3-carboxylic acid




To a solution of lithium hydroxide monohydrate (316 mg) in water (10 ml) was added dropwise a 30 wt% aqueous solution of hydrogen peroxide (1.8 ml) under ice-cooling and the mixture was stirred for 10 minutes. To this reaction solution was added tetrahydrofuran (10 ml), and then a solution of the low-polarity component of 5-((R)-4-benzyl-2-oxooxazolidine-3-carbonyl)-1-(2,4-dimethoxybenzyl)piperidin-2-one (3.37 g) in tetrahydrofuran (20 ml) was added dropwise thereto, and the mixture was stirred for additional 1 hour. To this reaction solution was added an aqueous solution of sodium hydrogen sulfite (1.81 g) was added dropwise, and then the mixture was warmed to room temperature, and stirred for 30 minutes. This mixed solution was extracted with ethyl acetate, and the resulting organic layer was washed with a saturated aqueous solution of sodium chloride, and then dried over anhydrous sodium sulfate, and concentrated under reduced pressure. To the resulting residue was added ethyl acetate, and the mixture was stirred at room temperature. The insoluble substance was collected by filtration, and dried under reduced pressure to give the titled compound (1.36 g). An analysis of this solid by HPLC analysis condition 2 showed that an isomer with shorter retention time was a main product.

An isomer with shorter retention time (retention time 4.5 minutes)

An isomer with longer retention time (retention time 6.6 minutes)

[0271]

(6) Optically active compound of 6-oxopiperidine-3-carboxylic acid



To an optically active compound of 1-(2,4-dimethoxybenzyl)-6-oxopiperidine-3-carboxylic acid (1.36 g) were added anisole (758 μ l) and trifluoroacetic acid (10 ml), and the mixture was stirred at 80°C for 6 hours. This reaction solution was cooled to room temperature, and then concentrated under reduced pressure. To the resulting residue was added diisopropyl ether, and the mixture was stirred at room temperature. The insoluble substance was collected by filtration, and dried under reduced pressure to give the titled compound (628 mg).


$^1\text{H-NMR}$ (DMSO- D_6) δ : 1.75-1.88 (m, 1H), 1.91-2.01 (m, 1H), 2.11-2.24 (m, 2H), 2.66-2.73 (m, 1H), 3.21-3.32 (m, 2H), 7.45 (s, 1H), 12.51 (br s, 1H).

Also, the high-polarity component of 5-((R)-4-benzyl-2-oxoxazolidinone-3-carbonyl)-1-(2,4-dimethoxybenzyl)piperidin-2-on was subjected to the same reactions and post-treatment as described above to give an optical isomer of the titled compound (485 mg).


¹H-NMR (DMSO-D₆) δ: 1.75-1.88 (m, 1H), 1.91-2.01 (m, 1H), 2.11-2.24 (m, 2H), 2.66-2.73 (m, 1H), 3.21-3.32 (m, 2H), 7.45 (s, 1H), 12.51 (br s, 1H).

[0272]

[Preparation 8]: Example 5: Synthesis of optically active compound of (6-oxopiperidine-3-carboxylic acid [1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-yl]amide)



(1) 1-Bromo-3-propylbenzene




To a solution of 3-bromopropiophenone (25 g) in trifluoroacetic acid (250 ml) was added dropwise triethylsilane (94 ml) under ice-cooling over 20 minutes, and the mixture was stirred for additional 20 minutes. This reaction solution was heated to 80°C, and stirred overnight. This reaction solution was cooled to room temperature, concentrated under reduced pressure, and again concentrated under reduced pressure after adding toluene thereto. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane) to give the titled compound (16.6 g).

¹H-NMR (CDCl₃) δ: 0.95 (t, 3H, J = 7.4 Hz), 1.64 (tq, 2H, J = 7.7, 7.4 Hz), 2.57 (t, 2H, J = 7.7 Hz), 7.08-7.13 (m, 1H), 7.13-7.18 (m, 1H), 7.30-7.35 (m, 2H).

[0273]

(2) 3-Propylbenzaldehyde




To a solution of 1-bromo-3-propylbenzene (16.6 g) in tetrahydrofuran (83 ml) cooled to -78°C was added dropwise n-butyllithium (2.66M solution in n-hexane, 34 ml) over 9 minutes, and then the mixture was stirred for 1 hour. To this reaction solution was added dropwise N,N-dimethylformamide (7.7 ml) over 6 minutes, and the mixture was stirred for additional 20 minutes. To this reaction solution was added dropwise a 4M solution of hydrogen chloride in 1,4-dioxane (23 ml) over 5 minutes, and then the mixture was stirred after warming to room temperature. To this mixed solution was added water (83 ml), followed by a 6M aqueous solution of hydrochloric acid (12 ml), and then the mixture was extracted with n-hexane (80 ml). The resulting organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure to give a crude product of the titled compound (14.0

g).

¹H-NMR (CDCl₃) δ: 0.95 (t, 3H, J = 7.3 Hz), 1.68 (tq, 2H, J = 7.7, 7.3 Hz), 2.67 (t, 2H, J = 7.7 Hz), 7.42-7.47 (m, 2H), 7.67-7.72 (m, 2H), 9.98-10.01 (m, 1H).

[0274]

5 (3) Mixture of (E)-3-(3-propylphenyl)acrylonitrile and (Z)-3-(3-propylphenyl)acrylonitrile




To a suspension of (cyanomethyl)triphenylphosphonium chloride (34 g) in tetrahydrofuran (110 ml) was added portionwise potassium tert-butoxide (13 g) under ice-cooling, and then the mixture was stirred for 1 hour. To this reaction solution was added dropwise a solution of a crude product of 3-propyl benzaldehyde (14.0 g) in tetrahydrofuran (30 ml) over 7 minutes under ice-cooling, and then the mixture was stirred at room temperature overnight. To this mixture was added n-hexane (280 ml), and the mixture was stirred for 1 hour under ice-cooling, and then the insoluble substance was filtered off through Celite, and eluted with n-hexane (840 ml). To the resulting residue by concentrating the filtrate was added a mixed solvent of n-hexane/ethyl acetate (5/1, 120 ml), and the mixture was stirred. The insoluble substance was filtered off through Celite, and eluted with a mixed solvent of n-hexane/ethyl acetate (10/1). The filtrate was concentrated, and the resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/0 to 25/1) to give the titled compound (11.4 g).

¹H-NMR (CDCl₃) (E)-isomer δ: 0.95 (t, 3H, J = 7.5 Hz), 1.65 (tq, 2H, J = 7.7, 7.5 Hz), 2.60 (t, 2H, J = 7.7 Hz), 5.87 (d, 1H, J = 16.5 Hz), 7.22-7.41 (m, 5H). (Z)-isomer δ: 0.95 (t, 3H, J = 7.5 Hz), 1.67 (tq, 2H, J = 7.7, 7.5 Hz), 2.64 (t, 2H, J = 7.7 Hz), 5.42 (d, 1H, J = 12.1 Hz), 7.11 (d, 1H, J = 12.1 Hz), 7.23-7.42 (m, 2H), 7.58-7.66 (m, 2H).

[0275]

(4) 1-Phenyl-5-(3-propylphenyl)-4,5-dihydro-1H-pyrazol-3-ylamine




To a solution of sodium ethoxide (20 wt% in ethanol, 31 ml) in ethanol (66 ml) was stirred and added dropwise phenylhydrazine (6.4 ml) at room temperature, and then the mixture was stirred for 40 minutes. To this reaction solution was added dropwise a solution of the mixture of (E)-3-(3-propylphenyl)acrylonitrile and (Z)-3-(3-propylphenyl)acrylonitrile (11.1 g) in ethanol (33 ml), and then the mixture was stirred overnight at reflux. This reaction solution was cooled to room temperature, and water (30 ml) was added thereto. Next, pH of the reaction solution was adjusted to 7 by adding a 6M aqueous solution of hydrochloric acid under ice-cooling. This mixed solution was extracted with ethyl acetate, and the resulting organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. To the resulting residue was added toluene, and the mixture was dried over anhydrous magnesium sulfate, and then concentrated under reduced pressure. To the resulting residue was added toluene, and again concentrated under reduced pressure to give a crude product of the titled compound

(18.6 g).

¹H-NMR (CDCl₃) δ: 0.91 (t, 3H, J = 7.5 Hz), 1.61 (tq, 2H, J = 7.7, 7.5 Hz), 2.56 (t, 2H, J = 7.7 Hz), 2.78 (dd, 1H, J = 16.3, 8.5 Hz), 3.41 (dd, 1H, J = 16.3, 11.3 Hz), 3.96 (s, 2H), 4.90 (dd, 1H, J = 11.3, 8.5 Hz), 6.67-6.72 (m, 1H), 6.80-6.87 (m, 2H), 7.02-7.37 (m, 6H).

[0276]

(5) 1-Phenyl-5-(3-propylphenyl)-1H-pyrazol-3-ylamine




To a solution of the crude product of 1-phenyl-5-(3-propylphenyl)-4,5-dihydro-1H-pyrazol-3-ylamine (18.6 g) in toluene (185 ml) was added activated carbon (pH5 to 8, 9.29 g), and the mixture was stirred overnight under oxygen atmosphere at reflux. This reaction solution was cooled to room temperature, and then the activated carbon was filtered off through Celite, and eluted with ethyl acetate. This filtrate was concentrated under reduced pressure, the resulting residue was dissolved in toluene (185 ml), and to the solution was added fresh activated carbon (pH5 to 8, 9.29 g), and the mixture was stirred overnight under oxygen atmosphere at reflux. This reaction solution was cooled to room temperature, and then the activated carbon was filtered off through Celite, and eluted with ethyl acetate. This filtrate was concentrated under reduced pressure, and the resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 4/1 to 1/1) to give the titled compound (8.68 g).

¹H-NMR (CDCl₃) δ: 0.82 (t, 3H, J = 7.3 Hz), 1.49 (tq, 2H, J = 7.6, 7.3 Hz), 2.47 (t, 2H, J = 7.6 Hz), 3.75 (s, 2H), 5.90 (s, 1H), 6.96-7.00 (m, 1H), 7.02-7.12 (m, 2H), 7.15-7.30 (m, 6H).

[0277]

(6) Optically active compound of 6-oxopiperidine-3-carboxylic acid [1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-yl]amide




To a solution of 1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-ylamine (56 mg) in tetrahydrofuran (1.5 ml) were sequentially added an optically active compound of 6-oxopiperidine-3-carboxylic acid (43 mg) synthesized in Preparation 7 (derived from the low-polarity component of 5-((R)-4-benzyl-2-oxoxazolidine-3-carbonyl)-1-(2,4-dimethoxybenzyl)piperidin-2-one), HOBt · H₂O (46 mg) and WSC · HCl (58 mg), and the mixture was stirred at room temperature overnight. To this reaction solution was added a saturated aqueous solution of sodium hydrogen carbonate, and the mixture was extracted with ethyl acetate. The separated organic layer was concentrated under reduced pressure, the resulting residue was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 10/1) to give the titled compound (50 mg).

¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.5, 7.3 Hz), 1.83-2.03 (m, 2H),


2.12-2.30 (m, 2H), 2.46 (t, 2H, $J = 7.5$ Hz), 2.84 (tt, 1H, $J = 10.7, 3.6$ Hz), 3.24-3.31 (m, 2H), 6.89 (s, 1H), 7.00 (s, 1H), 7.06 (d, 1H, $J = 7.7$ Hz), 7.17 (d, 1H, $J = 7.7$ Hz), 7.19-7.43 (m, 6H), 7.51 (s, 1H), 10.83 (s, 1H).

[0278]

- 5 [Preparation 9]: Synthesis of (R)-3-methyl-2-oxoimidazolidine-4-carboxylic acid and (S)-3-methyl-2-oxoimidazolidine-4-carboxylic acid




- (1) (R)-2-Oxoimidazolidine-1,5-dicarboxylic acid 1-benzyl ester



- 10 To a solution of sodium hydroxide (743 mg) in water (15 ml) was added dropwise bromine (318 μ l) under ice-cooling over 30 minutes. To this reaction solution was added Z-D-asparagine (1.5 g), and then the mixture was heated to 55°C, and stirred for 3 hours. This reaction solution was cooled to room temperature, and washed twice with diethyl ether (15 ml). To the resulting aqueous layer was added dropwise a 6M aqueous solution of hydrochloric acid until the pH reached 1. This mixed solution was left to stand at 4°C overnight. The precipitated solid was collected by filtration, washed with water, and dried under reduced pressure to give the titled compound (1.1 g).
- 15 $^1\text{H-NMR}$ (DMSO- D_6) δ : 3.22 (dd, 1H, $J = 10.0, 3.3$ Hz), 3.65 (dd, 1H, $J = 10.0, 10.2$ Hz), 4.68 (dd, 1H, $J = 10.2, 3.3$ Hz), 5.16 (d, 1H, $J = 14.8$ Hz), 5.19 (d, 1H, $J = 14.8$ Hz), 7.29-7.39 (m, 5H), 7.57 (s, 1H), 13.25 (br s, 1H).

[0279]

- 20 (2) (R)-2-Oxoimidazolidine-1,5-dicarboxylic acid 1-benzyl ester 5-tert-butyl ester




- 25 To a solution of (R)-2-oxoimidazolidine-1,5-dicarboxylic acid 1-benzyl ester (534 mg) in chloroform (2.6 ml) were sequentially added pyridine (1.0 ml) and tert-butylalcohol (1.5 ml) at room temperature, and the mixture was stirred. This reaction solution was iced, and phosphorous oxychloride (0.23 ml) was added thereto, and then the mixture was stirred at room temperature for 5 hours. This reaction solution was iced, and a 20 wt% aqueous solution of sodium acetate (20 ml) was added thereto, and then chloroform (5 ml) was further added at room temperature, and the mixture was stirred for 30 minutes. This mixed solution was extracted with chloroform, and the resulting organic layer was washed with a 0.1M aqueous solution of hydrochloric acid, water, a saturated
- 30

aqueous solution of sodium hydrogen carbonate and a saturated aqueous solution of sodium chloride, and then dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a crude product of the titled compound (467 mg).

¹H-NMR (CDCl₃) δ: 1.39 (s, 9H), 3.38 (dd, 1H, J = 9.8, 3.3 Hz), 3.71 (dd, 1H, J = 9.8, 9.8 Hz), 4.62 (dd, 1H, J = 9.8, 3.3 Hz), 5.24 (d, 1H, J = 12.6 Hz), 5.29 (d, 1H, J = 12.6 Hz), 6.00 (br s, 1H), 7.28-7.41 (m, 5H).

[0280]

(3) (R)-3-Methyl-2-oxoimidazolidine-1,4-dicarboxylic acid 1-benzyl ester 4-tert-butyl ester




To a solution of the crude product of (R)-2-oxoimidazolidine-1,5-dicarboxylic acid 1-benzyl ester 5-tert-butyl ester (318 mg) in N,N-dimethylformamide (3 ml) was added sodium hydride (42 mg) under ice-cooling, and the mixture was stirred for 40 minutes. To this reaction solution was added methyl iodide (309 μl), and then the mixture was warmed to room temperature, and stirred for 4 hours. This reaction solution was iced, and water and ethyl acetate were added thereto, and the mixture was stirred. This mixed solution was extracted with ethyl acetate, and the resulting organic layer was washed with a saturated aqueous solution of sodium chloride, and then dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/1) to give the titled compound (167 mg).

¹H-NMR (CDCl₃) δ: 1.48 (s, 9H), 2.91 (s, 3H), 3.81-3.85 (m, 1H), 3.96-4.02 (m, 2H), 5.27 (d, 1H, J = 13.1 Hz), 5.30 (d, 1H, J = 13.1 Hz), 7.29-7.45 (m, 5H).

[0281]

(4) (R)-3-Methyl-2-oxoimidazolidine-4-carboxylic acid tert-butyl ester




To a solution of (R)-3-methyl-2-oxoimidazolidine-1,4-dicarboxylic acid 1-benzyl ester 4-tert-butyl ester (167 mg) in methanol (1.7 ml) was added 10 wt% palladium on carbon at room temperature, and the mixture was stirred for 15 hours under oneatm. of hydrogen. Under nitrogen atmosphere, this reaction solution was filtered through Celite, and eluted with methanol. The filtrate was concentrated under reduced pressure to give the titled compound (94 mg).

¹H-NMR (CDCl₃) δ: 1.50 (s, 9H), 2.86 (s, 3H), 3.41-3.46 (m, 1H), 3.62-3.67 (m, 1H), 4.05 (dd, 1H, J = 9.8, 6.1 Hz), 4.38 (br s, 1H).

[0282]

(5) (R)-3-Methyl-2-oxoimidazolidine-4-carboxylic acid




To a solution of (R)-3-methyl-2-oxoimidazolidine-4-carboxylic acid tert-butyl ester (94 mg) in chloroform (2 ml) was added trifluoroacetic acid (1 ml) at room temperature, and the mixture was stirred for 3 hours. This reaction solution was concentrated under reduced pressure, additional toluene (3 ml) was added thereto, and the mixture was again concentrated. To the resulting residue was added diethyl ether (0.5 ml) at room temperature, and the mixture was stirred. Next, n-hexane (2 ml) was added thereto, and the mixture was further stirred. The insoluble substance was collected by filtration, and dried under reduced pressure to give the titled compound (52 mg).

¹H-NMR (DMSO-D₆) δ: 2.66 (s, 3H), 3.18 (dd, 1H, J = 9.2, 5.7 Hz), 3.51 (dd, 1H, J = 9.2, 9.9 Hz), 4.12 (dd, 1H, J = 9.9, 5.7 Hz), 6.42 (br s, 1H), 12.99 (br s, 1H).


Also, Z-L-asparagine was subjected to the same reactions and post-treatment as described above to give (S)-3-methyl-2-oxoimidazolidine-4-carboxylic acid (686 mg) as the optical isomer of the titled compound.

¹H-NMR (DMSO-D₆) δ: 2.66 (s, 3H), 3.18 (dd, 1H, J = 9.2, 5.7 Hz), 3.51 (dd, 1H, J = 9.2, 9.9 Hz), 4.12 (dd, 1H, J = 9.9, 5.7 Hz), 6.42 (br s, 1H), 12.99 (br s, 1H).




[0283]

[Preparation 10]: Example 6: Synthesis of ((R)-3-methyl-2-oxoimidazolidine-4-carboxylic acid [1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-yl]amide)



(1) (R)-3-methyl-2-oxoimidazolidine-4-carboxylic acid [1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-yl]amide




To a solution of 1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-ylamine (56 mg) prepared according to the same procedures as Preparation 8 in N,N-dimethylformamide (0.6 ml) were sequentially added (R)-3-methyl-2-oxoimidazolidine-4-carboxylic acid (38 mg) prepared in

Preparation 9, HOBt·H₂O (46 mg) and WSC·HCl (58 mg), and the mixture was stirred at room temperature for 2 hours. To this reaction solution were added a saturated aqueous solution of sodium hydrogen carbonate and water, and the mixture was extracted with ethyl acetate. The separated organic layer was washed with water and a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 10/1) to give the titled compound (40 mg).


¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.5, 7.3 Hz), 2.46 (t, 3H, J = 7.5 Hz), 2.64 (s, 3H), 3.21 (dd, 1H, J = 8.8, 6.3 Hz), 3.54 (dd, 1H, J = 9.4, 8.8 Hz), 4.28 (dd, 1H, J = 9.4, 6.3 Hz), 6.44 (s, 1H), 6.92 (s, 1H), 7.01 (s, 1H), 7.08 (d, 1H, J = 7.0 Hz), 7.17 (d, 1H, J = 7.0 Hz), 7.20-7.44 (m, 6H), 11.02 (s, 1H).

[0284]

[Preparation 11]: Synthesis of 5-oxotetrahydrofuran-3-carboxylic acid



(1) Sodium 5-oxo-2,5-dihydrofuran-3-carboxylate




To a solution of itaconic acid (26.0 g) in water (35 ml) was added dropwise bromine (10.3 ml) at room temperature while keeping a temperature of the reaction solution below 45°C, and then the mixture was stirred at 30°C for 15 minutes. To this reaction solution was added portionwise sodium hydrogen carbonate (33.6 g), and then the mixture was heated to 55°C. To this reaction solution was added dropwise a solution of sodium carbonate (10.6 g) in water (15 ml), and then the mixture was stirred for 20 minutes. This reaction solution was iced, and stirred for 1 hour. The precipitated solid was collected by filtration, washed with cold water and ethanol, and then dried under reduced pressure to give the titled compound (16.8 g).

¹H-NMR (DMSO-d₆) δ: 4.83 (d, 2H, J = 2.0 Hz), 5.93-5.95 (m, 1H).

[0285]

(2) 5-Oxotetrahydrofuran-3-carboxylic acid




To a solution of sodium 5-oxo-2,5-dihydrofuran-3-carboxylate (150 mg) in water (3 ml) was added 10 wt% palladium on carbon (30 mg), and the mixture was stirred for 18 hours at room temperature under one atm. of hydrogen. Under nitrogen atmosphere, an appropriate amount of amberlite IR-120 was added to the reaction solution, the mixture was stirred at room temperature for 10 minutes, and then filtered through Celite. The filtrate was concentrated under reduced pressure, ethyl acetate was added to the resulting residue, the mixture was dried over sodium sulfate, and then

again concentrated under reduced pressure to give the titled compound (83 mg).


¹H-NMR (DMSO-d₆) δ: 2.64 (dd, 1H, J = 17.6, 6.2 Hz), 2.75 (dd, 1H, J = 17.6, 9.4 Hz), 3.39-3.49 (m, 1H), 4.34 (dd, 1H, J = 8.8, 5.4 Hz), 4.44 (dd, 1H, J = 8.8, 8.2 Hz), 12.82 (br s, 1H).

[0286]

- 5 [Preparation 12]: Example 7: Synthesis of (5-oxotetrahydrofuran-3-carboxylic acid [1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-yl]amide)



- (1) 5-Oxotetrahydrofuran-3-carboxylic acid [1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-yl]amide



10

To a solution of 1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-ylamine (109 mg) prepared according to the same procedures as Preparation 8 in N,N-dimethylformamide/tetrahydrofuran (1/1, 2 ml) were sequentially added 5-oxotetrahydrofuran-3-carboxylic acid (82 mg) prepared in Preparation 9, HOBT·H₂O (96 mg) and WSC·HCl (121 mg), and the mixture was stirred at room temperature overnight. To this reaction solution was added a saturated aqueous solution of sodium hydrogen carbonate, and the mixture was extracted with ethyl acetate. The separated organic layer was concentrated under reduced pressure, and the resulting residue was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 10/1) to give the titled compound (96 mg).


15

¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.38-1.50 (m, 2H), 2.46 (t, 2H, J = 7.5 Hz), 2.69 (dd, 1H, J = 17.4, 5.9 Hz), 2.81 (dd, 1H, J = 17.4, 9.0 Hz), 3.56-3.65 (m, 1H), 4.34 (dd, 1H, J = 8.8, 5.0 Hz), 4.47-4.55 (m, 1H), 6.89 (s, 1H), 7.00 (s, 1H), 7.07 (d, 1H, J = 7.5 Hz), 7.17 (d, 1H, J = 7.1 Hz), 7.20-7.30 (3H, m), 7.31-7.43 (3H, m), 11.00 (s, 1H).


20

[0287]

- 25 [Preparation 13]: Example 8: Synthesis of ((R)-2-oxoimidazolidine-4-carboxylic acid [1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-yl]amide)



- (1) (R)-2,3-Bis-tert-butoxycarbonylaminopropionic acid




To a solution of di-tert-butyl dicarbonate (513 mg) in 1,4-dioxane (4 ml) was added N-Boc-(R)-2,3-diaminopropionic acid (400 mg) at room temperature, and the mixture was stirred for 20 minutes, and then heated to 50°C, and stirred for 20 minutes. The mixture was again cooled to room temperature, and then triethylamine (546 μ l) was added thereto, and the mixture was stirred for 2 hours. To this reaction solution were added water and a 10 wt% aqueous solution of citric acid, and the mixture was extracted with ethyl acetate. The resulting organic layer was washed twice with water, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: chloroform/ethanol = 1/0 to 10/1) to give the titled compound (557 mg).

$^1\text{H-NMR}$ (DMSO- D_6) δ : 1.37 (s, 9H), 1.38 (s, 9H), 3.23 (dd, 2H, $J = 6.2, 6.0$ Hz), 3.99 (dt, 1H, $J = 8.2, 6.2$ Hz), 6.78 (t, 1H, $J = 6.0$ Hz), 6.86 (d, 1H, $J = 8.2$ Hz), 12.52 (br s, 1H).

[0288]

(2) {(R)-2-tert-Butoxycarbonylamino-2-[1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-ylcarbamoyl]-ethyl}carbamic acid tert-butyl ester




To a solution of 1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-ylamine (177 mg) prepared according to the same procedures as Preparation 8 in DMF (1.8 ml) were added (R)-2,3-bis-tert-butoxycarbonylamino propionic acid (292 mg), HOBt $\cdot$ H $_2$ O (147 mg) and WSC $\cdot$ HCl (184 mg), and the mixture was stirred at room temperature for 2 hours. To this reaction solution were added water and a saturated aqueous solution of sodium hydrogen carbonate, and the mixture was extracted with ethyl acetate. The separated organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 2/1) to give the titled compound (329 mg).

$^1\text{H-NMR}$ (CDCl_3) δ : 0.83 (t, 3H, $J = 7.3$ Hz), 1.45 (s, 9H), 1.48 (s, 9H), 1.50 (tq, 2H, $J = 7.3, 7.5$ Hz), 2.49 (t, 2H, $J = 7.5$ Hz), 3.61-3.64 (br m, 2H), 4.38-4.41 (br m, 1H), 5.06 (br s, 1H), 5.94 (br s, 1H), 7.01 (s, 1H), 7.05-7.35 (m, 9H), 8.94 (br s, 1H).

[0289]

(3) (R)-2,3-Diamino-N-[1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-yl]-propionamide dihydrochloride




To a solution of [(R)-2-tert-butoxycarbonylamino-2-[1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-yl]carbonyl]-ethyl]carbamic acid tert-butyl ester (328 mg) in 1,4-dioxane (3 ml) was added a 4M solution of hydrogen chloride in 1,4-dioxane (3 ml) at room temperature, and the mixture was stirred for 2 hours. This reaction solution was concentrated, and then n-hexane (4 ml) was added thereto, and the mixture was stirred. The insoluble substance was collected by filtration, washed with n-hexane, and dried under reduced pressure to give the titled compound (181 mg).

¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.5, 7.3 Hz), 2.46 (t, 2H, J = 7.5 Hz), 3.41-3.46 (m, 2H), 3.57 (s, 1H), 4.38-4.41 (m, 1H), 6.91 (s, 1H), 7.11-7.31 (m, 9H), 8.56 (br s, 4H).

[0290]

(4) (R)-2-Oxoimidazolidine-4-carboxylic acid [1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-yl]amide




To a solution of (R)-2,3-diamino-N-[1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-yl]-propionamide dihydrochloride (181 mg) in N,N-dimethylformamide (3 ml) were added 1,1'-carbonyldiimidazole (67 mg), and triethylamine (116 μl) at room temperature, and the mixture was stirred for 16 hours. To this reaction solution was added water, and the mixture was extracted with ethyl acetate. The resulting organic layer was washed with water and a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 10/1) to give the titled compound (76 mg).

¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 1.44 (tq, 2H, J = 7.5, 7.4 Hz), 2.46 (t, 2H, J = 7.5 Hz), 3.35 (dd, 1H, J = 9.4, 5.6 Hz), 3.59 (dd, 1H, J = 9.4, 9.4 Hz), 4.32 (dd, 1H, J = 9.4, 5.6 Hz), 6.34 (s, 1H), 6.61 (s, 1H), 6.90 (s, 1H), 7.00 (s, 1H), 7.06-7.41 (m, 8H), 10.71 (s, 1H).

[0291]

[Preparation 14]: Synthesis of (S)-5-oxopyrrolidine-3-carboxylic acid



(1) (S)-4-Benzyl-3-[(S)-1-(2,4-dimethoxybenzyl)-5-oxopyrrolidine-3-carbonyl]-2-oxazolidinone




To a solution of 1-(2,4-dimethoxybenzyl)-5-oxopyrrolidine-3-carboxylic acid (20 g) prepared according to the same procedures as Preparation 5 in chloroform (200 ml) were sequentially added (S)-4-benzyl-2-oxazolidinone (15.2 g) and WSC·HCl (16.4 g), 4-(dimethylamino)pyridine (4.4 g) at room temperature, and the mixture was stirred for 3 hours. This reaction solution was washed with a 5 wt% aqueous solution of sodium hydrogen carbonate and a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/1 to 1/3), subjected to azeotropic drying with toluene, and dried under reduced pressure to give the titled compound (17.2 g).

¹H-NMR (CDCl₃) δ: 2.72-2.83 (m, 3H), 3.27 (dd, 1H, J = 13.4, 3.5 Hz), 3.46-3.57 (m, 2H), 3.80 (s, 3H), 3.80 (s, 3H), 4.15-4.29 (m, 3H), 4.40-4.48 (m, 2H), 4.61-4.69 (m, 1H), 6.42-6.47 (m, 2H), 7.12-7.21 (m, 3H), 7.27-7.37 (m, 3H).

[0292]

(2) (S)-1-(2,4-dimethoxybenzyl)-5-oxopyrrolidine-3-carboxylic acid



To a solution of lithium hydroxide monohydrate (2.0 g) in water (45 ml) was added dropwise 30 wt% aqueous solution of hydrogen peroxide solution (11 ml) in an ice-sodium chloride cooling bath, and the mixture was stirred for 15 minutes. To this reaction solution was added tetrahydrofuran (36 ml), and then a solution of (S)-4-benzyl-3-[(S)-1-(2,4-dimethoxybenzyl)-5-oxopyrrolidine-3-carbonyl]-2-oxazolidinone (17.2 g) in tetrahydrofuran (144 ml) was added dropwise thereto, and the mixture was stirred for additional 20 minutes. To this reaction solution was slowly added a 10 wt% aqueous solution of sodium hydrogen sulfite (131 ml), and the mixture was stirred at room temperature for 1 hour after removing the bath, and then extracted with ethyl acetate (180 ml). The resulting organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: ethyl acetate to chloroform/methanol = 9/1). To the resulting product (6.7 g) was added diisopropyl ether (13.4 ml), the mixture was stirred at 70°C, and then cooled to 0°C with stirring. A solid was collected from this suspension by filtration, dried under reduced pressure to give the titled compound (5.5 g). An analysis of this solid by HPLC analysis condition 1 showed that an isomer with longer retention time was a main product.


An isomer with shorter retention time (retention time 4.7 minutes)

An isomer with longer retention time (retention time 5.4 minutes)

$^1\text{H-NMR}$ (CDCl_3) δ : 2.66-2.82 (m, 2H), 3.20 (dt, 1H, $J = 17.3, 7.6$ Hz), 3.49-3.53 (m, 2H), 3.80 (s, 6H), 4.40 (d, 1H, $J = 14.6$ Hz), 4.46 (d, 1H, $J = 14.6$ Hz), 6.42-6.47 (m, 2H), 7.14 (t, 1H, $J = 4.5$ Hz).

[0293]

(3) (S)-5-oxopyrrolidine-3-carboxylic acid




To (S)-1-(2,4-dimethoxybenzyl)-5-oxopyrrolidine-3-carboxylic acid (5.3 g) were added anisole (3.1 ml) and trifluoroacetic acid (26.5 ml) at room temperature, and the mixture was stirred for 4 hours at reflux. This reaction solution was cooled to room temperature, concentrated under reduced pressure, azeotropic-dried with toluene. To the resulting residue was added diisopropyl ether (53 ml), and the mixture was stirred at room temperature. A solid was collected from this suspension by filtration, washed with diisopropyl ether, and dried under reduced pressure to give the titled compound (2.4 g).


$^1\text{H-NMR}$ (DMSO-D_6) δ : 2.26-2.40 (m, 2H), 3.18-3.28 (m, 1H), 3.30-3.35 (m, 1H), 3.40-3.47 (m, 1H), 7.64 (s, 1H), 12.58 (s, 1H)

[0294]

[Preparation 15]: Example 235: Synthesis of ((S)-5-oxopyrrolidine-3-carboxylic acid {5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1-phenyl-1H-pyrazol-3-yl} amide)



(1) 1-Fluoro-3-((R)-2,2,2-trifluoro-1-methylethoxymethyl)benzene




Under argon atmosphere, a suspension of sodium hydride (76mg, 60 wt% oil dispersion) in N,N-dimethylformamide (2.0 ml) was cooled to 0°C , a solution of N,N-dimethylformamide (2.0 ml) in (R)-1,1,1-trifluoropropan-2-ol (219 mg) was added thereto, and the mixture was stirred for 25 minutes. Then, a solution of 1-bromomethyl-3-fluorobenzene (297 mg) in N,N-dimethylformamide (2.0 ml) was added thereto, and then the mixture was stirred for 2 hours. To this reaction mixture was added water, and the mixture was extracted with ethyl acetate. This organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel thin-layer chromatography (eluent: n-hexane/ethyl acetate = 9/1) to give the titled compound (242 mg).

$^1\text{H-NMR}$ (CDCl_3) δ : 1.35 (d, 3H, $J = 6.47$ Hz), 3.84 (qq, 1H, $J = 6.47, 9.40$ Hz), 4.63 (d, 1H, $J =$

12.00Hz), 4.76 (d, 1H, $J = 12.00$ Hz), 6.98-7.03 (m, 1H), 7.06-7.12 (m, 2H), 7.29-7.35 (m, 1H).

[0295]

(2) 2-[3-Fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane



5

Under argon atmosphere, to a solution of 1-fluoro-3-((R)-2,2,2-trifluoro-1-methylethoxymethyl)benzene (159 mg) in tetrahydrofuran (3.0 ml) were sequentially added bispinacolate diboron (202 mg), 4,4'-di-tert-butyl-2,2'-dipyridyl (10 mg) and di-μ-methoxybis(1,5-cyclooctadiene)diiridium (I) (12 mg) at room temperature, and then the mixture was stirred for 16 hours at reflux. This reaction mixture was cooled to room temperature, and then a saturated aqueous solution of sodium hydrogen carbonate and ethyl acetate were added thereto, the mixture was filtered through Celite, and extracted with ethyl acetate. The filtrate was extracted with ethyl acetate, the separated organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel thin-layer chromatography (eluent: n-hexane/ethyl acetate = 85/15) to give the titled compound (49 mg).

10


15

¹H-NMR (CDCl₃) δ: 1.35 (m, 15H), 3.83 (qq, 1H, $J = 6.47, 9.40$ Hz), 4.62 (d, 1H, $J = 12.00$ Hz), 4.75 (d, 1H, $J = 12.00$ Hz), 7.19 (d, 1H, $J = 9.60$ Hz), 7.43 (d, 1H, $J = 9.60$ Hz), 7.49 (s, 1H).

[0296]

20

(3) 5-[3-Fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1-phenyl-1H-pyrazol-3-ylamine




To 2-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane (49 mg) were sequentially added a solution of 5-iodo-1-phenyl-1H-pyrazol-3-ylamine (36 mg) prepared according to the same procedures as Preparation 2 in 1,2-dimethoxyethane (1.0 ml), a 2M aqueous solution of sodium carbonate (0.5 ml), tricyclohexylphosphine (7.3 mg) and palladium (II) acetate (3.0 mg) at room temperature, and the mixture was stirred at 100°C for 2 hours. This reaction mixture was cooled to room temperature, a saturated aqueous solution of sodium hydrogen carbonate and ethyl acetate were added thereto, the mixture was filtered through Celite, and extracted with ethyl acetate. The filtrate was extracted with ethyl acetate, the separated organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 9/1) to give the titled compound (37 mg).

25

30

[0297]

(4) (S)-5-Oxopyrrolidine-3-carboxylic acid {5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1-phenyl-1H-pyrazol-3-yl}amide




To a solution of 5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1-phenyl-1H-pyrazol-3-ylamine (37 mg) in *N,N*-dimethylacetamide (1.0 ml) were sequentially added (S)-5-oxopyrrolidine-3-carboxylic acid (16 mg) prepared in Preparation 14 and WSC·HCl (23 mg) at room temperature, the mixture was stirred for 5.2 hours. To this reaction mixture was added water, and then the precipitated solid was collected by filtration. This solid was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 9/1). To the resulting solid were added water and a small amount of methanol, and the mixture was stirred. A solid was collected from this suspension by filtration, and was dried under reduced pressure to give the titled compound (20 mg).


¹H-NMR (DMSO-D₆) δ: 1.18 (d, 3H, *J* = 6.5 Hz), 2.36-2.38 (m, 2H), 3.32-3.33 (m, 1H), 3.37-3.41 (m, 1H), 3.46-3.49 (m, 1H), 4.10 (qq, 1H, *J* = 6.5, 9.4 Hz), 4.57 (d, 1H, *J* = 12.7 Hz), 4.61 (d, 1H, *J* = 12.7 Hz), 6.96-3.98 (m, 2H), 7.09 (s, 1H), 7.15-7.18 (m, 1H), 7.25-7.27 (m, 2H), 7.35-7.44 (m, 3H), 7.63 (s, 1H), 10.88 (s, 1H).

[0298]

[Preparation 16]: Example 236: Synthesis of ((S)-5-oxopyrrolidine-3-carboxylic acid {1-(4-chlorophenyl)-5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-yl}amide)



(1) 1-(4-Chlorophenyl)-3-(2,5-dimethylpyrrol-1-yl)-1H-pyrazole




Under argon atmosphere, to a solution of 3-(2,5-dimethylpyrrol-1-yl)-1H-pyrazole (3.06 g) prepared according to the same procedures as Preparation 4 in *N*-methylpyrrolidone (31 ml) were sequentially added cesium carbonate (12.37 g), 8-quinolinol (0.55 g), copper (I) oxide (0.27 g) and 1-chloro-4-iodobenzene (6.79 g) at room temperature, the mixture was stirred at 110°C for 2 hours. This reaction mixture was cooled to room temperature, and then toluene (30 ml) was added thereto, the mixture was filtered through Celite, and further subjected to elution with toluene. The filtrate was sequentially washed with a 1N aqueous solution of sodium hydroxide, water, a saturated aqueous

solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/0 to 9/1) to give the titled compound (4.98 g).

¹H-NMR (CDCl₃) δ: 2.19 (s, 6H), 5.90 (s, 2H), 6.40 (d, 1H, J = 2.65 Hz), 7.43 (d, 2H, J = 8.82 Hz), 7.66 (d, 2H, J = 8.82 Hz), 7.94 (d, 1H, J = 2.65 Hz).

[0299]

(2) 1-(4-Chlorophenyl)-3-(2,5-dimethylpyrrol-1-yl)-5-iodo-1H-pyrazole



Under argon atmosphere, a solution of 3-(2,5-dimethylpyrrol-1-yl)-1-(4-chlorophenyl)-1H-pyrazole (4.98 g) in tetrahydrofuran (40 ml) was cooled to -78°C, n-butyllithium (13.7 ml, 1.6M solution in n-hexane) was added dropwise thereto, and the mixture was stirred for 30 minutes. To this reaction solution was added dropwise a solution of iodine (5.58 g) in tetrahydrofuran (10 ml), the mixture was stirred for 30 minutes. To this reaction solution were added a 20 wt% aqueous solution of sodium sulfite and a saturated aqueous solution of ammonium chloride, the mixture was extracted with ethyl acetate. The organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. To the resulting solid residue was added a mixed solvent of ethyl acetate /n-hexane=1/2, the mixture was stirred. A solid was collected from this suspension by filtration, and was dried under reduced pressure to give the titled compound (4.19 g).

¹H-NMR (CDCl₃) δ: 2.18 (s, 6H), 5.88 (s, 2H), 6.55 (s, 1H), 7.47 (d, 2H, J = 9.04 Hz), 7.55 (d, 2H, J = 9.04 Hz).

[0300]

(3) 1-(4-Chlorophenyl)-5-iodo-1H-pyrazol-3-ylamine




To a suspension of 1-(4-chlorophenyl)-3-(2,5-dimethylpyrrol-1-yl)-5-iodo-1H-pyrazole (4.19 g) in ethanol/water (2/1, 72 ml) were sequentially added hydroxylammonium chloride (14.64 g) and triethylamine (2.9 ml) at room temperature, and the mixture was stirred at 95°C for 86 hours. This reaction mixture was cooled to room temperature, and concentrated under reduced pressure. To the resulting residue was added an 8M aqueous solution of sodium hydroxide, and then the mixture was extracted with ethyl acetate. This organic layer was sequentially washed with water and a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 2/1) to give the titled compound (1.87 g).

¹H-NMR (CDCl₃) δ: 3.77 (br s, 2H), 6.01 (s, 1H), 7.39-7.50 (m, 4H).

[0301]


(4) 1-(4-Chlorophenyl)-5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-ylamine



Under argon atmosphere, to a solution of 1-(4-chlorophenyl)-5-iodo-1H-pyrazol-3-ylamine (35 mg) in 1,2-dimethoxyethane (0.7ml) were sequentially added 2-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane (46 mg) prepared according to the same procedures as Preparation 15, a 2M aqueous solution of sodium carbonate (0.35ml), tricyclohexylphosphine (6.1 mg) and palladium (II) acetate (2.5 mg) at room temperature, and the mixture was stirred at 100°C for 2 hours. This reaction mixture was cooled to room temperature, filtered through Celite, was subjected to elution with ethyl acetate. The filtrate was dried over anhydrous magnesium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel thin-layer chromatography (eluent: n-hexane/ ethyl acetate = 1/1) to give the titled compound.

[0302]

(5) (S)-5-Oxopyrrolidine-3-carboxylic acid {1-(4-chlorophenyl)-5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-yl}amide




To a solution of 1-(4-chlorophenyl)-5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-ylamine (about 0.110 mol) prepared in the previous step in N,N-dimethylacetamide (0.5 ml) were sequentially added (S)-5-oxopyrrolidine-3-carboxylic acid (17 mg) prepared in Preparation 14 and WSC·HCl (32 mg) at room temperature, and the mixture was stirred for 0.5 hours. To this reaction mixture was added water, and then the precipitated solid was collected by filtration. The resulting solid was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 9/1) to give the titled compound (18 mg).


¹H-NMR (DMSO-D₆) δ: 1.20 (d, 3H, J = 6.5 Hz), 2.38-2.40 (m, 2H), 3.33-3.35 (m, 1H), 3.40-3.44 (m, 1H), 3.48-3.51 (m, 1H), 4.10-4.15 (m, 1H), 4.63 (s, 2H), 6.99 (s, 1H), 7.06 (s, 1H), 7.08 (d, 1H, J = 9.3 Hz), 7.19 (d, 1H, J = 9.3 Hz), 7.27 (dd, 2H, J = 6.6, 2.0 Hz), 7.48 (dd, 2H, J = 6.6, 2.0 Hz), 7.63 (s, 1H), 10.91 (s, 1H).

[0303]

[Preparation 17]: Example 237: Synthesis of ((S)-5-oxopyrrolidine-3-carboxylic acid {1-(2-fluorophenyl)-5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-yl}amide)



(1) 3-(2,5-Dimethylpyrrol-1-yl)-1-(2-fluorophenyl)-1H-pyrazole




To a solution of 3-(2,5-dimethylpyrrol-1-yl)-1H-pyrazole (1.00 g) prepared according to the same procedures as Preparation 4 in N-methylpyrrolidone (10 ml) were sequentially added 1-fluoro-2-iodobenzene (1.08 ml), copper (I) oxide (89 mg), 8-quinolinol (180 mg) and cesium carbonate (4.04 g) at room temperature, and the mixture was stirred at 110°C for 3 hours. This reaction mixture was cooled to room temperature, and then toluene was added thereto, and the mixture was filtered through Celite. The filtrate was sequentially washed with a 2N aqueous solution of sodium hydroxide, water and a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/0 to 10/1) to give the titled compound (1.19 g). [0304]

(2) 3-(2,5-Dimethylpyrrol-1-yl)-1-(2-fluorophenyl)-5-iodo-1H-pyrazole



Under nitrogen atmosphere, a solution of 3-(2,5-dimethylpyrrol-1-yl)-1-(2-fluorophenyl)-1H-pyrazole (1.19 g) in tetrahydrofuran (10 ml) was cooled to -78°C, n-butyllithium (3.5 ml, 1.6M solution in n-hexane) was added dropwise thereto, and the mixture was stirred for 30 minutes. To this reaction solution was added dropwise a solution of iodine (1.42 g) in tetrahydrofuran (5 ml), and the mixture was stirred for 30 minutes. To this reaction mixture was added a saturated aqueous solution of sodium sulfite and a saturated aqueous solution of ammonium chloride, and the mixture was extracted with ethyl acetate. This organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. To the resulting residue was added a mixed solvent of n-hexane/ethyl acetate (=6/1), the mixture was stirred. A solid was collected from this suspension by filtration, and was dried under reduced pressure to give the titled compound (1.23 g). [0305]

(3) 1-(2-Fluorophenyl)-5-iodo-1H-pyrazol-3-ylamine




To a solution of 3-(2,5-dimethylpyrrol-1-yl)-1-(2-fluorophenyl)-5-iodo-1H-pyrazole (1.23 g)

in ethanol/water (2/1, 18 ml) were sequentially added hydroxylammonium chloride (4.47 g) and triethylamine (0.9 ml) at room temperature, and the mixture was stirred at 90°C for 24 hours. This reaction mixture was cooled to room temperature, concentrated under reduced pressure. To the resulting residue was added a 8M aqueous solution of sodium hydroxide, and then the mixture was extracted with ethyl acetate. This organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 4/1 to 1/1) to give the titled compound (0.61 g).

¹H-NMR (CDCl₃) δ: 3.76 (br s, 2H), 6.03 (s, 1H), 7.19-7.25 (m, 2H), 7.38-7.46 (m, 2H).

[0306]


(4) 1-(2-Fluorophenyl)-5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-ylamine



To a solution of 1-(2-fluorophenyl)-5-iodo-1H-pyrazol-3-ylamine (38 mg) in 1,4-dioxane (0.4 ml) were sequentially added 2-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (52 mg) prepared according to the same procedures as Preparation 15, palladium (II) acetate (3 mg), tricyclohexylphosphine (7 mg), 2M aqueous solution of potassium carbonate (0.2ml) at room temperature, and the mixture was stirred at 105°C for 4 hours. After removing the aqueous layer of this reaction mixture, the organic layer was filtered through silica gel (5 g), and subjected to elution with ethyl acetate. The filtrate was concentrated under reduced pressure, and the resulting residue was purified by silica gel thin-layer chromatography (eluent: n-hexane/ethyl acetate 1/1) to give the titled compound.

[0307]

(5) (S)-5-Oxopyrrolidine-3-carboxylic acid {1-(2-fluorophenyl)-5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-yl}amide




To a solution of 1-(2-fluorophenyl)-5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-ylamine (about 0.125 mmol) prepared in the previous step in dimethylformamide (0.5 ml) were sequentially added (S)-5-oxopyrrolidine-3-carboxylic acid (18 mg) prepared in Preparation 14 and WSC·HCl (29 mg) at room temperature, and the mixture was stirred for 3 hours. To this reaction mixture was added water, and the mixture was extracted with ethyl acetate. This organic layer was concentrated under reduced pressure, and the resulting residue was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 10/1). To the

resulting solid were added water and a small amount of methanol, and the mixture was stirred. A solid was collected from this suspension by filtration, and was dried under reduced pressure to give the titled compound (32 mg).


¹H-NMR (CDCl₃) δ: 1.29 (d, 3H, J = 6.4 Hz), 2.59 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.8 Hz), 3.29-3.40 (m, 1H), 3.60 (dd, 1H, J = 9.2 Hz), 3.67-3.80 (m, 2H), 4.52 (d, 1H, J = 12.0 Hz), 4.63 (d, 1H, J = 12.0 Hz), 6.08 (s, 1H), 6.85 (dt, 1H, J = 9.6, 2.0 Hz), 6.98-7.04 (m, 2H), 7.09-7.17 (m, 2H), 7.22 (t, 1H, J = 8.0 Hz), 7.34-7.22 (m, 2H), 8.69 (s, 1H).

[0308]

[Preparation 18]: Example 238: Synthesis of ((S)-5-oxopyrrolidine-3-carboxylic acid {1-(4-fluorophenyl)-5-[3-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-yl} amide)



(1) 3-(2,5-Dimethylpyrrol-1-yl)-1-(4-fluorophenyl)-1H-pyrazole




To a solution of 3-(2,5-dimethylpyrrol-1-yl)-1H-pyrazole (500 mg) prepared according to the same procedures as Preparation 4, 4-fluorophenyl boronic acid (651 mg) and copper (II) acetate (563 mg) in chloroform (5 ml) was added pyridine (0.50 ml) at room temperature, the mixture was stirred for 40 hours. This reaction solution was filtered through silica gel, and subjected to elution with ethyl acetate. The filtrate was concentrated under reduced pressure, the resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 9/1) to give the titled compound (599 mg).

¹H-NMR (CDCl₃) δ: 2.19 (s, 6H), 5.90 (s, 2H), 6.39 (d, 1H, J = 2.56 Hz), 7.15-7.17 (m, 2H), 7.67-7.70 (m, 2H), 7.91 (d, 1H, J = 2.56 Hz).

[0309]

(2) 3-(2,5-Dimethylpyrrol-1-yl)-1-(4-fluorophenyl)-5-iodo-1H-pyrazole



Under argon atmosphere, a solution of 3-(2,5-dimethylpyrrol-1-yl)-1-(4-fluorophenyl)-1H-pyrazole (599 mg) in tetrahydrofuran (4 ml) was cooled to -78°C, n-butyllithium (1.8ml, 1.6M solution in n-hexane) was added dropwise thereto, and the mixture was stirred for 30 minutes. To this reaction solution was added dropwise a solution of iodine (715 mg) in tetrahydrofuran (2 ml), the mixture was stirred for additional 30 minutes. To this reaction mixture was added a saturated aqueous solution of sodium sulfite, and the mixture was extracted with ethyl acetate. The resulting organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. To the resulting solid residue was added a

mixed solvent of ethyl acetate /n-hexane (=1/5), and the mixture was stirred. A solid was collected from this suspension by filtration, dried under reduced pressure to give the titled compound (625 mg). ¹H-NMR (CDCl₃) δ: 2.18 (s, 6H), 5.88 (s, 2H), 6.54 (s, 1H), 7.18-7.20 (m, 2H), 7.54-7.57 (m, 2H).

[0310]

5 (3) 1-(4-Fluorophenyl)-5-iodo-1H-pyrazol-3-ylamine




To a suspension of 3-(2,5-dimethylpyrrol-1-yl)-1-(4-fluorophenyl)-5-iodo-1H-pyrazole (625 mg) in ethanol/water (2/1, 12.5 ml) were sequentially added hydroxylammonium chloride (2.28 g) and triethylamine (0.46 ml) at room temperature, and the mixture was stirred at 95°C for 15 hours.

10 This reaction mixture was cooled to room temperature, concentrated under reduced pressure. To the resulting residue was added a 8M aqueous solution of sodium hydroxide, and then the mixture was extracted with ethyl acetate. This organic layer was sequentially washed with water and a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/1) to give the titled compound (318 mg).

¹H-NMR (CDCl₃) δ: 3.75 (br s, 2H), 6.00 (s, 1H), 7.11-7.15 (m, 2H), 7.44-7.47 (m, 2H).

[0311]


(4) (R)-3-[1,1,1-Trifluoropropan-2-yloxy]phenyl boronic acid



20 A solution of (R)-1,1,1-trifluoropropan-2-ol (212 mg) in N,N-dimethylformamide (1.5 ml) was cooled to 0°C, sodium hydride (74mg, 60 wt.% oil dispersion) and 3-(bromomethyl)phenyl boronic acid (100 mg) were sequentially added thereto, and the mixture was stirred at room temperature for 2 hours. To this reaction mixture were added water and an 1M aqueous solution of hydrochloric acid, and the mixture was extracted with ethyl acetate. This organic layer was washed with water and a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure to give the titled compound (147 mg).

[0312]

(5) 1-(4-Fluorophenyl)-5-[3-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-ylamine



30


To a solution of 1-(4-fluorophenyl)-5-iodo-1H-pyrazol-3-ylamine (70 mg) in 1,2-dimethoxyethane (0.7ml) were sequentially added (R)-3-(1,1,1-trifluoropropan-2-yloxy)phenyl boronic acid (104 mg), a 2M aqueous solution of sodium carbonate (0.35 ml), tricyclohexylphosphine

(13.0 mg) and palladium (II) acetate (5.2 mg) at room temperature, and the mixture was stirred at 100°C for 15 hours. This reaction mixture was cooled to room temperature, filtered through Celite, and subjected to elution with ethyl acetate. The filtrate was extracted with ethyl acetate, the resulting organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/1) to give the titled compound (52 mg).

¹H-NMR (CDCl₃) δ: 1.28 (d, 3H, J = 6.51 Hz), 3.74-3.80 (m, 2H), 3.76-3.78 (m, 1H), 4.56 (d, 1H, J = 11.86 Hz), 4.66 (d, 1H, J = 11.86 Hz), 5.92 (s, 1H), 6.96-6.99 (m, 2H), 7.15-7.21 (m, 4H), 7.29-7.29 (m, 2H).

[0313]

(6) (S)-5-Oxopyrrolidine-3-carboxylic acid {1-(4-fluorophenyl)-5-[3-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-yl} amide




To a solution of 1-(4-fluorophenyl)-5-[3-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-ylamine (52 mg) in N,N-dimethylacetamide (0.5 ml) were sequentially added (S)-5-oxopyrrolidine-3-carboxylic acid (20 mg) prepared according to the same procedures as Preparation 5 and WSC·HCl (40 mg) at room temperature, and the mixture was stirred for 0.5 hours. To this reaction mixture was added water, and then the precipitated solid was collected by filtration. This solid was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 9/1). To the resulting solid were added water and a small amount of methanol, and the mixture was stirred. A solid was collected by filtration from this suspension, and dried under reduced pressure to give the titled compound (37 mg).

¹H-NMR (DMSO-D₆) δ: 1.20 (d, 3H, J = 6.5 Hz), 2.38-2.39 (m, 2H), 3.33-3.34 (m, 1H), 3.39-3.43 (m, 1H), 3.49-3.51 (m, 1H), 4.09-4.11 (m, 1H), 4.59 (d, 1H, J = 12.1 Hz), 4.63 (d, 1H, J = 12.1 Hz), 6.92 (s, 1H), 7.18-7.39 (m, 8H), 7.63 (s, 1H), 10.85 (s, 1H).


[0314]

[Preparation 19]: Example 239: Synthesis of ((S)-5-oxopyrrolidine-3-carboxylic acid {1-(4-fluorophenyl)-5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-yl} amide)



(1) 1-(4-Fluorophenyl)-5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-


pyrazol-3-ylamine



To a solution of 1-(4-fluorophenyl)-5-iodo-1H-pyrazol-3-ylamine (38 mg) prepared in Preparation 18 in 1,4-dioxane (0.4 ml) were sequentially added 2-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane (52 mg) prepared according to the same procedures as Preparation 15, palladium (II) acetate (3 mg), tricyclohexylphosphine (7 mg) and a 2M aqueous solution of potassium carbonate (0.2ml) at room temperature, and the mixture was stirred at 105°C for 4 hours. After removing an aqueous layer of the reaction mixture, an organic layer was filtered through silica gel (5 g), and subjected to elution with ethyl acetate. The filtrate was concentrated under reduced pressure, the resulting residue was purified by silica gel thin-layer chromatography (eluent: n-hexane/ ethyl acetate = 1/1) to give the titled compound.

[0315]

(2) (S)-5-Oxopyrrolidine-3-carboxylic acid {1-(4-fluorophenyl)-5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-yl} amide




To a solution of 1-(4-fluorophenyl)-5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-ylamine prepared in the previous step (about 0.125 mmol) in dimethylformamide (0.5 ml) were sequentially added (S)-5-oxopyrrolidine-3-carboxylic acid (18 mg) prepared according to the same procedures as Preparation 14 and WSC·HCl (29 mg) at room temperature, and the mixture was stirred for 3 hours. To this reaction mixture was added water, and the mixture was extracted with ethyl acetate. This organic layer was concentrated under reduced pressure, the resulting residue was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 10/1). To the resulting solid were added water and a small amount of methanol, and the mixture was stirred. A solid was collected from this suspension by filtration, and was dried under reduced pressure to give the titled compound (31 mg).


¹H-NMR (CDCl₃) δ: 1.31 (d, 3H, J = 6.4 Hz), 2.61 (dd, 1H, J = 16.8, 9.6 Hz), 2.78 (dd, 1H, J = 16.8, 8.4 Hz), 3.31-3.42 (m, 1H), 3.61 (t, 1H, J = 8.8 Hz), 3.67-3.83 (m, 2H), 4.55 (d, 1H, J = 12.0 Hz), 4.65 (d, 1H, J = 12.0 Hz), 6.04 (s, 1H), 6.85 (d, 1H, J = 9.2 Hz), 6.98-7.10 (m, 5H), 7.19-7.25 (m, 2H), 8.48 (s, 1H).

[0316]

[Preparation 20]: Synthesis of (R)-2-oxoimidazolidine-4-carboxylic acid



(1) (R)-2-Oxoimidazolidine-1,5-dicarboxylic acid 1-benzyl ester




Under ice-cooling, to a 1M aqueous solution of sodium hydroxide (124 ml) was added dropwise bromine (6.60 ml), and then after adding (R)-2-benzyloxycarbonylamino succinamic acid (10.0 g), the mixture was stirred at 55°C for 3 hours. This reaction solution was cooled to room temperature, washed twice with diethyl ether, and then a 6M aqueous solution of hydrochloric acid (21 ml) was added thereto. This reaction mixture was left to stand at 4°C for 3 days, and a precipitated solid was collected by filtration to give the titled compound (6.16 g).

¹H-NMR (DMSO-D₆) δ: 3.19-3.22 (m, 1H), 3.63 (dd, 1H, J = 10.2, 5.0 Hz), 4.67 (dd, 1H, J = 10.2, 3.3 Hz), 5.14-5.18 (m, 2H), 7.28-7.40 (m, 5H), 7.56 (s, 1H), 13.24 (br s, 1H).

[0317]

(2) (R)-2-Oxoimidazolidine-4-carboxylic acid




Under nitrogen atmosphere, to a solution of (R)-2-oxoimidazolidine-1,5-dicarboxylic acid 1-benzyl ester (1.61 g) in methanol/tetrahydrofuran (4/1, 20 ml) was added 10 wt% palladium on carbon (161 mg) at room temperature, and the mixture was stirred for 24 hours under one atmosphere of hydrogen. Under nitrogen atmosphere, palladium on carbon in this reaction solution was filtered off through Celite, and the filtrate was concentrated under reduced pressure. To the resulting residue were added diisopropyl ether and n-hexane, and the mixture was stirred. A solid was collected from this suspension by filtration, and dried under reduced pressure to give the titled compound (565 mg).


¹H-NMR (DMSO-D₆) δ: 3.27 (ddd, 1H, J = 9.8, 4.8, 0.8 Hz), 3.54 (dd, 1H, J = 9.8, 9.8 Hz), 4.10 (dd, 1H, J = 9.8, 4.8 Hz), 6.24 (s, 1H), 6.56 (s, 1H), 12.83 (br s, 1H).

[0318]

[Preparation 21]: Example 240: Synthesis of ((R)-2-oxoimidazolidine-4-carboxylic acid {1-(4-fluorophenyl)-5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-yl}amide)




(1) 1-(4-Fluorophenyl)-5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-ylamine



To a solution of 1-(4-fluorophenyl)-5-iodo-1H-pyrazol-3-ylamine (84 mg) prepared according to the same procedures as Preparation 18 in 1,4-dioxane (1.0ml) were sequentially added 2-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane (106 mg) prepared according to the same procedures as Preparation 15, palladium (II) acetate (7 mg), tricyclohexylphosphine (17 mg) and a 2M aqueous solution of potassium carbonate (0.5ml) at room temperature, and the mixture was stirred at 105°C for 2 hours. This reaction mixture was cooled to room temperature and after removing an aqueous layer, an organic layer was filtered through silica gel (5 g), and subjected to elution with ethyl acetate. The filtrate was concentrated under reduced pressure, and the resulting residue was purified by silica gel thin-layer chromatography (eluent: n-hexane/ ethyl acetate = 3/2) to give the titled compound.

[0319]

(2) (R)-2-Oxoimidazolidine-4-carboxylic acid {1-(4-fluorophenyl)-5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-yl}amide




To a solution of 1-(4-fluorophenyl)-5-[3-fluoro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-ylamine (about 0.277 mmol) prepared in the previous step in chloroform (1 ml) were sequentially added (R)-2-oxoimidazolidine-4-carboxylic acid (40 mg) and WSC·HCl (64 mg) at room temperature, and the mixture was stirred for 3 hours. This reaction mixture was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 10/1). To the resulting solid were added water and a small amount of methanol, and the mixture was stirred. A solid was collected from this suspension by filtration and was dried under reduced pressure to give the titled compound (56 mg).

¹H-NMR (DMSO-D₆) δ: 1.20 (d, 3H, J = 6.0 Hz), 3.35 (dd, 1H, J = 8.6, 6.2 Hz), 3.58 (dd, 1H, J = 9.3, 8.6 Hz), 4.12 (tq, 1H, J = 9.7, 6.0 Hz), 4.31 (dd, 1H, J = 9.3, 6.2 Hz), 4.60 (d, 1H, J = 12.5 Hz), 4.63


(d, 1H, J = 12.5 Hz), 6.33 (s, 1H), 6.59 (s, 1H), 6.98 (s, 1H), 7.02-7.07 (m, 2H), 7.18 (d, 1H, J = 9.6 Hz), 7.23-7.35 (m, 4H), , 10.72 (s, 1H).

[0320]

[Preparation 22]: Example 241: Synthesis of ((S)-5-oxopyrrolidine-3-carboxylic acid {5-[3-chloro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1-phenyl-1H-pyrazol-3-yl} amide)




(1) 1-Chloro-3-((R)-2,2,2-trifluoro-1-methylethoxymethyl)benzene



Under argon atmosphere, a suspension of sodium hydride (77mg, 60 wt% oil dispersion) in N,N-dimethylformamide (2.0 ml) was cooled to 0°C, a solution of (R)-1,1,1-trifluoropropan-2-ol (213 mg) in N,N-dimethylformamide (2.0 ml) was added thereto, and the mixture was stirred for 15 minutes. Then, a solution of 1-bromomethyl-3-chlorobenzene (311 mg) in N,N-dimethylformamide (2.0 ml) was added thereto, and then the mixture was stirred at room temperature for 2.5 hours. To this reaction mixture was added water, and the mixture was extracted with ethyl acetate. This organic layer was washed with a saturated aqueous solution of sodium chloride, dried over magnesium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel chromatography (eluent: n-hexane/ ethyl acetate = 1/0 to 19/1) to give the titled compound (307 mg). ¹H-NMR (CDCl₃) δ: 1.35 (d, 3H, J = 6.47 Hz), 3.84 (qq, 1H, J = 6.47, 9.40 Hz), 4.62 (d, 1H, J = 12.00Hz), 4.73 (d, 1H, J = 12.00 Hz), 7.21-7.24 (m, 1H), 7.27-7.30 (m, 2H), 7.34 (s 1H)

[0321]

(2) 2-[3-Chloro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-4,4,5,5-tetramethyl[1,3,2]dioxaborolane




Under argon atmosphere, to a solution of 1-chloro-3-((R)-2,2,2-trifluoro-1-methylethoxymethyl)benzene (307 mg) in tetrahydrofuran (6.0 ml) were sequentially added bispinacolate diboron (362 mg), 4,4'-di-tert-butyl-2,2'-dipyridyl (18 mg) and di-μ-methoxobis(1,5-cyclooctadiene)diiridium (I) (22 mg) at room temperature, and then the mixture was heated to reflux for 13 hours. After cooling this reaction mixture to room temperature, a saturated aqueous solution of sodium hydrogen carbonate and ethyl acetate were added thereto, the mixture was filtered through Celite. The filtrate was extracted with ethyl acetate. The resulting organic layer was washed with a

saturated aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel thin-layer chromatography (eluent: n-hexane/ethyl acetate = 9/1) to give the titled compound (374 mg).

¹H-NMR (CDCl₃) δ: 1.35 (m, 15H), 3.83 (qq, 1H, J = 6.47, 9.40 Hz), 4.59 (d, 1H, J = 12.00Hz), 4.73 (d, 1H, J = 12.00 Hz), 7.46 (s, 1H), 7.59 (d, 1H, J = 9.60 Hz), 7.72 (s, 1H).

[0322]


(3) 5-[3-Chloro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1-phenyl-1H-pyrazol-3-ylamine



Under argon atmosphere, to 2-[3-chloro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-4,4,5,5-tetramethyl[1,3,2]dioxaborolane (269 mg) were sequentially added a solution of 5-iodo-1-phenyl-1H-pyrazol-3-ylamine (181 mg) prepared according to the same procedures as Preparation 2 in 1,2-dimethoxyethane (2.0 ml), a 2M aqueous solution of sodium carbonate (1.0 ml), tricyclohexylphosphine (36 mg) and palladium (II) acetate (15 mg) at room temperature, and the mixture was stirred at 100°C for 2 hours. This reaction mixture was cooled to room temperature, a saturated aqueous solution of sodium hydrogen carbonate and ethyl acetate were added thereto, the mixture was filtered through Celite, and subjected to elution with ethyl acetate. The filtrate was extracted with ethyl acetate. The resulting organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel thin-layer chromatography (eluent: n-hexane/ethyl acetate = 3/7) to give the titled compound (196 mg).

[0323]

(4) (S)-5-Oxopyrrolidine-3-carboxylic acid {5-[3-chloro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1-phenyl-1H-pyrazol-3-yl} amide




To a solution of 5-[3-chloro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1-phenyl-1H-pyrazol-3-ylamine (79 mg) in N,N-dimethylacetamide (1.0 ml) were sequentially added (S)-5-oxopyrrolidine-3-carboxylic acid (31 mg) prepared according to the same procedures as Preparation 14 and WSC·HCl (47 mg) at room temperature, and the mixture was stirred overnight. To this reaction mixture was added water, and then the precipitated solid was collected by filtration. This solid was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 9/1). To the resulting solid were added water and a small amount of methanol, and the mixture was stirred. A solid was collected from this suspension by filtration and was dried under reduced pressure to give the titled compound (60 mg).


¹H-NMR (DMSO-D₆) δ: 1.18 (d, 3H, J = 6.5 Hz), 2.36-2.38 (m, 2H), 3.32-3.33 (m, 1H), 3.37-3.41 (m, 1H), 3.46-3.49 (m, 1H), 4.10 (qq, 1H, J = 6.5, 9.4 Hz), 4.57 (d, 1H, J = 12.7 Hz), 4.61 (d, 1H, J = 12.7 Hz), 6.98 (s, 1H), 7.19-7.20 (m, 2H), 7.25-7.27 (m, 2H), 7.35-7.45 (m, 4H), 7.63 (s, 1H), 10.88 (s, 1H).

[0324]

[Preparation 23]: Example 242: Synthesis of ((S)-5-oxopyrrolidine-3-carboxylic acid {5-[3-chloro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1-(4-fluorophenyl)-1H-pyrazol-3-yl} amide)



(1) 5-[3-Chloro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1-(4-fluorophenyl)-1H-pyrazol-3-ylamine




To a solution of 1-(4-fluorophenyl)-5-iodo-1H-pyrazol-3-ylamine (50 mg) prepared according to the same procedures as Preparation 18 in 1,2-dimethoxyethane (0.5 ml) were sequentially added 2-[3-chloro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (72 mg) prepared according to the same procedures as Preparation 22, a 2M aqueous solution of sodium carbonate (0.25 ml), tricyclohexylphosphine (9.3 mg) and palladium (II) acetate (3.7 mg) at room temperature, and the mixture was stirred at 100°C for 2 hours. This reaction mixture was cooled to room temperature, filtered through Celite, and subjected to elution with ethyl acetate. The filtrate was extracted with ethyl acetate, the resulting organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/1) to give the titled compound (79 mg).

¹H-NMR (CDCl₃) δ: 1.29 (d, 3H, J = 6.47 Hz), 3.75 (qq, 1H, J = 6.47, 9.40 Hz), 3.78 (br s, 2H), 4.50 (d, 1H, J = 12.02 Hz), 4.61 (d, 1H, J = 12.02 Hz), 5.93 (s, 1H), 6.99-7.02 (m, 3H), 7.18-7.20 (m, 3H), 7.26-7.28 (m, 1H).

[0325]

(2) (S)-5-Oxopyrrolidine-3-carboxylic acid {5-[3-chloro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1-(4-fluorophenyl)-1H-pyrazol-3-yl} amide




To a solution of 5-[3-chloro-5-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1-(4-fluorophenyl)-1H-pyrazol-3-ylamine (79 mg) in chloroform (0.4 ml) were sequentially added (S)-5-oxopyrrolidine-3-carboxylic acid (26 mg) prepared according to the same procedures as Preparation 14 and WSC·HCl (47 mg) at room temperature, and the mixture was stirred for 2 hours. This reaction mixture was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 9/1). To the resulting solid were added water and a small amount of methanol, and the mixture was stirred. A solid was collected from this suspension by filtration and was dried under reduced pressure to give the titled compound (43 mg).


¹H-NMR (DMSO-D₆) δ: 1.18 (d, 3H, J = 6.5 Hz), 2.36-2.38 (m, 2H), 3.32-3.33 (m, 1H), 3.37-3.41 (m, 1H), 3.46-3.49 (m, 1H), 4.10 (qq, 1H, J = 6.5, 9.4 Hz), 4.57 (d, 1H, J = 12.7 Hz), 4.61 (d, 1H, J = 12.7 Hz), 6.97 (s, 1H), 7.13 (s, 1H), 7.23-7.32 (m, 5H), 7.37 (s, 1H), 7.61 (s, 1H), 10.86 (s, 1H).

[0326]

[Preparation 24]: Example 243 Synthesis of (S)-5-oxopyrrolidine-3-carboxylic acid [5-[3-chloro-5-(2,2,2-trifluoro-1,1-dimethylethoxymethyl)phenyl]-1-phenyl-1H-pyrazol-3-yl]amide)



(1) 1-Chloro-3-(2,2,2-trifluoro-1,1-dimethylethoxymethyl)benzene




To a suspension cooled to 0°C of sodium hydride (126mg, 60 wt.% oil dispersion) in N,N-dimethylformamide (5 ml) was added dropwise 1,1,1-trifluoro-2-methylpropan-2-ol (320 μL) under argon atmosphere, and the mixture was stirred for 15 minutes. To this reaction mixture was added dropwise 1-bromomethyl-3-chlorobenzene (320 μL), and the mixture was stirred for 3 hours. To this reaction mixture was added water, and the mixture was extracted with ethyl acetate. This organic layer was washed with water and a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/0 to 40/1) to give the titled compound (389 mg).

¹H-NMR (CDCl₃) δ: 1.45 (s, 6H), 4.50 (s, 2H), 7.15-7.30 (m, 3H), 7.34 (s, 1H).

[0327]


(2) 2-[3-Chloro-5-(2,2,2-trifluoro-1,1-dimethylethoxymethyl)phenyl]-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane



Under argon atmosphere, to a solution of 1-chloro-3-(2,2,2-trifluoro-1,1-dimethylethoxymethyl)benzene (389 mg) in tetrahydrofuran (4 ml) were sequentially added bispinacolate diboron (430 mg), 4,4'-di-tert-butyl-2,2'-dipyridyl (21 mg), di- μ -methoxobis(1, 5-cyclooctadiene)diiridium (I) (29 mg) at room temperature, and the mixture was heated to reflux for 13 hours. This reaction mixture was cooled to room temperature, a saturated aqueous solution of sodium hydrogen carbonate was added thereto, and the mixture was extracted with ethyl acetate. This organic layer was dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 100/1 to 20/1) to give the titled compound (481 mg).

$^1\text{H-NMR}$ (CDCl_3) δ : 1.34 (s, 12H), 1.44 (s, 6H), 4.57 (s, 2H), 7.45 (s, 1H), 7.57 (s, 1H), 7.69 (s, 1H). [0328]


(3) 5-[3-Chloro-5-(2,2,2-trifluoro-1,1-dimethylethoxymethyl)phenyl]-1-phenyl-1H-pyrazol-3-ylamine



Under argon atmosphere, to a solution of 5-iodo-1-phenyl-1H-pyrazol-3-ylamine (40 mg) prepared according to the same procedures as Preparation 2 in 1,2-dimethoxyethane (1.0ml) were sequentially added 2-[3-chloro-5-(2,2,2-trifluoro-1,1-dimethylethoxymethyl)phenyl]-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane (64 mg), palladium (II) acetate (3 mg), tricyclohexylphosphine (8 mg) and a 2M aqueous solution of sodium carbonate (0.5ml) at room temperature, and the mixture was stirred at 100°C for 13 hours. This reaction mixture was cooled to room temperature, and then purified by silica gel column chromatography (eluent: n-hexane/ ethyl acetate 3/2 to 1/1) to give the titled compound (62 mg).

$^1\text{H-NMR}$ (CDCl_3) δ : 1.35 (s, 6H), 3.77 (br s, 2H), 4.46 (s, 2H), 5.93 (s, 1H), 7.00 (s, 1H), 7.15 (s, 1H), 7.20-7.26 (m, 4H), 7.28-7.33 (m, 2H). [0329]

(4) (S)-5-Oxopyrrolidine-3-carboxylic acid {5-[3-chloro-5-(2,2,2-trifluoro-1,1-dimethylethoxymethyl)phenyl]-1-phenyl-1H-pyrazol-3-yl} amide




Under argon atmosphere, to a solution of 5-[3-chloro-5-(2,2,2-trifluoro-1,1-dimethylethoxymethyl)phenyl]-1-phenyl-1H-pyrazol-3-ylamine (62 mg) in N,N-dimethylacetamide (1 ml) were sequentially added (S)-5-oxopyrrolidine-3-carboxylic acid (26 mg) prepared according to the same procedures as Preparation 14, WSC·HCl (40 mg) at room temperature, and the mixture was stirred for 1 hour. To this reaction mixture was added water, and the mixture was extracted with ethyl acetate. This organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 9/1). To the resulting solid were added water and a small amount of methanol, and the mixture was stirred. A solid was collected from this suspension by filtration, and was dried under reduced pressure to give the titled compound (35 mg).


¹H-NMR (DMSO-D₆) δ: 1.32 (s, 6H), 2.37-2.42 (m, 2H), 3.33-3.53 (m, 3H), 4.53 (s, 2H), 6.98 (s, 1H), 7.16 (s, 1H), 7.19 (s, 1H), 7.23-7.28 (m, 2H), 7.33-7.45 (m, 4H), 7.63 (s, 1H), 10.89 (s, 1H).

[0330]

[Preparation 25]: Example 244: Synthesis of ((S)-5-oxopyrrolidine-3-carboxylic acid {1-(4-chlorophenyl)-5-[3-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-yl} amide)



(1) 2-(3-Bromomethylphenyl)-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane




To a solution of 3-(bromomethyl)phenyl boronic acid (3.13 g) in ethyl acetate (45 ml) were sequentially added pinacol (1.72 g) and anhydrous magnesium sulfate (15 g) at room temperature, and the mixture was stirred for 8 hours. The insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure to give the titled compound (4.43 g).

¹H-NMR (CDCl₃) δ: 1.33 (s, 12H), 4.49 (s, 2H), 7.31-7.37 (m, 1H), 7.46-7.50 (m, 1H), 7.70-7.74 (m, 1H), 7.79-7.82 (m, 1H).

[0331]

(2) 4,4,5,5-Tetramethyl-2-[3-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-[1,3,2]dioxaborolane




Under argon stream, to a solution of (R)-1,1,1-trifluoropropan-2-ol (3.32 g) in N,N-

dimethylformamide (43 ml) was added portionwise sodium hydride (1.16 g, 60 wt% oil dispersion) at 0°C, and the mixture was stirred for 20 minutes. Then, 2-(3-bromomethylphenyl)-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane (4.32 g) was added thereto, and then the mixture was stirred at room temperature overnight. To this reaction mixture were sequentially added water and a 2M aqueous solution of hydrochloric acid, and the mixture was extracted with ethyl acetate. This organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/0 to 19/1) to give the titled compound (3.43 g).

¹H-NMR (CDCl₃) δ: 1.31 (d, 3H, J = 6.6 Hz), 1.35 (s, 12H), 3.77-3.89 (m, 1H), 4.62 (d, 1H, J = 11.7 Hz), 4.76 (d, 1H, J = 11.7 Hz), 7.35-7.43 (m, 1H), 7.46-7.54 (m, 1H), 7.71-7.81 (m, 2H).

[0332]


(3) 1-(4-Chlorophenyl)-5-[3-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-ylamine



To a solution of 1-(4-chlorophenyl)-5-iodo-1H-pyrazol-3-ylamine (40 mg) prepared in Preparation 16 in 1,4-dioxane (0.4 ml) were sequentially added 4,4,5,5-tetramethyl-2-[3-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-[1,3,2]dioxaborolane (50 mg), palladium (II) acetate (3 mg), tricyclohexylphosphine (7 mg) and a 2M aqueous solution of potassium carbonate (0.2 ml) at room temperature, and the mixture was stirred at 105°C for 4 hours. After removing an aqueous layer of the reaction mixture, an organic layer was filtered through silica gel (5 g), and subjected to elution with ethyl acetate. The filtrate was concentrated under reduced pressure, the resulting residue was purified by silica gel thin-layer chromatography (eluent: n-hexane/ethyl acetate = 1/1) to give the titled compound.

[0333]

(4) (S)-5-Oxopyrrolidine-3-carboxylic acid {1-(4-chlorophenyl)-5-[3-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-yl} amide




To a solution of 1-(4-chlorophenyl)-5-[3-((R)-2,2,2-trifluoro-1-methylethoxymethyl)phenyl]-1H-pyrazol-3-ylamine (about 0.125 mmol) prepared in the previous step in N,N-dimethylformamide (1 ml) were sequentially added (S)-5-oxopyrrolidine-3-carboxylic acid prepared according to the same procedures as preparation 14 (18 mg) and WSC·HCl (29 mg) at room temperature, and the mixture was stirred for 3 hours. To this reaction mixture was added water, and the mixture was

extracted with ethyl acetate. The resulting organic layer was concentrated under reduced pressure, the resulting residue was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 10/1). To the resulting solid were added water and a small amount of methanol, and the mixture was stirred. A solid was collected from this suspension by filtration, and was dried under reduced pressure to give the titled compound (33 mg).


¹H-NMR (CDCl₃) δ: 1.29 (d, 3H, J = 6.8 Hz), 2.61 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.8 Hz), 3.30-3.40 (m, 1H), 3.61 (t, 1H, J = 8.8 Hz), 3.68-3.82 (m, 2H), 4.57 (d, 1H, J = 12.0 Hz), 4.68 (d, 1H, J = 12.0 Hz), 5.90 (s, 1H), 7.04 (s, 1H), 7.15-7.20 (m, 3H), 7.24 (br s, 1H), 7.28-7.35 (m, 4H), 8.38 (s, 1H).

[0334]

[Preparation 26]: Example 245: Synthesis of ((R)-2-oxoimidazolidine-4-carboxylic acid [5-(3-butylphenyl)-1-phenyl-1H-pyrazol-3-yl]amide)



(1) 1-Bromo-3-butylbenzene



Under nitrogen stream, to a solution of 1-bromo-3-bromomethylbenzene (25 g) in tetrahydrofuran (200 ml) were sequentially added dropwise n-propylmagnesium bromide (60 ml, 2M solution in tetrahydrofuran) and lithium tetrachlorocuprate (5 ml, 0.1M solution in tetrahydrofuran) at -78°C. Then, this reaction mixture was stirred at 0°C for 2 hours. Under ice-cooling, to this reaction mixture were sequentially added a saturated aqueous solution of ammonium chloride (100 ml) and water (100 ml), and the mixture was extracted with ethyl acetate (200 ml). The resulting aqueous layer was again extracted with ethyl acetate (100 ml), the combined organic layer was washed with a saturated aqueous solution of sodium chloride (100 ml), and dried over anhydrous sodium sulfate. This organic layer was concentrated under reduced pressure, the resulting residue was purified by silica gel chromatography (eluent: n-hexane/ ethyl acetate = 1/0 to 100/1) to give the titled compound (8.5 g).

¹H-NMR (CDCl₃) δ: 0.91 (t, 3H, J = 7.4 Hz), 1.31-1.35 (m, 2H), 1.55-1.59 (m, 2H), 2.56 (t, 2H, J = 7.8 Hz), 7.06-7.13 (m, 2H), 7.27-7.32 (m, 2H).

[0335]

(2) 2-(3-Butylphenyl)-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane




Under argon atmosphere, to a solution of 1-bromo-3-butylbenzene (1.44 g) in 1,4-dioxane (15 ml) were added bispinacolate diboron (1.97 g), [1,1'-

bis(diphenylphosphino)ferrocene]dichloropalladium (II) dichloromethane adduct (287 mg) and potassium acetate (2.07 g) at room temperature, and the mixture was stirred at 80°C for 15 hours. The mixture was cooled to room temperature, and the insoluble substance was filtered off through Celite. The filtrate was washed with water and a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 30/1 to 10/1) to give the titled compound (1.37 g).

¹H-NMR (CDCl₃) δ: 0.92 (t, 3H, J = 7.3 Hz), 1.31-1.41 (m, 14H), 1.59-1.61 (m, 2H), 2.61 (t, 2H, J = 7.9 Hz), 7.24-7.30 (m, 2H), 7.59-7.65 (m, 2H).

[0336]

(3) 5-(3-Butylphenyl)-1-phenyl-1H-pyrazol-3-ylamine




To a solution of 5-iodo-1-phenyl-1H-pyrazol-3-ylamine (1.00 g) prepared according to the same procedures as Preparation 2 in 1,2-dimethoxyethane (20 ml) were sequentially added 2-(3-butylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.37 g), a 2M aqueous solution of sodium carbonate (10 ml), tricyclohexylphosphine (197 mg) and palladium acetate (79 mg) at room temperature, and the mixture was stirred at reflux for 3 hours. This reaction mixture was cooled to room temperature, a saturated aqueous solution of sodium hydrogen carbonate was added thereto, and then the insoluble substance was filtered off through Celite. The filtrate was extracted with ethyl acetate, the resulting organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 2/1 to 2/3) to give the titled compound (880 mg).

¹H-NMR (CDCl₃) δ: 0.85 (t, 3H, J = 7.3 Hz), 1.17-1.27 (m, 2H), 1.40-1.47 (m, 2H), 2.49 (t, 2H, J = 7.6 Hz), 3.75 (br s, 2H), 5.90 (s, 1H), 6.97 (s, 1H), 7.02-7.12 (m, 2H), 7.13-7.29 (m, 6H).

[0337]

(4) (R)-2-Oxoimidazolidine-4-carboxylic acid [5-(3-butylphenyl)-1-phenyl-1H-pyrazol-3-yl]amide




To a solution of 5-(3-butylphenyl)-1-phenyl-1H-pyrazol-3-ylamine (40 mg) in N,N-dimethylformamide (0.4 ml) were sequentially added (R)-2-oxoimidazolidine-4-carboxylic acid (21 mg) prepared by the same procedures as Preparation 20, HATU (63 mg), diisopropylethylamine (29 μL) at room temperature, and the mixture was stirred for 2 hours. To this reaction mixture was added water, and the mixture was extracted with ethyl acetate. The resulting organic layer was washed with water and a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and

then concentrated under reduced pressure. The resulting residue was purified by silica gel thin-layer chromatography (eluent:n-hexane/acetone=1/5). To the resulting solid was added diisopropyl ether, and the mixture was stirred. A solid was collected from this suspension by filtration, and was dried under reduced pressure to give the titled compound (12 mg).


¹H-NMR (DMSO-D₆) δ: 0.82 (t, 3H, J = 7.4 Hz), 1.15 (tq, 2H, J = 7.4 Hz, 7.4Hz), 1.39 (tt, 2H, J = 7.4 Hz, 7.4Hz), 2.48 (t, 2H, J = 7.4 Hz), 3.36 (dd, 1H, J = 9.2 Hz, 5.6Hz), 3.59 (dd, 1H, J = 9.2 Hz, 9.2Hz), 4.32 (dd, 1H, J = 9.2 Hz, 5.6Hz), 6.33 (s, 1H), 6.60 (s, 1H), 6.90 (s, 1H), 6.98 (s, 1H), 7.09 (d, 1H, J = 7.6 Hz), 7.17 (d, 1H, J = 7.6 Hz), 7.20-7.30 (m, 3H), 7.32-7.43 (m, 3H), 10.70 (s, 1H).

[0338]

[Preparation 27]: Example 246: Synthesis of ((S)-5-oxopyrrolidine-3-carboxylic acid {1-phenyl-5-[3-(2,2,2-trifluoro-1-trifluoromethylethoxymethyl)phenyl]-1H-pyrazol-3-yl} amide)




(1) 3-(1,1,1,3,3,3-Hexafluoropropan-2-yloxy)phenyl boronic acid



A suspension of sodium hydride (149mg, 60 wt.% oil dispersion) in N,N-dimethylformamide (3.0 ml) was cooled to 0°C, (R)-1,1,1,3,3,3-hexafluoropropan-2-ol (392 mg) was added thereto, and the mixture was stirred for 15 minutes. Then, 3-(bromomethyl)phenyl boronic acid (200 mg) was added thereto, and then the mixture was stirred at room temperature for 2 hours. To this reaction mixture were sequentially added water and a 1M aqueous solution of hydrochloric acid, and the mixture was extracted with ethyl acetate. This organic layer was washed with a saturated aqueous solution of sodium chloride, dried over magnesium sulfate, and then concentrated under reduced pressure to give a crude product of the titled compound (1.0 g).

[0339]

(2) 1-Phenyl-5-[3-(2,2,2-trifluoro-1-trifluoromethylethoxymethyl)phenyl]-1H-pyrazol-3-ylamine




To 3-(1,1,1,3,3,3-hexafluoropropan-2-yloxy)phenyl boronic acid (1.0 g) were sequentially added a solution of 5-iodo-1-phenyl-1H-pyrazol-3-ylamine (180 mg) prepared according to the same procedures as Preparation 2 in 1,2-dimethoxyethane (2.0 ml), a 2M aqueous solution of sodium carbonate (1.0 ml), tricyclohexylphosphine (35 mg) and palladium (II) acetate (18 mg) at room

temperature, and the mixture was stirred at 100°C for 2 hours. This reaction mixture was cooled to room temperature, filtered through Celite, and subjected to elution with ethyl acetate. The filtrate was extracted with ethyl acetate, the resulting organic layer was washed with a saturated aqueous solution of sodium chloride, dried over magnesium sulfate, and then concentrated under reduced pressure.

The resulting residue was purified by silica gel thin-layer chromatography (eluent: n-hexane/ethyl acetate = 3/7) to give the titled compound (151 mg).

[0340]

(3) (S)-5-Oxopyrrolidine-3-carboxylic acid {1-phenyl-5-[3-(2,2,2-trifluoro-1-trifluoromethylethoxymethyl)phenyl]-1H-pyrazol-3-yl} amide




To a solution of 1-phenyl-5-[3-(2,2,2-trifluoro-1-trifluoromethyl-ethoxymethyl)phenyl]-1H-pyrazol-3-ylamine (78 mg) in N,N-dimethylacetamide (1.0 ml) were sequentially added (S)-5-oxopyrrolidine-3-carboxylic acid (29 mg) prepared according to the same procedures as Preparation 5 and WSC·HCl (44 mg) at room temperature, and the mixture was stirred overnight. To this reaction mixture was added water, and then the precipitated solid was collected by filtration. The resulting solid was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 9/1). To the resulting solid were added water and a small amount of methanol, and the mixture was stirred. A solid was collected from this suspension by filtration, and was dried under reduced pressure to give the titled compound (54 mg).


¹H-NMR (DMSO-D₆) δ: 2.36-2.38 (m, 2H), 3.32-3.33 (m, 1H), 3.37-3.41 (m, 1H), 3.46-3.49 (m, 1H), 4.85 (m, 2H), 5.53-5.59 (m, 1H), 6.94 (s, 1H), 7.17-7.18 (m, 1H), 7.22-7.24 (m, 2H), 7.34- 7.41 (m, 6H), 7.63 (s, 1H), 10.88 (s, 1H).

[0341]

[Preparation 28]: Example 247: Synthesis of ((S)-5-oxopyrrolidine-3-carboxylic acid {1-(4-fluorophenyl)-5-[3-(2,2,2-trifluoroethoxymethyl)phenyl]-1H-pyrazol-3-yl} amide)



(1) 1-Bromo-3-(2,2,2-trifluoroethoxymethyl)benzene




Under nitrogen atmosphere, a suspension of sodium hydride (352mg, 60 wt.% oil dispersion)

in tetrahydrofuran (10 ml) was cooled to 0°C, 2,2,2-trifluoroethanol (0.63 ml) was added thereto, and the mixture was stirred for 5 minutes. Then, 3-bromobenzyl bromide (2.00 g) was added thereto, and then the mixture was stirred at reflux for 1 hour. This reaction mixture was cooled to room temperature, a saturated aqueous solution of ammonium chloride was added thereto, and the mixture was extracted with ethyl acetate. This organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/0 to 10/1) to give the titled compound (2.05 g).

¹H-NMR (CDCl₃) δ: 3.84 (q, 2H, J = 8.68 Hz), 4.65 (s, 2H), 7.22-7.28 (m, 2H), 7.45-7.47 (m, 1H), 7.49-7.51 (m, 1H).

[0342]

(2) 4,4,5,5-Tetramethyl-2-[3-(2,2,2-trifluoroethoxymethyl)phenyl]-[1,3,2]dioxaborolane




Under argon atmosphere, to a solution of 1-bromo-3-(2,2,2-trifluoroethoxymethyl)benzene (2.05 g) in 1,4-dioxane (21 ml) were sequentially added bispinacolate diboron (2.12 g), potassium acetate (2.24 g) and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium (II) dichloromethane adduct (310 mg) at room temperature, and the mixture was stirred at 80°C for 19 hours. This reaction mixture was cooled to room temperature, filtered through Celite, and then subjected to elution with ethyl acetate. The filtrate was concentrated under reduced pressure, the resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 30/1) to give the titled compound (1.69 g).

¹H-NMR (CDCl₃) δ: 1.35 (s, 12H), 3.81 (q, 2H, J = 8.67 Hz), 4.68 (s, 2H), 7.38-7.40 (m, 1H), 7.47-7.48 (m, 1H), 7.76-7.78 (m, 2H).

[0343]

(3) 1-(4-Fluorophenyl)-5-[3-(2,2,2-trifluoroethoxymethyl)phenyl]-1H-pyrazol-3-ylamine




Under argon atmosphere, to a solution of 1-(4-fluorophenyl)-5-iodo-1H-pyrazol-3-ylamine (70 mg) prepared in Preparation 18 in 1,2-dimethoxyethane (0.7ml) were sequentially added 4,4,5,5-tetramethyl-2-[3-(2,2,2-trifluoroethoxymethyl)phenyl]-[1,3,2]dioxaborolane (80 mg), a 2M aqueous solution of sodium carbonate (0.35 ml), tricyclohexylphosphine (13.0 mg) and palladium (II) acetate (5.2 mg) at room temperature, and the mixture was stirred at 100°C for 2 hours. This reaction mixture was cooled to room temperature, filtered through Celite, and subjected to elution with ethyl acetate. The filtrate was extracted with ethyl acetate, the resulting organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then

concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/1) to give the titled compound (43 mg).

¹H-NMR (CDCl₃) δ: 3.74 (q, 2H, J = 8.61 Hz), 3.77 (br s, 2H), 4.60 (s, 2H), 5.93 (s, 1H), 6.97-6.99 (m, 2H), 7.16-7.21 (m, 4H), 7.29-7.31 (m, 2H).

[0344]

(4) (S)-5-Oxopyrrolidine-3-carboxylic acid {1-(4-fluorophenyl)-5-[3-(2,2,2-trifluoroethoxymethyl)phenyl]-1H-pyrazol-3-yl}amide




To a solution of 1-(4-fluorophenyl)-5-[3-(2,2,2-trifluoroethoxymethyl)phenyl]-1H-pyrazol-3-ylamine (43 mg) in N,N-dimethylacetamide (0.4 ml) were sequentially added (S)-5-oxopyrrolidine-3-carboxylic acid (17 mg) prepared according to the same procedures as Preparation 5 and WSC·HCl (34 mg) at room temperature, and the mixture was stirred for 0.5 hours. To this reaction mixture was added water, and then the precipitated solid was collected by filtration. This solid was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 9/1). To the resulting solid were added water and a small amount of methanol, and the mixture was stirred. A solid was collected from this suspension by filtration, and was dried under reduced pressure to give the titled compound (26 mg).


¹H-NMR (DMSO-D₆) δ: 2.38-2.40 (m, 2H), 3.32-3.35 (m, 1H), 3.40-3.45 (m, 1H), 3.48-3.52 (m, 1H), 3.98 (q, 2H, J = 9.3 Hz), 4.61 (s, 2H), 6.93 (s, 1H), 7.18-7.40 (m, 8H), 7.63 (s, 1H), 10.86 (s, 1H).

[0345]

[Preparation 29]: Example 248: Synthesis of ((S)-5-oxopyrrolidine-3-carboxylic acid {1-(4-fluorophenyl)-5-[3-fluoro-5-(2,2,2-trifluoroethoxymethyl)phenyl]-1H-pyrazol-3-yl}amide)



(1) 1-Chloro-3-fluoro-5-(2,2,2-trifluoroethoxymethyl)benzene




Under argon stream, to a suspension of sodium hydride (601mg, 60 wt.% oil dispersion) in dimethylformamide (13 ml) was added dropwise trifluoroethanol (1.08 ml) at 0°C, and the mixture was stirred for 15 minutes. To this reaction solution was added dropwise a solution of 1-chloro-3-fluoro-5-chloromethylbenzene (2.23 g) in dimethylformamide (9 ml), and then the mixture was stirred

at room temperature for 1 hour. To this reaction mixture was added water, and the mixture was extracted with a mixed solvent of n-hexane/ ethyl acetate = 1/1. This organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/0 to 9/1) to give the titled compound (2.85 g).
¹H-NMR (CDCl₃) δ: 3.87 (q, 2H, J = 8.8 Hz), 4.64 (s, 2H), 6.97 (d, 1H, J = 8.8 Hz), 7.05 (dt, 1H, J = 8.8, 2.4 Hz), 7.13 (br s, 1H).

[0346]

(2) 2-[3-Fluoro-5-(2,2,2-trifluoroethoxymethyl)phenyl]-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane




Under argon atmosphere, to a solution of 1-chloro-3-fluoro-5-(2,2,2-trifluoroethoxymethyl)benzene (2.65 g) in 1,4-dioxane (60 ml) were sequentially added bispinacolate diboron (3.41 g), potassium acetate (3.33 g), 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl (521 mg) and tris(dibenzylideneacetone) dipalladium (0) (253 mg) at room temperature, and the mixture was stirred at reflux for 6 hours. This reaction mixture was cooled to room temperature, and then filtered through Celite, and subjected to elution with ethyl acetate. The filtrate was concentrated under reduced pressure, the resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/0 to 19/1). The resulting crude product was suspended in n-hexane, and the insoluble substance was filtered off. The filtrate was concentrated under reduced pressure to give the titled compound (2.71 g).

¹H-NMR (CDCl₃) δ: 1.35 (s, 12H), 3.83 (q, 2H, J = 8.8 Hz), 4.67 (s, 2H), 7.18 (d, 1H, J = 8.8 Hz), 7.44 (dd, 1H, J = 8.8, 2.4 Hz), 7.50 (br s, 1H).

[0347]

(3) 1-(4-Chlorophenyl)-5-[3-fluoro-5-(2,2,2-trifluoroethoxymethyl)phenyl]-1H-pyrazol-3-ylamine




To a solution of 1-(4-fluorophenyl)-5-iodo-1H-pyrazol-3-ylamine (38 mg) prepared in Preparation 18 in 1,4-dioxane (0.4 ml) were sequentially added 2-[3-fluoro-5-(2,2,2-trifluoroethoxymethyl)phenyl]-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane (50 mg), palladium (II) acetate (3 mg), tricyclohexylphosphine (7 mg) and a 2M aqueous solution of potassium carbonate (0.2 ml) at room temperature, and the mixture was stirred at 105°C for 4 hours. After removing an aqueous layer of the reaction mixture, an organic layer was filtered through silica gel (5 g), and subjected to elution with ethyl acetate. The filtrate was concentrated under reduced pressure, the resulting residue was purified by silica gel thin-layer chromatography (eluent: n-hexane/ ethyl acetate

1/1) to give the titled compound.

[0348]

(4) Synthesis of (S)-5-oxopyrrolidine-3-carboxylic acid {1-(4-fluorophenyl)-5-[3-fluoro-5-(2,2,2-trifluoroethoxymethyl)phenyl]-1H-pyrazol-3-yl} amide




To a solution of 1-(4-chlorophenyl)-5-[3-fluoro-5-(2,2,2-trifluoroethoxymethyl)phenyl]-1H-pyrazol-3-ylamine (about 0.125 mmol) prepared in the previous step in *N,N*-dimethylformamide (0.5 ml) were sequentially added (S)-5-oxopyrrolidine-3-carboxylic acid (18 mg) prepared according to the same procedures as preparation 14 and WSC·HCl (29 mg) at room temperature, and the mixture was stirred for 3 hours. To this reaction mixture was added water, and the mixture was extracted with ethyl acetate. This organic layer was concentrated under reduced pressure, the resulting residue was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 10/1). To the resulting solid were added water and a small amount of methanol, and the mixture was stirred. A solid was collected from this suspension by filtration and was dried under reduced pressure to give the titled compound (33 mg).


¹H-NMR (CDCl₃) δ: 2.61 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.8 Hz), 3.32-3.42 (m, 1H), 3.61 (t, 1H, J = 8.8 Hz), 3.71 (dd, 1H, J = 9.6, 7.2 Hz), 3.79 (q, 2H, J = 8.8 Hz), 4.59 (s, 2H), 6.04 (s, 1H), 6.89 (d, 1H, J = 9.2 Hz), 6.99-7.10 (m, 5H), 7.19-7.24 (m, 2H), 8.48 (s, 1H).

[0349]

[Preparation 30]: Example 249: Synthesis of (4-oxo-3-azabicyclo[3.1.0]hexane-1-carboxylic acid [1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-yl]amide)



(1) 2-(tert-Butyldimethylsilanyloxymethyl)-2-propen-1-ol




Under argon atmosphere, to a suspension of sodium hydride (490mg, 60 wt.% oil dispersion) in tetrahydrofuran (10 ml) were sequentially added 2-methylene-1,3-propanediol (1 ml) and tert-butyldimethylchlorosilane (2.22 g) at 0°C, and then the mixture was stirred at room temperature for 1 hour. To this reaction mixture was added a saturated aqueous solution of ammonium chloride, and the mixture was extracted with ethyl acetate. This organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-

hexane/ethyl acetate = 3/1) to give the titled compound (1.62 g).

$^1\text{H-NMR}$ (CDCl_3) δ : 0.09 (s, 6H), 0.91 (s, 9H), 1.96 (t, 1H, $J = 3.09$ Hz), 4.18 (d, 2H, $J = 3.09$ Hz), 4.25 (s, 2H), 5.08 (d, 1H, $J = 1.32$ Hz), 5.10 (d, 1H, $J = 1.32$ Hz).

[0350]

- 5 (2) [2-(tert-Butyldimethylsilanyloxymethyl)allyl]-(2,4-dimethoxybenzyl) amine




To a solution of 2-(tert-butyldimethylsilyloxy)prop-1-en-1-ol (1.58 g) in chloroform (16 ml) were sequentially added triethylamine (2.2 ml) and methanesulfonyl chloride (0.73 ml) at 0°C , and then the mixture was stirred for 2 hours. To this reaction solution were sequentially added triethylamine (1.2 ml) and 2,4-dimethoxybenzylamine (1.3 ml), and then the mixture was stirred at room temperature for 15 hours. To this reaction mixture was added water, and an organic layer was separated from the mixture, and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/1) to give the titled compound (1.14 g).

$^1\text{H-NMR}$ (CDCl_3) δ : 0.07 (s, 6H), 0.91 (s, 9H), 3.21 (s, 2H), 3.69 (s, 2H), 3.80 (s, 6H), 4.18 (s, 2H), 5.02 (s, 1H), 5.13 (s, 1H), 6.41-6.45 (m, 2H), 7.12 (d, 1H, $J = 7.91$ Hz).

[0351]

- (3) (E)-2-(2-Tosylhydrazono)acetic acid




To glyoxylic acid monohydrate (644 mg) was added water (7 ml). The resulting suspension was dissolved in water by heating to 65°C , a solution of p-toluenesulfonylhydrazide (1.30 g) in a 2.5M aqueous solution of hydrochloric acid (4.2 ml) was added thereto, and the mixture was stirred for 15 minutes. This reaction mixture was cooled to room temperature, the precipitated solid was collected by filtration, and washed with water. The resulting solid was dried under reduced pressure to give the titled compound (1.42 g).

$^1\text{H-NMR}$ (CDCl_3) δ : 2.46 (s, 3H), 7.14 (s, 1H), 7.38 (d, 2H, $J = 8.14$ Hz), 7.83 (d, 2H, $J = 8.14$ Hz), 8.46 (s, 1H).

[0352]

- (4) Diazoacetic acid 2,5-dioxopyrrolidin-1-yl ester




Under argon atmosphere, to a solution of (E)-2-(2-tosylhydrazono)acetic acid (1.42 g) in 1,4-dioxane (14 ml) were sequentially added N-hydroxysuccinimide (677 mg) and dicyclohexylcarbodiimide (1.21 g) at room temperature, and the mixture was stirred for 17 hours. This reaction mixture was filtered through Celite, and eluted with 1,4-dioxane. This filtrate was

concentrated under reduced pressure, the resulting residue was purified by silica gel column chromatography (eluent: n-hexane/diethyl ether=1/3) to give the titled compound (579 mg).

$^1\text{H-NMR}$ (CDCl_3) δ : 2.85 (s, 4H), 5.11 (s, 1H).

[0353]

- 5 (5) N-[2-(tert-Butyldimethylsilyloxymethyl)allyl]-2-diazo-N-(2,4-dimethoxybenzyl) acetamide




To a solution of [2-(tert-butyl(dimethyl)silyloxymethyl)allyl]-(2,4-dimethoxybenzyl)amine (384 mg) in chloroform (3.8 ml) were sequentially added triethylamine (0.23 ml) and diazoacetic acid 2,5-dioxopyrrolidin-1-yl ester (200 mg) at room temperature, and the mixture was stirred for 16 hours. This reaction mixture was concentrated under reduced pressure, the resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 4/1) to give the titled compound (223 mg).

$^1\text{H-NMR}$ (CDCl_3) δ : 0.06 (s, 6H), 0.90 (s, 9H), 3.79 (s, 3H), 3.80 (s, 3H), 3.88-3.91 (m, 2H), 4.08 (s, 2H), 4.93 (s, 1H), 5.04 (s, 1H), 5.18 (s, 1H), 6.43-6.47 (m, 3H).

[0354]

- 15 (6) 5-(tert-Butyldimethylsilyloxymethyl)-3-(2,4-dimethoxybenzyl)-3-azabicyclo[3.1.0]hexan-2-one




Under argon atmosphere, to a solution of N-[2-(tert-butyl(dimethyl)silyloxymethyl)allyl]-2-diazo-N-(2,4-dimethoxybenzyl)acetamide (43 mg) in benzonitrile (0.4 ml) was added rhodium (II) acetate dimer dihydrate (2.4 mg) at room temperature, and the mixture was stirred at 50°C for 24 hours. This reaction mixture was concentrated under reduced pressure, the resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 2/1) to give the titled compound (14.2 mg).

$^1\text{H-NMR}$ (CDCl_3) δ : 0.01 (s, 6H), 0.72 (dd, 1H, $J = 4.42, 3.61$ Hz), 0.85 (s, 9H), 1.11 (dd, 1H, $J = 8.84, 4.42$ Hz), 1.81 (dd, 1H, $J = 8.84, 3.61$ Hz), 3.16 (d, 1H, $J = 10.47$ Hz), 3.36 (d, 1H, $J = 10.47$ Hz), 3.62 (d, 1H, $J = 10.70$ Hz), 3.70 (d, 1H, $J = 10.70$ Hz), 3.79 (s, 3H), 3.79 (s, 3H), 4.25 (d, 1H, $J = 14.65$ Hz), 4.38 (d, 1H, $J = 14.65$ Hz), 6.40-6.44 (m, 2H), 7.06-7.09 (m, 1H).

[0355]

- 30 (7) 3-(2,4-Dimethoxybenzyl)-5-hydroxymethyl-3-azabicyclo[3.1.0]hexan-2-one




To a solution of 5-(tert-butyl dimethylsilyloxymethyl)-3-(2,4-dimethoxybenzyl)-3-azabicyclo[3.1.0]hexan-2-one (14.2 mg) in tetrahydrofuran (0.3 ml) was added n-butylammonium fluoride (54 μ l, 1.0M solution in tetrahydrofuran) at room temperature, and the mixture was stirred for 1 hour. This reaction mixture was concentrated under reduced pressure, the resulting residue was purified by silica gel column chromatography (eluent: ethyl acetate, to chloroform/methanol =9/1) to give the titled compound (9.2 mg).

$^1\text{H-NMR}$ (CDCl_3) δ : 0.77 (dd, 1H, $J = 4.84, 3.43$ Hz), 1.12 (dd, 1H, $J = 8.87, 4.84$ Hz), 1.85 (dd, 1H, $J = 8.87, 3.43$ Hz), 3.23 (d, 1H, $J = 10.07$ Hz), 3.45 (d, 1H, $J = 10.07$ Hz), 3.66 (d, 1H, $J = 11.69$ Hz), 3.70 (d, 1H, $J = 11.69$ Hz), 3.79 (s, 3H), 3.79 (s, 3H), 4.21 (d, 1H, $J = 14.51$ Hz), 4.43 (d, 1H, $J = 14.51$ Hz), 6.41-6.44 (m, 2H), 7.07-7.10 (m, 1H).

[0356]


(8) 3-(2,4-Dimethoxybenzyl)-4-oxo-3-azabicyclo[3.1.0]hexane-1-carbaldehyde



Under argon atmosphere, to a solution of 3-(2,4-dimethoxybenzyl)-5-hydroxymethyl-3-azabicyclo[3.1.0]hexan-2-one (9.2 mg) in dimethylsulfoxide/chloroform (2/1, 135 μ l) were sequentially added triethylamine (23 μ l) and sulfur trioxide-pyridine complex (10.6 mg) at room temperature, and the mixture was stirred for 2 hours. To this reaction mixture was added water, and the mixture was extracted with ethyl acetate. This organic layer was washed with water, a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure to give a crude product of the titled compound (10.8 mg).

[0357]

(9) 3-(2,4-Dimethoxybenzyl)-4-oxo-3-azabicyclo[3.1.0]hexane-1-carboxylic acid




Under argon atmosphere, to a solution of 3-(2,4-dimethoxybenzyl)-4-oxo-3-azabicyclo[3.1.0]hexane-1-carbaldehyde (10.8 mg) in tert-butanol/tetrahydrofuran/water (4/3/1, 0.5 ml) was added sodium dihydrogenphosphate dihydrate (20.2 mg) at room temperature. This reaction mixture was cooled to 0°C, 2-methyl-2-butene (18 μ l) and sodium chlorite (9.0 mg) were sequentially added thereto, and then the mixture was stirred at room temperature for 24 hours. This reaction mixture was cooled to 0°C, and then water was added thereto, and the mixture was extracted with

ethyl acetate. This organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 9/1) to give the titled compound (3.7 mg).

¹H-NMR (CDCl₃) δ: 0.69-0.71 (m, 1H), 1.68-1.71 (m, 1H), 2.25-2.28 (m, 1H), 3.11 (d, 1H, J = 11.28 Hz), 3.49 (s, 1H), 3.71 (s, 3H), 3.73 (s, 3H), 3.75 (d, 1H, J = 11.28 Hz), 3.98 (d, 1H, J = 13.30 Hz), 4.42 (d, 1H, J = 13.30 Hz), 6.33-6.38 (m, 2H), 6.95-6.98 (m, 1H).

[0358]


(10) 3-(2,4-Dimethoxybenzyl)-4-oxo-3-azabicyclo[3.1.0]hexane-1-carboxylic acid [1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-yl]amide



To a solution of 1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-ylamine (3.5 mg) prepared according to the same procedures as Preparation 2-2 in N,N-dimethylformamide (0.5 ml) were sequentially added 3-(2,4-dimethoxybenzyl)-4-oxo-3-azabicyclo[3.1.0]hexane-1-carboxylic acid (3.7 mg) and WSC·HCl (3.6 mg) at room temperature, and the mixture was stirred for 14 hours. To this reaction mixture were added diisopropylethylamine (3.3 μl) and HATU (7.3 mg), and the mixture was stirred at room temperature for additional 72 hours. To this reaction mixture was added water, and the mixture was extracted with ethyl acetate. This organic layer was washed with water and a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure to give the titled compound (6.3 mg).

[0359]

(11) 4-Oxo-3-azabicyclo[3.1.0]hexane-1-carboxylic acid [1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-yl]amide




To 3-(2,4-dimethoxybenzyl)-4-oxo-3-azabicyclo[3.1.0]hexane-1-carboxylic acid [1-phenyl-5-(3-propylphenyl)-1H-pyrazol-3-yl]amide (6.3 mg) were sequentially added anisole (2 μl) and trifluoroacetic acid (0.5 ml) at room temperature, and the mixture was stirred at 80°C for 3 hours. This reaction mixture was concentrated under reduced pressure, the resulting residue was purified by silica gel thin-layer chromatography (eluent: chloroform/methanol = 9/1) to give the titled compound (0.3 mg).

¹H-NMR (CDCl₃) δ: 0.83-0.88 (m, 4H), 1.22-1.28 (m, 3H), 1.95 (dd, 1H, J = 9.3, 4.8 Hz), 2.49 (t, 2H, J = 8.5 Hz), 3.60 (d, 1H, J = 10.1 Hz), 4.01 (d, 1H, J = 10.1 Hz), 5.14 (s, 1H), 6.99 (s, 1H), 7.04-7.34

(m, 9H), 7.83 (s, 1H).


[0360]

[Preparation 31]: Example 250: Synthesis of (4-oxo-5-azabisp[2.4]heptane-7-carboxylic acid {1-phenyl-5-[3-(2,2,2-trifluoroethoxymethyl)phenyl]-1H-pyrazol-3-yl} amide)



5

(1) 1-Acetylcyclopropanecarboxylic acid methyl ester



To a solution of methyl acetoacetate (3 ml) in acetone (30 ml) were sequentially added potassium carbonate (11.54 g) and 1,2-dibromoethane (2.4 ml) at room temperature, and the mixture was stirred at 65°C for 17 hours. This reaction mixture was filtered, the filtrate was concentrated under reduced pressure, and the resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 10/1) to give the titled compound (2.17 g).

¹H-NMR (CDCl₃) δ: 1.48 (s, 4H), 2.47 (s, 3H), 3.75 (s, 3H).

[0361]

(2) 1-(1-Trifluoromethanesulfonyloxyvinyl)cyclopropane carboxylic acid methyl ester



Under argon atmosphere, to a solution of diisopropylamine (2.37 ml) in tetrahydrofuran (11 ml) was added n-butyllithium (10.4ml, 1.6M solution in n-hexane) at 0°C, and the mixture was stirred for 0.5 hours. At -78°C, a solution of 1-acetylcyclopropane carboxylic acid methyl ester (2.17 g) in tetrahydrofuran (11 ml) was added thereto, and the mixture was stirred for 0.5 hours. Then, N-phenylbis(trifluoromethanesulfonimide) (6.00 g) was added thereto, and the mixture was stirred for 0.5 hours, and then stirred at 0°C for additional 1 hour. To this reaction mixture was added water, and the mixture was extracted with ethyl acetate. This organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure to give the titled compound (4.36 g).

[0362]

(3) 1-(1-Methoxycarbonylvinyl)cyclopropanecarboxylic acid methyl ester



To a solution of 1-(1-trifluoromethanesulfonyloxyvinyl)cyclopropane carboxylic acid methyl ester (4.36 g) prepared in the previous step in N,N-dimethylformamide/methanol (1/2, 66 ml) were sequentially added triethylamine (6.4 ml) and bis (triphenylphosphine)palladium (II) dichloride (1.07 g) at room temperature, and the air in reaction vessel was replaced with carbon monoxide at normal


30

pressure. This reaction solution was stirred at 50°C for 18 hours, and then cooled to room temperature, and carbon monoxide in the reaction vessel was replaced with nitrogen. To this reaction mixture were added water and ethyl acetate, the mixture was filtered through Celite, and subjected to elution with ethyl acetate. The filtrate was extracted with ethyl acetate, the separated organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 8/1) to give the titled compound (720 mg).

¹H-NMR (CDCl₃) δ: 1.03 (dd, 2H, J = 7.33, 4.07 Hz), 1.51 (dd, 2H, J = 7.33, 4.07 Hz), 3.65 (s, 3H), 3.78 (s, 3H), 5.64 (d, 1H, J = 0.93 Hz), 6.31 (d, 1H, J = 0.93 Hz).

[0363]

(4) 5-(2,4-Dimethoxybenzyl)-4-oxo-5-azaspiro[2.4]heptane-7-carboxylic acid methyl ester




To a solution of 1-(1-methoxycarbonylvinyl)cyclopropanecarboxylic acid methyl ester (720 mg) in toluene (3.6 ml) was added 2,4-dimethoxybenzylamine (587 μl) at room temperature, and then the mixture was stirred at reflux for 20 hours. This reaction mixture was concentrated under reduced pressure, the resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 1/1) to give the titled compound (567 mg).

¹H-NMR (CDCl₃) δ: 0.81-0.84 (m, 2H), 1.09-1.12 (m, 1H), 1.22-1.28 (m, 1H), 3.10 (dd, 1H, J = 8.84, 5.23 Hz), 3.48 (dd, 1H, J = 10.00, 8.84 Hz), 3.66 (dd, 1H, J = 10.00, 5.23 Hz), 3.67 (s, 3H), 3.80 (s, 3H), 3.81 (s, 3H), 4.45 (d, 1H, J = 14.77 Hz), 4.51 (d, 1H, J = 14.77 Hz), 6.45-6.47 (m, 2H), 7.16-7.18 (m, 1H).

[0364]

(5) 5-(2,4-Dimethoxybenzyl)-4-oxo-5-azaspiro[2.4]heptane-7-carboxylic acid




To a solution of 5-(2,4-dimethoxybenzyl)-4-oxo-5-azaspiro[2.4]heptane-7-carboxylic acid methyl ester (2.02 g) in tetrahydrofuran /methanol (1/1, 12 ml) was added a 1M aqueous solution of sodium hydroxide (12 ml) at room temperature, and the mixture was stirred for 1 hour. To this reaction solution was added a 2M aqueous solution of hydrochloric acid (6 ml), and the mixture was stirred, and then concentrated under reduced pressure. The resulting residue was dissolved in ethyl acetate, the solution was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. To the resulting solid was added diisopropyl ether (8 ml), and the mixture was stirred. A solid was collected from this suspension by filtration, and was dried under reduced pressure to give the titled compound (1.59 g).

¹H-NMR (CDCl₃) δ: 0.83-0.88 (m, 1H), 0.95-1.00 (m, 1H), 1.12-1.18 (m, 1H), 1.21-1.28 (m, 1H), 3.10 (dd, 1H, J = 8.61, 5.12 Hz), 3.49 (dd, 1H, J = 10.23, 8.61 Hz), 3.64 (dd, 1H, J = 10.23, 5.12 Hz),

3.78 (s, 3H), 3.79 (s, 3H), 4.43 (d, 1H, $J = 14.65$ Hz), 4.50 (d, 1H, $J = 14.65$ Hz), 6.42-6.45 (m, 2H), 7.13-7.16 (m, 1H).

[0365]

(6) Optically active compound of 7-((R)-4-Benzyl-2-oxooxazolidine-3-carbonyl)-5-(2,4-dimethoxybenzyl)-5-azaspiro[2.4]heptan-4-one




To a solution of 5-(2,4-dimethoxybenzyl)-4-oxo-5-azaspiro[2.4]heptane-7-carboxylic acid (1.59 g) in chloroform (16 ml) were sequentially added (R)-4-benzyl-2-oxazolidinone (1.11 g), WSC·HCl (1.20 g) and *N,N*-dimethyl-4-aminopyridine (318 mg) at room temperature, and the mixture was stirred for 4 hours. To this reaction mixture was added water, and the organic layer was separated from the solution, and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: *n*-hexane/ethyl acetate = 2/1) to separate a low-polarity component as the titled compound (1.39 g).

$^1\text{H-NMR}$ (CDCl_3) δ : 0.77-0.82 (m, 1H), 1.11-1.17 (m, 2H), 1.26-1.30 (m, 1H), 2.67 (dd, 1H, $J = 13.23$, 10.37 Hz), 3.38 (dd, 1H, $J = 13.23$, 3.42 Hz), 3.49-3.55 (m, 1H), 3.65 (dd, 1H, $J = 10.37$, 3.42 Hz), 3.81 (s, 3H), 3.81 (s, 3H), 4.17-4.19 (m, 2H), 4.34-4.36 (m, 1H), 4.48 (d, 1H, $J = 15.00$ Hz), 4.57 (d, 1H, $J = 15.00$ Hz), 4.64-4.68 (m, 1H), 6.45-6.50 (m, 2H), 7.21-7.24 (m, 3H), 7.29-7.37 (m, 3H).

[0366]

(7) Optically active compound of 5-(2,4-Dimethoxybenzyl)-4-oxo-5-azaspiro[2.4]heptane-7-carboxylic acid




To a solution of lithium hydroxide monohydrate (142 mg) in water (5 ml) was added 30 wt% aqueous solution of hydrogen peroxide (0.8ml) at -5°C , and the mixture was stirred for 15 minutes. To this reaction mixture were sequentially added tetrahydrofuran (5 ml) and a solution of an optically active compound of 7-((R)-4-benzyl-2-oxooxazolidine-3-carbonyl)-5-(2,4-dimethoxybenzyl)-5-azaspiro[2.4]heptan-4-one (1.39 g) in tetrahydrofuran (10 ml), and the mixture was stirred for additional 1 hour. To this reaction mixture was added a solution of sodium hydrogen sulfite (815 mg) in water (5 ml), and the mixture was stirred at room temperature for 1 hour, and then extracted with ethyl acetate. This organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: ethyl acetate only). To the resulting solid was added diisopropyl ether/ethyl acetate/*n*-hexane (10/1/5, 10 ml), and the mixture was stirred. A solid was collected from this suspension by filtration, and dried under reduced pressure to give the titled compound (372 mg).

$^1\text{H-NMR}$ ($\text{DMSO-}d_6$) δ : 0.83-0.89 (m, 3H), 0.94-0.97 (m, 1H), 3.07-3.11 (m, 1H), 3.42-3.51 (m, 2H),

3.75 (s, 3H), 3.78 (s, 3H), 4.28 (d, 1H, J = 14.89 Hz), 4.34 (d, 1H, J = 14.89 Hz), 6.49 (dd, 1H, J = 8.37, 2.33 Hz), 6.57 (d, 1H, J = 2.33 Hz), 7.04 (d, 1H, J = 8.37 Hz), 12.56 (br s, 1H).

[0367]

(8) Optically active compound of 4-Oxo-5-azaspiro[2.4]heptane-7-carboxylic acid




To an optically active compound of 5-(2,4-dimethoxybenzyl)-4-oxo-5-azaspiro[2.4]heptane-7-carboxylic acid (372 mg) were sequentially added anisole (159 μ l) and trifluoroacetic acid (5.6 ml) at room temperature, and the mixture was stirred at 80°C for 4 hours. This reaction mixture was concentrated under reduced pressure, and diisopropyl ether (10 ml) was added thereto. The precipitated solid was collected by filtration to give the titled compound (144 mg).

$^1\text{H-NMR}$ (DMSO- D_6) δ : 0.77-0.82 (m, 3H), 0.86-0.91 (m, 1H), 3.12 (t, 1H, J = 6.75 Hz), 3.48 (d, 2H, J = 6.75 Hz), 7.70 (br s, 1H), 12.51 (br s, 1H).

[0368]

(9) 1-Phenyl-5-[3-(2,2,2-trifluoroethoxymethyl)phenyl]-1H-pyrazol-3-ylamine




To a solution of 5-iodo-1-phenyl-1H-pyrazol-3-ylamine (221 mg) prepared according to the same procedures as Preparation 2 in 1,2-dimethoxyethane (2.2 ml) were sequentially added 4,4,5,5-tetramethyl-2-[3-(2,2,2-trifluoroethoxymethyl)phenyl]-[1,3,2]dioxaborolane (269 mg) prepared in Preparation 28, a 2M aqueous solution of sodium carbonate (1.1 ml), tricyclohexylphosphine (43.4 mg) and palladium (II) acetate (17.4 mg) at room temperature, and the mixture was stirred at 100°C for 12 hours. To this reaction mixture was added a saturated aqueous solution of sodium hydrogen carbonate at room temperature, and then the mixture was filtered through Celite, and subjected to elution with ethyl acetate. The filtrate was extracted with ethyl acetate, the resulting organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate = 2/1) to give the titled compound (258 mg).

$^1\text{H-NMR}$ (CDCl_3) δ : 3.66 (q, 2H, J = 8.7 Hz), 3.77 (br s, 2H), 4.57 (s, 2H), 5.92 (s, 1H), 7.12-7.33 (m, 9H).

[0369]

(10) Optically active compound of 4-Oxo-5-aza bispiro[2.4]heptane-7-carboxylic acid {1-phenyl-5-[3-(2,2,2-trifluoroethoxymethyl)phenyl]-1H-pyrazol-3-yl} amide





To a solution of 1-phenyl-5-[3-(2,2,2-trifluoroethoxymethyl)phenyl]-1H-pyrazol-3-ylamine (19.5 mg) in N,N-dimethylacetamide (0.2 ml) were sequentially added an optically active compound of 4-oxo-5-azaspiro[2.4]heptane-7-carboxylic acid (10.5 mg) and WSC·HCl (16.1 mg) at room temperature, and the mixture was stirred for 0.5 hours. To this reaction mixture was added water, and then the precipitated solid was collected by filtration, and dried under reduced pressure to give the titled compound (15.4 mg).




¹H-NMR (DMSO-D₆) δ: 0.66-0.71 (m, 1H), 0.78-0.82 (m, 1H), 0.88-0.99 (m, 2H), 3.33-3.37 (m, 1H), 3.50-3.54 (m, 1H), 3.56-3.60 (m, 1H), 3.97 (q, 2H, J = 9.30 Hz), 4.61 (s, 2H), 6.93 (s, 1H), 7.15-7.17 (m, 1H), 7.22-7.26 (m, 3H), 7.31-7.41 (m, 5H), 7.72 (s, 1H), 10.73 (s, 1H).



[0370] Compounds of Examples 9 to 234 and 251 to 605 were prepared according to the same procedures as Preparations described in above. The compounds of Examples 9 to 234 are shown in the following Tables 1 together with Compounds of Examples 1 - 8. The compounds of Examples 251 to 605 are shown in the following Tables 2 together with Compounds of Examples 235 - 250.




[0371]




[Table 1]



Ex. No.	Molecular Structure	NMR	MS (M+H)	MS (M-H)	Comments
1		¹ H-NMR (DMSO-D ₆) δ : 1.08 (d, 3H, J = 7.3 Hz), 2.55 (dq, 1H, J = 9.2, 7.3 Hz), 3.04 (ddd, 1H, J = 9.2, 8.7, 8.5 Hz), 3.23 (dd, 1H, J = 9.2, 8.7 Hz), 3.45 (dd, 1H, J = 9.2, 8.5 Hz), 7.03 (s, 1H), 7.09 (s, 1H), 7.25-7.27 (m, 2H), 7.35-7.45 (m, 5H), 7.52 (t, 1H, J = 8.0 Hz), 7.68 (s, 1H), 10.99 (s, 1H).	445	443	(3R, 4R)-form
2		¹ H-NMR (DMSO-D ₆) δ : 1.08 (d, 3H, J = 7.1 Hz), 2.54 (dq, 1H, J = 9.3, 7.1 Hz), 3.04 (ddd, 1H, J = 9.3, 9.2, 8.4 Hz), 3.23 (dd, 1H, J = 9.2, 8.6 Hz), 3.44 (dd, 1H, J = 8.6, 8.4 Hz), 7.03 (s, 1H), 7.11 (s, 1H), 7.25-7.38 (m, 6H), 7.53 (dd, 1H, J = 8.2, 7.9 Hz), 7.68 (s, 1H), 10.98 (s, 1H).	463	461	(3R, 4R)-form




3		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 1.08 (d, 3H, J = 7.3 Hz), 2.54 (dq, 1H, J = 9.3, 7.3 Hz), 3.04 (ddd, 1H, J = 9.3, 8.9, 8.5 Hz), 3.23 (dd, 1H, J = 8.9, 8.7 Hz), 3.45 (dd, 1H, J = 8.7, 8.5 Hz), 7.04 (s, 1H), 7.05-7.07 (m, 1H), 7.14-7.18 (m, 2H), 7.22-7.26 (m, 1H), 7.39-7.48 (m, 3H), 7.56 (t, 1H, J = 8.0 Hz), 7.68 (s, 1H), 11.03 (s, 1H).</p>	463	461	(3R, 4R)-form
4		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 2.35-2.42 (m, 2H), 3.31-3.54 (m, 3H), 7.01 (s, 1H), 7.08 (s, 1H), 7.22-7.28 (m, 2H), 7.32-7.46 (m, 5H), 7.52 (t, 1H, J = 8.0 Hz), 7.64 (s, 1H), 10.90 (s, 1H).</p>	431	429	R-form
5		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.5, 7.3 Hz), 1.83-2.03 (m, 2H), 2.12-2.30 (m, 2H), 2.46 (t, 2H, J = 7.5 Hz), 2.84 (tt, 1H, J = 10.7, 3.6 Hz), 3.24-3.31 (m, 2H), 6.89 (s, 1H), 7.00 (s, 1H), 7.06 (d, 1H, J = 7.7 Hz), 7.17 (d, 1H, J = 7.7 Hz), 7.19-7.43 (m, 6H), 7.51 (s, 1H), 10.83 (s, 1H).</p>	403	401	<p>Optically active substance</p> <p>Optical isomer of Example 96</p>



6		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.5, 7.3 Hz), 2.46 (t, 3H, J = 7.5 Hz), 2.64 (s, 3H), 3.21 (dd, 1H, J = 8.8, 6.3 Hz), 3.54 (dd, 1H, J = 9.4, 8.8 Hz), 4.28 (dd, 1H, J = 9.4, 6.3 Hz), 6.44 (s, 1H), 6.92 (s, 1H), 7.01 (s, 1H), 7.08 (d, 1H, J = 7.0 Hz), 7.17 (d, 1H, J = 7.0 Hz), 7.20-7.44 (m, 6H), 11.02 (s, 1H).</p>	404	402	R-form
7		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 0.77 (t, 3H, J = 7.3 Hz), 1.38-1.50 (m, 2H), 2.46 (t, 2H, J = 7.5 Hz), 2.69 (dd, 1H, J = 17.4, 5.9 Hz), 2.81 (dd, 1H, J = 17.4, 9.0 Hz), 3.56-3.65 (m, 1H), 4.34 (dd, 1H, J = 8.8, 5.0 Hz), 4.47-4.55 (m, 1H), 6.89 (s, 1H), 7.00 (s, 1H), 7.07 (d, 1H, J = 7.5 Hz), 7.17 (d, 1H, J = 7.1 Hz), 7.20-7.30 (3H, m), 7.31-7.43 (3H, m), 11.00 (s, 1H).</p>	390	388	Racemate



8		¹ H-NMR (DMSO-D ₆) δ : 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.5, 7.3 Hz), 2.46 (t, 2H, J = 7.5 Hz), 3.35 (dd, 1H, J = 8.8, 6.2 Hz), 3.59 (dd, 1H, J = 9.4, 8.8 Hz), 4.32 (dd, 1H, J = 9.4, 6.2 Hz), 6.34 (s, 1H), 6.61 (s, 1H), 6.90 (s, 1H), 7.00 (s, 1H), 7.08 (d, 1H, J = 7.7 Hz), 7.15-7.43 (m, 7H), 10.71 (s, 1H).	390	388	R-form
9		¹ H-NMR (DMSO-D ₆) δ : 2.36-2.42 (m, 2H), 3.30-3.36 (m, 1H), 3.37-3.47 (m, 1H), 3.47-3.54 (m, 1H), 6.90 (s, 1H), 7.20-7.26 (m, 4H), 7.31-7.42 (m, 6H), 7.64 (s, 1H), 10.86 (s, 1H).	347	345	Racemate
10		¹ H-NMR (DMSO-D ₆) δ : 1.83-2.03 (m, 2H), 2.12-2.29 (m, 2H), 2.79-2.90 (m, 1H), 3.25-3.37 (m, 2H), 6.89 (s, 1H), 7.20-7.26 (m, 4H), 7.31-7.42 (m, 6H), 7.50 (s, 1H), 10.84 (s, 1H).	361	359	Racemate




11		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.3, 7.5 Hz), 2.37-2.42 (m, 2H), 2.46 (t, 2H, J = 7.5 Hz), 3.30-3.37 (m, 1H), 3.38-3.47 (m, 1H), 3.47-3.54 (m, 1H), 6.90 (s, 1H), 7.00 (s, 1H), 7.07 (d, 1H, J = 7.7 Hz), 7.17 (d, 1H, J = 7.7 Hz), 7.20-7.24 (m, 2H), 7.27 (dd, 1H, J = 7.7, 7.7 Hz), 7.31-7.42 (m, 3H), 7.63 (s, 1H), 10.85 (s, 1H).</p>	389	387	Racemate
12		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 0.73 (t, 3H, J = 7.5 Hz), 1.70 (tq, 2H, J = 7.1, 7.5 Hz), 2.33-2.38 (m, 2H), 3.27-3.33 (m, 1H), 3.34-3.43 (m, 1H), 3.44-3.50 (m, 1H), 3.95 (t, 2H, J = 7.1 Hz), 6.59 (s, 1H), 7.43-7.54 (m, 5H), 7.60 (s, 1H), 10.65 (s, 1H).</p>	313	311	Racemate
13		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 0.78 (t, 3H, J = 7.1 Hz), 1.06-1.20 (m, 6H), 1.62-1.71 (m, 2H), 2.33-2.38 (m, 2H), 3.27-3.33 (m, 1H), 3.34-3.43 (m, 1H), 3.44-3.50 (m, 1H), 3.99 (t, 2H, J = 7.1 Hz), 6.58 (s, 1H), 7.43-7.53 (m, 5H), 7.60 (s, 1H), 10.65 (s, 1H).</p>	355	353	Racemate





14		<p>¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.1 Hz), 0.90 (t, 3H, J = 7.5 Hz), 1.07-1.20 (m, 6H), 1.62 (tq, 2H, J = 7.1, 7.7 Hz), 1.64-1.70 (m, 2H), 2.32-2.38 (m, 2H), 2.62 (t, 2H, J = 7.7 Hz), 3.27-3.33 (m, 1H), 3.33-3.43 (m, 1H), 3.44-3.50 (m, 1H), 3.98 (t, 2H, J = 7.1 Hz), 6.57 (s, 1H), 7.23-7.29 (m, 3H), 7.40 (dd, 1H, J = 7.7, 7.7 Hz), 7.60 (s, 1H), 10.64 (s, 1H).</p>	397	395	Racemate
15		<p>¹H-NMR (DMSO-D₆) δ: 0.73 (t, 3H, J = 7.5 Hz), 0.90 (t, 3H, J = 7.3 Hz), 1.63 (tq, 2H, J = 7.5, 7.5 Hz), 1.70 (tq, 2H, J = 7.3, 7.5 Hz), 2.34-2.38 (m, 2H), 2.62 (t, 2H, J = 7.5 Hz), 3.27-3.33 (m, 1H), 3.34-3.43 (m, 1H), 3.44-3.50 (m, 1H), 3.95 (t, 2H, J = 7.1 Hz), 6.57 (s, 1H), 7.23-7.29 (m, 3H), 7.41 (dd, 1H, J = 7.7, 7.7 Hz), 7.60 (s, 1H), 10.64 (s, 1H).</p>	355	353	Racemate




16		¹ H-NMR (DMSO-D ₆) δ : 1.35 (d, 6H, J = 6.4 Hz), 2.29-2.42 (m, 2H), 3.27-3.33 (m, 1H), 3.34-3.43 (m, 1H), 3.44-3.50 (m, 1H), 4.46 (sep, 1H, J = 6.4Hz), 6.55 (s, 1H), 7.40-7.54 (m, 5H), 7.60 (s, 1H), 10.71 (s, 1H).	313	311	Racemate
17		¹ H-NMR (DMSO-D ₆) δ : 0.91 (t, 3H, J = 7.5 Hz), 1.34 (d, 6H, J = 6.4 Hz), 1.63 (tq, 2H, J = 7.5, 7.7 Hz), 2.30-2.41 (m, 2H), 2.62 (t, 2H, J = 7.7 Hz), 3.27-3.33 (m, 1H), 3.34-3.43 (m, 1H), 3.44-3.50 (m, 1H), 4.46 (sep, 1H, J = 6.4 Hz), 6.53 (s, 1H), 7.20-7.25 (m, 2H), 7.29 (d, 1H, J = 7.7 Hz), 7.41 (dd, 1H, J = 7.7, 7.7 Hz), 7.60 (s, 1H), 10.70 (s, 1H).	355	353	Racemate
18		¹ H-NMR (DMSO-D ₆) δ : 0.71 (d, 6H, J = 6.6 Hz), 0.89 (t, 3H, J = 7.4 Hz), 1.62 (tq, 2H, J = 7.4, 7.4 Hz), 1.94-2.08 (m, 1H), 2.28-2.42 (m, 2H), 2.62 (t, 2H, J = 7.5 Hz), 3.23-3.51 (m, 3H), 3.82 (d, 2H, J = 7.3 Hz), 6.58 (s, 1H), 7.22-7.30 (m, 3H), 7.37-7.43 (m, 1H), 7.61 (s, 1H), 10.64 (s, 1H)	369	367	Racemate



19		¹ H-NMR (CDCl ₃) δ: 0.83 (t, 3H, J = 7.3 Hz), 1.11 (d, 3H, J = 6.0 Hz), 1.50 (tq, 2H, J = 7.5, 7.3 Hz), 2.45-2.63 (m, 3H), 2.68-2.81 (m, 2H), 3.96-4.06 (m, 1H), 6.66 (s, 1H), 7.02-7.08 (m, 3H), 7.12 (d, 1H, J = 7.4 Hz), 7.19 (d, 1H, J = 7.9 Hz), 7.22-7.26 (m, 3H), 7.28-7.38 (m, 3H), 9.55 (s, 1H).	403	401	Single diastereomer (Racemate) cis-trans isomer of Example 20
20		¹ H-NMR (CDCl ₃) δ: 0.83 (t, 3H, J = 7.3 Hz), 1.09 (d, 3H, J = 6.5 Hz), 1.44-1.56 (tq, 2H, J = 7.5, 7.3 Hz), 2.32 (dd, 1H, J = 17.0, 8.8 Hz), 2.49 (t, 2H, J = 7.5 Hz), 2.95 (dd, 1H, J = 17.0, 8.7 Hz), 3.21 (q, 1H, J = 8.6 Hz), 3.83-3.92 (m, 1H), 6.30 (s, 1H), 7.02-7.07 (m, 3H), 7.12 (d, 1H, J = 7.7 Hz), 7.17-7.26 (m, 3H), 7.29-7.37 (m, 3H), 9.32 (s, 1H).	403	401	Single diastereomer (Racemate) cis-trans isomer of Example 19


21		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 0.77 (t, 3H, J = 7.3 Hz), 1.38-1.49 (m, 5H), 2.15 (d, 1H, J = 16.5 Hz), 2.46 (t, 2H, J = 7.5 Hz), 2.74 (d, 1H, J = 16.5 Hz), 3.14 (d, 1H, J = 10.0 Hz), 3.65 (d, 1H, J = 10.0 Hz), 6.90 (s, 1H), 7.00 (s, 1H), 7.07 (dt, 1H, J = 7.7, 1.4 Hz), 7.17 (d, 1H, J = 7.7 Hz), 7.21-7.30 (m, 3H), 7.31-7.43 (m, 3H), 7.62 (s, 1H), 10.58 (s, 1H).</p>	403	401	Racemate
22		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 0.77 (t, 3H, J = 7.3 Hz), 1.38-1.50 (tq, 2H, J = 7.5, 7.3 Hz), 2.36-2.49 (m, 4H), 3.33-3.54 (m, 3H), 6.90 (s, 1H), 7.00 (s, 1H), 7.07 (d, 1H, J = 7.9 Hz), 7.17 (d, 1H, J = 7.9 Hz), 7.19-7.43 (m, 6H), 7.64 (s, 1H), 10.85 (s, 1H).</p>	389	387	R-form



23		¹ H-NMR (DMSO-D ₆) δ : 0.77 (t, 3H, J = 7.3 Hz), 1.38-1.50 (tq, 2H, J = 7.5, 7.3 Hz), 2.36-2.49 (m, 4H), 3.33-3.54 (m, 3H), 6.90 (s, 1H), 7.00 (s, 1H), 7.07 (d, 1H, J = 7.9 Hz), 7.17 (d, 1H, J = 7.9 Hz), 7.19-7.43 (m, 6H), 7.64 (s, 1H), 10.85 (s, 1H).	389	387	S-form
24		¹ H-NMR (DMSO-D ₆) δ : 0.90 (t, 3H, J = 7.3 Hz), 1.29 (t, 3H, J = 7.2 Hz), 1.62 (tq, 2H, J = 7.3, 7.3 Hz), 2.31-2.39 (m, 2H), 2.62 (t, 2H, J = 7.5 Hz), 3.24-3.52 (m, 3H), 4.01 (q, 2H, J = 7.1 Hz), 6.57 (s, 1H), 7.23-7.30 (m, 3H), 7.37-7.43 (m, 1H), 7.60 (s, 1H), 10.64 (s, 1H)	341	339	Racemate
25		¹ H-NMR (DMSO-D ₆) δ : 1.09 (s, 9H), 2.35-2.42 (m, 2H), 3.32-3.55 (m, 3H), 6.92 (s, 1H), 7.06 (s, 1H), 7.18 (d, 1H, J = 7.3 Hz), 7.21-7.26 (m, 2H), 7.28-7.44 (m, 5H), 7.63 (s, 1H), 10.84 (s, 1H).	403	401	R-form





26		¹ H-NMR (DMSO-D ₆) δ: 1.09 (s, 9H), 2.35-2.42 (m, 2H), 3.32-3.55 (m, 3H), 6.92 (s, 1H), 7.06 (s, 1H), 7.18 (d, 1H, J = 7.3 Hz), 7.21-7.26 (m, 2H), 7.28-7.44 (m, 5H), 7.63 (s, 1H), 10.84 (s, 1H).	403	401	S-form
27		¹ H-NMR (DMSO-D ₆) δ: 2.35-2.43 (m, 2H), 3.32-3.54 (m, 3H), 7.05 (s, 1H), 7.23-7.29 (m, 2H), 7.34-7.75 (m, 8H), 10.91 (s, 1H).	415	413	R-form
28		¹ H-NMR (DMSO-D ₆) δ: 2.35-2.43 (m, 2H), 3.32-3.54 (m, 3H), 7.05 (s, 1H), 7.23-7.29 (m, 2H), 7.34-7.75 (m, 8H), 10.91 (s, 1H).	415	413	S-form
29		¹ H-NMR (DMSO-D ₆) δ: 2.35-2.42 (m, 2H), 3.31-3.54 (m, 3H), 7.01 (s, 1H), 7.08 (s, 1H), 7.22-7.28 (m, 2H), 7.32-7.46 (m, 5H), 7.52 (t, 1H, J = 8.0 Hz), 7.64 (s, 1H), 10.90 (s, 1H).	431	429	S-form
30		¹ H-NMR (DMSO-D ₆) δ: 1.24 (t, 3H, J = 7.0 Hz), 2.35-2.42 (m, 2H), 3.34-3.54 (m, 3H), 3.91 (q, 2H, J = 7.0 Hz), 6.72-6.80 (m, 2H), 6.88-6.93 (m, 2H), 7.22-7.28 (m, 3H), 7.31-7.44 (m, 3H), 7.64 (s, 1H), 10.86 (s, 1H).	391	389	R-form




31		¹ H-NMR (DMSO-D ₆) δ : 1.24 (t, 3H, J = 7.0 Hz), 2.35-2.42 (m, 2H), 3.34-3.54 (m, 3H), 3.91 (q, 2H, J = 7.0 Hz), 6.72-6.80 (m, 2H), 6.88-6.93 (m, 2H), 7.22-7.28 (m, 3H), 7.31-7.44 (m, 3H), 7.64 (s, 1H), 10.86 (s, 1H).	391	389	S-form
32		¹ H-NMR (DMSO-D ₆) δ : 0.90 (t, 3H, J = 7.4 Hz), 1.03-1.27 (m, 3H), 1.54-1.69 (m, 3H), 1.71-1.89 (m, 6H), 2.28-2.37 (m, 2H), 2.62 (t, 2H, J = 7.42 Hz), 3.22-3.53 (m, 3H), 3.94-4.08 (m, 1H), 6.52 (s, 1H), 7.18-7.24 (m, 2H), 7.25-7.31 (m, 1H), 7.37-7.44 (m, 1H), 7.60 (s, 1H), 10.68 (s, 1H)	395	393	Racemate
33		¹ H-NMR (DMSO-D ₆) δ : 0.77 (t, 3H, J = 7.3 Hz), 1.08 (d, 3H, J = 7.2 Hz), 1.38-1.50 (tq, 2H, J = 7.5, 7.3 Hz), 2.46 (t, 2H, J = 7.5 Hz), 2.52-2.58 (m, 1H), 3.03 (ddd, 1H, J = 9.2, 8.8, 8.7 Hz), 3.23 (dd, 1H, J = 9.2, 8.8 Hz), 3.44 (dd, 1H, J = 9.2, 8.7 Hz), 6.92 (s, 1H), 7.01 (s, 1H), 7.07 (d, 1H, J = 7.9 Hz), 7.17 (d, 1H, J = 7.9 Hz), 7.20-7.43 (m, 6H), 7.67 (s, 1H), 10.94 (s, 1H).	403	401	trans-form(Racemate)




34		¹ H-NMR (DMSO-D ₆) δ : 0.78 (t, 3H, J = 7.5 Hz), 1.46 (tq, 2H, J = 7.3, 7.5 Hz), 2.37-2.41 (m, 2H), 2.47 (t, 2H, J = 7.3 Hz), 3.30-3.35 (m, 1H), 3.37-3.46 (m, 1H), 3.47-3.53 (m, 1H), 6.89 (s, 1H), 7.00 (s, 1H), 7.07 (d, 1H, J = 7.7 Hz), 7.18 (d, 1H, J = 7.7 Hz), 7.22-7.30 (m, 5H), 7.64 (s, 1H), 10.84 (s, 1H).	407	405	R-form
35		¹ H-NMR (DMSO-D ₆) δ : 0.78 (t, 3H, J = 7.3 Hz), 1.46 (tq, 2H, J = 7.3, 7.3 Hz), 2.37-2.41 (m, 2H), 2.47 (t, 2H, J = 7.3 Hz), 3.30-3.36 (m, 1H), 3.37-3.46 (m, 1H), 3.47-3.53 (m, 1H), 6.89 (s, 1H), 7.00 (s, 1H), 7.07 (d, 1H, J = 7.7 Hz), 7.18 (d, 1H, J = 7.7 Hz), 7.21-7.30 (m, 5H), 7.64 (s, 1H), 10.84 (s, 1H).	407	405	S-form




36		¹ H-NMR (DMSO-D ₆) δ : 0.78 (t, 3H, J = 7.3 Hz), 1.47 (tq, 2H, J = 7.3, 7.3 Hz), 2.36-2.41 (m, 2H), 2.48 (t, 2H, J = 7.3 Hz), 3.28-3.37 (m, 1H), 3.37-3.47 (m, 1H), 3.47-3.55 (m, 1H), 6.91 (s, 1H), 7.02 (s, 1H), 7.08 (d, 1H, J = 8.1 Hz), 7.19	423	421	R-form
----	---	--	-----	-----	--------





		(d, 1H, J = 7.7 Hz), 7.24 (d, 2H, J = 8.9 Hz), 7.30 (dd, 1H, J = 7.7, 8.1 Hz), 7.46 (d, 2H, J = 8.9 Hz), 7.64 (s, 1H), 10.88 (s, 1H).			
37		¹ H-NMR (DMSO-D ₆) δ : 0.78 (t, 3H, J = 7.3 Hz), 1.47 (tq, 2H, J = 7.3, 7.3 Hz), 2.37-2.41 (m, 2H), 2.48 (t, 2H, J = 7.3 Hz), 3.31-3.36 (m, 1H), 3.38-3.46 (m, 1H), 3.47-3.53 (m, 1H), 6.91 (s, 1H), 7.02 (s, 1H), 7.08 (d, 1H, J = 8.1 Hz), 7.19 (d, 1H, J = 7.7 Hz), 7.24 (d, 2H, J = 8.9 Hz), 7.30 (dd, 1H, J = 7.7, 8.1 Hz), 7.46 (d, 2H, J = 8.9 Hz), 7.64 (s, 1H), 10.88 (s, 1H).	423	421	S-form
38		¹ H-NMR (DMSO-D ₆) δ : 2.35-2.42 (m, 2H), 3.32-3.54 (m, 3H), 6.97 (s, 1H), 7.17 (d, 1H, J = 7.5 Hz), 7.23-7.31 (m, 3H), 7.34-7.46 (m, 5H), 7.64 (s, 1H), 10.89 (s, 1H).	381	379	R-form




39		¹ H-NMR (DMSO-D ₆) δ: 2.35-2.42 (m, 2H), 3.32-3.54 (m, 3H), 6.97 (s, 1H), 7.17 (d, 1H, J = 7.5 Hz), 7.23-7.31 (m, 3H), 7.34-7.46 (m, 5H), 7.64 (s, 1H), 10.89 (s, 1H).	381	379	S-form
40		¹ H-NMR (DMSO-D ₆) δ: 1.09-1.37 (m, 5H), 1.55-1.83 (m, 5H), 2.31- 2.40 (m, 2H), 2.60-2.70 (m, 1H), 3.27-3.50 (m, 3H), 6.59 (s, 1H), 7.40- 7.56 (m, 5H), 7.62 (s, 1H), 10.67 (s, 1H).	353	351	R-form
41		¹ H-NMR (DMSO-D ₆) δ: 1.09-1.37 (m, 5H), 1.55-1.83 (m, 5H), 2.31- 2.40 (m, 2H), 2.60-2.70 (m, 1H), 3.27-3.50 (m, 3H), 6.59 (s, 1H), 7.40- 7.56 (m, 5H), 7.62 (s, 1H), 10.67 (s, 1H).	353	351	S-form
42		¹ H-NMR (DMSO-D ₆) δ: 0.88 (t, 3H, J = 7.3 Hz), 1.55-1.67 (m, 2H), 2.29- 2.37 (m, 2H), 2.60 (t, 2H, J = 7.5 Hz), 3.24-3.51 (m, 3H), 6.38 (s, 1H), 7.14-7.22 (m, 2H), 7.23- 7.29 (m, 1H), 7.30-7.38 (m, 1H), 7.59 (s, 1H), 10.57 (s, 1H)	369	367	Racemate



43		¹ H-NMR (DMSO-D ₆) δ: 1.05 (t, 3H, J = 7.5 Hz), 2.36-2.41 (m, 2H), 2.52 (q, 2H, J = 7.5 Hz), 3.32-3.54 (m, 3H), 6.90 (s, 1H), 7.01-7.07 (m, 2H), 7.16-7.29 (m, 4H), 7.31-7.43 (m, 3H), 7.63 (s, 1H), 10.85 (s, 1H).	375	373	R-form
44		¹ H-NMR (DMSO-D ₆) δ: 1.05 (t, 3H, J = 7.5 Hz), 2.36-2.41 (m, 2H), 2.52 (q, 2H, J = 7.5 Hz), 3.32-3.54 (m, 3H), 6.90 (s, 1H), 7.01-7.07 (m, 2H), 7.16-7.29 (m, 4H), 7.31-7.43 (m, 3H), 7.63 (s, 1H), 10.85 (s, 1H).	375	373	S-form
45		¹ H-NMR (DMSO-D ₆) δ: 0.82 (t, 3H, J = 7.3 Hz), 1.15 (tq, 2H, J = 7.5, 7.3 Hz), 1.39 (tt, 2H, J = 7.6, 7.5 Hz), 2.36-2.49 (m, 4H), 3.31-3.54 (m, 3H), 6.89 (s, 1H), 6.98 (s, 1H), 7.08 (d, 1H, J = 7.7 Hz), 7.17 (d, 1H, J = 7.7 Hz), 7.20-7.42 (m, 6H), 7.63 (s, 1H), 10.85 (s, 1H).	403	401	R-form




46		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 0.82 (t, 3H, J = 7.3 Hz), 1.15 (tq, 2H, J = 7.5, 7.3 Hz), 1.39 (tt, 2H, J = 7.6, 7.5 Hz), 2.36-2.49 (m, 4H), 3.31-3.54 (m, 3H), 6.89 (s, 1H), 6.98 (s, 1H), 7.08 (d, 1H, J = 7.7 Hz), 7.17 (d, 1H, J = 7.7 Hz), 7.20-7.42 (m, 6H), 7.63 (s, 1H), 10.85 (s, 1H).</p>	403	401	S-form
47		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 1.49-1.61 (m, 4H), 1.92-2.10 (m, 4H), 2.33-2.38 (m, 2H), 3.25-3.51 (m, 3H), 5.75-5.81 (m, 1H), 6.63 (s, 1H), 7.33-7.51 (m, 5H), 7.62 (s, 1H), 10.74 (s, 1H).</p>	351	349	R-form
48		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 0.77 (t, 3H, J = 7.3 Hz), 1.45 (tq, 2H, J = 7.3, 7.3 Hz), 2.31 (s, 3H), 2.36-2.41 (m, 2H), 2.47 (t, 2H, J = 7.3 Hz), 3.30-3.35 (m, 1H), 3.37-3.46 (m, 1H), 3.47-3.53 (m, 1H), 6.87 (s, 1H), 7.01 (s, 1H), 7.04 (d, 1H, J = 7.7 Hz), 7.10 (d, 2H, J = 8.5 Hz), 7.16 (d, 1H, J = 8.1 Hz), 7.19 (d, 2H, J = 8.5 Hz), 7.26 (dd, 1H, J = 7.7, 8.1 Hz), 7.63 (s, 1H), 10.83 (s, 1H).</p>	403	401	R-form




49		¹ H-NMR (DMSO-D ₆) δ : 0.77 (t, 3H, J = 7.3 Hz), 1.45 (tq, 2H, J = 7.3, 7.3 Hz), 2.31 (s, 3H), 2.36-2.41 (m, 2H), 2.47 (t, 2H, J = 7.3 Hz), 3.30-3.35 (m, 1H), 3.37-3.46 (m, 1H), 3.47-3.53 (m, 1H), 6.87 (s, 1H), 7.01 (s, 1H), 7.04 (d, 1H, J = 7.3 Hz), 7.10 (d, 2H, J = 8.5 Hz), 7.16 (d, 1H, J = 7.7 Hz), 7.19 (d, 2H, J = 8.5 Hz), 7.26 (dd, 1H, J = 7.7, 7.3 Hz), 7.63 (s, 1H), 10.83 (s, 1H).	403	401	S-form
50		¹ H-NMR (DMSO-D ₆) δ : 0.77 (t, 3H, J = 7.3 Hz), 1.43 (tq, 2H, J = 7.5, 7.3 Hz), 2.35-2.41 (m, 2H), 2.50 (t, 2H, J = 7.5 Hz), 3.33-3.53 (m, 3H), 6.90 (s, 1H), 7.08-7.44 (m, 8H), 7.63 (s, 1H), 10.85 (s, 1H).	407	405	R-form
51		¹ H-NMR (DMSO-D ₆) δ : 0.77 (t, 3H, J = 7.3 Hz), 1.43 (tq, 2H, J = 7.5, 7.3 Hz), 2.35-2.41 (m, 2H), 2.50 (t, 2H, J = 7.5 Hz), 3.33-3.53 (m, 3H), 6.90 (s, 1H), 7.08-7.44 (m, 8H), 7.63 (s, 1H), 10.85 (s, 1H).	407	405	S-form




52		1H-NMR (DMSO-D6) δ : 0.46 (t, 3H, J = 7.4 Hz), 1.05 (s, 6H), 1.41 (q, 2H, J = 7.4 Hz), 2.34-2.43 (m, 2H), 3.32-3.55 (m, 3H), 6.91 (s, 1H), 6.96 (s, 1H), 7.19-7.44 (m, 8H), 7.63 (s, 1H), 10.84 (s, 1H).	417	415	R-form
53		1H-NMR (DMSO-D6) δ : 0.46 (t, 3H, J = 7.4 Hz), 1.05 (s, 6H), 1.41 (q, 2H, J = 7.4 Hz), 2.34-2.43 (m, 2H), 3.32-3.55 (m, 3H), 6.91 (s, 1H), 6.96 (s, 1H), 7.19-7.44 (m, 8H), 7.63 (s, 1H), 10.84 (s, 1H).	417	415	S-form
54		1H-NMR (DMSO-D6) δ : 1.04 (d, 6H, J = 7.1 Hz), 2.36-2.42 (m, 2H), 2.77-2.79 (m, 1H), 3.32-3.54 (m, 3H), 6.91 (s, 1H), 6.99 (s, 1H), 7.11 (d, 1H, J = 7.7 Hz), 7.18-7.44 (m, 7H), 7.63 (s, 1H), 10.84 (s, 1H).	389	387	R-form
55		1H-NMR (DMSO-D6) δ : 1.04 (d, 6H, J = 7.1 Hz), 2.36-2.42 (m, 2H), 2.77-2.79 (m, 1H), 3.32-3.54 (m, 3H), 6.91 (s, 1H), 6.99 (s, 1H), 7.11 (d, 1H, J = 7.7 Hz), 7.18-7.44 (m, 7H), 7.63 (s, 1H), 10.84 (s, 1H).	389	387	S-form




56		¹ H-NMR (DMSO-D ₆) δ: 1.95 (s, 3H), 2.38-2.40 (m, 2H), 3.33-3.36 (m, 1H), 3.40-3.45 (m, 1H), 3.47-3.52 (m, 1H), 6.77 (s, 1H), 7.13-7.15 (m, 2H), 7.22-7.35 (m, 7H), 7.63 (s, 1H), 10.90 (s, 1H).	361	359	R-form
57		¹ H-NMR (DMSO-D ₆) δ: 2.25 (s, 3H), 2.38-2.39 (m, 2H), 3.32-3.35 (m, 1H), 3.39-3.46 (m, 1H), 3.48-3.52 (m, 1H), 6.89 (s, 1H), 6.94 (d, 1H, J = 7.7 Hz), 7.13 (s, 1H), 7.16-7.24 (m, 4H), 7.32-7.41 (m, 3H), 7.63 (s, 1H), 10.85 (s, 1H).	361	359	R-form
58		¹ H-NMR (DMSO-D ₆) δ: 2.30 (s, 3H), 2.38-2.39 (m, 2H), 3.32-3.35 (m, 1H), 3.39-3.46 (m, 1H), 3.48-3.52 (m, 1H), 6.86 (s, 1H), 7.11 (d, 2H, J = 8.2 Hz), 7.17 (d, 2H, J = 8.2 Hz), 7.22-7.23 (m, 2H), 7.31-7.41 (m, 3H), 7.63 (s, 1H), 10.84 (s, 1H).	361	359	R-form




59		<p>¹H-NMR (DMSO-D₆) δ: 0.78 (t, 3H, J = 7.3 Hz), 0.95 (s, 3H), 1.17 (s, 3H), 1.39-1.50 (tq, 2H, J = 7.5, 7.3 Hz), 2.46 (t, 2H, J = 7.5 Hz), 3.13 (dd, 1H, J = 7.3, 6.8 Hz), 3.29-3.35 (m, 1H), 3.43 (dd, 1H, J = 9.6, 6.8 Hz), 6.91 (s, 1H), 7.02 (s, 1H), 7.07 (d, 1H, J = 7.9 Hz), 7.17 (d, 1H, J = 7.9 Hz), 7.20-7.29 (m, 3H), 7.31-7.43 (m, 3H), 7.64 (s, 1H), 10.79 (s, 1H).</p>	417	415	Racemate
60		<p>¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 0.90 (t, 3H, J = 7.4 Hz), 1.42-1.46 (m, 3H), 1.63-1.76 (m, 1H), 2.46 (t, 2H, J = 7.5 Hz), 2.52-2.57 (m, 1H), 3.11-3.22 (m, 2H), 3.39-3.48 (m, 1H), 6.92 (s, 1H), 7.01 (s, 1H), 7.07 (d, 1H, J = 7.7 Hz), 7.17 (d, 1H, J = 7.7 Hz), 7.20-7.30 (m, 3H), 7.31-7.43 (m, 3H), 7.68 (s, 1H), 10.98 (s, 1H).</p>	417	415	trans-form(Racemate)




61		¹ H-NMR (DMSO-D ₆) δ : 0.77 (t, 3H, $J = 7.3$ Hz), 1.44 (tq, 2H, $J = 7.5, 7.3$ Hz), 1.83-2.03 (m, 2H), 2.12-2.30 (m, 2H), 2.46 (t, 2H, $J = 7.5$ Hz), 2.84 (tt, 1H, $J = 10.7, 3.6$ Hz), 3.24-3.31 (m, 2H), 6.89 (s, 1H), 7.00 (s, 1H), 7.06 (d, 1H, $J = 7.7$ Hz), 7.17 (d, 1H, $J = 7.7$ Hz), 7.19-7.43 (m, 6H), 7.51 (s, 1H), 10.83 (s, 1H).	403	401	Racemate
62		¹ H-NMR (DMSO-D ₆) δ : 1.25-1.87 (m, 12H), 2.31-2.38 (m, 2H), 2.81-2.90 (m, 1H), 3.26-3.50 (m, 3H), 6.60 (s, 1H), 7.38-7.56 (m, 5H), 7.62 (s, 1H), 10.67 (s, 1H).	367	365	R-form
63		¹ H-NMR (DMSO-D ₆) δ : 1.44-1.59 (m, 4H), 1.63-1.76 (m, 2H), 1.81-1.96 (m, 2H), 2.31-2.38 (m, 2H), 2.99-3.11 (m, 1H), 3.26-3.51 (m, 3H), 6.60 (s, 1H), 7.40-7.55 (m, 5H), 7.62 (s, 1H), 10.68 (s, 1H).	339	-	R-form





64		¹ H-NMR (DMSO-D ₆) δ: 0.90 (t, 3H, J = 7.3 Hz), 1.48-1.68 (m, 4H), 1.74-2.03 (m, 6H), 2.27- 2.43 (2H, m), 2.62 (2H, t, J = 7.5 Hz), 3.25-3.50 (3H, m), 4.61 (1H, quin, J = 7.4 Hz), 6.53 (1H, s), 7.20-7.30 (3H, m), 7.38- 7.44 (1H, m), 7.61 (1H, s), 10.66 (1H, s)	381	379	R-form
65		¹ H-NMR (DMSO-D ₆) δ: 0.90 (t, 3H, J = 7.3 Hz), 1.48-1.68 (m, 4H), 1.74-2.03 (m, 6H), 2.27- 2.41 (2H, m), 2.62 (2H, t, J = 7.5 Hz), 3.25-3.50 (3H, m), 4.61 (1H, quin, J = 7.5 Hz), 6.53 (1H, s), 7.20- 7.30 (3H, m), 7.37- 7.44 (1H, m), 7.60 (1H, s), 10.66 (1H, s)	381	379	S-form
66		¹ H-NMR (DMSO-D ₆) δ: 0.89 (3H, t, J = 7.3 Hz), 1.62 (2H, tq, J = 7.5, 7.6 Hz), 2.37 (2H, d, J = 8.4 Hz), 2.63 (2H, t, J = 7.5 Hz), 3.27-3.53 (3H, m), 4.88 (2H, q, J = 8.8 Hz), 6.73 (1H, s), 7.25-7.34 (3H, m), 7.40-7.46 (1H, m), 7.62 (1H, s), 10.83 (1H, s)	395	393	R-form




67		¹ H-NMR (DMSO-D ₆) δ: 0.89 (3H, t, J = 7.4 Hz), 1.62 (2H, tq, J = 7.5, 7.6 Hz), 2.37 (2H, d, J = 8.2 Hz), 2.63 (2H, t, J = 7.5 Hz), 3.28- .52 (3H, m), 4.88 (2H, q, J = 8.8 Hz), 6.73 (1H, s), 7.25-7.34 (3H, m), 7.40- 7.46 (1H, m), 7.62 (1H, s), 10.83 (1H, s)	395	393	S-form
68		¹ H-NMR (DMSO-D ₆) δ: 0.91 (3H, t, J = 7.4 Hz), 1.28-1.42 (2H, m), 1.46- 1.80 (8H, m), 1.82-2.06 (4H, m), 2.28-2.42 (2H, m), 2.63 (2H, t, J = 7.5 Hz), 3.26-3.52 (3H, m), 4.19-4.29 (1H, m), 6.53 (1H, s), 7.18- 7.24 (2H, m), 7.26-7.32 (1H, m), 7.39-7.45 (1H, m), 7.60 (1H, s), 10.68 (1H, s)	409	407	R-form
69		¹ H-NMR (DMSO-D ₆) δ: 0.91 (3H, t, J = 7.3 Hz), 1.28-1.42 (2H, m), 1.46- 1.80 (8H, m), 1.82-2.06 (4H, m), 2.28-2.42 (2H, m), 2.63 (2H, t, J = 7.5 Hz), 3.26-3.52 (3H, m), 4.19-4.29 (1H, m), 6.53 (1H, s), 7.18- 7.24 (2H, m), 7.26-7.32 (1H, m), 7.39-7.45 (1H, m), 7.60 (1H, s), 10.68 (1H, s)	409	407	S-form



70		¹ H-NMR (DMSO-D ₆) δ : 0.74 (d, 6H, J = 6.6 Hz), 1.64 (tsep, 1H, J = 6.6, 7.1 Hz), 2.34 (d, 2H, J = 7.1 Hz), 2.39-2.41 (m, 2H), 3.34-3.36 (m, 2H), 3.39-3.44 (m, 1H), 3.47-3.51 (m, 1H), 6.89 (s, 1H), 6.93 (s, 1H), 7.10-7.40 (m, 8H), 7.63 (s, 1H), 10.84 (s, 1H).	403	401	R-form
71		¹ H-NMR (DMSO-D ₆) δ : 0.74 (d, 6H, J = 6.6 Hz), 1.64 (tsep, 1H, J = 6.6, 7.1 Hz), 2.34 (d, 2H, J = 7.1 Hz), 2.39-2.41 (m, 2H), 3.34-3.36 (m, 2H), 3.39-3.44 (m, 1H), 3.47-3.51 (m, 1H), 6.89 (s, 1H), 6.93 (s, 1H), 7.10-7.40 (m, 8H), 7.63 (s, 1H), 10.84 (s, 1H).	403	401	S-form
72		¹ H-NMR (DMSO-D ₆) δ : 0.76 (t, 3H, J = 7.6 Hz), 2.06 (tq, 2H, J = 16.5, 7.6 Hz), 2.38-2.40 (m, 2H), 3.33-3.37 (m, 1H), 3.39-3.48 (m, 1H), 3.49-3.53 (m, 1H), 6.99 (s, 1H), 7.22-7.25 (m, 3H), 7.34-7.52 (m, 6H), 7.64 (s, 1H), 10.89 (s, 1H).	425	423	R-form



73		¹ H-NMR (DMSO-D ₆) δ: 0.76 (t, 3H, J = 7.6 Hz), 2.06 (tq, 2H, J = 16.5, 7.6 Hz), 2.38-2.40 (m, 2H), 3.33-3.37 (m, 1H), 3.39-3.48 (m, 1H), 3.49-3.53 (m, 1H), 6.99 (s, 1H), 7.22-7.25 (m, 3H), 7.34-7.52 (m, 6H), 7.64 (s, 1H), 10.89 (s, 1H).	425	423	S-form
74		¹ H-NMR (DMSO-D ₆) δ: 0.76 (t, 3H, J = 7.4 Hz), 1.44 (tq, 2H, J = 7.7, 7.4 Hz), 2.38-2.40 (m, 2H), 2.46 (d, 2H, J = 7.7 Hz), 3.34-3.36 (m, 1H), 3.39-3.46 (m, 1H), 3.48-3.53 (m, 1H), 6.84 (s, 1H), 6.88 (d, 1H, J = 9.5 Hz), 6.96 (s, 1H), 7.04 (d, 1H, J = 9.9 Hz), 7.24-7.25 (m, 2H), 7.35-7.44 (m, 3H), 7.64 (s, 1H), 10.88 (s, 1H).	407	405	R-form
75		¹ H-NMR (DMSO-D ₆) δ: 0.76 (t, 3H, J = 7.4 Hz), 1.44 (tq, 2H, J = 7.7, 7.4 Hz), 2.38-2.40 (m, 2H), 2.46 (d, 2H, J = 7.7 Hz), 3.34-3.36 (m, 1H), 3.39-3.46 (m, 1H), 3.48-3.53 (m, 1H), 6.84 (s, 1H), 6.88 (d, 1H, J = 9.5 Hz), 6.96 (s, 1H), 7.04 (d, 1H, J = 9.9 Hz), 7.24-7.25 (m, 2H), 7.35-7.44 (m, 3H), 7.64 (s, 1H), 10.88 (s, 1H).	407	405	S-form


76		¹ H-NMR (DMSO-D ₆) δ: 0.76 (3H, t, J = 7.3 Hz), 1.43 (2H, tq, J = 7.4, 7.4 Hz), 2.24 (3H, s), 2.35-2.41 (2H, m), 2.46 (2H, t, J = 7.4 Hz), 3.28-3.53 (3H, m), 6.88 (1H, s), 6.93-7.02 (2H, m), 7.04-7.10 (2H, m), 7.12-7.19 (2H, m), 7.21-7.30 (2H, m), 7.64 (1H, s), 10.84 (1H, s)	403	401	R-form
77		¹ H-NMR (DMSO-D ₆) δ: 0.76 (3H, t, J = 7.3 Hz), 1.43 (2H, tq, J = 7.4, 7.4 Hz), 2.24 (3H, s), 2.35-2.41 (2H, m), 2.46 (2H, t, J = 7.4 Hz), 3.28-3.53 (3H, m), 6.88 (1H, s), 6.93-7.02 (2H, m), 7.04-7.10 (2H, m), 7.12-7.19 (2H, m), 7.21-7.30 (2H, m), 7.64 (1H, s), 10.84 (1H, s)	403	401	S-form
78		¹ H-NMR (DMSO-D ₆) δ: 0.82 (s, 6H), 1.29 (t, 2H, J = 6.4 Hz), 1.75-1.80 (m, 2H), 2.02-2.10 (m, 2H), 2.32-2.39 (m, 2H), 3.27-3.51 (m, 3H), 5.71-5.77 (m, 1H), 6.62 (s, 1H), 7.33-7.50 (m, 5H), 7.63 (s, 1H), 10.75 (s, 1H).	379	377	R-form



79		¹ H-NMR (DMSO-D ₆) δ: 0.86 (s, 6H), 1.34-1.42 (m, 2H), 1.55-1.65 (m, 2H), 1.98-2.06 (m, 2H), 2.31-2.39 (m, 2H), 3.27-3.51 (m, 3H), 5.34 (1H, s), 6.64 (s, 1H), 7.32-7.51 (m, 5H), 7.63 (s, 1H), 10.75 (s, 1H).	379	377	R-form
80		¹ H-NMR (DMSO-D ₆) δ: 1.27 (s, 6H), 2.38-2.40 (m, 2H), 3.33-3.35 (m, 1H), 3.38-3.46 (m, 1H), 3.48-3.53 (m, 1H), 4.99 (s, 1H), 6.92 (s, 1H), 7.09 (d, 1H, J = 7.7 Hz), 7.21-7.46 (m, 8H), 7.64 (s, 1H), 10.86 (s, 1H).	405	403	R-form
81		¹ H-NMR (DMSO-D ₆) δ: 1.27 (s, 6H), 2.38-2.40 (m, 2H), 3.33-3.35 (m, 1H), 3.38-3.46 (m, 1H), 3.48-3.53 (m, 1H), 4.99 (s, 1H), 6.92 (s, 1H), 7.09 (d, 1H, J = 7.7 Hz), 7.21-7.46 (m, 8H), 7.64 (s, 1H), 10.86 (s, 1H).	405	403	S-form
82		¹ H-NMR (DMSO-D ₆) δ: 1.27 (s, 6H), 2.38-2.40 (m, 2H), 2.77 (s, 3H), 3.34-3.35 (m, 1H), 3.41-3.44 (m, 1H), 3.49-3.53 (m, 1H), 6.93 (s, 1H), 7.05 (s, 1H), 7.22-7.28 (m, 3H), 7.33-7.42 (m, 5H), 7.64 (s, 1H), 10.86 (s, 1H).	419	417	R-form



83		¹ H-NMR (DMSO-D ₆) δ: 1.27 (s, 6H), 2.38-2.40 (m, 2H), 2.77 (s, 3H), 3.34-3.35 (m, 1H), 3.41-3.44 (m, 1H), 3.49-3.53 (m, 1H), 6.93 (s, 1H), 7.05 (s, 1H), 7.22-7.28 (m, 3H), 7.33-7.42 (m, 5H), 7.64 (s, 1H), 10.86 (s, 1H).	419	417	S-form
84		¹ H-NMR (DMSO-D ₆) δ: 1.28-1.42 (11H, m), 1.46-1.56 (4H, m), 1.65-1.80 (2H, m), 1.82-2.06 (4H, m), 2.28-2.41 (2H, m), 3.25-3.50 (3H, m), 4.18-4.29 (1H, m), 6.53 (1H, s), 7.21 (1H, d, J = 7.4 Hz), 7.36 (1H, s), 7.43 (1H, t, J = 7.7 Hz), 7.61 (1H, s), 10.70 (1H, s)	423	421	R-form
85		¹ H-NMR (DMSO-D ₆) δ: 1.28-1.42 (2H, m), 1.45-1.60 (4H, m), 1.65-1.79 (2H, m), 1.83-2.04 (4H, m), 2.28-2.41 (2H, m), 3.25-3.50 (3H, m), 4.15-4.25 (1H, m), 6.61 (1H, s), 7.39 (1H, s), 7.43-7.52 (1H, m), 7.68-7.71 (2H, m), 10.74 (1H, m)	451	449	R-form



86		¹ H-NMR (DMSO-D ₆) δ : 0.89 (t, 3H, J = 7.3 Hz), 1.61 (tq, 2H, J = 7.3, 7.5 Hz), 2.37-2.40 (m, 2H), 2.62 (t, 2H, J = 7.5 Hz), 3.30-3.35 (m, 1H), 3.37-3.45 (m, 1H), 3.47-3.53 (m, 1H), 6.93 (s, 1H), 7.30 (d, 1H, J = 7.9 Hz), 7.31 (s, 1H), 7.36 (d, 1H, J = 7.7 Hz), 7.42 (dd, 1H, J = 7.7, 7.9 Hz), 7.65 (s, 1H), 11.17 (s, 1H).	381	379	R-form
87		¹ H-NMR (DMSO-D ₆) δ : 0.89 (t, 3H, J = 7.3 Hz), 1.61 (tq, 2H, J = 7.3, 7.5 Hz), 2.36-2.41 (m, 2H), 2.62 (t, 2H, J = 7.5 Hz), 3.30-3.35 (m, 1H), 3.37-3.45 (m, 1H), 3.47-3.53 (m, 1H), 6.93 (s, 1H), 7.30 (d, 1H, J = 8.1 Hz), 7.31 (s, 1H), 7.36 (d, 1H, J = 8.1 Hz), 7.42 (dd, 1H, J = 8.1, 8.1 Hz), 7.65 (s, 1H), 11.18 (s, 1H).	381	379	S-form



88		¹ H-NMR (DMSO-D ₆) δ : 0.75 (s, 3H), 0.85 (s, 3H), 1.08-1.47 (m, 6H), 1.49-1.60 (m, 1H), 1.74-1.83 (m, 1H), 2.33 (dd, 1H, $J = 16.5, 8.9$ Hz), 2.38 (dd, 1H, $J = 16.5, 7.7$ Hz), 2.87 (tt, 1H, $J = 12.1, 3.2$ Hz), 3.30 (dd, 1H, $J = 8.5, 6.0$ Hz), 3.32-3.42 (m, 1H), 3.47 (dd, 1H, $J = 8.5, 8.5$ Hz), 6.58 (s, 1H), 7.41-7.47 (m, 3H), 7.49-7.56 (m, 2H), 7.62 (s, 1H), 10.68 (s, 1H).	381	379	Optically active substance Diastereomer of Example 89 Retention time under HPLC analysis condition 2 (17.02 minutes)
89		¹ H-NMR (DMSO-D ₆) δ : 0.75 (s, 3H), 0.85 (s, 3H), 1.08-1.48 (m, 6H), 1.50-1.59 (m, 1H), 1.73-1.83 (m, 1H), 2.33 (dd, 1H, $J = 16.5, 8.9$ Hz), 2.38 (dd, 1H, $J = 16.5, 8.1$ Hz), 2.87 (tt, 1H, $J = 12.1, 3.0$ Hz), 3.30 (dd, 1H, $J = 8.7, 6.2$ Hz), 3.32-3.43 (m, 1H), 3.47 (dd, 1H, $J = 8.7, 8.3$ Hz), 6.58 (s, 1H), 7.40-7.47 (m, 3H), 7.50-7.56 (m, 2H), 7.62 (s, 1H), 10.68 (s, 1H).	381	379	Optically active substance Diastereomer of Example 88 Retention time under HPLC analysis condition 2 (20.77 minutes)




90		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 0.77 (t, 3H, J = 7.5 Hz), 1.08 (d, 3H, J = 6.9 Hz), 1.44 (tq, 2H, J = 7.5, 7.5 Hz), 2.46 (t, 2H, J = 7.5 Hz), 2.54 (dq, 1H, J = 8.9, 6.9 Hz), 3.04 (ddd, 1H, J = 8.9, 8.9, 8.9 Hz), 3.23 (dd, 1H, J = 8.9, 8.9 Hz), 3.44 (dd, 1H, J = 8.9, 8.9 Hz), 6.92 (s, 1H), 7.01 (dd, 1H, J = 1.6, 1.6 Hz), 7.07 (ddd, 1H, J = 7.7, 1.6, 1.6 Hz), 7.17 (ddd, 1H, J = 7.7, 1.6, 1.6 Hz), 7.20-7.24 (m, 2H), 7.27 (dd, 1H, J = 7.7, 7.7 Hz), 7.31-7.42 (m, 3H), 7.68 (s, 1H), 10.94 (s, 1H).</p>	403	401	(3S, 4S)-form
----	---	--	-----	-----	---------------




91		<p>¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.5 Hz), 1.08 (d, 3H, J = 6.9 Hz), 1.44 (tq, 2H, J = 7.5, 7.5 Hz), 2.46 (t, 2H, J = 7.5 Hz), 2.54 (dq, 1H, J = 8.9, 6.9 Hz), 3.04 (ddd, 1H, J = 8.9, 8.9, 8.9 Hz), 3.23 (dd, 1H, J = 8.9, 8.9 Hz), 3.44 (dd, 1H, J = 8.9, 8.9 Hz), 6.92 (s, 1H), 7.01 (dd, 1H, J = 1.6, 1.6 Hz), 7.07 (ddd, 1H, J = 7.7, 1.6, 1.6 Hz), 7.17 (ddd, 1H, J = 7.7, 1.6, 1.6 Hz), 7.20-7.24 (m, 2H), 7.27 (dd, 1H, J = 7.7, 7.7 Hz), 7.31-7.42 (m, 3H), 7.68 (s, 1H), 10.94 (s, 1H).</p>	403	401	(3R, 4R)-form
92		<p>¹H-NMR (DMSO-D₆) δ: 0.46-0.48 (m, 2H), 0.85-0.90 (m, 2H), 1.81-1.87 (m, 1H), 2.38-2.39 (m, 2H), 3.35-3.38 (m, 1H), 3.41-3.45 (m, 1H), 3.49-3.51 (m, 1H), 6.82 (s, 1H), 6.90 (s, 1H), 6.99 (d, 1H, J = 7.7 Hz), 7.09 (d, 1H, J = 7.7 Hz), 7.21-7.22 (m, 3H), 7.37-7.41 (m, 3H), 7.64 (s, 1H), 10.84 (s, 1H).</p>	387	385	R-form




93		<p>¹H-NMR (DMSO-D₆) δ: 0.46-0.48 (m, 2H), 0.85-0.90 (m, 2H), 1.81-1.87 (m, 1H), 2.38-2.39 (m, 2H), 3.35-3.38 (m, 1H), 3.41-3.45 (m, 1H), 3.49-3.51 (m, 1H), 6.82 (s, 1H), 6.90 (s, 1H), 6.99 (d, 1H, $J = 7.7$ Hz), 7.09 (d, 1H, $J = 7.7$ Hz), 7.21-7.22 (m, 3H), 7.37-7.41 (m, 3H), 7.64 (s, 1H), 10.84 (s, 1H).</p>	387	385	S-form
94		<p>¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, $J = 7.3$ Hz), 0.98 (d, 3H, $J = 7.4$ Hz), 1.44 (tq, 2H, $J = 7.5$, 7.3 Hz), 2.46 (t, 2H, $J = 7.5$ Hz), 2.59-2.68 (m, 1H), 3.28-3.35 (m, 1H), 3.41-3.50 (m, 1H), 6.89 (s, 1H), 7.02 (s, 1H), 7.07 (d, 1H, $J = 7.7$ Hz), 7.17 (d, 1H, $J = 7.7$ Hz), 7.20-7.43 (6H, m), 7.63 (s, 1H), 10.81 (s, 1H).</p>	403	401	cis-form (Racemate)



95		¹ H-NMR (DMSO-D ₆) δ: 0.77 (3H, t, J = 7.3 Hz), 1.46 (2H, tq, J = 7.4, 7.4 Hz), 2.38 (2H, d, J = 8.1 Hz), 2.45-2.52 (2H, m), 3.29-3.53 (3H, m), 6.91 (1H, s), 7.03 (1H, s), 7.09-7.24 (3H, m), 7.27-7.35 (2H, s), 7.36-7.43 (2H, m), 7.65 (1H, s), 10.90 (1H, s)	423	421	R-form
96		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.5, 7.3 Hz), 1.83-2.03 (m, 2H), 2.12-2.30 (m, 2H), 2.46 (t, 2H, J = 7.5 Hz), 2.84 (tt, 1H, J = 10.7, 3.6 Hz), 3.24-3.31 (m, 2H), 6.89 (s, 1H), 7.00 (s, 1H), 7.06 (d, 1H, J = 7.7 Hz), 7.17 (d, 1H, J = 7.7 Hz), 7.19-7.43 (m, 6H), 7.51 (s, 1H), 10.83 (s, 1H).	403	401	Optically active substance optical isomer of Example 5



97		¹ H-NMR (DMSO-D ₆) δ : 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.5, 7.3 Hz), 2.46 (t, 2H, J = 7.5 Hz), 3.35 (dd, 1H, J = 8.8, 6.2 Hz), 3.59 (dd, 1H, J = 9.4, 8.8 Hz), 4.32 (dd, 1H, J = 9.4, 6.2 Hz), 6.34 (s, 1H), 6.61 (s, 1H), 6.90 (s, 1H), 7.00 (s, 1H), 7.08 (d, 1H, J = 7.7 Hz), 7.15-7.43 (m, 7H), 10.71 (s, 1H).	390	388	S-form
98		¹ H-NMR (DMSO-D ₆) δ : 0.86 (t, 3H, J = 7.3 Hz), 1.56 (tq, 2H, J = 7.3, 7.5 Hz), 2.30-2.41 (m, 2H), 2.56 (t, 2H, J = 7.5 Hz), 3.27-3.34 (m, 1H), 3.34-3.41 (m, 1H), 3.44-3.49 (m, 1H), 5.25 (s, 2H), 6.69 (s, 1H), 6.99-7.04 (m, 2H), 7.21-7.32 (m, 6H), 7.37 (dd, 1H, J = 7.7, 7.7 Hz), 7.62 (s, 1H), 10.71 (s, 1H).	403	401	R-form



99		¹ H-NMR (DMSO-D6) δ : 0.86 (t, 3H, J = 7.3 Hz), 1.56 (tq, 2H, J = 7.3, 7.5 Hz), 2.30-2.41 (m, 2H), 2.56 (t, 2H, J = 7.5 Hz), 3.27-3.34 (m, 1H), 3.34-3.41 (m, 1H), 3.44-3.49 (m, 1H), 5.25 (s, 2H), 6.69 (s, 1H), 6.99-7.04 (m, 2H), 7.21-7.32 (m, 6H), 7.37 (dd, 1H, J = 7.9, 7.7 Hz), 7.62 (s, 1H), 10.71 (s, 1H).	403	401	S-form
100		¹ H-NMR (DMSO-D6) δ : 0.71 (3H, t, J = 7.4 Hz), 1.39 (2H, tq, J = 7.4, 7.5 Hz), 2.36-2.46 (4H, m), 3.31-3.54 (3H, m), 6.93-6.99 (2H, m), 7.04 (2H, d, J = 7.7 Hz), 7.11 (2H, d, J = 7.7 Hz), 7.22 (1H, t, J = 7.6 Hz), 7.43-7.62 (4H, m), 7.65 (1H, s), 10.82 (1H, s)	423	421	R-form
101		¹ H-NMR (DMSO-D6) δ : 0.71 (3H, t, J = 7.4 Hz), 1.39 (2H, tq, J = 7.4, 7.5 Hz), 2.36-2.46 (4H, m), 3.31-3.54 (3H, m), 6.93-6.99 (2H, m), 7.04 (2H, d, J = 7.7 Hz), 7.11 (2H, d, J = 7.7 Hz), 7.22 (1H, t, J = 7.7 Hz), 7.43-7.62 (4H, m), 7.65 (1H, s), 10.82 (1H, s)	423	421	S-form



102		¹ H-NMR (DMSO-D6) δ: 0.71 (3H, t, J = 7.4 Hz), 1.38 (2H, tq, J = 7.2, 7.5 Hz), 1.89 (3H, s), 2.36-2.45 (4H, m), 3.31-3.54 (3H, m), 6.91-6.97 (2H, m), 7.01 (1H, d, J = 7.7 Hz), 7.10 (1H, d, J = 7.5 Hz), 7.20 (1H, t, J = 7.7 Hz), 7.65 (1H, s), 10.77 (1H, s)	403	401	R-form
103		¹ H-NMR (DMSO-D6) δ: 0.78 (3H, t, J = 7.3 Hz), 1.46 (2H, tq, J = 7.1, 7.4 Hz), 2.39 (2H, d, J = 8.4 Hz), 2.45-2.53 (3H, m), 3.30-3.55 (3H, m), 6.91 (1H, s), 7.00-7.15 (4H, m), 7.16-7.24 (2H, m), 7.31 (1H, t, J = 7.6 Hz), 7.38-7.46 (1H, m), 7.65 (1H, s), 10.90 (1H, s)	407	405	R-form
104		¹ H-NMR (DMSO-D6) δ: 0.78 (3H, t, J = 7.4 Hz), 1.46 (2H, tq, J = 7.1, 7.4 Hz), 2.39 (2H, d, J = 8.4 Hz), 2.45-2.53 (3H, m), 3.30-3.55 (3H, m), 6.91 (1H, s), 7.00-7.15 (4H, m), 7.16-7.24 (2H, m), 7.31 (1H, t, J = 7.6 Hz), 7.38-7.46 (1H, m), 7.65 (1H, s), 10.90 (1H, s)	407	405	S-form




105		¹ H-NMR (DMSO-D ₆) δ : 2.36-2.42 (m, 2H), 3.30-3.54 (m, 3H), 7.01 (s, 1H), 7.15 (s, 1H), 7.27 (dt, 2H, J = 9.3, 2.6 Hz), 7.35 (d, 1H, J = 8.0 Hz), 7.39 (d, 1H, J = 8.0 Hz), 7.49 (dt, 2H, J = 9.3, 2.6 Hz), 7.55 (t, 1H, J = 8.0 Hz), 7.65 (s, 1H), 10.94 (s, 1H).	465	463	R-form
106		¹ H-NMR (DMSO-D ₆) δ : 2.35-2.43 (m, 2H), 3.33-3.54 (m, 3H), 7.02 (s, 1H), 7.13-7.21 (m, 2H), 7.34-7.47 (m, 5H), 7.56 (t, 1H, J = 8.0 Hz), 7.65 (s, 1H), 10.94 (s, 1H).	465	463	R-form
107		¹ H-NMR (DMSO-D ₆) δ : 0.88 (t, 3H, J = 7.3 Hz), 1.55 (tq, 2H, J = 6.9, 7.3 Hz), 2.34-2.40 (m, 2H), 2.51 (t, 2H, J = 6.9 Hz), 3.06 (t, 2H, J = 7.1 Hz), 3.29-3.34 (m, 1H), 3.36-3.44 (m, 1H), 3.46-3.52 (m, 1H), 4.15 (t, 2H, J = 7.1 Hz), 6.53 (s, 1H), 6.80 (s, 1H), 6.95-7.00 (m, 2H), 7.00 (d, 1H, J = 7.7 Hz), 7.17-7.26 (m, 4H), 7.31 (dd, 1H, J = 7.7, 7.7 Hz), 7.63 (s, 1H), 10.74 (s, 1H).	417	415	R-form




108		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 0.88 (t, 3H, J = 7.3 Hz), 1.55 (tq, 2H, J = 6.9, 7.3 Hz), 2.34-2.40 (m, 2H), 2.51 (t, 2H, J = 6.9 Hz), 3.06 (t, 2H, J = 7.1 Hz), 3.29-3.34 (m, 1H), 3.36-3.44 (m, 1H), 3.46-3.52 (m, 1H), 4.15 (t, 2H, J = 7.1 Hz), 6.53 (s, 1H), 6.80 (s, 1H), 6.95-7.00 (m, 2H), 7.00 (d, 1H, J = 7.7 Hz), 7.17-7.26 (m, 4H), 7.31 (dd, 1H, J = 7.7, 7.7 Hz), 7.63 (s, 1H), 10.74 (s, 1H).</p>	417	415	S-form
109		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 1.26 (s, 9H), 2.35-2.39 (m, 2H), 3.06 (t, 2H, J = 7.1 Hz), 3.29-3.34 (m, 1H), 3.36-3.45 (m, 1H), 3.46-3.52 (m, 1H), 4.17 (t, 2H, J = 7.1 Hz), 6.54 (s, 1H), 6.94-6.99 (m, 2H), 6.99 (d, 1H, J = 7.7 Hz), 7.14-7.24 (m, 4H), 7.34 (dd, 1H, J = 7.7, 8.1 Hz), 7.44 (d, 1H, J = 8.1 Hz), 7.63 (s, 1H), 10.74 (s, 1H).</p>	431	429	R-form




110		¹ H-NMR (DMSO-D ₆) δ : 2.35-2.40 (m, 2H), 3.03 (t, 2H, J = 6.9 Hz), 3.29-3.35 (m, 1H), 3.36-3.45 (m, 1H), 3.46-3.52 (m, 1H), 4.19 (t, 2H, J = 6.9 Hz), 6.59 (s, 1H), 6.90-6.95 (m, 2H), 6.98 (s, 1H), 7.14-7.23 (m, 4H), 7.40 (d, 1H, J = 8.1 Hz), 7.53 (dd, 1H, J = 7.7, 8.1 Hz), 7.64 (s, 1H), 10.79 (s, 1H).	459	457	R-form
111		¹ H-NMR (DMSO-D ₆) δ : 0.67-0.81 (m, 2H), 0.89 (t, 3H, J = 7.3 Hz), 0.99-1.14 (m, 3H), 1.37-1.46 (m, 2H), 1.48-1.67 (m, 3H), 1.62 (tq, 2H, J = 7.3, 7.3 Hz), 1.69-1.80 (m, 1H), 2.30-2.41 (m, 2H), 2.62 (t, 2H, J = 7.3 Hz), 3.27-3.35 (m, 1H), 3.35-3.43 (m, 1H), 3.44-3.50 (m, 1H), 3.84 (d, 2H, J = 7.3 Hz), 6.57 (s, 1H), 7.22-7.29 (m, 3H), 7.40 (dd, 1H, J = 8.1, 7.7 Hz), 7.62 (s, 1H), 10.65 (s, 1H).	409	407	R-form




112		¹ H-NMR (DMSO-D ₆) δ : 0.67-0.80 (m, 2H), 0.89 (t, 3H, $J = 7.3$ Hz), 0.99-1.13 (m, 3H), 1.37-1.46 (m, 2H), 1.48-1.67 (m, 3H), 1.62 (tq, 2H, $J = 7.3, 7.5$ Hz), 1.69-1.79 (m, 1H), 2.30-2.41 (m, 2H), 2.62 (t, 2H, $J = 7.5$ Hz), 3.27-3.35 (m, 1H), 3.35-3.43 (m, 1H), 3.43-3.50 (m, 1H), 3.84 (d, 2H, $J = 7.3$ Hz), 6.57 (s, 1H), 7.22-7.29 (m, 3H), 7.40 (dd, 1H, $J = 7.7, 8.1$ Hz), 7.62 (s, 1H), 10.65 (s, 1H).	409	407	S-form
113		¹ H-NMR (DMSO-D ₆) δ : 0.71-0.83 (m, 2H), 0.90 (t, 3H, $J = 7.3$ Hz), 0.97-1.13 (m, 4H), 1.41-1.48 (m, 2H), 1.49-1.67 (m, 5H), 1.63 (tq, 2H, $J = 7.3, 7.5$ Hz), 2.30-2.41 (m, 2H), 2.62 (t, 2H, $J = 7.5$ Hz), 3.26-3.34 (m, 1H), 3.35-3.42 (m, 1H), 3.44-3.50 (m, 1H), 4.01 (t, 2H, $J = 7.3$ Hz), 6.56 (s, 1H), 7.23-7.30 (m, 3H), 7.41 (dd, 1H, $J = 7.7, 8.1$ Hz), 7.62 (s, 1H), 10.66 (s, 1H).	423	421	R-form





114		¹ H-NMR (DMSO-D ₆) δ : 0.71-0.84 (m, 2H), 0.90 (t, 3H, $J = 7.3$ Hz), 0.99-1.11 (m, 4H), 1.41-1.49 (m, 2H), 1.50-1.68 (m, 5H), 1.63 (tq, 2H, $J = 7.3, 7.7$ Hz), 2.30-2.41 (m, 2H), 2.62 (t, 2H, $J = 7.7$ Hz), 3.26-3.35 (m, 1H), 3.35-3.42 (m, 1H), 3.44-3.50 (m, 1H), 4.01 (t, 2H, $J = 7.3$ Hz), 6.56 (s, 1H), 7.23-7.30 (m, 3H), 7.41 (dd, 1H, $J = 7.7, 8.1$ Hz), 7.62 (s, 1H), 10.66 (s, 1H).	423	421	S-form
115		¹ H-NMR (DMSO-D ₆) δ : 0.77 (t, 3H, $J = 7.3$ Hz), 1.38-1.59 (m, 1H), 1.80-1.93 (m, 1H), 2.46 (t, 3H, $J = 7.5$ Hz), 2.58-2.70 (m, 1H), 3.12-3.24 (m, 2H), 3.41-3.53 (m, 3H), 4.57 (t, 1H, $J = 5.0$ Hz), 6.92 (s, 1H), 7.01 (s, 1H), 7.07 (d, 1H, $J = 7.7$ Hz), 7.17 (d, 1H, $J = 7.5$ Hz), 7.20-7.43 (m, 6H), 7.72 (s, 1H), 10.93 (s, 1H).	433	431	trans-form (Racemate)




116		¹ H-NMR (DMSO-D ₆) δ: 0.73 (3H, t, J = 7.4 Hz), 1.41 (2H, tq, J = 7.3, 7.4 Hz), 2.36-2.47 (4H, m), 3.30-3.55 (3H, m), 6.93-7.01 (2H, m), 7.06 (1H, d, J = 7.7 Hz), 7.14 (1H, d, J = 7.7 Hz), 7.24 (1H, t, J = 7.6 Hz), 7.28-7.37 (2H, m), 7.45-7.57 (2H, m), 7.65 (1H, s), 10.84 (1H, s)	407	405	S-form
117		¹ H-NMR (DMSO-D ₆) δ: 0.74 (3H, t, J = 7.4 Hz), 1.41 (2H, tq, J = 7.3, 7.4 Hz), 2.35-2.48 (4H, m), 3.30-3.55 (3H, m), 6.93-7.01 (2H, m), 7.06 (1H, d, J = 7.5 Hz), 7.14 (1H, d, J = 7.5 Hz), 7.24 (1H, t, J = 7.6 Hz), 7.28-7.37 (2H, m), 7.45-7.58 (2H, m), 7.65 (1H, s), 10.84 (1H, s)	407	405	R-form
118		¹ H-NMR (DMSO-D ₆) δ: 2.31-2.41 (m, 2H), 3.30-3.53 (m, 3H), 6.60-6.66 (m, 2H), 6.72-6.77 (m, 1H), 6.83 (s, 1H), 7.14 (t, 1H, J = 8.2 Hz), 7.23 (d, 2H, J = 7.1 Hz), 7.29-7.44 (m, 3H), 7.64 (s, 1H), 9.58 (s, 1H), 10.85 (s, 1H).	363	361	R-form



119		¹ H-NMR (DMSO-D ₆) δ : 2.36-2.42 (m, 2H), 3.33-3.54 (m, 3H), 4.45 (d, 2H, $J = 5.7$ Hz), 5.21 (t, 1H, $J = 5.7$ Hz), 6.90 (s, 1H), 6.99 (d, 1H, $J = 6.6$ Hz), 7.20-7.44 (m, 8H), 7.64 (s, 1H), 10.86 (s, 1H).	377	375	R-form
120		¹ H-NMR (DMSO-D ₆) δ : 2.26 (s, 3H), 2.36-2.41 (m, 2H), 3.33-3.54 (m, 3H), 6.97-7.03 (m, 2H), 7.08 (s, 1H), 7.10 (s, 1H), 7.19 (d, 1H, $J = 7.5$ Hz), 7.29 (t, 1H, $J = 7.7$ Hz), 7.34-7.40 (m, 2H), 7.53 (t, 1H, $J = 8.0$ Hz), 7.65 (s, 1H), 10.89 (s, 1H).	445	443	R-form
121		¹ H-NMR (DMSO-D ₆) δ : 0.77 (3H, t, $J = 7.3$ Hz), 1.45 (2H, tq, $J = 7.3, 7.6$ Hz), 2.36-2.41 (2H, m), 2.42-2.50 (2H, t, $J = 7.5$ Hz), 3.29-3.54 (3H, m), 3.76 (3H, s), 6.86 (1H, s), 6.91-6.98 (2H, m), 7.00 (1H, s), 7.05 (1H, d, $J = 7.6$ Hz), 7.12-7.19 (3H, m), 7.25 (1H, t, $J = 7.6$ Hz), 7.64 (1H, s), 10.8 (1H, m)	419	417	R-form




122		¹ H-NMR (DMSO-D ₆) δ : 0.76 (t, 3H, J = 7.3 Hz), 1.41 (tq, 2H, J = 7.5, 7.3 Hz), 2.31-2.46 (m, 4H), 3.36-3.54 (m, 3H), 4.64 (s, 2H), 6.77-6.92 (m, 3H), 7.06 (dd, 1H, J = 8.6, 2.2 Hz), 7.20-7.44 (m, 5H), 7.64 (s, 1H), 10.82 (s, 1H).	463	461	R-form
123		¹ H-NMR (DMSO-D ₆) δ : 0.78 (3H, t, J = 7.4 Hz), 1.35-1.55 (2H, m), 2.35-2.55 (4H, m), 3.25-3.6 (3H, m), 3.65 (3H, s), 6.7-6.83 (2H, m), 6.85-6.95 (2H, m), 7.03 (1H, s), 7.09 (1H, d, J = 8.0 Hz), 7.19 (1H, d, J = 8.0 Hz), 7.22-7.38 (2H, m), 7.64 (1H, s), 10.87 (1H, s)	419	417	R-form
124		¹ H-NMR (DMSO-D ₆) δ : 0.79 (3H, t, J = 7.3 Hz), 1.47 (2H, tq, J = 7.3, 7.4 Hz), 2.36-2.42 (2H, m), 2.45-2.52 (2H, m), 2.99-3.55 (3H, m), 4.59 (2H, s), 6.75 (1H, dd, J = 8.0, 1.1 Hz), 6.80 (1H, t, J = 2.2Hz), 6.86-6.92 (2H, m), 7.03 (1H, s), 7.07 (1H, d, J = 7.7 Hz), 7.18 (1H, d, J = 7.7 Hz), 7.23-7.32 (2H, m), 7.64 (1H, s), 10.88 (1H, s), 13.01 (1H, s)	463	461	R-form




125		¹ H-NMR (DMSO-D ₆) δ : 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.5, 7.3 Hz), 2.46 (t, 3H, J = 7.5 Hz), 2.64 (s, 3H), 3.21 (dd, 1H, J = 8.8, 6.3 Hz), 3.54 (dd, 1H, J = 9.4, 8.8 Hz), 4.28 (dd, 1H, J = 9.4, 6.3 Hz), 6.44 (s, 1H), 6.92 (s, 1H), 7.01 (s, 1H), 7.08 (d, 1H, J = 7.0 Hz), 7.17 (d, 1H, J = 7.0 Hz), 7.20-7.44 (m, 6H), 11.02 (s, 1H).	404	402	S-form
126		¹ H-NMR (DMSO-D ₆) δ : 1.13 (9H, s), 2.39 (2H, d, J = 7.9 Hz), 3.30-3.55 (3H, m), 6.94 (1H, s), 7.09 (1H, s), 7.17-7.26 (2H, m), 7.28 (1H, s), 7.33-7.48 (4H, m), 7.65 (1H, s), 10.90 (1H, s)	437	435	R-form
127		¹ H-NMR (DMSO-D ₆) δ : 1.10 (9H, s), 2.25 (3H, s), 2.36-2.42 (2H, m), 3.3-3.55 (3H, m), 6.91 (1H, s), 7.00 (1H, d, J = 7.9 Hz), 7.04-7.10 (2H, m), 7.14-7.23 (2H, m), 7.24-7.41 (3H, m), 7.65 (1H, s), 10.84 (1H, s)	417	415	R-form




128		¹ H-NMR (DMSO-D ₆) δ: 0.69-0.85 (m, 2H), 0.94-1.14 (m, 4H), 1.38-1.61 (m, 7H), 2.31-2.40 (m, 2H), 3.26-3.51 (m, 3H), 4.04 (t, 2H, J = 7.2 Hz), 6.66 (s, 1H), 7.45-7.69 (m, 5H), 10.71 (s, 1H).	465	463	R-form
129		¹ H-NMR (DMSO-D ₆) δ: 0.69-0.84 (m, 2H), 0.95-1.13 (m, 4H), 1.32 (s, 9H), 1.40-1.64 (m, 7H), 2.32-2.39 (m, 2H), 3.26-3.51 (m, 3H), 4.01 (t, 2H, J = 7.3 Hz), 6.57 (s, 1H), 7.25 (d, 1H, J = 7.3 Hz), 7.38-7.52 (m, 3H), 7.62 (s, 1H), 10.67 (s, 1H).	437	435	R-form
130		¹ H-NMR (DMSO-D ₆) δ: 2.40 (d, 2H, J = 8.5 Hz), 2.40 (d, 2H, J = 8.4 Hz), 3.34 (dd, 1H, J = 8.5, 8.6 Hz), 3.42 (dddd, 1H, J = 8.4, 8.5, 8.5, 8.6 Hz), 3.51 (dd, 1H, J = 8.5, 8.5 Hz), 7.01 (s, 1H), 7.09 (s, 1H), 7.31-7.35 (m, 2H), 7.47-7.56 (m, 3H), 7.60-7.66 (m, 3H), 10.88 (s, 1H).	465	463	R-form
131		¹ H-NMR (DMSO-D ₆) δ: 2.40 (2H, d, J = 7.9Hz), 3.30-3.55 (3H, m), 7.07 (2H, s), 7.29-7.40 (4H, m), 7.47-7.56 (2H, m), 7.60 (1H, dt, J = 7.8, 1.5 Hz), 7.65 (1H, s), 10.90 (1H, s)	449	447	R-form





132		<p>¹H-NMR (DMSO-D₆) δ:</p> <p>1.06 (9H, s), 2.36-2.43 (2H, m), 3.30-3.55 (3H, m), 6.96-7.03 (2H, m), 7.16-7.23 (1H, m), 7.25-7.36 (2H, m), 7.43-7.58 (3H, m), 7.59-7.69 (2H, m), 10.82 (1H, s)</p>	437	435	R-form
133		<p>¹H-NMR (DMSO-D₆) δ:</p> <p>0.72 (3H, t, J = 7.2 Hz), 1.38 (2H, tq, J = 7.2, 7.4 Hz), 2.30-2.45 (4H, m), 3.29-3.53 (6H, m), 6.85 (1H, s), 6.93 (1H, s), 6.96-7.11 (4H, m), 7.18 (1H, t, J = 7.6 Hz), 7.33 (1H, d, J = 7.7 Hz), 7.40 (1H, t, d = 7.9 Hz), 7.63 (1H, s), 10.72 (1H, s)</p>	419	417	R-form
134		<p>¹H-NMR (DMSO-D₆) δ:</p> <p>0.82 (s, 9H), 0.89 (t, 3H, J = 7.3 Hz), 1.57-1.69 (m, 4H), 2.32-2.39 (m, 2H), 2.62 (t, 2H, J = 7.5 Hz), 3.26-3.40 (m, 2H), 3.47 (t, 1H, J = 8.5 Hz), 3.96-4.05 (m, 2H), 6.57 (s, 1H), 7.24-7.31 (m, 3H), 7.41 (t, 1H, J = 7.8 Hz), 7.62 (s, 1H), 10.66 (s, 1H).</p>	397	395	R-form




135		<p>¹H-NMR (DMSO-D₆) δ: 0.75 (d, 6H, J = 6.6 Hz), 0.89 (t, 3H, J = 7.4 Hz), 1.36-1.43 (m, 1H), 1.52-1.69 (m, 4H), 2.32-2.39 (m, 2H), 2.62 (t, 2H, J = 7.4 Hz), 3.26-3.50 (m, 3H), 4.01 (t, 2H, J = 7.3 Hz), 6.57 (s, 1H), 7.24-7.30 (m, 3H), 7.41 (t, 1H, J = 7.8 Hz), 7.62 (s, 1H), 10.65 (s, 1H).</p>	383	381	R-form
136		<p>¹H-NMR (DMSO-D₆) δ: 0.91 (t, 3H, J = 7.5 Hz), 1.63 (tq, 2H, J = 7.5, 7.7 Hz), 1.75-1.82 (m, 2H), 2.00-2.12 (m, 2H), 2.30-2.41 (m, 2H), 2.63 (t, 2H, J = 7.7 Hz), 3.25-3.34 (m, 3H), 3.35-3.43 (m, 1H), 3.44-3.50 (m, 1H), 3.88-3.95 (m, 2H), 4.24-4.33 (m, 1H), 6.57 (s, 1H), 7.23-7.27 (m, 2H), 7.30 (d, 1H, J = 7.7 Hz), 7.42 (dd, 1H, J = 7.7, 8.1 Hz), 7.62 (s, 1H), 10.72 (s, 1H).</p>	397	395	R-form




137		¹ H-NMR (DMSO-D ₆) δ : 0.87-0.98 (m, 2H), 0.89 (t, 3H, $J = 7.3$ Hz), 1.32-1.43 (m, 2H), 1.43-1.72 (m, 7H), 1.62 (tq, 2H, $J = 7.3, 7.5$ Hz), 2.30-2.41 (m, 2H), 2.62 (t, 2H, $J = 7.5$ Hz), 3.27-3.33 (m, 1H), 3.33-3.42 (m, 1H), 3.44-3.50 (m, 1H), 4.00 (t, 2H, $J = 7.1$ Hz), 6.57 (s, 1H), 7.23-7.30 (m, 3H), 7.41 (dd, 1H, $J = 7.7, 8.1$ Hz), 7.62 (s, 1H), 10.65 (s, 1H).	409	407	R-form
138		¹ H-NMR (DMSO-D ₆) δ : 0.78 (d, 6H, $J = 6.7$ Hz), 2.15-2.30 (m, 1H), 2.36-2.42 (m, 2H), 3.32-3.54 (m, 3H), 6.98 (s, 1H), 7.13 (s, 1H), 7.24 (d, 2H, $J = 7.2$ Hz), 7.32-7.55 (m, 6H), 7.64 (s, 1H), 10.89 (s, 1H).	439	437	R-form
139		¹ H-NMR (DMSO-D ₆) δ : 0.78 (t, 3H, $J = 7.3$ Hz), 2.10 (tq, 2H, $J = 16.5, 7.3$ Hz), 2.36-2.42 (m, 2H), 3.32-3.55 (m, 3H), 7.01 (s, 1H), 7.15-7.21 (m, 1H), 7.28 (s, 1H), 7.33 (s, 1H), 7.39-7.58 (m, 5H), 7.64 (s, 1H), 10.92 (s, 1H).	459	457	R-form




140		¹ H-NMR (DMSO-D ₆) δ : 0.69 (t, 3H, J = 7.4 Hz), 2.02 (tq, 2H, J = 16.0, 7.4 Hz), 2.38-2.41 (m, 2H), 3.33-3.37 (m, 1H), 3.38-3.46 (m, 1H), 3.49-3.54 (m, 1H), 7.06 (s, 1H), 7.16 (s, 1H), 7.41-7.65 (m, 8H), 10.86 (s, 1H).	459	457	R-form
141		¹ H-NMR (DMSO-D ₆) δ : 0.78 (t, 3H, J = 7.4 Hz), 2.10 (tq, 2H, J = 16.2, 7.4 Hz), 2.38-2.40 (m, 2H), 3.33-3.35 (m, 1H), 3.39-3.44 (m, 1H), 3.49-3.52 (m, 1H), 7.00 (s, 1H), 7.24-7.28 (m, 3H), 7.42-7.54 (m, 5H), 7.64 (s, 1H), 10.92 (s, 1H).	459	457	R-form
142		¹ H-NMR (DMSO-D ₆) δ : 0.74 (d, 6H, J = 6.4 Hz), 0.92 (t, 3H, J = 7.5 Hz), 1.39 (tsep, 1H, J = 6.4, 7.0 Hz), 1.57 (dt, 2H, J = 7.0, 7.5 Hz), 2.27 (tq, 2H, J = 16.5, 7.5 Hz), 2.33-2.41 (m, 2H), 3.27-3.33 (m, 1H), 3.34-3.43 (m, 1H), 3.44-3.50 (m, 1H), 4.02 (t, 2H, J = 7.5 Hz), 6.65 (s, 1H), 7.55 (s, 1H), 7.58-7.67 (m, 4H), 10.70 (s, 1H).	419	417	R-form




143		¹ H-NMR (DMSO-D ₆) δ: 0.82 (s, 9H), 0.92 (t, 3H, J = 7.5 Hz), 1.58-1.66 (m, 2H), 2.27 (tq, 2H, J = 16.5, 7.5 Hz), 2.33-2.39 (m, 2H), 3.27-3.34 (m, 1H), 3.34-3.42 (m, 1H), 3.44-3.51 (m, 1H), 3.97-4.05 (m, 2H), 6.66 (s, 1H), 7.56 (s, 1H), 7.58-7.67 (m, 4H), 10.70 (s, 1H).	433	431	R-form
144		¹ H-NMR (DMSO-D ₆) δ: 0.69 (3H, t, J = 7.5 Hz), 1.89 (3H, s), 1.92-2.08 (2H, m), 2.36-2.43 (2H, m), 3.3-3.55 (3H, m), 7.05 (1H, s), 7.13 (1H, s), 7.25-7.50 (7H, m), 7.65 (1H, s), 10.81 (1H, s)	439	437	R-form
145		¹ H-NMR (DMSO-D ₆) δ: 0.76 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.5, 7.3 Hz), 2.36-2.43 (m, 2H), 2.44-2.49 (m, 2H), 3.35-3.54 (m, 3H), 6.92 (s, 1H), 7.10 (s, 1H), 7.23 (d, 2H, J = 7.3 Hz), 7.30-7.44 (m, 3H), 7.59-7.72 (m, 3H), 10.87 (s, 1H).	433	431	R-form



146		¹ H-NMR (DMSO-D ₆) δ: 0.74 (t, 3H, J = 7.3 Hz), 1.40 (tq, 2H, J = 7.5, 7.3 Hz), 2.35-2.45 (m, 4H), 3.33-3.54 (m, 3H), 4.43 (d, 2H, J = 5.7 Hz), 5.17 (t, 1H, J = 5.7 Hz), 6.76 (s, 1H), 6.89 (s, 1H), 7.12 (d, 2H, J = 5.3 Hz), 7.23 (d, 2H, J = 7.3 Hz), 7.31-7.43 (m, 3H), 7.64 (s, 1H), 10.85 (s, 1H).	419	417	R-form
147		¹ H-NMR (DMSO-D ₆) δ: 2.35-2.42 (m, 2H), 3.33-3.54 (m, 3H), 7.00 (s, 1H), 7.10 (s, 1H), 7.24-7.40 (m, 6H), 7.53 (t, 1H, J = 7.9 Hz), 7.64 (s, 1H), 10.90 (s, 1H).	449	447	R-form
148		¹ H-NMR (DMSO-D ₆) δ: 2.39 (2H, d, J = 8.6 Hz), 3.30-3.55 (3H, m), 4.73 (2H, q, J = 8.9 Hz), 6.80 (1H, d, J = 7.9 Hz), 6.97(1H, s), 6.98-7.08 (2H, m), 7.20-7.45 (6H, m), 7.64 (1H, s), 10.87 (1H, s)	445	443	R-form
149		¹ H-NMR (DMSO-D ₆) δ: 2.39 (2H, d, J = 8.4 Hz), 2.61-2.76 (2H, m), 3.30-3.55 (3H, m), 4.11 (2H, t, J = 5.7 Hz), 6.77-6.84 (2H, m), 6.91-6.98 (2H, m), 7.21-7.44 (6H, m), 7.64 (1H, s), 10.86 (1H, s)	459	457	R-form




150		¹ H-NMR (DMSO-D ₆) δ: 1.84-1.98 (m, 2H), 2.14-2.29 (m, 2H), 2.81-2.88 (m, 1H), 3.27-3.33 (m, 2H), 6.99 (s, 1H), 7.08 (s, 1H), 7.24-7.26 (m, 2H), 7.34-7.54 (m, 7H), 10.88 (s, 1H).	445	443	Optically active substance (The same configuration as Example 5)
151		¹ H-NMR (DMSO-D ₆) δ: 0.76 (t, 3H, J = 7.4 Hz), 1.88-2.28 (m, 6H), 2.82-2.87 (m, 1H), 3.30-3.32 (m, 2H), 6.98 (s, 1H), 7.22-7.25 (m, 3H), 7.34-7.52 (m, 7H), 10.87 (s, 1H).	439	437	Optically active substance (The same configuration as Example 5)
152		¹ H-NMR (DMSO-D ₆) δ: 1.08 (d, 3H, J = 7.2 Hz), 2.54 (dq, 1H, J = 9.6, 7.2 Hz), 3.04 (ddd, 1H, J = 8.6, 9.0, 9.6 Hz), 3.23 (dd, 1H, J = 9.0, 8.4 Hz), 3.44 (dd, 1H, J = 8.6, 8.4 Hz), 7.03 (s, 1H), 7.15 (s, 1H), 7.26-7.29 (m, 2H), 7.34-7.40 (m, 2H), 7.48-7.56 (m, 3H), 7.68 (s, 1H), 11.02 (s, 1H).	479	477	(3R, 4R)-form




153		¹ H-NMR (DMSO-D6) δ : 1.13 (s, 9H), 2.38-2.40 (m, 2H), 3.33-3.36 (m, 1H), 3.39-3.45 (m, 1H), 3.49-3.51 (m, 1H), 6.93 (s, 1H), 7.09 (s, 1H), 7.17 (d, 1H, $J = 7.4$ Hz), 7.25 (d, 2H, $J = 8.8$ Hz), 7.34 (dd, 1H, $J = 7.4, 8.1$ Hz), 7.39 (d, 1H, $J = 8.1$ Hz), 7.48 (d, 2H, $J = 8.8$ Hz), 7.64 (s, 1H), 10.88 (s, 1H).	437	435	R-form
154		¹ H-NMR (DMSO-D6) δ : 0.76 (t, 3H, $J = 7.5$ Hz), 1.09 (d, 3H, $J = 7.3$ Hz), 2.06 (tq, 2H, $J = 16.5, 7.5$ Hz), 2.55 (dq, 1H, $J = 9.7, 7.3$ Hz), 3.04 (ddd, 1H, $J = 8.5, 9.1, 9.7$ Hz), 3.23 (dd, 1H, $J = 8.5, 9.1$ Hz), 3.45 (dd, 1H, $J = 8.5, 8.5$ Hz), 7.02 (s, 1H), 7.22-7.27 (m, 3H), 7.34-7.45 (m, 4H), 7.46-7.53 (m, 2H), 7.68 (s, 1H), 10.98 (s, 1H).	439	437	(3R, 4R)-form
155		¹ H-NMR (DMSO-D6) δ : 0.94 (t, 3H, $J = 7.5$ Hz), 1.75-1.85 (m, 2H), 2.00-2.13 (m, 2H), 2.28 (tq, 2H, $J = 16.5, 7.5$ Hz), 2.33-2.39 (m, 2H), 3.26-3.34 (m, 3H), 3.35-3.44 (m, 1H), 3.45-3.51 (m, 1H), 3.88-3.95 (m, 2H), 4.21-4.31 (m, 1H), 6.65 (s, 1H), 7.53 (s, 1H), 7.57-7.68 (m, 4H), 10.76 (s, 1H).	433	431	R-form




156		¹ H-NMR (DMSO-D ₆) δ: 1.91 (3H, s), 2.36-2.43 (2H, m), 3.30-3.55 (3H, m), 6.97 (1H, s), 7.07 (1H, s), 7.25-7.43 (6H, m), 7.47 (1H, t, J = 8.1 Hz), 7.64 (1H, s), 10.82 (1H, s)	445	443	R-form
157		¹ H-NMR (DMSO-D ₆) δ: 2.38-2.40 (m, 2H), 3.32-3.35 (m, 1H), 3.38-3.45 (m, 1H), 3.49-3.53 (m, 1H), 7.02 (s, 1H), 7.05-7.07 (m, 1H), 7.13-7.26 (m, 3H), 7.38-7.48 (m, 3H), 7.56 (dd, 1H, J = 8.0, 8.0 Hz), 7.65 (s, 1H), 10.94 (s, 1H).	449	447	R-form
158		¹ H-NMR (DMSO-D ₆) δ: 0.80 (d, 6H, J = 7.0 Hz), 2.23-2.40 (m, 3H), 3.34-3.36 (m, 1H), 3.39-3.45 (m, 1H), 3.48-3.53 (m, 1H), 6.99 (s, 1H), 7.14 (s, 1H), 7.26 (d, 2H, J = 8.6 Hz), 7.46-7.56 (m, 5H), 7.64 (s, 1H), 10.92 (s, 1H).	473	471	R-form




159		¹ H-NMR (DMSO-D ₆) δ : 2.38-2.40 (m, 2H), 3.33-3.36 (m, 1H), 3.39-3.46 (m, 1H), 3.48-3.52 (m, 1H), 4.76 (q, 2H, J = 8.8 Hz), 6.81 (d, 1H, J = 7.4 Hz), 6.98 (s, 1H), 7.05 (s, 1H), 7.07 (d, 1H, J = 8.0 Hz), 7.25 (d, 2H, J = 8.8 Hz), 7.32 (dd, 1H, J = 7.4, 8.0 Hz), 7.47 (d, 2H, J = 8.8 Hz), 7.64 (s, 1H), 10.90 (s, 1H).	479	477	R-form
160		¹ H-NMR (DMSO-D ₆) δ : 2.37-2.40 (m, 2H), 2.71 (tq, 2H, J = 5.9, 11.0 Hz), 3.32-3.35 (m, 1H), 3.38-3.46 (m, 1H), 3.48-3.52 (m, 1H), 4.15 (t, 2H, J = 5.9 Hz), 6.82 (d, 1H, J = 7.9 Hz), 6.84 (s, 1H), 6.94 (s, 1H), 6.97 (d, 1H, J = 8.4 Hz), 7.26 (d, 2H, J = 8.6 Hz), 7.30 (dd, 1H, J = 7.9, 8.4 Hz), 7.47 (d, 2H, J = 8.6 Hz), 7.64 (s, 1H), 10.89 (s, 1H).	493	491	R-form
161		¹ H-NMR (DMSO-D ₆) δ : 1.84 (t, 3H, J = 18.8 Hz), 2.38-2.40 (m, 2H), 3.33-3.35 (m, 1H), 3.39-3.46 (m, 1H), 3.47-3.53 (m, 1H), 7.00 (s, 1H), 7.23-7.26 (m, 2H), 7.33-7.54 (m, 7H), 7.64 (s, 1H), 10.89 (s, 1H).	411	409	R-form



162		<p>¹H-NMR (DMSO-D6) δ: 1.89 (t, 3H, J = 19.0 Hz), 2.39 (d, 2H, J = 8.1 Hz), 3.33 (dd, 1H, J = 8.6, 8.6 Hz), 3.42 (ddt, 1H, J = 8.1, 8.6, 8.6 Hz), 3.51 (dd, 1H, J = 8.6, 8.6 Hz), 7.01 (s, 1H), 7.27 (d, 2H, J = 8.6 Hz), 7.36 (d, 1H, J = 7.4 Hz), 7.40 (s, 1H), 7.49 (d, 2H, J = 8.6 Hz), 7.50 (dd, 1H, J = 7.4, 7.7 Hz), 7.56 (d, 1H, J = 7.7 Hz), 7.64 (s, 1H), 10.92 (s, 1H).</p>	445	443	R-form
163		<p>¹H-NMR (DMSO-D6) δ: 2.39 (2H, d, J = 8.2 Hz), 3.30-3.55 (3H, m), 3.67 (3H, s), 6.76 (1H, dd, J = 7.6, 1.4 Hz), 6.83 (1H, t, J = 2.2 Hz), 6.94 (1H, dd, J = 8.1, 2.2 Hz), 6.99 (1H, s), 7.13 (1H, s), 7.30 (1H, t, J = 8.1 Hz), 7.34-7.41 (2H, m), 7.54 (1H, t, J = 8.1 Hz), 7.64 (1H, s), 10.92 (1H, s)</p>	461	459	R-form



164		¹ H-NMR (DMSO-D ₆) δ: 0.77 (3H, t, J = 7.4 Hz), 2.08 (2H, m), 2.39 (2H, d, J = 8.4 Hz), 3.30-3.55 (3H, m), 3.66 (3H, s), 6.76 (1H, dd, J = 7.8, 1.2 Hz), 6.81 (1H, t, J=2.2 Hz), 6.92 (1H, dd, J = 8.2, 2.2 Hz), 6.98 (1H, s), 7.24-7.32 (2H, m), 7.41- 7.56 (3H, m), 7.64 (1H, s), 10.91 (1H, s)	455	453	R-form
165		¹ H-NMR (DMSO-D ₆) δ: 1.86-1.99 (m, 2H), 2.13-2.28 (m, 2H), 2.80- 2.88 (m, 1H), 3.28-3.31 (m, 2H), 6.99 (s, 1H), 7.10 (s, 1H), 7.25-7.37 (m, 6H), 7.51-7.54 (m, 2H), 10.87 (s, 1H).	463	461	Optically active substance (The same configurati on as Example 5)
166		¹ H-NMR (DMSO-D ₆) δ: 1.09 (d, 3H, J = 7.0 Hz), 2.55 (dq, 1H, J = 9.3, 7.0 Hz), 3.03 (ddd, 1H, J = 9.1, 9.2, 9.3 Hz), 3.23 (dd, 1H, J = 9.2, 8.8 Hz), 3.45 (dd, 1H, J = 9.1, 8.8 Hz), 7.08 (s, 1H), 7.09 (s, 1H), 7.31-7.37 (m, 4H), 7.49-7.62 (m, 3H), 7.68 (s, 1H), 10.99 (s, 1H).	463	461	(3R, 4R)- form




167		¹ H-NMR (DMSO-D ₆) δ: 1.87-1.99 (m, 2H), 2.19-2.24 (m, 2H), 2.83-2.85 (m, 1H), 3.27-3.32 (m, 2H), 7.06-7.07 (m, 2H), 7.31-7.38 (m, 4H), 7.50-7.59 (m, 4H), 10.88 (s, 1H).	463	461	Optically active substance (The same configuration as Example 5)
168		¹ H-NMR (DMSO-D ₆) δ: 1.88-1.98 (m, 2H), 2.13-2.33 (m, 2H), 2.83-2.86 (m, 1H), 3.28-3.32 (m, 2H), 7.01 (s, 1H), 7.04-7.06 (m, 1H), 7.14-7.26 (m, 3H), 7.39-7.58 (m, 5H), 10.92 (s, 1H).	463	461	Optically active substance (The same configuration as Example 5)
169		¹ H-NMR (DMSO-D ₆) δ: 0.93 (t, 3H, J = 7.4 Hz), 1.55-1.57 (m, 4H), 2.23-2.36 (m, 8H), 2.77 (t, 2H, J = 6.4 Hz), 3.28-3.32 (m, 1H), 3.36-3.42 (m, 1H), 3.47-3.49 (m, 1H), 4.07 (t, 2H, J = 6.4 Hz), 6.65 (s, 1H), 7.61-7.66 (m, 5H), 10.71 (s, 1H).	446	444	R-form




170		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, J = 7.4 Hz), 1.09 (d, 3H, J = 7.3 Hz), 2.10 (tq, 2H, J = 16.4, 7.4 Hz), 2.54 (dq, 2H, J = 9.3, 7.3 Hz), 3.04 (ddd, 1H, J = 8.7, 8.8, 9.3 Hz), 3.23 (dd, 1H, J = 8.7, 8.8 Hz), 3.45 (dd, 1H, J = 8.7, 8.7 Hz), 7.03-7.28 (m, 5H), 7.41-7.53 (m, 4H), 7.68 (s, 1H), 11.01 (s, 1H).	457	455	(3R, 4R)-form
171		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 1.90-2.23 (m, 6H), 2.83-2.86 (m, 1H), 3.28-3.31 (m, 2H), 6.99 (s, 1H), 7.04-7.24 (m, 3H), 7.27 (s, 1H), 7.41-7.56 (m, 5H), 10.90 (s, 1H).	457	455	Optically active substance (The same configuration as Example 5)
172		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 1.08 (d, 3H, J = 7.3 Hz), 2.09 (tq, 2H, J = 16.5, 7.4 Hz), 2.54 (dq, 1H, J = 9.3, 7.4 Hz), 3.04 (ddd, 1H, J = 9.3, 8.8, 8.4 Hz), 3.23 (dd, 1H, J = 8.8, 8.6 Hz), 3.45 (dd, 1H, J = 8.6, 8.4 Hz), 7.02 (s, 1H), 7.24-7.32 (m, 5H), 7.41-7.53 (m, 3H), 7.68 (s, 1H), 10.97 (s, 1H).	457	455	(3R, 4R)-form



173		¹ H-NMR (DMSO-D ₆) δ : 0.77 (t, 3H, J = 7.5 Hz), 2.09 (tq, 2H, J = 16.5, 7.5 Hz), 2.38-2.40 (m, 2H), 3.34-3.35 (m, 1H), 3.38-3.46 (m, 1H), 3.50-3.52 (m, 1H), 6.99 (s, 1H), 7.23-7.32 (m, 5H), 7.42-7.54 (m, 3H), 7.65 (s, 1H), 10.89 (s, 1H).	443	441	R-form
174		¹ H-NMR (DMSO-D ₆) δ : 2.31-2.33 (m, 2H), 3.23-3.28 (m, 1H), 3.36-3.38 (m, 1H), 3.43-3.45 (m, 1H), 6.56 (s, 1H), 7.18 (s, 1H), 7.25-7.30 (m, 2H), 7.39-7.41 (m, 1H), 7.48-7.55 (m, 2H), 7.60 (s, 1H), 7.82 (dd, 1H, J = 8.4, 8.4 Hz), 10.71 (s, 1H).	483	481	R-form
175		¹ H-NMR (DMSO-D ₆) δ : 0.93 (t, 3H, J = 7.5 Hz), 0.99 (d, 6H, J = 6.0 Hz), 2.26 (tq, 2H, J = 16.5, 7.5 Hz), 2.34-2.39 (m, 2H), 3.28-3.33 (m, 1H), 3.34-3.51 (m, 3H), 3.75 (t, 2H, J = 5.2 Hz), 4.07 (t, 2H, J = 5.2 Hz), 6.65 (s, 1H), 7.57-7.64 (m, 3H), 7.68 (d, 1H, J = 6.9 Hz), 7.77 (s, 1H), 10.72 (s, 1H).	435	433	R-form




176		<p>¹H-NMR (DMSO-D₆) δ: 0.93 (t, 3H, J = 7.5 Hz), 0.99 (d, 6H, J = 6.0 Hz), 1.07 (d, 3H, J = 6.9 Hz), 2.26 (tq, 2H, J = 16.5, 7.5 Hz), 2.51 (dq, 1H, J = 6.9, 9.7 Hz), 3.01 (ddd, 1H, J = 8.3, 8.9, 9.7 Hz), 3.20 (dd, 1H, J = 8.9, 8.9 Hz), 3.42 (dd, 1H, J = 8.3, 8.3 Hz), 3.44 (sep, 1H, J = 6.0 Hz), 3.75 (t, 2H, J = 5.2 Hz), 4.08 (t, 2H, J = 5.2 Hz), 6.68 (s, 1H), 7.57-7.71 (m, 4H), 7.77 (s, 1H), 10.80 (s, 1H).</p>	449	447	(3R, 4R)-form
177		<p>¹H-NMR (DMSO-D₆) δ: 1.32 (s, 9H), 1.76-1.85 (m, 2H), 2.03-2.15 (m, 2H), 2.30-2.41 (m, 2H), 3.26-3.35 (m, 3H), 3.35-3.44 (m, 1H), 3.44-3.51 (m, 1H), 3.89-3.96 (m, 2H), 4.24-4.34 (m, 1H), 6.59 (s, 1H), 7.26 (ddd, 1H, J = 7.7, 1.3, 1.0 Hz), 7.40 (dd, 1H, J = 1.6, 1.3 Hz), 7.45 (dd, 1H, J = 7.7, 7.7 Hz), 7.51 (ddd, 1H, J = 7.7, 1.6, 1.0 Hz), 7.62 (s, 1H), 10.72 (s, 1H).</p>	411	409	R-form



178		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 1.06 (d, 3H, J = 6.9 Hz), 1.33 (s, 9H), 1.76-1.85 (m, 2H), 2.02-2.16 (m, 2H), 2.52 (dq, 1H, J = 9.7, 6.9 Hz), 3.01 (ddd, 1H, J = 8.5, 9.1, 9.7 Hz), 3.18 (dd, 1H, J = 9.1, 9.1 Hz), 3.26-3.35 (m, 2H), 3.41 (dd, 1H, J = 8.5, 8.5 Hz), 3.88-3.97 (m, 2H), 4.24-4.35 (m, 1H), 6.61 (s, 1H), 7.27 (ddd, 1H, J = 7.7, 1.6, 1.0 Hz), 7.40 (dd, 1H, J = 1.6, 1.6 Hz), 7.45 (dd, 1H, J = 7.7, 7.7 Hz), 7.51 (ddd, 1H, J = 7.7, 1.6, 1.0 Hz), 7.65 (s, 1H), 10.81 (s, 1H).</p>	425	423	(3R, 4R)-form
179		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 0.94 (t, 3H, J = 7.5 Hz), 1.06 (d, 3H, J = 7.3 Hz), 1.75-1.85 (m, 2H), 2.01-2.14 (m, 2H), 2.28 (tq, 2H, J = 16.5, 7.5 Hz), 2.52 (dq, 1H, J = 9.7, 7.3 Hz), 3.02 (ddd, 1H, J = 8.5, 8.9, 9.7 Hz), 3.18 (dd, 1H, J = 8.9, 8.9 Hz), 3.25-3.35 (m, 2H), 3.42 (dd, 1H, J = 8.5, 8.5 Hz), 3.87-3.97 (m, 2H), 4.21-4.32 (m, 1H), 6.68 (s, 1H), 7.54 (s, 1H), 7.58-7.69 (m, 4H), 10.85 (s, 1H).</p>	447	445	(3R, 4R)-form



180		¹ H-NMR (DMSO-D ₆) δ : 1.03 (d, 3H, J = 7.0 Hz), 2.46 (dq, 1H, J = 9.3, 7.0 Hz), 2.97 (ddd, 1H, J = 9.7, 9.3, 8.7 Hz), 3.15 (dd, 1H, J = 8.3, 8.7 Hz), 3.38 (dd, 1H, J = 8.3, 9.7 Hz), 6.59 (s, 1H), 7.17 (s, 1H), 7.27-7.32 (m, 2H), 7.40 (d, 1H, J = 8.1 Hz), 7.49-7.57 (m, 2H), 7.64 (s, 1H), 7.83 (dd, 1H, J = 8.8, 8.1 Hz), 10.79 (s, 1H).	497	495	(3R, 4R)-form
181		¹ H-NMR (DMSO-D ₆) δ : 0.78 (t, 3H, J = 7.4 Hz), 2.10 (tq, 2H, J = 16.4, 7.4 Hz), 2.38-2.40 (m, 2H), 3.33-3.35 (m, 1H), 3.39-3.47 (m, 1H), 3.50-3.52 (m, 1H), 7.01 (s, 1H), 7.04-7.24 (m, 3H), 7.27 (s, 1H), 7.41-7.54 (m, 4H), 7.64 (s, 1H), 10.93 (s, 1H).	443	441	R-form
182		¹ H-NMR (DMSO-D ₆) δ : 0.71 (t, 3H, J = 7.4 Hz), 2.02 (tq, 2H, J = 15.7, 7.4 Hz), 2.37-2.39 (m, 2H), 3.33-3.34 (m, 1H), 3.37-3.45 (m, 1H), 3.48-3.50 (m, 1H), 7.04 (s, 1H), 7.19 (s, 1H), 7.29-7.34 (m, 2H), 7.42-7.63 (m, 6H), 10.86 (s, 1H).	443	441	R-form



183		¹ H-NMR (DMSO-D ₆) δ : 2.37-2.39 (m, 2H), 3.33-3.36 (m, 1H), 3.40-3.44 (m, 1H), 3.47-3.51 (m, 1H), 6.84 (s, 1H), 7.41-7.43 (m, 1H), 7.51-7.53 (m, 1H), 7.60-7.63 (m, 3H), 7.69 (dd, 1H, J = 8.1, 8.1 Hz), 7.79-7.82 (m, 1H), 8.12-8.13 (m, 1H), 10.91 (s, 1H).	467	465	R-form
184		¹ H-NMR (DMSO-D ₆) δ : 1.08 (d, 3H, J = 7.2 Hz), 2.54 (dq, 1H, J = 9.3, 7.2 Hz), 3.03 (ddd, 1H, J = 8.5, 9.0, 9.3 Hz), 3.22 (dd, 1H, J = 8.5, 9.0 Hz), 3.44 (dd, 1H, J = 8.5, 9.0 Hz), 6.86 (s, 1H), 7.41-7.43 (m, 1H), 7.51-7.53 (m, 1H), 7.60-7.61 (m, 2H), 7.65-7.71 (m, 2H), 7.78-7.83 (m, 1H), 8.12-8.15 (m, 1H), 10.99 (s, 1H).	481	479	(3R, 4R)-form
185		¹ H-NMR (DMSO-D ₆) δ : 0.94 (t, 3H, J = 7.5 Hz), 1.75-1.99 (m, 4H), 2.01-2.35 (m, 4H), 2.28 (tq, 2H, J = 16.5, 7.5 Hz), 2.76-2.86 (m, 1H), 3.23-3.35 (m, 4H), 3.87-3.96 (m, 2H), 4.21-4.31 (m, 1H), 6.64 (s, 1H), 7.48 (s, 1H), 7.53 (s, 1H), 7.57-7.68 (m, 3H), 10.74 (s, 1H).	447	445	Optically active substance (The same configuration as Example 5)



186		¹ H-NMR (DMSO-D ₆) δ : 1.32 (s, 9H), 1.75-1.99 (m, 4H), 2.02-2.28 (m, 4H), 2.76-2.85 (m, 1H), 3.22-3.36 (m, 4H), 3.89-3.97 (m, 2H), 4.24-4.33 (m, 1H), 6.57 (s, 1H), 7.26 (ddd, 1H, $J = 7.7, 1.3, 1.4$ Hz), 7.39 (dd, 1H, $J = 1.5, 1.4$ Hz), 7.45 (dd, 1H, $J = 7.7, 7.7$ Hz), 7.48 (s, 1H), 7.51 (ddd, 1H, $J = 7.7, 1.5, 1.3$ Hz), 10.71 (s, 1H).	425	423	Optically active substance (The same configuration as Example 5)
187		¹ H-NMR (DMSO-D ₆) δ : 1.79-1.82 (m, 2H), 2.01-2.10 (m, 2H), 2.35 (d, 1H, $J = 8.6$ Hz), 2.36 (d, 1H, $J = 7.9$ Hz), 3.30-3.49 (m, 3H), 3.90-3.93 (m, 2H), 4.25-4.30 (m, 1H), 6.65 (s, 1H), 7.46 (s, 1H), 7.48-7.52 (m, 2H), 7.62 (s, 1H), 7.67 (dd, 1H, $J = 8.0, 8.0$ Hz), 10.77 (s, 1H).	439	437	R-form



188		¹ H-NMR (DMSO-D ₆) δ : 1.06 (d, 3H, J = 7.1 Hz), 1.78-1.84 (m, 2H), 2.01-2.10 (m, 2H), 2.53 (dq, 1H, J = 9.3, 7.1 Hz), 3.01 (ddd, 1H, J = 8.6, 9.3, 9.4 Hz), 3.18 (dd, 1H, J = 9.4, 8.5 Hz), 3.28-3.34 (m, 2H), 3.41 (dd, 1H, J = 8.6, 8.5 Hz), 3.90-3.93 (m, 2H), 4.25-4.30 (m, 1H), 6.68 (s, 1H), 7.46-7.53 (m, 3H), 7.65-7.69 (m, 2H), 10.85 (s, 1H).	453	451	(3R, 4R)-form
189		¹ H-NMR (DMSO-D ₆) δ : 1.79-2.27 (m, 8H), 2.79-2.83 (m, 1H), 3.23-3.36 (m, 4H), 3.90-3.93 (m, 2H), 4.26-4.29 (m, 1H), 6.64 (s, 1H), 7.45-7.52 (m, 4H), 7.66 (dd, 1H, J = 7.9, 7.9 Hz), 10.75 (s, 1H).	453	451	Optically active substance (The same configuration as Example 5)
190		¹ H-NMR (DMSO-D ₆) δ : 2.36 (d, 2H, J = 8.5 Hz), 2.82 (tq, 2H, J = 6.9, 11.3 Hz), 3.30 (dd, 1H, J = 8.5, 5.6 Hz), 3.39 (ddt, 1H, J = 8.5, 5.6, 8.5 Hz), 3.48 (dd, 1H, J = 8.5, 8.5 Hz), 4.25 (t, 2H, J = 6.9 Hz), 6.70 (s, 1H), 7.50 (ddd, 1H, J = 7.5, 1.3, 1.3 Hz), 7.51 (dd, 1H, J = 1.3, 1.3 Hz), 7.54 (ddd, 1H, J = 7.5, 1.3, 1.3 Hz), 7.63 (s, 1H), 7.67 (dd, 1H, J = 7.5, 7.5 Hz), 10.76 (s, 1H).	451	449	R-form



191		¹ H-NMR (DMSO-D ₆) δ : 1.06 (d, 3H, J = 7.3 Hz), 2.52 (dq, 1H, J = 9.7, 7.3 Hz), 2.83 (tq, 2H, J = 6.9, 11.3 Hz), 3.01 (ddd, 1H, J = 8.3, 8.9, 9.7 Hz), 3.20 (dd, 1H, J = 8.9, 8.9 Hz), 3.42 (dd, 1H, J = 8.3, 8.3 Hz), 4.25 (t, 2H, J = 6.9 Hz), 6.72 (s, 1H), 7.51 (ddd, 1H, J = 7.7, 1.3, 1.3 Hz), 7.52 (dd, 1H, J = 1.3, 1.3 Hz), 7.54 (ddd, 1H, J = 7.7, 1.3, 1.3 Hz), 7.66 (s, 1H), 7.67 (dd, 1H, J = 7.7, 7.7 Hz), 10.84 (s, 1H).	465	463	(3R, 4R)-form
192		¹ H-NMR (DMSO-D ₆) δ : 2.37 (d, 1H, J = 8.6 Hz), 2.38 (d, 1H, J = 7.7 Hz), 3.32 (dd, 1H, J = 6.5, 8.5 Hz), 3.41 (dddd, 1H, J = 6.5, 7.7, 8.5, 8.6 Hz), 3.49 (dd, 1H, J = 8.5, 8.5 Hz), 6.80 (s, 1H), 7.47-7.57 (m, 4H), 7.63 (s, 1H), 7.66-7.71 (m, 2H), 8.01 (s, 1H), 10.82 (s, 1H).	483	481	R-form





193		<p>¹H-NMR (DMSO-D₆) δ: 1.09 (d, 3H, J = 7.2 Hz), 2.56 (dq, 1H, J = 9.3, 7.2 Hz), 3.03 (ddd, 1H, J = 9.0, 9.3, 9.3 Hz), 3.23 (dd, 1H, J = 8.4, 9.0 Hz), 3.45 (dd, 1H, J = 8.4, 9.3 Hz), 7.02 (s, 1H), 7.11 (s, 1H), 7.31-7.35 (m, 2H), 7.47-7.56 (m, 3H), 7.60-7.65 (m, 2H), 7.68 (s, 1H), 10.96 (s, 1H).</p>	479	477	(3R, 4R)-form
194		<p>¹H-NMR (DMSO-D₆) δ: 1.06 (s, 9H), 1.09 (d, 3H, J = 7.2 Hz), 2.56 (dq, 1H, J = 9.3, 7.2 Hz), 3.03 (ddd, 1H, J = 9.0, 9.2, 9.3 Hz), 3.23 (dd, 1H, J = 8.5, 9.2 Hz), 3.45 (dd, 1H, J = 8.5, 9.0 Hz), 7.01 (s, 1H), 7.01-7.02 (m, 1H), 7.19-7.21 (m, 1H), 7.27-7.33 (m, 2H), 7.45-7.56 (m, 3H), 7.61-7.63 (m, 1H), 7.67 (s, 1H), 10.90 (s, 1H).</p>	451	449	(3R, 4R)-form



195		¹ H-NMR (DMSO-D ₆) δ : 1.91 (s, 3H), 2.39 (d, 1H, J = 8.6 Hz), 2.39 (d, 1H, J = 8.3 Hz), 3.34 (dd, 1H, J = 8.5, 6.5 Hz), 3.42 (dddd, 1H, J = 6.5, 8.3, 8.5, 8.6 Hz), 3.50 (dd, 1H, J = 8.5, 8.5 Hz), 7.00 (s, 1H), 7.06 (s, 1H), 7.14 (td, 1H, J = 8.5, 2.9 Hz), 7.24 (dd, 1H, J = 9.7, 2.9 Hz), 7.30-7.33 (m, 2H), 7.37 (dd, 1H, J = 8.7, 5.5 Hz), 7.49 (dd, 1H, J = 8.7, 8.0 Hz), 7.64 (s, 1H), 10.83 (s, 1H).	463	461	R-form
196		¹ H-NMR (DMSO-D ₆) δ : 1.08 (d, 3H, J = 7.2 Hz), 1.91 (s, 3H), 2.55 (dq, 1H, J = 9.2, 7.2 Hz), 3.03 (ddd, 1H, J = 8.5, 9.1, 9.2 Hz), 3.23 (dd, 1H, J = 8.7, 9.1 Hz), 3.45 (dd, 1H, J = 8.5, 8.7 Hz), 7.01 (s, 1H), 7.09 (s, 1H), 7.15 (dt, 1H, J = 2.8, 8.0 Hz), 7.24 (dd, 1H, J = 9.7, 2.8 Hz), 7.31-7.34 (m, 2H), 7.38 (dd, 1H, J = 8.0, 9.7 Hz), 7.49 (t, 1H, J = 8.0 Hz), 7.68 (s, 1H), 10.92 (s, 1H).	477	475	(3R, 4R)-form




197		¹ H-NMR (DMSO-D ₆) δ : 1.93-2.13 (m, 8H), 2.34-2.36 (m, 2H), 3.27- 3.49 (m, 3H), 4.28-4.33 (m, 1H), 6.65 (s, 1H), 7.45 (s, 1H), 7.48-7.52 (m, 2H), 7.61 (s, 1H), 7.67 (dd, 1H, J = 7.9, 7.9 Hz), 10.77 (s, 1H).	473	471	R-form
198		¹ H-NMR (DMSO-D ₆) δ : 1.05 (d, 3H, J = 7.2 Hz), 1.92-2.13 (m, 8H), 2.53 (dq, 1H, J = 9.3, 7.2 Hz), 3.00 (ddd, 1H, J = 8.6, 9.0, 9.3 Hz), 3.18 (dd, 1H, J = 8.6, 9.0 Hz), 3.41 (dd, 1H, J = 8.4, 8.6 Hz), 4.28-4.33 (m, 1H), 6.67 (s, 1H), 7.46 (s, 1H), 7.49-7.52 (m, 2H), 7.65 (s, 1H), 7.67-7.69 (m, 1H), 10.86 (s, 1H).	487	485	(3R, 4R)- form




199		<p>¹H-NMR (DMSO-D₆) δ: 0.96-1.10 (m, 2H), 1.27-1.35 (m, 2H), 1.90-2.03 (m, 1H), 2.36 (d, 2H, $J = 8.5$ Hz), 3.11-3.20 (m, 2H), 3.30 (dd, 1H, $J = 8.3, 5.6$ Hz), 3.38 (ddt, 1H, $J = 8.5, 5.6, 8.3$ Hz), 3.47 (dd, 1H, $J = 8.3, 8.3$ Hz), 3.69-3.77 (m, 2H), 3.92 (d, 2H, $J = 7.3$ Hz), 6.67 (s, 1H), 7.48 (dd, 1H, $J = 1.2, 1.2$ Hz), 7.48 (ddd, 1H, $J = 8.9, 1.2, 1.2$ Hz), 7.52 (ddd, 1H, $J = 7.7, 1.2, 1.2$ Hz), 7.62 (s, 1H), 7.65 (dd, 1H, $J = 8.9, 7.7$ Hz), 10.71 (s, 1H).</p>	453	451	R-form
200		<p>¹H-NMR (DMSO-D₆) δ: 0.98-1.10 (m, 2H), 1.06 (d, 3H, $J = 6.9$ Hz), 1.27-1.36 (m, 2H), 1.90-2.03 (m, 1H), 2.51 (dq, 1H, $J = 9.7, 6.9$ Hz), 3.00 (ddd, 1H, $J = 8.5, 9.1, 9.7$ Hz), 3.11-3.20 (m, 2H), 3.19 (dd, 1H, $J = 9.1, 9.1$ Hz), 3.41 (dd, 1H, $J = 8.5, 8.5$ Hz), 3.70-3.77 (m, 2H), 3.92 (d, 2H, $J = 6.9$ Hz), 6.70 (s, 1H), 7.47-7.51 (m, 2H), 7.52 (ddd, 1H, $J = 7.7, 1.3, 1.3$ Hz), 7.66 (dd, 1H, $J = 8.9, 7.7$ Hz), 7.66 (s, 1H), 10.80 (s, 1H).</p>	467	465	(3R, 4R)-form



201		<p>¹H-NMR (DMSO-D₆) δ: 0.98-1.11 (m, 2H), 1.20-1.35 (m, 3H), 1.61 (dt, 2H, $J = 7.1, 6.7$ Hz), 2.36 (d, 2H, $J = 8.5$ Hz), 3.07-3.18 (m, 2H), 3.30 (dd, 1H, $J = 8.7, 5.6$ Hz), 3.38 (ddt, 1H, $J = 8.5, 5.6, 8.7$ Hz), 3.47 (dd, 1H, $J = 8.7, 8.7$ Hz), 3.67-3.75 (m, 2H), 4.06 (t, 2H, $J = 7.1$ Hz), 6.67 (s, 1H), 7.48 (ddd, 1H, $J = 7.3, 1.2, 1.2$ Hz), 7.49 (dd, 1H, $J = 1.2, 1.2$ Hz), 7.53 (ddd, 1H, $J = 7.9, 1.2, 1.2$ Hz), 7.62 (s, 1H), 7.66 (dd, 1H, $J = 7.9, 7.3$ Hz), 10.71 (s, 1H).</p>	467	465	R-form
202		<p>¹H-NMR (DMSO-D₆) δ: 1.08 (d, 3H, $J = 7.0$ Hz), 2.54-2.57 (m, 1H), 3.04 (ddd, 1H, $J = 18.1, 8.4, 4.2$ Hz), 3.23 (dd, 1H, $J = 9.0, 4.5$ Hz), 3.45 (dd, 1H, $J = 8.5, 4.2$ Hz), 7.07 (s, 1H), 7.26-7.34 (m, 4H), 7.53 (s, 1H), 7.55 (d, 1H, $J = 7.9$ Hz), 7.61 (dd, 1H, $J = 7.7, 3.8$ Hz), 7.68 (s, 1H), 7.73 (d, 1H, $J = 7.7$ Hz), 10.99 (s, 1H).</p>	447	445	(3R, 4R)-form



203		¹ H-NMR (DMSO-D ₆) δ: 1.09 (d, 3H, J = 7.2 Hz), 2.54-2.60 (m, 1H), 3.04 (ddd, 1H, J = 18.1, 8.6, 4.3 Hz), 3.24 (dd, 1H, J = 8.8, 4.4 Hz), 3.46 (dd, 1H, J = 8.7, 4.4 Hz), 7.15 (s, 1H), 7.41 (s, 1H), 7.49-7.68 (m, 8H), 10.97 (s, 1H).	463	461	(3R, 4R)-form
204		¹ H-NMR (DMSO-D ₆) δ: 2.39 (d, 2H, J = 8.4 Hz), 3.34-3.35 (m, 1H), 3.43-3.48 (m, 2H), 7.04 (s, 1H), 7.26-7.34 (m, 4H), 7.52 (s, 1H), 7.55 (d, 1H, J = 7.9 Hz), 7.60-7.64 (m, 2H), 7.73 (d, 1H, J = 7.7 Hz), 10.91 (s, 1H).	433	431	R-form
205		¹ H-NMR (DMSO-D ₆) δ: 2.39-2.40 (m, 2H), 3.35-3.53 (m, 3H), 7.13 (s, 1H), 7.40 (s, 1H), 7.48-7.69 (m, 8H), 10.89 (s, 1H).	449	447	R-form
206		¹ H-NMR (DMSO-D ₆) δ: 2.38-2.40 (m, 2H), 3.33-3.53 (m, 3H), 4.77 (q, 2H, J = 8.8 Hz), 6.84 (d, 1H, J = 7.9 Hz), 6.98 (s, 1H), 7.02-7.15 (m, 4H), 7.19-7.21 (m, 1H), 7.32-7.34 (m, 1H), 7.42-7.44 (m, 1H), 7.64 (s, 1H), 10.91 (s, 1H).	463	461	R-form



207		¹ H-NMR (DMSO-D ₆) δ : 1.09 (d, 3H, J = 7.0 Hz), 2.57 (dq, 1H, J = 9.3, 7.0 Hz), 3.04 (ddd, 1H, J = 9.3, 9.3, 9.3 Hz), 3.23 (dd, 1H, J = 8.5, 9.3 Hz), 3.45 (dd, 1H, J = 8.5, 9.3 Hz), 7.07 (s, 1H), 7.25-7.27 (m, 2H), 7.38-7.45 (m, 3H), 7.50 (s, 1H), 7.55-7.62 (m, 2H), 7.67-7.73 (m, 2H), 11.00 (s, 1H).	429	427	(3R, 4R)-form
208		¹ H-NMR (DMSO-D ₆) δ : 0.69 (t, 3H, J = 7.4 Hz), 1.09 (d, 3H, J = 7.2 Hz), 2.02 (tq, 2H, J = 16.2, 7.4 Hz), 2.56 (dq, 1H, J = 9.3, 7.2 Hz), 3.03 (ddd, 1H, J = 8.5, 9.1, 9.3 Hz), 3.24 (dd, 1H, J = 8.8, 9.1 Hz), 3.45 (dd, 1H, J = 8.5, 8.8 Hz), 7.09 (s, 1H), 7.17 (s, 1H), 7.41-7.54 (m, 5H), 7.58-7.63 (m, 2H), 7.68 (s, 1H), 10.94 (s, 1H).	473	471	(3R, 4R)-form



209		<p>¹H-NMR (DMSO-D₆) δ: 1.09 (d, 3H, J = 7.2 Hz), 2.56 (dq, 1H, J = 9.3, 7.2 Hz), 3.05 (ddd, 1H, J = 8.6, 9.2, 9.3 Hz), 3.23 (dd, 1H, J = 9.0, 9.2 Hz), 3.45 (dd, 1H, J = 8.6, 9.0 Hz), 7.04-7.06 (m, 1H), 7.08 (s, 1H), 7.18-7.27 (m, 2H), 7.42-7.47 (m, 1H), 7.56-7.65 (m, 3H), 7.68 (s, 1H), 7.75 (d, 1H, J = 7.7 Hz), 11.03 (s, 1H).</p>	447	445	(3R, 4R)-form
210		<p>¹H-NMR (DMSO-D₆) δ: 1.08 (d, 3H, J = 7.2 Hz), 2.55 (dq, 1H, J = 9.3, 7.2 Hz), 3.04 (ddd, 1H, J = 8.6, 9.2, 9.3 Hz), 3.23 (dd, 1H, J = 8.6, 8.7 Hz), 3.44 (dd, 1H, J = 8.7, 9.2 Hz), 4.73 (q, 2H, J = 8.8 Hz), 6.80 (d, 1H, J = 8.1 Hz), 6.99 (s, 1H), 7.01-7.07 (m, 2H), 7.22-7.31 (m, 3H), 7.33-7.43 (m, 3H), 7.67 (s, 1H), 10.95 (s, 1H).</p>	459	457	(3R, 4R)-form
211		<p>¹H-NMR (DMSO-D₆) δ: 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.75 (q, 2H, J = 8.8 Hz), 6.80 (d, 1H, J = 7.9 Hz), 6.97 (s, 1H), 7.01-7.06 (m, 2H), 7.23-7.32 (m, 5H), 7.64 (s, 1H), 10.86 (s, 1H).</p>	463	461	R-form




212		¹ H-NMR (DMSO-D ₆) δ : 1.08 (d, 3H, J = 7.2 Hz), 2.55 (dq, 1H, J = 9.3, 7.2 Hz), 3.03 (ddd, 1H, J = 8.6, 9.3, 9.3 Hz), 3.22 (dd, 1H, J = 8.8, 9.3 Hz), 3.44 (dd, 1H, J = 8.6, 8.8 Hz), 4.75 (q, 2H, J = 8.8 Hz), 6.79-6.81 (m, 1H), 6.99 (s, 1H), 7.03-7.06 (m, 2H), 7.24-7.32 (m, 5H), 7.67 (s, 1H), 10.95 (s, 1H).	477	475	(3R, 4R)-form
213		¹ H-NMR (DMSO-D ₆) δ : 2.39 (d, 2H, J = 8.1 Hz), 3.33-3.35 (m, 1H), 3.39-3.53 (m, 2H), 7.04 (s, 1H), 7.06 (s, 1H), 7.19-7.26 (m, 2H), 7.41-7.47 (m, 1H), 7.56-7.66 (m, 4H), 7.75 (d, 1H, J = 7.7 Hz), 10.95 (s, 1H).	433	431	R-form
214		¹ H-NMR (DMSO-D ₆) δ : 2.38-2.41 (m, 2H), 3.34-3.52 (m, 3H), 4.70 (q, 2H, J = 8.8 Hz), 6.76 (dd, 1H, J = 7.9, 2.0 Hz), 6.94 (dd, 1H, J = 2.0, 2.0 Hz), 7.00 (dd, 1H, J = 8.1, 2.1 Hz), 7.05 (s, 1H), 7.24 (dd, 1H, J = 7.9, 8.1 Hz), 7.46-7.54 (m, 2H), 7.57-7.61 (m, 2H), 7.64 (s, 1H), 10.83 (s, 1H).	479	477	R-form




215		<p>¹H-NMR (DMSO-D₆) δ: 1.09 (d, 3H, J = 7.2 Hz), 2.57 (dq, 1H, J = 9.3, 7.2 Hz), 3.03 (ddd, 1H, J = 8.5, 8.8, 9.3 Hz), 3.23 (dd, 1H, J = 8.6, 8.8 Hz), 3.45 (dd, 1H, J = 8.5, 8.6 Hz), 4.71 (q, 2H, J = 8.8 Hz), 6.75-6.77 (m, 1H), 6.95-7.01 (m, 2H), 7.07 (s, 1H), 7.24 (dd, 1H, J = 8.3, 8.3 Hz), 7.46-7.54 (m, 2H), 7.57-7.62 (m, 2H), 7.68 (s, 1H), 10.92 (s, 1H).</p>	493	491	(3R, 4R)-form
216		<p>¹H-NMR (DMSO-D₆) δ: 2.31 (s, 3H), 2.38 (d, 2H, J = 8.6 Hz), 3.33 (dd, 1H, J = 8.4, 8.5 Hz), 3.41 (ddt, 1H, J = 8.4, 8.5, 8.6 Hz), 3.50 (dd, 1H, J = 8.4, 8.4 Hz), 4.73 (q, 2H, J = 8.9 Hz), 6.79 (d, 1H, J = 7.7 Hz), 6.95 (s, 1H), 7.00 (d, 1H, J = 2.4 Hz), 7.03 (dd, 1H, J = 8.0, 2.4 Hz), 7.12 (d, 2H, J = 8.4 Hz), 7.20 (d, 2H, J = 8.4 Hz), 7.28 (dd, 1H, J = 7.7, 8.0 Hz), 7.63 (s, 1H), 10.85 (s, 1H).</p>	459	457	R-form



217		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 1.08 (d, 3H, J = 7.7 Hz), 2.31 (s, 3H), 2.55 (dq, 1H, J = 9.3, 7.7 Hz), 3.03 (ddd, 1H, J = 9.3, 9.4, 9.5 Hz), 3.22 (dd, 1H, J = 8.7, 9.4 Hz), 3.44 (dd, 1H, J = 8.7, 9.5 Hz), 4.74 (q, 2H, J = 9.0 Hz), 6.78-6.80 (m, 1H), 6.97 (s, 1H), 7.02-7.05 (m, 2H), 7.12 (d, 2H, J = 8.4 Hz), 7.21 (d, 2H, J = 8.4 Hz), 7.28 (dd, 1H, J = 7.9, 7.9 Hz), 7.67 (s, 1H), 10.94 (s, 1H).</p>	473	471	(3R, 4R)-form
218		<p>¹H-NMR (DMSO-D₆)</p> <p>δ: 1.08 (d, 6H, J = 6.0 Hz), 1.57-1.65 (m, 2H), 1.79-1.82 (m, 2H), 2.38-2.43 (m, 2H), 3.32-3.56 (m, 3H), 4.26-4.29 (m, 2H), 4.67-4.69 (m, 1H), 6.82 (s, 1H), 7.29 (d, 1H, J = 7.7 Hz), 7.55 (dd, 1H, J = 7.7, 7.7 Hz), 7.67 (s, 1H), 7.69 (s, 1H), 7.82 (d, 1H, J = 7.7 Hz), 10.04 (s, 1H).</p>	467	465	<p>Lactam ring: R-form</p> <p>Tetrahydro pyrane ring: (2R, 4S, 6S)-form</p>



219		<p>¹H-NMR (DMSO-D₆) δ: 1.15 (d, 6H, J = 6.0 Hz), 1.63-1.72 (m, 2H), 1.82-1.85 (m, 2H), 2.39-2.44 (m, 2H), 3.35-3.37 (m, 1H), 3.42-3.49 (m, 1H), 3.53-3.57 (m, 3H), 4.36-4.41 (m, 1H), 6.75 (s, 1H), 7.27 (d, 1H, J = 8.1 Hz), 7.51 (dd, 1H, J = 7.7, 8.1 Hz), 7.67 (s, 1H), 7.68 (s, 1H), 7.79 (d, 1H, J = 7.7 Hz), 10.07 (s, 1H).</p>	467	465	<p>Lactam ring: R-form</p> <p>Tetrahydro pyrane ring: (2R, 4R, 6S)-form</p>
220		<p>¹H-NMR (DMSO-D₆) δ: 1.07 (d, 6H, J = 6.0 Hz), 1.13 (d, 3H, J = 6.9 Hz), 1.58-1.66 (m, 2H), 1.80-1.83 (m, 2H), 2.53 (dq, 1H, J = 9.3, 6.9 Hz), 3.05 (ddd, 1H, J = 8.5, 9.1, 9.3 Hz), 3.26 (dd, 1H, J = 8.5, 8.9 Hz), 3.48 (dd, 1H, J = 8.9, 9.1 Hz), 4.24-4.30 (m, 2H), 4.66-4.69 (m, 1H), 6.84 (s, 1H), 7.29 (d, 1H, J = 8.1 Hz), 7.55 (dd, 1H, J = 8.1, 8.1 Hz), 7.70 (s, 1H), 7.71 (s, 1H), 7.83 (d, 1H, J = 8.1 Hz), 10.09 (s, 1H).</p>	481	479	<p>Lactam ring: (3R, 4R)-form</p> <p>Tetrahydro pyrane ring: (2R, 4S, 6S)-form</p>



221		¹ H-NMR (DMSO-D ₆) δ : 1.14 (d, 3H, J = 7.3 Hz), 1.17 (d, 6H, J = 6.0 Hz), 1.64-1.74 (m, 2H), 1.82-1.88 (m, 2H), 2.55 (dq, 1H, J = 9.3, 7.3 Hz), 3.06 (ddd, 1H, J = 8.5, 8.9, 9.3 Hz), 3.27 (dd, 1H, J = 8.5, 9.3 Hz), 3.50 (dd, 1H, J = 8.9, 9.3 Hz), 3.54-3.60 (m, 2H), 4.35-4.40 (m, 1H), 6.79 (s, 1H), 7.29 (d, 1H, J = 7.7 Hz), 7.53 (dd, 1H, J = 7.7, 7.7 Hz), 7.71 (s, 1H), 7.72 (s, 1H), 7.81 (d, 1H, J = 7.7 Hz), 10.14 (s, 1H).	481	479	Lactam ring: (3R, 4R)-form Tetrahydropyrane ring: (2R, 4R, 6S)-form
222		¹ H-NMR (DMSO-D ₆) δ : 1.08 (d, 3H, J = 7.2 Hz), 2.55-2.57 (m, 1H), 3.04 (ddd, 1H, J = 8.8, 4.4, 2.2 Hz), 3.22 (dd, 1H, J = 9.0, 4.5 Hz), 3.45 (dd, 1H, J = 8.8, 4.4 Hz), 4.77 (q, 2H, J = 8.8 Hz), 6.84 (d, 1H, J = 7.7 Hz), 7.00 (s, 1H), 7.16-7.32 (m, 7H), 7.67 (s, 1H), 10.99 (s, 1H).	477	475	(3R, 4R)-form

223		¹ H-NMR (DMSO-D ₆) δ: 1.07 (3H, d, J = 7.2 Hz), 1.90 (3H, s), 2.48-2.60 (1H, m), 3.02 (1H, ddd, J = 9.3, 9.0, 8.8 Hz), 3.22 (1H, dd, J = 9.0, 8.8 Hz), 3.44 (1H, dd, J = 9.0, 8.8 Hz), 6.97 (1H, s), 7.08 (1H, s), 7.24-7.42 (6H, m), 7.45 (1H, t, J = 8.0 Hz), 7.66 (1H, s), 10.90 (1H, s)	459	457	(3R, 4R)- form
224		¹ H-NMR (DMSO-D ₆) δ: 1.90 (3H, s), 2.34-2.44 (2H, m), 3.30-3.55 (3H, m), 7.10 (1H, s), 7.24- 7.42 (5H, m), 7.49-7.58 (2H, m), 7.59-7.68 (2H, m), 10.82 (1H, s)	429	427	R-form
225		¹ H-NMR (DMSO-D ₆) δ: 1.08 (3H, d, J = 7.2 Hz), 1.91 (3H, s), 2.48-2.60 (1H, m), 3.03 (1H, ddd, J = 8.8, 8.8, 8.8 Hz), 3.23 (1H, dd, J = 9.0, 8.6 Hz), 3.44 (1H, dd, J = 9.0, 8.6 Hz), 7.12 (1H, s), 7.26- 7.42 (5H, m), 7.50-7.58 (2H, m), 7.60-7.69 (2H, m), 10.91 (1H, s)	443	441	(3R, 4R)- form

226		¹ H-NMR (DMSO-D ₆) δ: 1.90 (s, 3H), 2.37-2.39 (br m, 2H), 3.32-3.51 (m, 3H), 4.65 (q, 2H, J = 8.9 Hz), 6.74 (d, 1H, J = 7.9 Hz), 6.88 (s, 1H), 6.96 (dd, 1H, J = 8.1, 2.6 Hz), 7.02 (s, 1H), 7.23-7.34 (m, 5H), 7.63 (s, 1H), 10.76 (s, 1H)	459	457	R-form
227		¹ H-NMR (DMSO-D ₆) δ: 1.90 (s, 3H), 2.37-2.39 (m, 2H), 3.32-3.51 (m, 3H), 4.65 (q, 2H, J = 8.9 Hz), 6.74 (d, 1H, J = 7.9 Hz), 6.88 (s, 1H), 6.96 (dd, 1H, J = 8.1, 2.6 Hz), 7.02 (s, 1H), 7.23-7.34 (m, 5H), 7.63 (s, 1H), 10.76 (s, 1H)	473	471	(3R, 4R)-form
228		¹ H-NMR (DMSO-D ₆) δ: 1.90 (s, 3H), 2.37-2.39 (m, 2H), 3.32-3.51 (m, 3H), 4.65 (q, 2H, J = 8.9 Hz), 6.74 (d, 1H, J = 7.9 Hz), 6.88 (s, 1H), 6.96 (dd, 1H, J = 8.1, 2.6 Hz), 7.02 (s, 1H), 7.23-7.34 (m, 5H), 7.63 (s, 1H), 10.76 (s, 1H)	433	431	R-form




229		¹ H-NMR (DMSO-D ₆) δ: 1.09 (d, 3H, J = 7.3 Hz), 2.52-2.59 (m, 1H), 3.04 (ddd, 1H, J = 8.5, 8.5, 8.5 Hz), 3.24 (dd, 1H, J = 8.9, 8.9 Hz), 3.45 (dd, 1H, J = 8.5, 8.5 Hz), 7.14 (s, 1H), 7.31-7.39 (m, 2H), 7.47 (s, 1H), 7.49- 7.64 (m, 4H), 7.65-7.75 (m, 2H), 11.00 (s, 1H)	447	445	(3R, 4R)- form
230		¹ H-NMR (DMSO-D ₆) δ: 2.31 (s, 3H), 2.37 (d, 2H, J = 8.4 Hz), 3.32 (dd, 5H, J = 8.5, 6.0 Hz), 3.36-3.45 (m, 1H), 3.49 (dd, 1H, J = 8.5, 8.5 Hz), 7.00 (s, 1H), 7.12 (dd, 2H, J = 6.5, 1.9 Hz), 7.21 (dd, 2H, J = 6.5, 2.1 Hz), 7.50 (d, 1H, J = 7.7 Hz), 7.51 (s, 1H), 7.58 (t, 1H, J = 7.7 Hz), 7.62 (s, 1H), 7.69 (d, 1H, J = 7.7 Hz), 10.88 (s, 1H).	429	427	R-form

231		<p>¹H-NMR (DMSO-D₆) δ: 1.07 (d, 3H, J = 7.2 Hz), 2.53 (dq, 1H, J = 9.5, 7.2 Hz), 3.02 (ddd, 1H, J = 9.5, 9.0, 9.0 Hz), 3.21 (dd, 1H, J = 9.0, 9.0 Hz), 3.43 (dd, 1H, J = 9.0, 9.0 Hz), 7.03 (s, 1H), 7.13 (dd, 2H, J = 6.4, 2.0 Hz), 7.21 (dd, 2H, J = 6.4, 1.9 Hz), 7.50 (d, 1H, J = 7.7 Hz), 7.52 (s, 1H), 7.58 (t, 1H, J = 7.7 Hz), 7.66 (s, 1H), 7.70 (d, 1H, J = 7.7 Hz), 10.97 (s, 1H).</p>	443	441	(3R, 4R)-form
232		<p>¹H-NMR (DMSO-D₆) δ: 0.97-1.11 (m, 2H), 1.06 (d, 3H, J = 6.9 Hz), 1.20-1.35 (m, 3H), 1.61 (dt, 2H, J = 7.3, 6.5 Hz), 2.51 (dq, 1H, J = 9.7, 6.9 Hz), 3.00 (ddd, 1H, J = 8.5, 9.1, 9.7 Hz), 3.08-3.16 (m, 2H), 3.19 (dd, 1H, J = 9.1, 9.1 Hz), 3.41 (dd, 1H, J = 8.5, 8.5 Hz), 3.67-3.75 (m, 2H), 4.06 (t, 2H, J = 7.3 Hz), 6.69 (s, 1H), 7.48 (ddd, 1H, J = 7.1, 1.4, 1.4 Hz), 7.49 (dd, 1H, J = 1.4, 1.4 Hz), 7.54 (ddd, 1H, J = 7.9, 1.4, 1.4 Hz), 7.66 (dd, 1H, J = 7.9, 7.1 Hz), 7.66 (s, 1H), 10.80 (s, 1H).</p>	481	479	(3R, 4R)-form




233		¹ H-NMR (DMSO-D ₆) δ : 2.32 (s, 3H), 2.39 (d, 2H, $J = 8.5$ Hz), 3.33 (dd, 1H, $J = 8.7, 5.6$ Hz), 3.42 (ddt, 1H, $J = 8.5, 5.6, 8.7$ Hz), 3.50 (dd, 1H, $J = 8.7, 8.7$ Hz), 6.98 (s, 1H), 7.11 (dd, 1H, $J = 1.2, 1.2$ Hz), 7.13 (d, 2H, $J = 8.1$ Hz), 7.22 (d, 2H, $J = 8.1$ Hz), 7.32 (ddd, 1H, $J = 8.1, 1.2, 1.2$ Hz), 7.35 (ddd, 1H, $J = 8.1, 1.2, 1.2$ Hz), 7.51 (dd, 1H, $J = 8.1, 8.1$ Hz), 7.64 (s, 1H), 10.89 (s, 1H).	445	443	R-form
234		¹ H-NMR (DMSO-D ₆) δ : 1.08 (d, 3H, $J = 7.3$ Hz), 2.32 (s, 3H), 2.54 (dq, 1H, $J = 9.7, 7.3$ Hz), 3.03 (ddd, 1H, $J = 8.7, 9.1, 9.7$ Hz), 3.23 (dd, 1H, $J = 9.1, 9.1$ Hz), 3.44 (dd, 1H, $J = 8.7, 8.7$ Hz), 7.01 (s, 1H), 7.11 (dd, 1H, $J = 1.1, 1.3$ Hz), 7.14 (d, 2H, $J = 8.1$ Hz), 7.22 (d, 2H, $J = 8.1$ Hz), 7.32 (ddd, 1H, $J = 8.1, 1.1, 1.3$ Hz), 7.35 (ddd, 1H, $J = 8.1, 1.1, 1.3$ Hz), 7.51 (dd, 1H, $J = 8.1, 8.1$ Hz), 7.68 (s, 1H), 10.98 (s, 1H).	459	457	(3R, 4R)-form




[0372]



[Table 2]





Example No.	Molecular Structure	NMR (400MHz)	MS(M +H)	MS(M -H)	Comments
235		¹ H-NMR (DMSO-D ₆) δ: 1.18 (d, 3H, J = 6.5 Hz), 2.36-2.38 (m, 2H), 3.32-3.33 (m, 1H), 3.37-3.41 (m, 1H), 3.46-3.49 (m, 1H), 4.10 (qq, 1H, J = 6.5, 9.4 Hz), 4.57 (d, 1H, J = 12.7 Hz), 4.61 (d, 1H, J = 12.7 Hz), 6.96-3.98 (m, 2H), 7.09 (s, 1H), 7.15-7.18 (m, 1H), 7.25-7.27 (m, 2H), 7.35-7.44 (m, 3H), 7.63 (s, 1H), 10.88 (s, 1H).	491	489	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form
236		¹ H-NMR (DMSO-D ₆) δ: 1.20 (d, 3H, J = 6.5 Hz), 2.38-2.40 (m, 2H), 3.33-3.35 (m, 1H), 3.40-3.44 (m, 1H), 3.48-3.51 (m, 1H), 4.10-4.15 (m, 1H), 4.63 (s, 2H), 6.99 (s, 1H), 7.06 (s, 1H), 7.08 (d, 1H, J = 9.3 Hz), 7.19 (d, 1H, J = 9.3 Hz), 7.27 (dd, 2H, J = 6.6, 2.0 Hz), 7.48 (dd, 2H, J = 6.6, 2.0 Hz), 7.63 (s, 1H), 10.91 (s, 1H).	525	523	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form
237		¹ H-NMR (CDCl ₃) δ: 1.29 (d, 3H, J = 6.4 Hz), 2.59 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.8 Hz), 3.29-3.40 (m, 1H), 3.60 (dd, 1H, J = 9.2 Hz), 3.67-3.80 (m, 2H), 4.52 (d, 1H, J = 12.0 Hz), 4.63 (d, 1H, J = 12.0 Hz), 6.08 (s, 1H), 6.85 (dt, 1H, J = 9.6, 2.0 Hz), 6.98-7.04 (m, 2H), 7.09-7.17 (m, 2H), 7.22 (t, 1H, J = 8.0 Hz), 7.34-7.22 (m, 2H), 8.69 (s, 1H).	509	507	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form




238		¹ H-NMR (DMSO-D ₆) δ: 1.20 (d, 3H, J = 6.5 Hz), 2.38-2.39 (m, 2H), 3.33-3.34 (m, 1H), 3.39-3.43 (m, 1H), 3.49-3.51 (m, 1H), 4.09-4.11 (m, 1H), 4.59 (d, 1H, J = 12.1 Hz), 4.63 (d, 1H, J = 12.1 Hz), 6.92 (s, 1H), 7.18-7.39 (m, 8H), 7.63 (s, 1H), 10.85 (s, 1H).	491	489	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
239		¹ H-NMR (CDCl ₃) δ: 1.31 (d, 3H, J = 6.4 Hz), 2.61 (dd, 1H, J = 16.8, 9.6 Hz), 2.78 (dd, 1H, J = 16.8, 8.4 Hz), 3.31-3.42 (m, 1H), 3.61 (t, 1H, J = 8.8 Hz), 3.67-3.83 (m, 2H), 4.55 (d, 1H, J = 12.0 Hz), 4.65 (d, 1H, J = 12.0 Hz), 6.04 (s, 1H), 6.85 (d, 1H, J = 9.2 Hz), 6.98-7.10 (m, 5H), 7.19-7.25 (m, 2H), 8.48 (s, 1H).	509	507	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
240		¹ H-NMR (DMSO-D ₆) δ: 1.20 (d, 3H, J = 6.0 Hz), 3.35 (dd, 1H, J = 8.6, 6.2 Hz), 3.58 (dd, 1H, J = 9.3, 8.6 Hz), 4.12 (tq, 1H, J = 9.7, 6.0 Hz), 4.31 (dd, 1H, J = 9.3, 6.2 Hz), 4.60 (d, 1H, J = 12.5 Hz), 4.63 (d, 1H, J = 12.5 Hz), 6.33 (s, 1H), 6.59 (s, 1H), 6.98 (s, 1H), 7.02-7.07 (m, 2H), 7.18 (d, 1H, J = 9.6 Hz), 7.23-7.35 (m, 4H), 10.72 (s, 1H).	510	509	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Alpha position of carbonyl group (amide): R-form




241		¹ H-NMR (DMSO-D ₆) δ: 1.18 (d, 3H, J = 6.5 Hz), 2.36-2.38 (m, 2H), 3.32-3.33 (m, 1H), 3.37-3.41 (m, 1H), 3.46-3.49 (m, 1H), 4.10 (qq, 1H, J = 6.5, 9.4 Hz), 4.57 (d, 1H, J = 12.7 Hz), 4.61 (d, 1H, J = 12.7 Hz), 6.98 (s, 1H), 7.19-7.20 (m, 2H), 7.25-7.27 (m, 2H), 7.35-7.45 (m, 4H), 7.63 (s, 1H), 10.88 (s, 1H).	507	505	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
242		¹ H-NMR (DMSO-D ₆) δ: 1.18 (d, 3H, J = 6.5 Hz), 2.36-2.38 (m, 2H), 3.32-3.33 (m, 1H), 3.37-3.41 (m, 1H), 3.46-3.49 (m, 1H), 4.10 (qq, 1H, J = 6.5, 9.4 Hz), 4.57 (d, 1H, J = 12.7 Hz), 4.61 (d, 1H, J = 12.7 Hz), 6.97 (s, 1H), 7.13 (s, 1H), 7.23-7.32 (m, 5H), 7.37 (s, 1H), 7.61 (s, 1H), 10.86 (s, 1H).	525	523	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
243		¹ H-NMR (DMSO-D ₆) δ: 1.32 (s, 6H), 2.37-2.42 (m, 2H), 3.33-3.53 (m, 3H), 4.53 (s, 2H), 6.98 (s, 1H), 7.16 (s, 1H), 7.19 (s, 1H), 7.23-7.28 (m, 2H), 7.33-7.45 (m, 4H), 7.63 (s, 1H), 10.89 (s, 1H).	521	519	S-form





244		¹ H-NMR (CDCl ₃) δ: 1.29 (d, 3H, J = 6.8 Hz), 2.61 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.8 Hz), 3.30-3.40 (m, 1H), 3.61 (t, 1H, J = 8.8 Hz), 3.68-3.82 (m, 2H), 4.57 (d, 1H, J = 12.0 Hz), 4.68 (d, 1H, J = 12.0 Hz), 5.90 (s, 1H), 7.04 (s, 1H), 7.15-7.20 (m, 3H), 7.24 (br s, 1H), 7.28-7.35 (m, 4H), 8.38 (s, 1H).	507	505	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form
245		¹ H-NMR (DMSO-D ₆) δ: 0.82 (t, 3H, J = 7.4 Hz), 1.15 (tq, 2H, J = 7.4 Hz, 7.4Hz), 1.39 (tt, 2H, J = 7.4 Hz, 7.4Hz), 2.48 (t, 2H, J = 7.4 Hz), 3.36 (dd, 1H, J = 9.2 Hz, 5.6Hz), 3.59 (dd, 1H, J = 9.2 Hz, 9.2Hz), 4.32 (dd, 1H, J = 9.2 Hz, 5.6Hz), 6.33 (s, 1H), 6.60 (s, 1H), 6.90 (s, 1H), 6.98 (s, 1H), 7.09 (d, 1H, J = 7.6 Hz), 7.17 (d, 1H, J = 7.6 Hz), 7.20-7.30 (m, 3H), 7.32-7.43 (m, 3H), 10.70 (s, 1H).	404	402	R-form
246		¹ H-NMR (DMSO-D ₆) δ: 2.36-2.38 (m, 2H), 3.32-3.33 (m, 1H), 3.37-3.41 (m, 1H), 3.46-3.49 (m, 1H), 4.85 (m, 2H), 5.53-5.59 (m, 1H), 6.94 (s, 1H), 7.17-7.18 (m, 1H), 7.22-7.24 (m, 2H), 7.34-7.41 (m, 6H), 7.63 (s, 1H), 10.88 (s, 1H).	527	525	S-form





247		¹ H-NMR (DMSO-D ₆) δ: 2.38-2.40 (m, 2H), 3.32-3.35 (m, 1H), 3.40-3.45 (m, 1H), 3.48-3.52 (m, 1H), 3.98 (q, 2H, J = 9.3 Hz), 4.61 (s, 2H), 6.93 (s, 1H), 7.18-7.40 (m, 8H), 7.63 (s, 1H), 10.86 (s, 1H).	477	475	S-form
248		¹ H-NMR (CDCl ₃) δ: 2.61 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.8 Hz), 3.32-3.42 (m, 1H), 3.61 (t, 1H, J = 8.8 Hz), 3.71 (dd, 1H, J = 9.6, 7.2 Hz), 3.79 (q, 2H, J = 8.8 Hz), 4.59 (s, 2H), 6.04 (s, 1H), 6.89 (d, 1H, J = 9.2 Hz), 6.99-7.10 (m, 5H), 7.19-7.24 (m, 2H), 8.48 (s, 1H).	495	493	S-form




249		¹ H-NMR (CDCl ₃) δ: 0.83-0.88 (m, 4H), 1.22-1.28 (m, 3H), 1.95 (dd, 1H, J = 9.3, 4.8 Hz), 2.49 (t, 2H, J = 8.5 Hz), 3.60 (d, 1H, J = 10.1 Hz), 4.01 (d, 1H, J = 10.1 Hz), 5.14 (s, 1H), 6.99 (s, 1H), 7.04-7.34 (m, 9H), 7.83 (s, 1H).	401	399	Single Diastereoisomer (racemate)
250		¹ H-NMR (DMSO-D ₆) δ: 0.66-0.71 (m, 1H), 0.78-0.82 (m, 1H), 0.88-0.99 (m, 2H), 3.33-3.37 (m, 1H), 3.50-3.54 (m, 1H), 3.56-3.60 (m, 1H), 3.97 (q, 2H, J = 9.3 Hz), 4.61 (s, 2H), 6.93 (s, 1H), 7.15-7.17 (m, 1H), 7.22-7.26 (m, 3H), 7.31-7.41 (m, 5H), 7.72 (s, 1H), 10.73 (s, 1H).	485	483	Optically active substance
251		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 1.44 (tq, 2H, J = 7.4, 7.4 Hz), 2.46 (t, 2H, J = 7.4 Hz), 2.87-2.92 (m, 1H), 3.27-3.30 (m, 4H), 6.23 (s, 2H), 6.88 (s, 1H), 6.99 (s, 1H), 7.05-7.07 (m, 1H), 7.17-7.27 (m, 4H), 7.33-7.41 (m, 3H), 10.91 (s, 1H).	404	401	None
252		¹ H-NMR (DMSO-D ₆) δ: 2.88-2.91 (m, 1H), 3.27-3.31 (m, 4H), 6.23 (s, 2H), 6.99 (s, 1H), 7.08 (s, 1H), 7.24-7.26 (m, 2H), 7.33-7.54 (m, 6H), 10.96 (s, 1H).	446	444	None





253		¹ H-NMR (DMSO-D ₆) δ: 2.63-2.70 (m, 1H), 3.20-3.30 (m, 1H), 3.40-3.52 (m, 1H), 3.58-3.72 (m, 2H), 4.86 (t, 1H, J = 4.8 Hz), 7.02 (s, 1H), 7.08 (s, 1H), 7.23-7.28 (m, 2H), 7.33-7.45 (m, 5H), 7.52 (t, 1H, J = 8.0 Hz), 7.70 (s, 1H), 10.87 (s, 1H).	461	459	trans-form (racemate)
254		¹ H-NMR (DMSO-D ₆) δ: 1.13 (d, 3H, J = 6.3 Hz), 2.52-2.55 (m, 1H), 2.71-2.77 (m, 1H), 3.69-3.73 (m, 1H), 4.21-4.23 (m, 1H), 4.33-4.36 (m, 1H), 6.99 (s, 1H), 7.09 (s, 1H), 7.25-7.27 (m, 2H), 7.34-7.44 (m, 6H), 7.51-7.53 (m, 1H), 11.16 (s, 1H).	461	459	Single Diastereoisomer (racemate)
255		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 1.39-1.47 (m, 1H), 1.44 (tq, 2H, J = 7.4, 7.5 Hz), 1.76-2.06 (m, 3H), 2.19-2.22 (m, 1H), 2.45 (t, 2H, J = 7.5 Hz), 2.53-2.56 (m, 1H), 3.12-3.18 (m, 1H), 3.35-3.38 (m, 2H), 6.87 (s, 1H), 7.00 (s, 1H), 7.05-7.07 (m, 1H), 7.16-7.28 (m, 4H), 7.31-7.43 (m, 4H), 10.72 (s, 1H).	417	415	Optically active substance (Structure Information A) Optical isomer of Example 256




256		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 1.39-1.47 (m, 1H), 1.44 (tq, 2H, J = 7.4, 7.5 Hz), 1.76-2.06 (m, 3H), 2.19-2.22 (m, 1H), 2.45 (t, 2H, J = 7.5 Hz), 2.53-2.56 (m, 1H), 3.12-3.18 (m, 1H), 3.35-3.38 (m, 2H), 6.87 (s, 1H), 7.00 (s, 1H), 7.05-7.07 (m, 1H), 7.16-7.28 (m, 4H), 7.31-7.43 (m, 4H), 10.72 (s, 1H).	417	415	Optically active substance (Structure Information A) Optical isomer of Example 255
257		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.3, 7.9 Hz), 2.45 (d, 2H, J = 7.9 Hz), 3.52 (dd, 1H, J = 9.0, 5.9 Hz), 3.75 (dd, 1H, J = 9.2, 9.0 Hz), 5.13 (dd, 1H, J = 9.2, 5.9 Hz), 6.89 (s, 1H), 7.01-7.10 (m, 2H), 7.19-7.41 (m, 7H), 7.75 (s, 1H), 11.04 (s, 1H).	391	389	S-form
258		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.3, 7.9 Hz), 2.45 (d, 2H, J = 7.9 Hz), 3.52 (dd, 1H, J = 9.0, 5.9 Hz), 3.75 (dd, 1H, J = 9.2, 9.0 Hz), 5.13 (dd, 1H, J = 9.2, 5.9 Hz), 6.89 (s, 1H), 7.01-7.10 (m, 2H), 7.19-7.41 (m, 7H), 7.75 (s, 1H), 11.04 (s, 1H).	391	389	R-form

259		¹ H-NMR (DMSO-D ₆) δ: 1.13 (d, 3H, J = 6.4 Hz), 2.74 (ddd, 1H, J = 9.6, 9.0, 4.0 Hz), 3.72 (dq, 1H, J = 12.8, 8.8 Hz), 4.22 (dd, 1H, J = 10.0, 10.0 Hz), 4.35 (dd, 1H, J = 10.8, 4.0 Hz), 6.99 (s, 1H), 7.10 (br s, 1H), 7.24-7.39 (m, 6H), 7.42 (br s, 1H), 7.53 (t, 1H, J = 8.4 Hz), 11.16 (s, 1H).	479	477	Single Diastereoisomer (racemate)
260		¹ H-NMR (DMSO-D ₆) δ: 2.39 (d, 2H, J = 8.5 Hz), 3.30-3.54 (m, 3H), 4.73 (q, 2H, J = 8.9 Hz), 6.80 (d, 1H, J = 7.7 Hz), 6.97 (s, 1H), 6.99-7.07 (m, 2H), 7.22-7.44 (m, 6H), 7.64 (s, 1H), 10.87 (s, 1H).	445	443	S-form
261		¹ H-NMR (DMSO-D ₆) δ: 0.71 (t, 3H, J = 7.5 Hz), 1.33-1.44 (m, 2H), 1.89 (s, 3H), 2.35-2.46 (m, 4H), 3.31-3.53 (m, 3H), 6.92-6.96 (m, 2H), 6.98-7.03 (m, 1H), 7.07-7.12 (m, 1H), 7.16-7.40 (m, 5H), 7.64 (s, 1H), 10.76 (s, 1H).	403	401	S-form
262		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, J = 7.5 Hz), 1.41-1.52 (m, 2H), 2.39 (d, 2H, J = 8.5 Hz), 6.91 (s, 1H), 7.02-7.05 (m, 1H), 7.10-7.24 (m, 3H), 7.28-7.34 (m, 2H), 7.38-7.42 (m, 2H), 7.64 (s, 1H), 10.89 (s, 1H).	423	421	S-form





263		¹ H-NMR (DMSO-D ₆) δ: 3.52 (dd, 1H, J = 9.1, 6.0 Hz), 3.76 (dd, 1H, J = 9.2, 9.1 Hz), 5.13 (dd, 1H, J = 9.2, 6.0 Hz), 7.01 (s, 1H), 7.09 (s, 1H), 7.25-7.28 (m, 2H), 7.36-7.45 (m, 5H), 7.53 (t, 1H, J = 8.0 Hz), 7.75 (s, 1H), 11.09 (s, 1H).	433	431	S-form
264		¹ H-NMR (DMSO-D ₆) δ: 3.52 (dd, 1H, J = 9.1, 6.0 Hz), 3.76 (dd, 1H, J = 9.2, 9.1 Hz), 5.13 (dd, 1H, J = 9.2, 6.0 Hz), 7.01 (s, 1H), 7.09 (s, 1H), 7.25-7.28 (m, 2H), 7.36-7.45 (m, 5H), 7.53 (t, 1H, J = 8.0 Hz), 7.75 (s, 1H), 11.09 (s, 1H).	433	431	R-form
265		¹ H-NMR (DMSO-D ₆) δ: 2.37-2.40 (m, 2H), 3.18 (s, 3H), 3.34-3.36 (m, 1H), 3.40-3.44 (m, 1H), 3.48-3.52 (m, 1H), 4.35 (s, 2H), 6.91 (s, 1H), 7.11-7.13 (m, 1H), 7.20-7.25 (m, 3H), 7.28-7.41 (m, 5H), 7.63 (s, 1H), 10.86 (s, 1H).	391	389	S-form
266		¹ H-NMR (DMSO-D ₆) δ: 3.36 (dd, 1H, J = 9.0, 6.4 Hz), 3.60 (dd, 1H, J = 9.4, 9.0 Hz), 4.32 (dd, 1H, J = 9.4, 6.4 Hz), 6.33 (s, 1H), 6.60 (s, 1H), 7.01 (s, 1H), 7.09 (s, 1H), 7.24-7.28 (m, 2H), 7.35-7.46 (m, 5H), 7.53 (t, 1H, J = 7.9 Hz), 10.74 (s, 1H).	432	430	S-form




267		<p>¹H-NMR (DMSO-D₆) δ: 3.36 (dd, 1H, J = 9.0, 6.4 Hz), 3.60 (dd, 1H, J = 9.4, 9.0 Hz), 4.32 (dd, 1H, J = 9.4, 6.4 Hz), 6.33 (s, 1H), 6.60 (s, 1H), 7.01 (s, 1H), 7.09 (s, 1H), 7.24-7.28 (m, 2H), 7.35-7.46 (m, 5H), 7.53 (t, 1H, J = 7.9 Hz), 10.74 (s, 1H).</p>	432	430	R-form
268		<p>¹H-NMR (DMSO-D₆) δ: 0.82 (t, 3H, J = 7.4 Hz), 1.15 (tq, 2H, J = 7.4 Hz, 7.4Hz), 1.39 (tt, 2H, J = 7.4 Hz, 7.4Hz), 2.48 (t, 2H, J = 7.4 Hz), 3.36 (dd, 1H, J = 9.2 Hz, 5.6Hz), 3.59 (dd, 1H, J = 9.2 Hz, 9.2Hz), 4.32 (dd, 1H, J = 9.2 Hz, 5.6Hz), 6.33 (s, 1H), 6.60 (s, 1H), 6.90 (s, 1H), 6.98 (s, 1H), 7.09 (d, 1H, J = 7.6 Hz), 7.17 (d, 1H, J = 7.6 Hz), 7.20-7.30 (m, 3H), 7.32-7.43 (m, 3H), 10.70 (s, 1H).</p>	404	402	S-form
269		<p>¹H-NMR (DMSO-D₆) δ: 1.27 (s, 6H), 2.77 (s, 3H), 3.32-3.37 (br m, 1H), 3.60 (t, 1H, J = 9.3 Hz), 4.33 (dd, 1H, J = 9.3, 5.3 Hz), 6.33 (s, 1H), 6.60 (s, 1H), 6.93 (s, 1H), 7.05 (s, 1H), 7.22-7.42 (m, 8H), 10.69 (s, 1H).</p>	420	418	R-form




270		¹ H-NMR (DMSO-D ₆) δ: 1.27 (s, 6H), 2.77 (s, 3H), 3.32-3.37 (br m, 1H), 3.60 (t, 1H, J = 9.3 Hz), 4.33 (dd, 1H, J = 9.3, 5.3 Hz), 6.33 (br s, 1H), 6.60 (br s, 1H), 6.93 (s, 1H), 7.05 (br s, 1H), 7.22-7.42 (m, 8H), 10.69 (br s, 1H).	420	418	S-form
271		¹ H-NMR (DMSO-D ₆) δ: 1.27 (s, 6H), 2.77 (s, 3H), 2.85-2.94 (m, 1H), 3.23-3.34 (m, 4H), 6.25 (br s, 2H), 6.91 (s, 1H), 7.05 (br s, 1H), 7.22-7.27 (m, 3H), 7.32-7.43 (m, 5H), 10.92 (br s, 1H).	434	432	None
272		¹ H-NMR (DMSO-D ₆) δ: 2.39 (d, 2H, J = 8.4 Hz), 2.62-2.75 (m, 2H), 3.30-3.54 (m, 3H), 4.11 (t, 2H, J = 6.0 Hz), 6.78-6.83 (m, 2H), 6.92-6.97 (m, 2H), 7.22-7.43 (m, 6H), 7.64 (s, 1H), 10.86 (s, 1H).	459	457	S-form
273		¹ H-NMR (DMSO-D ₆) δ: 1.81-1.90 (m, 2H), 2.29-2.43 (m, 4H), 3.29-3.53 (m, 3H), 3.95 (t, 2H, J = 6.2 Hz), 6.75-6.82 (m, 2H), 6.90-6.96 (m, 2H), 7.22-7.29 (m, 3H), 7.31-7.44 (m, 3H), 7.64 (s, 1H), 10.86 (s, 1H).	473	471	S-form




274		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.39-1.47 (m, 2H), 1.52-1.60 (m, 2H), 1.86-1.94 (m, 2H), 2.26-2.31 (m, 1H), 2.44-2.48 (m, 3H), 2.66-2.69 (m, 1H), 3.12-3.16 (m, 2H), 6.88 (s, 1H), 7.00 (s, 1H), 7.05-7.07 (m, 1H), 7.15-7.17 (m, 1H), 7.20-7.28 (m, 3H), 7.31-7.41 (m, 3H), 7.49-7.50 (m, 1H), 10.69 (s, 1H).	417	415	racemate
275		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 1.44 (tq, 2H, J = 7.5, 7.4 Hz), 2.46 (t, 2H, J = 7.5 Hz), 3.43-3.47 (m, 2H), 4.13 (d, 1H, J = 16.1 Hz), 4.22 (d, 1H, J = 16.1 Hz), 4.46 (dd, 1H, J = 8.4, 4.4 Hz), 6.87 (s, 1H), 7.01 (s, 1H), 7.06-7.08 (m, 1H), 7.16-7.18 (m, 1H), 7.22-7.29 (m, 3H), 7.35-7.42 (m, 3H), 8.14 (s, 1H), 10.61 (s, 1H).	405	403	Optically active substance (Structure Information B) Optical isomer of Example 277
276		¹ H-NMR (DMSO-D ₆) δ: 3.43-3.47 (m, 2H), 4.13 (d, 1H, J = 16.3 Hz), 4.22 (d, 1H, J = 16.3 Hz), 4.47 (dd, 1H, J = 8.6, 4.4 Hz), 6.98 (s, 1H), 7.09 (s, 1H), 7.25-7.28 (m, 2H), 7.35-7.45 (m, 5H), 7.52 (t, 1H, J = 7.9 Hz), 8.14 (s, 1H), 10.68 (s, 1H).	447	445	Optically active substance (The same configuration as Example 275) Optical isomer of Example 278




277		¹ H-NMR (CDCl ₃) δ: 0.84 (t, 3H, J = 7.2 Hz), 1.51 (tq, 2H, J = 7.2, 8.0 Hz), 2.49 (t, 2H, J = 8.0 Hz), 3.61 (t, 1H, J = 12 Hz), 3.82-3.87 (m, 1H), 4.33 (d, 1H, J = 17.2 Hz), 4.42-4.48 (m, 2H), 6.13 (s, 1H), 7.00 (s, 1H), 7.04-7.08 (m, 2H), 7.13 (d, 1H, J = 7.6 Hz), 7.20 (d, 1H, J = 8.0 Hz), 7.23-7.26 (m, 2H), 7.28-7.36 (m, 3H), 8.90 (s, 1H).	405	403	Optically active substance (Structure Information B) Optical isomer of Example 275
278		¹ H-NMR (CDCl ₃) δ: 3.61 (t, 1H, J = 12 Hz), 3.85 (m, 1H), 4.33 (d, 1H, J = 17.2 Hz), 4.42-4.48 (m, 2H), 6.12 (s, 1H), 7.06 (m, 2H), 7.17 (d, 1H, J = 8.4 Hz), 7.21-7.25 (m, 3H), 7.33-7.40 (m, 4H), 8.91 (s, 1H).	447	445	Optically active substance (The same configuration as Example 277) Optical isomer of Example 276
279		¹ H-NMR (DMSO-D ₆) δ: 2.37-2.39 (m, 2H), 3.34-3.36 (m, 1H), 3.38-3.44 (m, 1H), 3.48-3.52 (m, 1H), 3.95 (d, 1H, J = 9.5 Hz), 4.00 (d, 1H, J = 9.5 Hz), 4.61 (s, 2H), 6.93 (s, 1H), 7.15-7.17 (m, 1H), 7.22-7.26 (m, 3H), 7.32-7.41 (m, 6H), 7.64 (s, 1H), 10.87 (s, 1H).	459	457	S-form




280		¹ H-NMR (DMSO-D ₆) δ: 2.37-2.39 (m, 2H), 3.34-3.36 (m, 1H), 3.38-3.44 (m, 1H), 3.48-3.52 (m, 1H), 3.95 (d, 1H, J = 9.5 Hz), 4.00 (d, 1H, J = 9.5 Hz), 4.61 (s, 2H), 6.93 (s, 1H), 7.15-7.17 (m, 1H), 7.22-7.26 (m, 3H), 7.32-7.41 (m, 6H), 7.64 (s, 1H), 10.87 (s, 1H).	459	457	R-form
281		¹ H-NMR (DMSO-D ₆) δ: 2.39 (d, 2H, J = 8.4 Hz), 2.48-2.57 (m, 2H), 3.32-3.52 (m, 3H), 3.54 (t, 2H, J = 6.0 Hz), 4.45 (s, 2H), 6.91 (s, 1H), 7.12 (d, 1H, J = 7.6 Hz), 7.22-7.24 (m, 3H), 7.29-7.41 (m, 5H), 7.64 (s, 1H), 10.87 (s, 1H).	473	471	S-form
282		¹ H-NMR (DMSO-D ₆) δ: 2.39 (d, 2H, J = 8.4 Hz), 2.48-2.57 (m, 2H), 3.32-3.52 (m, 3H), 3.54 (t, 2H, J = 6.0 Hz), 4.45 (s, 2H), 6.91 (s, 1H), 7.12 (d, 1H, J = 7.6 Hz), 7.22-7.24 (m, 3H), 7.29-7.41 (m, 5H), 7.64 (s, 1H), 10.87 (s, 1H).	473	471	R-form
283		¹ H-NMR (DMSO-D ₆) δ: 1.07 (t, 3H, J = 6.8 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.31-3.37 (m, 3H), 3.38-3.53 (m, 2H), 4.39 (s, 2H), 6.91 (s, 1H), 7.13 (d, 1H, J = 7.6 Hz), 7.19 (s, 1H), 7.22-7.24 (m, 2H), 7.28-7.41 (m, 5H), 7.63 (s, 1H), 10.87 (s, 1H).	405	403	S-form




284		¹ H-NMR (CDCl ₃) δ: 0.98 (t, 3H, J = 7.6 Hz), 1.72 (tq, 2H, J = 7.6, 6.8 Hz), 2.43 (dd, 1H, J = 16.4, 9.2 Hz), 2.72 (dd, 1H, J = 16.4, 8.4 Hz), 3.10-3.20 (m, 1H), 3.43 (t, 1H, J = 8.8 Hz), 3.60 (dd, 1H, J = 9.6, 7.2 Hz), 3.78 (t, 2H, J = 6.8 Hz), 6.34 (s, 1H), 6.74-6.80 (m, 2H), 6.85 (dd, 1H, J = 8.0, 2.0 Hz), 7.05 (s, 1H), 7.18 (t, 1H, J = 8.0 Hz), 7.26-7.38 (m, 5H), 9.37 (s, 1H).	405	403	S-form
285		¹ H-NMR (CDCl ₃) δ: 0.98 (t, 3H, J = 7.6 Hz), 1.72 (tq, 2H, J = 7.6, 6.8 Hz), 2.43 (dd, 1H, J = 16.4, 9.2 Hz), 2.72 (dd, 1H, J = 16.4, 8.4 Hz), 3.10-3.20 (m, 1H), 3.43 (t, 1H, J = 8.8 Hz), 3.60 (dd, 1H, J = 9.6, 7.2 Hz), 3.78 (t, 2H, J = 6.8 Hz), 6.34 (s, 1H), 6.74-6.80 (m, 2H), 6.85 (dd, 1H, J = 8.0, 2.0 Hz), 7.05 (s, 1H), 7.18 (t, 1H, J = 8.0 Hz), 7.26-7.38 (m, 5H), 9.37 (s, 1H).	405	403	R-form
286		¹ H-NMR (CDCl ₃) δ: 0.97 (d, 6H, J = 6.4 Hz), 1.93-2.04 (m, 1H), 2.43 (dd, 1H, J = 16.8, 9.6 Hz), 2.72 (dd, 1H, J = 16.8, 8.4 Hz), 3.11-3.19 (m, 1H), 3.43 (t, 1H, J = 9.2 Hz), 3.54-3.64 (m, 3H), 6.35 (s, 1H), 6.72-6.89 (m, 3H), 7.05 (s, 1H), 7.17 (t, 1H, J = 8.0 Hz), 7.23-7.39 (m, 5H), 9.38 (s, 1H).	419	417	S-form




287		¹ H-NMR (CDCl ₃) δ: 0.97 (d, 6H, J = 6.4 Hz), 1.93-2.04 (m, 1H), 2.43 (dd, 1H, J = 16.8, 9.6 Hz), 2.72 (dd, 1H, J = 16.8, 8.4 Hz), 3.11-3.19 (m, 1H), 3.43 (t, 1H, J = 9.2 Hz), 3.54-3.64 (m, 3H), 6.35 (s, 1H), 6.72-6.89 (m, 3H), 7.05 (s, 1H), 7.17 (t, 1H, J = 8.0 Hz), 7.23-7.39 (m, 5H), 9.38 (s, 1H).	419	417	R-form
288		¹ H-NMR (CDCl ₃) δ: 0.99 (s, 9H), 2.43 (dd, 1H, J = 16.8, 9.6 Hz), 2.72 (dd, 1H, J = 16.8, 8.4 Hz), 3.10-3.20 (m, 1H), 3.47-3.58 (m, 3H), 3.60 (dd, 1H, J = 9.6, 7.2 Hz), 6.34 (s, 1H), 6.71 (d, 1H, J = 7.6 Hz), 6.82-6.89 (m, 2H), 7.06 (s, 1H), 7.15 (t, 1H, J = 7.6 Hz), 7.24-7.39 (m, 5H), 9.39 (s, 1H).	433	431	S-form
289		¹ H-NMR (CDCl ₃) δ: 0.99 (s, 9H), 2.43 (dd, 1H, J = 16.8, 9.6 Hz), 2.72 (dd, 1H, J = 16.8, 8.4 Hz), 3.10-3.20 (m, 1H), 3.47-3.58 (m, 3H), 3.60 (dd, 1H, J = 9.6, 7.2 Hz), 6.34 (s, 1H), 6.71 (d, 1H, J = 7.6 Hz), 6.82-6.89 (m, 2H), 7.06 (s, 1H), 7.15 (t, 1H, J = 7.6 Hz), 7.24-7.39 (m, 5H), 9.39 (s, 1H).	433	431	R-form




290		¹ H-NMR (DMSO-D ₆) δ: 1.11 (s, 9H), 2.36-2.41 (m, 2H), 3.30-3.53 (m, 3H), 6.67 (t, 1H, J = 1.8 Hz), 6.89 (s, 1H), 6.93-6.97 (m, 1H), 7.06-7.10 (m, 1H), 7.22-7.44 (m, 6H), 7.63 (s, 1H), 10.85 (s, 1H).	419	417	S-form
291		¹ H-NMR (DMSO-D ₆) δ: 0.81 (t, 3H, J = 7.5 Hz), 1.02 (s, 6H), 1.41 (q, 2H, J = 7.5 Hz), 2.37-2.41 (m, 2H), 3.30-3.53 (m, 3H), 6.63 (t, 1H, J = 2.0 Hz), 6.89 (s, 1H), 6.91-6.95 (m, 1H), 7.07-7.11 (m, 1H), 7.22-7.26 (m, 2H), 7.28-7.37 (m, 2H), 7.38-7.44 (m, 2H), 7.63 (s, 1H), 10.85 (s, 1H).	433	431	S-form
292		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.5 Hz), 1.12 (d, 3H, J = 7.3 Hz), 1.39-1.50 (m, 2H), 1.66-1.74 (m, 1H), 2.08-2.17 (m, 1H), 2.36 (q, 1H, J = 6.9 Hz), 2.46 (t, 2H, J = 7.5 Hz), 2.93-3.01 (m, 1H), 3.26-3.39 (m, 2H), 6.89 (s, 1H), 6.99-7.01 (m, 1H), 7.04-7.09 (m, 1H), 7.14-7.19 (m, 1H), 7.20-7.45 (m, 7H), 10.84 (s, 1H).	417	415	Single diastereoisomer (racemate) Geometrical isomer of Example 293




293		<p>¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.5 Hz), 1.09 (d, 3H, J = 7.3 Hz), 1.39-1.49 (m, 2H), 1.62 (q, 1H, J = 12.4 Hz), 2.05-2.13 (m, 1H), 2.18-2.28 (m, 1H), 2.46 (t, 2H, J = 7.5 Hz), 2.88-2.98 (m, 1H), 3.24-3.37 (m, 2H), 6.88 (s, 1H), 6.98-7.01 (m, 1H), 7.05-7.09 (m, 1H), 7.15-7.19 (m, 1H), 7.20-7.29 (m, 3H), 7.31-7.44 (m, 4H), 10.81 (s, 1H).</p>	417	415	Single diastereoisomer (racemate) Geometrical isomer of Example 292
294		<p>¹H-NMR (DMSO-D₆) δ: 0.90 (s, 9H), 1.55 (t, 2H, J = 7.5 Hz), 2.37-2.42 (m, 2H), 3.30-3.54 (m, 3H), 3.85 (t, 2H, J = 7.5 Hz), 6.68-6.71 (m, 1H), 6.80-6.84 (m, 1H), 6.87-6.93 (m, 2H), 7.23-7.28 (m, 3H), 7.31-7.37 (m, 1H), 7.38-7.44 (m, 2H), 7.63 (s, 1H), 10.85 (s, 1H).</p>	447	445	S-form
295		<p>¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.01 (t, 3H, J = 7.2 Hz), 1.44 (tq, 2H, J = 7.5, 7.3 Hz), 2.46 (t, 2H, J = 7.5 Hz), 2.87 (dq, 1H, J = 7.2, 14.0 Hz), 3.22 (dd, 1H, J = 9.2, 6.4 Hz), 3.36 (dq, 1H, J = 7.2, 14.0 Hz), 3.52 (dd, 1H, J = 9.5, 9.2 Hz), 4.41 (dd, 1H, J = 9.5, 6.4 Hz), 6.42 (s, 1H), 6.92 (s, 1H), 7.02 (s, 1H), 7.07-7.09 (m, 1H), 7.16-7.19 (m, 1H), 7.22-7.29 (m, 3H), 7.33-7.43 (m, 3H), 11.06 (s, 1H).</p>	418	416	S-form





296		¹ H-NMR (DMSO-D ₆) δ: 1.01 (t, 3H, J = 7.2 Hz), 2.87 (dq, 1H, J = 14.2, 7.2 Hz), 3.22 (dd, 1H, J = 8.9, 6.4 Hz), 3.36 (dq, 1H, J = 14.2, 7.2 Hz), 3.52 (dd, 1H, J = 9.6, 8.9 Hz), 4.41 (dd, 1H, J = 9.6, 6.4 Hz), 6.42 (s, 1H), 7.03 (s, 1H), 7.10 (s, 1H), 7.24-7.27 (m, 2H), 7.35-7.45 (m, 5H), 7.52 (t, 1H, J = 8.0 Hz), 11.12 (s, 1H).	460	458	S-form
297		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 0.82 (t, 3H, J = 7.4 Hz), 1.35-1.52 (m, 4H), 2.46 (t, 2H, J = 7.4 Hz), 2.76-2.83 (m, 1H), 3.20-3.29 (m, 2H), 3.53 (dd, 1H, J = 9.5, 8.8 Hz), 4.39 (dd, 1H, J = 9.5, 6.0 Hz), 6.40 (s, 1H), 6.92 (s, 1H), 7.02 (s, 1H), 7.06-7.08 (m, 1H), 7.16-7.18 (m, 1H), 7.22-7.29 (m, 3H), 7.33-7.42 (m, 3H), 11.05 (s, 1H).	432	430	S-form
298		¹ H-NMR (DMSO-D ₆) δ: 0.82 (t, 3H, J = 7.4 Hz), 1.35-1.51 (m, 2H), 2.76-2.83 (m, 1H), 3.22-3.28 (m, 2H), 3.53 (dd, 1H, J = 9.5, 9.0 Hz), 4.39 (dd, 1H, J = 9.5, 6.0 Hz), 6.41 (s, 1H), 7.02 (s, 1H), 7.10 (s, 1H), 7.24-7.28 (m, 2H), 7.35-7.45 (m, 5H), 7.52 (t, 1H, J = 8.0 Hz), 11.10 (s, 1H).	474	472	S-form




299		¹ H-NMR (DMSO-D ₆) δ: 1.06 (d, 6H, J = 6.4 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.33-3.35 (m, 1H), 3.38-3.55 (m, 3H), 4.39 (s, 2H), 6.90 (s, 1H), 7.12 (d, 1H, J = 7.6 Hz), 7.20-7.24 (m, 3H), 7.27-7.41 (m, 5H), 7.63 (s, 1H), 10.86 (s, 1H).	419	417	S-form
300		¹ H-NMR (DMSO-D ₆) δ: 0.84 (t, 3H, 7.2 Hz), 1.47 (tq, 2H, 6.4, 7.2 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.26 (t, 2H, J = 6.4 Hz), 3.32-3.52 (m, 3H), 4.39 (s, 2H), 6.91 (s, 1H), 7.14 (d, 1H, J = 7.6 Hz), 7.19 (s, 1H), 7.22-7.24 (m, 2H), 7.28-7.41 (m, 5H), 7.63 (s, 1H), 10.86 (s, 1H).	419	417	S-form
301		¹ H-NMR (DMSO-D ₆) δ: 1.13 (d, 6H, J = 5.6 Hz), 2.36-2.42 (m, 2H), 3.30-3.36 (m, 1H), 3.37-3.46 (m, 1H), 3.47-3.54 (m, 1H), 4.43 (quint, 1H, J = 5.6 Hz), 6.67-6.69 (m, 1H), 6.79-6.82 (m, 1H), 6.86-6.90 (m, 1H), 6.90 (s, 1H), 7.21-7.28 (m, 3H), 7.32-7.37 (m, 1H), 7.38-7.43 (m, 2H), 7.63 (s, 1H), 10.84 (s, 1H).	405	403	S-form




302		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.5 Hz), 1.39-1.49 (m, 2H), 2.04-2.23 (m, 4H), 2.46 (t, 2H, J = 7.5 Hz), 2.72-2.80 (m, 1H), 3.13-3.20 (m, 4H), 6.89 (s, 1H), 6.99-7.02 (m, 1H), 7.04-7.08 (m, 1H), 7.15-7.19 (m, 1H), 7.20-7.29 (m, 3H), 7.31-7.42 (m, 3H), 10.83 (s, 1H).	438	436	None
303		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.01 (t, 3H, J = 7.2 Hz), 1.44 (tq, 2H, J = 7.5, 7.3 Hz), 2.46 (t, 2H, J = 7.5 Hz), 2.87 (dq, 1H, J = 7.2, 14.0 Hz), 3.22 (dd, 1H, J = 9.2, 6.4 Hz), 3.36 (dq, 1H, J = 7.2, 14.0 Hz), 3.52 (dd, 1H, J = 9.5, 9.2 Hz), 4.41 (dd, 1H, J = 9.5, 6.4 Hz), 6.42 (s, 1H), 6.92 (s, 1H), 7.02 (s, 1H), 7.07-7.09 (m, 1H), 7.16-7.19 (m, 1H), 7.22-7.29 (m, 3H), 7.33-7.43 (m, 3H), 11.06 (s, 1H).	418	416	R-form
304		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 0.82 (t, 3H, J = 7.4 Hz), 1.35-1.52 (m, 4H), 2.46 (t, 2H, J = 7.4 Hz), 2.76-2.83 (m, 1H), 3.20-3.29 (m, 2H), 3.53 (dd, 1H, J = 9.5, 8.8 Hz), 4.39 (dd, 1H, J = 9.5, 6.0 Hz), 6.40 (s, 1H), 6.92 (s, 1H), 7.02 (s, 1H), 7.06-7.08 (m, 1H), 7.16-7.18 (m, 1H), 7.22-7.29 (m, 3H), 7.33-7.42 (m, 3H), 11.05 (s, 1H).	432	430	R-form




305		¹ H-NMR (DMSO-D ₆) δ: 3.52 (dd, 1H, J = 8.7, 5.7 Hz), 3.75 (dd, 1H, J = 9.3, 8.7 Hz), 3.98 (q, 2H, J = 9.3 Hz), 4.61 (s, 2H), 5.13 (dd, 1H, J = 9.3, 5.7 Hz), 6.93 (s, 1H), 7.16-7.18 (m, 1H), 7.24-7.26 (m, 3H), 7.32-7.42 (m, 5H), 7.76 (s, 1H), 11.07 (s, 1H).	461	459	R-form
306		¹ H-NMR (DMSO-D ₆) δ: 0.82 (t, 3H, J = 7.4 Hz), 1.15 (tq, 2H, J = 7.4 Hz, 7.4Hz), 1.39 (tt, 2H, J = 7.4 Hz, 7.4Hz), 2.49 (t, 2H, J = 7.4 Hz), 3.52 (dd, 1H, J = 9.2 Hz, 5.6Hz), 3.75 (dd, 1H, J = 9.2 Hz, 9.2Hz), 5.13 (dd, 1H, J = 9.2 Hz, 5.6Hz), 6.90 (s, 1H), 6.99 (s, 1H), 7.09 (d, 1H, J = 7.6 Hz), 7.17 (d, 1H, J = 7.6 Hz), 7.22-7.30 (m, 3H), 7.32-7.43 (m, 3H), 7.75 (s, 1H), 11.04 (s, 1H).	405	403	R-form
307		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 0.90 (t, 3H, J = 7.4 Hz), 1.39-1.50 (m, 4H), 1.65-1.75 (m, 1H), 2.46 (t, 2H, J = 7.4 Hz), 2.50-2.60 (m, 1H), 3.12-3.21 (m, 2H), 3.40-3.46 (m, 1H), 6.92 (s, 1H), 7.01 (s, 1H), 7.07 (d, 1H, J = 7.6 Hz), 7.17 (d, 1H, J = 7.6 Hz), 7.20-7.28 (m, 3H), 7.32-7.42 (m, 3H), 7.66 (s, 1H), 10.96 (s, 1H).	417	415	Optically active substance (transform, Structure Information C) Optical isomer of Example 309




308		<p>¹H-NMR (DMSO-D₆) δ: 0.90 (t, 3H, J = 7.6 Hz), 1.39-1.49 (m, 1H), 1.65-1.75 (m, 1H), 2.50-2.60 (m, 1H), 3.12-3.21 (m, 2H), 3.40-3.48 (m, 1H), 7.02 (s, 1H), 7.09 (s, 1H), 7.26 (m, 2H), 7.33-7.45 (m, 5H), 7.52 (t, 1H, J = 8.0Hz), 7.67 (s, 1H), 11.02 (s, 1H).</p>	459	457	Optically active substance (transform, the same configuration as Example 307) Optical isomer of Example 310
309		<p>¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 0.90 (t, 3H, J = 7.4 Hz), 1.39-1.50 (m, 4H), 1.65-1.75 (m, 1H), 2.46 (t, 2H, J = 7.4 Hz), 2.50-2.60 (m, 1H), 3.12-3.21 (m, 2H), 3.40-3.46 (m, 1H), 6.92 (s, 1H), 7.01 (s, 1H), 7.07 (d, 1H, J = 7.6 Hz), 7.17 (d, 1H, J = 7.6 Hz), 7.20-7.28 (m, 3H), 7.32-7.42 (m, 3H), 7.66 (s, 1H), 10.96 (s, 1H).</p>	417	415	Optically active substance (transform, Structure Information C) Optical isomer of Example 307
310		<p>¹H-NMR (DMSO-D₆) δ: 0.90 (t, 3H, J = 7.6 Hz), 1.39-1.49 (m, 1H), 1.65-1.75 (m, 1H), 2.50-2.60 (m, 1H), 3.12-3.21 (m, 2H), 3.40-3.48 (m, 1H), 7.02 (s, 1H), 7.09 (s, 1H), 7.26 (m, 2H), 7.33-7.45 (m, 5H), 7.52 (t, 1H, J = 8.0Hz), 7.67 (s, 1H), 11.02 (s, 1H).</p>	459	457	Optically active substance (transform, the same configuration as Example 309) Optical isomer of Example 308



311		¹ H-NMR (DMSO-D ₆) δ: 1.06 (d, 6H, J = 6.4 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.33-3.35 (m, 1H), 3.38-3.55 (m, 3H), 4.39 (s, 2H), 6.90 (s, 1H), 7.12 (d, 1H, J = 7.6 Hz), 7.20-7.24 (m, 3H), 7.27-7.41 (m, 5H), 7.63 (s, 1H), 10.86 (s, 1H).	419	417	R-form
312		¹ H-NMR (DMSO-D ₆) δ: 0.84 (t, 3H, 7.2 Hz), 1.47 (tq, 2H, 6.4, 7.2 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.26 (t, 2H, J = 6.4 Hz), 3.32-3.52 (m, 3H), 4.39 (s, 2H), 6.91 (s, 1H), 7.14 (d, 1H, J = 7.6 Hz), 7.19 (s, 1H), 7.22-7.24 (m, 2H), 7.28-7.41 (m, 5H), 7.63 (s, 1H), 10.86 (s, 1H).	419	417	R-form
313		¹ H-NMR (DMSO-D ₆) δ: 1.14 (s, 9H), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.33 (s, 2H), 6.90 (s, 1H), 7.10 (d, 1H, J = 7.6 Hz), 7.20 (s, 1H), 7.23-7.42 (m, 7H), 7.64 (s, 1H), 10.86 (s, 1H).	433	431	S-form
314		¹ H-NMR (DMSO-D ₆) δ: 0.83 (d, 6H, J = 6.8 Hz), 1.70-1.81 (m, 1H), 2.39 (d, 2H, J = 8.4 Hz), 3.08 (d, 2H, J = 10. Hz), 3.32-3.52 (m, 3H), 4.39 (s, 2H), 6.90 (s, 1H), 7.14 (d, 1H, J = 7.6 Hz), 7.19 (s, 1H), 7.28-7.41 (m, 7H), 7.64 (s, 1H), 10.86 (s, 1H).	433	431	S-form




315		¹ H-NMR (DMSO-D ₆) δ: 0.83 (s, 9H), 2.39 (d, 2H, J = 8.4 Hz), 2.99 (s, 2H), 3.32-3.52 (m, 3H), 4.42 (s, 2H), 6.90 (s, 1H), 7.15 (d, 1H, J = 7.6 Hz), 7.19 (s, 1H), 7.28-7.41 (m, 7H), 7.64 (s, 1H), 10.86 (s, 1H).	447	445	S-form
316		¹ H-NMR (DMSO-D ₆) δ: 0.89 (t, 3H, J = 7.3 Hz), 1.36 (sext, 2H, J = 7.3 Hz), 1.59 (quint, 2H, J = 7.3 Hz), 2.36-2.42 (m, 2H), 3.31-3.36 (m, 1H), 3.37-3.47 (m, 1H), 3.47-3.54 (m, 1H), 3.83 (t, 2H, J = 7.3 Hz), 6.71-6.75 (m, 1H), 6.75-6.81 (m, 1H), 6.87-6.93 (m, 1H), 6.91 (s, 1H), 7.21-7.29 (m, 3H), 7.31-7.38 (m, 1H), 7.38-7.45 (m, 2H), 7.63 (s, 1H), 10.84 (s, 1H).	419	417	S-form
317		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.5 Hz), 1.38-1.49 (m, 2H), 1.75-1.86 (m, 2H), 2.16-2.25 (m, 2H), 2.46 (t, 2H, J = 7.5 Hz), 2.58-2.72 (m, 3H), 3.23-3.33 (m, 2H), 6.87 (s, 1H), 6.98-7.01 (m, 1H), 7.04-7.08 (m, 1H), 7.14-7.18 (m, 1H), 7.19-7.29 (m, 3H), 7.30-7.42 (m, 3H), 10.74 (s, 1H).	422	420	Single diastereo isomer Diastereo isomer of Example 318




318		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, J = 7.5 Hz), 1.39-1.49 (m, 2H), 1.73-1.83 (m, 2H), 2.22-2.35 (m, 2H), 2.46 (t, 2H, J = 7.5 Hz), 2.60-2.74 (m, 3H), 2.88-2.96 (m, 2H), 6.90 (s, 1H), 7.00-7.02 (m, 1H), 7.05-7.09 (m, 1H), 7.15-7.19 (m, 1H), 7.20-7.29 (m, 3H), 7.31-7.42 (m, 3H), 10.72 (s, 1H).	422	420	Principal component is a diastereoisomer of Example 317.
319		¹ H-NMR (DMSO-D ₆) δ: 0.88 (d, 6H, J = 6.9 Hz), 1.51 (q, 2H, J = 6.9 Hz), 1.62-1.75 (m, 1H), 2.36-2.42 (m, 2H), 3.31-3.36 (m, 1H), 3.36-3.47 (m, 1H), 3.47-3.53 (m, 1H), 3.84 (t, 2H, J = 6.9 Hz), 6.70-6.74 (m, 1H), 6.77-6.82 (m, 1H), 6.87-6.93 (m, 1H), 6.91 (s, 1H), 7.22-7.28 (m, 3H), 7.31-7.37 (m, 1H), 7.37-7.44 (m, 2H), 7.63 (s, 1H), 10.85 (s, 1H).	433	431	S-form
320		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.5 Hz), 1.39-1.49 (m, 2H), 2.16-2.26 (m, 1H), 2.46 (t, 2H, J = 7.5 Hz), 3.08-3.32 (m, 3H), 3.38-3.52 (m, 2H), 6.89 (s, 1H), 6.99-7.01 (m, 1H), 7.05-7.09 (m, 1H), 7.15-7.19 (m, 1H), 7.21-7.29 (m, 3H), 7.32-7.42 (m, 3H), 10.99 (s, 1H).	424	422	racemate




321		¹ H-NMR (DMSO-D6) δ : 1.13 (d, 6H, J = 5.6 Hz), 2.36-2.42 (m, 2H), 3.30-3.36 (m, 1H), 3.37-3.46 (m, 1H), 3.47-3.54 (m, 1H), 4.43 (quint, 1H, J = 5.6 Hz), 6.67-6.69 (m, 1H), 6.79-6.82 (m, 1H), 6.86-6.90 (m, 1H), 6.90 (s, 1H), 7.21-7.28 (m, 3H), 7.32-7.37 (m, 1H), 7.38-7.43 (m, 2H), 7.63 (s, 1H), 10.84 (s, 1H).	405	403	R-form
322		¹ H-NMR (CDCl ₃) δ : 0.83 (t, 3H, J = 7.6 Hz), 1.45-1.56 (m, 2H), 2.24 (s, 6H), 2.38 (dd, 1H, J = 16.8, 9.6 Hz), 2.50 (t, 2H, J = 7.6 Hz), 2.71 (dd, 1H, J = 16.8, 8.8 Hz), 3.05-3.15 (m, 1H), 3.40 (t, 1H, J = 8.8 Hz), 3.59 (dd, 1H, J = 9.6, 7.6 Hz), 6.29 (s, 1H), 6.85 (s, 2H), 6.94 (s, 1H), 7.01-7.14 (m, 4H), 7.20 (t, 1H, J = 7.6 Hz), 9.46 (s, 1H).	417	415	S-form
323		¹ H-NMR (CDCl ₃) δ : 0.77 (t, 3H, J = 7.2 Hz), 1.44 (tq, 2H, J = 7.2, 7.2 Hz), 1.79 (br s, 3H), 2.24 (s, 3H), 2.29-2.41 (m, 1H), 2.44 (t, 2H, J = 7.2 Hz), 2.71 (dd, 1H, J = 16.8, 8.8 Hz), 2.97-3.11 (m, 1H), 3.34-3.46 (m, 1H), 3.56-3.63 (m, 1H), 5.89 (s, 1H), 6.95-7.24 (m, 8H), 9.19 (br s, 1H).	417	415	S-form




324		¹ H-NMR (CDCl ₃) δ: 0.78 (t, 3H, J = 7.2 Hz), 1.46 (tq, 2H, J = 7.2, 7.2 Hz), 1.88 (s, 3H), 2.31 (s, 3H), 2.40-2.50 (m, 3H), 2.74 (dd, 1H, J = 16.8, 8.8 Hz), 3.10-3.21 (m, 1H), 3.50 (t, 1H, J = 9.2 Hz), 3.66 (dd, 1H, J = 9.2, 7.2 Hz), 5.72 (s, 1H), 6.98-7.17 (m, 8H), 8.70 (s, 1H).	417	415	S-form
325		¹ H-NMR (CDCl ₃) δ: 0.78 (t, 3H, J = 7.2 Hz), 1.45 (tq, 2H, J = 7.2, 7.2 Hz), 1.94 (s, 3H), 1.97 (s, 3H), 2.41-2.52 (m, 3H), 2.76 (dd, 1H, J = 17.2, 8.8 Hz), 3.11-3.21 (m, 1H), 3.51 (t, 1H, J = 9.2 Hz), 3.67 (dd, 1H, J = 9.2, 8.0 Hz), 5.73 (s, 1H), 6.93-7.26 (m, 8H), 8.70 (s, 1H).	417	415	S-form
326		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.5 Hz), 1.28 (s, 6H), 1.33 (tq, 2H, J = 7.5, 6.6 Hz), 2.38-2.40 (m, 2H), 2.89 (t, 2H, J = 6.6 Hz), 3.32-3.53 (m, 3H), 6.92 (s, 1H), 7.08 (s, 1H), 7.21-7.27 (m, 3H), 7.32-7.41 (m, 5H), 7.63 (br s, 1H), 10.85 (br s, 1H).	447	445	S-form




327		<p>1H-NMR (DMSO-D6) δ: 0.77 (t, 3H, J = 7.4 Hz), 1.05 (d, 3H, J = 7.1 Hz), 1.12 (d, 3H, J = 7.1 Hz), 1.44 (tq, 2H, J = 7.4, 7.5 Hz), 2.46 (t, 2H, J = 7.5 Hz), 3.16 (dd, 1H, J = 9.0, 6.6 Hz), 3.50 (dd, 1H, J = 9.0, 9.7 Hz), 3.88 (tt, 1H, J = 7.1, 7.1 Hz), 4.42 (dd, 1H, J = 9.7, 6.6 Hz), 6.39 (s, 1H), 6.91 (s, 1H), 7.02 (s, 1H), 7.06-7.08 (m, 1H), 7.16-7.18 (m, 1H), 7.22-7.29 (m, 3H), 7.32-7.42 (m, 3H), 11.04 (s, 1H).</p>	432	430	S-form
328		<p>1H-NMR (DMSO-D6) δ: 0.77 (t, 3H, J = 7.4 Hz), 1.05 (d, 3H, J = 7.1 Hz), 1.12 (d, 3H, J = 7.1 Hz), 1.44 (tq, 2H, J = 7.4, 7.5 Hz), 2.46 (t, 2H, J = 7.5 Hz), 3.16 (dd, 1H, J = 9.0, 6.6 Hz), 3.50 (dd, 1H, J = 9.0, 9.7 Hz), 3.88 (tt, 1H, J = 7.1, 7.1 Hz), 4.42 (dd, 1H, J = 9.7, 6.6 Hz), 6.39 (s, 1H), 6.91 (s, 1H), 7.02 (s, 1H), 7.06-7.08 (m, 1H), 7.16-7.18 (m, 1H), 7.22-7.29 (m, 3H), 7.32-7.42 (m, 3H), 11.04 (s, 1H).</p>	432	430	R-form




329		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 1.00 (t, 3H, J = 7.4 Hz), 1.44 (tq, 2H, J = 7.5, 7.4 Hz), 2.46 (t, 2H, J = 7.5 Hz), 3.09 (q, 2H, J = 7.4 Hz), 3.39 (dd, 1H, J = 9.2, 6.0 Hz), 3.62 (dd, 1H, J = 9.6, 9.2 Hz), 4.26 (dd, 1H, J = 9.6, 6.0 Hz), 6.72 (s, 1H), 6.90 (s, 1H), 7.00 (s, 1H), 7.07-7.09 (m, 1H), 7.16-7.18 (m, 1H), 7.22-7.29 (m, 3H), 7.32-7.42 (m, 3H), 10.76 (s, 1H).	418	416	S-form
330		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 1.00 (t, 3H, J = 7.4 Hz), 1.44 (tq, 2H, J = 7.5, 7.4 Hz), 2.46 (t, 2H, J = 7.5 Hz), 3.09 (q, 2H, J = 7.4 Hz), 3.39 (dd, 1H, J = 9.2, 6.0 Hz), 3.62 (dd, 1H, J = 9.6, 9.2 Hz), 4.26 (dd, 1H, J = 9.6, 6.0 Hz), 6.72 (s, 1H), 6.90 (s, 1H), 7.00 (s, 1H), 7.07-7.09 (m, 1H), 7.16-7.18 (m, 1H), 7.22-7.29 (m, 3H), 7.32-7.42 (m, 3H), 10.76 (s, 1H).	418	416	R-form
331		¹ H-NMR (DMSO-D ₆) δ: 0.93 (t, 3H, J = 6.9 Hz), 1.27 (s, 6H), 2.38-2.40 (m, 2H), 2.97 (q, 2H, J = 7.0 Hz), 3.32-3.53 (m, 3H), 6.92 (s, 1H), 7.06 (br s, 1H), 7.21-7.27 (m, 3H), 7.32-7.41 (m, 5H), 7.63 (br s, 1H), 10.86 (br s, 1H).	433	431	S-form




332		<p>¹H-NMR (DMSO-D₆) δ: 0.79 (t, 6H, J = 7.6 Hz), 1.39-1.46 (m, 4H), 2.39 (d, 2H, J = 8.4 Hz), 3.13-3.19 (m, 1H), 3.32-3.52 (m, 3H), 4.39 (s, 2H), 6.90 (s, 1H), 7.12 (d, 1H, J = 7.6 Hz), 7.22-7.24 (m, 3H), 7.31-7.41 (m, 5H), 7.64 (s, 1H), 10.86 (s, 1H).</p>	447	445	S-form
333		<p>¹H-NMR (DMSO-D₆) δ: 0.83 (s, 9H), 1.86-1.98 (m, 2H), 2.13-2.26 (m, 2H), 2.83-2.85 (m, 1H), 2.99 (s, 2H), 3.29-3.35 (m, 2H), 4.41 (s, 2H), 6.89 (s, 1H), 7.15-7.18 (m, 2H), 7.22-7.24 (m, 2H), 7.28-7.40 (m, 5H), 7.50 (s, 1H), 10.84 (s, 1H).</p>	461	459	Optically active substance (The same configuration as Example 5)
334		<p>¹H-NMR (DMSO-D₆) δ: 1.19 (d, 3H, J = 6.4 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.06-4.13 (m, 1H), 4.61 (s, 2H), 6.92 (s, 1H), 7.15 (d, 1H, J = 6.8 Hz), 7.23-7.25 (m, 3H), 7.32-7.41 (m, 5H), 7.64 (s, 1H), 10.87 (s, 1H).</p>	473	471	2,2,2-trifluoroethyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form Diastereoisomer of Example 335





335		<p>¹H-NMR (DMSO-D₆) δ: 1.19 (d, 3H, J = 6.4 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.06-4.13 (m, 1H), 4.61 (s, 2H), 6.92 (s, 1H), 7.15 (d, 1H, J = 6.8 Hz), 7.23-7.25 (m, 3H), 7.32-7.41 (m, 5H), 7.64 (s, 1H), 10.87 (s, 1H).</p>	473	471	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: S-form Lactam ring: S-form Diastereo isomer of Example 334
336		<p>¹H-NMR (DMSO-D₆) δ: 0.94 (s, 9H), 1.85-2.30 (m, 2H), 2.13-2.28 (m, 2H), 2.50-2.58 (m, 1H), 2.80-2.90 (m, 1H), 3.29-3.38 (m, 1H), 3.49 (s, 2H), 6.72-6.78 (m, 2H), 6.88-6.92 (m, 2H), 7.20-7.27 (m, 3H), 7.33-7.43 (m, 3H), 7.49 (s, 1H), 10.82 (s, 1H).</p>	447	445	Optically active substance (The same configuration as Example 5)
337		<p>¹H-NMR (DMSO-D₆) δ: 2.50-2.60 (m, 2H), 3.49-3.56 (m, 3H), 3.75 (dd, 1H, J = 9.2 Hz, 9.2 Hz), 4.45 (s, 2H), 5.13 (dd, 1H, J = 9.2 Hz, 5.8 Hz), 6.91 (s, 1H), 7.12-7.15 (m, 1H), 7.22-7.27 (m, 3H), 7.29-7.43 (m, 5H), 7.75 (s, 1H), 11.06 (s, 1H).</p>	475	473	R-form





338		¹ H-NMR (DMSO-D ₆) δ: 1.85-2.30 (m, 2H), 2.13-2.28 (m, 2H), 2.50-2.58 (m, 3H), 2.80-2.90 (m, 1H), 3.29-3.38 (m, 1H), 3.54 (t, 2H, J = 6.0 Hz), 4.45 (s, 2H), 6.90 (s, 1H), 7.12 (d, 1H, J = 7.6 Hz), 7.20-7.23 (m, 3H), 7.28-7.42 (m, 5H), 7.49 (s, 1H), 10.83 (s, 1H).	487	485	Optically active substance (The same configuration as Example 5)
339		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 1.43 (tq, 2H, J = 7.4 Hz, 7.4Hz), 2.46 (t, 2H, J = 7.4 Hz), 3.07 (m, 1H), 3.35-3.46 (m, 2H), 4.28-4.41 (m, 2H), 6.87 (s, 1H), 7.00 (s, 1H), 7.07 (d, 1H, J = 8.0 Hz), 7.17 (d, 1H, J = 8.0 Hz), 7.20-7.42 (m, 7H), 11.01 (s, 1H).	405	403	racemate
340		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, J = 7.3 Hz), 1.41-1.51 (m, 2H), 2.36-2.41 (m, 2H), 2.45-2.52 (m, 2H), 3.32-3.53 (m, 3H), 4.98 (s, 2H), 6.76-6.80 (m, 1H), 6.86 (t, 1H, J = 2.0 Hz), 6.88 (s, 1H), 6.98 (dd, 1H, J = 8.3, 2.0 Hz), 7.01-7.04 (m, 1H), 7.07-7.11 (m, 1H), 7.17-7.21 (m, 1H), 7.26-7.41 (m, 7H), 7.63 (s, 1H), 10.86 (s, 1H).	495	493	S-form





341		¹ H-NMR (DMSO-D ₆) δ: 0.79 (t, 3H, J = 7.5 Hz), 1.42-1.52 (m, 2H), 2.36-2.41 (m, 2H), 2.45-2.51 (m, 2H), 3.31-3.53 (m, 3H), 6.53-6.58 (m, 1H), 6.64 (t, 1H, J = 2.0 Hz), 6.68-6.73 (m, 1H), 6.86 (s, 1H), 7.01-7.04 (m, 1H), 7.06-7.18 (m, 3H), 7.27 (t, 1H, J = 7.7 Hz), 7.63 (s, 1H), 9.88 (br s, 1H), 10.82 (s, 1H).	405	403	S-form
342		¹ H-NMR (DMSO-D ₆) δ: 0.90 (s, 9H), 1.54 (t, 2H, J = 7.3 Hz), 1.83-2.03 (m, 2H), 2.12-2.30 (m, 2H), 2.79-2.87 (m, 1H), 3.25-3.35 (m, 2H), 3.85 (t, 2H, J = 7.3 Hz), 6.70 (t, 1H, J = 2.0 Hz), 6.80-6.84 (m, 1H), 6.87-6.92 (m, 2H), 7.23-7.28 (m, 3H), 7.32-7.44 (m, 3H), 7.50 (br s, 1H), 10.82 (s, 1H).	461	459	Optically active substance (The same configuration as Example 5)
343		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, J = 7.5 Hz), 1.41-1.52 (m, 2H), 2.36-2.41 (m, 2H), 2.45-2.53 (m, 2H), 3.31-3.53 (m, 3H), 3.65 (s, 3H), 6.73-6.77 (m, 1H), 6.79 (t, 1H, J = 2.2 Hz), 6.87-6.92 (m, 2H), 7.02-7.04 (m, 1H), 7.07-7.11 (m, 1H), 7.16-7.20 (m, 1H), 7.24-7.31 (m, 2H), 7.64 (s, 1H), 10.87 (s, 1H).	419	417	S-form




344		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, J = 7.3 Hz), 1.13 (d, 6H, J = 6.0 Hz), 1.40-1.50 (m, 2H), 2.36-2.41 (m, 2H), 2.44-2.53 (m, 2H), 3.31-3.53 (m, 2H), 4.38-4.46 (m, 3H), 6.68 (t, 1H, J = 2.2 Hz), 6.78-6.82 (m, 1H), 6.84-6.88 (m, 2H), 7.01-7.03 (m, 1H), 7.07-7.11 (m, 1H), 7.15-7.20 (m, 1H), 7.24-7.31 (m, 1H), 7.63 (s, 1H), 10.84 (s, 1H).	447	445	S-form
345		¹ H-NMR (CDCl ₃) δ: 0.84 (t, 3H, J = 7.3 Hz), 1.50 (tq, 2H, J = 7.3, 7.5 Hz), 2.49 (t, 2H, J = 7.5 Hz), 3.27-3.33 (m, 1H), 3.52 (s, 3H), 3.54-3.65 (m, 3H), 3.85-3.94 (m, 2H), 4.37 (dd, 1H, J = 10.3, 7.4 Hz), 4.51 (s, 1H), 7.03-7.14 (m, 5H), 7.19-7.33 (m, 5H), 10.07 (s, 1H).	448	446	S-form
346		¹ H-NMR (CDCl ₃) δ: 0.84 (t, 3H, J = 7.3 Hz), 1.50 (tq, 2H, J = 7.3, 7.5 Hz), 2.49 (t, 2H, J = 7.5 Hz), 3.27-3.33 (m, 1H), 3.52 (s, 3H), 3.54-3.65 (m, 3H), 3.85-3.94 (m, 2H), 4.37 (dd, 1H, J = 10.3, 7.4 Hz), 4.51 (s, 1H), 7.03-7.14 (m, 5H), 7.19-7.33 (m, 5H), 10.07 (s, 1H).	448	446	R-form




347		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.3, 7.3 Hz), 2.45 (t, 2H, J = 7.3 Hz), 3.23-3.27 (m, 1H), 3.50-3.55 (m, 1H), 3.60-3.63 (m, 1H), 3.90-3.96 (m, 1H), 4.48-4.53 (m, 1H), 6.58 (s, 1H), 6.89 (s, 1H), 6.99-7.02 (m, 1H), 7.06-7.08 (m, 1H), 7.16-7.18 (m, 1H), 7.21-7.29 (m, 1H), 7.33-7.42 (m, 3H), 12.77 (br s, 1H).	448	446	S-form
348		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.3, 7.3 Hz), 2.45 (t, 2H, J = 7.3 Hz), 3.23-3.27 (m, 1H), 3.50-3.55 (m, 1H), 3.60-3.63 (m, 1H), 3.90-3.96 (m, 1H), 4.48-4.53 (m, 1H), 6.58 (s, 1H), 6.89 (s, 1H), 6.99-7.02 (m, 1H), 7.06-7.08 (m, 1H), 7.16-7.18 (m, 1H), 7.21-7.29 (m, 1H), 7.33-7.42 (m, 3H), 12.77 (br s, 1H).	448	446	R-form
349		¹ H-NMR (DMSO-D ₆) δ: 3.33-3.38 (m, 1H), 3.60 (dd, 1H, J = 10.1, 8.3 Hz), 3.97 (q, 2H, J = 9.2 Hz), 4.32 (dd, 1H, J = 8.8, 5.6 Hz), 4.61 (s, 2H), 6.33 (s, 1H), 6.59 (s, 1H), 6.93 (s, 1H), 7.16-7.17 (m, 1H), 7.22-7.26 (m, 3H), 7.34-7.40 (m, 5H), 10.71 (s, 1H).	460	458	R-form




350		¹ H-NMR (DMSO-D ₆) δ: 0.80 (t, 3H, J = 7.6 Hz), 1.03 (d, 3H, J = 6.4 Hz), 1.32-1.50 (m, 2H), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.35 (d, 1H, J = 12.8 Hz), 4.44 (d, 1H, J = 12.8 Hz), 6.90 (s, 1H), 7.11 (d, 1H, J = 6.8 Hz), 7.21-7.24 (m, 3H), 7.28-7.41 (m, 5H), 7.63 (s, 1H), 10.86 (s, 1H).	433	431	sec-butoxymethyl group: R-form Lactam ring: S-form Diastereo isomer of Example 351
351		¹ H-NMR (DMSO-D ₆) δ: 0.80 (t, 3H, J = 7.6 Hz), 1.03 (d, 3H, J = 6.4 Hz), 1.32-1.50 (m, 2H), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.35 (d, 1H, J = 12.8 Hz), 4.44 (d, 1H, J = 12.8 Hz), 6.90 (s, 1H), 7.11 (d, 1H, J = 6.8 Hz), 7.21-7.24 (m, 3H), 7.28-7.41 (m, 5H), 7.63 (s, 1H), 10.86 (s, 1H).	433	431	sec-butoxymethyl group: S-form Lactam ring: S-form Diastereo isomer of Example 350
352		¹ H-NMR (DMSO-D ₆) δ: 1.34 (s, 6H), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.53 (s, 2H), 6.91 (s, 1H), 7.13 (d, 1H, J = 7.6 Hz), 7.22-7.25 (m, 3H), 7.28-7.42 (m, 5H), 7.63 (s, 1H), 10.86 (s, 1H).	487	485	S-form
353		¹ H-NMR (DMSO-D ₆) δ: 1.72 (s, 3H), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.78 (s, 2H), 6.92 (s, 1H), 7.17 (d, 1H, J = 7.6 Hz), 7.22-7.26 (m, 3H), 7.31-7.41 (m, 5H), 7.63 (s, 1H), 10.87 (s, 1H).	541	539	S-form




354		<p>¹H-NMR (DMSO-D₆) δ: 2.47-2.57 (m, 2H), 3.34-3.38 (m, 1H), 3.54 (t, 2H, J = 6.0 Hz), 3.59 (t, 1H, 9.2 Hz), 4.30-4.34 (m, 1H), 4.45 (s, 2H), 6.33 (s, 1H), 6.60 (s, 1H), 6.92 (s, 1H), 7.12 (d, 1H, J = 7.6 Hz), 7.22-7.25 (m, 3H), 7.29-7.41 (m, 5H), 10.71 (s, 1H).</p>	474	472	R-form
355		<p>¹H-NMR (DMSO-D₆) δ: 0.67 (s, 9H), 2.37-2.41 (m, 2H), 3.30-3.52 (m, 3H), 4.12 (d, 1H, J = 4.0 Hz), 5.13 (d, 1H, J = 4.0 Hz), 6.86 (s, 1H), 7.07-7.10 (m, 1H), 7.14-7.18 (m, 1H), 7.20-7.40 (m, 7H), 7.63 (s, 1H), 10.85 (s, 1H).</p>	433	431	1-hydroxy-2,2-dimethyl-propyl group: racemate Lactam ring: S-form
356		<p>¹H-NMR (DMSO-D₆) δ: 1.07 (s, 9H), 2.38-2.41 (m, 2H), 3.31-3.54 (m, 3H), 6.97 (s, 1H), 7.24-7.28 (m, 2H), 7.32-7.66 (m, 8H), 10.88 (s, 1H).</p>	431	429	S-form
357		<p>¹H-NMR (DMSO-D₆) δ: 1.06 (s, 6H), 2.37-2.42 (m, 2H), 3.13 (s, 3H), 3.16 (s, 2H), 3.31-3.54 (m, 3H), 6.92 (s, 1H), 7.04-7.06 (m, 1H), 7.17-7.25 (m, 3H), 7.28-7.44 (m, 5H), 7.63 (s, 1H), 10.84 (s, 1H).</p>	433	431	S-form




358		¹ H-NMR (CDCl ₃) δ: 0.78 (t, 3H, J = 7.2 Hz), 0.97 (t, 3H, J = 7.6 Hz), 1.44 (tq, 2H, J = 7.2, 7.2 Hz), 2.33 (q, 2H, J = 7.6 Hz), 2.41-2.53(m, 3H), 2.75 (dd, 1H, J = 16.8, 8.4 Hz), 3.16-3.26 (m, 1H), 3.53 (t, 1H, J = 9.2 Hz), 3.68 (dd, 1H, J = 9.6, 7.6Hz), 5.69 (s, 1H), 6.97-7.40 (m, 9H), 8.54 (s, 1H).	417	415	S-form
359		¹ H-NMR (CDCl ₃) δ: 0.65-1.20 (m, 9H), 1.43 (tq, 2H, J = 7.2, 7.2 Hz), 2.35-2.54 (m, 3H), 2.60-2.69(m, 1H), 2.74 (dd, 1H, J = 17.2, 8.8 Hz), 3.12-3.25 (m, 1H), 3.44-3.54 (m, 1H), 3.61-3.70 (m, 1H), 5.80 (s, 1H), 6.95-7.44 (m, 9H), 8.76 (br s, 1H).	431	429	S-form
360		¹ H-NMR (CDCl ₃) δ: 0.76-0.87 (m, 4H), 0.92-1.00 (m, 1H), 1.12-1.27 (m, 2H), 1.51 (tq, 2H, J = 7.6, 7.6 Hz), 2.49 (t, 2H, J = 7.6 Hz), 3.03 (dd, 1H, J = 8.4, 5.2 Hz), 3.46 (dd, 1H, J = 9.6, 8.4 Hz), 3.80 (dd, 1H, J = 9.6, 5.2 Hz), 6.46 (s, 1H), 7.03-7.37 (m, 10H), 9.29 (s, 1H).	415	413	Optically active substance Optical isomer of Example 361(The same configuration as Example 250)
361		¹ H-NMR (CDCl ₃) δ: 0.76-0.87 (m, 4H), 0.92-1.00 (m, 1H), 1.12-1.27 (m, 2H), 1.51 (tq, 2H, J = 7.6, 7.6 Hz), 2.49 (t, 2H, J = 7.6 Hz), 3.03 (dd, 1H, J = 8.4, 5.2 Hz), 3.46 (dd, 1H, J = 9.6, 8.4 Hz), 3.80 (dd, 1H, J = 9.6, 5.2 Hz), 6.46 (s, 1H), 7.03-7.37 (m, 10H), 9.29 (s, 1H).	415	413	Optically active substance Optical isomer of Example 360




362		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 1.45 (tq, 2H, J = 7.4 Hz, 7.4Hz), 2.35-2.40 (m, 2H), 2.46 (t, 2H, J = 7.4 Hz), 3.30-3.53 (m, 3H), 3.76 (s, 3H), 6.86 (s, 1H), 6.92-6.96 (m, 2H), 7.00 (s, 1H), 7.05 (d, 1H, J = 7.6 Hz), 7.12-7.18 (m, 3H), 7.25 (dd, 1H, J = 7.6 Hz, 7.6 Hz), 7.63 (s, 1H), 10.80 (s, 1H).	419	417	S-form
363		¹ H-NMR (DMSO-D ₆) δ: 0.74 (t, 3H, J = 7.4 Hz), 1.19 (d, 6H, J = 6.8 Hz), 1.37 (tq, 2H, J = 7.4 Hz, 7.4Hz), 2.36-2.44 (m, 4H), 2.91 (sep, 1H, J = 6.8 Hz), 3.30-3.53 (m, 3H), 6.88 (s, 2H), 7.10-7.18 (m, 4H), 7.23-7.31 (m, 3H), 7.62 (s, 1H), 10.81 (s, 1H).	431	429	S-form
364		¹ H-NMR (DMSO-D ₆) δ: 1.72 (s, 3H), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.78 (s, 2H), 6.92 (s, 1H), 7.17 (d, 1H, J = 7.6 Hz), 7.22-7.26 (m, 3H), 7.31-7.41 (m, 5H), 7.63 (s, 1H), 10.87 (s, 1H).	541	539	R-form




365		<p>¹H-NMR (DMSO-D₆) δ: 1.19 (d, 3H, J = 6.4 Hz), 3.34-3.38 (m, 1H), 3.60 (t, 1H, 9.2 Hz), 4.05-4.13 (m, 1H), 4.31-4.34 (m, 1H), 4.62 (s, 2H), 6.34 (s, 1H), 6.60 (s, 1H), 6.93 (s, 1H), 7.15 (d, 1H, J = 6.8 Hz), 7.23-7.25 (m, 3H), 7.32-7.42 (m, 5H), 10.72 (s, 1H).</p>	474	472	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Alpha position of carbonyl group (amide): R-form
366		<p>¹H-NMR (DMSO-D₆) δ: 3.50-3.54 (m, 1H), 3.76 (t, 1H, J = 9.2 Hz), 4.86 (s, 2H), 5.11-5.15 (m, 1H), 5.52-5.59 (m, 1H), 6.93 (s, 1H), 7.16-7.18 (m, 1H), 7.24-7.26 (m, 2H), 7.34-7.42 (m, 6H), 7.76 (s, 1H), 11.08 (s, 1H).</p>	529	527	R-form
367		<p>¹H-NMR (DMSO-D₆) δ: 1.19 (d, 3H, J = 6.4 Hz), 3.50-3.54 (m, 1H), 3.75 (t, 1H, J = 9.2 Hz), 4.06-4.13 (m, 1H), 4.62 (s, 2H), 5.11-5.15 (m, 1H), 6.92 (s, 1H), 7.15 (d, 1H, J = 7.6 Hz), 7.24-7.26 (m, 3H), 7.32-7.42 (m, 5H), 7.75 (s, 1H), 11.06 (s, 1H).</p>	475	473	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Alpha position of carbonyl group (amide): R-form




368		<p>¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.4, 7.3 Hz), 2.46 (t, 2H, J = 7.4 Hz), 4.27-4.35 (m, 1H), 4.47-4.54 (m, 2H), 6.91 (s, 1H), 7.00 (s, 1H), 7.08 (d, 1H, J = 7.7 Hz), 7.18 (d, 1H, J = 7.7 Hz), 7.22-7.29 (m, 3H), 7.34-7.42 (m, 3H), 8.04 (br s, 1H), 10.98 (br s, 1H).</p>	391	389	S-form
369		<p>¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.4, 7.3 Hz), 2.46 (t, 2H, J = 7.4 Hz), 4.27-4.34 (m, 1H), 4.47-4.54 (m, 2H), 6.91 (s, 1H), 7.00 (s, 1H), 7.08 (d, 1H, J = 7.9 Hz), 7.18 (d, 1H, J = 7.7 Hz), 7.22-7.29 (m, 3H), 7.33-7.42 (m, 3H), 8.04 (br s, 1H), 10.99 (br s, 1H).</p>	391	389	R-form
370		<p>¹H-NMR (DMSO-D₆) δ: 0.76 (t, 3H, J = 7.3 Hz), 1.43 (tq, 2H, J = 7.3, 7.3 Hz), 2.35-2.39 (m, 2H), 2.44 (t, 2H, J = 7.3 Hz), 3.32-3.35 (m, 1H), 3.39-3.42 (m, 1H), 3.46-3.50 (m, 1H), 5.08 (s, 2H), 6.85 (s, 1H), 6.98-7.05 (m, 4H), 7.13-7.15 (m, 3H), 7.22-7.26 (m, 1H), 7.30-7.44 (m, 5H), 7.61 (s, 1H), 10.77 (s, 1H).</p>	495	493	S-form




371		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.45 (tq, 2H, J = 7.3, 7.4 Hz), 2.35-2.37 (m, 2H), 2.45 (t, 2H, J = 7.4 Hz), 3.33-3.35 (m, 1H), 3.37-3.43 (m, 1H), 3.46-3.50 (m, 1H), 6.71-6.73 (m, 2H), 6.83 (s, 1H), 6.98-7.04 (m, 4H), 7.11-7.13 (m, 1H), 7.23 (t, 1H, J = 7.7 Hz), 7.61 (s, 1H), 9.72 (br s, 1H), 10.74 (s, 1H).	405	403	S-form
372		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, J = 7.3 Hz), 1.32 (t, 3H, J = 6.9 Hz), 1.45 (tq, 2H, J = 14.9, 7.3 Hz), 2.38-2.40 (m, 2H), 2.46 (t, 2H, J = 7.7 Hz), 3.33-3.36 (m, 1H), 3.38-3.44 (m, 1H), 3.48-3.52 (m, 1H), 4.02 (q, 2H, J = 6.9 Hz), 6.86 (s, 1H), 6.91-6.93 (m, 2H), 6.99-7.01 (m, 1H), 7.04-7.06 (m, 1H), 7.12-7.16 (m, 3H), 7.24-7.26 (m, 1H), 7.62 (s, 1H), 10.79 (s, 1H).	433	431	S-form
373		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.26 (d, 6H, J = 6.0 Hz), 1.42 (tq, 2H, J = 7.4, 7.3 Hz), 2.37-2.40 (m, 2H), 2.45 (t, 2H, J = 7.4 Hz), 3.32-3.35 (m, 1H), 3.40-3.44 (m, 1H), 3.48-3.52 (m, 1H), 4.60 (sep, 1H, J = 6.0 Hz), 6.85-6.97 (m, 4H), 7.08-7.15 (m, 4H), 7.27 (t, 1H, J = 7.6 Hz), 7.63 (s, 1H), 10.78 (s, 1H).	447	445	S-form



374		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.3, 7.2 Hz), 2.38-2.40 (m, 2H), 2.46 (t, 2H, J = 7.2 Hz), 2.74-2.82 (m, 2H), 3.32-3.35 (m, 1H), 3.39-3.42 (m, 1H), 3.48-3.51 (m, 1H), 4.20 (t, 2H, J = 5.8 Hz), 6.87 (s, 1H), 6.96-6.99 (m, 3H), 7.06-7.08 (m, 1H), 7.14-7.17 (m, 3H), 7.26 (t, 1H, J = 7.6 Hz), 7.63 (s, 1H), 10.80 (s, 1H).	501	499	S-form
375		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, J = 7.3 Hz), 1.46 (tq, 2H, J = 7.7, 7.3 Hz), 2.37-2.39 (m, 2H), 2.46 (t, 2H, J = 7.7 Hz), 3.32-3.34 (m, 1H), 3.39-3.43 (m, 1H), 3.49-3.52 (m, 1H), 4.78 (q, 2H, J = 8.9 Hz), 6.87 (s, 1H), 7.00-7.09 (m, 4H), 7.15-7.28 (m, 4H), 7.63 (s, 1H), 10.81 (s, 1H).	487	485	S-form
376		¹ H-NMR (CDCl ₃) δ: 0.80 (t, 3H, J = 7.2 Hz), 1.46 (tq, 2H, J = 7.2, 7.2 Hz), 2.15-2.33 (m, 1H), 2.46 (t, 2H, J = 7.2 Hz), 2.67 (dd, 1H, J = 16.8, 8.8 Hz), 2.90-3.04 (m, 1H), 3.31 (t, 1H, J = 8.8 Hz), 3.40 (s, 3H), 3.52 (t, 1H, J = 8.8 Hz), 6.03 (s, 1H), 6.87 (d, 1H, J = 8.4 Hz), 6.99-7.08 (m, 5H), 7.11-7.17 (m, 1H), 7.33-7.44 (m, 2H), 9.76 (s, 1H).	419	417	S-form




377		¹ H-NMR (CDCl ₃) δ: 0.80 (t, 3H, J = 7.6 Hz), 0.93 (t, 3H, J = 6.8 Hz), 1.46 (tq, 2H, J = 7.6, 7.6 Hz), 2.01-2.33 (m, 1H), 2.45 (t, 2H, J = 7.6 Hz), 2.65 (dd, 1H, J = 16.8, 8.8 Hz), 2.85-3.01 (m, 1H), 3.13-3.87 (m, 4H), 6.00 (s, 1H), 6.83 (d, 1H, J = 8.4 Hz), 6.99-7.16 (m, 6H), 7.34 (t, 1H, J = 7.2 Hz), 7.47 (d, 1H, J = 7.6 Hz), 9.99 (br s, 1H).	433	431	S-form
378		¹ H-NMR (CDCl ₃) δ: 0.38-1.18 (m, 9H), 1.46 (tq, 2H, J = 7.6, 7.6 Hz), 2.45 (t, 2H, J = 7.6 Hz), 2.66 (dd, 1H, J = 16.4, 8.8 Hz), 2.81-3.41 (m, 2H), 3.50 (t, 1H, J = 9.2 Hz), 4.20-4.31 (m, 1H), 6.11 (s, 1H), 6.80 (d, 1H, J = 8.0 Hz), 6.98-7.16 (m, 6H), 7.28-7.55 (m, 1H), 10.16 (br s, 1H).	447	445	S-form
379		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, J = 7.3 Hz), 1.23 (t, 3H, J = 7.1 Hz), 1.40-1.51 (m, 2H), 2.37-2.41 (m, 2H), 2.44-2.51 (m, 2H), 3.31-3.53 (m, 3H), 3.89 (q, 2H, J = 7.1 Hz), 6.72-6.79 (m, 2H), 6.85-6.90 (m, 2H), 7.01-7.04 (m, 1H), 7.07-7.11 (m, 1H), 7.16-7.20 (m, 1H), 7.24-7.31 (m, 2H), 7.63 (s, 1H), 10.86 (s, 1H).	433	431	S-form




380		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, J = 7.5 Hz), 0.90 (t, 3H, J = 7.5 Hz), 1.41-1.51 (m, 2H), 1.56-1.67 (m, 2H), 2.37-2.42 (m, 2H), 2.45-2.51 (m, 2H), 3.31-3.53 (m, 2H), 3.78 (t, 2H, J = 6.4 Hz), 6.73 (t, 1H, J = 2.2 Hz), 6.76-6.80 (m, 1H), 6.85-6.90 (m, 2H), 7.02-7.04 (m, 1H), 7.07-7.11 (m, 1H), 7.16-7.20 (m, 1H), 7.24-7.31 (m, 1H), 7.63 (s, 1H), 10.86 (s, 1H).	447	445	S-form
381		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, J = 7.3 Hz), 0.89 (d, 6H, J = 6.9 Hz), 1.41-1.51 (m, 2H), 1.83-1.94 (m, 1H), 2.37-2.41 (m, 2H), 2.45-2.52 (m, 2H), 3.31-3.54 (m, 2H), 3.58 (d, 2H, J = 6.9 Hz), 6.72 (t, 1H, J = 2.0 Hz), 6.76-6.81 (m, 1H), 6.85-6.90 (m, 2H), 7.02-7.04 (m, 1H), 7.07-7.12 (m, 1H), 7.17-7.21 (m, 1H), 7.24-7.31 (m, 2H), 7.63 (s, 1H), 10.86 (s, 1H).	461	459	S-form
382		¹ H-NMR (DMSO-D ₆) δ: 0.79 (t, 3H, J = 7.3 Hz), 1.40-1.76 (m, 10H), 2.36-2.41 (m, 2H), 2.44-2.52 (m, 2H), 3.29-3.53 (m, 3H), 4.57-4.63 (m, 1H), 6.64 (t, 1H, J = 2.2 Hz), 6.81-6.86 (m, 2H), 6.88 (s, 1H), 7.01-7.03 (m, 1H), 7.07-7.11 (m, 1H), 7.16-7.20 (m, 1H), 7.24-7.31 (m, 2H), 7.63 (s, 1H), 10.84 (s, 1H).	473	471	S-form




383		<p>¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 0.97 (t, 3H, J = 7.3 Hz), 1.45 (tq, 2H, J = 7.3, 7.2 Hz), 1.72 (tq, 2H, J = 6.9, 7.3 Hz), 2.38-2.39 (m, 2H), 2.45-2.46 (m, 5H), 3.32-3.35 (m, 1H), 3.39-3.39 (m, 1H), 3.48-3.50 (m, 1H), 3.92 (t, 2H, J = 6.5 Hz), 6.86 (s, 1H), 6.92-6.94 (m, 2H), 6.99-7.00 (m, 1H), 7.05-7.07 (m, 1H), 7.12-7.17 (m, 3H), 7.25 (t, 1H, J = 7.6 Hz), 7.62 (s, 1H), 10.79 (s, 1H).</p>	447	445	S-form
384		<p>¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.3, 7.2 Hz), 2.47 (t, 2H, J = 7.2 Hz), 3.32-3.34 (m, 1H), 3.55-3.67 (m, 2H), 4.15-4.23 (m, 1H), 4.51-4.55 (m, 1H), 6.90 (s, 1H), 6.99-7.01 (m, 2H), 7.06-7.08 (m, 1H), 7.16-7.18 (m, 1H), 7.21-7.29 (m, 3H), 7.34-7.42 (m, 3H), 11.10 (s, 1H).</p>	472	470	S-form
385		<p>¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.44 (tq, 2H, J = 7.3, 7.2 Hz), 2.47 (t, 2H, J = 7.2 Hz), 3.32-3.34 (m, 1H), 3.55-3.67 (m, 2H), 4.15-4.23 (m, 1H), 4.51-4.55 (m, 1H), 6.90 (s, 1H), 6.99-7.01 (m, 2H), 7.06-7.08 (m, 1H), 7.16-7.18 (m, 1H), 7.21-7.29 (m, 3H), 7.34-7.42 (m, 3H), 11.10 (s, 1H).</p>	472	470	R-form




386		¹ H-NMR (DMSO-D ₆) δ: 0.75 (t, 3H, J = 7.3 Hz), 1.41 (tq, 2H, J = 7.3, 7.7 Hz), 2.38-2.40 (m, 2H), 2.45 (t, 2H, J = 7.7 Hz), 3.33-3.35 (m, 1H), 3.40-3.45 (m, 1H), 3.49-3.52 (m, 1H), 6.92-6.94 (m, 2H), 7.15-7.20 (m, 2H), 7.30-7.36 (m, 3H), 7.40-7.42 (m, 2H), 7.63 (s, 1H), 10.87 (s, 1H).	473	471	S-form
387		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, J = 7.5 Hz), 1.41-1.51 (m, 2H), 2.37-2.41 (m, 2H), 2.45-2.52 (m, 2H), 3.30-3.53 (m, 3H), 4.68 (q, 2H, J = 8.9 Hz), 6.83-6.87 (m, 1H), 6.90 (s, 1H), 6.96 (t, 1H, J = 2.0 Hz), 7.02-7.06 (m, 2H), 7.07-7.11 (m, 1H), 7.17-7.21 (m, 1H), 7.26-7.36 (m, 2H), 7.63 (s, 1H), 10.86 (s, 1H).	487	485	S-form
388		¹ H-NMR (DMSO-D ₆) δ: 0.79 (t, 3H, J = 7.5 Hz), 1.43-1.51 (m, 1H), 2.37-2.41 (m, 2H), 2.45-2.52 (m, 2H), 3.31-3.54 (m, 3H), 6.90 (s, 1H), 7.02-7.12 (m, 4H), 7.15-7.21 (m, 2H), 7.29 (t, 1H, J = 7.7 Hz), 7.40 (t, 1H, J = 8.1 Hz), 7.64 (s, 1H), 10.89 (s, 1H).	505	503	S-form




389		¹ H-NMR (CDCl ₃) δ: 0.86 (t, 3H, J = 7.6 Hz), 1.54 (tq, 2H, J = 7.6, 7.6 Hz), 2.52 (t, 2H, J = 7.6 Hz), 2.60 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.4 Hz), 3.30-3.40 (m, 1H), 3.59 (t, 1H, J = 9.2 Hz), 3.67-3.75 (m, 4H), 6.00 (s, 1H), 6.73-6.85 (m, 2H), 6.98-7.08 (m, 4H), 7.12-7.25 (m, 2H), 8.48 (s, 1H).	437	435	S-form
390		¹ H-NMR (CDCl ₃) δ: 0.86 (t, 3H, J = 7.2 Hz), 1.34 (t, 3H, J = 7.2 Hz), 1.54 (tq, 2H, J = 7.2, 7.2 Hz), 2.52 (t, 2H, J = 7.2 Hz), 2.59 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.4 Hz), 3.29-3.41 (m, 1H), 3.62 (t, 1H, J = 8.8 Hz), 3.74 (dd, 1H, J = 9.6, 7.6 Hz), 3.89 (q, 2H, J = 7.2 Hz), 5.71 (s, 1H), 6.73-6.83 (m, 2H), 6.98-7.08 (m, 4H), 7.12-7.25 (m, 2H), 8.17 (s, 1H).	451	449	S-form



391		¹ H-NMR (CDCl ₃) δ: 0.86 (t, 3H, J = 7.6 Hz), 1.18 (d, 3H, J = 6.0 Hz), 1.20 (d, 3H, J = 6.0 Hz), 1.54 (tq, 2H, J = 7.6, 7.6 Hz), 2.51 (t, 2H, J = 7.6 Hz), 2.59 (dd, 1H, J = 17.2, 10.0 Hz), 2.77 (dd, 1H, J = 17.2, 8.8 Hz), 3.28-3.38 (m, 1H), 3.57 (t, 1H, J = 8.8 Hz), 3.70 (dd, 1H, J = 9.6, 7.2 Hz), 4.22-4.32 (m, 1H), 6.13 (s, 1H), 6.73-6.86 (m, 2H), 6.99-7.09 (m, 4H), 7.13 (d, 1H, J = 7.6 Hz), 7.22 (t, 1H, J = 7.6 Hz), 8.67 (s, 1H).	465	463	S-form
392		¹ H-NMR (CDCl ₃) δ: 0.84 (t, 3H, J = 7.6 Hz), 1.10 (d, 6H, J = 7.2 Hz), 1.50 (tq, 2H, J = 7.6, 7.6 Hz), 2.38-2.52 (m, 3H), 2.67-2.87 (m, 2H), 3.07-3.19 (m, 1H), 3.44 (t, 1H, J = 8.8 Hz), 3.62 (dd, 1H, J = 9.6, 7.6 Hz), 4.22-4.32 (m, 1H), 6.07 (s, 1H), 7.00-7.30 (m, 9H), 9.19 (s, 1H).	431	429	S-form
393		¹ H-NMR (CDCl ₃) δ: 0.83 (t, 3H, J = 7.2 Hz), 1.11 (t, 3H, J = 7.6 Hz), 1.50 (tq, 2H, J = 7.2, 7.2 Hz), 2.36-2.76 (m, 6H), 3.07-3.17 (m, 1H), 3.38-3.66 (m, 2H), 6.18 (s, 1H), 7.01-7.25 (m, 9H), 9.33 (s, 1H).	417	415	S-form




394		¹ H-NMR (CDCl ₃) δ: 0.84 (t, 3H, J = 7.6 Hz), 1.15 (s, 9H), 1.51 (tq, 2H, J = 7.6, 7.6 Hz), 2.42-2.53 (m, 3H), 2.73 (dd, 1H, J = 17.2, 8.4 Hz), 3.13-3.24 (m, 1H), 3.50 (t, 1H, J = 8.8 Hz), 3.65 (dd, 1H, J = 9.6, 7.6 Hz), 5.88 (s, 1H), 7.00-7.33 (m, 9H), 8.89 (s, 1H).	445	443	S-form
395		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, 7.2 Hz), 1.17 (d, 3H, 6.4 Hz), 1.32-1.41 (m, 2H), 2.39 (d, 2H, J = 8.4 Hz), 2.95-3.00 (m, 1H), 3.06-3.11 (m, 1H), 3.32-3.52 (m, 3H), 4.29 (q, 1H, J = 6.4 Hz), 6.92 (s, 1H), 7.04 (s, 1H), 7.21-7.41 (m, 8H), 7.64 (s, 1H), 10.86 (s, 1H).	433	431	1-propoxy-ethyl group: R-form Lactam ring: S-form Diastereo isomer of Example 396
396		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, 7.2 Hz), 1.17 (d, 3H, 6.4 Hz), 1.32-1.41 (m, 2H), 2.39 (d, 2H, J = 8.4 Hz), 2.95-3.00 (m, 1H), 3.06-3.11 (m, 1H), 3.32-3.52 (m, 3H), 4.29 (q, 1H, J = 6.4 Hz), 6.92 (s, 1H), 7.04 (s, 1H), 7.21-7.41 (m, 8H), 7.64 (s, 1H), 10.86 (s, 1H).	433	431	1-propoxy-ethyl group: S-form Lactam ring: S-form Diastereo isomer of Example 395




397		¹ H-NMR (CDCl ₃) δ: 0.78 (t, 3H, J = 7.2 Hz), 0.95 (s, 3H), 1.17 (s, 3H), 1.44 (tq, 2H, J = 7.2, 7.6 Hz), 2.46 (t, 2H, J = 7.6 Hz), 3.13 (t, 1H, J = 7.2 Hz), 3.30-3.37 (m, 1H), 3.41-3.45 (m, 1H), 6.91 (s, 1H), 7.02 (s, 1H), 7.06 (d, 1H, J = 7.6 Hz), 7.16 (d, 1H, J = 8.0 Hz), 7.22-7.28 (m, 3H), 7.32-7.41 (m, 3H), 7.64 (s, 1H), 10.79 (s, 1H).	417	415	Optically active substance (Structure Information D) Optical isomer of Example 398
398		¹ H-NMR (CDCl ₃) δ: 0.78 (t, 3H, J = 7.2 Hz), 0.95 (s, 3H), 1.17 (s, 3H), 1.44 (tq, 2H, J = 7.2, 7.6 Hz), 2.46 (t, 2H, J = 7.6 Hz), 3.13 (t, 1H, J = 7.2 Hz), 3.30-3.37 (m, 1H), 3.41-3.45 (m, 1H), 6.91 (s, 1H), 7.02 (s, 1H), 7.06 (d, 1H, J = 7.6 Hz), 7.16 (d, 1H, J = 8.0 Hz), 7.22-7.28 (m, 3H), 7.32-7.41 (m, 3H), 7.64 (s, 1H), 10.79 (s, 1H).	417	415	Optically active substance (Structure Information D) Optical isomer of Example 397
399		¹ H-NMR (DMSO-D ₆) δ: 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 5.12 (s, 2H), 6.95 (s, 1H), 7.19-7.25 (m, 3H), 7.33-7.42 (m, 6H), 7.64 (s, 1H), 10.88 (s, 1H).	445	444	S-form



400		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, J = 7.3 Hz), 1.46 (tq, 2H, J = 7.0, 7.3 Hz), 2.38-2.40 (m, 2H), 2.47 (t, 2H, J = 7.0 Hz), 3.33-3.35 (m, 1H), 3.41-3.45 (m, 1H), 3.49-3.53 (m, 1H), 6.92 (s, 1H), 7.02-7.05 (m, 1H), 7.10-7.12 (m, 1H), 7.18-7.26 (m, 3H), 7.30-7.34 (m, 2H), 7.51 (t, 1H, J = 8.3 Hz), 7.63 (s, 1H), 10.92 (s, 1H).	473	471	S-form
401		¹ H-NMR (DMSO-D ₆) δ: 0.76 (t, 3H, J = 7.3 Hz), 1.45 (tq, 2H, J = 7.0, 7.3 Hz), 2.38-2.40 (m, 2H), 2.49 (t, 2H, J = 7.0 Hz), 3.33-3.36 (m, 1H), 3.39-3.47 (m, 1H), 3.49-3.53 (m, 1H), 6.93 (s, 1H), 7.05 (s, 1H), 7.12-7.14 (m, 1H), 7.22-7.23 (m, 1H), 7.30-7.34 (m, 1H), 7.49-7.55 (m, 2H), 7.59-7.69 (m, 3H), 10.93 (s, 1H).	457	455	S-form
402		¹ H-NMR (DMSO-D ₆) δ: 0.73 (t, 3H, J = 7.3 Hz), 1.41 (tq, 2H, J = 7.0, 7.3 Hz), 2.37-2.39 (m, 2H), 2.46 (t, 2H, J = 7.0 Hz), 3.32-3.34 (m, 1H), 3.40-3.44 (m, 1H), 3.48-3.51 (m, 1H), 6.93 (s, 1H), 6.97 (s, 1H), 7.13 (d, 1H, J = 7.7 Hz), 7.20 (d, 1H, J = 7.7 Hz), 7.31 (dd, 1H, J = 7.7, 7.7 Hz), 7.40 (d, 2H, J = 8.6 Hz), 7.62 (s, 1H), 7.75 (d, 2H, J = 8.6 Hz), 10.92 (s, 1H).	457	455	S-form




403		<p>¹H-NMR (DMSO-D₆) δ: 0.76 (t, 3H, J = 7.4 Hz), 1.17 (t, 3H, J = 7.6 Hz), 1.41 (tq, 2H, J = 7.4 Hz, 7.4Hz), 2.36-2.41 (m, 2H), 2.45 (t, 2H, J = 7.4 Hz), 2.61 (q, 2H, J = 7.6 Hz), 3.30-3.53 (m, 3H), 6.88 (s, 1H), 6.95 (s, 1H), 7.09 (d, 1H, J = 7.6 Hz), 7.12 (d, 2H, J = 8.4 Hz), 7.15 (d, 1H, J = 7.6 Hz), 7.22 (d, 2H, J = 8.4 Hz), 7.28 (dd, 1H, J = 7.6 Hz, 7.6 Hz), 7.62 (s, 1H), 10.82 (s, 1H).</p>	417	415	S-form
404		<p>¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 1.20 (d, 3H, J = 6.4 Hz), 1.44 (tq, 2H, J = 7.4 Hz, 7.4Hz), 2.37-2.50 (m, 4H), 2.91-2.98 (m, 1H), 3.73 (dq, 1H, J = 6.4 Hz), 6.91 (s, 1H), 7.00 (s, 1H), 7.07 (d, 1H, J = 7.6 Hz), 7.17 (d, 1H, J = 7.6 Hz), 7.20-7.30 (m, 3H), 7.31-7.42 (m, 3H), 7.77 (s, 1H), 10.91 (s, 1H).</p>	403	401	Optically active substance (transform, Structure Information E) Optical isomer of Example 405
405		<p>¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 1.20 (d, 3H, J = 6.4 Hz), 1.44 (tq, 2H, J = 7.4 Hz, 7.4Hz), 2.37-2.50 (m, 4H), 2.91-2.98 (m, 1H), 3.73 (dq, 1H, J = 6.4 Hz), 6.91 (s, 1H), 7.00 (s, 1H), 7.07 (d, 1H, J = 7.6 Hz), 7.17 (d, 1H, J = 7.6 Hz), 7.20-7.30 (m, 3H), 7.31-7.42 (m, 3H), 7.77 (s, 1H), 10.91 (s, 1H).</p>	403	401	Optically active substance (transform, Structure Information E) Optical isomer of Example 404




406		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, J = 7.3 Hz), 1.46 (tq, 2H, J = 7.3, 7.7 Hz), 2.38-2.40 (m, 2H), 2.48 (t, 2H, J = 7.7 Hz), 3.26 (s, 3H), 3.33-3.34 (m, 1H), 3.40-3.44 (m, 1H), 3.49-3.51 (m, 1H), 3.55-3.58 (m, 2H), 3.94-3.98 (m, 2H), 6.76-6.79 (m, 2H), 6.88-6.92 (m, 2H), 7.02-7.04 (m, 1H), 7.07-7.09 (m, 1H), 7.17-7.19 (m, 1H), 7.25-7.30 (m, 2H), 7.63 (s, 1H), 10.85 (s, 1H).	463	461	S-form
407		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.2 Hz), 1.45 (tq, 2H, J = 7.2, 7.2 Hz), 2.37-2.40 (m, 2H), 2.46 (t, 2H, J = 7.2 Hz), 3.34-3.36 (m, 1H), 3.38-3.43 (m, 1H), 3.49-3.51 (m, 1H), 3.68-3.70 (m, 2H), 4.00-4.03 (m, 2H), 4.51 (s, 2H), 6.76-6.80 (m, 2H), 6.88-6.93 (m, 2H), 7.02-7.09 (m, 2H), 7.16-7.18 (m, 1H), 7.25-7.36 (m, 7H), 7.63 (s, 1H), 10.85 (s, 1H).	539	537	S-form




408		¹ H-NMR (DMSO-D6) δ : 0.77 (t, 3H, J = 7.3 Hz), 1.45 (tq, 2H, J = 7.2, 7.3 Hz), 2.36-2.38 (m, 2H), 2.46 (t, 2H, J = 7.2 Hz), 3.32-3.35 (m, 1H), 3.39-3.43 (m, 1H), 3.48-3.50 (m, 1H), 3.62-3.64 (m, 2H), 3.84-3.86 (m, 2H), 4.81 (t, 1H, J = 5.6 Hz), 6.72-6.78 (m, 2H), 6.87-6.89 (m, 2H), 7.01-7.08 (m, 2H), 7.16-7.18 (m, 1H), 7.24-7.28 (m, 2H), 7.61 (s, 1H), 10.83 (s, 1H).	449	447	S-form
409		¹ H-NMR (DMSO-D6) δ : 2.38-2.40 (m, 2H), 3.33-3.35 (m, 1H), 3.40-3.44 (m, 1H), 3.48-3.51 (m, 1H), 4.84 (s, 2H), 5.54 (sep, 1H, J = 6.6 Hz), 6.94 (s, 1H), 7.21-7.29 (m, 6H), 7.35-7.42 (m, 2H), 7.63 (s, 1H), 10.86 (s, 1H).	545	543	S-form
410		¹ H-NMR (DMSO-D6) δ : 1.17 (d, 3H, J = 6.4 Hz), 2.25 (s, 3H), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.02-4.08 (m, 1H), 4.54 (s, 2H), 6.90 (s, 1H), 6.97 (s, 1H), 7.04 (s, 1H), 7.14 (s, 1H), 7.22-7.25 (m, 2H), 7.32-7.41 (m, 3H), 7.63 (s, 1H), 10.85 (s, 1H).	487	485	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form




411		¹ H-NMR (DMSO-D ₆) δ: 1.06 (s, 9H), 2.29 (s, 3H), 2.33-2.45 (m, 2H), 3.30-3.53 (m, 3H), 6.79 (s, 1H), 6.91 (s, 1H), 7.06 (s, 1H), 7.17 (s, 1H), 7.20-7.25 (m, 2H), 7.32-7.43 (m, 3H), 7.63 (s, 1H), 10.83 (s, 1H).	417	415	S-form
412		¹ H-NMR (DMSO-D ₆) δ: 0.79 (t, 3H, J = 7.3 Hz), 1.42-1.59 (m, 3H), 1.65-1.76 (m, 1H), 1.84-1.97 (m, 2H), 2.14-2.24 (m, 2H), 2.37-2.41 (m, 2H), 2.45-2.52 (m, 2H), 3.30-3.53 (m, 3H), 4.42-4.51 (m, 1H), 6.55 (t, 1H, J = 2.2 Hz), 6.77-6.89 (m, 3H), 7.03 (s, 1H), 7.06-7.10 (m, 1H), 7.17-7.20 (m, 1H), 7.26-7.31 (m, 2H), 7.63 (s, 1H), 10.85 (s, 1H).	459	457	S-form
413		¹ H-NMR (DMSO-D ₆) δ: 1.47 (s, 9H), 2.38-2.41 (m, 2H), 3.31-3.54 (m, 3H), 6.99 (s, 1H), 7.24-7.28 (m, 2H), 7.35-7.54 (m, 5H), 7.64 (s, 1H), 7.67-7.70 (m, 1H), 7.84-7.88 (m, 1H), 10.89 (s, 1H).	447	445	S-form





414		¹ H-NMR (CDCl ₃) δ: 0.86 (t, 3H, J = 7.6 Hz), 1.54 (tq, 2H, J = 7.6, 7.6 Hz), 2.49-2.62 (m, 3H), 2.77 (dd, 1H, J = 17.2, 8.8 Hz), 3.28-3.38 (m, 1H), 3.40 (s, 3H), 3.56 (t, 1H, J = 8.8 Hz), 3.65-3.71 (m, 3H), 3.96-4.00 (m, 2H), 6.24 (s, 1H), 6.77 (ddd, 1H, J = 8.8, 4.0, 2.8 Hz), 6.86 (dd, 1H, J = 7.6, 2.8 Hz), 6.98-7.08 (m, 4H), 7.14 (d, 1H, J = 7.6 Hz), 7.22 (t, 1H, J = 7.6 Hz), 8.82 (s, 1H).	481	479	S-form
415		¹ H-NMR (CDCl ₃) δ: 0.86 (t, 3H, J = 7.6 Hz), 1.54 (tq, 2H, J = 7.6, 7.6 Hz), 2.48-2.58 (m, 3H), 2.76 (dd, 1H, J = 16.8, 8.4 Hz), 3.20-3.30 (m, 1H), 3.49 (t, 1H, J = 9.2 Hz), 3.65 (dd, 1H, J = 9.6, 7.6 Hz), 3.85-3.99 (m, 4H), 6.25 (s, 1H), 6.80 (ddd, 1H, J = 8.4, 4.0, 2.4 Hz), 6.86 (dd, 1H, J = 7.2, 2.4 Hz), 7.00-7.08 (m, 4H), 7.15 (d, 1H, J = 7.6 Hz), 7.23 (t, 1H, J = 7.6 Hz), 9.08 (s, 1H).	467	465	S-form

416		¹ H-NMR (CDCl ₃) δ: 2.61 (dd, 1H, J = 16.8, 9.6 Hz), 2.78 (dd, 1H, J = 16.8, 8.8 Hz), 3.32-3.42 (m, 1H), 3.61 (t, 1H, J = 9.6 Hz), 3.69-3.81 (m, 6H), 4.61 (s, 2H), 6.00 (s, 1H), 6.72 (ddd, 1H, J = 8.8, 4.0, 2.8 Hz), 6.87 (dd, 1H, J = 7.6, 2.8 Hz), 6.98-7.05 (m, 2H), 7.18-7.36 (m, 4H), 8.49 (s, 1H).	507	505	S-form
417		¹ H-NMR (CDCl ₃) δ: 1.35 (t, 3H, J = 7.2 Hz), 2.61 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.4 Hz), 3.30-3.40 (m, 1H), 3.61 (t, 1H, J = 8.8 Hz), 3.69-3.81 (m, 3H), 3.92 (q, 2H, J = 6.8 Hz), 4.61 (s, 2H), 5.93 (s, 1H), 6.72 (ddd, 1H, J = 9.2, 4.0, 2.8 Hz), 6.84 (dd, 1H, J = 7.6, 2.4 Hz), 6.98-7.05 (m, 2H), 7.17-7.36 (m, 4H), 8.41 (s, 1H).	521	519	S-form
418		¹ H-NMR (CDCl ₃) δ: 1.21 (d, 3H, J = 6.0 Hz), 1.22 (d, 1H, J = 6.0 Hz), 2.60 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.4 Hz), 3.29-3.40 (m, 1H), 3.59 (t, 1H, J = 9.2 Hz), 3.68-3.81 (m, 3H), 4.27-4.37 (m, 1H), 4.61 (s, 2H), 6.14 (s, 1H), 6.76-6.83 (m, 2H), 7.01-7.07 (m, 2H), 7.17-7.36 (m, 4H), 8.65 (s, 1H).	535	533	S-form




419		¹ H-NMR (CDCl ₃) δ: 0.97 (t, 3H, J = 7.6 Hz), 1.73 (tq, 2H, J = 7.6, 7.6 Hz), 2.61 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.4 Hz), 3.31-3.40 (m, 1H), 3.61 (t, 1H, J = 8.8 Hz), 3.70-3.84 (m, 5H), 4.61 (s, 2H), 5.91 (s, 1H), 6.71 (ddd, 1H, J = 8.8, 4.0, 2.4 Hz), 6.84 (dd, 1H, J = 7.6, 2.4 Hz), 6.98-7.04 (m, 2H), 7.17-7.35 (m, 4H), 8.38 (s, 1H).	535	533	S-form
420		¹ H-NMR (CDCl ₃) δ: 2.62 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.4 Hz), 3.32-3.42 (m, 1H), 3.60 (t, 1H, J = 9.2 Hz), 3.70-3.76 (m, 4H), 6.22 (s, 1H), 6.74 (ddd, 1H, J = 8.8, 4.0, 2.8 Hz), 6.83 (dd, 1H, J = 7.6, 2.4 Hz), 7.01-7.10 (m, 3H), 7.17-7.24 (m, 2H), 7.38 (t, 1H, J = 8.0 Hz), 8.68 (s, 1H).	479	477	S-form
421		¹ H-NMR (CDCl ₃) δ: 1.35 (t, 3H, J = 7.2 Hz), 2.61 (dd, 1H, J = 17.2, 9.6 Hz), 2.77 (dd, 1H, J = 17.2, 8.4 Hz), 3.31-3.41 (m, 1H), 3.58 (t, 1H, J = 8.8 Hz), 3.69 (dd, 1H, J = 9.6, 7.2 Hz), 3.91 (q, 2H, 7.2 Hz), 6.25 (s, 1H), 6.72-6.81 (m, 2H), 7.02-7.09 (m, 3H), 7.17-7.22 (m, 2H), 7.37 (t, 1H, J = 8.0 Hz), 8.71 (s, 1H).	493	491	S-form




422		¹ H-NMR (CDCl ₃) δ: 1.20 (d, 3H, J = 6.0 Hz), 1.22 (d, 3H, J = 6.0 Hz), 2.61 (dd, 1H, J = 17.2, 9.6 Hz), 2.77 (dd, 1H, J = 17.2, 8.4 Hz), 3.31-3.41 (m, 1H), 3.58 (t, 1H, J = 9.2 Hz), 3.69 (dd, 1H, J = 9.6, 7.2 Hz), 4.25-4.36 (m, 1H), 6.40 (s, 1H), 6.73 (dd, 1H, J = 7.2, 2.4 Hz), 6.81 (ddd, 1H, J = 8.8, 4.0, 2.8 Hz), 7.04-7.11 (m, 3H), 7.16-7.23 (m, 2H), 7.37 (t, 1H, J = 8.4 Hz), 8.90 (s, 1H).	507	505	S-form
423		¹ H-NMR (CDCl ₃) δ: 0.98 (t, 3H, J = 7.6 Hz), 1.74 (tq, 2H, J = 7.6, 7.6 Hz), 2.62 (dd, 1H, J = 17.2, 10.0 Hz), 2.77 (dd, 1H, J = 17.2, 8.8 Hz), 3.32-3.42 (m, 1H), 3.62 (t, 1H, J = 8.8 Hz), 3.73 (dd, 1H, J = 9.6, 7.2 Hz), 3.81 (t, 2H, J = 7.6 Hz), 5.90 (s, 1H), 6.73 (ddd, 1H, J = 8.8, 4.0, 2.8 Hz), 6.80 (dd, 1H, J = 7.6, 2.8 Hz), 7.01-7.10 (m, 3H), 7.17-7.23 (m, 2H), 7.37 (t, 1H, J = 8.0 Hz), 8.32 (s, 1H).	507	505	S-form
424		¹ H-NMR (CDCl ₃) δ: 2.57 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.8 Hz), 3.30-3.42 (m, 4H), 3.53 (t, 1H, J = 9.2 Hz), 3.60-3.73 (m, 3H), 3.78 (q, 2H, J = 8.4 Hz), 3.95-4.03 (m, 2H), 4.60 (s, 2H), 6.72-6.90 (m, 3H), 6.98-7.08 (m, 2H), 7.15-7.26 (m, 2H), 7.28-7.36 (m, 2H), 9.39 (s, 1H).	551	549	S-form




425		¹ H-NMR (CDCl ₃) δ: 1.27-1.37 (m, 6H), 2.59 (dd, 1H, J = 17.2, 9.6 Hz), 2.77 (dd, 1H, J = 17.2, 8.4 Hz), 3.29-3.40 (m, 1H), 3.56 (t, 1H, J = 8.8 Hz), 3.68 (dd, 1H, J = 9.6, 7.2 Hz), 3.72-3.82 (m, 1H), 3.90 (q, 2H, J = 7.2 Hz), 4.57 (d, 1H, J = 11.6 Hz), 4.66 (d, 1H, J = 11.6 Hz), 6.39 (s, 1H), 6.73-6.83 (m, 2H), 6.99-7.06 (m, 2H), 7.16-7.25 (m, 2H), 7.29-7.35 (m, 2H), 8.90 (s, 1H).	535	533	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
426		¹ H-NMR (CDCl ₃) δ: 1.13-1.32 (m, 9H), 2.57 (dd, 1H, J = 17.2, 9.6 Hz), 2.77 (dd, 1H, J = 17.2, 8.4 Hz), 3.28-3.82 (m, 4H), 4.23-4.33 (m, 1H), 4.56 (d, 1H, J = 12.4 Hz), 4.66 (d, 1H, J = 12.4 Hz), 6.71-6.86 (m, 2H), 7.00-7.35 (m, 7H), 9.69 (s, 1H).	549	547	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
427		¹ H-NMR (DMSO-D ₆) δ: 1.21 (d, 3H, J = 6.4 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.10-4.17 (m, 1H), 4.63 (s, 2H), 6.93 (s, 1H), 7.20-7.25 (m, 3H), 7.28-7.38 (m, 4H), 7.42-7.46 (m, 1H), 7.63 (s, 1H), 10.92 (s, 1H).	491	489	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form




428		¹ H-NMR (DMSO-D6) δ : 1.62 (brs, 4H), 2.27 (brs, 4H), 2.33-2.45 (m, 2H), 3.30-3.53 (m, 5H), 6.89 (s, 1H), 7.11 (s, 1H), 7.17 (d, 1H, J = 7.6 Hz), 7.20-7.29 (m, 3H), 7.30-7.42 (m, 4H), 7.32-7.43 (m, 3H), 7.63 (s, 1H), 10.85 (s, 1H).	430	428	S-form
429		¹ H-NMR (DMSO-D6) δ : 0.87 (t, 3H, J = 7.3 Hz), 1.47-1.57 (m, 2H), 2.37-2.42 (m, 2H), 3.20 (q, 2H, J = 6.6 Hz), 3.32-3.55 (m, 3H), 7.02 (s, 1H), 7.19-7.25 (m, 3H), 7.31-7.42 (m, 4H), 7.64 (s, 1H), 7.78-7.82 (m, 1H), 7.85-7.87 (m, 1H), 8.52 (t, 1H, J = 5.7 Hz), 10.88 (s, 1H).	432	430	S-form
430		¹ H-NMR (DMSO-D6) δ : 0.55-0.90 (m, 3H), 1.30-1.57 (m, 2H), 2.37-2.42 (m, 2H), 2.60-2.65 (m, 1H), 2.82-2.98 (m, 2H), 3.30-3.54 (m, 5H), 6.95 (s, 1H), 7.09 (s, 1H), 7.22-7.28 (m, 2H), 7.31-7.48 (m, 6H), 7.63 (s, 1H), 10.87 (s, 1H).	446	444	S-form
431		¹ H-NMR (DMSO-D6) δ : 1.15 (d, 6H, J = 6.7 Hz), 2.38-2.42 (m, 2H), 3.30-3.54 (m, 3H), 4.03-4.12 (m, 1H), 7.02 (s, 1H), 7.17-7.26 (m, 3H), 7.32-7.42 (m, 4H), 7.64 (s, 1H), 7.78-7.83 (m, 1H), 7.87 (s, 1H), 8.28 (d, 1H, J = 7.7 Hz), 10.88 (s, 1H).	432	430	S-form




432		¹ H-NMR (DMSO-D6) δ: 0.90-1.15 (m, 6H), 2.37-2.42 (m, 2H), 2.69-2.81 (m, 1H), 3.28-3.53 (m, 6H), 6.94 (s, 1H), 7.00-7.15 (m, 1H), 7.21-7.27 (m, 2H), 7.31-7.50 (m, 6H), 7.63 (s, 1H), 10.88 (s, 1H).	446	444	S-form
433		¹ H-NMR (DMSO-D6) δ: 2.38-2.42 (m, 2H), 3.30-3.55 (m, 3H), 4.03-4.15 (m, 2H), 7.05 (s, 1H), 7.22-7.27 (m, 3H), 7.32-7.45 (m, 4H), 7.64 (s, 1H), 7.85 (d, 1H, J = 7.9 Hz), 7.93 (s, 1H), 9.18 (t, 1H, J = 6.2 Hz), 10.89 (s, 1H).	472	470	S-form
434		¹ H-NMR (DMSO-D6) δ: 0.55 (s, 3H), 0.61 (s, 3H), 0.73 (t, 3H, J = 7.4 Hz), 0.94-1.05 (m, 1H), 1.11-1.22 (m, 1H), 2.36-2.42 (m, 2H), 3.31-3.53 (m, 3H), 4.18 (d, 1H, J = 4.2 Hz), 5.08 (d, 1H, J = 4.2 Hz), 6.86 (s, 1H), 7.07 (s, 1H), 7.15-7.40 (m, 8H), 7.63 (s, 1H), 10.84 (s, 1H).	447	445	1-hydroxy-2,2-dimethyl - butyl group: racemate Lactam ring: S-form
435		¹ H-NMR (DMSO-D6) δ: 0.80 (t, 3H, J = 7.3 Hz), 1.13 (s, 9H), 1.42-1.51 (m, 2H), 2.37-2.42 (m, 2H), 2.44-2.52 (m, 3H), 3.31-3.53 (m, 3H), 6.70-6.72 (m, 1H), 6.87 (s, 1H), 6.91-6.95 (m, 1H), 7.01-7.10 (m, 3H), 7.15-7.19 (m, 1H), 7.24-7.34 (m, 2H), 7.63 (s, 1H), 10.83 (s, 1H).	461	459	S-form




436		1H-NMR (DMSO-D6) δ : 2.38-2.40 (m, 2H), 3.33-3.36 (m, 1H), 3.40-3.44 (m, 1H), 3.48-3.51 (m, 1H), 5.01 (s, 2H), 6.78-6.81 (m, 1H), 6.90-6.92 (m, 1H), 6.99-7.03 (m, 2H), 7.11-7.13 (m, 1H), 7.28-7.40 (m, 8H), 7.53 (t, 1H, J = 7.9 Hz), 7.63 (s, 1H), 10.90 (s, 1H).	537	535	S-form
437		1H-NMR (DMSO-D6) δ : 2.38-2.40 (m, 2H), 3.32-3.36 (m, 1H), 3.40-3.45 (m, 1H), 3.48-3.51 (m, 1H), 3.99 (q, 2H, J = 9.3 Hz), 4.63 (s, 2H), 5.00 (s, 2H), 6.76-6.78 (m, 1H), 6.89-6.92 (m, 2H), 6.97-6.99 (m, 1H), 7.16-7.18 (m, 1H), 7.25-7.40 (m, 9H), 7.63 (s, 1H), 10.87 (s, 1H).	565	563	S-form
438		1H-NMR (DMSO-D6) δ : 1.19 (d, 3H, J = 6.4 Hz), 2.37-2.41 (m, 2H), 3.33-3.35 (m, 1H), 3.39-3.45 (m, 1H), 3.46-3.52 (m, 1H), 4.05-4.16 (m, 1H), 4.63 (s, 2H), 4.99 (s, 2H), 6.77-6.79 (m, 1H), 6.89-6.92 (m, 2H), 6.97-6.99 (m, 1H), 7.15-7.17 (m, 1H), 7.25-7.48 (m, 9H), 7.63 (s, 1H), 10.87 (s, 1H).	579	577	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form




439		¹ H-NMR (DMSO-D ₆) δ: 2.38-2.40 (m, 2H), 3.33-3.36 (m, 1H), 3.41-3.45 (m, 1H), 3.48-3.52 (m, 1H), 6.60-6.62 (m, 1H), 6.66-6.67 (m, 1H), 6.75-6.77 (m, 1H), 6.97 (s, 1H), 7.10-7.12 (m, 1H), 7.16-7.20 (m, 1H), 7.34-7.36 (m, 2H), 7.53 (t, 1H, J = 8.2 Hz), 7.63 (s, 1H), 9.78 (br s, 1H), 10.87 (s, 1H).	447	445	S-form
440		¹ H-NMR (DMSO-D ₆) δ: 2.37-2.40 (m, 2H), 3.32-3.34 (m, 1H), 3.38-3.46 (m, 1H), 3.48-3.52 (m, 1H), 3.99 (q, 2H, J = 9.3 Hz), 4.62 (s, 2H), 6.57-6.60 (m, 1H), 6.65-6.67 (m, 1H), 6.71-6.73 (m, 1H), 6.89 (s, 1H), 7.12-7.18 (m, 2H), 7.26-7.28 (m, 1H), 7.32-7.39 (m, 2H), 7.63 (s, 1H), 9.78 (br s, 1H), 10.84 (s, 1H).	475	473	S-form
441		¹ H-NMR (DMSO-D ₆) δ: 1.21 (d, 3H, J = 6.2 Hz), 2.37-2.40 (m, 2H), 3.33-3.35 (m, 1H), 3.38-3.46 (m, 1H), 3.48-3.52 (m, 1H), 4.08-4.15 (m, 1H), 4.63 (s, 2H), 6.57-6.60 (m, 1H), 6.66-6.68 (m, 1H), 6.70-6.74 (m, 1H), 6.89 (s, 1H), 7.12-7.17 (m, 2H), 7.26-7.28 (m, 1H), 7.32-7.38 (m, 2H), 7.63 (s, 1H), 9.79 (br s, 1H), 10.84 (s, 1H).	489	487	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form




442		¹ H-NMR (DMSO-D ₆) δ: 2.38-2.40 (m, 2H), 3.32-3.35 (m, 1H), 3.40-3.45 (m, 1H), 3.48-3.51 (m, 1H), 3.67 (s, 3H), 6.76-6.78 (m, 1H), 6.83-6.84 (m, 1H), 6.92-6.95 (m, 1H), 6.99 (s, 1H), 7.12 (s, 1H), 7.30 (t, 1H, J = 8.0 Hz), 7.36-7.39 (m, 2H), 7.53 (t, 1H, J = 8.0 Hz), 7.63 (s, 1H), 10.91 (s, 1H).	461	459	S-form
443		¹ H-NMR (DMSO-D ₆) δ: 1.24 (t, 3H, J = 7.0 Hz), 2.38-2.40 (m, 2H), 3.33-3.35 (m, 1H), 3.40-3.45 (m, 1H), 3.48-3.53 (m, 1H), 3.92 (q, 2H, J = 7.0 Hz), 6.75-6.79 (m, 2H), 6.91-6.93 (m, 1H), 6.99 (s, 1H), 7.11-7.13 (m, 1H), 7.27-7.30 (m, 1H), 7.35-7.38 (m, 2H), 7.53 (t, 1H, J = 8.0 Hz), 7.63 (s, 1H), 10.90 (s, 1H).	475	473	S-form
444		¹ H-NMR (DMSO-D ₆) δ: 0.90 (t, 3H, J = 7.4 Hz), 1.62 (tq, 2H, J = 6.6, 7.4 Hz), 2.38-2.40 (m, 2H), 3.33-3.35 (m, 1H), 3.40-3.44 (m, 1H), 3.48-3.53 (m, 1H), 3.80 (t, 2H, J = 6.6 Hz), 6.76-6.80 (m, 2H), 6.91-6.94 (m, 1H), 6.98 (s, 1H), 7.11-7.13 (m, 1H), 7.29 (t, 1H, J = 8.3 Hz), 7.36-7.38 (m, 2H), 7.54 (t, 1H, J = 8.3 Hz), 7.63 (s, 1H), 10.89 (s, 1H).	489	487	S-form




445		<p>¹H-NMR (DMSO-D₆) δ: 1.13 (d, 6H, J = 6.0 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.33-3.35 (m, 1H), 3.39-3.44 (m, 1H), 3.48-3.52 (m, 1H), 4.45 (sep, 1H), 6.71-6.73 (m, 1H), 6.81-6.82 (m, 1H), 6.89-6.91 (m, 1H), 6.98 (s, 1H), 7.11 (s, 1H), 7.30 (t, 1H, J = 8.0 Hz), 7.34-7.38 (m, 2H), 7.53 (t, 1H, J = 8.0 Hz), 7.63 (s, 1H), 10.89 (s, 1H).</p>	489	487	S-form
446		<p>¹H-NMR (DMSO-D₆) δ: 2.38-2.40 (m, 2H), 3.33-3.34 (m, 1H), 3.40-3.45 (m, 1H), 3.48-3.52 (m, 1H), 3.66 (s, 3H), 3.98 (q, 2H, J = 9.3 Hz), 4.62 (s, 2H), 6.73-6.75 (m, 1H), 6.81-6.84 (m, 1H), 6.88-6.91 (m, 2H), 7.18-7.20 (m, 1H), 7.25-7.28 (m, 2H), 7.33-7.40 (m, 2H), 7.63 (s, 1H), 10.88 (s, 1H).</p>	489	487	S-form
447		<p>¹H-NMR (DMSO-D₆) δ: 1.24 (t, 3H, J = 7.0 Hz), 2.38-2.40 (m, 2H), 3.33-3.35 (m, 1H), 3.39-3.44 (m, 1H), 3.49-3.52 (m, 1H), 3.90 (q, 2H, J = 7.0 Hz), 3.98 (q, 2H, J = 9.3 Hz), 4.62 (s, 2H), 6.74-6.79 (m, 2H), 6.86-6.88 (m, 1H), 6.91 (s, 1H), 7.17-7.19 (m, 1H), 7.23-7.29 (m, 2H), 7.33-7.40 (m, 2H), 7.63 (s, 1H), 10.87 (s, 1H).</p>	503	501	S-form




448		¹ H-NMR (DMSO-D ₆) δ: 0.90 (t, 3H, J = 7.3 Hz), 1.62 (tq, 2H, J = 6.6, 7.3 Hz), 2.37-2.41 (m, 2H), 3.34-3.36 (m, 1H), 3.39-3.43 (m, 1H), 3.48-3.52 (m, 1H), 3.80 (t, 2H, J = 6.6 Hz), 3.98 (q, 2H, J = 9.3 Hz), 4.62 (s, 2H), 6.74-6.79 (m, 2H), 6.86-6.89 (m, 1H), 6.91 (s, 1H), 7.18-7.20 (m, 1H), 7.23-7.28 (m, 1H), 7.33-7.40 (m, 2H), 7.63 (s, 1H), 10.87 (s, 1H).	517	515	S-form
449		¹ H-NMR (DMSO-D ₆) δ: 1.13 (d, 6H, J = 6.0 Hz), 2.37-2.39 (m, 2H), 3.33-3.35 (m, 1H), 3.41-3.45 (m, 1H), 3.48-3.52 (m, 1H), 3.98 (q, 2H, J = 9.3 Hz), 4.43 (sep, 1H, J = 6.0 Hz), 4.62 (s, 2H), 6.72-6.73 (m, 1H), 6.77-6.79 (m, 1H), 6.85-6.87 (m, 1H), 6.91 (s, 1H), 7.17-7.19 (m, 1H), 7.25-7.28 (m, 2H), 7.33-7.41 (m, 2H), 7.63 (s, 1H), 10.85 (s, 1H).	517	515	S-form
450		¹ H-NMR (DMSO-D ₆) δ: 2.38-2.40 (m, 2H), 3.26 (s, 3H), 3.34-3.36 (m, 1H), 3.40-3.44 (m, 1H), 3.48-3.52 (m, 1H), 3.55-3.58 (m, 2H), 3.96-4.02 (m, 4H), 4.62 (s, 2H), 6.76-6.81 (m, 2H), 6.89-6.91 (m, 2H), 7.17-7.18 (m, 1H), 7.24-7.29 (m, 2H), 7.33-7.40 (m, 2H), 7.62 (s, 1H), 10.86 (s, 1H).	533	531	S-form






451		¹ H-NMR (DMSO-D ₆) δ: 1.20 (d, 3H, J = 6.5 Hz), 2.36-2.40 (m, 2H), 3.33-3.36 (m, 1H), 3.39-3.44 (m, 1H), 3.48-3.52 (m, 1H), 3.66 (s, 3H), 4.06-4.13 (m, 1H), 4.63 (s, 2H), 6.73-6.75 (m, 1H), 6.81-6.84 (m, 1H), 6.88-6.92 (m, 2H), 7.17-7.19 (m, 1H), 7.24-7.28 (m, 2H), 7.33-7.39 (m, 2H), 7.63 (s, 1H), 10.87 (s, 1H).	503	501	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
452		¹ H-NMR (DMSO-D ₆) δ: 1.20 (d, 3H, J = 6.3 Hz), 1.24 (t, 3H, J = 7.0 Hz), 2.38-2.40 (m, 2H), 3.33-3.36 (m, 1H), 3.40-3.44 (m, 1H), 3.47-3.53 (m, 1H), 3.91 (q, 2H, J = 7.0 Hz), 4.06-4.13 (m, 1H), 4.62 (s, 2H), 6.74-6.79 (m, 2H), 6.87-6.90 (m, 2H), 7.17-7.19 (m, 1H), 7.23-7.28 (m, 2H), 7.33-7.38 (m, 2H), 7.63 (s, 1H), 10.86 (s, 1H).	517	515	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
453		¹ H-NMR (DMSO-D ₆) δ: 0.90 (t, 3H, J = 7.4 Hz), 1.20 (d, 3H, J = 6.5 Hz), 1.62 (tq, 2H, J = 7.0, 6.6 Hz), 2.38-2.40 (m, 2H), 3.33-3.36 (m, 1H), 3.40-3.45 (m, 1H), 3.48-3.52 (m, 1H), 3.80 (t, 2H, J = 6.6 Hz), 4.06-4.15 (m, 1H), 4.63 (s, 2H), 6.74-6.78 (m, 2H), 6.87-6.90 (m, 2H), 7.17-7.19 (m, 1H), 7.23-7.28 (m, 2H), 7.33-7.39 (m, 2H), 7.63 (s, 1H), 10.86 (s, 1H).	531	529	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form





454		¹ H-NMR (DMSO-D ₆) δ: 1.14 (d, 6H, J = 6.0 Hz), 1.20 (d, 3H, J = 6.5 Hz), 2.37-2.40 (m, 2H), 3.33-3.35 (m, 1H), 3.40-3.44 (m, 1H), 3.48-3.52 (m, 1H), 4.07-4.14 (m, 1H), 4.43 (sep, 1H, J = 6.0 Hz), 4.62 (s, 2H), 6.71-6.73 (m, 1H), 6.76-6.79 (m, 1H), 6.85-6.87 (m, 1H), 6.90 (s, 1H), 7.17-7.19 (m, 1H), 7.24-7.28 (m, 1H), 7.32-7.39 (m, 2H), 7.63 (s, 1H), 10.85 (s, 1H).	531	529	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form
455		¹ H-NMR (DMSO-D ₆) δ: 1.18 (d, 3H, J = 6.3 Hz), 2.36-2.38 (m, 2H), 3.24 (s, 3H), 3.32-3.33 (m, 1H), 3.40-3.44 (m, 1H), 3.47-3.49 (m, 1H), 3.53-3.57 (m, 2H), 3.94-3.97 (m, 2H), 4.06-4.13 (m, 1H), 4.61 (s, 2H), 6.74-6.76 (m, 1H), 6.79-6.81 (m, 1H), 6.88-6.91 (m, 2H), 7.14-7.16 (m, 1H), 7.23-7.28 (m, 2H), 7.33-7.37 (m, 2H), 7.61 (s, 1H), 10.85 (s, 1H).	547	545	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form
456		¹ H-NMR (DMSO-D ₆) δ: 1.20 (d, 3H, J = 6.4 Hz), 2.33-2.45 (m, 2H), 2.46-2.58 (m, 4H), 3.30-3.53 (m, 3H), 3.70 (s, 3H), 4.10 (tt, 1H, J = 6.4 Hz, 6.4 Hz), 4.59-4.66 (m, 2H), 6.68-6.72 (m, 1H), 6.92 (s, 1H), 7.10 (dd, 1H, J = 7.8 Hz, 2.2 Hz), 7.16-7.26 (m, 3H), 7.32-7.42 (m, 2H), 7.63 (s, 1H), 10.89 (s, 1H).	565	563	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form





457		¹ H-NMR (DMSO-D ₆) δ: 1.20 (d, 3H, J = 6.4 Hz), 2.33-2.45 (m, 2H), 3.30-3.53 (m, 3H), 3.70 (s, 3H), 4.10 (tt, 1H, J = 6.4 Hz, 6.4 Hz), 4.59-4.66 (m, 2H), 6.68-6.72 (m, 1H), 6.92 (s, 1H), 7.10 (dd, 1H, J = 7.8 Hz, 2.6 Hz), 7.15-7.26 (m, 3H), 7.32-7.42 (m, 2H), 7.63 (s, 1H), 10.89 (s, 1H).	521	519	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form
458		¹ H-NMR (CDCl ₃) δ: 0.78 (t, 3H, J = 7.2 Hz), 1.45 (tq, 2H, J = 7.2, 7.6 Hz), 2.39 (d, 2H, J = 8.4 Hz), 2.47 (t, 2H, J = 7.6 Hz), 3.19 (s, 3H), 3.32-3.52 (m, 3H), 4.37 (s, 2H), 6.89 (s, 1H), 7.02 (s, 1H), 7.06-7.10 (m, 2H), 7.17 (d, 1H, J = 7.6 Hz), 7.22 (s, 1H), 7.25-7.29 (m, 2H), 7.34 (t, 1H, J = 8.0 Hz), 7.63 (s, 1H), 10.85 (s, 1H).	433	431	S-form
459		¹ H-NMR (CDCl ₃) δ: 0.78 (t, 3H, J = 7.2 Hz), 1.06 (d, 6H, J = 6.0 Hz), 1.45 (tq, 2H, J = 7.2, 7.6 Hz), 2.39 (d, 2H, J = 8.4 Hz), 2.47 (t, 2H, J = 7.6 Hz), 3.32-3.52 (m, 4H), 4.41 (s, 2H), 6.89 (s, 1H), 7.02 (s, 1H), 7.06-7.10 (m, 2H), 7.16-7.20 (m, 2H), 7.25-7.29 (m, 2H), 7.33 (t, 1H, J = 8.0 Hz), 7.63 (s, 1H), 10.85 (s, 1H).	461	459	S-form





460		¹ H-NMR (DMSO-D ₆) δ: 1.28 (t, 3H, J = 7.2 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 3.94-4.01 (m, 4H), 4.60 (s, 2H), 6.97-7.02 (m, 2H), 7.06-7.08 (m, 1H), 7.16-7.22 (m, 2H), 7.27 (s, 1H), 7.27-7.37 (m, 2H), 7.63 (s, 1H), 10.86 (s, 1H).	521	519	S-form
461		¹ H-NMR (DMSO-D ₆) δ: 1.18 (d, 6H, J = 6.0 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 3.98 (q, 2H, J = 9.2 Hz), 4.53 (sep, 1H, J = 6.0 Hz), 4.60 (s, 2H), 6.95-7.00 (m, 2H), 7.02-7.04 (m, 1H), 7.17-7.22 (m, 2H), 7.26 (s, 1H), 7.30-7.37 (m, 2H), 7.64 (s, 1H), 10.86 (s, 1H).	535	533	S-form
462		¹ H-NMR (DMSO-D ₆) δ: 2.39 (d, 2H, J = 8.4 Hz), 3.28 (s, 3H), 3.32-3.52 (m, 3H), 3.59-3.61 (m, 2H), 3.98 (q, 2H, J = 9.2 Hz), 4.04-4.07 (m, 2H), 4.61 (s, 2H), 6.97 (s, 1H), 7.01-7.05 (m, 1H), 7.10-7.12 (m, 1H), 7.15-7.23 (m, 2H), 7.28 (s, 1H), 7.30-7.37 (m, 2H), 7.64 (s, 1H), 10.86 (s, 1H).	551	549	S-form





463		¹ H-NMR (DMSO-D ₆) δ: 1.19 (d, 3H, J = 6.4 Hz), 1.28 (t, 3H, J = 7.2 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 3.98 (q, 2H, 7.2 Hz), 4.06-4.12 (m, 1H), 4.61 (s, 2H), 6.96 (s, 1H), 6.98-7.02 (m, 1H), 7.06-7.09 (m, 1H), 7.16-7.22 (m, 2H), 7.26 (s, 1H), 7.29-7.36 (m, 2H), 7.64 (s, 1H), 10.86 (s, 1H).	535	533	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
464		¹ H-NMR (DMSO-D ₆) δ: 1.14-1.20 (m, 9H), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.06-4.12 (m, 1H), 4.51-4.57 (m, 1H), 4.63 (s, 2H), 6.96 (s, 1H), 6.97-7.00 (m, 1H), 7.03-7.05 (m, 1H), 7.17-7.22 (m, 2H), 7.25 (s, 1H), 7.29-7.36 (m, 2H), 7.64 (s, 1H), 10.86 (s, 1H).	549	547	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
465		¹ H-NMR (DMSO-D ₆) δ: 1.19 (d, 3H, J = 6.4 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.28 (s, 3H), 3.32-3.52 (m, 3H), 3.59-3.62 (m, 2H), 4.05-4.13 (m, 3H), 4.61 (s, 2H), 6.97 (s, 1H), 7.01-7.05 (m, 1H), 7.10-7.12 (m, 1H), 7.16-7.23 (m, 2H), 7.27 (s, 1H), 7.29-7.36 (m, 2H), 7.64 (s, 1H), 10.86 (s, 1H).	565	563	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form




466		¹ H-NMR (CDCl ₃) δ: 2.56 (dd, 1H, J = 17.2, 9.6 Hz), 2.77 (dd, 1H, J = 17.2, 8.8 Hz), 3.28-3.39 (m, 1H), 3.58 (t, 1H, J = 8.8 Hz), 3.68 (dd, 1H, J = 9.2, 7.2 Hz), 6.29 (s, 1H), 7.01 (br s, 1H), 7.10-7.25 (m, 5H), 7.31-7.44 (m, 3H), 8.93 (s, 1H).	449	447	S-form
467		¹ H-NMR (CDCl ₃) δ: 2.54-2.87 (m, 2H), 3.33-3.75 (m, 3H), 6.39 (s, 1H), 6.93-7.11 (m, 5H), 7.19-7.42 (m, 4H), 8.90 (s, 1H).	449	447	S-form
468		¹ H-NMR (CDCl ₃) δ: 2.56-2.83 (m, 2H), 3.32-3.79 (m, 5H), 4.61 (s, 2H), 5.96 (s, 1H), 6.95-7.06 (m, 4H), 7.21-7.38 (m, 5H), 8.44 (s, 1H).	477	475	S-form
469		¹ H-NMR (CDCl ₃) δ: 1.28 (d, 3H, J = 6.4 Hz), 2.59 (dd, 1H, J = 17.2, 9.6 Hz), 2.76 (dd, 1H, J = 17.2, 8.4 Hz), 3.31-3.41 (m, 1H), 3.56-3.81 (m, 3H), 4.57 (d, 1H, J = 12.0 Hz), 4.68 (d, 1H, J = 12.0 Hz), 6.29 (s, 1H), 6.95-7.07 (m, 4H), 7.17-7.36 (m, 5H), 8.82 (s, 1H).	491	489	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form
470		¹ H-NMR (DMSO-D ₆) δ: 0.62 (t, 3H, J = 7.5 Hz), 1.02 (s, 6H), 1.44 (q, 2H, J = 7.5 Hz), 2.37-2.41 (m, 2H), 3.31-3.54 (m, 3H), 6.97 (s, 1H), 7.24-7.29 (m, 3H), 7.33-7.44 (m, 3H), 7.48-7.59 (m, 3H), 7.63 (s, 1H), 10.88 (s, 1H).	445	443	S-form




471		¹ H-NMR (CDCl ₃) δ: 2.57 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.8 Hz), 3.29-3.39 (m, 1H), 3.58-3.75 (m, 4H), 4.57 (s, 2H), 5.88 (s, 1H), 7.07-7.42 (m, 9H), 8.46 (s, 1H).	477	475	S-form
472		¹ H-NMR (CDCl ₃) δ: 1.26 (d, 3H, J = 6.4 Hz), 2.55 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.4 Hz), 3.26-3.37 (m, 1H), 3.55-3.76 (m, 3H), 4.52 (d, 1H, J = 12.4 Hz), 4.64 (d, 1H, J = 12.4 Hz), 6.05 (s, 1H), 7.07-7.42 (m, 9H), 8.68 (s, 1H).	491	489	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form
473		¹ H-NMR (DMSO-D ₆) δ: 2.39-2.41 (m, 2H), 3.34-3.37 (m, 1H), 3.39-3.44 (m, 1H), 3.48-3.53 (m, 1H), 7.01 (s, 1H), 7.08 (s, 1H), 7.30-7.34 (m, 2H), 7.47-7.55 (m, 3H), 7.59-7.63 (m, 3H), 10.86 (s, 1H).	465	463	S-form
474		¹ H-NMR (DMSO-D ₆) δ: 2.37-2.40 (m, 2H), 3.32-3.34 (m, 1H), 3.38-3.42 (m, 1H), 3.46-3.51 (m, 1H), 3.94 (q, 2H, J = 9.3 Hz), 4.56 (s, 2H), 6.98 (s, 1H), 7.09-7.11 (m, 1H), 7.20-7.29 (m, 3H), 7.40-7.62 (m, 5H), 10.81 (s, 1H).	493	491	S-form




475		¹ H-NMR (DMSO-D ₆) δ: 1.19 (d, 3H, J = 5.8 Hz), 2.38-2.40 (m, 2H), 3.35-3.37 (m, 1H), 3.40-3.44 (m, 1H), 3.48-3.52 (m, 1H), 4.05-4.10 (m, 1H), 4.57 (s, 2H), 6.99 (s, 1H), 7.11-7.12 (m, 1H), 7.21 (s, 1H), 7.26-7.32 (m, 2H), 7.46-7.60 (m, 4H), 7.63 (s, 1H), 10.82 (s, 1H).	507	505	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form
476		¹ H-NMR (CDCl ₃) δ: 2.51 (dd, 1H, J = 17.2, 9.6 Hz), 2.76 (dd, 1H, J = 17.2, 8.4 Hz), 3.23-3.34 (m, 1H), 3.55-3.72 (m, 2H), 3.87 (s, 3H), 5.97 (s, 1H), 6.91-7.37 (m, 8H), 8.77 (s, 1H).	479	477	S-form
477		¹ H-NMR (CDCl ₃) δ: 1.40 (t, 3H, J = 6.8 Hz), 2.50 (dd, 1H, J = 17.2, 9.6 Hz), 2.76 (dd, 1H, J = 17.2, 8.4 Hz), 3.21-3.32 (m, 1H), 3.52-3.70 (m, 2H), 4.08 (q, 2H, J = 6.8 Hz), 6.16 (s, 1H), 6.89-7.38 (m, 8H), 8.99 (s, 1H).	493	491	S-form
478		¹ H-NMR (CDCl ₃) δ: 1.29 (d, 6H, J = 6.0 Hz), 2.56 (dd, 1H, J = 17.2, 9.6 Hz), 2.77 (dd, 1H, J = 17.2, 8.4 Hz), 3.27-3.39 (m, 1H), 3.56-3.74 (m, 2H), 4.43-4.52 (m, 1H), 6.01 (s, 1H), 6.90-7.37 (m, 8H), 8.65 (s, 1H).	507	505	S-form

479		¹ H-NMR (CDCl ₃) δ: 2.48 (dd, 1H, J = 17.2, 9.6 Hz), 2.76 (dd, 1H, J = 17.2, 8.4 Hz), 3.18-3.29 (m, 1H), 3.41 (s, 3H), 3.52-3.76 (m, 4H), 4.11-4.22 (m, 2H), 5.98 (s, 1H), 6.92-7.37 (m, 8H), 8.83 (s, 1H).	523	521	S-form
480		¹ H-NMR (CDCl ₃) δ: 2.49 (dd, 1H, J = 17.2, 9.6 Hz), 2.76 (dd, 1H, J = 17.2, 8.4 Hz), 3.22-3.31 (m, 1H), 3.54-3.71 (m, 4H), 3.86 (s, 3H), 4.58 (s, 2H), 5.97 (s, 1H), 6.90-7.34 (m, 8H), 8.84 (s, 1H).	507	505	S-form
481		¹ H-NMR (CDCl ₃) δ: 1.39 (t, 3H, J = 6.8 Hz), 2.49 (dd, 1H, J = 17.2, 9.6 Hz), 2.76 (dd, 1H, J = 17.2, 8.4 Hz), 3.21-3.32 (m, 1H), 3.51-3.71 (m, 4H), 4.07 (q, 2H, J = 6.8 Hz), 4.58 (s, 2H), 6.11 (s, 1H), 6.88-7.12 (m, 4H), 7.20-7.34 (m, 4H), 9.01 (s, 1H).	521	519	S-form
482		¹ H-NMR (CDCl ₃) δ: 1.29 (d, 6H, J = 6.4 Hz), 2.56 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.4 Hz), 3.28-3.38 (m, 1H), 3.56-3.73 (m, 4H), 4.42-4.52 (m, 1H), 4.58 (s, 2H), 6.09 (s, 1H), 6.88-7.12 (m, 4H), 7.19-7.33 (m, 4H), 8.78 (s, 1H).	535	533	S-form





483		¹ H-NMR (CDCl ₃) δ: 2.43 (dd, 1H, J = 17.2, 9.6 Hz), 2.75 (dd, 1H, J = 17.2, 8.4 Hz), 3.13-3.24 (m, 1H), 3.41 (s, 1H), 3.50-3.76 (m, 6H), 4.10-4.21 (m, 2H), 4.58 (s, 2H), 6.04 (s, 1H), 6.92-7.13 (m, 4H), 7.19-7.33 (m, 4H), 9.01 (s, 1H).	551	549	S-form
484		¹ H-NMR (CDCl ₃) δ: 1.25 (d, 3H, J = 6.0 Hz), 2.48 (dd, 1H, J = 17.2, 9.6 Hz), 2.76 (dd, 1H, J = 17.2, 8.4 Hz), 3.20-3.30 (m, 1H), 3.52-3.76 (m, 3H), 3.86 (s, 1H), 4.53 (d, 1H, J = 12.0 Hz), 4.65 (d, 1H, J = 12.0 Hz), 6.05 (s, 1H), 6.88-7.31 (m, 8H), 8.95 (s, 1H).	521	519	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
485		¹ H-NMR (CDCl ₃) δ: 1.25 (d, 3H, J = 6.4 Hz), 1.39 (t, 3H, J = 7.2 Hz), 2.49 (dd, 1H, J = 17.2, 9.6 Hz), 2.76 (dd, 1H, J = 17.2, 8.4 Hz), 3.22-3.32 (m, 1H), 3.53-3.76 (m, 3H), 4.07 (q, 2H, J = 7.2 Hz), 4.53 (d, 1H, J = 12.0 Hz), 4.65 (d, 1H, J = 12.0 Hz), 6.04 (s, 1H), 6.87-7.11 (m, 4H), 7.17-7.31 (m, 4H), 8.92 (s, 1H).	535	533	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
486		¹ H-NMR (CDCl ₃) δ: 1.24-1.31 (m, 9H), 2.56 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.4 Hz), 3.28-3.38 (m, 1H), 3.56-3.77 (m, 3H), 4.42-4.68 (m, 3H), 6.01 (s, 1H), 6.88-7.10 (m, 4H), 7.16-7.31 (m, 4H), 8.65 (s, 1H).	549	547	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form





487		¹ H-NMR (CDCl ₃) δ: 1.26 (d, 3H, J = 6.8 Hz), 2.42 (dd, 1H, J = 17.2, 9.6 Hz), 2.75 (dd, 1H, J = 17.2, 8.4 Hz), 3.12-3.23 (m, 1H), 3.41 (s, 3H), 3.49-3.78 (m, 5H), 4.10-4.22 (m, 2H), 4.53 (d, 1H, J = 12.0 Hz), 4.66 (d, 1H, J = 12.0 Hz), 6.10 (s, 1H), 6.91-7.31 (m, 8H), 9.08 (s, 1H).	565	563	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form
488		¹ H-NMR (DMSO-D ₆) δ: 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 3.99 (q, 2H, J = 9.2 Hz), 4.62 (s, 2H), 6.97-6.99 (m, 2H), 7.09 (s, 1H), 7.17 (d, 1H, J = 9.6 Hz), 7.25-7.27 (m, 2H), 7.35-7.44 (m, 3H), 7.64 (s, 1H), 10.89 (s, 1H).	477	475	S-form
489		¹ H-NMR (DMSO-D ₆) δ: 2.39 (d, 2H, J = 8.4 Hz), 3.18 (s, 1H), 3.32-3.52 (m, 3H), 3.98 (q, 2H, J = 9.6 Hz), 4.37 (s, 2H), 4.61 (s, 2H), 6.92 (s, 1H), 7.10 (d, 1H, J = 8.4 Hz), 7.17 (d, 1H, J = 7.2 Hz), 7.23-7.26 (m, 3H), 7.32-7.39 (m, 3H), 7.63 (s, 1H), 10.87 (s, 1H).	503	501	S-form

490		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, J = 7.3 Hz), 1.42-1.51 (m, 2H), 1.84-1.92 (m, 2H), 2.04 (s, 3H), 2.36-2.41 (m, 2H), 2.46-2.56 (m, 4H), 3.31-3.53 (m, 3H), 3.92 (t, 2H, J = 6.2 Hz), 6.75-6.78 (m, 2H), 6.87-6.92 (m, 2H), 7.03 (s, 1H), 7.07-7.31 (m, 4H), 7.63 (s, 1H), 10.85 (s, 1H).	493	491	S-form
491		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, J = 7.3 Hz), 1.30 (s, 3H), 1.40-1.51 (m, 2H), 2.37-2.42 (m, 2H), 2.45-2.55 (m, 2H), 3.31-3.53 (m, 3H), 3.91 (s, 2H), 4.27 (d, 2H, J = 5.8 Hz), 4.41 (d, 2H, J = 5.8 Hz), 6.76-6.96 (m, 4H), 7.03-7.05 (m, 1H), 7.09-7.13 (m, 1H), 7.17-7.21 (m, 1H), 7.25-7.32 (m, 2H), 7.63 (s, 1H), 10.85 (s, 1H).	489	487	S-form
492		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 3H, J = 7.3 Hz), 1.42-1.51 (m, 2H), 2.04-2.13 (m, 2H), 2.36-2.42 (m, 2H), 2.44-2.54 (m, 2H), 3.00 (s, 3H), 3.18-3.24 (m, 2H), 3.30-3.53 (m, 3H), 3.99 (t, 2H, J = 6.2 Hz), 6.71-6.76 (m, 1H), 6.82-6.93 (m, 3H), 7.03 (s, 1H), 7.07-7.10 (m, 1H), 7.16-7.20 (m, 1H), 7.23-7.31 (m, 2H), 7.63 (s, 1H), 10.86 (s, 1H).	525	523	S-form





493		<p>¹H-NMR (DMSO-D₆) δ: 0.77 (t, 3H, J = 7.3 Hz), 1.38-1.48 (m, 2H), 1.53 (s, 3H), 1.81 (s, 3H), 2.36-2.41 (m, 2H), 2.43-2.48 (m, 2H), 3.31-3.53 (m, 3H), 6.17 (s, 1H), 6.88 (s, 1H), 6.97-7.03 (m, 2H), 7.07-7.18 (m, 4H), 7.27 (t, 1H, J = 7.8 Hz), 7.36 (t, 1H, J = 7.8 Hz), 7.63 (s, 1H), 10.83 (s, 1H).</p>	443	441	S-form
494		<p>¹H-NMR (DMSO-D₆) δ: 0.76 (t, 3H, J = 7.3 Hz), 1.39-1.49 (m, 2H), 1.80 (dd, 3H, J = 6.5, 1.2 Hz), 2.37-2.41 (m, 2H), 2.44-2.50 (m, 2H), 3.32-3.53 (m, 3H), 6.12-6.21 (m, 1H), 6.32-6.38 (m, 1H), 6.89 (s, 1H), 7.00-7.10 (m, 3H), 7.16-7.35 (m, 5H), 7.63 (s, 1H), 10.84 (s, 1H).</p>	429	427	S-form
495		<p>¹H-NMR (DMSO-D₆) δ: 0.79 (t, 3H, J = 7.3 Hz), 1.25 (s, 9H), 1.42-1.52 (m, 2H), 2.37-2.41 (m, 2H), 2.45-2.51 (m, 2H), 3.31-3.53 (m, 3H), 6.89 (s, 1H), 7.01-7.04 (m, 1H), 7.09-7.14 (m, 2H), 7.18-7.22 (m, 2H), 7.26-7.35 (m, 3H), 7.63 (s, 1H), 10.84 (s, 1H).</p>	469	467	S-form





496		¹ H-NMR (DMSO-D ₆) δ: 0.72-0.81 (m, 9H), 1.39-1.50 (m, 2H), 1.60-1.71 (m, 1H), 2.35-2.41 (m, 4H), 2.45 (t, 2H, J = 7.4 Hz), 3.31-3.53 (m, 3H), 6.88 (s, 1H), 6.95-7.00 (m, 2H), 7.03-7.18 (m, 4H), 7.22-7.32 (m, 2H), 7.63 (s, 1H), 10.84 (s, 1H).	445	443	S-form
497		¹ H-NMR (DMSO-D ₆) δ: 0.78 (t, 6H, J = 7.2 Hz), 1.39-1.50 (m, 4H), 2.37-2.41 (m, 2H), 2.43-2.51 (m, 4H), 3.31-3.53 (m, 3H), 6.88 (s, 1H), 6.98-7.08 (m, 4H), 7.13-7.18 (m, 2H), 7.23-7.31 (m, 2H), 7.63 (s, 1H), 10.84 (s, 1H).	431	429	S-form
498		¹ H-NMR (DMSO-D ₆) δ: 0.79 (t, 3H, J = 7.3 Hz), 0.87 (s, 9H), 1.21-1.28 (m, 2H), 1.40-1.51 (m, 2H), 2.36-2.50 (m, 6H), 3.31-3.53 (m, 3H), 6.88 (s, 1H), 6.95-7.10 (m, 4H), 7.12-7.20 (m, 2H), 7.25-7.31 (m, 2H), 7.63 (s, 1H), 10.84 (s, 1H).	473	471	S-form
499		¹ H-NMR (DMSO-D ₆) δ: 1.26 (s, 9H), 2.37-2.41 (m, 2H), 3.31-3.52 (m, 3H), 6.92 (s, 1H), 7.06-7.10 (m, 1H), 7.22-7.44 (m, 8H), 7.63 (s, 1H), 10.86 (s, 1H).	427	425	S-form





500		¹ H-NMR (DMSO-D ₆) δ: 0.77 (t, 3H, J = 7.4 Hz), 1.44 (sext, 2H, J = 7.4 Hz), 2.46 (t, 2H, J = 7.4 Hz), 3.61-3.91 (m, 4H), 4.41-4.51 (m, 1H), 6.89 (s, 1H), 6.99-7.02 (m, 1H), 7.07-7.12 (m, 1H), 7.16-7.21 (m, 1H), 7.21-7.26 (m, 2H), 7.26-7.31 (m, 1H), 7.34-7.45 (m, 3H), 8.45-8.50 (m, 1H), 9.85 (br s, 2H), 11.54 (s, 1H).	440	438	S-form
501		¹ H-NMR (DMSO-D ₆) δ: 0.92-1.02 (m, 2H), 2.33-2.45 (m, 2H), 2.85-2.95 (m, 1H), 2.98-3.10 (m, 2H), 3.30-3.53 (m, 3H), 3.72 (d, 2H, J = 5.6 Hz), 6.92 (s, 1H), 7.05-7.10 (m, 1H), 7.20-7.43 (m, 8H), 7.63 (s, 1H), 10.85 (s, 1H).	458	456	S-form
502		¹ H-NMR (DMSO-D ₆) δ: 2.38-2.40 (m, 2H), 3.32-3.35 (m, 1H), 3.40-3.46 (m, 1H), 3.48-3.52 (m, 1H), 7.00 (s, 1H), 7.10 (s, 1H), 7.25-7.39 (m, 6H), 7.53 (t, 1H, J = 8.0 Hz), 7.63 (s, 1H), 10.88 (s, 1H).	449	447	S-form
503		¹ H-NMR (DMSO-D ₆) δ: 2.28-2.35 (m, 2H), 3.20-3.45 (m, 3H), 3.90-4.00 (m, 2H), 4.58-4.63 (m, 2H), 6.48 (d, 1H, J = 2.4 Hz), 7.12-7.15 (m, 2H), 7.28-7.38 (m, 2H), 7.55-7.60 (m, 3H), 7.82 (dd, 1H, J = 9.0 Hz, 5.0 Hz), 10.63 (s, 1H).	511	509	S-form




504		¹ H-NMR (CDCl ₃) δ: 2.62 (dd, 1H, J = 17.2, 9.6 Hz), 2.79 (dd, 1H, J = 17.2, 8.4 Hz), 3.35-3.45 (m, 1H), 3.58-3.74 (m, 2H), 6.37 (s, 1H), 6.99-7.11 (m, 3H), 7.19-7.42 (m, 6H), 8.90 (s, 1H).	465	463	S-form
505		¹ H-NMR (CDCl ₃) δ: 2.62 (dd, 1H, J = 17.2, 9.6 Hz), 2.79 (dd, 1H, J = 17.2, 8.4 Hz), 3.33-3.43 (m, 1H), 3.60-3.79 (m, 4H), 4.62 (s, 2H), 5.85 (s, 1H), 7.01-7.07 (m, 2H), 7.19-7.38 (m, 7H), 8.31 (s, 1H).	493	491	S-form
506		¹ H-NMR (CDCl ₃) δ: 1.29 (d, 3H, J = 6.4 Hz), 2.61 (dd, 1H, J = 17.2, 9.6 Hz), 2.79 (dd, 1H, J = 17.2, 8.4 Hz), 3.32-3.43 (m, 1H), 3.59-3.81 (m, 3H), 4.57 (d, 1H, J = 12.0 Hz), 4.69 (d, 1H, J = 12.0 Hz), 5.98 (s, 1H), 7.01-7.07 (m, 2H), 7.18-7.37 (m, 7H), 8.47 (s, 1H).	507	505	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form
507		¹ H-NMR (CDCl ₃) δ: 2.61 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.4 Hz), 3.32-3.42 (m, 1H), 3.56-3.74 (m, 2H), 6.26 (s, 1H), 7.05-7.10 (m, 2H), 7.13-7.23 (m, 4H), 7.31-7.47 (m, 3H), 8.76 (s, 1H).	465	463	S-form




508		¹ H-NMR (CDCl ₃) δ: 2.61 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.4 Hz), 3.31-3.41 (m, 1H), 3.58-3.81 (m, 4H), 4.62 (s, 2H), 5.86 (s, 1H), 7.04 (s, 1H), 7.15-7.36 (m, 8H), 8.34 (s, 1H).	493	491	S-form
509		¹ H-NMR (DMSO-D ₆) δ: 2.38-2.40 (m, 2H), 3.34-3.36 (m, 1H), 3.38-3.44 (m, 1H), 3.49-3.54 (m, 1H), 4.00 (q, 2H, J = 9.4 Hz), 4.38 (t, 2H, J = 5.8 Hz), 4.62 (s, 2H), 4.70 (t, 2H, J = 6.4 Hz), 5.12-5.14 (m, 1H), 6.50-6.53 (m, 1H), 6.79-6.81 (m, 1H), 6.88-6.92 (m, 2H), 7.16-7.19 (m, 1H), 7.25 (s, 1H), 7.30-7.41 (m, 3H), 7.63 (s, 1H), 10.87 (s, 1H).	531	529	S-form
510		¹ H-NMR (DMSO-D ₆) δ: 2.39-2.41 (m, 2H), 3.35-3.36 (m, 1H), 3.40-3.44 (m, 1H), 3.48-3.52 (m, 1H), 3.99 (q, 2H, J = 9.3 Hz), 4.59 (s, 2H), 6.90 (d, 1H, J = 11.4 Hz), 7.06-7.08 (m, 2H), 7.13 (d, 1H, J = 11.4 Hz), 7.49-7.55 (m, 2H), 7.59-7.64 (m, 3H), 10.86 (s, 1H).	511	509	S-form




511		¹ H-NMR (DMSO-D ₆) δ: 2.38-2.40 (m, 2H), 3.32-3.35 (m, 1H), 3.39-3.46 (m, 1H), 3.49-3.53 (m, 1H), 4.01 (q, 2H, J = 9.2 Hz), 4.64 (s, 2H), 7.00 (s, 1H), 7.07-7.23 (m, 4H), 7.38-7.44 (m, 3H), 7.63 (s, 1H), 10.92 (s, 1H).	511	509	S-form
512		¹ H-NMR (DMSO-D ₆) δ: 2.38-2.40 (m, 2H), 3.33-3.35 (m, 1H), 3.38-3.46 (m, 1H), 3.48-3.52 (m, 1H), 4.01 (q, 2H, J = 9.3 Hz), 4.63 (s, 2H), 6.99 (s, 1H), 7.07 (d, 1H, J = 8.8 Hz), 7.09 (s, 1H), 7.20 (d, 1H, J = 8.8 Hz), 7.26-7.29 (m, 2H), 7.47-7.49 (m, 2H), 7.63 (s, 1H), 10.92 (s, 1H).	511	509	S-form
513		¹ H-NMR (DMSO-D ₆) δ: 1.19 (d, 3H, J = 6.5 Hz), 2.39-2.41 (m, 2H), 3.35-3.37 (m, 1H), 3.39-3.45 (m, 1H), 3.48-3.52 (m, 1H), 4.08-4.15 (m, 1H), 4.59 (s, 2H), 6.92 (d, 1H, J = 9.1 Hz), 7.05-7.07 (m, 2H), 7.12 (d, 1H, J = 9.1 Hz), 7.47-7.56 (m, 2H), 7.59-7.65 (m, 3H), 10.86 (s, 1H).	525	523	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form
514		¹ H-NMR (DMSO-D ₆) δ: 1.20 (d, 3H, J = 6.5 Hz), 2.38-2.40 (m, 2H), 3.33-3.35 (m, 1H), 3.41-3.44 (m, 1H), 3.48-3.52 (m, 1H), 4.10-4.16 (m, 1H), 4.64 (s, 2H), 6.99 (s, 1H), 7.07-7.10 (m, 2H), 7.14-7.22 (m, 2H), 7.38-7.43 (m, 3H), 7.63 (s, 1H), 10.92 (s, 1H).	525	523	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form




515		¹ H-NMR (DMSO-D6) δ : 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 3.75 (s, 3H), 7.03-7.07 (m, 2H), 7.13 (s, 1H), 7.16-7.18 (m, 1H), 7.23 (t, 1H, J = 9.6 Hz), 7.35-7.37 (m, 2H), 7.52 (t, 1H, J = 8.0 Hz), 7.63 (s, 1H), 10.90 (s, 1H).	479	477	S-form
516		¹ H-NMR (DMSO-D6) δ : 1.28 (t, 3H, J = 6.8 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.00 (q, 2H, J = 6.8 Hz), 7.01-7.05 (m, 2H), 7.12-7.14 (m, 2H), 7.22 (t, 1H, J = 9.6 Hz), 7.35-7.37 (m, 2H), 7.52 (t, 1H, J = 8.0 Hz), 7.64 (s, 1H), 10.89 (s, 1H).	493	491	S-form
517		¹ H-NMR (DMSO-D6) δ : 1.19 (d, 6H, J = 6.0 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.55 (sep, 1H, 6.0 Hz), 7.01 (dt, 1H, J = 4.0, 8.8 Hz), 7.05 (s, 1H), 7.08-7.10 (m, 2H), 7.22 (t, 1H, J = 9.6 Hz), 7.34-7.39 (m, 2H), 7.52 (t, 1H, J = 8.0 Hz), 7.63 (s, 1H), 10.89 (s, 1H).	507	505	S-form
518		¹ H-NMR (DMSO-D6) δ : 2.39 (d, 2H, J = 8.4 Hz), 3.28 (s, 3H), 3.32-3.52 (m, 3H), 3.60-3.63 (m, 2H), 4.06-4.09 (m, 2H), 7.04-7.08 (m, 2H), 7.13 (s, 1H), 7.16-7.18 (m, 1H), 7.23 (t, 1H, J = 9.6 Hz), 7.35-7.37 (m, 2H), 7.51 (t, 1H, J = 8.0 Hz), 7.63 (s, 1H), 10.89 (s, 1H).	523	521	S-form




519		¹ H-NMR (CDCl ₃) δ: 2.60 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.8 Hz), 3.30-3.40 (m, 1H), 3.59-3.76 (m, 4H), 4.56 (s, 2H), 5.78 (s, 1H), 6.88 (d, 1H, J = 9.2 Hz), 6.98-7.03 (m, 2H), 7.10-7.17 (m, 2H), 7.23 (t, 1H, J = 8.0 Hz), 7.34-7.44 (m, 2H), 8.29 (s, 1H).	495	493	S-form
520		¹ H-NMR (CDCl ₃) δ: 2.62 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.8 Hz), 3.29-3.43 (m, 1H), 3.60-3.82 (m, 4H), 4.60 (s, 2H), 5.92 (s, 1H), 6.88-7.09 (m, 7H), 7.27-7.34 (m, 1H), 8.37 (s, 1H).	495	493	S-form
521		¹ H-NMR (CDCl ₃) δ: 1.31 (d, 3H, J = 6.4 Hz), 2.62 (dd, 1H, J = 17.2, 9.6 Hz), 2.78 (dd, 1H, J = 17.2, 8.8 Hz), 3.33-3.43 (m, 1H), 3.61-3.82 (m, 3H), 4.56 (d, 1H, J = 12.8 Hz), 4.67 (d, 1H, J = 12.8 Hz), 5.81 (s, 1H), 6.86-7.10 (m, 7H), 7.27-7.34 (m, 1H), 8.26 (s, 1H).	509	507	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form
522		¹ H-NMR (DMSO-D ₆) δ: 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 3.74 (s, 3H), 3.98 (q, 2H, J = 9.2 Hz), 4.61 (s, 1H), 6.98 (s, 1H), 7.02 (dt, 1H, J = 4.0, 8.8 Hz), 7.10-7.13 (m, 1H), 7.16-7.23 (m, 2H), 7.27 (s, 1H), 7.30-7.37 (m, 2H), 7.64 (s, 1H), 10.87 (s, 1H).	507	505	S-form




523		¹ H-NMR (DMSO-D ₆) δ: 1.19 (d, 3H, J = 6.8 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 3.74 (s, 3H), 4.02-4.12 (m, 1H), 4.61 (s, 2H), 6.97 (s, 1H), 7.00-7.04 (m, 1H), 7.11-7.13 (m, 1H), 7.17-7.23 (m, 2H), 7.27 (s, 1H), 7.29-7.36 (m, 2H), 7.63 (s, 1H), 10.86 (s, 1H).	521	519	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form
524		¹ H-NMR (DMSO-D ₆) δ: 2.35-2.45 (m, 2H), 3.28-3.53 (m, 3H), 3.68 (s, 3H), 6.62-6.68 (m, 2H), 6.84 (dt, 1H, J = 11.2 Hz, 2.4 Hz), 7.00 (s, 1H), 7.17 (s, 1H), 7.39-7.43 (m, 2H), 7.55-7.60 (m, 1H), 7.64 (s, 1H), 10.94 (s, 1H).	479	477	S-form
525		¹ H-NMR (DMSO-D ₆) δ: 1.22 (t, 3H, J = 7.0 Hz), 2.35-2.45 (m, 2H), 3.28-3.53 (m, 3H), 3.92 (q, 2H, J = 7.0 Hz), 6.59 (s, 1H), 6.67 (dt, 1H, J = 9.6 Hz, 2.0 Hz), 6.84 (dt, 1H, J = 10.8 Hz, 2.0 Hz), 7.00 (s, 1H), 7.17 (s, 1H), 7.39-7.43 (m, 2H), 7.55-7.59 (m, 1H), 7.64 (s, 1H), 10.93 (s, 1H).	493	491	S-form





526		¹ H-NMR (DMSO-D ₆) δ: 1.12 (d, 6H, J = 6.0 Hz), 2.35-2.58 (m, 2H), 3.28-3.53 (m, 3H), 4.46 (sep, 1H, J = 6.0 Hz), 6.53 (s, 1H), 6.71 (dt, 1H, J = 9.2 Hz, 2.2 Hz), 6.81 (dt, 1H, J = 10.8 Hz, 2.2 Hz), 6.99 (s, 1H), 7.16 (s, 1H), 7.38-7.43 (m, 2H), 7.54-7.59 (m, 1H), 7.63 (s, 1H), 10.92 (s, 1H).	507	505	S-form
527		¹ H-NMR (DMSO-D ₆) δ: 2.35-2.60 (m, 2H), 3.28-3.53 (m, 3H), 3.66 (s, 3H), 4.00 (q, 2H, J = 9.4 Hz), 4.64 (s, 1H), 6.62-6.65 (m, 2H), 6.81 (dt, 1H, J = 11.2 Hz, 2.4 Hz), 6.93 (s, 1H), 7.22-7.26 (m, 1H), 7.29 (s, 1H), 7.36-7.45 (m, 2H), 7.63 (s, 1H), 10.92 (s, 1H).	507	505	S-form
528		¹ H-NMR (DMSO-D ₆) δ: 1.22 (t, 3H, J = 7.0 Hz), 2.35-2.63 (m, 2H), 3.28-3.53 (m, 3H), 3.90 (q, 2H, J = 7.0 Hz), 4.00 (q, 2H, J = 9.2 Hz), 4.64 (s, 2H), 6.58 (s, 1H), 6.64 (dt, 1H, J = 9.6 Hz, 2.0 Hz), 6.79 (dt, 1H, J = 11.2 Hz, 2.0 Hz), 6.92 (s, 1H), 7.22-7.25 (m, 1H), 7.29 (s, 1H), 7.36-7.45 (m, 2H), 7.63 (s, 1H), 10.91 (s, 1H).	521	519	S-form

529		¹ H-NMR (DMSO-D ₆) δ: 1.12 (d, 6H, J = 6.0 Hz), 2.35-2.58 (m, 2H), 3.28-3.53 (m, 3H), 4.00 (q, 2H, J = 9.2 Hz), 4.44 (sep, 1H, J = 6.0 Hz), 4.64 (s, 2H), 6.52 (s, 1H), 6.67 (dt, 1H, J = 10.0 Hz, 2.0 Hz), 6.76 (dt, 1H, J = 11.0 Hz, 2.0 Hz), 6.92 (s, 1H), 7.24 (d, 1H, J = 7.6 Hz), 7.29 (s, 1H), 7.35-7.45 (m, 2H), 7.63 (s, 1H), 10.89 (s, 1H).	535	533	S-form
530		¹ H-NMR (DMSO-D ₆) δ: 2.39-2.41 (m, 3H), 3.15 (s, 3H), 3.32-3.35 (m, 1H), 3.37-3.45 (m, 1H), 3.49-3.53 (m, 1H), 4.02 (q, 2H, J = 9.4 Hz), 4.64 (s, 2H), 6.98 (s, 1H), 7.22-7.24 (m, 1H), 7.31 (s, 1H), 7.38-7.43 (m, 2H), 7.48-7.51 (m, 1H), 7.63 (t, 2H, J = 7.9 Hz), 7.81-7.86 (m, 2H), 10.97 (s, 1H).	537	535	S-form
531		¹ H-NMR (DMSO-D ₆) δ: 1.06 (s, 9H), 2.35-2.42 (m, 2H), 3.32-3.53 (m, 3H), 6.18 (d, 1H, J = 16.4 Hz), 6.27 (d, 1H, J = 16.4), 6.94 (s, 1H), 6.96-7.01 (m, 1H), 7.22-7.28 (m, 3H), 7.31-7.38 (m, 3H), 7.39-7.45 (m, 2H), 7.63 (s, 1H), 10.85 (s, 1H).	429	427	S-form





532		¹ H-NMR (DMSO-D ₆) δ: 0.67-0.73 (m, 2H), 0.83-0.90 (m, 2H), 1.45-1.56 (m, 1H), 2.35-2.40 (m, 2H), 3.32-3.55 (m, 3H), 6.91 (s, 1H), 7.09-7.14 (m, 1H), 7.19-7.26 (m, 3H), 7.27-7.46 (m, 5H), 7.63 (s, 1H), 10.86 (s, 1H).	411	409	S-form
533		¹ H-NMR (DMSO-D ₆) δ: 1.20 (d, 3H, J = 6.4 Hz), 2.35-2.60 (m, 2H), 3.28-3.53 (m, 3H), 3.66 (s, 3H), 4.07-4.15 (m, 1H), 4.61-4.68 (m, 2H), 6.60-6.65 (m, 2H), 6.82 (dt, 1H, J = 10.8 Hz, 2.4 Hz), 6.92 (s, 1H), 7.24 (dt, 1H, J = 7.2 Hz, 1.6 Hz), 7.29 (s, 1H), 7.36-7.45 (m, 2H), 7.63 (s, 1H), 10.91 (s, 1H).	521	519	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form
534		¹ H-NMR (DMSO-D ₆) δ: 1.19-1.24 (m, 6H), 2.35-2.58 (m, 2H), 3.28-3.53 (m, 3H), 3.90 (q, 2H, J = 7.0 Hz), 4.07-4.15 (m, 1H), 4.61-4.68 (m, 2H), 6.58 (s, 1H), 6.64 (dt, 1H, J = 9.2 Hz, 2.2 Hz), 6.79 (dt, 1H, J = 10.8 Hz, 2.2 Hz), 6.92 (s, 1H), 7.23 (d, 1H, J = 7.6 Hz), 7.29 (s, 1H), 7.36-7.45 (m, 2H), 7.63 (s, 1H), 10.90 (s, 1H).	535	533	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form




535		<p>¹H-NMR (DMSO-D₆) δ: 1.12 (d, 6H, J = 6.0 Hz), 1.21(d, 3H, J = 6.4 Hz), 2.35-2.58 (m, 2H), 3.28-3.53 (m, 3H), 4.08-4.16 (m, 1H), 4.44 (sep, 1H, J = 6.0 Hz), 4.61-4.68 (m, 2H), 6.52 (s, 1H), 6.66 (dt, 1H, J = 9.4 Hz, 2.0 Hz), 6.77 (dt, 1H, J = 11.0 Hz, 2.0 Hz), 6.91 (s, 1H), 7.24 (d, 1H, J = 7.6 Hz), 7.28 (s, 1H), 7.35-7.45 (m, 2H), 7.63 (s, 1H), 10.89 (s, 1H).</p>	549	547	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
536		<p>¹H-NMR (DMSO-D₆) δ: 1.81-1.93 (m, 2H), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 3.58-3.73 (m, 4H), 4.05-4.09 (m, 1H), 4.41 (s, 2H), 6.91 (s, 1H), 7.14 (d, 1H, J = 7.2 Hz), 7.20-7.25 (m, 3H), 7.29-7.42 (m, 5H), 7.63 (s, 1H), 10.86 (s, 1H).</p>	447	445	Tetrahydrofuran ring: R-form Lactam ring: S-form
537		<p>¹H-NMR (DMSO-D₆) δ: 1.81-1.93 (m, 2H), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 3.58-3.73 (m, 4H), 4.05-4.09 (m, 1H), 4.41 (s, 2H), 6.91 (s, 1H), 7.14 (d, 1H, J = 7.2 Hz), 7.20-7.25 (m, 3H), 7.29-7.42 (m, 5H), 7.63 (s, 1H), 10.86 (s, 1H).</p>	447	445	Tetrahydrofuran ring: S-form Lactam ring: S-form





538		¹ H-NMR (DMSO-D ₆) δ: 1.44-1.53 (m, 1H), 1.72-1.90 (m, 3H), 2.39 (d, 2H, J = 8.4 Hz), 3.30-3.32 (m, 2H), 3.32-3.52 (m, 3H), 3.58-3.64 (m, 1H), 3.69-3.73 (m, 1H), 3.87-3.93 (m, 1H), 4.43 (s, 2H), 6.91 (s, 1H), 7.16 (d, 1H, J = 7.6 Hz), 7.22-7.24 (m, 3H), 7.28-7.41 (m, 5H), 7.63 (s, 1H), 10.86 (s, 1H).	461	459	Tetrahydropyran ring: R-form Lactam ring: S-form
539		¹ H-NMR (DMSO-D ₆) δ: 1.36-1.46 (m, 2H), 1.63-1.69 (m, 1H), 1.84-1.90 (m, 1H), 2.39 (d, 2H, J = 8.4 Hz), 3.16-3.21 (m, 1H), 3.25-3.31 (m, 1H), 3.32-3.52 (m, 4H), 3.56-3.61 (m, 1H), 3.66-3.70 (m, 1H), 4.46 (s, 2H), 6.91 (s, 1H), 7.12 (d, 1H, J = 7.6 Hz), 7.21-7.25 (m, 3H), 7.28-7.42 (m, 5H), 7.63 (s, 1H), 10.86 (s, 1H).	461	459	Tetrahydropyran ring: S-form Lactam ring: S-form
540		¹ H-NMR (CDCl ₃) δ: 1.15 (s, 9H), 2.59 (dd, 1H, J = 16.8, 9.6 Hz), 2.79 (dd, 1H, J = 16.8, 8.4 Hz), 2.86 (s, 2H), 3.29-3.39 (m, 1H), 3.62 (t, 1H, J = 8.8 Hz), 3.72 (dd, 1H, J = 9.6, 7.6 Hz), 5.74 (s, 1H), 7.13 (s, 1H), 7.20-7.25 (m, 2H), 7.35-7.41 (m, 3H), 7.49-7.55 (m, 2H), 7.71-7.75 (m, 1H), 7.84-7.90 (m, 1H), 8.34 (s, 1H).	481	479	S-form




541		¹ H-NMR (CDCl ₃) δ: 1.28 (t, 3H, J = 7.2 Hz), 2.54 (dd, 1H, J = 16.8, 9.6 Hz), 2.65-2.79 (m, 3H), 3.22-3.33 (m, 1H), 3.55 (t, 1H, J = 8.8 Hz), 3.68 (dd, 1H, J = 9.6, 7.6 Hz), 4.10 (s, 2H), 5.93 (s, 1H), 7.05 (s, 1H), 7.18-7.25 (m, 3H), 7.31-7.44 (m, 6H), 8.64 (s, 1H).	453	451	S-form
542		¹ H-NMR (DMSO-D ₆) δ: 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.33-4.37 (m, 4H), 4.47-4.52 (m, 1H), 4.55-4.59 (m, 2H), 6.92 (s, 1H), 7.13-7.16 (m, 1H), 7.23-7.25 (m, 3H), 7.30-7.43 (m, 5H), 7.63 (s, 1H), 10.86 (s, 1H).	433	431	S-form
543		¹ H-NMR (DMSO-D ₆) δ: 1.32-1.41 (m, 2H), 1.76-1.80 (m, 2H), 2.39 (d, 2H, J = 8.4 Hz), 3.27-3.53 (m, 6H), 3.73-3.78 (m, 2H), 4.46 (s, 2H), 6.91 (s, 1H), 7.13-7.16 (m, 1H), 7.21-7.25 (m, 3H), 7.29-7.42 (m, 5H), 7.63 (s, 1H), 10.86 (s, 1H).	461	459	S-form
544		¹ H-NMR (DMSO-D ₆) δ: 1.22 (d, 3H, J = 6.4 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.15-4.21 (m, 1H), 4.71 (s, 2H), 7.07 (s, 1H), 7.26-7.28 (m, 2H), 7.36-7.45 (m, 4H), 7.57 (s, 1H), 7.64-7.66 (m, 2H), 10.91 (s, 1H).	541	539	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form




545		¹ H-NMR (DMSO-D6) δ: 1.22 (d, 3H, J = 6.4 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.13-4.23 (m, 1H), 4.69 (s, 2H), 6.98 (s, 1H), 7.02 (s, 1H), 7.25-7.27 (m, 2H), 7.31 (s, 1H), 7.36-7.47 (m, 4H), 7.64 (s, 1H), 10.90 (s, 1H).	557	555	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form
546		¹ H-NMR (CDCl ₃) δ: 0.96 (d, 6H, J = 6.8 Hz), 1.66-1.77 (m, 1H), 2.52 (dd, 1H, J = 16.8, 9.6 Hz), 2.60 (d, 2H, J = 6.8 Hz), 2.75 (dd, 1H, J = 16.8, 8.4 Hz), 3.20-3.30 (m, 1H), 3.54 (t, 1H, J = 9.2 Hz), 3.68 (dd, 1H, J = 9.6, 7.2 Hz), 4.10 (s, 2H), 5.93 (s, 1H), 7.02-7.39 (m, 10H), 8.77 (s, 1H).	435	433	S-form
547		¹ H-NMR (DMSO-D6) δ: 1.19 (d, 3H, J = 6.4 Hz), 1.95-2.03 (m, 1H), 2.09-2.26 (m, 2H), 2.29-2.39 (m, 1H), 4.06-4.13 (m, 1H), 4.25-4.28 (m, 1H), 4.61 (s, 2H), 6.92 (s, 1H), 7.15 (d, 1H, J = 6.8 Hz), 7.23-7.25 (m, 3H), 7.32-7.41 (m, 5H), 7.88 (s, 1H), 10.86 (s, 1H).	473	471	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: S-form
548		¹ H-NMR (DMSO-D6) δ: 1.95-2.03 (m, 1H), 2.09-2.26 (m, 2H), 2.29-2.39 (m, 1H), 3.97 (q, 2H, J = 9.2 Hz), 4.25-4.28 (m, 1H), 4.61 (s, 2H), 6.92 (s, 1H), 7.15 (d, 1H, J = 6.8 Hz), 7.23-7.25 (m, 3H), 7.32-7.41 (m, 5H), 7.88 (s, 1H), 10.86 (s, 1H).	459	457	R-form

549		¹ H-NMR (DMSO-D ₆) δ: 1.95-2.03 (m, 1H), 2.09-2.26 (m, 2H), 2.29-2.39 (m, 1H), 3.97 (q, 2H, J = 9.2 Hz), 4.25-4.28 (m, 1H), 4.61 (s, 2H), 6.92 (s, 1H), 7.15 (d, 1H, J = 6.8 Hz), 7.23-7.25 (m, 3H), 7.32-7.41 (m, 5H), 7.88 (s, 1H), 10.86 (s, 1H).	459	457	S-form
550		¹ H-NMR (DMSO-D ₆) δ: 2.40 (d, 2H, J = 8.4 Hz), 3.32-3.54 (m, 3H), 4.95 (q, 2H, J = 10.0 Hz), 7.11 (s, 1H), 7.23-7.28 (m, 2H), 7.35-7.45 (m, 3H), 7.54-7.69 (m, 3H), 7.85-7.94 (m, 2H), 10.93 (s, 1H).	493	491	S-form
551		¹ H-NMR (CDCl ₃) δ: 2.33-2.83 (m, 4H), 3.08-3.15 (m, 2H), 3.30-3.41 (m, 1H), 3.60-3.78 (m, 2H), 5.68 (s, 1H), 7.15 (s, 1H), 7.19-7.25 (m, 2H), 7.36-7.42 (m, 3H), 7.57-7.70 (m, 3H), 7.87 (d, 1H, J = 8.0 Hz), 8.21 (s, 1H).	507	505	S-form
552		¹ H-NMR (CDCl ₃) δ: 2.56 (dd, 1H, J = 16.8, 9.6 Hz), 2.76 (dd, 1H, J = 16.8, 8.4 Hz), 3.25-3.35 (m, 1H), 3.44-3.74 (m, 4H), 4.29 (s, 2H), 5.84 (s, 1H), 7.06 (s, 1H), 7.20-7.26 (m, 3H), 7.30-7.45 (m, 6H), 8.51 (s, 1H).	507	505	S-form





553		¹ H-NMR (CDCl ₃) δ: 1.02 (d, 6H, J = 6.4 Hz), 2.08-2.19 (m, 1H), 2.58 (dd, 1H, J = 16.8, 9.6 Hz), 2.77 (dd, 1H, J = 16.8, 8.4 Hz), 2.84 (d, 2H, J = 6.8 Hz), 3.28-3.38 (m, 1H), 3.60 (t, 1H, J = 8.8 Hz), 3.72 (dd, 1H, J = 9.6, 7.6 Hz), 5.82 (s, 1H), 7.13 (s, 1H), 7.20-7.25 (m, 2H), 7.33-7.42 (m, 3H), 7.49-7.56 (m, 2H), 7.73 (br s, 1H), 7.87 (d, 1H, J = 6.4 Hz), 8.43 (s, 1H).	467	465	S-form
554		¹ H-NMR (DMSO-D ₆) δ: 1.19 (d, 3H, J = 6.4 Hz), 1.95-2.03 (m, 1H), 2.09-2.26 (m, 2H), 2.29-2.39 (m, 1H), 4.07-4.17 (m, 1H), 4.25-4.28 (m, 1H), 4.62 (s, 2H), 6.97-6.98 (m, 2H), 7.10 (s, 1H), 7.17 (d, 1H, J = 6.8 Hz), 7.25-7.27 (m, 2H), 7.35-7.45 (m, 3H), 7.89 (s, 1H), 10.89 (s, 1H).	491	489	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
555		¹ H-NMR (DMSO-D ₆) δ: 1.19 (d, 3H, J = 6.4 Hz), 1.95-2.03 (m, 1H), 2.09-2.26 (m, 2H), 2.29-2.39 (m, 1H), 4.07-4.17 (m, 1H), 4.25-4.28 (m, 1H), 4.62 (s, 2H), 6.97-6.98 (m, 2H), 7.10 (s, 1H), 7.17 (d, 1H, J = 6.8 Hz), 7.25-7.27 (m, 2H), 7.35-7.45 (m, 3H), 7.89 (s, 1H), 10.89 (s, 1H).	491	489	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form




556		¹ H-NMR (DMSO-D ₆) δ: 1.19 (d, 3H, J = 6.4 Hz), 1.95-2.03 (m, 1H), 2.09-2.26 (m, 2H), 2.29-2.39 (m, 1H), 4.06-4.13 (m, 1H), 4.25-4.28 (m, 1H), 4.61 (s, 2H), 6.92 (s, 1H), 7.15 (d, 1H, J = 6.8 Hz), 7.23-7.25 (m, 3H), 7.32-7.41 (m, 5H), 7.88 (s, 1H), 10.86 (s, 1H).	473	471	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: R-form
557		¹ H-NMR (DMSO-D ₆) δ: 1.18 (d, 3H, J = 6.4 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.52 (m, 3H), 4.09-4.16 (m, 1H), 4.66 (s, 2H), 6.90 (s, 1H), 7.22-7.25 (m, 4H), 7.33-7.42 (m, 4H), 7.63 (s, 1H), 10.87 (s, 1H).	491	489	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
558		¹ H-NMR (DMSO-D ₆) δ: 3.33-3.38 (m, 1H), 3.60 (dd, 1H, J = 9.2 Hz, 9.2 Hz), 3.97 (q, 2H, J = 9.2 Hz), 4.30-4.35 (m, 1H), 4.61 (s, 2H), 6.32 (s, 1H), 6.59 (s, 1H), 6.93 (s, 1H), 7.16 (dd, 1H, 9.2 Hz, 2.0 Hz), 7.22-7.26 (m, 3H), 7.31-7.42 (m, 5H), 10.71 (s, 1H).	460	458	S-form
559		¹ H-NMR (DMSO-D ₆) δ: 1.08 (d, 3H, J = 7.6 Hz), 2.50-2.60 (m, 1H), 3.00-3.08 (m, 1H), 3.20-3.25 (m, 1H), 3.42-3.47 (m, 1H), 3.97 (q, 2H, J = 9.2 Hz), 4.61 (s, 2H), 6.93 (s, 1H), 7.16 (dd, 1H, 11.2 Hz, 2.0 Hz), 7.22-7.26 (m, 3H), 7.31-7.42 (m, 5H), 7.66 (s, 1H), 10.95 (s, 1H).	473	471	(3R, 4R) Body





560		¹ H-NMR (DMSO-D6) δ : 1.85-2.20 (m, 2H), 2.31-2.40 (m, 2H), 2.48-2.52 (m, 1H), 2.80-2.90 (m, 1H), 3.27-3.50 (m, 1H), 3.97 (q, 2H, J = 9.2 Hz), 4.60 (s, 2H), 6.92 (s, 1H), 7.15 (dd, 1H, 7.2 Hz, 1.6 Hz), 7.22-7.26 (m, 3H), 7.31-7.42 (m, 5H), 7.49 (s, 1H), 10.83 (s, 1H).	473	471	Optically active substance (The same configuration as Example 5)
561		¹ H-NMR (DMSO-D6) δ : 1.19(d, 3H, J = 6.0 Hz), 2.35-2.58 (m, 2H), 3.28-3.55 (m, 3H), 4.06-4.13 (m, 1H), 4.58-4.65 (m, 2H), 6.91 (s, 1H), 7.15 (dt, 1H, J = 6.8 Hz, 1.6 Hz), 7.22-7.26 (m, 3H), 7.31-7.42 (m, 5H), 7.62 (s, 1H), 10.86 (s, 1H).	473	471	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: R-form
562		¹ H-NMR (DMSO-D6) δ : 1.19(d, 3H, J = 6.0 Hz), 3.34-3.38 (m, 1H), 3.57-3.62 (m, 1H), 4.06-4.13 (m, 1H), 4.30-4.34 (m, 1H), 4.58-4.65 (m, 2H), 6.32 (s, 1H), 6.59 (s, 1H), 6.93 (s, 1H), 7.15 (dt, 1H, J = 8.0 Hz, 2.0 Hz), 7.22-7.26 (m, 3H), 7.31-7.42 (m, 5H), 10.71 (s, 1H).	474	472	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Alpha position of carbonyl group (amide): S-form




563		1H-NMR (DMSO-D6) δ : 1.09 (d, 3H, J = 6.8 Hz), 1.19(d, 3H, J = 6.0 Hz), 2.50-2.60 (m, 1H), 3.00-3.08 (m, 1H), 3.20-3.25 (m, 1H), 3.42-3.47 (m, 1H), 4.06-4.13 (m, 1H), 4.58-4.65 (m, 2H), 6.95 (s, 1H), 7.15 (dt, 1H, J = 7.2 Hz, 1.8 Hz), 7.22-7.26 (m, 3H), 7.31-7.42 (m, 5H), 7.66 (s, 1H), 10.95 (s, 1H).	487	485	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: (3R, 4R) Body
564		1H-NMR (DMSO-D6) δ : 1.09 (d, 3H, J = 7.6 Hz), 1.19(d, 3H, J = 6.0 Hz), 2.50-2.60 (m, 1H), 3.00-3.08 (m, 1H), 3.20-3.25 (m, 1H), 3.42-3.47 (m, 1H), 4.06-4.13 (m, 1H), 4.58-4.65 (m, 2H), 6.95 (s, 1H), 7.15 (dt, 1H, J = 7.2 Hz, 1.8 Hz), 7.22-7.26 (m, 3H), 7.31-7.42 (m, 5H), 7.66 (s, 1H), 10.95 (s, 1H).	487	485	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: (3S, 4S) Body
565		1H-NMR (DMSO-D6) δ : 1.19(d, 3H, J = 6.0 Hz), 1.85-2.05 (m, 2H), 2.15-2.30 (m, 2H), 2.48-2.52 (m, 1H), 2.80-2.90 (m, 1H), 3.27-3.50 (m, 1H), 4.06-4.13 (m, 1H), 4.58-4.65 (m, 2H), 6.91 (s, 1H), 7.15 (dd, 1H, 6.8 Hz, 2.0 Hz), 7.22-7.26 (m, 3H), 7.31-7.42 (m, 5H), 7.49 (s, 1H), 10.83 (s, 1H).	487	485	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: The same configuration as Example 5

566		¹ H-NMR (DMSO-D6) δ: 1.18 (d, 6H, J = 6.8), 2.35-2.42 (m, 2H), 2.87-2.98 (m, 1H), 3.32-3.55 (m, 3H), 4.43 (s, 2H), 6.92 (s, 1H), 7.18-7.28 (m, 3H), 7.32-7.45 (m, 6H), 7.63 (s, 1H), 10.88 (s, 1H).	467	465	S-form
567		¹ H-NMR (DMSO-D6) δ: 1.32(s, 6H), 2.35-2.58 (m, 2H), 3.28-3.55 (m, 3H), 4.54 (s, 2H), 6.95-6.97 (m, 2H), 7.06 (s, 1H), 7.12 (d, 1H, J = 9.6 Hz), 7.24-7.27 (m, 2H), 7.35-7.45 (m, 3H), 7.63 (s, 1H), 10.88 (s, 1H).	505	503	S-form
568		¹ H-NMR (DMSO-D6) δ: 1.01 (d, 6H, J = 6.8 Hz), 2.40 (d, 2H, J = 8.4 Hz), 3.23-3.55 (m, 4H), 7.06 (s, 1H), 7.24-7.85 (m, 10H), 10.91 (s, 1H).	453	451	S-form
569		¹ H-NMR (DMSO-D6) δ: 1.07 (s, 9H), 2.39 (d, 2H, J = 8.4 Hz), 3.33-3.55 (m, 3H), 7.02 (s, 1H), 7.25-7.46 (m, 6H), 7.62-7.85 (m, 4H), 10.91 (s, 1H).	467	465	S-form
570		¹ H-NMR (DMSO-D6) δ: 1.20 (d, 3H, J = 6.2 Hz), 3.36 (dd, 1H, J = 10.2, 5.0 Hz), 3.58 (dd, 1H, J = 10.2, 10.0 Hz), 4.09-4.16 (m, 1H), 4.32 (dd, 1H, J = 10.0, 5.0 Hz), 4.63 (s, 2H), 6.33 (s, 1H), 6.59 (s, 1H), 6.99 (s, 1H), 7.07 (s, 1H), 7.08 (d, 1H, J = 9.0 Hz), 7.20 (d, 1H, J = 9.0 Hz), 7.28 (dd, 2H, J = 6.2, 2.5 Hz), 7.48 (dd, 2H, J = 6.2, 2.5 Hz), 10.76 (s, 1H).	526	524	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Alpha position of carbonyl group (amide): R-form




571		1H-NMR (DMSO-D6) δ : 1.20 (d, 3H, J = 6.2 Hz), 3.51 (dd, 1H, J = 8.8, 5.5 Hz), 3.75 (dd, 1H, J = 9.2, 8.8 Hz), 4.09-4.17 (m, 1H), 4.63 (s, 2H), 5.13 (dd, 1H, J = 9.2, 5.5 Hz), 6.99 (s, 1H), 7.07 (s, 1H), 7.09 (d, 1H, J = 9.5 Hz), 7.20 (d, 1H, J = 9.5 Hz), 7.29 (dd, 2H, J = 6.7, 2.1 Hz), 7.49 (dd, 2H, J = 6.7, 2.1 Hz), 7.75 (s, 1H), 11.11 (s, 1H).	527	525	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Alpha position of carbonyl group (amide): R-form
572		1H-NMR (DMSO-D6) δ : 1.20 (d, 3H, J = 6.2 Hz), 1.86-1.99 (m, 2H), 2.14-2.24 (m, 2H), 2.53-2.61 (m, 2H), 2.78-2.88 (m, 1H), 4.10-4.15 (m, 1H), 4.63 (s, 2H), 6.98 (s, 1H), 7.06 (s, 1H), 7.08 (d, 1H, J = 9.7 Hz), 7.19 (d, 1H, J = 9.7 Hz), 7.27 (dd, 2H, J = 6.7, 2.1 Hz), 7.47-7.49 (m, 3H), 10.89 (s, 1H).	539	537	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Lactam ring: The same configuration as Example 5
573		1H-NMR (DMSO-D6) δ : 2.39 (d, 2H, J = 8.4), 3.33-3.55 (m, 3H), 4.85 (s, 2H), 5.50-5.64 (m, 1H), 7.00 (s, 1H), 7.05-7.13 (m, 2H), 7.18-7.41 (m, 5H), 7.63 (s, 1H), 10.89 (s, 1H).	563	561	S-form
574		1H-NMR (DMSO-D6) δ : 1.34 (s, 6H), 2.34-2.42 (m, 2H), 3.32-3.55 (m, 3H), 4.53 (s, 2H), 6.91 (s, 1H), 7.15-7.21 (m, 2H), 7.22-7.32 (m, 5H), 7.33-7.42 (m, 1H), 7.63 (s, 1H), 10.85 (s, 1H).	505	503	S-form




575		¹ H-NMR (DMSO-D ₆) δ: 1.34 (s, 6H), 2.38-2.40 (m, 2H), 3.33-3.35 (m, 1H), 3.39-3.45 (m, 1H), 3.48-3.53 (m, 1H), 4.54 (s, 2H), 6.92 (s, 1H), 7.19-7.21 (m, 2H), 7.24-7.26 (m, 2H), 7.31 (d, 1H, J = 6.9 Hz), 7.36-7.40 (m, 1H), 7.44-7.47 (m, 2H), 7.63 (s, 1H), 10.89 (s, 1H).	521	519	S-form
576		¹ H-NMR (DMSO-D ₆) δ: 2.36-2.38 (m, 2H), 3.32-3.33 (m, 1H), 3.39-3.43 (m, 1H), 3.47-3.51 (m, 1H), 4.85 (s, 2H), 5.54 (sep, 1H, J = 6.5 Hz), 6.93 (s, 1H), 7.20-7.25 (m, 3H), 7.30 (s, 1H), 7.36-7.45 (m, 4H), 7.61 (s, 1H), 10.88 (s, 1H).	561	559	S-form
577		¹ H-NMR (DMSO-D ₆) δ: 1.19 (d, 3H, J = 6.7 Hz), 2.38-2.41 (m, 2H), 3.33-3.36 (m, 1H), 3.41-3.45 (m, 1H), 3.49-3.53 (m, 1H), 4.10-4.14 (m, 1H), 4.59 (d, 1H, J = 12.7 Hz), 4.61 (d, 1H, J = 12.7 Hz), 7.05 (s, 1H), 7.18-7.19 (m, 2H), 7.32-7.36 (m, 3H), 7.50-7.58 (m, 2H), 7.63 (s, 1H), 10.88 (s, 1H).	525	523	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form




578		¹ H-NMR (DMSO-D ₆) δ: 1.20 (d, 3H, J = 6.5 Hz), 2.38-2.40 (m, 2H), 3.33-3.36 (m, 1H), 3.40-3.44 (m, 1H), 3.49-3.52 (m, 1H), 4.10-4.16 (m, 1H), 4.61 (d, 1H, J = 12.5 Hz), 4.65 (d, 1H, J = 12.5 Hz), 7.00 (s, 1H), 7.04-7.06 (m, 1H), 7.16-7.29 (m, 4H), 7.42-7.47 (m, 2H), 7.63 (s, 1H), 10.92 (s, 1H).	525	523	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
579		¹ H-NMR (DMSO-D ₆) δ: 2.37-2.41 (m, 2H), 3.30-3.37 (m, 1H), 3.37-3.47 (m, 1H), 3.47-3.54 (m, 1H), 4.85 (s, 2H), 5.57 (t, 1H, J = 6.4 Hz), 7.00 (s, 1H), 7.20-7.22 (m, 1H), 7.23-7.30 (m, 3H), 7.34-7.46 (m, 4H), 7.63 (s, 1H), 10.89 (s, 1H).	561	559	S-form
580		¹ H-NMR (DMSO-D ₆) δ: 1.33 (s, 6H), 2.39 (d, 2H, J = 8.4 Hz), 3.31-3.53 (m, 3H), 4.55 (s, 2H), 6.98 (s, 1H), 7.02 (br s, 1H), 7.07 (d, 1H, J = 9.2 Hz), 7.15 (d, 1H, J = 9.2 Hz), 7.27 (dt, 2H, J = 9.2, 2.4 Hz), 7.48 (dt, 2H, J = 9.2, 2.4 Hz), 7.63 (br s, 1H), 10.91 (s, 1H).	539	537	S-form
581		¹ H-NMR (DMSO-D ₆) δ: 1.32 (s, 6H), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.53 (m, 3H), 4.53 (s, 2H), 6.98 (s, 1H), 7.10 (br s, 1H), 7.23-7.37 (m, 6H), 7.63 (br s, 1H), 10.88 (s, 1H).	539	537	S-form




582		¹ H-NMR (DMSO-D ₆) δ: 1.30 (d, 3H, J = 6.4 Hz), 1.78 (d, 2H, J = 10.8 Hz), 2.08 (ddd, 2H, J = 24.1, 12.5, 4.6 Hz), 2.36 (dd, 2H, J = 8.3, 2.1 Hz), 3.20-3.54 (m, 5H), 3.92 (dd, 2H, J = 11.5, 4.0 Hz), 4.20-4.38 (m, 2H), 4.76 (dd, 2H, J = 20.4, 12.0 Hz), 6.59 (s, 1H), 7.39-7.45 (m, 3H), 7.49-7.56 (m, 1H), 7.60 (s, 1H), 10.72 (s, 1H).	481	479	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
583		¹ H-NMR (DMSO-D ₆) δ: 1.31 (d, 3H, J = 6.6 Hz), 1.79 (d, 2H, J = 10.8 Hz), 2.06 (ddd, 2H, J = 4.6, 12.8, 24.8 Hz), 2.30-2.40 (m, 2H), 3.25-3.53 (m, 5H), 3.91 (dd, 2H, J = 3.9, 11.8 Hz), 4.23-4.36 (m, 2H), 4.77 (dd, 2H, J = 12.7, 20.0 Hz), 6.63 (s, 1H), 7.24-7.32 (m, 3H), 7.61 (s, 1H), 10.75 (s, 1H).	499	497	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form
584		¹ H-NMR (DMSO-D ₆) δ: 1.30 (d, 3H, J = 6.4 Hz), 1.79 (d, 2H, J = 9.0 Hz), 1.99-2.13 (m, 2H), 2.29-2.40 (m, 2H), 3.26-3.53 (m, 5H), 3.87-3.3.95 (m, 2H), 4.20-4.34 (m, 2H), 4.77 (dd, 2H, J = 12.5, 21.3 Hz), 6.63 (s, 1H), 7.34-7.40 (m, 1H), 7.46-7.49 (m, 1H), 7.50-7.53 (m, 1H), 7.61 (s, 1H), 10.75 (s, 1H).	515	513	2,2,2-trifluoro methyl-1-methyl-ethoxym ethyl group: R-form Lactam ring: S-form




585		¹ H-NMR (DMSO-D6) δ: 2.38-2.40 (m, 2H), 3.33-3.36 (m, 1H), 3.41-3.46 (m, 1H), 3.49-3.53 (m, 1H), 4.86 (s, 2H), 5.57 (sep, 1H, J = 6.4 Hz), 6.97-7.01 (m, 2H), 7.18-7.27 (m, 4H), 7.35-7.44 (m, 3H), 7.63 (s, 1H), 10.89 (s, 1H).	545	543	S-form
586		¹ H-NMR (DMSO-D6) δ: 2.38-2.40 (m, 2H), 3.33-3.36 (m, 1H), 3.39-3.46 (m, 1H), 3.49-3.53 (m, 1H), 3.99 (q, 2H, J = 9.2 Hz), 4.61 (s, 2H), 7.00 (s, 1H), 7.20-7.27 (m, 4H), 7.35-7.44 (m, 4H), 7.63 (s, 1H), 10.89 (s, 1H).	493	491	S-form
587		¹ H-NMR (DMSO-D6) δ: 2.38-2.40 (m, 2H), 3.34-3.36 (m, 1H), 3.40-3.44 (m, 1H), 3.48-3.52 (m, 1H), 3.99 (q, 2H, J = 9.2 Hz), 4.59 (s, 2H), 7.06 (s, 1H), 7.17-7.20 (m, 2H), 7.32-7.38 (m, 3H), 7.49-7.63 (m, 3H), 10.88 (s, 1H).	511	509	S-form
588		¹ H-NMR (DMSO-D6) δ: 2.38-2.40 (m, 2H), 3.33-3.36 (m, 1H), 3.41-3.46 (m, 1H), 3.48-3.53 (m, 1H), 4.01 (q, 2H, J = 9.3 Hz), 4.62 (s, 2H), 7.00-7.06 (m, 2H), 7.17-7.29 (m, 4H), 7.42-7.46 (m, 2H), 7.63 (s, 1H), 10.92 (s, 1H).	511	509	S-form



589		¹ H-NMR (DMSO-D ₆) δ: 2.38-2.40 (m, 2H), 3.33-3.36 (m, 1H), 3.41-3.44 (m, 1H), 3.48-3.52 (m, 1H), 3.99 (q, 2H, J = 9.3 Hz), 4.60 (s, 2H), 6.99 (s, 1H), 7.15 (s, 1H), 7.24-7.33 (m, 5H), 7.40 (s, 1H), 7.63 (s, 1H), 10.88 (s, 1H).	511	509	S-form
590		¹ H-NMR (DMSO-D ₆) δ: 1.20 (d, 3H, J = 6.2 Hz), 3.35 (dd, 1H, J = 8.8, 5.3 Hz), 3.59 (dd, 1H, J = 9.0, 8.8 Hz), 4.07-4.14 (m, 1H), 4.32 (dd, 1H, J = 9.0, 5.3 Hz), 4.61 (d, 1H, J = 12.6 Hz), 4.65 (d, 1H, J = 12.6 Hz), 6.33 (s, 1H), 6.59 (s, 1H), 6.94 (s, 1H), 7.20-7.27 (m, 4H), 7.34-7.48 (m, 4H), 10.74 (s, 1H).	508	506	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Alpha position of carbonyl group (amide): R-form
591		¹ H-NMR (DMSO-D ₆) δ: 1.20 (d, 3H, J = 6.0 Hz), 3.36 (dd, 1H, J = 8.8, 5.2 Hz), 3.59 (dd, 1H, J = 8.9, 8.8 Hz), 4.07-4.13 (m, 1H), 4.32 (dd, 1H, J = 8.9, 5.2 Hz), 4.60 (d, 1H, J = 12.1 Hz), 4.63 (d, 1H, J = 12.1 Hz), 6.33 (s, 1H), 6.59 (s, 1H), 6.93 (s, 1H), 7.19-7.40 (m, 8H), 10.70 (s, 1H).	492	490	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Alpha position of carbonyl group (amide): R-form

592		¹ H-NMR (DMSO-D ₆) δ : 1.20 (d, 3H, J = 6.8 Hz), 3.36 (dd, 1H, J = 8.6, 6.2 Hz), 3.60 (dd, 1H, J = 9.3, 8.6 Hz), 4.12 (tq, 1H, J = 9.7, 6.0 Hz), 4.32 (dd, 1H, J = 9.3, 6.2 Hz), 4.62 (d, 1H, J = 12.5 Hz), 4.63 (d, 1H, J = 12.5 Hz), 6.33 (s, 1H), 6.59 (s, 1H), 6.95-7.00 (m, 2H), 7.10 (s, 1H), 7.17 (d, 1H, J = 9.2 Hz), 7.24-7.29 (m, 2H), 7.34-7.46 (m, 3H), 10.73 (s, 1H).	492	490	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Alpha position of carbonyl group (amide): R-form
593		¹ H-NMR (DMSO-D ₆) δ : 1.20 (d, 3H, J = 6.4 Hz), 2.39 (d, 2H, J = 8.4 Hz), 3.32-3.54 (m, 3H), 4.08-4.17 (m, 1H), 4.61 (d, 1H, J = 13.2 Hz), 4.63 (d, 1H, J = 13.2 Hz), 6.94-7.00 (m, 2H), 7.09 (s, 1H), 7.17 (d, 1H, J = 9.6 Hz), 7.23-7.29 (m, 2H), 7.34-7.45 (m, 3H), 7.63 (br s, 1H), 10.88 (s, 1H).	491	489	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: S-form Lactam ring: S-form
594		¹ H-NMR (DMSO-D ₆) δ : 1.32 (s, 6H), 3.36 (dd, 1H, J = 8.6, 6.2 Hz), 3.60 (dd, 1H, J = 9.3, 8.6 Hz), 4.32 (dd, 1H, J = 9.3, 6.2 Hz), 4.54 (s, 2H), 6.33 (s, 1H), 6.60 (s, 1H), 6.98 (s, 1H), 7.16 (s, 1H), 7.20 (t, 1H, J = 2.0 Hz), 7.24-7.29 (m, 2H), 7.33-7.46 (m, 4H), 10.74 (s, 1H).	522	520	R-form

595		<p>¹H-NMR (DMSO-D₆) δ: 1.33(s, 6H), 2.35-2.58 (m, 2H), 3.28-3.55 (m, 3H), 4.54 (s, 2H), 6.97 (s, 1H), 7.00 (s, 1H), 7.04 (d, 1H, J = 9.2 Hz), 7.13(d, 1H, J = 9.2 Hz), 7.24-7.35 (m, 4H), 7.63 (s, 1H), 10.87 (s, 1H).</p>	523	521	S-form
596		<p>¹H-NMR (DMSO-D₆) δ: 3.35 (dd, 1H, J = 9.1, 5.4 Hz), 3.59 (dd, 1H, J = 10.1, 9.1 Hz), 4.01 (q, 2H, J = 9.3 Hz), 4.32 (dd, 1H, J = 10.1, 5.4 Hz), 4.63 (s, 2H), 6.33 (s, 1H), 6.59 (s, 1H), 7.00 (s, 1H), 7.05-7.11 (m, 2H), 7.18-7.23 (m, 1H), 7.25-7.31 (m, 2H), 7.46-7.51 (m, 2H), 10.77 (s, 1H).</p>	512	510	R-form
597		<p>¹H-NMR (DMSO-D₆) δ: 1.20 (d, 3H, J = 6.4 Hz), 3.36 (dd, 1H, J = 9.1, 5.4 Hz), 3.60 (dd, 1H, J = 9.7, 9.1 Hz), 4.07-4.18 (m, 1H), 4.33 (dd, 1H, J = 9.7, 5.4 Hz), 4.60 (d, 1H, J = 12.1 Hz), 4.63 (d, 1H, J = 12.1 Hz), 6.33 (s, 1H), 6.59 (s, 1H), 6.99 (s, 1H), 7.19-7.22 (m, 2H), 7.25-7.29 (m, 2H), 7.35-7.46 (m, 4H), 10.73 (s, 1H).</p>	508	506	2,2,2-trifluoro methyl-1-methyl-ethoxymethyl group: R-form Alpha position of carbonyl group (amide): R-form

598		¹ H-NMR (DMSO-D ₆) δ: 1.20 (d, 3H, J = 6.0 Hz), 3.35 (dd, 1H, J = 8.5, 6.0 Hz), 3.60 (dd, 1H, J = 9.3, 8.5 Hz), 4.06-4.17 (m, 1H), 4.32 (dd, 1H, J = 9.3, 6.0 Hz), 4.59 (d, 1H, J = 12.1 Hz), 4.63 (d, 1H, J = 12.1 Hz), 6.33 (s, 1H), 6.59 (s, 1H), 6.99 (s, 1H), 7.13-7.17 (m, 1H), 7.23-7.36 (m, 5H), 7.38-7.41 (m, 1H), 10.73 (s, 1H).	526	524	2,2,2-trifluoroethyl-1-methyl-ethoxymethyl group: R-form Alpha position of carbonyl group (amide): R-form
599		¹ H-NMR (DMSO-D ₆) δ: 1.34 (s, 6H), 3.35 (dd, 1H, J = 9.7, 6.0 Hz), 3.60 (dd, 1H, J = 9.7, 8.9 Hz), 4.32 (dd, 1H, J = 8.9, 6.0 Hz), 4.54 (s, 2H), 6.33 (s, 1H), 6.59 (s, 1H), 6.93 (s, 1H), 7.19-7.23 (m, 2H), 7.23-7.29 (m, 2H), 7.30-7.34 (m, 1H), 7.35-7.41 (m, 1H), 7.44-7.49 (m, 2H), 10.74 (s, 1H).	522	520	R-form
600		¹ H-NMR (DMSO-D ₆) δ: 1.32 (s, 6H), 3.35 (dd, 1H, J = 8.6, 6.2 Hz), 3.60 (dd, 1H, J = 9.3, 8.6 Hz), 4.32 (dd, 1H, J = 9.3, 6.2 Hz), 4.53 (s, 2H), 6.33 (s, 1H), 6.60 (s, 1H), 6.99 (s, 1H), 7.10 (s, 1H), 7.23-7.37 (m, 6H), 10.73 (s, 1H).	540	538	R-form

601		¹ H-NMR (DMSO-D ₆) δ: 1.33 (s, 6H), 3.36 (dd, 1H, J = 9.1, 5.8 Hz), 3.59 (dd, 1H, J = 9.1, 8.7 Hz), 4.32 (dd, 1H, J = 8.7, 5.8 Hz), 4.55 (s, 2H), 6.33 (s, 1H), 6.59 (s, 1H), 6.96-6.99 (m, 2H), 7.06 (s, 1H), 7.13 (d, 1H, J = 8.5 Hz), 7.25-7.28 (m, 2H), 7.37-7.45 (m, 3H), 10.73 (s, 1H).	506	504	R-form
602		¹ H-NMR (DMSO-D ₆) δ: 1.33 (s, 6H), 3.36 (dd, 1H, J = 9.2, 5.5 Hz), 3.60 (dd, 1H, J = 9.2, 9.5 Hz), 4.32 (dd, 1H, J = 9.5, 5.5 Hz), 4.54 (s, 2H), 6.33 (s, 1H), 6.60 (s, 1H), 6.98 (s, 1H), 7.01 (s, 1H), 7.05 (d, 1H, J = 9.7 Hz), 7.14 (d, 1H, J = 9.7 Hz), 7.25-7.33 (m, 4H), 10.73 (s, 1H).	524	522	R-form
603		¹ H-NMR (DMSO-D ₆) δ: 1.32 (s, 6H), 3.35 (dd, 1H, J = 8.7, 5.8 Hz), 3.60 (dd, 1H, J = 9.5, 8.7 Hz), 4.32 (dd, 1H, J = 9.5, 5.8 Hz), 4.55 (s, 2H), 6.32 (s, 1H), 6.59 (s, 1H), 6.99 (s, 1H), 7.02 (s, 1H), 7.08 (d, 1H, J = 10.1 Hz), 7.15 (d, 1H, J = 10.1 Hz), 7.28 (dd, 2H, J = 6.6, 2.2 Hz), 7.48 (dd, 2H, J = 6.6, 2.2 Hz), 10.76 (s, 1H).	540	538	R-form


604		¹ H-NMR (DMSO-D ₆) δ: 1.35 (s, 3H), 3.12-3.17 (m, 2H), 3.47-3.51 (m, 1H), 5.38 (s, 1H), 7.00 (s, 1H), 7.08 (s, 1H), 7.24-7.27 (m, 2H), 7.34-7.44 (m, 5H), 7.52 (t, 1H, J = 8.0 Hz), 7.78 (s, 1H), 10.62 (s, 1H).	461	459	Single Diastereoisomer (racemate) Geometrical isomer of Example 605
605		¹ H-NMR (DMSO-D ₆) δ: 1.14 (s, 3H), 3.27-3.29 (m, 1H), 3.33-3.34 (m, 2H), 5.65 (s, 1H), 7.00 (s, 1H), 7.09 (s, 1H), 7.25-7.27 (m, 2H), 7.34-7.45 (m, 5H), 7.51-7.54 (m, 1H), 7.78 (s, 1H), 10.46 (s, 1H).	461	459	Single Diastereoisomer (racemate) Geometrical isomer of Example 604

[0373] In the table 2, Structure Information A, B, C, D and E (Examples 255, 256, 275, 277, 307, 309, 397, 398, 404 and 405) are the information on the enantiomer derived from an asymmetric carbon atom of the alphaposition of carbonyl group. Said enantiomer was isolated and purified according to the similar method as the Auxiliary step 2 of the Preparation Method 3-1 above.

[0374]

Structure Information A (Examples 255 and 256)



Example 255 was synthesized using a high-polarity component obtained by silicagel thin-layer chromatography (eluent: ethyl acetate) between the following Compound [60] or Compound [61]. Example 256 was synthesized using a low-polarity component.



[0375]

Structure Information B (Examples 275 and 277)

Example 275 was synthesized using a low-polarity component obtained by silicagel thin-layer chromatography (eluent: n-hexane/ethyl acetate=1/3) between the following Compound [62] or Compound [63]. Example 277 was synthesized using a high-polarity component.



**[62]****[63]**

5

[0376]

Structure Information C (Examples 307 and 309)

Example 307 was synthesized using a low-polarity component obtained by silicagel thin-layer chromatography (eluent: n-hexane/ethyl acetate=1/1) between the following Compound [64] or Compound [65]. Example 309 was synthesized using a high-polarity component.

**[64]****[65]**


10

[0377]


Structure Information D (Examples 397 and 398)

Example 397 was synthesized using a low-polarity component obtained by silicagel thin-layer chromatography (eluent: n-hexane/ethyl acetate=3/7) between the following Compound [66] or Compound [67]. Example 398 was synthesized using a high-polarity component.

15

**[66]**


or

**[67]**


[0378]

Structure Information E (Examples 404 and 405)

5 Example 404 was synthesized using a low-polarity component obtained by silicagel thin-layer chromatography (eluent: chloroform/ethyl acetate=1/1) between the following Compound [68] or Compound [69]. Example 405 was synthesized using a high-polarity component.

**[68]**

or

**[69]**

[0379]

[Test Example]

10 SGLT1 inhibitory activity of a test compound (IC_{50} value) was calculated based on an amount of intake into a cell of labeled α -methyl-D-glucopyranoside (^{14}C -AMG) which is transported by SGLT1.

1) Construction of human SGLT1 expression plasmid

15 A DNA fragment containing human SGLT1, wherein NheI recognition and cleavage sequence was added upstream of a Kozac consensus sequence of the vector; and a stop codon (TAG) and a SalI recognition and cleavage sequence was added just downstream of a coding region of the human SGLT1, was amplified by PCR using pCMV6-hSGLT1 (OriGene) as a template. The purified DNA fragment was cleaved by restriction enzymes NheI and SalI, and the obtained fragment was ligated into pcDNA3.1 (+) which was cleaved by NheI and XhoI to construct human SGLT1
20 expression plasmid. The nucleic acid sequence of human SGLT1 inserted into the vector was completely identical to the coding region of human SGLT1 sequence (Accession number NM_000343) registered in GenBank. In addition, the sequence which connects the inserted nucleic

acid sequence to the vector was those to be expected.

[0380]

2) Establishment of stable cell lines expressing human SGLT1

A human SGLT1 expression plasmid pcDNA-hSGLT1 was transfected into CHO-K1 cells by Lipofectamine2000 (Invitrogen). The cells were cultured in the presense of G418 (Nakalai tesque) and drug-resgistant cells were selected. A cell line having the highest S/B ratio of an amount of ^{14}C -AMG uptake per cell treated by solvent to an amount of ^{14}C -AMG uptake per cell treated by phlorizin (SGLT inhibitor) was selected from the drug-resistant cells as a stable cell lines expressing human SGLT1.

[0381]

3) Evaluation of SGLT1 inhibitory activity

The stable cell line expressing human SGLT1 was seeded at 5×10^4 cells/well on BioCoat™ Poly-D-Lysine 96 well plate with Lid (Becton Dickinson and Company) and cultured at 37°C under 5 % CO₂ overnight. The medium was replaced with 100 μL /well of a Na (-) buffer (140mM choline chloride, 2mM KCl, 1mM MgCl₂, 1mM CaCl₂, 10mM HEPES, 5mM Tris, pH7.4) followed by incubation for 20 minutes at 37°C under 5 % CO₂. After removing the Na (-) buffer, a test compound solution which was prepared from Na (+) buffer (140mM NaCl, 2mM KCl, 1mM MgCl₂, 1mM CaCl₂, 10mM HEPES, 5mM Tris, pH7.4) containing BSA was added thereto at 40 μL /well. In addition, Na (+) buffer containing 8kBq of ^{14}C -AMG and 2mM AMG was added thereto at 40 μL /well, and was mixed well. Na (-) buffer containing BSA was added to a blank well at 40 μL /well and additionally adding a Na (+) buffer containing 8kBq of ^{14}C -AMG and 2mM AMG, and was mixed well. Cells were incubated for 1 hour at 37°C under 5 % CO₂. After the reaction, the cells were washed twice with 100 μL /well of iced wash buffer (100mM AMG, 140mM choline chloride, 2mM KCl, 1mM MgCl₂, 1mM CaCl₂, 10mM HEPES, 5mM Tris, pH7.4), then lysed in 50 μL /well of a 0.2N aqueous solution of NaOH. The cell lysate was transferred to OptiPlate96 (Perkin-Elmer) and 100 μL /well of MicroScint-40 (Perkin-Elmer) was dispensed. The radioactivity was counted by TOPCOUNT NXT (Perkin-Elmer).

A data was obtained by deducting the average value (CPM) of blank wells from the average value (CPM) of each treated well. The inhibition ratio (%) was calculated by the following formula: inhibition rate (%) = $[(A-B)/A] \times 100$, where A is a data of the solvent control, and B is a data of a test compound. IC₅₀ values (50% Inhibition Concentration) of a test ccompound was calculated based on two concentrations which have the inhibition rate before as well as after 50% and their inhibition rates. The each result of Examples 1 to 234 and Examples 249 to 605 is shown in the following tables 3 and 4.

[0382]

[Table 3]

Example No.	human SGLT1 IC ₅₀ (uM)	% inhibition
1	0.004	
2	0.0034	
3	0.0062	
4	0.0086	

5	0.0069	
6	0.0072	
7	0.033	
8	0.02	
9	1.6	
10	5.5	
11	0.0046	
12	4.4	
13	0.25	
14	0.012	
15	0.04	
16	11.7	
17	0.066	
18	0.031	
19	0.033	
20	0.15	
21	0.68	
22	0.0047	
23	0.0074	
24	0.13	
25	0.0018	
26	0.0035	
27	0.023	
28	0.052	
29	0.024	
30	0.095	
31	0.27	
32	0.012	
33	0.0061	
34	0.0068	
35	0.015	
36	0.0094	
37	0.016	
38	0.13	
39	0.35	
40	0.65	
41	1.9	
42	0.19	
43	0.011	
44	0.034	
45	0.0043	
46	0.0037	

47	0.42	
48	0.0035	
49	0.012	
50	0.022	
51	0.08	
52	0.0037	
53	0.0038	
54	0.01	
55	0.018	
56	14.3	
57	0.23	
58	3.5	
59	0.044	
60	0.017	
61	0.021	
62	0.47	
63	2.4	
64	0.012	
65	0.04	
66	0.21	
67		29% (0.1 uM)
68	0.0067	
69	0.018	
70	0.0016	
71	0.0035	
72	0.0047	
73	0.013	
74	0.0031	
75	0.0039	
76	0.0017	
77	0.002	
78	0.0077	
79	0.022	
80	0.22	
81		43% (0.3 uM)
82	0.012	
83	0.011	
84	0.0072	
85	0.05	
86	0.054	

87	0.11	
88	0.017	
89	0.14	
90	0.034	
91	0.0018	
92	0.025	
93	0.031	
94	0.01	
95	0.0024	
96	0.37	
97	0.01	
98	0.03	
99	0.073	
100	0.0019	
101	0.0024	
102	0.0022	
103	0.005	
104	0.0098	
105	0.037	
106	0.0068	
107	0.044	
108		45% (0.1 uM)
109	0.065	
110	0.085	
111	0.046	
112	0.083	
113	0.012	
114	0.027	
115	0.041	
116	0.0061	
117	0.0014	
118	9.3	
119	0.57	
120	0.0045	
121	0.011	
122		9% (1 uM)
123	0.005	
124	0.17	
125	0.017	
126	0.0012	

127	0.0014	
128	0.017	
129	0.028	
130	0.0013	
131	0.0063	
132	0.0009	
133	0.011	
134	0.0068	
135	0.011	
136	0.028	
137	0.014	
138	0.0015	
139	0.0017	
140	0.0011	
141	0.0055	
142	0.0092	
143	0.0065	
144	0.0013	
145		43% (1 uM)
146		38% (0.1 uM)
147	0.015	
148	0.01	
149	0.0095	
150	0.0077	
151	0.019	
152	0.0093	
153	0.0033	
154	0.0011	
155	0.019	
156	0.0027	
157	0.013	
158	0.0034	
159	0.017	
160	0.011	
161	0.022	
162	0.019	
163	0.0071	
164	0.0026	
165	0.022	
166	0.0029	

167	0.015	
168	0.036	
169		4% (0.1 uM)
170	0.003	
171	0.0083	
172	0.0055	
173	0.013	
174		9% (1 uM)
175		40% (0.1 uM)
176	0.055	
177	0.056	
178	0.014	
179	0.019	
180		-10% (0.3 uM)
181	0.0043	
182	0.0038	
183	20.1	
184		3% (0.03 uM)
185	0.066	
186	0.075	
187	0.042	
188	0.023	
189	0.2	
190	0.13	
191	0.054	
192		6% (0.3 uM)
193	0.0013	
194	0.0007	
195	0.0083	
196	0.0048	
197	0.18	
198	0.067	
199		40% (0.3 uM)
200	0.12	
201	0.093	

202	0.012	
203	0.0014	
204	0.018	
205	0.0036	
206	0.0048	
207	0.0047	
208	0.0013	
209	0.0092	
210	0.0059	
211	0.017	
212	0.0058	
213	0.014	
214	0.0023	
215	0.0014	
216	0.0087	
217	0.0062	
218		26% (0.1 uM)
219		20% (0.1 uM)
220		16% (0.1 uM)
221		15% (0.1 uM)
222	0.0041	
223	0.0023	
224	0.0063	
225	0.0039	
226	0.0033	
227	0.0021	
228	0.0086	
229	0.0031	
230	0.012	
231	0.0067	
232	0.032	
233	0.02	
234	0.0074	

[0383]

[Table 4]

Example No.	human SGLT1 IC ₅₀ (uM)	% Inhibition
235	0.0034	

236	0.00535	
237	0.00385	
238	0.0079	
239	0.007	
240	0.02	
241	0.00995	
242	0.009	
243	0.014	
244	0.008	
245	0.019	
246	0.0115	
247	0.022	
248	0.025	
249	0.039	
250	0.0034	
251		41% (0.1 uM)
252		7% (0.03 uM)
253	0.014	
254		49% (0.3 uM)
255		7% (0.3 uM)
256		3% (0.3 uM)
257	0.076	
258	0.023	
259		20% (0.1 uM)
260	0.023	
261	0.00377	
262	0.006	
263	0.37	
264	0.039	
265		17% (0.1 uM)
266	0.045	
267	0.081	
268	0.0085	
269	0.092	
270	0.045	
271	0.29	
272	0.037	
273	0.027	
274	0.18	
275		48% (0.1 uM)
276		30% (0.1 uM)
277		48% (0.1 uM)
278		34% (0.1 uM)
279	0.0175	
280	0.0098	
281	0.0215	
282	0.0099	
283	0.11	
284	0.075	
285	0.021	
286	0.03	
287	0.012	

288	0.0115	
289	0.0045	
290	0.0055	
291	0.0051	
292		43% (0.1 uM)
293		27% (0.1 uM)
294	0.0245	
295	0.029	
296		48% (0.1 uM)
297	0.071	
298		38% (0.1 uM)
299	0.033	
300	0.038	
301	0.063	
302	0.0015	
303	0.0047	
304	0.01	
305	0.029	
306	0.013	
307	0.006	
308	0.016	
309	0.049	
310		42% (0.1 uM)
311	0.0175	
312	0.022	
313	0.0195	
314	0.015	
315	0.0155	
316	0.039	
317		40% (0.1 uM)
318	0.0032	
319	0.028	
320	0.003	
321	0.022	
322	0.0047	
323	0.0048	
324	0.0022	
325	0.047	
326	0.015	
327		15% (0.3 uM)
328	0.029	
329		35% (0.3 uM)
330		11% (0.3 uM)
331	0.022	
332	0.022	
333	0.02	
334	0.0063	
335	0.016	
336	0.025	
337	0.032	
338	0.017	
339	0.057	

340	0.016	
341	0.06	
342	0.045	
343	0.0072	
344	0.00335	
345	0.0079	
346	0.0027	
347		11% (0.3 uM)
348		26% (0.3 uM)
349	0.072	
350	0.025	
351	0.032	
352	0.014	
353	0.016	
354	0.079	
355	0.0315	
356	0.011	
357	0.031	
358	0.017	
359	0.039	
360	0.0015	
361		33% (0.1 uM)
362	0.03	
363	0.013	
364	0.0083	
365	0.018	
366	0.02	
367	0.0115	
368	0.056	
369	0.015	
370	0.00635	
371	0.036	
372	0.014	
373	0.0053	
374	0.0088	
375	0.013	
376	0.031	
377	0.0097	
378	0.021	
379	0.00525	
380	0.00675	
381	0.0115	
382	0.012	
383	0.014	
384	0.048	
385	0.007	
386	0.035	
387	0.0125	
388	0.00535	
389	0.015	
390	0.0094	
391	0.00375	
392	0.0057	

393	0.00525	
394	0.015	
395	0.013	
396	0.031	
397	0.016	
398		7% (0.1 uM)
399	0.019	
400	0.015	
401	0.013	
402	0.017	
403	0.012	
404	0.018	
405		29% (0.1 uM)
406	0.00785	
407	0.0076	
408	0.008	
409	0.012	
410	0.022	
411	0.0044	
412	0.01095	
413		48% (0.1 uM)
414	0.01145	
415	0.013	
416	0.015	
417	0.0043	
418	0.00235	
419	0.00475	
420	0.047	
421	0.02	
422	0.014	
423	0.016	
424	0.0071	
425	0.00185	
426	0.00205	
427		49% (0.1 uM)
428		31% (0.1 uM)
429		40% (0.1 uM)
430		8% (0.1 uM)
431		21% (0.1 uM)
432		14% (0.1 uM)
433		8% (0.1 uM)
434	0.021	
435	0.00345	
436	0.019	
437	0.0045	
438	0.0031	
439		45% (0.1 uM)
440	0.037	
441	0.009	
442	0.015	
443	0.011	
444	0.01	

445	0.0067	
446	0.0056	
447	0.0047	
448	0.0043	
449	0.0023	
450	0.0081	
451	0.0033	
452	0.0017	
453	0.0017	
454	0.002	
455	0.0031	
456	0.0034	
457	0.0037	
458	0.0059	
459	0.0048	
460	0.0021	
461	0.0014	
462	0.0034	
463	0.0011	
464	0.0014	
465	0.0017	
466	0.016	
467	0.026	
468	0.017	
469	0.0052	
470	0.007	
471	0.015	
472	0.0047	
473	0.0075	
474	0.0073	
475	0.0031	
476	0.021	
477	0.025	
478	0.036	
479		46% (0.1 uM)
480	0.0068	
481	0.0087	
482	0.012	
483	0.059	
484	0.0042	
485	0.0041	
486	0.0065	
487	0.023	
488	0.018	
489	0.011	
490	0.0076	
491	0.0065	
492	0.019	
493	0.0082	
494	0.0091	
495	0.022	
496	0.0069	
497	0.0033	

498	0.07	
499		4% (0.1 uM)
500		19% (0.1 uM)
501	0.069	
502	0.081	
503		-17% (0.1 uM)
504	0.023	
505	0.011	
506	0.0047	
507		50% (0.1 uM)
508	0.02	
509	0.038	
510	0.0083	
511	0.0079	
512	0.016	
513	0.0032	
514	0.004	
515	0.012	
516	0.0083	
517	0.0038	
518	0.011	
519	0.013	
520	0.018	
521	0.00515	
522	0.0053	
523	0.00215	
524	0.027	
525	0.018	
526	0.0092	
527	0.0085	
528	0.0036	
529	0.0023	
530		40% (0.1 uM)
531	0.03	
532		26% (0.1 uM)
533	0.0021	
534	0.0018	
535	0.0028	
536		15% (0.1 uM)
537		1% (0.1 uM)
538		33% (0.1 uM)
539		35% (0.1 uM)
540	0.048	
541		24% (0.1 uM)
542		18% (0.1 uM)
543		15% (0.1 uM)
544		21% (0.1 uM)
545		9% (0.1 uM)
546	0.0057	
547	0.043	
548		3% (0.1 uM)
549		41% (0.1 uM)

550		29% (0.1 uM)
551		42% (0.1 uM)
552		8% (0.1 uM)
553		31% (0.1 uM)
554		32% (0.1 uM)
555	0.024	
556		23% (0.1 uM)
557		48% (0.1 uM)
558	0.038	
559	0.0077	
560	0.024	
561	0.0055	
562	0.013	
563		57% (0.00265 uM)
564	0.045	
565	0.011	
566		22% (0.3 uM)
567	0.014	
568		27% (0.3 uM)
569		29% (0.1 uM)
570	0.0087	
571	0.0093	
572	0.0051	
573	0.013	
574	0.017	
575	0.012	
576	0.019	
577	0.0067	
578	0.0085	
579	0.023	
580	0.0076	
581	0.012	
582		47% (0.1 uM)
583		48% (0.1 uM)
584	0.084	
585	0.013	
586	0.046	
587	0.04	
588	0.04	
589	0.055	
590	0.013	
591	0.019	
592	0.015	
593	0.01	
594	0.066	
595	0.01	
596	0.035	
597	0.06	
598	0.047	
599	0.027	
600	0.041	
601	0.031	

602	0.028	
603	0.016	
604	N.T	
605	N.T	

[0384]

[Formulations]

The formulation examples of the present invention include the following formulations.

5 However, the present invention is not intended to be limited thereto.

Formulation 1 (Preparation of Capsule)

- | | |
|-------------------------------|-------|
| 1) Compound 1 | 30 mg |
| 2) Microcrystalline cellulose | 10 mg |
| 10 3) Lactose | 19 mg |
| 4) Magnesium stearate | 1 mg |

1), 2), 3) and 4) are mixed to fill in a gelatin capsule.

[0385]

Formulation 2 (Preparation of Tablet)

- | | |
|-----------------------|------|
| 15 1) Compound 1 | 10 g |
| 2) Lactose | 50 g |
| 3) Cornstarch | 15 g |
| 4) Carmellose calcium | 44 g |
| 5) Magnesium stearate | 1 g |

20 The whole amount of 1), 2) and 3) and 30 g of 4) are combined with water, dried in vacuo, and then granulated. The resulting granules are mixed with 14 g of 4) and 1 g of 5), and pressed into tablets by a tableting machine. Then, 1000 tablets wherein Compound 1 (10 mg) is comprised in each tablet are obtained.


25 INDUSTRIAL APPLICABILITY

[0386] The compound or a pharmaceutically acceptable salt thereof in the present invention is useful for the treatment and/or prevention of various diseases or conditions which are expected to be improved by controlling SGLT1 activity because it has SGLT1 inhibitory activity. Such diseases or conditions include, for example, diabetes such as type II diabetes, obesity, diabetic complications (e.g.,

30 retinopathy, nephropathy and neuropathy which are known as microangiopathy, and cerebrovascular disease, ischemic heart disease and arteriosclerosis obliterans which are known as macroangiopathy), hypertrophic cardiomyopathy, ischemic heart disease, cancer and constipation.

CLAIMS

1. A compound or a pharmaceutically acceptable salt thereof represented by the general formula [Ib]:



[Ib]

5

, wherein


Ring Cy is

- (1) C_{6-10} aryl,
 - (2) C_{3-8} cycloalkyl or
 - 10 (3) C_{3-8} cycloalkenyl,
- $n1$ is 0, 1, 2, 3 or 4,

R^{1a} is

- (1) a halogen atom,
- (2) hydroxy,
- 15 (3) carboxy,
- (4) a C_{1-6} alkyl group,
- (5) a C_{2-8} alkenyl group,
- (6) a C_{2-8} alkynyl group,

- (7) a C₁₋₆ alkoxy group,
- (8) a C₃₋₆ cycloalkyl group,
- (9) a halo C₁₋₆ alkyl group,
- (10) a hydroxy C₁₋₆ alkyl group,
- 5 (11) a C₁₋₆ alkoxy C₁₋₆ alkyl group,
- (12) a halo C₁₋₆ alkoxy C₁₋₆ alkyl group,
- (13) a C₁₋₆ alkylsulfonyl C₁₋₆ alkyl group,
- (14) a halo C₁₋₆ alkylsulfonyl C₁₋₆ alkyl group,
- (15) a halo C₁₋₆ alkylamino C₁₋₆ alkyl group,
- 10 (16) a C₃₋₆ cycloalkyl C₂₋₆ alkynyl group,
- (17) a halo C₁₋₆ alkoxy group,
- (18) a carboxy C₁₋₆ alkoxy group,
- (19) a C₁₋₆ alkylsulfanyl group,
- (20) a C₁₋₆ alkylsulfonyl group,
- 15 (21) a halo C₁₋₆ alkylsulfonyl group,
- (22) a C₁₋₈ alkylcarbonyl group,
- (23) a C₁₋₆ alkyloxycarbonyl group,
- (24) a group represented by the Formula:



20 , wherein

R₅ is

- (a) a hydrogen atom or
- (b) a C₁₋₆ alkyl group, and

R₆ is

- (a) a C₁₋₆ alkyl group or
 - 5 (b) a halo C₁₋₆ alkyl group,
- (25) a saturated heterocyclo C₁₋₆ alkyl group (the saturated heterocycle is of 4-membered to 6-membered saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur) or
- (26) a saturated heterocyclo oxy C₁₋₆ alkyl group (the saturated
- 10 heterocycle is of 4-membered to 6-membered saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur),
- , provided when n₁ is 2, 3 or 4, R^{1a} each is the same or different,
- R^{2a} is
- (1) a C₁₋₈ alkyl group,
 - 15 (2) a C₃₋₈ cycloalkyl C₁₋₆ alkyl group,
 - (3) a C₆₋₁₀ aryl C₁₋₆ alkyl group,
 - (4) a saturated heterocyclo C₁₋₆ alkyl group (the saturated heterocycle is of 4-membered to 6-membered saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur),
 - 20 (5) a C₃₋₈ cycloalkyl group,
 - (6) a halo C₁₋₆ alkyl group,
 - (7) a C₁₋₆ alkoxy C₁₋₆ alkyl group,
 - (8) a halo C₃₋₈ cycloalkyl group,

- (9) a C₆₋₁₀ aryl group (the C₆₋₁₀ aryl group is optionally substituted by 1 to 4 substituents selected from
- (a) a halogen atom,
 - (b) hydroxy,
 - 5 (c) a C₁₋₆ alkyl group,
 - (d) a C₂₋₈ alkenyl group,
 - (e) a C₂₋₆ alkynyl group,
 - (f) a C₁₋₆ alkoxy group,
 - (g) a halo C₁₋₆ alkyl group,
 - 10 (h) a C₁₋₆ alkoxy C₁₋₆ alkyl group,
 - (i) a halo C₁₋₆ alkoxy group,
 - (j) a hydroxy C₁₋₆ alkoxy group,
 - (k) a C₁₋₆ alkoxy C₁₋₆ alkoxy group,
 - (l) a carboxy C₁₋₆ alkoxy group,
 - 15 (m) a C₆₋₁₀ aryl C₁₋₆ alkoxy group,
 - (n) a C₁₋₆ alkylsulfanyl C₁₋₆ alkoxy group,
 - (o) a C₁₋₆ alkylsulfonyl C₁₋₆ alkoxy group,
 - (p) a C₆₋₁₀ aryl C₁₋₆ alkoxy C₁₋₆ alkoxy group,
 - (q) a saturated heterocyclo C₁₋₆ alkoxy group (the saturated
 - 20 heterocycle is of 4-membered to 6-membered saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur, and is optionally substituted by 1 to 2 C₁₋₆ alkyl groups),

(r) a saturated heterocyclo oxy group (the saturated heterocycle is a 4-membered to 6-membered saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur),

(s) a C₃₋₈ cycloalkyloxy group,

5 (t) a C₁₋₆ alkoxy carbonyloxy group and


(u) a C₁₋₆ alkylsulfonyl group), or

(10) a saturated heterocycle group (the saturated heterocycle is of 4-membered to 6-membered saturated heterocycle having 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulfur, and is optionally substituted by 1 to 4 C₁₋₆ alkyl groups), and

10

Ring Cy^a is a group selected from

(1) a group represented by the Formula:




, wherein

15 m₂ is 1, 2 or 3,

m₃ is 1, 2 or 3,

n₃ is 1 or 2,

(2) a group represented by the Formula:




, wherein

m4 is 0, 1, 2 or 3,

m5 is 0, 1 or 2,

5 , provided sum of m4 and m5 is 1 or more,

(3) a group represented by the Formula:




, wherein

10 m4 is 0, 1, 2 or 3,

m5 is 0, 1 or 2,

, provided sum of m4 and m5 is 1 or more,

(4) a group represented by the Formula:



, wherein


m6 is 0, 1 or 2,

m7 is 0, 1 or 2,

, provided sum of m6 and m7 is 1 or more,

5

(5) a group represented by the Formula:




, wherein

m6 is 0, 1 or 2,

10 m7 is 0, 1 or 2,

, provided sum of m6 and m7 is 1 or more,

(6) a group represented by the Formula:




15 , wherein

m8 is 1 or 2,

m9 is 1 or 2, and

(7) a group represented by the Formula:



5 , wherein

m8 is 1 or 2,

m9 is 1 or 2,

n2 is 0, 1, 2, 3 or 4,

10 R^{3a} is

(1) hydroxy,

(2) a C₁₋₆ alkyl group or

(3) a hydroxy C₁₋₆ alkyl group,

, provided when n2 is 2, 3 or 4, R^{3a} each is the same or different,

15

R^{3b} is

(1) hydroxy,

(2) a C₁₋₆ alkyl group (which optionally form a C₃₋₆ cycloalkyl group

together with the carbon to which it is attached and the carbon adjacent

20 thereto) or

(3) a hydroxy C₁₋₆ alkyl group, or

(4) when two R^{3b} are attached to the same carbon, they optionally form a C₃₋₆ cycloalkyl group together with the carbon to which they are attached, provided, when n₂ is 2, 3 or 4, each R^{3b} is the same or different, and

5

R^{4b} is

(1) a hydrogen atom,

(2) a C₁₋₆ alkyl group,

(3) a carboxy C₁₋₆ alkyl group,


10 (4) a halo C₁₋₆ alkyl group or

(5) a C₁₋₆ alkoxy C₁₋₆ alkyl group.

2. The compound or a pharmaceutically acceptable salt thereof according to claim 1, wherein

15 Ring Cy^a is selected from

(1) a group represented by the Formula:




, wherein

m₄ is 0, 1, 2 or 3,

20 m₅ is 0, 1 or 2,

, provided sum of m₄ and m₅ is 1 or more,

(2) a group represented by the Formula:




, wherein

5 m6 is 0, 1 or 2,

m7 is 0, 1 or 2,

, provided sum of m6 and m7 is 1 or more,

(3) a group represented by the Formula:



10

, wherein


m6 is 0, 1 or 2,

m7 is 0, 1 or 2,

, provided sum of m6 and m7 is 1 or more,

15

(4) a group represented by the Formula:




, wherein

m8 is 1 or 2,

m9 is 1 or 2, and

5

(5) a group represented by the Formula:



, wherein

m8 is 1 or 2,

10 m9 is 1 or 2,

n2 is 0, 1, 2, 3 or 4,

R^{3a} is

(1) hydroxy,

15 (2) a C₁₋₆ alkyl group or

(3) a hydroxy C₁₋₆ alkyl group,

, provided when n_2 is 2, 3 or 4, each R^{3a} is the same or different,

R^{3b} is

- (1) hydroxy,
- 5 (2) a C_{1-6} alkyl group (which optionally form a C_{3-6} cycloalkyl group together with the carbon to which it is attached and the carbon adjacent thereto) or
- (3) a hydroxy C_{1-6} alkyl group, or
- (4) when two R^{3b} are attached to the same carbon, they optionally form a
- 10 C_{3-6} cycloalkyl group together with the carbon to which they are attached, provided, when n_2 is 2, 3 or 4, each R^{3b} is the same or different, and


R^{4b} is

- (1) a hydrogen atom,
- 15 (2) a C_{1-6} alkyl group,
- (3) a carboxy C_{1-6} alkyl group,
- (4) a halo C_{1-6} alkyl group or
- (5) a C_{1-6} alkoxy C_{1-6} alkyl group.

- 20 3. The compound or a pharmaceutically acceptable salt thereof according to claim 1, wherein

Ring Cy^a is selected from

- (1) a group represented by the Formula:




, wherein,

m4 is 0, 1, 2 or 3,

m5 is 0, 1 or 2,

5 , provided sum of m4 and m5 is 1 or more, and

(2) a group represented by the Formula:



, wherein

10 m6 is 0, 1 or 2,

m7 is 0, 1 or 2,

, provided sum of m6 and m7 is 1 or more,

n2 is 0, 1, 2, 3 or 4,

15 R^{3a} is

(1) hydroxy,

(2) a C₁₋₆ alkyl group or

(3) a hydroxy C₁₋₆ alkyl group,

, provided, when n₂ is 2, 3 or 4, each R^{3a} is the same or different,

R^{3b} is

5 (1) hydroxy,

(2) a C₁₋₆ alkyl group (which optionally form a C₃₋₆ cycloalkyl group together with the carbon to which it is attached and the carbon adjacent thereto) or

(3) a hydroxy C₁₋₆ alkyl group, or

10 (4) when two R^{3b} are attached to the same carbon, they optionally form a C₃₋₆ cycloalkyl group together with the carbon to which they are attached, provided, when n₂ is 2, 3 or 4, each R^{3b} is the same or different, and

R^{4b} is

15 (1) a hydrogen atom,

(2) a C₁₋₆ alkyl group,

(3) a carboxy C₁₋₆ alkyl group,

(4) a halo C₁₋₆ alkyl group or


(5) a C₁₋₆ alkoxy C₁₋₆ alkyl group.

20

4. The compound or a pharmaceutically acceptable salt thereof according to claim 1, wherein

Ring Cy^a is selected from


(1) a group represented by the Formula:



, wherein

$n2$ is 0, 1, 2, 3 or 4, and

5 (2) a group represented by the Formula:



, wherein

$n2$ is 0 or 1,

10 R^{3a} is

(1) hydroxy,

(2) a C₁₋₆ alkyl group or

(3) a hydroxy C₁₋₆ alkyl group, and

15 R^{3b} is

(1) hydroxy,

(2) a C₁₋₆ alkyl group (which optionally form a C₃₋₆ cycloalkyl group together with the carbon to which it is attached and the carbon adjacent thereto) or

(3) a hydroxy C₁₋₆ alkyl group, or

- 5 (4) when two R^{3b} are attached to the same carbon, they optionally form a C₃₋₆ cycloalkyl group together with the carbon to which they are attached, provided, when n₂ is 2, 3 or 4, each R^{3b} is the same or different, and

R^{4b} is

- 10 (1) a hydrogen atom,
(2) a C₁₋₆ alkyl group,
(3) a carboxy C₁₋₆ alkyl group,
(4) a halo C₁₋₆ alkyl group or
(5) a C₁₋₆ alkoxy C₁₋₆ alkyl group.

15

5. The compound or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 4, wherein the ring Cy is C₆₋₁₀ aryl.

6. The compound or a pharmaceutically acceptable salt thereof according
20 to claim 5, wherein the ring Cy is phenyl.

7. The compound or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 6 wherein R^{1a} is

- (1) a halogen atom,

- (2) a C₁₋₆ alkyl group,
 - (3) a C₁₋₆ alkoxy group,
 - (4) a halo C₁₋₆ alkyl group,
 - (5) a C₁₋₆ alkoxy C₁₋₆ alkyl group,
 - 5 (6) a halo C₁₋₆ alkoxy C₁₋₆ alkyl group or
 - (7) a halo C₁₋₆ alkoxy group,
- , provided when n₁ is 2, 3, or 4, R^{1a} each is the same or different.




8. The compound or a pharmaceutically acceptable salt thereof according
- 10 to any one of claims 1 to 7
- , wherein R^{2a} is a C₆₋₁₀ aryl group (the C₆₋₁₀ aryl group is optionally substituted by 1 to 4 substituents selected from
- (a) a halogen atom,
 - (b) a C₁₋₆ alkyl group,
 - 15 (c) a C₁₋₆ alkoxy group and
 - (d) a carboxy C₁₋₆ alkoxy group).

9. The compound or a pharmaceutically acceptable salt thereof according
- to claim 8
- 20 , wherein R^{2a} is a phenyl group (the phenyl group is optionally substituted by 1 to 4 substituents selected from
- (a) a halogen atom,
 - (b) a C₁₋₆ alkyl group,
 - (c) a C₁₋₆ alkoxy group and





(d) a carboxy C₁₋₆ alkoxy group).





10. The compound or pharmaceutically acceptable salt thereof of claim 1, wherein the compound is selected from the following Formulae:

5









385






387







11. The compound or a pharmaceutically acceptable salt thereof of
5 claim 1 represented by the following formula:



12. The compound or a pharmaceutically acceptable salt thereof of
claim 1 represented by the following formula:




13. The compound or a pharmaceutically acceptable salt thereof of claim 1 represented by the following formula:




5

14. The compound or a pharmaceutically acceptable salt thereof of claim 1 represented by the following formula:




10

15. The compound or a pharmaceutically acceptable salt thereof of claim 1 represented by the following formula:




16. The compound or a pharmaceutically acceptable salt thereof of claim 1 represented by the following formula:



5

17. The compound or a pharmaceutically acceptable salt thereof of claim 1 represented by the following formula:



10

18. A pharmaceutical composition which comprises a compound or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 17 and a pharmaceutically acceptable carrier.
- 5 19. A SGLT1 inhibitor which comprises a compound or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 17.
20. An agent for treating or preventing diabetes which comprises a
10 compound or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 17.
21. The agent for treating or preventing diabetes according to claim 20, wherein diabetes is type II diabetes mellitus.
- 15 22. Use of a compound or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 17 for manufacturing a SGLT1 inhibitor.
- 20 23. Use of a compound or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 17 for manufacturing an agent for treating or preventing diabetes.

24. The use according to claim 23 wherein the diabetes is type II diabetes mellitus.