

DESCRIPTION

[Title of Invention] PROCESS FOR PRODUCING PHENYL-SUBSTITUTED HETEROCYCLIC DERIVATIVE THROUGH COUPLING USING TRANSITION METAL CATALYST

[Technical Field]

[0001]

The present invention relates to a process for producing a phenyl-substituted heterocyclic derivative by using a novel coupling process between a phenyl derivative and a heterocyclic derivative by the use of a transition metal catalyst. More particularly, the present invention relates to an excellent process for producing a phenyl-substituted heterocyclic derivative or an intermediate thereof, which is useful, for example, as a xanthine oxidase inhibitor used as a therapeutic agent for gout, hyperuricemia, and the like.

[Background Art]

[0002]

The underlying condition of gout is hyperuricemia and, after remission of the gout attack, remedies to improve hyperuricemia are carried out. The therapeutic agents for hyperuricemia are broadly divided into a uricosuric agent and uric acid synthesis inhibitor (xanthine oxidase inhibitor), and are selected appropriately depending on the state and degree of the condition.

[0003]

The xanthine oxidase (XOD) inhibitors include 2-phenylthiazole derivatives (PTLs 1 to

6, NPL 1), 3-phenylisothiazole derivatives (PTLs 7 and 8), phenylpyrazole derivatives (PTL 9 to 11), 2-phenyloxazole derivatives (PTL 12), and phenyl-heteroaryl derivatives (PTL 13). The production processes described in the patent literatures cited in PTLs 1 to 12 are those wherein heterocyclic rings are formed by production processes comprising a series of consecutive reactions and, thus, involve a large number of steps. The production process described in PLT 13 is a process wherein the skeleton is formed by direct coupling of a phenyl ring with a heterocyclic ring and, thus, involves a small number of steps. However, in this process, it is necessary to prepare boron compounds and therefore results in a higher cost. Thus, this process is not yet satisfactory from a standpoint of a low-cost production process involving a small number of steps.

[0004]

As a process to bind a C-H bond on a heterocyclic ring directly to a phenyl ring without the use of boron compounds, there have been reported coupling reactions by using, as a catalyst, palladium (NPLs 2 to 10), rhodium (NPL 11), iridium (NPL 12), copper (NPL 13), nickel (NPL 14 and 15), cobalt (NPL 16), palladium-copper (NPL 17 to 19), and palladium-silver (NPL 20). Among them, the production process using a nickel catalyst is related to a synthetic process for a phenyl-substituted heterocyclic derivative which is a xanthine oxidase (XOD) inhibitor (NPL 9). However, there has been reported no example where a phenyl-substituted heterocyclic derivative of the present invention was synthesized by using a metal catalyst other than the nickel catalyst. In addition, none of the reactions is satisfactory from a standpoint of substrate restriction, cost, and yield.

[Citation List]

[Patent Literature]

[0005]

[PTL 1] International Publication No. WO 92/09279

[PTL 2] Japanese Patent Laid-Open Publication No. H6-293746

[PTL 3] Japanese Patent Laid-Open Publication No. H6-329647

[PTL 4] Japanese Patent Laid-Open Publication No. H6-345724

[PTL 5] Japanese Patent Laid-Open Publication No. H10-139770

[PTL 6] Japanese Patent Laid-Open Publication No. H11-60552

[PTL 7] Japanese Patent Laid-Open Publication No. S57-85379

[PTL 8] Japanese Patent Laid-Open Publication No. H6-211815

[PTL 9] Japanese Patent Laid-Open Publication No. S59-95272

[PTL 10] International Publication No. WO 98/18765

[PTL 11] Japanese Patent Laid-Open Publication No. H10-310578

[PTL 12] Japanese Patent Laid-Open Publication No. H6-65210

[PTL 13] International Publication No. WO 2007/097403

[Non Patent Literature]

[NPL 1] Heterocycles, 1998:47, 857

[NPL 2] J. Am. Chem. Soc., 2003:125, 1700

[NPL 3] J. Am. Chem. Soc., 2006:128, 16496

[NPL 4] Angew. Chem., Int. Ed. 2007:46, 7996

[NPL 5] J. Org. Chem., 2009:74, 1826

[NPL 6] Org. Lett., 2009:10(13), 2909

[NPL 7] Tetrahedron Letters, 2008:49(6), 1045

[NPL 8] Tetrahedron Letters, 2003:59(30), 5685

[NPL 9] Bull. Chem. Soc. Jpn., 1998:71, 467

- [NPL 10] Chem. A. Eur. J., 2009:15(6), 1337
- [NPL 11] J. Am. Chem. Soc., 2008:130, 14926
- [NPL 12] Chem. Comm., 2004, 1926
- [NPL 13] J. Am. Chem. Soc., 2007:129(41), 12404
- [NPL 14] Org. Lett., 2009:11(8), 1733
- [NPL 15] Org. Lett., 2009:11(8), 1737
- [NPL 16] Org. Lett., 2003:5(20), 3607
- [NPL 17] Tetrahedron, 2007:63(9), 1970
- [NPL 18] Org. Lett., 2004:6(12), 2011
- [NPL 19] J. Am. Chem. Soc., 2003:125, 1700
- [NPL 20] Angew. Chem. Int. Ed., 2007:46, 7996

[Summary of Invention]

[Technical Problem]

[0007]

It is an object of the present invention proposes to provide an excellent short-step production process, which is different from the heretofore known production processes mentioned above, for a phenyl-substituted heterocyclic derivative or its intermediate, the derivative being a xanthine oxidase inhibitor used as a therapeutic agent for gout, hyperuricemia, and the like.

[0008]

[Solution to Problem]

The present inventors conducted a diligent research with the above object and, as a result, found that the phenyl ring of a phenyl derivative and a C-H bond on a heterocyclic

derivative can be coupled directly with high selectivity by use of a transition metal compound.


[0009]

That is, the present invention relates to the following:

[1] A process comprising reacting a compound represented by the following formula (1):

[0010]

[Formula 1]




[0011]

(In the formula (1), R¹ represents a hydrogen atom or halogen atom; R² represents a hydrogen atom, cyano group, nitro group, halogen atom, formyl group, or halomethyl group; A represents a hydrogen atom, C₁ to C₈ alkyl group, C₃ to C₆ cycloalkyl group, phenyl group, fluorine atom (only when X is a bond), or protecting group for a hydroxyl group (only when X is an oxygen atom), wherein A may be substituted by 1 to 3 substituents, such substituent representing a group selected from the group consisting of a halogen atom, C₁ to C₄ alkyl group, C₁ to C₄ alkoxy group, C₁ to C₄ alkylthio group, C₃ to C₆ cycloalkyl group, phenyl group, phenoxy group, and pyridyl group; X represents a bond (only when A is a phenyl group or fluorine atom) or oxygen atom; and Y represents a leaving group)

and a compound represented by the following formula (2):

[0012]

[Formula 2]




[0013]

(In the formula (2), H represents a hydrogen atom; and B represents a group selected from the following formulae:

[0014]

[Formula 3]




[0015]

R^3 represents COOR^{3a} or COR^{3b} ; R^{3a} represents a hydrogen atom, C_1 to C_4 alkyl group, or ester-type protecting group for a carboxyl group; R^{3b} represents an amide-type protecting group for a carboxyl group, the protecting group forming an amide with a neighboring carbonyl group; R^4 represents a hydrogen atom, halogen atom, or C_1 to C_4 alkyl group; and W represents an oxygen atom or sulfur atom)

in the presence of a transition metal compound to produce a phenyl-substituted heterocyclic derivative represented by the following formula (3):

[0016]

[Formula 4]



[0017]

(in the formula (3), A , X , R^1 , and R^2 are the same as defined in the formula (1); and B and R^3 are the same as defined in the formula (2)).

[2] The production process according to [1], wherein A is a C_1 to C_5 alkyl group.

[3] The production process according to [1], wherein A is an isobutyl group.

[4] The production process according to any of [1] to [3], wherein X is an oxygen atom.

[5] The production process according to any of [1] to [4], wherein R^1 is a hydrogen atom.

[6] The production process according to any of [1] to [5], wherein R^2 is a cyano group.


[7] The production process according to any of [1] to [6], wherein Y represents a halogen atom, $-\text{OCO}_2-(C_1 \text{ to } C_4 \text{ alkyl group})$, $-\text{OCO}_2-(\text{phenyl group})$, $-\text{OSO}_2-(C_1 \text{ to } C_4 \text{ alkyl group})$, $-\text{OSO}_2-(\text{phenyl group})$, or a diazonium group, wherein, in Y , the C_1 to C_4 alkyl group may be substituted with 1 to 3 halogen atoms and the phenyl group may be substituted with 1 to 3 halogen atoms or C_1 to C_4 alkyl groups.

[8] The production process according to any of [1] to [7], wherein B is represented by the

following group:

[0018]

[Formula 5]



[0019]

[9] The production process according to any of [1] to [8], wherein R^4 is a methyl group.

[10] The production process according to any of [1] to [9], wherein the transition metal compound is zero-valent copper or a salt of mono-valent copper.

[11] The production process according to any of [1] to [9], wherein the transition metal compound is zero-valent palladium or a salt of mono- or di-valent palladium.

[12] The production process according to any of [1] to [9], wherein the transition metal compound is zero-valent palladium or a salt of di-valent palladium.

[13] The production process according to any of [1] to [9], wherein the transition metal compound is zero-valent cobalt or a salt of di-valent cobalt.

[14] The production process according to any of [1] to [9], wherein the transition metal compound is copper (I) iodide (CuI).

[15] The production process according to any of [1] to [9], wherein the transition metal compound is palladium (II) acetate ($\text{Pd}(\text{OAc})_2$), palladium (II) propionate ($\text{Pd}(\text{O}(\text{C}=\text{O})\text{CH}_2\text{CH}_3)_2$), palladium (II) 2-methylpropanoate ($\text{Pd}(\text{O}(\text{C}=\text{O})\text{CH}(\text{CH}_3)_2)_2$), palladium pivalate ($\text{Pd}(\text{OPiv})_2$), palladium (II) chloride (PdCl_2), palladium (I) bromide

(Pd₂Br₂), or palladium (II) hydroxide (Pd(OH)₂).

[16] The production process according to any of [1] to [9], wherein the transition metal compound is palladium (II) acetate (Pd(OAc)₂), palladium (II) propionate (Pd(O(C=O)CH₂CH₃)₂), palladium (II) 2-methylpropanoate (Pd(O(C=O)CH(CH₃)₂)₂), or palladium pivalate (Pd(OPiv)₂).

[17] The production process according to any of [1] to [9], wherein the transition metal compound is palladium (II) acetate (Pd(OAc)₂), palladium (II) chloride (PdCl₂), or palladium (II) hydroxide (Pd(OH)₂).

[18] The production process according to any of [1] to [9], wherein the transition metal compound is cobalt (II) acetate (Co(OAc)₂).

[19] The production process according to any of [1] to [18], wherein a ligand capable of coordinating to the transition metal compound is additionally present during the course of the reaction.

[20] The production process according to [19], wherein the ligand is triphenylphosphine, tri(tert-butyl)phosphine, di(tert-butyl)methylphosphine, tert-butyldicyclohexylphosphine, di(tert-butyl)cyclohexylphosphine, tri(cyclohexyl)phosphine, 2-dicyclohexylphosphino-2',6'-diisopropoxy-1,1'-biphenyl, 2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl, phenanthroline, 1,1'-bis(diphenylphosphino)ferrocene, or salts thereof.

[21] The production process according to [19], wherein the ligand is triphenylphosphine, tri(tert-butyl)phosphine, di(tert-butyl)methylphosphine, tert-butyldicyclohexylphosphine, di(tert-butyl)cyclohexylphosphine, tri(cyclohexyl)phosphine, phenanthroline, or 1,1'-bis(diphenylphosphino)ferrocene.

[22] The production process according to [19], wherein the ligand is triphenylphosphine,

phenanthroline, or 1,1'-bis(diphenylphosphino)ferrocene.

[23] The production process according to [19], wherein the ligand is a phosphine-type ligand.

[24] The production process according to [23], wherein the ligand is tri(tert-butyl)phosphine, tert-butyldicyclohexylphosphine, di(tert-butyl)methylphosphine, di(tert-butyl)cyclohexylphosphine, tri(cyclohexyl)phosphine, 2-dicyclohexylphosphino-2',6'-diisopropoxy-1,1'-biphenyl, 2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl, or a salt thereof.

[25] The production process according to [23], wherein the ligand is tri(tert-butyl)phosphine, tert-butyldicyclohexylphosphine, di(tert-butyl)methylphosphine, di(tert-butyl)cyclohexylphosphine, or tri(cyclohexyl)phosphine.

[26] The production process according to any of [1] to [25], wherein a base is additionally present during the course of the reaction.

[27] The production process according to [26], wherein the base is lithium tert-butoxide.

[28] The production process according to [26], wherein the base is potassium carbonate or cesium carbonate.

[29] The production process according to any of [1] to [28], wherein a silver salt is additionally present during the course of the reaction.

[30] The production process according to [29], wherein the silver salt is silver carbonate.

[31] The production process according to any of [1] to [30], wherein a C₁ to C₁₂ carboxylic acid or salt thereof is additionally present during the course of the reaction.

[32] The production process according to any of [1] to [30], wherein a C₁ to C₆ carboxylic acid or a salt thereof is additionally present during the course of the reaction.


[33] The production process according to [32], wherein the carboxylic acid or salt thereof is

2-methylpropanoic acid, pivalic acid, or a salt thereof.

[34] The production process according to [32], wherein the carboxylic acid or a salt thereof is pivalic acid.


[0019a]

In one embodiment, the present invention relates to a process comprising reacting a compound represented by the following formula (1):




(In the formula (1), R¹ represents a hydrogen atom or halogen atom; R² represents a hydrogen atom, cyano group, nitro group, halogen atom, formyl group, or halomethyl group; A represents a hydrogen atom, C₁ to C₈ alkyl group, C₃ to C₆ cycloalkyl group, phenyl group, fluorine atom (only when X is a bond), or protecting group for a hydroxyl group (only when X is an oxygen atom), wherein A may be substituted by 1 to 3 substituents, such substituent representing a group selected from the group consisting of a halogen atom, C₁ to C₄ alkyl group, C₁ to C₄ alkoxy group, C₁ to C₄ alkylthio group, C₃ to C₆ cycloalkyl group, phenyl group, phenoxy group, and pyridyl group; X represents a bond (only when A is a phenyl group or fluorine atom) or oxygen atom; and Y represents a leaving group)

and a compound represented by the following formula (2):



(In the formula (2), H represents a hydrogen atom;

R^3 represents COOR^{3a} or COR^{3b} ; R^{3a} represents a hydrogen atom, C_1 to C_4 alkyl group, or ester-type protecting group for a carboxyl group; R^{3b} represents an amide-type protecting group for a carboxyl group, the protecting group forming an amide with a neighboring carbonyl group; R^4 represents a halogen atom, or C_1 to C_4 alkyl group in the presence of (i) a transition metal compound selected from the group consisting of zero-valent palladium and a salt of mon- and di-valent palladium, and (ii) a phosphine-type ligand capable of coordinating to the transition metal compound to produce a phenyl-substituted heterocyclic derivative represented by the following formula (3):



(in the formula (3), A, X, R^1 , and R^2 are the same as defined in the formula (1); and

R^3 and R^4 are the same as defined in the formula (2)).

[Advantageous Effects of Invention]

[0020]

According to the present invention, a phenyl-substituted heterocyclic derivative (a compound represented by the formula (3)) can be obtained in a process involving a small number of steps by subjecting a phenyl derivative (a compound represented by the formula (1)) and a heterocyclic derivative (a compound represented by the formula (2)) to a selective coupling reaction by using a transition metal catalyst.

[0021]

Furthermore, since the process involves a small number of steps, it is possible to produce the phenyl-substituted heterocyclic derivative (a compound represented by the formula (3)) in high yield and at low cost.

[Description of Embodiments]

[0022]

The terms used singly or in combination in the present description will be explained in the following. Unless otherwise noted, explanation of each substituent shall be common to each position. In addition, a combination of substituents and variables is allowed only when such a combination results in a chemically stable compound. When the substituent itself is substituted with two or more groups, these many groups can exist on the same or different carbon atom as long as a stable structure is formed.

[0023]

In the present invention, the “halogen atom” means a fluorine atom, chlorine atom, bromine atom, or iodine atom.

[0024]

In the present invention, the “C₁ to C₈ alkyl group” means a linear or branched saturated aliphatic hydrocarbon group having 1 to 8 carbon atoms and includes, for example, a methyl group, ethyl group, n-propyl group, isopropyl group, n-butyl group, isobutyl group, tert-butyl group, n-pentyl group, isopentyl group, neopentyl group, 1-methylpropyl group, n-hexyl group, isohexyl group, 1,1-dimethylbutyl group, 2,2-dimethylbutyl group, 3,3-dimethylbutyl group, n-heptyl group, n-octyl group, and the like.

[0025]

In the present invention, the “C₁ to C₄ alkoxy group” means a group consisting of the “C₁ to C₄ alkyl group” and an oxy group. The examples include a methoxy group, ethoxy group, n-propyloxy group, isopropyloxy group, n-butyloxy group, isobutyloxy group, tert-butyloxy group, and the like.

[0026]

In the present invention, the “C₃ to C₆ cycloalkyl group” means a cyclic alkyl group having 3 to 6 carbon atoms and includes, for example, a cyclopropyl group, cyclobutyl group, cyclopentyl group, cyclohexyl group, and the like.

[0027]

In the present invention, the “C₁ to C₄ alkylthio group” means a group consisting of the “C₁ to C₄ alkyl group” and a thio group. The examples include a methylthio group, ethylthio group, n-propylthio group, isopropylthio group, n-butylthio group, isobutylthio group, tert-butylthio group, and the like.

[0028]

In the present invention, the “halomethyl group” means a methyl group substituted with one or more halogen atoms and includes, for example, a trifluoromethyl group, difluoromethyl group, fluoromethyl group, trichloromethyl group, dichloromethyl group, chloromethyl group, tribromomethyl group, dibromomethyl group, bromomethyl group, and the like.

[0029]

In the present invention, the “leaving group” means an atom or a group of atoms which departs from the reaction substrate in a substitution reaction, elimination reaction, or the like. Such a leaving group includes, for example, a halogen atom, $-\text{OCO}_2-(\text{C}_1 \text{ to } \text{C}_4 \text{ alkyl group})$, $-\text{OCO}_2-(\text{phenyl group})$, $-\text{OSO}_2-(\text{C}_1 \text{ to } \text{C}_4 \text{ alkyl group})$, $-\text{OSO}_2-(\text{phenyl group})$, diazonium group ($-\text{N}^+\equiv\text{N}$), or the like. Furthermore, the C_1 to C_4 alkyl group and the phenyl group that constitute the leaving group may be substituted with 1 to 3 halogen atoms and with 1 to 3 halogen atoms or C_1 to C_4 alkyl groups, respectively. However, the present invention is not limited to these.

[0030]

The “protecting group for a hydroxyl group” means a group which protects the hydroxyl group. Such a “protecting group for a hydroxyl group” is well known in the art and is classified into an ether-type protecting group, silyl ether-type protecting group, ester-type protecting group, carbonate-type protecting group, phosphine-type protecting group, sulfonate-type protecting group, and the like. Examples include the groups described as a protecting group for phenol and the like in T. W. Greene and P. G. M. Wuts, “Protective Groups in Organic Synthesis (3rd Ed., 1994), (4th Ed., 2006)” and the like, such as a benzyloxymethyl group, methoxyethoxymethyl group, phenylthiomethyl group,

phenacylmethyl group, 4-bromophenacylmethyl group, cyclopropylmethyl group, allyl group, propargyl group, cyclohexyl group, benzyl group, ortho-nitrobenzyl group, 4-(dimethylamino)carbonylbenzyl group, 4-methylsulfinylbenzyl group, 9-anthranylmethyl group, 4-picoryl group, trimethylsilyl group, tert-butyldimethylsilyl group, tert-butyldiphenylsilyl group, triisopropylsilyl group, formyl group, $-(C=O)-(C_1 \text{ to } C_4 \text{ alkyl group})$, benzoyl group, 4-oxopentanoyl group, pivaloyl group, methyl ester group, 1-adamantyloxycarbonyl group, tert-butoxycarbonyl group, 4-methylsulfinylbenzyloxycarbonyl group, 2,4-dimethylpent-3-yloxycarbonyl group, 2,2,2-trichloroethoxycarbonyl group, vinyloxycarbonyl group, benzyloxycarbonyl group, $-(C=O)NH-(C_1 \text{ to } C_4 \text{ alkyl group})$, methanesulfonyl group, toluenesulfonyl group, and the like. However, the present invention is not limited to the groups exemplified herein and any group may be selected as long as it is used as a protecting group for the hydroxyl group. Here, the protecting group for the hydroxyl group as A is used as such when X is an oxygen atom. For example, when a benzyl group is the protecting group, A-X- corresponds to $PhCH_2-O-$.

[0031]

The “ester-type protecting group for a carboxyl group” in the present invention means a group which protects the carboxyl group by combining with the oxygen atom of the carboxyl group to be protected and forming an ester. Such an “ester-type protecting group for a carboxyl group” includes the groups described as an ester-type protecting group for the carboxyl group in T. W. Greene and P. G. M. Wuts, “Protective Groups in Organic Synthesis (3rd Ed., 1994), (4th Ed., 2006),” such as a C_1 to C_6 alkyl group, 9-fluorenylmethyl group, methoxymethyl group, methylthiomethyl group, tetrahydropyranyl group, tetrahydrofuranyl group, methoxyethoxymethyl group,

2-(trimethylsilyl)ethoxymethyl group, benzyloxymethyl group, pivaloyloxymethyl group, phenylacetoxymethyl group, triisopropylsilylmethyl group, para-bromophenacyl group, α -methylphenacyl group, para-methoxyphenacyl group, decyl group, carboxamidomethyl group, para-azobenzenecarboxamidomethyl group, N-phthalimidomethyl group, 2,2,2-trichloroethyl group, 2-haloethyl group, ω -chloroalkyl group, 2-(triethylsilyl)ethyl group, 2-methylthioethyl group, 1,3-dithianyl-2-methyl group, 2-(para-nitrophenylsulfenyl)ethyl group, 2-(para-toluenesulfonyl)ethyl group, 2-(2'-pyridyl)ethyl group, 2-(para-methoxyphenyl)ethyl group, 2-(diphenylphosphino)ethyl group, 1-methyl-1-phenylethyl group, 2-(4-acetyl-2-nitrophenyl)ethyl group, 2-cyanoethyl group, dicyclopropylmethyl group, cyclopentyl group, cyclohexyl group, allyl group, methallyl group, 2-methylbut-3-en-2-yl group, 3-methylbut-2-(prenyl) group, 3-buten-1-yl group, 4-(trimethylsilyl)-2-buten-1-yl group, cinnamyl group, α -methylcinnamyl group, prop-2-ynyl (propargyl) group, phenyl group, 2,6-dimethylphenyl group, 2,6-diisopropylphenyl group, 2,6-di(tert-butyl)-4-methylphenyl group, 2,6-di(tert-butyl)-4-methoxyphenyl group, para-(methylthio)phenyl group, pentafluorophenyl group, benzyl group, triphenylmethyl group, diphenylmethyl group, bis(ortho-nitrophenyl)methyl group, 9-anthranylmethyl group, 2-(9,10-dioxo)anthranylmethyl group, 5-dibenzosuberyl group, 1-pyrenylmethyl group, 2-(trifluoromethyl)-6-chromonylmethyl group, 2,4,6-trimethylbenzyl group, para-bromobenzyl group, ortho-nitrobenzyl group, para-nitrobenzyl group, para-methoxybenzyl group, 2,6-dimethoxybenzyl group, 4-(methylsulfinyl)benzyl group, 4-sulfobenzyl group, 4-azidomethoxybenzyl group, piperonyl group, 4-piconyl group, para-benzyl group, trimethylsilyl group, triethylsilyl group, tert-butyldimethylsilyl group, isopropyldimethylsilyl group, phenyldimethylsilyl group, di(tert-butyl)methylsilyl group,

triisopropylsilyl group, C₁ to C₆ alkylthio group, oxazole group, 2-alkyl-1,3-oxazoline group, 4-alkyl-5-oxo-1,3-oxazolidine group, 2,2-bistrifluoromethyl-4-alkyl-5-oxo-1,3-oxazolidine group, 5-alkyl-4-oxo-1,3-dioxolane group, dioxanone group, and the like. However, the present invention is not limited to the groups exemplified herein and any group may be selected as long as it is used as a protecting group for the carboxyl group.

[0032]

The “amide-type protecting group for a carboxyl group” in the present invention means a group which protects the carboxyl group by combining with the carbonyl carbon atom of the carboxyl group to be protected and forming an amide. Such an “amide-type protecting group for a carboxyl group” includes the groups described as a protecting group for the carboxyl group in T. W. Greene and P. G. M. Wuts, “Protective Groups in Organic Synthesis (3rd Ed., 1994), (4th Ed., 2006),” such as an N,N-dimethylamino group, pyrrolidinyl group, piperidinyl group, 5,6-dihydrophenanthridyl group, ortho-nitrophenylamino group, N-7-nitroindolyl group, N-8-nitro-1,2,3,4-tetrahydroquinolyl group, N-phenylhydrazyl group, N,N'-diisopropylhydrazyl group, and the like. However, the present invention is not limited to the groups exemplified herein and any amino group may be selected as long as it is used as a protecting group for the carboxyl group.

[0033]

In the present invention, “C” such as in “C₁” and the like represents a carbon atom and the numeral which follows it represents the number of the carbon atoms. For example, “C₁ to C₆” represents a range of the number of carbon atoms from 1 to 6. Naturally, in the present invention, when the number of carbon atoms is different, the corresponding group is different. For example, a “C₁ to C₄ alkyl group” means a group having 1 to 4 carbon atoms

among the alkyl groups defined by the “C₁ to C₈ alkyl group.” Treatment of the number of carbon atoms in other groups is the same.

[0034]

The “diazonium group” in the present invention may form a salt. Such a salt includes a fluoride salt, chloride salt, bromide salt, iodide salt, tetrafluoroborate salt, and the like.

[0035]


Abbreviations used in the present invention are as follows:

TfO: trifluoromethanesulfonyloxy group, MsO: methanesulfonyloxy group, TsO: toluenesulfonyloxy group, Me: methyl group, Et: ethyl group, n-Pr: n-propyl group, i-Pr: isopropyl group, i-Bu: isobutyl group, t-Bu: tert-butyl group, MeO: methoxy group, Ph: phenyl group, OAc: acetyloxy group, 4-MeO-Ph: 4-methoxy-phenyl group, Cy: cyclohexyl group, Piv: pivaloyl group.

The present invention relates to a process comprising reacting a compound represented by the following formula (1):

[0036]

[Formula 6]




(1)

[0037]

and a compound represented by the following formula (2):

[0038]

[Formula 7]




[0039]

in the presence of a transition metal compound to produce a phenyl-substituted heterocyclic derivative represented by the following formula (3):

[0040]

[Formula 8]



[0002]

In the formulae (1) and (3), R^1 represents a hydrogen atom or halogen atom.

[0003]

The "halogen atom" in R^1 is preferably a chlorine atom or fluorine atom, more preferably a fluorine atom.

[0043]

As a whole, R^1 is preferably a hydrogen atom.

[0044]

In the formulae (1) and (3), R^2 represents a hydrogen atom, cyano group, nitro group, halogen atom, formyl group, or halomethyl group.

[0045]

The "halogen atom" in R^2 is preferably a bromine atom.

[0046]

The "halomethyl group" in R^2 is preferably a chloromethyl group, dichloromethyl group, trichloromethyl group, or trifluoromethyl group.

[0047]

As a whole, R^2 is preferably a cyano group, nitro group, or formyl group and, above all, a cyano group is preferable.

[0048]

In the formulae (1) and (3), A represents a hydrogen atom, C_1 to C_8 alkyl group, C_3 to C_6 cycloalkyl group, phenyl group, fluorine atom (only when X is a bond), or protecting group for a hydroxyl group (only when X is an oxygen atom). Here, the protecting group for a hydroxyl group as A is used as such when X is an oxygen atom. For example, when the protecting group is a benzyl group, A-X- represents $PhCH_2-O-$.

[0049]

Furthermore, A may be substituted with 1 to 3 substituents, such substituent representing a group selected from the group consisting of a halogen atom, C_1 to C_4 alkyl

group, C₁ to C₄ alkoxy group, C₁ to C₄ alkylthio group, C₃ to C₆ cycloalkyl group, phenyl group, phenoxy group, and pyridyl group.

[0050]

The "C₁ to C₈ alkyl group" in A is preferably a methyl group, ethyl group, n-propyl group, isopropyl group, n-butyl group, isobutyl group, tert-butyl group, n-pentyl group, isopentyl group, or neopentyl group. Above all, preferable is an isobutyl group or neopentyl group; more preferable is an isobutyl group.

[0051]

As a whole, A is preferably a C₁ to C₅ alkyl group.

[0052]

In the formulae (1) and (3), X represents a bond (only when A is a phenyl group or fluorine atom) or an oxygen atom. Above all, an oxygen atom is preferable.

[0053]

In the formula (1), Y represents a leaving group. Above all, preferable is a halogen atom, -OCO₂-(C₁ to C₄ alkyl group), -OCO₂-(phenyl group), -OSO₂-(C₁ to C₄ alkyl group), -OSO₂-(phenyl group), or a diazonium group.

[0054]

When the leaving group as Y is "-OCO₂-(C₁ to C₄ alkyl group)" or "-OSO₂-(C₁ to C₄ alkyl group)," such a "C₁ to C₄ alkyl group" in Y is preferably a methyl group.

[0055]

When the leaving group as Y is "-OCO₂-(C₁ to C₄ alkyl group)" or "-OSO₂-(C₁ to C₄ alkyl group)," such a "C₁ to C₄ alkyl group" in Y may be substituted with 1 to 3 halogen atoms. Such a "halogen atom" is preferably a fluorine atom. It is particularly preferable that the group is substituted with three fluorine atoms.

[0056]

When the leaving group as Y is “-OCO₂-(phenyl group)” or “-OSO₂-(phenyl group),” such a “phenyl group” in Y may be substituted with 1 to 3 halogen atoms or C₁ to C₄ alkyl groups. Such a “C₁ to C₄ alkyl group” is preferably a methyl group.

[0057]

When the leaving group as Y is a “halogen atom,” the “halogen atom” is preferably an iodine atom, bromine atom, or chlorine atom. Above all, an iodine atom or bromine atom is preferable.

[0058]

The “diazonium group” may form a salt. When the leaving group as Y represents a “diazonium group,” the salt of the “diazonium group” is preferably a tetrafluoroborate.

[0059]

As a whole, Y is preferably an iodine atom, bromine atom, trifluoromethanesulfonyloxy group, or the like.

[0060]


In the formula (2), H represents a hydrogen atom.

[0061]

In the formulae (2) and (3), B represents a group selected from the following formulae. Meanwhile, the bond at the right-hand side of each of the following formulae is for bonding to R³; and the bond at the left-hand side is for bonding to a hydrogen atom in the formula (2) and to a phenyl group in the formula (3):

[0062]

[Formula 9]




[0063]

Above all, the following group is preferable:

[0064]

[Formula 10]



[0065]

In the formulae (2) and (3), R³ represents COOR^{3a} or COR^{3b}.

[0066]

R^{3a} represents a hydrogen atom, C₁ to C₄ alkyl group, or ester-type protecting group for a carboxyl group. Here, the ester-type protecting group for a carboxyl group as R^{3a} protects the carboxyl group which R^{3a} substitutes.

[0067]

As R^{3a}, preferable is a hydrogen atom or C₁ to C₄ alkyl group.

[0068]

R^{3b} represents an amide-type protecting group for a carboxyl group which forms an amide with a neighboring carbonyl group.

[0069]

As a whole, R^3 is preferably $COOR^{3a}$.

[0070]

In the formulae (2) and (3), R^4 represents a hydrogen atom, halogen atom, or C_1 to C_4 alkyl group.

[0071]

The "halogen atom" in R^4 is preferably a fluorine atom.

[0072]

The " C_1 to C_4 alkyl group" in R^4 is preferably a methyl group.

[0073]

As a whole, R^4 is preferably a C_1 to C_4 alkyl group. Above all, a methyl group is preferable.

[0074]

In the formulae (2) and (3), W represents an oxygen atom or sulfur atom.

[0075]

In the formula (3), definition of A , X , R^1 , and R^2 and preferable groups thereof are respectively the same as those in the formula (1); and definition of B and R^3 and preferable groups thereof are respectively the same as those in the formula (2).


[0076]

Specific examples of the compounds represented by the formula (1) and specific examples of the compounds represented by formula (2) are listed in Tables 1 to 4 and

Tables 5 to 7, respectively. However, the compounds represented by the formula (1) and formula (3) are not limited to such specific examples.

[0077]


[Table 1]



No.	A	X	R ¹	R ²	Y
1	i-Bu	O	H	CN	I
2	i-Bu	O	H	CN	Br
3	i-Bu	O	H	CN	Cl
4	i-Bu	O	H	CN	OTf
5	i-Bu	O	H	CN	OMs
6	i-Bu	O	H	CN	OTs
7	i-Bu	O	H	CHO	I
8	i-Bu	O	H	CHO	Br
9	i-Bu	O	H	CHO	Cl
10	i-Bu	O	H	CHO	OTf
11	i-Bu	O	H	CHO	OMs
12	i-Bu	O	H	CHO	OTs
13	i-Bu	O	H	NO ₂	I
14	i-Bu	O	H	NO ₂	Br
15	i-Bu	O	H	NO ₂	Cl
16	i-Bu	O	H	NO ₂	OTf
17	i-Bu	O	H	NO ₂	OMs
18	i-Bu	O	H	NO ₂	OTs
19	i-Bu	O	H	H	I
20	i-Bu	O	H	H	Br
21	i-Bu	O	H	H	Cl
22	i-Bu	O	H	H	OTf
23	i-Bu	O	H	H	OMs
24	i-Bu	O	H	H	OTs
25	i-Bu	O	H	Br	I
26	i-Bu	O	H	Br	Br
27	i-Bu	O	H	Br	Cl
28	i-Bu	O	H	Br	OTf
29	i-Bu	O	H	Br	OMs
30	i-Bu	O	H	Br	OTs
31	(CH ₃) ₃ CCH ₂	O	H	CN	I
32	(CH ₃) ₃ CCH ₂	O	H	CN	Br
33	(CH ₃) ₃ CCH ₂	O	H	CN	Cl
34	(CH ₃) ₃ CCH ₂	O	H	CN	OTf
35	(CH ₃) ₃ CCH ₂	O	H	CN	OMs
36	(CH ₃) ₃ CCH ₂	O	H	CN	OTs
37	(CH ₃) ₃ CCH ₂	O	H	CHO	I
38	(CH ₃) ₃ CCH ₂	O	H	CHO	Br
39	(CH ₃) ₃ CCH ₂	O	H	CHO	Cl
40	(CH ₃) ₃ CCH ₂	O	H	CHO	OTf
41	(CH ₃) ₃ CCH ₂	O	H	CHO	OMs
42	(CH ₃) ₃ CCH ₂	O	H	CHO	OTs

[0078]


[Table 2]



No.	A	X	R ¹	R ²	Y
43	(CH ₃) ₂ CCH ₂	O	H	NO ₂	I
44	(CH ₃) ₂ CCH ₂	O	H	NO ₂	Br
45	(CH ₃) ₂ CCH ₂	O	H	NO ₂	Cl
46	(CH ₃) ₂ CCH ₂	O	H	NO ₂	OTf
47	(CH ₃) ₂ CCH ₂	O	H	NO ₂	OMs
48	(CH ₃) ₂ CCH ₂	O	H	NO ₂	OTs
49	(CH ₃) ₂ CCH ₂	O	H	H	I
50	(CH ₃) ₂ CCH ₂	O	H	H	Br
51	(CH ₃) ₂ CCH ₂	O	H	H	Cl
52	(CH ₃) ₂ CCH ₂	O	H	H	OTf
53	(CH ₃) ₂ CCH ₂	O	H	H	OMs
54	(CH ₃) ₂ CCH ₂	O	H	H	OTs
55	(CH ₃) ₂ CCH ₂	O	H	Br	I
56	(CH ₃) ₂ CCH ₂	O	H	Br	Br
57	(CH ₃) ₂ CCH ₂	O	H	Br	Cl
58	(CH ₃) ₂ CCH ₂	O	H	Br	OTf
59	(CH ₃) ₂ CCH ₂	O	H	Br	OMs
60	(CH ₃) ₂ CCH ₂	O	H	Br	OTs
61	Benzyl	O	H	CN	I
62	Benzyl	O	H	CN	Br
63	Benzyl	O	H	CN	Cl
64	Benzyl	O	H	CN	OTf
65	Benzyl	O	H	CN	OMs
66	Benzyl	O	H	CN	OTs
67	Benzyl	O	H	CHO	I
68	Benzyl	O	H	CHO	Br
69	Benzyl	O	H	CHO	Cl
70	Benzyl	O	H	CHO	OTf
71	Benzyl	O	H	CHO	OMs
72	Benzyl	O	H	CHO	OTs
73	Benzyl	O	H	NO ₂	I
74	Benzyl	O	H	NO ₂	Br
75	Benzyl	O	H	NO ₂	Cl
76	Benzyl	O	H	NO ₂	OTf
77	Benzyl	O	H	NO ₂	OMs
78	Benzyl	O	H	NO ₂	OTs
79	Benzyl	O	H	H	I
80	Benzyl	O	H	H	Br
81	Benzyl	O	H	H	Cl
82	Benzyl	O	H	H	OTf
83	Benzyl	O	H	H	OMs
84	Benzyl	O	H	H	OTs

[0079]


[Table 3]



No.	A	X	R ¹	R ²	Y
85	Benzyl	O	H	Br	I
86	Benzyl	O	H	Br	Br
87	Benzyl	O	H	Br	Cl
88	Benzyl	O	H	Br	OTf
89	Benzyl	O	H	Br	OMs
90	Benzyl	O	H	Br	OTs
91	Ph	bond	H	CN	I
92	Ph	bond	H	CN	Br
93	Ph	bond	H	CN	Cl
94	Ph	bond	H	CN	OTf
95	Ph	bond	H	CN	OMs
96	Ph	bond	H	CN	OTs
97	Ph	bond	H	CHO	I
98	Ph	bond	H	CHO	Br
99	Ph	bond	H	CHO	Cl
100	Ph	bond	H	CHO	OTf
101	Ph	bond	H	CHO	OMs
102	Ph	bond	H	CHO	OTs
103	Ph	bond	H	NO ₂	I
104	Ph	bond	H	NO ₂	Br
105	Ph	bond	H	NO ₂	Cl
106	Ph	bond	H	NO ₂	OTf
107	Ph	bond	H	NO ₂	OMs
108	Ph	bond	H	NO ₂	OTs
109	Ph	bond	H	H	I
110	Ph	bond	H	H	Br
111	Ph	bond	H	H	Cl
112	Ph	bond	H	H	OTf
113	Ph	bond	H	H	OMs
114	Ph	bond	H	H	OTs
115	Ph	bond	H	Br	I
116	Ph	bond	H	Br	Br
117	Ph	bond	H	Br	Cl
118	Ph	bond	H	Br	OTf
119	Ph	bond	H	Br	OMs
120	Ph	bond	H	Br	OTs
121	4-MeO-Ph	bond	H	CN	I
122	4-MeO-Ph	bond	H	CN	Br
123	4-MeO-Ph	bond	H	CN	Cl
124	4-MeO-Ph	bond	H	CN	OTf
125	4-MeO-Ph	bond	H	CN	OMs
126	4-MeO-Ph	bond	H	CN	OTs

[0080]


[Table 4]



No.	A	X	R ¹	R ²	Y
127	4-MeO-Ph	bond	H	CHO	I
128	4-MeO-Ph	bond	H	CHO	Br
129	4-MeO-Ph	bond	H	CHO	Cl
130	4-MeO-Ph	bond	H	CHO	OTf
131	4-MeO-Ph	bond	H	CHO	OMs
132	4-MeO-Ph	bond	H	CHO	OTs
133	4-MeO-Ph	bond	H	NO ₂	I
134	4-MeO-Ph	bond	H	NO ₂	Br
135	4-MeO-Ph	bond	H	NO ₂	Cl
136	4-MeO-Ph	bond	H	NO ₂	OTf
137	4-MeO-Ph	bond	H	NO ₂	OMs
138	4-MeO-Ph	bond	H	NO ₂	OTs
139	4-MeO-Ph	bond	H	H	I
140	4-MeO-Ph	bond	H	H	Br
141	4-MeO-Ph	bond	H	H	Cl
142	4-MeO-Ph	bond	H	H	OTf
143	4-MeO-Ph	bond	H	H	OMs
144	4-MeO-Ph	bond	H	H	OTs
145	4-MeO-Ph	bond	H	Br	I
146	4-MeO-Ph	bond	H	Br	Br
147	4-MeO-Ph	bond	H	Br	Cl
148	4-MeO-Ph	bond	H	Br	OTf
149	4-MeO-Ph	bond	H	Br	OMs
150	4-MeO-Ph	bond	H	Br	OTs

[0081]


[[Table 5]



No.	R^{3a}	R^4
151	t-Bu	CH_3
152	n-Pr	CH_3
153	i-Pr	CH_3
154	Et	CH_3
155	Me	CH_3
156	H	CH_3
157	t-Bu	H
158	n-Pr	H
159	i-Pr	H
160	Et	H
161	Me	H
162	H	H

[0082]


[Table 6]



No.	R ^{3a}	R ⁴
163	t-Bu	5-CH ₃
164	n-Pr	5-CH ₃
165	i-Pr	5-CH ₃
166	Et	5-CH ₃
167	Me	5-CH ₃
168	H	5-CH ₃
169	t-Bu	H
170	n-Pr	H
171	i-Pr	H
172	Et	H
173	Me	H
174	H	H
175	t-Bu	3-F
176	n-Pr	3-F
177	i-Pr	3-F
178	Et	3-F
179	Me	3-F
180	H	3-F
181	t-Bu	5-Cl
182	n-Pr	5-Cl
183	i-Pr	5-Cl
184	Et	5-Cl
185	Me	5-Cl
186	H	5-Cl

[0083]

[Table 7]



No.	W	R ^{3a}	R ⁴
187	O	t-Bu	CH ₃
188	O	n-Pr	CH ₃
189	O	i-Pr	CH ₃
190	O	Et	CH ₃
191	O	Me	CH ₃
192	O	H	CH ₃
193	O	t-Bu	H
194	O	n-Pr	H
195	O	i-Pr	H
196	O	Et	H
197	O	Me	H
198	O	H	H
199	S	t-Bu	CH ₃
200	S	n-Pr	CH ₃
201	S	i-Pr	CH ₃
202	S	Et	CH ₃
203	S	Me	CH ₃
204	S	H	CH ₃
205	S	t-Bu	H
206	S	n-Pr	H
207	S	i-Pr	H
208	S	Et	H
209	S	Me	H
210	S	H	H

[0084]

The production process of the present invention is characterized in that a transition metal compound is used as the catalyst. In the production process of the present invention, the transition metals in the transition metal compounds used are those other than nickel and include copper, palladium, cobalt, iron, rhodium, ruthenium, iridium, and the like. Above

all, copper, palladium, or cobalt is preferable. Copper includes zero-valent Cu(0), mono-valent Cu(I), and di-valent Cu(II), wherein preferable is zero-valent Cu(0) or mono-valent Cu(I). Palladium is preferably zero-valent Pd(0), mono-valent Pd(I), or di-valent Pd(II). Cobalt includes zero-valent Co(0), mono-valent Co(I), di-valent Co(II), and tri-valent Co(III), wherein preferable is zero-valent Co(0), mono-valent Co or di-valent Co (II). Iron includes zero-valent Fe(0), di-valent Fe(II), and tri-valent Fe(III), wherein preferable is di-valent Fe(II) or trivalent Fe(III). Rhodium is preferably zero-valent Rh(0) or mono-valent Rh(I). Ruthenium is preferably zero-valent Ru(0) or di-valent Rh(II). Iridium includes zero-valent Ir(0), mono-valent Ir(I), di-valent Ir(II), tri-valent Ir(III), and tetra-valent Ir(IV), wherein preferable is trivalent Ir(III).

[0085]

The salt of Cu(I) includes copper(I) chloride, copper(I) bromide, copper(I) iodide, copper(I) acetate, copper tetrafluoroborate, copper thiophene-2-carboxylate, hydrates thereof, mixtures of these, and the like.

[0086]

The salt of Cu(II) includes copper(II) fluoride, copper(II) chloride, copper(II) bromide, copper(II) iodide, copper(II) acetate, copper(II) formate, copper(II) hydroxide, copper(II) nitrate, copper(II) carbonate, copper(II) acetylacetonate, copper(II) borate, copper(II) oxalate, copper(II) phthalate, copper(II) tartrate, copper (II) trifluoromethanesulfonate, copper(II) benzoate, hydrates thereof, mixtures of these, and the like.

[0087]

Above all, preferable is copper(I) iodide (CuI).

[0088]

The salt of Pd(I) includes dibromo-dipalladium(I), a hydrate thereof, and the like.

[0089]

The salt of Pd(II) includes palladium(II) acetate, palladium(II) propionate, palladium(II) butanoate, palladium(II) 2-methylpropanoate, palladium(II) 3-methylbutanoate, palladium(II) 2-methylbutanoate, palladium(II) 2-ethylbutanoate, palladium(II) pivalate, palladium(II) 3,3-dimethylbutanoate, palladium(II) 2,2,3,3-tetramethylbutanoate, palladium(II) trifluoroacetate, palladium(II) nitrate, palladium(II) chloride, palladium(II) bromide, palladium(II) iodide, palladium(II) acetylacetonate, palladium(II) perchlorate, palladium(II) citrate, palladium(II) oxalate, palladium(II) cyclohexanecarboxylate, palladium(II) benzoate, palladium(II) stearate, palladium(II) sulfamate, palladium(II) carbonate, palladium(II) thiocyanate, palladium(II) trifluoromethanesulfonate, palladium(II) bis(4-diethylaminodithiobenzyl)palladium(II), palladium(II) cyanide, palladium(II) fluoride, palladium(II) boride, palladium(II) borate, palladium(II) hypophosphite, palladium(II) ammonium sulfate, palladium(II) hydroxide, cyclopentadienyl palladium(II), hydrates thereof, mixtures of these, and the like. Above all, preferable is palladium(II) acetate ($\text{Pd}(\text{OAc})_2$), palladium(II) propionate ($\text{Pd}(\text{O}(\text{C}=\text{O})\text{CH}_2\text{CH}_3)_2$), palladium(II) 2-methylpropanoate ($\text{Pd}(\text{O}(\text{C}=\text{O})\text{CH}(\text{CH}_3)_2)_2$), palladium(II) pivalate ($\text{Pd}(\text{OPiv})_2$), palladium(II) chloride (PdCl_2), palladium(I) bromide (Pd_2Br_2), or palladium(II) hydroxide ($\text{Pd}(\text{OH})_2$); especially preferable is palladium(II) acetate ($\text{Pd}(\text{OAc})_2$), palladium(II) propionate ($\text{Pd}(\text{O}(\text{C}=\text{O})\text{CH}_2\text{CH}_3)_2$), palladium(II) 2-methylpropanoate ($\text{Pd}(\text{O}(\text{C}=\text{O})\text{CH}(\text{CH}_3)_2)_2$), or palladium pivalate ($\text{Pd}(\text{OPiv})_2$).

[0090]

The salt of cobalt(II) includes cobalt(II) acetate, cobalt(II) nitrate, cobalt(II) chloride, cobalt(II) bromide, cobalt(II) iodide, cobalt(II) acetylacetonate, cobalt(II) perchlorate, cobalt(II) citrate, cobalt(II) oxalate, cobalt(II) fumarate, cobalt(II) gluconate, cobalt(II)

benzoate, cobalt(II) lactate, cobalt(II) stearate, cobalt(II) sulfamate, cobalt(II) carbonate, cobalt(II) thiocyanate, cobalt(II) fluoride, cobalt(II) phosphate, cobalt(II) sulfate, cobalt(II) hydroxide, cobalt(II) sulfide, hydrates thereof, mixtures of these, and the like. Above all, preferable is cobalt(II) acetate ($\text{Co}(\text{OAc})_2$).

[0091]

The salt of cobalt(III) includes cobalt(III) fluoride, cobalt(III) chloride, cobalt(III) bromide, cobalt(III) iodide, cobalt(III) acetylacetonate, cobalt(III) sulfate, cobalt(III) nitrate, cobalt(III) phosphate, cobalt(III) perchlorate, cobalt(III) citrate, hydrates thereof, mixtures of these, and the like.

[0092]

The salt of iron(II) includes iron(II) fluoride, iron(II) chloride, iron(II) bromide, iron(II) iodide, iron(II) sulfate, iron(II) nitrate, iron(II) oxalate, iron(II) fumarate, iron(II) acetate, iron(II) lactate, iron(II) gluconate, iron(II) benzoate, iron(II) stearate, iron(II) acetylacetonate, iron(II) sulfide, hydrates thereof, mixtures of these, and the like.

[0093]

The salt of iron(III) includes iron(III) fluoride, iron(III) chloride, iron(III) bromide, iron(III) iodide, iron(III) sulfate, iron(III) phosphate, iron(III) perchlorate, hydrates thereof, mixtures of these, and the like.

[0094]

The salt of rhodium(I) includes rhodium(I) chloride, hydrates thereof, mixtures of these, and the like.

[0095]

The salt of ruthenium(II) includes ruthenium(II) chloride, hydrates thereof, mixtures of these, and the like.

[0096]

The salt of iridium(III) includes iridium(III) chloride, iridium(III) bromide, iridium(III) acetate, iridium(III) carbonyl, (acetylacetonato)iridium(III), potassium hexachloroiridate(III), potassium pentachloronitrosyliridate(III), iridium(III) 2,4-pentanedionate, (pentamethylcyclopentadienyl)iridium(III) dichloride dimer, dichloro(pentamethylcyclopentadienyl)iridium(III) dimer, (pentamethylcyclopentadienyl)iridium hydrochloride dimer, hydrates thereof, mixtures of these, and the like.

[0097]

These transition metal compounds may be used as a mixture.

[0098]


Among these transition metal compounds, the particularly preferable metal species is palladium.

[0099]

As these transition metal compounds, there may be used compounds to which ligands were coordinated in advance. As such transition metal compounds having the ligands coordinated, there may be mentioned, for example, the following transition metal compounds. However, the present invention is not limited to these.

[0100]

[Formula 11]



[0101]

In the production process of the present invention, a ligand which can coordinate to the transition metal may be present together with the transition metal compound. The presence of a ligand which can coordinate to the transition metal compound in the reaction facilitate the coupling of the phenyl ring of a phenyl derivative and the C-H bond on the heterocyclic derivative can be coupled with high selectivity, resulting in a higher yield of the compound represented by the formula (3). Such a ligand used in the production process of the present invention includes a carboxylic acid type, amide type, phosphine type, oxime type, sulfide type, sulfonic acid type, 1,3-diketone type, Schiff's base type, oxazoline type, diamine type,

hydrocarbon type, carbon monoxide, a carbene type, and the like. However, the present invention is not limited to these. The coordinating atom in the ligand includes a nitrogen atom, phosphorous atom, oxygen atom, sulfur atom, and the like. As the ligand, there are a monodentate ligand having a coordinating atom at one position and a polydentate ligand having coordinating atoms at two or more positions. As for the hydrocarbon type, carbon monoxide, and carbene type, the coordinating atom is the carbon atom. These ligands may be used as salts.

[0102]

The monodentate ligand includes a phosphine-type ligand represented by $PR^5R^6R^7$ (wherein R^5 , R^6 , and R^7 each independently represent a C_1 to C_8 alkyl group, C_1 to C_4 alkoxy group, C_3 to C_8 cycloalkyl group, phenyl group, biphenyl group, phenoxy group, and furyl group. The C_3 to C_8 cycloalkyl group may further be substituted with a C_1 to C_4 alkyl group. The phenyl group may further be substituted with a methyl group, sulfonic acid group, or a salt thereof. The biphenyl group may further be substituted each independently with a C_1 to C_4 alkyl group, C_1 to C_4 alkoxy group, and dimethylamino group), triethylamine, pyridine, and the like.

[0103]

The phosphine-type ligand represented by $PR^5R^6R^7$ includes, for example,


tert-butylidicyclohexylphosphine,	isobutylidicyclohexylphosphine,
(n-butyl)dicyclohexylphosphine,	isopropylidicyclohexylphosphine,
(n-propyl)dicyclohexylphosphine,	ethylidicyclohexylphosphine,
methylidicyclohexylphosphine,	cyclopropylidicyclohexylphosphine,
cyclobutylidicyclohexylphosphine,	tert-butylidicyclooctylphosphine,
tert-butylidicycloheptylphosphine,	tert-butylidicyclopentylphosphine,

tert-butylidicyclobutylphosphine, tert-butylidicyclopropylphosphine, triethylphosphine,
 tri(n-propyl)phosphine, tri(isopropyl)phosphine, tri(tert-butyl)phosphine,
 tri(n-butyl)phosphine, tri(n-octyl)phosphine, tri(cyclooctyl)phosphine,
 tri(cycloheptyl)phosphine, tri(cyclohexyl)phosphine, tri(cyclopentyl)phosphine,
 tri(cyclobutyl)phosphine, tri(cyclopropyl)phosphine, di(tert-butyl)methylphosphine,
 di(tert-butyl)ethylphosphine, di(tert-butyl)n-propylphosphine,
 di(tert-butyl)isopropylphosphine, di(tert-butyl)n-butylphosphine,
 di(tert-butyl)isobutylphosphine, di(tert-butyl)neopentylphosphine, triphenylphosphine,
 tri(ortho-tolyl)phosphine, tri(mesityl)phosphine, tri(phenoxy)phosphine,
 tri(2-furyl)phosphine, trimethoxyphosphine, triethoxyphosphine, tri(n-propyloxy)phosphine,
 tri(isopropyloxy)phosphine, tri(n-butyloxy)phosphine, tri(isobutyloxy)phosphine,
 tri(tert-butyloxy)phosphine, di(tert-butyl)cyclohexylphosphine,
 di(isobutyl)cyclohexylphosphine, di(n-butyl)cyclohexylphosphine,
 di(isopropyl)cyclohexylphosphine, di(n-propyl)cyclohexylphosphine,
 diethylcyclohexylphosphine, dimethylcyclohexylphosphine,
 di(tert-butyl)cyclopentylphosphine, di(isobutyl)cyclopentylphosphine,
 di(n-butyl)cyclopentylphosphine, di(isopropyl)cyclopentylphosphine,
 di(n-propyl)cyclopentylphosphine, diethylcyclopentylphosphine,
 dimethylcyclopentylphosphine, di(tert-butyl)cyclooctylphosphine,
 di(tert-butyl)cycloheptylphosphine, di(tert-butyl)cyclopentylphosphine,
 di(tert-butyl)cyclobutylphosphine, di(tert-butyl)cyclopropylphosphine,
 dimethylphenylphosphine, diethylphenylphosphine, di(n-propyl)phenylphosphine,
 di(isopropyl)phenylphosphine, di(n-butyl)phenylphosphine, di(isobutyl)phenylphosphine,
 di(tert-butyl)phenylphosphine, di(cyclooctyl)phenylphosphine,

dicycloheptylphenylphosphine, dicyclohexylphenylphosphine,
 dicyclopentylphenylphosphine, dicyclobutylphenylphosphine,
 dicyclopropylphenylphosphine, dicyclohexyl(para-tolyl)phosphine,
 dicyclohexyl(ortho-tolyl)phosphine, dicyclohexyl(para-tolyl)phosphine,
 dicyclohexyl(2,4,6-trimethylphenyl)phosphine, methyldiphenylphosphine,
 ethyldiphenylphosphine, (n-propyl)diphenylphosphine, isopropyldiphenylphosphine,
 (n-butyl)diphenylphosphine, isobutyldiphenylphosphine, (tert-butyl)diphenylphosphine,
 cyclooctyldiphenylphosphine, cycloheptyldiphenylphosphine,
 cyclohexyldiphenylphosphine, cyclopentylphenylphosphine,
 cyclobutyldiphenylphosphine, cyclopropyldiphenylphosphine,
 bis(para-sulfonatophenyl)phenylphosphine potassium, cBRIDP, BippyPhos, TrippyPhos,
 XPhos (2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl), t-Bu-XPhos,
 JohnPhos, Cy-JohnPhos, MePhos, t-Bu-MePhos, DavePhos, t-Bu-DavePhos, SPhos,
 RuPhos (2-dicyclohexylphosphino-2',6'-diisopropoxy-1,1'-biphenyl), cataCXium A,
 cataCXium ABn, cataCXium PtB, cataCXium PCy, cataCXium POMetB, cataCXium
 POMeCy, cataCXium PIntB, cataCXium PInCy, cataCXium PICy, Q-Phos, JOSIPHOS,
 and the like; and mixtures of these.

[0104]

(Formula 12)



[0105]














The bidendate ligand includes 2,2'-bipyridyl, 4,4'-(tert-butyl)bipyridyl, phenanthroline,
 2,2'-bipyrimidyl, 1,4-diazabicyclo[2.2.2]octane, 2-(dimethylamino)ethanol,
 tetramethylethylenediamine, N,N-dimethylethylenediamine,
 N,N'-dimethylethylenediamine, 2-aminomethylpyridine,
 (NE)-N-(pyridin-2-ylmethylidene)aniline, 1,1'-bis(diphenylphosphino)ferrocene,
 1,1'-bis(tert-butyl)ferrocene, diphenylphosphinomethane,

1,2-bis(diphenylphosphino)ethane, 1,3-bis(diphenylphosphino)propane,
 1,5-bis(diphenylphosphino)pentane, 1,2-bis(dipentafluorophenylphosphino)ethane,
 1,2-bis(dicyclohexylphosphino)ethane, 1,3-(dicyclohexylphosphino)propane,
 1,2-bis(di-tert-butylphosphino)ethane, 1,3-bis(di-tert-butylphosphino)propane,
 1,2-bis(diphenylphosphino)benzene, 1,5-cyclooctadiene, BINAP, BIPHEMP, PROPHOS,
 DIOP, DEGUPHOS, DIPAMP, DuPHOS, NORPHOS, PNNP, SKEWPHOS, BPPFA,
 SEGPHOS, CHIRAPHOS, DPEphos, Xantphos, and the like; and mixtures of these.

[0106]

[Formula 13]

DPEphos



[0107]

BINAP includes derivatives thereof and specific examples include 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl,

2,2'-bis(di-para-tolylphosphino)-1,1'-binaphthyl,
 2,2'-bis(di-para-tert-butylphenylphosphino)-1,1'-binaphthyl,
 2,2'-bis(di-meta-tolylphosphino)-1,1'-binaphthyl,
 2,2'-bis(di-3,5-dimethylphenylphosphino)-1,1'-binaphthyl,
 2,2'-bis(di-para-methoxyphenylphosphino)-1,1'-binaphthyl,
 2,2'-bis(dicyclopentylphosphino)-1,1'-binaphthyl,
 2,2'-bis(dicyclohexylphosphino)-1,1'-binaphthyl,
 2-di(β -naphthyl)phosphino-2'-diphenylphosphino-1,1'-binaphthyl,
 2-diphenylphosphino-2'-di(para-trifluoromethylphenyl)phosphino-1,1'-binaphthyl, and the like.

[0108]

BIPHEMP includes derivatives thereof and specific examples include
 2,2'-dimethyl-6,6'-bis(diphenylphosphino)-1,1'-biphenyl,
 2,2'-dimethyl-6,6'-bis(dicyclohexylphosphino)-1,1'-biphenyl,
 2,2'-dimethyl-4,4'-bis(dimethylamino)-6,6'-bis(diphenylphosphino)-1,1'-biphenyl,
 2,2',4,4'-tetramethyl-6,6'-bis(diphenylphosphino)-1,1'-biphenyl,
 2,2'-dimethoxy-6,6'-bis(diphenylphosphino)-1,1'-biphenyl,
 2,2',3,3'-tetramethoxy-6,6'-bis(diphenylphosphino)-1,1'-biphenyl,
 2,2',4,4'-tetramethyl-3,3'-dimethoxy-6,6'-bis(diphenylphosphino)-1,1'-biphenyl,
 2,2'-dimethyl-6,6'-bis(di-para-tolylphosphino)-1,1'-biphenyl,
 2,2'-dimethyl-6,6'-bis(di-para-tert-butylphenylphosphino)-1,1'-biphenyl,
 2,2',4,4'-tetramethyl-3,3'-dimethoxy-6,6'-bis(di-para-methoxyphenylphosphino)-1,1'-biphenyl, and the like.

[0109]

The ligand used in the reaction of the present invention may be used as a salt. Such a salt includes, for example, a hydrochloric acid salt, hydrobromic acid salt, tetrafluoroboric acid salt, and the like.

[0110]

When a palladium catalyst is used, the ligand is preferably a phosphine-type ligand. Above all, the phosphine-type ligand represented by $PR^5R^6R^7$ is preferable. Specifically, preferable is tri(tert-butyl)phosphine, tri(cyclohexyl)phosphine, tert-butylidicyclohexylphosphine, di(tert-butyl)cyclohexylphosphine, di(tert-butyl)methylphosphine, 2-dicyclohexylphosphino-2',6'-diisopropoxy-1,1'-biphenyl, 2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl, or salts thereof; more preferable is tri(tert-butyl)phosphine, di(tert-butyl)cyclohexylphosphine, or a salt thereof; and particularly preferable is di(tert-butyl)cyclohexylphosphine or a salt thereof.

[0111]

When the ligand is coordinated in advance, a preferred ligand may be used by coordinating the same to palladium.

[0112]

The ligand may be used as a mixture. Furthermore, the ligand may be used by coordinating the same to a transition metal compound in advance. In addition, the ligand used in the reaction of the present invention may not be used depending on the circumstances.

[0113]

In the production process of the present invention, a base may be used together with the transition metal compound. By using the base together, the yield of the compound represented by the formula (3) can be improved. Such a base used in the production process

of the present invention is not particularly limited but preferable above all is lithium hydride, sodium hydride, potassium hydride, sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, cesium carbonate, potassium fluoride, cesium fluoride, tripotassium phosphate, sodium acetate, potassium acetate, and the like; a C₁ to C₆ alkoxide metal salt (lithium salt, sodium salt, potassium salt, and magnesium salt); C₁ to C₆ alkyl anion metal salt (lithium salt, sodium salt, potassium salt, and magnesium salt); tetra(C₁ to C₄ alkyl)ammonium salt (fluoride salt, chloride salt, and bromide salt), diisopropylethylamine, tributylamine, N-methylmorpholine, diazabicycloundecene, diazabicyclooctane, imidazole; or the like.

[0114]

In the “C₁ to C₆ alkoxide metal salt (lithium salt, sodium salt, potassium salt, and magnesium salt)” used as a base in the production process of the present invention, the “C₁ to C₆ alkoxide” includes methoxide, ethoxide, n-propoxide, isopropoxide, n-butoxide, isobutoxide, tert-butoxide, n-pentoxide, isopentoxide, neopentoxide, 1-methylpropoxide, n-hexoxide, isohexoxide, 1,1-dimethylbutoxide, 2,2-dimethylbutoxide, 3,3-dimethylbutoxide, and the like. Furthermore, there may be used mixtures of these.

[0115]

In the “C₁ to C₆ alkyl anion metal salt (lithium salt, sodium salt, potassium salt, and magnesium salt)” used as a base in the reaction of the present invention, the “C₁ to C₆ alkyl anion” includes a methyl anion, ethyl anion, n-propyl anion, isopropyl anion, n-butyl anion, isobutyl anion, tert-butyl anion, n-pentyl anion, isopentyl anion, neopentyl anion, 1-methylpropyl anion, n-hexyl anion, isohexyl anion, 1,1-dimethylbutyl anion, 2,2-dimethylbutyl anion, 3,3-dimethylbutyl anion, and the like. Furthermore, there may be used mixtures of these.

[0116]

When a palladium catalyst is used, the base is preferably potassium carbonate, potassium bicarbonate, cesium carbonate, or tetra(n-butyl)ammonium fluoride; particularly preferably potassium carbonate or cesium carbonate.

[0117]

When a copper catalyst is used, the preferable base is potassium phosphate. When a cobalt catalyst is used, the preferable base is cesium fluoride.

[0118]

Depending on the circumstances, the base used in the present reaction may not be used.

[0119]

In the production process of the present invention, a reducing agent which reduces the transition metal may be used together with the transition metal compound. For example, there may be mentioned zinc and the like.

[0120]

In the production process of the present invention, there may be added a silver salt. By addition of the silver salt may further improve the yield of the compound represented by the formula (3). Such a silver salt includes, for example, silver carbonate and the like.

[0121]

In the production process of the present invention, there may be added a C₁ to C₁₂ carboxylic acid or salt thereof. The addition of the C₁ to C₁₂ carboxylic acid or salt thereof may further improve the yield of the compound represented by the formula (3) and/or the rate of the reaction. These C₁ to C₁₂ carboxylic acids and salts thereof may be used as a mixture. The C₁ to C₁₂ carboxylic acid has 1 to 12 carbon atoms including the carbon atom of the carboxyl group and may contain a halogen atom, oxo group, and ether bond. The

examples include formic acid, acetic acid, propionic acid, butanoic acid, 2-methylpropanoic acid, pentanoic acid, 3-methylbutanoic acid, 2-methylbutanoic acid, pivalic acid, 3,3-dimethylbutanoic acid, 2-methylpentanoic acid, 2-methylhexanoic acid, 2-methylheptanoic acid, pentanecarboxylic acid, hexanoic acid, 4-methylpentanoic acid, 3,3-dimethylbutanoic acid, 2-ethylbutanoic acid, 2-methylpentanoic acid, 3-methylpentanoic acid, 2,2-dimethylbutanoic acid, 2,3-dimethylbutanoic acid, heptanoic acid, 2-methylhexanoic acid, 3-methylhexanoic acid, 4-methylhexanoic acid, 5-methylhexanoic acid, 2,2-dimethylpentanoic acid, 2,3,3-trimethylbutanoic acid, octanoic acid, 2-propylpentanoic acid, 2-ethylhexanoic acid, 2-methylheptanoic acid, 3-methylheptanoic acid, 4-methylheptanoic acid, 6-methylheptanoic acid, 2,2-dimethylheptanoic acid, 3-methylheptanoic acid, 2,2-diethylbutanoic acid, 2,2,4-trimethylpentanoic acid, 2-methyloctanoic acid, 2-methylundecanoic acid, 2-methylnonanoic acid, α -methylcinnamyl acid, cyclopropylacetic acid, 3-cyclopropylpropionic acid, cyclobutylacetic acid, cyclopentylacetic acid, cyclohexylacetic acid, cyclopentylpropionic acid, (2-methylcyclopentyl)acetic acid, cyclopentanecarboxylic acid, 3-oxocyclopentanecarboxylic acid, cyclopropanecarboxylic acid, cyclobutanecarboxylic acid, cyclohexanecarboxylic acid, cycloheptanecarboxylic acid, 1-methylcyclopropanecarboxylic acid, 2-methylcyclopropanecarboxylic acid, 2,2-dimethylcyclopropanecarboxylic acid, 2,2,3,3-tetramethylcyclopropanecarboxylic acid, 2-octylcyclopropanecarboxylic acid, 1-(4-methylphenyl)-1-cyclopropanecarboxylic acid, 2-para-tolylcyclopropanecarboxylic acid, 1-(2-fluorophenyl)cyclopropanecarboxylic acid, 1-(3-fluorophenyl)cyclopropanecarboxylic acid, 1-(4-fluorophenyl)cyclopropanecarboxylic acid, 1-(4-chlorophenyl)cyclopropanecarboxylic acid, 1-(3-chlorophenyl)cyclopropanecarboxylic acid, 2-(4-chlorophenyl)cyclopropanecarboxylic

acid,	1-(2,4-dichlorophenyl)cyclopropanecarboxylic	acid,
1-(3,4-dichlorophenyl)cyclopropanecarboxylic		acid,
2-fluoro-2-phenylcyclopropanecarboxylic		acid,
1-(4-methoxyphenyl)cyclopropanecarboxylic		acid,
2-(4-(tert-butyl)phenyl)cyclopropanecarboxylic	acid,	2,2-difluorocyclopropanecarboxylic
acid,	2,2-dichlorocyclopropanecarboxylic	acid,
2-chloro-2-fluorocyclopropanecarboxylic		
acid,	1-trifluoromethylcyclopropanecarboxylic	acid,
2,2-dichloro-1-methylcyclopropanecarboxylic	acid,	cyclopropane-1,1-dicarboxylic
acid,		
2,2'-oxydiacetic	acid,	1,2-dimethylcyclopropanedicarboxylic
		acid,
4-methylcyclobutanecarboxylic	acid,	4-ethylcyclopropanecarboxylic
		acid,
3-methoxycyclobutanecarboxylic	acid,	3-chlorocyclobutanecarboxylic
		acid,
4-chlorobutanecarboxylic	acid,	3-oxo-cyclobutanecarboxylic
		acid,
3,3-dimethylcyclobutanecarboxylic	acid,	1-methylcyclopentanecarboxylic
		acid,
3-cyclopentenecarboxylic	acid,	1-methylcyclopentanecarboxylic
		acid,
1-methylcyclohexanecarboxylic	acid,	4-methylcyclohexanecarboxylic
		acid,
2-methylcyclohexanecarboxylic	acid,	3-methylcyclohexanecarboxylic
		acid,
cyclooctanecarboxylic	acid,	spiro[2.2]pentane-1-carboxylic
		acid,
spiro[2.3]hexane-1-carboxylic	acid,	bicyclo[4.1.0]heptane-7-carboxylic
		acid,
tricyclo[3.2.1.0*2, 4*]octane-3-carboxylic	acid,	bicyclo[6.1.0]nonane-9-carboxylic
		acid,
bicyclo[2.2.1]heptane-1-carboxylic	acid,	bicyclo[2.2.1]heptane-2-carboxylic
		acid,
7,7-dimethyltricyclo[2.2.1.0(2,6)]heptane-1-carboxylic	acid,	5-norbornene-2-carboxylic
		acid,
norbornane-2-carboxylic	acid,	1-adamantanecarboxylic
		acid,
3-methyladamantane-1-carboxylic	acid,	3-fluoroadamantane-1-carboxylic
		acid,
3,5-dimethyladamantane-1-carboxylic	acid,	3-ethyladamantane-1-carboxylic
		acid,

3-chloroadamantane-1-carboxylic acid, 3,5,7-trimethyladamantane-1-carboxylic acid, 3-bromoadamantane-1-carboxylic acid, 5-bromo-3-methyladamantane-1-carboxylic acid, 5-bromo-3-ethyladamantane-1-carboxylic acid, tetrahydrofuran-2-carboxylic acid, tetrahydrofuran-3-carboxylic acid, tetrahydropyran-4-carboxylic acid, tetrahydropyran-3-carboxylic acid, methoxyacetic acid, trichloroacetic acid, dichloroacetic acid, chloroacetic acid, fluoroacetic acid, 2-fluoro-2-methylpropanoic acid, difluoroacetic acid, 2-chloropropanoic acid, 3-fluoropropionic acid, 2-fluoropropionic acid, 2-chloropropionic acid, 3-chloropropionic acid, 2-chlorobutanoic acid, 3-chlorobutanoic acid, 4-chlorobutanoic acid, 2-chloro-2-methylpropanoic acid, 3-chloro-2,2-dimethylpropanoic acid, 5-chloropentanoic acid, 2-chloro-3-methylbutanoic acid, dichloroacetic acid, 1-fluoro-1-chloroacetic acid, 2,2-difluoropropionic acid, 2,2-difluorobutanoic acid, 2,2-dichloropropionic acid, 2,3-dichloropropionic acid, chlorodifluoroacetic acid, trifluoroacetic acid, 3,3,3-trifluoropropionic acid, 2-methyl-4,4,4-trifluorobutanoic acid, 4,4,4-trifluorobutanoic acid, 2,2,3,3-tetrafluoropropionic acid, 2,3,3,3-tetrafluoropropionic acid, and the like. However, the present invention is not limited to these.

[0122]

As the C₁ to C₁₂ carboxylic acid, preferable is a carboxylic acid wherein the carbon atom at the α -position of the carboxyl group is not a carbon atom on an aromatic ring, more preferable being a carboxylic acid which may contain a halogen atom or ether bond. Examples include acetic acid, propionic acid, 2-methylpropanoic acid, 2-ethylbutanoic acid, pivalic acid, cyclopropanoic acid, 2,2,3,3-tetramethylcyclopropanoic acid, cyclopentanoic acid, 1-adamantanecarboxylic acid, 2-chloro-2-methylpropanoic acid, tetrahydrofuran-2-carboxylic acid, 2,2'-oxydiacetic acid, cyclopropane-1,1-dicarboxylic

acid, and the like.

[0123]

Above all, preferable is a carboxylic acid having one carboxyl group. Examples include acetic acid, propionic acid, 2-methylpropanoic acid, 2-ethylbutanoic acid, pivalic acid, cyclopropanoic acid, 2,2,3,3-tetramethylcyclopropanoic acid, cyclopentanoic acid, 1-adamantanecarboxylic acid, 2-chloro-2-methylpropanoic acid, tetrahydrofuran-2-carboxylic acid, and the like.

[0124]

Further, more preferable is a carboxylic acid wherein the number of hydrogen atoms bonded to the carbon atom at the α -position of the carboxyl group is 0 or 1. Examples include 2-methylpropanoic acid, 2-ethylbutanoic acid, pivalic acid, cyclopropanoic acid, 2,2,3,3-tetramethylcyclopropanoic acid, cyclopentanoic acid, 1-adamantanecarboxylic acid, 2-chloro-2-methylpropanoic acid, tetrahydrofuran-2-carboxylic acid, and the like.

[0125]

Particularly, more preferable is a carboxylic acid consisting of only carbon atoms and hydrogen atoms except the carboxyl group. Examples include 2-methylpropanoic acid, 2-ethylbutanoic acid, pivalic acid, cyclopropanoic acid, 2,2,3,3-tetramethylcyclopropanoic acid, cyclopentanoic acid, 1-adamantanecarboxylic acid, and the like.

[0126]

More particularly, preferable is a carboxylic acid having 1 to 8 carbon atoms, more preferable being a carboxylic acid having 1 to 6 carbon atoms. Examples include 2-methylpropanoic acid, 2-ethylbutanoic acid, pivalic acid, cyclopropanoic acid, cyclopentanoic acid, and the like. Even more particularly, preferable is 2-methylpropanoic acid or pivalic acid.

[0127]

The C₁ to C₁₂ carboxylic acid used in the reaction of the present invention may be used as a salt. Such a salt includes, for example, alkali metal salts such as a sodium salt, potassium salt, lithium salt, and the like; alkaline earth metal salts such as a calcium salt, magnesium salt, and the like; metal salts such as an aluminum salt, iron salt, and the like; inorganic salts such as an ammonium salt and the like; organic salts such as amine salts including a t-octylamine salt, dibenzylamine salt, morpholine salt, glucosamine salt, phenylglycyl alkyl ester salt, ethylenediamine salt, N-methylglucamine salt, guanidine salt, diethylamine salt, triethylamine salt, dicyclohexylamine salt, N,N'-dibenzylethylenediamine salt, chloroprocaine salt, procaine salt, diethanolamine salt, N-benzylphenylamine salt, piperazine salt, tetramethylammonium salt, tris(hydroxymethyl)aminomethane salt, and the like.

[0128]

The C₁ to C₁₂ carboxylic acid and salt thereof used in the reaction of the present invention can usually be used as an additive. However, if they are commercially available as or easy to prepare as carboxylic acid salts of transition metal compounds (for example, palladium(II) propionate), the transition metal carboxylates may be used.

[0129]

The production process of the present invention may be carried out in a wide range of temperature. The range is generally 0°C to 200°C, preferably 0°C to 150°C. Furthermore, the reaction is preferably carried out under ordinary pressure but may also be carried out under increased pressure or reduced pressure. The reaction time is 0.1 to 72 hours, preferably 0.1 to 48 hours. The reaction can be carried out in air but it is desirable that the reaction is carried out under a gas atmosphere having no ill effect on the reaction, such as

argon gas, nitrogen gas, and the like. Furthermore, in the present reaction, microwave may be irradiated.

[0130]

The solvent used in the production process of the present invention includes aliphatic hydrocarbons (hexane, cyclohexane, heptane, and the like), aliphatic halogenated hydrocarbons (dichloromethane, chloroform, carbon tetrachloride, dichloroethane, and the like), aromatic hydrocarbons (benzene, toluene, mesitylene, chlorobenzene, and the like), ethers (diethyl ether, dibutyl ether, dimethoxyethane (DME), cyclopentyl methyl ether (CPME), tert-butyl methyl ether, tetrahydrofuran, dioxane, and the like), esters (ethyl acetate, ethyl propionate, and the like), acid amides (dimethylformamide (DMF), dimethylacetamide (DMA), N-methylpyrrolidone (NMP), and the like), nitriles (acetonitrile, propionitrile, and the like), dimethyl sulfoxide (DMSO), mixed solvents thereof, and the like.

[0131]

In the production process of the present invention, the amount used of the compound of the formula (2) may be in a range from 1 mol% to 1000 mol% relative to the compound of the formula (1). The range is preferably 50 mol% to 200 mol%, more preferably 80 mol% to 120 mol%.

[0132]

In the production process of the present invention, the amount used of the transition metal compound and ligand may be in a range of 100 mol% or less relative to the compound of the formula (1) or the compound of the formula (2). Preferably, the amount is in a range of 20 mol% or less. The ligand may not be used depending on the circumstances.

[0133]

In the production process of the present invention, the amount used of the base may be in a range of 1000 mol% or less relative to the compound of the formula (1) or the compound of the formula (2). Preferably, the amount is in a range of 500 mol% or less.

[0134]

In the production process of the present invention, the amount used of the solvent may be 1000 times or less the weight of the compound of the formula (1) or the compound of the formula (2). The amount is preferably 100 times or less, more preferably 20 times or less.

[0135]

The addition order of the compound of the formula (1), compound of the formula (2), transition metal compound, ligand, base, C₁ to C₁₂ carboxylic acid, and solvent, which are used in the production process of the present invention, is optional. The best order may be selected depending on the combination of the reagents used.

[0136]

In the production process of the present invention, the amount used of the C₁ to C₁₂ carboxylic acid may be 50000 mol% or less relative to the transition metal compound used. The amount is preferably 5000 mol% or less, more preferably 1000 mol% or less, particularly preferably 500 mol% or less.

[0137]

In the production process of the present invention, the amount used of the silver salt is 500 mol% or less relative to the compound of the formula (1). Preferably, the amount is 200 mol% or less.

[0138]

The “mol%” represents a concentration of a certain material, obtained by dividing the

number of moles of the certain material by 100 mol of a related material.


[0139]

The compound represented by the formula (1), which is used in the production process of the present invention, can be produced by the following methods.

Synthetic method (1):

[0140]

[Formula 14]



[0141]

In the reaction equation, X represents an oxygen atom; R^1 , R^2 , A, and Y are the same as defined in the formula (1); L^1 represents a leaving group and includes a halogen atom, methanesulfonyloxy group, trifluoromethanesulfonyloxy group, para-toluenesulfonyloxy group, and the like.

Specifically, the compound represented by the formula (1) can be produced by reacting the compound (a) with the compound (b) in the presence of a suitable base in a suitable solvent under a suitable temperature condition.

[0142]

The solvent used is not particularly limited and includes, for example, aliphatic hydrocarbons (hexane, cyclohexane, heptane, and the like), aliphatic halogenated

hydrocarbons (dichloromethane, chloroform, carbon tetrachloride, dichloroethane, and the like), aromatic hydrocarbons (benzene, toluene, xylene, chlorobenzene, mesitylene, and the like), ethers (diethyl ether, dibutyl ether, dimethoxyethane (DME), cyclopentyl methyl ether (CPME), tetrahydrofuran, dioxane, and the like), esters (ethyl acetate, ethyl propionate, and the like), acid amides (dimethylformamide (DMF), dimethylacetamide (DMA), N-methylpyrrolidone (NMP), and the like), nitriles (acetonitrile, propionitrile, and the like), dimethyl sulfoxide (DMSO), water, mixed solvents thereof, and the like.

[0143]

The base used includes, for example, lithium hydride, sodium hydride, potassium hydride, sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, cesium carbonate, potassium fluoride, cesium fluoride, tripotassium phosphate, sodium acetate, potassium acetate, and the like; a C₁ to C₆ alkoxide metal salt (lithium salt, sodium salt, and potassium salt); C₁ to C₆ alkyl anion metal salt (lithium salt, sodium salt, and potassium salt); diisopropylethylamine, tributylamine, N-methylmorpholine, diazabicycloundecene, diazabicyclooctane, imidazole; and the like.

[0144]


For example, the synthesis can be carried out by referring to the Reference Example of the present invention or "Bioorg. Med. Chem. Lett., 2004:14, pp. 2547-2550," etc.

[0145]

Synthetic method (2):

[0146]

[Formula 15]



[0147]

In the reaction equation, X represents an oxygen atom; R^1 , R^2 , A, and Y are the same as defined in the formula (1). This reaction can be carried out by using the Mitsunobu reaction. For example, in the presence of diethyl azodicarboxylate (DEAD), diisopropyl azodicarboxylate (DIAD), 1,1'-(azodicarbonyl)dipiperidine (ADDP), 1,1'-azobis(N,N-dimethylformamide) (TMAD), or the like and in the presence of triphenylphosphine, tributylphosphine, or the like, the reaction can be carried out in a solvent such as tetrahydrofuran, diethyl ether, 1,2-dimethoxyethane, dichloromethane, toluene, or the like at a temperature range of from 0°C to 150°C.

[0148]

The compound represented by the formula (1) can be produced by a reaction using the Mitsunobu reaction and its related reactions described in Bull. Chem. Soc. Jpn., 1967:40, p. 2380; Synthesis, 1981, p. 1; and Org. React., 1992:42, p. 335.

[0149]

Besides the above, the compound represented by the formula (1) can also be synthesized by using an existing general ether synthesis method. For example, the synthesis can be carried out by referring to general text books on synthetic organic chemistry such as Jikken Kagaku Koza 4th ed., No. 20, "Organic Synthesis II Alcohol/Amine," Ed. Chemical


Society of Japan (Maruzen Co., Ltd.), pp. 187-205, and the like.

[0150]

Among the compounds represented by the formula (2), those having a thiazole ring as B are commercially available in some cases but they can be synthesized by referring, for example, to the following scheme:

[0151]

[Formula 16]



[0152]

In the reaction scheme described above, R³ and R⁴ are the same as defined in the formula (2) of the present invention. X' represents a halogen atom.

[0153]

The 2-aminothiazole derivative according to the thiazole cyclization reaction of Step 1 can be synthesized by referring to Pharmaceutical Chemistry Journal, 2007:41, pp. 105-108; Pharmaceutical Chemistry Journal, 2001:35, pp. 96-98; WO 2005/075435; WO 2005/026137; and the like. The reaction of Step 2 can be carried out by reference to Journal of Heterocyclic Chemistry, 1985:22, pp. 1621-1630; Journal of the Chemical Society, Perkin Transactions 1: Organic and Bio-Organic Chemistry, 1982:1, pp. 159-164; and Bioorganic & Medicinal Chemistry Letters, 2008:18, pp. 6231-6235. In addition, the thiazole derivative represented by the formula (2) can be synthesized by referring to, for example, WO 2002/051849 and WO 2001/062250.

[0154]

Among the compounds represented by the formula (2), various compounds having a pyridine ring as B are commercially available. As well as being purchasable, synthetic

methods of these compounds are widely reported and, thus, they can be synthesized by using those techniques.

[0155]

Among the compounds represented by the formula (2), those having an isoxazole ring or isothiazole ring [in the formula (2), W is an oxygen atom or sulfur atom] are commercially available in some cases. However, they can be synthesized by referring, for example, to the methods described in Tetrahedron Letters, 1968, pp. 5209-5213; Synthesis, 1970, pp. 344-350; Angewandte Chemie, 1967:79, pp. 471-472; and Chemische Berichte, 1973:106, pp. 3291-3311.

[Examples]

[0156]

In the following, the present invention will be described specifically with reference to Examples, etc. However, it should be understood that the scope of the present invention is not limited in any sense by these Examples.

[0157]

In the present Examples, the analysis and purification were carried out by using the following instruments and the like:

[0158]

TLC: E. Merck silica gel 60 F₂₅₄ (0.25 mm)

Flash column chromatography: Biotage Flash, Si40

Preparative thin-layer chromatography (PTLC): Merck silica gel 60 F₂₅₄ (1 mm)

Liquid Chromatography/Mass Spectrometry (LC/MS):

Analytical System: SHIMAZU LCMS-2010A

Software: LCMS Solution

Experimental Conditions:

Column: Phenomenex Gemini 3 μm 4.6 mm $\times$ 30 mm

Flow Rate: 1.2 mL/min

Measurement Temperature: 40°C

A-Solvent: 5% MeCN/95% H₂O + 0.05% TFA

B-Solvent: 95% MeCN/5% H₂O + 0.05% TFA

MS-mode: ESI+

ESI Voltage: 4.5 KV

Source Temp: 130°C

Desolvation Temp: 320°C

[0159]

[Table 8]

	Time [min]	A [%]	B [%]	Flow [ml/min]
1	0.01	95	5	1.2
2	0.3	60	40	1.2
3	2.3	0	100	1.2
4	3.8	0	100	1.2
5	4.0	95	5	1.2
6	4.5	95	5	0

A dual column system was employed.

[0160]

Nuclear magnetic resonance (NMR): JEOL JNM-AL400 (¹H 400 MHz)

The ¹H-NMR shift values are shown in ppm with the shift value of tetramethylsilane (δ

0.0 ppm). A dual column system was employed. The following abbreviations:

s = singlet, d = doublet, dd = doublet of doublets, t = triplet, q = quartet, m = multiplet, br = broad signal.


In addition, in the ^1H -NMR spectra in Reference Examples and Examples, the proton signal of a carboxylic acid may not be able to be confirmed in some cases, depending on the measurement conditions such as the solvent and the like.

[Reference Example 1]

Synthesis of tert-butyl 4-methylthiazole-5-carboxylate

[0161]

[Formula 17]



[0162]

A mixture of 4-methyl-5-thiazolecarboxylic acid (1.36 g, 9.48 mmol) and thionyl chloride (28.7 mL) was stirred at 80°C for 1 hour. The reaction mixture was concentrated under reduced pressure by removing thionyl chloride and the crude product obtained was dried under reduced pressure. To a solution of this crude material in dichloromethane (5.68 mL), there were added tert-butanol (2.84 mL) and pyridine (16.9 mL) and the mixture was stirred at 60°C overnight. After the reaction was complete, the reaction mixture was concentrated under reduced pressure. To the crude material obtained were added a saturated

aqueous sodium carbonate solution and ethyl acetate. After separating ethyl acetate, extraction was performed by adding ethyl acetate again to the saturated aqueous sodium bicarbonate solution. The combined organic phases were washed with a saturated aqueous sodium chloride solution and dried over anhydrous magnesium sulfate. After removing magnesium sulfate by filtration, the solvent was concentrated under reduced pressure. The crude product obtained was purified by silica gel chromatography (hexane/ethyl acetate = 85/15) to obtain the title compound (964 mg). Yield, 51%.


$^1\text{H-NMR}$ (400MHz, CDCl_3): δ 8.72 (s, 1H), 2.74 (s, 3H), 1.58 (s, 9H).

[Reference Example 2]

Synthesis of 5-iodo-2-isobutoxybenzonitrile

[0163]

[Formula 18]



[0164]

A solution of 2-methyl-1-propanol (0.56 mL, 6.06 mmol) in *N,N*-dimethylformamide (10 mL) was cooled to 0°C and to this was added sodium hydride (242 mg, a 60% suspension in mineral oil, 6.06 mmol) in small portions. The turbid reaction mixture was stirred at 0°C for 5 minutes and the temperature was raised to 23°C. Thereafter, the mixture

was stirred at room temperature for 10 minutes and cooled again to 0°C. To the reaction mixture was added 2-fluoro-5-iodobenzonitrile (1.0 g, 4.04 mmol) and, after being warmed to room temperature, the reaction mixture was stirred for 1.5 hours. After the reaction was complete, water (20 mL) was added to the reaction mixture, which was extracted with ethyl acetate (3 × 30 mL). The organic phases were combined, washed with a saturated aqueous sodium chloride solution (3 × 30 mL), and thereafter dried over anhydrous magnesium sulfate. After removing magnesium sulfate by filtration, the solvent was concentrated under reduced pressure. The crude product obtained was purified by silica gel chromatography (hexane/ethyl acetate = 98/2) to obtain the title compound (950 mg). Yield, 78%.


¹H-NMR (400MHz, CDCl₃): δ 7.81 (d, J = 2.2Hz, 1H), 7.76 (dd, J = 8.8Hz, 2.2Hz, 1H), 6.72 (d, J = 8.8Hz, 1H), 3.80 (d, J = 6.3Hz, 2H), 2.21-2.11 (m, 1H), 1.06 (d, J = 6.8Hz, 6H).

[Reference Example 3]

Synthesis of 5-bromo-2-isobutoxybenzonitrile

[0165]

[Formula 19]



[0166]

A suspension of sodium hydride (1.64 g, a 60% suspension in mineral oil, 37.5 mmol) in N,N-dimethylformamide (50 mL) was cooled to 0°C and thereafter 2-methyl-1-propanol

(3.47 mL, 37.5 mmol) was added thereto in small portions. The reaction mixture was stirred at room temperature for 20 minutes. The mixture was cooled again to 0°C, 2-fluoro-5-bromobenzonitrile (5.00 g, 25.0 mmol) was added thereto in small portions, and thereafter the reaction mixture was stirred at room temperature for 15 hours. After the reaction was complete, water (100 mL) was added to the reaction mixture, which was then extracted with ethyl acetate (3 × 100 mL). The organic phases were combined, washed with a saturated aqueous sodium chloride solution (2 × 50 mL), and thereafter dried over anhydrous sodium sulfate. After removing sodium sulfate by filtration, the solvent was concentrated under reduced pressure. The crude product obtained was purified by silica gel chromatography (hexane/ethyl acetate = 9/1) to obtain the title compound (6.04 g). Yield, 95%.


¹H-NMR (400MHz, CDCl₃): δ 7.65 (d, J = 2.4Hz, 1H), 7.60 (d, J = 9.0Hz, 2.4Hz, 1H), 6.84 (d, J = 8.8Hz, 1H), 3.81 (d, J = 6.6Hz, 2H), 2.22-2.12 (m, 1H), 1.06 (d, J = 6.6Hz, 6H).

[Example 1]

Synthesis of tert-butyl 2-(3-cyano-4-isobutoxyphenyl)-4-methylthiazole-5-carboxylate

[0167]

[Formula 20]



[0168]

To a reaction vessel were added tert-butyl 4-methylthiazole-5-carboxylate (49.8 mg,

0.25 mmol) obtained in Reference Example 1, 5-iodo-2-isobutoxybenzonitrile (112.9 mg, 0.375 mmol) obtained in Reference Example 2, and anhydrous N,N-dimethylformamide (1.25 mL). After adding tert-butoxy lithium (40.0 mg, 0.5 mmol) and copper(I) iodide (9.5 mg, 0.05 mmol) to the mixture under a nitrogen atmosphere, the reaction mixture was heated to 140°C and stirred for 30 minutes. After the reaction was complete, the reaction mixture was cooled to room temperature. Water was added to the reaction mixture, which was then extracted with ethyl acetate. The combined organic phases were washed with a saturated aqueous sodium chloride solution and dried over magnesium sulfate. After removing magnesium sulfate by filtration, the organic solvent was concentrated under reduced pressure. The crude product obtained was purified by thin-layer silica gel chromatography (hexane/ethyl acetate = 4/1) to obtain the title compound (29.2 mg). Yield, 31%.


¹H-NMR (400MHz, CDCl₃): δ 8.16 (d, J = 2.4Hz, 1H), 8.08 (dd, J = 8.8Hz, 2.4Hz, 1H), 7.00 (d, J = 8.8Hz, 1H), 3.89 (d, J = 6.8Hz, 2H), 2.73 (s, 3H), 2.24-2.16 (m, 1H), 1.59 (s, 9H), 1.09 (d, J = 6.8Hz, 6H).

[Example 2]

Synthesis of tert-butyl 2-(3-cyano-4-isobutoxyphenyl)-4-methylthiazole-5-carboxylate

[0169]

[Formula 21]



[0170]

To a reaction vessel were added tert-butyl 4-methylthiazole-5-carboxylate (49.8 mg, 0.25 mmol) obtained in Reference Example 1, 5-iodo-2-isobutoxybenzonitrile (112.9 mg, 0.375 mmol) obtained in Reference Example 2, and water (0.5 mL). Under a nitrogen atmosphere, a complex of palladium (II) chloride with bidentate 1,1'-bis(diphenylphosphino)ferrocene [PdCl₂(dppf)] (20.7 mg, 0.025 mmol), triphenylphosphine (39.3 mg, 0.15 mmol), and silver carbonate (138.4 mg, 0.5 mmol) were added and, thereafter, the reaction mixture was heated to 60°C and stirred for 24 hours. After the reaction was complete, the reaction mixture was cooled to room temperature. Ethyl acetate (2.5 mL) was added to the reaction mixture, and insoluble material was separated by filtration and the filtrate was washed with ethyl acetate. The filtrate was extracted with ethyl acetate twice. The combined organic phases were washed with a saturated aqueous sodium chloride solution and dried over magnesium sulfate. After removing magnesium sulfate by filtration, the organic solvent was concentrated under reduced pressure. The crude product obtained was purified by thin-layer silica gel chromatography (hexane/ethyl acetate = 3/1) to obtain the title compound (87.6 mg). Yield, 94%.

¹H-NMR (400MHz, CDCl₃): δ 8.16 (d, J = 2.4Hz, 1H), 8.08 (dd, J = 8.8Hz, 2.4Hz, 1H), 7.00 (d, J = 8.8Hz, 1H), 3.89 (d, J = 6.8Hz, 2H), 2.73 (s, 3H), 2.24-2.16 (m, 1H), 1.59 (s,


9H), 1.09 (d, $J = 6.8\text{Hz}$, 6H).

[Example 3]

Synthesis of tert-butyl 2-(3-cyano-4-isobutoxyphenyl)-4-methylthiazole-5-carboxylate

[0171]

[Formula 22]



[0172]

To a test tube-type reaction vessel (50 mL) were added tert-butyl 4-methylthiazole-5-carboxylate (598 mg, 3.0 mmol) obtained in Reference Example 1, 5-bromo-2-isobutoxybenzonitrile (762 mg, 3.0 mmol) obtained in Reference Example 3, palladium acetate (67.4 mg, 0.30 mmol), tri(cyclohexyl)phosphine (168 mg, 0.60 mmol), cesium carbonate (1.95 g, 6.0 mmol), and toluene (11 mL). Thereafter, the reaction vessel was filled with nitrogen, heated to 120°C under airtight seal, and the reaction mixture was stirred for 19 hours. After the reaction was complete, ethyl acetate (30 mL) was added to the reaction mixture and insoluble matter was removed by filtration. To the filtrate was added 0.1 mol/L hydrochloric acid (20 mL) and the organic phase was extracted and separated. Further, the aqueous phase was extracted with ethyl acetate (20 mL). The combined organic phases were washed with water (30 mL) and a saturated aqueous sodium chloride solution (30 mL) and dried over sodium sulfate. After removing sodium sulfate by filtration, the organic solvent was concentrated under reduced pressure. The crude product

obtained was purified by silica gel chromatography (hexane/ethyl acetate = 7/1) to obtain the title compound (930 mg). Yield, 83%.


$^1\text{H-NMR}$ (400MHz, CDCl_3): δ 8.16 (d, $J = 2.44\text{Hz}$, 1H), 8.08 (dd, $J = 8.78\text{Hz}$, 2.20Hz, 1H), 7.00 (d, $J = 8.78\text{Hz}$, 1H), 3.90 (d, $J = 6.59\text{Hz}$, 2H), 2.73 (s, 3H), 2.25-2.16 (m, 1H), 1.59 (s, 9H), 1.09 (d, $J = 6.83\text{Hz}$, 6H).

[Example 4]

Synthesis of tert-butyl 2-(3-cyano-4-isobutoxyphenyl)-4-methylthiazole-5-carboxylate

[0173]

[Formula 23]



[0174]

To a test tube-type reaction vessel were added tert-butyl 4-methylthiazole-5-carboxylate (180 mg, 0.903 mmol) obtained in Reference Example 1, 5-bromo-2-isobutoxybenzonitrile (230 mg, 0.903 mmol) obtained in Reference Example 3, palladium hydroxide (31.7 mg, 0.045 mmol), tri(cyclohexyl)phosphine (12.7 mg, 0.045 mmol), potassium carbonate (250 mg, 1.81 mmol), copper iodide (172 mg, 0.903 mmol), and dimethyl sulfoxide (3.0 mL). Thereafter, the reaction vessel was filled with nitrogen, heated to 120°C under airtight seal, and the reaction mixture was stirred for 20 hours. After the reaction was complete, ethyl acetate (10 mL) and water (10 mL) were added to the reaction mixture, and the resultant

mixture was stirred at room temperature for 30 minutes. The solution was filtered through Celite and the Celite layer was washed with ethyl acetate (20 mL) and water (10 mL). The organic phase was separated from the filtrate and the aqueous phase was extracted further with ethyl acetate (20 mL). The combined organic phases were washed with a saturated aqueous sodium chloride solution (10 mL) and dried over sodium sulfate. After removing sodium sulfate by filtration, the organic solvent was concentrated under reduced pressure. The crude product obtained was purified by silica gel chromatography (hexane/ethyl acetate = 49/1 to 4/1) to obtain a crude product of the title compound (188 mg). After dissolving this again in ethanol (3 mL) by heating (80°C), the solution was cooled to 10°C and the precipitated solid was separated by filtration and washed with ethanol (2 mL). The solid was dried at room temperature under reduced pressure to obtain the title compound (132 mg). Yield, 39%.


$^1\text{H-NMR}$ (400MHz, CDCl_3): δ 8.17 (d, $J = 2.20\text{Hz}$, 1H), 8.08 (dd, $J = 8.90\text{Hz}$, 2.32Hz , 1H), 7.00 (d, $J = 9.02\text{Hz}$, 1H), 3.90 (d, $J = 6.59\text{Hz}$, 2H), 2.73 (s, 3H), 2.24-2.15 (m, 1H), 1.59 (s, 9H), 1.09 (d, $J = 6.83\text{Hz}$, 6H).

[Example 5]

Synthesis of tert-butyl 2-(3-cyano-4-isobutoxyphenyl)-4-methylthiazole-5-carboxylate

[0175]

[Formula 24]



[0176]

To a reaction vessel were added tert-butyl 4-methylthiazole-5-carboxylate (598 mg, 3.0 mmol) obtained in Reference Example 1, 5-bromo-2-isobutoxybenzonitrile (762 mg, 3.0 mmol) obtained in Reference Example 3, palladium acetate (67.4 mg, 0.30 mmol), tri(cyclohexyl)phosphine (168 mg, 0.60 mmol), potassium carbonate (829 mg, 6.0 mmol), toluene (10 mL), and pivalic acid (92 mg, 0.90 mmol). Thereafter, the reaction mixture was stirred at room temperature for 30 minutes under a nitrogen atmosphere and for further 9 hours with heating under reflux. After the reaction was complete, water (20 mL) and ethyl acetate (20 mL) were added to the reaction mixture and the organic phase was extracted and separated. Further, the aqueous phase was extracted with ethyl acetate (20 mL). The combined organic phases were dried over magnesium sulfate. After removing magnesium sulfate by filtration, the organic solvent was concentrated under reduced pressure. The crude product obtained was purified by silica gel chromatography (hexane/ethyl acetate = 100/0 to 0/100) to obtain the title compound (880 mg). Yield, 79%.


¹H-NMR (400MHz, CDCl₃): δ 8.16 (d, J = 2.20Hz, 1H), 8.08 (dd, J = 8.78Hz, 2.20Hz, 1H), 7.00 (d, J = 8.78Hz, 1H), 3.90 (d, J = 6.34Hz, 2H), 2.73 (s, 3H), 2.25-2.15 (m, 1H), 1.59 (s, 9H), 1.09 (d, J = 6.59Hz, 6H).

[Example 6]

Synthesis of ethyl 2-(3-cyano-4-isobutoxyphenyl)-4-methylthiazole-5-carboxylate

[0177]

[Formula 25]



[0178]

To a reaction vessel were added ethyl 4-methylthiazole-5-carboxylate (514 mg, 3.0 mmol), 5-bromo-2-isobutoxybenzonitrile (762 mg, 3.0 mmol) obtained in Reference Example 3, palladium acetate (67.4 mg, 0.30 mmol), tri(cyclohexyl)phosphine (168 mg, 0.60 mmol), potassium carbonate (829 mg, 6.0 mmol), toluene (10 mL), and pivalic acid (92 mg, 0.90 mmol). Thereafter, the reaction mixture was stirred at room temperature for 30 minutes under a nitrogen atmosphere and for further 10 hours with heating under reflux. After the reaction was complete, water (20 mL) and ethyl acetate (20 mL) were added to the reaction mixture and the organic phase was extracted and separated. Further, the aqueous phase was extracted with ethyl acetate (20 mL). The combined organic phases were dried over magnesium sulfate. After removing magnesium sulfate by filtration, the solvent was concentrated under reduced pressure. The crude product obtained was purified by silica gel chromatography (hexane/ethyl acetate = 100/0 to 0/100) to obtain the title compound 734 mg). Yield, 71%.


¹H-NMR (400MHz, CDCl₃): δ 8.18 (d, J = 2.44Hz, 1H), 8.09 (dd, J = 8.78Hz, 2.20Hz, 1H), 7.01 (d, J = 8.78Hz, 1H), 4.36 (q, J = 7.07Hz, 2H), 3.90 (d, J = 6.34Hz, 2H), 2.77 (s, 3H), 2.26-2.16 (m, 1H), 1.39 (t, J = 7.19Hz, 3H), 1.09 (d, J = 6.83Hz, 6H).

[Example 7]

Synthesis of tert-butyl 2-(3-cyano-4-isobutoxyphenyl)-4-methylthiazole-5-carboxylate

[0179]

[Formula 26]



[0180]

To a reaction vessel were added tert-butyl 4-methylthiazole-5-carboxylate (598 mg, 3.0 mmol) obtained in Reference Example 1, 5-bromo-2-isobutoxybenzonitrile (801 mg, 3.15 mmol) obtained in Reference Example 3, palladium acetate (67.4 mg, 0.30 mmol), di(tert-butyl)cyclohexylphosphine (137 mg, 0.60 mmol), potassium carbonate (829 mg, 6.0 mmol), and toluene (10 mL). Thereafter, the reaction mixture was stirred at room temperature for 30 minutes and for further 24 hours under a nitrogen atmosphere with heating under reflux. After the reaction was complete, water (15 mL) and ethyl acetate (20 mL) were added to the reaction mixture and the organic phase was extracted and separated. Further, the aqueous phase was extracted with ethyl acetate (20 mL). The combined organic phases were dried over magnesium sulfate. After removing magnesium sulfate by filtration, the organic solvent was concentrated under reduced pressure to obtain a crude product. After dissolving the crude product by adding toluene (1 mL) and heating to 70°C, heptane (9 mL) was added thereto at 70°C and the resultant mixture was allowed to cool to room temperature and further cooled to 0°C. The precipitated solid was separated by filtration, washed with heptane (20 mL) to obtain the title compound (611 mg, 1.64 mmol). Further,

the filtrate was concentrated under reduced pressure and the crude product obtained was purified by silica gel chromatography (hexane/ethyl acetate = 100/0 to 0/100) to obtain the title compound (405 mg, 1.09 mmol). Yield, 91%.


$^1\text{H-NMR}$ (400MHz, CDCl_3): δ 8.17(d, $J = 2.20\text{Hz}$, 1H), 8.09 (dd, $J = 8.90\text{Hz}$, 2.32Hz, 1H), 7.00 (d, $J = 8.78\text{Hz}$, 1H), 3.90 (d, $J = 6.34\text{Hz}$, 2H), 2.73 (s, 3H), 2.24-2.17 (m, 1H), 1.59 (s, 9H), 1.09 (d, $J = 6.83\text{Hz}$, 6H).

[Example 8]

Synthesis of ethyl 2-(3-cyano-4-isobutoxyphenyl)-4-methylthiazole-5-carboxylate

[0181]

[Formula 27]



[0182]

To a reaction vessel were added ethyl 4-methylthiazole-5-carboxylate (1.71 g, 10.0 mmol), 5-bromo-2-isobutoxybenzonitrile (2.54 mg, 10.0 mmol) obtained in Reference Example 3, palladium acetate (22.4 mg, 0.10 mmol), a tetrafluoroboric acid salt of tri(tert-butyl)phosphine (87.0 mg, 0.30 mmol), potassium carbonate (1.45 g, 10.5 mmol), and xylene (10 mL). Thereafter, the reaction mixture was stirred at room temperature for 30 minutes and for further 17 hours under a nitrogen atmosphere, with heating under reflux. After the reaction was complete, the reaction mixture was filtered while hot and the filtration bed was washed with toluene and dichloromethane. The filtrate was concentrated

under reduced pressure and the crude product obtained was purified to obtain the title compound (2.69 g). Yield, 78%.


$^1\text{H-NMR}$ (400MHz, CDCl_3): δ 8.18 (d, $J = 2.44\text{Hz}$, 1H), 8.09 (dd, $J = 8.78\text{Hz}$, 2.20Hz, 1H), 7.01 (d, $J = 8.78\text{Hz}$, 1H), 4.36 (q, $J = 7.07\text{Hz}$, 2H), 3.90 (d, $J = 6.34\text{Hz}$, 2H), 2.77 (s, 3H), 2.26-2.16 (m, 1H), 1.39 (t, $J = 7.19\text{Hz}$, 3H), 1.09 (d, $J = 6.83\text{Hz}$, 6H).

[Example 9]

Synthesis of ethyl 2-(3-cyano-4-isobutoxyphenyl)-4-methylthiazole-5-carboxylate

[0183]

[Formula 28]



[0184]

To a reaction vessel were added ethyl 4-methylthiazole-5-carboxylate (1.71 g, 10.0 mmol), 5-bromo-2-isobutoxybenzonitrile (2.69 g, 10.5 mmol) obtained in Reference Example 3, palladium acetate (22.4 mg, 0.10 mmol), di(tert-butyl)cyclohexylphosphine (68.5 mg, 0.30 mmol), potassium carbonate (1.45 g, 10.5 mmol), and xylene (10 mL). Thereafter, the reaction mixture was stirred at room temperature for 30 minutes and for further 24 hours under a nitrogen atmosphere with heating under reflux. After the reaction was complete, the reaction mixture was filtered while hot and the filtration bed was washed with toluene and dichloromethane. The filtrate was concentrated under reduced pressure and the crude product obtained was purified to obtain the title compound (2.83 g). Yield,

82%.


$^1\text{H-NMR}$ (400MHz, CDCl_3): δ 8.18 (d, $J = 2.44\text{Hz}$, 1H), 8.09 (dd, $J = 8.78\text{Hz}$, 2.20Hz, 1H), 7.01 (d, $J = 8.78\text{Hz}$, 1H), 4.36 (q, $J = 7.07\text{Hz}$, 2H), 3.90 (d, $J = 6.34\text{Hz}$, 2H), 2.77 (s, 3H), 2.26-2.16 (m, 1H), 1.39 (t, $J = 7.19\text{Hz}$, 3H), 1.09 (d, $J = 6.83\text{Hz}$, 6H).

[Example 10]

Synthesis of tert-butyl 2-(3-cyano-4-isobutoxyphenyl)-4-methylthiazole-5-carboxylate

[0185]

[Formula 29]



[0186]

To a test tube-type reaction vessel (10 mL) were added tert-butyl 4-methylthiazole-5-carboxylate (59.8 mg, 0.3 mmol) obtained in Reference Example 1, 5-bromo-2-isobutoxybenzonitrile (76.2 mg, 0.3 mmol) obtained in Reference Example 3, palladium acetate (6.7 mg, 0.030 mmol), a ligand, a base (0.60 mmol), and a solvent (1 mL).

Thereafter, the reaction vessel was filled with nitrogen, heated to 120°C under airtight seal, and the reaction mixture was stirred. After the reaction was complete, a portion of the reaction mixture was diluted with DMSO and the solution obtained was measured by HPLC. The total HPLC area% of the compounds A to D and TM was set as 100% and the calculated yield of the target material was computed from the HPLC area% of the target material. Computation of the calculated yield of the target material from the HPLC area% thereof was carried out by putting the corresponding values in the following calculation formula.

[0187]

$$\text{Yield of TM (\%)} = \frac{\text{total amount of TM (mol)}}{[\{\text{total amount of B (mol)} + \text{total amount of D (mol)} \times 2 + \text{total amount of TM (mol)}\} / 2 + \{\text{total amount of A (mol)} + \text{total amount of (C)} \times 2 + \text{total amount of TM (mol)}\} / 2]} \times 100$$
$$\text{Total amount of each compound (mol)} = \text{HPLC area value (mAU)} / \text{HPLC area value per mol of each compound (mAU/mol)}$$

High Performance Liquid Chromatography:

Analytical System: G1315A Hewlett Packard Series 1100

Software: ChemStation for LC 3D

Experimental Conditions:

Column: Intakt Cadenza CD-C18 4.6 × 100 mm

Flow: 1.0 mL/min

Wavelength: 254 nm

Temperature: 40°C

A-Solvent: 5% MeCN/95% H₂O + 0.05% TFA

B-Solvent: 95% MeCN/5% H₂O + 0.05%TFA

Gradient:

0-1 min 10% B-Solv.

1-7 min 10-50% B-Solv.

7-14 min 0-80% B-Solv.

14-16 min 80-100% B-Solv.

16-20 min 100% B-Solv.

20-22 min 100-10% B-Solv.

22-25 min 10% B-Solv.

The results in the present Example are shown in the following.

[0188]

Experiment No.	Reaction Conditions							HPLC area %					Calculated Yield % TM
	Solvent 0.3M	Base 2 eq.	Pd 0.1 eq.	Ligand 0.2 eq.	Additive 0.3 eq.	Temperature °C	Reaction Time hr	TM	A	B	C	D	
1	DME	K ₂ CO ₃	Pd(OAc) ₂	P(t-Bu) ₃ ·HBF ₄ (0.4 eq.)	-	120	24	63.0	4.1	21.6	N.D.	N.D.	98.8
2	NMP	K ₃ PO ₄	Pd(OAc) ₂	P(t-Bu) ₃ ·HBF ₄	-	120	24	20.8	8.2	62.1	3.7	N.D.	89.2
3	NMP	CS ₂ CO ₃	Pd(OAc) ₂	P(t-Bu) ₃ ·HBF ₄	-	120	24	38.2	7.3	38.4	2.7	N.D.	95.0
4	Toluene	CS ₂ CO ₃	Pd(OAc) ₂	PPh ₃	-	120	24	19.5	6.4	59.6	5.6	N.D.	65.5
5	Toluene	CS ₂ CO ₃	Pd(OAc) ₂	dppp	-	120	24	20.8	5.7	56.9	11.6	N.D.	65.8
6	Toluene	CS ₂ CO ₃	Pd(OAc) ₂	dppe	-	120	24	20.5	5.4	55.5	13.5	N.D.	64.4
7	Toluene	CS ₂ CO ₃	Pd(OAc) ₂	dppf	-	120	24	13.5	5.8	68.8	6.4	N.D.	61.7
8	Toluene	CS ₂ CO ₃	Pd(OAc) ₂	dppb	-	120	24	24.9	6.0	53.7	7.8	N.D.	71.0
9	DME	CS ₂ CO ₃	Pd(OAc) ₂	PCy ₃	-	120	19	15.3	5.9	71.6	5.2	N.D.	90.0
10	EA	K ₂ CO ₃	Pd(OAc) ₂	PCy ₃	PivOH	120	24	60.6	0.0	26.5	3.1	9.8	81.9
11	CPME	K ₂ CO ₃	Pd(OAc) ₂	PCy ₃	PivOH	120	24	63.6	0.0	35.7	0.7	0.0	87.9
12	Toluene	K ₂ CO ₃	Pd(OAc) ₂	P(n-Bu) ₃	PivOH	120	24	54.7	0.0	30.0	6.5	8.7	78.8
13	Toluene	K ₂ CO ₃	Pd(OAc) ₂	P(n-Oct.) ₃	PivOH	120	24	55.4	0.0	28.7	7.9	8.1	79.0
14	Toluene	K ₂ CO ₃	Pd(OAc) ₂	PCy ₃	PivOH	80	90	82.7	0.0	13.1	0.0	4.2	92.9

[Table 9]

[0189]

Abbreviations in the Table represent the following:

DME: Dimethoxyethane

NMP: N-Methylpyrrolidone

EA: Ethyl acetate

CPME: Cyclopentyl methyl ether

HB F_4 : Tetrafluoroboric acid

dppp: 1,1'-Bis(diphenylphosphino)propane

dppe: 1,1'-Bis(diphenylphosphino)ethane

dppb: 1,1'-Bis(diphenylphosphino)butane

dppf: 1,1'-Bis(diphenylphosphino)ferrocene

PCy $_3$: Tricyclohexylphosphine

PivOH: Pivalic acid


n-Oct.: n-Octyl

[Example 11]

Synthesis of ethyl 2-(3-cyano-4-isobutoxyphenyl)-4-methylthiazole-5-carboxylate

[0190]

[Formula 30]



[0191]

To a test tube-type reaction vessel (10 mL) were added ethyl 4-methylthiazole-5-carboxylate (85.5 mg, 0.5 mmol), 5-bromo-2-isobutoxybenzonitrile (127.1 mg, 0.5 mmol) obtained in Reference Example 3, a palladium species, a ligand, a base (1.0 mmol), an additive (0.15 mmol), and a solvent (1.7 mL). Thereafter, the reaction vessel was filled with nitrogen, heated to 120°C under airtight seal, and the reaction mixture was stirred. After the reaction was complete, a portion of the reaction mixture was diluted with DMSO and the solution obtained was measured by HPLC. The total HPLC area % of the compounds A to D and TM was set as 100% and the calculated yield of the target material was computed from the HPLC area % of the target material. The results are shown in Table 10.

[0192]

High Performance Liquid Chromatography:

Analytical System: G1315A Hewlett Packard Series 1100

Software: ChemStation for LC 3D

Experimental Conditions:

Column: Phenomenex Luna Phenyl-Hexyl 5um 4.6×100 mm

Flow: 1.0 mL/min

Wavelength: 240nm

Temperature: 40°C A-Solvent: 5%MeCN/95% H_2O + 0.05%TFA

B-solvent: 95%MeCN/5% H_2O + 0.05%TFA

Gradient:

0-1 min 10% B-Solv.

1-14 min 10-70% B-Solv.

14-24 min 70-80% B-Solv.

24-25 min 80-100% B-Solv.

25-30 min 100% B-Solv.

30-32 min 100-10% B-Solv.

32-35 min 10% B-Solv

[Table 10]

Experiment No.	Reaction Conditions							HPLC area %					Calculated Yield % TM
	Solvent 0.3M	Base 2 eq.	Pd	Ligand	Additive 0.3 eq.	Temperature °C	Reaction Time hr	TM	A	B	C	D	
1	Toluene	K ₂ CO ₃	Pd(OAc) ₂ (0.1 eq.)	P(t-Bu) ₃ ·HBF ₄ (0.2 eq.)	-	120	5.5	31.4	39.2	29.4	N.D.	N.D.	26.5
2	Toluene	K ₂ CO ₃	Pd(OAc) ₂ (0.1 eq.)	XPhos (0.2 eq.)	-	120	5.5	31.5	38.0	30.6	N.D.	N.D.	26.5
3	Toluene	K ₂ CO ₃	Pd(OAc) ₂ (0.1 eq.)	RuPhos (0.2 eq.)	-	120	5.5	39.4	35.7	24.9	N.D.	N.D.	34.0
4	Toluene	K ₂ CO ₃	Pd(OPiv) ₂ (0.02 eq.)	P(t-Bu) ₃ ·HBF ₄ (0.04 eq.)	-	120	15	90.1	3.1	4.1	0.7	2.1	90.7
5	Toluene	KHCO ₃	Pd(OAc) ₂ (0.02 eq.)	P(t-Bu) ₃ ·HBF ₄ (0.04 eq.)	-	120	15	41.0	34.6	24.1	0.0	0.3	35.7
6	Toluene	K ₂ CO ₃	Pd(OAc) ₂ (0.02 eq.)	P(t-Bu) ₃ ·HBF ₄ (0.04 eq.)	-	120	15	46.0	30.3	22.2	0.8	0.7	40.9
7	Toluene	K ₂ CO ₃	Pd(OAc) ₂ (0.02 eq.)	P(t-Bu) ₃ ·HBF ₄ (0.04 eq.)	AcOH	120	15	47.5	29.6	21.9	0.6	0.4	42.1
8	Toluene	K ₂ CO ₃	Pd(OAc) ₂ (0.02 eq.)	P(t-Bu) ₃ ·HBF ₄ (0.04 eq.)	PivOH	120	15	79.3	10.3	8.2	0.3	1.9	77.1
9	Toluene	K ₂ CO ₃	Pd(OAc) ₂ (0.02 eq.)	P(t-Bu) ₃ ·HBF ₄ (0.04 eq.)	EtCO ₂ H	120	15	62.7	20.0	15.7	0.6	0.9	57.9
10	Toluene	K ₂ CO ₃	Pd(OAc) ₂ (0.02 eq.)	P(t-Bu) ₃ ·HBF ₄ (0.04 eq.)	i-PrCO ₂ H	120	15	93.4	1.0	3.8	0.6	1.2	93.7
11	Toluene	K ₂ CO ₃	Pd(P(t-Bu) ₃) ₂ (0.1 eq.)	-	-	120	6	8.9	50.2	40.9	0.0	0.0	7.1
12	Toluene	K ₂ CO ₃	(Pd(P(t-Bu) ₃)Br) ₂ (0.1 eq.)	-	-	120	6	11.9	49.7	36.8	1.6	0.0	9.8
13	Toluene	K ₂ CO ₃	Pd(P(t-Bu) ₃) ₂ (0.1 eq.)	-	PivOH	120	6	90.7	0.1	6.7	0.0	2.5	91.4
14	Toluene	K ₂ CO ₃	(Pd(P(t-Bu) ₃)Br) ₂ (0.1 eq.)	-	PivOH	120	6	78.0	10.5	0.5	3.5	7.5	87.0

[0193]

Abbreviations in the Table represent the following:

AcOH: acetic acid

EtCO₂H: propionic acid

i-PrCO₂H: 2-methylpropanoic acid


[0194]

[Example 12]

Synthesis of ethyl 2-(3-cyano-4-isobutoxyphenyl)-4-methylthiazole-5-carboxylate

[0195]

[Formula 31]



[0196]

To a test tube-type reaction vessel (20 mL) were added ethyl 4-methylthiazole-5-carboxylate (381.2 mg, 1.5 mmol), 5-bromo-2-isobutoxybenzonitrile (256.8 mg, 1.5 mmol) obtained in Reference Example 3, palladium acetate (6.7 mg, 0.030 mmol), di(*tert*-butyl)cyclohexylphosphine (13.7 mg, 0.060 mmol), potassium carbonate (414.6 mg, 3.0 mmol), an additive (0.45 mmol), and xylene (5.0 mL). Thereafter, the reaction vessel was filled with nitrogen, heated to 120°C under airtight seal, and the reaction mixture was stirred for 5 hours. After the reaction was complete, a portion of the reaction mixture was diluted with DMSO and the solution obtained was measured by HPLC. The total area % of the compounds A to D and TM was set as 100% and the calculated yield of the target material was computed from the HPLC area % of the target material. The results are shown in Table 11.

[0197]

High Performance Liquid Chromatography:

Analytical System: G1315A Hewlett Packard Series 1100

Software: ChemStation for LC 3D

Experimental Conditions:

Column: Phenomenex Luna Phenyl-Hexyl 5 μ m 4.6 $\times$ 100 mm

Flow: 1.0 mL/min

Wavelength: 240 nm

Temperature: 40°C

A-Solvent: 5%MeCN/95%H₂O + 0.05%TFA

B-solvent: 95%MeCN/5%H₂O + 0.05%TFA

Gradient:

0-1 min	10%	B-Solv.
1-14 min	10-70%	B-Solv.
14-24 min	70-80%	B-Solv.
24-25 min	80-100%	B-Solv.
25-30 min	100%	B-Solv.
30-32 min	100-10%	B-Solv.
32-35 min	10% B-Solv.	

Experiment No.	Reaction Conditions							HPLC area %					Calculated Yield % TM
	Solvent 0.3M	Base 2 eq.	Pd 0.02 eq.	Ligand 0.04 eq.	Additive 0.3 eq.	Temperature °C	Reaction Time hr	TM	A	B	C	D	
1	Xylene	-	Pd(OAc) ₂	(t-Bu) ₂ PCy	-	120	5	1.1	59.2	39.5	0.0	0.2	0.9
2	Xylene	K ₂ CO ₃	Pd(OAc) ₂	(t-Bu) ₂ PCy	-	120	5	31.0	40.5	27.4	0.3	0.8	26.7
3	Xylene	K ₂ CO ₃	Pd(OAc) ₂	(t-Bu) ₂ PCy	PivOH	120	5	74.6	13.2	9.1	0.0	3.1	72.5
4	Xylene	K ₂ CO ₃	Pd(OAc) ₂	(t-Bu) ₂ PCy	CyPrCO ₂ H	120	5	28.7	42.5	28.7	0.0	0.1	24.3
5	Xylene	K ₂ CO ₃	Pd(OAc) ₂	(t-Bu) ₂ PCy	tetraMeCyPrCO ₂ H	120	5	45.5	32.5	19.1	0.0	2.8	41.6
6	Xylene	K ₂ CO ₃	Pd(OAc) ₂	(t-Bu) ₂ PCy	(CH ₃ CH ₂) ₂ CHCO ₂ H	120	5	61.8	21.2	12.7	0.5	3.8	59.4
7	Xylene	K ₂ CO ₃	Pd(OAc) ₂	(t-Bu) ₂ PCy	CyPnCO ₂ H	120	5	52.0	28.2	18.7	0.0	1.1	46.8
8	Xylene	K ₂ CO ₃	Pd(OAc) ₂	(t-Bu) ₂ PCy	1-ad-CO ₂ H	120	5	49.9	29.5	20.1	0.0	0.6	44.4
9	Xylene	K ₂ CO ₃	Pd(OAc) ₂	(t-Bu) ₂ PCy	CyPenteneCO ₂ H	120	5	37.0	37.1	25.7	0.0	0.3	31.8
10	Xylene	K ₂ CO ₃	Pd(OAc) ₂	(t-Bu) ₂ PCy	3-THFCO ₂ H	120	5	50.5	29.0	20.1	0.0	0.3	44.9
11	Xylene	K ₂ CO ₃	PdCl ₂	(t-Bu) ₂ PCy	PivOH	120	5	86.2	6.1	5.6	0.0	2.1	85.2

[Table 11]

[0198]

Abbreviations in the Table represent the following:

CyPrCO₂H: cyclopropanecarboxylic acid

tetraMeCyPrCO₂H: 2,2,3,3-tetramethylcyclopropanecarboxylic acid

CyPnCO₂H: cyclopentanecarboxylic acid

1-ad-CO₂H: 1-adamantanecarboxylic acid

CyPenteneCO₂H: 3-cyclopentenecarboxylic acid

3-THFCO₂H: tetrahydrofuran-3-carboxylic acid


[Industrial Applicability]

[0199]

The novel coupling process of the present invention, wherein a phenyl derivative represented by the formula (1) and a heterocyclic derivative represented by the formula (2) are coupled in the presence of a transition metal compound to obtain a phenyl-substituted heterocyclic derivative represented by the formula (3), is useful for producing a xanthine oxidase inhibitor, which is a therapeutic agent for hyperuricemia, or an intermediate thereof in a process involving a small number of steps and, consequently, in high yield and at low cost.


CLAIMS

[Claim 1] A process comprising reacting a compound represented by the following formula (1):




(In the formula (1), R¹ represents a hydrogen atom or halogen atom; R² represents a hydrogen atom, cyano group, nitro group, halogen atom, formyl group, or halomethyl group; A represents a hydrogen atom, C₁ to C₈ alkyl group, C₃ to C₆ cycloalkyl group, phenyl group, fluorine atom (only when X is a bond), or protecting group for a hydroxyl group (only when X is an oxygen atom), wherein A may be substituted by 1 to 3 substituents, such substituent representing a group selected from the group consisting of a halogen atom, C₁ to C₄ alkyl group, C₁ to C₄ alkoxy group, C₁ to C₄ alkylthio group, C₃ to C₆ cycloalkyl group, phenyl group, phenoxy group, and pyridyl group; X represents a bond (only when A is a phenyl group or fluorine atom) or oxygen atom; and Y represents a leaving group)

and a compound represented by the following formula (2):



(In the formula (2), H represents a hydrogen atom;

R^3 represents COOR^{3a} or COR^{3b} ; R^{3a} represents a hydrogen atom, C_1 to C_4 alkyl group, or ester-type protecting group for a carboxyl group; R^{3b} represents a amide-type protecting group for a carboxyl group, the protecting group forming an amide with a neighboring carbonyl group; R^4 represents a halogen atom, or C_1 to C_4 alkyl group in the presence of (i) a transition metal compound selected from the group consisting of zero-valent palladium and a salt of mon- and di-valent palladium, and (ii) a phosphine-type ligand capable of coordinating to the transition metal compound to produce a phenyl-substituted heterocyclic derivative represented by the following formula (3):



(in the formula (3), A, X, R¹, and R² are the same as defined in the formula (1); and R³ and R⁴ are the same as defined in the formula (2)).

[Claim 2] The production process according to claim 1, wherein A is a C₁ to C₈ alkyl group.

[Claim 3] The production process according to claim 1, wherein A is an isobutyl group.

[Claim 4] The production process according to any of claims 1 to 3, wherein X is an oxygen atom.

[Claim 5] The production process according to any of claims 1 to 4, wherein R¹ is a hydrogen atom.

[Claim 6] The production process according to any of claims 1 to 5, wherein R is a cyano group.

[Claim 7] The production process according to any of claims 1 to 6, wherein Y represents a halogen atom, -OC(=O)-(C₁ to C₄ alkyl group), -OC(=O)-(phenyl group), -OS(=O)₂-(C₁ to C₄ alkyl group), -OS(=O)₂-(phenyl group), or a diazonium group wherein, in Y, the C₁ to C₄ alkyl group may be substituted with 1 to 3 halogen atoms and the phenyl group may be substituted with 1 to 3 halogen atoms or C₁ to C₄ alkyl groups.

[Claim 8] The production process according to any of claims 1 to 7, wherein R^4 is a methyl group.

[Claim 9] The production process according to any of claims 1 to 8, wherein the transition metal compound is palladium (II) acetate ($Pd(OAc)_2$), palladium (II) propionate ($Pd(O(C=O)CH_2CH_3)_2$), palladium (II) 2-methylpropanoate ($Pd(O(C=O)CH(CH_3)_2)_2$), palladium pivalate ($Pd(OPiv)_2$), palladium (II) chloride ($PdCl_2$), palladium (I) bromide ($PdBr$), or palladium (II) hydroxide ($Pd(OH)_2$).

[Claim 10] The production process according to any of claims 1 to 8, wherein the transition metal compound is palladium (II) acetate ($Pd(OAc)_2$), palladium (II) propionate ($Pd(O(C=O)CH_2CH_3)_2$), palladium (II) 2-methylpropanoate ($Pd(O(C=O)CH(CH_3)_2)_2$), or palladium pivalate ($Pd(OPiv)_2$).

[Claim 11] The production process according to claims 1 to 10, wherein the phosphine-type

ligand is tri(tert-butyl)phosphine, di(tert-butyl)methylphosphine, tert-butylidicyclohexylphosphine, di(tert-butyl)cyclohexylphosphine, tri(cyclohexyl)phosphine, 2-dicyclohexylphosphino-2',6'-diisopropoxy-1,1'-biphenyl, 2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl, or a salt thereof.

[Claim 12] The production process according to any of claims 1 to 11, wherein a base is additionally present during the course of the reaction.

[Claim 13] The production process according to claim 12, wherein the base is potassium carbonate, potassium bicarbonate, cesium carbonate, or tetra-n-butylammonium fluoride.

[Claim 14] The production process according to any of claims 1 to 13, wherein a silver salt is additionally present during the course of the reaction.

[Claim 15] The production process according to claim 14, wherein the silver salt is silver carbonate.

[Claim 16] The production process according to any of claims 1 to 15, wherein a C₁ to C₁₂ carboxylic acid or salt thereof is additionally present during the course of the reaction.

[Claim 17] The production process according to claim 16, wherein the carboxylic acid or salt thereof is 2-methylpropanoic acid, pivalic acid, or a salt thereof.