

METHODS OF PRODUCING HETERODIMERIC ANTIBODIES

CROSS-REFERENCE TO RELATED APPLICATIONS

5 This application claims the benefit of United States Provisional Application Serial number 62/781,180, filed 18 December 2018, the entire contents of which are incorporated herein by reference.

FIELD OF THE INVENTION

The invention relates to methods of producing heterodimeric antibodies.

SEQUENCE LISTING

10 This application contains a Sequence Listing submitted via EFS-Web, the entire content of which is incorporated herein by reference. The ASCII text file, created on 10 December 2019, is named JBI6029WOPCT1_ST25.txt and is 21 kilobytes in size.

BACKGROUND OF THE INVENTION

15 Monoclonal antibodies have demonstrated success as therapeutic molecules, especially in the treatment of cancer. Bispecific antibodies are expected to increase the potency and efficacy of monoclonal antibody therapies as they may be used to direct a drug or toxic compound to target cells, to redirect effector mechanisms to disease-associated sites or to increase specificity for tumor cells, for example by binding to one or more target molecules that are expressed on tumor cells. Further, by combining the specificity of two monoclonal antibodies in one, bispecific antibodies could potentially engage a greater array of mechanisms of action.

25 Different formats of bispecific antibodies and methods of producing them have been described, however a challenge exists to produce the bispecific antibodies in large scale by optimizing manufacturing processes.

BRIEF SUMMARY OF THE INVENTION

30 The invention provides a method of producing a heterodimeric antibody, comprising providing a first homodimeric antibody comprising a first Fc region of an immunoglobulin comprising a first CH3 region and a second homodimeric antibody comprising a second Fc region of an immunoglobulin comprising a second CH3 region, wherein the amino acid sequences of the first CH3 region and the second CH3 regions are different and are such

35

that a heterodimeric interaction between the first CH3 region and the second CH3 region is stronger than a homodimeric interaction between the first CH3 region or a homodimeric interaction between the second CH3 region;

combining the first homodimeric antibody and the second homodimeric antibody into a mixture; and

incubating the mixture in the presence of a reducing agent and ambient dissolved oxygen (DO₂) to produce the heterodimeric antibody, wherein the method lacks one or more steps of measuring a percentage (%) DO₂ in the mixture, controlling the % DO₂ in the mixture, or adding oxygen into the mixture during producing the heterodimeric antibody.

The invention also provides a method of producing a heterodimeric antibody, comprising

providing a first homodimeric antibody comprising a first Fc region of an immunoglobulin comprising a first CH3 region and a second homodimeric antibody comprising a second Fc region of an immunoglobulin comprising a second CH3 region, wherein the amino acid sequences of the first CH3 region and the second CH3 regions are different and are such that the heterodimeric interaction between the first CH3 region and the second CH3 region is stronger than a homodimeric interaction between the first CH3 region or a homodimeric interaction between the second CH3 region;

combining the first homodimeric antibody and the second homodimeric antibody into a mixture;

incubating the mixture in the presence of a reducing agent; and

removing the reducing agent to produce the heterodimeric antibody, wherein

the percentage (%) of dissolved oxygen (DO₂) is controlled to be about 30% or lower

during step of incubating the mixture in the presence of the reducing agent, during step of

removing the reducing agent to produce the heterodimeric antibody, or during steps of

incubating the mixture in the presence of the reducing agent and removing the reducing agent to produce the heterodimeric antibody.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows a summary of Fab-arm exchange.

FIG. 2 shows process steps during reduction and ultrafiltration/diafiltration (UF/DF) during manufacturing of bispecific antibodies using Fab-arm exchange.

FIG. 3 shows molecular details of the thiol-disulfide exchange reactions

FIG. 4 shows a summary of the reactions that may take place during Fab-arm exchange in the presence of 2-MEA.

FIG. 5 shows a cartoon of the UF/DF setup and the location of DO₂ and pH sensors where RmV = relative millivolts.

FIG. 6 shows percent (%) DO₂ during UF/DF measured in the retentate, permeate and inlet lines during manufacturing of bispecific antibody A in low DO₂ conditions during UF/DF. a.s.: air saturated.

FIG. 7 shows percent (%) DO₂ during UF/DF measured in the retentate, permeate and inlet lines during manufacturing of bispecific antibody B in low DO₂ conditions during UF/DF. a.s.: air saturated.

FIG. 8 shows percent (%) DO₂ over time during reduction in low DO₂ conditions (Low DO reduction) or at ambient DO₂ (target reduction). a.s.: air saturated.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as though fully set forth.

It is to be understood that the terminology used herein is for describing particular embodiments only and is not intended to be limiting. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains.

Although any methods and materials similar or equivalent to those described herein may be used in the practice for testing of the present invention, exemplary materials and methods are described herein. In describing and claiming the present invention, the following terminology will be used.

As used in this specification and the appended claims, the singular forms “a,” “an,” and “the” include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to “a cell” includes a combination of two or more cells, and the like.

The transitional terms “comprising,” “consisting essentially of,” and “consisting of” are intended to connote their generally accepted meanings in the patent vernacular; that is, (i) “comprising,” which is synonymous with “including,” “containing,” or “characterized by,” is inclusive or open-ended and does not exclude additional, unrecited elements or method steps; (ii) “consisting of” excludes any element, step, or ingredient not specified in the claim; and (iii) “consisting essentially of” limits the scope of a claim to the specified materials or steps “and those that do not materially affect the basic and novel characteristic(s)” of the claimed invention. Embodiments described in terms of the phrase

“comprising” (or its equivalents) also provide as embodiments those independently described in terms of “consisting of” and “consisting essentially of.”

“**Antibodies**” refer to immunoglobulin molecules having two heavy chains (HC) and two light chains (LC) interconnected by disulfide bonds. Each heavy chain is comprised of a heavy chain variable region (VH) and a heavy chain constant region, the heavy chain constant region divided into regions CH1, hinge, CH2 and CH3. Each light chain is comprised of a light chain variable region (VL) and a light chain constant region (CL). The VH and the VL may be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with framework regions (FR). Each VH and VL is composed of three CDRs and four FR segments, arranged from amino-to-carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. Antibodies include monoclonal antibodies including murine, human, humanized and chimeric antibodies, bispecific or multispecific antibodies.

“**Complementarity determining regions (CDR)**” are antibody regions that bind an antigen. There are three CDRs in the VH (HCDR1, HCDR2, HCDR3) and three CDRs in the VL (LCDR1, LCDR2, LCDR3). CDRs may be defined using various delineations such as Kabat (Wu *et al.* (1970) *J Exp Med* 132: 211-50) (Kabat *et al.*, Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md., 1991), Chothia (Chothia *et al.* (1987) *J Mol Biol* 196: 901-17), IMGT (Lefranc *et al.* (2003) *Dev Comp Immunol* 27: 55-77) and AbM (Martin and Thornton (1996) *J Biol Biol* 263: 800-15). The correspondence between the various delineations and variable region numbering are described (see e.g. Lefranc *et al.* (2003) *Dev Comp Immunol* 27: 55-77; Honegger and Pluckthun, *J Mol Biol* (2001) 309:657-70; International ImMunoGeneTics (IMGT) database; Web resources, [http://www_imgt_org](http://www.imgt.org)). Available programs such as abYsis by UCL Business PLC may be used to delineate CDRs. The term “CDR”, “HCDR1”, “HCDR2”, “HCDR3”, “LCDR1”, “LCDR2” and “LCDR3” as used herein includes CDRs defined by any of the methods described supra, Kabat, Chothia, IMGT or AbM, unless otherwise explicitly stated in the specification.

Immunoglobulins may be assigned to five major classes, IgA, IgD, IgE, IgG and IgM, depending on the heavy chain constant region amino acid sequence. IgA and IgG are further sub-classified as isotypes IgA1, IgA2, IgG1, IgG2, IgG3 and IgG4. Antibody light chains of any vertebrate species may be assigned to one of two clearly distinct types, namely kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

“**Antigen-binding fragment**” refers to a portion of an immunoglobulin molecule that retains the antigen binding properties of the parental full length antibody. Exemplary antigen-binding fragments are heavy chain complementarity determining regions (HCDR) 1, 2 and/or 3, light chain complementarity determining regions (LCDR) 1, 2 and/or 3, the VH, the VL, the VH and the VL, Fab, F(ab')₂, Fd and Fv fragments as well as domain antibodies (dAb) consisting of either one VH domain or one VL domain. The VH and the VL domains may be linked together via a synthetic linker to form various types of single chain antibody designs in which the VH/VL domains pair intramolecularly, or intermolecularly in those cases when the VH and VL domains are expressed by separate chains, to form a monovalent antigen binding site, such as single chain Fv (scFv) or diabody; described for example in Int. Pat. Publ. No. WO1998/44001, Int. Pat. Publ. No. WO1988/01649; Int. Pat. Publ. No. WO1994/13804; Int. Pat. Publ. No. WO1992/01047.

“**Monoclonal antibody**” refers to an antibody obtained from a substantially homogenous population of antibody molecules, i.e., the individual antibodies comprising the population are identical except for possible well-known alterations such as removal of C-terminal lysine from the antibody heavy chain or post-translational modifications such as amino acid isomerization or deamidation, methionine oxidation or asparagine or glutamine deamidation. Monoclonal antibodies typically bind one antigenic epitope. A bispecific monoclonal antibody binds two distinct antigenic epitopes. Monoclonal antibodies may have heterogeneous glycosylation within the antibody population. Monoclonal antibody may be monospecific or multispecific such as bispecific, monovalent, bivalent or multivalent.

“**Fab-arm**” refers to one heavy chain-light chain pair of an antibody.

“**Homodimerization**” refers to an interaction of two heavy chains having identical CH3 amino acid sequences.

“**Homodimer**” refers to an antibody having two heavy chains which have identical CH3 region amino acid sequences.

“**Heterodimerization**” refers to an interaction of two heavy chains having non-identical CH3 amino acid sequences.

“**Heterodimer**” refers to an antibody having two heavy chains which differ in their amino acid sequence in the CH3 region by one or more amino acids.

“**Fc region**” or “**Fc domain**” refers to an antibody region comprising at least a portion of a hinge region, a CH2 region and a CH3 region. The Fc region may be generated by digestion of an antibody with papain, or pepsing where the Fc region is the fragment obtained thereby, which includes one or both CH2-CH3 regions of an

immunoglobulin and a portion of the hinge region. The constant domain of an antibody heavy chain defines the antibody isotype, e.g. IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, IgE. The Fc-region mediates the effector functions of antibodies with cell surface Fc receptors and proteins of the complement system.

5 “**CH1 region**” or “**CH1 domain**” refers to the CH1 region of an immunoglobulin. The CH1 region of a human IgG1 antibody corresponds to amino acid residues 118-215. However, the CH1 region may also be any of the other antibody isotypes as described herein.

10 “**CH2 region**” or “**CH2 domain**” refers to the CH2 region of an immunoglobulin. The CH2 region of a human IgG1 antibody corresponds to amino acid residues 231-340. However, the CH2 region may also be any of the other antibody isotypes as described herein.

15 “**CH3 region**” or “**CH3 domain**” refers to the CH3 region of an immunoglobulin. The CH3 region of human IgG1 antibody corresponds to amino acid residues 341-446. However, the CH3 region may also be any of the other antibody isotypes as described herein.

20 “**Hinge**” or “**hinge region**” refers to the hinge region of an immunoglobulin. The hinge region of human IgG1 antibody generally corresponds to amino acids 216-230 according to the EU numbering system. “Hinge” may also be considered to include additional residues termed the upper and lower hinge regions, such as from amino acid residues 216 to 239.

 “**Mixture**” refers to an aqueous solution of two or more antibodies.

 “**Reducing agent**” refers to an agent that is capable of reducing the inter-chain disulfide bonds in the hinge region of an antibody.

25 “**GMP-compliant conditions**” refers to manufacturing under good manufacturing practice (CGMP) regulations enforced by the FDA. CGMPs provide for systems that assure proper design, monitoring, and control of manufacturing processes and facilities. Adherence to the CGMP regulations assures the identity, strength, quality, and purity of drug products by requiring that manufacturers of medications adequately control
30 manufacturing operations. This includes establishing strong quality management systems, obtaining appropriate quality raw materials, establishing robust operating procedures, detecting and investigating product quality deviations, and maintaining reliable testing laboratories. This formal system of controls at a pharmaceutical company, if adequately put into practice, helps to prevent instances of contamination, mix-ups, deviations,
35 failures, and errors. This assures that drug products meet their quality standards.

“**Drug substance**” or “**DS**” refers to Any substance or mixture of substances intended to be used in the manufacture of a drug (medicinal) product and that, when used in the production of a drug, becomes an active ingredient of the drug product. Such substances are intended to furnish pharmacological activity or other direct effect in the diagnosis, cure, mitigation, treatment, or prevention of disease or to affect the structure or function of the body.

“**Drug product**” or “**DP**” refers to a finished dosage form, for example, a tablet, capsule or solution that contains an active pharmaceutical ingredient (e.g. drug substance), generally, but not necessarily, in association with inactive ingredients.

“**Reference product**” refers to an approved biological product against which a biosimilar product is compared. A reference product is approved based on, among other things, a full complement of safety and effectiveness data and is approved in at least one of the U.S., Europe, or Japan.

“**Biosimilar**” (of an approved reference product/biological drug) refers to a biological product that is highly similar to the reference product notwithstanding minor differences in clinically inactive components with no clinically meaningful differences between the biosimilar and the reference product in terms of safety, purity and potency, based upon data derived from (a) analytical studies that demonstrate that the biological product is highly similar to the reference product notwithstanding minor differences in clinically inactive components; (b) animal studies (including the assessment of toxicity); and/or (c) a clinical study or studies (including the assessment of immunogenicity and pharmacokinetics or pharmacodynamics) that are sufficient to demonstrate safety, purity, and potency in one or more appropriate conditions of use for which the reference product is licensed and intended to be used and for which licensure is sought for the biosimilar.

The biosimilar may be an interchangeable product that may be substituted for the reference product at the pharmacy without the intervention of the prescribing healthcare professional. To meet the additional standard of “interchangeability,” the biosimilar is expected to produce the same clinical result as the reference product in any given patient and, if the biosimilar is administered more than once to an individual, the risk in terms of safety or diminished efficacy of alternating or switching between the use of the biosimilar and the reference product is not greater than the risk of using the reference product without such alternation or switch. The biosimilar utilizes the same mechanisms of action for the proposed conditions of use to the extent the mechanisms are known for the reference product. The condition or conditions of use prescribed, recommended, or suggested in the labeling proposed for the biosimilar must have been previously approved for the reference

product. The route of administration, the dosage form, and/or the strength of the biosimilar must be the same as those of the reference product and the biosimilar must be manufactured, processed, packed or held in a facility that meets standards designed to assure that the biosimilar continues to be safe, pure and potent. The biosimilar may include minor modifications in the amino acid sequence when compared to the reference product, such as N- or C-terminal truncations that are not expected to change the biosimilar performance. The reference product may be approved in at least one of the U.S., Europe, or Japan.

“**Isolated**” refers to a homogenous population of molecules (such as synthetic polynucleotides or a protein such as an antibody) which have been substantially separated and/or purified away from other components of the system the molecules are produced in, such as a recombinant cell, as well as a protein that has been subjected to at least one purification or isolation step. “Isolated antibody” refers to an antibody that is substantially free of other cellular material and/or chemicals and encompasses antibodies that are isolated to a higher purity, such as to 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% purity.

“**Humanized antibody**” refers to an antibody in which at least one CDR is derived from non-human species and at least one framework is derived from human immunoglobulin sequences. Humanized antibody may include substitutions in the frameworks so that the frameworks may not be exact copies of expressed human immunoglobulin or human immunoglobulin germline gene sequences.

“**Human antibody**” refers to an antibody that is optimized to have minimal immune response when administered to a human subject. Variable regions of human antibody are derived from human immunoglobulin sequences. If human antibody contains a constant region or a portion of the constant region, the constant region is also derived from human immunoglobulin sequences. Human antibody comprises heavy and light chain variable regions that are “derived from” sequences of human origin if the variable regions of the human antibody are obtained from a system that uses human germline immunoglobulin or rearranged immunoglobulin genes. Such exemplary systems are human immunoglobulin gene libraries displayed on phage, and transgenic non-human animals such as mice or rats carrying human immunoglobulin loci. “Human antibody” typically contains amino acid differences when compared to the immunoglobulins expressed in humans due to differences between the systems used to obtain the human antibody and human immunoglobulin loci, introduction of somatic mutations or intentional introduction of substitutions into the frameworks or CDRs, or both. Typically,

“human antibody” is at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical in amino acid sequence to an amino acid sequence encoded by human germline immunoglobulin or rearranged immunoglobulin genes. In some cases, “human antibody” may contain
5 consensus framework sequences derived from human framework sequence analyses, for example as described in Knappik *et al.*, (2000) *J Mol Biol* 296:57-86, or synthetic HCDR3 incorporated into human immunoglobulin gene libraries displayed on phage, for example as described in Shi *et al.*, (2010) *J Mol Biol* 397:385-96, and in WO2009/085462. Antibodies in which at least one CDR is derived from a non-human species are not
10 included in the definition of “human antibody”.

“**Recombinant**” refers to DNA, antibodies and other proteins that are prepared, expressed, created or isolated by recombinant means when segments from different sources are joined to produce recombinant DNA, antibodies or proteins. “Recombinant antibody” includes all antibodies that are prepared, expressed, created or isolated by
15 recombinant means, such as antibodies isolated from an animal (e.g., a mouse) that is transgenic or transchromosomal for human immunoglobulin genes or a hybridoma prepared therefrom (described further below), antibodies isolated from a host cell transformed to express the antibody, antibodies isolated from a recombinant, combinatorial antibody library, and antibodies prepared, expressed, created or isolated by
20 any other means that involve splicing of human immunoglobulin gene sequences to other DNA sequences, or antibodies that are generated *in vitro* using Fab arm exchange such as bispecific antibodies.

“**Bispecific**” refers to an antibody that specifically binds two distinct antigens or two distinct epitopes within the same antigen. The bispecific antibody may have cross-
25 reactivity to other related antigens or can bind an epitope that is shared between two or more distinct antigens.

“**Multispecific**” refers to an antibody that specifically binds at least two distinct antigens or at least two distinct epitopes within the same antigen. Multispecific antibody may bind for example two, three, four or five distinct antigens or distinct epitopes within
30 the same antigen.

“**About**” means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, i.e., the limitations of the measurement system. Unless explicitly stated otherwise within the Examples or elsewhere in the Specification in the

context of a particular assay, result or embodiment, “about” means within one standard deviation per the practice in the art, or a range of up to 5%, whichever is larger.

“**Variant**” refers to a polypeptide or a polynucleotide that differs from a reference polypeptide or a reference polynucleotide by one or more modifications for example, substitutions, insertions or deletions.

“**Mutation**” refers to an engineered or naturally occurring alteration in a polypeptide or polynucleotide sequence when compared to a reference sequence. The alteration may be a substitution, insertion or deletion of one or more amino acids or polynucleotides.

The numbering of amino acid residues in the antibody constant region throughout the specification is according to the EU index as described in Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991), unless otherwise explicitly stated. Antibody constant chain numbering can be found for example at ImMunoGeneTics website, at IMGT Web resources at IMGT Scientific charts.

The mutations in the CH3 region described herein are expressed as modified position(s) in the first CH3 domain of the first heavy chain/ modified position(s) in the second CH3 domain of the second heavy chain. For example, F405L/K409R refers to a F405L mutation in the first CH3 region and K409R mutation in the second CH3 region. L351Y_F405A_Y407V/T394W refers to L351Y, F405A and Y407V mutations in the first CH3 region and T394W mutation in the second CH3 region. D399FHKRQ/K409AGR refers to mutation in which D399 may be replaced by F, H, K, R or Q, and K409 may be replaced by A, G, R or H.

Conventional one and three-letter amino acid codes are used herein as shown in **Table 1.**

Table 1.

Amino acid	Three-letter code	One-letter code	Amino acid	Three-letter code	One-letter code
Alanine	Ala	A	Leucine	Leu	L
Arginine	Arg	R	Lysine	Lys	K
Asparagine	Asn	N	Methionine	Met	M
Aspartate	Asp	D	Phenylalanine	Phe	F
Cysteine	Cys	C	Proline	Pro	P
Glutamate	Gln	E	Serine	Ser	S
Glutamine	Glu	Q	Threonine	Thr	T
Glycine	Gly	G	Tryptophan	Trp	W
Histidine	His	H	Tyrosine	Tyr	Y

Isoleucine	Ile	I		Valine	Val	V
------------	-----	---	--	--------	-----	---

Methods of the invention

The invention provides methods of producing heterodimeric antibodies using Fab-arm exchange. The invention is based on, at least in part, on the identification that, contrary to what has been disclosed in literature (see, e.g., US2014/0303356), oxygen is not required for reformation of the disulfide bonds after their reduction to form stable heterodimeric antibodies during Fab-arm exchange, hence providing alternative approaches for process control during large scale production of heterodimeric antibodies (e.g., bispecific antibodies).

The invention provides a method of producing a heterodimeric antibody, comprising providing a first homodimeric antibody comprising a first Fc region of an immunoglobulin comprising a first CH3 region and a second homodimeric antibody comprising a second Fc region of an immunoglobulin comprising a second CH3 region, wherein the amino acid sequences of the first CH3 region and the second CH3 regions are different and are such that a heterodimeric interaction between the first CH3 region and the second CH3 region is stronger than a homodimeric interaction between the first CH3 region or a homodimeric interaction between the second CH3 region; combining the first homodimeric antibody and the second homodimeric antibody into a mixture; and incubating the mixture in the presence of a reducing agent and ambient dissolved oxygen (DO₂) to produce the heterodimeric antibody, wherein the method lacks one or more steps of measuring a percentage (%) DO₂ in the mixture, controlling the % DO₂ in the mixture, or adding oxygen into the mixture during producing the heterodimeric antibody.

In some embodiments, the % DO₂ in the mixture is between about 10% and about 90% during producing the heterodimeric antibody.

In some embodiments, % DO₂ in the mixture is about 30% or more in the mixture prior to addition of the denaturing agent.

The invention also provides a method of producing a heterodimeric antibody, comprising providing a first homodimeric antibody comprising a first Fc region of an immunoglobulin comprising a first CH3 region and a second homodimeric antibody comprising a second Fc region of an immunoglobulin comprising a second CH3 region, wherein the amino acid sequences of the first CH3 region and the second CH3 regions are different and are such that the heterodimeric interaction between the first CH3 region and the second CH3 region

is stronger than a homodimeric interaction between the first CH3 region or a homodimeric interaction between the second CH3 region;

combining the first homodimeric antibody and the second homodimeric antibody into a mixture;

- 5 incubating the mixture in the presence of a reducing agent; and
removing the reducing agent to produce the heterodimeric antibody, wherein

the percentage (%) of dissolved oxygen (DO_2) is controlled to be about 30% or lower in step c), step d) or both in step c) and step d).

- 10 In some embodiments, the % DO_2 the mixture is about 25% or less, 20% or less, about 15% or less, about 10% or less, about 9% or less, about 8% or less, about 7% or less, about 6% or less, about 5% or less, about 4% or less, about 3% or less, about 2% or less or about 1% or less.

In some embodiments, % DO_2 is controlled by displacing oxygen from the mixture by overlaying the mixture with an inert gas such as nitrogen. The concentration of dissolved oxygen may be monitored by known methods, such as utilizing a dissolved oxygen probe.

- 15 “Stronger” in terms of the heterodimeric interaction of the first CH3 region and the second CH3 region may be more than two times stronger, for example more than three times stronger, more than four times stronger or more than five times stronger than the strongest of the homodimeric interaction between the first CH3 region or a homodimeric interaction between the second CH3 region. The strength of the interaction of the CH3 domains may be measured using mass spectrometry. In an exemplary assay, constructs comprising the first CH3 region and the second CH3 region or alternatively both including the CH2 region are made using standard molecular biology techniques.
- 25 Samples are prepared that contain the first CH3 domain, the second CH3 domain or the first CH3 domain and the second CH3 domain and buffer-exchanged to 100 mM ammonium acetate pH 7, using 10 kDa MWCO spin-filter columns. Aliquots (~1 μL) of serial diluted samples (20 μM -25 nM; monomer equivalent) are loaded into gold-plated borosilicate capillaries for analysis on a LCT mass spectrometer (Waters). The monomer signal, M_s , is defined as the area of the monomer peaks as a fraction of the area of all peaks in the spectrum ($M_s/(M_s+D_s)$ where D_s =the dimer signal). The concentration of monomer at equilibrium, $[M]_{\text{eq}}$, is defined as $M_s/[M]_0$ where $[M]_0$ is the overall protein concentration in terms of monomer. The dimer concentration at equilibrium, $[D]_{\text{eq}}$, is defined as $([M]_0-[M]_{\text{eq}})/2$. The K_D , for the homodimeric CH3 interactions and the
- 30

heterodimeric CH3 interactions is then extracted from the gradient of a plot of $[D]_{eq}$ versus $[M]_{eq}^2$.

In some embodiments, the first homodimeric antibody and the second homodimeric antibody are combined into the mixture at a molar ratio of about 1:1.

5 In some embodiments, the first homodimeric antibody and the second homodimeric antibody are combined into the mixture at the molar ratio of about 1.05:1.

In some embodiments, the first homodimeric antibody and the second homodimeric antibody are combined into the mixture at the molar ratio of between about 1:1.03 to about 1:2. In some embodiments, the first homodimeric antibody and the second
10 homodimeric antibody are combined into the mixture at the molar ratio of between about 1:1.05 to 1:1.5. In some embodiments, the first homodimeric antibody and the second homodimeric antibody are combined into the mixture at the molar ratio of between about 1:1.1 to 1:1.5. In some embodiments, the first homodimeric antibody and the second homodimeric antibody are combined into the mixture at the molar ratio of between about
15 1:1.1 to 1:1.4. In some embodiments, the first homodimeric antibody and the second homodimeric antibody are combined into the mixture at the molar ratio of between about 1:1.15 to 1:1.35. In some embodiments, the first homodimeric antibody and the second homodimeric antibody are combined into the mixture at the molar ratio of between about 1:1.2 to 1:1.3.

20 In some embodiments, total concentration of immunoglobulin in the mixture is between about 1 g/L and 70 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is between about 8 g/L and about 50 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is between about 8 g/L and about 13 g/L. In some embodiments, total concentration of immunoglobulin in
25 the mixture is about 9.0 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 9.5 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 10 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 10.5 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 11.0 g/L. In
30 some embodiments, total concentration of immunoglobulin in the mixture is about 11.5 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 12.0 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 12.5 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 13.0 g/L. In some embodiments, total concentration of immunoglobulin
35 in the mixture is about 13.5 g/L. In some embodiments, total concentration of

immunoglobulin in the mixture is about 14.0 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 14.5 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 15.0 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 20.0 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 25.0 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 30.0 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 35.0 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 40.0 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 45.0 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 50.0 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 55.0 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 60.0 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 65.0 g/L. In some embodiments, total concentration of immunoglobulin in the mixture is about 70.0 g/L.

In some embodiments, the mixture is incubated in the presence of the reducing agent for about 10 minutes or longer. In some embodiments, the mixture is incubated in the presence of the reducing agent from about 10 minutes to about 30 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent from about 10 minutes to about 24 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 15 minutes. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 20 minutes. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 30 minutes. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 40 minutes. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 50 minutes. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 1 hour. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 2 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 3 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 4 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 4 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 5 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 6 hours. In some embodiments,

the mixture is incubated in the presence of the reducing agent for about 7 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 8 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 9 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 10 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 11 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 12 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 13 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 14 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 15 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 16 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 17 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 18 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 19 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 20 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 21 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 22 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 23 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 24 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 25 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 26 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 27 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 28 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 29 hours. In some embodiments, the mixture is incubated in the presence of the reducing agent for about 30 hours.

30 In some embodiments, the reducing agent is 2-mercaptoethylamine (2-MEA). In some embodiments, the reducing agent is a chemical derivative of 2-MEA. In some embodiments, the reducing agent is L-cysteine. In some embodiments, the reducing agent is D-cysteine. In some embodiments, the reducing agent is glutathione. In some embodiments, the reducing agent is tris(2-carboxyethyl)phosphine.

In some embodiments, the concentration of the reducing agent in the mixture is between about 0.1 mM to about 1 M. In some embodiments, the concentration of the reducing agent in the mixture is between about 1.0 mM to about 500 mM. In some embodiments, the concentration of the reducing agent in the mixture is between about 5.0 mM to about 100 mM. In some embodiments, the concentration of the reducing agent in the mixture is about 10 mM. In some embodiments, the concentration of the reducing agent in the mixture is about 15 mM. In some embodiments, the concentration of the reducing agent in the mixture is about 20 mM. In some embodiments, the concentration of the reducing agent in the mixture is about 25 mM. In some embodiments, the concentration of the reducing agent in the mixture is about 30 mM. In some embodiments, the concentration of the reducing agent in the mixture is about 35 mM. In some embodiments, the concentration of the reducing agent in the mixture is about 40 mM. In some embodiments, the concentration of the reducing agent in the mixture is about 50 mM. In some embodiments, the concentration of the reducing agent in the mixture is about 60 mM. In some embodiments, the concentration of the reducing agent in the mixture is about 70 mM. In some embodiments, the concentration of the reducing agent in the mixture is about 80 mM. In some embodiments, the concentration of the reducing agent in the mixture is about 90 mM. In some embodiments, the concentration of the reducing agent in the mixture is about 100 mM.

In some embodiments, the concentration of 2-MEA in the mixture is between about 10 mM and about 100 mM. In some embodiments, the concentration of 2-MEA in the mixture is between about 20 mM and about 90 mM. In some embodiments, the concentration of 2-MEA in the mixture is between about 20 mM and about 80 mM. In some embodiments, the concentration of 2-MEA in the mixture is between about 20 mM and about 70 mM. In some embodiments, the concentration of 2-MEA in the mixture is between about 20 mM and about 60 mM. In some embodiments, the concentration of 2-MEA in the mixture is between about 20 mM and about 50 mM. In some embodiments, the concentration of 2-MEA in the mixture is between about 20 mM and about 40 mM. In some embodiments, the concentration of 2-MEA in the mixture is about 20 mM. In some embodiments, the concentration of 2-MEA in the mixture is about 25 mM. In some embodiments, the concentration of 2-MEA in the mixture is about 30 mM. In some embodiments, the concentration of 2-MEA in the mixture is about 35 mM. In some embodiments, the concentration of 2-MEA in the mixture is about 40 mM.

In some embodiments the mass ratio of the total immunoglobulin in the mixture to the total reducing agent is between about 1.0 and about 5.0. In some embodiments the

mass ratio of the total immunoglobulin in the mixture to the total reducing agent is between about 1.4 and about 3.8. In some embodiments the mass ratio of the total immunoglobulin in the mixture to the total reducing agent is between about 1.4 and about 3.5. In some embodiments, the mass ratio is between about 1.8 and about 3.8. In some
 5 embodiments, the mass ratio is between about 2.3 and about 3.0. In some embodiments, the mass ratio is between about 1.6 and about 2.1. In some embodiments, the mass ratio is about 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3.0, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 4.0, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, or 5.0. “**Mass ratio**” refers to the total immunoglobulin in gram per liter to the total
 10 reducing agent in gram per liter. “**Total immunoglobulin**” refers to the amount of the two parental antibodies in the mixture at the beginning of the reduction step.

In some embodiments, the mixture comprises a buffer. In some embodiments, the buffer comprises a sodium acetate buffer. In some embodiments, the buffer further comprises NaCl. In some embodiments, the buffer comprises about 100 mM sodium
 15 acetate and about 30 mM NaCl. In some embodiments, pH of the buffer is about 7.3. Other buffers may be used, such as 1 x Dulbecco’s phosphate-buffered saline (DPBS), sodium phosphate buffer, potassium phosphate buffer, Tris buffer, histidine buffer or citrate buffer.

In some embodiments, the method further comprising a step of removing the
 20 reducing agent from the mixture.

In some embodiments, the reducing agent is removed by filtration.

In some embodiments, filtration is diafiltration.

In some embodiments, the first homodimeric antibody and the second homodimeric antibody are an IgG1, IgG2 or IgG4 isotype.

25 In some embodiments, the first homodimeric antibody and the second homodimeric antibody are an IgG1 isotype. In some embodiments, the first homodimeric antibody and the second homodimeric antibody are an IgG2 isotype. In some embodiments, the first homodimeric antibody and the second homodimeric antibody are an IgG4 isotype.

30 In some embodiments, the first CH3 domain and the second CH3 domain comprise following mutations when compared to the wild-type IgG1 of **SEQ ID NO: 1**, F405L/K409R, T350I_K370T_F405L/K409R, K370W/K409R, D399AFGHILMNRSTVWY/K409R, T366ADEFHILMQVY/K409R, L368ADEGHNRSTVQ/K409AGRH, D399FHKRQ/K409AGRH,
 35 F405IKLSTVW/K409AGRH, Y407LWQ/K409AGRH, T366Y/F405A, T366W/F405W,

- F405W/Y407A, T394W/Y407T, T394S/Y407A, T366W/T394S, F405W/T394S, T366W/T366S_L368A_Y407V, L351Y_F405A_Y407V/T394W, T366I_K392M_T394W/F405A_Y407V, T366L_K392M_T394W/F405A_Y407V, L351Y_Y407A/T366A_K409F, L351Y_Y407A/T366V_K409F, Y407A/T366A_K409F,
- 5 T350V_L351Y_F405A_Y407V/T350V_T366L_K392L_T394W, K409D/D399K, K409E/D399R, K409D_K360D/D399K_E356K, K409D_K360D/D399E_E356K, K409D_K370D/D399K_E357K, K409D_K370D/D399E_E357K, K409D_K392D/D399K_E356K_E357K or K409D_K392D/D399E_E356K_E357K. The mutations may also be introduced to the wild-type IgG2 of **SEQ ID NO: 2** or wild-type
- 10 IgG4 of **SEQ ID NO: 3**, in which case, as is well-known, the wild-type residue at the particular mutation site may not correspond to that of IgG1 due to sequence differences between wild-type IgG1, IgG2 and IgG4. For example, IgG4 mutations wild-type/F405L_R409K correspond to IgG1 mutations F405L/K409R, and IgG4 mutations R409D/D399K correspond to IgG1 mutations K409D/D399K. The mutations are
- 15 compared to the reference wild-type IgG1 of SEQ ID NO: 1, IgG2 of **SEQ ID NO: 2** and IgG4 of **SEQ ID NO: 3**.

- In some embodiment,s the first Fc region and/or the second Fc region comprise one or more mutations when compared to the wild-type IgG1 of **SEQ ID NO: 1**, wild-type IgG2 of **SEQ ID NO: 2**, wild-type IgG4 of **SEQ ID NO: 3** that modulate binding of the
- 20 first Fc region and/or the second Fc region to an Fc γ receptor (Fc γ R), an FcRn, or to protein A.

Wild-type IgG1 (**SEQ ID NO: 1**)

- ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAV
 25 LQSSGLYSLSSVVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPC
 PAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEV
 HNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISK
 AKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKT
 30 TPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

Wild-type IgG2 (**SEQ ID NO: 2**)

- ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAV
 LQSSGLYSLSSVVPSSNFGTQTYTCNVDHKPSNTKVDKTVERKCCVECPAP
 35 PVAGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVQFNWYVDGVEVHNAK
 TKPREEQFNSTFRVVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQP
 REPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDISVEWESNGQPENNYKTTPPML
 DSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

Wild-type IgG4 (SEQ ID NO: 3)

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAV
 LQSSGLYSLSVVTPSSSLGKTYTCNVDPKPSNTKVDKRVESKYGPPCPSCPAP
 EFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNA
 5 KTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQ
 PREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPV
 LDDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLSLGLGK

In some embodiments, the Fc γ R is Fc γ RI, Fc γ RIIa, Fc γ RIIb or Fc γ RIII.

10 In some embodiments, the one or more substitutions that modulate binding of the first Fc region and/or the second Fc region to the Fc γ are L234A_L235A, F234A_L235A, S228P_F234A_L235A, S228P_L234A_L235A, V234A_G237A_P238S_H268A_V309L_A330S_P331S, V234A_G237A, H268Q_V309L_A330S_P331S, S267E_L328F, L234F_L235E_D265A, 15 L234A_L235A_G237A_P238S_H268A_A330S_P331S or S228P_F234A_L235A_G237A_P238S.

In some embodiments, the one or more substitutions that modulate binding of the first Fc region and/or the second Fc region to the FcRn are M428L_N434S, M252Y_S254T_T256E, T250Q_M428L, N434A and T307A_E380A_N434A, H435A, 20 P257I_N434H, D376V_N434H, M252Y_S254T_T256E_H433K_N434F, T308P_N434A or H435R.

In some embodiments, the one or more substitutions that modulate binding of the first Fc region and/or the second Fc region to protein A are Q311R, Q311K, T307P_L309Q, T307P_V309Q, T307P_L309Q_Q311R, T307P_V309Q_Q311R, H435R 25 or H435R_Y436F.

In some embodiments, the steps of producing the heterodimeric antibody are conducted under GMP-compliant conditions.

In some embodiments the steps of producing the heterodimeric antibody are conducted during manufacture of a drug substance comprising the heterodimeric antibody.

30 In some embodiments, the heterodimeric antibody is a bispecific antibody.

In some embodiments, the bispecific antibody binds CD3, BCMA or CD123.

In some embodiments, the steps of producing the heterodimeric antibody are conducted during manufacture of an innovator drug product.

In some embodiments, the steps of producing the heterodimeric antibody are 35 conducted during manufacture of a generic drug product.

Fab-arm exchange and Fc region mutations promoting heterodimerization

Heterodimeric antibodies may be produced utilizing Fab-arm exchange, wherein one heavy chain and its attached light chain (half-arm) of one parental homodimeric antibody is exchanged with one heavy chain and its attached light chain of another parental homodimeric antibody to form a heterodimeric antibody composed of two heavy chains and two attached light chains (van der Neut Kolfshoten *et al.*, (2007) *Science* 317:1554-1557).

The two homodimeric parental antibodies are engineered to have asymmetric mutations in their CH3 regions that favor Fab-arm exchange and heterodimeric antibody formation upon reduction and reformation of disulfide bridges in the hinge region of antibodies. An illustration of the Fab-arm exchange reaction is represented in **FIG. 1**. Upon introduction of a reducing agent into a mixture of two parental homodimeric antibodies the half-arms dissociate and the asymmetric CH3 mutations favor reformation of the heterodimeric antibodies. Subsequent reformation of the disulfide bridges between the half-arms stabilize the formed heterodimeric antibody (Gramer *et al.* (2013) *MAbs* 5:962-973).

In the methods of the invention any CH3 region mutation that promotes CH3 heterodimer formation may be used, such as those described herein. Several approaches are known for modifications in the CH3 region to promote heterodimerization. Typically, in all such approaches the first CH3 region and the second CH3 region are engineered in a complementary manner so that each CH3 region (or the heavy chain comprising it) can no longer homodimerize with itself but is forced to heterodimerize with the complementarily engineered other CH3 region (so that the first and second CH3 region heterodimerize and no homodimers between the two first or the two second CH3 regions are formed).

CH3 mutations that favor Fab-arm exchange include Duobody® mutations (Genmab), Knob-in-Hole mutations (Genentech), electrostatically-matched mutations (Chugai, Amgen, NovoNordisk, Oncomed), the Strand Exchange Engineered Domain body (SEEDbody) (EMD Serono), and other asymmetric mutations (e.g. Zymeworks).

Duobody® mutations (Genmab) are disclosed for example in US9,150,663 and US2014/0303356 and include mutations F405L/K409R, wild-type/F405L_R409K, T350I_K370T_F405L/K409R, K370W/K409R, D399AFGHILMNRSTVWY/K409R, T366ADEFHILMQVY/K409R, L368ADEGHNRSTVQ/K409AGRH, D399FHKRQ/K409AGRH, F405IKLSTVW/K409AGRH and Y407LWQ/K409AGRH.

Knob-in-hole mutations are disclosed for example in WO1996/027011 and include mutations on the interface of CH3 region in which an amino acid with a small side chain (hole) is introduced into the first CH3 region and an amino acid with a large side chain (knob) is introduced into the second CH3 region, resulting in preferential interaction between

the first CH3 region and the second CH3 region. Exemplary CH3 region mutations forming a knob and a hole are T366Y/F405A, T366W/F405W, F405W/Y407A, T394W/Y407T, T394S/Y407A, T366W/T394S, F405W/T394S and T366W/T366S_L368A_Y407V.

Heavy chain heterodimer formation may be promoted by using electrostatic interactions by substituting positively charged residues on the first CH3 region and negatively charged residues on the second CH3 region as described in US2010/0015133, US2009/0182127, US2010/028637 or US2011/0123532.

Other asymmetric mutations that can be used to promote heavy chain heterodimerization are L351Y_F405A_Y407V/T394W,

T366I_K392M_T394W/F405A_Y407V, T366L_K392M_T394W/F405A_Y407V, L351Y_Y407A/T366A_K409F, L351Y_Y407A/T366V_K409F, Y407A/T366A_K409F, or T350V_L351Y_F405A_Y407V/T350V_T366L_K392L_T394W as described in US2012/0149876 or US2013/0195849.

SEEDbody mutations involve substituting select IgG residues with IgA residues to promote heavy chain heterodimerization as described in US20070287170.

Other exemplary mutations that may be used are R409D_K370E/D399K_E357K, S354C_T366W/Y349C_T366S_L368A_Y407V, Y349C_T366W/S354C_T366S_L368A_Y407V, T366K/L351D, L351K/Y349E, L351K/Y349D, L351K/L368E, L351Y_Y407A/T366A_K409F, L351Y_Y407A/T366V_K409F, K392D/D399K, K392D/E356K, K253E_D282K_K322D/D239K_E240K_K292D, K392D_K409D/D356K_D399K as described in WO2007/147901, WO 2011/143545, WO2013157954, WO2013096291 and US2018/0118849.

These different approaches for Fab-arm exchange may be combined with various bispecific antibody format, such as those involving VH/VL engineering such as VH/VL domain swaps, CH1/CL domain swaps, utilizing common light chain as described in WO98050431 or utilizing tethered light chains including inside-out tethered light chains as describe in US9062120.

30 Further engineering of heterodimeric antibodies

Fc engineering

In addition to the CH3 region mutations that promote heterodimerization, antibodies used in the methods of the invention may comprise mutations in the Fc region that modulate antibody effector functions or half-life. Further, antibodies used in the

methods of the invention may comprise Fc mutations that modulate binding of the antibodies to protein A, hence facilitating purification of the antibodies.

Fc positions that may be mutated to modulate antibody half-life (e.g. binding to FcRn) include positions 250, 252, 253, 254, 256, 257, 307, 376, 380, 428, 434 and 435.

5 Exemplary mutations that may be made singularly or in combination are mutations T250Q, M252Y, I253A, S254T, T256E, P257I, T307A, D376V, E380A, M428L, H433K, N434S, N434A, N434H, N434F, H435A and H435R. Exemplary singular or combination mutations that may be made to increase the half-life of the antibodies are mutations M428L/N434S, M252Y/S254T/T256E, T250Q/M428L, N434A and

10 T307A/E380A/N434A. Exemplary singular or combination mutations that may be made to reduce the half-life of the antibodies are mutations H435A, P257I/N434H, D376V/N434H, M252Y/S254T/T256E/H433K/N434F, T308P/N434A and H435R.

Mutations may be introduced to the Fc region which reduce binding of the antibody to an activating Fc γ receptor (Fc γ R) and reduce Fc effector functions such as C1q binding, complement dependent cytotoxicity (CDC), antibody-dependent cell-mediated cytotoxicity (ADCC) or phagocytosis (ADCP).

Fc positions that may be mutated to reduce binding of the antibody to the activating Fc γ R and subsequently to reduce effector function include positions 214, 233, 234, 235, 236, 237, 238, 265, 267, 268, 270, 295, 297, 309, 327, 328, 329, 330, 331 and 365. Exemplary mutations that may be made singularly or in combination are mutations K214T, E233P, L234V, L234A, deletion of G236, V234A, F234A, L235A, G237A, P238A, P238S, D265A, S267E, H268A, H268Q, Q268A, N297A, A327Q, P329A, D270A, Q295A, V309L, A327S, L328F, A330S and P331S in IgG1, IgG2, IgG3 or IgG4. Exemplary combination mutations that result in antibodies with reduced ADCC are mutations L234A/L235A on IgG1, V234A/G237A/ P238S/H268A/V309L/A330S/P331S on IgG2, F234A/L235A on IgG4, S228P/F234A/ L235A on IgG4, N297A on all Ig isotypes, V234A/G237A on IgG2, K214T/E233P/ L234V/L235A/G236-deleted/A327G/P331A/D365E/L358M on IgG1, H268Q/V309L/A330S/P331S on IgG2, S267E/L328F on IgG1, L234F/L235E/D265A on IgG1, 25 L234A/L235A/G237A/P238S/H268A/A330S/P331S on IgG1, 30 S228P/F234A/L235A/G237A/P238S on IgG4, and S228P/F234A/L235A/G236-deleted/G237A/P238S on IgG4. Hybrid IgG2/4 Fc domains may also be used, such as Fc with residues 117-260 from IgG2 and residues 261-447 from IgG4.

Exemplary mutation that result in antibodies with reduced CDC is a K322A 35 mutation.

Well-known S228P mutation may be made in IgG4 antibodies to enhance IgG4 stability.

Mutations may be introduced to the Fc region which enhance binding of the antibody to an Fc γ receptor (Fc γ R) and enhance Fc effector functions such as C1q binding, complement dependent cytotoxicity (CDC), antibody-dependent cell-mediated cytotoxicity (ADCC) and/or phagocytosis (ADCP).

Fc positions that may be mutated to increase binding of the antibody to the activating Fc γ R and enhance antibody effector functions include positions 236, 239, 243, 256, 290, 292, 298, 300, 305, 312, 326, 330, 332, 333, 334, 345, 360, 339, 378, 396 or 430 (residue numbering according to the EU index). Exemplary mutations that may be made singularly or in combination are G236A, S239D, F243L, T256A, K290A, R292P, S298A, Y300L, V305L, K326A, A330K, I332E, E333A, K334A, A339T and P396L. Exemplary combinations that result in antibodies with increased ADCC or ADCP are a S239D/I332E, S298A/E333A/K334A, F243L/R292P/Y300L, F243L/R292P/Y300L/P396L, F243L/R292P/Y300L/V305I/P396L and G236A/S239D/I332E on IgG1.

Fc positions that may be mutated to enhance CDC of the antibody include positions 267, 268, 324, 326, 333, 345 and 430. Exemplary mutations that may be made singularly or in combination are S267E, F1268F, S324T, K326A, K326W, E333A, E345K, E345Q, E345R, E345Y, E430S, E430F and E430T. Exemplary combinations that result in antibodies with increased CDC are K326A/E333A, K326W/E333A, H268F/S324T, S267E/H268F, S267E/S324T and S267E/H268F/S324T on IgG1.

"Antibody-dependent cellular cytotoxicity", "antibody-dependent cell-mediated cytotoxicity" or "ADCC" is a mechanism for inducing cell death that depends upon the interaction of antibody-coated target cells with effector cells possessing lytic activity, such as natural killer cells (NK), monocytes, macrophages and neutrophils via Fc gamma receptors (Fc γ R) expressed on effector cells. For example, NK cells express Fc γ RIIIa, whereas monocytes express Fc γ RI, Fc γ RII and Fc γ RIIIa. ADCC activity of the antibodies may be assessed using an *in vitro* assay using cells expressing the protein the antibody binds to as target cells and NK cells as effector cells. Cytolysis may be detected by the release of label (e.g. radioactive substrates, fluorescent dyes or natural intracellular proteins) from the lysed cells. In an exemplary assay, target cells are used with a ratio of 1 target cell to 4 effector cells. Target cells are pre-labeled with BATDA and combined with effector cells and the test antibody. The samples are incubated for 2 hours and cell

lysis measured by measuring released BATDA into the supernatant. Data is normalized to maximal cytotoxicity with 0.67% Triton X-100 (Sigma Aldrich) and minimal control determined by spontaneous release of BATDA from target cells in the absence of any antibody.

5 **"Antibody-dependent cellular phagocytosis"** ("ADCP") refers to a mechanism of elimination of antibody-coated target cells by internalization by phagocytic cells, such as macrophages or dendritic cells. ADCP may be evaluated by using monocyte-derived macrophages as effector cells and cells that express the protein the antibody binds to as target cells also engineered to express GFP or another labeled molecule. In an exemplary
10 assay, effector:target cell ratio may be for example 4:1. Effector cells may be incubated with target cells for 4 hours with or without the antibody of the invention. After incubation, cells may be detached using accutase. Macrophages may be identified with anti-CD11b and anti-CD14 antibodies coupled to a fluorescent label, and percent phagocytosis may be determined based on % GFP fluorescence in the CD11⁺CD14⁺
15 macrophages using standard methods.

"Complement-dependent cytotoxicity", or "CDC", refers to a mechanism for inducing cell death in which the Fc effector domain of a target-bound antibody binds and activates complement component C1q which in turn activates the complement cascade leading to target cell death. Activation of complement may also result in deposition of
20 complement components on the target cell surface that facilitate CDC by binding complement receptors (e.g., CR3) on leukocytes. CDC of cells may be measured for example by plating Daudi cells at 1×10^5 cells/well (50 μ L/well) in RPMI-B (RPMI supplemented with 1% BSA), adding 50 μ L of test antibodies to the wells at final concentration between 0-100 μ g/mL, incubating the reaction for 15 min at room
25 temperature, adding 11 μ L of pooled human serum to the wells, and incubation the reaction for 45 min at 37° C. Percentage (%) lysed cells may be detected as % propidium iodide stained cells in FACS assay using standard methods.

 Binding of the antibody to Fc γ R or FcRn may be assessed on cells engineered to express each receptor using flow cytometry. In an exemplary binding assay, 2×10^5 cells
30 per well are seeded in 96-well plate and blocked in BSA Stain Buffer (BD Biosciences, San Jose, USA) for 30 min at 4°C. Cells are incubated with a test antibody on ice for 1.5 hour at 4°C. After being washed twice with BSA stain buffer, the cells are incubated with R-PE labeled anti-human IgG secondary antibody (Jackson ImmunoResearch Laboratories) for 45 min at 4°C. The cells are washed twice in stain buffer and then resuspended in 150
35 μ L of Stain Buffer containing 1:200 diluted DRAQ7 live/dead stain (Cell Signaling

Technology, Danvers, USA). PE and DRAQ7 signals of the stained cells are detected by Miltenyi MACSQuant flow cytometer (Miltenyi Biotec, Auburn, USA) using B2 and B4 channel, respectively. Live cells are gated on DRAQ7 exclusion and the geometric mean fluorescence signals are determined for at least 10,000 live events collected. FlowJo software (Tree Star) is used for analysis. Data is plotted as the logarithm of antibody concentration versus mean fluorescence signals. Nonlinear regression analysis is performed.

“**Enhance**” or “**enhanced**” refers to enhanced effector function (e.g. ADCC, CDC and/or ADCP) or enhanced binding to an Fc γ receptor (Fc γ R) or FcRn of the antibody of the invention having at least one mutation in the Fc region when compared to the parental antibody without the mutation. “Enhanced” may be an enhancement of about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100% or more, or a statistically significant enhancement.

“**Reduce**” or “**reduced**” refers to reduced effector function (e.g. ADCC, CDC and/or ADCP) or reduced binding to an Fc γ receptor (Fc γ R) or FcRn of the antibody of the invention having at least one mutation in the Fc region when compared to the parental antibody without the mutation. “Reduced” may be a reduction of about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100% or more, or a statistically significant reduction.

“**Modulate**” refers to either enhanced or reduced effector function (e.g. ADCC, CDC and/or ADCP) or enhanced or reduced binding to an Fc γ receptor (Fc γ R) or FcRn of the antibody of the invention having at least one mutation in the Fc region when compared to the parental antibody without the mutation.

Mutations may be introduced to the antibodies used in the methods of the invention that modulate binding of the antibody to protein A. Production and purification of full length bispecific therapeutic antibodies require efficient separation of the bispecific antibodies from excess parental and/or intermediate molecules. Bispecific antibodies having protein A binding-modulating Fc mutations in asymmetric manner (e.g. in one heavy chain only) can therefore be purified from the parental antibodies based on their differential elution profile from protein A affinity columns. Exemplary mutations that may be introduced to modulate protein A binding into the antibodies used in the methods of the invention are Q311R, Q311K, T307P_L309Q, T307P_V309Q, T307P_L309Q_Q311R, T307P_V309Q_Q311R, H435R or H435R_Y436F.

Glycoengineering

The ability of the antibodies used in the methods of the invention to induce ADCC may be enhanced by engineering their oligosaccharide component. Human IgG1 is N-glycosylated at Asn297 with the majority of the glycans in the well-known biantennary G0, G0F, G1, G1F, G2 or G2F forms. Antibodies produced by non-engineered CHO cells typically have a glycan fucose content of about at least 85%. The removal of the core fucose from the biantennary complex-type oligosaccharides attached to the Fc regions enhances the ADCC of antibodies via improved FcγRIIIa binding without altering antigen binding or CDC activity. Such mAbs may be achieved using different methods reported to lead to the successful expression of relatively high defucosylated antibodies bearing the biantennary complex-type of Fc oligosaccharides such as control of culture osmolality, application of a variant CHO line Lec13 as the host cell line, application of a variant CHO line EB66 as the host cell line, application of a rat hybridoma cell line YB2/0 as the host cell line, introduction of small interfering RNA specifically against the α 1,6-fucosyltransferase (*FUT8*) gene, or coexpression of β -1,4-N-acetylglucosaminyltransferase III and Golgi α -mannosidase II or a potent alpha-mannosidase I inhibitor, kifunensine.

The heterodimeric antibodies used in the methods of the invention may have a biantennary glycan structure with fucose content of about between 0% to about 15%, for example 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or 0%.

The heterodimeric antibodies used in the methods of the invention may have a biantennary glycan structure with fucose content of about 50%, 40%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or 0%.

“**Fucose content**” means the amount of the fucose monosaccharide within the sugar chain at Asn297. The relative amount of fucose is the percentage of fucose-containing structures related to all glycostructures. These may be characterized and quantified by multiple methods, for example: 1) using MALDI-TOF of N-glycosidase F treated sample (e.g. complex, hybrid and oligo- and high-mannose structures); 2) by enzymatic release of the Asn297 glycans with subsequent derivatization and detection/quantitation by HPLC (UPLC) with fluorescence detection and/or HPLC-MS (UPLC-MS); 3) intact protein analysis of the native or reduced mAb, with or without treatment of the Asn297 glycans with Endo S or other enzyme that cleaves between the first and the second GlcNAc monosaccharides, leaving the fucose attached to the first GlcNAc; 4) digestion of the mAb to constituent peptides by enzymatic digestion (e.g., trypsin or endopeptidase Lys-C), and subsequent separation, detection and quantitation by HPLC-

MS (UPLC-MS) or 5) separation of the mAb oligosaccharides from the mAb protein by specific enzymatic deglycosylation with PNGase F at Asn 297. The oligosaccharides released may be labeled with a fluorophore, separated and identified by various complementary techniques which allow fine characterization of the glycan structures by matrix-assisted laser desorption ionization (MALDI) mass spectrometry by comparison of the experimental masses with the theoretical masses, determination of the degree of sialylation by ion exchange HPLC (GlycoSep C), separation and quantification of the oligosaccharide forms according to hydrophilicity criteria by normal-phase HPLC (GlycoSep N), and separation and quantification of the oligosaccharides by high performance capillary electrophoresis-laser induced fluorescence (HPCE-LIF).

“**Low fucose**” or “**low fucose content**” refers to antibodies with fucose content of about 0% - 15%.

“**Normal fucose**” or “**normal fucose content**” refers to antibodies with fucose content of about over 50%, typically about over 60%, 70%, 80% or over 85%.

The heterodimeric antibodies used in the methods of the invention may be post-translationally modified by processes such as glycosylation, isomerization, deglycosylation or non-naturally occurring covalent modification such as the addition of polyethylene glycol moieties (pegylation) and lipidation. Such modifications may occur *in vivo* or *in vitro*. For example, the heterodimeric antibodies used in the methods of the invention may be conjugated to polyethylene glycol (PEGylated) to improve their pharmacokinetic profiles. Conjugation may be carried out by techniques known to those skilled in the art. Conjugation of therapeutic antibodies with PEG has been shown to enhance pharmacodynamics while not interfering with function.

25 **C-terminal lysine**

The heterodimeric antibodies used in the methods of the invention may have their C-terminal lysine (CTL) partially removed during manufacturing. During manufacturing, CTL removal may be controlled to less than the maximum level by control of concentration of extracellular Zn^{2+} , EDTA or EDTA – Fe^{3+} as described in U.S. Patent Publ. No. US20140273092. CTL content in antibodies can be measured using known methods.

The heterodimeric antibodies used in the methods of the invention may have a C-terminal lysine content of about 10% to about 90%, about 20% to about 80%, about 40% to about 70%, about 55% to about 70%, or about 60%.

The heterodimeric antibodies used in the methods of the invention may have a C-terminal lysine content of about 0%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 100%.

5 Antibody allotypes

The heterodimeric antibodies used in the methods of the invention may be of any allotype. It is expected that allotype has no influence on Fab-arm exchange.

Immunogenicity of therapeutic antibodies is associated with increased risk of infusion reactions and decreased duration of therapeutic response (Baert *et al.*, (2003) *N Engl J*

10 *Med* 348:602-08). The extent to which therapeutic antibodies induce an immune response in the host may be determined in part by the allotype of the antibody (Stickler *et al.*, (2011) *Genes and Immunity* 12:213-21). Antibody allotype is related to amino acid sequence variations at specific locations in the constant region sequences of the antibody.

Table 2 shows select IgG1, IgG2 and IgG4 allotypes.

15

Table 2.

Allotype	Amino acid residue at position of diversity (residue numbering: EU Index)							
	IgG2		IgG4		IgG1			
	189	282	309	422	214	356	358	431
G2m(n)	T	M						
G2m(n-)	P	V						
G2m(n)/(n-)	T	V						
nG4m(a)			L	R				
G1m(17)					K	E	M	A
G1m(17,1)					K	D	L	A

Target antigens

20 The methods of the invention may be used to generate heterodimeric antibodies with any specificity using Fab-arm exchange, as the reduction and reformation of the disulfide bonds during manufacturing is not expected to be influenced by the VH/VL regions of the antibodies, as is also demonstrated herein. The heterodimeric antibodies may bind one target antigen at two distinct epitopes, or may bind two or more target antigens, depending on the specificity of each VH/VL pair. When the heterodimeric

25 antibody binds two target antigens, the target antigens may be located on the same cell or on two different cells. The target antigen may be a tumor-associated antigen, an antigen playing a role in inflammation, regulation of T or B cell activity, or in general any target

whose biological activity is desired to be modulated, or the number of cells expressing the target are desired to be reduced.

Exemplary antigens the heterodimeric antibodies used in the methods of the invention may bind are one or more of ABCF1, ACVR1, ACVR1B, ACVR2, ACVR2B, 5 ACVRL1, ADORA2A, Aggrecan, AGR2, AICDA, AIF1, AIG1, AKAP1, AKAP2, albumin, AMH, AMHR2, ANGPT1, ANGPT2, ANGPTL3, ANGPTL4, ANPEP, APC, APOC1, APOE, AR, AZGP1 (zinc- α -glycoprotein), B7.1, B7.2, BAD, BAFF, BAG1, BAI1, BCL2, BCL6, BCMA, BDNF, BLNK, BLR1 (MDR15), BlyS, BMP1, BMP2, BMP3B (GDF10), BMP4, BMP6, BMP8, BMPR1A, BMPR1B, BMPR2, BPAG1 10 (plectin), BRCA1, BTLA, C19orf10 (IL27w), C3, C4A, C5, C5R1, CANT1, CASP1, CASP4, CAV1, CCBP2 (D6/JAB61), CCL1 (1-309), CCL11 (eotaxin), CCL13 (MCP-4), CCL15 (MIP-1d), CCL16 (HCC-4), CCL17 (TARC), CCL18 (PARC), CCL19 (MIP-3b), CCL2 (MCP-1), MCAF, CCL20 (MIP-3a), CCL21 (MIP-2), SLC, exodus-2, CCL22 (MDC/STC-1), CCL23 (MPIF-1), CCL24 (MPIF-2/eotaxin-2), CCL25 (TECK), CCL26 15 (eotaxin-3), CCL27 (CTACK/ILC), CCL28, CCL3 (MIP-1a), CCL4 (MIP-1b), CCL5 (RANTES), CCL7 (MCP-3), CCL8 (mcp-2), CCNA1, CCNA2, CCND1, CCNE1, CCNE2, CCR1 (CKR1/HM145), CCR2 (mcp-1RB/RA), CCR3 (CKR3/CMKBR3), CCR4, CCR5 (CMKBR5/ChemR13), CCR6 (CMKBR6/CKR-L3/STRL22/DRY6), CCR7 (CKR7/EBI1), CCR8 (CMKBR8/TER1/CKR-L1), CCR9 (GPR-9-6), CCRL1 (VSHK1), 20 CCRL2 (L-CCR), CD123, CD137, CD164, CD16a, CD16b, CD19, CD1C, CD20, CD200, CD-22, CD24, CD28, CD3, CD3 ϵ , CD30, CD32a, CD32b, CD33, CD37, CD38, CD39, CD4, CD40, CD40L, CD44, CD45RB, CD47, CD52, CD69, CD72, CD73, CD74, CD79A, CD79B, CD8, CD80, CD81, CD83, CD86, CD89, CD96, CDH1 (E-cadherin), CDH10, CDH12, CDH13, CDH18, CDH19, CDH20, CDH5, CDH7, CDH8, CDH9, 25 CDK2, CDK3, CDK4, CDK5, CDK6, CDK7, CDK9, CDKN1A (p21Wap1/Cip1), CDKN1B (p27Kip1), CDKN1C, CDKN2A (p16INK4a), CDKN2B, CDKN2C, CDKN3, CEBPB, CER1, CHGA, CHGB, Chitinase, CHST10, CKLFSF2, CKLFSF3, CKLFSF4, CKLFSF5, CKLFSF6, CKLFSF7, CKLFSF8, CLDN3, CLDN7 (claudin-7), CLN3, CLU (clusterin), CMKLR1, CMKOR1 (RDC1), CNR1, COL18A1, COL1A1, COL4A3, 30 COL6A1, CR2, CRP, CSF1 (M-CSF), CSF2 (GM-CSF), CSF3 (G-CSF), CTLA4, CTNNB1 (b-catenin), CTSB (cathepsin B), CX3CL1 (SCYD1), CX3CR1 (V28), CXCL1 (GRO1), CXCL10 (IP-10), CXCL11 (I-TAC/IP-9), CXCL12 (SDF1), CXCL13, CXCL14, CXCL16, CXCL2 (GRO2), CXCL3 (GRO3), CXCL5 (ENA-78/LIX), CXCL6 (GCP-2), CXCL9 (MIG), CXCR3 (GPR9/CKR-L2), CXCR4, CXCR6 (TYMSTR/STRL33/Bonzo), 35 CYB5, CYC1, CYSLTR1, DAB2IP, DES, DKFZp451J0118, DNAM-1, DNCL1, DPP4,

E2F1, ECGF1, EDG1, EFNA1, EFNA3, EFNB2, EGF, EGFR, ELAC2, ENG, ENO1, ENO2, ENO3, EPHB4, EPO, ERBB2 (Her-2), EREG, ERK8, ESR1, ESR2, F3 (TF), FADD, FasL, FASN, FCER1A, FCER2, FCGR3A, FGF, FGF1 (aFGF), FGF10, FGF11, FGF12, FGF12B, FGF13, FGF14, FGF16, FGF17, FGF18, FGF19, FGF2 (bFGF),

5 FGF20, FGF21, FGF22, FGF23, FGF3 (int-2), FGF4 (HST), FGF5, FGF6 (HST-2), FGF7 (KGF), FGF8, FGF9, FGFR, FGFR3, FIGF (VEGFD), FIL1 (EPSILON), FIL1 (ZETA), FLJ12584, FLJ25530, FLRT1 (fibronectin), FLT1, FOS, FOSL1 (FRA-1), FY (DARC), GABRP (GABAa), GAGEB1, GAGEC1, GALNAC4S-6ST, GATA3, GDF5, GF11, GGT1, GITR, GITRL, GM-CSF, GNAS1, GNRH1, GPR2 (CCR10), GPR31, GPR44,

10 GPR81 (FKSG80), GRCC10 (C10), GRP, GSN (Gelsolin), GSTP1, HAVCR2, HDAC4, HDAC5, HDAC7A, HDAC9, HGF, HIF1A, HIP1, histamine and histamine receptors, HLA, HLA-A, HLA-DRA, HM74, HMOX1, HUMCYT2A, HVEM, ICEBERG, ICOS, ICOSL, IDO, ID2, IFN-a, IFNA1, IFNA2, IFNA4, IFNA5, IFNA6, IFNA7, IFNB1, IFNgamma, IFNW1, IGBP1, IGF1, IGF1R, IGF2, IGFBP2, IGFBP3, IGFBP6, IL-1, IL10,

15 IL10RA, IL10RB, IL11, IL11RA, IL-12, IL12A, IL12B, IL12RB1, IL12RB2, IL13, IL13RA1, IL13RA2, IL14, IL15, IL15RA, IL16, IL17, IL17B, IL17C, IL17R, IL18, IL18BP, IL18R1, IL18RAP, IL19, IL1A, IL1B, IL1F10, IL1F5, IL1F6, IL1F7, IL1F8, IL1F9, IL1HY1, IL1R1, IL1R2, IL1RAP, IL1RAPL1, IL1RAPL2, IL1RL1, IL1RL2, IL1RN, IL2, IL20, IL20RA, IL21R, IL22, IL22R, IL22RA2, IL23, IL24, IL25, IL26,

20 IL27, IL28A, IL28B, IL29, IL2RA, IL2RB, IL2RG, IL3, IL30, IL3RA, IL4, IL4R, IL5, IL5RA, IL6, IL6R, IL6ST (glycoprotein 130), IL7, IL7R, IL8, IL8RA, IL8RB, IL8RB, IL9, IL9R, ILK, INHA, INHBA, INSL3, INSL4, insulin, insulin receptor, IRAK1, IRAK2, ITGA1, ITGA2, ITGA3, ITGA6 (a6 integrin), ITGAV, ITGB3, ITGB4 (b 4 integrin), JAG1, JAK1, JAK3, JUN, K6HF, KAI1, KDR, KITLG, KIR, KLF5 (GC Box BP), KLF6,

25 KLK10, KLK12, KLK13, KLK14, KLK15, KLK3, KLK4, KLK5, KLK6, KLK9, KRT1, KRT19 (Keratin 19), KRT2A, KRTHB6 (hair-specific type II keratin), LAG-3, LAMA5, LDL, LEP (leptin), LFA, Lingo-p75, Lingo-Troy, LPS, LTA (TNF-b), LTB, LTB4R (GPR16), LTB4R2, LTBR, MACMARCKS, MAG or Omgp, MAP2K7 (c-Jun), MDK, mesothelin, c-Met, MIB1, midkine, MIF, MIP-2, MKI67 (Ki-67), MMP2, MMP9,

30 MS4A1, MSMB, MT3 (metallothionein-III), MTSS1, MUC1 (mucin), c-MYC, MYD88, NCK2, neurocan, NFKB1, NFKB2, NGFB (NGF), NGFR, NgR-Lingo, NgR-Nogo66 (Nogo), NgR-p75, NgR-Troy, NKG2D, NKp46, NME1 (NM23A), NOX5, NPPB, NR0B1, NR0B2, NR1D1, NR1D2, NR1H2, NR1H3, NR1H4, NR1I2, NR1I3, NR2C1, NR2C2, NR2E1, NR2E3, NR2F1, NR2F2, NR2F6, NR3C1, NR3C2, NR4A1, NR4A2,

35 NR4A3, NR5A1, NR5A2, NR6A1, NRP1, NRP2, NT5E, NTN4, ODZ1, OPRD1, OX-40,

OX-40L, P2RX7, PAP, PART1, PATE, PAWR, PCA3, PCNA, PD-1, PD-L1, PDGFA, PDGFB, PECAM1, PF4 (CXCL4), PGF, PGR, phosphacan, PIAS2, PIK3CG, PLAUI (uPA), PLG, PLXDC1, PPBP (CXCL7), PPID, PR1, PRKCQ, PRKD1, PRL, PROC, PROK2, PSAP, PSCA, PSMA, PTAFR, PTEN, PTGS2 (COX-2), PTN, RAC2

5 (p21Rac2), RARB, RGS1, RGS13, RGS3, RNF110 (ZNF144), ROBO2, ROR1, SI00A2, SCGB1D2 (lipophilin B), SCGB2A1 (mammaglobin 2), SCGB2A2 (mammaglobin 1), SCYE1 (endothelial Monocyte-activating cytokine), SDF2, SERPINA1, SERPINA3, SERPINB5 (maspin), SERPINE1 (PAI-1), SERPINF1, SHBG, SLA2, SLC2A2, SLC33A1, SLC43A1, SLIT2, SPP1, SPRR1B (Spr1), ST6GAL1, STAB1, STAT6,

10 STEAP, STEAP2, TB4R2, TBX21, TCP10, TDGF1, TEK, TF (transferrin receptor), TGFA, TGFB1, TGFB111, TGFB2, TGFB3, TGFB1, TGFB1, TGFB2, TGFB3, TH1L, THBS1 (thrombospondin-1), THBS2, THBS4, THPO, TIE (Tie-1), TIGIT, TIM-3, TIMP3, tissue factor, TLR10, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR8, TLR9, TNF, TNF-a, TNFAIP2 (B94), TNFAIP3, TNFRSF11A, TNFRSF1A, TNFRSF1B,

15 TNFRSF21, TNFRSF5, TNFRSF6 (Fas), TNFRSF7, TNFRSF8, TNFRSF9, TNFRSF10 (TRAIL), TNFRSF11 (TRANSE), TNFRSF12 (APO3L), TNFRSF13 (April), TNFRSF13B, TNFRSF14 (HVEM-L), TNFRSF15 (VEGI), TNFRSF18, TNFRSF4 (OX40 ligand), TNFRSF5 (CD40 ligand), TNFRSF6 (FasL), TNFRSF7 (CD27 ligand), TNFRSF8 (CD30 ligand), TNFRSF9 (4-1BB ligand), TOLLIP, Toll-like receptors, TOP2A (topoisomerase Iia), TP53,

20 TPM1, TPM2, TRADD, TRAF1, TRAF2, TRAF3, TRAF4, TRAF5, TRAF6, TREM1, TREM2, TRPC6, TSLP, TWEAK, VEGF, VEGFB, VEGFC, versican, VHL C5, VISTA, VLA-4, XCL1 (lymphotactin), XCL2 (SCM-1b), XCR1 (GPR5/CCXCR1), YY1, and ZFPM2.

25 In some embodiments, the heterodimeric antibody used in the methods of the invention binds CD3.

In some embodiments, the heterodimeric antibody used in the methods of the invention binds CD3 and a tumor-associated antigen (TAA).

In some embodiments, the heterodimeric antibody used in the methods of the invention binds CD123 and CD3 and BCMA and CD3.

30 In some embodiments, the heterodimeric antibody used in the methods of the invention binds EGFR and c-Met.

In some embodiments, the heterodimeric antibody that binds EGFR and c-Met comprises a first heavy chain (HC1) of SEQ ID NO: 4, a first light chain (LC1) of SEQ ID NO: 5, a second heavy chain (HC2) of SEQ ID NO: 6 and a second light chain (LC2) of

35 SEQ ID NO: 7.

The invention also provides a bispecific antibody produced by the methods of the invention.

The invention also provides a bispecific antibody that binds EGFR and c-Met produced by the methods of the invention.

- 5 The invention also provides a bispecific antibody that binds EGFR and c-Met comprising a first heavy chain (HC1) of SEQ ID NO: 4, a first light chain (LC1) of SEQ ID NO: 5, a second heavy chain (HC2) of SEQ ID NO: 6 and a second light chain (LC2) of SEQ ID NO: 7 produced by the methods of the invention.

10 **Production of homodimeric and heterodimeric antibodies**

- The first homodimeric antibody and the second homodimeric antibody may be produced together or separately in a culture vessel such as bioreactor. The first homodimeric antibody and the second homodimeric antibody may be produced by expression in a host cell by co-expression or by using a separate host cell to produce each
15 homodimeric antibody. In the latter case the host cell may be of same or different origin.

- “Host cell” refers to a cell into which one or more vectors has been introduced which express at least one antibody heavy chain and one antibody light chain. “Host cell” refers not only to the particular subject cell but to the progeny of such a cell, and also to a stable cell line generated from the particular subject cell. Because certain modifications
20 may occur in succeeding generations due to either mutation or environmental influences, such progeny may not be identical to the parent cell but are still included within the scope of the term “host cell” as used herein.

- Such host cells may be eukaryotic cells, prokaryotic cells, plant cells or archeal cells. *Escherichia coli*, bacilli, such as *Bacillus subtilis*, and other enterobacteriaceae,
25 such as *Salmonella*, *Serratia*, and various *Pseudomonas* species are examples of prokaryotic host cells. Other microbes, such as yeast, are also useful for expression. *Saccharomyces* (for example, *S. cerevisiae*) and *Pichia* are examples of suitable yeast host cells. Exemplary eukaryotic cells may be of mammalian, insect, avian or other animal origins. Mammalian eukaryotic cells include immortalized cell lines such as hybridomas
30 or myeloma cell lines such as SP2/0 (American Type Culture Collection (ATCC), Manassas, VA, CRL-1581), NS0 (European Collection of Cell Cultures (ECACC), Salisbury, Wiltshire, UK, ECACC No. 85110503), FO (ATCC CRL-1646) and Ag653 (ATCC CRL-1580) murine cell lines. An exemplary human myeloma cell line is U266 (ATCC CRL-TIB-196). Other useful cell lines include those derived from Chinese

Hamster Ovary (CHO) cells such as CHOK1SV (Lonza Biologics, Walkersville, MD), Potelligent® CHOK2SV (Lonza), CHO-K1 (ATCC CRL-61) or DG44.

Exemplary vectors that may be used to express one or more antibody heavy chains and light chains include pWLneo, pSV2cat, pOG44, PXR1, pSG (Stratagene) pSVK3, pBPV, pMSG and pSVL (Pharmacia), pEE6.4 (Lonza) and pEE12.4 (Lonza) as well as
5 vectors used in Lonza Xceed system.

The first homodimeric antibody and the second homodimeric antibody may be produced by expressing them in a separate host cell. In this case, the first homodimeric antibody and the second homodimeric antibody may be purified prior to mixing them in
10 the presence of a reducing agent to initiate the Fab-arm exchange. Purification may be accomplished using known methods such as protein A chromatography. Alternatively, culture media containing the first homodimeric antibody and the second homodimeric antibody may be combined into which a reducing agent may be added to initiate the Fab-arm exchange. If the first homodimeric antibody and the second homodimeric antibody
15 are produced by co-expression, the antibodies may also be purified using protein A chromatography.

The Fab-arm exchange to generate heterodimeric antibodies is described herein. Process conditions such as concentration of the first homodimeric antibody and the second homodimeric antibody, reducing agent, concentration of the reducing agent, time of
20 reduction, concentration of dissolved oxygen, temperature in which the reaction is conducted and buffers used are optimized to produce heterodimeric antibodies with high yield and purity as described herein.

After Fab-arm exchange, the produced heterodimeric antibody may be further purified. Methods for purification include combination of protein A, protein G
25 chromatography, other ways of affinity chromatography, such as affinity chromatography based on antigen binding or binding to anti-idiotypic antibodies, thioaffinity, ionic exchange, hydrophobic interaction, hydroxyapatite chromatography and other mixed mode resins. Additional methods may use precipitation with for example salts or polyethylene glycol to obtain purified heterodimeric antibodies.

Equipment suitable for the process of the method of the present invention are known. Expression of homodimeric antibodies by a host cell may for example typically be performed in a reaction vessel, such as a bioreactor. The reducing and oxidizing steps may take place in the same bioreactor as expression of the first homodimeric antibody and/or the second homodimeric antibody or it may take place in separate reactor vessel.
35 The reaction vessel and supporting process piping may be disposable or re-usable and

made from standard materials (plastic, glass, stainless steel etc). The reaction vessel may be equipped with mixing, sparging, headspace gassing, temperature control and/or be monitored with probes for measurement of temperature, weight/volume, pH, dissolved oxygen (DO), and redox potential. All such techniques are common within standard unit
5 operations of a manufacturing plant and well known to a person skilled in the art.

While having described the invention in general terms, the embodiments of the invention will be further disclosed in the following examples that should not be construed as limiting the scope of the claims.

10

Example 1. General methods

cSDS

Capillary sodium dodecyl sulfate-polyacrylamide gel electrophoresis separates proteins on the basis of molecular weight. Analysis employed a commercial capillary
15 electrophoresis system with a fused silica capillary in a temperature-controlled cartridge. The test articles were mixed with internal standards and alkylating reagent, heated for a defined time and temperature to denature the protein. Samples were then injected electro-kinetically by voltage and then analyzed by application of a greater electric field for a predefined duration. Detection was accomplished by absorbance at 220 nm and purity determined by
20 calculating the corrected peak area of the main peak relative to all other peaks in the electrophoregram after the internal standard.

IEX-HPLC

Percent bispecific antibody in relation to residual parental homodimers antibodies
25 were determined by Ion Exchange High Performance Liquid Chromatography (IEX-HPLC). Ion exchange chromatography achieves separation by exploiting differences in the surface charge on the molecules. IEX-HPLC was performed using Thermo Scientific, Propac WCX-10, (4 mm ID x 150 mm, packed 10 um particles). Separation of species were resolved using a salt gradient and/or a pH gradient. As the gradient concentration increased, the overall
30 surface interaction of each antibody with the column is neutralized and elution is detected at 280 nm. The relative amount of each IgG was measured by comparing peak area count.

HIC-HPLC

Hydrophobic interaction chromatography method to separate proteins based on their
35 hydrophobic nature. Proteins are retained on the column under high salt conditions and are

eluted by decreasing the salt concentration. High salt buffer conditions expose the hydrophobic regions due to decreased solvation of the protein and promotes binding to the HIC stationary phase. Samples were injected onto a Tosoh TSKgel Butyl-NPR (4.6 mm x 10 cm, 2.5 µm non-porous resin based bead) column and eluted with a ammonium sulfate gradient. The absorbance monitored at 280 nm and peaks identified compared to reference standard injections.

RP-HPLC (to detect cystamine)

Reverse Phased High Performance Chromatography separates solute molecule in the mobile phase based the hydrophobic binding interactions to the stationary phase media. The differences in hydrophobicity of the 2-MEA and oxidized dimeric form cystamine were exploited to resolve and quantitate by RP-HPLC by comparing measured values to standard curves. A Phenomenex Luna 18 (4.6 mm x 150 mm, 3µm) column using a hexane sulfonate/acetonitrile mobile phase system was used to separate the two species with step gradients of increasing organic solvent. The species were detected by monitoring absorbance at 220 nm (2-MEA) and 255 nm (cystamine) and the peak areas compared to their relative standard curves.

General antibody production scheme for mono and bispecific (the one done at 200 ml scale) and scale

Each monospecific antibody was thawed from individual mammalian cell banks, expanded, and expressed separately in bioreactors. The production bioreactors were supplemented with enriched medias and harvested prior to significant cell death. The bioreactors were harvested by means of centrifugation and/or filtration, and the monospecific antibodies were captured separately by affinity protein A chromatography. The resulting captured monospecific antibodies were then stored frozen until combined in the FAE stage.

Example 2. Process improvements during manufacturing of bispecific antibodies using Fab-arm exchange

Bispecific antibodies were generated using Fab-arm exchange by introduction of asymmetrical mutations in the CH3 domain of the two homodimer parental antibodies followed by reduction and reformation of intermolecular disulfide bonds. The asymmetric CH3 domain mutations drive preferential formation of the heterodimeric bispecific antibody over the homodimeric parental antibodies.

To ensure process robustness during manufacturing, experiments were designed to gain understanding of possible thresholds and ranges for various parameters during manufacturing. Antibody disulfide bond reformation has been shown to be dependent on the presence of oxygen and free metals, as EDTA and oxygen-free conditions inhibited re-oxidation of cysteines in an antibody to disulfide bonds (US2014/0303356). Hence studies were initiated to investigate the necessity to control the amount of dissolved oxygen (DO₂) and metals during manufacturing of bispecific antibodies utilizing Fab-arm exchange.

FIG. 2 shows the manufacturing steps during reduction and reformation of the disulfide bonds during manufacturing of bispecific antibodies by Fab-arm exchange.

In theory, antibody disulfide bond reformation after reduction with a reducing agent such as 2-Mercaptoethylamine (2-MEA) may occur via various routes. For example, thiol-disulfide exchange may account for the reduction and reformation of the disulfide bonds. In order for the thiol-disulfide exchange to proceed, the reducing agent such as 2-MEA needs to be in its thiolate (RS⁻) form for the thiol-disulfide exchange to proceed. **FIG. 3** shows the molecular details of the thiol-disulfide exchange reaction. To reduce the disulfide bonds in the parental antibodies, the thiolate anion (circled in the **FIG. 3**) attacks the sulfur atom of the disulfide bond, displacing one of the sulfur atom and forming a new disulfide bond with the 2-MEA thiolate.

Using the Henderson-Hasselbalch equation shown below, the ratio of the 2-MEA thiol (SH) and thiolate anion (S⁻) is highly dependent on the pH and 2-MEA thiol pK_a.

$$\frac{[S^-]}{[SH]} = 10^{pH - pK_a}$$

Hence, the reduction of disulfide bonds and overall Fab-arm exchange becomes inhibited at low pH, where the protonated thiol form of 2-MEA is favored relative to its deprotonated thiolate anion form. At higher pH, the equilibrium shifts towards thiolates enabling the reduction of disulfide bonds and thiol-disulfide exchange.

A summary of the reactions that may take place during Fab-arm exchange in the presence of 2-MEA is presented in **FIG. 4**. During Fab-arm exchange, the removal of a reducing agent such as 2-MEA by buffer exchange (during UF/DF) enables the reformation of disulfide bonds in the heterodimer bispecific antibody, however multiple reaction pathways allow for the disulfide reformation process to occur. In the presence of oxygen, the oxygen may react with protonated thiolates facilitating reformation of the disulfide bonds in the heterodimeric bispecific antibody as illustrated by Reaction 2 in **FIG. 4**. The presence of oxygen may also deplete the residual 2-MEA thiols in solution

forming a cystamine dimer and water as shown by Reaction 4 in FIG. 4. Thus, sufficiently high concentration of 2-MEA should be used during the Fab-arm exchange to improve efficiency and yield of the bispecific antibody.

However, thiolate disulfide exchange and reformation of disulfide bonds may still be possible in the absence of oxygen. Under oxygen deprived conditions, as the concentration of 2-MEA decreases during the buffer exchange process, antibody thiol groups and 2-aminoethyl disulfide Fab intermediate may drive Reaction 1 of FIG. 4 in the reverse direction reforming the disulfide bonds in the bispecific heterodimeric antibodies with concomitant production of 2-MEA thiolate. This reaction would continue to be pushed into the reverse direction as more 2-MEA is removed during the buffer exchange process.

Example 3. Bispecific antibody A manufacturing in low DO₂ conditions during UF/DF

Experimental Method

In this experiment, ambient DO₂ levels were present during reduction and up to 23 hours post-2-MEA addition, after which a nitrogen overlay was included to minimize % DO₂ during UF/DF. The approach reflected conditions in which oxygen was present during the reduction phase and absent during reformation of the disulfide bonds.

Bispecific antibody A binds BCMA and CD3 and is an IgG4 isotype with S228P, F234A, L235A substitutions ("PAA substitutions") in both heavy chains, F at position 405 and R at position 409 in one heavy chain and L at position 405 and K at position 409 in the second heavy chain to drive heterodimer formation. The parental antibodies were named p1A-IgG4PAAF405R409 and p2A-IgG4PAAL405K409.

Parental antibodies p1A-IgG4PAAF405R409 and p2A-IgG4PAAL405K409 were harvested from cell culture bioreactors and purified by protein A affinity chromatography.

A solution was prepared using p1A-IgG4PAAF405R409 and p2A-IgG4PAAL405K409 at a steered molar ratio of 1:1.06 to intentionally limit one of the parental antibodies. The mixture was then adjusted to pH 7.3 and diluted to a total IgG concentration of 10.5 g/L using 101 mM sodium acetate, 105 mM Tris Base, pH 7.3. A 50 mM Sodium Acetate, 800 mM 2-MEA pH 5.0 stock solution was added to the parental mAb mixture. The final reduction buffer composition prior to UF/DF was about 100 mM sodium acetate, about 35 mM 2-MEA, about 30 mM NaCl, pH 7.3. The reduced parental mAb solution was incubated for 23 hrs at 24°C and subsequently transferred to the UF/DF retentate vessel. Prior to the start of ultrafiltration (UF) and diafiltration (DF), an overlay

of nitrogen was delivered to the retentate vessel via the headspace of the vessel to deplete DO₂ during reformation of the disulfide bridges and was maintained throughout the entirety of the UF and DF process. To prevent the addition of DO₂ through the diafiltration buffer, the diafiltration buffer (100 mM Tris-acetate 30 mM NaCl pH 7.5) was sparged with nitrogen throughout processing. In-line DO₂ sensors were used to verify that the retentate DO₂ had reached levels below 5 % DO₂ and the diafiltration buffer had reached levels below 1 % DO₂ prior to the start of the UF and DF steps.

Throughout the UF and DF process, the DO₂ levels were measured in the feed inlet, retentate, and permeate lines during the UF/DF and were all maintained at < 4 % DO₂. The diafiltration buffer was maintained at levels < 1 % DO₂. The UF/DF system was maintained at a 54 mL/min crossflow rate, targeting a transmembrane pressure of 14.5 psi for 11 diafiltration volumes (DV). All processes following the incubation were performed at room temperature (18 - 22°C). Upon completion of UF/DF, the recovered retentate was quenched with 1.0 M Acetic Acid and adjusted to pH 5.0 to minimize further disulfide bond formation and subsequently stored at 2 - 8°C. No additional buffer recovery flush was performed following the completion of UF/DF to prevent any possible oxidation of the material from the introduction of the additional diafiltration buffer.

The UF/DF setup was equipped to measure DO₂ at 3 locations while processing. In line monitoring of % DO₂ in the feed inlet, retentate, and permeate lines is shown in **FIG. 5**. Upon lowering the DO₂, the levels remained below 5% in the permeate and below 3% in the inlet/retentate. **Table 3** shows the summary of UF/DF process parameters and performance.

Table 3.

Final 2-MEA Concentration (mM)	35.0
TMP (psi)	14.5 - 16.0
Crossflow (mL/min)	53 – 55
Diafiltration Concentration (g/L)	25.0
Total Diafiltration Volumes (DVs)	11
Mean Permeate Flux (L/m ² /hr)	23.2
Load ratio (g/m ²)	320
Yield (%)	88

Table 4 shows the summary of offline %DO₂, pH and conductivity measurements during pre- and post 2-MEA addition. **FIG. 6** shows the percent (%) DO₂ during UF/DF measured in the retentate, permeate and inlet lines. The % DO₂ stabilized to 3.2% in the permeate, 1.5% in the retentate and 1.4% in the feed over time after nitrogen overlay in retentate in recirculation.

Non-reduced cSDS analysis was used to measure disulfide bond integrity of the formed heterodimeric bispecific antibody (measured as % purity of the heterodimeric antibody on non-reduced cSDS). The purity of antibody A was measured to be 97.59%.

This study demonstrated that under ambient % DO₂ (ranging from 89 - 13%) during reduction and a low % DO₂ environment (< 4% DO₂ during the UF/DF) the bispecific antibody A formed with high levels of disulfide bond formation (reflected by % purity on non-reduced cSDS). Previously, it was believed that DO₂ would be required to oxidize thiols into disulfide bonds to stabilize the bispecific antibody.

Table 4.

Step	Dissolved Oxygen (%)	pH	Conductivity (mS/cm)
Parental IgG mixture	88.1	4.99	3.37
pH Adjusted Parental Pool	87.5	7.35	4.43
Diluted pH Adjusted Parental Pool	88.9	7.33	4.39
Post 2-MEA Addition (T=0)	36.9	6.99	7.73
Post 2-MEA Addition (T=2hr)	18.7		
Post 2-MEA Addition (T=23hr)	13.2		

The levels of oxygen present during this study was calculated not to be sufficient to contribute to oxidation of the thiols during UF/DF. The calculation below demonstrated that DO₂ concentration above 30% in solution was needed for oxidation of thiols to occur using the reaction represented by Reaction 2 in **FIG. 4**.

For this study, the following assumptions were made to calculate theoretical percentage DO₂ saturated solution to oxidize all thiols in the amount of antibody present:

- IgG solution: 10 g/L

- moles S-S (disulfide bonds to be reduced and oxidized) / mole mAb
- $\frac{1}{2}$ mole O_2 / mole S-S If oxygen is required to generate disulfide bonds from thiols, refer to Reaction 2 of **FIG. 4**
- Oxygen solubility in air saturated water = 7 ppm (g O_2 / 106 g solution) = 100% saturation
- mAb MW: 150kDa

To calculate % dissolved DO_2 concentration to oxidize 10 mg/mL antibody solution when 4 disulfides/mole mAb is oxidized and O_2 required to oxidize all thiols:

1. (10 mg mAb / mL solution) * (mole mAb / 150000 g mAb) * (2 moles S-S / mole mAb) * ($\frac{1}{2}$ mole O_2 / mole S-S) * (1000 mL / 1 L) (1 g / 1000 mg) = 6.67×10^{-4} moles O_2 / L solution
2. (6.67×10^{-5} moles O_2 / L solution) * (31.998 g O_2 / 1 mole O_2) * (1 L / 1000 g) = 2.13×10^{-6} g O_2 / g solution
3. (2.13×10^{-6} g O_2 / g solution) * (10^6 g solution) = 2.13 g O_2 / 10^6 g solution = 2.13 ppm
4. Air saturated water about 7 g O_2 / 10^6 g solution = 7 ppm
5. 2.13 ppm / 7 ppm = 30 % O_2 saturated solution needed

2% DO_2 at operating temperatures was equivalent to 0.192 mg/mL or 6.00 μ M, 3% DO_2 was equivalent to 0.288 mg/L or 8.99 μ M, 4% DO_2 was equivalent to 0.384 mg/L or 11.29 μ M, 5% DO_2 was equivalent to 0.480 mg/mL or 14.99 μ M. 30% DO_2 was equivalent to 2.88 mg/mL or 90 μ M.

This indicates that oxygen independent chemical reaction pathways existed that allowed reformation of the disulfide bonds in the absence of oxygen during the UF/DF step. It is possible that as 2-MEA is removed from solution the reverse reaction of Reaction 1 of **FIG. 4** is driving the reformation of the disulfide bridges in the heterodimeric bispecific antibody.

Conclusion: Maintaining levels of DO_2 at < 4% during UF/DF did not significantly affect disulfide bond formation. At completion of the UF/DF, purity of the bispecific antibody was > 97%.

Example 4. Bispecific antibody B manufacturing in low DO_2 during UF/DF

In this experiment, ambient DO_2 levels were present during reduction and up to 23 hours post-2-MEA addition, after which a nitrogen overlay was included to minimize % DO_2 during UF/DF. The approach reflected conditions in which oxygen was present during the reduction phase and absent during reformation of the disulfide bonds.

Bispecific antibody B binds CD123 and CD3 and is an IgG4 isotype with S228P, F234A, L235A substitutions ("PAA substitutions") in both heavy chains, F at position 405 and R at position 409 in one heavy chain and L at position 405 and K at position 409 in the second heavy chain to drive heterodimer formation. The parental antibodies were named
5 p1B-IgG4PAAF405R409 and p2B-IgG4PAAL405K409.

The parental antibodies p1B-IgG4PAAF405R409 and p2B-IgG4PAAL405K409 were harvested from cell culture bioreactors and purified by protein A affinity chromatography.

A solution of p1B-IgG4PAAF405R409 and p2B-IgG4PAAL405K409 was
10 prepared and adjusted to pH 7.3 and diluted to a total IgG concentration of 10.5 g/L using 101 mM sodium acetate, 105 mM Tris Base, pH 7.3. A 50 mM Sodium Acetate, 800 mM 2-MEA pH 5.0 stock solution was added to the parental mixture. The final reduction buffer composition prior to UF/DF was about 100 mM sodium acetate, about 35 mM 2-MEA, about 30 mM NaCl, pH 7.3. The reduced parental solution was incubated for 23 hrs
15 at 24°C and subsequently transferred to the UF/DF retentate vessel. Prior to the start of ultrafiltration (UF) and diafiltration (DF), an overlay of nitrogen was delivered to the retentate vessel via the headspace of the vessel to deplete DO₂ during reformation of the disulfide bridges and was maintained throughout the entirety of the UF and DF process. To prevent the addition of DO₂ from the diafiltration buffer, the diafiltration buffer (100
20 mM Tris-acetate 30mM NaCl pH 7.5) was sparged with nitrogen throughout processing. In-line DO₂ sensors were used to verify that the retentate had reached levels below 5 % DO₂ and the diafiltration buffer had reached levels below 1 % prior to the start of the UF and DF steps.

Throughout the UF and DF process, the DO₂ levels were measured in the feed
25 inlet, retentate, and permeate lines during the UF/DF and were all maintained at < 4 % DO₂. The diafiltration buffer was maintained at levels < 1 % DO₂. The UF/DF system was maintained at a 42 mL/min crossflow rate, targeting a transmembrane pressure of 15.0 psi for 11 diafiltration volumes (DV). All processes following the incubation were performed at room temperature (18 - 22°C). Upon completion of UF/DF, a 3 mL sample
30 was taken, remained unadjusted at pH 7.5, and immediately stored at -70°C to prevent further oxidation. The remaining bulk of recovered retentate was quenched with 1.0 M Acetic Acid and adjusted to pH 5.0, to minimize further disulfide bond formation. The bulk material was subsequently stored at -70°C. No additional buffer recovery flush was performed following the completion of UF/DF to prevent any possible oxidation of the

material from the introduction of the additional diafiltration buffer, however yields were lower than typically observed.

The UF/DF setup was equipped to measure DO₂ at 3 locations while processing. In line monitoring of % DO₂ in the feed inlet, retentate, and permeate lines is shown in

5 **FIG. 5.** Upon lowering the DO₂, the levels remained below 5% in the permeate and below 3% in the inlet/retentate.

Table 5 shows the summary of UF/DF process parameters. **FIG. 7** shows the % DO₂ during UF/DF measured in the retentate, permeate and inlet lines. **Table 6** shows the summary of offline %DO₂, pH and conductivity measurements.

10

Table 5.

Final 2-MEA Concentration (mM)	35.0
TMP (psi)	15.0-16.5
Crossflow (mL/min)	42-45
Diafiltration Concentration (g/L)	25.0
Total Diafiltration Volumes (DVs)	11
Mean Permeate Flux (L/m ² /hr)	17.7
Load ratio (g/m ²)	320
Yield (%)	85

Table 6.

Step	%DO ₂	pH	Conductivity (mS/cm)	RmV
Parental Pool	82.9	4.61	2.57	189
pH Adjusted Parental Pool	80.3	7.26	4.55	106
Post 2-MEA Addition (T=0)	78.1	7.22	7.55	-314
Post 2-MEA Addition (T=23hr)	24.4			

15

Non-Reduced cSDS analysis was used to measure the disulfide bond integrity in the formed bispecific antibody for both the frozen samples at pH 7.5 and the bulk material at

pH 5.0. Purity of the pH 7.5 bispecific antibody preparation was measured to be 97.16 % and purity of the antibody preparation at pH 5.0 was measured to be 97.27 %.

This study demonstrated that under ambient % DO₂ (ranging from 83 - 24%) during the Reduction and a low % DO₂ environment (< 4 %DO₂) during the UF/DF the bispecific antibody B formed with high level of disulfide bond formation.

Both this study and the study described in Example 2 demonstrated that oxygen was not necessary during the UF/DF step for reformation of the disulfide bridges during Fab-arm exchange.

Conclusion: Maintaining levels of DO₂ at < 4% during UF/DF step of manufacturing the bispecific antibody B using Fab-arm exchange did not significantly affect disulfide bond formation. At completion of the UF/DF, Non-Reduced cSDS demonstrated % Purity > 97%.

Example 5. Bispecific antibody A manufacturing in low DO₂ in the presence of EDTA during UF/DF

During manufacturing of bispecific antibodies there are opportunities for free metal ions to be introduced to the manufacturing process. During the preparation of buffers via raw materials and the leaching from metallic components, these free metal ions may become involved in the redox reactions of the Fab-arm exchange. To investigate the possible effects of trace metals in the Fab-arm exchange, addition of EDTA during the Fab-arm exchange was studied. EDTA addition sequestered possible free metal ions in the solution which may catalyze the oxidative reaction for the reformation of disulfide bonds.

A solution was prepared using p1A-IgG4PAAF405R409 and p2A-IgG4PAAL405K409 at a molar ratio of 1.05:1.00. The mixture was then adjusted to pH 7.3 and diluted to a total IgG concentration of 10.5 g/L using 101 mM sodium acetate, 105 mM Tris Base, pH 7.3. A 50 mM Sodium Acetate, 800 mM 2-MEA pH 5.0 stock solution was added to the parental mAb mixture. The final reduction buffer composition prior to UF/DF was about 100 mM sodium acetate, about 35 mM 2-MEA, about 30 mM NaCl, pH 7.3. The reduced parental solution was incubated for 23.5 hrs at 24°C. At the completion of the incubation, 500 mM EDTA, pH 8.0 stock solution was added to the reduced parental mixture to a target of 2 mM EDTA to chelate free metal ions that would be present in solution prior to the start of UF/DF. EDTA was also added to the diafiltration buffer to target of 100 mM Tris-Acetate, 30 mM NaCl, 2 mM EDTA, pH 7.5. This was done to ensure that additional free metal ions could not be introduced during the UF/DF buffer exchange. EDTA was not added during reduction step to allow any possible

oxidation of disulfides and cystamine formation which may occur spontaneously during the step.

The reduced parental solution containing EDTA was subsequently transferred to the UF/DF retentate vessel and prior to the start of the UF, an overlay of nitrogen was delivered to the retentate vessel via the headspace of the vessel to deplete DO₂ from the retentate and was maintained throughout the entirety of the UF and DF process. To prevent the addition of DO₂ through the diafiltration buffer, the diafiltration buffer was sparged with nitrogen throughout processing. In-line DO₂ sensors verified that the retentate DO₂ levels reached levels below 2 % DO₂ and the diafiltration buffer reached DO₂ levels below 1 % DO₂ prior to the start of the UF and DF steps.

Throughout the UF and DF process, DO₂ levels were measured in the feed inlet, retentate, and permeate lines during the UF/DF and were all maintained < 4 % DO₂. The diafiltration buffer was maintained at levels < 1 % DO₂. The UF/DF system was maintained at a 58 mL/min crossflow rate, targeting a transmembrane pressure of 14.5 psi for 11 diafiltration volumes (DV). All processes following the incubation were performed at room temperature (18 - 22°C). Upon completion of the UF/DF, a 9 mL sample was taken, remained unadjusted at pH 7.5, and immediately stored at -70°C to prevent further oxidation. The remaining bulk of the recovered retentate was quenched with 1.0 M Acetic Acid and adjusted to pH 5.0, to minimize further disulfide bond formation. The bulk bispecific antibody sample at pH 5.0 was subsequently stored at -70°C. No buffer recovery flush was performed following the completion of UF/DF to prevent possible oxidation of the material from the introduction of the additional diafiltration buffer.

Table 7 summarizes the UF/DF process parameters and performance.

Table 7.

Final 2-MEA Concentration (mM)	35.0
TMP (psi)	14.8 - 15.5
Crossflow (mL/min)	58
Diafiltration Concentration (g/L)	25.0
Total Diafiltration Volumes (DVs)	11
Mean Permeate Flux (L/m ² /hr)	30.9
Load ratio (g/m ²)	318
Yield (%)	N/A

Non-Reduced cSDS analysis was used to measure the disulfide bond integrity in the formed bispecific antibody for both the frozen sample at pH 7.5 and the bulk material at pH 5.0. Purity of the pH 7.5 bispecific antibody preparation was 97.44% and purity of the bispecific antibody preparation at pH 5.0 was measured to be 97.39%.

5 This study demonstrated that under low % DO₂ environment during UF/DF and minimal available free metal ions the bispecific antibody A formed with high levels of disulfide bond formation. The study demonstrated that free metal ions for catalyzing the oxidative reaction may not be needed during UF/DF of Fab-arm exchange.

10 Conclusion: Experimental results demonstrated that the addition of EDTA to the reduced parental mixture prior to UF/DF in Fab-arm exchange did not affect disulfide bond formation. At completion of UF/DF, Non-Reduced cSDS resulted in % purity > 97%.

Example 6. Comparison of Fab-arm exchange in ambient and low DO₂ conditions during reduction and UF/DF

15 This study was designed to evaluate the hypothesis that some DO₂ mediated oxidation could be occurring during the 2-MEA Reduction.

For the study, Fab-arm exchange was conducted using parental antibodies p1B-IgG4PAAF405R409 and p2B-IgG4PAAL405K409. A mixture of p1B-IgG4PAAF405R409 and p2B-IgG4PAAL405K409 in solution was prepared and adjusted to pH 7.3 and diluted to a total IgG concentration of 10.5 g/L using 101 mM sodium acetate, 105 mM Tris Base, pH 7.3. The mixture containing the parental antibodies was divided into two separate vessels. In one vessel, an overlay of nitrogen was delivered to the vessel via the headspace to deplete DO₂ from the parental mixture until a % DO₂ < 5% was achieved. The other parental mixture vessel remained under ambient air conditions as a control for the reduction. A 50 mM Sodium Acetate, 800 mM 2-MEA pH 5.0 stock solution was added to each of the parental mixtures. The final reduction buffer composition prior to UF/DF was about 100 mM sodium acetate, about 35 mM 2-MEA, about 30 mM NaCl, pH 7.3. Both reduced parental solutions were incubated for 24 hrs at 24°C and the % DO₂ was measured for both vessels, the results of which are shown in **FIG. 8**. The low % DO₂ reduction vessel was maintained under an overlay of nitrogen throughout the entirety of the incubation and remained at < 2 % DO₂ for the 24 hrs. There was an initial drop in % DO₂ following 2-MEA spike to approximately 40% DO₂ in the vessel which remained under ambient air, after which % DO₂ gradually increased linearly back to > 90 % DO₂. The summary of UF/DF process parameters and performance is shown in **Table 8**.

20

25

30

35

Table 8.

Final 2-MEA Concentration (mM)	35.0
TMP (psi)	14.5 - 16.0
Crossflow (mL/min)	55
Diafiltration Concentration (g/L)	25.0
Total Diafiltration Volumes (DVs)	12
Mean Permeate Flux (L/m ² /hr)	19.5
Load ratio (g/m ²)	251
Yield (%)	N/A

During reduction, samples were obtained from both vessels through a sampling port on the vessel. The port was flushed prior to each sample draw and samples were immediately quenched by an addition of 10 % v/v of cystamine RP-HPLC mobile phase, 20 mM Hexane Sulfonate, pH 2.0, buffer and each sample was verified to have been reduced to a pH < 5 prior to analysis. All timepoints were assayed for disulfide bond integrity by Non-Reduced cSDS and residual 2-MEA/Cystamine concentration by RP-HPLC.

Additional sample preparation was required for residual 2-MEA/Cystamine analysis involving removal of the antibody by 5 kDa molecular weight retention filter. Non-Reduced cSDS % purity and residual 2-MEA/Cystamine results are reported in **Table 9**.

At the completion of the 24 hr reduction incubation, the low % DO₂ vessel was transferred to the UF/DF system. An overlay of nitrogen was continued to be delivered to the retentate vessel via the headspace of the vessel to continue depleting DO₂ from the retentate and was maintained throughout the entirety of the UF and DF process. The diafiltration buffer was also sparged with nitrogen throughout processing, to prevent the addition of DO₂ through the buffer. In-line DO₂ sensors were used to verify that the retentate had reached levels below 5 % DO₂ and the diafiltration buffer had reached levels below 1 % DO₂ prior to the start of the UF and DF steps.

Throughout the UF and DF process, the DO₂ levels were measured in the feed inlet, retentate, and permeate lines during the UF/DF and were all maintained at < 4 % DO₂. The diafiltration buffer was maintained at levels < 1 % DO₂. The UF/DF system was maintained at a 55 mL/min crossflow rate, targeting a transmembrane pressure of 14.5 psi for 12 diafiltration volumes (DV) using a 100 mM Tris-Acetate, 30 mM NaCl, pH 7.5 buffer. All processes following the incubation were performed at room temperature (18 -

22°C). Upon completion of UF/DF, the recovered retentate was quenched with 1.0 M acetic acid and adjusted to pH 5.0, to minimize further disulfide bond formation. The bulk material was subsequently stored at -70°C. No additional buffer recovery flush was performed following the completion of UF/DF to prevent any possible oxidation of the material from the introduction of the additional diafiltration buf.

Disulfide integrity by Non-Reduced cSDS and residual 2-MEA/Cysteamine by RP-HPLC was measured at various timepoints throughout reduction and UF/DF for both the low DO₂ % vessel and the control vessel with ambient DO₂.

Summary of data is shown in **Table 9**. Only the Fab-arm exchange reactions under oxygen deprived conditions was continued through UF/DF in this experiment. Residual 2-MEA and cystamine was measured only through the reduction for both Fab-arm exchange conditions, however during the UF/DF for the oxygen deprived conditions, samples were obtained to measure the possible reformation of disulfide bonds by Non-Reduced cSDS.

15 **Table 9.**

Timepoint	Low DO ₂				Ambient DO ₂			
	cSDS (%)	2-MEA (mM)	Cystamine (mM)	DO ₂ (%)	cSDS (%)	2-MEA (mM)	Cystamine (mM)	DO ₂ %
t=0*	98.34	N/a	N/a	2.5	98.08	N/a	N/a	90
t=3min	23.5	33.3	0.5	4	44.85	35.3	0.4	85
t=10min	10.9	31.8	0.5	1.4	24.19	35.5	0.6	77
t=30min	4.57	31.1	0.5	1	14.22	34.8	N/a	63
t=1hr	2.86	30.6	0.5	<1	19.13	33.7	1.4	38
t=2hr	2.74	31.8	0.6	<1	41.16	31.1	2.7	37
t=4hr	3.37	31.9	0.8	<1	53.07	27.6	4.0	50
t=6hr	4.17	32.2	0.8	<1	56.02	25.1	5.1	63
t=8hr	4.92	32.2	0.8	<1	59.66	23.3	6.0	70
t=10hr	5.85	33.1	0.9	<1	61.32	21.9	6.9	75
t=12hr	6.96	32.2	0.9	<1	65.15	20.0	7.5	80
t=22hr	14.64	32.4	1.2	<1	62.82	15.4	10.5	85
t=24hr	14.74	32.1	1.6	<1	71.00	14.3	10.4	92
DV0	27.60							
DV2	75.69							
DV4	85.64							

DV6	82.54
DV8	90.38
DV12	89.90
FINAL	90.59

*Initial t=0 sample was taken immediately following the 2-MEA spike of the parental mixture. Sample time points were taken to determine the rate of reduction and reformation of disulfide bonds during both the reduction and UF/DF

Conclusion: This data supports the hypothesis that oxygen independent pathways mediate disulfide bond formation in the absence of oxygen during the reduction and UF/DF.

5 When Fab-arm exchange was performed under oxygen-deprived conditions, minimal cystamine formation and disulfide bond reformation was observed over the course of the reduction. By the completion of the 24 hr incubation in the presence of 2-MEA, only 1.6 mM Cystamine and 14.74 % intact antibody was observed. Nevertheless, once the antibody sample was processed through UF/DF, increasing levels of disulfide
10 bond reformation was evident as 2-MEA was removed. At the end of DV (at DV12), the formed bispecific antibody preparation at pH 5.0 was measured to be % Purity of 90.59 %.

 In ambient DO₂, disulfide bond reduction occurred quickly, 30 min after start of the reduction about 86% of the parental antibodies were reduced. Reformation of the disulfide bonds was observed during the remainder of the reduction, even in the continued
15 presence of residual 2-MEA in solution. At the completion of reduction phase, 71% of the antibodies had intact disulfide bonds. Cystamine dimer formation as well as increasing levels of formation of the bispecific antibody during the reduction incubation indicated that oxygen was being consumed by Reactions 4 and 2, respectively, illustrated in **FIG. 4**. This was supported by the RP-HPLC data which showed significant depletion of 2-MEA
20 and conversion to the dimer compound cystamine (Reaction 4 in **FIG. 4**).

 This study demonstrated that DO₂ during the reduction stage catalyzed the formation of IgG HC-HC and LC-HC interchain disulfide bonds, but was not essential for the formation of an intact bispecific antibody as the disulfide bonds were reformed even under oxygen-deprived conditions. Removal of 2-MEA, even under the absence of
25 oxygen, during diafiltration accounted for the disulfide bond reformation, driven by the reverse reaction in Reaction 1 in **FIG. 4**.

Example 7. Generation of bispecific EGFR/c-Met antibody using Fab-arm exchange in ambient DO₂ utilizing high concentrations of parental antibodies

This study was designed to evaluate the hypothesis that higher homodimeric mixture protein concentrations and varying ratio of reducing agent to protein mass produces

5 oxidized heterodimer antibody.

A bispecific EGFR/cMet antibody was used in the study. The EGFR/cMet antibody comprises a first heavy chain (HC1) of SEQ ID NO: 4, a first light chain (LC) of **SEQ ID NO: 5**, a second HC (HC2) of **SEQ ID NO: 6** and a second LC (LC2) of **SEQ ID NO: 7**.

For the study, Fab-arm exchange was conducted using parental EGFR and c-Met
10 antibodies (both were IgG1 isotypes). A mixture of the parental antibodies filtered neutralized protein A eluates in solution was prepared and adjusted to pH 7.9 and diluted to a total IgG concentration of 10 - 35 g/L using 209.5 mM sodium acetate, 300 mM NaCl, pH 7.9. A 50 mM Sodium Acetate, 800 mM 2-MEA pH 5.0 stock solution was added to each of the parental mixtures to a target reductant concentration of 35, 50, or 100 mM.
15 Both reduced parental solutions were incubated for 3 hrs controlled at either 21.5°C, 24°C, or left uncontrolled at ambient room temperature. After 3 hours the incubated mixtures were ultrafiltered and diafiltered against 10 mM Tris, 7.8 mM Acetic acid, pH 7.5. The summary of operating conditions and analytical results are shown in **Table 10**. This study demonstrated oxidized heterodimer under these conditions for a temperature range of 21.5
20 - 24°C, reductant concentrations of 35 – 100 mM, and homodimeric mixture protein concentration of 10-35 g/L. Bispecific antibody integrity and purity was assessed by both NR-cSDS and IHC-HPLC using methods described in Example 1.

Table 10.

Run number	Reduction Incubation Time (hours)	Reduction Incubation Temp	[2MEA] (mM)	[Protein] g/L	NR-cSDS (%)	HIC-HPLC (%)
1	24	24	35	10.5	96.01	
2	24	24	35	10.5	94.63	
3	3	24	50	10.5	96.74	92.2
4	3	24	50	10.5	96.95	90.1
5	3	24	100	10.5	95.85	93.2
6	3	21.5	100	30	96.46	92.2
7	3	21.5	100	35	96.33	92.1

8	3	24	100	30	96.77	
9	3	RT	100	35	95.96	
10	3	RT	100	35	96.86	
11	8	RT	100	35	96.66	
12	3	RT	100	35	97.18	93.2
13	3	RT	100	35	97.01	93.2

SEQ ID NO: 4 (HC1)

QVQLVESGGGVVQPGRSLRLSCAASGFTFSTYGMHWVRQAPGKGLEWVAVIWD
 DGSYKYYGDSVKGRFTISRDNSKNTLYLQMNSLRAEDTAVYYCARDGITMVRGV
 5 MKDYFDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPV
 TVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVPSSSLGTQTYICNVNHKPSNTK
 VDKRVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDV
 SHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEY
 KCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPS
 10 DIAVEWESNGQPENNYKTTPPVLDSDGSFLLYSKLTVDKSRWQQGNVFSCSVMH
 EALHNHYTQKSLSLSPGK

SEQ ID NO: 5 (LC1)

AIQLTQSPSSLSASVGDRVTITCRASQDISSALVWYQQKPGKAPKLLIYDASSLESG
 15 VPSRFGSGESGTDFTLTISSLQPEDFATYYCQQFNISYPLTFGGGTKVEIKRTVAAPS
 VFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSK
 DSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

SEQ ID NO: 6 (HC2)

20 QVQLVQSGAEVKKPGASVKVSCETSGYTFTSYGISWVRQAPGHGLEWMGWISAY
 NGYTNYAQLQGRVTMTTDTSTSTAYMELRSLRSDDTAVYYCARDLRGTNYFD
 YWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNS
 GALTSGVHTFPAVLQSSGLYSLSSVVPSSSLGTQTYICNVNHKPSNTKVDKRV
 PKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEV
 25 KFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSN
 KALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEW
 ESNGQPENNYKTTPPVLDSDGSFLLYSRLTVDKSRWQQGNVFSCSVMHEALHNH
 YTQKSLSLSPGK

SEQ ID NO: 7 (LC2)

DIQMTQSPSSVSASVGDRVITICRASQGISNWLAWFQHKPGKAPKLLIYAASSLLS
GVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQANSFPITFGQGRLEIKRTVAAPS
VFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSK
5 DSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

We claim

- 1) A method of producing a heterodimeric antibody, comprising
 - a) providing a first homodimeric antibody comprising a first Fc region of an immunoglobulin comprising a first CH3 region and a second homodimeric antibody comprising a second Fc region of an immunoglobulin comprising a second CH3 region, wherein the amino acid sequences of the first CH3 region and the second CH3 regions are different and are such that a heterodimeric interaction between the first CH3 region and the second CH3 region is stronger than a homodimeric interaction between the first CH3 region or a homodimeric interaction between the second CH3 region;
 - b) combining the first homodimeric antibody and the second homodimeric antibody into a mixture; and
 - c) incubating the mixture in the presence of a reducing agent and ambient dissolved oxygen (DO₂) to produce the heterodimeric antibody, wherein the method lacks one or more steps of measuring a percentage (%) DO₂ in the mixture, controlling the % DO₂ in the mixture, or adding oxygen into the mixture during producing the heterodimeric antibody.
- 2) The method of claim 1, wherein the % DO₂ in the mixture is about 30% or more at step 1b).
- 3) The method of claim 1 or 2, wherein the % DO₂ is between about 10% and about 90% during producing the heterodimeric antibody.
- 4) The method of any one of claims 1-3, wherein the first homodimeric antibody and the second homodimeric antibody are combined into the mixture in step 1b) at a molar ratio of between about 1:1 to about 1:2.
- 5) The method of any one of claims 1-3, wherein the first homodimeric antibody and the second homodimeric antibody are combined into the mixture in step 1b) at the molar ratio of about 1.05:1.
- 6) The method of any one of claims 1-5, wherein total concentration of immunoglobulin in the mixture is between about 8 g/L and about 13 g/L, such as about 10.5 g/L.
- 7) The method of any one of claims 1-6, wherein the mixture is incubated in the presence of the reducing agent for about 10 minutes or longer.
- 8) The method of claim 7, wherein the mixture is incubated in the presence of the reducing agent from about 10 minutes to about 24 hours.

- 9) The method of any one of claims 1-8, wherein the reducing agent is 2-mercaptoethylamine (2-MEA), a chemical derivative of 2-MEA, L-cysteine or D-cysteine.
- 10) The method of claim 9, wherein concentration of 2-MEA in the mixture is between about 20 mM and 40 mM, such as about 35 mM.
- 11) The method of any one of claims 1-10, wherein the mixture comprises a buffer.
- 12) The method of claim 11, wherein the buffer comprises a sodium acetate buffer.
- 13) The method of claim 12, wherein the buffer further comprises NaCl.
- 14) The method of claim 13, wherein the buffer comprises about 100 mM sodium acetate and about 30 mM NaCl.
- 15) The method of claim 14, wherein the pH of the buffer is about 7.3.
- 16) The method of any one of claims 1-15, comprising a step of removing the reducing agent from the mixture.
- 17) The method of claim 16, wherein the reducing agent is removed by filtration.
- 18) The method of claim 17, wherein filtration is diafiltration.
- 19) The method of any one of claims 1-18, wherein the first homodimeric antibody and the second homodimeric antibody are an IgG1, IgG2 or IgG4 isotype.
- 20) The method of claim 19, wherein the first CH3 domain and the second CH3 domain comprise following mutations when compared to the wild-type IgG1 of SEQ ID NO: 1, wild-type IgG2 of SEQ ID NO: 2 or wild-type IgG4 of SEQ ID NO: 3:
F405L/K409R, wild-type/F405L_R409K, T350I_K370T_F405L/K409R,
K370W/K409R, D399AFGHILMNRSTVWY/K409R,
T366ADEFHILMQVY/K409R, L368ADEGHNRSTVQ/K409AGRH,
D399FHKRQ/K409AGRH, F405IKLSTVW/K409AGRH, Y407LWQ/K409AGRH,
T366Y/F405A, T366W/F405W, F405W/Y407A, T394W/Y407T, T394S/Y407A,
T366W/T394S, F405W/T394S, T366W/T366S_L368A_Y407V,
L351Y_F405A_Y407V/T394W, T366I_K392M_T394W/F405A_Y407V,
T366L_K392M_T394W/F405A_Y407V, L351Y_Y407A/T366A_K409F,
L351Y_Y407A/T366V_K409F, Y407A/T366A_K409F,
T350V_L351Y_F405A_Y407V/T350V_T366L_K392L_T394W, K409D/D399K,
K409E/D399R, K409D_K360D/D399K_E356K, K409D_K360D/D399E_E356K,
K409D_K370D/D399K_E357K, K409D_K370D/D399E_E357K,
K409D_K392D/D399K_E356K_E357K or K409D_K392D/D399E_E356K_E357K,
- 21) The method of claim 20, wherein the first Fc region and/or the second Fc region comprise one or more mutations when compared to the wild-type IgG1 of SEQ ID

- NO: 1, wild-type IgG2 of SEQ ID NO: 2 or wild-type IgG4 of SEQ ID NO: 3 that modulate binding of the first Fc region and/or the second Fc region to an Fc γ receptor (Fc γ R), an FcRn, or to protein A.
- 22) The method of claim 21, wherein the Fc γ R is Fc γ RI, Fc γ RIIa, Fc γ RIIb or Fc γ RIII.
- 23) The method of claim 21, wherein the one or more substitutions that modulate binding of the first Fc region and/or the second Fc region to the Fc γ are L234A_L235A, F234A_L235A, S228P_F234A_L235A, S228P_L234A_L235A, V234A_G237A_P238S_H268A_V309L_A330S_P331S, V234A_G237A, H268Q_V309L_A330S_P331S, S267E_L328F, L234F_L235E_D265A, L234A_L235A_G237A_P238S_H268A_A330S_P331S or S228P_F234A_L235A_G237A_P238S.
- 24) The method of claim 21, wherein the one or more substitutions that modulate binding of the first Fc region and/or the second Fc region to the FcRn are M428L_N434S, M252Y_S254T_T256E, T250Q_M428L, N434A and T307A_E380A_N434A, H435A, P257I_N434H, D376V_N434H, M252Y_S254T_T256E_H433K_N434F, T308P_N434A or H435R.
- 25) The method of claim 21, wherein the one or more substitutions that modulate binding of the first Fc region and/or the second Fc region to protein A are Q311R, Q311K, T307P_L309Q, T307P_V309Q, T307P_L309Q_Q311R, T307P_V309Q_Q311R, H435R or H435R_Y436F.
- 26) The method of any one of claims 1-25, wherein the steps of producing the heterodimeric antibody are conducted under GMP-compliant conditions.
- 27) The method of claim 26, wherein the steps of producing the heterodimeric antibody are conducted during manufacture of a drug substance comprising the heterodimeric antibody.
- 28) The method of any one of claims 1-27, wherein the heterodimeric antibody is a bispecific antibody.
- 29) The method of claim 28, wherein the bispecific antibody binds CD3, CD123, BCMA, EGFR or c-Met, or any combination thereof.
- 30) The method of any one of claims 1-29, wherein a mass ratio of the reducing agent to total immunoglobulin in the mixture is between about 1.0 and about 5.0.
- 31) The method of claim 30, wherein the mass ratio is between about 1.4 and about 3.8.
- 32) The method of claim 30, wherein the mass ratio is between about 3.3 and about 4.4.
- 33) A method of producing a heterodimeric antibody, comprising

- a) providing a first homodimeric antibody comprising a first Fc region of an immunoglobulin comprising a first CH3 region and a second homodimeric antibody comprising a second Fc region of an immunoglobulin comprising a second CH3 region, wherein the amino acid sequences of the first CH3 region and the second CH3 regions are different and are such that the heterodimeric interaction between the first CH3 region and the second CH3 region is stronger than a homodimeric interaction between the first CH3 region or a homodimeric interaction between the second CH3 region;
 - b) combining the first homodimeric antibody and the second homodimeric antibody into a mixture;
 - c) incubating the mixture in the presence of a reducing agent; and
 - d) removing the reducing agent to produce the heterodimeric antibody, wherein the percentage (%) of dissolved oxygen (DO₂) is controlled to be about 30% or lower in step c), step d) or both in step c) and step d).
- 34) The method of claim 33, wherein the % DO₂ the mixture is about 25% or less, 20% or less, about 15% or less, about 10% or less, about 9% or less, about 8% or less, about 7% or less, about 6% or less, about 5% or less, about 4% or less, about 3% or less, about 2% or less or about 1% or less.
- 35) The method of claim 34, wherein % DO₂ is controlled by an overaly of nitrogen.
- 36) The method of claim 34 or 35, wherein the first homodimeric antibody and the second homodimeric antibody are combined into the mixture in step b) at a molar ratio of between about 1:1 to about 1:2.
- 37) The method of claim 34 or 35, wherein the first homodimeric antibody and the second homodimeric antibody are combined into the mixture in step b) at the molar ratio of about 1.05:1.
- 38) The method of any one of claims 33-37, wherein total concentration of immunoglobulin in the mixture is about 10.5 g/L.
- 39) The method of any one of claims 33-38, wherein the mixture is incubated in the presence of the reducing agent for about 10 minutes or longer.
- 40) The method of any one of claims 33-39, wherein the mixture is incubated in the presence of the reducing agent from about 10 minutes to about 24 hours.
- 41) The method of any one of claims 33-40, wherein the reducing agent is 2-mercaptoethylamine (2-MEA), a chemical derivative of 2-MEA, L-cysteine or D-cysteine.

- 42) The method of claim 41, wherein concentration of 2-MEA in the mixture is about 35 mM.
- 43) The method of any one of claims 33-42, wherein the mixture comprises a buffer.
- 44) The method of claim 43, wherein the buffer comprises a sodium acetate buffer.
- 45) The method of claim 44, wherein the buffer further comprises NaCl.
- 46) The method of claim 45, wherein the buffer comprises about 100 mM sodium acetate and about 30 mM NaCl.
- 47) The method of claim 46, wherein the pH of the buffer is about 7.3.
- 48) The method of any one of claims 33-47, comprising a step of removing the reducing agent from the mixture.
- 49) The method of claim 48, wherein the reducing agent is removed by filtration.
- 50) The method of claim 49, wherein filtration is diafiltration.
- 51) The method of any one of claims 33-50, wherein the first homodimeric antibody and the second homodimeric antibody are an IgG1, IgG2 or IgG4 isotype.
- 52) The method of claim 51, wherein the first CH3 domain and the second CH3 domain comprise following mutations when compared to the wild-type IgG1 of SEQ ID NO: 1, wild-type IgG2 of SEQ ID NO: 2 or wild-type IgG4 of SEQ ID NO: 3:
F405L/K409R, wild-type/F405L_R409K, T350I_K370T_F405L/K409R,
K370W/K409R, D399AFGHILMNRS TVWY/K409R,
T366ADEF GHILMQVY/K409R, L368ADEF GHNRS TVQ/K409AGRH,
D399FHKRQ/K409AGRH, F405IKLSTVW/K409AGRH, Y407LWQ/K409AGRH,
T366Y/F405A, T366W/F405W, F405W/Y407A, T394W/Y407T, T394S/Y407A,
T366W/T394S, F405W/T394S, T366W/T366S_L368A_Y407V,
L351Y_F405A_Y407V/T394W, T366I_K392M_T394W/F405A_Y407V,
T366L_K392M_T394W/F405A_Y407V, L351Y_Y407A/T366A_K409F,
L351Y_Y407A/T366V_K409F, Y407A/T366A_K409F,
T350V_L351Y_F405A_Y407V/T350V_T366L_K392L_T394W, K409D/D399K,
K409E/D399R, K409D_K360D/D399K_E356K, K409D_K360D/D399E_E356K,
K409D_K370D/D399K_E357K, K409D_K370D/D399E_E357K,
K409D_K392D/D399K_E356K_E357K or K409D_K392D/D399E_E356K_E357K,
- 53) The method of claim 52, wherein the first Fc region and/or the second Fc region comprise one or more mutations when compared to the wild-type IgG1 of SEQ ID NO: 1, wild-type IgG2 of SEQ ID NO: 2 or wild-type IgG4 of SEQ ID NO: 3 that modulate binding of the first Fc region and/or the second Fc region to an Fc γ receptor (Fc γ R), an FcRn, or to protein A.

- 54) The method of claim 53, wherein the Fc γ R is Fc γ RI, Fc γ RIIa, Fc γ RIIb or Fc γ RIII.
- 55) The method of claim 53, wherein the one or more substitutions that modulate binding of the first Fc region and/or the second Fc region to the Fc γ are L234A_L235A, F234A_L235A, S228P_F234A_L235A, S228P_L234A_L235A, V234A_G237A_P238S_H268A_V309L_A330S_P331S, V234A_G237A, H268Q_V309L_A330S_P331S, S267E_L328F, L234F_L235E_D265A, L234A_L235A_G237A_P238S_H268A_A330S_P331S or S228P_F234A_L235A_G237A_P238S.
- 56) The method of claim 53, wherein the one or more substitutions that modulate binding of the first Fc region and/or the second Fc region to the FcRn are M428L_N434S, M252Y_S254T_T256E, T250Q_M428L, N434A and T307A_E380A_N434A, H435A, P257I_N434H, D376V_N434H, M252Y_S254T_T256E_H433K_N434F, T308P_N434A or H435R.
- 57) The method of claim 53, wherein the one or more substitutions that modulate binding of the first Fc region and/or the second Fc region to protein A are Q311R, Q311K, T307P_L309Q, T307P_V309Q, T307P_L309Q_Q311R, T307P_V309Q_Q311R, H435R or H435R_Y436F.
- 58) The method of any one of claims 33-57, wherein the steps of producing the heterodimeric antibody are conducted under GMP-compliant conditions.
- 59) The method of any one of claims 33-57, wherein the steps of producing the heterodimeric antibody are conducted during manufacture of a drug substance comprising the heterodimeric antibody.
- 60) The method of any one of claims 33-59, wherein the heterodimeric antibody is a bispecific antibody.
- 61) The method of claim 60, wherein the bispecific antibody binds CD3, CD123, BCMA, EGFR or c-Met, or any combination thereof.
- 62) The method of any one of claims 33-61, wherein a mass ratio of the reducing agent to total immunoglobulin in the mixture is between about 1.0 and about 5.0.
- 63) The method of claim 62, wherein the mass ratio is between about 1.4 and about 3.8.
- 64) The method of claim 62, wherein the mass ratio is between about 3.3 and about 4.4.
- 65) An isolated bispecific antibody produced by a method of claim 1.
- 66) The bispecific antibody of claim 65, wherein the bispecific antibody binds EGFR and c-Met.

- 67) The bispecific antibody of claim 65 or 66, comprising a first heavy chain (HC1) of SEQ ID NO: 4, a first light chain (LC1) of SEQ ID NO: 5, a second heavy chain (HC2) of SEQ ID NO: 6 and a second light chain (LC2) of SEQ ID NO: 7.
- 68) The bispecific antibody of claim 65, wherein the bispecific antibody binds CD3.
- 69) An isolated bispecific antibody produced by a method of claim 33.
- 70) The bispecific antibody of claim 69, wherein the bispecific antibody binds EGFR and c-Met.
- 71) The bispecific antibody of claim 69 or 70, comprising a first heavy chain (HC1) of SEQ ID NO: 4, a first light chain (LC1) of SEQ ID NO: 5, a second heavy chain (HC2) of SEQ ID NO: 6 and a second light chain (LC2) of SEQ ID NO: 7.
- 72) The bispecific antibody of claim 69, wherein the bispecific antibody binds CD3.