


PROCESS FOR PREPARING AMINOBENZOFURAN DERIVATIVES

The present invention relates generally to the preparation of aminobenzofuran derivatives.

- 5 More specifically, the invention relates to a process for preparing 5-aminobenzofuran derivatives of general formula:



and also to the addition salts thereof, in which R_1 represents hydrogen or an alkyl group and R_2 represents an alkyl or dialkylaminoalkyl group.

- 15 In formula I above:

- R_1 represents, in particular, a linear or branched C_1 - C_8 alkyl group, especially a linear or branched C_1 - C_4 alkyl group such as methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl or tert-butyl,
- R_2 represents, in particular, a linear or branched C_1 - C_8 alkyl group, especially a linear or branched C_1 - C_4 alkyl group such as methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl or tert-butyl or alternatively a dialkylaminoalkyl group in which each linear or branched alkyl group is of C_1 - C_8 and especially in which each linear or branched alkyl group is of C_1 - C_4 , such as methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl or tert-butyl.

25 In particular, R_1 represents n-butyl and R_2 represents 3-(di-n-butylamino)propyl.

- 30 Among the compounds of formula I above, 2-n-butyl{4-[3-(di-n-butylamino)propoxy]benzoyl}-5-aminobenzofuran described in patent EP 0 471 609 proves to be particularly useful as an intermediate product for the final preparation of aminoalkoxybenzoylbenzofuran derivatives, in particular for the preparation of 2-n-butyl-3-{4-[3-(di-n-butylamino)propoxy]benzoyl}-5-methanesulfonamidobenzofuran, commonly known as dronedarone, and also

the pharmaceutically acceptable salts thereof. This methanesulfonamidobenzofuran derivative was described in the patent mentioned previously, along with its therapeutic applications, especially in the cardiovascular field where it proved to be particularly advantageous, for example as an antiarrhythmic agent.

In addition, that same patent EP 0 471 609 disclosed a process for synthesizing 2-n-butyl-3-{4-[3-(di-n-butylamino)propoxy]benzoyl}-5-aminobenzofuran using 2-n-butyl-3-{4-[3-(di-n-butylamino)propoxy]benzoyl}-5-nitrobenzofuran, which is reduced, under pressure, with hydrogen in the presence of platinum oxide as catalyst, to give the desired compound.


However, this process is not without drawbacks inherent especially to the type of reaction used, namely hydrogenation under pressure, which entails an industrial risk.

The search for a process for preparing 2-n-butyl-3-{4-[3-(di-n-butylamino)propoxy]benzoyl}-5-aminobenzofuran that is capable of overcoming this drawback and disadvantage thus remains of fundamental interest.

Patent application US 4 666 931 and the articles Chemistry of Heterocyclic Compounds (1975), vol. 11, pp. 1361-1364, Journal of Organometallic Chemistry (1998), vol. 560, pp. 163-167 and Bulletin of the Chemical Society of Japan (1967), vol. 40, pp. 1724-1726 describe N-phenylalkylamide and 5-N-alkylamidobenzofuran derivatives, but do not in any way suggest their use in the preparation of compounds of formula (I) according to the invention.

Furthermore, WO 03/040 120 describes a process for preparing dronedarone, in six steps, using a Friedel-Crafts reaction. However, that document does not describe the preparation of dronedarone comprising a simple step of treating a 5-N-alkylamidobenzofuran derivative of formula (IIa) according to the invention with a strong acid.

According to the invention, the 5-aminobenzofuran derivatives of formula I may be prepared by treating, with a strong acid such as a hydracid, for example hydrofluoric acid, a 5-N-alkylamidobenzofuran derivative of general formula:



5

II

in which R_1 and R_2 have the same meaning as previously and R_3 represents a linear or branched C_1 - C_4 alkyl group, for example methyl, to form an acid-addition salt of the compound of formula I, which salt is itself treated, if necessary, with a basic agent such as an alkali metal hydroxide, to give this compound of formula I in free base form.

Hereinabove and hereinbelow, unless specifically mentioned:

The term "strong acid" means any chemical compound which has a very high capacity for introducing protons into the reaction medium and which is characterized, in aqueous solution, by a pK_a of less than or equal to 1. The term "strong acid" especially means any hydracid such as chosen from hydrochloric acid, hydrobromic acid and hydrofluoric acid.


The term "basic agent" means any chemical compound which has a high affinity for protons H^+ and which is characterized, in aqueous solution, by a pK_a of greater than 7. The term "basic agent" especially means bases of any type such as chosen from organic bases, weak bases and strong bases, chosen especially from tertiary amines, alkali metal carbonates and alkali metal hydroxides.


The term "strong base" means any chemical compound which has a very high affinity for protons H^+ and which is characterized, in aqueous solution, by a pK_a of greater than 14. The term "strong base" especially means any alkali metal hydroxide, such as chosen from sodium hydroxide and potassium hydroxide.

The acid treatment may be undertaken in a polar solvent such as an alcohol, for example ethanol, using an acid generally in excess, for example from 1 to 6 equivalents of this acid per equivalent of compound of formula II.

Moreover, the acid-addition salt of the compound of formula I may be treated with a basic agent, after isolation from the reaction medium in which it is formed or, on the contrary, *in situ*, i.e. within this same reaction medium.

The starting compounds of formula II may be prepared according to the following reaction scheme:





i.e. starting with a compound of formula III, in which R_3 has the same meaning as previously, which compound is heated in the presence of trifluoroacetic acid in a suitable solvent, usually an amine, for example hexamethylenetetramine, to form an N-phenylalkylamide derivative of formula IV in which R_3 has the same meaning as previously.

This amide of formula IV is then reacted with an ester of formula V in which R_1 has the same meaning as previously, R_4 represents a linear or branched C_1-C_4 alkyl group and Hal represents a halogen, for example bromine, in the presence of a basic agent, generally a weak base such as an alkali metal carbonate, and by heating in a polar solvent so as to form an ester of formula VI in which R_1 , R_3 and R_4 have the same meaning as previously.

The ester of formula VI is then saponified in the presence of a strong base, generally an alkali metal hydroxide, the reaction usually taking place at

room temperature and in a suitable solvent, for example an ether, to give a salt of a carboxylic acid derivative, which is treated with a strong acid, such as a hydracid, for example hydrofluoric acid, which gives a carboxylic acid derivative of formula VII in which R_1 and R_3 have the same meaning as previously.


The compound of formula VII thus produced is then cyclized to a benzofuran derivative of formula VIII in which R_1 and R_3 have the same meaning as previously, this being done in the presence of an organic base, generally a tertiary amine, and of a benzenesulfonyl chloride. The reaction is usually performed by heating in a suitable solvent, in general an aprotic solvent such as an aromatic hydrocarbon or an ether.

The benzofuran derivative of formula VIII thus obtained is then coupled with an acyl halogen of formula IX in which R_5 represents a linear or branched C_1 - C_4 alkyl group, for example methyl, and Hal has the same meaning as previously, for example chlorine, this being done in the presence of a Lewis acid, for example ferric chloride, and in a non-polar solvent, for example a halogenated compound. The reaction medium thus obtained is then hydrolyzed in the presence of a strong acid, for example a hydracid, to produce a ketone of formula X in which R_1 , R_3 and R_5 have the same meaning as previously.

This ketone of formula X is then dealkylated by heating in the presence of aluminum chloride and in a non-polar solvent, usually a halogenated solvent such as chlorobenzene, to form a 4-hydroxyphenyl derivative of formula XI in which R_1 and R_3 have the same meaning as previously.


Thereafter, the compound of formula XI is reacted with an alkyl halide of formula XII in which R_2 has the same meaning as previously and Hal has the same meaning as previously, for example chlorine, the reaction taking place in the presence of a basic agent such as an alkali metal carbonate and by heating usually in a polar solvent such as a ketone, to give the desired compound of formula II.

Another subject of the present invention relates to the N-phenylalkylamide derivatives of general formula:



XIII


in which R_3 has the same meaning as previously and Y represents hydrogen
 5 or a group of general formula:



XIV

in which $\text{R}_{1'}$ and R_6 represent, independently of each other, hydrogen
 10 or a linear or branched $\text{C}_1\text{-C}_4$ alkyl group,
 to the exclusion of the compounds for which $\text{R}_3 = \text{Me}$, Et or isopropyl
 and $\text{Y} = \text{H}$.

Among the N-phenylalkylamide derivatives of general formula XIII,
 mention may be made of those in which R_3 represents a linear or branched
 15 $\text{C}_1\text{-C}_4$ alkyl group and Y represents a group of general formula XIV:



XIV

in which $\text{R}_{1'}$ and R_6 represent, independently of each other, hydrogen
 20 or a linear or branched $\text{C}_1\text{-C}_4$ alkyl group.


Among the compounds of formula XIII, mention may be made of the
 following subgroup in which Y represents hydrogen or the group XIV in which
 R_1 represents n-butyl, and also of the compounds of formula XIII in which R_3
 represents methyl.

Moreover, among the compounds of formula XIII, mention may further be made of the subgroups in which R_3 represents methyl, Y represents hydrogen or the group XIV in which R_1 represents n-butyl and R_6 represents hydrogen or methyl.

5 Consequently, the subgroups of the N-phenylalkylamide derivatives of formula XIII may be those in which:

- a) R_3 represents methyl and Y represents the group XIV in which R_1' represents n-butyl and R_6 represents methyl,
 - b) R_3 represents methyl and Y represents the group XIV in which R_1' represents n-butyl and R_6 represents hydrogen.
- 10


In addition, another subject of the present invention concerns the 5-N-alkylamidobenzofuran derivatives of general formula:



XV

15


in which R_1' and R_3 have the same meaning as previously and Z represents hydrogen or a group of general formula:



XVI

20 in which R_2' represents hydrogen, a linear or branched C_1 - C_4 alkyl group or a dialkylaminoalkyl group in which each linear or branched alkyl group is of C_1 - C_4 ,
to the exclusion of the compounds for which R_3 = Me, Z = H and R_1' = H, Me, Et or n-Bu.

25 Among the 5-N-alkylamidobenzofuran derivatives of general formula XV, mention may be made of those in which R_1' represents hydrogen or a linear or branched C_1 - C_4 alkyl group, R_3 represents a linear or branched C_1 - C_4 alkyl group and Z represents a group of general formula XVI:



XVI

in which R_2' represents hydrogen, a linear or branched C_1 - C_4 alkyl group or a dialkylaminoalkyl group in which each linear or branched alkyl group is of C_1 - C_4 .

Among the compounds of formula XV above, mention may be made of those in which R_1' represents n-butyl, and also of the compounds of formula XV in which R_3 represents methyl.

Moreover, among the compounds of formula XV, mention may be made of the following subgroups in which Z represents hydrogen or a group XVI in which R_2' represents hydrogen, methyl or 3-(di-n-butylamino)propyl.


Moreover, among the compounds of formula XV, mention may also be made of those in which R_1' represents n-butyl, R_3 represents methyl and R_2' represents hydrogen, methyl or 3-(di-n-butylamino)propyl.

Consequently, subgroups of the 5-N-alkylamidobenzofuran derivatives of formula XV may be those in which:

- a) R_1' represents n-butyl, R_3 represents methyl and Z represents the group XVI in which R_2' represents hydrogen,
- b) R_1' represents n-butyl, R_3 represents methyl and Z represents the group XVI in which R_2' represents methyl,
- c) R_1' represents n-butyl, R_3 represents methyl and Z represents the group XVI in which R_2' represents 3-(di-n-butylamino)propyl.


An additional subject of the present invention concerns the use of compounds of formula II for the preparation of dronedarone and of pharmaceutically acceptable salts thereof.

Thus, according to another characteristic of the present invention, dronedarone of formula:




and pharmaceutically acceptable salts thereof, may be obtained:

a) by treating a 5-N-alkylamidobenzofuran derivative of general formula:



in which R₃ has the same meaning as previously, for example methyl, with a strong acid such as a hydracid, for example hydrofluoric acid, to form an addition salt (also known as an "acid-addition salt") of 2-n-butyl-3-{4-[3-(di-n-butylamino)propoxy]benzoyl}-5-aminobenzofuran of formula:




which salt is itself treated with a basic agent such as an alkali metal hydroxide, to give this compound of formula I_a in free base form,


20 b) by coupling this 5-aminobenzofuran derivative of formula I_a with methanesulfonyl chloride to form the dronedarone compound in free base form, which may be reacted, if necessary, with an acid to produce a pharmaceutically acceptable salt of this compound.

25 The 5-N-alkylamidobenzofuran derivatives of formula II_a may themselves be obtained by performing a sequence of steps according to which:

a) an N-phenylalkylamide derivative of general formula:




30 in which R₃ has the same meaning as previously, is reacted with an ester of general formula:



Va


in which R_4 and Hal have the same meaning as previously, and in doing so in the presence of a basic agent and by heating in a polar solvent to form an ester of general formula:



VIa

in which R_3 and R_4 have the same meaning as previously,


- b) the ester of formula VI_a is saponified in the presence of a strong base, the reaction proceeding at room temperature and in a suitable solvent, to give a carboxylic acid derivative, which is treated with a strong acid to form the carboxylic acid derivative of general formula:



VIIa

in which R_3 has the same meaning as previously,


- c) the carboxylic acid derivative of formula VII_a is cyclized by heating in an aprotic solvent and in the presence of an organic base and a benzenesulfonyl halide to form a benzofuran derivative of general formula:



VIIIa

in which R_3 has the same meaning as previously,


- d) the benzofuran derivative of formula VIII_a is coupled with an acyl halide of general formula:



5

IX

in which R_5 and Hal have the same meaning as previously, this being done in the presence of a Lewis acid and in a non-polar solvent, the reaction medium thus formed then being hydrolyzed in the presence of a strong acid to form a ketone of general formula:




10

Xa

in which R_3 and R_5 have the same meaning as previously,

- e) the ketone of formula X_a is dealkylated in the presence of aluminum chloride and in a non-polar solvent to form a 4-hydroxyphenyl derivative of general formula:




15

XIa

in which R_3 has the same meaning as previously,

- f) the 4-hydroxyphenyl derivative of formula XI_a is reacted with an alkyl halide of general formula:




in which Hal has the same meaning as previously, the reaction taking place in the presence of a basic agent and by heating in a polar solvent, to form the desired compound of formula II_a.

10 The nonlimiting example that follows illustrates the invention.

PREPARATIONS

A. N-(3-Formyl-4-hydroxyphenyl)acetamide (compound IV: R₃= CH₃)

25 g of N-(4-hydroxyphenyl)acetamide (0.165 mol; 1 equivalent) and 92.7 g of trifluoroacetic acid (0.661 mol; 4 equivalents) are placed in a 500 ml reactor. The mixture is stirred, and 92.7 g of hexamethylenetetramine (0.661 mol; 4 equivalents) are then added portionwise. The reaction is exothermic. A temperature of 70°C is maintained over 18 hours, while monitoring the reaction progress by thin-layer chromatography (eluent: 8/2 toluene/methanol). The reaction medium is allowed to warm to room temperature. It becomes thick, and is diluted by addition of 200 ml of water to facilitate the stirring. 100 ml of ethyl acetate are then added. Stirring is continued and the phases are allowed to separate by settling. 300 ml of water and 200 ml of ethyl acetate are added, the mixture is stirred and the phases are then allowed to separate by settling. The aqueous phase is removed first, then the organic phase. The organic phase is then washed twice with 150 ml of water in total to collect, after phase separation, a new organic phase and a new aqueous phase. This new organic phase is then concentrated on a rotary evaporator (T = 45°C; 50 mmHg). 14 g of the desired compound are thus obtained in the form of crystals recovered from an orange-red solution.

¹H NMR (DMSO-d₆)

Chemical shifts $\delta \pm 0.01$ ppm	Multiplicity	Integration	Coupling constants ($ J \pm 0.5$ Hz)	Assignment
2.00	Singlet	3	-	CH ₃
6.94	Doublet	1	$^3J_{H6-H5}=9.0$ Hz	H(6)
7.64	Doubled doublet	1	$^3J_{H5-H6}=9.0$ Hz $^4J_{H5-H3}=2.5$ Hz	H(5)
7.88	Doublet	1	$^4J_{H3-H5}=2.5$ Hz	H(3)
9.86	Broad singlet	1	-	NH
10.24	Singlet	1	-	CH(7)
10.44	Broad singlet	1	-	OH

5


¹³C NMR (DMSO-d₆)

Chemical shifts $\delta \pm 0.1$ ppm	Number of carbons	Assignment
23.6	1	CH ₃
117.3	1	CH(6)
118.5	1	CH(3)
121.8	1	Quaternary C (2)
127.9	1	CH(5)

131.5	1	Quaternary C (4)
156.6	1	Quaternary C (1)
167.9	1	CH-C=O
191.0	1	CH(7) -aldehyde-

B. Methyl 2-(2-formyl-4-N-acetamidophenoxy)hexanoate (compound VI:
 $R_1 = n\text{-C}_4\text{H}_9$; $R_3 = \text{CH}_3$; $R_4 = \text{CH}_3$)

- 5 5.1 g of N-(3-formyl-4-hydroxyphenyl)acetamide (compound IV) (0.028 mmol; 1 equivalent) and 10 ml of N,N-dimethylformamide (2 volumes) are placed in a conical flask. The reaction mixture is stirred at 50°C, which gives a first dark orange solution.
- Potassium carbonate (0.6 equivalent) and 8 ml of N,N-
- 10 dimethylformamide (1.5 volumes) are placed in a 50 ml reactor. The mixture is stirred at 50°C and the first solution is then added. Stirring is continued at 50°C for 1 hour, which gives a second solution.
- 6.5 g of methyl 2-bromoheptanoate (compound V) (1.05 equivalents) are then prepared in 5 ml of N,N-dimethylformamide, and are added to the
- 15 second solution, and the whole is then stirred at 75°C for 5 hours. The reaction mixture is taken up in 15 ml of dichloromethane and 15 ml of water, stirred, and the two phases are separated by settling. The dichloromethane phase is washed with water (3 × 10 ml) and this phase is then concentrated on a rotary evaporator to give 9 g of brown crystals. These crystals are
- 20 reslurried in 20 ml of water and then crushed by spatula to recover finely divided crystals suspended in the water. The mixture is then filtered, which gives 8.9 g of wet crystals, which are taken up in 25 ml of methyl tert-butyl ether. 5 ml of methanol are added and the mixture is then refluxed for 1 hour in the presence of carbon black (0.1 g). The hot reaction mixture is filtered
- 25 through Celite and then concentrated to 50%. 4 g of a precipitate with an organic purity of 98% are thus recovered. The filtrate is again concentrated and then stirred at room temperature, which brings about the appearance of a new precipitate, which is filtered off and dried. 1.6 g of the desired compound are thus recovered.

¹H NMR (CDCl₃)

Chemical shifts $\delta \pm 0.01$ ppm	Multiplicity	Integration	Coupling constants ($ J \pm 0.5$ Hz)	Assignment
0.92	Triplet	3	$^3J_{\text{CH}_3\text{-CH}_2}=7.5$ Hz	CH ₃ (12)
1.38	Sextet	2	$^3J_{\text{CH}_2\text{-CH}_3}=7.5$ Hz $^3J_{\text{CH}_2\text{-CH}_2}=7.5$ Hz	CH ₂ (11)
1.49	Quintet	2	$^3J_{\text{CH}_2\text{-CH}_2}=7.5$ Hz	CH ₂ (10)
2.01	Multiplet	2	-	CH ₂ (9)
2.15	Singlet	3	-	CH ₃ (16)
3.74	Singlet	3	-	CH ₃ (14)
4.74	Triplet	2	$^3J_{\text{CH-CH}_2}=6.0$ Hz	CH(8)
6.79	Doublet	1	$^3J_{\text{H}_6\text{-H}_5}=9.0$ Hz	H(6)
7.62	Doublet	1	$^4J_{\text{H}_3\text{-H}_5}=3.0$ Hz	H(3)
8.09	Doubled doublet	1	$^3J_{\text{H}_5\text{-H}_6}=9.0$ Hz $^4J_{\text{H}_5\text{-H}_3}=3.0$ Hz	H(5)
8.11	Broad singlet	1	-	NH
10.51	Singlet	1	-	CH(7)

5

¹³C NMR (CDCl₃)

Chemical shifts $\delta \pm 0.1$ ppm	Number of carbons	Assignment
13.9	1	CH ₃ (12)

22.3	1	CH ₂ (11)
24.3	1	CH ₃ (16)
27.3	1	CH ₂ (10)
32.4	1	CH ₂ (9)
52.5	1	CH ₃ (14)
77.4	1	CH(7)
114.1 – 119.2 – 128.4	3	CH(3, 5, 6)
125.4	1	Quaternary C (2)
132.7	1	Quaternary C (4)
156.8	1	Quaternary C (1)
168.9 – 171.6	2	C=O (13, 15)
189.5	1	CH(7) –aldehyde-

C. 2-(2-Formyl-4-N-acetamidophenoxy)hexanoic acid

(compound VII: R₁= n-C₄H₉; R₃= CH₃)

5 3.5 g of methyl 2-(2-formyl-4-N-acetamidophenoxy)hexanoate (compound VI) (0.0114 mol; 1 equivalent) and then about 15 ml of methyl tert-butyl ether (about 3 volumes) are placed in a round-bottomed flask. The mixture is stirred at room temperature so as to form a suspension of this ester.


10 A sodium hydroxide solution (0.57 g/5 ml of water) is then added and the mixture is stirred for 1 hour. 5 ml of water are then added and stirring is continued until total dissolution of the ester compound is obtained. Two phases are thus recovered, one of methyl tert-butyl ether, the other an aqueous phase containing a carboxylic acid. These two phases are separated by settling and 5 ml of 36% hydrochloric acid/5 ml of water are then added to

15 the aqueous phase (pH = 14). The aqueous phase is brought to pH 1, which brings about the appearance of a yellow precipitate, and is then stirred for 30 minutes at room temperature. The precipitate is filtered off through a sinter, which gives 3.1 g of an orange-yellow mass with an organic purity of 99.7%. This mass is taken up in 20 ml of acetone, brought to reflux and then filtered

20 while hot. 1.7 g of wet white compound are thus obtained, and are dried to

give 1.5 g of the desired compound in the form of white crystals with an organic purity of 99.7%.

^1H NMR (DMSO- d_6)



5

Chemical shifts $\delta \pm 0.01$ ppm	Multiplicity	Integration	Coupling constants ($ J \pm 0.5$ Hz)	Assignment
0.89	Triplet	3	$^3J_{\text{CH}_3\text{-CH}_2}=7.0$	CH ₃ (11)
1.35	Sextet	2	$^3J_{\text{CH}_2\text{-CH}_3}=7.0$ $^3J_{\text{CH}_2\text{-CH}_2}=7.0$	CH ₂ (10)
1.48	Quintet	2	$^3J_{\text{CH}_2\text{-CH}_2}=7.0$	CH ₂ (9)
1.94	Broad doublet of triplets	2	$^3J_{\text{CH}_2\text{-CH}_2}=7.0$ $^3J_{\text{CH}_2\text{-CH}}=6.0$	CH ₂ (8)
2.02	Singlet	3	-	CH ₃ (13)
4.90	Triplet	1	$^3J_{\text{H}_7\text{-H}_8}=6.0$	CH(7)
7.07	Doublet	1	$^3J_{\text{H}_6\text{-H}_5}=9.0$	CH(6)
7.76	Doubled doublet	1	$^3J_{\text{H}_5\text{-H}_5}=9.0$ $^4J_{\text{H}_5\text{-H}_3}=2.5$	CH(5)
7.93	Doublet	1	$^4J_{\text{H}_3\text{-H}_5}=2.5$	CH(3)
9.98	Singlet	1	-	NH
10.43	Singlet	1	-	CH (aldehyde)
13.12	Broad singlet	1	-	COOH

¹³C NMR (DMSO-d₆)


Chemical shifts $\delta \pm 0.1$ ppm	Number of carbons	Assignment
13.7	1	CH ₃ (11)
21.7	1	CH ₂ (10)
23.7	1	CH ₃ (13)
26.7	1	CH ₂ (9)
31.5	1	CH ₂ (8)
76.4	1	CH(7)
114.9 – 117.3 – 126.9	3	CH aromatic (3, 5, 6)
124.6	1	Quaternary C aromatic (2)
133.2	1	Quaternary C aromatic (4)
156.0	1	Quaternary C aromatic (1)
168.1 – 172.0	2	C=O (12, 14)
189.0	1	CH (aldehyde)

D. 2-n-Butyl-5-N-acetamidobenzofuran (compound VIII: R₁= n-C₄H₉; R₃= CH₃)

- 5 1 g of benzenesulfonyl chloride and 1 ml of toluene are placed in a round-bottomed flask. The mixture is stirred, 1.3 g of triethylamine (3.6 equivalents) are introduced and stirring is continued at 80°C for 20 minutes, during which time the reaction medium becomes increasingly dark. A solution of 1 g of 2-(2-formyl-4-N-acetamidophenoxy)hexanoic acid (compound VII)
- 10 (0.00355 mol; 1 equivalent) in 3 ml of toluene and 2 ml of methyl tert-butyl ether is then introduced. The mixture is heated at 80°C for 2 hours while monitoring the reaction kinetics by gas chromatography. The reaction mixture is cooled to about 50°C and hydrolyzed by adding 4 ml of water. The toluene phase and the aqueous phase are separated by settling. 2 ml of water and 0.2
- 15 ml of 36% hydrochloric acid are added to this organic phase. The mixture is stirred for 5 minutes and the two phases are then separated by settling. The toluene phase is washed with 2 ml of water, and the two phases are separated by settling. The organic phase is washed with a solution of 0.9 g of

sodium hydroxide at 23% in 1.5 ml of water. The mixture is stirred, the toluene phase is separated out by settling and washed with 2 g of 10% sodium chloride solution. The two phases are separated by settling and the toluene phase is concentrated on a rotary evaporator to recover 1.1 g of desired compound in the form of a brown oil.

^1H NMR (CDCl_3)



Chemical shifts $\delta \pm 0.01$ ppm	Multiplicity	Integration	Coupling constants ($ J \pm 0.5$ Hz)	Assignment
0.94	Triplet	3	$^3J_{\text{CH}_3-\text{CH}_2}=7.5$ Hz	CH_3 (12)
1.40	Sextet	2	$^3J_{\text{CH}_2-\text{CH}_3}=^3J_{\text{CH}_2-\text{CH}_2}=7.5$ Hz	CH_2 (11)
1.70	Quintet	2	$^3J_{\text{CH}_2-\text{CH}_2}=7.5$ Hz	CH_2 (10)
2.14	Singlet	3	-	CH_3 (13)
2.73	Triplet	2	$^3J_{\text{CH}_2-\text{CH}_2}=7.5$ Hz	CH_2 (9)
6.29	Singlet	1	-	H(2)
7.12	Doubled doublet	1	$^3J_{\text{H}_5-\text{H}_6}=8.5$ Hz $^4J_{\text{H}_5-\text{H}_3}=2.0$ Hz	H(5)
7.28	Doublet	1	$^3J_{\text{H}_6-\text{H}_5}=8.5$ Hz	H(6)
7.64	Broad singlet	1	-	NH
7.72	Doublet	1	$^4J_{\text{H}_3-\text{H}_5}=2.0$ Hz	H(3)

10

^{13}C NMR (CDCl_3)


Chemical shifts $\delta \pm 0.1$ ppm	Number of carbons	Assignment
13.8	1	CH ₃ (12)
22.3	1	CH ₂ (11)
24.4	1	CH ₃ (13)
28.2	1	CH ₂ (10)
29.8	1	CH ₂ (9)
102.1	1	CH(2)
110.6 – 112.5 – 116.5	3	CH(3), CH(5), CH(6)
129.5 – 132.8 – 151.8 – 160.8 – 168.6	5	Aromatic quaternary carbons (1, 4, 7, 8) and C=O (14)

E. 2-n-Butyl-3-(4-methoxybenzoyl)-5-N-acetamidobenzofuran (compound X:
R₁= n-C₄H₉; R₃= CH₃; R₅=CH₃)

10 g of 2-n-butyl-5-N-acetamidobenzofuran (compound VIII) (0.04 mol;
5 1 equivalent) and then a solution of 29.4 g of 4-methoxybenzoyl chloride
(compound IX) (0.054 mol; 1.25 g equivalents) in dichloroethane are placed in
a 250 ml reactor. The whole is stirred at 40°C until dissolution is complete,
and 8.8 g of ferric chloride (0.054 mol; 1.25 equivalents) are then added
portionwise. The temperature is maintained at 40°C for 1 hour while
10 monitoring the reaction progress. The reaction medium is hydrolyzed by
adding 100 ml of water, and is heated to 50°C. The two phases are then
separated by settling, and the organic phase is recovered and concentrated
under vacuum on a rotary evaporator. 21 g of an orange oil which forms
crystals are thus obtained. This oil is taken up in 40 ml of ethyl acetate
15 (2 volumes) and brought to reflux in a round-bottomed flask, which brings
about dissolution of the crystals. The reaction medium is allowed to cool to
room temperature with stirring, which brings about the appearance of a
precipitate, which is kept in contact with an ice bath (5°C) for 10 minutes. The
reaction medium is filtered and pale yellow crystals are recovered. These
20 crystals are dried in a vacuum oven at 50°C to give 10.1 g of crystals. These
crystals are taken up in ethyl acetate (4 volumes) and the whole is then

refluxed until dissolution is complete. The reaction medium is allowed to cool to room temperature, and the crystals formed are then filtered off through a sinter. The crystals obtained are rinsed with 10 ml of ethyl acetate and dried in an oven at 50°C to give 8.2 g of desired compound in the form of a first crop of crystals (organic purity: 99.7%) and 1.3 g of the same compound in the form of a second crop of crystals (organic purity: 99.1%).

^1H NMR (CDCl_3)



Chemical shifts $\delta \pm 0.01$ ppm	Multiplicity	Integration	Coupling constants ($ J \pm 0.5$ Hz)	Assignment
0.86	Triplet	3	$^3J_{\text{CH}_3-\text{CH}_2}=7.5$ Hz	CH_3 (12)
1.32	Sextet	2	$^3J_{\text{CH}_2-\text{CH}_3}=^3J_{\text{CH}_2-\text{CH}_2}=7.5$ Hz	CH_2 (11)
1.71	Quintet	2	$^3J_{\text{CH}_2-\text{CH}_2}=7.5$ Hz	CH_2 (10)
2.08	Singlet	3	-	CH_3 (13)
2.84	Triplet	2	$^3J_{\text{CH}_2-\text{CH}_2}=7.5$ Hz	CH_2 (9)
3.86	Singlet	3	-	CH_3 (20)
6.93	Doublet (AB system)	2	$^3J_{\text{H}_{18}-\text{H}_{17}}=9.0$ Hz $^3J_{\text{H}_{18'}-\text{H}_{17'}}=9.0$ Hz	H(18, 18')
7.32	Doublet	1	$^4J_{\text{H}_3-\text{H}_5}=2.0$ Hz	H(3)
7.36	Doublet	1	$^3J_{\text{H}_6-\text{H}_5}=9.0$ Hz	H(6)
7.54	Doubled doublet	1	$^3J_{\text{H}_5-\text{H}_6}=9.0$ Hz $^4J_{\text{H}_5-\text{H}_3}=2.0$ Hz	H(5)
7.57	Broad singlet	1	-	NH
7.80	Doublet (AB system)	2	$^3J_{\text{H}_{17}-\text{H}_{18}}=9.0$ Hz $^3J_{\text{H}_{17'}-\text{H}_{18'}}=9.0$ Hz	H(17, 17')

¹³C NMR (CDCl₃)

Chemical shifts $\delta \pm 0.1$ ppm	Number of carbons	Assignment
13.7	1	CH ₃ (12)
22.4	1	CH ₂ (11)
24.3	1	CH ₃ (13)
28.0	1	CH ₂ (10)
30.1	1	CH ₂ (9)
55.5	1	CH ₃ (20)
111.0 – 113.0 – 118.2	3	CH(3), CH(5), CH(6)
113.8	2	CH (18, 18')
131.7	2	CH (17, 17')
116.9 – 127.5 – 131.7 – 133.8 – 150.7 – 163.6 – 165.4 – 168.5	8	Aromatic quaternary carbons (1,2,4,7,8,16,19) and C=O(14)
190.4	1	C=O(15)


5

F. 2-n-Butyl-3-(4-hydroxybenzoyl)-5-N-acetamidobenzofuran (compound XI:
R₁= n-C₄H₉; R₃= CH₃)

10 5 g of 2-n-butyl-3-(4-methoxybenzoyl)-5-N-acetamidobenzofuran
(compound X) (0.0137 mol; 1 equivalent) and 15 ml of chlorobenzene (3
volumes) are placed in a 250 ml reactor. The mixture is stirred at 60°C until
the methoxy derivative is partially dissolved, followed by addition, in a single
portion, of 5.5 g of aluminum chloride (0.0411 mol; 3 equivalents), which
15 brings about a change in the reaction medium. The methoxy derivative is
dissolved and a temperature of 60°C is maintained for 4 hours. The mixture is

then hydrolyzed by adding 15 ml of water (3 volumes) while stirring at about 45°C ± 5°C. The reaction medium is then extracted with n-butanol at a temperature of about 45°C. The two phases are separated by settling while hot. A butanol phase is thus recovered, in which is observed a precipitate which appears when the temperature lowers. The organic phase is filtered and 2 g of a white product are thus recovered. The filtrate is concentrated under vacuum on a rotary evaporator and the 4 g of oil thus obtained and the precipitate are then taken up in ethyl acetate (2 volumes). The medium is then heated to reflux to dissolve the particles, and is allowed to cool to room temperature. The crystals obtained are filtered off and 1.3 g of a slightly yellow product are thus recovered (organic purity: 97.9%). A further recrystallization from ethyl acetate is performed using the filtrate obtained, which makes it possible to recover 0.8 g of desired compound with an organic purity of 96.6%.

¹H NMR (DMSO-d₆)



Chemical shifts $\delta \pm 0.01$ ppm	Multiplicity	Integration	Coupling constants ($ J \pm 0.5$ Hz)	Assignment
0.77	Triplet	3	$^3J_{\text{CH}_3\text{-CH}_2}=7.5$ Hz	CH ₃ (12)
1.21	Sextet	2	$^3J_{\text{CH}_2\text{-CH}_3}=^3J_{\text{CH}_2\text{-CH}_2}=7.5$ Hz	CH ₂ (11)
1.61	Quintet	2	$^3J_{\text{CH}_2\text{-CH}_2}=7.5$ Hz	CH ₂ (10)
1.98	Singlet	3	-	CH ₃ (13)
2.75	Triplet	2	$^3J_{\text{CH}_2\text{-CH}_2}=7.5$ Hz	CH ₂ (9)
6.87	Doublet	2	$^3J_{\text{H}18\text{-H}17}=8.5$ Hz $^3J_{\text{H}18'\text{-H}17'}=8.5$ Hz	H(18, 18')

7.51	Multiplet	2	-	H(3), H(5),
7.64	Doublet	1	-	H(6)
7.66	Doublet	2	$^3J_{H17-H18}=8.5$ Hz $^3J_{H17'-H18'}=8.5$ Hz	H(17, 17')
9.90	Broad singlet	1	-	NH or OH
10.43	Very broad singlet	1	-	NH or OH

 ^{13}C NMR (DMSO- d_6)

Chemical shifts $\delta \pm 0.1$ ppm	Number of carbons	Assignment
13.3	1	CH ₃ (12)
21.5	1	CH ₂ (11)
23.8	1	CH ₃ (13)
27.0	1	CH ₂ (10)
29.4	1	CH ₂ (9)
110.7 – 116.5	3	CH(3), CH(5), CH(6)
115.3	2	CH (18, 18')
131.5	2	CH (17, 17')
116.5 – 126.7 – 129.4 – 135.4 – 149.1 – 162.2 – 163.4 – 167.9	8	Aromatic quaternary carbons (1,2,4,7,8,16,19) and C=O(14)
189.1	1	C=O(15)


5

G. 2-n-Butyl-3-{4-[3-(di-n-butylamino)propoxy]benzoyl}-5-N-acetamidobenzofuran (compound II: $R_1 = n\text{-C}_4\text{H}_9$; $R_2 = 3\text{-(di-n-butylamino)propyl}$; $R_3 = \text{CH}_3$)

10

4 g of 2-n-butyl-3-(4-hydroxybenzoyl)-5-N-acetamidobenzofuran (compound XI) (0.0114 mol; 1 equivalent), 2.1 g of potassium carbonate (1.3 equivalents) and 15 ml of methyl ethyl ketone are placed in a three-necked round-bottomed flask. The reaction medium is stirred, and 2.8 g of 1-chloro-3-(di-n-butylamino)propane (compound XII) (0.0137 mol; 1.2 equivalents) are added at a temperature of 80°C and the whole is heated at the reflux temperature of the methyl ethyl ketone for at least 8 hours. The white deposit formed around the flask is recovered and is then stirred again by adding a further 0.1 equivalent of chloramine. Heating is continued for about 5 hours and the reaction medium is then concentrated under vacuum on a rotary evaporator. The oil thus obtained is taken up in 15 ml of water and 15 ml of methyl tert-butyl ether, and the two phases are separated by settling. The organic phase is washed with 0.5 equivalent of acetic acid in 15 ml of water. The mixture is stirred, the phases are separated by settling and the organic phase is washed with 15 ml of water. This organic phase is concentrated, and 5.2 g of desired compound are thus recovered in the form of a whitish oil.

^1H NMR (CDCl_3)



Chemical shifts $\delta \pm 0.01$ ppm	Multiplicity	Integration	Assignment
0.87	Triplet	3	CH_3 (a)
0.89	Triplet	6	$2 \times \text{CH}_3$ (k)
1.31	Sextet	6	CH_2 (b) $2 \times \text{CH}_2$ (j)
1.48	Multiplet	4	$2 \times \text{CH}_2$ (i)
1.72	Quintet	2	CH_2 (c)
2.00	Multiplet	2	CH_2 (f)

2.09	Singlet	3	CH ₃ -C=O
2.54	Triplet	4	2x CH ₂ (h)
2.72	Triplet	2	CH ₂ (g)
2.85	Triplet	2	CH ₂ (d)
4.07	Triplet	2	CH ₂ (e)
6.91	Doublet	2	2x CH(12)
7.26	Doublet	1	CH(2)
7.37	Doublet	1	CH(5)
7.61	Doubled doublet	1	CH(4)

EXAMPLE

- 5 2-n-Butyl-3-{4-[3-(di-n-butylamino)propoxy]benzoyl}-5-aminobenzofuran hydrochloride (compound I: R₁= n-C₄H₉; R₂= 3-(di-n-butylamino)propyl)

1 1 equivalent of 2-n-butyl-3-{4-[3-(di-n-butylamino)propoxy]benzoyl}-5-N-acetamidobenzofuran (compound II) and 4 volumes of ethanol are placed in a
10 reactor. 6 equivalents of 36% hydrochloric acid are then added and the reaction medium is heated to reflux, which brings about formation of the desired hydrochloride. The reaction medium is then washed, at room temperature, with aqueous sodium carbonate solution so as to liberate this hydrochloride, which passes into the aqueous phase, from which it is
15 extracted.

In this manner, the desired compound is obtained in an organic purity of 93%.


Degree of conversion of compound II: 100%

20 The conversion of 2-n-butyl-3-{4-[3-(di-n-butylamino)propoxy]benzoyl}-5-aminobenzofuran hydrochloride into dronedarone hydrochloride is then performed as described in patent EP 0 471 609.

CLAIMS

1. Process for preparing 5-aminobenzofuran derivatives of general formula:


5



I

10 and also the addition salts thereof, in which R_1 represents hydrogen or an alkyl group and R_2 represents an alkyl or dialkylaminoalkyl group, characterized in that a 5-N-alkylamidobenzofuran derivative of general formula:

15



II

20 in which R_1 and R_2 have the same meaning as previously and R_3 represents a linear or branched C_1 - C_4 alkyl group, is treated with a strong acid to form an acid-addition salt of the compound of formula I, which salt is itself treated, if necessary, with a basic agent to form this compound of formula I in free base form.

2. Process according to Claim 1, characterized in that:

25

- R_1 represents a linear or branched C_1 - C_8 alkyl group,
- R_2 represents a linear or branched C_1 - C_8 alkyl group or a dialkylaminoalkyl group in which each linear or branched alkyl group is of C_1 - C_8 .


3. Process according to Claim 1 or 2, characterized in that:

30

- R_1 represents a linear or branched C_1 - C_4 alkyl group,
- R_2 represents a linear or branched C_1 - C_4 alkyl group or a dialkylaminoalkyl group in which each linear or branched alkyl group is of C_1 - C_4 .


4. Process according to one of Claims 1 to 3, characterized in that the 5-N-acetamidobenzofuran derivative of formula II is obtained by reaction between a 4-hydroxyphenyl derivative of general formula:

5



XI

10 in which R_1 represents hydrogen or an alkyl group and R_3 represents a linear or branched C_1 - C_4 alkyl group, and an alkyl halide of general formula:




XII

15 in which R_2 represents an alkyl or dialkylaminoalkyl group and Hal represents a halogen, the reaction taking place in the presence of a basic agent and by heating in a polar solvent, to form the desired compound.

5. Process according to Claim 4, characterized in that the 4-hydroxyphenyl derivative of formula XI is obtained by dealkylation of a ketone of general formula:

20




X

25 in which R_1 represents hydrogen or an alkyl group and R_3 and R_5 each represent a linear or branched C_1 - C_4 alkyl group, this being done in the presence of aluminum chloride and in a non-polar solvent, to form the desired compound.

6. Process according to Claim 5, characterized in that the ketone of formula X is obtained by coupling between, on the one hand, a benzofuran derivative of general formula:

30



VIII


- 5 in which R_1 represents hydrogen or an alkyl group and R_3 represents a linear or branched C_1 - C_4 alkyl group, and, on the other hand, an acyl halide of general formula:



IX

- in which R_5 represents a linear or branched C_1 - C_4 alkyl group and Hal represents a halogen, this being done in the presence of a Lewis acid and in a
 15 non-polar solvent, the reaction medium thus formed then being hydrolyzed in the presence of a strong acid to form the desired compound.

7. Process according to Claim 6, characterized in that the benzofuran derivative of formula VIII is obtained by cyclization of a carboxylic acid derivative of general formula:




25

VII

- in which R_1 represents hydrogen or an alkyl group and R_3 represents a linear or branched C_1 - C_4 alkyl group, this being done by heating in an aprotic solvent and in the presence of an organic base and a benzenesulfonyl halide, to form the desired compound.


- 30 8. Process according to Claim 7, characterized in that the carboxylic acid derivative of formula VII is obtained by saponification of an ester of general formula:




5

in which R₁ represents hydrogen or an alkyl group and R₃ and R₄ each represent a linear or branched C₁-C₄ alkyl group, this being done in the presence of a strong base, the reaction taking place at room temperature and in a suitable solvent, to give a salt of the carboxylic acid derivative, which is then treated with a strong acid to form the desired compound.

9. Process according to Claim 8, characterized in that the ester of formula VI is obtained by reacting an N-phenylalkylamide derivative of general formula:




in which R₃ represents a linear or branched C₁-C₄ alkyl group, with an ester of
20 general formula:




in which R₁ represents hydrogen or an alkyl group, R₄ represents a linear or branched C₁-C₄ alkyl group and Hal represents a halogen, this being done in the presence of a basic agent and by heating in a polar solvent, to form the
30 desired compound.

10. Process for preparing 2-n-butyl-3-{4-[3-(di-n-butylamino)propoxy]benzoyl}-5-methanesulfonamidobenzofuran or dronedarone of formula:




and also pharmaceutically acceptable salts thereof, characterized in that:

a) a 5-N-alkylamidobenzofuran derivative of general formula:



IIa

- 5 in which R_3 represents a linear or branched C_1 - C_4 alkyl group, is treated with a strong acid to form an acid-addition salt of 2-n-butyl-3-{4-[3-(di-n-butylamino)propoxy]benzoyl}-5-aminobenzofuran of formula:




Ia

- 10 which salt is itself treated with a basic agent to produce this compound of formula Ia in free base form,
 b) the 2-n-butyl-3-{4-[3-(di-n-butylamino)propoxy]benzoyl}-5-aminobenzofuran thus obtained is treated with methanesulfonyl chloride to form the dronedarone compound in free base form, which may be reacted, if
 15 necessary, with an acid to produce a pharmaceutically acceptable salt of this compound.


11. Process according to Claim 10, characterized in that the 5-N-alkylamidobenzofuran derivative of formula IIa is obtained by performing a sequence of steps according to which:

- 20 a) an N-phenylalkylamide derivative of general formula:




IV

in which R_3 represents a linear or branched C_1 - C_4 alkyl group, is reacted with an ester of general formula:



Va


in which R_4 represents a linear or branched C_1 - C_4 alkyl group and Hal represents a halogen, this being done in the presence of a basic agent and by heating in a polar solvent, to form an ester of general formula:



Vla

in which R_3 and R_4 have the same meaning as previously,


b) the ester of formula Vla is saponified in the presence of a strong base, the reaction proceeding at room temperature and in a suitable solvent, to give a salt of the carboxylic acid derivative, which is treated with a strong acid to form the carboxylic acid derivative of general formula:



VIIa

in which R_3 has the same meaning as previously,


c) the carboxylic acid derivative of formula VII_a is cyclized by heating in an aprotic solvent and in the presence of an organic base and a benzenesulfonyl halide to form a benzofuran derivative of general formula:



VIIIa

in which R_3 has the same meaning as previously,


d) the benzofuran derivative of formula VIII_a is coupled with an acyl halide of general formula:



5

IX

in which R_5 represents a C_1 - C_4 alkyl group and Hal represents a halogen, this being done in the presence of a Lewis acid and in a non-polar solvent, the reaction medium thus formed then being hydrolyzed in the presence of a strong acid to form a ketone of general formula:




10

Xa

in which R_3 and R_5 have the same meaning as previously,

e) the ketone of formula X_a is dealkylated in the presence of aluminum chloride and in a non-polar solvent to form a 4-hydroxyphenyl derivative of general formula:




15

Xla

in which R_3 has the same meaning as previously,

f) the 4-hydroxyphenyl derivative of formula XI_a is reacted with an alkyl halide of general formula:




5

in which Hal has the same meaning as previously, the reaction taking place in the presence of a basic agent and by heating in a polar solvent, to form the desired compound of formula II_a.


12. Process according to Claim 10 or 11, characterized in that R₃, R₄ and R₅ each represent methyl and Hal represents chlorine or bromine.

13. N-Phenylalkylamide derivatives of general formula:



XIII

in which R₃ represents a linear or branched C₁-C₄ alkyl group and Y represents hydrogen or a group of general formula:



XIV


in which R₁' and R₆ each independently represent hydrogen or a linear or branched C₁-C₄ alkyl group,

with the exclusion of the compounds for which R₃ = Me, Et, and isopropyl, and Y = H.


14. N-Phenylalkylamide derivatives according to Claim 13, characterized in that R₃ represents methyl and Y represents the group XIV in which R₁' represents n-butyl and R₆ represents methyl.

15. N-Phenylalkylamide derivatives according to Claim 13, characterized in that R₃ represents methyl and Y represents the group XIV in which R₁' represents n-butyl and R₆ represents hydrogen.

16. 5-N-Alkylamidobenzofuran derivatives of general formula:



in which R_1' represents hydrogen or a linear or branched C_1 - C_4 alkyl group, R_3 represents a linear or branched C_1 - C_4 alkyl group and Z represents hydrogen or a group of general formula:



5

XVI

in which R_2' represents hydrogen, a linear or branched C_1 - C_4 alkyl group or a dialkylaminoalkyl group in which each linear or branched alkyl group is of C_1 - C_4 ,

10 with the exclusion of the compounds for which $R_3 = \text{Me}$, $Z = \text{H}$, and $R_1' = \text{H}$, Me , Et and $n\text{-Bu}$.

17. 5-N-Alkylamidobenzofuran derivative according to Claim 16, characterized in that R_1' represents n -butyl, R_3 represents methyl and Z represents the group XVI in which R_2' represents hydrogen.

15 18. 5-N-Alkylamidobenzofuran derivative according to Claim 16, characterized in that R_1' represents n -butyl, R_3 represents methyl and Z represents the group XVI in which R_2' represents methyl.

19. 5-N-Alkylamidobenzofuran derivative according to Claim 16, characterized in that R_1' represents n -butyl, R_3 represents methyl and Z represents the group XVI in which R_2' represents 3-(di- n -butylamino)propyl.

20