


BRIEF SUMMARY OF THE INVENTION

[0016] The present ~~invention-disclosure~~ provides, in one aspect, compounds having formula I,



(I)

or pharmaceutically acceptable salts, hydrates or N-oxides thereof. The various groups (e.g., R¹, R², R³, R⁴, Z, X^a, X^b, X^c and the subscript n) are described in the Detailed Description of the Invention.

[0017] The compounds provided herein are useful for binding to CXCR7, and treating diseases that are dependent, at least in part, on CXCR7 activity. Accordingly, the present ~~invention-disclosure~~ provides in further-another aspects, a pharmaceutical compositions containing one or more of the above-noted compounds in admixture with a pharmaceutically acceptable excipient.

[0018] ~~In still another aspect, t~~The present ~~invention-disclosure~~ also provides methods for treating various diseases, discussed further herein, comprising administering to a subject in need to such treatment a therapeutically effective amount of a compound of the above formula for a period of time sufficient to treat the disease. In another aspect, the present disclosure provides a compound as defined herein for use in treating a disease or disorder in a mammal. In another aspect, the present disclosure provides use of a compound as defined herein in the manufacture of a medicament for the treatment of a disease or disorder selected from the group consisting of cancer, inflammation and neural or progenitor/stem cell disorders.

[0019] ~~In yet another aspect, t~~The present ~~invention-disclosure~~ also provides methods of diagnosing disease in an individual. In these methods, the compounds provided herein are administered in labeled form to a subject, followed by diagnostic imaging to determine the presence or absence of CXCR7. ~~In a related aspect, a~~The method of diagnosing disease is may be carried out by contacting a tissue or blood sample with a labeled compound as

provided herein and determining the presence, absence, or amount of CXCR7 in the sample.

In another aspect, the present disclosure provides a method for imaging a tumor, organ, or tissue, said method comprising detecting a radiolabeled or detectable form of a compound as defined herein in a subject who had been administered the compound to determine where the

5 compound is concentrated in the subject. In another aspect, the present disclosure provides a method for detecting elevated levels of CXCR7 in a sample, said method comprising (a) contacting a sample suspected of having elevated levels of CXCR7 with a radiolabeled or detectable form of a compound as defined herein; (b) determining a level of compound that is bound to CXCR7 present in the sample to determine the level of CXCR7 present in the
10 sample; and (c) comparing the level determined in step (b) with a control sample to determine if elevated levels of CXCR7 are present in the sample.

[0020] In some embodiments, an amount of a chemotherapeutic agent or radiation is administered to the subject prior to, subsequent to or in combination with the compounds of the present ~~invention~~disclosure. In some embodiments, the amount is sub-therapeutic when
15 the chemotherapeutic agent or radiation is administered alone.

[0020a] In another aspect, the present disclosure provides an *in vitro* method of inhibiting the binding of chemokines I-TAC or SDF-1 to a CXCR7 receptor, comprising contacting a compound as defined herein with a cell that expresses the CXCR7 receptor for a time sufficient to inhibit the binding of the chemokines to the CXCR7 receptor.

1 17. The compound of claim 1, selected from the compounds of the
2 Examples.

1 18. The compound of claim 1, selected from those in Table 1.

1 19. A pharmaceutical composition comprising a compound of claim 1 and
2 a pharmaceutically acceptable excipient.

1 20. The pharmaceutical composition of claim 19, wherein the compound is
2 a compound of Table 1.

1 21. A compound of claim 1 for use in ~~method of treating~~ a disease or
2 disorder in a mammal, ~~said method comprising administering to said subject a therapeutically~~
3 ~~effective amount of a compound of claim 1, for a period of time sufficient to treat said~~
4 ~~disease or disorder.~~

1 22. The ~~method compound~~ compound of claim 21, wherein the compound is a
2 compound of Table 1.

1 23. The ~~method compound~~ compound of claim 21, wherein said disease or disorder is
2 selected from the group consisting of cancer, inflammation and neural or progenitor/stem cell
3 disorders.

1 24. An in vitro method of inhibiting the binding of chemokines I-TAC or
2 SDF-1 to a CXCR7 receptor, comprising contacting a compound of claim 1 with a cell that
3 expresses the CXCR7 receptor for a time sufficient to inhibit the binding of the chemokines
4 to the CXCR7 receptor.

1 25. The in vitro method of claim 24, wherein the compound is a compound
2 of Table 1.

1 26. A method for imaging a tumor, organ, or tissue, said method
2 comprising:

3 ~~(a) administering to a subject in need of such imaging, a radiolabeled or~~
4 ~~detectable form of a compound of any of claims 1 to 15; and~~

5 ~~(b) detecting a radiolabeled or detectable form of a compound of any of claims~~
6 ~~1 to 15 in a subject who had been administered~~ said compound to determine where said
7 compound is concentrated in said subject.
8

1 27. A method in accordance with claim 26, wherein said compound is
2 radiolabeled.

1 28. A method for detecting elevated levels of CXCR7 in a sample, said
2 method comprising:

3 (a) contacting a sample suspected of having elevated levels of CXCR7 with a
4 radiolabeled or detectable form of a compound of any of claims 1 to 15 ;

5 (b) determining a level of compound that is bound to CXCR7 present in said
6 sample to determine the level of CXCR7 present in said sample; and

7 (c) comparing the level determined in step (b) with a control sample to
8 determine if elevated levels of CXCR7 are present in said sample.

1 29. A method in accordance with claim 28, wherein said compound is
2 radiolabeled.

3 30. Use of a compound of claim 1 in the manufacture of a medicament for
4 the treatment of a disease or disorder selected from the group consisting of cancer,
5 inflammation and neural or progenitor/stem cell disorders.