

- 1 -

TWO-and THREE-COMPONENT FUNGICIDAL MIXTURES

The present invention relates to fungicidal two- and three-component mixtures based on metalaxyl having an R-enantiomer content of more than 70 % by weight, and to the use thereof in controlling and preventing Oomycetes infestation. The metalaxyl component is called active ingredient I.

The following fungicides may be used as the second component II of the mixture:

IIA) mancozeb;

IIB) chlorothalonil;

IIC) a copper salt, for example CuCO_3 , CuSO_4 , $\text{Cu}(\text{OH})_2 \cdot \text{CuCl}_2$, especially $\text{Cu}(\text{OH})_2$;

IID) folpet;


IIIE) fluazinam [= 3-chloro-N-(3-chloro-5-trifluoromethyl-2-pyridyl)- α,α,α -trifluoro-2,6-dinitro-p-toluidine];

and

IIIF) cymoxanil, which can be used either on its own or in admixture with one of the above-mentioned components IIA to IIIE (three-component mixtures).

More specifically, the invention relates to mixtures comprising metalaxyl having an R-enantiomer content of more than 85 % by weight, preferably of more than 92 % by weight, and especially containing pure R-enantiomer that is essentially free of S-enantiomer.

Metalaxyl of the formula



has an asymmetric *C atom and can be resolved into the enantiomers in customary manner (GB-P.1 500 581). Since 1975 it has been known to those skilled in the art that the R-enantiomer is far superior to the S-enantiomer in terms of fungicidal action and is in practice regarded as the true mechanism of action. Commercial metalaxyl is available in the form of the racemate. Likewise, mixtures of metalaxyl racemate with mancozeb, chlorothalonil, copper preparations, folpet, fluazinam or cymoxanil have become known

- 2 -

commercially or otherwise. There has in the past never been any practical necessity to resolve the racemate, of which half consists of the desired R-enantiomer.

It has now been found, completely surprisingly, that R-metalaxyl in pure or more than 70 % form, in admixture with the fungicidal components IIA to IIF, achieves a synergistically enhanced action which in some cases exceeds that of the prior-known mixtures based on the racemate by a factor of 10. Given that half of the racemate consists of R-enantiomer, factors of approximately 2 or, at most, 3 were to be expected.

With this completely unexpected result, the present invention constitutes a very considerable enrichment of the art and represents a possible means of reducing in an environmentally protective manner the total amount of fungicides used for controlling Oomycetes on plants.

In addition to the two-component mixture I:II, the present invention relates also to a method of controlling fungi, which comprises treating a site, for example a plant, that is infested by or threatened with infestation by fungi with, in any desired sequence or simultaneously, a) component I and b) the active ingredient of formula II.

Advantageous mixing ratios of the two active ingredients are I:II = from 10:1 to 1:100, preferably I:II = from 5:1 to 1:30. In many cases, mixtures in which the mixing ratio of the active substances I:II is from 1:1 to 1:20, for example 2:5, 1:4, 1:8 or 1:10, are advantageous.

The active ingredient mixtures I+II according to the invention have very advantageous curative, preventive and systemic fungicidal properties for protecting cultivated plants. The active ingredient mixtures of the invention can be used to inhibit or destroy the microorganisms which occur on plants or on parts of plants (the fruit, blossom, leaves, stems, tubers or roots) of different crops of useful plants, while at the same time parts of plants that grow later are also protected against such microorganisms. This applies especially also to microorganisms that have developed reduced sensitivity to metalaxyl.

The active ingredient mixtures of formulae I and II are generally used in the form of compositions. R-metalaxyl (formula I) and the active ingredient of formula II can be applied to the area or plant to be treated simultaneously or in immediate succession, if desired together with further carriers, surfactants or other application-promoting adjuvants

customarily employed in formulation technology.

Suitable carriers and adjuvants can be solid or liquid and are the substances ordinarily employed in formulation technology, e.g. natural or regenerated mineral substances, solvents, dispersants, wetting agents, tackifiers, thickeners, binders or fertilisers.

A preferred method of applying an active ingredient mixture comprising at least one of each of the active ingredients I and II is application to the parts of the plants that are above the soil, especially to the leaves (foliar application). The frequency and rate of application depend upon the biological and climatic living conditions of the pathogen. The active ingredients can, however, also penetrate the plant through the roots via the soil (systemic action) if the locus of the plant is impregnated with a liquid formulation or if the substances are introduced in solid form into the soil, e.g. in the form of granules (soil application).

The compounds of the combination are used in unmodified form or, preferably, together with the adjuvants conventionally employed in formulation technology, and are therefore formulated in known manner e.g. into emulsifiable concentrates, coatable pastes, directly sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts, granules, or by encapsulation in e.g. polymer substances. As with the nature of the compositions, the methods of application, such as spraying, atomising, dusting, scattering, coating or pouring, are chosen in accordance with the intended objectives and the prevailing circumstances. Advantageous rates of application of the active ingredient mixture are normally from 50 g to 1800 g of active ingredient (a.i.) per hectare, preferably from 100 g to 1000 g a.i./ha. Mixtures of R-metalaxyl (I) with mancozeb (IIA) are advantageously employed at rates of application of from 100 g to 120 g of I and 1600 g of IIA.

The formulations are prepared in known manner, e.g. by homogeneously mixing and/or grinding the active ingredients with extenders, e.g. solvents, solid carriers and, where appropriate, surface-active compounds (surfactants).

The agrochemical compositions generally comprise 0.1 to 99 %, preferably 0.1 to 95 %, of active ingredients of formulae I and II, 99.9 to 1 %, preferably 99.9 to 5 %, of a solid or liquid adjuvant and 0 to 25 %, preferably 0.1 to 25 %, of a surfactant.

Whereas commercial products will preferably be formulated as concentrates, the end user

- 4 -

will normally employ dilute formulations.

Suitable target crops are especially potatoes, vines, hops, maize, sugar beet, tobacco, vegetables (tomatoes, paprika, lettuce, etc.), and also bananas, natural rubber plants, as well as lawn areas and ornamentals. Other plants threatened by downy mildew have become known inter alia from the literature relating to metalaxyl.

The Examples which follow serve to illustrate the invention, "active ingredient" denoting a mixture of R-metalaxyl and compound II in a specific mixing ratio.

Wettable powders

	a)	b)	c)
active ingredient			
[I:IIA = 1:4(a), 1:16(b), 2:13(c)]	27 %	51 %	75 %
sodium lignosulfonate	5 %	5 %	-
sodium laurylsulfate	3 %	-	5 %
sodium diisobutyl-naphthalene-sulfonate	-	6 %	10 %
octylphenol polyethylene glycol ether (7-8 mol of ethylene oxide)	-	2 %	-
highly dispersed silicic acid	5 %	10 %	10 %
kaolin	60 %	26 %	-

The active ingredient is thoroughly mixed with the adjuvants and the mixture is thoroughly ground in a suitable mill, affording wettable powders which can be diluted with water to give suspensions of the desired concentration.

Emulsifiable concentrate

active ingredient	
(I:IIA, IIB, IID or IIE = 3:7)	10 %
octylphenol polyethylene glycol ether (4-5 mol of ethylene oxide)	3 %
calcium dodecylbenzenesulfonate	3 %
castor oil polyglycol ether (35 mol of ethylene oxide)	4 %
cyclohexanone	30 %
xylene mixture	50 %

- 5 -

Emulsions of any required dilution, which can be used in plant protection, can be obtained from this concentrate by dilution with water.

Extruder granules

active ingredient (I:IIIC = 1:4)	15 %
sodium lignosulfonate	2 %
carboxymethylcellulose	1 %
kaolin	82 %

The active ingredient is mixed and ground with the adjuvants, and the mixture is moistened with water. The mixture is extruded and then dried in a stream of air.

Coated granules

active ingredient (I:IID = 3:5)	8 %
polyethylene glycol (mol. wt. 200)	3 %
kaolin	89 %
(mol. wt. = molecular weight)	

The finely ground active ingredient is uniformly applied, in a mixer, to the kaolin moistened with polyethylene glycol. Non-dusty coated granules are obtained in this manner.

Suspension concentrate

active ingredient(I:IIIF:IIA = 3:1:7)	44 %
propylene glycol	10 %
nonylphenol polyethylene glycol ether (15 mol of ethylene oxide)	6 %
sodium lignosulfonate	10 %
carboxymethylcellulose	1 %
silicone oil (in the form of a 75 % aqueous emulsion)	1 %
water	28 %

The finely ground active ingredient is intimately mixed with the adjuvants, giving a suspension concentrate from which suspensions of any desired dilution can be obtained by

dilution with water. Such dilutions can be used to treat living plants by spraying, pouring or immersion and to protect them against infestation by microorganisms.

Biological Examples

Fungicides have a synergistic effect whenever the fungicidal action of the active ingredient combination is greater than the sum of the actions of the active ingredients when applied individually.

The action E to be expected for a given active ingredient combination, for example of two fungicides, obeys the so-called COLBY formula and can be calculated as follows (COLBY, L.R. "Calculating synergistic and antagonistic responses of herbicide combinations". Weeds 15, pages 20-22, 1967) (LIMPEL *et al.*, 1962 "Weeds control by ... certain combinations". Proc. NEWCL, Vol. 16, pp. 48-53):

(mg a.i./litre = milligrams of active ingredient per litre or a multiple thereof)

X = % action by fungicide I at p mg a.i./litre

Y = % action by fungicide II at q mg a.i./litre

E = the expected action of fungicides I+II at a rate of application of p+q mg a.i./litre (additive action),

then according to Colby:
$$E = X + Y - \frac{X \cdot Y}{100}$$

If the action (O) actually observed is greater than the expected action, then the action of the combination is superadditive, i.e. there is a synergistic effect.

1) Action against *Plasmopara viticola* on vines

Small vine plants of the variety "Gutedel" are each grown in a plant-pot (diameter = 6 cm) under greenhouse conditions and are sprayed in the two-leaf stage with an active ingredient mixture prepared from an emulsifiable concentrate. The following concentrations of active ingredient are each applied to four plants:

metalaxyl racemate: 2 mg and 6 mg a.i./litre;

R-metalaxyl (100 %): 0.06 mg and 0.6 mg a.i./litre;

mancozeb: 2 mg and 6 mg a.i./litre.

These dilutions are prepared in demineralised water immediately before the leaves are sprayed. In order to eliminate the effect of the gas phase of the metalaxyl in the vicinity, all the plants are separated from one another laterally by means of transparent plastics films and are kept in the dark for one day at 20-22°C and about 100 % relative humidity.

- 7 -

The entire surface of the leaves of the plants is then uniformly sprayed to drip point with a freshly prepared sporangia suspension (120 000/ml) of a strain of Plasmopara viticola that is sensitive to metalaxyl. The plants are then kept at 20-22°C and about 100 % relative humidity for 7 days under artificial daylight of 16 hours' duration. Evaluation of the infestation is then carried out. The Tables show in each case the average of four parallel sprayings. The expected action in the case of mixtures is calculated according to COLBY.

Level of action of racemic metalaxyl and R-metalaxyl in admixture with the residual component mancozeb

Metalaxyl mg a.i./l	R-enantiomer of metalaxyl mg a.i./l	Mancozeb mg a.i./l	Mixing ratio	E Expected action [COLBY]	O Observed action
2 6					12 24
	0.06 0.6				12 18
		2 6			0 20
2 6		2 2	1:1 3:1	12 24	27 71
2 6		6 6	1:3 1:1	40 30	69 90
	0.06 0.6	2 2	1:33 1:3	12 18	43 56
	0.06 0.6	6 6	1:100 1:10	30 35	64 91

The action of 2 mg of racemic metalaxyl in admixture with 2 mg or 6 mg of mancozeb corresponds to that of 0.06 mg of R-metalaxyl in admixture with 2 mg or 6 mg of mancozeb. This shows that the action is improved about 30-fold when the R-enantiomer is used

instead of racemic metalaxyl.

2) Action against Phytophthora infestans on tomato plants

Residual-protective action

After a cultivation period of 3 weeks, tomato plants are sprayed with a series of concentrations of a spray mixture prepared from an emulsifiable concentrate of the active ingredient mixture. The treated plants are infected 48 hours later with a sporangia suspension of the fungus. Fungus infestation is evaluated after incubation of the infected plants for 4 days at 90-100 % relative humidity and 20°C.

The following results are obtained:

Mancozeb [ppm a.i.]	Metalaxyl [ppm a.i.]	R-enantiomer of metalaxyl [ppm a.i.]	Percent action
20	2		18
6	0.6		0
20		2	75
6		0.6	7

Copper oxychloride [ppm a.i.]	Metalaxyl [ppm a.i.]	R-enantiomer of metalaxyl [ppm a.i.]	Percent action
60	6		35
60	2		17.5
60		6	75
60		2	50

3) Action against Phytophthora on potato plants

Residual-protective action

After a cultivation period of 3 weeks, 2- to 3-week-old potato plants (variety Bintje) are sprayed with a series of concentrations of a spray mixture prepared from an emulsifiable concentrate of the active ingredient mixture. The treated plants are infected 24 hours later with a sporangia suspension of the fungus. Fungus infestation is evaluated after incubation of the infected plants for 5 days at 90-100 % relative humidity and 20°C.

The following results are obtained:

Mancozeb [ppm a.i.]	Metalaxyl [ppm a.i.]	R-enantiomer of metalaxyl [ppm a.i.]	Percent action
20	2		70
6	0.6		15
20		2	100
6		0.6	50

Copper oxychloride [ppm a.i.]	Metalaxyl [ppm a.i.]	R-enantiomer of metalaxyl [ppm a.i.]	Percent action
60	6		17.5
60	2		0
60		6	70
60		2	35

Similarly markedly increased actions are achieved also with the other components IIB, IID, IIE and IIF of the mixture.

- 10 -

Preferred mixing ratios (in amounts by weight) are:

R-metalaxyl(I):IIA (mancozeb) = 1:1 to 1:20

I:II B (chlorothalonil) = 2:1 to 1:12

I:II C (copper preparation) = 5:1 to 1:30

I:II D (folpet) = 3:1 to 1:10

I:II E (fluazinam) = 5:1 to 1:20

I:II F (cymoxanil) = 6:1 to 1:6

I:II F:II A = 1 to 7 : 1 : 4 to 10

I:II F:II D = 1 to 7 : 1 : 2 to 8.

These marked increases in action with R-metalaxyl are observed also against other Oomycetes, especially Peronosporales, Pseudoperonospora, Albugo occidentalis, Phytophthora spp., Pythium, Bremia and other pathogens.



What is claimed is:

1. A fungicidal two- or three-component composition based on metalaxyl, wherein more than 70 % by weight of the metalaxyl is the R-enantiomer (I) and the composition comprises as the further component:
IIA) mancozeb, IIB) chlorothalonil, IIC) a copper salt, IID) folpet, IIE) fluazinam or IIF) cymoxanil, it being possible when IIF) is used for a third component selected from IIA to IIE also to be present.
2. A composition according to claim 1, wherein the ratio by weight of I:II is from 10:1 to 1:100, preferably from 5:1 to 1:30.
3. A composition according to claim 1, wherein the metalaxyl has an R-enantiomer content of more than 85 % by weight.
4. A composition according to claim 3, wherein the metalaxyl has an R-enantiomer content of more than 92 % by weight.
5. A composition according to claim 4, wherein there is used pure R-metalaxyl that is essentially free of S-enantiomer.
6. A composition according to claim 1, wherein mancozeb (IIA) is used as the second component of the mixture with R-metalaxyl in a ratio by weight of from 1:1 to 1:20.
7. A composition according to claim 1, wherein chlorothalonil (IIB) is used as the second component of the mixture with R-metalaxyl in a ratio by weight of from 2:1 to 1:12.
8. A composition according to claim 1, wherein a fungicidal copper salt (IIC) is used as the second component of the mixture with R-metalaxyl in a ratio by weight of from 5:1 to 1:30.
9. A composition according to claim 8, wherein the copper salt is copper hydroxide [Cu(OH)₂].
10. A composition according to claim 1, wherein folpet (IID) is used as the second component of the mixture with R-metalaxyl in a ratio by weight of from 3:1 to 1:10.



11. A composition according to claim 1, wherein fluazinam (IIE) is used as the second component of the mixture with R-metalaxyl in a ratio by weight of from 5:1 to 1:20.
12. A composition according to claim 1, wherein cymoxanil (IIF) is used as the second component of the mixture with R-metalaxyl in a ratio by weight of from 6:1 to 1:6.
13. A composition according to claim 12, wherein in addition to R-metalaxyl (I) and cymoxanil (IIF) there is used as a third component mancozeb (IIA) in ratios by weight of I:IIF:IIA = 1 to 7 : 1 : 4 to 10.
14. A composition according to claim 12, wherein in addition to R-metalaxyl (I) and cymoxanil (IIF) there is used as a third component folpet (IID) in ratios by weight of I:IIF:IID = 1 to 7 : 1 : 2 to 8.
15. A method of controlling and preventing Oomycetes infestation in plants, parts of plants or at the site of their growth, which comprises applying in any desired sequence, simultaneously or in immediate succession, metalaxyl having an R-enantiomer content of more than 70 % by weight and a further fungicidal component selected from mancozeb (IIA), chlorothalonil (IIB), a copper salt (IIC), folpet (IID), fluazinam (IIE) and cymoxanil (IIF).
16. A method according to claim 15, wherein when cymoxanil (IIF) is used, there is additionally used as a third component one of the components mentioned for IIA to IIE.
17. A method according to claim 15, wherein the R-enantiomer content of the metalaxyl used is more than 85 % by weight.
18. A method according to claim 17, wherein the R-enantiomer content of the metalaxyl used is more than 92 % by weight.
19. A method according to claim 18, wherein pure R-metalaxyl that is essentially free of S-enantiomer is used as the metalaxyl component.
20. A method according to claim 15, wherein Phytophthora spp., Plasmopara, Pythium, Pseudoperonospora, Albugo occidentalis and/or Bremia are controlled.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A01N 37/46 // (A01N 37/46, 59:20, 47:34, 47:14, 47:04, 43:40, 37:34)	A1	(11) International Publication Number: WO 96/01560 (43) International Publication Date: 25 January 1996 (25.01.96)
(21) International Application Number: PCT/EP95/02545 (22) International Filing Date: 30 June 1995 (30.06.95) (30) Priority Data: 2208/94-3 11 July 1994 (11.07.94) CH (71) Applicant (for all designated States except US): CIBA-GEIGY AG [CH/CH]; Klybeckstrasse 141, CH-4002 Basle (CH). (72) Inventors; and (75) Inventors/Applicants (for US only): NUMINGER, Cosima [FR/FR]; 4, rue de la 1ère Armée, F-68790 Morschwiller- le-Bas (FR). GOGGIN, John, Edward, Nicholas [GB/CH]; Rebgasse 30, CH-4102 Binningen (CH). SOZZI, Dino [CH/CH]; Teichweg 19, CH-4450 Sissach (CH). (74) Common Representative: CIBA-GEIGY AG; Patentabteilung, Klybeckstrasse 141, CH-4002 Basle (CH).	(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MX, NO, NZ, PL, RO, RU, SG SI, SK, TJ, TM, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments. <i>ATMP's copy</i>	
(54) Title: TWO-AND THREE-COMPONENT FUNGICIDAL MIXTURES (57) Abstract When used in admixture with mancozeb, chlorothalonil, a copper salt, folpet, fluazinam or cymoxanil (this being possible for the latter to be used also together with one of the other five components), metalaxyl having a high R-enantiomer content of more than 70 % by weight, or pure R-metalaxyl, exhibits a markedly increased fungicidal action against plant diseases as compared with a similar mixture in which metalaxyl is used in the form of the racemate.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

9700202-6