CLAIMS

1. Use of an immunoconjugate comprising an anti-CEACAM5-antibody in the manufacture of a medicament for treating cancer, in combination with folinic acid, 5-fluoro-uracil and irinotecan (FOLFIRI).

2. The use of claim 1, wherein the anti-CEACAM5-antibody comprises a CDR-H1 consisting of SEQ ID NO: 1, CDR-H2 consisting of SEQ ID NO: 2, CDR-H3 consisting of SEQ ID NO: 3, CDR-L1 consisting of SEQ ID NO: 4, CDR-L2 consisting of amino acid sequence NTR, and CDR-L3 consisting of SEQ ID NO: 5.

3. The use of claim 1 or 2, wherein the anti-CEACAM5-antibody comprises a variable domain of a heavy chain (VH) consisting of SEQ ID NO: 6 and a variable domain of a light chain (VL) consisting of SEQ ID NO: 7.

4. The use of any of claims 1 to 3, wherein the anti-CEACAM5-antibody comprises a heavy chain (VH) consisting of SEQ ID NO: 8 and a light chain (VL) consisting of SEQ ID NO: 9.

5. The use of any of claims 1 to 4, wherein the immunoconjugate comprises at least one cytostatic agent.

6. The use of claim 5, wherein the cytostatic agent is selected from the group consisting of radioisotopes, protein toxins, small molecule toxins, and combinations thereof.

7. The use of claim 6, wherein the small molecule toxins are selected from antimetabolites, DNA-alkylating agents, DNA-cross-linking agents, DNA-intercalating agents, anti-microtubule agents, topoisomerase inhibitors, and combinations thereof.

8. The use of claim 7, wherein the anti-microtubule agent is selected from the group consisting of taxanes, vinca alkaloids, maytansinoids, colchicine, podophyllotoxin, gruseofulvin, and combinations thereof.

9. The use of claim 8, wherein the maytansinoids are selected from the group consisting of N2'-deacetyl-N2'-(3-mercapto-1-oxopropyl)-maytansine (DM1) or N2'-

deacetyl-N-2'(4-methyl-4-mercapto-1-oxopentyl)-maytansine (DM4), and combinations thereof.

10. The use of any of claims 1 to 9, wherein the anti-CEACAM5-antibody is covalently attached via a cleavable or non-cleavable linker to the at least one cytotoxic agent.

11. The use of claim 10, wherein said linker is selected from the group consisting of N-succinimidyl pyridyldithiobutyrate (SPDB), 4-(pyridin-2-yldisulfanyl)-2-sulfo-butyric acid (sulfo-SPDB), and succinimidyl(N-maleimidomethyl) cyclohexane-1-carboxylate (SMCC).

12. The use of any of claims 1 to 11, comprising an CEACAM5-antibody, which comprises a heavy chain (VH) consisting of SEQ ID NO: 8 and a light chain (VL) consisting of SEQ ID NO: 9 (huMAb2-3), and which is covalently linked to N2'-deacetyl-N-2'(4-methyl-4-mercapto-1-oxopentyl)-maytansine (DM4) via N-succinimidyl pyridyldithiobutyrate (SPDB).

13. The use of any of claims 1 to 12, wherein the immunoconjugate is characterised by a drug-to-antibody ratio (DAR) ranging from 1 to 10.

14. The use of any of claims 1 to 13, wherein the cancer is selected from the group consisting of colorectal, stomach, pancreas, and oesophagus cancer.

15. The use of any of claims 1 to 14, wherein the immunoconjugate and FOLFIRI are to be administered simultaneously to a subject in need thereof.

16. The immunoconjugate for the use of claim 15, wherein the immunoconjugate and FOLFIRI are formulated (i) in a single pharmaceutical composition comprising the immunoconjugate and FOLFIRI, or (ii) in the form of at least two separate pharmaceutical compositions, wherein at least one pharmaceutical composition comprises the immunoconjugate, and one or more pharmaceutical compositions comprise folinic acid, 5-fluoro-uracil and irinotecan, in separate or combined formulations.

17. The immunoconjugate for the use of any of claims 1 to 14, wherein the immunoconjugate and FOLFIRI are to be administered separately or sequentially to a subject in need thereof.

18. The immunoconjugate for the use of claim 17, wherein the immunoconjugate and FOLFIRI are formulated in the form of at least two separate pharmaceutical compositions, wherein (i) at least one pharmaceutical composition comprises the immunoconjugate, and (ii) one or more pharmaceutical compositions comprise folinic acid, 5-fluoro-uracil and irinotecan, in separate or combined formulations.

19. The immunoconjugate for the use of any of claims 1 to 18, wherein the immunoconjugate comprising an anti-CEACAM5-antibody, and folinic acid, 5-fluoro-uracil and irinotecan (FOLFIRI) are to be administered in 8 to 16 cycles, wherein in one cycle,
the immunoconjugate is to be administered at a dosage of from 60 to 210 mg/m^2, at least once in the cycle;
folinic acid is to be administered at a dosage of from 100 to 300 mg/m^2 or L-folinic acid is to be administered at a dose of 100 to 200 m/m^2, at least once in the cycle;
5-fluoro-uracil is to be administered at a dosage of from 1000 to 2000 mg/m^2, at least once in the cycle, and
irinotecan is to be administered at a dosage of from 100 to 300 mg/m^2, at least once in the cycle.

20. A pharmaceutical composition comprising the immunoconjugate of any of claims 1 to 14, and folinic acid, 5-fluoro-uracil and irinotecan.

21. A kit comprising (i) a pharmaceutical composition of the immunoconjugate of any of claims 1 to 14 and (ii) one or more pharmaceutical compositions comprising folinic acid, 5-fluoro-uracil and irinotecan, in separate or combined formulations.

22. The pharmaceutical composition according to claim 20 or the kit according to claim 21 for the use for treating cancer.