CLAIMS

What is claimed is:

1. Use of a TOR kinase inhibitor in the manufacture of a medicament for treating non-small cell lung carcinoma, in a patient having non-small cell lung carcinoma characterized by a LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation, relative to wild type, wherein the TOR kinase inhibitor is 7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or a pharmaceutically acceptable salt, stereoisomer, or tautomer thereof.

2. An *in vitro* method for predicting the likelihood of a patient having non-small cell lung carcinoma being responsive to TOR kinase inhibitor therapy, comprising screening said patient's carcinoma for the presence of a LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation, relative to wild type, wherein the presence of LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation predicts an increased likelihood that TOR kinase inhibitor therapy will treat said non-small cell lung carcinoma or cervical cancer, wherein the TOR kinase inhibitor is 7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or a pharmaceutically acceptable salt, stereoisomer, or tautomer thereof.

3. An *in vitro* method for predicting therapeutic efficacy of treatment of a patient having non-small cell lung carcinoma with a TOR kinase inhibitor, comprising screening said patient's carcinoma or cancer for the presence of a LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation, relative to wild type, wherein the presence of LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation is predictive of therapeutic efficacy of treatment with a TOR kinase inhibitor, wherein the TOR kinase inhibitor is 7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or a pharmaceutically acceptable salt, stereoisomer, or tautomer thereof.


4. Use of a TOR kinase inhibitor and use of one or more agents that module AMP levels, glucose uptake, metabolism or a stress response in the manufacture of a medicament for treating non- small cell lung carcinoma, wherein the TOR kinase inhibitor is 7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or a pharmaceutically acceptable salt, stereoisomer, or tautomer thereof.

5. A kit comprising one or more containers filled with a TOR kinase inhibitor or a pharmaceutical composition thereof, reagents for detecting a LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation, an AMPK gene loss, AMPK protein loss, AMPK gene mutation, or AMPK protein mutation, or both, relative to wild type, in a patient's cancer or in a patient having a tumor syndrome and instructions for detecting LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation, AMPK gene loss, AMPK protein loss, AMPK gene mutation, or AMPK protein mutation, or both, in a patient's cancer or in a patient having a tumor syndrome, wherein the TOR kinase inhibitor is 7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or a pharmaceutically acceptable salt, stereoisomer, or tautomer thereof.

CLAIMS

What is claimed is:

1. Use of a TOR kinase inhibitor in the manufacture of a medicament for treating non-small cell lung carcinoma, cervical cancer or Peutz-Jeghers Syndrome, in a patient having non-small cell lung carcinoma, cervical cancer or Peutz-Jeghers Syndrome characterized by a LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation, relative to wild type,
wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)


or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,
or a compound having the formula (IV):



(IV)


or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

2. Use of a TOR kinase inhibitor in the manufacture of a medicament for treating non-small cell lung carcinoma or cervical cancer, in a patient having non-small cell lung carcinoma or cervical cancer characterized by a LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation, relative to wild type,
wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)


or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,
or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

3. An *in vitro* method for predicting the likelihood of a patient having non-small cell lung carcinoma or cervical cancer being responsive to TOR kinase inhibitor therapy, comprising screening said patient's carcinoma or cancer for the presence of a LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation, relative to wild type, wherein the presence of LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation predicts an increased likelihood that TOR kinase inhibitor therapy will treat said non-small cell lung carcinoma or cervical cancer,

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

4. An *in vitro* method for predicting therapeutic efficacy of treatment of a patient having non-small cell lung carcinoma or cervical cancer with a TOR kinase inhibitor, comprising screening said patient's carcinoma or cancer for the presence of a LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation, relative to wild type, wherein the presence of LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation is predictive of therapeutic efficacy of treatment with a TOR kinase inhibitor,

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

5. Use of a TOR kinase inhibitor in the manufacture of a medicament for treating Peutz-Jeghers Syndrome, in a patient having Peutz-Jeghers Syndrome characterized by a LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation wherein the use comprises screening the patient for the presence of LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation, relative to wild type,

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

6. An in vitro method for predicting the likelihood of a patient having Peutz-Jeghers Syndrome being responsive to TOR kinase inhibitor therapy, comprising screening said patient for the presence of a LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation or LKB1 protein mutation, relative to wild type, wherein the presence of LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation predicts an increased likelihood that TOR kinase inhibitor therapy will treat said Peutz-Jeghers Syndrome, wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted


heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

7. An *in vitro* method for predicting therapeutic efficacy of treatment of a patient having Peutz-Jeghers Syndrome with a TOR kinase inhibitor, comprising screening said patient for the presence of a LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation, relative to wild type, wherein the presence of LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation is predictive of therapeutic efficacy of treatment with a TOR kinase inhibitor,

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

8. Use of a TOR kinase inhibitor and use of one or more agents that module AMP levels, glucose uptake, metabolism or a stress response in the manufacture of a medicament for treating non- small cell lung carcinoma or cervical cancer or Peutz-Jeghers Syndrome,
wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with

the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

9. A pharmaceutical composition comprising a TOR kinase inhibitor and one or more agents that modulate AMP levels, glucose uptake, metabolism or a stress response, and a pharmaceutically acceptable carrier, excipient or diluent,

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted

heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

10. A kit comprising one or more containers filled with a TOR kinase inhibitor or a pharmaceutical composition thereof, reagents for detecting a LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation, an AMPK gene loss, AMPK protein loss, AMPK gene mutation, or AMPK protein mutation, or both, relative to wild type, in a patient's cancer or in a patient having a tumor syndrome and instructions for detecting LKB1 gene loss, LKB1 protein loss, LKB1 gene mutation, or LKB1 protein mutation, AMPK gene loss, AMPK protein loss, AMPK gene mutation, or AMPK protein mutation, or both, in a patient's cancer or in a patient having a tumor syndrome,

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

11. Use of a TOR kinase inhibitor in the manufacture of a medicament for treating non-small cell lung carcinoma, cervical cancer or Peutz -Jeghers Syndrome, in a patient having non-small cell lung carcinoma, cervical cancer or Peutz- Jeghers Syndrome characterized by an AMPK gene loss, AMPK protein loss, AMPK gene mutation, or AMPK protein mutation, relative to wild type,

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted


heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

12. Use of a TOR kinase inhibitor in the manufacture of a medicament for treating non-small cell lung carcinoma or cervical cancer, in a patient having non-small cell lung carcinoma or cervical cancer characterized by an AMPK gene loss, AMPK protein loss, AMPK gene mutation, or AMPK protein mutation wherein the use comprises screening a patient's carcinoma or cancer for the presence of AMPK gene loss, AMPK protein loss, AMPK gene mutation, or AMPK protein mutation, relative to wild type,

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

13. An *in vitro* method for predicting the likelihood of a patient having non-small cell lung carcinoma or cervical cancer being responsive to TOR kinase inhibitor therapy, comprising screening said patient's carcinoma or cancer for the presence of an AMPK gene loss, AMPK protein loss, AMPK gene mutation, or AMPK protein mutation, relative to wild type, wherein the presence of AMPK gene loss, AMPK protein loss, AMPK gene mutation, or AMPK protein mutation predicts an increased likelihood that TOR kinase inhibitor therapy will treat said non-small cell lung carcinoma or cervical cancer,

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

14. An *in vitro* method for predicting therapeutic efficacy of treatment of a patient having non-small cell lung carcinoma or cervical cancer with a TOR kinase inhibitor, comprising screening said patient's carcinoma or cancer for the presence of an AMPK gene loss, AMPK protein loss, AMPK gene mutation, or AMPK protein mutation, relative to wild type, wherein the presence of AMPK gene loss, AMPK protein loss, AMPK gene mutation, or AMPK protein mutation is predictive of therapeutic efficacy of treatment with a TOR kinase inhibitor,

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

15. Use of a TOR kinase inhibitor in the manufacture of a medicament for treating Peutz-Jeghers Syndrome, in a patient having Peutz-Jeghers Syndrome characterized by an AMPK gene loss, AMPK protein loss, AMPK gene mutation, or AMPK protein mutation comprises screening a patient for the presence of AMPK gene loss, AMPK protein loss, AMPK gene mutation, or AMPK protein mutation, relative to wild type,

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV);



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

16. An *in vitro* method for predicting the likelihood of a patient having Peutz-Jeghers Syndrome being responsive to TOR kinase inhibitor therapy, comprising screening said patient for the presence of an AMPK gene loss, AMPK protein loss, AMPK gene mutation, or AMPK protein mutation, relative to wild type, wherein the presence of AMPK gene loss, AMPK protein loss, AMPK gene mutation, or AMPK protein mutation predicts an increased likelihood that TOR kinase inhibitor therapy will treat said Peutz- Jeghers Syndrome, wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted


heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

17. An *in vitro* method for predicting therapeutic efficacy of treatment of a patient having Peutz-Jeghers Syndrome with a TOR kinase inhibitor, comprising screening said patient for the presence of an AMPK gene loss, AMPK protein loss, AMPK gene mutation, or AMPK protein mutation, relative to wild type, wherein the presence of AMPK gene loss, AMPK protein loss, AMPK gene mutation, or AMPK protein mutation is predictive of therapeutic efficacy of treatment with a TOR kinase inhibitor,

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

18. Use of a TOR kinase inhibitor in the manufacture of a medicament for treating non-small cell lung carcinoma, cervical cancer or Peutz-Jeghers Syndrome, in a patient having non-small cell lung carcinoma, cervical cancer or Peutz-Jeghers Syndrome characterized by a reduced level of pAMPK protein, AMPK activity, or both, relative to wild type,
wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or

unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

19. Use of a TOR kinase inhibitor in the manufacture of a medicament for treating non-small cell lung carcinoma or cervical cancer, in a patient having non-small cell lung carcinoma or cervical cancer characterized by a reduced level of pAMPK protein, AMPK activity, or both wherein the use comprises screening a patient's carcinoma or cancer for the presence of a reduced level of pAMPK protein, AMPK activity, or both, relative to wild type, wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

20. An *in vitro* method for predicting the likelihood of a patient having non- small cell lung carcinoma or cervical cancer being responsive to TOR kinase inhibitor therapy, comprising screening said patient's carcinoma or cancer for the presence of a reduced level of pAMPK protein, AMPK activity, or both, relative to wild type, wherein the presence of a reduced level of pAMPK protein, AMPK activity, or both, predicts an increased likelihood that TOR kinase inhibitor therapy will treat said non-small cell lung carcinoma or cervical cancer,

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

21. An *in vitro* method for predicting therapeutic efficacy of treatment of a patient having non-small cell lung carcinoma or cervical cancer with a TOR kinase inhibitor, comprising screening said patient's carcinoma or cancer for the presence of a reduced level of pAMPK protein, AMPK activity, or both, relative to wild type, wherein the presence of a reduced level of pAMPK protein, AMPK activity, or both, is predictive of therapeutic efficacy of treatment with a TOR kinase inhibitor,

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted


heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

22. Use of a TOR kinase inhibitor in the manufacture of a medicament for treating Peutz-Jeghers Syndrome, in a patient having Peutz-Jeghers Syndrome characterized by a reduced level of pAMPK protein, AMPK activity, or both, wherein the use comprises screening a patient for the presence of a reduced level of pAMPK protein, AMPK activity, or both, relative to wild type,

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

23. An *in vitro* method for predicting the likelihood of a patient having Peutz-Jeghers Syndrome being responsive to TOR kinase inhibitor therapy, comprising screening said patient for the presence of a reduced level of pAMPK protein, AMPK activity, or both, relative to wild type, wherein the presence of a reduced level of pAMPK protein, AMPK activity, or both, predicts an increased likelihood that TOR kinase inhibitor therapy will treat said Peutz-Jeghers Syndrome,

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

24. An *in vitro* method for predicting therapeutic efficacy of treatment of a patient having Peutz-Jeghers Syndrome with a TOR kinase inhibitor, comprising screening said patient for the presence of a reduced level of pAMPK protein, AMPK activity, or both, relative to wild type, wherein the presence of a reduced level of pAMPK protein, AMPK activity, or both, is predictive of therapeutic efficacy of treatment with a TOR kinase inhibitor, and

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or

unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;


R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

25. A kit comprising one or more containers filled with a TOR kinase inhibitor or a pharmaceutical composition thereof, reagents for detecting a reduced level of pAMPK protein, AMPK activity, or both, relative to wild type, in a patient's cancer or in a patient having a tumor syndrome and instructions for detecting a reduced level of pAMPK protein and/or AMPK activity in a patient's cancer or in a patient having a tumor syndrome,

wherein the TOR kinase inhibitor is a compound having the formula (III):



(III)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:


R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4, together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

or a compound having the formula (IV):



(IV)

or a pharmaceutically acceptable salt, clathrate, solvate, stereoisomer, or tautomer thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;

R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl

and wherein the TOR kinase inhibitor is not 6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one, (R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

26. The use of any one of claims 1, 2, 5, 8, 11, 12, 15, 18, 19 or 22; or the method of any one of claims 3, 4, 6, 7, 13, 14, 16, 17, 20, 21, 23 or 24; or the pharmaceutical composition of claim 9; or the kit of claim 10 or 25, wherein the TOR kinase inhibitor is selected from the group consisting of:

6-(1H-pyrrolo[2,3-b]pyridin-3-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(4-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-((trans-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-((cis-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((trans-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-((trans-4-hydroxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((cis-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((trans-4-hydroxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-(cis-4-hydroxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((cis-4-hydroxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-(trans-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-(trans-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-(trans-4-hydroxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-((cis-4-hydroxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-(cis-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-isopropyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-(cis-4-hydroxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-(cis-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-ethyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-(trans-4-hydroxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-isopropyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
4-ethyl-6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(3-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(3-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-(cis-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(3-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-(trans-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
4-(2-methoxyethyl)-6-(4-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(3-(1H-1,2,4-triazol-5-yl)phenyl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
5-(8-(2-methoxyethyl)-6-oxo-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)-4-methylpicolinamide;
3-(6-oxo-8-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)benzamide;
3-(6-oxo-8-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)benzonitrile;
5-(8-(trans-4-methoxycyclohexyl)-6-oxo-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)-4-methylpicolinamide;
6-(1H-imidazo[4,5-b]pyridin-6-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(1H-indazol-6-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
4-((1R,3S)-3-methoxycyclopentyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
4-((1S,3R)-3-methoxycyclopentyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
4-((1R,3R)-3-methoxycyclopentyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
4-((1S,3S)-3-methoxycyclopentyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
4-ethyl-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(1H-pyrrolo[2,3-b]pyridin-5-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(1H-indol-6-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(1H-indol-5-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
4-(((1R,3S)-3-methoxycyclopentyl)methyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
4-(((1S,3R)-3-methoxycyclopentyl)methyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(3-fluoro-2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(3-fluoro-2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
3,3-dimethyl-6-(4-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((1R,3S)-3-methoxycyclopentyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((1S,3R)-3-methoxycyclopentyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(((1S,3S)-3-methoxycyclopentyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(((1R,3R)-3-methoxycyclopentyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((1S,3S)-3-methoxycyclopentyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((1R,3R)-3-methoxycyclopentyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(((1R,3S)-3-methoxycyclopentyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(((1S,3R)-3-methoxycyclopentyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(3-fluoro-4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(3-fluoro-4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7'-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-1'-((tetrahydro-2H-pyran-4-yl)methyl)-1'H-spiro[cyclopentane-1,2'-pyrazino[2,3-b]pyrazin]-3'(4'H)-one;
7'-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-1'-((tetrahydro-2H-pyran-4-yl)methyl)-1'H-spiro[cyclobutane-1,2'-pyrazino[2,3-b]pyrazin]-3'(4'H)-one;
4-(cyclopropylmethyl)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7'-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-1'H-spiro[cyclopentane-1,2'-pyrazino[2,3-b]pyrazin]-3'(4'H)-one;
7'-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-1'H-spiro[cyclobutane-1,2'-pyrazino[2,3-b]pyrazin]-3'(4'H)-one;
7'-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-1'H-spiro[cyclopropane-1,2'-pyrazino[2,3-b]pyrazin]-3'(4'H)-one;
(R)-6-(4-(4H-1,2,4-triazol-3-yl)phenyl)-4-((tetrahydrofuran-2-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
(S)-6-(4-(4H-1,2,4-triazol-3-yl)phenyl)-4-((tetrahydrofuran-2-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(1H-indazol-5-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
4-(6-oxo-8-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)benzamide;
4-(2-methoxyethyl)-3,3-dimethyl-6-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
4-ethyl-3,3-dimethyl-6-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
3,3-dimethyl-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
(R)-6-(6-(1-hydroxyethyl)pyridin-3-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
3,3-dimethyl-6-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)-4-methylpyridin-3-yl)-4-(trans-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)-4-methylpyridin-3-yl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

3,3-dimethyl-6-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
3,3-dimethyl-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)-2-methylpyridin-3-yl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)-2-methylpyridin-3-yl)-4-(trans-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
(S)-6-(6-(1-hydroxyethyl)pyridin-3-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
3,3-dimethyl-6-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,3-dimethyl-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(4-(2-hydroxypropan-2-yl)phenyl)-4-(trans-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(4-(2-hydroxypropan-2-yl)phenyl)-4-((trans-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
4-(cis-4-methoxycyclohexyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
4-(trans-4-methoxycyclohexyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(4-(2-hydroxypropan-2-yl)phenyl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
4-(2-methoxyethyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
9-(6-(4H-1,2,4-triazol-3-yl)-3-pyridyl)-6,11,4a-trihydromorpholino[4,3-e]pyrazino[2,3-b]pyrazin-5-one;
6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
5-(8-(cis-4-methoxycyclohexyl)-6-oxo-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)-6-methylpicolinonitrile;
6-(6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
9-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)-3-(2-methoxyacetyl)-6,11,4a-trihydropiperazino[1,2-e]pyrazino[2,3-b]pyrazin-5-one;
9-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)-6,11,4a-trihydropiperazino[1,2-e]pyrazino[2,3-b]pyrazin-5-one;
9-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)-3-(2-methoxyethyl)-6,11,4a-trihydropiperazino[1,2-e]pyrazino[2,3-b]pyrazin-5-one;
4-(cyclopentylmethyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
9-(6-(4H-1,2,4-triazol-3-yl)-2-methyl-3-pyridyl)-6,11,4a-trihydromorpholino[4,3-e]pyrazino[2,3-b]pyrazin-5-one;
4-(trans-4-hydroxycyclohexyl)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
4-(cis-4-hydroxycyclohexyl)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((tetrahydrofuran-3-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-(cyclopentylmethyl)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-neopentyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-isobutyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
3-methyl-6-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(piperidin-4-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(2-(tetrahydro-2H-pyran-3-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
8-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)(3aS,2R)-2-methoxy-5,10,3a-trihydropyrazino[2,3-b]pyrrolidino[1,2-e]pyrazin-4-one;
8-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)(2R,3aR)-2-methoxy-5,10,3a-trihydropyrazino[2,3-b]pyrrolidino[1,2-e]pyrazin-4-one;
8-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)(2S,3aR)-2-methoxy-5,10,3a-trihydropyrazino[2,3-b]pyrrolidino[1,2-e]pyrazin-4-one;
8-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)(2S,3aS)-2-methoxy-5,10,3a-trihydropyrazino[2,3-b]pyrrolidino[1,2-e]pyrazin-4-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(3-methoxypropyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
(S)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((tetrahydrofuran-2-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
(R)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((tetrahydrofuran-2-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
9-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)-3-methyl-6,11,4a-trihydropiperazino[1,2-e]pyrazino[2,3-b]pyrazin-5-one;
9-(4-(4H-1,2,4-triazol-3-yl)phenyl)-6,11,4a-trihydromorpholino[4,3-e]pyrazino[2,3-b]pyrazin-5-one;
9-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)-6,11,4a-trihydropiperidino[1,2-e]pyrazino[2,3-b]pyrazin-5-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(trans-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(cis-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(2-morpholinoethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-phenethyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
4-(cyclohexylmethyl)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((trans-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((cis-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

(R)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(tetrahydrofuran-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
(S)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(tetrahydrofuran-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-phenyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
(S)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3-methyl-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
9-[6-(1-hydroxy-isopropyl)-3-pyridyl]-6,11,4a-trihydromorpholino[4,3-e]pyrazino[2,3-b]pyrazin-5-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(2-amino-7-methyl-1H-benzo[d]imidazol-5-yl)-4-(3-(trifluoromethyl)benzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(3-(trifluoromethyl)benzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
9-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)-6,11,4a-trihydromorpholino[4,3-e]pyrazino[2,3-b]pyrazin-5-one;
6-(4-methyl-2-(methylamino)-1H-benzo[d]imidazol-6-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
8-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)-5,10,3a-trihydropyrazino[2,3-b]pyrrolidino[1,2-e]pyrazin-4-one;
6-(4-(4H-1,2,4-triazol-3-yl)phenyl)-4-ethyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(4-(4H-1,2,4-triazol-3-yl)phenyl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(3-(trifluoromethyl)benzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(4-methyl-1H-benzo[d]imidazol-6-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
6-(4-(2-hydroxypropan-2-yl)phenyl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one; and
6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one,
and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, and tautomers thereof.

27. The use of any one of claims 1, 2, 5, 8, 11, 12, 15, 18, 19 or 22; or the method of any one of claims 3, 4, 6, 7, 13, 14, 16, 17, 20, 21, 23 or 24; or the pharmaceutical composition of claim 9; or the kit of claim 10 or 25, wherein the TOR kinase inhibitor is selected from the group consisting of:
7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-((*trans*-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(*cis*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(1H-pyrrolo[2,3-b]pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-((*cis*-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-ethyl-7-(1H-pyrrolo[3,2-b]pyridin-5-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-((*cis*-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(1H-benzo[d]imidazol-4-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(1H-pyrrolo[2,3-b]pyridin-4-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-((*trans*-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-((*trans*-4-hydroxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(*cis*-4-hydroxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(*cis*-4-hydroxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-ethyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-((*cis*-4-hydroxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(1H-indol-4-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-((*trans*-4-hydroxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-((*cis*-4-hydroxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(*trans*-4-hydroxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-isopropyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(*trans*-4-hydroxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-isopropyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-ethyl-7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(2-hydroxypyridin-4-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-isopropyl-7-(4-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
5-(8-isopropyl-7-oxo-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)-4-methylpicolinamide;
7-(1H-indazol-4-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(2-aminopyrimidin-5-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(2-aminopyridin-4-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(methylamino)pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-hydroxypyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(4-(1H-pyrazol-3-yl)phenyl)-1-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(1H-indazol-4-yl)-1-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(1H-indazol-6-yl)-1-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(pyrimidin-5-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-methoxypyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-(2-methoxyethyl)-7-(1H-pyrrolo[2,3-b]pyridin-5-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-ethyl-7-(1H-pyrrolo[2,3-b]pyridin-5-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-ethyl-7-(1H-indazol-4-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(pyridin-4-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-aminopyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-methyl-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
2-(2-hydroxypropan-2-yl)-5-(8-(*trans*-4-methoxycyclohexyl)-7-oxo-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)pyridine 1-oxide;
4-methyl-5-(7-oxo-8-((tetrahydro-2H-pyran-4-yl)methyl)-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)picolinamide;
5-(8-((*cis*-4-methoxycyclohexyl)methyl)-7-oxo-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)-4-methylpicolinamide;
7-(1H-pyrazol-4-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-(*trans*-4-methoxycyclohexyl)-7-(4-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

3-((7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-2-oxo-3,4-dihydropyrazino[2,3-b]pyrazin-1(2H)-yl)methyl)benzonitrile;
1-((*trans*-4-methoxycyclohexyl)methyl)-7-(4-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
3-(7-oxo-8-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)benzamide;
5-(8-((*trans*-4-methoxycyclohexyl)methyl)-7-oxo-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)-4-methylpicolinamide;
3-((7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-2-oxo-3,4-dihydropyrazino[2,3-b]pyrazin-1(2H)-yl)methyl)benzonitrile;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-((1R,3R)-3-methoxycyclopentyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-((1S,3R)-3-methoxycyclopentyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-((1S,3S)-3-methoxycyclopentyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-((1R,3S)-3-methoxycyclopentyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(1H-indazol-6-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(2-morpholinoethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-(*trans*-4-hydroxycyclohexyl)-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-(*cis*-4-hydroxycyclohexyl)-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(2-morpholinoethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-isopropyl-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(1H-imidazo[4,5-b]pyridin-6-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-((*cis*-4-methoxycyclohexyl)methyl)-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-(*trans*-4-hydroxycyclohexyl)-7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-(*cis*-4-hydroxycyclohexyl)-7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
4-(7-oxo-8-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)benzamide;
7-(1H-indazol-5-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(1H-pyrrolo[2,3-b]pyridin-5-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-((1S,3R)-3-methoxycyclopentyl)-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-((1R,3R)-3-methoxycyclopentyl)-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

1-((1R,3S)-3-methoxycyclopentyl)-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-((1S,3S)-3-methoxycyclopentyl)-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(1H-indol-5-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-ethyl-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(1H-indol-6-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(4-(2-hydroxypropan-2-yl)phenyl)-1-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-((*trans*-4-methoxycyclohexyl)methyl)-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-((*cis*-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-(2-methoxyethyl)-7-(4-methyl-2-(methylamino)-1H-benzo[d]imidazol-6-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(7-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-(2-methoxyethyl)-7-(4-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-benzyl-7-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(3-fluoro-4-(4H-1,2,4-triazol-3-yl)phenyl)-1-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(3-fluoro-4-(4H-1,2,4-triazol-3-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(3-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-(*trans*-4-methoxycyclohexyl)-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(5-fluoro-2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(3-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-(2-methoxyethyl)-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-((*trans*-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-(cyclopentylmethyl)-7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(4-(2-hydroxypropan-2-yl)phenyl)-1-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

(S)-7-(6-(1-hydroxyethyl)pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
(R)-7-(6-(1-hydroxyethyl)pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(4-(2-hydroxypropan-2-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(4-(trifluoromethyl)benzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(3-(trifluoromethyl)benzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(3-methoxypropyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(4-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(4-methyl-2-(methylamino)-1H-benzo[d]imidazol-6-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(2-amino-4-methyl-1H-benzo[d]imidazol-6-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
(R)-7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3-methyl-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
(S)-7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3-methyl-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,3-dimethyl-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(2-amino-4-methyl-1H-benzo[d]imidazol-6-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(4-(1H-1,2,4-triazol-5-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
1-(1-hydroxypropan-2-yl)-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one; and
1-(2-hydroxyethyl)-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one,
and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, and tautomers thereof.