

C7-FLUORO SUBSTITUTED TETRACYCLINE COMPOUNDS

RELATED APPLICATION

This application claims the benefit of U.S. Provisional Application No. 61/188,307, filed on August 8, 2008. The entire teachings of the above application are incorporated herein by reference.


BACKGROUND OF THE INVENTION

The tetracyclines are broad spectrum anti-microbial agents that are widely used in human and veterinary medicine. The total production of tetracyclines by fermentation or semi-synthesis is measured in the thousands of metric tons per year.

The widespread use of tetracyclines for therapeutic purposes has led to the emergence of resistance to these antibiotics, even among highly susceptible bacterial species. Therefore, there is need for new tetracycline analogs with improved antibacterial activities and efficacies against other tetracycline responsive diseases or disorders.

SUMMARY OF THE INVENTION

The present invention is directed to a compound represented by Structural Formula (A):



or a pharmaceutically acceptable salt thereof, wherein:

X is selected from hydrogen, -(C₁-C₇)alkyl, carbocyclyl, aryl and heteroaryl;

Y is selected from hydrogen, -(C₁-C₇)alkyl, carbocyclyl,

-(C₁-C₄)alkylene-N(R^A)(R^B),


-(C₁-C₄)alkylene-N(R^F)-C(O)-[C(R^D)(R^E)]₀₋₄-N(R^A)(R^B), -CH=N-OR^A, -N(R^A)(R^B),

- N(R^F)-C(O)-[C(R^D)(R^E)]₁₋₄-N(R^A)(R^B), -N(R^F)-C(O)-N(R^A)(R^B),
 -N(R^F)-C(O)-(C₁-C₆)alkyl, -N(R^F)-C(O)-heterocyclyl, -N(R^F)-C(O)-heteroaryl,
 -N(R^F)-C(O)-carbocyclyl, -N(R^F)-C(O)-aryl,
 -N(R^F)-S(O)_m-(C₁-C₄)alkylene-N(R^A)(R^B),
 5 -N(R^F)-S(O)_m-(C₁-C₄)alkylene-carbocyclyl, and -N(R^F)-S(O)_m-(C₁-C₄)alkylene-aryl;
 at least one of X and Y is not hydrogen;
 each R^A and R^B are independently selected from hydrogen, (C₁-C₇)alkyl,
 -O-(C₁-C₇)alkyl, -(C₀-C₆)alkylene-carbocyclyl, -(C₀-C₆)alkylene-aryl,
 -(C₀-C₆)alkylene-heterocyclyl, -(C₀-C₆)alkylene-heteroaryl,
 10 -(C₁-C₆)alkylene-O-carbocyclyl, -(C₁-C₆)alkylene-O-aryl,
 -(C₁-C₆)alkylene-O-heterocyclyl, -(C₁-C₆)alkylene-O-heteroaryl,
 -S(O)_m-(C₁-C₆)alkyl, -(C₀-C₄)alkylene-S(O)_m-carbocyclyl,
 -(C₀-C₄)alkylene-S(O)_m-aryl, -(C₀-C₄)alkylene-S(O)_m-heterocyclyl and
 -(C₀-C₄)alkylene-S(O)_m-heteroaryl; or
 15 R^A and R^B taken together with the nitrogen atom to which they are bound
 form a heterocyclyl or heteroaryl, wherein the heterocycle or heteroaryl optionally
 comprises 1 to 4 additional heteroatoms independently selected from N, S and O;
 each R^D and each R^E is independently selected from hydrogen, (C₁-C₆)alkyl,
 carbocyclyl, aryl, heterocyclyl, heteroaryl, or a naturally occurring amino acid side
 20 chain moiety, or
 R^D and R^E taken together with the carbon atom to which they are bound form
 a 3-7 membered carbocyclyl, or a 4-7 membered heterocyclyl, wherein the
 heterocyclyl formed by R^D and R^E optionally comprises one to two additional
 heteroatoms independently selected from N, S and O;
 25 R^F is selected from hydrogen, (C₁-C₇)alkyl, carbocyclyl, aryl and heteroaryl;
 and
 m is 1 or 2, wherein:
 each carbocyclyl, aryl, heterocyclyl or heteroaryl is optionally and
 independently substituted with one or more substituents independently selected from
 30 halo, -(C₁-C₄)alkyl, -OH, =O, -O-(C₁-C₄)alkyl, -(C₁-C₄)alkyl-O-(C₁-C₄)alkyl,
 halo-substituted -(C₁-C₄)alkyl, halo-substituted -O-(C₁-C₄)alkyl, -C(O)-(C₁-C₄)alkyl,
 -C(O)-(fluoro-substituted-(C₁-C₄)alkyl), -S(O)_m-(C₁-C₄)alkyl, -N(R^G)(R^H), and CN;

each alkyl in the group represented by R^A , R^B , R^D and R^E is optionally and independently substituted with one or more substituents independently selected from halo, $-(C_1-C_4)alkyl$, $-OH$, $-O-(C_1-C_7)alkyl$, $-(C_1-C_4)alkyl-O-(C_1-C_4)alkyl$, fluoro-substituted- $(C_1-C_4)alkyl$, $-S(O)_m-(C_1-C_4)alkyl$, and $-N(R^G)(R^G)$, wherein

5 each R^G is hydrogen or $(C_1-C_4)alkyl$, wherein each alkyl in the group represented by R^G is optionally and independently substituted with one or more substituents independently selected from $-(C_1-C_4)alkyl$, $(C_3-C_6)cycloalkyl$, halo, $-OH$, $-O-(C_1-C_4)alkyl$, and $(C_1-C_4)alkyl-O-(C_1-C_4)alkyl$.

Another embodiment of the present invention is directed to a compound
 10 represented by Structural Formula (II)



or a pharmaceutically acceptable salt thereof, wherein:

R^1 and R^2 are each independently selected from hydrogen, $(C_1-C_7)alkyl$, $(C_3-C_6)cycloalkyl$, $(C_1-C_4)alkyl$, $(C_1-C_7)alkoxy$, $(C_1-C_4)alkyl$, $(C_3-C_6)cycloalkoxy$, $(C_1-C_4)alkyl$, $(C_3-C_6)cycloalkyl$, aryl, aryl- $(C_1-C_4)alkyl$, aryl- oxy , $(C_1-C_4)alkyl$, arylthio- $(C_1-C_4)alkyl$, arylsulfinyl- $(C_1-C_4)alkyl$, arylsulfonyl- $(C_1-C_4)alkyl$, and $-O-(C_1-C_7)alkyl$, or

15 R^1 and R^2 taken together with the nitrogen atom to which they are bonded form a monocyclic or bicyclic heteroaryl, or a monocyclic, fused bicyclic, bridged bicyclic or spiro bicyclic heterocycle, wherein the heteroaryl or heterocycle optionally contains one or two additional heteroatoms independently selected from N, O and S; and

wherein each alkyl, cycloalkyl, alkoxy and cycloalkoxy moiety in the groups represented by R^1 and R^2 and each heterocycle represented by NR^1R^2 taken together

25 is optionally substituted with one or more substituents independently selected from the group consisting of $(C_1-C_4)alkyl$, halo, $-OH$, $(C_1-C_4)alkoxy$, $(C_1-C_4)alkylthio$, $(C_1-C_4)alkylsulfinyl$, $(C_1-C_4)alkylsulfonyl$, $(C_1-C_4)alkoxy$, $(C_1-C_4)alkyl$, and $-N(R^3)(R^4)$; and


each aryl, aryloxy, arylthio, arylsulfinyl and arylsulfonyl moiety in the groups represented by R^1 and R^2 and each heteroaryl represented by NR^1R^2 taken together is optionally substituted with one or more substituents independently selected from the group consisting of (C₁-C₄)alkyl, halo, -OH, (C₁-C₄)alkoxy, -S-(C₁-C₄)alkyl,

5 -S(O)(C₁-C₄)alkyl, -S(O)₂(C₁-C₄)alkyl, (C₁-C₄)alkoxy(C₁-C₄)alkyl, -N(R³)(R⁴), -CN, halo(C₁-C₄)alkyl, and halo(C₁-C₄)alkoxy, and

R^3 and R^4 are each independently selected from the group consisting of -H and (C₁-C₄)alkyl, wherein the (C₁-C₄)alkyl represented by R^3 and R^4 is optionally substituted with one or more substituents independently selected from the group

10 consisting of (C₁-C₄)alkyl, halo, -OH, (C₁-C₄)alkoxy, and (C₁-C₄)alkoxy(C₁-C₄)alkyl. Values for X and R^V are as described above for Structural Formula (A).

Another embodiment of the present invention is directed to a compound represented by Structural Formula (I):



or a pharmaceutically acceptable salt thereof, wherein:

R^1 and R^2 are each independently selected from hydrogen, (C₁-C₇)alkyl, (C₃-C₆)cycloalkyl(C₁-C₄)alkyl, (C₁-C₇)alkoxy(C₁-C₄)alkyl, (C₃-C₆)cycloalkoxy(C₁-C₄)alkyl, (C₃-C₆)cycloalkyl, aryl, aryl(C₁-C₄)alkyl, 20 aryloxy(C₁-C₄)alkyl, arylthio(C₁-C₄)alkyl, arylsulfinyl(C₁-C₄)alkyl, and arylsulfonyl(C₁-C₄)alkyl, or

R^1 and R^2 taken together with the nitrogen atom to which they are bonded form a monocyclic or bicyclic heteroaryl, or a monocyclic, fused bicyclic, bridged bicyclic or spiro bicyclic heterocycle, wherein the heteroaryl or heterocycle 25 optionally contains one additional heteroatom independently selected from N, O and S; and the remainder of the variables are as described above for Structural Formula (II).

Another embodiment of the present invention is directed to a pharmaceutical composition comprising a pharmaceutically acceptable carrier or diluent and a compound represented by Structural Formula (A), (II) or (I) or a pharmaceutically acceptable salt thereof. The pharmaceutical composition is used in therapy, such as treating an infection in a subject.

Another embodiment of the present invention is a method of treating an infection in a subject comprising administering to the subject an effective amount of a compound represented by Structural Formula (A), (II) or (I) or a pharmaceutically acceptable salt thereof.

Another embodiment of the present invention is a method of preventing an infection in a subject comprising administering to the subject an effective amount of a compound represented by Structural Formula (A), (II) or (I) or a pharmaceutically acceptable salt thereof.

Another embodiment of the present invention is the use of a compound represented by Structural Formula (A), (II) or (I) or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for treating an infection in a subject.

Another embodiment of the present invention is the use of a compound represented by Structural Formula (A), (II) or (I) or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for preventing an infection in a subject.

Another embodiment of the present invention is the use of a compound represented by Structural Formula (A), (II) or (I) or a pharmaceutically acceptable salt thereof for therapy, such as treating or preventing an infection in a subject.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to a compound represented by Structural Formula (A) or a pharmaceutically acceptable salt thereof. Values and alternative values for the variables in Structural Formula (A) are defined as the following:

X is selected from hydrogen, -(C₁-C₇)alkyl, carbocyclyl, aryl and heteroaryl.

In one embodiment, X is hydrogen. In an alternative embodiment, X is -(C₁-C₇)alkyl. Alternatively, X is -(C₁-C₄)alkyl. In another embodiment, X is

carbocyclyl. In another alternative embodiment, X is aryl or heteroaryl. In another alternative embodiment, X is phenyl.

Y is selected from hydrogen, $-(C_1-C_7)\text{alkyl}$, carbocyclyl,

$-(C_1-C_4)\text{alkylene-N}(R^A)(R^B)$,

5 $-(C_1-C_4)\text{alkylene-N}(R^F)-C(O)-[C(R^D)(R^E)]_{0-4}-N(R^A)(R^B)$, $-\text{CH=N-OR}^A$, $-N(R^A)(R^B)$,

$-N(R^F)-C(O)-[C(R^D)(R^E)]_{1-4}-N(R^A)(R^B)$, $-N(R^F)-C(O)-N(R^A)(R^B)$,

$-N(R^F)-C(O)-(C_1-C_6)\text{alkyl}$, $-N(R^F)-C(O)\text{-heterocyclyl}$, $-N(R^F)-C(O)\text{-heteroaryl}$,

$-N(R^F)-C(O)\text{-carbocyclyl}$, $-N(R^F)-C(O)\text{-aryl}$,

$-N(R^F)-S(O)_m-(C_1-C_4)\text{alkylene-N}(R^A)(R^B)$,

10 $-N(R^F)-S(O)_m-(C_1-C_4)\text{alkylene-carbocyclyl}$, and $-N(R^F)-S(O)_m-(C_1-C_4)\text{alkylene-aryl}$,

provided at least one of X and Y is not hydrogen. In one embodiment, Y is selected

from hydrogen, $-(C_1-C_7)\text{alkyl}$, $-(C_1-C_4)\text{alkylene-N}(R^A)(R^B)$,

$-(C_1-C_4)\text{alkylene-N}(R^F)-C(O)-[C(R^D)(R^E)]_{0-4}-N(R^A)(R^B)$, $-\text{CH=N-OR}^A$, $-N(R^A)(R^B)$,

$-N(R^F)-C(O)-[C(R^D)(R^E)]_{1-4}-N(R^A)(R^B)$, $-N(R^F)-C(O)-N(R^A)(R^B)$,

15 $-N(R^F)-C(O)-(C_1-C_6)\text{alkyl}$, $-N(R^F)-C(O)\text{-heterocyclyl}$, $-N(R^F)-C(O)\text{-heteroaryl}$,

$-N(R^F)-C(O)\text{-carbocyclyl}$, $-N(R^F)-C(O)\text{-aryl}$,

$-N(R^F)-S(O)_m-(C_1-C_4)\text{alkylene-N}(R^A)(R^B)$,

$-N(R^F)-S(O)_m-(C_1-C_4)\text{alkylene-carbocyclyl}$, and $-N(R^F)-S(O)_m-(C_1-C_4)\text{alkylene-aryl}$,

provided at least one of X and Y is not hydrogen. In one embodiment, Y is selected

20 from hydrogen, $-(C_1-C_7)\text{alkyl}$, $-(C_1-C_4)\text{alkylene-N}(R^A)(R^B)$,

$-(C_1-C_4)\text{alkylene-N}(R^F)-C(O)-[C(R^D)(R^E)]_{0-4}-N(R^A)(R^B)$, $-\text{CH=N-OR}^A$, $-N(R^A)(R^B)$,

$-N(R^F)-C(O)-[C(R^D)(R^E)]_{1-4}-N(R^A)(R^B)$, $-\text{NH-C(O)-C}(R^{D'}) (R^E)-N(R^A)(R^B)$,

$-N(R^F)-C(O)-N(R^A)(R^B)$, $-N(R^F)-C(O)-(C_1-C_6)\text{alkyl}$, $-N(R^F)-C(O)\text{-heterocyclyl}$,

$-N(R^F)-C(O)\text{-heteroaryl}$, $-N(R^F)-C(O)\text{-carbocyclyl}$, $-N(R^F)-C(O)\text{-aryl}$

25 $-N(R^F)-S(O)_m-(C_1-C_4)\text{alkylene-N}(R^A)(R^B)$,

$-N(R^F)-S(O)_m-(C_1-C_4)\text{alkylene-carbocyclyl}$, and $-N(R^F)-S(O)_m-(C_1-C_4)\text{alkylene-aryl}$.

In another embodiment, Y is selected from hydrogen, $-(C_1-C_7)\text{alkyl}$,

$-(C_1-C_4)\text{alkylene-N}(R^A)(R^B)$, $-(C_1-C_4)\text{alkylene-NH-C(O)-(CH}_2\text{)}_{0-1}-N(R^A)(R^B)$,

$-N(R^A)(R^B)$, $-\text{NH-C(O)-carbocyclyl}$, $-\text{NH-C(O)-aryl}$, $-\text{NH-C(O)-heterocyclyl}$,

30 $-\text{NH-C(O)-heteroaryl}$, $-\text{NH-C(O)-N}(R^A)(R^A)$, $-N(R^F)-C(O)-\text{CH}_2-N(R^A)(R^B)$,

$-\text{NH-C(O)-C}(R^{D'}) (R^E)-N(R^A)(R^B)$ and $-\text{NH-S(O)}_m-(C_1-C_4)\text{alkylene-N}(R^A)(R^B)$.

Alternatively, the $-(C_1-C_7)\text{alkyl}$ represented by Y described above is a $-(C_1-C_4)\text{alkyl}$.

In yet another embodiment, Y is selected from $-N(R^A)(R^B)$,

-N(H)-C(O)-carbocyclyl, -N(H)-C(O)-aryl, -N(H)-C(O)-heterocycle, and
-N(H)-C(O)-heteroaryl. Alternatively, Y is -NH-C(O)-CH₂-N(R^A)(R^B). More
specifically, R^A and R^B in -NH-C(O)-CH₂-N(R^A)(R^B) are R¹ and R², respectively.

Each R^A and R^B are independently selected from hydrogen, (C₁-C₇)alkyl,
5 -O-(C₁-C₇)alkyl, -(C₀-C₆)alkylene-carbocyclyl, -(C₀-C₆)alkylene-aryl,
-(C₀-C₆)alkylene-heterocyclyl, -(C₀-C₆)alkylene-heteroaryl,
-(C₁-C₆)alkylene-O-carbocyclyl, -(C₁-C₆)alkylene-O-aryl,
-(C₁-C₆)alkylene-O-heterocyclyl, -(C₁-C₆)alkylene-O-heteroaryl,
-S(O)_m-(C₁-C₆)alkyl, -(C₀-C₄)alkylene-S(O)_m-carbocyclyl,
10 -(C₀-C₄)alkylene-S(O)_m-aryl, -(C₀-C₄)alkylene-S(O)_m-heterocyclyl and
-(C₀-C₄)alkylene-S(O)_m-heteroaryl; or
R^A and R^B taken together with the nitrogen atom to which they are bound form a
heterocyclyl or heteroaryl, wherein the heterocycle or heteroaryl optionally
comprises 1 to 4 additional heteroatoms independently selected from N, S and O. In
15 one embodiment, each R^A is independently selected from hydrogen and methyl; R^B
is selected from hydrogen, (C₁-C₇)alkyl, -(C₀-C₆)alkylene-carbocyclyl,
-(C₀-C₆)alkylene-aryl, -(C₀-C₆)alkylene-heteroaryl, -S(O)_m-(C₁-C₆)alkyl,
-(C₀-C₄)alkylene-S(O)_m-carbocyclyl, -(C₀-C₄)alkylene-S(O)_m-aryl,
-(C₀-C₄)alkylene-S(O)_m-heterocycle and -(C₀-C₄)alkylene-S(O)_m-heteroaryl; or R^A
20 and R^B taken together with the nitrogen atom to form a heterocycle, wherein the
heterocycle is optionally substituted with -O and -N(R^G)(R^G). In another
embodiment, R^A is hydrogen; and R^B is selected from (C₁-C₄)alkyl, and -S(O)₂-CH₃;
or R^A and R^B taken together to form 4-7 membered heterocyclic ring.

Each R^D and each R^E is independently selected from hydrogen, (C₁-C₆)alkyl,
25 carbocyclyl, aryl, heterocyclyl, heteroaryl, or a naturally occurring amino acid side
chain moiety, or R^D and R^E taken together with the carbon atom to which they are
bound form a 3-7 membered carbocyclyl, or a 4-7 membered heterocyclyl, wherein
the heterocyclyl formed by R^D and R^E optionally comprises one to two additional
heteroatoms independently selected from N, S and O. In one embodiment, R^D and
30 R^E are both -H.

R^F is selected from hydrogen, (C₁-C₇)alkyl, carbocyclyl, aryl and heteroaryl.
In one embodiment, R^F is hydrogen. In another embodiment, R^F is selected from
hydrogen, (C₁-C₇)alkyl, aryl and heteroaryl. In another embodiment, R^F is selected

from hydrogen, (C₁-C₇)alkyl and phenyl. In another embodiment, R^F is selected from hydrogen, (C₁-C₄)alkyl and phenyl

R^{D'} is selected from (C₁-C₆)alkyl, carbocyclyl, aryl, heteroaryl, heterocyclyl, and a naturally occurring amino acid side chain moiety, or R^{D'} and R^E taken together with the carbon atom to which they are bound form a 3-7 membered carbocyclyl, or a 4-7 membered heterocyclyl, wherein the heterocyclyl formed by R^{D'} and R^E optionally comprises one to two additional heteroatoms independently selected from N, S and O. In one embodiment, R^{D'} and R^E taken together with the carbon atom to which they are bound form a (C₃-C₇)cycloalkyl.

10 m is 1 or 2. In one embodiment, m is 2.

Each carbocyclyl, aryl, heterocyclyl or heteroaryl described above (*e.g.*, in the groups represented by Y, R^A, R^B, R^D, R^{D'} and R^E) is optionally and independently substituted with one or more substituents independently selected from halo,


-(C₁-C₄)alkyl, -OH, =O, -O-(C₁-C₄)alkyl, -(C₁-C₄)alkyl-O-(C₁-C₄)alkyl, halo-substituted -(C₁-C₄)alkyl, halo-substituted -O-(C₁-C₄)alkyl, -C(O)-(C₁-C₄)alkyl, -C(O)-(fluoro-substituted-(C₁-C₄)alkyl), -S(O)_m-(C₁-C₄)alkyl, -N(R^G)(R^G), and CN. In one embodiment, each carbocyclyl, heterocyclyl or heteroaryl is optionally and independently substituted with one or more substituents independently selected from halo, -(C₁-C₄)alkyl, halo-substituted -(C₁-C₄)alkyl, -O-(C₁-C₄)alkyl, and -N(R^G)(R^G).
15 In another embodiment, each carbocyclyl, aryl, heteroaryl, or heterocycle is optionally and independently substituted with one or more substituents independently selected from -CH₃, fluoro, and -N(CH₃)₂.

Each alkyl described above (*e.g.*, in the groups represented by Y, R^A, R^B, R^D, R^{D'}, R^E, R^F and R^{F'}) is optionally and independently substituted with one or more substituents independently selected from halo, -(C₁-C₄)alkyl, -OH, -O-(C₁-C₇)alkyl, -(C₁-C₄)alkyl-O-(C₁-C₄)alkyl, fluoro-substituted-(C₁-C₄)alkyl, -S(O)_m-(C₁-C₄)alkyl, and -N(R^G)(R^G). In one embodiment, each alkyl group (*e.g.*, in the group represented by Y or R^B) is optionally and independently substituted with one or more substituents independently selected from halo, -OH, and -N(R^G)(R^G).
25

Each R^G is hydrogen or (C₁-C₄)alkyl, wherein each alkyl in the group represented by R^G is optionally and independently substituted with one or more substituents independently selected from -(C₁-C₄)alkyl, (C₃-C₆)cycloalkyl, halo, -OH, -O-(C₁-C₄)alkyl, and (C₁-C₄)alkyl-O-(C₁-C₄)alkyl. In one embodiment, each
30

alkyl in the group represented by R^G is optionally and independently substituted with (C₃-C₆)cycloalkyl.

As used herein, when R^A and R^B taken together with the nitrogen atom to which they are bound form a heterocyclyl or heteroaryl, the heterocyclyl or heteroaryl represented by $-NR^A R^B$ can include a ring system that has a heteroatom adjacent to the nitrogen atom to which R^A and R^B are bound. For example, $-NR^A R^B$ can be, but is not limited to, the following ring systems:



Similarly, when R^D and R^E or $R^{D'}$ and R^E taken together with the carbon atom to which they are bound form a heterocyclyl, the heterocyclyl can include a ring system that has a heteroatom adjacent to the carbon atom to which R^D and R^E or $R^{D'}$ and R^E are bound.

The present invention is directed to a compound represented by Structural Formula (I) or (II) or a pharmaceutically acceptable salt thereof. Values and alternative values for the variables in Structural Formula (I) or (II) are defined as the following:

R^1 is selected from hydrogen, (C₁-C₇)alkyl, (C₃-C₆)cycloalkyl, (C₁-C₄)alkyl, (C₁-C₇)alkoxy, (C₁-C₄)alkyl, (C₃-C₆)cycloalkoxy, (C₁-C₄)alkyl, (C₃-C₆)cycloalkyl, aryl, aryl(C₁-C₄)alkyl, aryloxy, (C₁-C₄)alkyl, arylthio, (C₁-C₄)alkyl, arylsulfinyl, (C₁-C₄)alkyl, arylsulfonyl, (C₁-C₄)alkyl and -O-(C₁-C₇)alkyl. Each alkyl, cycloalkyl, alkoxy, cycloalkoxy, aryl, aryloxy, arylthio, arylsulfinyl and arylsulfonyl moiety in the groups represented by R^1 can be optionally substituted with one or more independently selected substituents defined above for Structural Formula (I). Alternatively, R^1 is selected from hydrogen, (C₁-C₇)alkyl, (C₃-C₆)cycloalkyl, (C₁-C₄)alkyl, (C₁-C₇)alkoxy, (C₁-C₄)alkyl, (C₃-C₆)cycloalkoxy, (C₁-C₄)alkyl, (C₃-C₆)cycloalkyl, aryl, aryl(C₁-C₄)alkyl, aryloxy, (C₁-C₄)alkyl, arylthio, (C₁-C₄)alkyl, arylsulfinyl, (C₁-C₄)alkyl and arylsulfonyl, (C₁-C₄)alkyl. In another alternative, R^1 is -H, (C₁-C₇)alkyl, or -O-(C₁-C₄)alkyl. In another alternative, R^1 is -H or (C₁-C₇)alkyl. In another alternative, R^1 is -H, methyl or ethyl. In yet another alternative, R^1 is -OCH₃ or -OC(CH₃)₃.

R^2 is selected from hydrogen, (C₁-C₇)alkyl, (C₃-C₆)cycloalkyl(C₁-C₄)alkyl, (C₁-C₇)alkoxy(C₁-C₄)alkyl, (C₃-C₆)cycloalkoxy(C₁-C₄)alkyl, (C₃-C₆)cycloalkyl, aryl, aryl(C₁-C₄)alkyl, aryloxy(C₁-C₄)alkyl, arylthio(C₁-C₄)alkyl, arylsulfinyl(C₁-C₄)alkyl, arylsulfonyl(C₁-C₄)alkyl and -O-(C₁-C₇)alkyl. Each alkyl, cycloalkyl, alkoxy, cycloalkoxy, aryl, and aryloxy moiety in the groups represented by R^2 can be optionally substituted with one or more independently selected substituents defined above for Structural Formula (I). Alternatively, R^2 is selected from hydrogen, (C₁-C₇)alkyl, (C₃-C₆)cycloalkyl(C₁-C₄)alkyl, (C₁-C₇)alkoxy(C₁-C₄)alkyl, (C₃-C₆)cycloalkoxy(C₁-C₄)alkyl, (C₃-C₆)cycloalkyl, aryl, aryl(C₁-C₄)alkyl, aryloxy(C₁-C₄)alkyl, arylthio(C₁-C₄)alkyl, arylsulfinyl(C₁-C₄)alkyl and arylsulfonyl(C₁-C₄)alkyl. Alternatively, R^2 is selected from (C₁-C₇)alkyl, (C₃-C₆)cycloalkyl(C₁-C₄)alkyl, (C₁-C₇)alkoxy(C₁-C₄)alkyl, phenyl, phenyl(C₁-C₄)alkyl, (C₃-C₆)cycloalkyl and halo(C₁-C₄)alkyl, wherein each alkyl, alkoxy and cycloalkyl moiety in the groups represented by R^2 is optionally substituted with one or more substituents independently selected from the group consisting of (C₁-C₄)alkyl and halo; and each phenyl moiety in the groups represented by R^2 is optionally substituted with one or more substituents independently selected from the group consisting of (C₁-C₄)alkyl, halo, (C₁-C₄)alkoxy, (C₁-C₄)alkoxy(C₁-C₄)alkyl, -CN, halo(C₁-C₄)alkyl, and halo(C₁-C₄)alkoxy. In another alternative, R^2 is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclopropylmethyl, cyclobutylmethyl, phenyl, benzyl, -(CH₂)₂-O-CH₃, -(CH₂)₃-OCH₃, -C(CH₃)₃, -CH(CH₃)₂, -CH₂C(CH₃)₃, -CH₂CH(CH₃)₂, -CH₂-CF₃, -(CH₂)₂-CH₂F, and -(CH₂)_nCH₃, wherein n is 0, 1, 2, 3, 4, 5 or 6; and wherein the phenyl or benzyl group represented by R^2 is optionally substituted with one or more substituents independently selected from the group consisting of (C₁-C₄)alkyl, halo, (C₁-C₄)alkoxy, (C₁-C₄)alkoxy(C₁-C₄)alkyl, -CN, halo(C₁-C₄)alkyl, and halo(C₁-C₄)alkoxy. In another alternative, R^2 is a phenyl or benzyl group optionally substituted with one or more substituents independently selected from the group consisting of (C₁-C₄)alkyl, halo, (C₁-C₄)alkoxy, (C₁-C₄)alkoxy(C₁-C₄)alkyl, -CN, halo(C₁-C₄)alkyl, and halo(C₁-C₄)alkoxy. In another alternative, R^2 is unsubstituted phenyl or benzyl. In another alternative, R^2 is selected from cyclopropyl, cyclopropylmethyl, cyclobutyl, cyclopentyl,

cyclohexyl, $-(CH_2)_2-O-CH_3$, $-C(CH_3)_3$, $-CH(CH_3)_2$, $-CH_2-CF_3$, $-CH_2CH(CH_3)_2$, $-CH_3$ and $-CH_2CH_3$.

- Alternatively, R^1 and R^2 taken together with the nitrogen atom to which they are bonded can also form a monocyclic or bicyclic heteroaryl, or a monocyclic, fused bicyclic, bridged bicyclic or spiro bicyclic heterocycle, wherein the heteroaryl or heterocycle optionally contains one or two additional heteroatoms independently selected from N, O and S in addition to the N atom to which R^1 and R^2 are bonded. The heteroaryl or heterocycle can be optionally substituted with one or more independently selected substituents described above for Structural Formula (I).
- Alternatively, the heteroaryl or heterocycle contains one additional heteroatom selected from N, O and S. Alternatively, R^1 and R^2 taken together with the nitrogen atom to which they are bonded form a heterocycle selected from the group consisting of azetidine, pyrrolidine, morpholine, piperidine, octahydrocyclopenta[c]pyrrol, isoindoline, and azabicyclo[3.1.0]hexane, wherein the heterocycle is optionally substituted with one or more substituents independently selected from the group consisting of (C_1-C_4) alkyl, halogen, $-OH$, (C_1-C_4) alkoxy, (C_1-C_4) alkylthio, (C_1-C_4) alkylsulfinyl, (C_1-C_4) alkylsulfonyl, (C_1-C_4) alkoxy (C_1-C_4) alkyl, and $-N(R^3)(R^4)$. In a more specific embodiment, these heterocycles are optionally substituted with one or more substituents independently selected from the group consisting of halogen, (C_1-C_4) alkoxy, hydroxy, (C_1-C_4) alkoxy (C_1-C_4) alkyl and $-N(R^3)(R^4)$. In another alternative embodiment, these heterocycles are optionally substituted with one or more substituents independently selected from the group consisting of halogen, methoxy, hydroxy, methoxymethyl and dimethylamino. Alternatively, R^1 and R^2 taken together with the nitrogen atom to which they are bonded form a ring selected from pyrrolidinyl, morpholinyl, azetidiny, piperidinyl, octahydrocyclopenta[c]pyrrolyl, isoindolinyl, indazolyl, imidazolyl, pyrazolyl, triazolyl, and tetrazolyl, wherein the ring formed by R^1 and R^2 taken together with the nitrogen atom to which they are bonded is optionally substituted with halogen, (C_1-C_4) alkoxy, hydroxy, (C_1-C_4) alkoxy (C_1-C_4) alkyl and $-N(R^3)(R^4)$. More specifically, the ring formed by R^1 and R^2 taken together with the nitrogen atom to which they are bonded is optionally substituted with fluoro, $-OH$, $-OCH_3$, or $N(CH_3)_2$.

R^3 and R^4 are each independently selected from the group consisting of -H and (C₁-C₄)alkyl, wherein the (C₁-C₄)alkyl represented by R^3 and R^4 is optionally substituted with one or more substituents independently selected from the group consisting of (C₁-C₄)alkyl, halo, -OH, (C₁-C₄)alkoxy, and

5 (C₁-C₄)alkoxy(C₁-C₄)alkyl. Alternatively, R^3 and R^4 are both methyl. In another alternative, R^3 and R^4 are both -H. In yet another alternative, R^3 and R^4 are each unsubstituted (C₁-C₄)alkyl.

In a first alternative embodiment, the compound of the present invention is represented by Structural Formula (I) or (II), or a pharmaceutically acceptable salt thereof, wherein:

R^1 is -H or (C₁-C₇)alkyl; and

R^2 is selected from (C₁-C₇)alkyl, (C₃-C₆)cycloalkyl(C₁-C₄)alkyl, (C₁-C₇)alkoxy(C₁-C₄)alkyl, phenyl, phenyl(C₁-C₄)alkyl, (C₃-C₆)cycloalkyl and halo(C₁-C₄)alkyl, wherein each alkyl, alkoxy or cycloalkyl moiety in the groups
15 represented by R^2 is optionally substituted with one or more substituents independently selected from the group consisting of (C₁-C₄)alkyl and halo; and each phenyl moiety in the groups represented by R^2 is optionally substituted with one or more substituents independently selected from the group consisting of (C₁-C₄)alkyl, halo, (C₁-C₄)alkoxy, (C₁-C₄)alkoxy(C₁-C₄)alkyl, -CN, halo(C₁-C₄)alkyl, and
20 halo(C₁-C₄)alkoxy. Alternatively, R^2 is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclopropylmethyl, cyclobutylmethyl, phenyl, benzyl, -(CH₂)₂-O-CH₃, -(CH₂)₃-OCH₃, -C(CH₃)₃, -CH(CH₃)₂, -CH₂C(CH₃)₃, -CH₂CH(CH₃)₂, -CH₂-CF₃, -(CH₂)₂-CH₂F, and -(CH₂)_nCH₃, wherein n is 0, 1, 2, 3, 4, 5 or 6; and wherein the phenyl or benzyl group represented by R^2 is optionally
25 substituted with one or more substituents independently selected from the group consisting of (C₁-C₄)alkyl, halo, (C₁-C₄)alkoxy, (C₁-C₄)alkoxy(C₁-C₄)alkyl, -CN, halo(C₁-C₄)alkyl, and halo(C₁-C₄)alkoxy. In another alternative, the phenyl or benzyl group represented by R^2 is unsubstituted. In yet another alternative, R^2 is selected from cyclopropyl, cyclopropylmethyl, cyclobutyl, cyclopentyl, cyclohexyl,
30 -(CH₂)₂-O-CH₃, -C(CH₃)₃, -CH(CH₃)₂, -CH₂-CF₃, -CH₂CH(CH₃)₂, -CH₃ and -CH₂CH₃.

In a second alternative embodiment, for compounds represented by Structural Formula (I) or (II), R¹ is hydrogen, methyl or ethyl; and values and alternative values for R² are as described above for the first alternative embodiment.

In a third alternative embodiment, for compounds represented by Structural Formula (I) or (II), R¹ is hydrogen, (C₁-C₄)alkyl or -O-(C₁-C₄)alkyl; R² is selected from (C₁-C₇)alkyl, (C₃-C₆)cycloalkyl(C₁-C₄)alkyl, (C₁-C₇)alkoxy(C₁-C₄)alkyl, phenyl, (C₃-C₆)cycloalkyl, and fluoro(C₁-C₄)alkyl; or R¹ and R² taken together with the nitrogen atom to which they are bonded form a ring selected from pyrrolidinyl, morpholinyl, azetidiny, piperidinyl, octahydrocyclopenta[c]pyrrolyl, isoindolinyl, indazolyl, imidazolyl, pyrazolyl, triazolyl, and tetrazolyl, wherein the ring formed by R¹ and R² taken together with the nitrogen atom to which they are bonded is optionally substituted with fluoro, -OH, -OCH₃, or N(CH₃)₂. More specifically, R¹ is hydrogen, methyl, ethyl, methoxy or tert-butoxy.

In a fourth alternative embodiment, for compounds represented by Structural Formula (I) or (II), R¹ hydrogen, methyl, or ethyl; R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, 2,2-dimethylpropyl, t-butyl, isobutyl, n-pentyl, (C₄-C₆)cycloalkyl, (C₃-C₅)cycloalkylmethyl, methoxyethyl, and 2-fluoroethyl; or R¹ and R² taken together with the nitrogen atom to which they are bonded form a ring selected from azetidiny, pyrrolidinyl, piperidinyl, tetrazolyl, or octahydrocyclopenta[c]pyrrolyl, and wherein the ring formed by R¹ and R² taken together with the nitrogen atom to which they are bonded is optionally substituted with fluoro.

In a fifth alternative embodiment, for compounds represented by Structural Formula (A), when X is hydrogen, Y is selected from hydrogen, -(C₁-C₄)alkyl, -(C₁-C₄)alkylene-N(R^A)(R^B), -(C₁-C₄)alkylene-N(R^F)-C(O)-[C(R^D)(R^E)]₀₋₄-N(R^A)(R^B), -CH-N-OR^A, -N(R^A)(R^B), -N(R^F)-C(O)-[C(R^D)(R^E)]₁₋₄-N(R^A)(R^B), -NH-C(O)-C(R^D)(R^E)-N(R^A)(R^B), -N(R^F)-C(O)-N(R^A)(R^A), -N(R^F)-C(O)-(C₁-C₆)alkyl, -N(R^F)-C(O)-heterocyclyl, -N(R^F)-C(O)-heteroaryl, -N(R^F)-C(O)-carbocyclyl, -N(R^F)-C(O)-aryl, -N(R^F)-S(O)_m-(C₁-C₄)alkylene-N(R^A)(R^B), -N(R^F)-S(O)_m-(C₁-C₄)alkylene-carbocyclyl, -N(R^F)-S(O)_m-(C₁-C₄)alkylene-aryl;

R^{D'} is selected from (C₁-C₆)alkyl, carbocyclyl, aryl, heteroaryl, heterocyclyl, and a naturally occurring amino acid side chain moiety, or

$R^{D'}$ and R^E taken together with the carbon atom to which they are bound form a 3-7 membered carbocyclyl, or a 4-7 membered heterocyclyl, wherein the heterocyclyl formed by $R^{D'}$ and R^E optionally comprises one to two additional heteroatoms independently selected from N, S and O; and

5 $R^{F'}$ is selected from (C₁-C₇)alkyl, carbocyclyl, aryl and heteroaryl.

Values and alternative values for the remainder of the variables are as described above for Structural Formula (A). Alternatively, $R^{F'}$ is selected from (C₁-C₄)alkyl and phenyl and the remainder of the variables are as described above in the fifth alternative embodiment.

10 In a sixth alternative embodiment, for compounds represented by Structural Formula (A), X is selected from hydrogen, methyl, ethyl and phenyl; and Y is selected from hydrogen, -(C₁-C₄ alkyl), -(C₁-C₄)alkylene-N(R^A)(R^B), -(C₁-C₄)alkylene-NH-C(O)-[CH₂]₀₋₁-N(R^A)(R^B), -N(R^A)(R^B), -NH-C(O)-carbocyclyl, -NH-C(O)-aryl, -NH-C(O)-heterocyclyl, 15 -NH-C(O)-heteroaryl, -NH-C(O)-N(R^A)(R^A), -N(R^{F'})-C(O)-CH₂-N(R^A)(R^B), -NH-C(O)-C(R^{D'})(R^E)-N(R^A)(R^B) and -NH-S(O)_m-(C₁-C₄)alkylene-N(R^A)(R^B); or X is selected from methyl, ethyl and phenyl; and Y is -NH-C(O)-CH₂-N(R^A)(R^B), wherein:

each R^A is independently selected from hydrogen and methyl;

20 R^B is selected from hydrogen, (C₁-C₇)alkyl, -(C₀-C₆)alkylene-carbocyclyl, -(C₀-C₆)alkylene-aryl, -(C₀-C₆)alkylene-heteroaryl, -S(O)_m-(C₁-C₆)alkyl, -(C₀-C₄)alkylene-S(O)_m-carbocyclyl, -(C₀-C₄)alkylene-S(O)_m-aryl, -(C₀-C₄)alkylene-S(O)_m-heterocycle and -(C₀-C₄)alkylene-S(O)_m-heteroaryl; or

R^A and R^B when bound to a common nitrogen atom are taken together with 25 the nitrogen atom to form a heterocycle, wherein the heterocycle is optionally substituted with =O and -N(R^G)(R^G);

$R^{D'}$ and R^E are taken together with the carbon atom to which they are bound form a (C₃-C₇)cycloalkyl; and

m is 1 or 2;

30 each carbocyclyl, aryl, heterocyclyl or heteroaryl is optionally and independently substituted with one or more substituents independently selected from halo, -(C₁-C₄)alkyl, halo-substituted -(C₁-C₄)alkyl, -O-(C₁-C₄)alkyl, and -N(R^G)(R^G);

each alkyl portion in the group represented by Y or R^B is optionally and independently substituted with one or more substituents independently selected from halo, -OH, and -N(R^G)(R^G), wherein

R^G is hydrogen or (C₁-C₄)alkyl, and wherein each alkyl in the group
5 represented by R^G is optionally and independently substituted with (C₃-C₆) cycloalkyl. The remainder of the variables are as described above in the fifth alternative embodiment.

In a seventh alternative embodiment, for compounds represented by Structural Formula (A), X is selected from hydrogen and methyl; and

10 Y is selected from -N(R^A)(R^B), -N(H)-C(O)-carbocyclyl, -N(H)-C(O)-aryl, -N(H)-C(O)-heterocycle, and -N(H)-C(O)-heteroaryl; or

X is methyl; and


Y is -NH-C(O)-CH₂-N(R^A)(R^B),

wherein:

15 R^A is hydrogen; and

R^B is selected from (C₁-C₄)alkyl, and -S(O)₂-CH₃; or R^A and R^B taken together to form 4-7 membered heterocyclic ring; wherein each carbocyclyl, aryl, heteroaryl, or heterocycle is optionally and independently substituted with one or more substituents independently selected from -CH₃, fluoro, and -N(CH₃)₂.

20 In a eighth alternative embodiment, for compounds represented by Structural



Formula (A), Y is , wherein ring A represents a 4-7 membered heterocyclyl; and R^{3'} is hydrogen or (C₁-C₆)alkyl. Values and alternative values for the remainder of the variables are as described above for Structural Formula (A).

More specifically, ring A is selected from the group consisting of azetidiny, pyrrolidiny, piperidiny, or octahydrocyclopenta[c]pyrrolyl, each of which is
25 optionally substituted with one or more substituents independently selected from the group consisting of halo, -(C₁-C₄)alkyl, halo-substituted-(C₁-C₄)alkyl (e.g., -CF₃), -OH, -O-(C₁-C₄)alkyl, or -N(R^G)(R^G), wherein R^G is hydrogen or (C₁-C₄)alkyl. Even more specifically, ring A described above is optionally substituted
30 with one or more fluoro.

In a ninth alternative embodiment, for compounds represented by Structural Formula (A), Y is -NH-C(O)-heteroaryl. Values and alternative values for the remainder of the variables are as described above for Structural Formula (A). More specifically, the heteroaryl in -NH-C(O)-heteroaryl is selected from the group
5 consisting of thienyl, pyridinyl, pyrrolyl, oxazolyl, pyrazolyl and thiazolyl, each of which is optionally substituted with one or more substituents independently selected from -(C₁-C₄)alkyl, halo-substituted-(C₁-C₄)alkyl (*e.g.*, -CF₃), -OH, -O-(C₁-C₄)alkyl, and -N(R^G)(R^G), wherein R^G is hydrogen or (C₁-C₄)alkyl. More specifically, the pyrrolyl and pyrazolyl are optionally substituted with a methyl group on the N atom
10 in the ring.

In a tenth alternative embodiment, for compounds represented by Structural Formula (A), Y is -NH-C(O)-phenyl. Values and alternative values for the remainder of the variables are as described above for Structural Formula (A). More specifically, the phenyl in -NH-C(O)-phenyl is optionally substituted with one or
15 more substituents independently selected from -(C₁-C₄)alkyl, halo-substituted-(C₁-C₄)alkyl (*e.g.*, -CF₃), -OH, -O-(C₁-C₄)alkyl, and -N(R^G)(R^G), wherein R^G is hydrogen or (C₁-C₄)alkyl. More specifically, the phenyl in -NH-C(O)-phenyl is optionally substituted with one or more substituents independently selected from -CF₃, -OCH₃ and -N(CH₃)₂.

In an eleventh alternative embodiment, for compounds represented by Structural Formula (A), Y is represented by -NH-S(O)₂-(C₁-C₆)alkyl, -NH-S(O)₂-phenyl, -NH-S(O)₂-heteroaryl. Values and alternative values for the remainder of the variables are as described above for Structural Formula (A). More specifically, the phenyl, heteroaryl or alkyl in the group represented by Y is
20 optionally substituted with one or more substituents independently selected from -(C₁-C₄)alkyl, halo-substituted-(C₁-C₄)alkyl (*e.g.*, -CF₃), -OH, -O-(C₁-C₄)alkyl, and -N(R^G)(R^G), wherein R^G is hydrogen or (C₁-C₄)alkyl.
25








In a twelfth alternative embodiment, for compounds represented by Structural Formula (A), Y is represented by -N(R^A)(R^B), wherein R^A and R^B are
30 each independently selected from hydrogen, (C₁-C₇)alkyl, -(C₁-C₄)alkyl-(C₃-C₆)cycloalkyl, wherein the (C₁-C₇)alkyl is optionally substituted with -N(R^G)(R^G), wherein R^G is hydrogen or (C₁-C₄)alkyl. Values and alternative values for the remainder of the variables are as described above for Structural Formula (A).








In a thirteenth alternative embodiment, for compounds represented by Structural Formula (A), Y is represented by $-\text{CH}_2-\text{N}(\text{R}^{\text{A}})(\text{R}^{\text{B}})$, wherein R^{A} and R^{B} are each independently selected from hydrogen, $(\text{C}_1-\text{C}_7)\text{alkyl}$, $-(\text{C}_1-\text{C}_4)\text{alkyl}-(\text{C}_3-\text{C}_6)\text{cycloalkyl}$, or R^{A} and R^{B} taken together with the nitrogen atom to which they are bound form a heterocyclyl, wherein the $(\text{C}_1-\text{C}_7)\text{alkyl}$ represented by R^{A} or R^{B} is optionally and independently substituted with $-\text{N}(\text{R}^{\text{G}})(\text{R}^{\text{G}})$, wherein R^{G} is hydrogen or $(\text{C}_1-\text{C}_4)\text{alkyl}$, and the $(\text{C}_1-\text{C}_4)\text{alkyl}$ represented by R^{G} is optionally substituted with $-\text{F}$. Values and alternative values for the remainder of the variables are as described above for Structural Formula (A).








In a fourteenth alternative embodiment, for compounds represented by Structural Formula (A), Y is represented by $-\text{CH}_2-\text{NH}-\text{C}(\text{O})-(\text{CH}_2)_{0-1}-\text{N}(\text{R}^{\text{A}})(\text{R}^{\text{B}})$, wherein R^{A} and R^{B} are each independently selected from hydrogen, $(\text{C}_1-\text{C}_7)\text{alkyl}$, $-(\text{C}_1-\text{C}_4)\text{alkyl}-(\text{C}_3-\text{C}_6)\text{cycloalkyl}$. Values and alternative values for the remainder of the variables are as described above for Structural Formula (A).

The compound of the present invention is exemplified by the compounds shown in the Table below or a pharmaceutically acceptable salt thereof and compounds described in Examples 1-12 below or a pharmaceutically acceptable salt thereof:







COMPOUND NO.	CHEMICAL STRUCTURE
11	
12	

13	
14	
15	
16	
17	
18	
19	

20	
21	
22	
23	
24	
25	
26	

27	
28	
29	
30	
31	
32	
33	

34	
35	
36	
37	
38	
39	
40	

41	
42	
43	
44	
45	
46	

“Alkyl” means a saturated aliphatic branched or straight-chain monovalent hydrocarbon radical having the specified number of carbon atoms. Thus, “(C₁-C₇)alkyl” means a radical having from 1-7 carbon atoms in a linear or branched arrangement. “(C₁-C₇)alkyl” includes methyl, ethyl, propyl, butyl, pentyl, hexyl

and heptyl. Suitable substitutions for a "substituted alkyl" include, but are not limited to, -halogen, -OH, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, (C₁-C₄)alkylthio, (C₁-C₄)alkylsulfinyl, (C₁-C₄)alkylsulfonyl, (C₁-C₄)alkoxy(C₁-C₄)alkyl, and -N(R³)(R⁴), wherein R³ and R⁴ are as described above.

- 5 "Cycloalkyl" means a saturated aliphatic cyclic hydrocarbon radical having the specified number of carbon atoms. (C₃-C₆)cycloalkyl includes cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl. Suitable substituents for a "substituted cycloalkyl" include halogen, -OH, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, (C₁-C₄)alkylthio, (C₁-C₄)alkylsulfinyl, (C₁-C₄)alkylsulfonyl, (C₁-C₄)alkoxy(C₁-C₄)alkyl, and
10 -N(R³)(R⁴), wherein R³ and R⁴ are as described above.

- "Heterocycle" means a 4-12 membered partially unsaturated or saturated heterocyclic ring containing 1, 2, or 3 heteroatoms independently selected from N, O or S. When one heteroatom is S, it can be optionally mono- or di-oxygenated (i.e. -S(O)- or -S(O)₂-). The heterocycle can be monocyclic, fused bicyclic, bridged
15 bicyclic, or spiro bicyclic.

- Examples of monocyclic heterocycle include, but not limited to, azetidine, pyrrolidine, piperidine, piperazine, hexahydropyrimidine, tetrahydrofuran, tetrahydropyran, morpholine, thiomorpholine, thiomorpholine 1,1-dioxide, tetrahydro-2H-1,2-thiazine, tetrahydro-2H-1,2-thiazine 1,1-dioxide, isothiazolidine,
20 isothiazolidine 1,1-dioxide.

- A fused bicyclic heterocycle has two rings which have two adjacent ring atoms in common. The first ring is a monocyclic heterocycle and the second ring is a cycloalkyl, partially unsaturated carbocycle, phenyl, heteroaryl or a monocyclic heterocycle. For example, the second ring is a (C₃-C₆)cycloalkyl, such as
25 cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. Alternatively, the second ring is phenyl. Example of fused bicyclic heterocycles includes, but not limited to, indoline, isoindoline, 2,3-dihydro-1H-benzo[d]imidazole, 2,3-dihydrobenzo[d]oxazole, 2,3-dihydrobenzo[d]thiazole, octahydrobenzo[d]oxazole, octahydro-1H-benzo[d]imidazole, octahydrobenzo[d]thiazole,
30 octahydrocyclopenta[c]pyrrole, 3-azabicyclo[3.1.0]hexane, and 3-azabicyclo[3.2.0]heptane.

 A spiro bicyclic heterocycle has two rings which have only one ring atom in common. The first ring is a monocyclic heterocycle and the second ring is a

cycloalkyl, partially unsaturated carbocycle or a monocyclic heterocycle. For example, the second ring is a (C₃-C₆)cycloalkyl. Example of spiro bicyclic heterocycle includes, but not limited to, azaspiro[4.4]nonane, 7-azaspiro[4.4]nonane, azaspiro[4.5]decane, 8-azaspiro[4.5]decane,

5 azaspiro[5.5]undecane, 3-azaspiro[5.5]undecane and 3,9-diazaspiro[5.5]undecane.

A bridged bicyclic heterocycle has two rings which have three or more adjacent ring atoms in common. The first ring is a monocyclic heterocycle and the other ring is a cycloalkyl (such as (C₃-C₆)cycloalkyl), partially unsaturated carbocycle or a monocyclic heterocycle. Examples of bridged bicyclic heterocycles
10 include, but are not limited to, azabicyclo[3.3.1]nonane, 3-azabicyclo[3.3.1]nonane, azabicyclo[3.2.1]octane, 3-azabicyclo[3.2.1]octane, 6-azabicyclo[3.2.1]octane and azabicyclo[2.2.2]octane, 2-azabicyclo[2.2.2]octane.

When the heterocycle contains a N atom other than the nitrogen atom to which R¹ and R² are bonded, the N atom can be substituted with H, alkyl,
15 cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, each of which can be optionally substituted with halogen, hydroxy, alkoxy, haloalkyl, alkyl, etc. The heterocycle can be optionally substituted with an oxo group (C=O) and oxo substituted heterocyclic rings include, but are not limited to, thiomorpholine 1-oxide, thiomorpholine 1,1-dioxide, tetrahydro-2H-1,2-thiazine 1,1-dioxide, and
20 isothiazolidine 1,1-dioxide, pyrrolidin-2-one, piperidin-2-one, piperazin-2-one, and morpholin-2-one. Other optional substituents for a heterocycle include (C₁-C₄)alkyl, halo, -OH, (C₁-C₄)alkoxy, (C₁-C₄)alkylthio, (C₁-C₄)alkylsulfinyl, (C₁-C₄)alkylsulfonyl, (C₁-C₄)alkoxy(C₁-C₄)alkyl, -N(R³)(R⁴), -CN, halo(C₁-C₄)alkyl, and halo(C₁-C₄)alkoxy.

25 "Heteroaryl" means a 5-12 membered monovalent heteroaromatic monocyclic or bicyclic ring radical. A heteroaryl contains 1, 2 or 3 heteroatoms independently selected from N, O, and S. Heteroaryls include, but are not limited to pyrrole, imidazole, pyrazole, oxazole, isoxazole, thiazole, isothiazole, 1,2,3-triazole, 1,2,4-triazole, 1,3,4-oxadiazole, 1,2,5-thiadiazole, 1,2,5-thiadiazole 1-oxide, 1,2,5-thiadiazole 1,1-dioxide, 1,3,4-thiadiazole, pyridine, pyrazine,
30 pyrimidine, pyridazine, 1,2,4-triazine, 1,3,5-triazine, and tetrazole. Bicyclic heteroaryl rings include, but are not limited to, bicyclo[4.4.0] and bicyclo[4.3.0] fused ring systems such as indolizine, indole, isoindole, indazole, benzimidazole,

benzthiazole, purine, quinoline, isoquinoline, cinnoline, phthalazine, quinazoline, quinoxaline, 1,8-naphthyridine, and pteridine.

“Carbocycle” means 4-12 membered saturated or unsaturated aliphatic cyclic hydrocarbon ring.

- 5 “Alkoxy” means an alkyl radical attached through an oxygen linking atom. “Alkoxy” can also be depicted as -O-alkyl. For example, (C₁-C₄)-alkoxy can also be depicted as -O-(C₁-C₄)alkyl. “(C₁-C₄)-alkoxy” includes methoxy, ethoxy, propoxy, and butoxy.

- 10 “Alkylthio” means an alkyl radical attached through a sulfur linking atom. “Alkylthio” can also be depicted as -S-alkyl. For example, “(C₁-C₄)alkylthio” can be depicted as -S-(C₁-C₄)alkyl. “(C₁-C₄)alkylthio” include methylthio, ethylthio, propylthio and butylthio.

- 15 “Alkylsulfinyl” means an alkyl radical attached through a -S(O)- linking group. “Alkylsulfinyl” can be depicted as -S(O)-alkyl. For example, “(C₁-C₄)alkylsulfinyl” can be depicted as -S(O)-(C₁-C₄)alkyl. “(C₁-C₄)alkylsulfinyl” include methylsulfinyl, ethylsulfinyl, propylsulfinyl and butylsulfinyl.

- 20 “Alkylsulfonyl” means an alkyl radical attached through a -S(O)₂- linking group. “Alkylsulfonyl” can be depicted as -S(O)₂-alkyl. For example, “(C₁-C₄)alkylsulfonyl” can be depicted as -S(O)₂-(C₁-C₄)alkyl. “(C₁-C₄)alkylsulfonyl” include methylsulfonyl, ethylsulfonyl, propylsulfonyl and butylsulfonyl.

- 25 Haloalkyl and halocycloalkyl include mono, poly, and perhaloalkyl groups where each halogen is independently selected from fluorine, chlorine, and bromine. Haloalkyl and halocycloalkyl can also be referred as halo-substituted alkyl and halo-substituted cycloalkyl, respectively.

- 30 “Cycloalkoxy” means a cycloalkyl radical attached through an oxygen linking atom. “Cycloalkoxy” can also be depicted as -O-cycloalkyl. For example, “(C₃-C₆)cycloalkoxy” can be depicted as -O-(C₃-C₆)cycloalkyl. “(C₃-C₆)cycloalkoxy” includes cyclopropyloxy, cyclobutyloxy, cyclopentyloxy and cyclohexyloxy.

“Aryl” means an aromatic monocyclic or polycyclic (e.g. bicyclic or tricyclic) carbocyclic ring system. In one embodiment, “aryl” is a 6-12 membered monocyclic or bicyclic systems. Aryl systems include, but not limited to, phenyl, naphthalenyl, fluorenyl, indenyl, azulenyl, and anthracenyl.

5 “Aryloxy” means an aryl moiety attached through an oxygen linking atom. “Aryloxy” can be also depicted as -O-aryl. Aryloxy includes, but not limited to, phenoxy.

“Arylthio” means an aryl moiety attached through a sulfur linking atom. “Arylthio” can be also depicted as -S-aryl. Arylthio includes, but not limited to, phenylthio.

10

“Arylsulfinyl” means an aryl moiety attached through a -S(O)- linking group. “Arylsulfinyl” can be also depicted as -S(O)-aryl. Arylsulfinyl includes, but not limited to, phenylsulfinyl.

“Arylsulfonyl” means an aryl moiety attached through a -S(O)₂- linking group. “Arylsulfonyl” can be also depicted as -S(O)₂-aryl. Arylsulfonyl includes, but not limited to, phenylsulfonyl.

15

“Hetero” refers to the replacement of at least one carbon atom member in a ring system with at least one heteroatom selected from N, S, and O. “Hetero” also refers to the replacement of at least one carbon atom member in a acyclic system. A hetero ring system or a hetero acyclic system may have 1, 2, or 3 carbon atom members replaced by a heteroatom.

20

“Halogen” or “halo” used herein refers to fluorine, chlorine, bromine, or iodine.

As used herein, cycloalkylalkyl can be depicted as -alkylene-cycloalkyl. For example, (C₃-C₆)cycloalkyl(C₁-C₄)alkyl can be depicted as -(C₁-C₄)alkylene-(C₃-C₆)cycloalkyl.

25

As use herein, alkoxyalkyl can be depicted as -alkylene-O-alkyl. For example, (C₁-C₇)alkoxy(C₁-C₄)alkyl can be depicted as -(C₁-C₄)alkylene-O-(C₁-C₇)alkyl.

As use herein, cycloalkoxyalkyl can be depicted as -alkylene-O-cycloalkyl. For example, (C₃-C₆)cycloalkoxy(C₁-C₄)alkyl can be depicted as -(C₁-C₄)alkylene-O-(C₃-C₅)alkyl.

30

As used herein, arylalkyl can be depicted as -alkylene-aryl. For example, aryl(C₁-C₄)alkyl can be depicted as -(C₁-C₄)alkylene-aryl.

As used herein, aryloxyalkyl can be depicted as -alkylene-O-aryl. For example, aryloxy(C₁-C₄)alkyl can be depicted as -(C₁-C₄)alkylene-O-aryl.

5 As used herein, arylthioalkyl can be depicted as -alkylene-S-aryl. For example, arylthio(C₁-C₄)alkyl can be depicted as -(C₁-C₄)alkylene-S-aryl.

As used herein, arylsulfinylalkyl can be depicted as -alkylene-S(O)-aryl. For example, arylsulfinyl(C₁-C₄)alkyl can be depicted as -(C₁-C₄)alkylene-S(O)-aryl.

10 As used herein, arylsulfonylalkyl can be depicted as -alkylene-S(O)₂-aryl. For example, arylsulfonyl(C₁-C₄)alkyl can be depicted as -(C₁-C₄)alkylene-S(O)₂-aryl.

Another embodiment of the present invention is a pharmaceutical composition comprising one or more pharmaceutically acceptable carrier and/or diluent and a compound disclosed herein or a pharmaceutically acceptable salt thereof.

20 "Pharmaceutically acceptable carrier" and "pharmaceutically acceptable diluent" means non-therapeutic components that are of sufficient purity and quality for use in the formulation of a composition of the invention that, when appropriately administered to an animal or human, typically do not produce an adverse reaction, and that are used as a vehicle for a drug substance (i.e. a compound of the present invention).

Pharmaceutically acceptable salts of the compounds of the present invention are also included. For example, an acid salt of a compound of the present invention containing an amine or other basic group can be obtained by reacting the compound with a suitable organic or inorganic acid, resulting in pharmaceutically acceptable anionic salt forms. Examples of anionic salts include the acetate, benzenesulfonate, benzoate, bicarbonate, bitartrate, bromide, calcium edetate, camsylate, carbonate, chloride, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, glyceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrobromide, hydrochloride, hydroxynaphthoate, iodide, isethionate, lactate, lactobionate, malate, maleate, mandelate, mesylate, methylsulfate, mucate, napsylate, nitrate, pamoate, pantothenate, phosphate/diphosphate, polygalacturonate,

salicylate, stearate, subacetate, succinate, sulfate, tannate, tartrate, teoclate, tosylate, and triethiodide salts.

Salts of the compounds of the present invention containing a carboxylic acid or other acidic functional group can be prepared by reacting with a suitable base.

5 Such a pharmaceutically acceptable salt may be made with a base which affords a pharmaceutically acceptable cation, which includes alkali metal salts (especially sodium and potassium), alkaline earth metal salts (especially calcium and magnesium), aluminum salts and ammonium salts, as well as salts made from physiologically acceptable organic bases such as trimethylamine, triethylamine,
10 morpholine, pyridine, piperidine, picoline, dicyclohexylamine, N,N'-dibenzylethylenediamine, 2-hydroxyethylamine, bis-(2-hydroxyethyl)amine, tri-(2-hydroxyethyl)amine, procaine, dibenzylpiperidine, dehydroabietylamine, N,N'-bisdehydroabietylamine, glucamine, N-methylglucamine, collidine, quinine, quinoline, and basic amino acids such as lysine and arginine.

15 The invention also includes various isomers and mixtures thereof. Certain of the compounds of the present invention may exist in various stereoisomeric forms. Stereoisomers are compounds which differ only in their spatial arrangement. Enantiomers are pairs of stereoisomers whose mirror images are not superimposable, most commonly because they contain an asymmetrically
20 substituted carbon atom that acts as a chiral center. "Enantiomer" means one of a pair of molecules that are mirror images of each other and are not superimposable. Diastereomers are stereoisomers that are not related as mirror images, most commonly because they contain two or more asymmetrically substituted carbon atoms. "R" and "S" represent the configuration of substituents around one or more
25 chiral carbon atoms. When a chiral center is not defined as R or S, either a pure enantiomer or a mixture of both configurations is present.

"Racemate" or "racemic mixture" means a compound of equimolar quantities of two enantiomers, wherein such mixtures exhibit no optical activity; i.e., they do not rotate the plane of polarized light.

30 The compounds of the invention may be prepared as individual isomers by either isomer-specific synthesis or resolved from an isomeric mixture. Conventional resolution techniques include forming the salt of a free base of each isomer of an isomeric pair using an optically active acid (followed by fractional

crystallization and regeneration of the free base), forming the salt of the acid form of each isomer of an isomeric pair using an optically active amine (followed by fractional crystallization and regeneration of the free acid), forming an ester or amide of each of the isomers of an isomeric pair using an optically pure acid, amine
5 or alcohol (followed by chromatographic separation and removal of the chiral auxiliary), or resolving an isomeric mixture of either a starting material or a final product using various well known chromatographic methods.

When the stereochemistry of a disclosed compound is named or depicted by structure, the named or depicted stereoisomer is at least 60%, 70%, 80%, 90%, 99%
10 or 99.9% by weight pure relative to the other stereoisomers. When a single enantiomer is named or depicted by structure, the depicted or named enantiomer is at least 60%, 70%, 80%, 90%, 99% or 99.9% by weight optically pure. Percent optical purity by weight is the ratio of the weight of the enantiomer that is present divided by the combined weight of the enantiomer that is present and the weight of
15 its optical isomer.

The present invention also provides a method of treating or preventing a subject with a tetracycline-responsive disease or disorder comprising administering to the subject an effective amount of a compound of the present invention or a pharmaceutically acceptable salt thereof.

20 "Tetracycline-responsive disease or disorder" refers to a disease or disorder that can be treated, prevented, or otherwise ameliorated by the administration of a tetracycline compound of the present invention. Tetracycline-responsive disease or disorder includes infections, cancer, inflammatory disorders, autoimmune disease, arteriosclerosis, corneal ulceration, emphysema, arthritis, osteoporosis,
25 osteoarthritis, multiple sclerosis, osteosarcoma, osteomyelitis, bronchiectasis, chronic pulmonary obstructive disease, skin and eye diseases, periodontitis, osteoporosis, rheumatoid arthritis, ulcerative colitis, prostatitis, tumor growth and invasion, metastasis, diabetes, diabetic proteinuria, panbronchiolitis; aortic or vascular aneurysms, skin tissue wounds, dry eye, bone, cartilage degradation,
30 malaria, senescence, diabetes, vascular stroke, neurodegenerative disorders, cardiac disease, juvenile diabetes, acute and chronic bronchitis, sinusitis, and respiratory infections, including the common cold; Wegener's granulomatosis; neutrophilic dermatoses and other inflammatory diseases such as dermatitis herpetiformis,

leukocytoclastic vasculitis, bullous lupus erythematosus, pustular psoriasis, erythema elevatum diutinum; vitiligo; discoid lupus erythematosus; pyoderma gangrenosum; pustular psoriasis; blepharitis, or meibomianitis; Alzheimer's disease; degenerative maculopathy; acute and chronic gastroenteritis and colitis; acute and chronic cystitis and urethritis; acute and chronic dermatitis; acute and chronic conjunctivitis; acute and chronic serositis; uremic pericarditis; acute and chronic cholecystitis; cystic fibrosis, acute and chronic vaginitis; acute and chronic uveitis; drug reactions; insect bites; burns and sunburn, bone mass disorder, acute lung injury, chronic lung disorders, ischemia, stroke or ischemic stroke, skin wound, aortic or vascular aneurysm, diabetic retinopathy, hemorrhagic stroke, angiogenesis, and other states for which tetracycline compounds have been found to be active (see, for example, U. S. Patent Nos. 5,789,395; 5,834,450; 6,277,061 and 5,532,227, each of which is expressly incorporated herein by reference).

In addition, a method to treat any disease or disease state that could benefit from modulating the expression and/or function of nitric oxide, metalloproteases, proinflammatory mediators and cytokines, reactive oxygen species, components of the immune response, including chemotaxis, lymphocyte transformation, delayed hypersensitivity, antibody production, phagocytosis, and oxidative metabolism of phagocytes. A method to treat any disease or disease state that could benefit from modulating the expression and/or function of C-reactive protein, signaling pathways (e.g., FAK signaling pathway), and/or augment the expression of COX-2 and PGE₂ production is covered. A method to treat any disease or disease state that could benefit from inhibition of neovascularization is covered.

Compounds of the invention can be used to prevent or treat important mammalian and veterinary diseases such as diarrhea, urinary tract infections, infections of skin and skin structure, ear, nose and throat infections, wound infection, mastitis and the like. In addition, methods for treating neoplasms using tetracycline compounds of the invention are also included (van der Bozert et al., Cancer Res., 48: 6686-6690 (1988)).

Infections that can be treated using compounds of the invention or a pharmaceutically acceptable salt thereof include, but are not limited to, skin infections, GI infections, urinary tract infections, genito-urinary infections, respiratory tract infections, sinuses infections, middle ear infections, systemic

infections, cholera, influenza, bronchitis, acne, malaria, sexually transmitted disease including syphilis and gonorrhea, Legionnaires' disease, Lyme disease, Rocky Mountain spotted fever, Q fever, typhus, bubonic plague, gas gangrene, hospital acquired infections, leptospirosis, whooping cough, anthrax and infections caused by the agents responsible for lymphogranuloma venereum, inclusion conjunctivitis, or psittacosis. Infections can be bacterial, fungal, parasitic and viral infections (including those which are resistant to other tetracycline compounds).

In one embodiment, the infection can be caused bacteria. In another embodiment, the infection is caused by a Gram-positive bacteria. In a specific aspect of this embodiment, the infection is caused by a Gram-positive bacterium selected from *Staphylococcus* spp., *Streptococcus* spp., *Propionibacterium* spp., *Enterococcus* spp., *Bacillus* spp., *Corynebacterium* spp., *Nocardia* spp., *Clostridium* spp., *Actinobacteria* spp., and *Listeria* spp.

In another embodiment, the infection is caused by a Gram-negative bacterium. In one aspect of this embodiment, the infection is caused by a proteobacteria (e.g., Betaproteobacteria and Gammaproteobacteria), including *Escherichia coli*, *Salmonella*, *Shigella*, other *Enterobacteriaceae*, *Pseudomonas*, *Moraxella*, *Helicobacter*, *Stenotrophomonas*, *Bdellovibrio*, acetic acid bacteria, *Legionella* or alpha-proteobacteria such as *Wolbachia*. In another aspect, the infection is caused by a Gram-negative bacteria selected from cyanobacteria, spirochaetes, green sulfur or green non-sulfur bacteria. In a specific aspect of this embodiment, the infection is caused by a Gram-negative bacteria selected from *Enterobacteriaceae* (e.g., *E. coli*, *Klebsiella pneumoniae* including those containing extended-spectrum β -lactamases and/or carbapenemases), *Bacteroidaceae* (e.g., *Bacteroides fragilis*), *Vibrionaceae* (*Vibrio cholerae*), *Pasteurellae* (e.g., *Haemophilus influenza*), *Pseudomonadaceae* (e.g., *Pseudomonas aeruginosa*), *Neisseriaceae* (e.g. *Neisseria meningitidis*), *Rickettsiae*, *Moraxellaceae* (e.g., *Moraxella catarrhalis*), any species of *Proteaceae*, *Acinetobacter* spp., *Helicobacter* spp., and *Campylobacter* spp.

In a particular embodiment, the infection is caused by Gram-negative bacterium selected from the group consisting of *Enterobacteriaceae* (e.g., *E. coli*, *Klebsiella pneumoniae*), *Pseudomonas*, and *Acinetobacter* spp.

In another embodiment, the infection is caused by an organism selected from the group consisting of *K. pneumoniae*, *Salmonella*, *E. hirae*, *A. baumanni*, *M. catarrhalis*, *H. influenzae*, *P. aeruginosa*, *E. faecium*, *E. coli*, *S. aureus*, and *E. faecalis*.

5 In another embodiment, the infection is caused by an organism selected from the group consisting of rickettsiae, chlamydiae, *Legionella* spp. and *Mycoplasma* spp. In another embodiment, the infection is caused by an organism resistant to tetracycline or any member of first and second generation of tetracycline antibiotics (e.g., doxycycline or minocycline).

10 In another embodiment, the infection is caused by an organism resistant to methicillin.

In another embodiment, the infection is caused by an organism resistant to vancomycin.

15 In another embodiment, the infection is caused by an organism resistant to a quinolone or fluoroquinolone.

In another embodiment, the infection is caused by an organism resistant to tigecycline.

20 In another embodiment, the infection is caused by a multidrug-resistant pathogen (having intermediate or full resistance to any two or more antibiotics). In another embodiment the infection is a *Bacillus anthracis* infection. "*Bacillus anthracis* infection" includes any state, diseases, or disorders caused or which result from exposure or alleged exposure to *Bacillus anthracis* or another member of the *Bacillus cereus* group of bacteria. In another embodiment, the infection is caused by *Bacillus anthracis* (anthrax), *Yersinia pestis* (plague), or *Francisella tularensis* (tularemia).

25 In yet another embodiment, the infection can be caused by more than one organism described above. Examples of such infections include, but are not limited to, intra-abdominal infections (often a mixture of a gram-negative species like *E. coli* and an anaerobe like *B. fragilis*), diabetic foot (various combinations of *Streptococcus*, *Serratia*, *Staphylococcus* and *Enterococcus* spp., anaerobes (S.E. Dowd, et al., PloS one 2008;3:e3326) and respiratory disease (especially in patients that have chronic infections like cystic fibrosis – e.g., *S. aureus* plus *P. aeruginosa* or *H. influenza*, atypical pathogens), wounds and abscesses (various gram-negative

and gram-positive bacteria, notably MSSA/MRSA, coagulase-negative staphylococci, enterococci, *Acinetobacter*, *P. aeruginosa*, *E. coli*, *B. fragilis*), and bloodstream infections (13% were polymicrobial (H. Wisplinghoff, et al., Clin. Infect. Dis. 2004;39:311-317)).

5 In a further embodiment, the tetracycline responsive disease or disorder is not a bacterial infection. In another embodiment, the tetracycline compounds of the invention are essentially non-antibacterial. For example, non-antibacterial compounds of the invention may have MIC values greater than about 4 µg/ml (as measured by assays known in the art and/or the assay given in Example 14. In
10 another embodiment, the tetracycline compounds of the invention have both antibacterial and non-antibacterial effects.

Tetracycline responsive disease or disorder also includes diseases or disorders associated with inflammatory process associated states (IPAS). The term "inflammatory process associated state" includes states in which inflammation or
15 inflammatory factors (e.g., matrix metalloproteinases (MMPs), nitric oxide (NO), TNF, interleukins, plasma proteins, cellular defense systems, cytokines, lipid metabolites, proteases, toxic radicals, adhesion molecules, etc.) are involved or are present in an area in aberrant amounts, e.g., in amounts which may be advantageous to alter, e.g., to benefit the subject. The inflammatory process is the response of
20 living tissue to damage. The cause of inflammation may be due to physical damage, chemical substances, micro-organisms, tissue necrosis, cancer or other agents. Acute inflammation is short-lasting, lasting only a few days. If it is longer lasting however, then it may be referred to as chronic inflammation.

IPASs include inflammatory disorders. Inflammatory disorders are generally
25 characterized by heat, redness, swelling, pain and loss of function. Examples of causes of inflammatory disorders include, but are not limited to, microbial infections (e.g., bacterial and fungal infections), physical agents (e.g., burns, radiation, and trauma), chemical agents (e.g., toxins and caustic substances), tissue necrosis and various types of immunologic reactions.

30 Examples of inflammatory disorders can be treated using the compounds of the invention or a pharmaceutically acceptable salt thereof include, but are not limited to, osteoarthritis, rheumatoid arthritis, acute and chronic infections (bacterial and fungal, including diphtheria and pertussis); acute and chronic bronchitis,

sinusitis, and upper respiratory infections, including the common cold; acute and chronic gastroenteritis and colitis; inflammatory bowel disorder; acute and chronic cystitis and urethritis; vasculitis; sepsis; nephritis; pancreatitis; hepatitis; lupus; inflammatory skin disorders including, for example, eczema, dermatitis, psoriasis, pyoderma gangrenosum, acne rosacea, and acute and chronic dermatitis; acute and chronic conjunctivitis; acute and chronic serositis (pericarditis, peritonitis, synovitis, pleuritis and tendinitis); uremic pericarditis; acute and chronic cholecystitis; acute and chronic vaginitis; acute and chronic uveitis; drug reactions; insect bites; burns (thermal, chemical, and electrical); and sunburn.

- 10 IPASs also include matrix metalloproteinase associated states (MMPAS). MMPAS include states characterized by aberrant amounts of MMPs or MMP activity. Examples of matrix metalloproteinase associated states ("MMPAS's") can be treated using compounds of the invention or a pharmaceutically acceptable salt thereof,
- 15 include, but are not limited to, arteriosclerosis, corneal ulceration, emphysema, osteoarthritis, multiple sclerosis (Liedtke et al., Ann. Neurol. 1998, 44: 35-46; Chandler et al., J. Neuroimmunol. 1997, 72: 155-71), osteosarcoma, osteomyelitis, bronchiectasis, chronic pulmonary obstructive disease, skin and eye diseases, periodontitis, osteoporosis, rheumatoid arthritis, ulcerative colitis, inflammatory disorders, tumor growth and invasion (Stetler-Stevenson et al., Annu. Rev. Cell Biol. 1993, 9: 541-73; Tryggvason et al., Biochim. Biophys. Acta 1987, 907: 191-217; Li et al., Mol. Carcinog. 1998, 22: 84-89), metastasis, acute lung injury, stroke, ischemia, diabetes, aortic or vascular aneurysms, skin tissue wounds, dry eye, bone and cartilage degradation (Greenwald et al., Bone 1998, 22: 33-38; Ryan et al., Curr. Op. Rheumatol. 1996, 8: 238-247). Other MMPAS include those described in U. S. Pat. Nos. 5,459,135; 5,321,017; 5,308,839; 5,258,371; 4,935,412; 4,704,383, 4,666,897, and RE 34,656, incorporated herein by reference in their entirety.
- 25

- In a further embodiment, the IPAS includes disorders described in U. S. Patents Nos. 5,929,055; and 5,532,227, incorporated herein by reference in their entirety.
- 30

Tetracycline responsive disease or disorder also includes diseases or disorders associated with NO associated states. The term "NO associated states" includes states which involve or are associated with nitric oxide (NO) or inducible

nitric oxide synthase (iNOS). NO associated state includes states which are characterized by aberrant amounts of NO and/or iNOS. Preferably, the NO associated state can be treated by administering tetracycline compounds of the invention. The disorders, diseases and states described in U. S. Patents Nos. 5 6,231,894; 6,015,804; 5,919,774; and 5,789,395 are also included as NO associated states. The entire contents of each of these patents are hereby incorporated herein by reference.

Examples of diseases or disorders associated with NO associated states can be treated using the compounds of the present invention or a pharmaceutically acceptable salt thereof include, but are not limited to, malaria, senescence, diabetes, 10 vascular stroke, neurodegenerative disorders (Alzheimer's disease and Huntington's disease), cardiac disease (reperfusion-associated injury following infarction), juvenile diabetes, inflammatory disorders, osteoarthritis, rheumatoid arthritis, acute, recurrent and chronic infections (bacterial, viral and fungal); acute and chronic 15 bronchitis, sinusitis, and respiratory infections, including the common cold; acute and chronic gastroenteritis and colitis; acute and chronic cystitis and urethritis; acute and chronic dermatitis; acute and chronic conjunctivitis; acute and chronic serositis (pericarditis, peritonitis, synovitis, pleuritis and tendonitis); uremic pericarditis; acute and chronic cholecystitis; cystic fibrosis, acute and chronic vaginitis; acute and 20 chronic uveitis; drug reactions; insect bites; burns (thermal, chemical, and electrical); and sunburn.

In another embodiment, the tetracycline responsive disease or disorder is cancer. Examples of cancers that can be treated using the compounds of the invention or a pharmaceutically acceptable salt thereof include all solid tumors, i.e., 25 carcinomas e.g., adenocarcinomas, and sarcomas. Adenocarcinomas are carcinomas derived from glandular tissue or in which the tumor cells form recognizable glandular structures. Sarcomas broadly include tumors whose cells are embedded in a fibrillar or homogeneous substance like embryonic connective tissue. Examples of carcinomas which may be treated using the methods of the invention include, but are 30 not limited to, carcinomas of the prostate, breast, ovary, testis, lung, colon, and breast. The methods of the invention are not limited to the treatment of these tumor types, but extend to any solid tumor derived from any organ system. Examples of treatable cancers include, but are not limited to, colon cancer, bladder cancer, breast

cancer, melanoma, ovarian carcinoma, prostate carcinoma, lung cancer, and a variety of other cancers as well. The methods of the invention also cause the inhibition of cancer growth in adenocarcinomas, such as, for example, those of the prostate, breast, kidney, ovary, testes, and colon. In one embodiment, the cancers
5 treated by methods of the invention include those described in U. S. Patent Nos. 6,100,248; 5,843,925; 5,837,696; or 5,668,122, incorporated herein by reference in their entirety.

Alternatively, the tetracycline compounds may be useful for preventing or reducing the likelihood of cancer recurrence, for example, to treat residual cancer
10 following surgical resection or radiation therapy. The tetracycline compounds useful according to the invention are especially advantageous as they are substantially non-toxic compared to other cancer treatments.

In a further embodiment, the compounds of the invention are administered in combination with standard cancer therapy, such as, but not limited to, chemotherapy.
15 Examples of tetracycline responsive states can be treated using the compounds of the invention or a pharmaceutically acceptable salt thereof also include neurological disorders which include both neuropsychiatric and neurodegenerative disorders, but are not limited to, such as Alzheimer's disease, dementias related to Alzheimer's disease (such as Pick's disease), Parkinson's and
20 other Lewy diffuse body diseases, senile dementia, Huntington's disease, Gilles de la Tourette's syndrome, multiple sclerosis, amyotrophic lateral sclerosis (ALS), progressive supranuclear palsy, epilepsy, and Creutzfeldt-Jakob disease; autonomic function disorders such as hypertension and sleep disorders, and neuropsychiatric disorders, such as depression, schizophrenia, schizoaffective disorder, Korsakoff's
25 psychosis, mania, anxiety disorders, or phobic disorders; learning or memory disorders, e. g., amnesia or age-related memory loss, attention deficit disorder, dysthymic disorder, major depressive disorder, mania, obsessive-compulsive disorder, psychoactive substance use disorders, anxiety, phobias, panic disorder, as well as bipolar affective disorder, e. g., severe bipolar affective (mood) disorder
30 (BP-I), bipolar affective neurological disorders, e. g., migraine and obesity.

Further neurological disorders include, for example, those listed in the American Psychiatric Association's Diagnostic and Statistical manual of Mental

Disorders (DSM), the most current version of which is incorporated herein by reference in its entirety.

In another embodiment, the tetracycline responsive disease or disorder is diabetes. Diabetes that can be treated using the compounds of the invention or a pharmaceutically acceptable salt thereof include, but are not limited to, juvenile
5 diabetes, diabetes mellitus, diabetes type I, or diabetes type II. In a further embodiment, protein glycosylation is not affected by the administration of the tetracycline compounds of the invention. In another embodiment, the tetracycline compound of the invention is administered in combination with standard diabetic
10 therapies, such as, but not limited to insulin therapy.

In another embodiment, the tetracycline responsive disease or disorder is a bone mass disorder. Bone mass disorders that can be treated using the compounds of the invention or a pharmaceutically acceptable salt thereof include disorders where a subjects bones are disorders and states where the formation, repair or
15 remodeling of bone is advantageous. For examples bone mass disorders include osteoporosis (e. g. , a decrease in bone strength and density), bone fractures, bone formation associated with surgical procedures (e. g. , facial reconstruction), osteogenesis imperfecta (brittle bone disease), hypophosphatasia, Paget's disease, fibrous dysplasia, osteopetrosis, myeloma bone disease, and the depletion of calcium
20 in bone, such as that which is related to primary hyperparathyroidism. Bone mass disorders include all states in which the formation, repair or remodeling of bone is advantageous to the subject as well as all other disorders associated with the bones or skeletal system of a subject which can be treated with the tetracycline compounds of the invention. In a further embodiment, the bone mass disorders include those
25 described in U. S. Patents Nos. 5,459,135; 5,231,017; 5,998,390; 5,770,588; RE 34,656; 5,308,839; 4,925,833; 3,304,227; and 4,666,897, each of which is hereby incorporated herein by reference in its entirety.

In another embodiment, the tetracycline responsive disease or disorder is acute lung injury. Acute lung injuries that can be treated using the compounds of
30 the invention or a pharmaceutically acceptable salt thereof include adult respiratory distress syndrome (ARDS), post-pump syndrome (PPS), and trauma. Trauma includes any injury to living tissue caused by an extrinsic agent or event. Examples

of trauma include, but are not limited to, crush injuries, contact with a hard surface, or cutting or other damage to the lungs.

The tetracycline responsive disease or disorders of the invention also include chronic lung disorders. Examples of chronic lung disorders that can be treated using the compounds of the invention or a pharmaceutically acceptable salt thereof
5 include, but are not limited, to asthma, cystic fibrosis, chronic obstructive pulmonary disease (COPD), and emphysema. In a further embodiment, the acute and/or chronic lung disorders that can be treated using the compounds of the invention or a pharmaceutically acceptable salt thereof include those described in U. S. Patents No.
10 5,977,091; 6,043,231; 5,523,297, and 5,773,430, each of which is hereby incorporated herein by reference in its entirety.

In yet another embodiment, the tetracycline responsive disease or disorder is ischemia, stroke, or ischemic stroke.

In a further embodiment, the tetracycline compounds of the invention or a
15 pharmaceutically acceptable salt thereof can be used to treat such disorders as described above and in U. S. Patents No. 6,231,894; 5,773,430; 5,919,775 and 5,789,395, incorporated herein by reference.

In another embodiment, the tetracycline responsive disease or disorder is a skin wound. The invention also provides a method for improving the healing
20 response of the epithelialized tissue (e.g., skin, mucosae) to acute traumatic injury (e.g., cut, burn, scrape, etc.). The method includes using a tetracycline compound of the invention or a pharmaceutically acceptable salt thereof to improve the capacity of the epithelialized tissue to heal acute wounds. The method may increase the rate of collagen accumulation of the healing tissue. The method may also decrease the
25 proteolytic activity in the epithelialized tissue by decreasing the collagenolytic and/or gellatinolytic activity of MMPs. In a further embodiment, the tetracycline compound of the invention or a pharmaceutically acceptable salt thereof is administered to the surface of the skin (e. g., topically). In a further embodiment, the tetracycline compound of the invention or a pharmaceutically acceptable salt
30 thereof is used to treat a skin wound, and other such disorders as described in, for example, U. S. Patent Nos. 5,827,840; 4,704,383; 4,935,412; 5,258,371; 5,308,839, 5,459,135; 5,532,227; and 6,015,804; each of which is incorporated herein by reference in its entirety.

In yet another embodiment, the tetracycline responsive disease or disorder is an aortic or vascular aneurysm in vascular tissue of a subject (e.g., a subject having or at risk of having an aortic or vascular aneurysm, etc.). The tetracycline compound or a pharmaceutically acceptable salt thereof may be effective to reduce the size of the vascular aneurysm or it may be administered to the subject prior to the onset of the vascular aneurysm such that the aneurysm is prevented. In one embodiment, the vascular tissue is an artery, e.g., the aorta, e.g., the abdominal aorta. In a further embodiment, the tetracycline compounds of the invention are used to treat disorders described in U. S. Patent Nos. 6,043,225 and 5,834,449, incorporated herein by reference in their entirety.

The compounds of the invention or a pharmaceutically acceptable salt thereof can be used alone or in combination with one or more therapeutic agent in the methods of the invention disclosed herein.

The language "in combination with" another therapeutic agent or treatment includes co-administration of the tetracycline compound and with the other therapeutic agent or treatment as either a single combination dosage form or as multiple, separate dosage forms, administration of the tetracycline compound first, followed by the other therapeutic agent or treatment and administration of the other therapeutic agent or treatment first, followed by the tetracycline compound.

The other therapeutic agent may be any agent that is known in the art to treat, prevent, or reduce the symptoms of a tetracycline-responsive disease or disorder. The choice of additional therapeutic agent(s) is based upon the particular tetracycline-responsive disease or disorder being treated. Such choice is within the knowledge of a treating physician. Furthermore, the other therapeutic agent may be any agent of benefit to the patient when administered in combination with the administration of a tetracycline compound.

As used herein, the term "subject" means a mammal in need of treatment or prevention, e.g., companion animals (e.g., dogs, cats, and the like), farm animals (e.g., cows, pigs, horses, sheep, goats and the like) and laboratory animals (e.g., rats, mice, guinea pigs and the like). Typically, the subject is a human in need of the specified treatment.

As used herein, the term "treating" or "treatment" refers to obtaining desired pharmacological and/or physiological effect. The effect can include achieving,

partially or substantially, one or more of the following results: partially or totally reducing the extent of the disease, disorder or syndrome; ameliorating or improving a clinical symptom or indicator associated with the disorder; delaying, inhibiting or decreasing the likelihood of the progression of the disease, disorder or syndrome.

5 As used herein, "preventing" or "prevention" refers to reducing the likelihood of the onset or development of disease, disorder or syndrome.

 "Effective amount" means that amount of active compound agent that elicits the desired biological response in a subject. In one embodiment, the effective amount of a compound of the invention is from about 0.01 mg/kg/day to
10 about 1000 mg/kg/day, from about 0.1 mg/kg/day to about 100 mg/kg/day, or from about 0.5 mg/kg/day to about 50 mg/kg/day.

 The invention further includes the process for making the composition comprising mixing one or more of the present compounds and an optional pharmaceutically acceptable carrier; and includes those compositions resulting from
15 such a process, which process includes conventional pharmaceutical techniques.

 The compositions of the invention include ocular, oral, nasal, transdermal, topical with or without occlusion, intravenous (both bolus and infusion), inhalable, and injection (intraperitoneally, subcutaneously, intramuscularly, intratumorally, or parenterally) formulations. The composition may be in a dosage unit such as a
20 tablet, pill, capsule, powder, granule, liposome, ion exchange resin, sterile ocular solution, or ocular delivery device (such as a contact lens and the like facilitating immediate release, timed release, or sustained release), parenteral solution or suspension, metered aerosol or liquid spray, drop, ampoule, auto-injector device, or suppository; for administration ocularly, orally, intranasally, sublingually,
25 parenterally, or rectally, or by inhalation or insufflation.

 Compositions of the invention suitable for oral administration include solid forms such as pills, tablets, caplets, capsules (each including immediate release, timed release, and sustained release formulations), granules and powders; and, liquid forms such as solutions, syrups, elixirs, emulsions, and suspensions. Forms
30 useful for ocular administration include sterile solutions or ocular delivery devices. Forms useful for parenteral administration include sterile solutions, emulsions, and suspensions.

The compositions of the invention may be administered in a form suitable for once-weekly or once-monthly administration. For example, an insoluble salt of the active compound may be adapted to provide a depot preparation for intramuscular injection (e.g., a decanoate salt) or to provide a solution for
5 ophthalmic administration.

The dosage form containing the composition of the invention contains an effective amount of the active ingredient necessary to provide a therapeutic effect. The composition may contain from about 5,000 mg to about 0.5 mg (preferably, from about 1,000 mg to about 0.5 mg) of a compound of the invention or salt form
10 thereof and may be constituted into any form suitable for the selected mode of administration. The composition may be administered about 1 to about 5 times per day. Daily administration or post-periodic dosing may be employed.

For oral administration, the composition is preferably in the form of a tablet or capsule containing, e.g., 500 to 0.5 milligrams of the active compound. Dosages will
15 vary depending on factors associated with the particular patient being treated (e.g., age, weight, diet, and time of administration), the severity of the condition being treated, the compound being employed, the mode of administration, and the strength of the preparation.

The oral composition is preferably formulated as a homogeneous
20 composition, wherein the active ingredient is dispersed evenly throughout the mixture, which may be readily subdivided into dosage units containing equal amounts of a compound of the invention. Preferably, the compositions are prepared by mixing a compound of the invention (or pharmaceutically acceptable salt thereof) with one or more optionally present pharmaceutical carriers (such as a
25 starch, sugar, diluent, granulating agent, lubricant, glidant, binding agent, and disintegrating agent), one or more optionally present inert pharmaceutical excipients (such as water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents, and syrup), one or more optionally present conventional tableting ingredients (such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid,
30 magnesium stearate, dicalcium phosphate, and any of a variety of gums), and an optional diluent (such as water).

Binder agents include starch, gelatin, natural sugars (e.g., glucose and beta-lactose), corn sweeteners and natural and synthetic gums (e.g., acacia and

tragacanth). Disintegrating agents include starch, methyl cellulose, agar, and bentonite.

Tablets and capsules represent an advantageous oral dosage unit form. Tablets may be sugarcoated or filmcoated using standard techniques. Tablets may
5 also be coated or otherwise compounded to provide a prolonged, control-release therapeutic effect. The dosage form may comprise an inner dosage and an outer dosage component, wherein the outer component is in the form of an envelope over the inner component. The two components may further be separated by a layer
10 which resists disintegration in the stomach (such as an enteric layer) and permits the inner component to pass intact into the duodenum or a layer which delays or sustains release. A variety of enteric and non-enteric layer or coating materials (such as polymeric acids, shellacs, acetyl alcohol, and cellulose acetate or combinations thereof) may be used.

Compounds of the invention may also be administered via a slow release
15 composition; wherein the composition includes a compound of the invention and a biodegradable slow release carrier (e.g., a polymeric carrier) or a pharmaceutically acceptable non-biodegradable slow release carrier (e.g., an ion exchange carrier).

Biodegradable and non-biodegradable slow release carriers are well known in the art. Biodegradable carriers are used to form particles or matrices which retain
20 an active agent(s) and which slowly degrade/dissolve in a suitable environment (e.g., aqueous, acidic, basic and the like) to release the agent. Such particles degrade/dissolve in body fluids to release the active compound(s) therein. The particles are preferably nanoparticles or nanoemulsions (e.g., in the range of about 1 to 500 nm in diameter, preferably about 50-200 nm in diameter, and most
25 preferably about 100 nm in diameter). In a process for preparing a slow release composition, a slow release carrier and a compound of the invention are first dissolved or dispersed in an organic solvent. The resulting mixture is added into an aqueous solution containing an optional surface-active agent(s) to produce an emulsion. The organic solvent is then evaporated from the emulsion to provide a
30 colloidal suspension of particles containing the slow release carrier and the compound of the invention.

The compound disclosed herein may be incorporated for administration orally or by injection in a liquid form such as aqueous solutions, suitably flavored

5 syrups, aqueous or oil suspensions, flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil and the like, or in elixirs or similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions, include synthetic and natural gums such as tragacanth, acacia,
10 alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone, and gelatin. The liquid forms in suitably flavored suspending or dispersing agents may also include synthetic and natural gums. For parenteral administration, sterile suspensions and solutions are desired. Isotonic preparations, which generally contain suitable preservatives, are employed when intravenous
15 administration is desired.

The compounds may be administered parenterally *via* injection. A parenteral formulation may consist of the active ingredient dissolved in or mixed with an appropriate inert liquid carrier. Acceptable liquid carriers usually comprise aqueous solvents and other optional ingredients for aiding solubility or preservation.
20 Such aqueous solvents include sterile water, Ringer's solution, or an isotonic aqueous saline solution. Other optional ingredients include vegetable oils (such as peanut oil, cottonseed oil, and sesame oil), and organic solvents (such as solketal, glycerol, and formyl). A sterile, non-volatile oil may be employed as a solvent or suspending agent. The parenteral formulation is prepared by dissolving or
25 suspending the active ingredient in the liquid carrier whereby the final dosage unit contains from 0.005 to 10% by weight of the active ingredient. Other additives include preservatives, isotonizers, solubilizers, stabilizers, and pain-soothing agents. Injectable suspensions may also be prepared, in which case appropriate liquid carriers, suspending agents and the like may be employed.

25 Compounds of the invention may be administered intranasally using a suitable intranasal vehicle.

In another embodiment, the compounds of this invention may be administered directly to the lungs by inhalation.

30 Compounds of the invention may also be administered topically or enhanced by using a suitable topical transdermal vehicle or a transdermal patch.

For ocular administration, the composition is preferably in the form of an ophthalmic composition. The ophthalmic compositions are preferably formulated as eye-drop formulations and filled in appropriate containers to facilitate administration

to the eye, for example a dropper fitted with a suitable pipette. Preferably, the compositions are sterile and aqueous based, using purified water. In addition to the compound of the invention, an ophthalmic composition may contain one or more of:

a) a surfactant such as a polyoxyethylene fatty acid ester; b) a thickening agents such as cellulose, cellulose derivatives, carboxyvinyl polymers, polyvinyl polymers, and polyvinylpyrrolidones, typically at a concentration in the range of about 0.05 to about 5.0% (wt/vol); c) (as an alternative to or in addition to storing the composition in a container containing nitrogen and optionally including a free oxygen absorber such as Fe), an anti-oxidant such as butylated hydroxyanisol, ascorbic acid, sodium thiosulfate, or butylated hydroxytoluene at a concentration of about 0.00005 to about 0.1% (wt/vol); d) ethanol at a concentration of about 0.01 to 0.5% (wt/vol); and e) other excipients such as an isotonic agent, buffer, preservative, and/or pH-controlling agent. The pH of the ophthalmic composition is desirably within the range of 4 to 8.

In certain embodiments, the composition of this invention includes one or more additional agents. The other therapeutic agent may be any agent that is capable of treating, preventing or reducing the symptoms of a tetracycline-responsive disease or disorder. Alternatively, the other therapeutic agent may be any agent of benefit to a patient when administered in combination with the tetracycline compound in this invention.

While this invention has been particularly shown and described with references to example embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

EXEMPLIFICATION

The following abbreviations and the terms have the indicated meanings:


Abbreviation/Term	Meaning
Ac	acetyl
AIBN	2,2'-azobis(2-methylpropionitrile)
aq	aqueous
Bn	benzyl
brine	saturated aqueous sodium chloride
Boc	<i>tert</i> -butoxy carbonyl or <i>t</i> -butoxy carbonyl

(Boc) ₂ O	di- <i>tert</i> -butyl dicarbonate
BBr ₃	boron tribromide
Bu	butyl
Cbz	benzyloxycarbonyl
CH ₂ Cl ₂	methylene chloride
CH ₃ CN or MeCN	acetonitrile
Cy	tricyclohexylphosphine
dba	dibenzylideneacetone
DIBAL-H	diisobutylaluminum hydride
DIEA	<i>N,N</i> -diisopropylethylamine
DMAP	4-(dimethylamino)pyridine
DME	1,2-dimethoxyethane
DMF	<i>N,N</i> -dimethylformamide
DMPU	1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidine
DMSO	dimethyl sulfoxide
EDC	<i>N</i> -(3-dimethylaminopropyl)- <i>N'</i> -ethylcarbodiimide
ESI	electrospray ionization
equiv.	equivalent
Et	ethyl
Et ₂ O	ethyl ether
EtOAc	ethyl acetate
h, hr	hour
HCl	hydrochloric acid
KHPO ₄	potassium hydrogenphosphate
HPLC	high performance liquid chromatography
HOBt	1-hydroxybenzotriazole
<i>i</i>	iso
IBX	2-iodoxybenzoic acid
LDA	lithium diisopropylamide
LHMDS	lithium bis(trimethylsilyl)amide
LTMP	lithium 2,2,6,6-tetramethylpiperidide
Me	methyl
MeOH	methanol
MeI	methyl iodide

min	minute
Ms	methanesulfonyl
MS	mass spectrum
MTBE	methyl <i>tert</i> -butyl ether
MW	molecular weight
NaHCO ₃	sodium bicarbonate
NaOH	sodium hydroxide
Na ₂ SO ₄	sodium sulfate
NBS	<i>N</i> -bromosuccinimide
NCS	<i>N</i> -chlorosuccinimide
NMR	nuclear magnetic resonance spectrometry
Ph	phenyl
Pr	propyl
<i>s</i>	secondary
<i>t</i>	tertiary
RP	reverse phase
TMEDA	tetramethylethylenediamine
TBS	<i>tert</i> -butyldimethylsilyl
TEA	triethylamine
Tf	trifluoromethanesulfonyl
TFA	trifluoroacetic acid
TFAA	trifluoroacetic anhydride
THF	tetrahydrofuran
TLC	thin layer chromatography
Ts	<i>para</i> -toluenesulfonyl
TsOH	<i>para</i> -toluenesulfonic acid
Xantphos	4,5-bis(diphenylphosphino)-9,9-dimethylxanthene

Example 1. Synthesis of Compounds of Structural Formula (I).


The compounds of the invention can be prepared according the synthetic scheme shown in Scheme 1.



Scheme 1


Specific conditions for the reactions depicted in Scheme 1 are provided in
 5 the following examples.

Compound **1**




To a THF solution of 5-fluoro-2-methoxybenzoic acid (500 mg, 2.94 mmol, Aldrich 523097) cooled at -78 °C was added a THF solution of *s*-BuLi (4.60 mL, 1.40 M, 6.44 mmol, 2.2 equiv) and TMEDA (0.97 mL, 6.47 mmol, 2.2 equiv). The reaction was stirred at -78 °C for 2 h. MeI (1.10 mL, 17.64 mmol, 6.0 equiv) was added to the reaction mixture dropwise. The reaction was allowed to warm to 25 °C over 1 h and stirred at 25 °C for 1 h. NaOH (6 N, 20 mL) was added. The resulting mixture was extracted with *t*-butylmethyl ether (20 mL × 2). The aqueous layer was acidified with HCl (6 N) to pH 1 and extracted with EtOAc (20 mL × 4). The combined EtOAc extracts were dried (Na₂SO₄) and concentrated to give 510 mg of crude product **1**: ¹H NMR (400 MHz, CDCl₃) δ 7.06 (dd, *J* = 9.8, 8.5 Hz, 1 H), 6.75 (dd, *J* = 9.8, 3.7 Hz, 1 H), 3.86 (s, 3 H), 2.34 (d, *J* = 2.4 Hz, 3 H); MS (ESI) *m/z* 185.12 (M+H).

Compound **2**




Oxalyl chloride (0.95 mL, 11.10 mmol, 5.5 equiv) was added to CH₂Cl₂ solution (15 mL, anhydrous) of **1** (510 mg, 2.00 mmol). DMF (0.1 mL) was added to the resulting mixture. The reaction was stirred at 25 °C for 1 h and concentrated. The resulting solid was re-dissolved in 15 mL of anhydrous CH₂Cl₂. Phenol (520 mg, 5.50 mmol, 2.8 equiv), DMAP (670 mg, 5.6 mmol, 2.8 equiv), and triethylamine (1.90 mL, 13.90 mmol, 7.0 equiv) were added to the reaction mixture. The reaction was stirred at 25 °C for 12 h and concentrated. EtOAc and H₂O were added to the residue. The organic layer was washed with NaOH (1 N), H₂O, and brine, dried (Na₂SO₄), and concentrated. Flash chromatography on silica gel (40:1 hexanes/EtOAc) yielded 400 mg of compound **2** (52% for 2 steps): ¹H NMR (400 MHz, CDCl₃) δ 7.47-7.41 (m, 2 H), 7.31-7.24 (m, 3 H), 7.08 (dd, *J* = 9.2, 9.2 Hz, 1 H), 6.77 (dd, *J* = 9.2, 3.7 Hz, 1 H), 3.88 (s, 3 H), 2.36 (d, *J* = 2.3 Hz, 3 H); MS (ESI) *m/z* 261.12 (M+H).

Compound 3




BBR₃ (1.85 mL, 1 M, 1.85 mmol, 1.2 equiv) was added to a CH₂Cl₂ solution
5 (8 mL) of **2** (400 mg, 1.54 mmol) at -78 °C. The reaction was stirred from -78 °C to
25 °C for 1.5 h, quenched with saturated NaHCO₃ and concentrated. EtOAc and
H₂O were added to the reaction mixture. The aqueous layer was extracted with
EtOAc. The combined EtOAc extracts were dried (Na₂SO₄) and concentrated to
yield 360 mg of crude **3**; ¹H NMR (400 MHz, CDCl₃) δ 10.66 (s, 1 H), 7.50-7.44
10 (m, 2 H), 7.36-7.31 (m, 1 H), 7.26-7.18 (m, 3 H), 6.86 (dd, *J* = 9.3, 4.9 Hz, 1 H),
2.60 (d, *J* = 2.4 Hz, 3 H); MS (ESI) *m/z* 245.11 (M-H).

Compound 4



15 Boc₂O (350 mg, 1.60 mmol, 1.1 equiv) and DMAP (20 mg, 0.16 mmol, 0.1
equiv) were added to a CH₂Cl₂ solution of crude **3** (360 mg). The reaction was
stirred at 25 °C for 1.5 h and concentrated. Flash chromatography on silica gel (35:1
hexanes/EtOAc) yielded 400 mg of compound **4** (94% for 2 steps); ¹H NMR (400
20 MHz, CDCl₃) δ 7.46-7.41 (m, 2 H), 7.31-7.23 (m, 3 H) 7.18 (dd, *J* = 8.8, 8.7 Hz, 1
H), 7.10 (dd, *J* = 8.8, 4.4 Hz, 1 H), 2.41 (d, *J* = 2.3 Hz, 3 H), 1.44 (s, 9 H); MS (ESI)
m/z 345.18 (M-H).


Compound 5



25

A THF solution (6 mL) of **4** (487 mg, 1.40 mmol, 2.0 equiv) was added to a THF solution (5 mL) of LDA (6.30 mL, 10% wt, 4.20 mmol, 6.0 equiv) and TMEDA (1.70 mL, 11.20 mmol, 16.0 equiv) at -78 °C. The reaction was stirred at
5 -78 °C for 5 min. A THF solution of enone (**3**) (339 mg, 0.70 mmol, 1.0 equiv) was added to the reaction mixture dropwise. The reaction was stirred from -78 °C to 25 °C for 1 h, quenched with saturated NH₄Cl, and extracted with EtOAc. The combined EtOAc extracts were dried (Na₂SO₄) and concentrated to yield the crude product. Preparative reverse phase HPLC purification on a Waters Autopurification
10 system using a Sunfire Prep C18 OBD column [5 µm, 19 × 50 mm; flow rate, 20 mL/min; Solvent A: H₂O with 0.1% HCO₂H; Solvent B: CH₃CN with 0.1% HCO₂H, injection volume: 4.0 mL (CH₃CN); gradient: 80→100% B over 15 min; mass-directed fraction collection] Fractions with the desired MW, eluting at 6.3-8.0 min, were collected and concentrated on a RotaVap at rt to remove most of the acetonitrile. The resulting mostly aqueous solution was extracted with EtOAc. The combined EtOAc extracts were dried (Na₂SO₄) and concentrated to give 185 mg of pure **5** (35%): ¹H NMR (400 MHz, CDCl₃) δ 15.67 (s, 1 H), 7.51-7.46 (m, 2 H), 7.39-7.29 (m, 3 H), 7.21 (dd, *J* = 8.9, 8.9 Hz, 1 H), 7.03 (dd, *J* = 8.9, 4.0 Hz, 1 H), 5.34 (s, 2 H), 3.93 (d, *J* = 10.4 Hz, 1 H), 3.30-3.21 (m, 1 H), 3.10-3.00 (m, 1 H),
20 2.57-2.41 (m, 3 H), 2.48 (s, 6 H), 2.17-2.12 (m, 1 H), 1.53 (s, 9 H), 0.82 (s, 9 H), 0.26 (s, 3 H), 0.12 (s, 3 H); MS (ESI) *m/z* 735.45 (M+H).

Compound **6**




25

Aqueous HF (3 mL, 48%) and TFA (4 µL) were added to a CH₃CN solution (7 mL) of **5** (210 mg, 0.29 mmol) in a polypropylene tube at 25 °C. The reaction was stirred at 25 °C for 18 h. The resulting mixture was poured into an aqueous solution of K₂HPO₄ (21 g, dissolved in 150 mL water). The mixture was extracted with EtOAc.
30 The combined EtOAc extracts were dried (Na₂SO₄) and concentrated to yield


180 mg of crude **6**: ^1H NMR (400 MHz, CDCl_3) δ 14.64 (s, 1 H), 11.47 (s, 1 H), 7.49-7.45 (m, 2 H), 7.39-7.32 (m, 3 H), 7.14 (dd, $J = 9.2, 8.8$ Hz, 1 H), 6.77 (dd, $J = 9.2, 4.3$ Hz, 1 H), 5.36 (s, 2 H), 3.68 (d, $J = 3.7$ Hz, 1 H), 3.09 (dd, $J = 15.6, 4.6$ Hz, 1 H), 3.02-2.92 (m, 1 H), 2.84-2.79 (m, 1 H), 2.49 (s, 6 H), 2.34-2.22 (m, 1 H), 2.09-2.02 (m, 1 H), 1.55-1.44 (m, 1 H); MS (ESI) m/z 521.30 (M+H).

Compound 7




Palladium on carbon (35 mg, 10 wt%) was added to a MeOH/dioxane solution (4 mL/4 mL) of crude **6** (180 mg). The reaction was purged with hydrogen and stirred under H_2 (balloon) at 25 °C for 1 h. The reaction mixture was filtered through a small Celite plug. The filtrate was concentrated to yield the crude product. Preparative reverse phase HPLC purification on a Waters Autopurification system using a Phenomenex Polymerx 10 μ RP-1 100A column [10 μm , 150 $\times$ 21.20 mm; flow rate, 20 mL/min; Solvent A: 0.05 N HCl/water; Solvent B: CH_3CN ; injection volume: 4.0 mL (0.05 N HCl/water), gradient: 0 $\rightarrow$ 100% B over 15 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 6.4-8.2 min, were collected and freeze-dried to yield 51 mg of compound **7** (41% for 2 steps): ^1H NMR (400 MHz, CD_3OD) δ 7.26 (dd, $J = 9.2, 9.2$ Hz, 1 H), 6.80 (dd, $J = 9.2, 4.3$ Hz, 1 H), 4.09 (br s, 1 H), 3.14 (dd, $J = 15.0, 4.6$ Hz, 1 H), 3.04 (s, 3 H), 2.96 (s, 3 H), 3.09-2.91 (m, 2 H), 2.31-2.18 (m, 2 H), 1.68-1.56 (m, 1 H); MS (ESI) m/z 433.28 (M+H).

Compound 8




A mixture of HNO₃ (8.5 μ L, 69%) and H₂SO₄ (0.5 mL) was added to a H₂SO₄ solution (1 mL) of **7** (51 mg, 0.12 mmol) at 0 °C. The reaction was stirred at 0 °C for 30 min. The resulting mixture was added dropwise to vigorously stirred diethyl ether (60 mL). The suspension was filtered through a small Celite pad and washed several times with more diethyl ether. The Celite pad was then eluted with MeOH until the eluent became colorless. The yellow MeOH eluent was collected and concentrated under reduced pressure to afford crude **8**: ¹H NMR (400 MHz, CD₃OD) δ 8.03 (d, J = 8.5 Hz, 1 H), 4.09 (br s, 1 H), 3.50-2.97 (m, 3 H), 3.04 (s, 3 H), 2.96 (s, 3 H), 2.46-2.36 (m, 1 H), 2.29-2.20 (m, 1 H), 1.71-1.59 (m, 1 H); MS (ESI) m/z 478.20 (M+H).

Compound **9**




Palladium on carbon (12 mg, 10 wt%) was added to a MeOH solution (4 mL) of crude **8**. The reaction was purged with hydrogen and stirred under H₂ (balloon) at 25 °C for 2 h. The catalyst was filtered off with a small Celite pad. The filtrate was concentrated to yield crude **9**. Preparative reverse phase HPLC purification on a Waters Autopurification system using a Phenomenex Polymerx 10 μ RP-1 100A column [10 μ m, 150 $\times$ 21.20 mm; flow rate, 20 mL/min; Solvent A: 0.05 N HCl; Solvent B: CH₃CN; injection volume: 4.0 mL (0.05 N HCl/water); gradient: 0 $\rightarrow$ 100% B over 15 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 5.0-6.6 min, were collected and freeze-dried to yield 43 mg of pure **9** (81% for 2 steps): ¹H NMR (400 MHz, CD₃OD) δ 7.43 (d, J = 8.5 Hz, 1 H), 4.11 (br s, 1 H), 3.22-3.16 (m, 1 H), 3.15-3.08 (m, 1 H), 3.06-2.95 (m, 1 H), 3.04 (s, 3 H), 2.96 (s, 3 H), 2.40-2.31 (m, 1 H), 2.28-2.21 (m, 1 H), 1.71-1.59 (m, 1 H); MS (ESI) m/z 448.24 (M+H).

Compound **11**



2-*t*-Butylaminoacetylchloride hydrochloride (4.2 mg, 0.022 mmol, 2.0 equiv) was added to a DMF solution (0.1 mL) of **9** (5 mg, 0.011 mmol) at 25 °C. The reaction was stirred at 25 °C for 30 min. The reaction mixture was diluted with 0.05 N HCl (2 mL) and injected into a Waters Autopurification system equipped with a Phenomenex Polymerx 10 μ RP-1 100A column [10 μ m, 150 $\times$ 21.20 mm; flow rate, 20 mL/min; Solvent A: 0.05 N HCl; Solvent B: CH_3CN ; gradient: 0 $\rightarrow$ 100% B over 20 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 6.4-7.0 min, were collected and freeze-dried to yield 3.9 mg of pure **11** (62%); ^1H NMR (400 MHz, CD_3OD) δ 8.25 (d, J = 11.0 Hz, 1 H), 4.11 (br s, 1 H), 4.09 (s, 2 H), 3.22-2.86 (m, 3 H), 3.05 (s, 3 H), 2.97 (s, 3 H), 2.33-2.20 (m, 2 H), 1.69-1.57 (m, 1 H), 1.42 (s, 9 H); MS (ESI) m/z 561.39 (M+H).

Compound **32**




Anhydrous Na_2CO_3 (16 mg, 0.15 mmol, 5.5 equiv) was added to an anhydrous DMPU/acetonitrile (150 μL /50 μL) solution of **9** (12 mg, 0.027 mmol). Bromoacetyl bromide (2.8 μL , 0.032 mmol, 1.2 equiv) was added to the mixture. The reaction was stirred at 25 °C for 10 min. LC/MS analysis indicated complete formation of intermediate **10**. Azetidine (36 μL , 0.54 mmol, 20 equiv) was added to the reaction mixture. The reaction was stirred at 25 °C for 2 h. The reaction mixture was concentrated and acidified with HCl (0.5 N in MeOH, 0.7 mL). The resulting mixture was added dropwise to vigorously stirred diethyl ether (10 mL). The suspension was filtered through a small Celite pad and washed several times

with more diethyl ether. The Celite pad was then eluted with MeOH until the eluent became colorless. The yellow MeOH eluent was collected and concentrated under reduced pressure to afford crude **32**. Preparative reverse phase HPLC purification on a Waters Autopurification system using a Phenomenex Polymerx 10 μ RP-1
5 100A column [10 μ m, 150 $\times$ 21.20 mm; flow rate, 20 mL/min; Solvent A: 0.05 N HCl; Solvent B: CH₃CN; injection volume: 2.0 mL (0.05 N HCl/water); gradient: 10 $\rightarrow$ 20% B over 30 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 10.8-12.5 min, were collected and freeze-dried to yield
10 2.0 mg of pure **32**: ¹H NMR (400 MHz, CD₃OD) δ 8.18 (d, J = 11.0 Hz, 1 H), 4.41-4.31 (m, 2 H), 4.32 (s, 2 H), 4.24-4.13 (m, 2), 4.08 (br s, 1 H), 3.18-2.86 (m, 3 H), 3.03 (s, 3 H), 2.95 (s, 3 H), 2.71-2.57 (m, 1 H), 2.54-2.42 (m, 1 H), 2.33-2.16 (m, 2 H), 1.69-1.57 (m, 1 H); MS (ESI) m/z 545.20 (M+H).


Compounds 12 -31 and **Compounds 33-46** are prepared similarly to **Compounds**
15 **11** or **32**, substituting the appropriate acyl halide for 2-*t*-Butylaminoacetylchloride in the synthesis of **Compound 11** or cyclic amine for azetidine in the synthesis of **Compound 32**.

Compound **12**
20




¹H NMR (400 MHz, CD₃OD) δ 8.23 (d, J = 11.2 Hz, 1 H), 4.08 (s, 3 H),
3.17-2.97 (m, 11 H), 2.31 (dd, J = 14.8, 14.8 Hz, 1 H), 2.24 (ddd, J = 14.0, 5.2, 2.8
25 Hz, 1H), 1.79-1.72 (m, 2 H), 1.66 (ddd, J = 13.6, 13.6, 13.6 Hz, 1 H), 1.05 (t, J =
7.2 Hz, 3 H); MS (ESI) m/z 547.2 (M+H).

Compound **13**



^1H NMR (400 MHz, CD_3OD) δ 8.23 (d, $J = 11.2$ Hz, 1 H), 4.08 (s, 3 H),
3.16-2.97 (m, 11 H), 2.30 (dd, $J = 14.8, 14.8$ Hz, 1 H), 2.24 (ddd, $J = 14.4, 5.2, 2.8$
5 Hz, 1H), 1.75-1.69 (m, 2 H), 1.66 (ddd, $J = 13.6, 13.6, 13.6$ Hz, 1 H), 1.49-1.41 (m,
2 H), 1.01 (t, $J = 7.2$ Hz, 3 H); MS (ESI) m/z 561.2 (M+H).


Compound 14



10

^1H NMR (400 MHz, CD_3OD) δ 8.21 (d, $J = 11.2$ Hz, 1 H), 4.08 (s, 1 H),
4.06 (s, 2 H), 3.16-2.96 (m, 11 H), 2.28 (dd, $J = 14.8, 14.8$ Hz, 1 H), 2.22 (ddd, $J =$
14.4, 5.2, 2.8 Hz, 1H), 1.77-1.71 (m, 2 H), 1.66 (ddd, $J = 14.0, 14.0, 14.0$ Hz, 1 H),
15 1.43-1.35 (m, 6 H), 0.93 (t, $J = 7.2$ Hz, 3 H); MS (ESI) m/z 589.2 (M+H).


Compound 15



20


^1H NMR (400 MHz, CD_3OD) δ 8.23 (d, $J = 10.8$ Hz, 1 H), 4.09 (s, 2 H),
4.07 (s, 1 H), 3.15-2.95 (m, 11 H), 2.29 (dd, $J = 14.4, 14.4$ Hz, 1 H), 2.25 (ddd, $J =$
14.4, 5.2, 2.8 Hz, 1H), 1.66 (ddd, $J = 13.2, 13.2, 13.2$ Hz, 1 H), 1.10 (s, 9 H); MS
(ESI) m/z 575.2 (M+H).

Compound 16




¹H NMR (400 MHz, CD₃OD) δ 8.24 (d, *J* = 11.0 Hz, 1H), 4.08 (s, 2H), 4.01-3.89 (m, 1H), 3.50-3.42 (m, 1H), 3.20-2.84 (m, 9H), 2.30 (at, *J* = 14.7 Hz, 1H), 2.23-2.15 (m, 1H), 1.70-1.58 (m, 1H), 1.37 (d, *J* = 6.7 Hz, 6H); MS (ESI) *m/z* 547.25 (M+H).

Compound 17



¹H NMR (400 MHz, CD₃OD) δ 8.24 (d, *J* = 11.0 Hz, 1 H), 4.20 (s, 2 H), 4.09 (br s, 1 H), 3.19-3.13 (m, 1 H), 3.12-2.89 (m, 2 H), 2.89-2.38 (m, 1 H), 3.04 (s, 3 H), 2.96 (s, 3 H), 2.35-2.19 (m, 2 H), 1.71-1.59 (m, 1 H), 0.95 (br s, 2), 0.94 (br s, 2); MS (ESI) *m/z* 545.37 (M+H).

Compound 18




¹H NMR (400 MHz, CD₃OD) δ 8.23 (d, *J* = 10.8 Hz, 1 H), 4.09 (s, 3 H), 3.68-3.61 (m, 1 H), 3.16-2.97 (m, 9 H), 2.29 (dd, *J* = 14.4, 14.4 Hz, 1 H), 2.25 (ddd,

$J = 14.4, 5.2, 2.8$ Hz, 1H), 2.20-2.12 (m, 2 H), 1.98-1.91 (m, 2 H), 1.75-1.68 (m, 4 H), 1.66 (ddd, $J = 13.6, 13.6, 13.6$ Hz, 1 H); MS (ESI) m/z 573.1 (M+H).


Compound 19

5



¹H NMR (400 MHz, CD₃OD) δ 8.26 (d, *J* = 11.0 Hz, 1 H), 4.09 (br s, 3 H), 3.19-2.93 (m, 5 H), 3.04 (s, 3 H), 2.96 (s, 3 H), 2.35-2.26 (m, 1 H), 2.25-2.18 (m, 1 H), 2.14-2.02 (m, 1 H), 1.71-1.59 (m, 1 H), 1.07 (d, *J* = 6.7, 6 H); MS (ESI) *m/z* 561.24 (M+H).


Compound 20



15

¹H NMR (400 MHz, CD₃OD) δ 8.24 (d, *J* = 11.0 Hz, 1 H), 4.11 (s, 2 H), 4.08 (br s, 1 H), 3.22-2.92 (m, 5 H), 3.03 (s, 3 H), 2.95 (s, 3 H), 2.33-2.24 (m, 1 H), 2.24-2.17 (m, 1 H), 1.69-1.58 (m, 1 H), 1.17-1.07 (m, 1 H), 0.77-0.71 (m, 2 H), 0.46-0.40 (m, 2 H); MS (ESI) *m/z* 559.23 (M+H).


Compound **21**.



^1H NMR (400 MHz, CD_3OD) δ 8.25 (d, $J = 11.0$ Hz, 1H), 4.12 (s, 2H), 4.09 (s, 1H), 3.72-3.67 (m, 2H), 3.43 (s, 3H), 3.19-2.92 (m, 11H), 2.35-2.18 (m, 2H) 1.71-1.58 (m, 1H); MS (ESI) m/z 563.23 (M+H).

5


Compound **22**



10 ^1H NMR (400 MHz, CD_3OD) δ 8.25 (d, $J = 11.0$ Hz, 1H), 4.22 (s, 2H), 4.14-4.05 (m, 3H), 3.18-2.84 (m, 9H), 2.34-2.17 (m, 2H), 1.70-1.57 (m, 1H); MS (ESI) m/z 587.28 (M+H).


Compound **23**

15



20 ^1H NMR (400 MHz, CD_3OD) δ 8.24 (d, $J = 11.0$ Hz, 1H), 4.24 (s, 2H), 4.09 (s, 1H), 3.14-2.93 (m, 15H), 2.24-2.18 (m, 2H), 1.65 (dt, $J = 13.4, 11.6$ Hz, 1H); MS (ESI) m/z 533.17 (M+H).


Compound **24**



^1H NMR (400 MHz, CD_3OD) δ 8.23 (d, $J = 10.4$ Hz, 1H), 4.29 (d, $J = 16.5$ Hz, 1H), 4.18 (d, $J = 15.9$ Hz, 1H), 4.09 (s, 1H), 3.19-2.89 (m, 14H), 2.36-2.17 (m, 2H), 1.70-1.58 (m, 1H), 1.38 (t, $J = 7.32$ Hz, 3H); MS (ESI) m/z 547.25 (M+H).

5


Compound 25



10 ^1H NMR (400 MHz, CD_3OD) δ 8.21 (d, $J = 10.8$ Hz, 1 H), 4.25 (s, 2 H), 4.10 (s, 1 H), 3.35 (t, $J = 7.2$ Hz, 3 H), 3.34 (t, $J = 7.2$ Hz, 3 H), 3.13-2.99 (m, 9 H), 2.31 (dd, $J = 14.8, 14.8$ Hz, 1 H), 2.27 (ddd, $J = 14.8, 5.2, 2.8$ Hz, 1H), 1.78-1.74 (m, 2 H), 1.68 (ddd, $J = 13.6, 13.6, 13.6$ Hz, 1 H), 1.38 (t, $J = 7.2$ Hz, 6 H); MS (ESI) m/z 561.2 (M+H).


15

Compound 26




20 ^1H NMR (400 MHz, CD_3OD) δ 8.23 (d, $J = 11.2$ Hz, 1 H), 4.10 (s, 3 H), 3.16-2.96 (m, 11 H), 2.31 (dd, $J = 14.4, 14.4$ Hz, 1 H), 2.24 (ddd, $J = 14.4, 5.2, 2.8$ Hz, 1H), 1.78-1.71 (m, 2 H), 1.66 (ddd, $J = 14.0, 14.0, 14.0$ Hz, 1 H), 1.45-1.38 (m, 4 H), 0.98 (t, $J = 7.2$ Hz, 3 H); MS (ESI) m/z 575.2 (M+H).

25 Compound 27



¹H NMR (400 MHz, CD₃OD) δ 8.24 (d, *J* = 10.8 Hz, 1 H), 4.09 (s, 3 H),
3.59 (t, *J* = 5.6 Hz, 2 H), 3.40 (s, 3 H), 3.23 (t, *J* = 5.6 Hz, 2 H), 3.15-2.94 (m, 9 H),
2.32 (dd, *J* = 15.2, 15.2 Hz, 1 H), 2.24 (ddd, *J* = 14.0, 5.2, 2.8 Hz, 1H), 2.08-2.02
5 2.32 (dd, *J* = 15.2, 15.2 Hz, 1 H), 1.66 (ddd, *J* = 15.2, 15.2, 15.2 Hz, 1 H); MS (ESI) *m/z* 577.2 (M+H).


Compound 28



10

¹H NMR (400 MHz, CD₃OD) δ 8.24 (d, *J* = 10.8 Hz, 1 H), 4.32 (d, *J* = 8.0
Hz, 1 H), 4.21 (d, *J* = 8.0 Hz, 1 H), 4.10 (s, 1 H), 3.18-2.99 (m, 9 H), 3.01 (s, 3 H),
2.33 (dd, *J* = 14.8, 14.8 Hz, 1 H), 2.29 (ddd, *J* = 15.2, 5.2, 2.8 Hz, 1H), 1.78-1.74
15 (m, 2 H), 1.88-1.81 (m, 2 H), 1.68 (ddd, *J* = 15.6, 15.6, 15.6 Hz, 1 H), 1.08 (t, *J* =
7.2 Hz, 3 H); MS (ESI) *m/z* 561.2 (M+H).

Compound 29




20

¹H NMR (400 MHz, CD₃OD) δ 8.25 (d, *J* = 10.8 Hz, 1 H), 4.10 (s, 3 H),
3.18-2.98 (m, 11 H), 2.31 (dd, *J* = 14.8, 14.8 Hz, 1 H), 2.26 (ddd, *J* = 14.4, 5.2, 2.8

Hz, 1H), 1.78-1.74 (m, 2 H), 1.66 (ddd, $J = 13.6, 13.6, 13.6$ Hz, 1 H), 1.42-1.30 (m, 8 H), 0.94 (t, $J = 6.8$ Hz, 3 H); MS (ESI) m/z 603.2 (M+H).

Compound **30**


5



^1H NMR (400 MHz, CD_3OD) δ 8.32 (d, $J = 10.4$ Hz, 1 H), 7.38-7.34 (m, 2 H), 7.10-7.06 (m, 3 H), 4.17 (s, 2 H), 4.10 (s, 1 H), 3.18-2.99 (m, 11 H), 2.29 (dd, $J = 15.6, 15.6$ Hz, 1 H), 2.25 (ddd, $J = 14.8, 5.2, 2.8$ Hz, 1 H), 1.66 (ddd, $J = 14.8, 14.8, 14.8$ Hz, 1 H); MS (ESI) m/z 581.1 (M+H).

Compound **31**


15



^1H NMR (400 MHz, CD_3OD) δ 8.25 (d, $J = 10.8$ Hz, 1 H), 4.36 (d, $J = 8.0$ Hz, 1 H), 4.21 (d, $J = 8.0$ Hz, 1 H), 4.10 (s, 1 H), 3.68-3.61 (m, 1 H), 3.18-2.98 (m, 9 H), 3.00 (s, 3 H), 2.29 (dd, $J = 14.4, 14.4$ Hz, 1 H), 2.20-2.10 (m, 3 H), 1.96-1.89 (m, 2 H), 1.78-1.68 (m, 4 H), 1.66 (ddd, $J = 14.4, 14.4, 14.4$ Hz, 1 H); MS (ESI) m/z 587.2 (M+H).


Compound **33**

20



¹H NMR (400 MHz, CD₃OD) δ 8.20 (d, *J* = 11.0 Hz, 1 H), 5.54-5.33 (m, 2 H), 4.71-4.37 (m, 4 H), 4.40 (s, 2 H), 4.06 (br s, 1 H), 3.17-2.92 (m, 3 H), 2.99 (s, 6 H), 2.33-2.24 (m, 1 H), 2.23-2.16 (m, 1 H), 1.70-1.58 (m, 1 H); MS (ESI) *m/z* 563.20 (M+H).

Compound 34




10

¹H NMR (400 MHz, CD₃OD) δ 8.22 (d, *J* = 11.0 Hz, 1H), 4.33 (s, 2H), 4.10 (s, 1H), 3.83-3.72 (m, 2H), 3.25-2.89 (m, 12H), 2.32-2.00 (m, 6H), 1.69-1.56 (m, 1H); MS (ESI) *m/z* 559.39 (M+H).

15


Compound 35



20


¹H NMR (400 MHz, CD₃OD) δ 8.25 (d, *J* = 11.0 Hz, 1H), 5.54-5.31 (m, 1H), 4.39-4.20 (m, 2H), 4.09-4.01 (m, 1H), 3.40-3.30 (m, 2H), 3.09-2.89 (m, 12H), 2.50-2.34 (m, 2H), 2.34-2.25 (m, 1H), 2.24-2.16 (m, 1H), 1.71-1.58 (m, 1H); MS (ESI) *m/z* 577.32 (M+H).

Compound **36**




5 ^1H NMR (400 MHz, CD_3OD) δ 8.23 (d, $J = 10.4$ Hz, 1H), 5.57-5.37 (m, 1H), 4.47-4.33 (m, 2H), 4.15-3.87 (m, 2H), 3.72-3.40 (m, 1H), 3.17-2.83 (m, 12H), 2.55-2.34 (m, 2H), 2.33-2.18 (m, 2H), 1.69-1.57 (m, 1H); MS (ESI) m/z 577.37 (M+H).

10 Compound **37**



15 ^1H NMR (400 MHz, CD_3OD) δ 8.28 (d, $J = 10.7$ Hz, 1H), 4.08 (s, 1H), 4.00-3.91 (m, 2H), 3.09-2.57 (m, 18H), 3.26-3.18 (m, 3H), 2.49-2.34 (m, 2H), 2.35-2.06 (m, 2H), 1.72-1.59 (m, 1H); MS (ESI) m/z 602.37 (M+H).

Compound **38**




20

^1H NMR (400 MHz, CD_3OD) δ 8.24 (d, $J = 11.0$ Hz, 1H), 4.46-4.32 (m, 2H), 4.26-4.16 (m, 1H), 4.08 (s, 2H), 4.00-3.72 (m, 2H), 3.18-2.91 (m, 16H), 2.68-

2.56 (m, 1H), 2.51-2.39 (m, 1H), 2.34-2.24 (m, 1H), 2.23-2.17 (m, 1H), 1.70-1.57 (m, 1H); MS (ESI) m/z 602.37 (M+H).

Compound **39**


5



^1H NMR (400 MHz, CD_3OD) δ 8.23 (d, $J = 11.0$ Hz, 1H), 4.62-4.54 (m, 1H), 4.48-4.24 (m, 2H), 4.08 (s, 1H), 3.99-3.69 (m, 3H), 3.50-3.40 (m, 1H), 3.17-2.90 (m, 9H), 2.44-2.11 (m, 4H), 2.10-2.00 (m, 1H), 1.69-1.56 (m, 1H); MS (ESI) m/z 575.27 (M+H).


Compound **40**

15




^1H NMR (400 MHz, CD_3OD) δ 8.23 (d, $J = 11.0$ Hz, 1H), 4.62-4.54 (m, 1H), 4.50-4.38 (m, 1H), 4.37-4.27 (m, 1H), 4.10 (s, 1H), 3.99-3.70 (m, 3H), 3.50-3.40 (m, 1H), 3.24-2.84 (m, 9H), 2.40-2.11 (m, 4H), 2.10-2.01 (m, 1H), 1.70-1.57 (m, 1H); MS (ESI) m/z 575.33 (M+H).

Compound **41**



¹H NMR (400 MHz, CD₃OD) δ 8.25 (d, *J* = 11.0 Hz, 1H), 4.54 (d, *J* = 16.5 Hz, 1H), 4.26 (d, *J* = 15.9 Hz, 1H), 4.09 (s, 1H), 3.95-3.81 (m, 2H), 3.81-3.75 (m, 1H), 3.69-3.62 (m, 1H), 3.35 (s, 3H), 3.23-2.92 (m, 9H), 2.35-2.04 (m, 6H), 1.91-1.80 (m, 1H), 1.71-1.59 (m, 1H); MS (ESI) *m/z* 603.35 (M+H).

Compound **42**




10

¹H NMR (400 MHz, CD₃OD) δ 8.24 (d, *J* = 11.0 Hz, 1H), 4.55 (d, *J* = 16.5 Hz, 1H), 4.27 (d, *J* = 16.5 Hz, 1H), 4.10 (s, 1H), 3.95-3.82 (m, 2H), 3.81-3.75 (m, 1H), 3.70-3.63 (m, 1H), 3.38 (s, 3H), 3.20-2.92 (m, 9H), 2.35-2.02 (m, 6H), 1.92-1.80 (m, 1H), 1.70-1.58 (m, 1H); MS (ESI) *m/z* 603.41 (M+H).

15


Compound **43**



20


¹H NMR (400 MHz, CD₃OD) δ 8.22 (d, *J* = 11.0 Hz, 1H), 4.19 (s, 2H), 4.09 (s, 1H), 3.65-3.58 (m, 2H), 3.19-2.92 (m, 10H), 2.34-2.18 (m, 2H), 2.02-1.79 (m, 6H), 1.69-1.50 (m, 2H); MS (ESI) *m/z* 573.35 (M+H).

25 Compound **44**



¹H NMR (400 MHz, CD₃OD) δ 8.24 (d, *J* = 11.0 Hz, 1H), 4.28 (s, 2H), 4.03-4.00 (m, 2H), 3.94-3.81 (m, 2H), 3.68-3.55 (m, 2H), 3.20-2.88 (m, 12H), 2.36-2.18 (m, 2H), 1.71-1.57 (m, 1H); MS (ESI) *m/z* 575.37 (M+H).

Compound 45




10

¹H NMR (400 MHz, CD₃OD, 2:1 mixture of diastereomers) δ 8.25 (d + d, *J* = 11.0 Hz, 1H), 4.29, 4.24 (s + s, 2H), 4.08 (s - s, 1H), 4.01-3.92 (m + m, 3H), 3.20-2.62 (m + m, 13H), 2.35-2.16 (m + m, 3H), 1.83-1.46 (m + m, 5H); MS (ESI) *m/z* 599.36 (M+H).

15


Compound 46



20


¹H NMR (400 MHz, CD₃OD) δ 8.29 (d, *J* = 11.0 Hz, 1H), 7.41 (s, 5H), 4.50-4.37 (m, 2H), 4.05 (s, 1H), 3.95-3.81 (m, 2H), 3.40-3.37 (m, 1H), 3.24-3.15 (m, 3H), 3.10-2.70 (m, 9H), 2.36-2.25 (m, 1H), 2.25-2.16 (m, 1H), 1.72-1.59 (m, 1H); MS (ESI) *m/z* 607.34 (M+H).

Compound 47




^1H NMR (400 MHz, CD_3OD) δ 8.15 (d, $J = 10.8$ Hz, 1 H), 4.00 (s, 1 H),
 3.99 (s, 2 H), 3.10-2.87 (m, 11 H), 2.32-2.12 (m, 2 H), 1.59-1.51 (m, 1 H), 1.26 (t, J
 5 = 7.2 Hz, 3 H); MS (ESI) m/z 533.1 ($\text{M}+\text{H}$).

Compound 48




^1H NMR (400 MHz, CD_3OD) δ 8.15 (d, $J = 11.2$ Hz, 1 H), 4.00 (s, 1 H),
 10 3.96 (s, 2 H), 3.08-2.87 (m, 11 H), 2.70-2.61 (m, 1 H), 2.23-2.09 (m, 4 H), 1.97-1.75
 (m, 4 H), 1.59-1.51 (m, 1 H); MS (ESI) m/z 572.2 ($\text{M}+\text{H}$).

Compound 49




15 ^1H NMR (400 MHz, CD_3OD) δ 8.26 (d, $J = 10.8$ Hz, 1 H), 4.10 (s, 3 H),
 3.21-2.97 (m, 11 H), 2.35-2.20 (m, 2 H), 2.15-2.05 (m, 1 H), 1.98-1.82 (m, 2 H),
 1.77-1.61 (m, 5 H), 1.35-1.26 (m, 2 H); MS (ESI) m/z 587.1 ($\text{M}+\text{H}$).

Compound 50



¹H NMR (400 MHz, CD₃OD) δ 8.26 (d, *J* = 10.8 Hz, 1 H), 4.16 (s, 1 H), 4.14 (s, 2 H), 3.20-2.95 (m, 11 H), 2.32-2.20 (m, 2 H), 1.88-1.59 (m, 6 H), 1.39-1.21 (m, 4 H), 1.12-1.02 (m, 2 H); MS (ESI) *m/z* 601.1 (M+H).


5 Compound 51



¹H NMR (400 MHz, CD₃OD) δ 8.14 (d, *J* = 10.8 Hz, 1 H), 4.00 (s, 1 H), 3.88 (s, 2 H), 3.77-3.73 (m, 1 H), 3.09-2.87 (m, 9 H), 2.29-2.10 (m, 6 H), 1.88-1.81 (m, 2 H), 1.59-1.50 (m, 1 H); MS (ESI) *m/z* 559.1 (M-H).

10


Compound 52



¹H NMR (400 MHz, CD₃OD) δ 8.24 (d, *J* = 10.8 Hz, 1 H), 4.10 (s, 3 H), 3.17-2.97 (m, 9 H), 2.32-2.09 (m, 4 H), 1.92-1.85 (m, 2 H), 1.75-1.63 (m, 2 H), 1.43-1.26 (m, 6 H); MS (ESI) *m/z* 587.2 (M+H).

15


Compound 53



¹H NMR (400 MHz, CD₃OD) δ 8.20 (d, *J* = 11.2 Hz, 1 H), 8.16 (d, *J* = 2.4
20 Hz, 1 H), 8.06 (d, *J* = 5.2 Hz, 1 H), 7.85-7.78 (m, 2 H), 4.27 (s, 2 H), 4.11 (s, 1 H),
3.18-2.98 (m, 9 H), 2.32-2.21 (m, 2 H), 1.70-1.60 (m, 1 H); MS (ESI) *m/z* 582.2
(M+H)

20


Compound 54



^1H NMR (400 MHz, CD_3OD) δ 8.24 (d, $J = 11.0$ Hz, 1 H), 4.31 (s, 2 H), 4.11 (s, 1 H), 3.22-2.88 (m, 9 H), 2.36-2.16 (m, 2 H), 1.70-1.56 (m, 1 H), 1.44 (s, 9 H); MS (ESI) m/z 577.41 (M+H).

5


Compound 55



^1H NMR (400 MHz, CD_3OD) δ 8.15 (d, $J = 10.8$ Hz, 1 H), 4.65 (t, $J = 4.8$ Hz, 2 H), 4.08 (s, 2 H), 4.00 (s, 1 H), 3.45 (t, $J = 4.4$ Hz, 1 H), 3.38 (t, $J = 5.6$ Hz, 1 H), 3.20-2.87 (m, 9 H), 2.25-2.09 (m, 2 H), 1.59-1.50 (m, 1 H); MS (ESI) m/z 551.0 (M+H).

10


Compound 56



^1H NMR (400 MHz, CD_3OD) δ 8.25 (d, $J = 11.2$ Hz, 1 H), 6.39 (tt, $J = 53.6$, 3.2 Hz, 1 H), 4.24 (s, 2 H), 4.13 (s, 1 H), 3.71 (td, $J = 15.2$, 2.8 Hz, 2 H), 3.19-2.91 (m, 9 H), 2.33-2.24 (m, 2 H), 1.70-1.60 (m, 1 H); MS (ESI) m/z 569.0 (M+H).

15


Compound 57



20

^1H NMR (400 MHz, CD_3OD) δ 8.21 (d, $J = 10.8$ Hz, 1 H), 4.01 (s, 1 H), 3.85 (s, 2 H), 3.73 (s, 3 H), 3.59-3.51 (m, 1 H), 3.12-2.87 (m, 9 H), 2.23-2.12 (m, 2 H), 1.88-1.50 (m, 9 H); MS (ESI) m/z 559.1 (M+H).

5 Compound 58




^1H NMR (400 MHz, CD_3OD) δ 8.24 (d, $J = 10.8$ Hz, 1 H), 4.48 (s, 2 H), 4.12 (s, 1 H), 4.10-4.07 (m, 2 H), 3.93-3.86 (m, 2 H), 3.19-2.90 (m, 9 H), 2.79-2.67 (m, 2 H), 2.37-2.21 (m, 2 H), 1.59-1.51 (m, 1 H); MS (ESI) m/z 595.0 (M+H).

10

Example 2. Synthesis of Compounds of Structural Formula (I), wherein R^1 and R^2 taken together with the nitrogen to which they are bonded form a monocyclic or bicyclic heteroaryl.


Scheme 2

15




The following compounds were prepared according to Scheme 2.

Compound **2-1**



5 To a 250 mL round bottom flask was added compound **3** (14.47g, 56.30 mmol, 1.0 equiv, crude), tetrabutylammonium bromide (0.90 g, 2.80 mmol, 0.05 equiv), 1,2-dichloroethane (60 mL), and water (60 mL). The clear bi-layer was cooled in a 20 °C water bath. Nitric acid (7.2 mL, 70 wt%, 112.60 mmol, 2.0 equiv)
10 was added. After the addition, the reaction temperature slowly rose to 26 °C. The reaction was stirred at room temperature overnight (19 hrs). TLC (heptane/EtOAc = 9.5/0.5) showed the reaction was complete. The organic layer was separated, washed with water (60mL x 2) and brine, and dried over anhydrous sodium sulfate. The solvent was removed to give compound **2-1** as a brown oil, which solidified on
15 standing (17.71 g, quantitative). The crude product was used directly for the next step.


Compound **2-2**



20 To a 250 mL round bottom flask was added compound **2-1** (17.7 g, 56.30 mmol 1.0 equiv), acetone (177 mL), anhydrous potassium carbonate (15.6 g, 113.00 mmol, 2.0 equiv), and potassium iodide (0.47 g, 2.80 mmol, 0.05 equiv). To the stirred suspension at room temperature was added benzyl bromide (7.03 mL, 59.10 mmol, 1.05 equiv). The suspension was then heated to 56 °C for 4 hrs. TLC
25 (heptane/EtOAc = 9/1) showed the reaction was complete. The solid was removed by filtration and washed with acetone (30 mL). The filtrate was concentrated to give a paste. The paste was partitioned between methyl t-butyl ether (MTBE, 120 mL) and water (80 mL). The organic layer was washed with water (80 mL) and


brine, dried over anhydrous sodium sulfate, and concentrated to give compound **2-2** as a brown oil (21.09 g, 98%). The crude product was used directly for the next step.

Compound **2-3**



5 To a 1 L round bottom flask was added compound **2-2** (21.08 g, 55.40 mmol, 1.0 equiv) and THF (230 mL). The solution was cooled in a cold water bath to 10 °C. To another 500 mL round bottom flask containing water (230 mL), sodium hydrosulfite (Na₂S₂O₄, 56.7 g, 276.80 mmol, 5.0 equiv) was added slowly with
10 stirring. The aqueous solution of sodium hydrosulfite was added to the THF solution of compound **2-2**. The temperature quickly rose from 10 °C to 20.4 °C after the addition. The yellow suspension was stirred while the cold water bath slowly warmed up to room temperature overnight to give an orange cloudy solution. The reaction temperature during this period was between 15 °C to 19 °C. TLC
15 (heptane/EtOAc = 9/1) showed the reaction was complete. The orange cloudy solution was diluted with EtOAc (460 mL). The organic layer was washed with water (150 mL x 2) and brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the crude product as a brown oil. The crude product was purified by flash silica gel column eluted with heptane/EtOAc 9/1
20 to yield the desired product **2-3** (15.83 g, 80%, 3 steps).


Compound **2-4**



25 To an NMP solution (50 mL) aniline **2-3** (10.02 g, 28.5 mmol, 1 equiv) was added allyl bromide (7.65 mL, 85.5 mmol, 3 equiv) and potassium carbonate (11.79 g, 85.5 mmol, 3 equiv). Potassium iodide (994.8 mg, 6 mmol, 0.2 equiv) was added and the reaction mixture was placed under nitrogen and heated to 100 °C. After 16 h, the reaction was cooled, diluted with water (60 mL), and extracted with EtOAc

(75 mL, then 2 x 50 mL). The combined organic extracts were washed with water (2 x 35 mL), were dried (Na₂SO₄), filtered, and concentrated to yield the crude product. Purification via flash column chromatography on silica gel (RediSep, 125 g, gradient 1-6% EtOAc in hexanes) gave 10.97 g of pure **2-4** (89%): ¹H NMR (400 MHz, CDCl₃) δ 7.42-7.30 (m, 7 H), 7.42-7.20 (m, 1 H), 7.00 (d, *J* = 8.5 Hz, 2 H), 6.72 (d, *J* = 11.0 Hz, 1 H), 5.77-5.70 (m, 2 H), 5.20-5.12 (m, 6 H), 3.81 (d, *J* = 6.1 Hz, 4 H), 2.26 (s, 3 H); MS (ESI) *m/z* 432.34 (M+H).

Compound **2-5**



10

A THF solution (6.5 mL) of **2-4** (875 mg, 2.03 mmol, 1.25 equiv) was added to a freshly-prepared solution of LDA in THF (0.051 M, 2.03 mmol, 40 mL, 1.25 equiv) and TMEDA (304 μL, 2.03 mmol, 1.25 equiv) at -78 °C. The reaction was stirred at -78 °C for 15 min. A THF solution (6.5 mL) of enone (784 mg, 1.62 mmol, 1.0 equiv) was added to the reaction mixture dropwise, followed by addition of LHMDS solution (1.0 M in THF, 2.03 mL, 2.03 mmol, 1.25 equiv). The reaction was stirred from -78 °C to -10 °C for 1 h, quenched with saturated NH₄Cl (6 mL), and warmed to 25 °C. The solution was poured into saturated NH₄Cl (20 mL) and extracted with EtOAc (2 x 75 mL). The combined EtOAc extracts were dried (Na₂SO₄), filtered, and concentrated to yield the crude product. Preparative reverse phase HPLC purification on a Waters Autopurification system using a Sunfire Prep C18 OBD column [5 μm, 19 x 50 mm; flow rate, 20 mL/min; Solvent A: H₂O with 0.1% HCO₂H; Solvent B: CH₃CN with 0.1% HCO₂H; injection volume: 4x 3.6-4.2 mL (CH₃CN); gradient: 88→100% B over 12 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 6.6-10.6 min, were collected and lyophilized to give 552 mg of pure **2-5** (41%): ¹H NMR (400 MHz, CDCl₃) δ 16.22 (s, 1 H), 7.49-7.47 (m, 4 H), 7.37-7.31 (m, 6 H), 6.80 (d, *J* = 11.0 Hz, 1 H), 5.76-5.64 (m, 2 H), 5.35 (s, 2 H), 5.17-5.11 (m, 4 H), 4.98 (d, *J* = 9.2, 1 H), 4.87 (d, *J* = 9.8 Hz, 1 H), 3.96 (m, *J* = 10.4 Hz, 1 H), 3.83-3.71 (m, 4 H), 3.14 (dd, *J* = 14.7,


15

20

25

4.3 Hz, 1 H), 3.0-2.87 (m, 1 H), 2.55-2.35 (m, 9 H), 2.11 (d, $J = 14.7$ Hz, 1 H), 0.82 (s, 9 H), 0.26 (s, 3 H), 0.13 (s, 3 H); MS (ESI) m/z 820.55 (M+H).

Compound 2-6



5


A solution of **2-5** (550 mg, 0.671 mmol, 1.0 equiv) in degassed CH_2Cl_2 (2.5 mL, with 1 and 1.5 mL rinse) was added under nitrogen via syringe to a flame-dried flask containing *N,N*-dimethylbarbituric acid (324 mg, 2.07 mmol, 3.0 equiv), and Tetrakis(triphenylphosphine)palladium(0) (56.9 mg, 0.0492 mmol, 0.07 equiv). The resulting solution was heated to 35 °C for 4 h, then concentrated to remove the solvent. The resulting crude mixture was purified via preparative reverse phase HPLC on a Waters Autopurification system using a Sunfire Prep C18 OBD column [5 μm , 19 $\times$ 50 mm; flow rate, 20 mL/min; Solvent A: H_2O with 0.1% HCO_2H ; Solvent B: CH_3CN with 0.1% HCO_2H ; injection volume: 3 $\times$ 3.1 mL (CH_3CN); gradient: 80 $\rightarrow$ 100% B over 17 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 6.1-10.1 min, were collected and freeze-dried to give 352 mg of pure **2-6** (71%): ^1H NMR (400 MHz, CDCl_3) δ 16.10 (s, 1 H), 7.51-7.43 (m, 4 H), 7.9-7.29 (m, 6 H), 6.61 (d, $J = 9.8$ Hz, 1 H), 5.35 (s, 2 H), 4.87 (dd, $J = 22.6, 10.4$ Hz, 2 H), 3.96 (d, $J = 10.4$ Hz, 1 H), 3.91 (s, 2 H), 3.12 (dd, $J = 15.3, 10.1$ Hz, 1 H), 3.04-2.92 (m, 1 H), 2.55-2.31 (m, 9 H), 2.11 (d, $J = 14.7$ Hz, 1 H), 0.82 (s, 9 H), 0.27 (s, 3 H), 0.12 (s, 3 H); MS (ESI) m/z 740.44 (M+H).

10

15

20

Compound 59



To a solution of aniline **2-6** (30 mg, 0.041 mmol, 1 equiv) in THF (600 μL) was added bromoacetyl bromide (3.7 μL , 0.043 mmol, 1.05 equiv). After 15 minutes, indazole (53 mg, 0.45 mmol, 10 equiv) was added. After 15 h the reaction was


25

heated to 80 °C. After an additional 26 h, another 20 mg indazole (0.17 mmol, 4 equiv) was added and the reaction heated at 80 °C. After 20 h, the solvent was removed *in vacuo* and the resulting crude mixture dried under vacuum.

The above crude intermediate was transferred to a plastic vial in dioxane (1.2 mL) and an aqueous solution of hydrogen fluoride (50%, 300 µL) was added. After five hours, the reaction solution was diluted with an aqueous solution of K₂HPO₄ (3.6 g in 30 mL) and extracted with EtOAc (2 x 30 mL). The combined organic layers were dried (Na₂SO₄), filtered, and concentrated to yield the crude product.

Palladium on carbon (10%, 10 mg) was added to solution of the above crude intermediate in dioxane:methanol (1:1, 1 mL). The flask was fitted with a septum and evacuated and back-filled three times with hydrogen gas. Hydrogen gas was bubbled through the reaction solution for two minutes, then the reaction was stirred under an atmosphere (balloon) of hydrogen gas for 1.5 h. The reaction mixture was filtered through celite to remove the palladium catalyst and concentrated under reduced pressure. Preparative reverse phase HPLC purification of the resulting oil was performed on a Waters Autopurification system using a Polymerx 10 µ RP-γ 100 R column [30 x 21.20 mm, 10 micron, solvent A: 0.05N HCl in water, solvent B: CH₃CN; injection volume: 4.8 mL (0.05N HCl in water); gradient elution with 10→60% B over 15 min, then held at 100% for 5 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 14-14.65 min, were collected and freeze-dried to yield 3.6 mg of compound **59** (15%): ¹H NMR (400 MHz, CD₃OD) δ 8.71 (s, 1 H), 8.19 (d, *J* = 11.0 Hz, 1 H), 7.92-7.90 (m, 1 H), 7.72-7.57 (m, 2 H), 7.35-7.29 (m, 1 H), 5.65 (s, 2 H), 4.08 (s, 1 H), 3.16-2.92 (m, 9 H), 2.31-2.18 (m, 2 H), 1.67-1.60 (m, 1 H); MS (ESI) *m/z* 606.41 (M+H).

Compound **60**




To a solution of aniline **2-6** (22 mg, 0.030 mmol, 1 equiv) in THF (500 µL) was added bromoacetyl bromide (2.7 µL, 0.031 mmol, 1.05 equiv). After 30 minutes, pyrazole (36 mg, 0.53 mmol, 18 equiv) was added. After 20 min the reaction was

heated to 80 °C for 1.5 h, cooled to room temperature for 15 h, then heated at 80 °C for 4.5 h. The solvent was removed *in vacuo* and the resulting crude mixture dried under vacuum.

The above crude intermediate was transferred to a plastic vial in acetonitrile (1.0 mL) and an aqueous solution of hydrogen fluoride (50%, 200 µL) was added. After 20 h, the reaction solution was diluted with an aqueous solution of K₂HPO₄ (2.4 g in 20 mL) and extracted with EtOAc (2 x 25 mL). The combined organic layers were dried (Na₂SO₄), filtered, and concentrated to yield the crude product.

Palladium on carbon (10%, 10 mg) was added to solution of the above crude intermediate in dioxane:methanol (1:1, 1 mL). The flask was fitted with a septum and evacuated and back-filled three times with hydrogen gas. Hydrogen gas was bubbled through the reaction solution for two minutes and the reaction was stirred under an atmosphere (balloon) of hydrogen gas for 1.5 h. The reaction mixture was filtered through celite to remove the palladium catalyst and concentrated under reduced pressure. Preparative reverse phase HPLC purification of the resulting oil was performed on a Waters Autopurification system using a Polymerx 10 µ RP-γ 100 R column [30 x 21.20 mm, 10 micron, solvent A: 0.05N HCl in water, solvent B: CH₃CN; injection volume: 3.0 mL (10% CH₃CN in 0.05N HCl in water); gradient elution with 10→60% B over 10 min, then held at 100% for 5 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 8.8-10.2 min were collected and freeze-dried. The resulting yellow solid was purified a second time via preparative reverse phase HPLC purification using the above procedure with a gradient over 20 minutes, with the fractions with the desired MW eluting at 10.7-12.4 min were collected and freeze-dried to give 8.2 mg of pure **60** (50%): ¹H NMR (400 MHz, CD₃OD) δ 8.19 (d, *J* = 11.0 Hz, 1 H), 8.05-7.92 (m, 2 H), 6.62-6.57 (m, 1 H), 5.33 (d, *J* = 4.9 Hz, 2 H), 4.08 (s, 1 H), 3.16-2.90 (m, 9 H), 2.31-2.17 (m, 2 H), 1.69-1.55 (m, 1 H); MS (ESI) *m/z* 556.42 (M+H).

Compound 61




To a solution of aniline **2-6** (23 mg, 0.032 mmol, 1 equiv) in THF (500 μ L) was added bromoacetyl bromide (2.9 μ L, 0.034 mmol, 1.05 equiv). After 30 minutes, imidazole (32 mg, 0.47 mmol, 15 equiv) was added and the solution was heated to 80 °C. After 2 h, the solution was cooled and the solvent was removed *in vacuo*.

5 The above crude intermediate was transferred to a plastic vial in dioxane (1.2 mL) and an aqueous solution of hydrogen fluoride (50%, 200 μ L) was added. After 1.5 h, the reaction solution was diluted with an aqueous solution of K_2HPO_4 (2.4 g in 30 mL) and extracted with EtOAc (2 x 25 mL). The combined organic layers were dried (Na_2SO_4), filtered, and concentrated to yield the crude product.

10 Palladium on carbon (10%, 8 mg) was added to solution of the above crude intermediate in dioxane:methanol (1:1, 1 mL). The flask was fitted with a septum and evacuated and back-filled three times with hydrogen gas. Hydrogen gas was bubbled through the reaction solution and the reaction was stirred under an atmosphere (balloon) of hydrogen gas for 1.5 h. More palladium catalyst was added
15 and the evacuation and backfilling with hydrogen was performed twice more at 1.5 h and 5 h. The reaction mixture was filtered through celite to remove the palladium catalyst and concentrated under reduced pressure. Preparative reverse phase HPLC purification of the resulting oil was performed on a Waters Autopurification system using a Polymerx 10 μ RP- γ 100 R column [30 x 21.20 mm, 10 micron, solvent A:
20 0.05N HCl in water, solvent B: CH_3CN ; injection volume: 2.8 mL (0.05N HCl in water); gradient elution with 10 $\rightarrow$ 60% B over 15 min, then held at 100% for 5 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 7.0-7.8 min were collected and freeze-dried to give 4.1 mg of pure **61** (23%): 1H NMR (400 MHz, CD_3OD) δ 9.02 (s, 1 H), 8.17 (d, J = 11.0 Hz, 1 H), 7.67 (s, 1 H), 7.61 (s, 1
25 H), 5.34 (s, 2 H), 4.09 (s, 1 H), 3.18-2.90 (m, 9 H), 2.34-2.1.7 (m, 2 H), 1.71-1.56 (m, 1 H); MS (ESI) m/z 556.45 (M+H).

Compound **62**




To a solution of aniline **2-6** (20.2 mg, 0.027 mmol, 1 equiv) in THF (500 μ L) was added bromoacetyl bromide (2.5 μ L, 0.029 mmol, 1.05 equiv). After 30 minutes, 1H-1,2,3-triazole (31 μ L, 0.54 mmol, 20 equiv) was added and the solution was heated to 80 °C. After 17 h, an additional 31 μ L (20 equiv) of 1H-1,2,3-triazole was added and the solution was heated for 22 h. The solution was cooled and the solvent was removed *in vacuo*.

The above crude intermediate was transferred to a plastic vial in dioxane (1.0 mL) and an aqueous solution of hydrogen fluoride (50%, 200 μ L) was added. After 17 h, the reaction solution was diluted with an aqueous solution of K₂HPO₄ (2.4 g in 20 mL) and extracted with EtOAc (2 x 25 mL). The combined organic layers were dried (Na₂SO₄), filtered, and concentrated to yield the crude product.

Palladium on carbon (10%, 7 mg) was added to solution of the above crude intermediate in dioxane:methanol (1:1, 1 mL). The flask was fitted with a septum and evacuated and back-filled three times with hydrogen gas. Hydrogen gas was bubbled through the reaction solution and the reaction was stirred under an atmosphere (balloon) of hydrogen gas for 1.5 h. The reaction mixture was filtered through celite to remove the palladium catalyst and concentrated under reduced pressure. Preparative reverse phase HPLC purification of the resulting oil was performed on a Waters Autopurification system using a Polymerx 10 μ RP- γ 100 R column [30 x 21.20 mm, 10 micron, solvent A: 0.05N HCl in water, solvent B: CH₃CN; injection volume: 2.5 mL (0.05N HCl in water); gradient elution with 10→60% B over 15 min, then held at 100% for 5 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 9.25-10.5 min were collected and freeze-dried. Purification was performed a second time as above and the fractions with the desired MW, eluting at 9.75-10.25 min were collected and freeze-dried to give 1.5 mg of pure **62** (10%): ¹H NMR (400 MHz, CD₃OD) δ 8.24 (s, 1 H), 8.17 (d, *J* = 11.0 Hz, 1 H), 8.00 (s, 1 H), 5.57 (s, 2 H), 4.09 (s, 1 H), 3.16-2.92 (m, 9 H), 2.34-2.16 (m, 2 H), 1.71-1.67 (m, 1 H); MS (ESI) *m/z* 557.44 (M+H).

Compound **63**




To a solution of aniline **2-6** (16.7 mg, 0.023 mmol, 1 equiv) in THF (500 μ L) was added bromoacetyl bromide (2.0 μ L, 0.024 mmol, 1.05 equiv). After 20 minutes, a tetrazole solution (0.45M in CH₃CN, 500 μ L, 0.23 mmol, 10 equiv) was added and the solution was heated to 80 °C. After 4 h, potassium carbonate (35 mg, 0.25 mmol, 11 equiv) was added and the reaction heated for 35 minutes. The solution was cooled and filtered through celite, and the solvent was removed *in vacuo*.

The above crude intermediate was transferred to a plastic vial in dioxane (1.0 mL) and an aqueous solution of hydrogen fluoride (50%, 200 μ L) was added. After 18 h, the reaction solution was diluted with an aqueous solution of K₂HPO₄ (2.4 g in 20 mL) and extracted with EtOAc (2 x 25 mL). The combined organic layers were dried (Na₂SO₄), filtered, and concentrated to yield the crude product.

Palladium on carbon (10%, 7 mg) was added to solution of the above crude intermediate in dioxane:methanol (1:1, 1 mL). The flask was fitted with a septum and evacuated and back-filled three times with hydrogen gas. Hydrogen gas was bubbled through the reaction solution and the reaction was stirred under an atmosphere (balloon) of hydrogen gas for 1 h. The reaction mixture was filtered through celite to remove the palladium catalyst and concentrated under reduced pressure. Preparative reverse phase HPLC purification of the resulting oil was performed on a Waters Autopurification system using a Polymerx 10 μ RP- γ 100 R column [30 x 21.20 mm, 10 micron, solvent A: 0.05N HCl in water, solvent B: CH₃CN; injection volume: 2.5 mL (10% CH₃CN in 0.05N HCl in water); gradient elution with 10 $\rightarrow$ 60% B over 15 min, then held at 100% for 5 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 11.2-12.1 min were collected and freeze-dried. Purification was performed a second time as above with the gradient extended over 20 min. Fractions with the desired MW, eluting at 13.7-14.5 min were collected and freeze-dried to give 1.6 mg of pure **63** (13%); ¹H NMR (400 MHz, CD₃OD) δ 8.78 (s, 1 H), 8.14 (d, *J* = 11.0 Hz, 1 H), 5.78 (s, 2 H), 4.07 (s, 1 H), 3.17-2.81 (m, 9 H), 2.36-2.16 (m, 2 H), 1.70-1.52 (m, 1 H); MS (ESI) *m/z* 558.43 (M+H).


Example 3. Synthesis of Compounds of Structural Formula (A), wherein X is hydrogen and Y is -NH-C(O)-heterocyclyl, -NH-C(O)-heteroaryl, -NH-C(O)-[C(R^D)(R^E)]₀₋₁-N(R^A)(R^B), -NH-C(O)-carbocyclyl, -NH-C(O)-aryl, -NH-SO₂-(C₁-C₆)alkyl, -NH-SO₂-carbocyclyl, -NH-SO₂-aryl, -NH-SO₂-heterocyclyl or -NH-SO₂-heteroaryl.

Scheme 3



In Scheme 3, R' represents heterocyclyl, heteroaryl, carbocyclyl, aryl, (C₁-C₆)alkyl, or -[(C(R^D)(R^E)]₀₋₁-N(R^A)(R^B); and R^{2'} represents hydrogen, (C₁-C₆)alkyl, -(C₀-C₅)alkylene-carbocyclyl, -(C₀-C₅)alkylene-aryl, -(C₀-C₅)alkylene-heteroaryl, or -(C₀-C₅)alkylene-heterocyclyl. For certain compounds made by Scheme 3 and described below, R^Z is hydrogen and R^X and R^Y are taken together with the carbon and nitrogen atoms to which they are bound to form an optionally substituted 4-7 membered saturated heterocyclyl. It will be readily apparent to those of skill in the art, however, that this Scheme 3 will also be useful to synthesize compounds where R^X, R^Y and R^Z are R^B, R^D and R^E, respectively, as defined in structural formula (A). The following compounds were prepared according to Scheme 3.


Compound **64**



To a solution of aniline **9** (17.0 mg, 0.038 mmol, 1 equiv) in DMF (200 μ L) was added N-Benzyloxycarbonyl-L-proline acid chloride (1.0 M in toluene, 57 μ L, 1.5 equiv). After 50 min the reaction mixture was diluted to 3 mL with 0.05 N HCl in H₂O and filtered to remove any solids. Preparative reverse phase HPLC purification of the resulting solution was performed on a Waters Autopurification system using a Polymerx 10 μ RP- γ 100 R column [30 $\times$ 21.20 mm, 10 micron, solvent A: 0.05N HCl in water, solvent B: CH₃CN; injection volume: 3.5 mL (0.05N HCl in water); gradient elution with 10 $\rightarrow$ 20% B over 25 min, then held at 100% for 5 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 27.1-28.4 min, were collected and freeze-dried.

Palladium on carbon (10%, 10 mg) was added to a solution of the above intermediate in dioxane:MeOH (1:3, 2.3 mL). The flask was fitted with a septum and evacuated and back-filled three times with hydrogen gas. The reaction solution was stirred under an atmosphere (balloon) of hydrogen gas for 1.7 h, then was filtered through celite to remove the palladium catalyst and concentrated under reduced pressure. Half of the resulting residue was purified via preparative reverse phase HPLC purification on a Waters Autopurification system using a Polymerx 10 μ RP- γ 100 R column [30 $\times$ 21.20 mm, 10 micron, solvent A: 0.05N HCl in water, solvent B: CH₃CN; injection volume: 1.8 mL (0.05 N HCl in water); gradient elution with 0 $\rightarrow$ 35% B over 10 min, then held at 100% for 5 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 7.8-8.5 min, were collected and freeze-dried to yield 1.9 mg of compound **64** (30%): ¹H NMR (400 MHz, CD₃OD) δ 8.16 (d, *J* = 11.0 Hz, 1 H), 4.59-4.56 (m, 1 H), 4.10 (s, 1 H), 3.48-3.33 (m, 2 H), 3.18-2.95 (m, 9 H), 2.59-2.50 (m, 1 H), 2.34-2.05 (m, 5 H), 1.70-1.60 (m, 1 H); MS (ESI) *m/z* 545.38 (M+H).


Compound **65**



To a solution of aniline **9** (15.7 mg, 0.035 mmol, 1 equiv) in DMF (200 μ L) was added N-Benzyloxycarbonyl-D-proline acid chloride (1.0 M in toluene, 53 μ L, 1.5 equiv). After 50 min, the reaction was complete. The reaction mixture was
5 diluted to 3 mL with 0.05 N HCl in H₂O and filtered to remove any solids. Preparative reverse phase HPLC purification of the resulting solution was performed on a Waters Autopurification system using a Polymerx 10 μ RP- γ 100 R column [30 $\times$ 21.20 mm, 10 micron, solvent A: 0.05 N HCl in water, solvent B: CH₃CN; injection volume: 3.5 mL (0.05 N HCl in water); gradient elution with 15 $\rightarrow$ 80% B
10 over 10 min, then held at 100% for 5 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 6.95-8.10 min, were collected and freeze-dried.


Palladium on carbon (10%, 15 mg) was added to a solution of the above intermediate in dioxane:MeOH (1:3, 2.3 mL). The flask was fitted with a septum
15 and evacuated and back-filled three times with hydrogen gas, and the reaction was stirred under an atmosphere (balloon) of hydrogen gas for 1.5 h. The reaction mixture was filtered through celite to remove the palladium catalyst and concentrated under reduced pressure. Half of the resulting residue was purified via preparative reverse phase HPLC purification on a Waters Autopurification system
20 using a Polymerx 10 μ RP- γ 100 R column [30 $\times$ 21.20 mm, 10 micron, solvent A: 0.05 N HCl in water, solvent B: CH₃CN; injection volume: 1.8 mL (0.05 N HCl in water); gradient elution with 0 $\rightarrow$ 35% B over 10 min, then held at 100% for 5 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 8.35-8.85 min, were collected and freeze-dried to yield 0.93 mg of compound **65** (24%):
25 ¹H NMR (400 MHz, CD₃OD) δ 8.17 (d, J = 11.0 Hz, 1 H), 4.59-4.53 (m, 1 H), 4.09 (s, 1 H), 3.48-3.37 (m, 2 H), 3.18-2.90 (m, 9 H), 2.59-2.50 (m, 1 H), 2.34-2.05 (m, 5 H), 1.70-1.59 (m, 1 H); MS (ESI) m/z 545.37 (M+H).

Compound **66**



The second half of crude **64** (0.012 mmol, 1.0 equiv) was dissolved in DMF (500 μ L), and formaldehyde (37% aqueous solution, 5.3 μ L, 0.072 mmol, 6 equiv), triethylamine (5.0 μ L, 0.036 mmol, 3 equiv), and sodium triacetoxyborohydride (8.4 mg, 0.039 mmol, 3.2 equiv) were added sequentially. After 2 h, the reaction mixture was diluted to 1.8 mL with 0.05N HCl in H₂O and purified via preparative reverse phase HPLC on a Waters Autopurification system using a Polymerx 10 μ RP- γ 100 R column [30 $\times$ 21.20 mm, 10 micron, solvent A: 0.05 N HCl in water, solvent B: CH₃CN; injection volume: 1.8 mL (0.05 N HCl in water); gradient: 0 $\rightarrow$ 30% B over 10 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 8.6-9.35 min, were collected and freeze-dried to provide a mixture of the desired compound and an over-formylated product. The resulting compound mixture was dissolved in 4 N aqueous HCl solution (1.5 mL) and stirred for 50 h, then freeze-dried to provide 1.0 mg of the desired compound **66** (15%): ¹H NMR (400 MHz, CD₃OD) δ 8.17 (d, J = 10.4 Hz, 1 H), 4.36 (t, J = 8.6 Hz, 1 H), 4.08 (s, 1 H), 3.82-3.73 (m, 1 H), 3.20-2.90 (m, 12 H), 2.73-2.68 (m, 1 H), 2.35-2.10 (m, 5 H), 1.70-1.60 (m, 1 H); MS (ESI) m/z 559.38 (M+H).

Compound 67




The second half of crude **65** (0.007 mmol, 1.0 equiv) was dissolved in DMF (500 μ L), and formaldehyde (37% aqueous solution, 3.1 μ L, 0.042 mmol, 6 equiv) and TEA (3.0 μ L, 0.021 mmol, 3 equiv), and sodium triacetoxyborohydride (4 mg, 0.026 mmol, 2.6 equiv) were added sequentially. After 2.2 h, the reaction mixture was diluted to 1.8 mL with 0.05N HCl in H₂O and purified via preparative reverse phase HPLC on a Waters Autopurification system using a using a Polymerx 10 μ RP- γ 100 R column [30 $\times$ 21.20 mm, 10 micron, solvent A: 0.05 N HCl in water,

solvent B: CH₃CN; injection volume: 2.0 mL (0.05 N HCl in water); gradient: 0→30% B over 10 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 8.9-9.6 min, were collected and freeze-dried to provide a mixture of the desired compound and an over-formylated product. The resulting compound mixture was dissolved in 6 N aqueous HCl solution and stirred for 50 h, then freeze-dried to provide 1.5 mg of the desired compound **67** (38%) : ¹H NMR (400 MHz, CD₃OD) δ 8.17 (d, *J* = 10.4 Hz, 1 H), 4.45-4.34 (m, 1 H), 4.08 (s, 1 H), 3.84-3.74 (m, 1 H), 3.20-2.90 (m, 12 H), 2.79-2.65 (m, 1 H), 2.33-2.05 (m, 5 H), 1.70-1.58 (m, 1 H); MS (ESI) *m/z* 559.40 (M+H).

10

Compound **68**




To a solution of (*S*)-(-)-1-Cbz-piperidinecarboxylic acid (34.2 mg, 0.13 mmol, 3 equiv), and (2-(7-aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluoro-phosphate) (50.0 mg, 0.13 mol, 3 equiv) in DMF (200 μL) was added triethylamine (18 μL, 0.13 mmol, 3 equiv). After 30 min, aniline **9** (17.5 mg, 0.039 mmol, 1 equiv) was added. After 16 h, the reaction mixture was diluted to 3 mL with 0.05 N HCl in H₂O and purified via preparative reverse phase HPLC on a Waters Autopurification system using a Polymerx 10 μ RP-γ 100 R column [30 × 21.20 mm, 10 micron, solvent A: 0.05 N HCl in water, solvent B: CH₃CN; injection volume: 3.5 mL (0.05 N HCl in water); gradient elution with 15→70% B over 10 min, then held at 100% for 5 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 9.07-10.0 min, were collected and freeze-dried. Palladium on carbon (10%, 4 mg) was added to a solution of this foam in dioxane:MeOH (1:3, 1.2 mL). The flask was fitted with a septum and evacuated and back-filled three times with hydrogen gas. The reaction mixture was stirred under an atmosphere (balloon) of hydrogen gas for 1.5 h, then was filtered through celite to remove the palladium catalyst and concentrated under reduced pressure. Preparative reverse phase HPLC purification of the resulting oil was performed on a Waters Autopurification system using a Polymerx 10 μ RP-γ 100 R column

15
20
25
30

[30 × 21.20 mm, 10 micron, solvent A: 0.05 N HCl in water, solvent B: CH₃CN; injection volume: 2.0 mL (0.05 N HCl in water); gradient elution with 0→35% B over 10 min, then held at 100% for 5 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 8.15-8.58 min, were collected and freeze-dried to yield 0.75 mg of compound **68** (4%): ¹H NMR (400 MHz, CD₃OD) δ 8.15 (d, *J* = 11.0 Hz, 1 H), 4.12-4.06 (m, 2 H), 3.48-3.40 (m, 2 H), 3.20-2.90 (m, 9 H), 2.36-2.18 (m, 3 H), 2.02-1.90 (m, 2 H), 1.82-1.60 (m, 4 H); MS (ESI) *m/z* 559.37 (M+H).


10 Compound **69**



To a solution of (*R*)-(+)-1-Cbz-piperidinecarboxylic acid (35.0 mg, 0.13 mmol, 3 equiv), and (2-(7-aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluoro-phosphate) (50.0 mg, 0.13 mol, 3 equiv) in DMF (200 μL) was added TEA (18 μL, 0.13 mmol, 3 equiv). After 30 min, aniline **9** (16.6 mg, 0.037 mmol, 1 equiv) was added. After 16 h, the reaction mixture was diluted to 3 mL with 0.05 N HCl in H₂O and purified via preparative reverse phase HPLC on a Waters Autopurification system using a Polymerx 10 μ RP-γ 100 R column [30 × 21.20 mm, 10 micron, solvent A: 0.05 N HCl in water, solvent B: CH₃CN; injection volume: 3.5 mL (0.05 N HCl in water); gradient elution with 10→50% B over 10 min, then held at 100% for 5 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 12.1-12.9 min, were collected and freeze-dried. Palladium on carbon (10%, 5 mg) was added to a solution of this foam in dioxane:MeOH (1:3, 800 μL). The flask was fitted with a septum and evacuated and back-filled three times with hydrogen gas. The reaction mixture was stirred under an atmosphere (balloon) of hydrogen gas for 1.75 h, then was filtered through celite to remove the palladium catalyst and concentrated under reduced pressure. Preparative reverse phase HPLC purification of the resulting oil was performed on a Waters Autopurification system using a Polymerx 10 μ RP-γ 100 R column [30 × 21.20 mm, 10 micron, solvent A: 0.05 N HCl in water, solvent B: CH₃CN;

injection volume: 2.0 mL (0.05 N HCl in water); gradient elution with 0→35% B over 10 min, then held at 100% for 5 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 8.75-9.16 min, were collected and freeze-dried to yield 0.55 mg of compound **69** (3%): ¹H NMR (400 MHz, CD₃OD) δ 8.16 (d, *J* = 11.0 Hz, 1 H), 4.13-4.06 (m, 2 H), 3.50-3.43 (m, 2 H), 3.20-2.90 (m, 9 H), 2.38-2.18 (m, 3 H), 2.04-1.88 (m, 2 H), 1.83-1.60 (m, 4 H); MS (ESI) *m/z* 559.38 (M+H).

Compound **70**




10

To a solution of compound **68** (0.0138 mmol, 1 equiv) in DMF (750 μL), were added formaldehyde (37% aqueous solution, 6.2 μL, 0.083 mmol, 6 equiv), TEA (5.8 μL, 0.041 mmol, 3 equiv), and sodium triacetoxyborohydride (11 mg, 0.051 mmol, 3.7 equiv) sequentially. After 17 h, the reaction mixture was concentrated to remove amine and 6 N aqueous HCl (500 μL) was added. After 19 days, the reaction solution was purified via preparative reverse phase HPLC on a Waters Autopurification system using a Polymerx 10 μ RP-γ 100 R column [30 × 21.20 mm, 10 micron, solvent A: 0.05 N HCl in water, solvent B: CH₃CN; injection volume: 2.5 mL (0.05 N HCl in water); gradient: 15→50% B over 15 min, mass-directed fraction collection]. Fractions with the desired MW, eluting at 5.75-6.2 min, were collected and freeze-dried to provide 2.4 mg of the desired compound **70** (31%): ¹H NMR (400 MHz, CD₃OD) δ 8.16 (d, *J* = 11.0 Hz, 1 H), 4.08-4.04 (m, 1 H), 3.59-3.53 (m, 1 H), 3.20-3.10 (m, 5 H), 3.06-2.96 (m, 5 H), 2.90m (s, 3 H), 2.36-2.25 (m, 2 H), 2.11-2.05 (m, 1 H), 2.02-1.94 (m, 2 H), 1.90-1.74 (m, 2 H), 1.71-1.58 (m, 2 H); MS (ESI) *m/z* 573.33 (M+H).

20

25


Compound **71**



Compound **9** (20 mg, 0.045 mmol, 1.0 equiv) in THF was added Na_2CO_3 (9.5 mg, 0.089 mmol, 2.0 equiv), (4R)-4-fluoro-1-methyl-L-proline (9.8 mg, 0.067 mmol, 1.5 equiv) and HATU (34.6 mg, 0.047 mmol, 2.0 equiv). The reaction mixture was stirred at room temperature for 20 hour. LC-MS analysis indicated the starting material was consumed completely. HCl/MeOH (1 mL, 4 N) was added to the mixture at 0 °C and stirred for 2 min. The mixture was concentrated under vacuum, the residue was purified by reverse phase HPLC to afford product **71** (6.1 mg): ^1H NMR (400 MHz, CD_3OD) δ 8.18 (d, J = 10.8 Hz, 1 H), 5.51 (d, J = 51.6 Hz, 1 H), 4.76-4.72 (m, 1 H), 4.22-4.16 (m, 1 H), 4.10 (s, 1 H), 3.74-3.63 (m, 1 H), 3.21-2.97 (m, 14 H), 2.35-2.21 (m, 2 H), 1.69-1.60 (m, 1 H); MS (ESI) m/z 577.1 (M+H).


Compounds **72** and **73** were prepared similarly to compound **71** using the corresponding amino acids.

Compound **72**




Prepared similarly to compound **71**: ^1H NMR (400 MHz, CD_3OD) δ 8.16 (d, J = 10.8 Hz, 1 H), 5.48 (d, J = 51.2 Hz, 1 H), 4.60-4.56 (m, 1 H), 4.11 (s, 1 H), 4.05-3.98 (m, 1 H), 3.67-3.54 (m, 1 H), 3.24-2.96 (m, 13 H), 2.55-2.44 (m, 1 H), 2.34-2.22 (m, 2 H), 1.70-1.66 (m, 1 H); MS (ESI) m/z 577.1 (M+H).

Compound **73**



Prepared similarly to compound 71: ^1H NMR (400 MHz, CD_3OD) δ 8.18 (d, $J = 10.8$ Hz, 1 H), 4.76-4.71 (m, 1 H), 4.17-4.12 (m, 1 H), 4.09 (s, 1 H), 3.96-3.86 (m, 1 H), 3.67-3.53 (m, 1 H), 3.55-3.53 (m, 1 H), 3.25-2.73 (m, 12 H), 2.33-2.19 (m, 2 H), 1.68-1.59 (m, 1 H); MS (ESI) m/z 595.3 (M+H)


Compound 3-1-1



1-(Bocamino)cyclopropanecarboxylic acid (67.4 mg, 0.335 mmol), O-(7-azabenzotriazol-1-yl)- N,N,N',N' -tetramethyluronium hexafluorophosphate (127 mg, 0.335 mmol), and triethylamine (0.078 mL, 0.56 mmol) were stirred in DMF (1 mL) for 30 minutes. Compound 9 (50 mg, 0.112 mmol) was added. After stirring overnight, the reaction mixture was purified directly by preparative reverse phase HPLC purification on a Waters Autopurification system using a Polymerx 10 μ RP- γ 100 R column [30 $\times$ 21.20 mm, 10 micron, solvent A: 0.05N HCl, solvent B: CH_3CN , gradient elution with 0 $\rightarrow$ 50% B over 10 min; mass-directed fraction collection]. Fractions with the desired MW were collected and freeze-dried. The material was repurified by preparative reverse phase HPLC purification on a Waters Autopurification system using a Polymerx 10 μ RP- γ 100 R column [30 $\times$ 21.20 mm, 10 micron, solvent A: 0.05N HCl, solvent B: CH_3CN , gradient elution with 0 $\rightarrow$ 100% B over 15 min; mass-directed fraction collection]. Fractions with the desired MW were collected and freeze-dried. This gave 42 mg of compound 3-1-1 (59%, ~80% pure) which was used without further purification.: MS (ESI) m/z 631.41 (M+H).


25

Compound 74



Compound **3-1-1** (42 mg, 0.067 mmol, ~80% pure) was stirred in 4M HCl in 1,4-dioxane (5 mL) overnight. The reaction mixture was concentrated under reduced pressure and was purified by preparative reverse phase HPLC purification on a Waters Autopurification system using a Polymerx 10 μ RP- γ 100 R column [30 $\times$ 21.20 mm, 10 micron, solvent A: 0.05N HCl, solvent B: CH₃CN, gradient elution with 0 $\rightarrow$ 50% B over 10 min; mass-directed fraction collection]. Fractions with the desired MW were collected and freeze-dried. The material was dissolved in MeOH (1 mL), and the solution was added dropwise to vigorously stirring diethyl ether (200 mL). The resulting solid was collected by filtration on a pad of Celite. This was washed with diethyl ether (3 times), and the solid was dissolved in MeOH and concentrated under reduced pressure. The material was freeze-dried, yielding 25.8 mg of compound **74**: ¹H NMR (400 MHz, CD₃OD with 1 drop DCl) δ 8.00 (d, *J* = 7.0 Hz, 1 H), 4.05 (s, 1 H), 3.20-2.85 (m, 9 H), 2.36-2.06 (m, 2 H), 1.70-1.52 (m, 3 H), 1.35-1.22 (m, 2 H); MS (ESI) *m/z* 531.33 (M+H).


Compound **75**



To a dichloromethane (5 mL) suspension of compound **9** (0.260 g, 0.50 mmol, 1.0 equiv) at rt was added triethylamine (0.139 mL, 1.00 mmol, 2.0 equiv). The reaction was stirred at rt until form a clear solution. Methylisocyanate (89.4 μ L, 1.50 mmol, 3.0 equiv) was added to the reaction mixture dropwise. The reaction was allowed to stir at 25 $^{\circ}$ C for 1 h. Additional methylisocyanate (45 μ L, 0.75 mmol, 1.5 equiv) was added and stirred overnight. LCMS indicate there are still starting material present. The solvent was removed under vacuum to give the crude **75**. The crude product was purified by HPLC on a Polymerx 10 μ RP- γ 100 R column

[30 × 21.20 mm, 10 micron, solvent A: 0.05 N HCl, solvent B: CH₃CN, sample in 2.0 mL (0.05 N HCl), gradient elution with 15→65 % B over 15 min, mass-directed fraction collection] to yield the desired product **75** as a yellow solid (80 mg, 31.7 %): ¹H NMR (400 MHz, CD₃OD) δ 8.12 (d, *J* = 11.4 Hz, 1 H), 4.07 (s, 1 H), 3.04 (s, 3 H), 2.96 (s, 3 H), 3.13 -2.93 (m, 3 H), 2.77 (s, 3 H), 2.27-2.15 (m, 2 H), 1.69-1.57 (m, 1 H); MS (ESI) *m/z* 505.41 (M+H).

Compound **76**




10

Compound **9** (20 mg, 0.045 mmol, 1.0 equiv) in THF was added Na₂CO₃ (9.5 mg, 0.089 mmol, 2.0 equiv) and 0.1 mL benzoyl chloride solution (54 μL in 1 mL THF, 0.047 mmol, 1.05 equiv). The reaction mixture was stirred at room temperature for 1 hour. LC-MS analysis indicated the starting material was consumed completely. HCl/MeOH (1 mL, 4 N) was added to the mixture at 0 °C and stirred for 2 min. The mixture was concentrated under vacuum, the residue was purified by reverse phase HPLC to afford product **76** (5.5 mg): ¹H NMR (400 MHz, CD₃OD) δ 8.23 (d, *J* = 10.8 Hz, 1 H), 7.97 (d, *J* = 7.6 Hz, 2 H), 7.66-7.54 (m, 3 H), 4.11 (s, 1 H), 3.21-2.90 (m, 9 H), 2.37-2.24 (m, 2 H), 1.72-1.66 (m, 1 H); MS (ESI) *m/z* 552.1 (M+H).

20

Compounds **77** – **83** were prepared similarly to compound **76** using the corresponding acid chlorides.


25 Compound **77**



^1H NMR (400 MHz, CD_3OD) δ 8.25 (s, 1 H), 8.21 (d, $J = 8.0$ Hz, 1 H), 8.14 (d, $J = 10.4$ Hz, 1 H), 7.92 (d, $J = 8.0$ Hz, 1 H), 7.76 (t, $J = 8.0$ Hz, 1 H), 4.08 (s, 1 H), 3.21-2.89 (m, 9 H), 2.35-2.22 (m, 2 H), 1.71-1.61 (m, 1 H); MS (ESI) m/z 620.1 (M+H).

5


Compound 78



^1H NMR (400 MHz, CD_3OD) δ 8.10 (d, $J = 10.8$ Hz, 1 H), 7.41-7.33 (m, 3 H), 7.09-7.07 (m, 1 H), 4.00 (s, 1 H), 3.78 (s, 3 H), 3.12-2.86 (m, 9 H), 2.23-2.13 (m, 2 H), 1.60-1.50 (m, 1 H); MS (ESI) m/z 582.1 (M+H).

10


Compound 79



^1H NMR (400 MHz, CD_3OD) δ 8.12 (d, $J = 10.8$ Hz, 1 H), 7.89 (d, $J = 3.2$ Hz, 1 H), 7.78 (d, $J = 4.8$ Hz, 1 H), 7.22 (t, $J = 8.8$ Hz, 1 H), 4.10 (s, 1 H), 3.20-2.98 (m, 9 H), 2.36-2.20 (m, 2 H), 1.68-1.61 (m, 1 H); MS (ESI) m/z 558.1 (M+H).

15


Compound 80



^1H NMR (400 MHz, CD_3OD) δ 9.34 (s, 1 H), 9.04-9.00 (m, 2 H), 8.20-8.15 (m, 2 H), 4.07 (s, 1 H), 3.27-2.94 (m, 9 H), 2.34-2.18 (m, 2 H), 1.68-1.59 (m, 1 H); MS (ESI) m/z 553.1 (M+H).


20

Compound 81




¹H NMR (400 MHz, CD₃OD) δ 8.13-8.06 (m, 2 H), 7.98 (d, *J* = 7.6 Hz, 1 H), 7.77 (d, *J* = 7.2 Hz, 1 H), 7.67 (t, *J* = 8.0 Hz, 1 H), 4.01 (s, 1 H), 3.26 (s, 6 H), 3.14-2.83 (m, 9 H), 2.27-2.13 (m, 2 H), 1.64-1.52 (m, 1 H); MS (ESI) *m/z* 595.1 (M+H).

Compound 82




¹H NMR (400 MHz, CD₃OD) δ 8.08 (d, *J* = 10.8 Hz, 1 H), 7.98 (d, *J* = 8.4 Hz, 2 H), 7.49 (d, *J* = 8.4 Hz, 2 H), 4.02 (s, 1 H), 3.19 (s, 6 H), 3.12-2.88 (m, 9 H), 2.24-2.13 (m, 2 H), 1.60-1.51 (m, 1 H); MS (ESI) *m/z* 595.1 (M+H).

Compound 83



¹H NMR (400 MHz, CD₃OD) δ 8.19-8.14 (m, 2 H), 8.05 (d, *J* = 8.4 Hz, 1 H), 7.91-7.89 (m, 1 H), 7.76-7.74 (m, 1 H), 4.12 (s, 1 H), 3.32 (s, 6 H), 3.21-2.96 (m, 9 H), 2.41-1.98 (m, 2 H), 1.72-1.59 (m, 1 H); MS (ESI) *m/z* 595.0 (M+H).


Compound 84



Compound **9** (20 mg, 0.045 mmol, 1.0 equiv) in THF was added DIEA (11.5 mg, 0.089 mmol, 2.0 equiv) and 2-thiophenesulfonyl chloride (12.2 mg, 0.067 mmol, 1.5 equiv). The reaction mixture was stirred at room temperature for 20 hour. LC-MS analysis indicated the starting material was consumed completely.
5 HCl/MeOH (1 mL, 4 N) was added to the mixture at 0°C and stirred for 2 min. The mixture was concentrated under vacuum, the residue was purified by reverse phase HPLC to afford compound **84** (2.0 mg): ¹H NMR (400 MHz, CD₃OD) δ 7.75 (dd, *J* = 5.2, 1.2 Hz, 1 H), 7.59 (d, *J* = 2.8 Hz, 1 H), 7.52 (d, *J* = 10.4 Hz, 1 H), 7.09 (t, *J* = 4.4 Hz, 1 H), 4.07 (s, 1 H), 3.11-2.92 (m, 9 H), 2.30-2.18 (m, 2 H), 1.68-1.58 (m, 1
10 H); MS (ESI) *m/z* 593.9 (M+H).

Compounds **85** – **87** were prepared similarly to compound **84** using the corresponding sulfonyl chlorides.


15 Compound **85**



¹H NMR (400 MHz, CD₃OD) δ 7.44 (d, *J* = 10.0 Hz, 1 H), 4.10 (s, 1 H), 3.21-2.90 (m, 12 H), 2.34-2.22 (m, 2 H), 1.67-1.61 (m, 1 H); MS (ESI) *m/z* 526.1 (M+H).


20

Compound **86**



¹H NMR (400 MHz, CD₃OD) δ 7.82 (d, *J* = 7.6 Hz, 2 H), 7.58-7.46 (m, 4 H), 4.07 (s, 1 H), 3.10-2.92 (m, 9 H), 2.35-2.25 (m, 2 H), 1.65-1.55 (m, 1 H); MS
25 (ESI) *m/z* 552.1 (M+H).

Compound **87**




^1H NMR (400 MHz, CD_3OD) δ 7.72 (t, $J = 5.6$ Hz, 1 H), 7.62 (d, $J = 7.6$ Hz, 1 H), 7.50 (d, $J = 8.4$ Hz, 1 H), 7.41-7.38 (m, 2 H), 3.97 (s, 1 H), 3.03-2.82 (m, 9 H), 2.19-2.06 (m, 2 H), 1.53-1.50 (m, 1 H); MS (ESI) m/z 622.1 (M+H).

Example 4. Synthesis of Compounds of Structural Formula (I), wherein X is hydrogen and Y is -NH-C(O)-heterocycly or -NH-C(O)-heteroaryl

Scheme 4

10



In Scheme 4, R represents heteroaryl and R^2 is R^A as defined in Structural Formula (A). For certain compounds made by Scheme 4 and described below, R^Z is hydrogen and R^X and R^Y are taken together with the carbon and nitrogen atoms to which they are respectively bound to form an optionally substituted 4-7 membered


15

saturated heterocyclyl. It will be readily apparent to those of skill in the art, however, that this Scheme 4 will also be useful to synthesize compounds where R^X , R^Y and R^Z are R^B , R^D and R^E , respectively, as defined in structural formula (A).

The following compounds were prepared according to Scheme 4.

5

Compound **88**



To a suspension of 1-Fmoc-L-azetidine-2-carboxylic acid (135 mg, 0.42 mmol, 2.9 equiv), and (2-(7-aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluoro-phosphate) (164 mg, 0.43 mol, 3 equiv) in THF (1.5 mL) was added triethylamine (60 μL , 0.43 mmol, 3 equiv). After 30 min, aniline **2-6** (106 mg, 0.14 mmol, 1 equiv) was added. After 18 h, the reaction mixture was concentrated under reduced pressure. Preparative reverse phase HPLC of the resulting oil was performed on a Waters Autopurification system using a Sunfire Prep C18 OBD column [5 μm , 19 $\times$ 50 mm; flow rate, 20 mL/min; Solvent A: H_2O with 0.1% HCO_2H ; Solvent B: CH_3CN with 0.1% HCO_2H ; injection volume: 3 $\times$ 2.0 mL (CH_3CN); gradient: 80 $\rightarrow$ 100% B over 15 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 10.35-12.0 min, were collected and freeze-dried to provide 131 mg of a yellow powder.

20

To a solution of the above intermediate in CH_2Cl_2 (2 mL) was added piperidine (500 μL). After 30 min, the reaction solution was poured into aqueous pH 7 phosphate buffer and extracted with EtOAc (3 $\times$ 20 mL). The combined organic layers were dried (Na_2SO_4), filtered, and concentrated under reduced pressure. Purification of the resulting crude oil via flash column chromatography on silica gel (Silicycle, 5 g, 0 to 5 to 10 to 50 % EtOAc in hexane gradient) provided 47.6 mg of the intermediate.

25


Half of the above intermediate (24 mg) was dissolved in acetonitrile (1 mL), and an aqueous solution of HF (50%, 200 μL) was added. After 18.5 h, the reaction

30

solution was poured into an aqueous K_2HPO_4 solution (2.5 g in 20 mL) and extracted with EtOAc (2 x 25 mL). The combined organic layers were dried (Na_2SO_4), filtered, and concentrated under reduced pressure.

5 Palladium on carbon (10%, 12.5 mg) was added to a solution of the above intermediate in dioxane:MeOH (1:1, 1 mL). The flask was fitted with a septum and evacuated and back-filled three times with hydrogen gas. Hydrogen gas was bubbled through the reaction solution for three minutes, and the reaction mixture was stirred under an atmosphere (balloon) of hydrogen gas for 4.5 h. The reaction mixture was
10 filtered through celite to remove the palladium catalyst and concentrated under reduced pressure. Preparative reverse phase HPLC purification of the resulting oil was performed on a Waters Autopurification system using a Polymerx 10 μ RP- γ 100 R column [30 x 21.20 mm, 10 micron, solvent A: 0.05N HCl in water, solvent B: CH_3CN ; injection volume: 3.0 mL (0.05N HCl in water); gradient elution
15 with 0→30% B over 10 min, then held at 100% for 5 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 9.8-11.25 min, were collected and freeze-dried. The resulting impure powder was purified via preparative reverse phase HPLC as above with gradient elution with 15→50% B over 12 min, then held at 100% for 3 min; mass-directed fraction collection. Fractions with the desired
20 MW, eluting at 6.5-8.0 min, were collected and freeze-dried to yield 2.0 mg of compound **88** (5%): 1H NMR (400 MHz, CD_3OD) δ 8.25 (d, J = 11.0 Hz, 1 H), 5.29-5.24 (m, 1 H), 4.20-4.11 (m, 1 H), 4.09 (s, 1 H), 3.19-2.89 (m, 10 H), 2.69-2.56 (m, 1 H), 2.33-2.19 (m, 2 H), 1.68-1.56 (m, 1 H); MS (ESI) m/z 531.30 (M+H).

25 *N*-methyl-L-azetidine-2-carboxylic acid




To a suspension of L-azetidine-2-carboxylic acid (290 mg, 2.87 mmol, 1 equiv) in MeOH (3.6 mL), was added aqueous formaldehyde solution (37%, 235 μ L, 3.15 mmol, 1.1 equiv) and palladium on carbon (10%, 76 mg). The flask was fitted
30 with a septum and evacuated and back-filled three times with hydrogen gas. The reaction was stirred under an atmosphere (balloon) of hydrogen gas for 19 h, and was filtered through celite to remove the palladium catalyst. The resulting solution

was concentrated under reduced pressure, concentrated from toluene three times and dried under vacuum to afford *N*-methyl-L-azetidine-2-carboxylic acid: ^1H NMR (400 MHz, CD_3OD) δ 4.50 (t, $J = 9.5$ Hz, 1 H), 3.96 (dt, $J = 4.3, 9.8$ Hz, 1 H), 3.81 (q, $J = 9.8$ Hz, 1 H), 2.86 (s, 3 H), 2.71-2.60 (m, 1 H), 2.50-2.38 (m, 1 H).

5


Compound 4-1-1



To a suspension of aniline **2-6** (302 mg, 0.408 mmol, 1 equiv) and *N*-methyl-L-azetidine-2-carboxylic acid (148 mg, 1.28 mmol, 3.1 equiv) in CH_2Cl_2 (6 mL) was added O-(Benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate (395 mg, 1.23 mmol, 3 equiv) and DIEA (285 μL , 1.64 mmol, 4 equiv). After 16.5 h, the resulting orange solution was concentrated under reduced pressure and purified via preparative reverse phase HPLC on a Waters Autopurification system using a Sunfire Prep C18 OBD column [5 μm , 19 $\times$ 50 mm; flow rate, 20 mL/min; Solvent A: H_2O with 0.1% HCO_2H ; Solvent B: CH_3CN with 0.1% HCO_2H ; injection volume: 4 $\times$ 2.5-3.0 mL (CH_3CN); gradient: 50 $\rightarrow$ 90% B over 15 min; mass-directed fraction collection]. Two sets of fractions with the desired MW, eluting at 4.6-6.5 min and 6.5-9.4 min, were collected separately and freeze-dried to provide 147 mg of **4-1-1** (43%): ^1H NMR (400 MHz, CDCl_3) δ 16.04 (s, 1 H), 10.10 (s, 1 H), 8.48 (d, $J = 11.0$ Hz, 1 H), 7.54-7.48 (m, 4 H), 7.40-7.32 (m, 5 H), 5.36 (s, 2 H), 4.99 (d, $J = 9.8$ Hz, 1 H), 4.90 (d, $J = 9.8$ Hz, 1 H), 3.96 (d, $J = 10.4$ Hz, 1 H), 3.54 (t, $J = 7.9$ Hz, 1 H), 3.39-3.34 (m, 1 H), 3.25-3.19 (m, 1 H), 3.05-2.92 (m, 2 H), 2.58-2.36 (m, 10 H), 2.23-2.06 (m, 4 H), 0.81 (s, 9 H), 0.28 (s, 3 H), 0.11 (s, 3 H); MS (ESI) m/z 837.37 ($\text{M}+\text{H}$).

25

Compound **89**




To a solution of **4-1-1** (147 mg, 0.175 mmol, 1 equiv) in dioxane (3.5 mL) was added an aqueous solution of HF (50%, 750 μ L). After 4 h, the reaction solution was poured into an aqueous K_2HPO_4 solution (9 g in 90 mL) and extracted with EtOAc (2 x 50 mL). The combined organic layers were dried (Na_2SO_4), filtered, and concentrated under reduced pressure to provide 128.4 mg of a crude yellow foam.

The HF deprotection product (144 mg, 0.199 mmol, 1 equiv) was dissolved in dioxane:MeOH (1:1, 4 mL), and palladium on carbon (10%, 43.5 mg) was added. The flask was fitted with a septum and evacuated and back-filled three times with hydrogen gas. Hydrogen gas was bubbled through the reaction solution for three minutes, and the reaction mixture was stirred under an atmosphere (balloon) of hydrogen gas for 3.25 h. The reaction mixture was filtered through celite to remove the palladium catalyst and concentrated under reduced pressure. Preparative reverse phase HPLC of this oil was performed on a Waters Autopurification system using a Polymerx 10 μ RP- γ 100 R column [30 x 21.20 mm, 10 micron, solvent A: 0.05 N HCl in water, solvent B: CH_3CN ; injection volume: 2 x 3.2 mL (0.05 N HCl in water); gradient: 10 $\rightarrow$ 35% B over 20 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 6.10-8.40 min and 6.9-9.4 min, respectively for each run, were combined. The pH of the solution at 0 $^{\circ}C$ was adjusted (pH 1.8 to pH 7.4) via dropwise addition of 0.5 M aqueous NaOH solution (approximately 7.8 mL) and careful monitoring with an electronic pH meter. The aqueous solution was extracted with CH_2Cl_2 (3 x 60 mL) and the combined organic layers were dried (Na_2SO_4), filtered, and concentrated under reduced pressure to provide 79.7 mg of compound **89** as the free base (0.146 mmol, 73%). This yellow solid was dissolved in MeOH (3 mL), and $MeSO_3H$ (19 μ L, 0.292 mmol, 2 equiv) was added. The solution was concentrated under reduced pressure, dried under vacuum, and freeze-dried from water to provide 105 mg of **89** as the dimesylate salt. 1H NMR (400 MHz, CD_3OD) δ 8.22 (d, J = 11.0 Hz, 1 H), 5.16 (t, J = 8.6 Hz, 1 H), 4.21-4.12 (m, 1 H), 4.09-4.02 (m, 2 H), 3.17-2.85 (m, 10 H), 2.68 (s, 6 H, mesylate H), 2.64-2.59 (m, 1 H), 2.34-2.15 (m, 2 H), 1.70-1.58 (m, 1 H); MS (ESI) m/z 545.18 (M+H).

General Procedures for the Preparation of Compounds **90** - **94**

To a solution of aniline **2-6** (1 equiv) in THF (0.05-0.09M) was added an acid chloride (3 equiv). The reaction solution was filtered through celite and concentrated under reduced pressure. The resulting oil was dissolved in dioxane (1 mL) and an aqueous solution of HIF (50%, 200 μ L) was added. Upon completion, the reaction was poured into an aqueous K_2HPO_4 solution (2.6 g in 30 mL) and extracted with EtOAc (2 x 25 mL). The combined organic layers were dried (Na_2SO_4), filtered, and concentrated under reduced pressure. Palladium on carbon (10%) was added to a solution of this crude oil in dioxane:MeOH (1:1, 1 mL). The flask was fitted with a septum and evacuated and back-filled three times with hydrogen gas, and then the solution was degassed with bubbling hydrogen for 2 minutes. The reaction was stirred under an atmosphere (balloon) of hydrogen gas for 2 h. The reaction mixture was filtered through celite to remove the palladium catalyst and concentrated under reduced pressure. The crude products were purified by preparative reverse phase HPLC.


Compound **90**



Prepared by above general procedure with the following reagents: aniline **2-6** (21.1 mg, 0.028 mmol, 1 equiv), picolinoyl chloride hydrochloride (15.8 mg, 0.088, 3 equiv), with triethylamine (11.7 μ L, 0.084 mmol, 3 equiv), and 10% Pd-C (10 mg), provided a crude oil. Preparative reverse phase HPLC of the crude product was performed on a Waters Autopurification system using a Polymerx 10 μ RP- γ 100 R column [30 x 21.20 mm, 10 micron, solvent A: 0.05 N HCl in water, solvent B: CH₃CN; injection volume: 2.5 mL (0.05 N HCl in water); gradient: 10 $\rightarrow$ 60% B over 20 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 14.8-16.4 min, were collected and freeze-dried to provide 5.8 mg of the desired compound **90** (37%): ¹H NMR (400 MHz, CD₃OD) δ 8.73-8.69 (m, 1 H), 8.58-8.52 (m, 1 H), 8.27-8.21 (m, 1 H), 8.08-8.00 (m, 1 H), 7.66-7.60 (m, 1 H), 4.09 (s, 1 H),

3.29-2.92 (m, 9 H), 2.38-2.18 (m, 2 H), 1.72-1.60 (m, 1 H); MS (ESI) m/z 553.27 (M+H).


Compound **91**



5

Prepared by above general procedure with the following reagents: aniline **2-6** (31.0 mg, 0.042 mmol, 1 equiv), 1-methylpyrrole-2-carbonyl chloride (22 mg, 0.15 mmol, 3 equiv), and 10% Pd-C (10 mg). Preparative reverse phase HPLC of the crude product was performed on a Waters Autopurification system using a Polymerx
10 10 μ RP- γ 100 R column [30 $\times$ 21.20 mm, 10 micron, solvent A: 0.05 N HCl in water, solvent B: CH₃CN; injection volume: 2.0 mL (0.05 N HCl in water); gradient: 20 $\rightarrow$ 70% B over 20 min; mass-directed fraction collection]. Fractions with the desired MW were collected and freeze-dried and repurified via the same system with
15 the gradient: 10 $\rightarrow$ 60% B over 20 min. Fractions with the desired MW, eluting at 15.5-16.5 min, were collected and freeze-dried to provide 2.5 mg of the desired compound **91** (11%): ¹H NMR (400 MHz, CD₃OD) δ 8.20 (d, J = 11.6 Hz, 1 H), 6.98-6.86 (m, 2 H), 6.17-6.10 (m, 1 H), 4.08 (s, 1 H), 3.94 (s, 3 H), 3.19-2.90 (m, 9 H), 2.33-2.18 (m, 2 H), 1.80-1.56 (m, 1 H); MS (ESI) m/z 555.32 (M+H).


20 Compound **92**



Prepared by above general procedure with the following reagents: aniline **2-6** (31.0 mg, 0.042 mmol, 1 equiv), 5-methylisoxazole-3-carbonyl chloride (19.0 mg, 0.13 mmol, 3 equiv), and 10% Pd-C (10 mg). Preparative reverse phase HPLC of the
25 crude product was performed on a Waters Autopurification system using a Polymerx 10 μ RP- γ 100 R column [30 $\times$ 21.20 mm, 10 micron, solvent A: 0.05 N HCl in water, solvent B: CH₃CN; injection volume: 2.8 mL (0.05 N HCl in water); gradient:


10→60% B over 20 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 14.5-15.5 min, were collected and freeze-dried to provide 4.0 mg of the desired compound **92** (17%): ¹H NMR (400 MHz, CD₃OD) δ 8.32 (d, *J* = 11.0 Hz, 1 H), 6.59 (s, 1 H), 4.09 (s, 1 H), 3.19-2.90 (m, 9 H), 2.52 (s, 3 H), 2.34-2.18 (m, 2 H), 1.71-1.58 (m, 1 H); MS (ESI) *m/z* 557.26 (M+H).

Compound **93**



Prepared by above general procedure with the following reagents: aniline **2-6** (30.0 mg, 0.041 mmol, 1 equiv), 1-methyl-1H-pyrazole-3-carbonyl chloride (16.8 mg, 0.12 mmol, 3 equiv), and 10% Pd-C (20 mg). Preparative reverse phase HPLC of the crude product was performed on a Waters Autopurification system using a Polymerx 10 μ RP-γ 100 R column [30 × 21.20 mm, 10 micron, solvent A: 0.05 N HCl in water, solvent B: CH₃CN; injection volume: 3.2 mL (0.05 N HCl in water); gradient: 10→60% B over 20 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 12.5-14.5 min, were collected and freeze-dried to provide 11.2 mg of the desired compound **93** (49%): ¹H NMR (400 MHz, CD₃OD) δ 8.38 (d, *J* = 11.0 Hz, 1 H), 7.68 (s, 1 H), 6.82-6.76 (m, 1 H), 4.09 (s, 1 H), 3.99 (s, 3 H), 3.16-2.90 (m, 9 H), 2.31-2.16 (m, 2 H), 1.70-1.56 (m, 1 H); MS (ESI) *m/z* 556.31 (M+H).

Compound **94**




Prepared by above general procedure with the following reagents: aniline **2-6** (30.0 mg, 0.041 mmol, 1 equiv), 1,3-thiazole-2-carbonyl chloride (17.8 mg, 0.12 mmol, 3 equiv), and 10% Pd-C (15 mg). Preparative reverse phase HPLC of the crude product was performed on a Waters Autopurification system using a Polymerx

10 μ RP- γ 100 R column [30 $\times$ 21.20 mm, 10 micron, solvent A: 0.05 N HCl in water, solvent B: CH₃CN; injection volume: 3.2 mL (0.05 N HCl in water), gradient: 10 $\rightarrow$ 60% B over 20 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 14.6-17.0 min, were collected and freeze-dried to provide 5.4 mg of the desired compound **94** (23%): ¹H NMR (400 MHz, CD₃OD) δ 8.38 (d, J = 11.0 Hz, 1 H), 8.02 (d, J = 3.0 Hz, 1 H), 7.95 (d, J = 2.4 Hz, 1 H), 4.09 (s, 1 H), 3.20-2.90 (m, 9 H), 2.34-2.17 (m, 2 H), 1.70-1.56 (m, 1 H); MS (ESI) m/z 559.23 (M+H).

Example 5. Synthesis of Compounds of Structural Formula (A), wherein Y is -N(R^A)(R^B), or -NH-SO₂-(CH₂)₂-N(R^A)(R^B).


Scheme 5



In Scheme 5, R represents $-(C_1-C_6)$ alkyl, $-(C_0-C_5)$ alkylene-carbocyclyl, $-(C_0-C_5)$ alkylene-aryl, $-(C_0-C_5)$ alkylene-heterocyclyl, $-(C_0-C_5)$ alkylene-heteroaryl, $-(C_1-C_3)$ alkylene- $N(R^A)(R^B)$; Ar represents an aryl or a heteroaryl group; and R^U and R^V are R^A and R^B , respectively, as defined in Structural Formula (B).

5 The following compounds were prepared according to Scheme 5.


Compound **5-1-1**



Compound **2-6** (150 mg, 0.203 mmol, 1.0 equiv) was dissolved in 1,2-dichloroethane (3 mL). HOAc (58.1 μ L, 1.01 mmol, 5 equiv.) and isovaleraldehyde (32.9 μ L, 0.304 mmol, 1.5 equiv) were added. The mixture was stirred for 1 h. Na(OAc)₃BH (129 mg, 0.609 mmol, 3.0 equiv) was added and the resulting mixture was stirred for another hour. The mixture was washed with H₂O (10 mL) and concentrated to give crude **5-1-1** (250 mg), which was used for the next step without further purification: MS (ESI) m/z 810.59 (M+H).

10
15

Compound **95**



Aqueous HF (0.3 mL, 48-50%) was added to a CH₃CN solution (1.5 mL) of **5-1-1** (250 mg crude) in a plastic vial at 25 °C. The reaction was stirred at 25 °C for 18 hrs. The resulting mixture was poured into an aqueous solution (10 mL) of K₂HPO₄ (2 g). The solution was extracted with EtOAc (3 x 15 mL). The combined EtOAc extracts were dried over sodium sulfate and concentrated to give the crude intermediate (155 mg).

20


10% Pd-C (20 mg) was added to a dioxane/MeOH solution (4 mL, 1:1) of the above crude intermediate. HCl/MeOH (0.5 mL, 0.5 N) was also added. The reaction mixture was stirred under H₂ (balloon) at 25 °C for 2 hrs and filtered

25

through a pad of Celite. The filtrate was concentrated to give the crude product 144 mg. The crude product was purified by HPLC on a Polymerx 10 μ RP- γ 100 R column [30 $\times$ 21.20 mm, 10 micron, solvent A: 0.05 N HCl, solvent B: CH₃CN, sample in 2.0 mL (0.05 N HCl), gradient elution with 10 $\rightarrow$ 100% B over 15 min, mass-directed fraction collection] to yield the desired product **95** as a yellow solid (82 mg, 78 %, 2 steps): ¹H NMR (400 MHz, CD₃OD) δ 7.44 (d, J = 9.2 Hz, 1 H), 4.12 (s, 1 H), 3.42-3.37 (m, 2 H), 3.05 (s, 3 H), 2.97 (s, 3 H), 3.21 -2.97 (m, 3 H), 2.39-2.30 (m, 1 H), 2.29-2.22 (m, 1 H), 1.79-1.59 (m, 4 H), 0.98 (d, J = 6.4 Hz, 6 H); MS (ESI) m/z 518.43 (M+H).


- 10 Compounds **96** – **101** were prepared similarly to compound **95** using the corresponding aldehydes in the reductive alkylation step.

Compound **96**




- 15 ¹H NMR (400 MHz, CD₃OD) δ 7.39 (d, J = 9.2 Hz, 1 H), 4.10 (s, 1 H), 3.34 (t, J = 7.8 Hz, 2 H), 3.04 (s, 3 H), 2.96 (s, 3 H), 3.21 -2.95 (m, 3 H), 2.35 (t, J = 13.7 Hz, 1 H), 2.27-2.20 (m, 1 H), 1.82-1.72 (m, 2 H), 1.71-1.60 (m, 1 H), 1.05 (t, J = 7.4 Hz, 3 H); MS (ESI) m/z 490.32 (M+H).

- 20 Compound **97**




- 25 ¹H NMR (400 MHz, CD₃OD) δ 7.34 (d, J = 9.2 Hz, 1 H), 4.10 (s, 1 H), 3.34 (t, J = 7.8 Hz, 2 H), 3.04 (s, 3 H), 2.96 (s, 3 H), 3.24 -2.95 (m, 11 H), 2.33 (t, J = 13.7 Hz, 1 H), 2.27-2.20 (m, 1 H), 2.11-1.98 (m, 1 H), 1.71-1.60 (m, 1 H), 1.08 (d, J = 6.9 Hz, 6 H); MS (ESI) m/z 504.46 (M+H).

Compound **98**




¹H NMR (400 MHz, CD₃OD) δ 7.43 (d, *J* = 8.7 Hz, 1 H), 4.10 (s, 1 H), 3.34 (t, *J* = 7.8 Hz, 2 H), 3.04 (s, 3 H), 2.96 (s, 3 H), 3.28-2.95 (m, 11 H), 2.41-2.31 (m, 1 H), 2.27-2.20 (m, 1 H), 2.11-1.98 (m, 1 H), 1.72-1.60 (m, 1 H), 1.20-1.11 (m, 1 H), 0.74-0.68 (m, 2 H), 0.43-0.38 (m, 2 H); MS (ESI) *m/z* 502.40 (M+H).

Compound 99




¹H NMR (400 MHz, CD₃OD) δ 6.97-6.89 (m, 1 H), 4.07 (s, 1 H), 3.34 (t, *J* = 7.8 Hz, 2 H), 3.03 (s, 3 H), 2.95 (s, 3 H), 3.14-2.92 (m, 11 H), 2.30-2.15 (m, 2 H), 1.89-1.59 (m, 7 H), 1.38-1.20 (m, 3 H), 1.11-1.00 (m, 2 H); MS (ESI) *m/z* 544.50 (M+H).

Compound 100




¹H NMR (400 MHz, CD₃OD) δ 6.83 (d, *J* = 10.5 Hz, 1 H), 4.06 (s, 1 H), 3.34 (t, *J* = 7.8 Hz, 2 H), 3.03 (s, 3 H), 2.95 (s, 3 H), 3.11-2.93 (m, 5 H), 2.27-2.14 (m, 2 H), 1.67-1.57 (m, 1 H), 1.04 (s, 9 H); MS (ESI) *m/z* 518.48 (M+H).

Compound 101




^1H NMR (400 MHz, CD_3OD) δ 7.46-7.42 (m, 1 H), 4.15 (s, 1 H), 3.33 (s, 6 H), 3.04 (s, 3 H), 2.96 (s, 3 H), 3.17-2.95 (m, 3 H), 2.44-2.34 (m, 1 H), 2.29-2.22 (m, 1 H), 1.71-1.60 (m, 1 H); MS (ESI) m/z 476.29 ($\text{M}+\text{H}$).

5 Compound 102




Prepared similarly to **95** using $\text{tBuN}(\text{Cbz})\text{CH}_2\text{CHO}$: ^1H NMR (400 MHz, CD_3OD) δ 6.72 (d, $J = 11.0$ Hz, 1 H), 4.07 (s, 1 H), 3.54-3.46 (m, 2 H), 3.26-3.19 (m, 2 H), 3.03 (s, 3 H), 2.95 (s, 3 H), 3.14-2.92 (m, 3 H), 2.23-2.14 (m, 2 H), 1.67-1.55 (m, 1 H), 1.38 (s, 9 H); MS (ESI) m/z 547.51 ($\text{M}+\text{H}$).

Compound 103



Compound **103** was also isolated from the preparation of **102**: ^1H NMR (400 MHz, CD_3OD) δ 6.71 (d, $J = 11.0$ Hz, 1 H), 4.07 (s, 1 H), 3.47 (t, $J = 6.0$ Hz, 2 H), 3.17 (t, $J = 6.0$ Hz, 2 H), 3.03 (s, 3 H), 2.95 (s, 3 H), 3.13-2.92 (m, 3 H), 2.23-2.12 (m, 2 H), 1.66-1.54 (m, 1 H); MS (ESI) m/z 491.42 ($\text{M}+\text{H}$).

Compound 5-3-1




20

A vessel containing aniline **2-6** (18.2 mg, 0.024 mmol, 1 equiv), Pd_2dba_3 (3.0 mg, 0.0033 mmol, 0.13 equiv), Xantphos (3.4 mg, 0.0059 mmol, 0.25 equiv), K_3PO_4 (40 mg, 0.188 mmol, 7.8 equiv) and 4,6-dichloropyrimidine (6.5 mg, 0.044 mmol, 1.8 equiv) was evacuated and back-filled with nitrogen gas three times. Dioxane (500 μL) was added, and the reaction mixture stirred vigorously and heated at 80 $^\circ\text{C}$

25

for 4.5 h. The reaction mixture was filtered through celite and concentrated under reduced pressure. Preparative reverse phase HPLC of the resulting yellow oil was performed on a Waters Autopurification system using a Sunfire Prep C18 OBD column [5 μ m, 19 $\times$ 50 mm; flow rate, 20 mL/min; Solvent A: H₂O with 0.1% HCO₂H; Solvent B: CH₃CN with 0.1% HCO₂H; injection volume: 1.8 mL (CH₃CN); gradient: 80 $\rightarrow$ 100% B over 15 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 9.2-9.8 min, were collected and freeze-dried to provide 7.5 mg of compound **5-3-1** (37%). ¹H NMR (400 MHz, CDCl₃) δ 15.97 (s, 1 H), 8.48 (s, 1 H), 8.33 (d, *J* = 5.5 Hz, 1 H), 7.52-7.46 (m, 2 H), 7.40-7.28 (m, 8 H), 7.07 (s, 1 H), 6.11 (s, 1 H), 5.34 (s, 2 H), 4.97 (d, *J* = 11.6 Hz, 1 H), 4.88 (d, *J* = 11.0 Hz, 1 H), 3.95 (d, *J* = 10.4 Hz, 1 H), 3.28-3.19 (m, 1 H), 3.09-2.98 (m, 1 H), 2.61-2.54 (m, 1 H), 2.54-2.39 (m, 8 H), 2.16 (d, *J* = 14.6 Hz, 1 H), 0.83 (s, 9 H), 0.28 (s, 3 H), 0.14 (s, 3 H); MS (ESI) *m/z* 852.57 (M+H).


Compound 104



To a solution of **5-3-1** (7.5 mg, 0.0088 mmol, 1 equiv) in dioxane (1.4 mL) was added an aqueous solution of HF (50%, 200 μ L). After 15.5 h, the reaction solution was poured into an aqueous K₂HPO₄ solution (2.4 g in 20 mL) and extracted with EtOAc (2 $\times$ 20 mL). The combined organic layers were dried (Na₂SO₄), filtered, and concentrated under reduced pressure. Palladium on carbon (10%, 10 mg) was added to a solution of this oil in dioxane:MeOH (1:1, 1 mL). The flask was fitted with a septum and evacuated and back-filled three times with hydrogen gas. Hydrogen gas was bubbled through the reaction solution for three minutes, and the reaction mixture was stirred under an atmosphere (balloon) of hydrogen gas for 2.5 h. The reaction mixture was filtered through celite to remove the palladium catalyst and concentrated under reduced pressure. Preparative reverse phase HPLC purification of the resulting oil was performed on a Waters Autopurification system using a Polymerx 10 μ RP- γ 100 R column [30 $\times$ 21.20 mm, 10 micron, solvent A: 0.05 N HCl in water, solvent B: CH₃CN;

injection volume: 2.0 mL (0.05 N HCl in water), gradient elution with 10→50% B over 10 min, then held at 100% for 5 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 6.90-7.80 min, were collected and freeze-dried to provide 2.2 mg of **104** (48%). ¹H NMR (400 MHz, CD₃OD) δ 8.83 (s, 1 H), 8.37-8.25 (m, 1 H), 8.18-8.05 (m, 1 H), 7.30-7.20 (m, 1 H), 4.10 (s, 1 H), 3.20-2.90 (m, 9 H), 2.40-2.29 (m, 1 H), 2.27-2.19 (m, 1 H), 1.72-1.58 (m, 1 H); MS (ESI) *m/z* 526.31 (M+H).

Compound **105**



10

To a solution of aniline **2-6** (30.0 mg, 0.041 mmol, 1 equiv) in 1,2-dichloroethane (500 μL) was added pyridine (16.3 μL, 0.20 mmol, 5 equiv) and 2-chloroethanesulfonyl chloride (21 μL, 0.20 mmol, 5 equiv). The reaction vessel was sealed and heated to 45 °C. After one hour, the reaction was a solid yellow gel, and another 500 μL 1,2-dichloroethane was added to form a suspension and the reaction was sealed and heated to 45 °C. After 18.5 h pyrrolidine (68 μL, 0.82 mmol, 20 equiv) was added and the reaction heated to 45 °C for 2.5 hours. The solution was poured into aqueous pH 7 phosphate buffer (8 mL) and extracted with EtOAc (2 x 25 mL). The combined organic layers were dried (Na₂SO₄), filtered, and concentrated under reduced pressure. To a solution of this crude oil in CH₃CN (1.8 mL) was added an aqueous solution of HIF (50%, 300 μL). After 15 h, the reaction solution was poured into an aqueous K₂HPO₄ solution (3.6 g in 30 mL) and extracted with EtOAc (3 x 15 mL). The combined organic layers were dried (Na₂SO₄), filtered, and concentrated under reduced pressure. Palladium on carbon (10%, 8.4 mg) was added to a solution of this oil in dioxane:MeOH (1:1, 1.2 mL). The flask was fitted with a septum and evacuated and back-filled three times with hydrogen gas, and the reaction mixture was stirred under an atmosphere (balloon) of hydrogen gas for 1.5 h. Another 10 mg palladium catalyst was added and the reaction was evacuated and back-filled with hydrogen gas as before. After 6 h, the reaction mixture was filtered through celite to remove the palladium catalyst and

15


20

25

30

concentrated under reduced pressure. Preparative reverse phase HPLC purification of the resulting oil was performed on a Waters Autopurification system using a Polymerx 10 μ RP- γ 100 R column [30 $\times$ 21.20 mm, 10 micron, solvent A: 0.05 N HCl in water, solvent B: CH₃CN; injection volume: 3.5 mL (0.05 N HCl in water); gradient elution with 0 $\rightarrow$ 100% B over 10 min, then held at 100% for 5 min; mass-directed fraction collection]. Two sets of fractions with the desired MW, eluting at 6.3-7.1 min and 8.7-9.3 min, were collected separately and freeze-dried to provide 9.7 mg of crude compounds **105**. Purification via preparative reverse phase HPLC with gradient elution with 20 $\rightarrow$ 70% B over 20 min; mass-directed fraction collection] provided 3.3 mg of pure **105** (13%): ¹H NMR (400 MHz, CD₃OD) δ 7.44 (d, J = 9.8 Hz, 1 H), 4.09 (s, 1 H), 3.79-3.65 (m, 4 H), 3.63-3.56 (m, 2 H), 3.18-2.90 (m, 11 H), 2.35-2.26 (m, 1 H), 2.26-2.10 (m, 3 H), 2.10-1.96 (m, 2 H), 1.69-1.59 (m, 1 H); MS (ESI) m/z 609.36 (M+H).


15 Compound **106**



Compound **106** (1.7 mg, 7%) was also isolated from the preparation of compound **105**: ¹H NMR (400 MHz, CD₃OD) δ 6.71 (d, J = 11.0 Hz, 1 H), 4.06 (s, 1 H), 3.67-3.60 (m, 2 H), 3.38-3.33 (m, 4 H), 3.09-2.90 (m, 9 H), 2.24-2.13 (m, 2 H), 1.95-1.91 (m, 5 H), 1.90-1.85 (m, 1 H), 1.68-1.55 (m, 1 H); MS (ESI) m/z 609.36 (M+H).


Example 6. Synthesis of Compounds 107 and 108

Scheme 6




The following compounds were prepared according to Scheme 6.

5 Compound 6-1




Compound 2-3 (5.0 g, 14.25 mmol, 1.0 equiv) in MeOH (20 mL) was added a aqueous solution of HCHO (2.3 g, 37%, 28.50 mmol, 2.0 equiv) and palladium on carbon (0.5 g, 10 wt%). The reaction was purged with hydrogen and stirred under H₂ (balloon) at room temperature for 2 hours. The reaction mixture was filtered through celite and concentrated to afford 1.3 g crude compound 6-1 as a yellow solid.

Compound 6-2



To compound **6-1** (0.9 g, 3.27 mmol, 1.0 equiv) in DCM was added Boc_2O (2.14 g, 9.81 mmol, 3.0 equiv) dropwise. DMAP (135 mg, 15 wt%) was added to the mixture and the reaction was stirred at room temperature for 1 hour. Then the reaction mixture was heated to reflux for 1 hour. The reaction mixture was concentrated. The crude compound was purified by column chromatography on silica gel eluted with (PE:EA= 200:1→100:1) to yield compound **6-2** (1.16 g, 73.4%) as a light yellow solid. ^1H NMR (400 MHz, DMSO) δ 7.60 (d, J = 10.0 Hz, 1 H), 7.53-7.44 (m, 2 H), 7.36-7.31 (m, 1 H), 7.28-7.22 (m, 2 H), 3.06 (s, 3 H), 2.33 (d, J = 2.0 Hz, 3 H), 1.38 (s, 9 H), 1.34 (s, 9 H); MS (ESI) m/z 476.2 (M+H).


Compound **6-3**



To diisopropylamine (0.28 mL, 3.2 mmol, 10.0 equiv) in THF at -78 °C was added $n\text{BuLi}$ (0.8 mL, 2.50 M/hexane, 3.2 mmol, 10.0 equiv) and TMEDA (0.40 mL, 5.0 mmol, 10.0 equiv) at -78 °C dropwise. The reaction was stirred at -78 °C for 40 min. Compound **6-2** (480 mg, 1.0 mmol, 3.0 equiv) in THF was added to the reaction mixture dropwise at -78 °C. The resulting deep-red solution was stirred at -78 °C 60 min, the enone (160 mg 0.33 mmol, 1.0 equiv) in THF was added to the mixture dropwise at -78 °C. The deep-red solution was gradually warmed up with stirring from -78 °C to -20 °C over a period of 1 h. The resulting orange solution was brought to 0 °C, and quenched with aqueous saturated ammonium chloride (100 mL). The yellow-green mixture was extracted with EtOAc two times. The combined EtOAc extracts were dried (Na_2SO_4) and concentrated to yield the crude product. Flash column chromatography on silica gel with 0%, 5%, 10%, EtOAc/hexane sequentially yielded the desired product **6-3** as a light-yellow solid (42 mg, 14.8%). ^1H NMR (400 MHz, CDCl_3) δ 15.70 (s, 1 H), 7.52-7.50 (m, 2 H), 7.42-7.33 (m, 3 H), 7.16 (d, J = 8.4 Hz, 1 H), 5.37 (s, 2 H), 3.95 (d, J = 10.8 Hz, 1 H), 3.28-3.23 (m,

1 H), 3.14 (s, 3 H), 3.10-3.05 (m, 1 H), 2.58-2.47 (m, 9 H), 2.16 (d, $J = 14.0$ Hz, 1 H), 1.53 (s, 9 H), 1.42 (s, 9 H), 0.89 (s, 9 H), 0.29 (s, 3 H), 0.15 (s, 3 H); MS (ESI) m/z 864.43 (M+H).


5 Compound **107**



Compound **6-3** (120 mg, 0.14 mmol) was dissolved in THF (5 mL) and aqueous HIF (40%, 2 mL) was added dropwise. The yellow solution was stirred at room temperature overnight. The resulting deep-red solution was slowly added into an aqueous K_2HPO_4 solution with stirring. The pH of the mixture was adjusted by aqueous K_2HPO_4 solution to about 8. The yellow mixture was extracted with EtOAc two times. The combined EtOAc extracts were dried (Na_2SO_4) and concentrated to yield the crude product.


The above crude compound (120 mg, crude, ~0.14 mmol, 1.0 equiv) was dissolved in HPLC grade MeOH(10 mL) and 10% Pd-C (25 mg, 0.03 mmol, 0.2 equiv) were added. The mixture was purged with hydrogen by bubbling hydrogen through with gentle stirring for 5 min. The reaction was then vigorously stirred under hydrogen balloon at room temperature for 2 hr. LC-MS analysis indicated the reaction completed. The catalyst was filtered off and the mixture was concentrated, the residue was purified by reverse phase HPLC to afford the desired compound **107** (50 mg, 78%) as a yellow solid. ^1H NMR (400 MHz, CD_3OD) δ 7.462 (d, $J = 8.4$ Hz, 1 H), 4.14 (s, 1 H), 3.21-2.93 (m, 9 H), 3.10 (s, 3 H), 2.38-2.25 (m, 2 H), 1.68-1.62 (m, 1 H); MS (ESI) m/z 462.2 (M+H).

25 Compound **108**



The following compounds were prepared according to Scheme 7.

Compound 7-2




5 To Boc-L-glutamic acid-1-benzyl ester (**7-1**) (3.00 g, 8.89 mmol, 1.0 eq) in DMF (20 mL) at rt was added potassium carbonate (1.84 g, 13.33 mmol, 1.5 eq) and methyl iodide (0.67 mL, 10.74 mmol, 1.2 eq). The mixture was diluted with EtOAc (200 mL), washed with water (200 mL), saturated aqueous sodium bicarbonate (100 mL x 2), and brine (100 mL x 1). The EtOAc solution was dried over sodium sulfate and concentrated in vacuo: R_f 0.33 (20% EtOAc/hexane).

10 Boc₂O (2.91 g, 13.33 mmol, 1.5 eq), DMAP (54 mg, 0.44 mmol, 0.05 eq), and DIEA (3.10 mL, 17.80 mmol, 2 eq) were added to the above intermediate in acetonitrile (20 mL). The solution was stirred at rt for 60 hrs, added with saturated aqueous sodium bicarbonate (100 mL), and extracted with EtOAc (100 mL x 1, 50 mL x 2). The EtOAc extracts were combined, dried over sodium sulfate, and concentrated in vacuo to yield the desired product **7-2** as a pale liquid (quantitative): R_f 0.45 (20% EtOAc/hexane); ¹H NMR (400 MHz, CDCl₃) δ 7.25-7.35 (m, 5 H), 5.14 (s, 2 H), 4.95 (dd, J = 4.9, 9.8 Hz, 1 H), 3.65 (s, 3 H), 2.43-2.52 (m, 1 H), 2.37-2.42 (m, 2 H), 2.15-2.25 (m, 1 H), 1.42 (s, 18 H); MS (ESI) m/z 452.3 (M+H).

20

Compound 7-3



To compound **7-2** (8.89 mmol, 1 eq) in anhydrous diethyl ether (40 mL) at -78 °C was added DIBAL-H (12.33 mL, 1 M/hexane, 12.33 mmol, 1.25 eq) dropwise. The reaction was stirred at -78 °C for 2 hrs. Additional DIBAL-H (1.20 mL, 1 M/hexane, 1.20 mmol) was added. The reaction was stirred at -78 °C for another 1 hr and quenched with HOAc (2.80 mL) at -78 °C. The reaction was warmed to rt and added with 10% aqueous sodium carbonate (75 mL). The mixture was stirred for 15 min and extracted with methylene chloride (200 mL x 1, 50 mL x 2). The methylene chloride extracts were combined, dried over sodium sulfate, and

30


Compound 7-5



15


25

Compound 7-6



To a suspension of compound **7-5** (0.044 mmol, 1 eq) and sodium carbonate (7 mg, 0.066 mmol, 1.5 eq) in THF at rt was added HATU (20 mg, 0.053 mmol, 1.2 eq). The mixture was rapidly stirred at rt for 2 hrs. Methanol (5 mL) was added. The solids were filtered off. The filtrate was concentrated under reduced pressure to yield crude **7-6** as a yellow solid; MS (ESI) m/z 645.1 (M+H).

Compound **109**




Compound **7-6** (0.044 mmol) was treated with 4 N HCl/dioxane (5 mL) at rt for overnight and concentrated in vacuo. The residue was re-dissolved in methanol (1 mL) and added dropwise into diethyl ether (50 mL) with rapid stirring. The yellow precipitates were collected, washed with more diethyl ether (5 mL x 3), and dried under vacuo to afford crude **109** as a brown solid.

One fifth of the above crude product was purified by reverse phase HPLC to yield pure **109** as a yellow solid (1.5 mg, 31%); ^1H NMR (400 MHz, CD_3OD) δ 7.36 (d, 9.2 Hz, 1 H), 4.09-4.15 (m, 1 H), 4.08 (s, 1 H), 3.70-3.80 (m, 1 H), 3.58-3.68 (m, 1 H), 2.90-3.50 (m, 12 H), 2.30-2.45 (m, 2 H), 2.10-2.25 (m, 3 H), 1.95-2.10 (m, 1 H), 1.58-1.70 (m, 1 H); MS (ESI) m/z 545.1 (M+H).

20


Compound **110**



To 2/5 of crude **109** (0.018 mmol, 1 eq) in DMF (1 mL) was added aqueous formaldehyde (0.007 mL, 36.5%/water, 0.094 mmol, 5 eq), InCl_3 (0.4 mg, 0.002


mmol, 0.1 eq), and Na(OAc)₃BH (15 mg, 0.071 mmol, 4 eq). The reaction was stirred at rt for 2 hrs and quenched with 0.5 N HCl/methanol (1 mL). The solution was added dropwise into diethyl ether (100 mL) with rapid stirring. The precipitates were collected, washed with more diethyl ether (2 mL x 4), and purified by reverse phase HPLC to afford the desired compound **110** as a yellow solid (1.8 mg, 18%):
¹H NMR (400 MHz, CD₃OD) δ 7.744 (d, *J* = 9.1 Hz, 1 H), 4.37 (dd, *J* = 6.1, 11.6 Hz, 1 H), 4.09 (s, 1 H), 3.60-3.75 (m, 2 H), 2.92-3.50 (m, 15 H), 2.86 (s, 3 H), 2.10-2.50 (m, 6 H), 1.60-1.72 (m, 1 H); MS (ESI) *m/z* 573.3 (M+H).

10 Compound **111**



To 2/5 of crude **109** (0.018 mmol, 1 eq) in DMF (1 mL) was added cyclopropanecarboxaldehyde (1.4 μL, 0.018 mmol, 1 eq), InCl₃ (0.4 mg, 0.002 mmol, 0.1 eq), and Na(OAc)₃BH (6 mg, 0.028 mmol, 1.5 eq). The reaction was stirred at rt for overnight and quenched with 0.5 N HCl/methanol (1 mL). The solution was added dropwise into diethyl ether (100 mL) with rapid stirring. The precipitates were collected, washed with more diethyl ether (2 mL x 4), and purified by reverse phase HPLC to afford the desired compound **111** as a yellow solid (1.3 mg, 12%):
¹H NMR (400 MHz, CD₃OD) δ 7.38 (d, *J* = 9.2 Hz, 1 H), 4.22 (dd, *J* = 6.1, 11.6 Hz, 1 H), 4.09 (d, 1 H), 3.60-3.78 (m, 2 H), 2.85-3.50 (m, 12 H), 2.00-2.50 (m, 6 H), 1.60-1.72 (m, 1 H), 1.10-1.20 (m, 1 H), 0.70-0.75 (m, 2 H), 0.40-0.50 (m, 2 H); MS (ESI) *m/z* 599.4 (M+H).

Compound **112**




25

Dialkylated product **112** was also isolated from the preparation of compound **111** (1.0 mg, yellow solid, 9%): ¹H NMR (400 MHz, CD₃OD) δ 7.42 (d, *J* = 9.2 Hz,

1 H), 4.70-4.80 (m, 1 H), 4.09 (s, 1 H), 3.55-3.80 (m, 3 H), 2.95-3.50 (m, 13 H), 2.10-2.50 (m, 6 H), 1.55-1.75 (m, 1 H), 1.20-1.30 (m, 2 H), 0.68-0.90 (m, 4 H), 0.38-0.58 (m, 4 H); MS (ESI) m/z 653.3 (M+H).

- 5 **Example 8. Synthesis of Compounds of Structural Formula (A), wherein Y is $-(C_1-C_4)\text{alkylene}-N(R^A)(R^B)$, or $-(C_1-C_4)\text{alkylene}-N(R^B)-C(O)-[C(R^D)(R^E)]_{0-4}-N(R^A)(R^B)$,**


Scheme 8



In Scheme 8, R and R' is R^B and R^A , respectively, as defined in Structural Formula (A) and R^w represents $-[C(R^D)(R^E)]_{1-4}-N(R^A)(R^B)$.

The following compounds were prepared according to Scheme 8.


- 15 **Compound 113**



Benzyl N-(hydroxymethyl)carbamate (92 mg, 0.51 mmol, 2.0 equiv) was added to a TFA/ CH_3SO_3H (1 mL/1 mL) solution of compound 7 (110 mg,

0.25 mmol) at 25 °C. The reaction was stirred at 25 °C for 30 min. Preparative reverse phase HPLC purification on a Waters Autopurification system using a Phenomenex Polymerx 10 μ RP-1 100A column [10 μ m, 150 $\times$ 21.20 mm; flow rate, 20 mL/min; Solvent A: 0.05 N HCl; Solvent B: CH₃CN; injection volume: 4.0 mL (0.05 N HCl/water); gradient: 0 $\rightarrow$ 30% B over 20 min; mass-directed fraction collection]. Fractions with the desired MW were collected and freeze-dried to yield 23 mg of pure **113**: ¹H NMR (400 MHz, CD₃OD) δ 7.47 (d, J = 9.2 Hz, 1 H), 4.16 (s, 2 H), 4.13 (s, 1 H), 3.21-2.94 (m, 3 H), 3.06 (s, 3 H), 2.97 (s, 3), 2.37-2.22 (m, 2 H), 1.70-1.58 (m, 1 H); MS (ESI) m/z 462.26 (M-H).


Compound **114**



Et₃N (2 μ L, 0.0136 mmol, 2.0 equiv) was added to a mixture of **113** (3 mg, 0.0065 mmol) and pivaldehyde (0.8 μ L, 0.00715 mmol, 1.1 equiv) in DMF (0.1 mL) at 25 °C. The reaction was stirred at 25 °C for 15 min. NaBH(OAc)₃ (3 mg, 0.013 mmol) and HOAc (2 μ L) was added to the resulting mixture. The reaction was stirred at 25 °C for 1 h. Preparative reverse phase HPLC purification on a Waters Autopurification system using a Phenomenex Polymerx 10 μ RP-1 100A column [10 μ m, 150 $\times$ 21.20 mm; flow rate, 20 mL/min, Solvent A: 0.05 N HCl; Solvent B: CH₃CN; injection volume: 4.0 mL (0.05 N HCl/water); gradient: 0 $\rightarrow$ 100% B over 15 min; mass-directed fraction collection]. Fractions with the desired MW were collected and freeze-dried to yield 1 mg of **114**: ¹H NMR (400 MHz, CD₃OD) δ 7.52 (d, J = 9.1 Hz, 1 H), 4.30 (s, 2 H), 4.09 (s, 1 H), 3.23-2.93 (m, 5 H), 3.04 (s, 3 H), 2.95 (s, 3 H), 2.40-2.19 (m, 2 H), 1.71-1.60 (m, 1 H), 1.05 (s, 9 H); MS (ESI) m/z 532.27 (M+H).

Compounds **115** - **118** were prepared similarly to compound **114** using the corresponding aldehydes.


Compound **115**



^1H NMR (400 MHz, CD_3OD) δ 7.51 (d, $J = 8.8$ Hz, 1 H), 4.25 (s, 2 H), 4.10 (s, 1 H), 3.25-2.90 (m, 5 H), 3.05 (s, 3 H), 2.96 (s, 3 H), 2.40-2.21 (m, 2 H), 1.90-1.60 (m, 7 H), 1.42-0.95 (m, 5 H); MS (ESI) m/z 558.31 ($\text{M}+\text{H}$).

5


Compound **116**



^1H NMR (400 MHz, CD_3OD) δ 7.50 (d, $J = 9.0$ Hz, 1 H), 4.24 (s, 2 H), 4.09 (s, 1 H), 3.25-2.90 (m, 5 H), 3.07 (s, 3 H), 2.94 (s, 3 H), 2.40-2.21 (m, 2 H), 1.82-1.58 (m, 3 H), 1.01 (t, $J = 6.7$ Hz, 3 H); MS (ESI) m/z 504.22 ($\text{M}+\text{H}$).

10


Compound **117**



^1H NMR (400 MHz, CD_3OD) δ 7.51 (d, $J = 8.9$ Hz, 1 H), 4.23 (s, 2 H), 4.09 (s, 1 H), 3.25-2.92 (m, 4), 3.02 (s, 3 H), 2.95 (s, 3 H), 2.40-2.19 (m, 2 H), 1.71-1.60 (m, 1 H), 1.40 (d, $J = 7.0$ Hz, 6 H); MS (ESI) m/z 504.23 ($\text{M}+\text{H}$).

15


Compound **118**



^1H NMR (400 MHz, CD_3OD) δ 7.54 (d, $J = 9.1$ Hz, 1 H), 4.37 (s, 2 H), 4.10 (s, 1 H), 3.20-2.85 (m, 3 H), 3.05 (s, 3 H), 2.97 (s, 3 H), 2.91 (s, 3 H), 2.90 (s, 3 H), 2.42-2.20 (m, 2 H), 1.72-1.60 (m, 1 H); MS (ESI) m/z 490.19 ($\text{M}+\text{H}$).


20

Compound **119**




Prepared from compound **114** by reductive alkylation using formaldehyde
 5 under similar conditions: ^1H NMR (400 MHz, CD_3OD) δ 7.57 (d, $J = 9.1$ Hz, 1 H), 4.61 (d, $J = 12.8$ Hz, 1 H), 4.27 (dd, $J = 12.8, 6.4$ Hz, 1 H), 4.10 (s, 1 H), 3.25-2.90 (m, 5), 3.03 (s, 3 H), 2.96 (s, 3 H), 2.95 (s, 3 H), 2.42-2.33 (m, 1 H), 2.29-2.20 (m, 1 H), 1.72-1.61 (m, 1 H), 1.10 (d, $J = 6$ Hz, 9 H); MS (ESI) m/z 546.30 ($\text{M}+\text{H}$).

10 Compound **120**



Prepared similarly to **114** by reductive alkylation of **113** with $t\text{-Bu-N}(\text{Cbz})\text{-CH}_2\text{CHO}$ followed by hydrogenation: ^1H NMR (400 MHz, CD_3OD) δ 7.59 (d, $J = 8.6$ Hz, 1 H), 4.38 (s, 2 H), 4.09 (s, 1 H), 3.60-2.95 (m, 7 H), 3.03 (s, 3 H), 2.96 (s, 3 H),
 15 H), 2.41-2.30 (m, 1 H), 2.28-2.20 (m, 1 H), 1.72-1.60 (m, 1 H), 1.44 (s, 9 H); MS (ESI) m/z 561.31 ($\text{M}+\text{H}$).


Compound **121**



20 $2\text{-}t\text{-Butylaminoacetylchloride}$ hydrochloride (5.8 mg, 0.031 mmol, 1.2 equiv) was added to a DMF solution (0.2 mL) of **113** (12 mg, 0.026 mmol) at 25°C . The reaction was stirred at 25°C for 30 min. The reaction mixture was diluted with 0.05 N HCl (2 mL) and injected into a Waters Autopurification system equipped with a Phenomenex Polymerx 10 μ RP-1 100A column [$10\ \mu\text{m}$, 150×21.20 mm; flow rate,


20 mL/min; Solvent A: 0.05 N HCl; Solvent B: CH₃CN; gradient: 0→100% B over 15 min; mass-directed fraction collection]. Fractions with the desired MW were collected and freeze-dried to yield 3.0 mg of pure **121**: ¹H NMR (400 MHz, CD₃OD) δ 7.34 (d, *J* = 9.6 Hz, 1 H), 4.46 (s, 2 H), 4.08 (s, 1 H), 3.81 (s, 2 H), 3.18-2.92 (m, 3 H), 3.03 (s, 3 H), 2.96 (s, 3 H), 2.32-2.18 (m, 2 H), 1.69-1.60 (m, 1 H), 1.38 (s, 9 H); MS (ESI) *m/z* 575.30 (M+H).

Compound **122**



10 Prepared similarly to compound **121**: ¹H NMR (400 MHz, CD₃OD) δ 7.33 (d, *J* = 9.9 Hz, 1 H), 4.46 (s, 2 H), 4.08 (s, 1 H), 4.00 (s, 2 H), 3.23-2.91 (m, 3), 3.04 (s, 3 H), 2.97 (s, 3 H), 2.95 (s, 6 H), 2.32-2.18 (m, 2 H), 1.70-1.58 (m, 1 H); MS (ESI) *m/z* 547.23 (M+H).


15 Compound **123**



20 Prepared similarly to **121** using n-propyl isocyanate: ¹H NMR (400 MHz, CD₃OD) δ 7.24 (d, *J* = 9.8 Hz, 1 H), 4.31 (s, 2 H), 4.08 (s, 1 H), 3.18-2.93 (m, 3 H), 3.10 (t, *J* = 6.7 Hz, 2 H), 3.03 (s, 3 H), 2.96 (s, 3 H), 2.32-2.18 (m, 2 H), 1.69-1.58 (m, 1 H), 1.55-1.46 (m, 2 H), 0.92 (t, *J* = 6.7 Hz, 3 H); MS (ESI) *m/z* 584.01 (M+H).

Example 9. Synthesis of Compounds of Structural Formula (A), wherein Y is -(CH₂)₃-N(R^A)(R^B).

Scheme 9




In Scheme 9, R and R' are R^A and R^B respectively, as defined in Structural Formula (A).

The following compounds were prepared according to Scheme 9.

5


Compound 9-1



Br₂ (2.7 mL, 52.0 mmol, 1.2 equiv) was added to a solution of **3** (10.6 g, 43.3 mmol) in acetic acid (100 mL) at 25 °C. The reaction was stirred at 25 °C for 12 h. The resulting mixture was added dropwise to ice-water (400 mL). The mixture


was allowed to warm to 25 °C over 1 h. The resulting suspension was filtered through a pad of Celite. The solid was washed off with EtOAc. The combined organic layer was dried (Na₂SO₄) and concentrated to give 14 g of crude **9-1**.

5 Compound **9-2**




Potassium carbonate (8.9 g, 64.5 mmol, 1.5 equiv) and benzyl bromide (11.5 mL, 96.8 mmol, 2.25 equiv) were added to an acetone solution (100 mL) of crude **9-1** (14 g, 43 mmol) at 25 °C. The reaction was stirred at 25 °C for 12 h and concentrated. The resulting mixture was partitioned between H₂O and EtOAc. The aqueous layer was extracted with EtOAc. The combined EtOAc extracts were dried (Na₂SO₄) and concentrated to yield crude **9-2**. Flash chromatography on silica gel (100:1 to 30:1 hexanes/EtOAc) yielded 15.4 g of compound **9-2** (87% for 2 steps).

15 Compound **9-3**




Pd(OAc)₂ (227 mg, 1.0 mmol, 0.2 equiv) and P(O-Tol)₃ (462 mg, 1.5 mmol, 0.3 equiv) were added to a DMF solution (10 mL) of **9-2** (2.1 g, 5.06 mmol). The reaction was purged with N₂ for 5 min. Et₃N (3.5 mL, 25.3 mmol, 5 equiv) and allyloxy-t-butyldimethylsilane (2.2 mL, 10.1 mmol, 2 equiv) were added to the reaction. The reaction was heated to 88 °C and stirred at 88 °C for 5 h. The reaction was allowed to cool to 25 °C and quenched with H₂O. The resulting mixture was extracted with EtOAc. The combined EtOAc extracts were dried (Na₂SO₄) and concentrated to give crude **9-3**. Flash chromatography on silica gel (100:0 to 100:1 hexanes/EtOAc) yielded 1.2 g of compound **9-3** (47%).

Compound **9-4**



n-BuLi (1.3 mL, 2.07 mmol, 5.5 equiv) was added to a THF solution (5 mL) of diisopropylamine (0.3 mL, 2.07 mmol, 5.5 equiv) at 0 °C. The reaction was stirred at 0 °C for 30 min and cooled to -78 °C. TMEDA (0.8 mL, 5.64 mmol, 15 equiv) was added to the mixture. To the resulting solution was added a THF solution (5 mL) of **9-3** (475 mg, 0.94 mmol, 2.5 equiv). The reaction was stirred at -78 °C for 10 min. A THF solution (5 mL) of enone (181 mg, 0.376 mmol) was added to the reaction at -78 °C. The reaction was stirred at -78 °C for 30 min and allowed to warm to 25 °C over 1 h, quenched with saturated NH₄Cl, and extracted with EtOAc. The combined EtOAc extracts were dried (Na₂SO₄) and concentrated to yield the crude product. Preparative reverse phase HPLC purification on a Waters Autopurification system using a Sunfire Prep C18 OBD column [5 μm, 19 × 50 mm; flow rate, 20 mL/min; Solvent A: H₂O with 0.1% HCO₂H; Solvent B: CH₃CN with 0.1% HCO₂H; injection volume: 4.0 mL (CH₃CN); gradient: 100→100% B over 15 min; mass-directed fraction collection]. Fractions with the desired MW were collected and concentrated on a RotaVac at 25 °C to remove most of the acetonitrile. The resulting mostly aqueous solution was extracted with EtOAc. The combined EtOAc extracts were dried (Na₂SO₄) and concentrated to give 200 mg of **9-4** (59%).


20 Compound **9-5**



TFA (0.5 mL) was added to a THF/H₂O (2 mL/0.5 mL) solution of **9-4** at 25 °C. The reaction was stirred at 25 °C for 1 h. The reaction was quenched with sat. NaHCO₃ solution. The reaction was extracted with EtOAc. The combined EtOAc extracts were dried (Na₂SO₄) and concentrated to give crude **9-5**. Preparative reverse phase HPLC purification on a Waters Autopurification system using a Sunfire Prep C18 OBD column [5 μm, 19 × 50 mm; flow rate, 20 mL/min; Solvent A: H₂O with

0.1% HCO₂H; Solvent B: CH₃CN with 0.1% HCO₂H; injection volume: 4.0 mL (CH₃CN); gradient: 80→100% B over 15 min; mass-directed fraction collection]. Fractions with the desired MW were collected and concentrated on a RotaVap at 25 °C to remove most of the acetonitrile. The resulting mostly aqueous solution was
5 extracted with EtOAc. The combined EtOAc extracts were dried (Na₂SO₄) and concentrated to give 80 mg of **9-5** (46%)


Compound **9-6**



10 Dess-Martin periodinane (18 mg, 0.043 mmol, 1.2 equiv) was added to a CH₂Cl₂ solution (1 mL) of **9-5** (28 mg, 0.036 mmol) at 25 °C. The reaction was stirred at 25 °C for 30 min and diluted with H₂O. The resulting mixture was extracted with CH₂Cl₂. The combined CH₂Cl₂ extracts were dried (Na₂SO₄) and concentrated to give crude **9-6**.

15

Compound **9-7-1**




Pyrrolidine (15 μL, 0.18 mmol, 5 equiv) was added to a dichloroethane solution (1 mL) of crude **9-6** (0.036 mmol) at 25 °C. The reaction was stirred at
20 25 °C for 10 min. HOAc (15 μL) and NaBH(OAc)₃ (15 mg, 0.072 mmol, 2 equiv) were added to the reaction. The reaction mixture was stirred at 25 °C for 1 h and quenched with H₂O. The resulting mixture was extracted with CH₂Cl₂. The combined CH₂Cl₂ extracts were dried (Na₂SO₄) and concentrated to give crude **9-7-1**. Preparative reverse phase HPLC purification on a Waters Autopurification system
25 using a Sunfire Prep C18 OBD column [5 μm, 19 × 50 mm; flow rate, 20 mL/min; Solvent A: H₂O with 0.1% HCO₂H; Solvent B: CH₃CN with 0.1% HCO₂H; injection volume: 4.0 mL (CH₃CN); gradient: 0→100% B over 15 min; mass-

directed fraction collection]. Fractions with the desired MW were collected and concentrated on a RotaVap at 25 °C to remove most of the acetonitrile. The resulting mostly aqueous solution was extracted with EtOAc. The combined EtOAc extracts were dried (Na₂SO₄) and concentrated to give 6 mg of **9-7-1** (20% for 2 steps).

5

Compound **124**




Aqueous HF (0.3 mL, 48%) was added to a CH₃CN solution (2 mL) of **9-7-1** (6 mg, 0.007 mmol) in a polypropylene tube at 25 °C. The reaction was stirred at 25 °C for 18 h. The resulting mixture was poured into an aqueous solution of K₂HPO₄ (2 g, dissolved in 15 mL water). The mixture was extracted with EtOAc. The combined EtOAc extracts were dried (Na₂SO₄) and concentrated to yield crude desilyl product.

Palladium on carbon (2 mg, 10 wt%) was added to a HCl/MeOH solution (0.5N, 2 mL) of the crude desilyl product. The reaction was purged with hydrogen and stirred under H₂ (balloon) at 25 °C for 4 h. The reaction mixture was filtered through a small Celite plug. The filtrate was concentrated to yield the crude product. Preparative reverse phase HPLC purification on a Waters Autopurification system using a Phenomenex Polymerx 10 μ RP-1 100A column [10 μm, 150 × 21.20 mm; flow rate, 20 mL/min; Solvent A: 0.05 N HCl/water, Solvent B: CH₃CN; injection volume: 4.0 mL (0.05 N HCl/water); gradient: 0→50% B over 7 min, 50→100% over 3 min, and 100% over 5 min; mass-directed fraction collection]. Fractions with the desired MW, eluting at 6.4-8.2 min, were collected and freeze-dried to yield 1.5 mg of compound **124**: ¹H NMR (400 MHz, CD₃OD) δ 7.28 (d, *J* = 9.7 Hz, 1 H), 4.08 (s, 1 H), 3.71-3.63 (m, 2 H), 3.32-2.95 (m, 7 H), 3.04 (s, 3 H), 2.96 (s, 3 H), 2.81-2.73 (m, 2 H), 2.32-1.98 (m, 8 H), 1.70-1.59 (m, 1 H); MS (ESI) *m/z* 544.18 (M+H).


Compounds **125** – **127** were prepared similarly to compound **124** using the corresponding amines in the reductive amination step.

Compound **125**




^1H NMR (400 MHz, CD_3OD) δ 7.28 (d, $J = 9.7$ Hz, 1 H), 4.08 (s, 1 H), 3.25-2.94 (m, 5 H), 3.04 (s, 3 H), 2.96 (s, 3 H), 2.89 (s, 6 H), 2.80-2.70 (m, 2 H), 2.32-2.18 (m, 2 H), 2.10-2.00 (m, 2 H), 1.70-1.58 (m, 1 H); MS (ESI) m/z 518.26 ($\text{M}+\text{H}$).

Compound **126**




^1H NMR (400 MHz, CD_3OD) δ 7.27 (d, $J = 9.6$ Hz, 1 H), 4.08 (s, 1 H), 3.20-2.93 (m, 5 H), 3.04 (s, 3 H), 2.96 (s, 3 H), 2.82-2.72 (m, 2 H), 2.33-2.19 (m, 2 H), 2.04-1.94 (m, 2 H), 1.70-1.58 (m, 2 H), 1.37 (s, 9 H); MS (ESI) m/z 546.20 ($\text{M}+\text{H}$).

Compound **127**



^1H NMR (400 MHz, CD_3OD) δ 7.28 (d, $J = 9.7$ Hz, 1 H), 4.09 (s, 1 H), 4.04 (q, $J = 9.0$ Hz, 2 H), 3.25-2.95 (m, 5 H), 3.04 (s, 3 H), 2.97 (s, 3 H), 2.84-2.75 (m, 2 H), 2.32-2.20 (m, 2 H), 2.13-2.03 (m, 2 H), 1.70-1.58 (m, 1 H); MS (ESI) m/z 572.22 ($\text{M}+\text{H}$).

Compound **128**




Compound **128** was prepared from compound **9-5** by HF treatment following by hydrogenation under similar conditions: ^1H NMR (400 MHz, CD_3OD) δ 7.21 (d, $J = 9.87$ Hz, 1 H), 4.07 (s, 1 H), 3.63-3.57 (m, 2 H), 3.20-2.90 (m, 5 H), 3.04 (s, 3 H), 2.96 (s, 3 H), 2.75-2.68 (m, 2 H), 2.32-2.17 (m, 2 H), 1.89-1.79 (m, 2 H), 1.70-1.57 (m, 1 H), 1.25 (d, $J = 7.2$ Hz, 1 H); MS (ESI) m/z 491.18 (M+H).

Example 10. Synthesis of Compound 129


Scheme 10

10



The following compounds were prepared according to Scheme 10.


15 Compound 10-1



$i\text{PrMgCl}\cdot\text{LiCl}$ (0.68 mL, 1.2 M, 0.82 mmol, 2 equiv) was added to a THF solution (5 mL) of **9-2** (170 mg, 0.41 mmol) at 0 °C. The reaction was stirred at 0 °C for 30 min. MeI (0.2 mL, 1.64 mmol, 4 equiv) was added to the reaction mixture.


The reaction was stirred at 0 °C for 30 min and allowed to warm to 25 °C over 1 h. The reaction was quenched with NH₄Cl solution and extracted with EtOAc. The combined EtOAc extracts were dried (Na₂SO₄) and concentrated to give crude **10-1**. Flash chromatography on silica gel (30:1 hexanes/EtOAc) yielded 31 mg of compound **10-1** (22%).

Compound **10-2**




A THF solution (1 mL) of **10-1** (31 mg, 0.088 mmol, 1.7 equiv) was added to a THF solution (1 mL) of LDA (0.13 mL, 1.3 M, 0.176 mmol, 3.3 equiv) and TMEDA (39 μ L, 0.26 mmol, 4.9 equiv). The reaction was stirred at -78 °C for 10 min. A THF solution (1 mL) of enone (26 mg, 0.053 mmol) was added to the reaction at -78 °C. The reaction was stirred at -78 °C for 30 min and allowed to warm to 25 °C over 1 h, quenched with saturated NH₄Cl solution, and extracted with EtOAc. The combined EtOAc extracts were dried (Na₂SO₄) and concentrated to yield the crude **10-2**. Preparative reverse phase HPLC purification on a Waters Autopurification system using a Sunfire Prep C18 OBD column [5 μ m, 19 $\times$ 50 mm; flow rate, 20 mL/min; Solvent A: H₂O with 0.1% HCO₂H; Solvent B: CH₃CN with 0.1% HCO₂H; injection volume: 4.0 mL (CH₃CN); gradient: 80 $\rightarrow$ 100% B over 15 min; mass-directed fraction collection]. Fractions with the desired MW were collected and concentrated on a RotaVac at 25 °C to remove most of the acetonitrile. The resulting mostly aqueous solution was extracted with EtOAc. The combined EtOAc extracts were dried (Na₂SO₄) and concentrated to give 10 mg of **10-2** (26%).

Compound **10-3**



Aqueous HF (0.3 mL, 48%) was added to a CH₃CN solution (2 mL) of **10-2** (6 mg, 0.008 mmol) in a polypropylene tube at 25 °C. The reaction was stirred at 25 °C for 18 h. The resulting mixture was poured into an aqueous solution of K₂HPO₄ (2 g, dissolved in 15 mL water). The mixture was extracted with EtOAc.
5 The combined EtOAc extracts were dried (Na₂SO₄) and concentrated to yield crude **10-3**.


Compound **129**



10 Palladium on carbon (2 mg, 10 wt%) was added to a HCl/MeOH solution (0.5N, 2 mL) of the crude **10-3**. The reaction was purged with hydrogen and stirred under H₂ (balloon) at 25 °C for 4 h. The reaction mixture was filtered through a small Celite plug. The filtrate was concentrated to yield the crude product. Preparative reverse phase HPLC purification on a Waters Autopurification system
15 using a Phenomenex Polymerx 10 μ RP-1 100A column [10 μm, 150 × 21.20 mm, flow rate, 20 mL/min; Solvent A: 0.05 N HCl/water; Solvent B: CH₃CN; injection volume: 4.0 mL (0.05 N HCl/water); gradient: 0→70% B over 7 min, 70→100% over 3 min, and 100% over 5 min; mass-directed fraction collection]. Fractions with the desired MW were collected and freeze-dried to yield 1.5 mg of compound **129**:
20 ¹H NMR (400 MHz, CD₃OD) δ 7.19 (d, *J* = 9.7 Hz, 1 H), 4.07 (s, 1 H), 3.20-2.93 (m, 3 H), 3.03 (s, 3 H), 2.96 (s, 3 H), 2.31-2.17 (m, 2 H), 2.22 (s, 3 H), 1.69-1.58 (m, 1 H); MS (ESI) *m/z* 447.23 (M+H).

Example 11. Synthesis of Compounds 130-132

25 *Scheme 11*




In Scheme 11, R⁶, R⁹, and R^{9'} is X, R^B, and R^A, respectively, as defined in Structural Formula (A).

5


Compounds 130 – 132 were prepared according to Scheme 11.

Compound **130**




10

Compound **131**



15


Compound **132**



Example 12. Synthesis of Compounds 133-135

Scheme 12

5




In Scheme 12, R⁹ and R^{9'} is R^B, and R^A, respectively, as defined in Structural Formula (A).

10


Compounds 133 – 135 were prepared according to Scheme 12.

Compound **133**




15

Compound **134**



Compound 135



5

Example 13

The antibacterial activities for the compounds of the invention were studied according to the following protocols.

10 Minimum Inhibitory Concentration Assay

Frozen bacterial strains were thawed and subcultured onto Mueller Hinton Broth (MHB) or other appropriate media (*Streptococcus* requires blood and *Haemophilus* requires hemin and NAD). Following incubation overnight, the strains were subcultured onto Mueller Hinton Agar and again incubated overnight.

- 15 Colonies were observed for appropriate colony morphology and lack of contamination. Isolated colonies were selected to prepare a starting inoculum equivalent to a 0.5 McFarland standard. The starting inoculum was diluted 1:125 using MHB for further use. Test compounds were prepared by dilution in sterile water to a final concentration of 5.128mg/mL. Antibiotics (stored frozen, thawed
- 20 and used within 3 hours of thawing) and compounds were further diluted to the desired working concentrations.

- The assays were run as follows. Fifty μ L of MHB was added to wells 2 – 12 of a 96-well plate. One hundred μ L of appropriately diluted antibiotics was added to well 1. Fifty μ L of antibiotics was removed from well 1 and added to well 2 and the
- 25 contents of well 2 mixed by pipetting up and down five times. Fifty μ L of the mixture in well 2 was removed and added to well 3 and mixed as above. Serial dilutions were continued in the same manner through well 12. Fifty μ L was removed from well 12 so that all contained 50 μ L. Fifty μ L of the working inoculum was then added to all test wells. A growth control well was prepared by adding 50 μ L of

working inoculum and 50 µL of MHB to an empty well. The plates were then incubated at 37 °C overnight, removed from the incubator and each well was read on a plate reading mirror. The lowest concentration (MIC) of test compound that inhibited the growth of the bacteria was recorded.

5 **Example:**

	1	2	3	4	5	6	7	8	9	10	11	12
[Abt]	32	16	8	4	2	1	0.5	0.25	0.125	0.06	0.03	0.015
grow	-	-	-	-	-	+	+	+	+	+	+	+

[abt] = antibiotic concentration in the well

Grow = bacterial growth (cloudiness)

Interpretation: MIC = 2 µg/mL

10

Protocol for Determining Inoculum Concentration (Viable Count)

Ninety µL of sterile 0.9% NaCl was pipetted into wells 2–6 of a 96-well microtiter plate. Fifty 50µl of the inoculum was pipetted into well 1. Ten µL was removed from well 1 and added it to well 2 followed by mixing. Ten µL was removed from well two and mixed with the contents of well 3 and so on creating serial dilutions through well 6. Ten µL was removed from each well and spotted onto an appropriate agar plate. The plate was placed into a CO₂ incubator overnight. The colonies in spots that contain distinct colonies were counted. Viable count was calculated by multiplying the number of colonies by the dilution factor.

20

Spot from Well	1	2	3	4	5	6
Dilution Factor	10 ²	10 ³	10 ⁴	10 ⁵	10 ⁶	10 ⁷

Bacterial Strains

Fifteen bacterial strains, listed below, were examined in minimum inhibitory concentration (MIC) assays.

ID	Organism	Source	Resistance	Comments	Gram Rx
SA100	<i>S. aureus</i>	ATCC 13709	MSSA	Smith strain (animal model)	positive
SA101	<i>S. aureus</i>	ATCC 29213	MSSA	control	positive
SA158	<i>S. aureus</i>	MR, SK75	tet resistant: tetK (efflux)		positive
SA161	<i>S. aureus</i>	Micromyx, LLC	tet resistant: tet(M) ribosomal protection		positive
EF103	<i>E. faecalis</i>	ATCC 29212	tet	control	positive

			intermediate/resistant - mechanism not specified		
EF159	<i>E. faecalis</i>	MR, DS160	tet resistant:tetM (rib protect)	cip-R, ery-I	positive
SP106	<i>S. pneumoniae</i>	ATCC 49619	wt	control	positive
SP160	<i>S. pneumoniae</i>	MR, 54	tet resistant:tet M (rib protect)	pen-R, ery-R	positive
EC107	<i>E. coli</i>	ATCC 25922	wt	control	negative
EC155	<i>E. coli</i>	MR, 10	tet resistant:tetA (efflux)		negative
KP109	<i>K. pneumoniae</i>	ATCC 13883	wt		negative
KP153	<i>K. pneumoniae</i>	MR, 1	tet resistant:tetA (efflux)	cip-R, gen-R	negative
EC108	<i>E. cloacae</i>	ATCC 13047	wt		negative
AB110	<i>A. baumannii</i>	ATCC 19606	wt		negative
PA111	<i>P. aeruginosa</i>	ATCC 27853	wt	control	negative

MSSA = methicillin susceptible *S. aureus*

wt = wild type

ATCC = American Type Culture Collection

MR = Marilyn Roberts, University of Washington

tet = tetracycline

cip = ciprofloxacin

R = resistant

gen = gentamicin

ery = erythromycin

pen = penicillin

Results

Values of minimum inhibition concentration (MIC) for the compounds of the invention are provided in Tables 1a, 1b, 2a, 2b and 3.

5

Compound ID	MICs (ug/mL)							
	SA101	SA100	SA161	SA153	EF103	EF159	SP106	SP160
11	0.125	0.25	0.25	0.0625	0.0625	0.125	≤0.0156	≤0.0156
12	≤0.0156	≤0.0156	0.125	0.5	≤0.0156	0.0625	≤0.0156	≤0.0156
13	≤0.0156	≤0.0156	0.0625	0.25	≤0.0156	0.0625	≤0.0156	≤0.0156
14	0.5	0.25	0.5	0.25	0.125	0.25	0.03	0.125
15	0.5	0.5	0.5	0.25	0.5	0.5	0.016	0.016
16	0.125	0.25	0.25	0.5	0.0625	0.0625	≤0.0156	≤0.0156
17	0.25	0.5	1	2	0.25	1	0.0625	0.0625
18	0.0625	≤0.0156	0.0625	0.125	≤0.0156	0.0313	≤0.0156	≤0.0156

19	0.125	0.25	0.25	0.25	≤0.0156	0.125	≤0.0156	≤0.0156
20	≤0.0156	0.25	0.25	0.5	≤0.0156	0.0625	≤0.0156	≤0.0156
21	0.25	0.25	0.5	4	0.125	1	≤0.0156	0.0625
22	1	1	2	4	2	4	0.5	1
23	≤0.0156	0.5	0.125	0.25	≤0.0156	0.0313	≤0.0156	≤0.0156
24	0.25	0.25	0.125	0.125	≤0.0156	≤0.0156	≤0.0156	≤0.0156
25	0.125	0.125	0.125	0.0313	≤0.0156	0.0625	≤0.0156	≤0.0156
26	1	1	1	1	0.25	0.5	≤0.0156	≤0.0156
27	2	2	4	16	2	4	0.125	0.5
28	1	2	2	1	1	1	0.125	0.0625
29	4	4	4	2	2	2	1	1
30	2	1	4	2	2	2	2	4
31	1	2	2	1	0.5	1	0.25	0.25

Table Ia

Compound ID	MICs (ug/mL)						
	EC107	EC155	AB110	PA111	EC108	KP109	KP153
11	0.25	2	0.5	16	1	1	2
12	0.25	8	1	16	1	1	4
13	0.125	4	0.25	16	1	0.5	2
14	0.5	4	0.25	16	2	1	2
15	1	4	0.125	16	4	2	4
16	0.5	8	1	16	1	1	4
17	2	32	0.5	32	8	4	32
18	0.125	4	0.25	16	0.5	0.5	4
19	0.25	4	0.25	16	2	1	4
20	0.25	8	1	8	1	1	4
21	1	16	2	16	4	2	8
22	16	>32	2	>32	>32	32	>32
23	0.125	8	0.125	8	1	0.5	8
24	0.5	8	0.0313	16	2	1	8
25	0.5	8	0.125	32	2	2	8
26	0.5	4	0.25	16	2	2	4
27	4	32	16	>32	16	8	32
28	2	16	0.5	>32	8	4	16
29	8	8	8	>32	8	8	8
30	>32	>32	8	>32	>32	>32	>32
31	2	8	0.5	>32	8	4	8

Table Ib

Compound ID	MICs (ug/mL)							
	SA101	SA100	SA161	SA158	EF103	EF159	SP106	SP160
32	0.125	0.5	0.25	0.5	0.0625	0.125	≤0.0156	≤0.0156
33	0.25	0.5	1	2	0.25	1	≤0.0156	0.125
34	≤0.0156	0.0625	0.0625	0.125	≤0.0156	≤0.0156	≤0.0156	≤0.0156
35	0.25	0.25	0.5	0.5	0.125	0.25	0.016	0.016
36	0.25	0.5	0.5	1	0.25	0.25	≤0.0156	≤0.0156
37	8	8	>32	>32	16	16	2	4
38	8	8	16	32	8	16	2	8
39	2	2	2	16	2	2	0.25	0.5
40	1	1	1	16	1	1	0.0625	0.5
41	1	1	2	1	1	2	0.125	0.125
42	0.5	1	2	1	0.5	1	0.125	0.0625
43	0.5	0.5	1	0.5	0.25	0.5	≤0.0156	≤0.0156
44	4	4	8	8	8	8	0.5	1
45	0.5	0.5	1	0.5	0.25	0.5	0.125	0.06
46	0.25	0.25	0.5	0.25	0.25	0.5	0.125	0.125

Table 2a

Compound ID	MICs (ug/mL)						
	EC107	EC155	AB110	PA111	EC108	KP109	KP153
32	0.25	8	2	8	1	1	4
33	2	>32	4	>32	16	4	>32
34	0.25	2	0.125	16	1	0.5	2
35	1	16	0.25	>32	8	4	8
36	2	32	0.125	>32	4	4	32
37	>32	>32	>32	>32	>32	>32	>32
38	>32	>32	>32	>32	>32	>32	>32
39	4	>32	32	>32	16	16	>32
40	4	>32	8	32	4	8	32
41	4	16	0.5	>32	16	8	16
42	4	16	0.5	>32	16	8	16
43	1	4	0.125	32	4	2	4
44	32	>32	8	>32	>32	>32	>32
45	1	2	0.06	>32	4	2	4
46	2	4	0.125	>32	32	4	32

Table 2b

Table 3. MIC Values for Compounds of the Invention Compared to Sanycline, Minocycline and Tigecycline. A = lower than or equal to lowest MIC among three control compounds; B = greater than lowest MIC among three control compounds, but lower than highest MIC among three control compounds; C = greater than MIC of all three control compounds. The lowest concentration limit of the assay is 0.016. Thus, reported values in the assay of 0.016 represent an MIC of 0.016 or lower.

Cmpd	SA101 29213	SA100 13709	SA161 MRSA, tetM	SA158 tetK	FF103 29212	FF159 tetM	SP106 49619	SP160 tetM	PC107 25922	FC155 tetA	AR110 19606	PA111 27853	EC108 13047	KP109 13883	KP153 tetA
11	B	B	B	B	B	B	A	A	B	B	C	B	B	B	B
12	A	A	A	B	A	B	A	A	B	B	C	B	B	B	B
13	A	A	A	B	A	B	A	A	B	B	B	B	B	B	B
14	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
15	B	B	B	B	B	B	A	A	B	B	B	B	B	B	B
16	B	B	B	B	B	B	A	A	B	B	C	B	B	B	B
17	B	B	B	B	B	B	B	B	B	B	C	C	B	B	B
18	B	A	A	B	A	A	A	A	B	B	B	B	B	B	B
19	B	B	B	B	A	B	A	A	B	B	B	B	B	B	B
20	A	B	B	B	A	B	A	A	B	B	C	A	B	B	B
21	B	B	B	B	B	B	A	B	B	B	C	B	B	B	B
22	C	B	B	B	B	B	C	B	C	C	C	C	C	C	C
23	A	B	A	B	A	A	A	A	B	B	B	A	B	B	B
24	B	B	A	B	A	A	A	A	B	B	A	B	B	B	B
25	B	B	A	B	A	B	A	A	B	B	B	C	B	B	B
26	C	B	B	B	B	B	A	A	B	B	B	B	B	B	B
27	C	C	B	C	B	B	B	B	B	B	C	C	C	B	B
28	C	C	B	B	B	B	B	B	B	B	C	C	B	B	B
29	C	C	B	B	B	B	C	B	B	B	C	C	B	B	B
30	C	B	B	B	B	B	C	B	C	C	C	C	C	C	C
31	C	C	B	B	B	B	B	B	B	B	C	C	B	B	B
32	B	B	B	B	B	B	A	A	B	B	C	A	B	B	B
33	B	B	B	B	B	B	A	B	B	C	C	C	C	B	C
34	A	B	A	B	A	A	A	A	B	B	B	B	B	B	B
35	B	B	B	B	B	B	A	A	B	B	B	C	B	B	B
36	B	B	B	B	B	B	A	A	B	B	B	C	B	B	B
37	C	C	C	C	C	B	C	B	C	C	C	C	C	C	C

Cmpd	SA101 29213	SA100 13709	SA161 MRSA, tetM	SA158 tetK	EF103 29212	EF159 tetM	SP106 49619	SP160 tetM	EC107 25922	EC155 tetA	AB110 19606	EA111 27853	EC108 13047	KP109 13883	KP153 tetA
38	C	C	C	C	B	B	C	B	C	C	C	C	C	C	C
39	C	C	B	C	B	B	B	B	B	C	C	C	C	C	C
40	C	B	B	C	B	B	B	B	B	C	C	C	B	B	B
41	C	D	B	B	B	B	B	B	B	B	C	C	C	B	B
42	B	B	B	B	B	B	B	B	B	B	C	C	C	B	B
43	B	B	B	B	B	B	A	A	B	B	B	C	B	B	B
44	C	C	D	C	D	B	C	B	C	C	C	C	C	C	C
45	B	B	B	B	B	B	B	B	B	B	A	C	B	B	B
46	B	B	B	B	B	B	B	B	B	B	B	C	C	B	B
47	D	D	B	B	D	B	A	A	D	B	C	A	B	B	B
48	B	B	B	B	A	B	A	A	A	D	A	A	A	A	A
49	B	B	B	B	B	B	A	B	B	B	B	C	B	B	A
50	B	B	B	B	D	B	B	B	B	B	B	C	B	B	B
51	A	A	B	B	A	A	A	A	B	B	B	B	B	B	B
52	B	B	B	B	B	B	A	A	B	B	B	C	B	B	A
53	C	C	C	C	B	C	B	D	C	C	C	C	C	C	C
54	B	B	B	B	B	B	C	B	C	C	C	C	C	C	C
55	B	B	B	B	B	B	A	B	B	C	C	C	B	B	C
56	B	B	B	B	B	B	B	B	B	C	C	C	C	B	C
57	B	B	B	B	B	B	C	B	C	C	C	C	C	C	C
58	C	B	B	B	B	B	C	B	C	C	C	C	C	C	C
59	B	B	B	B	B	B	B	B	B	C	C	C	C	C	C
60	B	B	B	B	B	B	B	B	B	C	C	C	B	B	B
61	C	C	C	C	B	B	B	B	C	C	C	C	C	C	C
62	C	B	D	C	B	B	B	B	C	C	C	C	C	C	C
63	B	B	B	C	B	B	A	B	B	C	C	C	B	D	C
64	B	B	B	B	B	B	A	A	B	B	C	B	B	B	B
65	C	C	B	B	B	B	B	B	B	B	C	C	B	B	B
66	B	B	B	B	B	B	A	A	B	B	B	C	B	B	B
67	C	C	B	B	B	B	B	B	B	B	C	C	B	B	B
68	B	B	B	B	B	B	B	B	B	B	C	B	B	B	B
69	C	C	B	B	B	B	C	B	B	C	C	C	D	C	B
70	C	C	B	B	B	B	B	B	B	C	C	C	B	B	C

Cmpd	SA101 29213	SA100 13709	SA161 MRSA, tetM	SA158 tetK	EF103 29212	EF159 tetM	SP106 49619	SP160 tetM	EC107 25922	EC155 tetA	AB110 19606	PA111 27853	EC108 13047	KPI09 13885	KPI53 tetA
71	B	B	B	B	B	B	B	B	B	C	C	C	C	B	C
72	B	B	B	B	B	B	A	A	B	C	C	C	C	C	C
73	C	B	B	B	B	B	C	B	C	C	C	C	C	C	C
74	C	C	C	C	C	B	C	B	C	C	C	C	C	C	C
75	B	B	B	C	B	B	B	B	B	C	C	C	B	B	C
76	B	B	B	B	B	B	C	C	C	C	C	C	C	C	C
77	B	B	B	B	B	B	C	C	C	C	C	C	C	C	C
78	B	B	B	B	B	B	C	C	C	C	C	C	C	C	C
79	B	B	B	B	B	B	C	C	C	C	C	C	C	C	C
80	C	C	B	C	B	B	C	B	C	C	C	C	C	C	C
81	B	B	A	B	B	B	C	C	C	C	C	C	C	C	C
82	B	B	A	B	B	B	C	B	C	C	C	C	C	C	C
83	B	B	A	B	B	B	C	C	C	C	C	C	C	C	C
84	B	B	B	B	B	B	B	B	C	C	C	C	C	C	C
85	B	B	B	B	B	B	A	B	B	C	C	C	B	B	C
86	B	B	B	B	B	B	B	B	C	C	C	C	C	C	C
87	C	B	B	B	B	B	C	C	C	C	C	C	C	C	C
88	B	B	A	B	B	B	A	B	B	B	C	A	B	B	B
89	A	A	A	A	A	A	A	A	A	B	A	A	A	A	A
90	B	B	A	B	B	B	C	B	C	C	C	C	C	C	C
91	B	B	B	B	B	B	C	B	C	C	C	C	C	C	C
92	B	B	B	B	B	B	C	B	C	C	C	C	C	C	C
93	B	B	B	B	B	B	C	B	C	C	C	C	C	C	C
94	B	B	A	B	B	B	C	B	C	C	C	C	C	C	C
95	B	B	B	B	B	B	C	B	C	C	C	C	C	C	C
96	B	B	B	B	B	B	C	B	C	C	C	C	C	C	C
97	B	B	B	B	B	B	C	C	C	C	C	C	C	C	C
98	B	B	B	B	B	B	C	B	C	C	C	C	C	C	C
99	C	B	B	B	B	B	C	C	C	C	C	C	C	C	C
100	B	B	B	B	B	B	C	C	C	C	C	C	C	C	C
101	B	B	B	B	B	B	B	B	B	B	C	C	B	B	C
102	C	B	B	B	B	B	B	B	B	B	C	C	B	B	B
103	C	C	C	C	B	C	C	B	B	C	C	C	B	C	C

Cmpd	SA101 29213	SA100 13709	SA161 MRSA, tetM	SA158 tetK	EF103 29212	EF159 tetM	SP106 49619	SP160 tetM	EC107 25922	EC155 tetA	AB110 19605	PA111 27853	EC108 13047	KP109 13883	KP153 tetA
104	B	B	B	B	B	B	B	B	B	C	C	C	B	B	C
105	C	B	B	C	B	B	B	B	B	C	C	C	C	C	C
106	C	B	B	B	B	B	C	B	C	C	C	C	C	C	C
107	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT
108	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
109	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
110	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
111	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
112	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
113	C	C	C	C	B	C	B	B	B	C	C	C	B	B	C
114	C	C	B	B	B	B	B	B	B	B	C	C	B	B	B
115	C	C	B	B	B	B	C	B	B	B	C	C	C	B	B
116	C	C	C	B	B	B	B	B	B	C	C	C	B	B	B
117	C	C	C	B	B	B	B	B	B	C	C	C	B	B	C
118	C	C	B	B	B	B	B	B	B	C	C	C	B	B	B
119	C	C	B	B	B	B	C	B	C	B	C	C	C	C	B
120	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
121	C	C	C	C	B	C	C	B	C	C	C	C	C	C	C
122	C	C	C	C	C	C	C	B	C	C	C	C	C	C	C
123	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
124	C	C	C	C	B	C	C	B	C	C	C	C	C	C	C
125	C	C	C	B	B	B	C	B	C	C	C	C	C	C	C
126	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
127	C	B	B	C	B	B	C	B	C	C	C	C	C	C	C
128	C	C	C	C	B	B	C	C	C	C	C	C	C	C	C
129	C	B	B	B	B	B	C	B	C	C	C	C	C	C	B
131	C	C	B	B	C	B	C	C	C	B	C	C	C	C	B
132	C	C	C	B	B	B	C	B	C	B	C	C	C	C	B
133	B	B	B	B	B	B	B	B	B	B	C	C	C	B	B
130	C	B	B	B	B	B	C	B	C	C	C	C	C	C	C
134	B	B	B	B	B	B	A	A	B	B	C	C	B	B	B
135	B	B	B	B	B	B	A	A	B	B	C	C	B	B	B
Mincycline	0.06	0.06	8	0.03	1	16	0.016	2	0.5	8	0.06	16	2	1	8

Compd	SA101 29213	SA100 13709	SA161 MRSA, tetM	SA158 tetK	EF103 29212	EF159 tetM	SP106 49619	SP160 tetM	EC107 25922	EC155 tetA	AB110 19606	PA111 27853	EC108 13047	KP109 13883	KP153 tetA
Sancycline	0.5	1		4	8	8	0.25	8	8	32	0.25	>32	8	8	32
Tigecycline	0.06	0.06	0.125	0.06	0.03	0.06	0.016	0.016	0.03	0.5	0.25	8	0.25	0.125	1

ED₅₀ IV study

Compounds of the present invention were evaluated for efficacy in a mouse systemic infection model against a susceptible *S. aureus* isolate, ATCC 13709.

5 Mice were infected via intraperitoneal injection with a bacterial load that would result in 0% survival within 48 hours after infection. Mice received the test compounds 60 minutes post infection via intravenous injection. Infection control groups did not receive treatment. Survival was assessed over a 48 hour period. Percent survival was calculated and PD₅₀ values determined using Probit analysis.

10

Tet-R sepsis study

The protocol for tetracycline-resistance sepsis study was similar to the protocol for EV₅₀ IV study described above except the infection model used *S. aureus* SA161, a tetracycline-resistant strain.

15

GN sepsis

The protocol for GN sepsis study was similar to the protocol for EV₅₀ IV study described above except the infection model used *E. coli* ATCC 25922.

20 Metabolic stability

A stock solution of the test compounds were prepared by dissolving the compound in DMSO to a final concentration of 1.0 mg/mL.

The analytes and internal standard were diluted and infused into the LC/MS system to determine optimal ionization, polarity and MS/MS fragmentation for
25 selection of specific MRM (multiple reaction monitoring) transitions. Generic chromatographic conditions were developed with a less than 5 minute cycle time.

Pooled human microsome assays were conducted at 0.2 mg/mL protein with an NADPH generating cofactor system (1.3mM NADP+, 3.3mM glucose-6-phosphate, 0.4U/mL glucose-6-phosphate dehydrogenase, and 3.3mM magnesium
30 chloride).

Test compounds were diluted from DMSO stocks to 1 μ M reaction concentrations. Time points were removed at 0 and 60 minutes. After a protein crash

with acetonitrile, samples are analyzed by LC/MS/MS. Peak areas are compared to the time 0 sample, and the percent analyte remaining is calculated. A reaction containing no cofactor was used to control for analyte loss due to non-specific binding, thermal degradation and solubility.

5

SA MIC₉₀

Twenty randomly selected clinical isolates of *S. aureus* were used to determine the minimal inhibitory concentration (MIC) of test compounds for 90% of the isolates (MIC₉₀). MICs were performed by microtiter broth dilution in a 96-well format according to Clinical Laboratory Standards Institute (CLSI) guidelines, as described above.

Viable counts were determined by 10-fold serial dilution. Dilutions were prepared in sterile 0.9% NaCl. Ten microliters of the inoculum and of each of 5 dilutions were plated onto blood or Mueller Hinton agar plates, incubated overnight at 37 °C with 5% CO₂, and counted.

15

TetR MIC₉₀

Ten isolates selected based on resistance to tetracycline were used to determine the MIC₉₀ as described above.

20

EC MIC₉₀

Twenty randomly selected clinical isolates of *E. coli* were used to determine the MIC₉₀ as described above.

25 Protein binding

Test compounds were prepared as 1.0 mg/mL stock solutions in DMSO. The analytes and internal standard were diluted and infused into the LC/MS system to determine optimal ionization, polarity and MS/MS fragmentation for selection of specific MRM (multiple reaction monitoring) transitions. Generic chromatographic conditions were developed with a less than 5 minute cycle time.

30

The DMSO stocks were diluted to 1 and 10 µg/mL in human plasma and incubated in RED devices for 4 hours at 37 °C. The time point was removed at the end of the incubation period. After a protein crash with acetonitrile, the samples were analyzed by LC/MS/MS. Peak areas for the buffer (receiver) and sample
5 (donor) chambers were compared and the protein bound fraction is calculated. Analysis was conducted in duplicate.



Thigh burden

Female CD-1 mice were pre-treated with cyclophosphamide to render the
10 mice neutropenic. Mice were infected with *S. aureus* ATCC 13709 via injection into the right thigh muscle of 0.1 ml per mouse. One and a half hours post infection mice were treated IV with test compounds in doses ranging from 0.3 to 30 mg/kg or 0.3 to 20 mg/kg. Four mice were treated with each drug concentration. Twenty-four
15 hours post treatment, mice were euthanized by CO₂ inhalation. The right thighs of the mice were aseptically removed, weighed, homogenized, serially diluted, and plated on TSA medium. The plates were incubated overnight at 37 °C in 5% CO₂. Colony forming units per gram of thigh was calculated by enumerating the plated colonies then adjusting for serial dilutions and the weight of the thigh.

20 The results for the biological activity studies described above are listed in Table 4.

Table 4.

Chemical Structure	ED50 IV	Meta Stab %	SA MIC 90	TetR MIC90	EC MIC90	Prot Bind%	TetR sepsis	GN seps is	Thigh Burden
	0.3	104%	0.03	2	0.5	90	0.4	6.2	4.41 log10 decrease
	3.55	72%	0.03	4	1	95.6	3.5	14.3	5.01 log10 decrease
	0.07	85%	0.03	4	0.5	87.1			
	0.3	86%	0.03	2	0.5	79.4	1	4.36	5.24 log10 decrease
	0.55	65%	0.03	8	2	87.9			
	0.3	96%	0.06	2	1	96.9			

	0.073	100%	0.06	4	4	93.6	0.62	4.3	4.12 log10 decrease
	0.97	55%	0.06	1	0.5	69.1	0.55	2.1	4.89 log10 decrease

ED₅₀: mg/kg required to protect 50% of mice in septicemia model with *S. aureus* Smith strain.

Meta Stab %: % compound remaining after 60 min, 37 °C incubation with human liver microsomes.

SA MIC90: concentration (ug/mL) required to inhibit 90% of *S. aureus* clinical isolates (n=20).

TetR MIC90: concentration (ug/mL) required to inhibit 90% of tetracycline-resistant *S. aureus* (n=10).

5 EC MIC90: concentration (ug/mL) required to inhibit 90% of *E. coli* clinical isolates (n=20).

Prot Bind: % Protein binding in human plasma with compound at 10uM

TetR sepsis: mg/kg required to protect 50% of mice in septicemia model with tetracycline-resistant *S. aureus* strain.

GtN sepsis: mg/kg required to protect 50% of mice in septicemia model with *E. coli* strain.


Thigh Burden: maximum log10 reduction in bacteria compared to untreated control

10 Maximum log10 reduction: maximum bacterial count (log10) reduction achieved at tested doses.

While this invention has been particularly shown and described with
references to example embodiments thereof, it will be understood by those skilled in
5 the art that various changes in form and details may be made therein without
departing from the scope of the invention encompassed by the appended claims.

CLAIMS

1. A compound represented by the following structural formula:



- or a pharmaceutically acceptable salt thereof.
2. A pharmaceutical composition comprising a compound as defined in Claim 1, or a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable carrier or diluent, and optionally one or more additional therapeutic agents.
3. A compound as defined in Claim 1, or a pharmaceutically acceptable salt thereof, for use in treating or preventing infections caused by bacteria.
4. The use of a compound as defined in Claim 1, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment or prevention of infections caused by bacteria.
5. Compound for use according to Claim 3, or use according to Claim 4, wherein the infection is caused by a Gram-positive bacterium selected from *Staphylococcus spp.*, *Streptococcus spp.*, *Propionibacterium spp.*, *Enterococcus spp.*, *Bacillus spp.*, *Corynebacterium spp.*, *Nocardia spp.*, *Clostridium spp.*, *Actinobacteria spp.*, and *Listeria spp.*.
6. Use or compound for use according to Claim 5, wherein bacteria is *Bacillus anthracis*.

7. Compound for use according to Claim 3, or use according to Claim 4, wherein the infection is caused by a Gram-negative bacterium selected from *Enterobacteriaceae*, *Bacteroidaceae*, *Vibrionaceae*, *Pasteurellae*, *Pseudomonadaceae*, *Neisseriaceae*, *Rickettsiae*, *Moraxellaceae*, any species of *Proteae*, *Acinetobacter spp.*, *Helicobacter spp.*, and *Campylobacter spp.*
8. Use or compound for use according to Claim 7, wherein the bacteria is *E. coli*, *Klebsiella pneumonia*, *Yersinia pestis* or *Francisella tularensis*.
9. Compound for use according to Claim 3, or use according to Claim 4, wherein the infection is caused by *K. pneumoniae*, *Salmonella*, *E. hirae*, *A. baumannii*, *M. catarrhalis*, *H. influenzae*, *P. aeruginosa*, *E. faecium*, *E. coli*, *S. aureus* or *E. faecalis*.
10. Compound for use according to Claim 3, or use according to Claim 4, wherein the infection is caused by rickettsiae, chlamydiae, *Legionella spp.* or *Mycoplasma spp.*
11. Compound for use according to Claim 3, or use according to Claim 4, wherein the infection is a skin infection, a respiratory tract infection, a hospital acquired infection, an intra-abdominal infection or a urinary tract infection.
12. Compound for use according to Claim 3, or use according to Claim 4, wherein the compound as defined in Claim 1, or a pharmaceutically acceptable salt thereof, is administered alone.
13. Compound for use according to Claim 3, or use according to Claim 4, wherein the compound as defined in Claim 1, or a pharmaceutically acceptable salt thereof, is administered in combination with one or more additional therapeutic agents.

14. A process for making a composition as defined in Claim 2, which process comprises mixing the compound as defined in Claim 1 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or diluent.