Claims

1. Use of *N,N-bis*-2-mercaptoethyl isophthalamide (NBMI), or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the prevention of paracetamol toxicity.

2. Use of *N,N-bis*-2-mercaptoethyl isophthalamide (NBMI), or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the prevention of acute liver failure associated with paracetamol toxicity.

3. A pharmaceutical formulation comprising:

(a) *N,N-bis*-2-mercaptoethyl isophthalamide (NBMI), or a pharmaceutically-acceptable salt thereof; and

(b) paracetamol,

and optionally one or more pharmaceutically-acceptable excipients.

4. A kit-of-parts comprising components:

(A) a pharmaceutical formulation comprising *N,N-bis*-2-mercaptoethyl isophthalamide (NBMI), or a pharmaceutically-acceptable salt thereof, and optionally one or more pharmaceutically-acceptable excipients; and

(B) a pharmaceutical formulation comprising paracetamol, and optionally one or more pharmaceutically-acceptable excipients,

which components (A) and (B) are each provided in a form that is suitable for administration in conjunction with the other for prevention of paracetamol toxicity and/or prevention of acute liver failure associated with paracetamol toxicity.