PCT WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : C12Q 1/02, C12N 5/00, C07D 207/00, 209/00, 295/00

A1

(11) International Publication Number: WO 00/34507

(43) International Publication Date: 15 June 2000 (15.06.00)

(21) International Application Number: PCT/US99/28769

(22) International Filing Date: 7 December 1999 (07.12.99)

(30) Priority Data:
60/111,195 7 December 1998 (07.12.98) US

(71) Applicant *(for all designated States except US)*: DUKE UNIVERSITY [US/US]; 230 North Building, P.O. Box 90083, Durham, NC 27708–0083 (US).

(72) Inventors; and
(75) Inventors/Applicants *(for US only)*: SMITH, Clayton, A. [US/US]; Duke University, Hematology & Medical Oncology, Dept. of Medicine, 421 Medical Science Research Building, Durham, NC 27710 (US). COLVIN, Michael [US/US]; 208 Arcadia Lane, Chapel Hill, NC 27514 (US). STORMS, Robert, W. [US/US]; 30 Dansey Circle, Durham, NC 27713 (US). LUDEMAN, Susan, M. [US/US]; 7313 208 Calibre Park Drive, Durham, NC 27707 (US).

(74) Agent: WILSON, Mary, J.; Nixon & Vanderhye P.C., Suite 800, 1100 North Glebe Road, Arlington, VA 22201–4714 (US).

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published
With international search report.

(54) Title: A METHOD OF ISOLATING STEM CELLS

(57) Abstract

The present invention relates, in general, to stem cells, and in particular, to a method of isolating stem cells and to reagents suitable for use in such a method. The invention further relates to stem cell populations isolatable in accordance with the present method.