




FIG. 1




FIG. 2




FIG. 3A




FIG. 3B




FIG. 4




FIG. 5A




FIG. 5B




FIG. 6




FIG. 7




FIG. 8





FIG. 9

**FIG. 10**




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 18




FIG. 19




FIG. 20




FIG. 21




FIG. 22




FIG. 23




FIG. 24

Mean Cumulative Key Secondary End Points – Overall




FIG. 25




FIG. 26

Mean Cumulative Key Secondary End Points – Secondary Prevention




FIG. 27




FIG. 28

Mean Cumulative Key Secondary End Points – Primary Prevention




FIG. 29




FIG. 30




FIG. 31




FIG. 32






FIG. 33

**FIG. 34A**

**FIG. 34B**




FIG. 35




FIG. 36




FIG. 37A




FIG. 37B




FIG. 38




FIG. 39




FIG. 40




FIG. 41




FIG. 42




FIG. 43




FIG. 44




FIG. 45




FIG. 46




FIG. 47





FIG. 48



Prespecified exploratory analysis with no adjustment for multiple comparisons. Repeated-measurements analysis of change from baseline blood pressure over time by mixed-effects model. ITT population. Icosapent ethyl n=4089, Placebo n=4091. Maximum number of observations per patient = 6.

FIG. 49




FIG. 50