


TRIAZOLOPYRIDINE COMPOUNDS AS PDE10A INHIBITORS

FIELD OF THE INVENTION

The present invention relates to novel compounds of formula (I)



5

wherein

R¹ is aryl, heteroaryl or NR⁴R⁵, wherein said aryl and said heteroaryl can be substituted by 1 to 3 substituents independently selected from the group consisting of hydroxyl, halogen, C₁-C₇ alkyl, C₁-C₇ alkoxy and C₁-C₇ haloalkyl;

10

R² is hydrogen, halogen or C₁-C₇ alkyl;

R³ is aryl or heteroaryl, wherein said aryl and said heteroaryl can be substituted by 1 to 3 substituents independently selected from the group consisting of C₁-C₇ alkyl, hydroxyl, halogen, -C(O)-NR⁶R⁷ and -C(O)-O-R⁸;

15

R⁴ and R⁵ are independently C₁-C₇ alkyl, C₃-C₈ cycloalkyl, together with the nitrogen atom to which they are attached, form a heterocycloalkyl which can be substituted by 1 to 3 substituents independently selected from the group consisting of hydroxyl, halogen, C₁-C₇ alkyl, C₁-C₇ alkoxy and C₁-C₇ haloalkyl;

20

R⁶ and R⁷, are independently C₁-C₇ alkyl, C₃-C₈ cycloalkyl, together with the nitrogen atom to which they are attached, form a heterocycloalkyl which can be substituted by 1 to 3 substituents independently selected from the group consisting of hydroxyl, halogen, C₁-C₇ alkyl, C₁-C₇ alkoxy and C₁-C₇ haloalkyl;

R⁸ is hydrogen, C₁-C₇ alkyl, cycloalkyl;

or a pharmaceutically acceptable salt thereof.

Further, the invention is concerned with a process for the manufacture of the above compounds, pharmaceutical preparations which contain such compounds as well as the use of these compounds for the production of pharmaceutical preparations.

5 SUMMARY OF THE INVENTION

Schizophrenia is a progressive and devastating neurological disease characterized by episodic positive symptoms such as delusions, hallucinations, thought disorders and psychosis and persistent negative symptoms such as flattened affect, impaired attention and social withdrawal, and cognitive impairments (Lewis DA and Lieberman JA, *Neuron*, , 28:325-33, 10 2000). For decades research has focused on the "dopaminergic hyperactivity" hypothesis which has led to therapeutic interventions involving blockade of the dopaminergic system (Vandenberg RJ and Aubrey KR., *Exp. Opin. Ther. Targets*, 5(4): 507-518, 2001; Nakazato A and Okuyama S, et al., *Exp. Opin. Ther. Patents*, 10(1): 75-98, 2000). This pharmacological approach, besides ameliorating positive symptoms in schizophrenic patients, poorly addresses negative and 15 cognitive symptoms which are the best predictors of functional outcome (Sharma T., *Br.J. Psychiatry*, 174(suppl. 28): 44-51, 1999). In addition, current antipsychotic treatment is associated with adverse effects like weight gain, extrapyramidal symptoms or effects on glucose and lipid metabolism, related to their unspecific pharmacology.

In conclusion there is still a need for developing new antipsychotics with improved 20 efficacy and safety profile. A complementary model of schizophrenia was proposed in the mid-1960' based upon the psychotomimetic action caused by the blockade of the glutamate system by compounds like phencyclidine (PCP) and related agents (ketamine) which are non-competitive NMDA receptor antagonists. Interestingly, in healthy volunteers PCP-induced psychotomimetic action incorporates positive and negative symptoms as well as cognitive 25 dysfunction, thus closely resembling schizophrenia in patients (Javitt DC et al., *Biol. Psychiatry*, 45: 668-679, 1999).

Cyclic nucleotides cyclic adenosine monophosphate (cAMP) and cyclic guanosine monophosphate (cGMP) are ubiquitous second messengers responsible for mediating the biological response of a variety of extracellular signals, including neurotransmitters, light and 30 hormones. cAMP and cGMP regulate a variety of intracellular processes particularly in neurons of the central nervous system by activating cAMP- and cGMP-dependent kinases which then phosphorylate proteins involved in the regulation of synaptic transmission, neuronal differentiation and survival.

A crucial mechanism for controlling intracellular cyclic nucleotide levels and therefore 35 cyclic nucleotide signaling is via hydrolysis of the 3',5'-phosphodiester bond by

phosphodiesterases. Phosphodiesterases (PDEs) are a family of widely expressed enzymes encoded by 21 different genes in humans, with each gene encoding several splice variants (Beavo, J., *Physiol. Rev.* 1995, 75, 725–748; Conti, M., Jin, S.L., *Prog. Nucleic Acid Res. Mol. Biol.* 1999, 63, 1–38; Soderling, S.H., Beavo, J.A., *Curr. Opin. Cell Biol.* 2000, 12, 174–179, 5 Manallack, D.T. et al. *J. Med.Chem.* 2005, 48 (10), 3449-3462).

The PDE families differ in their substrate specificity for the cyclic nucleotides, their mechanism of regulation and their sensitivity to inhibitors. Moreover, they are differentially localized in the organism, among the cells of an organ and even within the cells. These differences lead to a differentiated involvement of the PDE families in the various physiological 10 functions.

PDE10A is a dual substrate PDE encoded by a single gene as reported in 1999 by three separate research groups (Fujishige K., et al., *Eur J Biochem* (1999) 266(3):1118-1127, Soderling S.H., et al., *ProcNatl Acad Sci USA* (1999) 96(12):7071-7076, Loughney K., et al., *Gene* (1999) 234(1):109-117). PDE10A is unique from other members of the multigene family 15 with respect to amino acid sequence (779 aa), tissue-specific pattern of expression, affinity for cAMP and cGMP and the effect on PDE activity by specific and general inhibitors.

PDE10A has one of the most restricted distribution of any PDE family being primarily expressed in the brain particularly in the nucleus accumbens and the caudate putamen. Additionally thalamus, olfactory bulb, hippocampus and frontal cortex show moderate levels of 20 PDE10A expression. All these brain areas have been suggested to be involved in the pathophysiology of schizophrenia and psychosis, suggesting a central role of PDE10A in this devastating mental illness. Outside the central nervous system PDE10A transcript expression is also observed in peripheral tissues like thyroid gland, pituitary gland, insulin secreting pancreatic cells and testes (Fujishige, K. et al., *J. Biol. Chem.* 1999, 274, 18438–18445, Sweet, L. (2005) 25 WO 2005/012485). On the other hand expression of PDE10A protein has been observed only in enteric ganglia, in testis and epididymal sperm (Coskran T.M, et al., *J. Histochem. Cytochem.* 2006, 54 (11), 1205-1213).

In the striatum both mRNA and protein are expressed only in the GABA (- aminobutyric acid)-containing medium spiny projection neurons making it an intriguing target 30 for the treatment of diseases of the central nervous system (Fujishige, K. et al., *Eur. J. Biochem.* 1999, 266, 1118-1127; Seeger, T.F. et al., *Brain Res.* 2003, 985, 113-126). The striatal medium spiny neurons are the principal input site and first site for information integration in the basal ganglia circuit of the mammalian brain. The basal ganglia are a series of interconnected subcortical nuclei that integrate widespread cortical input with dopaminergic signaling to plan

and execute relevant motor and cognitive patterns while suppressing unwanted or irrelevant patterns (Graybiel, A.M. *Curr. Biol.* 2000, 10, R509–R511 (2000)).

Papaverine, a relatively specific PDE10A inhibitor, and PDE10A-knockout mice have been used to explore the physiology of this enzyme and the possible therapeutic utility of PDE10A inhibition. Inhibition of this enzyme pharmacologically or through gene disruption causes a reduction in activity and a reduced response to psychomotor stimulants. Inhibition also reduces the conditioned avoidance response, a behavioural response that is predictive of clinical antipsychotic activity (Siuciak, J.A.; et al., *Neuropharmacology* 2006, 51 (2), 386-396; Siuciak, J.A.; et al., *Neuropharmacology* 2006, 51 (2), 374-385).

In addition PDE10A inhibition bears the potential to improve the negative and cognitive symptoms associated to schizophrenia. Indeed papaverine have been shown to attenuate the deficits in the extra-dimensional shift learning induced in rats by sub-chronic treatment with PCP, an animal paradigm of NMDA receptor hypofunction (Rodefer, J.S., et al., *Eur. J. Neuroscience* 2005, 2, 1070-1076). In addition increased social interaction in PDE10A2-deficient mice have been observed (Sano, H. *J. Neurochem.* 2008, 105, 546-556).

Diseases that can be treated with PDE10A inhibitors include, but are not limited to, diseases thought to be mediated in part by dysfunction of the basal ganglia, of other parts of the central nervous system and of other PDE10A expressing tissues. In particular, diseases can be treated, where inhibition of PDE10A can have therapeutic effects.

These diseases include, but are not limited to, certain psychotic disorders such as schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia, delusional disorder or substance-induced psychotic disorder, anxiety disorders such as panic disorder, obsessive-compulsive disorder, acute stress disorder or generalized anxiety disorder, obsessive/compulsive disorders, drug addictions, movement disorders such as Parkinson's disease or restless leg syndrome, cognition deficiency disorders such as Alzheimer's disease or multi-infarct dementia, mood disorders such as depression or bipolar disorders, or neuropsychiatric conditions such as psychosis, attention-deficit/hyperactivity disorder (ADHD) or related attentional disorders.

The compounds of the present invention are also suitable for the treatment of diabetes and related disorders such as obesity by regulating the cAMP signaling system.

PDE10A inhibitors might also be useful in preventing neurons from undergoing apoptosis by raising cAMP and cGMP levels and, thus, might possess anti-inflammatory properties. Neurodegenerative disorders treatable with PDE10A inhibitors include, but are not

limited to, as Alzheimer's disease, Huntington's disease, Parkinson's disease, multiple sclerosis, stroke or spinal cord injury.

The growth of cancer cells is inhibited by cAMP and cGMP. Thus by raising cAMP and cGMP, PDE10A inhibitors can also be used for the treatment of different solid tumors and
5 hematological malignancies such as renal cell carcinoma or breast cancer.

DETAILED DESCRIPTION OF THE INVENTION

Unless otherwise indicated, the following definitions are set forth to illustrate and define the meaning and scope of the various terms used to describe the invention herein.

It must be noted that, as used in the specification and the claims, the singular forms "a",
10 "an" and "the" include plural referents unless the context clearly dictates otherwise.

When indicating the number of substituents, the term "one or more" means from one substituent to the highest possible number of substitution, i.e. replacement of one hydrogen up to replacement of all hydrogens by substituents.

The term "optional" or "optionally" denotes that a subsequently described event or
15 circumstance may but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not.

The term "halogen" refers to fluorine, chlorine, bromine and iodine, more specifically fluorine, chlorine and bromine.

The term "alkyl" denotes a monovalent linear or branched saturated hydrocarbon group
20 of 1 to 12 carbon atoms. In particular embodiments, alkyl has 1 to 7 carbon atoms, and in more particular embodiments 1 to 4 carbon atoms. Examples of alkyl include methyl, ethyl, propyl, isopropyl, n-butyl, iso-butyl, sec-butyl, or tert-butyl.

The term "aromatic" denotes the conventional idea of aromaticity as defined in the literature, in particular in IUPAC - Compendium of Chemical Terminology, 2nd, A. D.
25 McNaught & A. Wilkinson (Eds). Blackwell Scientific Publications, Oxford (1997).

The term "cycloalkyl" denotes a monovalent saturated monocyclic or bicyclic hydrocarbon group of 3 to 10 ring carbon atoms. In particular embodiments cycloalkyl denotes a monovalent saturated monocyclic hydrocarbon group of 3 to 8 ring carbon atoms. Bicyclic means consisting of two saturated carbocycles having one or more carbon atoms in common.
30 Particular cycloalkyl groups are monocyclic. Examples for monocyclic cycloalkyl are

cyclopropyl, cyclobutanyl, cyclopentyl, cyclohexyl or cycloheptyl. Examples for bicyclic cycloalkyl are bicyclo[2.2.1]heptanyl, or bicyclo[2.2.2]octanyl.

The term "alkoxy" denotes a group of the formula -O-R', wherein R' is an alkyl group. Examples of alkoxy moieties include methoxy, ethoxy, isopropoxy, and tert-butoxy.

5 The term "haloalkyl" denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by same or different halogen atoms, particularly fluoro atoms. Examples of haloalkyl include monofluoro-, difluoro- or trifluoro-methyl, -ethyl or -propyl, for example 3,3,3-trifluoropropyl, 2-fluoroethyl, 2,2,2-trifluoroethyl, fluoromethyl, or trifluoromethyl. The term "perhaloalkyl" denotes an alkyl group where all hydrogen atoms of the
10 alkyl group have been replaced by the same or different halogen atoms.

The term "aryl" denotes a monovalent aromatic carbocyclic mono- or bicyclic ring system comprising 6 to 10 carbon ring atoms. Examples of aryl moieties include phenyl and naphthyl

15 The term "amino" refers to a monovalent group that has a nitrogen atom with two hydrogen atoms (represented by -NH₂).

The term "oxo" when referring to substituents on heterocycloalkyl means that an oxygen atom is attached to the heterocycloalkyl ring. Thereby, the "oxo" may either replace two hydrogen atoms on a carbon atom, or it may simply be attached to sulfur, so that the sulfur exists in oxidized form, i.e. bearing one or two oxygens.

20 The term "heterocycloalkyl" denotes a monovalent saturated or partly unsaturated mono- or bicyclic ring system of 3 to 9 ring atoms, comprising 1, 2, or 3 ring heteroatoms selected from N, O and S, the remaining ring atoms being carbon. In particular embodiments, heterocycloalkyl is a monovalent saturated monocyclic ring system of 4 to 7 ring atoms, comprising 1, 2, or 3 ring heteroatoms selected from N, O and S, the remaining ring atoms being carbon. Examples for
25 monocyclic saturated heterocycloalkyl are aziridinyl, oxiranyl, azetidiny, oxetanyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydro-thienyl, pyrazolidinyl, imidazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, morpholinyl, thiomorpholinyl, 1,1-dioxo-thiomorpholin-4-yl, azepanyl, diazepanyl, homopiperazinyl, or oxazepanyl. Examples for bicyclic saturated heterocycloalkyl are 8-aza-bicyclo[3.2.1]octyl,
30 quinuclidinyl, 8-oxa-3-aza-bicyclo[3.2.1]octyl, 9-aza-bicyclo[3.3.1]nonyl, 3-oxa-9-aza-bicyclo[3.3.1]nonyl, or 3-thia-9-aza-bicyclo[3.3.1]nonyl. Examples for partly unsaturated heterocycloalkyl are dihydrofuryl, imidazolinyl, dihydro-oxazolyl, tetrahydro-pyridinyl, or dihydropyranyl.


The term "heteroaryl" denotes a monovalent aromatic heterocyclic mono- or bicyclic ring system of 5 to 12 ring atoms, comprising 1, 2, 3 or 4 heteroatoms selected from N, O and S, the remaining ring atoms being carbon. Examples of heteroaryl moieties include pyrrolyl, furanyl, thienyl, imidazolyl, oxazolyl, thiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridinyl, pyrazinyl, pyrazolyl, pyridazinyl, pyrimidinyl, triazinyl, azepinyl, diazepinyl, isoxazolyl, benzofuranyl, isothiazolyl, benzothienyl, indolyl, isoindolyl, isobenzofuranyl, benzimidazolyl, benzoxazolyl, benzoisoxazolyl, benzothiazolyl, benzoisothiazolyl, benzooxadiazolyl, benzothiadiazolyl, benzotriazolyl, purinyl, quinoliny, isoquinoliny, quinazolinyl, or quinoxalinyl.

Compounds of formula (I) can form pharmaceutically acceptable salts. Examples of such pharmaceutically acceptable salts are salts of compounds of formula (I) with physiologically compatible mineral acids, such as hydrochloric acid, sulphuric acid, sulphurous acid or phosphoric acid; or with organic acids, such as methanesulphonic acid, p-toluenesulphonic acid, acetic acid, lactic acid, trifluoroacetic acid, citric acid, fumaric acid, maleic acid, tartaric acid, succinic acid or salicylic acid. The term "pharmaceutically acceptable salts" refers to such salts. Compounds of formula (I) which comprise an acidic group, such as e.g. a COOH group, can further form salts with bases. Examples of such salts are alkaline, earth-alkaline and ammonium salts such as e.g. Na-, K-, Ca- and trimethylammonium salt. The term "pharmaceutically acceptable salts" also refers to such salts. Particular salts are those obtained by the addition of an acid.

The term "pharmaceutically acceptable esters" embraces derivatives of the compounds of formula (I), in which a carboxy group has been converted to an ester. Lower alkyl, lower hydroxyalkyl, lower-alkoxy-lower-alkyl, amino-lower-alkyl, mono- or di-lower-alkyl-amino-lower-alkyl, morpholino-lower-alkyl, pyrrolidino-lower-alkyl, piperidino-lower-alkyl, piperazino-lower-alkyl, lower-alkyl-piperazino-lower-alkyl and aralkyl esters are examples of suitable esters. Particular esters are methyl, ethyl, propyl, butyl and benzyl esters. The term "pharmaceutically acceptable esters" furthermore embraces compounds of formula (I) in which hydroxy groups have been converted to the corresponding esters with inorganic or organic acids such as, nitric acid, sulphuric acid, phosphoric acid, citric acid, formic acid, maleic acid, acetic acid, succinic acid, tartaric acid, methanesulphonic acid, p-toluenesulphonic acid and the like, which are non toxic to living organisms.

It will be appreciated that the compounds of general formula (I) in this invention may be derivatised at functional groups to provide derivatives which are capable of conversion back to the parent compound in vivo.

The present invention relates to novel compounds of formula (I)



wherein

5 R^1 is aryl, heteroaryl or NR^4R^5 , wherein said aryl and said heteroaryl can be substituted by 1 to 3 substituents independently selected from the group consisting of hydroxyl, halogen, C_1 - C_7 alkyl, C_1 - C_7 alkoxy and C_1 - C_7 haloalkyl;

R^2 is hydrogen, halogen or C_1 - C_7 alkyl;

10 R^3 is aryl or heteroaryl, wherein said aryl and said heteroaryl can be substituted by 1 to 3 substituents independently selected from the group consisting of C_1 - C_7 alkyl, hydroxyl, halogen, $-C(O)-NR^6R^7$ and $-C(O)-O-R^8$;

15 R^4 and R^5 are independently C_1 - C_7 alkyl, C_3 - C_8 cycloalkyl, together with the nitrogen atom to which they are attached, form a heterocycloalkyl which can be substituted by 1 to 3 substituents independently selected from the group consisting of hydroxyl, halogen, C_1 - C_7 alkyl, C_1 - C_7 alkoxy and C_1 - C_7 haloalkyl;

R^6 and R^7 , are independently C_1 - C_7 alkyl, cycloalkyl, together with the nitrogen atom to which they are attached, form a heterocycloalkyl which can be substituted by 1 to 3 substituents independently selected from the group consisting of hydroxyl, halogen, C_1 - C_7 alkyl, C_1 - C_7 alkoxy and C_1 - C_7 haloalkyl;

20 R^8 is hydrogen, C_1 - C_7 alkyl, cycloalkyl;

or a pharmaceutically acceptable salt thereof.


In a particular embodiment the invention relates to compounds of formula (I), wherein R^1 is phenyl, pyridinyl, thiazolyl or NR^4R^5 , wherein R^4 and R^5 are independently selected from C_1 - C_3 alkyl, R^4 and R^5 together with the nitrogen atom to which they are attached, form a pyrrolidinyl or morpholinyl ring.

25

In a further particular embodiment the invention relates to compounds of formula (I), wherein R^2 is hydrogen.

In yet another embodiment the invention relates to compounds of formula (I), wherein R^3 is :

5



wherein:

A_1 is NR^9 ,

A_2 is $NR^{9'}$,

10 R^6 and R^7 are independently selected from C_1 - C_3 alkyl, C_3 - C_8 cycloalkyl, both optionally substituted by 1 to 3 substituents independently selected from halogen, C_1 - C_3 alkyl, C_1 - C_3 alkoxy and C_1 - C_3 haloalkyl;

R^6 and R^7 together with the nitrogen atom to which they are attached form a heterocycloalkyl of 4 to 7 ring atoms, comprising 1 or 2 ring heteroatoms selected from N and O,
15 the heterocycloalkyl optionally substituted by 1 to 3 substituents independently selected from halogen, C_1 - C_3 alkyl, C_1 - C_3 alkoxy and C_1 - C_3 haloalkyl,

R^9 and $R^{9'}$ are independently selected from hydrogen, C_1 - C_3 alkyl, C_1 - C_3 alkoxy and C_1 - C_3 haloalkyl.

Another particular embodiment relates to compounds of formula (I), wherein:

20 R^6 and R^7 are independently selected from C_1 - C_3 alkyl, C_3 - C_8 cycloalkyl,

R^6 and R^7 together with the nitrogen atom to which they are attached form azetidiny, pyrrolidinyl and morpholinyl which are optionally substituted by 1 to 3 substituents independently selected from the group consisting of halogen, C_1 - C_3 alkyl, C_1 - C_3 alkoxy and C_1 - C_3 haloalkyl.

25 R^9 is selected from C_1 - C_3 alkyl, C_1 - C_3 alkoxy and C_1 - C_3 haloalkyl, preferably C_1 - C_3 alkyl,

R^{9'} is selected from hydrogen and C₁-C₃ alkyl, preferably hydrogen.

Particular compounds of formula (I) are selected from the group consisting of:

2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid (2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

5 1-Methyl-5-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid

4-(Azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

10 4-(Azetidine-1-carbonyl)-1-methyl-N-(2-(thiazol-4-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-1H-pyrazole-5-carboxamide

1-Methyl-4-(morpholine-4-carbonyl)-N-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-1H-pyrazole-5-carboxamide

4-(3-Fluoro-azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

15 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(ethyl-methyl-amide) 3-[(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

20 2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

4-(Azetidine-1-carbonyl)-1-methyl-N-(2-(pyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-1H-pyrazole-5-carboxamide

N4-Cyclopropyl-N4,1-dimethyl-N5-(2-(pyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-1H-pyrazole-4,5-dicarboxamide

25 1-Methyl-N-(2-(pyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-4-(pyrrolidine-1-carbonyl)-1H-pyrazole-5-carboxamide

2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid (2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-4-(azetidine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

5 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(ethyl-methyl-amide) 3-[(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

4-(3-Fluoro-azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

10 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

4-(Azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

15 2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid (2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 3-[(2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide] 4-(ethyl-methyl-amide)

20 4-(3-Fluoro-azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

6-cyclopropyl-3-methoxy-N-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)pyrazine-2-carboxamide

25 2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-4-(pyrrolidine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-diethylamino-[1,2,4]triazolo [1,5-a]pyridin-6-yl)-amide

2-Methyl-4-(azetidine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-diethylamino-[1,2,4]triazolo [1,5-a]pyridin-6-yl)-amide

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 3-[(2-diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide] 4-(ethyl-methyl-amide)

5 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-diethylamide 3-[(2-diethylamino-1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

10 2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

4-(Azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-pyrrolidin-1-yl-1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid (2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

15 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(ethyl-methyl-amide) 3-[(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

20 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-diethylamide 3-[(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-diethylamide 3-{[2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide}

2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid [2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide

25 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(ethyl-methyl-amide) 3-{[2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide}

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-{[2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide}

2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid [2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide

4-(Azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid [2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide.

5 Yet particular compounds of formula (I) are those selected from the group consisting of:

4-(Azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

1-Methyl-4-(morpholine-4-carbonyl)-N-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-1H-pyrazole-5-carboxamide

10 4-(3-Fluoro-azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

15 2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid (2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]


4-(Azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

20 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

The invention further relates to a process for the manufacture of compounds of formula (I) as defined above, which process comprises:


a) reacting a compound of formula (2)

25



with

b) a compound of formula (3)



5 wherein R^1 , R^2 , R^3 are as defined above and, if desired, converting the compounds into pharmaceutically acceptable salts thereof.


The reaction described above can be carried out under conditions as described in the description and examples or under conditions well known to the person skilled in the art.

10 The compounds of formula (2) and (3) can be prepared by methods known in the art or as described below or in analogy thereto.

The present invention also relates to compounds of formula (I) as defined above, when prepared by a process as described above.

15 Compounds of formula 1 can be prepared from building blocks 2 and 3 according to Scheme 1. The conversion, commonly known as amide coupling, can be achieved in several ways. In one method, the acid 2 is activated with a coupling reagent, such as 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate (TBTU), or propylphosphonic anhydride, and converted by addition of amine 3 to the desired product, 1. In another method, the acid 2 is activated by transformation into an acid chloride, e.g. by reaction with thionyl chloride. The acid chloride is then converted by addition of the amine 3 to the desired product, 1. A base, 20 e.g. diisopropylethylamine (DIPEA), is usually added to bind liberated HCl.


Scheme 1



25 Compounds of formula 3 can be prepared according to Scheme 2: A suitably substituted 2-aminopyridine 4 is reacted with a suitably substituted nitrile 5, 1,10-phenanthroline monohydrate and copper (I) bromide to form compound 6. Compound 6 is then reacted with a

compound **7** such as a tert-butyl carbamate to give compound **8**. Compound **8** is then converted to amine **3** by methods well known in the art. Compounds **4**, **5** and **7** are either commercially available, or can be prepared by methods well known in the art.

Scheme 2




Step 1: 1,10-phenantroline monohydrate, copper (I) bromide

Step 2: Cesium carbonate, tris(dibenzylideneacetone)dipalladium(0), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene, dioxane


Step 3: Hydrochloric acid (5M) in Diethylether

10 Compounds of formula 2, with R³ being a pyrazolyl carboxylic acid derivative, can be prepared according to Scheme 3: Compound 9 is reacted with a hydrazine 10, or a salt thereof, to give a pyrazole 11 (similar to the method of A. Hanzlowsky, B. Jelencic, S. Recnik, J. Svete, A. Golobic, B. Stanovnik J. Heterocyclic Chem. 2003, 40(3), 487-498). Selective mono-saponification of the diester 11 yields, depending on the reaction conditions, compound 2a or its
15 isomer, compound 2b.

Scheme 3




Compounds of formula 1, with R^3 being a C_1 - C_7 alkoxy-carbonyl-substituted hetero-aromatic ring, can be further transformed according to Scheme 4. For instance, compounds of the general formula 1-COOEt can be saponified by suitable methods, e.g. by reaction with LiOH, to give 1-COOH. Upon activation with a suitable reagent such as TBTU, 1-COOH can be converted with a primary or secondary amine to 1-CONR₂. Alternatively, 1-COOEt can be directly converted into 1-CONR₂, e.g. by reaction with an amine such as methylamine.

Scheme 4

Alternatively, compounds of formula 3, with R^1 being NR^4R^5 , can be prepared according to Scheme 5: A suitably substituted 2-aminopyridine 4 is reacted with ethoxycarbonyl-isothiocyanate 17 to form the thiourea 18, which can be cyclized using hydroxylamine and a suitable base as diisopropylethylamine to a compound of formula 19. This can be converted by methods well known in the art to compound 22 and further to the desired amine of formula 3c. For example, compounds of formula 19 can be treated with a nitrite such as sodium nitrite or alkyl nitrite such as tert.butyl nitrite and a bromide such as copper(II)bromide or benzyl-triethylammonium bromide in a suitable solvent such as acetonitrile or bromoform to form bromide 20. Bromide 20 can then be reacted with an amine of formula 21 in a suitable solvent such as THF or ethanol. A base, e.g. diisopropylethylamine (DIPEA), may be added to the reaction.

2-Amino-5-bromopyrimidine, ethoxycarbonyl-isothiocyanate and hydroxylamine are commercially available; amines 21 are either commercially available, or can be prepared by methods well known in the art.

Scheme 5



All reactions are typically performed in a suitable solvent and under an atmosphere of argon or nitrogen.

5 The corresponding salts with acids can be obtained by standard methods known to the person skilled in the art, e.g. by dissolving the compound of formula (I) in a suitable solvent such as e.g. dioxane or THF and adding an appropriate amount of the corresponding acid. The products can usually be isolated by filtration or by chromatography. The conversion of a compound of formula (I) into a pharmaceutically acceptable salt with a base can be carried out
 10 by treatment of such a compound with such a base. One possible method to form such a salt is e.g. by addition of 1/n equivalents of a basic salt such as e.g. $\text{M}(\text{OH})_n$, wherein M is metal or ammonium cation and n is number of hydroxide anions, to a solution of the compound in a suitable solvent (e.g. ethanol, ethanol-water mixture, tetrahydrofuran-water mixture) and to remove the solvent by evaporation or lyophilisation.

15 The conversion of compounds of formula (I) into pharmaceutically acceptable esters can be carried out e.g. by treatment of a suitable carboxy group present in the molecule with a suitable alcohol using e.g. a condensating reagent such as benzotriazol-1-yl-oxytris(dimethylamino)phosphonium hexafluorophosphate (BOP), N,N-dicyclohexylcarbodiimide (DCC), N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide
 20 hydrochloride (EDCI) or O-(1,2-dihydro-2-oxo-1-pyridyl)-N,N,N,N-tetra-methyluronium-tetrafluoroborate (TPTU), or by direct reaction with a suitable alcohol under acidic conditions, as for example in the presence of a strong mineral acid like hydrochloric acid, sulfuric acid and the like. Compounds having a hydroxyl group can be converted to esters with suitable acids by analogous methods.

25 Insofar as their preparation is not described in the examples, the compounds of formula (I) as well as all intermediate products can be prepared according to analogous methods or according to the methods set forth above. Starting materials are commercially available, known in the art or can be prepared by methods known in the art or in analogy thereto.

As described above, the novel compounds of the present invention have been found to inhibit PDE10A activity. The compounds of the present invention can therefore be used, either alone or in combination with other drugs, for the treatment and/or prophylaxis of diseases which are modulated by PDE10A inhibitors. These diseases include, but are not limited to, certain

5 psychotic disorders such as schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia, delusional disorder or substance-induced psychotic disorder, anxiety disorders such as panic disorder, obsessive/compulsive disorders, acute stress disorder or generalized anxiety disorder, drug addictions, movement disorders such as Parkinson's disease or restless leg syndrome, cognition deficiency disorders such as Alzheimer's disease or multi-

10 infarct dementia, mood disorders such as depression or bipolar disorders, or neuropsychiatric conditions such as psychosis, attention-deficit/hyperactivity disorder (ADHD) or related attentional disorders. Other disorders are diabetes and related disorders, such as type 2 diabetes mellitus, neurodegenerative disorders such as Alzheimer's disease, Huntington's disease, Parkinson's disease, multiple sclerosis, stroke or spinal cord injury, solid tumors and

15 hematological malignancies such as renal cell carcinoma or breast cancer.

The invention therefore also relates to compounds as described above for use as therapeutically active substance.

The invention also relates to pharmaceutical compositions comprising a compound as described above and a therapeutically inert carrier.

20 In another embodiment, the invention relates to the use of a compound as described above for the treatment or prophylaxis of psychotic disorders, schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia, delusional disorder, substance-induced psychotic disorder, anxiety disorders, panic disorder, obsessive/compulsive disorders, acute stress disorder, generalized anxiety disorder, drug addictions, movement disorders,

25 Parkinson's disease, restless leg syndrome, cognition deficiency disorders, Alzheimer's disease, multi-infarct dementia, mood disorders, depression, bipolar disorders, neuropsychiatric conditions, psychosis, attention-deficit/hyperactivity disorder, attentional disorders, diabetes and related disorders, type 2 diabetes mellitus, neurodegenerative disorders, Huntington's disease, multiple sclerosis, stroke, spinal cord injury, solid tumors, hematological malignancies, renal cell

30 carcinoma or breast cancer.

Yet in another embodiment, the invention relates to the use of a compound as described above for the preparation of a medicament for the treatment or prophylaxis of psychotic disorders, schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia, delusional disorder, substance-induced psychotic disorder, anxiety disorders,

35 panic disorder, obsessive/compulsive disorders, acute stress disorder, generalized anxiety

disorder, drug addictions, movement disorders, Parkinson's disease, restless leg syndrome, cognition deficiency disorders, Alzheimer's disease, multi-infarct dementia, mood disorders, depression, bipolar disorders, neuropsychiatric conditions, psychosis, attention-deficit/hyperactivity disorder, attentional disorders, diabetes and related disorders, type 2
5 diabetes mellitus, neurodegenerative disorders, Huntington's disease, multiple sclerosis, stroke, spinal cord injury, solid tumors, hematological malignancies, renal cell carcinoma or breast cancer.

The invention also relates to a compound as described above for the treatment or prophylaxis of psychotic disorders, schizophrenia, positive, negative and/or cognitive symptoms
10 associated with schizophrenia, delusional disorder, substance-induced psychotic disorder, anxiety disorders, panic disorder, obsessive/compulsive disorders, acute stress disorder, generalized anxiety disorder, drug addictions, movement disorders, Parkinson's disease, restless leg syndrome, cognition deficiency disorders, Alzheimer's disease, multi-infarct dementia, mood disorders, depression, bipolar disorders, neuropsychiatric conditions, psychosis, attention-
15 deficit/hyperactivity disorder, attentional disorders, diabetes and related disorders, type 2 diabetes mellitus, neurodegenerative disorders, Huntington's disease, multiple sclerosis, stroke, spinal cord injury, solid tumors, hematological malignancies, renal cell carcinoma or breast cancer.

The invention further relates to a compound as described above, when manufactured
20 according to a process as described above.

In another embodiment, the invention relates to a method for the treatment or prophylaxis of psychotic disorders, schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia, delusional disorder, substance-induced psychotic disorder, anxiety disorders, panic disorder, obsessive/compulsive disorders, acute stress disorder, generalized anxiety
25 disorder, drug addictions, movement disorders, Parkinson's disease, restless leg syndrome, cognition deficiency disorders, Alzheimer's disease, multi-infarct dementia, mood disorders, depression, bipolar disorders, neuropsychiatric conditions, psychosis, attention-deficit/hyperactivity disorder, attentional disorders, diabetes and related disorders, type 2 diabetes mellitus, neurodegenerative disorders, Huntington's disease, multiple sclerosis, stroke,
30 spinal cord injury, solid tumors, hematological malignancies, renal cell carcinoma or breast cancer, which method comprises administering an effective amount of a compound as described above.

Prevention and/or treatment of schizophrenia is a particular indication. Yet particular indication is prevention and/or treatment of positive, negative and/or cognitive symptoms
35 associated with schizophrenia.

The invention further relates to a pharmaceutical composition comprising compounds of formula (I) as defined above and a therapeutically inert carrier.

As described above, the novel compounds of the present invention have been found to inhibit PDE10A activity. The compounds of the present invention can therefore be used, either alone or in combination with other drugs, for the treatment or prophylaxis of psychotic disorders, schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia, delusional disorder, substance-induced psychotic disorder, anxiety disorders, panic disorder, obsessive/compulsive disorders, acute stress disorder, generalized anxiety disorder, drug addictions, movement disorders, Parkinson's disease, restless leg syndrome, cognition deficiency disorders, Alzheimer's disease, multi-infarct dementia, mood disorders, depression, bipolar disorders, neuropsychiatric conditions, psychosis, attention-deficit/hyperactivity disorder, attentional disorders, diabetes and related disorders, type 2 diabetes mellitus, neurodegenerative disorders, Huntington's disease, multiple sclerosis, stroke, spinal cord injury, solid tumors, hematological malignancies, renal cell carcinoma or breast cancer.

Yet in another embodiment, the invention relates to the use of a compound of the present invention for the preparation of a medicament for the treatment or prophylaxis of psychotic disorders, schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia, delusional disorder, substance-induced psychotic disorder, anxiety disorders, panic disorder, obsessive/compulsive disorders, acute stress disorder, generalized anxiety disorder, drug addictions, movement disorders, Parkinson's disease, restless leg syndrome, cognition deficiency disorders, Alzheimer's disease, multi-infarct dementia, mood disorders, depression, bipolar disorders, neuropsychiatric conditions, psychosis, attention-deficit/hyperactivity disorder, attentional disorders, diabetes and related disorders, type 2 diabetes mellitus, neurodegenerative disorders, Huntington's disease, multiple sclerosis, stroke, spinal cord injury, solid tumors, hematological malignancies, renal cell carcinoma or breast cancer.

The invention also relates to a compound as described above for the treatment or prophylaxis of psychotic disorders, schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia, delusional disorder, substance-induced psychotic disorder, anxiety disorders, panic disorder, obsessive/compulsive disorders, acute stress disorder, generalized anxiety disorder, drug addictions, movement disorders, Parkinson's disease, restless leg syndrome, cognition deficiency disorders, Alzheimer's disease, multi-infarct dementia, mood disorders, depression, bipolar disorders, neuropsychiatric conditions, psychosis, attention-deficit/hyperactivity disorder, attentional disorders, diabetes and related disorders, type 2 diabetes mellitus, neurodegenerative disorders, Huntington's disease, multiple sclerosis, stroke,

spinal cord injury, solid tumors, hematological malignancies, renal cell carcinoma or breast cancer.

The compounds of formula (I) can have one or more asymmetric C atoms and can therefore exist as an enantiomeric mixture, mixture of stereoisomers or as optically pure
5 compounds. The compounds of formula (I) include all diastereomers, tautomers, racemates and mixtures thereof.

Particular compounds of formula (I) are described in the examples as individual compounds as well as pharmaceutically acceptable salts as well as pharmaceutically acceptable esters thereof. Furthermore, the substituents as found in the specific examples described below,
10 individually constitute particular embodiments of the present invention.

PHARMACEUTICAL COMPOSITIONS AND ADMINISTRATION

Another embodiment provides pharmaceutical compositions or medicaments containing the compounds of the invention and a therapeutically inert carrier, diluent or excipient, as well as methods of using the compounds of the invention to prepare such compositions and
15 medicaments. In one example, compounds of formula (I) may be formulated by mixing at ambient temperature at the appropriate pH, and at the desired degree of purity, with physiologically acceptable carriers, i.e., carriers that are non-toxic to recipients at the dosages and concentrations employed into a galenical administration form. The pH of the formulation depends mainly on the particular use and the concentration of compound, but preferably ranges
20 anywhere from about 3 to about 8. In one example, a compound of formula (I) is formulated in an acetate buffer, at pH 5. In another embodiment, the compounds of formula (I) are sterile. The compound may be stored, for example, as a solid or amorphous composition, as a lyophilized formulation or as an aqueous solution.

Compositions are formulated, dosed, and administered in a fashion consistent with good
25 medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the agent, the method of administration, the scheduling of administration, and other factors known to medical practitioners. The “effective amount” of the compound to be administered will be governed by such considerations, and is the
30 minimum amount necessary to inhibit PDE10 and to control the cAMP signaling pathway. For example, such amount may be below the amount that is toxic to normal cells, or the mammal as a whole.

In one example, the pharmaceutically effective amount of the compound of the invention administered parenterally per dose will be in the range of about 0.01-100 mg/kg, alternatively

about 0.1 to 20 mg/kg of patient body weight per day, with the typical initial range of compound used being 0.3 to 15 mg/kg/day. In another embodiment, oral unit dosage forms, such as tablets and capsules, preferably contain from about 25-100 mg of the compound of the invention.

5 The compounds of the invention may be administered by any suitable means, including oral, topical (including buccal and sublingual), rectal, vaginal, transdermal, parenteral, subcutaneous, intraperitoneal, intrapulmonary, intradermal, intrathecal and epidural and intranasal, and, if desired for local treatment, intralesional administration. Parenteral infusions include intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration.

10 The compounds of the present invention may be administered in any convenient administrative form, e.g., tablets, powders, capsules, solutions, dispersions, suspensions, syrups, sprays, suppositories, gels, emulsions, patches, etc. Such compositions may contain components conventional in pharmaceutical preparations, e.g., diluents, carriers, pH modifiers, sweeteners, bulking agents, and further active agents.

15 A typical formulation is prepared by mixing a compound of the present invention and a carrier or excipient. Suitable carriers and excipients are well known to those skilled in the art and are described in detail in, e.g., Ansel, Howard C., et al., *Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems*. Philadelphia: Lippincott, Williams & Wilkins, 2004; Gennaro, Alfonso R., et al. *Remington: The Science and Practice of Pharmacy*. Philadelphia: Lippincott, Williams & Wilkins, 2000; and Rowe, Raymond C. *Handbook of Pharmaceutical*
20 *Excipients*. Chicago, Pharmaceutical Press, 2005. The formulations may also include one or more buffers, stabilizing agents, surfactants, wetting agents, lubricating agents, emulsifiers, suspending agents, preservatives, antioxidants, opaquing agents, glidants, processing aids, colorants, sweeteners, perfuming agents, flavoring agents, diluents and other known additives to provide an elegant presentation of the drug (i.e., a compound of the present invention or
25 pharmaceutical composition thereof) or aid in the manufacturing of the pharmaceutical product (i.e., medicament).

An example of a suitable oral dosage form is a tablet containing about 25mg, 50mg, 100mg, 250mg, or 500mg of the compound of the invention compounded with about 90-30 mg anhydrous lactose, about 5-40 mg sodium croscarmellose, about 5-30mg polyvinylpyrrolidone
30 (PVP) K30, and about 1-10 mg magnesium stearate. The powdered ingredients are first mixed together and then mixed with a solution of the PVP. The resulting composition can be dried, granulated, mixed with the magnesium stearate and compressed to tablet form using conventional equipment. An example of an aerosol formulation can be prepared by dissolving the compound, for example 5-400 mg, of the invention in a suitable buffer solution, e.g. a

phosphate buffer, adding a tonicifier, e.g. a salt such sodium chloride, if desired. The solution may be filtered, e.g., using a 0.2 micron filter, to remove impurities and contaminants.

An embodiment, therefore, includes a pharmaceutical composition comprising a compound of Formula (I), or a stereoisomer or pharmaceutically acceptable salt thereof. In a further embodiment includes a pharmaceutical composition comprising a compound of Formula (I), or a stereoisomer or pharmaceutically acceptable salt thereof, together with a pharmaceutically acceptable carrier or excipient.

The following test was carried out in order to determine the activity of the compounds of the present invention. PDE10 activity of the compounds of the present invention was determined using a Scintillation Proximity Assay (SPA)-based method similar to the one previously described (Fawcett, L. et al., Proc Natl Acad Sci USA (2000) 97(7):3702-3707).

The human PDE10A full length assay was performed in 96-well micro titer plates. The reaction mixture of 50 μ l contained 20 mM HEPES pH=7.5 /10 mM MgCl₂/0.05 mg/ml BSA (Sigma cat. # A-7906), 50 nM cGMP (Sigma, cat. # G6129) and 50 nM [³H]-cGMP (GE Healthcare, cat. # TRK392 S.A. 13.2Ci/mmol), 3.75 ng/well PDE10A enzyme (Enzo Life Science, Lausen, Switzerland cat # SE-534) with or without a specific test compound. A range of concentrations of the potential inhibitor was used to generate data for calculating the concentration of inhibitor resulting in 50% of the effect (e.g. IC₅₀, the concentration of the competitor inhibiting PDE10A activity by 50%). Non-specific activity was tested without the enzyme. The reaction was initiated by the addition of the substrate solution (cGMP and [³H]-cGMP) and allowed to progress for 20 minutes at room temperature. The reaction was terminated by adding 25 μ l of YSi-SPA scintillation beads (GE Healthcare, cat. # RPNQ0150) in 18 mM zinc sulphate solution (stop reagent). After 1 h under shaking, the plate was centrifuged one minute at 170 g to allow beads to settle. Afterwards, radioactive counts were measured on a Perkin Elmer Top-Count Scintillation plate reader.

The compounds according to formula (I) have an IC₅₀ value below 10 μ M, more specifically below 5 μ M, yet more specifically below 1 μ M. The following table shows data for some examples.

Example	PDE10A inhibition IC ₅₀ [nM]	Example	PDE10A inhibition IC ₅₀ [nM]
1	47.3	23	25.0

Example	PDE10A inhibition IC50 [nM]	Example	PDE10A inhibition IC50 [nM]
2	1.2	24	5.3
3	0.6	25	17
4	7.9	26	21.8
5	2.5	27	3.3
6	1.5	28	3.3
7	2.2	29	9.9
8	0.9	30	19.1
9	13.2	31	2.3
10	43.6	32	2.5
11	6.7	33	0.8
12	17.9	34	1.2
13	15.4	35	1.0
14	63.8	36	1.9
15	23.1	37	3.5
16	30.8	38	29.4
17	26.5	39	23.1
18	7.2	40	17.0
19	36.0	41	3.5
20	8.2	42	7.5
21	28.6	43	4.9
22	33.5		


Ames Test was performed as a downscaled version of the standard preincubation Ames test and adapted from the method described by Kado et al. with some modifications e.g. Maron, D.M. and B.N. Ames (1983). Revised methods for the Salmonella mutagenicity test. Mutation Research, 113, 173 – 215. N.Y. Kado, D. Langley and E. Eisenstadt A simple modification of the Salmonella liquid-incubation assay. Increased sensitivity for detecting mutagens in human urine, Mutat Res 121 (1983) 25-32.

The following examples serve to illustrate the present invention in more detail. They are, however, not intended to limit its scope in any manner.


EXAMPLES

Example 1

1-Methyl-5-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid




a) 6-bromo-2-phenyl-[1,2,4]triazolo[1,5-a]pyridine




A mixture of 5-bromopyridin-2-amine (2 g, 11.6 mmol), benzonitrile (26.2 g, 26.1 ml, 254 mmol), 1,10-phenanthroline monohydrate (115 mg, 578 μ mol) and copper (I) bromide (82.9 mg, 578 μ mol) was placed in a 100 ml 2 necked flask. The mixture was heated to 135 °C while air was bubbled gently through the mixture for 2 days. Another portion of copper (I) bromide (82.9 mg, 578 μ mol) and 1,10-phenanthroline monohydrate (115 mg, 578 μ mol) was added and bubbling and heating was continued for further 8 hours. The solvent was evaporated. Purification by flash chromatography on a 20 g silicagel column using heptane/ethyl acetate 10-50 % as eluent and evaporation of the solvents afforded 6-bromo-2-phenyl-[1,2,4]triazolo[1,5-a]pyridine (1.19 g, 38%) as light yellow solid. Mp.: 146°C. MS: m/z=274/276 (M+H⁺).

b) 2-Phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-carbamic acid tert-butylester




A mixture of 6-bromo-2-phenyl-[1,2,4]triazolo[1,5-a]pyridine (1.19 g, 4.34 mmol), tert-butyl carbamate (610 mg, 5.21 mmol) and cesium carbonate (1.98 g, 6.08 mmol) in 1,4-dioxane
5 (30 ml) was purged with argon. Then tris(dibenzylideneacetone)dipalladium(0) (239 mg, 260 μ mol) and 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (301 mg, 521 μ mol) were added and the purging was repeated. The mixture was heated to reflux and stirred for 18 hours under argon atmosphere. The reaction mixture was evaporated to dryness and the solid purified by flash chromatography over a 50 g silicagel column using heptane/ethyl acetate 10-50 % as eluent.
10 Final evaporation afforded 2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-carbamic acid tert-butylester (1.09 g, 81%) as yellow solid. Mp.: 204°C. MS: m/z =311.3 (M+H+).

c) 2-Phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-ylamine



A mixture of 2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-carbamic acid tert-butylester
15 (1.04 g, 3.35 mmol) and hydrochloric acid (5 molar in diethyl ether, 20 ml, 100 mmol) was stirred for 18 hours at 25 °C. The solvent was evaporated, the yellow residue was made basic using saturated aqueous sodium hydrogencarbonate and the mixture was extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate, filtrated and evaporated to dryness affording 2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-ylamine (715 mg, quant.) as
20 yellow solid. Mp.: 201°C. MS: m/z =210.9 (M+H+).


d) 1-Methyl-5-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid ethyl ester



A mixture of 2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-ylamine (300 mg, 1.43 mmol), 4-(ethoxycarbonyl)-1-methyl-1H-pyrazole-5-carboxylic acid (339 mg, 1.71 mmol), propylphosphonic anhydride in ethyl acetate 50% (2.27 g, 2.1 ml, 3.57 mmol) and diisopropylethyl amine (553 mg, 748 μ l, 4.28 mmol) in tetrahydrofuran (10 ml) was refluxed for 2.5 days.

- 5 The mixture was evaporated and the residue was treated with saturated aqueous sodiumhydrogen carbonate and stirred for 30 minutes. The solid was filtered off, washed with water and dried, affording 1-methyl-5-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid ethyl ester (483 mg, 87%) as brown solid. Mp.: 251-254°C. MS: m/z=391.1 (M+H+).


- 10 e) 1-Methyl-5-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid



- 15 A mixture of ethyl 1-methyl-5-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylate (480 mg, 1.23 mmol) and lithium hydroxide hydrate (155 mg, 3.69 mmol) in methanol (10 ml) and water (3.0 ml) was stirred for 18 hours at 25 °C. The solvent was evaporated, the residue was diluted with water and the solution was acidified with hydrochloric acid to pH= 0, the precipitated solid was collected, washed with water and dried in vacuo affording 1-methyl-5-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid (435 mg, 98%) as light brown solid. Mp.: 259°C. MS: m/z=361.2 (M-H+).

20 Example 2

2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid (2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide




-28-

A mixture of 1-methyl-5-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid (112 mg, 309 μ mol), pyrrolidine (220 mg, 256 μ l, 3.09 mmol), propylphosphonic anhydride in ethyl acetate 50% (492 mg, 455 μ l, 773 μ mol) and N,N-diisopropylethylamine (120 mg, 158 μ l, 927 μ mol) in tetrahydrofuran (7.00 ml) was refluxed for 18 hours. The solvent was evaporated, the residue was triturated with saturated aqueous sodium hydrogencarbonate. The precipitated solid was filtered off, washed with water and dried in vacuo, affording 2-methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid (2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide (45 mg, 35%) as brown solid. Mp.: 210°C. MS: m/z=416.1 (M+H+).

Example 3


4-(Azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide



Using azetidine, the title compound was prepared in the same manner as described for example 2. Off-white solid (70 mg, 63.2%). Mp.: 235-236°C. MS: m/z=402.1 (M+H+). Ames test negative.

Example 4


4-(Azetidine-1-carbonyl)-1-methyl-N-(2-(thiazol-4-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-1H-pyrazole-5-carboxamide



Using azetidine and 1-methyl-5-(2-(thiazol-4-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Light brown solid (25 mg, 18%). MS: $m/z=409.2$ ($M+H^+$).

Example 5


- 5 1-Methyl-4-(morpholine-4-carbonyl)-N-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-1H-pyrazole-5-carboxamide



- 10 Using morpholine, the title compound was prepared in the same manner as described for example 2. Brown solid (106 mg, 80%). Mp.: 183-185°C. MS: $m/z=432.5$ ($M+H^+$). Ames test negative.

Example 6


4-(3-Fluoro-azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide



15 Using 3-fluoroazetidine, the title compound was prepared in the same manner as described for example 2. White solid (83 mg, 64%). Mp.: 243°C. MS: $m/z=420.0$ ($M+H^+$).

Example 7


2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(ethyl-methyl-amide) 3-[(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]



- 5 Using ethyl-methyl-amine, the title compound was prepared in the same manner as described for example 2. Light yellow solid (63 mg, 57%). MS: $m/z=404.1$ ($M+H^+$).

Example 8

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]




10


Using cyclopropyl-methyl-amine, the title compound was prepared in the same manner as described for example 2. Light yellow solid (63 mg, 57%). Mp.: 106°C. MS: $m/z=416.1$ ($M+H^+$).

Example 9

- 15 2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide




a) 6-Bromo-2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridine



- 5 A mixture of 5-bromopyridin-2-amine (2 g, 11.6 mmol), nicotinonitrile (24.1 g, 231 mmol), 1,10-phenanthroline monohydrate (115 mg, 578 μ mol) and copper (I) bromide (82.9 mg, 578 μ mol) was placed in a 100 ml 2 necked flask and heated to 135 °C. Heating was continued while air was bubbled gently through the mixture for 1.5 days. The excess nicotinonitrile was distilled off under reduced pressure and the black residue purified by chromatography (70 g sili-
- 10 cagel column using heptane/ethyl acetate 10-100 % as eluent). Final evaporation of the solvents afforded 6-bromo-2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridine (1.66 g, 52%) as yellow solid. Mp.: 176°C. MS: m/z=275/277 (M+H+).


b) (2-Pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-carbamic acid tert-butyl ester



- 15 A mixture of 6-bromo-2-(pyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyridine (1.65 g, 6.00 mmol), tert-butyl carbamate (843 mg, 7.2 mmol) and cesium carbonate (2.74 g, 8.4 mmol) in 1,4-dioxane (37.8 ml) was purged with argon. Then tris(dibenzylideneacetone)dipalladium(0) (330 mg, 360 μ mol) and 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (416 mg, 720 μ mol) were added and the purging was repeated. The mixture was heated to reflux and stirred for 18 hours
- 20 under argon atmosphere. After evaporation to dryness under reduced pressure, the crude material was purified by flash chromatography (50 g silicagel column using heptane / ethyl acetate 20-100 % as eluent). Final evaporation of the solvents afforded (2-pyridin-3-yl-[1,2,4]triazolo[1,5-


a)pyridin-6-yl)-carbamic acid tert-butyl ester (1.35 g, 72.5%) as yellow solid. Mp.: 296°C. MS: m/z=312.3 (M+H+).

c) 2-Pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylamine




5 A mixture of tert-butyl 2-(pyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamate (1.34 g, 4.3 mmol), dichloromethane (10 ml) and hydrochloric acid (5 molar in diethyl ether (36.0 g, 30 ml, 987 mmol) was stirred for 18 hours at 25 °C. The solvents were evaporated under reduced pressure and the residue treated with sodium hydroxide solution (2 molar in water). The mixture was extracted with ethyl acetate, the organic layers were washed with water and brine, dried over
10 magnesium sulfate, filtrated and evaporated to dryness, affording 2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylamine (482 mg, 53%) as light brown solid. Mp.: 230°C. MS: m/z=212.1 (M+H+).

d) 1-Methyl-5-(2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid ethyl ester




15 A mixture of 2-(pyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-ylamine (480 mg, 2.27 mmol), 4-(ethoxycarbonyl)-1-methyl-1H-pyrazole-5-carboxylic acid (495 mg, 2.5 mmol), propylphosphonic anhydride in ethyl acetate 50% (3.62 g, 3.35 ml, 5.68 mmol) and N,N-diisopropylethylamine (1.17 g, 1.55 ml, 9.09 mmol) in tetrahydrofuran (10 ml) was stirred for 18
20 hours at 70 °C. The solvent was evaporated under reduced pressure, the residue was triturated with saturated aqueous sodium hydrogencarbonate. The precipitated solid was filtered off, washed with water and dried in vacuo, affording 1-methyl-5-(2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid ethyl ester (825 mg, 93%) as light brown solid. Mp.: 217°C. MS: m/z=392.1 (M+H+).

25 e) 1-Methyl-5-(2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid



A mixture of ethyl 1-methyl-5-(2-(pyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylate (810 mg, 2.07 mmol) and lithium hydroxide hydrate (434 mg, 10.3 mmol) in methanol (30 ml) and water (5 ml) was stirred for 18 hours at 25 °C. The mixture was adjusted to pH=7 using hydrochloric acid (2molar, 5.15 ml) and then evaporated to dryness under reduced pressure. 1-Methyl-5-(2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid (1.38 g) as light brown solid was used without further purification. MS: m/z=364.1 (M+H⁺).


f) 2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide



Using morpholine and 1-methyl-5-(2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. White solid (53 mg, 32%). Mp.: 229°C. MS: m/z=433.3 (M+H⁺).

Example 10


4-(Azetidine-1-carbonyl)-1-methyl-N-(2-(pyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-1H-pyrazole-5-carboxamide



Using azetidine and 1-methyl-5-(2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. White solid (75 mg, 48%). Mp.: 272°C. MS: m/z=403.2 (M+H+).

5 **Example 11**

N4-Cyclopropyl-N4,1-dimethyl-N5-(2-(pyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-1H-pyrazole-4,5-dicarboxamide




Using cyclopropyl-methyl-amine and 1-methyl-5-(2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. White solid (71 mg, 44%). Mp.: 237°C. MS: m/z=417.3 (M+H+). Ames test negative.

Example 12

1-Methyl-N-(2-(pyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-4-(pyrrolidine-1-carbonyl)-1H-pyrazole-5-carboxamide


-35-




Using pyrrolidine and 1-methyl-5-(2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. White solid (53 mg, 33%). Mp.: 251°C. MS: m/z=417.3 (M+H+).

5 Example 13

2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid (2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide




10 a) 1-Ethoxycarbonyl-3-(5-bromo-pyridin-2-yl)-thiourea




A mixture of 5-bromopyridin-2-amine (14.7 g, 85.0 mmol) and ethoxycarbonyl isothiocyanate (11.1 g, 10.0 ml, 85.0 mmol) in dioxane (300 ml) was stirred for 18 hours at 25 °C under nitrogen atmosphere. The solvent was evaporated under reduced pressure, the solid yellow residue was diluted with ethyl acetate and washed with water and brine, the organic layer was separated, dried over magnesium sulfate, filtrated and evaporated to dryness. 26.2 g (quant.) light yellow solid. Mp.: 159-161°C. MS: m/z=304.0/305.9 (M+H+).

b) 6-Bromo-[1,2,4]triazolo[1,5-a]pyridin-2-ylamine




A mixture of hydroxylamine hydrochloride (40.0 g, 575 mmol) and N-ethyl-diisopropylamine (44.6 g, 60.3 ml, 345 mmol) in ethanol (734 ml) was stirred for a few minutes at 25 °C, then the mixture was added to 1-ethoxycarbonyl-3-(5-bromo-pyridin-2-yl)-thiourea (35 g, 115 mmol) and the resulting mixture was refluxed for 1 day. The solvent was evaporated under reduced pressure and the white solid was treated with 400 ml water. The suspension was stirred for 1 hour, the solid was filtered off, washed with water 3 times and dried under reduced pressure, affording 6-bromo-[1,2,4]triazolo[1,5-a]pyridin-2-ylamine (21 g, 85%) as white solid. Mp.: 185°C. MS: m/z=213.0/215.1 (M+H+).

c) 2,6-Dibromo-[1,2,4]triazolo[1,5-a]pyridine




A mixture of 6-bromo-[1,2,4]triazolo[1,5-a]pyridin-2-amine (20.86 g, 97.9 mmol), sodium nitrite (67.6 g, 979 mmol) and benzyltriethylammonium bromide (53.3 g, 196 mmol) in bromoform (1.47 kg, 508 ml, 5.81 mol) was stirred for 30 minutes at 25 °C. Then dichloroacetic acid (25.3 g, 16.2 ml, 196 mmol) was added and the mixture was stirred for 20 hours at 25 °C under exclusion of light. 600 ml of water were added to the mixture and stirred for 30 minutes. The mixture was diluted with water and dichloromethane, filtrated over dicalite and the filtrate extracted 3 times with dichloromethane. The organic layers were combined, washed with water, dried over magnesium sulfate, filtrated and evaporated to dryness under reduced pressure. The crude material was purified by flash chromatography in 2 portions over 2 x 70 g silica columns using dichloromethane/methanol 5 % as eluent, affording 2,6-dibromo-[1,2,4]triazolo[1,5-a]pyridine (7 g, 26%) as light brown solid. Mp.: 201°C. MS: m/z=277.7/279.9 (M+H+).

d) 6-Bromo-2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridine




A mixture of 2,6-dibromo-[1,2,4]triazolo[1,5-a]pyridine (1 g, 3.61 mmol) and morpholine (10 g, 10 ml, 115 mmol) was refluxed for 4 hours under argon atmosphere. The mixture was evaporated to dryness under reduced pressure. Flash chromatography (silica gel, 50 g, 30-100% heptane/ethyl acetate) afforded 6-bromo-2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridine (672 mg, 66%) as a light yellow solid. Mp.: 136°C. MS: m/z=283.0/285.0 (M+H+).

e) (2-Morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-carbamic acid tert-butyl ester




A mixture of 4-(6-bromo-[1,2,4]triazolo[1,5-a]pyridin-2-yl)morpholine (200 mg, 706 μ mol), tert-butyl carbamate (99.3 mg, 848 μ mol) and cesium carbonate (322 mg, 989 μ mol) was purged several times with argon, then tris(dibenzylideneacetone)dipalladium(0) (12.9 mg, 14.1 μ mol) and 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (16.3 mg, 28.3 μ mol) were added and the mixture was purged with argon again, the vessel was then capped. The resulting mixture was refluxed for 20 hours. The mixture was evaporated to dryness under reduced pressure. Final purification by flash chromatography (20 g silicagel column using heptane / ethyl acetate 70-100 % as eluent) afforded (2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-carbamic acid tert-butyl ester (46 mg, 20.4%) as yellow solid. Mp.: 166°C. MS: m/z=320.0 (M+H+).

f) 2-Morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylamine




A mixture of tert-butyl 2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamate (363 mg, 1.14 mmol) and hydrochloric acid (5M in diethyl ether, 24.0 g, 20 ml, 100 mmol) was stirred for 5 hours at 25 °C and then the solvent was evaporated under reduced pressure. The light yellow residue was made alkaline using sodium hydroxide solution (2M in water). The mixture was extracted 2 times with ethyl acetate, the organic layers were combined, washed with water and brine, dried over magnesium sulfate, filtrated and evaporated to dryness, affording 2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylamine (227 mg, 91%) as grey solid. Mp.: 165°C. MS: m/z=220.1 (M+H+).

g) 1-Methyl-5-(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid ethyl ester




A mixture of 2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-amine (225 mg, 1.03 mmol), 4-(ethoxycarbonyl)-1-methyl-1H-pyrazole-5-carboxylic acid (224 mg, 1.13 mmol), propylphosphonic anhydride in ethyl acetate 50% (1.63 g, 1.51 ml, 2.57 mmol) and N,N-diisopropylethylamine (398 mg, 524 μ l, 3.08 mmol) in tetrahydrofuran (7 ml) was refluxed for 18 hours. The solvent was evaporated, the residue was triturated with saturated aqueous sodium hydrogen carbonate. The precipitated solid was filtered off, washed with water and dried in vacuo, affording 1-methyl-5-(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid ethyl ester (292 mg, 71%) as light brown solid. Mp.: 233.5°C. MS: m/z=400.1 (M+H+).

h) 1-Methyl-5-(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid



A mixture of 1-methyl-5-(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid ethyl ester (282 mg, 0.71 mmol) and lithium hydroxide hydrate (88.9 mg, 2.12 mmol) in methanol (10 ml) and water (3 ml) was stirred for 18 hours at 55°C. The solvents were evaporated under reduced pressure and the residue treated with water (5 ml) and hydrogen chloride (1 molar in water, 2.12 ml). The precipitated solid was filtered off, washed with water and dried under reduced pressure, affording 1-methyl-5-(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid (187 mg, 71%) as light brown solid. Mp.: 294°C. MS: m/z=372.0 (M+H+).


i) 2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid (2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide



Using pyrrolidine and 1-methyl-5-(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Brown solid (110 mg, 64%). MS: $m/z=425$ ($M+H^+$).

5 Example 14


2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide



10 Using morpholine and 1-methyl-5-(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. White solid (130 mg, 65%). MS: $m/z=441$ ($M+H^+$).

Example 15


2-Methyl-4-(azetidine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide



Using azetidine and 1-methyl-5-(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Light yellow solid (99 mg, 53%). MS: $m/z=411$ ($M+H^+$). Ames test negative.

Example 16


2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(ethyl-methyl-amide) 3-[(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]



Using ethyl-methyl-amine and 1-methyl-5-(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. White solid (100 mg, 53%). MS: $m/z=413$ ($M+H^+$).

Example 17


4-(3-Fluoro-azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide



Using 3-fluoro-azetidine and 1-methyl-5-(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Off-white solid (150 mg, 76%). MS: $m/z=429$ ($M+H^+$).

5 **Example 18**


2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]




Using cyclopropyl-methyl-amine and 1-methyl-5-(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Off-white solid (114 mg, 100%). Mp.: 172°C. MS: $m/z=425.0$ ($M+H^+$).

Example 19

2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide




a) (6-Bromo-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-dimethyl-amine




- 5 Using dimethylamine, this compound was prepared following the same method as for the synthesis of 6-bromo-2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridine White solid (0.65 g, 75%). MS: $m/z=242$ ($M+H^+$).

b) (2-Dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-carbamic acid tert-butyl ester



- 10 Using (6-bromo-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-dimethyl-amine, this compound was prepared following the same method as for the synthesis of (2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-carbamic acid tert-butyl ester. Reddish yellow solid (250 mg, 33%). MS: $m/z=278$ ($M+H^+$).


c) N²,N²-Dimethyl-[1,2,4]triazolo[1,5-a]pyridine-2,6-diamine



15


Using (2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-carbamic acid tert-butyl ester, this compound was prepared following the same method as for the synthesis of 2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylamine. White solid (64 mg, 83%). MS: $m/z=178$ ($M+H^+$).

d) 5-(2-Dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid ethyl ester



5 Using N²*,N²*-dimethyl-[1,2,4]triazolo[1,5-a]pyridine-2,6-diamine, this compound was prepared following the same method as for the synthesis of 1-methyl-5-(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid ethyl ester. White solid (64 mg, 48%). Mp.: 194°C. MS: m/z=358 (M+H⁺).


e) 5-(2-Dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid



10

Using 5-(2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid ethyl ester, this compound was prepared following the same method as for the synthesis of 1-Methyl-5-(2-morpholino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid. Brown solid (41 mg, 70%). MS: m/z=330 (M+H⁺).


15 f) 2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-dimethylamino-[1,2,4]triazolo [1,5-a]pyridin-6-yl)-amide



Using morpholine and 5-(2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Off-white solid (117 mg, 58%). MS: $m/z=399$ ($M+H^+$). Ames test negative.

Example 20


4-(Azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide



Using azetidine and 5-(2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Brown solid (93 mg, 42%). MS: $m/z=369$ ($M+H^+$). Ames test negative.

Example 21


2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid (2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide



Using pyrrolidine and 5-(2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Light brown solid (120 mg, 61%). MS: $m/z=383$ ($M+H^+$).

Example 22


2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 3-[(2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide] 4-(ethyl-methyl-amide)



- 5 Using ethyl-methyl-amine and 5-(2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Light yellow solid (129 mg, 57%). MS: $m/z=371$ ($M+H^+$).

Example 23

- 10 4-(3-Fluoro-azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide




- 15 Using 3-fluoro-azetidine and 5-(2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Light yellow solid (160 mg, 68%). MS: $m/z=387$ ($M+H^+$).

Example 24

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

-46-




Using cyclopropyl-methyl-amine and 5-(2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Light brown oil (30 mg, 49%). MS: $m/z=383.0$

5 (M+H+).

Example 25

6-cyclopropyl-3-methoxy-N-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)pyrazine-2-carboxamide




10 In a 5 ml round bottomed flask 6-cyclopropyl-3-methoxypyrazine-2-carboxylic acid (40 mg, 206 μmol), 2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-amine (47.6 mg, 227 μmol) and THF (1 ml) were mixed to form a solution. Then propylphosphonic anhydride in ethyl acetate 50% (328 mg, 301 μl , 515 μmol) and N,N-diisopropylethylamine (106 mg, 144 μl , 824 μmol) were added and the mixture heated to 70°C for 2 hours. The reaction mixture was allowed to cool to room


15 temperature, then 1.5 ml of acetonitrile:water (1:1) was added and the mixture stirred for 10 minutes. The precipitate formed was then filtered off, washed with 1 ml of acetonitrile:water (1:1) and 1 ml acetonitrile, then dried in vacuo to give 6-cyclopropyl-3-methoxy-N-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)pyrazine-2-carboxamide (45.5 mg, 57.2 %) as a yellow solid. MS: $m/z=387.2$ (M+H+).

20 Example 26

2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide




a) (6-Bromo-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-diethyl-amine




Using diethylamine, this compound was prepared following the same method as for the
5 synthesis of 6-bromo-2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridine Light brown solid (2.0 g, 82%). MS: $m/z=270$ ($M+H^+$).

b) N²*,N²*-Diethyl-[1,2,4]triazolo[1,5-a]pyridine-2,6-diamine



Using (6-bromo-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-diethyl-amine, this compound was
10 prepared following the same methods as for the synthesis of 2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylamine. Brown solid (500 mg, 66%). MS: $m/z=206$ ($M+H^+$).


c) 5-(2-Diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid ethyl ester



Using N²*,N²*-diethyl-[1,2,4]triazolo[1,5-a]pyridine-2,6-diamine, this compound was
15 prepared following the same method as for the synthesis of 1-methyl-5-(2-morpholin-4-yl-

[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid ethyl ester. Off white solid (1.2 g, 45%). MS: $m/z=386$ ($M+H^+$).


d) 5-(2-Diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid



5

Using 5-(2-diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid ethyl ester, this compound was prepared following the same method as for the synthesis of 1-Methyl-5-(2-morpholino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid. Off white solid (500 mg, 54%). MS: $m/z=358$ ($M+H^+$).


10 e) 2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-diethylamino-[1,2,4]triazolo [1,5-a]pyridin-6-yl)-amide



15 Using morpholine and 5-(2-diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. White solid (65 mg, 94%). MS: $m/z=357$ ($M+H^+$).

Example 27


2-Methyl-4-(pyrrolidine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-diethylamino-[1,2,4]triazolo [1,5-a]pyridin-6-yl)-amide



Using pyrrolidine and 5-(2-diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. White solid (60 mg, 75%). MS: $m/z=411$ ($M+H^+$).

5 Example 28


2-Methyl-4-(azetidine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-diethylamino-[1,2,4]triazolo [1,5-a]pyridin-6-yl)-amide



Using azetidine and 5-(2-diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Off white solid (60 mg, 77%). MS: $m/z=397$ ($M+H^+$).

Example 29


2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 3-[(2-diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide] 4-(ethyl-methyl-amide)



Using ethyl-methyl-amine and 5-(2-diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. White solid (50 mg, 64%). MS: $m/z=399$ ($M+H^+$).

Example 30


- 5 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-diethylamide 3-[(2-diethylamino-1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide



- 10 Using diethylamine and 5-(2-diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. White solid (70 mg, 87%). MS: $m/z=413$ ($M+H^+$).

Example 31


2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide




- 15 Using cyclopropyl-methyl-amine and 5-(2-diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. White solid (65 mg, 81%). MS: $m/z=411$ ($M+H^+$).

Example 32

2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide




a) 6-Bromo-2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridine



5

Using pyrrolidine, this compound was prepared following the same method as for the synthesis of 6-bromo-2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridine. Off white solid (1.0 g, 41%). MS: $m/z=268$ ($M+H^+$).


b) 2-Pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylamine



10

Using 6-Bromo-2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridine, this compound was prepared following the same method as for the synthesis of 2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylamine. Brown solid (1.1 g, 81%). MS: $m/z=204$ ($M+H^+$).


c) 1-Methyl-5-(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid ethyl ester



15


Using 2-Pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylamine, this compound was prepared following the same method as for the synthesis of 1-methyl-5-(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid ethyl ester. Brown solid (1.1 g, 65%). MS: $m/z=384$ ($M+H^+$).

- 5 d) 1-Methyl-5-(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid



- 10 Using 1-Methyl-5-(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid ethyl ester, this compound was prepared following the same method as for the synthesis of 1-Methyl-5-(2-morpholino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid. White solid (430 mg, 93%). MS: $m/z=356$ ($M+H^+$).

- e) 2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide




15

Using morpholine and 1-Methyl-5-(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. White solid (68 mg, 95%). MS: $m/z=425$ ($M+H^+$).

Example 33


4-(Azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-pyrrolidin-1-yl-1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide



Using azetidine and 1-Methyl-5-(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. White solid (40 mg, 51%). MS: $m/z=395$ ($M+H^+$).

Example 34

2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid (2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide




10

Using pyrrolidine and 1-Methyl-5-(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. White solid (46 mg, 58%). MS: $m/z=409$ ($M+H^+$).

Example 35

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide


15



Using cyclopropyl-methyl-amine and 1-Methyl-5-(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. White solid (50 mg, 43%). MS: $m/z=409$ ($M+H^+$).

5 Example 36


2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(ethyl-methyl-amide) 3-[(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide



Using ethyl-methyl-amine and 1-Methyl-5-(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. White solid (59 mg, 53%). MS: $m/z=397$ ($M+H^+$).

Example 37


2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-diethylamide 3-[(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide




Using diethylamine and 1-Methyl-5-(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. White solid (65 mg, 56%). MS: $m/z=411$ ($M+H^+$).

Example 38

- 5 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-diethylamide 3-{[2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide




- a) 6-Bromo-2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridine



- 10 Using pyrrolidine, this compound was prepared following the same method as for the synthesis of 6-bromo-2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridine Brown solid (2.5 g, 47%). MS: $m/z=298$ ($M+H^+$).


- b) 2-(3-Methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-ylamine



- 15 Using 6-Bromo-2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridine, this compound was prepared following the same method as for the synthesis of 2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylamine. Brown solid (1.4 g, 71%). MS: $m/z=234$ ($M+H^+$).


c) 5-[2-(3-Methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl]-1-methyl-1H-pyrazole-4-carboxylic acid ethyl ester

-56-




Using 2-(3-Methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-ylamine, this compound was prepared following the same method as for the synthesis of 1-methyl-5-(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid ethyl ester. Off white solid (1.2 g, 59%). MS: $m/z=414$ ($M+H^+$).

d) 5-[2-(3-Methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl]-1-methyl-1H-pyrazole-4-carboxylic acid



Using 5-[2-(3-Methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl]-1-methyl-1H-pyrazole-4-carboxylic acid ethyl ester, this compound was prepared following the same method as for the synthesis of 1-Methyl-5-(2-morpholino-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid. Off white solid (690 mg, 82%). MS: $m/z=386$ ($M+H^+$).


e) 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-diethylamide 3-{[2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide



Using diethylamine and 5-[2-(3-Methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl]-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Off white solid (115 mg, 77%). MS: $m/z=441$ ($M+H^+$).

Example 39


- 5 2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid [2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide



- 10 Using morpholine and 5-[2-(3-Methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl]-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Off white solid (42 mg, 51%). MS: $m/z=455$ ($M+H^+$).

Example 40


2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(ethyl-methyl-amide) 3-{[2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide}



- 15 Using ethyl-methyl-amine and 5-[2-(3-Methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl]-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Off white solid (85 mg, 59%). MS: $m/z=427$ ($M+H^+$).

Example 41


2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-{[2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide}



Using cyclopropyl-methyl-amine and 5-[2-(3-Methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl]-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Off white solid (80 mg, 54%). MS: m/z=439 (M+H+).

Example 42

2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid [2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide




Using pyrrolidine and 5-[2-(3-Methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl]-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Off white solid (72 mg, 53%). MS: m/z=439 (M+H+).

Example 43

4-(Azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid [2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide


-59-



Using azetidine and 5-[2-(3-Methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl]-1-methyl-1H-pyrazole-4-carboxylic acid, the title compound was prepared in the same manner as described for example 2. Off white solid (57 mg, 43%). MS: $m/z=425$ ($M+H^+$).

Claims

1. A compound of formula (I)



wherein

R^1 is aryl, heteroaryl or NR^4R^5 , wherein said aryl and said heteroaryl can be substituted by 1 to 3 substituents independently selected from the group consisting of hydroxyl, halogen, C_1 - C_7 alkyl, C_1 - C_7 alkoxy and C_1 - C_7 haloalkyl;

R^2 is hydrogen, halogen or C_1 - C_7 alkyl;

R^3 is aryl or heteroaryl, wherein said aryl and said heteroaryl can be substituted by 1 to 3 substituents independently selected from the group consisting of C_1 - C_7 alkyl, hydroxyl, halogen, $-C(O)-NR^6R^7$ and $-C(O)-O-R^8$;

R^4 and R^5 are independently C_1 - C_7 alkyl, C_3 - C_8 cycloalkyl, together with the nitrogen atom to which they are attached, form a heterocycloalkyl which can be substituted by 1 to 3 substituents independently selected from the group consisting of hydroxyl, halogen, C_1 - C_7 alkyl, C_1 - C_7 alkoxy and C_1 - C_7 haloalkyl;

R^6 and R^7 , are independently C_1 - C_7 alkyl, C_3 - C_8 cycloalkyl, together with the nitrogen atom to which they are attached, form a heterocycloalkyl which can be substituted by 1 to 3 substituents independently selected from the group consisting of hydroxyl, halogen, C_1 - C_7 alkyl, C_1 - C_7 alkoxy and C_1 - C_7 haloalkyl;


R^8 is hydrogen, C_1 - C_7 alkyl, cycloalkyl;

or a pharmaceutically acceptable salt thereof.

2. The compound according to claim 1, wherein R^1 is phenyl, pyridinyl, thiazolyl or NR^4R^5 , wherein R^4 and R^5 are independently selected from C_1 - C_3 alkyl, R^4 and R^5 together with the nitrogen atom to which they are attached, form a pyrrolidinyl or morpholinyl ring.

-61-

3. The compound according to claim 1 or 2, wherein R^2 is hydrogen.
4. The compound of any one of claims 1 to 3, wherein R^3 is :



5 wherein:

A_1 is NR^9 ,

A_2 is $NR^{9'}$,

R^6 and R^7 are independently selected from C_1 - C_3 alkyl, C_3 - C_8 cycloalkyl, both optionally substituted by 1 to 3 substituents independently selected from halogen, C_1 - C_3 alkyl, C_1 - C_3 alkoxy and C_1 - C_3 haloalkyl;

R^6 and R^7 together with the nitrogen atom to which they are attached form a heterocycloalkyl of 4 to 7 ring atoms, comprising 1 or 2 ring heteroatoms selected from N and O, the heterocycloalkyl optionally substituted by 1 to 3 substituents independently selected from halogen, C_1 - C_3 alkyl, C_1 - C_3 alkoxy and C_1 - C_3 haloalkyl,

R^9 and $R^{9'}$ are independently selected from hydrogen, C_1 - C_3 alkyl, C_1 - C_3 alkoxy and C_1 - C_3 haloalkyl.

5. The compound according to claim 4, wherein:

R^6 and R^7 are independently selected from C_1 - C_3 alkyl, C_3 - C_8 cycloalkyl,

R^6 and R^7 together with the nitrogen atom to which they are attached form azetidiny, pyrrolidinyl and morpholinyl which are optionally substituted by 1 to 3 substituents independently selected from the group consisting of halogen, C_1 - C_3 alkyl, C_1 - C_3 alkoxy and C_1 - C_3 haloalkyl.

R^9 is selected from C_1 - C_3 alkyl, C_1 - C_3 alkoxy and C_1 - C_3 haloalkyl, preferably C_1 - C_3 alkyl,

$R^{9'}$ is selected from hydrogen and C_1 - C_3 alkyl, preferably hydrogen.

6. The compound according to any of claims 1 to 5, selected from the group consisting of:

- 2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid (2-phenyl-
5 [1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide
- 1-Methyl-5-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-ylcarbamoyl)-1H-pyrazole-4-carboxylic acid
- 4-(Azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide
- 10 4-(Azetidine-1-carbonyl)-1-methyl-N-(2-(thiazol-4-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-1H-pyrazole-5-carboxamide
- 1-Methyl-4-(morpholine-4-carbonyl)-N-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-1H-pyrazole-5-carboxamide
- 4-(3-Fluoro-azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-phenyl-
15 [1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide
- 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(ethyl-methyl-amide) 3-[(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]
- 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]
- 20 2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide
- 4-(Azetidine-1-carbonyl)-1-methyl-N-(2-(pyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-1H-pyrazole-5-carboxamide
- N4-Cyclopropyl-N4,1-dimethyl-N5-(2-(pyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-
25 1H-pyrazole-4,5-dicarboxamide
- 1-Methyl-N-(2-(pyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-4-(pyrrolidine-1-carbonyl)-1H-pyrazole-5-carboxamide
- 2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid (2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-4-(azetidine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

5 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(ethyl-methyl-amide) 3-[(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

4-(3-Fluoro-azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

10 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

4-(Azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

15 2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid (2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 3-[(2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide] 4-(ethyl-methyl-amide)

20 4-(3-Fluoro-azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

6-cyclopropyl-3-methoxy-N-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)pyrazine-2-carboxamide

25 2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-4-(pyrrolidine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-diethylamino-[1,2,4]triazolo [1,5-a]pyridin-6-yl)-amide

2-Methyl-4-(azetidine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-diethylamino-[1,2,4]triazolo [1,5-a]pyridin-6-yl)-amide

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 3-[(2-diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide] 4-(ethyl-methyl-amide)

5 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-diethylamide 3-[(2-diethylamino-1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-diethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

10 2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

4-(Azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-pyrrolidin-1-yl-1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid (2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

15 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(ethyl-methyl-amide) 3-[(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

20 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-diethylamide 3-[(2-pyrrolidin-1-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-diethylamide 3-{[2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide}

2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid [2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide

25 2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(ethyl-methyl-amide) 3-{[2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide}

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-{[2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide}

2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid [2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide

4-(Azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid [2-(3-methoxy-pyrrolidin-1-yl)-[1,2,4]triazolo[1,5-a]pyridin-6-yl]-amide.

5 7. The compound according to any of claims 1 to 6, selected from the group consisting of:

4-(Azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

10 1-Methyl-4-(morpholine-4-carbonyl)-N-(2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-1H-pyrazole-5-carboxamide

4-(3-Fluoro-azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-phenyl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-4-(morpholine-4-carbonyl)-2H-pyrazole-3-carboxylic acid (2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

15 2-Methyl-4-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carboxylic acid (2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-morpholin-4-yl-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]


20 4-(Azetidine-1-carbonyl)-2-methyl-2H-pyrazole-3-carboxylic acid (2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide

2-Methyl-2H-pyrazole-3,4-dicarboxylic acid 4-(cyclopropyl-methyl-amide) 3-[(2-dimethylamino-[1,2,4]triazolo[1,5-a]pyridin-6-yl)-amide]

8. A pharmaceutical composition comprising a compound in accordance with any one of claims 1 - 7 and a therapeutically inert carrier.


25 9. A process for the manufacture of compounds of formula I as defined in any one of claims 1 – 7, which process comprises

a) reacting a compound of formula (2)



with

b) a compound of formula (3)



wherein R^1 , R^2 , R^3 are as defined above and, if desired, converting the compounds into pharmaceutically acceptable salts thereof.

10. The use of a compound according to any one of claims 1 - 7 for the treatment or prophylaxis of psychotic disorders, schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia, delusional disorder, substance-induced psychotic disorder, anxiety disorders, panic disorder, obsessive/compulsive disorders, acute stress disorder, generalized anxiety disorder, drug addictions, movement disorders, Parkinson's disease, restless leg syndrome, cognition deficiency disorders, Alzheimer's disease, multi-infarct dementia, mood disorders, depression, bipolar disorders, neuropsychiatric conditions, psychosis, attention-deficit/hyperactivity disorder, attentional disorders, diabetes and related disorders, type 2 diabetes mellitus, neurodegenerative disorders, Huntington's disease, multiple sclerosis, stroke, spinal cord injury, solid tumors, hematological malignancies, renal cell carcinoma or breast cancer.

11. The use of a compound according to any one of claims 1 - 7 for the preparation of a medicament for the treatment or prophylaxis of psychotic disorders, schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia, delusional disorder, substance-induced psychotic disorder, anxiety disorders, panic disorder, obsessive/compulsive disorders, acute stress disorder, generalized anxiety disorder, drug addictions, movement disorders, Parkinson's disease, restless leg syndrome, cognition deficiency disorders, Alzheimer's disease, multi-infarct dementia, mood disorders, depression, bipolar disorders, neuropsychiatric conditions, psychosis, attention-deficit/hyperactivity disorder, attentional disorders, diabetes and related disorders, type 2 diabetes mellitus, neurodegenerative disorders, Huntington's disease, multiple sclerosis, stroke, spinal cord injury, solid tumors, hematological malignancies, renal cell carcinoma or breast cancer.

12. A compound according to any one of claims 1 - 7 for the treatment or prophylaxis of psychotic disorders, schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia, delusional disorder, substance-induced psychotic disorder, anxiety disorders, panic disorder, obsessive/compulsive disorders, acute stress disorder, generalized anxiety disorder, drug addictions, movement disorders, Parkinson's disease, restless leg syndrome, cognition deficiency disorders, Alzheimer's disease, multi-infarct dementia, mood disorders, depression, bipolar disorders, neuropsychiatric conditions, psychosis, attention-deficit/hyperactivity disorder, attentional disorders, diabetes and related disorders, type 2 diabetes mellitus, neurodegenerative disorders, Huntington's disease, multiple sclerosis, stroke, spinal cord injury, solid tumors, hematological malignancies, renal cell carcinoma or breast cancer.

13. A method for the treatment or prophylaxis of psychotic disorders, schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia, delusional disorder, substance-induced psychotic disorder, anxiety disorders, panic disorder, obsessive/compulsive disorders, acute stress disorder, generalized anxiety disorder, drug addictions, movement disorders, Parkinson's disease, restless leg syndrome, cognition deficiency disorders, Alzheimer's disease, multi-infarct dementia, mood disorders, depression, bipolar disorders, neuropsychiatric conditions, psychosis, attention-deficit/hyperactivity disorder, attentional disorders, diabetes and related disorders, type 2 diabetes mellitus, neurodegenerative disorders, Huntington's disease, multiple sclerosis, stroke, spinal cord injury, solid tumors, hematological malignancies, renal cell carcinoma or breast cancer, which method comprises administering an effective amount of a compound as defined in any one of claims 1 - 7.

14. The invention as hereinbefore described.