

LACRIMAL IMPLANTS AND RELATED METHODS

CLAIM OF PRIORITY

Benefit of priority is hereby claimed to U.S. Provisional Patent Application
5 Serial No. 60/970,696 filed on September 7, 2007, U.S. Provisional Patent
Application Serial No. 60/970,720 filed on September 7, 2007, U.S. Provisional
Patent Application Serial No. 60/974,367 filed on September 21, 2007, U.S.
Provisional Patent Application Serial No. 61/033,211 filed on March 3, 2008, U.S.
Provisional Patent Application Serial No. 61/036,816 filed on March 14, 2008, U.S.
10 Provisional Patent Application Serial No. 61/049,360 filed on April 30, 2008, U.S.
Provisional Patent Application Serial No. 61/052,595 filed on May 12, 2008, and
U.S. Provisional Patent Application Serial No. 61/075,309 filed on June 24, 2008,
the specifications of which are herein incorporated by reference in their entirety.

15 TECHNICAL FIELD

This patent document pertains generally to ophthalmic devices, and
particularly to ocular implants. More particularly, but not by way of limitation, this
patent document pertains to lacrimal implants, methods of making such implants,
and methods of treating ocular, respiration, inner ear or other diseases or disorders
20 (e.g., pulmonary or immunological disorders) using such implants.

BACKGROUND

Dry eye, including keratoconjunctivitis sicca, is a common ocular condition
that can require therapy. Dry eye has been experienced by a broad demographic
25 band, and is common in elderly individuals. A variety of current treatment
modalities target physiological conditions that contribute to dry eye, including
augmentation of normal tear fluid, enhancement of tear film component production,
and methods to enhance the residence time of tears, such as blocking the tear flow
from an eye into and through a lacrimal canaliculus.

Many current tear flow blockage techniques have drawbacks, including being irreversible in nature. For instance, some tear flow blockage techniques involve closing the canalicular canal by stitching the punctal opening shut or by using electrical or laser cauterization to seal the punctal opening. Although such procedures can provide the desired result of blocking tear flow to treat a dry eye, they are unfortunately not reversible without reconstructive surgery.

In addition to dry eye symptom relief, a variety of challenges face patients and physicians in the area of ocular, respiration and inner ear disease or disorder management, including adequate drug or other therapeutic agent delivery to the eyes, nasal passage or inner ear. In ocular management, for example, many current ocular drug delivery systems require repetitive manual administration and are often ineffective due to a lack of patient compliance or inadequate drug concentrations reaching the eye.

In order to treat eye infection, inflammation of an eye, glaucoma and other ocular diseases or disorders, drugs or other therapeutic agents are often required to be administered to the eye. A conventional method of drug delivery is by topical drop application to the eye's surface. Topical eye drops, though effective, can be inefficient. As one example, when an eye drop is instilled in an eye, it often overfills the conjunctival sac (i.e., the pocket between the eye and the lids) causing a substantial portion of the drop to be lost due to overflow of the lid margin and spillage onto the cheek. In addition, a large portion of the drop remaining on the ocular surface can be washed away into and through a lacrimal canaliculus, thereby diluting the concentration of the drug before it can absorbably treat the eye. Moreover, topically applied drugs often have a peak ocular effect for about two hours post-application, after which additional applications of the drugs should be, but are often not, administered to maintain the desired drug therapeutic benefit.

To compound ocular management difficulty, patients often do not use their eye drops as prescribed. This poor compliance can be due to, for example, an initial stinging or burning sensation caused by the eye drop and experienced by a patient. Instilling eye drops in one's own eye can be difficult, in part because of the normal

reflex to protect the eye. Therefore, one or more drops may miss the eye. Older patients may have additional problems instilling drops due to arthritis, unsteadiness, and decreased vision. Pediatric and psychiatric populations pose difficulties as well.

5 In a field different from ocular management, control of respiration-related (e.g., allergies) and inner ear diseases or disorders often requires repetitive manual digestion or other intake of a medication (e.g., drugs or other therapeutic agents), and as such, can be ineffective due to a lack of patient compliance or non-localized drug delivery.

10 EXEMPLARY ASPECTS AND EMBODIMENTS OF THE INVENTION

The present inventors have recognized various promising techniques to increase the residence time of tears on an eye and delivery of drug or other therapeutic agent to the eye, nasal passage, inner ear or other system. These techniques can include placing a removable, and optionally drug releasing, lacrimal
15 implant through a lacrimal punctum and into the associated canaliculus. It is believed that by designing lacrimal implants that utilize the features of the nasolacrimal drainage system, patient comfort and implant retention in the ocular anatomy can be satisfied. In this way, the present lacrimal implants can overcome some of the drawbacks associated with current dry eye relief, such as being
20 irreversible in nature, and ocular drug administration (e.g., manual drop instillation or digestion), such as poor patient compliance, waste, untimely application, or non-localized delivery.

Further yet, the present inventors have recognized that a lacrimal implant can benefit from one or more of: the ability to be easily implanted and removed
25 without much biasing of the lacrimal punctum or associated canaliculus, the ability to be securely retainable in the lacrimal canaliculus upon implantation, optionally without being pre-sized to a particular lacrimal punctum or canaliculus, the ability to permit tear fluid, drug or other agent to flow into the nasolacrimal system, and, when made and used as a drug delivery system, the ability to allow for the

sustained, localized release of one or more drugs or other therapeutic agents at a desired therapeutic level for an extended period of time.

Lacrimal implants for treating diseases or disorders are disclosed. More particularly, lacrimal implants, methods of making such implants, and methods of
5 treating ocular, respiration, inner ear, pulmonary or immunological diseases or disorders using such implants are disclosed.

To better illustrate the subject matter described herein, a non-limiting list of exemplary aspects and embodiments is provided here:

- 10 1. A lacrimal implant insertable into a lacrimal canaliculus, comprising:
an implant body, including first and second portions, the implant body
extending from a proximal end of the first portion to a distal end of the second
portion;
the proximal end of the first portion defining a longitudinal proximal axis
15 and the distal end of the second portion defining a longitudinal distal axis;
the implant body configured such that, when implanted in the lacrimal
canaliculus, an angled intersection exists between the proximal axis and the distal
axis for biasing at least a portion of the implant body against at least a portion of the
lacrimal canaliculus located at or more distal to a canalicular curvature; and
20 wherein the second portion of the implant body includes a longitudinal
length having a magnitude less than four times a longitudinal length of the first
portion of the implant body.
2. The lacrimal implant according to aspect 1, wherein the implant body is
25 configured such that an angled intersection exists between the proximal axis and the
distal axis prior to being implanted in the lacrimal canaliculus.
3. The lacrimal implant according to any of aspects 1 or 2, wherein the implant
body is configured to partially or completely inhibit fluid flow into and through the
30 lacrimal canaliculus.

4. The lacrimal implant according to any of aspects 1-3, wherein a distal end of the first portion is integral with the second portion at or near a proximal end of the second portion.
- 5 5. The lacrimal implant according to any of aspects 1-4, wherein one or both of the first portion or the second portion includes a fluid swellable retention element configured to expand.
6. The lacrimal implant according to aspect 5, wherein the second portion
10 includes the fluid swellable retention element, the fluid swellable retention element configured to expand laterally, relative to the proximal axis of the first portion, when the implant body is implanted.
7. The lacrimal implant according to any of aspects 5 or 6, wherein the fluid
15 swellable retention element includes a portion configured to expand laterally in a direction away from a lacrimal canaliculus ampulla when the implant body is implanted.
8. The lacrimal implant according to any of aspects 5-7, wherein the fluid
20 swellable retention element includes a portion configured to expand laterally in a direction toward a lacrimal canaliculus ampulla when the implant body is implanted.
9. The lacrimal implant according to any of aspects 1-8, wherein the second
portion includes an expandable retention element comprising at least one of a coil, a
25 braid, a stent, a mesh tube, a suture, a thermoset polymer, a thermoplastic, a heat activatable material, or a shape memory material, the expandable retention element configured to expand laterally, to form the angled intersection, when the implant body is implanted.

10. The lacrimal implant according to any of aspects 1-9, comprising an expandable retention element disposed around a portion of the second portion, the expandable retention element configured to bias the second portion away from a wall of the lacrimal canaliculus upon expansion.

5

11. The lacrimal implant according to any of aspects 1-10, wherein the second portion includes an arm member movable between a first configuration and a second configuration; the arm member, in the first configuration, disposable along the implant body for insertion into the lacrimal canaliculus and, in the second configuration, laterally extendable from one side of the implant body.

10

12. The lacrimal implant according to any of aspects 1-11, wherein the second portion includes an integral dilator, the integral dilator generally narrowing from a location near a proximal end of the second portion to the distal end of the second portion to facilitate implantation of the implant body into the lacrimal canaliculus.

15

13. The lacrimal implant according to aspect 12, wherein a diameter of an integral dilator tip is between about 0.2 millimeters and about 0.5 millimeters.

14. The lacrimal implant according to any of aspects 12 or 13, wherein an outer surface slope of the integral dilator, as measured from the location near the proximal end of the second portion to the distal end of the second portion, is between about 1 degree and about 10 degrees with respect to the distal axis.

20

15. The lacrimal implant according to any of aspects 1-14, wherein the second portion includes at least one undulation.

25

16. The lacrimal implant according to any of aspects 1-15, wherein at least one of the first portion or the second portion comprises at least one intermediately-disposed annular, semi-annular, column-like, or barrel-like projection, the

30

intermediately-disposed projection having a cross-sectional size greater than an adjacent implant body portion.

17. The lacrimal implant according to any of aspects 1-16, comprising a
5 graspable projection extending at least partially from the proximal end of the first portion, the graspable projection configured to seat against or near a lacrimal punctum when the implant body is implanted.

18. The lacrimal implant according to aspect 17, wherein the second portion
10 includes an element extending or expanding laterally into a lacrimal canaliculus ampulla when the implant body is implanted.

19. The lacrimal implant according to any of aspects 17 or 18, wherein the
15 graspable projection extends laterally from the proximal end of the first portion, in a direction that is parallel to or away from an eye, when the implant body is implanted.

20. The lacrimal implant according to any of aspects 1-19, wherein the
20 implanted angled intersection of the proximal axis and the distal axis is at least about 45 degrees.

21. The lacrimal implant according to any of aspects 1-20, comprising a therapeutic agent.

22. The lacrimal implant according to aspect 21, comprising at least one drug
25 insert including a drug core, the drug core comprising the therapeutic agent.

23. The lacrimal implant according to aspect 22, wherein the drug core
30 comprises at least one exposed surface to deliver a sustained release.

24. A kit comprising the lacrimal implant according to any of aspects 1-23, and an instruction for using the lacrimal implant to treat an eye disease.
25. A kit comprising the lacrimal implant according to any of aspects 1-23, and
5 an instruction for using the lacrimal implant to treat a respiration-related disorder.
26. A kit comprising the lacrimal implant according to any of aspects 1-23, and an instruction for using the lacrimal implant to treat an inner ear disorder.
- 10 27. A lacrimal implant for insertion into a lacrimal canaliculus, comprising:
an implant body non-linearly extending from a proximal end portion
positionable within a vertical section of the lacrimal canaliculus to a distal end
portion positionable within a horizontal section of the lacrimal canaliculus and
having an intermediate portion therebetween;
- 15 the intermediate portion partially extending in a first direction toward the proximal end portion and partially extending in a second direction toward the distal end portion such that, when implanted in the lacrimal canaliculus, the implant body directionally biases laterally against at least a portion of the lacrimal canaliculus located at or more distal to a canalicular curvature; and
- 20 wherein the implant body inhibits fluid flow into and through the lacrimal canaliculus.
28. The lacrimal implant according to aspect 27, wherein a longitudinal length of the implant body positionable within the vertical section of the lacrimal
25 canaliculus is less than four times a longitudinal length of the implant body positionable within the horizontal section of the lacrimal canaliculus.
29. The lacrimal implant according to any of aspects 27 or 28, wherein the first direction extension of the intermediate portion is at an angle between about 45

degrees and about 135 degrees relative to the second direction extension of the intermediate portion.

30. The lacrimal implant according to any of aspects 27-29, wherein the
5 intermediate portion partially extends in a third direction, substantially opposite the second direction, toward a lacrimal canaliculus ampulla when the implant body is implanted.

31. The lacrimal implant according to any of aspects 27-30, wherein the second
10 direction extension includes a longitudinal dilator having a generally concave shape relative to the first direction extension; and wherein a radius of the generally concave shape is less than the radius of the canaliculus curvature.

32. The lacrimal implant according to any of aspects 27-31, wherein the second
15 direction extension includes a longitudinal dilator having a generally convex shape relative to the first direction extension.

33. The lacrimal implant according to any of aspects 27-32, wherein the second
20 direction extension includes a longitudinal dilator having an axis substantially perpendicular to an axis of the first direction extension.

34. The lacrimal implant according to any of aspects 27-33, wherein at least one
of the proximal end portion or the distal end portion comprises at least one
intermediately-disposed annular, semi-annular, column-like, or barrel-like
25 projection, the intermediately-disposed projection having a cross-sectional size greater than an adjacent implant body portion.

35. The lacrimal implant according to any of aspects 27-34, comprising a
graspable projection, the graspable projection extending laterally from the proximal
30 end portion.

36. The lacrimal implant according to any of aspects 27-35, comprising a fluid swellable material disposed on an outer surface portion of the implant body, the fluid swellable material configured to expand an outer surface diameter portion of the implant body when implanted.

5

37. The lacrimal implant according to any of aspects 27-36, comprising at least one of a first drug insert disposed in the proximal end portion or a second drug insert disposed in the distal end portion, one or both of the first or second drug inserts inhibiting fluid flow through the implant body and including at least one exposed surface configured to deliver a sustained release.

10

38. The lacrimal implant according to aspect 37, comprising a cavity in the proximal end portion, the cavity configured to house the first drug insert in the form of a drug core, the drug core including a first agent configured to treat an eye.

15

39. The lacrimal implant according to any of aspects 37 or 38, comprising a cavity in the distal end portion, the cavity configured to house the second drug insert in the form of a drug core, the drug core including a second agent configured to be received by a nasal passage.

20

40. A method of manufacturing a lacrimal implant insertable into a lacrimal canaliculus, the method comprising:

forming an implant body extending from a proximal end of a first body portion to a distal end of a second body portion, including

25

extending the second body portion to a longitudinal length which is less than four times a longitudinal length of the first body portion; and

configuring the proximal end and the distal end to respectively define, when implanted in the lacrimal canaliculus, a longitudinal proximal axis and a longitudinal distal axis that intersect at an angle such that the implant body is configured to directionally bias laterally against at least a

30

portion of the lacrimal canaliculus located at or more distal to a canaliculus curvature.

41. The method according to aspect 40, wherein forming one or both of the first
5 body portion or the second body portion includes forming at least one
intermediately-disposed annular, semi-annular, column-like, or barrel-like
projection, the intermediately-disposed projection having a cross-sectional size
greater than an adjacent implant body portion.

10 42. The method according to any of aspects 40 or 41, wherein forming the
second body portion includes forming a dilator generally narrowing from a location
near a proximal end of the second body portion to the distal end of the second body
portion.

15 43. The method according to any of aspects 40-42, wherein forming the dilator
includes forming an outer surface slope of the implant body, as measured from the
location near the proximal end of the second body portion to the distal end of the
second body portion, between about 1 degree and about 10 degrees with respect to
the longitudinal distal axis.

20

44. The method according to any of aspects 40-43, wherein forming the second
portion includes disposing a fluid swellable retention element near a distal end of the
first body portion, including disposing the fluid swellable retention element such
that a lateral expansion thereof, relative to the proximal axis, is configured to bias
25 against at least a portion of the lacrimal canaliculus or a lacrimal canaliculus
ampulla anatomy when the implant body is implanted.

45. The method according to any of aspects 40-44, comprising coating an outer
surface portion of the implant body with a fluid swellable material.

30

46. The method according to any of aspects 40-45, comprising disposing at least one of a first drug insert in the first body portion or a second drug insert in the second body portion, including positioning at least one of an exposed surface of the first drug insert adjacent to the proximal end or an exposed surface of the second drug insert adjacent to the distal end to provide a sustained release of a first agent or a second agent, respectively.

47. A method of treating a patient having at least one of an eye disorder, a respiration-related disorder, a pulmonary disorder, an immunological disorder or an inner ear disorder, the method comprising:

inserting a lacrimal implant into at least one lacrimal canaliculus of the patient, the lacrimal implant comprising,

an implant body, including first and second portions, the implant body extending from a proximal end of the first portion to a distal end of the second portion, the proximal end defining a longitudinal proximal axis and the distal axis defining a longitudinal distal axis;

the implant body configured such that, when implanted in a lacrimal canaliculus, an angled intersection exists between the proximal axis and the distal axis for biasing at least a portion of the implant body against at least a portion of the lacrimal canaliculus located at or more distal to a canalicular curvature; and

at least one of a first drug insert disposed in the first portion or a second drug insert disposed in the second portion, the first and second drug inserts configured to respectively provide a sustained release of a first agent or a second agent.

25

48. The method according to aspect 47, wherein inserting the lacrimal implant includes concurrently dilating the lacrimal canaliculus, using an integral dilator of the implant body, as the distal end of the second portion is moved into the lacrimal canaliculus.

30

49. The method according to any of aspects 47 or 48, comprising removing the inserted implant body from the lacrimal canaliculus.

50. The method according to any of aspects 47-49, comprising replacing the
5 lacrimal implant that has been inserted with a second lacrimal implant including at least one of the same first or second agent following an interval of time.

These and other embodiments, advantages, and aspects of the present lacrimal implants and methods will be set forth in part in following Detailed
10 Description. This Summary is intended to provide an overview of subject matter of the present patent application. It is not intended to provide an exclusive or exhaustive explanation of the present invention. The Detailed Description is included to provide further information about the present patent application.

15 BRIEF DESCRIPTION OF THE DRAWINGS

In the drawings, like numerals can be used to describe similar components throughout the several views. Like numerals having different letter suffixes can be used to represent different instances of similar components. The drawings illustrate generally, by way of example, but not by way of limitation, various embodiments
20 discussed in the present document.

FIGS. 1-2 illustrate examples of schematic views of anatomical tissue structures associated with the eye, such tissue structures providing a suitable environment in which a lacrimal implant can be used.

25 FIG. 3A illustrates an example of an isometric view of a lacrimal implant configured to be retained within a lacrimal punctum and associated canalicular anatomy, the lacrimal implant including a substantially perpendicular angled intersection between first and second implant body portions.

- FIG. 3B illustrates an example of a cross-sectional view of a lacrimal implant taken along a line parallel to a longitudinal axis of the implant, such as along line 3B-3B, and a dilation of an implant-receiving anatomical tissue structure.
- 5 FIG. 4 illustrates an example of a side view of a lacrimal implant configured to be retained within a lacrimal punctum and associated canalicular anatomy, the lacrimal implant including an integral dilator.
- 10 FIG. 5 illustrates an example of a schematic view of a lacrimal implant retained within a lacrimal punctum and associated canalicular anatomy, the lacrimal implant including at least one drug insert.
- 15 FIG. 6A illustrates an example of an isometric view of a lacrimal implant configured to be retained within a lacrimal punctum and associated canalicular anatomy, the lacrimal implant including a portion disposable within a lacrimal canaliculus ampulla.
- 20 FIG. 6B illustrates an example of a cross-sectional view of a lacrimal implant taken along a line parallel to a longitudinal axis of the implant, such as along line 6B-6B.
- FIG. 7A illustrates an example of an isometric view of a lacrimal implant configured to be retained within a lacrimal punctum and associated canalicular anatomy, the lacrimal implant including an annular graspable projection.
- 25 FIG. 7B illustrates an example of a cross-sectional view of a lacrimal implant taken along a line parallel to a longitudinal axis of the implant, such as along line 7B-7B.
- 30 FIG. 8A illustrates an example of an isometric view of a lacrimal implant configured to be retained within a lacrimal punctum and associated canalicular anatomy, the lacrimal implant

including a portion disposable within a lacrimal canaliculus ampulla and including an insertion-facilitating depression.

FIG. 8B illustrates an example of a cross-sectional view of a lacrimal implant taken along a line parallel to a longitudinal axis of the implant, such as along line 8B-8B.

FIG. 9 illustrates an example of side view of a lacrimal implant configured to be retained within a lacrimal punctum and associated canalicular anatomy, the lacrimal implant including an insertion-facilitating depression.

FIG. 10A illustrates an example of an isometric view of a lacrimal implant configured to be retained within a lacrimal punctum and associated canalicular anatomy, the lacrimal implant including a portion disposable within a lacrimal canaliculus ampulla.

FIG. 10B illustrates an example of a cross-sectional view of a lacrimal implant taken along a line parallel to a longitudinal axis of the implant, such as along line 10B-10B.

FIGS. 11-13 illustrate examples of a side view of various lacrimal implants configured to be retained within a lacrimal punctum and associated canalicular anatomy, each lacrimal implant including at least one intermediately-disposed retainment projection.

FIGS. 14-17 illustrate examples of an isometric view of various lacrimal implants configured to be retained within a lacrimal punctum and associated canalicular anatomy, each lacrimal implant including at least one intermediately-disposed retainment projection.

FIGS. 18-19 illustrate examples of a side view of lacrimal implants configured to be retained within a lacrimal punctum and associated canalicular anatomy, each lacrimal implant

including a non-perpendicular angled intersection between first and second implant body portions.

FIG. 20 illustrates an example of an isometric view of a lacrimal implant configured to be retained within a lacrimal punctum and associated canalicular anatomy, the lacrimal implant including one or more material cutouts.

FIGS. 21A-22B illustrate examples of a side view of lacrimal implants configured to be retained within a lacrimal punctum and associated canalicular anatomy, each lacrimal implant including one or more laterally extendable arms.

FIGS. 23A-23B illustrate examples of a side view of a lacrimal implant configured to be retained within a lacrimal punctum and associated canalicular anatomy, the lacrimal implant including an expandable retention element disposed around a portion of the implant body.

FIG. 24 illustrates an example of a schematic view of a lacrimal implant retained within a lacrimal punctum and associated canalicular anatomy.

FIGS. 25A-25B illustrate examples of an isomeric view of lacrimal implants configured to be retained within a lacrimal punctum and associated canalicular anatomy, each lacrimal implant including an implant body portion having a generally concave shape.

FIG. 26 illustrates an example of an isometric view of a lacrimal implant configured to be retained within a lacrimal punctum and associated canalicular anatomy, the lacrimal implant including an implant body portion having a generally convex shape.

FIG. 27 illustrates an example of a side view of a lacrimal implant configured to be retained within a lacrimal punctum and

associated canalicular anatomy, the lacrimal implant including an implant body portion having an undulating shape.

5 FIG. 28 illustrates an example of a side view of a lacrimal implant configured to be retained within a lacrimal punctum and associated canalicular anatomy, the lacrimal implant including at least one intermediately-disposed retainment projection.

10 FIGS. 29-32 illustrate examples of a side view of various lacrimal implants configured to be retained within a lacrimal punctum and associated canalicular anatomy, each lacrimal implant including a fluid swellable retention element.

15 FIG. 33 illustrates an example of a side view of a lacrimal implant configured to be retained within a lacrimal punctum and associated canalicular anatomy, the lacrimal implant including an expandable retention element.

20 FIGS. 34A-34B illustrate examples of a schematic view of lacrimal implants retained within a lacrimal punctum and associated canalicular anatomy, each lacrimal implant including an oriented graspable projection.

 FIGS. 35-38 illustrate examples of an isomeric view of various lacrimal implant proximal end portions, each proximal end portion including a graspable projection or void.

25 FIGS. 39A-39B illustrate examples of an isomeric view of drug inserts and a removal-facilitating filament.

 FIG. 40 illustrates an example of a method of manufacturing a lacrimal implant configured to be retained within a lacrimal punctum and associated canalicular anatomy.

DETAILED DESCRIPTION

In this patent document, lacrimal implants and related methods providing secure, wedgable retention within a lacrimal punctum and associated canaliculus of an eye are described. The lacrimal implants can comprise an implant body
5 configured for at least partial insertion through the lacrimal punctum and into the associated canaliculus. The implant body can include first and second portions, and can extend from a proximal end of the first portion defining a longitudinal proximal axis to a distal end of the second portion defining a longitudinal distal axis. The implant body can be configured such that, when implanted using an integral dilator,
10 an at least 45 degree angled intersection, for example, exists between the proximal axis and the distal axis. In this way, at least a portion of the implant body can be biased against at least a portion of the lacrimal canaliculus located at or more distal to a canalicular curvature, thereby retaining an implanted position of the lacrimal implant using anatomical structures. In various examples, the lacrimal implant can
15 further comprise a drug insert including a drug insert disposed in at least one of the first portion or the second portion of the implant body, providing a sustained release of a drug or other therapeutic agent to one or more of an eye, nasal passage or inner ear system.

FIGS. 1-2 illustrate examples of schematic views of anatomical tissue
20 structures associated with an eye 100. The anatomical tissue structures shown are suitable for treatment using the various lacrimal implants and methods discussed herein. As shown, the eye 100 is a spherical structure including a wall having three layers: an outer sclera 102, a middle choroid layer 104 and an inner retina 106. The sclera 102 includes a tough fibrous coating that protects the inner layers. It is
25 mostly white except for the transparent area at the front, commonly known as the cornea 108, which allows light to enter the eye 100.

The choroid layer 104, situated inside the sclera 102, contains many blood vessels and is modified at the front of the eye 100 as a pigmented iris 110. A biconvex lens 112 is situated just behind the pupil. A chamber 114 behind the lens
30 112 is filled with vitreous humour, a gelatinous substance. Anterior and posterior

chambers 116 are situated between the cornea 108 and iris 110, respectively and filled with aqueous humour. At the back of the eye 100 is the light-detecting retina 106.

5 The cornea 108 is an optically transparent tissue that conveys images to the back of the eye 100. It includes avascular tissue to which nutrients and oxygen are supplied via bathing with lacrimal fluid and aqueous humour as well as from blood vessels that line the junction between the cornea 108 and sclera 102. The cornea 108 includes a pathway for the permeation of drugs into the eye 100.

10 Turing to FIG. 2, other anatomical tissue structures associated with the eye 100 including the lacrimal drainage system, which includes a secretory system 230, a distributive system and an excretory system, are shown. The secretory system 230 comprises secretors that are stimulated by blinking and temperature change due to tear evaporation and reflex secretors that have an efferent parasympathetic nerve supply and secrete tears in response to physical or emotional stimulation. The
15 distributive system includes the eyelids 202 and the tear meniscus around the lid edges of an open eye, which spread tears over the ocular surface by blinking, thus reducing dry areas from developing.

The excretory part of the lacrimal drainage system includes, in order of flow drainage, the lacrimal puncta, the lacrimal canaliculi, the lacrimal sac 204 and the
20 lacrimal duct 206. From the lacrimal duct 206, tears and other flowable materials drain into a passage of the nasolacrimal system. The lacrimal canaliculi include an upper (superior) lacrimal canaliculus 208 and a lower (inferior) lacrimal canaliculus 210, which respectively terminate in an upper 212 and lower 214 lacrimal punctum. The upper 212 and lower 214 punctum are slightly elevated at the medial end of a
25 lid margin at the junction 216 of the ciliary and lacrimal portions near a conjunctival sac 218. The upper 212 and lower 214 punctum are generally round or slightly ovoid openings surrounded by a connective ring of tissue. Each of puncta 212, 214 leads into a vertical portion 220, 222 of their respective canaliculus before turning more horizontal at a canaliculus curvature 250 to join one another at the entrance of
30 the lacrimal sac 204. The canaliculi 208, 210 are generally tubular in shape and

lined by stratified squamous epithelium surrounded by elastic tissue, which permits them to be dilated. As shown, a lacrimal canaliculus ampulla **252** exists near an outer edge of each canaliculus curvature **250**.

FIG. **3A** illustrates an example of a lacrimal implant **300** that can be
5 insertable through a lacrimal punctum **212**, **214** and into the associated canaliculus **208**, **210** (FIG. **2**). The insertion of the lacrimal implant **300** through the lacrimal punctum **212**, **214** and into the associated canaliculus **208**, **210** can allow for one or more of: inhibition or blockage of tear flow therethrough (e.g., to treat dry eyes) or the sustained delivery of a drug or other therapeutic agent to an eye (e.g., to treat an
10 infection, inflammation, glaucoma or other ocular disease or disorder), a nasal passage (e.g., to treat a sinus or allergy disorder) or an inner ear system (e.g., to treat dizziness or a migraine).

As shown in this example, the lacrimal implant **300** can comprise an implant body **302** including first **304** and second **306** portions, and can extend from a
15 proximal end **308** of the first portion **304** to a distal end **310** of the second portion **306**. In various examples, the proximal end **308** can define a longitudinal proximal axis **312** and the distal end **310** can define a longitudinal distal axis **314**. The implant body **300** can be configured such that, when implanted within the lacrimal punctum and associated canaliculus, an at least 45 degree angled intersection **316**
20 exists between the proximal axis **312** and the distal axis **314** for biasing at least a portion of the implant body **302** against at least a portion of a lacrimal canaliculus **208**, **210** (FIG. **2**) located at or more distal to a canaliculus curvature **250** (FIG. **2**). In some examples, the implant body **302** can be configured such that the angled intersection **316** is between about 45 degrees and about 135 degrees. In this
25 example, the implant body **302** is configured such that the angled intersection **316** is about 90 degrees (i.e., the intersection **316** is about perpendicular). In various examples, a distal end **326** of the first portion **304** can be integral with the second portion **306** at or near a proximal end **328** of the second portion **306**.

In certain examples, the implant body **302** can include angularly disposed
30 cylindrical-like structures comprising one or both of a first cavity **318** disposed near

the proximal end 308 or a second cavity 320 disposed near the distal end 310. In this example, the first cavity 318 extends inward from the proximal end 308 of the first portion 304, and the second cavity 320 extends inward from the distal end 310 of the second portion 306. A first drug-releasing or other agent-releasing insert (e.g., drug core) 322 can be disposed in the first cavity 318 to provide a sustained drug or other therapeutic agent release to an eye, while a second drug-releasing or other agent-releasing insert (e.g., drug core) 324 can be disposed in the second cavity 320 to provide a sustained drug or other therapeutic agent release to a nasal passage or inner ear system, for example. An implant body septum 330 can be positioned between the first cavity 318 and the second cavity 320, and can be used to inhibit or prevent communication of a material (e.g., agent) between the first drug insert 322 and the second drug insert 324. In some examples, the implant body 302 is solid and does not include one or more cavities or other voids.

In some examples, the drug or other therapeutic agent release can occur, at least in part, via an exposed, non-sheath covered, surface of the drug inserts 322, 324. In some examples, by controlling geometry of the exposed surface, a predetermined drug or agent release rate can be achieved. For instance, the exposed surface can be constructed with a specific geometry or other technique appropriate to control the release rate of the drug or other therapeutic agent onto an eye 100, such as on an acute basis or on a chronic basis, between outpatient doctor visits. Further description regarding effective release rates of one or more drugs or other therapeutic agents from a drug insert 322, 324 can be found in commonly-owned DeJuan et al., U.S. Patent Application No. 11/695,545, entitled "NASOLACRIMAL DRAINAGE SYSTEM IMPLANTS FOR DRUG THERAPY".

In some examples, such as is shown in FIG. 3B, the exposed surface of the drug insert 322, 324 can be flush or slightly below the proximal end 308 of the first portion 304 or the distal end 310 of the second portion 306, respectively, such that the drug insert does not protrude outside of the implant body 302. In some examples, such as is shown in FIG. 4, the exposed surface of the first drug insert

322, for instance, can be positioned above the proximal end 308 such that the first drug insert 322 at least partially protrudes outside of the implant body 302.

The implant body 302 can include a graspable or other projection 332, such as one or more projections extending laterally at least partially from or around a proximal end 308 of the first implant body portion 304. In some examples, the graspable or other projection 332 can include a set of wings for use in inserting the lacrimal implant 300 into, or removing the lacrimal implant 300 from, an implanted position. The set of wings or other projection 332 can be configured without migration in mind, as the non-linear configuration of the implant body 302 can prevent implant 300 migration by assuming a size or shape of the canaliculus curvature 250 and optionally, the lacrimal canaliculus ampulla 252 (FIG. 2). In some examples, the graspable or other projection 332 can be configured to seat against or near the punctal opening 212, 214, such as for inhibiting or preventing the lacrimal implant 300 from passing completely within the lacrimal canaliculus 208, 210, or for providing tactile or visual feedback information to an implanting user, e.g., as to whether the implant is fully implanted.

As shown in FIGS. 34A-34B, and discuss below, the graspable or other projection 332 can extend laterally in a direction parallel to or away from an eye 100 when implanted. This may reduce irritation to the eye 100 as compared to a case in which a portion of the projection extends toward the eye 100. In addition, a lateral extension direction of the projection 332 from the proximal end 308 can be substantially the same as a lateral extension direction of the second implant body portion 306 relative to the distal end 326 of the first implant body portion 304, as shown in FIGS. 3A-3B, for example. This can also avoid extension toward the eye. The first drug insert 322 can partially extend through the region of the projection 332, such as to provide sustained release of a first drug or other therapeutic agent onto an eye.

In various examples, the implant body 302 can be molded using an elastic material, such as silicone, polyurethane or other urethane-based polymer or copolymer, NuSil (e.g., NuSil 4840 with 2% 6-4800) or an acrylic of a non-

biodegradable, partially biodegradable or biodegradable nature (i.e., erodeable within the body) allowing an implant body 302 configured such that, when implanted in a lacrimal canaliculus 208, 210, an angled intersection 316 exists between a proximal 312 and distal 314 axis to be formed. In some examples, the biodegradable elastic materials can include cross-linked polymers, such as poly (vinyl alcohol). In some examples, the implant body 302 can comprise a silicone/polyurethane co-polymer. Other co-polymers that can be used to form the implant body 302 include, but are not limited to, silicone/urethane, silicone/poly (ethylene glycol) (PEG), and silicone/2hydroxyethyl methacrylate (HEMA). As discussed in commonly-owned Utkhede et al., U.S. Patent Application No.

_____, Attorney Docket No. 2755.045US1, entitled "DRUG CORES FOR SUSTAINED RELEASE OF THERAPEUTIC AGENTS," filed even date herewith, urethane-based polymer and copolymer materials allow for a variety of processing methods and bond well to one another.

FIG. 3B illustrates an example of a cross-sectional view of the lacrimal implant 300 taken along a line parallel to a longitudinal axis of the implant, such as along line 3B-3B of FIG. 3A. As shown in FIG. 3B, the lacrimal implant 300 can include an implant body 302 including first 304 and second 306 portions, and can extend from a proximal end 308 of the first portion 304 to a distal end 310 of the second portion 306. In various examples, the proximal end 308 can define a longitudinal proximal axis 312 and the distal end 310 can define a longitudinal distal axis 314. The implant body 300 can be configured such that, when implanted, an at least 45 degree angled intersection 316 exists between the proximal axis 312 and the distal axis 314 for biasing at least a portion of the implant body 302 against at least a portion of a lacrimal canaliculus 208, 210 (FIG. 2) located at or more distal to a canaliculus curvature 250 (FIG. 2). In this example, the implant body 300 is configured such that the angled intersection 316 is approximately about 90 degrees.

In various examples, a distal end 326 of the first portion 304 can be integral with the second portion 306 at or near a proximal end 328 of the second portion 306. In some examples, the second portion 306 can include a length having a

magnitude less than four times a length of the first portion 304. In one example, the second portion 306 can include a length of less than about 10 millimeters and have a configured similar to that shown in FIG. 3B. In another example, the second portion 306 can include a length less than about 2 millimeters and have a
5 configuration similar to that shown in FIG. 24.

In various examples, the second portion 306 can comprise an integral dilator 350 to dilate anatomical tissue 352, such as one or both of a lacrimal punctum 212, 214 (FIG. 2) or associated canaliculus 208, 210, to a sufficient diameter as the lacrimal implant 300 is being implanted. In this way, the lacrimal implant 300 can
10 be implanted in various size ocular anatomies without the need for pre-dilation via a separate enlarging tool. The dilator 350 can be formed so as to not be traumatic to an inner lining of the punctum 212, 214 and the canaliculus 208, 210. In some examples, a lubricious coating disposed on, or impregnated in, an outer surface of the implant body 302 can be used to further aid insertion of the lacrimal implant 300
15 into the anatomical tissue 352. In one example, the lubricious coating can include a silicone lubricant.

As shown, the dilator 350 can generally narrow from a location near the proximal end 328 of the second portion 306 to the distal end 310 of the second portion 306, such as from a diameter of about 0.6 millimeters to a diameter of about
20 0.2 millimeters. In some examples, an outer surface slope of the dilator 350, as measured from the location near the proximal end 328 of the second portion 306 to the distal end 310 of the second portion 306, can be between about 1 degree and about 10 degrees (e.g., 2 degrees, 3 degrees, 4 degrees, or 5 degrees) with respect to the longitudinal distal axis 314. In some examples, the slope of the dilator 350 can
25 be less than 45 degrees with respect to the longitudinal distal axis 314. Among other factors, a determination of a desirable dilator 350 slope for a given implant situation can be made by balancing an implant body 302 strength desirable for implantation with a desire to have a soft, flexible and conforming implant body (e.g., to conform to a lacrimal canaliculus anatomy) upon implantation. In some

examples, a diameter of a dilator tip **354** can be between about 0.2 millimeters and about 0.5 millimeters.

In certain examples, the proximal end **328** of the second implant body portion **306** can include a retention element **356** configured to bias against at least a portion of a lacrimal canaliculus ampulla **252** (FIG. 2) when implanted. In this example, the retention element **356** projects proximally from the intersection between the first **304** and second **306** implant body portions, such as in an opposite direction as the extension of the dilator **350**. When present and implanted in the ampulla **252**, the retention element **356** can help secure a seated position of the graspable or other projection **332** against the punctal opening **212**, **214**

In certain examples, the implant body **302** includes a first cavity **318** disposed near the proximal end **308**. In this example, the first cavity **318** extends inward about 2 millimeters or less from the proximal end **308**, and houses a first drug-releasing or other agent-releasing drug insert **322** to provide a sustained drug or other agent release to an eye. In some examples, the drug insert **322** can include a plurality of therapeutic agent inclusions **360**, which can be distributed in a matrix **362**. In some examples, the inclusions **360** can comprise a concentrated (e.g., crystalline) form of the therapeutic agent. In some examples, the matrix **362** can comprise a silicone matrix or the like, and the distribution of inclusions **360** within the matrix can be substantially homogenous or non-homogeneous. In some examples, the agent inclusions **360** can include droplets of oil, such as Latanoprost oil. In still other examples, the agent inclusions **360** can comprise solid particles, such as Bimatoprost particles in crystalline form. In some examples, the drug insert **322** comprises a urethane-based (e.g., polyurethane) polymer or copolymer comprising therapeutic agent inclusions deliverable into the eye or surrounding tissues. The inclusions can be of many sizes and shapes. For instance, the inclusions can include microparticles having dimensions on the order of about 1 micrometer to about 100 micrometers. Further discussion of drug-releasing or other agent-releasing drug inserts can be found in commonly-owned Utkhede et al., U.S. Patent Application No. _____, Attorney Docket No.

2755.045US1, entitled "DRUG CORES FOR SUSTAINED RELEASE OF THERAPEUTIC AGENTS," filed even date herewith.

In various examples, the drug insert **322** can include a sheath body **366** disposed over at least a portion of the insert to define at least one insert exposed surface **368**. The exposed surface **368** can be located at or near the proximal end **308** of the implant body **302**, for example, thereby allowing direct contact with a tear or a tear film fluid and release of a drug or other therapeutic agent from the drug insert **322** over a sustained time period when the lacrimal implant **300** is inserted through the lacrimal punctum **212**, **214** and into the associated canaliculus **208**, **210**.

FIG. 4 illustrates an example of a side view of another integral dilator **450** of an implant body **402** second portion **406** of a lacrimal implant **400**. In this example, the dilator **450** narrows abruptly near a distal end **410** of the second portion **406**. As shown, an implant body first portion **404** can include a first cavity **418** disposed near the proximal end **408**. In this example, the first cavity **418** extends inward from the proximal end **408**, and houses a first drug-releasing or other agent-releasing drug insert **422** to provide a sustained drug or other therapeutic agent release to an eye, for instance. In some examples, the drug or other therapeutic agent can be released to an eye via an exposed, non-sheath covered surface **468** of the drug insert **422**. In this example, the exposed surface **468** of the drug insert **422** is positioned above the proximal end **408** such that the drug insert **422** at least partially protrudes outside of the implant body **402**.

In various examples, the outer surface **482** of the implant body **402** can be formed, or surface treated to be, generally smooth to inhibit bacteria from attaching to the lacrimal implant **400** and incubating. The generally smooth outer surface **482** can also prevent damage to the inner lining of the receiving anatomical tissue, such as a lacrimal punctum **212**, **214** (FIG. 2) or associated canaliculus **208**, **210** (FIG. 2), during implantation. As further discussed in commonly-owned Rapacki et al., U.S. Patent Application No. _____, Attorney Docket No. 2755.036US1, entitled "SURFACE TREATMENT OF IMPLANTS AND RELATED METHODS," filed

even date herewith, the outer surface **482** of the implant body **402** can be surface treated to be generally smooth via a polishing process. The polishing process can include causing a molded implant body **402** to be impacted with polishing media during an ongoing period of time in which the body **402** is in an enlarged, swelled state. This can smooth one or more surfaces or edges of the implant body **402**. In various examples, the polishing media can include at least some granules that are greater than about 3 millimeters in diameter.

In various examples, an antimicrobial coating **484** can be disposed on or impregnated in at least a portion of the outer surface **482** to further prevent bacteria growth on the implant body **402**. In some examples, the antimicrobial coating **484** can include an agent selected from the group consisting of 2-bromo-2-nitropropane-1,3-diol, 5-bromo-5-nitro-1,3-dioxane, 7-ethyl bicycloodiazolidine, benzalkonium chloride, benzethonium chloride, benzoic acid, benzyl alcohol, boric acid, bronopol, cetylpyridinium chloride, chlorhexidine digluconate, chloroacetamide, chlorobutanol, chloromethyl isothiazolinone and methyl isothiazoline, dimethoxane, dimethyl oxazolidine, dimethyl hydroxymethyl pyrazole, chloroxylonol, dehydroacetic acid, diazolidinyl urea, dichlorobenzyl alcohol, DMDM hydantoin, ethyl alcohol, formaldehyde, glutaraldehyde, hexachlorophene, hexetidine, hexamethylenetetramine, imidazolidinyl urea, iodopropynyl butylcarbamate, isothiazolinones, methenammonium chloride, methyldibromo glutaronitrile, MDM hydantoin, minocycline, ortho phenylphenol, p-chloro-m-cresol, parabens (butylparaben, ethylparaben, methylparaben), phenethyl alcohol, phenoxyethanol, piroctane olamine, polyaminopropyl biguanide, polymethoxy bicyclic oxazolidine, polyoxymethylene, polyquaternium-42, potassium benzoate, potassium sorbate, propionic acid, quaternium-15, rifampin, salicylic acid, selenium disulfide, sodium borate, sodium iodate, sodium hydroxymethylglycinate, sodium propionate, sodium pyrithione, sorbic acid, thimerosal, triclosan, triclocarban, undecylenic acid, zinc phenosulfonate, and zinc pyrithione. In some examples, the antimicrobial coating **484** can include a material selected from the group consisting of silver lactate, silver phosphate, silver citrate, silver acetate, silver benzoate, silver chloride, silver iodide,

silver iodate, silver nitrate, silver sulfadiazine, silver palmitate, or one or more mixtures thereof. In some examples, the antimicrobial coating **484** can include at least one of an antibiotic or an antiseptic. For instance, the antimicrobial coating **484** can include a temporary anesthetic lasting, on average, between a few hours and a day. In still other examples, the antimicrobial coating **484** can include a drug or other therapeutic agent used to treat an underlying disease, such as a bolus, for immediate effect.

FIG. 5 illustrates an example of a schematic view of a lacrimal implant, such as the lacrimal implant **300** shown in FIG. 3, implanted in a lower lacrimal punctum **214** and associated canaliculus **210**. In some examples, a lacrimal implant **300** can be implanted in an upper lacrimal punctum **212** and canaliculus **208**. As discussed above, the lacrimal implant **300** can comprise an implant body **302** including first **304** and second **306** portions. In various examples, the implant body **302** can be configured such that, when implanted, at least a portion of the implant body **302** is biased against at least a portion of the lacrimal canaliculus **210** located at or more distal to a canaliculus curvature **250** to securely retain an implanted position of the implant **300**. As shown, the first portion **304** can be configured to be inserted through the lacrimal punctum **214** and into the associated canaliculus **210** and rest between the punctal opening and a lacrimal canaliculus ampulla **252**, while the second portion **306** can be configured to insert through the lacrimal punctum **214** and into the canaliculus **210** and rest between the ampulla **252** and the lacrimal sac **204**. In certain examples, a retention element **356** projecting from a proximal end of the second portion **306** can be configured to bias into and against at least a portion of the ampulla **252** when implanted. In various examples, the first **304** and second **306** portions can be configured to bend, stretch or collapse, as desired, to maintain an adequate anatomical implanted fit without unduly stretching ocular anatomy.

In certain examples, to further secure an implant **300** within the lacrimal punctum **214** and canaliculus **210** or to make the implant body **302** adjustable in size, a hydrogel or other fluid swellable material can be disposed (e.g., coated) on an outer surface portion of the implant body **302**. The fluid swellable material can

effectively expand an outer surface diameter portion of the implant body 302 when implanted. In certain examples, the outer surface of the implant body 302 can include longitudinal channels or grooves or coatings of a wicking material so as to allow fluid flow around the implant body 302. Using one or a combination of these techniques, a lacrimal implant 300 can be configured to completely occlude or only partially occlude the lacrimal canaliculus 208, 210 when implanted therein. For instance, using the longitudinal channels or grooves in one or both of the first 304 or second 306 portions of the implant body 302 can allow diminished volumes or tear drainage can occur, potentially facilitating the release of a drug or other therapeutic agent from a drug insert.

Forceps or another insertion tool can be used to implant the lacrimal implant 300 through the lacrimal punctum 212, 214 and into the associated canaliculus 208, 210. In some examples, an insertion tool as discussed in commonly-owned De Juan, et al., U.S. Patent Application No. _____, Attorney Docket No. 2755.018US1, entitled "INSERTION AND EXTRACTION TOOLS FOR LACRIMAL IMPLANTS," filed even date herewith, can be used to implant the lacrimal implant 300. In various examples, the second portion 306 of the implant body 302 can be advanced into the depth of the lacrimal canaliculus 208, 210 by manipulation of the inserter tool until a graspable or other projection 332, if present, can be seated against the punctal opening 212, 214. When it is desired to remove the lacrimal implant 300, the projection 332 can be grasped with the forceps, for example, and withdrawn from the punctal opening 212, 214.

In certain examples, the implant body 302 can include one or both of a first cavity 318 disposed near the proximal end 308 or a second cavity 320 disposed near the distal end 310. In this example, the first cavity 318 extends inward from the proximal end 308 of the first portion 304, and the second cavity 320 extends inward from the distal end 310 of the second portion 306. A first drug-releasing or other agent-releasing drug insert 322 can be disposed in the first cavity 318 to provide a sustained drug or other therapeutic agent release to the eye (e.g., to treat an infection, inflammation, glaucoma or other ocular disease or disorder), while a

second drug-releasing or other agent-releasing drug insert **324** can be disposed in the second cavity **320** to provide a sustained drug or other therapeutic agent release to the nasal passage (e.g., to treat a sinus or allergy disorder) or inner ear system (e.g., to treat dizziness or a migraine), for example.

5 FIGS. **6A-6B** illustrate an example of another lacrimal implant **600** that is insertable through a lacrimal punctum **212, 214** and into the associated canaliculus **208, 210** (FIG. **2**). In this example, the lacrimal implant **600** can comprises an implant body **602** including first **604** and second **606** portions, and can extend from a proximal end **608** of the first portion **604** to a distal end **610** of the second portion **606**. The proximal end **608** can define a longitudinal proximal axis **612** and the distal end **610** can define a longitudinal distal axis **614**. The implant body **600** can be configured such that, when implanted, an angled intersection of approximately 90 degrees exists between the proximal axis **612** and the distal axis **614** for biasing at least a portion of the implant body against at least a portion of the lacrimal
10 canaliculus **208, 210** (FIG. **2**) located at or more distal to a canaliculus curvature **250** (FIG. **2**).

In this example, a proximal end **628** of the second implant body portion **606** can include a retention element **656** configured to bias against at least a portion of a lacrimal canaliculus ampulla **252** (FIG. **2**) when implanted. In this example, the
20 implant body **602** includes a first cavity **618**, configured to receive a first drug-releasing or other agent-releasing drug insert, disposed near the proximal end **608** of the first implant body portion **604**. Also in this example, the implant body **602** can include a graspable or other projection **632**, such as a set of wings having a combined length of about 1 millimeter, for example, and extending laterally from
25 the proximal end **608**.

FIG. **6B** illustrates an example of a cross-sectional view of the lacrimal implant **600** taken along a line parallel to a longitudinal axis of the implant, such as along line **6B-6B** of FIG. **6A**. As shown in FIG. **6B**, a distal end **626** of the first portion **604** can be integral with the second portion **606** at or near a proximal end
30 **628** of the second portion **606**. In various examples, the second portion **606** can

include a longitudinal length, as measured from the proximal axis **612** to the distal end **610**, having a magnitude less than four times a longitudinal length of the first portion **604**, as measured from the proximal end **608** to the distal axis **614**. In some examples, the first portion can include a longitudinal length of about 1.54
5 millimeters and the second portion can include a longitudinal length of between about 4.5 millimeters to about 5.42 millimeters.

In various examples, the second portion **606** can comprise an integral dilator **650** to dilate anatomical tissue, such as one or both of the lacrimal punctum **212**, **214** (FIG. 2) or associated canaliculus **208**, **210**, to a sufficient diameter as the
10 lacrimal implant **600** is being implanted. In some examples, the second portion **606** tapers from a diameter of the proximal end of between about 0.50 millimeters to about 0.75 millimeters to a dilator tip **654** diameter of about 0.36 millimeters.

FIGS. 7A-7B illustrate an example of another lacrimal implant **700** that is insertable through a lacrimal punctum **212**, **214** and into the associated canaliculus
15 **208**, **210** (FIG. 2). In this example, the lacrimal implant **700** can comprises an implant body **702** including first **704** and second **706** portions, and can extend from a proximal end **708** of the first portion **704** to a distal end **710** of the second portion **706**. The proximal end **708** can define a longitudinal proximal axis **712** and the distal end **710** can define a longitudinal distal axis **714**. The implant body **700** can
20 be configured such that, when implanted, an angled intersection of approximately 90 degrees exists between the proximal axis **712** and the distal axis **714** for biasing at least a portion of the implant body against at least a portion of the lacrimal canaliculus **208**, **210** (FIG. 2) located at or more distal to a canaliculus curvature **250** (FIG. 2). As shown in the example of FIG. 7A, a smooth transition can exist
25 between the first **704** and second **706** portions.

In this example, the implant body **702** includes a first cavity **718** configured to receive a first drug-releasing or other agent-releasing drug insert, disposed near the proximal end **708** of the first implant body portion **704**. Also in this example, the implant body **702** can include a graspable or other projection **732**, such as an
30 annular projection extending laterally from, and completely around, the proximal

end **708**. In some examples, the graspable or other projection **732** includes a partially trimmed projection having a trimmed width of about 0.75 millimeters and extending varying amounts around the proximal end **708**.

FIG. **7B** illustrates an example of a cross-sectional view of the lacrimal implant **700** taken along a line parallel to a longitudinal axis of the implant, such as along line **7B-7B** of FIG. **7A**. As shown in FIG. **7B**, a distal end **726** of the first portion **704** can be integral with the second portion **706** at or near a proximal end **728** of the second portion **706**. In various examples, the second portion **706** can include a longitudinal length, as measured from the proximal axis **712** to the distal end **710**, having a magnitude less than four times a longitudinal length of the first portion **704**, as measured from the proximal end **708** to the distal axis **714**. In some examples, the first portion can include a longitudinal length of about 1.5 millimeters and the second portion can include a longitudinal length of about 5 millimeters.

In various examples, the second portion **706** can comprise an integral dilator **750** to dilate anatomical tissue, such as one or both of the lacrimal punctum **212**, **214** (FIG. **2**) or associated canaliculus **208**, **210**, to a sufficient diameter as the lacrimal implant **700** is being implanted. In some examples, the second portion **706** tapers from a diameter of the proximal end of about 0.46 millimeters to a dilator tip **754** diameter of about 0.36 millimeters.

FIGS. **8A-8B** illustrate an example of another lacrimal implant **800** that is insertable through a lacrimal punctum **212**, **214** and into the associated canaliculus **208**, **210** (FIG. **2**). In this example, the lacrimal implant **800** can comprises an implant body **802** including first **804** and second **806** portions, and can extend from a proximal end **808** of the first portion **804** to a distal end **810** of the second portion **806**. The proximal end **808** can define a longitudinal proximal axis **812** and the distal end **810** can define a longitudinal distal axis **814**. The implant body **800** can be configured such that, when implanted, an angled intersection of approximately 90 degrees exists between the proximal axis **812** and the distal axis **814** for biasing at least a portion of the implant body against at least a portion of the lacrimal

canaliculus **208, 210** (FIG. 2) located at or more distal to a canaliculus curvature **250** (FIG. 2).

In this example, a proximal end **828** of the second implant body portion **806** can include a retention element **856** configured to bias against at least a portion of a lacrimal canaliculus ampulla **252** (FIG. 2) when implanted. The retention element **856** can include an insertion-facilitating depression **875** or other gripping means to aid in one or both of implant insertion or removal. In this example, the implant body **802** includes a first cavity **818** configured to receive a first drug-releasing or other agent-releasing drug insert, disposed near the proximal end **808** of the first implant body portion **804**. Also in this example, the implant body **802** can include a graspable or other projection **832**, such as an annular projection extending laterally from, and completely around, the proximal end **808**. In some examples, the graspable or other projection **832** includes a partially trimmed projection extending varying amounts around the proximal end **808**.

FIG. **8B** illustrates an example of a cross-sectional view of the lacrimal implant **800** taken along a line parallel to a longitudinal axis of the implant, such as along line **8B-8B** of FIG. **8A**. As shown in FIG. **8B**, a distal end **826** of the first portion **804** can be integral with the second portion **806** at or near the proximal end **828** of the second portion **806**. In various examples, the second portion **806** can include a longitudinal length, as measured from the proximal axis **812** to the distal end **810**, having a magnitude less than four times a longitudinal length of the first portion **804**, as measured from the proximal end **808** to the distal axis **814**. In some examples, the first portion can include a longitudinal length of between about 1.725 millimeters to about 1.77 millimeters and the second portion can include a longitudinal length of between about 4.77 millimeters to about 5 millimeters.

In various examples, the second portion **806** can comprise an integral dilator **850** to dilate anatomical tissue, such as one or both of the lacrimal punctum **212, 214** (FIG. 2) or associated canaliculus **208, 210**, to a sufficient diameter as the lacrimal implant **800** is being implanted. In some examples, the second portion **806**

tapers from a diameter of the proximal end **828** of about 0.46 millimeters to a dilator tip **854** diameter of about 0.36 millimeters.

FIG. 9 illustrates an example of another lacrimal implant **900** that is insertable through a lacrimal punctum **212, 214** and into the associated canaliculus **208, 210** (FIG. 2). In this example, the lacrimal implant **900** can comprises an implant body **902** including first **904** and second **906** portions, and can extend from a proximal end **908** of the first portion **904** to a distal end **910** of the second portion **906**. The proximal end **908** can define a longitudinal proximal axis **912** and the distal end **910** can define a longitudinal distal axis **914**. The implant body **900** can be configured such that, when implanted, an angled intersection of approximately 90 degrees exists between the proximal axis **912** and the distal axis **914** for biasing at least a portion of the implant body against at least a portion of the lacrimal canaliculus **208, 210** (FIG. 2) located at or more distal to a canaliculus curvature **250** (FIG. 2).

As shown, a smooth transition can exist between the first **904** and second **906** portions. In this example, the smooth transition can include an insertion-facilitating depression **975** or other gripping means to aid in one or both of implant insertion or removal. Also in this example, the implant body **902** can include a graspable or other projection **932**, such as an annular projection extending laterally from, and completely around, the proximal end **908**. In some examples, the graspable or other projection **932** includes a partially trimmed projection extending varying amounts around the proximal end **908**.

FIGS. 10A-10B illustrate an example of another lacrimal implant **1000** that is insertable through a lacrimal punctum **212, 214** and into the associated canaliculus **208, 210** (FIG. 2). In this example, the lacrimal implant **1000** can comprises an implant body **1002** including first **1004** and second **1006** portions, and can extend from a proximal end **1008** of the first portion **1004** to a distal end **1010** of the second portion **1006**. The proximal end **1008** can define a longitudinal proximal axis **1012** and the distal end **1010** can define a longitudinal distal axis **1014**. The implant body **1000** can be configured such that, when implanted, an

angled intersection of approximately 90 degrees exists between the proximal axis 1012 and the distal axis 1014 for biasing at least a portion of the implant body against at least a portion of the lacrimal canaliculus 208, 210 (FIG. 2) located at or more distal to a canaliculus curvature 250 (FIG. 2).

5 In this example, a proximal end 1028 of the second implant body portion 1006 can include a retention element 1056 configured to bias against at least a portion of a lacrimal canaliculus ampulla 252 (FIG. 2) when implanted. The retention element 1056 can include an insertion-facilitating depression 1075 or other gripping means to aid in one or both of implant insertion or removal. In this
10 example, the implant body 1002 includes a first cavity 1018 configured to receive a first drug-releasing or other agent-releasing drug insert, disposed near the proximal end 1008 of the first implant body portion 1004. Also in this example, the implant body 1002 can include a graspable or other projection 1032, such as an annular projection having a diameter of about 1.3 millimeters extending laterally from, and
15 completely around, the proximal end 1008. In some examples, the graspable or other projection 1032 includes a partially trimmed projection extending varying amounts around the proximal end 1008.

FIG. 10B illustrates an example of a cross-sectional view of the lacrimal implant 1000 taken along a line parallel to a longitudinal axis of the implant, such as
20 along line 10B-10B of FIG. 10A. As shown in FIG. 10B, a distal end 1026 of the first portion 1004 can be integral with the second portion 1006 at or near a proximal end 1028 of the second portion 1006. In various examples, the second portion 1006 can include a longitudinal length, as measured from the proximal axis 1012 to the distal end 1010, having a magnitude less than four times a longitudinal length of the
25 first portion 1004, as measured from the proximal end 1008 to the distal axis 1014. In some examples, the first portion can include a longitudinal length of about 1.5 millimeters and the second portion can include a longitudinal length of about 5 millimeters.

30 In various examples, the second portion 1006 can comprise an integral dilator 1050 to dilate anatomical tissue, such as one or both of the lacrimal punctum

212, 214 (FIG. 2) or associated canaliculus 208, 210, to a sufficient diameter as the lacrimal implant 1000 is being implanted. In some examples, the second portion 1006 tapers from a proximal end 1028 diameter of about 0.46 millimeters to a dilator tip 1054 diameter of about 0.36 millimeters.

5 FIGS. 11-17 illustrate examples of other lacrimal implants 1100, 1200, 1300, 1400, 1500, 1600, 1700 that are insertable through a lacrimal punctum 212, 214 and into the associated canaliculus 208, 210 (FIG. 2). In these examples, each lacrimal implant 1100, 1200, 1300, 1400, 1500, 1600, 1700 can comprises an implant body 1102, 1202, 1302, 1402, 1502, 1602, 1702 including first 1104, 1204, 1304, 1404, 1504, 1604, 1704 and second 1106, 1206, 1306, 1406, 1506, 1606, 1706 portions, and can extend from a proximal end 1108, 1208, 1308, 1408, 1508, 1608, 1708 of the first portion 1104, 1204, 1304, 1404, 1504, 1604, 1704 to a distal end 1110, 1210, 1310, 1410, 1510, 1610, 1710 of the second portion 1106, 1206, 1306, 1406, 1506, 1606, 1706. Each implant body 1102, 1202, 1302, 1402, 1502, 1602, 1702 can include at least one intermediately-disposed retainment projection 1192, 1292, 1392, 1492, 1592, 1692, 1792 to potentially further secure an implanted position of the lacrimal implants. The intermediately-disposed retainment projections 1192, 1292, 1392, 1492, 1592, 1692, 1792 can be positioned on one or both of the first 1104, 1204, 1304, 1404, 1504, 1604, 1704 or second 1106, 1206, 1306, 1406, 1506, 1606, 1706 implant body portions, and can take the form of annular, semi-annular, column-like or barrel-like projection. The intermediately-disposed retainment projections 1192, 1292, 1392, 1492, 1592, 1692, 1792 can include a cross-sectional size greater than an adjacent implant body portion and can slightly deform a portion of a canalicular wall to provide the added securement.

25 It is believed that the occlusion of the lower lacrimal canaliculus 210, for example, by a lacrimal implant may cause back pressure to build-up within the canaliculus 210, thereby urging the implant from an implanted position. It is thought that this back pressure could, for example, occur during a blink (where tears are being pumped from an anterior surface of the eye down a drainage system) or a
30 sneeze (where pressure is emanating up from the pulmonary system). Accordingly,

one of more of the additional retention features now shown in the form of at least one intermediately-disposed retainment projection **1192, 1292, 1392, 1492, 1592, 1692, 1792** may be used to prevent implant migration and further secure an implanted lacrimal implant position. These additional retention features can be
5 designed to prevent migration in the proximal direction while not increasing implant implantation difficulty.

FIGS. **18-19** illustrate examples of other lacrimal implants **1800, 1900** that are insertable through a lacrimal punctum **212, 214** and into the associated canaliculus **208, 210** (FIG. **2**). In these examples, each lacrimal implant **1800, 1900**
10 can comprise an implant body **1802, 1902** including first **1804, 1904** and second **1806, 1906** portions, and can extend from a proximal end **1808, 1908** of the first portion **1804, 1904** to a distal end **1810, 1910** of the second portion **1806, 1906**. As shown, an intermediate portion **1896, 1996** of each implant body **1802, 1902** can be angled relative to one or both of the first **1804, 1904** or second **1806, 1906** implant
15 body portions to potentially further secure an implanted position of the lacrimal implants.

It is believed that the angling of the intermediate portion **1896, 1996** may help capture the anatomy of the lacrimal punctum **212, 214** and canaliculus **208, 210** to keep the lacrimal implants **1800, 1900** in an implanted position, such as via a
20 directional force applied by the angling against the lacrimal canaliculus. This directional force can be designed to continuously urge a feedback or other projection **1832, 1932** flush with the punctum **212, 214**.

FIG. **20** illustrates an example of another lacrimal implant **2000** that is insertable through a lacrimal punctum **212, 214** and into the associated canaliculus
25 **208, 210** (FIG. **2**). In this example, the lacrimal implant **2000** can comprises an implant body **2002** including first **2004** and second **2006** portions, and can extend from a proximal end **2008** of the first portion **2004** to a distal end **2010** of the second portion **2006**. The proximal end **2008** can define a longitudinal proximal axis **2012** and the distal end **2010** can define a longitudinal distal axis **2014**. The
30 implant body **2000** can be configured such that, when implanted, an angled

intersection of approximately 90 degrees exists between the proximal axis **2012** and the distal axis **2014** for biasing at least a portion of the implant body against at least a portion of a lacrimal canaliculus **208, 210** (FIG. 2) located at or more distal to a canaliculus curvature **250** (FIG. 2). In various examples, a distal end **2026** of the first portion **2004** can be integral with the second portion **2006** at or near a proximal end **2028** of the second portion **2006**.

In this example, one or more material cutouts **2080** are made in an outer surface of the implant body **2002**. As a result, the angled intersection between the proximal axis **2012** and the distal axis **2014** can become more linearly aligned during implant, as shown in phantom, to facilitate insertion through the lacrimal punctum **212, 214** and into the associated canaliculus **208, 210**.

FIGS. **21A-21B** and **22A-22B** illustrate examples of a side view of other lacrimal implants **2100, 2200** that are insertable through a lacrimal punctum **212, 214** and into the associated canaliculus **208, 210** (FIG. 2). In these examples, each lacrimal implant **2100, 2200** can comprises an implant body **2102, 2202** including first **2104, 2204** and second **2106, 2206** portions, and can extend from a proximal end **2108, 2208** of the first portion **2104, 2204** to a distal end **2110, 2210** of the second portion **2106, 2206**. Each second portion **2106, 2206** can include one or more arm members **2170, 2270** movable between a first configuration, in which the one or more arm members **2170, 2270** are adjacent the implant body, and a second configuration, in which the one or more arm members **2170, 2270** laterally extend from a side of the implant body. In the first configuration, the one or more arm members **2170, 2270** facilitate insertion of the lacrimal implant through the lacrimal punctum **212, 214** and into the associated canaliculus **208, 210** by providing a narrow profile. In the second configuration, the one or more arm members **2170, 2270** laterally extend to fill at least one of a lacrimal canaliculus ampulla **252** (FIG. 2) or the canaliculus **208, 210** when implanted. Optionally, the one or more arm members **2170, 2270** can include a fluid swellable material, such as hydrogel, to further secure an implanted lacrimal implant within the lacrimal ampulla **252** or canaliculus **208, 210** when hydrated.

In some examples, the one or more arm members **2170**, **2270** can be incorporated into a mold that is also used to form the implant body **2102**, **2202**. The one or more arm members **2170**, **2270** can alternatively be attached by molding or gluing onto an existing implant body **2102**, **2202**. Different thicknesses and shapes for the one or more arm members **2170**, **2270** can be employed for different stiffness and securing/removal characteristics. Beyond hydrogel, the one or more arm members **2170**, **2270** can be made of other materials, such as those used for the haptics on the intraocular lenses or the like.

FIG. **23A-23B** illustrate an example of a side view of another lacrimal implant **2300** that is insertable through a lacrimal punctum **212**, **214** and into the associated canaliculus **208**, **210** (FIG. **2**). In this example, the lacrimal implant **2300** can comprises an implant body **2302** including first **2304** and second **2306** portions, and can extend from a proximal end **2308** of the first portion **2304** to a distal end **2310** of the second portion **2306**. The second portion **2306** can be surrounded, at least in part, by an expandable retention element (e.g., an inflatable balloon) **2372**, which is configured to bias the second portion **2306** away from a lacrimal canaliculus wall upon expansion.

In some examples, the expandable retention element **2372** contains or can be inflated by an agent to be delivered to a tissue of the eye or nasolacrimal system. In some examples, the expandable retention element **2372** can employ one or more balloons which are separate from any drug insert or other agent retaining structure. The one or more balloons may optionally be similar to those used on balloon catheters, with an inflation lumen or the like optionally being included in an implant insertion tool so as to allow controlled inflation of the balloon. In such an example, the lacrimal implant **2300** may be inserted with the balloons deflated, as shown in FIG. **23A**. Once the lacrimal implant **2300** is in place, the balloons can then inflated to secure an implanted position of the implant, as shown in FIG. **23B**.

The balloons can also be deflatable to make removal of the lacrimal implant **2300** easier. The balloons can optionally partially or substantially conform to variations in the size and shape of the canaliculus **208**, **210**. Alternative examples of

balloons may be inflated by swelling of a material disposed within the balloon, such as swelling of a hydrogel by absorption of water through perforations or openings in the balloon. The one or more balloons can be annular structures disposed around the supporting implant body, or may be disposed eccentrically about an axis of the implant body. As illustrated in **FIG. 23B**, the balloons may be disposed sufficiently distal to reside within or adjacent a horizontal portion of the tear drainage duct, within or adjacent a lacrimal ampulla of the tear drainage system, or the like. Alternative examples can include one or more balloons which are more proximal.

FIG. 24 illustrates an example of a schematic view of another lacrimal implant **2400** implanted through a lower lacrimal punctum **214** and into the associated canaliculus **210**. The lacrimal implant **2400** can comprise an implant body **2402** including first **2404** and second **2406** portions. In various examples, the implant body **2402** can be configured such that, when implanted, at least a portion of the implant body **2402** is biased against at least a portion of the lacrimal canaliculus **210** located at or more distal to a canaliculus curvature **250** to securely retain an implanted position of the implant **2400**. In this example, the second portion **2406** includes a longitudinal length less than about 2 millimeters, such as a size greater than a diameter of the first portion **2404**, but less than 2 millimeters. Also in this example, the implant body **2402** can include a graspable or other projection **2432**, such as extending laterally at least partially around a proximal end of the first implant body portion **2404**.

FIGS. 25A-25B illustrate examples of another lacrimal implant **2500** that is insertable through a lacrimal punctum **212**, **214** and into the associated canaliculus **208**, **210** (**FIG. 2**). In these examples, the lacrimal implant **2500** can comprise an implant body **2502** including first **2504** and second **2506** portions, and can extend from a proximal end **2508** of the first portion **2504** to a distal end **2510** of the second portion **2506**. The implant body can include a general shape, which can generally match the anatomical features of a canaliculus **208**, **210** to provide patient comfort and secure retainment, for example. The proximal end **2508** can define a longitudinal proximal axis **2512** and the distal end **2510** can define a longitudinal

distal axis **2514**. The implant body **2502** can be configured such that, when implanted, an angled intersection of between 45-90 degrees exists between the proximal axis **2512** and the distal axis **2514** such as for biasing at least a portion of the implant body **2502** against at least a portion of a lacrimal canaliculus **208**, **210** (FIG. 2) located at or more distal to a canaliculus curvature **250** (FIG. 2).

In the examples of FIGS. **25A-25B**, the implant body **2502** includes both of a first cavity **2518** disposed near the proximal end **2508** and a second cavity **2520** disposed near the distal end **2510**. The first cavity **2518** extends inward from the proximal end **2508** of the first portion **2504**, and the second cavity **2520** extends inward from the distal end **2510** of the second portion **2506**. A first drug-releasing or other agent-releasing drug insert can be disposed in the first cavity **2518** to provide a sustained drug or other therapeutic agent release to an eye, while a second drug-releasing or other agent-releasing drug insert can be disposed in the second cavity **2520** to provide a sustained drug or other therapeutic agent release to a nasal passage or inner ear system, for example. In some examples, the first cavity **2518** can extend inward from the proximal end **2508** of the first portion **2504** to a position near the distal end **2510** of the second portion **2506**, such as is shown in FIG. **26**, and is filled with a first drug-releasing or other agent-releasing drug insert. In some examples, the second cavity **2520** can extend inward from the distal end **2510** of the second portion **2506** to a position near the proximal end **2508** of the first portion **2504** and is filled with a second drug-releasing or other agent-releasing drug insert.

In certain examples, the second portion **2506** comprises an integral dilator **2550** to dilate anatomical tissue, such one or both of the lacrimal punctum **212**, **214** or canaliculus **208**, **210**, to a sufficient diameter as the lacrimal implant **2500** is being implanted. In this way, the lacrimal implant **2500** can be implanted in various size ocular anatomies without the need for pre-dilation via a separate enlarging tool. In these examples, the integral dilator **2550** includes a generally concave shape related to the first portion **2504**. In some examples, the concave shape includes a radius less than a radius of the canaliculus curvature **250**. In some examples, the concave shape includes a radius substantially the same as the radius of the

canaliculus curvature **250**. As shown in the example of FIG. **25B**, a smooth transition can exist between the first **2504** and second **2506** portions.

In certain examples, a proximal end **2528** of the second implant body portion **2506** can include a retention element **2556** configured to bias against at least a
5 portion of a lacrimal canaliculus ampulla **252** (FIG. **2**) when implanted. In the example of FIG. **25A**, the retention element **2556** projects proximally from the intersection between the first **2504** and second **2506** implant body portions.

FIG. **26** illustrates an example of another lacrimal implant **2600** that is insertable through a lacrimal punctum **212**, **214** and into the associated canaliculus
10 **208**, **210** (FIG. **2**). In this example, the lacrimal implant **2600** comprises an implant body **2602** including first **2604** and second **2606** portions, and extends from a proximal end **2608** of the first portion **2604** to a distal end **2610** of the second portion **2606**. The proximal end **2608** can define a longitudinal proximal axis **2612** and the distal end **2610** can define a longitudinal distal axis **2614**. The implant body
15 **2600** can be configured such that, when implanted, an angled intersection of between 90-135 degrees exists between the proximal axis **2612** and the distal axis **2614** for biasing at least a portion of the implant body against at least a portion of a lacrimal canaliculus **208**, **210** located at or more distal to a canaliculus curvature **250** (FIG. **2**).

In certain examples, the implant body **2602** can include a first cavity **2618**
20 disposed near the proximal end **2608**. In this example, the first cavity **2618** extends inward from the proximal end **2608** of the first portion **2604** to a position near the distal end **2610** of the second portion **2606**. A first drug-releasing or other agent-releasing drug insert having a volume between about 0.2 cubic centimeters to about
25 0.25 cubic centimeters, for example, can be disposed in the first cavity **2618** to provide a extended sustained drug or other therapeutic agent release to an eye.

In certain examples, the second portion **2606** comprises an integral dilator **2650** to dilate anatomical tissue, such one or both of the lacrimal punctum **212**, **214** or canaliculus **208**, **210**, to a sufficient diameter as the lacrimal implant **2600** is
30 being implanted. In this way, the lacrimal implant **2600** can be implanted in various

size ocular anatomies without the need for pre-dilation via a separate enlarging tool. In this example, the dilator **2650** includes a generally convex shape relative to the first portion **2604**. In some examples, the convex shape includes a radius less than a radius of the canaliculus curvature **250**. In some examples, the convex shape
5 includes a radius substantially the same as the radius of the canaliculus curvature **250**.

In certain examples, a proximal end **2628** of the second implant body portion **2606** can include a retention element **2656** configured to bias against at least a portion of a lacrimal canaliculus ampulla **252** (FIG. 2) when implanted. In this
10 example, the retention element **2656** projects proximally from the intersection between the first **2604** and second **2606** implant body portions. In some examples, such as is shown in FIGS. 29-30, a proximal end **2628** of the second implant body portion **2606** can include a retention element **2656** comprising a hydrogel retention element, which is configured to expand into the ampulla **252** when the implant body
15 **2602** is implanted.

FIG. 27 illustrates an example of a side view of another lacrimal implant **2700** that is insertable through a lacrimal punctum **212**, **214** and into the associated canaliculus **208**, **210** (FIG. 2). In this example, the lacrimal implant **2700** comprises an implant body **2702** including first and second portions, which prior to implant, are linear relative to one another. The implant body **2702** extends from a proximal
20 end **2708** of the first portion to a distal end **2710** of the second portion. The proximal end **2708** can define a longitudinal proximal axis **2712** and the distal end **2710** can define a longitudinal distal axis **2714**. The implant body **2702** can be configured such that, when implanted, an angled intersection of between 45-135
25 degrees exists between the proximal axis **2712** and the distal axis **2714** such as for biasing at least a portion of the implant body **2702** against at least a portion of a lacrimal canaliculus **208**, **210** (FIG. 2) located at or more distal to a canaliculus curvature **250** (FIG. 2). In this example, the second portion of the implant body **2702** includes at least one undulation **2790** to facilitate the biasing of the implant
30 body **2702** against the portion of the lacrimal canaliculus **208**, **210**.

FIG. 28 illustrates an example of a side view of another lacrimal implant 2800 that is insertable through a lacrimal punctum 212, 214 and into the associated canaliculus 208, 210 (FIG. 2). In this example, the lacrimal implant 2800 comprises an implant body 2802 including first and second portions, which prior to implant, are linear relative to one another. The implant body 2802 extends from a proximal end 2808 of the first portion to a distal end 2810 of the second portion. The proximal end 2808 can define a longitudinal proximal axis 2812 and the distal end 2810 can define a longitudinal distal axis 2814. The implant body 2802 can be configured such that, when implanted, an angled intersection of between 45-135 degrees exists between the proximal axis 2812 and the distal axis 2814 for biasing at least a portion of the implant body 2802 against at least a portion of a lacrimal canaliculus 208, 210 (FIG. 2) located at or more distal to a canaliculus curvature 250 (FIG. 2). In this example, the second portion of the implant body 2802 includes at least one intermediately-disposed retainment projection 2892, such an annular rib-like projection. The retainment projection 2892 includes a cross-sectional size greater than an adjacent implant body portion and can facilitate the securement of an implanted position of the implant body 2802, while the adjacent narrower implant body portion can facilitate the biasing of the implant body 2802 against the portion of the lacrimal canaliculus 208, 210.

FIGS. 29-32 illustrate examples of a side view of other lacrimal implants 2900, 3000, 3100, 3200 that are insertable through a lacrimal punctum 212, 214 and into the associated canaliculus 208, 210 (FIG. 2). In these examples, each lacrimal implant 2900, 3000, 3100, 3200 can comprise an implant body 2902, 3002, 3102, 3202 including first 2904, 3004, 3104, 3204 and second 2906, 3006, 3106, 3206 portions, and can extend from a proximal end 2908, 3008, 3108, 3208 of the first portion 2904, 3004, 3104, 3204 to a distal end 2910, 3010, 3110, 3210 of the second portion 2906, 3006, 3106, 3206. The proximal end 2908, 3008, 3108, 3208 can define a longitudinal proximal axis 2912, 3012, 3112, 3212.

The second portion 2906, 3006, 3106, 3206 can include a fluid swellable retention element 2994, 3094, 3194, 3294 configured to expand laterally, relative to

the proximal axis **2912, 3012, 3112, 3212**, when the implant body **2902, 3002, 3102, 3202** is implanted. In various examples, the fluid swellable retention element **2994, 3094, 3194, 3294** can be formed such that one or both of expansion direction or expansion amount can be controlled. For instance, the fluid swellable retention
5 element **2994, 3094, 3194, 3294** can expand more in one plane than another to securely anchor the lacrimal implants. In some examples, the fluid swellable retention element **2994, 3094, 3194, 3294** includes a portion configured to expand laterally, relative to the proximal axis **2912, 3012, 3112, 3212**, in a direction away from a lacrimal canaliculus ampulla **252** (FIG. 2) when the implant body is
10 implanted. In some examples, as shown in FIGS. 29-30, the fluid swellable retention element **2994, 3094, 3194, 3294** includes a portion configured to expand laterally, relative to the proximal axis **2912, 3012, 3112, 3212**, in a direction toward the lacrimal canaliculus ampulla **252** (FIG. 2) when the implant body is implanted.

In some examples, the fluid swellable retention element **2994, 3094, 3194, 3294** can comprise hydrogel, which is insertable through the lacrimal punctum **212, 214** and into the associated canaliculus **208, 210** in a narrow profile. After
15 insertion, the hydrogel or other fluid swellable retention element can hydrate and expand to a wide configuration. Protrusions, such as at least one intermediately-disposed retainment projection **2992, 3092, 3192, 3292**, can be used to retain to an
20 implanted position of the lacrimal implants while the hydrogel or other swellable element expands.

FIG. 33 illustrates an example of a side view of another lacrimal implant **3300** that is insertable through a lacrimal punctum **212, 214** and into the associated canaliculus **208, 210** (FIG. 2). In this example, the lacrimal implant **3300** can
25 comprise an implant body **3302** including first **3304** and second **3306** portions, and can extend from a proximal end **3308** of the first portion **3304** to a distal end **3310** of the second portion **3306**. As shown, the second portion **3306** can include an expandable retention element **3393** comprising at least one of a coil, a braid, a stent, a mesh tube, a suture, a thermoset polymer, a thermoplastic, a heat activatable
30 material, or a shape memory material. The expandable retention element **3393** can

be configured to expand laterally, relative to a proximal axis 3312 defined by the first portion 3304, when the implant body is implanted. Protrusions, such as at least one intermediately-disposed retainment projection 3392, can be used to potentially further secure an implanted position of the lacrimal implant.

5 FIGS. 34A-34B illustrate examples of a schematic view of another lacrimal implant 3400 and an implant environment. In various examples, the implant body 3402 can include a graspable or other projection 3432, such as one or more projections extending laterally at least partially from or around a proximal end 3408 of a first implant body portion. In some examples, such as is shown in FIG. 34B, 10 the projections 3432 can include a set of wings for use in inserting the lacrimal implant 3400 into, or removing the implant from, an implanted position. The set of wings can be configured without migration in mind, as the implanted, non-linear configuration of the implant body 3402 can prevent migration by assuming a size or shape of a canaliculus curvature 250 and optionally, a lacrimal canaliculus ampulla 15 252.

In the examples of FIGS. 34A-34B, the one or more projections 3432 extend laterally in a direction parallel to or away from an eye 100 when implanted. In this way, the projections 3432 can still act as a graspable or feedback feature, but can limit patient discomfort when the lacrimal implant 3400 is implanted. In addition, 20 the projections 3432, by extending away from the eye 100, may not be buried in tissue and may be easily recognized by the patient or physician. This can allow for a quick determination if the lacrimal implant 3400 is being retained in its proper place without having to dig and search in the soft tissue surrounding the eye 100. In some instances, a simple pull on the lower eyelid can expose the projection 3432 25 pointed in a direction away from the eye 100. In the example of FIG. 34B, a lateral extension of at least one projection 3432 from the proximal end 3408 is substantially the same as a lateral extension direction of a second implant body portion relative to a distal end of the first implant body portion.

FIGS. 35-38 illustrate examples of an isometric view of various graspable 30 projections or other gripping means 3532, 3632, 3732, 3832 extending from a

proximal end of a lacrimal implant **3500, 3600, 3700, 3800**. The graspable or other projections **3532, 3632, 3732, 3832** can be used for various functions, including providing a structure to which a user can grasp onto during implant insertion or removal, inhibiting or preventing the associated lacrimal implant from passing
5 completely within a lacrimal punctum **212, 214** and associated canaliculus **208, 210** (FIG. 2), or for providing tactile or visual feedback information to the user, e.g., as to whether the implant is fully implanted.

In some examples, as shown in FIG. 35, the graspable projection **3532** can include two or more expandable arm members, which are sized to rest on an exterior
10 of the lacrimal punctum. The arm members can be affixed to an implant body **3502**, for example, via molding, adhesion or welding. The expandable arm members are capable of expanding so as to limit penetration of the lacrimal implant **3500** through the lacrimal punctum **212, 214** and into the associated canaliculus **208, 210**. While two arm members are shown, some include more than two arm members, such as
15 four arm members. The expandable arm members can assume an expanded profile separation distance **3505** that corresponds to about twice a diameter of the implant body, such that proximal ends of the proximal expandable arm members remain on the exterior of the punctum. The expandable arm members can expand in many ways from the narrow profile configuration to the expanded profile configuration,
20 and can include at least one of a coil, a braid, a suture, a thermoset polymer, a thermoplastic, a heat activated material, Nitinol, a shape memory material, a polymer, polypropylene, polyester, nylon, natural fibers, stainless steel, polymethylmethacrylate or polyimide. In some examples, the expandable arm members can be expanded manually, for example by a physician, after the lacrimal
25 implant has been positioned in the canalicular lumen **208, 210**.

In some examples, as shown in FIG. 36, the graspable projection **3632** can include a loop of a filament embedded in the proximal end of the lacrimal implant **3600** to permit removal of the implant with proximal tension to the loop, for example with forceps. In some examples, the loop of filament assumes a shape
30 similar to a purse handle that extends from the lacrimal implant with a loop so as to

facilitate removal of the lacrimal implant. The filament can comprise at least one of a heat activated material, Nitinol, a shape memory material, a polymer, polypropylene, polyester, nylon, natural fibers, stainless steel, polymethylmethacrylate or polyimide. In some embodiments, the filament may
5 comprise an absorbable thermo plastic polymer, for example at least one of polylactic acid (PLA), poly glycolic acid (PGA) or polylactic co-glycolic acid (PLGA). A distal end of the filament can be embedded in, molded to or other affixed to an implant body **3602** so as to secure the filament to the lacrimal implant.

In some examples, as shown in FIG. **37**, the graspable projection **3732** can
10 include at least one axially extending projection coupled with an implant body **3702**, which is configured to bias an outer most portion of the lacrimal canaliculus **208**, **210**. Due to the natural constriction against outward biasing of the canaliculus, the interplay between the axially extending projections and the canaliculus inhibits over insertion of an associated lacrimal implant **3700**.

In some examples, as shown in FIG. **38**, a longitudinal indentation, channel
15 or other recess **3832** in an implant body **3802** can be used in lieu of a graspable projection to permit insertion or removal of a lacrimal implant **3800**. The indentation, channel or other recess **3832** may extend axially along only a portion of an implant body a sufficient distance to facilitate removal of an associated lacrimal
20 implant. In further examples, a lacrimal implant can include a filament molded into an implant body and extending proximally for removal of the implant from the punctum.

FIGS. **39A-39B** illustrate examples of an isometric view of a drug insert **322**
and a removal facilitating filament **3999**. In some examples, as shown in FIG. **39A**,
25 the filament **3999** can extend from the drug insert **322** and is molded therein for removal purposes. Among other things, the filament **3999** can comprise a suture, a thermoset polymer, or a shape memory alloy. In some examples, as shown in FIG. **39B**, the filament **3999** extends along the drug insert **322** adjacent an implant body **3902** and is bonded to a distal end of the insert for removal purposes. Filament can

be bonded to the distal end of the drug core insert with an adhesive, such as cyanoacrylate, acrylic, epoxy, urethane or a hot melt adhesive.

FIG. 40 is a block diagram illustrating an example of a method 4000 of manufacturing a lacrimal implant configured to be at least partially insertable
5 through a lacrimal punctum and into the associated canaliculus. At 4002, an implant body extending from a proximal end of a first body portion to a distal end of a second body portion is formed. In some examples, various sizes of implant bodies are formed to fit various patient anatomies. In various examples, the proximal end is formed to define a longitudinal proximal axis and the distal end is formed to
10 define a longitudinal distal axis. A formation of the implant body can be configured such that, when implanted, the proximal axis and the distal axis intersect at an angle of at least 45 degrees to laterally bias at least a portion of the implant body against at least a portion of a lacrimal canaliculus located at or more distal to a canaliculus curvature.

15 In some examples, the second body portion is formed to include a dilator generally narrowing from a location near a proximal end of the second body portion to the distal end of the second body portion. In some examples, the dilator is formed by sloping an outer surface of the second portion of the implant body between about 1 degree and about 10 degrees with respect to the longitudinal distal
20 axis. In some examples, the outer surface of the second implant body portion is sloped to a dilator tip of between about 0.2 millimeters and about 0.5 millimeters.

In some examples, the implant body is formed to include a graspable or other projection extending laterally from the proximal end of the first body portion. In certain examples, the projection is formed to substantially align with a lateral
25 extension direction of the second body portion relative to the first body portion. In certain examples, the projection is formed such that, when implanted, it laterally extends from the proximal end of the first body portion in a direction that is parallel to or away from an eye.

At 4004, a drug insert is disposed in at least one of the first body portion or
30 the second body portion. In various examples, the drug insert is positioned such that

an exposed drug insert surface sits adjacent at least one of the proximal end or the distal end for providing a sustained drug or other therapeutic agent release to an eye, nasal passage or inner ear, for example. In certain examples, a first drug insert is disposed in the first body portion and a second drug insert is disposed in the second
5 body portion. In various examples, the one or more drug inserts comprise drug cores including the drug or other therapeutic agent.

At 4006, an outer surface portion of the implant body is coated with at least one of a fluid swellable material, a lubricious coating or an antimicrobial coating. In various examples, the outer surface portion of the implant body is polished using
10 a polishing process.

Sheath Body Examples:

In various ways, the sheath body can comprise appropriate shapes and materials to control migration of drug or other therapeutic agents from a drug insert. In some examples, the sheath body is configured to be conformable to an implant
15 anatomy, such as an anatomy of a lacrimal punctum or associated canaliculus. As discussed, in some examples, the sheath body at least partially covers or surrounds the drug insert and can fit snugly against an outer surface of a matrix/agent mixture. The sheath body can be made from a material that is substantially impermeable to the drug or other therapeutic agent so that the rate of migration of the drug or agent
20 is largely controlled by an exposed surface area of the drug insert that is not covered by the sheath body. In many examples, migration of the agents through the sheath body can be about one tenth of the migration of the agent through the exposed surface of the drug insert, or less. Suitable sheath body materials can include, among others, polyimide, polyethylene terephthalate (PET). The sheath body can
25 have a thickness, as defined from the sheath surface adjacent the outer matrix/agent mixture surface to an opposing sheath surface away from the outer surface, of about 0.00025 inches to about 0.0015 inches. The total diameter of the sheath that extends across the drug insert ranges from about 0.2 millimeters to about 1.2 millimeters. The drug insert can be formed by dip coating the matrix in the sheath body. In some
30 examples, the sheath body can comprise a tube into which the matrix/agent mixture

is introduced. The sheath body can also be dip coated around the matrix/agent mixture, for example dip coated around a pre-formed matrix/agent core.

The sheath body can be provided with one or more additional features such as to facilitate clinical use of the lacrimal implants discussed herein. For example,
5 the sheath can receive a drug insert that is exchangeable *in situ*, while the implant body remains implanted in the patient, or after its removal. In some examples, the sheath body can be provided with one or more external protrusions that apply force to the sheath body when squeezed, which cause the matrix/agent mixture to be ejected from the sheath body. A replacement drug insert can then be positioned in
10 the sheath body.

Therapeutic Agent Examples:

A therapeutic agent (or simply "agent") can comprise, among other things, a drug made from one or any combination of the following or their equivalents, derivatives or analogs, including, anti-glaucoma medications, (e.g. adrenergic
15 agonists, adrenergic antagonists (beta blockers), carbonic anhydrase inhibitors (CAIs, systemic and topical), parasympathomimetics, prostaglandins and hypotensive lipids, and combinations thereof), antimicrobial agent (e.g., antibiotic, antiviral, antiparasitic, antifungal, etc.), a corticosteroid or other anti-inflammatory (e.g., an NSAID or other analgesic and pain management compounds), a
20 decongestant (e.g., vasoconstrictor), an agent that prevents or modifies an allergic response (e.g., an antihistamine, cytokine inhibitor, leukotriene inhibitor, IgE inhibitor, immunomodulator), a mast cell stabilizer, cycloplegic, mydriatic or the like.

Example available agents include, but are not limited to, thrombin inhibitors;
25 antithrombogenic agents; thrombolytic agents; fibrinolytic agents; vasospasm inhibitors; vasodilators; antihypertensive agents; antimicrobial agents, such as antibiotics (such as tetracycline, chlortetracycline, bacitracin, neomycin, polymyxin, gramicidin, cephalixin, oxytetracycline, chloramphenicol, rifampicin, ciprofloxacin, tobramycin, gentamycin, erythromycin, penicillin, sulfonamides, sulfadiazine,
30 sulfacetamide, sulfamethizole, sulfisoxazole, nitrofurazone, sodium propionate),

antifungals (such as amphotericin B and miconazole), and antivirals (such as idoxuridine trifluorothymidine, acyclovir, gancyclovir, interferon); inhibitors of surface glycoprotein receptors; antiplatelet agents; antimitotics; microtubule inhibitors; anti-secretory agents; active inhibitors; remodeling inhibitors; antisense

5 nucleotides; anti-metabolites; antiproliferatives (including antiangiogenesis agents); anticancer chemotherapeutic agents; anti-inflammatories (such as hydrocortisone, hydrocortisone acetate, dexamethasone 21-phosphate, fluocinolone, medrysone, methylprednisolone, prednisolone 21-phosphate, prednisolone acetate, fluoromethalone, betamethasone, triamcinolone, triamcinolone acetonide); non

10 steroidal anti-inflammatories (NSAIDs) (such as salicylate, indomethacin, ibuprofen, diclofenac, flurbiprofen, piroxicam indomethacin, ibuprofen, naxopren, piroxicam and nabumetone). Examples of such anti-inflammatory steroids contemplated for use with the present lacrimal implants, include triamcinolone acetonide (generic name) and corticosteroids that include, for example,

15 triamcinolone, dexamethasone, fluocinolone, cortisone, prednisolone, flumetholone, and derivatives thereof.); antiallergenics (such as sodium chromoglycate, antazoline, methapyrilene, chlorpheniramine, cetirizine, pyrilamine, prophenyridamine); anti proliferative agents (such as 1,3-cis retinoic acid, 5-fluorouracil, taxol, rapamycin, mitomycin C and cisplatin); decongestants (such as phenylephrine, naphazoline, tetrahydrazoline); miotics and anti-cholinesterase (such as pilocarpine, salicylate,

20 carbachol, acetylcholine chloride, physostigmine, eserine, diisopropyl fluorophosphate, phospholine iodine, demecarium bromide); antineoplastics (such as carmustine, cisplatin, fluorouracil); immunological drugs (such as vaccines and immune stimulants); hormonal agents (such as estrogens,--estradiol, progestational,

25 progesterone, insulin, calcitonin, parathyroid hormone, peptide and vasopressin hypothalamus releasing factor); immunosuppressive agents, growth hormone antagonists, growth factors (such as epidermal growth factor, fibroblast growth factor, platelet derived growth factor, transforming growth factor beta, somatotrapin, fibronectin); inhibitors of angiogenesis (such as angiostatin,

30 anecortave acetate, thrombospondin, anti-VEGF antibody); dopamine agonists;

radiotherapeutic agents; peptides; proteins; enzymes; extracellular matrix; components; ACE inhibitors; free radical scavengers; chelators; antioxidants; anti polymerases; photodynamic therapy agents; gene therapy agents; and other therapeutic agents such as prostaglandins, antiprostaglandins, prostaglandin
5 precursors, including antiglaucoma drugs including beta-blockers such as Timolol, betaxolol, levobunolol, atenolol, and prostaglandin analogues such as bimatoprost, travoprost, latanoprost etc; carbonic anhydrase inhibitors such as acetazolamide, dorzolamide, brinzolamide, methazolamide, dichlorphenamide, diamox; and neuroprotectants such as lubezole, nimodipine and related compounds; and
10 parasympathomimetics such as pilocarpine, carbachol, physostigmine and the like.

Additional agents that can be used with the present lacrimal implants include, but are not limited to, drugs that have been approved under Section 505 of the United States Federal Food, Drug, and Cosmetic Act or under the Public Health Service Act, some of which can be found at the U.S. Food and Drug Administration
15 (FDA) website <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index>. The present lacrimal implants can also be used with drugs listed in the Orange Book, either in paper or in electronic form, which can be found at the FDA Orange Book website (<http://www.fda.gov/cder/ob/>)), that has or records the same date as, earlier date than, or later date than, the filing date of this patent document. For example,
20 these drugs can include, among others, dorzolamide, olopatadine, travoprost, bimatoprost, cyclosporin, brimonidine, moxifloxacin, tobramycin, brinzolamide, aciclovir timolol maleate, ketorolac tromethamine, prednisolone acetate, sodium hyaluronate, nepafenac, bromfenac, diclofenac, flurbiprofen, suprofenac, binoxan, patanol, dexamethasone/tobramycin combination, moxifloxacin, or acyclovir.

25 Examples of diseases or disorders that can be treated with above-listed agents include, but are not limited to, glaucoma, pre- and post-surgical ocular treatments, dry eye, anti-eye allergy, anti-infective, post-surgical inflammation or pain, respiration-related disorders, such as allergies, inner ear disorders, such as dizziness or migraines, or other systemic disorders, such as hypertension,
30 cholesterol management, pulmonary disorders or immunological disorders. In some

examples, the therapeutic agent can include a lubricant or a surfactant, for example a lubricant to treat dry eye. In other examples, the therapeutic agent can include an absorbent capable of absorbing tear from an eye.

Drug Insert Examples:

5 The drug insert can comprise one or more drugs or other therapeutic agents, and in some examples, one or more matrix materials to provide sustained release of the drug or other agents. The one or more drugs or other therapeutic agents can migrate from an exposed surface of the drug insert to the target tissue (e.g., ciliary muscles of an eye) based, at least in part, on a solubility of the drugs or agents in the
10 matrix. The rate of migration of the drugs or agents from the exposed surface can also be related to the concentration of drugs or agents dissolved in the matrix. In some examples, the concentration of drugs or agents dissolved in the drug insert can be controlled to provide the desired release rate of the drugs or agents. In addition or in combination, the rate of migration of drugs or agents from the exposed surface
15 can be related to one or more properties of the matrix in which the drugs or agents dissolve, such as the properties of a silicone matrix formulation. In some examples, the drugs or agents included in the drug insert can include liquid, solid, solid gel, solid crystalline, solid amorphous, solid particulate, or dissolved forms. In one such example, liquid Latanoprost droplets or solid Bimatoprost particles are dispersed in
20 a silicone matrix.

 The drug insert can comprise one or more biocompatible materials capable of providing a sustained release of the one or more drugs or agents. Although the drug insert is primarily discussed above with respect to an example comprising a matrix including a substantially non-biodegradable silicone matrix with dissolvable
25 inclusions of the drugs or agents located therein, the drug insert can include other structures that provide sustained release of the drugs or agents, for example a biodegradable matrix, a porous drug insert, a liquid drug insert or a solid drug insert. A matrix that includes the drugs or agents can be formed from either biodegradable or non-biodegradable polymers. In some examples, a non-biodegradable drug insert
30 can include silicone, acrylates, polyethylenes, polyurethane, polyurethane, hydrogel,

polyester (e.g., DACRON.RTM. from E. I. Du Pont de Nemours and Company, Wilmington, Del.), polypropylene, polytetrafluoroethylene (PTFE), expanded PTFE (ePTFE), polyether ether ketone (PEEK), nylon, extruded collagen, polymer foam, silicone rubber, polyethylene terephthalate, ultra high molecular weight

5 polyethylene, polycarbonate urethane, polyurethane, polyimides, stainless steel, nickel-titanium alloy (e.g., Nitinol), titanium, stainless steel, cobalt-chrome alloy (e.g., ELGILOY.RTM. from Elgin Specialty Metals, Elgin, Ill.; CONICHROME.RTM. from Carpenter Metals Corp., Wyomissing, Pa.). In some examples, a biodegradable drug insert can comprise one or more biodegradable

10 polymers, such as protein, hydrogel, polyglycolic acid (PGA), polylactic acid (PLA), poly(L-lactic acid) (PLLA), poly(L-glycolic acid) (PLGA), polyglycolide, poly-L-lactide, poly-D-lactide, poly(amino acids), polydioxanone, polycaprolactone, polygluconate, polylactic acid-polyethylene oxide copolymers, modified cellulose, collagen, polyorthoesters, polyhydroxybutyrate, polyanhydride, polyphosphoester,

15 poly(alpha-hydroxy acid) and combinations thereof. In some examples, the drug insert can comprise a hydrogel polymer.

Closing Notes:

Among other things, lacrimal implants and related methods providing secure retention within a lacrimal punctum and canaliculus of an eye are discussed herein.

20 The lacrimal implants can comprise an implant body configured for at least partial insertion through the lacrimal punctum and into the canaliculus. The implant body can include first and second portions, and can extend from a proximal end of the first portion defining a longitudinal proximal axis to a distal end of the second portion defining a longitudinal distal axis. The implant body can be configured such

25 that, when implanted using an integral dilator, an at least 45 degree angled intersection exists between the proximal axis and the distal axis. In this way, at least a portion of the implant body can be biased against at least a portion of the lacrimal canaliculus located at or more distal to a canalicular curvature, thereby retaining an implanted position of the lacrimal implant using anatomical structures.

30 In various examples, the lacrimal implant can further comprise a drug insert

disposed in at least one of the first portion or the second portion of the implant body to provide a sustained release of a drug or other therapeutic agent to an eye, nasal passage, or inner ear system, for instance.

Advantageously, in some examples, the present lacrimal implants can
5 successfully block the flow of tears or provide sustained delivery of a drug or other therapeutic agent to an eye, nasal passage, or inner ear for varying periods of time, such as from days to months to years. In addition, by including first and second implant body cavities, a dual drug or other agent releasing profile can be possible. For instance, two separate drugs can be released from two different implant
10 locations. Further, the canalicular curve retaining configuration of the present implant body can reduce over-stretching of the lacrimal punctum and canaliculus and inadvertent fall out of implants. Even further, it is believe the present lacrimal implants can be implemented so as to provide a one-size-fits-all (or many) regime, as an expandable coating or other expandable retention member can be applied to
15 the implant body, such as an outer surface portion of the implant body, to fit in hollow tissue structures of varying sizes. The present lacrimal implant may also be better tolerated by a patient due to, for example, an orientation of a graspable or other projection located at the proximal end of the implant body.

The above Detailed Description includes references to the accompanying
20 drawings, which form a part of the Detailed Description. The drawings show, by way of illustration, specific embodiments in which the invention can be practiced. These embodiments are also referred to herein as “examples.” All publications, patents, and patent documents referred to in this document are incorporated by reference herein in their entirety, as though individually incorporated by reference.
25 In the event of inconsistent usages between this document and those documents so incorporated by reference, the usage in the incorporated reference(s) should be considered supplementary to that of this document; for irreconcilable Inconsistencies, the usage in this document controls.

In this document, the terms “a” or “an” are used, as is common in patent
30 documents, to include one or more than one, independent of any other instances or

usages of “at least one” or “one or more.” In this document, the term “or” is used to refer to a nonexclusive or, such that “A or B” includes “A but not B,” “B but not A,” and “A and B,” unless otherwise indicated. In this document, the term “about” is used to refer to an amount that is approximately, nearly, almost, or in the vicinity of being equal to a stated amount.

In this document, the term “proximal” refers to a location relatively closer to a hand of a physician implanting a lacrimal implant into a patient, and the term “distal” refers to a location relatively further from the hand of the physician, particularly during the implanting of the implant into the patient.

In this document, the term “hydrogel” is used to refer to an absorbing or otherwise retaining material (e.g., adsorbing material), such as super-absorbent polymers, hydrocolloids, and water-absorbent hydrophilic polymers, for example. In some examples, the term “hydrogel” refers to super-absorbent polymer particles in a “dry” state, more specifically, particles containing from no water up to an amount of water less than the weight of the particles, such as less than about 5%, by weight, water. In some examples, the term “hydrogel” refers to a super-absorbent polymer in the “dry” state when the hydrogel is not expandable and also refers to its hydrated or expanded state, more specifically, hydrogels that have absorbed at least their weight in water, such as several times their weight in water. As the hydrogel material absorbs fluid, its size can increase (swell) and its shape can change to bias against at least a portion of a lacrimal canaliculus ampulla or lacrimal canaliculus wall, for example.

In the appended claims, the terms “including” and “in which” are used as the plain-English equivalents of the respective terms “comprising” and “wherein.” Also, in the following claims, the terms “including” and “comprising” are open-ended, that is, a system, assembly, device, article, or process that includes elements in addition to those listed after such a term in a claim are still deemed to fall within the scope of that claim. Moreover, in the following claims, the terms “first,” “second,” and “third,” etc. are used merely as labels, and are not intended to impose numerical requirements on their objects.

The above description is intended to be illustrative, and not restrictive. For example, the above-described examples (or one or more features thereof) can be used in combination with each other. Other embodiments can be used, such as by one of ordinary skill in the art upon reviewing the above description. Also, in the
5 above Detailed Description, various features can be grouped together to streamline the disclosure. This should not be interpreted as intending that an unclaimed disclosed feature is essential to any claim. Rather, inventive subject matter can lie in less than all features of a particular disclosed embodiment. Thus, the following claims are hereby incorporated into the Detailed Description, with each claim
10 standing on its own as a separate embodiment. The scope of the invention should be determined with reference to the appended claims, along with the full scope of equivalents to which such claims are entitled.

The Abstract is provided to allow the reader to quickly ascertain the nature of the technical disclosure. It is submitted with the understanding that it will not be
15 used to interpret or limit the scope or meaning of the claims.

WHAT IS CLAIMED IS:

1. A lacrimal implant insertable into a lacrimal canaliculus, comprising:
an implant body, including first and second portions, the implant body extending from a proximal end of the first portion to a distal end of the second portion;
the proximal end of the first portion defining a longitudinal proximal axis and the distal end of the second portion defining a longitudinal distal axis;
the implant body configured such that, when implanted in the lacrimal canaliculus, an angled intersection exists between the proximal axis and the distal axis for biasing at least a portion of the implant body against at least a portion of the lacrimal canaliculus located at or more distal to a canalicular curvature; and
wherein the second portion of the implant body includes a longitudinal length having a magnitude less than four times a longitudinal length of the first portion of the implant body.
2. The lacrimal implant of claim 1, wherein the implant body is configured such that an angled intersection exists between the proximal axis and the distal axis prior to being implanted in the lacrimal canaliculus.
3. The lacrimal implant of any of claims 1 or 2, wherein the implant body is configured to partially or completely inhibit fluid flow into and through the lacrimal canaliculus.
4. The lacrimal implant of any of claims 1-3, wherein a distal end of the first portion is integral with the second portion at or near a proximal end of the second portion.
5. The lacrimal implant of any of claims 1-4, wherein one or both of the first portion or the second portion includes a fluid swellable retention element configured to expand.
6. The lacrimal implant of claim 5, wherein the second portion includes the fluid swellable retention element, the fluid swellable retention element configured to expand

laterally, relative to the proximal axis of the first portion, when the implant body is implanted.

7. The lacrimal implant of claim 6, wherein the fluid swellable retention element includes a portion configured to expand laterally in a direction away from a lacrimal canaliculus ampulla when the implant body is implanted.

8. The lacrimal implant of any of claims 6 or 7, wherein the fluid swellable retention element includes a portion configured to expand laterally in a direction toward a lacrimal canaliculus ampulla when the implant body is implanted.

9. The lacrimal implant of any of claims 1-8, wherein the second portion includes an expandable retention element comprising at least one of a coil, a braid, a stent, a mesh tube, a suture, a thermoset polymer, a thermoplastic, a heat activatable material, or a shape memory material, the expandable retention element configured to expand laterally, to form the angled intersection, when the implant body is implanted.

10. The lacrimal implant of any of claims 1-9, comprising an expandable retention element disposed around a portion of the second portion, the expandable retention element configured to bias the second portion away from a wall of the lacrimal canaliculus upon expansion.

11. The lacrimal implant of any of claims 1-10, wherein the second portion includes an arm member movable between a first configuration and a second configuration;
the arm member, in the first configuration, disposable along the implant body for insertion into the lacrimal canaliculus and, in the second configuration, laterally extendable from one side of the implant body.

12. The lacrimal implant of any of claims 1-11, wherein the second portion includes an integral dilator, the integral dilator generally narrowing from a location near a

proximal end of the second portion to the distal end of the second portion to facilitate implantation of the implant body into the lacrimal canaliculus.

13. The lacrimal implant of claim 12, wherein a diameter of an integral dilator tip is between about 0.2 millimeters and about 0.5 millimeters.

14. The lacrimal implant of any of claims 12 or 13, wherein an outer surface slope of the integral dilator, as measured from the location near the proximal end of the second portion to the distal end of the second portion, is between about 1 degree and about 10 degrees with respect to the distal axis.

15. The lacrimal implant of any of claims 1-14, wherein the second portion includes at least one undulation.

16. The lacrimal implant of any of claims 1-15, wherein at least one of the first portion or the second portion comprises at least one intermediately-disposed annular, semi-annular, column-like, or barrel-like projection, the intermediately-disposed projection having a cross-sectional size greater than an adjacent implant body portion.

17. The lacrimal implant of any of claims 1-16, comprising a graspable projection extending at least partially from the proximal end of the first portion, the graspable projection configured to seat against or near a lacrimal punctum when the implant body is implanted.

18. The lacrimal implant of claim 17, wherein the second portion includes an element extending or expanding laterally into a lacrimal canaliculus ampulla when the implant body is implanted.

19. The lacrimal implant of any of claims 17 or 18, wherein the graspable projection extends laterally from the proximal end of the first portion, in a direction that is parallel to or away from an eye, when the implant body is implanted.

20. The lacrimal implant of any of claims 1-19, wherein the implanted angled intersection of the proximal axis and the distal axis is at least about 45 degrees.
21. The lacrimal implant of any of claims 1-20, comprising a therapeutic agent.
22. The lacrimal implant of claim 21, comprising at least one drug insert including a drug core, the drug core comprising the therapeutic agent.
23. The lacrimal implant of claim 22, wherein the drug core comprises at least one exposed surface to deliver a sustained release.
24. A kit comprising the lacrimal implant of any of claims 1-23, and an instruction for using the lacrimal implant to treat an eye disease.
25. A kit comprising the lacrimal implant of any of claims 1-23, and an instruction for using the lacrimal implant to treat a respiration-related disorder.
26. A kit comprising the lacrimal implant of any of claims 1-23, and an instruction for using the lacrimal implant to treat an inner ear disorder.
27. A lacrimal implant for insertion into a lacrimal canaliculus, comprising:
an implant body non-linearly extending from a proximal end portion positionable within a vertical section of the lacrimal canaliculus to a distal end portion positionable within a horizontal section of the lacrimal canaliculus and having an intermediate portion therebetween;
the intermediate portion partially extending in a first direction toward the proximal end portion and partially extending in a second direction toward the distal end portion such that, when implanted in the lacrimal canaliculus, the implant body directionally biases laterally against at least a portion of the lacrimal canaliculus located at or more distal to a canalicular curvature; and

wherein the implant body inhibits fluid flow into and through the lacrimal canaliculus.

28. The lacrimal implant of claim 27, wherein a longitudinal length of the implant body positionable within the vertical section of the lacrimal canaliculus is less than four times a longitudinal length of the implant body positionable within the horizontal section of the lacrimal canaliculus.

29. The lacrimal implant of any of claims 27 or 28, wherein the first direction extension of the intermediate portion is at an angle between about 45 degrees and about 135 degrees relative to the second direction extension of the intermediate portion.

30. The lacrimal implant of any of claims 27-29, wherein the intermediate portion partially extends in a third direction, substantially opposite the second direction, toward a lacrimal canaliculus ampulla when the implant body is implanted.

31. The lacrimal implant of any of claims 27-30, wherein the second direction extension includes a longitudinal dilator having a generally concave shape relative to the first direction extension; and wherein a radius of the generally concave shape is less than the radius of the canaliculus curvature.

32. The lacrimal implant of any of claims 27-31, wherein the second direction extension includes a longitudinal dilator having a generally convex shape relative to the first direction extension.

33. The lacrimal implant of any of claims 27-32, wherein the second direction extension includes a longitudinal dilator having an axis substantially perpendicular to an axis of the first direction extension.

34. The lacrimal implant of any of claims 27-33, wherein at least one of the proximal end portion or the distal end portion comprises at least one intermediately-disposed

annular, semi-annular, column-like, or barrel-like projection, the intermediately-disposed projection having a cross-sectional size greater than an adjacent implant body portion.

35. The lacrimal implant of any of claims 27-34, comprising a graspable projection, the graspable projection extending laterally from the proximal end portion.

36. The lacrimal implant of any of claims 27-35, comprising a fluid swellable material disposed on an outer surface portion of the implant body, the fluid swellable material configured to expand an outer surface diameter portion of the implant body when implanted.

37. The lacrimal implant of any of claims 27-36, comprising at least one of a first drug insert disposed in the proximal end portion or a second drug insert disposed in the distal end portion, one or both of the first or second drug inserts inhibiting fluid flow through the implant body and including at least one exposed surface configured to deliver a sustained release.

38. The lacrimal implant of claim 37, comprising a cavity in the proximal end portion, the cavity configured to house the first drug insert in the form of a drug core, the drug core including a first agent configured to treat an eye.

39. The lacrimal implant of any of claims 37 or 38, comprising a cavity in the distal end portion, the cavity configured to house the second drug insert in the form of a drug core, the drug core including a second agent configured to be received by a nasal passage.

40. A method of manufacturing a lacrimal implant insertable into a lacrimal canaliculus, the method comprising:

forming an implant body extending from a proximal end of a first body portion to a distal end of a second body portion, including

extending the second body portion to a longitudinal length which is less than four times a longitudinal length of the first body portion; and

configuring the proximal end and the distal end to respectively define, when implanted in the lacrimal canaliculus, a longitudinal proximal axis and a longitudinal distal axis that intersect at an angle such that the implant body is configured to directionally bias laterally against at least a portion of the lacrimal canaliculus located at or more distal to a canaliculus curvature.

41. The method of claim 40, wherein forming one or both of the first body portion or the second body portion includes forming at least one intermediately-disposed annular, semi-annular, column-like, or barrel-like projection, the intermediately-disposed projection having a cross-sectional size greater than an adjacent implant body portion.

42. The method of any of claims 40 or 41, wherein forming the second body portion includes forming a dilator generally narrowing from a location near a proximal end of the second body portion to the distal end of the second body portion.

43. The method of any of claims 40-42, wherein forming the dilator includes forming an outer surface slope of the implant body, as measured from the location near the proximal end of the second body portion to the distal end of the second body portion, between about 1 degree and about 10 degrees with respect to the longitudinal distal axis.

44. The method of any of claims 40-43, wherein forming the second portion includes disposing a fluid swellable retention element near a distal end of the first body portion, including disposing the fluid swellable retention element such that a lateral expansion thereof, relative to the proximal axis, is configured to bias against at least a portion of the lacrimal canaliculus or a lacrimal canaliculus ampulla anatomy when the implant body is implanted.

45. The method of any of claims 40-44, comprising coating an outer surface portion of the implant body with a fluid swellable material.

46. The method of any of claims 40-45, comprising disposing at least one of a first drug insert in the first body portion or a second drug insert in the second body portion, including positioning at least one of an exposed surface of the first drug insert adjacent to the proximal end or an exposed surface of the second drug insert adjacent to the distal end to provide a sustained release of a first agent or a second agent, respectively.

47. A method of treating a patient having at least one of an eye disorder, a respiration-related disorder, a pulmonary disorder, an immunological disorder or an inner ear disorder, the method comprising:

inserting a lacrimal implant into at least one lacrimal canaliculus of the patient, the lacrimal implant comprising,

an implant body, including first and second portions, the implant body extending from a proximal end of the first portion to a distal end of the second portion, the proximal end defining a longitudinal proximal axis and the distal axis defining a longitudinal distal axis;

the implant body configured such that, when implanted in a lacrimal canaliculus, an angled intersection exists between the proximal axis and the distal axis for biasing at least a portion of the implant body against at least a portion of the lacrimal canaliculus located at or more distal to a canalicular curvature; and at least one of a first drug insert disposed in the first portion or a second drug insert disposed in the second portion, the first and second drug inserts configured to respectively provide a sustained release of a first agent or a second agent.

48. The method of claim 47, wherein inserting the lacrimal implant includes concurrently dilating the lacrimal canaliculus, using an integral dilator of the implant body, as the distal end of the second portion is moved into the lacrimal canaliculus.

49. The method of any of claims 47 or 48, comprising removing the inserted implant body from the lacrimal canaliculus.

50. The method of any of claims 47-49, comprising replacing the lacrimal implant that has been inserted with a second lacrimal implant including at least one of the same first or second agent following an interval of time.