

Method for producing milk like products

Field of the invention

5 The present invention concerns a method for producing mammary gland cells. The present invention also concerns a method for producing in vitro a mammalian milk like product, for example human milk like product, which comprises generating lactocytes derived from mammalian induced pluripotent stem cells (hiPSC), for example human induced pluripotent stem cells (hiPSC), through culture and
10 differentiation and/or mammary-like gland organoids comprising such lactocytes and expressing the mammalian milk like product, for example the human milk like product from such lactocytes and/or mammary-like gland organoids. The present invention also relates to the mammalian milk like product, for example the human milk like product obtainable from such method.

15

Background of the invention

Mammalian and especially human milk is a complex fluid with a multitude of components, each of which may contribute substantially to infant and perhaps
20 maternal health. It is becoming increasingly clear that human breastmilk is the most appropriate source of nutrition at least up to the age of 6 months. Many components of human milk are simply not found or poorly found or less active in cow's milk upon which infant formula manufacture is based. This includes for instance protein lactoferrin, growth factors, long chain polyunsaturated fatty acids
25 or oligosaccharides. Human milk composition has been used as a gold standard to develop current infant formula, despite recent major development in infant formula composition, it is illusory to think that human milk replication will be achieved with current manufacturing processes.

Today the only source of human milk is human donors (breastfeeding mothers). Donation are reported for non-commercial use (human milk biobank) and commercial use. However, this is limited and has strong regulatory, safety and
5 sometime ethical or religious constraints.

Stem cells were found in mammalian and especially human milk called human breastmilk stem cells (hBSC). hBSCs were shown to be highly plastic and to differentiate in culture into multiple cell types and more importantly into the three
10 lineages required to shape the lobulo-alveolar structure of the human mammary gland (Hassiotou F. et al. Stem Cells. 2012). However, the use of hBSC to produce human breast milk is neither practical nor sustainable, as it requires human donors.

A technology based on a cell line with stem cell functionality - called induced
15 pluripotent stem cells (iPSC) is known. A reliable two step protocol to generate human mammary like organoids from human iPSC (hiPSC) was developed (Ying Qu et al, Stem Cell Report vol 8, 205-215, February 14th 2017).

Accordingly, it is an object of the present invention to provide improved methods
20 for producing mammary gland cells and to reproduce expression of mammalian milk, for example human milk in cultured cells. It is also an object of the present invention to prepare customized mammalian milk like product, for example human milk like product, in cultured cells which could be adapted to specific needs of the recipient and/or to produce human milk bioactives to complement existing cow-
25 based solutions for infant nutrition.

Summary of the invention

The present invention solves the above-mentioned technical problem.

Provided herein is a method of producing a population of mammary gland cells, comprising:

- i) culturing mammalian induced pluripotent stem cells (miPSCs) in a culture medium comprising bone morphogenic protein 4 (BMP4) to generate embryoid bodies (EBs), and
- ii) growing the EBs to generate a population of mammary cells.

Also provided herein is use of BMP4 for increasing the differentiation efficiency of mammalian induced pluripotent stem cell (miPSCs) into mammary-gland progenitor cells in a differentiation protocol.

Further provided herein is a method for producing a mammalian milk like product, comprising:

- A) Generating lactocyte mammary-like gland organoids derived from mammalian induced pluripotent stem cells (miPSC);
- B) Secreting the mammalian milk like product from said lactocytes, wherein step A) comprises culturing the miPSCs in a culture medium comprising BMP4.

Further provided herein is a human milk like product which is obtainable according to the methods as described anywhere herein.

Further provided herein is a human milk like product for use in therapy.

Finally, provided herein is use of a human milk like product as described anywhere herein as a human milk substitute, optionally a breast-feeding substitute.

Detailed description of the invention

Definitions

- 5 Within the context of the present invention, the term “in vitro” means performed or taking place in a test tube, culture dish, bioreactor or elsewhere outside a living organism.

- Within the context of the present invention, the term “mammalian” identify an
10 animal belonging to the mammalian species, for example human, cow, monkey, camel, sheep, goat etc.

- Within the context of the present invention, the terms “lactocytes” or “mammary-like cells” identify secretory epithelial cells expressing CK18 cell marker and derived
15 from mammalian induced pluripotent stem cells (miPSC) and in particular from human induced pluripotent stem cells (hiPSC). Human induced pluripotent stem cells (hiPSC) as used herein are commercially available and may be selected from any suitable hiPSC line. A suitable human induced pluripotent stem cell line in the context of the current invention is e.g., hiPSC line 603, commercially available from
20 Fujifilm Cellular Dynamics, Inc (FCDI), as used according to the invention. Further suitable hiPSCs may be selected as described e.g., in Ying Qu et al., (2017, supra). In one embodiment of the present invention, the hiPSC are not engineered. In one embodiment, they are not engineered to comprise an exogenous nucleic acid and/or an inducible gene expression system which includes an exogenous nucleic acid
25 acid, where the inducible gene expression system is configured to express a hormone or a signaling factor. In one embodiment, the exogenous nucleic acid and/or inducible gene expression system which includes an exogenous nucleic acid is promoting the cell differentiation towards lactocytes.

Within the context of the present invention the terms “mammary gland like organoids” or “mammary like organoids” mean a miniaturized and simplified version of a mammary gland which develops in two or three dimensions (2D/3D) and which comprises lactocytes as above defined.

5

Within the context of the present invention, the term “human milk like product” is a cell cultured milk product. It is an edible product which is expressed by the lactocytes and/or mammary gland like organoids generated according to the process of the present invention.

10

The “human milk like product” according to the present invention can have the same components as human breast milk of a well-nourished mother (for example in terms of bioactives, macro and micronutrients and levels thereof). This is referred to herein as a “standard human milk product”. Alternatively, “human milk like product”

15 according to the present invention can have altered ratios and concentrations of components found naturally in human breast milk of a well-nourished mother. This is referred to herein as a “non-standard milk like product”. A “human milk like product” according to the present invention can be modified such that it includes components that are not found naturally in human breast milk of a well-nourished
20 mother (a “modified milk like product”). Non-limiting examples of human milk like products are selected from the group consisting of: supplement, fortifier, human breast milk substitute (or replacer) and ingredient enriched in only one and/or a portion of bioactives, macro- and micronutrients which can be typically found in human breast milk of a well-nourished mother.

25

The “human milk like product” can be used to replace consumption of naturally lactated milk (a “human milk substitute”). The milk substitute product can be used

as a supplement (a “human milk supplement”) or as a fortifier (a “human milk fortifier”) to be consumed in combination with naturally lactated milk.

In an embodiment, the standard human milk like product according to the present invention comprises at least macro- and micronutrients which can be typically found in human breast milk of a well-nourished mother. In one embodiment, the standard human milk like product according to the present invention comprises: proteins, peptides, lipids (including linoleic acid and alpha-linolenic acid), carbohydrates, Vitamins (including Vitamin A, Vitamin D3, Vitamin E, Vitamin K, Thiamin, Riboflavin, Niacin, Vitamin B6, Vitamin B12, Pantothenic acid, folic acid, Vitamin C and Biotin), minerals (including iron, calcium, phosphorus, magnesium, sodium, chloride, potassium, manganese, iodine, selenium, copper and zinc), choline, myoinositol and L-carnitine. In one embodiment, the standard human milk like product according to the present invention also comprises at least one bioactive selected from the group consisting of: growth factors, cytokines, probiotics, extracellular vesicles (e.g. milk fat globules and or exosomes), bioactives from exosome (for example miRNA) and secretory IgA. The standard human milk like product according to the present invention is not the product of human breast lactation as occurring in nature.

In another embodiment, the human milk like product according to the present invention can be adapted to specific needs of the infant who will receive it. It may comprise only one and/or a portion of bioactives, macro and micro nutrients which can be typically found in human breast milk of a well-nourished mother. In such embodiment, the human breast milk like product may also be referred to with the term “non-standard human milk like product”. In one embodiment, the non-standard human milk like product according to the present invention comprises one or more of the nutrients or bioactives selected from the group consisting of proteins, peptides, lipids (including linoleic acid and alpha-linolenic acid), carbohydrates

(including human milk oligosaccharides), Vitamins (including Vitamin A, Vitamin D3, Vitamin E, Vitamin K, Thiamin, Riboflavin, Niacin, Vitamin B6, Vitamin B12, Pantothenic acid, folic acid, Vitamin C and Biotin), minerals (including iron, calcium, phosphorus, magnesium, sodium, chloride, potassium, manganese, iodine, selenium, copper and zinc), choline, myoinositol, L-carnitine, growth factors, cytokines, probiotics, extracellular vesicles (e.g. milk fat globules and or exosomes), bioactives from exosome (for example miRNA) and secretory IgA.

Within the context of the present invention, the term “non-modified human milk like product” indicates a human milk like product which is expressed by lactocytes and/or by the mammary gland like organoids generated according to steps A) and B) of the process of the present invention and which is not subject to the further treatment according to optional step C) of the process of present invention. Non-modified human milk like product may comprise both standard and non-standard human milk like products. Non limiting examples of non-standard human milk like products are selected from the group consisting of: supplement, fortifier, and ingredient enriched in only one and/or a portion of bioactives, macro and micro nutrients which can be typically found in human breast milk of a well-nourished mother.

20

Within the context of the present invention, the term “modified human milk like product” indicates a human milk like product which is expressed by lactocytes and/or by the mammary gland like organoids generated according to steps A) and B) of the process of the present invention and which is subject to the further treatment according to optional step C) of the process of present invention.

25

Modified human milk like products may comprise both standard and non-standard human milk like products.

Within the context of the present invention the term “EBs” means “embryoid bodies”.

- 5 Within the context of the present invention the term “mEBs” means “MammoCult medium-cultured embryoid bodies”.

MammoCult medium refers to a serum-free culture medium comprising basal medium, at least one proliferation supplement, heparin and hydrocortisone.

10

Within the context of the present invention the terms “embryoid bodies (EBs)”, “MammoCult medium-cultured embryoid bodies (mEBs)”, “mammospheres” and/or “spheroids” refer to three-dimensional aggregates formed in suspension by pluripotent stem cells (PSC) under step A) of the process of the present invention.

15

The term “infant” in the context of the present invention identifies a child under the age of 12 months, such as under the age of 9 months, particularly under the age of 6 months.

- 20 In the context of the present invention the infant may be any term infant or preterm infant. In an embodiment of the invention, the infant is selected from the group of preterm infants and term infants.

25 The term “term infant” refers to infants born at term or at a gestational age of 37 weeks or more.

The term “preterm infant” refers to infants who are born at a gestational age of less than 37 weeks.

In the context of the present invention, the term “birth weight” means the first weight of the fetus or newborn obtained after birth.

- 5 Within the context of the present invention, the term “low birth weight” means a birth weight of less than 2500 g (up to and including 2499 g).

Within the context of the present invention, the term “very low birth weight” means a birth weight of less than 1500 g (up to and including 1499 g).

10

Within the context of the present invention, the term “extremely low birth weight” means a birth weight of less than 1000 g (up to and including 999 g).

- The term “small for gestational age infant” refers to infants having a birth weight
15 that is more than 2 standard deviations below the mean reference to a birth weight for gestational growth chart or having a birth weight that is less than the 10th percentile of population-based weight data obtained from infants at the same gestational age. The term “small for gestational age infants” includes infants who are small at birth either from a constitutive or genetic origin or, as a consequence
20 of intrauterine growth restriction.

Within the context of the present invention, the term “young children” or “toddler” indicates a child between the age of 1 and 3 years.

- 25 The term “infant formula” as used herein refers to a nutritional composition intended for infants and as defined in *Codex Alimentarius*, (Codex STAN 72-1981) and Infant Specialities (incl. Food for Special Medical Purpose) as defined in *Codex Alimentarius*, (Codex STAN 72-1981). It also refers to a foodstuff intended for particular nutritional use by infants during the first months of life and satisfying by

itself the nutritional requirements of this category of person (Article 2(c) of the European Commission Directive 91/321/EEC 2006/141/EC of 22 December 2006 on infant formulae and follow-on formulae). The infant formulas encompass the starter infant formulas and the follow-up or follow-on formulas. Generally, a starter
5 formula is for infants from birth as breast-milk substitute, and a follow-up or follow-on formula from the 6th month onwards.

The “growing-up milks” (or GUMs) are given from one year onwards. It is generally a milk-based beverage adapted for the specific nutritional needs of young children.
10 They are nutritional compositions used for feeding children from 12 months to 2-3 years old in combination with other foods.

Within the context of the present invention, the term “fortifier” refers to a composition which comprises one or more nutrients having a nutritional benefit for
15 infants or young children.

By the term “milk fortifier”, it is meant any composition used to fortify or supplement either human breast milk, infant formula, growing-up milk or human breast milk fortified with other nutrients. Accordingly, the human milk fortifier of
20 the present invention can be administered after dissolution in human breast milk, infant formula, growing-up milk or human breast milk fortified with other nutrients or otherwise it can be administered as a standalone composition.

When administered as a stand-alone composition, the human milk fortifier of the present invention can be also identified as being a “supplement”. In one embodiment, the milk fortifier of the present invention is a supplement.
25

By the term “human milk fortifier”, it is meant any composition used to fortify or supplement human breast milk, or human breast milk fortified with other nutrients.

The “human milk fortifier” according to the present invention may be intended to be administered to infants who were born preterm, with very low birth weight (VLBW) or with extremely low birth weight (ELBW).

- 5 The milk fortifier according to the present invention may be in powder or liquid form.

Milk fortifier compositions having a liquid form presents some particular benefits. For example, liquid formulations might be more convenient if coupled with a packaging that delivers calibrated drops of a certain weight or volume.

- 10 In addition, liquid formulations are easier to mix with the compositions to be fortified, whereas the powder ones can, in some cases, form lumps.

- Within the context of the present invention, the term ‘increasing differentiation efficiency and maturation of mammalian induced pluripotent stem cell (miPSCs) into
- 15 mammary-gland progenitor cells’ means increasing the proportion of mammary-gland progenitor cells compared to non-mammary-gland progenitor cells, generated from a starting population of miPSCs which have undergone a differentiation protocol. In the present invention, said differentiation protocol includes the use of BMP4. Thus, said increase of the proportion of mammary-gland progenitor cells may
- 20 be compared to the proportion of mammary-gland progenitor cells relative to non-mammary-gland progenitor cells, generated from a starting population of miPSCs which have undergone a differentiation protocol that does not include the use of BMP4 but is otherwise the same.

- 25 Within the context of the present invention, the term ‘mammary-gland progenitor cells’ or similar means cells that express at least two mammary-gland progenitor markers. Said markers include but are not limited to CD49f, EpCAM, MUC1 and GATA3. Conversely, within the context of the present invention, the term ‘non

mammary-gland progenitor cells' or similar means cells that do not express at least two mammary-gland progenitor markers.

Reference herein to EpiCult or EpiCultB medium refers to a serum free culture
5 medium comprising hydrocortisone, insulin, FGF10 and HGF.

A culture medium as disclosed anywhere herein refers to a solid, semi-solid or liquid comprising essential nutrients, designed to support the growth and differentiation of microorganisms. MammoCult medium is one example of a culture medium that
10 may be used in the present invention.

Methods and Uses according to the present invention

Mammalian mammary cell production 15

The present invention relates to methods of producing mammary gland cells using iPSCs that are cultured and differentiated in specific conditions.

It has been surprisingly demonstrated in the present invention that the addition of
20 Bone morphogenetic protein 4 (BMP4) in the methods as described anywhere herein, provides a more homogenous mixture of non-neural ectoderm progenitor cells by increasing the proportion of non-neural ectoderm progenitor cells and decreasing the proportion of neural ectoderm progenitor cells.

25 It has also been demonstrated in the present invention that the addition of BMP4 in the methods as described anywhere herein increases the differentiation efficiency of iPSCs to mammary gland progenitor cells, and thus increasing the number of mammary gland progenitor cells produced by the methods as described anywhere herein. The benefits of this include a higher yield of mammary gland cells.

Thus, the invention provides a method of producing a population of mammary gland cells, comprising:

- i) culturing mammalian induced pluripotent stem cells (miPSCs) in a culture medium comprising bone morphogenic protein 4 (BMP4) to generate embryoid bodies (EBs), and
- ii) growing the EBs to generate a population of mammary cells.

The invention also provides use of BMP4 for increasing the differentiation efficiency of mammalian induced pluripotent stem cell (miPSCs) into mammary-gland progenitor cells in a differentiation protocol.

In some embodiments, the mammary gland cells are human mammary gland cells. In some embodiments, the mammary gland cells form lactocyte mammary-like gland organoids. Said lactocyte mammary-like gland organoids are lactogenic, i.e. are capable of producing milk.

BMP4 is added to the culture medium in the early stages of the differentiation methods as described anywhere herein. In some embodiments, BMP4 is added to the culture medium between day 0 and day 10. In some embodiments, BMP4 is added to the culture medium between day 0 and day 3. Day 0 is the time point where the iPSCs are first added to the culture medium, i.e. when the iPSCs are first induced for differentiation.

In some embodiments, BMP4 is added to the culture medium for 3 days.

In some embodiments, BMP4 is added to the culture medium in a concentration of 5 to 20 ng/mL. In some embodiments, BMP4 is added to the culture medium in a

concentration of 5 ng/mL. In some embodiments, BMP4 is added to the culture medium in a concentration of 10 ng/mL. In some embodiments, BMP4 is added to the culture medium in a concentration of 20 ng/mL.

- 5 In some embodiments, BMP4 is added to the culture medium between day 0 and day 3, and in a concentration of between 5 to 20 ng/mL.

The EBs generated in the methods as described anywhere herein express one or more mammary gland positive progenitor-cell markers. In some embodiments, said
10 one or more mammary gland positive progenitor-cell markers are selected from EpCAM, CD49f, MUC1 and GATA3.

In some embodiments, the EBs have increased expression of one or more mammary gland positive progenitor-cell markers compared to the expression level
15 of said mammary gland positive progenitor-cell markers in EBs not treated with BMP4.

In some embodiments, at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80% or more of the EBs express one or more mammary gland positive progenitor-cell markers.
20 In some embodiments, at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80% or more of the EBs express two or more mammary gland positive progenitor-cell markers. In some embodiments, said mammary gland positive progenitor-cell markers are selected from EpCAM, CD49f, MUC1 and GATA3.

25 In some embodiments, at least 50% of EBs express EpCAM and CD49f mammary gland positive progenitor-cell markers. In some embodiments, BMP4 is added to the culture medium in a concentration of at least 5 ng/mL and at least 50% of EBs express EpCAM and CD49f mammary gland positive progenitor-cell markers. In

some embodiments, at least 75% of EBs express EpCAM and CD49f mammary gland positive progenitor-cell markers. In some embodiments, BMP4 is added to the culture medium in a concentration of at least 20 ng/mL and at least 75% of EBs express EpCAM and CD49f mammary gland positive progenitor-cell markers.

5

In some embodiments, at least 15% of EBs express MUC1 and CD49f mammary gland positive progenitor-cell markers. In some embodiments, BMP4 is added to the culture medium in a concentration of at least 5 ng/mL and at least 15% of EBs express MUC1 and CD49f mammary gland positive progenitor-cell markers. In

10 some embodiments, at least 20% of EBs express MUC1 and CD49f mammary gland positive progenitor-cell markers. In some embodiments, BMP4 is added to the culture medium in a concentration of at least 20 ng/mL and at least 20% of EBs express MUC1 and CD49f mammary gland positive progenitor-cell markers.

15 In some embodiments, at least 20% of EBs express MUC1 and EpCAM mammary gland positive progenitor-cell markers. In some embodiments, BMP4 is added to the culture medium in a concentration of at least 5 ng/mL and at least 20% of EBs express MUC1 and EpCAM mammary gland positive progenitor-cell markers. In some embodiments, at least 35% of EBs express MUC1 and EpCAM mammary gland positive progenitor-cell markers. In some embodiments, BMP4 is added to the culture medium in a concentration of at least 20 ng/mL and at least 35% of EBs express MUC1 and EpCAM mammary gland positive progenitor-cell markers.

20 In some embodiments, at least 15% of EBs express GATA3 mammary gland positive progenitor-cell markers. In some embodiments, BMP4 is added to the culture medium in a concentration of at least 20 ng/mL and at least 15% of EBs express GATA3 mammary gland positive progenitor-cell markers.

The EBs also express non-neuronal ectodermal markers, thereby demonstrating enrichment towards the non-neuronal lineage. In some embodiments, said one or more non-neuronal ectodermal markers are selected from TFAP2A and TFAP2C.

- 5 In some embodiments, the EBs have increased expression of one or more non-neuronal ectodermal markers of at least 2-fold compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4.

- 10 In some embodiments, the EBs have increased expression of one or more non-neuronal ectodermal markers of at least 3 to 15-fold compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4.

- 15 In some embodiments, the EBs have increased expression of the non-neuronal ectodermal marker TFAP2A of at least 2-fold compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4. In some embodiments, the EBs have increased expression of the non-neuronal ectodermal marker TFAP2C of at least 2-fold compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4.

- 20 In some embodiments, the EBs have decreased expression of one or more neuronal ectodermal markers. In some embodiments, said one or more neuronal ectodermal markers are selected from PAX6, OTX2 and SOX11.

- 25 The methods as described anywhere herein therefore provide a more homogeneous cell population in terms of the types of cells present in the cell population, i.e., the cell population comprises a more homogenous population of non-neuronal ectodermal lineage cells.

In some embodiments, the EBs have decreased expression of one or more neuronal ectodermal markers of at least 0.5-fold compared to the expression level of said neuronal ectodermal markers in EBs not treated with BMP4, i.e., the expression level of these markers is reduced by half. In some embodiments, said
5 neuronal ectodermal markers are selected from PAX6, OXT2 and SOX11.

In some embodiments, the EBs express one or more milk-specific bioactive markers. In some embodiments, said milk-specific bioactive marker is osteopontin (OPN).

10

In some embodiments, the EBs have increased expression of one or more milk-specific bioactive markers compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4.

15 In some embodiments, the EBs have increased expression of one or more milk-specific bioactive markers of at least 2-fold compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4.

In some embodiments, the EBs have increased expression of OPN of at least 2-fold
20 compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4. In some embodiments, BMP4 is added to the culture medium in a concentration of 5 to 20 ng/mL and the EBs have increased expression of OPN of at least 2-fold compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4.

25

In some embodiments, the EBs have increased expression of OPN of at least 4-fold compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4.

In some embodiments, the EBs have increased expression of OPN of at least 18-fold compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4.

5

In some embodiments, the mammary gland cells generate increased expression of milk-specific bioactive markers compared to mammary gland cells that are not treated with BMP4. In some embodiments, the mammary gland cells generate at least a 2-fold increase in expression of milk-specific bioactive markers compared to
10 mammary gland cells that are not treated with BMP4. In some embodiments, said markers are selected from estrogen related receptor alpha (ESRRA), Keratin 14 (KRT14) and MUC1. In some embodiments, the mammary gland cells generate at least a 2-fold increase in expression ESRRA compared to mammary gland cells that are not treated with BMP4. In some embodiments, the mammary gland cells
15 generate at least a 2-fold increase in expression KRT14 compared to mammary gland cells that are not treated with BMP4. In some embodiments, the mammary gland cells generate at least a 2-fold increase in expression MUC1 compared to mammary gland cells that are not treated with BMP4.

20 The methods of producing a population of mammary gland cells as described anywhere herein, may be combined and/or utilized in the methods for producing a mammalian milk like product as described anywhere herein, particular as part of Step A) as described anywhere herein.

25 For example, in some embodiments of the method of producing a population of mammary gland cells as described anywhere herein, the culturing step i) comprises culturing the miPSCs in MammoCult medium and BMP4, in a 3D-suspension culture

system, for example 3D-suspension condition, thereby directing the iPSCs to differentiate towards non-neural ectoderm cells, optionally for at least 12 days.

5 In some embodiments in some embodiments of the method of producing a population of mammary gland cells as described anywhere herein, the growing step ii) comprises growing the formed EBs in a 3D embedding system, for example a mixed floating gel composed of matrix protein such as Matrigel and/or Collagen I for at least 30 days, for example for 32 days, to generate lactocytes.

10 In some embodiments of the method of producing a population of mammary gland cells as described anywhere herein, the culturing step i) comprises culturing the miPSCs in MammoCult medium and BMP4, in a 3D-suspension culture system, for example 3D-suspension condition, thereby directing the iPSCs to differentiate towards non-neural ectoderm cells, optionally for at least 12 days, and the growing
15 step ii) comprises growing the formed EBs in a 3D embedding system, for example a mixed floating gel composed of matrix protein such as Matrigel and/or Collagen I for at least 30 days, for example for 32 days, to generate lactocytes.

Mammalian milk like product production

20

The present invention also relates to methods for producing a mammalian milk like product as defined herein, including any of steps A) and B) as defined herein and optional step C) as defined herein. Said methods for producing a mammalian milk like product as defined herein may also include the methods of producing
25 mammalian mammary cells, as part of step A). In particular, BMP4 may be added to any of the methods for producing a mammalian milk like product as defined herein, particularly as part of step A).

Step A - Generating lactocytes and/or mammary like organoids from hiPSCs

According to the method of the present invention, mammary like cells and/or organoid structures are generated under step A).

- 5 Such mammary like cells and/or organoid structures can be generated according to any reported method making use of iPSC.

In one embodiment, such mammary like cells and/or organoid structures may be generated according to the procedure described in Ying Qu et al., Stem Cell Reports,
10 Vol.8, 205-215 which is hereby incorporated in its entirety.

More precisely, the methodology described in the above-mentioned scientific publication (herein-below also referred to as "Ying Qu publication", or Ying Qu, et al. (2017)) represents a two-step protocol to generate human mammary like cells
15 and/or organoids from iPSCs.

It preferably includes as a first step (step 1) the differentiation and enrichment of non-neural ectoderm-cell-containing spheres (mEBs/mammospheres) from iPSCs and as a second step (step 2) the generation of mammary like organoids from 10-day mEBs (mammospheres) using 3D floating mixed gel-culture of Matrigel and
20 Collagen I.

In step 1 differentiation and enrichment of non-neural ectoderm-cell-containing spheres (mEBs/mammospheres) from hiPSCs occurs by culturing hiPSCs in complete
25 MammoCult medium (StemCell Technologies). Complete MammoCult medium is preferably composed of the basal medium, proliferation supplements, heparin (typically 4µg/mL), and hydrocortisone (typically 0.48µg/mL). Medium is usually

changed every three days. mEBs (mammospheres) obtained in said step are then enriched for non-neural ectoderm cells.

In some embodiments, BMP4 is added to the culture medium in step 1. This is to
5 increase the proportion of mammary gland progenitor cells as described herein.

BMP4 is added to the culture medium as described in anywhere herein.

BMP4 is added to the culture conditions in the early stages of the differentiation
10 methods as described anywhere herein. In some embodiments, BMP4 is added to the culture medium between day 0 and day 10. In some embodiments, BMP4 is added to the culture medium between day 0 and day 3. Day 0 is the time point where the iPSCs are first added to the culture medium, i.e. when the iPSCs are first induced for differentiation.

15

In some embodiments, BMP4 is added to the culture medium for 3 days.

In some embodiments, BMP4 is added to the culture medium in a concentration of 5 to 20 ng/mL. In some embodiments, BMP4 is added to the culture medium in a
20 concentration of 5 ng/mL. In some embodiments, BMP4 is added to the culture medium in a concentration of 10 ng/mL. In some embodiments, BMP4 is added to the culture medium in a concentration of 20 ng/mL.

In some embodiments, BMP4 is added to the culture medium between day 0 and
25 day 3, and in a concentration of between 5 to 20 ng/mL.

In step 2, following the protocol of Ying Qu, et al. (2017), mammary-like organoids are generated by firstly preparing a 3D culture on basis of a floating mixed gel (e.g.

Matrigel and Collagen I). 10 day mEBs (mammospheres) are then grown for 5 days in the mixed gel floated in complete EpiCultB medium supplemented with Parathyroid hormone (pTHrP). For induction of branch and alveolar differentiation for preparation of mammary like organoids/lactocytes, cells are then cultured in
5 complete EpiCultB medium supplemented with hydrocortisone, insulin, FGF10 and HGF. Milk protein expression is typically induced at day 35 by adding prolactin, hydrocortisone and insulin to complete EpiCultB medium supplemented with BSA (lactogenic medium) and culturing for 5 days. The process of Ying Qu, et al. (2017) is typically completed at day 40.

10

In one embodiment of the present invention, a method for producing a human milk like product is provided comprising generating lactocytes under step A) from human induced pluripotent stem cells (hiPSC), where such step A) comprises:

- i) directing iPSCs to differentiate towards non-neural ectoderm cells by culturing
15 them in an appropriate culture medium (for example MammoCult medium) and after 10 days collecting mammospheres formed thereof; and
- ii) growing such mammospheres in an appropriate system (for example a floating mixed gel culture system as described in Hassiotou F. et al. Stem Cells. 2012) for at least 10 days to generate lactocytes.

20

In one embodiment of the present invention, a method for producing a human milk like product is provided comprising generating lactocytes under step A) from human induced pluripotent stem cells (hiPSC), where such step A) comprises:

- i) directing iPSCs to differentiate towards non-neural ectoderm cells by culturing
25 them in an appropriate culture medium (for example MammoCult medium) and BMP4 as described anywhere herein, and after 10 days collecting mammospheres formed thereof; and

ii) growing such mammospheres in an appropriate system (for example a floating mixed gel culture system as described in Hassiotou F. et al. Stem Cells. 2012) for at least 10 days to generate lactocytes.

5 In another embodiment, a method for producing a human milk like product is provided comprising generating lactocytes under step A) from human induced pluripotent stem cells (hiPSC), where such step A) comprises:

- i) directing iPSCs to differentiate towards non-neural ectoderm cells by culturing them in an appropriate culture medium (for example MammoCult medium) in non-
10 adherent conditions for mammospheres formation; and
- ii) growing such mammospheres in a 3D appropriate system (for example a mixed floating gel composed of matrix protein such as Matrigel and/or Collagen or in suspension cultures in non-adherent plates) for at least 10 days to generate lactocytes.

15

In another embodiment, a method for producing a human milk like product is provided comprising generating lactocytes under step A) from human induced pluripotent stem cells (hiPSC), where such step A) comprises:

- i) directing iPSCs to differentiate towards non-neural ectoderm cells by culturing
20 them in an appropriate culture medium (for example MammoCult medium) and BMP4 as described anywhere herein, in non-adherent conditions for mammospheres formation; and
- ii) growing such mammospheres in a 3D appropriate system (for example a mixed floating gel composed of matrix protein such as Matrigel and/or Collagen or in
25 suspension cultures in non-adherent plates) for at least 10 days to generate lactocytes.

In one embodiment, mammary commitment under step A) is obtained by applying a conditioned medium (for example EpiCultB) supplemented with specific factors (for example Parathyroid hormone (pThrP), hydrocortisone, insulin, FGF10, and HGF).

5

In one embodiment of the present invention, the method comprises generating mammary - like organoids under step A).

In one embodiment of the present invention, the method to generate mammary - like organoids under step A) includes culturing the cells under conditions selected from the group consisting of: 2D monolayers of cells, 2D with attached EBs, in suspension in non-adherent plates and in mixed floating gel.

10

In a preferred embodiment, mixed floating gel comprises Matrigel and Collagen I.

In another preferred embodiment, mammospheres (mEBs) in step A) are grown in an appropriate system (for example a floating mixed gel culture system as described in Hassiotou F. et al. Stem Cells. 2012) for at least 15 days.

In a more preferred embodiment, mammospheres (mEBs) in step A) are grown in an appropriate system (for example a floating mixed gel culture system as described in Hassiotou F. et al. Stem Cells. 2012) for 20 days.

In one embodiment, the method according to the present invention provides for culture conditions according to step A) [for example under step A)i) and /or under step A)ii)], which are adapted to generate lactocytes derived from human induced pluripotent stem cells (hiPSC) capable to secrete a human milk like product.

In a preferred embodiment, a method for producing a human milk like product is provided comprising generating lactocytes under step A) from human induced pluripotent stem cells (hiPSC), where such step A) comprises directing hiPSCs to
5 differentiate towards mammary gland cells (for example lactocytes) in an appropriate 3D culture system (for example 3D-suspension condition) for at least 42 days.

In a preferred embodiment, a method for producing a human milk like product is
10 provided comprising generating lactocytes under step A) from human induced pluripotent stem cells (hiPSC), where such step A) comprises directing hiPSCs to differentiate towards mammary gland cells (for example lactocytes) in an appropriate 3D culture system comprising BMP4 as described anywhere herein (for example 3D-suspension condition) for at least 42 days.

15

In another preferred embodiment, a method for producing a human milk like product is provided comprising generating lactocytes under step A) from human induced pluripotent stem cells (hiPSC), where such step A) comprises:

- i) directing hiPSCs to differentiate towards non-neural ectoderm cells by culturing
20 them in an appropriate culture medium (for example MammoCult medium) in an appropriate 3D culture system (for example 3D-suspension condition) for at least 12 days (day -2 to day 10), and
- ii) growing the formed mEBs (mammospheres) in an appropriate 3D embedding system (for example a mixed floating gel composed of matrix protein such as
25 Matrigel and/or Collagen I) for at least 30 days, preferably for 32 days, to generate lactocytes.

In another preferred embodiment, a method for producing a human milk like product is provided comprising generating lactocytes under step A) from human induced pluripotent stem cells (hiPSC), where such step A) comprises:

- i) directing hiPSCs to differentiate towards non-neural ectoderm cells by culturing
5 them in an appropriate culture medium (for example MammoCult medium) and BMP4 as described anywhere herein, in an appropriate 3D culture system (for example 3D-suspension condition) for at least 12 days (day -2 to day 10), and
- ii) growing the formed mEBs (mammospheres) in an appropriate 3D embedding system (for example a mixed floating gel composed of matrix protein such as
10 Matrigel and/or Collagen I) for at least 30 days, preferably for 32 days, to generate lactocytes.

In a particularly preferred embodiment of the present invention, a method of producing a human milk like product is provided comprising generating lactocytes
15 under step A) from human induced pluripotent stem cells (hiPSCs), wherein step A)i) is defined as follows:

- i) generation of embryoid bodies (EBs) from hiPSCs by incubation in standard iPSC medium E8 (comprising DMEM/F12, L-ascorbic acid-2-phosphate magnesium, sodium selenium, FGF2, insulin, NaHCO₃ and transferrin, TGFβ1 or NODAL as
20 described in Chen et al., Nat Methods, 2011) or mTeSR™ for two days (day -2-day 0), and producing mEBs (mammospheres) highly enriched in non-neural ectodermal cells by incubation of EBs in complete MammoCult medium (StemCell Technologies) comprising the basal medium, proliferation supplement and supplemented with heparin (typically 4μg/mL), and hydrocortisone (typically 0.48μg/mL) for 10 days
25 (day 0-day 10), and wherein
step A)ii) is distinguished into further substeps and comprises the following steps:
ii), iii) and iv):

- ii) incubation of mEBs (mammospheres) in complete EpiCultB medium supplemented with EpiCult proliferation supplement and Parathyroid hormone (pThrP) for 5 days (day 10-day 15),
- iii) promotion of branch and alveolar differentiation and mammary cell specification
- 5 by incubating mEBs (mammospheres) in EpiCultB medium supplemented with EpiCult proliferation supplement, hydrocortisone, insulin, FGF10 and HGF for 20 days (day 15-day 35), and
- iv) induction of milk protein expression by incubating mEBs (mammospheres) in EpiCultB medium supplemented with EpiCult proliferation supplement,
- 10 hydrocortisone, insulin, FBS, prolactin, progesterone and β -estradiol for 7 days (day 35-day 42).

In a particularly preferred embodiment of the present invention, a method of producing a human milk like product is provided comprising generating lactocytes

15 under step A) from human induced pluripotent stem cells (hiPSCs), wherein step A)i) is defined as follows:

- i) generation of embryoid bodies (EBs) from hiPSCs by incubation in standard iPSC medium E8 (comprising DMEM/F12, L-ascorbic acid-2-phosphate magnesium, sodium selenium, FGF2, insulin, NaHCO_3 and transferrin, TGF β 1 or NODAL as
- 20 described in Chen et al., Nat Methods, 2011) or mTeSRTM for two days (day -2-day 0), and producing mEBs (mammospheres) highly enriched in non-neural ectodermal cells by incubation of EBs in complete MammoCult medium (StemCell Technologies) comprising the basal medium, proliferation supplement and supplemented with heparin (typically 4 $\mu\text{g}/\text{mL}$), hydrocortisone (typically 0.48 $\mu\text{g}/\text{mL}$) and BMP4 as
- 25 described anywhere herein, for 10 days (day 0-day 10), and wherein step A)ii) is distinguished into further substeps and comprises the following steps: ii), iii) and iv):

- ii) incubation of mEBs (mammospheres) in complete EpiCultB medium supplemented with EpiCult proliferation supplement and Parathyroid hormone (pThrP) for 5 days (day 10-day 15),
- iii) promotion of branch and alveolar differentiation and mammary cell specification
- 5 by incubating mEBs (mammospheres) in EpiCultB medium supplemented with EpiCult proliferation supplement, hydrocortisone, insulin, FGF10 and HGF for 20 days (day 15-day 35), and
- iv) induction of milk protein expression by incubating mEBs (mammospheres) in EpiCultB medium supplemented with EpiCult proliferation supplement,
- 10 hydrocortisone, insulin, FBS, prolactin, progesterone and β -estradiol for 7 days (day 35-day 42).

Step iv) preferably leads to differentiation into milk protein expressing cells, particularly lactocytes, and/or mammary like gland organoids.

15

In a further particularly preferred embodiment of the present invention, a method of producing a human milk like product is provided comprising generating lactocytes under step A) from human induced pluripotent stem cells (hiPSCs), wherein step A)i) is defined as follows:

- 20 i) generation of embryoid bodies (EBs) from hiPSCs by incubation in standard iPSC medium E8 (comprising DMEM/F12, L-ascorbic acid-2-phosphate magnesium, sodium selenium, FGF2, insulin, NaHCO_3 and transferrin, TGF β 1 or NODAL as described in Chen et al., Nat Methods, 2011) mTeSRTM for two days (day-2-day 0), and producing mEBs (mammospheres) highly enriched in non-neural ectodermal
- 25 cells by incubation of EBs in MammoCultB medium supplemented with MammoCult proliferation supplement, hydrocortisone and heparin for 10 days (day 0-day 10), and wherein step A)ii) is distinguished into further substeps and comprises the following steps ii), iii) and iv):

- ii) embedding the formed mEBs (mammospheres) in a mixture of Matrigel and Collagen I floated in EpiCultB medium supplemented with EpiCult proliferation supplement and Parathyroid hormone (pTHrP) for 5 days (day 10-day 15),
- iii) promotion of branch and alveolar differentiation and mammary cell specification
- 5 by incubating embedded mEBs (mammospheres) in EpiCultB medium supplemented with EpiCult proliferation supplement, hydrocortisone, insulin, FGF10 and HGF for 20 days (day 15 to day 35), and
- iv) induction of milk protein expression by incubating mEBs (mammospheres) in EpiCultB medium supplemented with EpiCult proliferation supplement,
- 10 hydrocortisone, insulin, FBS, prolactin, progesterone and β -estradiol for 7 days (day 35 to day 42).

In a further particularly preferred embodiment of the present invention, a method of producing a human milk like product is provided comprising generating lactocytes

15 under step A) from human induced pluripotent stem cells (hiPSCs), wherein step A)i) is defined as follows:

- i) generation of embryoid bodies (EBs) from hiPSCs by incubation in standard iPSC medium E8 (comprising DMEM/F12, L-ascorbic acid-2-phosphate magnesium, sodium selenium, FGF2, insulin, NaHCO_3 and transferrin, TGF β 1 or NODAL as
- 20 described in Chen et al., Nat Methods, 2011) mTeSRTM for two days (day-2-day 0), and producing mEBs (mammospheres) highly enriched in non-neural ectodermal cells by incubation of EBs in MammoCultB medium supplemented with MammoCult proliferation supplement, hydrocortisone, heparin and BMP4 as described anywhere herein, for 10 days (day 0-day 10), and wherein step A)ii) is distinguished
- 25 into further substeps and comprises the following steps ii), iii) and iv):
- ii) embedding the formed mEBs (mammospheres) in a mixture of Matrigel and Collagen I floated in EpiCultB medium supplemented with EpiCult proliferation supplement and Parathyroid hormone (pTHrP) for 5 days (day 10-day 15),

- iii) promotion of branch and alveolar differentiation and mammary cell specification by incubating embedded mEBs (mammospheres) in EpiCultB medium supplemented with EpiCult proliferation supplement, hydrocortisone, insulin, FGF10 and HGF for 20 days (day 15 to day 35), and
- 5 iv) induction of milk protein expression by incubating mEBs (mammospheres) in EpiCultB medium supplemented with EpiCult proliferation supplement, hydrocortisone, insulin, FBS, prolactin, progesterone and β -estradiol for 7 days (day 35 to day 42).
- 10 Step iv) preferably leads to differentiation into milk protein expressing cells, particularly lactocytes, and/or mammary like gland organoids.

Standard iPSC medium E8 (comprising DMEM/F12, L-ascorbic acid-2-phosphate magnesium, sodium selenium, FGF2, insulin, NaHCO₃ and transferrin, TGF β 1 or

15 NODAL as described in Chen et al., Nat Methods, 2011) as mentioned herein is commercially available, e.g., as “Essential 8™ Medium” from ThermoFischer Scientific, catalogue number A1517001 (see also <https://www.thermofisher.com/order/catalog/product/A1517001#/A1517001>).

20 mTeSR™ medium is commercially available from STEMCELL Technologies, catalogue number 85850 (see also <https://www.stemcell.com/mtesr1.html>). Such medium is also described in “Defined, Feeder- Independent medium for human embryonic stem cell culture”, Current protocol in Stem Cell Biology, Volume 2, Issue 1, Sept 2007.

25

In one embodiment, steps iii) and/or iv) as defined above for the particularly preferred embodiments, preferably lead to formation of/differentiation into at least breast cells, luminal cells, and basal cells. In this context, breast cells preferably

express one or more, preferably all of markers selected from the group consisting of: β -Casein, milk protein, and hormone receptors. Moreover, luminal cells preferably express one or more, preferably all markers selected from the group consisting of: EpCAM, MUC1, CD49F, GATA3, CK8, and CK18. Moreover, basal cells
5 preferably express one or more markers selected from the group consisting of: CK14, α -smooth muscle actin and P63.

In one further embodiment, after induction of mEBs (mammospheres) in step ii) and/or iv) as defined above for the particularly preferred embodiments, mammary
10 like gland organoids may be obtained, that express one or more markers selected from the group consisting of: β -Casein, milk protein, and hormone receptors, luminal cells that express one or more markers selected from the group consisting of: EpCAM, MUC1, CD49F, GATA3, CK8, CK18, and basal cells that express one or more markers selected from the group consisting of: CK14, α -smooth muscle actin
15 and P63.

In one embodiment of the invention, the methods above described are provided for producing a human milk like product.

20 In one embodiment (of step A), delivery of nutrients and biomimetic stimuli is controlled to influence cell growth, differentiation and tissue formation. In one embodiment (of step A), such control is performed in a bioreactor.

Where BMP4 is added to the culture medium in the methods for producing a
25 mammalian milk like product as described anywhere herein, the EBs generated in the methods as described anywhere herein express one or more mammary gland positive progenitor-cell markers. In some embodiments, said one or more mammary

gland positive progenitor-cell markers are selected from EpCAM, CD49f, MUC1 and GATA3.

In some embodiments, the EBs have increased expression of one or more
5 mammary gland positive progenitor-cell markers compared to the expression level of said mammary gland positive progenitor-cell markers in EBs not treated with BMP4.

In some embodiments, at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80% or more
10 of the EBs express one or more mammary gland positive progenitor-cell markers. In some embodiments, at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80% or more of the EBs express two or more mammary gland positive progenitor-cell markers. In some embodiments, said mammary gland positive progenitor-cell markers are selected from EpCAM, CD49f, MUC1 and GATA3.

15 In some embodiments, at least 50% of EBs express EpCAM and CD49f mammary gland positive progenitor-cell markers. In some embodiments, BMP4 is added to the culture medium in a concentration of at least 5 ng/mL and at least 50% of EBs express EpCAM and CD49f mammary gland positive progenitor-cell markers. In
20 some embodiments, at least 75% of EBs express EpCAM and CD49f mammary gland positive progenitor-cell markers. In some embodiments, BMP4 is added to the culture medium in a concentration of at least 20 ng/mL and at least 75% of EBs express EpCAM and CD49f mammary gland positive progenitor-cell markers.

25 In some embodiments, at least 15% of EBs express MUC1 and CD49f mammary gland positive progenitor-cell markers. In some embodiments, BMP4 is added to the culture medium in a concentration of at least 5 ng/mL and at least 15% of EBs express MUC1 and CD49f mammary gland positive progenitor-cell markers. In

some embodiments, at least 20% of EBs express MUC1 and CD49f mammary gland positive progenitor-cell markers. In some embodiments, BMP4 is added to the culture medium in a concentration of at least 20 ng/mL and at least 20% of EBs express MUC1 and CD49f mammary gland positive progenitor-cell markers.

5

In some embodiments, at least 20% of EBs express MUC1 and EpCAM mammary gland positive progenitor-cell markers. In some embodiments, BMP4 is added to the culture medium in a concentration of at least 5 ng/mL and at least 20% of EBs express MUC1 and EpCAM mammary gland positive progenitor-cell markers. In

10 some embodiments, at least 35% of EBs express MUC1 and EpCAM mammary gland positive progenitor-cell markers. In some embodiments, BMP4 is added to the culture medium in a concentration of at least 20 ng/mL and at least 35% of EBs express MUC1 and EpCAM mammary gland positive progenitor-cell markers.

15 In some embodiments, at least 15% of EBs express GATA3 mammary gland positive progenitor-cell markers. In some embodiments, BMP4 is added to the culture medium in a concentration of at least 20 ng/mL and at least 15% of EBs express GATA3 mammary gland positive progenitor-cell markers.

20 The EBs also express non-neuronal ectodermal markers, thereby demonstrating enrichment towards the non-neuronal lineage. In some embodiments, said one or more non-neuronal ectodermal markers are selected from TFAP2A and TFAP2C.

In some embodiments, the EBs have increased expression of one or more non-
25 neuronal ectodermal markers of at least 2-fold compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4.

In some embodiments, the EBs have increased expression of one or more non-neuronal ectodermal markers of at least 3 to 15-fold compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4.

- 5 In some embodiments, the EBs have increased expression of the non-neuronal ectodermal marker TFAP2A of at least 2-fold compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4. In some embodiments, the EBs have increased expression of the non-neuronal ectodermal marker TFAP2C of at least 2-fold compared to the expression level of said non-
10 neuronal ectodermal markers in EBs not treated with BMP4.

In some embodiments, the EBs have decreased expression of one or more neuronal ectodermal markers. In some embodiments, said one or more neuronal ectodermal markers are selected from PAX6, OTX2 and SOX11.

15

The methods as described anywhere herein therefore provide a more homogeneous cell population in terms of the types of cells present in the cell population, i.e., the cell population comprises a more homogenous population of non-neuronal ectodermal lineage cells.

20

In some embodiments, the EBs have decreased expression of one or more neuronal ectodermal markers of at least 0.5-fold compared to the expression level of said neuronal ectodermal markers in EBs not treated with BMP4, i.e., the expression level of these markers is reduced by half. In some embodiments, said
25 neuronal ectodermal markers are selected from PAX6, OXT2 and SOX11.

In some embodiments, the EBs express one or more milk-specific bioactive markers. In some embodiments, said milk-specific bioactive marker is osteopontin (OPN).

- 5 In some embodiments, the EBs have increased expression of one or more milk-specific bioactive markers compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4.

- 10 In some embodiments, the EBs have increased expression of one or more milk-specific bioactive markers of at least 2-fold compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4.

- 15 In some embodiments, the EBs have increased expression of OPN of at least 2-fold compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4. In some embodiments, BMP4 is added to the culture medium in a concentration of 5 to 20 ng/mL and the EBs have increased expression of OPN of at least 2-fold compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4.

- 20 In some embodiments, the EBs have increased expression of OPN of at least 4-fold compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4.

- 25 In some embodiments, the EBs have increased expression of OPN of at least 18-fold compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4.

In some embodiments, the mammary gland cells generate increased expression of milk-specific bioactive markers compared to mammary gland cells that are not treated with BMP4. In some embodiments, the mammary gland cells generate at least a 2-fold increase in expression of milk-specific bioactive markers compared to

5 mammary gland cells that are not treated with BMP4. In some embodiments, said markers are selected from estrogen related receptor alpha (ESRRA), Keratin 14 (KRT14) and MUC1. In some embodiments, the mammary gland cells generate at least a 2-fold increase in expression ESRRA compared to mammary gland cells that are not treated with BMP4. In some embodiments, the mammary gland cells

10 generate at least a 2-fold increase in expression KRT14 compared to mammary gland cells that are not treated with BMP4. In some embodiments, the mammary gland cells generate at least a 2-fold increase in expression MUC1 compared to mammary gland cells that are not treated with BMP4.

15 Thus, as an example, provided herein is a method for producing a human milk like product, comprising generating lactocytes under step A) from human induced pluripotent stem cells (hiPSC), where such step A) comprises:

i) directing iPSCs to differentiate towards non-neural ectoderm cells by culturing them in an appropriate culture medium (for example MammoCult medium) and

20 BMP4 as described anywhere herein, and after 10 days collecting mammospheres formed thereof; and

ii) growing such mammospheres in an appropriate system (for example a floating mixed gel culture system as described in Hassiotou F. et al. Stem Cells. 2012) for at least 10 days to generate lactocytes,

25 wherein the mammospheres (EBs) have increased expression of one or more mammary gland positive progenitor-cell markers compared to the expression level of said mammary gland positive progenitor-cell markers in mammospheres (EBs) not treated with BMP4.

It will be understood that any method or method step disclosed herein may be conducted in 3D suspension culture rather than using a membrane matrix as a support. Thus, in some embodiments, during all the differentiation procedure, cells are maintained in suspension culture.

5

Step B - Expressing a human breast milk like product

In one embodiment of the present invention, the methods comprise expressing the human milk like product from mammary like organoids derived from human induced pluripotent stem cells (hiPSCs), preferably prepared according to step A). Expressing
10 human milk like products preferably occurs upon induction of expression of the human milk like product from such lactocytes and/or mammary-like gland organoids.

In one embodiment, lactating lactocytes are induced by applying a specific medium (for example EpiCultB) supplemented with lactogenic factors (for example prolactin,
15 hydrocortisone, and insulin).

Particularly, the human milk like product obtained from mammary like organoids derived from human induced pluripotent stem cells (hiPSCs), preferably prepared according to step A), contains bioactives of human milk, selected from the group
20 comprising or consisting of proteins, lipids or oligosaccharides, preferably human milk oligosaccharides, etc. Inventors particularly managed to identify with the particularly preferred protocol according to steps A i) to iv) as carried out above inter alia oligosaccharides (including lactose and some HMOs), lipids (including 4 fatty acids), proteins (7 detected including caseins), and miRNA (75 detected, including
25 11 typically detected in HBM).

In one embodiment, the human milk like product obtained from mammary like organoids derived from human induced pluripotent stem cells (hiPSCs), preferably

prepared according to step A), contains bioactives of human milk, selected from the group comprising or consisting of: oligosaccharides, lipids, proteins, exosomes and miRNA.

- 5 In another embodiment, human milk like product obtained from mammary like organoids derived from human induced pluripotent stem cells (hiPSCs), preferably prepared according to step A), contains bioactives of human milk, selected from the group comprising or consisting of: lactose, 6'SL, C-4:0 fatty acid, C-8:0 fatty acid, C-10:0 fatty acid, C-14:0 fatty acid, C-15:0 fatty acid, C-16:0 fatty acid, C-16:1n7 fatty acid, C-17:0 fatty acid, C-18:0 fatty acid, C-18:1 n9 fatty acid, C-18:1 fatty acid, C-18:2 n6 fatty acid, C-20:0 fatty acid, C-20:1 n9 fatty acid, C-18:3 n3 fatty acid, C-22:0 fatty acid, lactoferrin, albumin, prolactin, Alpha S1-casein, Hemoglobin subunit beta, Hemoglobin subunit alpha, α -lactalbumin, Alpha-2-macroglobulin, β -casein, bile salt-activated lipase, κ -casein, lactadherin, CD14, fatty acid synthase, IgA, plgR, Serum albumin, Xanthine dehydrogenase, exosomes, miR-21-5p, miR-181a-5p, miR-30d-5p, miR-30b-5p, miR-22-3p, miR-146b-3p, miR-30c-5p, miR-30a-5p, miR-30e-5p and miR-148b-3p.

Step C - Further treatments to produce modified human breast milk like product

- 20 In one optional embodiment of the present invention, the herein-described methods comprise an additional step C) which is performed on the human milk like product obtainable from step B) and which comprises performing an additional treatment on such product to provide a modified human milk like product.
- 25 In one particular embodiment, the additional treatment step C) performed on the inventive human breast milk like product may be selected from the group consisting of: a purification step, an isolation process, an extraction process, a fractionation step, an enrichment process, an enzymatic treatment, the addition of further

components (for example which can't be expressed by the human mammary gland organoid (such as for example Immunoglobulins, probiotic and/or minerals) or combinations thereof.

5 Human milk like products

'Standard' human breast milk like product

In one embodiment of the present invention, the human breast milk like product is a 'standard' human breast milk like product, i.e., comprises the same components
10 as human breast milk of a well-nourished mother.

The benefits of breast feeding are well known in the scientific literature and the possibility to have access to human breast milk like product allows its use for a number of equally well-known health benefits.

15

In such an embodiment, the human breast milk like product can be used as a substitute of breastfeeding under circumstances where real breastfeeding is not possible.

20 In such embodiment, the human breast milk like product is intended to be used for example to support longer breastfeeding experience for women who have less milk or who stop to produce milk after 6 months from birth.

Similarly, the human breast milk like product is intended to be used for example to
25 allow breastfeeding even under circumstances where sicknesses compromise real breastfeeding from the mother.

In another embodiment, the human breast milk like product is intended to be used under circumstances whereby breastmilk production would not naturally be initiated, for example if an infant is adopted.

- 5 In one embodiment, the human milk like product according to the present invention is not the product of human breast milk lactation as occurring in nature.

In one embodiment, the human breast milk like product is for use in providing optimal nutrition for infant.

10

In one embodiment, the human breast milk like product is for use in providing healthy growth in infants.

- 15 In one embodiment, the human breast milk like product is for use in preventing infection, obesity and promoting immunity development in infants.

In one embodiment, the human breast milk like product is a non-modified human breast milk like product.

- 20 In another embodiment, the human breast milk like product is a modified human breast milk like product.

In one embodiment, the human milk like product according to the present invention comprises: proteins, lipids, carbohydrates, vitamins and minerals.

25

In another embodiment, the human milk like product according to the present invention comprises: proteins, lipids, carbohydrates, vitamins, minerals and bioactives.

In one embodiment, the human milk like product according to the present invention comprises: proteins, lipids (including linoleic acid and alpha-linolenic acid), carbohydrates, Vitamins (including Vitamin A, Vitamin D3, Vitamin E, Vitamin K, Thiamin, Riboflavin, Niacin, Vitamin B6, Vitamin B12, Pantothenic acid, folic acid, Vitamin C and Biotin), minerals (including iron, calcium, phosphorus, magnesium, sodium, chloride, potassium, manganese, iodine, selenium, copper and zinc), choline, myoinositol and L-carnitine.

10 In a further embodiment, the human milk like product according to the present invention also comprises at least one bioactive selected in the group consisting of: growth factors, cytokines, probiotics, extracellular vesicles (e.g. milk fat globules and or exosomes), bioactives from exosomes (for example miRNA) and secretory IgA.

15 Such human breast milk like product may be prepared according to the method of the present invention for example by including a step C) of addition of growth factors, cytokines, probiotics, extracellular vesicles (e.g. milk fat globules and or exosomes), bioactives from exosomes (for example miRNA) and secretory IgA.

20 In one embodiment, the human breast milk like product contains probiotics.

Such human breast milk like product may be prepared according to the method of the present invention for example by including a step C) of addition of probiotics (for example *B.Lactis*, *B.Infantis*, *L. Ramnhusus*) which can be obtained from several commercially available sources.

In such embodiment, the human breast milk like product may be used for optimizing gastro intestinal function and/or promoting Immunity.

In one embodiment, the human breast milk like product contains secretory IgA and probiotics.

- 5 Such human breast milk like product may be prepared according to the method of the present invention for example by including a step C) of addition of a combination of probiotics and secretory IgA which may be prepared as described for example in patent applications WO2009/156301 and WO2009/156367 which are hereby incorporated by reference. In such embodiment, the human breast milk like product
- 10 may be used for preventing Immunoglobulin deficiency and/or in the prevention of recurrent infection in infants and young children.

'Non-standard' human breast milk like product

- 15 In one embodiment of the present invention, the human milk like product can have altered ratios and concentrations of components found naturally in human breast milk of a well-nourished mother. This is referred to herein as a "non-standard milk like product".
- 20 In one embodiment, the human milk like product according to the present invention may be selected from the group consisting of a milk fortifier, a supplement, and/or a human breast milk replacer adapted for special purposes.

Human Milk fortifiers and Human milk bioactive supplements

- 25 In one embodiment, the method of the present invention provides for a human breast milk like product which may be used to fortify human breast milk naturally obtained from a nursing mother or to fortify infant formulas.

In another embodiment, the method of the present invention provides for a human breast milk like product which may be used as a supplement for infants or young children in need thereof.

- 5 In such embodiments the human breast milk like products may be used for providing healthy growth and/or to reduce the risk of developing a disease typically associated to specific conditions in an infant or young child (such as for example asthma, allergy, cognitive alterations) and /or to promote catch up growth, development of immunity, protection from infections.

10

Remarkably, the human origin of the constituents (especially bioactive constituents) in such fortifiers or supplements combined with the fact that they are according to the method of the invention, is supposed to provide to such constituents an intact or higher functionality.

15

The human breast milk like product is preferably intended to be used as a fortifier. Such human breast milk like product intended to be used as a fortifier and may be prepared according to the method of the present invention for example by including a step C) of isolation and/or enrichment of (certain) bioactives from the human
20 breast milk like product obtainable from step B). Such isolation step may be performed via classical fractionation, enrichment and/or purification of the non-modified human breast milk like product obtainable from step B).

- 25 The human breast milk like product intended to be used as a supplement may comprise one or more bioactives selected from the group consisting of: human milk oligosaccharides (for example 2FL, 3FL, LNT, LnNT, DiFI, 6SL and /or 3SL), lipids, growth factors (for example epidermal growth factor (EGF), heparin binding epidermal growth factor), cytokines (for example transforming growth factor -beta

2 (TGFbeta-2), IL-1. IL-2, IL-6, IL-10, IL-18, interferon gamma (INF-gamma), TNF-alpha), extracellular vesicles (e.g. milk fat globules and or exosomes), exosome comprising microRNAs and antimicrobial/protecting bioactives (for example IgA, lactoferrin, lysozyme, lactadherine). Such human breast milk like product intended
5 to be used as a supplement may be prepared according to the method of the present invention for example by including a step C) of isolation of the bioactives from the non-modified human breast milk like product obtainable from step B). Such isolation step may be performed via classical fractionation, enrichment and/or purification of the non -modified human breast milk like product obtainable from step B).

10

In one embodiment, the human breast milk like product is a supplement or milk fortifier which contains fucosylated human milk oligosaccharides, for example 2FL and/or 3FL. Such supplement or milk fortifier is for use in completing the profile of human breast milk of women who do not secrete fucosylated oligosaccharides
15 because of the inactivity of their FUT2 gene.

Such human breast milk like product intended to be used as a fortifier or supplement may be prepared according to the method of the present invention for example by including a step C) of isolation and/or enrichment of fucosylated oligosaccharides
20 (for example 2FL and or 3FL) from the non-modified human breast milk like product obtainable from step B).

In such an embodiment, the human breast milk like product may be used for optimizing gastro intestinal function and/or promoting Immunity.

25

Human breast milk like product for infants with genetic diseases

In one embodiment, the human breast milk like product according to the present invention may be adapted to address the specific need of infants who are born with a genetic disease.

5 Galactossemia

In such embodiment, the human breast milk like product may be adapted to the needs on infants suffering from Galactossemia. Galactossemia is a rare genetic disease that affects babies' ability to metabolize galactose.

10

In such embodiment, the human breast milk like product should be deprived of lactose and/or lactose containing saccharides. In such embodiment human breast milk like product may be used for providing healthy growth to the infants affected by galactossemia.

15

In one embodiment, a human breast milk like product deprived of lactose and/or lactose containing saccharides may be obtained according to the method of the present invention by including a step C) of enzymatic treatment (lactase treatment), or of membrane fractionation and ultrafiltration of the non-modified human breast milk like product obtainable from step B).

20

In another embodiment, a human breast milk like product deprived of lactose and/or lactose containing saccharides may be obtained according to the method of the present invention by using under step A) GMO alpha-lactalbumin deficient human cells to generate hiPSCs.

25

Phenyl Keturonia

In such an embodiment, the human breast milk like product may be adapted to the needs on infants suffering from Phenyl Keturonia (PKU). PKU is due to absent or dysfunctional phenylalanine hydroxylase, which converts phenylalanine to tyrosine. Untreated, it leads to severe mental retardation due to brain toxicity.

5

In such an embodiment, the human breast milk like product should be deprived or depleted of phenylalanine.

In such an embodiment, the human breast milk like product may be used for providing healthy growth to the infants affected by PKU.

10

In one embodiment, the human breast milk like product is depleted of phenylalanine in such a way that phenylalanine content is kept below 20 mg/kg body weight of the subject receiving it.

15

In one embodiment, a human breast milk like product depleted or deprived of phenylalanine may be obtained according to the method of the present invention by including a step C) of enzymatic treatment (protein hydrolysis) or of filtration of the non -modified human breast milk like product obtainable from step B).

20

In one embodiment, a human breast milk like product depleted of phenylalanine may be obtained according to the method of the present invention by including a step C) of enzymatic treatment (protein hydrolysis) or of filtration of the non -modified human breast milk like product obtainable from step B).

25

In another embodiment, a human breast milk like product depleted of phenylalanine may be obtained according to the method of the present invention by providing in step B) a culture medium providing limited or zero amounts of

phenylalanine, such as for example a culture medium containing Glycomacropeptide (GMP) from whey.

Additional Embodiments of the Invention

5

There is provided:

S1. A method of producing a population of mammary gland cells, comprising:

10 i) culturing mammalian induced pluripotent stem cells (miPSCs) in a culture medium comprising bone morphogenic protein 4 (BMP4) to generate embryoid bodies (EBs), and

ii) growing the EBs to generate a population of mammary cells.

15 S2. The method of statement 1, wherein the culturing step i) comprises culturing the miPSCs in MammoCult medium and BMP4, in a 3D-suspension culture system, for example 3D-suspension condition, thereby directing the iPSCs to differentiate towards non-neural ectoderm cells, optionally for at least 12 days.

20 S3. The method of statements 1 or 2, wherein the growing step ii) comprises growing the formed EBs in a 3D embedding system, for example a mixed floating gel composed of matrix protein such as Matrigel and/or Collagen I for at least 30 days, for example for 32 days, to generate lactocytes.

25 S4. The method of any one of statements 1 to 3, wherein the mammary gland cells are human mammary gland cells.

S5. The method of any one of statements 1 to 4, wherein BMP4 is added to the culture medium between day 0 and day 10, preferably between day 0 and day 3, where day 0 is the time point where the iPSCs are first added to the culture medium.

5 S6. The method of any one of statements 1 to 5, wherein BMP4 is added to the culture medium for 3 days.

S7. The method of any one of statements 1 to 6, wherein BMP4 is added to the culture medium in a concentration of 5 to 20 ng/mL, preferably 5 ng/mL, 10 ng/mL
10 or 20 ng/mL.

S8. The method of any one of statements 1 to 7, wherein the EBs express one or more mammary gland positive progenitor-cell markers, optionally selected from EpCAM, CD49f, MUC1 and GATA3.

15

S9. The method of any one of statements 1 to 8, wherein the EBs have increased expression of one or more mammary gland positive progenitor-cell markers compared to the expression level of said mammary gland positive progenitor-cell markers in EBs not treated with BMP4.

20

S10. The method of any one of statements 1 to 9, wherein at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80% or more of the EBs express one or more mammary gland positive progenitor-cell markers, optionally wherein at least 50% of EBs express EpCAM and CD49f mammary gland positive progenitor-cell markers,
25 optionally wherein at least 15% of EBs express MUC1 and CD49f mammary gland positive progenitor-cell markers, optionally wherein at least 20% of EBs express MUC1 and EpCAM mammary gland positive progenitor-cell markers, and/or

optionally wherein at least 15% of EBs express GATA3 mammary gland positive progenitor-cell markers.

5 S11. The method of any one of statements 1 to 10, wherein the EBs express one or more non-neuronal ectodermal markers, optionally selected from TFAP2A and TFAP2C.

10 S12. The method of any one of statements 1 to 11, wherein the EBs have increased expression of one or more non-neuronal ectodermal markers of at least 2-fold compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4, optionally at least a 3 to 15-fold increase.

15 S13. The method of any one of statements 1 to 12, wherein the EBs have decreased expression of one or more neuronal ectodermal markers of at least 0.5-fold compared to the expression level of said neuronal ectodermal markers in EBs not treated with BMP4, optionally wherein said neuronal ectodermal markers are selected from PAX6, OXT2 and SOX11.

20 S14. The method of any one of statements 1 to 13, wherein the EBs express one or more milk-specific bioactive markers, optionally osteopontin (OPN).

25 S15. The method of any one of statements 1 to 14, wherein the EBs have increased expression of OPN of at least 2-fold compared to the expression level of OPN in EBs not treated with BMP4, optionally at least a 4 to 18-fold increase.

S16. The method of any one of statements 1 to 15, wherein the mammary gland cells form lactocyte mammary-like gland organoids.

S17. The method of any one of statements 1 to 16, wherein the mammary gland cells generate increased expression of milk-specific bioactive markers compared to mammary gland cells that are not treated with BMP4, optionally wherein the mammary gland cells generate at least a 2-fold increase in expression of milk-specific bioactive markers compared to mammary gland cells that are not treated with BMP4, optionally wherein said markers are selected from estrogen related receptor alpha (ESRRA), Keratin 14 (KRT14) and MUC1.

S18. Use of BMP4 for increasing the differentiation efficiency of mammalian induced pluripotent stem cell (miPSCs) into mammary-gland progenitor cells in a differentiation protocol.

S19. A method for producing a mammalian milk like product, comprising:
A) Generating lactocyte mammary-like gland organoids derived from mammalian induced pluripotent stem cells (miPSC);
B) Secreting the mammalian milk like product from said lactocytes,
wherein step A) comprises culturing the miPSCs in a culture medium comprising BMP4.

S20. The method of statement 19, wherein the method is for producing a human milk like product,
wherein step A) further comprises:
i) directing hiPSCs to differentiate towards non-neural ectoderm cells by culturing them in an appropriate culture medium comprising BMP4, for example MammoCult medium and BMP4, in an appropriate 3D culture system, for example 3D-suspension condition, for at least 12 days and
ii) growing the formed mEBs (mammospheres) in an appropriate 3D embedding system, for example a mixed floating gel composed of matrix protein

such as Matrigel and/or Collagen I for at least 30 days, for example for 32 days, to generate lactocytes.

S21. The method according to statement 20 wherein step A)i) is defined as follows:

- 5 i) generation of embryoid bodies (EBs) from hiPSCs by incubation in standard iPSC medium E8 comprising DMEM/F12, L-ascorbic acid-2-phosphate magnesium, sodium selenium, FGF2, insulin, NaHCO₃ and transferrin, TGFβ1 or NODAL or in medium mTeSR™ for two days, and producing mEBs (mammospheres) highly enriched in non-neural ectodermal cells by incubation of EBs in complete
- 10 MammoCult medium comprising the basal medium, proliferation supplement and supplemented with BMP4, heparin, and hydrocortisone for 10 days, and wherein step A)ii) is distinguished into further substeps and comprises the following steps: ii), iii) and iv):
- ii) incubation of mEBs (mammospheres) in complete EpiCultB medium
- 15 supplemented with EpiCult proliferation supplement and Parathyroid hormone (pThrP) for 5 days,
- iii) promotion of branch and alveolar differentiation and mammary cell specification by incubating mEBs (mammospheres) in EpiCultB medium supplemented with EpiCult proliferation supplement, hydrocortisone, insulin, FGF10 and HGF for 20
- 20 days, and
- iv) induction of milk protein expression by incubating mEBs (mammospheres) in EpiCultB medium supplemented with EpiCult proliferation supplement, hydrocortisone, insulin, FBS, prolactin, progesterone and β-estradiol for 7 days.

25 S22. The method according to statement 20 wherein step A)i) is defined as follows:

- i) generation of embryoid bodies (EBs) from hiPSCs by incubation in standard iPSC medium E8 comprising DMEM/F12, L-ascorbic acid-2-phosphate magnesium, sodium selenium, FGF2, insulin, NaHCO₃ and transferrin, TGFβ1 or NODAL for two

- days, and producing mEBs (mammospheres) highly enriched in non-neural ectodermal cells by incubation of EBs in MammoCultB medium supplemented with MammoCult proliferation supplement, hydrocortisone, heparin and BMP4 for 10 days, and wherein step A)ii) is distinguished into further substeps and comprises the
- 5 following steps ii), iii) and iv):
- ii) embedding the formed mEBs (mammospheres) in a mixture of Matrigel and Collagen I floated in EpiCultB medium supplemented with EpiCult proliferation supplement and Parathyroid hormone (pTHrP) for 5 days,
- iii) promotion of branch and alveolar differentiation and mammary cell specification
- 10 by incubating embedded mEBs (mammospheres) in EpiCultB medium supplemented with EpiCult proliferation supplement, hydrocortisone, insulin, FGF10 and HGF for 20 days, and
- iv) induction of milk protein expression by incubating mEBs (mammospheres) in EpiCultB medium supplemented with EpiCult proliferation supplement,
- 15 hydrocortisone, insulin, FBS, prolactin, progesterone and β -estradiol for 7 days.

S23. The method of any one of statements 19 to 22, wherein Step A) comprises a method of any one of statements 1 to 17.

- 20 S24. The method of statement 19 or 23, wherein Step A) is conducted in 3D suspension culture conditions.

S25. A human milk like product which is obtainable according to the method described in anyone of statements 19 to 24.

25

S26. A human milk like product according to statement 25 for use in therapy.

S27. Use of a human milk like product according to statement 25 as a human milk substitute, optionally a breast-feeding substitute.

It should be appreciated that the various aspects and embodiments of the detailed description as disclosed herein are illustrative of the specific ways to make and use the invention and do not limit the scope of invention when taken into consideration with the claims and the detailed description. It will also be appreciated that features from aspects and embodiments of the invention may be combined with further features from the same or different aspects and embodiments of the invention.

10

As used in this detailed description and the appended claims, the singular forms “a,” “an” and “the” include plural referents unless the context clearly dictates otherwise.

15

Figures

Fig. 1: shows the differentiation of human induced pluripotent stem cells (hiPSCs) according to the protocol outlined in Ying Qu and as applied in one alternative in step A) of the inventive methods.

20

Fig. 2: shows the differentiation of human induced pluripotent stem cells (hiPSCs) according to step A).

Fig. 3: shows the differentiation of human induced pluripotent stem cells (hiPSCs) according to the preferred and particularly preferred embodiments for step A) of the inventive methods.

25

- Fig. 4:** shows that three-dimensional organotypic cultures of hiPSCs as produced according to the methods of **Fig. 2** are highly permissive for mammary glands specification. mRNA expression of Nanog, TUBB3, FOXA2, TP63, KR-14, EpCAM, KRT8 and CSN2 for 3D-differentiation (42 days) protocol are shown. Markers from left to right: Stages of Pluripotency (Nanog), Lineage (ectoderm & endoderm) (TUBB3, FOXA2), Basal-cell/myoepithelial markers (TP63, KR-14), Luminal epithelial markers (EpCAM, KRT8), and milk proteins (CSN2 (Casein Beta)).
- Fig. 5:** shows the two-dimensional organotypic culture of hiPSCs produced as a comparative example. mRNA expression of Nanog, TUBB3, FOXA2, TP63, KR-14, EpCAM, KRT8 and CSN2 for 2D-differentiation (31 days) protocols are shown. Markers from left to right: Stages of Pluripotency (Nanog), Lineage (ectoderm & endoderm) (TUBB3, FOXA2), Basal-cell/myoepithelial markers (TP63, KR-14), Luminal epithelial markers (EpCAM, KRT8), and milk proteins (CSN2 (Casein Beta)).
- Fig. 6:** shows the effect of bone morphogenetic protein 4 (BMP4) on non-neural ectoderm lineage differentiation of induced pluripotent stem cells (iPSCs) using the 3D-organoid setting. **a**, Scheme summarizing the procedure for the generation of mammary gland progenitors and BMP4 treatment between day 0 and day 3. **b**, Microscopic analysis of morphological changes after exposure to different concentration of BMP4. **c-f**, Flow cytometry quantification of the number of mammary gland positive progenitor-cells using different identified markers: EpCAM, CD49f, MUC1 and GATA3. **g-h**, RNA expression level of human TFAP2A and TFAP2C as predominant non-neural ectodermal markers. **i-k**, human PAX6, OTX2 and SOX11 as neural

ectoderm markers. **l**, human KRT18 as a luminal marker. **m**, human OPN as a secreted phosphorylated milk glycoprotein.

Fig. 7: shows the effect of BMP4 treated samples. **a**, showed a higher expression of double mammary gland markers EpCAM+/CD49f+ and double mature luminal lactocyte markers MUC1+/EpCAM+ compared to the control samples. **b**, effect of bone morphogenetic protein 4 (BMP4) on estrogen related receptor alpha (ESRRA), Keratin 14 (KRT14) and MUC1 RNA expression level between day 30 and 35 of mammary-gland 3D-differentiation.

10

Experimental section

Example 1

Cultivation and differentiation of hiPSCs into lactocytes to obtain a human milk 5 like product

Lactocytes are cultured starting from ihPSCs according to the procedure described in Ying Qu et al, Stem Cell Report vol 8, 205-215 February 14th 2017 and the human milk like product thereby secreted is collected and can be used in therapy and/or as
10 a breastfeeding substitute according to the present invention.

Example 2

Cultivation and differentiation of hiPSCs into 3D-lactocytes to obtain a human milk 15 like product

Lactocytes are cultured starting from hiPSCs according to the method of the present invention following steps A) and B) as described above) and the human milk like product thereby secreted is collected and can be used in therapy and/or as a breastfeeding substitute according to the present invention.
20

Example 3

Alternative methods of cultivation and differentiation of hiPSCs into lactocytes to 25 obtain a human milk like product

Efficient lactocytes differentiation from hiPSCs can be obtained from alternative culture conditions including conditions 1 to 4 as below described:

1. 2D culture on vitronectin coated plates as monolayer of cells derived from the EBs and cultured for at least 28 days in a medium containing (RPMI 1640 with L-glutamine; Fetal bovine serum (FBS); Insulin; Epidermal growth factor (EGF); hydrocortisone; Pen-Strep (penicillin/streptomycin : antibiotic-
30 antimycotic solution).

2. 2D culture on vitronectin coated plates of attached aggregates (EBs) of cells derived from the EBs and cultured for at least 28 days in a medium containing (RPMI 1640 with L-glutamine; Fetal bovine serum (FBS); Insulin; Epidermal growth factor (EGF); hydrocortisone; Pen-Strep (antibiotic-antimycotic solution).
3. 3D culture in suspension in MammoCult medium for at least 10 days and then culture in mixed floating gels (for example Matrigel and Collagen 1) for another 5 days in a specific medium (for example EpiCultB) in presence of Parathyroid hormone followed by 25 days in presence of insulin, HGF, hydrocortisone and FGF10;
4. 3D culture of EBs in suspension (ultra low adherent plate) in MammoCult medium for at least 10 days and then in suspension culture for another 5 days in a specific medium (for example EpiCultB) in presence of Parathyroid hormone followed by 25 days in presence of insulin, HGF, hydrocortisone and FGF10.

Example 4

2D- and 3D-lactocyte differentiation based on human-induced pluripotent stem cell (hiPSC) line 603

(a) 3D-lactocyte differentiation based on human-induced pluripotent stem cell (hiPSC) line 603:

The human-induced pluripotent stem cell (hiPSC) line 603 was used for 3D-lactocyte differentiation. The human-induced pluripotent stem cell (hiPSC) line 603 was purchased from Fujifilm Cellular Dynamics, Inc (FCDI).

(i) For the 3D differentiation protocol (according to the invention), EBs (spheroids) were formed by incubating single cells of hiPSC in E8 medium with 10uM rock inhibitor at 37°C, 5% CO₂ in rotation at 95 rpm overnight.

Second day, medium was replaced with E8 (day -2-day 0).

Next day, medium was replaced with Mammo1 medium (MammoCult - medium with proliferation supplements, heparin (4µg/mL), and hydrocortisone (0.48µg/mL) with penicillin/streptomycin) for 10 days (day 0-day 10).

5 Medium was changed every second day.

(ii) The differentiation was followed by 5 days in Mammo2 medium (EpiCultB + supplements, pThrP 100ng/ml plus penicillin/streptomycin). Culture medium was changed every 3 days (day 10-day 15).

(iii) In order to induce branching epithelial structure, alveolar
10 differentiation and mammary cell specification, mEBs (spheroids/mammospheres) were fed with Mammo3 medium (complete EpiCultB, hydrocortisone (1 µg/ml), insulin (10 µg/ml), FGF10 (50 ng/ml), HGF (50 ng/ml) and penicillin/streptomycin) for 20 days. Medium was changed every 3 days (day 15-day 35).

(iv) Finally, to induce the milk bioactive production (3D), we used
15 the Mammo4 medium (complete EpiCultB, 10% FBS, prolactin (10 µg/ml), hydrocortisone (1 µg/ml), insulin (10 µg/ml), progesterone, β-estradiol and penicillin/streptomycin for 7 days and medium was changed every 3 days (day 35-day 42). During all the differentiation procedure, spheroids were maintained in the suspension culture (rotating at 95 rpm). The differentiation procedure ended at day
20 42. Results are displayed in Figure 4.

(b) 2D-lactocyte differentiation based on human-induced pluripotent stem cell (hiPSC) line 603

The human-induced pluripotent stem cell (hiPSC) line 603 was
25 used also for 2D-lactocyte differentiation. The human-induced pluripotent stem cell (hiPSC) line 603 was purchased from Fujifilm Cellular Dynamics, Inc (FCDI).

For the 2D-differentiation protocol (used for comparison), we used the Lacto medium during all the differentiation stages (RPMI 1640, 20% FBS,

1mM glutamine, 4 µg/ml insulin, 20 ng/ml EGF, 0.5µg/ml hydrocortisone with penicillin/streptomycin). Cells were incubated at 37°C, 5% CO₂. Medium was replaced every second day. Results are displayed in Figure 5.

5 (c) Results

The different differentiation stages during lactocyte derivation were captured using quantitative RT-PCR (Figure 4, 3D-differentiation, Figure 5, 2D-differentiation). In both 2D- and 3D-settings, Nanog expression as a marker for pluripotency is decreased while cells are passing towards the maturation and differentiation. The neuroectodermal and endodermal markers, TUBB3 (Tubulin Beta 3 Class III) and Forkhead box protein A2 (FOXA2) were not expressed significantly in 3D-format and TUBB3 elevation is only captured in 2D-setting. This demonstrates that hiPSCs are patterned towards the non-neural ectodermal lineage, thus enriching mammary progenitors in 3D-format. We investigated the expression pattern of commonly used basal cell/myoepithelial markers, such as p63 (a p53-homologous nuclear protein) and cytokeratin 14 (KRT-14). Both markers are detectable significantly in both systems. Additionally, the epithelial cell adhesion molecule (EpCAM) and cytokeratin 8 (KRT8) were tracked only in the 3D-system and KRT8 was only partially expressed in the 2D-format. Consequently, 3D-platform in an organotypic setting expressed common breast tissue, luminal, and basal markers. Such mammary like organoids express human breast specific proteins including CSN2 (casein beta), milk protein peptides, and hormone receptors. The luminal cells specifically express EpCAM, MUC1, CD49F, GATA3, CK8, and CK18 while basal cells will specifically express CK14, α-smooth muscle actin and P63. Eventually EpCAM and CD49F double positive cells can be detected at an earlier progenitor stage between D10 and D35. Interestingly, CSN2 expression is only captured at the last time point (D42) of the 3D-organotypic system and not in the 2D-directed differentiation platform.

Analysis of the mammary like organoids secretome showed secretion of human milk specific bioactives including oligosaccharides (including lactose and some HMOs), lipids (including 4 fatty acids), proteins (7 detected including caseins), and miRNA (75 detected, including 11 typically detected in HBM) as below described.

Primary cell supernatant was analyzed for presence of lactose or human milk oligosaccharides following the procedure described in “Austin and Benet, Quantitative determination of non-lactose milk oligosaccharides, *Analytica Chimica Acta* 2018, 1010, 86-96” with minor modification. The samples were analysed with UHPLC and detected lactose or human milk oligosaccharides (HMOs) were quantified against a calibration curve of lactose and a mix of 7 HMOs (2’FL, 3FL, DFL, LNT, LNnT, 3’SL and 6’SL). The method had an estimated limit of 0.1mg/L. In the primary cell supernatants, Lactose (0.22 mg/l) and 6’SL (0.32 mg/l) were detected at day 42.

Fatty acids were analysed in media and cell supernatants by gas chromatography coupled with flame ionization detector. Briefly, the supernatants obtained at day 42 is analysed to investigate the presence of fatty acids contained in several lipid classes. A 7890A gas-chromatograph with a 7693 autosampler with preparative station module equipped with a fused-silica CP-Sil 88 capillary column (100% cyanopropylpolysiloxane; 100 m, 0.25 mm id, 0.25 mm film thickness is used with a split injector (1:25 ratio) heated at 250°C and a flame-ionization detector operated at 300°C. Preparation of FAMES (fatty acids methyl esters) is performed by direct transesterification of sample with methanolic chloridric acid. Separation of FAMES is performed using capillary gas chromatography-FID (GC). Identification of FAMES is done by retention time (RT) and comparison with an external standard. Quantification of fatty acids is done by calculation using methyl C11:0 as internal

standard. Transesterification performance of the method is controlled with TAG C13:0 as second internal standard. After addition of internal standards, the solution was mixed with 2 mL of methanol, 2 mL of Methanol/HCl (3N) and 1 mL of hexane. After heating at 100°C/60min, the sample is cooled down to room temperature
 5 (about 15 min) and the reaction is stopped by adding 2mL of water. After centrifugation the organic phase is directly injected into the GC.

Fatty acid results from protocol of Example 4a at time day 42 are reported in table 1 (differences observed between media and supernatant).

10 The table 1 below lists the expressed fatty acids in cell supernatant sample.

Fatty acid	Detected amount in cell supernatant (mg/100 mL)
C-4:0	2.53
C-8:0	0.49
C-10:0	0.38
C-14:0	0.44
C-15:0	0.41
C-16:0	1.85
C-16:1n7	0.08
C-17:0	0.09
C-18:0	0.97
C-18:1 n9	18.82
C-18:1	0.28
C-18:2 n6	2.16
C-20:0	0.13
C-20:1 n9	0.11

C-18:3 n3	0.08
C-22:0	0.29
Other fatty acids	1.38

- Proteins in the cell supernatant were analysed using SDS-PAGE profiling and then band isolation for identity confirmation by LC-MSMS. For SDS-PAGE analysis, the total volume of the prepared sample was loaded on the gel. A human milk sample was added for comparison as control. Selected gel regions (bands) were cut to look for human proteins by LC-MSMS. Eventually, bands were submitted to in-gel trypsin digestion and analyzed by LC-MSMS. LC-MSMS data were analyzed with Peaks Studio and matched against the UniProt database for human proteins.
- 10 The table 2 below lists the best candidates for all the excised bands.

Name of expressed proteins in the cell supernatant
Lactoferrin
Albumin
Prolactin
Alpha S1-casein
Hemoglobin subunit beta
Hemoglobin subunit alpha
α -lactalbumin
Alpha-2-macroglobulin
β -casein

bile salt-activated lipase
κ -casein
lactadherin
CD14
fatty acid synthase
IgA
pIgR
Serum albumin
Xanthine dehydrogenase

Exosome isolation and miRNA profiling was performed using ExoQuick polymer nets. ExoQuick polymer works to precipitate exosomes by forming a network and collects all exosomes of a certain size. Once the ExoQuick mesh is formed, a simple, low-speed centrifugation easily precipitates the exosomes as a pellet. The exosomes are intact, ready for protein or RNA analysis and are bioactive for functional studies. Precipitation buffer was added in a ration 0.25X to the sample then vortex. The mix was incubated overnight at 4°C. After incubation, samples were centrifuged 30 min at 1,500xg. The exosome pellet was re-suspended by vertexing in initial volume with Buffer XE (QIAGEN) for QC or Lysis Buffer from HTG EdgeSeq miRNA Whole Transcriptome Assay for miRNA profiling. In order to assess the extracellular vesicles (EVs) isolation, the supernatant was first centrifuged at 3000g for 15 min to remove cell pellet and debris. Then 100 microliters of media was used for an overnight precipitation at 4°C with ExoQuick buffer (ratio 0.25X). EV precipitates were recovered by centrifugation for 30 min at 1500g. Two precipitations were performed for each sample, one EV precipitation was resuspended in Buffer XE (QIAGEN) for potential further analysis, and a second one in only 50 ul HTG Lysis buffer in order to concentrate by 10-fold before miRNA profiling with HTG.

For miRNA profiling, samples were used directly in the first step of lysis. Thus, Whole sample was used directly and was lysed with Plasma lysis buffer in a ratio 1:1. Next, proteinase K (1/10) was added and the samples were incubated 3h at 50°C at 600rpm on Thermomixer. EVs were resuspended in Lysis buffer and lysed in the same conditions, with an incubation step at 95°C for 10min added before the lysis incubation. 26 µl of lysate was process with 70 µl of oil on the HTG processor following the HTG EdgeSeq miRNA Whole Transcriptome Assay V2 procedure. For indexing and amplification libraries, samples were tagged with Illumina adaptors and indexes by PCR with OneTaq® Hot Start 2X Master Mix GC Buffer (95°C - 4 min; 16 cycles: 95°C-15 sec, 56°C-45 sec, 68°C- 45 sec; 68°C10 min; Hold at 4°C) and AMPure cleaned (ratio 2.5) on a robotic liquid handler SciClone NGS WorkStation (Perkin Elmer). Pools were obtained with our custom pooling program on Hamilton robot. The samples were pooled based on GX touch Chip HS quantification. The pools were purified manually a second time with AMPure Bead (ratio 1.8) to remove potential remaining traces of primer-dimer and quantified with Qubit to adjust the final concentration to 2 nM. And as a last step, for MiSeq sequencing, pools were loaded on MiSeq at 20pM with a 5% PhiX spike and sequenced for 50 base Single read on MiSeq with 150V3 kit.

Briefly, 974 miRNAs detected in the in the cell supernatant which more than 75 of them are highly expressed miRNAs in the milk samples.

The table 3 below lists the top ten highly expressed miRNAs.

miRNA name	log2 counts	CV
miR-21-5p	9.76	0.01
miR-181a-5p	9.07	0.03
miR-30d-5p	8.63	0.01
miR-30b-5p	8.63	0.01
miR-22-3p	8.49	0.01
miR-146b-3p	8.40	0.01

miR-30c-5p	8.12	0.04
miR-30a-5p	7.63	0.02
miR-30e-5p	7.26	0.01
miR-148b-3p	6.77	0.04

Our findings provide a novel iPSC-based 3D-organotypic model for studying the regulation and development of normal mammary cell fate and function as well as breast milk bioactives production.

Example 5

Effect of bone morphogenetic protein 4 (BMP4) on non-neural ectoderm lineage differentiation of induced pluripotent stem cells (iPSCs) using the 3D-organoid setting

Methods: The human-induced pluripotent stem cell (hiPSC) line 603 was purchased from Fujifilm Cellular Dynamics, Inc (FCDI) and used for the 3D-differentiaion of mammary gland progenitors. Briefly, 5.5 million dissociated iPSCs were plated in the 6 well-plates (ultra-low attachment) containing 4.5 mL aggregation media using a planar shaker platform (set at 95 RPM). BMP4 (314-BP-050, Bio-Techne AG) at different concentrations was added to the culture media between day 0 and day 3.

Results: iPSC cells were tested with different concentrations of BMP4 to induce non-neural ectodermal differentiation (Figure 6a, b). 20 ng/mL of BMP4 significantly induced the expression of mammary gland progenitor markers such as EpCAM (CD326), CD49f, MUC1 (CD227) and GATA3 using flow cytometry quantification (Figure 6c-f). Expression of human TFAP2A and TFAP2C (AP2 Gamma) as predominant non-neural ectodermal markers were increased at day 10 of

differentiation using Nanostring analysis compared to the control sample (Figure 6g-h). Moreover, expression levels of human specific neural ectoderm markers, such as paired box gene 6 (PAX6), orthodenticle homeobox 2 (OTX2) and SRY-box transcription factor 11 (SOX11) were decreased during differentiation (Figure 6i-k).

- 5 Lastly, BMP4 can induce the expression of the human luminal specific markers cytokeratin 18 (KRT18) and secreted phosphorylated milk glycoprotein osteopontin (OPN) (Figure 6l-m).

- These results demonstrate that BMP4 steers iPSCs commitment towards the non-
10 neural ectodermal lineage and surprisingly significantly increases mammary gland progenitor cell specification.

Example 6

- 15 The BMP4 treated samples showed a higher expression of double mammary gland markers EpCAM+/CD49f+ and double mature luminal lactocyte markers MUC1+/EpCAM+ compared to the control samples (Figure 7a).

- Moreover, the effect of BMP4 at the end-stage of 3D-mammary gland
20 differentiation of iPSC cells was assessed using estrogen related receptor alpha (ESRRA), Keratin 14 (KRT14) and MUC1 for their expression profiles. Surprisingly, BMP4 increased the RNA expression levels of all the genes at day 30, 33 and 35 compared to the control sample (Figure 7b).

- 25 It should be understood that various changes and modifications to the presently preferred embodiments described herein will be apparent to those skilled in the art. Such changes and modifications can be made without departing from the spirit and scope of the present invention and without diminishing its attendant advantages. It

is therefore intended that such changes and modifications be covered by the appended claims.

Claims

1. A method of producing a population of mammary gland cells, comprising:
 - i) culturing mammalian induced pluripotent stem cells (miPSCs) in a culture
5 medium comprising bone morphogenic protein 4 (BMP4) to generate embryoid bodies (EBs), and
 - ii) growing the EBs to generate a population of mammary cells.
2. The method of claim 1, wherein the culturing step i) comprises culturing the
10 miPSCs in MammoCult medium and BMP4, in a 3D-suspension culture system, for example 3D-suspension condition, thereby directing the iPSCs to differentiate towards non-neural ectoderm cells, optionally for at least 12 days.
3. The method of claims 1 or 2, wherein the growing step ii) comprises growing
15 the formed EBs in a 3D embedding system, for example a mixed floating gel composed of matrix protein such as Matrigel and/or Collagen I for at least 30 days, for example for 32 days, to generate lactocytes.
4. The method of any one of claims 1 to 3, wherein the mammary gland cells are
20 human mammary gland cells.
5. The method of any one of claims 1 to 4, wherein BMP4 is added to the culture medium between day 0 and day 10, preferably between day 0 and day 3, where day 0 is the time point where the iPSCs are first added to the culture medium.
25
6. The method of any one of claims 1 to 5, wherein BMP4 is added to the culture medium for 3 days.

7. The method of any one of claims 1 to 6, wherein BMP4 is added to the culture medium in a concentration of 5 to 20 ng/mL, preferably 5 ng/mL, 10 ng/mL or 20 ng/mL.
- 5 8. The method of any one of claims 1 to 7, wherein the EBs express one or more mammary gland positive progenitor-cell markers, optionally selected from EpCAM, CD49f, MUC1 and GATA3.
9. The method of any one of claims 1 to 8, wherein the EBs have increased
10 expression of one or more mammary gland positive progenitor-cell markers compared to the expression level of said mammary gland positive progenitor-cell markers in EBs not treated with BMP4.
10. The method of any one of claims 1 to 9, wherein at least 10%, 20%, 30%, 40%,
15 50%, 60%, 70%, 80% or more of the EBs express one or more mammary gland positive progenitor-cell markers, optionally wherein at least 50% of EBs express EpCAM and CD49f mammary gland positive progenitor-cell markers, optionally wherein at least 15% of EBs express MUC1 and CD49f mammary gland positive progenitor-cell markers, optionally wherein at least 20% of EBs express MUC1 and
20 EpCAM mammary gland positive progenitor-cell markers, and/or optionally wherein at least 15% of EBs express GATA3 mammary gland positive progenitor-cell markers.
11. The method of any one of claims 1 to 10, wherein the EBs express one or more non-neuronal ectodermal markers, optionally selected from TFAP2A and TFAP2C.
- 25 12. The method of any one of claims 1 to 11, wherein the EBs have increased expression of one or more non-neuronal ectodermal markers of at least 2-fold

compared to the expression level of said non-neuronal ectodermal markers in EBs not treated with BMP4, optionally at least a 3 to 15-fold increase.

13. The method of any one of claims 1 to 12, wherein the EBs have decreased
5 expression of one or more neuronal ectodermal markers of at least 0.5-fold compared to the expression level of said neuronal ectodermal markers in EBs not treated with BMP4, optionally wherein said neuronal ectodermal markers are selected from PAX6, OXT2 and SOX11.
- 10 14. The method of any one of claims 1 to 13, wherein the EBs express one or more milk-specific bioactive markers, optionally osteopontin (OPN).
- 15 15. The method of any one of claims 1 to 14, wherein the EBs have increased expression of OPN of at least 2-fold compared to the expression level of OPN in EBs not treated with BMP4, optionally at least a 4 to 18-fold increase.
16. The method of any one of claims 1 to 15, wherein the mammary gland cells form lactocyte mammary-like gland organoids.
- 20 17. The method of any one of claims 1 to 16, wherein the mammary gland cells generate increased expression of milk-specific bioactive markers compared to mammary gland cells that are not treated with BMP4, optionally wherein the mammary gland cells generate at least a 2-fold increase in expression of milk-specific bioactive markers compared to mammary gland cells that are not treated with BMP4,
25 optionally wherein said markers are selected from estrogen related receptor alpha (ESRRA), Keratin 14 (KRT14) and MUC1.

18. Use of BMP4 for increasing the differentiation efficiency of mammalian induced pluripotent stem cell (miPSCs) into mammary-gland progenitor cells in a differentiation protocol.
- 5 19. A method for producing a mammalian milk like product, comprising:
- C) Generating lactocyte mammary-like gland organoids derived from mammalian induced pluripotent stem cells (miPSC);
 - D) Secreting the mammalian milk like product from said lactocytes,
- 10 wherein step A) comprises culturing the miPSCs in a culture medium comprising BMP4.
20. The method of claim 19, wherein the method is for producing a human milk like product,
- wherein step A) further comprises:
- 15 i) directing hiPSCs to differentiate towards non-neural ectoderm cells by culturing them in an appropriate culture medium comprising BMP4, for example MammoCult medium and BMP4, in an appropriate 3D culture system, for example 3D-suspension condition, for at least 12 days and
- 20 ii) growing the formed mEBs (mammospheres) in an appropriate 3D embedding system, for example a mixed floating gel composed of matrix protein such as Matrigel and/or Collagen I for at least 30 days, for example for 32 days, to generate lactocytes.
21. The method according to claim 20 wherein step A)i) is defined as follows:
- i) generation of embryoid bodies (EBs) from hiPSCs by incubation in standard iPSC
- 25 medium E8 comprising DMEM/F12, L-ascorbic acid-2-phosphate magnesium, sodium selenium, FGF2, insulin, NaHCO₃ and transferrin, TGFβ1 or NODAL or in medium mTeSR™ for two days, and producing mEBs (mammospheres) highly enriched in non-neural ectodermal cells by incubation of EBs in complete MammoCult medium

comprising the basal medium, proliferation supplement and supplemented with BMP4, heparin, and hydrocortisone for 10 days, and wherein

step A)ii) is distinguished into further substeps and comprises the following steps: ii), iii) and iv):

- 5 ii) incubation of mEBs (mammospheres) in complete EpiCultB medium supplemented with EpiCult proliferation supplement and Parathyroid hormone (pThrP) for 5 days,
- iii) promotion of branch and alveolar differentiation and mammary cell specification by incubating mEBs (mammospheres) in EpiCultB medium supplemented with EpiCult proliferation supplement, hydrocortisone, insulin, FGF10 and HGF for 20 days, and
- 10 iv) induction of milk protein expression by incubating mEBs (mammospheres) in EpiCultB medium supplemented with EpiCult proliferation supplement, hydrocortisone, insulin, FBS, prolactin, progesterone and β -estradiol for 7 days.

22. The method according to claim 20 wherein step A)i) is defined as follows:

- 15 i) generation of embryoid bodies (EBs) from hiPSCs by incubation in standard iPSC medium E8 comprising DMEM/F12, L-ascorbic acid-2-phosphate magnesium, sodium selenium, FGF2, insulin, NaHCO_3 and transferrin, TGF β 1 or NODAL for two days, and producing mEBs (mammospheres) highly enriched in non-neural ectodermal cells by incubation of EBs in MammoCultB medium supplemented with MammoCult proliferation supplement, hydrocortisone, heparin and BMP4 for 10 days, and wherein
- 20 step A)ii) is distinguished into further substeps and comprises the following steps ii), iii) and iv):
- ii) embedding the formed mEBs (mammospheres) in a mixture of Matrigel and Collagen I floated in EpiCultB medium supplemented with EpiCult proliferation supplement and
- 25 Parathyroid hormone (pThrP) for 5 days,
- iii) promotion of branch and alveolar differentiation and mammary cell specification by incubating embedded mEBs (mammospheres) in EpiCultB medium supplemented

with EpiCult proliferation supplement, hydrocortisone, insulin, FGF10 and HGF for 20 days, and

- iv) induction of milk protein expression by incubating mEBs (mammospheres) in EpiCultB medium supplemented with EpiCult proliferation supplement,
5 hydrocortisone, insulin, FBS, prolactin, progesterone and β -estradiol for 7 days.

23. The method of any one of claims 19 to 22, wherein Step A) comprises a method of any one of claims 1 to 17.

- 10 24. The method of claim 19 or 23, wherein Step A) is conducted in 3D suspension culture conditions.

25. A human milk like product which is obtainable according to the method described in anyone of claims 19 to 24.

15

26. A human milk like product according to claim 25 for use in therapy.

27. Use of a human milk like product according to claim 25 as a human milk substitute, optionally a breast-feeding substitute.

20