

INHIBITORS OF RNA HELICASE DHX9 AND USES THEREOF

RELATED APPLICATIONS

This application claims priority to U.S. Provisional Application No. 63/309,917, filed
5 on February 14, 2022. The entire contents of the foregoing application are expressly
incorporated herein by reference.

FIELD OF THE INVENTION

The present disclosure relates to inhibitors of RNA helicase DHX9, and
10 pharmaceutically acceptable salts thereof, compositions of these compounds, processes for
their preparation, and their use in the treatment of diseases.

BACKGROUND OF THE INVENTION

15 DHX9, also known as RNA Helicase A (RHA) or Nuclear DNA Helicase II (NDH
II), is a DExH-box RNA helicase, which shuttles between nucleus and cytoplasm, and can
use all four NTPs to power cycles of directional movement from 3' to 5'. Functionally,
DHX9 can bind to and unwind or resolve dsDNA/RNA, ssDNA/RNA, DNA:RNA hybrids
(such as R-loops), circular RNA, and DNA/RNA G quadruplexes. As such, DHX9 has
20 regulatory roles in various RNA and DNA related cellular processes, such as transcription,
translation, RNA splicing, editing, RNA transport and processing, microRNA genesis, and
maintenance of genomic stability (Pan et al., 2021, *Current Protein & Peptide Science* (22),
29-40).


Due to its regulatory role in processes such as transcription and maintenance of
25 genomic stability, DHX9 has shown to be a key regulator in a variety of cancer types
(Gulliver et al., 2020, *Future Science OA* (2), FSO650). DHX9 has been shown to be
involved in the regulation of genes associated with sustained proliferative signaling, evasion
of growth suppressors, evasion of apoptosis, angiogenesis, and metastasis, all of which are
hallmarks of cancer. Specifically, microsatellite instable cancers, such as Microsatellite
30 Instable (MSI) colorectal cancer, and tumors with defective MisMatch Repair (MMR) exhibit
a strong dependence on DHX9.

In addition to its role in cancer, DHX9 has been implicated in other diseases involving
gene replication, translation or regulation. These diseases include viral infections and
autoimmune disease.

Thus, there is a need for DHX9 inhibitors as potential therapeutic agents for treating diseases or disorders that are responsive to DHX9 inhibition.

SUMMARY OF THE INVENTION

- 5 The present disclosure provides compounds that are DHX9 inhibitors. In a first aspect, the present disclosure relates to compounds having the Formula I:



or a pharmaceutically acceptable salt thereof, wherein:

X is halo;

- 10 Y is O or NR^y;

R^y is H or C₁₋₄alkyl;

R¹ is C₁₋₄alkyl or C₃₋₆cycloalkyl; wherein the C₁₋₄alkyl or C₃₋₆cycloalkyl are each optionally and independently substituted with 1 to 3 halo or -OH,

- 15 Ring A is a C₃₋₆cycloalkyl, 4 to 6-membered monocyclic heterocyclyl, 6 to 10 membered bicyclic heterocyclyl; phenyl, 5 to 6-membered monocyclic heteroaryl, or 8 to 10-membered bicyclic heteroaryl;

m is 0 to 3;

n is 0 or 2;

- 20 each R² is independently selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, halo, OR^{2a}, cyano, -NR^{2b}R^{2c}, -SO₂R^{2a}, -C(O)R^{2d}, and -C(O)NR^{2b}R^{2c} wherein the C₁₋₄alkyl and C₃₋₆cycloalkyl are optionally substituted with 1 to 4 R^{2e}; or 2 R² together form oxo;

R^{2a} is H, C₁₋₄alkyl, C₁₋₄haloalkyl or C₁₋₄alkoxyC₁₋₄alkyl;

- 25 R^{2d} is H, C₁₋₄alkyl, C₃₋₆cycloalkyl, OR^{2a}, phenyl, 4 to 6-membered monocyclic heterocyclyl, 6 to 8-membered bicyclic heterocyclyl, or 5 to 6-membered monocyclic heteroaryl, wherein the C₁₋₄alkyl is optionally substituted with 1 to 3 halo and the 5- to 6-membered monocyclic heteroaryl is optionally substituted with 1 to 3 C₁₋₄alkyl, C₁₋₄haloalkyl, or phenyl;

R^{2b} and R^{2c} are each independently selected from H, C₁₋₄alkyl, C₃₋₆cycloalkyl, phenyl, 4 to 6-membered monocyclic heterocyclyl and 5 to 6-membered monocyclic heteroaryl,

wherein the C₁₋₄alkyl is optionally substituted with 1 to 3 substituents independently selected from halo and C₁₋₃alkoxy; or R^{2b} and R^{2c}, together with the nitrogen to which they are attached form a 4 to 6 membered monocyclic heterocyclyl or 6 to 10 membered bicyclic heterocyclyl; wherein the 4 to 6 membered monocyclic heterocyclyl or 6 to 10 membered bicyclic heterocyclyl are each optionally substituted with 1 to 3 R^{2d};

each R^{2e} is independently selected from halo, cyano, NR^{2b}R^{2c}, OR^{2a}, phenyl, and 4 to 6 membered monocyclic heterocyclyl;

Z is a bond, -CH₂-, -O-, -O-C₁₋₄alkylene-*, -C₁₋₄alkylene-O-*, -C(O)-, -C(O)O-*, -OC(O)-*, -S(O)₂-, -S(O)₂N(Z^a)-*, -N(Z^a)S(O)₂-, -N(Z^a)-, -N(Z^a)-C₁₋₄alkylene-*, -C₁₋₄alkylene-N(Z^a)-*, -C(O)N(Z^a)-*, -N(Z^a)C(O)-*, or -C(O)N(Z^a)-C₁₋₃alkylene-*, wherein * indicates the attachment point to R³;

Z^a is H or C₁₋₄alkyl;

R³ is C₃₋₆cycloalkyl, 7 to 10-membered bicyclic carbocyclyl, 4 to 6-membered monocyclic heterocyclyl, 6 to 10-membered bicyclic heterocyclyl, phenyl, 5 to 6-membered monocyclic heteroaryl, or 8 to 10-membered bicyclic heteroaryl, wherein the C₃₋₆cycloalkyl, 7 to 10-membered bicyclic carbocyclyl, 4 to 6-membered monocyclic heterocyclyl, 6 to 10-membered bicyclic heterocyclyl, phenyl, 5 to 6-membered monocyclic heteroaryl, or 8 to 10-membered bicyclic heteroaryl are each optionally and independently substituted with 1 to 4 R⁴;

each R⁴ is independently selected from C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, halo, OR^{4a}, cyano, -NR^{4b}R^{4c}, -C(O)R^{4a}, -C(O)OR^{4a}, -C(O)NR^{4b}R^{4c}, -NR^{4b}C(O)R^{4a}, -NR^{4b}C(O)OR^{4a}, -NR^{4b}SO₂R^{4a}, -SR^{4a}, -S(O)R^{4a}, -SO₂R^{4a}, SO₂NR^{4b}R^{4c}, -P(O)R^{4b}R^{4c}, phenyl, 5 to 6-membered monocyclic heteroaryl, 4 to 6-membered monocyclic heterocyclyl, and 6 to 10-membered bicyclic heterocyclyl, wherein the C₁₋₄alkyl, C₂₋₄alkenyl, or C₂₋₄alkynyl is optionally substituted with 1 to 4 R^{4d}, and wherein the phenyl, 5 to 6-membered monocyclic heteroaryl, 4 to 6-membered monocyclic heterocyclyl and 6 to 10-membered bicyclic heterocyclyl are each optionally substituted with 1 to 3 R^{4e} and further optionally substituted with 1 or 2 oxo; or two R⁴ together form oxo;

R^{4a} is H, C₁₋₄alkyl optionally substituted with 1 to 4 R^{4d}, -NR^{4b}R^{4c}, C₃₋₆cycloalkyl, 4 to 6-membered monocyclic heterocyclyl, phenyl, or 5 to 6-membered monocyclic heteroaryl; wherein the C₃₋₆cycloalkyl, 4 to 6-membered monocyclic heterocyclyl, phenyl or 5 to 6-membered monocyclic heteroaryl are each optionally and independently substituted with 1 to 3 R^{4e};

R^{4b} and R^{4c} are each independently selected from H, phenyl, 4 to 6-membered monocyclic heterocyclyl, 5 to 6-membered monocyclic heteroaryl, and C_{1-4} alkyl optionally substituted with 1 to 4 R^{4d} ; or R^{4b} and R^{4c} together with the nitrogen atom to which they are attached to form a 4 to 6-membered monocyclic heterocyclyl;

5 each R^{4d} is independently selected from halo, OR^{4f} , $-C(O)C_{1-4}$ alkyl, $-C(O)NR^{4b}R^{4c}$, $-C(O)C_{1-4}$ haloalkyl, $-C(O)OR^{4f}$, $-NR^{4b}R^{4c}$, phenyl, and 5 to 6-membered monocyclic heteroaryl, wherein the phenyl and 5 to 6-membered monocyclic heteroaryl are each optionally substituted with 1 to 3 R^{4g} ;

each R^{4e} is independently selected from halo, C_{1-4} alkyl, cyano, OR^{4f} , $-NR^{4b}R^{4c}$, $-C(O)H$, $-C(O)R^{4h}$, $-SO_2C_{1-3}$ alkyl, and $-C(O)NR^{4b}R^{4c}$, wherein C_{1-4} alkyl is optionally substituted with 1 to 3 substitutes independently selected from halo, $-SO_2C_{1-3}$ alkyl and $-C(O)NR^{4b}R^{4c}$; and $-C(O)NR^{4b}R^{4c}$; or two R^{4e} together form oxo;

10

R^{4f} is H, C_{1-4} alkyl, C_{1-4} haloalkyl, phenyl or 5 to 6-membered monocyclic heteroaryl, wherein the phenyl and 5 to 6-membered monocyclic heteroaryl are each optionally substituted with one to three halo;

15

each R^{4g} is independently selected from halo, OR^{4f} , C_{1-4} alkyl, C_{1-4} haloalkyl, halo, cyano, $-NR^{4b}R^{4c}$, $-C(O)H$, and $-C(O)OR^{4f}$; and

each R^{4h} is independently C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-3} alkoxy or $-N(R^{4b})_2$.

Another aspect of the disclosure relates to pharmaceutical compositions comprising compounds of Formula (I) or pharmaceutically acceptable salts thereof, and a pharmaceutical carrier.

20

In yet another aspect, the present disclosure provides a method of treating a disease or disorder that is responsive to inhibition of DHX9 in a subject comprising administering to said subject an effective amount of at least one compound described herein or a pharmaceutically acceptable salt thereof. In some embodiments, the method is for the treatment of cancer.

25

Another aspect of the present disclosure relates to the use of at least one compound described herein or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of a disease or disorder responsive to inhibition of DHX9. Also provided is a compound described herein or a pharmaceutically acceptable salt thereof for use in treating a disease or disorder responsive to inhibition of DHX9.

30

BREIF DESCRIPTION OF THE DRAWINGS

FIG. 1a and FIG. 1b show tumor volume over time for an *in vivo* efficacy study of compound of Example 31 in LS411N Xenografts.

FIG. 1c shows percentage of tumor volume change over time for the *in vivo* efficacy study of compound of Example 31 in LS411N Xenografts.

FIG. 2 shows percentage of body weight change over time for the *in vivo* efficacy studies of compound of Example 31 in LS411N Xenografts.

FIG 3a. shows tumor volume over time during and after dosing period for the *in vivo* efficacy study of compound of Example 31 in LS411N Xenografts.

FIG. 3b shows percentage of tumor volume change over time during and after 28 days dosing period for the *in vivo* efficacy study of compound of Example 31 in LS411N Xenografts.

FIG. 4 shows anti-proliferation activity of 290 DHX9 inhibitor compounds described herein in CRC-MSI (LS411N) cells versus CRC-MSS (NCI-H747) cells.

FIG . 5a shows 10-day proliferation screen for compound of Example 18 in various cancer cell line panels.


FIG. 5b shows proliferation EC₅₀ for compound of Example 18 in MSI cell lines vs. MSS cell lines as represented in colorectal, endometrial, and hematopoietic cancers.

DETAILED DESCRIPTION OF THE INVENTION

The present disclosure provides compounds and pharmaceutical compositions thereof that may be useful in the treatment of diseases or disorders through mediation of DHX9 function/activity. In some embodiments, the compounds of present disclosure are DHX9 inhibitors.

COMPOUNDS AND COMPOSITIONS

In a first embodiment, the present disclosure provides a compound of Formula (I):



or a pharmaceutically acceptable salt thereof, wherein the variables in Formula (I) are as defined as follows:

X is halo;

Y is O or NR^y;

5 R^y is H or C₁₋₄alkyl;

R¹ is C₁₋₄alkyl or C₃₋₆cycloalkyl; wherein the C₁₋₄alkyl or C₃₋₆cycloalkyl are each optionally and independently substituted with 1 to 3 halo;

Ring A is a C₃₋₆cycloalkyl, 4 to 6-membered monocyclic heterocyclyl, 6 to 10 membered bicyclic heterocyclyl; phenyl, 5 to 6-membered monocyclic heteroaryl, or 8 to 10-
10 membered bicyclic heteroaryl;

m is 0 to 3;

n is 0 or 1;

each R² is independently selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, halo, OR^{2a}, cyano, -NR^{2b}R^{2c}, -SO₂R^{2a}, -C(O)R^{2d}, and -C(O)NR^{2b}R^{2c}, wherein the C₁₋₄alkyl and C₃₋₆cycloalkyl are
15 optionally substituted with 1 to 4 R^{2e}; or 2 R² together form oxo;

R^{2a} is H, C₁₋₄alkyl, C₁₋₄haloalkyl or C₁₋₄alkoxyC₁₋₄alkyl;

R^{2d} is H, C₁₋₄alkyl, C₃₋₆cycloalkyl, OR^{2a}, phenyl, 4 to 6-membered monocyclic heterocyclyl or 5 to 6-membered monocyclic heteroaryl, wherein the C₁₋₄alkyl is optionally substituted with 1 to 3 halo;

20 R^{2b} and R^{2c} are each independently selected from H, C₁₋₄alkyl, C₃₋₆cycloalkyl, phenyl, 4 to 6-membered monocyclic heterocyclyl and 5 to 6-membered monocyclic heteroaryl; or R^{2b} and R^{2c}, together with the nitrogen to which they are attached form a 4 to 6 membered monocyclic heterocyclyl or 6 to 10 membered bicyclic heterocyclyl; wherein the 4 to 6 membered monocyclic heterocyclyl or 6 to 10 membered bicyclic heterocyclyl are each
25 optionally substituted with 1 to 3 R^{2d};

each R^{2e} is independently selected from halo, cyano, NR^{2b}R^{2c}, OR^{2a}, phenyl, and 4 to 6 membered monocyclic heterocyclyl;

Z is a bond, -CH₂-, -O-, -O-C₁₋₄alkylene-*, -C₁₋₄alkylene-O-*, -C(O)-, -C(O)O-*, -OC(O)-*, -S(O)₂-, -S(O)₂N(Z^a)-*, -N(Z^a)S(O)₂-, -N(Z^a)-, -N(Z^a)-C₁₋₄alkylene-*, -C₁₋₄alkylene-N(Z^a)-*, -C(O)N(Z^a)-*, or -C(O)N(Z^a)-C₁₋₃alkylene-*, wherein * indicates the
30 attachment point to R³;

Z^a is H or C₁₋₄alkyl;

R^3 is C_{3-6} cycloalkyl, 7 to 10-membered bicyclic carbocyclyl, 4 to 6-membered monocyclic heterocyclyl, 6 to 10-membered bicyclic heterocyclyl, phenyl, 5 to 6-membered monocyclic heteroaryl, or 8 to 10-membered bicyclic heteroaryl, wherein the C_{3-6} cycloalkyl, 7 to 10-membered bicyclic carbocyclyl, 4 to 6-membered monocyclic heterocyclyl, 6 to 10-membered bicyclic heterocyclyl, phenyl, 5 to 6-membered monocyclic heteroaryl, or 8 to 10-membered bicyclic heteroaryl are each optionally and independently substituted with 1 to 4 R^4 ;

each R^4 is independently selected from C_{1-4} alkyl, halo, OR^{4a} , cyano, $-NR^{4b}R^{4c}$, $-C(O)R^{4a}$, $-C(O)OR^{4a}$, $-C(O)NR^{4b}R^{4c}$, $-NR^{4b}C(O)R^{4a}$, $-NR^{4b}C(O)OR^{4a}$, $-SR^{4a}$, $-SO_2R^{4a}$, $-SO_2NR^{4b}R^{4c}$, $-P(O)R^{4b}R^{4c}$, phenyl, 5 to 6-membered monocyclic heteroaryl, 4 to 6-membered monocyclic heterocyclyl, and 6 to 10-membered bicyclic heterocyclyl, wherein the C_{1-4} alkyl is optionally substituted with 1 to 4 R^{4d} , and wherein the phenyl, 5 to 6-membered monocyclic heteroaryl, 4 to 6-membered monocyclic heterocyclyl and 6 to 10-membered bicyclic heterocyclyl are each optionally substituted with 1 to 3 R^{4e} ; or two R^4 together form oxo;

R^{4a} is H, C_{1-4} alkyl optionally substituted with 1 to 4 R^{4d} , C_{3-6} cycloalkyl, 4 to 6-membered monocyclic heterocyclyl, phenyl, or 5 to 6-membered monocyclic heteroaryl; wherein the C_{3-6} cycloalkyl, 4 to 6-membered monocyclic heterocyclyl, phenyl or 5 to 6-membered monocyclic heteroaryl are each optionally and independently substituted with 1 to 3 R^{4e} ;

R^{4b} and R^{4c} are each independently selected from H, phenyl, 4 to 6-membered monocyclic heterocyclyl, 5 to 6-membered monocyclic heteroaryl, and C_{1-4} alkyl optionally substituted with 1 to 4 R^{4d} ;

each R^{4d} is independently selected from halo, OR^{4f} , $-C(O)C_{1-4}$ alkyl, $-C(O)NR^{4b}R^{4c}$, $-C(O)C_{1-4}$ haloalkyl, phenyl, and 5 to 6-membered monocyclic heteroaryl, wherein the phenyl and 5 to 6-membered monocyclic heteroaryl are each optionally substituted with 1 to 3 R^{4g} ;

each R^{4e} is independently selected from halo, C_{1-4} alkyl optionally substituted with 1 to 3 halo, cyano, OR^{4f} , $-NR^{4b}R^{4c}$, $-C(O)H$, $-C(O)C_{1-4}$ alkyl, $-C_{1-4}$ alkylene- $C(O)NR^{4b}R^{4c}$, and $-C(O)NR^{4b}R^{4c}$; or two R^{4e} together form oxo;

R^{4f} is H, C_{1-4} alkyl, C_{1-4} haloalkyl, phenyl or 5 to 6-membered monocyclic heteroaryl, wherein the phenyl and 5 to 6-membered monocyclic heteroaryl are each optionally substituted with one to three halo;

each R^{4g} is independently selected from halo, OR^{4f} , C_{1-4} alkyl, halo, cyano, $-NR^{4b}R^{4c}$, $-C(O)H$, and $-C(O)OR^{4f}$. In some embodiments, Y is O.


In a second embodiment, for the compound of Formula (I) described in the first aspect or the first embodiment, or a pharmaceutically acceptable salt thereof, X is Cl, Br, or F; and the remaining variables are as described in the first aspect or the first embodiment.


In a third embodiment, for the compound of Formula (I) described in the first aspect
5 or the first embodiment, or a pharmaceutically acceptable salt thereof, X is Cl or Br; and the remaining variables are as described in the first aspect or the first embodiment.


In a fourth embodiment, for the compound of Formula (I) described in the first aspect or the first embodiment, or a pharmaceutically acceptable salt thereof, X is Cl; and the remaining variables are as described in the first aspect or the first embodiment.


10 In a fifth embodiment, for the compound of Formula (I) described in the first aspect or the first embodiment, or a pharmaceutically acceptable salt thereof, ring A is selected from phenyl, thiophenyl, pyrrolyl, pyrazoyl, furanyl, isothiazoyl, thiazoyl, imidazoyl, cyclobutyl, benzofuranyl, 2-oxo-2,3-dihydro-1H-benzo[d]imidazolyl, imidazo[1,2-a]pyridin-6-yl, 1,4,5,6-tetrahydrocyclopenta[c]pyrazolyl, 2-oxo-2,3-dihydro-1H-imidazolyl, indolizinyl,
15 pyrrolo[1,2-a]pyrimidinyl, pyrrolo[1,2-c]pyrimidinyl, pyrrolo[1,2-a]pyrazinyl, 5,6-dihydro-4H-cyclopenta[b]thiophenyl, 5-oxo-5H-thiazolo[3,2-a]pyridinyl, thieno[3,2-b]pyridinyl, thieno[3,2-c]pyridinyl, thieno[2,3-c]pyridinyl, benzothiophenyl, thieno[3,2-d]pyrimidinyl, pyrrolo[1,2-b]pyridazinyl, pyrrolo[1,2-a]pyrazinyl, pyrrolo[1,2-a]pyrimidinyl, pyrrolo[1,2-b]pyrimidinyl, pyrrolo[1,2-c]pyrimidinyl, 1-oxo-1,2-dihydropyrrolo[1,2-a]pyrazinyl,
20 pyrrolo[2,1-f][1,2,4]triazinyl, 4,5,6,7-tetrahydrothieno[2,3-c]pyridinyl, 5,6-dihydro-4H-cyclopenta[b]thiophenyl, and 4,7-dihydro-5H-thieno[2,3-c]pyranyl; and the remaining variables are as described in the first aspect or in the first, second, third, or fourth embodiment. In an alternative fifth embodiment, ring A is phenyl, thiophenyl, pyrrolyl, pyrazoyl, furanyl, isothiazoyl, thiazoyl, imidazoyl, cyclobutyl, benzofuranyl, 2-oxo-2,3-dihydro-1H-benzo[d]imidazolyl, imidazo[1,2-a]pyridin-6-yl, 1,4,5,6-
25 tetrahydrocyclopenta[c]pyrazolyl, 2-oxo-2,3-dihydro-1H-imidazolyl, indolizinyl, pyrrolo[1,2-a]pyrimidinyl, pyrrolo[1,2-c]pyrimidinyl, pyrrolo[1,2-a]pyrazinyl, 5-oxo-5H-thiazolo[3,2-a]pyridinyl, thieno[3,2-b]pyridinyl, thieno[3,2-c]pyridinyl, thieno[2,3-c]pyridinyl, benzothiophenyl, thieno[3,2-d]pyrimidinyl, pyrrolo[1,2-b]pyridazinyl, pyrrolo[1,2-a]pyrazinyl, pyrrolo[1,2-a]pyrimidinyl, pyrrolo[1,2-b]pyrimidinyl, pyrrolo[1,2-c]pyrimidinyl,
30 1-oxo-1,2-dihydropyrrolo[1,2-a]pyrazinyl, pyrrolo[2,1-f][1,2,4]triazinyl, 4,5,6,7-tetrahydrothieno[2,3-c]pyridinyl, 5,6-dihydro-4H-cyclopenta[b]thiophenyl, and 4,7-dihydro-5H-thieno[2,3-c]pyranyl; and the remaining variables are as described in the first aspect or in the first, second, third, or fourth embodiment.



In a sixth embodiment, for the compound of Formula (I) described in the first aspect or the first embodiment, or a pharmaceutically acceptable salt thereof, ring A is represented by

the following formula:  ;


 ;


5  ;


 ;


 ; or  ; each of which is substituted with 0 to 2 R^2 and 0 to 1 $Z-R^3$; and the



remaining variables are as described in the first aspect or the first, second, third, or fourth

embodiment. In an alternative sixth embodiment, ring A is  ;

10  ;

 ;

 ;

 ; or  ; each of which is substituted with 0 to 2 R^2 and

0 to 1 $Z-R^3$; and the remaining variables are as described in the first aspect or the first,
15 second, third, or fourth embodiment.


In a seventh embodiment, for the compound of Formula (I) described in the first aspect or in the first embodiment, or a pharmaceutically acceptable salt thereof, ring A is

represented by the following formula:

The formula shows four variations of a benzene ring substituted with R³, Z, and R² groups, separated by semicolons. The first three variations show a benzene ring with R³ and Z groups at the 1 and 2 positions, and a wavy line at the 4 position. The fourth variation shows a benzene ring with R³ and Z groups at the 1 and 2 positions, and an R² group at the 4 position.


5


10



remaining variables are as described in the first aspect or in the first, second, third, or fourth

5 embodiment. In an alternative seventh embodiment, ring A is







In an eighth embodiment, for the compound of Formula (I) described in the first



In an eighth embodiment, for the compound of Formula (I) described in the first aspect or the first embodiment, or a pharmaceutically acceptable salt thereof, ring A is phenyl, thiophenyl, pyrrolyl, pyrazolyl, furanyl, isothiazoyl, and imidazolyl; and the remaining variables are as described in the first aspect or in the first, second, third, or fourth embodiment.

In a ninth embodiment, for the compound of Formula (I) described in the first aspect



In a tenth embodiment, for the compound of Formula (I) described in the first aspect or


in the first embodiment, or a pharmaceutically acceptable salt thereof, ring A is  ;



5 first aspect or in the first, second, third, or fourth embodiment.


In an eleventh embodiment, for the compound of Formula (I) described in first aspect or the first embodiment, R² is selected from -OCH₃, cyclopropyl, -CH₃, -CF₃, -CH₂CH₃, -CH(CH₃)₂, -CH(OH)CH₃, -C(CH₃)₂OH, -CH₂OCH₃, -CH₂OH, -CH₂CF₃, -CH₂CH₂OH, -CH₂CH₂CH₂OH, -CH₂CN, -CH₂CHF₂, -CH₂NH₂, -N(CH₃)₂, -SO₂CH₃, cyano, halo, -


10 C(O)OH, -C(O)N(CH₃)₂, and -C(O)NHCH₃, or the following structural formula:



are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, or tenth embodiment or any alternative embodiments described therein . In

15 alternative eleventh embodiment, R² is -CH₃, -CF₃, -CH₂CH₃, -CH(CH₃)₂, -CH(OH)CH₃, -CH₂OCH₃, -CH₂OH, -CH₂CF₃, -CH₂CH₂OH, -CH₂CH₂CH₂OH, -CH₂CN, -CH₂CHF₂, -

CH₂NH₂, -N(CH₃)₂, cyano, halo, -C(O)OH, -C(O)N(CH₃)₂, -C(O)NHCH₃,  ,



; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, or tenth embodiment or any alternative embodiments described therein.

In a twelfth embodiment, for the compound of Formula (I) described in the first aspect or in the first embodiment, or a pharmaceutically acceptable salt thereof, n is 1; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, eighth, or eleventh embodiment or any alternative embodiments described therein.

In a thirteenth embodiment, for the compound of Formula (I) described in the first aspect or in the first embodiment, or a pharmaceutically acceptable salt thereof, n is 1 and Z is a bond; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, eleventh, or twelfth embodiment.

In a fourteenth embodiment, for the compound of Formula (I) described in the first aspect or in the first embodiment, or a pharmaceutically acceptable salt thereof, n is 1 and Z is $-\text{CH}_2-$, $-\text{CH}(\text{CH}_3)-$, $-\text{O}-$, $-\text{N}(\text{H})-$, $-\text{N}(\text{C}_{1-4}\text{alkyl})-$, or $-\text{C}(\text{O})\text{NH}-*$, wherein $*$ indicates the attachment point to R^3 ; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, eleventh or twelfth embodiment.

In a fifteenth embodiment, for the compound of Formula (I) described in the first aspect or in the first embodiment, or a pharmaceutically acceptable salt thereof, R^3 is selected from cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, piperidinyl, piperazinyl, morpholinyl, pyrrolidinyl, phenyl, pyrrolyl, pyrazolyl, pyridyl, pyrimidyl, pyrazinyl, pyrazinyl, thiophenyl, tetrahydropyranyl, tetrahydrofuranyl, oxabicyclo[3.2.1]octanyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, oxadiazolyl, 3-oxo-2,3-dihydro-1H-pyrazolyl, benzamidazolyl, indazolyl, indoyl, imidazolidinyl, azetidiny, 2,3-dihydrobenzofuranyl, imidazo[1,2-a]pyridinyl, 2,3-dihydro-1H-indenyl, 2-oxo-2,3-dihydro-1H-benzo[d]imidazolyl, spiro[3.3]heptanyl, 2-azaspiro[3.3]heptanyl, and 2-oxaspiro[3.3]heptanyl, each of which is optionally substituted with 1 to 3 R^4 ; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, or fourteenth embodiment or any alternative embodiments described therein. In an alternative fifteenth embodiment, R^3 is cyclopropyl, cyclobutyl, cyclohexyl, cyclopentyl, piperidinyl, piperazinyl, morpholinyl, pyrrolidinyl, phenyl, pyrrolyl, pyrazolyl, pyridyl, pyrimidyl,


pyrazinyl, pyrazinyl, thiophenyl, tetrahydropyranyl, tetrahydrofuranyl, oxabicyclo[3.2.1]octanyl, thiazolyl, triazolyl, tetrazolyl, oxadiazolyl, 3-oxo-2,3-dihydro-1H-pyrazolyl, benzamidazolyl, indazolyl, indolyl, 2,3-dihydrobenzofuranyl, imidazo[1,2-a]pyridinyl, 2,3-dihydro-1H-indenyl, 2-oxo-2,3-dihydro-1H-benzo[d]imidazolyl, spiro[3.3]heptanyl, 2-azaspiro[3.3]heptanyl, and 2-oxaspiro[3.3]heptanyl, each of which is optionally substituted with 1 to 3 R⁴; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, or fourteenth embodiment or any alternative embodiments described therein. In some embodiments, R³ is optionally substituted with 1 to 2 R⁴. In some


10 embodiments, R³ is substituted with 1 R⁴. In some embodiments, R³ is unsubstituted.

In a sixteenth embodiment, for the compound of Formula (I) described in the first aspect or in the first embodiment, or a pharmaceutically acceptable salt thereof, Z is -CH₂-, R³ is thiophenyl, pyrrolyl, and 2-oxo-2,3-dihydro-1H-benzo[d]imidazolyl, each of which is optionally substituted with 1 to 2 R⁴; and the remaining variables are as described in the first

15 aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, or fourteenth embodiment or any alternative embodiments described therein.

In a seventeenth embodiment, for the compound of Formula (I) described in the first aspect or in the first embodiment, or a pharmaceutically acceptable salt thereof, R³ is












optionally substituted with 1 to 3 R⁴; and the remaining variables are as described in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth,

5 fourteenth, fifteenth, or sixteenth embodiment or any alternative embodiments described

therein. In an alternative seventh embodiment,. R^3 is





HN-; and O-; each of which is optionally substituted with 1 to 3 R⁴; and

the remaining variables are as described in the first or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, or

15 sixteenth embodiment or any alternative embodiments described therein.. In some
embodiments, R³ is optionally substituted with 1 to 2 R⁴. In some embodiments, R³ is
substituted with 1 R⁴. In some embodiments, R³ is unsubstituted.


In an eighteenth embodiment, for the compound of Formula (I) described in the first aspect or in the first embodiment, or a pharmaceutically acceptable salt thereof, R^3 is






R⁴ ; or ; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, or sixteenth embodiment or any alternative

embodiments described therein. In an alternative eighteenth embodiment, R³ is ;








third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, or fourteenth embodiment or any alternative embodiments described therein.



In a twentieth embodiment, for the compound of Formula (I) described in the first aspect or in the first embodiment, or a pharmaceutically acceptable salt thereof, R^3 is phenyl, pyrazolyl, pyridyl, and pyrimidyl, each of which is optionally substituted with 1 to 2 R^4 ; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, or fourteenth embodiment or any alternative embodiments described therein.

In a twenty-first embodiment, for the compound of Formula (I) described in the first aspect or in the first embodiment, or a pharmaceutically acceptable salt thereof, R^3 is

; each of which is optionally substituted with 1 to 2 R^4 ; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, or fourteenth embodiment or any alternative embodiments described therein.

In a twenty-second embodiment, for the compound of Formula (I) described in the first aspect or in the first embodiment, or a pharmaceutically acceptable salt thereof, R^3 is

represented by the following structural formula: ; ; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, or fourteenth embodiment or any alternative embodiments described therein. In an alternative twenty-


second embodiment, R^3 is ; ; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh,

twelfth, thirteenth, or fourteenth embodiment or any alternative embodiments described therein.


In a twenty-third embodiment, for the compound of Formula (I) described in the first embodiment or in the first embodiment, or a pharmaceutically acceptable salt thereof, R^4 is selected from C_{1-3} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, $-OR^{4a}$, $-CN$, $-NH_2$, $-NR^{4b}R^{4c}$, halo, $-C(O)R^{4a}$, $-NHC(O)R^{4a}$, $-NHC(O)OR^{4a}$, $-NR^{4b}SO_2R^{4a}$, $-SR^{4a}$, $-S(O)R^{4a}$, $-SO_2R^{4a}$, $-SO_2NR^{4b}R^{4c}$, $-P(O)R^{4b}R^{4c}$, $-C(O)O-C_{1-4}$ alkyl, benzyl, phenyl, 5 to 6-membered monocyclic heteroaryl, 6 to 10-membered bicyclic heterocyclyl, and 4 to 6-membered monocyclic heterocyclyl, wherein the C_{1-3} alkyl, C_{2-4} alkenyl, or C_{2-4} alkynyl represented by R^4 is optionally substituted with 1 to 3 substituents independently selected from halo, phenyl, 5 to 6-membered monocyclic heteroaryl optionally substituted with 1 or 2 R^{4g} , and 4 to 6-membered monocyclic heterocyclyl optionally substituted with 1 or 2 R^{4g} , $-C(O)NR^{4b}R^{4c}$, $-C(O)R^{4a}$, and OR^{4f} ; and the benzyl, phenyl, 5 to 6-membered monocyclic heteroaryl, 6 to 10-membered bicyclic heterocyclyl, and 4 to 6-membered monocyclic heterocyclyl represented by R^4 are each optionally substituted with 1 to 3 R^{4e} and further optionally substituted with one or two oxo, or two R^4 together form an oxo; R^{4a} is H, C_{1-4} alkyl optionally substituted with 1 to 4 R^{4d} , $-NR^{4b}R^{4c}$, C_{3-6} cycloalkyl, phenyl, 4- to 6-membered monocyclic heterocyclyl, or 5 to 6-membered monocyclic heteroaryl, wherein the phenyl, 4- to 6-membered monocyclic heterocyclyl, and 5 to 6-membered monocyclic heteroaryl are optionally substituted with 1 to 3 R^{4e} ; each R^{4b} is independently H or C_{1-4} alkyl; R^{4c} is H, C_{1-4} alkyl optionally substituted with 1 to 4 R^{4d} , 4 to 6-membered monocyclic heterocyclyl, or 5 to 6-membered monocyclic heteroaryl; each R^{4d} is independently selected from halo, $-OR^{4f}$, C_{3-6} cycloalkyl, $-C(O)OH$, $-NR^{4b}R^{4c}$, phenyl, and 5 to 6-membered monocyclic heteroaryl, wherein the phenyl and 5 to 6-membered monocyclic heteroaryl are each optionally substituted with 1 to 3 substituents independently selected from C_{1-3} alkyl, C_{1-3} haloalkyl, halo, CN and OH; each R^{4e} is independently selected from halo, $-OH$, C_{1-4} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy, $-C(O)R^{4h}$, $-SO_2C_{1-3}$ alkyl, $-C_{1-3}$ alkyl SO_2C_{1-3} alkyl, $-C(O)N(C_{1-3}alkyl)_2$, and cyano; or two R^{4e} together form an oxo; R^{4f} is H, C_{1-4} alkyl, phenyl or 5 to 6-membered monocyclic heteroaryl, wherein phenyl and 5 to 6-membered monocyclic heteroaryl are each optionally substituted with 1 to 3 halo; each R^{4g} is independently C_{1-3} alkyl or halo; and each R^{4h} is independently C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-3} alkoxyl or $-N(R^{4b})_2$; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, or twenty second embodiment or any

In a twenty-fourth embodiment, for the compound of Formula (I) described in the first
25 aspect or in the first embodiment, or a pharmaceutically acceptable salt thereof, R^4 is

22




; each of which is optionally substituted with 1 to 3 R^{4e} and further optionally substituted with 1 or 2 oxo; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, or twenty-second embodiment or any alternative embodiments described therein.




10 ; each of which is optionally substituted with 1 to 3 R^{4e}; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, or twenty-second embodiment or any alternative embodiments described therein.


15 In a twenty-fifth embodiment, for the compound of Formula (I) described in the first aspect or in the first embodiment, or a pharmaceutically acceptable salt thereof, R^4 is





10






5 and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, or twenty-second embodiment or any alternative embodiments described therein.


In a twenty-sixth embodiment, for the compound of Formula (I) described in the first aspect or in the first embodiment, or a pharmaceutically acceptable salt thereof, each R^{4e} is independently $-F$, $-CH_3$, $-CF_3$, $-CH_2CHF_2$, $-CH_2CF_3$, $-CH_2SO_2CH_3$, $-OH$, $-OCH_3$, $-SO_2CH_3$,
 10 $-C(O)CH_3$, $-C(O)cyclopropyl$, $-C(O)cyclobutyl$, $-C(O)cyclopentyl$, $-C(O)CH(CH_3)_2$, $-C(O)C(CH_3)_3$, $-C(O)N(CH_3)_2$, $-C(O)OCH_3$, or cyano; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, twenty-third, twenty-fourth or
 15 twenty-fifth embodiment or any alternative embodiments described therein. In an alternative twenty-sixth embodiment, R^{4e} is $-CH_3$, $-C(O)CH_3$, $-C(O)CH(CH_3)_2$, $-C(O)N(CH_3)_2$, or cyano; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-
 20 second, twenty-third, twenty-fourth or twenty-fifth embodiment or any alternative embodiments described therein.


In a twenty-seventh embodiment, for the compound of Formula (I) described in the first aspect or in the first embodiment, or a pharmaceutically acceptable salt thereof, R^4 is $-OR^{4a}$; R^{4a} is C_{1-3} alkyl optionally substituted with 1 to 3 R^{4d} or 4- to 6-membered monocyclic
 25 heterocyclyl optionally substituted with 1 to 3 R^{4e} ; each R^{4d} is independently halo, C_{1-2} alkyl, phenyl or 5 to 6-membered monocyclic heteroaryl, provided when two or three R^{4d} are present, one of the R^{4d} is phenyl or 5 to 6-membered monocyclic heteroaryl, and the other R^{4d}

are each independently halo or C₁₋₂alkyl; wherein the phenyl and 5 to 6-membered monocyclic heteroaryl represented by R^{4d} are each optionally substituted with 1 to 3 substituents independently selected from C₁₋₃alkyl, C₁₋₃haloalkyl and halo; each R^{4e} is independently C₁₋₄alkyl, C₁₋₄haloalkyl, -SO₂C₁₋₃alkyl, or -C(O)C₃₋₆cycloalkyl; the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, or twenty-third embodiment or any alternative embodiments described therein. In an alternative twenty-seventh embodiment, R⁴ is -OR^{4a}; R^{4a} is C₁₋₃alkyl substituted with 1 to 3 R^{4d}; one of the R^{4d} is phenyl or 5 to 6-membered monocyclic heteroaryl, and the other R^{4d} are each independently halo or C₁₋₂alkyl; wherein the phenyl and 5 to 6-membered monocyclic heteroaryl represented by R^{4d} are each optionally substituted with 1 to 3 halo; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, or twenty-third embodiment or any alternative embodiments described therein.

In a twenty-eighth embodiment, for the compound of Formula (I) described in the first aspect or in the first embodiment, or a pharmaceutically acceptable salt thereof, R⁴ is -CH₃, CH₂OH, -CHF₂, -CF₃, -CH(CF₃)OH, -CH₂OCH₃, -CH₂CN, -CH₂CF₃, -CH₂CH₂OH, -CH₂C(O)N(CH₃)₂, -C≡CC(CH₃)₂OH, -OH, -OCH₃, -OCH₂CH₃, -OCH₂CH₂OCH₃, -OCH(CH₃)₂, -OCF₃, -OCHF₂, -OCH₂CF₃, -O-cyclopropyl, -O-cyclobutyl, -OC(CH₃)₃, -OC(CH₃)₂CH₂OH, -OCH(CH₃)C(O)OH, -OC(CH₃)₂C(O)OH, -OCH(CH₃)CH₂OH, -OCH₂CH₂N(CH₃)₂, -CN, -NH₂, -NHCH₃, halo, -C(O)H, -C(O)CH(CH₃)₂, -C(O)OCH₃, -C(O)O-*t*-butyl, -N(CH₃)SO₂CH₃, -NHC(O)CH₃, -NHC(O)cyclopropyl, -NHC(O)OC(CH₃)₃, -C(O)CH₃, -C(O)OH, -C(O)NH₂, -C(O)NHCH₃, -C(O)N(CH₃)₂, -C(O)NC(CH₃)₃, -SCH₃, -S(O)CH₃, -SO₂CH₃, -SO₂N(CH₃)₂, -P(O)(CH₃)₂, phenyl, or R⁴ is represented by the following structural formula:








the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth,


- 10 nineteenth, twentieth, twenty-first, twenty-second, or twenty-third embodiment or any alternative embodiments described therein. In an alternative twenty-eighth embodiment, R^4 is $-\text{CH}_3$, CH_2OH , $-\text{CHF}_2$, $-\text{CF}_3$, $-\text{CH}(\text{CF}_3)\text{OH}$, $-\text{CH}_2\text{OCH}_3$, $-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{CF}_3$, $-\text{CH}_2\text{CH}_2\text{OH}$,

-CH₂C(O)N(CH₃)₂, -OH, -OCH₃, -OCH₂CH₃, -OCH(CH₃)₂, -OCF₃, -CN, -NH₂, -NHR^{4c}, halo, -C(O)H, -C(O)CH(CH₃)₂, -C(O)OCH₃, -C(O)O-*t*-butyl, -NHC(O)CH₃, -NHC(O)cyclopropyl, -NHC(O)OC(CH₃)₃, -C(O)CH₃, -C(O)OH, -C(O)NH₂, -C(O)NHCH₃, -C(O)N(CH₃)₂, -C(O)NC(CH₃)₃, -SCH₃, -SO₂CH₃, -SO₂N(CH₃)₂, -P(O)(CH₃)₂, phenyl,




second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, or twenty-third embodiment or any alternative embodiments described therein.

- 5 In a twenty-ninth embodiment, the compound of the present disclosure is represented by Formula (IIA), (IIB), or (IIC):




- or a pharmaceutically acceptable salt thereof, wherein m is 0 or 1; and the variables X , R^1 , R^2 , and R^3 depicted in Formula (IIA), (IIB), or (IIC) are as described in the first aspect or in the first, second, third, fourth, eleventh, fifteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, twenty-third, twenty-fourth, twenty-fifth, twenty-sixth, twenty-seventh, or twenty-eighth embodiment or any alternative embodiments described therein.

- 15 In a thirtieth embodiment, the compound of the present disclosure is represented by Formula (IIIA) or (IIIB):



- or a pharmaceutically acceptable salt thereof; where the variables X , R^1 , R^2 , and R^3 depicted in Formula (IIIA) or (IIIB) are as described in the first aspect or in the first, second, third, fourth, eleventh, fifteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, twenty-third, twenty-fourth, twenty-fifth, twenty-sixth, twenty-seventh, or twenty-eighth embodiment or any alternative embodiments described therein.

In a thirty-first embodiment, the compound of the present disclosure is represented by Formula (IVA), (IVB), (IVC), or (IVD):




5 or a pharmaceutically acceptable salt thereof, wherein m is 0 or 1; and the variables X , R^1 , R^2 , and R^3 depicted in Formula (IVA), (IVB), (IVC), or (IVD) are as described in the first aspect or in the first, second, third, fourth, eleventh, fifteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, twenty-third, twenty-fourth, twenty-fifth, twenty-sixth, twenty-seventh, or twenty-eighth embodiment or any alternative embodiments
10 described therein.

In a thirty-second embodiment, for compounds of Formula (I), (IIA), (IIB), (IIC), (IIIA), (IIIB), (IVA), (IVB), (IVC), or (IVD), or a pharmaceutically acceptable salt thereof, R^1 is $-\text{CH}_3$ or $-\text{CH}_2\text{CH}_3$; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth,
15 thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, twenty-third, twenty-fourth, twenty-fifth, twenty-sixth, twenty-seventh, twenty-eighth, twenty-ninth, thirtieth, or thirty-first embodiment or any alternative embodiments described therein.

In a thirty-third embodiment, for compounds of Formula (I), (IIA), (IIB), (IIC), (IIIA), (IIIB), (IVA), (IVB), (IVC), or (IVD), or a pharmaceutically acceptable salt thereof, R^1 is $-\text{CH}_3$; and the remaining variables are as described in the first aspect or in the first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, twenty-third, twenty-fourth, twenty-fifth, twenty-sixth, twenty-seventh, twenty-
20 eighth, twenty-ninth, thirtieth, or thirty-first embodiment or any alternative embodiments described therein.

In a thirty-fourth embodiment, the compound of the present disclosure is represented by Formula (V):



or a pharmaceutically acceptable salt thereof, wherein:

5 X is Cl or Br;

R¹ is -CH₃ or -CH₂CH₃;

ring A is selected from phenyl, thiophenyl, pyrrolyl, pyrazolyl, furanyl, isothiazolyl, and imidazolyl;

m is 0 or 1;

10 each R² is independently selected from C₁₋₃alkyl and -C(O)R^{2d}, wherein the C₁₋₃alkyl is optionally substituted with R^{2g};

R^{2d} is 4 to 6-membered monocyclic heterocyclyl;

R^{2g} is cyano or OH;

15 R³ is phenyl or 5 to 6-membered monocyclic heteroaryl, wherein the phenyl or 5 to 6-membered monocyclic heteroaryl are each optionally and independently substituted with 1 or 2 R⁴;

each R⁴ is independently selected from halo, C₁₋₄alkyl, -O-benzyl, -O-C₁₋₃alkyl, cyano, -NH₂, and 4 to 6-membered monocyclic heterocyclyl, wherein the benzyl is optionally substituted with 1 to 3 halo or C₁₋₃haloalkyl.

20 In an alternative thirty-fourth embodiment, for the compound of formula (V) or a pharmaceutically acceptable salt thereof, the definitions of variables are:

X is Cl or Br;

R¹ is -CH₃ or -CH₂CH₃;

25 ring A is selected from phenyl, thiophenyl, pyrrolyl, pyrazolyl, furanyl, isothiazolyl, and imidazolyl;

m is 0 or 1;

each R² is independently selected from C₁₋₃alkyl and -C(O)R^{2d}, wherein the C₁₋₃alkyl is optionally substituted with R^{2g};

R^{2d} is 4 to 6-membered monocyclic heterocyclyl;

R^{2g} is cyano or OH;

R³ is phenyl or 5 to 6-membered monocyclic heteroaryl, wherein the phenyl or 5 to 6-membered monocyclic heteroaryl are each optionally and independently substituted with 1 or 2 R⁴;


5 each R⁴ is independently selected from C₁₋₄alkyl, -O-benzyl, cyano, -NH₂, and 4 to 6-membered monocyclic heterocyclyl, wherein the benzyl is optionally substituted with halo.

In a thirty-fifth embodiment, for compounds of Formula (V), or a pharmaceutically

acceptable salt thereof, ring A is

10 ; and the remaining variables are as described in the thirty-fourth embodiment or any alternative embodiments described therein.

In a thirty-sixth embodiment, the compound of the present disclosure is represented by Formula (IIB), (IIIA), (IVA); or (IVB):





or a pharmaceutically acceptable salt thereof, wherein m is 0 or 1; and the variables X, R¹, R², and R³ depicted in Formula (IIIA), (IVA), or (IVB) are as described in the thirty-fourth embodiment or any alternative embodiments described therein.

In a thirty-seventh embodiment, for compounds of Formula (V), (IIB), (IIIA), (IVA),
20 or (IVB), or a pharmaceutically acceptable salt thereof, R³ is phenyl, pyrazolyl, pyridyl, and


pyrimidyl, each of which is optionally substituted with 1 to 2 R^4 ; and the remaining variables are as described in the thirty-fourth, thirty-fifth, or thirty-sixth embodiment or any alternative embodiments described therein.


- In a thirty-eighth embodiment, for compounds of Formula (V), (IIB), (IIIA), (IVA), or (IVB) or a pharmaceutically acceptable salt thereof, R^3 is represented by the following




- 10 eighth embodiment, R^3 is ; and the remaining variables are as described in the thirty-fourth, thirty-fifth, or thirty-sixth embodiment or any alternative embodiments described therein.

In a thirty-ninth embodiment, for compounds of Formula (V), (IIB), (IIIA), (IVA); or (IVB) or a pharmaceutically acceptable salt thereof, R^4 is $-F$, $-OCH(CH_3)_2$, $-OC(CH_3)_3$, $-$


- 15 CH_3 , cyano, $-NH_2$, or R^4 is represented by the following structural formula: 

; and the remaining variables are as described in the thirty-fourth, thirty-fifth, thirty-sixth, thirty-seventh, or thirty-eighth embodiment or any alternative embodiments described therein. In an alternative thirty-ninth embodiment, R^4 is $-$




 CH_3 , cyano, $-\text{NH}_2$, ; and the remaining variables are as described in the thirty-fourth, thirty-fifth, thirty-sixth, thirty-seventh, or thirty-eighth embodiment or any alternative embodiments described therein.

In a fortieth embodiment, for compounds of Formula (V), (IIB), (IIIA), (IVA), or
 5 (IVB) or a pharmaceutically acceptable salt thereof, R^2 is $-\text{CH}_3$, $-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{OH}$,




; and the remaining variables are as described in the thirty-fourth, thirty-fifth, thirty-sixth, thirty-seventh, thirty-eighth, or thirty-ninth embodiment or any alternative embodiments described therein.

In a forty-first embodiment, the present disclosure provides a compound represented
 10 by Formula (IIB) or (IVB):



(IIB);



(IVB);

or a pharmaceutically acceptable salt thereof, wherein:

m is 1;

R^1 is $-\text{CH}_3$;



15 X is Cl ;

R^2 is $-\text{CH}_3$;


R^3 is pyrimidinyl or pyridinyl, each of which is optionally substituted with 1 or 2 R^4 ;

20 R^4 is halo, $-\text{OC}_{1-3}\text{alkyl}$, or 4- to 6-membered monocyclic heterocyclyl, wherein the 4- to 6-membered monocyclic heterocycl is optionally substituted with 1 or 2 halo or $\text{C}_{1-3}\text{haloalkyl}$.

In a forty-second embodiment, for compounds of Formula (IIB) or (IVB), or a pharmaceutically acceptable salt thereof, R^3 is represented by the following structural

formula: ; or ; and the remaining variables are as described in the forty-first embodiment.

In a forty-third embodiment, for compounds of Formula (IIB) or (IVB), or a pharmaceutically acceptable salt thereof, each R^4 is independently $-F$, $-OCH(CH_3)_2$, or $-$

- 5 $OC(CH_3)_3$, or R^4 is represented by the following structural formula: ; and the remaining variables are as described in the forty-first or forty-second embodiment.

In a forty-fourth embodiment, the present disclosure provides a compound described herein (*e.g.*, a compound of any one of Examples 1-736), or a pharmaceutically acceptable salt thereof.

- 10 In a forty-fifth embodiment, the present disclosure provides a compound selected from the group consisting of:
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-phenyl-1H-imidazole-4-carboxamide;
 - 5-bromo-N-(3-fluoro-5-(methylsulfonamido)phenyl)-1H-pyrrole-3-carboxamide;
 - N-(3-fluoro-5-(methylsulfonamido)phenyl)-5-methyl-1H-pyrazole-3-carboxamide;
 - 15 5-bromo-N-(3-fluoro-5-(methylsulfonamido)phenyl)-1H-pyrazole-3-carboxamide;
 - N-(3-fluoro-5-(methylsulfonamido)phenyl)-5-methyl-1H-pyrrole-3-carboxamide;
 - N-(3-chloro-5-(methylsulfonamido)phenyl)-2-phenyl-1H-imidazole-4-carboxamide;
 - N-(3-chloro-5-(methylsulfonamido)phenyl)-4-phenylfuran-2-carboxamide;
 - N-(3-chloro-5-(methylsulfonamido)phenyl)-4-phenyl-1H-pyrrole-2-carboxamide;
 - 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-phenyl-1H-imidazole-2-carboxamide;
 - N-(3-bromo-5-(methylsulfonamido)phenyl)-4-methyl-3-(1H-pyrazol-1-yl)benzamide;
 - N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-phenylthiophene-2-carboxamide;
 - N-(3-bromo-5-(methylsulfonamido)phenyl)-2-phenylcyclobutane-1-carboxamide;
 - 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-3'-cyano-[1,1'-biphenyl]-3-carboxamide;
 - N-(3-chloro-5-(methylsulfonamido)phenyl)-3-(pyridin-4-yl)benzamide;
 - N-(3-bromo-5-(methylsulfonamido)phenyl)-4-phenylthiophene-2-carboxamide;
 - N-(3-iodo-5-(methylsulfonamido)phenyl)-4-phenylthiophene-2-carboxamide;

- N-(3-bromo-5-(methylsulfonamido)phenyl)-4-(pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-4-(pyrazin-2-yl)thiophene-2-carboxamide;
- 5 N-(3-bromo-5-(methylsulfonamido)phenyl)-4-(2-hydroxy-6-methylphenyl)thiophene-2-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-4-(2-hydroxyphenyl)-5-methylthiophene-2-carboxamide;
- 10 N-(3-bromo-5-(methylsulfonamido)phenyl)-4-(2-hydroxyphenyl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-(pyrimidin-2-yl)thiophene-2-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-5-methyl-4-(pyridin-2-yl)thiophene-2-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-(pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-4-(pyrimidin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(pyrimidin-2-yl)thiophene-2-carboxamide;
- 20 N-(3-chloro-5-(ethylsulfonamido)phenyl)-5-methyl-4-(pyrimidin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(ethylsulfonamido)phenyl)-5-methyl-4-(pyridin-2-yl)thiophene-2-carboxamide;
- 25 4-(3-(benzyloxy)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)thiophene-2-carboxamide;
- 4-(2-(benzyloxy)phenyl)-N-(3-bromo-5-(methylsulfonamido)phenyl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-methylpyridin-2-yl)thiophene-2-carboxamide;
- 30 4-(3-(benzyloxy)pyridin-2-yl)-N-(3-bromo-5-(methylsulfonamido)phenyl)thiophene-2-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-4-(3-methylpyridin-2-yl)thiophene-2-carboxamide;

- 4-(3-(benzyloxy)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-fluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 5 N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(4-cyanopyridin-2-yl)-1H-pyrazole-4-
- 10 carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(5-fluoropyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(pyrazin-2-yl)-1H-pyrazole-4-carboxamide;
- 15 N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(5-cyanopyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(5-methylpyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(5-(trifluoromethyl)pyridin-2-yl)-1H-
- 20 pyrazole-4-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(2-cyano-6-methylphenyl)-1H-pyrazole-4-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-5-methyl-1-(3-methylpyridin-2-yl)-1H-pyrrole-3-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-cyanopyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(trifluoromethyl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-1-cyclohexyl-1H-pyrazole-4-
- 30 carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(3-methylpyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(pyrimidin-2-yl)-1H-pyrazole-4-carboxamide;

- 1-(2-(benzyloxy)phenyl)-N-(3-bromo-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(3-cyanophenyl)-1H-pyrazole-4-carboxamide;
- 5 N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(5-cyano-2-methylphenyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-methylpyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 10 N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(5-morpholinopyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(6-(difluoromethyl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(6-(difluoromethoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 15 N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(2-formylphenyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(pyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-4-(1H-pyrazol-1-yl)thiophene-2-
- 20 carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(pyridin-3-ylmethoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(1-phenylethoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(pyridin-2-ylmethoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(pyridin-4-ylmethoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-((3-fluorobenzyl)oxy)pyridin-2-yl)-
- 30 1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(oxazol-5-ylmethoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 1-(2-(benzyloxy)-5-cyanophenyl)-N-(3-bromo-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;

- 1-(3-(benzyloxy)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-((4-fluorobenzyl)oxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-((3-chlorobenzyl)oxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 4-bromo-N-(3-fluoro-5-(methylsulfonamido)phenyl)thiophene-2-carboxamide;
- 4-bromo-N-(3-(ethylsulfonamido)-5-fluorophenyl)thiophene-2-carboxamide;
- N-(3-fluoro-5-(methylsulfonamido)phenyl)-4-phenylthiophene-2-carboxamide;
- 10 N-(3-(ethylsulfonamido)-5-fluorophenyl)-4-phenylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-phenylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(pyridin-2-yl)thiophene-2-carboxamide;
- 15 N-(3-bromo-5-(methylsulfonamido)phenyl)-4-(pyridin-3-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-ethyl-5-phenyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-phenyl-1-(2,2,2-trifluoroethyl)-1H-pyrrole-3-carboxamide;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(3-methylpyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(pyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
- 25 5-bromo-N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-1-methyl-5-(3-methylpyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-1-cyclohexyl-5-methyl-1H-pyrrole-3-carboxamide;
- 30 1-(5-(4-acetylpiperazin-1-yl)pyridin-2-yl)-N-(3-bromo-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;
- 1-(5-(4-acetylpiperazin-1-yl)-3-methylpyridin-2-yl)-N-(3-bromo-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;

- N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(5-(4-methyl-3-oxopiperazin-1-yl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 4-(6-(4-((3-chloro-5-(methylsulfonamido)phenyl)carbamoyl)-1H-pyrazol-1-yl)pyridin-3-yl)-N,N-dimethylpiperazine-1-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(2-oxopyrrolidin-1-yl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(4-morpholinopyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(2-methylmorpholino)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 10 1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-morpholinopyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(4-methyl-3-oxopiperazin-1-yl)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- 15 N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(cyanomethyl)-5-(pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-1-methyl-2-(3-methylpyridin-2-yl)-1H-imidazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(difluoro(phenyl)methoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 20 1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-cyano-1-(pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- 1-benzyl-N-(3-bromo-5-(methylsulfonamido)phenyl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazole-5-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-3-(3-cyano-1H-pyrazol-1-yl)benzamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-2-oxo-1-phenyl-2,3-dihydro-1H-imidazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-3-(2-chloropyridin-4-yl)benzamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(2-(2,2,2-trifluoro-1-hydroxyethyl)phenyl)-1H-pyrazole-4-carboxamide;
- 30 4-((1H-pyrrol-2-yl)methyl)-N-(3-bromo-5-(methylsulfonamido)phenyl)thiophene-2-carboxamide;
- N-(3-fluoro-5-(methylsulfonamido)phenyl)-4-(pyridin-3-yl)thiophene-2-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(2-hydroxyphenyl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(2-methoxyphenyl)thiophene-2-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-fluorophenyl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-phenyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(pyridin-3-yl)thiophene-2-carboxamide;
- 10 4-(2-(benzyloxy)phenyl)-N-(3-chloro-5-(methylsulfonamido)phenyl)thiophene-2-carboxamide;
- N-(3-fluoro-5-(methylsulfonamido)phenyl)-4-(pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-(ethylsulfonamido)-5-fluorophenyl)-4-(pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-fluoro-5-(methylsulfonamido)phenyl)-4-(2-hydroxyphenyl)thiophene-2-
- 15 carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(2,6-difluorophenyl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(pyridin-3-yl)thiophene-2-carboxamide;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-3-(1H-pyrazol-1-yl)benzamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-3-(1H-pyrazol-1-yl)benzamide;
- 1-ethyl-N-(3-fluoro-5-(methylsulfonamido)phenyl)-5-methyl-1H-pyrazole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(hydroxymethyl)-4-phenylthiophene-2-
- 25 carboxamide;
- N-(3-bromo-5-(methylsulfonylamidimidamido)phenyl)-4-(pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-bromo-5-(methylsulfonylamidimidamido)phenyl)-4-(3-methylpyridin-2-yl)thiophene-2-carboxamide;
- 30 N-(3-bromo-5-(methylsulfonylamidimidamido)phenyl)-5-methyl-4-(pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-phenylthiophene-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-phenylfuran-3-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-hydroxypropyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-hydroxyethyl)-1H-pyrazole-4-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-oxaspiro[3.3]heptan-6-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-phenyl-1H-pyrrole-3-carboxamide;
- 10 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(3-methylpyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(tetrahydro-2H-pyran-4-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-2-ethylimidazo[1,2-a]pyridine-6-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-3-(dimethylamino)-2-ethylimidazo[1,2-a]pyridine-6-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)imidazo[1,2-a]pyridine-6-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)benzofuran-7-carboxamide;
- tert-butyl 4-(4-((3-chloro-5-(methylsulfonamido)phenyl)carbamoyl)-1H-pyrazol-1-yl)piperidine-1-carboxylate;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-3-(pyridin-2-yl)isothiazole-5-carboxamide;
- 5-(2-(benzyloxy)phenyl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-1H-pyrazole-3-carboxamide;
- 25 3-(2-(benzyloxy)phenyl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-1H-pyrazole-5-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(pyridin-2-yl)-1H-pyrazole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-3-(pyridin-2-yl)-1H-pyrazole-5-
- 30 carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(1-phenylethyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2,2-difluoroethyl)-5-(3-methylpyridin-2-yl)-1H-pyrrole-3-carboxamide;

- tert-butyl 6-(4-((3-chloro-5-(methylsulfonamido)phenyl)carbamoyl)-1H-pyrazol-1-yl)-2-azaspiro[3.3]heptane-2-carboxylate;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(piperidin-4-yl)-1H-pyrazole-4-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-azaspiro[3.3]heptan-6-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(4-(piperazin-1-yl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 10 1-(4-(4-acetylpiperazin-1-yl)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(4-isobutyrylpiperazin-1-yl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-hydroxycyclohexyl)-1H-pyrazole-4-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(4-hydroxycyclohexyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-hydroxycyclohexyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-2-phenyl-1H-imidazole-4-carboxamide;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-2-(3-methylpyridin-2-yl)-1H-imidazole-4-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(2-oxopiperidin-1-yl)-1H-pyrrole-3-carboxamide;
- 25 1-(2-(benzyloxy)cyclobutyl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-3-(3-(hydroxymethyl)-1H-pyrazol-1-yl)benzamide;
- 30 N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(2-(hydroxymethyl)phenyl)-1H-pyrazole-4-carboxamide;
- 1-(3-aminopyridin-2-yl)-N-(3-bromo-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(hydroxymethyl)-4-(pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-3-(phenylamino)-1H-pyrazole-5-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(piperazine-1-carbonyl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N2-(3-chloro-5-(methylsulfonamido)phenyl)-N4-phenylthiophene-2,4-dicarboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(phenoxymethyl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 10 1-(3-(benzylamino)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(hydroxymethyl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoropyrimidin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-ethoxypyrimidin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 20 N-(3-chloro-5-(ethylsulfonamido)phenyl)-1-methyl-5-(pyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-((5-fluoropyridin-3-yl)methoxy)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-((5-fluoropyridin-3-yl)methoxy)pyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(2-oxopyrrolidin-1-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-cyclopropyl-4-(pyridin-2-yl)thiophene-2-carboxamide;
- 30 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(pyrimidin-2-yloxy)pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-((3-hydroxybenzyl)oxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(3-oxomorpholino)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 5 5-(aminomethyl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(tetrahydrofuran-3-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(4-methyl-3-oxopiperazin-1-yl)pyrimidin-2-yl)-1H-pyrazole-4-carboxamide;
- 10 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(4-methyl-3-oxopiperazin-1-yl)pyrazin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-(pyridin-3-ylmethoxy)pyridin-2-yl)-1-(2,2,2-trifluoroethyl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-((5-fluoropyridin-3-yl)methoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-((3,5-difluorobenzyl)oxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(pyrimidin-5-ylmethoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-((2-fluorobenzyl)oxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 20 1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(tetrahydro-2H-pyran-4-yl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 1-(5-(8-oxa-3-azabicyclo[3.2.1]octan-3-yl)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(5,6-dihydroimidazo[1,2-a]pyrazin-7(8H)-yl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(4-methyl-3-oxopiperazin-1-yl)-3-(pyridin-3-ylmethoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-morpholino-3-(pyridin-3-ylmethoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 30 1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(2-hydroxy-2-methylpropoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(pyrimidin-2-yloxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-methoxypyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(methylsulfonyl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(methylsulfonyl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(N,N-dimethylsulfamoyl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(dimethylphosphoryl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 10 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-fluoro-3-((3-fluorobenzyl)oxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-((3-cyanobenzyl)oxy)-5-fluoropyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-((3-fluorobenzyl)oxy)-6-methylpyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(thiazol-2-ylmethoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 1-(3-((1H-pyrazol-4-yl)methoxy)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-fluoropyrimidin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-ethoxypyrimidin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)indolizine-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-((3-fluorobenzyl)oxy)phenyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-fluoro-6-((3-fluorobenzyl)oxy)phenyl)-1H-pyrazole-4-carboxamide;
- 30 3-(3-(benzyloxy)pyridin-2-yl)-N5-(3-chloro-5-(methylsulfonamido)phenyl)-N2-methylthiophene-2,5-dicarboxamide;
- 3-(3-(benzyloxy)pyridin-2-yl)-N5-(3-chloro-5-(methylsulfonamido)phenyl)-N2,N2-dimethylthiophene-2,5-dicarboxamide;

- 4-(3-(benzyloxy)pyridin-2-yl)-N-(3-fluoro-5-(methylsulfonamido)phenyl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(oxazol-5-ylmethoxy)pyridin-2-yl)thiophene-2-carboxamide;
- 5 4-(3-(benzyloxy)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(hydroxymethyl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(pyridin-3-ylmethoxy)pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(methylthio)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 10 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(methylthio)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(cyclopropanecarboxamido)pyridin-2-yl)thiophene-2-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(2-(dimethylamino)-2-oxoethyl)pyridin-2-yl)thiophene-2-carboxamide;
- 2-(5-((3-chloro-5-(methylsulfonamido)phenyl)carbonyl)thiophen-3-yl)-N,N-dimethylnicotinamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(trifluoromethyl)pyridin-2-yl)thiophene-2-carboxamide;
- 20 N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(3-methylpyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-fluoro-3-((5-fluoropyridin-3-yl)methoxy)pyridin-2-yl)thiophene-2-carboxamide;
- 25 N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(5-fluoro-3-methylpyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-hydroxypyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-((3,5-difluorophenoxy)methyl)pyridin-2-yl)thiophene-2-carboxamide;
- 30 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5'-fluoro-[3,3'-bipyridin]-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)thieno[2,3-c]pyridine-2-carboxamide;

- 1-(3-((3-bromobenzyl)oxy)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-ethyl-4-(pyridin-2-yl)thiophene-2-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-isopropyl-4-(pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(methoxymethyl)-4-(pyridin-2-yl)thiophene-2-carboxamide;
- 4-(1-benzyl-1H-1,2,3-triazol-4-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methylthiophene-2-carboxamide;
- 10 methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-(1-phenyl-1H-1,2,3-triazol-4-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(pyrimidin-2-ylamino)pyridin-2-yl)thiophene-2-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(methyl(pyrimidin-2-yl)amino)pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(pyridin-3-yloxy)pyridin-2-yl)thiophene-2-carboxamide;
- 4-(2-(benzyloxy)-6-fluorophenyl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methylthiophene-2-carboxamide;
- 20 methylthiophene-2-carboxamide;
- 4-bromo-N-(3-chloro-5-(methylsulfonamido)phenyl)thiophene-2-carboxamide;
- methyl (5-((3-chloro-5-(methylsulfonamido)phenyl)carbonyl)thiophen-3-yl)prolinate;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(piperidin-4-yl)thiophene-2-carboxamide;
- 25 carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(piperazin-1-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-morpholinothiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(4-(2,2,2-trifluoroethyl)piperazin-1-yl)thiophene-2-carboxamide;
- 30 N-(3-chloro-5-(cyclopropanesulfonamido)phenyl)-5-methyl-1-(5-morpholinopyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-((difluoromethyl)sulfonamido)phenyl)-5-methyl-1-(5-morpholinopyridin-2-yl)-1H-pyrrole-3-carboxamide;

- N-(3-chloro-5-(ethylsulfonamido)phenyl)-5-methyl-1-(5-morpholinopyridin-2-yl)-1H-pyrrole-3-carboxamide;
- 1-(8-oxabicyclo[3.2.1]octan-3-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-fluoro-3-(3-fluorophenethyl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 1-(3-((1H-imidazol-2-yl)methoxy)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;
- 1-(3-((1H-pyrazol-5-yl)methoxy)pyridin-2-yl)-N-(3-chloro-5-
- 10 (methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;
- 1-(3-((1H-imidazol-4-yl)methoxy)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-((3-fluorobenzyl)oxy)-6-methylphenyl)-1H-pyrazole-4-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-((3,5-difluorophenoxy)methyl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-(5-fluoropyridin-3-yl)phenyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-((3,5-difluorobenzyl)oxy)-5-(2-oxa-
- 20 6-azaspiro[3.3]heptan-6-yl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-phenoxycyclopentyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-phenylcyclopentyl)-1H-pyrazole-4-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-methyl-3-(oxazol-5-ylmethoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(((5-fluoropyridin-3-yl)oxy)methyl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-(pyrimidin-2-yloxy)phenyl)-1H-
- 30 pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-(pyridin-3-yloxy)phenyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-(3-fluorophenoxy)phenyl)-1H-pyrazole-4-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(1-methyl-2-oxopiperidin-4-yl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-fluoropyridin-2-yl)-5-(trifluoromethyl)-1H-pyrrole-3-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoro-3-(oxazol-5-ylmethoxy)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(methyl(tetrahydrofuran-3-yl)amino)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(pyrrolidin-1-yl)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- 10 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-morpholinopyridin-2-yl)-5-(trifluoromethyl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(2,2-dimethylmorpholino)pyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(5-morpholinopyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-ethyl-5-(5-morpholinopyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-morpholinopyridin-2-yl)-1-(2,2,2-trifluoroethyl)-1H-pyrrole-3-carboxamide;
- 20 1-(5-(2-oxa-6-azaspiro[3.3]heptan-6-yl)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1H-pyrrole-3-carboxamide;
- 5-(5-(2-oxa-6-azaspiro[3.3]heptan-6-yl)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-1H-pyrrole-3-carboxamide;
- 25 5-(5-(2-oxa-6-azaspiro[3.3]heptan-6-yl)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2,2,2-trifluoroethyl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(3-(thiazol-2-ylmethoxy)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(3-(thiazol-2-ylmethoxy)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- 30 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-hydroxypyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-cyclopentyl-5-methyl-1H-pyrrole-3-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(4-isobutyrylpiperazin-1-yl)pyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- 4-(6-(4-((3-chloro-5-(methylsulfonamido)phenyl)carbamoyl)-2-methyl-1H-pyrrol-1-yl)pyridin-3-yl)-N,N-dimethylpiperazine-1-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(1,1-dioxidothiomorpholino)pyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2,2-difluoroethyl)-5-(5-fluoropyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(1-hydroxyethyl)-1-(pyridin-2-yl)-1H-
10 pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)thieno[3,2-c]pyridine-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)thieno[3,2-b]pyridine-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-7-methylthieno[3,2-c]pyridine-2-
carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-5,6-dimethoxybenzo[b]thiophene-2-
carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)pyrrolo[1,2-b]pyridazine-6-carboxamide;
- 6-chloro-N-(3-chloro-5-(methylsulfonamido)phenyl)benzo[b]thiophene-2-
carboxamide;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-6-fluorobenzo[b]thiophene-2-
carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-fluorobenzo[b]thiophene-2-
carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-fluorobenzo[b]thiophene-2-
25 carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-7-fluorobenzo[b]thiophene-2-
carboxamide;
- 5-chloro-N-(3-chloro-5-(methylsulfonamido)phenyl)benzo[b]thiophene-2-
carboxamide;
- 30 6-chloro-N-(3-chloro-5-(methylsulfonamido)phenyl)indolizine-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-6-methylindolizine-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-oxo-1,2-dihydropyrrolo[1,2-a]pyrazine-
7-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-8-(trifluoromethyl)indolizine-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-7-methoxybenzo[b]thiophene-2-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methylindolizine-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-8-fluoroindolizine-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-hydroxycyclobutyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(1-methyl-2-oxopiperidin-4-yl)-1H-
- 10 pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-phenethyl-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-cyclopentyl-1H-pyrazole-4-carboxamide;
- 1-(6-acetamidospiro[3.3]heptan-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-
- 15 1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-methyl-2-azaspiro[3.3]heptan-6-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-(2,2,2-trifluoroethyl)-2-azaspiro[3.3]heptan-6-yl)-1H-pyrazole-4-carboxamide;
- 20 1-(2-acetyl-2-azaspiro[3.3]heptan-6-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;
- N-(3-bromo-5-(methylsulfonamido)phenyl)-2,4,6-d₃-4-phenylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2,2-dimethylpiperidin-4-yl)-1H-
- 25 pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)-1H-pyrazole-4-carboxamide;
- 1-(1-acetylpiperidin-4-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;
- 30 tert-butyl (6-(4-((3-chloro-5-(methylsulfonamido)phenyl)carbamoyl)-1H-pyrazol-1-yl)spiro[3.3]heptan-2-yl)carbamate;
- 1-(6-aminospiro[3.3]heptan-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide; and

- 1-(3-fluoro-5-morpholinopyridin-2-yl)-5-methyl-N-(3-(methylsulfonamido)phenyl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-fluoro-5-morpholinopyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoropyridin-2-yl)-1-(2-hydroxyethyl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(4-methyl-3-oxopiperazin-1-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-morpholinopyrimidin-2-yl)-1H-pyrazole-4-carboxamide;
- 10 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5'-fluoro-[3,3'-bipyridin]-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-methyl-2-oxoimidazolidin-1-yl)thiophene-2-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-methoxybenzo[b]thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1,3-dimethylpyrrolo[1,2-a]pyrazine-7-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(4-phenylpiperazine-1-carbonyl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(trifluoromethyl)benzo[b]thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4,5-difluorobenzo[b]thiophene-2-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-4,7-difluorobenzo[b]thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(trifluoromethyl)benzo[b]thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(pyrimidin-5-yl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 30 N-(3-chloro-5-(methylsulfonamido)phenyl)-6-methoxybenzo[b]thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(piperidine-1-carbonyl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(morpholine-4-carbonyl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3,4,4-trimethyl-2,5-dioxoimidazolidin-1-yl)thiophene-2-carboxamide;
- 5 4-(3-(2-(azetidin-1-yl)-2-oxoethyl)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(2-oxooxazolidin-3-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-morpholinopyrazin-2-yl)-1H-pyrazole-4-carboxamide;
- 10 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(pyrimidin-2-ylmethyl)pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(pyridin-2-yl)-5-(4-(2,2,2-trifluoroethyl)piperazine-1-carbonyl)-1H-pyrrole-3-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-(2-oxopyrrolidin-1-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-hydroxy-5-(2-oxa-6-azaspiro[3.3]heptan-6-yl)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(2-oxo-2-(pyrrolidin-1-yl)ethyl)pyridin-2-yl)thiophene-2-carboxamide;
- 20 N-(3-fluoro-5-(methylsulfonamido)phenyl)-5-methyl-4-(3-methylpyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(4-methylpiperazine-1-carbonyl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(pyridin-3-yloxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(3-(pyridin-3-yloxy)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(2-(dimethylamino)-2-oxoethyl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 30 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(2-hydroxypropan-2-yl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-6,8-dimethylindolizine-2-carboxamide;

- N-(3-fluoro-5-(methylsulfonamido)phenyl)-4-(3-methylpyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(pyridin-2-yl)-5-(2,6-diazaspiro[3.3]heptane-2-carbonyl)-1H-pyrrole-3-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-8-methylindolizine-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)pyrrolo[1,2-a]pyrazine-7-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3,5-difluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoro-3-methoxypyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 10 methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(5-(trifluoromethyl)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-cyanopyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methylbenzo[b]thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-ethoxypyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-(1-(tetrahydro-2H-pyran-4-yl)-1H-1,2,3-triazol-4-yl)thiophene-2-carboxamide;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(5-(trifluoromethyl)pyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-ethoxypyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-methoxypyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(methoxymethyl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-fluoropyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- 30 methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-ethoxypyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-fluoropyrimidin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-ethoxypyridin-2-yl)thiophene-2-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-chloropyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-methoxypyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(pyridin-3-yloxy)-5,6-dihydro-4H-
- 10 cyclopenta[b]thiophene-2-carboxamide;
- 4-chloro-N-(3-chloro-5-(methylsulfonamido)phenyl)benzo[b]thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(3-fluorophenoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(4-methoxy-1H-pyrazol-1-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-fluoropyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-(1-phenethyl-1H-imidazol-2-
- 20 yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(4-methoxy-1H-pyrazol-1-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(methoxymethyl)pyrimidin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-cyanopyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-chloropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-chloropyrimidin-2-yl)-1-methyl-1H-
- 30 pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(3-fluoroazetidin-1-yl)pyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-fluoropyridin-2-yl)-5-(methoxymethyl)-1H-pyrrole-3-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-ethoxypyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(4-ethoxy-1H-pyrazol-1-yl)-5-methylthiophene-2-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(4-fluoro-1H-pyrazol-1-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(4-fluoro-1H-pyrazol-1-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-methoxypyrimidin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- 10 1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(methoxymethyl)pyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-methoxypyrimidin-2-yl)thiophene-2-carboxamide;
- 15 5-(5-(azetidin-1-yl)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-methoxypyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(3,3-difluoropyrrolidin-1-yl)pyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- 20 2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(methyl(tetrahydrofuran-3-yl)amino)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(methyl(tetrahydrofuran-3-yl)amino)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(trifluoromethyl)pyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(2-methoxyethoxy)pyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- 30 methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-ethoxypyrimidin-2-yl)thiophene-2-carboxamide;
- 1-(5-(azetidin-1-yl)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1H-pyrrole-3-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-(cyclopropylmethoxy)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-cyclopropyl-4-(5-fluoropyrimidin-2-yl)thiophene-2-carboxamide;
- 5 1-(5-(azetidin-1-yl)pyrimidin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(3-fluoroazetidin-1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(5-fluoropyrimidin-2-yl)thiophene-2-
- 10 carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-chloropyrimidin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- 5-chloro-N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-fluoropyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(2-oxopyrrolidin-1-yl)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5,5'-difluoro-[3,3'-bipyridin]-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-fluoro-1'-methyl-2'-oxo-1',2'-
- 20 dihydro-[3,4'-bipyridin]-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-fluoro-1'-methyl-6'-oxo-1',6'-dihydro-[3,3'-bipyridin]-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5,5'-difluoro-[3,3'-bipyridin]-2-yl)-5-methylthiophene-2-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(3,3-difluoroazetidin-1-yl)pyrimidin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(methylsulfonyl)pyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(trifluoromethyl)pyridin-2-
- 30 yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(cyclopropylmethoxy)pyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- tert-butyl 3-(4-((3-chloro-5-(methylsulfonamido)phenyl)carbamoyl)-2-(5-fluoropyrimidin-2-yl)-1H-pyrrol-1-yl)azetidine-1-carboxylate;

- 1-(5-(3-azabicyclo[3.1.0]hexan-3-yl)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1H-pyrrole-3-carboxamide;
N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(5-fluoropyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
5 N-(3-chloro-5-(ethylsulfonamido)phenyl)-5-cyclopropyl-4-(5-fluoropyrimidin-2-yl)thiophene-2-carboxamide;
N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(methylthio)pyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(methylsulfinyl)pyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
10 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(5-(pyrrolidin-1-yl)pyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
N-(3-chloro-5-(methylsulfonamido)phenyl)-5-cyclopropyl-1-(5-fluoropyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
15 5-(5-(azetidin-1-yl)pyrimidin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-1H-pyrrole-3-carboxamide;
N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(4,4-dimethyl-2,5-dioxo-3-(pyridin-2-yl)imidazolidin-1-yl)-5-methylthiophene-2-carboxamide;
N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-fluoro-2'-methyl-[3,3'-bipyridin]-2-yl)-5-methylthiophene-2-carboxamide;
20 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-fluoro-3-(oxazol-5-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(oxetan-3-yloxy)pyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
25 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-ethoxypyrimidin-2-yl)-5-(trifluoromethyl)-1H-pyrrole-3-carboxamide;
N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(difluoromethoxy)pyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
30 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-fluoro-3-methoxypyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3,5-difluoropyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;

- 5-chloro-N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-fluoropyrimidin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(ethylsulfonamido)phenyl)-5-(5-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-ethyl-4-(5-fluoropyrimidin-2-yl)thiophene-2-carboxamide;
- 1-(5-(5-azaspiro[2.4]heptan-5-yl)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(3-fluoroazetidin-1-yl)pyrimidin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- 10 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-(5-(N-methylmethylsulfonamido)pyrimidin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-cyclopropyl-5-(5-fluoropyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(2-oxopyrrolidin-1-yl)pyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoropyrimidin-2-yl)-1-isopropyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-(5-(methylsulfonyl)pyrimidin-2-yl)thiophene-2-carboxamide;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5,5'-difluoro-[3,3'-bipyridin]-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3,4-dimethyl-2,5-dioxo-4-(pyridin-2-yl)imidazolidin-1-yl)-5-methylthiophene-2-carboxamide;
- 25 N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(5-fluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-fluoro-5-methylpyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- methyl 4-(6-(4-((3-chloro-5-(methylsulfonamido)phenyl)carbamoyl)-1-methyl-1H-pyrrol-2-yl)pyridin-3-yl)piperazine-1-carboxylate;
- 30 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(3,3-difluoroazetidin-1-yl)pyrimidin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(5-((1-methylazetidin-3-yl)oxy)pyridin-2-yl)-1H-pyrrole-3-carboxamide;

- methyl 4-(6-(5-((3-chloro-5-(methylsulfonamido)phenyl)carbamoyl)-2-methylthiophen-3-yl)-5-methylpyridin-3-yl)piperazine-1-carboxylate;
N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(3,5-difluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 5 N-(3-chloro-5-(ethylsulfonamido)phenyl)-5-(5-fluoropyrimidin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(3-fluoroazetidin-1-yl)pyrimidin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-fluoro-3-methylpyridin-2-yl)-5-
- 10 methylthiophene-2-carboxamide;
N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3,5-difluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;
N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(5-fluoro-3-methylpyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-fluoro-3-methoxypyridin-2-yl)-5-methylthiophene-2-carboxamide;
N-(3-chloro-5-(ethylsulfonamido)phenyl)-1-(3-chloro-5-fluoropyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-methoxypyridin-2-yl)-5-
- 20 methylthiophene-2-carboxamide;
N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(oxetan-3-yloxy)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
N-(3-chloro-5-(ethylsulfonamido)phenyl)-5-(3,5-difluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(5-(pyrrolidin-1-yl)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(5-(tetrahydro-2H-pyran-4-yl)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-chloro-5-fluoropyridin-2-yl)-1-
- 30 methyl-1H-pyrrole-3-carboxamide;
N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoro-3-methylpyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
N-(3-chloro-5-(ethylsulfonamido)phenyl)-1-(3,5-difluoropyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;

- 1-(azetidin-3-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoropyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-cyano-5-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 5 N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(5-methoxypyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(5-fluoro-3-methoxypyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(3-chloro-5-fluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 10 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-(5-(methylamino)pyrimidin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(2-oxooxazolidin-3-yl)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(2,4,6-trifluorophenyl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(ethylsulfonamido)phenyl)-5-(3-fluoro-5-methylpyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-(2,4,6-trifluorophenyl)thiophene-2-carboxamide;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(2-oxooxazolidin-3-yl)pyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(4-(cyclopropanecarbonyl)piperazin-1-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(5-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-(5-(trifluoromethyl)pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(ethylsulfonamido)phenyl)-5-(5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 30 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-(5-(trifluoromethyl)pyrimidin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-fluoro-5-methoxypyridin-2-yl)-5-methylthiophene-2-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-chloro-5-fluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(5-(4-pivaloylpiperazin-1-yl)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-methoxypyrimidin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(4-(cyclopentanecarbonyl)piperazin-1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 10 1-(5-(7-azabicyclo[2.2.1]heptan-7-yl)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(3-methoxyazetidin-1-yl)pyrimidin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(3,3-difluoroazetidin-1-yl)-3-fluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(2-cyano-4-fluoro-6-methylphenyl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(3,3-difluoroazetidin-1-yl)-3-methylpyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(3-fluoro-5-methoxypyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 20 N-(3-chloro-5-(ethylsulfonamido)phenyl)-5-methyl-4-(3-methyl-5-(trifluoromethyl)pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-methoxy-5-(trifluoromethyl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(4-(cyclobutanecarbonyl)piperazin-1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(4,4-difluoropiperidin-1-yl)-3-methylpyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(tetrahydro-1H-furo[3,4-c]pyrrol-5(3H)-yl)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- 30 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(6-(3,3-difluoroazetidin-1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-methoxy-3-methylpyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3,5-difluoropyridin-4-yl)-5-methylthiophene-2-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(3,3-difluoroazetidin-1-yl)-3-methylpyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-cyano-5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(4-(cyclopropanecarbonyl)piperazin-10 1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(3,3-difluoroazetidin-1-yl)-3-fluoropyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(3,3-difluoroazetidin-1-yl)-3-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 15 1-(3-chloro-5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-isopropoxypyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(4,4-difluoropiperidin-1-yl)-3-20 methylpyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-(difluoromethoxy)-5-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(ethylsulfonamido)phenyl)-1-(5-(3,3-difluoroazetidin-1-yl)pyrimidin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(4-(3,3-difluoroazetidin-1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-chloro-5-methoxypyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(3,3-difluoroazetidin-1-yl)-3-methoxypyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- 30 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-(5-(2,2,2-trifluoroethoxy)pyrimidin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(3,3-difluoroazetidin-1-yl)-3-(trifluoromethyl)pyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(3,3-difluoroazetidin-1-yl)-3-isopropoxy-pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-cyclobutoxypyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- 5 N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(5-ethoxypyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(3,3-difluoroazetidin-1-yl)-3-methoxy-pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(3,3-difluoroazetidin-1-yl)-3-methoxy-pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 10 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(5-(3,4,4-trimethyl-2,5-dioxoimidazolidin-1-yl)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(difluoromethoxy)pyrimidin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 15 N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(5-(3,3-difluoroazetidin-1-yl)pyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-fluoro-5-(trifluoromethyl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(5-(3-fluoroazetidin-1-yl)pyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- 20 5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-ethoxy-3-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(ethylsulfonamido)phenyl)-5-methyl-4-(5-(trifluoromethyl)pyrimidin-2-yl)thiophene-2-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(3-fluorooxetan-3-yl)pyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoropyridin-2-yl)-1-(methylsulfonyl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-fluoro-5-methoxypyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 30 methyl-1H-pyrrole-3-carboxamide;
- 4-(5-(1-acetylpiperidin-4-yl)pyrimidin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(cyclopropanesulfonamido)phenyl)-5-(5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;

- N-(3-chloro-5-(propylsulfonamido)phenyl)-5-(5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- methyl 4-(6-(4-((3-chloro-5-(methylsulfonamido)phenyl)carbamoyl)-1-methyl-1H-pyrrol-2-yl)-5-fluoropyridin-3-yl)piperazine-1-carboxylate;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(4-methoxypiperidin-1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-cyano-5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(3,3-difluoroazetidin-1-yl)-3-(2,2,2-trifluoroethoxy)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 10 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(3,3-difluoroazetidin-1-yl)-3-(2,2,2-trifluoroethoxy)pyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-cyclobutoxy-5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-(5-((1-methylazetidin-3-yl)oxy)pyrimidin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(3,3-difluoroazetidin-1-yl)-3-(difluoromethoxy)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 4-(5-(tert-butoxy)pyrimidin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methylthiophene-2-carboxamide;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(4-(cyclopropanecarbonyl)piperazin-1-yl)-3-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-ethoxy-5-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(3-fluoro-3-methylazetidin-1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-cyclobutoxy-5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(5-(3-(methylsulfonyl)azetidin-1-yl)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- 30 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(3-methoxyazetidin-1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-fluoro-5-(3-fluoroazetidin-1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-fluoro-5-(6-azaspiro[2.5]octan-6-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-fluoro-5-(6-azaspiro[2.5]octan-6-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-fluoro-5-(2-oxa-6-azaspiro[3.4]octan-6-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(4,4-difluoropiperidin-1-yl)-3-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(4,4-dimethyl-1,4-azasilinan-1-yl)-3-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 10 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(6,6-difluoro-2-azaspiro[3.3]heptan-2-yl)-3-fluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(4,4-difluoropiperidin-1-yl)-3-fluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(3,3-difluoroazetidin-1-yl)pyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(5-(3-((methylsulfonyl)methyl)azetidin-1-yl)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-fluoro-5-(4-(trifluoromethyl)piperidin-1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(4,4-dimethyl-1,4-azasilinan-1-yl)-3-fluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-fluoro-5-(2-oxa-7-azaspiro[3.5]nonan-7-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-fluoro-5-(2-oxa-7-azaspiro[3.5]nonan-7-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(6,6-dimethyl-3-azabicyclo[3.1.0]hexan-3-yl)-3-fluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(3,3-difluoroazetidin-1-yl)-3-isopropoxypyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- 30 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-fluoro-5-(3-(trifluoromethyl)azetidin-1-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(1,1-difluoro-5-azaspiro[2.4]heptan-5-yl)-3-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(3,3-difluoropyrrolidin-1-yl)-3-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(6,6-dimethyl-3-azabicyclo[3.1.0]hexan-3-yl)-3-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 5 N-(3-chloro-5-((2-hydroxyethyl)sulfonamido)phenyl)-5-(5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(6,6-difluoro-2-azaspiro[3.3]heptan-2-yl)-3-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(ethylsulfonamido)phenyl)-5-(5-(3,3-difluoroazetidin-1-yl)-3-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 10 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(1,1-difluoro-6-azaspiro[2.5]octan-6-yl)-3-fluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(3,3-difluoropyrrolidin-1-yl)pyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-((1-(cyclopropanecarbonyl)azetidin-3-yl)oxy)-3-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(1,1-difluoro-5-azaspiro[2.4]heptan-5-yl)-3-fluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(3,3-difluoroazetidin-1-yl)-3-isopropoxy-pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-fluoro-5-(4-(trifluoromethyl)piperidin-1-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(3,3-difluoropiperidin-1-yl)-3-fluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(1,1-difluoro-6-azaspiro[2.5]octan-6-yl)-3-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(3,3-difluoropiperidin-1-yl)-3-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-(5-(1-(2,2,2-trifluoroethyl)azetidin-3-yl)pyridin-2-yl)thiophene-2-carboxamide;
- 30 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(1-(2,2-difluoroethyl)piperidin-4-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(3,3-difluoropyrrolidin-1-yl)-3-fluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-cyano-5-(trifluoromethyl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(3,3-difluoropyrrolidin-1-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-fluoro-5-(3-(trifluoromethyl)azetidin-1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-fluoro-5-(trifluoromethyl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-fluoro-5-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 10 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-fluoro-5-((1-methylazetidin-3-yl)oxy)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-cyanopyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-fluoro-5-isopropoxypyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-ethoxy-3-fluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(3,3-difluoroazetidin-1-yl)-3-methylpyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-isopropoxypyrimidin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-fluoro-5-isopropoxypyridin-2-yl)thiophene-2-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(trifluoromethyl)pyrimidin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-cyano-3-methylpyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-cyanopyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- 30 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(trifluoromethyl)pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-cyano-3-fluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(3,3-difluoroazetidin-1-yl)pyrimidin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-isopropoxypyridin-2-yl)thiophene-2-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(3,3-difluoropyrrolidin-1-yl)-3-fluoropyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(3,3-difluoropyrrolidin-1-yl)pyrimidin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- 10 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(3,3-difluoroazetidin-1-yl)-3-fluoropyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-(5-(3-(trifluoromethyl)azetidin-1-yl)pyrimidin-2-yl)thiophene-2-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(3,3-difluoropyrrolidin-1-yl)pyrimidin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(3,3-difluoropyrrolidin-1-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(6,6-difluoro-2-azaspiro[3.3]heptan-2-yl)pyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(6,6-difluoro-2-azaspiro[3.3]heptan-2-yl)pyrimidin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(1,1-difluoro-5-azaspiro[2.4]heptan-5-yl)pyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(1,1-difluoro-5-azaspiro[2.4]heptan-5-yl)-3-fluoropyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(6,6-difluoro-2-azaspiro[3.3]heptan-2-yl)pyrimidin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-((1-hydroxy-2-methylpropan-2-yl)oxy)pyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- 30 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-ethyl-4-(3-fluoropyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-fluoro-5-(trifluoromethyl)pyridin-2-yl)thiophene-2-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(5-(3-(trifluoromethyl)azetidin-1-yl)pyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
4-(5-(tert-butoxy)-3-fluoropyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methylthiophene-2-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(dimethylphosphoryl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3,5-difluoropyridin-2-yl)-5-ethylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(1,1-difluoro-5-azaspiro[2.4]heptan-5-yl)pyrimidin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 10 N-(3-chloro-5-((2-hydroxyethyl)sulfonamido)phenyl)-4-(5-(3,3-difluoroazetidin-1-yl)-3-methoxypyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(1,1-difluoro-5-azaspiro[2.4]heptan-5-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-4-(5-(3-(trifluoromethyl)azetidin-1-yl)pyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(5-(3-(trifluoromethyl)azetidin-1-yl)pyridin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-ethyl-4-(5-fluoropyridin-2-yl)thiophene-2-carboxamide;
- 20 4-(5-(tert-butoxy)pyrimidin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(dimethylphosphoryl)pyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- 25 4-(5-(tert-butoxy)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(6,6-difluoro-2-azaspiro[3.3]heptan-2-yl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(6,6-difluoro-2-azaspiro[3.3]heptan-2-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 30 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-(1,1-difluoro-5-azaspiro[2.4]heptan-5-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 1-(5-(tert-butoxy)-3-fluoropyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1H-pyrrole-3-carboxamide;

- 1-(5-(tert-butoxy)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1H-pyrrole-3-carboxamide;
- 5-(5-(tert-butoxy)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-1H-pyrrole-3-carboxamide;
- 5 N-(3-chloro-5-(propylsulfonamido)phenyl)-4-(3-methylpyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3,5-difluoropyridin-2-yl)-1H-pyrrole-3-carboxamide;
- 4-(5-(tert-butoxy)pyrimidin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-
- 10 (hydroxymethyl)thiophene-2-carboxamide;
- N-(3-chloro-5-((1-methylethyl)sulfonamido)phenyl)-4-(3-methylpyridin-2-yl)thiophene-2-carboxamide;
- 5-(5-(tert-butoxy)pyrimidin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-1H-pyrrole-3-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-methylpyridin-2-yl)thiophene-2-carboxamide;
- N-(3-chloro-5-(N-methylmethylsulfonamido)phenyl)-4-(3-methylpyridin-2-yl)thiophene-2-carboxamide;
- 1-(5-(tert-butoxy)pyrimidin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-
- 20 methyl-1H-pyrrole-3-carboxamide;
- 5-(5-(tert-butoxy)-3-fluoropyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-1H-pyrrole-3-carboxamide;
- 5-(3,5-difluoropyridin-2-yl)-N-(3-fluoro-5-(methylsulfonamido)phenyl)-1-methyl-1H-pyrrole-3-carboxamide;
- 25 N-(3-chloro-5-((2-hydroxyethyl)sulfonamido)phenyl)-5-(3,5-difluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-3-fluoro-4-(5-isopropoxypyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-3-methyl-4-(3-methylpyridin-2-
- 30 yl)thiophene-2-carboxamide;
- 4-(3-fluoro-5-(3-(trifluoromethyl)azetidin-1-yl)pyridin-2-yl)-N-(3-fluoro-5-(methylsulfonamido)phenyl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-fluoropyridin-2-yl)-5-methylthiazole-2-carboxamide;

- N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(3-fluoro-5-(3-(trifluoromethyl)azetidin-1-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-((2-hydroxyethyl)sulfonamido)phenyl)-4-(3-fluoro-5-(3-(trifluoromethyl)azetidin-1-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-fluoro-5-((1-methylazetidin-3-yl)oxy)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 4-(5-(azetidin-3-yloxy)-3-fluoropyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(1,1-dioxidothiomorpholino)-5-fluoropyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 10 4-(3-(4-acetylpiperazin-1-yl)-5-fluoropyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-fluoro-3-(3-methoxypyrrolidin-1-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 15 N-(3-chloro-5-((methyl)sulfonamido-1,1,1-d₃)phenyl)-4-(5-isopropoxypyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-fluoro-3-(4-(methylsulfonyl)piperazin-1-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-fluoro-3-(6-azaspiro[2.5]octan-6-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(5-(3-(trifluoromethyl)azetidin-1-yl)pyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
- 2-(5-((3-chloro-5-(methylsulfonamido)phenyl)carbonyl)-2-methylthiophen-3-yl)-3,5-difluoropyridine 1-oxide;
- 25 4-(5-(azetidin-3-yloxy)pyrimidin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-fluoro-3-(3-hydroxy-3-methylbut-1-yn-1-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-fluoro-5-(3-hydroxy-1-methylazetidin-3-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 30 N-(2-fluoro-3-(methylsulfonamido)phenyl)-4-(3-(pyridin-3-ylmethoxy)pyridin-2-yl)thiophene-2-carboxamide;
- 1-(2-(benzyloxy)cyclopentyl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;

- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-(((5-fluoropyridin-3-yl)oxy)methyl)phenyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-fluoro-6-((5-fluoropyridin-3-yl)methoxy)phenyl)-1H-pyrazole-4-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-(dimethylcarbamoyl)-6-((3-fluorobenzyl)oxy)phenyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2,4-difluoro-6-(oxazol-5-yl)methoxy)phenyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-fluoro-3-(oxazol-5-yl)methoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide;
- 10 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-((5-fluoropyridin-3-yl)methoxy)pyridin-2-yl)thiophene-2-carboxamide;
- 1-(2-(benzyloxy)phenyl)-N-(3-fluoro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-fluoro-6-(oxazol-5-yl)methoxy)phenyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-((3-fluorobenzyl)oxy)-4-(2-oxa-6-azaspiro[3.3]heptan-6-yl)phenyl)-1H-pyrazole-4-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-((5-fluoropyridin-3-yl)methoxy)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-((5-fluoropyridin-3-yl)methoxy)pyridin-2-yl)-5-(hydroxymethyl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(5-fluoro-3-(thiazol-2-yl)methoxy)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoro-3-((2-methylthiazol-4-yl)methoxy)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-((3,5-dimethylisoxazol-4-yl)methoxy)-5-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(5-fluoro-3-(pyrimidin-5-yl)methoxy)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 30 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(2-((5-fluoropyridin-3-yl)methoxy)-4-methylpyridin-3-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoro-3-((1-methyl-1H-1,2,3-triazol-5-yl)methoxy)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;

- (E)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(2-(2-(5-fluoropyridin-3-yl)vinyl)phenyl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoro-3-((1-methyl-1H-tetrazol-5-yl)methoxy)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoro-3-(2-(5-fluoropyridin-3-yl)ethyl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- (E)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoro-3-(2-(5-fluoropyridin-3-yl)vinyl)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 4-(3-((5-fluoropyridin-3-yl)methoxy)pyridin-2-yl)-5-methyl-N-(3-
- 10 (methylsulfonamido)phenyl)thiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-(1-(3-fluoro-5-(trifluoromethyl)phenyl)ethoxy)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-((1S)-1-(3-fluoro-5-(trifluoromethyl)phenyl)ethoxy)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-((1R)-1-(3-fluoro-5-(trifluoromethyl)phenyl)ethoxy)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-((5-ethylpyridin-3-yl)methoxy)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(1-(5-fluoropyridin-3-
- 20 yl)ethoxy)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- (R)-N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(1-(5-fluoropyridin-3-yl)ethoxy)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- (S)-N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(1-(5-fluoropyridin-3-yl)ethoxy)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 25 N-(3-chloro-5-((2-hydroxyethyl)sulfonamido)phenyl)-4-(3-((5-fluoropyridin-3-yl)methoxy)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-fluoro-3-(oxazol-5-ylmethoxy)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(3,3-difluoroazetidin-1-yl)-3-
- 30 (oxazol-5-ylmethoxy)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-((3,5-difluorobenzyl)oxy)-2-methylpyrimidin-4-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoro-3-(pyrimidin-5-ylmethoxy)pyridin-2-yl)-1-(2,2,2-trifluoroethyl)-1H-pyrrole-3-carboxamide;

- 5-(3-(((6-(2-hydroxypropan-2-yl)pyridin-2-yl)methoxy)pyridin-2-yl)-1-methyl-N-(3-(methylsulfonamido)phenyl)-1H-pyrrole-3-carboxamide;
- 5-(3-(((3-hydroxybenzyl)oxy)pyridin-2-yl)-1-methyl-N-(3-(methylsulfonamido)phenyl)-1H-pyrrole-3-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(1-(3,5-difluorophenyl)ethoxy)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- (S)-N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(1-(3,5-difluorophenyl)ethoxy)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- (R)-N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(1-(3,5-difluorophenyl)ethoxy)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- 10 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(((3,5-difluorobenzyl)oxy)pyrimidin-4-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(((1-isopropylazetidin-3-yl)oxy)pyrimidin-2-yl)-1-(2,2,2-trifluoroethyl)-1H-pyrrole-3-carboxamide;
- 15 (S)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(((1-methylpyrrolidin-3-yl)oxy)pyrimidin-2-yl)-1-(2,2,2-trifluoroethyl)-1H-pyrrole-3-carboxamide;
- (R)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(((1-methylpyrrolidin-3-yl)oxy)pyrimidin-2-yl)-1-(2,2,2-trifluoroethyl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2,2-difluoroethyl)-5-(5-(((1-methylazetidin-3-yl)oxy)pyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
- 20 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(((1-(methylsulfonyl)azetidin-3-yl)oxy)pyrimidin-2-yl)-1-(2,2,2-trifluoroethyl)-1H-pyrrole-3-carboxamide;
- (R)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-fluoro-5-(((1-methylpyrrolidin-3-yl)oxy)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 25 (S)-N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-fluoro-5-(((1-methylpyrrolidin-3-yl)oxy)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 2-((2-(5-(((3-chloro-5-(methylsulfonamido)phenyl)carbamoyl)-2-methylthiophen-3-yl)pyrimidin-5-yl)oxy)propanoic acid;
- N-(3-chloro-5-((2-hydroxyethyl)sulfonamido)phenyl)-4-(5-isopropoxypyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- 30 N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-hydroxypyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- 4-(5-(tert-butoxy)pyrimidin-2-yl)-N-(3-chloro-5-(ethylsulfonamido)phenyl)-5-methylthiophene-2-carboxamide;

- 2-((2-(5-((3-chloro-5-(methylsulfonamido)phenyl)carbamoyl)-2-methylthiophen-3-yl)pyrimidin-5-yl)oxy)-2-methylpropanoic acid;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-((1-hydroxypropan-2-yl)oxy)pyrimidin-2-yl)-5-methylthiophene-2-carboxamide;
- 5 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-(2-(dimethylamino)ethoxy)-3-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-fluoro-5-(1-hydroxy-1-(1-methylazetidin-3-yl)ethyl)pyridin-2-yl)-5-methylthiophene-2-carboxamide;
- N-(3-chloro-5-(ethylsulfonamido)phenyl)-4-(5-isopropoxypyrimidin-2-yl)-5-
- 10 methylthiophene-2-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-fluoro-5-((1-(2,2,2-trifluoroethyl)azetidin-3-yl)oxy)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- 2-(4-((3-chloro-5-(methylsulfonamido)phenyl)carbamoyl)-1-methyl-1H-pyrrol-2-yl)-3,5-difluoropyridine 1-oxide;
- 15 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-fluoro-5-((1-isopropylazetidin-3-yl)oxy)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-((1-(2,2-difluoroethyl)azetidin-3-yl)oxy)-3-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2,2-difluoroethyl)-5-(5-(3-
- 20 (trifluoromethyl)azetidin-1-yl)pyrimidin-2-yl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-fluoro-6-methoxypyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(3-fluoropyridin-2-yl)-1-(2,2,2-trifluoroethyl)-1H-pyrrole-3-carboxamide;
- 25 N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-methoxypyrimidin-2-yl)-1-(2,2,2-trifluoroethyl)-1H-pyrrole-3-carboxamide;
- N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoropyrimidin-2-yl)-1-(2,2,2-trifluoroethyl)-1H-pyrrole-3-carboxamide; and N-(3-chloro-5-(methylsulfonamido)phenyl)-5-
- (5-((1-methylazetidin-3-yl)oxy)pyrimidin-2-yl)-1-(2,2,2-trifluoroethyl)-1H-pyrrole-3-
- 30 carboxamide;
- or a pharmaceutically acceptable salt thereof.

The compounds and intermediates described herein may be isolated and used as the compound per se. Alternatively, when a moiety is present that is capable of forming a salt, the compound or intermediate may be isolated and used as its corresponding salt. As used

herein, the terms “salt” or “salts” refers to an acid addition or base addition salt of a compound described herein. “Salts” include in particular “pharmaceutical acceptable salts”. The term “pharmaceutically acceptable salts” refers to salts that retain the biological effectiveness and properties of the compounds described herein and, which typically are not
5 biologically or otherwise undesirable. In many cases, the compounds of the present disclosure are capable of forming acid and/or base salts by virtue of the presence of amino and/or carboxyl groups or groups similar thereto.

Pharmaceutically acceptable acid addition salts can be formed with inorganic acids or organic acids, e.g., acetate, aspartate, benzoate, besylate, bromide/hydrobromide,
10 bicarbonate/carbonate, bisulfate/sulfate, camphorsulfonate, chloride/hydrochloride, chlorthalophyllonate, citrate, ethandisulfonate, fumarate, gluceptate, gluconate, glucuronate, hippurate, hydroiodide/iodide, isethionate, lactate, lactobionate, laurylsulfate, malate, maleate, malonate, mandelate, mesylate, methylsulphate, naphthoate, napsylate, nicotinate, nitrate, octadecanoate, oleate, oxalate, palmitate, pamoate, phosphate/hydrogen
15 phosphate/dihydrogen phosphate, polygalacturonate, propionate, stearate, succinate, sulfate, sulfosalicylate, tartrate, tosylate and trifluoroacetate salts.

Inorganic acids from which salts can be derived include, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like.

Organic acids from which salts can be derived include, for example, acetic acid,
20 propionic acid, glycolic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, toluenesulfonic acid, sulfosalicylic acid, and the like.

Pharmaceutically acceptable base addition salts can be formed with inorganic and organic bases.

25 Inorganic bases from which salts can be derived include, for example, ammonium salts and metals from columns I to XII of the periodic table. In certain embodiments, the salts are derived from sodium, potassium, ammonium, calcium, magnesium, iron, silver, zinc, and copper; particularly suitable salts include ammonium, potassium, sodium, calcium and magnesium salts.

30 Organic bases from which salts can be derived include, for example, primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, basic ion exchange resins, and the like. Certain organic amines include isopropylamine, benzathine, choline, diethanolamine, diethylamine, lysine, meglumine, piperazine and tromethamine.

The salts can be synthesized by conventional chemical methods from a compound containing a basic or acidic moiety. Generally, such salts can be prepared by reacting free acid forms of these compounds with a stoichiometric amount of the appropriate base (such as Na, Ca, Mg, or K hydroxide, carbonate, bicarbonate or the like), or by reacting free base forms of these compounds with a stoichiometric amount of the appropriate acid. Such reactions are typically carried out in water or in an organic solvent, or in a mixture of the two. Generally, use of non-aqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile is desirable, where practicable. Lists of additional suitable salts can be found, e.g., in "Remington's Pharmaceutical Sciences", 20th ed., Mack Publishing Company, Easton, Pa., (1985); and in "Handbook of Pharmaceutical Salts: Properties, Selection, and Use" by Stahl and Wermuth (Wiley-VCH, Weinheim, Germany, 2002).

Isotopically-labeled compounds of Formula (I) can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the accompanying Examples and Preparations using an appropriate isotopically-labeled reagents in place of the non-labeled reagent previously employed. In one embodiment, the present disclosure provides deuterated compounds described herein or a pharmaceutically acceptable salt thereof.

Pharmaceutically acceptable solvates in accordance with the disclosure include those wherein the solvent of crystallization may be isotopically substituted, e.g. D₂O, d₆-acetone, d₆-DMSO.

It will be recognized by those skilled in the art that the compounds of the present disclosure may contain chiral centers and as such may exist in different stereoisomeric forms. As used herein, the term "an optical isomer" or "a stereoisomer" refers to any of the various stereo isomeric configurations which may exist for a given compound of the present disclosure. It is understood that a substituent may be attached at a chiral center of a carbon atom. Therefore, the disclosure includes enantiomers, diastereomers or racemates of the compound.

"Enantiomers" are a pair of stereoisomers that are non-superimposable mirror images of each other. A 1:1 mixture of a pair of enantiomers is a "racemic" mixture. The term "racemic" or "rac" is used to designate a racemic mixture where appropriate. When designating the stereochemistry for the compounds of the present disclosure, a single stereoisomer with known relative and absolute configuration of the two chiral centers is designated using the conventional RS system (e.g., (1S,2S)). "Diastereoisomers" are stereoisomers that have at least two asymmetric atoms, but which are not mirror-images of

each other. The absolute stereochemistry is specified according to the Cahn-Ingold-Prelog R-S system. When a compound is a pure enantiomer the stereochemistry at each chiral carbon may be specified by either R or S. Resolved compounds whose absolute configuration is unknown can be designated (+) or (-) depending on the direction (dextro- or levorotatory) which they rotate plane polarized light at the wavelength of the sodium D line. Alternatively, the resolved compounds can be defined by the respective retention times for the corresponding enantiomers/diastereomers via chiral HPLC.

Certain of the compounds described herein contain one or more asymmetric centers or axes and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that may be defined, in terms of absolute stereochemistry, as (R)- or (S)-.

Unless specified otherwise, the compounds of the present disclosure are meant to include all such possible stereoisomers, including racemic mixtures, optically pure forms and intermediate mixtures. Optically active (R)- and (S)-stereoisomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques (e.g., separated on chiral SFC or HPLC chromatography columns, such as CHIRALPAK^{RTM} and CHIRALCEL^{RTM} available from DAICEL Corp. using the appropriate solvent or mixture of solvents to achieve good separation). If the compound contains a double bond, the substituent may be E or Z configuration. If the compound contains a disubstituted cycloalkyl, the cycloalkyl substituent may have a cis- or trans-configuration. All tautomeric forms are also intended to be included.

The present disclosure also provides a pharmaceutical composition comprising a compound described herein (e.g., a compound according to any one of the preceding embodiments), or a pharmaceutically acceptable salt thereof, and one or more pharmaceutically acceptable carriers.

METHODS OF USE

The compounds described herein have DHX9 inhibitory activity. As used herein, "DHX9 inhibitory activity" refers to the ability of a compound or composition to induce a detectable decrease in DHX9 activity *in vivo* or *in vitro* (e.g., at least 10% decrease in DHX9 activity as measured by a given assay such as the bioassay described in the examples and known in the art).

In certain embodiments, the present disclosure provides a method of treating a disease or disorder responsive to inhibition of DHX9 activity (referred herein as "DHX9 mediated disease or disorder") in a subject in need of the treatment. The method comprises

administering to the subject a compound described herein (*e.g.*, a compound described in any one of the first to forty-fifth embodiments) or a pharmaceutically acceptable salt thereof or a pharmaceutical composition thereof.

5 In certain embodiments, the present disclosure provides the use of a compound described herein (*e.g.*, a compound described in any one of the first to forty-fifth embodiments) or a pharmaceutically acceptable salt thereof or a pharmaceutical composition comprising a compound described herein or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of a DHX9 mediated disorder or disease in a subject in need of the treatment.

10 In certain embodiments, the present disclosure provides a compound described herein (*e.g.*, a compound described in any one of the first to forty-fifth embodiments) or a pharmaceutically acceptable salt thereof or a pharmaceutical composition comprising a compound described herein or a pharmaceutically acceptable salt thereof for use in the treatment of a DHX9 mediated disorder or disease in a subject in need of the treatment.

15 In certain embodiments, the DHX9 mediated disease or disorder is selected from cancer, viral infections, and autoimmune disease.

In some embodiments, the present disclosure provides a method of treating cancer. In some embodiments, the cancer is selected from colorectal, endometrial, ovarian, gastric, hematopoietic, breast, brain, skin, lung, blood, prostate, head and neck, pancreatic, bladder,
20 bone, soft-tissue, kidney, and liver cancer. In some embodiments the cancer is selected from colorectal, endometrial, ovarian, hematopoietic, and gastric cancer. In some embodiments, the cancer is colorectal cancer. In some embodiments, the cancer is Ewing's Sarcoma.

In some embodiments, the cancer is a microsatellite instable (MSI) cancer. In some embodiments, the cancer is MSI-high cancer. In other embodiments, the cancer is MSI-low cancer. MSI is determined by PCR analysis of 5 different nucleotide repeats, which is
25 dependent on the cancer type. MSI-low cancers are characterized by instability at only 1 of the 5 sites; while MSI-high cancers are characterized by instability at 2 or more of the 5 sites (G. Yang *et al.* Correlations between microsatellite instability and the biological behavior of tumors. *Journal of Cancer Research and Clinical Oncology*, 2019) .

30 MSI-high cancer is additionally characterized by defective mismatch repair (dMMR) (M. Lorenzi, *et al.* Epidemiology of Microsatellite Instability High (MSI-H) and Deficient Mismatch Repair (dMMR) in Solid Tumors: A Structured Literature Review. *Journal of Oncology*, Volume 2020, Article ID 1897929). For example, dMMR in colorectal cancer can be determined by MLH1 promoter hypermethylation, rendering MLH1 inactive, which

comprises about 80-90% of MSI-high colorectal cancers. MSI-high colorectal cancer with inactivated MLH1 is also named sporadic MSI-high colorectal cancer. Alternatively, dMMR can be determined by immunohistochemistry mutation status of MLH1, MSH2, MSH6, MSH3, PMS1 and/or PMS2 mismatch repair (MMR) proteins. In MSI-high colorectal cancer, 5 MLH1 and MSH2 are the 2 predominantly mutated MMR proteins. They are mutated in about 10-20% of MSI-high colorectal cancers. MSI-high colorectal cancers with these mutations are also known as Lynch Syndrome cancers.

In some embodiments, the cancer is a MSI cancer and/or has mutations or defects in DNA mis-match repair (MMR), and/or mutations or defects in RNA splicing and the 10 kinetochore complex.

The compounds, or pharmaceutically acceptable salts thereof described herein may be used to decrease the expression or activity of DHX9, or to otherwise affect the properties and/or behavior of DHX9 in a cell.

One embodiment of the present disclosure includes a method of decreasing the 15 expression or activity of DHX9, or to otherwise affect the properties and/or behavior of DHX9 in a subject comprising administering to said subject an effective amount of at least one compound described herein, or a pharmaceutically acceptable salt thereof.

In certain embodiments, the present disclosure relates to the aforementioned methods, wherein said subject is a mammal.

20 In certain embodiments, the present disclosure relates to the aforementioned methods, wherein said subject is a primate.

In certain embodiments, the present disclosure relates to the aforementioned methods, wherein said subject is a human.

25 As used herein, an “effective amount” and a “therapeutically effective amount” can be used interchangeably. It means an amount effective for treating or lessening the severity of one or more of the diseases, disorders or conditions as recited herein. In some embodiments, the effective dose can be between 10 µg and 500 mg.

The compounds and compositions, according to the methods of the present disclosure, may be administered using any amount and any route of administration effective for treating 30 or lessening the severity of one or more of the diseases, disorders or conditions recited above.

In certain embodiments, the present disclosure relates to the aforementioned methods, wherein said compound is administered parenterally.

In certain embodiments, the present disclosure relates to the aforementioned methods, wherein said compound is administered intramuscularly, intravenously, subcutaneously, orally, pulmonary, rectally, intrathecally, topically or intranasally.

In certain embodiments, the present disclosure relates to the aforementioned methods,
5 wherein said compound is administered systemically.

The compounds of the present disclosure are typically used as a pharmaceutical composition (e.g., a compound of the present disclosure and at least one pharmaceutically acceptable carrier). As used herein, the term “pharmaceutically acceptable carrier” includes generally recognized as safe (GRAS) solvents, dispersion media, surfactants, antioxidants,
10 preservatives (e.g., antibacterial agents, antifungal agents), isotonic agents, salts, preservatives, drug stabilizers, buffering agents (e.g., maleic acid, tartaric acid, lactic acid, citric acid, acetic acid, sodium bicarbonate, sodium phosphate, and the like), and the like and combinations thereof, as would be known to those skilled in the art (see, for example, Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990, pp. 1289-
15 1329). Except insofar as any conventional carrier is incompatible with the active ingredient, its use in the therapeutic or pharmaceutical compositions is contemplated. For purposes of this disclosure, solvates and hydrates are considered pharmaceutical compositions comprising a compound of the present disclosure and a solvent (i.e., solvate) or water (i.e., hydrate).

The formulations may be prepared using conventional dissolution and mixing
20 procedures. For example, the bulk drug substance (i.e., compound of the present disclosure or stabilized form of the compound (e.g., complex with a cyclodextrin derivative or other known complexation agent)) is dissolved in a suitable solvent in the presence of one or more of the excipients described above. The compound of the present disclosure is typically formulated into pharmaceutical dosage forms to provide an easily controllable dosage of the drug and to
25 give the patient an elegant and easily handleable product.

The pharmaceutical composition (or formulation) for application may be packaged in a variety of ways depending upon the method used for administering the drug. Generally, an article for distribution includes a container having deposited therein the pharmaceutical formulation in an appropriate form. Suitable containers are well-known to those skilled in the
30 art and include materials such as bottles (plastic and glass), sachets, ampoules, plastic bags, metal cylinders, and the like. The container may also include a tamper-proof assemblage to prevent indiscreet access to the contents of the package. In addition, the container has deposited thereon a label that describes the contents of the container. The label may also include appropriate warnings.

The pharmaceutical composition comprising a compound of the present disclosure is generally formulated for use as a parenteral or oral administration or alternatively suppositories.

For example, the pharmaceutical oral compositions of the present disclosure can be made up in a solid form (including without limitation capsules, tablets, pills, granules, powders or suppositories), or in a liquid form (including without limitation solutions, suspensions or emulsions). The pharmaceutical compositions can be subjected to conventional pharmaceutical operations such as sterilization and/or can contain conventional inert diluents, lubricating agents, or buffering agents, as well as adjuvants, such as preservatives, stabilizers, wetting agents, emulsifiers and buffers, etc.

Typically, the pharmaceutical compositions are tablets or gelatin capsules comprising the active ingredient together with

- a) diluents, e.g., lactose, dextrose, sucrose, mannitol, sorbitol, cellulose and/or glycine;
- b) lubricants, e.g., silica, talcum, stearic acid, its magnesium or calcium salt and/or polyethylene glycol; for tablets also
- c) binders, e.g., magnesium aluminum silicate, starch paste, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose and/or polyvinylpyrrolidone; if desired
- d) disintegrants, e.g., starches, agar, alginic acid or its sodium salt, or effervescent mixtures; and/or
- e) absorbents, colorants, flavors and sweeteners.

Tablets may be either film coated or enteric coated according to methods known in the art.

Suitable compositions for oral administration include a compound of the disclosure in the form of tablets, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use are prepared according to any method known in the art for the manufacture of pharmaceutical compositions and such compositions can contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets may contain the active ingredient in admixture with nontoxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients are, for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium

phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example, starch, gelatin or acacia; and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets are uncoated or coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate can be employed. Formulations for oral use can be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example, peanut oil, liquid paraffin or olive oil.

The parenteral compositions (e.g, intravenous (IV) formulation) are aqueous isotonic solutions or suspensions. The parenteral compositions may be sterilized and/or contain adjuvants, such as preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for regulating the osmotic pressure and/or buffers. In addition, they may also contain other therapeutically valuable substances. The compositions are generally prepared according to conventional mixing, granulating or coating methods, respectively, and contain about 0.1-75%, or contain about 1-50%, of the active ingredient.

The compound of the present disclosure or pharmaceutical composition thereof for use in a subject (e.g., human) is typically administered orally or parenterally at a therapeutic dose. When administered intravenously via infusion, the dosage may depend upon the infusion rate at which an IV formulation is administered. In general, the therapeutically effective dosage of a compound, the pharmaceutical composition, or the combinations thereof, is dependent on the species of the subject, the body weight, age and individual condition, the disorder or disease or the severity thereof being treated. A physician, pharmacist, clinician or veterinarian of ordinary skill can readily determine the effective amount of each of the active ingredients necessary to prevent, treat or inhibit the progress of the disorder or disease.

The above-cited dosage properties are demonstrable in vitro and in vivo tests using advantageously mammals, e.g., mice, rats, dogs, monkeys or isolated organs, tissues and preparations thereof. The compounds of the present disclosure can be applied in vitro in the form of solutions, e.g., aqueous solutions, and in vivo either enterally, parenterally, advantageously intravenously, e.g., as a suspension or in aqueous solution. The dosage in vitro may range between about 10^{-3} molar and 10^{-9} molar concentrations.

DEFINITIONS

As used herein, a “patient,” “subject” or “individual” are used interchangeably and refer to either a human or non-human animal. The term includes mammals such as humans.

5 Typically, the animal is a mammal. A subject also refers to for example, primates (e.g., humans, male or female), cows, sheep, goats, horses, dogs, cats, rabbits, rats, mice, fish, birds and the like. In certain embodiments, the subject is a primate. In some embodiments, the subject is a human.

10 As used herein, the term “inhibit”, “inhibition” or “inhibiting” refers to the reduction or suppression of a given condition, symptom, or disorder, or disease, or a significant decrease in the baseline activity of a biological activity or process.

As used herein, the term “treat”, “treating” or “treatment” of any disease, condition or disorder, refers to the management and care of a patient for the purpose of combating the disease, condition, or disorder and includes the administration of a compound of the present disclosure to obtaining desired pharmacological and/or physiological effect. The effect can be therapeutic, which includes achieving, partially or substantially, one or more of the following results: partially or totally reducing the extent of the disease, condition or disorder; ameliorating or improving a clinical symptom, complications or indicator associated with the disease, condition or disorder; or delaying, inhibiting or decreasing the likelihood of the progression of the disease, condition or disorder; or eliminating the disease, condition or disorder. In certain embodiments, the effect can be to prevent the onset of the symptoms or complications of the disease, condition or disorder.

15
20

As used herein the term “cancer” has the meaning normally accepted in the art. The term can broadly refer to abnormal cell growth.

25 As used herein, a subject is “in need of” a treatment if such subject would benefit biologically, medically or in quality of life from such treatment (in some embodiments, a human).

As used herein, the phrase “optionally substituted” is used interchangeably with the phrase “substituted or unsubstituted.” In general the term “optionally substituted” refers to the replacement of hydrogen radicals in a given structure with the radical of a specified substituent. Specific substituents are described in the definitions and in the description of compounds and examples thereof. Unless otherwise indicated, an optionally substituted group can have a substituent at each substitutable position of the group, and when more than one

30

position in any given structure can be substituted with more than one substituent selected from a specified group, the substituent can be either the same or different at every position.

As used herein, the term “alkyl” refers to a fully saturated branched or unbranched hydrocarbon moiety. The term “C₁₋₄alkyl” refers to an alkyl having 1 to 4 carbon atoms. The terms “C₁₋₃alkyl” and “C₁₋₂alkyl” are to be construed accordingly. Representative examples of “C₁₋₄alkyl” include, but are not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, iso-butyl, and tert-butyl. Similarly, the alkyl portion (i.e., alkyl moiety) of an alkoxy have the same definition as above. When indicated as being “optionally substituted”, the alkane radical or alkyl moiety may be unsubstituted or substituted with one or more substituents (generally, one to three substituents except in the case of halogen substituents such as perchloro or perfluoroalkyls).

As used herein, the term “alkylene” refers to a fully saturated branched or unbranched divalent hydrocarbon radical. The term “C₁₋₄alkylene” refers to an alkylene having 1 to 4 carbon atoms. The terms “C₁₋₃alkylene” and “C₁₋₂alkylene” are to be construed accordingly. Representative examples of “C₁₋₄alkylene” include, but are not limited to, methylene, ethylene, n-propylene, iso-propylene, n-butylene, sec-butylene, iso-butylene, and tert-butylene.

As used herein, the term “alkoxy” refers to a fully saturated branched or unbranched alkyl moiety attached through an oxygen bridge (i.e. a --O-- C₁₋₄ alkyl group wherein C₁₋₄ alkyl is as defined herein). Representative examples of alkoxy include, but are not limited to, methoxy, ethoxy, propoxy, 2-propoxy, butoxy, tert-butoxy and the like. In some embodiments, alkoxy groups have about 1-4 carbons, and in some embodiments about 1-2 carbons. The term “C₁₋₂ alkoxy” is to be construed accordingly.

As used herein, the term “C₁₋₄ alkoxyC₁₋₄ alkyl” refers to a C₁₋₄ alkyl group as defined herein, wherein at least of the hydrogen atoms is replaced by an C₁₋₄ alkoxy. The C₁₋₄ alkoxyC₁₋₄ alkyl group is connected through the rest of the molecule described herein through the alkyl group.

The number of carbon atoms in a group is specified herein by the prefix “C_{x-xx}”, wherein x and xx are integers. For example, “C₁₋₃ alkyl” is an alkyl group which has from 1 to 3 carbon atoms.

“Halogen” or “halo” may be fluorine, chlorine, bromine or iodine.

As used herein, the term “halo-substituted-C₁₋₄alkyl” or “C₁₋₄haloalkyl” refers to a C₁₋₄alkyl group as defined herein, wherein at least one of the hydrogen atoms is replaced by a halo atom. The C₁₋₄haloalkyl group can be monohalo-C₁₋₄alkyl, dihalo-C₁₋₄alkyl or polyhalo-

C₁₋₄ alkyl including perhalo-C₁₋₄alkyl. A monohalo-C₁₋₄alkyl can have one iodo, bromo, chloro or fluoro within the alkyl group. Dihalo-C₁₋₄alkyl and polyhalo-C₁₋₄alkyl groups can have two or more of the same halo atoms or a combination of different halo groups within the alkyl. Typically the polyhalo-C₁₋₄alkyl group contains up to 9, or 8, or 7, or 6, or 5, or 4, or 3,
5 or 2 halo groups. Non-limiting examples of C₁₋₄haloalkyl include fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl and dichloropropyl. A perhalo-C₁₋₄alkyl group refers to a C₁₋₄alkyl group having all hydrogen atoms replaced with halo atoms.

10 The term “aryl” refers to an aromatic carbocyclic single ring or two fused ring system containing 6 to 10 carbon atoms. Examples include phenyl, indanyl, tetrahydronaphthalene, and naphthyl.

The term “heteroaryl” refers to a 5- to 12-membered aromatic radical containing 1-4 heteroatoms selected from N, O, and S. In some instances, nitrogen atoms in a heteroaryl may
15 be quaternized. The term “heteroaryl” may be used interchangeably with the terms “heteroaryl ring”, “heteroaryl group”, or “heteroaromatic”. A heteroaryl group may be mono- or bi-cyclic. Monocyclic heteroaryl includes, for example, pyrrolyl, furanyl, thiophenyl (or thienyl), imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, furazanyl, oxadiazolyl, thiadiazolyl, dithiazolyl, triazolyl, tetrazolyl, pyridinyl, pyranyl, thiopyranyl,
20 pyrazinyl, pyrimidinyl, pyridazinyl, oxazinyl, thiazinyl, dioxinyl, dithiinyl, oxathianyl, triazinyl, tetrazinyl, and the like. Bi-cyclic heteroaryls include groups in which a monocyclic heteroaryl ring is fused to one or more aryl or heteroaryl rings. Non-limiting examples include indolyl, indazolyl, benzofuranyl, benzimidazolyl, and imidazo[1,2-a]pyridine.

The term “carbocyclic ring” or “carbocyclyl” refers to a 4- to 12-membered saturated
25 or partially unsaturated hydrocarbon ring and may exist as a single ring, bicyclic ring (including fused, spiro or bridged carbocyclic rings) or a spiro ring. Bi-cyclic carbocyclyl groups include, *e.g.*, unsaturated carbocyclic radicals fused to another unsaturated carbocyclic radical, cycloalkyl, or aryl, such as, for example, cyclohexyl, cyclohexenyl, 2,3-dihydroindenyl, decahydronaphthalenyl, and 1,2,3,4-tetrahydronaphthalenyl. Unless specified
30 otherwise, the carbocyclic ring generally contains 4- to 10- ring members.

The term “C₃₋₆ cycloalkyl” refers to a carbocyclic ring which is fully saturated (*e.g.*, cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl).

The term “heterocycle” or “heterocyclyl” refers to a 4- to 12-membered saturated or partially unsaturated heterocyclic ring containing 1 to 4 heteroatoms independently selected

from N, O, and S. A heterocyclyl group may be mono- or bicyclic (e.g., a bridged, fused, or spiro bicyclic ring). Examples of monocyclic saturated or partially unsaturated heterocyclic radicals include, without limitation, piperdinyl, piperazinyl, tetrahydropyranyl, morpholinyl, and pyrrolidinyl. Bi-cyclic heterocyclyl groups include, *e.g.*, unsaturated heterocyclic radicals fused to another unsaturated heterocyclic radical, cycloalkyl, aryl, or heteroaryl ring, such as, for example, 2-oxo-2,3-dihydro-1*H*-benzo[*d*]imidazolyl, 1,4,5,6-tetrahydrocyclopenta[*c*]pyrazolyl, 4,5,6,7-tetrahydrothieno[2,3-*c*]pyridinyl, 5,6-dihydro-4*H*-cyclopenta[*b*]thiophenyl, and 4,7-dihydro-5*H*-thieno[2,3-*c*]pyranyl. In some embodiments, the heterocyclyl group is a 4 to 6 membered monocyclic heterocyclyl group. In some

5

10

embodiments, the heterocyclyl group is a 8 to 10 membered bicyclic heterocyclyl group.

As used herein the term “spiro” ring means a two-ring system wherein both rings share one common atom. Examples of spiro rings include 5-oxaspiro[2.3]hexane, oxaspiro[2.4]heptanyl, 5-oxaspiro[2.4]heptanyl, 4-oxaspiro[2.4]heptane, 4-oxaspiro[2.5]octanyl, 6-oxaspiro[2.5]octanyl, oxaspiro[2.5]octanyl, oxaspiro[3.4]octanyl, oxaspiro[bicyclo[2.1.1]hexane-2,3'-oxetan]-1-yl, oxaspiro[bicyclo[3.2.0]heptane-6,1'-cyclobutan]-7-yl, 2,6-diazaspiro[3.3]heptanyl, -oxa-6-azaspiro[3.3]heptane, 2,2,6-diazaspiro[3.3]heptane, 3-azaspiro[5.5]undecanyl, 3,9-diazaspiro[5.5]undecanyl, 7-azaspiro[3.5]nonane, 2,6-diazaspiro[3.4]octane, 8-azaspiro[4.5]decane, 1,6-diazaspiro[3.3]heptane, 5-azaspiro[2.5]octane, 4,7-diazaspiro[2.5]octane, 5-oxa-2-azaspiro[3.4]octane, 6-oxa-1-azaspiro[3.3]heptane, 3-azaspiro[5.5]undecanyl, 3,9-diazaspiro[5.5]undecanyl, and the like.

15

20

The term “fused” ring refers to two ring systems share two adjacent ring atoms. Fused heterocycles have at least one the ring systems contain a ring atom that is a heteroatom selected from O, N and S (e.g., 3-oxabicyclo[3.1.0]hexane).

As used herein the term “bridged” refers to a 5 to 10 membered cyclic moiety connected at two non-adjacent ring atoms (e.g. bicyclo[1.1.1]pentane, bicyclo [2.2.1] heptane and bicyclo [3.2.1] octane).

25

The phrase “pharmaceutically acceptable” indicates that the substance, composition or dosage form must be compatible chemically and/or toxicologically, with the other ingredients comprising a formulation, and/or the mammal being treated therewith.

30

Unless specified otherwise, the term “compounds of the present disclosure” refers to compounds of Formula (I), as well as all stereoisomers (including diastereoisomers and enantiomers), rotamers, tautomers, isotopically labeled compounds (including deuterium

substitutions). When a moiety is present that is capable of forming a salt, then salts are included as well, in particular pharmaceutically acceptable salts.

As used herein, the term “a,” “an,” “the” and similar terms used in the context of the present disclosure (especially in the context of the claims) are to be construed to cover both
5 the singular and plural unless otherwise indicated herein or clearly contradicted by the context. The use of any and all examples, or exemplary language (e.g. “such as”) provided herein is intended merely to better illuminate the disclosure and does not pose a limitation on the scope of the disclosure otherwise claimed.

It is also possible that the intermediates and compounds of the present disclosure may
10 exist in different tautomeric forms, and all such forms are embraced within the scope of the disclosure. The term “tautomer” or “tautomeric form” refers to structural isomers of different energies which are interconvertible via a low energy barrier. For example, proton tautomers (also known as prototropic tautomers) include interconversions via migration of a proton, such as keto-enol and imine-enamine isomerizations. A specific example of a proton
15 tautomer is the imidazole moiety where the proton may migrate between the two ring nitrogens. Valence tautomers include interconversions by reorganization of some of the bonding electrons.

In one embodiment, the present disclosure relates to a compound of the Formula (I) as defined herein, in free form. In another embodiment, the present disclosure relates to a
20 compound of the Formula (I) as defined herein, in salt form. In another embodiment, the present disclosure relates to a compound of the Formula (I) as defined herein, in acid addition salt form. In a further embodiment, the present disclosure relates to a compound of the Formula (I) as defined herein, in pharmaceutically acceptable salt form. In yet a further embodiment, the present disclosure relates to a compound of the Formula (I) as defined
25 herein, in pharmaceutically acceptable acid addition salt form. In yet a further embodiment, the present disclosure relates to any one of the compounds of the Examples in free form. In yet a further embodiment, the present disclosure relates to any one of the compounds of the Examples in salt form. In yet a further embodiment, the present disclosure relates to any one of the compounds of the Examples in acid addition salt form. In yet a further embodiment,
30 the present disclosure relates to any one of the compounds of the Examples in pharmaceutically acceptable salt form. In still another embodiment, the present disclosure relates to any one of the compounds of the Examples in pharmaceutically acceptable acid addition salt form.

Compounds of the present disclosure may be synthesized by synthetic routes that include processes analogous to those well-known in the chemical arts, particularly in light of the description contained herein. The starting materials are generally available from commercial sources such as Sigma-Aldrich or are readily prepared using methods well known to those skilled in the art (e.g., prepared by methods generally described in Louis F. Fieser and Mary Fieser, *Reagents for Organic Synthesis*, v. 1-19, Wiley, New York (1967-1999 ed.), or *Beilsteins Handbuch der organischen Chemie*, 4, Aufl. ed. Springer-Verlag, Berlin, including supplements (also available via the Beilstein online database)).

For illustrative purposes, the reaction schemes depicted below provide potential routes for synthesizing the compounds of the present disclosure as well as key intermediates. For a more detailed description of the individual reaction steps, see the Examples section below. Although specific starting materials and reagents are depicted in the schemes and discussed below, other starting materials and reagents can be easily substituted to provide a variety of derivatives and/or reaction conditions.

EXEMPLIFICATION

Abbreviation:

PE = petroleum ether

EtOAc = EA = ethyl acetate

ESI = electrospray ionisation

MeOH = methanol

EtOH = ethanol

DCM = dichloromethane

DIAD = Diisopropyl azodicarboxylate

DMF = dimethylformamide

DPPA = Diphenylphosphoryl azide

EDCI = 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide

HATU = Hexafluorophosphate Azabenzotriazole Tetramethyl Uronium

HCl = hydrochloric acid

H₂O = water

NBS = N-bromosuccinimide

LCMS = liquid chromatography mass spectrometry

HPLC = high pressure liquid chromatography

THF = tetrahydrofuran

- MeCN = ACN = acetonitrile
DMSO = dimethylsulfoxide
PhMe = Toluene
AcOH = acetic acid
- 5 CDI = carbonyldiimidazole
TFA = trifluoroacetic acid
DIPEA = DIEA = diisopropylethyl amine
TLC = Thin Layer Chromatography
N₂ = Nitrogen
- 10 KOH = Potassium Hydroxide
NH₄HCO₃ = Ammonium Bicarbonate
t-BuOH = tert-butanol
NH₄Cl = ammonium chloride
t-BuOK = potassium tert-butoxide
- 15 NaH = sodium hydride
Xantphos = 4,5-Bis(diphenylphosphino)-9,9-dimethylxanthene
Na₂SO₄ = sodium sulfate
K₂CO₃ = potassium carbonate
K₃PO₄ = potassium phosphate
- 20 Cs₂CO₃ = cesium carbonate
NaHCO₃ = sodium bicarbonate
Pd(OAc)₂ = Palladium (II) Acetate
Pd₂(dba)₃ = tris(dibenzylideneacetone)dipalladium(0)
NaOH = Sodium Hydroxide
- 25 TCFH = N'-tetramethylformamidinium hexafluorophosphate
NMI = 1-Methylimidazole
Xantphos = 4,5-Bis(diphenylphosphino)-9,9-dimethylxanthene
XantPhosPdG₃ = [(4,5-Bis(diphenylphosphino)-9,9-dimethylxanthene)-2-(2'-amino-1,1'-biphenyl)]palladium(II) methanesulfonate
- 30 RuPhos = 2-Dicyclohexylphosphino-2',6'-diisopropoxybiphenyl
RuPhos Pd G₃ = (2-Dicyclohexylphosphino-2',6'-diisopropoxy-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) methanesulfonate
Pd/C = Palladium on Carbon

Pd(dppf)Cl₂ DCM = [1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium(II),
complex with dichloromethane

Pd(dppf)Cl₂ = [1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium(II)

Pd(PPh₃)₄ = Tetrakis(triphenylphosphine)palladium(0)

5 Pd(PPh₃)Cl₂ = Bis(triphenylphosphine)palladium(II) dichloride

STAB = NaBH(AcO)₃ = sodium triacetoxymethylborohydride

AlMe₃ = Al(CH₃)₃ = trimethylaluminum

DIAD = diisopropyl azodicarboxylate

10 PEPSI-IHeptCl = (SP-4-1)-[1,3-Bis[2,6-bis(1-propylbutyl)phenyl]-4,5-dichloro-1,3-dihydro-2H-imidazol-2-ylidene]dichloro(3-chloropyridine-κN)-Palladium

Er(OTf)₃ = erbium(III) triflate

Br₂ = bromine

KOAc = potassium acetate

TEBAc = benzyltriethylammonium chloride

15 B₂pin₂ = bis(pinacolato)diboron

BINAP = 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl

BPO = benzoyl peroxide

Py = pyridine

LiAlH₄ = lithium aluminum hydride

20

GENERAL METHODS

1. ¹H NMR spectra were recorded on:

NMR10 Bruker AVANCE III HD 300MHz

NMR16 Bruker AVANCE III HD 300MHz

25 NMR19 Bruker AVANCE III HD 400MHz

NMR24 Bruker AVANCE NEO 400MHz

NMR30 Bruker AVANCE NEO 400MHz

30 2. LCMS measurement was run on SHIMADZU LCMS-2020 using the following conditions:

Method A: Mobile Phase: A: Water (0.05%TFA) B: Acetonitrile (0.05%TFA);

Gradient Phase: 5%B to 100%B within 2.0 min, 100%B with 0.7 min (total runtime:

2.8 min); Flow Rate: 1.5 mL/min; Column: HALO C18, 3.0*30mm, 2.0 μ m; Column Temperature: 40 °C. Detectors: AD2 ELSD, PDA (220 nm and 254 nm), ESI.


Method B: Mobile Phase: A: Water (0.1%FA) B: Acetonitrile (0.1%FA); Gradient Phase: 5%B to 100%B within 2.0 min, 100%B with 0.7 min (total runtime: 2.8 min); Flow Rate: 1.5 mL/min; Column: HALO C18, 3.0*30mm, 2.0 μ m; Column Temperature: 40 °C. Detectors: AD2 ELSD, PDA (220 nm and 254 nm), ESI.

Method C: Mobile Phase: A: Water (5mM NH₄HCO₃) B: Acetonitrile; Gradient Phase: 10%B to 95%B within 2.0 min, 100%B with 0.6 min (total runtime: 2.8 min); Flow Rate: 1.5 mL/min; Column: Poroshell HPH-C18, 3.0*50mm, 4.0 μ m; Column Temperature: 40 °C. Detectors: AD2 ELSD, PDA (220 nm and 254 nm), ESI.

PREPARATION OF INTERMEDIATES

The compounds claimed herein were prepared following the procedures outlined in the following schemes. Compound names were generated using the software built into ChemDraw. To the extent that there are discrepancies between the name of a compound and its depicted structure, the depicted chemical structure is to be taken as the appropriate compound.

Intermediate A: *N*-(3-amino-5-chlorophenyl)methanesulfonamide




Step 1: To a stirred mixture of 1-chloro-3-fluoro-5-nitrobenzene (45 g, 256.35 mmol, 1.00 equiv) and methane sulfonamide (24.38 g, 256.35 mmol, 1.00 equiv), Cs₂CO₃ (250.57 g, 769.05 mmol, 3.00 equiv) in 500 mL of DMSO, the resulted solution was stirred for 2h at 60 °C. The mixture was cooled then quenched with 1000 mL of water and extracted with 3 x 1000 mL of ethyl acetate and dried over anhydrous Na₂SO₄ and concentrated and the residue was purified onto silica gel column eluted with 50% of ethyl acetate in petroleum ether to afford *N*-(3-chloro-5-nitrophenyl) methane sulfonamide (41 g, 63.81%) as a light yellow solid. LCMS (ESI) [M + H]⁺: 249.9

Step 2: To a stirred solution of *N*-(3-chloro-5-nitrophenyl) methanesulfonamide (41.00 g, 163.57 mmol, 1.00 equiv) and Fe (91.35 g, 1635.75 mmol, 10.00 equiv), NH₄Cl (87.50 g,

1635.75 mmol, 10.00 equiv) in 500 mL of ethanol and 300 mL of water, this was stirred for 2h at 90 °C. The resulting mixture was filtered, the filter cake was washed with 6 x 50 mL of methanol. The filtrate was concentrated under reduced pressure. The residue was purified onto silica gel column eluted with 50% of ethyl acetate in petroleum ether to afford N-(3-amino-5-chlorophenyl) methanesulfonamide (25.3 g, 70.1%) as a light yellow solid.

LCMS (ESI) $[M + H]^+$: 220.01. ^1H NMR (300 MHz, DMSO- d_6) δ 9.64 (s, 1H), 6.41 – 6.31 (m, 3H), 5.54 (s, 2H), 2.98 (s, 3H)


Intermediate E: N-(3-amino-5-chlorophenyl)propane-1-sulfonamide



- 10 A solution of 3-bromo-5-chloroaniline (1 g, 4.843 mmol, 1 equiv), propane-1-sulfonamide (1.19 g, 9.686 mmol, 2 equiv), (1S,2S)-N¹,N²-dimethylcyclohexane-1,2-diamine (2.07 g, 14.529 mmol, 3 equiv), CuI (2.77 g, 14.529 mmol, 3 equiv) and K₂CO₃ (2.01 g, 14.529 mmol, 3 equiv) in ACN (10 mL) was stirred for 2 hours at 110°C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (1:1) to afford N-(3-amino-5-chlorophenyl)propane-1-sulfonamide (540 mg, 45% yield) as a yellow solid.

LCMS (ESI) $[M + H]^+$: 249

Intermediate H: N-(3-amino-5-chlorophenyl)cyclopropanesulfonamide









- 20 Step 1: A solution of 3-chloro-5-nitroaniline (300 mg, 1.738 mmol, 1 equiv) and cyclopropanesulfonyl chloride (244 mg, 1.738 mmol, 1 equiv) in pyridine (5 ml) was stirred for 2 hours at room temperature. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (5:1) to afford N-(3-chloro-5-nitrophenyl)cyclopropanesulfonamide (255 mg, 53% yield) as a white solid.
- 25 LCMS (ESI) $[M+H]^+$: 277

Step 2: A solution of N-(3-chloro-5-nitrophenyl)cyclopropanesulfonamide (235 mg, 0.849 mmol, 1 equiv), Fe (474 mg, 8.490 mmol, 10 equiv) and NH₄Cl (136 mg, 2.547 mmol, 3 equiv) in ethanol (5 mL) and H₂O (1 mL) was stirred for 2 hours at 80 °C under nitrogen


- atmosphere. The mixture was allowed to cool down to room temperature. The resulting mixture was filtered, the filter cake was washed with ethanol (3x10 mL). The filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (1:1) to afford *N*-(3-amino-5-chlorophenyl)cyclopropanesulfonamide (190 mg, 91% yield) as a white solid.

LCMS (ESI) [M+H]⁺: 247

The following intermediates were prepared using experimental procedures similar for intermediates **A** **E** or **H**:


Structure	Intermediate	MS (ESI) [M + H] ⁺	¹ H NMR
	B	205	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.63 (s, 1H), 6.26 (t, J = 1.9 Hz, 1H), 6.12 (dt, J = 10.8, 2.1 Hz, 1H), 6.04 (dt, J = 11.5, 2.1 Hz, 1H), 5.52 (s, 2H), 2.98 (s, 3H).
	C	265	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.63 (s, 1H), 6.58 – 6.36 (m, 3H), 5.53 (s, 2H), 2.98 (s, 3H)
	D	313	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.39 (s, 1H), 6.66 (q, J = 1.6 Hz, 2H), 6.46 (t, J = 1.6 Hz, 1 H), 5.42 (s, 2H) 2.96 (s, 3H)
	F	249	
	G	220	
	I	251	

Intermediate J: 2-bromo-5-(tert-butoxy)pyrimidine



A mixture of 2-bromopyrimidin-5-ol (2.5 g, 14.3 mmol, 1 equiv) and Er(OTf)₃ (0.44 g, 0.7 mmol, 0.05 equiv) in Boc₂O (60 mL) was stirred for overnight at 50 °C. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (5:1) to afford 2-bromo-5-(tert-butoxy)pyrimidine (400 mg, 6% yield) as a yellow oil. This reaction was repeated 13 times and to afford 2.9 g of the desired product in total. LCMS (ESI) [M + H]⁺: 231

10 *Intermediate K: 5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxylate*




Step 1: A mixture of 5-methylthiophene-2-carboxylic acid (100 g, 703.383 mmol, 1 equiv) and Br₂ (123.65 g, 773.721 mmol, 1.1 equiv) in AcOH (1 L) was stirred for 2 hours at 60°C. The product was precipitated by the addition of water (500 mL) at 0°C. The precipitated solids were collected by filtration and washed with water to afford 4-bromo-5-methylthiophene-2-carboxylic acid (130 g, 84%) as a yellow solid. LCMS (ESI) [M+H]⁺: 221

Step 2: A mixture of 4-bromo-5-methylthiophene-2-carboxylic acid (100 g, 452.345 mmol, 1 equiv) and H₂SO₄ (100 mL, 1876.211 mmol, 4.15 equiv) in MeOH (1 L) was stirred for overnight at 50°C. The resulting mixture was concentrated under vacuum. The resulting mixture was diluted with water (200 mL). The aqueous layer was extracted with EtOAc (3x300 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column

chromatography, eluted with PE / EA (10:1) to afford methyl 4-bromo-5-methylthiophene-2-carboxylate (80 g, 74% yield) as a light yellow solid. GCMS: 234. ^1H NMR (300 MHz, DMSO- d_6) δ 7.70 (s, 1H), 3.82 (s, 3H), 2.43 (s, 3H).

Step 3: A mixture of methyl 4-bromo-5-methylthiophene-2-carboxylate (80 g, 340.281 mmol, 1 equiv), bis(pinacolato)diboron (172.82 g, 680.562 mmol, 2 equiv) Pd(dppf)Cl₂ (12.45 g, 17.014 mmol, 0.05 equiv) and KOAc (100.19 g, 1020.843 mmol, 3 equiv) in 1,4-dioxane (800 mL) was stirred for overnight at 80°C under nitrogen atmosphere. The resulting mixture was filtered and the filter cake was washed with EtOAc. The filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (15:1) to afford methyl 5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxylate (90 g, crude). The crude product was re-crystallized from hexane to afford methyl 5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxylate (35 g, 36% yield) as a white solid. LCMS (ESI) [M+H]⁺: 283. ^1H NMR (400 MHz, DMSO- d_6) δ 7.70 (s, 1H), 3.79 (s, 3H), 2.65 (s, 3H), 1.28 (s, 12H)


Intermediate L: methyl 5-methyl-1H-pyrrole-3-carboxylate



Step 1: Tosyl methylisocyanide (200g, 1.03 mol, 1 equiv) was dissolved in dichloromethane 1.5 L and cooled to 0 °C. Benzyltriethylammonium chloride (46.62 g, 2.05 mol, 2 equiv.), methyl iodide 126.3mL (2.05 mol, 2 equiv) and 1.5 L of 30% aqueous sodium hydroxide solution were added. The mixture was stirred for 3 hours at 0°C. Water (4 L) was added to the reaction mixture, extracted with dichloromethane (1.5 L × 3). The organic layer was dried over sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (4:1) to afford 1-(1-isocianoethylsulfonyl)-4-methylbenzene (167g, 78.0% yield) as a brown oil. Step 2: A solution of methyl acrylate (138.2 mL, 1.55 mol, 0.97 equiv) and 1-(1-isocianoethylsulfonyl)-4-methylbenzene (335 g, 1.60 mol, 1 equiv) in diethyl ether (3 L) and DMSO (1.5L) were added dropwise to a stirred suspension of sodium hydride (60% purity) (99.7 g, 2.49 mol, 1.55 equiv) in diethyl ether (4L) under nitrogen. The resulting suspension was stirred at room temperature for 4 hours. The reaction mixture was poured into sodium

chloride solution (5 L), layers separated and the aqueous extracted with EA (3 x 5 L), the organic layers were combined and washed with water (5 L), dried over Na₂SO₄, filtered and evaporated to afford beige solid. The crude product was purified by flash silica chromatography, elution gradient 0 to 40% ethyl acetate to afford methyl 5-methyl-1H-pyrrole-3-carboxylate (105.6g, 47.4% yield) as a white solid. LCMS (ESI) [M+H]⁺: 140. ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.18 (s, 1H), 7.24 (dd, *J* = 3.0, 1.7 Hz, 1H), 6.10 (dt, *J* = 2.7, 1.4 Hz, 1H), 3.66 (s, 3H), 2.15 (d, *J* = 1.0 Hz, 3H).

Intermediate M: 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrole-3-carboxylate



Step 1: To a stirred solution of methyl 1H-pyrrole-3-carboxylate (100 g, 800 mmol, 1 equiv) and NaH (60%) (64 g, 1.6 mol, 2 equiv) in DMF (1 L) was added CH₃I (227.2 g, 1.6 mol, 2 equiv) dropwise at 0 °C. The resulting mixture was stirred for additional 3 hours at room temperature. The reaction was quenched by the addition of water (5 L) at 0°C. The resulting mixture was extracted with EtOAc (3 x 5 L). The combined organic layers were washed with brine (3x5L), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (5:1) to afford methyl 1-methylpyrrole-3-carboxylate (95g, 85.6%) as a white solid. LCMS (ESI) [M+H]⁺: 140

Step 2: A solution of methyl 1-methylpyrrole-3-carboxylate (95 g, 683.5 mmol, 1 equiv), B₂Pin₂ (138.9 g, 546.7 mmol, 0.8 equiv), Di-μ-methoxybis(1,5-cyclooctadiene)diiridium(I) (6.7g, 10.3 mmol, 0.015 equiv) and 4,4'-Di-tert-butyl-2,2'-bipyridine (5.49 g, 20.5 mmol, 0.03 equiv) in 1,4-dioxane (1L) was stirred for 3 hours at 100°C under nitrogen atmosphere. The mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (2/1) to afford crude product. The crude product was triturated with PE for 12 hours at RT and filtered to afford 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrole-3-carboxylate (105 g, 57.8% yield) as a white solid. LCMS (ESI) [M+H]⁺: 266. ¹H NMR (300 MHz, Chloroform-*d*) δ 7.37 (d, *J* = 1.6 Hz, 1H), 7.21 (d, *J* = 1.6 Hz, 1H), 3.82 (s, 3H), 3.78 (s, 3H), 1.31 (s, 12H).


The following intermediates were prepared using experimental procedures similar for intermediate M:

Structure	Intermediate	MS (ESI) [M + H] ⁺	¹ H NMR
	N	252	
	O	316	
	P	334	¹ H NMR (400 MHz, DMSO-d ₆) δ 7.80 (d, 1H), 6.99 (d, 1H), 5.08 (m, 2H), 3.72 (s, 3H), 1.26 (s, 12H).

5 GENERAL SCHEMES


Compounds described below can be prepared through, for example, the following general schemes, among other methods known to those skilled in the art:

General Scheme A:



A carboxylic acid ester, either commercially available or prepared through various means (e.g. Suzuki coupling, Buchwald coupling, alkylation, S_NAr, ring condensation, or multistep syntheses) is hydrolyzed with various reagents (e.g. lithium hydroxide) to yield a carboxylic acid. The carboxylic acid is then coupled with an appropriate reagent (e.g. one of the above-described anilines or other amine) using various reagents to provide the desired product.

General Scheme B:




A carboxylic acid ester, either commercially available or prepared through various means

- 5 (e.g. Suzuki coupling, Buchwald coupling, alkylation, S_NAr , ring condensation, or multistep syntheses) is reacted with an appropriate reagent (e.g. one of the above-described anilines or other amine) in the presence of trimethylaluminum or other appropriate reagent to provide the desired product.

- 10 In general, either method is suitable for obtaining the examples shown below and can be considered interchangeable. For the examples shown below, one skilled in the art can identify alternative conditions for individual steps, such as alkylations, hydrolysis, displacements, cross-coupling or other chemical transformations.

15 EXAMPLES

Example 1: N-(3-chloro-5-(methanesulfonamido)phenyl)-1-phenyl-1H-imidazole-4-carboxamide












- 20 To a mixture of 1-phenyl-1H-imidazole-4-carboxylic acid (70 mg, 0.371 mmol) and N-(3-amino-5-chlorophenyl)methanesulfonamide (81.8 mg, 0.371 mmol) in ACN (1 mL) was added TCFH (103 mg, 0.371 mmol) and NMI (30.4 mg, 0.371 mmol), the resulting mixture was stirred for two hours at room temperature. Then it was concentrated, the residue was purified by reverse phase flash chromatography eluting with 60% of acetonitrile in water (0.1% NH_4HCO_3) to afford N-(3-chloro-5-methanesulfonamidophenyl)-1-phenyl-1H-





imidazole-4-carboxamide (47.2 mg, 32.4%) as a white solid. LCMS $[M + H]^+$: 391. ^1H NMR (400 MHz, DMSO- d_6) δ 10.27 (s, 1H), 10.05 (s, 1H), 8.53 – 8.44 (m, 2H), 7.83 – 7.74 (m, 4H), 7.57 (dd, J = 8.6, 7.2 Hz, 2H), 7.48 – 7.39 (m, 1H), 6.92 (t, J = 2.0 Hz, 1H), 3.10 (s, 3H).





Example 2-14 and 321-332






- 5 The compounds listed in the following table were prepared using a procedure similar to that described for example 1:




Structure	Example No.	MS (ESI) $[M + H]^+$	^1H NMR
	2	376	^1H NMR (300 MHz, DMSO- d_6) δ 12.10 (s, 1H), 10.02 (s, 1H), 9.78 (s, 1H), 7.56 (s, 1H), 7.52 – 7.39 (m, 2H), 6.73 (s, 1H), 6.67 (dt, J = 10.4, 2.2 Hz, 1H), 3.06 (s, 3H).
	3	313	^1H NMR (300 MHz, DMSO- d_6) δ 13.12 (s, 1H), 10.24 (s, 1H), 10.04 (s, 1H), 7.61 (s, 1H), 7.48 (dt, J = 11.4, 2.1 Hz, 1H), 6.69 (dt, J = 10.5, 2.2 Hz, 1H), 6.53 (s, 1H), 3.09 (s, 3H), 2.30 (s, 3H).
	4	377	^1H NMR (300 MHz, DMSO- d_6) δ 10.45 (s, 1H), 7.45 (d, J = 11.1 Hz, 2H), 7.17 (s, 1H), 6.75 (d, J = 10.6 Hz, 1H), 3.08 (s, 3H).
	5	312	^1H NMR (300 MHz, DMSO- d_6) δ 11.11 (s, 1H), 9.99 (s, 1H), 9.62 (s, 1H), 7.58 – 7.29 (m, 3H), 6.64 (dt, J = 10.5, 2.2 Hz, 1H), 6.33 (dt, J = 2.6, 1.3 Hz, 1H), 3.06 (s, 3H), 2.19 (s, 3H).

	6	391	¹ H NMR (300 MHz, DMSO-d ₆) δ 13.19 (s, 1H), 10.03 (s, 1H), 8.09 (d, J = 7.6 Hz, 2H), 7.99 (s, 1H), 7.77 (s, 2H), 7.52 (t, J = 7.5 Hz, 2H), 7.43 (t, J = 7.2 Hz, 1H), 6.92 (t, J = 2.0 Hz, 1H), 3.10 (s, 3H).
	7	391	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.49 (s, 1H), 10.08 (s, 1H), 8.49 (d, J = 1.0 Hz, 1H), 7.84 (d, J = 1.0 Hz, 1H), 7.74 – 7.65 (m, 4H), 7.44 (t, J = 7.7 Hz, 2H), 7.37 – 7.29 (m, 1H), 6.96 (t, J = 2.0 Hz, 1H), 3.08 (s, 3H).
	8	390	¹ H NMR (300 MHz, DMSO-d ₆) δ 11.93 (s, 1H), 10.06 (d, J = 3.7 Hz, 2H), 7.70 (dt, J = 13.2, 1.9 Hz, 2H), 7.60 (dt, J = 8.2, 1.7 Hz, 2H), 7.49 (q, J = 2.2, 1.7 Hz, 2H), 7.37 (t, J = 7.7 Hz, 2H), 7.24 – 7.12 (m, 1H), 6.92 (t, J = 1.9 Hz, 1H), 3.09 (s, 3H).
	9	391	¹ H NMR (300 MHz, DMSO-d ₆) δ 13.56 (d, J = 74.4 Hz, 1H), 10.60 (d, J = 58.3 Hz, 1H), 10.10 (s, 1H), 7.99 – 7.87 (m, 3H), 7.74 (dd, J = 31.7, 18.6 Hz, 2H), 7.41 (t, J = 7.5 Hz, 2H), 7.27 (t, J = 7.4 Hz, 1H), 6.95 (s, 1H), 3.11 (s, 3H).
	10	449	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.68 (s, 1H), 9.64 (d, J = 0.8 Hz, 1H), 8.27 (dd, J = 8.2, 7.2 Hz, 1H), 8.26 – 8.17 (m, 1H), 8.17 – 8.09 (m, 2H), 7.82 – 7.69 (m, 4H), 3.23 (s, 3H), 3.07 – 3.00 (m, 3H).


	11	421	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.33 (s, 1H), 10.07 (s, 1H), 8.09 (s, 1H), 7.67 (t, J = 1.9 Hz, 1H), 7.61 (t, J = 1.9 Hz, 1H), 7.54 – 7.45 (m, 4H), 7.40 (dq, J = 8.8, 4.4 Hz, 1H), 6.95 (t, J = 2.0 Hz, 1H), 3.06 (s, 3H), 2.53 (s, 3H).
	12	423	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.91 (s, 1H), 9.74 (s, 1H), 7.35 (t, J = 1.8 Hz, 1H), 7.26 – 7.18 (m, 4H), 7.18 – 7.13 (m, 1H), 7.08 (ddd, J = 8.5, 5.3, 2.4 Hz, 1H), 6.92 (t, J = 1.9 Hz, 1H), 3.98 (q, J = 8.7 Hz, 1H), 3.62 – 3.54 (m, 1H), 2.98 (s, 3H), 2.49 – 2.43 (m, 1H), 2.33 – 2.21 (m, 2H), 2.19 – 2.09 (m, 1H).
	13	426	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.56 (s, 1H), 10.10 (s, 1H), 8.30 (dt, J = 4.1, 1.8 Hz, 2H), 8.15 (dt, J = 8.1, 1.4 Hz, 1H), 8.06 – 7.94 (m, 2H), 7.90 (dt, J = 7.7, 1.3 Hz, 1H), 7.81 – 7.63 (m, 4H), 6.99 (t, J = 1.9 Hz, 1H), 3.09 (s, 3H).
	14	402	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.60 (s, 1H), 10.10 (s, 1H), 8.79 – 8.63 (m, 2H), 8.33 (t, J = 1.8 Hz, 1H), 8.05 (ddt, J = 8.2, 3.6, 1.5 Hz, 2H), 7.89 – 7.79 (m, 2H), 7.72 (q, J = 3.4, 2.9 Hz, 3H), 6.99 (t, J = 1.9 Hz, 1H), 3.09 (s, 3H).

	321	411.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.71 (d, J = 11.3 Hz, 1H), 10.11 (d, J = 11.3 Hz, 1H), 8.54 (d, J = 11.5 Hz, 1H), 7.78 – 7.66 (m, 2H), 7.62 (dd, J = 11.5, 8.0 Hz, 1H), 7.48 (dt, J = 11.6, 8.0 Hz, 1H), 7.04 – 6.94 (m, 2H), 4.00 (d, J = 11.6 Hz, 3H), 3.10 (d, J = 11.6 Hz, 3H)
	322	393.2	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.24 (s, 1H), 10.05 (s, 1H), 8.19 (d, J = 1.6 Hz, 1H), 8.02 (s, 1H), 7.73 (d, J = 1.9 Hz, 1H), 7.66 (d, J = 2.0 Hz, 1H), 7.39 (s, 1H), 6.93 (d, J = 2.0 Hz, 1H), 3.07 (s, 3H), 2.59 (s, 3H), 2.29 (d, J = 1.1 Hz, 3H).
	323	446.9 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.89 (d, J = 3.3 Hz, 1H), 10.10 (d, J = 3.0 Hz, 1H), 8.62 – 8.56 (m, 1H), 8.44 (dd, J = 8.3, 3.6 Hz, 1H), 7.89 (dd, J = 7.5, 3.6 Hz, 1H), 7.68 (tq, J = 5.8, 3.9, 3.0 Hz, 3H), 7.01 (dt, J = 3.8, 1.9 Hz, 1H), 3.08 (d, J = 3.2 Hz, 3H)
	324	414.9 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.84 (s, 1H), 10.17 (s, 1H), 8.58 (d, J = 3.3 Hz, 1H), 7.71 (t, J = 1.8 Hz, 1H), 7.65 (t, J = 1.9 Hz, 1H), 7.46 (td, J = 9.0, 3.6 Hz, 1H), 7.37 (td, J = 9.1, 3.5 Hz, 1H), 7.00 (t, J = 1.9 Hz, 1H), 3.09 (s, 3H)

	325	414.9 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.84 (s, 1H), 10.13 (s, 1H), 8.58 (d, J = 3.4 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.65 (t, J = 1.9 Hz, 1H), 7.46 (td, J = 9.0, 3.7 Hz, 1H), 7.37 (td, J = 9.1, 3.6 Hz, 1H), 7.00 (t, J = 2.0 Hz, 1H), 3.08 (s, 3H)
	326	446.9 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.89 (s, 1H), 10.11 (s, 1H), 8.48 (d, J = 10.0 Hz, 2H), 8.34 (d, J = 8.6 Hz, 1H), 7.80 (dd, J = 8.6, 1.9 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.67 (t, J = 2.0 Hz, 1H), 7.00 (t, J = 2.0 Hz, 1H), 3.09 (s, 3H)
	327	408.9 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.61 (s, 1H), 10.08 (s, 1H), 8.30 (s, 1H), 7.90 (d, J = 8.8 Hz, 1H), 7.70 (d, J = 1.7 Hz, 1H), 7.68 – 7.60 (m, 2H), 7.09 (dd, J = 8.9, 2.4 Hz, 1H), 6.96 (q, J = 1.7 Hz, 1H), 3.86 (s, 3H), 3.08 (s, 3H)
	328	391.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.09 (s, 1H), 9.97 (s, 1H), 8.06 (d, J = 1.8 Hz, 1H), 8.01 – 7.96 (m, 1H), 7.75 (t, J = 1.8 Hz, 1H), 7.69 (q, J = 1.6 Hz, 1H), 6.94 (dt, J = 12.8, 1.6 Hz, 2H), 6.48 (t, J = 1.4 Hz, 1H), 3.07 (s, 3H), 2.35 (s, 3H), 2.17 (d, J = 1.2 Hz, 3H)
	329	378	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.14 (s, 1H), 10.03 (s, 1H), 8.18 (dd, J = 14.1, 3.3 Hz, 2H), 7.75 (d, J = 2.0 Hz, 1H), 7.69 (t, J = 2.0 Hz, 1H), 7.00 (s,

			1H), 6.92 (t, J = 2.0 Hz, 1H), 6.63 – 6.55 (m, 2H), 3.07 (s, 3H), 2.38 (s, 3H)
	330	364.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.33 (s, 1H), 10.06 (s, 1H), 8.96 (d, J = 1.5 Hz, 1H), 8.35 – 8.29 (m, 2H), 7.74 (t, J = 1.9 Hz, 1H), 7.67 (t, J = 1.9 Hz, 1H), 7.57 (d, J = 4.9 Hz, 1H), 7.43 (d, J = 1.3 Hz, 1H), 6.95 (t, J = 2.0 Hz, 1H), 3.07 (s, 3H).
	331	394.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.69 (s, 1H), 10.09 (s, 1H), 8.31 (d, J = 2.0 Hz, 1H), 7.94 (d, J = 8.2 Hz, 1H), 7.81 (s, 1H), 7.75 – 7.69 (m, 1H), 7.73 – 7.64 (m, 1H), 7.35 (d, J = 8.3 Hz, 1H), 6.97 (q, J = 2.1 Hz, 1H), 3.11 – 3.06 (m, 3H), 2.46 (d, J = 1.9 Hz, 3H)
	332	414.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.84 (s, 1H), 10.10 (s, 1H), 8.54 (d, J = 0.9 Hz, 1H), 8.08 (dd, J = 8.0, 1.0 Hz, 1H), 7.70 (dt, J = 9.2, 1.8 Hz, 2H), 7.59 (dd, J = 7.7, 1.0 Hz, 1H), 7.52 (t, J = 7.9 Hz, 1H), 7.00 (t, J = 1.9 Hz, 1H), 3.08 (s, 3H)

Example 15: N-(3-bromo-5-(methanesulfonamido)phenyl)-4-phenylthiophene-2-carboxamide



Step 1: To a mixture of methyl 4-bromothiophene-2-carboxylate (30 g, 135 mmol),
 5 phenylboronic acid (32.9 g, 270 mmol), Pd(dppf)Cl₂ (11.0 g, 13.5 mmol) and K₃PO₄ (85.8 g, 405 mmol) was added dioxane (300 mL) and H₂O (30 mL). The resulting mixture was stirred for 2 hours at 80 °C under N₂ atmosphere. The reaction mixture was extracted with ethyl acetate. The organic phase was concentrated under reduced pressure. The residue was purified by flash chromatography on silica gel eluting with 20% of ethyl acetate in petroleum ether to afford methyl 4-phenylthiophene-2-carboxylate (27.1 g, 124 mmol) as a light yellow solid.


Step 2: To a stirred solution of methyl 4-phenylthiophene-2-carboxylate (26.1 g, 119 mmol) in 260 mL of THF and 260 mL of water was added LiOH (24.9 g, 595 mmol). The resulting mixture was stirred overnight at room temperature. Then it was concentrated, and the pH
 15 value of the residue solution was adjusted to 3 with 1M HCl (aq). The product was precipitated from the solution. After filtration, obtained 4-phenylthiophene-2-carboxylic acid (24.4 g, 119 mmol) as a white solid.

Step 3: To a mixture of 4-phenylthiophene-2-carboxylic acid (10.1 g, 49.6 mmol) and N-(3-amino-5-bromophenyl) methanesulfonamide (11 g, 41.4 mmol) in ACN (110 mL) was added
 20 TCFH (17.3 g, 62.1 mmol) and NMI (10.1 g, 124 mmol) at room temperature. The resulting mixture was stirred overnight, Then it was concentrated, the residue was purified by reverse phase flash chromatography eluting with 60% of acetonitrile in water (0.1%FA) to afford N-(3-bromo-5-methanesulfonamidophenyl)-4-phenylthiophene-2-carboxamide (11.0519 g, 24.486 mmol) as a light yellow solid. LCMS (ESI) [M - H]⁻: 449. ¹H NMR (300 MHz,


DMSO-d₆) δ 10.46 (s, 1H), 10.09 (s, 1H), 8.49 (d, J = 1.5 Hz, 1H), 8.23 (d, J = 1.4 Hz, 1H), 7.96 – 7.60 (m, 4H), 7.57 – 7.24 (m, 3H), 7.11 (t, J = 1.9 Hz, 1H), 3.08 (s, 3H).

Examples 16

The compounds listed in the following table were prepared using a procedure similar to that described for example 15:

Structure	Example No.	MS (ESI) [M + H] ⁺	¹ H NMR
	16	499	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.39 (s, 1H), 10.00 (s, 1H), 8.49 (d, J = 1.5 Hz, 1H), 8.22 (d, J = 1.4 Hz, 1H), 7.96 (t, J = 1.7 Hz, 1H), 7.75 (dt, J = 5.8, 1.4 Hz, 3H), 7.49 (t, J = 7.5 Hz, 2H), 7.36 (t, J = 7.4 Hz, 1H), 7.28 (t, J = 1.8 Hz, 1H), 3.07 (s, 3H).


Example 17: N-(3-bromo-5-(methylsulfonamido)phenyl)-4-(pyridin-2-yl)thiophene-2-carboxamide



- 10 Step 1: A mixture of ethyl 4-bromothiophene-2-carboxylate (250 g, 1063.37 mmol, 1.00 equiv) and bis(pinacolato)diboron (810.10 g, 3190.13 mmol, 3.00 equiv) and Pd(dppf)Cl₂ (38.90 g, 53.17 mmol, 0.05 equiv) and AcOK (313.09 g, 3190.13 mmol, 3.00 equiv) in 1500 mL of 1,4-dioxane, the resulted solution was stirred for 2h at 80 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was
- 15 purified onto silica gel column eluted with 50% of ethyl acetate in petroleum ether to afford ethyl 4-(4,4,5-trimethyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxylate (231.02 g, 81.0%) as a yellow solid.

- Step 2: A mixture of methyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxylate (231.00 g, 861.52 mmol, 1.00 equiv) and 2-bromopyridine (163.34 g, 1033.83 mmol, 1.20 equiv) and Pd(dppf)Cl₂ (31.52 g, 43.076 mmol, 0.05 equiv) and K₂CO₃ (357.20 g, 2584.57 mmol, 3.00 equiv) in 1000 mL of 1,4-dioxane and 100 mL of water, the resulted solution was stirred for 2 h at 90 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified onto silica gel column eluted with 50% of ethyl acetate in petroleum ether to afford methyl 4-(pyridin-2-yl)thiophene-2-carboxylate (196.12 g, 103.8%) as a white solid. LCMS (ESI). [M + H]⁺: 219
- Step 3: A mixture of methyl 4-(pyridin-2-yl)thiophene-2-carboxylate (186.00 g, 848.31 mmol, 1.00 equiv) and LiOH (60.95 g, 2544.924 mmol, 3.00 equiv) in 1000 mL of ethanol and 100 mL of water, this was stirred for 1h at 60°C. The mixture was adjusted pH to 6 with HCl (aq.). The precipitated solids were collected by filtration and washed with 3 x 100 mL of water. This resulted in 4-(pyridin-2-yl)thiophene-2-carboxylic acid (133.12 g, 76.4%) as a white solid. LCMS (ESI) [M + H]⁺: 206
- Step 4: A mixture of 4-(pyridin-2-yl)thiophene-2-carboxylic acid (133.12 g, 648.05 mmol, 1.00 equiv) and N-(3-amino-5-bromophenyl)methanesulfonamide (171.82 g, 648.05 mmol, 1.00 equiv) and TCFH (272.75 g, 972.08 mmol, 1.50 equiv) and NMI (159.63 g, 1944.159 mmol, 3 equiv) in 800 mL of acetonitrile, this was stirred for 2h at room temperature. The residue was purified onto silica gel column eluted with 10% of methanol in dichloromethane to afford 75g (95% purity) of the desired product. The desired product was purified by trituration with 300 mL of 30% of methanol in dichloromethane. This resulted in N-(3-bromo-5-methanesulfonamidophenyl)-4-(pyridin-2-yl)thiophene-2-carboxamide (54.8097 g, 18.7%) as a white solid. LCMS (ESI) [M + H]⁺: 452. ¹H NMR (300 MHz, DMSO-d₆) δ 10.61 (s, 1H), 10.08 (s, 1H), 8.75 (d, J = 1.4 Hz, 1H), 8.65 (m, J = 4.8, 1.4 Hz, 1H), 8.52 (d, J = 1.3 Hz, 1H), 7.92 (m, J = 2.9, 1.4 Hz, 2H), 7.84 (t, J = 1.8 Hz, 1H), 7.75 (t, J = 1.9 Hz, 1H), 7.36 (td, J = 5.3, 3.1 Hz, 1H), 7.10 (t, J = 1.8 Hz, 1H), 3.08 (s, 3H).

Example 31: N-(3-chloro-5-methanesulfonamidophenyl)-4-(3-methylpyridin-2-yl)thiophene-2-carboxamide



Step 1: A solution of methyl 4-bromothiophene-2-carboxylate (150 g, 678.52 mmol, 1.00 equiv), Pd(dppf)Cl₂ (24.82 g, 33.93 mmol, 0.05 equiv), AcOK (199.77 g, 2035.55 mmol, 3.00 equiv) and bis(pinacolato)diboron (206.76 g, 814.22 mmol, 1.2 equiv) in 1000 mL of dioxane. This was stirred for 2 hours at 80 °C under nitrogen atmosphere. The resulting mixture was cooled and concentrated. The residue was purified onto silica gel column eluted with 10% of ethyl acetate in petroleum ether to afford methyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl) thiophene-2-carboxylate (170 g, 92.5%) as a yellow solid.

Step 2: A solution of methyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxylate (170 g, 634.02 mmol, 1.00 equiv), Pd(dppf)Cl₂ (23.20 g, 31.70 mmol, 0.05 equiv), K₂CO₃ (262.87 g, 1902.06 mmol, 3.00 equiv) and 2-bromo-3-methylpyridine (109.07 g, 634.02 mmol, 1.00 equiv) in 1000 mL of 1,4-dioxane and 200 mL of water, this was stirred for 2 hours at 90 °C under nitrogen atmosphere. The resulting mixture was cooled and concentrated. The residue was purified onto silica gel column chromatography, eluted with 50% of ethyl acetate in petroleum ether to afford methyl 4-(3-methylpyridin-2-yl)thiophene - 2-carboxylate (115 g, 77%) as a yellow solid. LCMS (ESI) [M + H]⁺: 234

Step 3: To a stirred solution of methyl 4-(3-methylpyridin-2-yl)thiophene-2-carboxylate (115 g, 492.95 mmol, 1.00 equiv), LiOH (118.06 g, 4929.49 mmol, 10.00 equiv) in 250 mL of ethanol and 250 mL of water. Then the mixture was stirred for 2 hours at room temperature. The mixture was adjusted pH to 5 with citric acid. The precipitated solids were collected by

filtration and washed with water. The resulted solid was dried under vacuum to afford 4-(3-methylpyridin-2-yl)thiophene-2-carboxylic acid (60 g, 54.9%) as a yellow solid.



LCMS (ESI) [M + H]⁺: 220





Step 4: To a stirred solution of 4-(3-methylpyridin-2-yl)thiophene-2-carboxylic acid (60 g, 273.65 mmol, 1.00 equiv), TCFH (115.17 g, 410.47 mmol, 1.50 equiv) and NMI (67.40 g, 820.94 mmol, 3.00 equiv) in 1000 mL of acetonitrile, to this was added N-(3-amino-5-chlorophenyl)methanesulfonamide (50 g, 226.58 mmol, 0.83 equiv). Then the mixture was stirred for 2 hours at room temperature. The mixture was concentrated and purified by reverse flash chromatography eluting with 56% acetonitrile in water (0.05% FA) to afford N-(3-chloro-5-methanesulfonamidophenyl)-4-(3-methylpyridin-2-yl)thiophene-2-carboxamide (49.6634 g, 42.8%) as a white solid. LCMS (ESI) [M + H]⁺: 422. ¹H NMR (300 MHz, Methanol-d₄) δ 8.49 – 8.41 (m, 1H), 8.21 (d, J = 1.4 Hz, 1H), 7.99 (d, J = 1.4 Hz, 1H), 7.85 – 7.75 (m, 1H), 7.69 (t, J = 1.9 Hz, 1H), 7.60 (t, J = 1.9 Hz, 1H), 7.35 (dd, J = 7.7, 4.9 Hz, 1H), 7.05 (t, J = 2.0 Hz, 1H), 3.06 (s, 3H), 2.50 (s, 3H).






15





Example 18-35 and 333-386





The compounds listed in the following table were prepared using a procedure similar to that described for example 17:







Structure	Example No.	MS (ESI) [M + H] ⁺	¹ H NMR
	18	453	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.63 (s, 1H), 9.24 (s, 1H), 8.74 (d, J = 21.9 Hz, 3H), 8.61 (d, J = 2.5 Hz, 1H), 7.83 (s, 1H), 7.73 (s, 1H), 7.10 (s, 1H), 3.08 (s, 3H)
	19	479	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.32 (s, 1H), 9.35 (s, 1H), 8.02 (s, 1H), 7.80 (s, 1H), 7.73 – 7.63 (m, 2H), 7.06 (d, J = 8.4 Hz, 2H), 6.76 (t, J = 8.4 Hz, 3H), 3.05 (s, 3H), 2.12 (s, 3H).





	20	481	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.26 (s, 1H), 10.04 (s, 1H), 9.61 (s, 1H), 7.98 (s, 1H), 7.81 (t, J = 1.8 Hz, 1H), 7.67 (t, J = 2.0 Hz, 1H), 7.27 – 7.14 (m, 2H), 7.07 (t, J = 1.8 Hz, 1H), 6.97 (d, J = 8.0 Hz, 1H), 6.88 (t, J = 7.4 Hz, 1H), 3.06 (s, 3H), 2.37 (s, 3H).
	21	467	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 9.95 (s, 2H), 8.47 (d, J = 1.4 Hz, 1H), 8.14 (d, J = 1.4 Hz, 1H), 7.81 (t, J = 1.8 Hz, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.53 (dd, J = 7.7, 1.7 Hz, 1H), 7.17 (td, J = 7.7, 1.7 Hz, 1H), 7.09 (t, J = 1.9 Hz, 1H), 6.98 (dd, J = 8.1, 1.2 Hz, 1H), 6.90 (td, J = 7.5, 1.2 Hz, 1H), 3.07 (s, 3H).
	22	423	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.60 (s, 1H), 10.08 (s, 1H), 8.93 (d, J = 4.9 Hz, 2H), 8.73 (s, 1H), 7.70 (dt, J = 5.1, 1.9 Hz, 2H), 7.44 (t, J = 4.9 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 3.08 (s, 3H), 2.88 (s, 3H).
	23	466	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 10.04 (s, 1H), 8.72 – 8.66 (m, 1H), 8.40 (s, 1H), 7.93 (td, J = 7.7, 1.9 Hz, 1H), 7.81 (t, J = 1.8 Hz, 1H), 7.73 – 7.66 (m, 1H), 7.41 – 7.33 (m, 2H), 7.08 (t, J = 1.9 Hz, 1H), 3.06 (s, 3H), 2.71 (s, 3H).






	24	422	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.45 (s, 1H), 10.09 (s, 1H), 8.69 (s, 1H), 8.40 (s, 1H), 7.93 (s, 1H), 7.73 – 7.53 (m, 3H), 7.37 (s, 1H), 6.95 (s, 1H), 3.07 (s, 3H), 2.71 (s, 3H).
	25	453	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.66 (s, 1H), 10.07 (s, 1H), 8.90 (d, J = 4.9 Hz, 2H), 8.84 (d, J = 1.3 Hz, 1H), 8.67 (s, 1H), 7.84 (d, J = 1.8 Hz, 1H), 7.75 (t, J = 1.9 Hz, 1H), 7.45 (t, J = 4.9 Hz, 1H), 7.10 (t, J = 1.9 Hz, 1H), 3.08 (s, 3H).
	26	409	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.68 (s, 1H), 10.09 (s, 1H), 8.90 (d, J = 4.9 Hz, 2H), 8.85 (d, J = 1.3 Hz, 1H), 8.67 (d, J = 1.3 Hz, 1H), 7.71 (dt, J = 4.8, 1.9 Hz, 2H), 7.46 (t, J = 4.9 Hz, 1H), 6.97 (t, J = 1.9 Hz, 1H), 3.08 (s, 3H).
	27	437	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.59 (s, 1H), 10.14 (s, 1H), 8.93 (d, J = 4.9 Hz, 2H), 8.73 (s, 1H), 7.69 (d, J = 2.0 Hz, 2H), 7.44 (t, J = 4.9 Hz, 1H), 6.96 (t, J = 1.9 Hz, 1H), 3.18 (q, J = 7.3 Hz, 2H), 2.88 (s, 3H), 1.22 (t, J = 7.3 Hz, 3H).
	28	436	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.45 (s, 1H), 10.18 (s, 1H), 8.84 – 8.56 (m, 1H), 8.41 (s, 1H), 8.07 – 7.84 (m, 1H), 7.68 (dd, J = 14.5, 7.8 Hz, 3H), 7.49 – 7.20 (m, 1H), 6.96 (d, J = 2.2 Hz, 1H), 3.17 (q, J = 7.2





			Hz, 2H), 2.71 (s, 3H), 1.22 (t, J = 7.2 Hz, 3H).
	29	514	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.65 (s, 1H), 10.09 (s, 1H), 8.84 (d, J = 1.3 Hz, 1H), 8.47 (d, J = 1.3 Hz, 1H), 8.33 – 8.18 (m, 1H), 7.77 – 7.64 (m, 3H), 7.54 (d, J = 7.2 Hz, 2H), 7.47 – 7.31 (m, 4H), 6.96 (t, J = 1.9 Hz, 1H), 5.37 (s, 2H), 3.08 (s, 3H).
	30	557	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 10.09 (s, 1H), 8.45 (d, J = 1.4 Hz, 1H), 8.10 (d, J = 1.3 Hz, 1H), 7.80 (t, J = 1.8 Hz, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.60 (dd, J = 7.7, 1.7 Hz, 1H), 7.49 (d, J = 7.3 Hz, 2H), 7.42 – 7.20 (m, 5H), 7.15 – 7.02 (m, 2H), 5.24 (s, 2H), 3.08 (s, 3H).
	31	422	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.57 (s, 1H), 10.08 (s, 1H), 8.58 – 8.45 (m, 2H), 8.20 (d, J = 1.3 Hz, 1H), 7.75 (dd, J = 7.7, 1.6 Hz, 1H), 7.69 (dt, J = 10.2, 1.9 Hz, 2H), 7.31 (dd, J = 7.7, 4.7 Hz, 1H), 6.97 (t, J = 1.9 Hz, 1H), 3.08 (s, 3H), 2.51 (s, 3H).
	32	558	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.63 (s, 1H), 10.06 (s, 1H), 8.83 (d, J = 1.4 Hz, 1H), 8.47 (d, J = 1.3 Hz, 1H), 8.27 (dd, J = 4.6, 1.3 Hz, 1H), 7.83 (t, J = 1.8 Hz, 1H), 7.75 (t, J = 1.9 Hz, 1H), 7.69 (dd, J = 8.5, 1.2 Hz, 1H), 7.59 – 7.49 (m, 2H), 7.46 – 7.39 (m, 2H), 7.39 –






			7.31 (m, 2H), 7.09 (t, J = 1.9 Hz, 1H), 3.07 (s, 3H).
	33	466	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.55 (s, 1H), 10.06 (s, 1H), 8.51 (d, J = 1.5 Hz, 2H), 8.19 (d, J = 1.3 Hz, 1H), 7.83 (t, J = 1.8 Hz, 1H), 7.76 (dd, J = 8.1, 1.5 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.31 (dd, J = 7.7, 4.7 Hz, 1H), 7.10 (t, J = 1.9 Hz, 1H), 3.07 (s, 3H), 2.50 (s, 3H).
	34	528	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.36 (s, 1H), 10.10 (s, 1H), 8.31 – 8.25 (m, 1H), 8.18 (s, 1H), 7.73 – 7.60 (m, 3H), 7.46 – 7.38 (m, 3H), 7.33 (q, J = 7.4, 6.7 Hz, 3H), 6.95 (t, J = 2.0 Hz, 1H), 5.22 (s, 2H), 3.07 (s, 3H), 2.41 (s, 3H).
	35	440	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 10.06 (s, 1H), 8.69 (d, J = 2.9 Hz, 1H), 8.36 (s, 1H), 7.88 (td, J = 8.7, 3.0 Hz, 1H), 7.77 (dd, J = 8.8, 4.4 Hz, 1H), 7.68 (t, J = 1.8 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 3.07 (s, 3H), 2.69 (s, 3H).
	333	420.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.34 (s, 1H), 10.11 (s, 1H), 8.50 (dd, J = 4.9, 1.7 Hz, 1H), 8.00 (s, 1H), 7.81 – 7.75 (m, 1H), 7.46 (dd, J = 9.9, 2.1 Hz, 2H), 7.35 (dd, J = 7.7, 4.8 Hz, 1H), 6.72 (dt, J = 10.5, 2.2 Hz, 1H), 3.07 (s, 3H), 2.35 (s, 3H), 2.24 (s, 3H).






	334	405.8	¹ H NMR (300 MHz, Methanol-d ₄) δ 8.45 (d, J = 4.5 Hz, 1H), 8.21 (d, J = 1.4 Hz, 1H), 7.99 (d, J = 1.4 Hz, 1H), 7.80 (d, J = 7.7 Hz, 1H), 7.55 – 7.49 (m, 1H), 7.42 – 7.31 (m, 2H), 6.81 (dt, J = 10.3, 2.2 Hz, 1H), 3.06 (s, 3H), 2.50 (s, 3H).
	335	467	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.57 (s, 1H), 10.07 (s, 1H), 8.65 (s, 3H), 7.72 – 7.63 (m, 2H), 6.95 (s, 1H), 4.27 (q, J = 6.9 Hz, 2H), 3.08 (s, 3H), 2.84 (s, 3H), 1.40 (t, J = 7.0 Hz, 3H).
	336	440.85	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.58 (s, 1H), 10.08 (s, 1H), 9.01 (s, 2H), 8.68 (s, 1H), 7.68 (dt, J = 7.0, 2.0 Hz, 2H), 6.95 (t, J = 2.0 Hz, 1H), 3.07 (s, 3H), 2.85 (s, 3H).
	337	465.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.42 (s, 2H), 8.42 – 8.30 (m, 2H), 7.70 – 7.60 (m, 2H), 7.50 (dd, J = 8.7, 3.0 Hz, 1H), 6.94 (t, J = 2.0 Hz, 1H), 4.16 (q, J = 6.9 Hz, 2H), 3.06 (s, 3H), 2.67 (s, 3H), 1.38 (t, J = 6.9 Hz, 3H).
	338	426.85	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.68 (s, 1H), 10.09 (s, 1H), 8.99 (s, 2H), 8.80 (s, 1H), 8.63 (s, 1H), 7.74 – 7.66 (m, 2H), 6.97 (d, J = 2.1 Hz, 1H), 3.08 (s, 3H).
	339	452.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.57 (s, 1H), 10.08 (s, 1H), 8.67 (d, J = 1.4 Hz, 1H), 8.33 (dd, J = 5.8, 2.1 Hz, 2H), 7.85 (d, J





			= 8.7 Hz, 1H), 7.69 (dt, J = 6.3, 1.9 Hz, 2H), 7.50 (dd, J = 8.8, 3.0 Hz, 1H), 6.96 (t, J = 2.0 Hz, 1H), 4.16 (q, J = 6.9 Hz, 2H), 3.08 (s, 3H), 1.37 (t, J = 7.0 Hz, 3H).
	340	439.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.65 (s, 1H), 10.06 (s, 1H), 8.76 (d, J = 1.3 Hz, 1H), 8.63 (s, 2H), 8.50 (d, J = 1.3 Hz, 1H), 7.70 (t, J = 1.8 Hz, 2H), 6.96 (t, J = 2.0 Hz, 1H), 3.97 (s, 3H), 3.07 (s, 3H).
	341	453.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.56 (s, 1H), 10.05 (s, 1H), 8.65 (d, J = 8.1 Hz, 3H), 7.68 (dt, J = 6.0, 1.8 Hz, 2H), 6.95 (t, J = 1.9 Hz, 1H), 3.97 (s, 3H), 3.07 (d, J = 1.3 Hz, 3H), 2.83 (s, 3H).
	342	452.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.64 (s, 1H), 10.08 (s, 1H), 8.76 (s, 1H), 8.62 (s, 2H), 8.49 (s, 1H), 7.70 (s, 2H), 6.96 (d, J = 2.3 Hz, 1H), 4.26 (q, J = 6.9 Hz, 2H), 3.08 (s, 3H), 1.39 (t, J = 6.9 Hz, 3H).
	343	440.9	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.67 (s, 1H), 10.13 (s, 1H), 8.99 (d, J = 0.8 Hz, 2H), 8.80 (d, J = 1.3 Hz, 1H), 8.63 (d, J = 1.3 Hz, 1H), 7.70 (d, J = 2.0 Hz, 2H), 6.98 (t, J = 2.0 Hz, 1H), 3.19 (q, J = 7.3 Hz, 2H), 1.23 (t, J = 7.3 Hz, 3H).






	344	454.85	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.58 (s, 1H), 10.12 (s, 1H), 9.00 (s, 2H), 8.67 (s, 1H), 7.67 (d, J = 2.0 Hz, 2H), 6.95 (t, J = 1.7 Hz, 1H), 3.18 (q, J = 7.3 Hz, 2H), 2.85 (s, 3H), 1.22 (t, J = 7.3 Hz, 3H).
	345	488.9	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.61 (s, 1H), 10.07 (s, 1H), 8.89 (s, 2H), 8.70 (s, 1H), 7.74 – 7.17 (m, 3H), 6.96 (t, J = 1.9 Hz, 1H), 3.08 (s, 3H), 2.86 (s, 3H).
	346	454	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 10.13 (s, 1H), 8.69 (d, J = 3.0 Hz, 1H), 8.37 (s, 1H), 7.89 (td, J = 8.7, 3.0 Hz, 1H), 7.77 (dd, J = 8.9, 4.4 Hz, 1H), 7.65 (dt, J = 8.3, 1.9 Hz, 2H), 6.96 (t, J = 1.9 Hz, 1H), 3.18 (q, J = 7.3 Hz, 2H), 2.69 (s, 3H), 1.22 (t, J = 7.3 Hz, 3H).
	347	472	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 10.15 (s, 1H), 8.66 (d, J = 2.5 Hz, 1H), 8.24 – 8.07 (m, 2H), 7.71 – 7.57 (m, 2H), 6.95 (t, J = 2.0 Hz, 1H), 3.16 (q, J = 7.2 Hz, 2H), 2.52 (d, J = 1.5 Hz, 3H), 1.21 (t, J = 7.3 Hz, 3H).
	348	453.85	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.31 (s, 1H), 10.07 (s, 1H), 8.50 (d, J = 2.8 Hz, 1H), 7.97 (s, 1H), 7.78 (dd, J = 9.5, 2.8 Hz, 1H), 7.66 (t, J = 1.8 Hz, 1H), 7.59 (t, J = 1.9 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 3.05 (s,





			3H), 2.33 (s, 3H), 2.26 (s, 3H).
	349	458	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 10.07 (s, 1H), 8.66 (d, J = 2.4 Hz, 1H), 8.18 (d, J = 1.7 Hz, 1H), 8.14 (ddd, J = 10.4, 8.9, 2.4 Hz, 1H), 7.66 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 3.06 (s, 3H), 2.52 (s, 3H).
	350	468	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.30 (s, 1H), 10.13 (s, 1H), 8.50 (d, J = 2.9 Hz, 1H), 7.97 (s, 1H), 7.78 (dd, J = 9.7, 2.9 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 7.59 (t, J = 1.9 Hz, 1H), 6.94 (t, J = 2.0 Hz, 1H), 3.16 (q, J = 7.3 Hz, 2H), 2.33 (s, 3H), 2.26 (s, 3H), 1.20 (t, J = 7.3 Hz, 3H).
	351	470	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.37 (s, 1H), 10.08 (s, 1H), 8.28 (d, J = 2.3 Hz, 1H), 8.09 (s, 1H), 7.67 (ddd, J = 7.4, 4.8, 2.4 Hz, 2H), 7.63 (t, J = 1.9 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 3.89 (s, 3H), 3.07 (s, 3H), 2.41 (s, 3H).
	352	452	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 10.11 (s, 1H), 8.40 (d, J = 2.9 Hz, 1H), 8.34 (s, 1H), 7.71 – 7.61 (m, 3H), 7.53 (dd, J = 8.7, 3.0 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 3.89 (s, 3H), 3.07 (s, 3H), 2.67 (s, 3H).






	353	467	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.57 (s, 1H), 10.11 (s, 1H), 8.66 (d, J = 6.6 Hz, 3H), 7.68 (d, J = 2.0 Hz, 2H), 6.95 (t, J = 2.0 Hz, 1H), 3.98 (s, 3H), 3.18 (q, J = 7.3 Hz, 2H), 2.84 (s, 3H), 1.22 (t, J = 7.3 Hz, 3H).
	354	484	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.36 (s, 1H), 10.11 (s, 1H), 8.28 (d, J = 2.3 Hz, 1H), 8.09 (s, 1H), 7.72 – 7.56 (m, 3H), 6.95 (t, J = 2.0 Hz, 1H), 3.89 (s, 3H), 3.17 (q, J = 7.3 Hz, 2H), 2.40 (s, 3H), 1.21 (t, J = 7.3 Hz, 3H).
	355	487.95	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.38 (s, 1H), 10.11 (s, 1H), 8.73 (d, J = 2.6 Hz, 1H), 8.30 (dd, J = 8.5, 2.6 Hz, 1H), 8.08 (s, 1H), 7.63 (dt, J = 16.4, 1.9 Hz, 2H), 6.95 (t, J = 2.0 Hz, 1H), 3.17 (q, J = 7.3 Hz, 2H), 2.40 (s, 3H), 1.20 (t, J = 7.3 Hz, 3H).
	356	489.9	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.51 (s, 1H), 10.09 (s, 1H), 9.07 (s, 1H), 8.48 (s, 1H), 8.40 – 8.33 (m, 1H), 7.93 (d, J = 8.4 Hz, 1H), 7.68 (d, J = 1.9 Hz, 1H), 7.64 (s, 1H), 6.96 (t, J = 2.0 Hz, 1H), 3.08 (s, 3H), 2.77 (s, 3H).
	357	490.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.67 (s, 1H), 10.11 (s, 1H), 9.35 (s, 2H), 8.78 (s, 1H), 7.68 (dt, J = 10.4, 1.9 Hz, 2H), 6.95 (t, J = 2.0 Hz, 1H), 3.07 (s, 3H), 2.91 (s, 3H).





	358	469.95	¹ H NMR (400 MHz, Chloroform-d) δ 8.33 (s, 1H), 8.23 (s, 1H), 7.85 (s, 1H), 7.63 (s, 1H), 7.50 (d, J = 19.4 Hz, 2H), 7.19 (d, J = 10.8 Hz, 1H), 6.97 (t, J = 1.7 Hz, 1H), 3.94 (s, 3H), 3.04 (s, 3H), 2.52 (d, J = 1.8 Hz, 3H).
	359	473.9	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.39 (s, 1H), 10.07 (s, 1H), 8.73 (d, J = 2.6 Hz, 1H), 8.30 (dd, J = 8.5, 2.6 Hz, 1H), 8.08 (s, 1H), 7.63 (dt, J = 19.9, 1.9 Hz, 2H), 6.94 (t, J = 2.0 Hz, 1H), 3.06 (s, 3H), 2.40 (s, 3H).
	360	483.9	¹ H NMR (400 MHz, Chloroform-d) δ 8.25 (s, 2H), 7.85 (s, 1H), 7.54 (d, J = 1.8 Hz, 1H), 7.47 (s, 1H), 7.22 (s, 1H), 7.19 (s, 1H), 6.98 (t, J = 1.8 Hz, 1H), 3.95 (s, 3H), 3.17 (q, J = 7.3 Hz, 2H), 2.54 (d, J = 1.8 Hz, 3H), 1.37 (t, J = 7.3 Hz, 3H).
	361	465.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.29 (s, 1H), 10.09 (s, 1H), 8.21 (d, J = 2.9 Hz, 1H), 7.95 (s, 1H), 7.71 – 7.54 (m, 2H), 7.39 (d, J = 2.9 Hz, 1H), 6.94 (t, J = 2.0 Hz, 1H), 3.87 (s, 3H), 3.05 (s, 3H), 2.32 (s, 3H), 2.22 (s, 3H).
	362	458	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 10.09 (s, 1H), 8.72 (s, 2H), 8.06 (s, 1H), 7.67 (t, J = 1.9 Hz, 1H), 7.60 (t, J = 1.9 Hz, 1H), 6.96 (t, J = 1.9 Hz, 1H), 3.07 (s, 3H), 2.41 (s, 3H).



	363	488.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.87 (s, 1H), 8.60 (d, J = 2.4 Hz, 1H), 7.90 (dd, J = 9.6, 2.5 Hz, 1H), 7.77 – 7.69 (m, 2H), 7.63 (t, J = 1.9 Hz, 1H), 7.37 (s, 1H), 7.08 (d, J = 1.9 Hz, 1H), 6.91 (t, J = 1.9 Hz, 1H), 3.78 (s, 3H), 3.06 (s, 3H).
	364	485.95	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.37 (s, 1H), 10.06 (s, 1H), 8.39 (d, J = 2.6 Hz, 1H), 8.05 (s, 1H), 7.76 (d, J = 2.6 Hz, 1H), 7.67 (t, J = 1.8 Hz, 1H), 7.61 (t, J = 1.9 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 3.92 (s, 3H), 3.06 (s, 3H), 2.38 (s, 3H).
	365	481	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.55 (s, 1H), 10.12 (s, 1H), 8.64 (d, J = 2.1 Hz, 3H), 7.67 (d, J = 2.0 Hz, 2H), 6.94 (t, J = 2.0 Hz, 1H), 4.26 (q, J = 7.0 Hz, 2H), 3.17 (q, J = 7.3 Hz, 2H), 2.83 (s, 3H), 1.39 (t, J = 6.9 Hz, 3H), 1.21 (t, J = 7.3 Hz, 3H).
	366	505	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.65 (s, 1H), 10.11 (s, 1H), 9.35 (s, 2H), 8.78 (s, 1H), 7.67 (dd, J = 4.6, 2.2 Hz, 2H), 6.96 (t, J = 2.0 Hz, 1H), 3.18 (q, J = 7.3 Hz, 2H), 2.91 (s, 3H), 1.22 (t, J = 7.3 Hz, 3H).

	367	514.8	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.54 (s, 1H), 10.07 (s, 1H), 9.38 (dd, J = 2.2, 0.9 Hz, 1H), 9.09 (d, J = 2.3 Hz, 1H), 8.27 (s, 1H), 7.75 – 7.34 (m, 2H), 6.96 (t, J = 1.9 Hz, 1H), 3.07 (s, 3H), 2.58 (s, 3H).
	368	507.8	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.51 (s, 1H), 10.08 (s, 1H), 8.98 (d, J = 1.8 Hz, 1H), 8.51 (dd, J = 10.3, 1.9 Hz, 1H), 8.27 (d, J = 1.9 Hz, 1H), 7.65 (dt, J = 17.9, 1.9 Hz, 2H), 6.96 (t, J = 1.9 Hz, 1H), 3.06 (s, 3H), 2.60 (d, J = 1.5 Hz, 3H).
	369	476.85	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.74 (s, 1H), 10.11 (s, 1H), 9.34 (t, J = 0.9 Hz, 2H), 8.87 (dd, J = 16.7, 1.2 Hz, 2H), 7.71 (dt, J = 6.6, 1.9 Hz, 2H), 6.97 (t, J = 1.9 Hz, 1H), 3.09 (s, 3H).
	370	476	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.67 (s, 1H), 10.10 (s, 1H), 9.08 – 8.98 (m, 1H), 8.80 (d, J = 1.4 Hz, 1H), 8.72 (d, J = 1.3 Hz, 1H), 8.34 (dd, J = 8.5, 2.4 Hz, 1H), 8.15 (d, J = 8.4 Hz, 1H), 7.69 (dt, J = 11.0, 1.9 Hz, 2H), 6.97 (t, J = 1.9 Hz, 1H), 3.08 (s, 3H).
	371	493.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.74 (s, 1H), 10.09 (s, 1H), 8.94 (s, 1H), 8.81 (d, J = 1.4 Hz, 1H), 8.63 (t, J = 1.6 Hz, 1H), 8.48 (dd, J = 11.4, 1.9 Hz, 1H), 7.69 (dt, J = 7.5, 1.9 Hz, 2H), 6.97 (t, J


			= 1.9 Hz, 1H), 3.08 (s, 3H).
	372	450	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.49 – 8.40 (m, 1H), 8.18 (d, J = 1.4 Hz, 1H), 7.97 (d, J = 1.4 Hz, 1H), 7.78 (ddd, J = 7.8, 1.8, 0.9 Hz, 1H), 7.66 (t, J = 1.9 Hz, 1H), 7.55 (t, J = 1.9 Hz, 1H), 7.33 (dd, J = 7.7, 4.9 Hz, 1H), 7.02 (t, J = 1.9 Hz, 1H), 3.21 – 3.09 (m, 2H), 2.48 (s, 3H), 1.91 – 1.71 (m, 2H), 1.03 (t, J = 7.5 Hz, 3H).
	373	450	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.43 (dd, J = 4.7, 1.7 Hz, 1H), 8.18 (d, J = 1.4 Hz, 1H), 7.97 (d, J = 1.4 Hz, 1H), 7.78 (dd, J = 7.7, 1.4 Hz, 1H), 7.68 (t, J = 1.9 Hz, 1H), 7.54 (t, J = 1.8 Hz, 1H), 7.33 (dd, J = 7.7, 4.9 Hz, 1H), 7.04 (t, J = 1.9 Hz, 1H), 3.40 – 3.33 (m, 1H), 2.48 (s, 3H), 1.36 (d, J = 6.8 Hz, 6H).
	374	436	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.61 (s, 1H), 8.59 – 8.45 (m, 2H), 8.21 (d, J = 1.3 Hz, 1H), 7.92 (d, J = 2.0 Hz, 1H), 7.79 – 7.68 (m, 2H), 7.31 (dd, J = 7.7, 4.7 Hz, 1H), 7.28 (d, J = 1.9 Hz, 1H), 3.26 (s, 3H), 3.02 (s, 3H), 2.51 (s, J = 2.1 Hz, 3H).
	375	465	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.51 (s, 1H), 10.07 (s, 1H), 9.03 (t, J = 1.5 Hz, 1H), 8.60 (dd, J = 10.4, 1.7 Hz, 1H), 8.28 (d, J = 2.1 Hz, 1H), 7.65 (dt, J = 15.0, 1.9 Hz, 2H), 6.96 (t, J =

			1.9 Hz, 1H), 3.07 (s, 3H), 2.60 (d, J = 1.5 Hz, 3H).
	376	461	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.35 (s, 1H), 10.08 (s, 1H), 8.95 (d, J = 2.0 Hz, 1H), 8.35 (d, J = 2.1 Hz, 1H), 8.03 (s, 1H), 7.67 – 7.62 (m, 1H), 7.59 (t, J = 2.0 Hz, 1H), 6.95 (t, J = 2.0 Hz, 1H), 3.06 (s, 3H), 2.38 (s, 3H), 2.31 (s, 3H).
	377	447	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.50 (s, 1H), 10.09 (s, 1H), 9.12 (dd, J = 2.2, 0.9 Hz, 1H), 8.60 – 8.30 (m, 2H), 7.90 (dd, J = 8.3, 0.9 Hz, 1H), 7.66 (dt, J = 15.7, 1.9 Hz, 2H), 6.96 (t, J = 2.0 Hz, 1H), 3.08 (s, 3H), 2.76 (s, 3H).
	378	475	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.37 (s, 1H), 10.09 (s, 1H), 7.98 (s, 1H), 7.67 (t, J = 1.8 Hz, 1H), 7.59 (t, J = 1.9 Hz, 1H), 7.41 (t, J = 8.6 Hz, 2H), 6.95 (t, J = 1.9 Hz, 1H), 3.06 (s, 3H), 2.35 (s, 3H).
	379	481	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.66 (s, 1H), 10.09 (s, 1H), 8.78 (d, J = 1.3 Hz, 1H), 8.62 (s, 2H), 8.55 (d, J = 1.3 Hz, 1H), 7.70 (dd, J = 4.7, 2.9 Hz, 2H), 6.96 (t, J = 2.0 Hz, 1H), 3.08 (s, 3H), 1.38 (s, 9H).
	380	494	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 10.07 (s, 1H), 8.56 – 8.25 (m, 2H), 7.72 – 7.61 (m, 3H), 7.57 (dd, J = 8.6, 2.8 Hz, 1H), 6.94 (t, J = 2.0 Hz, 1H), 3.06 (s,

			3H), 2.69 (s, 3H), 1.36 (s, 9H).
	381	512	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 10.07 (s, 1H), 8.28 (dd, J = 2.3, 1.3 Hz, 1H), 8.19 (d, J = 1.6 Hz, 1H), 7.70 – 7.59 (m, 3H), 6.94 (t, J = 2.0 Hz, 1H), 3.06 (s, 3H), 2.53 (d, J = 1.5 Hz, 3H), 1.40 (s, 9H).
	382	505	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.54 (s, 1H), 8.60 – 8.49 (m, 1H), 8.34 (d, J = 1.4 Hz, 1H), 8.05 (d, J = 1.3 Hz, 1H), 7.74 (d, J = 6.3 Hz, 1H), 7.68 (s, 1H), 7.63 (s, 1H), 7.38 (dd, J = 7.7, 4.6 Hz, 1H), 6.94 (s, 1H), 4.08 (t, J = 7.5 Hz, 2H), 3.85 (t, J = 7.7 Hz, 2H), 3.58 (s, 2H), 3.05 (s, 3H), 2.22 – 2.10 (m, 2H).
	383	519	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.54 (s, 1H), 8.55 (dd, J = 4.8, 1.7 Hz, 1H), 8.33 (d, J = 1.4 Hz, 1H), 7.97 (d, J = 1.3 Hz, 1H), 7.76 – 7.62 (m, 3H), 7.37 (dd, J = 7.8, 4.7 Hz, 1H), 6.95 (t, J = 2.0 Hz, 1H), 3.80 (s, 2H), 3.40 (t, J = 6.7 Hz, 2H), 3.28 (d, J = 6.7 Hz, 2H), 3.06 (s, 3H), 1.80 (dp, J = 26.7, 6.8 Hz, 4H).
	384	500	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.51 (s, 1H), 10.02 (s, 1H), 8.73 (d, J = 4.9 Hz, 2H), 8.56 (dd, J = 4.7, 1.7 Hz, 1H), 8.39 (d, J = 1.4 Hz, 1H), 8.19 (d, J = 1.3 Hz, 1H), 7.78 (dd, J = 7.8, 1.7 Hz, 1H), 7.66 (dt, J = 14.8, 1.9 Hz, 2H), 7.40 – 7.32 (m, 2H), 6.95 (t, J

			= 2.0 Hz, 1H), 4.48 (s, 2H), 3.06 (s, 3H).
	385	509	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.33 (s, 1H), 10.06 (s, 1H), 8.54 (dd, J = 4.8, 1.7 Hz, 1H), 7.90 (s, 1H), 7.77 – 7.66 (m, 2H), 7.60 (t, J = 1.9 Hz, 1H), 7.40 (dd, J = 7.8, 4.7 Hz, 1H), 6.94 (t, J = 2.0 Hz, 1H), 3.62 (s, 2H), 3.06 (s, 3H), 2.82 (s, 3H), 2.75 (s, 3H), 2.28 (s, 3H).
	386	422	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.47 (s, 1H), 10.07 (s, 1H), 8.54 – 8.41 (m, 1H), 8.05 (d, J = 4.1 Hz, 1H), 7.83 – 7.54 (m, 4H), 7.32 (dd, J = 7.7, 4.7 Hz, 1H), 6.97 (t, J = 2.0 Hz, 1H), 3.08 (s, 3H), 2.59 (s, 3H).

Example 36: N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(pyridin-2-yl)-1H-pyrazole-4-carboxamide



- 5 Step 1: To a stirred solution of 2-fluoropyridine (150 g, 1.54 mol), KOH (260 g, 4.62 mol), ethyl 1H-pyrazole-4-carboxylate (258.7 g, 1.85 mol) in 3500 mL of DMSO. Then the mixture was stirred for 2h at 100 °C. The mixture was cooled then quenched with 3000 mL of water. The aqueous layer was extracted with 3 x 1000 mL of ethyl acetate. The mixture was


concentrated and purified onto silica gel eluting with 35% of ethyl acetate in petroleum ether to afford ethyl 1-(pyridin-2-yl)-1H-pyrazole-4-carboxylate (168 g, 49.5 %) as a yellow solid. LCMS (ESI) [M + H]⁺: 218






Step 2: To a stirred solution of ethyl 1-(pyridin-2-yl)-1H-pyrazole-4-carboxylate (168g, 0.77mol), LiOH (53 g, 2.32mol) in 1000 mL of water and 3000 mL of tetrahydrofuran Then the mixture was stirred for 2h at room temperature. the mixture was adjusted pH 6 with conc. HCl. The aqueous layer was extracted with 3 x 500 mL of ethyl acetate. This was concentrated and purified by reverse flash chromatography eluting with 50% acetonitrile in water (0.05% FA) to afford 1-(pyridin-2-yl)-1H-pyrazole-4- carboxylic acid (116 g, 80.5 %) as an off-white solid. LCMS (ESI) [M + H]⁺: 190






Step 4: To a stirred solution of 1-(pyridin-2-yl)-1H-pyrazole-4-carboxylic acid (95g, 0.51mol), TCFH (211 g, 0.75mol), NMI (246g, 3.03mol), N-(3-amino-5-bromophenyl)methanesulfonamide (134g, 0.51mol) in 300 mL of acetonitrile. Then the mixture was stirred for 2h at room temperature. The mixture was concentrated and purified by reverse flash chromatography eluting with 50% acetonitrile in water (0.05% FA) to afford N-(3-bromo-5-methanesulfonamidophenyl)-1- (pyridin-2-yl)-1H-pyrazole-4 –carbo xamide (51.5 g , 23.6 %) as a white solid. LCMS (ESI) [M + H]⁺: 436. ¹H NMR (300 MHz, Acetone-d₆) δ 9.67 (s, 1H), 9.29 (d, J = 0.9 Hz, 1H), 8.83 (s, 1H), 8.54 (dt, J = 4.8, 1.4 Hz, 1H), 8.26 (d, J = 0.8 Hz, 1H), 8.13 – 8.01 (m, 2H), 7.99 (dt, J = 2.5, 1.2 Hz, 1H), 7.80 (dt, J = 2.9, 1.4 Hz, 1H), 7.44 (h, J = 4.7 Hz, 1H), 7.26 (t, J = 1.9 Hz, 1H), 3.11 (s, 3H).




Examples 37-49 and 387

The compounds listed in the following table were prepared using a procedure similar to that described for example 36:


Structure	Example No.	MS (ESI) [M + H] ⁺	¹ H NMR
	37	392	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.31 (s, 1H), 10.08 (s, 1H), 9.43 (d, J = 0.8 Hz, 1H), 8.56 (ddd, J = 4.8, 1.8, 0.9 Hz, 1H), 8.30 (d, J = 0.8 Hz, 1H), 8.16 – 7.90 (m, 2H), 7.73 (t, J = 1.9 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 7.47 (ddd, J = 7.2, 4.8, 1.2 Hz, 1H), 6.95 (t,

			J = 1.9 Hz, 1H), 3.07 (s, 3H).
	38	461	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.34 (s, 1H), 10.08 (s, 1H), 9.44 (s, 1H), 8.81 (d, J = 5.1 Hz, 1H), 8.37 (d, J = 3.8 Hz, 2H), 7.93 (dd, J = 5.0, 1.4 Hz, 1H), 7.85 (t, J = 1.8 Hz, 1H), 7.68 (t, J = 1.9 Hz, 1H), 7.09 (t, J = 1.9 Hz, 1H), 3.07 (s, 3H).
	39	454	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.29 (s, 1H), 10.06 (s, 1H), 9.36 (d, J = 0.8 Hz, 1H), 8.61 (t, J = 1.7 Hz, 1H), 8.31 (d, J = 0.8 Hz, 1H), 8.17 – 7.91 (m, 2H), 7.85 (t, J = 1.8 Hz, 1H), 7.69 (t, J = 1.9 Hz, 1H), 7.08 (t, J = 1.9 Hz, 1H), 3.07 (s, 3H).
	40	437	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.34 (s, 1H), 10.07 (s, 1H), 9.42 (s, 1H), 9.29 (d, J = 1.5 Hz, 1H), 8.74 (d, J = 2.6 Hz, 1H), 8.66 (dd, J = 2.6, 1.4 Hz, 1H), 8.40 (s, 1H), 7.85 (t, J = 1.8 Hz, 1H), 7.68 (t, J = 1.9 Hz, 1H), 7.09 (t, J = 1.9 Hz, 1H), 3.08 (s, 3H).
	41	459 [M-H]	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.35 (s, 1H), 10.08 (s, 1H), 9.49 (s, 1H), 9.08 (dd, J = 2.2, 0.8 Hz, 1H), 8.55 (dd, J = 8.6, 2.2 Hz, 1H), 8.39 (s, 1H), 8.14 (d, J = 8.6 Hz, 1H), 7.85 (t, J = 1.8 Hz, 1H), 7.69 (d, J = 2.0 Hz, 1H), 7.09 (t, J = 1.9 Hz, 1H), 3.08 (s, 3H).
	42	450	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.27 (s, 1H), 10.06 (s, 1H), 9.40 (s, 1H), 8.39 (s, 1H), 8.27 (s, 1H), 8.03 – 7.79 (m, 3H), 7.69 (d, J = 1.9 Hz, 1H), 7.08 (t, J = 1.9

			Hz, 1H), 3.07 (s, 3H), 2.38 (s, 3H).
	43	504	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.37 (s, 1H), 10.08 (s, 1H), 9.50 (d, J = 0.8 Hz, 1H), 9.14 – 8.84 (m, 1H), 8.56 – 8.42 (m, 1H), 8.38 (d, J = 0.8 Hz, 1H), 8.18 (d, J = 8.6 Hz, 1H), 7.86 (t, J = 1.8 Hz, 1H), 7.69 (t, J = 1.9 Hz, 1H), 7.09 (t, J = 1.9 Hz, 1H), 3.08 (s, 3H).
	44	474	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.24 (s, 1H), 10.10 (s, 1H), 8.78 (s, 1H), 8.41 (s, 1H), 7.94 (d, J = 7.6 Hz, 1H), 7.83 (dd, J = 5.0, 3.1 Hz, 2H), 7.73 – 7.62 (m, 2H), 7.08 (t, J = 1.9 Hz, 1H), 3.07 (s, 3H), 2.14 (s, 3H).
	45	463	¹ H NMR (400 MHz, Chloroform-d) δ 8.45 (d, J = 4.6 Hz, 1H), 7.89 – 7.73 (m, 2H), 7.58 (t, J = 9.8 Hz, 3H), 7.48 – 7.34 (m, 2H), 7.16 (d, J = 1.8 Hz, 1H), 6.37 (s, 1H), 3.00 (s, 3H), 2.16 (s, 3H), 2.07 (s, 3H).
	46	417	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.38 (s, 1H), 10.10 (s, 1H), 9.39 (s, 1H), 8.84 (dd, J = 4.8, 1.7 Hz, 1H), 8.57 (dd, J = 7.9, 1.7 Hz, 1H), 8.40 (s, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.67 (dd, J = 7.9, 4.8 Hz, 1H), 7.63 (t, J = 1.9 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 3.07 (s, 3H).
	47	460	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.32 (s, 1H), 10.08 (s, 1H), 9.09 (s, 1H), 8.93 – 8.84 (m, 1H), 8.54 (dd, J = 7.9, 1.6 Hz, 1H), 8.32 (s, 1H), 7.82 (dd, J = 7.9, 4.8 Hz, 1H), 7.71 (t, J = 1.8 Hz, 1H), 7.61 (t, J =

			1.9 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 3.07 (s, 3H).
	48	441	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.03 (s, 1H), 9.97 (s, 1H), 8.40 (s, 1H), 8.03 (s, 1H), 7.80 (t, J = 1.8 Hz, 1H), 7.63 (t, J = 1.9 Hz, 1H), 7.05 (t, J = 1.9 Hz, 1H), 4.31 – 4.06 (m, 1H), 3.06 (s, 3H), 2.14 – 1.94 (m, 2H), 1.81 (t, J = 10.7 Hz, 2H), 1.69 (td, J = 11.9, 3.5 Hz, 3H), 1.41 (q, J = 12.7 Hz, 2H), 1.22 (q, J = 13.5, 13.0 Hz, 1H).
	49	450	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.25 (s, 1H), 10.06 (s, 1H), 9.10 (d, J = 0.8 Hz, 1H), 8.43 (dd, J = 4.8, 1.7 Hz, 1H), 8.28 (d, J = 0.8 Hz, 1H), 8.02 – 7.91 (m, 1H), 7.85 (t, J = 1.8 Hz, 1H), 7.68 (t, J = 1.9 Hz, 1H), 7.47 (dd, J = 7.6, 4.7 Hz, 1H), 7.08 (t, J = 1.9 Hz, 1H), 3.07 (s, 3H), 2.51 (dd, J = 3.9, 2.0 Hz, 3H).
	387	470.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.27 (s, 1H), 10.06 (s, 1H), 9.15 (d, J = 8.9 Hz, 2H), 8.82 – 8.64 (m, 3H), 8.23 (dd, J = 7.7, 1.7 Hz, 1H), 8.00 (s, 1H), 7.80 – 7.67 (m, 2H), 7.60 (t, J = 1.9 Hz, 1H), 6.94 (t, J = 2.0 Hz, 1H), 3.07 (s, 3H).

Example 50: N-(3-bromo-5-(methanesulfonamido)phenyl)-1-(pyrimidin-2-yl)-1H-pyrazole-4-carboxamide



Step 1: A solution of ethyl 1H-pyrazole-4-carboxylate (500 mg, 3.568 mmol, 1 equiv) and
 5 pyrimidine, 2-bromo-(567.23 mg, 3.568 mmol, 1equiv) and (1S,2S)-N1, N2-
 dimethylcyclohexane-1,2-diamine (101.50 mg, 0.714 mmol, 0.2 equiv) and CuI (67.95 mg,
 0.357 mmol, 0.1 equiv) and K₂CO₃ (1479.27 mg, 10.704 mmol, 3 equiv) in Toluene (5 mL)
 was stirred for 2 h at 110 °C under nitrogen atmosphere. The aqueous layer was extracted
 with EtOAc (3x30 mL). The residue was purified by silica gel column chromatography,
 10 eluted with Petroleum ether / EtOAc (1:1) to afford ethyl 1-(pyrimidin-2-yl) pyrazole-4-
 carboxylate (440 mg, 56.52%) as a white solid. LCMS (ESI) [M + H]⁺: 219

Step 2: A solution of ethyl 1-(pyrimidin-2-yl) pyrazole-4-carboxylate (440 mg, 2.016 mmol,
 1 equiv) and LiOH (144.87 mg, 6.048 mmol, 3 equiv) in EtOH (4 mL) and H₂O (1.5 mL) was
 stirred for 1 h at 60 °C. The mixture was acidified to pH 6 with HCl (aq.). The residue was
 15 purified by reverse phase flash chromatography with the following conditions: column, silica
 gel; mobile phase, MeCN in water, 0% to 100% gradient in 20 min; detector, UV 254 nm.
 This resulted in 1-(pyrimidin-2-yl) pyrazole-4-carboxylic acid (380 mg, 99.10%) as a white
 solid. LCMS (ESI) [M + H]⁺: 191




Step 3: To a stirred solution of 1-(pyrimidin-2-yl) pyrazole-4-carboxylic acid (80 mg, 0.421
 20 mmol, 1 equiv) and N-(3-amino-5-bromophenyl) methanesulfonamide (111.54 mg, 0.421
 mmol, 1 equiv) in DMSO (4 mL) were added NMI (103.62 mg, 1.263 mmol, 3 equiv) and
 NMI (103.62 mg, 1.263 mmol, 3 equiv) in portions at room temperature. The residue was





purified by reverse phase flash chromatography with the following conditions: column, silica gel; mobile phase, MeCN in water, 0% to 100% gradient in 20 min; detector, UV 254 nm.





This resulted in N-(3-bromo-5-methanesulfonamidophenyl)-1-(pyrimidin-2-yl) pyrazole-4-carboxamide (19.3 mg, 10.49%) as a white solid. LCMS (ESI) $[M + H]^+$: 436. ^1H NMR (300 MHz, DMSO- d_6) δ 10.34 (s, 1H), 10.07 (s, 1H), 9.46 (s, 1H), 8.96 (d, $J = 4.8$ Hz, 2H), 8.33 (s, 1H), 7.86 (t, $J = 1.8$ Hz, 1H), 7.69 (t, $J = 1.8$ Hz, 1H), 7.59 (t, $J = 4.8$ Hz, 1H), 7.09 (t, $J = 1.9$ Hz, 1H), 3.08 (s, 3H).





Examples 51-60 and 388-393


The compounds listed in the following table were prepared using a procedure similar to that described for example 50:

Structure	Example No.	MS (ESI) $[M + H]^+$	^1H NMR
	51	541	^1H NMR (300 MHz, DMSO- d_6) δ 10.14 (s, 1H), 10.06 (s, 1H), 8.83 (d, $J = 0.6$ Hz, 1H), 8.29 (d, $J = 0.6$ Hz, 1H), 7.82 (t, $J = 1.8$ Hz, 1H), 7.72 – 7.59 (m, 2H), 7.39 (dddd, $J = 18.5, 11.4, 8.6, 4.3$ Hz, 7H), 7.20 – 7.02 (m, 2H), 5.29 (s, 2H), 3.07 (s, 3H).
	52	460	^1H NMR (300 MHz, DMSO- d_6) δ 10.23 (s, 1H), 10.09 (s, 1H), 9.23 (s, 1H), 8.41 (d, $J = 2.6$ Hz, 2H), 8.33 – 8.18 (m, 1H), 7.95 – 7.71 (m, 3H), 7.65 (t, $J = 1.9$ Hz, 1H), 7.09 (t, $J = 1.9$ Hz, 1H), 3.07 (s, 3H).
	53	474	^1H NMR (300 MHz, DMSO- d_6) δ 10.18 (s, 1H), 10.07 (s, 1H), 8.76 (s, 1H), 8.34 (s, 1H), 8.02 (d, $J = 1.7$ Hz, 1H), 7.91 (dd, $J = 7.9, 1.8$ Hz, 1H), 7.83 (d, $J = 1.9$


			Hz, 1H), 7.72 – 7.60 (m, 2H), 7.08 (t, J = 1.9 Hz, 1H), 3.07 (s, 3H), 2.35 (s, 3H).
	54	406	¹ H NMR (300 MHz, Methanol-d ₄) δ 8.87 (d, J = 0.8 Hz, 1H), 8.40 (d, J = 4.8 Hz, 1H), 8.29 (d, J = 0.8 Hz, 1H), 7.99 – 7.84 (m, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.61 (t, J = 1.9 Hz, 1H), 7.45 (dd, J = 7.6, 4.7 Hz, 1H), 7.04 (t, J = 2.0 Hz, 1H), 3.06 (s, 3H), 2.51 (s, 3H).
	55	521	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.24 (s, 1H), 10.06 (s, 1H), 9.28 (d, J = 0.8 Hz, 1H), 8.32 – 8.14 (m, 2H), 7.91 – 7.76 (m, 2H), 7.74 – 7.53 (m, 2H), 7.08 (t, J = 1.8 Hz, 1H), 3.78 (t, J = 4.7 Hz, 4H), 3.25 (t, J = 4.9 Hz, 4H), 3.07 (s, 3H).
	56	486	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 10.07 (s, 1H), 9.39 (d, J = 0.8 Hz, 1H), 8.35 (d, J = 0.8 Hz, 1H), 8.32 – 8.22 (m, 1H), 8.17 (dd, J = 8.3, 1.0 Hz, 1H), 7.86 (t, J = 1.8 Hz, 1H), 7.76 (d, J = 7.4 Hz, 1H), 7.69 (t, J = 1.9 Hz, 1H), 7.31 – 6.86 (m, 2H), 3.08 (s, 3H).
	57	502	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.34 (s, 1H), 10.08 (s, 1H), 9.31 (s, 1H), 8.39 (s, 1H), 8.28 – 7.97 (m, 2H), 7.89 – 7.72 (m, 2H), 7.68 (d, J = 1.9 Hz, 1H), 7.19 – 7.01 (m, 2H), 3.07 (s, 3H).

	58	463	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.24 (s, 1H), 10.08 (s, 1H), 9.97 (s, 1H), 9.02 (s, 1H), 8.40 (s, 1H), 7.94 (dd, J = 7.7, 1.5 Hz, 1H), 7.91 – 7.77 (m, 3H), 7.72 – 7.63 (m, 2H), 7.09 (t, J = 1.9 Hz, 1H), 3.08 (s, 3H).
	59	406	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.03 (s, 2H), 8.89 (d, J = 4.8 Hz, 2H), 8.51 (d, J = 2.1 Hz, 1H), 7.72 (t, J = 1.8 Hz, 1H), 7.66 (t, J = 1.8 Hz, 1H), 7.48 (t, J = 4.9 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H), 6.58 (dd, J = 1.8, 1.0 Hz, 1H), 3.06 (s, 3H), 2.61 (d, J = 1.0 Hz, 3H).
	60	441	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.60 (s, 1H), 10.08 (s, 1H), 8.53 (d, J = 1.5 Hz, 1H), 8.40 (d, J = 2.5 Hz, 1H), 8.06 (d, J = 1.5 Hz, 1H), 7.82 (t, J = 1.8 Hz, 1H), 7.78 – 7.62 (m, 2H), 7.10 (t, J = 1.9 Hz, 1H), 6.71 – 6.31 (m, 1H), 3.07 (s, 3H).
	388	478.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.24 (s, 1H), 10.07 (s, 1H), 9.18 (s, 1H), 8.72 (d, J = 1.3 Hz, 1H), 8.36 – 8.09 (m, 2H), 7.70 (t, J = 1.9 Hz, 1H), 7.61 (t, J = 1.9 Hz, 1H), 6.93 (t, J = 2.0 Hz, 1H), 3.75 (dd, J = 5.9, 3.8 Hz, 4H), 3.60 (t, J = 4.9 Hz, 4H), 3.06 (s, 3H).

	389	438.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.88 (s, 1H), 8.64 (d, J = 2.6 Hz, 1H), 8.16 (dd, J = 8.7, 2.6 Hz, 1H), 7.99 (d, J = 2.0 Hz, 1H), 7.73 – 7.65 (m, 2H), 7.62 (t, J = 1.9 Hz, 1H), 6.90 (t, J = 2.0 Hz, 1H), 6.59 (dd, J = 2.1, 1.1 Hz, 1H), 3.05 (s, 3H), 2.41 – 2.36 (m, 3H).
	390	422.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.92 (d, J = 57.4 Hz, 2H), 8.59 (d, J = 3.0 Hz, 1H), 8.04 – 7.90 (m, 2H), 7.76 – 7.68 (m, 2H), 7.63 (t, J = 2.0 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H), 6.58 (s, 1H), 3.06 (s, 3H), 2.35 (s, 3H).
	391	435	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.81 (s, 1H), 8.28 (d, J = 3.0 Hz, 1H), 7.86 (d, J = 2.0 Hz, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.66 – 7.52 (m, 3H), 6.88 (t, J = 2.0 Hz, 1H), 6.54 (t, J = 1.6 Hz, 1H), 3.90 (s, 3H), 3.04 (s, 3H), 2.31 (d, J = 1.0 Hz, 3H).
	392	487	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.20 (s, 1H), 9.07 (s, 1H), 8.68 (dd, J = 4.8, 1.7 Hz, 1H), 8.55 (d, J = 2.8 Hz, 1H), 8.27 (t, J = 1.8 Hz, 1H), 8.18 (dd, J = 7.7, 1.7 Hz, 1H), 7.99 (s, 1H), 7.78 – 7.64 (m, 2H), 7.59 (t, J = 1.9 Hz, 1H), 7.47 (t, J = 1.9 Hz, 1H), 6.87 (t, J = 2.0 Hz, 1H), 2.96 (s, 3H).

	393	453	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.99 (s, 1H), 9.78 (s, 1H), 8.19 (d, J = 2.5 Hz, 1H), 7.86 (dd, J = 10.3, 2.4 Hz, 1H), 7.71 (d, J = 1.9 Hz, 1H), 7.66 – 7.59 (m, 2H), 6.89 (t, J = 2.0 Hz, 1H), 6.53 (d, J = 1.7 Hz, 1H), 3.90 (s, 3H), 3.06 (s, 3H), 2.07 (d, J = 1.0 Hz, 3H).
---	-----	-----	--

Example 61: N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(pyridin-3-ylmethoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide



- 5 Step 1: To a mixture of 3-(bromomethyl)pyridine (520 mg, 3.02 mmol), 2-bromopyridin-3-ol (629 mg, 3.62 mmol), K₂CO₃ (1.25 g, 9.06 mmol) was added DMF (5 mL). The resulting mixture was stirred for 2 hours at 60 °C under N₂ atmosphere. The reaction mixture was extracted with ethyl acetate. The organic phase was concentrated under reduced pressure. The residue was purified by flash chromatography on silica gel eluting with 15% of ethyl acetate
- 10 in petroleum ether to afford 2-bromo-3-[(pyridin-3-yl)methoxy]pyridine (550 mg, 68.6%) as a white solid. LCMS [M + H]⁺: 265

- Step 2: To a mixture of 2-bromo-3-[(pyridin-3-yl)methoxy]pyridine (550 mg, 2.07 mmol), (1S,2S)-N1,N2-di methylcyclohexane-1,2-diamine (117 mg, 0.828 mmol), ethyl 1H-pyrazole-4-carboxylate (434 mg, 3.10 mmol), CuI (78.8 mg, 0.414 mmol) and K₂CO₃ (855
- 15 mg, 6.20 mmol) was added ACN (6 mL). The resulting mixture was stirred for 2 hours at 90 °C under N₂ atmosphere. The reaction mixture was extracted with ethyl acetate. The




organic phase was concentrated under reduced pressure. The residue was purified by flash chromatography on silica gel eluting with 95% of ethyl acetate in petroleum ether to afford ethyl 1-{3-[(pyridin-3-yl)methoxy]pyridin-2-yl}-1H-pyrazole-4-carboxylate (450 mg, 66.9%) as a yellow solid. LCMS [M + H]⁺: 325





- 5 Step 3: To a stirred solution of ethyl 1-{3-[(pyridin-3-yl)methoxy]pyridin-2-yl}-1H-pyrazole-4-carboxylate (450 mg, 1.38 mmol) in 4 mL of EtOH and 4 mL of water was added LiOH (289 mg, 6.89 mmol). The resulting mixture was stirred for two hours at room temperature. Then it was concentrated, and the PH value of the residue solution was adjusted to 3 with 1M HCl(aq). The product was precipitated from the solution. After filtration, the filter cake was
- 10 washed with water and dried under vacuum. This resulted in 1-{3-[(pyridin-3-yl)methoxy]pyridin-2-yl}-1H-pyrazole-4-carboxylic acid (260 mg, 63.2%) as a white solid. LCMS [M + H]⁺: 297



- Step 4: To a mixture of 1-{3-[(pyridin-3-yl)methoxy]pyridin-2-yl}-1H-pyrazole-4-carboxylic acid (120 mg, 0.405 mmol) and N-(3-amino-5-chlorophenyl)methanesulfonamide (107 mg, 0.485 mmol) in ACN (1 mL) was added TCFH (169 mg, 0.607 mmol) and NMI (99.2 mg, 1.21 mmol) at room temperature for two hours. Then it was concentrated, the residue was purified by reverse phase flash chromatography eluting with 65% of acetonitrile in water (0.1% NH₄HCO₃) to afford N-(3-chloro-5-methanesulfonamidophenyl)-1-{3-[(pyridin-3-yl)methoxy]pyridin-2-yl}-1H-pyrazole-4-carboxamide (27.0 mg, 13.3%) as a white solid.
- 20 LCMS [M + H]⁺: 499. ¹H NMR (300 MHz, DMSO-d₆) δ 10.21 (s, 1H), 10.08 (s, 1H), 8.91 (d, J = 0.7 Hz, 1H), 8.70 – 8.63 (m, 1H), 8.53 (dd, J = 4.8, 1.7 Hz, 1H), 8.30 (d, J = 0.7 Hz, 1H), 8.21 (dd, J = 4.6, 1.3 Hz, 1H), 7.96 – 7.81 (m, 2H), 7.70 (t, J = 1.9 Hz, 1H), 7.65 – 7.53 (m, 2H), 7.47 – 7.37 (m, 1H), 6.94 (t, J = 2.0 Hz, 1H), 5.36 (s, 2H), 3.07 (s, 3H).

Examples 62-70


The compounds listed in the following table were prepared using a procedure similar to that described for example 61:

Structure	Example No.	MS (ESI) [M + H] ⁺	¹ H NMR
	62	512	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.23 (s, 1H), 10.07 (s, 1H), 8.89 (s, 1H), 8.33 (s, 1H), 8.10 (d, J = 4.6 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.69 – 7.60 (m, 2H), 7.46 – 7.36 (m, 3H), 7.37 – 7.30 (m, 2H), 7.29 – 7.23 (m, 1H), 6.95 (t, J = 2.0 Hz, 1H), 3.08 (s, 3H), 1.53 (d, J = 6.3 Hz, 3H).
	63	499	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.23 (s, 1H), 10.07 (s, 1H), 8.98 (s, 1H), 8.58 (dt, J = 4.7, 1.5 Hz, 1H), 8.32 (s, 1H), 8.19 (dd, J = 4.6, 1.3 Hz, 1H), 7.93 – 7.79 (m, 2H), 7.72 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.59 – 7.49 (m, 2H), 7.34 (ddd, J = 7.6, 4.9, 1.2 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 5.39 (s, 2H), 3.08 (s, 3H).
	64	499	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.24 (s, 1H), 10.08 (s, 1H), 8.96 (d, J = 0.7 Hz, 1H), 8.73 – 8.46 (m, 2H), 8.34 (d, J = 0.7 Hz, 1H), 8.21 (dd, J = 4.6, 1.3 Hz, 1H), 7.83 (dd, J = 8.4, 1.4 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.57 (dd, J = 8.3, 4.7 Hz, 1H), 7.46 – 7.35 (m, 2H), 6.95 (t, J = 1.9 Hz, 1H), 5.39 (s, 2H), 3.08 (s, 3H).

	65	516	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.22 (s, 1H), 10.08 (s, 1H), 8.93 (d, J = 0.7 Hz, 1H), 8.32 (d, J = 0.7 Hz, 1H), 8.20 (dd, J = 4.6, 1.3 Hz, 1H), 7.86 (dd, J = 8.4, 1.4 Hz, 1H), 7.71 (t, J = 1.8 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.57 (dd, J = 8.3, 4.6 Hz, 1H), 7.43 (td, J = 8.0, 6.0 Hz, 1H), 7.29 (t, J = 6.8 Hz, 2H), 7.15 (td, J = 8.5, 2.1 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 5.34 (s, 2H), 3.08 (s, 3H).
	66	489	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.20 (s, 1H), 10.09 (s, 1H), 8.81 (s, 1H), 8.40 (s, 1H), 8.27 (s, 1H), 8.21 (dd, J = 4.6, 1.3 Hz, 1H), 7.98 (dd, J = 8.4, 1.4 Hz, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.62 – 7.54 (m, 2H), 7.35 (s, 1H), 6.94 (t, J = 2.0 Hz, 1H), 5.39 (s, 2H), 3.07 (s, 3H).
	67	566	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.19 (s, 1H), 10.06 (s, 1H), 8.87 (s, 1H), 8.35 (s, 1H), 8.16 (d, J = 2.1 Hz, 1H), 7.95 (dd, J = 8.7, 2.2 Hz, 1H), 7.81 (t, J = 1.8 Hz, 1H), 7.65 (d, J = 1.9 Hz, 1H), 7.56 (d, J = 8.8 Hz, 1H), 7.46 (d, J = 7.2 Hz, 2H), 7.43 – 7.27 (m, 3H), 7.08 (t, J = 1.9 Hz, 1H), 5.42 (s, 2H), 3.07 (s, 3H).
	68	498	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.21 (s, 1H), 10.08 (s, 1H), 8.90 (d, J = 0.7 Hz, 1H), 8.30 (d, J = 0.7 Hz, 1H), 8.18 (dd, J = 4.6, 1.3 Hz, 1H), 7.86 (dd, J = 8.4, 1.4 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.61 (t, J = 1.9 Hz, 1H), 7.55 (dd, J = 8.3, 4.7 Hz, 1H), 7.49 – 7.42 (m, 2H), 7.42 – 7.27 (m, 3H), 6.95 (t, J = 1.9 Hz, 1H), 5.31 (s, 2H), 3.08 (s, 3H).

	69	516	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.21 (s, 1H), 10.08 (s, 1H), 8.89 (s, 1H), 8.29 (s, 1H), 8.18 (dd, J = 4.6, 1.3 Hz, 1H), 7.86 (dd, J = 8.4, 1.3 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.61 (t, J = 1.9 Hz, 1H), 7.56 (dd, J = 8.3, 4.6 Hz, 1H), 7.53 – 7.43 (m, 2H), 7.30 – 7.14 (m, 2H), 6.94 (t, J = 2.0 Hz, 1H), 5.29 (s, 2H), 3.07 (s, 3H).
	70	532	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.21 (s, 1H), 10.05 (s, 1H), 8.93 (s, 1H), 8.30 (s, 1H), 8.20 (d, J = 4.6 Hz, 1H), 7.85 (d, J = 8.3 Hz, 1H), 7.70 (t, J = 1.8 Hz, 1H), 7.61 (t, J = 1.9 Hz, 1H), 7.57 (dd, J = 8.3, 4.7 Hz, 1H), 7.54 (s, 1H), 7.46 – 7.31 (m, 3H), 6.94 (t, J = 1.9 Hz, 1H), 5.32 (s, 2H), 3.07 (s, 3H).

Example 71: 4-bromo-N-(3-fluoro-5-(methylsulfonamido)phenyl)thiophene-2-carboxamide



Step 1: To a stirred solution of methyl 3-amino-5-fluorobenzoate (1 g, 5.91 mmol), HATU (2.92 g, 7.68 mmol) and DIEA (2.28 g, 17.7 mmol) in DMF (10 mL) was added 4-bromothiophene-2-carboxylic acid (1.34 g, 6.50 mmol). Then the mixture was stirred for 1 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford methyl 3-(4-bromothiophene-2-amido)-5-fluorobenzoate (1.28 g, 3.59 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 358

Step 2: To a stirred solution of methyl 3-(4-bromothiophene-2-amido)-5-fluorobenzoate (1.25 g, 3.48 mmol) in EtOH (10 mL) and H₂O (10 mL) was added LiOH (416 mg, 17.4 mmol). Then the mixture was stirred for 2 h at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 3-(4-bromothiophene-2-amido)-5-fluorobenzoic acid (1.02 g, 2.98 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 344

Step 3: To a stirred solution of 3-(4-bromothiophene-2-amido)-5-fluorobenzoic acid (1 g, 2.906 mmol, 1 equiv) and DPPA (1.04 g, 3.778 mmol, 1.3 equiv) in t-BuOH (20 mL) was added Et₃N (0.88 g, 8.718 mmol, 3 equiv) at 80 °C under nitrogen atmosphere for 6 h. The


resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with Petroleum ether / EtOAc (2:1) to afford tert-butyl N-[3-(4-bromothiophene-2-amido)-5-fluorophenyl]carbamate (750 mg, 61.53%) as a light yellow solid. LCMS (ESI) [M + H]⁺: 415







- 5 Step 4: To a stirred solution of tert-butyl N-[3-(4-bromothiophene-2-amido)-5-fluorophenyl]carbamate (700 mg, 1.68 mmol) in DCM (10 mL) was added TFA (1.91 g, 16.8 mmol) dropwise at 0 °C for 1 h. The resulting mixture was concentrated under vacuum. The mixture was basified to pH 9 with NaOH. The mixture was concentrated and purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (0-100%) to afford N-(3-amino-5-fluorophenyl)-4-bromothiophene-2-carboxamide (513 mg, 1.62 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 315
- 10

- Step 5: To a stirred solution of N-(3-amino-5-fluorophenyl)-4-bromothiophene-2-carboxamide (100 mg, 0.317 mmol) in Pyridine (5 mL) was added MsCl (39.8 mg, 0.348 mmol) dropwise at 0 °C. Then the mixture was stirred for 1 h at room temperature. The mixture was concentrated and purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (0-100%) to afford 4-bromo-N-(3-fluoro-5-methanesulfonamidophenyl)thiophene-2-carboxamide (25.3 mg, 0.0643 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 393. ¹H NMR (300 MHz, DMSO-d₆) δ 10.52 (s, 1H), 10.12 (s, 1H), 8.11 (d, J = 1.5 Hz, 1H), 8.05 (d, J = 1.4 Hz, 1H), 7.46 (dd, J = 10.4, 2.1 Hz, 2H), 6.75 (dt, J = 10.5, 2.2 Hz, 1H), 3.08 (s, 3H).
- 15
- 20

Examples 72-78


The compounds listed in the following table were prepared using a procedure similar to that described for example 71:

Structure	Example No.	MS (ESI) [M + H] ⁺	¹ H NMR
	72	407	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.51 (s, 1H), 10.15 (s, 1H), 8.11 (d, J = 1.5 Hz, 1H), 8.04 (d, J = 1.4 Hz, 1H), 7.45 (dt, J = 9.7, 2.2 Hz, 2H), 6.76 (dt, J = 10.5, 2.2 Hz, 1H), 3.19 (q, J = 7.3 Hz, 2H), 1.22 (t, J = 7.3 Hz, 3H).

	73	391	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.51 (s, 1H), 10.12 (s, 1H), 8.49 (d, J = 1.5 Hz, 1H), 8.22 (d, J = 1.4 Hz, 1H), 7.85 – 7.67 (m, 2H), 7.57 – 7.42 (m, 4H), 7.42 – 7.29 (m, 1H), 6.76 (dt, J = 10.5, 2.2 Hz, 1H), 3.09 (s, 3H).
	74	405	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.50 (s, 1H), 10.15 (s, 1H), 8.49 (d, J = 1.5 Hz, 1H), 8.22 (d, J = 1.4 Hz, 1H), 7.82 – 7.70 (m, 2H), 7.55 – 7.43 (m, 4H), 7.42 – 7.32 (m, 1H), 6.76 (dt, J = 10.4, 2.2 Hz, 1H), 3.20 (q, J = 7.3 Hz, 2H), 1.23 (t, J = 7.3 Hz, 3H).
	75	407	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.47 (s, 1H), 10.11 (s, 1H), 8.49 (d, J = 1.5 Hz, 1H), 8.23 (d, J = 1.4 Hz, 1H), 7.80 – 7.72 (m, 2H), 7.66 (dt, J = 12.1, 1.9 Hz, 2H), 7.49 (t, J = 7.6 Hz, 2H), 7.42 – 7.30 (m, 1H), 6.98 (t, J = 1.9 Hz, 1H), 3.08 (s, 3H).
	76	408	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.61 (s, 1H), 10.14 (s, 1H), 8.75 (d, J = 1.3 Hz, 1H), 8.64 (dt, J = 4.8, 1.5 Hz, 1H), 8.51 (d, J = 1.3 Hz, 1H), 7.96 – 7.86 (m, 2H), 7.72 – 7.66 (m, 2H), 7.36 (ddd, J = 6.2, 4.8, 2.5 Hz, 1H), 6.97 (t, J = 2.0 Hz, 1H), 3.18 (q, J = 7.3 Hz, 2H), 1.22 (t, J = 7.3 Hz, 3H).
	77	422	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.61 (s, 1H), 10.14 (s, 1H), 8.75 (d, J = 1.4 Hz, 1H), 8.65 (dt, J = 4.8, 1.4 Hz, 1H), 8.52 (d, J = 1.3 Hz, 1H), 7.97 – 7.85 (m, 2H), 7.69 (d, J = 2.0 Hz, 2H), 7.36 (td, J = 5.3, 2.9 Hz, 1H), 6.97 (t, J = 1.9 Hz, 1H), 3.18 (dd, J = 8.1, 6.6 Hz, 2H), 1.23 (t, J = 7.3 Hz, 3H).
	78	452	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.46 (s, 1H), 9.00 (d, J = 2.3 Hz, 1H), 8.60 – 8.50 (m, 2H), 8.38 (d, J = 1.4 Hz, 1H), 8.14 (dt, J = 8.0, 2.0 Hz, 1H), 7.78 (t, J = 1.8 Hz, 1H),

			7.66 (t, J = 1.9 Hz, 1H), 7.52 (dd, J = 8.0, 4.7 Hz, 1H), 7.09 (t, J = 1.9 Hz, 1H), 3.05 (s, 3H).
--	--	--	---

Example 79: N-(3-chloro-5-(methylsulfonamido)phenyl)-1-ethyl-5-phenyl-1H-pyrrole-3-carboxamide



- 5 Step 1: To a solution of methyl 1H-pyrrole-3-carboxylate (2.0 g, 15.9 mmol) in DMF (20 mL) was added NaH (791 mg, 19.8 mmol, 60% purity) at 0 °C. The mixture was stirred at 0 - 5°C for 15 min, MeI (3.37 g, 23.8 mmol) was added to the above mixture at 0-5°C. The mixture was stirred at 25°C for 1.5 h. The mixture was poured into water (20 mL) and extracted with EA (30 mL*3). The combined organic layers were washed with brine (15 mL*2) and dried over Na₂SO₄. The mixture was filtered, and the filtrate was concentrated in vacuo to give a crude. The crude product was pre-purified by column chromatography on silica gel eluted with PE:EA = 100:1 to 5:1. Methyl 1-ethyl-1H-pyrrole-3-carboxylate (1.2 g, 7.83 mmol, 98.3% yield) was obtained as yellow oil. LCMS: MS (ESI) Retention time: 0.654 min, (M+1) + = 154.0. ¹H NMR (400 MHz, CHLOROFORM-d) δ = 7.24 (t, J=2.0 Hz, 1 H) 6.52 - 6.56 (m, 1 H) 6.50 (dd, J=2.8, 1.6 Hz, 1 H) 3.86 (q, J=7.2 Hz, 2 H) 3.72 (s, 3 H) 1.37 (t, J=7.2 Hz, 3 H).
- 10
- 15

Step 2: To a solution of Methyl 1-ethyl-1H-pyrrole-3-carboxylate (1.2 g, 7.83 mmol) in THF (20 mL) was added NBS (1.43 g, 8.06 mmol) at 0 °C. The mixture was stirred at 25 °C for 1 h. The mixture was poured into water (25 mL), and was extracted with EtOAc (30 mL*3). The combined organic layers were washed with brine (20 mL*3), dried over Na₂SO₄, and concentrated in vacuum to give a residue. The residue was purified by silica gel column chromatography (Petroleum ether/Ethyl acetate = 10/1). Methyl 5-bromo-1-ethyl-1H-pyrrole-3-carboxylate (1.8 g, 7.75 mmol, 99.4% yield) was obtained as colorless oil. LCMS: MS (ESI) Retention time: 0.783 min, (M+1) + = 233.9. ¹H NMR (400 MHz, CHLOROFORM-d) δ = 7.37 (d, J=1.6 Hz, 1 H) 6.59 (d, J=1.6 Hz, 1 H) 3.96 (q, J=7.2 Hz, 2 H) 3.75 - 3.84 (m, 3 H) 1.40 (t, J=7.2 Hz, 3 H).

Step 3: To a solution of methyl 5-bromo-1-ethyl-1H-pyrrole-3-carboxylate (1.8 g, 7.75 mmol) and phenylboronic acid (1.2 g, 9.84 mmol) in MeOH (4 mL) and toluene (30 mL) was added tetrakis (triphenylphosphane) palladium (716 mg, 620 μmol) and 2.0 M aq. Solution of disodium carbonate (7.75 ul, 15.5 mmol). The mixture was stirred under a nitrogen atmosphere at 80 °C for 3 h. The mixture was cooled to 25 °C poured into water (20 mL) and extracted with EA (30 mL*3). The combined organic layers were washed with brine (15 mL*2) and dried over Na₂SO₄. The mixture was filtered, and the filtrate was concentrated in vacuo to give residue. The resulting oil was purified by flash chromatography eluting with 5% ethyl acetate/ hexanes. Compound 4 (1.1 g, 4.79 mmol, 62.1% yield) was obtained as colorless oil. LCMS: MS (ESI) Retention time: 0.830 min, (M+1) + = 230.0. ¹H NMR (400 MHz, CHLOROFORM-d) δ = 7.26 - 7.40 (m, 6 H) 6.50 - 6.52 (m, 1 H) 3.86 - 3.93 (m, 2 H) 3.73 - 3.79 (m, 3 H) 1.26 (t, J=7.2 Hz, 3 H).



Step 4: To a solution of methyl 1-ethyl-5-phenyl-1H-pyrrole-3-carboxylate (1 g, 3.53 mmol) in THF (10 mL), MeOH (10 mL) and H₂O (10 mL) was added LiOH · H₂O (545 mg, 13.0 mmol). The mixture was stirred at 25 °C for 2 h. The mixture was cooled to 25°C, and poured into water (20 mL). The pH of mixture was adjusted to 3 - 4 by 1 N HCl. And the mixture was extracted with EA (30 mL*3). The combined organic layers were washed with brine (15 mL*2) and dried over Na₂SO₄. The mixture was filtered, and the filtrate was concentrated in vacuo to give 1-ethyl-5-phenyl-1H-pyrrole-3-carboxylic acid (850 mg, 3.94 mmol, 90.6% yield) was obtained as white solid. LCMS: MS (ESI) Retention time: 0.766 min, (M+1) + = 216.0. ¹H NMR (400 MHz, METHANOL-d₄) δ = 7.51 (d, J=2.0 Hz, 1 H) 7.35 - 7.48 (m, 5 H) 6.42 - 6.55 (m, 1 H) 4.01 - 4.10 (m, 2 H) 1.28 (t, J=7.3 Hz, 3 H)


Step 5: To a solution of 1-ethyl-5-phenyl-1H-pyrrole-3-carboxylic acid (250 mg, 1.16 mmol) and N-(3-amino-5-chlorophenyl)methanesulfonamide (300 mg, 1.35 mmol) in pyridine (5

mL) was added EDCI (667 mg, 3.48 mmol). The mixture was stirred at 25 °C for 15 h. The mixture was partitioned between water (30 mL) and EtOAc (30 mL). The aqueous layer was separated and extracted with EtOAc (30 mL*2). The combined organic phases were washed with 1N HCl (20 mL*4) and brine (25 mL*2), dried over anhydrous Na₂SO₄, filtered, and the filtrate was evaporated in vacuo to give a residue. The residue was purified by Prep-HPLC (Column: Unisil 3 - 100 C18 Ultra 150*50 mm*3 um, mobile phase: [water (0.225%FA) - ACN]; B%: 36%-66%, 10 min) and lyophilized. N-(3-chloro-5-(methylsulfonamido)phenyl)-1-ethyl-5-phenyl-1H-pyrrole-3-carboxamide (286.74 mg, 98.48% Purity, 0.675 mmol, 58.2% yield) was obtained as off-white solid. LCMS: MS (ESI) Retention time: 0.920 min, (M+1) + = 417.9. ¹H NMR (400 MHz, DMSO-d₆) δ = 9.93 - 10.26 (m, 1 H) 9.77 (s, 1 H) 7.72 (d, J=1.6 Hz, 2 H) 7.58 - 7.66 (m, 1 H) 7.34 - 7.54 (m, 5 H) 6.88 (s, 1 H) 6.74 (d, J=1.6 Hz, 1 H) 4.02 (q, J=7.2 Hz, 2 H) 2.94 - 3.12 (m, 3 H) 1.25 (t, J=7.2 Hz, 3 H)


Examples 80-82

The compounds listed in the following table were prepared using a procedure similar to that described for example 79:

Structure	Example No.	MS (ESI) [M + H] ⁺	¹ H NMR
	80	472	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.91 (s, 1 H) 7.74 - 7.83 (m, 1 H) 7.65 - 7.71 (m, 1 H) 7.57 - 7.63 (m, 1 H) 7.37 - 7.54 (m, 5 H) 6.79 - 6.95 (m, 2 H) 4.86 - 5.08 (m, 2 H) 3.03 (s, 3 H).
	81	419	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.80 (s, 2H), 8.50 (br d, J = 4.6 Hz, 1H), 8.27 (s, 1H), 7.80 - 7.62 (m, 4H), 7.34 - 7.22 (m, 1H), 6.91 (dd, J = 1.6, 6.4 Hz, 2H), 3.69 (d, J = 0.8 Hz, 2H), 3.06 (d, J = 1.2 Hz, 3H), 2.40 (s, 3H)

	82	406	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.96 (s, 2H), 8.81 (d, J = 4.8 Hz, 2H), 7.79 - 7.71 (m, 3H), 7.68 (s, 1H), 7.30 (t, J = 4.8 Hz, 1H), 6.90 (s, 1H), 4.10 (s, 3H), 3.07 (s, 3H)
---	----	-----	--


Example 83: 5-bromo-N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-1H-pyrrole-3-carboxamide



- 5 Step 1: To a stirred solution of methyl 5-bromo-1H-pyrrole-3-carboxylate (500 mg, 2.45 mmol), CH₃I (521 mg, 3.67 mmol) and Cs₂CO₃ (2.38 g, 7.35 mmol) was added DMSO (10 mL). Then the mixture was stirred for 2 h at room temperature. The mixture was concentrated and purified by flash chromatography eluting with Petroleum ether/EtOAc (0-100%, acidic system) to afford methyl 5-bromo-1-methyl-1H-pyrrole-3-carboxylate (500 mg, 2.29 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 217.97
- 10 Step 2: To a stirred solution of methyl 5-bromo-1-methyl-1H-pyrrole-3-carboxylate (500 mg, 2.29 mmol) in EtOH (6 mL) and H₂O (2 mL) was added LiOH (164 mg, 6.86 mmol). Then the mixture was stirred for 2 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (0-100%, acidic system) to afford 5-bromo-1-methyl-1H-pyrrole-3-carboxylic acid (453 mg, 2.22 mmol) as an off-white solid. LCMS (ESI) [M + H]⁺: 203.96
- 15 Step 3: To a stirred solution of 5-bromo-1-methyl-1H-pyrrole-3-carboxylic acid (203 mg, 0.994 mmol), TCFH (417 mg, 1.49 mmol) and N-(3-amino-5-chlorophenyl)methanesulfonamide (219 mg, 0.993 mmol), NMI (407 mg, 4.97 mmol) was

added ACN (10 mL). Then the mixture was stirred for 2 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (0-100%, acidic system) to afford 5-bromo- N-(3-chloro-5-methanesulfonamidophenyl)-1-methyl-1H-pyrrole-3-carboxamide (31.9 mg, 0.0782 mmol) as an off-white solid. LCMS (ESI) [M + H]⁺: 405.95. ¹H NMR (300 MHz, DMSO-d₆) δ 10.04 (s, 1H), 9.78 (s, 1H), 7.72 – 7.65 (m, 2H), 7.59 (t, J = 1.9 Hz, 1H), 6.89 (t, J = 2.0 Hz, 1H), 6.83 (d, J = 2.1 Hz, 1H), 3.63 (s, 3H), 3.05 (s, 3H).

Example 84: N-(3-bromo-5-(methylsulfonamido)phenyl)-1-methyl-5-(3-methylpyridin-2-yl)-1H-pyrrole-3-carboxamide



10

Step 1: To a stirred solution of methyl 5-bromo-1H-pyrrole-3-carboxylate (450 mg, 2.20 mmol) and iodomethane (374 mg, 2.64 mmol) in DMSO (10 mL). Then the mixture was stirred for 2h at room temperature. The mixture was concentrated and purified by reverse flash chromatography eluting with ACN/H₂O (5-95%,acidic system) to afford methyl 5-bromo-1-methyl-1H-pyrrole-3-carboxylate (380 mg, 1.74 mmol) as a white solid.

Step 2: To a stirred solution of methyl 5-bromo-1-methyl-1H-pyrrole-3-carboxylate (380 mg, 1.74 mmol) and 3-methyl-2-(tributylstannyl)pyridine (1.99 g, 5.22 mmol) in DMSO (5 mL) was added CuI (66.2 mg, 348 μmol) , Pd(PPh₃)₄ (402 mg, 348 μmol) , PPh₃ (684 mg, 2.61 mmol) and LiCl (95.8 mg, 2.26 mmol) at 100°C for 12h .The reaction mixture was then cooled to RT, diluted with ethyl acetate (75 ml), then transferred to a separatory funnel, and washed with 10% aqueous ammonium chloride. The aqueous phase was extracted with ethyl acetate (3 times 60 ml). The combined organic phases were washed with water (3 times 75 ml), brine, and dried over magnesium sulfate.The mixture was concentrated and purified by

20


flash chromatography on silica gel eluting with EA/PE (0-100%) to afford methyl 1-methyl-5-(3-methylpyridin-2-yl)-1H-pyrrole-3-carboxylate (346 mg, 1.50 mmol) as a white solid.

Step 3: To a stirred solution of methyl 1-methyl-5-(3-methylpyridin-2-yl)-1H-pyrrole-3-carboxylate (200 mg, 868 μ mol) in EtOH (3 mL) and H₂O (3 mL) was added LiOH (207

5 mg, 8.68 mmol). Then the mixture was stirred for 2h at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 1-methyl-5-(3-methylpyridin-2-yl)-1H-pyrrole-3-carboxylic acid (150 mg, 693 μ mol) as a white solid.

Step 4: To a stirred solution of 1-methyl-5-(3-methylpyridin-2-yl)-1H-pyrrole-3-carboxylic acid (100 mg, 462 μ mol), TCFH (168 mg, 600 μ mol) and NMI (113 mg, 1.38 mmol) in ACN (5 mL) was added N-(3-amino-5-bromophenyl)methanesulfonamide (134 mg, 508 μ mol). Then the mixture was stirred for 1h at room temperature. The mixture was concentrated and purified by reverse flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford N-(3-bromo-5-methanesulfonamidophenyl)-1-methyl-5-(3-methylpyridin-2-yl)-1H-pyrrole-3-carboxamide (24.8 mg, 53.6 μ mol) as a white solid. LCMS (ESI): [M + H]⁺: 463. ¹H NMR (300 MHz, DMSO-d₆) δ 9.99 (s, 1H), 9.78 (s, 1H), 8.56 – 8.44 (m, 1H), 7.84 (t, J = 1.8 Hz, 1H), 7.77 (d, J = 7.8 Hz, 1H), 7.72 – 7.64 (m, 2H), 7.29 (dd, J = 7.7, 4.7 Hz, 1H), 7.03 (t, J = 1.9 Hz, 1H), 6.91 (d, J = 1.9 Hz, 1H), 3.69 (s, 3H), 3.06 (s, 3H), 2.40 (s, 3H).

Example 85: N-(3-bromo-5-(methylsulfonamido)phenyl)-1-cyclohexyl-5-methyl-1H-pyrrole-3-carboxamide



Step 1: To a mixture of methyl 5-methyl-1H-pyrrole-3-carboxylate (210 mg, 1.50 mmol), t-BuOK (252 mg, 2.25 mmol), 3-bromocyclohex-1-ene (288 mg, 1.79 mmol) was added NMP (2 mL). The resulting mixture was stirred for 2 hours at room temperature under N₂ atmosphere. The reaction mixture was extracted with ethyl acetate. The organic phase was concentrated under reduced pressure. The residue was purified by flash chromatography on silica gel eluting with 30% of ethyl acetate in petroleum ether to afford methyl 1-(cyclohex-2-en-1-yl)-5-methyl-1H-pyrrole-3-carboxylate (175 mg, 52.8%) as a light yellow solid.


LCMS [M + H]⁺: 220

Step 2: To a mixture of methyl 1-(cyclohex-2-en-1-yl)-5-methyl-1H-pyrrole-3-carboxylate (175 mg, 0.798 mmol), Pd/C (84.9 mg, 0.798 mmol) was added EtOAc (2 mL). The resulting mixture was stirred for 2 hours at room temperature under H₂ atmosphere. The reaction mixture was filtered. Then the reaction mixture was concentrated under reduced pressure. The residue was purified by flash chromatography on silica gel eluting with 30% of ethyl acetate in petroleum ether to afford methyl 1-cyclohexyl-5-methyl-1H-pyrrole-3-carboxylate (145 mg, 82.1%) as a yellow solid. LCMS [M + H]⁺: 222

Step 3: To a stirred solution of methyl 1-cyclohexyl-5-methyl-1H-pyrrole-3-carboxylate (145 mg, 0.656 mmol) in 2 mL of THF and 2 mL of water was added LiOH (137.7 mg, 3.27 mmol). The resulting mixture was stirred overnight at 50 °C. Then it was concentrated, and the PH value of the residue solution was adjusted to 3 with 1M HCl (aq). The product was precipitated from the solution. After filtration, the filter cake was washed with water (2 mL) and dried under vacuum. This resulted in 1-cyclohexyl-5-methyl-1H-pyrrole-3-carboxylic acid (60 mg, 43.9%) as a white solid. LCMS [M + H]⁺: 207

Step 4: To a mixture of 1-cyclohexyl-5-methyl-1H-pyrrole-3-carboxylic acid (60 mg, 0.289 mmol) and N-(3-amino-5-bromophenyl)methanesulfonamide (91.7 mg, 0.346 mmol) in ACN (1 mL) was added TCFH (121 mg, 0.433 mmol) and NMI (71.0 mg, 0.867 mmol) at room temperature. The resulting mixture was stirred for two hours, Then it was concentrated, the residue was purified by reverse phase flash chromatography eluting with 93% of acetonitrile in water (0.1%FA) to afford N-(3-bromo-5-methanesulfonamidophenyl)-1-cyclohexyl-5-methyl-1H-pyrrole-3-carboxamide (45.6 mg, 34.6%) as a white solid. LCMS [M + H]⁺: 454. ¹H NMR (300 MHz, DMSO-d₆) δ 9.96 (s, 1H), 9.56 (s, 1H), 7.81 (t, J = 1.8 Hz, 1H), 7.66 (t, J = 1.9 Hz, 1H), 7.57 (d, J = 2.0 Hz, 1H), 7.00 (t, J = 1.9 Hz, 1H), 6.34 (s, 1H), 3.87 (s, 1H), 3.05 (s, 3H), 2.26 – 2.13 (m, 3H), 1.88 (dd, J = 25.6, 12.1 Hz, 4H), 1.51 (dd, J = 46.0, 12.5 Hz, 6H).

Example 86: 1-(5-(4-acetylpiperazin-1-yl)pyridin-2-yl)-N-(3-bromo-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide



Step 1: A solution of 5-bromo-2-fluoropyridine (200 mg, 1.136 mmol, 1 equiv) and ethyl 1H-pyrazole-4-carboxylate (238.9 mg, 1.705 mmol, 1.50 equiv) in DMSO (5 mL) was added Cs₂CO₃ (1110.8 mg, 3.409 mmol, 3.00 equiv) at room temperature. The resulting solution was stirred for two hours at 60 °C. The reaction was quenched with water. The reaction mixture was diluted with water (50 mL). The resulting solution was extracted with EtOAc (50*3 mL) and the organic layers were combined. The organic layer was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by flash chromatography on silica gel eluting with EtOAc /Petroleum ether (42%) to afford ethyl 1-(5-bromopyridin-2-yl)pyrazole-4-carboxylate (300 mg, 84.69%) as a yellow solid LCMS (ESI): [M + H]⁺:296

Step 2: To a solution of ethyl 1-(5-bromopyridin-2-yl) pyrazole-4-carboxylate (200 mg, 0.675 mmol, 1 equiv), 1-(piperazin-1-yl) ethanone (129.8 mg, 1.013 mmol, 1.50 equiv), Pd(AcO)₂ (15.2 mg, 0.068 mmol, 0.10 equiv) and BINAP (84.1 mg, 0.135 mmol, 0.20 equiv) in toluene (5 mL) was added Cs₂CO₃ (660.2 mg, 2.026 mmol, 3.00 equiv) at room temperature under nitrogen. The resulting solution was stirred for two hours at 100 °C. The resulting solution




was concentrated under vacuum. The residue was purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (45%) to afford ethyl 1-[5-(4-acetylpiperazin-1-yl)pyridin-2-yl]pyrazole-4-carboxylate (130 mg, 53.81%) as a yellow solid. LCMS (ESI): [M + H]⁺:344




- 5 Step 3: To a solution of ethyl 1-[5-(4-acetylpiperazin-1-yl)pyridin-2-yl]pyrazole-4-carboxylate (135 mg, 0.393 mmol, 1 equiv) in THF (2 mL) and H₂O (2 mL) was added LiOH (47.1 mg, 1.967 mmol, 5.00 equiv) at room temperature. The resulting solution was stirred for one hour at room temperature. The residue was purified by flash chromatography on C18 column gel eluting with ACN/water(0.1%FA) (35%) to afford 1-[5-(4-acetylpiperazin-1-yl)pyridin-2-yl]pyrazole-4-carboxylic acid (95 mg, 70.50%) as a white solid. LCMS (ESI): [M + H]⁺:316




- Step 4: To a solution of 1-[5-(4-acetylpiperazin-1-yl)pyridin-2-yl]pyrazole-4-carboxylic acid (95 mg, 0.301 mmol, 1 equiv), NMI (49.5 mg, 0.603 mmol, 2.00 equiv) and TCFH (126.8 mg, 0.452 mmol, 1.50 equiv) in ACN (2 mL) at room temperature. The resulting solution was stirred for 10 minutes at room temperature. Then N-(3-amino-5-bromophenyl) methanesulfonamide (119.8 mg, 0.452 mmol, 1.50 equiv) was added and stirred for one hour at room temperature. The residue was purified by Pre-HPLC on condition: Column: Sunfire prep C18 column, 30*150 mm, 5μm; Mobile Phase A: Water (0.1%FA), Mobile Phase B: ACN; Flow rate: 60 mL/min; Gradient: 33% B to 47% B in 7 min, 47% B; Wave Length: 254/220 nm; RT1(min): 6.28 to afford 1-[5-(4-acetylpiperazin-1-yl)pyridin-2-yl]-N-(3-bromo-5-methanesulfonamidophenyl)pyrazole-4-carboxamide (19.7 mg, 11.37%) as a white solid. LCMS (ESI): [M + H]⁺:562. ¹H NMR (300 MHz, DMSO-d₆) δ 10.24 (s, 1H), 10.06 (s, 1H), 9.28 (d, J = 0.8 Hz, 1H), 8.33 – 8.16 (m, 2H), 7.96 – 7.78 (m, 2H), 7.76 – 7.54 (m, 2H), 7.08 (t, J = 1.9 Hz, 1H), 3.62 (t, J = 5.2 Hz, 4H), 3.25 (t, J = 5.3 Hz, 4H), 3.07 (s, 3H), 2.07 (s, 3H).




Example 87-94 and 394-423




The compounds listed in the following table were prepared using a procedure similar to that described for example 86:




Structure	Example No.	MS (ESI) [M+H] ⁺	¹ H NMR
	87	576	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.16 (s, 1H), 10.05 (s, 1H), 8.87 (s, 1H), 8.21 (s, 1H), 8.10 (d, J = 2.8 Hz, 1H), 7.80 (t, J = 1.8 Hz, 1H), 7.63 (t, J = 1.9 Hz, 1H), 7.48 (d, J = 2.8 Hz, 1H), 7.05 (t, J = 1.9 Hz, 1H), 3.62 (t, J = 5.3 Hz, 4H), 3.27 (t, J = 5.2 Hz, 4H), 3.04 (s, 3H), 2.37 (s, 3H), 2.07 (s, 3H).
	88	548	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.22 (s, 1H), 9.28 (d, J = 0.8 Hz, 1H), 8.38 – 8.12 (m, 2H), 7.95 – 7.73 (m, 2H), 7.63 (dd, J = 9.1, 2.8 Hz, 2H), 7.05 (t, J = 1.9 Hz, 1H), 3.92 (s, 2H), 3.67 – 3.54 (m, 2H), 3.48 (dd, J = 6.6, 4.3 Hz, 2H), 3.04 (s, 3H), 2.93 (s, 3H).
	89	547	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.26 (s, 1H), 10.08 (s, 1H), 9.29 (d, J = 0.8 Hz, 1H), 8.23 (s, 2H), 7.84 (d, J = 9.0 Hz, 1H), 7.73 (t, J = 1.8 Hz, 1H), 7.68 – 7.60 (m, 2H), 6.94 (t, J = 2.0 Hz, 1H), 3.28




			(s, 8H), 3.08 (s, 3H), 2.80 (s, 6H).
	90	475	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.31 (s, 1H), 10.09 (s, 1H), 9.37 (s, 1H), 8.84 (d, J = 2.6 Hz, 1H), 8.39 (dd, J = 9.0, 2.7 Hz, 1H), 8.29 (s, 1H), 8.01 (d, J = 9.0 Hz, 1H), 7.73 (t, J = 1.9 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 3.94 (t, J = 7.0 Hz, 2H), 3.08 (s, 3H), 2.60 – 2.52 (m, 2H), 2.15 (q, J = 7.5 Hz, 2H).
	91	477	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.27 (s, 1H), 10.08 (s, 1H), 9.37 (s, 1H), 8.25 (s, 1H), 8.15 (d, J = 5.9 Hz, 1H), 7.73 (t, J = 1.9 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 7.36 (d, J = 2.3 Hz, 1H), 6.93 (dt, J = 6.2, 2.2 Hz, 2H), 3.75 (t, J = 4.7 Hz, 4H), 3.40 (t, J = 4.9 Hz, 4H), 3.08 (s, 3H).
	92	491	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.26 (s, 1H), 10.08 (s, 1H), 9.28 (s, 1H), 8.28 – 8.17 (m, 2H), 7.84 (d, J = 9.1 Hz, 1H), 7.72 (t, J = 1.8 Hz, 1H), 7.67 – 7.57 (m, 2H), 6.94 (t, J = 1.9 Hz, 1H), 3.96 (d, J = 11.7 Hz, 1H), 3.69 (q, J = 13.0, 11.4 Hz, 4H), 3.07 (s, 3H), 2.82 – 2.72 (m, 1H), 2.48 –




			2.41 (m, 1H), 1.18 (d, J = 6.1 Hz, 3H).
	93	490	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.82 (s, 1H), 8.26 (d, J = 2.9 Hz, 1H), 7.84 (d, J = 2.0 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 7.56 (dd, J = 8.9, 3.0 Hz, 1H), 7.46 (d, J = 8.8 Hz, 1H), 6.89 (t, J = 1.9 Hz, 1H), 6.54 (dd, J = 2.1, 1.1 Hz, 1H), 3.78 (t, J = 4.9 Hz, 4H), 3.25 (t, J = 4.9 Hz, 4H), 3.06 (s, 3H), 2.30 (s, 3H).
	94	517	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.82 (s, 1H), 8.25 (d, J = 2.9 Hz, 1H), 7.84 (d, J = 2.0 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 7.57 (dd, J = 9.1, 3.0 Hz, 1H), 7.46 (d, J = 8.8 Hz, 1H), 6.89 (t, J = 1.9 Hz, 1H), 6.54 (s, 1H), 3.92 (s, 2H), 3.63 (t, J = 5.4 Hz, 2H), 3.48 (t, J = 5.5 Hz, 2H), 3.06 (s, 3H), 2.93 (s, 3H), 2.30 (s, 3H).
	394	477.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.27 (s, 1H), 10.12 (s, 1H), 9.29 (s, 1H), 8.60 (s, 2H), 8.24 (s, 1H), 7.66 (dt, J = 34.4, 1.9 Hz, 2H), 6.94 (d, J = 2.0 Hz, 1H), 3.79 (t, J =



			4.8 Hz, 4H), 3.33 (s, 4H), 3.06 (s, 3H).
	395	478.2	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.79 (s, 1H), 7.81 (dd, J = 7.7, 2.5 Hz, 2H), 7.71 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.42 (d, J = 8.6 Hz, 1H), 7.11 (dd, J = 8.7, 3.0 Hz, 1H), 6.88 (t, J = 1.9 Hz, 1H), 6.53 (d, J = 1.8 Hz, 1H), 5.84 – 5.32 (m, 1H), 4.29 (ddd, J = 21.0, 9.8, 5.6 Hz, 2H), 4.14 – 3.88 (m, 2H), 3.05 (s, 3H), 2.28 (d, J = 1.0 Hz, 3H).
	396	510	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.08 (s, 1H), 9.80 (s, 1H), 7.95 (d, J = 3.0 Hz, 1H), 7.85 – 7.58 (m, 3H), 7.44 (d, J = 8.7 Hz, 1H), 7.22 (dd, J = 8.9, 3.0 Hz, 1H), 6.88 (t, J = 2.0 Hz, 1H), 6.53 (s, 1H), 3.82 (t, J = 13.3 Hz, 2H), 3.59 (t, J = 7.2 Hz, 2H), 3.05 (s, 3H), 2.59 (dq, J = 14.4, 7.1 Hz, 2H), 2.27 (s, 3H).
	397	460	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.96 (s, 1H), 9.76 (s, 1H), 7.77 (d, J = 2.0 Hz, 1H), 7.73 (d, J = 2.9 Hz, 1H), 7.70 (s, 1H), 7.61 (s, 1H), 7.37 (d, J = 8.6 Hz, 1H), 7.00 (dd, J = 8.6, 2.9




			Hz, 1H), 6.87 (d, J = 2.0 Hz, 1H), 6.51 (d, J = 1.9 Hz, 1H), 3.94 (t, J = 7.3 Hz, 4H), 3.04 (s, 3H), 2.38 (p, J = 7.2 Hz, 2H), 2.26 (s, 3H).
	398	461	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.91 (s, 2H), 8.24 (d, J = 2.1 Hz, 1H), 8.09 (d, J = 2.0 Hz, 2H), 7.71 (t, J = 1.6 Hz, 1H), 7.65 (t, J = 1.7 Hz, 1H), 6.89 (t, J = 1.5 Hz, 1H), 6.51 (dd, J = 2.1, 1.1 Hz, 1H), 4.03 – 3.94 (m, 4H), 3.05 (s, 3H), 2.49 (s, 3H), 2.41 (p, J = 7.4 Hz, 2H).
	399	496.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.99 (s, 1H), 9.80 (s, 1H), 7.91 (d, J = 2.9 Hz, 1H), 7.81 (d, J = 2.0 Hz, 1H), 7.71 (d, J = 1.9 Hz, 1H), 7.63 (t, J = 2.0 Hz, 1H), 7.47 (d, J = 8.6 Hz, 1H), 7.21 (dd, J = 8.6, 3.0 Hz, 1H), 6.88 (d, J = 2.1 Hz, 1H), 6.53 (s, 1H), 4.43 (t, J = 12.3 Hz, 4H), 3.06 (s, 3H), 2.28 (s, 3H).
	400	468.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.92 (s, 1H), 9.80 (s, 1H), 8.27 (d, J = 2.1 Hz, 1H), 8.18 (d, J = 1.2 Hz, 2H), 7.67 (dd, J = 25.7, 1.9 Hz, 2H), 6.89 (t, J = 1.9 Hz, 1H), 6.52 (d, J = 2.1




			Hz, 1H), 5.54 (dt, J = 57.3, 2.9 Hz, 1H), 4.46 – 4.21 (m, 2H), 4.20 – 3.88 (m, 2H), 3.05 (d, J = 1.4 Hz, 3H), 2.51 (s, 3H).
	401	488.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.01 (s, 2H), 9.18 (s, 2H), 8.43 (d, J = 2.1 Hz, 1H), 7.69 (dt, J = 27.7, 1.9 Hz, 2H), 6.90 (t, J = 1.9 Hz, 1H), 6.60 – 6.54 (m, 1H), 3.94 (t, J = 7.1 Hz, 2H), 3.06 (s, 3H), 2.60 – 2.51 (m, 5H), 2.15 (p, J = 7.6 Hz, 2H).
	402	473.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.98 (s, 1H), 9.76 (s, 1H), 7.96 (d, J = 3.0 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.67 – 7.53 (m, 2H), 7.48 (d, J = 8.6 Hz, 1H), 7.07 – 6.75 (m, 3H), 3.90 (s, 3H), 3.29 (s, 4H), 3.06 (s, 3H), 2.02 – 1.94 (m, 4H).
	403	510.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.04 (s, 1H), 9.79 (s, 1H), 7.99 (d, J = 2.9 Hz, 1H), 7.71 (t, J = 2.0 Hz, 1H), 7.66 – 7.49 (m, 3H), 7.11 (dd, J = 8.8, 3.0 Hz, 1H), 7.03 (d, J = 2.0 Hz, 1H), 6.89 (t, J = 2.0 Hz, 1H), 4.40 (t, J = 12.3 Hz, 4H), 3.91 (s, 3H), 3.17 (q, J = 7.3




			Hz, 2H), 1.21 (t, J = 7.3 Hz, 3H).
	404	490.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.12 – 9.89 (m, 2H), 8.26 (d, J = 2.1 Hz, 1H), 8.13 (s, 2H), 7.68 (d, J = 26.7 Hz, 2H), 6.89 (t, J = 1.9 Hz, 1H), 6.55 – 6.49 (m, 1H), 4.42 – 4.35 (m, 1H), 4.21 (dd, J = 8.4, 6.2 Hz, 2H), 3.81 (dd, J = 8.5, 4.2 Hz, 2H), 3.30 (s, 3H), 3.26 (s, 3H), 3.05 (s, 3H).
	405	516	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.78 (s, 1H), 7.93 (d, J = 3.0 Hz, 1H), 7.78 (d, J = 2.1 Hz, 1H), 7.66 (dt, J = 34.9, 1.9 Hz, 2H), 7.39 (d, J = 8.7 Hz, 1H), 7.19 (dd, J = 8.9, 3.0 Hz, 1H), 6.88 (t, J = 2.0 Hz, 1H), 6.51 (dd, J = 2.1, 1.1 Hz, 1H), 3.87 (dd, J = 8.8, 6.5 Hz, 2H), 3.58 (dd, J = 8.8, 3.4 Hz, 2H), 3.45 (dd, J = 9.7, 7.2 Hz, 2H), 3.29 (dd, J = 10.2, 2.9 Hz, 2H), 3.05 (s, 5H), 2.29 – 2.24 (m, 3H).
	406	529.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.99 (s, 1H), 9.78 (s, 1H), 7.90 (d, J = 2.6 Hz, 1H), 7.70 (d, J = 2.2 Hz, 1H), 7.59 (d, J = 14.6 Hz, 2H), 7.42 (d, J = 2.5 Hz, 1H), 6.88 (d, J = 2.2 Hz,




			1H), 6.54 (s, 1H), 4.50 (t, J = 12.3 Hz, 4H), 3.05 (d, J = 1.7 Hz, 3H), 2.05 (d, J = 22.9 Hz, 3H).
	407	563.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.03 (s, 1H), 9.79 (d, J = 2.4 Hz, 1H), 8.17 (d, J = 2.9 Hz, 1H), 7.70 (q, J = 2.0 Hz, 1H), 7.63 (t, J = 2.3 Hz, 1H), 7.55 (s, 1H), 7.51 (d, J = 2.9 Hz, 1H), 6.90 (p, J = 1.9 Hz, 1H), 6.64 – 6.41 (m, 1H), 4.58 (t, J = 12.3 Hz, 4H), 3.06 (d, J = 1.6 Hz, 3H), 1.96 (s, 3H).
	408	528.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.81 (s, 1H), 7.84 (d, J = 2.4 Hz, 1H), 7.70 (d, J = 1.9 Hz, 1H), 7.65 – 7.55 (m, 2H), 7.26 (dd, J = 12.5, 2.5 Hz, 1H), 6.88 (d, J = 2.0 Hz, 1H), 6.55 (s, 1H), 3.86 (t, J = 13.1 Hz, 2H), 3.62 (t, J = 7.3 Hz, 2H), 3.05 (s, 3H), 2.70 – 2.56 (m, 2H), 2.10 (s, 3H).
	409	511.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.28 (d, J = 10.7 Hz, 3H), 7.69 (dq, J = 23.3, 2.2 Hz, 2H), 6.89 (t, J = 2.0 Hz, 1H), 6.53 (dd, J = 2.2, 1.2 Hz, 1H), 3.86 (t, J = 13.2 Hz, 2H), 3.63 (t, J = 7.3 Hz, 2H), 3.38 (s, 3H),

			3.06 (s, 3H), 2.59 (td, J = 14.5, 7.2 Hz, 2H).
	410	487.95	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.86 (s, 1H), 8.84 (d, J = 2.7 Hz, 1H), 8.36 (dd, J = 8.9, 2.8 Hz, 1H), 7.95 (d, J = 2.0 Hz, 1H), 7.75 – 7.59 (m, 3H), 6.89 (t, J = 1.9 Hz, 1H), 6.61 – 6.50 (m, 1H), 3.94 (t, J = 7.0 Hz, 2H), 3.05 (s, 3H), 2.57 (d, J = 7.9 Hz, 2H), 2.37 (d, J = 1.0 Hz, 3H), 2.13 (p, J = 7.5 Hz, 2H).
	411	486.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.03 (s, 1H), 9.78 (s, 1H), 7.86 (d, J = 3.0 Hz, 1H), 7.76 (d, J = 2.0 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.36 (d, J = 8.8 Hz, 1H), 7.12 (dd, J = 8.9, 3.0 Hz, 1H), 6.88 (t, J = 1.9 Hz, 1H), 6.51 (dd, J = 2.0, 1.1 Hz, 1H), 3.59 (d, J = 9.4 Hz, 2H), 3.33 – 3.24 (m, 2H), 3.05 (s, 3H), 2.26 (d, J = 1.0 Hz, 3H), 1.79 – 1.70 (m, 2H), 0.82 – 0.70 (m, 1H), 0.28 (d, J = 4.2 Hz, 1H).


	412	508.1	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.82 (s, 1H), 8.16 (d, J = 2.4 Hz, 1H), 7.80 – 7.43 (m, 4H), 6.89 (t, J = 2.0 Hz, 1H), 6.56 (dd, J = 1.9, 1.1 Hz, 1H), 3.77 (t, J = 4.9 Hz, 4H), 3.34 (s, 1H), 3.32 (s, 3H), 3.06 (s, 3H), 2.19 – 2.07 (m, 3H).
	413	537.2	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.19 (d, J = 2.1 Hz, 1H), 8.09 (s, 2H), 7.71 – 7.50 (m, 2H), 6.88 (s, 1H), 6.50 (s, 1H), 4.07 (s, 4H), 3.02 (s, 3H), 2.86 (t, J = 12.5 Hz, 5H), 2.46 (s, 3H).
	414	554.15	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.80 (s, 1H), 7.79 (d, J = 2.5 Hz, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.62 (dt, J = 8.4, 1.7 Hz, 2H), 7.17 (dd, J = 12.6, 2.5 Hz, 1H), 6.88 (t, J = 2.0 Hz, 1H), 6.54 (dd, J = 2.0, 1.1 Hz, 1H), 3.68 – 3.43 (m, 4H), 3.05 (s, 3H), 2.25 (dt, J = 13.2, 6.8 Hz, 1H), 2.09 (s, 4H), 1.74 – 1.62 (m, 2H).

	415	520.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.03 (s, 1H), 9.85 (s, 1H), 8.18 (d, J = 3.0 Hz, 1H), 7.77 (d, J = 3.0 Hz, 1H), 7.74 (d, J = 1.8 Hz, 1H), 7.70 (s, 1H), 7.62 (s, 1H), 6.90 (s, 1H), 6.62 (s, 1H), 4.52 (t, J = 12.2 Hz, 4H), 3.05 (s, 3H), 2.10 (d, J = 23.6 Hz, 3H).
	416	489.9	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.34 – 9.76 (m, 2H), 8.72 (d, J = 2.8 Hz, 1H), 8.26 (dd, J = 8.9, 2.8 Hz, 1H), 7.94 (d, J = 2.0 Hz, 1H), 7.76 – 7.56 (m, 3H), 6.89 (t, J = 2.0 Hz, 1H), 6.58 (dd, J = 2.0, 1.1 Hz, 1H), 4.52 (dd, J = 9.1, 6.8 Hz, 2H), 4.17 (dd, J = 9.1, 6.9 Hz, 2H), 3.06 (s, 3H), 2.39 – 2.33 (m, 3H).
	417	491.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.22 – 9.96 (m, 2H), 9.08 (s, 2H), 8.43 (d, J = 2.1 Hz, 1H), 7.73 (t, J = 1.9 Hz, 1H), 7.65 (t, J = 1.9 Hz, 1H), 6.91 (t, J = 1.9 Hz, 1H), 6.58 (dd, J = 2.1, 1.2 Hz, 1H), 4.56 (dd, J = 9.1, 6.8 Hz, 2H), 4.18 (dd, J = 9.1, 6.9 Hz, 2H), 3.06 (s, 3H), 2.59 (d, J = 1.0 Hz, 3H).

	418	497	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.96 (s, 2H), 8.30 (d, J = 2.1 Hz, 1H), 8.27 (s, 2H), 7.72 (t, J = 1.9 Hz, 1H), 7.66 (t, J = 1.9 Hz, 1H), 6.89 (t, J = 1.9 Hz, 1H), 6.53 (dd, J = 2.2, 1.1 Hz, 1H), 4.48 (t, J = 12.3 Hz, 4H), 3.06 (s, 3H), 2.52 (s, 3H).
	419	509.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.74 (s, 1H), 7.77 (d, J = 2.8 Hz, 1H), 7.69 (t, J = 1.9 Hz, 1H), 7.61 (t, J = 1.9 Hz, 1H), 7.51 (d, J = 2.0 Hz, 1H), 7.10 (d, J = 2.9 Hz, 1H), 6.88 (t, J = 1.9 Hz, 1H), 6.53 (t, J = 1.5 Hz, 1H), 4.43 (t, J = 12.3 Hz, 4H), 3.05 (s, 3H), 2.03 (s, 3H), 1.99 – 1.93 (m, 3H).
	420	513.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.81 (s, 1H), 7.79 (d, J = 2.5 Hz, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.64 (dt, J = 12.5, 1.8 Hz, 2H), 7.29 (dd, J = 11.7, 2.5 Hz, 1H), 6.89 (t, J = 2.0 Hz, 1H), 6.56 (d, J = 1.8 Hz, 1H), 4.48 (t, J = 12.3 Hz, 4H), 3.05 (s, 3H), 2.11 (s, 3H).

	421	526	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.98 (s, 1H), 9.73 (s, 1H), 7.66 (dd, J = 33.7, 2.1 Hz, 2H), 7.50 (dd, J = 19.8, 2.1 Hz, 2H), 6.88 (q, J = 2.3 Hz, 2H), 6.47 (d, J = 2.1 Hz, 1H), 4.46 (t, J = 12.3 Hz, 4H), 3.83 (s, 3H), 3.12 – 2.94 (m, 3H), 2.01 (s, 3H).
	422	554	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.79 (s, 1H), 7.98 (d, J = 2.9 Hz, 1H), 7.69 (t, J = 1.9 Hz, 1H), 7.64 – 7.51 (m, 3H), 7.09 (dd, J = 8.7, 2.9 Hz, 1H), 7.03 (d, J = 2.0 Hz, 1H), 6.88 (t, J = 2.0 Hz, 1H), 4.39 (t, J = 12.3 Hz, 4H), 3.91 (s, 3H), 3.76 (t, J = 6.6 Hz, 2H), 3.30 (t, J = 6.6 Hz, 2H).
	423	511	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.04 (s, 1H), 9.95 (s, 1H), 8.29 (d, J = 11.0 Hz, 3H), 7.75 – 7.55 (m, 2H), 6.90 (d, J = 2.0 Hz, 1H), 6.53 (s, 1H), 4.48 (t, J = 12.3 Hz, 4H), 3.16 (q, J = 7.3 Hz, 2H), 2.52 (s, 3H), 1.21 (t, J = 7.3 Hz, 3H).

Example 95: N-(3-bromo-5-(methylsulfonamido)phenyl)-1-methyl-2-(3-methylpyridin-2-yl)-1H-imidazole-4-carboxamide



Step 1: To a stirred solution of methyl 5-bromo-1H-pyrrole-3-carboxylate (600 mg, 2.94 mmol), K₂CO₃ (812 mg, 5.88 mmol) and 2-bromoacetonitrile (705 mg, 5.88 mmol) in DMSO (10 mL). Then the mixture was stirred for 4 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford methyl 5-bromo-1-(cyanomethyl)-1H-pyrrole-3-carboxylate (693 mg, 2.85 mmol) as a white solid.

- Step 2: A solution of methyl 5-bromo-1-(cyanomethyl)-1H-pyrrole-3-carboxylate (700 mg, 2.87 mmol), Pd(dppf)Cl₂ (420 mg, 0.574 mmol), KOAc (844 mg, 8.61 mmol) and B₂Pin₂ (3.63 g, 14.3 mmol) in dioxane (10 mL) was stirred for 2 h at 80 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with Petroleum ether / EtOAc (0-100%) to afford methyl 1-(cyanomethyl)-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrole-3-carboxylate (326 mg, 1.12 mmol) as a yellow solid.

Step 3: A solution of methyl 1-(cyanomethyl)-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrole-3-carboxylate (330 mg, 1.13 mmol), Pd(PPh₃)₂Cl₂ (158 mg, 0.226 mmol),


Na₂CO₃ (358 mg, 3.38 mmol) and 2-bromopyridine (213 mg, 1.35 mmol) in EtOH (4 mL) and H₂O (1 mL) was stirred for 2 h at 80 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with Petroleum ether / EtOAc (0-100%) to afford methyl 1-(cyanomethyl)-5-(pyridin-2-yl)-1H-pyrrole-3-carboxylate (148 mg, 0.615 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 242

Step 4: To a stirred solution of methyl 1-(cyanomethyl)-5-(pyridin-2-yl)-1H-pyrrole-3-carboxylate (150 mg, 0.621 mmol) in H₂O (5 mL) and EtOH (5 mL) was added LiOH (148 mg, 6.21 mmol). Then the mixture was stirred for 2 h at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 1-(cyanomethyl)-5-(pyridin-2-yl)-1H-pyrrole-3-carboxylic acid (108 mg, 0.479 mmol) as a yellow solid.

LCMS (ESI) [M + H]⁺: 228

Step 5: To a stirred solution of 1-(cyanomethyl)-5-(pyridin-2-yl)-1H-pyrrole-3-carboxylic acid (110 mg, 0.484 mmol), TCFH (176 mg, 0.629 mmol) and NMI (119 mg, 1.45 mmol) in ACN (5 mL) was added N-(3-amino-5-bromophenyl) methanesulfonamide (153 mg, 0.580 mmol). Then the mixture was stirred for 1 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford N-(3-bromo-5-methanesulfonamidophenyl)-1-(cyanomethyl)-5-(pyridin-2-yl)-1H-pyrrole-3-carboxamide (47.6 mg, 0.1004 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 474. ¹H NMR (300 MHz, DMSO-d₆) δ 10.09 (s, 1H), 10.01 (s, 1H), 8.62 (d, J = 4.8 Hz, 1H), 7.96 – 7.75 (m, 4H), 7.67 (t, J = 1.9 Hz, 1H), 7.42 (d, J = 1.9 Hz, 1H), 7.32 (ddd, J = 6.4, 5.0, 1.3 Hz, 1H), 7.05 (t, J = 1.9 Hz, 1H), 5.78 (s, 2H), 3.06 (s, 3H).


Example 96: N-(3-bromo-5-(methanesulfonamido)phenyl)-1-methyl-2-(3-methylpyridin-2-yl)-1H-imidazole-4-carboxamide



- Step 1: To a mixture of 2-bromo-3-methylpyridine (51 mg, 2.96 mmol), CuI (1.12 g, 5.92 mmol), Pd(AcO)₂ (132 mg, 0.592 mmol), PPh₃ (309 mg, 1.18 mmol), DBU (901 mg, 5.92 mmol), methyl 1-methyl-1H-imidazole-4-carboxylate (497 mg, 3.55 mmol) was added Dioxane (5 mL). The resulting mixture was stirred for 2 hours at 140 °C under N₂ atmosphere. The reaction mixture was concentrated under reduced pressure. The residue was purified by flash chromatography on silica gel eluting with 35% of ethyl acetate in petroleum ether to afford methyl 1-methyl-2-(3-methylpyridin-2-yl)-1H-imidazole-4-carboxylate (320 mg, 46.7%) as a white solid. LCMS [M + H]⁺: 232
- Step 2: To a stirred solution of methyl 1-methyl-2-(3-methylpyridin-2-yl)-1H-imidazole-4-carboxylate (150 mg, 0.648 mmol) in 2 mL of THF and 2 mL of water was added LiOH (135 mg, 3.23 mmol). The resulting mixture was stirred for two hours at room temperature. Then it was concentrated, and the PH value of the residue solution was adjusted to 3 with 1M HCl(aq). The product was precipitated from the solution. After filtration, the filter cake was washed with water and dried under vacuum. This resulted in 1-methyl-2-(3-methylpyridin-2-yl)-1H-imidazole-4-carboxylic acid (110 mg, 78.0%) as a white solid. LCMS [M + H]⁺: 218
- Step 3: To a mixture of 1-methyl-2-(3-methylpyridin-2-yl)-1H-imidazole-4-carboxylic acid (110 mg, 0.506 mmol) and N-(3-amino-5-bromophenyl)methanesulfonamide (160 mg, 0.607 mmol) in ACN (1 mL) was added TCFH (212 mg, 0.759 mmol) and NMI (123 mg, 1.51 mmol) at room temperature for two hours. Then it was concentrated, the residue was purified by reverse phase flash chromatography eluting with 60% of acetonitrile in water (0.1%

NH_4HCO_3) to afford N-(3-bromo-5-methanesulfonamidophenyl)-1-methyl-2-(3-methylpyridin-2-yl)-1H-imidazole-4-carboxamide (25.9 mg, 11.0%) as a light pink solid. LCMS $[\text{M} + \text{H}]^+$: 464¹H NMR (300 MHz, DMSO- d_6) δ 10.00 (s, 2H), 8.54 (d, J = 4.6 Hz, 1H), 8.03 (s, 1H), 7.91 – 7.76 (m, 3H), 7.43 (dd, J = 7.8, 4.8 Hz, 1H), 7.04 (s, 1H), 3.76 (s, 3H), 3.08 (s, 3H), 2.48 (s, 3H).

Example 97: N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(difluoro(phenyl)methoxy)pyridin-2-yl)-1H-pyrazole-4-carboxamide



Step 1: To a stirred solution of (difluoromethyl)benzene (1.2 g, 9.36 mmol) and NCS (1.36 g, 10.2 mmol) in CCl_4 (20 mL). The magnetically stirred mixture was exposed to sunlight during the daytime and a sunlamp at night for 50 h at room temperature. The resulting mixture was concentrated under reduced pressure at 0°C to afford (chlorodifluoromethyl)benzene (350 mg, 2.15 mmol, low boiling point) as a yellow oil.

Step 2: To a stirred solution of (chlorodifluoromethyl)benzene (350 mg, 2.15 mmol), Cs_2CO_3 (1.40 g, 4.30 mmol) and 2-bromopyridin-3-ol (410 mg, 2.36 mmol) in DMSO (5 mL). Then the mixture was stirred for 3 h at 120°C . The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/ H_2O (5-95%, acidic system) to afford

2-bromo-3-[difluoro(phenyl) methoxy]pyridine (50.0 mg, 0.166 mmol) as a yellow solid.

LCMS (ESI) $[M + H]^+$: 300

Step 3: A solution of 2-bromo-3-[difluoro(phenyl)methoxy]pyridine (50 mg, 0.166 mmol), CuI (6.32 mg, 0.0332 mmol), K_2CO_3 (68.8 mg, 0.498 mmol), ethyl 1H-pyrazole-4-

- 5 carboxylate (25.5 mg, 0.182 mmol) and (1S,2S)-N1, N2-dimethylcyclohexane-1,2-diamine (9.44 mg, 0.0664 mmol) in ACN (5 mL) was stirred for 2 h at 90 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with Petroleum ether / EtOAc (0-100%) to afford ethyl 1-{3-[difluoro(phenyl)methoxy]pyridin-2-yl}-1H-pyrazole-4-carboxylate
- 10 (40.0 mg, 0.111 mmol) as a yellow solid. LCMS (ESI) $[M + H]^+$: 360


Step 4: To a stirred solution of ethyl 1-{3-[difluoro(phenyl)methoxy]pyridin-2-yl}-1H-pyrazole-4-carboxylate (40 mg, 0.111 mmol) in EtOH (3 mL) and H_2O (3 mL) was added LiOH (26.5 mg, 1.11 mmol). Then the mixture was stirred for 2 h at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by

- 15 filtration and washed with water. The resulting solid was dried under vacuum to afford 1-{3-[difluoro(phenyl)methoxy]pyridin-2-yl}-1H-pyrazole-4-carboxylic acid (28.0 mg, 0.0845 mmol) as a yellow solid. LCMS (ESI) $[M + H]^+$: 332

Step 5: To a stirred solution of 1-{3-[difluoro(phenyl)methoxy]pyridin-2-yl}-1H-pyrazole-4-carboxylic acid (28 mg, 0.0845 mmol), TCFH (23.7 mg, 0.0845 mmol) and NMI (6.93 mg,

- 20 0.0845 mmol) in ACN (5 mL) was added N-(3-amino-5-chlorophenyl)methanesulfonamide (20.5 mg, 0.0929 mmol). Then the mixture was stirred for 1 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/ H_2O (5-95%, acidic system) to afford N-(3-chloro-5-methanesulfonamidophenyl)-1-{3-[difluoro(phenyl)methoxy]pyridin-2-yl}-1H-pyrazole-4-carboxamide (6.4 0 mg, 0.0119
- 25 mmol) as an orange solid. LCMS (ESI) $[M + H]^+$: 534. ^1H NMR (300 MHz, $DMSO-d_6$) δ 10.27 (s, 1H), 10.11 (s, 1H), 9.07 (s, 1H), 8.58 (dd, $J = 4.6, 1.5$ Hz, 1H), 8.34 (s, 1H), 8.24 – 8.15 (m, 1H), 7.79 – 7.66 (m, 4H), 7.66 – 7.50 (m, 4H), 6.94 (t, $J = 1.9$ Hz, 1H), 3.07 (s, 3H).


Example 98: N-(3-chloro-5-(methylsulfonamido)phenyl)-5-cyano-1-(pyridin-2-yl)-1H-pyrrole-3-carboxamide



- Step 1: Under nitrogen, a solution of methyl 1H-pyrrole-3-carboxylate (1 g, 7.992 mmol, 1 equiv), 2-bromo pyridine (1.89 g, 11.988 mmol, 1.5 equiv), CuI (304.4 mg, 1.598 mmol, 0.20 equiv) and (1S,2S)-N1, N2-dimethylcyclohexane-1,2-diamine (454.7 mg, 3.197 mmol, 0.40 equiv) in ACN (10 mL) was added K₂CO₃ (3.34 g, 23.992 mmol, 3.00 equiv) at room temperature. The resulting solution was stirred for two hours at 90 °C. The resulting solution was concentrated under vacuum. The residue was purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (26%) to afford methyl 1-(pyridin-2-yl) pyrrole-3-carboxylate (1.2 g, 68.69%) as a yellow solid. LCMS: (ESI): [M + H]⁺:203
- Step 2: To a solution of methyl 1-(pyridin-2-yl) pyrrole-3-carboxylate (850 mg, 4.203 mmol, 1 equiv) in THF (10 mL) was added NBS (897.79 mg, 5.044 mmol, 1.2 equiv) at room temperature. The resulting solution was stirred for two hours at room temperature. The residue was purified by flash chromatography on silica gel eluting with column gel eluting with EtOAc/Petroleum ether (28%) to afford methyl 5-bromo-1-(pyridin-2-yl) pyrrole-3-carboxylate (410 mg, 33.31%) as a yellow solid. LCMS: (ESI): [M + H]⁺:281

- Step 3: Under nitrogen, a solution of methyl 5-bromo-1-(pyridin-2-yl) pyrrole-3-carboxylate (410 mg, 1.459 mmol, 1 equiv) and CuCN (261.3 mg, 2.917 mmol, 2.00 equiv) in NMP (5 mL) was added KI (242.1 mg, 1.458 mmol, 1.00 equiv) at room temperature. The resulting solution was stirred for two hours at 120 °C. The reaction mixture was diluted with water (50 mL). The resulting solution was extracted with EtOAc (50×3 mL) and the organic layers were combined. The organic layer was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (35%) to afford methyl 5-cyano-1-(pyridin-2-yl) pyrrole-3-carboxylate (180 mg, 46.71%) as a yellow solid. LCMS: (ESI): [M + H]⁺:228
- Step 4: To a solution of methyl 5-cyano-1-(pyridin-2-yl) pyrrole-3-carboxylate (180 mg, 0.753 mmol, 1 equiv, 95%) in THF (4 mL) and H₂O (2 mL) was added LiOH (90.1 mg, 3.762 mmol, 5.00 equiv) at room temperature. The resulting solution was stirred for one hour at room temperature. The residue was purified by flash chromatography on C18 column gel eluting with ACN/water (0.1%FA) (45%) to afford 5-cyano-1-(pyridin-2-yl) pyrrole-3-carboxylic acid (150 mg, 86.01%) as a yellow solid. LCMS: (ESI): [M + H]⁺:214
- Step 5: A solution of 5-cyano-1-(pyridin-2-yl) pyrrole-3-carboxylic acid (150 mg, 0.704 mmol, 1 equiv), NMI (115.5 mg, 1.407 mmol, 2.00 equiv) and TCFH (296.1 mg, 1.055 mmol, 1.50 equiv) in ACN (5 mL) at room temperature. The resulting solution was stirred for 10 minutes at room temperature. Then N-(3-amino-5-chlorophenyl) methanesulfonamide (186.3 mg, 0.844 mmol, 1.20 equiv) was added and stirred for one hour at room temperature. The residue was purified by Pre-HPLC on condition: Column: Sunfire prep C18 column, 30*150 mm, 5μm; Mobile Phase A: Water(0.1%FA), Mobile Phase B: ACN; Flow rate: 60 mL/min; Gradient: 40% B to 60% B in 7 min, 60% B; Wave Length: 254/220 nm; RT1(min): 5.72 to afford N-(3-chloro-5-methanesulfonamidophenyl)-5-(phenylamino)-2H-pyrazole-3-carboxamide (29.6 mg, 48.73%) as a white solid. LCMS: (ESI): [M + H]⁺:416. ¹H NMR (300 MHz, DMSO-d₆) δ 10.21 (s, 1H), 10.10 (s, 1H), 8.63 (d, J = 4.8 Hz, 1H), 8.56 (d, J = 1.8 Hz, 1H), 8.19 – 8.08 (m, 1H), 7.89 (d, J = 8.2 Hz, 1H), 7.81 (d, J = 1.8 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.62 (d, J = 2.0 Hz, 1H), 7.57 (dd, J = 7.4, 4.7 Hz, 1H), 6.95 (t, J = 2.0 Hz, 1H), 3.08 (s, 3H).

Example 99: 1-benzyl-N-(3-bromo-5-(methylsulfonamido)phenyl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazole-5-carboxamide



Step 1: To a stirred solution of 4-fluoro-3-nitrobenzoic acid (1 g, 5.40 mmol) and 1-phenylmethanamine (578 mg, 5.40 mmol) in DMF (15 mL) was added Cs_2CO_3 (5.26 g, 16.2 mmol). Then the mixture was stirred for 2 h at 80 °C. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/ H_2O (0-100%, acidic system) to afford 4-(benzylamino)-3-nitrobenzoic acid (700 mg, 2.57 mmol) as a yellow solid.

LCMS (ESI) $[\text{M} + \text{H}]^+$: 273.08

Step 2: To a stirred solution of 4-(benzylamino)-3-nitrobenzoic acid (700 mg, 2.57 mmol) and NH_4Cl (1.08 g, 20.5 mmol) in EtOH (10 mL) and H_2O (3 mL) was added Fe (1.43 g, 25.6 mmol). Then the mixture was stirred for 2 h at 90 °C. The mixture was concentrated and purified by flash chromatography eluting with EtOAc/Petroleum ether (0-100%, acidic system) to afford 3-amino-4-(benzylamino)benzoic acid (330 mg, 1.36 mmol) as a yellow solid. LCMS (ESI) $[\text{M} + \text{H}]^+$: 243.11


Step 3: To a stirred solution of 3-amino-4-(benzylamino)benzoic acid (330 mg, 1.36 mmol) in THF (6 mL) and MeOH (2 mL) was added a solution of (trimethylsilyl)diazomethane (4.08 mmol) in hexane dropwise under N₂ atmosphere. Then the mixture was stirred for 2 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (0-100%, acidic system) to afford methyl 3-amino-4-(benzylamino) benzoate (265 mg, 1.03 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 357.12

Step 4: To a stirred solution of methyl 3-amino-4-(benzylamino)benzoate (265 mg, 1.03 mmol) in DMF (5 mL) was added CDI (225 mg, 3.09 mmol). Then the mixture was stirred for 2 h at 60 °C. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (0-100%, acidic system) to afford methyl 1-benzyl-2-oxo-2,3-dihydro-1H-1,3-benzodiazole-5-carboxylate (100 mg, 0.354 mmol) as an off-white solid. LCMS (ESI) [M + H]⁺: 283.02

Step 5: To a stirred solution of methyl 1-benzyl-2-oxo-2,3-dihydro-1H-1,3-benzodiazole-5-carboxylate (100 mg, 0.354 mmol) in THF (3 mL) and H₂O (1 mL) was added LiOH (25.3 mg, 1.06 mmol). Then the mixture was stirred for 2 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (0-100%, acidic system) to afford 1-benzyl-2-oxo-2,3-dihydro-1H-1,3-benzodiazole-5-carboxylic acid (76.0 mg, 0.283 mmol) as an off-white solid. LCMS (ESI) [M + H]⁺: 269.04

Step 6: To a stirred solution of 1-benzyl-2-oxo-2,3-dihydro-1H-1,3-benzodiazole-5-carboxylic acid (76 mg, 0.283 mmol), NMI (115 mg, 1.41 mmol) and TCFH (118 mg, 0.424 mmol) in ACN (5 mL) was added N-(3-amino-5-bromophenyl) methane sulfonamide (75.0 mg, 0.283 mmol). Then the mixture was stirred for 2 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (0-100%, acidic system) to afford 1-benzyl-N-(3-bromo-5-methanesulfonamido phenyl)-2-oxo-2,3-dihydro-1H-1,3-benzodiazole-5-carboxamide (21.1 mg, 0.0395 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 515.3. ¹H NMR (300 MHz, DMSO-d₆) δ 10.61 (s, 1H), 8.52 (d, J = 2.6 Hz, 1H), 8.34 (s, 1H), 8.03 (d, J = 8.1 Hz, 1H), 7.84 (d, J = 7.7 Hz, 1H), 7.71 (d, J = 1.9 Hz, 2H), 7.65 (t, J = 7.8 Hz, 1H), 6.98 (t, J = 2.0 Hz, 1H), 6.55 (d, J = 2.5 Hz, 1H), 5.26 (s, 1H), 4.55 (s, 2H), 3.08 (s, 3H).


Example 100: N-(3-chloro-5-(methanesulfonamido)phenyl)-3-(3-cyano-1H-pyrazol-1-yl)benzamide



Step 1: To a stirred solution of 1H-pyrazole-3-carbonitrile (260 mg, 2.79 mmol), CuI (106 mg, 0.558 mmol), (1S,2S)-N1,N2-dimethylcyclohexane-1,2-diamine (157 mg, 1.11 mmol) and 3-bromobenzoic acid (560 mg, 2.79 mmol) in DMSO (10 mL) was added Cs₂CO₃ (2.72 g, 8.37 mmol). Then the mixture was stirred for 2 h at 100 °C under N₂ atmosphere. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (0-100%, acidic system) to afford 3-(3-cyano-1H-pyrazol-1-yl)benzoic acid (85.0 mg, 0.398 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 214.05

Step 2: To a stirred solution of 3-(3-cyano-1H-pyrazol-1-yl)benzoic acid (85 mg, 0.398 mmol), TCFH (167 mg, 0.597 mmol) and NMI (163 mg, 1.99 mmol) in ACN (5 mL) was added N-(3-amino-5-chlorophenyl)methanesulfonamide (87.8 mg, 0.398 mmol). Then the mixture was stirred for 2 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (0-100%, acidic system) to afford N-(3-chloro-5-methane sulfonamidophenyl)-3-(3-cyano-1H-pyrazol-1-yl)benzamide (24.3 mg, 0.0570 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 416.05. ¹H NMR (300 MHz, DMSO-d₆) δ 10.66 (s, 1H), 10.12 (s, 1H), 8.92 (d, J = 2.7 Hz, 1H), 8.43 (d, J = 2.2 Hz, 1H), 8.19 – 8.09 (m, 1H), 8.01 (d, J = 7.8 Hz, 1H), 7.81 – 7.68 (m, 3H), 7.35 (d, J = 2.6 Hz, 1H), 6.99 (t, J = 2.0 Hz, 1H), 3.09 (s, 3H).


Example 101: N-(3-bromo-5-(methanesulfonamido)phenyl)-2-oxo-1-phenyl-2,3-dihydro-1H-imidazole-4-carboxamide



- Step 1: A mixture of phenylurea (500 mg, 3.67 mmol) and ethyl 3-bromo-2-oxopropanoate (715 mg, 3.67 mmol) in EtOH (10 mL) was stirred for 2 h at 80 °C. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (0-100%, acidic system) to afford ethyl 2-oxo-1-phenyl-2,3-dihydro-1H-imidazole-4-carboxylate (235 mg, 1.01 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 233.08
- Step 2: To a stirred solution of ethyl 2-oxo-1-phenyl-2,3-dihydro-1H-imidazole-4-carboxylate (235 mg, 1.01 mmol) in EtOH (3 mL) and H₂O (1mL) was added LiOH (72.5 mg, 3.03 mmol). Then the mixture was stirred for 2 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (0-100%, acidic system) to afford 2-oxo-1-phenyl-2,3-dihydro-1H-imidazole-4-carboxylic acid (180 mg, 0.881 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 205.05
- Step 3: To a stirred solution of 2-oxo-1-phenyl-2,3-dihydro-1H-imidazole-4-carboxylic acid (180 mg, 0.881 mmol), TCFH (369 mg, 1.32 mmol) and NMI (360 mg, 4.40 mmol) in ACN (10 mL) was added N-(3-amino-5-bromophenyl)methanesulfonamide (233 mg, 0.881 mmol). Then the mixture was stirred for 2 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (0-100%, acidic system) to afford N-(3-bromo-5-methanesulfonamidophenyl)-2-oxo-1-phenyl-2,3-dihydro-1H-imidazole-4-carboxamide (21.1 mg, 0.0466 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 451. ¹H NMR (300 MHz, DMSO-d₆) δ 10.38 (s, 1H), 10.05 (d, J = 7.7 Hz, 2H), 8.33 (s,

1H), 7.83 (s, 1H), 7.81 – 7.71 (m, 3H), 7.35 (t, J = 7.8 Hz, 2H), 7.12 (s, 1H), 7.00 (t, J = 7.3 Hz, 1H), 3.09 (s, 3H).

Example 102: N-(3-chloro-5-(methanesulfonamido)phenyl)-3-(2-chloropyridin-4-yl)benzamide




Step 1: To a stirred solution of (2-chloropyridin-4-yl)boronic acid (500 mg, 3.17 mmol), Pd(dppf)Cl₂ (347 mg, 0.475 mmol) and 3-bromobenzoic acid (637 mg, 3.17 mmol) in dioxane (15 mL) was added K₂CO₃ (1.32 g, 9.51 mmol). Then the mixture was stirred for 2 h at 80 °C. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (0-100%, acidic system) to afford 3-(2-chloropyridin-4-yl)benzoic acid (210 mg, 0.898 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 234.02

Step 2: To a stirred solution of 3-(2-chloropyridin-4-yl)benzoic acid (200 mg, 0.855 mmol), TCFH (358 mg, 1.28 mmol) and NMI (350 mg, 4.27 mmol) in ACN (5 mL) was added N-(3-amino-5-chlorophenyl)methanesulfonamide (188 mg, 0.855 mmol). Then the mixture was stirred for 2 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (0-100%, acidic system) to afford N-(3-chloro-5-methanesulfonamidophenyl)-3-(2-chloropyridin-4-yl)benzamide (24.8 mg, 0.0566 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 436.02. ¹H NMR (300 MHz, DMSO-d₆) δ 10.59 (s, 1H), 10.11 (s, 1H), 8.54 (d, J = 5.3 Hz, 1H), 8.37 (s, 1H), 8.10 (d, J = 7.7 Hz, 1H), 8.06 (d, J = 7.8 Hz, 1H), 8.00 (d, J = 1.6 Hz, 1H), 7.96 – 7.78 (m, 1H), 7.77 – 7.66 (m, 3H), 6.99 (t, J = 1.9 Hz, 1H), 3.09 (s, 3H).

20

Example 103: N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(2-(2,2,2-trifluoro-1-hydroxyethyl)phenyl)-1H-pyrazole-4-carboxamide



- 5 Step 1: To a stirred solution of ethyl 1H-pyrazole-4-carboxylate (310 mg, 2.21 mmol), 2-bromobenzaldehyde (490 mg, 2.65 mmol), (1S,2S)-N1,N2-dimethylcyclohexane-1,2-diamine (125 mg, 0.884 mmol), CuI (83.9 mg, 0.442 mmol) and K₂CO₃ (914 mg, 6.63 mmol) in ACN (10.00 mL). Then the mixture was stirred for 4 h at 90 °C under N₂. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford ethyl 1-(2-formylphenyl)-1H-pyrazole-4-carboxylate (332 mg, 1.36 mmol, 95 %) as a yellow solid. LCMS (ESI) [M + H]⁺: 245

- Step 2: To a stirred solution of ethyl 1-(2-formylphenyl)-1H-pyrazole-4-carboxylate (350 mg, 1.43 mmol), trimethyl(trifluoromethyl)silane (813 mg, 5.72 mmol) and K₂CO₃ (592 mg, 4.29 mmol) in DMF (10.00 mL). Then the mixture was stirred for 4 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford ethyl 1-[2-(2,2,2-trifluoro-1-hydroxyethyl)phenyl]-1H-pyrazole-4-carboxylate (209 mg, 95 %) as a yellow solid. LCMS (ESI) [M + H]⁺: 315

- Step 3: To a stirred solution of ethyl 1-[2-(2,2,2-trifluoro-1-hydroxyethyl)phenyl]-1H-pyrazole-4-carboxylate (220 mg, 0.700 mmol) and LiOH (10 mg, 2.5 mmol) in THF (10.00 mL) was added H₂O (2.00 mL). Then the mixture was stirred for 2 h at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 1-[2-

(2,2,2-trifluoro-1-hydroxyethyl)phenyl]-1H-pyrazole-4-carboxylic acid (104 mg, 95 %) as a white solid. LCMS (ESI) $[M + H]^+$: 310

Step 4: To a stirred solution of 1-[2-(2,2,2-trifluoro-1-hydroxyethyl)phenyl]-1H-pyrazole-4-carboxylic acid (110 mg, 0.384 mmol), TCFH (107 mg, 0.384 mmol) and NMI (31.3 mg,


5 0.384 mmol) in ACN (10.00 mL) was added N-(3-amino-5-

bromophenyl)methanesulfonamide (101 mg, 0.384 mmol). Then the mixture was stirred for 4 h at room temperature. The mixture was concentrated and purified by reverse phase flash

chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford N-(3-bromo-5-

10 methanesulfonamidophenyl)-1-[2-(2,2,2-trifluoro-1-hydroxyethyl)phenyl]-1H-pyrazole-4-carboxamide (34.4 mg, 96.918 %) as a white solid. LCMS (ESI) $[M + H]^+$: 534. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.17 (s, 1H), 10.09 – 10.04 (m, 1H), 8.70 (s, 1H), 8.34 (d, *J* = 0.6 Hz, 1H), 7.88 – 7.81 (m, 2H), 7.69 – 7.58 (m, 3H), 7.58 – 7.51 (m, 1H), 7.11 – 7.05 (m, 2H), 5.61 (p, *J* = 6.9 Hz, 1H), 3.07 (s, 3H).

Example 104: 4-((1H-pyrrol-2-yl)methyl)-N-(3-bromo-5-(methylsulfonamido)phenyl)-thiophene-2-carboxamide




Step 1: To a solution of methyl 4-methylthiophene-2-carboxylate (500 mg, 3.201 mmol, 1 equiv) and NBS (854.6 mg, 4.802 mmol, 1.50 equiv) in CCl₄ (10 mL) was added BPO

- (1640.6 mg, 6.402 mmol, 2.00 equiv) at room temperature under nitrogen. The resulting solution was stirred overnight at 80 °C. The resulting solution was concentrated under vacuum. The residue was purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (26%) to afford methyl 4-(bromomethyl) thiophene-2-carboxylate (200 mg, 26.5%) as a yellow oil. LCMS: (ESI): [M + H]⁺:235
- 5 Step 2: To a solution of methyl 4-(bromomethyl)thiophene-2-carboxylate (200 mg, 0.851 mmol, 1 equiv), Pd(PPh₃)₄ (98.3 mg, 0.085 mmol, 0.10 equiv) and 1-(tert-butoxycarbonyl)pyrrol-2-ylboronic acid (269.3 mg, 1.276 mmol, 1.50 equiv) in Dioxane (5 mL) and H₂O (0.5 mL) was added Cs₂CO₃ (831.5 mg, 2.552 mmol, 3.00 equiv) at room
- 10 temperature under nitrogen. The resulting solution was stirred for 2 hours at 90 °C. The reaction mixture was diluted with water (50 mL). The resulting solution was extracted with EtOAc (50×3 mL) and the organic layers were combined. The organic layer was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (42%) to afford tert-butyl
- 15 2-([5-(methoxycarbonyl) thiophen-3-yl]methyl)pyrrole-1-carboxylate (80 mg, 27.50%) as a yellow oil. LCMS: (ESI): [M + H]⁺:322
- Step 3: To a solution of tert-butyl 2-([5-(methoxycarbonyl)thiophen-3-yl]methyl)pyrrole-1-carboxylate (80 mg, 0.249 mmol, 1 equiv) in THF (2 mL) and H₂O (1 mL) was added LiOH (29.8 mg, 1.244 mmol, 5.00 equiv) at room temperature. The resulting solution was stirred
- 20 for one hour at room temperature. The residue was purified by flash chromatography on C18 column gel eluting with ACN/water (0.1%FA) (33%) to afford 4-([1-(tert-butoxycarbonyl)pyrrol-2-yl]methyl)thiophene-2-carboxylic acid (60 mg, 75.29%) as a white solid. LCMS: (ESI): [M + H]⁺:308
- Step 4: A solution of 4-([1-(tert-butoxycarbonyl)pyrrol-2-yl]methyl)thiophene-2-carboxylic acid (50 mg, 0.163 mmol, 1 equiv), NMI (26.7 mg, 0.325 mmol, 2.00 equiv) and TCFH (68.5 mg, 0.244 mmol, 1.50 equiv) in ACN (2 mL) at room temperature. The resulting solution was stirred for 10 min at room temperature. Then N-(3-amino-5-bromophenyl)
- 25 methanesulfonamide (64.7 mg, 0.244 mmol, 1.50 equiv) was added and stirred at room temperature for one hour. The residue was purified by C18 column gel eluting with
- 30 ACN/water (0.05 NH₄HCO₃) (65%) to afford tert-butyl 2-([5-([3-bromo-5-methanesulfonamidophenyl]carbonyl)thiophen-3-yl]methyl)pyrrole-1-carboxylate (55 mg, 59.93%) as a white solid. LCMS: (ESI): [M + H]⁺:554
- Step 5: To a solution of tert-butyl 2-([5-([3-bromo-5-methanesulfonamidophenyl]carbonyl)thiophen-3-yl]methyl)pyrrole-1-carboxylate (55 mg,

0.099 mmol, 1 equiv) in DCM (2 mL) was added TFA (1 mL) at 0 °C. The mixture was stirred for one hour at room temperature. The residue was purified by Pre-HPLC on condition: Column: YMC-Actus Triart C18 ExRS, 30*150 mm, 5µm; Mobile Phase A: Water (10mmol/L NH₄HCO₃), Mobile Phase B: ACN; Flow rate: 60mL/min; Gradient: 40% B to 60% B in 7 min, 60% B; Wave Length: 254/220 nm; RT1(min): 9.28/11.57 to afford N-(3-bromo-5-methanesulfonamidophenyl)-4-(1H-pyrrol-2-ylmethyl) thiophene-2-carboxamide (16.0 mg, 34.8 9%) as a white solid. LCMS: (ESI): [M + H]⁺:454

¹H NMR (300 MHz, DMSO-d₆) δ 10.65 (s, 1H), 10.36 (s, 1H), 10.08 (s, 1H), 7.89 (s, 1H), 7.77 (s, 1H), 7.65 (s, 1H), 7.50 (s, 1H) 7.06 (s, 1H), 6.63 (s, 1H), 5.93(d, 1H), 5.78(s, 1H), 3.90 (s, J = 5.2 Hz, 2H), 2.95 (s, 3H).






Example 105: N-(3-fluoro-5-(methanesulfonamido)phenyl)-4-(pyridin-3-yl)thiophene-2-carboxamide




Step 1: To a solution of 4-bromo-N-(3-fluoro-5-methanesulfonamidophenyl)thiophene-2-carboxamide (100 mg, 0.254 mmol, 1 equiv) and pyridin-3-ylboronic acid (46.89 mg, 0.381 mmol, 1.5 equiv) and Pd(PPh₃)₂Cl₂ (17.85 mg, 0.025 mmol, 0.1 equiv) and Na₂CO₃ (80.86 mg, 0.762 mmol, 3 equiv) in EtOH (3 mL) and H₂O (1 mL) was stirred for 2 h at 90 °C under nitrogen atmosphere. The residue was purified by reverse phase flash chromatography with the following conditions: column, silica gel; mobile phase, MeCN in water, 0% to 100% gradient in 20 min; detector, UV 254 nm. This resulted in N-(3-fluoro-5-methanesulfonamidophenyl)-4-(pyridin-3-yl)thiophene-2-carboxamide (27.8 mg, 27.93%) as a light green solid. LCMS: (ESI): LCMS (ESI) [M + H]⁺: 392. ¹H NMR (300 MHz, DMSO-d₆) δ 10.61 (s, 1H), 10.14 (s, 1H), 9.11 (s, 1H), 8.66 (d, J = 21.6 Hz, 2H), 8.48 (s, 2H), 7.75(s, 2H), 7.49 (s, 2H), 6.76 (d, J = 10.4 Hz, 1H), 3.09 (s, 3H).


Examples 108-111

The compounds listed in the following table were prepared using a procedure similar to that described for example 105:

Structure	Example No.	MS (ESI) [M+H] ⁺	¹ H NMR
	106	423	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.47 (s, 1H), 10.09 (s, 1H), 9.95 (s, 1H), 8.47 (d, J = 1.4 Hz, 1H), 8.15 (d, J = 1.3 Hz, 1H), 7.67 (dt, J = 11.3, 1.9 Hz, 2H), 7.54 (dd, J = 7.8, 1.7 Hz, 1H), 7.24 – 7.12 (m, 1H), 7.03 – 6.93 (m, 2H), 6.91 (td, J = 7.4, 1.2 Hz, 1H), 3.08 (s, 3H).
	107	437	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.45 (s, 1H), 10.08 (s, 1H), 8.41 (d, J = 1.4 Hz, 1H), 8.09 (d, J = 1.3 Hz, 1H), 7.66 (dt, J = 15.8, 1.9 Hz, 2H), 7.57 (dd, J = 7.5, 1.7 Hz, 1H), 7.36 (t, J = 8.1 Hz, 1H), 7.15 (d, J = 8.2 Hz, 1H), 7.06 (t, J = 7.4 Hz, 1H), 6.97 (t, J = 1.9 Hz, 1H), 3.87 (s, 3H), 3.07 (s, 3H).
	108	425	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.47 (s, 1H), 10.12 (s, 1H), 8.51 (d, J = 1.4 Hz, 1H), 8.33 (d, J = 1.4 Hz, 1H), 7.71 – 7.47 (m, 6H), 7.20 (dd, J = 9.7, 7.3 Hz, 1H), 6.98 (t, J = 1.9 Hz, 1H), 3.09 (s, 3H).
	109	404	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.80 (s, 1H), 7.73 (t, J = 1.9 Hz, 1H), 7.67 – 7.61 (m, 2H), 7.54 – 7.47 (m, 2H), 7.51 – 7.43 (m, 2H), 7.42 – 7.33 (m, 1H), 6.90 (t, J = 1.9 Hz, 1H), 6.81 (d, J = 2.0 Hz, 1H), 3.71 (s, 3H), 3.06 (s, 3H).
	110	422	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.49 (s, 1H), 10.15 (s, 1H), 9.00 (s, 1H), 8.59 – 8.50 (m, 2H), 8.37 (s, 1H), 8.13 (d, J = 8.0 Hz, 1H), 7.65 (d, J = 11.2 Hz, 2H), 7.51 (t, J = 6.6 Hz, 1H), 6.98 (s, 1H), 3.19 (q, J = 7.2 Hz, 2H), 1.22 (t, J = 7.2 Hz, 3H).

	111	513	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.45 (s, 1H), 10.09 (s, 1H), 8.45 (s, 1H), 8.10 (d, J = 1.4 Hz, 1H), 7.63 (dd, J = 18.6, 7.5 Hz, 3H), 7.48 (d, J = 7.2 Hz, 2H), 7.41 – 7.16 (m, 5H), 7.08 (s, 1H), 6.97 (s, 1H), 5.24 (s, 2H), 3.08 (s, 3H).
---	-----	-----	--

Example 112: N-(3-fluoro-5-(methanesulfonamido)phenyl)-4-(pyridin-2-yl)thiophene-2-carboxamide



- 5 Step 1: To a stirred solution of N-(3-amino-5-fluorophenyl)methanesulfonamide (100 mg, 0.489 mmol), TCFH (178 mg, 0.635 mmol) and NMI (119 mg, 1.46 mmol) in ACN (5 mL) was added 4-bromothiophene-2-carboxylic acid (111 mg, 0.537 mmol). Then the mixture was stirred for 1 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford 4-bromo-N-(3-fluoro-5-methanesulfonamidophenyl) thiophene-2-carboxamide (128 mg, 0.327 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 393
- 10 Step 2: A solution of 4-bromo-N-(3-fluoro-5-methanesulfonamidophenyl)thiophene-2-carboxamide (130 mg, 0.330 mmol), Pd(dppf)Cl₂ (24.1 mg, 0.0330 mmol), KOAc (97.1 mg, 0.990 mmol) and 4,4,5,5-tetramethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,2-dioxaborolane (335 mg, 1.32 mmol) in DMSO (10 mL) was stirred for 2 h at 80 °C under nitrogen atmosphere. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford N-(3-fluoro-5-
- 15

methanesulfonamidophenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxamide (69.3 mg, 0.157 mmol) as a yellow solid. LCMS (ESI) $[M + H]^+$: 441


Step 3: A solution of N-(3-fluoro-5-methanesulfonamidophenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxamide (70 mg, 0.158 mmol), Pd(dppf)Cl₂ (11.5 mg, 0.0158 mmol), K₂CO₃ (65.5 mg, 0.474 mmol) and 2-bromopyridine (49.9 mg, 0.316 mmol) in dioxane (5 mL) was stirred for 2 h at 80 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with Petroleum ether / EtOAc (0-100%) to afford N-(3-fluoro-5-methanesulfonamidophenyl)-4-(pyridin-2-yl)thiophene-2-carboxamide (26.7 mg, 0.0682 mmol) as a white solid. LCMS (ESI) $[M + H]^+$: 392. ¹H NMR (300 MHz, DMSO-d₆) δ 10.64 (s, 1H), 10.13 (s, 1H), 8.75 (d, J = 1.4 Hz, 1H), 8.65 (dt, J = 4.8, 1.5 Hz, 1H), 8.52 (d, J = 1.3 Hz, 1H), 7.98 – 7.85 (m, 2H), 7.56 – 7.43 (m, 2H), 7.41 – 7.31 (m, 1H), 6.75 (dt, J = 10.5, 2.2 Hz, 1H), 3.09 (s, 3H).

Examples 113 and 114

The compounds listed in the following table were prepared using a procedure similar to that described for example 112:

Structure	Example No.	MS (ESI) $[M+H]^+$	¹ H NMR
	113	406	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.63 (s, 1H), 10.13 (s, 1H), 8.75 (d, J = 1.4 Hz, 1H), 8.64 (dt, J = 4.7, 1.5 Hz, 1H), 8.51 (d, J = 1.3 Hz, 1H), 8.02 – 7.86 (m, 2H), 7.56 – 7.42 (m, 2H), 7.36 (ddd, J = 6.3, 4.8, 2.4 Hz, 1H), 6.75 (dt, J = 10.5, 2.2 Hz, 1H), 3.18 (t, J = 7.3 Hz, 2H), 1.22 (t, J = 7.3 Hz, 3H)
	114	407	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.49 (s, 1H), 10.10 (s, 1H), 9.95 (s, 1H), 8.47 (d, J = 1.4 Hz, 1H), 8.14 (d, J = 1.3 Hz, 1H), 7.54 (dd, J = 7.7, 1.7 Hz, 1H), 7.47 (d, J = 9.0 Hz, 2H), 7.22 – 7.13 (m, 1H), 6.99 (dd, J = 8.2, 1.2 Hz, 1H), 6.96 – 6.87 (m, 1H), 6.74 (dd, J = 10.5, 2.2 Hz, 1H), 3.08 (s, 3H).

Example 115: N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(2,6-difluorophenyl)thiophene-2-carboxamide



Step 1: To a stirred solution of 4-bromothiophene-2-carboxylic acid (2.02 g, 9.75 mmol),
 5 TCFH (4.11 g, 14.6 mmol) and NMI (2.3 g, 29.25 mmol) in ACN (50.00 mL) was added 3-chloro-5-nitroaniline (1.68 g, 9.75 mmol). Then the mixture was stirred for 4 h at room temperature. The mixture was concentrated and purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (0-100%) to afford 4-bromo-N-(3-chloro-5-nitrophenyl)thiophene-2-carboxamide (3.2 g, 95 %) as a yellow solid. LCMS (ESI) $[M + H]^+$:
 10 361


Step 2: To a solution of 4-bromo-N-(3-chloro-5-nitrophenyl) thiophene-2-carboxamide (3.2 g, 8.84 mmol) in a solution of methanol and water (50 mL) was added Fe (2.4 g, 44 mmol) and NH_4Cl (4.7 g, 88 mmol). The resulting mixture was stirred for 2 hours at 80 °C under N_2 atmosphere. The resulting mixture was filtered. The filtrate was concentrated under reduced
 15 pressure to give the crude product which was purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (0-100%) to afford N-(3-amino-5-chlorophenyl)-4-

bromothiophene-2-carboxamide (2.93 g, 95 %) as a yellow solid. LCMS (ESI) $[M + H]^+$: 331

Step 3: To a stirred solution of N-(3-amino-5-chlorophenyl)-4-bromothiophene-2-carboxamide (1.02 g, 3.07 mmol) in Pyridine (15.00 mL) was added methanesulfonyl chloride (703 mg, 6.14 mmol) dropwise at 0 °C. Then the mixture was stirred for 4 h at room temperature. The mixture was concentrated and purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (0-100%) to afford 4-bromo-N-(3-chloro-5-methanesulfonamidophenyl) thiophene-2-carboxamide (800 mg, 95 %) as a yellow solid. LCMS (ESI) $[M + H]^+$: 409

Step 4: To a stirred solution of 4-bromo-N-(3-chloro-5-methanesulfonamidophenyl) thiophene-2-carboxamide (100 mg, 0.244 mmol), 2-(2,6-difluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (117 mg, 0.488 mmol), XPhosPdG₃ (21 mg, 0.0244 mmol) and K₃PO₄ (155 mg, 0.732 mmol) in 1,4-dioxane (5.00 mL). Then the mixture was stirred for 8 h at 80 °C. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford N-(3-chloro-5-methanesulfonamidophenyl)-4-(2,6-difluorophenyl) thiophene-2-carboxamide (31.3 mg, 98.180 %) as an off-white solid. LCMS (ESI) $[M + H]^+$: 443. ¹H NMR (300 MHz, DMSO-d₆) δ 10.52 (s, 1H), 10.10 (s, 1H), 8.31 (s, 1H), 8.15 (s, 1H), 7.66 (d, J = 13.3 Hz, 2H), 7.50 (q, J = 7.4 Hz, 1H), 7.28 (t, J = 8.2 Hz, 2H), 6.98 (s, 1H), 3.08 (s, 3H).

Example 116: N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(pyridin-3-yl)thiophene-2-carboxamide




Step 1: To a mixture of 4-(pyridin-3-yl)thiophene-2-carboxylic acid (280 mg, 1.36 mmol) and 3-chloro-5-nitroaniline (281 mg, 1.63 mmol) in ACN (3 mL) was added TCFH (571 mg, 2.04 mmol) and NMI (334 mg, 4.08 mmol) at room temperature for two hours. Then it was concentrated, the residue was purified by reverse phase flash chromatography eluting with 50% of acetonitrile in water (0.1% NH_4HCO_3) to afford N-(3-chloro-5-nitrophenyl)-4-(pyridin-3-yl)thiophene-2-carboxamide (212 mg,) as a white solid.


Step 2: To a mixture of N-(3-chloro-5-nitrophenyl)-4-(pyridin-3-yl)thiophene-2-carboxamide (212 mg, 589 μmol), Fe powder (164 mg, 2.94 mmol), ethyl 1H-pyrazole-4-carboxylate (434 mg, 3.10 mmol), NH_4Cl (157 mg, 2.94 mmol) was added EtOH (2 mL) and H_2O (0.2 mL). The resulting mixture was stirred for 2 hours at 80 °C under N_2 atmosphere. The reaction mixture was extracted with ethyl acetate. The organic phase was concentrated under reduced pressure. The residue was purified by flash chromatography on silica gel eluting with 30% of ethyl acetate in petroleum ether to afford N-(3-amino-5-chlorophenyl)-4-(pyridin-3-yl)thiophene-2-carboxamide (156 mg) as a yellow solid.

Step 3: A solution of N-(3-amino-5-chlorophenyl)-4-(pyridin-3-yl)thiophene-2-carboxamide (156 mg, 473 μmol) and Et_3N (142 mg, 1.41 mmol) in DCM (1 mL) was stirred at room temperature. Then Ms_2O (123 mg, 709 μmol) was added into the solution at 0 °C and the resulting mixture was stirred at room temperature for 3 hours/ The solvent was concentrated under vacuum. The residue was purified onto silica gel column eluting with 60% of ethyl acetate in petroleum ether to afford N-(3-chloro-5-methanesulfonamidophenyl)-4-(pyridin-3-yl)thiophene-2-carboxamide (9.6 mg,) as a white solid. LCMS $[\text{M} + \text{H}]^+$: 408. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 10.49 (s, 1H), 9.00 (d, $J = 2.2$ Hz, 1H), 8.60 – 8.50 (m, 2H), 8.38 (d, $J = 1.4$ Hz, 1H), 8.14 (dt, $J = 7.9, 2.0$ Hz, 1H), 7.64 (dt, $J = 14.6, 1.9$ Hz, 2H), 7.52 (dd, $J = 7.9, 4.8$ Hz, 1H), 6.97 (t, $J = 1.9$ Hz, 1H), 3.07 (s, 3H).


Examples 117 and 118

The compounds listed in the following table were prepared using a procedure similar to that described for Example 116:

Structure	Example No.	MS (ESI) $[\text{M} + \text{H}]^+$	^1H NMR
	117	391	^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 10.62 (s, 1H), 10.10 (s, 1H), 8.63 (dd, $J = 2.5, 0.6$ Hz, 1H), 8.38 (t, $J = 2.0$ Hz, 1H), 8.08 (ddd, $J = 8.1, 2.3, 1.0$ Hz, 1H), 7.91 – 7.79 (m, 2H), 7.75 – 7.62 (m, 3H), 6.99 (t, $J = 2.0$ Hz,

			1H), 6.61 (dd, J = 2.5, 1.7 Hz, 1H), 3.09 (s, 3H).
	118	435	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.59 (s, 1H), 10.10 (s, 1H), 8.62 (d, J = 2.5 Hz, 1H), 8.38 (t, J = 2.0 Hz, 1H), 8.08 (ddd, J = 8.1, 2.3, 1.0 Hz, 1H), 7.85 (ddd, J = 9.7, 7.7, 1.6 Hz, 3H), 7.76 (t, J = 1.9 Hz, 1H), 7.67 (t, J = 7.9 Hz, 1H), 7.11 (t, J = 1.9 Hz, 1H), 6.61 (dd, J = 2.5, 1.8 Hz, 1H), 3.08 (s, 3H).

Example 119: 1-ethyl-N-(3-fluoro-5-(methylsulfonyl)phenyl)-5-methyl-1H-pyrazole-3-carboxamide



- 5 Step 1: To a stirred solution of methyl 3-amino-5-fluorobenzoate (1 g, 5.91 mmol), HATU (2.91 g, 7.68 mmol) and DIEA (2.28 g, 17.7 mmol) in DMF (15 mL) was added 1-ethyl-5-methyl-1H-pyrazole-3-carboxylic acid (1 g, 6.50 mmol). Then the mixture was stirred for 1 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford methyl 3-(1-ethyl-5-methyl-1H-pyrazole-3-amido)-5-fluorobenzoate (1.09 g, 3.59 mmol) as a white solid.

LCMS (ESI) [M + H]⁺: 306


- Step 2: To a stirred solution of methyl 3-(1-ethyl-5-methyl-1H-pyrazole-3-amido)-5-fluorobenzoate (1 g, 3.27 mmol) in EtOH (10 mL) and H₂O (10 mL) was added LiOH (782 mg, 32.6 mmol). Then the mixture was stirred for 2 h at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 3-(1-ethyl-5-methyl-1H-pyrazole-3-amido)-5-fluorobenzoic acid (848 mg, 2.91 mmol) as a yellow solid.

LCMS (ESI) [M + H]⁺: 292

Step 3: To a stirred mixture of 3-(1-ethyl-5-methylpyrazole-3-amido)-5-fluorobenzoic acid (500 mg, 1.717 mmol, 1 equiv) and DPPA (610 mg, 2.22 mmol) in Toluene (10 mL) was stirred for 2 h at 80 °C under nitrogen atmosphere. The mixture was allowed to cool down to room temperature, H₂O (1 mL) was added, and the mixture was stirred for another 1 h. The
5 resulting mixture was concentrated under vacuum. The mixture was concentrated and purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (0-100%) to afford N-(3-amino-5-fluorophenyl)-1-ethyl-5-methyl-1H-pyrazole-3-carboxamide (119 mg, 0.456 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 263

Step 4: To a stirred solution of N-(3-amino-5-fluorophenyl)-1-ethyl-5-methyl-1H-pyrazole-3-
10 carboxamide (100 mg, 0.381 mmol) and Et₃N (115 mg, 1.14 mmol) in DCM (10 mL) was added MsCl (47.7 mg, 0.419 mmol) dropwise at 0 °C. Then the mixture was stirred for 1 h at room temperature. The mixture was concentrated and purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (0-100%) to afford 1-ethyl-N-(3-fluoro-5-methanesulfonamidophenyl)-5-methyl-1H-pyrazole-3-carboxamide (54.7 mg, 0.1607 mmol)
15 as an off-white solid. LCMS (ESI) [M + H]⁺: 341. ¹H NMR (400 MHz, DMSO-d₆) δ 10.16 (d, J = 2.1 Hz, 1H), 10.04 (d, J = 2.2 Hz, 1H), 7.59 (d, J = 2.1 Hz, 1H), 7.48 (dt, J = 11.5, 2.2 Hz, 1H), 6.68 (dt, J = 10.4, 2.3 Hz, 1H), 6.57 (d, J = 2.1 Hz, 1H), 4.18 (qd, J = 7.2, 2.1 Hz, 2H), 3.09 (d, J = 2.1 Hz, 3H), 2.33 (d, J = 2.1 Hz, 3H), 1.37 (td, J = 7.3, 2.1 Hz, 3H).


Example 120: N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(hydroxymethyl)-4-phenylthiophene-2-carboxamide



- Step 1: To a stirred solution of N-(3-amino-5-chlorophenyl)methanesulfonamide (155 mg, 0.706 mmol), TCFH (297 mg, 1.059 mmol) and NMI (172 mg, 2.118 mmol) in ACN (15.00 mL) was added 5-bromo-4-phenylthiophene-2-carboxylic acid (200 mg, 0.706 mmol). Then the mixture was stirred for 4 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford 5-bromo-N-(3-chloro-5-methanesulfonamidophenyl)-4-phenylthiophene-2-carboxamide (350 mg, 95 %) as a white solid. LCMS (ESI) [M + H]⁺: 485
- Step 2: To a stirred solution of 5-bromo-N-(3-chloro-5-methanesulfonamidophenyl)-4-phenylthiophene-2-carboxamide (350 mg, 0.720 mmol), Ac₂O (140 mg, 1.37 mmol), Pd(OAc)₂ (16 mg, 0.071 mmol), Xantphos (80 mg, 0.138 mmol) and (COOH)₂ (129 mg, 1.44 mmol) and DIEA (266 mg, 2.06 mmol) in DMSO (10.00 mL). Then the mixture was stirred for 4 h at 90 °C under N₂ atmosphere. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford 5-[(3-chloro-5-methanesulfonamidophenyl) carbamoyl]-3-phenylthiophene-2-carboxylic acid (210 mg, 95 %) as a yellow solid. LCMS (ESI) [M + H]⁺: 451

- Step 3: To a stirred solution of 5-[(3-chloro-5-methanesulfonamidophenyl)carbamoyl]-3-phenylthiophene-2-carboxylic acid (210 mg, 0.465 mmol), TCFH (196 mg, 0.6975 mmol) and NMI (115 mg, 1.395 mmol) in ACN (5.00 mL) was added methanol (5 ml). Then the mixture was stirred for 4 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford methyl 5-[(3-chloro-5-methanesulfonamidophenyl)carbamoyl]-3-phenylthiophene-2-carboxylate (180 mg, 95 %) as a white solid. LCMS (ESI) [M + H]⁺: 465
- Step 4: To a stirred solution of methyl 5-[(3-chloro-5-methanesulfonamidophenyl)carbamoyl]-3-phenylthiophene-2-carboxylate (180 mg, 0.387 mmol) and NaBH₄ (44 mg, 1.16 mmol) in EtOH (10.00 mL) at 0 °C. Then the mixture was stirred for 1 h at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford N-(3-chloro-5-methanesulfonamidophenyl)-5-(hydroxymethyl)-4-phenylthiophene-2-carboxamide (37.2 mg, 98.340 %) as a white solid. LCMS (ESI) [M + H]⁺: 437. ¹H NMR (300 MHz, DMSO-d₆) δ 10.38 (s, 1H), 10.08 (s, 1H), 8.14 (s, 1H), 7.66 (dt, J = 15.7, 1.9 Hz, 2H), 7.54 – 7.44 (m, 4H), 7.41 (dt, J = 5.7, 3.4 Hz, 1H), 6.96 (t, J = 2.0 Hz, 1H), 5.88 (t, J = 5.4 Hz, 1H), 4.74 (d, J = 5.4 Hz, 2H), 3.08 (s, 3H).

Example 121: N-(3-bromo-5-(methylsulfonylamidimidamido)phenyl)-4-(pyridin-2-yl)thiophene-2-carboxamide



20

Step 1: A solution of N-(3-amino-5-bromophenyl)-4-(pyridin-2-yl) thiophene-2-carboxamide (180 mg, 0.481 mmol, 1 equiv) and 3-[[[(tert-butyl)dimethylsilyl]imino] (methyl)oxo-lambda6-sulfany]-1-methylimidazol-1-ium (158.41 mg, 0.577 mmol, 1.2 equiv) in MeCN (3



mL) was stirred for 1 h at room temperature. The residue was purified by reverse phase flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in Water (0.1% FA), 0% to 100% gradient in 10 min; detector, UV 254 nm. This resulted in N-{3-bromo-5-[N-(tert-butyldimethylsilyl)methanesulfonylimidamido]phenyl}-4-(pyridin-2-yl)thiophene-2-carboxamide (170 mg, 62.49%) as a white solid. LCMS: (ESI): LCMS (ESI) [M + H]⁺: 565



Step 2: N-{3-bromo-5-[N-(tert-butyldimethylsilyl)methanesulfonylimidamido]phenyl}-4-(pyridin-2-yl)thiophene-2-carboxamide (170 mg, 0.301 mmol, 1 equiv) was dissolved in a solution of HCl in 1,4-dioxane (3 mL, 1 N) and it was stirred for 1 h at room temperature.

The residue was purified by reverse phase flash chromatography with the following conditions: column, C₁₈ silica gel; mobile phase, MeCN in Water (0.1% FA), 0% to 100% gradient in 20 min; detector, UV 254 nm. This resulted in N-(3-bromo-5-methanesulfonylimidamidophenyl)-4-(pyridin-2-yl)thiophene-2-carboxamide (14.3 mg, 10.54%) as a white solid. LCMS: (ESI): LCMS (ESI) [M + H]⁺: 451. ¹H NMR (300 MHz, DMSO-d₆) δ 10.38 (s, 1H), 8.74 (d, J = 1.3 Hz, 1H), 8.64 (dt, J = 4.8, 1.4 Hz, 1H), 8.49 (d, J = 1.3 Hz, 1H), 7.95 – 7.88 (m, 2H), 7.58 (t, J = 1.8 Hz, 1H), 7.44 – 7.33 (m, 2H), 6.98 – 6.78 (m, 3H), 3.17 (s, 3H).

Examples 122 and 123

The compounds listed in the following table were prepared using a procedure similar to that described for example 120:

Structure	Example No.	MS (ESI) [M + H] ⁺	¹ H NMR
	122	465	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.33 (s, 1H), 8.64 – 8.47 (m, 2H), 8.17 (d, J = 1.4 Hz, 1H), 7.76 (dd, J = 7.8, 1.6 Hz, 1H), 7.57 (t, J = 1.9 Hz, 1H), 7.39 (t, J = 1.9 Hz, 1H), 7.32 (dd, J = 7.7, 4.7 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 6.87 (s, 2H), 3.16 (s, 3H), 2.51 (s, 3H).
	123	465	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.21 (s, 1H), 8.69 (ddd, J = 4.8, 1.8, 0.8 Hz, 1H), 8.39 (s, 1H), 7.93 (td, J = 7.7, 1.9 Hz, 1H), 7.69 (d, J = 8.0 Hz, 1H), 7.56 (t, J = 1.9 Hz, 1H), 7.37 (dt, J = 6.5, 2.9 Hz, 2H), 6.92 (t, J = 1.9 Hz, 1H), 6.87 (s, 2H), 3.16 (s, 3H), 2.71 (s, 3H).

Example 125: N-(3-chloro-5-(methanesulfonamido)phenyl)-5-phenylthiophene-3-carboxamide

- 5 Step 1: To a stirred solution of 5-bromothiophene-3-carboxylic acid (100 mg, 0.482 mmol), TCFH (204 mg, 0.723 mmol) and NMI (119 mg, 1.446 mmol) in ACN (5.00 mL) was added *N*-(3-amino-5-chlorophenyl)methanesulfonamide (106 mg, 0.482 mmol). Then the mixture was stirred for 4 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to
- 10 afford 5-bromo-*N*-(3-chloro-5-methanesulfonamidophenyl)thiophene-3-carboxamide (150 mg, 95%) as a yellow solid.
- LCMS (ESI) [M + H]⁺: 409.


- Step 2: To a stirred solution of 5-bromo-*N*-(3-chloro-5-methanesulfonamidophenyl)thiophene-3-carboxamide (90 mg, 0.219 mmol), phenylboronic acid (26.7 mg, 0.219 mmol), Pd(PPh₃)₄ (24 mg, 0.0219 mmol) and K₃PO₄ (139 mg, 0.657 mmol) in dioxane (10.00 mL) was added H₂O (3.00 mL). Then the mixture was stirred for 4 h at 90 °C under nitrogen. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford *N*-(3-chloro-5-methanesulfonamidophenyl)-5-phenylthiophene-3-carboxamide (45 mg, 99.350 %) as a
- 20 white solid. LCMS (ESI) [M + H]⁺: 407. ¹H NMR (300 MHz, DMSO-*d*₆) δ 10.31 (s, 1H), 10.09 (s, 1H), 8.35 (d, *J* = 1.4 Hz, 1H), 8.00 (d, *J* = 1.4 Hz, 1H), 7.71 (ddt, *J* = 10.6, 8.6, 1.7 Hz, 4H), 7.54 – 7.43 (m, 2H), 7.43 – 7.32 (m, 1H), 6.97 (t, *J* = 1.9 Hz, 1H), 3.08 (s, 3H).

Example 125

The compounds listed in the following table were prepared using a procedure similar to that described for example 123:

Structure	Example No.	MS (ESI) [M + H] ⁺	¹ H NMR
	125	391	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.24 (s, 1H), 10.09 (s, 1H), 8.48 (d, J = 0.8 Hz, 1H), 7.81 – 7.73 (m, 2H), 7.70 (t, J = 1.8 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.54 – 7.32 (m, 4H), 6.96 (t, J = 2.0 Hz, 1H), 3.08 (s, 3H).

5 **Example 126: N-(3-chloro-5-(methylsulfonylamino)phenyl)-1-(3-hydroxypropyl)-1H-pyrazole-4-carboxamide**





- Step 1: To a solution of ethyl 1H-pyrazole-4-carboxylate (200 mg, 1.42 mmol) and Cs₂CO₃ (693 mg, 2.13 mmol) in MeCN (5 mL) was added (3-bromopropoxy)(tert-butyl)dimethylsilane (377 mg, 1.49 mmol) at 0 °C, The mixture was stirred at 25°C for 5 h. The mixture was poured into water (20 mL) and extracted with EA (30 mL*3). The combined organic layers were washed with brine (15 mL*2) and dried over Na₂SO₄. The mixture was filtered, and the filtrate was concentrated in vacuo to give a crude. ethyl 1-((3-((tert-butyl)dimethylsilyloxy)propyl)-1H-pyrazol-4-yl)carboxylate (440 mg, 1.40 mmol, 99.3% yield) was obtained as colorless oil. LCMS: MS (ESI) Retention time: 0.972 min, (M) + = 313.2. ¹H NMR (400 MHz, CHLOROFORM-d) δ = 7.85 (d, J = 8.8 Hz, 2H), 4.33 - 4.14 (m,
- 10
- 15

4H), 3.52 (t, J = 5.6 Hz, 2H), 2.06 - 1.95 (m, 2H), 1.30 (t, J = 7.2 Hz, 3H), 0.86 (s, 9H), 0.08 - 0.11 (m, 6H).


- Step 2: To a solution of ethyl 1-(3-((tert-butyldimethylsilyl)oxy)propyl)-1H-pyrazole-4-carboxylate (440 mg, 1.47 mg) in THF (5 mL) and MeOH (5 mL) was added a solution of
- 5 LiOH.H₂O (185 mg, 4.41 mmol) in water (5 mL). The mixture was stirred at 25 °C for 14 h. The mixture was poured into water (20 mL) and extracted with EA (30 mL*3). The combined organic layers were washed with brine (15 mL*2) and dried over Na₂SO₄. The mixture was filtered, and the filtrate was concentrated in vacuo to give residue. The residue was purified by prep-HPLC and lyophilized. 1-(3-hydroxypropyl)-1H-pyrazole-4-carboxylic acid (160
- 10 mg, 0.592 mmol, 40.3% yield) was obtained as white solid. LCMS: MS (ESI) Retention time: 0.115 min, (M+1) + = 171.0. ¹H NMR (400 MHz, DMSO-d₆) δ = 8.22 (s, 1H), 7.78 (s, 1H), 4.60 (br s, 1H), 4.18 (t, J = 6.8 Hz, 2H), 3.40 - 3.35 (m, 2H), 1.91 (quin, J = 6.8 Hz, 2H)
- Step 3: To a solution of 1-(3-hydroxypropyl)-1H-pyrazole-4-carboxylic acid (75 mg, 0.441 mmol) and N-(3-amino-5-chlorophenyl)methanesulfonamide (106 mg, 484 μmol) in pyridine
- 15 (2 mL) was added EDCI (168 mg, 881 μmol). The mixture was stirred at 25 °C for 3 h. The mixture was poured into water (20 mL) and extracted with EA (25 mL*3). The combined organic layers were washed with 0.5 N HCl (15 mL*2), brine (15 mL*2) and dried over Na₂SO₄. The mixture was filtered, and the filtrate was concentrated in vacuo to give residue. The residue was purified by prep-HPLC and lyophilized. N-(3-chloro-5-
- 20 (methylsulfonamido)phenyl)-1-(3-hydroxypropyl)-1H-pyrazole-4-carboxamide (102.45 mg, 0.269 mmol, 97.96% purity, 60.9% yield) was obtained as white solid. LCMS: MS (ESI) Retention time: 0.718 min, (M+1) + = 373.0. ¹H NMR (400 MHz, DMSO-d₆) δ = 10.00 (s, 1H), 8.36 (s, 1H), 8.03 (s, 1H), 7.66 (t, J = 1.6 Hz, 1H), 7.56 (t, J = 1.6 Hz, 1H), 6.90 (t, J = 2.0 Hz, 1H), 4.63 (br s, 1H), 4.21 (t, J = 6.8 Hz, 2H), 3.41 - 3.37 (m, 2H), 3.05 (s, 3H), 1.93
- 25 (quin, J = 6.4 Hz, 2H)

Example 127-128

The compounds listed in the following table were prepared using a procedure similar to that described for example 125:

Structure	Example No.	MS (ESI) [M + H] ⁺	¹ H NMR
	127	359	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.05 (d, J = 4.8 Hz, 2H), 8.37 (s, 1H), 8.03 (s, 1H), 7.79 - 7.53 (m, 2H), 6.92 (t, J = 1.8 Hz, 1H), 4.99 (br t, J = 5.0 Hz, 1H), 4.20 (t, J = 5.2 Hz, 2H), 3.76 (q, J = 5.0 Hz, 2H), 3.07 (s, 3H)
	128	411	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.01 (s, 2H), 8.41 (s, 1H), 8.06 (s, 1H), 7.66 (t, J = 1.8 Hz, 1H), 7.56 (t, J = 1.8 Hz, 1H), 6.91 (t, J = 1.8 Hz, 1H), 4.79 (t, J = 8.0 Hz, 1H), 4.68 (s, 2H), 4.58 (s, 2H), 3.05 (s, 3H), 2.81 - 2.74 (m, 2H), 2.70 - 2.62 (m, 2H)

5 **Example 129: N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-phenyl-1H-pyrrole-3-carboxamide**



- Step 1: To a solution of ethyl 5-methyl-1H-pyrrole-3-carboxylate (100 mg, 652 μmol) and iodobenzene (199 mg, 978 μmol) in DMSO (2 mL) was added 1-hydroxypyrrolidine-2,5-dione (7.5 mg, 65.2 μmol), dipotassium carbonate (179 mg, 1.3 mmol) and λ¹-copper (1+) iodide (12.4 mg, 65.2 μmol) at 25 °C. The mixture was stirred at 90°C for 16 h. The mixture was poured to H₂O (50 mL). The mixture was extracted with EA (30 mL*2). The combined
- 10

organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column

chromatography (PE: EA = 10:1~1:1) to afford ethyl 5-methyl-1-phenyl-1H-pyrrole-3-carboxylate (430 mg, 1.87 mmol) as colorless oil. ¹H NMR (400 MHz, CHLOROFORM-d) δ = 7.58 - 7.38 (m, 4H), 7.35 - 7.29 (m, 2H), 6.48 (dd, J = 0.8, 1.8 Hz, 1H), 4.31 (q, J = 7.2 Hz, 2H), 1.36 (t, J = 7.2 Hz, 3H)


Step 2: To a solution of ethyl 5-methyl-1-phenyl-1H-pyrrole-3-carboxylate (420 mg, 1.83 mmol) in MeOH (3 mL), THF (2 mL) and H₂O (1 mL) was added lithium (1+) hydrate hydroxide (230 mg, 5.49 mmol) at 25 °C. The mixture was stirred at 60° for 16 h. The

mixture was poured to 1N HCl (20 mL). The mixture was extracted with EA (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 5-methyl-1-phenyl-1H-pyrrole-3-carboxylic acid (360 mg, 1.83 mmol, crude) as brown solid. LCMS: MS (ESI) Retention time: 0.747 min, (M) + = 202.0


Step 3: To a solution of 5-methyl-1-phenyl-1H-pyrrole-3-carboxylic acid (330 mg, 1.63 mmol) and N-(3-amino-5-chlorophenyl)methanesulfonamide (538 mg, 2.44 mmol) in pyridine (3 mL) was added EDCI (624 mg, 3.26 mmol) at 25°C. The mixture was stirred at 25°C for 2h. The mixture was poured to 1N HCl (20 mL). The mixture was extracted with EA (50 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC and lyophilized. 5-(2-(benzyloxy)phenyl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-1H-pyrazole-3-carboxamide (282.58 mg, 0.553 mmol, 98.80 % purity, 84.2% yield) was obtained as off-white solid. LCMS: MS (ESI) Retention time: 0.916 min, (M+1) + = 511.2. ¹H NMR (400 MHz, DMSO-d₆) δ = 10.33 (s, 1 H) 9.89 - 10.21 (m, 1 H) 7.68 - 7.78 (m, 2 H) 7.46 - 7.54 (m, 1 H) 7.25 - 7.41 (m, 7 H) 7.10 (t, J=7.6 Hz, 1 H) 6.91 (t, J=1.6 Hz, 1 H) 6.79 (s, 1 H) 5.18 (s, 2 H) 3.74 (s, 3 H) 3.08 (s, 3 H).

Example 130

The compounds listed in the following table were prepared using a procedure similar to that described for example 129:

Structure	Example No.	MS (ESI) [M+H] ⁺	¹ H NMR
	130	419	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.78 (s, 2 H) 8.35 - 8.55 (m, 1 H) 7.94 (d, J=7.6 Hz, 1 H) 7.70 (s, 1 H) 7.62 (d, J=1.6 Hz, 2 H) 7.50 (dd, J=7.8, 4.8 Hz, 1 H) 6.76 - 6.99 (m, 1 H) 6.57 (s, 1 H) 3.06 (s, 3 H) 2.12 (s, 3 H) 1.88 - 2.06 (m, 3 H).





5 **Example 131: N-(3-chloro-5-(methylsulfonylamino)phenyl)-1-(tetrahydro-2H-pyran-4-yl)-1H-pyrazole-4-carboxamide**




- To a solution of 1-(tetrahydro-2H-pyran-4-yl)-1H-pyrazole-4-carboxylic acid (50 mg, 254 μmol) and N-(3-amino-5-chlorophenyl)methanesulfonamide (61.5 mg, 279 μmol) in pyridine (1 mL) was added EDCI (97.3 mg, 508 μmol) at 25 °C. The mixture was stirred at 25°C for 12 h. The mixture was poured into H₂O (50 mL). The mixture was extracted with EA (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (FA) to afford N-(3-chloro-5-(methylsulfonylamino)phenyl)-1-(tetrahydro-2H-pyran-4-yl)-1H-pyrazole-4-carboxamide (54.68 mg, 137 μmol) as off-white solid. LCMS: MS (ESI) Retention time: 0.738 min, (M+1)⁺ = 399.1. ¹H NMR (400 MHz, DMSO-d₆) δ = 10.02 (s, 2H), 8.44 (s, 1H), 8.07 (s, 1H), 7.76 - 7.45 (m, 2H), 6.92 (t, J = 1.8 Hz, 1H), 4.60 - 4.39 (m, 1H), 3.98 (br dd, J = 2.4, 11.8 Hz, 2H), 3.50 - 3.44 (m, 2H), 3.06 (s, 3H), 2.06 - 1.89 (m, 4H).


Examples 132-136

The compounds listed in the following table were prepared using a procedure similar to that described for Example:

Structure	Example No.	MS (ESI) [M+H] ⁺	¹ H NMR
	132	393	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.51 (s, 1H), 10.22 - 9.93 (m, 1H), 9.17 (d, J = 0.6 Hz, 1H), 7.85 (s, 1H), 7.74 - 7.62 (m, 3H), 7.56 (d, J = 9.4 Hz, 1H), 6.97 (t, J = 1.8 Hz, 1H), 3.07 (s, 3H), 2.74 (q, J = 7.6 Hz, 2H), 1.28 (t, J = 7.6 Hz, 3H)
	133	436	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.58 (s, 1 H) 8.79 (s, 1 H) 8.26 (s, 1 H) 7.62 - 7.75 (m, 3 H) 7.49 (d, J=9.4 Hz, 1 H) 6.97 (t, J=2.0 Hz, 1 H) 3.01 - 3.12 (m, 3 H) 2.81 - 2.92 (m, 6 H) 2.76 (q, J=7.6 Hz, 2 H) 1.26 (t, J=7.6 Hz, 3 H)
	134	365	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.57 (s, 2H), 9.26 (s, 1H), 8.12 (s, 1H), 7.89 - 7.58 (m, 5H), 7.12 - 6.78 (m, 1H), 3.08 (s, 3H)
	135	365	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.57 (s, 1 H) 10.10 (br s, 1 H) 8.15 (d, J=2.0 Hz, 1 H) 7.88 (d, J=7.6 Hz, 1 H) 7.62 - 7.75 (m, 3 H) 7.40 (t, J=7.6 Hz, 1 H) 7.10 (d, J=2.0 Hz, 1 H) 6.98 (t, J=1.6 Hz, 1 H) 3.08 (s, 3 H)

	136	442	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.98 (s, 1H), 8.44 (s, 1H), 8.05 (s, 1H), 7.74 - 7.47 (m, 2H), 6.91 (t, J = 1.8 Hz, 1H), 4.50 - 4.37 (m, 1H), 4.05 (br d, J = 11.4 Hz, 2H), 3.04 (s, 3H), 2.99 - 2.78 (m, 2H), 2.09 - 2.00 (m, 2H), 1.86 - 1.72 (m, 2H), 1.43 (s, 9H)
---	-----	-----	--

Example 137: N-(3-chloro-5-(methylsulfonylamido)phenyl)-3-(pyridin-2-yl)isothiazole-5-carboxamide



- 5 Step 1: To a solution of picolinamide (1.5 g, 12.2 mmol) in toluene (15 mL) was added carbonochloridic hypochlorous thioanhydride (2.38 g, 18.2 mmol) at 25 °C. The mixture was stirred at 60°C for 16 h. The mixture was concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (PE: EA = 10:1~1:1) to afford desired compound. 5-(pyridin-2-yl)-1,3,4-oxathiazol-2-one (550 mg, 3.05 mmol) was
- 10 obtained as white solid. ¹H NMR (400 MHz, DMSO-d₆) δ = 8.85 - 8.72 (m, 1H), 8.16 - 7.98 (m, 2H), 7.67 (ddd, J = 1.8, 4.9, 6.9 Hz, 1H)


Step 2: To a solution of 5-(pyridin-2-yl)-1,3,4-oxathiazol-2-one (450 mg, 2.49 mmol) in xylene (5 mL) was added ethyl prop-2-ynoate (1.21 g, 12.4 mmol). The mixture was stirred at 160 °C for 16 h. The mixture was concentrated in vacuo to afford a residue. The residue was

purified by column chromatography (PE: EA = 10:1~1:1) to afford desired compound. Ethyl 3-(pyridin-2-yl)isothiazole-5-carboxylate (330 mg, 1.4 mmol) was obtained as brown solid. LCMS: MS (ESI) Retention time: 0.761 min, (M) + = 235.0. ¹H NMR (400 MHz, CHLOROFORM-d) δ = 8.78 - 8.65 (m, 1H), 8.56 (s, 1H), 8.24 (d, J = 7.8 Hz, 1H), 7.84 (dt, J = 1.8, 7.8 Hz, 1H), 7.35 (ddd, J = 1.0, 4.9, 7.5 Hz, 1H), 4.48 - 4.40 (m, 2H), 1.44 (t, J = 7.2 Hz, 3H)

Step 3: To a solution of ethyl 3-(pyridin-2-yl)isothiazole-5-carboxylate (320 mg, 1.36 mmol) in a mixture of MeOH (5 mL), THF (3 mL) and H₂O (1 mL) was added LiOH. H₂O (285 mg, 6.8 mmol). The mixture was stirred at 25 °C for 2 h. The mixture was poured into 1N HCl solution (10 mL). The mixture was extracted with EA (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give crude product. 3-(pyridin-2-yl)isothiazole-5-carboxylic acid (200 mg, 969 μ mol, crude) was obtained as yellow solid.

Step 4: To a solution of 3-(pyridin-2-yl)isothiazole-5-carboxylic acid (50 mg, 242 μ mol) and compound 7 (69.2 mg, 314 μ mol) in pyridine (1 mL) was added EDCI (92.7 mg, 484 μ mol) at 25 °C. The mixture was stirred at 25°C for 2 h. The mixture was poured into H₂O (50 mL) and extracted with EA (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (TFA) to afford N-(3-chloro-5-(methylsulfonamido)phenyl)-3-(pyridin-2-yl)isothiazole-5-carboxamide (57.79 mg, 141 μ mol) as yellow solid. LCMS: MS (ESI) Retention time: 0.793 min, (M+1) + = 409.0. ¹H NMR (400 MHz, DMSO-d₆) δ = 10.94 (s, 1H), 10.15 (s, 1H), 8.94 (s, 1H), 8.74 (d, J = 4.6 Hz, 1H), 8.23 (d, J = 7.8 Hz, 1H), 8.01 (dt, J = 1.8, 7.6 Hz, 1H), 7.78 - 7.67 (m, 2H), 7.54 (dd, J = 5.4, 7.0 Hz, 1H), 7.13 - 6.91 (m, 1H), 3.09 (s, 3H)

Example 138: N-(3-chloro-5-(methylsulfonamido)phenyl)-3-(pyridin-2-yl)isothiazole-5-carboxamide



Step 1: Sodium hydride (723 mg, 18.1 mmol) was added to EtOH (150 mL) at 0 °C, followed by diethyl oxalate (2.41 g, 16.5 mmol). Then, a solution of 1-(2-(benzyloxy)phenyl)ethan-1-one (3.75 g, 16.5 mmol) in EtOH (20 mL) was added to the reaction dropwise at 0 °C, after the addition, the reaction was stirred at 25 °C for 16 hr. The reaction mixture was concentrated to remove solvent, the residue was diluted with water (100 mL), extracted with EA (50 mL*3), washed with brine (50 mL), dried over Na₂SO₄, and concentrated. ethyl 4-(2-(benzyloxy)phenyl)-2,4-dioxobutanoate (4 g, 11.8 mmol) was obtained as colorless oil.




LCMS: MS (ESI) Retention time: 0.982, (M+1) + = 326.9

Step 2: To a solvent of EtOH (30 mL) was added NaH (183 mg, 60% purity, 4.59 mmol) at 0 °C, the mixture was stirred at 25 °C for 5 min. Then ethyl 4-(2-(benzyloxy)phenyl)-2,4-dioxobutanoate (1.5 g, 4.59 mmol) was added to above mixture, followed by methylhydrazine sulfate (661 mg, 4.59 mmol). The mixture was stirred at 25 °C for 25 min, and 60 °C for 2 h. The mixture was cooled to 25 °C and poured into water (35 mL). The pH value of mixture was adjusted to 4 - 5 by 1 N (HCl) and the mixture was extracted with EA (40


- mL*3). The combined organic layers were washed with brine (25 mL*2) and dried over Na₂SO₄. The mixture was filtered, and the filtrate was concentrated in vacuo to give a residue. The residue was purified by silica gel column chromatography (Petroleum ether/Ethyl acetate = 100/1 to 1/1). ethyl 3-(2-(benzyloxy)phenyl)-1-methyl-1H-pyrazole-5-carboxylate (0.9 g, 2.67 mmol, 58.4% yield) was obtained as lightly yellow oil. Also obtained ethyl 5-(2-(benzyloxy)phenyl)-1-methyl-1H-pyrazole-3-carboxylate. LCMS: MS (ESI) Retention time: 0.947 min, (M+1) + = 337.2. ¹H NMR (400 MHz, CHLOROFORM-d) δ = 7.32 - 7.39 (m, 1 H) 7.15 - 7.29 (m, 6 H) 6.95 - 7.04 (m, 2 H) 6.74 (s, 1 H) 5.01 (s, 2 H) 4.35 (q, J=7.1 Hz, 2 H) 3.71 (s, 3 H) 1.34 (t, J=7.2 Hz, 3 H).
- Step 3: To a solution of ethyl 3-(2-(benzyloxy)phenyl)-1-methyl-1H-pyrazole-5-carboxylate (500 mg, 1.48 mmol) in a mixed solvent (MeOH: THF: H₂O = 1:1:1, 15 mL) was added LiOH (185 mg, 4.43 mmol). The mixture was stirred at 25 °C for 2 h. The pH value of mixture was adjusted to 3 - 4. The mixture was partitioned between water (20 mL) and EtOAc (30 mL). The aqueous layer was separated and extracted with EtOAc (30 mL*2). The combined organic phases were washed with brine (25 mL*2), dried over anhydrous Na₂SO₄, filtered, and the filtrate was evaporated in vacuo to give 5-(2-(benzyloxy)phenyl)-1-methyl-1H-pyrazole-3-carboxylic acid (450 mg, 1.45 mmol, 98.6% yield) as yellow solid. ¹H NMR (400 MHz, METHANOL-d₄) δ = 7.45 - 7.56 (m, 1 H) 7.22 - 7.39 (m, 7 H) 7.07 - 7.15 (m, 1 H) 6.76 (s, 1 H) 5.16 (s, 2 H) 3.72 (s, 3 H).
- Step 4: To a solution of 5-(2-(benzyloxy)phenyl)-1-methyl-1H-pyrazole-3-carboxylic acid (220 mg) in Py (6 mL) was added N-(3-amino-5-chlorophenyl)methanesulfonamide (188 mg, 856 μ mol) and EDCI (272 mg, 1.42 mmol). The mixture was stirred at 25 °C for 15 h. The mixture was partitioned between water (30 mL) and EtOAc (30 mL). The aqueous layer was separated and extracted with EtOAc (30 mL*2). The combined organic phases were washed with 1 N HCl (20 mL*3) and brine (25 mL*2), dried over anhydrous Na₂SO₄, filtered, and the filtrate was evaporated in vacuo to give a residue. The residue was purified by Prep-HPLC and lyophilized. 5-(2-(benzyloxy)phenyl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-1H-pyrazole-3-carboxamide (282.58 mg, 0.553 mmol, 98.80 % purity, 84.2% yield) was obtained as off-white solid. LCMS: MS (ESI) Retention time: 0.916 min, (M+1) + = 511.2. ¹H NMR (400 MHz, DMSO-d₆) δ = 10.33 (s, 1 H) 9.89 - 10.21 (m, 1 H) 7.68 - 7.78 (m, 2 H) 7.46 - 7.54 (m, 1 H) 7.25 - 7.41 (m, 7 H) 7.10 (t, J=7.6 Hz, 1 H) 6.91 (t, J=1.6 Hz, 1 H) 6.79 (s, 1 H) 5.18 (s, 2 H) 3.74 (s, 3 H) 3.08 (s, 3 H).

Examples 139-141

The compounds listed in the following table were prepared using a procedure similar to that described for example 138:

Structure	Example No.	MS (ESI) [M + H] ⁺	¹ H NMR
	139	511	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.51 (s, 2H), 7.91 (dd, J = 1.4, 7.6 Hz, 1H), 7.64 (d, J = 1.6 Hz, 2H), 7.58 - 7.47 (m, 3H), 7.38 (t, J = 7.6 Hz, 2H), 7.33 - 7.25 (m, 2H), 7.17 (d, J = 8.4 Hz, 1H), 7.07 - 6.94 (m, 2H), 5.32 (s, 2H), 4.14 (s, 3H), 3.07 (s, 3H)
	140	406	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.41 (s, 1 H) 8.73 (d, J=4.4 Hz, 1 H) 7.86 - 8.09 (m, 2 H) 7.69 - 7.81 (m, 2 H) 7.42 - 7.51 (m, 1 H) 7.35 (s, 1 H) 6.93 (t, J=1.6 Hz, 1 H) 4.28 (s, 3 H) 3.10 (s, 3 H)
	141	406	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.54 (s, 2H), 8.63 (d, J = 4.4 Hz, 1H), 7.97 (d, J = 7.8 Hz, 1H), 7.92 - 7.84 (m, 1H), 7.77 (s, 1H), 7.71 (br d, J = 5.8 Hz, 2H), 7.40 - 7.33 (m, 1H), 6.98 (s, 1H), 4.19 (s, 3H), 3.08 (s, 3H)

Example 142: N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(1-phenylethyl)-1H-pyrazole-4-carboxamide




- Step 1: To a solution of ethyl 1H-pyrazole-4-carboxylate (1.0 g, 7.13 mmol) in DMF (15 mL) was added NaH (355 mg, 8.91 mmol, 60% purity) at 0 °C. The mixture was stirred at 0 °C for 0.5 h. (1-bromoethyl)benzene (1.64 g, 8.91 mmol) was added above mixture at 0 °C. The mixture was stirred at 25 °C for 2 h. The mixture was poured into water (80 mL) was extracted with EA (40 mL*3). The combined organic phase was washed with brine (25 mL*3), dried with anhydrous Na₂SO₄, filtered and concentrated in vacuum to give a residue.
- The residue was purified by silica gel column chromatography (Petroleum ether/Ethyl acetate = 5/1). Compound 3 (1.67 g, 6.83 mmol, 95.9% yield) was obtained as colorless oil. LCMS: MS (ESI) Retention time: 0.819 min, (M) + = 245.0. ¹H NMR (400 MHz, CHLOROFORM-d) δ = 7.85 (d, J = 16.4 Hz, 2H), 7.33 - 7.21 (m, 3H), 7.17 - 7.12 (m, 2H), 5.45 (q, J = 7.2 Hz, 1H), 4.20 (q, J = 7.2 Hz, 2H), 1.84 (d, J = 7.2 Hz, 3H), 1.26 (t, J = 7.2 Hz, 3H)
- Step 2: To a solution of ethyl 1-(1-phenylethyl)-1H-pyrazole-4-carboxylate (1.0 g, 4.09 mmol) in a mixture solvent (15 mL, THF/MeOH/H₂O = 1:1:1) was added lithium (1+) hydrate hydroxide (551 mg, 12.2 mmo). The mixture was stirred at 60 °C for 15 h. The mixture was cooled to 25°C and poured into water (20 mL). The pH value of mixture was adjusted to 3 - 4 by 1 N HCl and it was extracted with EA (30 mL*3). The combined organic layers were washed with brine (15 mL*2) and dried over Na₂SO₄. The mixture was filtered, and the filtrate was concentrated in vacuo to give a crude. The crude was used in next step without further purification. 1-(1-phenylethyl)-1H-pyrazole-4-carboxylic acid (880 mg, 4.06

mmol, 99.5% yield) was obtained as white solid. LCMS: MS (ESI) Retention time: 0.715 min, (M+1) + = 217.1

Step 3: To a solution of 1-(1-phenylethyl)-1H-pyrazole-4-carboxylic acid (150 mg, 0.694 mol) and in N-(3-amino-5-chlorophenyl)methanesulfonamide (183 mg, 832 ummol) in
 5 pyridine (5 mL) was added EDCI (264 mg, 1.38 mmol). The mixture was stirred at 25 °C for 5 h. The mixture was poured into water (40 mL), and was extracted with EtOAc (30 mL*3). The combined organic layers were washed with brine (15 mL*3) and dried over Na₂SO₄. The filtrate was filtered and concentrated in vacuum to give a residue. The residue was purified by
 silica gel column chromatography (Petroleum ether/Ethyl acetate = 50/1 - 1/1). The crude
 10 was re-purified by reversed-phase chromatography (H₂O-0.225% FA- MeCN condition) and lyophilized. N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(1-phenylethyl)-1H-pyrazole-4-carboxamide (188.2 mg, 0.454 mmol, 99.77 % purity, 64.4% yield) was obtained as white solid. LCMS: MS (ESI) Retention time: 0.812 min, (M+1) + = 419.2. ¹H NMR (400 MHz, DMSO-d₆) δ = 10.04 (s, 1H), 8.51 (s, 1H), 8.07 (s, 1H), 7.65 (d, J = 1.6 Hz, 1H), 7.55 (d, J =
 15 1.6 Hz, 1H), 7.40 - 7.24 (m, 5H), 6.91 (d, J = 1.6 Hz, 1H), 5.69 (q, J = 7.2 Hz, 1H), 3.05 (s, 3H), 1.94 - 1.73 (m, 3H)


Example 143: N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2,2-difluoroethyl)-5-(3-methylpyridin-2-yl)-1H-pyrrole-3-carboxamide



20 Step 1: To a solution of ethyl 2-chloro-5-(3-methylpyridin-2-yl)-1H-pyrrole-3-carboxylate (650 mg, 2.45 mmol) in DMF (6 mL) was added sodium hydride (146 mg, 3.67 mmol) at 0 °C. The mixture was stirred at 0 °C for 0.5 h. To the mixture was added 2, 2 -difluoroethyl trifluoromethanesulfonate (575 mg, 2.69 mmol) at 0 °C. The mixture was stirred at 25 °C for

- 1h. The mixture was poured to H₂O (50 mL). The mixture was extracted with EA (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (PE: EA = 10:1~0:1) to afford desired compound. 2-chloro-1-(2,2-difluoroethyl)-5-(3-methylpyridin-2-yl)-1H-pyrrole-3-carboxylic acid (300 mg, 1 mmol) was obtained as yellow solid. LCMS: MS (ESI) Retention time: 0.634 min, (M) + = 301.0. ¹H NMR (400 MHz, DMSO-d₆) δ = 12.77 - 12.24 (m, 1H), 8.49 (d, J = 3.6 Hz, 1H), 7.79 (d, J = 7.4 Hz, 1H), 7.32 (dd, J = 4.8, 7.7 Hz, 1H), 6.82 (s, 1H), 6.52 - 6.01 (m, 1H), 4.73 (dt, J = 3.2, 15.2 Hz, 2H), 2.37 (s, 3H)
- 10 Step 2: To a solution of 2-chloro-1-(2,2-difluoroethyl)-5-(3-methylpyridin-2-yl)-1H-pyrrole-3-carboxylic acid (300 mg, 997 μmol) in MeOH (5 mL) was added Pd/C (200 mg, 187 μmol) at 25 °C. The mixture was stirred at 25°C for 14 h under hydrogen (15 Psi). The mixture was filtered and filter cake was washed with MeOH (50 mL). The filtrate was concentrated in vacuo to afford crude product. 1-(2,2-difluoroethyl)-5-(3-methylpyridin-2-yl)-1H-pyrrole-3-
- 15 carboxylic acid (280 mg) was obtained as colorless oil. LCMS: MS (ESI) Retention time: 0.354 min, (M) + = 267.0
- Step 3: To a solution of 1-(2,2-difluoroethyl)-5-(3-methylpyridin-2-yl)-1H-pyrrole-3-carboxylic acid (100 mg, 375 μmol) and 1-(2,2-difluoroethyl)-5-(3-methylpyridin-2-yl)-1H-pyrrole-3-carboxylic acid (99.3 mg, 450 μmol) in pyridine (1 mL) was added EDCI (143 mg,
- 20 750 μmol) at 25 °C. The mixture was stirred at 25°C for 14 h. The mixture was poured to H₂O (50 mL). The mixture was extracted with EA (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC and lyophilized. N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(hydroxymethyl)-1-(pyridin-2-yl)-1H-pyrrole-3-
- 25 carboxamide (95.7 mg, 0.226 mmol, 99.4 purity, 61.7% yield) was obtained as white solid. LCMS: MS (ESI) Retention time: 0.683 min, (M-1) - = 419.0. ¹H NMR (400 MHz, DMSO-d₆) δ = 9.95 (s, 1H), 8.69 - 8.49 (m, 1H), 8.18 - 8.14 (m, 1H), 8.11 - 8.00 (m, 1H), 7.84 (d, J = 8.4 Hz, 1H), 7.77 - 7.70 (m, 1H), 7.65 (s, 1H), 7.51 - 7.41 (m, 1H), 6.91 (t, J = 1.6 Hz, 1H), 6.83 (d, J = 1.6 Hz, 1H), 5.32 (br t, J = 5.6 Hz, 1H), 4.60 (br d, J = 4.8 Hz, 2H), 3.06 (s, 3H)

Example 144: tert-butyl 6-(4-((3-chloro-5-(methylsulfonamido)phenyl)carbamoyl)-1H-pyrazol-1-yl)-2-azaspiro[3.3]heptane-2-carboxylate




- Step 1: To a solution of ethyl 1H-pyrazole-4-carboxylate (100 mg, 0.713 mmol) and tert-butyl 6-hydroxy-2-azaspiro[3.3]heptane-2-carboxylate (174 mg, 820 μ mol) in THF (2 mL) was added PPh₃ (243 mg, 927 μ mol) and DIAD (187 mg, 927 μ mol) at 0 °C slowly. The mixture was stirred at 25 °C for 15 h. The mixture was poured into water (20 mL) and extracted with EA (30 mL*3). The combined organic layers were washed with brine (15 mL*2) and dried over Na₂SO₄. The mixture was filtered, and the filtrate was concentrated in vacuo to give a residue. The residue was purified by column chromatography on silica gel using petroleum ether/ethylacetate (10:1 to 2:1) as eluent. tert-butyl 6-(4-(ethoxycarbonyl)-1H-pyrazol-1-yl)-2-azaspiro[3.3]heptane-2-carboxylate (170 mg, 0.507 mmol, 71.1% yield) was obtained as colorless oil. LCMS: MS (ESI) Retention time: 0.899 min, (M+1) + = 336.3. ¹H NMR (400 MHz, CHLOROFORM-d) δ = 7.98 (s, 1H), 7.85 (s, 1H), 7.81 (s, 1H), 4.65 - 4.50 (m, 1H), 4.30 - 4.18 (m, 3H), 3.96 (s, 2H), 3.89 (s, 2H), 2.67 (d, J = 8.0 Hz, 4H), 1.37 (s, 9H), 1.32 - 1.23 (m, 4H)

- Step 2: To a solution of tert-butyl 6-(4-(ethoxycarbonyl)-1H-pyrazol-1-yl)-2-azaspiro[3.3]heptane-2-carboxylate (170 mg, 0.506 mmol) in THF (2 mL) and MeOH (2 mL) was added a solution of lithium hydroxide hydrate (63.7 mg, 1.52 mmol) in water (2 mL). The mixture was stirred at 25 °C for 15 h. The mixture was poured into water (20 mL). The pH value of mixture was adjusted to 3 - 4 by 0.5 N HCl aqueous solution. It was extracted with EA (30 mL*3). The combined organic layers were washed with brine (15 mL*2) and

dried over Na_2SO_4 . The mixture was filtered, and the filtrate was concentrated in vacuo to give a crude. 1-(2-(tert-butoxycarbonyl)-2-azaspiro[3.3]heptan-6-yl)-1H-pyrazole-4-carboxylic acid (140 mg, 0.455 mmol, 90.3% yield) was obtained as white solid. LCMS: MS (ESI) Retention time: 0.818 min, $(\text{M}+1)^+ = 308.2$

- 5 Step 3: To a solution of 1-(2-(tert-butoxycarbonyl)-2-azaspiro[3.3]heptan-6-yl)-1H-pyrazole-4-carboxylic acid (140 mg, 0.455 mmol) and N-(3-amino-5-chlorophenyl)methanesulfonamide (110 mg, 501 μmol) in pyridine (4 mL) was added EDCI (200 mg, 1.04 mmol). The mixture was stirred at 25 °C for 3 h. The mixture was poured into water (20 mL) and extracted with EA (25 mL*3). The combined organic layers were washed
- 10 with 0.5 N HCl (15 mL*2), brine (15 mL*2) and dried over Na_2SO_4 . The mixture was filtered, and the filtrate was concentrated in vacuo to give residue. The residue was purified by prep-HPLC and lyophilized. tert-butyl 6-(4-((3-chloro-5-(methylsulfonamido)phenyl)carbamoyl)-1H-pyrazol-1-yl)-2-azaspiro[3.3]heptane-2-
- 15 LCMS: MS (ESI) Retention time: 0.840 min, $(\text{M}+1)^+ = 510.2$. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ = 10.01 (s, 1H), 8.42 (s, 1H), 8.06 (s, 1H), 7.64 (t, J = 1.6 Hz, 1H), 7.54 (t, J = 1.6 Hz, 1H), 6.90 (t, J = 2.0 Hz, 1H), 4.97 - 4.70 (m, 1H), 4.05 - 3.72 (m, 4H), 3.04 (s, 3H), 2.73 - 2.59 (m, 4H), 1.37 (s, 9H).

20 **Example 145: N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(piperidin-4-yl)-1H-pyrazole-4-carboxamide**





- To a solution of tert-butyl 4-(4-((3-chloro-5-(methylsulfonamido)phenyl)carbamoyl)-1H-pyrazol-1-yl)piperidine-1-carboxylate (200 mg, 401 μmol) in DCM (4 mL) was added trifluoroacetic acid (744 mg, 6.52 mmol) at 25 °C. The mixture was stirred at 25 °C for 2 h.
- 25 The mixture was concentrated in vacuo to afford a residue. The residue was purified by prep-HPLC (NH_4HCO_3) to afford desired compound. N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(piperidin-4-yl)-1H-pyrazole-4-carboxamide (41.59 mg, 104 μmol) was obtained as white solid. LCMS: MS (ESI) Retention time: 0.722 min, $(\text{M}+1)^+ = 425.2$. ^1H NMR (400 MHz,


DMSO-d₆) δ = 9.97 (s, 1H), 8.41 (s, 1H), 8.04 (s, 1H), 7.62 (t, J = 1.8 Hz, 1H), 7.52 (t, J = 1.8 Hz, 1H), 6.89 (t, J = 1.8 Hz, 1H), 4.29 (br s, 1H), 3.02 (s, 5H), 2.82 - 2.58 (m, 2H), 2.00 (br d, J = 10.6 Hz, 2H), 1.88 - 1.73 (m, 2H)

Examples 146-147

- 5 The compounds listed in the following table were prepared using a procedure similar to that described for example 145:

Structure	Example No.	MS (ESI) [M + H] ⁺	¹ H NMR
	146	410	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.12 (s, 1 H) 8.45 (s, 2 H) 8.06 (s, 1 H) 7.47 - 7.70 (m, 2 H) 6.89 (t, J=1.8 Hz, 1 H) 4.75 - 4.90 (m, 1 H) 4.01 (br s, 4 H) 3.02 (s, 3 H) 2.71 - 2.83 (m, 2 H) 2.57 - 2.70 (m, 2 H)
	147	476	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.28 (s, 1H), 9.36 (s, 1H), 8.25 (s, 1H), 8.21 (s, 1H), 8.13 (d, J = 6.0 Hz, 1H), 7.72 (t, J = 1.8 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 7.34 (d, J = 2.4 Hz, 1H), 6.94 (t, J = 2.0 Hz, 1H), 6.90 (dd, J = 6.1, 2.4 Hz, 1H), 3.43 (t, J = 5.2 Hz, 4H), 3.07 (s, 3H), 2.92 (t, J = 5.1 Hz, 4H).


Example 148: 1-(4-(4-acetylpiperazin-1-yl)pyridin-2-yl)-N-(3-chloro-5-methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide



To a stirred solution of N-(3-chloro-5-methanesulfonamidophenyl)-1-[4-(piperazin-1-yl)pyridin-2-yl]-1H-pyrazole-4-carboxamide (90 mg, 0.189 mmol) and Ac₂O (21.1 mg, 0.206 mmol) in DCM (5 mL) was added Et₃N (57.3 mg, 0.567 mmol) dropwise at 0 °C.

Then the mixture was stirred for 1 h at room temperature. The mixture was concentrated and purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (0-100%) to afford 1-[4-(4-acetylpiperazin-1-yl)pyridin-2-yl]-N-(3-chloro-5-methanesulfamidophenyl)-1H-pyrazole-4-carboxamide (14.6 mg, 0.0282 mmol) as a white solid. LCMS (ESI) $[M + H]^+$: 518. ^1H NMR (400 MHz, DMSO- d_6) δ 10.25 (s, 1H), 9.36 (s, 1H), 8.25 (s, 1H), 8.14 (d, J = 6.0 Hz, 1H), 7.69 (t, J = 1.9 Hz, 1H), 7.60 (t, J = 1.9 Hz, 1H), 7.34 (d, J = 2.4 Hz, 1H), 6.95 – 6.87 (m, 2H), 3.60 (s, 4H), 3.52 (d, J = 5.7 Hz, 2H), 3.46 (t, J = 5.5 Hz, 2H), 3.04 (s, 3H), 2.05 (s, 3H).


Example 149: N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(5-(4-isobutyrylpiperazin-1-yl)pyridin-2-yl)-1H-pyrazole-4-carboxamide



- Step 1: A solution of ethyl 1-(5-bromopyridin-2-yl)-1H-pyrazole-4-carboxylate (350 mg, 1.18 mmol), Pd(AcO)₂ (52.9 mg, 0.236 mmol), BINAP (293 mg, 0.472 mmol), Cs₂CO₃ (1.15 g, 3.54 mmol) and tert-butyl piperazine-1-carboxylate (240 mg, 1.29 mmol) in Toluene (5 mL) was stirred for 2 h at 110 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with Petroleum ether /EtOAc (0-100%) to afford tert-butyl 4-{6-[4-(ethoxycarbonyl)-1H-pyrazol-1-yl]pyridin-3-yl} piperazine-1-carboxylate (270 mg, 0.672 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 402
- Step 2: To a stirred solution of tert-butyl 4-{6-[4-(ethoxycarbonyl)-1H-pyrazol-1-yl]pyridin-3-yl}piperazine-1-carboxylate (270 mg, 0.672 mmol) in EtOH (3 mL) and H₂O (3 mL) was added LiOH (160 mg, 6.71 mmol). Then the mixture was stirred for 2 h at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 1-(5-{4-[(tert-butoxy) carbonyl]piperazin-1-yl} pyridin-2-yl)-1H-pyrazole-4-carboxylic acid (180 mg, 0.482 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 374
- Step 3: To a stirred solution of 1-(5-{4-[(tert-butoxy) carbonyl]piperazin-1-yl}pyridin-2-yl)-1H-pyrazole-4-carboxylic acid (180 mg, 0.482 mmol), TCFH (270 mg, 0.964 mmol) and NMI (118 mg, 1.44 mmol) in ACN (5 mL) was added N-(3-amino-5-chlorophenyl)methanesulfonamide (116 mg, 0.530 mmol). Then the mixture was stirred for 1 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford tert-butyl 4-(6-{4-[(3-chloro-5-methanesulfonamidophenyl)carbamoyl]-1H-pyrazol-1-yl}pyridin-3-yl) piperazine-1-carboxylate (220 mg, 0.381 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 576
- Step 4: To a stirred solution of tert-butyl 4-(6-{4-[(3-chloro-5-methanesulfonamidophenyl)carbamoyl]-1H-pyrazol-1-yl} pyridin-3-yl)piperazine-1-carboxylate (220 mg, 0.381 mmol) in HCl/dioxane (5 mL) at 0 °C for 1 h. The resulting mixture was concentrated under vacuum. The mixture was basified to pH 9 with NaOH. The mixture was concentrated and purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (0-100%) to afford N-(3-chloro-5-methanesulfonamidophenyl)-1-[5-(piperazin-1-yl)pyridin-2-yl]-1H-pyrazole-4-carboxamide (140 mg, 0.294 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 476
- Step 5: To a stirred solution of N-(3-chloro-5-methanesulfonamidophenyl)-1-[5-(piperazin-1-yl)pyridin-2-yl]-1H-pyrazole-4-carboxamide (140 mg, 0.294 mmol), TCFH (164 mg, 0.588

mmol) and NMI (72.4 mg, 0.882 mmol) in ACN (5 mL) was added 2-methylpropanoic acid (28.4 mg, 0.323 mmol). Then the mixture was stirred for 1 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford N-(3-chloro-5-methanesulfonamidophenyl)-1-{5-[4-(2-methylpropanoyl)piperazin-1-yl]pyridin-2-yl}-1H-pyrazole-4-carboxamide (13.5 mg, 0.0247 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 546. ¹H NMR (300 MHz, DMSO-d₆) δ 10.26 (s, 1H), 10.08 (s, 1H), 9.28 (s, 1H), 8.24 (d, J = 3.1 Hz, 2H), 7.84 (d, J = 8.9 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.69 – 7.60 (m, 2H), 6.94 (t, J = 1.9 Hz, 1H), 3.67 (s, 4H), 3.26 (s, 4H), 3.07 (s, 3H), 2.94 (p, J = 6.7 Hz, 1H), 1.03 (d, J = 6.7 Hz, 6H).

Example 150: N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(trans-2-hydroxycyclohexyl)-1H-pyrazole-4-carboxamide



stereochemistry arbitrarily assigned

Step 1: To a solution of ethyl 1H-pyrazole-4-carboxylate (2 g, 14.2 mmol) and 7-oxabicyclo[4.1.0]heptane (2.09 g, 21.3 mmol) in DMF (20 mL) was added cesium carbonate (13.8 g, 42.6 mmol) at 25 °C. The mixture was stirred at 100°C for 16 h. The mixture was poured to H₂O (50 mL). The mixture was extracted with EA (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (PE: EA = 10:1~1:1) to afford ethyl 1-(trans-2-hydroxycyclohexyl)-1H-pyrazole-4-carboxylate (1.8 g, 7.55 mmol) as colorless oil. LCMS: MS (ESI) Retention time: 0.706 min, (M) + = 239.0. ¹H NMR (400 MHz, DMSO-d₆) δ = 8.24 (s, 1H), 7.83 (s, 1H), 4.81

(d, J = 5.8 Hz, 1H), 4.21 (q, J = 7.0 Hz, 2H), 3.93 (ddd, J = 4.4, 9.8, 11.8 Hz, 1H), 3.68 (td, J = 4.6, 9.8 Hz, 1H), 2.05 - 1.61 (m, 6H), 1.30 - 1.24 (m, 5H)

Step 2: To a solution of ethyl 1-(trans-2-hydroxycyclohexyl)-1H-pyrazole-4-carboxylate (200 mg, 839 μ mol) in MeOH (3 mL) and THF (2 mL) and H₂O (1 mL) was added lithium

5 hydroxide hydrate (105 mg, 2.51 mmol) at 25 °C. The mixture was stirred at 25°C for 16 h.

The mixture was poured into 1N HCl (20 mL). The mixture was extracted with EA (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 1-(trans-2-hydroxycyclohexyl)-1H-pyrazole-4-carboxylic acid (150 mg, 713 μ mol) as white solid. LCMS: MS (ESI) Retention

10 time: 0.512 min, (M) + = 211.0


Step 3: To a solution of 1-trans-2-hydroxycyclohexyl)-1H-pyrazole-4-carboxylic acid (80 mg, 380 μ mol) and N-(3-amino-5-chlorophenyl)methanesulfonamide (100 mg, 455 μ mol) in pyridine (2 mL) was added EDCI (145 mg, 760 μ mol) at 25 °C. The mixture was stirred at 25°C for 14 h. The mixture was poured to H₂O (50 mL). The mixture was extracted with EA

15 (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC to afford desired compound. N-(3-chloro-5-

(methylsulfonamido)phenyl)-1-trans-2-hydroxycyclohexyl)-1H-pyrazole-4-carboxamide (113.15 mg, 274 μ mol) was obtained as yellow solid. LCMS: MS (ESI) Retention time: 0.749

20 min, (M+1) + = 413.1. ¹H NMR (400 MHz, DMSO-d₆) δ = 10.03 (d, J = 17.0 Hz, 2H), 8.34 (s, 1H), 8.04 (s, 1H), 7.80 - 7.47 (m, 2H), 6.92 (t, J = 1.8 Hz, 1H), 4.03 - 3.87 (m, 1H), 3.75 - 3.59 (m, 1H), 3.07 (s, 3H), 2.01 - 1.64 (m, 5H), 1.32 (br s, 3H)

Example 151: N-(3-chloro-5-(methylsulfonamido)phenyl)-1-((1s,4s)-4-hydroxycyclohexyl)-1H-pyrazole-4-carboxamide



- Step 1: To a solution of (1r,4r)-cyclohexane-1,4-diol (1 g, 8.6 mmol) and 1H-imidazole (614 mg, 9.03 mmol) in DMF (10 mL) and DCM (20 mL) was added a solution of tert-butyl (chloro) diphenylsilane (2.36 g, 8.6 mmol) in DCM (5 mL). The mixture was stirred at 25 °C for 16 h. The mixture was poured into water (20 mL) and extracted with EA (30 mL*3). The combined organic layers were washed with brine (15 mL*2) and dried over Na₂SO₄. The mixture was filtered, and the filtrate was concentrated in vacuo to give residue. The residue was purified by column chromatography on silica gel using petroleum ether/ethylacetate (10:1 to 2:1) as eluent. (1r,4r)-4-((tert-butyldiphenylsilyl)oxy)cyclohexan-1-ol (1.4 g, 3.94 mmol, 46% yield) was obtained as colorless oil. LCMS: MS (ESI) Retention time: 1.031min, (M+1) + = 337.4. ¹H NMR (400 MHz, CHLOROFORM-d) δ = 7.69 (d, J = 7.2 Hz, 4H), 7.50 - 7.34 (m, 6H), 3.79 - 3.62 (m, 2H), 1.97 - 1.87 (m, 2H), 1.82 (br dd, J = 3.6, 13.2 Hz, 2H), 1.51 - 1.39 (m, 2H), 1.32 - 1.15 (m, 3H), 1.07 (s, 9H).


- Step 2: To a solution of ethyl 1H-pyrazole-4-carboxylate (0.4 g, 2.85 mmol, 1 eq) and (1r,4r)-4-((tert-butyldiphenylsilyl)oxy)cyclohexan-1-ol (1.1 g, 3.13 mmol) in THF (15 mL) was added triphenylphosphine (1.15 g, 3.56 mol), DIAD (0.719 g, 3.56 mmol) at 0 °C. The mixture was stirred at 25 °C for 15 h. The mixture was extracted with EtOAc (40 mL*3). The combined organic layers were washed with brine (25 mL*2), dried over Na₂SO₄. The mixture was cooled to 25 °C and concentrated in vacuo to give a residue. The residue was purified by flash silica gel chromatography. ethyl 1-((1s,4s)-4-((tert-butyldiphenylsilyl)oxy)cyclohexyl)-1H-pyrazole-4-carboxylate (0.82 g, 1.72 mmol, 60.7 % yield) was obtained as colorless oil. ¹H NMR (400 MHz, CHLOROFORM-d) δ = 7.92 (s, 1 H) 7.82 - 7.88 (m, 1 H) 7.60 (dd, J=7.6, 1.2 Hz, 4 H) 7.25 - 7.43 (m, 6 H) 4.24 (q, J=7.2 Hz, 2 H) 3.92 - 4.04 (m, 1 H) 2.21 (qd, J=12.4, 3.2 Hz, 2 H) 1.91 (br dd, J=12.4, 3.2 Hz, 2 H) 1.76 (br dd, J=12.8, 2.8 Hz, 2 H) 1.36 - 1.47 (m, 1 H) 1.35 - 1.38 (m, 1 H) 1.29 (t, J=7.1 Hz, 3 H) 1.03 (s, 9 H).
- Step 3: To a solution of ethyl 1-((1s,4s)-4-((tert-butyldiphenylsilyl)oxy)cyclohexyl)-1H-pyrazole-4-carboxylate (810 mg, 1.69 mmol) in THF (5 mL) and MeOH (5 mL) was added a solution of LiOH.H₂O (212 mg, 5.06 mmol) in H₂O (5 mL). The mixture was stirred at 60 °C for 15 h. The mixture was cooled to 25 °C, the pH of the mixture was adjusted to 3 - 4 by 2 N HCl. The mixture was extracted with EtOAc (30 mL*3). The combined organic layers were washed with brine (15 mL*2), dried over Na₂SO₄. And the filtrate was concentrated in vacuo to give 1-((1s,4s)-4-((tert-butyldiphenylsilyl)oxy)cyclohexyl)-1H-pyrazole-4-carboxylic acid (730 mg, 1.62 mmol, 96.3% yield) as white solid. LCMS: MS (ESI) Retention time: 0.826 min, (M+1) + = 449.1. ¹H NMR (400 MHz, CHLOROFORM-d) δ = 8.24 (s, 1H), 7.83 (s, 1H), 7.69 - 7.56 (m, 4H), 7.52 - 7.33 (m, 6H), 4.21 (tt, J = 3.6, 11.2 Hz, 1H), 4.01 - 3.98 (m, 1H), 2.37 - 2.20 (m, 2H), 1.84 - 1.76 (m, 2H), 1.69 (br dd, J = 2.4, 13.2 Hz, 2H), 1.57 - 1.44 (m, 2H), 1.08 - 0.99 (m, 9H)
- Step 4: To a solution of 1-((1s,4s)-4-((tert-butyldiphenylsilyl)oxy)cyclohexyl)-1H-pyrazole-4-carboxylic acid (240 mg, 0.535 mmol) and N-(3-amino-5-chlorophenyl)methanesulfonamide (129 mg, 588 μmol) in pyridine (5 mL) was added EDCI (203 mg, 1.06 mmol). The mixture was stirred at 25 °C for 3 h. The mixture was poured into water (20 mL) and extracted with EA (25 mL*3). The combined organic layers were washed with 0.5 N HCl (15 mL*2), brine (15 mL*2) and dried over Na₂SO₄. The mixture was filtered, and the filtrate was concentrated in vacuo to give residue. The residue was purified by column chromatography on silica gel using petroleum ether/ethylacetate (50:1 to 2:1) as eluent. 1-((1s,4s)-4-((tert-butyldiphenylsilyl)oxy)cyclohexyl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide (300 mg, 0.461 mmol, 86.4%


yield) was obtained as colorless oil. LCMS: MS (ESI) Retention time: 1.075 min, (M+1) + = 651.3. ¹H NMR (400 MHz, DMSO-d₆) δ = 10.21 - 9.91 (m, 2H), 8.42 (s, 1H), 8.06 (s, 1H), 7.70 - 7.39 (m, 12H), 6.93 (t, J = 1.6 Hz, 1H), 4.24 (ddd, J = 3.6, 7.6, 11.1 Hz, 1H), 3.06 (s, 3H), 2.37 - 2.13 (m, 2H), 1.92 - 1.82 (m, 2H), 1.72 (br d, J = 10.6 Hz, 2H), 1.65 - 1.47 (m, 2H), 1.06 (s, 9H).

Step 5: To a solution of 1-((1s,4s)-4-((tert-butyldiphenylsilyl)oxy)cyclohexyl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide (200 mg, 0.307 mmol) in THF (5 mL) was added TBAF (399 ul, 1 M in THF) dropwise at 25 °C slowly. The mixture was stirred at 60°C for 18 h. The mixture was poured into water (20 mL) and extracted with EA (20 mL*3). The combined organic layers were washed with 0.5 N HCl (10 mL), brine (25 mL*2) and dried over Na₂SO₄. The mixture was filtered, and the filtrate was concentrated in vacuo to give residue. The residue was purified by flash chromatography (silica gel, PE/EA = 30:1 to 1:0) to give desired product. N-(3-chloro-5-(methylsulfonamido)phenyl)-1-((1s,4s)-4-hydroxycyclohexyl)-1H-pyrazole-4-carboxamide (94.13 mg, 0.228 mmol, 100% purity, 74.6% yield) was obtained as white solid. LCMS: MS (ESI) Retention time: 0.820 min, (M+1) + = 413.1. ¹H NMR (400 MHz, DMSO-d₆) δ = 9.85 - 10.19 (m, 2 H) 8.38 - 8.57 (m, 1 H) 8.01 (s, 1 H) 7.66 (t, J=1.8 Hz, 1 H) 7.57 (t, J=1.8 Hz, 1 H) 6.92 (t, J=1.8 Hz, 1 H) 4.51 (d, J=3.2 Hz, 1 H) 4.22 (tt, J=10.8, 3.6 Hz, 1 H) 3.85 (br d, J=2.4 Hz, 1 H) 3.06 (s, 3 H) 2.06 - 2.24 (m, 2 H) 1.68 - 1.86 (m, 4 H) 1.49 - 1.67 (m, 2 H)


Example 152

The compounds listed in the following table were prepared using a procedure similar to that described for example 151:

Structure	Example No.	MS (ESI) [M + H] ⁺	¹ H NMR
 *stereochemistry arbitrarily assigned*	152a	413	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.78 - 10.17 (m, 2 H) 8.40 (s, 1 H) 8.01 (s, 1 H) 7.66 (t, J=1.6 Hz, 1 H) 7.56 (t, J=1.6 Hz, 1 H) 6.91 (t, J=1.6 Hz, 1 H) 4.66 (d, J=3.2 Hz, 1 H) 4.45 - 4.59 (m, 1 H) 4.08 (br d, J=2.8 Hz, 1 H) 3.05 (s, 3 H) 1.87 - 2.07 (m, 3 H) 1.52 - 1.84 (m, 4 H) 1.38 - 1.50 (m, 1 H)

 *stereochemistry arbitrarily assigned*	152b	413	¹ H NMR (400 MHz, DMSO-d ₆) δ = 9.83 - 10.29 (m, 2 H) 8.39 (s, 1 H) 8.01 (s, 1 H) 7.65 (t, J=1.8 Hz, 1 H) 7.56 (t, J=1.8 Hz, 1 H) 6.91 (t, J=1.8 Hz, 1 H) 4.82 (d, J=4.4 Hz, 1 H) 4.17 - 4.32 (m, 1 H) 3.47 - 3.64 (m, 1 H) 3.05 (s, 3 H) 2.14 - 2.28 (m, 1 H) 1.73 - 2.02 (m, 3 H) 1.50 - 1.68 (m, 2 H) 1.27 - 1.47 (m, 1 H) 1.06 - 1.19 (m, 1 H)
---	------	-----	---

Example 153: N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(2-cis-hydroxycyclohexyl)-1H-pyrazole-4-carboxamide




stereochemistry arbitrarily assigned

- Step 1: To a solution of ethyl 1-(trans-2-hydroxycyclohexyl)-1H-pyrazole-4-carboxylate (1 g, 4.19 mmol) and 4-nitrobenzoic acid (1.04 g, 6.28 mmol) and PPh₃ (2.19 g, 8.38 mmol) was added DIAD (1.69 g, 8.38 mmol) at 0 °C. The mixture was stirred at 25°C for 16 h. The mixture was poured to H₂O (20 mL). The mixture was extracted with EA (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (PE: EA = 10:1~1:1) to afford ethyl 1-(cis-2-((4-nitrobenzoyl)oxy)cyclohexyl)-1H-pyrazole-4-carboxylate (1.6 g, 4.13 mmol) as white solid.
- LCMS: MS (ESI) Retention time: 0.883 min, (M+1) + = 388.1


- Step 2: To a solution of ethyl 1-(cis-2-((4-nitrobenzoyl)oxy)cyclohexyl)-1H-pyrazole-4-carboxylate (1.6 g, 4.13 mmol) in MeOH (20 mL) was added dipotassium carbonate (1.14 g, 8.26 mmol) at 25 °C. The mixture was stirred at 25°C for 2 h. The mixture was poured to H₂O (50 mL). The mixture was extracted with EA (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (PE: EA = 10:1~1:1) to afford ethyl 1-(cis-2-hydroxycyclohexyl)-1H-pyrazole-4-carboxylate (900 mg, 3.77 mmol) as white solid. LCMS: MS (ESI) Retention time: 0.802 min, (M+1) + = 239.2. ¹H NMR (400 MHz, DMSO-d₆) δ = 8.19 (s, 1H), 7.85 (s, 1H), 4.83 (d, J = 4.4 Hz, 1H), 4.81 - 4.77 (m, 2H), 4.24 (td, J = 3.0, 12.2 Hz, 1H), 2.10 (br dd, J = 3.6, 12.2 Hz, 1H), 1.84 - 1.71 (m, 3H), 1.67 - 1.51 (m, 2H), 1.47 - 1.33 (m, 2H), 1.12 (br s, 3H)
- Step 3: To a solution of ethyl 1-(cis-2-hydroxycyclohexyl)-1H-pyrazole-4-carboxylate (200 mg, 893 μmol) in MeOH (3 mL) and THF (2 mL) and H₂O (1 mL) was added LiOH.H₂O (105 mg, 2.51 mmol) at 25 °C. The mixture was stirred at 25°C for 16 h. The mixture was poured to 0.5N HCl (10 mL). The mixture was extracted with EA (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 1-(cis-2-hydroxycyclohexyl)-1H-pyrazole-4-carboxylic acid (150 mg, 713 μmol) as white solid. LCMS: MS (ESI) Retention time: 0.475 min, (M+1) + = 211.0
- Step 4: To a solution of 1-(cis-2-hydroxycyclohexyl)-1H-pyrazole-4-carboxylic acid (100 mg, 475 μmol) and N-(3-amino-5-chlorophenyl)methanesulfonamide (115 mg, 522 μmol) in pyridine (2 mL) was added EDCI (182 mg, 950 μmol) at 25 °C. The mixture was stirred at 2 h. The mixture was poured to H₂O (50 mL). The mixture was extracted with EA (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC to give N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(cis-2-hydroxycyclohexyl)-1H-pyrazole-4-carboxamide (60.96 mg, 147 μmol) as white solid. LCMS: MS (ESI) Retention time: 0.760 min, (M+1) + = 413.0. ¹H NMR (400 MHz, DMSO-d₆) δ = 9.99 (s, 1H), 8.41 (s, 1H), 7.99 (s, 1H), 7.76 - 7.47 (m, 2H), 6.91 (t, J = 1.8 Hz, 1H), 4.89 (br s, 1H), 4.29 - 4.18 (m, 1H), 4.04 (br s, 1H), 3.05 (s, 3H), 2.16 - 2.03 (m, 1H), 1.91 - 1.74 (m, 3H), 1.69 - 1.53 (m, 2H), 1.49 - 1.34 (m, 2H)

Example 154

The compounds listed in the following table were prepared using a procedure similar to that described for example 153:

Structure	Example No.	MS (ESI) [M + H] ⁺	¹ H NMR
 *stereochemistry arbitrarily assigned*	154	413	¹ H NMR (400 MHz, DMSO-d ₆) δ (400 MHz, DMSO-d ₆) δ 9.88 - 10.26 (m, 2 H) 8.38 (s, 1 H) 8.02 (s, 1 H) 7.65 (s, 1 H) 7.55 (s, 1 H) 6.90 (s, 1 H) 4.68 (d, J=4.0 Hz, 1 H) 4.19 (tt, J=11.6, 3.6 Hz, 1 H) 3.42 - 3.57 (m, 1 H) 3.04 (s, 3 H) 1.86 - 2.09 (m, 4 H) 1.69 - 1.85 (m, 2 H) 1.27 - 1.46 (m, 2 H).

Example 155: N-(3-chloro-5-(methanesulfonylamino)phenyl)-1-methyl-2-phenyl-1H-imidazole-4-carboxamide




- 5 Step 1: To a solution of methyl 1-methyl-1H-imidazole-4-carboxylate (100 mg, 713 μmol) and iodobenzene (188 mg, 926 μmol) in dioxane (2 mL) was added triphenylphosphine (37.2 mg, 142 μmol), DBU (216 mg, 1.42 mmol), Pd(OAc)₂ (16 mg, 71.3 μmol) and CuI (270 mg, 1.42 mmol) at 25 °C. The mixture was stirred at 120°C under microwave for 2 h. The mixture was filtered and filter cake was washed with EA (20 mL). The filtrate was concentrated in
- 10 vacuo to afford a residue. The residue was purified by column chromatography (PE: EA = 10:1~0:1) to afford desired compound. methyl 1-methyl-2-phenyl-1H-imidazole-4-carboxylate (900 mg, 4.16 mmol) was obtained as brown oil. LCMS: MS (ESI) Retention time: 0.816 min, (M) + = 217.1. ¹H NMR (400 MHz, CHLOROFORM-d) δ = 7.72 - 7.69 (m, 1H), 7.69 - 7.65 (m, 2H), 7.48 (dd, J = 1.8, 5.1 Hz, 3H), 3.93 (s, 3H), 3.79 (s, 3H)
- 15 Step 2: To a solution of methyl 1-methyl-2-phenyl-1H-imidazole-4-carboxylate (370 mg, 1.71 mmol) and N-(3-amino-5-chlorophenyl)methanesulfonamide (564 mg, 2.56 mmol) in toluene (5 mL) was added LiHMDS (4.27 mmol) at 20 °C. The mixture was stirred at 20°C for 2 h. The mixture was poured into H₂O (50 mL) and extracted with EA (50 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and
- 20 concentrated under reduced pressure to give a residue. The residue was purified by column


chromatography (PE: EA = 10:1~1:1) to afford desired compound. N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-2-phenyl-1H-imidazole-4-carboxamide (49.96 mg, 123 μ mol) was obtained as yellow solid. LCMS: MS (ESI) Retention time: 0.759 min, (M+1) + = 405.0. ^1H NMR (400 MHz, DMSO- d_6) δ = 10.17 (s, 1H), 10.04 (s, 1H), 8.08 (s, 1H), 7.86 - 7.69 (m, 4H), 7.59 - 7.51 (m, 3H), 6.92 (t, J = 1.8 Hz, 1H), 3.83 (s, 3H), 3.09 (s, 3H)

Example 156

The compounds listed in the following table were prepared using a procedure similar to that described for example 155:

Structure	Example No.	MS (ESI) [M + H] ⁺	^1H NMR
	156	420	^1H NMR (400 MHz, DMSO- d_6) δ 9.81 - 10.22 (m, 2 H) 8.54 (d, J=4.4 Hz, 1 H) 8.01 - 8.07 (m, 1 H) 7.84 (d, J=7.6 Hz, 1 H) 7.74 (d, J=1.2 Hz, 2 H) 7.43 (dd, J=7.6, 4.8 Hz, 1 H) 6.91 (t, J=1.6 Hz, 1 H) 3.76 (s, 3 H) 3.00 - 3.13 (m, 1 H) 3.08 (s, 2 H) 2.48 (s, 3 H)

10 Example 157: N-(3-bromo-5-(methylsulfonamido)phenyl)thiophene-2-carboxamide



Step 1: To a solution of 3-bromo-5-nitroaniline (500 mg, 2.3 mmol) and thiophene-2-carboxylic acid (589 mg, 4.6 mmol) in pyridine (5 mL) was added EDCI (881 mg, 4.6 mmol) at 20 °C. The mixture was stirred at 20°C for 2 h. The mixture was poured into 1 N HCl (100 mL). Then the aqueous layer was extracted with EA (2 $\times$ 100 mL). The combined organic layers were washed with brine (100 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give N-(3-bromo-5-nitrophenyl)thiophene-2-carboxamide (3 g, 9.17 mmol) as yellow solid.

LCMS: MS (ESI) Retention time: 0.876 min, (M) + = 328.9

Step 2: To a solution of N-(3-bromo-5-nitrophenyl)thiophene-2-carboxamide (3 g, 9.17 mmol) in EA (20 mL) and EtOH (10 mL) was added λ^2 -tin (2+) dichloride (13.8 g, 73.3 mmol) at 20 °C. The mixture was stirred at 60°C for 16 h. The mixture was poured to

5 aq. K_2CO_3 (100 mL). Then the aqueous layer was extracted with EA (2 × 200 mL). The combined organic layers were washed with brine (200 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (PE: EA = 5:1~0:1) to afford N-(3-amino-5-bromophenyl)thiophene-2-

10 carboxamide (1.7 g, 5.72 mmol) was obtained as brown oil. LCMS: MS (ESI) Retention

time: 0.728 min, (M) + = 264.0

^1H NMR (400 MHz, $DMSO-d_6$) δ = 10.03 (s, 1H), 8.00 (dd, J = 1.2, 3.8 Hz, 1H), 7.85 (dd, J = 1.2, 4.9 Hz, 1H), 7.22 (dd, J = 3.8, 5.0 Hz, 1H), 7.09 (t, J = 1.8 Hz, 1H), 7.00 (t, J = 1.8 Hz, 1H), 6.49 (t, J = 1.8 Hz, 1H), 5.73 - 5.15 (m, 2H)

Step 3: To a solution of N-(3-amino-5-bromophenyl)thiophene-2-carboxamide (300 mg, 1.00 mmol) in THF (3 mL) was added triethylamine (202 mg, 2.00 mmol) and methanesulfonyl chloride (598 mg, 5.22 mmol) at 0 °C. The mixture was stirred at 0 °C for 1 h. The mixture was poured to aq. $NaHCO_3$ (20 mL). Then the aqueous layer was extracted with EA (2 × 20 mL). The combined organic layers were washed with brine (20 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give N-(3-bromo-5-(N-

15 (methylsulfonyl)methylsulfonamido)phenyl)thiophene-2-carboxamide (300 mg, 661 μ mol) as white solid.


Step 4: To a solution of N-(3-bromo-5-(N-(methylsulfonyl)methylsulfonamido)phenyl)thiophene-2-carboxamide (270 mg, 595 μ mol) in THF (8 mL) was added a solution of KOH (220 mg, 3.92 mmol) in H_2O (4 mL) at 25 °C. The

25 mixture was stirred at 25°C for 3 h. To the mixture was added 1N HCl (10 mL), extracted with EA (20 mL*3), washed with brine (20 mL), dried over Na_2SO_4 , concentrated in vacuo to afford a residue. The residue was purified by prep-HPLC to afford N-(3-bromo-5-(methylsulfonamido)phenyl)thiophene-2-carboxamide (95.56 mg, 254 μ mol) as white solid.

LCMS: MS (ESI) Retention time: 0.777 min, (M+1)⁺ = 374.9

30 ^1H NMR (400 MHz, $DMSO-d_6$) δ = 10.41 (s, 1H), 10.08 (ddd, J = 2.2, 5.2, 11.2 Hz, 1H), 8.04 (dd, J = 1.2, 3.8 Hz, 1H), 7.90 (dd, J = 1.2, 4.8 Hz, 1H), 7.79 (t, J = 1.8 Hz, 1H), 7.67 (t, J = 1.8 Hz, 1H), 7.24 (dd, J = 3.8, 5.0 Hz, 1H), 7.09 (t, J = 1.8 Hz, 1H), 3.06 (s, 3H)

Example 158: N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(2-oxopiperidin-1-yl)-1H-pyrrole-3-carboxamide



Step 1: To a solution of methyl 1H-pyrrole-3-carboxylate (2 g, 15.9 mmol) in DMF (20 mL) was added sodium hydride (951 mg, 23.8 mmol) at 0 °C. The mixture was stirred at 0°C for 0.5 h. To the mixture was added iodomethane (3.37g, 23.8 mmol) at 0°C. The mixture was stirred at 25°C for 2h. The mixture was poured into H₂O (50 mL). Then the aqueous layer was extracted with EA (2 × 50 mL). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give methyl 1-methyl-1H-pyrrole-3-carboxylate (1.8 g, 12.9 mmol, crude) as colorless oil.

Step 2: To a solution of methyl 1-methyl-1H-pyrrole-3-carboxylate (1.8 g, 12.9 mmol) in THF (18 mL) was added 1-bromopyrrolidine-2,5-dione (2.27 g, 12.8 mmol) slowly at 0 °C. The mixture was stirred at 25°C for 0.5 h. The mixture was poured to aq. Na₂SO₃ (40 mL). Then the aqueous layer was extracted with EA (2 × 40 mL). The combined organic layers were washed with brine (40 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (PE: EA = 20:1~1:1) to afford methyl 5-bromo-1-methyl-1H-pyrrole-3-carboxylate (2.4 g, 11 mmol) as white solid.

Step 3: To a solution of methyl 5-bromo-1-methyl-1H-pyrrole-3-carboxylate (1.3 g, 5.96 mmol) and piperidin-2-one (1.17 g, 11.9 mmol) in DMF (10 mL) was added rac-(2R)-pyrrolidine-2-carboxylic acid (137 mg, 1.19 mmol) and λ¹-copper (1+) iodide (226 mg, 1.19 mmol) and dicaesium (1+) carbonate (5.79 g, 17.8 mmol) at 25 °C. The mixture was stirred at 140°C for 2 h under microwave. The mixture was poured into H₂O (50 mL). The mixture was extracted with EA (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (PE: EA = 10:1~1:1) to afford methyl 1-

methyl-5-(2-oxopiperidin-1-yl)-1H-pyrrole-3-carboxylate (300 mg, 1.26 mmol) was obtained as white solid.

LCMS: MS (ESI) Retention time: 0.636 min, (M) + = 237.0

Step 4: To a solution of methyl 1-methyl-5-(2-oxopiperidin-1-yl)-1H-pyrrole-3-carboxylate (200 mg, 846 μ mol) in MeOH (5 mL) and THF (3 mL) and H₂O (1 mL) was added lithium (1+) hydrate hydroxide (177 mg, 4.22 mmol) at 25°C. The mixture was stirred at 15 h. The mixture was poured into 1N HCl (20 mL). The mixture was extracted with EA (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 1-methyl-5-(2-oxopiperidin-1-yl)-1H-pyrrole-3-carboxylic acid (180 mg, 809 μ mol) was obtained as white solid.


LCMS: MS (ESI) Retention time: 0.372 min, (M) + = 222.9

Step 5: To a solution of 1-methyl-5-(2-oxopiperidin-1-yl)-1H-pyrrole-3-carboxylic acid (75 mg, 337 μ mol) and N-(3-amino-5-chlorophenyl)methanesulfonamide (81.6 mg, 370 μ mol) in pyridine (3 mL) was added EDCI (129 mg, 674 μ mol) at 25 °C. The mixture was stirred at 25°C for 1 h. The mixture was poured into H₂O (20 mL). The mixture was extracted with EA (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC to afford N-(3-chloro-5-(methylsulfonamido)phenyl)-1-methyl-5-(2-oxopiperidin-1-yl)-1H-pyrrole-3-carboxamide (68.57 mg, 161 μ mol) as off-white solid.

LCMS: MS (ESI) Retention time: 0.722 min, (M+1) + = 425.2

¹H NMR (400 MHz, DMSO-d₆) δ = 9.72 (s, 2H), 7.69 (t, J = 1.8 Hz, 1H), 7.58 (t, J = 1.8 Hz, 1H), 7.46 (d, J = 2.0 Hz, 1H), 6.87 (t, J = 1.8 Hz, 1H), 6.49 (d, J = 2.0 Hz, 1H), 3.52 (br t, J = 5.4 Hz, 2H), 3.41 (s, 3H), 3.05 (s, 3H), 2.42 (br t, J = 6.0 Hz, 2H), 1.94 - 1.80 (m, 4H)

Example 159: 1-(2-trans-(benzyloxy)cyclobutyl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide



stereochemistry arbitrarily assigned

- 5 Step 1: To a solution of 1,2-bis(trimethylsilyloxy)cyclobut-1-ene (5 g, 21.6 mmol) in HCl/dioxane (20 mL, 4M) was added phenylmethanol (2.8 g, 25.9 mmol) at 25 °C. The mixture was stirred at 80°C for 16 h. The mixture was concentrated in vacuo to afford a residue. The residue was purified by column chromatography (PE: EA = 10:1~1:1) to afford 2-(benzyloxy)cyclobutan-1-one (2.1 g, 11.9 mmol) as colorless oil. ¹H NMR (400 MHz, CHLOROFORM-d) δ = 7.52 - 7.29 (m, 5H), 4.83 - 4.71 (m, 2H), 4.69 - 4.60 (m, 1H), 2.88 - 2.70 (m, 2H), 2.42 - 2.23 (m, 1H), 2.04 - 1.87 (m, 1H)
- 10 Step 2: To a solution of 2-(benzyloxy)cyclobutan-1-one (2.1 g, 11.9 mmol) in MeOH (20 mL) was added sodium boranuide (900 mg, 23.8 mmol) at 0 °C. The mixture was stirred at 25°C for 2 h. The mixture was quenched with sat.NH₄Cl (20 mL) and stirred at 25°C for 15 min. The mixture was extracted with EA (30 mL*3). The combined organic phase was washed with brine (15 mL*2), dried with anhydrous Na₂SO₄, filtered and concentrated in vacuum to give a residue. The residue was purified by silica gel column chromatography (Petroleum ether/Ethyl acetate = 20/1 - 1/1) to afford desired compound. cis-2-(benzyloxy)cyclobutan-1-ol (690 mg, 3.87 mmol) was obtained as colorless oil, which was determined by ¹H NMR. Trans-2-(benzyloxy)cyclobutan-1-ol (690 mg, 3.87 mmol) was obtained as colorless oil, which was determined by ¹H NMR.

cis-2-(benzyloxy)cyclobutan-1-ol:

¹H NMR (400 MHz, CHLOROFORM-d) δ = 7.43 - 7.30 (m, 4H), 4.65 - 4.48 (m, 2H), 4.31 (br dd, J = 2.2, 4.6 Hz, 1H), 4.18 - 4.11 (m, 1H), 2.13 - 1.97 (m, 4H), 1.28 (t, J = 7.2 Hz, 1H)

trans-2-(benzyloxy)cyclobutan-1-ol ¹H NMR (400 MHz, CHLOROFORM-d) δ = 7.44 - 7.29 (m, 5H), 4.68 - 4.49 (m, 2H), 4.21 - 3.99 (m, 1H), 3.90 - 3.71 (m, 1H), 2.21 (s, 1H), 2.11 - 1.91 (m, 2H), 1.49 - 1.29 (m, 2H)

Step 3: To a solution of cis-2-(benzyloxy)cyclobutan-1-ol (690 mg, 3.87 mmol) and ethyl 1H-pyrazole-4-carboxylate (812 mg, 5.8 mmol) and triphenylphosphane (2.03 g, 7.74 mmol) in THF (10 mL) was added DIAD (1.56 g, 774 mmol) at 0 °C. The mixture was stirred at 25°C for 16 h. The mixture was poured to H₂O (50 mL). The mixture was extracted with EA (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (PE: EA = 10:1~1:1) to afford ethyl 1-(trans-2-(benzyloxy)cyclobutyl)-1H-pyrazole-4-carboxylate (690 mg, 2.13 mmol) as colorless oil. LCMS: MS (ESI) Retention time: 0.833 min, (M+1) + = 301.1 ¹H NMR (400 MHz, CHLOROFORM-d) δ = 8.01 - 7.92 (m, 1H), 7.79 (s, 1H), 7.35 - 7.19 (m, 5H), 4.66 - 4.56 (m, 1H), 4.48 (d, J = 1.8 Hz, 2H), 4.45 - 4.38 (m, 1H), 4.32 (q, J = 7.2 Hz, 2H), 2.39 - 2.18 (m, 2H), 1.88 - 1.74 (m, 2H), 1.41 - 1.33 (m, 3H)

Step 4: To a solution of ethyl 1-(trans-2-(benzyloxy)cyclobutyl)-1H-pyrazole-4-carboxylate (120 mg, 399 μmol) in THF (2 mL) and MeOH (3 mL) and H₂O (1 mL) was added lithium hydroxide hydrate (83.5 mg, 1.99 mmol) at 25 °C. The mixture was stirred at 25°C for 16 h. The mixture was poured to 1N HCl (15 mL). The mixture was extracted with EA (30 mL*2). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 1-(trans-2-(benzyloxy)cyclobutyl)-1H-pyrazole-4-carboxylic acid (100 mg, 367 μmol) as colorless oil. LCMS: MS (ESI) Retention time: 0.748 min, (M+1) + = 273.1


Step 5: To a solution of 1-(trans-2-(benzyloxy)cyclobutyl)-1H-pyrazole-4-carboxylic acid (70 mg, 257 μmol) and N-(3-amino-5-chlorophenyl)methanesulfonamide (62.2 mg, 282 μmol) in pyridine (2 mL) was added EDCI (98.5 mg, 514 μmol) at 25 °C. The mixture was stirred at 25°C for 16 h. The mixture was poured to H₂O (20 mL). Then the aqueous layer was extracted with EA (2 × 20 mL). The combined organic layers were washed with brine (20 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC to afford 1-(trans-2-(benzyloxy)cyclobutyl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide (28.48 mg, 59.9 μmol) as white solid.

LCMS: MS (ESI) Retention time: 0.845 min, (M+1) + = 475.2 ¹H NMR (400 MHz, DMSO-d₆) δ = 10.04 (s, 1H), 8.46 (s, 1H), 8.09 (s, 1H), 7.71 - 7.51 (m, 2H), 7.34 - 7.19 (m, 5H), 6.92


(t, J = 1.8 Hz, 1H), 4.80 (br d, J = 7.4 Hz, 1H), 4.47 - 4.31 (m, 3H), 3.05 (s, 3H), 2.28 - 2.13 (m, 2H), 1.99 (br t, J = 9.4 Hz, 1H), 1.76 - 1.62 (m, 1H)

Example 424

The compounds listed in the following table were prepared using a procedure similar to that described for example 159:

Structure	Example No.	MS (ESI) [M + H] ⁺	¹ H NMR
	424	489	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.03 (s, 2H), 8.47 (d, J = 3.5 Hz, 1H), 8.07 (d, J = 3.4 Hz, 1H), 7.75 - 7.55 (m, 3H), 7.37 - 7.13 (m, 6H), 6.97 - 6.71 (m, 1H), 4.74 (td, J = 8.0, 5.2 Hz, 1H), 4.44 (d, J = 3.8 Hz, 2H), 4.18 (q, J = 6.0 Hz, 2H), 3.06 (d, J = 3.0 Hz, 4H), 2.28 - 2.16 (m, 2H), 2.16 - 1.95 (m, 3H), 1.89 - 1.50 (m, 4H).

Example 160: N-(3-chloro-5-(methylsulfonyl)phenyl)-3-(3-(hydroxymethyl)-1H-pyrazol-1-yl)benzamide



- 10 Step 1: To a stirred solution of methyl 3-iodobenzoate (500 mg, 1.90 mmol) in EtOH (6 mL) and H₂O (2 mL) was added LiOH (136 mg, 5.70 mmol). Then the mixture was stirred for 2 h


at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (0-100%, acidic system) to afford 3-iodobenzoic acid (470 mg, 1.89 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 248.93

Step 2: To a stirred solution of 3-iodobenzoic acid (470 mg, 1.89 mmol), CuI (71.8 mg, 0.378 mmol) and (1S,2S)-N1,N2-dimethylcyclohexane-1,2-diamine (107 mg, 0.756 mmol) and methyl 1H-pyrazole-3-carboxylate (285 mg, 2.26 mmol) in DMSO (6 mL) was added Cs₂CO₃ (1.84 g, 5.67 mmol). Then the mixture was stirred for 2 h at 100 °C under N₂ atmosphere. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (0-100%, acidic system) to afford 3-[3-(methoxycarbonyl)-1H-pyrazol-1-yl]benzoic acid (283 mg, 1.14 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 247.06

Step 3: To a stirred solution of 3-[3-(methoxycarbonyl)-1H-pyrazol-1-yl]benzoic acid (283 mg, 1.14 mmol), TCFH (478 mg, 1.71 mmol) and NMI (467 mg, 5.70 mmol) in ACN (5 mL) was added N-(3-amino-5-chlorophenyl) methanesulfonamide (251 mg, 1.14 mmol). Then the mixture was stirred for 2 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (0-100%, acidic system) to afford methyl 1-{3-[(3-chloro-5-methanesulfonamidophenyl)carbamoyl]phenyl}-1H-pyrazole-3-carboxylate (230 mg, 0.512 mmol) as an off-white solid. LCMS (ESI) [M + H]⁺: 449.06

Step 4: To a stirred solution of methyl 1-{3-[(3-chloro-5-methanesulfonamidophenyl)carbamoyl]phenyl}-1H-pyrazole-3-carboxylate (230 mg, 0.512 mmol) in THF (5 mL) was added LiAlH₄ (110 mg, 1.53 mmol) under N₂ atmosphere. Then the mixture was stirred for 2 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (0-100%, acidic system) to afford N-(3-chloro-5-methanesulfonamidophenyl)-3-[3-(hydroxymethyl)-1H-pyrazol-1-yl] benzamide (14.6 mg, 0.0344 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 421.07 ¹H NMR (300 MHz, DMSO-d₆) δ 10.61 (s, 1H), 8.52 (d, J = 2.6 Hz, 1H), 8.34 (s, 1H), 8.03 (d, J = 8.1 Hz, 1H), 7.84 (d, J = 7.7 Hz, 1H), 7.71 (d, J = 1.9 Hz, 2H), 7.65 (t, J = 7.8 Hz, 1H), 6.98 (t, J = 2.0 Hz, 1H), 6.55 (d, J = 2.5 Hz, 1H), 5.26 (s, 1H), 4.55 (s, 2H), 3.08 (s, 3H).


Example 161: N-(3-bromo-5-(methylsulfonamido)phenyl)-1-(2-(hydroxymethyl)phenyl)-1H-pyrazole-4-carboxamide



To a stirred solution of N-(3-bromo-5-methanesulfonamidophenyl)-1-(2-formylphenyl)-1H-pyrazole-4-carboxamide (200 mg, 0.431 mmol) in MeOH (5 mL) was added NaBH₄ (21.1 mg, 0.560 mmol) at 0 °C. Then the mixture was stirred for 1 h at room temperature. The mixture was concentrated and purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (0-100%) to afford N-(3-bromo-5-methanesulfonamidophenyl)-1-[2-(hydroxymethyl)phenyl]-1H-pyrazole-4-carboxamide (25.6 mg, 0.055 mmol) as a white solid.

LCMS (ESI) [M + H]⁺: 465 ¹H NMR (300 MHz, DMSO-d₆) δ 10.15 (s, 1H), 10.09 (s, 1H), 8.73 (s, 1H), 8.29 (s, 1H), 7.82 (t, J = 1.8 Hz, 1H), 7.74 – 7.61 (m, 2H), 7.59 – 7.40 (m, 3H), 7.07 (t, J = 1.9 Hz, 1H), 5.34 (t, J = 5.6 Hz, 1H), 4.47 (d, J = 5.2 Hz, 2H), 3.06 (s, 3H).

Example 162: 1-(3-aminopyridin-2-yl)-N-(3-bromo-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide



Step 1: To a stirred solution of ethyl 1H-pyrazole-4-carboxylate (150 mg, 1.07 mmol), 2-bromo-3-nitropyridine (259 mg, 1.28 mmol) and K₂CO₃ (529 mg, 3.84 mmol) in DMSO (10.00 mL). Then the mixture was stirred for 4 h at 90 °C under nitrogen. The mixture was


concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford ethyl 1-(3-nitropyridin-2-yl)-1H-pyrazole-4-carboxylate (110 mg, 95 %) as a yellow solid. LCMS (ESI) [M + H]⁺: 263

Step 2: To a stirred solution of ethyl 1-(3-nitropyridin-2-yl)-1H-pyrazole-4-carboxylate (110 mg, 0.419 mmol), N-(3-amino-5-bromophenyl) methanesulfonamide (111 mg, 0.419 mmol) and Al(CH₃)₃ (0.5 ml) in toluene (5.00 mL). Then the mixture was stirred for 1.5 h at 80 °C under nitrogen. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford N-(3-bromo-5-methanesulfonamidophenyl)-1-(3-nitropyridin-2-yl)-1H-pyrazole-4-carboxamide (120 mg, 95%) as a white solid. LCMS (ESI) [M + H]⁺: 481

Step 3: To a solution of N-(3-bromo-5-methanesulfonamidophenyl)-1-(3-nitropyridin-2-yl)-1H-pyrazole-4-carboxamide (120 mg, 0.249 mmol) in a solution of methanol and water (15 mL) was added Fe (70 mg, 1.245 mmol) and NH₄Cl (131 mg, 2.49 mmol). The resulting mixture was stirred for 1 hour at 80 °C under N₂ atmosphere. The resulting mixture was filtered. The filtrate was concentrated under reduced pressure to give the crude product. The mixture was concentrated and purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (0-100%) to afford 1-(3-aminopyridin-2-yl)-N-(3-bromo-5-methanesulfonamidophenyl)-1H-pyrazole-4-carboxamide (45.9 mg, 97.661%) as a white solid. LCMS (ESI) [M + H]⁺: 451

¹H NMR (300 MHz, DMSO-d₆) δ 10.28 (s, 1H), 10.06 (s, 1H), 9.31 (d, J = 0.8 Hz, 1H), 8.29 (d, J = 0.8 Hz, 1H), 7.86 (t, J = 1.8 Hz, 1H), 7.77 (dd, J = 4.4, 1.5 Hz, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.35 (dd, J = 8.1, 1.6 Hz, 1H), 7.21 (dd, J = 8.1, 4.4 Hz, 1H), 7.08 (t, J = 1.9 Hz, 1H), 6.41 (s, 2H), 3.07 (s, 3H).

Example 163: N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(hydroxymethyl)-4-(pyridin-2-yl)thiophene-2-carboxamide



- Step 1: To a stirred solution of methyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxylate (600 mg, 2.23 mmol), 2-bromopyridine (527 mg, 3.34 mmol), Pd(PPh₃)Cl₂ (156 mg, 0.223 mmol) and Na₂CO₃ (709 mg, 6.69 mmol) in EtOH (10.00 mL) was added H₂O (2.00 mL). Then the mixture was stirred for 4 h at 80 °C under N₂ atmosphere. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford methyl 4-(pyridin-2-yl)thiophene-2-carboxylate (456 mg, 95 %) as a white solid. LCMS (ESI) [M + H]⁺: 220
- Step 2: To a solution of methyl 4-(pyridin-2-yl)thiophene-2-carboxylate (480 mg, 2.18 mmol) in DMF (15 mL) was added N-Bromosuccinimide (1.75 g, 9.81 mmol) and AcOH (392 mg, 6.54 mmol). The resulting mixture was stirred for 4 hours at 60 °C under N₂ atmosphere. The mixture was concentrated and purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (0-100%) to afford methyl 5-bromo-4-(pyridin-2-yl)thiophene-2-carboxylate (294 mg, 95%) as a yellow solid. LCMS (ESI) [M + H]⁺: 299

Step 3: To a stirred solution of methyl 5-bromo-4-(pyridin-2-yl)thiophene-2-carboxylate (310 mg, 1.03 mmol) in THF (15.00 mL) was added chloro(propan-2-yl) magnesium (126 mg, 1.23 mmol) dropwise at -60 °C under N₂ atmosphere. Then the mixture was stirred for 3 h at -60 °C. After that the mixture was added DMF (751 mg, 10.3 mmol) dropwise at -60 °C.

- 5 Then the mixture was stirred for 8 h at room temperature. The mixture was concentrated and purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (0-100%) to afford methyl 5-formyl-4-(pyridin-2-yl) thiophene-2-carboxylate (142 mg, 95 %) as a yellow solid.

LCMS (ESI) [M + H]⁺: 248


- 10 Step 4: To a stirred solution of methyl 5-formyl-4-(pyridin-2-yl)thiophene-2-carboxylate (150 mg, 0.606 mmol) and LiOH (10 mg, 2.5 mmol) in THF (10.00 mL) was added H₂O (2.00 mL) dropwise. Then the mixture was stirred for 2 h at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to 5-formyl-4-(pyridin-2-yl)thiophene-2-carboxylic acid (104 mg, 95%) as a yellow solid. LCMS (ESI) [M + H]⁺: 234

- 15 Step 5: To a stirred solution of 5-formyl-4-(pyridin-2-yl)thiophene-2-carboxylic acid (110 mg, 0.471 mmol), TCFH (198 mg, 0.706 mmol) and NMI (115 mg, 1.41 mmol) in ACN (5.00 mL) was added N-(3-amino-5-chlorophenyl)methanesulfonamide (103 mg, 0.471 mmol). Then the mixture was stirred for 4 h at room temperature. The mixture was
- 20 concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford N-(3-chloro-5-methanesulfonamidophenyl)-5-formyl-4-(pyridin-2-yl)thiophene-2-carboxamide (85.5 mg, 95 %) as a white solid. LCMS (ESI) [M + H]⁺: 437

- Step 6: To a stirred solution of N-(3-chloro-5-methanesulfonamidophenyl)-5-formyl-4-(pyridin-2-yl)thiophene-2-carboxamide (90 mg, 0.206 mmol) in DCM (10.00 mL) was added
- 25 NaBH₃CN (12.9 mg, 0.206 mmol) dropwise at 0 °C. Then the mixture was stirred for 30 minutes at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford N-(3-chloro-5-methanesulfonamidophenyl)-5-(hydroxymethyl)-4-(pyridin-2-yl)thiophene-2-carboxamide
- 30 (9.6 mg, 97.816%) as an off-white solid. LCMS (ESI) [M + H]⁺: 439

¹H NMR (400 MHz, DMSO-d₆) δ 10.44 (s, 1H), 10.07 (s, 1H), 8.67 (ddd, J = 4.8, 1.8, 0.9 Hz, 1H), 8.48 (s, 1H), 7.94 (td, J = 7.7, 1.8 Hz, 1H), 7.76 – 7.67 (m, 2H), 7.65 (t, J = 1.9 Hz, 1H), 7.35 (ddd, J = 7.5, 4.8, 1.1 Hz, 1H), 6.96 (t, J = 2.0 Hz, 1H), 6.04 (t, J = 5.6 Hz, 1H), 4.99 (d, J = 5.6 Hz, 2H), 3.07 (s, 3H).


Example 164: N-(3-chloro-5-(methylsulfonamido)phenyl)-3-(phenylamino)-1H-pyrazole-5-carboxamide



- Step 1: Under nitrogen, a solution of ethyl 5-bromo-2H-pyrazole-3-carboxylate (1 g, 4.565 mmol, 1 equiv) in DMF (15 mL) was added NaH (164 mg, 6.834 mmol, 1.50 equiv) at 0 °C. The resulting solution was stirred for half an hour at room temperature. Then [2-(chloromethoxy)ethyl]trimethylsilane (910 mg, 5.458 mmol, 1.20 equiv) was added and stirred for one hour at room temperature. The reaction was quenched with water (2 mL). The reaction mixture was diluted with water (50 mL). The resulting solution was extracted with EtOAc (3×50 mL) and the organic layers were combined. The organic layer was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (23%) to afford ethyl 5-bromo-2-[[2-(trimethylsilyl)ethoxy]methyl]pyrazole-3-carboxylate (1.2 g, 75.25%) as a yellow oil. LCMS: (ESI): [M + H]⁺:349
- Step 2: Under nitrogen, a solution of ethyl 5-bromo-2-[[2-(trimethylsilyl)ethoxy]methyl]pyrazole-3-carboxylate (500 mg, 1.431 mmol, 1 equiv),

- aniline (266.62 mg, 2.862 mmol, 2 equiv), Pd(OAc)₂ (16.07 mg, 0.072 mmol, 0.05 equiv) and XantPhos (82.8 mg, 0.143 mmol, 0.10 equiv) in Toluene (5 mL) was added Cs₂CO₃ (1.4 g, 4.297 mmol, 3.00 equiv) at room temperature. The resulting solution was stirred for two hours at 100 °C. The resulting solution was concentrated under vacuum. The residue was
- 5 purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (45%) to afford ethyl 5-(phenylamino)-2-[[2-(trimethylsilyl)ethoxy]methyl]pyrazole-3-carboxylate (230 mg, 44.45%) as a yellow oil. LCMS: (ESI): [M + H]⁺:362
- Step 3: To a solution of ethyl 5-(phenylamino)-2-[[2-(trimethylsilyl)ethoxy]methyl]pyrazole-3-carboxylate (230 mg, 0.636 mmol, 1 equiv) in THF (5 mL) and H₂O (2 mL) was added
- 10 LiOH (76.2 mg, 3.182 mmol, 5.00 equiv) at room temperature. The resulting solution was stirred for one hour at room temperature. The residue was purified by flash chromatography on C₁₈ column gel eluting with ACN/water (0.1%FA) (37%) to afford 5-(phenylamino)-2-[[2-(trimethylsilyl)ethoxy]methyl]pyrazole-3-carboxylic acid (130 mg, 61.28%) as a yellow solid. LCMS: (ESI): [M + H]⁺:334
- 15 Step 4: A solution of 5-(phenylamino)-2-[[2-(trimethylsilyl)ethoxy]methyl]pyrazole-3-carboxylic acid (130 mg, 0.390 mmol, 1 equiv), NMI (64 mg, 0.779 mmol, 2.00 equiv) and TCFH (164.1 mg, 0.585 mmol, 1.50 equiv) in ACN (5 mL) at room temperature. The resulting solution was stirred for 10 minutes at room temperature. Then N-(3-amino-5-chlorophenyl)methanesulfonamide (103.2 mg, 0.468 mmol, 1.20 equiv) was added and
- 20 stirred for one hour at room temperature. The residue was purified by C₁₈ column gel eluting with ACN/water (0.05 NH₄HCO₃) (68%) to afford N-(3-chloro-5-methanesulfonamidophenyl)-5-(phenylamino)-2-[[2-(trimethylsilyl)ethoxy]methyl]pyrazole-3-carboxamide (80 mg, 38.28%) as a yellow solid. LCMS: (ESI): [M + H]⁺:536
- Step 5: To a solution of N-(3-chloro-5-methanesulfonamidophenyl)-5-(phenylamino)-2-[[2-(trimethylsilyl)ethoxy]methyl]pyrazole-3-carboxamide (80 mg, 0.149 mmol, 1 equiv) in
- 25 Et₃SiH (1 mL, 6.192 mmol, 41.50 equiv) was added TFA (1 mL) at 0 °C. The mixture was stirred for one hour at room temperature. The residue was purified by Pre-HPLC on condition: Column: Sunfire prep C18 column, 30*150 mm, 5μm; Mobile Phase A: Water (0.1%FA), Mobile Phase B: ACN; Flow rate: 60 mL/min; Gradient: 40% B to 52% B in 7
- 30 min, 52% B; Wave Length: 254/220 nm; RT1(min): 6.62 to afford N-(3-chloro-5-methanesulfonamidophenyl)-5-(phenylamino)-2H-pyrazole-3-carboxamide (29.6 mg, 48.73%) as a white solid. LCMS: (ESI): [M + H]⁺:406. ¹H NMR (300 MHz, DMSO-d₆) δ 12.98 (s, 1H), 10.21 (d, J = 75.1 Hz, 2H), 8.57 (s, 1H), 7.68 (s, 2H), 7.32 (s, 1H), 7.22 (t, J = 7.6 Hz, 2H), 6.96 (s, 1H), 6.75 (d, J = 9.1 Hz, 2H), 3.08 (s, 3H).


Example 165: N-(3-chloro-5-(methanesulfonamido)phenyl)-5-(piperazine-1-carbonyl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxamide



- Step 1: To a stirred solution of 5-(ethoxycarbonyl)-1H-pyrrole-3-carboxylic acid (320 mg, 1.74 mmol), K_2CO_3 (721 mg, 5.22 mmol), CuBr (249 mg, 1.74 mmol) and 2-bromopyridine (301 mg, 1.91 mmol) in Pyridine (5 mL). Then the mixture was stirred for 5 h at 110 °C. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford 5-(ethoxycarbonyl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxylic acid (120 mg, 0.461 mmol) as a yellow solid. LCMS (ESI) $[M + H]^+$: 261
- Step 2: To a stirred solution of 5-(ethoxycarbonyl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxylic acid (120 mg, 0.461 mmol), TCFH (193 mg, 0.691 mmol) and NMI (113 mg, 1.3 mmol) in ACN (5 mL) was added N-(3-amino-5-chlorophenyl)methanesulfonamide (111 mg, 0.507 mmol). Then the mixture was stirred for 2 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford ethyl 4-[(3-chloro-5-methanesulfonamidophenyl)carbamoyl]-1-(pyridin-2-yl)-1H-pyrrole-2-carboxylate (130 mg, 0.280 mmol) as a yellow solid. LCMS (ESI) $[M + H]^+$: 463

- Step 3: To a stirred solution of ethyl 4-[(3-chloro-5-methanesulfonamidophenyl)carbamoyl]-1-(pyridin-2-yl)-1H-pyrrole-2-carboxylate (130 mg, 0.280 mmol) in EtOH (3 mL) and H₂O (3 mL) was added LiOH (66.8 mg, 2.79 mmol). Then the mixture was stirred for 2 h at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 4-[(3-chloro-5-methanesulfonamidophenyl)carbamoyl]-1-(pyridin-2-yl)-1H-pyrrole-2-carboxylic acid (90.0 mg, 0.206 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 435
- Step 4: To a stirred solution of 4-[(3-chloro-5-methanesulfonamidophenyl)carbamoyl]-1-(pyridin-2-yl)-1H-pyrrole-2-carboxylic acid (90 mg, 0.206 mmol), TCFH (86.6 mg, 0.309 mmol) and NMI (50.7 mg, 0.618 mmol) in ACN (5 mL) was added tert-butyl piperazine-1-carboxylate (42.0 mg, 0.226 mmol). Then the mixture was stirred for 1 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford tert-butyl 4-{4-[(3-chloro-5-methanesulfonamidophenyl) carbamoyl]-1-(pyridin-2-yl)-1H-pyrrole-2-carbonyl}piperazine-1-carboxylate (80.0 mg, 0.132 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 603
- Step 5: To a stirred solution of tert-butyl 4-{4-[(3-chloro-5-methanesulfonamidophenyl)carbamoyl]-1-(pyridin-2-yl)-1H-pyrrole-2-carbonyl}piperazine-1-carboxylate (80 mg, 0.132 mmol) in HCl/dioxane (5 mL) at 0 °C for 1 h. The resulting mixture was concentrated under vacuum. The mixture was basified to pH 9 with NaOH. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford N-(3-chloro-5-methanesulfonamidophenyl)-5-(piperazine-1-carbonyl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxamide (19.1 mg, 0.0379 mmol) as a light yellow solid. LCMS (ESI) [M + H]⁺: 503. ¹H NMR (300 MHz, DMSO-d₆) δ 10.05 (s, 1H), 8.55 – 8.46 (m, 1H), 8.31 – 8.20 (m, 2H), 8.02 (td, J = 7.8, 1.9 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.63 (t, J = 1.9 Hz, 1H), 7.55 (d, J = 8.2 Hz, 1H), 7.48 – 7.38 (m, 1H), 6.99 (d, J = 1.8 Hz, 1H), 6.92 (t, J = 1.9 Hz, 1H), 3.44 (s, 4H), 3.07 (s, 3H), 2.67 (s, 4H), 1.24 (s, 1H).


Example 166: N2-(3-chloro-5-(methanesulfonamido)phenyl)-N4-phenylthiophene-2,4-dicarboxamide



- Step 1: To a stirred solution of N-(3-amino-5-chlorophenyl)-4-bromothiophene-2-carboxamide (100.0 mg, 0.49 mmol, 1.00 equiv) in a mixture of 2 mL DCM and 2 mL C₅H₅N was added MsCl (83.8 mg, 0.74 mmol, 1.50 equiv) dropwise, the resulting solution was stirred for 2 h at room temperature. Then it was quenched with the addition of 10 mL of water and extracted with 3 x 20 mL of ethyl acetate and the organic phase was dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified onto silica gel column chromatography, eluted with 50% of ethyl acetate in petroleum ether to afford 4-bromo-N-(3-chloro-5-methanesulfonamidophenyl) thiophene-2-carboxamide (90.0 mg, 73.1%) as a white solid. LCMS (ESI) [M + H]⁺: 408.90
- Step 2: Into an 8-mL sealed tube, was placed 4-bromo-N-(3-chloro-5-methanesulfonamidophenyl) thiophene-2-carboxamide (90.0 mg, 0.40 mmol, 1.00 equiv), (Ac)₂O (20.0 mg, 0.08 mmol, 0.20 equiv), (COOH)₂ (20.0 mg, 0.08 mmol, 0.20 equiv), Pd(OAc)₂ (12.4 mg, 0.08 mmol, 0.20 equiv), Xantphos (16.3 mg, 0.08 mmol, 0.20 equiv), DIEA (100.0 mg, 1.20 mmol, 3.00 equiv) in 4 mL of DMSO, the mixture was stirred for 2 h at 100 °C under N₂ atmosphere. The mixture was purified onto silica gel column chromatography, eluted with 50% of ethyl acetate in petroleum ether to afford 5-[(3-chloro-5-methanesulfonamidophenyl) carbamoyl]thiophene-3-carboxylic acid (60.0 mg, 73.1%) as a white solid. LCMS (ESI) [M + H]⁺: 374.98

Step 3: Into an 8-mL sealed tube, was placed 5-[(3-chloro-5-methanesulfonamidophenyl) carbamoyl]thiophene-3-carboxylic acid (60.0 mg, 0.36 mmol, 1.00 equiv) and aniline (14.9 mg, 0.36 mmol, 1.00 equiv), NMI (54 mg, 1.08 mmol, 3.00 equiv) in 4 mL of ACN, to this was added TCFH (266.7 mg, 0.7 mmol, 1.30 equiv) and the resulting mixture was stirred for 1 h at room temperature. Then it was quenched with the addition of 10 mL of water and extracted with 3 x 20 mL of ethyl acetate and the combined organic phase was dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified onto silica gel column chromatography, eluted with 50% of ethyl acetate in petroleum ether to afford N2-(3-chloro-5-methanesulfonamidophenyl)-N4-phenylthiophene-2,4-dicarboxamide (10.8 mg, 14.8%) as a white solid. LCMS (ESI) [M + H]⁺: 450.03. ¹H NMR (400 MHz, DMSO-d₆) δ 10.64 (s, 1H), 10.25 (s, 1H), 10.10 (s, 1H), 8.66 (s, 1H), 8.53 (s, 1H), 7.74 (s, 3H), 7.68 (d, J = 17.7 Hz, 1H), 7.37 (s, 2H), 7.12 (s, 1H), 6.97 (s, 1H), 3.39 (s, 1H), 3.08 (s, 3H), 2.69 (s, 1H).

Example 167: N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-(phenoxyethyl)pyridin-2-yl)-1H-pyrazole-4-carboxamide



Step 1: To a stirred solution of (2-bromopyridin-3-yl)methanol (500 mg, 2.65 mmol) and Ms₂O (923 mg, 5.30 mmol) in DCM (20 mL) was added Et₃N (804 mg, 7.95 mmol) dropwise

at 0 °C. Then the mixture was stirred for 2 h at room temperature. The mixture was concentrated and purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (0-100%) to afford (2-bromopyridin-3-yl)methyl methanesulfonate (544 mg, 2.04 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 266

- 5 Step 2: To a stirred solution of (2-bromopyridin-3-yl)methyl methanesulfonate (550 mg, 2.06 mmol) and Phenoxysodium (262 mg, 2.26 mmol) in DMF (10 mL). Then the mixture was stirred for 2 h at room temperature. The mixture was concentrated and purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (0-100%) to afford 2-bromo-3-(phenoxymethyl)pyridine (465 mg, 1.76 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 264


- 10 Step 3: To a stirred solution of 2-bromo-3-(phenoxymethyl)pyridine (470 mg, 1.77 mmol), K₂CO₃ (733 mg, 5.31 mmol), CuI (67.4 mg, 0.354 mmol), ethyl 1H-pyrazole-4-carboxylate (322 mg, 2.30 mmol) and (1S,2S)-N1,N2-dimethylcyclohexane-1,2-diamine (100 mg, 0.708 mmol) in ACN (5 mL). Then the mixture was stirred for 2 h at 90 °C. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford ethyl 1-[3-(phenoxymethyl)pyridin-2-yl]-1H-pyrazole-4-carboxylate (356 mg, 1.10 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 324

- 15 Step 4: To a stirred solution of ethyl 1-[3-(phenoxymethyl)pyridin-2-yl]-1H-pyrazole-4-carboxylate (360 mg, 1.11 mmol) in EtOH (5 mL) and H₂O (5 mL) was added LiOH (265 mg, 11.1 mmol). Then the mixture was stirred for 2 h at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 1-[3-(phenoxymethyl)pyridin-2-yl]-1H-pyrazole-4-carboxylic acid (257 mg, 0.871 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 296

- 20 Step 5: To a stirred solution of 1-[3-(phenoxymethyl)pyridin-2-yl]-1H-pyrazole-4-carboxylic acid (170 mg, 0.575 mmol), TCFH (209 mg, 0.747 mmol) and NMI (141 mg, 1.72 mmol) in ACN (5 mL) was added N-(3-amino-5-chlorophenyl) methanesulfonamide (139 mg, 0.632 mmol). Then the mixture was stirred for 1 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford N-(3-chloro-5-methanesulfonamidophenyl)-1-[3-(phenoxymethyl)pyridin-2-yl]-1H-pyrazole-4-carboxamide (183.0 mg, 0.3675 mmol) as an off-white solid. LCMS (ESI) [M + H]⁺: 498. ¹H NMR (300 MHz, DMSO-d₆) δ 10.32 (s, 1H), 10.09 (s, 1H), 9.29 (s, 1H), 8.56 (dd, J = 4.7, 1.7 Hz, 1H), 8.31 (s, 1H), 8.23 (dd, J = 7.8, 1.7

Hz, 1H), 7.73 (t, J = 1.9 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 7.58 (dd, J = 7.8, 4.7 Hz, 1H), 7.30 (dd, J = 8.6, 7.3 Hz, 2H), 6.96 (ddd, J = 7.0, 4.2, 2.5 Hz, 4H), 5.52 (s, 2H), 3.08 (s, 3H).

Example 168: 1-(3-(benzylamino)pyridin-2-yl)-N-(3-chloro-5-(methylsulfonamido)phenyl)-1H-pyrazole-4-carboxamide



5

- Step 1: To a stirred solution of ethyl 1H-pyrazole-4-carboxylate (200 mg, 1.42 mmol), 2-bromo-3-nitropyridine (288 mg, 1.42 mmol), (1S,2S)-N1,N2-dimethylcyclohexane-1,2-diamine (40.3 mg, 0.284 mmol), CuI (107 mg, 0.568 mmol) and K₂CO₃ (587 mg, 4.26 mmol) in ACN (10.00 mL). Then the mixture was stirred for 4 h at 90 °C under N₂ atmosphere. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford ethyl 1-(3-nitropyridin-2-yl)-1H-pyrazole-4-carboxylate (237 mg, 95 %) as a yellow solid. LCMS (ESI) [M + H]⁺: 263
- Step 2: To a solution of ethyl 1-(3-nitropyridin-2-yl)-1H-pyrazole-4-carboxylate (250 mg, 0.953 mmol) in a solution of methanol and water (15 mL) was added Fe (266 mg, 4.76 mmol) and NH₄Cl (505 mg, 9.53 mmol). The resulting mixture was stirred for 2 hours at 80 °C under N₂ atmosphere. The resulting mixture was filtered. The filtrate was concentrated under
- 10
- 15


reduced pressure to give the crude product. The mixture was concentrated and purified by flash chromatography on silica gel eluting with EtOAc/Petroleum ether (0-100%) to afford ethyl 1-(3-aminopyridin-2-yl)-1H-pyrazole-4-carboxylate (209 mg, 90 %) as a yellow solid. LCMS (ESI) $[M + H]^+$: 233

- 5 Step 3: To a stirred solution of ethyl 1-(3-aminopyridin-2-yl)-1H-pyrazole-4-carboxylate (220 mg, 0.947 mmol), (bromomethyl)benzene (161 mg, 0.947 mmol), K_2CO_3 (130 mg, 0.947 mmol) in DMF (10.00 mL). Then the mixture was stirred for 4 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/ H_2O (5-95%, acidic system) to afford ethyl 1-[3-(benzylamino)pyridin-2-yl]-1H-pyrazole-4-carboxylate (142 mg, 95 %) as a yellow solid. LCMS (ESI) $[M + H]^+$: 323

- 10 Step 4: To a stirred solution of ethyl 1-[3-(benzylamino)pyridin-2-yl]-1H-pyrazole-4-carboxylate (150 mg, 0.465 mmol) and LiOH (10 mg, 2.5 mmol) in THF (10.00 mL) was added H_2O (2.00 mL). Then the mixture was stirred for 2 h at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 1-[3-(benzylamino)pyridin-2-yl]-1H-pyrazole-4-carboxylic acid (95.0 mg, 95 %) as a yellow solid. LCMS (ESI) $[M + H]^+$: 295

- Step 5: To a stirred solution of 1-[3-(benzylamino)pyridin-2-yl]-1H-pyrazole-4-carboxylic acid (100 mg, 0.339 mmol), TCFH (142 mg, 0.508 mmol) and NMI (82.8 mg, 1.01 mmol) in ACN (5.00 mL) was added N-(3-amino-5-chlorophenyl)methanesulfonamide (74.8 mg, 0.339 mmol). Then the mixture was stirred for 4 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/ H_2O (5-95%, acidic system) to afford 1-[3-(benzylamino)pyridin-2-yl]-N-(3-chloro-5-methanesulfonamidophenyl)-1H-pyrazole-4-carboxamide (38.5 mg, 98.283%) as a white solid. LCMS (ESI) $[M + H]^+$: 498. ^1H NMR (300 MHz, $DMSO-d_6$) δ 10.33 (s, 1H), 10.08 (s, 1H), 9.37 (d, $J = 0.9$ Hz, 1H), 8.33 (d, $J = 0.8$ Hz, 1H), 7.95 (t, $J = 6.0$ Hz, 1H), 7.82 – 7.70 (m, 2H), 7.65 (t, $J = 1.9$ Hz, 1H), 7.43 – 7.31 (m, 4H), 7.31 – 7.17 (m, 3H), 6.95 (t, $J = 1.9$ Hz, 1H), 4.53 (d, $J = 5.9$ Hz, 2H), 3.08 (s, 3H).

Example 169: N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(hydroxymethyl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxamide



- Step 1: To a solution of methyl 1H-pyrrole-3-carboxylate (1.0 g, 7.99 mmol) and Cs₂CO₃ (3.87 g, 11.9 mmol) in DMF (10 mL) was added 2-fluoropyridine (1.93 g, 19.9 mmol). The mixture was stirred at 100 °C for 2 h. The mixture was cooled to 25 °C and poured into water (40 mL), and was extracted with EtOAc (30 mL*3). The combined organic layers were washed with brine (20 mL*3), dried over Na₂SO₄. And the filtrate was concentrated in vacuo to give residue. The residue was purified by column chromatography on silica gel using petroleum ether/ethylacetate (100:1 to 3:1) as eluent. methyl 1-(pyridin-2-yl)-1H-pyrrole-3-carboxylate (1.57 g, 7.76 mmol, 97.5% yield) was obtained as white solid. ¹H NMR (400 MHz, CHLOROFORM-d) δ = 8.36 - 8.69 (m, 1 H) 8.07 - 8.25 (m, 1 H) 7.82 (ddd, J=8.4, 7.6, 2.0 Hz, 1 H) 7.51 (dd, J=3.2, 2.4 Hz, 1 H) 7.39 (d, J=8.4 Hz, 1 H) 7.22 (ddd, J=7.6, 4.8, 0.8 Hz, 1 H) 6.78 (dd, J=3.2, 1.6 Hz, 1 H) 3.87 (s, 3 H)
- Step 2: The Vilsmeier reagent was prepared by adding POCl₃ (1.13 g, 7.4 mmol) dropwise to ice cold dry DMF (4 mL) under stirring. The mixture was then stirred for 15 min at 0 °C. To the above Vilsmeier reagent was added methyl 1-(pyridin-2-yl)-1H-pyrrole-3-carboxylate (300 mg, 1.48 mmol) as a solution in DCE (15 mL). Then the mixture was allowed to warm

to 100 °C and was stirred for 1.0 h. The reaction mixture was poured into saturated sodium chloride aqueous (30 mL). The mixture was extracted with dichloromethane (3 × 30 mL), the combined organic phase was washed with water (3 × 20 mL), dried over Na₂SO₄, filtered, and concentrated in vacuo. The crude product was purified by flash chromatography (silica gel, PE/EA = 30:1 to 4:1) to give desired product. methyl 5-formyl-1-(pyridin-2-yl)-1H-pyrrole-3-carboxylate (400 mg, 0.98 mmol, 24.5% yield) was obtained as yellow solid. ¹H NMR (400 MHz, DMSO-d₆) δ = 9.80 (s, 1H), 8.64 - 8.53 (m, 1H), 8.20 (d, J = 1.6 Hz, 1H), 8.05 (dt, J = 1.6, 8.0 Hz, 1H), 7.75 (d, J = 8.0 Hz, 1H), 7.60 - 7.43 (m, 2H), 3.80 (s, 3H).

Step 3: To a solution of methyl 5-formyl-1-(pyridin-2-yl)-1H-pyrrole-3-carboxylate (400 mg, 1.73 mmol) in MeOH (10 mL) was added NaBH₄ (196 mg, 5.19 mmol) at 0 °C slowly. The mixture was stirred at 25 °C for 1 h. The reaction mixture was poured into saturated NH₄Cl aqueous (30 mL) and stirred at 25 °C for 0.5 h. The mixture was extracted with EA (3 × 30 mL), the combined organic phase was washed with brine (2 × 20 mL), dried over Na₂SO₄, filtered, and concentrated in vacuo to give a crude. The crude was purified by silica gel column chromatography (Petroleum ether/Ethyl acetate = 5/1). methyl 5-(hydroxymethyl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxylate (270 mg, 1.16 mmol, 67.3% yield) was obtained as white solid. ¹H NMR (400 MHz, CHLOROFORM-d) δ = 8.58 - 8.48 (m, 1H), 8.04 - 7.91 (m, 1H), 7.74 (d, J = 1.6 Hz, 1H), 7.47 (d, J = 8.0 Hz, 1H), 7.39 - 7.31 (m, 1H), 6.72 (d, J = 1.6 Hz, 1H), 4.54 (s, 2H), 3.85 (s, 3H).


Step 4: To a solution of methyl 5-(hydroxymethyl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxylate (270 mg, 1.16 mmol) in THF (5 mL) and MeOH (5 mL) was added a solution of LiOH.H₂O in water (5 mL). The mixture was stirred at 60 °C for 15 h. The mixture was cooled to 25 °C, the pH value of the mixture was adjusted to 3 - 4 by 2 N HCl. The mixture was extracted with EtOAc (30 mL*3). The combined organic layers were washed with brine (10 mL*2), dried over Na₂SO₄. And the filtrate was concentrated in vacuo to give a crude. 5-(hydroxymethyl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxylic acid (230 mg, 1.05 mmol, 90.9% yield) was obtained as white solid. LCMS: MS (ESI) Retention time: 0.508 min, (M-17) + = 201.2. ¹H NMR (400 MHz, DMSO-d₆) δ = 12.05 (br s, 1H), 8.53 (dd, J = 1.2, 4.8 Hz, 1H), 8.07 - 7.95 (m, 1H), 7.89 - 7.72 (m, 2H), 7.46 - 7.37 (m, 1H), 6.55 (d, J = 2.0 Hz, 1H), 5.22 (t, J = 5.6 Hz, 1H), 4.55 (d, J = 5.6 Hz, 2H)

Step 5: To a solution of 5-(hydroxymethyl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxylic acid (80 mg, 0.3666 mmol) and N-(3-amino-5-chlorophenyl)methanesulfonamide (88.9 mg, 403 μmol) in Py (2 mL) was added EDCI (140 mg, 733 μmol). The mixture was stirred at 25 °C for 3 h. The mixture was poured into water (20 mL) and extracted with EA (25 mL*3). The

combined organic layers were washed with 0.5 N HCl (15 mL*2), brine (15 mL*2) and dried over Na₂SO₄. The mixture was filtered, and the filtrate was concentrated in vacuo to give residue. The residue was purified by prep-HPLC and lyophilized. N-(3-chloro-5-

(methylsulfonamido)phenyl)-5-(hydroxymethyl)-1-(pyridin-2-yl)-1H-pyrrole-3-carboxamide
 5 (95.7 mg, 0.226 mmol, 99.4 purity, 61.7% yield) was obtained as white solid. LCMS: MS (ESI) Retention time: 0.683 min, (M-1) - = 419.0. ¹H NMR (400 MHz, DMSO-d₆) δ = 9.95 (s, 1H), 8.69 - 8.49 (m, 1H), 8.18 - 8.14 (m, 1H), 8.11 - 8.00 (m, 1H), 7.84 (d, J = 8.4 Hz, 1H), 7.77 - 7.70 (m, 1H), 7.65 (s, 1H), 7.51 - 7.41 (m, 1H), 6.91 (t, J = 1.6 Hz, 1H), 6.83 (d, J = 1.6 Hz, 1H), 5.32 (br t, J = 5.6 Hz, 1H), 4.60 (br d, J = 4.8 Hz, 2H), 3.06 (s, 3H)

10 **Example 170: N-(3-chloro-5-(methylsulfonamido)phenyl)-5-methyl-1-(pyridin-2-yl)-1H-pyrrole-3-carboxamide**



Step 1: A solution of 2-fluoropyridine (300 mg, 3.090 mmol, 1 equiv), methyl 5-methyl-1H-pyrrole-3-carboxylate (430 mg, 3.090 mmol, 1 equiv) and K₂CO₃ (1279 mg, 9.270 mmol, 3 equiv) in DMSO (5 ml) was stirred for 1 h at 60 °C. The mixture was allowed to cool down to room temperature. The reaction mixture was diluted with water (50 mL). The resulting mixture was extracted with EtOAc (3*50 mL). The combined organic layers were washed with EtOAc (100 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with Petroleum ether / EtOAc (1:1) to afford methyl 5-methyl-1-(pyridin-2-yl)pyrrole-3-carboxylate (280 mg, 41.91%) as a white solid. LCMS (ESI) [M+H]⁺: 217.09.

Step 2: To a stirred solution of methyl 5-methyl-1-(pyridin-2-yl)pyrrole-3-carboxylate (50 mg, 0.231 mmol, 1 equiv) and N-(3-amino-5-chlorophenyl)methanesulfonamide (51 mg, 0.231 mmol, 1 equiv) in PhMe (2 ml) was added Al(Me)₃ (0.6 ml) at 0 °C. The resulting mixture was


s stirred for 1 h at 80 °C. The mixture was allowed to cool down to room temperature. The reaction was quenched by the addition of Water (2 mL) at 0 °C. The precipitated solids were collected by filtration and washed with acetonitrile (3*6 mL). The resulting mixture was concentrated under reduced pressure. The residue was purified by reverse phase flash chromatography with the following conditions: C18 column; mobile phase, MeCN in Water (0.1% FA), 10% to 50% gradient in 10 min; detector, UV 254 nm. 35% to afford *N*-(3-chloro-5-methanesulfonylphenyl)-5-methyl-1-(pyridin-2-yl)pyrrole-3-carboxamide (30.7 mg, 32.75%) as a white solid. LCMS (ESI) $[M+H]^+$: 405.10. ^1H NMR (400 MHz, DMSO- d_6) δ 10.02 (s, 1H), 9.87 (s, 1H), 8.58 (dt, J = 4.8, 1.5 Hz, 1H), 8.03 (td, J = 7.8, 2.0 Hz, 2H), 7.72 (t, J = 1.9 Hz, 1H), 7.67 – 7.60 (m, 2H), 7.49 – 7.42 (m, 1H), 6.89 (t, J = 2.0 Hz, 1H), 6.58 (dd, J = 2.1, 1.1 Hz, 1H), 3.06 (s, 3H), 2.39 (d, J = 1.0 Hz, 3H).

Examples 425-426

The compounds listed in the following table were prepared using a procedure similar to that described for example 170:

Structure	Example No.	MS (ESI) $[M + H]^+$	^1H NMR
	425	429.95	^1H NMR (400 MHz, DMSO- d_6) δ 10.02 (s, 1H), 9.93 (s, 1H), 9.04 (d, J = 2.2 Hz, 1H), 8.52 (dd, J = 8.6, 2.3 Hz, 1H), 8.14 (d, J = 2.0 Hz, 1H), 7.87 (d, J = 8.5 Hz, 1H), 7.67 (dt, J = 29.8, 1.9 Hz, 2H), 6.91 (t, J = 2.0 Hz, 1H), 6.64 (d, J = 1.8 Hz, 1H), 3.06 (s, 3H), 2.46 (s, 3H).
	426	473.95	^1H NMR (400 MHz, DMSO- d_6) δ 10.07 (d, J = 37.7 Hz, 2H), 9.32 (d, J = 0.9 Hz, 2H), 8.57 (d, J = 2.1 Hz, 1H), 7.73 (t, J = 1.9 Hz, 1H), 7.66 (t, J = 1.9 Hz, 1H), 6.91 (t, J = 2.0 Hz, 1H), 6.63 (t, J = 1.7 Hz, 1H), 3.06 (s, 3H), 2.63 (d, J = 1.0 Hz, 3H).

Example 171: N-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoropyrimidin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide



Step 1: To a solution of methyl 1H-pyrrole-3-carboxylate (500 mg, 3.99 mmol) and B₂pin₂ (2.02 g, 7.98 mmol) in THF (10 mL) were added Di-μ-methoxybis(1,5-cyclooctadiene)diiridium(I) (264 mg, 0.399 mmol, CAS:12148-71-9) and 4,4'-Di-*tert*-butyl-2,2'-bipyridine (107 mg, 0.399 mmol). After stirring for 2 h at 100 °C under a nitrogen atmosphere, the resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with Petroleum ether/EtOAc (1/1) to afford methyl 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrole-3-carboxylate (350 mg, 1.39 mmol) as a brown solid. LCMS (ESI) [M + H]⁺:252.

Step 2: A mixture of methyl 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrole-3-carboxylate (200 mg, 0.796 mmol), 2-bromo-5-fluoropyrimidine (169 mg, 0.955 mmol), Pd(dppf)Cl₂ (65.0 mg, 0.0796 mmol) and K₃PO₄ (504 mg, 2.38 mmol) in THF (2 mL) and H₂O (0.2 mL) was stirred for 2 hours at 80 °C under N₂ atmosphere. The reaction mixture was extracted with ethyl acetate. The organic phase was concentrated under reduced pressure. The residue was purified by flash chromatography on silica gel eluting with 30% of ethyl acetate in petroleum ether to afford methyl 5-(5-fluoropyrimidin-2-yl)-1H-pyrrole-3-carboxylate (135 mg, 76.7%) as a white solid. LCMS [M+H]⁺: 222

Step 3: To a stirred solution of methyl 5-(5-fluoropyrimidin-2-yl)-1H-pyrrole-3-carboxylate (135 mg, 0.610 mmol), CH₃I (129 mg, 0.915 mmol) in DMF (1 ml) was added NaH (73.2 mg, 1.83 mmol) at 0 °C. The resulting mixture was stirred for 2 hours at room temperature. Then it was quenched with water, the mixture was concentrated and the residue was purified by


y reverse phase flash chromatography eluting with 60% of acetonitrile in water (0.1% NH₄HCO₃) to afford methyl 5-(5-fluoropyrimidin-2-yl)-1-methyl-1*H*-pyrrole-3-carboxylate (60.0 mg, 41.9%) as a white solid. LCMS [M+H]⁺: 236

Step 4: To a stirred solution of methyl 5-(5-fluoropyrimidin-2-yl)-1-methyl-1*H*-pyrrole-3-carboxylate (60 mg, 0.255 mmol) in EtOH (1 mL) and water (1 mL) was added LiOH (53.3 mg, 1.27 mmol). The resulting mixture was stirred for 2 hours at room temperature. Then it was concentrated, and the pH value of the residue solution was adjusted to 3 with 1M HCl (aq). The product was precipitated from the solution. The mixture was filtered and the filter cake was washed with water (2 mL) to afford 5-(5-fluoropyrimidin-2-yl)-1-methyl-1*H*-pyrrole-3-carboxylic acid (35.0 mg, 62.3%) as a white solid. LCMS [M+H]⁺: 222


Step 5: To a mixture of 5-(5-fluoropyrimidin-2-yl)-1-methyl-1*H*-pyrrole-3-carboxylic acid (35 mg, 0.158 mmol) and *N*-(3-amino-5-chlorophenyl)methanesulfonamide (41.7 mg, 0.189 mmol) in ACN (0.5 mL) was added TCFH (66.3 mg, 0.237 mmol) and NMI (38.8 mg, 0.474 mmol) at room temperature for 2 hours. Then it was concentrated, the residue was purified by reverse phase flash chromatography eluting with 50% of acetonitrile in water (0.1% FA) to afford *N*-(3-chloro-5-(methylsulfonamido)phenyl)-5-(5-fluoropyrimidin-2-yl)-1-methyl-1*H*-pyrrole-3-carboxamide (11.8 mg, 17.6%) as a white solid. LCMS [M+H]⁺: 424. ¹H NMR (300 MHz, DMSO-*d*₆) δ 10.02 (s, 1H), 8.98 (d, *J* = 0.8 Hz, 2H), 7.80 (t, *J* = 1.7 Hz, 2H), 7.77 – 7.69 (m, 2H), 6.97 (t, *J* = 1.9 Hz, 1H), 4.14 (s, 3H), 3.13 (s, 3H).

20 Example 172

The compounds listed in the following table were prepared using a procedure similar to that described for example 171:

Structure	Example No.	MS (ESI) [M + H] ⁺	¹ H NMR
	172	450	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.90 (s, 1H), 8.56 (s, 2H), 7.69 (d, <i>J</i> = 26.9 Hz, 3H), 7.55 (d, <i>J</i> = 2.1 Hz, 1H), 6.89 (s, 1H), 4.22 (q, <i>J</i> = 6.9 Hz, 2H), 4.05 (s, 3H), 3.06 (s, 3H), 1.38 (t, <i>J</i> = 6.9 Hz, 3H).

Example 173: N-(3-chloro-5-(ethylsulfonamido)phenyl)-1-methyl-5-(pyrimidin-2-yl)-1H-pyrrole-3-carboxamide




- Step 1: A solution of 2-bromopyrimidine (500 mg, 3.145 mmol, 1 equiv), methyl 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrole-3-carboxylate (790 mg, 3.145 mmol, 1 equiv), Pd(dppf)Cl₂ (257 mg, 0.315 mmol, 0.1 equiv) and K₃PO₄ (2003 mg, 9.435 mmol, 3 equiv) in 1,4-dioxane (10 ml) and H₂O (2 ml) was stirred for 2 h at 80 °C under nitrogen atmosphere. The mixture was allowed to cool down to room temperature. The reaction mixture was diluted with water (50 mL). The resulting mixture was extracted with EtOAc (3*50 mL). The combined organic layers were washed with (100 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with Petroleum ether / EtOAc (1:1) to afford methyl 5-(pyrimidin-2-yl)-1H-pyrrole-3-carboxylate (480 mg, 75.11%) as a white solid. LCMS (ESI) [M+H]⁺: 204.07.
- Step 2: To a stirred solution of methyl 5-(pyrimidin-2-yl)-1H-pyrrole-3-carboxylate (460 mg, 2.264 mmol, 1 equiv) and NaH (163 mg, 6.792 mmol, 3 equiv) in DMF (5 ml) was added CH₃I (643 mg, 4.528 mmol, 2 equiv) dropwise 0 °C. The reaction was quenched by the addition of Water (2 mL) at 0 °C. The resulting mixture was extracted with EtOAc (3*50 mL). The combined organic layers were washed with water (100 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with Petroleum ether / EtOAc (3:1) to afford methyl

1-methyl-5-(pyrimidin-2-yl)pyrrole-3-carboxylate (390 mg, 79.31%) as a white solid. LCM S (ESI) $[M+H]^+$: 218.09.

Step 3: To a stirred solution of methyl 1-methyl-5-(pyrimidin-2-yl)pyrrole-3-carboxylate (80 mg, 0.368 mmol, 1 equiv) and *N*-(3-amino-5-chlorophenyl)ethanesulfonamide (87 mg, 0.368 mmol, 1 equiv) in PhMe (2 ml) were added $Al(Me)_3$ at 0 °C. The resulting mixture was stirred for additional 1.5 h at 80 °C. The mixture was allowed to cool down to room temperature. The reaction was quenched by the addition of Water (1.5 mL) at 0 °C. The resulting mixture was concentrated under reduced pressure. The residue was purified by reverse phase flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in Water (0.1% FA), 10% to 80% gradient in 10 min; detector, UV 254 nm. 45% to afford *N*-(3-chloro-5-ethanesulfonamidophenyl)-1-methyl-5-(pyrimidin-2-yl)pyrrole-3-carboxamide (21.7 mg, 13.90%) as a white solid. LCMS (ESI) $[M + H]^+$: 419.95. ^1H NMR (400 MHz, DMSO-*d*₆) δ 9.95-10.04 (dd, 2H), 8.81 (dd, *J* = 4.8, 1.0 Hz, 2H), 7.67-7.73 (m, *J* = 1.5 Hz, 4H), 7.30 (td, *J* = 4.9, 1.0 Hz, 1H), 6.90 (q, *J* = 1.5, 1.1 Hz, 1H), 4.09 (s, 3H), 3.17 (q, *J* = 7.3 Hz, 2H), 1.22 (t, *J* = 7.3 Hz, 3H).

Example 174: *N*-(3-chloro-5-(methanesulfonamido)phenyl)-5-(3-((5-fluoropyridin-3-yl)methoxy)pyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide



Step 1: A solution of 2-bromopyridin-3-ol (370 mg, 2.126 mmol, 1 equiv), 3-(chloromethyl)-5-fluoropyridine (310 mg, 2.126 mmol, 1 equiv) and K_2CO_3 (880 mg, 6.378 mmol, 3 equiv)

in DMF (5 ml) was stirred for 2 hours at 80 °C. The mixture was allowed to cool down to room temperature. The reaction mixture was diluted with water (50 mL). The resulting mixture was extracted with EtOAc (3*50 mL). The combined organic layers were washed with water (100 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with Petroleum ether / EtOAc (5:1) to afford 2-bromo-3-[(5-fluoropyridin-3-yl)methoxy]pyridine (380 mg, 63.12%) as a white solid. LCMS (ESI) [M+H]⁺: 283.

Step 2: A solution of 2-bromo-3-[(5-fluoropyridin-3-yl)methoxy]pyridine (360 mg, 1.272 mmol, 1 equiv), methyl 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrole-3-carboxylate (319 mg, 1.272 mmol, 1 equiv), Pd(dppf)Cl₂ (104 mg, 0.127 mmol, 0.1 equiv) and K₃PO₄ (810 mg, 3.816 mmol, 3 equiv) in 1,4-dioxane (5 ml) and H₂O (1 ml) was stirred for 2 hours at 80 °C under nitrogen atmosphere. The mixture was allowed to cool down to room temperature. The reaction mixture was diluted with water (50 mL). The resulting mixture was extracted with EtOAc (3 *50 mL). The combined organic layers were washed with water (100 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with Petroleum ether / EtOAc (1:1) to afford methyl 5-{3-[(5-fluoropyridin-3-yl)methoxy]pyridin-2-yl}-1H-pyrrole-3-carboxylate (280 mg, 67.27%) as a white solid. LCMS (ESI) [M+H]⁺: 329.

Step 3: To a stirred solution of methyl 5-{3-[(5-fluoropyridin-3-yl)methoxy]pyridin-2-yl}-1H-pyrrole-3-carboxylate (260 mg, 0.794 mmol, 1 equiv) and NaH (57 mg, 2.382 mmol, 3 equiv) in DMF (5 ml) was added CH₃I (226 mg, 1.588 mmol, 2 equiv) dropwise at 0 °C. The reaction was quenched by the addition of water (2 mL) at 0 °C. The reaction mixture was diluted with water (50 mL). The resulting mixture was extracted with EtOAc (3*50 mL). The combined organic layers were washed with water (100 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with Petroleum ether/EtOAc (1:1) to afford methyl 5-{3-[(5-fluoropyridin-3-yl)methoxy]pyridin-2-yl}-1-methylpyrrole-3-carboxylate (112 mg, 41.31%) as a white solid. LCMS (ESI) [M+H]⁺: 342.

Step 4: To a stirred solution of methyl 5-{3-[(5-fluoropyridin-3-yl)methoxy]pyridin-2-yl}-1-methylpyrrole-3-carboxylate (102 mg, 0.299 mmol, 1 equiv) and N-(3-amino-5-chlorophenyl)methanesulfonamide (66 mg, 0.299 mmol, 1 equiv) in PhMe (2 ml) was added Al(Me)₃ (0.8 ml) dropwise at room temperature. The resulting mixture was stirred for


additional 1.5 hours at 80 °C. The mixture was allowed to cool down to room temperature.

The reaction was quenched by the addition of Water (1.5 mL) at 0 °C. The resulting mixture was concentrated under reduced pressure. The crude product (50 mg) was purified by Prep-

- 5 Mobile Phase A: Water(0.1%FA), Mobile Phase B: ACN; Flow rate: 60 mL/min; Gradient: 33% B to 49% B in 8 min, 49% B; Wave Length: 254/220 nm; RT1(min): 7.27 to afford N-(3-chloro-5-methanesulfonamidophenyl)-5-{3-[(5-fluoropyridin-3-yl)methoxy]pyridin-2-yl}-1-methylpyrrole-3-carboxamide (23.9 mg, 15.09%) as a white solid. LCMS (ESI) [M + H]⁺: 486. ¹H NMR (300 MHz, DMSO-d₆) δ 9.80 (s, 1H), 8.55 (d, J = 2.5 Hz, 2H), 8.28 (dd, J = 4.7, 1.3 Hz, 1H), 7.80 (dt, J = 9.6, 2.3 Hz, 1H), 7.72 – 7.59 (m, 4H), 7.36 (dd, J = 8.4, 4.7 Hz, 1H), 7.18 (d, J = 1.9 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H), 5.35 (s, 2H), 3.80 (s, 3H), 3.06 (s, 3H).
- 10


Example 427

The compounds listed in the following table were prepared using a procedure similar to that described for example 174:

Structure	Example No.	MS (ESI) [M + H] ⁺	¹ H NMR
	427	502	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.24 (s, 1H), 10.08 (s, 1H), 9.03 (s, 1H), 8.47 (dd, J = 4.6, 1.5 Hz, 1H), 8.21 (s, 1H), 7.82 (dd, J = 8.2, 1.5 Hz, 1H), 7.69 (t, J = 1.9 Hz, 1H), 7.66 – 7.56 (m, 2H), 7.40 (td, J = 8.2, 6.8 Hz, 1H), 7.06 – 6.91 (m, 3H), 6.85 (dd, J = 8.0, 2.3 Hz, 1H), 3.07 (s, 3H).

15

Example 175: N-(3-chloro-5-(methylsulfonamido)phenyl)-1-(3-((5-fluoropyridin-3-yl)methoxy)pyridin-2-yl)-5-methyl-1H-pyrrole-3-carboxamide



Step 1: A solution of 2-bromopyridin-3-ol (300 mg, 1.724 mmol, 1 equiv), 3-(chloromethyl)-5-fluoropyridine (251 mg, 1.724 mmol, 1 equiv) and K_2CO_3 (714 mg, 5.172 mmol, 3 equiv) in DMF (5 ml) was stirred for 1.5 hours at 80 °C. The mixture was allowed to cool down to room temperature. The reaction mixture was diluted with water (50 mL). The resulting mixture was extracted with EtOAc (3*50 mL). The combined organic layers were washed with water (100 mL), dried over anhydrous Na_2SO_4 . After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with EtOAc/ Petroleum ether (25%) to afford 2-bromo-3-[(5-fluoropyridin-3-yl)methoxy]pyridine (220 mg, 45.07%) as a white solid. LCMS (ESI) $[M+H]^+$: 282.98.


Step 2: A solution of 2-bromo-3-[(5-fluoropyridin-3-yl)methoxy]pyridine (200 mg, 0.706 mmol, 1 equiv), methyl 5-methyl-1H-pyrrole-3-carboxylate (98 mg, 0.706 mmol, 1 equiv), (1S,2S)-N1,N2-dimethylcyclohexane-1,2-diamine (40 mg, 0.282 mmol, 0.4 equiv) and CuI (27 mg, 0.141 mmol, 0.2 equiv) in acetonitrile (5 ml) was stirred for 2 hours at 90 °C under nitrogen atmosphere. The mixture was allowed to cool down to room temperature. The resulting mixture was extracted with EtOAc (3*50 mL). The combined organic layers were washed with water (100 mL), dried over anhydrous Na_2SO_4 . After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with EtOAc /Petroleum ether (35%) to afford methyl 1-[3-[(5-fluoropyridin-3-yl)methoxy]pyridin-2-yl]-5-methylpyrrole-3-carboxylate (115 mg, 47.69%) as a white solid. LCMS (ESI) $[M+H]^+$: 342.12.



Step 3: A solution of methyl 1-{3-[(5-fluoropyridin-3-yl)methoxy]pyridin-2-yl}-5-methylpyrrole-3-carboxylate (95 mg, 0.278 mmol, 1 equiv) and LiOH (20 mg, 0.834 mmol, 3 equiv) in THF (4 ml) and H₂O (1 ml) was stirred for 2 hours at room temperature. The mixture was acidified to pH 7 with HCl (aq.). The precipitated solids were collected by filtration and washed with water (50 mL). The residue was purified by silica gel column chromatography, eluted with EtOAc /Petroleum ether (75%) to afford 1-{3-[(5-fluoropyridin-3-yl)methoxy]pyridin-2-yl}-5-methylpyrrole-3-carboxylic acid (73 mg, 80.14%) as a white solid. LCMS (ESI) [M+H]⁺: 328.10.

Step 4: A solution of 1-{3-[(5-fluoropyridin-3-yl)methoxy]pyridin-2-yl}-5-methylpyrrole-3-carboxylic acid (54 mg, 0.165 mmol, 1 equiv), N-(3-amino-5-chlorophenyl)methanesulfonamide (37 mg, 0.165 mmol, 1 equiv), TCFH (69 mg, 0.247 mmol, 1.5 equiv) and NMI (68 mg, 0.825 mmol, 5 equiv) in acetonitrile (2 ml) was stirred for 2 hours at room temperature. The residue solution was purified by Prep-HPLC with the following conditions: Sunfire prep C18 column, 30*150 mm, 5 μ m; Mobile Phase A: Water(0.1%FA), Mobile Phase B: ACN; Flow rate: 60 mL/min; Gradient: 31% B to 55% B in 8 min, 55% B; Wave Length: 254/220 nm; RT1(min): 7.65 to afford N-(3-chloro-5-methanesulfonamidophenyl)-1-{3-[(5-fluoropyridin-3-yl)methoxy]pyridin-2-yl}-5-methylpyrrole-3-carboxamide (23.3 mg, 26.47%) as a white solid. LCMS (ESI) [M + H]⁺: 530.25. ¹H NMR (300 MHz, DMSO-d₆) δ 9.78 (s, 1H), 8.54 (d, J = 2.8 Hz, 1H), 8.44 (d, J = 1.8 Hz, 1H), 8.23 (dd, J = 4.7, 1.3 Hz, 1H), 7.89 (dd, J = 8.3, 1.4 Hz, 1H), 7.78 – 7.69 (m, 2H), 7.69 – 7.53 (m, 3H), 6.89 (t, J = 2.0 Hz, 1H), 6.55 (d, J = 1.7 Hz, 1H), 5.34 (s, 2H), 3.05 (s, 3H), 2.12 – 2.06 (m, 3H).


Examples 428-430

The compounds listed in the following table were prepared using a procedure similar to that described for example 175:

Structure	Example No.	MS (ESI) [M + H] ⁺	¹ H NMR
	428	516	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.13 (s, 1H), 10.07 (s, 1H), 8.75 (d, J = 2.3 Hz, 1H), 8.27 (d, J = 2.2 Hz, 1H), 8.23 – 8.01 (m, 2H), 7.81 – 7.68 (m, 2H), 7.59 (dq, J = 8.1, 1.9 Hz, 4H), 7.41 (dt, J = 11.1, 2.4 Hz, 1H), 6.94 (t, J = 1.9 Hz,

			1H), 5.27 (s, 2H), 3.07 (d, J = 2.1 Hz, 3H).
	429	524	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.13 (s, 2H), 8.50 (s, 1H), 8.39 (s, 1H), 8.26 (s, 1H), 7.66 (t, J = 1.9 Hz, 1H), 7.56 (t, J = 1.9 Hz, 1H), 7.36 (dt, J = 10.9, 2.2 Hz, 1H), 7.31 (s, 1H), 7.20 (td, J = 9.5, 2.6 Hz, 1H), 6.93 (t, J = 2.0 Hz, 1H), 5.34 (s, 2H), 3.05 (s, 3H).
	430	481	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.18 (s, 1H), 10.09 (s, 1H), 8.84 (s, 1H), 8.30 (s, 1H), 7.68 (dd, J = 7.9, 1.6 Hz, 1H), 7.50 – 7.26 (m, 9H), 7.24 – 7.04 (m, 1H), 6.73 (dt, J = 10.4, 2.2 Hz, 1H), 5.29 (s, 2H), 3.08 (s, 3H).

Example 176: N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(2-oxopyrrolidin-1-yl)thiophene-2-carboxamide



- 5 Step 1: A mixture of methyl 4-bromothiophene-2-carboxylate (1050 mg, 4.74 mmol), pyrrolidin-2-one (403 mg, 4.74 mmol), K₂CO₃ (1.95 g, 14.2 mmol), CuI (180 mg, 0.948 mmol) and (1S,2S)-N1,N2-dimethylcyclohexane-1,2-diamine (674 mg, 4.74 mmol) in ACN


(20.00 mL) was stirred for 4 hours at 90 °C under N₂ atmosphere. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford methyl 4-(2-oxopyrrolidin-1-yl)thiophene-2-carboxylate (665 mg, 2.95 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 226






- 5 Step 2: To a stirred solution of methyl 4-(2-oxopyrrolidin-1-yl)thiophene-2-carboxylate (250 mg, 1.10 mmol) and LiOH (2.5 mg, 0.5 mmol) in THF (2.50 mL) was added H₂O (0.50 mL). Then the mixture was stirred for 2 h at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 4-(2-oxopyrrolidin-1-yl)thiophene-2-carboxylic acid (171 mg, 0.809 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 212.24





- Step 3: To a stirred solution of 4-(2-oxopyrrolidin-1-yl)thiophene-2-carboxylic acid (120 mg, 0.568 mmol), TCFH (239 mg, 0.851 mmol) and NMI (139 mg, 1.70 mmol) in ACN (5.00 mL) was added N-(3-amino-5-chlorophenyl)methanesulfonamide (125 mg, 0.568 mmol). Then the mixture was stirred for 4 hours at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford N-(3-chloro-5-methanesulfonamidophenyl)-4-(2-oxopyrrolidin-1-yl)thiophene-2-carboxamide (58.9 mg, 99.018 %) as a white solid. LCMS (ESI) [M + H]⁺: 414.89. ¹H NMR (400 MHz, DMSO-d₆) δ 10.56 (s, 1H), 10.07 (s, 1H), 8.49 (d, J = 1.5 Hz, 1H), 7.67 (dd, J = 12.3, 1.7 Hz, 3H), 6.96 (t, J = 2.0 Hz, 1H), 3.84 (t, J = 7.1 Hz, 2H), 3.07 (s, 3H), 2.51 (d, J = 16.2 Hz, 1H), 2.18 – 2.05 (m, 2H).





Examples 431-447




The compounds listed in the following table were prepared using a procedure similar to that described for example 177:


Structure	Compound No.	MS (ESI) [M+H] ⁺	¹ H NMR
	431	427 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.53 (s, 1H), 10.11 (s, 1H), 8.41 (d, J = 1.6 Hz, 1H), 7.64 (d, J = 2.0 Hz, 2H), 7.30 (d, J = 1.5 Hz, 1H), 6.95 (t, J = 2.0 Hz, 1H), 3.78 (dd, J = 9.2, 6.7 Hz, 2H), 3.48 (dd, J = 9.2, 6.6 Hz, 2H), 3.06 (s, 3H), 2.78 (s, 3H).

	432	469 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.58 (s, 1H), 10.07 (s, 1H), 8.36 (d, J = 1.5 Hz, 1H), 8.01 (d, J = 1.4 Hz, 1H), 7.66 (dt, J = 6.7, 1.9 Hz, 2H), 6.96 (t, J = 2.0 Hz, 1H), 3.07 (s, 3H), 2.90 (s, 3H), 1.43 (s, 6H).
	433	414 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.59 (s, 1H), 10.06 (s, 1H), 8.39 (d, J = 1.6 Hz, 1H), 7.65 (d, J = 2.0 Hz, 2H), 7.50 (d, J = 1.5 Hz, 1H), 6.97 (d, J = 2.1 Hz, 1H), 4.49 (dd, J = 9.1, 6.9 Hz, 2H), 4.05 (dd, J = 9.1, 6.9 Hz, 2H), 3.06 (s, 3H).
	434	426 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.35 (s, 1H), 10.08 (s, 1H), 7.92 (s, 1H), 7.62 (dt, J = 22.1, 1.9 Hz, 2H), 6.94 (t, J = 1.9 Hz, 1H), 3.72 (t, J = 7.0 Hz, 2H), 3.06 (s, 3H), 2.43 (t, J = 8.0 Hz, 2H), 2.32 (s, 3H), 2.18 – 2.06 (m, 2H).
	435	427	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.61 (s, 1H), 10.12 (s, 1H), 8.49 (d, J = 1.5 Hz, 1H), 8.19 (s, 1H), 7.92 (d, J = 1.5 Hz, 1H), 7.67 (dt, J = 8.4, 1.9 Hz, 2H), 7.56 (s, 1H), 6.97 (t, J = 1.9 Hz, 1H), 3.78 (s, 3H), 3.08 (s, 3H).
	436	441	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 10.15 (s, 1H), 8.18 (s, 1H), 7.92 (s, 1H), 7.65 (d, J = 1.9 Hz, 1H), 7.60 (d, J = 2.0 Hz, 1H), 7.56 (s, 1H), 6.94 (t,

			J = 2.0 Hz, 1H), 3.76 (s, 3H), 3.04 (s, 3H), 2.54 (s, 3H).
	437	449	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.98 (s, 1H), 9.80 (s, 1H), 8.25 (d, J = 3.0 Hz, 1H), 7.86 (d, J = 1.9 Hz, 1H), 7.70 (s, 1H), 7.60 (dd, J = 8.8, 3.4 Hz, 2H), 7.54 (d, J = 8.8 Hz, 1H), 6.88 (d, J = 2.7 Hz, 1H), 6.54 (s, 1H), 4.18 (q, J = 6.9 Hz, 2H), 3.04 (d, J = 4.4 Hz, 2H), 2.30 (s, 3H), 1.38 (t, J = 7.0 Hz, 3H).
	438	455	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 10.06 (s, 1H), 8.17 (s, 1H), 7.90 (s, 1H), 7.67 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.55 (s, 1H), 6.96 (t, J = 2.0 Hz, 1H), 3.99 (q, J = 7.0 Hz, 2H), 3.06 (s, 3H), 2.53 (s, 3H), 1.32 (t, J = 7.0 Hz, 3H).
	439	429	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.46 (s, 1H), 10.07 (s, 1H), 8.36 (d, J = 4.5 Hz, 1H), 8.18 (s, 1H), 7.85 (d, J = 4.0 Hz, 1H), 7.66 (d, J = 1.9 Hz, 1H), 7.61 (t, J = 1.9 Hz, 1H), 6.96 (t, J = 1.9 Hz, 1H), 3.06 (s, 3H), 2.52 (s, 3H).
	440	415	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.61 (s, 1H), 10.08 (s, 1H), 8.60 (d, J = 4.5 Hz, 1H), 8.49 (d, J = 1.5 Hz, 1H), 8.01 (d, J = 1.6 Hz, 1H), 7.85 (d, J = 4.1 Hz, 1H), 7.67 (dt, J = 12.4, 1.9 Hz, 2H), 6.97 (t,

			J = 1.9 Hz, 1H), 3.07 (s, 3H).
	441	436	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.97 (s, 2H), 8.64 (s, 2H), 8.35 (d, J = 2.1 Hz, 1H), 7.71 (d, J = 1.9 Hz, 1H), 7.68 – 7.62 (m, 1H), 6.90 (t, J = 2.0 Hz, 1H), 6.55 (dd, J = 2.1, 1.1 Hz, 1H), 3.97 (s, 3H), 3.05 (s, 3H), 2.55 (s, 3H).
	442	479	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.11 – 9.73 (m, 2H), 8.28 (d, J = 3.0 Hz, 1H), 7.87 (d, J = 2.1 Hz, 1H), 7.72 (t, J = 1.8 Hz, 1H), 7.67 – 7.60 (m, 2H), 7.55 (d, J = 8.8 Hz, 1H), 6.89 (t, J = 1.9 Hz, 1H), 6.55 (d, J = 2.0 Hz, 1H), 4.29 – 4.22 (m, 2H), 3.74 – 3.67 (m, 2H), 3.33 (s, 3H), 3.06 (s, 3H), 2.31 (s, 3H).
	443	473	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.03 (s, 1H), 9.93 (s, 1H), 9.02 – 8.95 (m, 1H), 8.44 (dd, J = 9.0, 2.5 Hz, 1H), 8.13 (d, J = 2.0 Hz, 1H), 7.89 (d, J = 8.6 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.63 (t, J = 1.9 Hz, 1H), 6.91 (t, J = 2.0 Hz, 1H), 6.65 (dd, J = 2.0, 1.1 Hz, 1H), 3.06 (s, 3H), 2.47 (d, J = 1.0 Hz, 3H).
	444	441	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.03 (s, 1H), 9.88 (s, 1H), 8.59 (d, J = 2.5 Hz, 1H), 8.45 – 8.24 (m, 1H), 7.77 (t, J = 1.9 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H),

			6.66 – 6.59 (m, 1H), 3.06 (s, 3H), 2.18 (s, 3H).
	445	455	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.08 (s, 1H), 9.87 (s, 1H), 8.59 (d, J = 2.5 Hz, 1H), 8.35 (ddd, J = 9.9, 8.3, 2.5 Hz, 1H), 7.77 (t, J = 1.9 Hz, 1H), 7.69 (t, J = 1.8 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H), 6.62 (dd, J = 1.9, 1.1 Hz, 1H), 3.16 (q, J = 7.3 Hz, 2H), 2.18 (s, 3H), 1.21 (t, J = 7.3 Hz, 3H).
	446	471	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.05 (s, 1H), 9.84 (s, 1H), 8.69 (d, J = 2.7 Hz, 1H), 8.48 (dd, J = 8.0, 2.7 Hz, 1H), 7.74 – 7.67 (m, 2H), 7.62 (t, J = 1.9 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H), 6.60 (dd, J = 1.9, 1.1 Hz, 1H), 3.17 (q, J = 7.3 Hz, 2H), 2.07 (d, J = 1.0 Hz, 3H), 1.21 (t, J = 7.3 Hz, 3H).
	447	449	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.95 (d, J = 52.6 Hz, 2H), 8.52 (d, J = 2.3 Hz, 1H), 8.03 – 7.92 (m, 2H), 7.78 – 7.54 (m, 3H), 6.89 (t, J = 1.9 Hz, 1H), 6.58 (dd, J = 2.0, 1.1 Hz, 1H), 4.52 (s, 2H), 3.35 (s, 3H), 3.05 (s, 3H), 2.38 (d, J = 1.0 Hz, 3H).

Example 177: N-(3-chloro-5-(methanesulfonamido)phenyl)-5-cyclopropyl-4-(pyridin-2-yl)thiophene-2-carboxamide

Step 1: A solution of methyl 5-bromo-4-(pyridin-2-yl)thiophene-2-carboxylate (300 mg, 1.00 mmol), $\text{Pd}_2(\text{dba})_3$ (183 mg, 0.200 mmol), X-phos (190 mg, 0.400 mmol), K_2CO_3 (414 mg, 3.00 mmol) and cyclopropylboronic acid (128 mg, 1.50 mmol) in dioxane (5 mL) and H_2O (1 mL) was stirred for 2 hours at 80 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with Petroleum ether / EtOAc (0-100%) to afford methyl 5-cyclopropyl-4-(pyridin-2-yl)thiophene-2-carboxylate (165 mg, 0.636 mmol) as a white solid. LCMS (ESI) $[\text{M} + \text{H}]^+$: 260.





Step 2: To a stirred solution of methyl 5-cyclopropyl-4-(pyridin-2-yl)thiophene-2-carboxylate (165 mg, 0.636 mmol) in EtOH (3 mL) and H_2O (3 mL) was added LiOH (152 mg, 6.35 mmol). Then the mixture was stirred for 2 hours at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 5-cyclopropyl-4-(pyridin-2-yl)thiophene-2-carboxylic acid (130 mg, 0.529 mmol) as a white solid. LCMS (ESI) $[\text{M} + \text{H}]^+$: 246.

Step 3: To a stirred solution of 5-cyclopropyl-4-(pyridin-2-yl)thiophene-2-carboxylic acid (130 mg, 0.529 mmol), TCFH (294 mg, 1.05 mmol) and NMI (260 mg, 3.17 mmol) in ACN (5 mL) was added N-(3-amino-5-chlorophenyl)methanesulfonamide (128 mg, 0.581 mmol). Then the mixture was stirred for 1 hour at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/ H_2O (5-95%, acidic system) to afford N-(3-chloro-5-methanesulfonamidophenyl)-5-cyclopropyl-4-(pyridin-2-

- yl)thiophene-2-carboxamide (56.2 mg, 0.125 mmol) as a white solid. LCMS (ESI) $[M + H]^+$: 448. ^1H NMR (400 MHz, DMSO- d_6) δ 10.46 (s, 1H), 10.10 (s, 1H), 8.69 (dt, $J = 4.6, 1.6$ Hz, 1H), 8.40 (s, 1H), 7.94 (td, $J = 7.7, 1.9$ Hz, 1H), 7.85 (dd, $J = 8.0, 1.2$ Hz, 1H), 7.65 (dt, $J = 17.3, 1.9$ Hz, 2H), 7.38 (ddd, $J = 7.6, 4.8, 1.2$ Hz, 1H), 6.94 (t, $J = 2.0$ Hz, 1H), 3.07 (s, 3H),
- 5 2.68 (tt, $J = 8.4, 5.2$ Hz, 1H), 1.25 – 1.14 (m, 2H), 0.85 – 0.76 (m, 2H).


Examples 448-451

The compounds listed in the following table were prepared using a procedure similar to that described for example 177:

Structure	Compound No.	MS (ESI) $[M+H]^+$	^1H NMR
	448	466.9	^1H NMR (400 MHz, DMSO- d_6) δ 10.56 (s, 1H), 10.05 (s, 1H), 9.00 (s, 2H), 8.62 (s, 1H), 7.66 (dt, $J = 10.7, 2.0$ Hz, 2H), 6.94 (d, $J = 2.1$ Hz, 1H), 3.42 (tt, $J = 8.5, 5.2$ Hz, 1H), 3.06 (s, 3H), 1.24 (h, $J = 4.5$ Hz, 2H), 0.83 (q, $J = 5.1$ Hz, 2H).
	449	480.9	^1H NMR (400 MHz, DMSO- d_6) δ 10.56 (s, 1H), 10.11 (s, 1H), 9.00 (s, 2H), 8.62 (s, 1H), 7.69 – 7.62 (m, 2H), 6.95 (d, $J = 1.9$ Hz, 1H), 3.41 (td, $J = 8.5, 4.3$ Hz, 1H), 3.17 (q, $J = 7.3$ Hz, 2H), 1.22 (q, $J = 7.4$ Hz, 5H), 0.86 – 0.77 (m, 2H).
	450	460.85	^1H NMR (400 MHz, DMSO- d_6) δ 10.73 (s, 1H), 10.08 (s, 1H), 9.07 (s, 2H), 8.69 (s, 1H), 7.71 – 7.61 (m, 2H), 6.97 (t, $J = 1.9$ Hz, 1H), 3.07 (s, 3H).
	451	450	^1H NMR (300 MHz, DMSO- d_6) δ 10.34 – 9.92 (m, 2H), 9.01 (s, 2H), 8.35 (d, $J = 2.1$ Hz, 1H), 7.67 (dt, $J = 17.9, 1.9$ Hz, 2H), 6.90 (t, $J = 1.9$ Hz, 1H), 6.50 (dd, $J = 2.1, 1.0$ Hz, 1H), 3.06 (s, 3H), 2.61 – 2.51 (m,

			1H), 0.93 – 0.81 (m, 2H), 0.67 – 0.56 (m, 2H).
--	--	--	--

Example 178: N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-(pyrimidin-2-yloxy)pyridin-2-yl)thiophene-2-carboxamide



- 5 Step 1: A solution of 2-bromopyridin-3-ol (520 mg, 2.98 mmol), Pd(dppf)Cl₂ (218 mg, 0.298 mmol), K₃PO₄ (1.89 g, 8.94 mmol) and methyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxylate (957 mg, 3.57 mmol) in dioxane (10 mL) and H₂O (1 mL) was stirred for 2 h at 80 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with EtOAc / Petroleum ether (0-100%) to afford methyl 4-(3-hydroxypyridin-2-yl)thiophene-2-carboxylate (510 mg, 2.16 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 236.


- 15 Step 2: A mixture of methyl 4-(3-hydroxypyridin-2-yl)thiophene-2-carboxylate (250 mg, 1.06 mmol), K₂CO₃ (439 mg, 3.18 mmol) and 2-chloropyrimidine (121 mg, 1.06 mmol) in DMF (5 mL) was stirred for 5 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford methyl 4-(3-(pyrimidin-2-yloxy)pyridin-2-yl)thiophene-2-carboxylate (210 mg, 0.670 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 314.





Step 3: To a stirred solution of methyl 4-[3-(pyrimidin-2-yloxy)pyridin-2-yl]thiophene-2-carboxylate (150 mg, 0.478 mmol) in EtOH (4 mL) and H₂O (4 mL) was added LiOH (114 mg, 4.77 mmol). Then the mixture was stirred for 2 h at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 4-[3-(pyrimidin-2-yloxy)pyridin-2-yl]thiophene-2-carboxylic acid (108 mg, 0.360 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 300.




Step 4: To a stirred solution of 4-[3-(pyrimidin-2-yloxy)pyridin-2-yl]thiophene-2-carboxylic acid (110 mg, 0.367 mmol), TCFH (205 mg, 0.734 mmol) and NMI (90.2 mg, 1.10 mmol) in ACN (5 mL) was added *N*-(3-amino-5-chlorophenyl)methanesulfonamide (80.9 mg, 0.367 mmol). Then the mixture was stirred for 1 h at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford *N*-(3-chloro-5-methanesulfonamidophenyl)-4-[3-(pyrimidin-2-yloxy)pyridin-2-yl]thiophene-2-carboxamide (56.3 mg, 0.112 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 502. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.64 (s, 1H), 10.08 (s, 1H), 8.72 (d, *J* = 1.4 Hz, 1H), 8.63 (dd, *J* = 18.2, 4.7 Hz, 3H), 8.23 (d, *J* = 1.3 Hz, 1H), 7.85 (dd, *J* = 8.2, 1.4 Hz, 1H), 7.67 (dt, *J* = 6.6, 1.9 Hz, 2H), 7.51 (dd, *J* = 8.2, 4.6 Hz, 1H), 7.30 (t, *J* = 4.8 Hz, 1H), 6.96 (t, *J* = 2.0 Hz, 1H), 3.07 (s, 3H).

Examples 452-459


The compounds listed in the following table were prepared using a procedure similar to that described for example 178:

Structure	Compound No.	MS (ESI) [M+H] ⁺	¹ H NMR
	452	492.9	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 10.56 (s, 1H), 10.06 (s, 1H), 8.63 (s, 1H), 8.55 (s, 2H), 7.69 (dt, <i>J</i> = 7.0, 1.9 Hz, 2H), 6.94 (t, <i>J</i> = 2.0 Hz, 1H), 4.94 (p, <i>J</i> = 7.1 Hz, 1H), 3.07 (s, 3H), 2.83 (s, 3H), 2.54 (s, 1H), 2.50 (s, 1H), 2.11 (dd, <i>J</i> = 11.1, 8.7 Hz, 2H), 1.84 (q, <i>J</i> = 10.3 Hz, 1H), 1.68 (p, <i>J</i> = 9.3 Hz, 1H).

	453	466.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.85 (s, 1H), 8.31 – 8.16 (m, 1H), 7.75 – 7.67 (m, 2H), 7.64 (t, J = 1.8 Hz, 1H), 7.57 (dd, J = 12.6, 2.5 Hz, 1H), 7.07 (dd, J = 3.2, 1.9 Hz, 1H), 6.89 (t, J = 1.9 Hz, 1H), 4.18 (q, J = 6.9 Hz, 2H), 3.84 (s, 3H), 3.06 (s, 3H), 1.37 (t, J = 6.9 Hz, 3H).
	454	452.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.85 (s, 1H), 8.27 (dd, J = 2.5, 1.1 Hz, 1H), 7.77 – 7.66 (m, 2H), 7.66 – 7.51 (m, 2H), 7.08 (dd, J = 3.3, 2.0 Hz, 1H), 6.90 (t, J = 2.0 Hz, 1H), 3.87 (d, J = 23.9 Hz, 6H), 3.06 (s, 3H).
	455	497.85	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.17 (dd, J = 2.5, 1.0 Hz, 1H), 7.96 (d, J = 1.5 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 7.55 (t, J = 1.9 Hz, 1H), 7.39 (dd, J = 11.9, 2.5 Hz, 1H), 7.02 (t, J = 1.9 Hz, 1H), 4.75 (hept, J = 6.1 Hz, 1H), 3.03 (s, 3H), 2.53 (d, J = 1.3 Hz, 3H), 1.39 (d, J = 6.0 Hz, 6H).
	456	484.2	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.20 (dd, J = 2.4, 1.0 Hz, 1H), 7.96 (d, J = 1.5 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 7.55 (t, J = 1.9 Hz, 1H), 7.38 (dd, J = 11.7, 2.4 Hz, 1H), 7.01 (t, J = 1.9 Hz, 1H), 4.19 (q, J = 7.0 Hz, 2H), 3.03 (s, 3H), 2.52 (d, J = 1.3 Hz, 3H), 1.46 (t, J = 7.0 Hz, 3H).

	457	466.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.65 (s, 1H), 10.09 (s, 1H), 8.76 (d, J = 1.4 Hz, 1H), 8.61 (s, 2H), 8.49 (d, J = 1.4 Hz, 1H), 7.70 (p, J = 1.8 Hz, 2H), 6.96 (t, J = 1.9 Hz, 1H), 4.88 (hept, J = 6.0 Hz, 1H), 3.08 (s, 3H), 1.34 (d, J = 6.0 Hz, 6H).
	458	483.85	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.64 (s, 1H), 10.09 (s, 1H), 8.68 (d, J = 1.4 Hz, 1H), 8.37 – 8.20 (m, 2H), 7.68 (dt, J = 5.4, 1.8 Hz, 2H), 7.61 (dd, J = 13.5, 2.4 Hz, 1H), 6.95 (t, J = 2.0 Hz, 1H), 4.80 (hept, J = 6.1 Hz, 1H), 3.07 (s, 3H), 1.33 (d, J = 6.0 Hz, 6H).
	459	466.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.57 (s, 1H), 10.23 – 9.78 (m, 1H), 8.66 (d, J = 1.4 Hz, 1H), 8.31 (dd, J = 16.8, 2.1 Hz, 2H), 7.84 (d, J = 8.8 Hz, 1H), 7.69 (dt, J = 6.8, 1.9 Hz, 2H), 7.51 (dd, J = 8.8, 3.0 Hz, 1H), 6.96 (t, J = 1.9 Hz, 1H), 4.76 (hept, J = 6.1 Hz, 1H), 3.08 (s, 3H), 1.31 (d, J = 6.0 Hz, 6H).

Example 460: N-(3-chloro-5-(methylsulfonamido)phenyl)-4-(3-fluoro-5-(3-(trifluoromethyl)azetidin-1-yl)pyridin-2-yl)-5-methylthiophene-2-carboxamide



- Step 1: A solution of 2-bromo-5-chloro-3-fluoropyridine (14 g, 66.5 mmol, 1 equiv),
- 5 Pd(dppf)Cl₂ (3.40 g, 4.65 mmol, 0.07 equiv), K₃PO₄ (42.1 g, 199 mmol, 3 equiv) and methyl 5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxylate (16.8 g, 59.8 mmol, 0.9 equiv) in dioxane (140 mL) and H₂O (14 mL) was stirred for 2 hours at 80 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA
- 10 (2:1) to afford methyl 4-(5-chloro-3-fluoropyridin-2-yl)-5-methylthiophene-2-carboxylate (11.9 g, 62.6% yield) as a white solid. LCMS (ESI) [M + H]⁺: 286

- Step 2: A solution of methyl 4-(5-chloro-3-fluoropyridin-2-yl)-5-methylthiophene-2-carboxylate (11.9 g, 41.6 mmol, 1 equiv), Cs₂CO₃ (40.4 g, 124 mmol, 3 equiv), Xantphos (4.81 g, 8.32 mmol, 0.2 equiv), Pd₂(dba)₃ (3.80 g, 4.16 mmol, 0.1 equiv) and 3-
- 15 (trifluoromethyl)azetidine hydrochloride (8.06 g, 49.9 mmol, 1.2 equiv) in PhMe (120 mL) was stirred for 5 hours at 100 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (1:1) to afford methyl 4-(3-fluoro-5-(3-




(trifluoromethyl) azetidin-1-yl)pyridin-2-yl)-5-methylthiophene-2-carboxylate (8.20 g, 52.9% yield) as a white solid. LCMS (ESI) $[M + H]^+$: 375




Step 3: To a stirred solution of 4-(3-fluoro-5-(3-(trifluoromethyl)azetidin-1-yl)pyridin-2-yl)-5-methylthiophene-2-carboxylate (8.2 g, 21.9 mmol, 1 equiv) in EtOH (82 mL) and H₂O (82 mL) was added LiOH (5.24 g, 219 mmol, 10 equiv) at 0 °C. Then the mixture was stirred for 4 hours at room temperature. The mixture was acidified to pH 5 with 1 M HCl (aq.). The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 4-{3-fluoro-5-[3-(trifluoromethyl)azetidin-1-yl]pyridin-2-yl}-5-methylthiophene-2-carboxylic acid (5.60 g, 70.9% yield) as a white solid. LCMS (ESI) $[M + H]^+$: 361

Step 4: To a stirred solution of 4-{3-fluoro-5-[3-(trifluoromethyl)azetidin-1-yl]pyridin-2-yl}-5-methylthiophene-2-carboxylic acid (5.6 g, 15.5 mmol, 1 equiv), TCFH (13.0 g, 46.5 mmol, 3 equiv) and NMI (5.08 g, 62.0 mmol, 4 equiv) in ACN (56 mL) was added *N*-(3-amino-5-chlorophenyl)methanesulfonamide (4.08 g, 18.5 mmol, 1.2 equiv) at room temperature. Then the mixture was stirred for 3 hours at room temperature. The mixture was concentrated under reduced pressure. The residue was purified by reversed-phase flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in water (0.1% formic acid), 0% to 50% gradient in 20 min; detector, UV 254 nm. This resulted in *N*-(3-chloro-5-methanesulfonamidophenyl)-4-{3-fluoro-5-[3-(trifluoromethyl)azetidin-1-yl]pyridin-2-yl}-5-methylthiophene-2-carboxamide (4.8783 g, 55.8 % yield) as a white solid. LCMS (ESI) $[M + H]^+$: 563.10. ¹H NMR (300 MHz, DMSO-*d*₆) δ 10.42 (s, 1H), 10.07 (s, 1H), 8.12 (s, 1H), 7.87 (s, 1H), 7.65 (d, *J* = 12.0 Hz, 2H), 7.03 (dd, *J* = 12.4, 2.3 Hz, 1H), 6.94 (d, *J* = 2.0 Hz, 1H), 4.22 (t, *J* = 8.6 Hz, 2H), 4.09 – 3.98 (m, 2H), 3.79 (s, 1H), 3.06 (s, 3H), 2.48 (s, 3H).




Examples 461-514



The compounds listed in the following table were prepared using a procedure similar to that described for example A:




Structure	Compound No.	MS (ESI) [M+H] ⁺	¹ H NMR
	461	510.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.75 (s, 1H), 7.90 (d, J = 2.7 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.67 – 7.59 (m, 2H), 6.98 (d, J = 2.7 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H), 6.76 (d, J = 1.9 Hz, 1H), 4.40 (t, J = 12.3 Hz, 4H), 3.61 (s, 3H), 3.07 (s, 3H), 2.32 (s, 3H).
	462	525.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.98 (s, 1H), 9.76 (s, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.67 – 7.51 (m, 3H), 7.00 – 6.80 (m, 2H), 6.74 (d, J = 2.3 Hz, 1H), 4.42 (t, J = 12.3 Hz, 4H), 3.86 (s, 3H), 3.69 (s, 3H), 3.06 (s, 3H).
	463	527.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.55 (s, 1H), 10.09 (s, 1H), 8.61 (s, 1H), 8.29 (s, 2H), 7.68 (d, J = 2.0 Hz, 2H), 6.95 (t, J = 1.9 Hz, 1H), 4.50 (t, J = 12.3 Hz, 4H), 3.18 (q, J = 7.3 Hz, 2H).




			2.82 (s, 3H), 1.22 (t, J = 7.3 Hz, 3H).
	464	565.15	¹ H NMR (400 MHz, DMSO- d ₆) δ 9.98 (s, 1H), 9.82 (s, 1H), 8.25 (t, J = 2.0 Hz, 1H), 7.78 – 7.54 (m, 3H), 7.40 (dd, J = 14.1, 2.5 Hz, 1H), 7.02 (dd, J = 3.3, 2.0 Hz, 1H), 6.89 (t, J = 2.0 Hz, 1H), 3.84 (s, 3H), 3.64 (s, 3H), 3.52 (t, J = 5.2 Hz, 4H), 3.34 (d, J = 6.1 Hz, 4H), 3.06 (s, 3H).
	465	490.15	¹ H NMR (300 MHz, DMSO- d ₆) δ 10.01 (s, 1H), 9.79 (s, 1H), 7.86 (d, J = 2.8 Hz, 1H), 7.73 (t, J = 1.9 Hz, 1H), 7.68 – 7.56 (m, 2H), 7.50 (d, J = 8.5 Hz, 1H), 7.04 – 6.84 (m, 3H), 4.37 (ddd, J = 10.4, 6.1, 4.2 Hz, 1H), 4.15 (dd, J = 8.2, 6.2 Hz, 2H), 3.91 (s, 3H), 3.72 (dd, J = 8.5, 4.1 Hz, 2H), 3.26 (s, 3H), 3.07 (s, 3H).
	466	496	¹ H NMR (400 MHz, DMSO- d ₆) δ 9.99 (s, 1H), 9.82 (s, 1H), 7.84 – 7.76 (m, 1H), 7.75 – 7.54 (m, 3H), 7.04 – 6.81 (m, 3H), 5.73 – 5.28



			(m, 1H), 4.38 – 4.17 (m, 2H), 4.05 (dddd, J = 24.1, 9.8, 3.1, 1.4 Hz, 2H), 3.81 (s, 3H), 3.06 (s, 3H).
	467	549.1	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 10.07 (s, 1H), 8.30 (s, 1H), 8.15 (s, 1H), 7.72 – 7.59 (m, 2H), 7.46 – 7.33 (m, 1H), 6.95 (t, J = 2.0 Hz, 1H), 3.42 (t, J = 5.5 Hz, 4H), 3.07 (s, 3H), 2.52 (s, 3H), 1.46 (t, J = 5.5 Hz, 4H), 0.37 (s, 4H).
	468	532.15	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.83 (s, 1H), 8.25 (t, J = 2.0 Hz, 1H), 7.73 (t, J = 1.8 Hz, 1H), 7.68 – 7.57 (m, 2H), 7.38 (dd, J = 14.5, 2.4 Hz, 1H), 7.00 (dd, J = 3.2, 1.9 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H), 3.84 (s, 3H), 3.40 (dd, J = 6.7, 4.4 Hz, 4H), 3.07 (s, 3H), 1.50 – 1.41 (m, 4H), 0.37 (s, 4H).




	469	541.95	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.84 (s, 1H), 8.31 (t, J = 2.0 Hz, 1H), 7.80 – 7.59 (m, 3H), 7.49 (dd, J = 14.2, 2.5 Hz, 1H), 7.03 (dd, J = 3.3, 1.9 Hz, 1H), 6.89 (t, J = 1.9 Hz, 1H), 3.85 (s, 3H), 3.52 (t, J = 5.8 Hz, 4H), 3.06 (s, 3H), 2.13 (d, J = 6.3 Hz, 1H), 2.08 (s, 1H), 2.06 – 1.98 (m, 2H).
	470	550	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.82 (s, 1H), 8.14 (t, J = 2.1 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.65 (d, J = 1.8 Hz, 2H), 7.24 (dd, J = 14.5, 2.4 Hz, 1H), 6.97 (dd, J = 3.2, 1.9 Hz, 1H), 6.90 (t, J = 2.0 Hz, 1H), 3.85 (s, 3H), 3.73 (t, J = 6.3 Hz, 4H), 3.07 (s, 3H), 0.77 (dd, J = 7.4, 4.6 Hz, 4H), 0.11 (s, 6H).
	471	559.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 10.07 (s, 1H), 8.36 (t, J = 2.0 Hz, 1H), 8.15 (d, J = 1.6 Hz, 1H), 7.66 (dt, J = 10.7, 1.9 Hz, 2H), 7.50 (dd, J = 13.9, 2.4 Hz, 1H), 6.95 (t,




			J = 2.0 Hz, 1H), 3.54 (t, J = 5.8 Hz, 4H), 3.07 (s, 3H), 2.51 (s, 3H), 2.10 (td, J = 13.5, 12.8, 7.3 Hz, 4H).
	472	574.2	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.83 (s, 1H), 8.25 (d, J = 2.1 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.70 – 7.56 (m, 2H), 7.42 (dd, J = 14.3, 2.4 Hz, 1H), 7.02 (dd, J = 3.3, 1.9 Hz, 1H), 6.89 (t, J = 1.9 Hz, 1H), 4.00 (d, J = 13.0 Hz, 2H), 3.84 (s, 3H), 3.06 (s, 3H), 2.89 (t, J = 12.5 Hz, 2H), 2.60 (d, J = 8.7 Hz, 1H), 1.89 (d, J = 12.9 Hz, 2H), 1.55 (tt, J = 13.1, 6.5 Hz, 2H).
	473	567.2	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.29 (s, 1H), 9.95 (s, 1H), 8.22 – 7.96 (m, 2H), 7.54 (dt, J = 14.9, 1.9 Hz, 2H), 7.14 (dd, J = 14.5, 2.5 Hz, 1H), 6.84 (t, J = 1.9 Hz, 1H), 3.63 (t, J = 6.3 Hz, 4H), 2.96 (s, 3H), 2.42 (s, 3H), 0.66 (dd, J = 7.5, 4.9 Hz, 4H), 0.00 (s, 6H).




	474	565.15 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.41 (s, 1H), 10.06 (s, 1H), 8.28 (t, J = 2.1 Hz, 1H), 8.13 (d, J = 1.6 Hz, 1H), 7.65 (dt, J = 14.5, 1.9 Hz, 2H), 7.39 (dd, J = 14.1, 2.5 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 4.36 (s, 4H), 3.32 – 3.28 (m, 4H), 3.06 (s, 3H), 2.51 (s, 3H), 1.96 – 1.83 (m, 4H).
	475	548.15	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.82 (s, 1H), 8.23 (d, J = 2.1 Hz, 1H), 7.86 – 7.53 (m, 3H), 7.38 (dd, J = 14.4, 2.4 Hz, 1H), 7.06 – 6.78 (m, 2H), 4.35 (s, 4H), 3.83 (s, 3H), 3.31 – 3.16 (m, 4H), 3.05 (s, 3H), 2.01 – 1.68 (m, 4H).
	476	549.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.40 (s, 1H), 10.05 (s, 1H), 8.10 (s, 1H), 7.85 (s, 1H), 7.65 (d, J = 14.4 Hz, 2H), 6.94 (s, 1H), 6.87 (d, J = 13.5 Hz, 1H), 3.49 (d, J = 10.2 Hz, 2H), 3.28 (s, 2H), 3.06 (s, 3H), 2.48 (s, 3H), 1.59 (s, 2H), 1.07 (s, 3H), 0.83 (s, 3H).




	477	563.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.40 (s, 1H), 10.05 (s, 1H), 8.12 (d, J = 1.6 Hz, 1H), 7.87 (t, J = 2.0 Hz, 1H), 7.65 (dt, J = 14.8, 1.9 Hz, 2H), 7.02 (dd, J = 12.4, 2.3 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 4.22 (t, J = 8.6 Hz, 2H), 4.03 (dd, J = 8.7, 5.1 Hz, 2H), 3.79 (d, J = 9.3 Hz, 1H), 3.06 (s, 3H), 2.48 (d, J = 1.3 Hz, 3H).
	478	585.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 10.06 (s, 1H), 8.41 – 8.27 (m, 1H), 8.14 (d, J = 1.6 Hz, 1H), 7.65 (dt, J = 14.9, 1.9 Hz, 2H), 7.43 (dd, J = 14.1, 2.5 Hz, 1H), 6.94 (t, J = 2.0 Hz, 1H), 3.47 (ddd, J = 10.8, 6.7, 3.9 Hz, 2H), 3.43 – 3.35 (m, 2H), 3.06 (s, 3H), 2.51 (s, 3H), 1.83 – 1.71 (m, 2H), 1.64 (s, 2H), 1.37 (t, J = 8.7 Hz, 2H).
	479	528.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.55 (s, 1H), 10.06 (s, 1H), 8.60 (s, 1H), 8.32 (s, 2H), 7.69 (p, J = 1.9 Hz, 2H), 6.94 (t, J = 1.9 Hz, 1H), 3.87 (t, J = 13.2 Hz,



			2H), 3.64 (t, J = 7.3 Hz, 2H), 3.07 (s, 3H), 2.82 (s, 3H), 2.58 (dp, J = 14.8, 7.4 Hz, 2H).
	480	571.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.41 (s, 1H), 10.06 (s, 1H), 8.13 (d, J = 1.6 Hz, 1H), 7.95 (t, J = 2.1 Hz, 1H), 7.66 (dt, J = 14.2, 1.9 Hz, 2H), 7.02 (dd, J = 13.4, 2.4 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 3.63 – 3.50 (m, 3H), 3.46 (dd, J = 10.3, 4.9 Hz, 1H), 3.06 (s, 3H), 2.49 (s, 3H), 2.25 (dt, J = 13.4, 6.8 Hz, 1H), 2.15 (dt, J = 12.7, 6.4 Hz, 1H), 1.69 (dd, J = 11.2, 7.3 Hz, 2H).
	481	591.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 10.07 (s, 1H), 8.31 (d, J = 2.4 Hz, 1H), 8.15 (d, J = 1.6 Hz, 1H), 7.66 (dt, J = 11.4, 1.9 Hz, 2H), 7.43 (dd, J = 14.1, 2.4 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 4.03 (d, J = 13.1 Hz, 2H), 3.07 (s, 3H), 2.91 (t, J = 12.1 Hz, 2H), 2.70 – 2.57 (m, 1H), 2.52 (s, 3H), 1.91 (d, J = 12.8 Hz, 2H), 1.65 – 1.44 (m, 2H).




	482	559.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 10.07 (s, 1H), 8.34 (s, 1H), 8.15 (s, 1H), 7.73 – 7.61 (m, 2H), 7.50 (dd, J = 13.9, 2.4 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 3.75 (t, J = 11.9 Hz, 2H), 3.46 (s, 2H), 3.07 (s, 3H), 2.52 (s, 3H), 2.10 (dt, J = 14.4, 7.4 Hz, 2H), 1.81 (s, 2H).
	483	542.2	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.84 (s, 1H), 8.29 (d, J = 2.0 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.69 – 7.60 (m, 2H), 7.49 (dd, J = 14.3, 2.4 Hz, 1H), 7.02 (dd, J = 3.2, 1.9 Hz, 1H), 6.89 (t, J = 1.9 Hz, 1H), 3.84 (s, 3H), 3.81 – 3.66 (m, 2H), 3.44 (t, J = 5.4 Hz, 2H), 3.06 (s, 3H), 2.08 (dp, J = 18.1, 6.3 Hz, 2H), 1.79 (s, 2H).
	484	545	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.41 (s, 1H), 10.08 (s, 1H), 8.13 (d, J = 1.6 Hz, 1H), 7.99 (t, J = 2.0 Hz, 1H), 7.64 (dt, J = 15.3, 1.9 Hz, 2H), 7.10 (dd, J = 13.3, 2.4 Hz, 1H), 6.94 (t,



			J = 1.9 Hz, 1H), 3.85 (t, J = 13.2 Hz, 2H), 3.61 (t, J = 7.3 Hz, 2H), 3.05 (s, 3H), 2.71 – 2.52 (m, 2H), 2.49 (s, 3H).
	485	527.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.41 (s, 1H), 10.07 (s, 1H), 8.32 (s, 1H), 8.10 (d, J = 3.0 Hz, 1H), 7.66 (dt, J = 12.3, 1.9 Hz, 2H), 7.53 (d, J = 8.7 Hz, 1H), 7.14 (dd, J = 8.7, 3.0 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 3.82 (t, J = 13.3 Hz, 2H), 3.59 (t, J = 7.2 Hz, 2H), 3.07 (s, 3H), 2.66 (s, 3H), 2.58 (dq, J = 14.5, 7.2 Hz, 2H).
	486	500.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.64 (s, 1H), 10.08 (s, 1H), 8.74 (d, J = 1.3 Hz, 1H), 8.43 (d, J = 1.2 Hz, 1H), 8.26 (s, 2H), 7.70 (p, J = 1.9 Hz, 2H), 6.96 (t, J = 1.9 Hz, 1H), 4.50 (t, J = 12.3 Hz, 4H), 3.08 (s, 3H).
	487	498.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.58 (s, 1H), 10.09 (s, 1H), 8.64 (d, J = 1.4 Hz, 1H), 8.28 (d, J = 1.3 Hz, 1H), 7.99




			(d, J = 2.8 Hz, 1H), 7.78 (d, J = 8.5 Hz, 1H), 7.73 – 7.62 (m, 2H), 7.11 (dd, J = 8.6, 2.9 Hz, 1H), 6.96 (t, J = 2.0 Hz, 1H), 4.41 (t, J = 12.3 Hz, 4H), 3.08 (s, 3H).
	488	516.85	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.64 (s, 1H), 10.13 (s, 1H), 8.65 (d, J = 1.4 Hz, 1H), 8.20 (t, J = 1.5 Hz, 1H), 7.90 (t, J = 2.0 Hz, 1H), 7.68 (p, J = 1.9 Hz, 2H), 7.12 (dd, J = 13.3, 2.4 Hz, 1H), 6.95 (t, J = 2.0 Hz, 1H), 4.46 (t, J = 12.3 Hz, 4H), 3.07 (s, 3H).
	489	546.15	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.55 (s, 1H), 10.05 (s, 1H), 8.60 (s, 1H), 8.23 (s, 1H), 7.68 (dt, J = 4.8, 1.9 Hz, 2H), 6.94 (t, J = 2.0 Hz, 1H), 4.26 (t, J = 8.6 Hz, 2H), 4.07 (dd, J = 8.6, 5.1 Hz, 2H), 3.80 (dt, J = 9.4, 4.8 Hz, 1H), 3.07 (s, 3H), 2.81 (s, 3H).
	490	533.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.54 (s, 1H), 10.05 (s, 1H), 8.59 (s, 1H), 8.16 (s, 2H), 7.68 (dt, J = 5.4, 1.9 Hz,



			2H), 6.94 (t, J = 1.9 Hz, 1H), 4.11 (s, 4H), 3.07 (s, 3H), 2.89 (t, J = 12.5 Hz, 4H), 2.80 (s, 3H).
	491	544.15	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.55 (s, 1H), 10.05 (s, 1H), 8.60 (s, 1H), 8.27 (s, 2H), 7.69 (dt, J = 5.2, 1.9 Hz, 2H), 6.94 (t, J = 1.9 Hz, 1H), 3.68 – 3.53 (m, 3H), 3.47 (dd, J = 10.2, 4.8 Hz, 1H), 3.07 (s, 3H), 2.81 (s, 3H), 2.20 (ddt, J = 44.1, 12.2, 6.4 Hz, 2H), 1.69 (dd, J = 11.2, 7.4 Hz, 2H).
	492	537.15	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.99 (s, 1H), 9.88 (s, 1H), 8.19 – 7.94 (m, 2H), 7.84 – 7.53 (m, 3H), 7.45 (t, J = 1.7 Hz, 1H), 6.98 – 6.62 (m, 1H), 4.25 – 3.82 (m, 7H), 3.06 (d, J = 1.0 Hz, 3H), 2.88 (t, J = 12.5 Hz, 4H).
	493	529.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.90 (d, J = 2.2 Hz, 2H), 8.17 (d, J = 1.3 Hz, 2H), 7.75 (h, J = 1.3 Hz, 1H), 7.68 (q, J = 1.7 Hz, 1H), 7.62 (d, J = 2.1 Hz, 1H), 7.48 (d, J = 2.2 Hz,




			1H), 6.90 (q, J = 2.3 Hz, 1H), 4.23 (t, J = 8.6 Hz, 2H), 4.04 (d, J = 9.5 Hz, 5H), 3.80 (dt, J = 14.1, 9.3, 5.4 Hz, 1H), 3.07 (d, J = 2.3 Hz, 3H).
	494	537.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.87 (s, 1H), 8.22 (s, 2H), 7.70 (dt, J = 27.3, 1.9 Hz, 2H), 7.60 (d, J = 2.1 Hz, 1H), 7.43 (d, J = 2.1 Hz, 1H), 6.88 (t, J = 2.0 Hz, 1H), 4.03 (s, 3H), 3.61 – 3.49 (m, 3H), 3.47 – 3.40 (m, 1H), 3.06 (s, 3H), 2.25 (dt, J = 13.3, 6.9 Hz, 1H), 2.12 (ddd, J = 16.4, 11.6, 5.1 Hz, 1H), 1.77 – 1.56 (m, 2H).
	495	536.2	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.77 (s, 1H), 8.00 (d, J = 2.9 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.66 – 7.56 (m, 2H), 7.51 (d, J = 8.7 Hz, 1H), 7.06 (dd, J = 8.8, 2.9 Hz, 1H), 6.97 (d, J = 2.0 Hz, 1H), 6.88 (t, J = 1.9 Hz, 1H), 3.91 (s, 3H), 3.61 – 3.38 (m, 4H), 3.06 (s, 3H), 2.24 (dt, J = 13.3, 6.7 Hz, 1H), 2.16 – 2.08




			(m, 1H), 1.75 – 1.56 (m, 2H).
	496	545.15	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.41 (s, 1H), 10.07 (s, 1H), 8.31 (s, 1H), 7.98 (d, J = 2.9 Hz, 1H), 7.66 (dt, J = 13.1, 1.9 Hz, 2H), 7.52 (d, J = 8.5 Hz, 1H), 7.04 (dd, J = 8.6, 2.9 Hz, 1H), 6.94 (t, J = 2.0 Hz, 1H), 4.18 (t, J = 8.5 Hz, 2H), 4.00 (dd, J = 8.5, 5.1 Hz, 2H), 3.86 – 3.68 (m, 1H), 3.06 (s, 3H), 2.65 (s, 3H).
	497	528.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.78 (s, 1H), 7.92 (d, J = 2.9 Hz, 1H), 7.78 – 7.38 (m, 4H), 7.07 – 6.95 (m, 2H), 6.89 (t, J = 1.9 Hz, 1H), 4.16 (t, J = 8.4 Hz, 2H), 3.97 (dd, J = 8.4, 5.2 Hz, 2H), 3.91 (s, 3H), 3.76 (ddq, J = 14.1, 9.4, 5.0 Hz, 1H), 3.06 (s, 3H).
	498	536.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.77 (s, 1H), 7.86 (d, J = 2.8 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.65 – 7.56 (m, 2H), 7.49 (d, J = 8.6 Hz, 1H), 7.01 – 6.84 (m, 3H),




			4.01 (s, 4H), 3.90 (s, 3H), 3.06 (s, 3H), 2.88 (t, J = 12.5 Hz, 4H).
	499	553.05	¹ H NMR (400 MHz, DMSO- d6) δ 10.40 (s, 1H), 10.09 (s, 1H), 8.29 (s, 1H), 7.98 – 7.84 (m, 1H), 7.65 (dt, J = 12.9, 1.9 Hz, 2H), 7.53 – 7.44 (m, 1H), 7.04 – 6.87 (m, 2H), 4.03 (s, 4H), 3.06 (s, 3H), 2.89 (t, J = 12.5 Hz, 4H), 2.64 (s, 3H).
	500	533.05	¹ H NMR (400 MHz, DMSO- d6) δ 10.41 (s, 1H), 10.07 (s, 1H), 8.31 (s, 1H), 8.06 (d, J = 3.0 Hz, 1H), 7.66 (dt, J = 12.4, 1.9 Hz, 2H), 7.51 (d, J = 8.7 Hz, 1H), 7.09 (dd, J = 8.7, 3.0 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 3.61 – 3.48 (m, 3H), 3.43 (dd, J = 10.1, 4.9 Hz, 1H), 3.07 (s, 3H), 2.66 (s, 3H), 2.25 (dt, J = 12.8, 6.7 Hz, 1H), 2.13 (dq, J = 12.7, 6.4 Hz, 1H), 1.68 (dq, J = 13.1, 8.2, 6.5 Hz, 2H).

	501	547.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 10.07 (s, 1H), 8.12 (d, J = 1.6 Hz, 1H), 7.86 (d, J = 2.0 Hz, 1H), 7.45 (dd, J = 10.2, 2.1 Hz, 2H), 7.02 (dd, J = 12.5, 2.3 Hz, 1H), 6.72 (dt, J = 10.4, 2.2 Hz, 1H), 4.22 (t, J = 8.6 Hz, 2H), 4.03 (dd, J = 8.7, 5.1 Hz, 2H), 3.79 (ddq, J = 14.2, 9.5, 5.1 Hz, 1H), 3.06 (s, 3H), 2.49 – 2.47 (S, 3H).
	502	557.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.40 (s, 1H), 10.09 (s, 1H), 8.12 (s, 1H), 7.86 (d, J = 2.3 Hz, 1H), 7.72 – 7.59 (m, 2H), 7.02 (dd, J = 12.4, 2.3 Hz, 1H), 6.95 (d, J = 2.2 Hz, 1H), 4.22 (t, J = 8.6 Hz, 2H), 4.03 (dd, J = 8.7, 5.1 Hz, 2H), 3.79 (ddq, J = 14.2, 9.5, 5.1 Hz, 1H), 3.17 (q, J = 7.3 Hz, 2H), 2.48 (s, 3H), 1.21 (t, J = 7.3 Hz, 3H).
	503	573.3	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.36 (s, 1H), 10.08 (s, 1H), 8.35 (d, J = 2.4 Hz, 1H), 8.21 (s, 1H), 7.66 (tt, J = 11.6, 2.4 Hz, 3H), 6.94 (d, J = 2.2 Hz, 1H),

			3.38 – 3.31 (m, 4H), 3.10 (d, J = 5.3 Hz, 4H), 3.06 (s, 3H), 2.44 (s, 3H).
	504	566.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.36 (s, 1H), 10.07 (s, 1H), 8.31 (d, J = 2.4 Hz, 1H), 8.20 (s, 1H), 7.67 (t, J = 1.8 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.48 (dd, J = 10.9, 2.5 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 3.45 (d, J = 10.4 Hz, 1H), 3.38 (d, J = 9.9 Hz, 1H), 3.05 (s, 3H), 2.81 (p, J = 4.6 Hz, 4H), 2.44 (s, 3H), 1.96 (s, 3H).
	505	539.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.36 (s, 1H), 10.10 (s, 1H), 8.07 – 7.98 (m, 2H), 7.67 (t, J = 1.9 Hz, 1H), 7.61 (d, J = 2.0 Hz, 1H), 7.16 (dd, J = 12.1, 2.5 Hz, 1H), 6.93 (d, J = 2.0 Hz, 1H), 3.95 – 3.90 (m, 1H), 3.16 (s, 3H), 3.05 (s, 4H), 3.11 – 3.00 (m, 1H), 3.00 – 2.91 (m, 1H), 2.84 (dd, J = 10.9, 2.0 Hz, 1H), 2.23 (s, 3H), 1.92 (td, J = 8.5, 8.0, 4.3 Hz, 2H).


	506	602	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.32 (s, 1H), 9.99 (s, 1H), 8.26 (d, J = 2.4 Hz, 1H), 8.13 (s, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.54 (t, J = 1.9 Hz, 1H), 7.47 (dd, J = 10.9, 2.6 Hz, 1H), 6.87 (t, J = 2.0 Hz, 1H), 3.07 – 3.00 (m, 4H), 2.99 (s, 3H), 2.87 (t, J = 4.9 Hz, 4H), 2.80 (s, 3H), 2.37 (s, 3H).
	507	549.15	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.37 (s, 1H), 10.06 (s, 1H), 8.25 (d, J = 2.4 Hz, 1H), 8.19 (s, 1H), 7.65 (dt, J = 16.8, 1.7 Hz, 2H), 7.47 (dd, J = 11.1, 2.5 Hz, 1H), 6.94 (q, J = 1.7 Hz, 1H), 3.06 (s, 3H), 2.86 (t, J = 5.2 Hz, 4H), 2.45 (s, 3H), 1.30 (t, J = 5.0 Hz, 4H), 0.26 (s, 4H).
	508	579.1	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.99 (s, 2H), 8.18 (s, 2H), 7.74 (t, J = 1.9 Hz, 1H), 7.72 – 7.66 (m, 2H), 7.57 (d, J = 2.1 Hz, 1H), 6.91 (t, J = 1.9 Hz, 1H), 6.59 – 6.13 (m, 1H), 5.08 (td, J = 14.5, 3.9 Hz, 2H), 4.24 (t, J = 8.6 Hz, 2H),

			4.06 (dd, J = 8.6, 5.1 Hz, 2H), 3.80 (ddq, J = 13.9, 9.2, 5.0 Hz, 1H), 3.07 (s, 3H).
	509	510	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.54 (s, 1H), 10.08 (s, 1H), 8.59 (s, 1H), 8.20 (s, 1H), 7.68 (d, J = 1.9 Hz, 2H), 6.95 (t, J = 1.9 Hz, 1H), 5.54 (ddt, J = 60.6, 6.1, 3.1 Hz, 1H), 4.48 – 4.25 (m, 2H), 4.21 – 4.02 (m, 2H), 3.17 (q, J = 7.3 Hz, 2H), 2.81 (s, 3H), 1.22 (t, J = 7.3 Hz, 3H).
	510	514	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.56 (s, 1H), 10.06 (s, 1H), 8.61 (s, 1H), 8.29 (s, 1H), 7.69 (dt, J = 5.0, 1.9 Hz, 2H), 6.94 (t, J = 1.9 Hz, 1H), 4.50 (t, J = 12.3 Hz, 4H), 3.07 (s, 3H), 2.82 (s, 3H).
	511	571	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.41 (s, 1H), 10.07 (s, 1H), 8.12 (d, J = 1.6 Hz, 1H), 7.80 (t, J = 2.0 Hz, 1H), 7.66 (dt, J = 11.2, 1.9 Hz, 2H), 7.00 – 6.88 (m, 2H), 4.08 (s, 4H), 3.07 (s, 3H), 2.90 (t, J = 12.5 Hz, 4H), 2.48

			(d, J = 1.5 Hz, 3H).
	512	551	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 10.04 (s, 1H), 8.12 (d, J = 1.6 Hz, 1H), 7.92 (t, J = 2.1 Hz, 1H), 7.66 (dt, J = 10.5, 1.9 Hz, 2H), 7.04 – 6.82 (m, 2H), 4.75 – 4.43 (m, 4H), 3.63 (s, 2H), 3.38 (d, J = 7.0 Hz, 2H), 3.07 (s, 3H), 2.49 (d, J = 1.4 Hz, 3H), 2.30 (t, J = 6.9 Hz, 2H).
	513	554	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.99 (s, 1H), 9.79 (s, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.62 (q, J = 2.0 Hz, 3H), 6.89 (dd, J = 5.2, 2.0 Hz, 2H), 6.79 (s, 1H), 4.73 (p, J = 5.8 Hz, 1H), 4.43 (dd, J = 13.4, 10.6 Hz, 4H), 3.68 (s, 3H), 3.06 (s, 3H), 1.28 (d, J = 5.9 Hz, 6H).
	514	583	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.36 (s, 1H), 10.06 (s, 1H), 8.06 (s, 1H), 7.67 (t, J = 1.9 Hz, 1H), 7.63 (dd, J = 4.8, 2.2 Hz, 2H), 6.94 (t, J = 1.9 Hz, 1H), 6.51 (d, J = 2.3 Hz, 1H), 4.78 (p, J = 7.1 Hz, 1H), 4.42 (t, J = 12.3

			Hz, 4H), 3.07 (s, 3H), 2.45 (d, J = 7.7 Hz, 2H), 2.40 (s, 3H), 2.12 – 1.94 (m, 2H), 1.79 (q, J = 10.2 Hz, 1H), 1.65 (p, J = 9.1 Hz, 1H).
--	--	--	--

Example 515: *N*-(3-chloro-5-methanesulfonamidophenyl)-5-[5-(4-methoxypiperidin-1-yl)pyridin-2-yl]-1-methyl-1*H*-pyrrole-3-carboxamide







- 5 Step 1: To a stirred solution of 5-bromo-2-iodopyridine (320 mg, 1.12 mmol), Pd(dppf)Cl₂ (122 mg, 168 μmol) and K₂CO₃ (467 mg, 3.36 mmol) in dioxane (9 mL) and H₂O (3 mL) was added methyl 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrrole-3-carboxylate (296 mg, 1.12 mmol). Then the mixture was stirred for 1 hour at 90 °C. The mixture was concentrated and purified by normal phase flash chromatography eluting
- 10 with PE/EA (70%) to afford methyl 5-(5-bromopyridin-2-yl)-1-methyl-1*H*-pyrrole-3-carboxylate (255 mg, 864 μmol) as an off yellow solid. LCMS (ESI) [M + H]⁺: 295
- Step 2: To a stirred solution of methyl 5-(5-bromopyridin-2-yl)-1-methyl-1*H*-pyrrole-3-carboxylate (255 mg, 864 μmol) and Cs₂CO₃ (841 mg, 2.59 mmol) in dioxane (10 mL) was added 4-ethylpiperidine (97.8 mg, 864 μmol) and {1,3-bis[2,6-bis(heptan-4-yl)phenyl]-4,5-dichloro-2,3-dihydro-1*H*-imidazol-2-yl}dichloro(3-chloro-λ⁴-pyridin-1-yl)palladium (33.5
- 15 mg, 8.64 μmol). Then the mixture was stirred for 1 hour at 100°C. The mixture was concentrated and purified by normal phase flash chromatography eluting with PE/EA (81%) to afford methyl 5-[5-(4-methoxypiperidin-1-yl)pyridin-2-yl]-1-methyl-1*H*-pyrrole-3-carboxylate (150 mg, 455 μmol) as an off white solid. LCMS (ESI) [M + H]⁺: 330.17




- Step 3: To a stirred solution of methyl 5-[5-(4-methoxypiperidin-1-yl)pyridin-2-yl]-1-methyl-1*H*-pyrrole-3-carboxylate (130 mg, 394 μ mol) and *N*-(3-amino-5-chlorophenyl)methanesulfonamide (86.9 mg, 394 μ mol) in PhMe (6 mL) was added $\text{Al}(\text{CH}_3)_3$ (36 mg, 1.195 mmol). Then the mixture was stirred for 1 hour at 80 °C. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/ H_2O (42%, acidic system) to afford *N*-(3-chloro-5-methanesulfonamidophenyl)-5-[5-(4-methoxypiperidin-1-yl)pyridin-2-yl]-1-methyl-1*H*-pyrrole-3-carboxamide (122 mg, 229 μ mol) as a white solid. LCMS (ESI) $[\text{M} + \text{H}]^+$: 517.95. ^1H NMR (400 MHz, DMSO- d_6) δ 10.02 (s, 1H), 9.78 (s, 1H), 8.31 (d, $J = 2.9$ Hz, 1H), 7.72 (t, $J = 1.9$ Hz, 1H), 7.66 – 7.57 (m, 2H), 7.51 (d, $J = 8.8$ Hz, 1H), 7.41 (dd, $J = 8.9, 3.0$ Hz, 1H), 7.01 (d, $J = 2.0$ Hz, 1H), 6.89 (t, $J = 2.0$ Hz, 1H), 3.92 (s, 3H), 3.58 (dt, $J = 13.1, 4.8$ Hz, 2H), 3.38 (tt, $J = 8.1, 3.7$ Hz, 1H), 3.28 (s, 3H), 3.06 (s, 3H), 3.00 (ddd, $J = 12.7, 9.5, 3.2$ Hz, 2H), 1.99 – 1.90 (m, 2H), 1.53 (dtd, $J = 12.8, 9.0, 3.8$ Hz, 2H).


Example 516-523

- The compounds listed in the following table were prepared using a procedure similar to that described for example 515:


Structure	Compound No.	MS (ESI) $[\text{M} + \text{H}]^+$	^1H NMR
	516	532.2	^1H NMR (300 MHz, DMSO- d_6) δ 10.01 (s, 1H), 9.81 (s, 1H), 7.82 (s, 1H), 7.71 (t, $J = 1.9$ Hz, 1H), 7.63 (d, $J = 1.8$ Hz, 2H), 6.95 – 6.72 (m, 3H), 3.80 (s, 3H), 3.48 (d, $J = 10.3$ Hz, 2H), 3.28 (d, $J = 10.4$ Hz, 2H), 3.06 (s, 3H), 1.58 (d, $J = 3.6$ Hz, 2H), 1.07 (s, 3H), 0.83 (s, 3H).

	517	528.15	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.05 (s, 1H), 9.83 (s, 1H), 7.89 (dd, J = 2.4, 1.4 Hz, 1H), 7.75 – 7.53 (m, 3H), 7.12 (dd, J = 12.6, 2.4 Hz, 1H), 7.02 (dd, J = 3.3, 2.0 Hz, 1H), 6.89 (t, J = 1.9 Hz, 1H), 4.44 (t, J = 12.3 Hz, 4H), 3.82 (s, 3H), 3.16 (q, J = 7.2 Hz, 2H), 1.21 (t, J = 7.3 Hz, 3H).
	518	568.15	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.84 (s, 1H), 8.27 (t, J = 2.0 Hz, 1H), 7.78 – 7.68 (m, 1H), 7.70 – 7.56 (m, 2H), 7.42 (dt, J = 14.0, 1.9 Hz, 1H), 7.12 – 6.96 (m, 1H), 6.90 (t, J = 1.9 Hz, 1H), 3.85 (s, 3H), 3.44 (tdd, J = 12.7, 7.3, 4.2 Hz, 4H), 3.07 (s, 3H), 1.76 (ddd, J = 11.9, 7.4, 3.9 Hz, 2H), 1.70 – 1.50 (m, 2H), 1.37 (t, J = 8.7 Hz, 2H).
	519	538.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.05 (s, 1H), 9.83 (d, J = 5.8 Hz, 1H), 7.96 – 7.88 (m, 1H), 7.73 (p, J = 1.7 Hz, 1H), 7.63 (dt, J = 4.5, 2.3 Hz, 2H), 7.54 (dd, J = 8.6, 0.7 Hz, 1H), 7.11 – 6.95 (m,

			2H), 6.93 – 6.83 (m, 1H), 4.47 (td, J = 8.1, 4.1 Hz, 1H), 4.31 – 4.07 (m, 4H), 3.91 (s, 3H), 3.07 (d, J = 4.3 Hz, 6H).
	520	475.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.05 (s, 1H), 9.86 (s, 1H), 8.16 (s, 2H), 7.73 (t, J = 1.9 Hz, 1H), 7.66 (t, J = 1.9 Hz, 1H), 7.58 (d, J = 2.1 Hz, 1H), 7.39 (d, J = 2.2 Hz, 1H), 6.88 (t, J = 1.9 Hz, 1H), 4.02 (s, 3H), 3.25 (s, 4H), 3.05 (s, 3H), 2.03 – 1.93 (m, 4H).
	521	460.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.99 (s, 1H), 9.88 (s, 1H), 8.05 (s, 2H), 7.74 (d, J = 2.2 Hz, 1H), 7.68 (d, J = 2.2 Hz, 1H), 7.60 (d, J = 2.2 Hz, 1H), 7.44 (d, J = 2.2 Hz, 1H), 6.89 (t, J = 2.1 Hz, 1H), 4.02 (s, 3H), 3.97 (t, J = 7.3 Hz, 4H), 3.07 (s, 3H), 2.40 (p, J = 7.3 Hz, 2H).
	522	500	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.92 (s, 1H), 9.78 (s, 1H), 7.82 (d, J = 3.0 Hz, 1H), 7.77 (d, J = 2.0 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H),

			7.63 (t, J = 1.9 Hz, 1H), 7.36 (d, J = 8.8 Hz, 1H), 7.06 (dd, J = 8.8, 3.1 Hz, 1H), 6.88 (t, J = 1.9 Hz, 1H), 6.51 (dd, J = 2.0, 1.1 Hz, 1H), 3.48 (t, J = 6.8 Hz, 2H), 3.25 (s, 2H), 3.05 (s, 3H), 2.25 (d, J = 0.9 Hz, 3H), 1.95 (t, J = 6.7 Hz, 2H), 0.66 (dt, J = 6.9, 1.8 Hz, 4H).
	523	500.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.79 (s, 1H), 8.23 (d, J = 3.0 Hz, 1H), 7.83 (d, J = 2.0 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.63 (t, J = 1.9 Hz, 1H), 7.53 (dd, J = 8.8, 3.0 Hz, 1H), 7.38 (d, J = 8.7 Hz, 1H), 6.88 (t, J = 1.9 Hz, 1H), 6.51 (s, 1H), 4.41 (t, J = 2.6 Hz, 2H), 3.06 (s, 3H), 2.29 (s, 3H), 1.71 – 1.65 (m, 4H), 1.47 (t, J = 6.4 Hz, 4H).

Example 524: N-(3-chloro-5-methanesulfo namidophenyl)-5-{3-fluoro-5-[3-(trifluoromethyl)azetidin-1-yl]pyridin-2-yl}-1-methyl-1H-pyrrole-3-carboxamide



Step 1: To a mixture of 2-bromo-5-chloro-3-fluoropyridine (520 mg, 2.47 mmol), methyl 1-
 5 methyl-5-(4,4,5, 5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrole-3-carboxylate (784
 mg, 2.96 mmol), Pd(dppf)Cl₂ (201 mg, 247 μmol) and K₃PO₄ (1.57 g, 7.41 mmol) was added
 dioxane (5 mL) and H₂O (0.5 mL). The resulting mixture was stirred for 2 hours at 80 °C
 under N₂ atmosphere. The reaction mixture was extracted with ethyl acetate. The organic
 phase was concentrated under reduced pressure. The residue was purified by flash
 10 chromatography on silica gel eluting with 0-100% of ethyl acetate in petroleum ether to
 afford methyl 5-(5-chloro-3-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxylate (310 mg,
 47.6% yield) as a white solid. LCMS [M+H]⁺:269

Step 2: To a mixture of methyl 5-(5-chloro-3-fluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-
 carboxylate (310 mg, 1.15 mmol), 3-(trifluoromethyl)azetidine hydrochloride (277 mg, 1.72
 15 mmol), Pd₂(dba)₃ (105 mg, 115 μmol), Xantphos (132 mg, 230 μmol) and Cs₂CO₃ (1.12
 g, 3.44 mmol) was added PhMe (3 mL). The resulting mixture was stirred for 2 hours at
 100 °C under N₂ atmosphere. The reaction mixture was extracted with ethyl acetate. The
 organic phase was concentrated under reduced pressure. The residue was purified by flash
 chromatography on silica gel eluting with 0-100% of ethyl acetate in petroleum ether to
 20 afford methyl 5-{3-fluoro-5-[3-(trifluoromethyl)azetidin-1-yl]pyridin-2-yl}-1-methyl-1H-
 pyrrole-3-carboxylate (133 mg, 32.4% yield) as a yellow solid. LCMS [M+H]⁺: 358


Step 3: To a mixture of methyl 5-{3-fluoro-5-[3-(trifluoromethyl)azetidin-1-yl]pyridin-2-yl}-
 1-methyl-1H-pyrrole-3-carboxylate (133 mg, 372 μmol), N-(3-amino-5-
 chlorophenyl)methanesulfonamide (82.0 mg, 372 μmol) in toluene (0.8 mL) was stirred at
 25 room temperature. Then AlMe₃ (0.5 mL) in Toluene (0.7 mL) was added into the



solution and the resulting mixture was stirred at 80 °C for 2 hours. The reaction mixture was quenched with water. The reaction mixture was extracted with ethyl acetate. The organic phase was concentrated under reduced pressure, The residue was purified by reverse phase flash chromatography eluting with 70% of acetonitrile in water (0.1% NH₄HCO₃) to afford


- 5 *N*-(3-chloro-5-methanesulfonylamidophenyl)-5-{3-fluoro-5-[3-(trifluoromethyl)azetidin-1-yl]pyridin-2-yl}-1-methyl-1*H*-pyrrole-3-carboxamide (114.1 mg, 56.1% yield) as a light yellow solid. LCMS [M-H]⁻: 544.15. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.99 (s, 1H), 9.82 (s, 1H), 7.87 – 7.80 (m, 1H), 7.76 – 7.59 (m, 3H), 7.07 – 6.96 (m, 2H), 6.89 (t, *J* = 1.9 Hz, 1H), 4.21 (t, *J* = 8.6 Hz, 2H), 4.02 (dd, *J* = 8.7, 5.1 Hz, 2H), 3.81 (s, 4H), 3.06 (s, 3H).



10 Examples 525-531


The compounds listed in the following table were prepared using a procedure similar to that described for example 524:

Structure	Compound No.	MS (ESI) [M+H] ⁺	¹ H NMR
	525	527.9	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 10.03 (s, 1H), 9.83 (s, 1H), 7.95 (d, <i>J</i> = 2.1 Hz, 1H), 7.72 (d, <i>J</i> = 1.9 Hz, 1H), 7.70 – 7.49 (m, 2H), 7.09 (dd, <i>J</i> = 13.4, 2.4 Hz, 1H), 6.99 (dd, <i>J</i> = 3.4, 1.9 Hz, 1H), 6.89 (q, <i>J</i> = 1.9 Hz, 1H), 4.01 – 3.68 (m, 5H), 3.60 (t, <i>J</i> = 7.2 Hz, 2H), 3.06 (d, <i>J</i> = 1.7 Hz,


			3H), 2.59 (dq, J = 14.5, 7.1 Hz, 2H).
	526	511.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.88 (s, 1H), 8.26 (s, 2H), 7.74 (t, J = 1.9 Hz, 1H), 7.67 (t, J = 1.9 Hz, 1H), 7.61 (d, J = 2.1 Hz, 1H), 7.45 (d, J = 2.1 Hz, 1H), 6.89 (t, J = 1.9 Hz, 1H), 4.03 (s, 3H), 3.84 (t, J = 13.2 Hz, 2H), 3.62 (t, J = 7.2 Hz, 2H), 3.06 (d, J = 1.5 Hz, 3H), 2.58 (dq, J = 14.5, 7.1 Hz, 2H).
	527	510.15	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.99 (s, 1H), 9.78 (s, 1H), 8.04 (d, J = 2.9 Hz, 1H), 7.72 (d, J = 2.1 Hz, 1H), 7.64 – 7.57 (m, 2H), 7.54 (d, J = 8.7

			Hz, 1H), 7.11 (dd, J = 8.8, 3.0 Hz, 1H), 6.99 (d, J = 2.0 Hz, 1H), 6.88 (d, J = 2.2 Hz, 1H), 3.91 (s, 3H), 3.79 (t, J = 13.2 Hz, 2H), 3.57 (t, J = 7.2 Hz, 2H), 3.06 (s, 3H), 2.57 (dq, J = 14.6, 7.2 Hz, 2H).
	528	573.05	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.84 (s, 1H), 7.58 (dt, J = 36.1, 2.0 Hz, 3H), 7.02 (t, J = 2.0 Hz, 1H), 6.75 (d, J = 2.0 Hz, 1H), 4.39 (t, J = 11.9 Hz, 4H), 3.94 (t, J = 6.3 Hz, 2H), 3.88 (s, 3H), 3.34 (d, J = 6.3 Hz, 2H), 2.39 (s, 3H).

	529	593.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.41 (s, 1H), 10.04 (s, 1H), 8.12 (d, J = 1.6 Hz, 1H), 7.87 (t, J = 1.9 Hz, 1H), 7.67 (t, J = 1.9 Hz, 1H), 7.63 (t, J = 1.9 Hz, 1H), 7.02 (dd, J = 12.5, 2.3 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 4.94 (t, J = 5.7 Hz, 1H), 4.22 (t, J = 8.6 Hz, 2H), 4.03 (dd, J = 8.7, 5.1 Hz, 2H), 3.89 – 3.60 (m, 3H), 3.30 (s, 2H), 2.49 – 2.47 (s, 3H).
	530	522	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.79 (s, 1H), 7.98 (d, J = 2.9 Hz, 1H), 7.70 (dt, J = 4.3, 1.9 Hz, 2H), 7.61 (d, J

			= 2.0 Hz, 1H), 7.57 (d, J = 8.6 Hz, 1H), 7.06 (s, 1H), 7.03 (d, J = 2.1 Hz, 1H), 6.91 (t, J = 2.0 Hz, 1H), 4.39 (t, J = 12.3 Hz, 4H), 3.92 (s, 3H), 2.69 (tt, J = 7.8, 5.0 Hz, 1H), 0.99 (ddt, J = 10.7, 8.0, 2.5 Hz, 4H).
	531	524	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.08 (s, 1H), 9.81 (s, 1H), 7.99 (d, J = 2.9 Hz, 1H), 7.73 (t, J = 1.9 Hz, 1H), 7.62 (s, 2H), 7.57 (d, J = 8.6 Hz, 1H), 7.17 – 6.97 (m, 2H), 6.90 (t, J = 2.0 Hz, 1H), 4.40 (t, J = 12.3 Hz, 4H), 3.92 (s, 3H), 3.18 – 2.98 (m, 2H), 1.87 – 1.52 (m, 2H), 0.96 (t, J = 7.4 Hz, 3H).

Example 532: *N*-(3-chloro-5-methanesulfonamidophenyl)-4-[5-(3,3-difluoroazetidin-1-yl)-3-fluoropyridin-2-yl]-5-methylthiophene-2-carboxamide



- 5 Step 1: To a stirred solution of 5-bromo-2-chloro-3-fluoropyridine (1.03 g, 4.89 mmol), Xantphos (565 mg, 0.978 mmol), Pd₂(dba)₃ (447 mg, 0.489 mmol), 3,3-difluoroazetidine hydrochloride (695 mg, 5.37 mmol) and t-BuOK (1.63 g, 14.6 mmol) in PhMe (25 mL) at room temperature. Then the mixture was stirred for 5 hours at 110 °C under nitrogen atmosphere. The mixture was concentrated under reduced pressure. The residue was purified
- 10 by reversed-phase flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in water (0.1% formic acid), 0% to 45% gradient in 20 min; detector, UV 254 nm. This resulted in 2-chloro-5-(3,3-difluoroazetidin-1-yl)-3-fluoropyridine (180 mg, 0.808 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 223.




- Step 2: A solution of 2-chloro-5-(3,3-difluoroazetidin-1-yl)-3-fluoropyridine (180 mg, 0.808 mmol), Pd(dppf)Cl₂ (66 mg, 0.0808 mmol), K₃PO₄ (513 mg, 2.42 mmol) and methyl 5-methyl-4-(4,4,5,5-tetra methyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxylate (250 mg, 0.888 mmol) in dioxane (10 mL) and H₂O (1 mL) was stirred for 2 hours at 80 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (1:2) to
- 15





afford methyl 4-[5-(3,3-difluoroazetidin-1-yl)-3-fluoropyridin-2-yl]-5-methylthiophene-2-carboxylate (180 mg, 0.525 mmol) as a white solid. LCMS (ESI) $[M + H]^+$: 343.

- Step 3: To a stirred solution of methyl 4-[5-(3,3-difluoroazetidin-1-yl)-3-fluoropyridin-2-yl]-5-methylthiophene-2-carboxylate (160 mg, 0.467 mmol) in EtOH (5 mL) and H₂O (1 mL) was added LiOH (111 mg, 4.67 mmol) at 0 °C. Then the mixture was stirred for 2 hours at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 4-[5-(3,3-difluoroazetidin-1-yl)-3-fluoropyridin-2-yl]-5-methylthiophene-2-carboxylic acid (160 mg, 0.487 mmol) as a yellow solid. LCMS (ESI) $[M + H]^+$: 329.
- Step 4: To a stirred solution of 4-[5-(3,3-difluoroazetidin-1-yl)-3-fluoropyridin-2-yl]-5-methylthiophene-2-carboxylic acid (160 mg, 0.487 mmol), TCFH (408 mg, 1.46 mmol) and NMI (80 mg, 0.974 mmol) in ACN (5 mL) was added *N*-(3-amino-5-chlorophenyl)methanesulfonamide (128 mg, 0.584 mmol) at room temperature. Then the mixture was stirred for 1 hour at room temperature. The mixture was concentrated under reduced pressure. The residue was purified by reversed-phase flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in water (0.1% formic acid), 0% to 50% gradient in 20 min; detector, UV 254 nm. This resulted in *N*-(3-chloro-5-methanesulfonamidophenyl)-4-[5-(3,3-difluoroazetidin-1-yl)-3-fluoropyridin-2-yl]-5-methylthiophene-2-carboxamide (75.6 mg, 0.142 mmol) as a white solid. LCMS (ESI) $[M + H]^+$: 530.95. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.41 (s, 1H), 9.78 (s, 1H), 8.13 (d, *J* = 1.6 Hz, 1H), 7.93 (t, *J* = 2.0 Hz, 1H), 7.65 (dt, *J* = 14.9, 1.9 Hz, 2H), 7.12 (dd, *J* = 12.3, 2.3 Hz, 1H), 6.94 (t, *J* = 1.9 Hz, 1H), 4.46 (t, *J* = 12.3 Hz, 4H), 3.06 (s, 3H), 2.49 (s, 3H).

Examples 533-539


The compounds listed in the following table were prepared using a procedure similar to that described for example 532:

Structure	Compound No.	MS (ESI) [M+H] ⁺	¹ H NMR
	533	512.85	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 10.06 (s, 1H), 8.31 (s, 1H), 8.05 (d, J = 2.9 Hz, 1H), 7.66 (dt, J = 12.9, 1.9 Hz, 2H), 7.56 (d, J = 8.6 Hz, 1H), 7.12 (dd, J = 8.6, 3.0 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 4.42 (t, J = 12.3 Hz, 4H), 3.07 (s, 3H), 2.66 (s, 3H).
	534	571.15	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.33 (s, 1H), 10.05 (s, 1H), 8.02 (s, 1H), 7.63 (dd, J = 13.3, 2.2 Hz, 3H), 6.93 (t, J = 1.9 Hz, 1H), 6.73 (d, J = 2.2 Hz, 1H), 4.66 (p, J = 5.9 Hz, 1H), 4.41 (t, J = 12.3 Hz, 4H), 3.05 (s, 3H), 2.39 (s, 3H), 1.24 (d, J = 6.0 Hz, 6H).
	535	543.1	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.34 (s, 1H), 10.11 (s, 1H), 8.05 (s, 1H), 7.65 (dt, J = 13.3, 2.2 Hz, 3H), 6.93 (t, J = 1.9 Hz, 1H), 6.77 (d, J = 2.3 Hz, 1H), 4.44 (t, J = 12.3 Hz, 4H), 3.85 (s, 3H), 3.06 (s, 3H), 2.38 (s, 3H).

	536	579	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.39 (s, 1H), 10.07 (s, 1H), 8.04 (s, 1H), 7.94 (d, J = 2.3 Hz, 1H), 7.67 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.36 (d, J = 73.3 Hz, 1H), 7.01 – 6.91 (m, 2H), 4.48 (t, J = 12.3 Hz, 4H), 3.06 (s, 3H), 2.40 (s, 3H).
	537	526.85	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.30 (s, 1H), 10.06 (s, 1H), 7.93 (s, 1H), 7.88 (d, J = 2.7 Hz, 1H), 7.66 (d, J = 1.9 Hz, 1H), 7.61 (t, J = 1.9 Hz, 1H), 6.98 (d, J = 2.7 Hz, 1H), 6.94 (t, J = 2.0 Hz, 1H), 4.39 (t, J = 12.3 Hz, 4H), 3.06 (s, 3H), 2.31 (s, 3H), 2.18 (s, 3H).
	538	460	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.05 (s, 1H), 9.77 (s, 1H), 7.82 (d, J = 2.9 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.58 (d, J = 2.0 Hz, 1H), 7.48 (d, J = 8.6 Hz, 1H), 6.96 (d, J = 2.0 Hz, 1H), 6.93 – 6.85 (m, 2H), 3.95 – 3.87 (m, 7H), 3.06 (s, 3H), 2.37 (p, J = 7.3 Hz, 2H).
	539	478	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.77 (s, 1H), 7.90 (d, J = 2.8 Hz, 1H), 7.71 (t, J = 1.8 Hz, 1H), 7.65 – 7.43 (m, 3H), 7.03 – 6.94 (m, 2H), 6.88 (t, J = 1.9 Hz, 1H), 5.53 (ddd, J = 60.0, 5.8, 2.8 Hz, 1H), 4.26 (ddd, J = 20.8, 9.8,

			6.1 Hz, 2H), 4.04 (dd, J = 9.1, 2.5 Hz, 1H), 3.96 (dd, J = 10.2, 3.2 Hz, 1H), 3.90 (s, 3H), 3.05 (s, 3H).
--	--	--	---

Example 540: N-(3-chloro-5-methanesulfonamidophenyl)-5-(3,5-difluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxamide




- 5 Step 1: To a mixture of methyl 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrole-3-carboxylate (116 g, 440 mmol, 1 equiv), 2-bromo-3,5-difluoropyridine (85 g, 440 mmol, 1 equiv), Pd(dppf)Cl₂ (18 g, 22 mmol, 0.05 equiv) and K₃PO₄ (200 g, 880 mmol, 2 equiv) was added dioxane (1 L) and H₂O (100 mL). The resulting mixture was stirred for 2 hours at 80 °C under N₂ atmosphere. The reaction mixture was extracted with ethyl acetate.
- 10 The organic phase was concentrated under reduced pressure. The residue was purified by flash chromatography on silica gel eluting with 30% of ethyl acetate in petroleum ether to afford methyl 5-(3,5-difluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxylate (98g, 88.6% yield) as a white solid.

- Step 2: To a stirred solution of methyl 5-(3,5-difluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxylate (295 g, 1165 mmol, 1 equiv) in tBuOH (3 L) and water (2 L) was added LiOH (246g, 5853 mmol, 5 equiv, in 1L water). The mixture was stirred for 4 hours at 60 °C. The pH value of the solution was adjusted to 3 with 1M of HCl (aq). The product was precipitated from the solution. The mixture was filtered and the filter cake was washed with water (3 x 200 mL) to afford 5-(3,5-difluoropyridin-2-yl)-1-methyl-1H-pyrrole-3-carboxylic acid (212 g, 76.0% yield) as a white solid. LCMS [M+H]⁺: 239.
- 20

Step 3: A mixture of 5-(3,5-difluoropyridin-2-yl)-1-methyl-1*H*-pyrrole-3-carboxylic acid (212 g, 886 mmol, 1 equiv), TCFH (341 g, 1219 mmol, 1.37 equiv), NMI (199 g, 2438 mmol, 2.75 equiv) and *N*-(3-amino-5-chlorophenyl)methanesulfonamide (214 g, 487 mmol) in ACN (2 L) was stirred for two hours at room temperature. The mixture was quenched with water (2 L). The resulting mixture was extracted with DCM (3 x 3 L) and water. The combined organic layers were dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with DCM / ACN (5:1) to afford *N*-(3-chloro-5-methanesulfonamidophenyl)-5-(3,5-difluoropyridin-2-yl)-1-methyl-1*H*-pyrrole-3-carboxamide (155g, 39.5% yield) as a white solid. LCMS [M+H]⁺: 441. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.00 (s, 1H), 9.90 (s, 1H), 8.59 (d, *J* = 2.4 Hz, 1H), 8.08 (ddd, *J* = 11.1, 8.9, 2.5 Hz, 1H), 7.79 – 7.69 (m, 2H), 7.64 (t, *J* = 1.9 Hz, 1H), 7.21 (dd, *J* = 3.6, 1.9 Hz, 1H), 6.91 (t, *J* = 2.0 Hz, 1H), 3.89 (s, 3H), 3.06 (s, 3H).

Example 541: *N*-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-isopropoxythiophen-2-yl)-5-methylthiophene-2-carboxamide



Step 1: A mixture of 2-bromopyrimidin-5-ol (15.0 g, 85.7 mmol, 1 equiv), K₂CO₃ (35.48 g, 257.1 mmol, 3 equiv) and 2-bromopropane (13.71 g, 111.5 mmol, 1.3 equiv) in DMF (150

mL) was stirred for 5 hours at room temperature. The reaction mixture was diluted with water (1 L). The resulting solution was extracted with EA (3 x 800 mL) and the organic layers were combined. The organic layer was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by flash chromatography on silica gel column eluting with EA/PE (20%) to afford 2-bromo-5-isopropoxypyrimidine (15.0 g, 81% yield) as a white solid. LCMS (ESI) [M + H]⁺: 217

Step 2: A mixture of 2-bromo-5-isopropoxypyrimidine (14.0 g, 64.5 mmol, 1 equiv), Pd(dppf)Cl₂ (3.3 g, 4.5 mmol, 0.07 equiv), K₃PO₄ (41.1 g, 193.5 mmol, 3 equiv) and methyl 5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl) thiophene-2-carboxylate (23.66 g, 83.8 mmol, 1.3 equiv) in 1,4-dioxane (150 mL) and H₂O (15 mL) was stirred for 2 hours at 80 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (32%) to afford methyl 4-(5-isopropoxypyrimidin-2-yl)-5-methylthiophene-2-carboxylate (12.70 g, 67% yield) as a white solid. LCMS (ESI) [M + H]⁺: 293




Step 3: To a stirred solution of methyl 4-(5-isopropoxypyrimidin-2-yl)-5-methylthiophene-2-carboxylate (12.7 g, 43.4 mmol, 1 equiv) in EtOH (150 mL) and H₂O (150 mL) was added LiOH (10.4 g, 434.4 mmol, 10 equiv). Then the mixture was stirred for 4 hours at room temperature. The mixture was acidified to pH 5 with HCl (aq.). The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 4-(5-isopropoxypyrimidin-2-yl)-5-methylthiophene-2-carboxylic acid (10.4 g, 86% yield) as a white solid. LCMS (ESI) [M + H]⁺: 279





Step 4: To a stirred solution of 4-(5-isopropoxypyrimidin-2-yl)-5-methylthiophene-2-carboxylic acid (10.4 g, 37.4 mmol, 1 equiv), TCFH (20.97 g, 74.7 mmol, 2 equiv) and NMI (12.27 g, 149.5 mmol, 4 equiv) in ACN (180 mL) was added *N*-(3-amino-5-chlorophenyl)methanesulfonamide (8.25 g, 37.4 mmol, 1 equiv). Then the mixture was stirred for 2 hours at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (56%, 0.1%FA) to afford *N*-(3-chloro-5-(methylsulfonamido)phenyl)-4-(5-isopropoxypyrimidin-2-yl)-5-methylthiophene-2-carboxamide (10.0113 g, 55.7% yield) as a white solid. LCMS (ESI) [M + H]⁺: 481. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.57 (s, 1H), 10.06 (s, 1H), 8.64 (d, *J* = 2.7 Hz, 3H), 7.69 (dt, *J* = 6.5, 1.9 Hz, 2H), 6.95 (t, *J* = 2.0 Hz, 1H), 4.88 (p, *J* = 6.0 Hz, 1H), 3.08 (s, 3H), 2.83 (s, 3H), 1.34 (d, *J* = 6.0 Hz, 6H).





Examples 542-567




The compounds listed in the following table were prepared using a procedure similar to that described for examples 540 and 541:





5




Structure	Compound No.	MS (ESI) [M+H] ⁺	¹ H NMR
	542	437.95	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.70 (s, 2H), 7.59 (ddt, J = 25.9, 23.5, 2.0 Hz, 4H), 6.99 (d, J = 2.1 Hz, 1H), 4.11 (s, 3H), 3.17 (q, J = 7.4 Hz, 2H), 1.33 (t, J = 7.3 Hz, 3H).
	543	454.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.05 (s, 1H), 9.89 (s, 1H), 8.60 (d, J = 2.4 Hz, 1H), 8.09 (ddd, J = 11.1, 8.9, 2.5 Hz, 1H), 7.76 (d, J = 2.0 Hz, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 7.20 (dd, J = 3.6, 1.9 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H), 3.88 (s, 3H), 3.16 (q, J = 7.3 Hz, 2H), 1.21 (t, J = 7.3 Hz, 3H).
	544	456.85	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.22 – 9.81 (m, 2H), 8.70 (d, J = 2.6 Hz, 1H), 8.25 (dd, J = 8.5, 2.6 Hz, 1H), 7.84 – 7.49 (m, 3H), 7.15 (d, J = 1.9 Hz, 1H), 6.90 (t, J = 2.0 Hz, 1H), 3.70 (s, 3H), 3.06 (s, 3H).





	545	436.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.06 (s, 1H), 9.80 (s, 1H), 8.51 (d, J = 2.8 Hz, 1H), 7.77 (dd, J = 9.7, 2.9 Hz, 1H), 7.73 – 7.66 (m, 2H), 7.63 (t, J = 2.0 Hz, 1H), 6.90 (q, J = 1.9 Hz, 2H), 3.64 (s, 3H), 3.06 (s, 3H), 2.41 (s, 3H).
	546	448.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.03 (s, 1H), 9.94 (s, 1H), 8.96 (d, J = 2.9 Hz, 1H), 8.56 (dd, J = 8.6, 2.9 Hz, 1H), 7.84 (d, J = 1.9 Hz, 1H), 7.71 (d, J = 1.9 Hz, 1H), 7.62 (d, J = 2.0 Hz, 1H), 7.35 (d, J = 1.9 Hz, 1H), 6.91 (d, J = 2.0 Hz, 1H), 3.83 (s, 3H), 3.06 (s, 3H).
	547	458	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.82 (s, 1H), 7.75 (d, J = 1.9 Hz, 1H), 7.70 (d, J = 1.8 Hz, 1H), 7.60 (d, J = 1.9 Hz, 1H), 7.40 (dd, J = 9.2, 7.8 Hz, 2H), 6.89 (t, J = 2.0 Hz, 1H), 6.82 (d, J = 1.9 Hz, 1H), 3.51 (s, 3H), 3.05 (s, 3H).
	548	451	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.07 (s, 1H), 9.88 (s, 1H), 8.36 (s, 1H), 7.77 – 7.61 (m, 4H), 7.18 (dd, J = 3.6, 1.9 Hz, 1H), 6.91 (t, J = 1.9 Hz,


			1H), 3.90 (s, 3H), 3.17 (q, J = 7.3 Hz, 2H), 2.37 (s, 3H), 1.22 (t, J = 7.3 Hz, 3H).
	549	436.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.92 (s, 1H), 8.58 (s, 2H), 7.74 (d, J = 1.9 Hz, 1H), 7.66 (d, J = 2.1 Hz, 2H), 7.56 (d, J = 2.1 Hz, 1H), 6.89 (t, J = 2.0 Hz, 1H), 4.05 (s, 3H), 3.94 (s, 3H), 3.06 (s, 3H).
	550	519.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.41 (s, 1H), 10.06 (s, 1H), 8.63 (s, 1H), 8.16 (s, 1H), 7.92 (d, J = 1.8 Hz, 1H), 7.71 – 7.54 (m, 2H), 6.94 (t, J = 2.0 Hz, 1H), 3.97 (s, 3H), 3.06 (s, 3H), 2.46 (s, 3H).
	551	495.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.82 (s, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.69 – 7.56 (m, 3H), 7.19 (d, J = 2.0 Hz, 1H), 7.12 (d, J = 7.5 Hz, 1H), 6.89 (t, J = 1.9 Hz, 1H), 6.45 (d, J = 8.1 Hz, 1H), 4.42 (t, J = 12.4 Hz, 4H), 3.99 (s, 3H), 3.06 (s, 3H).
	552	496.05	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.31 – 8.11 (m, 2H), 7.65 (t, J = 1.9 Hz, 1H), 7.61 (d, J = 2.0 Hz, 1H), 7.55 (t, J = 1.8 Hz, 1H),

			7.05 (d, J = 2.0 Hz, 1H), 6.98 (t, J = 1.9 Hz, 1H), 6.76 (d, J = 2.4 Hz, 1H), 6.58 (dd, J = 6.2, 2.4 Hz, 1H), 4.51 (t, J = 11.9 Hz, 4H), 3.87 (s, 3H), 3.03 (s, 3H).
	553	471.85	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.99 (d, J = 18.0 Hz, 2H), 8.79 (s, 2H), 7.82 – 7.60 (m, 4H), 7.29 (d, J = 73.0 Hz, 1H), 6.90 (d, J = 2.0 Hz, 1H), 4.07 (s, 3H), 3.07 (s, 3H).
	554	490.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.97 (s, 1H), 8.87 (s, 1H), 8.47 – 8.31 (m, 1H), 7.86 (d, J = 1.9 Hz, 1H), 7.71 (q, J = 1.6 Hz, 1H), 7.65 (q, J = 1.7 Hz, 1H), 7.46 (dd, J = 4.1, 1.9 Hz, 1H), 6.92 (q, J = 1.6 Hz, 1H), 4.00 (s, 3H), 3.07 (d, J = 1.3 Hz, 3H).
	555	477	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.03 (s, 1H), 9.84 (d, J = 2.1 Hz, 1H), 8.28 (dd, J = 2.8, 0.7 Hz, 1H), 7.73 (dq, J = 3.7, 1.8 Hz, 1H), 7.68 – 7.61 (m, 3H), 7.50 (ddd, J = 8.7, 3.2, 1.6 Hz, 1H), 7.18 – 7.11 (m, 1H), 6.93 – 6.75 (m, 1H), 4.11 – 3.81 (m,


			3H), 3.16 – 3.04 (m, 3H), 1.34 (d, J = 0.9 Hz, 9H).
	556	476.15 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.94 (s, 1H), 8.55 (s, 2H), 7.85 – 7.42 (m, 4H), 6.90 (t, J = 2.0 Hz, 1H), 4.08 (s, 3H), 3.07 (s, 3H), 1.36 (s, 9H).
	557	493.1 [M-H] ⁻	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.88 (s, 1H), 8.23 (dd, J = 2.3, 1.2 Hz, 1H), 7.72 (q, J = 1.9 Hz, 2H), 7.67 – 7.53 (m, 2H), 7.15 (dd, J = 3.5, 1.9 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H), 3.89 (s, 3H), 3.07 (s, 3H), 1.39 (s, 9H).
	558	425.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.91 (s, 1H), 8.60 (d, J = 2.4 Hz, 1H), 8.09 (ddd, J = 11.1, 8.9, 2.4 Hz, 1H), 7.76 (d, J = 1.9 Hz, 1H), 7.54 – 7.34 (m, 2H), 7.20 (dd, J = 3.7, 1.9 Hz, 1H), 6.67 (dt, J = 10.4, 2.2 Hz, 1H), 3.88 (s, 3H), 3.06 (s, 3H).
	559	495	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.57 (s, 1H), 10.07 (s, 1H), 8.66 (d, J = 13.9 Hz, 3H), 7.69 (dt, J = 8.3, 1.8 Hz, 2H), 6.95 (t, J = 1.9 Hz, 1H), 3.08 (s, 3H), 2.86 (s, 3H), 1.38 (s, 9H).

	560	427.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 12.14 (s, 1H), 9.94 (s, 2H), 8.54 (d, J = 2.2 Hz, 1H), 8.05 (ddt, J = 11.2, 8.8, 2.4 Hz, 1H), 7.76 (dd, J = 3.3, 1.6 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.67 (t, J = 1.8 Hz, 1H), 7.31 (dt, J = 4.2, 2.1 Hz, 1H), 6.91 (t, J = 2.0 Hz, 1H), 3.07 (s, 3H).
	561	466.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.82 (s, 1H), 8.22 (d, J = 2.4 Hz, 1H), 7.76 – 7.55 (m, 4H), 7.10 (d, J = 2.0 Hz, 1H), 6.89 (t, J = 2.0 Hz, 1H), 4.18 (q, J = 6.9 Hz, 2H), 3.76 (s, 3H), 3.06 (s, 3H), 1.38 (t, J = 6.9 Hz, 3H).
	562	490 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.11 (s, 1H), 10.05 (s, 1H), 8.94 (q, J = 0.9 Hz, 2H), 7.87 (d, J = 2.0 Hz, 1H), 7.80 (d, J = 2.0 Hz, 1H), 7.73 (t, J = 1.9 Hz, 1H), 7.68 (q, J = 1.6 Hz, 1H), 6.93 (td, J = 2.0, 0.7 Hz, 1H), 5.75 (q, J = 9.0 Hz, 2H), 3.08 (d, J = 0.9 Hz, 3H).

	563	503.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.04 (d, J = 13.6 Hz, 2H), 8.60 (s, 2H), 7.79 (d, J = 2.0 Hz, 1H), 7.73 (t, J = 1.9 Hz, 1H), 7.70 – 7.66 (m, 2H), 6.92 (t, J = 2.0 Hz, 1H), 5.77 (q, J = 9.1 Hz, 2H), 3.95 (s, 3H), 3.06 (s, 3H).
	564	491	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.06 (s, 1H), 10.03 (s, 1H), 8.50 (dt, J = 4.7, 1.6 Hz, 1H), 7.98 – 7.83 (m, 2H), 7.67 (dt, J = 24.5, 2.0 Hz, 2H), 7.49 – 7.38 (m, 2H), 6.93 (t, J = 2.0 Hz, 1H), 5.68 (q, J = 9.2 Hz, 2H), 3.06 (s, 3H).
	565	495	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.55 (s, 1H), 10.09 (s, 1H), 8.63 (s, 3H), 7.68 (d, J = 2.0 Hz, 2H), 6.95 (t, J = 1.9 Hz, 1H), 4.88 (p, J = 6.0 Hz, 1H), 3.18 (q, J = 7.3 Hz, 2H), 2.83 (s, 3H), 1.34 (d, J = 6.0 Hz, 6H), 1.22 (t, J = 7.3 Hz, 3H).
	566	509.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.57 (s, 1H), 10.10 (s, 1H), 8.66 (d, J = 11.5 Hz, 3H), 7.68 (p, J = 1.9 Hz, 2H), 6.95 (t, J = 1.9 Hz, 1H), 3.18 (q, J = 7.3 Hz, 2H), 2.86 (s, 3H), 1.38

		(s, 9H), 1.22 (t, J = 7.3 Hz, 3H).
	567	457 ¹ H NMR (400 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.86 (s, 1H), 8.72 (dt, J = 4.0, 1.8 Hz, 1H), 7.87 – 7.78 (m, 2H), 7.71 (t, J = 1.8 Hz, 1H), 7.61 (t, J = 1.9 Hz, 1H), 7.01 – 6.72 (m, 2H), 3.57 (s, 3H), 3.06 (s, 3H).

Example 568: N-(3-chloro-5-methanesulfonamidophenyl)-5-(5-fluoro-3-methoxypyridin-2-yl)-1-methylpyrrole-3-carboxamide






- 5 Step 1: A solution of 2-bromo-5-fluoro-3-methoxypyridine (212 mg, 1.029 mmol, 1 equiv), methyl 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrole-3-carboxylate (273 mg, 1.029 mmol, 1 equiv), Pd(dppf)Cl₂ (84.04 mg, 0.103 mmol, 0.1 equiv) and K₂CO₃ (426.03 mg, 3.087 mmol, 3 equiv) in 1,4-dioxane (5 ml) and H₂O (1 ml) was stirred for 1.5
- 10 temperature. The resulting mixture was diluted with water and extracted with EtOAc (3 x 15 mL). The combined organic layers were washed with brine (3x10 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (1:1) to afford methyl 5-(5-fluoro-3-methoxypyridin-2-yl)-1-methylpyrrole-3-carboxylate (80 mg, 29% yield) as a
- 15 white solid. LCMS (ESI) [M+H]⁺: 265.09




Step 2: To a stirred solution of methyl 5-(5-fluoro-3-methoxypyridin-2-yl)-1-methylpyrrole-3-carboxylate (80 mg, 0.303 mmol, 1 equiv) and *N*-(3-amino-5-chlorophenyl)methanesulfonamide (67 mg, 0.303 mmol, 1 equiv) in PhMe (2 ml) was added $\text{Al}(\text{Me})_3$ (0.5 ml) dropwise at 0 °C. The resulting mixture was stirred for additional 1 hour at 80 °C. The mixture was allowed to cool down to room temperature. The reaction was quenched by the addition of water (1 mL) at 0 °C. The resulting mixture was concentrated under reduced pressure. The crude product (60 mg) was purified by Prep-HPLC to afford *N*-(3-chloro-5-methanesulfonamidophenyl)-5-(5-fluoro-3-methoxypyridin-2-yl)-1-methylpyrrole-3-carboxamide (48.8 mg, 35% yield) as a white solid. LCMS (ESI) $[\text{M} + \text{H}]^+$: 452.85. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.03 (s, 1H), 9.82 (s, 1H), 8.24 (d, $J = 2.4$ Hz, 1H), 7.71 (d, $J = 2.2$ Hz, 1H), 7.66 (d, $J = 2.0$ Hz, 1H), 7.65 – 7.60 (m, 2H), 7.09 (d, $J = 2.0$ Hz, 1H), 6.89 (t, $J = 2.0$ Hz, 1H), 3.92 (s, 3H), 3.75 (s, 3H), 3.06 (s, 3H).





Examples 569-585




The compounds listed in the following table were prepared using a procedure similar to that described for example 568:



Structure	Compound No.	MS (ESI) $[\text{M} + \text{H}]^+$	^1H NMR
	569	473.25	^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.02 (s, 1H), 9.91 (s, 1H), 8.95 (s, 1H), 8.21 (dd, $J = 8.9, 2.4$ Hz, 1H), 7.94 (d, $J = 8.4$ Hz, 1H), 7.78 (d, $J = 2.0$ Hz, 1H), 7.72 (s, 1H), 7.63 (s, 1H), 7.46 (d, $J = 2.0$ Hz, 1H), 6.91 (s, 1H), 4.05 (s, 3H), 3.07 (s, 3H).
	570	429.9	^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.01 (s, 1H), 9.93 (s, 1H), 8.99 (t, $J = 1.5$ Hz, 1H), 8.28 (dd, $J = 8.5, 2.2$ Hz, 1H), 7.95 – 7.88 (m,

			1H), 7.81 (d, J = 1.9 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.50 (d, J = 2.0 Hz, 1H), 6.91 (t, J = 1.9 Hz, 1H), 4.04 (s, 3H), 3.07 (s, 3H).
	571	473.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.04 (s, 2H), 9.21 (s, 2H), 7.88 (dd, J = 29.9, 2.1 Hz, 2H), 7.71 (dt, J = 28.6, 1.9 Hz, 2H), 6.91 (t, J = 2.0 Hz, 1H), 4.12 (s, 3H), 3.07 (s, 3H).
	572	449.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.03 (s, 1H), 9.81 (s, 1H), 8.31 (d, J = 2.9 Hz, 1H), 7.73 (t, J = 1.8 Hz, 1H), 7.63 (t, J = 2.6 Hz, 3H), 7.45 (dd, J = 8.8, 3.0 Hz, 1H), 7.07 (d, J = 1.9 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H), 4.15 (q, J = 6.9 Hz, 2H), 3.93 (s, 3H), 3.07 (s, 3H), 1.37 (t, J = 7.0 Hz, 3H).
	573	434.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.04 (s, 1H), 9.81 (s, 1H), 8.33 (d, J = 3.0 Hz, 1H), 7.73 (t, J = 1.9 Hz, 1H), 7.68 – 7.61 (m, 3H), 7.47 (dd, J = 8.8, 3.0 Hz, 1H), 7.08 (d, J = 2.0 Hz, 1H), 6.90 (t,


			J = 2.0 Hz, 1H), 3.93 (s, 3H), 3.87 (s, 3H), 3.07 (s, 3H).
	574	449	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.85 (s, 1H), 8.54 (d, J = 2.2 Hz, 1H), 7.79 (dd, J = 8.2, 2.2 Hz, 1H), 7.75 – 7.66 (m, 3H), 7.63 (t, J = 2.0 Hz, 1H), 7.24 (d, J = 2.0 Hz, 1H), 6.90 (t, J = 2.0 Hz, 1H), 4.47 (s, 2H), 4.00 (s, 3H), 3.34 (s, 3H), 3.07 (s, 3H).
	575	423	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.84 (s, 1H), 8.60 (d, J = 2.4 Hz, 1H), 7.85 – 7.74 (m, 2H), 7.74 – 7.66 (m, 2H), 7.61 (d, J = 2.2 Hz, 1H), 7.20 (d, J = 2.0 Hz, 1H), 6.89 (d, J = 2.0 Hz, 1H), 3.95 (s, 3H), 3.06 (s, 3H).
	576	438.85	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.86 (s, 1H), 8.63 (d, J = 2.5 Hz, 1H), 7.96 (dd, J = 8.6, 2.6 Hz, 1H), 7.79 – 7.59 (m, 4H), 7.28 (d, J = 2.0 Hz, 1H), 6.90 (d, J = 2.0 Hz, 1H), 3.98 (s, 3H), 3.06 (s, 3H).

	577	438 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.97 (s, 1H), 9.69 (s, 1H), 8.91 (s, 2H), 7.82 – 7.70 (m, 3H), 7.65 (t, J = 1.9 Hz, 1H), 6.89 (t, J = 2.0 Hz, 1H), 4.07 (s, 3H), 3.05 (s, 3H).
	578	439.85	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.03 (d, J = 12.0 Hz, 2H), 9.00 (s, 2H), 8.47 (d, J = 2.1 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.66 (t, J = 1.9 Hz, 1H), 6.91 (t, J = 1.9 Hz, 1H), 6.59 (dd, J = 2.1, 1.1 Hz, 1H), 3.06 (s, 3H), 2.58 (d, J = 1.1 Hz, 3H).
	579	437.15	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.06 (s, 1H), 9.85 (s, 1H), 8.60 (t, J = 1.7 Hz, 1H), 7.85 – 7.74 (m, 2H), 7.71 (t, J = 1.9 Hz, 1H), 7.69 (d, J = 2.0 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.20 (d, J = 2.0 Hz, 1H), 6.90 (t, J = 2.0 Hz, 1H), 3.95 (s, 3H), 3.17 (q, J = 7.3 Hz, 2H), 1.22 (t, J = 7.3 Hz, 3H).
	580	437.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.94 (d, J = 40.3 Hz, 2H), 8.35 (q, J = 1.5 Hz, 1H), 7.76 – 7.60 (m, 4H),

			7.17 (dd, J = 3.5, 1.9 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H), 3.90 (s, 3H), 3.06 (s, 3H), 2.37 (s, 3H).
	581	471.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.90 (s, 1H), 8.60 (d, J = 2.4 Hz, 1H), 8.09 (ddd, J = 11.2, 8.9, 2.4 Hz, 1H), 7.76 (d, J = 1.9 Hz, 1H), 7.71 (t, J = 1.8 Hz, 1H), 7.63 (t, J = 1.9 Hz, 1H), 7.20 (dd, J = 3.6, 1.9 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H), 4.95 (t, J = 5.7 Hz, 1H), 3.88 (s, 3H), 3.80 – 3.69 (m, 2H), 3.30 (d, J = 6.5 Hz, 2H).
	582	511.05	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.51 (s, 2H), 8.45 (s, 1H), 7.65 (d, J = 2.0 Hz, 1H), 7.56 (t, J = 1.9 Hz, 1H), 7.03 (d, J = 2.0 Hz, 1H), 4.84 – 4.75 (m, 1H), 3.95 (t, J = 6.3 Hz, 2H), 3.35 (t, J = 6.3 Hz, 2H), 2.85 (s, 3H), 1.40 (d, J = 6.0 Hz, 6H).
	583	438.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.56 (d, J = 9.8 Hz, 2H), 10.06 (s, 1H), 8.61 (s, 1H), 8.45 (s, 2H), 7.69 (dt, J = 6.8, 2.0 Hz, 2H),

			6.94 (t, J = 2.0 Hz, 1H), 3.07 (s, 3H), 2.81 (s, 3H).
	584	475	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.83 (s, 1H), 8.20 (dd, J = 4.6, 1.3 Hz, 1H), 7.75 – 7.65 (m, 2H), 7.62 (t, J = 1.9 Hz, 1H), 7.52 (dd, J = 8.5, 1.3 Hz, 1H), 7.29 (dd, J = 8.4, 4.6 Hz, 1H), 7.17 (d, J = 2.0 Hz, 1H), 6.89 (t, J = 1.9 Hz, 1H), 3.95 (d, J = 6.9 Hz, 2H), 3.81 (s, 3H), 3.05 (s, 3H), 1.34 – 1.18 (m, 1H), 0.64 – 0.52 (m, 2H), 0.41 – 0.30 (m, 2H).
	585	448	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.63 (s, 1H), 10.12 (s, 1H), 9.39 (s, 2H), 8.78 (s, 1H), 7.67 (dt, J = 7.2, 1.9 Hz, 2H), 6.95 (t, J = 2.0 Hz, 1H), 3.06 (s, 3H), 2.90 (s, 3H).

Example 586: *N*-(3-chloro-5-methanesulfonamidophenyl)-5-ethyl-4-(5-fluoropyrimidin-2-yl)thiophene-2-carboxamide



Step 1: A solution of methyl 4-(5-fluoropyrimidin-2-yl)thiophene-2-carboxylate (300 mg, 1.25 mmol) and NBS (1.33 g, 7.50 mmol) in DMF (10 mL) was stirred for 2 hours at 60 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (1:1) to afford methyl 5-bromo-4-(5-fluoropyrimidin-2-yl)thiophene-2-carboxylate (280 mg, 0.882 mmol) as a white solid. LCMS (ESI) $[M + H]^+$: 317

Step 2: A solution of methyl 5-bromo-4-(5-fluoropyrimidin-2-yl)thiophene-2-carboxylate (200 mg, 0.630 mmol), diethylzinc (1.88 mL, 1.88 mmol) and Pd(dppf)Cl₂ (51 mg, 0.0630 mmol) in THF (10 mL) was stirred for 2 hours at 70 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (1:1) to afford methyl 5-ethyl-4-(5-fluoropyrimidin-2-yl)thiophene-2-carboxylate (107 mg, 0.401 mmol) as a white solid. LCMS (ESI) $[M + H]^+$: 267


Step 3: To a stirred solution of methyl 5-ethyl-4-(5-fluoropyrimidin-2-yl)thiophene-2-carboxylate (107 mg, 0.401 mmol) in EtOH (5 mL) and H₂O (2 mL) was added LiOH (96 mg, 4.01 mmol) at 0 °C. Then the mixture was stirred for 2 hours at room temperature. The



mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 5-ethyl-4-(5-fluoropyrimidin-2-yl)thiophene-2-carboxylic acid (78.0 mg, 0.309 mmol) as a yellow solid. LCMS (ESI) $[M + H]^+$: 253

- 5 Step 4: To a stirred solution of 5-ethyl-4-(5-fluoropyrimidin-2-yl)thiophene-2-carboxylic acid (78 mg, 0.309 mmol), TCFH (173 mg, 0.618 mmol) and NMI (74 mg, 0.926 mmol) in ACN (5 mL) was added *N*-(3-amino-5-chlorophenyl)methanesulfonamide (68 mg, 0.309 mmol) at room temperature. Then the mixture was stirred for 1 hour at room temperature. The mixture was concentrated under reduced pressure. The residue was purified by reversed-
- 10 phase flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in water (0.1% formic acid), 0% to 50% gradient in 20 min; detector, UV 254 nm. This resulted in *N*-(3-chloro-5-methanesulfonamidophenyl)-5-ethyl-4-(5-fluoropyrimidin-2-yl)thiophene-2-carboxamide (43.7 mg, 0.0960 mmol) as a white solid. LCMS (ESI) $[M + H]^+$: 454.90. ^1H NMR (400 MHz, DMSO- d_6) δ 10.59 (s, 1H), 10.08 (s, 1H), 9.01 (s, 2H), 8.68 (s, 1H), 7.68 (dt, $J = 10.2, 2.0$ Hz, 2H), 6.95 (d, $J = 2.0$ Hz, 1H), 3.43
- 15 – 3.35 (m, 2H), 3.07 (s, 3H), 1.32 (t, $J = 7.4$ Hz, 3H).


Examples 587-589

The compounds listed in the following table were prepared using a procedure similar to that described for example 586:

Structure	Compound No.	MS (ESI) $[M+H]^+$	^1H NMR
	587	454.1	^1H NMR (300 MHz, DMSO- d_6) δ 10.45 (s, 1H), 10.08 (s, 1H), 8.57 (dq, $J = 4.6, 1.4$ Hz, 1H), 8.20 (d, $J = 1.9$ Hz, 1H), 7.91 (ddd, $J = 10.6, 8.4, 1.3$ Hz, 1H), 7.68 (t, $J = 1.9$ Hz, 1H), 7.63 (t, $J = 1.9$ Hz, 1H), 7.55 (dt, $J = 8.5, 4.3$ Hz, 1H), 6.96 (t, $J = 2.0$ Hz, 1H), 3.07 (s, 3H), 2.96 (q, $J = 7.5$ Hz, 2H), 1.25 (t, $J = 7.5$ Hz, 3H).

	588	471.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 10.07 (s, 1H), 8.66 (d, J = 2.4 Hz, 1H), 8.23 – 8.03 (m, 2H), 7.64 (dt, J = 21.4, 1.9 Hz, 2H), 6.95 (t, J = 1.9 Hz, 1H), 3.06 (s, 3H), 2.93 (q, J = 7.5 Hz, 2H), 1.24 (t, J = 7.5 Hz, 3H).
	589	454	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 10.07 (s, 1H), 8.68 (d, J = 3.0 Hz, 1H), 8.33 (s, 1H), 7.88 (td, J = 8.7, 3.0 Hz, 1H), 7.74 (dd, J = 8.8, 4.4 Hz, 1H), 7.65 (dt, J = 20.2, 1.9 Hz, 2H), 6.95 (t, J = 1.9 Hz, 1H), 3.14 (q, J = 7.4 Hz, 2H), 3.07 (s, 3H), 1.28 (t, J = 7.4 Hz, 3H).

Example 590: *N*-(3-chloro-5-methanesulfonamidophenyl)-4-[5-(4-cyclopropanecarbonylpiperazin-1-yl)pyridin-2-yl]-5-methylthiophene-2-carboxamide



Step 1: A solution of 2-bromo-5-chloropyridine (1.02 g, 5.30 mmol), Pd(dppf)Cl₂ (345 mg, 0.424 mmol), K₃PO₄ (3.37 g, 15.9 mmol) and methyl 5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxylate (2.99 g, 10.6 mmol) in 1,4-dioxane (15 mL) and H₂O (1 mL) was stirred for 2 hours at 80 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (1:1) to afford methyl 4-(5-chloropyridin-2-yl)-5-methylthiophene-2-carboxylate (800 mg, 2.98 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 268

Step 2: To a stirred solution of methyl 4-(5-chloropyridin-2-yl)-5-methylthiophene-2-carboxylate (800 mg, 2.98 mmol), Cs₂CO₃ (2.91 g, 8.94 mmol), RuPhos Pd G₃ (374 mg, 0.447 mmol), RuPhos (555 mg, 1.19 mmol) and *tert*-butyl piperazine-1-carboxylate (609 mg, 3.27 mmol) in PhMe (20 mL) at room temperature. Then the mixture was stirred for 5 hours at 100 °C under nitrogen atmosphere. The mixture was concentrated under reduced pressure. The residue was purified by reversed-phase flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in water (0.1% formic acid), 0% to 65% gradient in 20 min; detector, UV 254 nm. This resulted in *tert*-butyl 4-{6-[5-(methoxycarbonyl)-2-methylthiophen-3-yl]pyridin-3-yl}piperazine-1-carboxylate (830 mg, 1.98 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 418

Step 3: To a stirred solution of *tert*-butyl 4-{6-[5-(methoxycarbonyl)-2-methylthiophen-3-yl]pyridin-3-yl}piperazine-1-carboxylate (830 mg, 1.98 mmol) in DCM (20 mL) was added TFA (4 mL) dropwise at 0 °C. The reaction was stirred for 1 hour at room temperature. The resulting mixture was concentrated under vacuum. The mixture was basified to pH 9 with 1 M NaOH(aq). The mixture was concentrated under reduced pressure. The residue was purified by reversed-phase flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in water (0.1% formic acid), 0% to 40% gradient in 20 min; detector, UV 254 nm. This resulted in methyl 5-methyl-4-[5-(piperazin-1-yl)pyridin-2-yl]thiophene-2-carboxylate (500 mg, 1.57 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 318

Step 4: To a stirred solution of methyl 5-methyl-4-[5-(piperazin-1-yl)pyridin-2-yl]thiophene-2-carboxylate (500 mg, 1.57 mmol) and Et₃N (476 mg, 4.71 mmol) in DCM (15 mL) was added cyclopropanecarbonyl chloride (179 mg, 1.72 mmol) at 0 °C under nitrogen atmosphere. Then the mixture was stirred for 5 hours at room temperature. The mixture was

concentrated under reduced pressure. The residue was purified by reversed-phase flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in water (0.1% formic acid), 0% to 60% gradient in 20 min; detector, UV 254 nm. This resulted in methyl 4-[5-(4-cyclopropanecarbonylpiperazin-1-yl)pyridin-2-yl]-5-



- 5 methylthiophene-2-carboxylate (410 mg, 1.06 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 386



- Step 5: To a stirred solution of methyl 4-[5-(4-cyclopropanecarbonylpiperazin-1-yl)pyridin-2-yl]-5-methylthiophene-2-carboxylate (410 mg, 1.06 mmol) in EtOH (10 mL) and H₂O (2 mL) was added LiOH (253 mg, 10.6 mmol) at 0 °C. Then the mixture was stirred for 2 hours
- 10 at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 4-[5-(4-cyclopropanecarbonylpiperazin-1-yl)pyridin-2-yl]-5-methylthiophene-2-carboxylic acid (300 mg, 0.807 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 372

- 15 Step 6: To a stirred solution of 4-[5-(4-cyclopropanecarbonylpiperazin-1-yl)pyridin-2-yl]-5-methylthiophene-2-carboxylic acid (300 mg, 0.807 mmol), TCFH (450 mg, 1.61 mmol) and NMI (193 mg, 2.42 mmol) in ACN (10 mL) was added *N*-(3-amino-5-chlorophenyl)methanesulfonamide (195 mg, 0.887 mmol) at room temperature. Then the mixture was stirred for 1 hour at room temperature. The mixture was concentrated under
- 20 reduced pressure. The residue was purified by reversed-phase flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in water (0.1% formic acid), 0% to 50% gradient in 20 min; detector, UV 254 nm. This resulted in *N*-(3-chloro-5-methanesulfonamidophenyl)-4-[5-(4-cyclopropanecarbonylpiperazin-1-yl)pyridin-2-yl]-5-methylthiophene-2-carboxamide (102.9 mg, 0.179 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 574.15. ¹H NMR (300 MHz, DMSO-*d*₆) δ 10.43 (s, 1H), 10.07 (s, 1H), 8.42 (d, *J* = 2.8 Hz, 1H), 8.34 (s, 1H), 7.67 (d, *J* = 9.3 Hz, 2H), 7.60 – 7.45 (m, 2H), 6.95 (d, *J* = 2.0 Hz, 1H), 3.87 (s, 2H), 3.66 (s, 2H), 3.27 (s, 4H), 3.07 (s, 3H), 2.68 (s, 3H), 2.07 (d, *J* = 7.2 Hz, 1H), 0.76 (t, *J* = 6.3 Hz, 4H).

30 Examples 591-594


The compounds listed in the following table were prepared using a procedure similar to that described for example 590:

Structure	Compound No.	MS (ESI) [M+H] ⁺	¹ H NMR
	591	573.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.05 (s, 1H), 9.79 (s, 1H), 8.32 (d, J = 2.9 Hz, 1H), 7.72 (t, J = 1.8 Hz, 1H), 7.66 – 7.50 (m, 3H), 7.43 (dd, J = 8.9, 2.9 Hz, 1H), 7.04 (d, J = 2.0 Hz, 1H), 6.88 (t, J = 1.9 Hz, 1H), 3.93 (s, 3H), 3.72 (t, J = 5.1 Hz, 4H), 3.23 (t, J = 5.3 Hz, 4H), 3.06 (s, 3H), 1.23 (s, 9H).
	592	585.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.80 (s, 1H), 8.33 (d, J = 2.9 Hz, 1H), 7.73 (d, J = 1.9 Hz, 1H), 7.67 – 7.52 (m, 3H), 7.45 (dd, J = 8.9, 3.0 Hz, 1H), 7.04 (d, J = 2.0 Hz, 1H), 6.89 (t, J = 2.0 Hz, 1H), 3.93 (s, 3H), 3.65 (dt, J = 17.7, 5.4 Hz, 4H), 3.24 (dt, J = 19.8, 5.1 Hz, 4H), 3.06 (s, 4H), 1.78 (q, J = 7.6, 6.0 Hz, 2H), 1.75 – 1.57 (m, 2H), 1.64 (m, 2H).

			– 1.46 (m, 4H).
	593	571	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.79 (s, 1H), 8.32 (d, J = 2.9 Hz, 1H), 7.72 (d, J = 2.0 Hz, 1H), 7.66 – 7.58 (m, 2H), 7.55 (d, J = 8.8 Hz, 1H), 7.43 (dd, J = 8.9, 3.0 Hz, 1H), 7.03 (d, J = 2.0 Hz, 1H), 6.89 (d, J = 2.1 Hz, 1H), 3.93 (s, 3H), 3.64 – 3.57 (m, 2H), 3.48 (t, J = 5.2 Hz, 2H), 3.42 (q, J = 8.5 Hz, 1H), 3.21 (q, J = 5.1 Hz, 4H), 3.06 (s, 3H), 2.26 – 2.02 (m, 4H), 1.91 (dt, J = 10.9, 8.6 Hz, 1H), 1.79 – 1.72 (m, 1H).
	594	557.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.80 (s, 1H), 8.34 (d, J = 2.9 Hz, 1H), 7.73 (t, J = 1.8 Hz, 1H), 7.66 – 7.59 (m, 2H), 7.56 (d, J = 8.8 Hz, 1H), 7.45 (dd, J = 8.9, 3.0 Hz, 1H), 7.04 (d, J = 2.0 Hz, 1H), 6.89 (t, J = 2.0 Hz, 1H),

			3.93 (s, 3H), 3.85 (s, 2H), 3.64 (s, 2H), 3.31 (s, 2H), 3.23 (s, 2H), 3.06 (s, 3H), 2.05 (tt, J = 7.7, 4.6 Hz, 1H), 0.75 (tt, J = 7.9, 2.9 Hz, 4H).
--	--	--	---

Example 595: *N*-(3-chloro-5-methanesulfonamidophenyl)-5-[5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl]-1-methyl-1*H*-pyrrole-3-carboxamide



- Step 1: To a mixture of 5-bromo-2-iodopyridine (5.1 g, 17.9 mmol), methyl 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrrole-3-carboxylate (5.67 g, 21.4 mmol), Pd(dppf)Cl₂ (1.45 g, 1.78 mmol) and K₃PO₄ (11.3 g, 53.6 mmol) was added 1,4-dioxane (60 ml) and H₂O (6 ml). The resulting mixture was stirred for 2 hours at 70 °C under N₂ atmosphere. The reaction mixture was extracted with ethyl acetate. The organic phase was concentrated under reduced pressure. The residue was purified by flash chromatography on silica gel eluting with 15% of ethyl acetate in petroleum ether to afford methyl 5-(5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl)-1-methyl-1*H*-pyrrole-3-carboxylate (5.10 g, 96.5%) as a white solid.
- LCMS [M+H]⁺: 295



Step 2: To a mixture of methyl 5-(5-bromopyridin-2-yl)-1-methyl-1*H*-pyrrole-3-carboxylate (3.1 g, 10.5 mmol), 3,3-difluoroazetidine hydrochloride (4.08 g, 31.5 mmol), Cs₂CO₃ (10.28 g, 31.6 mmol) and PEPPSI-IHeptCl (510 mg, 525 μmol) was added dioxane (30 mL). The resulting mixture was stirred for 1 hour at 120 °C under N₂ atmosphere. The reaction mixture
5 was extracted with ethyl acetate. The organic phase was concentrated under reduced pressure. The residue was purified by flash chromatography on silica gel eluting with 30% of ethyl acetate in petroleum ether to afford methyl 5-[5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl]-1-methyl-1*H*-pyrrole-3-carboxylate (2.20 g, 68.3%) as a yellow solid. LCMS [M+H]⁺: 308




Step 3: To a stirred solution of methyl 5-[5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl]-1-methyl-1*H*-pyrrole-3-carboxylate (2.2 g, 7.15 mmol) in 20 mL of EtOH and 20 mL of water was
10 added LiOH (1.49 g, 35.7 mmol). The resulting mixture was stirred for 1 hour at 70 °C. Then it was concentrated, and the pH value of the residue solution was adjusted to 3 with 1M of HCl (aq). The product was precipitated from the solution. The mixture was filtered and the filter cake was washed with water (20 mL) to afford 5-[5-(3,3-difluoroazetidin-1-yl)pyridin-
15 2-yl]-1-methyl-1*H*-pyrrole-3-carboxylic acid (1.75 g, 83.7%) as a yellow solid. LCMS [M+H]⁺: 294




Step 4: To a mixture of 5-[5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl]-1-methyl-1*H*-pyrrole-3-carboxylic acid (1.6 g, 5.45 mmol) and *N*-(3-amino-5-chlorophenyl)methanesulfonamide (1.44 g, 6.54 mmol) in ACN (16 mL) was added TCFH (2.28 g, 8.17 mmol) and NMI (1.33
20 g, 16.3 mmol) at room temperature for two hours. Then it was concentrated, the residue was purified by reverse phase flash chromatography eluting with 60% of acetonitrile in water (0.1%FA) to afford *N*-(3-chloro-5-methanesulfonamidophenyl)-5-[5-(3,3-difluoroazetidin-1-yl)pyridin-2-yl]-1-methyl-1*H*-pyrrole-3-carboxamide (1.57 g, 58.1%) as a yellow solid.
LCMS [M+H]⁺: 496. ¹H NMR (300 MHz, DMSO-*d*₆) δ 10.00 (s, 1H), 9.79 (s, 1H), 7.98 (dd, *J* = 2.9, 0.7 Hz, 1H), 7.72 (t, *J* = 1.8 Hz, 1H), 7.67 – 7.58 (m, 2H), 7.61 – 7.52 (m, 1H), 7.09 (dd, *J* = 8.7, 2.9 Hz, 1H), 7.03 (d, *J* = 2.0 Hz, 1H), 6.89 (t, *J* = 1.9 Hz, 1H), 4.41 (d, *J* = 12.3 Hz, 3H), 4.35 (s, 1H), 3.92 (s, 3H), 3.06 (s, 3H).

Examples 596-603:

The compounds listed in the following table were prepared using a procedure similar to that described for example 595:


Structure	Compound No.	MS (ESI) [M+H] ⁺	¹ H NMR
	596	547.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.79 (s, 1H), 8.32 (d, J = 2.9 Hz, 1H), 7.72 (t, J = 1.8 Hz, 1H), 7.67 – 7.50 (m, 3H), 7.43 (dd, J = 8.9, 2.9 Hz, 1H), 7.04 (d, J = 2.0 Hz, 1H), 6.89 (t, J = 1.9 Hz, 1H), 3.93 (s, 3H), 3.64 (s, 3H), 3.58 – 3.49 (m, 4H), 3.24 (d, J = 10.5 Hz, 4H), 3.06 (s, 3H).
	597	496.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.90 (d, J = 14.2 Hz, 2H), 8.23 (s, 2H), 7.72 (t, J = 1.9 Hz, 1H), 7.65 (t, J = 1.9 Hz, 1H), 7.63 (d, J = 2.2 Hz, 1H), 7.50 (d, J = 2.1 Hz, 1H), 6.88 (t, J = 2.0 Hz, 1H), 4.46 (t, J = 12.3 Hz, 4H), 4.03 (s, 3H), 3.05 (s, 3H).

	598	479.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.04 (s, 1H), 9.88 (s, 1H), 8.14 (s, 1H), 7.73 (t, J = 1.9 Hz, 1H), 7.66 (t, J = 1.9 Hz, 1H), 7.61 (d, J = 2.1 Hz, 1H), 7.46 (d, J = 2.1 Hz, 1H), 6.88 (t, J = 1.9 Hz, 1H), 5.75 – 5.30 (m, 1H), 4.41 – 4.23 (m, 2H), 4.16 – 3.93 (m, 5H), 3.05 (s, 3H).
	599	514	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.83 (s, 1H), 7.89 (s, 1H), 7.67 (dd, J = 16.7, 14.7 Hz, 3H), 7.12 (dd, J = 12.4, 2.3 Hz, 1H), 7.02 (s, 1H), 6.89 (s, 1H), 4.44 (t, J = 12.3 Hz, 4H), 3.83 (s, 3H), 3.06 (s, 3H).
	600	538.1	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.74 (s, 1H), 8.29 (d, J = 2.9 Hz, 1H), 7.68 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.5 Hz, 2H), 7.38 (d, J = 2.8 Hz, 1H), 6.88 (t, J = 2.0 Hz, 1H), 6.76 (d, J = 1.9 Hz, 1H), 3.62 (s, 3H), 3.46 (t, J = 5.8 Hz, 4H), 3.05 (s, 3H), 2.33 (s, 3H).

			2.09 (td, J = 13.9, 6.7 Hz, 4H).
	601	578.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.29 (s, 1H), 10.04 (s, 1H), 8.23 (d, J = 2.8 Hz, 1H), 7.94 (s, 1H), 7.66 (t, J = 1.9 Hz, 1H), 7.60 (t, J = 1.9 Hz, 1H), 7.38 (s, 1H), 6.94 (t, J = 2.0 Hz, 1H), 3.64 (s, 3H), 3.54 (t, J = 5.2 Hz, 4H), 3.27 (d, J = 10.7 Hz, 4H), 3.06 (s, 3H), 2.32 (s, 3H), 2.19 (s, 3H).
	602	555	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.29 (s, 1H), 10.05 (s, 1H), 8.28 (d, J = 2.8 Hz, 1H), 7.94 (s, 1H), 7.66 (t, J = 1.9 Hz, 1H), 7.61 (t, J = 1.9 Hz, 1H), 7.39 (d, J = 2.7 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 3.45 (t, J = 5.9 Hz, 4H), 3.06 (s, 3H), 2.33 (s, 3H), 2.19 (s, 3H), 2.08 (tt, J = 13.7, 5.8 Hz, 4H).
	603	545	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.03 (s, 1H), 9.89 (s, 1H), 8.65 – 8.60 (m, 1H), 7.94 – 7.82 (m, 2H), 7.76 – 7.69 (m,

			2H), 7.63 (s, 1H), 7.31 (d, J = 2.0 Hz, 1H), 6.91 (t, J = 1.9 Hz, 1H), 4.02 (s, 3H), 3.07 (s, 3H), 2.91 (s, 3H), 1.46 (s, 6H).
--	--	--	--

Example 604: N-(3-chloro-5-methanesulfonamidophenyl)-5-{5,5'-difluoro-[3,3'-bipyridin]-2-yl}-1-methylpyrrole-3-carboxamide



Step 1: A solution of 2-bromo-3-chloro-5-fluoropyridine (300 mg, 2 mmol, 1 equiv), methyl
 5 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrole-3-carboxylate (410 mg, 2
 mmol, 1 equiv), Pd(dppf)Cl₂ (80 mg, 0.3 mmol, 0.1 equiv) and K₂CO₃ (400 mg, 8 mmol, 3
 equiv) in Dioxane (10 mL) was stirred for 1 hour at 80 °C under nitrogen atmosphere. The
 reaction mixture was concentrated under vacuum. The residue was purified by flash
 chromatography on silica gel eluting with EA/PE(32%) to afford methyl 5-(3-chloro-5-
 10 fluoropyridin-2-yl)-1-methylpyrrole-3-carboxylate (200 mg, 60% yield) as a white solid.
 LCMS (ESI) [M + H]⁺: 269

Step 2: A solution of methyl 5-(3-chloro-5-fluoropyridin-2-yl)-1-methylpyrrole-3-carboxylate
 (200 mg, 2 mmol, 1 equiv), 5-fluoropyridin-3-ylboronic acid (150 mg, 2 mmol, 1 equiv),
 Pd(dppf)Cl₂ (50 mg, 0.3 mmol, 0.1 equiv) and K₂CO₃ (200 mg, 8 mmol, 3 equiv) in Dioxane
 15 (10 mL) was stirred for 1h at 80°C under nitrogen atmosphere. The reaction mixture was
 concentrated under vacuum. The residue was purified by flash chromatography on silica gel
 eluting with EA/PE(22%) to afford methyl 5-{5,5'-difluoro-[3,3'-bipyridin]-2-yl}-1-


methylpyrrole-3-carboxylate (120 mg, 55% yield) as a white solid. LCMS (ESI) $[M + H]^+$: 330




Step 3: A solution of ethyl methyl 5-{5,5'-difluoro-[3,3'-bipyridin]-2-yl}-1-methylpyrrole-3-carboxylate (120 mg, 0.6 mmol, 1 equiv) and LiOH (86.8 mg, 3.624 mmol, 3.00 equiv) in
 5 ethyl alcohol (2 mL)/H₂O (1 mL) was stirred at RT for four hours. The mixture was acidified with HCl (0.1M). The solvent was concentrated under vacuum. The residue was purified by flash chromatography on C18 column gel eluting with ACN/water (0.1% FA) (33%) to afford methyl 5-{5,5'-difluoro-[3,3'-bipyridin]-2-yl}-1-methylpyrrole-3-carboxylic acid(100 mg, 66% yield) as a white solid. LCMS (ESI) $[M + H]^+$: 316

10 Step 4: A solution of 5-{5,5'-difluoro-[3,3'-bipyridin]-2-yl}-1-methylpyrrole-3-carboxylic acid (100 mg, 0.2 mmol, 1 equiv), *N*-(3-amino-5-chlorophenyl)methanesulfonamide (80 mg, 0.2 mmol, 1 equiv) and NMI (10 mg, 0.04 mmol, 0.1 equiv), TCFH (12 mg, 0.05 mmol, 0.1 equiv) in ACN (5 mL) was stirred for 1 hour at 30°C under nitrogen atmosphere. The reaction mixture was concentrated under vacuum. The residue (120mg) was purified by Prep-HPLC to
 15 afford *N*-(3-chloro-5-methanesulfonamidophenyl)-5-{5,5'-difluoro-[3,3'-bipyridin]-2-yl}-1-methylpyrrole-3-carboxamide (18.1 mg, 20% yield) as a white solid. LCMS (ESI) $[M + H]^+$: 518. ¹H NMR (300 MHz, DMSO-*d*₆) δ 9.67 (s, 1H), 8.79 (d, *J* = 2.8 Hz, 1H), 8.56 (d, *J* = 2.8 Hz, 1H), 8.32 (t, *J* = 1.8 Hz, 1H), 8.07 (dd, *J* = 9.3, 2.8 Hz, 1H), 7.77 (ddd, *J* = 9.9, 2.8, 1.8 Hz, 1H), 7.66 – 7.56 (m, 2H), 7.52 (t, *J* = 1.9 Hz, 1H), 6.86 (t, *J* = 1.9 Hz, 1H), 6.24 (d, *J* =
 20 1.9 Hz, 1H), 3.64 (s, 3H), 3.02 (s, 3H).


Examples 605-608:

The compounds listed in the following table were prepared using a procedure similar to that described for example 604:

Structure	Compound No.	MS (ESI) $[M+H]^+$	¹ H NMR
	605	546.9	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 10.35 (s, 1H), 10.10 (s, 1H), 8.76 (d, <i>J</i> = 2.8 Hz, 1H), 8.02 – 7.86 (m, 2H), 7.66 – 7.55 (m, 3H), 6.93 (t, <i>J</i> = 1.9 Hz, 1H), 6.41 (d, <i>J</i> = 2.0 Hz, 1H), 5.93 (dd, <i>J</i> = 7.0, 2.0 Hz, 1H), 3.39 (s, 3H), 3.05 (s, 3H), 2.13 (s, 3H).

	606	547.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.34 (s, 2H), 8.67 (d, J = 2.8 Hz, 1H), 8.00 – 7.86 (m, 3H), 7.61 (dt, J = 17.0, 1.9 Hz, 2H), 7.11 – 6.87 (m, 2H), 6.25 (d, J = 9.4 Hz, 1H), 3.44 (s, 3H), 3.05 (s, 3H), 2.08 (d, J = 5.0 Hz, 3H).
	607	534.95	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.33 (s, 1H), 10.06 (s, 1H), 8.80 (d, J = 2.8 Hz, 1H), 8.56 (d, J = 2.7 Hz, 1H), 8.29 (t, J = 1.8 Hz, 1H), 8.13 (dd, J = 9.4, 2.8 Hz, 1H), 7.87 (s, 1H), 7.74 (ddd, J = 9.9, 2.8, 1.8 Hz, 1H), 7.61 (dt, J = 15.5, 1.9 Hz, 2H), 6.94 (t, J = 1.9 Hz, 1H), 3.06 (s, 3H), 1.99 (s, 3H).
	608	531	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.25 (s, 1H), 8.78 (d, J = 2.8 Hz, 1H), 8.43 (dd, J = 4.8, 1.7 Hz, 1H), 7.97 (dd, J = 9.3, 2.8 Hz, 1H), 7.67 (s, 1H), 7.61 – 7.51 (m, 3H), 7.24 (dd, J = 7.7, 4.8 Hz, 1H), 6.93 (t, J = 2.0 Hz, 1H), 3.05 (s, 3H), 2.21 (s, 3H), 2.09 (s, 3H).

Example 609: *N*-(3-chloro-5-methanesulfonamidophenyl)-5-methyl-4-{5-[(1-methylazetidin-3-yl)oxy]pyrimidin-2-yl}thiophene-2-carboxamide



Step 1: To a stirred solution of 2-bromopyrimidin-5-ol (1.5 g, 8.57 mmol), methyl 5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxylate (2.41 g, 8.57 mmol), Pd(dppf)Cl₂ (699 mg, 0.857 mmol) and K₃PO₄ (5.44 g, 25.7 mmol) in 1,4-dioxane (20.00 mL) was added H₂O (3.00 mL). Then the mixture was stirred for 12 hours at 80 °C under N₂. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford methyl 4-(5-hydroxypyrimidin-2-yl)-5-methylthiophene-2-carboxylate (1.80 g, 7.19 mmol, 84.1% yield) as a white solid. LCMS (ESI) [M + H]⁺: 251.2

Step 2: To a stirred solution of methyl 4-(5-hydroxypyrimidin-2-yl)-5-methylthiophene-2-carboxylate (520 mg, 2.07 mmol), *tert*-butyl 3-iodoazetidine-1-carboxylate (586 mg, 2.07 mmol) and K₂CO₃ (855 mg, 6.20 mmol) in DMF (5 mL). Then the mixture was stirred for 6

hours at 80 °C under N₂. The mixture was concentrated and purified by flash chromatography on silica gel eluting with EA/PE (0-100%) to afford *tert*-butyl 3-({2-[5-(methoxycarbonyl)-2-methylthiophen-3-yl]pyrimidin-5-yl}oxy) azetidine-1-carboxylate (470 mg, 1.15 mmol, 56% yield) as a yellow solid. LCMS (ESI) [M + H]⁺: 406.4

- 5 Step 3: To a stirred solution of *tert*-butyl 3-({2-[5-(methoxycarbonyl)-2-methylthiophen-3-yl]pyrimidin-5-yl}oxy) azetidine-1-carboxylate (470 mg, 1.15 mmol) in DCM (5 ml) was added trifluoroacetic acid (655 mg, 5.75 mmol) dropwise. The reaction was stirred for 3 hours at room temperature under nitrogen atmosphere. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic
- 10 system) to afford methyl 4-[5-(340yridine340-3-yloxy)pyrimidin-2-yl]-5-methylthiophene-2-carboxylate (230 mg, 0.753 mmol, 65.5% yield) as a white solid. LCMS (ESI) [M + H]⁺: 306.3

- Step 4: A mixture of methyl 4-[5-(340yridine340-3-yloxy)pyrimidin-2-yl]-5-methylthiophene-2-carboxylate (230 mg, 0.753 mmol), formaldehyde (45.0 mg, 1.50 mmol)
- 15 and AcOH (135 mg, 2.25 mmol) in MeOH (3.00 mL) was stirred for 15 min at room temperature. Then sodium triacetoxyborohydride (477 mg, 2.25 mmol) was added into the mixture at room temperature. The resulting mixture was stirred for additional 1 hour at room temperature. The mixture was concentrated and purified by reverse phase flash
- 20 chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford methyl 5-methyl-4-{5-[(1-methylazetidin-3-yl)oxy]pyrimidin-2-yl}thiophene-2-carboxylate (200 mg, 0.626 mmol, 83.3% yield) as a white solid. LCMS (ESI) [M + H]⁺: 320.1

- Step 5: To a stirred solution of methyl 5-methyl-4-{5-[(1-methylazetidin-3-yl)oxy]pyrimidin-2-yl}thiophene-2-carboxylate (200 mg, 626 μmol) and LiOH (10 mg, 2.5 mmol) in THF
- 25 (10.00 mL) was added H₂O (2.00 mL) dropwise. Then the mixture was stirred for 2 hours at 60 °C. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 5-methyl-4-{5-[(1-methylazetidin-3-yl)oxy] pyrimidin-2-yl}thiophene-2-carboxylic acid (150 mg, 491 μmol, 78.5%yield) as a white solid. LCMS (ESI) [M + H]⁺: 306

- Step 6: To a stirred solution of 5-methyl-4-{5-[(1-methylazetidin-3-yl)oxy]pyrimidin-2-yl}thiophene-2-carboxylic acid (150 mg, 0.491 mmol), TCFH (206 mg, 0.736 mmol) and
- 30 NMI (120 mg, 1.47 mmol) in ACN (5.00 mL) was added *N*-(3-amino-5-chlorophenyl)methanesulfonamide (108 mg, 0.491 mmol). Then the mixture was stirred for 4

hours at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford *N*-(3-chloro-5-methanesulfonamidophenyl)-5-methyl-4-{5-[(1-methylazetidin-3-yl)oxy]pyrimidin-2-yl}thiophene-2-carboxamide (108.5 mg, 0.211 mmol, 42.9% yield) as an off-white solid.


- 5 LCMS (ESI) [M + H]⁺: 508.25. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.56 (s, 1H), 10.07 (s, 1H), 8.64 (s, 1H), 8.56 (s, 2H), 7.68 (dt, *J* = 8.0, 1.9 Hz, 2H), 6.94 (t, *J* = 1.9 Hz, 1H), 5.01 (p, *J* = 5.5 Hz, 1H), 3.78 (td, *J* = 6.1, 1.9 Hz, 2H), 3.12 – 2.99 (m, 5H), 2.83 (s, 3H), 2.31 (s, 3H).

Examples 610-611:

- 10 The compounds listed in the following table were prepared using a procedure similar to that described for example 609:

Structure	Compound No.	MS (ESI) [M+H] ⁺	¹ H NMR
	610	588.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 10.06 (s, 1H), 8.51 (s, 2H), 7.85 – 7.61 (m, 4H), 6.92 (s, 1H), 5.76 (q, <i>J</i> = 9.1 Hz, 2H), 4.96 (p, <i>J</i> = 5.7 Hz, 1H), 3.72 (t, <i>J</i> = 6.8 Hz, 2H), 3.06 (s, 3H), 3.00 (t, <i>J</i> = 6.4 Hz, 2H), 2.33 (q, <i>J</i> = 6.5 Hz, 1H), 0.88 (d, <i>J</i> = 6.1 Hz, 6H).
	611	508	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 10.00 (s, 1H), 9.85 (s, 1H), 8.18 (dd, <i>J</i> = 2.6, 1.0 Hz, 1H), 7.70 (d, <i>J</i> = 1.9 Hz, 2H), 7.63 (d, <i>J</i> = 2.0 Hz, 1H), 7.47 (dd, <i>J</i> = 12.3, 2.5 Hz, 1H), 7.08 (dd, <i>J</i> = 3.4, 1.9 Hz, 1H), 6.89 (t, <i>J</i> = 2.0 Hz, 1H), 4.92 (p, <i>J</i> = 5.5 Hz, 1H), 3.84 (s, 3H), 3.77 (td, <i>J</i> = 6.1, 1.9 Hz, 2H), 3.05 (s, 3H), 3.04 – 2.98 (m, 2H), 2.30 (s, 3H).

Example 612: *N*-(3-chloro-5-methanesulfonamidophenyl)-1-methyl-5-{5-[(1-methylazetidin-3-yl)oxy]pyridin-2-yl}pyrrole-3-carboxamide



Step 1: A solution of 6-bromopyridin-3-ol (320 mg, 1.839 mmol, 1 equiv), methyl 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrole-3-carboxylate (488 mg, 1.839 mmol, 1 equiv), Pd(dppf)Cl₂ (135 mg, 0.184 mmol, 0.1 equiv) and K₂CO₃ (508 mg, 3.678 mmol, 2 equiv) in dioxane (6 mL) and H₂O (2 mL) was stirred for 1.5 hours at 80 °C under nitrogen atmosphere. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (1:1) to afford methyl 5-(5-hydroxypyridin-2-yl)-1-methylpyrrole-3-carboxylate (280 mg, 66% yield) as a white solid. LCMS (ESI) [M + H]⁺: 233

Step 2: A solution of methyl 5-(5-hydroxypyridin-2-yl)-1-methylpyrrole-3-carboxylate (280 mg, 1.206 mmol, 1 equiv), *tert*-butyl 3-iodoazetidine-1-carboxylate (410 mg, 1.447 mmol, 1.2 equiv) and Cs₂CO₃ (786 mg, 2.412 mmol, 2 equiv) in DMSO (5 mL) was stirred for 1.5 hours at 80 °C under air atmosphere. The reaction was diluted with water. The resulting mixture was extracted with EtOAc (3 x 50 mL). The combined organic layers were washed with brine, dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (1:1) to afford methyl 5-(5-{[1-(*tert*-butoxycarbonyl)azetidin-3-yl]oxy}pyridin-2-yl)-1-methylpyrrole-3-carboxylate (270 mg, 58% yield) as an off-white solid. LCMS (ESI) [M + H]⁺: 388

Step 3: A solution of methyl 5-(5-{[1-(*tert*-butoxycarbonyl)azetidin-3-yl]oxy}pyridin-2-yl)-1-methylpyrrole-3-carboxylate (270 mg, 0.697 mmol, 1 equiv) in TFA (2 mL) and DCM (2 mL) was stirred for 1 hour at room temperature under air atmosphere. The mixture was

concentrated under reduced pressure. The residue was purified by reverse flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in water (10 mmol/L NH_4HCO_3), 0% to 100% gradient in 18 min. This resulted in methyl 5-[5-(azetidin-3-yloxy)pyridin-2-yl]-1-methylpyrrole-3-carboxylate (150 mg, 75% yield) as a white solid. LCMS (ESI) $[\text{M} + \text{H}]^+$: 288

Step 4: A solution of methyl 5-[5-(azetidin-3-yloxy)pyridin-2-yl]-1-methylpyrrole-3-carboxylate (130 mg, 0.452 mmol, 1 equiv) and formaldehyde (27 mg, 0.904 mmol, 2 equiv) in methanol (2 mL) was stirred for 1 hour at 50 °C under air atmosphere. To the above mixture was add $\text{NaBH}(\text{AcO})_3$ (144 mg, 0.678 mmol, 1.5 equiv) at room temperature. The mixture was stirred for 30 min at room temperature under air atmosphere. The mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (5:1) to afford methyl 1-methyl-5-{5-[(1-methylazetidin-3-yl)oxy]pyridin-2-yl}pyrrole-3-carboxylate (70 mg, 51% yield) as a white solid. LCMS (ESI) $[\text{M} + \text{H}]^+$: 302



Step 5: A solution of methyl 1-methyl-5-{5-[(1-methylazetidin-3-yl)oxy]pyridin-2-yl}pyrrole-3-carboxylate (70 mg, 0.232 mmol, 1 equiv) and LiOH (11 mg, 0.464 mmol, 2 equiv) in EtOH (2 mL) and H_2O (1 mL) was stirred for 1 hour at 50°C under air atmosphere. The residue was acidified to pH 5 with 1 M HCl (aq.). The mixture was concentrated under reduced pressure. The residue was purified by reverse flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in water (0.1% formic acid), 0% to 40% gradient in 18 min. This resulted in 1-methyl-5-{5-[(1-methylazetidin-3-yl)oxy]pyridin-2-yl}pyrrole-3-carboxylic acid (45 mg, 67% yield) as a white solid. LCMS (ESI) $[\text{M} + \text{H}]^+$: 288


Step 6: A solution of 1-methyl-5-{5-[(1-methylazetidin-3-yl)oxy]pyridin-2-yl}pyrrole-3-carboxylic acid (45 mg, 0.157 mmol, 1 equiv), *N*-(3-amino-5-chlorophenyl)methanesulfonamide (41 mg, 0.188 mmol, 1.2 equiv), TCFH (66 mg, 0.235 mmol, 1.5 equiv) and NMI (39 mg, 0.471 mmol, 3 equiv) in ACN (3 mL) was stirred for 2 hours at room temperature under air atmosphere. The mixture was concentrated under reduced pressure. The residue was purified by reverse flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in water (10 mmol/L NH_4HCO_3), 0% to 50% gradient in 18 min. This resulted in *N*-(3-chloro-5-methanesulfonamidophenyl)-1-methyl-5-{5-[(1-methylazetidin-3-yl)oxy]pyridin-2-

yl}pyrrole-3-carboxamide (23.1 mg, 30% yield) as a white solid. LCMS (ESI) $[M + H]^+$: 490.00. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 9.96 (s, 1H), 9.77 (s, 1H), 8.21 (d, $J = 2.8$ Hz, 1H), 7.71 – 7.54 (m, 4H), 7.35 (dd, $J = 8.8, 3.0$ Hz, 1H), 7.07 (d, $J = 2.0$ Hz, 1H), 6.86 (t, $J = 2.0$ Hz, 1H), 4.87 (p, $J = 5.6$ Hz, 1H), 3.92 (s, 3H), 3.76 (dd, $J = 8.3, 6.0$ Hz, 2H), 3.12 – 2.89 (m, 5H), 2.30 (s, 3H).


Examples 613-615:

The compounds listed in the following table were prepared using a procedure similar to that described for example 612:

Structure	Compound No.	MS (ESI) $[M+H]^+$	^1H NMR
	613	559.1	^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.07 (s, 1H), 8.50 (s, 2H), 7.84 – 7.79 (m, 1H), 7.73 (t, $J = 1.8$ Hz, 1H), 7.69 (p, $J = 1.6$ Hz, 2H), 6.93 (q, $J = 1.8$ Hz, 1H), 5.76 (q, $J = 9.0$ Hz, 2H), 5.01 (p, $J = 5.5$ Hz, 1H), 3.87 – 3.78 (m, 2H), 3.13 (qt, $J = 5.0, 3.5, 3.0$ Hz, 2H), 3.08 (d, $J = 1.2$ Hz, 3H), 2.34 (t, $J = 1.5$ Hz, 3H).
	614	558.35	^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 10.00 (s, 1H), 9.86 (s, 1H), 8.23 – 8.10 (m, 1H), 7.71 (t, $J = 1.9$ Hz, 2H), 7.64 (t, $J = 1.9$ Hz, 1H), 7.48 (dd, $J = 12.3, 2.5$ Hz, 1H), 7.08 (dd, $J = 3.4, 1.9$ Hz, 1H), 6.90 (t, $J = 1.9$ Hz, 1H), 5.99 (tt, $J =$

			55.7, 4.1 Hz, 1H), 5.00 (p, J = 5.4 Hz, 1H), 3.95 – 3.79 (m, 5H), 3.26 (dd, J = 8.8, 4.9 Hz, 2H), 3.06 (s, 3H), 2.91 (td, J = 16.1, 4.1 Hz, 2H).
	615	576.15	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.85 (s, 1H), 8.19 (d, J = 2.3 Hz, 1H), 7.73 – 7.67 (m, 2H), 7.63 (t, J = 2.0 Hz, 1H), 7.48 (dd, J = 12.3, 2.6 Hz, 1H), 7.08 (dd, J = 3.4, 2.0 Hz, 1H), 6.89 (t, J = 2.0 Hz, 1H), 5.03 (p, J = 5.5 Hz, 1H), 3.93 (dd, J = 8.3, 6.0 Hz, 2H), 3.84 (s, 3H), 3.40 (d, J = 4.6 Hz, 2H), 3.34 (d, J = 4.6 Hz, 2H), 3.05 (s, 3H).

Example 616: *N*-(3-chloro-5-methanesulfonamidophenyl)-5-methyl-4-[5-(methylamino)pyrimidin-2-yl]thiophene-2-carboxamide




Step 1: To a stirred solution of methyl 5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxylate (820 mg, 2.90 mmol), 2-bromopyrimidin-5-amine (504 mg, 2.90 mmol), Pd(dppf)Cl₂ (236 mg, 0.29 mmol) and K₂CO₃ (1.20 g, 8.70 mmol) in 1,4-dioxane (10 mL) was added H₂O (2 mL). Then the mixture was stirred for 12 hours at 80 °C under N₂. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford methyl 4-(5-aminopyrimidin-2-yl)-5-methylthiophene-2-carboxylate (700 mg, 2.80 mmol, 96% yield) as a white solid. LCMS (ESI) [M + H]⁺: 250.2

Step 2: To a stirred solution of acetic acid (201 mg, 3.36 mmol), TCFH (944 mg, 3.36 mmol) and NMI (551 mg, 6.72 mmol) in ACN (5.00 mL) was added methyl 4-(5-aminopyrimidin-2-yl)-5-methylthiophene-2-carboxylate (560 mg, 2.24 mmol). Then the mixture was stirred for 4 hours at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford methyl 4-(5-acetamidopyrimidin-2-yl)-5-methylthiophene-2-carboxylate (380 mg, 1.30 mmol, 58.2% yield) as a white solid. LCMS (ESI) [M + H]⁺: 292.3

Step 3: To a stirred mixture of methyl 4-(5-acetamidopyrimidin-2-yl)-5-methylthiophene-2-carboxylate (380 mg, 1.30 mmol) and Cs_2CO_3 (1.27 g, 3.90 mmol) in DMF (5 ml) was added CH_3I (276 mg, 1.95 mmol) dropwise at 80 °C under nitrogen atmosphere. Then the mixture was stirred for 4 hours at 80 °C. The mixture was concentrated and purified by reverse phase
5 flash chromatography eluting with ACN/ H_2O (5-95%, acidic system) to afford methyl 5-methyl-4-[5-(*N*-methylacetamido)pyrimidin-2-yl]thiophene-2-carboxylate (270 mg, 0.884 mmol, 68.1% yield) as a white solid. LCMS (ESI) $[\text{M} + \text{H}]^+$: 306.3

Step 4: To a stirred mixture of methyl 5-methyl-4-[5-(*N*-methylacetamido)pyrimidin-2-yl]thiophene-2-carboxylate (120 mg, 0.392 mmol) and *N*-(3-amino-5-chlorophenyl)methanesulfonamide (86.5 mg, 0.392 mmol) in toluene (3 mL) was added
10 $\text{Al}(\text{Me})_3$ (37.1 mg, 0.509 mmol) dropwise at 80 °C under nitrogen atmosphere. Then the mixture was stirred for 4 hours at 80 °C. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/ H_2O (5-95%, acidic system) to afford residue. The residue was purified by Prep-HPLC with the following conditions (Column:
15 Sunfire prep C18 column, 30*150 mm, 5 μm ; Mobile Phase A: Water(0.1%FA), Mobile Phase B: ACN; Flow rate: 60 mL/min; Gradient: 45% B to 62% B in 7 min; Wave Length: 254/220 nm; RT1(min): 5.43) to afford *N*-(3-chloro-5-methanesulfonamidophenyl)-5-methyl-4-[5-(methylamino)pyrimidin-2-yl]thiophene-2-carboxamide (5.5 mg, 0.0118 mmol, 3 %
20 yield) as a white solid. LCMS (ESI) $[\text{M} + \text{H}]^+$: 452.05. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.51 (s, 1H), 10.12 (s, 1H), 8.57 (s, 1H), 8.21 (s, 2H), 7.67 (dt, $J = 6.8, 1.9$ Hz, 2H), 6.93 (t, $J = 2.0$ Hz, 1H), 6.25 (q, $J = 5.0$ Hz, 1H), 3.05 (s, 3H), 2.79 (t, $J = 2.6$ Hz, 6H).

Example 617: *N*-(3-chloro-5-methanesulfonamidophenyl)-5-{5-[1-(2,2-difluoroethyl)piperidin-4-yl]pyridin-2-yl}-1-methylpyrrole-3-carboxamide



Step 1: A solution of 5-bromo-2-iodopyridine (1.5 g, 5.284 mmol, 1 equiv), methyl 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrole-3-carboxylate (1.40 g, 5.284 mmol, 1 equiv), Pd(dppf)Cl_2 (0.43 g, 0.528 mmol, 0.1 equiv) and K_3PO_4 (3.36 g, 15.852 mmol, 3 equiv) in 1,4-dioxane (30 mL) and H_2O (6 mL) was stirred for 1 hour at 60 °C under nitrogen atmosphere. The mixture was allowed to cool down to room temperature. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (5:1) to afford methyl 5-(5-bromopyridin-2-yl)-

1-methylpyrrole-3-carboxylate (1 g, 64% yield) as a white solid. LCMS (ESI) [M + H]⁺: 250.00.

Step 2: A solution of methyl 5-(5-bromopyridin-2-yl)-1-methylpyrrole-3-carboxylate (500 mg, 1.694 mmol, 1 equiv), *tert*-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydro-2*H*-pyridine-1-carboxylate (524 mg, 1.694 mmol, 1 equiv), Pd(dppf)Cl₂ (138 mg, 0.169 mmol, 0.1 equiv) and K₃PO₄ (1079 mg, 5.082 mmol, 3 equiv) in 1,4-dioxane (10 mL) and H₂O (2 mL) was stirred for 1 hour at 80 °C under nitrogen atmosphere. The mixture was allowed to cool down to room temperature. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (5:1) to afford *tert*-butyl 6-[4-(methoxycarbonyl)-1-methylpyrrol-2-yl]-3',6'-dihydro-2'*H*-[3,4'-bipyridine]-1'-carboxylate (350 mg, 52% yield) as a white solid. LCMS (ESI) [M + H]⁺: 398.20.

Step 3: A solution of *tert*-butyl 6-[4-(methoxycarbonyl)-1-methylpyrrol-2-yl]-3',6'-dihydro-2'*H*-[3,4'-bipyridine]-1'-carboxylate (330 mg, 0.830 mmol, 1 equiv) and Pd/C (44 mg, 0.415 mmol, 0.5 equiv) in THF (5 mL) was stirred for 1 day at room temperature under hydrogen atmosphere. The precipitated solids were collected by filtration and washed with EtOH (3x10 mL). The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (1:1) to afford *tert*-butyl 4-{6-[4-(methoxycarbonyl)-1-methylpyrrol-2-yl]pyridin-3-yl}piperidine-1-carboxylate (263 mg, 79% yield) as a white solid. LCMS (ESI) [M + H]⁺: 400.22.

Step 4: A solution of *tert*-butyl 4-{6-[4-(methoxycarbonyl)-1-methylpyrrol-2-yl]pyridin-3-yl}piperidine-1-carboxylate (263 mg, 0.658 mmol, 1 equiv) in DCM (3 mL) and TFA (1 mL) was stirred for 1 hour at room temperature. The resulting mixture was concentrated under reduced pressure to afford methyl 1-methyl-5-[5-(piperidin-4-yl)pyridin-2-yl]pyrrole-3-carboxylate (190 mg, 96% yield) as a white solid. LCMS (ESI) [M + H]⁺: 300.16.


Step 5: A solution of methyl 1-methyl-5-[5-(piperidin-4-yl)pyridin-2-yl]pyrrole-3-carboxylate (170 mg, 0.568 mmol, 1 equiv), 2-bromo-1,1-difluoroethane (123 mg, 0.852 mmol, 1.5 equiv) and K₂CO₃ (235 mg, 1.704 mmol, 3 equiv) in DMF (5 mL) was stirred for 1h at 80°C. The mixture was allowed to cool down to room temperature. The resulting mixture was diluted with water and extracted with EtOAc (3 x89 mL). The combined organic layers were washed with water (3x46 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel


column chromatography, eluted with PE / EA (1:1) to afford methyl 5-{5-[1-(2,2-difluoroethyl)piperidin-4-yl]pyridin-2-yl}-1-methylpyrrole-3-carboxylate (133 mg, 64% yield) as a white solid. LCMS (ESI) $[M + H]^+$: 364.18.


Step 6: To a stirred solution of methyl 5-{5-[1-(2,2-difluoroethyl)piperidin-2-yl]-1-methylpyrrole-3-carboxylate (113 mg, 0.311 mmol, 1 equiv) and *N*-(3-amino-5-chlorophenyl)methanesulfonamide (102.92 mg, 0.467 mmol, 1.5 equiv) in toluene (2 mL) was added $Al(Me)_3$ (0.5 mL) dropwise at 0 °C. The resulting mixture was stirred for 2 hours at 80 °C. The mixture was allowed to cool down to room temperature. The reaction was quenched by the addition of water (2 mL) at 0 °C. The resulting mixture was concentrated under reduced pressure. The residue (200 mg) was purified by Prep-HPLC with the following conditions (Column: YMC-Actus Triart C18 ExRS, 30*150 mm, 5 μ m; Mobile Phase A: Water (10 mmol/L NH_4HCO_3), Mobile Phase B: ACN; Flow rate: 60 mL/min; Gradient: 35% B to 65% B in 7 min, 65% B; Wave Length: 254/220 nm; RT1(min): 7.33; Number Of Runs: 0) to afford *N*-(3-chloro-5-methanesulfonamidophenyl)-5-{5-[1-(2,2-difluoroethyl)piperidin-4-yl]pyridin-2-yl}-1-methylpyrrole-3-carboxamide (106.6 mg, 62% yield) as a white solid. LCMS (ESI) $[M + H]^+$: 551.95. ^1H NMR (300 MHz, $DMSO-d_6$) δ 10.02 (s, 1H), 9.84 (s, 1H), 8.50 (d, $J = 2.3$ Hz, 1H), 7.80 – 7.69 (m, 2H), 7.69 – 7.56 (m, 3H), 7.17 (d, $J = 2.0$ Hz, 1H), 6.90 (t, $J = 2.0$ Hz, 1H), 6.17 (tt, $J = 55.9, 4.4$ Hz, 1H), 3.98 (s, 3H), 3.07 (s, 5H), 2.77 (td, $J = 15.7, 4.3$ Hz, 2H), 2.64 – 2.52 (m, 1H), 2.43 – 2.18 (m, 2H), 1.83 – 1.52 (m, 4H).

20 Examples 618-619:

The compounds listed in the following table were prepared using a procedure similar to that described for example 617:

Structure	Compound No.	MS (ESI) $[M+H]^+$	^1H NMR
	618	605	^1H NMR (400 MHz, $DMSO-d_6$) δ 10.42 (s, 1H), 10.07 (s, 1H), 8.53 – 8.39 (m, 1H), 8.19 (d, $J = 1.7$ Hz, 1H), 7.84 (dd, $J = 11.8, 1.8$ Hz, 1H), 7.64 (dt, $J = 16.6, 1.9$ Hz, 2H), 6.94 (t, $J = 2.0$ Hz, 1H), 3.22 (q, J

			= 10.3 Hz, 2H), 3.06 (s, 5H), 2.76 – 2.62 (m, 1H), 2.53 (d, J = 1.6 Hz, 3H), 2.47 (d, J = 8.0 Hz, 2H), 1.87 – 1.71 (m, 4H).
	619	570	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.83 (s, 1H), 8.50 (d, J = 2.3 Hz, 1H), 7.79 – 7.69 (m, 2H), 7.69 – 7.60 (m, 3H), 7.17 (d, J = 2.0 Hz, 1H), 6.89 (t, J = 2.0 Hz, 1H), 3.98 (s, 3H), 3.22 (q, J = 10.2 Hz, 2H), 3.06 (s, 5H), 2.58 (dq, J = 10.7, 5.6, 4.4 Hz, 1H), 2.46 (dd, J = 11.2, 3.3 Hz, 2H), 1.87 – 1.56 (m, 4H).

Example 620:

Step 1: A mixture of 3-(chloromethyl)-5-fluoropyridine (180 mg, 1.23 mmol), K₂CO₃ (508 mg, 3.68 mmol) and 2-bromopyridin-3-ol (234 mg, 1.35 mmol) in DMF (5 mL) was stirred for 2 hours at 80 °C. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, formic acid) to afford 2-bromo-3-[(5-fluoropyridin-3-yl) methoxy]pyridine (200 mg, 0.706 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 283




Step 2: A mixture of 2-bromo-3-[(5-fluoropyridin-3-yl)methoxy]pyridine (200 mg, 0.706 mmol), Pd(dppf)Cl₂ (51.6 mg, 0.0706 mmol), K₂CO₃ (291 mg, 2.11 mmol) and methyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxylate (245 mg, 0.917 mmol) in 1,4-dioxane (10 mL) was stirred for 2 hours at 80 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (0-100%) to afford methyl 4-{3-[(5-fluoropyridin-3-yl)methoxy]pyridin-2-yl}thiophene-2-carboxylate (114 mg, 0.331 mmol) as a yellow solid. LCMS (ESI) [M + H]⁺: 345





Step 3: To a stirred solution of methyl 4-{3-[(5-fluoropyridin-3-yl)methoxy]pyridin-2-yl}thiophene-2-carboxylate (140 mg, 0.406 mmol) in EtOH (3 mL) and H₂O (3 mL) was added LiOH (97.2 mg, 4.06 mmol). Then the mixture was stirred for 2 hours at room temperature. The mixture was acidified to pH 5 with citric acid. The precipitated solids were collected by filtration and washed with water. The resulting solid was dried under vacuum to afford 4-{3-[(5-fluoropyridin-3-yl) methoxy]pyridin-2-yl}thiophene-2-carboxylic acid (110 mg, 0.333 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 331

Step 4: To a stirred solution of 4-{3-[(5-fluoropyridin-3-yl)methoxy]pyridin-2-yl}thiophene-2-carboxylic acid (110 mg, 0.333 mmol), TCFH (1.96 g, 6.99 mmol) and NMI (163 mg, 1.99 mmol) in ACN (5 mL) was added *N*-(3-amino-5-chlorophenyl)methanesulfonamide (80.7 mg, 0.366 mmol). Then the mixture was stirred for 1 hour at room temperature. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, formic acid system) to afford *N*-(3-chloro-5-methanesulfonamidophenyl)-4-{3-[(5-fluoropyridin-3-yl)methoxy]pyridin-2-yl}thiophene-2-carboxamide (66.5 mg, 0.124 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 533.00. ¹H NMR (300 MHz, DMSO-*d*₆) δ 10.63 (s, 1H), 10.07 (s, 1H), 8.81 (d, *J* = 1.3 Hz, 1H), 8.68 – 8.56 (m, 2H), 8.45 (d, *J* = 1.2 Hz, 1H), 8.31 (dd, *J* = 4.6, 1.2 Hz, 1H), 7.93 (dt, *J* = 9.5, 2.3 Hz, 1H), 7.78 – 7.66 (m, 3H), 7.40 (dd, *J* = 8.4, 4.6 Hz, 1H), 6.96 (t, *J* = 1.9 Hz, 1H), 5.44 (s, 2H), 3.08 (s, 3H).


Examples 621-627:

The compounds listed in the following table were prepared using a procedure similar to that described for example 620:

Structure	Compound No.	MS (ESI) [M+H] ⁺	¹ H NMR
	621	547	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.34 (s, 1H), 10.11 (s, 1H), 8.59 – 8.47 (m, 2H), 8.32 (dd, J = 4.7, 1.2 Hz, 1H), 8.20 (s, 1H), 7.73 (d, J = 8.5 Hz, 2H), 7.66 – 7.57 (m, 2H), 7.46 (dd, J = 8.4, 4.7 Hz, 1H), 6.94 (t, J = 2.0 Hz, 1H), 5.31 (s, 2H), 3.06 (s, 3H), 2.41 (s, 3H).
	622	567	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.36 (s, 1H), 10.12 (s, 1H), 8.36 (d, J = 2.3 Hz, 1H), 8.13 (s, 1H), 7.89 (dd, J = 10.9, 2.3 Hz, 1H), 7.85 (d, J = 3.2 Hz, 1H), 7.77 (d, J = 3.2 Hz, 1H), 7.63 (d, J = 12.0 Hz, 2H), 6.94 (t, J = 1.9 Hz, 1H), 5.61 (s, 2H), 3.17 (q, J = 7.2 Hz, 2H), 2.41 (s, 3H), 1.20 (t, J = 7.3 Hz, 3H).
	623	562	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.30 (s, 1H), 10.11 (s, 1H), 9.15 (s, 1H), 8.84 (s, 2H), 8.36 (d, J = 2.3 Hz, 1H), 8.14 (s, 1H), 7.86 (dd, J = 10.8, 2.4 Hz, 1H), 7.61 (dt, J = 3.4, 1.8 Hz, 2H), 6.94 (t, J = 1.9 Hz, 1H), 5.33 (s, 2H), 3.16 (q, J = 7.3 Hz, 2H), 2.37 (s, 3H), 1.20 (t, J = 7.3 Hz, 3H).

	624	544	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.99 (s, 1H), 9.70 (s, 1H), 8.53 (d, J = 2.8 Hz, 1H), 8.46 (s, 1H), 7.91 (d, J = 7.0 Hz, 1H), 7.74 – 7.57 (m, 4H), 6.88 (t, J = 2.0 Hz, 1H), 6.52 (d, J = 1.9 Hz, 1H), 6.35 (d, J = 7.0 Hz, 1H), 5.28 – 5.11 (m, 2H), 3.36 (s, 3H), 3.06 (s, 3H), 2.08 (s, 3H).
	625	535	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.80 (s, 1H), 8.35 (d, J = 2.4 Hz, 1H), 7.88 (dd, J = 10.8, 2.4 Hz, 1H), 7.74 – 7.61 (m, 3H), 7.01 (d, J = 2.0 Hz, 1H), 6.89 (t, J = 2.0 Hz, 1H), 5.71 (s, 2H), 4.03 (s, 3H), 3.73 (s, 3H), 3.06 (s, 3H).
	626	577	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.33 (s, 1H), 10.04 (s, 1H), 8.54 – 8.45 (m, 2H), 8.31 (dd, J = 4.6, 1.2 Hz, 1H), 8.19 (s, 1H), 7.72 (dt, J = 9.1, 1.9 Hz, 2H), 7.63 (dt, J = 4.9, 1.9 Hz, 2H), 7.45 (dd, J = 8.4, 4.7 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 5.31 (s, 2H), 4.94 (s, 1H), 3.75 (t, J = 6.6 Hz, 2H), 3.30 (d, J = 6.6 Hz, 2H), 2.40 (s, 3H).
	627	548	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.93 (s, 2H), 8.83 (s, 1H), 8.66 (s, 1H), 7.83 (d, J = 1.9 Hz, 1H), 7.72 – 7.49 (m, 3H), 7.31 – 7.15 (m, 3H), 6.90 (t, J = 1.9 Hz, 1H), 5.44 (s, 2H), 4.00 (s, 3H), 3.05 (s, 3H).

Examples 628 and 629: *N*-(3-chloro-5-methane sulfonamidophenyl)-5-{3-[1-[3-fluoro-5-(trifluoromethyl)phenyl]ethoxy]pyridin-2-yl}-1-methyl-1*H*-pyrrole-3-carboxamide



- Step 1: To a stirred solution of 1-[3-fluoro-5-(trifluoromethyl)phenyl]ethan-1-one (2 g, 9.70 mmol) in THF (30 mL) was added NaBH₄ (1.10 g, 29.1 mmol) at room temperature. Then the mixture was stirred for 1 hour at 0 °C. The mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (1:2) to afford 1-[3-fluoro-5-(trifluoromethyl)phenyl]ethan-1-ol (1.55 g, 7.44 mmol) as an off yellow solid. LCMS (ESI) [M + H]⁺: 209.05
- Step 2: To a stirred solution of 1-[3-fluoro-5-(trifluoromethyl)phenyl]ethan-1-ol (1.55 g, 7.44 mmol) in DCM (5 mL) was added sulfurooyl dichloride (2.65 g, 22.3 mmol) dropwise at 0 °C under nitrogen atmosphere. Then the mixture was stirred for 1 hour at 0 °C. The mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (1:1) to afford 1-(1-chloroethyl)-3-fluoro-5-(trifluoromethyl)benzene (1.30 g, 5.73 mmol) as an off yellow solid. LCMS (ESI) [M + H]⁺: 227.06

Step 3: To a stirred solution of 1-(1-chloroethyl)-3-fluoro-5-(trifluoromethyl)benzene (420 mg, 1.85 mmol) and K₂CO₃ (765 mg, 5.55 mmol) in DMSO (10 mL) was added methyl 5-(3-hydroxypyridin-2-yl)-1-methyl-1*H*-pyrrole-3-carboxylate (429 mg, 1.85 mmol) at room temperature. Then the mixture was stirred for 1 hour at 60 °C. The mixture was diluted with

water. The resulting mixture was extracted with EA (3 x 100 mL). The combined organic layers were washed with brine, dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by reversed-phase flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in water (0.1% formic acid), 5% to 40% gradient in 20 min; detector, UV 254 nm to afford methyl 5-(3-{1-[3-fluoro-5-(trifluoromethyl)phenyl]ethoxy}pyridin-2-yl)-1-methyl-1*H*-pyrrole-3-carboxylate (375 mg, 887 μmol) as an off white solid. LCMS (ESI) [M + H]⁺: 423.13

Step 4: To a stirred solution of methyl 5-(3-{1-[3-fluoro-5-(trifluoromethyl)phenyl]ethoxy}pyridin-2-yl)-1-methyl-1*H*-pyrrole-3-carboxylate (375 mg, 887 μmol) and *N*-(3-amino-5-chlorophenyl)methanesulfonamide (195 mg, 887 μmol) in PhMe (10 mL) was added AlMe₃ (191 mg, 2.6 mmol) at room temperature. Then the mixture was stirred for 1 hour at 80 °C. The mixture was concentrated under reduced pressure. The residue was purified by reversed-phase flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in water (0.1% formic acid), 5% to 80% gradient in 20 min; detector, UV 254 nm to afford mixture products. The racemic mixture was purified Chiral-HPLC (Column: CHIRAL ART Amylose-SA, 3*25 cm, 5 μm; Mobile Phase A: hexanes with 10mM NH₃-MeOH, Mobile Phase B: EtOH; Flow rate: 40 mL/min; Gradient: 30% B to 30% B in 17.4 min; Sample Solvent: EtOH--HPLC; Injection Volume: 0.3 mL) to afford the separated enantiomers of *N*-(3-chloro-5-methane sulfonamidophenyl)-5-{3-[1-[3-fluoro-5-(trifluoromethyl)phenyl]ethoxy]pyridin-2-yl}-1-methyl-1*H*-pyrrole-3-carboxamide.

Example 628: *N*-(3-chloro-5-methane sulfonamidophenyl)-5-{3-[1-[3-fluoro-5-(trifluoromethyl)phenyl]ethoxy]pyridin-2-yl}-1-methyl-1*H*-pyrrole-3-carboxamide, enantiomer 1




126.7 mg, 206 μmol, white solid. Single enantiomer, arbitrarily assigned. Elution time: 11.6 min. LCMS (ESI) [M + H]⁺: 611.10. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.06 (s, 1H), 9.80 (s, 1H), 8.21 (d, *J* = 2.8 Hz, 1H), 7.77 (dd, *J* = 9.7, 2.9 Hz, 2H), 7.23 – 7.76 (m, 7H), 7.83 (t, *J* = 2.0 Hz, 1H), 5.80 (q, *J* = 1.9 Hz, 1H), 3.84 (s, 3H), 3.06 (s, 3H), 1.61 (s, 3H).


Example 629: *N*-(3-chloro-5-methanesulfonamidophenyl)-5-{3-[1-[3-fluoro-5-(trifluoromethyl)phenyl]ethoxy]pyridin-2-yl}-1-methyl-1*H*-pyrrole-3-carboxamide, enantiomer 2

122.0 mg, 199 μ mol, white solid. Single enantiomer, arbitrarily assigned. Elution time: 14.6 min. LCMS (ESI) $[M + H]^+$: 611.11. ^1H NMR (400 MHz, DMSO- d_6) δ 10.06 (s, 1H), 9.80 (s, 1H), 8.21 (d, $J = 2.8$ Hz, 1H), 7.77 (dd, $J = 9.7, 2.9$ Hz, 2H), 7.23 – 7.76 (m, 7H), 7.83 (t, $J = 2.0$ Hz, 1H), 5.80 (q, $J = 1.9$ Hz, 1H), 3.84 (s, 3H), 3.06 (s, 3H), 1.61 (s, 3H).


5 Examples 630-633:

The compounds listed in the following table were prepared using a procedure similar to that described for example 628:

Structure	Compound No.	MS (ESI) $[M+H]^+$	^1H NMR
 Enantiomer 1	630	561	^1H NMR (300 MHz, DMSO- d_6) δ 10.36 (s, 1H), 10.08 (s, 1H), 8.56 (d, $J = 1.9$ Hz, 1H), 8.35 (dd, $J = 4.9, 1.1$ Hz, 1H), 8.28 (dd, $J = 4.6, 1.2$ Hz, 1H), 8.14 (s, 1H), 7.69 (t, $J = 1.9$ Hz, 1H), 7.63 (t, $J = 1.9$ Hz, 1H), 7.54 (dd, $J = 8.6, 1.3$ Hz, 1H), 7.35 (dd, $J = 8.4, 4.6$ Hz, 1H), 7.27 (dd, $J = 6.3, 4.9$ Hz, 1H), 6.95 (t, $J = 1.9$ Hz, 1H), 5.80 (q, $J = 6.3$ Hz, 1H), 3.08 (s, 3H), 2.45 (s, 3H), 1.58 (d, $J = 6.4$ Hz, 3H).
 Enantiomer 2	631	561	^1H NMR (300 MHz, DMSO- d_6) δ 10.36 (s, 1H), 10.08 (s, 1H), 8.56 (d, $J = 1.9$ Hz, 1H), 8.35 (dd, $J = 4.9, 1.1$ Hz, 1H), 8.28 (dd, $J = 4.7, 1.3$ Hz, 1H), 8.14 (s, 1H), 7.69 (t, $J = 1.9$ Hz, 1H), 7.63 (t, $J = 1.9$ Hz, 1H), 7.54 (dd, $J = 8.5, 1.3$ Hz, 1H), 7.35 (dd, $J = 8.4, 4.6$ Hz, 1H), 7.27 (dd, $J = 6.3, 4.9$ Hz, 1H), 6.95 (t, $J = 1.9$ Hz, 1H), 5.80 (q, $J = 6.3$ Hz, 1H), 3.08 (s, 3H), 2.45 (s, 3H), 1.58 (d, $J = 6.4$ Hz, 3H).
	632	578.1	^1H NMR (400 MHz, DMSO- d_6) δ 10.36 (s, 1H), 10.06 (s, 1H), 8.25 (dd, $J = 4.7, 1.3$ Hz, 1H), 8.15 (s, 1H), 7.65 (dt, $J = 15.5, 1.9$ Hz, 2H), 7.52 (dd, $J = 8.6, 1.4$ Hz, 1H), 7.34 (dd, $J = 8.4, 4.6$ Hz, 1H), 7.10 (tt, $J = 9.3, 2.4$ Hz, 1H), 7.02 – 6.92 (m, 3H), 5.60 (q, $J = 6.3$ Hz, 1H), 3.06 (s, 3H), 2.44 (s, 3H), 1.52 (d, $J = 6.3$ Hz, 3H).

Enantiomer 1			
	633	578.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.36 (s, 1H), 10.06 (s, 1H), 8.25 (dd, J = 4.7, 1.3 Hz, 1H), 8.15 (s, 1H), 7.65 (dt, J = 15.5, 1.9 Hz, 2H), 7.52 (dd, J = 8.6, 1.4 Hz, 1H), 7.34 (dd, J = 8.4, 4.6 Hz, 1H), 7.10 (tt, J = 9.3, 2.4 Hz, 1H), 7.02 – 6.92 (m, 3H), 5.60 (q, J = 6.3 Hz, 1H), 3.06 (s, 3H), 2.44 (s, 3H), 1.52 (d, J = 6.3 Hz, 3H).

Example 634: *N*-(3-chloro-5-methanesulfonamidophenyl)-5-{3-fluoro-5-[(1-isopropylazetidin-3-yl)oxy]pyridin-2-yl}-1-methylpyrrole-3-carboxamide



Step 1: A solution of *tert*-butyl 3-iodoazetidine-1-carboxylate (1.5 g, 5.29 mmol, 1 equiv), K₂CO₃ (2.20 g, 15.89 mmol, 3 equiv) and 6-chloro-5-fluoropyridin-3-ol (0.94 g, 6.35 mmol, 1.2 equiv) in DMF (15 mL) was stirred for 1 hour at 80 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by
5 reversed-phase flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in Water (0.1% FA), 10% to 50% gradient in 10 min; detector, UV 254 nm. This resulted in *tert*-butyl 3-[(6-chloro-5-fluoropyridin-3-yl)oxy]azetidine-1-carboxylate (1.3 g, 81% yield) as a white solid. LCMS (ESI) [M+H]⁺: 303.00.

Step 2: A solution of *tert*-butyl 3-[(6-chloro-5-fluoropyridin-3-yl)oxy]azetidine-1-carboxylate
10 (1.3 g, 4.29 mmol, 1 equiv), methyl 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrole-3-carboxylate (1.37 g, 5.15 mmol, 1.2 equiv), Pd(dppf)Cl₂ (0.31 g, 0.42 mmol, 0.1 equiv) and K₂CO₃ (1.78 g, 12.88 mmol, 3 equiv) in dioxane (10 mL) and H₂O (1 mL) was stirred for 1 hour at 80 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column
15 chromatography, eluted with PE / EA (1:1) to afford methyl 5-(5-{[1-(*tert*-butoxycarbonyl)azetidin-3-yl]oxy}-3-fluoropyridin-2-yl)-1-methylpyrrole-3-carboxylate (1.08 g, 62% yield) as a white solid. LCMS (ESI) [M+H]⁺: 406.00.

Step 3: A solution of methyl 5-(5-{[1-(*tert*-butoxycarbonyl)azetidin-3-yl]oxy}-3-fluoropyridin-2-yl)-1-methylpyrrole-3-carboxylate (1.08 g, 2.66 mmol, 1 equiv) and TFA (5
20 mL) in DCM (15 mL) was stirred for 1 hour at room temperature under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. This resulted in methyl 5-[5-(azetidin-3-yloxy)-3-fluoropyridin-2-yl]-1-methylpyrrole-3-carboxylate (930 mg, 114% yield) as a white solid. LCMS (ESI) [M+H]⁺: 306.00.

Step 4: A solution of methyl 5-[5-(azetidin-3-yloxy)-3-fluoropyridin-2-yl]-1-methylpyrrole-3-carboxylate (200 mg, 0.65 mmol, 1 equiv) and acetone (2 mL) in MeOH (2 mL) was stirred
25 for overnight at 40 °C under nitrogen atmosphere. To the above mixture was added sodium triacetoxyborohydride (277 mg, 1.31 mmol, 2 equiv) in portions over 2 minutes at 0 °C. The resulting mixture was stirred for additional 10 min at room temperature. The residue was purified by reversed-phase flash chromatography with the following conditions: column, C18
30 silica gel; mobile phase, MeCN in Water (0.1% FA), 10% to 50% gradient in 10 min; detector, UV 254 nm. This resulted in methyl 5-{3-fluoro-5-[(1-isopropylazetidin-3-yl)oxy]pyridin-2-yl}-1-methylpyrrole-3-carboxylate (135 mg, 59% yield) as a white solid. LCMS (ESI) [M-H]⁺: 348.00.


- Step 5: A solution of methyl 5-{3-fluoro-5-[(1-isopropylazetidin-3-yl)oxy]pyridin-2-yl}-1-methylpyrrole-3-carboxylate (135 mg, 0.39 mmol, 1 equiv) and LiOH (74 mg, 3.11 mmol, 8 equiv) in EtOH (2 mL) and H₂O (0.4 mL) was stirred for 1 hour at room temperature under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by reversed-phase flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in Water (0.1% FA), 10% to 50% gradient in 10 min; detector, UV 254 nm. This resulted in 5-{3-fluoro-5-[(1-isopropylazetidin-3-yl)oxy]pyridin-2-yl}-1-methylpyrrole-3-carboxylic acid (108 mg, 83% yield) as a white solid. LCMS (ESI) [M-H]⁺: 334.00.
- Step 6: A solution of 5-{3-fluoro-5-[(1-isopropylazetidin-3-yl)oxy]pyridin-2-yl}-1-methylpyrrole-3-carboxylic acid (108 mg, 0.32 mmol, 1 equiv), *N*-(3-amino-5-chlorophenyl)methanesulfonamide (85 mg, 0.38 mmol, 1.2 equiv), TCFH (136 mg, 0.48 mmol, 1.5 equiv) and NMI (79 mg, 0.97 mmol, 3 equiv) in ACN (2 mL) was stirred for 1 hour at room temperature under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by reversed-phase flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeCN in Water (0.1% FA), 10% to 50% gradient in 10 min; detector, UV 254 nm. This resulted in *N*-(3-chloro-5-methanesulfonamidophenyl)-5-{3-fluoro-5-[(1-isopropylazetidin-3-yl)oxy]pyridin-2-yl}-1-methylpyrrole-3-carboxamide (103.5 mg, 60%) as a white solid. LCMS (ESI) [M-H]⁺: 536.20. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.00 (s, 1H), 9.86 (s, 1H), 8.19 (dt, *J* = 2.4, 1.2 Hz, 1H), 7.71 (dt, *J* = 4.0, 1.5 Hz, 2H), 7.64 (q, *J* = 1.6 Hz, 1H), 7.47 (dt, *J* = 12.2, 2.4 Hz, 1H), 7.08 (dd, *J* = 3.5, 1.8 Hz, 1H), 6.90 (dt, *J* = 2.2, 1.2 Hz, 1H), 4.89 (p, *J* = 5.5 Hz, 1H), 3.84 (s, 3H), 3.76 – 3.68 (m, 2H), 3.06 (s, 3H), 3.03 – 2.95 (m, 2H), 2.39 – 2.28 (m, 1H), 0.88 (dd, *J* = 6.1, 0.9 Hz, 6H).

Example 635:

The compounds listed in the following table were prepared using a procedure similar to that described for example 634:

Structure	Compound No.	MS (ESI) [M+H] ⁺	¹ H NMR
	635	525	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.39 (s, 1H), 8.20 (d, J = 2.4 Hz, 1H), 7.98 (d, J = 1.7 Hz, 1H), 7.64 (t, J = 2.0 Hz, 1H), 7.54 (t, J = 1.9 Hz, 1H), 7.35 (dd, J = 11.3, 2.5 Hz, 1H), 7.01 (t, J = 2.0 Hz, 1H), 5.14 (tt, J = 6.2, 4.4 Hz, 1H), 4.36 – 4.27 (m, 2H), 3.89 (dd, J = 11.1, 4.3 Hz, 2H), 3.03 (s, 3H), 2.78 (s, 3H), 2.53 (d, J = 1.2 Hz, 3H).

5 **Example 636:** 4-[5-(azetidin-3-yloxy)pyrimidin-2-yl]-N-(3-chloro-5-methanesulfonamidophenyl)-5-methylthiophene-2-carboxamide



Step 1: To a stirred solution of 2-bromopyrimidin-5-ol (1 g, 5.71 mmol), K₂CO₃ (3.62 g, 17.1 mmol) and *tert*-butyl 3-iodoazetidine-1-carboxylate (1.61 g, 5.71 mmol) in DMF (30 mL).

- 10 Then the mixture was stirred for 5 hours at 80 °C. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (5-95%, acidic system) to afford

tert-butyl 3-[(2-bromopyrimidin-5-yl)oxy]azetidine-1-carboxylate (1.20 g, 3.63 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 330

Step 2: A mixture of *tert*-butyl 3-[(2-bromopyrimidin-5-yl)oxy]azetidine-1-carboxylate (520 mg, 1.57 mmol), methyl 5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxylate (485 mg, 1.72 mmol), K₃PO₄ (0.9997 g, 4.71 mmol), Pd(dppf)Cl₂ (128 mg, 0.157 mmol) and H₂O (2 mL) in dioxane (20 mL) was stirred for 16 hours at 80 °C under nitrogen atmosphere. The residue was purified by silica gel column chromatography, eluted with PE / EA (35%) to afford *tert*-butyl 3-[(2-[5-(methoxycarbonyl)-2-methylthiophen-3-yl]pyrimidin-5-yl)oxy]azetidine-1-carboxylate (740 mg, 1.82 mmol) as a yellow solid.
LCMS (ESI) [M + H]⁺: 406

Step 3: To a stirred solution of *tert*-butyl 3-[(2-[5-(methoxycarbonyl)-2-methylthiophen-3-yl]pyrimidin-5-yl)oxy]azetidine-1-carboxylate (420 mg, 1.03 mmol) in EtOH (15 mL) and H₂O (15 mL) was added LiOH (49.3 mg, 2.06 mmol). Then the mixture was stirred for 4 hours at room temperature. The mixture was acidified to pH 5 with hydrochloric acid. The mixture was concentrated and purified by reverse phase flash chromatography eluting with ACN/H₂O (50%, acidic system) to afford 4-[5-[(1-[(*tert*-butoxy)carbonyl]azetidin-3-yl)oxy]pyrimidin-2-yl]-5-methylthiophene-2-carboxylic acid (366 mg, 0.935 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 426



Step 4: To a stirred solution of 4-[5-[(1-[(*tert*-butoxy)carbonyl]azetidin-3-yl)oxy]pyrimidin-2-yl]-5-methylthiophene-2-carboxylic acid (366 mg, 0.935 mmol), TCFH (785 mg, 2.80 mmol) in NMI (459 mg, 5.60 mmol) and ACN (20 mL) was added *N*-(3-amino-5-chlorophenyl)methanesulfonamide (247 mg, 1.12 mmol). Then the mixture was stirred for 2 hours at room temperature. The mixture was concentrated and purified by flash chromatography eluted with PE / EA (50%) to afford *tert*-butyl 3-[(2-{5-[(3-chloro-5-methanesulfonamidophenyl)carbamoyl]-2-methylthiophen-3-yl}pyrimidin-5-yl)oxy]azetidine-1-carboxylate (240 mg, 0.403 mmol) as a white solid. LCMS (ESI) [M + H]⁺: 594

Step 5: A mixture of *tert*-butyl 3-[(2-{5-[(3-chloro-5-methanesulfonamidophenyl)carbamoyl]-2-methylthiophen-3-yl}pyrimidin-5-yl)oxy]azetidine-1-carboxylate (240 mg, 0.403 mmol) and TFA (367 mg, 3.22 mmol) in DCM (20 mL) was stirred for 2 hours at ambient temperature and was concentrated under vacuum to afford 4-[5-(azetidin-3-yloxy)pyrimidin-2-yl]-*N*-(3-chloro-5-


methanesulfonamidophenyl)-5-methylthiophene-2-carboxamide (63.2 mg, 0.127 mmol) as a light yellow oil. LCMS (ESI) $[M + H]^+$: 494. ^1H NMR (300 MHz, Methanol- d_4) δ 8.45 (d, J = 9.2 Hz, 3H), 7.64 (t, J = 2.0 Hz, 1H), 7.55 (d, J = 1.9 Hz, 1H), 7.01 (t, J = 2.0 Hz, 1H), 5.28 – 5.18 (m, 1H), 4.04 (dd, J = 10.3, 6.7 Hz, 2H), 3.77 (dd, J = 10.3, 5.5 Hz, 2H), 3.03 (s, 3H), 2.84 (s, 3H).

Examples 637-638:

The compounds listed in the following table were prepared using a procedure similar to that described for example 636:

Structure	Compound No.	MS (ESI) $[M+H]^+$	^1H NMR
	637	511	^1H NMR (400 MHz, Methanol- d_4) δ 8.16 (d, J = 2.4 Hz, 1H), 7.97 (s, 1H), 7.63 (d, J = 2.1 Hz, 1H), 7.54 (d, J = 2.0 Hz, 1H), 7.28 (dd, J = 11.3, 2.5 Hz, 1H), 7.01 (d, J = 2.1 Hz, 1H), 5.20 (p, J = 5.9 Hz, 1H), 4.08 (dd, J = 10.2, 6.5 Hz, 2H), 3.79 (dd, J = 10.4, 5.2 Hz, 2H), 3.03 (s, 3H), 2.53 (s, 3H).
	638	576	^1H NMR (400 MHz, DMSO- d_6) δ 10.01 (s, 2H), 8.18 (d, J = 2.4 Hz, 1H), 7.84 (d, J = 1.9 Hz, 1H), 7.69 (t, J = 1.8 Hz, 1H), 7.63 (t, J = 1.9 Hz, 1H), 7.51 (dd, J = 12.4, 2.4 Hz, 1H), 7.23 (dd, J = 3.5, 1.8 Hz, 1H), 6.92 (t, J = 1.9 Hz, 1H), 5.55 (q, J = 9.1 Hz, 2H), 4.94 (p, J = 5.6 Hz, 1H), 3.77 (td, J = 6.2, 1.8 Hz, 2H), 3.12 – 2.96 (m, 5H), 2.30 (s, 3H).

Example 639: *N*-(3-chloro-5-methanesulfonamidophenyl)-5-(3-fluoro-5-[(3*S*)-1-methylpyrrolidin-3-yl]oxy}pyridin-2-yl)-1-methylpyrrole-3-carboxamide



- Step 1: A mixture of 6-chloro-5-fluoropyridin-3-ol (200 mg, 1.35 mmol, 1 equiv), methyl 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrole-3-carboxylate (359 mg, 1.35 mmol, 1 equiv), K_3PO_4 (863 mg, 4.06 mmol, 3 equiv) and $Pd(dppf)Cl_2$ (198 mg, 0.27 mmol, 0.2 equiv) in 1,4-dioxane (5 mL) was stirred for two hours at 80 °C under N_2 atmosphere. The reaction mixture was concentrated under vacuum. The residue was purified by flash chromatography on silica gel eluting with EA/PE (80%) to afford methyl 5-(3-fluoro-5-hydroxypyridin-2-yl)-1-methylpyrrole-3-carboxylate (230 mg, 68%) as a yellow solid.

LCMS (ESI) $[M + H]^+$: 251

- Step 2: Under nitrogen, a solution of methyl 5-(3-fluoro-5-hydroxypyridin-2-yl)-1-methylpyrrole-3-carboxylate (230 mg, 0.92 mmol, 1 equiv), *tert*-butyl (3*R*)-3-hydroxypyrrolidine-1-carboxylate (172 mg, 0.92 mmol, 1 equiv) and PPh_3 (723 mg, 2.76 mmol, 3 equiv) in THF (5 mL) was added DIAD (558 mg, 2.76 mmol, 3 equiv) at room temperature. The resulting solution was stirred for two hours at 80 °C. The reaction mixture

was concentrated under vacuum. The residue was purified by flash chromatography on silica gel eluting with EA/PE (40%) to afford methyl 5-(5-[(3*S*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-yl]oxy)-3-fluoropyridin-2-yl)-1-methylpyrrole-3-carboxylate (350 mg, 91%) as a white solid.

5 LCMS (ESI) [M + H]⁺: 420

Step 3: A solution of methyl 5-(5-[(3*S*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-yl]oxy)-3-fluoropyridin-2-yl)-1-methylpyrrole-3-carboxylate (350 mg, 0.83 mmol, 1 equiv), *N*-(3-amino-5-chlorophenyl)methanesulfonamide (184 mg, 0.83 mmol, 1 equiv) and Al(Me)₃ (0.5 mL) in Toluene (5 mL) was stirred for 1 hour at 80 °C under nitrogen atmosphere. The

10 reaction mixture was concentrated under vacuum. The residue (500 mg) was purified by Prep-HPLC with the following conditions (Column: Sunfire prep C18 column, 30*150 mm, 5µm; Mobile Phase A: Water(0.1%FA), Mobile Phase B: ACN; Flow rate: 60 mL/min; Gradient: 18% B to 40% B in 8 min; Wave Length: 254/220 nm; RT1(min): 5.27) to afford *tert*-butyl (3*S*)-3-[(6-{4-[(3-chloro-5-methanesulfonamidophenyl)carbamoyl]-1-methylpyrrol-2-yl}-5-fluoropyridin-3-yl)oxy] pyrrolidine-1-carboxylate (300 mg, 59%
15 yield) as a white solid. LCMS (ESI) [M + H]⁺: 608

Step 4: To a stirred solution of *tert*-butyl (3*S*)-3-[(6-{4-[(3-chloro-5-methanesulfonamidophenyl)carbamoyl]-1-methyl pyrrol-2-yl}-5-fluoropyridin-3-yl)oxy]pyrrolidine-1-carboxylate (300 mg, 0.49 mmol, 1 equiv) in 1,4-dioxane (5 mL) was
20 added a dioxane solution of HCl (5 mL, 4 M in dioxane) and the resulting mixture was stirred for 1 hour at 40 °C under nitrogen atmosphere. The reaction mixture was concentrated under vacuum. The residue was purified by flash chromatography on silica gel eluting with EA/PE (32%) to afford *N*-(3-chloro-5-methanesulfonamidophenyl)-5-{3-fluoro-5-[(3*S*)-pyrrolidin-3-yloxy] pyridin-2-yl}-1-methylpyrrole-3-carboxamide (220 mg, 88%) as a yellow solid.



25 LCMS (ESI) [M + H]⁺: 508


Step 5: A solution of *N*-(3-chloro-5-methanesulfonamidophenyl)-5-{3-fluoro-5-[(3*S*)-pyrrolidin-3-yloxy] pyridin-2-yl}-1-methylpyrrole-3-carboxamide (220 mg, 0.43 mmol, 1 equiv), sodium triacetoxymethylborohydride (184 mg, 0.87 mmol, 2 equiv) in formaldehyde (3 mL) was stirred for 1 hour at 60 °C under nitrogen atmosphere. The resulting solid was dried
30 under vacuum. The residue (150 mg) was purified by Prep-HPLC with the following conditions (Column: YMC-Actus Triart C18, 30*150 mm, 5µm; Mobile Phase A: Water(0.1%FA), Mobile Phase B: ACN; Flow rate: 60 mL/min; Gradient: 38% B to 52% B in 7 min, 52% B; Wave Length: 254/220 nm; RT1(min): 6.38) to afford *N*-(3-chloro-5-

methanesulfonamidophenyl)-5-(3-fluoro-5-[[[(3*S*)-1-methylpyrrolidin-3-yl]oxy]pyridin-2-yl)-1-methylpyrrole-3-carboxamide (91.7 mg, 41%) as a white solid. LCMS (ESI) $[M + H]^+$: 522.15. ^1H NMR (300 MHz, DMSO- d_6) δ 10.03 (s, 1H), 9.86 (s, 1H), 8.21 (d, $J = 2.3$ Hz, 1H), 7.72 (q, $J = 2.1$ Hz, 2H), 7.64 (t, $J = 1.9$ Hz, 1H), 7.53 (dd, $J = 12.6, 2.4$ Hz, 1H), 7.08 (t, $J = 2.6$ Hz, 1H), 6.90 (t, $J = 2.0$ Hz, 1H), 5.03 (s, 1H), 3.85 (s, 3H), 3.07 (s, 3H), 2.74 (ddd, $J = 24.9, 10.8, 7.0$ Hz, 3H), 2.32 (d, $J = 24.2$ Hz, 5H), 1.82 (d, $J = 10.2$ Hz, 1H).


Examples 640-642:




The compounds listed in the following table were prepared using a procedure similar to that described for example 639:




Structure	Example No.	MS (ESI) $[M+H]^+$	^1H NMR
	640	522	^1H NMR (300 MHz, DMSO- d_6) δ 9.86 (s, 1H), 8.23 (d, $J = 2.4$ Hz, 1H), 8.15 (s, 1H), 7.72 (q, $J = 1.7$ Hz, 2H), 7.68 – 7.62 (m, 1H), 7.57 (dd, $J = 12.5, 2.4$ Hz, 1H), 7.14 – 7.04 (m, 1H), 6.90 (t, $J = 2.0$ Hz, 1H), 5.10 (s, 1H), 3.85 (s, 3H), 3.06 (s, 3H), 3.02 – 2.89 (m, 3H), 2.78 – 2.57 (m, 1H), 2.46 (s, 4H), 1.93 (dd, $J = 14.3, 7.0$ Hz, 1H).
	641	573	^1H NMR (400 MHz, DMSO- d_6) δ 9.98 (s, 2H), 8.47 (s, 2H), 7.72 (d, $J = 2.0$ Hz, 1H), 7.65 (t, $J = 1.8$ Hz, 1H), 7.63 – 7.56 (m, 2H), 6.85 (d, $J = 2.0$ Hz, 1H), 5.69 (q, $J = 9.1$ Hz, 2H), 5.09 – 4.88 (m, 1H), 2.99 (s, 3H), 2.71 (dd, $J = 10.7, 5.8$ Hz, 1H), 2.63 (ddd, $J = 10.4, 7.0, 2.9$ Hz, 2H), 2.28 (dq, $J = 10.6, 7.7, 6.2$ Hz, 2H), 2.20 (s, 3H), 1.81 – 1.66 (m, 1H).




	642	573	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.05 (s, 2H), 8.55 (s, 2H), 7.80 (d, J = 2.1 Hz, 1H), 7.72 (t, J = 1.8 Hz, 1H), 7.71 – 7.60 (m, 2H), 6.92 (t, J = 2.0 Hz, 1H), 5.76 (q, J = 9.1 Hz, 2H), 5.08 (t, J = 6.6 Hz, 1H), 3.06 (s, 3H), 2.78 (dd, J = 10.7, 5.8 Hz, 1H), 2.70 (ddd, J = 10.4, 6.8, 2.9 Hz, 2H), 2.43 – 2.30 (m, 2H), 2.27 (s, 3H), 1.87 – 1.73 (m, 1H).
---	-----	-----	--



The following compounds were also synthesized using procedures similar to those described above.



Structure	Example No.	MS (ESI) [M+H] ⁺	¹ H NMR
	179	514	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.22 (s, 1H), 10.07 (s, 1H), 9.46 (s, 1H), 8.89 (s, 1H), 8.30 (s, 1H), 8.20 – 8.14 (m, 1H), 7.85 – 7.79 (m, 1H), 7.71 (t, J = 1.7 Hz, 1H), 7.64 – 7.60 (m, 1H), 7.54 (dd, J = 8.3, 4.6 Hz, 1H), 7.15 (t, J = 7.8 Hz, 1H), 6.95 (t,




	180	491	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.30 (s, 1H), 10.16 (s, 1H), 9.41 (s, 1H), 8.67 (d, J = 2.5 Hz, 1H), 8.31 (s, 1H), 8.15 (dd, J = 8.8, 2.5 Hz, 1H), 8.04 (d, J = 8.8 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 6.94 (t, J = 2.0 Hz, 1H), 4.28 (s
	181	420	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.97 (s, 1H), 8.62 – 8.55 (m, 1H), 8.33 (s, 1H), 8.15 (d, J = 1.9 Hz, 1H), 8.07 (td, J = 7.8, 1.8 Hz, 1H), 7.79 (d, J = 8.1 Hz, 1H), 7.71 (t, J = 1.8 Hz, 1H), 7.61 (t, J = 1.9 Hz, 1H), 7.47 (dd, J = 7.4, 4.8 Hz, 1H), 6.90 (t, J
	182	385	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.05 (s, 2H), 8.44 (s, 1H), 8.06 (s, 1H), 7.67 (s, 1H), 7.58 (s, 1H), 6.93 (s, 1H), 5.10 (s, 1H), 4.05 – 3.89 (m, 3H), 3.84 (q, J = 7.6 Hz, 1H), 3.06 (s, 3H), 2.42 (p, J = 7.8, 7.1 Hz, 1H), 2.34 – 2.20 (m, 1H).




	183	505	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.26 (s, 1H), 10.11 (s, 1H), 9.31 (d, J = 0.8 Hz, 1H), 8.62 (s, 2H), 8.24 (d, J = 0.8 Hz, 1H), 7.68 (t, J = 1.8 Hz, 1H), 7.59 (t, J = 1.9 Hz, 1H), 6.92 (t, J = 2.0 Hz, 1H), 4.00 (s, 2H), 3.69 (dd, J = 6.5, 4.3 Hz, 2H), 3.50 (d
	184	505	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.23 (s, 1H), 9.21 (s, 1H), 8.74 (d, J = 1.4 Hz, 1H), 8.34 (s, 1H), 8.30 (d, J = 1.5 Hz, 1H), 8.27 (s, 1H), 7.68 (t, J = 1.8 Hz, 1H), 7.58 (d, J = 1.9 Hz, 1H), 6.92 (t, J = 2.0 Hz, 1H), 4.21 (s, 2H), 3.94 (t, J = 5.4 Hz, 2H),
	185	580	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.03 (s, 1H), 9.97 (s, 1H), 8.67 (s, 1H), 8.53 (d, J = 4.0 Hz, 1H), 8.25 (d, J = 4.6 Hz, 1H), 7.86 (dt, J = 7.9, 2.0 Hz, 1H), 7.82 (s, 1H), 7.73 – 7.66 (m, 2H), 7.62 (t, J = 1.9 Hz, 1H), 7.41 (dd, J = 7.9, 4.8 Hz, 1H), 7.38 –




	186	517	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.22 (s, 1H), 10.08 (s, 1H), 8.94 (d, J = 0.7 Hz, 1H), 8.56 (d, J = 2.7 Hz, 2H), 8.32 (d, J = 0.7 Hz, 1H), 8.22 (dd, J = 4.7, 1.3 Hz, 1H), 7.91 (dd, J = 8.3, 1.0 Hz, 1H), 7.82 (dt, J = 9.5, 2.3 Hz, 1H), 7.71 (t, J = 1.8 Hz, 1H)
	187	534	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.23 (s, 1H), 10.08 (s, 1H), 8.96 (s, 1H), 8.33 (s, 1H), 8.21 (dd, J = 4.7, 1.3 Hz, 1H), 7.84 (dd, J = 8.4, 1.4 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.58 (dd, J = 8.3, 4.6 Hz, 1H), 7.20 (ddd, J = 7.5, 5
	188	500	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.19 (s, 1H), 9.17 (s, 1H), 8.93 (d, J = 0.7 Hz, 1H), 8.90 (s, 2H), 8.31 (d, J = 0.7 Hz, 1H), 8.23 (dd, J = 4.7, 1.3 Hz, 1H), 8.10 (s, 1H), 7.94 (dd, J = 8.4, 1.4 Hz, 1H), 7.67 (m, 1H), 7.61 (dd, J = 8.3, 4.7 Hz, 1H), 7.57 (m,



	189	516	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.20 (s, 1H), 10.08 (s, 1H), 8.86 (s, 1H), 8.29 (s, 1H), 8.21 (dd, J = 4.6, 1.3 Hz, 1H), 7.94 (dd, J = 8.4, 1.3 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.64 – 7.52 (m, 3H), 7.46 – 7.35 (m, 1H), 7.30 – 7.17 (m, 2H), 6.95 (t, J = 2.0
	190	476	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.30 (s, 1H), 10.09 (s, 1H), 9.40 (s, 1H), 8.48 (d, J = 2.1 Hz, 1H), 8.29 (s, 1H), 7.99 (dd, J = 8.6, 2.2 Hz, 1H), 7.93 (d, J = 8.5 Hz, 1H), 7.73 (t, J = 1.8 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 3.99 (d




	191	503	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.25 (s, 1H), 10.17 (s, 1H), 9.26 (s, 1H), 8.22 (s, 1H), 8.11 (d, J = 2.9 Hz, 1H), 7.80 (d, J = 9.0 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.51 (dd, J = 9.1, 3.0 Hz, 1H), 6.93 (t, J = 1.9 Hz, 1H), 4.47 (s
	192	513	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.26 (s, 1H), 9.30 (s, 1H), 8.35 (d, J = 2.9 Hz, 1H), 8.24 (s, 1H), 7.86 (d, J = 9.0 Hz, 1H), 7.76 (dd, J = 9.0, 3.0 Hz, 1H), 7.71 (s, 1H), 7.62 (s, 1H), 7.14 (s, 1H), 6.93 (d, J = 4.8 Hz, 2H), 4.55 (s, 2H), 4.23 – 4.07 (m, 2H




	193	611	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.16 (s, 1H), 10.07 (s, 1H), 8.71 (s, 1H), 8.62 (d, J = 2.2 Hz, 1H), 8.52 (dd, J = 4.8, 1.6 Hz, 1H), 8.23 (s, 1H), 7.86 (d, J = 2.4 Hz, 1H), 7.81 (dt, J = 8.0, 1.9 Hz, 1H), 7.70 (t, J = 1.8 Hz, 1H), 7.61 (t, J = 1.8 Hz, 1H), 7
	194	584	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.61 (s, 1H), 8.51 (s, 1H), 8.44 (d, J = 3.3 Hz, 1H), 8.20 (s, 1H), 7.79 (d, J = 5.7 Hz, 2H), 7.60 – 7.55 (m, 1H), 7.52 (t, J = 1.7 Hz, 1H), 7.43 – 7.35 (m, 1H), 7.24 (s, 1H), 6.97 – 6.90 (m, 1H), 5.24 (s, 2H), 3.84 – 3.65 (m,
	195	480	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.19 (s, 1H), 10.10 (s, 1H), 8.90 (s, 1H), 8.30 (s, 1H), 8.16 (dd, J = 4.6, 1.3 Hz, 1H), 7.81 (dd, J = 8.4, 1.4 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.53 (dd, J = 8.3, 4.6 Hz, 1H), 6.94 (t, J = 1.9 Hz,



	196	486	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.21 (s, 1H), 9.16 (s, 1H), 8.55 (dd, J = 14.8, 4.8 Hz, 3H), 8.12 – 8.00 (m, 1H), 7.96 (s, 1H), 7.71 – 7.58 (m, 2H), 7.57 (s, 1H), 7.23 (t, J = 4.8 Hz, 1H), 6.92 (s, 1H), 3.05 (s, 3H).
	197	422	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.28 (s, 1H), 10.09 (s, 1H), 9.31 (s, 1H), 8.26 (m, 2H), 7.94 (d, J = 9.0 Hz, 1H), 7.73 (t, J = 1.8 Hz, 1H), 7.68 (dd, J = 9.0, 3.0 Hz, 1H), 7.63 (t, J = 1.9 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 3.91 (s, 3H), 3.08 (s, 3H).
	198	470	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.45 (s, 1H), 9.60 (s, 1H), 9.12 (s, 1H), 8.62 (d, J = 8.3 Hz, 1H), 8.47 (s, 1H), 8.29 (d, J = 8.8 Hz, 1H), 7.73 (d, J = 30.3 Hz, 2H), 7.01 (s, 1H), 3.46 (s, 3H), 3.13 (s, 3H).




	199	470	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.34 (s, 1H), 10.12 (s, 1H), 9.08 (s, 1H), 8.90 (dd, J = 4.8, 1.7 Hz, 1H), 8.62 (dd, J = 8.0, 1.7 Hz, 1H), 8.36 (s, 1H), 7.88 (dd, J = 8.0, 4.8 Hz, 1H), 7.72 (t, J = 1.8 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 6.96 (t, J = 1.9 Hz,
	200	499	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.39 (s, 1H), 10.09 (s, 1H), 9.50 (s, 1H), 8.89 (dd, J = 2.4, 0.8 Hz, 1H), 8.44 – 8.36 (m, 2H), 8.21 (dd, J = 8.6, 0.8 Hz, 1H), 7.73 (t, J = 1.9 Hz, 1H), 7.63 (t, J = 1.9 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 3.07 (s, 3H), 2.71 (
	201	468	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.35 (s, 1H), 10.13 (s, 1H), 9.18 (s, 1H), 8.74 (dt, J = 4.4, 2.0 Hz, 1H), 8.65 (ddd, J = 12.3, 7.8, 1.9 Hz, 1H), 8.34 (s, 1H), 7.82 – 7.68 (m, 2H), 7.62 (t, J = 1.9 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 3.07 (s, 3H), 1.71 (d, J




	202	534	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.22 (s, 1H), 10.06 (s, 1H), 8.87 (s, 1H), 8.31 (s, 1H), 8.24 (d, J = 2.4 Hz, 1H), 7.97 (dd, J = 10.1, 2.4 Hz, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.61 (t, J = 1.9 Hz, 1H), 7.43 (td, J = 8.1, 6.1 Hz, 1H), 7.30 – 7.22 (m, 2H), 7.16
	203	541	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.21 (s, 1H), 10.05 (s, 1H), 8.87 (s, 1H), 8.30 (s, 1H), 8.25 (d, J = 2.4 Hz, 1H), 7.98 (dd, J = 10.1, 2.5 Hz, 1H), 7.89 (d, J = 1.8 Hz, 1H), 7.81 (dt, J = 7.7, 1.5 Hz, 1H), 7.77 (d, J = 7.9 Hz, 1H), 7.70 (t, J = 1.9 Hz, 1H),




	204	530	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.22 (s, 1H), 10.07 (s, 1H), 8.89 (s, 1H), 8.30 (s, 1H), 7.75 (d, J = 8.4 Hz, 1H), 7.72 (t, J = 1.8 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.40 (m, 2H), 7.30 – 7.23 (m, 2H), 7.14 (t, J = 8.1 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 5.29
	205	505	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.21 (s, 1H), 10.07 (s, 1H), 8.91 (s, 1H), 8.31 (s, 1H), 8.23 (d, J = 4.6 Hz, 1H), 7.98 (dd, J = 8.4, 1.2 Hz, 1H), 7.83 (d, J = 3.2 Hz, 1H), 7.78 (d, J = 3.2 Hz, 1H), 7.71 (d, J = 1.9 Hz, 1H), 7.64 – 7.53 (m, 2H), 6.94 (t, J =
	206	488	¹ H NMR (400 MHz, DMSO-d ₆) δ 12.84 (s, 1H), 10.18 (s, 1H), 10.06 (s, 1H), 8.82 (s, 1H), 8.26 (s, 1H), 8.15 (d, J = 4.6 Hz, 1H), 7.92 (d, J = 8.3 Hz, 1H), 7.80 (s, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.66 – 7.42 (m, 3H), 6.94 (t, J = 2.0 Hz, 1H), 5.18 (s, 2H), 3




	207	424	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.03 (s, 1H), 10.01 (s, 1H), 8.99 (s, 2H), 8.42 (d, J = 2.1 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.66 (t, J = 1.9 Hz, 1H), 6.91 (t, J = 1.9 Hz, 1H), 6.58 (dd, J = 2.1, 1.2 Hz, 1H), 3.06 (s, 3H), 2.57 (s, 3H).
	208	450	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.10 (s, 1H), 9.97 (s, 1H), 8.63 (s, 2H), 8.35 (d, J = 2.1 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 6.89 (t, J = 2.0 Hz, 1H), 6.55 (s, 1H), 4.26 (q, J = 6.9 Hz, 2H), 3.05 (s, 3H), 2.55 (s, 3H), 1.39 (t, J =
	209	364	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.16 (s, 1H), 10.08 (s, 1H), 8.32 (dd, J = 7.0, 1.1 Hz, 1H), 8.17 (d, J = 1.6 Hz, 1H), 7.75 (t, J = 1.8 Hz, 1H), 7.68 (t, J = 1.9 Hz, 1H), 7.50 (d, J = 9.1 Hz, 1H), 6.99 (t, J = 1.2 Hz, 1H), 6.93 (t, J = 2.0 Hz, 1H), 6.78 (ddd




	210	515	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.14 (s, 1H), 10.06 (s, 1H), 8.84 (s, 1H), 8.29 (s, 1H), 7.70 – 7.65 (m, 2H), 7.59 (t, J = 1.9 Hz, 1H), 7.47 – 7.37 (m, 2H), 7.35 (dd, J = 8.5, 1.3 Hz, 1H), 7.29 – 7.21 (m, 2H), 7.18 – 7.14 (m, 1H), 7.14 – 7.10 (m, 1H), 6.94 (
	211	533	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.17 (s, 1H), 10.07 (s, 1H), 8.66 (s, 1H), 8.32 (s, 1H), 7.69 (t, J = 1.9 Hz, 1H), 7.59 (t, J = 1.9 Hz, 1H), 7.57 – 7.50 (m, 1H), 7.39 (td, J = 8.0, 5.9 Hz, 1H), 7.13 (dt, J = 24.6, 8.6 Hz, 5H), 6.94 (t, J = 2.0 Hz, 1H), 5.26




	212	571	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.58 (s, 1H), 10.11 (s, 1H), 8.53 (q, J = 4.1 Hz, 1H), 8.29 (dd, J = 4.7, 1.2 Hz, 1H), 8.26 (s, 1H), 7.69 (t, J = 1.9 Hz, 1H), 7.65 (ddd, J = 4.4, 3.0, 1.3 Hz, 2H), 7.46 (dd, J = 8.4, 4.7 Hz, 1H), 7.42 – 7.25 (m, 5H), 6.98 (t,
	213	585	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.61 (s, 1H), 10.13 (s, 1H), 8.41 (s, 1H), 8.22 (d, J = 4.5 Hz, 1H), 7.70 – 7.63 (m, 2H), 7.60 (d, J = 8.3 Hz, 1H), 7.50 – 7.24 (m, 6H), 6.98 (s, 1H), 5.25 (s, 2H), 3.08 (s, 3H), 2.84 (s, 3H), 2.59 (s, 3H).
	214	498	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.67 (s, 1H), 10.09 (s, 1H), 8.83 (s, 1H), 8.46 (s, 1H), 8.27 (dd, J = 4.6, 1.2 Hz, 1H), 7.69 (d, J = 8.4 Hz, 1H), 7.53 (d, J = 7.4 Hz, 3H), 7.50 – 7.45 (m, 1H), 7.42 (t, J = 7.4 Hz, 2H), 7.38 – 7.32 (m, 2H), 6.74 (dt, J = 10.




	215	505	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.63 (s, 1H), 10.10 (s, 1H), 8.80 (d, J = 1.3 Hz, 1H), 8.44 (s, 1H), 8.38 (d, J = 1.3 Hz, 1H), 8.31 (dd, J = 4.6, 1.2 Hz, 1H), 7.80 (dd, J = 8.5, 1.3 Hz, 1H), 7.69 (dt, J = 5.0, 1.9 Hz, 2H), 7.45 – 7.36 (m, 2H), 6.95 (t, J = 2
	216	544	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.38 (s, 1H), 10.07 (s, 1H), 8.30 (s, 1H), 8.26 (dd, J = 4.6, 1.3 Hz, 1H), 7.70 – 7.62 (m, 3H), 7.47 – 7.42 (m, 2H), 7.40 (dd, J = 8.4, 4.7 Hz, 1H), 7.37 – 7.25 (m, 3H), 6.95 (t, J = 1.9 Hz, 1H), 5.80 (s, 1H), 5.23 (s, 2H), 4.
	217	515	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.63 (s, 1H), 10.06 (s, 1H), 8.81 (d, J = 1.4 Hz, 1H), 8.75 (d, J = 2.2 Hz, 1H), 8.57 (dd, J = 4.8, 1.7 Hz, 1H), 8.43 (d, J = 1.3 Hz, 1H), 8.29 (dd, J = 4.6, 1.3 Hz, 1H), 7.96 (dt, J = 7.9, 2.0 Hz, 1H), 7.74 (dd, J = 8.5, 1.3




	218	438	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.29 (s, 2H), 9.37 (s, 1H), 8.45 (d, J = 2.4 Hz, 1H), 8.29 (s, 1H), 7.99 (dd, J = 8.7, 2.4 Hz, 1H), 7.93 (d, J = 8.6 Hz, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.61 (d, J = 1.9 Hz, 1H), 6.93 (t, J = 2.0 Hz, 1H), 3.05 (s, 3H), 2.60 (s,
	219	438	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.33 (s, 1H), 9.17 (s, 1H), 8.41 (dd, J = 4.6, 1.5 Hz, 1H), 8.35 (s, 1H), 8.07 (dd, J = 8.1, 1.5 Hz, 1H), 7.75 (t, J = 1.9 Hz, 1H), 7.65 (t, J = 1.9 Hz, 1H), 7.61 (dd, J = 8.0, 4.6 Hz, 1H), 7.00 (t, J = 2.0 Hz, 1H), 3.11 (s, 3
	220	491	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.60 (s, 1H), 10.08 (s, 1H), 10.01 (s, 1H), 8.56 (d, J = 1.4 Hz, 1H), 8.49 (dd, J = 4.6, 1.6 Hz, 1H), 8.19 (d, J = 1.3 Hz, 1H), 7.90 (dd, J = 8.2, 1.6 Hz, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.65 (t, J = 1.9 Hz, 1H), 7.39 (dd, J =




	221	493	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.56 (s, 1H), 10.09 (s, 1H), 8.56 (dd, J = 4.7, 1.7 Hz, 1H), 8.36 (d, J = 1.4 Hz, 1H), 7.92 (d, J = 1.4 Hz, 1H), 7.74 – 7.63 (m, 3H), 7.37 (dd, J = 7.7, 4.7 Hz, 1H), 6.96 (t, J = 1.9 Hz, 1H), 3.88 (s, 2H), 3.08 (s, 3H), 2.99 (
	222	479	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.65 (s, 1H), 10.10 (s, 1H), 8.73 (dd, J = 4.8, 1.7 Hz, 1H), 8.56 (d, J = 1.4 Hz, 1H), 7.92 (d, J = 1.3 Hz, 1H), 7.81 (dd, J = 7.7, 1.7 Hz, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.67 (t, J = 1.9 Hz, 1H), 7.48 (dd, J = 7.7, 4.8 Hz, 1H
	223	476	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.56 (s, 1H), 10.08 (s, 1H), 8.94 (dt, J = 4.7, 1.3 Hz, 1H), 8.35 (dd, J = 8.1, 1.6 Hz, 1H), 8.32 (d, J = 1.4 Hz, 1H), 8.08 (s, 1H), 7.68 (ddd, J = 8.0, 4.4, 1.4 Hz, 2H), 7.63 (d, J = 1.9 Hz, 1H), 6.96 (t, J = 1.9 Hz, 1H), 3.0




	224	436	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.57 (s, 1H), 10.13 (s, 1H), 8.51 (dd, J = 4.7, 1.6 Hz, 2H), 8.20 (d, J = 1.3 Hz, 1H), 7.81 – 7.72 (m, 1H), 7.68 (p, J = 1.8 Hz, 2H), 7.31 (dd, J = 7.7, 4.7 Hz, 1H), 6.97 (t, J = 1.9 Hz, 1H), 3.19 (q, J = 7.3 Hz, 2H), 2.51 (s,
	225	551	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.62 (s, 1H), 10.10 (s, 1H), 8.74 (d, J = 1.3 Hz, 1H), 8.65 (d, J = 1.8 Hz, 1H), 8.61 (d, J = 2.8 Hz, 1H), 8.39 (d, J = 1.3 Hz, 1H), 8.34 (d, J = 2.2 Hz, 1H), 7.95 (dt, J = 9.6, 2.2 Hz, 1H), 7.84 (dd, J = 10.8, 2.3 Hz, 1H), 7.
	226	454	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.55 (s, 1H), 10.11 (s, 1H), 8.51 (d, J = 2.9 Hz, 1H), 8.45 (d, J = 1.4 Hz, 1H), 8.17 (d, J = 1.4 Hz, 1H), 7.76 (dd, J = 9.7, 2.9 Hz, 1H), 7.67 (dt, J = 5.2, 1.9 Hz, 2H), 6.97 (t, J = 1.9 Hz, 1H), 3.18 (q, J = 7.3 Hz, 2H), 2.5




	227	438	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.38 (s, 1H), 10.17 (s, 1H), 10.06 (s, 1H), 8.17 (s, 1H), 8.15 (dd, J = 4.6, 1.4 Hz, 1H), 7.69 (t, J = 1.9 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 7.36 (dd, J = 8.2, 1.5 Hz, 1H), 7.25 (dd, J = 8.2, 4.5 Hz, 1H), 6.94 (t, J = 1.9 Hz,
	228	550	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.57 (s, 1H), 10.07 (s, 1H), 8.70 (d, J = 4.8 Hz, 1H), 8.50 (s, 1H), 8.11 (s, 1H), 8.08 (d, J = 7.9 Hz, 1H), 7.67 (d, J = 2.0 Hz, 2H), 7.49 (dd, J = 7.8, 4.7 Hz, 1H), 6.96 (d, J = 2.1 Hz, 1H), 6.91 – 6.75 (m, 3H), 5.26 (s, 2H)
	229	517	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.34 (s, 1H), 10.06 (s, 1H), 8.78 (dd, J = 4.8, 1.6 Hz, 1H), 8.54 (d, J = 2.7 Hz, 1H), 8.29 (t, J = 1.7 Hz, 1H), 8.09 (dd, J = 7.8, 1.7 Hz, 1H), 7.91 (s, 1H), 7.70 (ddd, J = 9.9, 2.7, 1.7 Hz, 1H), 7.67 – 7.56 (m, 3H), 6.95 (t,




	230	382	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.89 (s, 1H), 10.24 (s, 1H), 9.36 (s, 1H), 8.57 (d, J = 5.5 Hz, 1H), 8.42 (s, 1H), 8.00 (d, J = 5.5 Hz, 1H), 7.70 (s, 1H), 7.65 (s, 1H), 6.99 (d, J = 2.1 Hz, 1H), 3.07 (s, 3H).
	231	576	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.22 (s, 1H), 10.08 (s, 1H), 8.94 (s, 1H), 8.31 (s, 1H), 8.20 (dd, J = 4.7, 1.3 Hz, 1H), 7.86 (dd, J = 8.4, 1.3 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.69 (d, J = 1.8 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.58 (dd, J = 8.3, 4.6 Hz,
	232	436	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 10.08 (s, 1H), 8.75 – 8.63 (m, 1H), 8.36 (s, 1H), 7.93 (td, J = 7.7, 1.8 Hz, 1H), 7.71 – 7.65 (m, 2H), 7.64 (t, J = 1.9 Hz, 1H), 7.37 (ddd, J = 7.5, 4.8, 1.1 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 3.18 (q, J = 7.4 Hz




	233	450	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.41 (s, 1H), 10.08 (s, 1H), 8.79 – 8.50 (m, 1H), 8.27 (s, 1H), 7.93 (td, J = 7.7, 1.9 Hz, 1H), 7.74 – 7.50 (m, 3H), 7.45 – 7.26 (m, 1H), 6.95 (t, J = 2.0 Hz, 1H), 3.99 (p, J = 6.7 Hz, 1H), 3.07 (s, 3H), 1.31 (d, J = 6.8 Hz, 6)
	234	452	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.46 (s, 1H), 10.06 (s, 1H), 8.68 (d, J = 4.7 Hz, 1H), 8.48 (s, 1H), 7.95 (t, J = 7.8 Hz, 1H), 7.77 – 7.58 (m, 3H), 7.37 (dd, J = 7.5, 4.8 Hz, 1H), 6.97 (d, J = 2.3 Hz, 1H), 4.98 (s, 2H), 3.44 (s, 3H), 3.07 (s, 3H).
	235	502	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.48 (s, 1H), 10.06 (s, 1H), 8.52 (s, 1H), 8.41 (s, 1H), 7.65 (d, J = 12.3 Hz, 2H), 7.46 – 7.25 (m, 5H), 6.94 (d, J = 2.1 Hz, 1H), 5.69 (s, 2H), 3.06 (s, 3H), 2.63 (s, 3H).




	236	488	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.56 (s, 1H), 10.10 (s, 1H), 9.08 (s, 1H), 8.51 (s, 1H), 8.16 – 7.93 (m, 2H), 7.75 – 7.59 (m, 4H), 7.57 – 7.48 (m, 1H), 6.96 (t, J = 2.0 Hz, 1H), 3.08 (s, 3H), 2.71 (s, 3H).
	237	501	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.59 (s, 1H), 10.08 (s, 1H), 9.19 (s, 1H), 8.59 (d, J = 1.4 Hz, 1H), 8.49 (dd, J = 4.6, 1.6 Hz, 1H), 8.35 (d, J = 4.8 Hz, 2H), 8.15 (d, J = 1.3 Hz, 1H), 7.98 (dd, J = 8.2, 1.6 Hz, 1H), 7.68 (dt, J = 9.1, 1.9 Hz, 2H), 7.41 (dd,
	238	515	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.60 (s, 1H), 10.10 (s, 1H), 8.67 – 8.52 (m, 2H), 8.33 (s, 2H), 7.86 (dd, J = 8.0, 1.6 Hz, 1H), 7.79 (d, J = 1.3 Hz, 1H), 7.67 (dt, J = 5.0, 1.9 Hz, 2H), 7.49 (dd, J = 8.0, 4.6 Hz, 1H), 6.96 (t, J = 2.0 Hz, 1H), 6.72 (t, J = 4




	239	501	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.67 (s, 1H), 10.09 (s, 1H), 8.82 (s, 1H), 8.49 (t, J = 16.8 Hz, 4H), 7.70 (s, 2H), 7.49 (d, J = 23.5 Hz, 4H), 6.97 (s, 1H), 3.08 (s, 3H).
	240	545	¹ H NMR (300 MHz, Methanol-d ₄) δ 7.81 (s, 1H), 7.65 (t, J = 1.9 Hz, 1H), 7.54 (t, J = 1.9 Hz, 1H), 7.44 – 7.34 (m, 1H), 7.28 (d, J = 4.0 Hz, 5H), 7.06 – 7.00 (m, 2H), 6.87 (t, J = 8.6 Hz, 1H), 5.13 (s, 2H), 3.05 (s, 3H), 2.32 (s, 3H).
	241	407	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.50 (s, 1H), 10.12 (s, 1H), 8.10 (d, J = 1.4 Hz, 1H), 8.05 (d, J = 1.4 Hz, 1H), 7.67 (t, J = 1.9 Hz, 1H), 7.59 (t, J = 1.9 Hz, 1H), 6.97 (t, J = 2.0 Hz, 1H), 3.07 (s, 3H).




	242	458	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.31 (s, 1H), 10.08 (s, 1H), 7.64 – 7.57 (m, 2H), 7.55 (d, J = 1.7 Hz, 1H), 6.95 (t, J = 2.0 Hz, 1H), 6.27 (d, J = 1.7 Hz, 1H), 4.22 (dd, J = 9.0, 2.4 Hz, 1H), 3.66 (s, 3H), 3.53 – 3.42 (m, 1H), 3.27 (d, J = 7.7 Hz, 1H), 3.06
	243	414	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.22 (s, 1H), 7.98 (s, 1H), 7.51 (dd, J = 5.0, 3.0 Hz, 2H), 7.44 (d, J = 2.0 Hz, 1H), 6.85 (t, J = 2.0 Hz, 1H), 3.07 (d, J = 12.1 Hz, 2H), 2.93 (s, 3H), 2.66 (q, J = 12.6, 11.9 Hz, 3H), 1.87 (d, J = 12.8 Hz, 2H), 1.51 (q, J =
	244	415	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.49 (s, 1H), 7.77 (d, J = 1.8 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 7.56 (t, J = 1.9 Hz, 1H), 7.01 (t, J = 1.9 Hz, 1H), 6.81 (d, J = 1.7 Hz, 1H), 3.34 (dd, J = 6.5, 3.5 Hz, 8H), 3.03 (s, 3H).



	245	416	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.28 (s, 1H), 10.07 (s, 1H), 7.88 (d, J = 1.7 Hz, 1H), 7.62 (dt, J = 15.9, 2.0 Hz, 2H), 6.95 (t, J = 2.0 Hz, 1H), 6.77 (d, J = 1.7 Hz, 1H), 3.82 – 3.64 (m, 4H), 3.06 (d, J = 7.9 Hz, 7H).
	246	497	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.26 (s, 1H), 10.06 (s, 1H), 7.87 (d, J = 1.7 Hz, 1H), 7.61 (dt, J = 16.3, 2.0 Hz, 2H), 6.94 (t, J = 2.0 Hz, 1H), 6.73 (d, J = 1.6 Hz, 1H), 3.25 (q, J = 10.3 Hz, 2H), 3.08 (d, J = 13.9 Hz, 7H), 2.77 (t, J = 5.0 Hz, 4H).
	247	516	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.97 (s, 1H), 9.79 (s, 1H), 8.25 (d, J = 3.0 Hz, 1H), 7.83 (d, J = 2.1 Hz, 1H), 7.69 (q, J = 2.0 Hz, 2H), 7.55 (dd, J = 9.0, 3.1 Hz, 1H), 7.45 (d, J = 8.8 Hz, 1H), 6.90 (t, J = 2.0 Hz, 1H), 6.53 (t, J = 1.5 Hz, 1H), 3.77 (dd, J



	248	526	¹ H NMR (400 MHz, DMSO-d ₆) δ 11.21 (s, 1H), 9.83 (s, 1H), 8.25 (d, J = 3.0 Hz, 1H), 7.83 (d, J = 2.0 Hz, 1H), 7.77 (s, 1H), 7.68 (s, 1H), 7.55 (dd, J = 8.9, 3.0 Hz, 1H), 7.45 (d, J = 8.8 Hz, 1H), 7.15 (t, J = 52.4 Hz, 1H), 6.92 (d, J = 2.0 Hz, 1H), 6.53 (d
	249	504	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.04 (s, 1H), 9.79 (s, 1H), 8.25 (d, J = 3.0 Hz, 1H), 7.83 (d, J = 2.0 Hz, 1H), 7.69 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.55 (dd, J = 8.9, 3.1 Hz, 1H), 7.45 (d, J = 8.8 Hz, 1H), 6.89 (t, J = 2.0 Hz, 1H), 6.53 (dd,
	250	425	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.30 (s, 1H), 8.04 (s, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.54 (t, J = 1.8 Hz, 1H), 6.99 (t, J = 1.9 Hz, 1H), 4.72 (dq, J = 11.6, 6.0 Hz, 1H), 4.53 (s, 2H), 3.02 (s, 3H), 2.18 (td, J = 12.3, 3.4 Hz, 2H), 2.02 (dd, J = 12.2, 5.4




	251	532	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.26 (s, 1H), 10.09 (s, 1H), 8.90 (s, 1H), 8.49 (s, 1H), 8.34 (s, 1H), 8.07 – 7.90 (m, 1H), 7.68 (d, J = 30.3 Hz, 2H), 7.43 – 7.14 (m, 1H), 6.98 (d, J = 12.0 Hz, 4H), 3.17 (t, J = 7.7 Hz, 2H), 3.08 (s, 3H), 2.86 (d, J = 8.3 Hz
	252	488	¹ H NMR (400 MHz, DMSO-d ₆) δ 12.33 (s, 1H), 10.20 (d, J = 4.7 Hz, 1H), 8.89 (d, J = 2.6 Hz, 1H), 8.31 – 8.12 (m, 3H), 8.05 (dd, J = 8.4, 1.4 Hz, 1H), 7.69 (d, J = 2.1 Hz, 1H), 7.62 – 7.46 (m, 2H), 6.93 (t, J = 2.0 Hz, 3H), 5.26 (s, 2H), 3.06 (s, 3H).
	253	488	¹ H NMR (400 MHz, DMSO-d ₆) δ 12.86 (s, 1H), 10.16 (s, 1H), 8.84 (s, 1H), 8.39 (s, 1H), 8.29 – 8.09 (m, 2H), 7.95 (d, J = 8.4 Hz, 1H), 7.76 – 7.47 (m, 4H), 6.90 (s, 1H), 6.32 (d, J = 2.2 Hz, 1H), 5.27 (s, 2H), 3.01 (s, 3H).




	254	488	¹ H NMR (400 MHz, DMSO-d ₆) δ 12.87 (s, 1H), 10.19 (s, 1H), 10.06 (s, 1H), 8.85 (s, 1H), 8.26 (s, 1H), 8.16 (dd, J = 4.7, 1.3 Hz, 1H), 8.07 – 7.93 (m, 2H), 7.71 (t, J = 1.9 Hz, 1H), 7.60 (t, J = 1.9 Hz, 1H), 7.54 (dd, J = 8.3, 4.6 Hz, 1H), 7.34 (s, 1H), 6.9
	255	529	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.07 (s, 2H), 8.54 (s, 1H), 8.29 (s, 1H), 7.65 (t, J = 1.9 Hz, 1H), 7.53 (t, J = 1.9 Hz, 1H), 7.45 – 7.26 (m, 2H), 7.15 – 6.97 (m, 5H), 6.90 (t, J = 1.9 Hz, 1H), 5.17 (s, 2H), 3.02 (s, 3H), 2.04 (s, 3H).
	256	534	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.30 (s, 1H), 10.06 (s, 1H), 9.28 (s, 1H), 8.58 (dd, J = 4.8, 1.7 Hz, 1H), 8.29 (s, 1H), 8.23 – 8.09 (m, 1H), 7.72 (d, J = 2.0 Hz, 1H), 7.58 (dd, J = 7.7, 4.7 Hz, 2H), 6.95 (d, J = 2.0 Hz, 1H), 6.86 – 6.59 (m, 3H), 5.55 (s, 2H)




	257	486	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.06 (d, J = 15.6 Hz, 2H), 8.51 (d, J = 2.7 Hz, 2H), 8.13 (s, 2H), 7.74 – 7.60 (m, 5H), 7.56 – 7.45 (m, 2H), 6.93 (t, J = 1.9 Hz, 1H), 3.05 (s, 3H).
	258	631	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.11 (d, J = 25.0 Hz, 2H), 8.70 (s, 1H), 8.23 (s, 1H), 7.69 (t, J = 1.9 Hz, 1H), 7.60 (t, J = 1.9 Hz, 1H), 7.37 (d, J = 2.3 Hz, 1H), 7.22 – 7.05 (m, 3H), 6.93 (t, J = 2.0 Hz, 1H), 6.77 (d, J = 2.3 Hz, 1H), 5.27 (s, 2H), 4.75 (
 *stereochemistry arbitrarily assigned*	259	475	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.95 (s, 1H), 8.43 (s, 1H), 7.98 (s, 1H), 7.63 (t, J = 1.9 Hz, 1H), 7.55 (t, J = 2.0 Hz, 1H), 7.29 – 7.16 (m, 2H), 6.97 – 6.86 (m, 2H), 6.86 – 6.65 (m, 2H), 5.01 – 4.74 (m, 2H), 3.04 (s, 3H), 2.39 – 2.19 (m, 2H), 2.19 – 2.10 (m




 *stereochemistry arbitrarily assigned*	260	459	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.82 (s, 1H), 7.90 (s, 1H), 7.60 (t, J = 1.8 Hz, 1H), 7.53 (t, J = 1.9 Hz, 1H), 7.16 – 7.03 (m, 3H), 7.03 – 6.82 (m, 3H), 5.02 (td, J = 7.2, 3.0 Hz, 1H), 3.50 – 3.40 (m, 1H), 3.04 (s, 3H), 2.44 – 2.03 (m, 4H), 1.
	261	503	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.74 (s, 1H), 8.27 (s, 1H), 8.21 (s, 1H), 8.04 (d, J = 1.8 Hz, 1H), 7.80 – 7.75 (m, 1H), 7.64 (t, J = 1.9 Hz, 1H), 7.57 (t, J = 1.9 Hz, 1H), 7.27 (s, 1H), 7.01 (t, J = 1.9 Hz, 1H), 5.35 (s, 2H), 3.03 (s, 3H), 2.46 (s, 3H).




	262	517	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.30 (s, 2H), 9.30 (d, J = 0.8 Hz, 1H), 8.59 (dd, J = 4.8, 1.7 Hz, 1H), 8.34 – 8.14 (m, 4H), 7.71 (t, J = 1.9 Hz, 1H), 7.64 – 7.55 (m, 2H), 7.50 (dt, J = 11.0, 2.4 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 5.63 (s, 2H), 3.07 (s, 3H).
	263	485	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.09 (d, J = 17.8 Hz, 2H), 8.68 (s, 1H), 8.59 (d, J = 4.8 Hz, 2H), 8.14 (s, 1H), 7.79 (dd, J = 8.2, 1.7 Hz, 1H), 7.65 (t, J = 1.9 Hz, 1H), 7.60 – 7.54 (m, 2H), 7.48 (ddd, J = 8.7, 3.7, 2.0 Hz, 2H), 7.24 (t, J = 4.8 Hz, 1H), 6.




	264	484	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.17 (s, 1H), 10.06 (s, 1H), 8.83 (s, 1H), 8.57 – 8.34 (m, 2H), 8.26 (s, 1H), 7.83 (dd, J = 7.9, 1.8 Hz, 1H), 7.67 (t, J = 1.8 Hz, 1H), 7.62 – 7.33 (m, 5H), 7.24 (dd, J = 8.2, 1.4 Hz, 1H), 6.94 (t, J = 2.0 Hz, 1H), 3.06 (s, 3H)
	265	501	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.17 (s, 1H), 10.06 (s, 1H), 8.80 (s, 1H), 8.27 (s, 1H), 7.94 – 7.74 (m, 1H), 7.67 (s, 1H), 7.62 – 7.35 (m, 4H), 7.22 (d, J = 8.1 Hz, 1H), 7.09 – 6.76 (m, 4H), 3.06 (s, 3H).
	266	503	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.28 (s, 1H), 9.39 (s, 1H), 8.49 (d, J = 2.3 Hz, 1H), 8.29 (s, 1H), 8.09 – 7.87 (m, 2H), 7.69 (t, J = 1.9 Hz, 1H), 7.59 (t, J = 1.9 Hz, 1H), 6.92 (t, J = 1.9 Hz, 1H), 3.48 – 3.37 (m, 1H), 3.31 – 3.17 (m, 2H), 3.04 (s, 3H), 2.8



	267	477	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.13 (s, 1H), 10.07 (s, 1H), 8.65 (d, J = 3.0 Hz, 1H), 8.20 (d, J = 2.0 Hz, 1H), 8.15 – 8.02 (m, 1H), 7.83 (dd, J = 8.9, 3.8 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.53 (s, 1H), 6.94 (t, J = 2.0 Hz, 1H),
	268	520	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.04 (s, 1H), 9.81 (s, 1H), 8.44 (s, 1H), 8.29 (d, J = 2.3 Hz, 1H), 7.85 (dd, J = 10.9, 2.4 Hz, 1H), 7.74 – 7.65 (m, 2H), 7.59 (t, J = 1.9 Hz, 1H), 7.40 (s, 1H), 6.99 (d, J = 2.0 Hz, 1H), 6.88 (t, J = 2.0 Hz, 1H), 5.39 (s, 2H)
	269	504	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.99 (s, 1H), 9.78 (s, 1H), 8.14 (d, J = 2.9 Hz, 1H), 7.79 (d, J = 2.0 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.63 (t, J = 1.9 Hz, 1H), 7.45 – 7.28 (m, 2H), 6.88 (t, J = 1.9 Hz, 1H), 6.52 (dd, J = 2.0, 1.1 Hz, 1H), 4.65 (dq, J = 8.




	270	474	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.78 (s, 1H), 7.85 (d, J = 3.0 Hz, 1H), 7.76 (d, J = 2.0 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.8 Hz, 1H), 7.36 (d, J = 8.7 Hz, 1H), 7.09 (dd, J = 8.8, 3.1 Hz, 1H), 6.88 (t, J = 2.0 Hz, 1H), 6.50 (dd,
	271	544	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.07 (d, J = 10.3 Hz, 2H), 8.26 (d, J = 2.9 Hz, 1H), 8.07 (d, J = 2.0 Hz, 1H), 7.71 (d, J = 2.0 Hz, 1H), 7.63 – 7.52 (m, 2H), 7.50 (d, J = 8.8 Hz, 1H), 7.44 (d, J = 1.8 Hz, 1H), 6.93 (t, J = 2.0 Hz, 1H), 3.77 (t, J = 4.9 Hz, 4
	272	518	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.81 (s, 1H), 8.23 (d, J = 3.0 Hz, 1H), 7.81 (d, J = 2.0 Hz, 1H), 7.71 (s, 1H), 7.63 (s, 1H), 7.54 (dd, J = 8.9, 2.9 Hz, 1H), 7.43 (d, J = 8.8 Hz, 1H), 6.89 (s, 1H), 6.53 (s, 1H), 3.82 – 3.67 (m, 2H), 3.25 – 3.16




	273	490	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.80 (s, 1H), 8.33 (d, J = 2.9 Hz, 1H), 7.73 (t, J = 1.8 Hz, 1H), 7.65 – 7.60 (m, 2H), 7.56 (d, J = 8.8 Hz, 1H), 7.43 (dd, J = 8.9, 3.0 Hz, 1H), 7.04 (d, J = 2.0 Hz, 1H), 6.89 (t, J = 1.9 Hz, 1H), 3.93 (s, 3H), 3
	274	504	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.04 (s, 1H), 9.79 (s, 1H), 8.32 (d, J = 2.8 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.67 (d, J = 2.0 Hz, 1H), 7.63 (t, J = 1.9 Hz, 1H), 7.55 (d, J = 8.8 Hz, 1H), 7.43 (dd, J = 8.9, 2.9 Hz, 1H), 7.02 (d, J = 2.0 Hz, 1H), 6.89 (t, J
	275	558	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.04 (s, 1H), 9.94 (s, 1H), 8.30 (s, 1H), 7.76 – 7.67 (m, 2H), 7.62 (d, J = 7.7 Hz, 2H), 7.49 – 7.33 (m, 1H), 7.17 (s, 1H), 6.91 (s, 1H), 5.73 (d, J = 9.4 Hz, 2H), 3.76 (d, J = 5.6 Hz, 4H), 3.25 (d, J = 5.0 Hz, 4H), 3.07 (s, 3




	276	502	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.94 (s, 1H), 9.78 (s, 1H), 7.77 (dd, J = 5.1, 2.5 Hz, 2H), 7.69 (t, J = 1.9 Hz, 1H), 7.60 (d, J = 2.2 Hz, 1H), 7.38 (d, J = 8.6 Hz, 1H), 7.04 (dd, J = 8.6, 3.0 Hz, 1H), 6.87 (t, J = 2.0 Hz, 1H), 6.51 (s, 1H), 4.74 (s, 4H), 4.1
	277	502	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.96 (s, 1H), 9.75 (s, 1H), 7.86 (d, J = 2.8 Hz, 1H), 7.69 (s, 1H), 7.58 (d, J = 2.0 Hz, 2H), 7.49 (d, J = 8.6 Hz, 1H), 7.03 – 6.91 (m, 2H), 6.87 (t, J = 1.9 Hz, 1H), 4.74 (s, 4H), 4.09 (s, 4H), 3.90 (s, 3H), 3.04 (s, 3H).
	278	570	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.93 (s, 1H), 7.83 (d, J = 2.9 Hz, 1H), 7.70 (d, J = 4.7 Hz, 2H), 7.63 (s, 1H), 7.55 (d, J = 8.6 Hz, 1H), 7.10 (d, J = 1.9 Hz, 1H), 6.95 (dd, J = 8.7, 2.9 Hz, 1H), 6.91 (d, J = 2.1 Hz, 1H), 5.68 (q, J = 9.2 Hz, 2H), 4.73 (s, 4H)




	279	518	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.82 (s, 2H), 8.38 – 8.20 (m, 1H), 7.85 (d, J = 3.2 Hz, 1H), 7.78 (d, J = 3.2 Hz, 1H), 7.75 – 7.67 (m, 3H), 7.62 (d, J = 2.0 Hz, 1H), 7.35 (dd, J = 8.4, 4.6 Hz, 1H), 7.18 (d, J = 2.0 Hz, 1H), 6.89 (d, J = 2.0 Hz, 1H), 5.59 (s,
	280	518	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.77 (s, 1H), 8.22 (d, J = 4.7 Hz, 1H), 7.93 (d, J = 8.3 Hz, 1H), 7.83 (d, J = 3.2 Hz, 1H), 7.76 (d, J = 3.2 Hz, 1H), 7.70 (s, 2H), 7.62 – 7.52 (m, 2H), 6.88 (s, 1H), 6.54 (s, 1H), 5.59 (s, 2H), 3.04 (s, 3H), 2.1
	281	421	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.73 (s, 1H), 8.00 (d, J = 4.6 Hz, 1H), 7.67 (d, J = 2.1 Hz, 2H), 7.57 (d, J = 2.1 Hz, 1H), 7.46 (d, J = 8.2 Hz, 1H), 7.35 (dd, J = 8.2, 4.6 Hz, 1H), 6.86 (d, J = 2.0 Hz, 1H), 6.49 (s, 1H), 3.02 (s, 3H), 2.12 (s, 3H).




	282	396	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.96 (s, 1H), 9.58 (s, 1H), 7.66 (t, J = 1.9 Hz, 1H), 7.60 (t, J = 1.9 Hz, 1H), 7.55 (d, J = 2.0 Hz, 1H), 6.86 (t, J = 1.9 Hz, 1H), 6.35 (dd, J = 2.0, 1.0 Hz, 1H), 4.44 (p, J = 7.1 Hz, 1H), 3.05 (s, 3H), 2.30 – 2.00 (m, 5H), 1.
	283	559	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.82 (s, 1H), 8.27 (d, J = 3.0 Hz, 1H), 7.84 (s, 1H), 7.72 (s, 1H), 7.68 – 7.52 (m, 2H), 7.46 (d, J = 8.9 Hz, 1H), 6.89 (s, 1H), 6.54 (s, 1H), 3.66 (s, 4H), 3.27 (s, 4H), 3.06 (s, 3H), 2.94 (p, J = 6.6 Hz, 1H), 2




	284	560	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.81 (s, 1H), 8.26 (d, J = 2.9 Hz, 1H), 7.84 (d, J = 2.0 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.63 (t, J = 1.9 Hz, 1H), 7.57 (dd, J = 8.9, 3.0 Hz, 1H), 7.45 (d, J = 8.8 Hz, 1H), 6.89 (t, J = 2.0 Hz, 1H), 6.53 (dd,
	285	538	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.99 (s, 1H), 9.81 (s, 1H), 8.33 (d, J = 3.1 Hz, 1H), 7.84 (d, J = 2.0 Hz, 1H), 7.71 (q, J = 4.7, 3.3 Hz, 1H), 7.68 – 7.58 (m, 2H), 7.47 (d, J = 8.9 Hz, 1H), 6.89 (t, J = 2.0 Hz, 1H), 6.54 (s, 1H), 3.90 (t, J = 5.3 Hz, 4H), 3.1
	286	473	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.99 (d, J = 35.0 Hz, 2H), 8.59 (t, J = 1.8 Hz, 1H), 7.83 (dd, J = 7.2, 2.2 Hz, 2H), 7.77 (d, J = 1.9 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.30 (d, J = 1.9 Hz, 1H), 6.91 (t, J = 2.0 Hz, 1H), 6.56 – 6.16




	287	433	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.91 (s, 2H), 8.58 (dd, J = 5.0, 1.9 Hz, 1H), 8.06 (td, J = 7.8, 1.9 Hz, 1H), 8.01 (d, J = 2.0 Hz, 1H), 7.79 (d, J = 8.1 Hz, 1H), 7.71 (d, J = 1.9 Hz, 1H), 7.64 (d, J = 2.0 Hz, 1H), 7.47 (dd, J = 7.4, 4.9 Hz, 1H), 6.90 (t, J =
	288	382	¹ H NMR (300 MHz, Methanol-d ₄) δ 9.23 (s, 1H), 8.51 (d, J = 5.8 Hz, 1H), 8.37 (s, 1H), 8.12 (d, J = 14.5 Hz, 1H), 7.73 (d, J = 1.9 Hz, 1H), 7.63 (s, 1H), 7.08 (d, J = 2.3 Hz, 1H), 3.07 (s, 3H).
	289	382	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.81 (s, 1H), 10.13 (s, 1H), 8.79 (dd, J = 4.5, 1.5 Hz, 1H), 8.65 – 8.49 (m, 2H), 7.71 (dt, J = 13.7, 1.9 Hz, 2H), 7.51 (dd, J = 8.2, 4.5 Hz, 1H), 7.00 (t, J = 1.9 Hz, 1H), 3.09 (s, 3H).



	290	396	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.86 (s, 1H), 10.14 (s, 1H), 9.15 (s, 1H), 8.55 (s, 1H), 8.39 (d, J = 1.1 Hz, 1H), 7.69 (dt, J = 28.2, 1.9 Hz, 2H), 6.99 (t, J = 1.9 Hz, 1H), 3.09 (s, 3H), 2.54 (s, 3H).
	291	441	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.60 (s, 1H), 10.10 (s, 1H), 8.23 (s, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.66 (t, J = 1.9 Hz, 1H), 7.62 (s, 1H), 7.46 (s, 1H), 6.97 (t, J = 2.0 Hz, 1H), 3.86 (d, J = 2.4 Hz, 6H), 3.09 (s, 3H).
	292	365	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.24 (s, 1H), 10.07 (s, 1H), 8.50 (dd, J = 1.8, 0.7 Hz, 1H), 8.29 (dd, J = 4.4, 1.7 Hz, 1H), 8.08 (dd, J = 9.3, 1.7 Hz, 1H), 7.72 (dt, J = 19.2, 1.9 Hz, 2H), 7.11 (d, J = 1.8 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 6.77 (dd, J = 9.




	293	415	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.76 (s, 1H), 10.11 (s, 1H), 8.41 – 8.32 (m, 1H), 8.26 (d, J = 1.9 Hz, 1H), 8.05 (d, J = 8.5 Hz, 1H), 7.68 (dt, J = 20.7, 1.9 Hz, 2H), 7.52 (dd, J = 8.6, 2.0 Hz, 1H), 6.98 (t, J = 1.9 Hz, 1H), 3.08 (s, 3H).
	294	399	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.72 (s, 1H), 10.12 (s, 1H), 8.38 (s, 1H), 8.04 (ddd, J = 26.0, 9.0, 3.9 Hz, 2H), 7.67 (dt, J = 21.9, 2.0 Hz, 2H), 7.37 (td, J = 9.0, 2.5 Hz, 1H), 6.97 (d, J = 2.0 Hz, 1H), 3.08 (s, 3H).
	295	399	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.79 (s, 1H), 10.12 (s, 1H), 8.36 (s, 1H), 8.13 (dd, J = 8.9, 4.9 Hz, 1H), 7.89 (dd, J = 9.6, 2.6 Hz, 1H), 7.69 (dt, J = 19.0, 2.0 Hz, 2H), 7.41 (td, J = 9.0, 2.6 Hz, 1H), 6.98 (d, J = 2.0 Hz, 1H), 3.09 (s, 3H).




	296	397	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.75 (s, 1H), 10.22 (s, 1H), 8.52 (s, 1H), 7.93 (d, J = 8.2 Hz, 1H), 7.67 (d, J = 23.6 Hz, 2H), 7.56 – 7.41 (m, 1H), 7.31 (dd, J = 10.5, 7.9 Hz, 1H), 6.97 (t, J = 1.8 Hz, 1H), 3.06 (s, 3H).
	297	397	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.83 (d, J = 3.7 Hz, 1H), 10.14 (s, 1H), 8.48 (t, J = 3.9 Hz, 1H), 7.91 (dd, J = 8.1, 3.6 Hz, 1H), 7.69 (dq, J = 19.1, 2.3 Hz, 2H), 7.54 (dq, J = 8.3, 4.2 Hz, 1H), 7.42 (ddd, J = 11.3, 7.9, 3.5 Hz, 1H), 6.99 (p, J = 2.1 Hz, 1H)
	298	415	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.82 (s, 1H), 10.12 (s, 1H), 8.34 (s, 1H), 8.18 – 8.02 (m, 2H), 7.69 (dt, J = 19.5, 1.9 Hz, 2H), 7.54 (dd, J = 8.7, 2.1 Hz, 1H), 6.98 (t, J = 1.9 Hz, 1H), 3.09 (d, J = 2.4 Hz, 3H).




	299	398	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.23 (s, 1H), 10.05 (s, 1H), 8.63 (dd, J = 1.8, 1.0 Hz, 1H), 8.22 – 8.12 (m, 1H), 7.74 (d, J = 2.0 Hz, 1H), 7.68 (d, J = 2.1 Hz, 1H), 7.58 (d, J = 9.6 Hz, 1H), 7.10 (d, J = 1.5 Hz, 1H), 6.93 (q, J = 1.5, 0.9 Hz, 1H), 6.81 (dd,
	300	378	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.10 (s, 1H), 10.03 (s, 1H), 8.11 (d, J = 1.9 Hz, 1H), 8.06 (d, J = 1.7 Hz, 1H), 7.74 (t, J = 1.9 Hz, 1H), 7.68 (t, J = 1.9 Hz, 1H), 7.41 (d, J = 9.2 Hz, 1H), 6.92 (dd, J = 5.0, 3.0 Hz, 2H), 6.65 (dd, J = 9.3, 1.5 Hz, 1H), 3.0
	301	379	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.63 (d, J = 5.7 Hz, 1H), 10.14 (s, 2H), 8.04 (d, J = 1.7 Hz, 1H), 7.73 (d, J = 2.2 Hz, 1H), 7.65 (d, J = 1.8 Hz, 1H), 7.60 (s, 1H), 7.39 (d, J = 5.8 Hz, 1H), 6.92 (d, J = 1.9 Hz, 1H), 6.69 (t, J = 5.6 Hz, 1H), 3.05 (s, 3H).




	302	432	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.29 (s, 1H), 10.06 (s, 1H), 8.61 (d, J = 7.0 Hz, 1H), 8.37 (d, J = 1.6 Hz, 1H), 7.72 (dt, J = 17.5, 1.9 Hz, 2H), 7.31 (d, J = 6.9 Hz, 1H), 7.23 (s, 1H), 6.95 (t, J = 2.0 Hz, 1H), 6.81 (t, J = 7.0 Hz, 1H), 3.07 (s, 3H).
	303	411	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.72 (s, 1H), 10.12 (s, 1H), 8.38 (s, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.67 (t, J = 1.9 Hz, 1H), 7.61 (d, J = 7.9 Hz, 1H), 7.45 (t, J = 7.9 Hz, 1H), 7.08 (d, J = 7.8 Hz, 1H), 6.98 (t, J = 1.9 Hz, 1H), 3.99 (s, 3H), 3.09 (s, 3H).
	304	378	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.15 (s, 1H), 9.88 (s, 1H), 8.05 (s, 1H), 7.72 (dt, J = 17.4, 2.0 Hz, 2H), 7.45 (d, J = 9.0 Hz, 1H), 7.06 (d, J = 1.6 Hz, 1H), 6.94 (d, J = 2.2 Hz, 1H), 6.80 (dd, J = 9.1, 6.5 Hz, 1H), 6.60 (d, J = 6.7 Hz, 1H), 3.08 (d, J = 2.




	305	382	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.23 (s, 1H), 10.07 (d, J = 2.4 Hz, 1H), 8.31 (dt, J = 3.5, 1.9 Hz, 1H), 8.25 – 8.13 (m, 1H), 7.77 (t, J = 2.0 Hz, 1H), 7.69 (t, J = 2.0 Hz, 1H), 7.25 – 7.13 (m, 1H), 7.02 – 6.83 (m, 1H), 6.81 – 6.52 (m, 2H), 3.08 (d, J = 1.9
 *stereochemistry arbitrarily assigned*	306a	385	¹ H NMR (400 MHz, DMSO-d ₆) δ = 10.03 (s, 1H), 8.46 (s, 1H), 8.07 (s, 1H), 7.67 (t, J = 1.8 Hz, 1H), 7.61 – 7.51 (m, 1H), 6.92 (t, J = 1.8 Hz, 1H), 5.70 (br d, J = 2.6 Hz, 1H), 4.51 (q, J = 8.4 Hz, 1H), 4.32 (br d, J = 7.4 Hz, 1H), 3.05 (s, 3H), 2.12 (br t,



 *stereochemistry arbitrarily assigned*	306b	385	¹ H NMR (400 MHz, DMSO-d ₆) δ = 10.26 - 9.85 (m, 2H), 8.47 (d, J = 8.8 Hz, 1H), 8.05 (d, J = 18.8 Hz, 1H), 7.69 (q, J = 1.8 Hz, 1H), 7.58 (td, J = 1.8, 7.6 Hz, 1H), 6.92 (t, J = 1.8 Hz, 1H), 5.71 (br d, J = 6.8 Hz, 1H), 5.30 (d, J = 5.4 Hz, 1H), 4.90 (br d,
	307	426	¹ H NMR (400 MHz, DMSO-d ₆) δ = 10.07 (s, 1H), 8.43 (s, 1H), 8.08 (s, 1H), 7.76 - 7.49 (m, 2H), 6.92 (t, J = 1.8 Hz, 1H), 4.89 - 4.69 (m, 1H), 3.31 - 3.18 (m, 2H), 3.05 (s, 3H), 2.84 (s, 3H), 2.78 - 2.69 (m, 2H), 2.31 - 2.14 (m, 2H)
	308	419	¹ H NMR (400 MHz, DMSO-d ₆) δ = 9.99 (s, 2H), 8.25 (s, 1H), 8.06 (s, 1H), 7.67 (s, 1H), 7.57 (s, 1H), 7.36 - 7.08 (m, 5H), 6.92 (t, J = 1.8 Hz, 1H), 4.41 (t, J = 7.2 Hz, 2H), 3.14 (t, J = 7.2 Hz, 2H), 3.06 (s, 3H)



	309	383	¹ H NMR (400 MHz, DMSO-d ₆) δ = 9.98 (s, 1 H), δ = 8.40 (s, 1 H), δ = 8.02 (s, 1 H), δ = 7.65 (t, J=1.7 Hz, 1 H), δ = 7.56 (t, J=1.7 Hz, 1 H), δ = 6.91 (t, J=1.8 Hz, 1 H), δ = 4.75 (quin, J=7.0 Hz, 1 H), δ = 3.05 (s, 3 H), δ = 2.04 - 2.18 (m, 2 H), δ = 1.86 -
	310	466	¹ H NMR (400 MHz, DMSO-d ₆) δ = 10.01 (s, 2H), 8.40 (s, 1H), 8.11 - 8.01 (m, 2H), 7.77 - 7.45 (m, 2H), 6.92 (t, J = 2.0 Hz, 1H), 4.82 (t, J = 8.4 Hz, 1H), 4.28 - 4.00 (m, 1H), 3.06 (s, 3H), 2.65 - 2.55 (m, 2H), 2.49 - 2.37 (m, 3H), 2.26 (br s, 1H), 2.06 - 1
	311	424	¹ H NMR (400 MHz, METHANOL-d ₄) δ = 8.53 (s, 1 H), δ = 8.25 (br s, 1 H), δ = 8.08 (s, 1 H), δ = 7.63 (br s, 1 H), δ = 7.53 (s, 1 H), δ = 6.99 (s, 1 H), δ = 4.76 - 4.86 (m, 1 H), δ = 4.24 (br d, J=14.2 Hz, 4 H), δ = 3.04 (s, 3 H), δ = 2.74 - 2.99 (m, 7 H)



	312	492	¹ H NMR (400 MHz, DMSO-d ₆) δ = 10.03 (s, 1 H), δ = 8.41 (s, 1 H), δ = 8.05 (s, 1 H), δ = 7.65 (t, J=1.8 Hz, 1 H), δ = 7.56 (t, J=1.8 Hz, 1 H), δ = 6.91 (t, J=1.9 Hz, 1 H), δ = 4.81 (t, J=8.1 Hz, 1 H), δ = 3.45 (s, 2 H), δ = 3.35 (s, 2 H), δ = 3.10 - 3.18 (m)
	313	452	¹ H NMR (400 MHz, DMSO-d ₆) δ = 9.96 (s, 1 H), δ = 8.42 (d, J=2.6 Hz, 1 H), δ = 8.06 (s, 1 H), δ = 7.59 (s, 1 H), δ = 7.48 (s, 1 H), δ = 6.86 (t, J=1.8 Hz, 1 H), δ = 4.85 (t, J=8.1 Hz, 1 H), δ = 4.08 - 4.26 (m, 2 H), δ = 3.76 - 4.01 (m, 2 H), δ = 2.99 (s, 3)
	314	454	¹ H NMR (400 MHz, DMSO-d ₆) δ = 10.46 (s, 1 H), δ = 9.91 - 10.23 (m, 1 H), δ = 8.49 (s, 1 H), δ = 8.22 (s, 1 H), δ = 7.75 (d, J=7.6 Hz, 2 H), δ = 7.49 (t, J=7.6 Hz, 2 H), δ = 7.32 - 7.41 (m, 1 H), δ = 3.07 (s, 3 H)



	315	426	¹ H NMR (400 MHz, DMSO-d ₆) δ = 10.08 (d, J = 5.6 Hz, 2H), 8.96 (br d, J = 10.0 Hz, 1H), 8.68 (br t, J = 9.6 Hz, 1H), 8.44 (s, 1H), 8.11 (s, 1H), 7.68 (t, J = 1.8 Hz, 1H), 7.58 (t, J = 1.8 Hz, 1H), 6.93 (t, J = 1.8 Hz, 1H), 4.75 (br t, J = 4.0 Hz, 1H), 3.38
	316	480	¹ H NMR (400 MHz, DMSO-d ₆) δ = 9.91 - 10.22 (m, 2 H), δ = 8.44 (s, 1 H), δ = 8.06 (s, 1 H), δ = 7.67 (t, J=1.8 Hz, 1 H), δ = 7.58 (s, 1 H), δ = 6.92 (t, J=1.8 Hz, 1 H), δ = 4.29 - 4.33 (m, 1 H), δ = 3.27 - 3.50 (m, 2 H), δ = 2.94 - 3.19 (m, 5 H), δ = 2.60
	317	440	¹ H NMR (400 MHz, DMSO-d ₆) δ = 10.00 (s, 2 H), δ = 8.42 (s, 1 H), δ = 8.05 (s, 1 H), δ = 7.66 (s, 1 H), δ = 7.56 (s, 1 H), δ = 6.91 (s, 1 H), δ = 4.43 - 4.61 (m, 2 H), δ = 3.92 (br d, J=14.1 Hz, 1 H), δ = 3.17 - 3.25 (m, 1 H), δ = 3.05 (s, 3 H), δ = 2.65 -



	318	468	¹ H NMR (400 MHz, DMSO-d ₆) δ = 10.01 (s, 1H), 8.45 - 8.40 (m, 1H), 8.05 (s, 1H), 7.79 - 7.40 (m, 2H), 7.11 (br d, J = 8.0 Hz, 1H), 6.90 (t, J = 1.8 Hz, 1H), 4.80 (t, J = 8.2 Hz, 1H), 3.87 (br d, J = 8.2 Hz, 1H), 3.02 (s, 3H), 2.49 - 2.30 (m, 5H), 2.27 - 2.
	319	424	¹ H NMR (400 MHz, DMSO-d ₆) δ = 10.08 (s, 1H), 8.43 (s, 1H), 8.06 (s, 1H), 7.73 - 7.47 (m, 2H), 6.91 (s, 1H), 4.81 (br t, J = 8.2 Hz, 1H), 3.53 (br t, J = 7.8 Hz, 1H), 3.04 (s, 3H), 2.59 (td, J = 3.8, 7.4 Hz, 1H), 2.50 - 2.38 (m, 4H), 2.34 - 2.24 (m, 1H), 2.
	320	508	¹ H NMR (300 MHz, DMSO-d ₆) δ = 9.99 (br s, 1H), 9.82 (s, 1H), 8.15 (d, J = 2.1 Hz, 1H), 7.55-7.71 (m, 4H), 6.89 (t, J = 2.0 Hz, 1H), 6.56 (d, J = 0.6 Hz), 3.77 (m, 4H), 3.32 (m, 4H), 3.06 (s, 3H), 2.12 (s, 3H)



	643	515.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.96 (s, 1H), 8.50 (dd, J = 5.1, 1.8 Hz, 1H), 8.13 (d, J = 1.8 Hz, 1H), 7.96 (td, J = 7.8, 1.9 Hz, 1H), 7.57 (d, J = 1.9 Hz, 1H), 7.53 – 7.39 (m, 3H), 7.18 (d, J = 1.8 Hz, 1H), 6.86 (d, J = 2.0 Hz, 1H), 4.44 (s, 2H), 4.06 (s, 2H), 3.72 (s, 4H), 2.97 (s, 3H).
	644	502	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.78 (s, 1H), 8.49 (d, J = 4.9 Hz, 1H), 8.21 (d, J = 1.9 Hz, 1H), 8.12 – 7.91 (m, 1H), 7.69 (q, J = 1.8 Hz, 1H), 7.64 – 7.49 (m, 2H), 7.42 (dd, J = 7.3, 5.1 Hz, 1H), 6.93 (dt, J = 18.8, 1.8 Hz, 2H), 3.47 (s, 4H), 3.04 (d, J = 1.6 Hz, 3H), 1.59 (s, 2H), 1.46 (s, 4H).




	645	565	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.97 (d, J = 12.8 Hz, 2H), 8.91 (d, J = 0.8 Hz, 2H), 8.19 (d, J = 2.0 Hz, 1H), 7.82 – 7.43 (m, 3H), 6.92 (t, J = 2.0 Hz, 1H), 5.85 (t, J = 6.5 Hz, 1H), 4.40 (t, J = 8.6 Hz, 2H), 4.11 (dd, J = 9.5, 5.5 Hz, 2H), 3.06 (s, 3H), 1.42 (s, 9H).
	646	510.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.55 (s, 1H), 9.85 (s, 1H), 8.27 (dd, J = 2.4, 1.1 Hz, 1H), 8.16 (s, 1H), 7.74 – 7.68 (m, 2H), 7.67 – 7.57 (m, 2H), 7.07 (dd, J = 3.2, 2.0 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H), 4.21 (t, J = 5.7 Hz, 2H), 3.84 (s, 3H), 3.06 (s, 3H), 2.68 (t, J = 5.6 Hz, 2H), 2.25 (s, 6H).




	647	498.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.48 (s, 1H), 10.12 (s, 1H), 9.19 – 8.77 (m, 1H), 8.45 (s, 1H), 8.27 (ddd, J = 10.5, 8.1, 2.1 Hz, 1H), 7.84 (dt, J = 7.9, 1.4 Hz, 1H), 7.65 (dt, J = 16.7, 1.9 Hz, 2H), 6.95 (t, J = 1.9 Hz, 1H), 3.06 (s, 3H), 2.75 (s, 3H), 1.76 (d, J = 13.5 Hz, 6H).
	648	496.9 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.65 (s, 1H), 10.07 (s, 1H), 9.19 (d, J = 5.7 Hz, 2H), 8.78 (s, 1H), 7.69 (dt, J = 7.9, 1.9 Hz, 2H), 6.96 (t, J = 1.9 Hz, 1H), 3.08 (s, 3H), 2.91 (s, 3H), 1.81 (d, J = 13.8 Hz, 6H).



	649	582.9 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.49(s,1H),8.50 (dd, J = 5.0, 1.8 Hz, 1H), 8.24 (d, J = 1.8 Hz, 1H), 8.02 (td, J = 7.8, 1.9 Hz, 1H), 7.68 (t, J = 1.9 Hz, 1H), 7.62 – 7.53 (m, 2H), 7.43 (dd, J = 7.4, 4.9 Hz, 1H), 6.99 (d, J = 1.8 Hz, 1H), 6.91 (t, J = 2.0 Hz, 1H), 3.53 (s, 4H), 3.24 (q, J = 10.2 Hz, 2H), 3.04 (s, 3H), 2.62 (s, 4H).
	650	447.15 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.99 (s, 1H), 9.86 (s, 1H), 8.59 (ddd, J = 4.9, 1.9, 0.8 Hz, 1H), 8.10 (td, J = 7.8, 1.9 Hz, 1H), 7.87 (d, J = 2.0 Hz, 1H), 7.79 (dt, J = 8.2, 1.0 Hz, 1H), 7.68 (t, J = 1.9 Hz, 1H), 7.61 (t, J = 1.9 Hz, 1H), 7.53 (ddd, J = 7.5, 4.9, 1.0 Hz, 1H), 6.90 (t, J = 2.0 Hz, 1H), 6.76 (d, J = 2.0 Hz, 1H), 6.28 (s, 1H), 3.05 (s, 3H), 1.37 (s, 6H).



	651	623.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.07 (s, 2H), 8.55 (s, 2H), 7.81 (s, 1H), 7.76 – 7.61 (m, 3H), 6.92 (s, 1H), 5.75 (t, J = 9.1 Hz, 2H), 5.26 (t, J = 5.7 Hz, 1H), 4.38 (dd, J = 9.5, 6.6 Hz, 2H), 4.03 (dd, J = 9.4, 4.6 Hz, 2H), 3.08 (d, J = 12.7 Hz, 6H), .
	652	464.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.25 – 9.75 (m, 2H), 7.72 (d, J = 1.8 Hz, 1H), 7.64 (s, 1H), 7.54 (s, 1H), 7.21 (d, J = 1.8 Hz, 1H), 6.85 (s, 1H), 6.63 (d, J = 5.0 Hz, 1H), 6.54 (s, 1H), 5.46 (s, 1H), 4.65 (d, J = 11.3 Hz, 1H), 4.20 (d, J = 13.0 Hz, 1H), 3.85 (d, J = 10.1 Hz, 1H), 3.51 (s, 1H), 3.00 (s, 4H).



	653	553.1	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.50 (d, J = 5.0 Hz, 1H), 8.46 (s, 1H), 7.99 (d, J = 1.8 Hz, 1H), 7.79 (t, J = 5.4 Hz, 1H), 7.65 (t, J = 1.9 Hz, 1H), 7.55 (t, J = 1.9 Hz, 1H), 7.01 (t, J = 1.9 Hz, 1H), 4.32 (d, J = 24.8 Hz, 2H), 3.91 (s, 2H), 3.46 (s, 1H), 3.03 (s, 3H), 2.88 (s, 3H), 2.53 (d, J = 1.3 Hz, 3H), 1.65 – 1.61 (m, 3H).
	654	579	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.01 (d, J = 21.2 Hz, 2H), 8.47 (dd, J = 5.3, 1.8 Hz, 1H), 8.25 (d, J = 1.8 Hz, 1H), 8.01 (td, J = 7.8, 1.9 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.63 (t, J = 1.9 Hz, 1H), 7.58 (d, J = 8.2 Hz, 1H), 7.40 (dd, J = 7.5, 4.9 Hz, 1H), 7.28 – 7.21 (m, 2H), 7.05 (d, J = 1.8 Hz, 1H), 6.99 – 6.91 (m, 3H), 6.82 (t, J = 7.3 Hz, 1H), 3.68 (s, 4H), 3.14 (q, J = 10.5, 9.0 Hz, 4H), 3.06 (s, 3H).



	655	497.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.82 (s, 1H), 8.54 – 8.48 (m, 1H), 8.38 (d, J = 15.5 Hz, 2H), 7.69 (s, 2H), 7.61 – 7.51 (m, 2H), 7.43 (s, 3H), 7.15 (d, J = 2.0 Hz, 1H), 6.88 (s, 1H), 3.87 (s, 3H), 3.05 (d, J = 6.7 Hz, 3H).
	656	483.85	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.13 (s, 1H), 10.03 (s, 1H), 9.29 (s, 2H), 8.61 (d, J = 2.1 Hz, 1H), 7.72 (t, J = 1.8 Hz, 1H), 7.65 (t, J = 2.0 Hz, 1H), 6.92 (t, J = 2.0 Hz, 1H), 6.64 (dd, J = 2.1, 1.2 Hz, 1H), 3.44 (s, 3H), 3.06 (s, 3H), 2.64 (d, J = 1.1 Hz, 3H).
	657	467.9	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.12 (s, 2H), 8.57 (d, J = 2.1 Hz, 1H), 7.58 (d, J = 27.8 Hz, 2H), 6.85 (s, 1H), 6.62 (d, J = 1.4 Hz, 1H), 3.00 (d, J = 10.1 Hz, 6H), 2.63 (d, J = 1.1 Hz, 3H).




	658	451.95	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.99 (d, J = 14.1 Hz, 2H), 8.82 (s, 2H), 8.44 (d, J = 2.1 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.65 (t, J = 1.9 Hz, 1H), 6.90 (t, J = 2.0 Hz, 1H), 6.58 (dd, J = 2.1, 1.1 Hz, 1H), 3.06 (s, 3H), 2.63 (s, 3H), 2.58 (d, J = 1.1 Hz, 3H).
	659	460.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.83 (s, 1H), 7.84 (dd, J = 8.3, 2.8 Hz, 1H), 7.75 (d, J = 1.8 Hz, 1H), 7.71 (s, 1H), 7.67 (d, J = 8.8 Hz, 1H), 7.61 (s, 1H), 6.89 (t, J = 1.9 Hz, 1H), 6.74 (d, J = 1.9 Hz, 1H), 3.41 (s, 3H), 3.05 (s, 3H), 2.16 (s, 3H).
	660	547.9	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.33 (s, 1H), 10.10 (s, 1H), 8.64 (ddd, J = 4.8, 1.9, 0.9 Hz, 1H), 7.83 (td, J = 7.8, 1.8 Hz, 1H), 7.54 (dt, J = 9.8, 1.9 Hz, 2H), 7.50 – 7.37 (m, 3H), 6.95 (t, J = 2.0 Hz, 1H), 3.07 (d, J = 8.2 Hz, 6H), 2.01 (s, 3H), 1.92 (s, 3H).



	661	441.15 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.26 (s, 1H), 10.07 (s, 1H), 7.91 (d, J = 1.8 Hz, 1H), 7.61 (dt, J = 6.7, 1.9 Hz, 2H), 6.95 (t, J = 1.9 Hz, 1H), 6.84 (d, J = 1.7 Hz, 1H), 3.72 (s, 2H), 3.49 – 3.37 (m, 4H), 3.06 (s, 3H), 2.90 (s, 3H).
	662	495.15	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.83 (s, 1H), 8.18 (d, J = 2.4 Hz, 1H), 7.83 (dd, J = 11.3, 2.4 Hz, 1H), 7.73 (dt, J = 10.3, 1.8 Hz, 2H), 7.62 (t, J = 1.9 Hz, 1H), 6.90 (t, J = 2.0 Hz, 1H), 6.65 – 6.49 (m, 1H), 3.06 (s, 3H), 2.16 (s, 3H), 1.41 (s, 9H).


	663	477.1	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.83 (s, 1H), 8.24 (d, J = 2.8 Hz, 1H), 7.93 (d, J = 2.0 Hz, 1H), 7.72 (t, J = 1.8 Hz, 1H), 7.70 – 7.60 (m, 2H), 7.55 (d, J = 8.7 Hz, 1H), 6.89 (t, J = 1.9 Hz, 1H), 6.56 (d, J = 2.0 Hz, 1H), 3.06 (s, 3H), 2.47 – 2.30 (m, 3H), 1.36 (s, 9H).
	664	478.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 2H), 8.61 (s, 2H), 8.42 (d, J = 2.1 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.65 (t, J = 1.9 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H), 6.66 – 6.43 (m, 1H), 3.06 (s, 3H), 2.68 – 2.54 (m, 3H), 1.37 (s, 9H).



	665	516.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.01 (d, J = 3.3 Hz, 1H), 9.89 (s, 1H), 8.49 (dd, J = 5.1, 1.8 Hz, 1H), 8.22 (t, J = 2.2 Hz, 1H), 8.01 (td, J = 7.8, 1.9 Hz, 1H), 7.69 (q, J = 2.1 Hz, 1H), 7.61 (q, J = 1.8 Hz, 1H), 7.54 (d, J = 8.1 Hz, 1H), 7.43 (dd, J = 7.5, 4.8 Hz, 1H), 6.99 (d, J = 1.8 Hz, 1H), 6.91 (p, J = 2.2 Hz, 1H), 3.49 (s, 4H), 3.05 (d, J = 1.4 Hz, 3H), 2.26 (s, 4H), 2.18 (s, 3H).
	666	515.85 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.75 (s, 2H), 8.75 (dd, J = 3.0, 1.3 Hz, 1H), 8.61 – 8.53 (m, 1H), 8.34 (ddd, J = 8.8, 3.0, 1.2 Hz, 1H), 8.24 – 8.18 (m, 1H), 7.65 (q, J = 1.7 Hz, 1H), 7.57 (dq, J = 9.7, 1.6 Hz, 2H), 7.52 (t, J = 1.7 Hz, 1H), 6.87 (q, J = 1.8 Hz, 1H), 6.49 (d, J = 1.9 Hz, 1H), 3.03 (d, J = 1.2 Hz, 3H), 1.89 (s, 3H).



	667	451.95	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.99 (s, 1H), 9.91 (s, 1H), 8.91 (d, J = 0.9 Hz, 2H), 7.93 (d, J = 2.1 Hz, 1H), 7.79 – 7.52 (m, 3H), 6.90 (t, J = 1.9 Hz, 1H), 5.76 (p, J = 6.6 Hz, 1H), 3.06 (s, 3H), 1.46 (d, J = 6.7 Hz, 6H).
	668	477	¹ H NMR (300 MHz, Methanol-d ₄) δ 8.11 (dd, J = 2.8, 0.9 Hz, 1H), 7.70 (d, J = 2.1 Hz, 1H), 7.65 (t, J = 1.9 Hz, 1H), 7.54 (t, J = 1.9 Hz, 1H), 7.50 – 7.33 (m, 2H), 6.98 (t, J = 1.9 Hz, 1H), 6.54 (dd, J = 2.1, 1.1 Hz, 1H), 5.41 (tt, J = 5.9, 4.8 Hz, 1H), 5.06 (ddd, J = 7.0, 5.9, 1.0 Hz, 2H), 4.74 (ddd, J = 7.4, 4.8, 1.0 Hz, 2H), 3.03 (s, 3H), 2.31 (d, J = 0.9 Hz, 3H).



	669	478.15	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.97 (s, 2H), 8.53 (s, 2H), 8.35 (d, J = 2.1 Hz, 1H), 7.72 (t, J = 1.8 Hz, 1H), 7.65 (t, J = 1.9 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H), 6.56 (dd, J = 2.1, 1.1 Hz, 1H), 5.51 (tt, J = 5.9, 4.7 Hz, 1H), 4.99 (ddd, J = 7.1, 5.9, 1.0 Hz, 2H), 4.63 (ddd, J = 7.5, 4.7, 1.0 Hz, 2H), 3.06 (s, 3H), 2.55 (d, J = 1.0 Hz, 3H).
	670	525.05	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.48 (d, J = 3.0 Hz, 3H), 7.65 (t, J = 2.0 Hz, 1H), 7.56 (t, J = 1.9 Hz, 1H), 7.02 (t, J = 2.0 Hz, 1H), 3.04 (s, 3H), 2.86 (s, 3H), 1.67 (s, 6H).
	671	511.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.55 (s, 1H), 10.05 (s, 1H), 8.63 (s, 1H), 8.56 (s, 2H), 7.68 (dt, J = 6.0, 1.8 Hz, 2H), 6.94 (t, J = 2.0 Hz, 1H), 5.06 (s, 1H), 3.07 (s, 3H), 2.82 (s, 3H), 1.54 (d, J = 6.7 Hz, 3H).




	672	554.15	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.99 (s, 1H), 9.74 (s, 1H), 7.70 (q, J = 1.7 Hz, 1H), 7.63 (t, J = 1.9 Hz, 1H), 7.51 (t, J = 1.7 Hz, 1H), 7.47 (d, J = 2.1 Hz, 1H), 6.88 (q, J = 1.7 Hz, 2H), δ 6.46 (s, 1H), 4.67 (hept, J = 6.0 Hz, 1H), 4.44 (t, J = 12.3 Hz, 4H), 3.06 (d, J = 1.3 Hz, 3H), 2.04 (s, 3H), 1.20 (dd, J = 6.1, 1.3 Hz, 6H).
	673	566.35	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.99 (s, 1H), 9.75 (s, 1H), 7.70 (s, 1H), 7.62 (s, 1H), 7.53 (d, J = 2.0 Hz, 1H), 7.47 (d, J = 2.4 Hz, 1H), 6.88 (t, J = 2.0 Hz, 1H), 6.64 (d, J = 2.4 Hz, 1H), 6.48 (s, 1H), 4.86 – 4.72 (m, 1H), 4.44 (t, J = 12.3 Hz, 4H), 3.05 (s, 3H), 2.42 (d, J = 8.7 Hz, 2H), 2.05 (d, J = 1.0 Hz, 3H), 1.99 – 1.86 (m, 2H), 1.76 (d, J = 10.1 Hz, 1H), 1.61 (d, J = 9.6 Hz, 1H).



	674	504.1	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.04 (s, 1H), 9.79 (s, 1H), 8.14 (d, J = 3.0 Hz, 1H), 7.80 (d, J = 2.0 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.62 (d, J = 2.0 Hz, 1H), 7.48 – 7.34 (m, 2H), 6.88 (t, J = 2.0 Hz, 1H), 6.52 (s, 1H), 4.66 (s, 1H), 3.97 (td, J = 8.4, 5.0 Hz, 1H), 3.86 – 3.71 (m, 2H), 3.65 (q, J = 8.0 Hz, 1H), 3.05 (s, 3H), 2.87 (s, 3H), 2.35 – 2.18 (m, 4H), 1.95 – 1.77 (m, 1H).
---	-----	-------	---



	675	504.1	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.80 (s, 1H), 8.14 (d, J = 2.7 Hz, 1H), 7.80 (d, J = 2.0 Hz, 1H), 7.76 – 7.68 (m, 1H), 7.63 (t, J = 1.9 Hz, 1H), 7.47 – 7.34 (m, 2H), 6.89 (t, J = 1.9 Hz, 1H), 6.56 – 6.49 (m, 1H), 4.66 (t, J = 7.3 Hz, 1H), 3.97 (td, J = 8.4, 5.1 Hz, 1H), 3.86 – 3.71 (m, 2H), 3.65 (q, J = 8.0 Hz, 1H), 3.06 (s, 3H), 2.87 (s, 3H), 2.35 – 2.17 (m, 4H), 1.94 – 1.77 (m, 1H).
	676	485.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.24 (s, 1H), 9.06 (d, J = 0.8 Hz, 1H), 8.48 (dd, J = 4.6, 1.4 Hz, 1H), 8.44 – 8.32 (m, 2H), 8.19 (d, J = 0.7 Hz, 1H), 7.85 (dd, J = 8.2, 1.5 Hz, 1H), 7.69 (t, J = 1.9 Hz, 1H), 7.65 – 7.57 (m, 2H), 7.50 – 7.35 (m, 2H), 6.94 (t, J = 1.9 Hz, 1H), 3.06 (s, 3H).




	677	497	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.57 (s, 1H), 10.06 (s, 1H), 8.65 (d, J = 6.9 Hz, 3H), 7.69 (dt, J = 6.4, 1.9 Hz, 2H), 6.94 (t, J = 2.0 Hz, 1H), 4.98 (t, J = 5.7 Hz, 1H), 4.71 (p, J = 5.8 Hz, 1H), 3.58 (t, J = 5.4 Hz, 2H), 3.07 (s, 3H), 2.83 (s, 3H), 1.27 (d, J = 6.2 Hz, 3H).
	678	510.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.56 (s, 1H), 10.08 (s, 1H), 8.65 (d, J = 9.1 Hz, 3H), 7.68 (dt, J = 7.4, 1.9 Hz, 2H), 6.94 (t, J = 1.8 Hz, 1H), 4.74 (s, 1H), 3.97 (s, 2H), 3.06 (s, 3H), 2.83 (s, 3H), 1.23 (s, 6H).



	679	541.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 8.48 (s, 2H), 7.77 – 7.70 (m, 2H), 7.67 (t, J = 1.9 Hz, 1H), 7.64 (d, J = 2.1 Hz, 1H), 6.91 (t, J = 2.0 Hz, 1H), 6.55 – 6.20 (m, 1H), 5.10 (td, J = 14.5, 3.9 Hz, 2H), 4.98 (p, J = 5.5 Hz, 1H), 3.81 – 3.73 (m, 2H), 3.10 – 2.99 (m, 5H), 2.30 (s, 3H).
	680	497.15	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.64 (s, 1H), 10.07 (s, 1H), 9.11 (d, J = 1.1 Hz, 2H), 8.76 (s, 1H), 7.69 (dt, J = 8.2, 1.9 Hz, 2H), 6.95 (t, J = 1.9 Hz, 1H), 5.17 – 4.96 (m, 4H), 3.07 (s, 3H), 2.90 (s, 3H).



	681	525.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.46 (s, 1H), 10.07 (s, 1H), 8.51 (d, J = 4.8 Hz, 1H), 8.22 (d, J = 1.7 Hz, 1H), 7.67 (t, J = 1.9 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 7.58 – 7.51 (m, 1H), 6.95 (t, J = 1.9 Hz, 1H), 6.27 (s, 1H), 3.78 – 3.72 (m, 2H), 3.38 (d, J = 7.9 Hz, 2H), 3.06 (s, 3H), 2.54 (d, J = 1.2 Hz, 3H), 2.31 (s, 3H).
	682	451.05 [M-H] ⁻	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.92 (s, 2H), 8.44 (d, J = 4.7 Hz, 1H), 8.06 (t, J = 9.3 Hz, 1H), 7.90 (t, J = 2.1 Hz, 1H), 7.69 (s, 1H), 7.66 – 7.59 (m, 2H), 6.90 (d, J = 1.8 Hz, 2H), 4.48 (s, 2H), 3.03 (d, J = 4.1 Hz, 6H).
	683	499.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.17 (d, J = 2.0 Hz, 1H), 10.06 (s, 1H), 8.68 (s, 2H), 7.59 (dt, J = 4.2, 1.9 Hz, 2H), 6.96 (t, J = 2.0 Hz, 1H), 4.89 (p, J = 6.0 Hz, 1H), 3.07 (s, 3H), 2.65 (s, 3H), 1.35 (d, J = 6.0 Hz, 6H).



	684	436.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.38 (s, 1H), 10.06 (s, 1H), 8.49 (dd, J = 4.7, 1.6 Hz, 1H), 7.76 (d, J = 9.2 Hz, 2H), 7.60 (dt, J = 15.0, 1.8 Hz, 2H), 7.33 (dd, J = 7.7, 4.8 Hz, 1H), 6.96 (t, J = 1.9 Hz, 1H), 3.07 (s, 3H), 2.20 (s, 6H).
	685	547.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.51 (s, 1H), 10.04 (s, 1H), 8.46 (dd, J = 4.9, 1.9 Hz, 1H), 8.12 – 8.05 (m, 1H), 8.02 (s, 1H), 7.90 (tt, J = 8.5, 1.5 Hz, 1H), 7.67 (q, J = 1.6 Hz, 1H), 7.61 (q, J = 1.7 Hz, 1H), 7.27 – 7.20 (m, 1H), 6.95 (q, J = 1.7 Hz, 1H), 3.06 (d, J = 1.0 Hz, 3H), 2.41 – 2.35 (m, 3H), 1.87 – 1.80 (m, 6H).



	686	510.95	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.58 (s, 1H), 10.08 (s, 1H), 8.71 (s, 1H), 8.64 (s, 2H), 7.70 (dt, J = 5.6, 1.9 Hz, 2H), 6.95 (t, J = 2.0 Hz, 1H), 6.29 – 5.82 (m, 1H), 5.13 (d, J = 5.5 Hz, 2H), 3.08 (s, 3H), 1.38 (s, 9H).
	687	463.9	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.40 (s, 1H), 10.06 (s, 1H), 8.37 (d, J = 2.9 Hz, 1H), 8.23 (dd, J = 4.6, 1.3 Hz, 1H), 7.98 (s, 1H), 7.67 (t, J = 1.8 Hz, 1H), 7.60 (t, J = 1.9 Hz, 1H), 7.55 – 7.50 (m, 1H), 7.39 (dd, J = 8.4, 4.5 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 5.92 (d, J = 5.7 Hz, 1H), 3.19 – 3.10 (m, 1H), 3.06 (s, 3H), 3.03 – 2.95 (m, 2H), 2.43 (d, J = 10.6 Hz, 1H).



	688	492.1	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.79 (s, 1H), 7.95 – 7.87 (m, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.65 – 7.57 (m, 2H), 7.52 (d, J = 8.6 Hz, 1H), 6.99 (q, J = 3.2 Hz, 2H), 6.89 (t, J = 1.9 Hz, 1H), 4.14 – 3.94 (m, 4H), 3.90 (s, 3H), 3.06 (s, 3H), 1.65 (d, J = 22.1 Hz, 3H).
	689	453.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.85 (s, 1H), 8.57 (d, J = 2.4 Hz, 1H), 7.83 – 7.76 (m, 2H), 7.75 – 7.69 (m, 2H), 7.63 (t, J = 2.0 Hz, 1H), 7.18 (d, J = 2.0 Hz, 1H), 6.90 (t, J = 2.0 Hz, 1H), 4.89 (t, J = 5.2 Hz, 1H), 4.47 (t, J = 5.6 Hz, 2H), 3.64 (q, J = 5.5 Hz, 2H), 3.06 (s, 3H).
	690	450	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.98 (d, J = 29.4 Hz, 2H), 8.76 (s, 2H), 7.95 – 7.47 (m, 4H), 6.90 (t, J = 2.0 Hz, 1H), 4.46 (s, 2H), 4.10 (s, 3H), 3.35 (s, 3H), 3.06 (s, 3H).



	691	505.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.72 (s, 1H), 10.25 (s, 1H), 10.09 (s, 1H), 9.06 (s, 1H), 8.28 (s, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.62 (t, J = 1.9 Hz, 1H), 7.30 (d, J = 2.4 Hz, 1H), 6.95 (t, J = 2.0 Hz, 1H), 6.54 (d, J = 2.4 Hz, 1H), 4.74 (s, 4H), 4.10 (s, 4H), 3.07 (s, 3H).
	692	496.25	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.51 (s, 1H), 10.07 (s, 1H), 8.52 (s, 1H), 8.42 (s, 1H), 7.68 (dt, J = 8.6, 1.9 Hz, 2H), 6.95 (t, J = 1.9 Hz, 1H), 4.85 (p, J = 8.0, 7.5 Hz, 1H), 4.06 – 3.95 (m, 2H), 3.54 (td, J = 11.3, 5.4 Hz, 2H), 3.08 (s, 3H), 2.64 (s, 3H), 2.18 – 2.02 (m, 4H).



	693	504.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.22 (s, 1H), 10.04 (s, 1H), 8.69 (s, 2H), 8.58 (d, J = 2.1 Hz, 1H), 7.71 (t, J = 1.9 Hz, 1H), 7.63 (t, J = 1.9 Hz, 1H), 7.48 (d, J = 2.2 Hz, 1H), 6.94 (t, J = 2.0 Hz, 1H), 4.28 (q, J = 6.9 Hz, 2H), 3.06 (s, 3H), 1.39 (t, J = 6.9 Hz, 3H).
	694	529.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.24 (d, J = 2.1 Hz, 1H), 8.19 (s, 2H), 7.68 (dt, J = 8.5, 1.9 Hz, 2H), 6.92 (t, J = 1.9 Hz, 1H), 6.54 (t, J = 1.7 Hz, 1H), 4.24 (t, J = 8.6 Hz, 2H), 4.07 (dd, J = 8.6, 5.1 Hz, 2H), 3.74 (dt, J = 9.4, 4.8 Hz, 1H), 3.06 (s, 3H), 2.50 (s, 3H).




	695	552.15	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.78 (s, 1H), 7.85 (d, J = 2.9 Hz, 1H), 7.72 (t, J = 1.9 Hz, 1H), 7.66 – 7.54 (m, 2H), 7.49 (d, J = 8.6 Hz, 1H), 7.02 – 6.91 (m, 2H), 6.88 (t, J = 1.9 Hz, 1H), 4.14 (t, J = 7.7 Hz, 2H), 3.90 (s, 3H), 3.75 (dd, J = 7.6, 6.2 Hz, 2H), 3.57 (d, J = 7.4 Hz, 2H), 3.27 (dd, J = 14.6, 7.4 Hz, 1H), 3.03 (d, J = 24.1 Hz, 6H).
	696	522.1	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.35 (s, 1H), 10.08 (s, 1H), 8.69 (d, J = 2.8 Hz, 1H), 8.23 (s, 1H), 7.96 (dd, J = 9.2, 2.9 Hz, 1H), 7.64 (dt, J = 14.4, 1.9 Hz, 2H), 6.95 (t, J = 2.0 Hz, 1H), 5.48 (s, 1H), 3.07 (s, 3H), 2.48 (s, 3H), 1.35 (s, 6H).



	697	562.05	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.87 (s, 1H), 8.32 – 8.14 (m, 1H), 7.72 (q, J = 1.8 Hz, 2H), 7.64 (t, J = 1.9 Hz, 1H), 7.54 (dd, J = 12.2, 2.4 Hz, 1H), 7.11 (dd, J = 3.4, 1.9 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H), 5.30 – 5.17 (m, 1H), 4.77 (t, J = 8.1 Hz, 1H), 4.45 – 4.32 (m, 1H), 4.28 (d, J = 9.6 Hz, 1H), 3.87 (s, 4H), 3.06 (s, 3H), 1.67 – 1.51 (m, 1H), 0.73 (t, J = 5.4 Hz, 4H).
	698	515.1	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.32 (s, 1H), 7.85 (s, 1H), 7.64 (dt, J = 16.5, 1.9 Hz, 2H), 7.34 (d, J = 1.3 Hz, 1H), 7.23 – 7.13 (m, 3H), 7.04 – 6.94 (m, 4H), 4.20 (t, J = 6.9 Hz, 2H), 3.07 (s, 3H), 2.92 (t, J = 6.9 Hz, 2H), 2.27 (s, 3H).




	699	548.25	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.59 (s, 1H), 10.07 (s, 1H), 8.84 (s, 1H), 8.71 (s, 2H), 8.71 (s, 1H), 7.70 (p, J = 1.9 Hz, 2H), 6.95 (t, J = 2.0 Hz, 1H), 4.58 (d, J = 12.9 Hz, 1H), 3.97 (d, J = 13.4 Hz, 1H), 3.16 (t, J = 12.3 Hz, 1H), 3.08 (s, 3H), 2.92 (d, J = 12.2 Hz, 1H), 2.87 (s, 3H), 2.62 (t, J = 12.5 Hz, 1H), 2.05 (s, 3H), 1.86 (d, J = 12.0 Hz, 2H), 1.79 – 1.54 (m, 2H).
	700	517.9	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.35 (s, 1H), 10.14 (s, 1H), 8.89 (s, 1H), 8.25 (d, J = 2.3 Hz, 1H), 8.03 (s, 1H), 7.64 (t, J = 1.9 Hz, 1H), 7.59 (t, J = 1.9 Hz, 1H), 6.95 (t, J = 1.9 Hz, 1H), 3.16 (q, J = 7.3 Hz, 2H), 2.37 (d, J = 8.1 Hz, 6H), 1.21 (t, J = 7.3 Hz, 3H).



	701	486.85	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.22 (s, 1H), 10.14 – 9.93 (m, 1H), 8.68 (s, 1H), 8.28 (t, J = 1.6 Hz, 1H), 7.92 – 7.76 (m, 2H), 7.71 (q, J = 1.6 Hz, 1H), 7.63 (q, J = 1.7 Hz, 1H), 7.27 – 7.09 (m, 1H), 6.94 (q, J = 1.8 Hz, 1H), 4.08 (s, 3H), 3.07 (d, J = 1.1 Hz, 3H).
	702	554	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.99 (s, 1H), 9.81 (s, 1H), 7.81 – 7.46 (m, 4H), 7.13 – 6.76 (m, 3H), 4.06 (s, 4H), 3.80 (s, 3H), 3.06 (s, 3H), 2.89 (t, J = 12.5 Hz, 4H).




	703	453.05	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.91 (s, 1H), 7.80 – 7.76 (m, 1H), 7.76 – 7.74 (m, 1H), 7.72 (q, J = 1.5 Hz, 1H), 7.65 (dt, J = 2.7, 1.3 Hz, 1H), 7.28 (dd, J = 3.6, 1.9 Hz, 1H), 6.90 (dt, J = 2.8, 1.4 Hz, 1H), 6.78 (ddd, J = 8.8, 2.7, 0.9 Hz, 1H), 4.01 (d, J = 1.0 Hz, 3H), 3.89 (d, J = 0.9 Hz, 3H), 3.06 (d, J = 1.0 Hz, 3H).
	704	489	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.85 (s, 1H), 8.54 – 8.47 (m, 1H), 7.84 – 7.51 (m, 5H), 7.18 (d, J = 1.9 Hz, 1H), 6.90 (t, J = 1.9 Hz, 1H), 3.98 (s, 5H), 3.53 – 3.38 (m, 2H), 3.07 (s, 3H), 2.91 – 2.80 (m, 1H), 1.78 – 1.66 (m, 4H).




	705	507	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.32 (s, 1H), 10.07 (s, 1H), 8.73 (d, J = 2.8 Hz, 1H), 8.53 (s, 1H), 8.18 (dd, J = 9.5, 2.9 Hz, 1H), 7.92 (s, 1H), 7.62 (d, J = 29.4 Hz, 2H), 6.93 (d, J = 2.1 Hz, 1H), 6.74 (s, 1H), 3.04 (s, 3H), 2.17 (s, 3H).
	706	520.85	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.58 (s, 1H), 10.07 (s, 1H), 8.79 (s, 2H), 8.67 (s, 1H), 7.69 (dt, J = 5.8, 1.9 Hz, 2H), 6.95 (t, J = 1.9 Hz, 1H), 5.05 (q, J = 8.8 Hz, 2H), 3.07 (s, 3H), 2.84 (s, 3H).
	707	611.1	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.35 (s, 1H), 10.06 (s, 1H), 8.00 (s, 1H), 7.74 (d, J = 2.2 Hz, 1H), 7.68 – 7.59 (m, 2H), 6.94 (s, 2H), 4.85 (q, J = 8.9 Hz, 2H), 4.44 (t, J = 12.3 Hz, 4H), 3.06 (s, 3H), 2.38 (s, 3H).



	708	593.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.00 (s, 1H), 9.76 (s, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.62 (s, 1H), 7.59 (d, J = 2.3 Hz, 1H), 7.51 (d, J = 2.0 Hz, 1H), 7.07 (d, J = 2.4 Hz, 1H), 6.88 (t, J = 2.0 Hz, 1H), 6.50 (dd, J = 2.1, 1.1 Hz, 1H), 4.89 (q, J = 8.8 Hz, 2H), 4.47 (t, J = 12.3 Hz, 4H), 3.05 (s, 3H), 2.03 (d, J = 0.9 Hz, 3H).
	709	443.85	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.11 (s, 1H), 10.02 (s, 1H), 9.08 (s, 2H), 8.35 (d, J = 2.2 Hz, 1H), 7.69 (d, J = 2.2 Hz, 1H), 7.62 (d, J = 2.4 Hz, 1H), 6.92 (d, J = 2.3 Hz, 2H), 3.06 (s, 3H).



	710	558.95	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.46 (s, 1H), 10.07 (s, 1H), 8.65 (d, J = 2.3 Hz, 1H), 8.39 (s, 1H), 8.00 (dd, J = 8.3, 2.4 Hz, 1H), 7.77 – 7.60 (m, 3H), 6.95 (t, J = 1.9 Hz, 1H), 3.95 – 3.71 (m, 3H), 3.47 – 3.36 (m, 3H), 3.30 (s, 1H), 3.07 (s, 3H), 2.71 (s, 3H).
	711	500.95	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.67 (s, 1H), 10.06 (s, 1H), 9.35 (s, 2H), 8.81 (s, 1H), 7.68 (dt, J = 8.4, 1.9 Hz, 2H), 6.95 (t, J = 1.9 Hz, 1H), 3.46 (s, 3H), 3.07 (s, 3H), 2.94 (s, 3H).
	712	450	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.95 (d, J = 42.4 Hz, 2H), 8.91 (s, 2H), 7.75 (d, J = 2.1 Hz, 1H), 7.70 (s, 1H), 7.66 (s, 1H), 7.54 (d, J = 2.1 Hz, 1H), 6.90 (s, 1H), 4.08 (tt, J = 7.4, 4.2 Hz, 1H), 3.06 (s, 3H), 1.18 – 0.68 (m, 4H).



	713	475.2	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.96 (s, 1H), 9.75 (s, 1H), 8.14 (dd, J = 4.7, 1.5 Hz, 1H), 7.73 (d, J = 8.2 Hz, 1H), 7.68 (dd, J = 6.4, 2.0 Hz, 2H), 7.60 (d, J = 2.6 Hz, 1H), 7.50 (dd, J = 8.3, 4.7 Hz, 1H), 6.87 (t, J = 2.0 Hz, 1H), 6.52 (d, J = 1.8 Hz, 1H), 3.96 (d, J = 6.9 Hz, 2H), 3.03 (d, J = 2.3 Hz, 3H), 2.12 (s, 3H), 1.16 – 1.14 (m, 1H), 0.56 – 0.49 (m, 2H), 0.29 (dd, J = 4.7, 1.8 Hz, 2H).
	714	575.25	¹ H NMR (300 MHz, Chloroform-d) δ 8.16 (s, 1H), 8.00 (d, J = 24.0 Hz, 1H), 7.54 (d, J = 11.7 Hz, 3H), 7.44 (s, 1H), 7.02 (s, 2H), 6.93 (s, 1H), 3.84 (s, 7H), 3.36 (s, 4H), 3.05 (s, 3H), 1.28 (s, 1H), 1.06 (t, J = 3.8 Hz, 2H), 0.85 (dt, J = 7.0, 3.7 Hz, 2H).



	715	538.25	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 10.08 (s, 1H), 8.31 (d, J = 2.9 Hz, 1H), 8.14 (s, 1H), 7.70 – 7.58 (m, 3H), 6.96 (t, J = 1.9 Hz, 1H), 4.51 (t, J = 12.3 Hz, 4H), 3.07 (s, 3H), 2.49 (s, 3H).
	716	441	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.78 (s, 1H), 10.12 (s, 1H), 8.69 (d, J = 3.0 Hz, 1H), 8.46 (dd, J = 8.9, 4.6 Hz, 1H), 7.95 (td, J = 8.8, 3.0 Hz, 1H), 7.79 (t, J = 1.9 Hz, 1H), 7.73 (t, J = 1.9 Hz, 1H), 7.00 (t, J = 1.9 Hz, 1H), 3.09 (s, 3H), 2.90 (s, 3H).
	717	586	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.12 (s, 2H), 8.64 (s, 1H), 8.20 (s, 1H), 7.67 (t, J = 1.9 Hz, 1H), 7.58 (t, J = 2.1 Hz, 1H), 7.53 (d, J = 8.0 Hz, 1H), 7.43 – 7.32 (m, 2H), 7.15 (dd, J = 24.1, 8.1 Hz, 3H), 7.05 (d, J = 7.6 Hz, 1H), 6.94 (t, J = 2.0 Hz, 1H), 5.27 (s, 2H), 3.06 (s, 3H), 2.79 (d, J = 9.1 Hz, 6H).



	718	534	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.16 (s, 2H), 8.68 (s, 1H), 8.52 (d, J = 2.8 Hz, 1H), 8.43 (d, J = 1.8 Hz, 1H), 8.33 (s, 1H), 7.68 (t, J = 1.9 Hz, 1H), 7.60 (ddt, J = 11.4, 9.2, 4.9 Hz, 3H), 7.23 (d, J = 8.6 Hz, 1H), 7.14 (t, J = 9.0 Hz, 1H), 6.94 (t, J = 1.9 Hz, 1H), 5.34 (s, 2H), 3.06 (s, 3H).
	719	507	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.20 (s, 1H), 10.06 (s, 1H), 8.75 (s, 1H), 8.42 (s, 1H), 8.26 (d, J = 2.9 Hz, 2H), 8.11 (dd, J = 10.2, 2.5 Hz, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.60 (d, J = 2.0 Hz, 1H), 7.38 (s, 1H), 6.95 (t, J = 1.9 Hz, 1H), 5.43 (s, 2H), 3.07 (s, 3H).
	720	530.15	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.62 (s, 1H), 10.08 (s, 1H), 8.96 (s, 2H), 8.72 (s, 1H), 7.69 (dt, J = 7.9, 1.9 Hz, 2H), 6.95 (t, J = 1.9 Hz, 1H), 3.38 (s, 3H), 3.14 (s, 3H), 3.07 (s, 3H), 2.88 (s, 3H).



	721	506	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.15 (s, 1H), 8.52 (s, 1H), 8.38 (s, 1H), 8.27 (s, 1H), 7.76 – 7.47 (m, 3H), 7.32 (d, J = 15.8 Hz, 2H), 7.12 (t, J = 9.0 Hz, 1H), 6.92 (d, J = 2.1 Hz, 1H), 5.31 (s, 2H), 3.04 (s, 3H).
	722	612	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.07 (d, J = 5.5 Hz, 2H), 8.63 (d, J = 0.7 Hz, 1H), 8.20 (d, J = 0.7 Hz, 1H), 7.68 (t, J = 1.8 Hz, 1H), 7.59 (t, J = 1.9 Hz, 1H), 7.50 – 7.33 (m, 2H), 7.31 – 7.06 (m, 3H), 6.93 (t, J = 1.9 Hz, 1H), 6.29 (d, J = 2.3 Hz, 1H), 6.15 (dd, J = 8.6, 2.3 Hz, 1H), 5.23 (s, 2H), 4.74 (s, 4H), 4.04 (s, 4H), 3.07 (s, 3H).



	723	546	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.98 (s, 1H), 9.82 (s, 1H), 8.52 (d, J = 2.8 Hz, 1H), 8.47 (d, J = 1.8 Hz, 1H), 8.19 (dd, J = 4.7, 1.3 Hz, 1H), 7.92 – 7.81 (m, 2H), 7.75 – 7.66 (m, 2H), 7.62 (d, J = 2.0 Hz, 1H), 7.54 (dd, J = 8.3, 4.7 Hz, 1H), 6.89 (t, J = 2.0 Hz, 1H), 6.77 (d, J = 1.9 Hz, 1H), 5.33 (s, 2H), 4.93 (t, J = 5.7 Hz, 1H), 4.41 (d, J = 5.7 Hz, 2H), 3.04 (s, 3H).
	724	550	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.89 (s, 1H), 9.80 (s, 1H), 8.27 (d, J = 2.4 Hz, 1H), 7.81 (dd, J = 11.0, 2.4 Hz, 1H), 7.72 – 7.54 (m, 4H), 7.08 (d, J = 2.0 Hz, 1H), 6.88 (t, J = 2.0 Hz, 1H), 5.29 (s, 2H), 3.74 (s, 3H), 3.05 (s, 3H), 2.65 (s, 3H).



	725	548	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.99 (s, 1H), 9.79 (s, 1H), 8.30 (d, J = 2.3 Hz, 1H), 7.82 (dd, J = 11.0, 2.4 Hz, 1H), 7.71 – 7.64 (m, 2H), 7.63 – 7.57 (m, 1H), 6.95 (dd, J = 2.0, 0.9 Hz, 1H), 6.89 (t, J = 1.9 Hz, 1H), 5.08 (s, 2H), 3.70 (s, 3H), 3.05 (s, 3H), 2.38 (s, 3H), 2.13 (s, 3H).
	726	534	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.98 (s, 1H), 9.80 (s, 1H), 8.32 (d, J = 2.4 Hz, 1H), 7.88 (dd, J = 10.8, 2.4 Hz, 1H), 7.80 (s, 1H), 7.68 (d, J = 2.1 Hz, 2H), 7.59 (s, 1H), 7.00 (d, J = 2.0 Hz, 1H), 6.88 (s, 1H), 5.46 (s, 2H), 3.97 (s, 3H), 3.70 (s, 3H), 3.04 (s, 3H).

	727	525	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.80 (s, 1H), 8.53 (d, J = 1.8 Hz, 1H), 8.46 (d, J = 2.7 Hz, 1H), 7.91 (d, J = 8.0 Hz, 1H), 7.82 (dt, J = 10.4, 2.2 Hz, 1H), 7.76 – 7.68 (m, 2H), 7.62 (t, J = 1.9 Hz, 1H), 7.52 – 7.35 (m, 3H), 7.24-7.11 (d, J = 16.5 Hz, 2H), 6.89 (t, J = 1.9 Hz, 1H), 6.70 (d, J = 1.9 Hz, 1H), 3.44 (s, 3H), 3.06 (s, 3H).
	728	546	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.02 (s, 1H), 9.80 (s, 1H), 8.54 (d, J = 2.8 Hz, 1H), 8.37 (d, J = 2.8 Hz, 1H), 8.15 (t, J = 1.9 Hz, 1H), 7.87 (dd, J = 9.8, 2.9 Hz, 1H), 7.74 (dd, J = 9.8, 2.9 Hz, 1H), 7.65 (m, 1H), 7.63 (t, J = 1.9 Hz, 1H), 7.45 (ddd, J = 10.1, 2.8, 1.7 Hz, 1H), 6.94 (dd, J = 9.8, 2.9 Hz, 1H), 6.81 (m, 1H), 3.45 (s, 3H), 3.14-3.06 (s, 4H), 2.91 (dd, J = 9.0, 6.3 Hz, 2H).

	729	544	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.97 (s, 1H), 9.82 (s, 1H), 8.66 (d, J = 2.7 Hz, 1H), 8.60 (d, J = 1.8 Hz, 1H), 8.50 (d, J = 2.7 Hz, 1H), 8.26 (dd, J = 10.1, 2.8 Hz, 1H), 7.98 – 7.87 (m, 1H), 7.77 (d, J = 1.9 Hz, 1H), 7.67 (t, J = 1.8 Hz, 1H), 7.59 (t, J = 1.9 Hz, 1H), 7.50 – 7.34 (m, 2H), 6.92 – 6.80 (m, 2H), 3.72 (s, 3H), 3.04 (s, 3H).
	730	540	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.88 (s, 2H), 8.46 (d, J = 2.0 Hz, 1H), 8.38 (d, J = 2.1 Hz, 1H), 8.31 – 8.19 (m, 1H), 7.75 (d, J = 2.2 Hz, 1H), 7.73 – 7.62 (m, 4H), 7.36 (dd, J = 8.4, 4.6 Hz, 1H), 7.22 (d, J = 2.0 Hz, 1H), 6.90 (t, J = 2.0 Hz, 1H), 5.27 (s, 2H), 3.79 (s, 3H), 3.06 (s, 3H), 2.59 (q, J = 7.6 Hz, 2H), 1.12 (t, J = 7.6 Hz, 3H).

	731	503.95	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.03 (s, 1H), 8.58 – 8.42 (m, 1H), 8.24 (d, J = 1.7 Hz, 1H), 8.03 (td, J = 7.8, 1.9 Hz, 1H), 7.70 (t, J = 1.8 Hz, 1H), 7.65 – 7.54 (m, 2H), 7.44 (dd, J = 7.5, 4.9 Hz, 1H), 7.02 (d, J = 1.8 Hz, 1H), 6.92 (t, J = 2.0 Hz, 1H), 3.54 (s, 8H), 3.06 (s, 3H).
	732	537	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.34 (s, 1H), 10.06 (s, 1H), 8.40 (s, 1H), 8.33 (d, J = 2.4 Hz, 1H), 8.03 (s, 1H), 7.89 (dd, J = 10.9, 2.4 Hz, 1H), 7.66 (t, J = 1.9 Hz, 1H), 7.60 (t, J = 1.9 Hz, 1H), 7.37 (s, 1H), 6.94 (t, J = 1.9 Hz, 1H), 5.37 (s, 2H), 3.06 (s, 3H), 2.34 (s, 3H).

	733	593	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.98 (s, 1H), 9.76 (s, 1H), 8.42 (s, 1H), 7.70 (t, J = 1.9 Hz, 1H), 7.65 (d, J = 2.2 Hz, 1H), 7.63 – 7.56 (m, 2H), 7.35 (s, 1H), 6.96 (d, J = 2.3 Hz, 1H), 6.88 (t, J = 2.0 Hz, 1H), 6.83 (d, J = 1.9 Hz, 1H), 5.31 (s, 2H), 4.43 (t, J = 12.2 Hz, 4H), 3.66 (s, 3H), 3.06 (s, 3H).
	734	562.15	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.01 (s, 1H), 9.92 (s, 1H), 8.54 (s, 1H), 7.81 (d, J = 2.0 Hz, 1H), 7.65 (dt, J = 16.7, 1.9 Hz, 2H), 7.54 (d, J = 2.0 Hz, 1H), 7.22 (ddt, J = 9.5, 7.6, 2.8 Hz, 3H), 6.90 (t, J = 2.0 Hz, 1H), 5.39 (s, 2H), 3.99 (s, 3H), 3.06 (s, 3H), 2.58 (s, 3H).

	735	599.1	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.04 (s, 1H), 9.98 (s, 1H), 9.18 (s, 1H), 8.90 (s, 2H), 8.32 (dd, J = 2.4, 0.6 Hz, 1H), 7.92 – 7.79 (m, 2H), 7.65 (dt, J = 10.5, 1.8 Hz, 2H), 7.25 (d, J = 1.9 Hz, 1H), 6.93 (t, J = 1.9 Hz, 1H), 5.51 – 5.36 (m, 4H), 3.07 (s, 3H).
	736	474	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.41 (s, 1H), 10.18 (s, 1H), 8.75 (dt, J = 4.8, 1.9 Hz, 1H), 8.06 (s, 1H), 7.84 (td, J = 8.8, 2.3 Hz, 1H), 7.64 (t, J = 1.9 Hz, 1H), 7.57 (t, J = 1.9 Hz, 1H), 6.93 (t, J = 2.0 Hz, 1H), 3.04 (s, 3H), 2.37 (d, J = 1.1 Hz, 3H).

Example 737. ASSAYS

Compounds of the disclosure were assessed for their ability to inhibit DHX9 activity. The inhibitory properties of the compounds of the disclosure described herein can be

5 evidenced by testing in any one of the following assays.

1. Biochemical Assay

Materials and methods

5 **Recombinant DHX9** - Human recombinant DHX9 protein was custom ordered from Viva.

RNA substrate - Oligos for double-stranded RNA substrate were custom ordered from IDT Technologies. (Oligo1 – GCCUGGUCCCUGUCCUUGUUAUUUCCUUGGUUAAUU (SEQ ID NO:1). Oligo2 –
10 GAAUUAACCAAGGAAAAUAACAAGGACAGGGACCAGG (SEQ ID NO:2)). Each oligo was reconstituted in RNase/DNase-free water to achieve a 100uM solution. The two oligo solutions were mixed in 1:1 ratio and annealed by heating at 70 °C for 5 minutes and cooling gradually to room temperature on the benchtop.

Chemicals and Assay Components - Aurintricarboxylic Acid and dimethylsulfoxide
15 were purchased from Sigma-Aldrich (St. Louis, MO). White 384-well assay plates (Catalog #781075) were obtained from Greiner Bio-One (Frickenhause, Germany). The ADP-Glo™ kinase assay kit from Promega Corporation is composed of ADP-Glo reagent, kinase detection reagent (made by mixing kinase detection buffer with a lyophilized kinase detection substrate), Ultra Pure ATP and ADP.

20

ATPase Assay

 DHX9 ATPase assay was performed in small-volume, nonbinding, 384-well white plates at a final volume of 10 µL/well. Test compounds (10 mM solution in DMSO; 100
25 nL/well) were serially diluted on Bravo (Agilent, Santa Clara, CA) and dispensed into wells of columns 3–22 of the plates using an Echo 555 acoustic dispenser (Labcyte, Sunnyvale, CA). 100nL of Aurintricarboxylic acid was dispensed into low control wells and 100 nL of DMSO was dispensed into high control wells. Then, a Multidrop Combi Reagent Dispenser (Thermo Fisher Scientific, Waltham, MA) was used to add a solution of DHX9 (1.25 nM, 5
30 µL/well) in assay buffer (40 mM HEPES [pH 7.5], 0.01% Tween 20, 0.01%BSA, 1 mM DTT, 5 mM MgCl₂, 0.004U/ml RNaseOUT). The reaction was initiated by the addition of 5 µL of substrate solution (30 nM double-stranded RNA substrate, 10 µM Ultra Pure ATP in assay buffer) into the wells. The plates were incubated at room temperature for 1 h. After the indicated incubation times, 10 µL ADP-Glo reagent was added to the reactions and the plate

was incubated at room temperature for 40 min. Then, 20 μ L of kinase detection reagent was added and after an incubation time of 40 min, luminescence was recorded on Envision plate reader (Perkin-Elmer, Billerica, MA).

5 *Data Processing and Analysis*

All data was analyzed by Accent Therapeutics internal data analysis software developed by Scigilian (Montreal, Canada). The percentage inhibition was calculated based on the high control (DMSO) as 0% inhibition, and low control (10 μ M Aurintricarboxylic acid) control as 100% inhibition and used for the calculation of IC₅₀ and IP (inflection point) values by fitting the dose–response curves to a four-parameter logistic model. Assay results reported IP values instead of IC₅₀ values since several compounds did not reach 100% inhibition at higher compound concentrations.

15 **2. DHX9 Cellular Proliferation Assay**

The CellTiter-Glo® Luminescent Cell Viability Assay is a homogeneous method to determine the number of viable cells in culture based on quantitation of the ATP present, which signals the presence of metabolically active cells.

The following cell lines were obtained from ATCC: HCT116 (CCL-247), Colo205 (CCL-222), LS411N (CRL-2159), and NCI-H747 (CCL-252). HCT116 cells were grown and assayed in McCoy's-5a growth media supplemented with 10%FBS, and Colo205, LS411N and NCI-H747 cells were grown in RPMI-1640 media supplemented with 10%FBS.

The cells are plated at pre-determined cell densities in 384-well solid white cell culture plates and incubated overnight in 37°C at 5% CO₂. In a separate plate, reference and test compounds were prepared in DMSO stock solution and serially diluted 3-fold for 10 points. Compound (0.1% final DMSO concentration) was transferred to the cell plate in designated wells, and incubated 37°C at 5% CO₂ for 120 hours. CellTiter-Glo reagent was prepared fresh according to manufacturer's (Promega) directions and added to each well. The plate is then shaken at 300rpm for 10 minutes at RT, and then read on an Envision plate reader using a Luminescence protocol.

Data analysis was performed by normalizing the raw data (raw luminescence units or RSample) to an average of the positive control values for wells containing culture media only (100% cell death or RLC) and the negative control values for 0.1% DMSO (0% cell death or RHC). An IC₅₀ was calculated using a 4-parameter logistic nonlinear regression model in

Scigilian Analyzer or GraphPad Prism, constraining the top parameter to 100 and the bottom parameter to 0.

$$\% \text{Inhibition} = 100\% \times \left(1 - \frac{R_{\text{Sample}} - R_{\text{LC}}}{R_{\text{HC}} - R_{\text{LC}}} \right)$$

IC₅₀ Fit formula = (A+((B-A)/(1+((x/C)^D)))) ... whereas A:Bottom; B:Top; C:IC₅₀;

5 D:Slope

3. DHX9 circBRIP1 Cellular Target Engagement SYBR Green Assay

Evaluation of the circular and linear format of BRIP1 mRNA in HCT116 cells by Quantitative Polymerase Chain Reaction (qPCR). Upon DHX9 inhibition circBRIP1 levels
10 are increased and linear BRIP1 mildly decreases or remains unchanged.

HCT116 cells were obtained from ATCC (CCL-247) and were grown and assayed in McCoy's 5a growth media supplemented with 10%FBS.

The cells are plated at pre-determined cell densities in 384-well cell culture plates and incubated overnight in 37°C at 5% CO₂. In a separate plate, reference and test compounds
15 were prepared in DMSO stock solution and serially diluted 3-fold for 10 points. Compound (0.1% final DMSO concentration) was transferred to the cell plate in designated wells, and incubated 37°C at 5% CO₂ for 72 hours.

RNA is extracted from the cells using the SYBR Green Fast Advanced Cell-to-CT kit according to manufacturer's (Invitrogen) protocol, and cDNA is reverse transcribed using the
20 High Capacity RNA-to-cDNA Kit according to the manufacturer's (Invitrogen) protocol. Using SYBR Green Master Mix and thermal cycler conditions for the QuantStudio Thermocycler, qPCR is carried out using the below custom primer sequences:

- BRIP1-Circular-Forward (TCT GTG TGC CAG ACT GTG AG (SEQ ID NO:3))
- BRIP1-Circular-Reverse (ACA CCA AGT TCT GAC GAA AAG G (SEQ ID NO:4))
- 25 • BRIP1-Linear-Forward (GTC CAT CCT GAG GTA GTC GG (SEQ ID NO:5))
- BRIP1-Linear-Reverse (TTC CCC AGG CTG ACA AGT TC (SEQ ID NO:6))
- GAPDH-Forward (GTCTCCTCTGACTTCAACAGCG (SEQ ID NO:7))
- GAPDH-Reverse (ACCACCCTGTTGCTGTAGCCAA (SEQ ID NO:8))

Note: Forward and Reverse primers are combined per gene, and each sample has to be run on a separate plate for each gene.

Data analysis was performed by using the QuantStudio Thermocycler software to determine threshold Ct values. ΔCt for circBRIP1 or linBRIP1 is calculated by normalizing the target gene Ct to the GAPDH housekeeping gene Ct for each sample. The inhibition activity is calculated by the following formula, whereas:

- ΔCt_s : Mean of Ct(target gene) – Mean of Ct(GAPDH) for each sample
- ΔCt_c : Mean of Ct(target gene) – Mean of Ct(GAPDH) for DMSO control.

$$\% \text{Inhibition} = 100\% \times \frac{2^{\Delta Ct_s}}{2^{\Delta Ct_c}}$$

An EC_{50} was then calculated using a 4-parameter logistic nonlinear regression model in Scigilian Analyzer or GraphPad Prism, floating both the top and bottom parameters.

EC_{50} Fit formula = $(A + ((B - A) / (1 + ((x/C)^D))))$... whereas A:Bottom; B:Top; C: IC_{50} ; D:Slope

4. DHX9 circBRIP1 Cellular Target Engagement TaqMan Multiplex Assay:

Evaluation of the circular and linear format of BRIP1 mRNA in HCT116 cells by Quantitative Polymerase Chain Reaction (qPCR). Upon DHX9 inhibition circBRIP1 levels are increased and linear BRIP1 mildly decreases or remains unchanged. HCT116 cells were obtained from ATCC (CCL-247) and were grown and assayed in McCoy's 5a growth media supplemented with 10% FBS.

The cells are plated at pre-determined cell densities in 384-well cell culture plates and incubated overnight in 37 °C at 5% CO₂. In a separate plate, reference and test compounds were prepared in DMSO stock solution and serially diluted 3-fold for 10 points. Compound (0.1% final DMSO concentration) was transferred to the cell plate in designated wells, and incubated 37 °C at 5% CO₂ for 72 hours.

RNA is extracted from the cells using the TaqMan Gene Expression Cells-to-CT™ kit according to manufacturer's (Invitrogen) protocol, and cDNA is reverse transcribed using the High Capacity RNA-to-cDNA Kit according to the manufacturer's (Invitrogen) protocol. Using TaqMan Multiplex Mastermix and thermal cycler conditions for the QuantStudio Thermocycler, qPCR is carried out using TaqMan Gene Expression Assay with VIC Dye for

Linear BRIP1 (Assay ID: Hs00908156_m1), TaqMan GAPDH primer limited assay with JUN dye/QSY probe, and the below custom TaqMan Gene Expression Assay with FAM Dye for circBRIP1:

- Forward Primer: GTGAGCCAAGGAATTTTGTGTTT (SEQ ID NO:9)
- Reverse Primer: GGTCTGAACCTTGCCATTAATATCTG (SEQ ID NO:10)
- Probe: CCATCTTACAAGGCCTTT (SEQ ID NO:11)

Note: All target genes can be combined in one well per sample.

Data analysis was performed by using the QuantStudio Thermocycler software to determine threshold Ct values. ΔCt for circBRIP1 or linBRIP1 is calculated by normalizing the target gene Ct to the GAPDH housekeeping gene Ct for each sample. The inhibition activity is calculated by the following formula, whereas:

- ΔCts: Mean of Ct(target gene) – Mean of Ct(GAPDH) for each sample
- ΔCtc: Mean of Ct(target gene) – Mean of Ct(GAPDH) for DMSO control.

$$\%Inhibition = 100\% \times \frac{2^{\Delta Cts}}{2^{\Delta Ctc}}$$

An EC₅₀ was then calculated using a 4-parameter logistic nonlinear regression model in Scigilian Analyzer or GraphPad Prism, floating both the top and bottom parameters.

IC50 Fit formula = (A+((B-A)/(1+((x/C)^D)))) ... whereas A:Bottom; B:Top; C:IC50; D:Slope

DATA FOR EXAMPLES

Table 1.

Example Number	DHX9 Biochemical Inflection Point (μM)	DHX9 Biochemical Max Inhibition (%)	DHX9 Cellular Target Engagement EC50 (μM)	**LS411N (CRC-MSI-H) Antiproliferative IC ₅₀	**H747 (CRC-MSS) Antiproliferative IC ₅₀
1	0.32	48.3	16.4 ⁺		
2	0.173	64.8	>30 ⁺		
3	0.648	76.4			
4	4.82	64.1			
5	0.215	74.6	>30 ⁺		
6	0.0917	52.5	9.34 ⁺		
7	0.38	46.4	24.2 ⁺		
8	0.395	54.4	5.52 ⁺		
9	0.39	44.5	30 ⁺		

10	0.133	61	12.3 ⁺		
11	0.123	62.7	0.851 ⁺		
12	84.7	58.6			
13	1.15	41.7	7.97 ⁺		
14	0.353	59.6			
15	0.1355	54.8	0.7283 ⁺	0.663	>10
16	0.251	46.3	3.51 ⁺		
17	0.0073	61	0.0571 ⁺	0.2783	9.6533
18	0.0086	61	0.0984 ⁺	0.546	>10
19	0.115	53.5	2.11 ⁺	1.08	>10
20	0.0572	62.5	0.1 ⁺	0.609	7.22
21	0.0332	61.7	0.26 ⁺	0.791	>10
22	0.0025	71	0.0358 ⁺	0.115	>10
23	0.0032	69	0.0542 ⁺	0.0531	>10
24	0.002	73.4	0.015 ⁺	0.117	>10
25	0.0056	61.3	0.048 ⁺	0.0991	>10
26	0.0018	65.6	0.0609 ⁺	0.217	>10
27	0.0064	68	0.0497 ⁺	0.117	>10
28	0.0042	68.3	0.0258 ⁺	0.0788	>10
29	0.0482	65.2	0.0519 ⁺	0.124	9.28
30	0.787	40.5	0.687 ⁺		
31	0.0038	72.5	0.0642 ⁺	0.0698	>10
32	0.0383	56.3	0.0482 ⁺	0.101	>10
33	0.0048	61.7	0.0215 ⁺	0.116	>10
34	0.0063	75.2	0.013 ⁺	0.0141	8.67
35	0.0039	75.7	0.0372 ⁺	0.0768	>10
36	0.028	56.86	0.3134 ⁺	1.085	>10
37	0.0479	62.6	0.83 ⁺	3.47	>10
38	0.0655	44.7	0.933 ⁺		
39	0.0701	52.3	0.884 ⁺		
40	0.0157	60	1.15 ⁺		
41	0.0474	40.3	4.34 ⁺		
42	0.0187	48.7	0.308 ⁺	0.917	>10
43	0.258	49.8	4.71 ⁺		
44	0.137	60.3	4.19 ⁺		
45	0.0164	67.4	0.563 ⁺		
46	0.0473	57.2	2.25 ⁺		
47	0.269	64.7	6.73 ⁺		
48	0.0931	54.1	1.41 ⁺		
49	0.0497	58.6	0.4335 ⁺	0.901	>10
50	0.045	55.5	0.895 ⁺		

51	0.145	61.1	0.188 ⁺	0.259	6.47
52	0.198	37.9	3.01 ⁺		
53	1.91	52.2	4.95 ⁺		
54	0.0656	66.6	0.953 ⁺		
55	0.0099	48.6	0.143 ⁺	1.06	>10
56	0.197	48.8	7.43 ⁺		
57	14.6	45.8			
58	0.0849	44.4	1.41 ⁺		
59	0.0127	77.5	0.104 ⁺	0.357	>10
60	0.0078	63	0.201 ⁺		
61	0.0145	66.6	1.52 ⁺		
62	0.211	58.4	6.8 ⁺		
63	0.0839	66.2	2.43 ⁺		
64	0.0599	61.8	5.64 ⁺		
65	0.0055	72.6	0.0631 ⁺	0.109	>10
66	0.0566	62.9	4.87 ⁺		
67	0.154	54	0.297 ⁺	1.05	>10
68	0.0238	68.7	0.278 ⁺		
69	0.113	62.3	0.661 ⁺		
70	0.0085	78.9	0.107 ⁺	0.375	>10
71	0.171	77.8	26.3 ⁺		
72	0.378	67.2			
73	0.189	63	11.2 ⁺		
74	0.255	57.7	8.32 ⁺		
75	0.0946	55.5	1.64 ⁺		
76	0.0092	71.8	0.0623 ⁺	0.253	>10
77	0.0154	57.8	0.154 ⁺		
78	0.0319	56.2	0.457 ⁺		
79	0.146	67.8	1.12 ⁺		
80	0.493	71.2	3.09 ⁺		
81	0.0326	66.7	0.29 ⁺	1.07	>10
82	0.0092	69.4	0.0732 ⁺	0.134	>10
83	0.0478	60.7	0.911 ⁺		
84	0.0101	66.7	0.194 ⁺	0.385	>10
85	0.099	59.5	2.76 ⁺		
86	0.0184	51.5	0.585 ⁺		
87	0.0261	55.7	2.46 ⁺		
88	0.0071	54.2	>10 ⁺		
89	0.0144	64.9	0.667 ⁺		
90	0.024	61.5	1.06 ⁺		
91	0.0279	59.7	0.382 ⁺		

92	0.0221	61.1	0.281 ⁺	1.1	>10
93	0.0017	70.6	0.03 ⁺	0.253	>10
94	0.0018	72.4	0.292 ⁺	3.2	>10
95	0.0205	59.2	0.179 ⁺	1.02	7.99
96	0.253	52.8	9.63 ⁺	>10	>10
97	0.488	61.5	3.16 ⁺		
98	0.0483	62.4	1.33 ⁺		
99	0.298	50.9	>10 ⁺		
100	13.8	49.9			
101	13	53.6			
102	0.762	61.8	>30 ⁺		
103	0.452	49.5	5.31 ⁺		
104	0.061	68	5.49 ⁺		
105	0.0803	71.7	8.33 ⁺		
106	0.078	58.7	0.414 ⁺		
107	0.0622	53	0.913 ⁺		
108	0.21	59.8	6.65 ⁺		
109	0.0176	57.2	0.437 ⁺		
110	0.04	53.7	2.71 ⁺		
111	0.25	68.2	0.161 ⁺	0.318	>10
112	0.0211	70.8	0.627 ⁺		
113	0.04	65.9	7.44 ⁺		
114	0.0435	66.4	1.48 ⁺		
115	0.0743	56.5	0.41 ⁺		
116	0.0097	56.6	0.734 ⁺		
117	0.104	55.5	3.87 ⁺		
118	0.0451	55.8	2.93 ⁺		
119	1.16	82.1			
120	0.0089	72.5	0.144 ⁺	1.07	6.98
121	0.0406	63.5	5.43 ⁺		
122	0.0288	67.7	1.29 ⁺		
123	0.0148	72.8	6.59 ⁺		
124	0.604	53.8	5.14 ⁺		
125	0.33	56.7	16.1 ⁺		
126	0.182	64.5	>10 ⁺		
127	0.138	66.4	>10 ⁺		
128	0.173	57.3	>10 ⁺		
129	0.0384	66.4	0.333 ⁺	1.09	>10
130	0.0396	70.4	0.489 ⁺	1.16	>10
131	0.151	55.9	>10 ⁺		
132	0.295	55.1	>10 ⁺	>10	>10

133	13.6	41.8			
134	0.808	40			
135	1.07	44.6			
136	0.234	59.6	6.41 ⁺		
137	0.0737	62.9	3.01 ⁺		
138	13.5	42.6	>10 ⁺		
139	6.65	59.1	+		
140	0.3	63.3	>10 ⁺		
141	0.442	54.9	>10 ⁺		
142	0.514	62	>10 ⁺		
143	0.0511	66.8	0.516 ⁺	1.09	8.54
144	0.224	65.1	5.85 ⁺		
145	0.058	67.9	>10 ⁺		
146	0.0304	71.9	>10 ⁺		
147	0.0063	76.5	0.834 ⁺		
148	0.0185	63.3	4.34 ⁺		
149	0.0199	65.4	1.8 ⁺		
150	0.147	65.9	6.56 ⁺		
151	0.179	58.9	>10 ⁺		
152a	0.138	63.2	>10 ⁺		
152b	0.126	69.9			
153	0.256	59.7	6.11 ⁺		
154	0.179	58.9	>10 ⁺		
155	0.126	66.4	3.61 ⁺		
156	0.695	51.2	>10 ⁺		
157	0.0525	61.7	4.32 ⁺		
158	0.0345	76.5	1.29 ⁺		
159	4.27	61	>10 ⁺		
160	0.154	59.6	6.94 ⁺	>10	>10
161	0.0484	44.9	1.42 ⁺		
162	0.0197	60.5	0.138 ⁺		
163	0.0043	71.7	0.07 ⁺	0.609	>10
164	9.3	50.5	+		
165	0.0539	57.4	>10 ⁺		
166	0.0285	64	2.39 ⁺		
167	0.386	58.6	1.75 ⁺		
168	0.772	50.7	2.13 ⁺		
169	0.0087	67.9	0.189 ⁺	2.17	>10
170	0.0057	75.4	0.0447 ⁺	0.122	>10
171	0.0145	61.8	0.0957 ⁺	0.372	>10
172	0.0159	72.4	0.135 ⁺	0.237	6.89

173	0.0086	61.3	0.118 ⁺		
174	0.0018	75.8	0.007 ⁺	0.0333	>10
175	0.0036	78.8	0.0445 ⁺	0.0863	>10
176	0.011	70.2	0.408 ⁺		
177	0.0216	77	0.0848 ⁺	0.251	>10
178	0.011	67.9	0.101 ⁺	0.0811	>10
179	0.0272	72.3	0.997 ⁺		
180	0.0202	67.4	1.2 ⁺		
181	0.0096	57.4	2.09 ⁺		
182	0.181	73.4	>10 ⁺		
183	0.0201	67.1	>10 ⁺		
184	0.0194	66.3	>10 ⁺		
185	0.0047	80.5	0.0733 ⁺	0.139	8.38
186	0.0039	87.9	0.161 ⁺	2.44	>10
187	0.0032	82.4	0.0504 ⁺	0.0773	>10
188	0.0188	66.2	1.46 ⁺		
189	0.0327	72.1	0.308 ⁺		
190	0.0191	68.8	0.373 ⁺		
191	0.0042	67.6	0.432 ⁺	>10	>10
192	0.0165	72.6	3.6 ⁺		
193	0.0111	65.8	>10 ⁺		
194	0.0068	64.3	2.68 ⁺		
195	0.0353	67.1	0.751 ⁺		
196	0.0786	68.9	6.73 ⁺		
197	0.026	72.7	2.37 ⁺		
198	0.0713	60.7	>10 ⁺		
199	0.0844	73.4	>10 ⁺		
200	0.0846	57.7	1.31 ⁺		
201	0.184	66.1	>10 ⁺		
202	0.0127	72.3	0.103 ⁺	0.111	8.56
203	0.0281	76.5	0.341 ⁺		
204	0.0096	68.9	0.0539 ⁺	0.0548	>10
205	0.0226	71.4	0.841 ⁺		
206	0.0168	74.9	1.06 ⁺		
207	0.0152	70	0.508 ⁺		
208	0.0155	79.9	0.125 ⁺	0.218	>10
209	0.0894	64	3.46 ⁺		
210	0.0036	71.3	0.0772 ⁺	0.111	>10
211	0.0624	65.4	0.189 ⁺		
212	0.0052	72.6	0.0326 ⁺	0.202	>10
213	0.0164	71.6	0.0746 ⁺	0.173	>10

214	0.088	75.9	0.106 ⁺	0.203	7.15
215	0.004	75.4	0.0237 ⁺	0.118	>10
216	0.0044	78	0.0127 ⁺	0.0385	>10
217	0.0021	69.6	0.0136 ⁺	0.0533	>10
218	0.0765	59	1.39 ⁺		
219	0.0406	62.3	0.265 ⁺		
220	0.016	79.7	0.303 ⁺		
221	0.0066	78.3	0.121 ⁺		
222	0.0177	78.7	0.2 ⁺		
223	0.0565	75.7	0.421 ⁺		
224	0.0064	67.3	0.0558 ⁺	0.122	>10
225	0.0069	74.2	0.0199 ⁺	0.0439	>10
226	0.0084	73.6	0.388 ⁺		
227	0.0072	70.6	0.0403 ⁺	0.879	>10
228	0.0655	67.2	0.0483 ⁺	0.117	>10
229	0.0096	72.5	0.0781 ⁺	0.051	>10
230	0.093	65.9	7.25 ⁺		
231	0.0491	66.3	0.203 ⁺		
232	0.0202	77.5	0.061 ⁺		
233	0.0455	80.3	0.46 ⁺	0.356	>10
234	0.0095	84.6	1.18 ⁺		
235	0.0358	78.1	0.645 ⁺		
236	0.0521	68.9	0.106 ⁺		
237	0.021	73.5	1.91 ⁺	0.249	3.78
238	0.0229	72.9	0.289 ⁺		
239	0.008	67.7	0.348 ⁺		
240	0.353	71	0.228 ⁺		
241	0.115	63.8	3.68 ⁺		
242	0.105	69.7	5.02 ⁺		
243	0.0923	73.2	>10 ⁺		
244	0.0733	71.9	>10 ⁺		
245	0.121	71.4	4.31 ⁺		
246	0.186	67.5	3.93 ⁺		
247	0.131	70	2.24 ⁺		
248	0.185	70.8	>10 ⁺		
249	0.0028	64.1	0.0295 ⁺		
250	0.185	64.6	3.55 ⁺		
251	5.2	64.7	>10 ⁺	0.432	>10
252	0.0489	72.1	1.79 ⁺		
253	0.11	70	9.23 ⁺		
254	0.0595	64.1	0.578 ⁺		

255	0.0724	75.8	0.378 ⁺	0.0743	>10
256	0.403	70.1	2.03 ⁺		
257	0.207	66.5	2.61 ⁺		
258	0.0047	72.9	0.291 ⁺		
259	1.94	69.2	>10 ⁺		
260	0.715	50	>10 ⁺		
261	0.0475	71	1.53 ⁺		
262	0.12	66.5	0.439 ⁺		
263	0.17	63.8	>10 ⁺		
264	0.119	56.7	3.54 ⁺		
265	8.05	51.8	4.65 ⁺		
266	0.0131	64.4	1.16 ⁺		
267	0.248	66.4	2.26 ⁺		
268	0.0094	73.1	0.182 ⁺		
269	0.0044	76.9	0.037 ⁺		
270	0.0097	74.1	0.101 ⁺		
271	0.0865	66.1	0.761 ⁺		
272	0.0054	71.6	0.0311 ⁺		
273	0.0065	71.4	0.0375 ⁺		
274	0.0322	72.1	0.0979 ⁺		
275	0.0762	75.1	0.19 ⁺		
276	0.0047	73.8	0.0642 ⁺	0.0617	>10
277	0.0043	72.5	0.0476 ⁺	0.0667	4.7
278	0.0442	74.1	0.251 ⁺		
279	0.004	73.2	0.0533 ⁺	0.0811	>10
280	0.326	76.3	0.0742 ⁺	0.116	>10
281	0.028	75.7	0.483 ⁺	0.367	7.34
282	0.0964	72.4	1.88 ⁺		
283	0.0292	78	0.131 ⁺	0.0923	>10
284	0.0041	79.6	0.103 ⁺	0.161	>10
285	0.0054	70.6	0.503 ⁺		
286	0.24	72.9	2.63 ⁺	0.131	>10
287	0.0191	82.6	0.232 ⁺	0.165	>10
288	0.132	58.7	7.8 ⁺		
289	0.0806	66.9	6.55 ⁺		
290	5.59	59.7	>10 ⁺		
291	0.144	59.7	0.803 ⁺	0.352	>10
292	0.076	68.5	2.3 ⁺		
293	0.0872	55	>10 ⁺		
294	1.61	59	>10 ⁺		
295	0.307	63.8	>10 ⁺		

296	0.125	69.9	1.14 ⁺		
297	0.997	57	>10 ⁺		
298	4.21	58.8	>10 ⁺		
299	0.335	56.7	>10 ⁺		
300	0.132	58.5	3.94 ⁺		
301	0.0377	69.5	2.41 ⁺		
302	0.64	54.6	1.93 ⁺		
303	4.02	57.2	>10 ⁺		
304	0.399	59.1	5.98 ⁺		
305	0.0953	63.8	3.33 ⁺		
306a	0.127	70.6	>10 ⁺		
306b	0.253	74.2	>10 ⁺		
307	0.143	70.4	>10 ⁺		
308	0.375	60.6	>10 ⁺		
309	0.162	60.8	3.73 ⁺		
310	0.102	70.3	>10 ⁺		
311	0.0385	77.7	3.98 ⁺		
312	0.22	69.2	9.97 ⁺		
313	0.147	63.6	>10 ⁺		
314	0.119	58.9	0.844 ⁺		
315	0.0753	66.9	>10 ⁺		
316	0.205	66.6	4.92 ⁺		
317	0.272	63.8	>10 ⁺		
318	0.207	64.4	4.9 ⁺		
319	0.0577	70.1	>10 ⁺		
320	0.0105	72.2	0.0961 ⁺		
321	0.0135	79.2	0.241 ⁺		
322	0.215	65.6	>10 ⁺		
323	0.282	58.2	>10 ⁺		
324	0.164	63.1	5.9 ⁺		
325	0.591	63.9	6.09 ⁺		
326	8.11	64.5	>10 ⁺		
327	0.264	63.3	3.54 ⁺		
328	0.422	67.6	2.67 ⁺		
329	0.139	66.5	4.98 ⁺		
330	0.088	61.9	4.87 ⁺		
331	0.194	57.8	3.75 ⁺		
332	4.83	61.5	3.34 ⁺		
333	0.0351	78.1	0.421 ⁺		
334	0.0109	81.5	0.251 ⁺		
335	0.0107	79	0.0511 ⁺	0.0526	>10

336	0.0096	84	0.0846 ⁺	0.146	>10
337	0.0107	81.2	0.0385 ⁺	0.119	>10
338	0.0157	78	0.126 ⁺		
339	0.0107	71	0.0918 ⁺	0.341	5.69
340	0.0236	65.6	0.106 ⁺	0.118	>10
341	0.0062	83.05	0.0346 ⁺	0.122	>10
342	0.0331	73.6	0.0478 ⁺	0.133	>10
343	0.0552	54.4	0.189 ⁺		
344	0.021	74.1	0.153 ⁺		
345	0.0336	77.1	0.108 ⁺	0.362	8.45
346	0.0257	73	0.0438 [*]	0.0716	7.34
347	0.034	72.1	0.0968 [*]	0.16	>10
348	0.0318	74.6	0.228 [*]		
349	0.0237	76.4	0.0582 [*]	0.154	>10
350	0.0539	67.1	0.257 [*]		
351	0.0155	80.6	0.0286 [*]	0.082	>10
352	0.0049	84.1	0.021 [*]	0.05	8.75
353	0.0242	69.2	0.094 [*]	0.111	8.49
354	0.0134	75.5	0.0345 [*]	0.0715	8.06
355	0.0341	69.9	0.362 [*]		
356	0.1484	77.85	0.0448 [*]	0.1325	2.545
357	0.0856	91.7	0.119 [*]		
358	0.0384	84.3	0.0533 [*]	0.129	6.87
359	0.0549	68.3	0.13 [*]		
360	0.0409	56.5	0.0521 [*]	0.235	7.07
361	0.0191	88.8	0.0432 [*]	0.361	6.25
362	0.0443	88.6	0.124 [*]		
363	0.0252	94.3	0.171 [*]		
364	0.0156	79.8	0.387 [*]		
365	0.0205	78.7	0.0339 [*]	0.116	>10
366	0.0467	76.2	0.128 [*]		
367	0.0559	63.5	0.219 [*]		
368	0.118	67.3	0.31 [*]		
369	0.0576	61.5	0.0868 [*]	0.227	>10
370	0.0652	64.3	0.171 [*]		
371	0.14	65.3	0.311 [*]		
372	0.402	49.2	2.12 [*]		
373	0.0743	49.9	0.506 [*]		
374	0.579	60	>10 [*]		
375	0.0346	64.6	0.68 [*]		
376	0.0302	64.1	0.0699 [*]	0.303	>10

377	0.0212	70.6	0.0374 [*]	0.14	>10
378	0.874	80.9	0.361 [*]		
379	0.0418	65.4	0.0566 [*]	0.083	4.71
380	0.0335	68.7	0.0251 [*]	0.0541	3.62
381	0.0552	67.4	0.183 [*]	0.347	0.965
382	0.0142	75.9	0.64 ⁺		
383	0.0101	78.3	0.557 ⁺		
384	0.039	75.5	0.122 ⁺		
385	0.0838	78.1	1.33 ⁺		
386	0.42	59.2	1.97 [*]		
387	0.186	58.2	>10 ⁺		
388	0.381	52.8	6.53 ⁺		
389	0.0195	82.3	0.11 ⁺	0.113	8.04
390	0.0015	80.4	0.167 ⁺	0.0631	>10
391	0.0089	78.5	0.063 ⁺	0.171	>10
392	0.0029	79.3	2.8 ⁺		
393	0.0023	77	0.133 ⁺		
394	0.0248	73.7	4.23 ⁺		
395	0.0066	77.1	0.0238 ⁺	0.117	9.22
396	0.0036	68.9	0.033 ⁺	0.122	>10
397	0.0102	77.2	0.0263 ⁺	0.0465	6.94
398	0.0088	65.9	0.0411 ⁺	0.116	>10
399	0.0114	72.1	0.0344 ⁺	0.0952	8.89
400	0.0237	74.2	0.0388 [*]	0.137	>10
401	0.0106	70.3	0.135 [*]		
402	0.0451	76.8	0.0686 [*]	0.167	7.05
403	0.186	84.5	0.1 [*]	0.23	7.41
404	0.0845	92	0.0356 [*]	0.138	>10
405	0.0052	93.9	0.0261 [*]	0.116	>10
406	0.0551	75.7	0.161 [*]		
407	0.203	105	0.517 [*]		
408	0.0416	63	0.0671 [*]	0.215	8.16
409	0.0297	65.7	0.177 [*]		
410	0.0205	69.3	0.0212 ⁺	0.373	>10
411	0.0416	74.2	0.0741 ⁺	0.142	>10
412	0.0105	72.2	0.0961 ⁺		
413	0.0133	64.5	0.0421 [*]	0.138	2.17
414	0.0506	58.3	0.128 [*]		
415	0.0154	99.9	0.0689 [*]	0.502	7.15
416	0.0135	81.8	0.133 [*]		
417	0.0331	70.1	0.223 [*]		

418	0.017	70.25	0.0674 ⁺	0.139	>10
419	0.0186	78.6	0.121 [*]		
420	0.0177	95	0.0506 [*]	0.353	3.87
421	0.0108	81.6	0.0194 [*]	0.114	>10
422	0.0753	56.2	0.125 [*]		
423	0.0216	74.2	0.0512 [*]	0.269	8.03
424	0.0051	81.7	4.11 ⁺		
425	0.0555	90	0.193 ⁺		
426	0.0585	74.7	0.226 ⁺		
427	0.122	58.4	0.305 ⁺		
428	0.0708	68.9	0.201 ⁺		
429	0.255	59.6	1.18 ⁺		
430	0.142	74.9	1.17 ⁺		
431	0.0086	73.5	0.447 ⁺		
432	0.0121	75.4	0.112 ⁺	0.141	>10
433	0.0241	71.2	1.41 ⁺		
434	0.0511	67.1	2.32 ⁺		
435	0.0087	79.1	1.15 ⁺		
436	0.0157	81.4	0.0895 ⁺	0.272	>10
437	0.0155	84.2	0.0468 ⁺	0.123	>10
438	0.0058	66.9	0.0912 ⁺	0.508	>10
439	0.0187	61.5	0.199 ⁺		
440	0.0376	68.5	0.37 ⁺		
441	0.0265	72.6	0.123 ⁺	0.13	>10
442	0.0093	72.7	0.046 ⁺	0.123	>10
443	0.059	76.6	0.096 ⁺	0.353	4.36
444	0.0391	72.9	0.16 [*]		
445	0.0808	70.5	0.319 [*]		
446	0.115	64.1	0.966 [*]		
447	0.009	74.9	0.0278 ⁺	0.11	>10
448	1.75	66.7	2.73 ⁺		
449	0.336	69.5	0.86 ⁺		
450	0.0791	62.8	0.266 [*]		
451	0.0543	73.2	1.05 ⁺		
452	0.0329	80.5	0.0653 [*]	0.223	>10
453	0.0122	90.7	0.0546 [*]	0.218	6.55
454	0.005	82.9	0.0182 [*]	0.218	5.81
455	0.0342	67.3	0.0576 [*]	0.0886	6.78
456	0.0162	61.5	0.0328 [*]	0.209	1.94
457	0.0202	58.7	0.0853 [*]	0.0761	4.8
458	0.0641	60.2	0.199 [*]		

459	0.0267	65.7	0.0582 [*]	0.0937	2.66
460	0.131	60.9	0.0586 [*]	0.221	3.8
461	0.0539	46.1	0.158 [*]		
462	0.0091	82.4	0.0257 [*]	0.257	>10
463	0.0085	77.1	0.018 [*]	0.0593	>10
464	0.0057	74.8	0.0172 [*]	0.247	>10
465	0.0068	85.3	0.0352 [*]	0.115	>10
466	0.0111	82.2	0.0524 [*]	0.35	4.85
467	0.155	99.3	0.164 [*]		
468	0.132	96	0.269 [*]		
469	0.0225	83.4	0.0838 [*]	0.522	3.79
470	0.125	93.8	0.62 [*]		
471	0.0292	72.1	0.0275 [*]	0.0725	8.87
472	0.211	60.9	0.127 [*]		
473	0.313	54.5	0.426 [*]		
474	0.0117	70.4	0.0135 [*]	0.0872	6.63
475	0.0136	64.7	0.0315 [*]	0.0812	8.77
476	0.312	61.1	0.108 [*]	0.14	4.14
477	0.0468	68	0.0242 [*]	0.0802	2.525
478	0.15	66.8	0.102 [*]	0.259	4.84
479	0.0175	69.6	0.0169 [*]	0.0374	7.41
480	0.146	67.4	0.0558 [*]	0.15	>10
481	0.204	62.1	0.128 [*]		
482	0.0665	63.3	0.0356 [*]	0.0668	1.14
483	0.0673	56.4	0.061 [*]	0.101	1.71
484	0.0335	66.5	0.0318 [*]	0.0662	2.6
485	0.0191	66.5	0.0124 [*]	0.0583	2.06
486	0.0179	61.7	0.0311 [*]	0.112	1.99
487	0.0253	64.3	0.654 [*]		
488	0.0679	61.1	0.285 [*]		
489	0.0299	68.7	0.0225 [*]	0.051	2.06
490	0.041	67.9	0.0198 [*]	0.0762	2.12
491	0.0108	72	0.0207 [*]	0.0591	2.16
492	0.0066	62.3	0.0444 [*]	0.209	>10
493	0.0248	67	0.022 [*]	0.131	>10
494	0.0269	55.9	0.0429 [*]	0.129	>10
495	0.0854	61.8	0.142 [*]		
496	0.0195	71.9	0.0233 [*]	0.0482	2.02
497	0.0316	65.2	0.0437 [*]	0.169	2.49
498	0.038	65.4	0.131 [*]		
499	0.0357	68.1	0.0248 [*]	0.0657	>10

500	0.0669	72	0.0361 [*]	0.0836	5.2
501	0.0709	73.3	0.0933 [*]	0.274	7.26
502	0.0983	63.6	0.175 [*]		
503	0.0397	63.7	9.03 [*]		
504	0.0235	65.9	6.86 [*]		
505	0.0464	54.3	0.219 [*]		
506	0.184	47.4	0.725 [*]		
507	1.51	52.9	1.86 [*]		
508	0.0965	67.1	0.0634 [*]	0.108	6.46
509	0.004	90.9	0.0177 [*]	0.0731	5.89
510	0.0082	83.7	0.0161 [*]	0.123	>10
511	0.0511	89.8	0.0372 [*]	0.149	7.41
512	0.0109	74.3	0.0176 [*]	0.107	6.72
513	0.0777	56.4	0.133 [*]		
514	0.0733	73.8	0.0455 [*]	0.149	>10
515	0.0016	73.9	0.0216 [*]	0.0893	8.78
516	0.204	57.7	0.22 [*]		
517	0.0237	54.1	0.0168 [*]	0.135	2.22
518	0.153	56.7	0.0573 [*]	0.203	>10
519	0.0081	71.2	0.191 [*]		
520	0.0137	70.8	0.0402 ⁺	0.118	>10
521	0.0035	71.8	0.0351 ⁺	0.069	>10
522	0.0408	71.8	0.109 [*]	0.211	>10
523	0.0314	64.1	0.0572 [*]	0.174	5.95
524	0.0329	62.65	0.0766 [*]	0.138	6.29
525	0.0433	63.3	0.043 [*]	0.144	1.35
526	0.0174	61.7	0.0293 [*]	0.133	>10
527	0.0424	65.3	0.109 [*]	0.0654	5.58
528	0.0262	61.5	0.0723 [*]	1.13	6.96
529	0.163	57.3	0.253 [*]		
530	0.427	83.7	>10 [*]		
531	2.6	72.3	>10 [*]		
532	0.0264	70.8	0.0301 [*]	0.1	4.2
533	0.0072	83.2	0.0145 [*]	0.0739	4.81
534	0.0199	78.3	0.0876 [*]	0.0554	>10
535	0.0034	84.2	0.0047 [*]	0.0933	>10
536	0.0453	69.7	0.049 [*]	0.0704	6.8
537	0.0208	63	0.03 [*]	0.0828	6.53
538	0.0092	66.9	0.0574 ⁺	0.124	3.62
539	0.0181	76	0.0162 ⁺	0.0983	7.01
540	0.0174	67.5	0.0741	0.175	7.91

541	0.0151	74.1	0.0377*	0.0498	5.9975
542	0.03	73	0.307*		
543	0.0486	60.2	0.183*		
544	0.059	66.5	0.538*		
545	0.0742	64.7	0.192*		
546	0.0131	73.6	0.109*	1.09	>10
547	0.126	81.1	0.127*	0.173	5.43
548	0.0273	64.6	0.0804*	0.226	6.46
549	0.019	73.6	0.0483*	0.369	>10
550	0.025	99.8	0.105*	0.352	3.64
551	0.105	87.4	0.127*	0.225	3.5
552	0.0166	89.7	0.238*		
553	0.0203	80.2	0.0716*	0.211	>10
554	0.0761	79.7	0.238*		
555	0.0436	61.9	0.0466*	0.121	4.47
556	0.0206	64.8	0.0545*	0.0749	4.52
557	0.0371	59.8	0.0926*	0.131	2.3
558	0.0458	78	2.91*		
559	0.0334	72.175	0.0425*	0.074	3.373333
560	0.051	59	0.148*	0.145	>10
561	0.0159	69.7	0.274*		
562	0.0645	62.3	0.543*		
563	0.0713	66.9	0.327*		
564	0.0701	77.8	0.761*		
565	0.0371	60	0.0565*	0.0582	3.01
566	0.186	66	0.156*		
567	0.0584	77	0.34*		
568	0.0079	77.3	0.0575 ⁺	0.349	>10
569	0.0556	75.9	0.146 ⁺		
570	0.0214	69.8	0.224 ⁺		
571	0.0211	71.3	0.0621 ⁺	0.159	3.56
572	0.0112	73.3	0.0551 ⁺	0.144	>10
573	0.0155	82.8	0.106 ⁺	0.103	>10
574	0.0075	80.6	0.0395 ⁺	0.0922	>10
575	0.0177	76.2	0.215 ⁺		
576	0.0285	68.1	0.128 ⁺		
577	0.0249	91.5	0.153 ⁺		
578	0.149	70.1	0.491 ⁺		
579	0.0311	65.2	0.103*	0.329	>10
580	0.0239	70.5	0.156*		
581	0.0991	50.5	1.68*		

582	0.287	49.9	0.21 [*]		
583	0.0049	72.5	0.0431 [*]		
584	0.017	63.5	0.0421 ⁺	0.122	>10
585	0.0229	70.9	0.0651 [*]	0.208	>10
586	0.149	72.2	0.353 [*]		
587	0.0167	70.9	0.128 [*]		
588	0.103	67.7	0.461 [*]		
589	0.0614	69.7	0.117 [*]		
590	0.0059	74.8	0.0337 [*]	0.533	>10
591	0.0183	65.9	0.13 [*]		
592	0.0541	77.5	0.303 [*]		
593	0.03	99.9	0.17 [*]		
594	0.0109	80.1	0.14 [*]		
595	0.0175	69.35	0.0352 ⁺	0.167	7.4
596	0.0181	72.2	0.038 [*]	0.172	9.88
597	0.022	69.1	0.0648 [*]	0.164	>10
598	0.0164	72.1	0.0506 [*]	0.174	>10
599	0.0192	88.7	0.0378 [*]	0.149	4.33
600	0.0416	79.3	0.186 [*]		
601	0.0212	72.2	0.106 [*]	0.0645	>10
602	0.0578	92.4	0.18 [*]		
603	0.0109	83.4	0.0458 [*]	1.09	8.37
604	0.037	69.1	0.16 ⁺		
605	0.0246	67.5	5.33 ⁺		
606	0.026	71.8	1.51 ⁺		
607	0.0131	75.9	0.126 ⁺		
608	0.0072	75.4	0.422 ⁺		
609	0.003	72.7	0.0047 [*]	0.658	>10
610	0.0287	64.1	0.121 [*]		
611	0.0038	65.3	0.0205 [*]	0.133	8.83
612	0.0101	70	0.0212 [*]	0.281	>10
613	0.0042	70.3	0.116 [*]	0.173	>10
614	0.0146	70.4	0.0269 [*]	0.0531	3.19
615	0.0526	73.9	0.133 [*]	0.0713	7.56
616	0.0114	84.2	0.0173 [*]	0.348	>10
617	0.0252	63.8	0.0321 [*]	0.0579	4.48
618	0.0592	66.6	0.0454 [*]	0.132	1.85
619	0.178	81.3	0.0845 [*]	0.178	3.62
620	0.0037	75.9	0.0062 ⁺	0.0264	>10
621	0.0068	74.4	0.002 ⁺	0.0087	>10
622	0.023	62.9	0.0559 [*]	0.0489	>10

623	0.0039	94.5	0.0112 [*]	0.0723	>10
624	0.0089	97.2	0.442 [*]		
625	0.0126	87.8	1.01 [*]		
626	0.0042	75.7	0.0241 [*]	0.137	>10
627	0.0294	75.8	0.0376 [*]	0.0294	>10
628	0.537	58.1	1.08 [*]		
629	0.0289	74.8	0.112 [*]	0.0692	>10
630	0.0229	67	0.0892 [*]	0.0983	>10
631	0.0312	64.4	0.0593 [*]	0.156	4.88
632	0.241	71.5	0.112 [*]	0.0968	>10
633	0.0526	71.2	0.046 [*]	0.0104	6.46
634	0.0133	68.2	0.0186 [*]	0.0247	8.98
635	0.002	68.5	0.0191 [*]	0.0792	>10
636	0.0122	70.3	0.3 [*]	1.21	>10
637	0.0034	71.4	0.427 [*]		
638	0.0139	70.5	0.0621 [*]		
639	0.0077	59.7	0.0266 [*]		
640	0.007	59	0.0162 [*]		
641	0.009	71	0.0552 [*]		
642	0.0198	67.3	0.0945 [*]		
643	0.0408	61.4	>10 ⁺		
644	0.095	71.2	8.72 ⁺		
645	0.366	70.8	1.26 ⁺		
646	0.0079	66.6	0.0453 [*]	0.0325	>10
647	0.0061	72.9	0.998 [*]		
648	0.0069	69.1	0.471 [*]		
649	0.0503	67.6	9.93 ⁺		
650	0.0524	80.6	0.859 ⁺		
651	0.0706	59.6	0.535 [*]		
652	0.343	68.1	3.77 [*]		
653	0.002	75.7	0.292 [*]		
654	0.0216	72.7	5.31 ⁺		
655	0.0114	67.8	4.52 ⁺		
656	0.0307	71.3	0.239 ⁺		
657	0.0147	71.8	0.817 ⁺		
658	0.0539	73.1	0.425 ⁺		
659	0.0688	50.7	>10 [*]		
660	2.12	67.1	>10 [*]		
661	0.0032	78.6	>10 ⁺		
662	0.068	66.1	0.14 [*]		
663	0.033	66.3	0.124 [*]		

664	0.0877	68.8	0.0719 [*]	0.163	>10
665	0.0309	70.1	4.99 ⁺		
666	0.121	78.4	0.427 [*]		
667	0.145	67.1	0.804 [*]		
668	0.0073	82.3	0.22 [*]		
669	0.0019	80.8	0.124 ⁺		
670	0.0062	73.4	>10 [*]		
671	0.0068	68.8	>10 [*]		
672	0.0978	65.1	0.154 [*]		
673	0.104	70.2	0.13 [*]		
674	0.0039	66.5	0.0289 ⁺	0.119	>10
675	0.0041	71.2	0.0232 ⁺	0.121	>10
676	0.07	68.4	0.806 ⁺		
677	0.0068	75.1	0.0137 [*]		
678	0.0057	71.3	0.0187 [*]	0.391	3.1
679	0.0125	70.2	0.196 [*]		
680	0.003	95	0.031 [*]	0.208	8.76
681	0.0024	77.5	0.133 [*]		
682	0.0835	85.4	0.149 ⁺		
683	1.71	62.5	>10 [*]		
684	8.31	50.6	>10 [*]		
685	0.117	64.3	0.769 ⁺		
686	0.0182	66.6	0.0477 [*]	0.132	4.63
687	0.119	68.1	3.56 ⁺		
688	0.0161	67.9	0.0475 [*]	0.122	6.36
689	0.0036	85	0.34 ⁺		
690	0.0073	75.8	0.0557 ⁺	0.183	>10
691	0.0359	68.5	0.291 ⁺		
692	0.0027	81.5	0.142 ⁺		
693	0.408	64.9	0.858 ⁺		
694	0.0293	73	0.09 [*]	0.0804	7.77
695	0.0085	69.7	0.143 [*]	2.97	>10
696	0.0115	75.4	0.0367 [*]	0.0451	7.7
697	0.0409	59.5	0.329 [*]		
698	1.05	75.4	2.49 ⁺		
699	0.0022	93.6	0.0219 [*]	0.517	>10
700	0.153	85.6	0.871 [*]		
701	0.115	66.9	0.638 [*]		
702	0.0955	70.8	0.0316 [*]	0.224	8.12
703	0.0444	53.6	0.387 [*]		
704	0.0125	78.1	0.0254 [*]	0.0569	7.14

705	0.0192	55.6	0.679 ⁺		
706	0.0196	96.1	0.0517 [*]	0.243	3.88
707	0.0283	67.1	0.0254 [*]	0.0685	5.36
708	0.0691	73	0.0926 [*]	0.14	6.46
709	0.1	60.9	1.24 ⁺		
710	0.0145	69	0.0107 [*]	0.0481	5.4
711	0.0321	69.3	0.17 [*]		
712	0.0562	74.2	0.312 [*]		
713	0.025	76	0.105 ⁺	0.105	>10
714	0.0053	68.7	0.181 [*]		
715	0.0103	67.8	0.0144 [*]	0.0699	9.76
716	0.128	55.9	>10 [*]		
717	0.199	68.1	>10 ⁺		
718	0.0157	72.7	0.137 ⁺	0.114	>10
719	0.0847	69.7	2.28 ⁺		
720	0.0252	72.6	0.0549 [*]	0.452	7.9
721	0.231	68.1	1.57 ⁺		
722	0.0191	74.4	0.0511 ⁺	0.0506	>10
723	0.0016	84.3	0.0735 ⁺	0.733	>10
724	0.0197	80.4	0.224 [*]		
725	0.0365	74	0.625 [*]		
726	0.0175	88.1	0.32 [*]		
727	9.03	55.3	0.71 [*]		
728	0.158	62.5	1.35 [*]		
729	0.114	45.5	0.382 [*]		
730	0.0117	72.6	0.171 [*]		
731	0.0635	69.1	>10 ⁺		
732	0.0105	70.8	0.0261 [*]	0.0341	4.57
733	0.0218	61.3	0.0883 [*]	0.206	>10
734	0.0066	71.2	0.003 [*]	0.0106	6.17
735	0.007	80.1	0.0772 [*]	0.306	>10
736	0.0549	70.3	0.372 [*]		

⁺ indicates the SyBr assay was used and ^{*} indicates the Taqman assay was used.

^{**} Anti-proliferation assay was performed at Pharmaron.

Table 2.

Example No.	circBRIP1 EC50 (μM)	CRC-MSI		CRC-MSS	
		*HCT116 Anti-proliferative IC50 (μM)	*LS411N Anti-proliferative Activity IC50 (μM)	*Colo205 Anti-proliferation IC50 (μM)	*NCI-H747 Anti-proliferative Activity (μM)
15	0.85	1.03	2.57	10.59	23.21

17	0.057	0.11	0.131	11.82	>30
31	0.054	0.045	0.041	7.05	>10

*Anti-proliferation assay was performed at Accent Therapeutics, Inc.

As shown in Table 1, 290 DHX9 inhibitor compounds were screened in the DHX9 cellular proliferation assay in both a CRC-MSI (LS411N) and a CRC-MSS (NCI-H747) cell line. Overall DHX9 compound sensitivity is significantly greater in the CRC-MSI cell line when compared to the CRC-MSS cell line using a unpaired two-tailed t-test, p-value= <0.0001 (FIG. 4).

5. *In Vivo* Efficacy Demonstration for DHX9 Compounds

Experiments were performed in female BALB/c nude mice (GenPharmatech Co.). Animals were allowed to acclimate for 7 days before the study. The general health of the animals were evaluated by a veterinarian, and complete health checks were performed prior to the study. General procedures for animal care and housing were in accordance with the standard, Commission on Life Sciences, National Research Council, Standard Operating Procedures (SOPs) of Pharmaron, Inc. The mice were kept in laminar flow rooms at constant temperature and humidity with 3-5 mice in each cage. Animals were housed in polycarbonate cage which is the size of 300 x 180 x 150 mm³ and in an environmentally monitored, well-ventilated room maintained at a temperature of (23 ± 3°C) and a relative humidity of 40% 70%. Fluorescent lighting provided illumination approximately 12 hours per day. Animals had free access to irradiation sterilized dry granule food during the entire study period except for time periods specified by the protocol, as well as sterile drinking water in a bottle was available ad libitum during the quarantine and study periods.

The LS411N (ATCC; CRL-2159) tumor cell lines were maintained *in vitro* as monolayer in RPMI 1640 medium supplemented with 10% heat inactivated FBS, at 37°C in an atmosphere of 5% CO₂ in air. The tumor cells were sub-cultured, not exceeding 4-5 passages, and cells growing in an exponential growth phase were harvested and counted for tumor inoculation. Each mouse was inoculated subcutaneously on the right flank with LS411N tumor cells (2 × 10⁶) in 0.1 mL of RPMI-1640 with Matrigel (1:1) for model development.

Depending on the target concentration, 14.4 – 144 mg of compound of Example 31 was dissolved in 0.96 mL Solutol while vortexing and sonicating for 1 hour to obtain a suspension. Then 2.4 mL of 1% MC (with 100mg/mL PVP VA64) was added and vortexed for 5 min to obtain homogeneous suspension. The mixture was kept stirring at room temperature for 5 min,

and then 1.44 mL of water was added while vortexing to make the final volume of 4.8 mL. The mixture was kept stirring at RT for 30 min and homogenized for 3 min. The resulting solution was used for *in vivo* studies.

Treatment was started when the mean tumor size reached approximately 100-150 mm³, at which time the mice were randomized into treatment groups. Animals were then treated with vehicle (20% Solutol/ 50% 1% MC (4000 cp) with 100 mg/mL PVP VA64/ 30% water) or indicated mg/kg (30-300) of Example 31 daily BID (12 hour schedule) by oral gavage at a final dosing volume of 10 mL/kg.

All study animals were monitored for not only tumor growth but also behavior such as mobility, food and water consumption (by cage side checking only), body weight (BW), eye/hair matting and any other abnormal effect. Body weights of all animals was measured and recorded twice per week throughout the study. Body weight change, expressed in %, was calculated using the following formula:

$$\text{BW change (\%)} = (\text{BWDay PG-DX} / \text{BWDay PG-D1}) \times 100;$$
 PG-D1 is the first day of dosing.

The measurement of tumor size was conducted with a caliper and recorded twice per week. The tumor volume (TV) (mm³) was estimated using the formula: $\text{TV} = a \times b^2 / 2$, where “a” and “b” are long and short diameters of a tumor, respectively.

The TVs were used for calculation of the tumor growth inhibition and tumor growth delay. For the tumor growth inhibition (TGI), the value using the formula:

$$\%T/C = (\text{TreatedTV}_{\text{final}} - \text{TreatedTV}_{\text{initial}}) / (\text{VehicleTV}_{\text{final}} - \text{VehicleTV}_{\text{initial}}) \times 100$$

$$\%TGI = [1 - (\text{TreatedTV}_{\text{final}} - \text{TreatedTV}_{\text{initial}}) / (\text{VehicleTV}_{\text{final}} - \text{VehicleTV}_{\text{initial}})] \times 100$$

The “TV_{final}” and “TV_{initial}” are the mean tumor volumes on the final day and initial day.

Tumor regression was calculated according to the following formula:

$$-(\text{TreatedTV}_{\text{final}} - \text{TreatedTV}_{\text{initial}} / \text{TreatedTV}_{\text{initial}}) \times 100$$

All statistical tests was conducted on GraphPad, and the level of significance was set at 5% or $P < 0.05$. The group means and standard deviations was calculated for all measurement parameters. One-way Ordinary and Two-way RM ANOVA followed by Tukeys post hoc comparisons of the means was applied among groups.

Treatment was initiated with, for example, compound of Example 31 treated at 30, 100, 200 and 300 mg/kg using BID oral application when the tumor volume was an average of 133 mm³ (n=8/group). The initial treatment period with compound of Example 31 was 21

days, after which overall efficacy and tolerability were evaluated based on tumor volume and body weight changes observed during the treatment period (Fig. 1a and Fig 1c).

The treatment was then continued for an additional 7 days for the vehicle, 300 mg/kg and 200 mg/kg groups. On day 28, Example 31 dosed orally at 300 mg/kg BID induced an antitumor response against LS411N xenografts in nude mice, where the %T/C value was -5.2% and the %TGI was 105.2%, with a p-value= <0.0001 when compared with vehicle control using a one way ordinary ANOVA test (Fig. 1b). The mean tumor volume on day 28 for 300 mg/kg BID showed a -56% regression (Fig. 1c). Based on body weight, BID dosing of all concentrations of compound of Example 31 was well tolerated (Fig. 2).

After 28 days of treatment with compound of Example 31, dosing was stopped and the animals and tumors were monitored for an additional 28 days (Fig. 3a). Vehicle animals were sacrificed on day 38 due to maximum tumor volume reaching above 2500 mm³. Regressed 300 mg/kg BID treated tumors showed little rebound with no compound treatment, going from -56% regression on day 28, to -17% regression on day 56 (Fig. 3b).

6. *Cell Line Panel Proliferation Screen*

Study was performed at Eurofins Discovery. The OncoPanel™ cell proliferation assay measures the proliferation response of cancer cell lines to drug treatments through bioluminescence CellTiter-Glo readout. Cell viability was measured by the bioluminescence signal generated by the production of ATP in viable cells.

Cells were grown in RPMI 1640, 10% FBS, 2 mM L-alanyl-L-glutamine, 1 mM Na pyruvate, or a special medium according to cell vendor recommendations. Cells were seeded into 384-well plates and incubated in a humidified atmosphere of 5% CO₂ at 37° C.

Compounds were added the day following cell seeding. At the same time, a time zero untreated cell plate was generated. After a 10-day incubation period, cells were lysed with cell viability detection reagent CellTiter-Glo® (Promega). At 7 days post-seeding, the growth media were replaced, and the plates were re-dosed with the test compound.

Compound of Example 17 was serially diluted in 3-fold steps from the highest test concentration of 10 μM, and assayed over 10 concentrations with a maximum assay concentration of 0.1% DMSO. Bioluminescence was read by a PerkinElmer Envision® microplate reader.

Cellular response parameters were calculated using nonlinear regression to a sigmoidal single-site dose response model:

$$y = A + \frac{B - A}{1 + (C/x)^D}$$

Where y is a response measured at concentration x, A and B are the lower and upper limits of the response, C is the concentration at the response midpoint (EC₅₀), and D is the Hill Slope1.


- 5 EC₅₀ is the concentration of Example 17 at the curve inflection point or half the effective response (parameter C of the fitted curve solution).

Curve-fitting, calculations, and report generation was performed using a custom data reduction engine and MathIQ based software (AIM).

- 10 A total of 214 cell lines representing 17 tumor indications were screened using Example 17. The results of this assay are shown by plotting the EC₅₀ values across the different indications (FIG. 5a). MSI status of the cell lines within the panel were annotated, DHX9 inhibitor Example 17 showed greater sensitivity in the MSI populations vs MSS populations as represented in colorectal, endometrial, and hematopoietic cancer. CRC-MSI sensitivity over CRC-MSS is statistically significant using an unpaired two-tailed t-test, p-
15 value= <0.00010 (FIG. 5b).

CLAIMS

1. A compound represented by Formula (I):



- 5 or a pharmaceutically acceptable salt thereof, wherein:

X is halo;

Y is O or NR^y;

R^y is H or C₁₋₄alkyl;

- 10 R¹ is C₁₋₄alkyl or C₃₋₆cycloalkyl; wherein the C₁₋₄alkyl or C₃₋₆cycloalkyl are each optionally and independently substituted with 1 to 3 halo or -OH;

Ring A is a C₃₋₆cycloalkyl, 4 to 6-membered monocyclic heterocyclyl, 6 to 10 membered bicyclic heterocyclyl; phenyl, 5 to 6-membered monocyclic heteroaryl, or 8 to 10-membered bicyclic heteroaryl;

m is 0 to 3;

- 15 n is 0 or 2;

each R² is independently selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, halo, OR^{2a}, cyano, -NR^{2b}R^{2c}, -SO₂R^{2a}, -C(O)R^{2d}, and -C(O)NR^{2b}R^{2c}, wherein the C₁₋₄alkyl and C₃₋₆cycloalkyl are optionally substituted with 1 to 4 R^{2e}; or 2 R² together form oxo;

R^{2a} is H, C₁₋₄alkyl, C₁₋₄haloalkyl or C₁₋₄alkoxyC₁₋₄alkyl;

- 20 R^{2d} is H, C₁₋₄alkyl, C₃₋₆cycloalkyl, OR^{2a}, phenyl, 4 to 6-membered monocyclic heterocyclyl, 6 to 8-membered bicyclic heterocyclyl, or 5 to 6-membered monocyclic heteroaryl, wherein the C₁₋₄alkyl is optionally substituted with 1 to 3 halo and the 5- to 6-membered monocyclic heteroaryl is optionally substituted with 1 to 3 C₁₋₄alkyl, C₁₋₄haloalkyl, or phenyl;

- 25 R^{2b} and R^{2c} are each independently selected from H, C₁₋₄alkyl, C₃₋₆cycloalkyl, phenyl, 4 to 6-membered monocyclic heterocyclyl and 5 to 6-membered monocyclic heteroaryl, wherein the C₁₋₄alkyl is optionally substituted with 1 to 3 substituents independently selected from halo and C₁₋₃alkoxy; or R^{2b} and R^{2c}, together with the nitrogen to which they are attached form a 4 to 6 membered monocyclic heterocyclyl

or 6 to 10 membered bicyclic heterocyclyl; wherein the 4 to 6 membered monocyclic heterocyclyl or 6 to 10 membered bicyclic heterocyclyl are each optionally substituted with 1 to 3 R^{2d};

each R^{2e} is independently selected from halo, cyano, NR^{2b}R^{2c}, OR^{2a}, phenyl, and 4 to 6 membered monocyclic heterocyclyl;

Z is a bond, -CH₂-, -O-, -O-C₁₋₄alkylene-*, -C₁₋₄alkylene-O-*, -C(O)-, -C(O)O-*, -OC(O)-*, -S(O)₂-, -S(O)₂N(Z^a)-*, -N(Z^a)S(O)₂-, -N(Z^a)-, -N(Z^a)-C₁₋₄alkylene-*, -C₁₋₄alkylene-N(Z^a)-*, -C(O)N(Z^a)-*, or -C(O)N(Z^a)-C₁₋₃alkylene-*, wherein * indicates the attachment point to R³;

Z^a is H or C₁₋₄alkyl;

R³ is C₃₋₆cycloalkyl, 7 to 10-membered bicyclic carbocyclyl, 4 to 6-membered monocyclic heterocyclyl, 6 to 10-membered bicyclic heterocyclyl, phenyl, 5 to 6-membered monocyclic heteroaryl, or 8 to 10-membered bicyclic heteroaryl, wherein the C₃₋₆cycloalkyl, 7 to 10-membered bicyclic carbocyclyl, 4 to 6-membered monocyclic heterocyclyl, 6 to 10-membered bicyclic heterocyclyl, phenyl, 5 to 6-membered monocyclic heteroaryl, or 8 to 10-membered bicyclic heteroaryl are each optionally and independently substituted with 1 to 4 R⁴;

each R⁴ is independently selected from C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, halo, OR^{4a}, cyano, -NR^{4b}R^{4c}, -C(O)R^{4a}, -C(O)OR^{4a}, -C(O)NR^{4b}R^{4c}, -NR^{4b}C(O)R^{4a}, -NR^{4b}C(O)OR^{4a}, -NR^{4b}SO₂R^{4a}, -SR^{4a}, -S(O)R^{4a}, -SO₂R^{4a}, -SO₂NR^{4b}R^{4c}, -P(O)R^{4b}R^{4c}, phenyl, 5 to 6-membered monocyclic heteroaryl, 4 to 6-membered monocyclic heterocyclyl, and 6 to 10-membered bicyclic heterocyclyl, wherein the C₁₋₄alkyl, C₂₋₄alkenyl, or C₂₋₄alkynyl is optionally substituted with 1 to 4 R^{4d}, and wherein the phenyl, 5 to 6-membered monocyclic heteroaryl, 4 to 6-membered monocyclic heterocyclyl and 6 to 10-membered bicyclic heterocyclyl are each optionally substituted with 1 to 3 R^{4e} and further optionally substituted with 1 or 2 oxo; or two R⁴ together form oxo;

R^{4a} is H, C₁₋₄alkyl optionally substituted with 1 to 4 R^{4d}, -NR^{4b}R^{4c}, C₃₋₆cycloalkyl, 4 to 6-membered monocyclic heterocyclyl, phenyl, or 5 to 6-membered monocyclic heteroaryl; wherein the C₃₋₆cycloalkyl, 4 to 6-membered monocyclic heterocyclyl, phenyl or 5 to 6-membered monocyclic heteroaryl are each optionally and independently substituted with 1 to 3 R^{4e};

R^{4b} and R^{4c} are each independently selected from H, phenyl, 4 to 6-membered monocyclic heterocyclyl, 5 to 6-membered monocyclic heteroaryl, and C₁₋₄alkyl

optionally substituted with 1 to 4 R^{4d} ; or R^{4b} and R^{4c} together with the nitrogen atom to which they are attached to form a 4 to 6-membered monocyclic heterocyclyl;

each R^{4d} is independently selected from halo, OR^{4f} , $-C(O)C_{1-4}alkyl$, $-C(O)NR^{4b}R^{4c}$, $-C(O)C_{1-4}haloalkyl$, $-C(O)OR^{4f}$, $-NR^{4b}R^{4c}$, phenyl, and 5 to 6-membered monocyclic heteroaryl, wherein the phenyl and 5 to 6-membered monocyclic heteroaryl are each optionally substituted with 1 to 3 R^{4g} ;


each R^{4e} is independently selected from halo, $C_{1-4}alkyl$, cyano, OR^{4f} , $-NR^{4b}R^{4c}$, $-C(O)H$, $-C(O)R^{4h}$, $-SO_2C_{1-3}alkyl$, and $-C(O)NR^{4b}R^{4c}$, wherein $C_{1-4}alkyl$ is optionally substituted with 1 to 3 substitutes independently selected from halo, $-SO_2C_{1-3}alkyl$ and $-C(O)NR^{4b}R^{4c}$; or two R^{4e} together form oxo;

R^{4f} is H, $C_{1-4}alkyl$, $C_{1-4}haloalkyl$, phenyl or 5 to 6-membered monocyclic heteroaryl, wherein the phenyl and 5 to 6-membered monocyclic heteroaryl are each optionally substituted with one to three halo;

each R^{4g} is independently selected from halo, OR^{4f} , $C_{1-4}alkyl$, $C_{1-4}haloalkyl$, halo, cyano, $-NR^{4b}R^{4c}$, $-C(O)H$, and $-C(O)OR^{4f}$; and

each R^{4h} is independently $C_{1-4}alkyl$, $C_{3-6}cycloalkyl$, $C_{1-3}alkoxy$ or $-N(R^{4b})_2$.

2. The compound of claim 1, wherein the compound represented by Formula (I):



or a pharmaceutically acceptable salt thereof, wherein:

X is halo;

Y is O or NR^y ;

R^y is H or $C_{1-4}alkyl$;

R^1 is $C_{1-4}alkyl$ or $C_{3-6}cycloalkyl$; wherein the $C_{1-4}alkyl$ or $C_{3-6}cycloalkyl$ are each optionally and independently substituted with 1 to 3 halo;

Ring A is a $C_{3-6}cycloalkyl$, 4 to 6-membered monocyclic heterocyclyl, 6 to 10 membered bicyclic heterocyclyl; phenyl, 5 to 6-membered monocyclic heteroaryl, or 8 to 10-membered bicyclic heteroaryl;

m is 0 to 3;

n is 0 or 1;

each R^2 is independently selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, halo, OR^{2a} , cyano, $-NR^{2b}R^{2c}$, $-SO_2R^{2a}$, $-C(O)R^{2d}$, and $-C(O)NR^{2b}R^{2c}$, wherein the C_{1-4} alkyl and C_{3-6} cycloalkyl are optionally substituted with 1 to 4 R^{2e} ; or 2 R^2 together form oxo;

5 R^{2a} is H, C_{1-4} alkyl, C_{1-4} haloalkyl or C_{1-4} alkoxy C_{1-4} alkyl;

R^{2d} is H, C_{1-4} alkyl, C_{3-6} cycloalkyl, OR^{2a} , phenyl, 4 to 6-membered monocyclic heterocyclyl or 5 to 6-membered monocyclic heteroaryl, wherein the C_{1-4} alkyl is optionally substituted with 1 to 3 halo;

10 R^{2b} and R^{2c} are each independently selected from H, C_{1-4} alkyl, C_{3-6} cycloalkyl, phenyl, 4 to 6-membered monocyclic heterocyclyl and 5 to 6-membered monocyclic heteroaryl; or R^{2b} and R^{2c} , together with the nitrogen to which they are attached form a 4 to 6 membered monocyclic heterocyclyl or 6 to 10 membered bicyclic heterocyclyl; wherein the 4 to 6 membered monocyclic heterocyclyl or 6 to 10 membered bicyclic heterocyclyl are each optionally substituted with 1 to 3 R^{2d} ;

15 each R^{2e} is independently selected from halo, cyano, $NR^{2b}R^{2c}$, OR^{2a} , phenyl, and 4 to 6 membered monocyclic heterocyclyl;

20 Z is a bond, $-CH_2-$, $-O-$, $-O-C_{1-4}$ alkylene-*, $-C_{1-4}$ alkylene-O-*, $-C(O)-$, $-C(O)O-*$, $-OC(O)-*$, $-S(O)_2-$, $-S(O)_2N(Z^a)-*$, $-N(Z^a)S(O)_2-*$, $-N(Z^a)-$, $-N(Z^a)-C_{1-4}$ alkylene-*, $-C_{1-4}$ alkylene- $N(Z^a)-*$, $-C(O)N(Z^a)-*$, or $-C(O)N(Z^a)-C_{1-3}$ alkylene-*, wherein * indicates the attachment point to R^3 ;

Z^a is H or C_{1-4} alkyl;

25 R^3 is C_{3-6} cycloalkyl, 7 to 10-membered bicyclic carbocyclyl, 4 to 6-membered monocyclic heterocyclyl, 6 to 10-membered bicyclic heterocyclyl, phenyl, 5 to 6-membered monocyclic heteroaryl, or 8 to 10-membered bicyclic heteroaryl, wherein the C_{3-6} cycloalkyl, 7 to 10-membered bicyclic carbocyclyl, 4 to 6-membered monocyclic heterocyclyl, 6 to 10-membered bicyclic heterocyclyl, phenyl, 5 to 6-membered monocyclic heteroaryl, or 8 to 10-membered bicyclic heteroaryl are each optionally and independently substituted with 1 to 4 R^4 ;

30 each R^4 is independently selected from C_{1-4} alkyl, halo, OR^{4a} , cyano, $-NR^{4b}R^{4c}$, $-C(O)R^{4a}$, $-C(O)OR^{4a}$, $-C(O)NR^{4b}R^{4c}$, $-NR^{4b}C(O)R^{4a}$, $-NR^{4b}C(O)OR^{4a}$, $-SR^{4a}$, $-SO_2R^{4a}$, $-SO_2NR^{4b}R^{4c}$, $-P(O)R^{4b}R^{4c}$, phenyl, 5 to 6-membered monocyclic heteroaryl, 4 to 6-membered monocyclic heterocyclyl, and 6 to 10-membered bicyclic heterocyclyl, wherein the C_{1-4} alkyl is optionally substituted with 1 to 4 R^{4d} , and wherein the phenyl, 5 to 6-membered monocyclic heteroaryl, 4 to 6-membered monocyclic heterocyclyl

and 6 to 10-membered bicyclic heterocyclyl are each optionally substituted with 1 to 3 R^{4e}; or two R⁴ together form oxo;

R^{4a} is H, C₁₋₄alkyl optionally substituted with 1 to 4 R^{4d}, C₃₋₆cycloalkyl, 4 to 6-membered monocyclic heterocyclyl, phenyl, or 5 to 6-membered monocyclic heteroaryl; wherein the C₃₋₆cycloalkyl, 4 to 6-membered monocyclic heterocyclyl, phenyl or 5 to 6-membered monocyclic heteroaryl are each optionally and independently substituted with 1 to 3 R^{4e};

R^{4b} and R^{4c} are each independently selected from H, phenyl, 4 to 6-membered monocyclic heterocyclyl, 5 to 6-membered monocyclic heteroaryl, and C₁₋₄alkyl optionally substituted with 1 to 4 R^{4d};

each R^{4d} is independently selected from halo, OR^{4f}, -C(O)C₁₋₄alkyl, -C(O)NR^{4b}R^{4c}, -C(O)C₁₋₄haloalkyl, phenyl, and 5 to 6-membered monocyclic heteroaryl, wherein the phenyl and 5 to 6-membered monocyclic heteroaryl are each optionally substituted with 1 to 3 R^{4g};

each R^{4e} is independently selected from halo, C₁₋₄alkyl optionally substituted with 1 to 3 halo, cyano, OR^{4f}, -NR^{4b}R^{4c}, -C(O)H, -C(O)C₁₋₄alkyl, -C₁₋₄alkylene-C(O)NR^{4b}R^{4c}, and -C(O)NR^{4b}R^{4c}; or two R^{4e} together form oxo;

R^{4f} is H, C₁₋₄alkyl, C₁₋₄haloalkyl, phenyl or 5 to 6-membered monocyclic heteroaryl, wherein the phenyl and 5 to 6-membered monocyclic heteroaryl are each optionally substituted with one to three halo;

each R^{4g} is independently selected from halo, OR^{4f}, C₁₋₄alkyl, halo, cyano, -NR^{4b}R^{4c}, -C(O)H, and -C(O)OR^{4f}.


3. The compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein X is Cl, Br, or F.
4. The compound of claim 3, or a pharmaceutically acceptable salt thereof, wherein X is Cl or Br.
5. The compound of claim 3, or a pharmaceutically acceptable salt thereof, wherein X is Cl.
6. The compound of any one of claims 1 to 5, or a pharmaceutically acceptable salt thereof, wherein ring A is selected from phenyl, thiophenyl, pyrrolyl, pyrazoyl,


furanyl, isothiazoyl, thiazoyl, imidazoyl, cyclobutyl, benzofuranyl, 2-oxo-2,3-
 dihydro-1*H*-benzo[*d*]imidazolyl, imidazo[1,2-*a*]pyridin-6-yl, 1,4,5,6-
 tetrahydrocyclopenta[*c*]pyrazolyl, 2-oxo-2,3-dihydro-1*H*-imidazolyl, indoliziny,
 pyrrolo[1,2-*a*]pyrimidinyl, pyrrolo[1,2-*c*]pyrimidinyl, pyrrolo[1,2-*a*]pyraziny, 5,6-
 5 dihydro-4*H*-cyclopenta[*b*]thiophenyl, 5-oxo-5*H*-thiazolo[3,2-*a*]pyridinyl, thieno[3,2-
 b]pyridinyl, thieno[3,2-*c*]pyridinyl, thieno[2,3-*c*]pyridinyl, benzothiophenyl,
 thieno[3,2-*d*]pyrimidinyl, pyrrolo[1,2-*b*]pyridazinyl, pyrrolo[1,2-*a*]pyrazinyl,
 pyrrolo[1,2-*a*]pyrimidinyl, pyrrolo[1,2-*b*]pyrimidinyl, pyrrolo[1,2-*c*]pyrimidinyl, 1-
 oxo-1,2-dihydropyrrolo[1,2-*a*]pyrazinyl, pyrrolo[2,1-*f*][1,2,4]triazinyl, 4,5,6,7-
 10 tetrahydrothieno[2,3-*c*]pyridinyl, 5,6-dihydro-4*H*-cyclopenta[*b*]thiophenyl, and 4,7-
 dihydro-5*H*-thieno[2,3-*c*]pyranyl.

7. The compound of any one of claims 1 to 5, or a pharmaceutically acceptable salt
 thereof, wherein ring A is selected from phenyl, thiophenyl, pyrrolyl, pyrazoyl,
 15 furanyl, isothiazoyl, thiazoyl, imidazoyl, cyclobutyl, benzofuranyl, 2-oxo-2,3-
 dihydro-1*H*-benzo[*d*]imidazolyl, imidazo[1,2-*a*]pyridin-6-yl, 1,4,5,6-
 tetrahydrocyclopenta[*c*]pyrazolyl, 2-oxo-2,3-dihydro-1*H*-imidazolyl, indoliziny,
 pyrrolo[1,2-*a*]pyrimidinyl, pyrrolo[1,2-*c*]pyrimidinyl, pyrrolo[1,2-*a*]pyraziny, 5-oxo-
 5*H*-thiazolo[3,2-*a*]pyridinyl, thieno[3,2-*b*]pyridinyl, thieno[3,2-*c*]pyridinyl,
 20 thieno[2,3-*c*]pyridinyl, benzothiophenyl, thieno[3,2-*d*]pyrimidinyl, pyrrolo[1,2-
 b]pyridazinyl, pyrrolo[1,2-*a*]pyrazinyl, pyrrolo[1,2-*a*]pyrimidinyl, pyrrolo[1,2-*b*]
 pyrimidinyl, pyrrolo[1,2-*c*]pyrimidinyl, 1-oxo-1,2-dihydropyrrolo[1,2-*a*]pyrazinyl,
 pyrrolo[2,1-*f*][1,2,4]triazinyl, 4,5,6,7-tetrahydrothieno[2,3-*c*]pyridinyl, 5,6-dihydro-
 4*H*-cyclopenta[*b*]thiophenyl, and 4,7-dihydro-5*H*-thieno[2,3-*c*]pyranyl.

25


8. The compound of any one of claims 1 to 5, or a pharmaceutically acceptable salt






- 5 9. The compound of any one of claims 1 to 5, or a pharmaceutically acceptable salt


thereof, wherein ring A is represented by the following formula:




10. The compound of claim 8, or a pharmaceutically acceptable salt thereof, wherein ring




5






5

11. The compound of claim 9, or a pharmaceutically acceptable salt thereof, wherein ring




10




12. The compound of claim 6 or 7, or a pharmaceutically acceptable salt thereof, wherein
10 ring A is selected from phenyl, thiophenyl, pyrrolyl, pyrazolyl, furanyl, isothiazoyl,
and imidazolyl.

13. The compound of claim 8 or 9, or a pharmaceutically acceptable salt thereof, wherein

ring A is represented by the following formula:







 ; or
 
 ; each of which is substituted with 0 to 1 R^2 and 1 R^3 .

5

14. The compound of claim 13, or a pharmaceutically acceptable salt thereof, wherein

ring A is represented by the following formula:






 ; or
 
 .

10



15. The compound of any one of claim 1 to 14, or a pharmaceutically acceptable salt thereof, wherein R^2 is selected from $-OCH_3$, cyclopropyl, $-CH_3$, $-CF_3$, $-CH_2CH_3$, $-CH(CH_3)_2$, $-CH(OH)CH_3$, $-C(CH_3)_2OH$, $-CH_2OCH_3$, $-CH_2OH$, $-CH_2CF_3$, $-CH_2CH_2OH$, $-CH_2CH_2CH_2OH$, $-CH_2CN$, $-CH_2CHF_2$, $-CH_2NH_2$, $-N(CH_3)_2$, $-SO_2CH_3$, cyano, halo, $-C(O)OH$, $-C(O)N(CH_3)_2$, and $-C(O)NHCH_3$, or the following structural

formula:





16. The compound of any one of claim 1 to 14, or a pharmaceutically acceptable salt thereof, wherein R^2 is selected from $-\text{CH}_3$, $-\text{CF}_3$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{OH})\text{CH}_3$, $-\text{CH}_2\text{OCH}_3$, $-\text{CH}_2\text{OH}$, $-\text{CH}_2\text{CF}_3$, $-\text{CH}_2\text{CH}_2\text{OH}$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$, $-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{CHF}_2$, $-\text{CH}_2\text{NH}_2$, $-\text{N}(\text{CH}_3)_2$, cyano, halo, $-\text{C}(\text{O})\text{OH}$, $-\text{C}(\text{O})\text{N}(\text{CH}_3)_2$, and –




17. The compound of any one of claims 1 to 9, 12, 13, 15 and 16, or a pharmaceutically acceptable salt thereof, wherein n is 1.
18. The compound of any one of claims 1 to 9, 12, 13, 15, 16 and 17, or a pharmaceutically acceptable salt thereof, wherein n is 1 and Z is a bond.
19. The compound of any one of claims 1 to 9, 12, 13, 15, 16 and 17, or a pharmaceutically acceptable salt thereof, wherein n is 1 and Z is $-\text{CH}_2-$, $-\text{CH}(\text{CH}_3)-$, $-\text{O}-$, $-\text{N}(\text{H})-$, $-\text{N}(\text{C}_{1-4}\text{alkyl})-$, or $-\text{C}(\text{O})\text{NH}-*$, wherein $*$ indicates the attachment point to R^3 .
20. The compound of any one of claims 1 to 19, or a pharmaceutically acceptable salt thereof, wherein R^3 is selected from cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, piperidinyl, piperazinyl, morpholinyl, pyrrolidinyl, phenyl, pyrrolyl, pyrazolyl,


pyridyl, pyrimidyl, pyrazinyl, pyrazinyl, thiophenyl, tetrahydropyranyl,
 tetrahydrofuranyl, oxabicyclo[3.2.1]octanyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl,
 oxadiazolyl, 3-oxo-2,3-dihydro-1H-pyrazolyl, benzamidazolyl, indazolyl, indoyl,
 imidazolidinyl, azetidyl, 2,3-dihydrobenzofuranyl, imidazo[1,2-a]pyridinyl, 2,3-
 5 dihydro-1H-indenyl, 2-oxo-2,3-dihydro-1H-benzo[d]imidazolyl, spiro[3.3]heptanyl,
 2-azaspiro[3.3]heptanyl, and 2-oxaspiro[3.3]heptanyl, each of which is optionally
 substituted with 1 to 3 R⁴.


21. The compound of any one of claims 1 to 19, or a pharmaceutically acceptable salt
 10 thereof, wherein R³ is selected from cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,
 piperidinyl, piperazinyl, morpholinyl, pyrrolidinyl, phenyl, pyrrolyl, pyrazolyl,
 pyridyl, pyrimidyl, pyrazinyl, pyrazinyl, thiophenyl, tetrahydropyranyl,
 tetrahydrofuranyl, oxabicyclo[3.2.1]octanyl, thiazolyl, triazolyl, tetrazolyl,
 oxadiazolyl, 3-oxo-2,3-dihydro-1H-pyrazolyl, benzamidazolyl, indazolyl, indoyl, 2,3-
 15 dihydrobenzofuranyl, imidazo[1,2-a]pyridinyl, 2,3-dihydro-1H-indenyl, 2-oxo-2,3-
 dihydro-1H-benzo[d]imidazolyl, spiro[3.3]heptanyl, 2-azaspiro[3.3]heptanyl, and 2-
 oxaspiro[3.3]heptanyl, each of which is optionally substituted with 1 to 3 R⁴.

22. The compound of claim 19, or a pharmaceutically acceptable salt thereof, wherein Z
 20 is -CH₂- and R³ is selected from thiophenyl, pyrrolyl, and 2-oxo-2,3-dihydro-1H-
 benzo[d]imidazolyl, each of which is optionally substituted with 1 to 2 R⁴.

23. The compound of any one of claims 1 to 22, or a pharmaceutically acceptable salt

thereof, wherein R³ is represented by the following structural formula:  ;







optionally substituted with 1 to 3 R^4 .

24. The compound of any one of claims 1 to 22, or a pharmaceutically acceptable salt

thereof, wherein R^3 is represented by the following structural formula: ;








25. The compound of claim 23, or a pharmaceutically acceptable salt thereof, wherein R³


5


is represented by the following structural formula:






26. The compound of claim 24, or a pharmaceutically acceptable salt thereof, wherein R^3








27. The compound of any one of claims 1 to 19, or a pharmaceutically acceptable salt thereof, wherein R³ is phenyl or 5 to 6-membered monocyclic heteroaryl, each of which is optionally substituted with 1 to 2 R⁴.
28. The compound of claim 27, or a pharmaceutically acceptable salt thereof, wherein R³ is selected from phenyl, pyrazolyl, pyridyl, and pyrimidyl, each of which is optionally substituted with 1 to 2 R⁴.

29. The compound of claim 28, or a pharmaceutically acceptable salt thereof, wherein R^3

is represented by the following structural formula:  ; or




; each of which is optionally substituted with 1 to 2 R⁴.


30. The compound of claim 28, or a pharmaceutically acceptable salt thereof, wherein R^3

is represented by the following structural formula:





$$R^4-\text{CH}=\text{N} \quad ; \text{ or } \quad \text{CH}=\text{N}^{\cdot}$$

31. The compound of claim 28, or a pharmaceutically acceptable salt thereof, wherein R^3



- 5 32. The compound of any one of claims 1 to 31, or a pharmaceutically acceptable salt thereof, wherein:

R^4 is selected from C_{1-3} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, $-OR^{4a}$, $-CN$, $-NH_2$, $-NR^{4b}R^{4c}$, halo, $-C(O)R^{4a}$, $-NHC(O)R^{4a}$, $-NHC(O)OR^{4a}$, $-NR^{4b}SO_2R^{4a}$, $-SR^{4a}$, $-S(O)R^{4a}$, $-SO_2R^{4a}$, $-SO_2NR^{4b}R^{4c}$, $-P(O)R^{4b}R^{4c}$, $-C(O)O-C_{1-4}$ alkyl, benzyl, phenyl, 5 to 6-membered monocyclic heteroaryl, 6 to 10-membered bicyclic heterocyclyl, and 4 to 6-membered monocyclic heterocyclyl, wherein the C_{1-3} alkyl, C_{2-4} alkenyl, or C_{2-4} alkynyl represented by R^4 is optionally substituted with 1 to 3 substituents independently selected from halo, phenyl, 5 to 6-membered monocyclic heteroaryl optionally substituted with 1 or 2 R^{4g} , and 4 to 6-membered monocyclic heterocyclyl optionally substituted with 1 or 2 R^{4g} , $-C(O)NR^{4b}R^{4c}$, $-C(O)R^{4a}$, and OR^{4f} , and the benzyl, phenyl, 5 to 6-membered monocyclic heteroaryl, 6 to 10-membered bicyclic heterocyclyl, and 4 to 6-membered monocyclic heterocyclyl represented by R^4 are each optionally substituted with 1 to 3 R^{4e} and further optionally substituted with one or two oxo, or two R^4 together form an oxo;

20 R^{4a} is H, C_{1-4} alkyl optionally substituted with 1 to 4 R^{4d} , $-NR^{4b}R^{4c}$, C_{3-6} cycloalkyl, phenyl, 4- to 6-membered monocyclic heterocyclyl, or 5 to 6-membered monocyclic heteroaryl, wherein the phenyl, 4- to 6-membered monocyclic heterocyclyl, and 5 to 6-membered monocyclic heteroaryl are optionally substituted with 1 to 3 R^{4e} ;

25 each R^{4b} is independently H or C_{1-4} alkyl;

R^{4c} is H, C_{1-4} alkyl optionally substituted with 1 to 4 R^{4d} , 4 to 6-membered monocyclic heterocyclyl, or 5 to 6-membered monocyclic heteroaryl;

each R^{4d} is independently selected from halo, $-OR^{4f}$, C_{3-6} cycloalkyl, $-C(O)OH$, $-NR^{4b}R^{4c}$, phenyl, and 5 to 6-membered monocyclic heteroaryl, wherein the phenyl

and 5 to 6-membered monocyclic heteroaryl are each optionally substituted with 1 to 3 substituents independently selected from C₁₋₃alkyl, C₁₋₃haloalkyl, halo, CN and OH;

each R^{4e} is independently selected from halo, -OH, C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkoxy, -C(O)R^{4h}, -SO₂C₁₋₃alkyl, -C₁₋₃alkylSO₂C₁₋₃alkyl, -C(O)N(C₁₋₃alkyl)₂, and cyano; or two R^{4e} together form an oxo;

R^{4f} is H, C₁₋₄alkyl, phenyl or 5 to 6-membered monocyclic heteroaryl, wherein phenyl and 5 to 6-membered monocyclic heteroaryl are each optionally substituted with 1 to 3 halo;

each R^{4g} is independently C₁₋₃alkyl or halo; and

each R^{4h} is independently C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₃alkoxyl or -N(R^{4b})₂.

33. The compound of any one of claims 1 to 31, or a pharmaceutically acceptable salt thereof, wherein:

R⁴ is selected from C₁₋₃alkyl, -OR^{4a}, -CN, -NH₂, -NHR^{4c}, halo, -C(O)R^{4a}, -NHC(O)R^{4a}, -NHC(O)OR^{4a}, -SR^{4a}, -SO₂R^{4a}, -SO₂NR^{4b}R^{4c}, -P(O)R^{4b}R^{4c}, -C(O)O-C₁₋₄alkyl, benzyl, phenyl, 5 to 6-membered monocyclic heteroaryl, and 4 to 6-membered monocyclic heterocyclyl, wherein the C₁₋₃alkyl represented by R⁴ is optionally substituted with 1 to 3 substituents independently selected from halo, phenyl, 5 to 6-membered monocyclic heteroaryl, -C(O)NR^{4b}R^{4c}, and OR^{4f}; the benzyl, phenyl, 5 to 6-membered monocyclic heteroaryl, and 4 to 6-membered monocyclic heterocyclyl represented by R⁴ are each optionally substituted with 1 to 3 R^{4e}, or two R⁴ together form an oxo;

R^{4a} is H, C₁₋₄alkyl optionally substituted with 1 to 4 R^{4d}, -NR^{4b}R^{4c} phenyl, or 5 to 6-membered monocyclic heteroaryl, wherein the phenyl and 5 to 6-membered monocyclic heteroaryl are optionally substituted with 1 to 3 R^{4e};

R^{4b} is H or C₁₋₄alkyl;

R^{4c} is H, C₁₋₄alkyl optionally substituted with 1 to 4 R^{4d}, 4 to 6-membered monocyclic heterocyclyl, or 5 to 6-membered monocyclic heteroaryl;


each R^{4d} is independently selected from C₁₋₃alkyl, halo, phenyl and 5 to 6-membered monocyclic heteroaryl, wherein the phenyl and 5 to 6-membered monocyclic heteroaryl are each optionally substituted with 1 to 3 substituents independently selected from halo, CN and OH;


each R^{4e} is independently selected from halo, C₁₋₄alkyl, -C(O)C₁₋₃alkyl, -C(O)N(C₁₋₃alkyl)₂, and cyano; or two R^{4e} together form an oxo;

R^{4f} is H, C_{1-4} alkyl, phenyl or 5 to 6-membered monocyclic heteroaryl, wherein phenyl and 5 to 6-membered monocyclic heteroaryl are each optionally substituted with 1 to 3 halo.

34. The compound of any one of claims 1 to 31, or a pharmaceutically acceptable salt

5

thereof, wherein R^4 is represented by the following structural formula: 





10

substituted with 1 to 3 R^{4e} and further optionally substituted with 1 or 2 oxo

35. The compound of any one of claims 1 to 31, or a pharmaceutically acceptable salt



15








thereof, wherein R^4 is represented by the following structural formula: 




substituted with 1 to 3 R^{4e} .


36. The compound of claim 34, or a pharmaceutically acceptable salt thereof, wherein R⁴

is represented by the following structural formula: ; 





37. The compound of claim 35, or a pharmaceutically acceptable salt thereof, wherein R^4




- 10 38. The compound of any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, wherein each R^{4e} is independently -F, -CH₃, -CF₃, -CH₂CHF₂, -CH₂CF₃, -CH₂SO₂CH₃, -OH, -OCH₃, -SO₂CH₃, -C(O)CH₃, -C(O)cyclopropyl, -C(O)cyclobutyl, -C(O)cyclopentyl, -C(O)CH(CH₃)₂, -C(O)C(CH₃)₃, -C(O)N(CH₃)₂, -C(O)OCH₃, or cyano.


39. The compound of any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, wherein R^{4e} is -CH₃, -C(O)CH₃, -C(O)CH(CH₃)₂ -C(O)N(CH₃)₂, or cyano.


40. The compound of any one of claims 1 to 33, or a pharmaceutically acceptable salt thereof, wherein:
- R^4 is $-OR^{4a}$;
- R^{4a} is C_{1-3} alkyl optionally substituted with 1 to 3 R^{4d} or 4- to 6-membered monocyclic heterocyclyl optionally substituted with 1 to 3 R^{4e} ;
- each R^{4d} is independently halo, C_{1-2} alkyl, phenyl or 5 to 6-membered monocyclic heteroaryl, provided when two or three R^{4d} are present, one of the R^{4d} is phenyl or 5 to 6-membered monocyclic heteroaryl, and the other R^{4d} are each independently halo or C_{1-2} alkyl; wherein the phenyl and 5 to 6-membered monocyclic heteroaryl represented by R^{4d} are each optionally substituted with 1 to 3 substituents independently selected from C_{1-3} alkyl, C_{1-3} haloalkyl and halo; and
- each R^{4e} is independently C_{1-4} alkyl, C_{1-4} haloalkyl, $-SO_2C_{1-3}$ alkyl, or $-C(O)C_{3-6}$ cycloalkyl.
41. The compound of any one of claims 1 to 33, or a pharmaceutically acceptable salt thereof, wherein:
- R^4 is $-OR^{4a}$;
- R^{4a} is C_{1-3} alkyl substituted with 1 to 3 R^{4d} ;
- each R^{4d} is independently halo, C_{1-2} alkyl, phenyl or 5 to 6-membered monocyclic heteroaryl, provided when two or three R^{4d} are present, one of the R^{4d} is phenyl or 5 to 6-membered monocyclic heteroaryl, and the other R^{4d} are each independently halo or C_{1-2} alkyl; wherein the phenyl and 5 to 6-membered monocyclic heteroaryl represented by R^{4d} are each optionally substituted with 1 to 3 halo.
42. The compound of any one of claims 1 to 33, wherein R^4 is $-CH_3$, CH_2OH , $-CHF_2$, $-CF_3$, $-CH(CF_3)OH$, $-CH_2OCH_3$, $-CH_2CN$, $-CH_2CF_3$, $-CH_2CH_2OH$, $-CH_2C(O)N(CH_3)_2$, $-C\equiv CC(CH_3)_2OH$, $-OH$, $-OCH_3$, $-OCH_2CH_3$, $-OCH_2CH_2OCH_3$, $-OCH(CH_3)_2$, $-OCF_3$, $-OCHF_2$, $-OCH_2CF_3$, $-O$ -cyclopropyl, $-O$ -cyclobutyl, $-OC(CH_3)_3$, $-OC(CH_3)_2CH_2OH$, $-OCH(CH_3)C(O)OH$, $-OC(CH_3)_2C(O)OH$, $-OCH(CH_3)CH_2OH$, $-OCH_2CH_2N(CH_3)_2$, $-CN$, $-NH_2$, $-NHCH_3$, halo, $-C(O)H$, $-C(O)CH(CH_3)_2$, $-C(O)OCH_3$, $-C(O)O$ -*t*-butyl, $-N(CH_3)SO_2CH_3$, $-NHC(O)CH_3$, $-NHC(O)$ cyclopropyl, $-NHC(O)OC(CH_3)_3$, $-C(O)CH_3$, $-C(O)OH$, $-C(O)NH_2$, $-C(O)NHCH_3$, $-C(O)N(CH_3)_2$, $-C(O)NC(CH_3)_3$, $-SCH_3$, $-S(O)CH_3$, $-SO_2CH_3$, $-SO_2N(CH_3)_2$, $-P(O)(CH_3)_2$, phenyl, or R^4 is represented by the following structural formula:

5




10







- 5 43. The compound of any one of claims 1 to 33, wherein R⁴ is -CH₃, CH₂OH, -CHF₂, -
 CF₃, -CH(CF₃)OH, -CH₂OCH₃, -CH₂CN, -CH₂CF₃, -CH₂CH₂OH, -CH₂C(O)N(CH₃)₂,
 -OH, -OCH₃, -OCH₂CH₃, -OCH(CH₃)₂, -OCF₃, -CN, -NH₂, halo, -C(O)H, -
 C(O)CH(CH₃)₂, -C(O)OCH₃, -C(O)O-*t*-butyl, -NHC(O)CH₃, -NHC(O)cyclopropyl, -
 NHC(O)OC(CH₃)₃, -C(O)CH₃, -C(O)OH, -C(O)NH₂, -C(O)NHCH₃, -C(O)N(CH₃)₂, -
 10 C(O)NC(CH₃)₃, -SCH₃, -SO₂CH₃, -SO₂N(CH₃)₂, -P(O)(CH₃)₂, phenyl, or R⁴ is
 represented by the following structural formula:



15




44. The compound of any one of claims 1 to 5, 15, 16, 20, 21, and 23 to 43, wherein the compound is represented by Formula (IIA), (IIB), or (IIC):



or a pharmaceutically acceptable salt thereof, wherein m is 0 or 1.


45. The compound of any one of claims 1 to 5, 15, 16, 20, 21, and 23 to 43, wherein the compound is represented by Formula (IIIA) or (IIIB):



or a pharmaceutically acceptable salt thereof.


5

46. The compound of any one of claims 1 to 5, 15, 16, 20, 21, and 23 to 43, wherein the compound is represented by Formula (IVA), (IVB), (IVC), or (IVD):



10 or a pharmaceutically acceptable salt thereof, wherein m is 0 or 1.

47. The compound of any one of claims 1 to 46, or a pharmaceutically acceptable salt thereof, wherein R^1 is $-\text{CH}_3$ or $-\text{CH}_2\text{CH}_3$.
- 15 48. The compound of any one of claims 1 to 47, or a pharmaceutically acceptable salt thereof, wherein R^1 is $-\text{CH}_3$.
49. The compound of claim 1, wherein the compound is represented by Formula (V):



or a pharmaceutically acceptable salt thereof, wherein:

X is Cl or Br;

R¹ is -CH₃ or -CH₂CH₃;

ring A is selected from phenyl, thiophenyl, pyrrolyl, pyrazolyl, furanyl,
isothiazolyl, and imidazolyl;

m is 0 or 1;

each R² is independently selected from C₁₋₃alkyl and -C(O)R^{2d}, wherein the
C₁₋₃alkyl is optionally substituted with R^{2g};


R^{2d} is 4 to 6-membered monocyclic heterocyclyl;

R^{2g} is cyano or OH;

R³ is phenyl or 5 to 6-membered monocyclic heteroaryl, wherein the phenyl or
5 to 6-membered monocyclic heteroaryl are each optionally and independently
substituted with 1 or 2 R⁴;

each R⁴ is independently selected from halo, C₁₋₄alkyl, -O-benzyl, -O-C₁₋₃alkyl,
cyano, -NH₂, and 4 to 6-membered monocyclic heterocyclyl, wherein the
benzyl is optionally substituted with 1 to 3 halo or C₁₋₃haloalkyl.

50. The compound of claim 2, wherein the compound is represented by Formula (V):



or a pharmaceutically acceptable salt thereof, wherein:

X is Cl or Br;

R¹ is -CH₃ or -CH₂CH₃;

ring A is selected from phenyl, thiophenyl, pyrrolyl, pyrazolyl, furanyl,
isothiazolyl, and imidazolyl;

m is 0 or 1;

each R² is independently selected from C₁₋₃alkyl and -C(O)R^{2d}, wherein the
C₁₋₃alkyl is optionally substituted with R^{2g};

R^{2d} is 4 to 6-membered monocyclic heterocyclyl;


R^{2g} is cyano or OH;

R^3 is phenyl or 5 to 6-membered monocyclic heteroaryl, wherein the phenyl or 5 to 6-membered monocyclic heteroaryl are each optionally and independently substituted with 1 or 2 R^4 ;


each R^4 is independently selected from C_{1-4} alkyl, -O-benzyl, cyano, -NH₂, and 4 to 6-membered monocyclic heterocyclyl, wherein the benzyl is optionally substituted with halo.

51. The compound of claim 49 or 50, or a pharmaceutically acceptable salt thereof,

wherein ring A is represented by the following formula:




52. The compound of claim 49 or 50, wherein the compound is represented by Formula (IIB), (IIIA), (IVA); or (IVB):




or a pharmaceutically acceptable salt thereof, wherein m is 0 or 1.

53. The compound of any one of claims 49 to 52, or a pharmaceutically acceptable salt thereof, wherein R^3 is selected from phenyl, pyrazolyl, pyridyl, and pyrimidyl, each of which is optionally substituted with 1 to 2 R^4 .


- 5 54. The compound of any one of claims 49 to 52, or a pharmaceutically acceptable salt



- 10 55. The compound of any one of claims 49 to 52, or a pharmaceutically acceptable salt





56. The compound of any one of claims 49 to 55, wherein R^4 is $-F$, $-OCH(CH_3)_2$, $-OC(CH_3)_3$, $-CH_3$, cyano, $-NH_2$, or R^4 is represented by the following structural
- 15




57. The compound of any one of claims 49 to 55, wherein R^4 is $-\text{CH}_3$, cyano, $-\text{NH}_2$, or R^4


is represented by the following structural formula:

58. The compound of any one of claims 49 to 57, or a pharmaceutically acceptable salt thereof, wherein R^2 is $-\text{CH}_3$, $-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{OH}$, or R^2 is represented by the formula:



59. The compound of claim 49, wherein the compound is represented by Formula (IIB) or (IVB):



or a pharmaceutically acceptable salt thereof, wherein:

m is 1;

R^1 is $-\text{CH}_3$;


X is Cl ;

R^2 is $-\text{CH}_3$;


R^3 is pyrimidinyl or pyridinyl, each of which is optionally substituted with 1 or 2 R^4 ;

R^4 is halo, $-\text{OC}_{1-3}\text{alkyl}$, or 4- to 6-membered monocyclic heterocyclyl, wherein the 4- to 6-membered monocyclic heterocyclyl is optionally substituted with 1 or 2 halo or $\text{C}_{1-3}\text{haloalkyl}$.

60. The compound of claim 59, or a pharmaceutically acceptable salt thereof, wherein R^3



61. The compound of claim 59 or 60, or a pharmaceutically acceptable salt thereof,
5 wherein each R^4 is independently $-F$, $-OCH(CH_3)_2$, or $-OC(CH_3)_3$, or R^4 is



62. The compound of claim 1, wherein the compound is selected from any one of
Examples 1-736, or a pharmaceutically acceptable salt thereof.
- 10 63. A pharmaceutical composition comprising a compound of any one of claims 1 to 62,
or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable
excipient.
- 15 64. A method of treating a DHX9 mediated disease or disorder in a subject, comprising
administering to the subject a compound, or pharmaceutically acceptable salt thereof,
of any one of claims 1 to 62, or the pharmaceutical composition of claim 63.
65. The method of claim 64, wherein the disease or disorder is cancer, viral infection or
20 autoimmune disease.
66. The method of claim 65, wherein the disease or disorder is cancer.
67. The method of claim 66, wherein the cancer is selected from colorectal, endometrial,
25 ovarian, gastric, hematopoietic, breast, brain, skin, lung, blood, prostate, head and
neck, pancreatic, bladder, bone, soft-tissue, kidney, and liver cancer.
68. The method of claim 67, wherein the cancer is selected from colorectal, endometrial,
ovarian, hematopoietic, and gastric cancer.

69. The method of any one of claims 66 to 68, wherein the cancer is a microsatellite instability (MSI) cancer.
- 5 70. The method of any one of claims 66 to 68, wherein the cancer has mutations or defects in DNA mis-match repair (MMR), and/or mutations or defects in RNA splicing and the kinetochore complex.
71. The method of claim 69, wherein the cancer is a microsatellite instability-high (MSI-
10 H) colorectal cancer.
72. The method of claim 66, wherein the cancer is Ewing's Sarcoma.