

DESCRIPTION

INTERLEUKIN-2 PROPROTEINS AND USES THEREOF

1. CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the priority benefit of U.S. provisional application no. 63/349,079, filed June 4, 2022, U.S. provisional application no. 63/355,382, filed June 24, 2022, U.S. provisional application no. 63/387,006, filed December 12, 2022, U.S. provisional application no. 63/481,096, filed January 23, 2023, U.S. provisional application no. 63/493,551, filed March 31, 2023, and U.S. provisional application no. 63/500,997, filed May 9, 2023, the contents of each of which are incorporated herein in their entireties by reference thereto.

2. SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing which has been submitted electronically and is hereby incorporated by reference in its entirety. Said copy, created on May 31, 2023, is named RGN-023WO_SL.xml and is 332,645 bytes in size.

3. BACKGROUND

[0003] Interleukin 2 (IL-2 or IL2) is a pluripotent cytokine produced primarily by CD4+ helper T cells. It stimulates the proliferation and differentiation of T cells, induces the generation of cytotoxic T lymphocytes (CTLs) and the differentiation of peripheral blood lymphocytes to cytotoxic cells and lymphokine-activated killer (LAK) cells, promotes cytokine and cytolytic molecule expression by T cells, facilitates the proliferation and differentiation of B-cells and the synthesis of immunoglobulin by B-cells, and stimulates the generation, proliferation and activation of natural killer (NK) cells (see Waldmann, 2009, Nat Rev Immunol 6:595-601 and Malek, 2008, Annu Rev Immunol 26:453-79).

[0004] Due to its pleotropic effects, IL2 is not optimal for inhibiting tumor growth. The use of IL2 as an antineoplastic agent has been limited by the serious toxicities that accompany the doses necessary for a tumor response. Proleukin® (marketed by Prometheus Laboratories, San Diego, Calif.), is a recombinant form of IL2 that is approved for the treatment of metastatic melanoma and metastatic renal cancer, but its side effects are so severe that its use is only recommended in a hospital setting with access to intensive care. Patients receiving high-dose IL2 treatment frequently experience severe cardiovascular, pulmonary, renal, hepatic, gastrointestinal, neurological, cutaneous, haematological and systemic adverse events, which require intensive monitoring and in-patient management. The major side effect of IL2 therapy is

vascular leak syndrome (VLS), which leads to the accumulation of interstitial fluid in the lungs and liver resulting in pulmonary edema and liver damage. There is no treatment for VLS other than withdrawal of IL2. Low-dose IL2 regimens have been tested in patients to avoid VLS, however, at the expense of suboptimal therapeutic results. It has been shown that IL2-induced pulmonary edema resulted from direct binding of IL2 to lung endothelial cells, which express low to intermediate levels of functional high affinity IL2 receptors (Krieg *et al.*, 2010, Proc Nat Acad Sci USA 107:11906-11).

[0005] A variety of IL2 variants and prodrugs have been generated with the aim of reducing the toxicity of IL2 cancer therapy. However, it has been surprisingly discovered that such molecules have poor therapeutic indices for cancer therapy. For example, the PEGylated IL2 prodrug bempegaldesleukin failed to improve on the therapeutic efficacy of a PD1 checkpoint inhibitor in melanoma patients in phase 3 clinical studies (Mullard, 2022, Nature Reviews Drug Discovery 21:327 (doi: <https://doi.org/10.1038/d41573-022-00069-3>)).

[0006] Thus, there is a need in the art for novel IL2 therapies with improved therapeutic efficacy and safety profiles.

4. SUMMARY

[0007] The present disclosure relates to IL2 proproteins that are activated by proteases, *e.g.*, proteases expressed in the tumor environment.

[0008] The IL2 proproteins comprise an IL2 moiety that is masked by an IL2R α moiety, configured so the mask is released following cleavage by a protease. The IL2 proproteins preferably further comprise a targeting moiety that directs the IL2 proprotein to a particular tissue or cell type.

[0009] In certain embodiments, the IL2 proproteins of the disclosure comprise two polypeptide chains, each comprising, from N- to C-terminus, an Fc domain, a first linker which may be cleavable or non-cleavable, an IL2 moiety, a second linker that is protease-cleavable, and an IL2R α moiety. The IL2 proproteins may further comprise, *e.g.*, N-terminal to one or both Fc domains, a targeting moiety (or a component thereof, *e.g.*, one chain of a Fab). The targeting moiety comprises an antigen-binding domain ("ABD") that can, for example, bind to a target molecule present on the tumor surface (*e.g.*, a tumor associated antigen) or other component in the tumor microenvironment (*e.g.*, extracellular matrix ("ECM") or tumor lymphocytes).

[0010] Exemplary IL2 moieties that can be used in the IL2 proproteins of the disclosure are described in Section 6.3.

[0011] Exemplary IL2R α moieties that can be used in the IL2 proproteins of the disclosure are described in Section 6.4.

5 [0012] Protease-cleavable linkers that can be used in the IL2 proproteins of the disclosure are described in Section 6.5.

[0013] Non-cleavable linkers that can be used in the IL2 proproteins of the disclosure are described in Section 6.6.

10 [0014] Targeting moieties that can be used in the IL2 proproteins of the disclosure are described in Section 6.7 and targeting moiety formats are disclosed in Section 6.8.

[0015] Fc domains that can be incorporated into the IL2 proproteins of the disclosure are described in Section 6.9.

[0016] Exemplary IL2 proproteins of the disclosure are described in Section 6.2 and numbered embodiments 1 to 165.

15 [0017] The disclosure further provides nucleic acids encoding the IL2 proproteins of the disclosure. The nucleic acids encoding the IL2 proproteins can be a single nucleic acid (*e.g.*, a vector encoding all polypeptide chains of an IL2 proprotein) or a plurality of nucleic acids (*e.g.*, two or more vectors encoding the different polypeptide chains of an IL2 proprotein). The disclosure further provides host cells and cell lines engineered to express the nucleic acids and
20 IL2 proproteins of the disclosure. The disclosure further provides methods of producing an IL2 proprotein of the disclosure. Exemplary nucleic acids, host cells, and cell lines, and methods of producing an IL2 proprotein are described in Section 6.10 and numbered embodiments 166 to 168.

25 [0018] The disclosure further provides pharmaceutical compositions comprising the IL2 proproteins of the disclosure. Exemplary pharmaceutical compositions are described in Section 6.11 and numbered embodiment 169.

[0019] Further provided herein are methods of using the IL2 proproteins and the pharmaceutical compositions of the disclosure, *e.g.*, for treating cancer. Exemplary methods are described in Section 6.12 and numbered embodiments 170 to 208.

5. BRIEF DESCRIPTION OF THE FIGURES

[0020] FIGS. 1A-1B. FIG. 1A is an illustration of an exemplary targeted IL2 proprotein comprising four protease-cleavable linkers. Although the targeting moieties in FIG. 1A are illustrated as Fabs, the targeting moieties can be in other formats, *e.g.*, scFvs or other formats described in Section 6.8. FIGS. 1A-1 through 1A-4 illustrate a close-up view of an embodiment of Linkers A, B, C and D, respectively comprising a spacer (A1, A2, B1, B2, C1, C2 and D1, D2, respectively) on either side of a cleavable substrate. The number of substrate and spacer sequences is for illustrative purposes only, and it is expected that the protease cleavable linkers will typically have multiple substrate and spacer sequences as detailed in Section 6.5. FIG. 1B illustrates the mechanism of activation of an exemplary targeted IL2 proprotein according to FIG. 1A for which the targeting moiety binds to a TAA. Targeting moieties that bind to other targets as disclosed herein may be used.

[0021] FIGS. 2A-2D. FIG. 2A is an illustration of an exemplary targeted IL2 proprotein comprising two protease-cleavable linkers. Although the targeting moieties in FIG. 2A are illustrated as Fabs, the targeting moieties can be in other formats, *e.g.*, scFvs or other formats described in Section 6.8. FIGS. 2A-1 and 2A-2 illustrate a close-up view of an embodiment of Linkers B and D, respectively comprising a spacer (B1, B2 and D1, D2, respectively) on either side of a cleavable substrate. The number of substrate and spacer sequences is for illustrative purposes only, and it is expected that the protease cleavable linkers will typically have multiple substrate and spacer sequences as detailed in Section 6.5. FIGS. 2B-2D illustrate the mechanism of activation of an exemplary targeted IL2 proprotein according to FIG. 2A for which the targeting moiety binds to a TCA. The resulting activated IL2 protein may bind to the IL2 receptor on a T-cell via one or both IL2 moieties (as illustrated in FIG. 2B), to the TCA on a T-cell via one or both targeting moieties (as illustrated in FIG. 2C), or may simultaneously bind to the IL2 receptor on a T-cell via one or both IL2 moieties and to the TCA on the same T-cell via one or both targeting moieties (as illustrated in FIG. 2D). Targeting moieties that bind to other targets as disclosed herein may be used.

[0022] FIGs. 3A-3E show the *in vivo* anti-tumor efficacy of EGFR-targeted or non-targeted IL2 proproteins comprising cleavable and non-cleavable linkers. FIG. 3A is a graph that compares mean anti-tumor efficacies of EGFR-targeted IL2 proproteins, non-targeted IL2 proproteins, and isotype controls. Each of FIGs. 3B-3E display the anti-tumor efficacy of a single IL2 proprotein or control in individual mice.

5 [0023] FIGs. 4A-4D show the *in vivo* anti-tumor efficacy of PD1-targeted or non-targeted IL2 proproteins comprising cleavable and non-cleavable linkers. FIG. 4A is a graph that compares mean anti-tumor efficacies of PD1-targeted IL2 proproteins, non-targeted IL2 proproteins, and isotype controls. Each of FIGs. 4B-4D display the anti-tumor efficacy of a single IL2 proprotein or control in individual mice.

[0024] FIGs. 5A-5B show an exemplary western blot displaying uPA-digested and non-digested IL2 proprotein samples. FIG. 5A is a western blot image with samples loaded as identified in FIG. 5B.

10 [0025] FIGs. 6A-6F show *in vitro* activity of tumor-targeted IL2 proproteins comprising protease-cleavable and non-cleavable linkers in engineered CD25 KO/ PD1 KO YT/STAT5-Luc reporter cells. FIGs. 6A-6D are graphs that show the luciferase activity associated with CA9-targeted IL2 proproteins, where each shows the activity of IL2 proproteins comprising a different CD9 targeting moiety (FIG. 6A - aCD9(Ab1), FIG. 6B - aCD9(Ab2), FIG. 6C - aCD9(Ab3), and FIG. 6D - aCD9(Ab4)). FIGs. 6E and 6F show the luciferase activity associated with EGFR-targeted and PD1-targeted IL2 proproteins, respectively.

20 [0026] FIGs. 7A-7F show *in vitro* activity of tumor-targeted IL2 proproteins comprising protease-cleavable and non-cleavable linkers in engineered CD25 OE/ PD1 KO YT/STAT5-Luc reporter cells. FIGs. 7A-7D are graphs that show the luciferase activity associated with CD9-targeted IL2 proproteins, where each shows the activity of IL2 proproteins comprising a different CD9 targeting moiety (FIG. 7A - aCD9(Ab1), FIG. 7B - aCD9(Ab2), FIG. 7C - aCD9(Ab3), FIG. 7D - aCD9(Ab4)). FIGs. 7E and 7F show the luciferase activity associated with EGFR-targeted and PD1-targeted IL2 proproteins, respectively.

25 [0027] FIGs. 8A-8D show *in vitro* activity of PD1-targeted IL2 proproteins comprising non-cleavable linkers of different lengths. FIG. 8A is a graph showing the activity of IL2 proproteins in PD1 OE/ CD25 KO YT/STAT5-Luc reporter cells. FIG. 8B is a graph showing the activity of IL2 proproteins in PD1 KO/ CD25 KO YT/STAT5-Luc reporter cells. FIG. 8C is a graph showing the activity of IL2 proproteins in PD1 OE/ CD25 OE YT/STAT5-Luc reporter cells. FIG. 8D is a graph showing the activity of IL2 proproteins in PD1 KO/ CD25 OE YT/STAT5-Luc reporter cells.

6. DETAILED DESCRIPTION

6.1. DEFINITIONS

[0028] As used herein, the following terms are intended to have the following meanings:

[0029] **ABD chain, targeting moiety chain:** Targeting moieties and antigen binding sites

5 (ABD's) within them can exist as one (e.g., in the case of an scFv or scFab) polypeptide chain or form through the association of more than one polypeptide chains (e.g., in the case of a Fab or an Fv). As used herein, the terms "ABD chain" and "targeting moiety chain" refer to all or a portion of an ABD or targeting moiety that exists on a single polypeptide chain. The use of the term "ABD chain" or "targeting moiety chain" is intended for convenience and descriptive
10 purposes only and does not connote a particular configuration or method of production. Further, the reference to an ABD or targeting moiety when describing an IL2 proprotein encompasses an ABD chain or targeting moiety chain unless the context dictates otherwise. Thus, when describing an IL2 proprotein in which an Fc domain is operably linked to a targeting moiety, the Fc domain may be covalently linked directly or indirectly (e.g., via a linker) through a peptide
15 bond to, e.g., (1) a first ABD or targeting moiety chain of a Fab or Fv (with the other components of the Fab or Fv on a second, associated ABD or targeting moiety chain) or (2) an ABD or targeting moiety chain containing an scFv or scFab.

[0030] **About, Approximately:** The terms "about", "approximately" and the like are used throughout the specification in front of a number to show that the number is not necessarily
20 exact (e.g., to account for fractions, variations in measurement accuracy and/or precision, timing, etc.). It should be understood that a disclosure of "about X" or "approximately X" where X is a number is also a disclosure of "X." Thus, for example, a disclosure of an embodiment in which one sequence has "about X% sequence identity" to another sequence is also a disclosure of an embodiment in which the sequence has "X% sequence identity" to the other sequence.

25 [0031] **Activate, activation:** The terms "activation", "activation", and the like in conjunction with an IL2 proprotein of the disclosure refers to the protease-mediated enzymatic cleavage of a protease-cleavable linker that results in the unmasking or release of an IL2 moiety from an IL2R α moiety.

[0032] **And, or:** Unless indicated otherwise, an "or" conjunction is intended to be used in its
30 correct sense as a Boolean logical operator, encompassing both the selection of features in the alternative (A or B, where the selection of A is mutually exclusive from B) and the selection of features in conjunction (A and B, where both A and B are selected). In some places in the text,

the term “and/or” is used for the same purpose, which shall not be construed to imply that “or” is used with reference to mutually exclusive alternatives.

[0033] Antibody: The term “antibody” as used herein refers to a polypeptide (or set of polypeptides) of the immunoglobulin family that is capable of binding an antigen non-covalently, reversibly and specifically. For example, a naturally occurring “antibody” of the IgG type is a tetramer comprising at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as VL) and a light chain constant region. The light chain constant region is comprised of one domain (abbreviated herein as CL). The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The variable regions of the heavy and light chains contain a binding domain that interacts with an antigen. The constant regions of the antibodies may mediate the binding of the immunoglobulin to host tissues or factors, including various cells of the immune system (*e.g.*, effector cells) and the first component (C1q) of the classical complement system. The term “antibody” includes, but is not limited to, monoclonal antibodies, human antibodies, humanized antibodies, camelized antibodies, chimeric antibodies, bispecific or multispecific antibodies and anti-idiotypic (anti-id) antibodies. The antibodies can be of any isotype/class (*e.g.*, IgG, IgE, IgM, IgD, IgA and IgY) or subclass (*e.g.*, IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2). Both the light and heavy chains are divided into regions of structural and functional homology. The terms “constant” and “variable” are used functionally. In this regard, it will be appreciated that the variable domains of both the light (VL) and heavy (VH) chain portions determine antigen recognition and specificity. Conversely, the constant domains of the light chain (CL) and the heavy chain (CH1, CH2 or CH3) confer important biological properties such as secretion, transplacental mobility, Fc receptor binding, complement binding, and the like. By convention the numbering of the constant region domains increases as they become more distal from the antigen-binding domain or amino-terminus of the antibody. The N-terminus is a variable region and at the C-terminus is a constant region; the CH3 and CL domains represent the carboxy-terminus of the heavy and light chain, respectively, of natural antibodies. For convenience, and unless the context dictates otherwise, the reference to an antibody also refers to antibody fragments as

well as engineered antibodies that include non-naturally occurring antigen-binding domains and/or antigen-binding domains having non-native configurations.

[0034] Antigen-binding domain: The term “antigen-binding domain” or “ABD” as used herein refers to a portion of an antibody or antibody fragment (*e.g.*, a targeting moiety) that has the ability to bind to an antigen non-covalently, reversibly and specifically. Examples of an antibody fragment that can comprise an ABD include, but are not limited to, a single-chain Fv (scFv), a Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains; a F(ab)₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; a Fd fragment consisting of the VH and CH1 domains; a Fv fragment consisting of the VL and VH domains of a single arm of an antibody; a dAb fragment (Ward *et al.*, 1989, Nature 341:544-546), which consists of a VH domain; and an isolated complementarity determining region (CDR). Thus, the term “antibody fragment” encompasses both proteolytic fragments of antibodies (*e.g.*, Fab and F(ab)₂ fragments) and engineered proteins comprising one or more portions of an antibody (*e.g.*, an scFv). Antibody fragments can also be incorporated into single domain antibodies, maxibodies, minibodies, intrabodies, diabodies, triabodies, tetrabodies, v-NAR and bis-scFv (see, *e.g.*, Hollinger and Hudson, 2005, Nature Biotechnology 23: 1126-1136).

[0035] Associated: The term “associated” in the context of an IL2 proprotein refers to a functional relationship between two or more polypeptide chains. In particular, the term “associated” means that two or more polypeptides are associated with one another, *e.g.*, non-covalently through molecular interactions or covalently through one or more disulfide bridges or chemical cross-linkages, so as to produce a functional IL2 proprotein. Examples of associations that might be present in an IL2 proprotein of the disclosure include (but are not limited to) associations between Fc domains to form an Fc region (homodimeric or heterodimeric as described in Section 6.9), associations between VH and VL regions in a Fab or Fv, and associations between CH1 and CL in a Fab.

[0036] Cancer: The term “cancer” refers to a disease characterized by the uncontrolled (and often rapid) growth of aberrant cells. Cancer cells can spread locally or through the bloodstream and lymphatic system to other parts of the body. Examples of various cancers are described herein and include but are not limited to, breast cancer, prostate cancer, ovarian cancer, cervical cancer, skin cancer, pancreatic cancer, colorectal cancer, renal cancer, liver cancer, brain cancer, adrenal gland cancer, autonomic ganglial cancer, biliary tract cancer, bone cancer, endometrial cancer, eye cancer, fallopian tube cancer, genital tract cancers, large

intestinal cancer, cancer of the meninges, oesophageal cancer, peritoneal cancer, pituitary cancer, penile cancer, placental cancer, pleura cancer, salivary gland cancer, small intestinal cancer, stomach cancer, testicular cancer, thymus cancer, thyroid cancer, upper aerodigestive cancers, urinary tract cancer, vaginal cancer, vulva cancer, lymphoma, leukemia, lung cancer and the like, e.g., any TAA-positive cancers of any of the foregoing types.

[0037] Complementarity Determining Region: The terms “complementarity determining region” or “CDR,” as used herein, refer to the sequences of amino acids within antibody variable regions which confer antigen specificity and binding affinity. For example, in general, there are three CDRs in each heavy chain variable region (e.g., CDR-H1, CDR-H2, and CDR-H3) and three CDRs in each light chain variable region (CDR-L1, CDR-L2, and CDR-L3). The precise amino acid sequence boundaries of a given CDR can be determined using any of a number of well-known schemes, including those described by Kabat *et al.*, 1991, “Sequences of Proteins of Immunological Interest,” 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (“Kabat” numbering scheme), Al-Lazikani *et al.*, 1997, JMB 273:927-948 (“Chothia” numbering scheme) and ImMunoGenTics (IMGT) numbering (Lefranc, 1999, The Immunologist 7:132-136; Lefranc *et al.*, 2003, Dev. Comp. Immunol. 27:55-77 (“IMGT” numbering scheme). For example, for classic formats, under Kabat, the CDR amino acid residues in the heavy chain variable domain (VH) are numbered 31-35 (CDR-H1), 50-65 (CDR-H2), and 95-102 (CDR-H3); and the CDR amino acid residues in the light chain variable domain (VL) are numbered 24-34 (CDR-L1), 50-56 (CDR-L2), and 89-97 (CDR-L3). Under Chothia, the CDR amino acids in the VH are numbered 26-32 (CDR-H1), 52-56 (CDR-H2), and 95-102 (CDR-H3); and the amino acid residues in VL are numbered 26-32 (CDR-L1), 50-52 (CDR-L2), and 91-96 (CDR-L3). By combining the CDR definitions of both Kabat and Chothia, the CDRs consist of amino acid residues 26-35 (CDR-H1), 50-65 (CDR-H2), and 95-102 (CDR-H3) in human VH and amino acid residues 24-34 (CDR-L1), 50-56 (CDR-L2), and 89-97 (CDR-L3) in human VL. Under IMGT the CDR amino acid residues in the VH are numbered approximately 26-35 (CDR-H1), 51-57 (CDR-H2) and 93-102 (CDR-H3), and the CDR amino acid residues in the VL are numbered approximately 27-32 (CDR-L1), 50-52 (CDR-L2), and 89-97 (CDR-L3) (numbering according to “Kabat”). Under IMGT, the CDR regions of an antibody can be determined using the program IMGT/DomainGap Align.

[0038] Effector Function: The term “effector function” refers to an activity of an antibody molecule that is mediated by binding through a domain of the antibody other than the antigen-binding domain, usually mediated by binding of effector molecules. Effector function includes

complement-mediated effector function, which is mediated by, for example, binding of the C1 component of the complement to the antibody. Activation of complement is important in the opsonization and lysis of cell pathogens. The activation of complement also stimulates the inflammatory response and may also be involved in autoimmune hypersensitivity. Effector

5 function also includes Fc receptor (FcR)-mediated effector function, which may be triggered upon binding of the constant domain of an antibody to an Fc receptor (FcR). Binding of antibody to Fc receptors on cell surfaces triggers a number of important and diverse biological responses including engulfment and destruction of antibody-coated particles, clearance of immune complexes, lysis of antibody-coated target cells by killer cells (called antibody- dependent cell-

10 mediated cytotoxicity, or ADCC), release of inflammatory mediators, placental transfer and control of immunoglobulin production. An effector function of an antibody may be altered by altering, e.g., enhancing or reducing, the affinity of the antibody for an effector molecule such as an Fc receptor or a complement component. Binding affinity will generally be varied by modifying the effector molecule binding site, and in this case it is appropriate to locate the site of

15 interest and modify at least part of the site in a suitable way. It is also envisaged that an alteration in the binding site on the antibody for the effector molecule need not alter significantly the overall binding affinity but may alter the geometry of the interaction rendering the effector mechanism ineffective as in non-productive binding. It is further envisaged that an effector function may also be altered by modifying a site not directly involved in effector molecule

20 binding, but otherwise involved in performance of the effector function.

[0039] Epitope: An epitope, or antigenic determinant, is a portion of an antigen recognized by an antibody or other antigen-binding moiety as described herein. An epitope can be linear or conformational.

[0040] Fab: The term "Fab" refers to a pair of polypeptide chains, the first comprising a variable

25 heavy (VH) domain of an antibody operably linked (typically N-terminal to) to a first constant domain (referred to herein as C1), and the second comprising variable light (VL) domain of an antibody N-terminal operably linked (typically N-terminal) to a second constant domain (referred to herein as C2) capable of pairing with the first constant domain. In a native antibody, the VH is N-terminal to the first constant domain (CH1) of the heavy chain and the VL is N-terminal to the

30 constant domain of the light chain (CL). The Fabs of the disclosure can be arranged according to the native orientation or include domain substitutions or swaps that facilitate correct VH and VL pairings. For example, it is possible to replace the CH1 and CL domain pair in a Fab with a CH3-domain pair to facilitate correct modified Fab-chain pairing in heterodimeric molecules. It is

also possible to reverse CH1 and CL, so that the CH1 is attached to VL and CL is attached to the VH, a configuration generally known as Crossmab. The term “Fab” encompasses single chain Fabs.

- 5 **[0041] Fc Domain and Fc Region:** The term “Fc domain” refers to a portion of the heavy chain that pairs with the corresponding portion of another heavy chain. The term “Fc region” refers to the region formed by association of two heavy chain Fc domains. The two Fc domains within the Fc region may be the same or different from one another. In a native antibody the Fc domains are typically identical, but one or both Fc domains might be modified to allow for heterodimerization, *e.g.*, via a knob-in-hole interaction.
- 10 **[0042] Fv:** The term “Fv” refers to the minimum antibody fragment derivable from an immunoglobulin that contains a complete target recognition and binding site. This region consists of a dimer of one heavy and one light chain variable domain in a tight, noncovalent association (VH-VL dimer). It is in this configuration that the three CDRs of each variable domain interact to define a target binding site on the surface of the VH-VL dimer. Often, the six
- 15 CDRs confer target binding specificity to the antibody. However, in some instances even a single variable domain (or half of an Fv comprising only three CDRs specific for a target) can have the ability to recognize and bind target. The reference to a VH-VL dimer herein is not intended to convey any particular configuration. When present on a single polypeptide chain (*e.g.*, a scFv), the VH and be N- terminal or C-terminal to the VL.
- 20 **[0043] Half Antibody:** The term “half antibody” refers to a molecule that comprises at least one Fc domain and can associate with another molecule comprising an Fc through, *e.g.*, a disulfide bridge or molecular interactions. A half antibody can be composed of one polypeptide chain or more than one polypeptide chains (*e.g.*, the two polypeptide chains of a Fab). An example of a half antibody is a molecule comprising a heavy and light chain of an antibody (*e.g.*, an IgG
- 25 antibody). Another example of a half antibody is a molecule comprising a first polypeptide comprising a VL domain and a CL domain, and a second polypeptide comprising a VH domain, a CH1 domain, a hinge domain, a CH2 domain, and a CH3 domain, wherein said VL and VH domains form an ABD. Yet another example of a half antibody is a polypeptide comprising an scFv domain, a CH2 domain and a CH3 domain. The IL2 proproteins of the disclosure typically
- 30 comprise two half antibodies, each comprising an Fc domain, an IL2R α moiety C-terminal to the Fc domain, a protease-cleavable linker C-terminal to the IL2R α moiety, and an IL2 moiety C-terminal to the protease-cleavable linker. One or both half antibodies in the IL2 proproteins may further comprise a targeting moiety, *e.g.*, N-terminal to the Fc domain.

[0044] The term “half antibody” is intended for descriptive purposes only and does not connote a particular configuration or method of production. Descriptions of a half antibody as a “first” half antibody, a “second” half antibody, a “left” half antibody, a “right” half antibody or the like are merely for convenience and descriptive purposes.

5 [0045] **Host cell or recombinant host cell:** The terms “host cell” or “recombinant host cell” refer to a cell that has been genetically-engineered, *e.g.*, through introduction of a heterologous nucleic acid. It should be understood that such terms are intended to refer not only to the particular subject cell but to the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny
10 may not, in fact, be identical to the parent cell, but are still included within the scope of the term “host cell” as used herein. A host cell may carry the heterologous nucleic acid transiently, *e.g.*, on an extrachromosomal heterologous expression vector, or stably, *e.g.*, through integration of the heterologous nucleic acid into the host cell genome. For purposes of expressing a IL2 proprotein of the disclosure, a host cell is preferably a cell line of mammalian origin or
15 mammalian-like characteristics, such as monkey kidney cells (COS, *e.g.*, COS-1, COS-7), HEK293), baby hamster kidney (BHK, *e.g.*, BHK21), Chinese hamster ovary (CHO), NSO, PerC6, BSC-1, human hepatocellular carcinoma cells (*e.g.*, Hep G2), SP2/0, HeLa, Madin-Darby bovine kidney (MDBK), myeloma and lymphoma cells, or derivatives and/or engineered variants thereof. The engineered variants include, *e.g.*, derivatives that grow at higher density
20 than the original cell lines and/or glycan profile modified derivatives and and/or site-specific integration site derivatives.

[0046] **Linker:** The term “linker” as used herein refers to a protease-cleavable linker or a non-cleavable linker.

25 [0047] **Non-cleavable linker:** A non-cleavable linker refers to a peptide whose amino acid sequence lacks a substrate sequence for a protease, *e.g.*, a protease as described in Section 6.5.1, that recognizes and cleaves a specific sequence motif, *e.g.*, a substrate as described in Section 6.5.2.

30 [0048] **Operably linked:** The term “operably linked” refers to a functional relationship between two or more peptide or polypeptide domains or nucleic acid (*e.g.*, DNA) segments. In the context of a fusion protein or other polypeptide, the term “operably linked” means that two or more amino acid segments are linked so as to produce a functional polypeptide. For example, in the context of a IL2 proprotein of the disclosure, separate components (*e.g.*, an Fc domain and an IL2R α moiety) can be operably linked directly or through peptide linker sequences. In the

context of a nucleic acid encoding a fusion protein, such as a half antibody of an IL2 proprotein of the disclosure, “operably linked” means that the two nucleic acids are joined such that the amino acid sequences encoded by the two nucleic acids remain in-frame. In the context of transcriptional regulation, the term refers to the functional relationship of a transcriptional regulatory sequence to a transcribed sequence. For example, a promoter or enhancer sequence is operably linked to a coding sequence if it stimulates or modulates the transcription of the coding sequence in an appropriate host cell or other expression system.

[0049] Polypeptide, Peptide and Protein: The terms “polypeptide”, “peptide” and “protein” are used interchangeably herein to refer to a polymer of amino acid residues.

10 **[0050] Proprotein:** A “proprotein” is a protein precursor that is inactive and which can be activated by proteolysis by a protease. Thus, proproteins are “protease activatable”.

[0051] Protease: The term “protease” as used herein refers to any enzyme that catalyzes hydrolysis of a peptide bond. Generally, the proteases useful in the present disclosure, *e.g.*, the proteases described in Section 6.5.1, recognize and cleaves a specific sequence motif, *e.g.*, a substrate as described in Section 6.5.2. Preferably, the proteases are expressed at higher levels in cancer tissues as compared to normal tissues.

15 **[0052] Protease-cleavable linker:** As used herein, the term “protease-cleavable linker” or “PCL” refers to a peptide whose amino acid sequence contains one or more (*e.g.*, two, three or more) substrate sequences for one or more proteases. Exemplary protease-cleavable linkers are described in Section 6.5 and exemplary protease-cleavable linker sequences are disclosed in Section 6.5.4.

[0053] Recognize: The term “recognize” as used herein refers to an antibody or antibody fragment (*e.g.*, a targeting moiety) that finds and interacts (*e.g.*, binds) with its epitope.

25 **[0054] Single Chain Fab or scFab:** The term “single chain Fab” or “scFab” as used herein refers an ABD comprising a VH domain, a CH1 domain, a VL domain, a CL domain and a linker. In some embodiments, the foregoing domains and linker are arranged in one of the following orders in a N-terminal to C-terminal orientation: (a) VH-CH1-linker-VL-CL, (b) VL-CL-linker-VH-CH1, (c) VH-CL-linker-VL-CH1 or (d) VL-CH1-linker-VH-CL. Linkers are suitably noncleavable linkers of at least 30 amino acids, preferably between 32 and 50 amino acids. Single chain Fab fragments are typically stabilized via the natural disulfide bond between the CL domain and the CH1 domain. In addition, these single chain Fab molecules might be further stabilized by

30

generation of interchain disulfide bonds via insertion of cysteine residues (e.g., at position 44 in the VH domain and position 100 in the VL domain according to Kabat numbering).

[0055] Single Chain Fv or scFv: The term “single-chain Fv” or “scFv” as used herein refers to ABDs comprising the VH and VL domains of an antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the VH and VL domains which enables the scFv to form the desired structure for antigen-binding. For a review of scFv see Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds. (1994), Springer-Verlag, New York, pp. 269-315. The VH and VL and be arranged in the N- to C-terminal order VH-VL or VL-VH, typically separated by a linker, for example a linker as set forth in Table E.

[0056] Spacer: As used herein, the term “spacer” refers to a peptide, the amino acid sequence of which is not a substrate for a protease, incorporated into a linker containing a substrate. A spacer can be used to separate the substrate from other domains in a molecule, for example an ABD. In some aspects, residues in the spacer minimize aminopeptidase and/or exopeptidase action to prevent cleavage of N-terminal amino acids.

[0057] Specifically (or selectively) binds: The term “specifically (or selectively) binds” to an antigen or an epitope refers to a binding reaction that is determinative of the presence of a cognate antigen or an epitope in a heterogeneous population of proteins and other molecules. The binding reaction can be but need not be mediated by an antibody or antibody fragment. The term “specifically binds” does not exclude cross-species reactivity. For example, an antigen-binding domain (e.g., an antigen-binding fragment of an antibody) that “specifically binds” to an antigen from one species may also “specifically bind” to that antigen in one or more other species. Thus, such cross-species reactivity does not itself alter the classification of an antigen-binding domain as a “specific” binder. In certain embodiments, an antigen-binding domain of the disclosure that specifically binds to a human antigen has cross-species reactivity with one or more non-human mammalian species, e.g., a primate species (including but not limited to one or more of *Macaca fascicularis*, *Macaca mulatta*, and *Macaca nemestrina*) or a rodent species, e.g., *Mus musculus*.

[0058] Subject: The term “subject” includes human and non-human animals. Non-human animals include all vertebrates, e.g., mammals and non-mammals, such as non-human primates, sheep, dog, cow, chickens, amphibians, and reptiles. In preferred embodiments, the subject is human.

[0059] Substrate: The term “substrate” refers to peptide sequence on which a protease will act and within which the protease will cleave a peptide bond.

[0060] Target Molecule: The term “target molecule” as used herein refers to any biological molecule (*e.g.*, protein, carbohydrate, lipid or combination thereof) expressed on a cell surface or in the extracellular matrix that can be specifically bound by a targeting moiety in an IL2 proprotein of the disclosure.

[0061] Targeting Moiety: The term “targeting moiety” as used herein refers to any molecule or binding portion (*e.g.*, an immunoglobulin or an antigen binding fragment) thereof that can bind to a cell surface or extracellular matrix molecule at a site to which an IL2 proprotein of the disclosure is to be localized, for example on tumor cells or on lymphocytes in the tumor microenvironment. In some embodiments, the targeting moiety binds to a TAA. In other embodiments, the targeting moiety binds to a TCA. The targeting moiety can also have a functional activity in addition to localizing an IL2 proprotein to a particular site. For example, a targeting moiety that binds to a checkpoint inhibitor such as PD1 can also exhibit anti-tumor activity or enhance the anti-tumor activity by IL2, for example by inhibiting PD1 signaling.

[0062] T-Cell Antigen, TCA: The term “T-cell antigen” or “TCA” refers to a molecule (typically a protein, carbohydrate, lipid or some combination thereof) that is expressed on the surface of a T-lymphocyte and is useful for the preferential targeting of a pharmacological agent to a particular site. In some embodiments, the site is cancer tissue and/or the T-cell antigen is a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, or a checkpoint inhibitor expressed on a T-lymphocyte.

[0063] Tumor: The term “tumor” is used interchangeably with the term “cancer” herein, *e.g.*, both terms encompass solid and liquid, *e.g.*, diffuse or circulating, tumors. As used herein, the term “cancer” or “tumor” includes premalignant, as well as malignant cancers and tumors.

[0064] Tumor-Associated Antigen, TAA: The term “tumor-associated antigen” or “TAA” refers to a molecule (typically a protein, carbohydrate, lipid or some combination thereof) that is expressed on the surface of a cancer cell, either entirely or as a fragment (*e.g.*, MHC/peptide), and which is useful for the preferential targeting of a pharmacological agent to the cancer cell. In some embodiments, a TAA is a marker expressed by both normal cells and cancer cells, *e.g.*, a lineage marker. In some embodiments, a TAA is a cell surface molecule that is overexpressed in a cancer cell in comparison to a normal cell, for instance, 1-fold over expression, 2-fold overexpression, 3-fold overexpression or more in comparison to a normal cell. In some

embodiments, a TAA is a cell surface molecule that is inappropriately synthesized in the cancer cell, for instance, a molecule that contains deletions, additions or mutations in comparison to the molecule expressed on a normal cell. In some embodiments, a TAA will be expressed exclusively on the cell surface of a cancer cell, entirely or as a fragment (e.g., MHC/peptide),
5 and not synthesized or expressed on the surface of a normal cell. Accordingly, the term "TAA" encompasses antigens that are specific to cancer cells, sometimes known in the art as tumor-specific antigens ("TSAs").

[0065] Treat, Treatment, Treating: As used herein, the terms "treat", "treatment" and "treating" refer to the reduction or amelioration of the progression, severity and/or duration of a
10 proliferative disorder, or the amelioration of one or more symptoms (preferably, one or more discernible symptoms) of a proliferative disorder resulting from the administration of one or more IL2 proproteins of the disclosure. In specific embodiments, the terms "treat", "treatment" and "treating" refer to the amelioration of at least one measurable physical parameter of a proliferative disorder, such as growth of a tumor, not necessarily discernible by the patient. In
15 other embodiments the terms "treat", "treatment" and "treating" refer to the inhibition of the progression of a proliferative disorder, either physically by, e.g., stabilization of a discernible symptom, physiologically by, e.g., stabilization of a physical parameter, or both. In other embodiments the terms "treat", "treatment" and "treating" refer to the reduction or stabilization of tumor size or cancerous cell count.

20 **[0066] Universal Light Chain, ULC:** The term "universal light chain" or "ULC" as used herein refers to a light chain variable region (VL) that can pair with more than one heavy chain variable region (VH). In the context of a targeting moiety, the term "universal light chain" or "ULC" refers to a light chain polypeptide capable of pairing with the heavy chain region of the targeting moiety and also capable of pairing with other heavy chain regions. ULCs can also include
25 constant domains, e.g., a CL domain of an antibody. Universal light chains are also known as "common light chains".

[0067] VH: The term "VH" refers to the variable region of an immunoglobulin heavy chain of an antibody, including the heavy chain of an Fv, scFv, dsFv or Fab.

[0068] VL: The term "VL" refers to the variable region of an immunoglobulin light chain,
30 including the light chain of an Fv, scFv, dsFv or Fab.

6.2. IL2 Proproteins

- [0069] The present disclosure relates to IL2 propoteins comprising an IL2 moiety, an IL2R α moiety, and a protease-cleavable linker, arranged so that the IL2R α diminishes or blocks the activity of the IL2 moiety. The IL2 propotein is configured such that upon encountering a protease, *e.g.*, a protease that is overexpressed in the tumor environment, the protease-cleavable linker is cleaved and IL2 is released and stimulates cytotoxic T-cell activity against tumor cells. Typically, the IL2 propoteins of the disclosure are dimeric and comprise two Fc domains that associate for form an Fc region, C-terminal to which are linkers which may be protease-cleavable or non-cleavable, the IL2 moieties, additional linkers that are protease-cleavable, and IL2R α moieties arranged in N- to C-terminal order.
- [0070] In some embodiments, the IL2 propoteins of the disclosure generally comprise: (a) a first Fc domain and a second Fc domain capable of associating to form an Fc region; (b) two linkers which may be protease-cleavable or non-cleavable C-terminal to the Fc domains which, in reference to the embodiments depicted in FIG. 1A and FIG. 2A, correspond to Linker A and Linker C and in reference to the numbered embodiments below correspond to the first linker and the third linker) (c) two (a first and a second) IL2 moieties C-terminal to the first and third linkers; (c) two further linkers C-terminal to the IL2 moieties that are protease-cleavable and which, in reference to the embodiments depicted in FIG. 1A and FIG. 2A, correspond to Linker B and Linker D and in reference to the numbered embodiments below correspond to the second linker and the fourth linker; and (d) two (a first and a second) IL2R α moieties C-terminal to the second and fourth linkers. The IL2 moiety in the IL2 propotein is in an inactive form by virtue of masking by the IL2R α moiety, but is released following protease-cleavage of one or more of the protease-cleavable linkers at a locale that expresses a protease capable of cleaving one or more of the protease-cleavable linkers, *e.g.*, in the tumor environment.
- [0071] The IL2 propoteins may further comprise one or more targeting moieties, and in some embodiments comprise two targeting moieties N-terminal to the Fc domains. Examples of targeting moieties are described in 6.7 and suitable targeting moiety formats are described in Section 6.8. In reference to the embodiments depicted in FIG. 1A and FIG. 2A, the IL2 propoteins may comprise two Fab domains at their N-termini. In some embodiments, such as depicted in FIG. 1A and FIG. 2A, the Fc domains comprise hinge domains at their N-termini.

[0072] Examples of suitable IL2 moieties for incorporation into the IL2 propoteins are described in Section 6.3, examples of suitable IL2R α moieties are described in Section 6.4, and examples of suitable protease-cleavable linkers are described in Section 6.5.

[0073] Generally, the IL2 proproteins of the disclosure contain multiple linkers. Preferably, when present, linkers other than the specified protease-cleavable linkers are non-cleavable. Examples of non-cleavable linkers are set forth in Section 6.6.

[0074] Suitable Fc domains with or without hinge sequences are described in Section 6.9.

5 **[0075]** One exemplary IL2 proprotein is depicted in FIG. 1A. The IL2 proprotein comprises:

- 10 a) a first polypeptide chain comprising the heavy chain of a Fab associated with the light chain of a Fab on a separate polypeptide chain, together forming a first targeting moiety, followed by an Fc domain comprising a hinge domain, followed by a first protease-cleavable linker ("Linker A"), followed by an IL2 moiety, followed by a second protease-cleavable linker ("Linker B"), followed by an IL2R α moiety; and
- 15 b) a second polypeptide chain comprising the heavy chain of a Fab associated with the light chain of a Fab on a separate polypeptide chain, together forming a second targeting moiety, followed by an Fc domain comprising a hinge domain, followed by a first protease cleavable linker ("Linker C"), followed by an IL2 moiety, followed by a second protease-cleavable linker ("Linker D"), followed by an IL2R α moiety.

[0076] An additional exemplary IL2 proprotein is depicted in FIG. 2A. The IL2 proprotein comprises:

- 20 a) a first polypeptide chain comprising the heavy chain of a Fab associated with the light chain of a Fab on a separate polypeptide chain, together forming a first targeting moiety, followed by an Fc domain comprising a hinge domain, followed by a non-cleavable linker ("Linker A"), followed by an IL2 moiety, followed by a protease-cleavable linker ("Linker B"), followed by an IL2R α moiety; and
- 25 b) a second polypeptide chain comprising the heavy chain of a Fab associated with the light chain of a Fab on a separate polypeptide chain, together forming a second targeting moiety, followed by an Fc domain comprising a hinge domain, followed by a non-cleavable linker ("Linker C"), followed by an IL2 moiety, followed by a protease-cleavable linker ("Linker D"), followed by an IL2R α moiety.

[0077] Accordingly, the IL2 proproteins may include four protease-cleavable linkers, as illustrated in FIG. 1A, or two protease-cleavable linkers, as illustrated in FIG. 2A.

30 **[0078]** Cleavage of all protease-cleavable linkers in IL2 proproteins with four protease-cleavable linkers results in release of an activated IL2 protein comprising the IL2 moiety and

lacking an Fc moiety, an IL2R α moiety, and, if present, a targeting moiety. In some embodiments, this configuration is advantageously utilized for IL2 proproteins comprising a targeting moiety that binds to a TAA or ECM target molecule that is expressed in the tumor environment. As illustrated in FIG. 1B, and without intending to be bound by theory, the

5 inventors believe that in this configuration, the targeting moiety targets the IL2 proprotein to the tumor environment, where proteases cleave the protease-cleavable linkers resulting in the release of an IL2 protein comprising the IL2 moiety and linker sequences. This locally activated IL2 protein then induces an immune response against the cancer cells by stimulating the T-lymphocytes in the tumor environment.

10 **[0079]** Cleavage of both protease-cleavable linkers in IL2 proproteins with two protease-cleavable linkers results in release of an activated IL2 protein comprising the IL2 moiety, the Fc moiety, and, if present, the targeting moiety, but lacking the IL2R α moiety. In some embodiments, this configuration is advantageously utilized for IL2 proproteins comprising a targeting moiety that binds to a TCA, particularly a TCA that is expressed on an antigen

15 activated T cell (e.g. PD1, Lag3, 41BB, etc.). As illustrated in FIGs. 2B-2D, and without intending to be bound by theory, the inventors believe that in this configuration, cleavage of the protease-cleavable linkers in the tumor environment results in the release of an IL2 protein comprising the IL2 moiety and a T-cell targeting moiety. This locally activated, T-cell-targeted IL2 protein then induces an enhanced immune response against the cancer cells by stimulating

20 the T-lymphocytes in the tumor environment.

[0080] Importantly, without being bound by theory, the inventors believe inclusion of a protease-cleavable linker between the IL2 moieties and IL2R α moieties of IL2 proproteins of the disclosure to be important for optimal stimulation of cytotoxic T-cell activity against tumor cells and induction of an enhanced immune response by IL2. In contrast, a molecule having

25 components arranged as in the IL2 proproteins of the disclosure but having a non-cleavable linker separating the IL2 moieties and IL2R α moieties showed low tumor growth control (see U.S. 2022/0402989 at paragraph [0253] and FIGs. 6G and 6H).

6.3. The IL2 Moiety

[0081] The IL2 moiety of the IL2 proproteins of the disclosure comprises a wild type or variant

30 IL2 moiety.

[0082] In eukaryotic cells human IL2 is synthesized as a precursor polypeptide of 153 amino acids, from which 20 amino acids are removed to generate mature secreted IL2 (Taniguchi *et al.*, 1983, Nature 302(5906):305-10). Mature human IL2 has the following amino acid sequence:

5 APTSSSTKKTQLQLEHLLLDLQMILNGINNYKNPKLTRMLTFKFYMPKKATELKHLCLE
 EELKPLEEVNLNAQSKNFHLRPRDLISNINVIVLELKGSETTFMCEYADETATIVEFLNRW
 ITFCQSIISTLT (SEQ ID NO: 15)

[0083] In some embodiments, the IL2 moieties of the disclosure are not CD122 directed, *e.g.*, they do not have amino acid substitutions in the IL2 moiety that make them preferentially bind to IL2R β as compared to IL2R α .

10 **[0084]** In some embodiments, the IL2 moieties of the disclosure are CD25 directed, *e.g.*, they have one or more amino acid substitutions in the IL2 moiety that make them preferentially bind to IL2R α as compared to IL2R β .

[0085] In certain embodiments, the IL2 proproteins of the disclosure have one or more amino acid substitutions in the IL2 moiety that reduce binding to IL2R β . For example, in some
 15 embodiments, the IL2 moiety can have up to 50-fold (and in some embodiments up to 100-fold) to 1,000-fold attenuated binding to human IL2R β as compared to wild-type human IL2.

[0086] The IL2 moiety with reduced binding to IL2R β can retain its affinity to IL2R α , or have reduced binding to IL2R α . For example, in some embodiments, the IL2 moiety can have up to 50-fold attenuated binding to human IL2 as compared to wild-type human IL2.

20 **[0087]** Other characteristics of useful IL2 variants may include the ability to induce proliferation of IL2R α -bearing CD8⁺ T cells in tumors, the ability to induce IL2 signaling in IL2R α -bearing CD8⁺ T cells in tumors, and an improved therapeutic index.

[0088] In one embodiment, the IL2 moiety comprises one or more amino acid substitutions that reduce affinity to IL2R β and preserve affinity to IL2R α . An exemplary amino acid substitution is
 25 N88D. Other amino acid substitutions that reduce or abolish the affinity of IL2 to IL2R β are D20T, N88R, N88D or Q126D (see *e.g.*, US Patent Publication No. US 2007/0036752).

[0089] In one embodiment, the IL2 moiety comprises one or more amino acid substitutions that reduce affinity to IL2R α and preserve, or reduces affinity to a lesser degree, to IL2R β , resulting in CD122 directed IL2 moieties. Exemplary CD122 directed IL2 moieties are those comprising
 30 both H16A and F42A substitutions. Accordingly, in some embodiments, the IL2 moiety

comprises the amino acid sequence of human IL2 with H16A and F42A substitutions, as shown below:

5 SAPTSSSTKKTQLQLEALLDLQMILNGINNYKNPKLTRMLTAKFYMPKKATELKHLCCL
 EEELKPLEEVNLNAQSKNFHLRPRDLISINIVLELKGSETTFMCEYADETATIVEFLNR
 WITFCQSIISTLT (SEQ ID NO:124)

[0090] In certain embodiments, the IL2 moiety comprises an amino acid substitution which eliminates the O-glycosylation site of IL2 at a position corresponding to residue 3 of human IL2. Exemplary amino acid substitutions at T3 are T3A, T3G, T3Q, T3E, T3N, T3D, T3R, T3K, and T3P. In a specific embodiment, the substitution is T3A.

10 [0091] The IL2 moiety is preferably essentially a full-length IL2 molecule, *e.g.*, a human IL2 molecule. In certain embodiments the IL2 moiety is a human IL-2 molecule.

[0092] C125 can be substituted with S, V, or A to reduce protein aggregation, as described in U.S. Patent No. 4,518,584.

15 [0093] As described therein, one may also delete the N-terminal alanine residue of IL2, resulting in des-A1 IL2.

[0094] Further, the IL2 moiety may include a substitution of methionine 104 with a neutral amino acid such as alanine, as described in U.S. Patent No. 5,206,344.

20 [0095] Accordingly, the IL2 moieties of the disclosure can have amino acid deletions and / or substitutions selected from des-A1 M104A IL2, des-A1 M104A C125S IL2, M104A IL2, M104A C125A IL2, des-A1 M104A C125A IL2, or M104A C125S IL2, in addition to other variations alter the binding of IL2 to its receptor. These and other mutants may be found in U.S. Patent No. 5,116,943 and in Weiger *et al.*, 1989, Eur J Biochem 180:295-300.

25 [0096] In various aspects, any of the foregoing IL2 moieties comprises an amino acid sequence having at least about 90%, at least about 91%, at least about 92%, about at least 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% or 100% sequence identity to mature human IL2.

6.4. The IL2R α Moiety

30 [0097] The IL2 proproteins of the disclosure comprise an IL2R α moiety, comprising or consisting of an IL2-binding domain of IL2R α , *e.g.*, the extracellular domain of an IL2R α . The sequence of the mature human IL2R α extracellular domain (corresponding to amino acids 22-272 of human IL2R α) is:

5 Glu Leu Cys Asp Asp Asp Pro Pro Glu Ile Pro His Ala Thr Phe Lys
 Ala Met Ala Tyr Lys Glu Gly Thr Met Leu Asn Cys Glu Cys Lys Arg
 Gly Phe Arg Arg Ile Lys Ser Gly Ser Leu Tyr Met Leu Cys Thr Gly
 Asn Ser Ser His Ser Ser Trp Asp Asn Gln Cys Gln Cys Thr Ser Ser
 10 Ala Thr Arg Asn Thr Thr Lys Gln Val Thr Pro Gln Pro Glu Glu Gln
 Lys Glu Arg Lys Thr Thr Glu Met Gln Ser Pro Met Gln Pro Val Asp
 Gln Ala Ser Leu Pro Gly His Cys Arg Glu Pro Pro Pro Trp Glu Asn
 Glu Ala Thr Glu Arg Ile Tyr His Phe Val Val Gly Gln Met Val Tyr
 Tyr Gln Cys Val Gln Gly Tyr Arg Ala Leu His Arg Gly Pro Ala Glu
 15 Ser Val Cys Lys Met Thr His Gly Lys Thr Arg Trp Thr Gln Pro Gln
 Leu Ile Cys Thr Gly Glu Met Glu Thr Ser Gln Phe Pro Gly Glu Glu
 Lys Pro Gln Ala Ser Pro Glu Gly Arg Pro Glu Ser Glu Thr Ser Cys
 Leu Val Thr Thr Thr Asp Phe Gln Ile Gln Thr Glu Met Ala Ala Thr
 Met Glu Thr Ser Ile Phe Thr Thr Glu Tyr Gln Val Ala Val Ala Gly
 20 Cys Val Phe Leu Leu Ile Ser Val Leu Leu Leu Ser Gly Leu Thr Trp
 Gln Arg Arg Gln Arg Lys Ser Arg Arg Thr Ile (SEQ ID NO: 8)

[0098] The sequence of an IL2 binding portion of the human IL2R α extracellular domain
 (comprising the two "sushi" domains, which corresponds to amino acids 22-186 of human
 20 IL2R α) is:

25 Glu Leu Cys Asp Asp Asp Pro Pro Glu Ile Pro His Ala Thr Phe Lys
 Ala Met Ala Tyr Lys Glu Gly Thr Met Leu Asn Cys Glu Cys Lys Arg
 Gly Phe Arg Arg Ile Lys Ser Gly Ser Leu Tyr Met Leu Cys Thr Gly
 Asn Ser Ser His Ser Ser Trp Asp Asn Gln Cys Gln Cys Thr Ser Ser
 Ala Thr Arg Asn Thr Thr Lys Gln Val Thr Pro Gln Pro Glu Glu Gln
 Lys Glu Arg Lys Thr Thr Glu Met Gln Ser Pro Met Gln Pro Val Asp
 Gln Ala Ser Leu Pro Gly His Cys Arg Glu Pro Pro Pro Trp Glu Asn
 Glu Ala Thr Glu Arg Ile Tyr His Phe Val Val Gly Gln Met Val Tyr
 Tyr Gln Cys Val Gln Gly Tyr Arg Ala Leu His Arg Gly Pro Ala Glu
 30 Ser Val Cys Lys Met Thr His Gly Lys Thr Arg Trp Thr Gln Pro Gln
 Leu Ile Cys Thr Gly (SEQ ID NO: 9)

[0099] The sequence of an alternative IL2 binding portion of the human IL2R α extracellular
 domain, which corresponds to amino acids 22-240 of human IL2R α , is:

35 Glu Leu Cys Asp Asp Asp Pro Pro Glu Ile Pro His Ala Thr Phe Lys
 Ala Met Ala Tyr Lys Glu Gly Thr Met Leu Asn Cys Glu Cys Lys Arg
 Gly Phe Arg Arg Ile Lys Ser Gly Ser Leu Tyr Met Leu Cys Thr Gly
 Asn Ser Ser His Ser Ser Trp Asp Asn Gln Cys Gln Cys Thr Ser Ser
 Ala Thr Arg Asn Thr Thr Lys Gln Val Thr Pro Gln Pro Glu Glu Gln
 40 Lys Glu Arg Lys Thr Thr Glu Met Gln Ser Pro Met Gln Pro Val Asp
 Gln Ala Ser Leu Pro Gly His Cys Arg Glu Pro Pro Pro Trp Glu Asn
 Glu Ala Thr Glu Arg Ile Tyr His Phe Val Val Gly Gln Met Val Tyr
 Tyr Gln Cys Val Gln Gly Tyr Arg Ala Leu His Arg Gly Pro Ala Glu
 Ser Val Cys Lys Met Thr His Gly Lys Thr Arg Trp Thr Gln Pro Gln
 Leu Ile Cys Thr Gly Glu Met Glu Thr Ser Gln Phe Pro Gly Glu Glu
 45 Lys Pro Gln Ala Ser Pro Glu Gly Arg Pro Glu Ser Glu Thr Ser Cys
 Leu Val Thr Thr Thr Asp Phe Gln Ile Gln Thr Glu Met Ala Ala Thr

Met Glu Thr Ser Ile Phe Thr Thr Glu Tyr Gln (SEQ ID NO: 10)

5 [0100] The IL2R α moiety preferably comprises an amino acid sequence with at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% or 100% sequence identity to any of the sequences above, *i.e.*, any one of amino acids 22-186 of IL2R α , amino acids 22-240 of IL2R α , or amino acids 22-272 of IL2R α , or any IL2 binding portion thereof.

10 [0101] In certain aspects, the IL2R α moiety can comprise or consist of an amino acid sequence having at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% or 100% sequence identity to an IL2 binding portion of human IL2R α , optionally wherein the binding portion has an amino acid sequence of (a) at least 160 amino acids, at least 161 amino acids, at least 162 amino acids, at least 164 amino acids or at least 165 amino acids and/or (b) up to 251, up to 240, up to 230, up to 220, up to 210, up to 200, up to 190, up to 180 or up to 170 amino acids of the extracellular domain of human IL2R α .
15 In particular embodiments, the portion of human IL2R α is bounded by any one of (a) and (b) in the preceding sentence, *e.g.*, at least 160 and up to 180 amino acids from human IL2R α , at least 162 and up to 200 amino acids from human IL2R α , at least 160 and up to 220 amino acids from human IL2R α , at least 164 and up to 190 amino acids from human IL2R α , and so on and so forth.

20 [0102] In some embodiments, the IL2R α moiety comprises or consists of an amino acid sequence having at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% or 100% sequence identity to amino acids 22-186, with or without an additional up to 5 amino acids, up to 10 amino acids, up to 15 amino acids, up to 20 amino acids, up to 30 amino acids, or up to 40 amino acids C-terminal to amino acid
25 residue 186, of IL2R α .

[0103] In certain embodiments, the IL2R α moiety has at least one fewer O-glycosylation and/or N-glycosylation compared to the extracellular domain of native IL2R α , for example by a substitution at one or more of amino acid N49, amino acid N68, amino acid T74, amino acid T85, amino acid T197, amino acid T203, amino acid T208, and amino acid T216. In some
30 embodiments, the one or more substitutions are from asparagine to an amino acid selected from the group consisting of alanine, threonine, serine, arginine, aspartic acid, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, tryptophan, tyrosine, and valine. In some embodiments, the one or more substitutions are from

threonine to an amino acid selected from the group consisting of alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, tryptophan, tyrosine, and valine. In some embodiments, the one or more substitutions are at amino acid S50 (*e.g.*, S50P), amino acid S51
5 (*e.g.*, S51R, S51N, S51D, S51C, S51Q, S51E, S51G, S51H, S51I, S51L, S51K, S51M, S51F, S51P, S51W, S51Y, or S51V), amino acid T69 (*e.g.*, T69P), amino acid T70 (*e.g.*, T70R, T70N, T70D, T70C, T70Q, T70E, T70G, T70H, T70I, T70L, T70K, T70M, T70F, T70P, T70W, T70Y, or T70V), amino acid C192 (*e.g.*, C192R, C192N, C192D, C192Q, C192E, C192G, C192H, C192I, C192L, C192K, C192M, C192F, C192P, C192W, C192Y, or C192V), or any combination
10 thereof.

6.5. Protease-Cleavable Linkers

[0104] The IL2 proproteins of the disclosure typically comprise four linkers, referred to in the numbered embodiments below as the first, second, third and fourth linkers, with the first and second linkers on one polypeptide chain and the third and fourth linkers on another polypeptide
15 chain. In the embodiments depicted in FIG. 1A and FIG. 2A, the first and second linkers are referred to as Linker A and Linker B and the third and fourth linkers are referred to as Linker C and Linker D.

[0105] In other embodiments, all four linkers (the first, second, third and fourth linkers, corresponding to Linker A, Linker B, Linker C and Linker D) are protease cleavable. An
20 exemplary IL2 proprotein configured according to such embodiments is illustrated in FIG. 1A.

[0106] In some embodiments, the second and fourth linkers (corresponding to Linker B and Linker D) are protease cleavable and the first and third linkers (corresponding to Linker A and Linker C) are non-cleavable. An exemplary IL2 proprotein configured according to such
embodiments is illustrated in FIG. 2A.

[0107] A protease-cleavable linker can range from 20 amino acids to 80 or more amino acids, and in certain aspects a non-cleavable peptide linker ranges from 20 amino acids to 60 amino acids, 20 amino acids to 40 amino acids, from 30 amino acids to 50 amino acids, from 20 amino acids to 80 amino acids, or from 30 amino acids to 70 amino acids in length.

[0108] The protease-cleavable linkers comprise one or more substrate sequences for one or
30 more proteases, for example one or more of the proteases set forth in Section 6.5.1. The one or more substrate sequences, *e.g.*, one or more of the substrate sequences set forth in Section 6.5.2, are typically flanked by one or more spacer sequences, *e.g.*, spacer sequences as

described in Section 6.5.3. Each protease-cleavable linker can include one, two, three or more substrate sequences. The spacer sequences can be adjoining, overlapping, or separated by spacer sequences. Preferably, the C- and N-termini of the protease-cleavable linkers contain spacer sequences.

- 5 **[0109]** In various aspects of IL2 proproteins comprising four protease-cleavable linkers, the first and third protease-cleavable linkers (corresponding to Linkers A and C in the embodiment of FIG. 1A) are cleavable by the same protease and/or the second and fourth protease-cleavable linkers (corresponding to Linkers B and D in the embodiment of FIG. 1A) are cleavable by the same protease. In some embodiments, the protease is a protease set forth in Table A.
- 10 **[0110]** In further aspects of IL2 proproteins comprising four protease-cleavable linkers, the first and third protease-cleavable linkers (corresponding to Linkers A and C in the embodiment of FIG. 1A) comprise the same substrate sequence(s) and/or the second and fourth protease-cleavable linkers (corresponding to Linkers B and D in the embodiment of FIG. 1A) comprise the same substrate sequence(s). In some embodiments, the substrate sequence(s) are set forth in
- 15 Table B. In further embodiments, the first and third protease-cleavable linkers (corresponding to Linkers A and C in the embodiment of FIG. 1A) also comprise the same spacer sequence(s) and/or the second and fourth protease-cleavable linkers (corresponding to Linkers B and D in the embodiment of FIG. 1A) also comprise the same spacer sequence(s). In some embodiments, the spacer sequence(s) are set forth in Table C.
- 20 **[0111]** In further aspects IL2 proproteins comprising four protease-cleavable linkers, the first and third protease-cleavable linkers (corresponding to Linkers A and C in the embodiment of FIG. 1A) comprise the same linker sequence(s) and/or the second and fourth protease-cleavable linkers (corresponding to Linkers B and D in the embodiment of FIG. 1A) comprise the same linker sequence(s). In some embodiments, the linker sequence(s) are set forth in Table
- 25 D.
- [0112]** In some embodiments of IL2 proproteins comprising four protease-cleavable linkers, the first and third protease-cleavable linkers (corresponding to Linkers A and C in the embodiment of FIG. 1A) are the same as the second and fourth protease-cleavable linkers (corresponding to Linkers B and D in the embodiment of FIG. 1A).
- 30 **[0113]** In other embodiments of IL2 proproteins comprising four protease-cleavable linkers, the first and third protease-cleavable linkers (corresponding to Linkers A and C in the embodiment

of FIG. 1A) are the different from the second and fourth protease-cleavable linkers (corresponding to Linkers B and D in the embodiment of FIG. 1A).

[0114] In some embodiments of IL2 proproteins comprising two protease-cleavable linkers (corresponding to Linkers B and D in the embodiment of FIG. 2A), both protease-cleavable linkers are the same. In other embodiments, the two protease-cleavable linkers are different.

[0115] In the foregoing aspects and embodiments of both IL2 proproteins comprising four protease-cleavable linkers and IL2 proproteins comprising two protease-cleavable linkers, the different linkers may be cleavable by the same protease, different proteases, or when a linker comprises multiple substrate sequences, the different linkers may be cleavable by multiple proteases, one or more of which are common and one or more of which are different.

[0116] Exemplary protease-cleavable linker sequences are set forth in Section 6.5.4.

6.5.1. Proteases

[0117] Exemplary protease whose substrate sequences can be incorporated into the protease-cleavable linkers are set forth in Table A below.

TABLE A Exemplary Proteases for Substrate Cleavage		
ADAMS, ADAMTS, <i>e.g.</i> ADAM8 ADAM9 ADAM10 ADAM12 ADAM15 ADAM17/TACE ADAMDEC1 ADAMTS1 ADAMTS4 ADAMTS5	Caspases, <i>e.g.</i> , Caspase 1 Caspase 2 Caspase 3 Caspase 4 Caspase 5 Caspase 6 Caspase 7 Caspase 8 Caspase 9 Caspase 10 Caspase 14	MMP24 MMP26 MMP27 Cysteine proteinases, <i>e.g.</i> , Cruzipain Legumain Otubain-2
Aspartate proteases, <i>e.g.</i> , BACE Renin	Cysteine cathepsins, <i>e.g.</i> , Cathepsin B Cathepsin C Cathepsin K Cathepsin L Cathepsin S Cathepsin V/L2 Cathepsin X/Z/P	KLKs, <i>e.g.</i> , KLK4 KLK5 KLK6 KLK7 KLK8 KLK10 KLK11 KLK13 KLK14
Aspartic cathepsins, <i>e.g.</i> , Cathepsin D Cathepsin E		
NS3/4A PACE4 Plasmin	MMPs, <i>e.g.</i> ,	

TABLE A Exemplary Proteases for Substrate Cleavage		
PSA	MMP1	Metallo proteinases, <i>e.g.</i> , Meprin Neprilysin PSMA BMP-1
tPA	MMP2	
Thrombin	MMP3	
Tryptase		
uPA	MMP7	
	MMP8	
Type II Transmembrane Serine Proteases (TTSPs), <i>e.g.</i> ,	MMP9	Serine proteases, <i>e.g.</i> , activated protein C Cathepsin A Cathepsin G Chymase coagulation factor proteases (<i>e.g.</i> , FVIIa, FIXa, FXa, FXIa, FXIIa) Human Neutrophil Elastase Lactoferrin
DESC1	MMP11	
DPP-4	MMP12	
FAP	MMP13	
Hepsin	MMP14	
Matriptase-2	MMP15	
MT/SP1/Matriptase	MMP16	
	MMP17	
TMPRSS2	MMP19	
TMPRSS3	MMP20	
TMPRSS4	MMP23	

- [0118] In particular embodiments, the protease is matrix metalloprotease (MMP)-2, MMP-9, legumain asparaginyl endopeptidase, thrombin, fibroblast activation protease (FAP), MMP-1, MMP-3, MMP-7, MMP-8, MMP-12, MMP-13, MMP-14, membrane type 1 matrix metalloprotease (MT1-MMP), plasmin, transmembrane protease, serine (TMPRSS-3/4), cathepsin A, cathepsin B, cathepsin D, cathepsin E, cathepsin F, cathepsin H, cathepsin K, cathepsin L, cathepsin L2, cathepsin O, cathepsin S, caspase 1, caspase 2, caspase 3, caspase 4, caspase 5, caspase 6, caspase 7, caspase 8, caspase 9, caspase 10, caspase 11, caspase 12, caspase 13, caspase 14, human neutrophil elastase, urokinase/urokinase-type plasminogen activator (uPA), a disintegrin and metalloprotease (ADAM)10, ADAM12, ADAM17, ADAM with thrombospondin motifs (ADAMTS), ADAMTS5, beta secretase (BACE), granzyme A, granzyme B, guanidinobenzoatase, hepsin, matriptase, matriptase 2, meprin, neprilysin, prostate-specific membrane antigen (PSMA), tumor necrosis factor-converting enzyme (TACE), kallikrein-related peptidase (KLK)3, KLK5, KLK7, KLK11, NS3/4 protease of hepatitis C virus (HCV-NS3/4), tissue plasminogen activator (tPA), calpain, calpain 2, glutamate carboxypeptidase II, plasma kallikrein, AMSH-like protease, AMSH, γ -secretase component, antiplasmin cleaving enzyme (APCE), decysin 1, apoptosis-related cysteine peptidase, or N-acetylated alpha-linked acidic dipeptidase-like 1.

6.5.2. Substrates

[0119] Exemplary substrate sequences that are cleavable by a tumor protease and can be incorporated into the protease-cleavable linkers are set forth in Table B below.

TABLE B Substrate Sequences for Protease-Cleavable Linkers		
Substrate Sequence	Designation	Cleaving Protease
(DE) ₈ RPLALWRS(DR) ₈ (SEQ ID NO: 16)	SU1	MMP7
AARGPAIH (SEQ ID NO: 17)	SU2	
AAYHLVSQ (SEQ ID NO: 18)	SU3	Collagenase
AGLGISST (SEQ ID NO: 19)	SU4	Collagenase
AGLGVVER (SEQ ID NO: 20)	SU5	Collagenase
ALAL (SEQ ID NO: 21)	SU6	Lysosomal Enzyme
ALFFSSPP (SEQ ID NO: 22)	SU7	
ALFKSSFP (SEQ ID NO: 23)	SU8	
ALLLALL (SEQ ID NO: 24)	SU9	TOP
AQFVLTEG (SEQ ID NO: 25)	SU10	Collagenase
AQNLLGMV (SEQ ID NO: 26)	SU11	
AVGLLAPP (SEQ ID NO: 27)	SU12	Serine protease
DAFK (SEQ ID NO: 28)	SU13	Urokinase plasminogen activator (uPA)
DEVD (SEQ ID NO: 29)	SU14	Caspase-3
DEVDP (SEQ ID NO: 30)	SU15	Caspase-3
DPRSFL (SEQ ID NO: 31)	SU16	Thrombin
DVAQFVLT (SEQ ID NO: 32)	SU17	Collagenase
DVLK (SEQ ID NO: 33)	SU18	Plasmin
DWLYWPGI (SEQ ID NO: 34)	SU19	
EDDDDKA (SEQ ID NO: 35)	SU20	Enterokinase
EP(Cit)G(Hof)YL (SEQ ID NO: 36)	SU21	MMP2, MMP9, MMP14
EPQALAMS (SEQ ID NO: 37)	SU22	Collagenase
ESLPVVAV (SEQ ID NO: 38)	SU23	Collagenase
ESPAYYTA (SEQ ID NO: 39)	SU24	MMP
F(Pip)RS	SU25	Thrombin
FK	SU26	Lysosomal Enzyme
FPRPLGITGL (SEQ ID NO: 40)	SU27	
FRLLDWQW (SEQ ID NO: 41)	SU28	
GFLG (SEQ ID NO: 42)	SU29	Lysosomal Enzyme
GGAANLVRGG (SEQ ID NO: 43)	SU30	MMP11
GGGRR (SEQ ID NO: 44)	SU31	Urokinase plasminogen activator (uPA)
GGPRGLPG (SEQ ID NO: 45)	SU32	Cathepsin K
GGQPSGMWGW (SEQ ID NO: 46)	SU33	
GGSIDGR (SEQ ID NO: 47)	SU34	Factor Xa
GGWHTGRN (SEQ ID NO: 48)	SU35	
GIAGQ (SEQ ID NO: 49)	SU36	Collagenase

TABLE B Substrate Sequences for Protease-Cleavable Linkers		
Substrate Sequence	Designation	Cleaving Protease
GKAFRR (SEQ ID NO: 50)	SU37	Kallikrein 2
GPAGLYAQ (SEQ ID NO: 51)	SU38	
GPAGMKGL (SEQ ID NO: 52)	SU39	
GPEGLRVG (SEQ ID NO: 53)	SU40	Collagenase
GPLGIAGI (SEQ ID NO: 54)	SU41	Collagenase
GPLGVRG (SEQ ID NO: 55)	SU42	
GPQGIAGQ (SEQ ID NO: 56)	SU43	Collagenase
GPQGLLGA (SEQ ID NO: 57)	SU44	Collagenase
GPRSFSG (SEQ ID NO: 58)	SU45	
GPRSFGL (SEQ ID NO: 59)	SU46	
GPSHLVLT (SEQ ID NO: 60)	SU47	
GVSQNYPIVG (SEQ ID NO: 61)	SU48	HIV Protease
GVVQASCRLA (SEQ ID NO: 62)	SU49	CMV Protease
GWEHDG (SEQ ID NO: 63)	SU50	Interleukin 1 β converting enzyme
HSSKLQ (SEQ ID NO: 64)	SU51	Prostate Specific Antigen
HSSKLQEDA (SEQ ID NO: 65)	SU52	Prostate Specific Antigen
HSSKLQL (SEQ ID NO: 66)	SU53	Prostate Specific Antigen
HTGRSGAL (SEQ ID NO: 67)	SU54	
IDGR (SEQ ID NO: 68)	SU55	Factor Xa
IEGR (SEQ ID NO: 69)	SU56	Factor Xa
ILPRSPAF (SEQ ID NO: 70)	SU57	
IPVSLRSG (SEQ ID NO: 71)	SU58	MMP
ISSGL (SEQ ID NO: 72)	SU59	MMP
ISSGLL (SEQ ID NO: 73)	SU60	MMP
ISSGLLS (SEQ ID NO: 74)	SU61	MMP
ISSGLLSS (SEQ ID NO: 75)	SU62	MMP
ISSGLSS (SEQ ID NO: 76)	SU63	MMP
KGSGDVEG (SEQ ID NO: 77)	SU64	Caspase-3
KQEQNPGST (SEQ ID NO: 78)	SU65	FAP
KRALGLPG (SEQ ID NO: 79)	SU66	MMP7
LAAPLGLL (SEQ ID NO: 80)	SU67	
LAPLGLQRR (SEQ ID NO: 81)	SU68	
LAQKLKSS (SEQ ID NO: 82)	SU69	
LAQRLRSS (SEQ ID NO: 83)	SU70	
LEATA (SEQ ID NO: 84)	SU71	MMP9
LKAAPRWA (SEQ ID NO: 85)	SU72	
LLAPSHRA (SEQ ID NO: 86)	SU73	
LPGGLSPW (SEQ ID NO: 87)	SU74	
LSGRSANI (SEQ ID NO: 88)	SU75	Serine protease
LSGRSANP (SEQ ID NO: 89)	SU76	Serine protease
LSGRSDDH (SEQ ID NO: 90)	SU77	Serine protease

TABLE B Substrate Sequences for Protease-Cleavable Linkers		
Substrate Sequence	Designation	Cleaving Protease
LSGRSDIH (SEQ ID NO: 91)	SU78	Serine protease
LSGRSDNH (SEQ ID NO: 92)	SU79	Serine protease
LSGRSDNI (SEQ ID NO: 93)	SU80	Serine protease
LSGRSDNP (SEQ ID NO: 94)	SU81	Serine protease
LSGRSDQG (SEQ ID NO: 95)	SU82	Serine protease
LSGRSDQH (SEQ ID NO: 96)	SU83	Serine protease
LSGRSDTH (SEQ ID NO: 97)	SU84	Serine protease
LSGRSDYH (SEQ ID NO: 98)	SU85	Serine protease
LSGRSGNH (SEQ ID NO: 99)	SU86	Serine protease
LVLASSSFGY (SEQ ID NO: 100)	SU87	Herpes Simplex Virus Protease
MDAFLESS (SEQ ID NO: 101)	SU88	Collagenase
MGLFSEAG (SEQ ID NO: 102)	SU89	
MIAPVAYR (SEQ ID NO: 103)	SU90	
MVLGRSLL (SEQ ID NO: 104)	SU91	
NLL	SU92	Cathepsin B
NTLSGRSENHSG (SEQ ID NO: 105)	SU93	
NTLSGRSGNHGS (SEQ ID NO: 106)	SU94	
PAGLWLDP (SEQ ID NO: 107)	SU95	
PGGPAGIG (SEQ ID NO: 108)	SU96	
PIC(Et)FF (SEQ ID NO: 109)	SU97	Cathepsin D
PLGC(me)AG (SEQ ID NO: 110)	SU98	MMP
PLGL (SEQ ID NO: 111)	SU99	
PLGLAG (SEQ ID NO: 112)	SU100	MMP
PLGLAX (SEQ ID NO: 113)	SU101	MMP
PLGLWA (SEQ ID NO: 114)	SU102	MMP
PLGLWSQ (SEQ ID NO: 115)	SU103	MMP
PLTGRSGG (SEQ ID NO: 116)	SU104	
PMACK (SEQ ID NO: 117)	SU105	
PPRSFL (SEQ ID NO: 118)	SU106	Thrombin
PR(S/T)(L/I)(S/T)	SU107	MMP9
PRFRIIGG (SEQ ID NO: 119)	SU108	Plasmin
PVGYTSSL (SEQ ID NO: 120)	SU109	
PVQPIGPQ (SEQ ID NO: 121)	SU110	Collagenase
QALAMSAI (SEQ ID NO: 122)	SU111	Collagenase
QGRAITFI (SEQ ID NO: 123)	SU112	
QNQALRMA (SEQ ID NO: 125)	SU113	
RGPA (SEQ ID NO: 126)	SU114	
RGPAFNPM (SEQ ID NO: 127)	SU115	
RGPATPIM (SEQ ID NO: 128)	SU116	
RKSSIIIRMRDVL (SEQ ID NO: 129)	SU117	Plasmin
RLQLKAC (SEQ ID NO: 130)	SU118	MMP

TABLE B Substrate Sequences for Protease-Cleavable Linkers		
Substrate Sequence	Designation	Cleaving Protease
RLQLKL (SEQ ID NO: 131)	SU119	MMP
RMHLRSLG (SEQ ID NO: 132)	SU120	
RPSPMWAY (SEQ ID NO: 133)	SU121	
RQARVVNG (SEQ ID NO: 134)	SU122	Matripase
SAGFSLPA (SEQ ID NO: 135)	SU123	
SAPAVESE (SEQ ID NO: 136)	SU124	Collagenase
SARGPSRW (SEQ ID NO: 137)	SU125	
SGEPAYYA (SEQ ID NO: 138)	SU126	
SGGPLGVR (SEQ ID NO: 139)	SU127	
SGRIGFLRTA (SEQ ID NO: 140)	SU128	MMP14
SGRSA (SEQ ID NO: 141)	SU129	Urokinase plasminogen activator (uPA)
SGRSANPRG (SEQ ID NO: 142)	SU130	
SMLRSMPL (SEQ ID NO: 143)	SU131	
SPLPLRVP (SEQ ID NO: 144)	SU132	
SPLTGRSG (SEQ ID NO: 145)	SU133	
SPRSIMLA (SEQ ID NO: 146)	SU134	
SSRGPAYL (SEQ ID NO: 147)	SU135	
SSRHRRALD (SEQ ID NO: 148)	SU136	Plasmin
SSSFDKGKYKKGDDA (SEQ ID NO: 149)	SU137	Plasmin
SSSFDKGKYKRGDDA (SEQ ID NO: 150)	SU138	Plasmin
STFPFGMF (SEQ ID NO: 151)	SU139	
TARGPSFK (SEQ ID NO: 152)	SU140	
TGRGPSWW (SEQ ID NO: 153)	SU141	
TSGRSANP (SEQ ID NO: 154)	SU142	
TSTSGRSANPRG (SEQ ID NO: 155)	SU143	
VAGRSMRP (SEQ ID NO: 156)	SU144	
VAQFVLTE (SEQ ID NO: 157)	SU145	Collagenase
VHMPGLFLGP (SEQ ID NO: 158)	SU146	
VPLSLYSG (SEQ ID NO: 159)	SU147	MMP9
VVPEGRRS (SEQ ID NO: 160)	SU148	
WATPRPMR (SEQ ID NO: 161)	SU149	
YGAGLGVV (SEQ ID NO: 162)	SU150	Collagenase
HPVGLLAR (SEQ ID NO: 163)	SU151	

6.5.3. Spacers

- [0120] Exemplary spacer sequences that can be incorporated into the protease-cleavable linkers are set forth in Table C below. In addition to the spacer sequences set forth in Table C,
- 5 any of the non-cleavable linker sequences described in Section 6.6, e.g., the non-cleavable linker sequences set forth in Table E, or portions thereof can be used as spacer sequences.

TABLE C Spacer Sequences for Protease-Cleavable Linkers	
Spacer Sequence	Designation
(GGGGS) _n (SEQ ID NO: 164)	SP1
(GGGS) _n (SEQ ID NO: 165)	SP2
(GGS) _n (SEQ ID NO: 166)	SP3
(GS) _n (SEQ ID NO: 167)	SP4
(GSGGS) _n (SEQ ID NO: 168)	SP5
GGGGSGGGGS (SEQ ID NO: 169)	SP6
GGGGSGGGSGGGGS (SEQ ID NO: 170)	SP7
GGGGSGGGSGGGSGGGGS (SEQ ID NO: 171)	SP8
GGGKSGGKSGGGKS (SEQ ID NO: 172)	SP9
GGGKSGGKSGGKGS (SEQ ID NO: 173)	SP10
GGGS (SEQ ID NO: 174)	SP11
GGGSG (SEQ ID NO: 175)	SP12
GGKSGGKSGGKGS (SEQ ID NO: 176)	SP13
GGSGGGSGGGGS (SEQ ID NO: 177)	SP14
GGSGGS (SEQ ID NO: 178)	SP15
GGSGGSGGSGS (SEQ ID NO: 179)	SP16
GSGGG (SEQ ID NO: 180)	SP17
GSGSG (SEQ ID NO: 181)	SP18
GSS	SP19
GSSG (SEQ ID NO: 182)	SP20
GSSGGSGSG (SEQ ID NO: 183)	SP21
GSSGGSGSGG (SEQ ID NO: 184)	SP22
GSSGGSGSGGS (SEQ ID NO: 185)	SP23
GSSGGSGSGGSG (SEQ ID NO: 186)	SP24
GSSGGSGSGSGGGS (SEQ ID NO: 187)	SP25
GSSGGSGSGSGS (SEQ ID NO: 188)	SP26
GSSGT (SEQ ID NO: 189)	SP27
GSSSG (SEQ ID NO: 190)	SP28
QGQSGQ (SEQ ID NO: 191)	SP29
QGQSGQG (SEQ ID NO: 192)	SP30
QGQSGS (SEQ ID NO: 193)	SP31
QSGQ (SEQ ID NO: 194)	SP32
QSGQG (SEQ ID NO: 195)	SP33
QSGS (SEQ ID NO: 196)	SP34
SGQ	SP35
SGQG (SEQ ID NO: 197)	SP36
SGS	SP37
(G) _n (SEQ ID NO: 198)	SP38

[0121] In some embodiments, as used in Table C above, n is an integer from 1 to 10, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10.

6.5.4. Exemplary Protease-Cleavable Linkers

[0122] Exemplary protease-cleavable linkers comprising one or more substrate sequences as well as spacer sequences are set forth in Table D below.

TABLE D Protease-Cleavable Linker Sequences		
Linker Sequence	Designation	Cleaving Protease(s)
GGGISSGLLSGRSDNHGGGISSGLLSGRSD NHGGS (SEQ ID NO: 199)	PCL1	
GGGISSGLLSGRSDNHGGGISSGLLSGRSD NHGGS GGGISSGLLSGRSDNHGGGISSGLLSGRSD NHGGS (SEQ ID NO: 200)	PCL2	
GGSGGSIPVSLRSGGGISSGLLSGRSDNH GGSGGS (SEQ ID NO: 201)	PCL3	
GGSGGSVPLSLYSGGGISSGLLSGRSDNH GGSGGS (SEQ ID NO: 202)	PCL4	
GGSHPVGLLARGGGHPVGLLARGGGHPV GLLARGS (SEQ ID NO: 203)	PCL5	
GGSHPVGLLARGGGHPVGLLARGGSGRS AGGSGRSA (SEQ ID NO: 204)	PCL6	
AVGLLAPPGGLSGRSANI (SEQ ID NO: 205)	PCL7	ADAM17_2, FAPa_1, CTSL1_1
AVGLLAPPGGLSGRSANP (SEQ ID NO: 206)	PCL8	FAPa_1, ADAM17_2, CTSL1_1
AVGLLAPPGGLSGRSDDH (SEQ ID NO: 207)	PCL9	MMP14_1, MMP14_1, MMP14_1
AVGLLAPPGGLSGRSDIH (SEQ ID NO: 208)	PCL10	MMP14_1, MMP14_1, MMP14_1
AVGLLAPPGGLSGRSDNH (SEQ ID NO: 209)	PCL11	MMP14_1, MMP14_1
AVGLLAPPGGLSGRSDNI (SEQ ID NO: 210)	PCL12	MMP14_1, CTSL1_1, ADAM17_2
AVGLLAPPGGLSGRSDNP (SEQ ID NO: 211)	PCL13	CTSL1_1, ADAM17_2, FAPa_1
AVGLLAPPGGLSGRSDQH (SEQ ID NO: 212)	PCL14	
AVGLLAPPGGLSGRSDTH (SEQ ID NO: 213)	PCL15	FAPa_1, CTSL1_1, ADAM17_2
AVGLLAPPGGLSGRSDYH (SEQ ID NO: 214)	PCL16	
AVGLLAPPGGTSTSGRSANPRG (SEQ ID NO: 215)	PCL17	
AVGLLAPPSGRSANPRG (SEQ ID NO: 216)	PCL18	
AVGLLAPPTSGRSANPRG (SEQ ID NO: 217)	PCL19	

TABLE D Protease-Cleavable Linker Sequences		
Linker Sequence	Designation	Cleaving Protease(s)
GGALFKSSFPGPAGLYAQPLAQKLKSSGG K (SEQ ID NO: 218)	PCL20	CTSL1_1, MMP14_1, ADAM17_2
GGGSGGGSGGGSGGGSFVGGTGGGGSGG GGSGGS (SEQ ID NO: 219)	PCL21	
GGGSGGGSGGGSGGGSISSGLLSGRSDNH GGSGGS (SEQ ID NO: 220)	PCL22	
GGGSGGGSGGGSGGGSVPLSLYSGGGSG GGSGGS (SEQ ID NO: 221)	PCL23	
GGGSGGGSGGGSGPLGLWSQGGGGSGGGG SGGGGSGG (SEQ ID NO: 222)	PCL24	
GGGSGGGSGGSKKAAPGGGGSGGGGSGG GGSGGS (SEQ ID NO: 223)	PCL25	
GGGSGGGSGGSKKAAPVNGGGGSGGGG SGGGGS (SEQ ID NO: 224)	PCL26	
GGGSGGGSGGSPMAKKGGGGSGGGGSGG GGSGGS (SEQ ID NO: 225)	PCL27	
GGGSGGGSGGSPMAKKVNGGGGSGGGG SGGGGS (SEQ ID NO: 226)	PCL28	
GGGSGGGSGGSQARAKGGGGSGGGGSGG GGSGGS (SEQ ID NO: 227)	PCL29	
GGGSGGGSGGSQARAKVNGGGGSGGGG SGGGGS (SEQ ID NO: 228)	PCL30	
GGGSGGGSGGSRQARVNGGGGSGGGG SGGGGS (SEQ ID NO: 229)	PCL31	
GGGSGGGSGGSRQARVNGGGGSGVPLSL YSGGGGSGGGGS (SEQ ID NO: 230)	PCL32	
GGGSGGGSGGSRQARVNSVPLSLYSGGG GGSGGGGS (SEQ ID NO: 231)	PCL33	
GGGSGGGSGGSHMPLGFLGPGGGSGG GGSGGS (SEQ ID NO: 232)	PCL34	
GGGSGVHMPLGFLGPGRSRGSPGGGGGS (SEQ ID NO: 233)	PCL35	
GGGSGVHMPLGFLGPPMAKKGGGGSGGG GGSGGS (SEQ ID NO: 234)	PCL36	
GGGSGVHMPLGFLGPRQARVNGGGGSG GGGS (SEQ ID NO: 235)	PCL37	
GGGSGVHMPLGFLGPRQARVNGGGGSG GGGSGG (SEQ ID NO: 236)	PCL38	
GGPLAQKLKSSALFKSSFPGPAGLYAQGG R (SEQ ID NO: 237)	PCL39	ADAM17_2, CTSL1_1, MMP14_1
GLSGRSDNHGGAVGLLAPP (SEQ ID NO: 238)	PCL40	

TABLE D Protease-Cleavable Linker Sequences		
Linker Sequence	Designation	Cleaving Protease(s)
GLSGRSDNHGGVHMPGLGFLGP (SEQ ID NO: 239)	PCL41	
ISSGLLSGRSANI (SEQ ID NO: 240)	PCL42	MMP, Serine protease
ISSGLLSGRSANP (SEQ ID NO: 241)	PCL43	MMP, Serine protease
ISSGLLSGRSANPRG (SEQ ID NO: 242)	PCL44	MMP, Serine protease
ISSGLLSGRSDDH (SEQ ID NO: 243)	PCL45	MMP, Serine protease
ISSGLLSGRSDIH (SEQ ID NO: 244)	PCL46	MMP, Serine protease
ISSGLLSGRSDNH (SEQ ID NO: 245)	PCL47	MMP, Serine protease
ISSGLLSGRSDNI (SEQ ID NO: 246)	PCL48	CTSL1_1, MMP14_1
ISSGLLSGRSDNP (SEQ ID NO: 247)	PCL49	MMP, Serine protease
ISSGLLSGRSDQH (SEQ ID NO: 248)	PCL50	MMP, Serine protease
ISSGLLSGRSDTH (SEQ ID NO: 249)	PCL51	MMP, Serine protease
ISSGLLSGRSDYH (SEQ ID NO: 250)	PCL52	MMP, Serine protease
ISSGLLSGRSGNH (SEQ ID NO: 251)	PCL53	MMP, Serine protease
ISSGLLSSGGSGGSLSGRSDNH (SEQ ID NO: 252)	PCL54	
ISSGLLSSGGSGGSLSGRSGNH (SEQ ID NO: 253)	PCL55	
KGGPGGPAGIGPLAQLRSSLFSSFPGR (SEQ ID NO: 254)	PCL56	FAPa_1, ADAM17_1, CTSL1_1
KSGPGGPAGIGALFFSSPPLAQKLKSSGGR (SEQ ID NO: 255)	PCL57	FAPa_1, CTSL1_2, ADAM17_2
LSGRSDNHGGAVGLLAPP (SEQ ID NO: 256)	PCL58	
LSGRSDNHGGSGGSISSGLLSS (SEQ ID NO: 257)	PCL59	
LSGRSDNHGGSGGSGNQALRMA (SEQ ID NO: 258)	PCL60	
LSGRSDNHGGVHMPGLGFLGP (SEQ ID NO: 259)	PCL61	
LSGRSGNHGGSGGSISSGLLSS (SEQ ID NO: 260)	PCL62	
LSGRSGNHGGSGGSGNQALRMA (SEQ ID NO: 261)	PCL63	
QNQALRMAGGSGGSLSGRSDNH (SEQ ID NO: 262)	PCL64	
QNQALRMAGGSGGSLSGRSGNH (SEQ ID NO: 263)	PCL65	
RGGALFKSSFPLAQKLKSSGPAGLYAQGG K (SEQ ID NO: 264)	PCL66	CTSL1_1, ADAM17_2, MMP14_1
RGGGPAGLYAQPLAQKLKSSALFKSSFP G (SEQ ID NO: 265)	PCL67	MMP14_1, ADAM17_2, CTSL1_1

TABLE D Protease-Cleavable Linker Sequences		
Linker Sequence	Designation	Cleaving Protease(s)
SGGFPRSGGSFNPRTFGSKRKRRGSRGG GG (SEQ ID NO: 266)	PCL68	thrombin, factor Xa, hepsin
SGPLAQKLKSSGPAGLYAQALFKSSFPGSK (SEQ ID NO: 267)	PCL69	ADAM17_2, MMP14_1, CTSL1_1
TSTSGRSANPRGGGAVGLLAPP (SEQ ID NO: 268)	PCL70	
TSTSGRSANPRGGGVHMLPLGFLGP (SEQ ID NO: 269)	PCL71	
VHMLPLGFLGPGGLSGRSDNH (SEQ ID NO: 270)	PCL72	
VHMLPLGFLGPGGTSTSGRSANPRG (SEQ ID NO: 271)	PCL73	
SGRSAGGGSGRSAGGGSGRSA (SEQ ID NO: 272)	PCL74	uPA
HPVGLLARGGGHPVGLLARGGGSGRSAG GGSGRSA (SEQ ID NO: 273)	PCL75	MPA (MMP-2 and uPA)
GPLGVRGK (SEQ ID NO: 274)	PCL76	MMP-2
HPVGLLAR (SEQ ID NO: 163)	PCL77	MMP-2
GPQGIAGQ (SEQ ID NO: 275)	PCL78	MMP-2, MMP-9, and to some degree MT1-MMP
VPMSMRGG (SEQ ID NO: 276)	PCL79	MMP-9 and MMP-2
IPVSLRSG (SEQ ID NO: 277)	PCL80	MMP-2, and to some degree MMP-9 or MMP-7
RPFSMIMG (SEQ ID NO: 278)	PCL81	MMP-9 and MMP-7, to some degree MMP-3
VPLSLTMG (SEQ ID NO: 279)	PCL82	MMP-7, to some degree MMP-9, MMP-2, MPT-1-MMP
VPLSLYSG (SEQ ID NO: 280)	PCL83	MMP-2, MMP-9, MMP-7
IPESLRAG (SEQ ID NO: 281)	PCL84	MMP-2, MMP-7, MMP-9, to some degree MPT-1-MMP
GISSGLLSGRSDNHG (SEQ ID NO: 282)	PCL85	
GGGSISSGLLSGRSDNHGGGS (SEQ ID NO: 283)	PCL86	
GGGISSGLLSGRSDNHGGGS (SEQ ID NO: 284)	PCL87	
GGGHPVGLLARGGGGS (SEQ ID NO: 285)	PCL88	
GGSGGGSGGGGISSGLLSGRSDNHGGG SGGGSGGS (SEQ ID NO: 286)	PCL89	
GGGGISSGLLSGRSDNHGGGISSGLLSGR SDNHGGGS (SEQ ID NO: 287)	PCL90	
GGSGGSIIPVSLRSGGGISSGLLSGRSDN HGGSGGS (SEQ ID NO: 288)	PCL91	

TABLE D Protease-Cleavable Linker Sequences		
Linker Sequence	Designation	Cleaving Protease(s)
GGSGGGSVPLSLYSGGGISSGLLSGRSDN HGGSGGS (SEQ ID NO: 289)	PCL92	
GGGSHPVGLLARGGGHPVGLLARGGGHP VGLLARGS (SEQ ID NO: 290)	PCL93	
GGGSHPVGLLARGGGHPVGLLARGGSGR SAGGSGRS (SEQ ID NO: 291)	PCL94	

[0123] In certain aspects, the protease-cleavable linker comprises an amino acid sequence having up to 5, up to 4, up to 3, up to 2 or up to 1 amino acid substitution(s) as compared to the sequence set forth in Table D. Thus, in some embodiments, the protease-cleavable linker comprises or consists of any amino acid sequence in Table D with 1-5 amino acid substitutions as compared to the sequence set forth in Table D.

6.6. Non-Cleavable Linkers

[0124] In certain aspects, the present disclosure provides IL2 proproteins in which two or more components of an IL2 proprotein are connected to one another by a peptide linker. By way of example and not limitation, linkers can be used to connect an Fc domain and a targeting moiety or different domains within a targeting moiety (e.g., VH and VL domains in an scFv).

[0125] Preferably, all linkers in the IL2 proprotein other than the protease-cleavable linkers whose cleavage results in activation of IL2 are non-cleavable linkers (NCLs).

[0126] A non-cleavable linker can range from 2 amino acids to 60 or more amino acids, and in certain aspects a non-cleavable peptide linker ranges from 3 amino acids to 50 amino acids, from 4 to 30 amino acids, from 5 to 25 amino acids, from 10 to 25 amino acids, 10 amino acids to 60 amino acids, from 12 amino acids to 20 amino acids, from 20 amino acids to 50 amino acids, or from 25 amino acids to 35 amino acids in length.

[0127] In particular aspects, a non-cleavable linker is at least 5 amino acids, at least 6 amino acids or at least 7 amino acids in length and optionally is up to 30 amino acids, up to 40 amino acids, up to 50 amino acids or up to 60 amino acids in length.

[0128] In some embodiments of the foregoing, the non-cleavable linker ranges from 5 amino acids to 50 amino acids in length, e.g., ranges from 5 to 50, from 5 to 45, from 5 to 40, from 5 to 35, from 5 to 30, from 5 to 25, or from 5 to 20 amino acids in length. In other embodiments of the foregoing, the non-cleavable linker ranges from 6 amino acids to 50 amino acids in length,

e.g., ranges from 6 to 50, from 6 to 45, from 6 to 40, from 6 to 35, from 6 to 30, from 6 to 25, or from 6 to 20 amino acids in length. In yet other embodiments of the foregoing, the non-cleavable linker ranges from 7 amino acids to 50 amino acids in length, e.g., ranges from 7 to 50, from 7 to 45, from 7 to 40, from 7 to 35, from 7 to 30, from 7 to 25, or from 7 to 20 amino acids in length.

[0129] Charged (e.g., charged hydrophilic linkers) and/or flexible non-cleavable linkers are particularly preferred.

[0130] Examples of flexible non-cleavable linkers that can be used in the IL2 proproteins of the disclosure include those disclosed by Chen *et al.*, 2013, Adv Drug Deliv Rev. 65(10): 1357-1369 and Klein *et al.*, 2014, Protein Engineering, Design & Selection 27(10): 325-330. Particularly useful flexible non-cleavable linkers are or comprise repeats of glycines and serines, e.g., a monomer or multimer of G_nS (SEQ ID NO: 292) or SG_n (SEQ ID NO: 293), where n is an integer from 1 to 10, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10. In one embodiment, the non-cleavable linker is or comprises a monomer or multimer of repeat of G₄S (SEQ ID NO: 294) e.g., (GGGGS)_n (SEQ ID NO: 295).

[0131] Polyglycine non-cleavable linkers can suitably be used in the IL2 proproteins of the disclosure. In some embodiments, a peptide non-cleavable linker comprises two consecutive glycines (2Gly), three consecutive glycines (3Gly), four consecutive glycines (4Gly) (SEQ ID NO: 296), five consecutive glycines (5Gly) (SEQ ID NO: 297), six consecutive glycines (6Gly) (SEQ ID NO: 298), seven consecutive glycines (7Gly) (SEQ ID NO: 299), eight consecutive glycines (8Gly) (SEQ ID NO: 300) or nine consecutive glycines (9Gly) (SEQ ID NO: 301).

[0132] Exemplary non-cleavable linker sequences are set forth in Table E below.

TABLE E Non-Cleavable Linker Sequences	
Linker Sequence	Designation
GGGGSLALGPGGGGSLALGPGGGG GSLALGPGG (SEQ ID NO: 302)	NCL1
GGGGSGGGSGGGSGGGSGGGSGGGG S (SEQ ID NO: 303)	NCL2
(GGGGS) _n (SEQ ID NO: 295)	NCL3
(GGGS) _n (SEQ ID NO: 304)	NCL4
(GGS) _n	NCL5
(GS) _n	NCL6
(GSGGS) _n (SEQ ID NO: 305)	NCL7
ADAAP (SEQ ID NO: 306)	NCL8
ADAAPTVSIFP (SEQ ID NO: 307)	NCL9

TABLE E Non-Cleavable Linker Sequences	
Linker Sequence	Designation
ADAAPTVSIFPP (SEQ ID NO: 308)	NCL10
AKTTAP (SEQ ID NO: 309)	NCL11
AKTTAPSVYPLAP (SEQ ID NO: 310)	NCL12
AKTTPKLEEGEFSEARV (SEQ ID NO: 311)	NCL13
AKTTPKLGG (SEQ ID NO: 312)	NCL14
AKTTPP (SEQ ID NO: 313)	NCL15
AKTTPPSVTPLAP (SEQ ID NO: 314)	NCL16
ASTKGP (SEQ ID NO: 315)	NCL17
ASTKGPSVFPLAPASTKGPSVFPLAP (SEQ ID NO: 316)	NCL18
EGKSSGSGSESKST (SEQ ID NO: 317)	NCL19
GEGESGEGESGEGES (SEQ ID NO: 318)	NCL20
GEGESGEGESGEGESGEGES (SEQ ID NO: 319)	NCL21
GEGGSGEGGSGEGGS (SEQ ID NO: 320)	NCL22
GENKVEYAPALMALS (SEQ ID NO: 321)	NCL23
GGEGSGGEGSGGEGS (SEQ ID NO: 322)	NCL24
GGGESGGEGSGGEGGS (SEQ ID NO: 323)	NCL25
GGGESGGGESGGGES (SEQ ID NO: 324)	NCL26
GGGGSGGGGS (SEQ ID NO: 169)	NCL27
GGGGSGGGGSGGGGS (SEQ ID NO: 170)	NCL28
GGGGSGGGGSGGGGSGGGGS (SEQ ID NO: 171)	NCL29
GGGKSGGGKSGGGKS (SEQ ID NO: 172)	NCL30
GGGKSGGKGSGKGGGS (SEQ ID NO: 173)	NCL31
GGGS (SEQ ID NO: 174)	NCL32
GGGSG (SEQ ID NO: 175)	NCL33
GGKSGGKGSGGKGS (SEQ ID NO: 176)	NCL34
GGG	NCL35
GGSG (SEQ ID NO: 325)	NCL36
GGSGG (SEQ ID NO: 326)	NCL37
GGSGG (SEQ ID NO: 327)	NCL38
GGSGGGGSG (SEQ ID NO: 328)	NCL39

TABLE E Non-Cleavable Linker Sequences	
Linker Sequence	Designation
GGSGGGGSGGGGS (SEQ ID NO: 177)	NCL40
GHEAAVMQVQYPAS (SEQ ID NO: 329)	NCL41
GKGGSGKGGSGKGG (SEQ ID NO: 330)	NCL42
GKGKSGKGKSGKGKS (SEQ ID NO: 331)	NCL43
GKGKSGKGKSGKGKSGKGKS (SEQ ID NO: 332)	NCL44
GKPGSGKPGSGKPGS (SEQ ID NO: 333)	NCL45
GKPGSGKPGSGKPGSGKPSGS (SEQ ID NO: 334)	NCL46
GPAKELTPLKEAKVS (SEQ ID NO: 335)	NCL47
GSAGSAAGSGEF (SEQ ID NO: 336)	NCL48
GSGGG (SEQ ID NO: 180)	NCL49
GSGSG (SEQ ID NO: 181)	NCL50
GSS	NCL51
GSSG (SEQ ID NO: 182)	NCL52
GSSGGSGGSG (SEQ ID NO: 183)	NCL53
GSSGGSGGSGG (SEQ ID NO: 184)	NCL54
GSSGGSGGSGGS (SEQ ID NO: 185)	NCL55
GSSGGSGGSGGSG (SEQ ID NO: 186)	NCL56
GSSGGSGGSGGSGGGS (SEQ ID NO: 187)	NCL57
GSSGGSGGSGGS (SEQ ID NO: 188)	NCL58
GSSGT (SEQ ID NO: 189)	NCL59
GSSSG (SEQ ID NO: 190)	NCL60
GSTSGSGKPGSGEGSTKG (SEQ ID NO: 337)	NCL61
GTAAAGAGAAGGAAAGAAG (SEQ ID NO: 338)	NCL62
GTSGSSGSGSGGSGSGGGG (SEQ ID NO: 339)	NCL63
IRPRAIGGSKPRVA (SEQ ID NO: 340)	NCL64
KESGSVSSEQLAQFRSLD (SEQ ID NO: 341)	NCL65
KTPKLEEGEFSEAR (SEQ ID NO: 342)	NCL66
PRGASKSGSASQTGSAPGS (SEQ ID NO: 343)	NCL67
QPKAAP (SEQ ID NO: 344)	NCL68
QPKAAPSVTLFPP (SEQ ID NO: 345)	NCL69
RADAAAA(G4S) ₄ (SEQ ID NO: 346)	NCL70

TABLE E Non-Cleavable Linker Sequences	
Linker Sequence	Designation
RADAAAAGGPGS (SEQ ID NO: 347)	NCL71
RADAAP (SEQ ID NO: 348)	NCL72
RADAAPTVS (SEQ ID NO: 349)	NCL73
SAKTHP (SEQ ID NO: 350)	NCL74
SAKTHPKLEEGEFSEARV (SEQ ID NO: 351)	NCL75
SAKTHPKLGG (SEQ ID NO: 352)	NCL76
STAGDTHLGGEDFD (SEQ ID NO: 353)	NCL77
TVAAP (SEQ ID NO: 354)	NCL78
TVAAPSVFIFPP (SEQ ID NO: 355)	NCL79
TVAAPSVFIFPPTVAAPSVFIFPP (SEQ ID NO: 356)	NCL80

[0133] In certain aspects, the IL2 proprotein of the disclosure may comprise a polypeptide chain comprising, in an N- to C-terminal orientation, a targeting moiety (or targeting moiety chain), a hinge domain and a CH2 domain, and a CH3 domain. Thus, the hinge domain connects the targeting moiety with the CH2 domain and can be said to constitute a type of linker. Exemplary hinge domains are set forth in Section 6.9.3.

6.7. Targeting Moiety

[0134] The incorporation of targeting moieties in the IL2 proproteins of the disclosure permits the delivery of high concentrations of IL2 into the tumor microenvironment with a concomitant reduction of systemic exposure, resulting in fewer side effects than obtained with unmasked IL2 molecules.

[0135] It is anticipated that any type of target molecule present or capable of driving the IL2 proprotein at a particular locale or tissue may be targeted by the IL2 proproteins of the disclosure. In some embodiments, the IL2 proproteins are intended to treat cancer, e.g., by inducing a local immune response against tumor tissue. Accordingly, the targeting molecule can be any local tumor and associated target molecule. The target molecules recognized by the targeting moieties of the IL2 proproteins of the disclosure are generally found, for example, on the surfaces of activated T cells, on the surfaces of tumor cells, on the surfaces of virus-infected cells, on the surfaces of other diseased cells, free in blood serum, in the extracellular matrix (ECM), or immune cells present in the target site, e.g., tumor reactive lymphocytes.

[0136] In various extracellular matrix ("ECM") antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen ("TCA"), a checkpoint inhibitor, or a tumor-associated antigen ("TAA"). The skilled artisan would recognize that the foregoing categories of target molecules are not mutually exclusive and thus a given target molecule may fall into more than one of the foregoing categories of target molecules. For example, some molecules may be considered both TAAs and ECM proteins, and other molecules may be considered both TCAs and checkpoint inhibitors.

[0137] Exemplary types of cancers that may be targeted include acute lymphoblastic leukemia, acute myelogenous leukemia, biliary cancer, B-cell leukemia, B-cell lymphoma, biliary cancer, bone cancer, brain cancer, breast cancer, triple-negative breast cancer, cervical cancer, Burkitt lymphoma, chronic lymphocytic leukemia, chronic myelogenous leukemia, colorectal cancer, endometrial cancer, esophageal cancer, gall bladder cancer, gastric cancer, gastrointestinal tract cancer, glioma, hairy cell leukemia, head and neck cancer, Hodgkin's lymphoma, liver cancer, lung cancer, medullary thyroid cancer, melanoma, multiple myeloma, ovarian cancer, non-Hodgkin's lymphoma, pancreatic cancer, prostate cancer, pulmonary tract cancer, renal cancer, sarcoma, skin cancer, testicular cancer, urothelial cancer, and other urinary bladder cancers. However, the skilled artisan will realize that TAAs and other target molecules associated with the tumor microenvironment are known for virtually any type of cancer.

[0138] Non-limiting examples of ECM antigens include syndecan, heparanase, integrins, osteopontin, link, cadherins, laminin, laminin type EGF, lectin, fibronectin, notch, nectin (e.g., nectin-4), tenascin, collagen (e.g., collagen type X) and matrixin.

[0139] Other target molecules are cell surface molecules of tumor or viral lymphocytes, for example T-cell co-stimulatory proteins such as CD27, CD28, 4-1BB (CD137), OX40, CD30, CD40, ICOS, lymphocyte function-associated antigen-1 (LFA-1), CD2, CD7, LIGHT, NKG2C, and B7-H3.

[0140] In particular embodiments, the target molecules are checkpoint inhibitors, for example CTLA-4, PD1, PDL1, PDL2, B7-H3, B7-H4, BTLA, HVEM, TIM3, GAL9, LAG3, VISTA, KIR, 2B4, CD160, CGEN-15049, CHK1, CHK2. In particular embodiments, the target molecule is PD1. In other embodiments, the target molecule is LAG3.

[0141] The antibodies and antigen-binding portions generally bind to specific antigenic determinants and are able to direct the IL2 proprotein to a target site, for example to a specific type of tumor cell or tumor stroma that bears the antigenic determinant. In particular

embodiments, the targeting moiety recognizes a tumor-associated antigen (TAA). Preferably, the TAA is a human TAA. The antigen may or may not be present on normal cells. In certain embodiments, the TAA is preferentially expressed or upregulated on tumor cells as compared to normal cells. In other embodiments, the TAA is a lineage marker. Exemplary TAAs include

5 Fibroblast Activation Protein (FAP), the A1 domain of Tenascin-C (TNC A1), the A2 domain of Tenascin-C (TNC A2), the Extra Domain B of Fibronectin (EDB), the Melanoma-associated Chondroitin Sulfate Proteoglycan (MCSP), MART-1/Melan-A, gp100, Dipeptidyl peptidase IV (DPPIV), adenosine deaminase-binding protein (ADA bp), cyclophilin b, colorectal associated antigen (CRC)-C017-1A/GA733, Carcinoembryonic Antigen (CEA) and its immunogenic

10 epitopes CAP-1 and CAP-2, etv6, aml1, Prostate Specific Antigen (PSA) and its immunogenic epitopes PSA-1, PSA-2, and PSA-3, prostate-specific membrane antigen (PSMA), T-cell receptor/CD3-zeta chain, MAGE-family of tumor antigens (e.g., MAGE-A1, MAGE-A2, MAGE-A3, MAGE-A4, MAGE-A5, MAGE-A6, MAGE-A7, MAGE-A8, MAGE-A9, MAGE-A10, MAGE-A11, MAGE-A12, MAGE-Xp2 (MAGE-B2), MAGE-Xp3 (MAGE-B3), MAGE-Xp4 (MAGE-B4),

15 MAGE-C1, MAGE-C2, MAGE-C3, MAGE-C4, MAGE-C5), GAGE-family of tumor antigens (e.g., GAGE-1, GAGE-2, GAGE-3, GAGE-4, GAGE-5, GAGE-6, GAGE-7, GAGE-8, GAGE-9), BAGE, RAGE, LAGE-1, NAG, GnT-V, MUM-1, CDK4, tyrosinase, p53, MUC family, HER2/neu, p21ras, RCAS1, α -fetoprotein, E-cadherin, α -catenin, β -catenin and γ -catenin, p120ctn, gp100 Pmel117, PRAME, NY-ESO-1, cdc27, adenomatous polyposis coli protein (APC), fodrin, Connexin 37, Ig-

20 idiotypic, p15, gp75, GM2 and GD2 gangliosides, viral products such as human papilloma virus proteins, Smad family of tumor antigens, Imp-1, P1A, EBV-encoded nuclear antigen (EBNA)-1, brain glycogen phosphorylase, SSX-1, SSX-2 (HOM-MEL-40), SSX-1, SSX-4, SSX-5, SCP-1 and CT-7, c-erbB-2, Her2, EGFR, IGF-1R, CD2 (T-cell surface antigen), CD3 (heteromultimer associated with the TCR), CD22 (B-cell receptor), CD23 (low affinity IgE receptor), CD30

25 (cytokine receptor), CD33 (myeloid cell surface antigen), CD40 (tumor necrosis factor receptor), IL-6R-(IL6 receptor), CD20, MCSP, PDGF β R (β -platelet-derived growth factor receptor), ErbB2 epithelial cell adhesion molecule (EpCAM), EGFR variant III (EGFRvIII), CD19, disialoganglioside GD2, ductal-epithelial mucine, gp36, TAG-72, glioma-associated antigen, β -human chorionic gonadotropin, α -fetoprotein (AFP), lectin-reactive AFP, thyroglobulin, MN-CA IX, human telomerase reverse transcriptase, RU1, RU2 (AS), intestinal carboxyl esterase, mut hsp70-2, M-CSF, prostase, prostase specific antigen (PSA), PAP, LAGA-1a, p53, prostein, PSMA, surviving and telomerase, prostate-carcinoma tumor antigen-1 (PCTA-1), ELF2M, neutrophil elastase, ephrin B2, insulin growth factor (IGF1)-I, IGF-II, IGF1 receptor, 5T4, ROR1,

30

Nkp30, NKG2D, tumor stromal antigens, the extra domain A (EDA) and extra domain B (EDB) of fibronectin and the A1 domain of tenascin-C(TnC A1).

[0142] Suitable targeting moiety formats are described in Section 6.8. The targeting moiety is preferably an antigen binding moiety, for example an antibody or an antigen-binding portion of an antibody, e.g., an scFv, as described in Section 6.8.2 or a Fab, as described in Section 6.8.1.

[0143] In some embodiments, the targeting moieties target the exemplary target molecules set forth in Table F below, together with references to exemplary antibodies or antibody sequences upon which the targeting moiety can be based.

TABLE F Exemplary Target Molecules	
Target	Antibody Name and/or Binding Sequences
1-92-LFA-3	Amevive™ (alefacept)
5T4	GEN1044
Activin Receptor Type II	Bimagrumab VH: SEQ ID NOs:107, 109 of US Patent No. 8,388,968 B2 VL: SEQ ID NOs:93, 95 of US Patent No. 8,388,968 B2
B7-H3	Obrindatamab (MGD009)
B7-H3 (CD276)	Enoblituzumab (MGA271)
B7-H3 (CD276)	MGC018
B7-H3 (CD276)	MGA012
B7-H3 (CD276)	8H9
B7-H3 (CD276)	VH: the VH sequence of the heavy chain of SEQ ID NO:21, 26 or 31 of US 2021/0171641 A1. VL: the VL sequence of the light chain of SEQ ID NO:20, 22 or 30 of US 2021/0171641 A1.
B7-H3 (CD276)	VH: the VH sequence of the heavy chain of SEQ ID NO:21, 29 or 37 of US 2019/0002563 A1. VL: the VL sequence of the light chain of SEQ ID NO:17, 25 or 33 of US 2019/0002563 A1.
B7-H3 (CD276)	VH: the VH sequence of the heavy chain of SEQ ID NO:146, 147 or 148 of US Patent No. 10,640,563. VL: the VL sequence of the light chain of SEQ ID NO:143, 144 or 145 of US Patent No. 10,640,563.
BAFF/B Lymphocyte Stimulator	Benlysta™ (velimumab)
BAFF/B Lymphocyte Stimulator	VH: amino acids 1-123 of SEQ ID NO:327 of US Patent No. 7,138,501 VL: amino acids 139-249 of SEQ ID NO:327 of US Patent No. 7,138,501.
BAFF/B Lymphocyte Stimulator	VH: amino acids 1-126 of SEQ ID NO:1321 of US Patent No. 7,605,236;

TABLE F Exemplary Target Molecules	
Target	Antibody Name and/or Binding Sequences
	VL: amino acids 143-251 of SEQ ID NO:1049 of US Patent No. 7,605,236.
BAFF/B Lymphocyte Stimulator	Belimumab
BCMA	VH: the VH sequence of the heavy chain of SEQ ID NO. 126 of US 2021/0206865 A1 VL: the VL sequence of the light chain of SEQ ID NO. 129 or SEQ ID NO. 132 of US 2021/0206865 A1
CA125	Igobumab
CA125	OvaRex™ (oregobumab)
Cadherin	The antibodies described in US Pub. No. US 2006/0039915.
N-cadherin	An antibody that binds to the amino acid sequence of SEQ ID NO: 10, 17 or 18 of US Pub. No. US 2010/0278821.
CD11a	Raptiva™ (efalizumab) Sequence in Werther <i>et al.</i> , 1996, The Journal of Immunology 157(11): 4986-4995.
CD19	Blincyto™ (blinatumomab)
CD19	SGN-CD19A
CD20	Bexxar™ (tositumomab) VH: the VH sequence of the heavy chain of SEQ ID NO:124 of US Patent Pub. US 2017/0002060 A1 VL: the VL sequence of the light chain of SEQ ID NO:125 of US Patent Pub. US 2017/0002060 A1
CD20	Zevalin™ (ibritumomab tiuxetan) VH: SEQ ID NO:9 of US Patent No. 5,736,137 VL: SEQ ID NO:6 of US Patent No. 5,736,137
CD20	Rituxan™ (rituximab) VH: SEQ ID NO:9 of US Patent No. 5,736,137 VL: SEQ ID NO:6 of US Patent No. 5,736,137
CD20	Ocrevus™ (ocrelizumab)
CD20	Okaratuzumab
CD20	Arzerra™ (ofatumumab) VH: SEQ ID NO:2 of US Patent No. 8,529,902 VL: SEQ ID NO:4 of US Patent No. 8,529,902
CD20	Gazyva™ (obinutuzumab)
CD20	VH: SEQ ID NO: 4 of US 2021/0206870 A1 VL of SEQ ID NO:6 of US 2021/0206870 A1
CD20	Epcoritamab
CD22	Belimumab
CD22	Epratuzumab

TABLE F Exemplary Target Molecules	
Target	Antibody Name and/or Binding Sequences
CD22	Besponsa [™] (inotuzumab ozogamicin)
CD22	Lumoxiti [™] (moxetumumab pasudox)
CD22	pinatuzumab vedotin
CD25	Zenapax [™] (daclizumab) VH: SEQ ID NO:9 of US Patent No. 7,060,269 VL: SEQ ID NO:10 of US Patent No. 7,060,269
CD30	Adcetris [™] (brentuximab vedotin) VH: SEQ ID NO:2 of US Patent No. 7,090,843 VL: SEQ ID NO:10 of US Patent No. 7,090,843
CD33	Myelotarg [™] (gemtuzumab) Sequence in Man Sung, <i>et al.</i> , 1993, Molecular immunology 30:1361-1367
CD33	Lintuzumab
CD38	Darzalex [™] (daratumumab)
CD40	Lukatumumab
CD40	Dacetuzumab
CD40L	Hu5c8 (ruplizumab)
CD44v6	vibatuzumab mertansine
CD52	Campath [™] (alemtuzumab) VH: SEQ ID NO:1 of US Patent Pub. US 2017/0002060 A1 VL: SEQ ID NO:2 of US Patent Pub. US 2017/0002060 A1
CD70	Blenrep [™] (borsetuzumab mafodotin)
CD123	Flotetuzumab
CD221	Tepezza [™] (teprotumumab)
CEA	Hybri-Ceaker® (altumomab pentetate)
CEA	Scintimun [™] (besilesomab)
CEA	CEA-CIDE [™] (labetuzumab))
CEA	CEA-Scan [™] (arcitumomab)
CEA	hMN-15 CDR-H1, CDR-H2 and CDR-H3 sequences of SEQ ID NOs:4-6 of US Patent No. 8,771,690 B2 CDR-L1, CDR-L2 and CDR-L3 sequences of SEQ ID NOs:1-3 of US Patent No. 8,771,690 B2
CEA	CEA binding portion of RO6958688/RG7802 from clinical trial NCT02324257
CEA	Cibisatamab
CEA	CEA binding portion of MEDI-565/MT110/AMG211 from clinical trials NCT01284231 and NCT02291614 VH: SEQ ID NO:49 or 51 of PCT Publication No. WO 2013/012414 A1

TABLE F Exemplary Target Molecules	
Target	Antibody Name and/or Binding Sequences
	VL: SEQ ID NO:48 of PCT Publication No. WO 2013/012414 A1.
CEA	Rabetuzumab
CEA	Atezolizumab
CEA	Cibisatamab
CEA	MEDI-565 (AMG211, MT111)
CEA	RO6958688
CEA	VH: SEQ ID No. 9 described in WO2022/048883A1 VL: SEQ ID No. 10 described in WO2022/048883A1
CLDN18.2	AMG910
Collagen alpha-4 chain	TRC093 (MT293)
Collagen	The collagen binding antibody fragment described in Liang, H. et al. A collagen-binding EGFR antibody fragment targeting tumors with a collagen-rich extracellular matrix. Sci. Rep. 5, 18205; doi: 10.1038/srep18205 (2016).
Collagen type I	Cetuximab (Erbiximab)
Collagen type X	The amino acid sequences of SEQ ID NO: 1 or 2 of PCT Pub No. WO 2019/020797.
Collagen type X	The amino acid sequences of SEQ ID NO: 1 of PCT Pub No. WO 2014/180992.
Collagen type X	Antibody X34 as described in I. Girkontaite et al., "Immunolocalization of type X collagen in normal fetal and adult osteoarthritic cartilage with monoclonal antibodies," Matrix Biol 15, 231-238 (1996).
Collagen type X	Antibodies X53 or 1H8 or ARC0659 or JF0961 collagen X polyclonal antibody sold under catalog number PA5-115039 or PA5-116871 or PA5-97603 or PA5-49198 from ThermoFisher Scientific.
Collagen type X	Antibody sold under catalog number RDI-COLL10abr from RDI.
Complement C5	Soliris™ (eculizumab) VH: amino acids 1-122 of SEQ ID NO: 10 of US Patent No. 6,355,245 VL: amino acids 3-110 of SEQ ID NO: 9 of US Patent No. 6,355,245
CTLA-4	Yervoy™ (ipilimumab) VH: SEQ ID NO: 17 of WO 2001/014424 A2 VL: SEQ ID NO: 7 of WO 2001/014424 A2
CTLA-4	(tremelimumab)
CTLA-4	Orencia™ (abatacept)
DLL3	AMG757
EGFR	Erbiximab™ (cetuximab) VH: SEQ ID NO: 11 of US Patent No. 6,217,866

TABLE F	
Exemplary Target Molecules	
Target	Antibody Name and/or Binding Sequences
	VL: SEQ ID NO: 13 of US Patent No. 6,217,866
EGFR	Vectibix™ (panitumumab) VH: SEQ ID NO: 37 of US Patent No. 6,235,883 VL: SEQ ID NO: 38 of US Patent No. 6,235,883
EGFR	Zalutumumab VH: SEQ ID NO: 64 of WO 2018/140831 A2 VL: SEQ ID NO: 69 of WO 2018/140831 A2
EGFR	Mapatumumab
EGFR	Matuzumab
EGFR	Nimotuzumab VH: SEQ ID NO: 51 of WO 2018/140831 A2 VL: SEQ ID NO: 56 of WO 2018/140831 A2
EGFR	ICR62
EGFR	mAb 528
EGFR	CH806
EGFRv3	AMG596
EGFRv3	AMG404
EGFR/CD64	MDX-447
EpCAM	Panorex™ (edrecolomab) VH: SEQ ID NO: 129 of WO 2018/140831 A2 VL: SEQ ID NO: 134 of WO 2018/140831 A2
EpCAM	Adecatumumab VH: SEQ ID NO: 142 of WO 2018/140831 A2 VL: SEQ ID NO: 147 of WO 2018/140831 A2
EpCAM	tucotuzumab celmoleukin
EpCAM	citatumumab bogatox
EpCAM	EP1629013 B1 VH: SEQ ID NOS: 80, 84, 88, 92 or 96 VL: SEQ ID NOS: 82, 86, 90, 94 or 98
EpCAM	G8.8 HC: SEQ ID NO: 4 of US Patent Pub. No. US 2020/0317806 A1 HL: SEQ ID NO: 3 of US Patent Pub. No. US 2020/0317806 A1
EpCAM	VH: SEQ ID NOS: 17-22 of WO 2021/211510 A2. VL: SEQ ID NO: 15-16 of WO 2021/211510 A2.
EpCAM	Removab™ (catumaxomab)
EpCAM	Vicineum™ (oportuzumab monatox)
EpCAM	M701
F protein of RSV	Synagic™ (palivizumab)
GD2	3F8
Glycoprotein receptor IIb/IIIa	ReoPro™ (abiximab)
gpA33	MGD007

TABLE F Exemplary Target Molecules	
Target	Antibody Name and/or Binding Sequences
GPC3	ERY974
GUCY2C	PF-07062119
Heparanase	An antibody selected from HP130, HP 239, HP 108.264, HP 115.140, HP 152.197, HP 110.662, HP 144.141, HP 108.371, HP 135.108, HP 151.316, HP 117.372, HP 37/33, HP3/17, HP 201 or HP 102 or an amino acid sequence of SEQ ID NO: 1-11 described in US Patent Pub. US 2004/0170631.
Her2	Herceptin™ (trastuzumab)
Her2	Aldesleukin (proleukine)
Her2	Sargramustim (leukine)
Her2	M802
Her2	Runimotamab (BTRC4017A, R07227780)
Her2	ISB1302
Her2-neu	Perjeta™ (pertuzumab) VH: SEQ ID NO:16 of WO 2013/096812 A1. VL: SEQ ID NO:15 of WO 2013/096812 A1.
Her2-neu	Rexomun™ (ertumaxomab)
IgE	Xolair™ (omalizumab)
IGFIR	(figitumumab)
IL1β	Ilaris™ (canakinumab) VH: SEQ ID NO:1 of US Patent No. 7,446,175. VL: SEQ ID NO: 2 of US Patent No. 7,446,175
IL12/IL23	Stelara™ (ustekinumab)
IL1Ra	Antril™, Kineret™ (ankinra)
IL2R	Simulect™ (basiliximab) VH: SEQ ID NO: 3 of US Patent No. 6,383,487 VL: SEQ ID NO: 6 of US Patent No. 6,383,487
IL6	Clazakizumab
IL6 receptor	Actemra™ (tocilizumab) VH: SEQ ID NO: 31 of US Patent No. 7,479,543 VL: SEQ ID NO: 29 of US Patent No. 7,479,543
IL12/IL23 p40 subunit	Stelara™ (ustekinumab) VH: SEQ ID NO: 7 of US Patent No. 6,902,734 VL: SEQ ID NO: 8 of US Patent No. 6,902,734
Integrin α4	Tysabri™ (natalizumab) VH: SEQ ID NOs: 11-13 of US Patent No. 5,840,299 VL: SEQ ID NOs: 7-8 of US Patent No. 5,840,299
Integrin α4 β7	Entyvio™ (vedolizumab) HC: SEQ ID NO: 2 of US Patent Pub. US 2012/0282249. LC: SEQ ID NO: 4 of US Patent Pub. US 2012/0282249.
Integrin α5 β1	VH: SEQ ID NO: 2 of European Patent No. 1 755 659. VL: SEQ ID NO: 4 of European Patent No. 1 755 659.

TABLE F Exemplary Target Molecules	
Target	Antibody Name and/or Binding Sequences
Integrin β 1	VH: SEQ ID NO:2, 6, 8, 10, 12, 14, 29-43 or 91-100 of US Patent Pub. US 2022/0089744. VL: SEQ ID NO:4, 16, 18, 20, 22, 44-57 or 107-116 of US Patent Pub. US 2022/0089744.
LAG-3	Relatlimab (BMS-98016)
LAG-3	Sym022
LAG-3	HLX26
LAG-3	TSR-033
LAG-3	ABL501
LAG-3	INCAGN02385
LAG-3	Fianlimab (REGN3767)
LAG-3	RO7247669
LAG-3	EMB-02
LAG-3	FS118
LAG-3	GSK2831781
LAG-3	IBI323
LAG-3	IBI110
LAG-3	LAG525
LAG-3	XmAb@22841
LAG-3	LBL-007
LAG-3	VH: SEQ ID NO:1, 8, 10 or 12 of US Patent No. 9,902,772. VL: SEQ ID NO:2, 3, 4, 5, 6, 7, 9, 11, 13 or 14 of US Patent No. 9,902,772.
LAG-3	VH: SEQ ID NO: 182 of US Patent Pub. US 2021/0095026. VL: SEQ ID NO: 88 of US Patent Pub. US 2021/0095026.
Laminin	Lam-89 from Sigma Aldrich
Mesothelin	Amatuximab
Mesothelin	HPN536
MUC1	civatuzumab tetraxetane
MUC1	Pankomab TM (gatipotuzumab)
MUC1	femtumumab
MUC1	Cantuzumab ravtansine
MUC16 (CA125)	Anti-MUC16 antibodies having VH and VL sequences having the amino acid sequences of any one of the following SEQ ID NO: pairs from US 2018/0118848A1: 18/26; 82/858; 98/170
MUC17	AMG199
Nectin-4	Enfortumab (ASP7465, ASG-22CE, ASG-22ME) VH: SEQ ID NO:3 of PCT Pub. WO 2021/151984. VL: SEQ ID NO: 4 of PCT Pub. WO 2021/151984.
Nectin-4	SBT290

TABLE F	
Exemplary Target Molecules	
Target	Antibody Name and/or Binding Sequences
Nectin-4	VH: SEQ ID NO:1 of US Patent No. 11,274,160. VL: SEQ ID NO:2 of US Patent No. 11,274,160.
NGF	(tanezumab)
Osteopontin	HC: SEQ ID NO: 22 of PCT Pub. WO 2021/030209. LC: SEQ ID NO: 24 of PCT Pub. WO 2021/030209.
PD1	MDX-1106 / BMS-936558 (nivolumab), a human IgG4 mAb with the structure described in WHO Drug Information, Vol. 27, No. 1, pages 68-69 (2013) and whose heavy and light chain sequences are disclosed in FIG. 7 of US Pub. No. US20190270812A1
PD1	MK-3475 (pembrolizumab), a humanized IgG4 mAb with the structure described in WHO Drug Information, Vol. 27, No. 2, pages 161-162 (2013) and whose heavy and light chain sequences are disclosed in FIG. 6 of US Pub. No. US20190270812A1
PD1	REGN2810 (disclosed as H4H7798N in US Pub No. 20150203579)
PD1	Anti-PD1 antibodies disclosed in Tables 1-3 of PCT Pub. WO2015112800A1, including but not limited to anti-PD1 antibodies having VH/VL pairs having SEQ ID NOs: 2/10, 18/26, 34/42, 50/58, 66/74, 82/90, 98/106, 114/122, 130/138, 146/154, 162/170, 178/186, 194/202, 210/202, 218/202, 226/202, 234/202, 242/202, 250/202, 258/202, 266/202, 274/202, 282/202, 290/202, 298/186, 306/186 and 314/186 of PCT Pub. WO2015112800A1.
PD1	MEDI-0680 (AMP-514)
PD1	PDR001
PD1	BGB-108
PD1	h409A11, described in WO2008/156712
PD1	h409A16, described in WO2008/156712
PD1	h409A17, described in WO2008/156712
PD1	Anti-PD1 antibodies described in US Patent No. 7488802
PD1	Anti-PD1 antibodies described in US Patent No. 7521051
PD1	Anti-PD1 antibodies described in US Patent No. 8008449
PD1	Anti-PD1 antibodies described in US Patent No. 8354509
PD1	Anti-PD1 antibodies described in US Patent No. 8168757
PD1	Anti-PD1 antibodies described in PCT Pub. No. W02004/004771
PD1	Anti-PD1 antibodies described in PCT Pub. No. W02004/056875
PD1	Anti-PD1 antibodies described in PCT Pub. No. W02004/072286

TABLE F	
Exemplary Target Molecules	
Target	Antibody Name and/or Binding Sequences
PD1	Anti-PD1 antibodies described in US Pub. No. US2011/0271358
PDL1	Durvalumab (MEDI4736)
PDL1	Atezolizumab (Tecentriq, MPDL3280A))
PDL1	MDX 1105 (BMS-936559)
PDL1	Avelumab
PDL1	ZKAB001 (Socazolimab)
PDL1	TQB2450
PDL1	MEDI4736
PDL1	HLX20
PDL1	KN035
PDL1	LY3434172
PDL1	LY3300054
PDL1	LDP
PDL1	EMB-09
PDL1	ABL501
PDL1	INBRX-105
PDL1	SHR-1210
PDL1	STI-3031 (IMC-001)
PDL1	MPDL3280A (RG7446)
PDL1	KN035
PDL1	BGB-A333
PDL1	HLX301
PDL1	Y101D
PDL1	ES101
PDL1	IBI322
PDL1	Envafolimab
PDL1	VH: SEQ ID NO:46, 48, 50 or 52 of US Patent No. 11,168,144. VL: SEQ ID NO:58, 137 or 12 of US Patent No. 11,168,144.
PDL1	VH: SEQ ID NO:23, 124, 126, 127, 128, 130, 140 or 145 of US Patent No. 11,208,486. VL: SEQ ID NO:24 or 125 of US Patent No. 11,208,486.
Phosphatidylserine	(bavituximab)
PSCA	GEM3PSCA
PSMA	huJ591
PSMA	Anti-PSMA antibodies having VH and VL sequences having the amino acid sequences of any one of the following SEQ ID NO: pairs from WO 2017/023761A1: 2/1642; 10/1642; 18/1642; 26/1642; 34/1642; 42/1642; 50/1642; 58/1642; 66/1642; 74/1642; 82/1642; 90/1642; 98/1642; 106/1642; 114/1642; 122/130; and 138/146.

TABLE F Exemplary Target Molecules	
Target	Antibody Name and/or Binding Sequences
PSMA	An antibody such as: PSMA 3.7, PSMA 3.8, PSMA 3.9, PSMA 3.11, PSMA 5.4, PSMA 7.1, PSMA 7.3, PSMA 10.3, PSMA 1.8.3, PSMA A3.1.3, PSMA A3.3.1, Abgenix 4.248.2, Abgenix 4.360.3, Abgenix 4.7.1, Abgenix 4.4.1, Abgenix 4.177.3, Abgenix 4.16.1, Abgenix 4.22.3, Abgenix 4.28.3, Abgenix 4.40.2, Abgenix 4.48.3, Abgenix 4.49.1, Abgenix 4.209.3, Abgemx 4.219.3, Abgenix 4.288.1, Abgenix 4.333.1, Abgemx 4.54.1, Abgenix 4.153.1, Abgenix 4.232.3, Abgenix 4.292.3, Abgenix 4.304.1, Abgenix 4.78.1 and Abgenix 4.152.1 described in WO2003034903A2 A hybridoma cell line such as: PSMA 3.7 (PTA-3257), PSMA 3.8, PSMA 3.9 (PTA- 3258), PSMA 3.11 (PTA-3269), PSMA 5.4 (PTA-3268), PSMA 7.1 (PTA-3292), PSMA 7.3 (PTA-3293), PSMA 10.3 (PTA-3247) , PSMA 1.8.3 (PTA-3906), PSMA A3.1.3 (PTA- 3904), PSMA A3.3.1 (PTA-3905), Abgenix 4.248.2 (PTA-4427), Abgenix 4.360.3 (PTA- 4428), Abgenix 4.7.1 (PTA-4429), Abgenix 4.4.1 (PTA-4556), Abgenix 4.177.3 (PTA-4557), Abgenix 4.16.1 (PTA-4357), Abgenix 4.22.3 (PTA-4358), Abgenix 4.28.3 (PTA-4359), Abgenix 4.40.2 (PTA-4360), Abgenix 4.48.3 (PTA-4361), Abgenix 4.49.1 (PTA-4362), Abgenix 4.209.3 (PTA-4365), Abgenix 4.219.3 (PTA-4366), Abgenix 4.288.1 (PTA-4367), Abgenix 4.333.1 (PTA-4368), Abgenix 4.54.1 (PTA-4363), Abgenix 4.153.1 (PTA-4388), Abgenix 4.232.3 (PTA-4389), Abgenix 4.292.3 (PTA-4390), Abgenix 4.304.1 (PTA-4391), Abgenix 4.78.1 (PTA-4652), and Abgemx 4.152.1(PTA-4653) described in WO 2003/034903A2. VH of SEQ ID NOs: 2-7 described in WO 2003/034903A2 VL of SEQ ID NOs: 8-13 described in WO 2003/034903A2
PMSA	VH: SEQ ID NOs: 225, 239, 253, 267, 281, 295, 309, 323, 337, 351, 365, 379, 393, 407, 421, 435, 449, 463, 477, 491, 505, 519, 533, 547, 561, 575, 589, 603 or 617 described in WO 2011/121110A1. VL SEQ ID NOs: 230, 244, 258, 272, 286, 300, 314, 328, 342, 356, 370, 384, 398, 412, 426, 440, 454, 468, 482, 496, 510, 524, 538, 552, 566, 580, 594, 608 or 622 described in WO 2011/121110A1. VH and VL SEQ ID Nos: 235, 249, 263, 277, 291, 305, 319, 333, 347, 361, 375, 389, 403, 417, 431, 445, 459, 473, 487, 501, 515, 529, 543, 557, 571, 585, 599, 613 or 627 described in WO 2011/121110A1.

TABLE F	
Exemplary Target Molecules	
Target	Antibody Name and/or Binding Sequences
PMSA	An anti-PMSA antibody having a VL amino acid sequence of any one of SEQ ID NOs: 229-312 of US 2022/0119525 A1 and a VH of SEQ ID NO: 217 of US 2022/0119525 A1.
PMSA	ES414
PMSA	BAY2010112 (pasotuxizumab)
PMSA	CCW702
PMSA	JNJ-63898081
PMSA	CC-1
PMSA	Acapatamab
PSMA	HPN424
RAAG12	RAV12
RANKL	Prolia™ (denosumab) VH: SEQ ID NO:51 of US Patent Pub. 2017/0002060 VL: SEQ ID NO:52 of US Patent Pub. 2017/0002060
SLAMF7	Empliciti™ (elotuzumab)
SSTR2	XmAb@18087
STEAP1	VHCDR1 SEQ ID NOs: 14, 33, 182, 184 or 185 described in US20210179731A1. VHCDR2 SEQ ID NOs: 15, 21, 34, 182, 184 or 185 described in US20210179731A1. VHCDR3 SEQ ID NOs: 16 and 35 described in US20210179731A1. VH SEQ ID NOs: 182 or 184 described in US20210179731A1. VLCDR1 SEQ ID NOs: 11 or 30 described in US20210179731A1. VLCDR2 SEQ ID NOs: 12 or 31 described in US20210179731A1. VLCDR3 SEQ ID NOs: 13 or 32 described in US20210179731A1. VL SEQ ID NOs: 183 or 186 described in US20210179731A1.
STEAP1	AMG509
STEAP2	Anti-STEAP 2 antibodies having CDR-H1, CDR-H2, CDR-H3, CDR-L1, CDR-L2 and CDR-L3 sequences selected from SEQ ID NOS: (1) 4-6-8-12-14-16; (2) 20-22-24-28-30-32; (3) 36-38-40-44-46-48; (4) 52-54-56-60-62-64; (5) 68-70-72-60-62-64; (6) 76-78-80-60-62-64; (7) 84-86-88-60-62-64; (8) 92-94-96-60-62-64; (9) 100-102-104-60-62-64; (10) 108-110-112-116-118-120; (11) 124-126-128-132-134-136; (12) 140-142-144-148-150-152; (13) 156-158-160-164-166-168; (14) 172-174-176-180-182-184; (15) 188-190-192-196-198-200; (16) 204-206-208-212-

TABLE F Exemplary Target Molecules	
Target	Antibody Name and/or Binding Sequences
	214-216; (17) 220-222-224-228-230-232; (18) 236-238-240-244-246-248; (19) 252-254-256-260-262-264; (20) 268-270-272-276-278-280; (21) 284-286-288-292-294-296; (22) 300-302-304-308-310-312; (23) 316-318-320-324-326-328; (24) 332-334-336-340-342-344; (25) 348-350-352-356-358-360; (26) 364-366-368-372-374-376; and (27) 380-382-384-388-390-392 of US Patent No. 10,772,972 B2. Anti-STEAP 2 antibodies having (a) a VH comprising the amino acid of any one of SEQ ID NOs: 2, 18, 34, 50, 66, 74, 82, 90, 98, 106, 122, 138, 154, 170, 186, 202, 218, 234, 250, 266, 282, 298, 314, 330, 346, 362, and 378 of US Patent No. 10,772,972 B2; and (b) a VL comprising the amino acid sequence of any one of SEQ ID NOs: 10; 26; 42; 58; 114; 130; 146; 162; 178; 194; 210; 226, 242; 258; 274; 290; 306; 322; 338; 354; 370; and 386 of US Patent No. 10,772,972 B2. Anti-STEAP 2 antibodies having a VH/VL pair comprising the amino acid sequences of any of the following pairs of SEQ ID NOs of US Patent No. 10,772,972 B2: 2/10; 18/26; 34/42; 50/58; 66/58; 74/58; 82/58; 90/58; 98/58; 106/114; 122/130; 138/146; 154/162; 170/178; 186/194; 202/210; 218/226; 234/242; 250/258; 266/274; 282/290; 298/306; 314/322; 330/338; 346/354; 362/370; and 378/386.
Syndecan-1 (CD 138)	The B-B4 antibody described in Wijdenes et al. (1996) Br. J. Haematol., 94: 318-323
Syndecan-4	The amino acid sequence of amino acids 93 and 121 of SEQ ID NO: 1 or the amino acid sequence of amino acids 92 and 122 of SEQ ID NO:2 described in European Patent Pub. EP 2 603 236.
TGFβ	GC1008
TNFR	Enbrel™ (etanercept)
TNFα	Remicade™ (infliximab) VH: SEQ ID NO:2 of Int. Patent Publication WO201/3087911 A1 VH: SEQ ID NO:3 of Int. Patent Publication WO2013/A1087911
TNFα	Humira™ (adalimumab) VH: SEQ ID NO:4 of US Patent No. 6,258,562 VL: SEQ ID NO:3 of US Patent No. 6,258,562

TABLE F Exemplary Target Molecules	
Target	Antibody Name and/or Binding Sequences
TNF α	Cimzia TM (certolizumab pegol) VH: SEQ ID NO:14 of US Patent No. 7,012,135 VL: SEQ ID NO:9 of US Patent No. 7,012,135
TNF α	Simponi TM (golimumab) VH: SEQ ID NO:7 of US Patent No. 7,250,165 VL: SEQ ID NO:8 of US Patent No. 7,250,165
VEGF	Avastin TM (bevacizumab) VH: SEQ ID NO:9 of US Patent No. 7,060,269 VL: SEQ ID NO:10 of US Patent No. 7,060,269
VEGF	Lucentis TM (ranibizumab) VH: SEQ ID NO:4 of US Patent No. 9,914,770 VL: SEQ ID NO:2 of US Patent No. 9,914,770

[0144] In some aspects, the targeting moiety competes with an antibody set forth in Table F for binding to the target molecule. In further aspects, the targeting moiety comprises CDRs having CDR sequences of an antibody set forth in Table F. In some embodiments, the targeting moiety comprises all 6 CDR sequences of the antibody set forth in Table F. In other embodiments, the targeting moiety comprises at least the heavy chain CDR sequences (CDR-H1, CDR-H2, CDR-H3 and the light chain CDR sequences of a universal light chain. In further aspects, a targeting moiety comprises a VH comprising the amino acid sequence of the VH of an antibody set forth in Table F. In some embodiments, the targeting moiety further comprises a VL comprising the amino acid sequence of the VL of the antibody set forth in Table F. In other embodiments, the targeting moiety further comprises a universal light chain VL sequence.

[0145] In some embodiments, the targeting moiety is non-blocking or poorly-blocking of ligand-receptor binding. Examples of non-blocking or poorly-blocking anti-PD1 antibodies includes antibodies having VH/VL amino acid sequences of SEQ ID Nos: 2/10 of PCT Pub. No. WO2015/112800A1; SEQ ID Nos: 16/17 of US Patent No. 11,034,765 B2; SEQ ID Nos. 164/178, 165/179, 166/180, 167/181, 168/182, 169/183, 170/184, 171/185, 172/186, 173/187, 174/188, 175/189, 176/190 and 177/190 of US Patent No. 10,294,299 B2. Examples of non-blocking or poorly-blocking anti-LAG3 antibodies includes antibodies having VH/VL amino acid sequences of SEQ ID Nos 23/24, 3/4 and 11/12 of US Pub. US2022/0056126A1.

[0146] Additional target molecules that can be targeted by the IL2 proproteins are disclosed in Table I below and in, e.g., Hafeez *et al.*, 2020, Molecules 25:4764,

doi:10.3390/molecules25204764, particularly in Table 1. Table 1 of Hafeez *et al.* is incorporated by reference in its entirety here.

6.8. Targeting Moiety Formats

[0147] In certain aspects, the targeting moiety of an IL2 proprotein of the disclosure can be any
 5 type of antibody or fragment thereof that retains specific binding to an antigenic determinant. In
 one embodiment the targeting moiety is an immunoglobulin molecule or fragment thereof,
 particularly an IgG class immunoglobulin molecule, more particularly an IgG₁ or IgG₄
 immunoglobulin molecule. Antibody fragments include, but are not limited to, VH (or V_H)
 fragments, VL (or V_L) fragments, Fab fragments, F(ab')₂ fragments, scFv fragments, Fv
 10 fragments, minibodies, diabodies, triabodies, and tetrabodies.

6.8.1. Fab

[0148] Fab domains were traditionally produced by proteolytic cleavage of immunoglobulin
 molecules using enzymes such as papain. The Fab domains can comprise constant domain
 and variable region sequences from any suitable species, and thus can be murine, chimeric,
 15 human or humanized.

[0149] Fab domains typically comprise a CH1 domain attached to a VH domain which pairs with
 a CL domain attached to a VL domain. In a wild-type immunoglobulin, the VH domain is paired
 with the VL domain to constitute the Fv region, and the CH1 domain is paired with the CL
 domain to further stabilize the binding site. A disulfide bond between the two constant domains
 20 can further stabilize the Fab domain.

[0150] For the IL2 proproteins of the disclosure, particularly when the light chains of the
 targeting moieties are not common or universal light chains, it is advantageous to use Fab
 heterodimerization strategies to permit the correct association of Fab domains belonging to the
 same targeting moiety and minimize aberrant pairing of Fab domains belonging to different
 25 targeting moieties. For example, the Fab heterodimerization strategies shown in Table G below
 can be used:

TABLE G Fab Heterodimerization Strategies					
STRATEGY	VH	CH1	VL	CL	REFERENCE
CrossMabCH1-CL	WT	CL domain	WT	CH1 domain	Schaefer <i>et al.</i> , 2011, Cancer Cell 2011; 20:472-86; PMID:22014573.

TABLE G Fab Heterodimerization Strategies					
STRATEGY	VH	CH1	VL	CL	REFERENCE
orthogonal Fab VHVRD1CH1C RD2 - VLVRD1C λ CR D2	39K, 62E	H172A, F174G	1R, 38D, (36F)	L135Y, S176W	Lewis <i>et al.</i> , 2014, Nat Biotechnol 32:191-8
orthogonal Fab VHVRD2CH1wt - VLVRD2C λ wt	39Y	WT	38R	WT	Lewis <i>et al.</i> , 2014, Nat Biotechnol 32:191-8
TCR C α C β	39K	TCR C α	38D	TCR C β	Wu <i>et al.</i> , 2015, MAbs 7:364-76
CR3	WT	T192E	WT	N137K, S114A	Golay <i>et al.</i> , 2016, J Immunol 196:3199-211.
MUT4	WT	L143Q, S188V	WT	V133T, S176V	Golay <i>et al.</i> , 2016, J Immunol 196:3199-211.
DuetMab	WT	F126C	WT	S121C	Mazor <i>et al.</i> , 2015, MAbs 7:377-89; Mazor <i>et al.</i> , 2015, MAbs 7:461-669.
Domain exchanged	WT	CH3 + knob or hole mutation	WT	CH3 + hole or knob mutation	Wozniak-Knopp <i>et al.</i> , 2018, PLoS ONE 13(4):e0195442

[0151] Accordingly, in certain embodiments, correct association between the two polypeptides of a Fab is promoted by exchanging the VL and VH domains of the Fab for each other or exchanging the CH1 and CL domains for each other, *e.g.*, as described in WO 2009/080251.

- 5 [0152] Correct Fab pairing can also be promoted by introducing one or more amino acid modifications in the CH1 domain and one or more amino acid modifications in the CL domain of the Fab and/or one or more amino acid modifications in the VH domain and one or more amino acid modifications in the VL domain. The amino acids that are modified are typically part of the VH:VL and CH1:CL interface such that the Fab components preferentially pair with each other
- 10 rather than with components of other Fabs.

- [0153] In one embodiment, the one or more amino acid modifications are limited to the conserved framework residues of the variable (VH, VL) and constant (CH1, CL) domains as indicated by the Kabat numbering of residues. Almagro, 2008, *Frontiers In Bioscience* 13:1619-1633 provides a definition of the framework residues on the basis of Kabat, Chothia, and IMGT
- 15 numbering schemes.

[0154] In one embodiment, the modifications introduced in the VH and CH1 and/or VL and CL domains are complementary to each other. Complementarity at the heavy and light chain interface can be achieved on the basis of steric and hydrophobic contacts, electrostatic/charge interactions or a combination of the variety of interactions. The complementarity between protein surfaces is broadly described in the literature in terms of lock and key fit, knob into hole, protrusion and cavity, donor and acceptor *etc.*, all implying the nature of structural and chemical match between the two interacting surfaces.

[0155] In one embodiment, the one or more introduced modifications introduce a new hydrogen bond across the interface of the Fab components. In one embodiment, the one or more introduced modifications introduce a new salt bridge across the interface of the Fab components. Exemplary substitutions are described in WO 2014/150973 and WO 2014/082179, the contents of which are hereby incorporated by reference.

[0156] In some embodiments, the Fab domain comprises a 192E substitution in the CH1 domain and 114A and 137K substitutions in the CL domain, which introduces a salt-bridge between the CH1 and CL domains (see, *e.g.*, Golay *et al.*, 2016, J Immunol 196:3199-211).

[0157] In some embodiments, the Fab domain comprises a 143Q and 188V substitutions in the CH1 domain and 113T and 176V substitutions in the CL domain, which serves to swap hydrophobic and polar regions of contact between the CH1 and CL domain (see, *e.g.*, Golay *et al.*, 2016, J Immunol 196:3199-211).

[0158] In some embodiments, the Fab domain can comprise modifications in some or all of the VH, CH1, VL, CL domains to introduce orthogonal Fab interfaces which promote correct assembly of Fab domains (Lewis *et al.*, 2014 Nature Biotechnology 32:191-198). In an embodiment, 39K, 62E modifications are introduced in the VH domain, H172A, F174G modifications are introduced in the CH1 domain, 1 R, 38D, (36F) modifications are introduced in the VL domain, and L135Y, S176W modifications are introduced in the CL domain. In another embodiment, a 39Y modification is introduced in the VH domain and a 38R modification is introduced in the VL domain.

[0159] Fab domains can also be modified to replace the native CH1:CL disulfide bond with an engineered disulfide bond, thereby increasing the efficiency of Fab component pairing. For example, an engineered disulfide bond can be introduced by introducing a 126C in the CH1 domain and a 121 C in the CL domain (see, *e.g.*, Mazor *et al.*, 2015, MAb 7:377-89).

[0160] Fab domains can also be modified by replacing the CH1 domain and CL domain with alternative domains that promote correct assembly. For example, Wu *et al.*, 2015, MAbs 7:364-76, describes substituting the CH1 domain with the constant domain of the T cell receptor and substituting the CL domain with the b domain of the T cell receptor, and pairing these domain replacements with an additional charge-charge interaction between the VL and VH domains by introducing a 38D modification in the VL domain and a 39K modification in the VH domain.

[0161] In lieu of, or in addition to, the use of Fab heterodimerization strategies to promote correct VH-VL pairings, the VL of common light chain (also referred to as a universal light chain) can be used for each unique ABD in the IL2 proproteins of the disclosure. In various embodiments, employing a common light chain as described herein reduces the number of inappropriate species in the IL2 proproteins as compared to employing original cognate VLs. In various embodiments, the VL domains of ABDs are identified from monospecific antibodies comprising a common light chain. In various embodiments, the VH regions of the ABDs in the IL2 proproteins comprise human heavy chain variable gene segments that are rearranged *in vivo* within mouse B cells that have been previously engineered to express a limited human light chain repertoire, or a single human light chain, cognate with human heavy chains and, in response to exposure with an antigen of interest, generate an antibody repertoire containing a plurality of human VHs that are cognate with one or one of two possible human VLs, wherein the antibody repertoire specific for the antigen of interest. Common light chains are those derived from a rearranged human Vk1-39Jk5 sequence or a rearranged human Vk3-20Jk1 sequence, and include somatically mutated (*e.g.*, affinity matured) versions. See, for example, U.S. Patent No. 10,412,940.

6.8.2. scFv

[0162] Single chain Fv or "scFv" antibody fragments comprise the VH and VL domains of an antibody in a single polypeptide chain, are capable of being expressed as a single chain polypeptide, and retain the specificity of the intact antibodies from which they are derived. Generally, the scFv polypeptide further comprises a polypeptide linker between the VH and VL domain that enables the scFv to form the desired structure for target binding. Examples of linkers suitable for connecting the VH and VL chains of an scFv are the non-cleavable linkers identified in Section v.

[0163] Unless specified, as used herein an scFv may have the VL and VH variable regions in either order, *e.g.*, with respect to the N-terminal and C-terminal ends of the polypeptide, the scFv may comprise VL-linker-VH or may comprise VH-linker-VL.

[0164] The scFv can comprise VH and VL sequences from any suitable species, such as murine, human or humanized VH and VL sequences.

[0165] To create an scFv-encoding nucleic acid, the VH and VL-encoding DNA fragments are operably linked to another fragment encoding a linker, *e.g.*, encoding any of the linkers described in Section 6.6 (typically a repeat of a sequence containing the amino acids glycine and serine, such as the amino acid sequence (Gly4~Ser)3 (SEQ ID NO: 170), such that the VH and VL sequences can be expressed as a contiguous single-chain protein, with the VL and VH regions joined by the flexible linker (see, *e.g.*, Bird *et al.*, 1988, Science 242:423-426; Huston *et al.*, 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; McCafferty *et al.*, 1990, Nature 348:552-554).

6.9. Fc Regions

[0166] The IL2 proproteins of the disclosure typically include a pair of Fc domains that associate to form an Fc region. In native antibodies, Fc regions comprise hinge regions at their N-termini to form a constant domain. Throughout this disclosure, the reference to an Fc domain encompasses an Fc domain with a hinge domain at its N-terminus unless specified otherwise.

[0167] The Fc domains can be derived from any suitable species operably linked to an ABD or component thereof. In one embodiment the Fc domain is derived from a human Fc domain. In preferred embodiments, the targeting moiety or component thereof is fused to an IgG Fc molecule. A targeting moiety or component thereof may be fused to the N-terminus or the C-terminus of the IgG Fc domain or both.

[0168] The Fc domains can be derived from any suitable class of antibody, including IgA (including subclasses IgA1 and IgA2), IgD, IgE, IgG (including subclasses IgG1, IgG2, IgG3 and IgG4), and IgM. In one embodiment, the Fc domain is derived from IgG1, IgG2, IgG3 or IgG4. In one embodiment the Fc domain is derived from IgG1. In one embodiment the Fc domain is derived from IgG4. Exemplary sequences of Fc domains from IgG1, IgG2, IgG3, and IgG4 are provided in Table Y, below.

TABLE Y Fc Sequences		
Fc	Sequence	SEQ ID NO
hIgG1 Fc (amino acids 99-330 of UniprotKB P01857-1)	EPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK	1
hIgG2 Fc (amino acids 99-326 of UniprotKB P01859-1)	ERKCCVECPPCAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDISVEWESNGQPENNYKTTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK	2
hIgG3 Fc (amino acids 99-377 of UniprotKB P01860-1)	ELKTPLGDTTHTCPRCPEPKSCDTPPPCPRCPEPKSCDTPPPCPRCP EPKSCDTPPPCPRCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFKWYVDGVEVHNAKTKPREEQYNSTFRVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESSGQPENNYNTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNRFTQKSLSLSPGK	3
hIgG4 Fc (amino acids 99-327 of UniprotKB P01861-1)	ESKYGPPCPSCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLSLSLGK	4

- [0169]** In some aspects, an Fc domain comprises an amino acid sequence having at least about 90%, at least about 91%, at least about 92%, about at least 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% sequence identity to SEQ ID NO:1. In cases where an Fc domain comprises at least 90% sequence identity and less than 100% sequence identity to SEQ ID NO:1 (e.g., between 90% and 99% sequence identity to SEQ ID NO:1), an Fc domain may also comprise one or more amino acid substitutions described herein, for example one or more substitutions

that reduce effector function (e.g., as described in Section 6.9.1) and/or one or more substitutions that promote Fc heterodimerization (e.g., as described in Section 6.9.2).

[0170] In some aspects, an Fc domain comprises an amino acid sequence having at least about 90%, at least about 91%, at least about 92%, about at least 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% sequence identity to SEQ ID NO:2. In cases where an Fc domain comprises at least 90% sequence identity and less than 100% sequence identity to SEQ ID NO:2 (e.g., between 90% and 99% sequence identity to SEQ ID NO:2), an Fc domain may also comprise one or more amino acid substitutions described herein, for example one or more substitutions that reduce effector function (e.g., as described in Section 6.9.1) and/or one or more substitutions that promote Fc heterodimerization (e.g., as described in Section 6.9.2).

[0171] In some aspects, an Fc domain comprises an amino acid sequence having at least about 90%, at least about 91%, at least about 92%, about at least 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% sequence identity to SEQ ID NO:3. In cases where an Fc domain comprises at least 90% sequence identity and less than 100% sequence identity to SEQ ID NO:3 (e.g., between 90% and 99% sequence identity to SEQ ID NO:3), an Fc domain may also comprise one or more amino acid substitutions described herein, for example one or more substitutions that reduce effector function (e.g., as described in Section 6.9.1) and/or one or more substitutions that promote Fc heterodimerization (e.g., as described in Section 6.9.2).

[0172] In some aspects, an Fc domain comprises an amino acid sequence having at least about 90%, at least about 91%, at least about 92%, about at least 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% sequence identity to SEQ ID NO:4. In cases where an Fc domain comprises at least 90% sequence identity and less than 100% sequence identity to SEQ ID NO:4 (e.g., between 90% and 99% sequence identity to SEQ ID NO:4), an Fc domain may also comprise one or more amino acid substitutions described herein, for example one or more substitutions that reduce effector function (e.g., as described in Section 6.9.1) and/or one or more substitutions that promote Fc heterodimerization (e.g., as described in Section 6.9.2).

[0173] The two Fc domains within the Fc region can be the same or different from one another. In a native antibody the Fc domains are typically identical, but for the purpose of producing multispecific binding molecules, e.g., the IL2 proproteins of the disclosure and MBMs produced

by their activation, the Fc domains might advantageously be different to allow for heterodimerization, as described in Section 6.9.2 below.

[0174] In native antibodies, the heavy chain Fc domain of IgA, IgD and IgG is composed of two heavy chain constant domains (CH2 and CH3) and that of IgE and IgM is composed of three heavy chain constant domains (CH2, CH3 and CH4). These dimerize to create an Fc region.

[0175] In IL2 proproteins of the present disclosure, the Fc region, and / or the Fc domains within it, can comprise heavy chain constant domains from one or more different classes of antibody, for example one, two or three different classes.

[0176] In one embodiment the Fc region comprises CH2 and CH3 domains derived from IgG1.

10 [0177] In one embodiment the Fc region comprises CH2 and CH3 domains derived from IgG2.

[0178] In one embodiment the Fc region comprises CH2 and CH3 domains derived from IgG3.

[0179] In one embodiment the Fc region comprises CH2 and CH3 domains derived from IgG4.

[0180] In one embodiment the Fc region comprises a CH4 domain from IgM. The IgM CH4 domain is typically located at the C-terminus of the CH3 domain.

15 [0181] In one embodiment the Fc region comprises CH2 and CH3 domains derived from IgG and a CH4 domain derived from IgM.

[0182] It will be appreciated that the heavy chain constant domains for use in producing an Fc region for the IL2 proproteins of the present disclosure may include variants of the naturally occurring constant domains described above. Such variants may comprise one or more amino acid variations compared to wild type constant domains. In one example the Fc region of the present disclosure comprises at least one constant domain that varies in sequence from the wild-type constant domain. It will be appreciated that the variant constant domains may be longer or shorter than the wild-type constant domain. Preferably the variant constant domains are at least 60% identical or similar to a wild-type constant domain. In another example the variant constant domains are at least 70% identical or similar. In another example the variant constant domains are at least 80% identical or similar. In another example the variant constant domains are at least 90% identical or similar. In another example the variant constant domains are at least 95% identical or similar.

25 [0183] IgM and IgA occur naturally in humans as covalent multimers of the common H2L2 antibody unit. IgM occurs as a pentamer when it has incorporated a J-chain, or as a hexamer when it lacks a J-chain. IgA occurs as monomer and dimer forms. The heavy chains of IgM and

IgA possess an 18 amino acid extension to the C-terminal constant domain, known as a tailpiece. The tailpiece includes a cysteine residue that forms a disulfide bond between heavy chains in the polymer, and is believed to have an important role in polymerization. The tailpiece also contains a glycosylation site. In certain embodiments, the IL2 proproteins of the present disclosure do not comprise a tailpiece.

[0184] The Fc domains that are incorporated into the IL2 proproteins of the present disclosure may comprise one or more modifications that alter the functional properties of the proteins, for example, binding to Fc-receptors such as FcRn or leukocyte receptors, binding to complement, modified disulfide bond architecture, or altered glycosylation patterns. Exemplary Fc modifications that alter effector function are described in Section 6.9.1.

[0185] The Fc domains can also be altered to include modifications that improve manufacturability of asymmetric IL2 proproteins, for example by allowing heterodimerization, which is the preferential pairing of non-identical Fc domains over identical Fc domains. Heterodimerization permits the production of IL2 proproteins in which different polypeptide components are connected to one another by an Fc region containing Fc domains that differ in sequence. Examples of heterodimerization strategies are exemplified in Section 6.9.2.

[0186] It will be appreciated that any of the modifications mentioned above can be combined in any suitable manner to achieve the desired functional properties and/or combined with other modifications to alter the properties of the IL2 proproteins.

6.9.1. Fc Domains with Altered Effector Function

[0187] In some embodiments, the Fc domain comprises one or more amino acid substitutions that reduces binding to an Fc receptor and/or effector function.

[0188] In a particular embodiment the Fc receptor is an Fcγ receptor. In one embodiment the Fc receptor is a human Fc receptor. In one embodiment the Fc receptor is an activating Fc receptor. In a specific embodiment the Fc receptor is an activating human Fcγ receptor, more specifically human FcγRIIIa, FcγRI or FcγRIIa, most specifically human FcγRIIIa. In one embodiment the effector function is one or more selected from the group of complement dependent cytotoxicity (CDC), antibody-dependent cell-mediated cytotoxicity (ADCC), antibody-dependent cellular phagocytosis (ADCP), and cytokine secretion. In a particular embodiment, the effector function is ADCC.

[0189] In one embodiment, the Fc domain (e.g., an Fc domain of an IL2 proprotein half antibody) or the Fc region (e.g., one or both Fc domains of an IL2 proprotein that can associate

to form an Fc region) comprises an amino acid substitution at a position selected from the group of E233, L234, L235, N297, P331 and P329 (numberings according to Kabat EU index). In a more specific embodiment, the Fc domain or the Fc region comprises an amino acid substitution at a position selected from the group of L234, L235 and P329 (numberings according to Kabat EU index). In some embodiments, the Fc domain or the Fc region comprises the amino acid substitutions L234A and L235A (numberings according to Kabat EU index). In one such embodiment, the Fc domain or region is an IgG Fc domain or region, particularly a human IgG Fc domain or region. In one embodiment, the Fc domain or the Fc region comprises an amino acid substitution at position P329. In a more specific embodiment, the amino acid substitution is P329A or P329G, particularly P329G (numberings according to Kabat EU index). In one embodiment, the Fc domain or the Fc region comprises an amino acid substitution at position P329 and a further amino acid substitution at a position selected from E233, L234, L235, N297 and P331 (numberings according to Kabat EU index). In a more specific embodiment, the further amino acid substitution is E233P, L234A, L235A, L235E, N297A, N297D or P331S. In particular embodiments, the Fc domain or the Fc region comprises amino acid substitutions at positions P329, L234 and L235 (numberings according to Kabat EU index). In more particular embodiments, the Fc domain comprises the amino acid mutations L234A, L235A and P329G ("P329G LALA", "PGLALA" or "LALAPG").

[0190] Typically, the same one or more amino acid substitution is present in each of the two Fc domains of an Fc region. Thus, in a particular embodiment, each Fc domain of the Fc region comprises the amino acid substitutions L234A, L235A and P329G (Kabat EU index numbering), *i.e.* in each of the first and the second Fc domains in the Fc region the leucine residue at position 234 is replaced with an alanine residue (L234A), the leucine residue at position 235 is replaced with an alanine residue (L235A) and the proline residue at position 329 is replaced by a glycine residue (P329G) (numbering according to Kabat EU index).

[0191] In one embodiment, the Fc domain is an IgG1 Fc domain, particularly a human IgG1 Fc domain. In some embodiments, the IgG1 Fc domain is a variant IgG1 comprising D265A, N297A mutations (EU numbering) to reduce effector function.

[0192] In another embodiment, the Fc domain is an IgG4 Fc domain with reduced binding to Fc receptors. Exemplary IgG4 Fc domains with reduced binding to Fc receptors may comprise an amino acid sequence selected from Table H below: In some embodiments, the Fc domain includes only the bolded portion of the sequences shown below:

TABLE H	
Fc Domain	Sequence
SEQ ID NO:1 of WO2014/121087	Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys (SEQ ID NO: 358)
SEQ ID NO:2 of WO2014/121087	Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys (SEQ ID NO: 359)
SEQ ID NO:30 of WO2014/121087	Ala Ser Thr Lys Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala

TABLE H	
Fc Domain	Sequence
	Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys (SEQ ID NO: 360)
SEQ ID NO:31 of WO2014/121087	Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys (SEQ ID NO: 361)
SEQ ID NO:37 of WO2014/121087	Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro

TABLE H	
Fc Domain	Sequence
	Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn Arg Phe Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys (SEQ ID NO: 362)
SEQ ID NO:38 of WO2014/121087	Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn Arg Phe Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys (SEQ ID NO: 363)

[0193] In a particular embodiment, the IgG4 with reduced effector function comprises the bolded portion of the amino acid sequence of SEQ ID NO:31 of WO2014/121087, sometimes referred to herein as IgG4s or hIgG4s.

[0194] For heterodimeric Fc regions, it is possible to incorporate a combination of the variant IgG4 Fc sequences set forth above, for example an Fc region comprising an Fc domain comprising the amino acid sequence of SEQ ID NO:30 of WO2014/121087 (or the bolded portion thereof) and an Fc domain comprising the amino acid sequence of SEQ ID NO:37 of
5 WO2014/121087 (or the bolded portion thereof) or an Fc region comprising an Fc domain comprising the amino acid sequence of SEQ ID NO:31 of WO2014/121087 (or the bolded portion thereof) and an Fc domain comprising the amino acid sequence of SEQ ID NO:38 of WO2014/121087 (or the bolded portion thereof).

6.9.2. Fc Heterodimerization Variants

- 10 [0195] Certain IL2 proproteins entail dimerization between two Fc domains that, unlike a native immunoglobulin, are operably linked to non-identical N-terminal or C-terminal regions. Inadequate heterodimerization of two Fc domains to form an Fc region can be an obstacle for increasing the yield of desired heterodimeric molecules and represents challenges for purification. A variety of approaches available in the art can be used in for enhancing
15 dimerization of Fc domains that might be present in the IL2 proproteins of the disclosure, for example as disclosed in EP 1870459A1; U.S. Patent No. 5,582,996; U.S. Patent No. 5,731,168; U.S. Patent No. 5,910,573; U.S. Patent No. 5,932,448; U.S. Patent No. 6,833,441; U.S. Patent No. 7,183,076; U.S. Patent Application Publication No. 2006204493A1; and PCT Publication No. WO 2009/089004A1.
- 20 [0196] In some embodiments, the present disclosure provides IL2 proproteins comprising Fc heterodimers, *i.e.*, Fc regions comprising heterologous, non-identical Fc domains. Typically, each Fc domain in the Fc heterodimer comprises a CH3 domain of an antibody. The CH3 domains are derived from the constant region of an antibody of any isotype, class or subclass, and preferably of IgG (IgG1, IgG2, IgG3 and IgG4) class, as described in the preceding section.
- 25 [0197] Heterodimerization of the two different heavy chains at CH3 domains give rise to the desired IL2 proprotein, while homodimerization of identical heavy chains will reduce yield of the desired IL2 proprotein. Thus, in a preferred embodiment, the polypeptides that associate to form an IL2 proprotein of the disclosure will contain CH3 domains with modifications that favor heterodimeric association relative to unmodified Fc domains.
- 30 [0198] In a specific embodiment said modification promoting the formation of Fc heterodimers is a so-called “knob-into-hole” or “knob-in-hole” modification, comprising a “knob” modification in one of the Fc domains and a “hole” modification in the other Fc domain. The knob-into-hole technology is described *e.g.*, in U.S. Patent No. 5,731,168; US 7,695,936; Ridgway *et al.*, 1996,

Prot Eng 9:617-621, and Carter, 2001, Immunol Meth 248:7-15. Generally, the method involves introducing a protuberance ("knob") at the interface of a first polypeptide and a corresponding cavity ("hole") in the interface of a second polypeptide, such that the protuberance can be positioned in the cavity so as to promote heterodimer formation and hinder homodimer formation. Protuberances are constructed by replacing small amino acid side chains from the interface of the first polypeptide with larger side chains (e.g., tyrosine or tryptophan). Compensatory cavities of identical or similar size to the protuberances are created in the interface of the second polypeptide by replacing large amino acid side chains with smaller ones (e.g., alanine or threonine).

10 **[0199]** Accordingly, in some embodiments, an amino acid residue in the CH3 domain of the first subunit of the Fc domain is replaced with an amino acid residue having a larger side chain volume, thereby generating a protuberance within the CH3 domain of the first subunit which is positionable in a cavity within the CH3 domain of the second subunit, and an amino acid residue in the CH3 domain of the second subunit of the Fc domain is replaced with an amino acid residue having a smaller side chain volume, thereby generating a cavity within the CH3 domain of the second subunit within which the protuberance within the CH3 domain of the first subunit is positionable. Preferably said amino acid residue having a larger side chain volume is selected from the group consisting of arginine (R), phenylalanine (F), tyrosine (Y), and tryptophan (W). Preferably said amino acid residue having a smaller side chain volume is selected from the group consisting of alanine (A), serine (S), threonine (T), and valine (V). The protuberance and cavity can be made by altering the nucleic acid encoding the polypeptides, e.g., by site-specific mutagenesis, or by peptide synthesis. An exemplary substitution is Y470T.

25 **[0200]** In a specific such embodiment, in the first Fc domain the threonine residue at position 366 is replaced with a tryptophan residue (T366W), and in the Fc domain the tyrosine residue at position 407 is replaced with a valine residue (Y407V) and optionally the threonine residue at position 366 is replaced with a serine residue (T366S) and the leucine residue at position 368 is replaced with an alanine residue (L368A) (numbering according to Kabat EU index). In a further embodiment, in the first Fc domain additionally the serine residue at position 354 is replaced with a cysteine residue (S354C) or the glutamic acid residue at position 356 is replaced with a cysteine residue (E356C) (particularly the serine residue at position 354 is replaced with a cysteine residue), and in the second Fc domain additionally the tyrosine residue at position 349 is replaced by a cysteine residue (Y349C) (numbering according to Kabat EU index). In a particular embodiment, the first Fc domain comprises the amino acid substitutions S354C and

T366W, and the second Fc domain comprises the amino acid substitutions Y349C, T366S, L368A and Y407V (numbering according to Kabat EU index).

[0201] In some embodiments, electrostatic steering (e.g., as described in Gunasekaran *et al.*, 2010, J Biol Chem 285(25): 19637-46) can be used to promote the association of the first and
5 the second Fc domains of the Fc region.

[0202] As an alternative, or in addition, to the use of Fc domains that are modified to promote heterodimerization, an Fc domain can be modified to allow a purification strategy that enables selections of Fc heterodimers. In one such embodiment, one polypeptide comprises a modified Fc domain that abrogates its binding to Protein A, thus enabling a purification method that yields
10 a heterodimeric protein. See, for example, U.S. Patent No. 8,586,713. As such, the IL2 proproteins comprise a first CH3 domain and a second Ig CH3 domain, wherein the first and second Ig CH3 domains differ from one another by at least one amino acid, and wherein at least one amino acid difference reduces binding of the IL2 proprotein to Protein A as compared to a corresponding IL2 proprotein lacking the amino acid difference. In one embodiment, the first
15 CH3 domain binds Protein A and the second CH3 domain contains a mutation/modification that reduces or abolishes Protein A binding such as an H95R modification (by IMGT exon numbering; H435R by EU numbering). The second CH3 may further comprise a Y96F modification (by IMGT; Y436F by EU). This class of modifications is referred to herein as “star” mutations.

20 [0203] In some embodiments, the Fc can contain one or more mutations (e.g., knob and hole mutations) to facilitate heterodimerization as well as star mutations to facilitate purification.

6.9.3. Hinge Domains

[0204] The IL2 proproteins of the disclosure can comprise an Fc domain comprising a hinge domain at its N-terminus. The hinge region can be a native or a modified hinge region. Hinge
25 regions are typically found at the N-termini of Fc regions. The term “hinge domain”, unless the context dictates otherwise, refers to a naturally or non-naturally occurring hinge sequence that in the context of a single or monomeric polypeptide chain is a monomeric hinge domain and in the context of a dimeric polypeptide (e.g., a homodimeric or heterodimeric IL2 proprotein formed by the association of two Fc domains) can comprise two associated hinge sequences on
30 separate polypeptide chains. Sometimes, the two associated hinge sequences are referred to as a “hinge region”.

[0205] A native hinge region is the hinge region that would normally be found between Fab and Fc domains in a naturally occurring antibody. A modified hinge region is any hinge that differs in length and/or composition from the native hinge region. Such hinges can include hinge regions from other species, such as human, mouse, rat, rabbit, shark, pig, hamster, camel, llama or goat hinge regions. Other modified hinge regions may comprise a complete hinge region derived from an antibody of a different class or subclass from that of the heavy chain Fc domain or Fc region. Alternatively, the modified hinge region may comprise part of a natural hinge or a repeating unit in which each unit in the repeat is derived from a natural hinge region. In a further alternative, the natural hinge region may be altered by converting one or more cysteine or other residues into neutral residues, such as serine or alanine, or by converting suitably placed residues into cysteine residues. By such means the number of cysteine residues in the hinge region may be increased or decreased. Other modified hinge regions may be entirely synthetic and may be designed to possess desired properties such as length, cysteine composition and flexibility.

[0206] A number of modified hinge regions have already been described for example, in U.S. Patent No. 5,677,425, WO 99/15549, WO 2005/003170, WO 2005/003169, WO 2005/003170, WO 98/25971 and WO 2005/003171 and these are incorporated herein by reference.

[0207] In one embodiment, an IL2 proprotein of the disclosure comprises an Fc region in which one or both Fc domains possesses an intact hinge domain at its N-terminus.

[0208] In various embodiments, positions 233-236 within a hinge region may be G, G, G and unoccupied; G, G, unoccupied, and unoccupied; G, unoccupied, unoccupied, and unoccupied; or all unoccupied, with positions numbered by EU numbering.

[0209] In some embodiments, the IL2 proproteins of the disclosure comprise a modified hinge region that reduces binding affinity for an Fcγ receptor relative to a wild-type hinge region of the same isotype (e.g., human IgG1 or human IgG4).

[0210] In one embodiment, the IL2 proproteins of the disclosure comprise an Fc region in which each Fc domain possesses an intact hinge domain at its N-terminus, where each Fc domain and hinge domain is derived from IgG4 and each hinge domain comprises the modified sequence CPPC. The core hinge region of human IgG4 contains the sequence CPSC compared to IgG1 that contains the sequence CPPC. The serine residue present in the IgG4 sequence leads to increased flexibility in this region, and therefore a proportion of molecules form disulfide bonds within the same protein chain (an intrachain disulfide) rather than bridging

to the other heavy chain in the IgG molecule to form the interchain disulfide. (Angel *et al.*, 1993, Mol Immunol 30(1):105-108). Changing the serine residue to a proline to give the same core sequence as IgG1 allows complete formation of inter-chain disulfides in the IgG4 hinge region, thus reducing heterogeneity in the purified product. This altered isotype is termed IgG4P.

5 **6.9.3.1. Chimeric Hinge Sequences**

[0211] The hinge domain can be a chimeric hinge domain.

[0212] For example, a chimeric hinge may comprise an “upper hinge” sequence, derived from a human IgG1, a human IgG2 or a human IgG4 hinge region, combined with a “lower hinge” sequence, derived from a human IgG1, a human IgG2 or a human IgG4 hinge region.

- 10 [0213] In particular embodiments, a chimeric hinge region comprises the amino acid sequence EPKSCDKTHTCPPCPAPPVA (SEQ ID NO: 364) (previously disclosed as SEQ ID NO:8 of WO2014/121087, which is incorporated by reference in its entirety herein) or ESKYGPPCPPCPAPPVA (SEQ ID NO: 365) (previously disclosed as SEQ ID NO:9 of WO2014/121087). Such chimeric hinge sequences can be suitably linked to an IgG4 CH2
- 15 region (for example by incorporation into an IgG4 Fc domain, for example a human or murine Fc domain, which can be further modified in the CH2 and/or CH3 domain to reduce effector function, for example as described in Section 6.9.1).

6.9.3.2. Hinge Sequences with Reduced Effector Function

- [0214] In further embodiments, the hinge region can be modified to reduce effector function, for example as described in WO2016161010A2, which is incorporated by reference in its entirety
- 20 herein. In various embodiments, the positions 233-236 of the modified hinge region are G, G, G and unoccupied; G, G, unoccupied, and unoccupied; G, unoccupied, unoccupied, and unoccupied; or all unoccupied, with positions numbered by EU numbering (as shown in FIG. 1 of WO2016161010A2). These segments can be represented as GGG-, GG--, G--- or ---- with “-”
- 25 representing an unoccupied position.

[0215] Position 236 is unoccupied in canonical human IgG2 but is occupied by in other canonical human IgG isotypes. Positions 233-235 are occupied by residues other than G in all four human isotypes (as shown in FIG. 1 of WO2016161010A2).

- [0216] The hinge modification within positions 233-236 can be combined with position 228
- 30 being occupied by P. Position 228 is naturally occupied by P in human IgG1 and IgG2 but is occupied by S in human IgG4 and R in human IgG3. An S228P mutation in an IgG4 antibody is advantageous in stabilizing an IgG4 antibody and reducing exchange of heavy chain light chain

pairs between exogenous and endogenous antibodies. Preferably positions 226-229 are occupied by C, P, P and C respectively.

[0217] Exemplary hinge regions have residues 226-236, sometimes referred to as middle (or core) and lower hinge, occupied by the modified hinge sequences designated GGG-(233-236), GG--(233-236), G---(233-236) and no G(233-236). Optionally, the hinge domain amino acid sequence comprises CPPCPAPGGG-GPSVF (SEQ ID NO: 366) (previously disclosed as SEQ ID NO:1 of WO2016161010A2), CPPCPAPGG--GPSVF (SEQ ID NO: 367) (previously disclosed as SEQ ID NO:2 of WO2016161010A2), CPPCPAPG---GPSVF (SEQ ID NO: 368) (previously disclosed as SEQ ID NO:3 of WO2016161010A2), or CPPCPAP----GPSVF (SEQ ID NO: 369) (previously disclosed as SEQ ID NO:4 of WO2016161010A2).

[0218] The modified hinge regions described above can be incorporated into a heavy chain constant region, which typically include CH2 and CH3 domains, and which may have an additional hinge segment (e.g., an upper hinge) flanking the designated region. Such additional constant region segments present are typically of the same isotype, preferably a human isotype, although can be hybrids of different isotypes. The isotype of such additional human constant regions segments is preferably human IgG4 but can also be human IgG1, IgG2, or IgG3 or hybrids thereof in which domains are of different isotypes. Exemplary sequences of human IgG1, IgG2 and IgG4 are shown in FIGS. 2-4 of WO2016161010A2.

[0219] In specific embodiments, the modified hinge sequences can be linked to an IgG4 CH2 region (for example by incorporation into an IgG4 Fc domain, for example a human or murine Fc domain, which can be further modified in the CH2 and/or CH3 domain to reduce effector function, for example as described in Section 6.9.1).

6.10. Nucleic Acids and Host Cells

[0220] In another aspect, the disclosure provides nucleic acids encoding the IL2 proproteins of the disclosure. In some embodiments, the IL2 proproteins are encoded by a single nucleic acid. In other embodiments, the IL2 proproteins can be encoded by a plurality (e.g., two, three, four or more) nucleic acids.

[0221] A single nucleic acid can encode an IL2 proprotein that comprises a single polypeptide chain, an IL2 proprotein that comprises two or more polypeptide chains, or a portion of an IL2 proprotein that comprises more than two polypeptide chains (for example, a single nucleic acid can encode two polypeptide chains of an IL2 proprotein comprising three, four or more polypeptide chains, or three polypeptide chains of an IL2 proprotein comprising four or more

polypeptide chains). For separate control of expression, the open reading frames encoding two or more polypeptide chains can be under the control of separate transcriptional regulatory elements (*e.g.*, promoters and/or enhancers). The open reading frames encoding two or more polypeptides can also be controlled by the same transcriptional regulatory elements, and
5 separated by internal ribosome entry site (IRES) sequences allowing for translation into separate polypeptides.

[0222] In some embodiments, an IL2 proprotein comprising two or more polypeptide chains is encoded by two or more nucleic acids. The number of nucleic acids encoding an IL2 proprotein can be equal to or less than the number of polypeptide chains in the IL2 proprotein (for
10 example, when more than one polypeptide chains are encoded by a single nucleic acid).

[0223] The nucleic acids of the disclosure can be DNA or RNA (*e.g.*, mRNA).

[0224] In another aspect, the disclosure provides host cells and vectors containing the nucleic acids of the disclosure. The nucleic acids may be present in a single vector or separate vectors present in the same host cell or separate host cell, as described in more detail herein below.

15 6.10.1. Vectors

[0225] The disclosure provides vectors comprising nucleotide sequences encoding an IL2 proprotein or a component thereof described herein, for example one or two of the polypeptide chains of a half antibody of an IL2 proprotein. The vectors include, but are not limited to, a virus, plasmid, cosmid, lambda phage or a yeast artificial chromosome (YAC).

20 [0226] Numerous vector systems can be employed. For example, one class of vectors utilizes DNA elements which are derived from animal viruses such as, for example, bovine papilloma virus, polyoma virus, adenovirus, vaccinia virus, baculovirus, retroviruses (Rous Sarcoma Virus, MMTV or MOMLV) or SV40 virus. Another class of vectors utilizes RNA elements derived from RNA viruses such as Semliki Forest virus, Eastern Equine Encephalitis virus and Flaviviruses.

25 [0227] Additionally, cells which have stably integrated the DNA into their chromosomes can be selected by introducing one or more markers which allow for the selection of transfected host cells. The marker may provide, for example, prototrophy to an auxotrophic host, biocide resistance (*e.g.*, antibiotics), or resistance to heavy metals such as copper, or the like. The selectable marker gene can be either directly linked to the DNA sequences to be expressed, or
30 introduced into the same cell by co-transformation. Additional elements may also be needed for optimal synthesis of mRNA. These elements may include splice signals, as well as transcriptional promoters, enhancers, and termination signals.

[0228] Once the expression vector or DNA sequence containing the constructs has been prepared for expression, the expression vectors can be transfected or introduced into an appropriate host cell. Various techniques may be employed to achieve this, such as, for example, protoplast fusion, calcium phosphate precipitation, electroporation, retroviral transduction, viral transfection, gene gun, lipid-based transfection or other conventional techniques. Methods and conditions for culturing the resulting transfected cells and for recovering the expressed polypeptides are known to those skilled in the art, and may be varied or optimized depending upon the specific expression vector and mammalian host cell employed, based upon the present description.

10 6.10.2. Cells

[0229] The disclosure also provides host cells comprising a nucleic acid of the disclosure.

[0230] In one embodiment, the host cells are genetically engineered to comprise one or more nucleic acids described herein.

15 [0231] In one embodiment, the host cells are genetically engineered by using an expression cassette. The phrase "expression cassette," refers to nucleotide sequences, which are capable of affecting expression of a gene in hosts compatible with such sequences. Such cassettes may include a promoter, an open reading frame with or without introns, and a termination signal. Additional factors necessary or helpful in effecting expression may also be used, such as, for example, an inducible promoter.

20 [0232] The disclosure also provides host cells comprising the vectors described herein.

[0233] The cell can be, but is not limited to, a eukaryotic cell, a bacterial cell, an insect cell, or a human cell. Suitable eukaryotic cells include, but are not limited to, Vero cells, HeLa cells, COS cells, CHO cells, HEK293 cells, BHK cells and MDCKII cells. Suitable insect cells include, but are not limited to, Sf9 cells.

25 6.11. Pharmaceutical Compositions

[0234] The IL2 proproteins of the disclosure may be in the form of compositions comprising the IL2 proprotein and one or more carriers, excipients and/or diluents. The compositions may be formulated for specific uses, such as for veterinary uses or pharmaceutical uses in humans. The form of the composition (*e.g.*, dry powder, liquid formulation, *etc.*) and the excipients, diluents and/or carriers used will depend upon the intended uses of the IL2 proprotein and, for therapeutic uses, the mode of administration.

30

- [0235] For therapeutic uses, the compositions may be supplied as part of a sterile, pharmaceutical composition that includes a pharmaceutically acceptable carrier. This composition can be in any suitable form (depending upon the desired method of administering it to a patient). The pharmaceutical composition can be administered to a patient by a variety of routes such as orally, transdermally, subcutaneously, intranasally, intravenously, intramuscularly, intratumorally, intrathecally, topically or locally. The most suitable route for administration in any given case will depend on the particular IL2 proprotein, the subject, and the nature and severity of the disease and the physical condition of the subject. Typically, the pharmaceutical composition will be administered intravenously or subcutaneously.
- [0236] Pharmaceutical compositions can be conveniently presented in unit dosage forms containing a predetermined amount of an IL2 proprotein of the disclosure per dose. The quantity of IL2 proprotein included in a unit dose will depend on the disease being treated, as well as other factors as are well known in the art. Such unit dosages may be in the form of a lyophilized dry powder containing an amount of IL2 proprotein suitable for a single administration, or in the form of a liquid. Dry powder unit dosage forms may be packaged in a kit with a syringe, a suitable quantity of diluent and/or other components useful for administration. Unit dosages in liquid form may be conveniently supplied in the form of a syringe pre-filled with a quantity of IL2 proprotein suitable for a single administration.
- [0237] The pharmaceutical compositions may also be supplied in bulk from containing quantities of IL2 proprotein suitable for multiple administrations.
- [0238] Pharmaceutical compositions may be prepared for storage as lyophilized formulations or aqueous solutions by mixing an IL2 proprotein having the desired degree of purity with optional pharmaceutically-acceptable carriers, excipients or stabilizers typically employed in the art (all of which are referred to herein as "carriers"), *i.e.*, buffering agents, stabilizing agents, preservatives, isotonifiers, non-ionic detergents, antioxidants, and other miscellaneous additives. See, Remington's Pharmaceutical Sciences, 16th edition (Osol, ed. 1980). Such additives should be nontoxic to the recipients at the dosages and concentrations employed.
- [0239] Buffering agents help to maintain the pH in the range which approximates physiological conditions. They may be present at a wide variety of concentrations, but will typically be present in concentrations ranging from about 2 mM to about 50 mM. Suitable buffering agents for use with the present disclosure include both organic and inorganic acids and salts thereof such as citrate buffers (*e.g.*, monosodium citrate-disodium citrate mixture, citric acid-trisodium citrate mixture, citric acid-monosodium citrate mixture, *etc.*), succinate buffers (*e.g.*, succinic acid-

monosodium succinate mixture, succinic acid-sodium hydroxide mixture, succinic acid-disodium succinate mixture, *etc.*), tartrate buffers (*e.g.*, tartaric acid-sodium tartrate mixture, tartaric acid-potassium tartrate mixture, tartaric acid-sodium hydroxide mixture, *etc.*), fumarate buffers (*e.g.*, fumaric acid-monosodium fumarate mixture, fumaric acid-disodium fumarate mixture, 5 monosodium fumarate-disodium fumarate mixture, *etc.*), gluconate buffers (*e.g.*, gluconic acid-sodium glyconate mixture, gluconic acid-sodium hydroxide mixture, gluconic acid-potassium glyconate mixture, *etc.*), oxalate buffer (*e.g.*, oxalic acid-sodium oxalate mixture, oxalic acid-sodium hydroxide mixture, oxalic acid-potassium oxalate mixture, *etc.*), lactate buffers (*e.g.*, lactic acid-sodium lactate mixture, lactic acid-sodium hydroxide mixture, lactic acid-potassium lactate mixture, *etc.*) and acetate buffers (*e.g.*, acetic acid-sodium acetate mixture, acetic acid-sodium hydroxide mixture, *etc.*). Additionally, phosphate buffers, histidine buffers and trimethylamine salts such as Tris can be used.

[0240] Preservatives may be added to retard microbial growth, and can be added in amounts ranging from about 0.2%-1 % (w/v). Suitable preservatives for use with the present disclosure 15 include phenol, benzyl alcohol, meta-cresol, methyl paraben, propyl paraben, octadecyldimethylbenzyl ammonium chloride, benzalconium halides (*e.g.*, chloride, bromide, and iodide), hexamethonium chloride, and alkyl parabens such as methyl or propyl paraben, catechol, resorcinol, cyclohexanol, and 3-pentanol. Isotonicifiers sometimes known as “stabilizers” can be added to ensure isotonicity of liquid compositions of the present disclosure 20 and include polyhydric sugar alcohols, for example trihydric or higher sugar alcohols, such as glycerin, erythritol, arabitol, xylitol, sorbitol and mannitol. Stabilizers refer to a broad category of excipients which can range in function from a bulking agent to an additive which solubilizes the therapeutic agent or helps to prevent denaturation or adherence to the container wall. Typical stabilizers can be polyhydric sugar alcohols (enumerated above); amino acids such as arginine, 25 lysine, glycine, glutamine, asparagine, histidine, alanine, ornithine, L-leucine, 2-phenylalanine, glutamic acid, threonine, *etc.*, organic sugars or sugar alcohols, such as lactose, trehalose, stachyose, mannitol, sorbitol, xylitol, ribitol, myoinisitol, galactitol, glycerol and the like, including cyclitols such as inositol; polyethylene glycol; amino acid polymers; sulfur containing reducing agents, such as urea, glutathione, thiocetic acid, sodium thioglycolate, thioglycerol, a- 30 monothioglycerol and sodium thio sulfate; low molecular weight polypeptides (*e.g.*, peptides of 10 residues or fewer); proteins such as human serum albumin, bovine serum albumin, gelatin or immunoglobulins; hydrophylic polymers, such as polyvinylpyrrolidone monosaccharides, such as xylose, mannose, fructose, glucose; disaccharides such as lactose, maltose, sucrose and

trehalose; and trisaccharides such as raffinose; and polysaccharides such as dextran.

Stabilizers may be present in amounts ranging from 0.5 to 10 wt % per wt of IL2 proprotein.

[0241] Non-ionic surfactants or detergents (also known as “wetting agents”) may be added to help solubilize the glycoprotein as well as to protect the glycoprotein against agitation-induced aggregation, which also permits the formulation to be exposed to shear surface stressed without causing denaturation of the protein. Suitable non-ionic surfactants include polysorbates (20, 80, etc.), polyoxamers (184, 188, etc.), and pluronic polyols. Non-ionic surfactants may be present in a range of about 0.05 mg/mL to about 1.0 mg/mL, for example about 0.07 mg/mL to about 0.2 mg/mL.

[0242] Additional miscellaneous excipients include bulking agents (e.g., starch), chelating agents (e.g., EDTA), antioxidants (e.g., ascorbic acid, methionine, vitamin E), and cosolvents.

[0243] The IL2 proproteins of the disclosure can be formulated as pharmaceutical compositions comprising the IL2 proproteins, for example containing one or more pharmaceutically acceptable excipients or carriers. To prepare pharmaceutical or sterile compositions comprising the IL2 proproteins of the present disclosure, an IL2 proprotein preparation can be combined with one or more pharmaceutically acceptable excipient or carrier.

[0244] For example, formulations of IL2 proproteins can be prepared by mixing IL2 proproteins with physiologically acceptable carriers, excipients, or stabilizers in the form of, e.g., lyophilized powders, slurries, aqueous solutions, lotions, or suspensions (see, e.g., Hardman *et al.*, 2001, Goodman and Gilman’s The Pharmacological Basis of Therapeutics, McGraw-Hill, New York, N.Y.; Gennaro, 2000, Remington: The Science and Practice of Pharmacy, Lippincott, Williams, and Wilkins, New York, N.Y.; Avis, *et al.* (eds.), 1993, Pharmaceutical Dosage Forms: General Medications, Marcel Dekker, NY; Lieberman, *et al.* (eds.), 1990, Pharmaceutical Dosage Forms: Tablets, Marcel Dekker, NY; Lieberman, *et al.* (eds.), 1990, Pharmaceutical Dosage Forms: Disperse Systems, Marcel Dekker, NY; Weiner and Kotkoskie, 2000, Excipient Toxicity and Safety, Marcel Dekker, Inc., New York, N.Y.).

[0245] An effective amount for a particular subject may vary depending on factors such as the condition being treated, the overall health of the subject, the method route and dose of administration and the severity of side effects (see, e.g., Maynard, *et al.* (1996) A Handbook of SOPs for Good Clinical Practice, Interpharm Press, Boca Raton, Fla.; Dent (2001) Good Laboratory and Good Clinical Practice, Urch Publ., London, UK).

[0246] A composition of the present disclosure may also be administered via one or more routes of administration using one or more of a variety of methods known in the art. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. Selected routes of administration for IL2 proproteins include

5 intravenous, intramuscular, intradermal, intraperitoneal, subcutaneous, spinal or other general routes of administration, for example by injection or infusion. General administration may represent modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous,

10 subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal, epidural and intrasternal injection and infusion. Alternatively, a composition of the disclosure can be administered via a non-general route, such as a topical, epidermal or mucosal route of administration, for example, intranasally, orally, vaginally, rectally, sublingually or topically. In one embodiment, the IL2 proproteins are administered by infusion. In another embodiment, the IL2 proprotein of the

15 disclosure is administered subcutaneously.

6.12. Therapeutic Indications and Methods of Use

[0247] The present disclosure provides methods for using and applications for the IL2 proproteins of the disclosure.

[0248] In certain aspects, the disclosure provides a method of treating cancer, comprising

20 administering to a subject in need thereof an IL2 proprotein or pharmaceutical composition as described herein. In some embodiments, an activated IL2 protein comprising the IL2 moiety is produced by cleavage of one or more protease-cleavable linkers in the IL2 proprotein by one or more proteases expressed by the cancer tissue. Accordingly, the IL2 proprotein is selectively activated in the cancer tissue.

[0249] In some embodiments, the disclosure provides a method of treating cancer with an IL2 protein that is selectively activated in cancer tissue, comprising administering to a subject in need thereof an IL2 proprotein or pharmaceutical composition as described herein, where the IL2 proprotein has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by cancer tissue to which the IL2 protein is

25 intended. Thus, an activated IL2 protein comprising the IL2 moiety is produced by cleavage of

30 one or more protease-cleavable linkers in the IL2 proprotein by one or more proteases in the cancer tissue.

[0250] The present disclosure further provides a method of localized delivery of an IL2 protein, comprising administering to a subject an IL2 proprotein or pharmaceutical composition as described herein, where the IL2 proprotein has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by a tissue to which the IL2 protein is to be locally delivered. As used herein, the term "locally delivered" does not require local administration but rather indicates that the active component of the IL2 proprotein refers to activation of the protein at a locale of interest by a protease active at the intended site, optionally in conjunction with targeting to the locale of interest with a targeting moiety that recognize a target molecule expressed by the tissue.

10 [0251] The present disclosure further provides a method of administering to the subject IL2 therapy with reduced systemic exposure and/or reduced systemic toxicity, comprising administering to a subject the IL2 therapy in the form of an IL2 proprotein or pharmaceutical composition as described herein, where the IL2 proprotein has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by a tissue for which IL2 therapy is desirable and/or intended.

[0252] Accordingly, the foregoing methods permit IL2 therapy with reduced off-target side effects by virtue of preferential activation of an IL2 proprotein at a locale intended for IL2 treatment.

20 [0253] In some embodiments of the foregoing methods, the IL2 proprotein is also targeted and comprises one or more targeting moieties that recognize a target molecule expressed in the locale (e.g., by the tissue) intended for treatment.

[0254] Accordingly, the present disclosure provides a method of targeted delivery of an activated IL2 protein to a locale intended for treatment, e.g., cancer tissue, comprising administering to a subject an IL2 proprotein or pharmaceutical composition as described herein, wherein the IL2 comprises one or more targeting moieties that recognize a target molecule expressed in the locale or by the tissue intended for treatment (e.g., cancer tissue) and which has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by a tissue for which IL2 therapy is desirable and/or intended.

30 [0255] The present disclosure further provides method of locally inducing an immune response in a target tissue, comprising administering to a subject IL2 proprotein or pharmaceutical composition as described herein which has one or more targeting moieties capable of binding a target molecule expressed in the target tissue and one or more protease-cleavable linkers, each

comprising one or more substrates for one or more proteases expressed in the target tissue. An activated IL2 protein comprising the IL2 moiety can then be produced by cleavage of one or more protease-cleavable linkers in the IL2 proprotein by one or more proteases in the target tissue. The resulting activated IL2 protein can then induce the immune response against at least one cell type in the target tissue.

[0256] In some embodiments, the administration is not local to the tissue. For example, when the target tissue is cancer tissue, the administration can be systemic or subcutaneous.

[0257] The IL2 proproteins of the disclosure can be used in the treatment of any proliferative disorder (e.g., cancer) that expresses a target molecule (either on the tumor cells or in the tumor microenvironment, e.g., the extracellular matrix or the tumor lymphocytes). In particular embodiments, the cancer is acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), adrenocortical carcinoma, anal cancer, appendix cancer, astrocytoma, basal cell carcinoma, brain tumor, bile duct cancer, bladder cancer, bone cancer, breast cancer, bronchial tumor, Burkitt Lymphoma, carcinoma of unknown primary origin, cardiac tumor, cervical cancer, chordoma, chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), chronic myeloproliferative neoplasm, colon cancer, colorectal cancer, craniopharyngioma, cutaneous T-cell lymphoma, ductal carcinoma, embryonal tumor, endometrial cancer, ependymoma, esophageal cancer, esthesioneuroblastoma, fibrous histiocytoma, Ewing sarcoma, eye cancer, germ cell tumor, gallbladder cancer, gastric cancer, gastrointestinal carcinoid tumor, gastrointestinal stromal tumor, gestational trophoblastic disease, glioma, head and neck cancer, hairy cell leukemia, hepatocellular cancer, histiocytosis, Hodgkin lymphoma, hypopharyngeal cancer, intraocular melanoma, islet cell tumor, Kaposi sarcoma, kidney cancer, Langerhans cell histiocytosis, laryngeal cancer, leukemia, lip and oral cavity cancer, liver cancer, lobular carcinoma in situ, lung cancer, lymphoma, macroglobulinemia, malignant fibrous histiocytoma, melanoma, Merkel cell carcinoma, mesothelioma, metastatic squamous neck cancer with occult primary, midline tract carcinoma involving NUT gene, mouth cancer, multiple endocrine neoplasia syndrome, multiple myeloma, mycosis fungoides, myelodysplastic syndrome, myelodysplastic/myeloproliferative neoplasm, nasal cavity and para-nasal sinus cancer, nasopharyngeal cancer, neuroblastoma, non-Hodgkin lymphoma, non-small cell lung cancer, oropharyngeal cancer, osteosarcoma, ovarian cancer, pancreatic cancer, papillomatosis, paraganglioma, parathyroid cancer, penile cancer, pharyngeal cancer, pheochromocytomas, pituitary tumor, pleuropulmonary blastoma, primary central nervous system lymphoma, prostate cancer, rectal cancer, renal cell cancer, renal pelvis and ureter cancer, retinoblastoma, rhabdoid

tumor, salivary gland cancer, Sezary syndrome, skin cancer, small cell lung cancer, small intestine cancer, soft tissue sarcoma, spinal cord tumor, stomach cancer, T-cell lymphoma, teratoid tumor, testicular cancer, throat cancer, thymoma and thymic carcinoma, thyroid cancer, urethral cancer, uterine cancer, vaginal cancer, vulvar cancer, or Wilms tumor.

- 5 **[0258]** Table I below shows exemplary indications for which IL2 proproteins targeting particular target molecules can be used.

TABLE I	
Examples of Target Molecule Indications	
Target	Exemplary Indication(s)
ADRB3	Ewing sarcoma
ALK	NSCLC, ALCL, IMT, neuroblastoma
B7H3	melanoma, osteosarcoma, leukemia, breast, prostate, ovarian, pancreatic, colorectal cancers
BCMA	multiple myeloma, leukemia (e.g., acute lymphoblastic leukemia ("ALL"), acute myeloid leukemia ("AML"), chronic lymphocytic leukemia ("CLL"), chronic myeloid leukemia ("CML") and hairy cell leukemia ("HCL")); lymphoma (e.g., Hodgkin's lymphoma, non-Hodgkin's lymphoma, including diffuse large B-cell lymphoma ("DLBCL"))
Cadherin 17	gastric, pancreatic, and colorectal adenocarcinomas
CAIX	clear-cell renal cell carcinoma, hypoxic solid tumors, head and neck squamous carcinoma
CD123	leukemia (e.g., ALL, CLL, AML, CML, HCL); lymphoma (e.g., Hodgkin's lymphoma, non-Hodgkin's lymphoma, e.g., DLBCL); multiple myeloma. In a preferred embodiment, the indication is AML.
CD171	neuroblastoma, paraganglioma
CD179a	B cell malignancies
CD19	leukemia (e.g., ALL, CLL, AML, CML, HCL); lymphoma (e.g., Hodgkin's lymphoma, non-Hodgkin's lymphoma, e.g., DLBCL); multiple myeloma.
CD20	leukemia (e.g., ALL, CLL, AML, CML, HCL); lymphoma (e.g., Hodgkin's lymphoma, non-Hodgkin's lymphoma, e.g., DLBCL); multiple myeloma.
CD22	leukemia (e.g., ALL, CLL, AML, CML, HCL); lymphoma (e.g., Hodgkin's lymphoma, non-Hodgkin's lymphoma, e.g., DLBCL); multiple myeloma; lung cancer
CD24	ovarian, breast, prostate, bladder, renal, non-small cell carcinomas
CD30	anaplastic large cell lymphoma, embryonal carcinoma, Hodgkin Lymphoma
CD32b	B cell malignancies, gastric, pancreatic, esophageal, glioblastoma, breast, colorectal
CD33	leukemia (e.g., ALL, CLL, AML, CML, HCL); lymphoma (e.g., Hodgkin's lymphoma, non-Hodgkin's lymphoma, e.g., DLBCL); multiple myeloma. In a preferred embodiment, the indication is AML.
CD38	leukemia (e.g., ALL, CLL, AML, CML, HCL); lymphoma (e.g., Hodgkin's lymphoma, non-Hodgkin's lymphoma, e.g., DLBCL); multiple myeloma
CD44v6	colon cancer, head and neck small cell carcinoma
CD97	B cell malignancies, gastric, pancreatic, esophageal, glioblastoma, breast, colorectal

TABLE I Examples of Target Molecule Indications	
Target	Exemplary Indication(s)
CEA (CEACAM5)	colorectal carcinoma, gastric carcinoma, pancreatic carcinoma, lung cancer, breast cancer, medullary thyroid carcinoma
CLDN6	ovarian, breast, lung cancer
CLL-1	leukemia (e.g., ALL, CLL, AML, CML, HCL); lymphoma (e.g., Hodgkin's lymphoma, non-Hodgkin's lymphoma, e.g., DLBCL); multiple myeloma. In a preferred embodiment, the indication is AML.
CS1 (SLAMF7)	multiple myeloma
EGFR	squamous cell carcinoma of lung, anal cancer, glioblastoma, epithelial tumors of head and neck, colon cancer
EGFRvIII	Glioblastoma
EPCAM	gastrointestinal carcinoma, colorectal cancer
EphA2	kaposi's sarcoma, glioblastoma, solid tumors, glioma
Ephrin B2	thyroid cancer, breast cancer, malignant melanoma
ERBB2 (Her2/neu)	breast, ovarian, gastric cancers, lung adenocarcinoma, non-small cell lung cancer, uterine cancer, uterine serous endometrial carcinoma, salivary duct carcinoma
FAP	pancreatic cancer, colorectal cancer, metastasis, epithelial cancers, soft tissue sarcomas
FCRL5	multiple myeloma
FLT3	leukemia (e.g., ALL, CLL, AML, CML, HCL), lymphoma (e.g., Hodgkin's lymphoma, non-Hodgkin's lymphoma, e.g., DLBCL), multiple myeloma
Folate receptor alpha	ovarian, breast, renal, lung, colorectal, brain cancers
Folate receptor beta	ovarian cancer
Fucosyl GM1	AML, myeloma
GD2	malignant melanoma, neuroblastoma
GD3	Melanoma
GloboH	ovarian, gastric, prostate, lung, breast, and pancreatic cancers
gp100	Melanoma
GPNCB	breast cancer, head and neck cancers
GPR20	GIST
GPR64	Ewing sarcoma, prostate, kidney and lung sarcomas
GPRC5D	multiple myeloma
HAVCR1	renal cancer
HER2	HER-2 (+) adenocarcinoma of gastroesophageal junction, HER-2 positive gastric adenocarcinoma, HER2 positive carcinoma of breast
HER3	colon and gastric cancers
HMWMAA	melanoma, glioblastoma, breast cancer
IGF-I receptor	breast, prostate, lung cancers
IL11R α	papillary thyroid cancer, osteosarcoma, colorectal adenocarcinoma, lymphocytic leukemia
IL13R α 2	renal cell carcinoma, prostate cancer, gliomas, head and neck cancer, astrocytoma
KIT	myeloid leukemia, kaposi's sarcoma, erythroleukemia, gastrointestinal stromal tumors

TABLE I	
Examples of Target Molecule Indications	
Target	Exemplary Indication(s)
KLRG2	breast cancers, lung cancers and ovarian cancers.
LewisY	squamous cell lung carcinoma, lung adenocarcinoma, ovarian carcinoma, and colorectal adenocarcinoma
LMP2	prostate cancer, Hodgkin's lymphoma, nasopharyngeal carcinoma
LRP6	breast cancer
LY6K	breast, lung, ovarian, and cervical cancer
LYPD8	colorectal and gastric cancers
Mesothelin	mesothelioma, pancreatic cancer, ovarian cancer, stomach cancer, lung cancer, endometrial cancer
MUC1	breast and ovarian cancers, lung, stomach, pancreatic, prostate cancers
NCAM	melanoma, Wilms' tumor, small cell lung cancer, neuroblastoma, myeloma, paraganglioma, pancreatic acinar cell carcinoma, myeloid leukemia
NY-BR-1	breast cancer
o-acetyl GD2	neuroblastoma, melanoma
OR51E2	prostate cancer
PANX3	Osteosarcoma
PLAC1	hepatocellular carcinoma
Polysialic acid	small cell lung cancer
PDGFR-beta	myelomonocytic leukemia, chronic myeloid leukemia, acute myelogenous leukemia, acute lymphoblastic leukemia
PRSS21	colon cancer, testicular cancer, ovarian cancer
PSCA	prostate cancer, gastric and bladder cancers
PSMA	prostate cancer
ROR1	metastatic cancers, chronic lymphocytic leukemia, solid tumors in lung, breast, ovarian, colon, pancreatic, sarcoma
SLC34A2	bladder cancer
SLC39A6	breast cancer, esophageal cancer
SLITRK6	breast cancer, urothelial cancer, lung cancer
SSEA-4	breast cancer, cancer stem cells, epithelial ovarian carcinoma
STEAP1	prostate cancer
STEAP2	prostate cancer (including castrate-resistant prostate cancer), bladder cancer, cervical cancer, lung cancer, colon cancer, kidney cancer, breast cancer, pancreatic cancer, stomach cancer, uterine cancer, ovarian cancer, preferably prostate cancer
TACSTD2	carcinomas, <i>e.g.</i> , non-small-cell lung cancer
TAG72	ovarian, breast, colon, lung, pancreatic cancers, gastric cancer
TEM1/CD248	colorectal cancer
TEM7R	colorectal cancer
Tn	colorectal, breast cancers, cervical, lung, stomach cancers
TSHR	thyroid cancer, multiple myeloma
Tyrosinase	prostate cancer, melanoma
UPK2	bladder cancer
VEGFR2	ovarian and pancreatic cancers, renal cell carcinoma, colorectal cancer, medullary thyroid carcinoma

[0259] Additional target molecules and corresponding indications are disclosed in, *e.g.*, Hafeez *et al.*, 2020, Molecules 25:4764, doi:10.3390/molecules25204764, particularly in Table 1. Table 1 is incorporated by reference in its entirety here.

7. SEQUENCES

- 5 **[0260]** Sequences of certain IL2 proproteins of the present disclosure are provided in Table S, below.

TABLE S Example IL2 Proprotein Component Sequences		
Component	Sequence	SEQ ID NO
hlgG1 Fc (amino acids 99-330 of UniprotKB P01857-1)	EPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK	1
hlgG2 Fc (amino acids 99-326 of UniprotKB P01859-1)	ERKCCVECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK	2
hlgG3 Fc (amino acids 99-377 of UniprotKB P01860-1)	ELKTPLGDTTHTCPRCPEPKSCDTPPPCPRCPEPKSCDTPPPCPRCPEPKSCDTPPCPRCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFKWYVDGVEVHNAKTKPREEQYNSTFRVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESSGQPENNYNTTPMLDSDGSFFLYSKLTVDKSRWQQGNIFSCSMHEALHNRTQKSLSLSPGK	3
hlgG4 Fc (amino acids 99-327 of UniprotKB P01861-1)	ESKYGPPCPSCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSDQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSMHEALHNHYTQKSLSLGLGK	4
hlgG4s Fc	ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSDQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSMHEALHNHYTQKSLSLGLGK	5
hlgG1 PVA Fc	EPKSCDKTHTCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK	6
hIL2	APTSSSTKKTQLQLEHLLLDLQMI LINGINNYKNPKLTRMLTFKFYMPKKATELKH LQCLEEELKPLEEVLNLAQSKNFHLRPDLISNINVIVLELKGSETTFMCEYADE TATIVEFLNRWITFCQSIISTLT	7
hIL2Ra extracellular domain	ELCDDDPPEI PHATFKAMAYKEGTM LNCECKRGFRRIKSGSLYMLCTGNSSHSW DNQCQCTSSATRNTTKQVTPQPEEQKERKTTEMQSPMQPVDQASLPGHCREPPPW ENEATERIYHFVVGQMVYYQCVQGYRALHRGPAESVCKMTHGKTRWTQPQLICTG EMETSQFPGEKPKQASPEGREPESETSCLVTTTDFQIQTEMAATMETSIFTTEYQV AVAGCVFLLISVLLSGLTWQRRQRKSRRTI	8
Amino acids 22-186 of hIL2Ra	ELCDDDPPEI PHATFKAMAYKEGTM LNCECKRGFRRIKSGSLYMLCTGNSSHSW DNQCQCTSSATRNTTKQVTPQPEEQKERKTTEMQSPMQPVDQASLPGHCREPPPW ENEATERIYHFVVGQMVYYQCVQGYRALHRGPAESVCKMTHGKTRWTQPQLICTG	9
Amino acids 22-240 of hIL2Ra	ELCDDDPPEI PHATFKAMAYKEGTM LNCECKRGFRRIKSGSLYMLCTGNSSHSW DNQCQCTSSATRNTTKQVTPQPEEQKERKTTEMQSPMQPVDQASLPGHCREPPPW ENEATERIYHFVVGQMVYYQCVQGYRALHRGPAESVCKMTHGKTRWTQPQLICTG EMETSQFPGEKPKQASPEGREPESETSCLVTTTDFQIQTEMAATMETSIFTTEYQ	10
hlgG1 CH1 (amino acids 1-98 of UniprotKB P01857-1)	ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPA VLQSSGLYSLSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKV	11

hIgG2 CH1 (amino acids 1-98 of UniprotKB P01859-1)	ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPA VLQSSGLYSLSSVVTVPSSNFGTQTYTCNVDHKPSNTKVDKTV	12
hIgG3 CH1 (amino acids 1-98 of UniprotKB P01860-1)	ASTKGPSVFPLAPCSRSTSGGTAAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPA VLQSSGLYSLSSVVTVPSSSLGTQTYTCNVNHKPSNTKVDKRV	13
hIgG4 CH1 (amino acids 1-98 of UniprotKB P01861-1)	ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPA VLQSSGLYSLSSVVTVPSSSLGTQTYTCNVDHKPSNTKVDKRV	14

8. NUMBERED EMBODIMENTS

[0261] While various specific embodiments have been illustrated and described, it will be appreciated that various changes can be made without departing from the spirit and scope of the disclosure(s). The present disclosure is exemplified by the numbered embodiments set forth below.

[0262] In the numbered embodiments that follow, the targeting moiety preferably binds to a mammalian target molecule, the IL2 and IL2R α moieties are preferably derived from a mammalian IL2 and IL2R α , the Fc domains are preferably derived from a mammalian antibody, and the subjects preferably mammals. More preferably, the mammal is human.

1. An IL2 proprotein comprising:

(a) a first polypeptide chain comprising:

- (i) a first Fc domain;
- (ii) a first linker which is a protease-cleavable linker (PCL) or a non-cleavable linker ("NCL");
- (iii) a first IL2R α moiety;
- (iv) a second linker which is a protease-cleavable linker (PCL); and
- (v) a first IL2 moiety; and

(b) a second polypeptide chain comprising:

- (i) a second Fc domain capable of associating with the first Fc domain to form an Fc region;
- (ii) a third linker which is a protease-cleavable linker (PCL) or a non-cleavable linker ("NCL");

- (iii) a second IL2R α moiety;
- (iv) a fourth linker which is a protease-cleavable linker (PCL); and
- (v) a second IL2 moiety.

2. The IL2 proprotein of embodiment 1, wherein the IL2 moiety comprises an amino
5 acid sequence having at least about 90% sequence identity to mature human IL2.

3. The IL2 proprotein of embodiment 1, wherein the IL2 moiety comprises an amino
acid sequence having about 95% sequence identity to mature human IL2.

4. The IL2 proprotein of any one of embodiments 1 to 3, wherein the IL2 moiety
comprises an amino acid sequence having an N-terminal alanine deletion as compared to
10 mature human IL2.

5. The IL2 proprotein of any one of embodiments 1 to 4, wherein the IL2 moiety
comprises an amino acid sequence having an amino acid substitution at position N88 as
compared to wild type IL2, optionally wherein the amino acid substitution is N88D.

6. The IL2 proprotein of any one of embodiments 1 to 5, wherein the IL2 moiety
15 comprises an amino acid sequence having the amino acid substitution C125S, C125A or C125V
as compared to wild type IL2.

7. The IL2 proprotein of any one of embodiments 1 to 6, wherein the IL2R α moiety
comprises or consists of an amino acid sequence having at least about 90% sequence identity
to an IL2 binding portion of human IL2R α .

20 8. The IL2 proprotein of any one of embodiments 1 to 6, wherein the IL2R α moiety
comprises or consists of an amino acid sequence having at least about 95% sequence identity
to an IL2 binding portion of human IL2R α .

9. The IL2 proprotein of any one of embodiments 1 to 6, wherein the IL2R α moiety
comprises or consists of an amino acid sequence having at least about 97% sequence identity
25 to an IL2 binding portion of human IL2R α .

10. The IL2 proprotein of any one of embodiments 1 to 6, wherein the IL2R α moiety comprises or consists of an amino acid sequence having at least about 98% sequence identity to an IL2 binding portion of human IL2R α .
- 5 11. The IL2 proprotein of any one of embodiments 1 to 6, wherein the IL2R α moiety comprises or consists of an amino acid sequence having at least about 99% sequence identity to an IL2 binding portion of human IL2R α .
12. The IL2 proprotein of any one of embodiments 1 to 6, wherein the IL2R α moiety comprises or consists of an amino acid sequence having 100% sequence identity to an IL2 binding portion of human IL2R α .
- 10 13. The IL2 proprotein of any one of embodiments 7 to 12, wherein the IL2R α moiety comprises an amino acid sequence having at least 90% sequence identity to (a) amino acids 22-186 of IL2R α , (b) amino acids 22-240 of IL2R α , and/or (c) amino acids 22-272 of human IL2R α .
- 15 14. The IL2 proprotein of any one of embodiments 7 to 12, wherein the IL2R α moiety comprises an amino acid sequence having at least 95% sequence identity to (a) amino acids 22-186 of IL2R α , (b) amino acids 22-240 of IL2R α , and/or (c) amino acids 22-272 of human IL2R α .
- 20 15. The IL2 proprotein of any one of embodiments 7 to 12, wherein the IL2R α moiety comprises an amino acid sequence having at least 96% sequence identity to (a) amino acids 22-186 of IL2R α , (b) amino acids 22-240 of IL2R α , and/or (c) amino acids 22-272 of human IL2R α .
- 25 16. The IL2 proprotein of any one of embodiments 7 to 12, wherein the IL2R α moiety comprises an amino acid sequence having at least 97% sequence identity to (a) amino acids 22-186 of IL2R α , (b) amino acids 22-240 of IL2R α , and/or (c) amino acids 22-272 of human IL2R α .
17. The IL2 proprotein of any one of embodiments 7 to 12, wherein the IL2R α moiety comprises an amino acid sequence having at least 98% sequence identity to (a) amino acids

22-186 of IL2R α , (b) amino acids 22-240 of IL2R α , and/or (c) amino acids 22-272 of human IL2R α .

18. The IL2 proprotein of any one of embodiments 7 to 12, wherein the IL2R α moiety comprises an amino acid sequence having at least 99% sequence identity to (a) amino acids
5 22-186 of IL2R α , (b) amino acids 22-240 of IL2R α , and/or (c) amino acids 22-272 of human IL2R α .

19. The IL2 proprotein of any one of embodiments 1 to 18, wherein the second linker, the fourth linker, optionally the first linker, optionally the third linker, or any combination of two or more or all of the foregoing (e.g., (i) first and third linkers, (ii) second and fourth linkers,
10 (iii) first and second linkers, (iv) third and fourth linkers, (v) first, second, third and fourth linkers) comprise(s) a substrate sequence cleavable by any protease set forth in Table A.

20. The IL2 proprotein of any one of embodiments 1 to 19, wherein the second linker, the fourth linker, optionally the first linker, optionally the third linker, or any combination of two or more or all of the foregoing (e.g., (i) first and third linkers, (ii) second and fourth linkers,
15 (iii) first and second linkers, (iv) third and fourth linkers, (v) first, second, third and fourth linkers) comprise(s) one or more substrate sequences selected from the substrate sequences set forth in Table B.

21. The IL2 proprotein of any one of embodiments 1 to 20, wherein the second linker, the fourth linker, optionally the first linker, optionally the third linker, or any combination of two or more or all of the foregoing (e.g., (i) first and third linkers, (ii) second and fourth linkers,
20 (iii) first and second linkers, (iv) third and fourth linkers, (v) first, second, third and fourth linkers) comprise(s) one or more spacer sequences selected from the substrate sequences set forth in Table C.

22. The IL2 proprotein of any one of embodiments 1 to 21, wherein the second
25 linker, the fourth linker, optionally the first linker, optionally the third linker, or any combination of two or more or all of the foregoing (e.g., (i) first and third linkers, (ii) second and fourth linkers, (iii) first and second linkers, (iv) third and fourth linkers, (v) first, second, third and fourth linkers) comprise(s) the amino acid sequence of any of the PCL sequences set forth in Table D or a variant thereof with up to 5 amino acid substitutions, e.g., a variant thereof with 1 amino acid

substitution, 2 amino acid substitutions, 3 amino acid substitutions, 4 amino acid substitutions, or 5 amino acid substitutions.

23. The IL2 proprotein of any one of embodiments 1 to 22, wherein the first and third linkers are identical and/or the third and fourth linkers are identical.

5 24. The IL2 proprotein of embodiment 23, wherein the first, second, third and fourth linkers are identical.

25. The IL2 proprotein of any one of embodiments 1 to 23, wherein the first and third linkers are non-cleavable linkers.

10 26. The IL2 proprotein of embodiment 25, wherein the non-cleavable linkers comprise or consist of any of the NCL sequences set forth in Table E.

27. The IL2 proprotein of embodiment 25, wherein the first and third linkers range from 2 to 60 amino acids in length.

28. The IL2 proprotein of embodiment 25, wherein the first and third linkers range from 5 to 25 amino acids in length.

15 29. The IL2 proprotein of embodiment 25, wherein the first and third linkers range from 7 to 20 amino acids in length.

30. The IL2 proprotein of embodiment 25, wherein the first and third linkers are at least 5 amino acids in length.

20 31. The IL2 proprotein of any one of embodiments 1 to 23, wherein the first, second, third, and fourth linkers are protease cleavable linkers.

32. The IL2 proprotein of embodiment 31, wherein the first, second, third, and fourth linkers range from 20 amino acids to 80 amino acids in length.

33. The IL2 proprotein of embodiment 31, wherein the first, second, third, and fourth linkers range from 20 amino acids to 60 amino acids in length.

34. The IL2 proprotein of any one of embodiments 1 to 33, wherein the first Fc domain and/or the second Fc domain comprises a hinge domain.

35. The IL2 proprotein of any one of embodiments 1 to 34, which further comprises one or more targeting moieties that bind to one or more target molecules.

5 36. The IL2 proprotein of embodiment 35, which comprises a first targeting moiety and/or a second targeting moiety.

37. The IL2 proprotein of embodiment 36, which comprises a first targeting moiety or component thereof (e.g., the VH of a Fab) N-terminal to the first Fc domain and/or a second targeting moiety or component thereof (e.g., the VH of a Fab) N-terminal to the second Fc
10 domain.

38. The IL2 proprotein of embodiment 35 or embodiment 37, wherein the first targeting moiety and/or the second targeting moiety is a Fab.

39. The IL2 proprotein of embodiment 35 or embodiment 37, wherein the first targeting moiety and/or the second targeting moiety is an scFv.

15 40. The IL2 proprotein of any one of embodiments 35 to 39, wherein the first targeting moiety and/or second targeting moiety is capable of binding to an extracellular matrix ("ECM") antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen ("TCA"), a checkpoint inhibitor, or a tumor-associated antigen ("TAA").

20 41. The IL2 proprotein of any one of embodiments 35 to 40, wherein the first targeting moiety and/or second targeting moiety is capable of binding to any target molecule identified in Section 6.7.

42. The IL2 proprotein of any one of embodiments 35 to 41, wherein the first targeting moiety and/or second targeting moiety (a) comprises the (i) CDR or (ii) VH and VL
25 sequences of antibody set forth in Table F or (b) competes with the antibody set forth in Table F for binding to the target molecule.

43. The IL2 proprotein of any one of embodiments 35 to 41, wherein the first targeting moiety and/or second targeting moiety is capable of binding to an ECM antigen which is optionally selected from syndecan, heparanase, integrins, osteopontin, link, cadherins, laminin, laminin type EGF, lectin, fibronectin, notch, nectin (*e.g.*, nectin-4), tenascin, collagen
5 (*e.g.*, collagen type X) and matrixin.

44. The IL2 proprotein of embodiment 43, wherein the first targeting moiety and/or second targeting moiety is capable of binding to a nectin, *e.g.*, nectin 4.

45. The IL2 proprotein of embodiment 43, wherein the first targeting moiety and/or second targeting moiety is capable of binding to a collagen, *e.g.*, collagen X.

10 46. The IL2 proprotein of any one of embodiments 35 to 41, wherein the first targeting moiety and/or second targeting moiety is capable of binding to a cell surface molecule of tumor or viral lymphocytes.

47. The IL2 proprotein of embodiment 46, wherein the antigen is a T-cell co-stimulatory protein.

15 48. The IL2 proprotein of embodiment 47, wherein the T-cell co-stimulatory protein is CD27, CD28, 4-1BB (CD137), OX40, CD30, CD40, ICOS, lymphocyte function-associated antigen-1 (LFA-1), CD2, CD7, LIGHT, NKG2C, or B7-H3.

49. The IL2 proprotein of embodiment 48, wherein the T-cell co-stimulatory protein is B7-H3.

20 50. The IL2 proprotein of any one of embodiments 35 to 41, wherein the first targeting moiety and/or second targeting moiety is capable of binding to a checkpoint inhibitor.

51. The IL2 proprotein of embodiment 50, wherein the checkpoint inhibitor is CTLA-4, PD1, PDL1, PDL2, B7-H3, B7-H4, BTLA, HVEM, TIM3, GAL9, LAG3, VISTA, KIR, 2B4, CD160, CGEN-15049, CHK1, or CHK2.

25 52. The IL2 proprotein of embodiment 51, wherein the checkpoint inhibitor is PDL1.

53. The IL2 proprotein of embodiment 51, wherein the checkpoint inhibitor is PD1.
54. The IL2 proprotein of embodiment 51, wherein the checkpoint inhibitor is LAG3.
55. The IL2 proprotein of any one of embodiments 35 to 41, wherein the first
targeting moiety and/or second targeting moiety is capable of binding to a tumor-associated
antigen ("TAA").
56. The IL2 proprotein of embodiment 55, wherein the first targeting moiety and/or
second targeting moiety is capable of binding to AFP, ALK, a BAGE protein, BIRC5 (survivin),
BIRC7, β -catenin, bcr-abl, BRCA1, BORIS, CA9, carbonic anhydrase IX, caspase-8, CALR,
CEACAM5 (also known as carcinoembryonic antigen or CEA), CCR5, CD19, CD20 (MS4A1),
CD22, CD30, CD40, CDK4, CEA, CTLA4, cyclin-B1, CYP1B1, EGFR, EGFRvIII, ErbB2/Her2,
ErbB3, ErbB4, ETV6-AML, EpCAM, EphA2, Fra-1, FOLR1, a GAGE protein (e.g., GAGE-1 or -
2), GD2, GD3, GloboH, glypican-3, GM3, gp100, Her2, HLA/B-raf, HLA/k-ras, HLA/MAGE-A3,
hTERT, LMP2, MAGE proteins (e.g., MAGE-1, -2, -3, -4, -6, and -12), MART-1, mesothelin, ML-
IAP, Muc1, Muc2, Muc3, Muc4, Muc5, Muc16 (CA-125), MUM1, NA17, NY-BR1, NY-BR62, NY-
BR85, NY-ESO1, OX40, p15, p53, PAP, PAX3, PAX5, PCTA-1, PLAC1, PRLR, PRAME, PSMA
(FOLH1), RAGE proteins, Ras, RGS5, Rho, SART-1, SART-3, STEAP1, STEAP2, TAG-72,
TGF- β , TMPRSS2, Thompson-nouvelle antigen (Tn), TRP-1, TRP-2, tyrosinase, or uroplakin-3.
57. The IL2 proprotein of embodiment 56, wherein the TAA is EGFR.
58. The IL2 proprotein of embodiment 56, wherein the TAA is HER2.
59. The IL2 proprotein of embodiment 56, wherein the TAA is EPCAM.
60. The IL2 proprotein of embodiment 56, wherein the TAA is CEACAM5.
61. The IL2 proprotein of embodiment 56, wherein the TAA is CD20.
62. The IL2 proprotein of any one of embodiments 1 to 61, wherein the Fc region is
homodimeric.

63. The IL2 proprotein of any one of embodiments 1 to 61, wherein the Fc region is heterodimeric.

64. The IL2 proprotein of any one of embodiments 1 to 63, wherein:

- (a) the first polypeptide chain comprises, in N- to C-terminal orientation:
- 5 (i) the first Fc domain;
- (ii) the first linker;
- (iii) the first IL2 moiety;
- (iv) the second linker; and
- (v) the first IL2R α moiety; and
- 10 (b) the second polypeptide chain comprises, in N- to C-terminal orientation;
- (i) the second Fc domain;
- (ii) the third linker;
- (iii) the second IL2 moiety;
- (iv) the fourth linker; and
- 15 (v) the second IL2R α moiety.

65. An IL2 proprotein, which is optionally an IL2 proprotein of any one of embodiments 1 to 64, wherein the IL2 proprotein comprises:

- (a) a first Fc domain and a second Fc domain capable of associating to form an Fc region;
- 20 (b) two linkers C-terminal to the Fc domains that are non-cleavable or protease cleavable;
- (c) two IL2 moieties C-terminal to the first and third linkers;
- (d) two further linkers C-terminal to the IL2 moieties that are protease-cleavable; and
- 25 (e) two IL2R α moieties C-terminal to the second and fourth linkers.

66. An IL2 proprotein according to any one of embodiments 1 to 63, wherein

(a) the first polypeptide chain comprises: the first Fc domain, followed by the first linker where the first linker is a protease-cleavable linker, followed by the first IL2 moiety, followed by the second linker, followed by the first IL2R α moiety; and

(b) the second polypeptide chain comprises: the second Fc domain, followed
5 by the third linker where the third linker is a protease-cleavable linker, followed by the second IL2 moiety, followed by the fourth linker, followed by the second IL2R α moiety.

67. An IL2 proprotein according to any one of embodiments 1 to 63, wherein

(a) the first polypeptide chain comprises: the first Fc domain, followed by the first linker where the first linker is a non-cleavable linker, followed by the first IL2 moiety,
10 followed by the second linker, followed by the first IL2R α moiety; and

(b) the second polypeptide chain comprises: the second Fc domain, followed by the third linker where the third linker is a non-cleavable linker, followed by the second IL2 moiety, followed by the fourth linker, followed by the second IL2R α moiety.

68. An IL2 proprotein according to any one of embodiments 1 to 63, which has the
15 configuration depicted in FIG. 1A.

69. An IL2 proprotein according to any one of embodiments 1 to 63, which has the configuration depicted in FIG. 2A.

70. An IL2 proprotein, which is optionally an IL2 proprotein of any one of embodiments 1 to 64, wherein the IL2 proprotein comprises:

20 (a) a first polypeptide chain comprising, in N- to C-terminal order:

(i) a first amino acid sequence having at least about 95% sequence identity to any one of SEQ ID NOs:1, 2, 3, 4, 5, and 6;

(ii) (1) a cleavable linker / cleavable means for connecting the first amino acid sequence to a third amino sequence, optionally wherein the cleavable linker /
25 cleavable means comprises or consists of a second amino acid sequence comprising one or more sequences set forth in Table B or Table D; or (2) a non-cleavable linker / non-cleavable means for connecting the first amino acid sequence to a third amino sequence, optionally

wherein the non-cleavable linker / non-cleavable means comprises or consists of a sequence set forth in Table E;

(iii) the third amino acid sequence, having at least about 95% sequence identity to SEQ ID NO:7;

5 (iv) a fourth amino acid sequence comprising a sequence set forth in Table B; and

(v) a fifth amino acid sequence having at least about 95% sequence identity to any one of SEQ ID NOs:8, 9, and 10; and

(b) a second polypeptide chain comprising, in N- to C-terminal order:

10 (i) a sixth amino acid sequence having at least about 95% sequence identity to any one of SEQ ID NOs:1, 2, 3, 4, 5, and 6;

(ii) (1) a cleavable linker / cleavable means for connecting the sixth amino acid sequence to an eighth amino sequence, optionally wherein the cleavable linker / cleavable means comprises or consists of a seventh amino acid sequence comprising one or
15 more sequences set forth in Table B or Table D; or (2) a non-cleavable linker / non-cleavable means for connecting the sixth amino acid sequence to an eighth amino sequence, optionally wherein the non-cleavable linker / non-cleavable means comprises or consists of a sequence set forth in Table E;

(iii) the eighth amino acid sequence, having at least about 95%
20 sequence identity to SEQ ID NO:7;

(iv) a ninth amino acid sequence comprising a sequence set forth in Table B; and

(v) a tenth amino acid sequence having at least about 95% sequence identity to SEQ ID NO:8, 9, or 10.

25 71. The IL2 proprotein of embodiment 70, wherein the first polypeptide comprises, N-terminal to the first amino acid sequence, an eleventh amino acid sequence having at least about 95% sequence identity to any one of SEQ ID NOs:11, 12, 13, or 14.

72. The IL2 proprotein of embodiment 71, wherein the eleventh amino acid sequence is the amino acid sequence of SEQ ID NO: 11.

73. The IL2 proprotein of embodiment 71, wherein the eleventh amino acid sequence is the amino acid sequence of SEQ ID NO: 12.

5 74. The IL2 proprotein of embodiment 71, wherein the eleventh amino acid sequence is the amino acid sequence of SEQ ID NO: 13.

75. The IL2 proprotein of embodiment 71, wherein the eleventh amino acid sequence is the amino acid sequence of SEQ ID NO: 14.

10 76. The IL2 proprotein of any one of embodiments 70 to 75, wherein the second polypeptide comprises, N-terminal to the first amino acid sequence, a twelfth amino acid sequence having at least about 95% sequence identity to any one of SEQ ID NOs:11, 12, 13, or 14.

77. The IL2 proprotein of embodiment 76, wherein the twelfth amino acid sequence is the amino acid sequence of SEQ ID NO: 11.

15 78. The IL2 proprotein of embodiment 76, wherein the twelfth amino acid sequence is the amino acid sequence of SEQ ID NO: 12.

79. The IL2 proprotein of embodiment 76, wherein the twelfth amino acid sequence is the amino acid sequence of SEQ ID NO: 13.

20 80. The IL2 proprotein of embodiment 76, wherein the twelfth amino acid sequence is the amino acid sequence of SEQ ID NO: 14.

81. The IL2 proprotein of any one of embodiments 70 to 80, wherein the first amino acid sequence has at least about 98% sequence identity to any one of SEQ ID NOs:1, 2, 3, 4, 5, or 6.

82. The IL2 proprotein of any one of embodiments 70 to 80, wherein the first amino acid sequence is the amino acid sequence of any one of SEQ ID NOs:1, 2, 3, 4, 5, or 6.

83. The IL2 proprotein of embodiment 82, wherein the first amino acid sequence is the amino acid sequence of SEQ ID NO:5.

5 84. The IL2 proprotein of embodiment 82, wherein the first amino acid sequence is the amino acid sequence of SEQ ID NO:6.

85. The IL2 proprotein of any one of embodiments 70 to 84, wherein the first amino acid sequence is 350 or fewer amino acids in length.

10 86. The IL2 proprotein of any one of embodiments 70 to 84, wherein the first amino acid sequence is 330 or fewer amino acids in length.

87. The IL2 proprotein of any one of embodiments 70 to 86, wherein the sixth amino acid sequence has at least about 98% sequence identity to any one of SEQ ID NOs:1, 2, 3, 4, 5, or 6.

15 88. The IL2 proprotein of any one of embodiments 70 to 80, wherein the sixth amino acid sequence is the amino acid sequence of any one of SEQ ID NOs:1, 2, 3, 4, 5, or 6.

89. The IL2 proprotein of embodiment 88, wherein the sixth amino acid sequence is the amino acid sequence of SEQ ID NO:5.

90. The IL2 proprotein of embodiment 88, wherein the sixth amino acid sequence is the amino acid sequence of SEQ ID NO:6.

20 91. The IL2 proprotein of any one of embodiments 70 to 90, wherein the sixth amino acid sequence is 350 or fewer amino acids in length.

92. The IL2 proprotein of any one of embodiments 70 to 90, wherein the sixth amino acid sequence is 330 or fewer amino acids in length.

93. The IL2 proprotein of any one of embodiments 70 to 92, wherein the second amino acid sequence comprises one or more amino acid sequences set forth in Table B.
94. The IL2 proprotein of embodiment 93, wherein the second amino acid sequence comprises the sequence HPVGLLAR (SEQ ID NO: 163).
- 5 95. The IL2 proprotein of embodiment 93, wherein the second amino acid sequence comprises the sequence VPLSLYSG (SEQ ID NO: 159).
96. The IL2 proprotein of embodiment 93, wherein the second amino acid sequence comprises the sequence ISSGLLS (SEQ ID NO: 370).
97. The IL2 proprotein of embodiment 93, wherein the second amino acid sequence
10 comprises the sequence PLGLWSQ (SEQ ID NO: 115).
98. The IL2 proprotein of any one of embodiments 70 to 93, wherein the second amino acid sequence is an amino acid sequence set forth in Table D.
99. The IL2 proprotein of embodiment 98, wherein the second amino acid sequence is the amino acid sequence GGGISSGLLSGRSDNHGGGISSGLLSGRSDNHGGS (SEQ ID NO:
15 199).
100. The IL2 proprotein of embodiment 98, wherein the second amino acid sequence is the amino acid sequence GGSHPVGLLARGGGHPVGLLARGGGHPVGLLARGS (SEQ ID NO: 203).
101. The IL2 proprotein of embodiment 98, wherein the second amino acid sequence
20 is the amino acid sequence GGGHPVGLLARGGGS (SEQ ID NO: 285).
102. The IL2 proprotein of embodiment 98, wherein the second amino acid sequence is the amino acid sequence GISSGLLSGRSDNHG (SEQ ID NO: 282).
103. The IL2 proprotein of embodiment 98, wherein the second amino acid sequence is the amino acid sequence GGGSISSGLLSGRSDNHGGS (SEQ ID NO: 283).

104. The IL2 proprotein of embodiment 98, wherein the second amino acid sequence is the amino acid sequence GGGISSGLLSGRSDNHGGGS (SEQ ID NO: 284).

105. The IL2 proprotein of embodiment 98, wherein the second amino acid sequence is the amino acid sequence GGGGSGGGGSGGGGSVPLSLYSGGGSGGSGGSGS (SEQ ID NO: 221).

106. The IL2 proprotein of any one of embodiments 70 to 92, wherein the second amino acid sequence is an amino acid sequence set forth in Table E.

107. The IL2 proprotein of embodiment 106, wherein the second amino acid sequence is the amino acid sequence (GGGGS)_n, wherein n is 1, 2, 3, 4, or 5 (SEQ ID NO: 357).

108. The IL2 proprotein of any one of embodiments 93 to 107, wherein the second amino acid sequence is 25 or fewer amino acids in length.

109. The IL2 proprotein of any one of embodiments 93 to 107, wherein the second amino acid sequence is 15 or fewer amino acids in length.

110. The IL2 proprotein of any one of embodiments 93 to 107, wherein the second amino acid sequence is 6 or fewer amino acids in length.

111. The IL2 proprotein of any one of embodiments 70 to 110, wherein the seventh amino acid sequence comprises one or more amino acid sequences set forth in Table B.

112. The IL2 proprotein of embodiment 111, wherein the seventh amino acid sequence comprises the sequence HPVGLLAR (SEQ ID NO: 163).

113. The IL2 proprotein of embodiment 111, wherein the seventh amino acid sequence comprises the sequence VPLSLYSG (SEQ ID NO: 159).

114. The IL2 proprotein of embodiment 111, wherein the seventh amino acid sequence comprises the sequence ISSGLLS (SEQ ID NO: 370).

115. The IL2 proprotein of embodiment 111, wherein the seventh amino acid sequence comprises the sequence PLGLWSQ (SEQ ID NO: 115).

116. The IL2 proprotein of any one of embodiments 70 to 111, wherein the seventh amino acid sequence is an amino acid sequence set forth in Table D.

5 117. The IL2 proprotein of embodiment 116, wherein the second amino acid sequence is the amino acid sequence GGGISSGLLSGRSDNHGGGISSGLLSGRSDNHGGS (SEQ ID NO: 199).

118. The IL2 proprotein of embodiment 116, wherein the second amino acid sequence is the amino acid sequence GGSHPVGLLARGGGHPVGLLARGGGHPVGLLARGS (SEQ ID
10 NO: 203).

119. The IL2 proprotein of embodiment 116, wherein the second amino acid sequence is the amino acid sequence GGGHPVGLLARGGGS (SEQ ID NO: 285).

120. The IL2 proprotein of embodiment 116, wherein the second amino acid sequence is the amino acid sequence GISSGLLSGRSDNHG (SEQ ID NO: 282).

15 121. The IL2 proprotein of embodiment 116, wherein the second amino acid sequence is the amino acid sequence GGGSISSGLLSGRSDNHGGGS (SEQ ID NO: 283).

122. The IL2 proprotein of embodiment 116, wherein the second amino acid sequence is the amino acid sequence GGGISSGLLSGRSDNHGGGS (SEQ ID NO: 284).

20 123. The IL2 proprotein of embodiment 116, wherein the second amino acid sequence is the amino acid sequence GGGSGGGSGGGSGVPLSLYSGGGSGGGSGGS (SEQ ID NO: 221).

124. The IL2 proprotein of any one of embodiments 70 to 110, wherein the seventh amino acid sequence is an amino acid sequence set forth in Table E.

125. The IL2 proprotein of embodiment 124, wherein the second amino acid sequence is the amino acid sequence (GGGGS)_n, wherein n is 1, 2, 3, 4, or 5 (SEQ ID NO: 357).

126. The IL2 proprotein of embodiment any one of embodiments 111 to 125, wherein the seventh amino acid sequence is 25 or fewer amino acids in length.

5 127. The IL2 proprotein of embodiment any one of embodiments 111 to 125, wherein the seventh amino acid sequence is 15 or fewer amino acids in length.

128. The IL2 proprotein of embodiment any one of embodiments 111 to 125, wherein the seventh amino acid sequence is 6 or fewer amino acids in length.

10 129. The IL2 proprotein of any one of embodiments 70 to 128, wherein the third amino acid sequence has at least about 98% sequence identity to SEQ ID NO:7.

130. The IL2 proprotein of any one of embodiments 70 to 128, wherein the third amino acid sequence is the amino acid sequence of SEQ ID NO:7.

131. The IL2 proprotein of any one of embodiments 70 to 130, wherein the third amino acid sequence is 150 or fewer amino acids in length.

15 132. The IL2 proprotein of any one of embodiments 70 to 130, wherein the third amino acid sequence is 135 or fewer amino acids in length.

133. The IL2 proprotein of any one of embodiments 70 to 132, wherein the eighth amino acid sequence has at least about 98% sequence identity to SEQ ID NO:7.

20 134. The IL2 proprotein of any one of embodiments 70 to 132, wherein the eighth amino acid sequence is the amino acid sequence of SEQ ID NO:7.

135. The IL2 proprotein of any one of embodiments 70 to 134, wherein the eighth amino acid sequence is 150 or fewer amino acids in length.

136. The IL2 proprotein of any one of embodiments 70 to 134, wherein the eighth amino acid sequence is 135 or fewer amino acids in length.

137. The IL2 proprotein of any one of embodiments 70 to 136, wherein the fourth amino acid sequence comprises one or more amino acid sequences set forth in Table B.

138. The IL2 proprotein of any one of embodiments 70 to 136, wherein the fourth amino acid sequence is an amino acid sequence set forth in Table D.

5 139. The IL2 proprotein of any one of embodiments 70 to 138, wherein the ninth amino acid sequence comprises one or more amino acid sequences set forth in Table B.

140. The IL2 proprotein of any one of embodiments 70 to 139, wherein the ninth amino acid sequence is an amino acid sequence set forth in Table D.

10 141. The IL2 proprotein of any one of embodiments 70 to 140, wherein the fifth amino acid sequence has at least about 98% sequence identity to SEQ ID NO:8, 9, or 10.

142. The IL2 proprotein of embodiment 141, wherein the fifth amino acid sequence is the amino acid sequence of SEQ ID NO:8.

143. The IL2 proprotein of embodiment 141, wherein the fifth amino acid sequence is the amino acid sequence of SEQ ID NO:9.

15 144. The IL2 proprotein of embodiment 141, wherein the fifth amino acid sequence is the amino acid sequence of SEQ ID NO:10.

145. The IL2 proprotein of any one of embodiments 70 to 144, wherein the fifth amino acid sequence is 255 or fewer amino acids in length.

20 146. The IL2 proprotein of any one of embodiments 70 to 144, wherein the fifth amino acid sequence is 225 or fewer amino acids in length.

147. The IL2 proprotein of any one of embodiments 70 to 144, wherein the fifth amino acid sequence is 170 or fewer amino acids in length.

148. The IL2 proprotein of any one of embodiments 70 to 147, wherein the tenth amino acid sequence has at least about 98% sequence identity to SEQ ID NO:8, 9, or 10.

149. The IL2 proprotein of embodiment 148, wherein the tenth amino acid sequence is the amino acid sequence of SEQ ID NO:8.

150. The IL2 proprotein of embodiment 148, wherein the tenth amino acid sequence is the amino acid sequence of SEQ ID NO:9.

5 151. The IL2 proprotein of embodiment 148, wherein the tenth amino acid sequence is the amino acid sequence of SEQ ID NO:10.

152. The IL2 proprotein of any one of embodiments 70 to 151, wherein the tenth amino acid sequence is 255 or fewer amino acids in length.

10 153. The IL2 proprotein of any one of embodiments 70 to 151, wherein the tenth amino acid sequence is 225 or fewer amino acids in length.

154. The IL2 proprotein of any one of embodiments 70 to 151, wherein the tenth amino acid sequence is 170 or fewer amino acids in length.

155. The IL2 proprotein of any one of embodiments 70 to 154, wherein the first polypeptide chain lacks additional sequences C-terminal to the first amino acid sequence.

15 156. The IL2 proprotein of any one of embodiments 70 to 155, wherein the first polypeptide chain lacks additional sequences between the first and second amino acid sequences.

20 157. The IL2 proprotein of any one of embodiments 70 to 156, wherein the first polypeptide chain lacks additional sequences between the second and third amino acid sequences.

158. The IL2 proprotein of any one of embodiments 70 to 157, wherein the first polypeptide chain lacks additional sequences between the third and fourth amino acid sequences.

159. The IL2 proprotein of any one of embodiments 70 to 158, wherein the first polypeptide chain lacks additional sequences between the fourth and fifth amino acid sequences.

160. The IL2 proprotein of any one of embodiments 70 to 159, wherein the second
5 polypeptide chain lacks additional sequences C-terminal to the sixth amino acid sequence.

161. The IL2 proprotein of any one of embodiments 70 to 160, wherein the second polypeptide chain lacks additional sequences between the sixth and seventh amino acid sequences.

162. The IL2 proprotein of any one of embodiments 70 to 161, wherein the second
10 polypeptide chain lacks additional sequences between the seventh and eighth amino acid sequences.

163. The IL2 proprotein of any one of embodiments 70 to 162, wherein the second polypeptide chain lacks additional sequences between the eighth and ninth amino acid sequences.

164. The IL2 proprotein of any one of embodiments 70 to 163, wherein the second
15 polypeptide chain lacks additional sequences between the ninth and tenth amino acid sequences.

165. The IL2 proprotein of any one of embodiments 70 to 164, wherein the first polypeptide and the second polypeptide are identical.

20 166. A nucleic acid or plurality of nucleic acids encoding the IL2 proprotein of any one of embodiments 1 to 165.

167. A host cell engineered to express the IL2 proprotein of any one of embodiments 1 to 165 or the nucleic acid(s) of embodiment 166.

168. A method of producing the IL2 proprotein of any one of embodiments 1 to 165, comprising culturing the host cell of embodiment 167 and recovering the IL2 proprotein expressed thereby.

169. A pharmaceutical composition comprising the IL2 proprotein of any one of
5 embodiments 1 to 165 and an excipient.

170. A method of treating cancer, comprising administering to a subject in need thereof the IL2 proprotein of any one of embodiments 1 to 165 or the pharmaceutical composition of embodiment 169.

171. The method of embodiment 170, wherein the IL2 proprotein comprises at least
10 one targeting moiety that is capable of binding to a target molecule and wherein the cancer is associated with expression of the target molecule, e.g., a TAA and associated cancer as set forth in Table I.

172. The method of embodiment 171, wherein an activated IL2 protein comprising the IL2 moiety is produced by cleavage of one or more protease-cleavable linkers in the IL2
15 proprotein by one or more proteases expressed by the cancer tissue.

173. The method of embodiment 172, wherein the IL2 protein is selectively activated in the cancer tissue.

174. A method of localized delivery of an IL2 protein, comprising administering to a subject an IL2 proprotein according to any one of embodiments 1 to 165 (or a pharmaceutical
20 composition comprising the IL2 proprotein and an excipient) which has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by a tissue to which the IL2 protein is to be locally delivered.

175. The method of embodiment 174, wherein the IL2 proprotein comprises one or more targeting moieties that recognize a target molecule expressed by the tissue.

25 176. The method of embodiment 175, wherein the IL2 proprotein comprises two targeting moieties that each recognize a target molecule expressed by the tissue.

177. The method of embodiment 175 or embodiment 176, wherein the tissue is cancer tissue.

178. The method of embodiment 177, wherein the one or more targeting moieties are capable of binding to an extracellular matrix ("ECM") antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen ("TCA"), a checkpoint inhibitor, or a tumor-associated antigen ("TAA").

179. The method of any one of embodiments 174 to 178, wherein an activated IL2 protein comprising the IL2 moiety is produced by cleavage of one or more protease-cleavable linkers in the IL2 proprotein by one or more proteases in the tissue.

180. A method of treating cancer with an IL2 protein that is selectively activated in cancer tissue, comprising administering to a subject in need thereof an IL2 proprotein according to any one of embodiments 1 to 165 (or a pharmaceutical composition comprising the IL2 proprotein and an excipient) which has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by cancer tissue to which the IL2 protein.

181. The method of embodiment 180, wherein the IL2 proprotein comprises one or more targeting moieties that recognize a target molecule expressed by the cancer tissue.

182. The method of embodiment 181, wherein the IL2 proprotein comprises two targeting moieties that each recognize a target molecule expressed by the cancer tissue.

183. The method of embodiment 181 or embodiment 182, wherein the one or more targeting moieties are capable of binding to an extracellular matrix ("ECM") antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen ("TCA"), a checkpoint inhibitor, or a tumor-associated antigen ("TAA").

184. The method of any one of embodiments 180 to 183, wherein an activated IL2 protein comprising the IL2 moiety is produced by cleavage of one or more protease-cleavable linkers in the IL2 proprotein by one or more proteases in the cancer tissue.

185. A method of administering to the subject IL2 therapy with reduced systemic exposure and/or reduced systemic toxicity, comprising administering to a subject the IL2 therapy in the form of an IL2 proprotein according to any one of embodiments 1 to 165 (or a pharmaceutical composition comprising the IL2 proprotein and an excipient) which has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by a tissue for which IL2 therapy is desirable and/or intended.

186. The method of embodiment 185, wherein the IL2 proprotein comprises one or more targeting moieties that recognize a target molecule expressed by the tissue.

187. The method of embodiment 186, wherein the IL2 proprotein comprises two targeting moieties that each recognize a target molecule expressed by the tissue.

188. The method of any one of embodiments 185 to 187, wherein the tissue is cancer tissue.

189. The method of embodiment 188, wherein the one or more targeting moieties are capable of binding to an extracellular matrix ("ECM") antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen ("TCA"), a checkpoint inhibitor, or a tumor-associated antigen ("TAA").

190. The method of any one of embodiments 184 to 189, wherein an activated IL2 protein comprising the IL2 moiety is produced by cleavage of one or more protease-cleavable linkers in the IL2 proprotein by one or more proteases in the tissue.

191. A method of treating cancer with an IL2 protein that is selectively activated in cancer tissue, comprising administering to a subject in need thereof an IL2 proprotein according to any one of embodiments 1 to 165 (or a pharmaceutical composition comprising the IL2 proprotein and an excipient) which has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by the cancer tissue.

192. The method of embodiment 191, wherein the IL2 proprotein comprises one or more targeting moieties that recognize a target molecule expressed by the cancer tissue.

193. The method of embodiment 192, wherein the IL2 proprotein comprises two targeting moieties that each recognize a target molecule expressed by the cancer tissue.

194. The method of embodiment 191 or embodiment 192, wherein the one or more targeting moieties are capable of binding to an extracellular matrix ("ECM") antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen ("TCA"), a checkpoint inhibitor, or a tumor-associated antigen ("TAA").

195. The method of any one of embodiments 191 to 194, wherein an activated IL2 protein comprising the IL2 moiety is produced by cleavage of one or more protease-cleavable linkers in the IL2 proprotein by one or more proteases in the cancer tissue.

196. A method of targeted delivery of an activated IL2 protein to cancer tissue, comprising administering to a subject an IL2 proprotein according to any one of embodiments 1 to 165 (or a pharmaceutical composition comprising the IL2 proprotein and an excipient), wherein the IL2 proprotein:

- (a) comprises one or more targeting moieties that recognize a target molecule expressed by the cancer tissue; and
- (b) has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by a tissue for which IL2 therapy is desirable and/or intended.

197. The method of embodiment 196, wherein the IL2 proprotein comprises two targeting moieties that each recognize a target molecule expressed by the cancer tissue.

198. The method of embodiment 196 or embodiment 197, wherein the one or more targeting moieties are capable of binding to an extracellular matrix ("ECM") antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen ("TCA"), a checkpoint inhibitor, or a tumor-associated antigen ("TAA").

199. The method of any one of embodiments 196 to 198, wherein an activated IL2 protein comprising the IL2 moiety is produced by cleavage of one or more protease-cleavable linkers in the IL2 proprotein by one or more proteases in the cancer tissue.

200. A method of locally inducing an immune response in a target tissue, comprising administering to a subject an IL2 proprotein according to any one of embodiments 1 to 165 (or a pharmaceutical composition comprising the IL2 proprotein and an excipient) which has one or more targeting moieties capable of binding a target molecule expressed in the target tissue and
5 one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed in the target tissue.

201. The method of embodiment 200, wherein the IL2 proprotein comprises two targeting moieties that each recognize a target molecule expressed in the target tissue.

202. The method of embodiment 200 or embodiment 201, wherein the target tissue is
10 cancer tissue.

203. The method of any one of embodiments 200 to 202, wherein the one or more targeting moieties are capable of binding to an extracellular matrix ("ECM") antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen ("TCA"), a checkpoint inhibitor, or a tumor-associated antigen ("TAA").

15 204. The method of any one of embodiments 200 to 203, wherein an activated IL2 protein comprising the IL2 moiety is produced by cleavage of one or more protease-cleavable linkers in the IL2 proprotein by one or more proteases in the target tissue.

205. The method of embodiment 204, wherein the IL2 protein induces the immune response against at least one cell type in the target tissue.

20 206. The method of any one of embodiments 170 to 205, wherein the administration is non-local.

207. The method of embodiment 206, wherein the administration is systemic.

208. The method of embodiment 206, wherein the administration is subcutaneous.

9. EXAMPLES

9.1. Example 1: Production of IL2 Proproteins

- [0263]** Constructs encoding IL2 proproteins comprising targeting moieties, Fc domains, IL2 and IL2R α moieties, and cleavable and/or noncleavable linkers were synthesized as DNA fragments and cloned into suitable expression vectors. A 29-amino acid signal sequence from murine inactive tyrosine-protein kinase transmembrane receptor ROR1 (mROR1) was added to the N-termini of the constructs. All IL2 proproteins were expressed as preproteins containing the signal sequence. The signal sequence was cleaved by intracellular processing to produce a mature protein.
- [0264]** The constructs were transiently expressed in in Expi293F™ cells (ThermoFisher) following the manufacturer's protocol. Proteins in Expi293F supernatant were purified using the ProteinMaker system (Protein BioSolutions, Gaithersburg, MD) with either HiTrap™ Protein G HP or MabSelect SuRe pcc columns (Cytiva). After single step elution, the antibodies were neutralized, dialyzed into a final buffer of phosphate buffered saline (PBS) with 5% glycerol, aliquoted and stored at -80 °C.
- [0265]** An overview of the IL2 proproteins encoded by the generated constructs is provided in Table 1, below. Table 1 describes a single half antibody of each IL2 proprotein, where each IL2 proprotein comprises two, identical half antibodies.

TABLE 1 IL2 Proproteins	
#	Structure
1	Fab – Fc(IgG4s)-3xG4S-hIL2-4XG4S-hIL2R α (22-186)
2	Fab – Fc(IgG4s)-3XG4S-hIL2-PCL(34aa)-hIL2R α (22-186)
3	Fab – Fc(IgG4s)-3xG4S-hIL2-2XG4S-hIL2R α (22-186)
4	Fab – Fc(IgG4s)-3xG4S-hIL2-3XG4S-hIL2R α (22-186)
5	Fab – Fc(IgG4s)-4xG4S-hIL2-3XG4S-hIL2R α (22-186)
6	Fab – Fc(IgG4s)-5xG4S-hIL2-3XG4S-hIL2R α (22-186)
7	Fab – Fc(IgG4s)-5xG4S-hIL2-5XG4S-hIL2R α (22-186)
8	Fab – Fc(IgG4s)-1xG4S-hIL2-1XG4S-hIL2R α (22-186)
9	Fab – Fc(IgG4s)-1xG4S-hIL2-2XG4S-hIL2R α (22-186)
10	Fab – Fc(IgG4s)-1xG4S-hIL2-3XG4S-hIL2R α (22-186)
11	Fab – Fc(IgG4s)-2xG4S-hIL2-2XG4S-hIL2R α (22-186)
12	Fab – Fc(IgG4s)-PCL(15AA)-hIL2-PCL(21AA)-hIL2R α (22-186)
13	Fab – Fc(IgG4s)-PCL(15AA)-hIL2-PCL(15AA)-hIL2R α (22-186)
14	Fab – Fc(IgG4s)-3xG4S-hIL2-PCL(15AA)-hIL2R α (22-186)
15	Fab – Fc(IgG4s)-3xG4S-hIL2-PCL(20AA)-hIL2R α (22-186)
16	Fab – Fc(IgG4s)-2xG4S-hIL2-PCL(15AA)-hIL2R α (22-186)
17	Fab – Fc(IgG4s)-1xG4S-hIL2-PCL(15AA)-hIL2R α (22-186)
18	Fab – Fc(IgG4s)-PCL(15AA)-hIL2-PCL(15AA)-hIL2R α (22-186)
19	Fab – Fc(IgG4s)-PCL(15AA)-hIL2-PCL(15AA)-hIL2R α (22-186)
20	Fab – Fc(IgG4s)-PCL(15AA)-hIL2-PCL(15AA)-hIL2R α (22-186)
21	Fab – Fc(IgG4s)-PCL(8AA)-hIL2-PCL(8AA)-hIL2R α (22-186)
22	Fab – Fc(IgG4s)-PCL(15AA)-hIL2-PCL(20AA)-hIL2R α (22-186)

9.2. Example 2: The Anti-Tumor Activity of EGFR-Targeted IL2 Proproteins

[0266] The anti-tumor activity of EGFR-targeted IL2 proproteins was evaluated in an MC38 tumor model. Briefly, 7×10^5 MC38 tumor cells were implanted subcutaneously into the right hind flanks of 8-10-week-old female mice that express humanized EGFR protein on day 0. Tumor-inoculated mice were randomized into treatment groups on day 9, when average tumor sizes reached 90 mm³. Mice in each randomized group received two total i.p. injections on days 9 and 12. Tumor sizes were measured semiweekly using a digital caliper and the tumor sizes were calculated as $\text{length} \times \text{width}^2 / 2$. Average tumor volumes (mm³ +/- SEM) after tumor

implantation in each treatment group are shown in FIG 3A. Individual tumor growth curves of each treatment group are depicted in FIGs. 3B-3E.

[0267] When tumor growth curves were averaged for each treatment group, EGFR-targeted IL2 proprotein constructs with cleavable linkers displayed more effective inhibition of tumor growth compared to both targeted IL2 proprotein constructs with non-cleavable linkers and non-targeted IL2 proproteins with cleavable linkers and isotype controls (FIG. 3A; arrows indicate the days of treatment). Individual tumor growth curves of each treatment group are shown in FIGs. 3B-3E. Overall, these results suggest that both TAA targeting and cleavable linkers significantly enhanced the *in vivo* anti-tumor efficacy of EGFR-targeted IL2 proprotein constructs.

9.3. Example 3: The Anti-Tumor Activity of PD1-Targeted IL2 Proproteins

[0268] The anti-tumor activity of PD1-targeted IL2 proproteins was evaluated in an MC38 tumor model. On day 0, 3×10^5 MC38 tumor cells (ACL8874) were implanted subcutaneously into the right hind flanks of 8-10-week-old female mice that express humanized PD1. Tumor-inoculated mice were randomized into treatment groups on day 9, when average tumor sizes reached 90 mm³. Mice in each randomized group received a total of two i.p. injections of 1.5 mg/kg of their assigned proprotein on days 9 and 12. Tumor sizes were measured semiweekly using a digital caliper and the tumor sizes were calculated as $\text{length} \times \text{width}^2 / 2$. Average tumor volumes (mm³ $\pm$ SEM) in each treatment group were plotted post dosing (FIG. 4A).

[0269] When tumor growth curves were averaged for each treatment group, tumor-inoculated mice treated with PD1-targeted IL2 proprotein constructs with cleavable linkers displayed diminished tumor growth, whereas mice treated with a non-targeted IL2 proprotein with a cleavable linker or an isotype control displayed no inhibition of tumor growth (FIG. 4A). Individual tumor growth curves of each treatment group revealed that none of the isotype-treated mice displayed tumor inhibition (FIG. 4B). Similarly, none of the mice treated with non-targeted IL2 proprotein displayed tumor inhibition (FIG. 4C). However, four out of five mice treated with PD1-targeted IL2 proprotein were tumor free for the duration of the assessments (FIG. 4D).

[0270] Compared to isotype control or non-targeted IL2 proproteins, PD1-targeted IL2 proproteins with cleavable linkers demonstrated robust tumor growth inhibition with a higher frequency of treated mice undergoing complete tumor regression, demonstrating that TCA targeting enhanced *in vivo* anti-tumor efficacy of PD1-targeted IL2 proproteins with cleavable linkers.

9.4. Example 4: Assessment of the Cleavability of the IL2 Proproteins Comprising Cleavable Linkers

- [0271] To evaluate whether the protease cleavable linkers in the IL2 proproteins were accessible for digestion by recombinant proteases, two constructs, mAb-PCL(15AA)-IL2-PCL(15AA)-IL2R α and mAb-PCL(15AA)-IL2-PCL(20AA)-IL2R α , were selected to be digested using a recombinant human protease. Briefly, each construct was incubated at 37 °C for 24 hours with 200 ng of uPA (R&D Cat # 1310-SE) in the assay buffer following manufacturer's protocol. No digestion controls were incubated in the same conditions for the same duration without the addition of the protease. After digestion, SDS sample loading buffer containing reducing agent was added to each sample. Next the samples were boiled and run on an Invitrogen 4-20% Tris-Glycine gel for SDS PAGE. The proteins were then transferred to a PVDF membrane using iBlot 2 dry transfer method. The membrane was then probed with Biotinylated anti-hIL2 (R&D Cat # BAF202) followed by streptavidin-HRP of detection (Cytiva (Cat # RPN1231)
- [0272] Protease digestion of the IL2 proproteins with uPA resulted in the release of free IL2 from both IL2 proproteins, although the IL2 proprotein construct comprising a 20AA linker between IL2 and IL2R α domains displayed a more complete linker digestion and IL2 release relative to the construct comprising a 15AA linker between IL2 and IL2R α (FIGs 5A and 5B). These observations suggest that the extent of protease digestion depends on the length of the protease cleavable linker.

9.5. Example 5: *In vitro* Activity of Tumor-Targeted IL2 Proproteins

- [0273] The *in vitro* activity of protease-digested and undigested IL2 proproteins comprising protease-cleavable or non-cleavable linkers was evaluated with a luciferase reporter assay as described below in Section 9.5.1.2, using one of the engineered reporter cell lines generated as described in Section 9.5.1.1.

9.5.1. Methods

9.5.1.1. Engineering of YT/STAT5-Luc Reporter Cells

- [0274] The human T/NK-like leukemia YT cell line was electroporated with a Signal Transducer and Activator of Transcription 5 (STAT5)-driven luciferase reporter construct and maintained in Iscove's modified Dulbecco's medium supplemented with 2 mM L-Glutamine/ Penicillin/ Streptomycin + 20% Fetal Bovine Serum (FBS) + 200 μ g/ml hygromycin. A single cell clone, having high responsiveness to IL2, was identified and renamed YT/STAT5-Luc cl.4. IL2R α (CD25) was knocked out in this clone using CRISPR-Cas9 technology, and the resulting cell

line, YT/STAT5-Luc/IL2R α KO, which is referred to as CD25 KO for simplicity, was validated by flow cytometry.

[0275] Human IL2R α was then stably reintroduced into the CD25 KO cell line (amino acids M1-I272 of accession number NP_000408.1), and the resulting cell line, CD25 OE, was validated by flow cytometry and maintained in Iscove's modified Dulbecco's medium supplemented with 2 mM L-Glutamine/Penicillin/Streptomycin + 20% FBS + 200 μ g/ml hygromycin + 15 μ g/mL blasticidin.

[0276] Since YT cells endogenously express PD1, CD25 KO and CD25 OE cells were engineered to knock out PD1 expression using CRISPR/Cas9 technology, and the resulting cell lines, CD25 KO/ PD1 KO and CD25 OE/ PD1 KO, were validated by flow cytometry.

9.5.1.2. Luciferase Reporter Assay

[0277] A day prior to screening, engineered YT/STAT5-Luc reporter cells, CD25 KO/ PD1 KO and CD25 OE/ PD1 KO, were diluted at 3×10^5 cells/mL in RPMI1640 media supplemented with 2 mM L-Glutamine/ Penicillin/ Streptomycin + 10% FBS.

[0278] IL2 proproteins were digested overnight with or without recombinant human uPA (R&D Cat # 1310-SE) in digestion buffer (50 mM Tris, 0.01% (v/v) Tween[®] 20, pH 8.5). 20 nM of enzyme was added per 200 nM of fusion protein and allowed to digest before diluting it with assay medium (RPMI1640 media supplemented with 2 mM L-Glutamine/ Penicillin/ Streptomycin + 10% FBS) for the bioassay.

[0279] On the day of the assay, cells were spun down, resuspended in assay medium, plated at 2.5×10^4 reporter cells/well in 96-well white flat bottom plates, and incubated with recombinant IL2, uPA-digested IL2 proproteins, or undigested IL2 proproteins. Each construct was serially diluted (1:5) over an 11-point titration range (50 nM to 5.12 fM) and a 12th point containing no protein. After plates were incubated for 4 hours and 30 minutes at 37 °C and 5% CO₂, 100 μ L ONE-Glo[™] (Promega) reagent was added to the wells to lyse the cells and detect luciferase activity. The emitted light was measured in RLU on an Envision multilabel plate reader (PerkinElmer).

9.5.2. Results

[0280] IL2 proproteins comprising different tumor-targeting moieties (e.g., anti-CA9, anti-EGFR, or anti-PD1 Fab moieties) were first evaluated using CD25 KO/ PD1 KO cells. For all constructs tested, each protease-cleavable linker (PCL) was 15 amino acids in length.

[0281] The uPA-digested IL2 proproteins resulted in reporter activity similar to the activity associated with recombinant IL2 in each instance, regardless of the targeting moiety (FIGs. 6A-6F). The undigested IL2 proproteins resulted in little to no luciferase activity (FIGs. 6A-6F). Similarly, none of the IL2 proproteins with non-cleavable linkers displayed luciferase activity (FIGs. 6A-6D).

[0282] Next, the same IL2 proproteins were evaluated using CD25 OE/ PD1 KO cells. In this assessment, all constructs were associated with detectable luciferase activity (FIGs. 7A-7F). Yet, recombinant IL2 and uPA-digested IL2 proproteins comprising protease-cleavable linkers displayed relatively high and comparable potency, which was orders of magnitude higher than the potency observed with undigested IL2 proproteins or non-cleavable linker comprising constructs (FIGs. 7A-7F).

[0283] Taken together, these results suggested that IL2 proproteins displayed minimal activity unless their IL2 moieties were released upon protease digestion. Furthermore, IL2 released from the IL2 proproteins was as potent as recombinant IL2.

9.6. Example 6: Effect of Linker Length on IL2 Proprotein Activity

[0284] To reduce treatment-associated side effects, it is important to minimize the activity of IL2 proproteins until they reach the target tissue. The role of linker length on attenuating IL2 proprotein activity was evaluated with a luciferase reporter assay using engineered YT/STAT5-Luc reporter cells.

[0285] RPMI1640 media supplemented with 2mM L-Glutamine/Penicillin/Streptomycin + 10% Fetal Bovine Serum (FBS) was used as the assay medium to prepare cell suspensions and protein dilutions. A day prior to screening, engineered YT/STAT5-Luc reporter cells (CD25 KO/ PD1 KO, CD25 KO/ PD1 OE, CD25 OE/ PD1 OE and CD25 OE/ PD1 KO) were diluted at 3×10^5 cells/mL. On the day of the assay, cells were spun down, resuspended in assay medium, plated at 2.5×10^4 reporter cells/well in 96-well white flat bottom plates, and incubated with recombinant IL2 or PD1-targeted IL2 proproteins with different lengths of non-cleavable linkers. Constructs were serially diluted (1:5) over an 11-point titration range (50 nM to 5.12 fM) and a 12th point containing no protein. Plates were incubated for 4 hours and 30 minutes at 37 °C and 5% CO₂. Next, 100mL ONE-Glo™ (Promega) reagent was added to the wells to lyse the cells and detect luciferase activity. The emitted light was measured in RLU on an Envision multilabel plate reader (PerkinElmer).

[0286] Generally, IL2 proproteins with longer linker lengths displayed greater luciferase activity than their counterparts with shorter linkers (FIGs. 8A-8D). This linker length-dependent attenuation of activity was most pronounced in PD1 OE/ CD25 KO cells (FIG. 8A).

10. CITATION OF REFERENCES

- 5 [0287] All publications, patents, patent applications and other documents cited in this application are hereby incorporated by reference in their entireties for all purposes to the same extent as if each individual publication, patent, patent application or other document were individually indicated to be incorporated by reference for all purposes. In the event that there is an inconsistency between the teachings of one or more of the references incorporated herein
- 10 and the present disclosure, the teachings of the present specification are intended.

CLAIMS

1. An IL2 proprotein comprising:
- (a) a first polypeptide chain comprising, in N- to C-terminal order:
- 5 (i) a first targeting moiety or component thereof;
- (ii) a first Fc domain;
- (iii) a first linker which is a protease-cleavable linker (PCL) or a non-cleavable linker ("NCL");
- (iv) a first IL2 moiety;
- (v) a second linker which is a protease-cleavable linker (PCL); and
- 10 (vi) a first IL2R α moiety; and
- (b) a second polypeptide chain comprising, in N- to C-terminal order:
- (i) a second targeting moiety or component thereof;
- (ii) a second Fc domain capable of associating with the first Fc domain to form an Fc region;
- 15 (iii) a third linker which is a protease-cleavable linker (PCL) or a non-cleavable linker ("NCL");
- (iv) a second IL2 moiety;
- (v) a fourth linker which is a protease-cleavable linker (PCL); and
- (vi) a second IL2R α moiety.
- 20 2. The IL2 proprotein of claim 1, wherein the IL2 moiety comprises an amino acid sequence having at least about 90% or about 95% sequence identity to mature human IL2.
3. The IL2 proprotein of claim 1 or 2, wherein the IL2 moiety comprises an amino acid sequence having an N-terminal alanine deletion as compared to mature human IL2.
4. The IL2 proprotein of any one of claims 1 to 3, wherein the IL2 moiety comprises
- 25 an amino acid sequence having an amino acid substitution at position N88 as compared to wild type IL2, optionally wherein the amino acid substitution is N88D, and/or (b) an amino acid sequence having the amino acid substitution C125S, C125A or C125V as compared to wild type IL2.

5. The IL2 proprotein of any one of claims 1 to 4, wherein the IL2R α moiety comprises or consists of an amino acid sequence having at least about 90%, at least about 95%, at least about 97%, at least about 98%, at least about 99%, or 100% sequence identity to an IL2 binding portion of human IL2R α .
- 5 6. The IL2 proprotein of claim 5, wherein the IL2R α moiety comprises an amino acid sequence having at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to (a) amino acids 22-186 of IL2R α , (b) amino acids 22-240 of IL2R α , and/or (c) amino acids 22-272 of human IL2R α .
- 10 7. The IL2 proprotein of any one of claims 1 to 6, wherein the second linker, the fourth linker, optionally the first linker, optionally the third linker, or any combination of two or more or all of the foregoing (e.g., (i) first and third linkers, (ii) second and fourth linkers, (iii) first and second linkers, (iv) third and fourth linkers, (v) first, second, third and fourth linkers) comprise(s) (a) a substrate sequence cleavable by any protease set forth in Table A, (b) one or more substrate sequences selected from the substrate sequences set forth in Table B, (c) one or
15 more spacer sequences selected from the substrate sequences set forth in Table C, and/or (d) the amino acid sequence of any of the PCL sequences set forth in Table D or a variant thereof with up to 5 amino acid substitutions, e.g., a variant thereof with 1 amino acid substitution, 2 amino acid substitutions, 3 amino acid substitutions, 4 amino acid substitutions, or 5 amino acid substitutions.
- 20 8. The IL2 proprotein of any one of claims 1 to 7, wherein the first and third linkers are identical and/or the third and fourth linkers are identical.
9. The IL2 proprotein of claim 8, wherein the first, second, third and fourth linkers are identical.
- 25 10. The IL2 proprotein of any one of claims 1 to 8, wherein the first and third linkers are non-cleavable linkers.
11. The IL2 proprotein of claim 10, wherein the non-cleavable linkers comprise or consist of any of the NCL sequences set forth in Table E.

12. The IL2 proprotein of claim 10, wherein the first and third linkers (a) range from 2 to 60 amino acids, from 5 to 25 amino acids, or from 7 to 20 amino acids in length, or (b) are at least 5 amino acids in length.
13. The IL2 proprotein of any one of claims 1 to 8, wherein the first, second, third,
5 and fourth linkers are protease cleavable linkers.
14. The IL2 proprotein of claim 13, wherein the first, second, third, and fourth linkers range from 20 amino acids to 80 amino acids or from 20 amino acids to 60 in length.
15. The IL2 proprotein of any one of claims 1 to 14, wherein the first Fc domain and/or the second Fc domain comprises a hinge domain.
- 10 16. The IL2 proprotein of any one of claims 1 to 15, wherein the first targeting moiety and/or the second targeting moiety is a Fab.
17. The IL2 proprotein of any one of claims 1 to 15, wherein the first targeting moiety and/or the second targeting moiety is an scFv.
18. The IL2 proprotein of any one of claims 1 to 17, wherein the first targeting moiety
15 and/or second targeting moiety is capable of binding to any target molecule identified in Section 6.7.
19. The IL2 proprotein of any one of claims 1 to 18, wherein the first targeting moiety and/or second targeting moiety is capable of binding to an ECM antigen which is optionally selected from syndecan, heparanase, integrins, osteopontin, link, cadherins, laminin, laminin
20 type EGF, lectin, fibronectin, notch, nectin (e.g., nectin-4), tenascin, collagen (e.g., collagen type X) and matrixin.
20. The IL2 proprotein of any one of claims 1 to 19, wherein the first targeting moiety and/or second targeting moiety is capable of binding to a cell surface molecule of tumor or viral lymphocytes.

21. The IL2 proprotein of claim 20, wherein the antigen is a T-cell co-stimulatory protein, optionally selected from CD27, CD28, 4-1BB (CD137), OX40, CD30, CD40, ICOS, lymphocyte function-associated antigen-1 (LFA-1), CD2, CD7, LIGHT, NKG2C, and B7-H3.

22. The IL2 proprotein of any one of claims 1 to 21, wherein the first targeting moiety and/or second targeting moiety is capable of binding to a checkpoint inhibitor, optionally selected from CTLA-4, PD1, PDL1, PDL2, B7-H3, B7-H4, BTLA, HVEM, TIM3, GAL9, LAG3, VISTA, KIR, 2B4, CD160, CGEN-15049, CHK1, and CHK2.

23. The IL2 proprotein of any one of claims 1 to 22, wherein the first targeting moiety and/or second targeting moiety is capable of binding to a tumor-associated antigen ("TAA"), optionally selected from AFP, ALK, a BAGE protein, BIRC5 (survivin), BIRC7, β -catenin, bcr-abl, BRCA1, BORIS, CA9, carbonic anhydrase IX, caspase-8, CALR, CEACAM5 (also known as carcinoembryonic antigen or CEA), CCR5, CD19, CD20 (MS4A1), CD22, CD30, CD40, CDK4, CEA, CTLA4, cyclin-B1, CYP1B1, EGFR, EGFRvIII, ErbB2/Her2, ErbB3, ErbB4, ETV6-AML, EpCAM, EphA2, Fra-1, FOLR1, a GAGE protein (e.g., GAGE-1 or -2), GD2, GD3, GloboH, glypican-3, GM3, gp100, Her2, HLA/B-raf, HLA/k-ras, HLA/MAGE-A3, hTERT, LMP2, MAGE proteins (e.g., MAGE-1, -2, -3, -4, -6, and -12), MART-1, mesothelin, ML-IAP, Muc1, Muc2, Muc3, Muc4, Muc5, Muc16 (CA-125), MUM1, NA17, NY-BR1, NY-BR62, NY-BR85, NY-ESO1, OX40, p15, p53, PAP, PAX3, PAX5, PCTA-1, PLAC1, PRLR, PRAME, PSMA (FOLH1), RAGE proteins, Ras, RGS5, Rho, SART-1, SART-3, STEAP1, STEAP2, TAG-72, TGF- β , TMPRSS2, Thompson-nouvelle antigen (Tn), TRP-1, TRP-2, tyrosinase, and uroplakin-3.

24. The IL2 proprotein of any one of claims 1 to 23, wherein the Fc region is homodimeric.

25. The IL2 proprotein of any one of claims 1 to 23, wherein the Fc region is heterodimeric.

26. An IL2 proprotein according to any one of claims 1 to 25, which has the configuration depicted in FIG. 1A.

27. An IL2 proprotein according to any one of claims 1 to 25, which has the configuration depicted in FIG. 2A.

28. An IL2 proprotein, which is optionally an IL2 proprotein of any one of claims 1 to 27, wherein the IL2 proprotein comprises:

- (a) a first polypeptide chain comprising, in N- to C-terminal order:
 - (i) a first amino acid sequence having at least about 95% sequence identity to any one of SEQ ID NOs:1, 2, 3, 4, 5, and 6;
 - (ii) a second amino acid sequence comprising: (1) one or more sequences set forth in Table B, or (2) a sequence set forth in Table E;
 - (iii) a third amino acid sequence having at least about 95% sequence identity to SEQ ID NO:7;
 - (iv) a fourth amino acid sequence comprising a sequence set forth in Table B; and
 - (v) a fifth amino acid sequence having at least about 95% sequence identity to any one of SEQ ID NOs:8, 9, and 10; and
- (b) a second polypeptide chain comprising, in N- to C-terminal order:
 - (i) a sixth amino acid sequence having at least about 95% sequence identity to any one of SEQ ID NOs:1, 2, 3, 4, 5, and 6;
 - (ii) a seventh amino acid sequence comprising: (1) one or more sequences set forth in Table B, or (2) a sequence set forth in Table E;
 - (iii) an eighth amino acid sequence having at least about 95% sequence identity to SEQ ID NO:7;
 - (iv) a ninth amino acid sequence comprising a sequence set forth in Table B; and
 - (v) a tenth amino acid sequence having at least about 95% sequence identity to SEQ ID NO:8, 9, or 10.

29. The IL2 proprotein of claim 28, wherein the first polypeptide comprises, N-terminal to the first amino acid sequence, an eleventh amino acid sequence having at least about 95%, at least about 98%, or 100% sequence identity to any one of SEQ ID NOs:11, 12, 13, or 14.

30. The IL2 proprotein of claim 28 or 29, wherein the second polypeptide comprises, N-terminal to the first amino acid sequence, a twelfth amino acid sequence having at least about 95%, at least about 98%, or 100% sequence identity to any one of SEQ ID NOs:11, 12, 13, or 14.

5 31. The IL2 proprotein of any one of claims 28 to 30, wherein the first amino acid sequence has at least about 98% or 100% sequence identity to any one of SEQ ID NOs:1, 2, 3, 4, 5, or 6.

32. The IL2 proprotein of any one of claims 28 to 31, wherein the sixth amino acid sequence has at least about 98% or 100% sequence identity to any one of SEQ ID NOs:1, 2, 3,
10 4, 5, or 6.

33. The IL2 proprotein of any one of claims 28 to 32, wherein the second amino acid sequence comprises (a) one or more amino acid sequences set forth in Table B, or (b) an amino acid sequence set forth in Table D.

34. The IL2 proprotein of any one of claims 28 to 32, wherein the second amino acid
15 sequence is an amino acid sequence set forth in Table E.

35. The IL2 proprotein of any one of claims 28 to 34, wherein the second amino acid sequence is 25 or fewer, 15 or fewer, or 6 or fewer amino acids in length.

36. The IL2 proprotein of any one of claims 28 to 35, wherein the seventh amino acid sequence comprises (a) one or more amino acid sequences set forth in Table B, or (b) an
20 amino acid sequence set forth in Table D.

37. The IL2 proprotein of any one of claims 28 to 35, wherein the seventh amino acid sequence is an amino acid sequence set forth in Table E.

38. The IL2 proprotein of claim any one of claims 28 to 37, wherein the seventh amino acid sequence is 25 or fewer, 15 or fewer, or 6 or fewer amino acids in length.

39. The IL2 proprotein of any one of claims 28 to 38, wherein the third amino acid sequence has at least about 98% or 100% sequence identity to SEQ ID NO:7.
40. The IL2 proprotein of any one of claims 28 to 39, wherein the eighth amino acid sequence has at least about 98% or 100% sequence identity to SEQ ID NO:7.
- 5 41. The IL2 proprotein of any one of claims 28 to 40, wherein the fourth amino acid sequence comprises (a) one or more amino acid sequences set forth in Table B or (b) an amino acid sequence set forth in Table D.
42. The IL2 proprotein of any one of claims 28 to 41, wherein the ninth amino acid sequence comprises (a) one or more amino acid sequences set forth in Table B, or (b) an amino acid sequence set forth in Table D.
- 10 43. The IL2 proprotein of any one of claims 28 to 42, wherein the ninth amino acid sequence is an amino acid sequence set forth in Table D.
44. The IL2 proprotein of any one of claims 28 to 43, wherein the fifth amino acid sequence has at least about 98% or 100% sequence identity to SEQ ID NO:8, 9, or 10.
- 15 45. The IL2 proprotein of any one of claims 28 to 44, wherein the tenth amino acid sequence has at least about 98% or 100% sequence identity to SEQ ID NO:8, 9, or 10.
46. The IL2 proprotein of any one of claims 28 to 45, wherein the first polypeptide chain lacks additional sequences C-terminal to the first amino acid sequence.
- 20 47. The IL2 proprotein of any one of claims 28 to 46, wherein the first polypeptide chain (a) lacks additional sequences between the first and second amino acid sequences, (b) lacks additional sequences between the second and third amino acid sequences, (c) lacks additional sequences between the third and fourth amino acid sequences, and/or (d) lacks additional sequences between the fourth and fifth amino acid sequences.
- 25 48. The IL2 proprotein of any one of claims 28 to 47, wherein the second polypeptide chain lacks additional sequences C-terminal to the sixth amino acid sequence.

49. The IL2 proprotein of any one of claims 28 to 48, wherein the second polypeptide chain (a) lacks additional sequences between the sixth and seventh amino acid sequences, (b) lacks additional sequences between the seventh and eighth amino acid sequences, (c) lacks additional sequences between the eighth and ninth amino acid sequences, and/or (d) lacks
5 additional sequences between the ninth and tenth amino acid sequences.

50. A nucleic acid or plurality of nucleic acids encoding the IL2 proprotein of any one of claims 1 to 49.

51. A host cell engineered to express the IL2 proprotein of any one of claims 1 to 49 or the nucleic acid(s) of claim 50.

10 52. A method of producing the IL2 proprotein of any one of claims 1 to 49, comprising culturing the host cell of claim 51 and recovering the IL2 proprotein expressed thereby.

53. A pharmaceutical composition comprising the IL2 proprotein of any one of claims 1 to 49 and an excipient.

15 54. A method of treating cancer, comprising administering to a subject in need thereof the IL2 proprotein of any one of claims 1 to 49 or the pharmaceutical composition of claim 53.

20 55. A method of localized delivery of an IL2 protein, of treating cancer with an IL2 protein that is selectively activated in cancer tissue, of administering to the subject IL2 therapy with reduced systemic exposure and/or reduced systemic toxicity, of targeted delivery of an activated IL2 protein to cancer tissue, or of locally inducing an immune response in a target tissue the method comprising administering to a subject an IL2 proprotein according to any one of claims 1 to 49 (or a pharmaceutical composition comprising the IL2 proprotein and an excipient) which has one or more protease-cleavable linkers, each comprising one or more
25 substrates for one or more proteases expressed by a tissue to which the IL2 protein is to be locally delivered (e.g., a cancer tissue).

56. The method of claim 54 or 55, wherein the administration is non-local.