

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
14 June 2001 (14.06.2001)

PCT

(10) International Publication Number
WO 01/42225 A2(51) International Patent Classification⁷: C07D 239/06, A61K 31/505, A61P 1/04, 19/02

(21) International Application Number: PCT/EP00/11884

(22) International Filing Date:
28 November 2000 (28.11.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/169,089 6 December 1999 (06.12.1999) US(71) Applicant: F. HOFFMANN-LA ROCHE AG [CH/CH];
124 Grenzacherstrasse, CH-4070 Basle (CH).

(72) Inventors: SIDDURI, Achyutharao; 22 Washington Court, Livingston, NJ 07039 (US). TILLEY, Jefferson, Wright; 19 Evergreen Drive, North Caldwell, NJ 07006 (US).

(74) Agent: LOESCHNER, Thomas; 124 Grenzacherstrasse, CH-4070 Basel (CH).


(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW.(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 4-PYRIMIDINYL-N-ACYL-L-PHENYLALANINES

(57) Abstract: Compounds of Formula (I) are disclosed, wherein R¹ to R⁶ are as defined in specification and which are inhibitors of binding between VCAM-1 and cells expressing VLA-4, and accordingly are useful for treating diseases whose symptoms and or damage are related to the binding of VCAM-1 to cells expressing VLA-4.

WO 01/42225 A2

- 1 -

4-PYRIMIDINYL-N-ACYL-L-PHENYLALANINES

5 Vascular cell adhesion molecule-1 (VCAM-1), a member of the
immunoglobulin (Ig) supergene family, is expressed on activated, but not resting,
endothelium. The integrin VLA-4 ($\alpha_4\beta_1$), which is expressed on many cell types
including circulating lymphocytes, eosinophils, basophils, and monocytes, but not
neutrophils, is the principal receptor for VCAM-1. Antibodies to VCAM-1 or VLA-4
10 can block the adhesion of these mononuclear leukocytes, as well as melanoma cells,
to activated endothelium *in vitro*. Antibodies to either protein have been effective at
inhibiting leukocyte infiltration and preventing tissue damage in several animal
models of inflammation. Anti-VLA-4 monoclonal antibodies have been shown to
block T-cell emigration in adjuvant-induced arthritis, prevent eosinophil
15 accumulation and bronchoconstriction in models of asthma, and reduce paralysis
and inhibit monocyte and lymphocyte infiltration in experimental autoimmune
encephalitis (EAE). Anti-VCAM-1 monoclonal antibodies have been shown to
prolong the survival time of cardiac allografts. Recent studies have demonstrated that
anti-VLA-4 mAbs can prevent insulinitis and diabetes in non-obese diabetic mice, and
20 significantly attenuate inflammation in the cotton-top tamarin model of colitis. It has
further been shown that VCAM is expressed on endothelial cells of inflamed colonic
tissue in a TNB/ethanol rat model of inflammatory bowel disease (*Gastroenterology*
1999, 116, 874-883).

25 Thus, compounds which inhibit the interaction between α_4 -containing
integrins and VCAM-1 will be useful as therapeutic agents for the treatment of
chronic inflammatory diseases such as rheumatoid arthritis (RA), multiple sclerosis
(MS), asthma, and inflammatory bowel disease (IBD). The compounds of the present
invention do have this effect.

As used in this specification, the term "halogen" means any of the four halogens, bromine, chlorine, fluorine, and iodine unless indicated otherwise. Preferred halogens are bromine, fluorine, and chlorine.

5 The term "lower alkyl", alone or in combination, means a straight-chain or branched-chain alkyl group containing a maximum of six carbon atoms, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, sec.butyl, isobutyl, tert.butyl, n-pentyl, n-hexyl and the like.

10 The term "Substituted lower alkyl" means a lower alkyl group as defined above which is substituted by one or more groups selected independently from cycloalkyl, nitro, aryloxy, aryl, hydroxy, halogen, cyano, lower alkoxy, lower alkanoyl, lower alkylthio, lower alkyl sulfinyl, lower alkyl sulfonyl, amino and mono or dilower alkyl amino. Examples of substituted lower alkyl groups include 2-hydroxyethyl, 3-
15 oxobutyl, cyanomethyl, and 2-nitropropyl.

 The term "perfluoro lower alkyl" for purposes of R_4 , R_{22} or R_{23} means a substituted lower alkyl group as defined above which is a methyl or ethyl group where all of the hydrogens are substituted by fluoro, i.e. trifluoromethyl and
20 pentafluoroethyl.

 The term "lower alkenyl" means an alkylene group having from 2 to 10 carbon atoms with a double bond located between any two adjacent carbon atoms.

25 The term "cycloalkyl" means an unsubstituted or substituted 3- to 7-membered carbacyclic ring. Substituents useful in accordance with the present invention are hydroxy, halogen, cyano, lower alkoxy, lower alkanoyl, lower alkyl, aroyl, lower alkylthio, lower alkyl sulfinyl, lower alkyl sulfonyl, aryl, heteroaryl and substituted amino.

30 The term "lower alkoxy" means a straight-chain or branched-chain alkoxy group containing a maximum of six carbon atoms, such as methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, tert-butoxy and the like.

5 The term "lower alkylthio" means a lower alkyl group bonded to the rest of the molecule through a divalent sulfur atom, for example, a methyl mercapto or a isopropyl mercapto group. The term "lower alkylsulfinyl" means a lower alkyl group as defined above bound to the rest of the molecule through the sulfur atom in the sulfinyl group. The term "lower alkyl sulfonyl" means a lower alkyl group as defined above bound to the rest of the molecule through the sulfur atom in the sulfonyl group.

10 The term "aryl" means a mono- or bicyclic aromatic group, such as phenyl or naphthyl, which is unsubstituted or substituted by conventional substituent groups. Preferred substituents are lower alkyl, lower alkoxy, hydroxy lower alkyl, hydroxy, hydroxyalkoxy, halogen, lower alkylthio, lower alkylsulfinyl, lower alkylsulfonyl, cyano, nitro, perfluoroalkyl, alkanoyl, aroyl, aryl alkynyl, lower alkynyl and lower
15 alkanoylamino. Examples of aryl groups that may be used in accordance with this invention are phenyl, p-tolyl, p-methoxyphenyl, p-chlorophenyl, m-hydroxy phenyl, m-methylthiophenyl, 2-methyl-5-nitrophenyl, 2,6-dichlorophenyl, 1-naphthyl and the like.

20 The term "arylalkyl" means a lower alkyl group as hereinbefore defined in which one or more hydrogen atoms is/are replaced by an aryl or heteroaryl group as herein defined. Any conventional aralkyl may be used in accordance with this invention, such as benzyl and the like.

25 The term "heteroaryl" means an unsubstituted or substituted 5- or 6-membered monocyclic heteroaromatic ring or a 9- or 10-membered bicyclic heteroaromatic ring containing 1, 2, 3 or 4 heteroatoms which are independently N, S or O. Examples of heteroaryl rings are pyridine, benzimidazole, indole, imidazole, thiophene, isoquinoline, quinzoline and the like. Substituents as defined
30 above for "aryl" are included in the definition of heteroaryl.

The term "lower alkoxycarbonyl" means a lower alkoxy group bonded to the rest of the molecule via a carbonyl group. Examples of alkoxycarbonyl groups are ethoxycarbonyl and the like.

5 The term "lower alkylcarbonyloxy" means a lower alkylcarbonyloxy group bonded to the rest of the molecule via an oxygen atom, for example an acetoxo group. The term "acyloxy" has the same meaning.

10 The term "lower alkanoyl" means a lower alkyl group bonded to the rest of the molecule via a carbonyl group and embraces in the sense of the foregoing definition groups such as acetyl, propionyl and the like. The term "perfluoro lower alkanoyl" means a perfluoro lower alkyl group (a substituted lower alkyl group where all of the hydrogens are substituted by fluoro, preferably trifluoromethyl or penta-

15

The term perfluoro lower alkanoylamino" means a perfluoro lower alkanoyl group bonded to the rest of the molecule via an amino group.


20 The term "lower alkylcarbonylamino" means lower alkylcarbonyl groups bonded to the rest of the molecule via a nitrogen atom, such as acetylamino. The term lower alkylaminocarbonyl" means lower alkyl amino groups bonded to the rest of the molecule via a carbonyl group.

25 The term "arylaminocarbonyl" means aryl groups bonded to an amino group further bonded to the rest of the molecule via a carbonyl group.

30 The term "aroyl" means a mono- or bicyclic aryl or heteroaryl group bonded to the rest of the molecule via a carbonyl group. Examples of aroyl groups are benzoyl, 3-cyanobenzoyl, 2-naphthoyl, nicotinoyl and the like.


Pharmaceutically acceptable salts are well known in the art and can be made by conventional methods taking into account the chemical nature of the compound. Examples of pharmaceutically acceptable salts for acidic compounds are alkali metal or alkaline earth metals such as sodium, potassium, calcium, magnesium, basic amines or basic amino acids, ammonium or alkyl ammonium salts. Particularly desirable salts for compounds of this invention are sodium salts, e.g. from the acidic compound where R_6 is H. The sodium salt of any acid of this invention is easily obtained from the acid by treatment with sodium hydroxide. For basic compounds, examples are salts of inorganic or organic acids such as hydrochloric, hydrobromic, sulphuric, nitric, phosphoric, citric, formic, fumaric, maleic, acetic, succinic, tartaric, methanesulfonic, and p-toluenesulfonic acid.

In one embodiment the present invention relates to a compound of the formula I:



and the pharmaceutically acceptable salts thereof. In accordance with the invention, R_1 is a group Y-1, Y-2 or Y-3 as described below:

R^1 is Y-1, a group of the formula:




wherein:

R_{22} and R_{23} are independently hydrogen, lower alkyl, lower alkoxy, cycloalkyl, aryl, arylalkyl, nitro, cyano, lower alkylthio, lower alkylsulfinyl, lower alkyl sulfonyl, lower alkanoyl, halogen or perfluorolower alkyl and at least one of R_{22} and

R_{23} is other than hydrogen, and R_{24} is hydrogen, lower alkyl, lower alkoxy, aryl, nitro, cyano, lower alkyl sulfonyl, or halogen,

R^1 is Y-2, a five or six membered heteroaromatic ring bonded via a carbon atom to the amide carbonyl wherein said ring contains one, two or three heteroatoms selected from the group consisting of N, O and S and one or two atoms of said ring are independently substituted by lower alkyl, cycloalkyl, halogen, cyano, perfluoroalkyl, or aryl and at least one of said substituted atoms is adjacent to the carbon atom bonded to the amide carbonyl,

R^1 is Y-3, a 3-7 membered ring of the formula:



wherein:

R_{25} is lower alkyl, unsubstituted or fluorine substituted lower alkenyl, or a group of formula $R_{26}-(CH_2)_e-$, R_{26} is aryl, heteroaryl, azido, cyano, hydroxy, lower alkoxy, lower alkoxy carbonyl, lower alkanoyl, lower alkylthio, lower alkyl sulfonyl, lower alkyl sulfinyl, perfluoro lower alkanoyl, nitro, or R_{26} is a group of formula - $NR_{28}R_{29}$, wherein R_{28} is hydrogen or lower alkyl, R_{29} is hydrogen, lower alkyl, lower alkoxy carbonyl, lower alkanoyl, aroyl, perfluoro lower alkanoylamino, lower alkyl sulfonyl, lower alkylaminocarbonyl, arylaminocarbonyl, or R_{28} and R_{29} , taken together with the attached nitrogen atom, form a 4, 5 or 6-membered saturated heterocyclic ring optionally containing one additional heteroatom selected from O, S, and N- R_{40} , Q is $-(CH_2)_fO-$, $-(CH_2)_fS-$, $-(CH_2)_fN(R_{27})-$, $-(CH_2)_f-$, R_{27} is H, lower alkyl, aryl, lower alkanoyl, aroyl or lower alkoxy carbonyl, R_{40} is H, lower alkyl, aryl, lower alkanoyl, aroyl or lower alkoxy carbonyl, the carbon atoms in the ring are unsubstituted or substituted by lower alkyl or halogen, e is an integer from 0 to 4, and f is an integer from 0 to 3;


R_2 is hydrogen or lower alkyl, substituted lower alkyl, arylalkyl, or aryl;

R_3 is hydrogen or lower alkyl, substituted lower alkyl, arylalkyl, or aryl;


R_4 is hydrogen, halogen, lower alkyl, substituted lower alkyl (such as perfluoro lower alkyl), or aryl;

5 R_5 is hydrogen, lower alkyl, chloro, or lower alkoxy;

R_6 is hydrogen, lower alkyl, lower alkylcarbonyloxy lower alkyl, substituted lower alkyl, or R_6 is a group of formula P-3:



10 wherein: R_{32} is hydrogen or lower alkyl, R_{33} is hydrogen, lower alkyl, aryl, R_{34} is hydrogen or lower alkyl, h is an integer from 0 to 2, g is an integer from 0 to 2, the sum of h and g is 1 to 3; or R_6 is a group of formula P-4:



15 wherein: R_{32} , g , and h are as previously defined, Q' is O, S, $\text{---}(\text{CH}_2)_j\text{---}$, or a group of the formula N- R_{35} ; wherein R_{35} is hydrogen, lower alkyl, lower alkanoyl, lower alkoxy, carbonyl, j is 0, 1 or 2

20 The compounds of the invention can exist as stereoisomers and diastereomers, all of which are encompassed within the scope of the present invention. Each of the compounds mentioned in the various embodiments above and below is also contemplated in its pharmaceutically acceptable salt form.

25 In a first particular embodiment of the compounds of formula I
 R^2 is hydrogen, lower alkyl, substituted lower alkyl, arylalkyl, or aryl;
 R^3 is hydrogen, lower alkyl, substituted lower alkyl, arylalkyl, or aryl; and
 R^4 is hydrogen, lower alkyl, perfluoro lower alkyl, or aryl.

In a second particular embodiment of the compounds of formula I
 R^2 is hydrogen, lower alkyl, substituted lower alkyl or aryl;

R^3 is hydrogen, lower alkyl, substituted lower alkyl or aryl; and

R^4 is hydrogen, halogen, lower alkyl, substituted lower alkyl or aryl.

5 In a third particular embodiment of the compounds of formula I and its first and second particular embodiments mentioned before, preferably the first particular embodiment,

R^2 is hydrogen, lower alkyl, substituted lower alkyl or aryl;

R^3 is hydrogen, lower alkyl, substituted lower alkyl, or aryl; and

R^4 is hydrogen, lower alkyl, perfluoro lower alkyl, or aryl.


10

In a preferred embodiment of the compounds of formula I and its three particular embodiments mentioned before, preferably the third one, R^4 is hydrogen, lower alkyl or perfluoro lower alkyl.

15 In a more preferred embodiment of Formula I and its three particular embodiments, especially the first particular one, R_1 is a group of the formula Y-1 as defined in formula I and R_{22} and R_{23} are independently lower alkyl or halogen; and R_{24} is hydrogen. In another more preferred embodiment, R_1 is a group of the formula Y-1 as defined in formula I and R_{22} and R_{23} are independently hydrogen or
20 halogen; and R_{24} is lower alkoxy, preferably methoxy.

Another more preferred embodiment of Formula I and its particular embodiments features R_1 as a group of formula Y-3 as defined in formula I where R_{25} is a group of formula $R_{26}-(CH_2)_e-$, wherein R_{26} is lower alkoxy, Q is $-(CH_2)_f$,
25 e is an integer from 0 to 4, and f is an integer from 0 to 3, preferably 2.

Another more preferred embodiment of Formula I and its particular embodiments is a compound of the formula Ib:



where R_1 is as defined in formula I, R_2 is lower alkyl; R_3 is lower alkyl; R_4 is hydrogen, perfluoro lower alkyl, or lower alkyl, R_5 is hydrogen or lower alkyl; and R_6 is hydrogen, lower alkyl, preferably methyl, lower alkyl-carbonyloxy lower alkyl, preferably 1-(acetoxy)ethyl, a group of formula P-3 as defined in formula I or a group of formula P-4 as defined in formula I. Within this more preferred embodiment, it is particular preferred that R_1 is a group of the formula Y-1 as defined in formula I where R_{22} and R_{23} are independently perfluoro lower alkyl, preferably trifluoromethyl; lower alkyl, preferably methyl, ethyl, propyl or isopropyl; or halogen, preferably fluoro, chloro or bromo; and R_{24} is hydrogen.

When R_1 is a particular preferred group of the formula Y-1 as defined in the preceding paragraph, it is preferred that i) R^2 and R^3 are lower alkyl; R^4 is hydrogen or lower alkyl, and R_5 and R_6 are hydrogen, or ii) R^2 and R^3 are lower alkyl; R^4 is hydrogen or lower alkyl, R_5 is hydrogen, and R_6 is hydrogen, lower alkyl or lower alkylcarbonyloxy lower alkyl, or R_6 is a group of formula P-3 as defined in formula I, or R_6 is a group of formula P-4 as defined in formula I (preferably R^{35} is hydrogen), especially where R_6 is lower alkyl; or R_6 is lower alkylcarbonyloxy lower alkyl; or R_6 is a group of the formula P-3 wherein R^{32} is hydrogen; R^{33} and R^{34} are lower alkyl; h is 1; and g is 0; or R_6 is a group of the formula P-4 wherein R^{32} is hydrogen; h is 1; g is 0; and Q' is O, or iii) R^2 and R^3 are lower alkyl; R^4 is perfluoro lower alkyl, and R^5 and R^6 are hydrogen, or iv) R^2 and R^3 are lower alkyl; R^4 is hydrogen; R^5 is lower alkyl, and R^6 is hydrogen.

Particular compounds in connection with preferred embodiment i) mentioned in the paragraph before are selected from

N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

N-[(2-bromo-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

5 N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

N-[(2-ethyl-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

10 N-[[2-(2-methylethyl)-6-methylphenyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

N-[[2-fluoro-6-(trifluoromethyl)phenyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

15 N-[[2,6-di-(2-methylethyl)phenyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

N-[(2-chloro-6-ethylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

20 N-[(2-chloro-6-propylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

N-[[2-chloro-6-(2-methylethyl)phenyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

25 N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine.

Others compounds are selected from

30 N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

N-[(2-bromo-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine; or

N-[(2-bromo-5-methoxyphenyl)carbonyl]-4-[1,3-dimethyl-2,4-dioxo-5-pyrimidinyl]-L-phenylalanine.

Particular compounds in connection with ii) above where R₅ is lower alkyl are selected from

5 N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine ethyl ester;

N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine ethyl ester; or

10 N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine ethyl ester;

where R₅ is lower alkylcarbonyloxy lower alkyl compounds are selected from

N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxylethyl ester);

15 N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxylethyl ester); or

N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxylethyl ester);

where R₅ is a group of formula P-3 compounds are selected from

20 N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-[(N,N-diethyl)amino]ethyl ester;

N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-[(N,N-diethyl)amino]ethyl ester; or

N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-[(N,N-diethyl)amino]ethyl ester;

25 where R⁶ is a group of formula P-4 compounds are selected from

N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-(4-morpholino)ethyl ester; or

N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-(4-morpholino)ethyl ester.

30

Particular compounds in connection with iii) above are selected from

N-[1-(2,6-dichlorophenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine;

N-[(2-chloro-6-methylphenyl)carbonyl]-4-[1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl]-L-phenylalanine; or

5 N-[[2-fluoro-6-(trifluoromethyl)phenyl]carbonyl]-4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine.

Particular compounds in connection with iv) above are selected from


10 N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine;

N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine; or

N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine.

15

In compounds of this invention of the formula Ib defined above,



where R_1 is as defined in formula I, R_2 is lower alkyl; R_3 is lower alkyl; R_4 is hydrogen, perfluoro lower alkyl, or lower alkyl, R_5 is hydrogen or lower alkyl; and R_6 is hydrogen, lower alkyl, lower alkylcarbonyloxy lower alkyl, a group of formula P-3 as defined in formula I or a group of formula P-4 as defined in formula I, it is also preferred that R_1 is a group of formula Y-3 as defined in formula I, where R_{25} is a group of formula $R_{26}-(CH_2)_e-$, wherein R_{26} is lower alkoxy, Q is $-(CH_2)_f$, e is an integer from 0 to 4, and f is an integer from 0 to 3. In such
20
25 compounds, it is preferred that R^2 and R^3 are lower alkyl, R^4 is hydrogen or lower alkyl; and R^5 and R^6 are hydrogen.


Particular compounds in connection with this are

4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-N-[[1-(2-methoxyethyl)cyclopentyl]carbonyl]-L-phenylalanine;

5 N-[[1-(2-methoxyethyl)cyclopentyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine; or

4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-N-[[1-(2-methoxyethyl)cyclopentyl]carbonyl]-L-phenylalanine.

In another group of compounds of this invention of the formula Ib:



10 where R_1 is as defined in formula I, R_2 is lower alkyl; R_3 is lower alkyl; R_4 is hydrogen, perfluoro lower alkyl, or lower alkyl, R_5 is hydrogen or lower alkyl; and R_6 is hydrogen, lower alkyl, lower alkylcarbonyloxy lower alkyl, a group of formula P-3 as defined in formula I or a group of formula P-4 as defined in formula I, it is particularly preferred that R_2 and R_3 are lower alkyl; and R_4 , R_5 and R_6 are hydrogen, especially where R_1 is a group of the formula Y-1 as defined in formula I, preferably where R_{22} and R_{23} are independently lower alkyl or halogen; and R_{24} is hydrogen or where R_{22} and R_{23} are independently hydrogen or halogen; and R_{24} is lower alkoxy.

20 In the compound of the preceding paragraph where R_2 and R_3 are lower alkyl; and R_4 , R_5 and R_6 are hydrogen it is also preferred that R_1 is a five or six membered heteroaromatic ring bonded via a carbon atom to the amide carbonyl wherein said ring contains one, two or three heteroatoms selected from the group consisting of N, O and S and one or two atoms of said ring are independently substituted by lower alkyl, cycloalkyl, halogen, cyano, perfluoroalkyl, or aryl and at least one of said
25 substituted atoms is adjacent to the carbon atom bonded to the amide carbonyl.

In the compound of the penultimate paragraph where R_2 and R_3 are lower alkyl; and R_4 , R_5 and R_6 are hydrogen it is also preferred that R_1 is a group of formula the Y-3 as defined in formula I, preferably where R_{25} is a group of formula $R_{26}-(CH_2)_e$, wherein R_{26} is lower alkoxy, Q is $-(CH_2)_f$, e is an integer from 0 to 4, and f is an integer from 0 to 3, preferably 2.

In another embodiment of the embodiment of the compounds of formula I, where R^2 is hydrogen, lower alkyl, substituted lower alkyl or aryl;

R^3 is hydrogen, lower alkyl, substituted lower alkyl or aryl; and

R^4 is hydrogen, halogen, lower alkyl, substituted lower alkyl or aryl, i.e. the second particular embodiment,

R^1 is Y-1 wherein R_{22} and R_{23} are as defined in formula I. Preferably R_{22} and R_{23} are independently lower alkyl or halogen; and R_{24} is hydrogen, or R_{22} and R_{23} are independently hydrogen or halogen; and R_{24} is lower alkoxy.

In another embodiment of the second particular embodiment,

R^1 is Y-2. In a further embodiment of the second particular embodiment, R^1 is Y-3, especially wherein R_{25} is a group of formula $R_{26}-(CH_2)_e$, wherein R_{26} is lower alkoxy, Q is $-(CH_2)_f$, e is an integer from 0 to 4, and f is an integer from 0 to 3, preferably 2.

In a more specific embodiment of the second particular embodiment

R^2 is lower alkyl;


R^3 is lower alkyl;

R^4 is hydrogen, lower alkyl or halogen;


R^5 is hydrogen and R^6 is hydrogen.

A preferred embodiment within this more specific embodiment is wherein R^4 is hydrogen. Within this preferred embodiment R_1 is Y-1 as defined in formula I, especially wherein R_{22} and R_{23} are independently lower alkyl or halogen; and R_{24} is hydrogen or wherein R_{22} and R_{23} are independently hydrogen or halogen; and R_{24} is lower alkoxy. With in this preferred embodiment R_1 is also optionally Y-2 as defined in formula I or R_1 is optionally also Y-3 as defined in formula I, especially wherein in Y-3 R_{25} is a group of formula $R_{26}-(CH_2)_e$, wherein R_{26} is lower alkoxy, Q is $-(CH_2)_f$, e is an integer from 0 to 4, and f is an integer from 0 to 3, preferably 2.

Additionally, a preferred embodiment of the present invention is a compound of the formula I:




5 wherein R_1 is a group of the formula Y-1



Y-1

10 wherein R_{22} and R_{23} are independently hydrogen, lower alkyl, lower alkoxy, cycloalkyl, aryl, arylalkyl, nitro, cyano, lower alkylthio, lower alkylsulfinyl, lower alkyl sulfonyl, lower alkanoyl, halogen, or perfluorolower alkyl and at least one of R_{22} and R_{23} is other than hydrogen; and R_{24} is hydrogen, lower alkyl, lower alkoxy, aryl, nitro, cyano, lower alkyl sulfonyl, or halogen; R_2 is lower alkyl; R_3 is lower alkyl; R_4 is hydrogen, or lower alkyl; R_5 is hydrogen; and R_6 is hydrogen.

15 A more preferred embodiment within the above embodiment of the present invention is a compound of formula I above wherein R_1 is a group of the formula Y-1




Y-1

20 wherein R_{22} and R_{23} are independently hydrogen, lower alkyl or halogen, R_{24} is hydrogen or lower alkoxy; R_2 is lower alkyl; R_3 is lower alkyl; R_4 is hydrogen, or lower

alkyl; R_5 is hydrogen; and R_6 is lower alkyl, lower alkylcarbonyloxy lower alkyl, or preferably hydrogen.

Another preferred embodiment of the present invention is a compound of Formula I above wherein R_1 is a five or six membered heteroaromatic ring bonded via a carbon atom to the amide carbonyl wherein said ring contains one, two or three heteroatoms selected from the group consisting of N, O and S and one or two atoms of said ring are independently substituted by lower alkyl, cycloalkyl, halogen, cyano, perfluoroalkyl, or aryl and at least one of said substituted atoms is adjacent to the carbon atom bonded to the amide carbonyl; R_2 is lower alkyl; R_3 is lower alkyl; R_4 is hydrogen, perfluoro lower alkyl, or lower alkyl; R_5 is hydrogen; and R_6 is hydrogen.


Another preferred embodiment of the present invention is a compound of Formula I above wherein R_1 is a group of formula Y-3 which is a 3-7 membered ring of the formula:



wherein R_{25} is lower alkyl, unsubstituted or fluorine substituted lower alkenyl, or a group of formula $R_{26}-(CH_2)_e$, R_{26} is aryl, heteroaryl, azido, cyano, hydroxy, lower alkoxy, lower alkoxycarbonyl, lower alkanoyl, lower alkylthio, lower alkyl sulfonyl, lower alkyl sulfinyl, perfluoro lower alkanoyl, nitro, or R_{26} is a group of formula - $NR_{28}R_{29}$, wherein R_{28} is hydrogen or lower alkyl, R_{29} is hydrogen, lower alkyl, lower alkoxycarbonyl, lower alkanoyl, aroyl, perfluoro lower alkanoylamino, lower alkyl sulfonyl, lower alkylaminocarbonyl, arylaminocarbonyl; or R_{28} and R_{29} , taken together with the attached nitrogen atom, form a 4, 5 or 6-membered saturated heterocyclic ring optionally containing one additional heteroatom selected from O, S, and N- R_{40} ; Q is $-(CH_2)_fO-$, $-(CH_2)_fS-$, $-(CH_2)_fN(R_{27})-$, $-(CH_2)_f$, R_{27} is H, lower alkyl, aryl, lower alkanoyl, aroyl or lower alkoxycarbonyl, R_{40} is H, lower alkyl, aryl, lower alkanoyl, aroyl or lower alkoxycarbonyl; the carbon atoms in the ring are unsubstituted or substituted by lower alkyl or halogen, e is an integer from 0 to 4, and


f is an integer from 0 to 3; R₂ is lower alkyl; R₃ is lower alkyl; R₄ is hydrogen, perfluoro lower alkyl, or lower alkyl; R₅ is hydrogen; and R₆ is hydrogen.

A more preferred embodiment of the present invention is a compound of formula I above wherein R₁ is a group of formula Y-3 which is a 3-7 membered ring of the formula:



wherein R₂₅ is a group of formula R₂₆-(CH₂)_e, wherein R₂₆ is lower alkoxy, Q is -(CH₂)_f, e is an integer from 0 to 4, and f is an integer from 0 to 3; R₂ is lower alkyl; R₃ is lower alkyl; R₄ is hydrogen, perfluoro lower alkyl, or lower alkyl; R₅ is hydrogen; and R₆ is hydrogen.

In another aspect the present invention relates to compounds of formula II



wherein R²-R⁵ are as defined in formula I and P₁ and P₂ each are a protecting group. More particular, P₁ is a standard nitrogen protecting group such as tert.butyloxy-carbonyl (Boc) or the carbobenzyloxy group and P₂ is a standard carboxy protecting group such as lower alkyl or substituted lower alkyl. These compounds are useful intermediates in the preparation of the compounds of formula I.

The compounds of the invention inhibit the binding of VCAM-1 and fibronectin to VLA-4 on circulating lymphocytes, eosinophils, basophils, and monocytes ("VLA-4-expressing cells"). The binding of VCAM-1 and fibronectin to VLA-4 on such cells is known to be implicated in certain disease states, such as

rheumatoid arthritis, multiple sclerosis, inflammatory bowel disease, and particularly in the binding of eosinophils to airway endothelium which contributes to the cause of the lung inflammation which occurs in asthma. Thus, the compounds of the present invention are useful for the treatment of asthma.

5

On the basis of their capability of inhibiting binding of VCAM-1 and fibronectin to VLA-4 on circulating lymphocytes, eosinophils, basophils, and monocytes, the compounds of the invention can be used as a medicament or as a pharmaceutical composition especially for the treatment of disorders which are known to be associated with or mediated by such binding. Examples of such disorders are rheumatoid arthritis, multiple sclerosis, asthma, and inflammatory bowel disease. The compounds of the invention are preferably used in the treatment of diseases which involve pulmonary inflammation, such as asthma. The pulmonary inflammation which occurs in asthma is related to the activation and lung infiltration of eosinophils, monocytes and lymphocytes which have been activated by some asthma-triggering event or substance.

10

15

Furthermore, compounds of the invention also inhibit the binding of VCAM-1 and MadCAM to the cellular receptor alpha4-beta7, also known as LPAM, which is expressed on lymphocytes, eosinophiles and T-cells. While the precise role of alpha4-beta7 interaction with various ligands in inflammatory conditions such as asthma is not completely understood, compounds of the invention which inhibit both alpha4-beta1 and alpha4-beta7 receptor binding are particularly effective in animal models of asthma. Furthermore work with monoclonal antibodies to alpha4-beta7 indicate that compounds which inhibit alpha4-beta7 binding to MadCAM or VCAM are useful for the treatment of inflammatory bowel disease. They would also be useful in the treatment of other diseases in which such binding is implicated as a cause of disease damage or symptoms.

20

25

The compounds of the invention can be administered orally, rectally, or parentally, e.g., intravenously, intramuscularly, subcutaneously, intrathecally or transdermally; or sublingually, or as ophthalmological preparations, or as an aerosol in the case of pulmonary inflammation. Capsules, tablets, suspensions or solutions for

30

oral administration, suppositories, injection solutions, eye drops, salves or spray solutions are examples of administration forms.

Intravenous, intramuscular, oral or inhalation administration is a preferred form of use. The dosages in which the compounds of the invention are administered in effective amounts depending on the nature of the specific active ingredient, the age and the requirements of the patient and the mode of administration. Dosages may be determined by any conventional means, e.g., by dose-limiting clinical trials. Thus, the invention further comprises a method of treating a host suffering from a disease in which VCAM-1 or fibronectin binding to VLA-4-expressing cells is a causative factor in the disease symptoms or damage by administering an amount of a compound of the invention sufficient to inhibit VCAM-1 or fibronectin binding to VLA-4-expressing cells so that said symptoms or said damage is reduced. In general, dosages of about 0.1-100 mg/kg body weight per day are preferred, with dosages of 1-25 mg/kg per day being particularly preferred, and dosages of 1-10 mg/kg body weight per day being especially preferred.

In another aspect, the invention relates to the use of a compound of the present invention or a pharmaceutically salt thereof for the preparation of a medicament for the treatment of disease states mediated by the binding of VCAM-1 or VLA-4 or VLA-4-expressing cells, especially in the treatment of rheumatoid arthritis, multiple sclerosis, inflammatory bowel disease and asthma.

In another aspect the invention further relates to medicaments or pharmaceutical compositions comprising a pharmaceutically effective amount of a compound of the invention and a pharmaceutically acceptable carrier. It also relates to a process for the preparation of a pharmaceutical composition which process comprises bringing a compound of the present invention or a pharmaceutically acceptable salt thereof, and a compatible pharmaceutical carrier into a galenical administration form. Such compositions may be formulated by any conventional means. Tablets or granulates can contain a series of binders, fillers, carriers or

diluents. Liquid compositions can be, for example, in the form of a sterile water-miscible solution. Capsules can contain a filler or thickener in addition to the active ingredient. Furthermore, flavour-improving additives as well as substances usually used as preserving, stabilizing, moisture-retaining and emulsifying agents as well as
5 salts for varying the osmotic pressure, buffers and other additives can also be present.

The previously mentioned carrier materials and diluents can comprise any conventional pharmaceutically acceptable organic or inorganic substances, e.g.,

water, gelatine, lactose, starch, magnesium stearate, talc, gum arabic, polyalkylene glycols and the like.


Oral unit dosage forms, such as tablets and capsules, preferably contain from
5 25 mg to 1000 mg of a compound of the invention.

Generally the compounds of the present invention can be prepared from suitable phenylalanine derivatives via a palladium catalyzed reaction with a 5-halo-2,4-dioxypyrimidone.

10


As shown in Reaction Scheme 1, a 4-iodo- or 4-bromophenylalanine derivative such as 1, is converted into a protected phenylalanine derivative 2 in which R_3' is hydrogen, chloro, lower alkyl or lower alkoxy, P_1 is a standard nitrogen protecting group such as a Boc, or carbobenzyloxy group and P_2 is lower alkyl or substituted
15 lower alkyl selected appropriately to serve as a protecting group or an element of a prodrug. The group P_2 can be introduced by conventional means familiar to those who practice peptide chemistry. The order of the addition of P_1 and P_2 is not critical and will depend on the particular choice of reagents. A discussion of the use and introduction of protecting groups is provided in Theodora W. Greene and Peter G.
20 M. Wuts., *Protecting Groups in Organic Synthesis*, Wiley Interscience, New York, 1991. Alternatively, a compound of formula 1 may be converted to a compound of formula 4, in which R_1' represents a component of an acyl group of the invention. A convenient method is to introduce the ester group P_2 first, followed by a coupling reaction of the free amine using conventional peptide coupling reagents, for example
25 HBTU in the presence of a tertiary amine base such as diethylisopropylamine. Again, the particular choice of reagents may dictate altering the sequence of the introduction of R_1' and P_2 . Conversion of compounds of formula 2 or 4 to derivatives 3 or 5, in which M represents a substituted tin or boron atom, can be effected by treatment with a suitable species, for example hexamethylditin, hexabutylditin or a
30 tetraalkoxydiboron in the presence of a source of palladium zero. The methodology is outlined and referenced in F. Diederich and P. J. Stang, ed, *Metal Catalyzed Cross Coupling Reactions*, Wiley-VCH, Weinheim, Germany, 1998.

Reaction Scheme 1.



Pyrimidine-2,4-diones (uracil derivatives) of formula 6 wherein R_4 is hydrogen, lower alkyl or perfluorolower alkyl are well known in the literature or can be made by known methods. 1,3-Disubstituted pyrimidin-2,4-diones of formula 7 wherein R_2 and R_3 are lower alkyl, aryl lower alkyl or aryl are also known compounds or can be prepared by standard procedures. Papers reporting synthetic methods for their construction include: Shigeo Senda, et al. Chem. Pharm. Bull. 1974, 22, 189-195, Chem Pharm Bull 1972, 20, 1389-1396, and Yasuo Morita, et al., Chem Comm. 1997 359-360. For the case where R_2' and R_3' are lower alkyl or aryl lower alkyl, compounds of formula 7 are available by alkylation of compounds of formula 6 by treatment with an alkylating agents such as iodomethane, benzylbromide, allyl bromide in the presence of a base such as potassium carbonate and optionally, a phase transfer catalyst. For less reactive alkylating agents, it may be necessary to use a stronger base such as an alkali metal hydroxide and to heat the reaction mixture. Compounds of formula 6 or 7 as defined above may be halogenated in the 5-position by treatment with conventional halogenating reagents such as bromine, N-iodosuccinimide or N-bromosuccinimide in a suitable solvent such as glacial acetic acid or aqueous acetic acid to give halopyrimidines of formula 8, $X = \text{Br}$ or I , R_2 and R_3 are independently hydrogen, lower alkyl, aryl lower alkyl or aryl, $R_4 =$ hydrogen, lower alkyl or perfluorolower alkyl.


Reaction Scheme 2.



As shown in Reaction Scheme 3, the compound of formula 8 can be used in a palladium catalyzed coupling reaction with a phenylalanine derivative of formula 3 or 5. For example, when M is a substituted tin, treatment of a mixture of 8 and the phenylalanine of formula 3 or 5 with a source of palladium zero such as tetraakis(triphenylphosphine)palladium or bis(triphenylphosphine)palladium dichloride in the presence of an inert solvent such as DMF at a temperature of between room temperature and 100 °C gives a compound of formula 9 or 10. Compounds of structure 9 may be converted into compounds of structure 10 by removal of the protecting group P_1 , which may be accomplished by conventional means depending on the selection of P_1 . For example, if P_1 is a Boc group, it may be removed by treatment with a strong acid, such as trifluoroacetic acid, optionally in the presence of a solvent such as dichloromethane and a scavenging agent. The resulting free amine may then be acylated with an acid of the formula $R_1'CO_2H$ using conventional peptide coupling techniques. For example, by treatment with HBTU in the presence of a tertiary amine base such as diethylisopropylamine in the presence of an aprotic solvent such as DMF to give the compound of structure 10.

If the free acid 11 is the desired end product, the ester group, P_2 may be removed by conventional means. For example, in the case that P_2 is lower alkyl, for example methyl, it may be removed by treatment with an alkali metal hydroxide, for example lithium hydroxide, in a suitable solvent, for example aqueous THF optionally containing methanol to assist with solubility. If P_2 were a benzyl or substituted benzyl group, it could also be removed by catalytic hydrogenation over a noble metal catalyst, for example palladium on carbon.


Reaction Scheme 3.



- 5 Alternatively, as shown in Reaction Scheme 4, a compound of structure 8, wherein X is bromide or iodide, may be converted to a species of formula 12, in which M' represents a substituted tin, boron or zinc atom. In the case of the tin or boron

derivatives, in which M' represents a substituted tin or boron atom, the conversion can be effected by treatment with a suitable species, for

Reaction Scheme 4.




example hexamethylditin, hexabutylditin or a tetraalkoxydiboron in the presence of a source of palladium zero. For the formation of the zinc derivative, 12, M' = Zn(halogen), conversion may be effected by treatment of the compound of formula 8, X = I with a source of activated zinc metal in a suitable inert solvent, for example dimethylacetamide at a temperature of from room temperature to 100 °C until conversion is complete to give a compound of formula 12, M' = Zn(halogen). These

compounds of formula 12 can be reacted with a 4-substituted phenylalanine derivative of formula 4' in which X' is iodo, bromo, or trifluoromethylsulfonyloxy in the presence of a source of palladium zero to give a compound of formula 10'. In the case where the ester group represented by P₂ is not part of the targeted compound, it can be removed using ester hydrolysis procedures appropriate to the particular P₂. For example, where P₂ is lower alkyl, for example methyl, it can be removed by standard base hydrolysis using an alkali metal hydroxide, for example, lithium hydroxide. In a variation on this procedure, it may be desirable to carry a protecting group through the coupling reaction and substitute it at a later time. In this case, a compound of formula 2', in which P₁' is lower alkoxycarbonyl or benzyloxycarbonyl and X' is as defined above, may be coupled with a pyrimidinedione of structure 12 to give a compound of structure 9' which in turn may be converted to a compound of the invention using the general procedures noted above in reaction scheme 3.

An alternative route to compounds of this invention, as shown in Reaction Scheme 5, which is particularly applicable to compounds in which R₅' is other than hydrogen, is to build an aldehyde of formula 14. This can be accomplished by reacting a compound of formula 12 with a compound of formula 13, in which R₅' represents lower alkyl or lower alkoxy, and X'' represents an iodide, bromide, of trifluoromethylsulfonyloxy moiety and R₈ represents a protected alcohol or aldehyde. For alcohols, suitable protecting groups include silyl ethers, benzyl ethers. Aldehydes, may be protected as their acetal derivatives. The compound of formula 12 can be converted to an aldehyde of formula 15 by conventional steps which, when R₈ is an alcohol, would involve protecting group removal, if necessary, followed by oxidation. Any of the common reagents for the selective oxidation of primary benzyl alcohols to aldehydes may be employed, for example, treatment with activated manganese dioxide in an inert solvent. In the case where R₈ represents a protected aldehyde, conversion to an aldehyde of formula 15 can be carried out by a suitable protecting group removal, for example hydrolysis of an acetal with dilute acid. Reaction of 15 to give a dehydroamino acid of formula 16 can be effected by treatment with a Wittig reagent of formula 17 in which P₁' is lower alkoxycarbonyl or benzyloxycarbonyl and P₂ is as defined above. For example treatment of 15 with (±)-N-(benzyloxycarbonyl)-

α -phosphonoglycine trimethyl ester in the presence of a suitable base for example tetramethylguanidine leads directly to a dehydroamino acid of formula 16, $P_2 =$ methyl and $P_1' =$ benzyloxycarbonyl. Enantioselective reduction of 16 to the L-amino acid 18 can be effected by use of a number of reducing agents suitable for the purpose, for example, the recently described ethyl-DuPHOS rhodium reagent (Burk, M. J., Feaster, J. E.; Nugent, W. A.; Harlow, R. L. *J. Am. Chem. Soc.* 1993, 115, 10125) using essentially the literature procedure. Further conversion of 18 to the compounds of the invention can be carried out using the general procedures discussed above.


Reaction Scheme 5.



In one embodiment, the N-acyl group, R_1' of structure 11, is derived from a 2-substituted benzoic acid. Appropriate 2-substituted benzoic acids are either commercially available or can be prepared by conventional means. For example ortho-substituted aryl iodides or triflates may be carbonylated in the presence of

carbon monoxide and a suitable palladium catalyst. The preparation of such iodide or triflate intermediates is dependent on the particular substitution pattern desired and they may be obtained by direct iodination or diazotization of an aniline followed by treatment with a source of iodide for example, potassium iodide. Triflates may be derived from the corresponding phenols by conventional means such as by treatment with trifluoromethane sulfonic anhydride in the presence of a base such as triethylamine or diisopropylethylamine in an inert solvent. As shown in Reaction Scheme 6, one other means of obtaining ortho-substituted benzoic acids involves treatment of an 2-methoxyphenyloxazoline derivative such as compound 19, Z₁ and Z₂ = hydrogen, alkyl, chloro, perfluoroalkyl, lower alkoxy with an alkyl Grignard reagent followed by hydrolysis of the oxazoline ring following the general procedure described by Meyers, A. I., Gabel, R., Mihelick, E. D, *J. Org. Chem.* 1978, 43, 1372-1379, to give an acid of formula 20. 2- or 2,6-Disubstituted benzonitriles also serve as convenient precursors to the corresponding benzoic acids. In the case of highly hindered nitriles, for example 2-chloro-6-methylbenzonitrile, conventional hydrolysis under acidic or basic conditions is difficult and better results are obtained by DIBAL reduction to the corresponding benzaldehyde followed by oxidation using a chromium based oxidizing reagent. Other methods are exemplified in Chen, et al., WO 99/10312.

Reaction Scheme 6



Referring now to Reaction Scheme 7, cyclic acids of formula 23 are known compounds or can be prepared using standard methodologies. For the preparation of substituted alkyl- or cycloalkylcarboxylic acids, alkylation reactions can be employed using an alkali metal dianion of the acid or monoanion of the corresponding ester.

For example, a cycloalkyl carboxylic acid ester of formula 21 can be treated with a strong base, for example, lithium diisopropylamide in an inert solvent, for example THF followed by addition of group R_{41} -Lv wherein R_{41} represents a desired side chain, such as a substituted benzyl, lower alkyl, lower alkoxy alkyl, azidolower alkyl and the like and Lv represents a leaving group such as a bromide, iodide, mesylate or similar group known to participate in ester enolate alkylation reactions. The product ester 22 may be hydrolyzed to the acid 23 using alkali metal hydroxide in a suitable solvent, for example aqueous alcohol. Depending on the nature of R_{41} and the eventual target, the compound 23 may be coupled to an amine such as compound 1 and converted to the target directly or R_{41} may be subject to further manipulation at a suitable point in the synthesis. For example, if R_{41} is an azido lower alkyl moiety, the azide may be reduced using for example a trialkyl phosphine reagent followed by functionalization of the product amine by alkylation, acylation, sulfonylation and related procedures well known to those skilled in the art. If R_{41} incorporates a leaving group, for example, a terminal bromine atom, this group may be displaced by an appropriate nucleophile, for example, sodium methyl mercaptide to give in this case, a thioether which may be the desired product or can be itself further manipulated, for example by oxidation to a sulfoxide or sulfone using standard reaction conditions. Other nucleophiles which may be employed to produce intermediates leading to compounds of this invention include: sodium cyanide, sodium methoxide, sodium azide, morpholine and others. When R_{41} incorporates a ketal group, this group may be hydrolyzed at a convenient point in the synthesis to provide a keto group. This group in turn may be further manipulated, for example by reduction to an alcohol or conversion to derivative such as an oxime.

Examples of the application of these methods to the synthesis of compounds of formula 23 are provided in Chen, et al. WO 99/10313.

5

10

20

25

esterification. Alternatively, a diimide mediated coupling between the carboxylic acid and an alcohol may be employed with the optional use of a promoter such as 4,4-dimethylaminopyridine. A typical diimide is dicyclohexylcarbodiimide. Another alternative is to treat the carboxylic acid with a reactive alkyl halide, for example, an alkyl iodide or an acyloxymethyl chloride in the presence of a base, for example sodium bicarbonate and an inert solvent, for example DMF. The particular choice of method will be determined by the nature of the particular combination of carboxylic acid and desired ester moiety and will be apparent to one skilled in the art. Ester groups which may constitute prodrugs may be introduced at any convenient point in the synthesis. For example the group P₂ in formula 1 may represent a desirable prodrug ester and be retained in the final product.

EXAMPLES

The Examples which follow are for purposes of illustration and are not intended to limit the invention in any way.

General Methods: Melting points were taken on a Thomas-Hoover apparatus and are uncorrected. Optical rotations were determined with a Perkin-Elmer model 241 polarimeter. ¹H-NMR spectra were recorded with Varian XL-200, Mercury-300 or Unityplus 400 MHz spectrometers, using tetramethylsilane (TMS) as internal standard. Electron impact (EI, 70 ev) and fast atom bombardment (FAB) mass spectra were taken on VG Autospec or VG 70E-HF mass spectrometers. Silica gel used for column chromatography was Mallinkrodt SiliCar 230-400 mesh silica gel for flash chromatography; columns were run under a 0-5 psi head of nitrogen to assist flow. Thin layer chromatograms were run on glass thin layer plates coated with silica gel as supplied by E. Merck (E. Merck # 1.05719) and were visualized by viewing under 254 nm UV light in a view box, by exposure to I₂ vapor, or by spraying with either phosphomolybdic acid (PMA) in aqueous ethanol, or after exposure to Cl₂, with a 4,4'-tetramethyldiaminodiphenylmethane reagent prepared according to E. Von Arx, M. Faupel and M Brugger, *J. Chromatography*, 1976, 120, 224-228.

Reversed phase high pressure liquid chromatography (RP-HPLC) was carried out using a Rainin HPLC employing a 41.4 x 300 mm, 8 μ M, Dynamax™ C-18 column at a flow of 49 mL/min employing a gradient of acetonitrile:water (each containing 0.75% TFA) typically from 5 to 95% acetonitrile over 35-40 min. HPLC conditions are typically described in the format (5-95-35-214); this refers to a linear gradient of from 5% to 95% acetonitrile in water over 35 min while monitoring the effluent with a UV detector at a wavelength of 214 nM.

Methylene chloride (dichloromethane), 2-propanol, DMF, THF, toluene, hexane, ether, and methanol, were Fisher or Baker reagent grade and were used without additional purification except as noted, acetonitrile was Fisher or Baker hplc grade and was used as is.

Definitions as used herein:

THF is tetrahydrofuran,
DMF is N,N-dimethylformamide,
DMA is N,N-dimethylacetamide
HOBT is 1-hydroxybenzotriazole,
BOP is [(benzotriazole-1-yl)oxy]tris-(dimethylamino)phosphonium
hexafluorophosphate,
HATU is O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium
hexafluorophosphate
HBTU is O-benzotriazole-N,N,N',N'-tetramethyluronium hexafluorophosphate,
DIPEA is diisopropylethylamine,
DMAP is 4-(N,N-dimethylamino)pyridine
DPPA is diphenylphosphoryl azide
DPPP is 1,3-bis(diphenylphosphino)propane
DBU is 1,8-diazabicyclo[5.4.0]undec-7-ene
NaH is sodium hydride
brine is saturated aqueous sodium chloride solution
TLC is thin layer chromatography
LDA is lithium diisopropylamide

BOP-Cl is bis(2-oxo-3-oxazolidinyl)phosphinic chloride

NMP is N-methylpyrrolidinone


Lawesson's reagent is [2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide]

5 NIS is N-iodosuccinic anhydride.

Silica gel chromatography on Biotage columns refers to use of a flash chromatography system supplied by the Biotage Division of the Dyax Corporation employing prepacked 40g (40s columns), 90g (40m columns) or 800g (75m columns). Elution is carried out with hexane-ethyl acetate mixtures under 10-15 psi nitrogen pressure.

Example 1.

N-[(1,1-dimethylethoxy)carbonyl]-4-[(tributyl)stannyl]-L-phenylalanine methyl ester.



15 A solution of N-[(1,1-dimethylethoxy)carbonyl]-4-iodo-L-phenylalanine methyl ester (5.3 g, 13 mmol) and hexabutylditin (27.5 mL, 54 mmol) in toluene (50 mL) was deoxygenated by alternately freezing the mixture in a liquid nitrogen bath under vacuum and thawing under argon (3 x). Tetrakis(triphenylphosphine)palladium was added (280 mg, 0.22 mmol) and the reaction mixture was heated to reflux for 45 min as the color changed from yellow to black. TLC (1:6 ethyl acetate:hexane) indicated the presence of some starting iodide and an additional portion (140 mg, 0.11 mmol) of the catalyst was added. Reflux was continued for 1 hr. The mixture was allowed to cool and was concentrated. The residue was taken up in hexane (200 mL) and triethylamine (30 mL), stirred for 30 min and was filtered. The filtrate was concentrated and was chromatographed over a dry silica gel column containing 150 g


of silica gel and eluting with hexane followed by 1:6 ethyl acetate:hexanes to give N-[(1,1-dimethylethoxy)carbonyl]-4-[(tributyl)stannyl]-L-phenylalanine methyl ester (5.7 g, 77%) as a clear oil. LR(+)-LSIMS (C₂₇H₄₇NO₄Sn): m/z 1081 (2M-C₄H₉) 570 (M+H).

5

Example 2.


N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine

10 a) Preparation of 5-iodo uracil




A mixture of uracil (28.6 mmol, 3.2 g) and iodine monochloride (49.2 mmol, 7.988 g) in methanol (120 mL) was refluxed for 40 hr. The solvent was removed under vacuum and the residue was crystallized from ethanol:water (1:1, 150 mL) and stored in the refrigerator for 3 h. The resulting needles were collected by filtration and were washed with ethanol:water (1:1 mixture, 50 mL), water (30 mL), hexane (30 mL) and then dried in air to obtain 5.47 g (80%) of 5-iodo uracil as a white needles (mp 278-282 °C, Lit. 274-276 °C, *Synthetic communication* 1988, 18, 855-867). EI-HRMS m/e calcd for C₄H₃IN₂O₂ (M⁺) 237.9239, found 237.9244.

b) Preparation of 1,3-dimethyl-5-iodo uracil



A mixture of 5-iodo uracil (22.2 mmol, 5.28 g) and powdered potassium carbonate (60 mmol, 10.3 g) in DMF (188 mL) was stirred for 24 h at room temperature and then methyl iodide (53.5 mmol, 3.33 mL) was added. Then, the reaction mixture was stirred for another 72 h at room temperature and was poured into water (150 mL) and ethyl acetate (150 mL). The layers were separated and the aqueous layer was extracted with ethyl acetate (2 x 100 mL). The combined extracts were washed with brine solution and dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the solvent gave crude solid which was crystallized from ethanol:water (3:1, 150:50 mL) and stored in the refrigerator overnight. The solids were collected by filtration and washed with ethanol:water mixture (3:1, 120 mL), water (30 mL), hexane (30 mL) and dried under vacuum to obtain 4.61 g (78%) of 1,3-dimethyl 5-iodo uracil as white needles (mp 226-228 °C, Lit. 225-227 °C, *Synthetic communication* 1988, 18, 855-867). EI-HRMS m/e calcd for C₆H₇IN₂O₂ (M⁺) 266.0021, found 266.0023.

c) Preparation of N-[(1,1-dimethylethoxy)carbonyl]-4-[1,3-dimethyl-2,4-dioxo-5-pyrimidinyl]-L-phenylalanine methyl ester




To a suspension of zinc dust (30 mmol, 2.0 g) in THF (3.0 mL) was added 1,2-dibromoethane (2.0 mmol, 0.174 mL) at room temperature. This suspension was heated to 60-65 °C with a heat gun until evolution of ethylene gas ceased (observed). This process was repeated three times. Then, the suspension was cooled to r.t. and trimethylchlorosilane (1.0 mmol, 0.15 mL) was added and the mixture was stirred for 10 min. A hot (the iodo compound was not very soluble in either solvent and it precipitated upon cooling to room temperature) solution of 5-iodo-1,3-dimethyluracil

(11.0 mmol, 2.92 g) in THF (3.0 mL) and DMA (10 mL) was added to the zinc suspension. After addition, the mixture was heated to 73 °C. The internal temperature of the reaction mixture rose to 77-78 °C due to the exothermic reaction. The reaction mixture was stirred at a bath temperature of 73 °C for 1.5 h and then was cooled to room temperature and stirred another 1.5 h. ¹H-NMR of a hydrolysate of a 0.25 mL aliquot of the mixture indicated the presence of traces of starting material and iodolysis of a 0.25 mL aliquot of the mixture gave the iodide back. The reaction mixture was heated to 70 °C and stirred for another 30 min. The reaction mixture was diluted with THF (5 mL) after cooling to room temperature and the excess zinc was allowed to settle for 30-60 min.


The above prepared zinc compound (5.5 mmol) was added to a suspension of Pd(dba)₂ (0.07 mmol, 40 mg), trifurylphosphine (TFP) (0.26 mmol, 66.6 mg) and Boc-4-iodo-L-phenylalanine methyl ester (4.5 mmol, 1.823 g) in THF (10 mL) at room temperature and the light yellow mixture was stirred for 15 h. The mixture was poured into a saturated ammonium chloride solution and was extracted with ethyl acetate (3 x 50 mL). The combined extracts were washed with brine solution and dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the solution gave the crude product which was purified by column chromatography to obtain 0.80 g (43%) of N-[(1,1-dimethylethoxy)carbonyl]-4-[1,3-dimethyl-2,4-dioxo-5-pyrimidinyl]-L-phenylalanine methyl ester as a white solid: mp 65-69 °C. EI-HRMS m/e calcd for C₂₁H₂₇N₃O₆ (M⁺) 418.1978, found 418.1965.

d) Preparation of 4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester.



A solution of N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester (1.94 mmol, 0.811 g) in dioxane (5 mL) was treated with 4.0 N (5 mL, 20 mmol) hydrochloric acid in dioxane at room temperature and the solution was stirred for 0.5 h. By this time, a light yellow precipitate slowly formed. The solids were collected by filtration and were washed with hexane to afford 412 mg (60% yield), mp 187-193 °C. The mother liquor was concentrated under vacuum and the residue was triturated with dichloromethane. The combined solids, which contained some starting material, were combined and dissolved in methanol. The major portion of the methanol was evaporated and dichloromethane was added to form a precipitate. The solids were collected and dried to obtain 330 mg (48%) of 4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester hydrochloride salt (mp 187-193 °C). EI-HRMS *m/e* calcd for C₁₆H₁₉N₃O₄ (M+H) 318.1454, found 318.1447.


e) Preparation of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester



To a suspension 4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester hydrochloride salt (0.5 mmol, 0.180 g), 2-chloro-6-methylbenzoic acid (0.55 mmol, 0.084 g) and HBTU (0.6 mmol, 0.227 g) in DMF (4 mL) was added diisopropylethylamine (3.0 mmol, 0.52 mL) at room temperature. After 1 min, everything went into solution and the yellow clear solution was stirred 15 h at room temperature. By this time, it turned to a brown solution was diluted with ethyl acetate

(25 mL). The ethyl acetate layer was washed successively with water (2 x 20 mL), saturated sodium bicarbonate solution (25 mL), and brine solution (25 mL) and was dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the solvent gave the crude product which was purified by silica gel column chromatography to afford 151.5 mg (72% yield) of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester as a white solid: mp 155-157 °C. EI-HRMS m/e calcd for $C_{24}H_{24}N_3O_5Cl$ (M^+) 470.1483, found 470.1484.

- 10 f) Preparation of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine




To a suspension of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester (0.278 mmol, 131 mg) in ethanol (4 mL) was added aqueous 1.0 N sodium hydroxide (3 mL) at room temperature. The mixture was heated to 40-45 °C and the resulting clear solution was stirred for 2-3 h. The ethanol was removed under reduced pressure and the residue was diluted with water (20 mL) and NaOH (3 mL, 1.0N) to dissolve the sodium salt. The aqueous solution was washed with ether (50 mL) to remove any neutral impurities. The aqueous layer was acidified with 1.0 N HCl and the product was extracted into ethyl acetate (2 x 50 mL). The combined organic extracts were washed with brine solution and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the filtrate afforded 107 mg (85%) of N-[(2-chloro-6-

methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine as a white solid: mp 234-236 °C. EI-HRMS m/e calcd for $C_{23}H_{22}N_3O_5Cl$ (M+H) 456.1326, found 456.1326.

5 Example 3.

Preparation of 4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)]-N-[[1-(2-methoxyethyl)cyclopentyl]carbonyl]-L-phenylalanine




10 4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)]-N-[[1-(2-methoxyethyl)cyclopentyl]-carbonyl]-L-phenylalanine was prepared from 4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)]-L-phenylalanine methyl ester and 1-(2-methoxyethyl)cyclopentane carboxylic acid (see WO 9910312) using the general procedures described in example 2. EI-HRMS m/e calcd for $C_{24}H_{31}N_3O_5$ (M+H) 458.2292, found 458.2279.

15

Example 4.


Preparation of N-[(2-bromo-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine



5 N-[(2-bromo-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine was prepared from 4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine and 2-bromo-6-methylbenzoic acid using the general procedures described in example 2. EI-HRMS m/e calcd for $C_{23}H_{22}N_3O_5Br$ (M+H) 500.0822, found 500.0825.

Example 5.


10 Preparation of N-[(2-bromo-5-methoxyphenyl)carbonyl]-4-[1,3-dimethyl-2,4-dioxo-5-pyrimidinyl]-L-phenylalanine



15 N-[(2-bromo-5-methoxyphenyl)carbonyl]-4-[1,3-dimethyl-2,4-dioxo-5-pyrimidinyl]-L-phenylalanine was prepared from 4-[1,3-dimethyl-2,4-dioxo-5-pyrimidinyl]-L-phenylalanine methyl ester and 2-bromo-5-methoxybenzoic acid using the general procedures described in example 2. EI-HRMS m/e calcd for $C_{23}H_{22}N_3O_6Br$ (M+H) 516.0770, found 516.0780.


Example 6. N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine

5 a) Preparation of 5-iodo-1,3,6-trimethyl uracil



A mixture of 5-iodo-6-methyl uracil (22.18 mmol, 5.58 g) and powdered potassium carbonate (60 mmol, 8.29 g) in DMF (188 mL) was stirred for 24 h at room temperature and then methyl iodide (90.6 mmol, 3.33 mL) was added. The reaction mixture was stirred for another 76 h at room temperature and was poured into water (150 mL) and ethyl acetate (150 mL). The layers were separated and the aqueous layer was extracted with ethyl acetate (2 x 100 mL). The combined extracts were washed with brine solution (150 mL) and dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the solvent gave crude solid which was crystallized from ethanol:water (3:1, 150:50 mL) and stored in the refrigerator overnight. The solids were collected by filtration and washed with ethanol:water (3:1, 120 mL), water (30 mL), hexanes (30 mL) and dried under high vacuum to obtain 5.8 g (93% yield) of 5-iodo-1,3,6-trimethyl uracil as a white solid: mp 155-157 °C. EI-HRMS m/e calcd for $\text{C}_7\text{H}_9\text{IN}_2\text{O}_2$ (M^+) 279.9709, found 279.9709.

20 b) Preparation of N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester



To a suspension of zinc dust (800 mmol, 52.29 g) in THF (26.0 mL) was added 1,2-dibromoethane (53.2 mmol, 4.58 mL) at room temperature. This suspension was heated to 60-65 °C with a heat gun until evolution of ethylene gas ceased (observed).


5 The suspension was cooled to room temperature, trimethylchlorosilane (26.6 mmol, 3.38 mL) was added and the mixture was stirred for 15 min. A suspension of 5-iodo-1,3,6-trimethyl uracil (266 mmol, 74.6 g) in DMA (225 mL) was warmed to obtain a clear solution and was added in one portion to the reaction mixture. After addition, the mixture was heated to 70 °C. The internal temperature of the reaction mixture rose to 80-85 °C due to the exothermic reaction. The reaction mixture was stirred at 10 70 °C for 3-4 h at which time TLC of an aliquot which had been quenched with saturated ammonium chloride indicated the absence of starting material. The reaction mixture was diluted with THF (140 mL), was cooled to room temperature and the excess zinc dust was allowed to settle over 2-3 h.

15 This solution containing the zinc compound (266 mmol) was added to a solution of Pd(dba)_2 (8 mmol, 4.6 g), tri-*o*-tolylphosphine [P(Tol)_3] (29.6 mmol, 9.0 g) and N-[(1,1-dimethylethoxy)carbonyl]-4-iodo-L-phenylalanine methyl ester (186 mmol, 75.56 g) in THF (280 mL) at room temperature and the light yellow mixture was stirred for 48 h at 50-55 °C. The reaction mixture was poured into a saturated ammonium chloride solution and was extracted with ethyl acetate (3 x 750 mL). The combined extracts were washed with brine solution (1.5 L) and dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration gave the crude product which was purified by silica gel column chromatography using a Biotage (75m) column to obtain 57.88 g (72% yield) of N-[(1,1-dimethylethoxy)carbonyl]-4-

20


(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester as an amorphous white solid. EI-HRMS m/e calcd for $C_{22}H_{29}N_3O_6$ (M^+) 431.2056, found 431.2054.

- 5 c) Preparation of 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester hydrochloride salt




10 portion of the solid N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester (17.15 mmol, 7.4 g) obtained above was treated with 4N hydrochloric acid in dioxane (68 mmol, 17 mL) at room temperature and the solution was stirred for 1 h as a white precipitate formed. The mixture was diluted with diethyl ether and the supernatant was decanted and the residue was dried first on the rotary evaporator and then under high vacuum to afford 6.28 g (99% yield) of 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester hydrochloride salt as an amorphous yellow solid. FAB-HRMS m/e calcd for $C_{17}H_{21}N_3O_4$ ($M+H$) 332.1610, found 332.1617.

- 20 d) Preparation of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester



To a suspension of 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester hydrochloride salt (3.12 mmol, 1.15 g) and 2,6-dichlorobenzoyl chloride (3.51 mmol, 0.735 g) in dichloromethane (40 mL) was added diisopropylethylamine (9.36 mmol, 1.63 mL) at room temperature. After 1 min, everything went into solution and the clear yellow solution was stirred for 20 h at room temperature. The resulting brown solution was diluted with dichloromethane (50 mL). The dichloromethane layer was washed successively with 1N hydrochloric acid (2 x 50 mL), saturated sodium bicarbonate solution (50 mL), and brine solution (50 mL) and was dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the solvent gave the crude product which was purified by silica gel chromatography using a Biotage (40m) column to afford 1.46 g (93% yield) of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester as an amorphous white solid. FAB-HRMS m/e calcd for $\text{C}_{24}\text{H}_{23}\text{Cl}_2\text{N}_3\text{O}_5$ (M+H) 504.1093, found 504.1083.


e) Preparation of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine



To a suspension of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester (2.2 mmol, 1.11 g) in ethanol (12 mL) was added aqueous 1.0 N sodium hydroxide (8.8 mL) at room temperature. The mixture was heated to 45-50 °C and the resulting clear solution was stirred for approximately 2 h. The ethanol was removed under reduced pressure and the residue was diluted with water (50 mL) and NaOH (5 mL, 1.0N) to dissolve the sodium salt. The aqueous solution was washed with diethyl ether (50 mL) to remove any neutral impurities. The aqueous layer was acidified with 1.0 N HCl and the product was extracted into ethyl acetate (2 x 75 mL). The combined organic extracts were washed with brine solution (100 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the filtrate afforded 970 mg (90% yield) of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine as a white solid: mp 225-227 °C. FAB-HRMS m/e calcd for $\text{C}_{23}\text{H}_{21}\text{Cl}_2\text{N}_3\text{O}_5$ (M+H) 490.0937, found 490.0940.


Example 7. Preparation of N-[[1-(2-methoxyethyl)cyclopentyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine

a) Preparation of N-[[1-(2-methoxyethyl)cyclopentyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester




To a suspension of 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester hydrochloride salt (0.4 mmol, 173 mg), HBTU (0.5 mmol, 189 mg) and 1-(2-methoxyethyl)cyclopentane carboxylic acid (0.5 mmol, 86 mg) in DMF (2 mL) was added diisopropylethylamine (1.2 mmol, 0.29 mL) at room temperature. After 5 min, everything went into solution and the clear yellow solution was stirred for 24 h at room temperature. The resulting dark-brown solution was diluted with ethyl acetate (30 mL). The ethyl acetate layer was washed successively with 1N hydrochloric acid (2 x 30 mL), saturated sodium bicarbonate solution (30 mL), and brine solution (30 mL) and was dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the solvent gave the crude product which was purified by silica gel column chromatography using a Biotage (40m) column to afford 139 mg (72% yield) of N-[[1-(2-methoxyethyl)cyclopentyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester as an amorphous white solid. FAB-HRMS m/e calcd for $\text{C}_{26}\text{H}_{35}\text{N}_3\text{O}_6$ (M+H) 486.2604, found 486.2602.

b) Preparation of N-[[1-(2-methoxyethyl)cyclopentyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine



To a suspension of N-[[1-(2-methoxyethyl)cyclopentyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester (0.273 mmol, 133 mg) in ethanol (3 mL) was added aqueous 1.0 N sodium hydroxide (1.5 mL) at room temperature. The mixture was heated to 40-45 °C and the resulting clear solution was stirred for 15 h. The ethanol was removed under reduced pressure and the residue was diluted with water (25 mL) and NaOH (3 mL, 1.0N) to dissolve the sodium salt. The aqueous solution was washed with diethyl ether (50 mL) to remove any neutral impurities. The aqueous layer was acidified with 1.0 N HCl and the product was extracted into ethyl acetate (2 x 25 mL). The combined organic extracts were washed with brine solution (50 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the filtrate afforded 121 mg (94% yield) of N-[[1-(2-methoxyethyl)cyclopentyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine as an amorphous white solid. FAB-HRMS m/e calcd for $\text{C}_{25}\text{H}_{33}\text{N}_3\text{O}_6$ (M+H) 472.2448, found 472.2467.


Example 8. Preparation of N-[(2-bromo-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine



$C_{24}H_{24}BrN_3O_5$
Mol. Wt.: 514.37

N-[(2-bromo-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine was prepared from 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester and 2-bromo-6-methylbenzoic acid using the general procedures described in example 7 and was obtained as a white solid: mp 240-242°C. FAB-HRMS m/e calcd for $C_{24}H_{24}BrN_3O_5$ (M+H) 514.0978, found 514.0965.

Example 9. Preparation of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine




$C_{24}H_{24}ClN_3O_5$
Mol. Wt.: 469.92

N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine was prepared from 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester and 2-chloro-6-methylbenzoic acid using the general procedures described in example 7 and was obtained as a white solid: mp

238-240°C. FAB-HRMS m/e calcd for $C_{24}H_{24}ClN_3O_5$ ($M+H$) 470.1483, found 470.1489.


5 Example 10. Preparation of N-[(2-ethyl-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine



$C_{26}H_{29}N_3O_5$
Mol. Wt.: 463.53

10 N-[(2-ethyl-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine was prepared from 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester and 2-ethyl-6-methylbenzoic acid using the general procedures described in example 7 and was obtained as a white solid: mp 127-133°C. ES-HRMS m/e calcd for $C_{26}H_{29}N_3O_5$ ($M+Na$) 494.1498, found 494.1501.


Example 11. Preparation of N-[[2-(2-methylethyl)-6-methylphenyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine



$C_{27}H_{31}N_3O_5$
Mol. Wt.: 477.55

5 N-[[2-(2-methylethyl)-6-methylphenyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine was prepared from 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester and 2-(2-methylethyl)-6-methylbenzoic acid using the general procedures described in example 7 and was obtained as an amorphous white solid. ES-HRMS m/e calcd for $C_{27}H_{31}N_3O_5$ (M+Na) 500.2156, found 500.2160.


Example 12. Preparation of N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine



$C_{23}H_{21}F_2N_3O_5$
Mol. Wt.: 457.43

10 N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine was prepared from 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester and 2,6-difluorobenzoic acid using the general procedures described in example 7 and was obtained as an amorphous white solid. ES-HRMS
15 m/e calcd for $C_{23}H_{21}F_2N_3O_5$ (M+Na) 480.1483, found 480.1489.


Example 13. Preparation of N-[[2-fluoro-6-(trifluoromethyl)phenyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine



$C_{24}H_{21}F_4N_3O_5$
Mol. Wt.: 507.43

5 N-[(2-fluoro-6-(trifluoromethyl)phenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine was prepared from 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester and 2-fluoro-6-(trifluoromethyl)benzoic acid using the general procedures described in example 7 and was obtained as a white solid: mp 218-220°C. ES-HRMS m/e calcd for $C_{25}H_{23}F_4N_3O_5$ (M+Na) 530.1310, found 530.1317.

10 Example 14. Preparation of N-[[2,6-di-(2-methylethyl)phenyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine




$C_{29}H_{35}N_3O_5$
Mol. Wt.: 505.61

15 N-[[2,6-di-(2-methylethyl)phenyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine was prepared from 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester and 2,6-di-(2-methylethyl)benzoic acid using the general procedures described in example 7 and was obtained as an

amorphous white solid. ES-HRMS m/e calcd for $C_{29}H_{35}N_3O_5$ ($M+Na$) 530.1310, found 530.1317.

5 Example 15. Preparation of N-[(2-chloro-6-ethylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine


a) Preparation of N-(2-chloro-6-fluorobenzylidene)butyl amine
(Ro 50-5007/000, 30935-229)



$C_{11}H_{13}ClFN$
Mol. Wt.: 213.68

10 To a suspension of 2-chloro-6-fluorobenzaldehyde (416 mmol, 66 g) in heptanes (200 mL) was added *n*-butylamine (460 mmol, 45.5 mL) at room temperature. After addition, an exothermic reaction as the solids dissolved completely. The solution was stirred for 3 h at room temperature, was transferred into a separatory funnel, was washed with brine solution (200 mL) and was dried over anhydrous magnesium
15 sulfate. Filtration of the drying agent and concentration gave a yellow oil which was purified by distillation under high vacuum (bp 95-98°C/4.5 mm Hg) to obtain 86.31 g (97% yield) of N-(2-chloro-6-fluorobenzylidene)butyl amine as a yellow oil. EI-HRMS m/e calcd for $C_{11}H_{13}ClFN$ (M^+) 213.0720, found 213.0714.

20 b) Preparation of 2-chloro-6-ethylbenzaldehyde




C_9H_9ClO
Mol. Wt.: 168.62

To a solution of N-(2-chloro-6-fluorobenzylidene)butyl amine (15 mmol, 3.21 g) in THF (20 mL) was added dropwise a solution of ethylmagnesium bromide (30 mmol, 30 mL, 1M) in THF by maintaining the temperature at 5-15°C. After addition, the

reaction mixture was allowed to warm to 20°C and was stirred for 5 h. Then, it was cooled to 0°C (ice bath) and 20% HCl in water (50 mL) was added dropwise while maintaining the temperature below 15°C with ice bath cooling. After addition, the mixture was allowed to warm to room temperature and was stirred for 15 h. Then, it was diluted with water (75 mL) and extracted with ethyl acetate (2 x 50 mL). The combined extracts were washed with brine solution (100 mL) and dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration gave 2.27 g (90% yield) of 2-chloro-6-ethylbenzaldehyde as a yellow oil. EI-HRMS m/e calcd for C₉H₉ClO (M⁺) 167.0264, found 167.0263.


c) Preparation of 2-chloro-6-ethylbenzoic acid



C₉H₉ClO₂
Mol. Wt.: 184.62

To a room temperature suspension of 2-chloro-6-ethylbenzaldehyde (13.5 mmol, 2.27 g) in acetonitrile (25 mL) was added a solution of monobasic sodium phosphate (3.4 mmol, 0.465 g) in water (7.5 mL) followed by hydrogen peroxide (1.8 mL, 30%). Then, a solution of sodium chlorite (23.7 mmol, 2.15 g) in water (20 mL) was added dropwise at 0 °C while maintaining the temperature below 3°C. After addition, the yellow suspension was stirred for 15 h at 0°C to room temperature. At this time TLC analysis of the mixture indicated the absence of starting material. Then, a solution of sodium bisulfite (20.5 mmol, 2.8 g) in water (10 mL) was added dropwise at 0 °C until the yellow color disappeared (KI-paper positive). Cooling is essential to control the exothermic reaction. After 1 h, the solvent was removed under vacuum. The neutral impurities were extracted with diethyl ether (200 mL). Then, the basic aqueous solution was neutralized with 10% HCl to pH ~1. The precipitated white solid was collected by filtration and dried at in air to afford 2.415 g (97% yield) of 2-chloro-6-ethylbenzoic acid as an amorphous white solid. EI-HRMS m/e calcd for C₉H₉ClO₂ (M⁺) 184.0291, found 184.0295.



d) Preparation of N-[(2-chloro-6-ethylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester



$C_{26}H_{28}ClN_3O_5$
Mol. Wt.: 497.98

To a suspension of 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester hydrochloride salt (0.5 mmol, 184 mg), HBTU (0.7 mmol, 265 mg) and 2-chloro-6-ethylbenzoic acid (0.7 mmol, 129 mg) in DMF (2 mL) was added diisopropylethylamine (1.25 mmol, 0.22 mL) at room temperature. After 5 min, everything went into solution and the clear yellow solution was stirred for 15 h at room temperature. The resulting dark-brown solution was diluted with ethyl acetate (30 mL). The ethyl acetate layer was washed successively with 1N hydrochloric acid (2 x 30 mL), saturated sodium bicarbonate solution (30 mL), and brine solution (30 mL) and was dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the solvent gave the crude product which was purified by silica gel column chromatography using a Biotage (40m) column to afford 175 mg (70% yield) of N-[(2-chloro-6-ethylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester as an amorphous white solid. ES-HRMS m/e calcd for $C_{26}H_{28}ClN_3O_5$ (M+Na) 520.1611, found 520.1613.


e) Preparation of N-[(2-chloro-6-ethylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine



To a suspension of N-[(2-chloro-6-ethylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester (0.33 mmol, 164 mg) in ethanol (2 mL) was added aqueous 1.0 N sodium hydroxide (0.7 mL) at room temperature. The mixture was stirred for 3 h at room temperature. The ethanol was removed under reduced pressure and the residue was diluted with water (30 mL). The aqueous solution was washed with diethyl ether (30 mL) to remove any neutral impurities. The aqueous layer was acidified with 1.0 N HCl and the product was extracted into ethyl acetate (2 x 35 mL). The combined organic extracts were washed with brine solution (50 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the filtrate afforded 138 mg (87% yield) of N-[(2-chloro-6-ethylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine as a white solid: mp 187-190°C. ES-HRMS m/e calcd for $C_{25}H_{26}ClN_3O_5$ (M+Na) 506.1459, found 506.1455.


Example 16. Preparation of N-[(2-chloro-6-propylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine

a. Preparation of 2-chloro-6-propylbenzoic acid.



2-chloro-6-propylbenzoic acid was prepared from 2-fluoro-6-chlorobenzilidine)-butylamine and propyl magnesium bromide using the general procedure described in example 15.


- 5 b. Preparation of N-[(2-chloro-6-propylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine



$C_{26}H_{28}ClN_3O_5$
Mol. Wt.: 497.97

- 10 N-[(2-chloro-6-propylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine was prepared from 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester and 2-chloro-6-propylbenzoic acid using the general procedures described in example 15 and was obtained as a white solid: mp 225-227°C. ES-HRMS m/e calcd for $C_{26}H_{28}ClN_3O_5$ (M+Na) 520.1611, found 520.1615.


- 15 Example 17. Preparation of N-[[2-chloro-6-(2-methylethyl)phenyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine
a. Preparation of 2-chloro-6-(2-methylethyl)benzoic acid.



Molecular Weight = 155.56
Molecular Formula = $C_7H_4ClO_2$

2-chloro-6-(2-methylethyl)benzoic acid was prepared from 2-fluoro-6-chlorobenzilidene)butylamine and isopropyl magnesium bromide using the general procedure described in example 15.

- 5 b. Preparation of N-[[2-chloro-6-(2-methylethyl)phenyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine




$C_{26}H_{28}ClN_3O_5$
Mol. Wt.: 497.97

- 10 N-[[2-chloro-6-(2-methylethyl)phenyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine was prepared from 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester and 2-chloro-6-(2-methylethyl)benzoic acid using the general procedures described in example 15 and was obtained as a white solid: mp 205-209°C. ES-HRMS m/e calcd for $C_{26}H_{28}ClN_3O_5$ (M+Na) 520.1611, found 520.1617.


- 15 Example 18. N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine

- a) Preparation of 5-iodo-1,3-diethyl-6-methyl uracil



To a suspension of 5-iodo-6-methyl uracil (20.97 mmol, 5.45 g) and powdered potassium carbonate (60 mmol, 8.29 g) in DMF (188 mL) was added ethyl iodide (83.88 mmol, 6.7 mL). The reaction mixture was stirred for 15 h at room temperature and was poured into water (150 mL) and ethyl acetate (150 mL). The layers were separated and the aqueous layer was extracted with ethyl acetate (2 x 100 mL). The combined extracts were washed with brine solution (150 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration gave a crude solid which was triturated with dichloromethane/diethyl ether/hexanes (1:1:1) to afford 3.89 g (60% yield) of 5-iodo-1,3-diethyl-6-methyl uracil as a white crystalline solid: mp 159-161.5 °C. EI-HRMS m/e calcd for $C_9H_{13}IN_2O_2$ (M^+) 308.0022, found 308.0018.

b) Preparation of N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester




To a suspension of zinc dust (33 mmol, 1.96 g) in THF (3 mL) was added 1,2-dibromoethane (3 mmol, 0.261 mL) at room temperature. This suspension was heated to 60-65 °C with a heat gun until evolution of ethylene gas ceased. The suspension was cooled to room temperature, trimethylchlorosilane (1.5 mmol, 0.19 mL) was added and the mixture was stirred for 15 min. A suspension of 5-iodo-1,3-diethyl-6-methyl uracil (11 mmol, 3.39 g) in DMA (6 mL) was warmed to obtain a clear solution and was added in one portion to the reaction mixture. After addition, the mixture was heated to 70 °C. The internal temperature of the reaction mixture rose to 75 °C due to the exothermic reaction. The reaction mixture was maintained at

70 °C for 15 h at which time the TLC analysis of an aliquot of the reaction mixture, which was quenched with saturated ammonium chloride solution, indicated the absence of starting material. The reaction mixture was diluted with THF (6 mL) and the reaction mixture was cooled to room temperature. The excess zinc dust was allowed to settle.

The solution containing the above prepared zinc compound (11 mmol) was added to a solution of Pd(dba)₂ (0.14 mmol, 80 mg), trifurylphosphine (TFP) (0.52 mmol, 134 mg) and N-[(1,1-dimethylethoxy)carbonyl]-4-iodo-L-phenylalanine methyl ester (9 mmol, 3.65 g) in THF (6 mL) at room temperature and the light yellow mixture was stirred for 72 h at 50-55 °C. The reaction mixture was poured into a saturated ammonium chloride solution (100 mL) and was extracted with ethyl acetate (3 x 75 mL). The combined extracts were washed with brine solution (150 mL) and dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration gave the crude product which was purified by silica gel column chromatography using a Biotage column (40m) to obtain 1.78 g (43% yield) of N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester as an amorphous white solid. ES-HRMS m/e calcd for C₂₄H₃₃N₃O₆ (M+Na) 482.2262, found 482.2262.


c) Preparation of 4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester hydrochloride salt



To a solution of N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester (3.87 mmol, 1.78 g) in dioxane (10 mL) was added 4N hydrochloric acid in dioxane (20 mmol, 5 mL) at room

temperature and the solution was stirred for 1 h. The solution was concentrated and was diluted with diethyl ether to form a white solid. The mother liquor was decanted and the residue was dried on a rotary evaporator and then under high vacuum to afford 0.72 g (47% yield) of 4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester hydrochloride salt as an amorphous solid. ES-HRMS m/e calcd for $C_{19}H_{25}N_3O_4$ (M+Na) 382.1737, found 382.1736.

d) Preparation of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester



$C_{26}H_{27}Cl_2N_3O_5$
Mol. Wt.: 532.42


10

15

20

To a suspension of 4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester hydrochloride salt (0.76 mmol, 0.3 g) and 2,6-dichlorobenzoyl chloride (0.84 mmol, 0.175 g) in dichloromethane (2 mL) was added diisopropylethylamine (3.03 mmol, 0.53 mL) at room temperature. After 5 min, everything went into solution and the clear yellow solution was stirred for 15 h at room temperature. The resulting brown solution was diluted with dichloromethane (25 mL). The dichloromethane layer was washed successively with 1N hydrochloric acid (2 x 25 mL), saturated sodium bicarbonate solution (25 mL), and brine solution (25 mL) and was dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the solvent gave the crude product which was purified by silica gel chromatography using a Biotage (40s) column to afford 0.40 g (99% yield) of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester as an amorphous white solid. ES-HRMS m/e calcd for $C_{26}H_{27}Cl_2N_3O_5$ (M+Na) 554.1221, found 554.1229.


e) Preparation of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine



To a suspension of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester (0.77 mmol, 0.41 g) in ethanol (2 mL) was added aqueous 1.0 N sodium hydroxide (1.5 mL) at room temperature. The mixture was heated to 50 °C and the resulting clear solution was stirred for 2 h. Then, the ethanol was removed under reduced pressure and the residue was diluted with water (25 mL) and NaOH (2 mL, 1.0N) to dissolve the sodium salt. The aqueous solution was washed with diethyl ether (30 mL) to remove any neutral impurities. The aqueous layer was acidified with 1.0 N HCl and the product was extracted into ethyl acetate (2 x 25 mL). The combined organic extracts were washed with brine solution (50 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the filtrate afforded 320 mg (80% yield) of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine as an amorphous white solid. ES-HRMS m/e calcd for $C_{25}H_{25}Cl_2N_3O_5$ (M+Na) 541.3921, found 541.3925.

Example 19. Preparation of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine


a) Preparation of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester



$C_{27}H_{30}ClN_3O_5$
Mol. Wt.: 512.00

To a suspension of 4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester hydrochloride salt (0.758 mmol, 300 mg), HBTU (0.84 mmol, 318 mg) and 2-chloro-6-methylbenzoic acid (0.84 mmol, 142 mg) in DMF (2 mL) was added diisopropylethylamine (1.9 mmol, 0.33 mL) at room temperature. After 5 min, everything went into solution and the clear yellow solution was stirred for 48 h at room temperature. The resulting dark-brown solution was diluted with ethyl acetate (30 mL). The ethyl acetate layer was washed successively with 1N hydrochloric acid (2 x 30 mL), saturated sodium bicarbonate solution (30 mL), and brine solution (30 mL) and was dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration gave the crude product which was purified by silica gel chromatography using a Biotage (40m) column to afford 380 mg (98% yield) of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester as an amorphous white solid. ES-HRMS m/e calcd for $C_{27}H_{30}ClN_3O_5$ (M+Na) 535.1026, found 535.1024.


b) Preparation of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine



$C_{26}H_{28}ClN_3O_5$
Mol. Wt.: 497.97

To a suspension of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester (0.82 mmol, 420 mg) in ethanol (2 mL) was added aqueous 1.0 N sodium hydroxide (1.6 mL) at room temperature. The mixture was heated to 50 °C and the resulting clear solution was stirred for 2 h. The ethanol was removed under reduced pressure and the residue was diluted with water (25 mL) and NaOH (3 mL, 1.0N) to dissolve the sodium salt. The aqueous solution was washed with diethyl ether (30 mL) to remove any neutral impurities. The aqueous layer was acidified with 1.0 N HCl and the product was extracted into ethyl acetate (2 x 25 mL). The combined organic extracts were washed with brine solution (50 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the filtrate afforded 277 mg (68% yield) of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine as an amorphous white solid. ES-HRMS m/e calcd for $C_{26}H_{28}ClN_3O_5$ (M+Na) 520.1611, found 520.1616.

Example 20. Preparation of 4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-N-[[1-(2-methoxyethyl)cyclopentyl]carbonyl]-L-phenylalanine




$C_{27}H_{37}N_3O_6$
Mol. Wt.: 499.60

4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-N-[[1-(2-methoxyethyl)-cyclopentyl]carbonyl]-L-phenylalanine was prepared from 4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester and 1-(2-methoxyethyl)-cyclopentane carboxylic acid using the general procedures described in example 19 and was obtained as an amorphous white solid. ES-HRMS m/e calcd for $C_{27}H_{37}N_3O_6$ (M+Na) 522.2575, found 522.2581.

Example 21. N-[1-(2,6-dichlorophenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine


a) Preparation of 3-methyl-6-(trifluoromethyl) uracil



To a pre-mixed solution of sodium methoxide (55 mmol, 2.97 g) and ethyl 3-amino-4,4,4-trifluorocrotonate (55 mmol, 10.0 g) in DMSO (19 mL, dried over molecular sieves) was added methyl isocyanate (55 mmol, 3.2 g) in DMSO (2.5 mL) over 15 min at 20 °C. The solution was stirred for 15 min and then another portion of sodium methoxide (27.5 mmol, 1.34 g) was added. After stirring for 15 min at 20 °C, methyl

isocyanate (14 mmol, 0.8 g) was added at this temperature. After a further 15 min, the reaction mixture was allowed to warm to room temperature and was stirred overnight. The resulting yellow suspension was poured into water (50 mL) to obtain a light yellow solution. The neutral impurities were extracted into diethyl ether (3 x 50 mL). The aqueous layer was acidified with concentrated hydrochloric acid to afford a white solid. The solids were collected by filtration and were washed with water. After air drying, 6.79 g (63% yield) of 3-methyl-6-(trifluoromethyl)uracil was obtained as a white solid: mp 235-237 °C. EI-HRMS m/e calcd for C₆H₅F₃N₂O₂ (M⁺) 194.0303, found 194.0303.


b) Preparation of 1,3-dimethyl-6-(trifluoromethyl)uracil



C₇H₇F₃N₂O₂
Mol. Wt.: 208.14


To a suspension of 3-methyl-6-(trifluoromethyl)uracil (20.6 mmol, 4.0 g) and powdered potassium carbonate (41.2 mmol, 5.7 g) in DME (30 mL) was added methyl iodide (82.4 mmol, 5.13 mL). Then, the reaction mixture was refluxed for 4 h at which time the TLC analysis of the reaction mixture indicated the absence of starting material. The reaction mixture was cooled to room temperature and was diluted with water (50 mL). Then, the DME was removed under reduced pressure to afford a white suspension. The solids were collected by filtration and washed with water. After air drying, 3.55 g (83% yield) of 1,3-dimethyl-6-(trifluoromethyl)uracil was obtained as a white solid: mp 85-87 °C. EI-HRMS m/e calcd for $C_7H_7F_3N_2O_2$ (M^+) 208.0459, found 208.0460.

c) Preparation of 1,3-dimethyl-5-iodo-6-(trifluoromethyl)uracil



A mixture of 1,3-dimethyl-6-(trifluoromethyl)uracil (16.91 mmol, 3.52 g), trifluoroacetic acid (20 mL) and trifluoroacetic anhydride (5 mL) was refluxed for 5 min. Then, NIS (16.91 mmol, 3.8 g) was added and the resulting mixture was stirred for 15 h at which time the TLC analysis of the reaction mixture indicated the presence of some starting material. Another portion of NIS (8.45 mmol, 1.9 g) was added and reflux was continued for another 5 h. The reaction mixture was cooled to room temperature and was poured slowly into a saturated potassium carbonate solution (100 mL). Then, sodium thiosulfite solution was added to remove the excess iodine color. The resulting solids were collected by filtration and washed with water. After air drying, 3.73 g (66% yield) of 1,3-dimethyl-5-iodo-6-(trifluoromethyl)uracil was obtained as a white solid: mp 149-151 °C. EI-HRMS m/e calcd for $\text{C}_7\text{H}_6\text{F}_3\text{IN}_2\text{O}_2$ (M^+) 333.9426, found 333.9436.


d) Preparation of N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine methyl ester



To a suspension of zinc dust (33 mmol, 1.96 g) in THF (3 mL) was added 1,2-dibromoethane (3 mmol, 0.261 mL) at room temperature. This suspension was heated to 60-65 °C with a heat gun until evolution of ethylene gas ceased. The suspension was cooled to room temperature, trimethylchlorosilane (1.5 mmol, 0.19 mL) was added and the mixture was stirred for 15 min. A suspension of 1,3-dimethyl-5-iodo-6-(trifluoromethyl)uracil (10 mmol, 3.34 g) in DMA (8 mL) was warmed to obtain a clear solution and was added in one portion to the reaction mixture. After addition, the mixture was heated to 70 °C. The internal temperature of the reaction mixture rose to 75 °C due to the exothermic reaction. The reaction mixture was stirred at 70 °C for approximately 3 h at which time TLC of an aliquot, which had been quenched with saturated ammonium chloride, indicated the absence of starting material. The reaction mixture was diluted with THF (5 mL), cooled to room temperature and excess zinc dust was allowed to settle.

The above solution of zinc compound (10 mmol) was added to a solution of Pd(dba)₂ (1.0 mmol, 520 mg), trifurylphosphine (TFP) (4.0 mmol, 0.93 g) and N-[(1,1-dimethylethoxy)carbonyl]-4-iodo-L-phenylalanine methyl ester (7 mmol, 2.84 g) in THF (10 mL) at room temperature and the light yellow mixture was stirred for 12 h at 45 °C. The reaction mixture was poured into a saturated ammonium chloride solution (100 mL) and was extracted with ethyl acetate (3 x 75 mL). The combined extracts were washed with brine solution (150 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration gave the crude product which was purified by silica gel chromatography using a Biotage (40m) column to obtain 1.45 g (42% yield) of N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine methyl ester as an amorphous white solid. ES-HRMS m/e calcd for C₂₂H₂₆NF₃N₃O₆ (M+Na) 508.1666, found 508.1670.

e) Preparation of 4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine methyl ester hydrochloride salt




H-Cl

 $C_{17}H_{19}ClF_3N_3O_4$

Mol. Wt.: 421.80

The solid N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3-diethyl-6)-(trifluoromethyl)-2,4-dioxo-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine methyl ester (2.92 mmol, 1.42 g) was treated with 4N hydrochloric acid in dioxane (28 mmol, 7 mL) at room temperature and the solution was stirred for 2 h. The reaction mixture was diluted with dichloromethane (5 mL) and was concentrated under reduced pressure on a rotoary evaporator. The residue was diluted with diethyl ether to form a light brown solid. The solids were collected by filtration and washed with diethyl ether. After drying, 1.21 g (91% yield) of 4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine methyl ester hydrochloride salt was obtained as a light brown solid: mp 244-247 °C. ES-HRMS m/e calcd for $C_{17}H_{18}F_3N_3O_4$ (M+H) 386.1322, found 386.1319.


f) Preparation of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine methyl ester



To a suspension of 4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine methyl ester hydrochloride salt (1.0 mmol, 421 mg) and 2,6-dichlorobenzoyl chloride (1.1 mmol, 0.235 g) in dichloromethane (3 mL) was added

diisopropylethylamine (4.4 mmol, 0.622 mL) at room temperature. After 1 min, everything went into solution and the light brown solution was stirred for 72 h at room temperature. The resulting dark brown solution was diluted with dichloromethane (25 mL). The dichloromethane layer was washed successively with 1N hydrochloric acid (2 x 25 mL), saturated sodium bicarbonate solution (25 mL), and brine solution (25 mL) and was dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the solvent gave a crude product, which was purified by silica gel chromatography using a Biotage (40s) column to afford 0.541 g (97% yield) of N-[1-(2,6-dichlorophenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine methyl ester as an amorphous white solid. ES-HRMS m/e calcd for $C_{24}H_{20}Cl_2F_3N_3O_5$ (M+Na) 580.0624, found 580.0629.

g) Preparation of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine




$C_{23}H_{18}Cl_2F_3N_3O_5$
Mol. Wt.: 544.31

To a suspension of N-[1-(2,6-dichlorophenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine methyl ester (0.422 mmol, 0.236 g) in pyridine (15 mL) was added lithium iodide (4.22 mmol, 0.571 g) at room temperature. The mixture was heated to reflux for 15 h. The reaction mixture was cooled to room temperature, diluted with 1N hydrochloric acid and extracted with ethyl acetate (2 x 25 mL). The combined organic extracts were washed with brine solution (50 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the filtrate afforded 201 mg (87% yield) of N-[1-

(2,6-dichlorophenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine as a light yellow solid: mp 125-128 °C. ES-HRMS m/e calcd for $C_{23}H_{18}Cl_2F_3N_3O_5$ ($M+Na$) 566.0469, found 566.0468.

5 Example 22. Preparation of N-[(2-chloro-6-methylphenyl)carbonyl]-4-[1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl]-L-phenylalanine



Molecular Weight = 523.89
Molecular Formula = $C_{24}H_{21}ClF_3N_3O_5$


A solution of 2-chloro-6-methylbenzoic acid (190 mg, 1.14 mmol) in dichloromethane (7 mL) containing DMF (4 drops) was treated with oxalyl chloride (0.42 mL, 4.8 mmol) and the mixture was stirred for 2 h. The mixture was concentrated, azeotroping with toluene to remove traces of oxalyl chloride and the residue was used directly in the next step.

A mixture of the above prepared acid chloride, 4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine methyl ester hydrochloride salt (423 mg, 1.003 mmol) in dichloromethane (5 mL) was treated with DIPEA (0.625 mL, 4.46 mmol) and the resulting light brown solution was stirred for 3 days. The mixture was concentrated, diluted with ethyl acetate, washed with 1 N HCl and brine solution and was dried over magnesium sulfate. Filtration and evaporation afforded a residue, which was purified by silica gel chromatography using a Biotage column (40s) to give N-[(2-chloro-6-methylphenyl)carbonyl]-4-[1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl]-L-phenylalanine methyl ester as a white foam (179 mg, 33%). ES-HRMS m/e calcd for $C_{25}H_{23}ClN_3O_5$ ($M+Na$) 560.1171, found 560.1172.

A solution of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine methyl ester (260 mg, 0.48 mmol),

obtained as in the above experiment, and lithium iodide (6534 mg, 4.8 mmol) in pyridine (14 mL) was heated to reflux overnight. The mixture was diluted with 1 N HCl and extracted with ethyl acetate. The combined extracts were washed with brine solution and dried over magnesium sulfate, filtered and evaporated. The residue was trituated with ether, hexane and dichloromethane to give N-[(2-chloro-6-methylphenyl)carbonyl]-4-[1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl]-L-phenylalanine (205 mg, 81%) as a white solid: mp 243-247°C. ES-HRMS m/e calcd for $C_{24}H_{21}ClF_3N_3O_5$ (M+Na) 546.1014, found 546.1013.

Example 23. Preparation of N-[[2-fluoro-6-(trifluoromethyl)phenyl]carbonyl]-4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine



A solution of 2-fluoro-6-trifluoromethylbenzoic acid (125 mg, 0.60 mmol) (Aldrich 33080-9) in dichloromethane (3 mL) containing DMF (2 drops) was treated with oxalyl chloride (0.21 mL, 2.4 mmol) and the mixture was stirred for 2 h. The mixture was concentrated, azeotroping with toluene to remove traces of oxalyl chloride and the residue was used directly in the next step.


A mixture of the above prepared acid chloride, 4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine methyl ester hydrochloride salt (210 mg, 0.50 mmol) in dichloromethane (3 mL) was treated with DIPEA (0.336 mL, 2.4 mmol) and the resulting light brown solution was stirred for 3 days. The mixture was concentrated, diluted with ethyl acetate, washed with 1 N HCl and brine solution and was dried over magnesium sulfate. Filtration and evaporation afforded a residue, which was purified by silica gel chromatography using a Biotage column (40s) to give N-[[2-fluoro-6-(trifluoromethyl)phenyl]carbonyl]-4-(1,3-dimethyl-2,4-dioxo-6-

(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine methyl ester as a white foam (179 mg, 62%). ES-HRMS m/e calcd for $C_{25}H_{20}F_7N_3O_5$ (M+Na) 598.1183, found 598.1186.

A solution of N-[[2-fluoro-6-(trifluoromethyl)phenyl]carbonyl]-4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine methyl ester (266 mg, 0.46 mmol), obtained as in the above experiment, and lithium iodide (624 mg, 4.6 mmol) in pyridine (14 mL) was heated to reflux overnight. The mixture was diluted with 1 N HCl and extracted with ethyl acetate. The combined extracts were washed with brine solution and dried over magnesium sulfate, filtered and evaporated. The residue was triturated with ether and dichloromethane to give N-[[2-fluoro-6-(trifluoromethyl)phenyl]carbonyl]-4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine (167 mg, 64%) as a white solid: mp 122-125°C. ES-HRMS m/e calcd for $C_{24}H_{18}F_7N_3O_5$ (M+Na) 584.1027, found 584.1028.


Example 24. N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine

a) Preparation of 4-[(1,1-dimethylethoxy)carbonyl]amino-3-methylbenzyl alcohol




A mixture of 4-nitro-3-methylbenzyl alcohol (56.53 mmol, 9.45 g), di-tert-butyl dicarbonate (63 mmol, 13.74 g) and palladium on charcoal (450 mg) in ethyl acetate (240 mL) was hydrogenated for 2 h at room temperature. Then, the reaction mixture was filtered through a pad of celite washing with ethyl acetate (50 mL). The filtrate was concentrated under vacuum to obtain 10.18 g (76% yield) of 4-[(1,1-dimethylethoxy)carbonyl]amino-3-methylbenzyl alcohol as a light yellow solid. EI-HRMS m/e calcd for $C_{13}H_{19}NO_3$ (M^+) 237.0126, found 237.0129.

b) Preparation of 4-[[[(1,1-dimethylethoxy)carbonyl]amino]-3-methylbenzaldehyde



To a solution of 4-[[[(1,1-dimethylethoxy)carbonyl]amino]-3-methylbenzyl alcohol (42.9 mmol, 10.18 g) in dichloromethane (85 mL) was added manganese dioxide (138 mmol, 12 g) and 4A molecular sieves (6 g) at room temperature. The reaction mixture was stirred for 76 h at room temperature and was filtered through a pad of celite washing with dichloromethane. The filtrate was concentrated under reduced pressure and the residue was purified by silica gel chromatography using a Biotage (40m) column to obtain 7.82 g (77% yield) of 4-[[[(1,1-dimethylethoxy)carbonyl]amino]-3-methylbenzaldehyde as a white solid: mp 109-111 °C. EI-HRMS m/e calcd for $\text{C}_{13}\text{H}_{17}\text{NO}_3$ (M^+) 235.1208, found 235.1207.


c) Preparation of N-(benzyloxycarbonyl)-4-[[[(1,1-dimethylethoxy)carbonyl]amino]-3-methyldehydrophenylalanine methyl ester



To a solution of N-(benzyloxycarbonyl)- α -phosphonoglycine trimethyl ester (18 mmol, 5.96 g) (Aldrich Chemical Company) in dichloromethane (30 mL) was added tetramethylguanidine (18 mmol, 2.07 g) at room temperature. The reaction mixture was stirred for 1 h at room temperature and it was cooled to -30 °C. Then, a solution of 4-[[[(1,1-dimethylethoxy)carbonyl]amino]-3-methylbenzaldehyde (15 mmol, 3.52 g) in dichloromethane (12.5 mL) was added in one portion. After 30 min at this

temperature, the reaction mixture was allowed to warm to room temperature and was stirred for 15 h. Then, the reaction mixture was diluted with diethyl ether (100 mL) and was washed successively with 0.5N hydrochloric acid (2 x 50 mL), saturated sodium bicarbonate solution (100 mL), brine solution (100 mL) and was dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration gave a crude product, which was purified by silica gel chromatography using a Biotage (40m) column to obtain 3.87 g (58% yield) of N-(benzyloxycarbonyl)-4-[[[(1,1-dimethylethoxy)carbonyl]amino]-3-methyl-dehydrophenylalanine methyl ester as a white solid. EI-HRMS m/e calcd for $C_{24}H_{28}N_2O_6$ (M^+) 440.1527, found 440.1524.


d) Preparation of N-(benzyloxycarbonyl)-4-[[[(1,1-dimethylethoxy)carbonyl]amino]-3-methyl-L-phenylalanine methyl ester



$C_{24}H_{30}N_2O_6$
Mol. Wt.: 442.50


A stream of argon was passed through a solution of N-(benzyloxycarbonyl)-4-[[[(1,1-dimethylethoxy)carbonyl]amino]-3-methyl-dehydrophenylalanine methyl ester (8.08 mmol, 3.56 g) in methanol (25 mL) in a Parr pressure vessel overnight. Then, the catalyst, (+)-1,2-bis((2S,5S)-2,5-dimethylphospholano)benzene(cyclooctadiene)-rhodium (I) trifluoromethanesulfonate [[Rh(COD)(S,S)-(me)DuPHOS]+TfO-] (~40 mg) was added under a stream of argon in a glove box. The solution was stirred under a hydrogen pressure (60 psi) at room temperature for 22 h. The resulting solution was concentrated and the crude product was purified by silica gel chromatography using a Biotage (40m) column to obtain 2 g (55% yield) of N-(benzyloxycarbonyl)-4-[[[(1,1-dimethylethoxy)carbonyl]amino]-3-methyl-L-phenylalanine methyl ester as an amorphous white solid. EI-HRMS m/e calcd for $C_{24}H_{30}N_2O_6$ (M^+) 442.1627, found 442.1629.

e) Preparation of N-(benzyloxycarbonyl)-4-amino-3-methyl-L-phenylalanine methyl ester hydrochloride salt



- 5 To a solution of N-(benzyloxycarbonyl)-4-[(1,1-dimethylethoxy)carbonyl]amino-3-methyl-L-phenylalanine methyl ester (4.52 mmol, 2 g) in dioxane (12 mL) was added 4N hydrochloric acid in dioxane (48 mmol, 2 mL) at room temperature and the solution was stirred for approximately 2 h as a white precipitate was formed. The solids were diluted with diethyl ether, the mother liquor was decanted and the residue
- 10 was dried first on a rotary evaporator and then under high vacuum to afford 1.487 g (87% yield) of N-(benzyloxycarbonyl)-4-amino-3-methyl-L-phenylalanine methyl ester hydrochloride salt as an amorphous yellow solid. FAB-HRMS m/e calcd for $C_{19}H_{22}N_2O_4$ (M+H) 343.0142, found 343.0144.


15 f) Preparation of N-(benzyloxycarbonyl)-4-iodo-3-methyl-L-phenylalanine methyl ester



A suspension of sulfuric acid (0.3 mL), water (36 mL) and N-(benzyloxycarbonyl)-4-amino-3-methyl-L-phenylalanine methyl ester hydrochloride salt (2.9 mmol, 1.1 g)

was heated to obtain a clear solution. Then, it was cooled to -1°C (ice-bath) and a solution of sodium nitrite (5.8 mmol, 400 mg) in water (8 mL) was added dropwise. The reaction mixture was stirred for 30 min, and a solution of potassium iodide (8.7 mmol, 1.5 g) in water (6 mL) was added to obtain a brown suspension. After stirring for 30 min, the reaction mixture was allowed to warm to room temperature and was stirred for 1 h. Then, the reaction mixture was diluted with water (100 mL) and was extracted with ethyl acetate (3 x 50 mL). The combined extracts were washed with saturated sodium bisulfite solution (100 mL) and brine solution (100 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration gave a crude product, which was purified by silica gel chromatography using a Biotage (40m) column to afford 0.84 g (64% yield) of N-(benzyloxycarbonyl)-4-iodo-3-methyl-L-phenylalanine methyl ester as an amorphous white solid. ES-HRMS m/e calcd for $\text{C}_{19}\text{H}_{20}\text{INO}_4$ ($\text{M}+\text{Na}$) 476.0329, found 476.0336.

g) Preparation of N-(benzyloxycarbonyl)-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester




$\text{C}_{25}\text{H}_{27}\text{N}_3\text{O}_6$
Mol. Wt.: 465.50

To a suspension of zinc dust (15 mmol, 0.98 g) in THF (1.5 mL) was added 1,2-dibromoethane (1 mmol, 0.13 mL) at room temperature. This suspension was heated to $60\text{--}65^{\circ}\text{C}$ with a heat gun until evolution of ethylene gas ceased. Then, the suspension was cooled to room temperature and trimethylchlorosilane (0.5 mmol, 70 μL) was added and the mixture was stirred for 15 min. A suspension of 5-iodo-1,3-dimethyl uracil (2.5 mmol, 665 mg) in DMA (2 mL) was warmed to obtain a clear

solution and was added in one portion to the reaction mixture. After addition, the mixture was heated to 70 °C. The reaction mixture was stirred at 70 °C for approximately 3 h, at which time TLC of an aliquot, which had been quenched with saturated ammonium chloride, indicated the absence of starting material. The mixture was diluted with THF (2 mL), allowed to cool room temperature and the excess zinc dust was allowed to settle.

The above prepared solution of zinc compound (2.5 mmol) was added to a solution of Pd(dba)₂ (0.05 mmol, 27 mg), trifurylphosphine (TFP) (0.2 mmol, 50 mg) and N-(benzyloxycarbonyl)-4-iodo-3-methyl-L-phenylalanine methyl ester (0.5 mmol, 227 mg) in THF (2 mL) at room temperature and the resulting light yellow mixture was stirred for 15 h at 45 °C. The reaction mixture was then poured into a saturated ammonium chloride solution and was extracted with ethyl acetate (3 x 30 mL). The combined extracts were washed with brine solution (50 mL) and dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration gave the crude product, which was purified by silica gel chromatography using a Biotage (40m) column to obtain 161 mg (69% yield) of N-(benzyloxycarbonyl)-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester as an amorphous white solid. ES-HRMS m/e calcd for C₂₅H₂₇N₃O₆ (M+Na) 488.1792, found 488.1801.

h) Preparation of 4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester




C₁₇H₂₁N₃O₄
Mol. Wt.: 331.37

A mixture of N-(benzyloxycarbonyl)-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester (0.34 mmol, 159 mg), cyclohexene (1 mL) and 10% palladium on carbon (100 mg) in ethanol (3 mL) was heated to reflux for 20

min. Then, it was filtered through a pad of celite and the pad was washed with ethanol (10 mL). The combined filtrate was concentrated and the residue was dried under high vacuum to afford 96 mg (85% yield) of 4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester as a sticky yellow solid. ES-HRMS m/e calcd for $C_{17}H_{21}N_3O_4$ (M+Na) 354.1424, found 354.1424.



i) Preparation of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester



$C_{25}H_{26}ClN_3O_5$
Mol. Wt.: 483.94

To a suspension of 4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester (0.125 mmol, 46 mg), HBTU (0.125 mmol, 47.5 mg) and 2-chloro-6-methylbenzoic acid (0.137 mmol, 25 mg) in DMF (2 mL) was added diisopropylethylamine (0.312 mmol, 44 μ L) at room temperature. After 5 min, everything went into solution and the clear yellow solution was stirred for 72 h at room temperature. The resulting dark-brown solution was diluted with ethyl acetate (30 mL). The ethyl acetate layer was washed successively with 1N hydrochloric acid (2 x 30 mL), saturated sodium bicarbonate solution (30 mL), and brine solution (30 mL) and was dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the solvent gave the crude product which was purified by silica gel chromatography using a Biotage (40s) column to afford 42 mg (70% yield) of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester as a oily residue. ES-HRMS m/e calcd for $C_{25}H_{26}ClN_3O_5$ (M+Na) 506.1454, found 506.1459.


j) Preparation of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine



To a suspension of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester (0.083 mmol, 40 mg) in ethanol (1 mL) was added aqueous 1.0 N sodium hydroxide (0.2 mL) at room temperature. The mixture was stirred for 2 h at room temperature. Then, the ethanol was removed under reduced pressure and the residue was diluted with water (5 mL). The aqueous solution was washed with diethyl ether (20 mL) to remove any neutral impurities. The aqueous layer was acidified with 1.0 N HCl and the product was extracted into ethyl acetate (2 x 25 mL). The combined organic extracts were washed with brine solution (50 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the filtrate afforded 34 mg (87% yield) of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine as an amorphous white solid. ES-HRMS m/e calcd for $C_{24}H_{24}ClN_3O_5$ (M+Na) 492.1295, found 492.1301.


Example 25. Preparation of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine

a) Preparation of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester



To a suspension of 4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester (0.128 mmol, 47 mg) and 2,6-dichlorobenzoyl chloride (0.153 mmol, 32 mg) in dichloromethane (1 mL) was added diisopropylethylamine (0.45 mmol, 77 μ L) at room temperature. After 5 min, everything went into solution and the clear yellow solution was stirred for 15 h at room temperature. The resulting brown solution was diluted with dichloromethane (25 mL). The dichloromethane layer was washed successively with 1N hydrochloric acid (2 x 25 mL), saturated sodium bicarbonate solution (25 mL), and brine solution (25 mL) and was dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration gave the crude product which was purified by silica gel chromatography using a Biotage (40m) column to afford 52 mg (81% yield) of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester as an amorphous white solid. ES-HRMS m/e calcd for $C_{24}H_{23}Cl_2N_3O_5$ ($M+Na$) 526.0907, found 526.0912.

b) Preparation of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine




$C_{23}H_{21}Cl_2N_3O_5$
Mol. Wt.: 490.34

To a suspension of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester (0.1 mmol, 59 mg) in ethanol (1 mL) was added aqueous 1.0 N sodium hydroxide (0.2 mL) at room temperature. The mixture was stirred for 2 h at room temperature. The ethanol was removed under reduced pressure and the residue was diluted with water (10 mL). The aqueous solution was washed with diethyl ether (20 mL) to remove any neutral impurities. The aqueous layer was acidified with 1.0 N HCl and the product was extracted into ethyl acetate (2 x 15 mL). The combined organic extracts were washed with brine solution (50 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the filtrate afforded 20 mg (41% yield) of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine as an amorphous white solid. ES-HRMS m/e calcd for $C_{23}H_{21}Cl_2N_3O_5$ (M+Na) 512.0752, found 512.0754.

Example 26. N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine



a) Preparation of N-(benzyloxycarbonyl)-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester



To a suspension of zinc dust (15 mmol, 0.98 g) in THF (1.5 mL) was added 1,2-dibromoethane (1 mmol, 0.13 mL) at room temperature. This suspension was heated to 60-65 °C with a heat gun until evolution of ethylene gas ceased. Then, the suspension was cooled to room temperature and trimethylchlorosilane (0.5 mmol, 70 uL) was added and the mixture was stirred for 15 min. A suspension of 5-iodo-1,3,6-trimethyl uracil (2.5 mmol, 700 mg) in DMA (2 mL) was warmed to obtain a clear solution and was added in one portion to the reaction mixture. The reaction mixture was stirred at 70 °C for 3-4 h at which time TLC of an aliquot which, had been quenched with saturated ammonium chloride, indicated the absence of starting material. Then, the reaction mixture was diluted with THF (3 mL) and the excess zinc dust was allowed to settle.



The above prepared solution containing the zinc compound (2.5 mmol) was added to a solution of $\text{Pd}(\text{dba})_2$ (0.05 mmol, 27 mg), trifurylphosphine (TFP) (0.2 mmol, 50 mg) and N-(benzyloxycarbonyl)-4-iodo-3-methyl-L-phenylalanine methyl ester (0.5 mmol, 227 mg) in THF (2 mL) at room temperature and the light yellow mixture was stirred for 15 h at 45 °C. Then, the reaction mixture was poured into a saturated ammonium chloride solution and was extracted with ethyl acetate (3 x 30 mL). The combined extracts were washed with brine solution (50 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration gave the crude product which was purified by silica gel chromatography using a Biotage (40m) column to obtain 71 mg (30% yield) of N-(benzyloxycarbonyl)-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester as a yellow oil. ES-HRMS m/e calcd for $\text{C}_{26}\text{H}_{29}\text{N}_3\text{O}_6$ ($M+\text{Na}$) 502.2173, found 502.2174.

b) Preparation of 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester




- 5 A mixture of N-(benzyloxycarbonyl)-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester (0.145 mmol, 70 mg), cyclohexene (1 mL) and 10% palladium on carbon (100 mg) in ethanol (2 mL) was heated to reflux for 15 h. Then, it was filtered through a pad of celite and the pad was washed with ethanol (10 mL). The combined filtrate was concentrated under reduced pressure. The residue
10 was dried under high vacuum to afford 36 mg (72% yield) of 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester as a light yellow solid. ES-HRMS m/e calcd for $C_{18}H_{23}N_3O_4$ (M+Na) 368.1327, found 368.1321.

15 c) Preparation of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester



To a suspension of 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester (0.089 mmol, 34 mg) and 2,6-dichlorobenzoyl chloride (0.1 mmol, 21 mg) in dichloromethane (2 mL) was added diisopropylethylamine (0.4 mmol, 70 μ L) at room temperature. After 5 min, everything went into solution and the clear yellow solution was stirred for 15 h at room temperature. The resulting brown solution was diluted with dichloromethane (25 mL). The dichloromethane layer was washed successively with 1N hydrochloric acid (2 x 25 mL), saturated sodium bicarbonate solution (25 mL), and brine solution (25 mL) and was dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the solvent gave the crude product which was purified by silica gel chromatography using a Biotage (40m) column to afford 22 mg (48% yield) of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester as a viscous oil. ES-HRMS m/e calcd for $C_{25}H_{25}Cl_2N_3O_5$ (M+Na) 541.1065, found 541.1063.


d) Preparation of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine



To a suspension of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine methyl ester (0.04 mmol, 22 mg) in ethanol (2 mL) was added aqueous 1.0 N sodium hydroxide (0.5 mL) at room temperature. The mixture was stirred for 2 h at room temperature. The ethanol was removed under reduced pressure and the residue was diluted with water (10 mL). The aqueous solution was washed with diethyl ether (20 mL) to remove any neutral


impurities. The aqueous layer was acidified with 1.0 N HCl and the product was extracted into ethyl acetate (2 x 15 mL). The combined organic extracts were washed with brine solution (50 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the filtrate afforded 20 mg (93% yield) of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine as an amorphous white solid. ES-HRMS m/e calcd for $C_{23}H_{21}Cl_2N_3O_5$ [(M-H)+2Na] 548.0725, found 548.0733.

Example 27. N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine ethyl ester



To a suspension of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine (0.6 mmol, 300 mg) and sodium bicarbonate (3.6 mmol, 302 mg) in DMF (4 mL) was added iodoethane (3.6 mmol, 0.29 mL) at room temperature. The mixture was stirred for 72 h at room temperature. Then, the reaction mixture was poured into water (50 mL) and was extracted with ethyl acetate (3 x 20 mL). The combined extracts were washed with brine solution (60 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the filtrate gave the crude product which was purified by silica gel chromatography using a Biotage (40m) column to obtain 155 mg (50% yield) of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine ethyl ester as a crystalline white solid: mp 262-265 °C. ES-HRMS m/e calcd for $C_{25}H_{25}Cl_2N_3O_5$ (M+Na) 540.1062, found 540.1049.


Example 28. Preparation of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-[(N,N-diethyl)amino]ethyl ester



5 A mixture of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine (320 mg, 0.65 mmol), 2-[(N,N-diethyl)amino]ethyl chloride hydrochloride (579 mg, 3.26 mmol) and potassium carbonate (451 mg, 3.27 mmol) in ethyl acetate (5 mL) and water (5 mL) was at room temperature overnight. The layers were separated and the aqueous layer was extracted with ethyl acetate. The combined extracts were washed with brine, dried over magnesium sulfate and
 10 concentrated to afford N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-[(N,N-diethyl)amino]ethyl ester (190 mg, 49%) as an amorphous white solid. ES-HRMS m/e calcd for $C_{29}H_{34}Cl_2N_4O_5$ (M+H) 589.1978, found 589.1980.

15 Acidification of the aqueous layer afforded recovered N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine (167 mg).


Example 29. Preparation of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-[(N,N-diethyl)amino]ethyl ester



$C_{30}H_{37}ClN_4O_5$
Mol. Wt.: 569.09

N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-[(N,N-diethyl)amino]ethyl ester was prepared from N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine and 2-[(N,N-diethyl)aminoethyl chloride hydrochloride using the general procedure described in example 28 and was obtained as an amorphous white solid. ES-HRMS m/e calcd for $C_{30}H_{37}ClN_4O_5$ (M+H) 569.2525, found 569.2530.

Example 30. Preparation of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine ethyl ester




$C_{26}H_{28}ClN_3O_5$
Mol. Wt.: 497.97

N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine ethyl ester was prepared from N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine and iodoethane using the general procedure described in example 27

and was obtained as an amorphous white solid. ES-HRMS m/e calcd for $C_{26}H_{28}ClN_3O_5$ ($M+Na$) 520.1610, found 520.1591.

Example 31. N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-(4-morpholino)ethyl ester


a) Preparation of N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine



$C_{21}H_{27}N_3O_6$
Mol. Wt.: 417.46


To a suspension of N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester (8.8 mmol, 3.79 g) in ethanol (20 mL) was added aqueous 1.0 N sodium hydroxide (17.57 mL) at room temperature. The mixture was stirred for 2 h at room temperature. Then, the mixture was diluted with water (50 mL) and the ethanol was removed under reduced pressure. The aqueous solution was washed with diethyl ether (100 mL) to remove any neutral impurities. The aqueous layer was acidified with 1.0 N HCl and the product was extracted into ethyl acetate (2 x 100 mL). The combined organic extracts were washed with brine solution (200 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the filtrate afforded 3.44 g (94% yield) of N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine as a light brown foam solid. ES-HRMS m/e calcd for $C_{21}H_{27}N_3O_6$ ($M+Na$) 440.1792, found 440.1792.

b) Preparation of N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-(4-morpholino)ethyl ester




To a solution of N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine (1.21 mmol, 505 mg) and 2-(4-morpholino)ethanol (2.42 mmol, 318 mg) in THF (8 mL) was added di-*iso*-propylcarbodiimide (DIC) (1.82 mmol, 0.287 mL) and 4-dimethylaminopyridine (0.6 mmol, 74 mg) at room temperature. The resulting solution was stirred for 72 h. Then, the reaction mixture was poured into water (50 mL) and was extracted with ethyl acetate (3 x 50 mL). The combined extracts were washed with water (2 x 50 mL) and brine solution (100 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the filtrate gave the crude product which was purified by silica gel chromatography using a Biotage (40m) column to obtain 428 mg (67% yield) of N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-(4-morpholino)ethyl ester as an amorphous white solid. ES-HRMS m/e calcd for $C_{27}H_{38}N_4O_7$ ($M+Na$) 553.2633, found 553.2636.

c) Preparation of 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-(4-morpholino)ethyl ester hydrochloride salt



The solid N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-(4-morpholino)ethyl ester (1.6 mmol, 0.85 g) was treated with 4N hydrochloric acid in dioxane (16.02 mmol, 4 mL) at room temperature and the solution was stirred for 3 h as a white precipitate formed. The solids were diluted with diethyl ether, the mother liquor was decanted and the residue was dried first on the rotary evaporator and then under high vacuum to afford 0.75 g (99% yield) of 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-(4-morpholino)ethyl ester hydrochloride salt as an amorphous yellow solid. ES-HRMS m/e calcd for $C_{22}H_{30}N_4O_5$ (M+H) 431.2289, found 431.2292.


d) Preparation of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-(4-morpholino)ethyl ester



To a suspension of 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-(4-morpholino)ethyl ester hydrochloride salt (0.85 mmol, 399 mg) and 2,6-dichlorobenzoyl chloride (0.95 mmol, 0.201 g) in dichloromethane (4 mL) was added diisopropylethylamine (4.75 mmol, 0.66 mL) at room temperature. After 5 min, everything went into solution and the clear yellow solution was stirred for 48 h at room temperature. The resulting light brown solution was diluted with dichloromethane (50 mL). The dichloromethane layer was washed successively with 1N hydrochloric acid (2 x 50 mL), saturated sodium bicarbonate solution (50 mL), and brine solution (50 mL) and was dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration gave the crude product which was purified by silica gel chromatography using a Biotage (40m) column to afford 0.427

g (75% yield) of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-(4-morpholino)ethyl ester as a white solid: mp 90-94 °C. ES-HRMS m/e calcd for $C_{29}H_{32}Cl_2N_4O_6$ (M+H) 603.1772, found 603.1782.


- 5 Example 32. Preparation of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-(4-morpholino)ethyl ester



- 10 N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-(4-morpholino)ethyl ester was prepared from 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-(4-morpholino)ethyl ester and 2-chloro-6-methylbenzoyl chloride using the general procedures described in example 31 and was obtained as an amorphous white solid. ES-HRMS m/e calcd for $C_{30}H_{35}ClN_4O_6$ (M+H) 583.2318, found 583.2326.

- 15 Example 33. Preparation of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxy)ethyl ester


a) Preparation of N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxy)ethyl ester



$C_{25}H_{33}N_3O_8$
Mol. Wt.: 503.56

To a suspension of N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine (1.006 mmol, 420 mg) and sodium bicarbonate (5.03 mmol, 422 mg) in DMF (8 mL) was added 1-chloroethyl acetate (5.03 mmol, 616 mg) at room temperature. The reaction mixture was stirred for 48 h. Then, it was poured into water (50 mL) and was extracted with ethyl acetate (3 x 50 mL). The combined extracts were washed with water (2 x 50 mL) and brine solution (100 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the filtrate gave the crude product which was purified by silica gel chromatography using a Biotage (40s) column to obtain 390 mg (77% yield) of N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxylethyl ester) as an amorphous white solid. ES-HRMS m/e calcd for $C_{25}H_{33}N_3O_8$ (M+Na) 526.2160, found 526.2143.


b) Preparation of 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxylethyl ester) hydrochloride salt



$C_{20}H_{26}ClN_3O_6$
Mol. Wt.: 439.90

The solid N-[(1,1-dimethylethoxy)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxylethyl ester) (1.4 mmol, 0.705 g) was treated with 4N hydrochloric acid in dioxane (20 mmol, 5 mL) at room temperature and the solution was stirred for 2 h as a white precipitate was formed. The mixture was diluted with diethyl ether and dichloromethane and the solids were collected by filtration washing with diethyl ether. After air drying, 0.63 g (99% yield) of 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxylethyl ester) hydrochloride was obtained as an amorphous gray solid. ES-HRMS m/e calcd for $C_{20}H_{26}N_3O_6$ (M+H) 404.1816, found 404.1818.

c) Preparation of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxylethyl ester)




To a suspension of 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxylethyl ester) hydrochloride salt (0.723 mmol, 399 mg) and 2,6-dichlorobenzoyl chloride (0.8 mmol, 0.17 g) in dichloromethane (5 mL) was added diisopropylethylamine (3.2 mmol, 0.45 mL) at room temperature. After 5 min, everything went into solution and the clear yellow solution was stirred for 48 h at room temperature. The resulting light brown solution was diluted with dichloromethane (50 mL). The dichloromethane layer was washed successively with 1N hydrochloric acid (2 x 50 mL), saturated sodium bicarbonate solution (50 mL), and brine solution (50 mL) and was dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the solvent gave the crude product, which was purified by silica gel chromatography using a Biotage (40s)

column to afford 0.312 g (67% yield) of N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxylethyl ester as a white solid: mp 168-170°C. ES-HRMS m/e calcd for $C_{27}H_{27}Cl_2N_3O_7$ (M+Na) 598.1118, found 598.1122.

5

Example 34. Preparation of N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxylethyl ester




N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxylethyl ester was prepared from 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxylethyl ester and 2-chloro-6-methylbenzoyl chloride using the general procedures described in example 33 and was obtained as a white solid: mp 84-88°C. ES-HRMS m/e calcd for $C_{28}H_{30}ClN_3O_7$ (M+Na) 578.1664, found 578.1665.

15

Example 35. Preparation of L-prolyl-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine


a) Preparation of L-[(1,1-dimethylethoxy)carbonyl]-prolyl-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester

20




To a suspension of 4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester hydrochloride salt (7 mmol, 2.57 g), HBTU (8.75 mmol, 3.32 g) and L-[(1,1-dimethylethoxy)carbonyl]-proline (8.75 mmol, 1.88 g) in DMF (28 mL) was added diisopropylethylamine (21 mmol, 3.65 mL) at room temperature. After 2 min, everything went into solution and the yellow clear solution was stirred for 48 h at room temperature. The resulting dark-brown solution was diluted with ethyl acetate (100 mL). The ethyl acetate layer was washed successively with 1N hydrochloric acid (2 x 50 mL), saturated sodium bicarbonate solution (100 mL), and brine solution (100 mL) and was dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the solvent gave the crude product which was purified by silica gel column chromatography using a Biotage (40m) column to afford 2.5 g (67% yield) of L-[(1,1-dimethylethoxy)carbonyl]-prolyl-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester as an amorphous white solid. ES-HRMS m/e calcd for $\text{C}_{27}\text{H}_{36}\text{N}_4\text{O}_7$ (M+Na) 551.2476, found 551.2476.

b) Preparation of L-[(1,1-dimethylethoxy)carbonyl]-prolyl-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine



To a suspension of L-[(1,1-dimethylethoxy)carbonyl]-prolyl-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine methyl ester (1.51 mmol, 800 mg) in ethanol (8 mL) was added aqueous 1.0 N sodium hydroxide (5 mL) at room temperature. The mixture was heated to 45-50°C and the resulting clear solution was stirred for 3 h. The ethanol was removed under reduced pressure and the residue was diluted with water (25 mL). The aqueous solution was washed with diethyl ether (50 mL) to remove any neutral impurities. The aqueous layer was acidified with 1.0 N HCl and the product was extracted into ethyl acetate (2 x 50 mL). The combined organic extracts were washed with brine solution (50 mL) and were dried over anhydrous magnesium sulfate. Filtration of the drying agent and concentration of the filtrate afforded 640 mg (83% yield) of L-[(1,1-dimethylethoxy)carbonyl]-prolyl-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine as an amorphous white solid. ES-HRMS m/e calcd for $C_{26}H_{34}N_4O_7$ (M+Na) 537.2320, found 537.2321.


c) Preparation of L-prolyl-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine



The solid L-[(1,1-dimethylethoxy)carbonyl]-prolyl-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine (0.89 mmol, 462 mg) was treated with 4N hydrochloric acid in dioxane (16 mmol, 4 mL) at room temperature and the solution was stirred for 3 h. Then, the solvent was removed under vacuum and the residue was dried under high vacuum to afford a crude residue which was triturated with dichloromethane, diethyl ether and acetonitrile to obtain 395 mg (99% yield) of L-prolyl-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine hydrochloride


salt as an amorphous light yellow solid. ES-HRMS m/e calcd for $C_{21}H_{26}N_4O_5$ ($M+H$) 415.1976, found 415.1976.

Example 36. N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine ethyl ester




To a suspension of N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine (0.743 mmol, 340 mg) and sodium bicarbonate (5.94 mmol, 499 mg) in DMF (3.8 mL) was added iodoethane (5.94 mmol, 0.475 mL) at room temperature. The mixture was stirred for 48 h at room temperature. Then, the reaction mixture was poured into water (100 mL) and was extracted with ethyl acetate (3 x 25 mL). The combined extracts were washed with brine solution (80 mL) and were dried over anhydrous sodium sulfate. Filtration of the drying agent and concentration of the filtrate gave the crude product which was purified by silica gel chromatography using a Biotage (40s) column to obtain 335 mg (93% yield) of N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine ethyl ester as a crystalline white solid: mp 218-219 °C. ES-HRMS m/e calcd for $C_{25}H_{25}F_2N_3O_5$ ($M+Na$) 508.1654, found 508.1660.

Example 37. Preparation of N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-[(N,N-diethyl)amino]ethyl ester



A mixture of N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine (1.0 mmol, 460 mg), 2-[(N,N-diethyl)amino]ethyl chloride hydrochloride (8.05 mmol, 1.43 g) and potassium carbonate (8.05 mmol, 1.11 g) in ethyl acetate (5 mL) and water (5 mL) was stirred at room temperature overnight. The layers were separated and the aqueous layer was extracted with ethyl acetate (2 x 40 mL). The combined extracts were washed with brine (50 mL) and dried over anhydrous sodium sulfate. Filtration of the drying agent and concentration of the filtrate gave the crude product which was purified by silica gel chromatography using a Biotage (40m) column to obtain 426 mg (76% yield) of N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-[(N,N-diethyl)amino]ethyl ester as an amorphous white solid. ES-HRMS m/e calcd for $C_{29}H_{34}F_2N_4O_5$ (M+H) 557.2570, found 557.2575.

Example 38. N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxo)ethyl ester



To a suspension of N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine (0.743 mmol, 340 mg) and sodium bicarbonate (5.94 mmol, 499 mg) in DMF (3.0 mL) was added 1-chloroethyl acetate (5.94 mmol, 0.73 g) at room temperature. The mixture was stirred for 48 h at room temperature. Then, the reaction mixture was poured into water (50 mL) and was extracted with ethyl acetate (3 x 25 mL). The combined extracts were washed with brine solution (80 mL) and were dried over anhydrous sodium sulfate. Filtration of the drying agent and concentration of the filtrate gave the crude product which was purified by silica gel chromatography using a Biotage (40m) column to obtain 265 mg (66% yield) of N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxylethyl ester) as an amorphous white solid. ES-HRMS m/e calcd for $C_{27}H_{27}F_2N_3O_7$ ($M+Na$) 566.1709, found 566.1710.

Bioassay Examples

Example A.

VLA-4 / VCAM-1 Screening Assay

VLA-4 antagonist activity, defined as ability to compete for binding to immobilized VCAM-1, was quantitated using a solid-phase, dual antibody ELISA. VLA-4 ($\alpha 4\beta 1$ integrin) bound to VCAM-1 was detected by a complex of anti-integrin $\beta 1$ antibody: HRP-conjugated anti-mouse IgG: chromogenic substrate (K-Blue). Initially, this entailed coating 96 well plates (Nunc Maxisorp) with recombinant human VCAM-1 (0.4 μ g in 100 μ l PBS), sealing each plate and then allowing the plates to stand at 4°C for ~18 hr. The VCAM-coated plates were subsequently blocked with 250 μ l of 1%

BSA/0.02% NaN₃ to reduce non-specific binding. On the day of assay, all plates were washed twice with VCAM Assay Buffer (200 µl/well of 50 mM Tris-HCl, 100 mM NaCl, 1 mM MnCl₂, 0.05% Tween 20; pH 7.4). Test compounds were dissolved in 100% DMSO and then diluted 1:20 in VCAM Assay Buffer supplemented with 1 mg/mL BSA (i.e., final DMSO = 5%). A series of 1:4 dilutions were performed to achieve a concentration range of 0.005 nM - 1.563 µM for each test compound. 100 µl per well of each dilution was added to the VCAM-coated plates, followed by 10 µl of Ramos cell-derived VLA-4. These plates were sequentially mixed on a platform shaker for 1 min, incubated for 2 hr at 37°C, and then washed four times with 200 µl/well VCAM Assay Buffer. 100 µl of mouse anti-human integrin β1 antibody was added to each well (0.6 µg/mL in VCAM Assay Buffer + 1 mg/mL BSA) and allowed to incubate for 1 hr at 37°C. At the conclusion of this incubation period, all plates were washed four times with VCAM Assay Buffer (200 µl/well). A corresponding second antibody, HRP-conjugated goat anti-mouse IgG (100 µl per well @ 1:800 dilution in VCAM Assay Buffer + 1 mg/mL BSA), was then added to each well, followed by a 1 hr incubation at room temperature and concluded by three washes (200 µl/well) with VCAM Assay Buffer. Color development was initiated by addition of 100 µl K-Blue per well (15 min incubation, room temp) and terminated by addition of 100 µl Red Stop Buffer per well. All plates were then read in a UV/Vis spectrophotometer at 650 nm. Results were calculated as % inhibition of total binding (i.e., VLA-4 + VCAM-1 in the absence of test compound). The results are provided in the following Table I (A = IC₅₀ < 1 nM, B = IC₅₀ < 10 nM):

Table I

Compound of Example	Activity in VCAM/VLA-4 ELISA Assay
2	A
3	B
4	A
5	A

Example B.

Ramos (VLA-4) / VCAM-1 Cell-Based Screening Assay Protocol

Materials:

5 Soluble recombinant human VCAM-1 (mixture of 5- and 7-Ig domain) was purified from CHO cell culture media by immunoaffinity chromatography and maintained in a solution containing 0.1 M Tris-glycine (pH 7.5), 0.1 M NaCl, 5 mM EDTA, 1 mM PMSF, 0.02% 0.02% NaN₃ and 10 µg/mL leupeptin. Calcein-AM was purchased from Molecular Probes Inc.

Methods:

10 VLA-4 ($\alpha 4\beta 1$ integrin) antagonist activity, defined as ability to compete with cell-surface VLA-4 for binding to immobilized VCAM-1, was quantitated using a Ramos-VCAM-1 cell adhesion assay. Ramos cells bearing cell-surface VLA-4, were labeled with a fluorescent dye (Calcein-AM) and allowed to bind VCAM-1 in the presence or absence of test compounds. A reduction in fluorescence intensity associated with adherent cells (% inhibition) reflected competitive inhibition of VLA-4 mediated cell adhesion by the test compound.

20 Initially, this entailed coating 96 well plates (Nunc Maxisorp) with recombinant human VCAM-1 (100 ng in 100 µl PBS), sealing each plate and allowing the plates to stand at 4°C for 18 hr. The VCAM-coated plates were subsequently washed twice with 0.05% Tween-20 in PBS, and then blocked for 1hr (room temperature) with 200 µl of Blocking Buffer (1% BSA/0.02% thimerosal) to reduce non-specific binding. Following the incubation with Blocking Buffer, plates were inverted, blotted and the remaining buffer aspirated. Each plate was then washed with 300 µl PBS, inverted and the remaining PBS aspirated.

30 Test compounds were dissolved in 100% DMSO and then diluted 1:25 in VCAM Cell Adhesion Assay Buffer (4 mM CaCl₂, 4 mM MgCl₂ in 50 mM TRIS-HCl, pH 7.5) (final DMSO = 4%). A series of eight 1:4 dilutions were performed for each

compound (general concentration range of 1 nM - 12,500 nM). 100 μ l/well of each dilution was added to the VCAM-coated plates, followed by 100 μ l of Ramos cells (200,000 cells/well in 1% BSA/PBS). Plates containing test compounds and Ramos cells were allowed to incubate for 45 min at room temperature, after which 165 μ l/well PBS was added. Plates were inverted to remove non-adherent cells, blotted and 300 μ l/well PBS added. Plates were again inverted, blotted and the remaining buffer gently aspirated. 100 μ l Lysis Buffer (0.1% SDS in 50 mM TRIS-HCl, pH 8.5) was added to each well and agitated for 2 min on a rotary shaking platform. The plates were then read for fluorescence intensity on a Cytofluor 2300 (Millipore) fluorescence measurement system (excitation = 485 nm, emission = 530 nm). The results are shown in the following Table II, where (A = IC₅₀ < 100 nM, B = IC₅₀ < 10000 nM):

Table II

Compound of Example	Activity in VCAM/VLA-4 Ramos Cell Assay
2	A
3	A
4	A
5	B
6	A
7	A
8	A
9	A
10	A
11	A
12	A
13	A
14	B
15	A
16	A
17	A
18	A
19	A

20	A
21	A
22	A
23	A
24	B
25	A
26	B
27	B
28	A
29	A
30	B
31	B
32	B
33	B
34	B
35	A

Table II

Example C

Alpha4-Beta7 Assay Protocol

Two weeks to one day prior to the assay, Nunc high-binding F96 Maxisorp immuno plates, #442404 or #439454, were coated with 25ng/well (0.25µg/ml) MadCAM in a volume of 100µl/well. The plates were covered with sealer and wrapped in saran wrap followed by incubation in the refrigerator for at least 24 hours. The coating buffer employed was: 10mM carbonate/bicarbonate buffer made up from: 0.8 g/L sodium carbonate and 1.55 g/L sodium bicarbonate adjusted to pH 9.6 with 1 N HCl.

Assay buffers consisted of the following:

- 10 Wash Buffer: 0.05% Tween 20 in PBS
- Blocking Buffer: 1% Nonfat Dry Milk in PBS
- Labeling Buffer: PBS
- Cell Buffer: RPMI 1640 medium (no additives)
- Binding Buffer: 1.5mM CaCl₂
- 15 0.5mM MnCl₂
- 50mM TRIS-HCl; add NaOH dropwise to pH 7.5
- Bring to volume in H₂O
- Adjust to pH 7.5
- Dilution Buffer: 4% DMSO in Binding Buffer

20 Plates were washed 2X with wash buffer and then blocked at room temperature for at least 1 hour with Blocking Buffer. Sealed plates were sometimes blocked overnight in the refrigerator. Plates were then washed with PBS and hand blotted dry. Remaining liquid was aspirated from the wells.

25 Sufficient RPMI 8866 cells were removed from stock for assay (2×10^6 cells/ml x 10ml/plate x number of plates) and placed in a centrifuge tube. The tubes were filled to volume with PBS and were spun at 200 x G for 8 minutes. The buffer was poured off and the cells were resuspended to 10×10^6 /ml in PBS and a stock solution of calcein in DMSO (5mg/mL) was added at 5µl/ml of cell suspension. The suspension was incubated for 30 minutes at 37° C in dark. The cells were then washed with PBS. The

30 PBS was poured off and the cells resuspended in cell buffer at a concentration of 2×10^6 cells/mL for plating in the assay.

Stock solution of test compounds at 25 x first dilution desired in 100% DMSO were prepared. First dilutions for the standard, as well as test compounds, were 1:25 into straight Binding Buffer, while the remaining serial dilutions were into Dilution Buffer (Binding Buffer/4% DMSO). Stock concentrations and dilutions of compounds for screening were determined by anticipated activity.

For the assay, 129µl Binding Buffer was plated into first row of wells and 100µl Dilution Buffer was plated into remaining wells. A 5.4µl aliquot of each compound was pipetted into appropriate, labeled wells, in triplicate. The compounds were next diluted down the plate (34µl + 100µl => 4-fold dilution). For controls, 100µl of Dilution Buffer + 100µl Cell Buffer were plated into the nonspecific background wells (no cells, no compound) and 100µl Dilution Buffer + 100µl cells were plated into the total binding wells (no compound = 100% binding). Labeled cells at 2×10^6 cells/ml, 100µl/well (= 2×10^5 cells/well) were added to each well containing compound. The plates were sealed and incubated in the dark for 45 minutes at room temperature.

Following incubation, unbound cells were removed by adding 150µl PBS/well. The plates were inverted, blotted onto paper towels and washed by gently adding 200µl PBS to wells and blotting again. Remaining buffer was carefully aspirated from the wells. A final 100µl PBS was added to each well.


The plates were then read for fluorescence intensity on a Cytofluor 2300 (Millipore) fluorescence measurement system (excitation = 485 nm, emission = 530 nm). IC_{50} s of each compound were determined by linear regression analysis. The results are shown in the following table III:

Compound of Example	Activity in MadCAM/RPMI Cell Assay (A = IC ₅₀ < 100 nM, B = IC ₅₀ < 10000 nM, C = IC ₅₀ < 5,000 nM)
6	A
7	A
8	A
9	A
10	B
11	C
12	A
13	B
14	C
15	B
16	B
17	B
18	A
19	B
20	B
21	B
22	B
23	B
24	B
25	B
26	B
27	B


Table III

Claims

1. A compound of the formula I:



wherein R₁ is a group of the formula Y-1




Y-1

wherein R₂₂ and R₂₃ are independently hydrogen, lower alkyl, lower alkoxy, cycloalkyl, aryl, arylalkyl, nitro, cyano, lower alkylthio, lower alkylsulfinyl, lower alkyl sulfonyl, lower alkanoyl, halogen, or perfluorolower alkyl and at least one of R₂₂ and R₂₃ is other than hydrogen; and R₂₄ is hydrogen, lower alkyl, lower alkoxy, aryl, nitro, cyano, lower alkyl sulfonyl, or halogen; or

R₁ is a group of the formula Y-2, which is a five or six membered heteroaromatic ring bonded via a carbon atom to the amide carbonyl wherein said ring contains one, two or three heteroatoms selected from the group consisting of N, O and S and one or two atoms of said ring are independently substituted by lower alkyl, cycloalkyl, halogen, cyano, perfluoroalkyl, or aryl and at least one of said substituted atoms is adjacent to the carbon atom bonded to the amide carbonyl; or

R₁ is a group of formula Y-3 which is a 3-7 membered ring of the formula:



Y-3

wherein R₂₅ is lower alkyl, unsubstituted or fluorine substituted lower alkenyl, or a group of formula R₂₆—(CH₂)_e, R₂₆ is aryl, heteroaryl, azido, cyano, hydroxy, lower alkoxy, lower alkoxy carbonyl, lower alkanoyl, lower alkylthio, lower alkyl sulfonyl, lower alkyl sulfinyl, perfluoro lower alkanoyl, nitro, or R₂₆ is a group of formula -NR₂₈R₂₉, wherein R₂₈ is hydrogen or lower alkyl, R₂₉ is hydrogen, lower alkyl, lower alkoxy carbonyl, lower alkanoyl, aroyl, perfluoro lower alkanoylamino, lower alkyl sulfonyl, lower alkylaminocarbonyl, arylaminocarbonyl; or R₂₈ and R₂₉, taken together with the attached nitrogen atom, form a 4, 5 or 6-membered saturated heterocyclic ring optionally containing one additional heteroatom selected from O, S, and N—R₄₀,

Q is -(CH₂)_fO-, -(CH₂)_fS-, -(CH₂)_fN(R₂₇)-, -(CH₂)_f,

R₂₇ is H, lower alkyl, aryl, lower alkanoyl, aroyl or lower alkoxy carbonyl,

R₄₀ is H, lower alkyl, aryl, lower alkanoyl, aroyl or lower alkoxy carbonyl

the carbon atoms in the ring are unsubstituted or substituted by lower alkyl or halogen,

e is an integer from 0 to 4, and

f is an integer from 0 to 3;

R₂ is hydrogen, lower alkyl, substituted lower alkyl, arylalkyl, or aryl;


R₃ is hydrogen, lower alkyl, substituted lower alkyl, arylalkyl, or aryl;

R₄ is hydrogen, halogen, lower alkyl, substituted lower alkyl, or aryl;

R₅ is hydrogen, lower alkyl, chloro, or lower alkoxy;

R₆ is hydrogen, lower alkyl, lower alkylcarbonyloxy lower alkyl, substituted lower alkyl,

or R₆ is a group of formula P-3:




P-3

5

wherein R₃₂ is hydrogen or lower alkyl; R₃₃ is hydrogen, lower alkyl, aryl;
R₃₄ is hydrogen or lower alkyl; h is an integer from 0 to 2; g is an integer from
0 to 2; the sum of h and g is 1 to 3; or

10

or R₆ is a group of formula P-4:



wherein R₃₂, g, and h are as previously defined; Q' is O, S, -(CH₂)_j-, or a
group of the formula N-R₃₅; wherein R₃₅ is hydrogen, lower alkyl, lower
alkanoyl, lower alkoxy carbonyl; j is 0, 1 or 2; or its pharmaceutically
acceptable salts.

15

2. A compound of claim 1 wherein

20

R² is hydrogen, lower alkyl, substituted lower alkyl, arylalkyl, or aryl;
R³ is hydrogen, lower alkyl, substituted lower alkyl, arylalkyl, or aryl; and
R⁴ hydrogen, lower alkyl, perfluoro lower alkyl, or aryl.

3. A compound of claim 1, wherein

25

R² is hydrogen, lower alkyl, substituted lower alkyl, or aryl;
R³ is hydrogen, lower alkyl, substituted lower alkyl, or aryl; and
R⁴ hydrogen, halogen, lower alkyl, substituted lower alkyl, or aryl.

4. A compound according to any one of claims 1-3, wherein

R_2 is hydrogen, lower alkyl, substituted lower alkyl or aryl;

R_3 is hydrogen, lower alkyl, substituted lower alkyl, or aryl; and


R_4 is hydrogen, lower alkyl, perfluoro lower alkyl, or aryl.

5. A compound of any one of claims 1-4, wherein R_4 is hydrogen, lower alkyl, or perfluoro lower alkyl.

6. A compound of any one of claims 1-4, wherein R_1 is a group of the formula Y-1 wherein R_{22} and R_{23} are independently lower alkyl or halogen; and R_{24} is hydrogen.

7. A compound of any one of claims 1-4, wherein R_1 is a group of the formula Y-1 wherein R_{22} and R_{23} are independently hydrogen or halogen; and R_{24} is lower alkoxy.

8. A compound of any one of claims 1-4, wherein R_1 is a group of formula Y-3 which is a 3-7 membered ring of the formula:




20 Y-3

wherein R_{25} is a group of formula $R_{26}-(CH_2)_e$, wherein R_{26} is lower alkoxy,

25 Q is $-(CH_2)_f$, e is an integer from 0 to 4, and f is an integer from 0 to 3.

9. A compound of any one of claims 1-4 of the formula I:



wherein R_1 is as defined in claim 1;

R_2 is lower alkyl;

5 R_3 is lower alkyl;

R_4 is hydrogen, perfluoro lower alkyl, or lower alkyl;


R_5 is hydrogen or lower alkyl; and

R_6 is hydrogen, lower alkyl, lower alkylcarbonyloxy lower alkyl,

or R_6 is a group of formula P-3 as defined in claim 1

10 or R_6 is a group of formula P-4 as defined in claim 1.

10. A compound of claim 9 wherein R_1 is a group of the formula Y-1



Y-1

15 wherein R_{22} and R_{23} are independently perfluoro lower alkyl, lower alkyl, or halogen; and R_{24} is hydrogen.

11. A compound of claim 10 wherein R^2 and R^3 are lower alkyl; R^4 is hydrogen or lower alkyl, and R_5 and R_6 are hydrogen.

20

12. A compound of claim 11 which is selected from

N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

N-[(2-bromo-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

5 N-[(2-ethyl-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

N-[[2-(2-methylethyl)-6-methylphenyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

10 N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

N-[[2-fluoro-6-(trifluoromethyl)phenyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

N-[[2,6-di-(2-methylethyl)phenyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

15 N-[(2-chloro-6-ethylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

N-[(2-chloro-6-propylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

20 N-[[2-chloro-6-(2-methylethyl)phenyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine; or

N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine.

25


13. A compound of claim 11 which is selected from

N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine;

30 N-[(2-bromo-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine; or

N-[(2-bromo-5-methoxyphenyl)carbonyl]-4-[1,3-dimethyl-2,4-dioxo-5-pyrimidinyl]-L-phenylalanine.

14. A compound of claim 10 wherein R^2 and R^3 are lower alkyl; R^4 is hydrogen or lower alkyl, R_5 is hydrogen, and R_6 is hydrogen, lower alkylcarbonyloxy lower alkyl, lower alkyl, or R_6 is a group of formula P-3:




5

P-3

wherein R_{32} is hydrogen or lower alkyl; R_{33} is hydrogen, lower alkyl, aryl; R_{34} is hydrogen or lower alkyl; h is an integer from 0 to 2; g is an integer from 0 to 2; the sum of h and g is 1 to 3; or

10

R_6 is a group of formula P-4:



15

wherein R_{32} , g , and h are as previously defined; Q' is O, S, $-(CH_2)_j-$, or a group of the formula N-R35; wherein R_{35} is hydrogen.

15. A compound of claim 14 wherein R_6 is lower alkyl.

20

16. A compound of claim 15 which is

N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine ethyl ester;

N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine ethyl ester; or

25

N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine ethyl ester.

17. A compound of claim 14 wherein R_6 is lower alkylcarbonyloxy lower alkyl.

18. A compound of claim 17 which is

N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxylethyl ester;

5 N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxylethyl ester; or

N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 1-(acetoxylethyl ester.

10 19. A compound of claim 14 wherein R^6 is a group of the formula P-3 wherein R^{32} is hydrogen; R^{33} and R^{34} are lower alkyl; h is 1; and g is 0.

20. A compound of claim 19 which is

N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-[(N,N-diethylamino)ethyl ester;

15 N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-[(N,N-diethylamino)ethyl ester; or

N-[(2,6-difluorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-[(N,N-diethylamino)ethyl ester.

20 21. A compound of claim 14 wherein R^6 is a group of the formula P-4 wherein R^{32} is hydrogen; h is 1; g is 0; and Q' is O.

22. A compound of claim 21 which is

25 N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-(4-morpholino)ethyl ester; or

N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-L-phenylalanine 2-(4-morpholino)ethyl ester.

30 23. A compound of claim 10 wherein R^2 and R^3 are lower alkyl; R^4 is perfluoro lower alkyl, and R^5 and R^6 are hydrogen.

24. A compound of claim 23 which is

N-[1-(2,6-dichlorophenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine;

N-[(2-chloro-6-methylphenyl)carbonyl]-4-[1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl]-L-phenylalanine; or

5 N-[[2-fluoro-6-(trifluoromethyl)phenyl]carbonyl]-4-(1,3-dimethyl-2,4-dioxo-6-(trifluoromethyl)-5-pyrimidinyl)-L-phenylalanine.

25. A compound of claim 10 wherein R^2 and R^3 are lower alkyl; R^4 is hydrogen; R^5 is lower alkyl, and R^6 is hydrogen.

10

26. A compound of claim 25 which is

N-[(2-chloro-6-methylphenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine;


15

N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine; or

N-[(2,6-dichlorophenyl)carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-5-pyrimidinyl)-3-methyl-L-phenylalanine.

27. A compound of claim 9 wherein R_1 is a group of formula Y-3 which is a 3-7 membered ring of the formula:

20



Y-3

wherein R_{25} is a group of formula $R_{26}-(CH_2)_e$, wherein R_{26} is lower alkoxy, Q is $-(CH_2)_f$, e is an integer from 0 to 4, and f is an integer from 0 to

25

3.

28. A compound of claim 27 wherein R^2 and R^3 are lower alkyl, R^4 is hydrogen or lower alkyl; and R^5 and R^6 are hydrogen.

30

29. A compound of claim 28 which is

4-(1,3-dimethyl-2,4-dioxo-5-pyrimidinyl)]-N-[[1-(2-methoxyethyl)-
cyclopentyl]carbonyl]-L-phenylalanine;

N-[[1-(2-methoxyethyl)cyclopentyl]carbonyl]-4-(1,3,6-trimethyl-2,4-dioxo-
5-pyrimidinyl)-L-phenylalanine; or

4-(1,3-diethyl-6-methyl-2,4-dioxo-5-pyrimidinyl)-N-[[1-(2-
methoxyethyl)cyclopentyl]carbonyl]-L-phenylalanine.

30. A compound of claim 9 wherein R_2 and R_3 are lower alkyl; and R_4 is
hydrogen, and R_5 and R_6 are hydrogen.

31. A compound of claim 30 wherein R_1 is a group of the formula Y-1.

32. A compound of claim 30 wherein R_1 is a group of the formula Y-1 wherein
 R_{22} and R_{23} are independently lower alkyl or halogen; and R_{24} is hydrogen.


33. A compound of claim 31 wherein R_1 is a group of the formula Y-1 wherein
 R_{22} and R_{23} are independently hydrogen or halogen; and R_{24} is lower alkoxy.

34. A compound of claim 30 wherein R_1 is a five or six membered heteroaromatic
ring bonded via a carbon atom to the amide carbonyl wherein said ring contains one,
two or three heteroatoms selected from the group consisting of N, O and S and one
or two atoms of said ring are independently substituted by lower alkyl, cycloalkyl,
halogen, cyano, perfluoroalkyl, or aryl and at least one of said substituted atoms is
adjacent to the carbon atom bonded to the amide carbonyl.

35. A compound of claim 30 wherein R_1 is a group of formula Y-3.

36. A compound of claim 35 wherein R_1 is a group of formula Y-3 wherein R_{25} is a
group of formula $R_{26}-(CH_2)_e$, wherein R_{26} is lower alkoxy, Q is $-(CH_2)_f$, e is an
integer from 0 to 4, and f is an integer from 0 to 3.

37. A compound of formula



wherein R^2 - R^5 are as defined in claim 1 and P_1 and P_2 each are a protecting group.

38. A compound according to any of claims 1-36 for use as a medicament.

39. A pharmaceutical composition comprising a compound according to any one of claims 1-36 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

40. A compound according to any one of claims 1-36 for use in the treatment of disease states mediated by the binding of VCAM-1 or VLA-4 or VLA-4-expressing cells, especially in the treatment of rheumatoid arthritis, multiple sclerosis, inflammatory bowel disease and asthma.

41. The use of a compound according to any one of claims 1-36 or a pharmaceutically acceptable salt thereof, for the preparation of a medicament for the treatment of disease states mediated by the binding of VCAM-1 or VLA-4 or VLA-4-expressing cells, especially in the treatment of rheumatoid arthritis, multiple sclerosis, inflammatory bowel disease and asthma.

42. A process for the preparation of a pharmaceutical composition which process comprises bringing a compound according to any one of claims 1-36, or a pharmaceutically acceptable salt thereof, and a compatible pharmaceutical carrier into a galenical administration form.

43. The use of a compound according to any one of claims 1-36, or a pharmaceutically acceptable salt thereof, in the preparation of a medicament for the treatment of rheumatoid arthritis, multiple sclerosis, inflammatory bowel disease and asthma.

5

44. The invention substantially as hereinbefore described, especially with reference to the new compounds, intermediates, pharmaceutical compositions and uses thereof.

10