







FIG. 1 (continued)

(c)




FIG. 1 (continued)





FIG. 2


(c)

Time point (pio)		Acute	Early convalescent	Late convalescent	Early recovery	Late recovery	Full recovery
		2-7 days	10-14 days	1 month	3 months	5-6 months	1 year
Anti-ZIKV IgM	No of patients	58	43	45	41	38	32
	Positive	43 (74%)	38 (88%)	36 (80%)	33 (80%)	33 (87%)	21 (66%)
Anti-ZIKV IgG	No of patients	58	43	45	41	38	32
	Positive	15 (26%)	5 (12%)	9 (20%)	8 (20%)	5 (13%)	11 (34%)
Anti-ZIKV IgG	No of patients	58	43	45	41	38	32
	Positive	52 (90%)	42 (98%)	43 (96%)	40 (98%)	38 (100%)	30 (94%)
Anti-ZIKV IgG	No of patients	58	43	45	41	38	32
	Negative	6 (10%)	1 (2%)	2 (4%)	1 (2%)	0 (0%)	2 (6%)

(d)



(e)



(f)




FIG. 2 (continued)




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 6 (continued)




FIG. 6 (continued)




FIG. 7




FIG. 8




FIG. 8 (continued)




FIG. 8 (continued)

(a)




FIG. 9





FIG. 9 (continued)

(c)



(d)




FIG. 9 (continued)