

(2-UREIDOACETAMIDO)ALKYL DERIVATIVES AS FORMYL PEPTIDE RECEPTOR 2 MODULATORS

By inventors: Richard L. Beard, Tien Duong, John E. Donello,
Veena Viswanath and Michael E. Garst

RELATED APPLICATION

This application claims the benefit of U.S. Provisional Application Serial No. 61/624,495, filed April 16, 2012, the disclosure of which is hereby incorporated in its entirety herein by reference.

FIELD OF THE INVENTION

The present invention relates to (2-ureidoacetamido)alkyl derivatives, processes for preparing them, pharmaceutical compositions containing them and their use as pharmaceuticals, as modulators of the N-formyl peptide receptor 2 (FPR2). The invention relates specifically to the use of these compounds and their pharmaceutical compositions to treat disorders associated with the N-formyl peptide receptor 2 modulation.

BACKGROUND OF THE INVENTION

The N-formyl peptide receptor 2 also known as N-formyl peptide receptor like-1 (FRL-1), is a G protein-coupled receptor that is expressed on inflammatory cells such as monocytes and neutrophils, as well as T cells and has been shown to play a critical role in leukocyte trafficking during inflammation and human pathology. FPR2 is an exceptionally promiscuous receptor that responds to a large array of exogenous and endogenous ligands, including Serum amyloid A (SAA), chemokine variant sCK β 8-1, the neuroprotective peptide, anti-inflammatory eicosanoid lipoxin A4 (LXA4) and glucocorticoid-modulated protein annexin A1. FPR2 transduces anti-inflammatory effects of LXA4 in many systems, but it also can mediate the pro-inflammatory signaling cascade of peptides such as SAA. The ability of the receptor to mediate two opposite effects is proposed to be a result of different receptor domains used by different agonists.

Parmentier, Marc et al. Cytokine & Growth Factor Reviews 17 (2006) 501–519.

Activation of FPR2 by lipoxin A4 or its analogs and by Annexin I protein has been shown to result in anti-inflammatory activity by promoting active resolution of inflammation which involves inhibition of polymorphonuclear neutrophils (PMNs) and eosinophils


migration and also stimulate monocyte migration enabling clearance of apoptotic cells from the site of inflammation in a nonphlogistic manner. In addition, FPR2 has been shown to inhibit NK cytotoxicity and promote activation of T cells which further contributes to down regulation of tissue damaging inflammatory signals. FPR2/ LXA4 interaction has been shown to be beneficial in experimental models of ischemia reperfusion, angiogenesis, dermal inflammation, chemotherapy-induced alopecia, ocular inflammation such as endotoxin-induced uveitis, corneal wound healing, re-epithelialization etc. FPR2 thus represents an important novel pro-resolventary molecular target for the development of new therapeutic agents in diseases with excessive inflammatory responses.

10 SUMMARY OF THE INVENTION

A group of (2-ureidoacetamido)alkyl derivatives, which are potent and selective FPR2 modulators has been discovered. As such, the compounds described herein are useful in treating a wide variety of disorders associated with modulation of the FPR2 receptor. The term “modulator” as used herein, includes but is not limited to: receptor agonist, antagonist, inverse agonist, inverse antagonist, partial agonist, and partial antagonist.

This invention describes compounds of Formula I, which have FPR2 receptor biological activity. The compounds in accordance with the present invention are thus of use in medicine, for example in the treatment of humans with diseases and conditions that are alleviated by FPR2 receptor modulation.

In one aspect, the invention provides a compound represented by **Formula I** or the individual enantiomers, the individual diastereoisomers, the individual zwitterions or the pharmaceutically acceptable salts thereof:



Formula I

wherein:

n is 0 or 1;

R¹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, -N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN or -NO₂;

5 R² is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN or -NO₂;


R³ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN, -NO₂, -CF₃, -S(O)R¹⁵ or -S(O)₂R¹⁶;

R⁴ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN or -NO₂;

10 R⁵ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, -N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN or -NO₂;

R⁶ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted heterocycle, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted C₆₋₁₀ aryl, substituted or unsubstituted C₃₋₈ cycloalkenyl or -CH₂R¹⁹;

15



S(O)R¹⁵, -C(O)N(H)(CN), -S(O)₂NHS(O)₂R²⁵, -C(O)N(H)S(O)₂R¹⁹, -S(O)₂R¹⁶ or -

20 P(O)R¹⁷R¹⁸;

R⁸ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R⁹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R¹⁰ is hydrogen, or substituted or unsubstituted C₁₋₈ alkyl ;

R¹¹ is hydrogen , substituted or unsubstituted C₁₋₈ alkyl or -CF₃;

R¹² is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, -OR²⁴ or -NR⁸R⁹;

10 R¹³ is -OR²²;

R¹⁴ is -OR²³;

R¹⁵ is substituted or unsubstituted C₁₋₈ alkyl;

R¹⁶ is substituted or unsubstituted C₁₋₈ alkyl, -NR⁸R⁹ , -NHS(O)₂R¹⁹ or hydroxyl;

R¹⁷ is -OR¹⁰ or -NR⁸R⁹;

15 R¹⁸ is -OR¹⁰ or -NR⁸R⁹;

R¹⁹ is substituted or unsubstituted heterocycle, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted C₆₋₁₀ aryl or substituted or unsubstituted C₃₋₈ cycloalkenyl;

R²⁰ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R²¹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R²² is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, or together with R²³ can form a substituted or unsubstituted cycle;

R²³ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, or together with R²² can form a substituted or unsubstituted cycle;

R²⁴ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl; and

R²⁵ is substituted or unsubstituted aromatic heterocycle, or substituted or unsubstituted aromatic C₆₋₁₀ aryl;


R^{26} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10} aryl;

R^{27} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10} aryl;

R^{28} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10} aryl;

R^{29} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10} aryl; with the proviso:

the compound of Formula I is not of structures:



In another aspect, the invention provides a compound represented by **Formula I**, wherein:

n is 0 or 1;

- R^1 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, halogen, $-NR^8R^9$, $-N(R^{21})C(O)R^{20}$, $-OR^{10}$, $-OC(O)R^{21}$, $-SR^{11}$, $-C(O)R^{12}$, $-CN$ or $-NO_2$;
- R^2 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, halogen, $-NR^8R^9$, $-N(R^{21})C(O)R^{20}$, $-OR^{10}$, $-OC(O)R^{21}$, $-SR^{11}$, $-C(O)R^{12}$, $-CN$ or $-NO_2$;
- 5 R^3 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, halogen, $-NR^8R^9$, $-N(R^{21})C(O)R^{20}$, $-OR^{10}$, $-OC(O)R^{21}$, $-SR^{11}$, $-C(O)R^{12}$, $-CN$, $-NO_2$, $-CF_3$, $-S(O)R^{15}$ or $-S(O)_2R^{16}$;
- R^4 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, halogen, $-NR^8R^9$, $-N(R^{21})C(O)R^{20}$, $-OR^{10}$, $-OC(O)R^{21}$, $-SR^{11}$, $-C(O)R^{12}$, $-CN$ or $-NO_2$;
- R^5 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, halogen, $-NR^8R^9$, $-N(R^{21})C(O)R^{20}$,
10 $-OR^{10}$, $-OC(O)R^{21}$, $-SR^{11}$, $-C(O)R^{12}$, $-CN$ or $-NO_2$;
- R^6 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted heterocycle, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted C_{6-10} aryl, substituted or unsubstituted C_{3-8} cycloalkenyl or $-CH_2R^{19}$;
- R^7 is $-P(O)R^{17}R^{18}$;
- 15 R^8 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10} aryl;
- R^9 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10}
20 aryl;
- R^{10} is hydrogen or substituted or unsubstituted C_{1-8} alkyl ;
- R^{11} is hydrogen, substituted or unsubstituted C_{1-8} alkyl or $-CF_3$;
- R^{12} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, $-OR^{24}$ or $-NR^8R^9$;
- R^{13} is $-OR^{22}$;
- 25 R^{14} is $-OR^{23}$;
- R^{15} is substituted or unsubstituted C_{1-8} alkyl;
- R^{16} is substituted or unsubstituted C_{1-8} alkyl, $-NR^8R^9$, $-NHS(O)_2R^{19}$ or hydroxyl;
- R^{17} is OR^{10} or NR^8R^9 ;
- R^{18} is OR^{10} or NR^8R^9 ;
- 30 R^{19} is substituted or unsubstituted heterocycle, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted C_{6-10} aryl or substituted or unsubstituted C_{3-8} cycloalkenyl;
- R^{20} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10} aryl;

R²¹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R²² is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, or together with R²³ can form a substituted or unsubstituted cycle;

R²³ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, or together with R²² can form a substituted or unsubstituted cycle;

R²⁴ is hydrogen substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl; and

R²⁵ is substituted or unsubstituted aromatic heterocycle, or substituted or unsubstituted aromatic C₆₋₁₀ aryl;

R²⁶ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R²⁷ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R²⁸ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R²⁹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl.

In another aspect, the invention provides a compound represented by **Formula I**, wherein:
n is 0 or 1;

R¹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, -N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN or -NO₂;

R² is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, -N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN or -NO₂;

R³ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, -N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN, -NO₂, -CF₃, -S(O)R¹⁵ or -S(O)₂R¹⁶;

- R^4 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, halogen, $-NR^8R^9$, $-N(R^{21})C(O)R^{20}$, $-OR^{10}$, $-OC(O)R^{21}$, $-SR^{11}$, $-C(O)R^{12}$, $-CN$ or $-NO_2$;
- R^5 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, halogen, $-NR^8R^9$, $-N(R^{21})C(O)R^{20}$, $-OR^{10}$, $-OC(O)R^{21}$, $-SR^{11}$, $-C(O)R^{12}$, $-CN$ or $-NO_2$;
- 5 R^6 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted heterocycle, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted C_{6-10} aryl, substituted or unsubstituted C_{3-8} cycloalkenyl or $-CH_2R^{19}$;
- R^7 is $-SR^{11}$, $-S(O)R^{15}$ or $-S(O)_2R^{16}$;
- R^8 is hydrogen, substituted or unsubstituted C_{1-8} alkyl substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10} aryl;
- 10 R^9 is hydrogen, substituted or unsubstituted C_{1-8} alkyl substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10} aryl;
- 15 R^{10} is hydrogen or substituted or unsubstituted C_{1-8} alkyl;
- R^{11} is hydrogen, substituted or unsubstituted C_{1-8} alkyl or $-CF_3$;
- R^{12} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, $-OR^{24}$ or $-NR^8R^9$;
- R^{13} is $-OR^{22}$;
- R^{14} is $-OR^{23}$;
- 20 R^{15} is substituted or unsubstituted C_{1-8} alkyl;
- R^{16} is substituted or unsubstituted C_{1-8} alkyl, $-NR^8R^9$, $-NHS(O)_2R^{19}$ or hydroxyl;
- R^{17} is $-OR^{10}$ or $-NR^8R^9$;
- R^{18} is $-OR^{10}$ or $-NR^8R^9$;
- R^{19} is substituted or unsubstituted heterocycle, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted C_{6-10} aryl or substituted or unsubstituted C_{3-8} cycloalkenyl;
- 25 R^{20} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10} aryl;
- R^{21} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10} aryl;
- 30 R^{22} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, or together with R^{23} can form a substituted or unsubstituted cycle;

R^{23} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, or together with R^{22} can form a substituted or unsubstituted cycle;

R^{24} is hydrogen substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10}

5 aryl; and

R^{25} is substituted or unsubstituted aromatic heterocycle, or substituted or unsubstituted aromatic C_{6-10} aryl;

R^{26} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10}

10 aryl;


R^{27} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10} aryl;

R^{28} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10}

15 aryl;

R^{29} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10} aryl; with the proviso:

20 the compound of Formula I is not of structures:



In another aspect, the invention provides a compound represented by **Formula I**, wherein:

n is 0 or 1;

R¹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, -N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹ -SR¹¹, -C(O)R¹², -CN or -NO₂;


5 R² is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, -N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN or -NO₂;

R³ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, -N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN, -NO₂, -CF₃, -S(O)R¹⁵ or -S(O)₂R¹⁶;

10 R^4 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, halogen, $-NR^8R^9$, $-N(R^{21})C(O)R^{20}$, $-OR^{10}$, $-OC(O)R^{21}$, $-SR^{11}$, $-C(O)R^{12}$, $-CN$ or $-NO_2$;

R⁵ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, -N(R²¹)C(O)R²⁰, OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN or -NO₂;

R⁶ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted heterocycle, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted C₆₋₁₀ aryl, substituted or unsubstituted C₃₋₈ cycloalkenyl or -CH₂R¹⁹;



R⁸ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R⁹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R¹⁰ is hydrogen or substituted or unsubstituted C₁₋₈ alkyl;

R¹¹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl or -CF₃;

R¹² is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, -OR²⁴ or -NR⁸R⁹;

R¹³ is -OR²²;

R¹⁴ is -OR²³;

R¹⁵ is substituted or unsubstituted C₁₋₈ alkyl;

R¹⁶ is substituted or unsubstituted C₁₋₈ alkyl, -NR⁸R⁹, -NHS(O)₂R¹⁹ or hydroxyl;

R¹⁷ is -OR¹⁰ or -NR⁸R⁹;

R¹⁸ is -OR¹⁰ or -NR⁸R⁹;

R¹⁹ is substituted or unsubstituted heterocycle, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted C₆₋₁₀ aryl or substituted or unsubstituted C₃₋₈ cycloalkenyl;

R²⁰ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R²¹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R²² is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, or together with R²³ can form a substituted or unsubstituted cycle;

R²³ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, or together with R²² can form a substituted or unsubstituted cycle;

R²⁴ is hydrogen substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl; and

R²⁵ is substituted or unsubstituted aromatic heterocycle, or substituted or unsubstituted aromatic C₆₋₁₀ aryl;


R²⁶ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R²⁷ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R²⁸ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R²⁹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl; with the proviso:

the compound of Formula I is not of structures:



The term "alkyl", as used herein, refers to saturated, monovalent or divalent hydrocarbon moieties having linear or branched moieties or combinations thereof and containing 1 to 8 carbon atoms, unless otherwise specified. One methylene (-CH₂-) group, of the alkyl can be replaced by oxygen, sulfur, sulfoxide, nitrogen, -NH-, carbonyl, carboxyl, sulfonyl, amido, sulfonamido, by a divalent C₃₋₆ cycloalkyl, by a divalent heterocycle, or by a divalent aryl group. Alkyl groups can be independently substituted by one or more halogen (as in CF₃), hydroxyl, cycloalkyl, amine, heterocyclic, carboxylic acid, -C₂₋₆ alkenyl, -C₂₋₆ alkynyl, phosphonic acid, phosphonate, sulfonate, sulfate, sulphonic acid, phosphoric acid, nitro, amide, sulfonamide, aldehyde or ester groups.

The term "halogen", as used herein, refers to an atom of chlorine, bromine, fluorine, iodine.

The term "cycloalkyl", as used herein, refers to a monovalent or divalent group of 3 to 8 carbon atoms derived from a saturated cyclic hydrocarbon. Cycloalkyl groups can be monocyclic or polycyclic. Cycloalkyl can be independently substituted by halogen, -SC₁₋₈ alkyl, -S(O)₂C₁₋₈ alkyl, -S(O)C₁₋₈ alkyl, sulfonamide, amide, nitro, cyano, -OC₁₋₈ alkyl, -

C₁₋₈ alkyl, -C₂₋₆ alkenyl, -C₂₋₆ alkynyl, ketone, amine, C₃₋₈ cycloalkyl, aldehydes, esters, carboxylic acids, phosphonic acids, sulfonic acids or hydroxyl groups.

The term "cycloalkenyl", as used herein, refers to a monovalent or divalent group of 3 to 8 carbon atoms derived from a saturated cycloalkyl having at least one double bond.

5 Cycloalkenyl groups can be monocyclic or polycyclic. Cycloalkenyl groups can be independently substituted by halogen, -SC₁₋₈ alkyl, -S(O)₂C₁₋₈ alkyl, -S(O)C₁₋₈ alkyl, sulfonamide, amide, nitro, cyano, -OC₁₋₈ alkyl, -C₁₋₈ alkyl, -C₂₋₆ alkenyl, -C₂₋₆ alkynyl, ketone, amine, C₃₋₈ cycloalkyl, aldehydes, esters, carboxylic acid, phosphonic acid, sulfonic acid or hydroxyl groups.

10 The term "aryl" as used herein, refers to an organic moiety derived from an aromatic hydrocarbon consisting of a ring containing 6 to 10 carbon atoms by removal of one hydrogen, which can be substituted by halogen, -SC₁₋₈ alkyl, -S(O)₂C₁₋₈ alkyl, -S(O)C₁₋₈ alkyl, sulfonamide, amide, nitro, cyano, -OC₁₋₈ alkyl, -C₁₋₈ alkyl, -C₂₋₆ alkenyl, -C₂₋₆ alkynyl, ketone, amine, C₃₋₈ cycloalkyl, aldehydes, esters, carboxylic acid, phosphonic acid, sulfonic acid or hydroxyl groups.

The term "heterocycle" as used herein, refers to a 5- or 6- membered ring, which can be aromatic or non-aromatic, saturated or unsaturated, containing at least one heteroatom selected from O or N or S or combinations of at least two thereof, interrupting the carbocyclic ring structure. The heterocyclic ring can be interrupted by a C=O; the S and N heteroatoms can be oxidized. Heterocyclic ring moieties can be substituted by

20 halogen, -SC₁₋₈ alkyl, -S(O)₂C₁₋₈ alkyl, -S(O)C₁₋₈ alkyl, sulfonamide, amide, nitro, cyano, -OC₁₋₈ alkyl, -C₁₋₈ alkyl, -C₂₋₆ alkenyl, -C₂₋₆ alkynyl, ketone, amine, C₃₋₈ cycloalkyl, aldehydes, esters, carboxylic acid, phosphonic acid, sulfonic acid or hydroxyl groups. Examples of heterocycles are, but not limited to: isoxazol-3-ol-5-yl, isoxazol-5-ol-3-yl, isothiazol-3-ol-5-yl, 1-methyl-1H-pyrazol-3-ol-5-yl, 1H-tetrazole-5-yl, 1H-imidazol-1-ol-2-yl, 1H-1,2,3-triazol-1-ol-5-yl, 1H-imidazol-1-ol-5-yl, 1H-pyrazol-1-ol-5-yl, 3-hydroxy-4H-pyran-4-one-5-yl, imidazoles, triazoles, tetrazoles, oxadiazoles, isothiazoles, pyranes, 4H-pyran-4-ones, pyrroles, pyrrolidine-ones, pyrrolidine-diones, pyrazoles, isoxazoles, imidazolidine-diones.

30 The term "hydroxyl" as used herein, represents a group of formula "-OH".

The term "carbonyl" as used herein, represents a group of formula "-C(O)-".

The term "aldehyde" as used herein, represents a group of formula $-\text{C}(\text{O})\text{H}$.

The term "ketone" as used herein, represents a group of formula $-\text{C}(\text{O})\text{R}^x$ wherein R^x can be alkyl, aryl, cycloalkyl, cycloalkenyl, or heterocycle as defined above.

5 The term "ester" as used herein, represents a group of formula $-\text{C}(\text{O})\text{OR}^x$ wherein R^x can be alkyl, aryl, cycloalkyl, cycloalkenyl, or heterocycle as defined above.

The term "amine" as used herein, represents a group of formula NR^xR^y , wherein R^x and R^y can be independently H, alkyl, aryl, cycloalkyl, cycloalkenyl, or heterocycle as defined above.

The term "carboxyl" as used herein, represents a group of formula $-\text{C}(\text{O})\text{O}-$.

10 The term "sulfonyl" as used herein, represents a group of formula $-\text{SO}_2-$.

The term "sulfate" as used herein, represents a group of formula $-\text{O}-\text{S}(\text{O})_2-\text{O}-$.

The term "sulfonate" as used herein, represents a group of the formula $-\text{S}(\text{O})_2-\text{OR}^x$, wherein R^x can be hydrogen, alkyl, aryl, cycloalkyl, cycloalkenyl, heterocycle as defined above.

15 The term "carboxylic acid" as used herein, represents a group of formula $-\text{C}(\text{O})\text{OH}$.

The term "nitro" as used herein, represents a group of formula $-\text{NO}_2$.

The term "cyano" as used herein, represents a group of formula $-\text{CN}$.

20 The term "amide" as used herein, represents a group of formula $-\text{C}(\text{O})\text{NR}^x\text{R}^y$, wherein R^x and R^y can be independently H, alkyl, aryl, cycloalkyl, cycloalkenyl, heterocycle as defined above.

The term "amido" as used herein, represents a group of formula $-\text{C}(\text{O})\text{NR}^x-$, wherein R^x can be H, alkyl, aryl, cycloalkyl, cycloalkenyl, heterocycle as defined above.

25 The term "sulfonamide" as used herein, represents a group of formula $-\text{S}(\text{O})_2\text{NR}^x\text{R}^y$ wherein R^x and R^y can independently be H, alkyl, aryl, cycloalkyl, cycloalkenyl, heterocycle as defined above.

The term "sulfonamido" as used herein, represents a group of formula $-\text{S}(\text{O})_2\text{NR}^x-$ wherein R^x can be H, alkyl, aryl, cycloalkyl, cycloalkenyl, heterocycle as defined above.

The term "sulfoxide" as used herein, represents a group of formula $-\text{S}(\text{O})-$.

30 The term "phosphonic acid" as used herein, represents a group of formula $-\text{P}(\text{O})(\text{OH})_2$.

The term "phosphoric acid" as used herein, represents a group of formula $-\text{OP}(\text{O})(\text{OH})_2$.

The term "phosphonate" as used herein, represents a group of formula " $\text{-P(O)(OR}^x\text{)(OR}^y\text{)}$," wherein R^x and R^y can be independently hydrogen, alkyl, aryl, cycloalkyl, cycloalkenyl, heterocycle as defined above.

The term "sulphonic acid" as used herein, represents a group of formula " $\text{-S(O)}_2\text{OH}$ ".

5 The formula " H ", as used herein, represents a hydrogen atom.

The formula " O ", as used herein, represents an oxygen atom.

The formula " N ", as used herein, represents a nitrogen atom.

The formula " S ", as used herein, represents a sulfur atom.

Generally, n is 0 or 1.

10 Generally, R^1 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, halogen, $\text{-NR}^8\text{R}^9$, $\text{-N(R}^{21}\text{)C(O)R}^{20}$, -OR^{10} , -OC(O)R^{21} , -SR^{11} , -C(O)R^{12} , -CN or -NO_2 . Usually, R^1 is hydrogen, substituted or unsubstituted C_{1-8} alkyl or halogen. Preferred R^1 is hydrogen.


Generally, R^2 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, halogen, $\text{-NR}^8\text{R}^9$, $\text{-N(R}^{21}\text{)C(O)R}^{20}$, -OR^{10} , -OC(O)R^{21} , -SR^{11} , -C(O)R^{12} , -CN or -NO_2 . Usually, R^2 is hydrogen, substituted or unsubstituted C_{1-8} alkyl or halogen. Preferred R^2 is hydrogen.

15 Generally, R^3 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, halogen, $\text{-NR}^8\text{R}^9$, $\text{-N(R}^{21}\text{)C(O)R}^{20}$, -OR^{10} , -OC(O)R^{21} , -SR^{11} , -C(O)R^{12} , -CN , -NO_2 , -CF_3 , -S(O)R^{15} or $\text{-S(O)}_2\text{R}^{16}$. Usually, R^3 is substituted or unsubstituted C_{1-8} alkyl, halogen, -SR^{11} , -CF_3 or $\text{-S(O)}_2\text{R}^{16}$. Preferred R^3 is halogen, -SR^{11} , -CF_3 or $\text{-S(O)}_2\text{R}^{16}$. Most preferred R^3 is bromo, -SMe , -CF_3 or $\text{-S(O)}_2\text{Me}$.


20 Generally, R^4 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, halogen, $\text{-NR}^8\text{R}^9$, $\text{-N(R}^{21}\text{)C(O)R}^{20}$, -OR^{10} , -OC(O)R^{21} , -SR^{11} , -C(O)R^{12} , CN or NO_2 . Usually, R^4 is hydrogen, substituted or unsubstituted C_{1-8} alkyl or halogen. Preferred R^4 is hydrogen.

25 Generally, R^5 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, halogen, $\text{-NR}^8\text{R}^9$, $\text{-N(R}^{21}\text{)C(O)R}^{20}$, -OR^{10} , -OC(O)R^{21} , -SR^{11} , -C(O)R^{12} , CN or NO_2 . Usually, R^5 is hydrogen, substituted or unsubstituted C_{1-8} alkyl or halogen. Preferred R^5 is hydrogen.

30 Generally, R^6 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted heterocycle, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted C_{6-10} aryl, substituted or unsubstituted C_{3-8} cycloalkenyl or $\text{-CH}_2\text{R}^{19}$. Usually, R^6 is substituted or unsubstituted C_{1-8} alkyl or $\text{-CH}_2\text{R}^{19}$.



5 $P(O)R^{17}R^{18}$. Usually, R^7 is



Generally, R^8 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10} aryl.

Generally, R^9 is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10} aryl.

5 Generally, R^{10} is hydrogen or is substituted or unsubstituted C_{1-8} alkyl. Usually, R^{10} is hydrogen, ethyl or iso-propyl.

Generally, R^{11} is hydrogen, substituted or unsubstituted C_{1-8} alkyl or $-CF_3$. Usually, R^{11} is substituted or unsubstituted C_{1-8} alkyl. Preferred R^{11} is methyl.

Generally, R^{12} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, $-OR^{24}$ or $-NR^8R^9$;

10 Generally, R^{13} is $-OR^{22}$.

Generally, R^{14} is $-OR^{23}$.

Generally, R^{15} is substituted or unsubstituted C_{1-8} alkyl.

Generally, R^{16} is substituted or unsubstituted C_{1-8} alkyl, $-NR^8R^9$, $-NHS(O)_2R^{19}$ or hydroxyl. Usually, R^{16} is substituted or unsubstituted C_{1-8} alkyl or hydroxyl. Preferred R^{16} is
15 methyl or hydroxyl.

Generally, R^{17} is $-OR^{10}$ or $-NR^8R^9$. Usually, R^{17} is $-OR^{10}$. Preferred R^{17} is $-OH$, $-OEt$ or $-OiPr$.

Generally, R^{18} is OR^{10} or NR^8R^9 . Usually, R^{18} is OR^{10} . Preferred R^{18} is OH , OEt or $OiPr$.

20 Generally, R^{19} is substituted or unsubstituted heterocycle, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted C_{6-10} aryl or substituted or unsubstituted C_{3-8} cycloalkenyl.

Generally, R^{20} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or
25 unsubstituted C_{6-10} aryl.

Generally, R^{21} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C_{6-10} aryl.

30 Generally, R^{22} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, or together with R^{23} can form a cycle.

Generally, R^{23} is hydrogen, substituted or unsubstituted C_{1-8} alkyl, or together with R^{22} can form a cycle

Generally, R²⁴ is hydrogen substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl.

5 Generally, R²⁵ is substituted or unsubstituted aromatic heterocycle, or substituted or unsubstituted aromatic C₆₋₁₀ aryl.

Generally, R²⁶ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl. Usually, R²⁶ is hydrogen.

10 Generally, R²⁷ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl. Usually, R²⁷ is hydrogen.

Generally, R²⁸ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl. Usually, R²⁸ is hydrogen.

15 Generally, R²⁹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl. Usually, R²⁹ is hydrogen.

Compounds of the invention are:

Diethyl ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-4-methylpentanoyl]
20 amino}methyl)phosphonate;
(2S)-2-({[(4-Bromophenyl)amino]carbonyl}amino)-4-methyl-N-(1H-tetrazol-5-ylmethyl)pentanamide;
(2S)-2-({[(4-Bromophenyl)amino]carbonyl}amino)-4-methyl-N-[2-(1H-tetrazol-5-yl)ethyl]pentanamide;
25 (2S)-2-({[(4-Bromophenyl)amino]carbonyl}amino)-N-[(3-hydroxyisoxazol-5-yl)methyl]-4-methylpentanamide;
Diethyl ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-4-methylpentanoyl]
amino}methyl)phosphonate;
Diethyl ({[(2S,3S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-3-methylpentanoyl]
30 amino}methyl)phosphonate;
Diethyl ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino) pentanoyl]amino}methyl)phosphonate;
Diethyl ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-3-phenylpropanoyl]
amino}methyl)phosphonate;

- Diethyl (2-{{{(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-4-methylpentanoyl]amino}ethyl)phosphonate;
- Ethyl hydrogen ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-4-methylpentanoyl]amino}methyl)phosphonate;
- 5 Ethyl hydrogen ({[(2S,3S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-3-methylpentanoyl]amino}methyl)phosphonate;
- Ethyl hydrogen ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino) pentanoyl]amino}methyl)phosphonate;
- {{[(2S)-4-Methyl-2-({[4-(trifluoromethyl)phenyl]carbamoyl}amino) pentanoyl]amino}methyl)phosphonic acid;
- 10 Ethyl hydrogen ({[(2S)-4-methyl-2-({[4-(methylsulfonyl)phenyl]carbamoyl}amino) pentanoyl]amino}methyl)phosphonate;
- Diethyl ({[(2S)-4-methyl-2-({[4-(methylsulfonyl)phenyl]carbamoyl}amino) pentanoyl]amino}methyl)phosphonate;
- 15 Diethyl ({[(2S)-4-methyl-2-({[4-(methylsulfonyl)phenyl]carbamoyl}amino)pentanoyl]amino}methyl)phosphonate;
- {{(2S)-4-Methyl-2-({[4-(trifluoromethyl)phenyl]carbamoyl}amino) pentanoyl]amino}methanesulfonic acid;
- (2S)-4-Methyl-N-(1H-tetrazol-5-ylmethyl)-2-({[4-(trifluoromethyl)phenyl] carbamoyl}amino)pentanamide;
- 20 Ethyl hydrogen ({[(2S)-4-methyl-2-({[4-(trifluoromethyl)phenyl]carbamoyl}amino) pentanoyl]amino}methyl)phosphonate;
- Diethyl ({[(2S)-4-methyl-2-({[4-(trifluoromethyl)phenyl]carbamoyl}amino) pentanoyl]amino}methyl)phosphonate;
- 25 Dipropan-2-yl ({[(2S)-2-{{[(4-bromophenyl)carbamoyl]amino}-4-methylpentanoyl] amino}methyl)phosphonate;
- Propan-2-yl hydrogen {{[2-{{[(4-bromophenyl)carbamoyl]amino}pentanoyl] amino}methyl}phosphonate;
- {{(2S)-2-{{[(4-Bromophenyl)carbamoyl]amino}-4-methylpentanoyl] amino} methanesulfonic acid;
- 30 {{[(2S)-2-{{[(4-Bromophenyl)carbamoyl]amino}pentanoyl]amino} methyl}phosphonic acid;
- Ethyl hydrogen ({[(2S)-2-{{[(4-bromophenyl)carbamoyl]amino}-3-phenylpropanoyl]amino}methyl)phosphonate;

Dipropyl 2-yl (((2S)-2-((4-bromophenyl)carbamoyl)amino)pentanoyl) amino methyl) phosphonate.

Some compounds of Formula I and some of their intermediates have at least one asymmetric center in their structure. This asymmetric center may be present in an R or S configuration, said R and S notation is used in correspondence with the rules described in Pure Appl. Chem. (1976), 45, 11-13.

The term "pharmaceutically acceptable salts" refers to salts or complexes that retain the desired biological activity of the above identified compounds and exhibit minimal or no undesired toxicological effects. The "pharmaceutically acceptable salts" according to the invention include therapeutically active, non-toxic base or acid salt forms, which the compounds of Formula I are able to form.

The acid addition salt form of a compound of Formula I that occurs in its free form as a base can be obtained by treating the free base with an appropriate acid such as an inorganic acid, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; or an organic acid such as for example, acetic acid, hydroxyacetic acid, propanoic acid, lactic acid, pyruvic acid, malonic acid, fumaric acid, maleic acid, oxalic acid, tartaric acid, succinic acid, malic acid, ascorbic acid, benzoic acid, tannic acid, pantoic acid, citric acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, formic and the like (Handbook of Pharmaceutical Salts, P.Heinrich Stahl & Camille G. Wermuth (Eds), Verlag Helvetica Chimica Acta- Zürich, 2002, 329-345).

The base addition salt form of a compound of Formula I that occurs in its acid form can be obtained by treating the acid with an appropriate base such as an inorganic base, for example, sodium hydroxide, magnesium hydroxide, potassium hydroxide, calcium hydroxide, ammonia and the like; or an organic base such as for example, L-Arginine, ethanolamine, betaine, benzathine, morpholine and the like. (Handbook of Pharmaceutical Salts, P.Heinrich Stahl & Camille G. Wermuth (Eds), Verlag Helvetica Chimica Acta- Zürich, 2002, 329-345).

Compounds of Formula I and their salts can be in the form of a solvate, which is included within the scope of the present invention. Such solvates include for example hydrates, alcoholates and the like.

5 With respect to the present invention reference to a compound or compounds, is intended to encompass that compound in each of its possible isomeric forms and mixtures thereof unless the particular isomeric form is referred to specifically.

Compounds according to the present invention may exist in different polymorphic forms. Although not explicitly indicated in the above formula, such forms are intended to be included within the scope of the present invention.

10 The compounds of the invention are indicated for use in treating or preventing conditions in which there is likely to be a component involving the N-formyl peptide receptor 2.

In another embodiment, there are provided pharmaceutical compositions including at least one compound of the invention in a pharmaceutically acceptable carrier.

15 In a further embodiment of the invention, there are provided methods for treating disorders associated with modulation of the N-formyl peptide receptor 2.

Such methods can be performed, for example, by administering to a subject in need thereof a pharmaceutical composition containing a therapeutically effective amount of at least one compound of the invention.

20 Therapeutic utilities of the N-formyl peptide receptor 2 modulators are ocular inflammatory diseases including, but not limited to, wet and dry age-related macular degeneration (ARMD), uveitis, dry eye, Keratitis, allergic eye disease and conditions affecting the posterior part of the eye, such as maculopathies and retinal degeneration including non-exudative age related macular degeneration, exudative age related macular
25 degeneration, choroidal neovascularization, diabetic retinopathy (proliferative), retinopathy of prematurity (ROP), acute macular neuroretinopathy, central serous chorioretinopathy, cystoid macular edema, and diabetic macular edema; infectious keratitis, post-surgery corneal wound healing, herpetic keratitis, corneal angiogenesis, lymphangiogenesis, retinitis, and choroiditis such as acute multifocal placoid pigment epitheliopathy, Behcet's

disease, birdshot retinochoroidopathy, infectious (syphilis, lyme, tuberculosis, toxoplasmosis), intermediate uveitis (pars planitis), multifocal choroiditis, multiple evanescent white dot syndrome (mewds), ocular sarcoidosis, posterior scleritis, serpiginous choroiditis, subretinal fibrosis, Vogt-Koyanagi-and Harada syndrome; vascular diseases/ exudative diseases such as retinal arterial occlusive disease, central retinal vein occlusion, cystoids macular edema, disseminated intravascular coagulopathy, branch retinal vein occlusion, hypertensive fundus changes, ocular ischemic syndrome, retinal arterial microaneurysms, Coat's disease, parafoveal telangiectasis, hemi-retinal vein occlusion, papillophlebitis, central retinal artery occlusion, branch retinal artery occlusion, carotid artery disease (CAD), frosted branch angiitis, sickle cell retinopathy and other hemoglobinopathies, angioid streaks, familial exudative vitreoretinopathy, and Eales disease; traumatic/ surgical conditions such as sympathetic ophthalmia, uveitic retinal disease, corneal wound healing, retinal detachment, trauma, conditions caused by laser, conditions caused by photodynamic therapy, photocoagulation, hypoperfusion during surgery, radiation retinopathy, and bone marrow transplant retinopathy; proliferative disorders such as proliferative vitreal retinopathy and epiretinal membranes, and proliferative diabetic retinopathy; infectious disorders such as ocular histoplasmosis, ocular toxocariasis, presumed ocular histoplasmosis syndrome (POHS), endophthalmitis, toxoplasmosis, retinal diseases associated with HIV infection, choroidal disease associated with HIV infection, uveitic disease associated with HIV infection, viral retinitis, acute retinal necrosis, progressive outer retinal necrosis, fungal retinal diseases, ocular syphilis, ocular tuberculosis, diffuse unilateral subacute neuroretinitis, and myiasis; genetic disorders such as retinitis pigmentosa, systemic disorders associated with retinal dystrophies, congenital stationary night blindness, cone dystrophies, Stargardt's disease and fundus flavimaculatus, Best's disease, pattern dystrophy of the retinal pigmented epithelium, X-linked retinoschisis, Sorsby's fundus dystrophy, benign concentric maculopathy, Bietti's crystalline dystrophy, and pseudoxanthoma elasticum; retinal tears/ holes such as retinal detachment, macular hole, and giant retinal tear; tumors such as retinal disease associated with tumors, congenital hypertrophy of the retinal pigmented epithelium, posterior uveal melanoma, choroidal hemangioma, choroidal osteoma, choroidal metastasis, combined hamartoma of the retina and retinal pigmented epithelium, retinoblastoma, vasoproliferative tumors of the ocular fundus, retinal astrocytoma, and intraocular lymphoid tumors; and miscellaneous other diseases affecting the posterior part of the eye such as punctate inner choroidopathy, acute posterior multifocal placoid

pigment epitheliopathy, myopic retinal degeneration, and acute retinal pigment epitheliitis, systemic inflammatory diseases such as stroke, coronary artery disease, obstructive airway diseases, HIV-mediated retroviral infections, cardiovascular disorders including coronary artery disease, neuroinflammation, neurological disorders, pain and
5 immunological disorders, asthma, allergic disorders, inflammation, systemic lupus erythematosus, psoriasis, CNS disorders such as Alzheimer's disease, arthritis, sepsis, inflammatory bowel disease, cachexia, angina pectoris, post-surgical corneal inflammation, blepharitis, MGD, dermal wound healing, burns, rosacea, atopic dermatitis, acne, psoriasis, seborrheic dermatitis, actinic keratoses, viral warts, photoaging
10 rheumatoid arthritis and related inflammatory disorders, alopecia, glaucoma, branch vein occlusion, Best's vitelliform macular degeneration, retinitis pigmentosa, proliferative vitreoretinopathy (PVR), and any other degenerative disease of either the photoreceptors or the RPE (Perretti, Mauro et al. Pharmacology & Therapeutics 127 (2010) 175-188.)

These compounds are useful for the treatment of mammals, including humans, with
15 a range of conditions and diseases that are alleviated by the N-formyl peptide receptor 2 modulation: including, but not limited to the treatment of ocular inflammatory diseases: wet and dry age-related macular degeneration (ARMD), uveitis, dry eye, Keratitis, allergic eye disease and conditions affecting the posterior part of the eye, such as maculopathies and retinal degeneration including non-exudative age related macular degeneration, exudative
20 age related macular degeneration, post-surgery corneal wound healing, choroidal neovascularization, diabetic retinopathy (proliferative), retinopathy of prematurity (ROP), acute macular neuroretinopathy, central serous chorioretinopathy, cystoid macular edema, and diabetic macular edema; infectious keratitis, herpetic keratitis, corneal angiogenesis, lymphangiogenesis, retinitis, and choroiditis such as acute multifocal placoid pigment
25 epitheliopathy, Behcet's disease, birdshot retinochoroidopathy, infectious (syphilis, lyme, tuberculosis, toxoplasmosis), intermediate uveitis (pars planitis), multifocal choroiditis, multiple evanescent white dot syndrome (mewds), ocular sarcoidosis, posterior scleritis, serpiginous choroiditis, subretinal fibrosis, Vogt-Koyanagi-and Harada syndrome; vascular diseases/ exudative diseases such as retinal arterial occlusive disease, central retinal vein
30 occlusion, cystoids macular edema, corneal wound healing, disseminated intravascular coagulopathy, branch retinal vein occlusion, hypertensive fundus changes, ocular ischemic syndrome, retinal arterial microaneurysms, Coat's disease, parafoveal telangiectasis, hemi-retinal vein occlusion, papillophlebitis, central retinal artery occlusion,

branch retinal artery occlusion, carotid artery disease (CAD), frosted branch angiitis, sickle cell retinopathy and other hemoglobinopathies, angioid streaks, familial exudative vitreoretinopathy, and Eales disease; traumatic/ surgical conditions such as sympathetic ophthalmia, uveitic retinal disease, retinal detachment, trauma, conditions caused by
5 laser, conditions caused by photodynamic therapy, photocoagulation, hypoperfusion during surgery, radiation retinopathy, and bone marrow transplant retinopathy; proliferative disorders such as proliferative vitreal retinopathy and epiretinal membranes, and proliferative diabetic retinopathy; infectious disorders such as ocular histoplasmosis, ocular toxocariasis, presumed ocular histoplasmosis syndrome (POHS), endophthalmitis,
10 toxoplasmosis, retinal diseases associated with HIV infection, choroidal disease associated with HIV infection, uveitic disease associated with HIV infection, viral retinitis, acute retinal necrosis, progressive outer retinal necrosis, fungal retinal diseases, ocular syphilis, ocular tuberculosis, diffuse unilateral subacute neuroretinitis, and myiasis; genetic disorders such as retinitis pigmentosa, systemic disorders with associated retinal
15 dystrophies, congenital stationary night blindness, cone dystrophies, Stargardt's disease and fundus flavimaculatus, Best's disease, pattern dystrophy of the retinal pigmented epithelium, X-linked retinoschisis, Sorsby's fundus dystrophy, benign concentric maculopathy, Bietti's crystalline dystrophy, and pseudoxanthoma elasticum; retinal tears/ holes such as retinal detachment, macular hole, and giant retinal tear; tumors such as
20 retinal disease associated with tumors, congenital hypertrophy of the retinal pigmented epithelium, posterior uveal melanoma, choroidal hemangioma, choroidal osteoma, choroidal metastasis, combined hamartoma of the retina and retinal pigmented epithelium, retinoblastoma, vasoproliferative tumors of the ocular fundus, retinal astrocytoma, and intraocular lymphoid tumors; and miscellaneous other diseases affecting the posterior part
25 of the eye such as punctate inner choroidopathy, acute posterior multifocal placoid pigment epitheliopathy, myopic retinal degeneration, and acute retinal pigment epitheliitis, systemic inflammatory diseases such as stroke, coronary artery disease, obstructive airway diseases, HIV-mediated retroviral infections, cardiovascular disorders including coronary artery disease, neuroinflammation, neurological disorders, pain and
30 immunological disorders, asthma, allergic disorders, inflammation, systemic lupus erythematosus, psoriasis, CNS disorders such as Alzheimer's disease, arthritis, sepsis, inflammatory bowel disease, cachexia, angina pectoris, post-surgical corneal inflammation, blepharitis, MGD, dermal wound healing, burns, rosacea, atopic dermatitis, acne, psoriasis, seborrheic dermatitis, actinic keratoses, viral warts, photoaging

rheumatoid arthritis and related inflammatory disorders, alopecia, glaucoma, branch vein occlusion, Best's vitelliform macular degeneration, retinitis pigmentosa, proliferative vitreoretinopathy (PVR), and any other degenerative disease of either the photoreceptors or the RPE.

5 In still another embodiment of the invention, there are provided methods for treating disorders associated with modulation of the FPR2 receptor. Such methods can be performed, for example, by administering to a subject in need thereof a therapeutically effective amount of at least one compound of the invention, or any combination thereof, or pharmaceutically acceptable salts, hydrates, solvates, crystal forms and individual
10 isomers, enantiomers, and diastereomers thereof.

The present invention concerns the use of a compound of Formula I or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of ocular inflammatory diseases including, but not limited to, uveitis, dry eye, Keratitis, allergic eye disease and conditions affecting the posterior part of the eye, such
15 as maculopathies and retinal degeneration including non-exudative age related macular degeneration, exudative age related macular degeneration, choroidal neovascularization, diabetic retinopathy, acute macular neuroretinopathy, central serous chorioretinopathy, cystoid macular edema, and diabetic macular edema; infectious keratitis, , herpetic keratitis, corneal angiogenesis, lymphangiogenesis, retinitis, and choroiditis such as
20 acute multifocal placoid pigment epitheliopathy, Behcet's disease, birdshot retinochoroidopathy, infectious (syphilis, lyme, tuberculosis, toxoplasmosis), intermediate uveitis (pars planitis), multifocal choroiditis, multiple evanescent white dot syndrome (mewds), ocular sarcoidosis, posterior scleritis, serpiginous choroiditis, subretinal fibrosis, Vogt-Koyanagi-and Harada syndrome; vascular diseases/ exudative diseases such as
25 retinal arterial occlusive disease, central retinal vein occlusion, disseminated intravascular coagulopathy, branch retinal vein occlusion, hypertensive fundus changes, ocular ischemic syndrome, retinal arterial microaneurysms, corneal wound healing, Coat's disease, parafoveal telangiectasis, hemi-retinal vein occlusion, papillophlebitis, central retinal artery occlusion, branch retinal artery occlusion, carotid artery disease (CAD),
30 frosted branch angiitis, sickle cell retinopathy and other hemoglobinopathies, angioid streaks, familial exudative vitreoretinopathy, and Eales disease; traumatic/ surgical conditions such as sympathetic ophthalmia, uveitic retinal disease, retinal detachment, trauma, conditions caused by laser, conditions caused by photodynamic therapy,

photocoagulation, hypoperfusion during surgery, radiation retinopathy, and bone marrow transplant retinopathy; proliferative disorders such as proliferative vitreal retinopathy and epiretinal membranes, and proliferative diabetic retinopathy; infectious disorders such as ocular histoplasmosis, ocular toxocariasis, presumed ocular histoplasmosis syndrome (POHS), endophthalmitis, toxoplasmosis, retinal diseases associated with HIV infection, choroidal disease associated with HIV infection, uveitic disease associated with HIV infection, viral retinitis, acute retinal necrosis, progressive outer retinal necrosis, fungal retinal diseases, ocular syphilis, ocular tuberculosis, diffuse unilateral subacute neuroretinitis, and myiasis; genetic disorders such as retinitis pigmentosa, systemic disorders with associated retinal dystrophies, congenital stationary night blindness, cone dystrophies, Stargardt's disease and fundus flavimaculatus, Best's disease, pattern dystrophy of the retinal pigmented epithelium, X-linked retinoschisis, Sorsby's fundus dystrophy, benign concentric maculopathy, Bietti's crystalline dystrophy, and pseudoxanthoma elasticum; retinal tears/ holes such as retinal detachment, macular hole, and giant retinal tear; tumors such as retinal disease associated with tumors, congenital hypertrophy of the retinal pigmented epithelium, posterior uveal melanoma, choroidal hemangioma, choroidal osteoma, choroidal metastasis, combined hamartoma of the retina and retinal pigmented epithelium, retinoblastoma, vasoproliferative tumors of the ocular fundus, retinal astrocytoma, and intraocular lymphoid tumors; and miscellaneous other diseases affecting the posterior part of the eye such as punctate inner choroidopathy, acute posterior multifocal placoid pigment epitheliopathy, myopic retinal degeneration, and acute retinal pigment epitheliitis, systemic inflammatory diseases such as stroke, coronary artery disease, obstructive airway diseases, HIV-mediated retroviral infections, cardiovascular disorders including coronary artery disease, neuroinflammation, neurological disorders, pain and immunological disorders, asthma, allergic disorders, inflammation, systemic lupus erythematosus, psoriasis, CNS disorders such as Alzheimer's disease, arthritis, sepsis, inflammatory bowel disease, cachexia, angina pectoris, post-surgical corneal inflammation, blepharitis, MGD, dermal wound healing, burns, rosacea, atopic dermatitis, acne, psoriasis, seborrheic dermatitis, actinic keratoses, viral warts, photoaging rheumatoid arthritis and related inflammatory disorders, alopecia, glaucoma, branch vein occlusion, Best's vitelliform macular degeneration, retinitis pigmentosa, proliferative vitreoretinopathy (PVR), and any other degenerative disease of either the photoreceptors or the RPE.

The actual amount of the compound to be administered in any given case will be determined by a physician taking into account the relevant circumstances, such as the severity of the condition, the age and weight of the patient, the patient's general physical condition, the cause of the condition, and the route of administration.

5 The patient will be administered the compound orally in any acceptable form, such as a tablet, liquid, capsule, powder and the like, or other routes may be desirable or necessary, particularly if the patient suffers from nausea. Such other routes may include, without exception, transdermal, parenteral, subcutaneous, intranasal, via an implant stent, intrathecal, intravitreal, topical to the eye, direct injection, application at the back of the eye
10 or formulations that may further enhance the long duration of actions such as a slow releasing pellet, suspension, gel, or sustained delivery devices such as any suitable drug delivery system (DDS) known in the art. While topical administration is preferred, this compound may also be used in an intraocular implant as described in U.S. Patent 4,521,210 intramuscular, intravenous, and intrarectal modes of delivery. Additionally, the
15 formulations may be designed to delay release of the active compound over a given period of time, or to carefully control the amount of drug released at a given time during the course of therapy.

In another embodiment of the invention, there are provided pharmaceutical compositions including at least one compound of the invention in a pharmaceutically
20 acceptable carrier thereof. The phrase "pharmaceutically acceptable" means the carrier, diluent or excipient must be compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

Pharmaceutical compositions of the present invention can be used in the form of a solid, a solution, an emulsion, a dispersion, a patch, a micelle, a liposome, and the like,
25 wherein the resulting composition contains one or more compounds of the present invention, as an active ingredient, in admixture with an organic or inorganic carrier or excipient suitable for enteral or parenteral applications. Invention compounds may be combined, for example, with the usual non-toxic, pharmaceutically acceptable carriers for tablets, pellets, capsules, suppositories, solutions, emulsions, suspensions, and any other
30 form suitable for use. The carriers which can be used include glucose, lactose, gum acacia, gelatin, mannitol, starch paste, magnesium trisilicate, talc, corn starch, keratin, colloidal silica, potato starch, urea, medium chain length triglycerides, dextrans, and other

carriers suitable for use in manufacturing preparations, in solid, semisolid, or liquid form. In addition auxiliary, stabilizing, thickening and coloring agents and perfumes may be used. Invention compounds are included in the pharmaceutical composition in an amount sufficient to produce the desired effect upon the process or disease condition.

5 Pharmaceutical compositions containing invention compounds may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known in the art for the manufacture of pharmaceutical compositions and such
10 compositions may contain one or more agents selected from the group consisting of a sweetening agent such as sucrose, lactose, or saccharin, flavoring agents such as peppermint, oil of wintergreen or cherry, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets containing invention compounds in admixture with non-toxic pharmaceutically acceptable excipients may also
15 be manufactured by known methods. The excipients used may be, for example, (1) inert diluents such as calcium carbonate, lactose, calcium phosphate or sodium phosphate; (2) granulating and disintegrating agents such as corn starch, potato starch or alginic acid; (3) binding agents such as gum tragacanth, corn starch, gelatin or acacia, and (4) lubricating agents such as magnesium stearate, stearic acid or talc. The tablets may be uncoated or
20 they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

 In some cases, formulations for oral use may be in the form of hard gelatin
25 capsules wherein the invention compounds are mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin. They may also be in the form of soft gelatin capsules wherein the invention compounds are mixed with water or an oil medium, for example, peanut oil, liquid paraffin or olive oil.

 The pharmaceutical compositions may be in the form of a sterile injectable
30 suspension. This suspension may be formulated according to known methods using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic

parenterally-acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides, fatty acids (including oleic acid), naturally occurring vegetable oils like
5 sesame oil, coconut oil, peanut oil, cottonseed oil, etc., or synthetic fatty vehicles like ethyl oleate or the like. Buffers, preservatives, antioxidants, and the like can be incorporated as required.

Pharmaceutical compositions containing invention compounds may be in a form
10 suitable for topical use, for example, as oily suspensions, as solutions or suspensions in aqueous liquids or nonaqueous liquids, or as oil-in-water or water-in-oil liquid emulsions.

Pharmaceutical compositions may be prepared by combining a therapeutically effective amount of at least one compound according to the present invention, or a
15 pharmaceutically acceptable salt thereof, as an active ingredient with conventional ophthalmically acceptable pharmaceutical excipients and by preparation of unit dosage suitable for topical ocular use. The therapeutically efficient amount typically is between about 0.0001 and about 5% (w/v), preferably about 0.001 to about 2.0% (w/v) in liquid formulations.

20 For ophthalmic application, preferably solutions are prepared using a physiological saline solution as a major vehicle. The pH of such ophthalmic solutions should preferably be maintained between 4.5 and 8.0 with an appropriate buffer system, a neutral pH being preferred but not essential. The formulations may also contain conventional
25 pharmaceutically acceptable preservatives, stabilizers and surfactants.

Preferred preservatives that may be used in the pharmaceutical compositions of the present invention include, but are not limited to, benzalkonium chloride, chlorobutanol, thimerosal, phenylmercuric acetate and phenylmercuric nitrate.
A preferred surfactant is, for example, Tween 80. Likewise, various preferred vehicles may
30 be used in the ophthalmic preparations of the present invention. These vehicles include, but are not limited to, polyvinyl alcohol, povidone, hydroxypropyl methyl cellulose, poloxamers, carboxymethyl cellulose, hydroxyethyl cellulose cyclodextrin and purified water.

Tonicity adjustors may be added as needed or convenient. They include, but are not limited to, salts, particularly sodium chloride, potassium chloride, mannitol and glycerin, or any other suitable ophthalmically acceptable tonicity adjustor.

Various buffers and means for adjusting pH may be used so long as the resulting preparation is ophthalmically acceptable. Accordingly, buffers include acetate buffers, citrate buffers, phosphate buffers and borate buffers. Acids or bases may be used to adjust the pH of these formulations as needed.

In a similar manner an ophthalmically acceptable antioxidant for use in the present invention includes, but is not limited to, sodium metabisulfite, sodium thiosulfate, acetylcysteine, butylated hydroxyanisole and butylated hydroxytoluene.

Other excipient components which may be included in the ophthalmic preparations are chelating agents. The preferred chelating agent is edentate disodium, although other chelating agents may also be used in place of or in conjunction with it.

The ingredients are usually used in the following amounts:

Ingredient	Amount (% w/v)
active ingredient	about 0.001-5
preservative	0-0.10
vehicle	0-40
tonicity adjustor	0-10
buffer	0.01-10
pH adjustor	q .s. pH 4.5-7.8
antioxidant	as needed
surfactant	as needed
purified water	to make 100%

The actual dose of the active compounds of the present invention depends on the specific compound, and on the condition to be treated; the selection of the appropriate dose is well within the knowledge of the skilled artisan.

The ophthalmic formulations of the present invention are conveniently packaged in forms suitable for metered application, such as in containers equipped with a dropper, to facilitate application to the eye. Containers suitable for dropwise application are usually made of suitable inert, non-toxic plastic material, and generally contain between about 0.5

and about 15 ml solution. One package may contain one or more unit doses. Especially preservative-free solutions are often formulated in non-resealable containers containing up to about ten, preferably up to about five units doses, where a typical unit dose is from one to about 8 drops, preferably one to about 3 drops. The volume of one drop usually is about 20-35 μ L.


The compounds of the invention may also be administered in the form of suppositories for rectal administration of the drug. These compositions may be prepared by mixing the invention compounds with a suitable non-irritating excipient, such as cocoa butter, synthetic glyceride esters of polyethylene glycols, which are solid at ordinary temperatures, but liquefy and/or dissolve in the rectal cavity to release the drug.

Since individual subjects may present a wide variation in severity of symptoms and each drug has its unique therapeutic characteristics, the precise mode of administration and dosage employed for each subject is left to the discretion of the practitioner.

The compounds and pharmaceutical compositions described herein are useful as medicaments in mammals, including humans, for treatment of diseases and/or alleviations of conditions which are responsive to treatment by agonists or functional antagonists of the N-formyl peptide receptor 2. Thus, in further embodiments of the invention, there are provided methods for treating a disorder associated with modulation of the N-formyl peptide receptor 2. Such methods can be performed, for example, by administering to a subject in need thereof a pharmaceutical composition containing a therapeutically effective amount of at least one invention compound. As used herein, the term "therapeutically effective amount" means the amount of the pharmaceutical composition that will elicit the biological or medical response of a subject in need thereof that is being sought by the researcher, veterinarian, medical doctor or other clinician. In some embodiments, the subject in need thereof is a mammal. In some embodiments, the mammal is human.

The present invention concerns also processes for preparing the compounds of Formula I. The compounds of Formula I according to the invention can be prepared analogously to conventional methods as understood by the person skilled in the art of synthetic organic chemistry. Synthetic Scheme 1 set forth below, illustrates how the compounds according to the invention can be made.

Scheme 1



Compounds within the scope of the invention may be prepared as depicted in Scheme 1. In general, a t-butyl ester of an amino acid can be treated with an aryl isocyanide in an inert solvent to give the corresponding amino acid urea. The t-butyl ester can be hydrolyzed with a strong acid (e.g., trifluoroacetic acid) to provide the free amino acid urea. The carboxylic acid is then treated with an activating agent (e.g., EDCI) in the presence of a substituted alkylamine and an organic base (e.g., triethylamine) to provide compounds of Formula I. At this stage, those skilled in the art will appreciate that many additional compounds that fall under the scope of the invention may be prepared by performing various common chemical reactions. Details of certain specific chemical transformations are provided in the examples.

DETAILED DESCRIPTION OF THE INVENTION

It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the

invention claimed. As used herein, the use of the singular includes the plural unless specifically stated otherwise.

It will be readily apparent to those skilled in the art that some of the compounds of the invention may contain one or more asymmetric centers, such that the compounds may exist in enantiomeric as well as in diastereomeric forms. Unless it is specifically noted otherwise, the scope of the present invention includes all enantiomers, diastereomers and racemic mixtures. Some of the compounds of the invention may form salts with pharmaceutically acceptable acids or bases, and such pharmaceutically acceptable salts of the compounds described herein are also within the scope of the invention.

The present invention includes all pharmaceutically acceptable isotopically enriched compounds. Any compound of the invention may contain one or more isotopic atoms enriched or different than the natural ratio such as deuterium ^2H (or D) in place of protium ^1H (or H) or use of ^{13}C enriched material in place of ^{12}C and the like. Similar substitutions can be employed for N, O and S. The use of isotopes may assist in analytical as well as therapeutic aspects of the invention. For example, use of deuterium may increase the in vivo half-life by altering the metabolism (rate) of the compounds of the invention. These compounds can be prepared in accord with the preparations described by use of isotopically enriched reagents.

The following examples are for illustrative purposes only and are not intended, nor should they be construed as limiting the invention in any manner. Those skilled in the art will appreciate that variations and modifications of the following examples can be made without exceeding the spirit or scope of the invention.

As will be evident to those skilled in the art, individual isomeric forms can be obtained by separation of mixtures thereof in conventional manner. For example, in the case of diastereoisomeric isomers, chromatographic separation may be employed.

Compound names were generated with ACD version 12.0 and Intermediates and reagent names used in the examples were generated with softwares such as Chem Bio Draw Ultra version 12.0, ACD version 12.0 or Auto Nom 2000 from MDL ISIS Draw 2.5 SP1.

In general, characterization of the compounds is performed according to the following methods:

NMR spectra are recorded on 300 and/or 600 MHz Varian and acquired at room temperature. Chemical shifts are given in ppm referenced either to internal TMS or to the solvent signal. The optical rotation was recorded on Perkin Elmer Polarimeter 341, 589 nm at 20 °C, Na/Hal lamp.

All the reagents, solvents, catalysts for which the synthesis is not described are purchased from chemical vendors such as Sigma Aldrich, Fluka, Bio-Blocks, Combi-blocks, TCI, VWR, Lancaster, Oakwood, Trans World Chemical, Alfa, Fisher, Maybridge, Frontier, Matrix, Ukrorgsynth, Toronto, Ryan Scientific, SiliCycle, Anaspec, Syn Chem, Chem-Impex, MIC-scientific, Ltd; however some known intermediates, were prepared according to published procedures.

Usually the compounds of the invention were purified by column chromatography (Auto-column) on an Teledyne-ISCO CombiFlash with a silica column, unless noted otherwise.

Those skilled in the art will be routinely able to modify and/or adapt the following examples to synthesize any compound of the invention covered by **Formula I**.

The following abbreviations are used in the examples:


EtOAc	ethyl acetate
Et ₃ N	triethylamine
CH ₂ Cl ₂	dichloromethane
CDCl ₃	deuterated chloroform
MeOH	methanol
CD ₃ OD	deuterated methanol
Na ₂ SO ₄	sodium sulfate
DMF	N,N dimethylformamide
EDCI	1-ethyl-3-(3-dimethylaminopropyl) carbodiimide
HOBt	Hydroxybenzotriazole
THF	tetrahydrofuran
EtOH	ethanol
DIEPA	diisopropylethylamine

HATU 2-(7-Aza-1H-Benzotriazole -1-yl)-1,1,3,3-tetramethyluronium
hexafluorophosphate
HCl hydrochloric acid

5 Example 1

Intermediate 1

tert-Butyl (2S)-2-([(4-Bromophenyl)amino]carbonyl)amino)-4-methylpentanoate




10 To a solution of L-leucine *tert*-butyl ester hydrochloride (1.75 g, 9.35 mmol) and 20 mL of methylene chloride at 25 °C was added 4-bromo-phenyl isocyanate (1.85 g, 9.35 mmol) and triethylamine (1.95 mL, 14.0 mmol). The resulting mixture was stirred at 25 °C for 30 minutes. The mixture was concentrated and the residue was purified by medium pressure liquid chromatography on silica gel using ethyl acetate : hexane (20:80) to yield





15 **Intermediate 1**, as a white solid.

¹H NMR (CDCl₃, 300MHz) δ: 7.20 - 7.33 (m, 2H), 7.04 - 7.15 (m, 2H), 4.44 (dd, J = 9.1, 5.3 Hz, 1H), 1.74 (dd, J = 12.9, 6.4 Hz, 1H), 1.54 - 1.68 (m, 1H), 1.50 (s, 9H), 1.40 - 1.47 (m, 1H), 0.97 (d, J = 3.5 Hz, 3H), 0.95 (d, 3H).

Intermediates 2 through 6 were prepared from the corresponding amino acids in a similar manner to the procedure described in **Example 1** for **Intermediate 1**. The results are described below in **Table 1**.

Table 1

Interm. No.	IUPAC name	Starting material	¹ H NMR δ (ppm)
2	tert-Butyl (2S,3S)-2-([(4-Bromophenyl)amino]carbonyl)amino)-3-methylpentanoate 	N-(aminocarbonyl)-L-isoleucine <i>tert</i> -butyl ester hydrochloride	¹ H NMR (CDCl ₃ , 300MHz) δ: 7.29 - 7.39 (m, 2H), 7.10 - 7.22 (m, 2H), 6.83 (br. s., 1H), 4.44 (d, J = 4.4 Hz, 1H), 1.81 - 1.99 (m, 1H), 1.36 - 1.46 (m, 1H), 1.08 - 1.31 (m, 1H), 0.86 - 1.02 (m, 6H).


3	<i>tert</i>-Butyl (2<i>S</i>)-2-({[(4-Bromophenyl)amino]carbonyl}amino)pentanoate 	N-(aminocarbonyl)-L-norvaline <i>tert</i> -butyl ester hydrochloride	¹ H NMR (CDCl ₃ , 300MHz) δ: 7.26 - 7.36 (m, 2H), 7.09 - 7.18 (m, 2H), 6.95 (br. s., NH), 4.40 - 4.50 (m, 1H), 1.73 - 1.89 (m, 1H), 1.52 - 1.72 (m, 1H), 1.25 - 1.46 (m, 2H), 0.95 (t, 2H).
4	<i>tert</i>-Butyl (2<i>S</i>)-2-({[(4-Bromophenyl)amino]carbonyl}amino)-3-phenylpropanoate 	N-(aminocarbonyl)-L-phenylalanine <i>tert</i> -butyl ester hydrochloride	¹ H NMR (CDCl ₃ , 300MHz) δ: 7.20 - 7.35 (m, 5H), 7.13 - 7.20 (m, 2H), 7.01 - 7.10 (m, 2H), 6.79 (br. s., NH), 5.52 (br. s., NH), 4.70 (t, J = 6.2 Hz, 1H), 2.91 (ddd, J = 19.0 Hz, J = 6.0 Hz, 2H), 1.47 (m, 9H).
5	<i>tert</i>-butyl (2<i>S</i>)-4-methyl-2-({[(4-(trifluoromethyl)phenyl)amino]carbonyl}amino)pentanoate 	L-leucine <i>tert</i> -butyl ester hydrochloride 1-isocyanato-4-(trifluoromethyl)-benzene	¹ H NMR (CD ₃ OD, 300MHz) δ: 7.50 (s, 4H), 4.27 (dd, J = 9.1, 5.6 Hz, 1H), 1.68 - 1.86 (m, 1H), 1.52 - 1.66 (m, 2H), 1.45 - 1.50 (s, 9H), 0.95 (t, J = 6.9 Hz, 6H).
6	<i>tert</i>-butyl (2<i>S</i>)-4-methyl-2-({[(4-(methylthio)phenyl)amino]carbonyl}amino)pentanoate 	L-leucine <i>tert</i> -butyl ester hydrochloride 1-isocyanato-4-(methylthio)-benzene	¹ H NMR (CD ₃ OD, 300MHz) δ: 7.27 - 7.32 (m, 2H), 7.18 - 7.23 (m, 2H), 4.22 - 4.29 (m, 1H), 2.42 (s, 3H), 1.70 - 1.79 (m, 1H), 1.51 - 1.61 (m, 2H), 1.47 (s, 9H), 0.97 (t, J = 6.7 Hz, 6H).

Example 2

Intermediate 7

5

(2*S*)-2-({[(4-Bromophenyl)amino]carbonyl}amino)-4-methylpentanoic acid








A solution of **Intermediate 1** (2.77 g, 7.20 mmol) and 20 mL of formic acid was stirred at 25 °C for 3 hours. The resulting mixture was quenched with water (1mL) then extracted with ethyl acetate. The organic layer was washed with water, brine, dried over Na₂SO₄, filtered, and the filtrate was concentrated under reduced pressure. The residue was rinsed 4 times with methylene chloride : hexane (1:1) to yield **Intermediate 7** as white solid.

¹H NMR (acetone-d₆, 300MHz) δ: 8.17 (s, NH), 7.43 - 7.51 (m, 2H), 7.35 - 7.41 (m, 2H), 6.04 (d, J = 9.1 Hz, NH), 4.42 - 4.53 (m, 1H), 1.73 - 1.88 (m, 1H), 1.53 - 1.73 (m, 2H), 0.97 (d, J = 2.1 Hz, 3H), 0.95 (d, 3H).

Intermediates 8 through **12** were prepared from the corresponding urea derivative in a similar manner to the procedure described in **Example 2** for **Intermediate 7**. The results are described below in **Table 2**.

Table 2

Interm. No.	IUPAC name	Starting Material	¹ H NMR δ (ppm)
8	(2S,3S)-2-({[(4-Bromophenyl)amino]carbonyl}amino)-3-methylpentanoic acid 	Intermediate 2	¹ H NMR (acetone-d ₆ , 300MHz) δ: 8.24 (br. s., 1H), 7.44 - 7.53 (m, 2H), 7.32 - 7.42 (m, 2H), 6.08 (d, J = 8.8 Hz, 1H), 4.44 (dd, J = 8.6, 4.8 Hz, 1H), 1.86 - 2.00 (m, J = 9.1, 6.9, 4.6, 4.6 Hz, 1H), 1.43 - 1.61 (m, 1H), 1.15 - 1.33 (m, 1H), 0.88 - 1.04 (m, 6H).
9	(2S)-2-({[(4-Bromophenyl)amino]carbonyl}amino)pentanoic acid 	Intermediate 3	¹ H NMR (acetone-d ₆ , 300MHz) δ: 8.20 (s, NH), 7.43 - 7.52 (m, 2H), 7.33 - 7.41 (m, 2H), 6.08 (d, J = 9.1 Hz, NH), 4.38 - 4.50 (m, 1H), 1.77 - 1.92 (m, 1H), 1.61 - 1.76 (m, 1H), 1.36 - 1.53 (m,


			2H), 0.89 - 1.00 (m, 3H).
10	(2S)-2-([(4-Bromophenyl)amino]carbonyl)amino-3-phenylpropanoic Acid 	Intermediate 4	¹ H NMR (acetone-d ₆ , 300MHz) δ: 8.29 (s, NH), 7.40 - 7.50 (m, 2H), 7.32 - 7.40 (m, 2H), 7.18 - 7.31 (m, 5H), 5.98 (d, J = 7.9 Hz, NH), 4.67 (m, 1H), 3.02 (ddd, J = 19.0 Hz, J = 6.0 Hz, 2H).
11	(2S)-4-methyl-2-([(4-(trifluoromethyl)phenyl)amino]carbonyl)amino] pentanoic acid 	Intermediate 5	¹ H NMR (CD ₃ OD, 300MHz) δ: 7.49 - 7.57 (m, 4H), 4.38 (dd, J = 9.4, 5.0 Hz, 1H), 1.69 - 1.87 (m, 1H), 1.51 - 1.69 (m, 2H), 0.92 - 1.01 (m, 6H).
12	(2S)-4-methyl-2-([(4-(methylthio)phenyl)amino]carbonyl)amino]pentanoic acid 	Intermediate 6	¹ H NMR (CD ₃ OD, 300MHz) δ: 7.25 - 7.31 (m, 2H), 7.14 - 7.20 (m, 2H), 4.37 (dd, J = 9.2, 5.1 Hz, 1H), 2.39 (s, 3H), 1.68 - 1.83 (m, 1H), 1.51 - 1.67 (m, 2H), 0.96 (dd, J = 6.2, 2.3 Hz, 6H).

Example 3

5

Compound 1

Diethyl ([[(2S)-2-([(4-bromophenyl)amino]carbonyl)amino]-4-methylpentanoyl]amino)methyl)phosphonate



10



To a solution of diethyl (aminomethyl)phosphonate oxalate (82 mg, 0.32 mmol) and 8 mL of anhydrous dichloromethane at 0 °C was added N-methylmorpholine (130 mg, 1.28 mmol), **Intermediate 7** (100 mg, 0.32 mmol), EDCI (92 mg, 0.48 mmol), and HOBT (65




mg, 0.48 mmol). The resulting mixture was stirred at 25 °C for 4 hours. The mixture was quenched with water (2 mL), and the product was extracted with ethyl acetate (20 mL). The layers were separated, and the organic layer was washed with water, brine, dried over Na₂SO₄, filtered, and the filtrate was concentrated under reduced pressure. The resulting product was purified by medium pressure liquid chromatography on silica gel using 100% ethyl acetate to yield **Compound 1** as white solid.




¹H NMR (CD₃OD, 300MHz) δ: 7.34 - 7.41 (m, 2H), 7.27 - 7.32 (m, 2H), 4.33 (dd, J = 9.1, 5.9 Hz, 1H), 4.06 - 4.20 (m, 4H), 3.56 - 3.86 (m, 2H), 1.67 - 1.82 (m, 1H), 1.46 - 1.66 (m, 2H), 1.30 (td, J = 7.1, 2.2 Hz, 6H), 0.97 (dd, J = 6.6, 4.8 Hz, 7H).





Compounds 2 through **14** were prepared from the corresponding urea derivative in a similar manner to the procedure described in **Example 3** for **Compound 1**. The results are described below in **Table 3**.


Table 3

Comp. No.	IUPAC name	Starting material	¹ H NMR δ (ppm)	Properties
2	(2S)-2-({[(4-Bromophenyl)amino]carbonyl}amino)-4-methyl-N-(1H-tetrazol-5-ylmethyl)pentanamide 	Intermediate 7 2H-Tetrazole-5-methanamine	¹ H NMR (CD ₃ OD, 300MHz) δ: 7.32 - 7.39 (m, 2H), 7.25 - 7.32 (m, 2H), 4.57 - 4.76 (m, 2H), 4.33 (dd, J = 9.5, 5.1 Hz, 1H), 1.65 - 1.81 (m, 1H), 1.50 - 1.65 (m, 2H), 0.96 (d, J = 3.2 Hz, 3H), 0.94 (d, 3H)	Yellow solid
3	(2S)-2-({[(4-Bromophenyl)amino]carbonyl}amino)-4-methyl-N-[2-(1H-tetrazol-5-yl)ethyl]pentanamide 	Intermediate 7 2H-Tetrazole-5-ethanamine	¹ H NMR (CD ₃ OD, 300MHz) δ: 7.35 - 7.39 (m, 2H), 7.27 - 7.32 (m, 2H), 4.17 (dd, J = 8.1, 6.9 Hz, 1H), 3.52 - 3.69 (m, 2H), 3.12 - 3.20 (m, 2H), 1.66 (dt, J = 13.3, 6.7 Hz, 1H), 1.45 - 1.54 (m, 1H), 1.33 - 1.40 (m, 1H), 0.90 - 0.99 (m, 6H).	Yellow solid
4	(2S)-2-({[(4-	Intermediate	¹ H NMR (CD ₃ OD,	Off-white

	Bromophenyl]amino]carbonyl]amino)-N-[(3-hydroxyisoxazol-5-yl)methyl]-4-methylpentanamide 	7 5-(aminomethyl)-3(2H)-isoxazolone	600MHz) δ : 7.35 - 7.38 (m, 2H), 7.28 - 7.31 (m, 2H), 5.83 (s, 1H), 4.33 - 4.44 (m, 2H), 4.26 - 4.34 (m, 1H), 1.69 - 1.79 (m, 1H), 1.52 - 1.64 (m, 2H), 0.93 - 1.00 (m, 6H).	solid
5	Diethyl ({[(2S,3S)-2-({[(4-bromophenyl)amino]carbonyl]amino)-3-methylpentanoyl]amino}methyl]phosphonate 	Intermediate 8 diethyl (aminomethyl) phosphonate oxalate	¹ H NMR (CD ₃ OD, 300MHz) δ : 7.34 - 7.39 (m, 2H), 7.27 - 7.32 (m, 2H), 6.27 (d, J = 8.5 Hz, NH), 4.05 - 4.25 (m, 5H), 3.75 - 3.89 (m, 1H), 3.56 - 3.70 (m, 1H), 1.76 - 1.92 (m, 1H), 1.49 - 1.67 (m, 1H), 1.31 (td, J = 7.0, 1.8 Hz, 6H), 0.89 - 1.03 (m, 6H).	White solid
6	Diethyl ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl]amino)pentanoyl]amino}methyl]phosphonate 	Intermediate 9 diethyl (aminomethyl) phosphonate oxalate	¹ H NMR (CD ₃ OD, 600MHz) δ : 7.35 - 7.38 (m, 2H), 7.28 - 7.31 (m, 2H), 4.28 (dd, J = 8.1, 5.7 Hz, 1H), 4.09 - 4.18 (m, 4H), 3.80 (dd, J = 15.8, 12.0 Hz, 2H), 3.64 (dd, J = 15.8, 11.4 Hz, 2H), 1.75 (ddt, J = 13.6, 9.9, 6.1 Hz, 1H), 1.58 - 1.67 (m, 1H), 1.37 - 1.50 (m, 2H), 1.31 (td, J = 7.0, 3.5 Hz, 6H), 0.96 (t, J = 7.3 Hz, 3H).	


7	Diethyl ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-3-phenylpropanoyl]amino}methyl)phosphonate 	Intermediate 10 diethyl (aminomethyl) phosphonate oxalate	¹ H NMR (CD ₃ OD, 600MHz) δ: 7.32 - 7.35 (m, 2H), 7.19 - 7.31 (m, 7H), 4.57 (dd, J = 7.9, 6.2 Hz, 1H), 4.05 - 4.18 (m, 4H), 3.63 - 3.77 (m, 2H), 3.13 (dd, J = 13.8, 5.9 Hz, 1H), 2.92 (dd, J = 13.9, 8.1 Hz, 1H), 1.24 - 1.35 (m, 6H).	White solid
8	Diethyl (2-({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-4-methylpentanoyl]amino}ethyl)phosphonate 	Intermediate 7 diethyl ester <i>P</i> -(2-aminoethyl) phosphonic acid	¹ H NMR (CD ₃ OD, 300MHz) δ: 7.35 - 7.39 (m, 2H), 7.27 - 7.32 (m, 2H), 4.24 (dd, J = 9.2, 5.4 Hz, 1H), 4.02 - 4.17 (m, 4H), 3.43 (dt, J = 12.2, 7.5 Hz, 2H), 1.99 - 2.14 (m, 2H), 1.66 - 1.79 (m, 1H), 1.49 - 1.63 (m, 2H), 1.25 - 1.36 (m, 6H), 0.92 - 1.01 (m, 6H).	
9	Diisopropyl ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)pentanoyl]amino}methyl)phosphonate 	Intermediate 9 bis(1-methylethyl) ester <i>P</i> -(2-aminomethyl) phosphonic acid	¹ H NMR (CD ₃ OD, 300MHz) δ : 7.33 - 7.39 (m, 2H), 7.27 - 7.32 (m, 2H), 4.61 - 4.75 (m, 2H), 4.29 (dd, J = 7.9, 5.9 Hz, 1H), 3.69 - 3.83 (m, 1H), 3.50 - 3.64 (m, 1H), 1.54 - 1.82 (m, 2H), 1.39 - 1.52 (m, 2H), 1.27 - 1.36 (m, 12H), 0.96 (d, J = 14.7 Hz, 3H).	White solid

10	Diisopropyl ({[2-({[4-bromophenyl]amino}carbonyl)amino]-4-methylpentanoyl]amino}methyl)phosphonate 	Intermediate 7 bis(1-methylethyl) ester <i>P</i> -(2-aminomethyl) phosphonic acid	¹ H NMR (CD ₃ OD, 300MHz) δ: 7.26 - 7.40 (m, 2H), 4.60 - 4.75 (m, 2H), 4.30 - 4.39 (m, 1H), 3.69 - 3.83 (m, 1H), 3.51 - 3.63 (m, 1H), 1.68 - 1.82 (m, 1H), 1.46 - 1.66 (m, 2H), 1.32 (t, J = 3.1 Hz, 9H), 0.94 - 1.01 (m, 6H).	White solid
11	Diethyl [({(2S)-4-methyl-2-({[4-(trifluoromethyl)phenyl]amino}carbonyl)amino}pentanoyl]amino}methyl]phosphonate 	Intermediate 11 diethyl (aminomethyl) phosphonate oxalate	¹ H NMR (CD ₃ OD, 600MHz) δ: 7.45 - 7.62 (m, 4H), 6.35 (d, J = 8.2 Hz, NH), 4.28 - 4.43 (m, 1H), 4.02 - 4.22 (m, 4H), 3.55 - 3.89 (m, 2H), 1.68 - 1.80 (m, 1H), 1.49 - 1.66 (m, 2H), 1.27 - 1.35 (m, 6H), 0.98 (dd, J = 6.6, 4.5 Hz, 6H).	White solid
12	(2S)-4-methyl-N-(1H-tetrazol-5-ylmethyl)-2-({[4-(trifluoromethyl)phenyl]amino}carbonyl)amino}pentanamide 	Intermediate 11 2 <i>H</i> -Tetrazole-5-methanamine	¹ H NMR (CD ₃ OD, 300MHz) δ: 7.54 (s, 4H), 4.60 - 4.82 (m, 2H), 4.28 - 4.40 (m, 1H), 1.71 (dd, J = 13.3, 6.0 Hz, 1H), 1.52 - 1.67 (m, 2H), 0.99 (d, J = 3.5 Hz, 3H), 0.97 (d, J = 3.2 Hz, 3H).	Yellow solid
13	(({(2S)-4-methyl-2-({[4-(trifluoromethyl)phenyl]amino}carbonyl)amino}pentanoyl]amino)methanesulfonic acid 	Intermediate 11 1-amino-methanesulfonic acid	¹ H NMR (CD ₃ OD, 300MHz) δ: 7.45 - 7.61 (m, 4H), 6.40 (d, J = 7.9 Hz, NH), 4.22 - 4.47 (m, 3H), 1.52 - 1.84 (m, 3H), 0.99 (d, J = 2.1 Hz, 3H), 0.97 (d, J = 1.8 Hz, 3H).	Off white solid

14	Diethyl [({(2S)-4-methyl-2-[[[4-(methylthio)phenyl]amino]carbonyl}amino]pentanoyl}amino)methyl]phosphonate 	Intermediate 12 diethyl (aminomethyl) phosphonate oxalate	¹ H NMR (CD ₃ OD, 300MHz) δ: 7.30 - 7.33 (m, 2H), 7.27 - 7.30 (m, 2H), 4.33 (dd, J = 9.1, 5.9 Hz, 1H), 4.05 - 4.20 (m, 4H), 3.57 - 3.87 (m, 2H), 1.67 - 1.84 (m, 1H), 1.45 - 1.65 (m, 2H), 1.31 (td, J = 7.1, 2.2 Hz, 6H), 0.97 (dd, J = 6.6, 4.8 Hz, 6H).	Off white solid
----	--	---	--	-----------------

Example 4**Compound 15**

5 **Ethyl hydrogen ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-4-methylpentanoyl]amino)methyl]phosphonate**



10 A solution of **Compound 1** (350 mg, 0.73 mmol), 8 mL EtOH, 2 mL of THF and 2M LiOH (1.5 mL, 2.93 mmol) was stirred at 25 °C for 5 hours. The resulting mixture was concentrated under vacuum, quenched with 10% HCl (3mL) then extracted with ethyl acetate. The organic layer was washed with water, brine, dried over Na₂SO₄, filtered, and

15 the filtrate was concentrated under reduced pressure. The residue was rinsed 4 times with acetone : hexane (25:75) to yield **Compound 15** as white solid.




¹H NMR (CD₃OD, 300MHz) δ: 7.33 - 7.41 (m, 2H), 7.27 - 7.33 (m, 2H), 4.32 (dd, J = 10.1, 4.5 Hz, 1H), 3.89 (quin, J = 7.0 Hz, 2H), 3.35 - 3.56 (m, 2H), 1.67 - 1.83 (m, 1H), 1.48 - 1.68 (m, 2H), 1.14 - 1.23 (m, 3H), 0.98 (d, J = 4.4 Hz, 3H), 0.96 (d, 3H). [α]_D = -18.1 °




20 (c=1.00, MeOH).

Compounds 16 through **21** were prepared from the corresponding urea derivative in a similar manner to the procedure described in **Example 4** for **Compound 15**. The results

25 are described below in **Table 4**.

Table 4

Comp. No.	IUPAC name	Starting material	¹ H NMR δ (ppm)	Properties
16	Ethyl hydrogen ({[(2S,3S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-3-methylpentanoyl]amino}methyl)phosphonate 	Compound 5	¹ H NMR (CD ₃ OD, 300MHz) δ : 7.34 - 7.39 (m, 2H), 7.27 - 7.32 (m, 2H), 4.18 (d, J = 6.4 Hz, 1H), 4.08 (quin, J = 7.2 Hz, 2H), 3.51 - 3.80 (m, 2H), 1.77 - 1.92 (m, 1H), 1.51 - 1.66 (m, 1H), 1.25 - 1.31 (m, 3H), 1.24 - 1.33 (m, 3H), 1.10 - 1.24 (m, 1H), 0.90 - 1.01 (m, 6H).	white solid
17	Ethyl hydrogen ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)pentanoyl]amino}methyl)phosphonate 	Compound 6	¹ H NMR (CD ₃ OD, 600MHz) δ : 7.26 - 7.42 (m, 4H), 4.28 (dd, J = 8.2, 5.3 Hz, 1H), 3.89 (quin, J = 7.0 Hz, 2H), 3.33 - 3.57 (m, 2H), 1.80 (m, 1H), 1.64 (m, 1H), 1.35 - 1.53 (m, 2H), 1.19 (t, J = 7.0 Hz, 3H), 0.96 (t, J = 7.3 Hz, 3H).	Yellow solid
18	Ethyl hydrogen ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-3-phenylpropanoyl]amino}methyl)phosphonate 	Compound 7	¹ H NMR (CD ₃ OD, 300MHz) δ : 7.31 - 7.36 (m, 3H), 7.19 - 7.30 (m, 6H), 4.57 (dd, J = 8.2, 5.3 Hz, 1H), 4.03 (quin, J = 7.2 Hz, 2H), 3.53 - 3.62 (m, 1H), 3.16 (dd, J = 13.9, 5.4 Hz, 2H), 2.92 (dd, J = 13.9, 8.4 Hz, 1H), 1.23 - 1.29 (m, 3H).	White solid


19	/iso-propyl hydrogen ({[2-({[4-bromophenyl]amino]carbonyl]amino)pentanoyl]amino)methyl]phosphonate 	Compound 9	¹ H NMR (CD ₃ OD, 300MHz) δ: 7.28 - 7.38 (m, 4H), 4.35 - 4.51 (m, 1H), 4.21 - 4.33 (m, 1H), 3.34 - 3.56 (m, 2H), 1.73 - 1.90 (m, 1H), 1.55 - 1.72 (m, 1H), 1.34 - 1.54 (m, 2H), 1.21 (s, 3H), 1.19 (s, 3H), 0.96 (t, J = 7.3 Hz, 3H).	Light yellow solid
20	Ethyl hydrogen [{(2S)-4-methyl-2-({[4-(trifluoromethyl)phenyl]amino]carbonyl]amino]pentanoyl]amino)methyl]phosphonate 	Compound 11	¹ H NMR (CD ₃ OD, 300MHz) δ: 7.50 - 7.58 (m, 4H), 4.36 (dd, J = 9.2, 5.7 Hz, 1H), 4.08 (quin, J = 7.2 Hz, 2H), 3.50 - 3.81 (m, 2H), 1.70 - 1.87 (m, 1H), 1.49 - 1.68 (m, 2H), 1.28 (t, J = 7.0 Hz, 3H), 0.98 (dd, J = 6.4, 3.8 Hz, 6H).	White solid
21	ethyl hydrogen [{(2S)-4-methyl-2-({[4-(methylsulfonyl)phenyl]amino]carbonyl]amino]pentanoyl]amino)methyl]phosphonate 	Compound 24	¹ H NMR (CD ₃ OD, 300MHz) δ: 7.76 - 7.86 (m, 2H), 7.57 - 7.68 (m, 2H), 4.36 (dd, J = 9.8, 4.8 Hz, 1H), 3.96 (quin, J = 7.0 Hz, 2H), 3.52 (d, J = 12.3 Hz, 2H), 3.07 (s, 3H), 1.49 - 1.85 (m, 3H), 1.18 - 1.27 (m, 3H), 0.93 - 1.03 (m, 6H).	Off white solid

Example 5

5

Compound 22

([{(2S)-2-({[4-bromophenyl]amino]carbonyl]amino)pentanoyl]amino)methyl]phosphonic acid




10 A solution of **Compound 6** (100 mg, 0.21 mmol), 5 mL dichloromethane, and bromotrimethylsilane (0.16 mL, 0.63 mmol) was stirred at 25 °C for 3 hours. The resulting

mixture was quenched with ice then filtered. The filter cake was washed with 4 times with acetone : hexane (25:75) to yield **Compound 22** as a yellow solid.

¹H NMR (CD₃OD, 300MHz) δ: 7.26 - 7.39 (m, 2H), 4.28 (dd, J = 8.4, 5.4 Hz, 1H), 3.49 - 3.72 (m, 2H), 1.71 - 1.86 (m, 1H), 1.54 - 1.70 (m, 1H), 1.34 - 1.54 (m, 2H), 0.92 - 1.01 (m, 3H).

Compound 23 was prepared from the corresponding urea derivative in a similar manner to the procedure described in **Example 5** for **Compound 22**. The results are described below in **Table 5**.


Table 5

Comp. No.	IUPAC name	Starting material	¹ H NMR δ (ppm)	Properties
23	[[{(2S)-4-methyl-2-[[{4-(trifluoromethyl)phenyl]amino}carbonyl]amino]pentanoyl]amino)methyl]phosphonic acid 	Compound 11	¹ H NMR (CD ₃ OD, 300MHz) δ: 7.53 (s, 4H), 4.36 (dd, J = 9.2, 4.8 Hz, 1H), 3.47 - 3.75 (m, 2H), 1.49 - 1.86 (m, 3H), 0.95 - 1.05 (m, 6H).	Off white solid

Example 6

Compound 24

Diethyl [[{(2S)-4-methyl-2-[[{4-(methylsulfonyl)phenyl]amino}carbonyl]amino]pentanoyl]amino)methyl]phosphonate



To a solution of **Compound 14** (162 mg, 0.36 mmol) and 8 mL of anhydrous dichloromethane at 0 °C was added mCPBA (157 mg, 0.9 mmol). The resulting mixture was stirred at 25 °C for 2 hours. The resulting product was purified by medium pressure liquid chromatography on silica gel using methanol : dichloromethane (1 : 9) to yield **Compound 24** as a white solid.

¹H NMR (CD₃OD, 300MHz) δ: 7.77 - 7.87 (m, 2H), 7.57 - 7.68 (m, 2H), 4.36 (dd, J = 8.8, 5.6 Hz, 1H), 4.05 - 4.22 (m, 4H), 3.56 - 3.87 (m, 2H), 3.07 (s, 3H), 1.74 (dd, J = 13.6, 6.9 Hz, 1H), 1.49 - 1.67 (m, 2H), 1.22 - 1.36 (m, 6H), 0.91 - 1.03 (m, 6H).

5

Biological Data

Biological activity of compounds according to Formula I is set forth in **Table 6**.

CHO-Ga16 cells stably expressing FPR2 were cultured in (F12, 10% FBS, 1% PSA, 400 μg/ml geneticin and 50 μg/ml hygromycin) and HEK- Gqi5 cells stable expressing FPR2 were cultured in (DMEM high glucose, 10% FBS, 1% PSA, 400 μg/ml geneticin and 50 μg/ml hygromycin). In general, the day before the experiment, 18,000 cells/well were placed in a 384-well clear bottom poly-d-lysine coated plate. The following day the screening compound-induced calcium activity was assayed on the FLIPR^{Tetra}. The drug plates were prepared in 384-well microplates using the EP3 and the MultiPROBE robotic liquid handling systems. Compounds were tested at concentrations ranging from 0.61 to 10,000 nM. Results are expressed as EC₅₀ (nM) and efficacy values.

Table 6

20


Compound IUPAC name	FPR2 Ga16-CHO EC ₅₀ (nM) (0.%eff)
(2S)-2-({[(4-Bromophenyl)amino]carbonyl}amino)-4-methyl-N-(1H-tetrazol-5-ylmethyl)pentanamide	2.3 (0.81)
Ethyl hydrogen ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-4-methylpentanoyl]amino}methyl)phosphonate	0.95 (0.88)
Diethyl ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-4-methylpentanoyl]amino}methyl)phosphonate	0.59 (1.0)
Diethyl ({[(2S,3S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-3-methylpentanoyl]amino}methyl)phosphonate	2.4 (1.0)
(2S)-2-({[(4-Bromophenyl)amino]carbonyl}amino)-4-methyl-N-[2-(1H-tetrazol-5-yl)ethyl]pentanamide	47.9 (1.0)
(2S)-2-({[(4-Bromophenyl)amino]carbonyl}amino)-N-[(3-hydroxyisoxazol-5-yl)methyl]-4-methylpentanamide	12 (1.0)
Ethyl hydrogen ({[(2S,3S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-3-methylpentanoyl]amino}methyl)phosphonate	3.2 (0.88)
Diethyl ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)pentanoyl]amino}methyl)phosphonate	0.29 (0.94)
Diethyl ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-3-phenylpropanoyl]amino}methyl)phosphonate	20 (0.98)

Diethyl (2-{{(2S)-2-{{(4-bromophenyl)amino}carbonyl}amino)-4-methylpentanoyl}amino}ethyl)phosphonate	1.8 (1.0)
Ethyl hydrogen ({{(2S)-2-{{(4-bromophenyl)amino}carbonyl}amino)pentanoyl}amino} methyl)phosphonate	1.0 (0.94)
Dipropyl ({{(2S)-2-{{(4-bromophenyl)carbonyl}amino}pentanoyl}amino}methyl)phosphonate	0.8 (0.94)
Ethyl hydrogen ({{(2S)-2-{{(4-bromophenyl)carbonyl}amino)-3-phenylpropanoyl}amino}methyl)phosphonate	16 (1.10)
{{(2S)-2-{{(4-Bromophenyl)carbonyl}amino}pentanoyl}amino}methyl)phosphonic acid	32 (0.94)
{{(2S)-2-{{(4-Bromophenyl)carbonyl}amino)-4-methylpentanoyl}amino}methanesulfonic acid	2.3 (0.91)
Propyl hydrogen {{(2-{{(4-bromophenyl)carbonyl}amino}pentanoyl}amino)methyl}phosphonate	1.87 (0.89)
Dipropyl ({{(2S)-2-{{(4-bromophenyl)carbonyl}amino)-4-methylpentanoyl}amino}methyl)phosphonate	4.0 (1.00)
Diethyl ({{(2S)-4-methyl-2-{{(4-(trifluoromethyl)phenyl)carbonyl}amino)pentanoyl}amino}methyl)phosphonate	8.0 (1.00)
Ethyl hydrogen ({{(2S)-4-methyl-2-{{(4-(trifluoromethyl)phenyl)carbonyl}amino)pentanoyl}amino}methyl)phosphonate	2.0 (0.95)
(2S)-4-Methyl-N-(1H-tetrazol-5-ylmethyl)-2-{{(4-(trifluoromethyl)phenyl)carbonyl}amino}pentanamide	4.0 (0.91)
{{(2S)-4-Methyl-2-{{(4-(trifluoromethyl)phenyl)carbonyl}amino)pentanoyl}amino}methanesulfonic acid	1.6 (1.00)
Diethyl ({{(2S)-4-methyl-2-{{(4-(methylsulfonyl)phenyl)carbonyl}amino)pentanoyl}amino}methyl)phosphonate	2.1 (0.90)
Diethyl ({{(2S)-4-methyl-2-{{(4-(methylsulfonyl)phenyl)carbonyl}amino)pentanoyl}amino}methyl)phosphonate	45 (1.12)
Ethyl hydrogen ({{(2S)-4-methyl-2-{{(4(methylsulfonyl)phenyl)carbonyl}amino)pentanoyl}amino}methyl)phosphonate	133 (0.99)
{{(2S)-4-Methyl-2-{{(4-(trifluoromethyl)phenyl)carbonyl}amino)pentanoyl}amino}methyl)phosphonic acid	64 (1.02)

What is claimed is:

1. A compound represented by Formula I, individual enantiomers, individual diastereoisomers, individual tautomers, individual zwitterions or a pharmaceutically acceptable salt thereof:

5



Formula I

wherein:

- 10 n is 0 or 1;

R¹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, -N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN or -NO₂;


R² is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN or -NO₂;

- 15 R³ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN, -NO₂, -CF₃, -S(O)R¹⁵ or -S(O)₂R¹⁶;

R⁴ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN or -NO₂;

- 20 R⁵ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, -N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN or -NO₂;

R⁶ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted heterocycle, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted C₆₋₁₀ aryl, substituted or unsubstituted C₃₋₈ cycloalkenyl or -CH₂R¹⁹;



S(O)R^{15} , $-\text{C(O)N(H)(CN)}$, $-\text{S(O)}_2\text{NHS(O)}_2\text{R}^{25}$, $-\text{C(O)N(H)S(O)}_2\text{R}^{19}$, $-\text{S(O)}_2\text{R}^{16}$ or $-\text{P(O)R}^{17}\text{R}^{18}$;

R⁸ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R⁹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R¹⁰ is hydrogen, or substituted or unsubstituted C₁₋₈ alkyl ;

R¹¹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl or -CF₃;

R¹² is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, -OR²⁴ or -NR⁸R⁹;

R¹³ is -OR²²;

R^{14} is $-OR^{23}$:

R¹⁵ is substituted or unsubstituted C₁₋₈ alkyl;

R¹⁶ is substituted or unsubstituted C₁₋₈ alkyl, -NR⁸R⁹, -NHS(O)₂R¹⁹ or hydroxyl;

R¹⁷ is -OR¹⁰ or -NR⁸R⁹;

R^{18} is $-OR^{10}$ or $-NR^8R^9$:

R¹⁹ is substituted or unsubstituted heterocycle, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted C₆₋₁₀ aryl or substituted or unsubstituted C₃₋₈ cycloalkenyl;

R²⁰ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀

5 aryl;

R²¹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R²² is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, or together with R²³ can form a substituted or unsubstituted cycle;

R²³ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, or together with R²² can form a substituted or unsubstituted cycle;

R²⁴ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀

15 aryl; and

R²⁵ is substituted or unsubstituted aromatic heterocycle, or substituted or unsubstituted aromatic C₆₋₁₀ aryl;

R²⁶ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀


20 aryl;


R²⁷ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R²⁸ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R²⁹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl; with the proviso:

30 the compound of Formula I is not of structures:






5

2. A compound according to claim 1, wherein:

R^3 is substituted or unsubstituted C_{1-8} alkyl, halogen, $-SR^{11}$, $-CF_3$ or $-S(O)_2R^{16}$.

3. A compound according to claim 1, wherein:



4. A compound according to claim 1, wherein:

R^6 is substituted or unsubstituted C_{1-8} alkyl, or $-CH_2R^{19}$.

15

5. A compound according to claim 1, wherein:

R^{26} is hydrogen;

R^{27} is hydrogen;

R^{28} is hydrogen; and

20 R^{29} is hydrogen;

6. A compound according to claim 1, wherein:

n is 0 or 1;

R¹ is hydrogen;

5 R² is hydrogen;


R³ is substituted or unsubstituted C₁₋₈ alkyl, halogen, -SR¹¹, -CF₃ or S(O)₂R¹⁶;

R⁴ is hydrogen;

R⁵ is hydrogen;

R⁶ is substituted or unsubstituted C₁₋₈ alkyl or -CH₂R¹⁹;

10



7. A compound according to claim 1, wherein:

n is 0;

20 R¹ is hydrogen;


R² is hydrogen;

R³ is substituted or unsubstituted C₁₋₈ alkyl, halogen, -SR¹¹, CF₃ or S(O)₂R¹⁶;

R⁴ is hydrogen;

R⁵ is hydrogen;

25 R⁶ is substituted or unsubstituted C₁₋₈ alkyl or -CH₂R¹⁹;



R^7 is , $-S(O)_2R^{16}$ or $-P(O)R^{17}R^{18}$;

R^{11} is substituted or unsubstituted C_{1-8} alkyl;

R^{16} is substituted or unsubstituted C_{1-8} alkyl or hydroxyl;

5 R^{17} is OR^{10} ;

R^{18} is OR^{10} ; and

R^{19} is substituted or unsubstituted C_{6-10} aryl.

8. A compound according to claim 1 selected from:

10 Diethyl ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-4-methylpentanoyl]
amino}methyl)phosphonate;

(2S)-2-({[(4-Bromophenyl)amino]carbonyl}amino)-4-methyl-N-(1H-tetrazol-5-
ylmethyl)pentanamide;

15 (2S)-2-({[(4-Bromophenyl)amino]carbonyl}amino)-4-methyl-N-[2-(1H-tetrazol-5-
yl)ethyl]pentanamide;

(2S)-2-({[(4-Bromophenyl)amino]carbonyl}amino)-N-[(3-hydroxyisoxazol-5-yl)methyl]-4-
methylpentanamide;

Diethyl ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-4-methylpentanoyl]
amino}methyl)phosphonate;

20 Diethyl ({[(2S,3S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-3-methylpentanoyl]
amino}methyl)phosphonate;

Diethyl ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino) pentanoyl]amino}
methyl)phosphonate;

25 Diethyl ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-3-phenylpropanoyl]
amino}methyl)phosphonate;

Diethyl (2-{{[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-4-methylpentanoyl]
amino}ethyl)phosphonate;

Ethyl hydrogen ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-4-methyl
pentanoyl]amino}methyl)phosphonate;

Ethyl hydrogen ({[(2S,3S)-2-({[(4-bromophenyl)amino]carbonyl}amino)-3-methyl
pentanoyl]amino}methyl)phosphonate;

Ethyl hydrogen ({[(2S)-2-({[(4-bromophenyl)amino]carbonyl}amino) pentanoyl]amino}
methyl)phosphonate;

5 ({[(2S)-4-Methyl-2-({[4-(trifluoromethyl)phenyl]carbamoyl}amino) pentanoyl]
amino}methyl)phosphonic acid;

Ethyl hydrogen ({[(2S)-4-methyl-2-({[4-(methylsulfonyl)phenyl]carbamoyl}amino)
pentanoyl]amino}methyl)phosphonate;

10 Diethyl ({[(2S)-4-methyl-2-({[4-(methylsulfonyl)phenyl]carbamoyl}amino) pentanoyl]
amino}methyl)phosphonate;

Diethyl ({[(2S)-4-methyl-2-({[4-(methylsulfonyl)phenyl]carbamoyl}amino)pentanoyl]
amino}methyl)phosphonate;

{[(2S)-4-Methyl-2-({[4-(trifluoromethyl)phenyl]carbamoyl}amino) pentanoyl]
amino}methanesulfonic acid;

15 (2S)-4-Methyl-N-(1H-tetrazol-5-ylmethyl)-2-({[4-(trifluoromethyl)phenyl] carbamoyl}
amino)pentanamide;

Ethyl hydrogen ({[(2S)-4-methyl-2-({[4-(trifluoromethyl)phenyl]carbamoyl}amino)
pentanoyl]amino}methyl)phosphonate;

20 Diethyl ({[(2S)-4-methyl-2-({[4-(trifluoromethyl)phenyl]carbamoyl}amino) pentanoyl]
amino}methyl)phosphonate;

Dipropyl ({[(2S)-2-({[(4-bromophenyl)carbamoyl]amino}-4-methylpentanoyl] amino}
methyl)phosphonate;

Propan-2-yl hydrogen {[2-({[(4-bromophenyl)carbamoyl]amino}pentanoyl) amino]
methyl}phosphonate;

25 {[2-({[(4-Bromophenyl)carbamoyl]amino}-4-methylpentanoyl] amino}
methanesulfonic acid;

{[(2S)-2-({[(4-Bromophenyl)carbamoyl]amino}pentanoyl]amino) methyl}phosphonic
acid;

30 Ethyl hydrogen ({[(2S)-2-({[(4-bromophenyl)carbamoyl]amino}-3-phenylpropanoyl]
amino}methyl)phosphonate; and

Dipropyl ({[(2S)-2-({[(4-bromophenyl)carbamoyl]amino}pentanoyl] amino) methyl}
phosphonate.

9. A pharmaceutical composition comprising as active ingredient a therapeutically effective amount of a compound according to claim 1 and a pharmaceutically acceptable adjuvant, diluent or carrier.

10. A pharmaceutical composition according to claim 9 wherein the compound is selected from:

Diethyl (((2S)-2-(((4-bromophenyl)amino)carbonyl)amino)-4-methylpentanoyl]amino)methyl)phosphonate;

(2S)-2-(((4-Bromophenyl)amino)carbonyl)amino)-4-methyl-N-(1H-tetrazol-5-ylmethyl)pentanamide;

(2S)-2-(((4-Bromophenyl)amino)carbonyl)amino)-4-methyl-N-[2-(1H-tetrazol-5-yl)ethyl]pentanamide;

(2S)-2-(((4-Bromophenyl)amino)carbonyl)amino)-N-[(3-hydroxyisoxazol-5-yl)methyl]-4-methylpentanamide;

Diethyl (((2S)-2-(((4-bromophenyl)amino)carbonyl)amino)-4-methylpentanoyl]amino)methyl)phosphonate;

Diethyl (((2S,3S)-2-(((4-bromophenyl)amino)carbonyl)amino)-3-methylpentanoyl]amino)methyl)phosphonate;

Diethyl (((2S)-2-(((4-bromophenyl)amino)carbonyl)amino) pentanoyl]amino)methyl)phosphonate;

Diethyl (((2S)-2-(((4-bromophenyl)amino)carbonyl)amino)-3-phenylpropanoyl]amino)methyl)phosphonate;

Diethyl (2-(((2S)-2-(((4-bromophenyl)amino)carbonyl)amino)-4-methylpentanoyl]amino)ethyl)phosphonate;

Ethyl hydrogen (((2S)-2-(((4-bromophenyl)amino)carbonyl)amino)-4-methylpentanoyl]amino)methyl)phosphonate;


Ethyl hydrogen (((2S,3S)-2-(((4-bromophenyl)amino)carbonyl)amino)-3-methylpentanoyl]amino)methyl)phosphonate;

Ethyl hydrogen (((2S)-2-(((4-bromophenyl)amino)carbonyl)amino) pentanoyl]amino)methyl)phosphonate;

((((2S)-4-Methyl-2-([4-(trifluoromethyl)phenyl]carbamoyl)amino) pentanoyl]amino)methyl)phosphonic acid;

Ethyl hydrogen (((2S)-4-methyl-2-([4-(methylsulfonyl)phenyl]carbamoyl)amino) pentanoyl]amino)methyl)phosphonate;


- Diethyl ({{{(2S)-4-methyl-2-({[4-(methylsulfonyl)phenyl]carbamoyl}amino) pentanoyl] amino}methyl)phosphonate;
- Diethyl ({{{(2S)-4-methyl-2-({[4-(methylsulfanyl)phenyl]carbamoyl}amino)pentanoyl] amino}methyl)phosphonate;
- 5 {{{(2S)-4-Methyl-2-({[4-(trifluoromethyl)phenyl]carbamoyl}amino) pentanoyl] amino}methanesulfonic acid;
- (2S)-4-Methyl-N-(1H-tetrazol-5-ylmethyl)-2-({[4-(trifluoromethyl)phenyl] carbamoyl} amino)pentanamide;
- Ethyl hydrogen ({{{(2S)-4-methyl-2-({[4-(trifluoromethyl)phenyl]carbamoyl}amino) pentanoyl]amino}methyl)phosphonate;
- 10 Diethyl ({{{(2S)-4-methyl-2-({[4-(trifluoromethyl)phenyl]carbamoyl}amino) pentanoyl] amino}methyl)phosphonate;
- Dipropyl 2-yl ({{{(2S)-2-({[4-(bromophenyl)carbamoyl]amino}-4-methylpentanoyl] amino} methyl)phosphonate;
- 15 Propan-2-yl hydrogen {{{(2S)-2-({[4-(bromophenyl)carbamoyl]amino}pentanoyl] amino} methyl)phosphonate;
- {{{(2S)-2-({[4-Bromophenyl]carbamoyl]amino}-4-methylpentanoyl] amino} methanesulfonic acid;
- {{{(2S)-2-({[4-Bromophenyl]carbamoyl]amino}pentanoyl]amino} methyl)phosphonic acid;
- 20 Ethyl hydrogen ({{{(2S)-2-({[4-(bromophenyl)carbamoyl]amino}-3-phenylpropanoyl] amino}methyl)phosphonate; and
- Dipropyl 2-yl ({{{(2S)-2-({[4-(bromophenyl)carbamoyl]amino}pentanoyl] amino} methyl) phosphonate.
- 25 11. A method of treating a disorder associated with N-formyl peptide receptor 2 modulation, which comprises administering to a mammal in need thereof, a pharmaceutical composition comprising a therapeutically effective amount of at least one compound of Formula I



Formula I

wherein:

- 5 n is 0 or 1;
- R¹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, -N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN or -NO₂;
- R² is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN or -NO₂;
- 10 R³ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN, -NO₂, -CF₃, -S(O)R¹⁵ or -S(O)₂R¹⁶;
- R⁴ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN or -NO₂;
- R⁵ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, halogen, -NR⁸R⁹, -N(R²¹)C(O)R²⁰, -OR¹⁰, -OC(O)R²¹, -SR¹¹, -C(O)R¹², -CN or -NO₂;
- 15 R⁶ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted heterocycle, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted C₆₋₁₀ aryl, substituted or unsubstituted C₃₋₈ cycloalkenyl or -CH₂R¹⁹;



S(O)R^{15} , $-\text{C(O)N(H)(CN)}$, $-\text{S(O)}_2\text{NHS(O)}_2\text{R}^{25}$, $-\text{C(O)N(H)S(O)}_2\text{R}^{19}$, $-\text{S(O)}_2\text{R}^{16}$ or $-\text{P(O)R}^{17}\text{R}^{18}$;

R⁸ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R⁹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R¹⁰ is hydrogen, or substituted or unsubstituted C₁₋₈ alkyl ;

R¹¹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl or -CF₃;

R¹² is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, -OR²⁴ or -NR⁸R⁹;

R¹³ is -OR²²;

R^{14} is $-OR^{23}$:

R¹⁵ is substituted or unsubstituted C₁₋₈ alkyl;

R¹⁶ is substituted or unsubstituted C₁₋₈ alkyl, -NR⁸R⁹, -NHS(O)₂R¹⁹ or hydroxyl;

R¹⁷ is -OR¹⁰ or -NR⁸R⁹:

R¹⁸ is -OR¹⁰ or -NR⁸R⁹:

R¹⁹ is substituted or unsubstituted heterocycle, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted C₆₋₁₀ aryl or substituted or unsubstituted C₃₋₈ cycloalkenyl;

R²⁰ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀

5 aryl;

R²¹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R²² is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, or together with R²³ can form a substituted or unsubstituted cycle;

R²³ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, or together with R²² can form a substituted or unsubstituted cycle;

R²⁴ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀

15 aryl; and

R²⁵ is substituted or unsubstituted aromatic heterocycle, or substituted or unsubstituted aromatic C₆₋₁₀ aryl;

R²⁶ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀


20 aryl;


R²⁷ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R²⁸ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl;

R²⁹ is hydrogen, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted heterocycle, or substituted or unsubstituted C₆₋₁₀ aryl; with the proviso:

30 the compound of Formula I is not of structures:





5

12. A method of claim 11, wherein the pharmaceutical composition is administered to the mammal to treat ocular inflammatory diseases including, but not limited to, uveitis, dry eye, Keratitis, allergic eye disease and conditions affecting the posterior part of the eye, such as maculopathies and retinal degeneration including non-exudative age related macular degeneration, exudative age related macular degeneration, choroidal neovascularization, diabetic retinopathy, acute macular neuroretinopathy, central serous chorioretinopathy, post-surgery corneal wound healing, cystoid macular edema, and diabetic macular edema; infectious keratitis, herpetic keratitis, corneal angiogenesis, lymphangiogenesis, retinitis, and choroiditis such as acute multifocal placoid pigment epitheliopathy, Behcet's disease, birdshot retinochoroidopathy, infectious (syphilis, lyme, tuberculosis, toxoplasmosis), intermediate uveitis (pars planitis), multifocal choroiditis, multiple evanescent white dot syndrome (mewds), ocular sarcoidosis, posterior scleritis, serpiginous choroiditis, subretinal fibrosis, Vogt-Koyanagi-and Harada syndrome; vasuclar diseases/ exudative diseases such as retinal arterial occlusive disease, central retinal vein occlusion, disseminated intravascular coagulopathy, branch retinal vein occlusion, hypertensive fundus changes, ocular ischemic syndrome, retinal arterial microaneurysms, Coat's disease, parafoveal telangiectasis, hemi-retinal vein occlusion, papillophlebitis, central retinal artery occlusion, branch retinal artery occlusion, carotid artery disease (CAD), frosted branch angiitis, sickle cell retinopathy and other hemoglobinopathies,

20

angioid streaks, familial exudative vitreoretinopathy, and Eales disease; traumatic/ surgical conditions such as sympathetic ophthalmia, uveitic retinal disease, retinal detachment, trauma, conditions caused by laser, conditions caused by photodynamic therapy, photocoagulation, hypoperfusion during surgery, radiation retinopathy, and bone marrow transplant retinopathy; proliferative disorders such as proliferative vitreal retinopathy and epiretinal membranes, and proliferative diabetic retinopathy; infectious disorders such as ocular histoplasmosis, ocular toxocariasis, presumed ocular histoplasmosis syndrome (POHS), endophthalmitis, toxoplasmosis, retinal diseases associated with HIV infection, choroidal disease associated with HIV infection, uveitic disease associated with HIV infection, viral retinitis, acute retinal necrosis, progressive outer retinal necrosis, fungal retinal diseases, ocular syphilis, ocular tuberculosis, diffuse unilateral subacute neuroretinitis, and myiasis; genetic disorders such as retinitis pigmentosa, systemic disorders with associated retinal dystrophies, congenital stationary night blindness, cone dystrophies, Stargardt's disease and fundus flavimaculatus, Best's disease, pattern dystrophy of the retinal pigmented epithelium, X-linked retinoschisis, Sorsby's fundus dystrophy, benign concentric maculopathy, Bietti's crystalline dystrophy, and pseudoxanthoma elasticum; retinal tears/ holes such as retinal detachment, macular hole, and giant retinal tear; tumors such as retinal disease associated with tumors, congenital hypertrophy of the retinal pigmented epithelium, posterior uveal melanoma, choroidal hemangioma, choroidal osteoma, choroidal metastasis, combined hamartoma of the retina and retinal pigmented epithelium, retinoblastoma, vasoproliferative tumors of the ocular fundus, retinal astrocytoma, and intraocular lymphoid tumors; and miscellaneous other diseases affecting the posterior part of the eye such as punctate inner choroidopathy, acute posterior multifocal placoid pigment epitheliopathy, myopic retinal degeneration, and acute retinal pigment epitheliitis; Systemic inflammatory diseases such as stroke, coronary artery disease, obstructive airway diseases, HIV-mediated retroviral infections, cardiovascular disorders including coronary artery disease, neuroinflammation, neurological disorders, pain and immunological disorders, asthma, allergic disorders, inflammation, systemic lupus erythematosus, psoriasis, CNS disorders such as Alzheimer's disease, arthritis, sepsis, inflammatory bowel disease, cachexia, angina pectoris.

13. The method of claim 12 wherein the mammal is a human.